




A Book Concerning Statistical Signal Processing



VERSION 0.14X



+




=




○ zero of $L^*(\frac{1}{z^*)}$

○ zero of $L(z)$


\Re




Daniel J. Greenhoe



Signal Processing ABCs series
volume 2









title: *A Book Concerning Statistical Signal Processing*
document type: book
series: *Signal Processing ABCs*
volume: 2
author: Daniel J. Greenhoe
version: VERSION 0.14X
time stamp: 2019 July 26 (Friday) 06:48pm UTC
copyright: Copyright © 2019 Daniel J. Greenhoe
license: Creative Commons license CC BY-NC-ND 4.0
typesetting engine: X_ET_EX
document url: <https://www.researchgate.net/project/Signal-Processing-ABCs>

This text was typeset using X_E^AT_EX, which is part of the T_EXfamily of typesetting engines, which is arguably the greatest development since the Gutenberg Press. Graphics were rendered using the *pstricks* and related packages, and L_AT_EX graphics support.

The main roman, *italic*, and **bold** font typefaces used are all from the *Heuristica* family of typefaces (based on the *Utopia* typeface, released by *Adobe Systems Incorporated*). The math font is XITS from the XITS font project. The font used in quotation boxes is adapted from *Zapf Chancery Medium Italic*, originally from URW++ Design and Development Incorporated. The font used for the text in the title is Adventor (similar to *Avant-Garde*) from the T_EX-Gyre Project. The font used for the ISBN in the footer of individual pages is LIQUID CRYSTAL (*Liquid Crystal*) from *FontLab Studio*. The Latin handwriting font is *Lavi* from the *Free Software Foundation*.

The ship appearing throughout this text is loosely based on the *Golden Hind*, a sixteenth century English galleon famous for circumnavigating the globe.¹



¹  Paine (2000) page 63 (Golden Hind)

“Here, on the level sand,
Between the sea and land,
What shall I build or write
Against the fall of night? ”



“Tell me of runes to grave
That hold the bursting wave,
Or bastions to design
For longer date than mine. ”

Alfred Edward Housman, English poet (1859–1936) ²



“The uninitiated imagine that one must await inspiration in order to create. That is a mistake. I am far from saying that there is no such thing as inspiration; quite the opposite. It is found as a driving force in every kind of human activity, and is in no wise peculiar to artists. But that force is brought into action by an effort, and that effort is work. Just as appetite comes by eating so work brings inspiration, if inspiration is not discernible at the beginning. ”

Igor Fyodorovich Stravinsky (1882–1971), Russian-born composer ³



“As I think about acts of integrity and grace, I realise that there is nothing in my knowledge to compare with Frege's dedication to truth. His entire life's work was on the verge of completion, much of his work had been ignored to the benefit of men infinitely less capable, his second volume was about to be published, and upon finding that his fundamental assumption was in error, he responded with intellectual pleasure clearly submerging any feelings of personal disappointment. It was almost superhuman and a telling indication of that of which men are capable if their dedication is to creative work and knowledge instead of cruder efforts to dominate and be known. ”

Bertrand Russell (1872–1970), British mathematician, in a 1962 November 23 letter to Dr. van Heijenoort. ⁴



² quote:  Housman (1936), page 64 (“Smooth Between Sea and Land”),  Hardy (1940) (section 7)

image: <http://en.wikipedia.org/wiki/Image:Housman.jpg>

³ quote:  Ewen (1961), page 408,  Ewen (1950)

image: http://en.wikipedia.org/wiki/Image:Igor_Stravinsky.jpg

⁴ quote:  Heijenoort (1967), page 127

image: <http://www-history.mcs.st-andrews.ac.uk/PictDisplay/Russell.html>

SYMBOLS

“*rugula XVI. Quae vero praesentem mentis attentionem non requirunt, etiamsi ad conclusionem necessaria sint, illa melius est per brevissimas notas designare quam per integras figuras: ita enim memoria non poterit falli, nec tamen interim cogitatio distrahetur ad haec retinenda, dum aliis deducendis incumbit.*”



“*Rule XVI. As for things which do not require the immediate attention of the mind, however necessary they may be for the conclusion, it is better to represent them by very concise symbols rather than by complete figures. It will thus be impossible for our memory to go wrong, and our mind will not be distracted by having to retain these while it is taken up with deducing other matters.*”

René Descartes (1596–1650), French philosopher and mathematician ⁵



“*In signs one observes an advantage in discovery which is greatest when they express the exact nature of a thing briefly and, as it were, picture it; then indeed the labor of thought is wonderfully diminished.*”

Gottfried Leibniz (1646–1716), German mathematician, ⁶

Symbol list

symbol	description
numbers:	
\mathbb{Z}	integers
\mathbb{W}	whole numbers
\mathbb{N}	natural numbers
\mathbb{Z}^+	non-positive integers

...continued on next page...

⁵quote: [Descartes \(1684a\)](#) (rugula XVI), translation: [Descartes \(1684b\)](#) (rule XVI), image: Frans Hals (circa 1650), <http://en.wikipedia.org/wiki/Descartes>, public domain

⁶quote: [Cajori \(1993\)](#) (paragraph 540), image: http://en.wikipedia.org/wiki/File:Gottfried_Wilhelm_von_Leibniz.jpg, public domain

symbol	description
\mathbb{Z}^-	negative integers $\dots, -3, -2, -1$
\mathbb{Z}_o	odd integers $\dots, -3, -1, 1, 3, \dots$
\mathbb{Z}_e	even integers $\dots, -4, -2, 0, 2, 4, \dots$
\mathbb{Q}	rational numbers $\frac{m}{n}$ with $m \in \mathbb{Z}$ and $n \in \mathbb{Z} \setminus 0$
\mathbb{R}	real numbers completion of \mathbb{Q}
\mathbb{R}^+	non-negative real numbers $[0, \infty)$
\mathbb{R}^+	non-positive real numbers $(-\infty, 0]$
\mathbb{R}^+	positive real numbers $(0, \infty)$
\mathbb{R}^-	negative real numbers $(-\infty, 0)$
\mathbb{R}^*	extended real numbers $\mathbb{R}^* \triangleq \mathbb{R} \cup \{-\infty, \infty\}$
\mathbb{C}	complex numbers
\mathbb{F}	arbitrary field (often either \mathbb{R} or \mathbb{C})
∞	positive infinity
$-\infty$	negative infinity
π	pi 3.14159265 ...
relations:	
\circledcirc	relation
$\circledcirc \circ$	relational and
$X \times Y$	Cartesian product of X and Y
(Δ, ∇)	ordered pair
$ z $	absolute value of a complex number z
$=$	equality relation
\triangleq	equality by definition
\rightarrow	maps to
\in	is an element of
\notin	is not an element of
$D(\circledcirc)$	domain of a relation \circledcirc
$I(\circledcirc)$	image of a relation \circledcirc
$R(\circledcirc)$	range of a relation \circledcirc
$N(\circledcirc)$	null space of a relation \circledcirc
set relations:	
\subseteq	subset
\subsetneq	proper subset
\supseteq	super set
\supsetneq	proper superset
$\not\subseteq$	is not a subset of
$\not\subsetneq$	is not a proper subset of
operations on sets:	
$A \cup B$	set union
$A \cap B$	set intersection
$A \Delta B$	set symmetric difference
$A \setminus B$	set difference
A^c	set complement
$ \cdot $	set order
$\mathbb{1}_A(x)$	set indicator function or characteristic function
logic:	
1	“true” condition
0	“false” condition
\neg	logical NOT operation

...continued on next page...

symbol	description
\wedge	logical AND operation
\vee	logical inclusive OR operation
\oplus	logical exclusive OR operation
\Rightarrow	“implies”;
\Leftarrow	“implied by”;
\Leftrightarrow	“if and only if”;
\forall	universal quantifier:
\exists	existential quantifier:
order on sets:	
\vee	join or least upper bound
\wedge	meet or greatest lower bound
\leq	reflexive ordering relation
\geq	reflexive ordering relation
$<$	irreflexive ordering relation
$>$	irreflexive ordering relation
measures on sets:	
$ X $	order or counting measure of a set X
distance spaces:	
d	metric or distance function
linear spaces:	
$\ \cdot\ $	vector norm
$\ \cdot\ $	operator norm
$\langle \Delta \nabla \rangle$	inner-product
$\text{span}(V)$	span of a linear space V
algebras:	
\Re	real part of an element in a $*$ -algebra
\Im	imaginary part of an element in a $*$ -algebra
set structures:	
T	a topology of sets
R	a ring of sets
A	an algebra of sets
\emptyset	empty set
2^X	power set on a set X
sets of set structures:	
$\mathcal{T}(X)$	set of topologies on a set X
$\mathcal{R}(X)$	set of rings of sets on a set X
$\mathcal{A}(X)$	set of algebras of sets on a set X
classes of relations/functions/operators:	
2^{XY}	set of <i>relations</i> from X to Y
Y^X	set of <i>functions</i> from X to Y
$S_j(X, Y)$	set of <i>surjective</i> functions from X to Y
$I_j(X, Y)$	set of <i>injective</i> functions from X to Y
$B_j(X, Y)$	set of <i>bijective</i> functions from X to Y
$B(X, Y)$	set of <i>bounded</i> functions/operators from X to Y
$L(X, Y)$	set of <i>linear bounded</i> functions/operators from X to Y
$C(X, Y)$	set of <i>continuous</i> functions/operators from X to Y
specific transforms/operators:	
\tilde{F}	<i>Fourier Transform</i> operator (Definition N.2 page 309)
\hat{F}	<i>Fourier Series</i> operator

...continued on next page...

symbol	description
$\tilde{\mathbf{F}}$	<i>Discrete Time Fourier Series operator</i> (Definition O.1 page 319)
\mathbf{Z}	<i>Z-Transform operator</i> (Definition P.4 page 330)
$\tilde{f}(\omega)$	<i>Fourier Transform of a function $f(x) \in L^2_{\mathbb{R}}$</i>
$\check{x}(\omega)$	<i>Discrete Time Fourier Transform of a sequence $(x_n \in \mathbb{C})_{n \in \mathbb{Z}}$</i>
$\check{x}(z)$	<i>Z-Transform of a sequence $(x_n \in \mathbb{C})_{n \in \mathbb{Z}}$</i>

SYMBOL INDEX

$\bar{\delta}_n$, 245	x^+ , 192	span , 197	$[\cdot : \cdot)$, 259, 260
$(A, \ \cdot\ , *)$, 231	d , 195	$*$, 228	$[\cdot : \cdot]$, 259, 260
$\ \cdot\ $, 249	I_m , 229	\geq , 308	Y^X , 282
\perp , 194	I , 282	$\ \cdot\ $, 286	ρ , 170, 227
\star , 329	R_e , 229	\odot , 256, 258	σ_c , 170
σ , 343	Z , 330	\star , 312	σ_p , 170
\tilde{F} , 310	$N_n(x)$, 343	$B(X, Y)$, 289	σ_r , 170
$ x $, 192	$\text{epi}(f)$, 261	$(\cdot : \cdot)$, 259, 260	σ , 170, 227
x^- , 192	$\text{hyp}(f)$, 261	$(\cdot : \cdot]$, 259, 260	r , 227

CONTENTS

Title page	v
Typesetting	vi
Quotes	vii
Symbol list	ix
Symbol index	xiii
Contents	xv
I Statistical Analysis	1
1 Expectation operator	3
1.1 Definitions	3
1.2 Expectation as a linear operator	4
1.3 Expectation inequalities	6
1.4 Joint and conditional probability spaces	7
1.5 Expectation inner product space	8
2 Random Sequences	11
2.1 Definitions	11
2.2 Properties	12
2.3 Wide Sense Stationary processes	12
2.4 Spectral density	14
2.5 Spectral Power	16
3 Continuous Random Processes	17
3.1 Definitions	17
3.2 Properties	18
3.3 Basis for random processes	20
II Statistical Processing	23
4 Operations on Random Variables	25
4.1 Functions of one random variable	25
4.2 Functions of two random variables	31
5 Operators on Discrete Random Sequences	35
5.1 LTI operators on random sequences	35
5.2 LTI operators on WSS random sequences	36
5.3 Parallel operators on WSS random sequences	37
5.4 Whitening discrete random sequences	38
6 Operators on Continuous Random Sequences	41
6.1 LTI Operations on non-stationary random processes	41
6.2 LTI Operations on WSS random processes	43
6.3 Whitening continuous random sequences	48

7 Additive noise on random sequences	51
7.1 Additive noise and correlation	51
7.2 Additive noise and operators	53
7.3 Additive noise and LTI operators	56
7.4 Additive noise and dual operators	58
III Statistical Estimation	61
8 Estimation Theory	63
8.1 Estimation criterion	64
8.2 Estimation techniques	65
8.2.1 Sequential decoding	65
8.2.2 Norm minimization	66
8.2.3 Gradient search techniques	71
8.2.4 Direct search	73
9 KL-expansion application	75
9.1 Sufficient statistics	75
9.2 Optimal symbol estimation	80
9.3 Colored noise	87
9.4 Signal matching	88
10 Moment Estimation	91
11 System Identification	93
11.1 Estimation techniques	93
11.2 Additive noise system models	94
11.3 Transfer function estimate definitions and interpretation	95
11.4 Estimator relationships	100
11.5 Alternate forms	105
11.6 Least squares estimates of non-linear systems	107
11.7 Least squares estimates of linear systems	111
11.8 Coherence	115
11.8.1 Application	115
11.8.2 Definitions	116
11.8.3 A warning	116
12 Estimating Noise	119
13 Communication channels	121
13.1 System model	121
13.1.1 Channel operator	122
13.1.2 Receive operator	122
13.2 Optimization in the case of additional operations	123
13.3 Channel Statistics	124
14 Optimal Symbol Detection	125
14.1 ML Estimation	125
14.2 Generalized coherent modulation	126
14.3 Frequency Shift Keying (FSK)	127
14.4 Quadrature Amplitude Modulation (QAM)	129
14.4.1 Receiver statistics	129
14.4.2 Detection	130
14.4.3 Probability of error	130
14.5 Phase Shift Keying (PSK)	131
14.5.1 Receiver statistics	131
14.5.2 Detection	133
14.5.3 Probability of error	133
14.6 Pulse Amplitude Modulation (PAM)	134
14.6.1 Receiver statistics	134
14.6.2 Detection	134


14.6.3 Probability of error	135
15 Network Detection	137
15.1 Detection	137
15.2 Bayesian Estimation	137
15.3 Joint Gaussian Model	138
15.4 2 hypothesis, 2 sensor detection	139
IV Appendices	143
A Probability Space	145
A.1 Probability functions	146
A.2 Probability Space	147
A.3 Properties	148
A.4 Examples	150
A.5 Probability subspaces	153
B Probability Density Functions	155
B.1 Random variables	155
B.2 Probability distributions	156
B.3 Properties	156
C Some Probability Density Functions	161
C.1 Discrete distributions	161
C.2 Continuous distributions	161
C.2.1 Uniform distribution	161
C.2.2 Gaussian distribution	162
C.2.3 Gamma distribution	163
C.2.4 Chi-squared distributions	164
C.2.5 Radial distributions	167
C.3 Joint Gaussian distributions	167
D Spectral Theory	169
D.1 Operator Spectrum	169
D.2 Fredholm kernels	171
E Matrix Calculus	173
E.1 First derivative of a vector with respect to a vector	173
E.2 First derivative of a matrix with respect to a scalar	180
E.3 Second derivative of a scalar with respect to a vector	182
E.4 Multiple derivatives of a vector with respect to a scalar	183
F Linear spaces	185
F.1 Definition and basic results	185
F.2 Order on Linear Spaces	190
G Linear Combinations	197
G.1 Linear combinations in linear spaces	197
G.2 Bases in topological linear spaces	201
G.3 Schauder bases in Banach spaces	202
G.4 Linear combinations in inner product spaces	204
G.5 Orthonormal bases in Hilbert spaces	207
G.6 Riesz bases in Hilbert spaces	215
G.7 Frames in Hilbert spaces	220
H Normed Algebras	227
H.1 Algebras	227
H.2 Star-Algebras	228
H.3 Normed Algebras	231
H.4 C* Algebras	231

I Inner Product Spaces	233
I.1 Definition and basic results	233
I.2 Relationship between norms and inner products	237
I.2.1 Norms induced by inner products	237
I.2.2 Inner products induced by norms	239
I.3 Orthogonality	245
J Normed Linear Spaces	249
J.1 Definition and basic results	249
J.2 Relationship between metrics and norms	251
J.2.1 Metrics generated by norms	251
J.2.2 Norms generated by metrics	254
J.3 Orthogonality on normed linear spaces	256
K Intervals and Convexity	259
K.1 Intervals	259
K.2 Convex sets	260
K.3 Convex functions	261
K.4 Literature	264
L Finite Sums	267
L.1 Summation	267
L.2 Means	269
L.2.1 Weighted ϕ -means	269
L.2.2 Power means	270
L.3 Power Sums	279
M Operators on Linear Spaces	281
M.1 Operators on linear spaces	281
M.1.1 Operator Algebra	281
M.1.2 Linear operators	282
M.2 Operators on Normed linear spaces	286
M.2.1 Operator norm	286
M.2.2 Bounded linear operators	289
M.2.3 Adjoints on normed linear spaces	291
M.2.4 More properties	292
M.3 Operators on Inner product spaces	293
M.3.1 General Results	293
M.3.2 Operator adjoint	294
M.4 Special Classes of Operators	296
M.4.1 Projection operators	296
M.4.2 Self Adjoint Operators	298
M.4.3 Normal Operators	299
M.4.4 Isometric operators	300
M.4.5 Unitary operators	303
M.5 Operator order	308
N Fourier Transform	309
N.1 Definitions	309
N.2 Operator properties	310
N.3 Convolution	312
N.4 Real valued functions	313
N.5 Moment properties	314
N.6 Examples	316
O Discrete Time Fourier Transform	319
O.1 Definition	319
O.2 Properties	319
O.3 Derivatives	327
P Operations on Sequences	329
P.1 Convolution operator	329

P.2	Z-transform	330
P.3	From z-domain back to time-domain	332
P.4	Zero locations	333
P.5	Pole locations	334
P.6	Mirroring for real coefficients	335
P.7	Rational polynomial operators	336
P.8	Filter Banks	337
P.9	Inverting non-minimum phase filters	341
Q	B-Splines	343
Q.1	Definitions	343
Q.2	Algebraic properties	347
Q.3	Projection properties	356
Q.4	Fourier analysis	357
Q.5	Basis properties	359
Q.5.1	Uniqueness properties	359
Q.5.2	Partition of unity properties	360
Q.5.3	Riesz basis properties	361
Q.6	Mutiresolution properties	367
Q.6.1	Introduction	367
Q.6.2	B-spline dyadic decomposition	367
Q.6.3	B-spline MRA scaling functions	369
Q.6.4	B-spline MRA coefficient sequences	370
R	Source Code	375
Back Matter		379
References		380
Reference Index		405
Subject Index		409
License		424
End of document		427

Part I

Statistical Analysis



CHAPTER 1

EXPECTATION OPERATOR

1.1 Definitions

In a *probability space* (Ω, \mathbb{E}, P) (Definition A.2 page 147), all probability information is contained in the *measure* P . Often times this information is overwhelming and a simpler statistic, which does not offer so much information, is sufficient. Some of the most common statistics can be conveniently expressed in terms of the *expectation operator* E .

Definition 1.1. Let (Ω, \mathbb{E}, P) be a PROBABILITY SPACE (Definition A.2 page 147) and X a RANDOM VARIABLE (Definition ?? page ??) on (Ω, \mathbb{E}, P) with PROBABILITY DENSITY FUNCTION p_x .

D E F The **expectation operator** E_x on X is defined as

$$E_x X \triangleq \int_{x \in \mathbb{F}} x p_x(x) dx.$$

We already said that a *random variable* X is neither random nor a variable, but is rather a function of an underlying process that does appear to be random. However, because it is a function of a process that does appear random, the *random variable* X also appears to be random. That is, if we don't know the outcome of the underlying experimental process, then we also don't know for sure what X is, and so X does indeed appear to be random. However, even though X appears to be random, the expected value $E_x X$ of X is **not random**. Rather it is a fixed value (like 0 or 7.9 or -2.6).

On the other hand, even though EX is **not random**, note that $E(X|Y)$ is **random**. This is because $E(X|Y)$ is a function of Y . That is, once we know that Y equals some fixed value y (like 0 or 2.7 or -5.1) then $E(X|Y = y)$ is also fixed. However, if we don't know the value of Y , then Y is still a *random variable* and the expression $E(X|Y)$ is also random (a function of *random variable* Y).

Two common statistics that are conveniently expressed in terms of the expectation operator are the *mean* and *variance*. The mean is an indicator of the “middle” of a probability distribution and the variance is an indicator of the “spread”.

Definition 1.2. Let X be a RANDOM VARIABLE on the PROBABILITY SPACE (Ω, \mathbb{E}, P) .

- D E F**
- (1). The **mean** μ_X of X is $\mu_X \triangleq E_x X$
 - (2). The **variance** $\text{Var}(X)$ or σ_X^2 of X is $\text{Var}(X) \triangleq E_x [(X - E_x X)^2]$

1.2 Expectation as a linear operator

The next theorem demonstrates that the operator E is a *linear operator* (Definition M.3 page 282)—which in turn makes E part of a distinguished club of operators along with fellow member operators differentiation $\frac{d}{dx}$, integration $\int dx$, Laplace L , Fourier \tilde{F} , z-transform Z , etc. Because E is a linear operator, it immediately inherits all the properties that its linear operator birthright grants it (Corollary 1.1 page 4).

Theorem 1.1 (Linearity of E). ¹ Let X be a RANDOM VARIABLE on a PROBABILITY SPACE (Ω, \mathbb{E}, P) .

T	$E_x(aX + bY + c) = (aE_x X) + (bE_y Y) + c \quad \forall a, b, c \in \mathbb{R} \quad (\text{LINEAR})$
---	---

PROOF:

$$\begin{aligned}
 E_{xy}(aX + bY + c) &\triangleq \int_{x \in \mathbb{R}} \int_{y \in \mathbb{R}} [ax + by + c] p_{xy}(x, y) dy dx \quad \text{by definition of } E \text{ (Definition 1.1 page 3)} \\
 &= \int_{x \in \mathbb{R}} \int_{y \in \mathbb{R}} ax p_{xy}(x, y) dy dx + \int_{x \in \mathbb{R}} \int_{y \in \mathbb{R}} by p_{xy}(x, y) dy dx + \int_{x \in \mathbb{R}} \int_{y \in \mathbb{R}} c p_{xy}(x, y) dy dx \\
 &= \int_{x \in \mathbb{R}} ax \underbrace{\int_{y \in \mathbb{R}} p_{xy}(x, y) dy}_{p_x(x)} dx + \int_{y \in \mathbb{R}} by \underbrace{\int_{x \in \mathbb{R}} p_{xy}(x, y) dx}_{p_y(y)} dy + c \underbrace{\int_{y \in \mathbb{R}} \int_{x \in \mathbb{R}} p_{xy}(x, y) dx dy}_1 \\
 &= a \underbrace{\int_{x \in \mathbb{R}} x p_x(x) dx}_{EX} + b \underbrace{\int_{y \in \mathbb{R}} y p_y(y) dy}_{EY} + c \\
 &= (aE_x X) + (bE_y Y) + c
 \end{aligned}$$

⇒

Corollary 1.1. Let E be the EXPECTATION OPERATOR over a PROBABILITY SPACE (Ω, \mathbb{E}, P) . Let $sPLL$ be a VECTOR SPACE OF RANDOM VARIABLES over (Ω, \mathbb{E}, P) .

C	O	R	(1). $E\emptyset = \emptyset$ and (2). $E(-X) = -(EX) \quad \forall X \in L^2_{\mathbb{F}}$ and (3). $E(X - Y) = EX - EY \quad \forall X, Y \in L^2_{\mathbb{F}}$ and	(4). $E\left(\sum_{n=1}^N \alpha_n X_n\right) = \sum_{n=1}^N \alpha_n (EX_n) \quad \forall \alpha_n \in \mathbb{F}, \quad \forall X \in L^2_{\mathbb{F}}$
---	---	---	---	---

PROOF: These all follow immediately from the fact that E is a *linear operator* and from Theorem M.1 (page 282). ⇒

Remark 1.1. Projecting a stochastic process onto a basis often yields valuable insights into the nature of the underlying data. Typical projection operators include the Fourier operator \tilde{F} , Laplace L , and z-transform Z ...not to mention wavelet operators. But note that any such projection on a random sequence simply produces another random sequence. For example, the Fourier transform $\tilde{F}x(n)$ of a random sequence $x(n)$ is another random sequence.

One way to overcome this difficulty is to simply invoke the *sampling* operator $Sx(n)$ (CHAPTER ?? page ??), yielding a deterministic sequence, and then take the Fourier transform of the resulting deterministic sequence. The problem here is that every time you resample the sequence, you will very likely get a different Fourier transform.

¹ Wilks (1963), page 73 §3.2 “Mean value of a random variable”

Arguably a better approach (and the standard one at that) is to first invoke the expectation operator $E(x)$, also yielding a deterministic sequence.

The good news here is that because E and all the above mentioned operators are *linear*, we can do all the standard arithmetic acrobatics associated with linear algebra operators (next corollary).

Corollary 1.2. Let M and N be LINEAR OPERATORS (Definition M.3 page 282).

C O R	1. $E(MN) = (EM)N \quad \forall E \in \mathcal{L}(Z, W), M \in \mathcal{L}(Y, Z), N \in \mathcal{L}(X, Y) \quad (\text{ASSOCIATIVE})$
	2. $E(M + N) = (EM) + (EN) \quad \forall E \in \mathcal{L}(Y, Z), M \in \mathcal{L}(X, Y), N \in \mathcal{L}(X, Y) \quad (\text{LEFT DISTRIBUTIVE})$
	3. $(E + M)N = (EN) + (MN) \quad \forall E \in \mathcal{L}(Y, Z), M \in \mathcal{L}(Y, Z), N \in \mathcal{L}(X, Y) \quad (\text{RIGHT DISTRIBUTIVE})$
	4. $\alpha(EM) = (\alpha E)M = E(\alpha M) \quad \forall E \in \mathcal{L}(Y, Z), M \in \mathcal{L}(X, Y), \alpha \in \mathbb{F} \quad (\text{HOMOGENEOUS})$

PROOF: These all follow immediately from the fact that E is a *linear operator* and from Theorem M.4 (page 285). \Rightarrow

Corollary 1.3. Let X be a RANDOM VARIABLE on a PROBABILITY SPACE (Ω, \mathbb{E}, P) .

C O R	$\text{Var}(aX + b) = a^2 \text{Var}(X) \quad \forall a, b \in \mathbb{R}$
	$\text{Var}(X) = E_x(X^2) - (E_x X)^2$

PROOF:

$$\begin{aligned} \text{Var}(X) &\triangleq E_x[(X - E_x X)^2] && \text{by definition of } \text{Var} && (\text{Definition 1.2 page 3}) \\ &= E_x[X^2 - 2XE_x X + (E_x X)^2] && \text{by Binomial Theorem} && (\text{Theorem ?? page ??}) \\ &= E_x X^2 - E_x[2XE_x X] + E_x(E_x X)^2 && \text{by linearity of } E && (\text{Theorem 1.1 page 4}) \\ &= E_x X^2 - 2(E_x X)[E_x X] + (E_x X)^2 \\ &= E_x(X^2) - (E_x X)^2 \end{aligned}$$

$$\begin{aligned} \text{Var}(aX + b) &= E_x(aX + b)^2 - [E_x(aX + b)]^2 \\ &= E_x(a^2 X^2 + 2abX + b^2) - [a(E_x X) + b]^2 \\ &= a^2 E_x X^2 + 2abE_x X + b^2 - [a^2 [E_x X]^2 + 2abE_x X + b^2] && \text{by linearity of } E && (\text{Theorem 1.1 page 4}) \\ &= a^2 [E_x X^2 - (E_x X)^2] \\ &\triangleq a^2 \text{Var}(X) && \text{by previous result} \end{aligned}$$




Figure 1.1: Jensen's inequality

Jensen's inequality is an extremely useful application of *convexity* (Definition K.9 page 262) to the *expectation* operator. Jensen's inequality is stated in Corollary 1.4 (next) and illustrated in Figure 1.1 (page 5).

Corollary 1.4 (Jensen's inequality). ² Let f be a function in $\mathbb{R}^{\mathbb{R}}$ and X be a RANDOM VARIABLE on (Ω, \mathbb{E}, P) .

C O R	$\{f \text{ is CONVEX}\} \implies \{f(\mathbb{E}X) \leq \mathbb{E}f(X)\}$
-------------	---

PROOF:

1. Proof 1: Let $mx + c$ be a “support line” under $f(x)$ (Figure 1.1 page 5) such that

$$\begin{aligned} mx + c &< f(x) \quad \text{for } x \neq \mathbb{E}X \\ mx + c &= f(x) \quad \text{for } x = \mathbb{E}X. \end{aligned}$$

Then

$$\begin{aligned} f(\mathbb{E}X) &= m[\mathbb{E}X] + c \\ &= \mathbb{E}[mX + c] \\ &\leq \mathbb{E}f(X) \end{aligned}$$

2. Proof 2 (alternate proof):

$$\begin{aligned} f(\mathbb{E}X) &\triangleq f\left(\sum_{x \in \mathbb{E}} xP(x)\right) \\ &\leq \sum_{x \in \mathbb{E}} f(x)P(x) \quad \text{by Jensen's inequality for convex sets} \quad (\text{Theorem K.1 page 262}) \end{aligned}$$

Example 1.1. ³ Some examples of *Jensen's Inequality* (Corollary ?? page ??) applied to the *expectation operator* are the following:

E X	$(\mathbb{E}X)^{-1} < \mathbb{E}(X^{-1})$ $E(\log X) < \log(\mathbb{E}X)$ $e^{-\mathbb{E}X} \leq \mathbb{E}[e^{-X}]$
--------	--

Theorem 1.2 (Law of the Unconscious Statistician). ⁴

T H M	$E[g(X)] = \int_{x \in \mathbb{R}} g(x)p_x(x) dx$
-------------	---

1.3 Expectation inequalities

Theorem 1.3 (Markov's inequality). ⁵ Let $X : \Omega \rightarrow [0, \infty)$ be a non-negative valued RANDOM VARIABLE and $a \in (0, \infty)$. Then

T H M	$P\{X \geq a\} \leq \frac{1}{a} \mathbb{E}X$
-------------	--

² Shao (2003) page 31 (“1.3 Distributions and Their Characteristics”), Cover and Thomas (1991), page 25, Jensen (1906), pages 179–180

³ Shao (2003) pages 31–32 (“Example 1.18”), Dekking et al. (2006) page 110 (“8.5 Solutions to the quick exercises”)

⁴ Suhov et al. (2005) page 145 ((2.69)), Allen (2018) page 490 (18.3.4 The Law of the Unconscious Statistician), Papoulis (1990) page 124 (Fundamental Theorem)

⁵ Ross (1998), page 395

PROOF:

$$\begin{aligned} I &\triangleq \begin{cases} 1 & \text{for } X \geq a \\ 0 & \text{for } X < a \end{cases} \\ aI &\leq X \\ I &\leq \frac{1}{a}X \\ EI &\leq E\left(\frac{1}{a}X\right) \end{aligned}$$

$$\begin{aligned} P\{X \geq a\} &= 1 \cdot P\{X \geq a\} + 0 \cdot P\{X < a\} \\ &= EI \\ &\leq E\left(\frac{1}{a}X\right) \\ &= \frac{1}{a}EX \end{aligned}$$



Theorem 1.4 (Chebyshev's inequality). ⁶ Let X be a RANDOM VARIABLE with mean μ and variance σ^2 .

T H M	$P\{ X - \mu \geq a\} \leq \frac{\sigma^2}{a^2}$
-------------	---

PROOF:

$$\begin{aligned} P\{|X - \mu| \geq a\} &= P\{(X - \mu)^2 \geq a^2\} \\ &\leq \frac{1}{a^2} E(X - \mu)^2 && \text{by Markov's inequality} && (\text{Theorem 1.3 page 6}) \\ &= \frac{\sigma^2}{a^2} \end{aligned}$$



Theorem 1.5 (Kolmogorov's inequality). ⁷ Let X be a RANDOM VARIABLE with mean μ and variance σ^2 .

T H M	$\left\{ \begin{array}{l} (A). \quad (\mathbf{x}_n) \text{ are INDEPENDENT and} \\ (B). \quad \text{Each } \mathbf{x}_n \text{ has ZERO-MEAN and} \\ (C). \quad \text{Each } \mathbf{x}_n \text{ has variance } \sigma^2 \end{array} \right\} \implies P\left[\left \sum_{n=1}^N \mathbf{x}_n\right < \lambda \sum_{n=1}^N \mathbf{x}_n^2\right] \geq 1 - \frac{1}{\lambda^2}$
-------------	---

1.4 Joint and conditional probability spaces

Sometimes the problem of finding the expected value of a *random variable* X can be simplified by “conditioning X on Y ”.

Theorem 1.6. Let X and Y be RANDOM VARIABLES. Then

T H M	$E_x X = E_y E_{x y}(X Y)$
-------------	----------------------------

⁶ Ross (1998), page 396

⁷ Wilks (1963), page 107 (§4.5 “Kolmogorov’s inequality”)

PROOF:

$$\begin{aligned}
 E_y E_{x|y}(X|Y) &\triangleq E_y \left[\int_{x \in \mathbb{R}} x p(X=x|Y) dx \right] \\
 &\triangleq \int_{y \in \mathbb{R}} \left[\int_{x \in \mathbb{R}} x p(x|Y=y) dx \right] p(y) dy \\
 &= \int_{y \in \mathbb{R}} \int_{x \in \mathbb{R}} x p(x|y) p(y) dx dy \\
 &= \int_{x \in \mathbb{R}} x \int_{y \in \mathbb{R}} p(x,y) dy dx \\
 &= \int_{x \in \mathbb{R}} x p(x) dx \\
 &\triangleq E_x X
 \end{aligned}$$



1.5 Expectation inner product space

When possible, we like to generalize any given mathematical structure to a more general mathematical structure and then take advantage of the properties of that more general structure. Such a generalization can be done with *random variables*. Random variables can be viewed as vectors in a vector space. Furthermore, the expectation of the product of two *random variables* (e.g. $E(XY)$) can be viewed as an *inner product* in an *inner product space*. Since we have an *inner product space*, we can then immediately use all the properties of *inner product spaces*, *normed spaces*, *vector spaces*, *metric spaces*, and *topological spaces*.

Theorem 1.7.⁸ Let R be a ring, (Ω, \mathbb{E}, P) be a PROBABILITY SPACE, E the expectation operator, and $\mathbf{V} = \{X|X : \Omega \rightarrow R\}$ be the set of all random vectors in PROBABILITY SPACE (Ω, \mathbb{E}, P) .

- | | |
|-------------|--|
| T
H
M | (1). $\mathbf{V} \triangleq \{X X : \Omega \rightarrow R\}$ is a VECTOR SPACE.
(2). $\langle X Y \rangle \triangleq E(XY^*)$ is an INNER PRODUCT.
(3). $\ X\ \triangleq \sqrt{E(XX^*)}$ is a NORM.
(4). $(\mathbf{V}, \langle \Delta \nabla \rangle)$ is an INNER PRODUCT SPACE. |
|-------------|--|

PROOF:

1. Proof that \mathbf{V} is a vector space:

- | | | |
|--|---|-------------------------------|
| 1) $\forall X, Y, Z \in \mathbf{V}$ | $(X + Y) + Z = X + (Y + Z)$ | (+ is associative) |
| 2) $\forall X, Y \in \mathbf{V}$ | $X + Y = Y + X$ | (+ is commutative) |
| 3) $\exists 0 \in \mathbf{V}$ such that $\forall X \in \mathbf{V}$ | $X + 0 = X$ | (+ identity) |
| 4) $\forall X \in \mathbf{V} \exists Y \in \mathbf{V}$ such that | $X + Y = 0$ | (+ inverse) |
| 5) $\forall \alpha \in S$ and $X, Y \in \mathbf{V}$ | $\alpha \cdot (X + Y) = (\alpha \cdot X) + (\alpha \cdot Y)$ | (· distributes over +) |
| 6) $\forall \alpha, \beta \in S$ and $X \in \mathbf{V}$ | $(\alpha + \beta) \cdot X = (\alpha \cdot X) + (\beta \cdot X)$ | (· pseudo-distributes over +) |
| 7) $\forall \alpha, \beta \in S$ and $X \in \mathbf{V}$ | $\alpha(\beta \cdot X) = (\alpha \cdot \beta) \cdot X$ | (· associates with ·) |
| 8) $\forall X \in \mathbf{V}$ | $1 \cdot X = X$ | (· identity) |

⁸ Lindquist and Picci (2015) pages 25–26 (2.1 Hilbert Space of Second-Order Random Variables. $\langle \xi | \eta \rangle = E\{\xi\bar{\eta}\}$), Caines (1988) page 21 $\langle Exy = \int_{\Omega} x(\omega)y(\omega)dP(\omega) \rangle$, Caines (2018) page 21 $\langle Exy = \int_{\Omega} x(\omega)y(\omega)dP(\omega) \rangle$, Moon and Stirling (2000), pages 105–106

2. Proof that $\langle X | Y \rangle \triangleq E(XY^*)$ is an *inner product*.

- 1) $E(XX^*) \geq 0 \quad \forall X \in V$ (non-negative)
- 2) $E(XX^*) = 0 \iff X = 0 \quad \forall X \in V$ (non-degenerate)
- 3) $E(\alpha XY^*) = \alpha E(XY^*) \quad \forall X, Y \in V, \forall \alpha \in \mathbb{C}$ (homogeneous)
- 4) $E[(X + Y)Z^*] = E(XZ^*) + E(YZ^*) \quad \forall X, Y, Z \in V$ (additive)
- 5) $E(XY^*) = E(YX^*) \quad \forall X, Y \in V$ (conjugate symmetric).

3. Proof that $\|X\| \triangleq \sqrt{E(XX^*)}$ is a *norm*: This *norm* is simply induced by the above *inner product*.

4. Proof that $(V, \langle \Delta | \nabla \rangle)$ is an *inner product space*: Because V is a vector space and $\langle \Delta | \nabla \rangle$ is an *inner product*, $(V, \langle \Delta | \nabla \rangle)$ is an *inner product space*.



The next theorem gives some results that follow directly from vector space properties:

Theorem 1.8. Let (Ω, \mathbb{E}, P) be a PROBABILITY SPACE with expectation functional E .

T H M	<ol style="list-style-type: none"> 1. $\sqrt{E\left(\sum_{n=1}^N X_n\right)} \leq \sum_{n=1}^N E(X_n X_n^*)$ (GENERALIZED TRIANGLE INEQUALITY) 2. $E(XY^*) ^2 \leq E(XX^*) E(YY^*)$ (CAUCHY-SCHWARTZ INEQUALITY) 3. $2E(XX^*) + 2E(YY^*) = E[(X + Y)(X + Y)^*] + E[(X - Y)(X - Y)^*]$ (PARALLELOGRAM LAW)
-------------	--

PROOF:

1. $(\mathbb{R}^\Omega, E(x, y))$ is an *inner product space*. Proof: Theorem 1.7 (page 8).

2. Because it is an *inner product space*, the other properties follow:

1. Generalized triangle inequality: Theorem J.1 page 249
2. Cauchy-Schwartz inequality: Theorem I.2 page 234
3. Parallelogram Law: Theorem I.7 page 241



CHAPTER 2

RANDOM SEQUENCES



“A likely impossibility is always preferable to an unconvincing possibility.”¹
Aristotle (384 BC – 322 BC)



“We are quite in danger of sending highly trained and highly intelligent young men out into the world with tables of erroneous numbers under their arms, and with a dense fog in the place where their brains ought to be. In this century, of course, they will be working on guided missiles and advising the medical profession on the control of disease, and there is no limit to the extent to which they could impede every sort of national effort.”

Ronald A. Fisher, (1890–1962), Statistician, at a lecture in 1958 at Michigan State University²

2.1 Definitions

Definition 2.1.

D E F A **random sequence** is a **SEQUENCE**
over a **PROBABILITY SPACE** (Definition A.2 page 147).

Definition 2.2.³ Let $x(n)$ and $y(n)$ be RANDOM SEQUENCES.

¹ quote: <http://en.wikiquote.org/wiki/Aristotle>
image: <http://en.wikipedia.org/wiki/Aristotle>

² quote: [Yates and Mather \(1963\)](#) page 107. image: <http://www.genetics.org/content/154/4/1419>

³ [Papoulis \(1984\)](#) page 263 $\langle R_{xy}(m) = E\{x(m)y^*(0)\} \rangle$, [Wilks \(1963\)](#), page 77 “Moments of two-dimensional random variables”, [Crozat \(1987\)](#) page 341 $\langle r_{xy}(m) = E[x(m)y^*(0)] \rangle$, [MatLab \(2018b\)](#) $\langle R_{xy}(m) = E\{x_{n+m}y_n^*\} \rangle$, [MatLab \(2018a\)](#) $\langle R_{xy}(m) = E\{x_{n+m}y_n^*\} \rangle$

D E F	The mean	$\mu_X(n)$	<i>of $x(n)$ is</i>	$\mu_X(n) \triangleq E[x(n)]$
	The variance	$\sigma_X^2(n)$	<i>of $x(n)$ is</i>	$\sigma_X^2(n) \triangleq E([x(n) - \mu_X(n)]^2)$
	The cross-correlation	$R_{xy}(n, m)$	<i>of $x(n)$ and $y(n)$ is</i>	$R_{xy}(n, m) \triangleq E[x(n + m)y^*(n)]$
	The auto-correlation	$R_{xx}(n, m)$	<i>of $x(n)$ is</i>	$R_{xx}(n, m) \triangleq R_{xy}(n, m) _{y=x}$

2.2 Properties

Theorem 2.1.

T H M	$R_{xx}(n, m) = R_{xx}^*(n + m, -m)$
	$R_{xy}(n, m) = R_{yx}^*(n + m, -m)$

PROOF:

$$\begin{aligned}
 R_{xy}(n, m) &\triangleq E[x(n + m)y^*(n)] && \text{by definition of } R_{xy}(n, m) && (\text{Definition 2.2 page 11}) \\
 &= E[y^*(n)x(n + m)] && \text{by commutative property of } (\mathbb{C}, +, \cdot, 0, 1) \\
 &= (E[y(n)x^*(n + m)])^* && \text{by distributive property of } *-\text{algebras} && (\text{Definition H.3 page 228}) \\
 &= (E[y(n + m - m)x^*(n + m)])^* && \text{by additive identity property of } (\mathbb{R}, +, \cdot, 0, 1) \\
 &\triangleq R_{yx}^*(n + m, -m) && \text{by definition of } R_{xy}(n, m) && (\text{Definition 2.2 page 11})
 \end{aligned}$$

$$\begin{aligned}
 R_{xx}(n, m) &= R_{xy}(n, m)|_{y=x} && \text{by } y = x \text{ constraint} \\
 &= R_{xy}^*(n + m, -m)|_{y=x} && \text{by previous result} \\
 &= R_{xx}^*(n + m, -m) && \text{by } y = x \text{ constraint}
 \end{aligned}$$



2.3 Wide Sense Stationary processes

Definition 2.3. Let $x(n)$ be a RANDOM SEQUENCE with MEAN $\mu_X(n)$ and VARIANCE $\sigma_X^2(n)$ (Definition 2.2 page 11).

D E F	$x(n)$ is wide sense stationary (WSS) if
	1. $\mu_X(n)$ is CONSTANT with respect to n (STATIONARY IN THE 1ST MOMENT) and
	2. $\sigma_X^2(n)$ is CONSTANT with respect to n (STATIONARY IN THE 2ND MOMENT)

Definition 2.4. ⁴ Let $x(n)$ be a RANDOM SEQUENCE with statistics $\mu_X(n)$, $\sigma_X^2(n)$, $R_{xx}(n, m)$, and $R_{xy}(n, m)$ (Definition 2.2 page 11).

D E F	$\left\{ \begin{array}{l} x \text{ and } y \text{ are WIDE SENSE STATIONARY} \end{array} \right\} \implies$
	$\left\{ \begin{array}{lll} (1). \text{ The mean} & \mu_X & \text{of } x(n) \text{ is} \\ (2). \text{ The variance} & \sigma_X^2 & \text{of } x(n) \text{ is} \\ (4). \text{ The cross-correlation} & R_{xy}(m) & \text{of } x(n) \text{ and } y(n) \text{ is} \\ (3). \text{ The auto-correlation} & R_{xx}(m) & \text{of } x(n) \text{ is} \end{array} \right. \begin{array}{ll} \mu_X & \triangleq E[x(0)] \\ \sigma_X^2 & \triangleq E([x(0) - \mu_X]^2) \\ R_{xy}(m) & \triangleq E[x(m)y^*(0)] \\ R_{xx}(m) & \triangleq R_{xy}(m) _{y=x} \end{array} \right\}$

⁴ Papoulis (1984) page 263 $\langle R_{xy}(\tau) = E\{x(t + \tau)y^*(t)\} \rangle$, Cadzow (1987) page 341 $\langle r_{xy}(n) = E[x(k + n)y^*(k)] \rangle$ (10.41))

Remark 2.1. The $R_{xy}(n, m)$ of Definition 2.2 (page 11) and the $R_{xy}(m)$ of Definition 2.4 (page 12) (etc.) are examples of *function overload*—that is, functions that use the same mnemonic but are distinguished by different domains. Perhaps a more common example of function overload is the “+” mnemonic. Traditionally it is used with domain of the natural numbers \mathbb{N} as in $3 + 2$. Later it was extended for domain real numbers \mathbb{R} as in $\sqrt{3} + \sqrt{2}$, or even complex numbers \mathbb{C} as in $(\sqrt{3} + i\sqrt{2}) + (e + i\pi)$. And it was even more dramatically extended for use with domain $\mathbb{R}^N \times \mathbb{R}^M$ in “linear algebra” as in

$$\begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} + \begin{bmatrix} 5 & 6 \\ 7 & 8 \end{bmatrix} = \begin{bmatrix} 6 & 8 \\ 10 & 12 \end{bmatrix}$$

Proposition 2.1. Let $y(n)$ be a RANDOM SEQUENCE, $x(n)$ a RANDOM SEQUENCE with AUTO-CORRELATION $R_{xx}(n, m)$, and R_{xy} the CROSS-CORRELATION of x and y .

P R P	$\left\{ \begin{array}{l} x \text{ and } y \text{ are} \\ \text{WIDE SENSE STATIONARY} \\ (\text{WSS}) \text{ (Definition 6.1 page 43)} \end{array} \right\} \Rightarrow \left\{ \begin{array}{ll} R_{xx}(n, m) &= R_{xx}(m) & \forall n \in \mathbb{Z} \\ R_{xy}(n, m) &= R_{xy}(m) & \forall n \in \mathbb{Z} \\ \quad (\text{Definition 2.2 page 11}) & \quad (\text{Definition 2.4 page 12}) \end{array} \right\}$
----------------------	--

PROOF:

$$\begin{aligned} R_{xy}(n, m) &\triangleq E[x[n+m]y^*[n]] && \text{by definition of } R_{xy}(n, m) && \text{(Definition 2.2 page 11)} \\ &= E[x[n-n+m]y^*[n-n]] && \text{by wide sense stationary hypothesis} \\ &= E[x[m]y^*[0]] \\ &\triangleq R_{xy}(m) && \text{by definition of } R_{xy}(m) && \text{(Definition 2.4 page 12)} \\ R_{xx}(n, m) &= R_{xy}(n, m)|_{y=x} \\ &= R_{xy}(m)|_{y=x} && \text{by previous result} \\ &= R_{xx}(m) \end{aligned}$$




Figure 2.1: auto-correlation $R_{xx}(m)$

Corollary 2.1. Let $x(n)$ be a RANDOM SEQUENCE with AUTO-CORRELATION $R_{xx}(n, m)$, $y(n)$ a RANDOM SEQUENCE with AUTO-CORRELATION $R_{yy}(n, m)$, and $R_{xy}(n, m)$ the CROSS-CORRELATION of x and y . Let S be a SYSTEM with input $x(n)$ and output $y(n)$.

C O R	$\left\{ \begin{array}{l} (A). \quad x \text{ is WSS} \quad \text{and} \\ (B). \quad y \text{ is WSS} \quad \text{and} \\ (C). \quad S \text{ is LTI} \end{array} \right\} \Rightarrow \left\{ \begin{array}{ll} (1). \quad R_{xy}(m) &= R_{yx}^*(-m) & \text{and} \\ (2). \quad R_{xx}(m) &= R_{xx}^*(-m) & (\text{CONJUGATE SYMMETRIC}) \quad \text{and} \\ (3). \quad R_e R_{xx}(m) &= R_e R_{xx}(-m) & (\text{SYMMETRIC}) \quad \text{and} \\ (4). \quad I_m R_{xx}(m) &= -I_m R_{xx}(-m) & (\text{ANTI-SYMMETRIC}) \quad \text{and} \\ (5). \quad R_{xx}(m) &= R_{xx}(-m) & (\text{SYMMETRIC}) \quad \text{and} \\ (6). \quad \angle R_{xx}(m) &= -\angle R_{xx}(-m) & (\text{ANTI-SYMMETRIC}) \quad \text{and} \end{array} \right\}$
----------------------	--

PROOF:

$$\begin{aligned}
 R_{xy}(m) &= R_{xy}(n, m) && \text{by Proposition 2.1 page 13} && \text{and hypotheses (A),(B)} \\
 &= R_{yx}^*(n + m, -m) && \text{by Theorem 2.1 page 12} && \text{and hypothesis (B)} \\
 &= R_{yx}^*(-m) && \text{by Proposition 2.1 page 13} && \text{and hypothesis (A)} \\
 R_{xx}(m) &= R_{xx}(n, m) && \text{by Proposition 2.1 page 13} && \text{and hypothesis (A)} \\
 &= R_{xx}^*(n + m, -m) && \text{by Theorem 2.1 page 12} && \text{and hypothesis (B)} \\
 &= R_{xx}^*(-m) && \text{by Proposition 2.1 page 13} && \text{and hypothesis (A)}
 \end{aligned}$$



2.4 Spectral density

Definition 2.5. Let $x(n)$ and $y(n)$ be WIDE SENSE STATIONARY RANDOM SEQUENCES with auto-correlation $R_{xx}(m)$ and cross-correlation $R_{xy}(m)$. Let \mathbf{Z} be the Z-TRANSFORM OPERATOR (Definition P.4 page 330).

D E F The z-domain cross spectral density (CSD) $\check{S}_{xy}(z)$ of x and y is

$$\check{S}_{xy}(z) \triangleq \mathbf{Z}R_{xy}(m) \triangleq \sum_{m \in \mathbb{Z}} R_{xy}(m)z^{-m}$$

The z-domain power spectral density (PSD) $\check{S}_{xx}(z)$ of x is

$$\check{S}_{xx}(z) \triangleq \check{S}_{xy}(z)|_{y(n)=x(n)}$$

Definition 2.6. Let $x(n)$ and $y(n)$ be WIDE SENSE STATIONARY RANDOM SEQUENCES with auto-correlation $R_{xx}(m)$ and cross-correlation $R_{xy}(m)$. Let $\check{\mathbf{F}}$ be the DISCRETE TIME FOURIER TRANSFORM (DTFT) operator (Definition O.1 page 319).

D E F The auto-spectral density

$$\check{S}_{xx}(z) \text{ of } x \text{ is } \check{S}_{xx}(z) \triangleq \check{\mathbf{F}}R_{xx}(m) \triangleq \sum_{m \in \mathbb{Z}} R_{xx}(m)e^{-i\omega m}$$

The cross spectral density

$$(CSD) \check{S}_{xy}(z) \text{ of } x \text{ and } y \text{ is } \check{S}_{xy}(z) \triangleq \check{\mathbf{F}}R_{xy}(m) \triangleq \sum_{m \in \mathbb{Z}} R_{xy}(m)e^{-i\omega m}$$

The auto-spectral density is also called power spectral density (PSD).

Theorem 2.2. Let S be a system with IMPULSE RESPONSE $h(n)$, INPUT $x(n)$, and OUTPUT $y(n)$.

$$\begin{array}{l} \text{T H M} \\ \left\{ x \text{ and } y \text{ are WIDE SENSE STATIONARY} \right\} \implies \left\{ \begin{array}{l} (1). \check{S}_{xx}(z) = \check{S}_{xx}^*\left(\frac{1}{z^*}\right) \text{ and} \\ (2). \check{S}_{yx}(z) = \check{S}_{xy}^*\left(\frac{1}{z^*}\right) \end{array} \right\} \end{array}$$

PROOF:

$$\begin{aligned}
 \check{S}_{yx}(z) &\triangleq \mathbf{Z}R_{yx}(m) && \text{by definition of } \check{S}_{xy}(z) && (\text{Definition 2.6 page 14}) \\
 &\triangleq \sum_{m \in \mathbb{Z}} R_{yx}(m)z^{-m} && \text{by definition of } \mathbf{Z} && (\text{Definition P.4 page 330}) \\
 &\triangleq \sum_{m \in \mathbb{Z}} R_{xy}^*(-m)z^{-m} && \text{by Corollary 2.1 page 13} \\
 &= \left[\sum_{m \in \mathbb{Z}} R_{xy}(-m)(z^*)^{-m} \right]^* && \text{by antiautomorphic property of } *-\text{algebras} && (\text{Definition H.3 page 228})
 \end{aligned}$$



$$\begin{aligned}
 &= \left[\sum_{p \in \mathbb{Z}} R_{xy}(p)(z^*)^p \right]^* \quad \text{where } p \triangleq -m \quad \Rightarrow m = -p \\
 &= \left[\sum_{p \in \mathbb{Z}} R_{xy}(p)(z^*)^p \right]^* \quad \text{by } \textit{absolutely summable} \text{ property} \quad (\text{Definition P.2 page 329}) \\
 &= \left[\sum_{p \in \mathbb{Z}} R_{xy}(p) \left(\frac{1}{z^*} \right)^{-p} \right]^* \\
 &= \check{S}_{xy}^* \left(\frac{1}{z^*} \right) \quad \text{by definition of } \mathbf{Z} \quad (\text{Definition P.4 page 330}) \\
 \check{S}_{xx}(z) &= \check{S}_{xy}(z) \Big|_{y=x} \\
 &= \check{S}_{yx}(z) \Big|_{y=x} \\
 &= \check{S}_{xy}^* \left(\frac{1}{z^*} \right) \Big|_{y=x} \quad \text{by (2)—previous result} \\
 &= \check{S}_{xx}^* \left(\frac{1}{z^*} \right)
 \end{aligned}$$



Corollary 2.2. Let \mathbf{S} be a system with IMPULSE RESPONSE $h(n)$, INPUT $x(n)$, and OUTPUT $y(n)$.

C O R	$\left\{ \begin{array}{l} (A). \ h \text{ is LTI and} \\ (B). \ x \text{ and } y \text{ are WSS} \end{array} \right\} \Rightarrow \left\{ \begin{array}{l} (1). \ \tilde{S}_{xy}^*(\omega) = \tilde{S}_{yx}(\omega) \text{ (CONJUGATE-SYMMETRIC) and} \\ (2). \ \tilde{S}_{xx}^*(\omega) = \tilde{S}_{xx}(\omega) \text{ (CONJUGATE SYMMETRIC) and} \\ (3). \ \tilde{S}_{xx}(\omega) \in \mathbb{R} \text{ (REAL-VALUED)} \end{array} \right\}$
-------------	---

PROOF:

$$\begin{aligned}
 \tilde{S}_{xy}^*(\omega) &= \check{S}_{xy}^*(z) \Big|_{z=e^{i\omega}} \quad \text{by definition of DTFT} \quad (\text{Definition O.1 page 319}) \\
 &= \check{S}_{yx}^{**} \left(\frac{1}{z^*} \right) \Big|_{z=e^{i\omega}} \quad \text{by Theorem 2.2 page 14} \\
 &= \check{S}_{yx} \left(\frac{1}{z^*} \right) \Big|_{z=e^{i\omega}} \quad \text{by } \textit{involutory} \text{ property of } *-\text{algebras} \quad (\text{Definition H.3 page 228}) \\
 &= \check{S}_{yx} \left(\frac{1}{e^{i\omega*}} \right) \\
 &= \check{S}_{yx} (e^{i\omega}) \\
 &= \tilde{S}_{yx}(\omega) \quad \text{by definition of DTFT} \quad (\text{Definition O.1 page 319}) \\
 \tilde{S}_{xx}^*(\omega) &= \check{S}_{xx}^*(z) \Big|_{z=e^{i\omega}} \quad \text{by definition of DTFT} \quad (\text{Definition O.1 page 319}) \\
 &= \check{S}_{xx}^{**} \left(\frac{1}{z^*} \right) \Big|_{z=e^{i\omega}} \quad \text{by Theorem 2.2 page 14} \\
 &= \check{S}_{xx} \left(\frac{1}{z^*} \right) \Big|_{z=e^{i\omega}} \quad \text{by } \textit{involutory} \text{ property of } *-\text{algebras} \quad (\text{Definition H.3 page 228}) \\
 &= \check{S}_{xx} \left(\frac{1}{e^{i\omega*}} \right) \\
 &= \check{S}_{xx} (e^{i\omega}) \\
 &= \tilde{S}_{xx}(\omega) \quad \text{by definition of DTFT} \quad (\text{Definition O.1 page 319}) \\
 \implies \tilde{S}_{xx}(\omega) &\text{ is } \textit{real-valued} \\
 \tilde{S}_{xx}^*(\omega) &= \tilde{S}_{xy}^*(\omega) \Big|_{y=x} \\
 &= \tilde{S}_{yx}(\omega) \Big|_{y=x} \quad \text{by previous result} \\
 &= \tilde{S}_{xx}(\omega)
 \end{aligned}$$



2.5 Spectral Power

The term “*spectral power*” is a bit of an oxymoron because “spectral” deals with leaving the time-domain for the frequency-domain, howbeit the concept of power is solidly founded on the concept of time in that power = energy per time.

However, *Parseval's Theorem* (Proposition G.2 page 208) demonstrates that power in time can also be calculated in frequency. So, it makes some sense to speak of the term “spectral power”. Moreover, one way to estimate this power is to average the Fourier Transforms of the product $|x(n)|^2 = x(n)x^*(n)$...that is, to use an estimate of the auto-spectral density $\tilde{S}_{xx}(\omega)$. Thus, an alternate name for *auto-spectral density* is **power spectral density** (PSD).

CHAPTER 3

CONTINUOUS RANDOM PROCESSES



“*A likely impossibility is always preferable to an unconvincing possibility.*”¹
Aristotle (384 BC – 322 BC)

3.1 Definitions

Definition 3.1. ² Let $(\Omega, \mathbb{E}, \mathbb{P})$ be a PROBABILITY SPACE.

D E F The function $x : \Omega \rightarrow \mathbb{R}$ is a **random variable**.
The function $y : \mathbb{R} \times \Omega \rightarrow \mathbb{R}$ is a **random process**.

The random process $x(t, \omega)$, where t commonly represents time and $\omega \in \Omega$ is an outcome of an experiment, can take on more specialized forms depending on whether t and ω are fixed or allowed to vary. These forms are illustrated in Figure 3.1 page 17³ and Figure 3.2 page 18.

$x(t, \omega)$	fixed t	variable t
fixed ω	number	time function
variable ω	random variable	random process


Figure 3.1: Specialized forms of a random process $x(t, \omega)$

¹ quote: <http://en.wikiquote.org/wiki/Aristotle>

image: <http://en.wikipedia.org/wiki/Aristotle>

² Papoulis (1991), page 63, Papoulis (1991), page 285

³ Papoulis (1991), pages 285–286

Figure 3.2: Example of a random process $x(t, \omega)$

Definition 3.2. ⁴ Let $x(t)$ and $y(t)$ be random processes.

DEF

The mean	$\mu_X(t)$ of $x(t)$ is	$\mu_X(t) \triangleq E[x(t)]$
The cross-correlation	$R_{xy}(t)$ of $x(t)$ and $y(t)$ is	$R_{xy}(t, u) \triangleq E[x(t)y^*(u)]$
The autocorrelation function	$R_{xx}(t)$ of $x(t)$ is	$R_{xx}(t, u) \triangleq E[x(t)x^*(u)]$
The autocorrelation operator	R_f off(t) is	$R_f \triangleq \int_{u \in \mathbb{R}} R_{xx}(t, u)f(u) du$

Remark 3.1. ⁵ The equation $\int_{u \in \mathbb{R}} R_{xx}(t, u)f(u) du$ is a *Fredholm integral equation of the first kind* and $R_{xx}(t, u)$ is the *kernel* of the equation.

3.2 Properties

Theorem 3.1. Let $x(t)$ and $y(t)$ be random processes with cross-correlation $R_{xy}(t, u)$ and let $R_{xx}(t, u)$ be the auto-correlation of $x(t)$.

THM

$R_{xx}(t, u) = R_{xx}^*(u, t)$ (CONJUGATE SYMMETRIC)
$R_{xy}(t, u) = R_{yx}^*(u, t)$

PROOF:

$$\begin{aligned} R_{xx}(t, u) &\triangleq E[x(t)x^*(u)] &= E[x^*(u)x(t)] = (E[x(u)x^*(t)])^* &\triangleq R_{xx}^*(u, t) \\ R_{xy}(t, u) &\triangleq E[x(t)y^*(u)] &= E[y^*(u)x(t)] = (E[y(u)x^*(t)])^* &\triangleq R_{yx}^*(u, t) \end{aligned}$$

⇒

⁴ Papoulis (1984) page 216 $\langle R_{xy}(t_1, t_2) = E\{x(t_1)y^*(t_2)\} \rangle$ (9-35)),

⁵ Fredholm (1900), Fredholm (1903), page 365, Michel and Herget (1993), page 97, Keener (1988), page 101

Theorem 3.2. Let $\mathbf{R} : \mathbf{X} \rightarrow \mathbf{X}$ be an auto-correlation operator.

T H M	$\langle \mathbf{Rx} x \rangle \geq 0 \quad \forall x \in \mathbf{X} \quad (\text{NON-NEGATIVE})$
	$\langle \mathbf{Rx} y \rangle = \langle x \mathbf{Ry} \rangle \quad \forall x, y \in \mathbf{X} \quad (\text{SELF-ADJOINT})$

PROOF:

1. Proof that \mathbf{R} is non-negative:

$$\begin{aligned}
 \langle \mathbf{Ry} | y \rangle &= \left\langle \int_{u \in \mathbb{R}} R_{xx}(t, u)y(u) du | y(t) \right\rangle && \text{by definition of } \mathbf{R} && (\text{Definition 3.2 page 18}) \\
 &= \left\langle \int_{u \in \mathbb{R}} E[x(t)x^*(u)]y(u) du | y(t) \right\rangle && \text{by definition of } R_{xx}(t, u) && (\text{Definition 3.2 page 18}) \\
 &= E \left[\left\langle \int_{u \in \mathbb{R}} x(t)x^*(u)y(u) du | y(t) \right\rangle \right] && \text{by linearity of } \langle \Delta | \nabla \rangle \text{ and } \int && (\text{Definition I.1 page 233}) \\
 &= E \left[\int_{u \in \mathbb{R}} x^*(u)y(u) du \langle x(t) | y(t) \rangle \right] && \text{by additivity property of } \langle \Delta | \nabla \rangle && (\text{Definition I.1 page 233}) \\
 &= E[\langle y(u) | x(u) \rangle \langle x(t) | y(t) \rangle] && \text{by local definition of } \langle \Delta | \nabla \rangle && (\text{Definition I.1 page 233}) \\
 &= E[\langle x(u) | y(u) \rangle^* \langle x(t) | y(t) \rangle] && \text{by conjugate symmetry prop.} && (\text{Definition I.1 page 233}) \\
 &= E|\langle x(t) | y(t) \rangle|^2 && \text{by definition of } |\cdot| && (\text{Definition ?? page ??}) \\
 &\geq 0
 \end{aligned}$$

2. Proof that \mathbf{R} is self-adjoint:

$$\begin{aligned}
 \langle [\mathbf{Rx}](t) | y \rangle &= \left\langle \int_{u \in \mathbb{R}} R_{xx}(t, u)x(u) du | y(t) \right\rangle && \text{by definition of } \mathbf{R} && (\text{Definition 3.2 page 18}) \\
 &= \int_{u \in \mathbb{R}} x(u) \langle R_{xx}(t, u) | y(t) \rangle du && \text{by additive property of } \langle \Delta | \nabla \rangle && (\text{Definition I.1 page 233}) \\
 &= \int_{u \in \mathbb{R}} x(u) \langle y(t) | R_{xx}(t, u) \rangle^* du && \text{by conjugate symmetry prop.} && (\text{Definition I.1 page 233}) \\
 &= \langle x(u) | \langle y(t) | R_{xx}(t, u) \rangle \rangle && \text{by local definition of } \langle \Delta | \nabla \rangle && (\text{Definition I.1 page 233}) \\
 &= \left\langle x(u) | \int_{t \in \mathbb{R}} y(t) R_{xx}^*(t, u) dt \right\rangle && \text{by local definition of } \langle \Delta | \nabla \rangle && (\text{Definition I.1 page 233}) \\
 &= \left\langle x(u) | \int_{t \in \mathbb{R}} y(t) R_{xx}(u, t) dt \right\rangle && \text{by property of } R_{xx} && (\text{Theorem 3.1 page 18}) \\
 &= \left\langle x(u) | \underbrace{\mathbf{R}y}_{\mathbf{R}^*} \right\rangle && \text{by definition of } \mathbf{R} && (\text{Definition 3.2 page 18}) \\
 \implies \mathbf{R} &= \mathbf{R}^* \quad \implies \mathbf{R} \text{ is selfadjoint}
 \end{aligned}$$

Theorem 3.3.⁶ Let $(\lambda_n)_{n \in \mathbb{Z}}$ be the eigenvalues and $(\psi_n)_{n \in \mathbb{Z}}$ be the eigenfunctions of operator \mathbf{R} such that $\mathbf{R}\psi_n = \lambda_n\psi_n$.

T H M	1. $\lambda_n \in \mathbb{R}$	(eigenvalues of \mathbf{R} are REAL)
	2. $\lambda_n \neq \lambda_m \implies \langle \psi_n \psi_m \rangle = 0$	(eigenfunctions associated with distinct eigenvalues are ORTHOGONAL)
	3. $\ \psi_n(t)\ ^2 > 0 \implies \lambda_n \geq 0$	(eigenvalues are NON-NEGATIVE)
	4. $\ \psi_n(t)\ ^2 > 0, \langle \mathbf{R}\psi_n \psi_n \rangle > 0 \implies \lambda_n > 0$	(if \mathbf{R} is POSITIVE DEFINITE, then eigenvalues are POSITIVE)

⁶ Keener (1988), pages 114–119

PROOF:

1. Proof that eigenvalues are *real-valued*: Because \mathbf{R} is self-adjoint, its eigenvalues are real (Theorem M.17 page 298).
2. eigenfunctions associated with distinct eigenvalues are orthogonal: Because \mathbf{R} is self-adjoint, this property follows (Theorem M.17 page 298).
3. Proof that eigenvalues are *non-negative*:

$$\begin{aligned}
 0 &\geq \langle \mathbf{R}\psi_n | \psi_n \rangle && \text{by definition of non-negative definite} \\
 &= \langle \lambda_n \psi_n | \psi_n \rangle && \text{by hypothesis} \\
 &= \lambda_n \langle \psi_n | \psi_n \rangle && \text{by definition of inner-products} \\
 &= \lambda_n \|\psi_n\|^2 && \text{by definition of norm induced by inner-product}
 \end{aligned}$$

4. Eigenvalues are *positive* if \mathbf{R} is *positive definite*:

$$\begin{aligned}
 0 &> \langle \mathbf{R}\psi_n | \psi_n \rangle && \text{by definition of } \textit{positive definite} \\
 &= \langle \lambda_n \psi_n | \psi_n \rangle && \text{by hypothesis} \\
 &= \lambda_n \langle \psi_n | \psi_n \rangle && \text{by definition of inner-products} \\
 &= \lambda_n \|\psi_n\|^2 && \text{by definition of norm induced by inner-product}
 \end{aligned}$$



3.3 Basis for random processes

If a random process $x(t)$ is white ⁷ and $\Psi = \{\psi_1(t), \psi_2(t), \dots, \psi_N(t)\}$ is **any** set of orthonormal basis functions, then the innerproducts $\langle n(t) | \psi_n(t) \rangle$ and $\langle n(t) | \psi_m(t) \rangle$ are *uncorrelated* for $m \neq n$. However, if $x(t)$ is colored (not white), then the innerproducts are not in general uncorrelated. But if the elements of Ψ are chosen to be the eigenfunctions of \mathbf{R} such that

$$\mathbf{R}\psi_n = \lambda_n \psi_n,$$

then by Theorem 3.1 page 18, $\{\psi_n(t)\}$ are orthogonal and the innerproducts **are** uncorrelated even though $x(t)$ is not white. This criterion is called the Karhunen-Loève criterion for $x(t)$.

Theorem 3.4 (Karhunen-Loève Expansion). ⁸ Let \mathbf{R} be the AUTOCORRELATION OPERATOR of a RANDOM PROCESS $x(t)$.

THM	$ \left\{ \begin{array}{l} \text{(A). } (\lambda_n)_{n \in \mathbb{Z}} \text{ are the eigenvalues of } \mathbf{R} \quad \text{and} \\ \text{(B). } (\psi_n)_{n \in \mathbb{Z}} \text{ are the eigenfunctions of } \mathbf{R} \quad \text{such that} \\ \text{(C). } \mathbf{R}\psi_n = \lambda_n \psi_n \end{array} \right\} $ $ \Rightarrow \left\{ \begin{array}{l} \left\{ \mathbf{E} \left\{ \left x(t) - \sum_{n \in \mathbb{Z}} \langle x(t) \psi_n(t) \rangle \psi_n(t) \right ^2 \right\} = 0 \right\} \\ \left(\{\psi_n(t)\} \text{ is a BASIS for } x(t) \right) \end{array} \right\} $
-----	---

⁷ white noise process: random process $x(t)$ with autocorrelation $R_{xx}(\tau) = \delta(\tau)$

⁸ Keener (1988), pages 114–119

PROOF:

1. $\{\psi_n(t)\}$ is a basis for $x(t)$

$$\mathbb{E}\left\{\left|x(t) - \sum_{n \in \mathbb{Z}} \dot{x}_n \psi_n(t)\right|^2\right\} = 0 \quad \text{where } \dot{x}_n \triangleq \langle x(t) | \psi_n(t) \rangle$$

$$\begin{aligned} \mathbb{E}\left[x(t)\left(\sum_{n \in \mathbb{Z}} \dot{x}_n \psi_n(t)\right)^*\right] &= \mathbb{E}\left[x(t)\left(\sum_{n \in \mathbb{Z}} \int_{u \in \mathbb{R}} x(u) \psi_n^*(u) du \psi_n(t)\right)^*\right] \\ &= \sum_{n \in \mathbb{Z}} \left(\int_{u \in \mathbb{R}} \mathbb{E}[x(t)x^*(u)] \psi_n(u) du \right) \psi_n^*(t) \\ &= \sum_{n \in \mathbb{Z}} \left(\int_{u \in \mathbb{R}} R_{xx}(t, u) \psi_n(u) du \right) \psi_n^*(t) \\ &= \sum_{n \in \mathbb{Z}} \lambda_n \psi_n(t) \psi_n^*(t) \\ &= \sum_{n \in \mathbb{Z}} \lambda_n |\psi_n(t)|^2 \end{aligned}$$

$$\begin{aligned} \mathbb{E}\left[\sum_{n \in \mathbb{Z}} \dot{x}_n \psi_n(t) \left(\sum_{m \in \mathbb{Z}} \dot{x}_m \psi_m(t)\right)^*\right] &= \mathbb{E}\left[\sum_{n \in \mathbb{Z}} \int_{u \in \mathbb{R}} x(u) \psi_n^*(u) du \psi_n(t) \left(\sum_{m \in \mathbb{Z}} \int_v x(v) \psi_m^*(v) dv \psi_m(t)\right)^*\right] \\ &= \sum_{n \in \mathbb{Z}} \sum_{m \in \mathbb{Z}} \int_u \left(\int_v \mathbb{E}[x(u)x^*(v)] \psi_m(v) dv \right) \psi_n^*(u) du \psi_n(t) \psi_m^*(t) \\ &= \sum_{n \in \mathbb{Z}} \sum_{m \in \mathbb{Z}} \int_u \left(\int_v R_{xx}(u, v) \psi_m(v) dv \right) \psi_n^*(u) du \psi_n(t) \psi_m^*(t) \\ &= \sum_{n \in \mathbb{Z}} \sum_{m \in \mathbb{Z}} \int_u (\lambda_m \psi_m(u)) \psi_n^*(u) du \psi_n(t) \psi_m^*(t) \\ &= \sum_{n \in \mathbb{Z}} \sum_{m \in \mathbb{Z}} \lambda_m \left(\int_u \psi_m(u) \psi_n^*(u) du \right) \psi_n(t) \psi_m^*(t) \\ &= \sum_{n \in \mathbb{Z}} \sum_{m \in \mathbb{Z}} \lambda_m \bar{\delta}_{mn} \psi_n(t) \psi_m^*(t) \\ &= \sum_{n \in \mathbb{Z}} \lambda_n \psi_n(t) \psi_n^*(t) \\ &= \sum_{n \in \mathbb{Z}} \lambda_n |\psi_n(t)|^2 \end{aligned}$$

2. Using the previous two results, we can prove the following:

$$\begin{aligned} \mathbb{E}\left\{\left|x(t) - \sum_{n \in \mathbb{Z}} \dot{x}_n \psi_n(t)\right|^2\right\} \\ &= \mathbb{E}\left[\left[x(t) - \sum_{n \in \mathbb{Z}} \dot{x}_n \psi_n(t)\right] \left[x(t) - \sum_{m \in \mathbb{Z}} \dot{x}_m \psi_m(t)\right]^*\right] \\ &= \mathbb{E}\left[x(t)x^*(t) - x(t)\left(\sum_{n \in \mathbb{Z}} \dot{x}_n \psi_n(t)\right)^* - x^*(t) \sum_{n \in \mathbb{Z}} \dot{x}_n \psi_n(t) + \sum_{n \in \mathbb{Z}} \dot{x}_n \psi_n(t) \left(\sum_{m \in \mathbb{Z}} \dot{x}_m \psi_m(t)\right)^*\right] \end{aligned}$$

$$= \mathbb{E}[x(t)x^*(t)] - \mathbb{E}\left[x(t)\left(\sum_{n \in \mathbb{Z}} \dot{x}_n \psi_n(t)\right)^*\right] - \mathbb{E}\left[x^*(t) \sum_{n \in \mathbb{Z}} \dot{x}_n \psi_n(t)\right] + \mathbb{E}\left[\sum_{n \in \mathbb{Z}} \dot{x}_n \psi_n(t) \left(\sum_{m \in \mathbb{Z}} \dot{x}_m \psi_m(t)\right)^*\right]$$


by *Mercer's Theorem*(Theorem D.5 page 172)

$$\begin{aligned} &= \sum_{n \in \mathbb{Z}} \lambda_n |\psi_n(t)|^2 - \sum_{n \in \mathbb{Z}} \lambda_n |\psi_n(t)|^2 - \left(\sum_{n \in \mathbb{Z}} \lambda_n |\psi_n(t)|^2 \right)^* + \sum_{n \in \mathbb{Z}} \lambda_n |\psi_n(t)|^2 \\ &= \sum_{n \in \mathbb{Z}} \lambda_n |\psi_n(t)|^2 - \sum_{n \in \mathbb{Z}} \lambda_n |\psi_n(t)|^2 - \sum_{n \in \mathbb{Z}} \lambda_n |\psi_n(t)|^2 + \sum_{n \in \mathbb{Z}} \lambda_n |\psi_n(t)|^2 \\ &= 0 \end{aligned}$$



Part II

Statistical Processing



CHAPTER 4

OPERATIONS ON RANDOM VARIABLES

4.1 Functions of one random variable

Proposition 4.1. Let $(\Omega, \mathbb{E}, \mathbb{P})$ be a probability space, and X a RANDOM VARIABLE with CUMULATIVE DISTRIBUTION FUNCTION $c_X(x)$.

P R P	$\left\{ \begin{array}{l} X \text{ is UNIFORMLY DISTRIBUTED} \\ (\text{Definition C.1 page 161}) \end{array} \right\} \iff c_X(x) = \begin{cases} 0 & \text{for } x < 0 \\ x & \text{for } 0 < x \leq 1 \\ 1 & \text{otherwise} \end{cases}$
----------------------	--

Theorem 4.1 (Probability integral transform). ¹ Let $(\Omega, \mathbb{E}, \mathbb{P})$ be a probability space. Let X be a RANDOM VARIABLE with PROBABILITY DENSITY FUNCTION $p_X(x)$ and CUMULATIVE DISTRIBUTION FUNCTION $c_X(x)$. Let Y be a RANDOM VARIABLE CUMULATIVE DISTRIBUTION FUNCTION $c_Y(y)$.

T H M	$\left\{ \begin{array}{l} (1). Y = c_X(X) \\ (2). p_X(x) \text{ is CONTINUOUS} \end{array} \text{ and } \right\} \implies \left\{ \begin{array}{l} Y \text{ is UNIFORMLY DISTRIBUTED} \\ (\text{Definition C.1 page 161}) \end{array} \right\}$
----------------------	---

PROOF:


$$\begin{aligned}
 c_Y(y) &\triangleq \mathbb{P}\{Y \leq y\} && \text{by definition of cdf} && (\text{Definition B.2 page 156}) \\
 &= \mathbb{P}\{c_X(X) \leq y\} && \text{by hypothesis (1)} \\
 &= \mathbb{P}\{X \leq c_X^{-1}(y)\} && \text{by hypothesis (2) and} && \text{Proposition A.2 page 150} \\
 &\triangleq c_X[c_X^{-1}(y)] && \text{by definition of cdf} && (\text{Definition B.2 page 156}) \\
 &= y \\
 \implies Y &\text{ is uniformly distributed} && \text{by} && \text{Proposition 4.1 page 25}
 \end{aligned}$$

Theorem 4.2 (Inverse probability integral transform). ² Let $(\Omega, \mathbb{E}, \mathbb{P})$ be a probability space. Let X be a RANDOM VARIABLE with PROBABILITY DENSITY FUNCTION $p_X(x)$ and CUMULATIVE DISTRIBUTION FUNCTION $c_X(x)$. Let Y be a RANDOM VARIABLE CUMULATIVE DISTRIBUTION FUNCTION $c_Y(y)$.

T H M	$\left\{ \begin{array}{l} (1). Y = c_z^{-1}(X) \\ (2). Y \text{ is UNIFORMLY DISTRIBUTED} \\ (3). p_z(z) \text{ is CONTINUOUS} \end{array} \text{ and } \right\} \implies \left\{ \begin{array}{l} p_Y(y) = p_z(y) \\ (Y \text{ has distribution } p_z(y)) \end{array} \right\}$
----------------------	--

¹ Angus (1994), Roussas (2014) page 232 (Theorem 10), Devroye (1986) page 28 (Theorem 2.1)

² Devroye (1986) page 28 (Theorem 2.1), Balakrishnan and Lai (2009) page 624 (14.2.1 Introduction)

Figure 4.1: $Y = f(X)$

PROOF:

$$\begin{aligned}
 c_Y(y) &\triangleq P\{Y \leq y\} && \text{by definition of } c_Y && (\text{Definition B.2 page 156}) \\
 &= P\{c_Z^{-1}(X) \leq y\} && \text{by hypothesis (1)} \\
 &= P\{X \leq c_Z(y)\} && \text{by hypothesis (3) and} \\
 &\triangleq c_X[c_Z(y)] && \text{by definition of } c_X \\
 &= c_Z(y) && \text{because } 0 \leq c_Z(y) \leq 1 \text{ and by} \\
 \implies p_Y(y) &= p_Z(y) && (\text{Y has the distribution of Z}) && \text{Proposition A.2 page 150} \\
 &&&&& (\text{Definition B.2 page 156}) \\
 &&&&& \text{Proposition 4.1 page 25}
 \end{aligned}$$

Definition 4.1.³ Let $f(x)$ be a DIFFERENTIABLE FUNCTION in $\mathbb{R}^{\mathbb{R}}$.

D E F A point $p \in \mathbb{R}$ is a **critical point** of $f(x)$ if
 $f'(p) = 0$.

Theorem 4.3.⁴ Let X and Y be RANDOM VARIABLES in $\mathbb{R}^{\mathbb{R}}$. Let f be a DIFFERENTIABLE FUNCTION in $\mathbb{R}^{\mathbb{R}}$ with N CRITICAL POINTS (Definition 4.1 page 26). Let the range of X be partitioned into $N + 1$ partitions $\{A_n | n = 1, 2, \dots, N + 1\}$ with partition boundaries set at the N CRITICAL POINTS of $f(x)$ —as illustrated in Figure 4.1 (page 26). Let $g_n(x) \triangleq f(x)$ but with domain restricted to $x \in A_n$.

T H M	$\left\{ \begin{array}{l} (1). \quad Y = f(X) \\ (2). \quad f \text{ is DIFFERENTIABLE} \end{array} \right. \text{ and } \right\} \implies \left\{ p_Y(y) = \sum_{n=1}^{N+1} \frac{p_X(g_n^{-1}(y))}{ f'(g_n^{-1}(y)) } \right\}$
--------------	---


PROOF:

1. The problem with a function $f(x)$ with at least $N = 1$ critical point is that $f^{-1}(y)$ is *not invertible*. That is, $f^{-1}(y)$ has more than one solution (and thus the *relation* $f^{-1}(y)$ is not a *function*). However, note that in each partition A_n , $f(x)$ is *invertible* and thus $f^{-1}(y)$ in that partition has a *unique* solution. Thus, each $g_n(x)$ is *invertible* in its domain (and each $g_n^{-1}(y)$ exists as a function).

³ Callahan (2010) page 189 (Definition 6.1)⁴ Papoulis (1984) pages 95–96 (“Fundamental Theorem”), Papoulis (1990) page 157 (“Fundamental Theorem”) Papoulis (1991), page 93, Proakis (2001), page 30

2. Using item (1), the remainder of the proof follows ...

$$\begin{aligned}
 p_Y(y) &\triangleq \frac{d}{dy} P\{Y \leq y\} && \text{by definition of } p_Y \text{ (Definition B.2 page 156)} \\
 &= \frac{d}{dy} P\{f(X) \leq y\} && \text{by hypothesis (1)} \\
 &= \frac{d}{dy} \sum_{n=1}^{N+1} P\{f(X) \leq y | X \in A_n\} && \text{by sum of products (Theorem A.3 page 149)} \\
 &= \frac{d}{dy} \sum_{n=1}^{N+1} P\{f(X) \leq y | X \in A_n\} P\{X \in A_n\} && \text{by definition of } P\{X|Y\} \text{ (Definition A.4 page 148)} \\
 &= \frac{d}{dy} \sum_{n=1}^{N+1} P\{g_n(X) \leq y | X \in A_n\} P\{X \in A_n\} && \text{by definition of } g_n(x) \\
 &= \frac{d}{dy} \left\{ \begin{array}{ll} \sum_{n=1}^{N+1} P\{X \leq g_n^{-1}(y) | X \in A_n\} P\{X \in A_n\} & \text{for } f'(x) \geq 0 \\ \sum_{n=1}^{N+1} P\{X \geq g_n^{-1}(y) | X \in A_n\} P\{X \in A_n\} & \text{otherwise} \end{array} \right\} && \text{by item (1)} \\
 &= \frac{d}{dy} \left\{ \begin{array}{ll} \sum_{n=1}^{N+1} P\{X \leq g_n^{-1}(y) | X \in A_n\} P\{X \in A_n\} & \text{for } f'(x) \geq 0 \\ \sum_{n=1}^{N+1} [1 - P\{X \leq g_n^{-1}(y) | X \in A_n\} P\{X \in A_n\}] & \text{otherwise} \end{array} \right\} \\
 &= \frac{d}{dy} \left\{ \begin{array}{ll} \sum_{n=1}^{N+1} P\{X \leq g_n^{-1}(y) | X \in A_n\} & \text{for } f'(x) \geq 0 \\ \sum_{n=1}^{N+1} [1 - P\{X \leq g_n^{-1}(y) | X \in A_n\}] & \text{otherwise} \end{array} \right\} && \text{by definition of } P\{X|Y\} \\
 &= \frac{d}{dy} \left\{ \begin{array}{ll} \sum_{n=1}^{N+1} (P\{X \leq g_n^{-1}(y)\} - P\{X < \min A_{n-1}\}) & \text{for } f'(x) \geq 0 \\ \sum_{n=1}^{N+1} [1 - (P\{X \leq g_n^{-1}(y)\} - P\{X < \min A_{n-1}\})] & \text{otherwise} \end{array} \right\} \\
 &= \frac{d}{dy} \left\{ \begin{array}{ll} \sum_{n=1}^{N+1} P\{X \leq g_n^{-1}(y)\} & \text{for } f'(x) \geq 0 \\ \sum_{n=1}^{N+1} [1 - P\{X \leq g_n^{-1}(y)\}] & \text{otherwise} \end{array} \right\} && \text{because } \frac{d}{dy} P\{X < \text{a constant}\} = 0 \\
 &= \left\{ \begin{array}{ll} \sum_{n=1}^{N+1} \frac{d}{dy} c_x[g_n^{-1}(y)] & \text{for } f'(x) \geq 0 \\ \sum_{n=1}^{N+1} \frac{d}{dy} [1 - c_x(g_n^{-1}(y))] & \text{otherwise} \end{array} \right\} && \text{by linearity of } \frac{d}{dy} \text{ operator} \\
 &= \left\{ \begin{array}{ll} \sum_{n=1}^{N+1} p_x[g_n^{-1}(y)] \frac{d}{dy}[g_n^{-1}(y)] & \text{for } f'(x) \geq 0 \\ \sum_{n=1}^{N+1} \left[-p_x[g_n^{-1}(y)] \frac{d}{dy}[g_n^{-1}(y)] \right] & \text{otherwise} \end{array} \right\} && \text{by definition of } p_x \text{ (Definition B.2 page 156) and the chain rule}
 \end{aligned}$$

Figure 4.2: $Y = \frac{1}{X}$

$$\begin{aligned}
 &= \sum_{n=1}^{N+1} p_x(g_n^{-1}(y)) \left| \frac{d}{dy} [g_n^{-1}(y)] \right| \\
 &= \sum_{n=1}^{N+1} \frac{p_x(g_n^{-1}(y))}{|f'(g_n^{-1}(y))|} \quad \text{by Lemma ?? page ??}
 \end{aligned}$$

Corollary 4.1. ⁵ Let X and Y be RANDOM VARIABLES in $\mathbb{R}^{\mathbb{R}}$. Let $a, b \in \mathbb{R}$.


C O R	$\left\{ \begin{array}{l} (1). \quad Y = aX + b \quad \text{and} \\ (2). \quad a \neq 0 \end{array} \right\} \implies \left\{ p_Y(y) = \frac{1}{ a } p_X\left(\frac{y-b}{a}\right) \right\}$
----------------------	--

PROOF:

1. Note that $f(x) = ax + b$ is a *differentiable function* with $N = 0$ *critical points* and $f'(x) = a$.
2. The inverse of $f(x)$ is $g_1(y) = f^{-1}(y) = \frac{y-b}{a}$.
3. It follows that

$$\begin{aligned}
 p_Y(y) &= \sum_{n=1}^{N+1} \frac{p_x(g_n^{-1}(y))}{|f'(g_n^{-1}(y))|} \quad \text{by Theorem 4.3 (page 26)} \\
 &= \frac{p_x(f^{-1}(y))}{|f'(f^{-1}(y))|} \quad \text{because } N = 0 \\
 &= \frac{p_x(f^{-1}(y))}{|a|} \quad \text{by item (1)} \\
 &= \frac{1}{|a|} p_X\left(\frac{y-b}{a}\right) \quad \text{by item (2)}
 \end{aligned}$$

⁵ Papoulis (1984) page 96 ("Illustrations" 1), Papoulis (1991), page 95, Proakis (2001), page 29

Figure 4.3: $Y = rVX^2$ **Corollary 4.2.**⁶

COR $\left\{ Y = \frac{1}{X} \right\} \Rightarrow \left\{ p_Y(y) = \frac{1}{y^2} p_X\left(\frac{1}{y}\right) \text{ for } y > 0 \right\}$

PROOF:

1. Note that $f(x) = 1/x$ is a *differentiable function* in $x > 0$ with $N = 0$ *critical points* and $f'(x) = -1/x^2$.
2. The inverse of $f(x)$ is $g_1(y) = f^{-1}(y) = \frac{1}{y}$.
3. It follows that

$$\begin{aligned} p_Y(y) &= \sum_{n=1}^{N+1} \frac{p_X(g_n^{-1}(y))}{|f'(g_n^{-1}(y))|} && \text{by Theorem 4.3 (page 26)} \\ &= \frac{p_X(f^{-1}(y))}{|f'(f^{-1}(y))|} && \text{because } N = 0 \\ &= \frac{1}{|-1/(1/y)^2|} p_X\left(\frac{1}{y}\right) \\ &= \frac{1}{y^2} p_X\left(\frac{1}{y}\right) \end{aligned}$$

**Corollary 4.3.**⁷ Let X and Y be RANDOM VARIABLES.


COR $\left\{ Y = X^2 \right\} \Rightarrow \left\{ p_Y(y) = \frac{1}{2\sqrt{y}} [p_X(-\sqrt{y}) + p_X(\sqrt{y})] \right\}$

PROOF:

1. The roots of $y = x^2$ are $x_1 = -\sqrt{y}$ and $x_2 = +\sqrt{y}$.
2. The derivative of $f(x) \triangleq y = x^2$ is $f'(x) = 2x$.

⁶ Papoulis (1984) page 97 (Example 5-10), Papoulis (1991), page 94

⁷ Papoulis (1984) page 95 (Example 5-9), Devroye (1986) page 27 (Example 4.4), Papoulis (1991), page 95, Proakis (2001), page 29

Figure 4.4: $Z = \tan \Theta$

3. And so it follows that ...

$$\begin{aligned}
 p_Y(y) &= \sum_{n=1}^N \frac{p_X(x_n)}{|f'(x_n)|} && \text{by Theorem 4.3 page 26} \\
 &= \frac{p_X(x_1)}{|f'(x_1)|} + \frac{p_X(x_2)}{|f'(x_2)|} && \text{by definition of } \sum \\
 &= \frac{p_X(-\sqrt{y})}{|f'(-\sqrt{y})|} + \frac{p_X(\sqrt{y})}{|f'(\sqrt{y})|} && \text{by item (1)} \\
 &= \frac{p_X(-\sqrt{y})}{2\sqrt{y}} + \frac{p_X(\sqrt{y})}{2\sqrt{y}} && \text{by item (2)} \\
 &= \frac{1}{2\sqrt{y}} \left[p_X(-\sqrt{y}) + p_X(\sqrt{y}) \right] && \text{by linearity of } + \text{ operation}
 \end{aligned}$$

Corollary 4.4.⁸ Let $Z = \tan \Theta$. Then

C O R	$\{Z = \tan \Theta\} \implies \left\{ p_Z(z) = \frac{1}{1+z^2} \sum_{n \in Z} p_\theta(\tan(z) + n\pi) \right\}$
-------------	--

PROOF:

1. The roots of $z = \tan \theta$ are $\{\theta_n = \arctan z + n\pi | n \in \mathbb{Z}\}$.
2. The derivative of $z = \tan \theta$ is $f'(\theta) = \sec^2 \theta$.
3. It follows that

$$\begin{aligned}
 p_Z(z) &= \sum_{n=1}^N \frac{p_\theta(\theta_n)}{|f'(\theta_n)|} \\
 &= \sum_n \frac{p_\theta(\arctan z + n\pi)}{|f'(\arctan z + n\pi)|} \\
 &= \sum_n \frac{p_\theta(\arctan z + n\pi)}{|\sec^2(\arctan z + n\pi)|} \\
 &= \sum_n \cos^2(\arctan z + n\pi) p_\theta(\arctan z + n\pi)
 \end{aligned}$$

⁸ Papoulis (1991), pages 99–100

$$\begin{aligned}
 &= \cos^2(\tan z) \sum_n p_\theta(\tan z + n\pi) \\
 &= \frac{1}{1+z^2} \sum_n p_\theta(\tan z + n\pi)
 \end{aligned}$$



4.2 Functions of two random variables

Theorem 4.4. ⁹ Let X , Y , and Z be RANDOM VARIABLES. Let \star be the CONVOLUTION operator (Definition N.3 page 312).

T H M	$\left\{ \begin{array}{l} (1). \quad Z \triangleq X + Y \\ (2). \quad X \text{ and } Y \text{ are INDEPENDENT} \end{array} \right. \quad \text{and} \quad \begin{array}{l} \text{(Definition A.3 page 147)} \\ \hline \end{array} \right\} \Rightarrow \{p_Z(z) = p_X(z) \star p_Y(z)\}$
----------------------	--

PROOF:

$$\begin{aligned}
 p_Z(z) &\triangleq \frac{d}{dz} c_Z(z) && \text{by definition of } p_Z && (\text{Definition B.2 page 156}) \\
 &\triangleq \frac{d}{dz} P\{Z \leq z\} && \text{by definition of } c_Z && (\text{Definition B.2 page 156}) \\
 &= \frac{d}{dz} P\{X + Y \leq z\} && \text{by hypothesis (1)} && \\
 &= \frac{d}{dz} \lim_{\varepsilon \rightarrow 0} \sum_{n \in \mathbb{Z}} P\{X + Y \leq z | y + n\varepsilon < Y \leq y + (n+1)\varepsilon\} && \text{by sum of products} && (\text{Theorem A.3 page 149}) \\
 &= \frac{d}{dz} \int_{y \in \mathbb{R}} P\{X + Y \leq z | Y = y\} p_Y(y) dy && \text{by definiton of } P\{X | Y\} && (\text{Definition A.4 page 148}) \\
 &= \frac{d}{dz} \int_{y \in \mathbb{R}} P\{X \leq z - y | Y = y\} p_Y(y) dy && \\
 &= \frac{d}{dz} \int_{y \in \mathbb{R}} P\{X \leq z - y\} p_Y(y) dy && \text{by hypothesis (2)} && \\
 &\triangleq \frac{d}{dz} \int_{y \in \mathbb{R}} c_X(z - y) p_Y(y) dy && \text{by definition of } c_X && (\text{Definition B.2 page 156}) \\
 &= \int_{y \in \mathbb{R}} \frac{d}{dy} [c_X(z - y) p_Y(y)] dy && \text{by linearity of } \frac{d}{dz} \\
 &= \int_{y \in \mathbb{R}} \left[\frac{d}{dy} c_X(z - y) \right] p_Y(y) dy && \text{because } y \text{ is fixed inside the integral} \\
 &\triangleq \int_y p_X(z - y) p_Y(y) dy && \text{by definition of } p_X && (\text{Definition B.2 page 156}) \\
 &= p_X(z) \star p_Y(z) && \text{by definition of } \star && (\text{Definition N.3 page 312})
 \end{aligned}$$



Theorem 4.5. Let

- X_1 and X_2 be random variables with joint distribution $p_{X_1, X_2}(x_1, x_2)$
- $Y_1 = f_1(x_1, x_2)$ and $Y_2 = f_2(x_1, x_2)$

⁹ Papoulis (1990) page 160 (Example 5.16)




Figure 4.5: Sum of random variables yields convolution of pdfs (Theorem 4.4 page 31)

Then the joint distribution of Y_1 and Y_2 is

$$\text{THM } p_{Y_1, Y_2}(y_1, y_2) = \frac{p_{X_1, X_2}(x_1, x_2)}{|J(x_1, x_2)|} = \frac{p_{X_1, X_2}(x_1, x_2)}{\begin{vmatrix} \frac{\partial f_1}{\partial x_1} & \frac{\partial f_1}{\partial x_2} \\ \frac{\partial f_2}{\partial x_1} & \frac{\partial f_2}{\partial x_2} \end{vmatrix}} = \frac{p_{X_1, X_2}(x_1, x_2)}{\frac{\partial f_1}{\partial x_1} \frac{\partial f_2}{\partial x_2} - \frac{\partial f_1}{\partial x_2} \frac{\partial f_2}{\partial x_1}}$$

Proposition 4.2. Let X and Y be random variables with joint distribution $p_{XY}(x, y)$ and

$$R^2 \triangleq X^2 + Y^2 \quad \theta \triangleq \tan^{-1} \frac{Y}{X}.$$

Then

$$\text{PRP } p_{R, \theta}(r, \theta) = r p_{XY}(r \cos \theta, r \sin \theta)$$

PROOF:

$$\begin{aligned} p_{R, \theta}(r, \theta) &= \frac{p_{XY}(x, y)}{|J(x, y)|} = \frac{p_{XY}(x, y)}{\begin{vmatrix} \frac{\partial R}{\partial x} & \frac{\partial R}{\partial y} \\ \frac{\partial \theta}{\partial x} & \frac{\partial \theta}{\partial y} \end{vmatrix}} = \frac{p_{XY}(x, y)}{\begin{vmatrix} \frac{x}{\sqrt{x^2+y^2}} & \frac{y}{\sqrt{x^2+y^2}} \\ \frac{-y}{x^2+y^2} & \frac{x}{x^2+y^2} \end{vmatrix}} \\ &= \frac{p_{XY}(x, y)}{\frac{x}{\sqrt{x^2+y^2}} \frac{x}{x^2+y^2} - \frac{y}{\sqrt{x^2+y^2}} \frac{-y}{x^2+y^2}} \\ &= \frac{p_{XY}(x, y)}{\frac{x^2+y^2}{(x^2+y^2)^{3/2}}} \\ &= p_{XY}(x, y) \frac{(x^2+y^2)^{3/2}}{x^2+y^2} \\ &= p_{XY}(r \cos \theta, r \sin \theta) \frac{r^3}{r^2} \\ &= r p_{XY}(r \cos \theta, r \sin \theta) \end{aligned}$$

Proposition 4.3. Let $X \sim N(0, \sigma^2)$ and $Y \sim N(0, \sigma^2)$ be independent random variables and

$$R^2 \triangleq X^2 + Y^2 \quad \theta \triangleq \tan^{-1} \frac{Y}{X}.$$



Then

- | | |
|----------------------------------|--|
| P
R
P | 1. R and Θ are independent with joint distribution $p_{R,\Theta}(r, \theta) = p_R(r)p_\theta(\theta)$
2. R has Rayleigh distribution $p_R(r) = \frac{r}{\sigma^2} \exp \frac{-r^2}{2\sigma^2}$
3. Θ has uniform distribution $p_\theta(\theta) = \frac{1}{2\pi}$ |
|----------------------------------|--|

PROOF:

$$\begin{aligned}
 p_{R,\Theta}(r, \theta) &= r p_{XY}(r\cos\theta, r\sin\theta) && \text{by Proposition 4.2 (page 32)} \\
 &= r p_X(r\cos\theta) p_Y(r\sin\theta) && \text{by independence hypothesis} \\
 &= r \frac{1}{\sqrt{2\pi\sigma^2}} \exp \frac{(r\cos\theta - 0)^2}{-2\sigma^2} \frac{1}{\sqrt{2\pi\sigma^2}} \exp \frac{(r\sin\theta - 0)^2}{-2\sigma^2} \\
 &= \frac{1}{2\pi\sigma^2} r \exp \frac{r^2(\cos^2\theta + \sin^2\theta)}{-2\sigma^2} \\
 &= \frac{1}{2\pi\sigma^2} r \exp \frac{r^2}{-2\sigma^2} \\
 &= \left[\frac{1}{2\pi} \right] \left[\frac{r}{\sigma^2} \exp \frac{r^2}{-2\sigma^2} \right]
 \end{aligned}$$

Proposition 4.4. Let X and Y be RANDOM VARIABLES with covariance σ_{xy} on a PROBABILITY SPACE (Ω, \mathbb{E}, P) .

P R P	$\left\{ \begin{array}{l} (A). X \text{ is GAUSSIAN with } N(\mu_X, \sigma_X^2) \text{ and} \\ (B). Y \text{ is GAUSSIAN with } N(\mu_Y, \sigma_Y^2) \text{ and} \\ (C). \sigma_{xy} = \text{cov}[X, Y] \end{array} \right\} \Rightarrow \left\{ P\{X > Y\} = Q\left(\frac{-\mu_X + \mu_Y}{\sigma_X^2 + \sigma_Y^2 - 2\sigma_{xy}}\right) \right\}$
----------------------------------	---

PROOF: Because X and Y are jointly Gaussian, their linear combination $Z = rvX - Y$ is also Gaussian. A Gaussian distribution is completely defined by its mean and variance. So, to determine the distribution of Z , we just have to determine the mean and variance of Z .

$$\begin{aligned}
 EZ &= EX - EY \\
 &= \mu_X - \mu_Y
 \end{aligned}$$

$$\begin{aligned}
 \text{var } Z &= EZ^2 - (EZ)^2 \\
 &= E(X - Y)^2 - (EX - EY)^2 \\
 &= E(X^2 - 2XY + Y^2) - [(EX)^2 - 2EXEY + (EY)^2] \\
 &= [EX^2 - (EX)^2] + [Y^2 - (EY)^2] - 2[EXY - EXEY] \\
 &= \text{var } X + \text{var } Y - 2\text{cov}[X, Y] \\
 &\triangleq \sigma_X^2 + \sigma_Y^2 - 2\sigma_{xy}
 \end{aligned}$$

$$\begin{aligned}
 P\{X > Y\} &= P\{X - Y > 0\} \\
 &= P\{Z > 0\} \\
 &= Q\left(\frac{z - EZ}{\text{var } Z}\right)\Big|_{z=0} \\
 &= Q\left(\frac{0 - \mu_X + \mu_Y}{\sigma_X^2 + \sigma_Y^2 - 2\sigma_{xy}}\right)
 \end{aligned}$$




Figure 4.6: The distributions of sums of independent uniformly distributed random variables (Example 4.1 page 34)


Example 4.1. Let (X_1, X_2, X_3, \dots) be a sequence of *independent* (Definition A.3 page 147) *uniformly distributed* random variables. Let $p_N(x)$ be the *probability density function* of $Y \triangleq \sum_{n=1}^N X_n$. Some of these distributions are illustrated in Figure 4.6 (page 34). Note that the distributions of the sequence (p_1, p_2, p_3, \dots) are all *B-splines* (Definition Q.2 page 343) and all form a *partition of unity*.

CHAPTER 5

OPERATORS ON DISCRETE RANDOM SEQUENCES

5.1 LTI operators on random sequences

Theorem 5.1. ¹ Let $x(n)$ be a RANDOM SEQUENCE with MEAN μ_X and $y(n)$ a RANDOM SEQUENCE with MEAN μ_Y . Let S be a system with IMPULSE RESPONSE $h(n)$, INPUT $x(n)$, and OUTPUT $y(n)$.



T H M

$$\{ \text{S is (LTI)} \} \implies \left\{ \begin{array}{l} (1). \quad \mu_Y(n) = \sum_{k \in \mathbb{Z}} h(k) \mu_X(n-k) \triangleq h(n) \star \mu_X(n) \text{ and} \\ (2). \quad R_{xy}(n, m) = \sum_{k \in \mathbb{Z}} h^*(k) R_{xx}(n-k, m+k) \\ (3). \quad R_{yy}(n, m) = \sum_{k \in \mathbb{Z}} h^*(k) R_{yx}(n-k, m+k) \end{array} \right\}$$

PROOF:

$$\begin{aligned}
 \mu_Y(n) &\triangleq E[y(n)] && \text{by definition of } \mu_Y && (\text{Definition 2.2 page 11}) \\
 &= E\left[\sum_{k \in \mathbb{Z}} h(k)x(n-k)\right] && \text{by LTI hypothesis} \\
 &= \sum_{k \in \mathbb{Z}} h(k)E[x(n-k)] && \text{by linear property} \\
 &= \sum_{k \in \mathbb{Z}} h(k)\mu_X(n-k) && \text{by definition of } \mu_X && (\text{Definition 2.2 page 11}) \\
 &\triangleq h(n) \star \mu_X(n) && \text{by definition of convolution} && (\text{Definition P.3 page 329})
 \end{aligned}$$

$$\begin{aligned}
 R_{xy}(n, m) &\triangleq E[x(n+m)y^*(n)] && \text{by definition of } R_{xy}(n, m) && (\text{Definition 2.2 page 11}) \\
 &= E[x(n+m)(h(n) \star x(n))^*] && \text{by LTI hypothesis} \\
 &\triangleq E\left[x(n+m)\left(\sum_{k \in \mathbb{Z}} h(k)x(n-k)\right)^*\right] && \text{by definition of convolution } \star && (\text{Definition P.3 page 329}) \\
 &= E\left[x(n+m) \sum_{k \in \mathbb{Z}} h^*(k)x^*(n-k)\right] && \text{by distributive property of } *-\text{algebras} && (\text{Definition H.3 page 228})
 \end{aligned}$$

¹ Papoulis (1991), page 310

$$\begin{aligned}
 &= E \left[\sum_{k \in \mathbb{Z}} h^*(k) x(n+m) x^*(n-k) \right] && \text{by } \textit{distributive property of } (\mathbb{C}, +, \cdot, 0, 1) \\
 &= \sum_{k \in \mathbb{Z}} h^*(k) E[x(n-k+k+m) x^*(n-k)] && \text{by } \textit{linear property of } E \quad (\text{Theorem 1.1 page 4}) \\
 &\triangleq \sum_{k \in \mathbb{Z}} h^*(k) R_{xx}(n-k, m+k) && \text{by definition of } R_{xx}(n, m) \quad (\text{Definition 2.2 page 11})
 \end{aligned}$$

$$\begin{aligned}
 R_{yy}(n, m) &\triangleq E[y(n+m) y^*(n)] && \text{by definition of } R_{xy}(n, m) \quad (\text{Definition 2.2 page 11}) \\
 &= E[y(n+m)(h(n) \star x(n))^*] && \text{by LTI hypothesis} \\
 &\triangleq E \left[y(n+m) \left(\sum_{k \in \mathbb{Z}} h(k) x(n-k) \right)^* \right] && \text{by definition of convolution} \quad (\text{Definition P.3 page 329}) \\
 &= E \left[y(n+m) \sum_{k \in \mathbb{Z}} h^*(k) x^*(n-k) \right] && \text{by distributive property of } *-\text{algebras} \quad (\text{Definition H.3 page 228}) \\
 &= E \left[\sum_{k \in \mathbb{Z}} h^*(k) y(n+m) x^*(n-k) \right] && \text{by distributive property of } (\mathbb{C}, +, \cdot, 0, 1) \\
 &= \sum_{k \in \mathbb{Z}} h^*(k) E[y(n-k+k+m) x^*(n-k)] && \text{by linear property of } E \quad (\text{Theorem 1.1 page 4}) \\
 &\triangleq \sum_{k \in \mathbb{Z}} h^*(k) R_{yx}(n-k, m+k) && \text{by definition of } R_{xy}(n, m) \quad (\text{Definition 2.2 page 11})
 \end{aligned}$$

⇒

5.2 LTI operators on WSS random sequences

Corollary 5.1. Let S be the system defined in Theorem 5.1 (page 35).

COR

$$\left. \begin{array}{l} (A). \quad S \text{ is LTI} \\ (B). \quad x(n) \text{ is WSS} \end{array} \right\} \implies \left\{ \begin{array}{l} (1). \quad \mu_Y = \mu_X \sum_{n \in \mathbb{Z}} h(k) \quad \text{and} \\ (2). \quad R_{xy}(m) = R_{xx}(m) \star h^*(-m) \triangleq \sum_{k \in \mathbb{Z}} h^*(k) R_{xx}(m+k) \quad \text{and} \\ (3). \quad R_{yy}(m) = R_{yx}(m) \star h^*(-m) \triangleq \sum_{k \in \mathbb{Z}} h^*(k) R_{xy}(m+k) \quad \text{and} \\ (4). \quad R_{yy}(m) = R_{xx}^*(m) \star h(-m) \star h^*(-m) \end{array} \right.$$

PROOF:

$$\begin{aligned}
 \mu_Y &= \mu_Y(n) && \text{by Proposition 2.1 page 13} && \text{and hypothesis (A)} \\
 &= \sum_{n \in \mathbb{Z}} h(k) \mu_X(n-k) && \text{by Theorem 2.1 page 12} && \text{and hypothesis (B)} \\
 &= \sum_{n \in \mathbb{Z}} h(k) \mu_X(0) && \text{by Definition 6.1 page 43} && \text{and hypothesis (B)} \\
 &= \mu_X(0) \sum_{n \in \mathbb{Z}} h(k) && \text{by linear property of } \sum && \\
 &= \mu_X \sum_{n \in \mathbb{Z}} h(k) && \text{by Proposition 2.1 page 13} &&
 \end{aligned}$$

¹  Papoulis (1991), page 323

$$\begin{aligned}
 R_{xy}(m) &\triangleq R_{xy}(0, m) && \text{by Proposition 2.1 page 13} && \text{and hypothesis (A)} \\
 &= \sum_{k \in \mathbb{Z}} h^*(k) R_{xx}(0 - k, m + k) && \text{by Theorem 5.1 page 35} && \text{and hypothesis (B)} \\
 &= \sum_{k \in \mathbb{Z}} h^*(k) R_{xx}(m + k) && \text{by Proposition 2.1 page 13} && \text{and hypothesis (A)} \\
 &= h^*(-m) \star R_{xx}(m) && \text{by Proposition P.2 page 330} && \\
 R_{yy}(m) &\triangleq R_{yy}(0, m) && \text{by Proposition 2.1 page 13} && \text{and hypothesis (A)} \\
 &= \sum_{k \in \mathbb{Z}} h^*(k) R_{yx}(n - k, m + k) && \text{by Theorem 5.1 page 35} && \text{and hypothesis (B)} \\
 &= \sum_{k \in \mathbb{Z}} h^*(k) R_{yx}(m + k) && \text{by Proposition 2.1 page 13} && \text{and hypothesis (A)} \\
 &= h^*(-m) \star R_{yx}(m) && \text{by Proposition P.2 page 330} && \\
 R_{yy}(m) &= h^*(-m) \star R_{yx}(m) && \text{by result (2)} && \\
 &= h^*(-m) \star R_{xy}^*(m) && \text{by Corollary 2.1 page 13} && \\
 &= h^*(-m) \star [h^*(-m) \star R_{xx}(m)]^* && \text{by result (1)} && \\
 &= h^*(-m) \star h(-m) \star R_{xx}^*(m) && \text{by } \textit{distributive property of } *-\textbf{algebras} && \text{(Definition H.3 page 228)}
 \end{aligned}$$

⇒

Corollary 5.2. ² Let \mathbf{S} be a system with IMPULSE RESPONSE $h(n)$, INPUT $x(n)$, and OUTPUT $y(n)$.

COR

$$\left\{ \begin{array}{l} (A). \quad h \text{ is LINEAR TIME INVARIANT and} \\ (B). \quad x \text{ and } y \text{ are WIDE SENSE STATIONARY} \end{array} \right\} \implies \left\{ \begin{array}{l} (1). \quad \check{S}_{xy}(z) = \check{S}_{xx}(z)\check{H}^*\left(\frac{1}{z^*}\right) \text{ and} \\ (2). \quad \check{S}_{yy}(z) = \check{S}_{yx}(z)\check{H}^*\left(\frac{1}{z^*}\right) \text{ and} \\ (3). \quad \check{S}_{yy}(z) = \check{S}_{xx}(z)\check{H}(z)\check{H}^*\left(\frac{1}{z^*}\right) \end{array} \right\}$$

⇒

PROOF: The proof is given in Proposition ?? (page ??) (1).

⇒

Corollary 5.3. Let \mathbf{S} be a system with IMPULSE RESPONSE $h(n)$, INPUT $x(n)$, and OUTPUT $y(n)$.

COR

$$\left\{ \begin{array}{l} (A). \quad h \text{ is LTI and} \\ (B). \quad x \text{ and } y \text{ are WSS} \end{array} \right\} \implies \left\{ \begin{array}{l} (1). \quad \tilde{S}_{xy}(\omega) = \tilde{S}_{xx}(\omega)\tilde{H}^*(\omega) \text{ and} \\ (2). \quad \tilde{S}_{yy}(\omega) = \tilde{S}_{xy}(\omega)\tilde{H}(\omega) \text{ and} \\ (3). \quad \tilde{S}_{yy}(\omega) = \tilde{S}_{xx}(\omega)|\tilde{H}(\omega)|^2 \end{array} \right\}$$

⇒

PROOF: The proof is given in Proposition ?? (page ??) (1).


⇒

5.3 Parallel operators on WSS random sequences

Theorem 5.2. Let \mathbf{S} be the SYSTEM illustrated to the right, where \mathbf{T} is NOT NECESSARILY LINEAR. Let

$$(\mathbf{h}(n)) \triangleq \mathbf{H}\delta(n) \triangleq \sum_{m \in \mathbb{Z}} h(m)\delta(n - m)$$

be the IMPULSE RESPONSE of \mathbf{H} .



² Papoulis (1991), page 323

THM

$$\left\{ \begin{array}{l} \text{(A). } \mathbf{x}(n) \text{ is WSS and} \\ \text{(B). } \mathbf{H} \text{ is LTI} \end{array} \right\} \Rightarrow \left\{ \begin{array}{ll} \text{(1). } R_{wy}(m) &= \sum_{n \in \mathbb{Z}} h(n)R_{xy}(m-n) \text{ (convolution)} \\ &\triangleq h(m) \star R_{xy}(m) \quad \text{and} \\ \text{(2). } \check{S}_{wy}(z) &= \check{H}(z)\check{S}_{xy}(z) \quad \text{and} \\ \text{(3). } \tilde{S}_{wy}(\omega) &= \tilde{H}(\omega)\tilde{S}_{xy}(\omega) \end{array} \right\}$$

PROOF:

$$\begin{aligned}
 R_{wy}(m) &\triangleq E[\mathbf{w}(m)\mathbf{y}^*(0)] && \text{by (A) and definition of } R_{wy} && \text{(Definition 2.4 page 12)} \\
 &\triangleq E[(\mathbf{H}\mathbf{x})(m)\mathbf{y}^*(0)] && \text{by definition of } \mathbf{S} \\
 &= \mathbf{H}E(\mathbf{x}(m)\mathbf{y}^*(0)) && \text{by LTI hypothesis} && \text{(B)} \\
 &\triangleq \mathbf{H}R_{xy}(m) && \text{by definition of } R_{xy} && \text{(Definition 2.4 page 12)} \\
 &= \sum_{n \in \mathbb{Z}} h(n)R_{xy}(m-n) && \text{by definition of } \mathbf{H} \text{ impulse response } (h(n)) \\
 &= [h(m) \star R_{xy}(m)] && \text{by definition of convolution} && \text{(Definition P.3 page 329)} \\
 \check{S}_{wy}(z) &\triangleq ZR_{wy}(m) && \text{by definition of } \check{S}_{wy} && \text{(Definition 2.5 page 14)} \\
 &= [h(m) \star R_{xy}(m)] && \text{by previous result} \\
 &= \check{H}(z)\check{S}_{xy}(z) && \text{by Convolution Theorem} && \text{(Theorem P.2 page 332)} \\
 \tilde{S}_{wy}(\omega) &\triangleq \check{F}R_{wy}(m) && \text{by definition of } \tilde{S}_{wy} && \text{(Definition 6.3 page 44)} \\
 &= [h(m) \star R_{xy}(m)] && \text{by previous result} \\
 &= \tilde{H}(\omega)\tilde{S}_{xy}(\omega) && \text{by Convolution Theorem} && \text{(Theorem P.2 page 332)}
 \end{aligned}$$

⇒

5.4 Whitening discrete random sequences




Figure 5.1: Poles (x) and zeros (o) of a *minimum phase* filter

Definition 5.1. Let $\check{H}(z)$ be the z-transform of the impulse response of a filter. If $\check{H}(z)$ can be expressed as a rational expression with poles and zeros $r_n e^{i\theta_n}$, then the filter is **minimum phase** if each $r_n < 1$ (all roots lie inside the unit circle in the complex z-plane).

See Figure 5.1 page 38.

Note that if $L(z)$ has a root at $z = re^{i\theta}$, then $L^*(1/z^*)$ has a root at

$$\frac{1}{z^*} = \frac{1}{(re^{i\theta})^*} = \frac{1}{re^{-i\theta}} = \frac{1}{r}e^{i\theta}.$$

That is, if $L(z)$ has a root inside the unit circle, then $L^*(1/z^*)$ has a root directly opposite across the unit circle boundary (see Figure 5.2 page 39). A causal stable filter $\check{H}(z)$ must have all of its poles inside the unit circle. A minimum phase filter is a filter with both its poles and zeros inside the unit circle. One advantage of a minimum phase filter is that its reciprocals (zeros become poles and poles become zeros) is also causal and stable.




Figure 5.2: Mirrored roots in complex-z plane




Figure 5.3: Innovations and whitening filters

The next theorem demonstrates a method for “whitening” a *random sequence* $x(n)$ with a filter constructed from a decomposition of $R_{xx}(m)$. The technique is stated precisely in Theorem 5.3 page 39 and illustrated in Figure 5.3 page 39. Both imply two filters with impulse responses $l(n)$ and $\gamma(n)$. Filter $l(n)$ is referred to as the **innovations filter** (because it generates or “innovates” $x(n)$ from a white noise process $w(n)$) and $\gamma(n)$ is referred to as the **whitening filter** because it produces a white noise sequence when the input sequence is $x(n)$.³

Theorem 5.3. Let $x(n)$ be a WSS RANDOM SEQUENCE with auto-correlation $R_{xx}(m)$ and spectral density $\check{S}_{xx}(z)$. If $\check{S}_{xx}(z)$ has a rational expression, then the following are true:

1. There exists a rational expression $L(z)$ with minimum phase such that

$$\check{S}_{xx}(z) = L(z)L^*\left(\frac{1}{z^*}\right).$$

2. An LTI filter for which the Laplace transform of the impulse response $\gamma(n)$ is

$$\Gamma(z) = \frac{1}{L(z)}$$

is both causal and stable.

3. If $x(n)$ is the input to the filter $\gamma(n)$, the output $y(n)$ is a **white noise sequence** such that

$$S_{yy}(z) = 1 \quad R_{yy}(m) = \bar{\delta}(m).$$

³ Papoulis (1991), pages 401–402

PROOF:

$$\begin{aligned} S_{ww}(z) &= \Gamma(z)\Gamma^*\left(\frac{1}{z^*}\right)\check{S}_{xx}(z) \\ &= \frac{1}{L(z)}\frac{1}{L^*\left(\frac{1}{z^*}\right)}\check{S}_{xx}(z) \\ &= \frac{1}{L(z)}\frac{1}{L^*\left(\frac{1}{z^*}\right)}L(z)L^*\left(\frac{1}{z^*}\right) \\ &= 1 \end{aligned}$$



CHAPTER 6

OPERATORS ON CONTINUOUS RANDOM SEQUENCES

6.1 LTI Operations on non-stationary random processes




Figure 6.1: Linear system with random process input and output

Theorem 6.1. ¹ Let $h : \mathbb{R} \rightarrow \mathbb{C}$ be the impulse response of a linear time-invariant system and let $y(t) = h(t) \star x(t) \triangleq \int_{u \in \mathbb{R}} h(u)x(t-u) du$ as illustrated in Figure 6.1 page 41. Then

Correlation functions

$$\begin{aligned} R_{xy}(t, u) &= R_{xx}(t, u) \star h^*(u) &\triangleq \int_v h^*(v)R_{xx}(t, u-v) dv \\ R_{yy}(t, u) &= R_{xy}(t, u) \star h(t) &\triangleq \int_v h(v)R_{xy}(t-v, u) dv \\ R_{yy}(t, u) &= R_{xx}(t, u) \star h(t) \star h^*(u) &\triangleq \int_w h^*(w) \int_v h(v)R_{xx}(t-v, u-w) dv dw \end{aligned}$$

Laplace power spectral density functions

$$\begin{aligned} \check{S}_{xy}(s, r) &= \check{S}_{xx}(s, r)\check{h}^*(r^*) \\ \check{S}_{yy}(s, r) &= \check{S}_{xy}(s, r)\check{h}(s) \\ \check{S}_{yy}(s, r) &= \check{S}_{xx}(s, r)\check{h}(s)\check{h}^*(r^*) \end{aligned}$$

Power spectral density functions

$$\begin{aligned} S_{xy}(f, g) &= S_{xx}(f, g)\tilde{h}^*(-g) \\ S_{yy}(f, g) &= S_{xy}(f, g)\tilde{h}(\omega) \\ S_{yy}(f, g) &= S_{xx}(f, g)\tilde{h}(\omega)\tilde{h}^*(-g) \end{aligned}$$

T
H
M

¹ Papoulis (1991), page 312

PROOF:

$$\begin{aligned}
 R_{xy}(t, u) &\triangleq E[x(t)y^*(u)] \\
 &= E\left[x(t)\left(\int_v h(v)x(u-v) dv\right)^*\right] \\
 &= E\left[x(t)\int_v h^*(v)x^*(u-v) dv\right] \\
 &= \int_v h^*(v)E[x(t)x^*(u-v)] dv \\
 &= \int_v h^*(v)R_{xx}(t, u-v) dv \\
 &\triangleq R_{xx}(t, u) \star h^*(u)
 \end{aligned}$$

$$\begin{aligned}
 R_{yy}(t, u) &\triangleq E[y(t)y^*(u)] \\
 &= E\left[\left(\int_v h(v)x(t-v) dv\right)y^*(u)\right] \\
 &= \int_v h(v)E[x(t-v)y^*(u)] dv \\
 &= \int_v h(v)R_{xy}(t-v, u) dv \\
 &\triangleq R_{xy}(t, u) \star h(t)
 \end{aligned}$$

$$\begin{aligned}
 R_{yy}(t, u) &\triangleq E[y(t)y^*(u)] \\
 &= E\left[\left(\int_v h(v)x(t-v) dv\right)\left(\int_w h(w)x(u-w) dw\right)^*\right] \\
 &= \int_w h^*(w) \int_v h(v)E[x(t-v)x^*(u-w)] dv dw \\
 &= \int_w h^*(w) \int_v h(v)R_{xx}(t-v, u-w) dv dw \\
 &= \int_w h^*(w)[R_{xx}(t, u-w) \star h(t)] dw \\
 &\triangleq R_{xx}(t, u) \star h(t) \star h^*(u)
 \end{aligned}$$

$$\begin{aligned}
 \check{S}_{xy}(s, r) &\triangleq L[R_{xy}(t, u)] \\
 &= L[R_{xx}(t, u) \star h^*(u)] \\
 &= L[R_{xx}(t, u)]L[h^*(u)] \\
 &= \check{S}_{xx}(s, r) \int_{u \in \mathbb{R}} h^*(u)e^{-ru} du \\
 &= \check{S}_{xx}(s, r) \left[\int_{u \in \mathbb{R}} h(u)e^{-r^*u} du \right]^* \\
 &= \check{S}_{xx}(s, r) \check{h}^*(r^*)
 \end{aligned}$$

$$\begin{aligned}
 \check{S}_{yy}(s, r) &\triangleq L[R_{yy}(t, u)] \\
 &= L[R_{xy}(t, u) \star h(t)] \\
 &= L[R_{xy}(t, u)]L[h(t)] \\
 &= \check{S}_{xy}(s, r) \check{h}(s)
 \end{aligned}$$

$$\begin{aligned}
 &= \check{S}_{xy}(s, r)\check{h}(s) \\
 &= \check{S}_{xx}(s, r)\check{h}^*(r^*)\check{h}(s) \\
 &= \check{S}_{xx}(s, r)\check{h}(s)\check{h}^*(r^*)
 \end{aligned}$$

$$\begin{aligned}
 S_{xy}(f, g) &\triangleq \tilde{\mathbf{F}}R_{xy}(t, u) \\
 &= \tilde{\mathbf{F}}[R_{xx}(t, u) \star h^*(u)] \\
 &= \tilde{\mathbf{F}}[R_{xx}(t, u)]\tilde{\mathbf{F}}[h^*(u)] \\
 &= S_{xx}(f, g) \int_{u \in \mathbb{R}} h^*(u)e^{-i2\pi gu} du \\
 &= S_{xx}(f, g) \left[\int_{u \in \mathbb{R}} h(u)e^{i2\pi gu} du \right]^* \\
 &= S_{xx}(f, g) \left[\int_{u \in \mathbb{R}} h(u)e^{-i2\pi(-g)u} du \right]^* \\
 &= S_{xx}(f, g)\tilde{h}^*(-g)
 \end{aligned}$$

$$\begin{aligned}
 S_{yy}(f, g) &\triangleq \tilde{\mathbf{F}}R_{yy}(t, u) \\
 &= \tilde{\mathbf{F}}[R_{xy}(t, u) \star h(t)] \\
 &= \tilde{\mathbf{F}}[R_{xy}(t, u)]\tilde{\mathbf{F}}[h(t)] \\
 &= S_{xy}(f, g)\tilde{h}(\omega) \\
 &= S_{xy}(f, g)\tilde{h}^*(-g)\tilde{h}(\omega)
 \end{aligned}$$



6.2 LTI Operations on WSS random processes

Definition 6.1.

DEF A random process $x(t)$ is **wide sense stationary (WSS)** if

- (1). $\mu_X(t)$ is CONSTANT with respect to t (STATIONARY IN THE MEAN) and
- (2). $R_{xx}(t + \tau, t)$ is CONSTANT with respect to t (STATIONARY IN CORRELATION)

If a process $x(t)$ is *wide sense stationary*, mean and correlation are often written μ_X and $R_{xx}(\tau)$, respectively. If a pair of processes $x(t)$ and $y(t)$ are WSS, then their cross-correlation is commonly written $R_{xy}(\tau)$.

Definition 6.2. Let $x(t)$ and $y(t)$ be WSS random processes.

D<small>EF</small>	$\check{S}_{xx}(s) \triangleq \mathbf{L}R_{xx}(\tau) \triangleq \int_{\tau \in \mathbb{R}} R_{xx}(\tau)e^{-s\tau} d\tau$ $\check{S}_{yy}(s) \triangleq \mathbf{L}R_{yy}(\tau) \triangleq \int_{\tau \in \mathbb{R}} R_{yy}(\tau)e^{-s\tau} d\tau$ $\check{S}_{xy}(s) \triangleq \mathbf{L}R_{xy}(\tau) \triangleq \int_{\tau \in \mathbb{R}} R_{xy}(\tau)e^{-s\tau} d\tau$ $\check{S}_{yx}(s) \triangleq \mathbf{L}R_{yx}(\tau) \triangleq \int_{\tau \in \mathbb{R}} R_{yx}(\tau)e^{-s\tau} d\tau$
---------------------------	---

Definition 6.3. Let $x(t)$ and $y(t)$ be WSS random processes.

DEF

$$\begin{aligned}\tilde{S}_{xx}(\omega) &\triangleq [\tilde{\mathbf{F}}R_{xx}(\tau)](\omega) \triangleq \int_{\tau \in \mathbb{R}} R_{xx}(\tau) e^{-i\omega\tau} d\tau \\ \tilde{S}_{yy}(\omega) &\triangleq [\tilde{\mathbf{F}}R_{yy}(\tau)](\omega) \triangleq \int_{\tau \in \mathbb{R}} R_{yy}(\tau) e^{-i\omega\tau} d\tau \\ \tilde{S}_{xy}(\omega) &\triangleq [\tilde{\mathbf{F}}R_{xy}(\tau)](\omega) \triangleq \int_{\tau \in \mathbb{R}} R_{xy}(\tau) e^{-i\omega\tau} d\tau \\ \tilde{S}_{yx}(\omega) &\triangleq [\tilde{\mathbf{F}}R_{yx}(\tau)](\omega) \triangleq \int_{\tau \in \mathbb{R}} R_{yx}(\tau) e^{-i\omega\tau} d\tau\end{aligned}$$

Definition 6.4. ² Let $x(t)$ be a random variable that is STATIONARY IN THE MEAN such that $E[x(t)]$ is constant with respect to t .

DEF

$x(t)$ is ergodic in the mean if

$$E[\underbrace{x(t)}_{\text{ENSEMBLE AVERAGE}}] = \lim_{\tau \rightarrow \infty} \frac{1}{2\tau} \underbrace{\int_{-\tau}^{+\tau} x(t) dt}_{\text{TIME AVERAGE}}$$

Proposition 6.1.

PRP

$$\{x(t) \text{ is NON-STATIONARY}\} \implies \{x(t) \text{ is NOT ERGODIC IN THE MEAN}\}$$

PROOF: If $x(t)$ is non-stationary, then $E[x(t)]$ is not constant with time. But $\lim_{\tau \rightarrow \infty} \frac{1}{2\tau} \int_{-\tau}^{+\tau} x(t) dt$ must be a constant (if it is convergent). \Rightarrow

Definition 6.5. ³ Let $x(t)$ be a WIDE SENSE STATIONARY random process.

DEF

- (1). The average power P_{avg} of $x(t)$ is $P_{avg}x(t) \triangleq \lim_{\tau \rightarrow \infty} \frac{1}{2\tau} \int_{t \in \mathbb{R}} |x(t)|^2 dt$
- (2). The energy spectral density $|\tilde{x}(\omega)|^2$ of $x(t)$ is $|\tilde{x}(\omega)|^2 \triangleq |\tilde{\mathbf{F}}x(t)|^2$

Remark 6.1 (spectral power). Why does $\tilde{S}_{xx}(\omega)$ deserve the name *power spectral density*? This is answered by Theorem 6.2 (next). But to elaborate further, note that \tilde{S}_{xx} is the spectral representation of the statistical relationship (the *variance*) between samples of $x(t)$. For example, if there is no relationship, then $\tilde{S}_{xx}(\omega) = 1$. But in the case that $x(t)$ is ergodic in the mean, then \tilde{S}_{xx} takes on an additional meaning—it describes the “spectral power” present in $x(t)$. This is demonstrated by the next theorem.

Theorem 6.2. Let $x(t)$ be a RANDOM PROCESS.

THM

$$\left\{ \begin{array}{l} (A). \quad x(t) \text{ IS ERGODIC IN THE MEAN} \quad \text{and} \\ (B). \quad \tilde{x}(\omega) \text{ EXISTS} \end{array} \right\} \implies \left\{ \begin{array}{l} (1). \quad \tilde{S}_{xx}(\omega) = \lim_{\tau \rightarrow \infty} \frac{1}{2\tau} \underbrace{\int_{t \in \mathbb{R}} |x(t)|^2 dt}_{(\text{ESD})} \quad \text{and} \\ (2). \quad P_{avg}[x(t)] = \int_{\omega \in \mathbb{R}} \tilde{S}_{xx}(\omega) d\omega \end{array} \right\}$$

² Papoulis (1984) page 246 (Mean-Ergodic processes), Papoulis (2002) page 523 (12-1 ERGODICITY), KAY (1988) PAGE 58 (3.6 ERGODICITY OF THE AUTOCORRELATION FUNCTION), MANOLAKIS ET AL. (2005) PAGE 106 (ERGODIC RANDOM PROCESSES), KOOPMANS (1995) PAGES 53–61, CADZOW (1987) PAGE 378 (11.13 ERGODIC TIME SERIES), HELSTROM (1991) PAGE 336

³ ? page 177

PROOF:

$$\begin{aligned}
 \tilde{S}_{xx}(\omega) &\triangleq \int_{\tau \in \mathbb{R}} R_{xx}(\tau) e^{-i\omega\tau} d\tau && \text{by definition of } \tilde{S}_{xx}(\omega) && (\text{Definition 6.3 page 44}) \\
 &= \int_{\tau \in \mathbb{R}} E[x(t + \tau)x^*(t)] e^{-i\omega\tau} d\tau && \text{by definition of } R_{xx}(t) && (\text{Definition 3.2 page 18}) \\
 &= E\left[x^*(t) \int_{\tau \in \mathbb{R}} x(t + \tau)e^{-i\omega\tau} d\tau\right] && \text{by linearity of } E \text{ operator} \\
 &= E\left[x^*(t) \int_{u \in \mathbb{R}} x(u)e^{-i\omega(u-t)} du\right] && \text{where } u \triangleq t + \tau \implies \tau = u - t \\
 &= E\left[x^*(t)e^{i\omega t} \int_{u \in \mathbb{R}} x(u)e^{-i\omega u} du\right] \\
 &= E[x^*(t)e^{i\omega t} \tilde{x}(\omega)] && \text{by definition of Fourier Transform} && (\text{Definition N.2 page 309}) \\
 &= E[x^*(t)e^{i\omega t}] \tilde{x}(\omega) && \text{by hypothesis (B)} \\
 &= \left[\lim_{\tau \rightarrow \infty} \frac{1}{2\tau} \int_{-\tau}^{+\tau} x^*(t)e^{i\omega t} dt \right] \tilde{x}(\omega) && \text{by ergodic in the mean hypothesis} && (\text{Definition 6.4 page 44}) \\
 &= \lim_{\tau \rightarrow \infty} \frac{1}{2\tau} \left[\int_{t \in \mathbb{R}} x(t)e^{-i\omega t} dt \right]^* \tilde{x}(\omega) \\
 &= \lim_{\tau \rightarrow \infty} \frac{1}{2\tau} \tilde{x}^*(\omega) \tilde{x}(\omega) && \text{by hypothesis (B)} \\
 &= \lim_{\tau \rightarrow \infty} \frac{1}{2\tau} |\tilde{x}(\omega)|^2
 \end{aligned}$$

$$\begin{aligned}
 \int_{\omega \in \mathbb{R}} \tilde{S}_{xx}(\omega) d\omega &= \int_{\omega \in \mathbb{R}} \lim_{\tau \rightarrow \infty} \frac{1}{2\tau} |\tilde{x}(\omega)|^2 d\omega \\
 &= \lim_{\tau \rightarrow \infty} \frac{1}{2\tau} \int_{\omega \in \mathbb{R}} |\tilde{x}(\omega)|^2 d\omega \\
 &= \lim_{\tau \rightarrow \infty} \frac{1}{2\tau} \int_{t \in \mathbb{R}} |x(t)|^2 dt && \text{by Plancheral's formula} && (\text{Theorem N.3 page 311, Theorem G.9 page 210}) \\
 &= P_{avg} && \text{by definition of } P_{avg} && (\text{Definition 6.5 page 44})
 \end{aligned}$$

Thus, $\tilde{S}_{xx}(\omega)$ is the power density of $x(t)$ in the frequency domain.

⇒




Figure 6.2: Linear system with WSS random process input and output

Theorem 6.3. ⁴ Let $h : \mathbb{R} \rightarrow \mathbb{C}$ be the impulse response of a linear time-invariant system and let $y(t) = h(t) \star x(t) \triangleq \int_{u \in \mathbb{R}} h(u)x(t-u) du$ as illustrated in Figure 6.1 page 41. Then

⁴ Papoulis (1991), pages 323–324

THM

$$\begin{aligned} R_{xy}(\tau) &= R_{xx}(\tau) \star h^*(-\tau) &\triangleq \int_{u \in \mathbb{R}} h^*(-u) R_{xx}(\tau - u) du \\ R_{yy}(\tau) &= R_{xy}(\tau) \star h(\tau) &\triangleq \int_{u \in \mathbb{R}} h(u) R_{xy}(\tau - u) du \\ R_{yy}(\tau) &= R_{xx}(\tau) \star h(\tau) \star h^*(-\tau) &\triangleq \int_v \int_{u \in \mathbb{R}} h(u - v) h^*(-v) R_{xx}(\tau - u - v) du dv \end{aligned}$$

$$\begin{aligned} S_{xy}(s) &= S_{xx}(s) \hat{h}^*(-s^*) \\ S_{yy}(s) &= S_{xy}(s) \hat{h}(s) \\ S_{yy}(s) &= S_{xx}(s) \hat{h}(s) \hat{h}^*(-s^*) \end{aligned}$$

$$\begin{aligned} \tilde{S}_{xy}(\omega) &= \tilde{S}_{xx}(\omega) \tilde{h}^*(\omega) \\ \tilde{S}_{yy}(\omega) &= \tilde{S}_{xy}(\omega) \tilde{h}(\omega) \\ \tilde{S}_{yy}(\omega) &= \tilde{S}_{xx}(\omega) |\tilde{h}(\omega)|^2 \end{aligned}$$

PROOF:

$$\begin{aligned} R_{xx}(\tau) \star h^*(-\tau) &\triangleq \int_{u \in \mathbb{R}} h^*(-u) R_{xx}(\tau - u) du \\ &= \int_{u \in \mathbb{R}} h^*(-u) E[x(t)x^*(t - \tau + u)] du \\ &= E \left[x(t) \int_{u \in \mathbb{R}} h^*(-u) x^*(t - \tau + u) du \right] \\ &= E \left[x(t) \int_{u \in \mathbb{R}} h^*(u') x^*(t - \tau - u') du' \right] \\ &= E[x(t)y^*(t - \tau)] \\ &\triangleq R_{xy}(\tau) \end{aligned}$$

$$\begin{aligned} R_{xy}(\tau) \star h(\tau) &\triangleq \int_{u \in \mathbb{R}} h(u) R_{xy}(\tau - u) du \\ &= \int_{u \in \mathbb{R}} h(u) E[x(t + \tau - u)y^*(t)] du \\ &= E \left[y^*(t) \int_{u \in \mathbb{R}} h(u) x(t + \tau - u) du \right] \\ &= E[y^*(t)y(t + \tau)] \\ &= E[y(t + \tau)y^*(t)] \\ &\triangleq R_{yy}(\tau) \end{aligned}$$

$$\begin{aligned} R_{yy}(\tau) &= R_{xy}(\tau) \star h(\tau) \\ &= R_{xx}(\tau) \star h^*(-\tau) \star h(\tau) \\ &= R_{xx}(\tau) \star h(\tau) \star h^*(-\tau) \end{aligned}$$

$$\begin{aligned} S_{xy}(s) &\triangleq LR_{xy}(\tau) \\ &\triangleq \int_{\tau \in \mathbb{R}} R_{xy}(\tau) e^{-s\tau} d\tau \\ &= \int_{\tau \in \mathbb{R}} [R_{xx}(\tau) \star h^*(-\tau)] e^{-s\tau} d\tau \\ &= \int_{\tau \in \mathbb{R}} \left[\int_{u \in \mathbb{R}} h^*(-u) R_{xx}(\tau - u) du \right] e^{-s\tau} d\tau \\ &= \int_{u \in \mathbb{R}} h^*(-u) \int_{\tau \in \mathbb{R}} R_{xx}(\tau - u) e^{-s\tau} d\tau du \end{aligned}$$



$$\begin{aligned}
&= \int_{u \in \mathbb{R}} h^*(-u) \int_v R_{xx}(v) e^{-s(v+u)} dv du \\
&= \int_{u \in \mathbb{R}} h^*(-u) e^{-su} du \int_v R_{xx}(v) e^{-sv} dv \\
&= \int_{u \in \mathbb{R}} h^*(u) e^{-s(-u)} du \int_v R_{xx}(v) e^{-sv} dv \\
&= \left(\int_{u \in \mathbb{R}} h(u) e^{-(s^*)u} du \right)^* \int_v R_{xx}(v) e^{-sv} dv \\
&\triangleq \hat{h}^*(-s^*) S_{xx}(s)
\end{aligned}$$

where $v = \tau - u \iff \tau = v + u$

$$\begin{aligned}
S_{yy}(s) &\triangleq \mathbf{L}R_{yy}(\tau) \\
&\triangleq \int_{\tau \in \mathbb{R}} R_{yy}(\tau) e^{-s\tau} d\tau \\
&= \int_{\tau \in \mathbb{R}} [R_{xy}(\tau) \star h(\tau)] e^{-s\tau} d\tau \\
&= \int_{\tau \in \mathbb{R}} \left[\int_{u \in \mathbb{R}} h(u) R_{xy}(\tau - u) du \right] e^{-s\tau} d\tau \\
&= \int_{u \in \mathbb{R}} h(u) \int_{\tau \in \mathbb{R}} R_{xy}(\tau - u) e^{-s\tau} d\tau du \\
&= \int_{u \in \mathbb{R}} h(u) \int_{\tau \in \mathbb{R}} R_{xy}(v) e^{-s(v+u)} d\tau du \\
&= \int_{u \in \mathbb{R}} h(u) e^{-su} du \int_{\tau \in \mathbb{R}} R_{xy}(v) e^{-sv} d\tau \\
&\triangleq \hat{h}(s) S_{xy}(s)
\end{aligned}$$


where $v = \tau - u \iff \tau = v + u$

$$\begin{aligned}
S_{yy}(s) &= \hat{h}(s) S_{xy}(s) \\
&= \hat{h}(s) \hat{h}^*(-s^*) S_{xx}(s)
\end{aligned}$$

$$\begin{aligned}
\tilde{S}_{xy}(\omega) &= S_{xy}(s) \Big|_{s=j\omega} \\
&= \hat{h}^*(-s^*) S_{xx}(s) \Big|_{s=j\omega} \\
&= \left(\int_{u \in \mathbb{R}} h(u) e^{-(s^*)u} du \right)^* \int_v R_{xx}(v) e^{-sv} dv \Big|_{s=j\omega} \\
&= \left(\int_{u \in \mathbb{R}} h(u) e^{(-j\omega)^*u} du \right)^* \int_v R_{xx}(v) e^{-j\omega v} dv \\
&= \left(\int_{u \in \mathbb{R}} h(u) e^{-j\omega u} du \right)^* \int_v R_{xx}(v) e^{-j\omega v} dv \\
&\triangleq \tilde{h}^*(\omega) \tilde{S}_{xx}(\omega)
\end{aligned}$$

$$\begin{aligned}
\tilde{S}_{yy}(\omega) &= S_{yy}(s) \Big|_{s=j\omega} \\
&= \hat{h}(s) S_{xy}(s) \Big|_{s=j\omega} \\
&= \int_{u \in \mathbb{R}} h(u) e^{-su} du \int_{\tau \in \mathbb{R}} R_{xy}(v) e^{-sv} d\tau \Big|_{s=j\omega} \\
&= \int_{u \in \mathbb{R}} h(u) e^{-j\omega u} du \int_{\tau \in \mathbb{R}} R_{xy}(v) e^{-j\omega v} d\tau \\
&= \tilde{h}(\omega) \tilde{S}_{xy}(\omega)
\end{aligned}$$

$$\begin{aligned}\tilde{S}_{yy}(\omega) &= \tilde{h}(\omega)\tilde{S}_{xy}(\omega) \\ &= \tilde{h}(\omega)\tilde{h}^*(\omega)\tilde{S}_{xx}(\omega) \\ &= |\tilde{h}(\omega)|^2\tilde{S}_{xx}(\omega)\end{aligned}$$

Figure 6.3: auto-correlation $R_{xx}(\tau)$

Theorem 6.4. Let $x : \mathbb{R} \rightarrow \mathbb{C}$ be a WSS random process with auto-correlation $R_{xx}(\tau)$. Then $R_{xx}(\tau)$ is conjugate symmetric such that (see Figure 6.3 page 48)

T H M	$R_{xx}(\tau) = R_{xx}^*(-\tau)$ (CONJUGATE SYMMETRIC) $\mathbf{R}_e [R_{xx}(\tau)] = \mathbf{R}_e [R_{xx}^*(-\tau)]$ (SYMMETRIC) $\mathbf{I}_m [R_{xx}(\tau)] = -\mathbf{I}_m [R_{xx}^*(-\tau)]$ (ANTI-SYMMETRIC) $ R_{xx}(\tau) = R_{xx}^*(-\tau) $ (SYMMETRIC) $\angle R_{xx}(\tau) = \angle R_{xx}^*(-\tau)$ (ANTI-SYMMETRIC).
-------------	--

PROOF:

$$\begin{aligned}R_{xx}^*(\tau) &\triangleq (\mathbb{E}[x(t-\tau)x^*(t)])^* \\ &= \mathbb{E}[x^*(t-\tau)x(t)] \\ &= \mathbb{E}[x(t)x^*(t-\tau)] \\ &= \mathbb{E}[x(u+\tau)x^*(u)] \\ &\triangleq R_{xx}(\tau) \quad \text{where } u \triangleq t-\tau \iff t=u+\tau \\ \mathbf{R}_e [R_{xx}(\tau)] &= \mathbf{R}_e [R_{xx}^*(-\tau)] \\ \mathbf{I}_m [R_{xx}(\tau)] &= \mathbf{I}_m [R_{xx}^*(-\tau)] \\ abs R_{xx}(\tau) &= |R_{xx}^*(-\tau)| \\ \angle R_{xx}(\tau) &= \angle R_{xx}^*(-\tau) \\ &= \mathbf{R}_e [R_{xx}(-\tau)] \\ &= -\mathbf{I}_m [R_{xx}(-\tau)] \\ &= |R_{xx}(-\tau)| \\ &= -\angle R_{xx}(-\tau)\end{aligned}$$

6.3 Whitening continuous random sequences

Simple algebraic operations on white noise processes (processes with autocorrelation $R_{xx}(\tau) = \delta(\tau)$) often produce *colored* noise (processes with autocorrelation $R_{xx}(\tau) \neq \delta(\tau)$). Sometimes we would like to process a colored noise process to produce a white noise process. This operation is known as *whitening*. Reasons for why we may want to whiten a noise process include

1. Samples from a white noise process are uncorrelated. If the noise process is Gaussian, then these samples are also independent which often greatly simplifies analysis.

2. Any orthonormal basis can be used to decompose a white noise process. This is not true of a colored noise process. Karhunen–Loëve expansion can be used to decompose colored noise.⁵

Definition 6.6. A **rational expression** $p(s)$ is a polynomial divided by a polynomial such that

D E F

$$p(s) = \frac{\sum_{n=0}^N b_n s^n}{\sum_{n=0}^M a_n s^n}.$$

The **zeros** of a rational expression are the roots of its numerator polynomial.

The **poles** of a rational expression are the roots of its denominator polynomial.

Definition 6.7. Let $\check{h}(s)$ be the Laplace transform of the impulse response of a filter. If $\check{h}(s)$ can be expressed as a rational expression with poles and zeros at $a_n + ib_n$, then the filter is **minimum phase** if each $a_n < 0$ (all roots lie in the left hand side of the complex s -plane).

Note that if $L(s)$ has a root at $s = -a + ib$, then $L^*(-s^*)$ has a root at

$$-s^* = -(-a + ib)^* = -(-a - ib) = a + ib.$$

That is, if $L(s)$ has a root in the left hand plane, then $L^*(-s^*)$ has a root directly opposite across the imaginary axis in the right hand plane (see Figure 6.4 page 49). A causal stable filter $\hat{h}(s)$ must have all of its poles in the left hand plane. A minimum phase filter is a filter with both its poles and zeros in the left hand plane. One advantage of a minimum phase filter is that its reciprocal (zeros become poles and poles become zeros) is also causal and stable.




Figure 6.4: Mirrored roots in complex-s plane




Figure 6.5: Innovations and whitening filters

The next theorem demonstrates a method for “whitening” a random process $x(t)$ with a filter constructed from a decomposition of $R_{xx}(\tau)$. The technique is stated precisely in Theorem 6.5 page 50

⁵ Karhunen–Loëve expansion: Section 3.3 page 20

and illustrated in Figure 6.5 page 49. Both imply two filters with impulse responses $l(t)$ and $\gamma(t)$. Filter $l(t)$ is referred to as the **innovations filter** (because it generates or “innovates” $x(t)$ from a white noise process $w(t)$) and $\gamma(t)$ is referred to as the **whitening filter** because it produces a white noise sequence when the input sequence is $x(t)$.⁶

Theorem 6.5. Let $x(t)$ be a WSS random process with autocorrelation $R_{xx}(\tau)$ and spectral density $S_{xx}(s)$. If $S_{xx}(s)$ has a **rational expression**, then the following are true:

1. There exists a rational expression $L(s)$ with minimum phase such that

$$S_{xx}(s) = L(s)L^*(-s^*).$$

2. An LTI filter for which the Laplace transform of the impulse response $\gamma(t)$ is

$$\Gamma(s) = \frac{1}{L(s)}$$

is both causal and stable.

3. If $x(t)$ is the input to the filter $\gamma(t)$, the output $y(t)$ is a **white noise sequence** such that

$$S_{yy}(s) = 1 \quad R_{yy}(\tau) = \delta(\tau).$$

PROOF:

$$\begin{aligned} S_{ww}(s) &= \Gamma(s)\Gamma^*(-s^*)S_{xx}(s) \\ &= \frac{1}{L(s)} \frac{1}{L^*(-s^*)} S_{xx}(s) \\ &= \frac{1}{L(s)} \frac{1}{L^*(-s^*)} L(s)L^*(-s^*) \\ &= 1 \end{aligned}$$




⁶ Papoulis (1991), pages 401–402

CHAPTER 7

ADDITIONAL NOISE ON RANDOM SEQUENCES

7.1 Additive noise and correlation

Theorem 7.1. Let S be the system illustrated to the right, where T is NOT NECESSARILY LINEAR.



T H M	(A). $x(n)$ is WSS	and	(1). $R_{yy}(m) = R_{vv}(m)$	and
	(B). $x(n)$ and $v(n)$ are uncorrelated	and	(2). $R_{xy}(m) = R_{xx}(m)$	and
	(C). $v(n)$ is zero-mean	(3). $R_{yy}(m) = R_{xx}(m) + R_{vv}(m)$	and	

PROOF:

$$\begin{aligned}
 R_{yy}(m) &\triangleq E[y(m)y^*(0)] && \text{by (A) and definition of } R_{yy} && (\text{Definition 2.4 page 12}) \\
 &\triangleq E[(x(m) + v(m))v^*(0)] && \text{by definition of } y \\
 &= E[x(m)v^*(0)] + E[v(m)v^*(0)] && \text{by linearity of } E && (\text{Theorem 1.1 page 4}) \\
 &= Ex(m)Ev^*(0) + E[v(m)v^*(0)] && \text{by uncorrelated hypothesis} && (\text{B}) \\
 &= Ex(m)Ev^*(0) + E[v(m)v^*(0)] && \text{by zero-mean hypothesis} && (\text{C}) \\
 &= R_{vv}(m) && \text{by definition of } R_{vv} && (\text{Definition 2.4 page 12}) \\
 R_{xy}(m) &\triangleq E[x(m)y^*(0)] && \text{by (A) and definition of } R_{xy} && (\text{Definition 2.4 page 12}) \\
 &\triangleq E(x(m)[x(0) + v(0)]^*) && \text{by definition of } y \\
 &= E[x(m)x^*(0)] + E[x(m)v^*(0)] && \text{by linearity of } E && (\text{Theorem 1.1 page 4}) \\
 &= E[x(m)x^*(0)] + E[x(m)]E[v^*(0)] && \text{by uncorrelated hypothesis} && (\text{B}) \\
 &= E[x(m)x^*(0)] + E[x(m)]E[v^*(0)] && \text{by zero-mean hypothesis} && (\text{C}) \\
 &= R_{xx}(m) && \text{by definition of } R_{xx} && (\text{Definition 2.4 page 12}) \\
 R_{yy}(m) &\triangleq E[y(m)y^*(0)] && \text{by (A) and definition of } R_{yy} && (\text{Definition 2.4 page 12}) \\
 &\triangleq E[(x(m) + v(m))(x(0) + v(0))^*] && \text{by definition of } y \\
 &= E[x(m)x^*(0)] + E[x(m)v^*(0)] + E[v(m)x^*(0)] + E[v(m)v^*(0)] && \\
 &= E[x(m)x^*(0)] + Ex(m)Ev^*(0) + Ev(m)Ex^*(0) + E[v(m)v^*(0)] && \text{by uncorrelated hypothesis (B)} \\
 &= E[x(m)x^*(0)] + Ex(m)Ev^*(0) + Ev(m)Ex^*(0) + E[v(m)v^*(0)] && \text{by zero-mean hypothesis (C)}
 \end{aligned}$$

$$\begin{aligned}
 &= R_{xx}(m) + R_{vv}(m) && \text{by definition of } R_{xx} \\
 R_{xx}(m) &\triangleq E[x(m)x^*(0)] \\
 &\triangleq E([y(m) - v(m)][y(0) - v(0)]^*) \\
 &= E[y(m)y^*(0)] - E[y(m)v^*(0)] - E[v(m)y^*(0)] + E[v(m)v^*(0)] \\
 &\triangleq R_{yy}(m) - R_{yv}(m) - R_{vy}(m) + R_{vv}(m) \\
 &= R_{yy}(m) + R_{vv}(m) - 2R_e R_{yv}(m)
 \end{aligned}$$

⇒

Remark 7.1. Because in Theorem 7.1 $y = x + v$ and $R_{yy} = R_{xx} + R_{vv}$, one might assume that R is a kind of *linear operator* (Definition M.3 page 282) and further assume that because $x = y - v$ and $R_{(-v)(-v)} = R_{vv}$, that $R_{xx} = R_{yy} + R_{vv}$. As Theorem 7.1 demonstrates, this is simply **not the case**. The problem here is that y and v are very much *correlated*—in fact y is obviously a *function* of v .

Corollary 7.1. Let S be the system illustrated in Theorem 7.1 (page 51).


C O R	<i>hypotheses of Theorem 7.1 (page 51)</i>	\Rightarrow	$ \left\{ \begin{array}{lcl} (1). & \check{S}_{yy}(z) &= \check{S}_{xx}(z) + \check{S}_{vv}(z) & \text{and} \\ (2). & \check{S}_{yv}(z) &= \check{S}_{vv}(z) & \text{and} \\ (3). & \check{S}_{yv}(z) &= \check{S}_{yy}(z) + \check{S}_{vv}(z) + \check{S}_{yv}(z) + \check{S}_{yv}^*(z^*) & \text{and} \\ (4). & \tilde{S}_{yy}(\omega) &= \tilde{S}_{xx}(\omega) + \tilde{S}_{vv}(\omega) & \text{and} \\ (5). & \tilde{S}_{yv}(\omega) &= \tilde{S}_{vv}(\omega) & \text{and} \\ (6). & \tilde{S}_{yv}(\omega) &= \tilde{S}_{yy}(\omega) + \tilde{S}_{vv}(\omega) + \tilde{S}_{yv}(\omega) + \tilde{S}_{yv}^*(-\omega) & \text{and} \end{array} \right. $
----------------------	--	---------------	---

PROOF:

$$\begin{aligned}
 \check{S}_{yy}(z) &\triangleq ZR_{yy}(m) && \text{by definition of } \check{S}_{yy} && (\text{Definition 2.5 page 14}) \\
 &= ZR_{qq}(m) + ZR_{vv}(m) && \text{by previous result} && (1) \\
 &= \check{S}_{qq}(z) + \check{S}_{vv}(z) && \text{by definition of } \check{S}_{yy} && (\text{Definition 6.3 page 44}) \\
 \tilde{S}_{yy}(\omega) &\triangleq \check{F}R_{yy}(m) && \text{by definition of } \tilde{S}_{yy} && (\text{Definition 6.3 page 44}) \\
 &= \check{F}R_{qq}(m) + \check{F}R_{vv}(m) && \text{by previous result} && (1) \\
 &= \tilde{S}_{qq}(\omega) + \tilde{S}_{vv}(\omega) && \text{by definition of } \tilde{S}_{yy} && (\text{Definition 6.3 page 44})
 \end{aligned}$$

⇒

Theorem 7.2. Let S be the system illustrated to the right:



T H M	$ \left\{ \begin{array}{ll} \text{(A). } x(n) \text{ and } r(n) \text{ are wide sense stationary} & \text{and} \\ \text{(B). } E[x(n)w(n)] = E[x(n)]E[w(n)] \text{ (uncorrelated)} & \text{and} \\ \text{(C). } E[r(n)v(n)] = E[r(n)]E[v(n)] \text{ (uncorrelated)} & \text{and} \\ \text{(D). } E[w(n)v(n)] = E[w(n)]E[v(n)] \text{ (uncorrelated)} & \text{and} \\ \text{(E). } E[v(n)] = E[w(n)] = 0 \text{ (zero-mean)} & \text{and} \end{array} \right\} \Rightarrow \left\{ \begin{array}{lcl} R_{sy}(m) &= R_{sx}(m) \\ &= R_{ry}(m) \\ &= R_{rx}(m) \end{array} \right\} $
----------------------	---



PROOF:

$$\begin{aligned}
 R_{sy}(m) &\triangleq E[s(m)y^*(0)] && \text{by definition of } R_{xy} && (\text{Definition 2.4 page 12}) \\
 &\triangleq E([r(m) + w(m)][x(0) + v(0)]^*) && \text{by definition of } S \\
 &= E[r(m)x^*(0)] + E[r(m)v^*(0)] + E[w(m)x^*(0)] + E[w(m)v^*(0)] \\
 &= E[r(m)x^*(0)] + Er(m)Ev^*(0) \\
 &\quad + Ew(m)Ex^*(0) + Ew(m)Ev^*(0) && \text{by uncorrelated hypotheses} && (\text{B), (C), and (D)}) \\
 &= E[r(m)x^*(0)] + Er(m)Ev^*(0) && \text{by zero-mean hypothesis} && (\text{E}) \\
 &= R_{rx}(m) && \text{by definition of } R_{rx} && (\text{Definition 2.4 page 12}) \\
 R_{sx}(m) &\triangleq E[s(m)x^*(0)] && \text{by definition of } R_{xy} && (\text{Definition 2.4 page 12}) \\
 &\triangleq E([r(m) + w(m)]x^*(0)) \\
 &= E[r(m)x^*(0)] + Ew(m)Ex^*(0) && \text{by uncorrelated hypothesis} && (\text{B}) \\
 &= E[r(m)x^*(0)] + Ew(m)Ex^*(0) && \text{by zero-mean hypothesis} && (\text{E}) \\
 &= R_{rx}(m) && \text{by definition of } R_{rx} && (\text{Definition 2.4 page 12}) \\
 R_{ry}(m) &\triangleq E[r(m)y^*(0)] && \text{by definition of } S && (\text{Definition 2.4 page 12}) \\
 &\triangleq E(r(m)[x(0) + v(0)]^*) \\
 &= E[r(m)x^*(0)] + Er(m)Ev^*(0) && \text{by uncorrelated hypothesis} && (\text{C}) \\
 &= E[r(m)x^*(0)] + Er(m)Ev^*(0) && \text{by zero-mean hypothesis} && (\text{E}) \\
 &= R_{rx}(m) && \text{by definition of } R_{rx} && (\text{Definition 2.4 page 12})
 \end{aligned}$$



Corollary 7.2. Let S be the system illustrated in Theorem 7.2 (page 52).

COR	$\left\{ \begin{array}{l} \text{hypotheses of} \\ \text{Theorem 7.2 (page 52)} \end{array} \right\} \Rightarrow \left\{ \begin{array}{l} (1). \check{S}_{sy}(z) = \check{S}_{sx}(z) = \check{S}_{ry}(z) = \check{S}_{rx}(z) \text{ and} \\ (2). \check{S}_{sy}(\omega) = \check{S}_{sx}(\omega) = \check{S}_{ry}(\omega) = \check{S}_{rx}(\omega) \end{array} \right\}$
-----	---

PROOF:

$$\begin{aligned}
 \check{S}_{sy}(\omega) &\triangleq ZR_{sy}(m) && \text{by definition of } \check{S}_{xy} && (\text{Definition 2.5 page 14}) \\
 &= ZR_{rx}(m) && \text{by previous result} && (1) \\
 &= \check{S}_{rx}(z) && \text{by definition of } \check{S}_{xy} && (\text{Definition 2.5 page 14}) \\
 \check{S}_{sy}(\omega) &\triangleq \check{F}R_{sy}(m) && \text{by definition of } \check{S}_{xy} && (\text{Definition 6.3 page 44}) \\
 &= \check{F}R_{rx}(m) && \text{by previous result} && (1) \\
 &= \check{S}_{rx}(\omega) && \text{by definition of } \check{S}_{xy} && (\text{Definition 6.3 page 44})
 \end{aligned}$$




7.2 Additive noise and operators

Lemma 7.1. Let S be the system illustrated in Figure 7.2 (page 55) (A).

LEM	$\left\{ \begin{array}{l} (A). R_{xx}(n_1, m) = R_{xx}(n_2, m) \text{ (WSS)} \\ (B). E[x(n)v(n)] = Ex(n)Ev(n) \text{ (UNCORRELATED)} \\ (E). Ev(n) = 0 \text{ (ZERO-MEAN)} \end{array} \right\} \Rightarrow \left\{ \begin{array}{l} (1). R_{xy}(m) = R_{xq}(m) \text{ and} \\ (2). \check{S}_{xy}(z) = \check{S}_{xq}(z) \text{ and} \\ (3). \check{S}_{xy}(\omega) = \check{S}_{xq}(\omega) \end{array} \right\}$
-----	---



Figure 7.1: Additive noise with *linear/non-linear* operator **T**

PROOF:

$$\begin{aligned}
 R_{xy}(m) &\triangleq E[x(m)y^*(0)] && \text{by definition of } R_{xy} && (\text{Definition 2.4 page 12}) \\
 &\triangleq E[x(m)(q(0) + v(0))^*] && \text{by definition of } S && (\text{Figure 7.2 page 55}) (A) \\
 &= E[x(m)q^*(0) + p(m)v^*(0)] \\
 &= E[x(m)q^*(0)] + E[x(m)v^*(0)] \\
 &= E[x(m)q^*(0)] + [Ex(m)][Ev^*(0)] && \text{by uncorrelated hypothesis} && (B) \\
 &= E[x(m)q^*(0)] + [Ep(m)][Ev^*(0)]^0 && \text{by zero-mean hypothesis} && (E) \\
 &= E[x(m)q^*(0)] && \text{by definition of } R_{xq} && (\text{Definition 2.4 page 12}) \\
 &= R_{xq}(m) \\
 \check{S}_{xy}(z) &\triangleq ZR_{xy}(m) && \text{by definition of } \check{S}_{xy} && (\text{Definition 2.5 page 14}) \\
 &= ZR_{xq}(m) && \text{by previous result} && (1) \\
 &= \check{S}_{xq}(z) && \text{by definition of } \check{S}_{xq} && (\text{Definition 2.5 page 14}) \\
 \check{S}_{xy}(\omega) &\triangleq \check{F}R_{xy}(m) && \text{by definition of } \check{S}_{xy} && (\text{Definition 6.3 page 44}) \\
 &= \check{F}R_{xq}(m) && \text{by previous result} && (1) \\
 &= \check{S}_{xq}(\omega) && \text{by definition of } \check{S}_{xq} && (\text{Definition 6.3 page 44})
 \end{aligned}$$

Lemma 7.2. Let **S** be the system illustrated in Figure 7.2 (page 55) (B).

L E M	$\left\{ \begin{array}{ll} (A). & x(n) \text{ is (WSS)} \\ (B). & u(n) \text{ is (ZERO-MEAN) and} \\ (C). & x(n) \text{ and } u(n) \text{ are (UNCORRELATED)} \end{array} \right.$	$\Rightarrow \left\{ \begin{array}{ll} (1). & R_{pq}(m) = R_{xy}(m) + R_{uy}(m) \text{ and} \\ (2). & \check{S}_{pq}(z) = \check{S}_{xy}(z) + \check{S}_{uy}(z) \text{ and} \\ (3). & \check{S}_{pq}(\omega) = \check{S}_{xy}(\omega) + \check{S}_{uy}(\omega) \end{array} \right.$
----------------------	--	---

PROOF:

$$\begin{aligned}
 R_{xy}(m) &\triangleq E[x(m)y^*(0)] && \text{by definition of } R_{xy} && (\text{Definition 2.4 page 12}) \\
 &\triangleq E([p(m) - u(m)]y^*(0)) && \text{by definition of } S \\
 &= E[p(m)y^*(0) - u(m)y^*(0)] \\
 &= E[p(m)y^*(0)] - E[u(m)y^*(0)] && \text{because } E \text{ is a linear operator} && (\text{Theorem 1.1 page 4}) \\
 &\triangleq R_{py}(m) - R_{uy}(m) && \text{by definition of } R_{xy} && (\text{Definition 2.4 page 12})
 \end{aligned}$$

Lemma 7.3. Let **S** be the system illustrated in Figure 7.2 (page 55) (C).

L E M	$\left\{ \begin{array}{ll} (A). & x(n) \text{ is (WSS)} \\ (B). & u(n) \text{ is (ZERO-MEAN) and} \\ (C). & x(n) \text{ and } u(n) \text{ are (UNCORRELATED)} \end{array} \right.$	$\Rightarrow \left\{ \begin{array}{ll} (1). & R_{xy}(m) = R_{py}(m) \text{ and} \\ (2). & \check{S}_{xy}(z) = \check{S}_{py}(z) \text{ and} \\ (3). & \check{S}_{xy}(\omega) = \check{S}_{py}(\omega) \end{array} \right.$
----------------------	--	--



PROOF:

$$\begin{aligned}
 R_{xy}(m) &\triangleq E[x(m)y^*(0)] && \text{by definition of } R_{py} \\
 &\triangleq E[p(m) + u(m)]y^*(0) && \text{by definition of } S \\
 &= E[p(m)y^*(0) + u(m)y^*(0)] && \text{by field properties of } (\mathbb{R}, +, \cdot, 0, 1) \\
 &= E[p(m)y^*(0)] + E[u(m)y^*(0)] && \text{because } E \text{ is a linear operator} \quad (\text{Theorem 1.1 page 4}) \\
 &= E[p(m)y^*(0)] + E[u(m)]E[y^*(0)] && \text{by uncorrelated hypothesis} \quad (C) \\
 &= E[p(m)y^*(0)] + E[u(m)]E[y^*(0)]^0 && \text{by zero-mean hypothesis} \quad (B) \\
 &\triangleq R_{py}(m) && \text{by definition of } R_{xy} \quad (\text{Definition 2.4 page 12})
 \end{aligned}$$




Figure 7.2: linear / non-linear additive noise systems

Theorem 7.3 (communications additive noise cross-correlation).

Let S be the system illustrated in Figure 7.2 page 55 (A).

T H M	(A). $x(n)$ is WSS (B). $u(n)$ is ZERO-MEAN and (C). $v(n)$ is ZERO-MEAN and (D). $x(n), u(n), v(n)$ are UNCORRELATED	$\left\{ \begin{array}{l} (1). R_{pq}(m) = R_{py}(m) = R_{xy}(m) + R_{uy}(m) \text{ and} \\ (2). \tilde{S}_{pq}(z) = \tilde{S}_{py}(z) = \tilde{S}_{xy}(z) + \tilde{S}_{uy}(z) \text{ and} \\ (3). \tilde{S}_{pq}(\omega) = \tilde{S}_{py}(\omega) = \tilde{S}_{xy}(\omega) + \tilde{S}_{uy}(\omega) \end{array} \right.$
-------------	--	---

PROOF:

$$\begin{aligned}
 R_{pq}(m) &= R_{py}(m) && \text{by Lemma 7.1 page 53} \\
 R_{pq}(m) &= R_{xq}(m) + R_{uq}(m) && \text{by Lemma 7.2 page 54} \\
 R_{py}(m) &\triangleq E[p(m)y^*(0)] && \text{by definition } R_{py} \quad (\text{Definition 2.4 page 12}) \\
 &\triangleq E[(x(m) + u(m))y^*(0)] && \text{by definition } S \quad (\text{Figure 7.2 page 55}) \text{ (A)} \\
 &= E[x(m)y^*(0) + u(m)y^*(0)] && \text{by linearity of } E \quad (\text{Theorem 1.1 page 4}) \\
 &= E[x(m)y^*(0)] + E[u(m)y^*(0)] && \text{by definitions } R_{xy} \text{ and } R_{uy} \quad (\text{Definition 2.4 page 12})
 \end{aligned}$$

Theorem 7.4 (measurement additive noise cross-correlation).

Let S be the system illustrated in Figure 7.2 page 55 (B).

T H M	(A). $x(n)$ is WSS (B). $u(n)$ is ZERO-MEAN and (C). $v(n)$ is ZERO-MEAN and (D). $x(n), u(n), v(n)$ are UNCORRELATED	$\left\{ \begin{array}{l} (1). R_{pq}(m) = R_{py}(m) = R_{xq}(m) = R_{xy}(m) \text{ and} \\ (2). \tilde{S}_{pq}(z) = \tilde{S}_{py}(z) = \tilde{S}_{xq}(z) = \tilde{S}_{xy}(z) \text{ and} \\ (3). \tilde{S}_{pq}(\omega) = \tilde{S}_{py}(\omega) = \tilde{S}_{xq}(\omega) = \tilde{S}_{xy}(\omega) \end{array} \right.$
-------------	--	---

PROOF:

$$\begin{aligned}
 R_{pq}(m) &= R_{py}(m) && \text{by Lemma 7.1 page 53} \\
 R_{pq}(m) &= R_{xq}(m) && \text{by Lemma 7.3 page 54} \\
 R_{xy}(m) &\triangleq E[x(m)y^*(0)] && \text{by definition } R_{xy} \\
 &\triangleq E([p(m) + u(m)]y^*(0)) && \text{(Definition 2.4 page 12)} \\
 &= E[p(m)y^*(0) + u(m)y^*(0)] && \text{by definition S} \\
 &= E[p(m)y^*(0)] + E[u(m)y^*(0)] && \text{(Figure 7.2 page 55) (B)} \\
 &= E[p(m)y^*(0)] + E[\cancel{u(m)y^*(0)}] \xrightarrow{0} && \text{by linearity of E} \\
 &= E[p(m)y^*(0)] && \text{by uncorrelated hypothesis} \\
 &= R_{py}(m) && \text{(Theorem 1.1 page 4)} \\
 & && \text{(D)} \\
 & && \text{by definition of } R_{py} \\
 & && \text{(Definition 2.4 page 12)}
 \end{aligned}$$

⇒

7.3 Additive noise and LTI operators




Figure 7.3: Additive noise with LTI operator \mathbf{H}

Corollary 7.3. Let \mathbf{S} be the system illustrated in Figure 7.3 (page 56) (A).

C O R
$$\left\{ \begin{array}{lll} (A). & x(n) \text{ is} & (\text{WSS}) \\ (B). & u(n) \text{ is} & (\text{ZERO-MEAN}) \\ (C). & x(n) \text{ and } u(n) \text{ are} & (\text{UNCORRELATED}) \\ (D). & \mathbf{H} \text{ is} & (\text{LTI}) \end{array} \right. \text{ and } \Rightarrow \left\{ \begin{array}{l} \tilde{S}_{xy}(\omega) = \tilde{H}^*(\omega)\tilde{S}_{xx}(\omega) \end{array} \right\}$$

PROOF:

$$\begin{aligned}
 \tilde{S}_{xy}(\omega) &= \tilde{S}_{xq}(\omega) && \text{by Lemma 7.1 page 53} \\
 &= \tilde{H}^*(\omega)\tilde{S}_{xx}(\omega) && \text{by Corollary 5.3 page 37}
 \end{aligned}$$

⇒

Theorem 7.5. Let \mathbf{S} be the system illustrated in Figure 7.3 (page 56) (B).

T H M
$$\left\{ \begin{array}{lll} (A). & x(n) \text{ is} & (\text{WSS}) \\ (B). & u(n) \text{ is} & (\text{ZERO-MEAN}) \\ (C). & x(n) \text{ and } u(n) \text{ are} & (\text{UNCORRELATED}) \\ (D). & \mathbf{H} \text{ is} & (\text{LTI}) \end{array} \right. \text{ and } \Rightarrow \left\{ \begin{array}{lll} (1). & R_{yx}(m) = h(m) \star R_{xx}(m) & \text{and} \\ (2). & \tilde{S}_{yx}(z) = \check{h}(z)\check{S}_{xx}(z) & \text{and} \\ (3). & \tilde{S}_{xy}(\omega) = \tilde{H}^*(\omega)\tilde{S}_{xx}(\omega) & \text{and} \end{array} \right\}$$

PROOF:

- definition: Let $(h(n))$ be the *impulse response* of operator \mathbf{H} such that

$$\mathbf{H}\delta(n) \triangleq \sum_{m \in \mathbb{Z}} h(m)\delta(n-m)$$

2. lemma: $\mathbf{H}\mathbf{x}(n) = \sum_{m \in \mathbb{Z}} h(n)\mathbf{x}(m-n) = h(n) \star R_{xx}(n)$.

Proof: by the *linear time-invariant* hypotheses (D) and definition of *convolution* operator \star (Definition N.3 page 312)

3. Proof that $R_{yx}(m) = h(m) \star R_{xx}(m)$:

$$\begin{aligned}
 R_{yx}(m) &\triangleq E[y(m)x^*(0)] && \text{by definition of } R_{py} && (\text{Definition 2.4 page 12}) \\
 &= E([\mathbf{H}\mathbf{x}(m) + \mathbf{H}\mathbf{u}(m)]x^*(0)) && \text{by linear hypothesis} && (\text{D}) \\
 &= E([\mathbf{H}\mathbf{x}^*(m)]x^*(0) + [\mathbf{H}\mathbf{u}(0)]x^*(0)) \\
 &= E([\mathbf{H}\mathbf{x}^*(m)]x^*(0)) + E([\mathbf{H}\mathbf{u}(0)]x^*(0)) && \text{by linearity of } E && (\text{Theorem ?? page ??}) \\
 &= \mathbf{H}E[x(m)x^*(0)] + \mathbf{H}E[u(m)x^*(0)] && \text{by LTI hypotheses} && (\text{D}) \\
 &= \mathbf{H}E[x(m)x^*(0)] + \mathbf{H}E\mathbf{u}(m)Ex^*(0) && \text{by uncorrelated hypothesis} && (\text{C}) \\
 &= \mathbf{H}E[x(m)x^*(0)] + \mathbf{H}E\mathbf{u}(m)Ex^*(0) && \text{by zero-mean hypothesis} && (\text{B}) \\
 &= \mathbf{H}R_{xx}(m) && \text{by definition of } R_{xx} && (\text{Definition 2.4 page 12}) \\
 &= h(m) \star R_{xx}(m) && \text{by (2) lemma} &&
 \end{aligned}$$



When \mathbf{H} is *LTI*, what effect does the additive uncorrelated noise sources have on the cross-statistical properties of x and y ? Corollary 7.5 (next) demonstrates that, amazingly, under very general conditions, the noise sources have **no effect**.

Corollary 7.4. Let \mathbf{S} be the system illustrated in Figure 7.3 (page 56) (C).

COR	$ \left\{ \begin{array}{ll} \text{(A). } \mathbf{x}(n) \text{ is} & (\text{WSS}) \\ \text{(B). } \mathbf{u}(n) \text{ is} & (\text{ZERO-MEAN}) \\ \text{(C). } \mathbf{x}(n) \text{ and } \mathbf{w}(n) \text{ are} & (\text{UNCORRELATED}) \\ \text{(D). } \mathbf{H} \text{ is} & (\text{LTI}) \end{array} \right. \text{ and and and} $	$\Rightarrow \{ \tilde{S}_{xy}(\omega) = \tilde{H}^*(\omega)\tilde{S}_{pp}(\omega) \}$
-----	---	--

PROOF:

$$\begin{aligned}
 \tilde{S}_{xy}(\omega) &= \tilde{S}_{py}(\omega) && \text{by Lemma 7.3 page 54} \\
 &= \tilde{H}^*(\omega)\tilde{S}_{pp}(\omega) && \text{by Corollary 5.3 page 37}
 \end{aligned}$$





Figure 7.4:

Corollary 7.5. Let \mathbf{S} be the system illustrated in Figure 7.4 page 57 (A).

COR	$ \left\{ \begin{array}{ll} \text{(A). } \mathbf{x}(n) \text{ is} & (\text{WSS}) \\ \text{(B). } \mathbf{u}(n) \text{ is} & (\text{ZERO-MEAN}) \\ \text{(C). } \mathbf{v}(n) \text{ is} & (\text{ZERO-MEAN}) \\ \text{(D). } \mathbf{x}(n), \mathbf{u}(n), \mathbf{v}(n) \text{ are} & (\text{UNCORRELATED}) \\ \text{(E). } \mathbf{H} \text{ is} & (\text{LTI}) \end{array} \right. \text{ and and and and} $	$\Rightarrow \{ \tilde{S}_{xy}(\omega) = \tilde{H}^*(\omega)\tilde{S}_{xx}(\omega) \}$
-----	---	--



PROOF:

$$\begin{aligned}\tilde{S}_{yx}(\omega) &= \tilde{S}_{qx}(\omega) && \text{by Lemma 7.1 page 53} \\ &= \tilde{H}(\omega)\tilde{S}_{xx}(\omega) && \text{by Corollary 5.3 page 37}\end{aligned}$$


Corollary 7.6. Let S be the system illustrated in Figure 7.4 page 57 (B).

COR	$\left\{ \begin{array}{lll} (A). & \mathbf{x}(n) \text{ is} & \text{WSS} \\ (B). & \mathbf{w}(n) \text{ is} & \text{ZERO-MEAN} \\ (C). & \mathbf{v}(n) \text{ is} & \text{ZERO-MEAN} \\ (D). & \mathbf{x}(n), \mathbf{w}(n), \mathbf{v}(n) \text{ are} & \text{UNCORRELATED} \\ (E). & \mathbf{H} \text{ is} & \text{LTI} \end{array} \right. \text{ and } \right\} \Rightarrow \left\{ \begin{array}{l} \tilde{S}_{xy}(\omega) = \tilde{H}^*(\omega)\tilde{S}_{pp}(\omega) \end{array} \right\}$
-----	---


PROOF:

$$\begin{aligned}\tilde{S}_{yx}(\omega) &= \tilde{S}_{qx}(\omega) && \text{by Lemma 7.1 page 53} \\ &= \tilde{S}_{qp}(\omega) && \text{by Lemma 7.1 page 53} \\ &= \tilde{H}(\omega)\tilde{S}_{pp}(\omega) && \text{by Corollary 5.3 page 37}\end{aligned}$$

7.4 Additive noise and dual operators



(A) dual communications additive noise system
(Corollary 7.7 page 58)



(B) dual measurement additive noise system
(Corollary 7.8 page 59)

Figure 7.5: Dual Additive Noise Systems

Corollary 7.7. Let S be the system illustrated in Figure 7.5 (page 58) (A).

COR	$\left\{ \begin{array}{lll} (A). & \mathbf{H} \text{ is} & \text{LTI} \\ (B). & \mathbf{x}(n) \text{ is} & \text{WSS} \\ (C). & \mathbf{u} \text{ and } \mathbf{v} \text{ are} & \text{ZERO-MEAN} \\ (D). & \mathbf{x}, \mathbf{u}, \mathbf{v} \text{ are} & \text{UNCORRELATED} \end{array} \right. \text{ and } \right\} \Rightarrow \left\{ \begin{array}{l} (1). \quad \check{S}_{sy}(z) = \check{H}(z)[\check{S}_{xy}(z) + \check{S}_{uy}(z)] \quad \text{and} \\ (2). \quad \tilde{S}_{sy}(\omega) = \tilde{H}(\omega)[\tilde{S}_{xy}(\omega) + \tilde{S}_{uy}(\omega)] \end{array} \right\}$
-----	---

PROOF:

$$\begin{aligned}\check{S}_{sy}(z) &= \check{S}_{rq}(z) && \text{by Corollary 7.2 page 53} && \text{and (B), (C) and (D)} \\ &= \check{H}(z)\check{S}_{pq}(z) && \text{by Theorem 5.2 page 37} && \text{and (A)} \\ &= \check{H}(z)[\check{S}_{xq}(z) + \check{S}_{uq}(z)] && \text{by Lemma 7.2 page 54} \\ &= \check{H}(z)[\check{S}_{xy}(z) + \check{S}_{uy}(z)] && \text{by Lemma 7.1 page 53} \\ \tilde{S}_{sy}(\omega) &= \check{S}_{sy}(z)|_{z=e^{i\omega}} && && \\ &= \check{H}(z)[\check{S}_{xy}(z) + \check{S}_{uy}(z)]|_{z=e^{i\omega}} && \text{by previous result} && (1) \\ &= \tilde{H}(\omega)[\tilde{S}_{xy}(\omega) + \tilde{S}_{uy}(\omega)] && &&\end{aligned}$$

Corollary 7.8. Let \mathbf{S} be the system illustrated in Figure 7.5 (page 58) (B).

COR	$\left\{ \begin{array}{lll} (A). & \mathbf{H} \text{ is} & \text{LTI} \\ (B). & \mathbf{x}(n) \text{ is} & \text{WSS} \\ (C). & \mathbf{u} \text{ and } \mathbf{v} \text{ are} & \text{ZERO-MEAN} \\ (D). & \mathbf{p}, \mathbf{u}, \mathbf{v} \text{ are} & \text{UNCORRELATED} \end{array} \right. \text{ and } \left\{ \begin{array}{lll} (1). & \check{\mathbf{S}}_{sy}(z) & = \check{\mathbf{H}}(z)\check{\mathbf{S}}_{xy}(z) \\ (2). & \tilde{\mathbf{S}}_{sy}(\omega) & = \tilde{\mathbf{H}}(\omega)\tilde{\mathbf{S}}_{xy}(\omega) \end{array} \right. \right\} \Rightarrow \left\{ \begin{array}{lll} (1). & \check{\mathbf{S}}_{sy}(z) & = \check{\mathbf{H}}(z)\check{\mathbf{S}}_{xy}(z) \text{ and} \\ (2). & \tilde{\mathbf{S}}_{sy}(\omega) & = \tilde{\mathbf{H}}(\omega)\tilde{\mathbf{S}}_{xy}(\omega) \end{array} \right\}$
-----	--


PROOF:

$$\begin{aligned}
 \check{\mathbf{S}}_{sy}(z) &= \check{\mathbf{S}}_{rq}(z) && \text{by Corollary 7.2 page 53} && \text{and (B), (C) and (D)} \\
 &= \check{\mathbf{H}}(z)\check{\mathbf{S}}_{pq}(z) && \text{by Theorem 5.2 page 37} && \text{and (A)} \\
 &= \check{\mathbf{H}}(z)\check{\mathbf{S}}_{xq}(z) && \text{by Lemma 7.3 page 54} \\
 &= \check{\mathbf{H}}(z)\check{\mathbf{S}}_{xy}(z) && \text{by Lemma 7.1 page 53} \\
 \tilde{\mathbf{S}}_{sy}(\omega) &= \check{\mathbf{S}}_{sy}(z)|_{z=e^{j\omega}} && \text{by definition of } \mathbf{Z} && (\text{Definition P.4 page 330}) \\
 &= \check{\mathbf{H}}(z)\check{\mathbf{S}}_{xy}(z)|_{z=e^{j\omega}} && \text{by previous result} && (1) \\
 &= \tilde{\mathbf{H}}(\omega)\tilde{\mathbf{S}}_{xy}(\omega)
 \end{aligned}$$



Part III

Statistical Estimation



CHAPTER 8

ESTIMATION THEORY

Estimation types. Let $x(t; \theta)$ be a waveform with parameter θ . There are three basic types of estimation on s :

1. *detection*:

- ➊ The waveform $x(t; \theta_n)$ is known except for the value of parameter θ_n .
- ➋ The parameter θ_n is one of a finite set of values.
- ➌ Estimate θ_n and thereby also estimate $x(t; \theta)$.

2. *parametric estimation*:

- ➊ The waveform $x(t; \theta)$ is known except for the value of parameter θ .
- ➋ The parameter θ is one of an infinite set of values.
- ➌ Estimate θ and thereby also estimate $x(t; \theta)$.

3. *nonparametric estimation*:

- ➊ The waveform $x(t)$ is unknown.
- ➋ Estimate $x(t)$.

Estimation criterion. Optimization requires a criterion against which the quality of an estimate is measured.¹ The most demanding and general criterion is the *Bayesian* criterion. The Bayesian criterion requires knowledge of the probability distribution functions and the definition of a *cost function*. Other criterion are special cases of the Bayesian criterion such that the cost function is defined in a special way, no cost function is defined, and/or the distribution is not known (Figure 8.2 page 66).

Estimation techniques. Estimation techniques can be classified into five groups (Figure 8.2 page 66):²

1. sequential decoding
2. norm minimization

¹ Mandyam D. Srinath (1996) (013125295X).

² Nelles (2001) page 26 ("Fig 2.2 Overview of linear and nonlinear optimization techniques"), Nelles (2001) page 33 ("Fig 2.5 The Bayes method is the most general approach but..."), Nelles (2001) page 63 ("Table 3.3 Relationship between linear recursive and nonlinear optimization techniques"), Nelles (2001) page 66

3. gradient search
 4. inner product analysis
 5. direct search

Sequential decoding is a non-linear estimation family. Perhaps the most famous of these is the Viterbi algorithm which uses a trellis to calculate the estimate. The Viterbi algorithm has been shown to yield an optimal estimate in the maximal likelihood (ML) sense. Norm minimization and gradient search algorithms are all linear algorithms. While this restriction to linear operations often simplifies calculations, it often yields an estimate that is not optimal in the ML sense.

8.1 Estimation criterion




Figure 8.1: Estimation criterion

Definition 8.1. Let

- (A). $x(t; \theta)$ be a function with unknown parameter θ
 (B). $y(t)$ a known function which is statistically dependent on $x(t; \theta)$
 (C). $C(\theta, p(x, y))$ be a cost function.

Then the following estimates are defined as follows:



(1). <i>Bayesian estimate</i>		$\hat{\theta}_B$	$\triangleq \arg \min_{\theta} C(\theta, p(x, y))$
(2). <i>Mean square estimate</i>	(“ <i>MS estimate</i> ”)	$\hat{\theta}_{mms}$	$\triangleq \arg \min_{\theta} E \ C(\theta, p(x, y))\ ^2$
(3). <i>mini-max estimate</i>	(“ <i>MM estimate</i> ”)	$\hat{\theta}_{mm}$	$\triangleq \arg \min_{\theta} \max_p C(\theta, p(x, y))$
(4). <i>maximum a-posteriori probability estimate</i>	(“ <i>MAP estimate</i> ”)	$\hat{\theta}_{map}$	$\triangleq \arg \max_{\theta} P\{x(t; \theta) y(t)\}$
(5). <i>maximum likelihood estimate</i>	(“ <i>ML estimate</i> ”)	$\hat{\theta}_{ml}$	$\triangleq \arg \max_{\theta} P\{y(t) x(t; \theta)\}$

Theorem 8.1. Let $x(t; \theta)$ be a function with unknown parameter θ .

$$\{P\{\theta\} = \text{CONSTANT}\} \implies \{\hat{\theta}_{\text{map}} = \hat{\theta}_{\text{ml}}\}$$

PROOF:

$$\begin{aligned}
 \hat{\theta}_{\text{map}} &\triangleq \arg \max_{\theta} P\{s(t; \theta) | r(t)\} && \text{by definition of } \hat{\theta}_{\text{map}} && (\text{Definition 8.1 page 64}) \\
 &= \arg \max_{\theta} \frac{P\{s(t; \theta) \wedge y(t)\}}{P\{r(t)\}} \\
 &= \arg \max_{\theta} \frac{P\{r(t) | x(t; \theta)\} P\{s(t; \theta)\}}{P\{r(t)\}} \\
 &= \arg \max_{\theta} P\{r(t) | x(t; \theta)\} P\{s(t; \theta)\} \\
 &= \arg \max_{\theta} P\{r(t) | x(t; \theta)\} \\
 &\triangleq \hat{\theta}_{\text{ml}} && \text{by definition of } \hat{\theta}_{\text{ml}} && (\text{Definition 8.1 page 64})
 \end{aligned}$$



Definition 8.2.

D E F The **mean square error** $\text{mse}(\hat{\theta})$ of an estimate $\hat{\theta}$ of a parameter θ is defined as

$$\text{mse}(\hat{\theta}) \triangleq E[(\hat{\theta} - \theta)^2]$$

The *mean square error* of $\hat{\theta}$ can be expressed as the sum of two components: the variance of $\hat{\theta}$ and the bias of $\hat{\theta}$ squared (next Theorem).

Theorem 8.2.³

T H M

$$\text{mse}(\hat{\theta}) = \underbrace{E[(\hat{\theta} - E\hat{\theta})^2]}_{\text{variance of } \hat{\theta}} + \underbrace{[E\hat{\theta} - \theta]^2}_{\text{bias of } \hat{\theta} \text{ squared}}$$

PROOF:

$$\begin{aligned}
 \text{mse}(\hat{\theta}) &\triangleq E[(\hat{\theta} - \theta)^2] && \text{by definition of mse} && (\text{Definition 8.2 page 65}) \\
 &= E[(\hat{\theta} - E\hat{\theta} + E\hat{\theta} - \theta)^2] \\
 &= E\left[(\hat{\theta} - E\hat{\theta})^2 + \underbrace{(E\hat{\theta} - \theta)^2}_{\text{constant}} - 2(\hat{\theta} - E\hat{\theta})(E\hat{\theta} - \theta)\right] \\
 &= E(\hat{\theta} - E\hat{\theta})^2 + (E\hat{\theta} - \theta)^2 - 2E[\hat{\theta}E\hat{\theta} - \hat{\theta}\theta - E\hat{\theta}\hat{\theta} + E\hat{\theta}\theta] \\
 &= E(\hat{\theta} - E\hat{\theta})^2 + (E\hat{\theta} - \theta)^2 - 2[E\hat{\theta}E\hat{\theta} - E\hat{\theta}E\theta - E\hat{\theta}E\hat{\theta} + E\hat{\theta}E\theta] \\
 &= E(\hat{\theta} - E\hat{\theta})^2 + (E\hat{\theta} - \theta)^2 - 0
 \end{aligned}$$



8.2 Estimation techniques

8.2.1 Sequential decoding

It has been shown that the Viterbi algorithm (trellis) produces an optimal estimate in the maximal likelihood (ML) sense. A Verterbi trellis is shown in Figure 8.3 (page 66).

³ Kay (1988) page 45 (§“3.3 ESTIMATION THEORY”)




Figure 8.2: Estimation techniques




Figure 8.3: Viterbi algorithm trellis

8.2.2 Norm minimization

Norm minimization techniques are very powerful in that an optimum solution can be computed in one step without iteration or recursion. In this section we present two types of norm minimization:⁴

1. minimum mean square estimation (MMSE):

The MMS estimate is a *stochastic* estimate. To compute the MMS estimate, we do not need to know the actual data values, but we must know certain system statistics which are the input data autocorrelation and input/output crosscorrelation. The cost function is the expected value of the norm squared error.

2. least square estimation (LSE):

The LS estimate is a *deterministic* estimate. To compute the LS estimate, we must know the actual data values (although these may be “noisy” measurements). The cost function is the norm squared error.

Solutions to both are given in terms of two matrices:

Y : Autocorrelation matrix

W : Crosscorrelation matrix.

⁴The Least Squares algorithm is nothing new to mathematics. It was first developed in 1795 by Gauss who was also the first to discover the FFT.




Figure 8.4: Estimation using gradient of cost function

Minimum mean square estimation

Definition 8.3. Let the following vectors, matrices, and functions be defined as follows:

DEF	$x \in \mathbb{C}^m$	<i>data vector</i>
	$y \in \mathbb{C}^n$	<i>processed data vector</i>
	$\hat{y} \in \mathbb{C}^n$	<i>processed data estimate vector</i>
	$e \in \mathbb{C}^n$	<i>error vector</i>
	$p \in \mathbb{R}^m$	<i>parameter vector</i>
	$U \in \mathbb{C}_{mn}$	<i>regression matrix</i>
	$R \in \mathbb{C}_{mm}$	<i>autocorrelation matrix</i>
	$W \in \mathbb{C}^m$	<i>cross-correlation vector</i>
	$C : \mathbb{R}^m \rightarrow \mathbb{R}^+$	<i>cost function</i>

Theorem 8.3 (Minimum mean square estimation). Let

$$\begin{aligned}
 \hat{y}(p) &\triangleq U^H p \\
 e(p) &\triangleq \hat{y} - y \\
 C(p) &\triangleq E \|e\|^2 \triangleq E [e^H e] \\
 \hat{\theta}_{\text{mms}} &\triangleq \arg \min_p C(p) \\
 R &\triangleq E [UU^H] \\
 W &\triangleq E [Uy].
 \end{aligned}$$

Then

$$\begin{aligned}
 \hat{\theta}_{\text{mms}} &= (\mathbf{R}_e Y)^{-1} (\mathbf{R}_e W) \\
 C(p) &= p^H R p - (W^H p)^* - W^H p + E y^H y \\
 \nabla_p C(p) &= 2\mathbf{R}_e [Y] p - 2\mathbf{R}_e W \\
 C(\hat{\theta}_{\text{mms}}) &= (\mathbf{R}_e W^H) (\mathbf{R}_e Y)^{-1} R (\mathbf{R}_e Y)^{-1} (\mathbf{R}_e W) - 2(\mathbf{R}_e W^H) (\mathbf{R}_e Y)^{-1} (\mathbf{R}_e W) + E y^H y \\
 C(\hat{\theta}_{\text{mms}})|_{R \text{ real}} &= E y^H y - (\mathbf{R}_e W^H) R^{-1} (\mathbf{R}_e W).
 \end{aligned}$$

PROOF: See APPENDIX E (page 173) for a Matrix Calculus reference.

$$\begin{aligned}
C(p) &\triangleq E \|e\|^2 \\
&= E [e^H e] \\
&= E [(\hat{y} - y)^H (\hat{y} - y)] \\
&= E \left[(U^H p - y)^H (U^H p - y) \right] \\
&= E [(p^H U - y^H) (U^H p - y)] \\
&= E [p^H U U^H p - p^H U y - y^H U^H p + y^H y] \\
&= p^H E [U U^H] p - p^H E [U y] - E [y^H U^H] p + E y^H y \\
&= p^H E [U U^H] p - (E [U y]^H p)^H - E [U y]^H p + E y^H y \\
&= p^H R p - (W^H p)^H - W^H p + E y^H y \\
&= p^H R p - (W^H p)^* - W^H p + E y^H y \\
&= p^H R p - (W^H)^* p - W^H p + E y^H y \\
&= p^H R p - 2R_e [W^H] p + E y^H y
\end{aligned}$$

$$\begin{aligned}
\nabla_p C(p) &= \nabla_p [p^H R p - (W^H)^* p - W^H p + E y^H y] \\
&= R p + R^T p - [(W^H)^*]^T - [W^H]^T + 0 \\
&= R p + (R^H)^* p - W - W^* \\
&= R p + R^* p - W - W^* \\
&= (R + R^*) p - (W + W^*) \\
&= 2(R_e Y) p - 2R_e W
\end{aligned}$$

$$p_{\text{opt}} = (R_e Y)^{-1} (R_e W)$$

$$\begin{aligned}
C(p_{\text{opt}}) &= p_{\text{opt}}^H R p_{\text{opt}} - 2R_e [W^H] p_{\text{opt}} + E y^H y \\
&= [(R_e Y)^{-1} (R_e W)]^H R [(R_e Y)^{-1} (R_e W)] - 2R_e [W^H] [(R_e Y)^{-1} (R_e W)] + E y^H y \\
&= (R_e W^H) (R_e Y)^{-H} R (R_e Y)^{-1} (R_e W) - 2R_e [W^H] (R_e Y)^{-1} (R_e W) + E y^H y \\
&= (R_e W^H) (R_e R^H)^{-1} R (R_e Y)^{-1} (R_e W) - 2R_e [W^H] (R_e Y)^{-1} (R_e W) + E y^H y \\
&= (R_e W^H) (R_e Y)^{-1} R (R_e Y)^{-1} (R_e W) - 2(R_e W^H) (R_e Y)^{-1} (R_e W) + E y^H y
\end{aligned}$$

$$\begin{aligned}
C(p_{\text{opt}})|_{R \text{ real}} &= (R_e W^H) (R_e Y)^{-1} R (R_e Y)^{-1} (R_e W) - 2(R_e W^H) (R_e Y)^{-1} (R_e W) + E y^H y \\
&= (R_e W^H) R^{-1} R R^{-1} (R_e W) - 2(R_e W^H) R^{-1} (R_e W) + E y^H y \\
&= (R_e W^H) R^{-1} (R_e W) - 2(R_e W^H) R^{-1} (R_e W) + E y^H y \\
&= E y^H y - (R_e W^H) R^{-1} (R_e W)
\end{aligned}$$

⇒

In many adaptive filter and equalization applications, the autocorrelation matrix U is simply the m -element random data vector $x(k)$ at time k , as in the *Wiener-Hopf equations* (next).

Corollary 8.1 (Wiener-Hopf equations). ⁵

⁵ ↗ Ifeachor and Jervis (1993) pages 547–549 (§“9.3 Basic Wiener filter theory”), ↗ Ifeachor and Jervis (2002) pages 651–654 (§“10.3 Basic Wiener filter theory”)




Figure 8.5: Adaptive filter example

COR

$$\left\{ U \triangleq \mathbf{x}(k) \triangleq \begin{bmatrix} x(k) \\ x(k-1) \\ x(k-2) \\ \vdots \\ x(k-m+1) \end{bmatrix} \right\} \Rightarrow \left\{ \begin{array}{lcl} \hat{\theta}_{\text{mms}} & = & R^{-1}W \\ C(\hat{\theta}_{\text{mms}}) & = & W^T R^{-1} R R^{-1} W - 2W^T R^{-1} W + E y^T y \end{array} \right\}$$

PROOF: This is a special case of the more general case discussed in Theorem 8.3 (page 67). Here, the dimension of U is $m \times 1$ ($n=1$). As a result, y , \hat{y} , and e are simply scalar quantities (not vectors). In this special case, we have the following results (Figure 8.5 page 69):

$$\begin{aligned} \hat{y}(p) &\triangleq x^T p \\ e(p) &\triangleq \hat{y} - y \\ C(p) &\triangleq E \|e\|^2 \triangleq E [e^2] \\ \hat{\theta}_{\text{mms}} &\triangleq \arg \min_p C(p) \\ R &\triangleq E [xx^T] \\ W &\triangleq E [xy] \\ C(p) &= p^T Rp - 2W^T p + E [y^T y] \\ \nabla_p C(p) &= 2Rp - 2W \\ C(\hat{\theta}_{\text{mms}})|_{R \text{ real}} &= E y^T y - W^T R^{-1} W. \end{aligned}$$

Least squares

Theorem 8.4 (Least squares). Let

$$\begin{aligned} \hat{y}(p) &\triangleq U^H p \\ e(p) &\triangleq \hat{y} - y \\ C(p) &\triangleq \|e\|^2 \triangleq e^H e \end{aligned}$$

$$\begin{aligned}\hat{\theta}_{ls} &\triangleq \arg \min_p C(p) \\ R &\triangleq UU^H \\ W &\triangleq Uy.\end{aligned}$$

Then

THM	$\begin{aligned}\hat{\theta}_{ls} &= (\mathbf{R}_e Y)^{-1}(\mathbf{R}_e W) \\ C(p) &= p^H R p - (W^H p)^* - W^H p + E y^H y \\ \nabla_p C(p) &= 2\mathbf{R}_e [Y]p - 2\mathbf{R}_e W \\ C(\hat{\theta}_{ls}) &= (\mathbf{R}_e W^H)(\mathbf{R}_e Y)^{-1}R(\mathbf{R}_e Y)^{-1}(\mathbf{R}_e W) - 2(\mathbf{R}_e W^H)(\mathbf{R}_e Y)^{-1}(\mathbf{R}_e W) + E y^H y \\ C(\hat{\theta}_{ls}) _{R \text{ real}} &= E y^H y - (\mathbf{R}_e W^H)R^{-1}(\mathbf{R}_e W).\end{aligned}$
-----	---

PROOF: See APPENDIX E (page 173) for a Matrix Calculus reference.

$$\begin{aligned}C(p) &\triangleq \|e\|^2 \\ &= e^H e \\ &= (\hat{y} - y)^H (\hat{y} - y) \\ &= (U^H p - y)^H (U^H p - y) \\ &= (p^H U - y^H) (U^H p - y) \\ &= p^H UU^H p - p^H Uy - y^H U^H p + y^H y \\ &= p^H Rp - (W^H p)^H - W^H p + y^H y \\ &= p^H Rp - (W^H p)^* - W^H p + y^H y \\ &= p^H Rp - (W^H)^* p - W^H p + y^H y \\ &= p^H Rp - 2\mathbf{R}_e [W^H]p + y^H y\end{aligned}$$

$$\begin{aligned}\nabla_p C(p) &= \nabla_p [p^H Rp - (W^H)^* p - W^H p + y^H y] \\ &= Rp + R^T p - [(W^H)^*]^T - [W^H]^T + 0 \\ &= Rp + (R^H)^* p - W - W^* \\ &= Rp + R^* p - W - W^* \\ &= (R + R^*)p - (W + W^*) \\ &= 2(\mathbf{R}_e Y)p - 2\mathbf{R}_e W\end{aligned}$$

$$p_{opt} = (\mathbf{R}_e Y)^{-1}(\mathbf{R}_e W)$$

$$\begin{aligned}C(p_{opt}) &= p_{opt}^H Rp_{opt} - 2\mathbf{R}_e [W^H]p_{opt} + y^H y \\ &= [(\mathbf{R}_e Y)^{-1}(\mathbf{R}_e W)]^H R[(\mathbf{R}_e Y)^{-1}(\mathbf{R}_e W)] - 2\mathbf{R}_e [W^H][(\mathbf{R}_e Y)^{-1}(\mathbf{R}_e W)] + y^H y \\ &= (\mathbf{R}_e W^H)(\mathbf{R}_e Y)^{-H} R(\mathbf{R}_e Y)^{-1}(\mathbf{R}_e W) - 2\mathbf{R}_e [W^H](\mathbf{R}_e Y)^{-1}(\mathbf{R}_e W) + y^H y \\ &= (\mathbf{R}_e W^H)(\mathbf{R}_e R^H)^{-1} R(\mathbf{R}_e Y)^{-1}(\mathbf{R}_e W) - 2\mathbf{R}_e [W^H](\mathbf{R}_e Y)^{-1}(\mathbf{R}_e W) + y^H y \\ &= (\mathbf{R}_e W^H)(\mathbf{R}_e Y)^{-1} R(\mathbf{R}_e Y)^{-1}(\mathbf{R}_e W) - 2(\mathbf{R}_e W^H)(\mathbf{R}_e Y)^{-1}(\mathbf{R}_e W) + y^H y\end{aligned}$$

$$\begin{aligned}C(p_{opt})|_{R \text{ real}} &= (\mathbf{R}_e W^H)(\mathbf{R}_e Y)^{-1} R(\mathbf{R}_e Y)^{-1}(\mathbf{R}_e W) - 2(\mathbf{R}_e W^H)(\mathbf{R}_e Y)^{-1}(\mathbf{R}_e W) + y^H y \\ &= (\mathbf{R}_e W^H)R^{-1}RR^{-1}(\mathbf{R}_e W) - 2(\mathbf{R}_e W^H)R^{-1}(\mathbf{R}_e W) + y^H y\end{aligned}$$



$$\begin{aligned}
 &= (\mathbf{R}_e W^H) R^{-1} (\mathbf{R}_e W) - 2(\mathbf{R}_e W^H) R^{-1} (\mathbf{R}_e W) + y^H y \\
 &= y^H y - (\mathbf{R}_e W^H) R^{-1} (\mathbf{R}_e W)
 \end{aligned}$$

⇒

Example 8.1 (Polynomial approximation).

Suppose we **know** the locations $\{(x_n, y_n) | n = 1, 2, 3, 4, 5\}$ of 5 data points. Let x and y represent the locations of these points such that

$$x \triangleq \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \end{bmatrix} \quad y \triangleq \begin{bmatrix} y_1 \\ y_2 \\ y_3 \\ y_4 \\ y_5 \end{bmatrix}$$

Suppose we want to find a second order polynomial

$$cx^2 + bx + a$$

that best approximates these 5 points in the least squares sense. We define the matrix U (known) and vector $\hat{\theta}$ (to be computed) as follows:

$$U^H \triangleq \underbrace{\begin{bmatrix} 1 & x_1 & x_1^2 \\ 1 & x_2 & x_2^2 \\ 1 & x_3 & x_3^2 \\ 1 & x_4 & x_4^2 \\ 1 & x_5 & x_5^2 \end{bmatrix}}_{\text{Vandermonde matrix } ^6} \quad \hat{\theta} \triangleq \begin{bmatrix} a \\ b \\ c \end{bmatrix}$$

Then, using Theorem 8.4 (page 69), the best coefficients $\hat{\theta}$ for the polynomial are

$$\begin{aligned}
 \hat{\theta} &= \begin{bmatrix} a \\ b \\ c \end{bmatrix} \\
 &= R^{-1} W \\
 &= (U U^H)^{-1} (U y) \\
 &= \left(\begin{bmatrix} 1 & x_1 & x_1^2 \\ 1 & x_2 & x_2^2 \\ 1 & x_3 & x_3^2 \\ 1 & x_4 & x_4^2 \\ 1 & x_5 & x_5^2 \end{bmatrix} \begin{bmatrix} 1 & x_1 & x_1^2 \\ 1 & x_2 & x_2^2 \\ 1 & x_3 & x_3^2 \\ 1 & x_4 & x_4^2 \\ 1 & x_5 & x_5^2 \end{bmatrix}^H \right)^{-1} \left(\begin{bmatrix} 1 & x_1 & x_1^2 \\ 1 & x_2 & x_2^2 \\ 1 & x_3 & x_3^2 \\ 1 & x_4 & x_4^2 \\ 1 & x_5 & x_5^2 \end{bmatrix}^H \begin{bmatrix} y_1 \\ y_2 \\ y_3 \\ y_4 \\ y_5 \end{bmatrix} \right)
 \end{aligned}$$

8.2.3 Gradient search techniques

One of the biggest advantages of using a gradient search technique is that they can be implemented *recursively* as shown in the next equation. The general form of the gradient search parameter estimation techniques is⁷

T
H
M

$$p_n = p_{n-1} - \eta_{n-1} R [\nabla_p C(p_n)] \quad \text{where at time } n$$

⁶ Horn and Johnson (1990) 29

⁷ Nelles (2001), page 90

p_n	is the <i>state</i>	(vector)
η_n	is the <i>step size</i>	(scalar)
Y	is the <i>direction</i>	(matrix)
$\nabla_p C(p_n)$	is the <i>gradient</i> of the cost function $C(p_n)$	(vector)

Two major categories of gradient search techniques are

- ➊ steepest descent (includes LMS)
- ➋ Newton's method (includes RLS and Kalman filters).

The key difference between the two is that **steepest descent uses only first derivative information**, while **Newton's method uses both first and second derivative information** making it converge much faster but with significantly higher complexity.

First derivative techniques

Steepest descent. In this algorithm, $R = I$ (identity matrix). First derivative information is contained in ∇C . Second derivative information, if present, is contained in Y . Thus, steepest descent algorithms do not use second derivative information.

T
H
M

$$p_n = p_{n-1} - \eta_{n-1} [\nabla_p C(p_n)]$$

Least Mean Squares (LMS).⁸ This is a special case of *steepest descent*. In minimum mean square estimation (Section 8.2.2 page 67), the cost function $C(p)$ is defined as a *statistical average* of the error vector such that $C(p) = E [e^H e]$. In this case the gradient ∇C is difficult to compute. However, the LMS algorithm greatly simplifies the problem by instead defining the cost function as a function of the *instantaneous error* such that

$$\begin{aligned} y &= y(n) \\ \hat{y} &= \hat{y}(n) \\ C(p) &= \|e(n)\|^2 \\ &= e^2(n) \\ &= (\hat{y}(n) - y(n))^2 \end{aligned}$$

Computing the gradient of this cost function is then just a special case of *least squares estimation* (Section 8.2.2 page 69). Using LS, we let $U = x^T$ and hence

$$\begin{aligned} \nabla_p C(p) &= 2U^T U p - 2U^T y \\ &= 2x x^T p - 2x y \\ &= 2x \hat{y} - 2x y \\ &= 2x(\hat{y} - y) \\ &= 2x e(n) \end{aligned} \quad \begin{aligned} &\text{by Theorem 8.4 page 69} \\ &\text{by above definitions} \end{aligned}$$

The LMS algorithm uses this instantaneous gradient for ∇C , lets $R = I$, and uses a constant step

⁸ Manolakis et al. (2000), page 526



size η to give

T
H
M

$$p_n = p_{n-1} - 2\eta \mathbf{x}_n e(n)$$

Second derivative techniques

Newton's Method. This algorithm uses the *Hessian* matrix H , which is the second derivative of the cost function $C(p)$, and lets $R = H^{-1}$.

$$\begin{aligned} H_n &\triangleq \nabla_p \nabla_p C(p_n) \\ p_n &= p_{n-1} - \eta_{n-1} H_n^{-1} [\nabla_p C(p_n)] \end{aligned}$$

Kalman filtering ⁹

$$\begin{aligned} \gamma(k) &= \frac{1}{x^T(k)P(k-1)x(k) + 1} P(k-1)x(k) \\ P(k) &= (I - \gamma(k)x^T(k))P(k-1) + V \\ e(k) &= y(k) - x^T(k)\hat{p}(k-1) \\ \hat{p}(k) &= \hat{p}(k-1) + \gamma(k)e(k) \end{aligned}$$

Recursive Least Squares (RLS) ¹⁰ This algorithm is a special case of either the RLS with forgetting or the Kalman filter.

$$\begin{aligned} \gamma(k) &= \frac{1}{x^T(k)P(k-1)x(k) + 1} P(k-1)x(k) \\ P(k) &= (I - \gamma(k)x^T(k))P(k-1) \\ e(k) &= y(k) - x^T(k)\hat{p}(k-1) \\ \hat{p}(k) &= \hat{p}(k-1) + \gamma(k)e(k) \end{aligned}$$

8.2.4 Direct search

A direct search algorithm may be used in cases where the cost function over p has several local minima, making convergence difficult. Furthermore, direct search algorithms can be very computationally demanding.

⁹ Nelles (2001), page 66

¹⁰ Nelles (2001), page 66

CHAPTER 9

KL-EXPANSION APPLICATION

9.1 Sufficient statistics

Theorem 9.1 (page 75) (next) shows that the finite set $Y \triangleq \{\dot{y}_n | n = 1, 2, \dots, N\}$ provides just as much information as having the entire $\mathbf{y}(t)$ waveform (an uncountably infinite number of values) with respect to the following cases:

1. the conditional probability of $\mathbf{x}(t; \hat{\theta})$ given $\mathbf{y}(t)$
2. the *MAP estimate* of the information sequence
3. the *ML estimate* of the information sequence.

That is, even with a drastic reduction in the amount of information from uncountably infinite to finite N , no information is lost with respect to the quantities listed above.

This amazing result is very useful in practical system implementation and also for proving other theoretical results (notably estimation and detection theorems which come later in this chapter).

Theorem 9.1 (Sufficient statistic theorem). ¹ Let \mathbf{S} be an additive White Gaussian noise system and Ψ an orthonormal basis for $\mathbf{x}(t; \hat{\theta})$ such that

$$\begin{aligned}\mathbf{y}(t) &= \mathbf{x}(t; \hat{\theta}) + \mathbf{v}(t) \\ \Psi &= \{\psi_n | n = 1, 2, \dots, N\}\end{aligned}$$

Then $Y \triangleq \{\dot{y}_n | n = 1, 2, \dots, N\}$ is a **sufficient statistic** for $\mathbf{y}(t)$ such that...

T H M

$$\left\{ \mathbf{v}(t) \text{ is AWGN} \right\} \implies \left\{ \begin{array}{lcl} (1). & \mathbb{P} \{ \mathbf{x}(t; \hat{\theta}) | \mathbf{y}(t) \} & = \mathbb{P} \{ \mathbf{x}(t; \hat{\theta}) | Y \} \\ (2). & \hat{\theta}_{\text{map}} \triangleq \arg \max_{\hat{\theta}} \mathbb{P} \{ \mathbf{x}(t; \hat{\theta}) | \mathbf{y}(t) \} & = \arg \max_{\hat{\theta}} \mathbb{P} \{ \mathbf{x}(t; \hat{\theta}) | Y \} \\ (3). & \hat{\theta}_{\text{ml}} \triangleq \arg \max_{\hat{\theta}} \mathbb{P} \{ \mathbf{y}(t) | \mathbf{x}(t; \hat{\theta}) \} & = \arg \max_{\hat{\theta}} \mathbb{P} \{ Y | \mathbf{x}(t; \hat{\theta}) \} \end{array} \right\}$$

PROOF: Let $\mathbf{v}'(t) \triangleq \mathbf{v}(t) - \sum_{n=1}^N \dot{v}_n \psi_n(t)$.

¹  Fisher (1922) page 316 (“Criterion of Sufficiency”)

1. The relationship between $\mathbf{y}(t)$ and $\mathbf{v}'(t)$ is given by

$$\begin{aligned}
 \mathbf{y}(t) &= \sum_{n=1}^N \langle \mathbf{y}(t) | \psi_n(t) \rangle \psi_n(t) + \left[\mathbf{y}(t) - \sum_{n=1}^N \langle \mathbf{y}(t) | \psi_n(t) \rangle \psi_n(t) \right] \\
 &= \sum_{n=1}^N \langle \mathbf{y}(t) | \psi_n(t) \rangle \psi_n(t) + \left[\mathbf{y}(t) - \sum_{n=1}^N \langle \mathbf{x}(t) + \mathbf{v}(t) | \psi_n(t) \rangle \psi_n(t) \right] \\
 &= \sum_{n=1}^N \dot{y}_n \psi_n(t) + \left[\mathbf{x}(t) + \mathbf{v}(t) - \sum_{n=1}^N \langle \mathbf{x}(t) | \psi_n(t) \rangle \psi_n(t) - \sum_{n=1}^N \langle \mathbf{v}(t) | \psi_n(t) \rangle \psi_n(t) \right] \\
 &= \sum_{n=1}^N \dot{y}_n \psi_n(t) + \mathbf{x}(t) + \mathbf{v}(t) - \mathbf{x}(t) - [\mathbf{v}(t) - \mathbf{v}'(t)] \\
 &= \sum_{n=1}^N \dot{y}_n \psi_n(t) + \mathbf{v}'(t).
 \end{aligned}$$

2. Proof that the set of statistics \mathbf{Y} and the random process $\mathbf{v}'(t)$ are *uncorrelated*:

$$\begin{aligned}
 \mathbb{E}[\dot{y}_n \mathbf{v}'(t)] &= \mathbb{E}\left[\langle \mathbf{y}(t) | \psi_n(t) \rangle \left(\mathbf{v}(t) - \sum_{n=1}^N \langle \mathbf{v}(t) | \psi_n(t) \rangle \psi_n(t) \right)\right] \\
 &= \mathbb{E}\left[\langle \mathbf{x}(t) + \mathbf{v}(t) | \psi_n(t) \rangle \left(\mathbf{v}(t) - \sum_{n=1}^N \langle \mathbf{v}(t) | \psi_n(t) \rangle \psi_n(t) \right)\right] \\
 &= \mathbb{E}\left[\left(\langle \mathbf{x}(t) | \psi_n(t) \rangle + \langle \mathbf{v}(t) | \psi_n(t) \rangle\right) \left(\mathbf{v}(t) - \sum_{n=1}^N \langle \mathbf{v}(t) | \psi_n(t) \rangle \psi_n(t) \right)\right] \\
 &= \mathbb{E}\left[\left(\dot{x}_n + \dot{v}_n\right) \left(\mathbf{v}(t) - \sum_{n=1}^N \dot{v}_n \psi_n(t) \right)\right] \\
 &= \mathbb{E}\left[\dot{x}_n \mathbf{v}(t) - \dot{x}_n \sum_{n=1}^N \dot{v}_n \psi_n(t) + \dot{v}_n \mathbf{v}(t) - \dot{v}_n \sum_{n=1}^N \dot{v}_n \psi_n(t)\right] \\
 &= \mathbb{E}[\dot{x}_n \mathbf{v}(t)] - \mathbb{E}\left[\dot{x}_n \sum_{n=1}^N \dot{v}_n \psi_n(t)\right] + \mathbb{E}[\langle \mathbf{v}(t) | \psi_n(t) \rangle \mathbf{v}(t)] - \mathbb{E}\left[\sum_{m=1}^N \dot{v}_n \dot{v}_m \psi_m(t)\right] \\
 &= \cancel{\dot{x}_n} \mathbb{E}[\mathbf{v}(t)] - \cancel{\dot{x}_n} \sum_{n=1}^N \cancel{\mathbb{E}[\dot{v}_n]} \psi_n(t) + \mathbb{E}[\langle \mathbf{v}(t) \mathbf{v}(\hat{\theta}) | \psi_n(\hat{\theta}) \rangle] - \sum_{m=1}^N \mathbb{E}[\dot{v}_n \dot{v}_m] \psi_m(t) \\
 &= 0 - 0 + \langle \mathbb{E}[\mathbf{v}(t) n(\hat{\theta})] | \psi_n(\hat{\theta}) \rangle - \sum_{m=1}^N N_o \bar{\delta}_{mn} \psi_m(t) \quad (\text{because } \dot{v}_n \text{ is white}) \\
 &= \langle N_o \delta(t - \hat{\theta}) | \psi_n(\hat{\theta}) \rangle - N_o \psi_n(t) \\
 &= N_o \psi_n(t) - N_o \psi_n(t) \\
 &= 0
 \end{aligned}$$

3. This implies \dot{y}_n and $\mathbf{v}'(t)$ are uncorrelated. Since they are Gaussian processes (due to channel operator hypothesis), they are also independent.

4. Proof that $P\{\mathbf{x}(t; \hat{\theta}) | \mathbf{y}(t)\} = P\{\mathbf{x}(t; \hat{\theta}) | \dot{y}_1, \dot{y}_2, \dots, \dot{y}_N\}$:

$$P\{\mathbf{x}(t; \hat{\theta}) | \mathbf{y}(t)\} = P\left\{\mathbf{x}(t; \hat{\theta}) | \sum_{n=1}^N \dot{y}_n \psi_n(t) + \mathbf{v}'(t)\right\}$$



$$\begin{aligned}
&= P\{\mathbf{x}(t; \hat{\theta}) | R, \mathbf{v}'(t)\} && \text{because } Y \text{ and } \mathbf{v}'(t) \text{ can be extracted by } \langle \dots | \psi_n(t) \rangle \\
&= \frac{P\{R, \mathbf{v}'(t) | \mathbf{x}(t; \hat{\theta})\} P\{\mathbf{x}(t; \hat{\theta})\}}{P\{R, \mathbf{v}'(t)\}} \\
&= \frac{P\{R | \mathbf{x}(t; \hat{\theta})\} P\{\mathbf{v}'(t) | \mathbf{x}(t; \hat{\theta})\} P\{\mathbf{x}(t; \hat{\theta})\}}{P\{Y\} P\{\mathbf{v}'(t)\}} && \text{by independence of } Y \text{ and } \mathbf{v}'(t) \\
&= \frac{P\{R | \mathbf{x}(t; \hat{\theta})\} P\{\mathbf{v}'(t)\} P\{\mathbf{x}(t; \hat{\theta})\}}{P\{Y\} P\{\mathbf{v}'(t)\}} \\
&= \frac{P\{R | \mathbf{x}(t; \hat{\theta})\} P\{\mathbf{x}(t; \hat{\theta})\}}{P\{Y\}} \\
&= \frac{P\{R, \mathbf{x}(t; \hat{\theta})\}}{P\{Y\}} \\
&= P\{\mathbf{x}(t; \hat{\theta}) | Y\}
\end{aligned}$$

5. Proof that Y is a sufficient statistic for the *MAP estimate*:

$$\begin{aligned}
\hat{\theta}_{\text{map}} &\triangleq \arg \max_{\hat{\theta}} P\{\mathbf{x}(t; \hat{\theta}) | y(t)\} && \text{by definition of MAP estimate} \\
&= \arg \max_{\hat{\theta}} P\{\mathbf{x}(t; \hat{\theta}) | R\} && \text{by result 4.}
\end{aligned}$$

6. Proof that Y is a sufficient statistic for the *ML estimate*:

$$\begin{aligned}
\hat{\theta}_{\text{ml}} &\triangleq \arg \max_{\hat{\theta}} P\{\mathbf{y}(t) | \mathbf{x}(t; \hat{\theta})\} && \text{by definition of ML estimate} \\
&= \arg \max_{\hat{\theta}} P\left\{ \sum_{n=1}^N \dot{y}_n \psi_n(t) + \mathbf{v}'(t) | \mathbf{x}(t; \hat{\theta}) \right\} \\
&= \arg \max_{\hat{\theta}} P\{R, \mathbf{v}'(t) | \mathbf{x}(t; \hat{\theta})\} && \text{because } Y \text{ and } \mathbf{v}'(t) \text{ can be extracted by } \langle \dots | \psi_n(t) \rangle \\
&= \arg \max_{\hat{\theta}} P\{R | \mathbf{x}(t; \hat{\theta})\} P\{\mathbf{v}'(t) | \mathbf{x}(t; \hat{\theta})\} && \text{by independence of } Y \text{ and } \mathbf{v}'(t) \\
&= \arg \max_{\hat{\theta}} P\{R | \mathbf{x}(t; \hat{\theta})\} P\{\mathbf{v}'(t)\} && \text{by independence of } \mathbf{x}(t) \text{ and } \mathbf{v}'(t) \\
&= \arg \max_{\hat{\theta}} P\{R | \mathbf{x}(t; \hat{\theta})\} && \text{by independence of } \mathbf{v}'(t) \text{ and } \hat{\theta}
\end{aligned}$$

Theorem 9.2. Let $\mathbf{C} = \mathbf{C}_a$ be an additive noise channel.

THM	$\mathbf{C} = \mathbf{C}_a \implies \underbrace{\mathbf{C}}$ in additive noise channel $\implies \{ E(\dot{y}_n \theta) = \dot{x}_n(\theta) + E\dot{v}_n$
-----	---

PROOF:

$$\begin{aligned}
E(\dot{y}_n | \theta) &\triangleq (\langle \mathbf{y}(t) | \psi_n(t) \rangle | \theta) \\
&= \langle \mathbf{x}(t; \theta) + \mathbf{n}(t) | \psi_n(t) \rangle \\
&= \langle \mathbf{x}(t; \theta) | \psi_n(t) \rangle + \langle \mathbf{n}(t) | \psi_n(t) \rangle \\
&= \left\langle \sum_{k=1}^N \dot{x}_k(\theta) \psi_k(t) | \psi_n(t) \right\rangle + \dot{v}_n
\end{aligned}$$

$$\begin{aligned}
 &= \sum_{k=1}^N \dot{x}_k(\theta) \langle \psi_k(t) | \psi_n(t) \rangle + \dot{v}_n \\
 &= \dot{x}_n(\theta) + \dot{v}_n
 \end{aligned}$$

$$\begin{aligned}
 E(\dot{y}_n | \theta) &= E[\dot{x}_n(\theta) + \dot{v}_n] \\
 &= E\dot{x}_n(\theta) + E\dot{v}_n \\
 &= \dot{x}_n(\theta)
 \end{aligned}$$




Figure 9.1: Additive Gaussian noise channel Statistics

Theorem 9.3. Let $\mathbf{C} = \mathbf{C}_{\text{agn}}$ be an additive gaussian noise channel with distribution $n(t) \sim N(0, \sigma^2)$ for all t .

T H M	$\underbrace{\mathbf{C} = \mathbf{C}_{\text{agn}}}_{\text{additive Gaussian channel}} \implies \begin{cases} E\dot{v}_n = 0 \\ E(\dot{y}_n \theta) = \dot{x}_n(\theta) \\ \dot{v}_n \sim N(0, \sigma^2) \quad (\text{noise projections are Gaussian}) \\ \dot{y}_n \theta \sim N(\dot{x}_n(\theta), \sigma^2) \quad (\text{receiver projections are Gaussian}) \end{cases}$
----------------------	---

PROOF:

$$\begin{aligned}
 E\dot{v}_n &= E\langle n(t) | \psi_n(t) \rangle \\
 &= \langle En(t) | \psi_n(t) \rangle \\
 &= \langle 0 | \psi_n(t) \rangle \\
 &= 0
 \end{aligned}$$

$$\begin{aligned}
 (\dot{y}_n | \theta) &\triangleq \langle y(t) | \psi_n(t) \rangle | \theta \\
 &= \langle x(t; \theta) + n(t) | \psi_n(t) \rangle \\
 &= \langle x(t; \theta) | \psi_n(t) \rangle + \langle n(t) | \psi_n(t) \rangle \\
 &= \left\langle \sum_{k=1}^N \dot{x}_k(\theta) \psi_k(t) | \psi_n(t) \right\rangle + \dot{v}_n \\
 &= \sum_{k=1}^N \dot{x}_k(\theta) \langle \psi_k(t) | \psi_n(t) \rangle + \dot{v}_n \\
 &= \dot{x}_n(\theta) + \dot{v}_n
 \end{aligned}$$

$$\begin{aligned}
 E(\dot{y}_n | \theta) &= E[\dot{x}_n(\theta) + \dot{v}_n] \\
 &= E\dot{x}_n(\theta) + E\dot{v}_n \\
 &= \dot{x}_n(\theta)
 \end{aligned}$$

The distributions follow because they are linear operations on Gaussian processes.

Theorem 9.4. Let $\mathbf{C} = \mathbf{C}_{\text{awn}}$ be an additive white noise channel.

T H M	$\underbrace{\mathbf{C} = \mathbf{C}_{\text{awn}}}_{\text{additive white channel}} \Rightarrow \begin{cases} \mathbb{E}\dot{v}_n &= 0 & (\text{noise projection is zero-mean}) \\ \mathbb{E}(\dot{y}_n \theta) &= \dot{x}_n(\theta) & (\text{expected receiver projection} = \text{transmitted projection}) \\ \text{cov}[\dot{v}_n, \dot{v}_m] &= \sigma^2 \bar{\delta}_{nm} & (\text{noise projections are uncorrelated}) \\ \text{cov}[\dot{y}_n \theta, \dot{y}_m \theta] &= \sigma^2 \bar{\delta}_{nm} & (\text{receiver projections are uncorrelated}) \end{cases}$
----------------------	---

PROOF: Because the noise is additive (see Theorem 9.2 (page 77))

$$\begin{aligned} \mathbb{E}\dot{v}_n &= 0 \\ \mathbb{E}(\dot{y}_n|\theta) &= \dot{x}_n(\theta) + \dot{v}_n \\ \mathbb{E}(\dot{y}_n|\theta) &= \dot{x}_n(\theta). \end{aligned}$$

Because the noise is also white,

$$\begin{aligned} \text{cov}[\dot{v}_m, \dot{v}_n] &= \text{cov}[\langle n(t) | \psi_m(t) \rangle, \langle n(t) | \psi_n(t) \rangle] \\ &= \mathbb{E}[\langle n(t) | \psi_m(t) \rangle \langle n(t) | \psi_n(t) \rangle] \\ &= \mathbb{E}[\langle n(t) | \psi_m(t) \rangle \langle n(\hat{\theta}) | \psi_n(\hat{\theta}) \rangle] \\ &= \mathbb{E}[\langle n(\hat{\theta}) \langle n(t) | \psi_m(t) \rangle | \psi_n(\hat{\theta}) \rangle] \\ &= \mathbb{E}[\langle \langle n(\hat{\theta}) n(t) | \psi_m(t) \rangle | \psi_n(\hat{\theta}) \rangle] \\ &= \langle \langle \mathbb{E}[n(\hat{\theta}) n(t)] | \psi_m(t) \rangle | \psi_n(\hat{\theta}) \rangle \\ &= \langle \langle \sigma^2 \delta(t - \hat{\theta}) | \psi_m(t) \rangle | \psi_n(\hat{\theta}) \rangle \\ &= \sigma^2 \langle \psi_n(t) | \psi_m(t) \rangle \\ &= \begin{cases} \sigma^2 & \text{for } n = m \\ 0 & \text{for } n \neq m. \end{cases} \end{aligned}$$

$$\begin{aligned} \text{cov}[\dot{y}_n|\theta, \dot{y}_m|\theta] &= \mathbb{E}[\dot{y}_n \dot{y}_m|\theta] - [\mathbb{E}\dot{y}_n|\theta][\mathbb{E}\dot{y}_m|\theta] \\ &= \mathbb{E}[(\dot{x}_n(\theta) + \dot{v}_n)(\dot{x}_m(\theta) + \dot{v}_m)] - \dot{x}_n(\theta)\dot{x}_m(\theta) \\ &= \mathbb{E}[(\dot{x}_n(\theta) + \dot{v}_n)(\dot{x}_m(\theta) + \dot{v}_m)] - \dot{x}_n(\theta)\dot{x}_m(\theta) \\ &= \mathbb{E}[\dot{x}_n(\theta)\dot{x}_m(\theta) + \dot{x}_n(\theta)\dot{v}_m + \dot{v}_n\dot{x}_m(\theta) + \dot{v}_n\dot{v}_m] - \dot{x}_n(\theta)\dot{x}_m(\theta) \\ &= \dot{x}_n(\theta)\dot{x}_m(\theta) + \dot{x}_n(\theta)\mathbb{E}[\dot{v}_m] + \mathbb{E}[\dot{v}_n]\dot{x}_m(\theta) + \mathbb{E}[\dot{v}_n\dot{v}_m] - \dot{x}_n(\theta)\dot{x}_m(\theta) \\ &= 0 + \dot{x}_n(\theta) \cdot 0 + 0 \cdot \dot{x}_m(\theta) + \text{cov}[\dot{v}_n, \dot{v}_m] + [\mathbb{E}\dot{v}_n][\mathbb{E}\dot{v}_m] \\ &= \sigma^2 \bar{\delta}_{nm} + 0 \cdot 0 \\ &= \begin{cases} \sigma^2 & \text{for } n = m \\ 0 & \text{for } n \neq m. \end{cases} \end{aligned}$$




Figure 9.2: Additive white Gaussian noise channel statistics

Theorem 9.5. Let $\mathbf{C} = \mathbf{C}_{\text{awgn}}$ be an additive gaussian noise channel with distribution $n(t) \sim N(0, \sigma^2)$ for all t .

T
H
M

$$\underbrace{\mathbf{C} = \mathbf{C}_{\text{awgn}}}_{\text{AWGN}} \implies \begin{cases} \dot{v}_n \sim \mathcal{N}(0, \sigma^2) & (\text{noise projections are Gaussian}) \\ \dot{y}_n | \theta \sim \mathcal{N}(\dot{x}_n(\theta), \sigma^2) & (\text{receiver projections are Gaussian}) \\ \text{cov}[\dot{v}_n, \dot{v}_m] = \sigma^2 \bar{\delta}_{nm} & (\text{noise projections are uncorrelated}) \\ \text{cov}[\dot{y}_n, \dot{y}_m] = \sigma^2 \bar{\delta}_{nm} & (\text{receiver projections are uncorrelated}) \\ P\{\dot{v}_n = a \wedge \dot{v}_m = b\} = P\{\dot{v}_n = a\}P\{\dot{v}_m = b\} & (\text{noise projections are independent}) \\ P\{\dot{y}_n = a \wedge \dot{y}_m = b\} = P\{\dot{y}_n = a\}P\{\dot{y}_m = b\} & (\text{receiver projections are independent}) \end{cases}$$

PROOF: The distributions follow because they are linear operations on Gaussian processes.

By Theorem 9.4 (page 79) (for AWN channel)

$$\begin{aligned} E\dot{v}_n &= 0 \\ \text{cov}[\dot{v}_m, \dot{v}_n] &= \sigma^2 \bar{\delta}_{mn} \\ \dot{y}_n &= \dot{x}_n + \dot{v}_n \\ E\dot{y}_n &= \dot{x}_n \\ \text{cov}[\dot{y}_n, \dot{y}_m] &= \sigma^2 \bar{\delta}_{mn} \end{aligned}$$

Because the processes are Gaussian, uncorrelatedness implies *independence*. ⇒

9.2 Optimal symbol estimation

The AWGN projection statistics provided by Theorem 9.5 (page 79) help generate the optimal ML-estimates for a number of communication systems. These ML-estimates can be expressed in either of two standard forms:

- **Spectral decomposition:** The optimal estimate is expressed in terms of *projections* of signals onto orthonormal basis functions.
- **Matched signal:** The optimal estimate is expressed in terms of the (noisy) received signal correlated with (“matched” with) the (noiseless) transmitted signal.

Theorem 9.6 (page 80) (next) expresses the general optimal *ML estimate* in both of these forms.

Parameter detection is a special case of parameter estimation. In parameter detection, the estimate is a member of a finite set. In parameter estimation, the estimate is a member of an infinite set (Section 9.2 page 80).

Theorem 9.6 (General ML estimation). *Let Ψ be an orthonormal set spanning $\mathbf{x}(t)$ such that*

$$\begin{aligned} \Psi &\triangleq (\psi_1(t), \psi_2(t), \dots, \psi_n(t)) \\ \dot{y}_n &\triangleq \langle \mathbf{y}(t) | \psi_n(t) \rangle \\ \dot{x}_n &\triangleq \langle \mathbf{x}(t) | \psi_n(t) \rangle \\ \mathbf{y}(t) &= \mathbf{x}(t; \hat{\theta}) + \mathbf{n}(t). \end{aligned}$$

Then the optimal ML-estimate $\hat{\theta}_{\text{ml}}$ of parameter θ is

$$\begin{aligned} \hat{\theta}_{\text{ml}} &= \arg \min_{\hat{\theta}} \left[\sum_{n=1}^N [\dot{y}_n - \dot{x}_n(\hat{\theta})]^2 \right] && (\text{spectral decomposition}) \\ &= \arg \max_{\hat{\theta}} \left[2 \langle \mathbf{y}(t) | \mathbf{x}(t; \hat{\theta}) \rangle - \|\mathbf{x}(t; \hat{\theta})\|^2 \right] && (\text{matched signal}) \end{aligned}$$

PROOF:

$$\begin{aligned}
 \hat{\theta}_{\text{ml}} &= \arg \max_{\hat{\theta}} P\{y(t) | \mathbf{x}(t; \hat{\theta})\} \\
 &= \arg \max_{\hat{\theta}} P\{\dot{y}_1, \dot{y}_2, \dots, \dot{y}_n | \mathbf{x}(t; \hat{\theta})\} && \text{by Theorem 9.1 (page 75)} \\
 &= \arg \max_{\hat{\theta}} \prod_{n=1}^N P\{\dot{y}_n | \mathbf{x}(t; \hat{\theta})\} \\
 &= \arg \max_{\hat{\theta}} \prod_{n=1}^N p[\dot{y}_n | \mathbf{x}(t; \hat{\theta})] \\
 &= \arg \max_{\hat{\theta}} \prod_{n=1}^N \frac{1}{\sqrt{2\pi\sigma^2}} \exp \frac{[\dot{y}_n - \dot{x}_n(\hat{\theta})]^2}{-2\sigma^2} && \text{by Theorem 9.5 (page 79)} \\
 &= \arg \max_{\hat{\theta}} \left(\frac{1}{\sqrt{2\pi\sigma^2}} \right)^N \exp \frac{-1}{2\sigma^2} \sum_{n=1}^N [\dot{y}_n - \dot{x}_n(\hat{\theta})]^2 \\
 &= \arg \max_{\hat{\theta}} \left[- \sum_{n=1}^N [\dot{y}_n - \dot{x}_n(\hat{\theta})]^2 \right]
 \end{aligned}$$

$$\begin{aligned}
 &= \arg \max_{\hat{\theta}} \left[- \lim_{N \rightarrow \infty} \sum_{n=1}^N [\dot{y}_n - \dot{x}_n(\hat{\theta})]^2 \right] && \text{by Theorem 9.1 (page 75)} \\
 &= \arg \max_{\hat{\theta}} \left[- \|y(t) - \mathbf{x}(t; \hat{\theta})\|^2 \right] && \text{by Plancheral's formula} && (\text{Theorem G.9 page 210}) \\
 &= \arg \max_{\hat{\theta}} \left[- \|y(t)\|^2 + 2\mathbf{R}_e \langle y(t) | \mathbf{x}(t; \hat{\theta}) \rangle - \|\mathbf{x}(t; \hat{\theta})\|^2 \right] \\
 &= \arg \max_{\hat{\theta}} \left[2 \langle y(t) | \mathbf{x}(t; \hat{\theta}) \rangle - \|\mathbf{x}(t; \hat{\theta})\|^2 \right] && \text{because } y(t) \text{ independent of } \hat{\theta}
 \end{aligned}$$



Theorem 9.7 (ML amplitude estimation). ² Let \mathbf{S} be an additive white gaussian noise system such that

$$\begin{aligned}
 y(t) &= [\mathbf{C}_{\text{awgn}} s](t) = \mathbf{x}(t; a) + \mathbf{n}(t) \\
 \mathbf{x}(t; a) &\triangleq a \lambda(t).
 \end{aligned}$$

Then

T H M	$\hat{a}_{\text{ml}} = \frac{1}{\ \lambda(t)\ ^2} \langle y(t) \lambda(t) \rangle$ (optimal ML-estimate of a) $= \frac{1}{\ \lambda(t)\ ^2} \sum_{n=1}^N \dot{y}_n \dot{\lambda}_n$ $E\hat{a}_{\text{ml}} = a$ (\hat{a}_{ml} is unbiased) $\text{var } \hat{a}_{\text{ml}} = \frac{\sigma^2}{\ \lambda(t)\ ^2}$ (variance of estimate \hat{a}_{ml}) $\text{var } \hat{a}_{\text{ml}} = CR \text{ lower bound}$ (\hat{a}_{ml} is an efficient estimate)
----------------------	---

PROOF:

² Mandyam D. Srinath (1996) pages 158–159

1. *ML estimate* in “matched signal” form:

$$\begin{aligned}
 \hat{a}_{\text{ml}} &= \arg \max_a [2 \langle \mathbf{y}(t) | \mathbf{x}(t; \hat{\theta}) \rangle - \|\mathbf{x}(t; \phi)\|^2] && \text{by Theorem 9.6 (page 80)} \\
 &= \arg \max_a [2 \langle \mathbf{y}(t) | a\lambda(t) \rangle - \|a\lambda(t)\|^2] && \text{by hypothesis} \\
 &= \arg_a \left[\frac{\partial}{\partial a} 2a \langle \mathbf{y}(t) | \lambda(t) \rangle - \frac{\partial}{\partial a} a^2 \|\lambda(t)\|^2 = 0 \right] \\
 &= \arg_a [2 \langle \mathbf{y}(t) | \lambda(t) \rangle - 2a \|\lambda(t)\|^2 = 0] \\
 &= \arg_a [\langle \mathbf{y}(t) | \lambda(t) \rangle = a \|\lambda(t)\|^2] \\
 &= \frac{1}{\|\lambda(t)\|^2} \langle \mathbf{y}(t) | \lambda(t) \rangle
 \end{aligned}$$

2. *ML estimate* in “spectral decomposition” form:

$$\begin{aligned}
 \hat{a}_{\text{ml}} &= \arg \min_a \left(\sum_{n=1}^N [\dot{y}_n - \dot{x}_n(a)]^2 \right) && \text{by Theorem 9.6 (page 80)} \\
 &= \arg_a \left(\frac{\partial}{\partial a} \sum_{n=1}^N [\dot{y}_n - \dot{x}_n(a)]^2 = 0 \right) \\
 &= \arg_a \left(2 \sum_{n=1}^N [\dot{y}_n - \dot{x}_n(a)] \frac{\partial}{\partial a} \dot{x}_n(a) = 0 \right) \\
 &= \arg_a \left(\sum_{n=1}^N [\dot{y}_n - \langle a\lambda(t) | \psi_n(t) \rangle] \frac{\partial}{\partial a} \langle a\lambda(t) | \psi_n(t) \rangle = 0 \right) \\
 &= \arg_a \left(\sum_{n=1}^N [\dot{y}_n - a \langle \lambda(t) | \psi_n(t) \rangle] \frac{\partial}{\partial a} (a \langle \lambda(t) | \psi_n(t) \rangle) = 0 \right) \\
 &= \arg_a \left(\sum_{n=1}^N [\dot{y}_n - a \lambda_n] \langle \lambda(t) | \psi_n(t) \rangle = 0 \right) \\
 &= \arg_a \left(\sum_{n=1}^N [\dot{y}_n - a \lambda_n] \lambda_n = 0 \right) \\
 &= \arg_a \left(\sum_{n=1}^N \dot{y}_n \lambda_n = \sum_{n=1}^N a \lambda_n^2 \right) \\
 &= \left(\frac{1}{\sum_{n=1}^N \lambda_n^2} \right) \sum_{n=1}^N \dot{y}_n \lambda_n \\
 &= \frac{1}{\|\lambda(t)\|^2} \sum_{n=1}^N \dot{y}_n \lambda_n
 \end{aligned}$$

3. Prove that the estimate \hat{a}_{ml} is **unbiased**:

$$\begin{aligned}
 E\hat{a}_{\text{ml}} &= E \frac{1}{\|\lambda(t)\|^2} \int_{t \in \mathbb{R}} \mathbf{y}(t) \lambda(t) dt && \text{by previous result} \\
 &= E \frac{1}{\|\lambda(t)\|^2} \int_{t \in \mathbb{R}} [a\lambda(t) + n(t)] \lambda(t) dt && \text{by hypothesis} \\
 &= \frac{1}{\|\lambda(t)\|^2} \int_{t \in \mathbb{R}} E[a\lambda(t) + n(t)] \lambda(t) dt && \text{by linearity of } \int \cdot dt \text{ and } E
 \end{aligned}$$

$$\begin{aligned}
 &= \frac{1}{\|\lambda(t)\|^2} a \int_{t \in \mathbb{R}} \lambda^2(t) dt && \text{by E operation} \\
 &= \frac{1}{\|\lambda(t)\|^2} a \|\lambda(t)\|^2 && \text{by definition of } \|\cdot\|^2 \\
 &= a
 \end{aligned}$$

4. Compute the variance of \hat{a}_{ml} :

$$\begin{aligned}
 \mathbb{E}\hat{a}_{ml}^2 &= \mathbb{E} \left[\frac{1}{\|\lambda(t)\|^2} \int_{t \in \mathbb{R}} y(t) \lambda(t) dt \right]^2 \\
 &= \mathbb{E} \left[\frac{1}{\|\lambda(t)\|^4} \int_{t \in \mathbb{R}} y(t) \lambda(t) dt \int_v y(v) \lambda(v) dv \right] \\
 &= \mathbb{E} \left[\frac{1}{\|\lambda(t)\|^4} \int_{t \in \mathbb{R}} \int_v [a\lambda(t) + n(t)][a\lambda(v) + n(v)] \lambda(t) \lambda(v) dv dt \right] \\
 &= \mathbb{E} \left[\frac{1}{\|\lambda(t)\|^4} \int_{t \in \mathbb{R}} \int_v [a^2 \lambda(t) \lambda(v) + a\lambda(t)n(v) + a\lambda(v)n(t) + n(t)n(v)] \lambda(t) \lambda(v) dv dt \right] \\
 &= \left[\frac{1}{\|\lambda(t)\|^4} \int_{t \in \mathbb{R}} \int_v [a^2 \lambda(t) \lambda(v) + 0 + 0 + \sigma^2 \delta(t-v)] \lambda(t) \lambda(v) dv dt \right] \\
 &= \frac{1}{\|\lambda(t)\|^4} \int_{t \in \mathbb{R}} \int_v a^2 \lambda^2(t) \lambda^2(v) dv dt + \frac{1}{\|\lambda(t)\|^4} \int_{t \in \mathbb{R}} \int_v \sigma^2 \delta(t-v) \lambda(t) \lambda(v) dv dt \\
 &= \frac{1}{\|\lambda(t)\|^4} a^2 \int_{t \in \mathbb{R}} \lambda^2(t) dt \int_v \lambda^2(v) dv + \frac{1}{\|\lambda(t)\|^4} \sigma^2 \int_{t \in \mathbb{R}} \lambda^2(t) dt \\
 &= a^2 \frac{1}{\|\lambda(t)\|^4} \|\lambda(t)\|^2 \|\lambda(v)\|^2 + \frac{1}{\|\lambda(t)\|^4} \sigma^2 \|\lambda(t)\|^2 \\
 &= a^2 + \frac{\sigma^2}{\|\lambda(t)\|^2}
 \end{aligned}$$

$$\begin{aligned}
 \text{var } \hat{a}_{ml} &= \mathbb{E}\hat{a}_{ml}^2 - (\mathbb{E}\hat{a}_{ml})^2 \\
 &= \left(a^2 + \frac{\sigma^2}{\|\lambda(t)\|^2} \right) - \left(a^2 \right) \\
 &= \frac{\sigma^2}{\|\lambda(t)\|^2}
 \end{aligned}$$

5. Compute the Cramér-Rao Bound:

$$\begin{aligned}
 p[y(t)|s(t; a)] &= p[\dot{y}_1, \dot{y}_2, \dots, \dot{y}_N | s(t; a)] \\
 &= \prod_{n=1}^N \frac{1}{\sqrt{2\pi\sigma^2}} \exp \frac{(\dot{y}_n - a\dot{\lambda}_n)^2}{-2\sigma^2} \\
 &= \left(\frac{1}{\sqrt{2\pi\sigma^2}} \right)^N \exp \frac{1}{-2\sigma^2} \sum_{n=1}^N (\dot{y}_n - a\dot{\lambda}_n)^2
 \end{aligned}$$

$$\begin{aligned}
 \frac{\partial}{\partial a} \ln p[y(t)|s(t; a)] &= \frac{\partial}{\partial a} \ln \left(\frac{1}{\sqrt{2\pi\sigma^2}} \right)^N \exp \frac{1}{-2\sigma^2} \sum_{n=1}^N (\dot{y}_n - a\dot{\lambda}_n)^2 \\
 &= \frac{\partial}{\partial a} \ln \left(\frac{1}{\sqrt{2\pi\sigma^2}} \right)^N + \frac{\partial}{\partial a} \ln \exp \frac{1}{-2\sigma^2} \sum_{n=1}^N (\dot{y}_n - a\dot{\lambda}_n)^2 \\
 &= \frac{\partial}{\partial a} \frac{1}{-2\sigma^2} \sum_{n=1}^N (\dot{y}_n - a\dot{\lambda}_n)^2
 \end{aligned}$$

$$\begin{aligned}
 &= \frac{1}{-2\sigma^2} \sum_{n=1}^N 2(\dot{y}_n - a\dot{\lambda}_n)(-\dot{\lambda}_n) \\
 &= \frac{1}{\sigma^2} \sum_{n=1}^N \dot{\lambda}_n(\dot{y}_n - a\dot{\lambda}_n)
 \end{aligned}$$

$$\begin{aligned}
 \frac{\partial^2}{\partial a^2} \ln p[y(t)|s(t; a)] &= \frac{\partial}{\partial a} \frac{\partial}{\partial a} \ln p[y(t)|s(t; a)] \\
 &= \frac{\partial}{\partial a} \frac{1}{\sigma^2} \sum_{n=1}^N \dot{\lambda}_n(\dot{y}_n - a\dot{\lambda}_n) \\
 &= \frac{1}{\sigma^2} \sum_{n=1}^N \dot{\lambda}_n(-\dot{\lambda}_n) \\
 &= \frac{-1}{\sigma^2} \sum_{n=1}^N \dot{\lambda}_n^2 \\
 &= \frac{-\|\lambda(t)\|^2}{\sigma^2}
 \end{aligned}$$

$$\begin{aligned}
 \text{var } \hat{a}_{\text{ml}} &\triangleq E[\hat{a}_{\text{ml}} - E\hat{a}_{\text{ml}}]^2 \\
 &= E[\hat{a}_{\text{ml}} - a]^2 \\
 &\geq \frac{-1}{E\left(\frac{\partial^2}{\partial a^2} \ln p[y(t)|s(t; a)]\right)} \\
 &= \frac{-1}{E\left(\frac{-\|\lambda(t)\|^2}{\sigma^2}\right)} \\
 &= \frac{\sigma^2}{\|\lambda(t)\|^2} \quad (\text{Cramér-Rao lower bound of the variance})
 \end{aligned}$$

6. Prove that \hat{a}_{ml} is an **efficient estimate**:

A estimate is *efficient* if $\text{var } \hat{a}_{\text{ml}} = \text{CR lower bound}$. We have already proven this, so \hat{a}_{ml} is an *efficient* estimate.

Also, even without explicitly computing the variance of \hat{a}_{ml} , the variance equals the *Cramér-Rao lower bound* (and hence \hat{a}_{ml} is an *efficient* estimate) if and only if

$$\begin{aligned}
 \hat{a}_{\text{ml}} - a &= \left(\frac{-1}{E\left(\frac{\partial^2}{\partial a^2} \ln p[y(t)|s(t; a)]\right)} \right) \left(\frac{\partial}{\partial a} \ln p[y(t)|s(t; a)] \right) \\
 &\left(\frac{-1}{E\left(\frac{\partial^2}{\partial a^2} \ln p[y(t)|s(t; a)]\right)} \right) \left(\frac{\partial}{\partial a} \ln p[y(t)|s(t; a)] \right) = \left(\frac{\sigma^2}{\|\lambda(t)\|^2} \right) \left(\frac{1}{\sigma^2} \sum_{n=1}^N \dot{\lambda}(\dot{y} - a\dot{\lambda}) \right) \\
 &= \frac{1}{\|\lambda(t)\|^2} \sum_{n=1}^N \dot{\lambda} \dot{y} - \frac{1}{\|\lambda(t)\|^2} \sum_{n=1}^N \dot{\lambda}^2 \\
 &= \hat{a}_{\text{ml}} - a
 \end{aligned}$$



Theorem 9.8 (ML phase estimation). ³ Let S be an additive white gaussian noise system such that

³ Mandyam D. Srinath (1996) pages 159–160

$$\begin{aligned}\mathbf{y}(t) &= [\mathbf{C}_{\text{awgn}} s](t) = \mathbf{x}(t; \phi) + \mathbf{n}(t) \\ \mathbf{x}(t; \phi) &= A \cos(2\pi f_c t + \phi).\end{aligned}$$

Then the optimal ML-estimate of parameter ϕ is

$$\boxed{\text{T H M} \quad \hat{\phi}_{\text{ml}} = -\text{atan} \left(\frac{\langle \mathbf{y}(t) | \sin(2\pi f_c t) \rangle}{\langle \mathbf{y}(t) | \cos(2\pi f_c t) \rangle} \right)}$$

PROOF:

$$\begin{aligned}\hat{\phi}_{\text{ml}} &= \arg \max_{\phi} [2 \langle \mathbf{y}(t) | \mathbf{x}(t; \hat{\theta}) \rangle - \|\mathbf{x}(t; \phi)\|^2] && \text{by Theorem 9.6 (page 80)} \\ &= \arg \max_{\phi} [2 \langle \mathbf{y}(t) | \mathbf{x}(t; \phi) \rangle] && \text{because } \|\mathbf{x}(t; \phi)\| \text{ does not depend on } \phi \\ &= \arg_{\phi} \left[\frac{\partial}{\partial \phi} \langle \mathbf{y}(t) | \mathbf{x}(t; \phi) \rangle = 0 \right] \\ &= \arg_{\phi} \left[\left\langle \mathbf{y}(t) | \frac{\partial}{\partial \phi} \mathbf{x}(t; \phi) \right\rangle = 0 \right] && \text{because } \langle \cdot | \cdot \rangle \text{ is a linear operator} \\ &= \arg_{\phi} \left[\left\langle \mathbf{y}(t) | \frac{\partial}{\partial \phi} A \cos(2\pi f_c t + \phi) \right\rangle = 0 \right] \\ &= \arg_{\phi} [\langle \mathbf{y}(t) | -A \sin(2\pi f_c t + \phi) \rangle = 0] \\ &= \arg_{\phi} [-A \langle \mathbf{y}(t) | \cos(2\pi f_c t) \sin \phi + \sin(2\pi f_c t) \cos \phi \rangle = 0] \\ &= \arg_{\phi} [\sin \phi \langle \mathbf{y}(t) | \cos(2\pi f_c t) \rangle = -\cos \phi \langle \mathbf{y}(t) | \sin(2\pi f_c t) \rangle] \\ &= \arg_{\phi} \left[\frac{\sin \phi}{\cos \phi} = -\frac{\langle \mathbf{y}(t) | \sin(2\pi f_c t) \rangle}{\langle \mathbf{y}(t) | \cos(2\pi f_c t) \rangle} \right] \\ &= \arg_{\phi} \left[\tan \phi = -\frac{\langle \mathbf{y}(t) | \sin(2\pi f_c t) \rangle}{\langle \mathbf{y}(t) | \cos(2\pi f_c t) \rangle} \right] \\ &= -\text{atan} \left(\frac{\langle \mathbf{y}(t) | \sin(2\pi f_c t) \rangle}{\langle \mathbf{y}(t) | \cos(2\pi f_c t) \rangle} \right)\end{aligned}$$



Theorem 9.9 (ML estimation of a function of a parameter). ⁴ Let \mathbf{S} be an additive white gaussian noise system such that $\mathbf{y}(t) = [\mathbf{C}_{\text{awgn}} s](t) = \mathbf{x}(t; \hat{\theta}) + \mathbf{n}(t)$
 $\mathbf{x}(t; \hat{\theta}) = \mathbf{g}(\hat{\theta})$

where \mathbf{g} is one-to-one and onto (invertible).

T
H
M

Then the optimal ML-estimate of parameter u is

$$\hat{\theta}_{\text{ml}} = \mathbf{g}^{-1} \left(\frac{1}{N} \sum_{n=1}^N \dot{y}_n \right).$$

If an ML ESTIMATE $\hat{\theta}_{\text{ml}}$ is unbiased ($E\hat{\theta}_{\text{ml}} = \theta$) then

$$\text{var } \hat{\theta}_{\text{ml}} \geq \frac{\sigma^2}{N} \frac{1}{\left[\frac{\partial \mathbf{g}(\theta)}{\partial \theta} \right]^2}.$$

If $\mathbf{g}(\theta) = \theta$ then $\hat{\theta}_{\text{ml}}$ is an **efficient** estimate such that $\text{var } \hat{\theta}_{\text{ml}} = \frac{\sigma^2}{N}$.

PROOF:

$$\hat{\theta}_{\text{ml}} = \arg \min_{\hat{\theta}} \left[\sum_{n=1}^N [\dot{y}_n - \mathbf{g}(\hat{\theta})]^2 \right]$$

⁴ Mandyam D. Srinath (1996) pages 142–143

$$\begin{aligned}
&= \arg_{\hat{\theta}} \left[\frac{\partial}{\partial \hat{\theta}} \sum_{n=1}^N [\dot{y}_n - g(\hat{\theta})]^2 = 0 \right] \\
&= \arg_{\hat{\theta}} \left[2 \sum_{n=1}^N [\dot{y}_n - g(\hat{\theta})] \frac{\partial}{\partial \hat{\theta}} g(\hat{\theta}) = 0 \right] \\
&= \arg_{\hat{\theta}} \left[2 \sum_{n=1}^N [\dot{y}_n - g(\hat{\theta})] = 0 \right] \\
&= \arg_{\hat{\theta}} \left[\sum_{n=1}^N \dot{y}_n = N g(\hat{\theta}) \right] \\
&= \arg_{\hat{\theta}} \left[g(\hat{\theta}) = \frac{1}{N} \sum_{n=1}^N \dot{y}_n \right] \\
&= \arg_{\hat{\theta}} \left[u = g^{-1} \left(\frac{1}{N} \sum_{n=1}^N \dot{y}_n \right) \right] \\
&= g^{-1} \left(\frac{1}{N} \sum_{n=1}^N \dot{y}_n \right)
\end{aligned}$$

If $\hat{\theta}_{\text{ml}}$ is unbiased ($E\hat{\theta}_{\text{ml}} = \theta$), we can use the *Cramér-Rao bound* to find a lower bound on the variance:

$$\begin{aligned}
\text{var } \hat{\theta}_{\text{ml}} &\triangleq E[\hat{\theta}_{\text{ml}} - E\hat{\theta}_{\text{ml}}]^2 \\
&= E[\hat{\theta}_{\text{ml}} - \theta]^2 \\
&\geq \frac{-1}{E \left(\frac{\partial^2}{\partial \theta^2} \ln p[y(t)|s(t; \theta)] \right)} \\
&= \frac{-1}{E \left(\frac{\partial^2}{\partial \theta^2} \ln p[\dot{y}_1, \dot{y}_2, \dots, \dot{y}_N | s(t; \theta)] \right)} \\
&= \frac{-1}{E \left(\frac{\partial^2}{\partial \theta^2} \ln \left[\left(\frac{1}{\sqrt{2\pi\sigma^2}} \right)^N \exp \frac{-1}{2\sigma^2} \sum_{n=1}^N [\dot{y}_n - g(\theta)]^2 \right] \right)} \\
&= \frac{-1}{E \left(\frac{\partial^2}{\partial \theta^2} \ln \left[\left(\frac{1}{\sqrt{2\pi\sigma^2}} \right)^N \right] + \frac{\partial^2}{\partial \theta^2} \ln \left[\exp \frac{-1}{2\sigma^2} \sum_{n=1}^N [\dot{y}_n - g(\theta)]^2 \right] \right)} \\
&= \frac{-1}{E \left(\frac{\partial^2}{\partial \theta^2} \left(\frac{-1}{2\sigma^2} \sum_{n=1}^N [\dot{y}_n - g(\theta)]^2 \right) \right)} \\
&= \frac{2\sigma^2}{E \left(\frac{\partial}{\partial \theta} \frac{\partial}{\partial \theta} \sum_{n=1}^N [\dot{y}_n - g(\theta)]^2 \right)} \\
&= \frac{2\sigma^2}{E \left(-2 \frac{\partial}{\partial \theta} \frac{\partial g(\theta)}{\partial \theta} \sum_{n=1}^N [\dot{y}_n - g(\theta)] \right)} \\
&= \frac{-\sigma^2}{E \left(\frac{\partial g^2(\theta)}{\partial \theta^2} \sum_{n=1}^N [\dot{y}_n - g(\theta)] + \frac{\partial g(\theta)}{\partial \theta} \frac{\partial}{\partial \theta} \sum_{n=1}^N [\dot{y}_n - g(\theta)] \right)}
\end{aligned}$$

$$\begin{aligned}
&= \frac{-\sigma^2}{\mathbb{E} \left(\frac{\partial g^2(\theta)}{\partial \theta^2} \sum_{n=1}^N [\dot{y}_n - g(\theta)] - N \frac{\partial g(\theta)}{\partial \theta} \frac{\partial g(\theta)}{\partial \theta} \right)} \\
&= \frac{-\sigma^2}{\frac{\partial g^2(\theta)}{\partial \theta^2} \sum_{n=1}^N \mathbb{E}[\dot{y}_n - g(\theta)] - N \frac{\partial g(\theta)}{\partial \theta} \frac{\partial g(\theta)}{\partial \theta}} \\
&= \frac{-\sigma^2}{-N \frac{\partial g(\theta)}{\partial \theta} \frac{\partial g(\theta)}{\partial \theta}} \\
&= \frac{\sigma^2}{N} \frac{1}{\left[\frac{\partial g(\theta)}{\partial \theta} \right]^2}
\end{aligned}$$

The inequality becomes equality (an *efficient* estimate) if and only if

$$\hat{\theta}_{\text{ml}} - \theta = \left(\frac{-1}{\mathbb{E} \left(\frac{\partial^2}{\partial \theta^2} \ln p[y(t)|s(t; \theta)] \right)} \right) \left(\frac{\partial}{\partial \theta} \ln p[y(t)|s(t; \theta)] \right).$$

$$\begin{aligned}
&\left(\frac{-1}{\mathbb{E} \left(\frac{\partial^2}{\partial \theta^2} \ln p[y(t)|s(t; \theta)] \right)} \right) \left(\frac{\partial}{\partial \theta} \ln p[y(t)|s(t; \theta)] \right) = \left(\frac{\sigma^2}{N} \frac{1}{\left[\frac{\partial g(\theta)}{\partial \theta} \right]^2} \right) \left(\frac{-1}{2\sigma^2} (2) \frac{\partial g(\theta)}{\partial \theta} \sum_{n=1}^N [\dot{y}_n - g(\theta)] \right) \\
&= -\frac{1}{N} \frac{1}{\frac{\partial g(\theta)}{\partial \theta}} \left(\sum_{n=1}^N [\dot{y}_n - g(\theta)] \right) \\
&= -\frac{1}{\frac{\partial g(\theta)}{\partial \theta}} \left(\frac{1}{N} \sum_{n=1}^N \dot{y}_n - g(\theta) \right) \\
&= -\frac{1}{\frac{\partial g(\theta)}{\partial \theta}} (\hat{\theta}_{\text{ml}} - g(\theta)) \\
&= -(\hat{\theta}_{\text{ml}} - \theta)
\end{aligned}$$



9.3 Colored noise

This chapter presented several theorems whose results depended on the noise being white. However if the noise is **colored**, then these results are invalid. But there is still hope for colored noise. Processing colored signals can be accomplished using two techniques:

1. Karhunen-Loëve basis functions ⁵
2. whitening filter ⁶

Karhunen-Loëve. If the noise is white, the set $\{\langle y(t) | \psi_n(t) \rangle\}$ is a sufficient statistic regardless of which set $\{\psi_n(t)\}$ of orthonormal basis functions are used. If the noise is colored, and if $\{\psi_n(t)\}$

⁵ Karhunen-Loëve: Section 3.3 (page 20)

⁶ Continuous data whitening: Section 6.3 page 48

Discrete data whitening: Section 5.4 page 38

satisfy the Karhunen-Loève criterion

$$\int_{t_2} R_{xx}(t_1, t_2) \psi_n(t_2) dt_2 = \lambda_n \psi_n(t_1)$$

then $\{\langle y(t) | \psi_n(t) \rangle\}$ is still a sufficient statistic.

Whitening filter. The whitening filter makes the received signal $y(t)$ statistically white (uncorrelated in time). In this case, any orthonormal basis set can be used to generate sufficient statistics.

9.4 Signal matching

Detection methods. There are basically two types of detection methods:

1. signal matching
2. orthonormal decomposition.

Let S be the set of transmitted waveforms and Y be a set of orthonormal basis functions that span S . *Signal matching* computes the innerproducts of a received signal $y(t)$ with each signal from S . *Orthonormal decomposition* computes the innerproducts of $y(t)$ with each signal from the set Y .

In the case where $|S|$ is large, often $|R| \ll |S|$ making orthonormal decomposition much easier to implement. For example, in a QAM-64 modulation system, signal matching requires $|S| = 64$ innerproduct calculations, while orthonormal decomposition only requires $|R| = 2$ innerproduct calculations because all 64 signals in S can be spanned by just 2 orthonormal basis functions.

Maximizing SNR. Theorem 9.1 (page 75) shows that the innerproducts of $y(t)$ with basis functions of Y is sufficient for optimal detection. Theorem 9.10 (page 88) (next) shows that a receiver can maximize the SNR of a received signal when signal matching is used.

Theorem 9.10. Let $x(t)$ be a transmitted signal, $n(t)$ noise, and $y(t)$ the received signal in an AWGN channel. Let the SIGNAL TO NOISE RATIO SNR be defined as

$$\text{SNR}[y(t)] \triangleq \frac{|\langle x(t) | x(t) \rangle|^2}{E[|\langle n(t) | x(t) \rangle|^2]}.$$

T H M $\text{SNR}[y(t)] \leq \frac{2 \|x(t)\|^2}{N_o}$ and is maximized (equality) when $x(t) = ax(t)$, where $a \in \mathbb{R}$.

PROOF:

$$\begin{aligned} \text{SNR}[y(t)] &\triangleq \frac{|\langle x(t) | x(t) \rangle|^2}{E[|\langle n(t) | x(t) \rangle|^2]} \\ &= \frac{|\langle x(t) | f(t) \rangle|^2}{E\left[\left[\int_{t \in \mathbb{R}} n(t)x^*(t) dt\right] \left[\int_{\hat{\theta}} n(\hat{\theta})f^*(\hat{\theta}) du\right]^*\right]} \\ &= \frac{|\langle x(t) | x(t) \rangle|^2}{E\left[\int_{t \in \mathbb{R}} \int_{\hat{\theta}} n(t)n^*(\hat{\theta})x^*(t)x(\hat{\theta}) dt du\right]} \end{aligned}$$



$$\begin{aligned}
&= \frac{|\langle \mathbf{x}(t) | f(t) \rangle|^2}{\int_{t \in \mathbb{R}} \int_{\hat{\theta}} \mathbb{E}[\mathbf{n}(t)\mathbf{n}^*(\hat{\theta})] \mathbf{x}^*(t)\mathbf{x}(\hat{\theta}) dt du} \\
&= \frac{|\langle \mathbf{x}(t) | \mathbf{x}(t) \rangle|^2}{\int_{t \in \mathbb{R}} \int_{\hat{\theta}} \frac{1}{2} N_o \delta(t - \hat{\theta}) \mathbf{x}^*(t)\mathbf{x}(\hat{\theta}) dt du} \\
&= \frac{|\langle \mathbf{x}(t) | \mathbf{x}(t) \rangle|^2}{\frac{1}{2} N_o \int_{t \in \mathbb{R}} \mathbf{x}^*(t)\mathbf{x}(t) dt} \\
&= \frac{|\langle \mathbf{x}(t) | \mathbf{x}(t) \rangle|^2}{\frac{1}{2} N_o \|\mathbf{x}(t)\|^2} \\
&\leq \frac{|\|\mathbf{x}(t)\| \|\mathbf{x}(t)\||^2}{\frac{1}{2} N_o \|\mathbf{x}(t)\|^2} \quad \text{by Cauchy-Schwarz Inequality (Theorem 1.2 page 234)} \\
&= \frac{2 \|\mathbf{x}(t)\|^2}{N_o}
\end{aligned}$$

The Cauchy-Schwarz Inequality becomes an equality (SNR is maximized) when $\mathbf{x}(t) = a\mathbf{x}(t)$. \Rightarrow

Implementation. The innerproduct operations can be implemented using either

1. a correlator or
2. a matched filter.

A correlator is simply an integrator of the form $\langle \mathbf{y}(t) | f(t) \rangle = \int_0^T \mathbf{y}(t)f(t) dt$.

A matched filter introduces a function $h(t)$ such that $h(t) = \mathbf{x}(T-t)$ (which implies $\mathbf{x}(t) = h(T-t)$) giving

$$\langle \mathbf{y}(t) | \mathbf{x}(t) \rangle = \underbrace{\int_0^T \mathbf{y}(t)\mathbf{x}(t) dt}_{\text{correlator}} = \underbrace{\int_0^\infty \mathbf{x}(\tau)h(t-\tau) d\tau \Big|_{t=T}}_{\text{matched filter}} = \mathbf{x}(t) \star h(t)|_{t=T}.$$

This shows that $h(t)$ is the impulse response of a filter operation sampled at time T . By Theorem 9.10 (page 88), the optimal impulse response is $h(T-t) = f(t) = \mathbf{x}(t)$. That is, the optimal $h(t)$ is just a “flipped” and shifted version of $\mathbf{x}(t)$.

CHAPTER 10

MOMENT ESTIMATION

Theorem 10.1. Let $\hat{\mu} \triangleq \sum_{n=1}^N \lambda_n \mathbf{x}_n$ with $\sum_{n=1}^N \lambda_n = 1$ be the ARITHMETIC MEAN (Definition L.4 page 273).

T
H
M

$$\left\{ \begin{array}{l} (A). (\mathbf{x}_n) \text{ is WIDE SENSE STATIONARY} \\ (B). \mu \triangleq E\mathbf{x}_n \\ (C). (\mathbf{x}_n) \text{ is UNCORRELATED} \\ (D). \hat{\mu} \triangleq \sum_{n=1}^N \lambda_n \mathbf{x}_n \text{ (ARITHMETIC MEAN)} \end{array} \right. \text{ and } \left\{ \begin{array}{l} (1). E\hat{\mu} = \mu \text{ (UNBIASED) and} \\ (2). \text{var}(\hat{\mu}) = \sigma^2 \sum_{n=1}^N \lambda_n^2 \text{ and} \\ (3). \text{mse}(\hat{\mu}) = \sigma^2 \sum_{n=1}^N \lambda_n^2 \end{array} \right. \Rightarrow$$

PROOF:

$$\begin{aligned}
 E\hat{\mu} &\triangleq E \sum_{n \in \mathbb{Z}} \lambda_n \mathbf{x}_n && \text{by definition of } \textit{arithmetic mean} \quad (\text{Definition L.4 page 273}) \\
 &= \sum_{n \in \mathbb{Z}} \lambda_n E\mathbf{x}_n && \text{by } \textit{linearity of } E \quad (\text{Theorem 1.1 page 4}) \\
 &= \mu \sum_{n \in \mathbb{Z}} \lambda_n && \text{by } \textit{WSS hypothesis} \quad (\text{A}) \\
 &= \mu && \text{by } \sum \lambda_n = 1 \text{ hypothesis} \quad (\text{Definition L.4 page 273}) \\
 \text{var}(\hat{\mu}) &\triangleq E(\hat{\mu} - E\hat{\mu})^2 && \text{by definition of } \textit{variance} \\
 &= E(\hat{\mu} - \mu)^2 && \text{by previous result} \\
 &= E \left(\sum_{n=1}^N \lambda_n \mathbf{x}_n - \mu \right)^2 && \text{by definition of } \hat{\mu} \\
 &= E \left[\sum_{n=1}^N \mathbf{x}_n - \mu \underbrace{\sum_{n=1}^N \lambda_n}_1 \right]^2 \\
 &= E \left[\sum_{n=1}^N \lambda_n (\mathbf{x}_n - \mu) \right]^2
 \end{aligned}$$

$$\begin{aligned}
&= E \left[\sum_{n=1}^N \lambda_n (\mathbf{x}_n - \mu) \sum_{m=1}^N \lambda_m (\mathbf{x}_m - \mu) \right] \\
&= \sum_{n=1}^N \sum_{m=1}^N \lambda_n \lambda_m (E[(\mathbf{x}_n - \mu)(\mathbf{x}_m - \mu)]) \\
&= \sum_{n=1}^N \sum_{m=1}^N \lambda_n \lambda_m (E[\mathbf{x}_n \mathbf{x}_m] - \mu E[\mathbf{x}_n] - \mu E[\mathbf{x}_m] + \mu^2) \\
&= \sum_{n=1}^N \sum_{m=1}^N \lambda_n \lambda_m (E[\mathbf{x}_n \mathbf{x}_m] - \mu^2 - \mu^2 + \mu^2) \\
&= \sum_{n=1}^N \sum_{m=1}^N \lambda_n \lambda_m (E[\mathbf{x}_n \mathbf{x}_m] - \mu^2) \\
&= \sum_{n=1}^N \lambda_n^2 E[\mathbf{x}_n^2] - \mu^2 + \sum_{n=1}^N \sum_{m \neq n} \lambda_n \lambda_m (E[\mathbf{x}_n \mathbf{x}_m] - \mu^2) \\
&= \sum_{n=1}^N \lambda_n^2 (E[\mathbf{x}_n^2] - \mu^2) + \sum_{n=1}^N \sum_{m \neq n} \lambda_n \lambda_m (E[\mathbf{x}_n \mathbf{x}_m] - \mu^2) \\
&= \sum_{n=1}^N \lambda_n^2 \sigma^2 + \frac{1}{N^2} \sum_{n=1}^N \sum_{m \neq n} \lambda_n \lambda_m (\mu \mu - \mu^2) \xrightarrow{0} 0 \\
&= \sigma^2 \sum_{n=1}^N \lambda_n^2
\end{aligned}$$

$$\text{mse}(\hat{\mu}) = E(\hat{\mu} - E\hat{\mu})^2 + (E\hat{\mu} - \mu)^2$$

by Theorem 8.2 page 65

$$= \sigma^2 \sum_{n=1}^N \lambda_n^2 + (\mu - \mu)^2$$

by previous results

$$= \sigma^2 \sum_{n=1}^N \lambda_n^2$$

⇒


Corollary 10.1.¹

C O R	$\left\{ \begin{array}{l} (A). \quad (\mathbf{x}_n) \text{ is WIDE SENSE STATIONARY} \\ (B). \quad \mu \triangleq E\mathbf{x}_n \\ (C). \quad (\mathbf{x}_n) \text{ is UNCORRELATED} \\ (D). \quad \hat{\mu} \triangleq \frac{1}{N} \sum_{n=1}^N \mathbf{x}_n \quad (\text{AVERAGE}) \end{array} \right. \quad \text{and} \quad \left\{ \begin{array}{l} (1). \quad E\hat{\mu} = \mu \quad (\text{UNBIASED}) \\ (2). \quad \text{var}(\hat{\mu}) = \frac{\sigma^2}{N} \\ (3). \quad \text{mse}(\hat{\mu}) = \frac{\sigma^2}{N} \quad (\text{CONSISTENT}) \end{array} \right.$
-------------	---

¹  Kay (1988) page 45 (§“3.3 ESTIMATION THEORY”)

CHAPTER 11

SYSTEM IDENTIFICATION



“I can therefore gladly admit that falsificationists like myself much prefer an attempt to solve an interesting problem by a bold conjecture, even (and especially) if it so turns out to be false, to any recital of a sequence of irrelevant truisms. We prefer this because we believe that this is the way in which we can learn from our mistakes and that in finding that our conjecture was false we shall have learned much about the truth, and shall have got nearer to the truth.”

Karl R. Popper (1902–1994)¹

11.1 Estimation techniques

Let \mathbf{S} be a system with *impulse response* $h(n)$ with *DTFT* $\tilde{H}(\omega)$, input $x(n)$, and output $y(n)$. Often in the field of “digital signal processing” (DSP), \mathbf{S} is a “filter” with known $h(n)$ and $\tilde{H}(\omega)$ because the filter \mathbf{S} was designed by a designer who had direct control over $h(n)$.

However in many other practical situations, \mathbf{S} is some other system for which $h(n)$ and $\tilde{H}(\omega)$ are *not* known...but which we may want to *estimate*. Examples of such \mathbf{S} is a device on an industrial shaker table, a communication channel, or the entire earth.

Determining $h(n)$ and/or $\tilde{H}(\omega)$ is part of an operation called “*system identification*”. Determining $\tilde{H}(\omega)$ in particular is referred to as “*Frequency Response Identification*”² or as “*Frequency Response Function*” (“*FRF*”) estimation.³ *FRF* estimation is a challenging problem and one that many have devoted much effort to. This chapter describes some of that effort.

In the early days, people used a rather obvious technique for determining $\tilde{H}(\omega)$ —the humble *sine sweep*. That is, they drove the input with a sine wave with slowly increasing (or decreasing) frequency while measuring the resulting output. This technique, although effective, was “very slow”.⁴


¹ quote: [Popper \(1962\)](#), page 231, [Popper \(1963\)](#) page 313

image: https://en.wikipedia.org/wiki/File:Karl_Popper.jpg, “no known copyright restrictions”


² [Shin and Hammond \(2008\)](#) page 292

³ [Cobb \(1988\)](#) page 1 (FRF “measurement”)

⁴ [Leuridan et al. \(1986\)](#) 911 “Stepped Sine Testing”, [Cobb \(1988\)](#) page 1 (Chapter 1—Introduction), [Ewins](#)



(A) communications additive noise model

The “input signal” is $x(n)$.

(B) measurement additive noise model

The “input signal” is $p(n)$.In each model, $x(n)$ and $y(n)$ are “known”, and $u(n)$, $v(n)$, and $w(n)$ are *not*.In definition, the two models are **equivalent** under the relation $u(n) = -w(n)$.In practice, they are **different**:in (A), x and u would be typically *uncorrelated*;in (B), x and $w = -u$ are very much *correlated* (x is a function of u).Figure 11.1: Additive noise systems with *linear/non-linear* operator \mathbf{T}

And there is another problem here—we don't always have control over the input signal. Examples of this include earthquake and volcanic activity analysis.

An alternative to the sine-sweep input is *random sequence* input. All the techniques that follow in this chapter are of this type. A problem with using random sequences directly for estimating $\hat{H}(\omega)$ is that the estimate $\hat{H}(\omega)$ is itself also random. This is not what we want. We want an estimate that we can actually write down on paper or at least plot on paper.

A solution to this is to not use the random sequences directly to estimate $\hat{H}(\omega)$, but instead to first use the *expectation* operator E (Definition 1.1 page 3). The expectation operator takes a quantity X that is inherently “random” (with some probability distribution $p(x)$) and turns it into a deterministic “constant” EX .

The operator E is also used by the spectral density functions $\tilde{S}_{xx}(\omega)$ and $\tilde{S}_{xy}(\omega)$ (Definition 6.3 page 44). And $\tilde{S}_{xx}(\omega)$ and $\tilde{S}_{xy}(\omega)$ are what are typically used to calculate an estimate $\hat{H}(\omega)$.

11.2 Additive noise system models

Consider the additive noise systems illustrated in Figure 12.1 (page 119).

- ➊ The illustration on the left is suitable for modeling a communications system where x is the transmitted signal, y is the received signal, u and v are thermal noise, and the “transfer function” H is the communications channel (air, water, wires, etc.) that one wishes to estimate.
- ➋ The illustration on the right is suitable for modeling a testing system where p is an input test signal (from an industrial shaker or from a naturally occurring signal originating from geophysical activity), w is measurement noise, x is the measured input contaminated by noise, and H is the device under test (a piece of equipment, a building, or the entire earth).

Note that the two models are an equivalent system S under the relation $u = -w$. But although one might expect such a sign difference to wreak mathematical havoc in resulting equations, this

(1986) pages 125–140 (3.7 USE OF DIFFERENT EXCITATION TYPES)



is simply not the case here because

$$\tilde{S}_{ww} = \tilde{\mathbf{F}}\mathbf{E}[\mathbf{w}(m)\mathbf{w}^*(0)] = \tilde{\mathbf{F}}\mathbf{E}[(-\mathbf{u}(m))(-\mathbf{u}^*(0))] = \tilde{\mathbf{F}}\mathbf{E}[(\mathbf{u}(m))(\mathbf{u}^*(0))] = \tilde{S}_{uu}$$

So the sign difference is not that big of a difference after all. But there are some key differences in practice:

- In the communications model (on the left), the “input signal” is $x(n)$ and the frequency-domain input *signal-to-noise ratio (SNR)* is $\tilde{S}_{xx}(\omega)/\tilde{S}_{uu}(\omega)$. In the measurement model (on the right), the “input signal” is $p(n)$ and the frequency-domain input *signal-to-noise ratio (SNR)* is $\tilde{S}_{pp}(\omega)/\tilde{S}_{ww}(\omega) = \tilde{S}_{pp}(\omega)/\tilde{S}_{uu}(\omega)$.
- On the left, x and u would be typically *uncorrelated*; on the right, x and $w = -u$ are very much *correlated* (x is a function of u).

11.3 Transfer function estimate definitions and interpretation

As a first attempt at estimating the transfer function \mathbf{H} of \mathbf{S} , or at least the magnitude squared of \mathbf{H} , we might assume \mathbf{H} to be *LTI*, take a cue from the relation $\tilde{S}_{yy} = \tilde{S}_{xx}|\tilde{\mathbf{H}}|^2$ of Corollary 5.3 (page 37), and arrive at a function called “*transmissibility*” (next definition).

Definition 11.1. ⁵ Let \mathbf{S} be a system with input $x(n)$ and output $y(n)$.


DEF **transmissibility** $\tilde{\tau}_{xy}(\omega)$ is defined as $\tilde{\tau}_{xy}(\omega) \triangleq \sqrt{\frac{\tilde{S}_{yy}(\omega)}{\tilde{S}_{xx}(\omega)}}$

Transmissibility is in essence the ratio of “*spectral power*” (Remark 6.1 page 44) output to *spectral power* input. Note that it is a real-valued function (because \tilde{S}_{xx} and \tilde{S}_{yy} are real-valued). We might suspect that we could attain better estimates of \mathbf{H} by allowing the estimates to be complex-valued. And in fact, all the remaining estimates in this section are in general complex-valued.

And so to start (again), and in the very special (a.k.a unrealistic) case of \mathbf{S} having *zero measurement noise (zero measurement error)* ($v = u = w = 0$), $\mathbf{h}(n)$ being *linear time invariant (LTI)*, and input $x(n)$ being *wide sense stationary*...then we can determine (a.k.a “identify”) $\mathbf{h}(n)$ or $\tilde{\mathbf{H}}(\omega)$ exactly by $\tilde{\mathbf{H}}(\omega) = \tilde{S}_{yx}(\omega)/\tilde{S}_{xx}(\omega)$ (Corollary 5.3 page 37).

However, in practical situations, there is measurement noise/error. Examples may include “road noise” from a test being performed in a moving vehicle or *quantization noise* from an *analog-to-digital converter (ADC)*.

If the measurement error is at the output only (and under the assumptions of *LTI* and *WSS*) then $\hat{\mathbf{H}}_1$ (next definition) is the ideal estimator in the sense that $\hat{\mathbf{H}}_1 = \tilde{\mathbf{H}}$ (Corollary 11.4 page 113).



Definition 11.2. ⁶ Let \mathbf{S} be a system with input $x(n)$ and output $y(n)$.

⁵ Bendat and Piersol (2010) page 469 $\langle |H(f)| = [G_{yy}(f)/G_{xx}(f)]^{1/2} \rangle$, Yan and Ren (2012) page 204 $\langle (1) [G_{YY}(s)] = [H(s)][G_{FF}(s)][H^*(s)]^T \rangle$, Goldman (1999) page 179 \langle Transmissibility ... $H'_{ab} = G_{bb}/G_{aa}$ (note: differs by $\sqrt{\cdot}$ from Bendat and Piersol), Zhang et al. (2016), Zhou and Wahab (2018) page 824, https://link.springer.com/chapter/10.1007/978-3-319-54109-9_4

⁶ Bendat and Piersol (1993) pages 106–109 \langle 5.1.1 Optimality of Calculations \rangle , Bendat and Piersol (2010) page 185 $\langle H_1(f) = G_{xy}(f)/G_{xx}(f) \rangle$ (6.37), Shin and Hammond (2008) page 293 $\langle H_1(f) = \tilde{S}_{xy}(f)/\tilde{S}_{xx}(f) \rangle$ (9.63); which dif-

D E F The Least Squares transfer function estimate $\hat{H}_1(\omega)$ of S is defined as $\hat{H}_1(\omega) \triangleq \frac{\tilde{S}_{yx}(\omega)}{\tilde{S}_{xx}(\omega)}$

The estimator \hat{H}_1 is a good start. However in the early 1980s, L. D. Mitchell pointed out that in the presence of input noise, \hat{H}_1 is far from ideal in that it is *biased* with respect to \tilde{H} ; in fact, \hat{H}_1 *under estimates* \tilde{H} (Corollary 11.4 page 113). Mitchell proposed a new estimator \hat{H}_2 (next definition).

This estimator has the special property that when there is input noise but no output noise (and under LTI, WSS, and *uncorrelated* assumptions), then it is ideal in the sense that $\hat{H}_2(\omega) = \tilde{H}(\omega)$ (Corollary 11.4 page 113).

Note also that in the case of both no input and no output noise, then $\hat{H}_1 = \hat{H}_2$ (Corollary 5.3 page 37).

Definition 11.3. ⁷ Let S be a system with input $x(n)$ and output $y(n)$.

D E F The Inverse Method transfer function estimate $\hat{H}_2(\omega)$ of S is defined as $\hat{H}_2(\omega) \triangleq \frac{\tilde{S}_{yy}(\omega)}{\tilde{S}_{xy}(\omega)}$

Mitchell's \hat{H}_2 contribution "generated a flurry of activity"⁸ and soon more \tilde{H} estimators appeared. So far we have

• \hat{H}_1 which is ideal when there is no input noise but *under estimates* \tilde{H} when there is (Corollary 11.4 page 113)

• \hat{H}_2 which is ideal when there is no output noise but *over estimates* \tilde{H} when there is (Corollary 11.4 page 113).

But what about estimators for when there is noise on both input and output? Armed with two estimators that between them account for both input and output noise, an "ad hoc" solution might be to somehow take mean values of \hat{H}_1 and \hat{H}_2 to induce new estimators—this approach summarizes the next three definitions. An arguably more mature approach is to find estimators that are optimal with respect to least squares measures—and this approach summarizes Definition 11.9 – Definition 11.7 (page 99).

Definition 11.4. Let S be a system with input $x(n)$ and output $y(n)$.

D E F The Arithmetic Mean transfer function estimate $\hat{H}_{am}(\omega)$ of S is defined as

$$\hat{H}_{am}(\omega) \triangleq \frac{|\tilde{S}_{xy}(\omega)|^2 + \tilde{S}_{xx}(\omega)\tilde{S}_{yy}(\omega)}{2\tilde{S}_{xx}(\omega)\tilde{S}_{xy}(\omega)}$$

fers from Definition 11.2, but see APPENDIX ?? page ??),  Bendat (1978)cited by Cobb(1988)—variance estimate for \hat{H}_1 ,  Allemang et al. (1979) (cited by Shin(2008)),  Leuridan et al. (1986) page 910 (Least Squares Technique; (8) $[G_{xx}](H) = [G_{xy}]$),  Abom (1986)cited by Cobb(1988)—variance estimate for \hat{H}_1 ,  Allemang et al. (1987) pages 54–55 (5.3.1 H_1 Technique; $[H] = [G_{XF}][G_{FF}]^{-1}$ (11)),  Cobb (1988) page 2 (${}^1\hat{H}(f) = \hat{G}_{yx}(f)/\hat{G}_{xx}(f)$ (1)),  Goyder (1984) page 438 ($H(i\omega) = S_{qp}/S_{pp}$ (3)),  Pintelon and Schoukens (2012) page 233 ($\hat{G}(\Omega_k) = S_{yu}(j\omega_k)S_{uu}^{-1}(j\omega_k)$ (7-30)),  White et al. (2006) page 678 ($H_1(f) = \hat{S}_{x_my_m}(f)/\hat{S}_{x_mx_m}(f)$ (1) which differs by conjugate, references Bendat and Piersol),

⁷  Shin and Hammond (2008) page 293 ($H_2(f) = \tilde{S}_{yy}(f)/\tilde{S}_{yx}(f)$ (9.65); which differs from Definition 11.3, but see APPENDIX ?? page ??),  Bendat and Piersol (2010) page 186 ($H_2(f) = G_{yy}(f)/G_{yx}(f)$ (6.42)),  Mitchell (1980) (cited by Cobb(1988)),  Mitchell (1982) page 278 ("Define what will be called an inverse method for calculation of a FRF as..."; $H_2(f) = G_{yy}/G_{yx}$ (6); Note this differs with Definition 11.3 by a conjugate, but note that Mitchell seems to follow Bendat (see his [3] and [4]), which would explain this difference (APPENDIX ?? page ??)),  Cobb (1988) page 3 (${}^2\hat{H}(f) = \hat{G}_{yy}(f)/\hat{G}_{xy}(f)$ (1)),  White et al. (2006) page 678 ($H_2(f) = \hat{S}_{y_my_m}(f)/\hat{S}_{y_mx_m}(f)$ (2) which differs by conjugate, references Bendat and Piersol)

⁸  Cobb (1988) page 3

Proposition 11.1. ⁹ Let \mathbf{S} be a system with input $x(n)$ and output $y(n)$.

P R P
$$\hat{H}_{\text{am}}(\omega) = \frac{\hat{H}_1(\omega) + \hat{H}_2(\omega)}{2} \quad (\text{arithmetic mean of } \hat{H}_1 \text{ and } \hat{H}_2)$$

PROOF:

$$\begin{aligned} \hat{H}_{\text{am}}(\omega) &\triangleq \frac{|\tilde{S}_{xy}(\omega)|^2 + \tilde{S}_{xx}(\omega)\tilde{S}_{yy}(\omega)}{2\tilde{S}_{xx}(\omega)\tilde{S}_{xy}(\omega)} \quad \text{by definition of } \hat{H}_{\text{am}} \quad (\text{Definition 11.4 page 96}) \\ &= \frac{\tilde{S}_{xy}(\omega)\tilde{S}_{xy}^*(\omega)}{2\tilde{S}_{xx}(\omega)\tilde{S}_{xy}(\omega)} + \frac{\tilde{S}_{xx}(\omega)\tilde{S}_{yy}(\omega)}{2\tilde{S}_{xx}(\omega)\tilde{S}_{xy}(\omega)} = \frac{\frac{\tilde{S}_{xy}^*(\omega)}{\tilde{S}_{xx}(\omega)} + \frac{\tilde{S}_{yy}(\omega)}{\tilde{S}_{xy}(\omega)}}{2} \\ &= \frac{\hat{H}_1(\omega) + \hat{H}_2(\omega)}{2} \quad \text{by definitions of } \hat{H}_1 \text{ and } \hat{H}_2 \quad (\text{Definition 11.2 page 95, Definition 11.3 page 96}) \end{aligned}$$



Definition 11.5. Let \mathbf{S} be a system with input $x(n)$ and output $y(n)$.

The Geometric mean transfer function estimate $\hat{H}_{\text{gm}}(\omega)$ of \mathbf{S} is defined as

D E F
$$\hat{H}_{\text{gm}}(\omega) \triangleq \sqrt{\frac{\tilde{S}_{xy}^*(\omega)}{|\tilde{S}_{xy}(\omega)|}} \sqrt{\frac{\tilde{S}_{yy}(\omega)}{\tilde{S}_{xx}(\omega)}}$$

Proposition 11.2. ¹⁰ Let \mathbf{S} be a system with input $x(n)$ and output $y(n)$.

P R P
$$\pm \hat{H}_{\text{gm}}(\omega) = \sqrt{\hat{H}_1(\omega)\hat{H}_2(\omega)} \quad (\text{geometric mean of } \hat{H}_1 \text{ and } \hat{H}_2)$$

PROOF:

$$\begin{aligned} \pm \hat{H}_{\text{gm}}(\omega) &\triangleq \pm \sqrt{\frac{\tilde{S}_{xy}^*(\omega)}{|\tilde{S}_{xy}(\omega)|}} \sqrt{\frac{\tilde{S}_{yy}(\omega)}{\tilde{S}_{xx}(\omega)}} \quad \text{by definition of } \hat{H}_{\text{gm}} \quad (\text{Definition 11.5 page 97}) \\ &= \sqrt{\frac{[\tilde{S}_{xy}^*(\omega)]^2 \tilde{S}_{yy}(\omega)}{|\tilde{S}_{xy}(\omega)|^2 \tilde{S}_{xx}(\omega)}} = \sqrt{\frac{\tilde{S}_{xy}^*(\omega) \tilde{S}_{yy}(\omega)}{\tilde{S}_{xy}(\omega) \tilde{S}_{xx}(\omega)}} = \sqrt{\frac{\tilde{S}_{xy}^*(\omega) \tilde{S}_{yy}(\omega)}{\tilde{S}_{xx}(\omega) \tilde{S}_{xy}(\omega)}} \\ &= \sqrt{\hat{H}_1(\omega)\hat{H}_2(\omega)} \quad \text{by definitions of } \hat{H}_1 \text{ and } \hat{H}_2 \quad (\text{Definition 11.2 page 95, Definition 11.3 page 96}) \\ &= \text{Geometric mean of } \hat{H}_1(\omega) \text{ and } \hat{H}_2(\omega) \end{aligned}$$

Note that for a complex number $z \triangleq |z|e^{i\phi}$, \sqrt{z} has two solutions:¹¹

$$\sqrt{z} = \sqrt{|z|e^{i\phi}} = \{z_1, z_2\} = \left\{ \sqrt{|z|}e^{i(\phi/2)}, \sqrt{|z|}e^{i(\phi/2+\pi)} \right\} = \pm \sqrt{|z|}e^{i(\phi/2)}$$

because $z_1^2 = z$ and $z_2^2 = z$.



Note that the geometric mean estimator (Definition 11.5 page 97) and transmissibility (Definition 11.1 page 95) are closely related (next).

⁹ Mitchell (1982) page 279 (“Frequency Response Calculation: The Average Method”), Zheng et al. (2002) page 918 (“1.3 Arithmetic Mean Estimator H_3 ”)



¹⁰ Zheng et al. (2002) page 918 (“1.4 Geometric Mean Estimator H_4 ”)

¹¹ Many many thanks to Ben Cleveland for his help with this!!!

Proposition 11.3. Let $\phi(\omega)$ be the PHASE of $\tilde{S}_{xy}(\omega)$ such that $\tilde{S}_{xy}(\omega) \triangleq |\tilde{S}_{xy}(\omega)|e^{-i\phi(\omega)}$

P R P	$\hat{H}_{gm}(\omega) = \tilde{T}_{xy}(\omega) e^{-i\phi(\omega)} \quad \left(\begin{array}{l} \hat{H}_{gm}(\omega) = \tilde{T}_{xy}(\omega) \text{ is the MAGNITUDE of } \hat{H}_{gm}(\omega) \text{ and} \\ \angle \hat{H}_{gm}(\omega) = -\angle \tilde{S}_{xy}(\omega) \text{ is the PHASE of } \hat{H}_{gm}(\omega) \end{array} \right)$
-------------	--

PROOF: Let $\phi(\omega)$ be the *phase* of

$$\begin{aligned}
 \hat{H}_{gm}(\omega) &\triangleq \sqrt{\hat{H}_1(\omega)\hat{H}_2(\omega)} && \text{by definition of } \hat{H}_{gm} && \text{(Definition 11.5 page 97)} \\
 &\triangleq \sqrt{\frac{\tilde{S}_{xy}^*(\omega)\tilde{S}_{yy}(\omega)}{\tilde{S}_{xx}(\omega)\tilde{S}_{xy}(\omega)}} && \text{by definitions of } \hat{H}_1 \text{ and } \hat{H}_2 && \text{(Definition 11.2 page 95, Definition 11.3 page 96)} \\
 &= \sqrt{\frac{\tilde{S}_{yy}(\omega)\tilde{S}_{xy}^*(\omega)}{\tilde{S}_{xx}(\omega)\tilde{S}_{xy}(\omega)}} \\
 &= \tilde{T}_{xy}(\omega) \sqrt{\frac{\tilde{S}_{xy}^*(\omega)}{\tilde{S}_{xy}(\omega)}} && \text{by definition of } \tilde{T}_{xy} && \text{(Definition 11.1 page 95)} \\
 &= \tilde{T}_{xy}(\omega) \sqrt{\frac{|\tilde{S}_{xy}(\omega)|e^{-i\phi(\omega)}}{|\tilde{S}_{xy}(\omega)|e^{i\phi(\omega)}}} && \text{where } \tilde{S}_{xy}(\omega) \triangleq |\tilde{S}_{xy}(\omega)|e^{-i\phi(\omega)} \\
 &= \tilde{T}_{xy}(\omega) \sqrt{e^{-i2\phi(\omega)}} \\
 &= \tilde{T}_{xy}(\omega) e^{-i\phi(\omega)}
 \end{aligned}$$

Remark 11.1. Transmissibility \tilde{T}_{xy} is a kind of “black sheep” of the system identification function family. All the other members of this family ($\hat{H}_1, \hat{H}_2, \hat{H}_v, \hat{H}_s$) are *complex-valued*, but \tilde{T}_{xy} is only *real-valued*—a seemingly ordinary Joe born into a super-hero family. But Proposition 11.3 suggests that \tilde{T}_{xy} is not simply a “black sheep”, but rather a “dark horse” with abilities that can easily be unleashed by slight redefinition. In particular, Proposition 11.3 demonstrates that \tilde{T}_{xy} is the *magnitude* of the geometric mean of \hat{H}_1 and \hat{H}_2 . We can thus justifiably define a **complex transmissibility** function as \hat{H}_{gm} ...and the magnitude of this *complex transmissibility* function is the *ordinary transmissibility* function of Definition 11.1 (page 95).

R E M	complex transmissibility $\tilde{T}'_{xy}(\omega) \triangleq \hat{H}_{gm}(\omega)$
-------------	---

Definition 11.6. Let S be a SYSTEM with input $x(n)$ and output $y(n)$.

The **Harmonic mean transfer function estimate** $\hat{H}_{hm}(\omega)$ of S is defined as

$$\hat{H}_{hm}(\omega) \triangleq \frac{2\tilde{S}_{yy}(\omega)\tilde{S}_{xy}^*(\omega)}{\tilde{S}_{xx}(\omega)\tilde{S}_{yy}(\omega) + |\tilde{S}_{xy}(\omega)|^2}$$

Proposition 11.4.¹² Let S be a SYSTEM with input $x(n)$ and output $y(n)$.

P R P	$\hat{H}_{hm}(\omega) = \frac{2}{\frac{1}{\hat{H}_1(\omega)} + \frac{1}{\hat{H}_2(\omega)}} \quad (\text{Harmonic mean of } \hat{H}_1 \text{ and } \hat{H}_2)$
-------------	--

¹² Carne and Dohrmann (2006) ($H_C = [H_A^{-1} + H_B^{-1}]^{-1}$)

PROOF:

$$\begin{aligned}
 \hat{H}_{\text{hm}}(\omega) &\triangleq \frac{2\tilde{S}_{yy}(\omega)\tilde{S}_{xy}^*(\omega)}{\tilde{S}_{xx}(\omega)\tilde{S}_{yy}(\omega) + |\tilde{S}_{xy}(\omega)|^2} \quad \text{by definition of } \hat{H}_{\text{hm}} \quad (\text{Definition 11.6 page 98}) \\
 &= \frac{2}{\frac{\tilde{S}_{xx}(\omega)\tilde{S}_{yy}(\omega) + |\tilde{S}_{xy}(\omega)|^2}{\tilde{S}_{yy}(\omega)\tilde{S}_{xy}^*(\omega)}} = \frac{2}{\frac{\tilde{S}_{xx}(\omega)}{\tilde{S}_{xy}^*(\omega)} + \frac{\tilde{S}_{xy}(\omega)}{\tilde{S}_{yy}(\omega)}} \\
 &= \frac{2}{\frac{1}{\hat{H}_1(\omega)} + \frac{1}{\hat{H}_2(\omega)}} \quad \text{by definitions of } \hat{H}_1 \text{ and } \hat{H}_2 \quad (\text{Definition 11.2 page 95, Definition 11.3 page 96}) \\
 &= \text{Harmonic mean of } \hat{H}_1(\omega) \text{ and } \hat{H}_2(\omega)
 \end{aligned}$$



A bit of review reveals \hat{H}_1 at the low end of the estimation problem, \hat{H}_2 at the high end, and \hat{H}_{hm} , \hat{H}_{gm} , and \hat{H}_{am} somewhere between. But these three “between” estimates are not shown to be optimal in any sense—they are just conceptually interesting. What we might really like is a family of estimators that

- ☛ include \hat{H}_1 and \hat{H}_2 as limiting cases
- ☛ include the between cases
- ☛ are optimal in some sense

The estimator $\hat{H}_\kappa(\omega; \kappa)$ is one such estimator (next definition) that

- ☛ has \hat{H}_1 and \hat{H}_2 as limiting cases (Theorem 11.1 page 101),
- ☛ is optimal in the least squares sense (Theorem 11.6 page 114), and
- ☛ allows for a system designer to specify an output-input spectral noise ratio $\kappa(\omega)$ that can vary with frequency ω .

Moreover, $\hat{H}_\kappa(\omega)$ includes some special cases:

- ☛ In the case of constant κ , \hat{H}_κ simplifies to the *Scaling transfer function estimate* \hat{H}_s (Definition 11.8 page 99).
- ☛ In the case of $\kappa = 1$, \hat{H}_κ and \hat{H}_s simplify to the *Total least squares transfer function estimate* \hat{H}_v (Definition 11.9 page 100).

Definition 11.7. ¹³ Let S be a system with input $x(n)$ and output $y(n)$.

D E F The transfer function estimate $\hat{H}_\kappa(\omega; \kappa)$ with scaling function $\kappa(\omega)$ is defined as

$$\hat{H}_\kappa(\omega; \kappa) \triangleq \frac{\tilde{S}_{yy}(\omega) - \kappa(\omega)\tilde{S}_{xx}(\omega) + \sqrt{[\tilde{S}_{yy}(\omega) - \kappa(\omega)\tilde{S}_{xx}(\omega)]^2 + 4\kappa(\omega)|\tilde{S}_{xy}(\omega)|^2}}{2\tilde{S}_{xy}(\omega)}$$

Definition 11.8. ¹⁴ Let S be a system with input $x(n)$ and output $y(n)$.

D E F The Scaling transfer function estimate $\hat{H}_s(\omega; s)$ of S with scaling parameter $s \in [0 : \infty)$ is defined as $\hat{H}_s(\omega; s) \triangleq \hat{H}_\kappa(\omega; \kappa)$ with $\kappa(\omega) \triangleq s^2$

¹³ ☐ White et al. (2006) page 679 ((6)), ☐ Shin and Hammond (2008) page 293 ((9.67))

¹⁴ ☐ Shin and Hammond (2008) page 293 ((9.67) with $\kappa(\omega) = s^2$), ☐ White et al. (2006) page 679 ((6) with $\kappa(\omega) = s^2$), ☐ Leclerc et al. (2014) ((10) $\kappa(f) = 1/s^2$ and x and y swapped), ☐ Wicks and Vold (1986) page 898 (has additional s in denominator), ☐ Zheng et al. (2002) page 918 ((10), seems to differ)

Definition 11.9. ¹⁵ Let \mathbf{S} be a system with input $x(n)$ and output $y(n)$.

D E F The Total Least Squares transfer function estimate $\hat{H}_v(\omega)$ of \mathbf{S} is defined as
 $\hat{H}_v(\omega) \triangleq \hat{H}_k(\omega; \kappa) \quad \text{with } \kappa(\omega) = 1$

The previous estimators all assumed two signals: an input $x(n)$ and an output $y(n)$. However, in many practical systems, there is a third signal that is “driving” the system. In 1984 Goyder proposed an estimator (next definition) that is based on three signals.

Definition 11.10 (Three channel estimate). ¹⁶ Let \mathbf{S} be a system with input $x(n)$, output $y(n)$, and a driver $p(n)$.

D E F The transfer function estimate $\hat{H}_c(\omega)$ is defined as
 $\hat{H}_c(\omega) \triangleq \frac{\tilde{S}_{py}(\omega)}{\tilde{S}_{px}(\omega)}$

11.4 Estimator relationships

Lemma 11.1.

L E M	$\frac{d}{dp} \left[\tilde{S}_{yy} - p\tilde{S}_{xx} + \sqrt{(\tilde{S}_{yy} - p\tilde{S}_{xx})^2 + 4p \tilde{S}_{xy} ^2} \right] = \frac{\tilde{S}_{xx}(p\tilde{S}_{xx} - \tilde{S}_{yy}) + 2 \tilde{S}_{xy} ^2 - \tilde{S}_{xx}\sqrt{(p\tilde{S}_{xx} - \tilde{S}_{yy})^2 + 4p \tilde{S}_{xy} ^2}}{\sqrt{(p\tilde{S}_{xx} - \tilde{S}_{yy})^2 + 4p \tilde{S}_{xy} ^2}}$
	$\frac{d}{dp} \left[p\tilde{S}_{yy} - \tilde{S}_{xx} + \sqrt{(p\tilde{S}_{yy} - \tilde{S}_{xx})^2 + 4p \tilde{S}_{xy} ^2} \right] = \frac{\tilde{S}_{yy}(p\tilde{S}_{yy} - \tilde{S}_{xx}) + 2 \tilde{S}_{xy} ^2 + \tilde{S}_{yy}\sqrt{(p\tilde{S}_{yy} - \tilde{S}_{xx})^2 + 4p \tilde{S}_{xy} ^2}}{\sqrt{(p\tilde{S}_{yy} - \tilde{S}_{xx})^2 + 4p \tilde{S}_{xy} ^2}}$

PROOF:

$$\begin{aligned} \frac{d}{dp} \left[\tilde{S}_{yy} - p\tilde{S}_{xx} + \sqrt{(\tilde{S}_{yy} - p\tilde{S}_{xx})^2 + 4p|\tilde{S}_{xy}|^2} \right] &= -\tilde{S}_{xx} + \frac{-2\tilde{S}_{xx}(\tilde{S}_{yy} - p\tilde{S}_{xx}) + 4|\tilde{S}_{xy}|^2}{2\sqrt{(\tilde{S}_{yy} - p\tilde{S}_{xx})^2 + 4p|\tilde{S}_{xy}|^2}} \\ &= \frac{4|\tilde{S}_{xy}|^2 - 2\tilde{S}_{xx}(\tilde{S}_{yy} - p\tilde{S}_{xx}) - 2\tilde{S}_{xx}\sqrt{(\tilde{S}_{yy} - p\tilde{S}_{xx})^2 + 4p|\tilde{S}_{xy}|^2}}{2\sqrt{(\tilde{S}_{yy} - p\tilde{S}_{xx})^2 + 4p|\tilde{S}_{xy}|^2}} \\ &= \frac{\tilde{S}_{xx}(p\tilde{S}_{xx} - \tilde{S}_{yy}) + 2|\tilde{S}_{xy}|^2 - \tilde{S}_{xx}\sqrt{(p\tilde{S}_{xx} - \tilde{S}_{yy})^2 + 4p|\tilde{S}_{xy}|^2}}{\sqrt{(p\tilde{S}_{xx} - \tilde{S}_{yy})^2 + 4p|\tilde{S}_{xy}|^2}} \end{aligned}$$

$$\begin{aligned} \frac{d}{dp} \left[p\tilde{S}_{yy} - \tilde{S}_{xx} + \sqrt{(p\tilde{S}_{yy} - \tilde{S}_{xx})^2 + 4p|\tilde{S}_{xy}|^2} \right] &= \tilde{S}_{yy} + \frac{2\tilde{S}_{yy}(p\tilde{S}_{yy} - \tilde{S}_{xx}) + 4|\tilde{S}_{xy}|^2}{2\sqrt{(p\tilde{S}_{yy} - \tilde{S}_{xx})^2 + 4p|\tilde{S}_{xy}|^2}} \\ &= \frac{4|\tilde{S}_{xy}|^2 + 2\tilde{S}_{yy}(p\tilde{S}_{yy} - \tilde{S}_{xx}) + 2\tilde{S}_{yy}\sqrt{(p\tilde{S}_{yy} - \tilde{S}_{xx})^2 + 4p|\tilde{S}_{xy}|^2}}{2\sqrt{(p\tilde{S}_{yy} - \tilde{S}_{xx})^2 + 4p|\tilde{S}_{xy}|^2}} \end{aligned}$$

¹⁵ White et al. (2006) page 679 ⟨(6)⟩, Shin and Hammond (2008) page 294 ⟨(9.69)⟩

¹⁶ Shin and Hammond (2008) page 297 ⟨ $H_3(f) = S_{ry}(f)/S_{rx}(f)$ (9.78)⟩, Cobb (1988) page 4 ⟨ $c \hat{H}(f) = \hat{G}_{ys}(f)/\hat{G}_{xs}(f)$ (1.4)⟩, Goyder (1984) page 440 ⟨ $H(i\omega) = S_{qz}/S_{pz}$ (5)⟩, Cobb and Mitchell (1990) page 450 ⟨(1)⟩, Pintelon and Schoukens (2012) page 241 ⟨ $\hat{G}(\Omega_k) = \hat{G}_{ry}(\Omega_k)\hat{G}_{ru}^{-1}(\Omega_k)$ (7-49)⟩

$$= \frac{\tilde{S}_{yy}(p\tilde{S}_{yy} - \tilde{S}_{xx}) + 2|\tilde{S}_{xy}|^2 + \tilde{S}_{yy}\sqrt{(p\tilde{S}_{yy} - \tilde{S}_{xx})^2 + 4p|\tilde{S}_{xy}|^2}}{\sqrt{(p\tilde{S}_{yy} - \tilde{S}_{xx})^2 + 4p|\tilde{S}_{xy}|^2}}$$



**Lemma 11.2.**

LEM	$\tilde{S}_{yy} - p\tilde{S}_{xx} + \sqrt{(\tilde{S}_{yy} - p\tilde{S}_{xx})^2 + 4p \tilde{S}_{xy} ^2} \geq 0$
	$p\tilde{S}_{yy} - \tilde{S}_{xx} + \sqrt{(p\tilde{S}_{yy} - \tilde{S}_{xx})^2 + 4p \tilde{S}_{xy} ^2} \geq 0$

PROOF:

$$\begin{aligned} & \tilde{S}_{yy} - p\tilde{S}_{xx} + \sqrt{(\tilde{S}_{yy} - p\tilde{S}_{xx})^2 + 4p|\tilde{S}_{xy}|^2} \geq 0 \\ \Leftrightarrow & \sqrt{(p\tilde{S}_{xx} - \tilde{S}_{yy})^2 + 4p|\tilde{S}_{xy}|^2} \geq p\tilde{S}_{xx} - \tilde{S}_{yy} \\ \Leftrightarrow & (p\tilde{S}_{xx} - \tilde{S}_{yy})^2 + 4p|\tilde{S}_{xy}|^2 \geq (p\tilde{S}_{xx} - \tilde{S}_{yy})^2 \\ \Leftrightarrow & 4p|\tilde{S}_{xy}|^2 \geq 0 \\ \Leftrightarrow & |\tilde{S}_{xy}| \geq 0 \end{aligned}$$

$$\begin{aligned} & p\tilde{S}_{yy} - \tilde{S}_{xx} + \sqrt{(p\tilde{S}_{yy} - \tilde{S}_{xx})^2 + 4p|\tilde{S}_{xy}|^2} \geq 0 \\ \Leftrightarrow & \sqrt{(\tilde{S}_{xx} - p\tilde{S}_{yy})^2 + 4p|\tilde{S}_{xy}|^2} \geq \tilde{S}_{xx} - p\tilde{S}_{yy} \\ \Leftrightarrow & (\tilde{S}_{xx} - p\tilde{S}_{yy})^2 + 4p|\tilde{S}_{xy}|^2 \geq (\tilde{S}_{xx} - p\tilde{S}_{yy})^2 \\ \Leftrightarrow & 4p|\tilde{S}_{xy}|^2 \geq 0 \\ \Leftrightarrow & |\tilde{S}_{xy}| \geq 0 \end{aligned}$$

 \hat{H}_s as a function of $p \triangleq s^2$  $\frac{d}{dp}\hat{H}_s$ where $p \triangleq s^2$ Figure 11.2: \hat{H}_s with $\tilde{S}_{xx} = \tilde{S}_{yy} = 1$ and $\tilde{S}_{xy} = \frac{1}{2}$ **Theorem 11.1.** Let \hat{H}_s be defined as in Definition 11.8 (page 99).

THM	$\{s_1 < s_2\} \implies \hat{H}_s(\omega; s_2) \leq \hat{H}_s(\omega; s_1)$ (\hat{H}_s is monotonically decreasing in s)
	$ \hat{H}_1(\omega) \leq \hat{H}_s(\omega; s) \leq \hat{H}_2(\omega) $
	$\hat{H}_s(\omega; s) _{s=0} = \hat{H}_2(\omega)$
	$\hat{H}_s(\omega; s) _{s=1} = \hat{H}_v(\omega)$
	$\lim_{s \rightarrow \infty} \hat{H}_s(\omega; s) = \hat{H}_1(\omega)$



PROOF: I. Proofs for equalities:

$$\begin{aligned}
 \hat{H}_s(\omega; s) \Big|_{s=0} &\triangleq \frac{\tilde{S}_{yy} - s^2 \tilde{S}_{xx} + \sqrt{[\tilde{S}_{yy} - s^2 \tilde{S}_{xx}]^2 + 4s^2 |\tilde{S}_{xy}|^2}}{2\tilde{S}_{xy}} \Big|_{s=0} && \text{by def. of } \hat{H}_s && \text{(Definition 11.8 page 99)} \\
 &= \frac{\tilde{S}_{yy} - 0 + \sqrt{[\tilde{S}_{yy} - 0]^2 + 0}}{2\tilde{S}_{xy}} = \frac{\tilde{S}_{yy}}{\tilde{S}_{xy}} && && \\
 &\triangleq \hat{H}_2 && \text{by def. of } \hat{H}_2 && \text{(Definition 11.3 page 96)} \\
 \hat{H}_s(\omega; s) \Big|_{s=1} &\triangleq \frac{\tilde{S}_{yy} - s^2 \tilde{S}_{xx} + \sqrt{[\tilde{S}_{yy} - s^2 \tilde{S}_{xx}]^2 + 4s^2 |\tilde{S}_{xy}|^2}}{2\tilde{S}_{xy}} \Big|_{s=1} && \text{by def. of } \hat{H}_s && \text{(Definition 11.8 page 99)} \\
 &= \frac{\tilde{S}_{yy} - \tilde{S}_{xx} + \sqrt{[\tilde{S}_{yy} - \tilde{S}_{xx}]^2 + 4|\tilde{S}_{xy}|^2}}{2\tilde{S}_{xy}} && && \\
 &\triangleq \hat{H}_v && \text{by def. of } \hat{H}_v && \text{(Definition 11.9 page 100)} \\
 \lim_{s \rightarrow \infty} \hat{H}_s(\omega; s) &\triangleq \lim_{s \rightarrow \infty} \frac{\tilde{S}_{yy} - s^2 \tilde{S}_{xx} + \sqrt{[\tilde{S}_{yy} - s^2 \tilde{S}_{xx}]^2 + 4s^2 |\tilde{S}_{xy}|^2}}{2\tilde{S}_{xy}} && \text{by def. of } \hat{H}_s && \text{(Definition 11.8 page 99)} \\
 &\triangleq \lim_{p \rightarrow 0} \frac{\tilde{S}_{yy} - \frac{1}{p} \tilde{S}_{xx} + \sqrt{[\tilde{S}_{yy} - \frac{1}{p} \tilde{S}_{xx}]^2 + 4\frac{1}{p} |\tilde{S}_{xy}|^2}}{2\tilde{S}_{xy}} && && \\
 &= \lim_{p \rightarrow 0} \frac{p\tilde{S}_{yy} - \tilde{S}_{xx} + \sqrt{[p\tilde{S}_{yy} - \tilde{S}_{xx}]^2 + 4p |\tilde{S}_{xy}|^2}}{2p\tilde{S}_{xy}} && \text{where } p \triangleq \frac{1}{s^2} && \\
 &= \lim_{p \rightarrow 0} \frac{\frac{d}{dp} \left[p\tilde{S}_{yy} - \tilde{S}_{xx} + \sqrt{[p\tilde{S}_{yy} - \tilde{S}_{xx}]^2 + 4p |\tilde{S}_{xy}|^2} \right]}{\frac{d}{dp} [2p\tilde{S}_{xy}]} && \text{by l'Hôpital's rule} && \\
 &= \lim_{p \rightarrow 0} \frac{\tilde{S}_{yy}(p\tilde{S}_{yy} - \tilde{S}_{xx}) + 2|\tilde{S}_{xy}|^2 + \tilde{S}_{yy} \sqrt{(p\tilde{S}_{yy} - \tilde{S}_{xx})^2 + 4p |\tilde{S}_{xy}|^2}}{\sqrt{(p\tilde{S}_{yy} - \tilde{S}_{xx})^2 + 4p |\tilde{S}_{xy}|^2}} && && \\
 &= \lim_{p \rightarrow 0} \frac{\tilde{S}_{yy}(-\tilde{S}_{xx}) + 2|\tilde{S}_{xy}|^2 + \tilde{S}_{yy} \sqrt{(-\tilde{S}_{xx})^2}}{2\tilde{S}_{xy}} && \text{by Lemma 11.1 page 100} && \\
 &= \frac{2|\tilde{S}_{xy}|^2}{2\tilde{S}_{xx}\tilde{S}_{xy}} = \frac{\tilde{S}_{xy}^*}{\tilde{S}_{xx}} \triangleq \hat{H}_1 && \text{by def. of } \hat{H}_1 && \text{(Definition 11.2 page 95)}
 \end{aligned}$$

II. Proof for monotonicity:

1. Let $p \triangleq s^2$

2. lemma:

$$\begin{aligned}
 &\boxed{[2|\tilde{S}_{xy}|^2 + \tilde{S}_{xx}(p\tilde{S}_{xx} - \tilde{S}_{yy})]^2} \\
 &= 4|\tilde{S}_{xy}|^4 + 4\tilde{S}_{xx}|\tilde{S}_{xy}|^2(p\tilde{S}_{xx} - \tilde{S}_{yy}) + \tilde{S}_{xx}^2(p\tilde{S}_{xx} - \tilde{S}_{yy})^2
 \end{aligned}$$



$$\begin{aligned}
 & \leq 4|\tilde{S}_{xy}|^2\tilde{S}_{xx}\tilde{S}_{yy} + 4p\tilde{S}_{xx}|\tilde{S}_{xy}|^2\tilde{S}_{xx} - 4\tilde{S}_{xx}|\tilde{S}_{xy}|^2\tilde{S}_{yy} + \tilde{S}_{xx}^2(p\tilde{S}_{xx} - \tilde{S}_{yy})^2 \\
 & = 4\tilde{S}_{xx}\tilde{S}_{yy}|\tilde{S}_{xy}|^2 + 4p\tilde{S}_{xx}|\tilde{S}_{xy}|^2\tilde{S}_{xx} - 4\tilde{S}_{xx}\tilde{S}_{yy}|\tilde{S}_{xy}|^2 + \tilde{S}_{xx}^2(p\tilde{S}_{xx} - \tilde{S}_{yy})^2 \\
 & = \tilde{S}_{xx}^2[(p\tilde{S}_{xx} - \tilde{S}_{yy})^2 + 4p|\tilde{S}_{xy}|^2] \\
 & = \left[\tilde{S}_{xx}\sqrt{(p\tilde{S}_{xx} - \tilde{S}_{yy})^2 + 4p|\tilde{S}_{xy}|^2}\right]^2
 \end{aligned}
 \quad \left(\begin{array}{l} \text{by Cauchy Schwartz inequality} \\ (\text{Theorem I.2 page 234}) \end{array} \right)$$

3. lemma: $2|\tilde{S}_{xy}|^2 + \tilde{S}_{xx}(p\tilde{S}_{xx} - \tilde{S}_{yy}) \leq \tilde{S}_{xx}\sqrt{(p\tilde{S}_{xx} - \tilde{S}_{yy})^2 + 4p|\tilde{S}_{xy}|^2}$. Proof:

$$\begin{aligned}
 & 2|\tilde{S}_{xy}|^2 + \tilde{S}_{xx}(p\tilde{S}_{xx} - \tilde{S}_{yy}) \leq \tilde{S}_{xx}\sqrt{(p\tilde{S}_{xx} - \tilde{S}_{yy})^2 + 4p|\tilde{S}_{xy}|^2} \\
 \iff & [2|\tilde{S}_{xy}|^2 + \tilde{S}_{xx}(p\tilde{S}_{xx} - \tilde{S}_{yy})]^2 \leq \left[\tilde{S}_{xx}\sqrt{(p\tilde{S}_{xx} - \tilde{S}_{yy})^2 + 4p|\tilde{S}_{xy}|^2}\right]^2 \quad \left(\begin{array}{l} \text{because } f(x) \triangleq x^2 \text{ is} \\ \text{strictly monotonic increasing} \end{array} \right)
 \end{aligned}$$

The previous inequality is true by (2) lemma, so (3) lemma also true.

4. Proof that $\frac{d}{dp}|\hat{H}_s| \leq 0$:

$$\begin{aligned}
 \frac{d}{dp}|\hat{H}_s| & \triangleq \frac{d}{dp} \left| \frac{\tilde{S}_{yy} - s^2\tilde{S}_{xx} + \sqrt{(\tilde{S}_{yy} - s^2\tilde{S}_{xx})^2 + 4s^2|\tilde{S}_{xy}|^2}}{2\tilde{S}_{xy}} \right| \quad \text{by def. of } \hat{H}_s \text{ (Definition 11.8 page 99)} \\
 & \triangleq \frac{d}{dp} \left| \frac{\tilde{S}_{yy} - p\tilde{S}_{xx} + \sqrt{(\tilde{S}_{yy} - p\tilde{S}_{xx})^2 + 4p|\tilde{S}_{xy}|^2}}{2\tilde{S}_{xy}} \right| \quad \text{by definition of } p \text{ (item (1) page 102)} \\
 & = \frac{d}{dp} \left| \frac{\tilde{S}_{yy} - p\tilde{S}_{xx} + \sqrt{(\tilde{S}_{yy} - p\tilde{S}_{xx})^2 + 4p|\tilde{S}_{xy}|^2}}{|2\tilde{S}_{xy}|} \right| \\
 & = \frac{d}{dp} \left| \frac{\tilde{S}_{yy} - p\tilde{S}_{xx} + \sqrt{(\tilde{S}_{yy} - p\tilde{S}_{xx})^2 + 4p|\tilde{S}_{xy}|^2}}{2|\tilde{S}_{xy}|} \right| \quad \text{by Lemma 11.2 page 101} \\
 & = \frac{\tilde{S}_{xx}(p\tilde{S}_{xx} - \tilde{S}_{yy}) + 2|\tilde{S}_{xy}|^2 - \tilde{S}_{xx}\sqrt{(p\tilde{S}_{xx} - \tilde{S}_{yy})^2 + 4p|\tilde{S}_{xy}|^2}}{\sqrt{(\tilde{S}_{yy} - p\tilde{S}_{xx})^2 + 4p|\tilde{S}_{xy}|^2}} \quad \text{by Lemma 11.1 page 100} \\
 & = \frac{\tilde{S}_{xx}(p\tilde{S}_{xx} - \tilde{S}_{yy}) + 2|\tilde{S}_{xy}|^2 - \tilde{S}_{xx}\sqrt{(p\tilde{S}_{xx} - \tilde{S}_{yy})^2 + 4p|\tilde{S}_{xy}|^2}}{2|\tilde{S}_{xy}|\sqrt{(\tilde{S}_{yy} - p\tilde{S}_{xx})^2 + 4p|\tilde{S}_{xy}|^2}} \\
 & \leq 0 \quad \text{by (3) lemma}
 \end{aligned}$$

Theorem 11.2. Let S be a SYSTEM with input $x(n)$ and output $y(n)$.

T
H
M

$$|\hat{H}_1(\omega)| \leq |\hat{H}_{hm}(\omega)| \leq |\hat{H}_{gm}(\omega)| \leq |\hat{H}_{am}(\omega)| \leq |\hat{H}_2(\omega)|$$

PROOF:

1. lemma: $\hat{H}_1(\omega) \leq \hat{H}_2(\omega)$. Proof:

$$\begin{aligned}
 |\hat{H}_1| &\triangleq \left| \frac{\tilde{S}_{yx}(\omega)}{\tilde{S}_{xx}(\omega)} \right| && \text{by definition of } \hat{H}_1 && (\text{Definition 11.2 page 95}) \\
 &= \left| \frac{\langle y | x \rangle}{\|x\|^2} \right| = \frac{|\langle y | x \rangle|}{\|x\|^2} \\
 &\leq \frac{|\langle y | x \rangle|}{\|x\|^2} \left| \frac{\|x\| \|y\|}{\langle y | x \rangle} \right|^2 && \text{by Cauchy Schwartz inequality} && \text{Theorem I.2 page 234} \\
 &= \frac{\|y\|^2}{|\langle y | x \rangle|} = \left| \frac{\|y\|^2}{\langle x | y \rangle} \right| = \left| \frac{\tilde{S}_{yy}(\omega)}{\tilde{S}_{xy}(\omega)} \right| \\
 &= |\hat{H}_2| && \text{by definition of } \hat{H}_2 && (\text{Definition 11.3 page 96})
 \end{aligned}$$

2. remainder of the proof:

$$\begin{aligned}
 |\hat{H}_1(\omega)| &= \min \{ \hat{H}_1(\omega), \hat{H}_2(\omega) \} && \text{by (1) lemma} \\
 &\leq |\hat{H}_{hm}(\omega)| && \text{by Corollary L.1 page 274} && \text{with } \lambda_1 = \lambda_2 = \frac{1}{2} \\
 &\leq |\hat{H}_{gm}(\omega)| && \text{by Corollary L.1 page 274} && \text{with } \lambda_1 = \lambda_2 = \frac{1}{2} \\
 &\leq |\hat{H}_{am}(\omega)| && \text{by Corollary L.1 page 274} && \text{with } \lambda_1 = \lambda_2 = \frac{1}{2} \\
 &\leq \max \{ \hat{H}_1(\omega), \hat{H}_2(\omega) \} && \text{by Corollary L.1 page 274} && \text{with } \lambda_1 = \lambda_2 = \frac{1}{2} \\
 &= |\hat{H}_2(\omega)| && \text{by (1) lemma}
 \end{aligned}$$



Theorem 11.2 (page 103) compared the magnitudes of several transfer function estimates and demonstrated a simple *linear* relationship. What about phase? The phase of those estimates is even simpler than the magnitude, as demonstrated next.

Proposition 11.5 (Estimator phase). Let $z \triangleq |z|e^{i\phi}$ be a COMPLEX number in the set of complex numbers \mathbb{C} . Let $\angle z \triangleq \phi$ be the PHASE of z .

P R P	$ \begin{aligned} \angle \hat{H}_1(\omega) &= \angle \hat{H}_{hm}(\omega) = \angle \hat{H}_{gm}(\omega) = \angle \hat{H}_{am}(\omega) = \angle \hat{H}_2(\omega) = \angle \hat{H}_s(\omega) = \angle \hat{H}_v(\omega) = \angle \hat{H}_k(\omega) \\ &= \angle C_{xy}(\omega) \\ &= -\angle \tilde{S}_{xy}(\omega) \end{aligned} $
-------------	--

PROOF:

$$\begin{aligned}
 \angle \hat{H}_1 &\triangleq \angle \frac{\tilde{S}_{yx}}{\tilde{S}_{xx}} & = & = \angle \tilde{S}_{xy}^* = -\angle \tilde{S}_{xy} \\
 &\uparrow \text{(Definition 11.2 page 95)} & & \\
 \angle \hat{H}_{hm} &\triangleq \angle \frac{2\tilde{S}_{yy}\tilde{S}_{xy}^*}{\tilde{S}_{xx}\tilde{S}_{yy} + |\tilde{S}_{xy}|^2} & = & = \angle \tilde{S}_{xy}^* = -\angle \tilde{S}_{xy} \\
 &\uparrow \text{(Definition 11.6 page 98)} & & \\
 \angle \hat{H}_{gm} &\triangleq \angle \frac{\tilde{S}_{xy}^*}{|\tilde{S}_{xy}|} \sqrt{\frac{\tilde{S}_{yy}}{\tilde{S}_{xx}}} & = & = \angle \tilde{S}_{xy}^* = -\angle \tilde{S}_{xy} \\
 &\uparrow \text{(Definition 11.5 page 97)} & & \\
 \angle \hat{H}_{am} &\triangleq \angle \frac{|\tilde{S}_{xy}|^2 + \tilde{S}_{xx}\tilde{S}_{yy}}{2\tilde{S}_{xx}\tilde{S}_{xy}} & = \angle \frac{1}{\tilde{S}_{xy}} & = \angle \tilde{S}_{xy}^* = -\angle \tilde{S}_{xy} \\
 &\uparrow \text{(Definition 11.4 page 96)} & & \\
 \angle \hat{H}_2 &\triangleq \angle \frac{\tilde{S}_{yy}}{\tilde{S}_{xy}} & = \angle \frac{1}{\tilde{S}_{xy}} & = \angle \tilde{S}_{xy}^* = -\angle \tilde{S}_{xy} \\
 &\uparrow \text{(Definition 11.3 page 96)} & & \\
 \angle \hat{H}_s &\triangleq \angle \frac{\tilde{S}_{yy} - s^2\tilde{S}_{xx} + \sqrt{[\tilde{S}_{yy} - s^2\tilde{S}_{xx}]^2 + 4s^2|\tilde{S}_{xy}|^2}}{2\tilde{S}_{xy}} & = \angle \frac{1}{\tilde{S}_{xy}} & = \angle \tilde{S}_{xy}^* = -\angle \tilde{S}_{xy} \\
 &\uparrow \text{(Definition 11.8 page 99)} & & \\
 \angle \hat{H}_v &\triangleq \angle \frac{\tilde{S}_{yy} - \tilde{S}_{xx} + \sqrt{[\tilde{S}_{yy} - \tilde{S}_{xx}]^2 + 4|\tilde{S}_{xy}|^2}}{2\tilde{S}_{xy}} & = \angle \frac{1}{\tilde{S}_{xy}} & = \angle \tilde{S}_{xy}^* = -\angle \tilde{S}_{xy} \\
 &\uparrow \text{(Definition 11.9 page 100)} & & \\
 \angle \hat{H}_\kappa &\triangleq \angle \frac{\tilde{S}_{yy} - \kappa\tilde{S}_{xx} + \sqrt{[\tilde{S}_{yy} - \kappa\tilde{S}_{xx}]^2 + 4\kappa|\tilde{S}_{xy}|^2}}{2\tilde{S}_{xy}} & = \angle \frac{1}{\tilde{S}_{xy}} & = \angle \tilde{S}_{xy}^* = -\angle \tilde{S}_{xy} \\
 &\uparrow \text{(Definition 11.7 page 99)} & & \\
 \angle C_{xy} &\triangleq \angle \frac{\tilde{S}_{xy}^*}{\sqrt{\tilde{S}_{xx}\tilde{S}_{yy}}} & = & = \angle \tilde{S}_{xy}^* = -\angle \tilde{S}_{xy} \\
 &\uparrow \text{(Definition 11.12 page 116)} & & \\
 \Rightarrow &&&
 \end{aligned}$$

11.5 Alternate forms

Any standard kit of algebraic tricks should arguably always include the ability to swap the location of a square root between numerator and denominator. If you are of this persuasion, after traveling from the definition of \hat{H}_s on page 99, you won't be disappointed when arriving at the next proposition (Proposition 11.6 page 105). But it has more use than just allowing you to entertain friends at social occasions. It also makes it very easy to see (using only algebra) what previously employed *l'Hôpital's rule* (using calculus) in the proof of Theorem 11.1—that $\lim_{s \rightarrow \infty} \hat{H}_s = \hat{H}_1$.

Proposition 11.6. ¹⁷ Let $\hat{H}_\kappa(\omega; \kappa)$ be defined as in Definition 11.7 (page 99).

¹⁷  Shin and Hammond (2008) page 293 ((9.67)),  Leclere et al. (2014) ((10) $\kappa(f) = 1/s^2$ and x and y swapped)

P
R
P

$$\begin{aligned}\hat{H}_\kappa(\omega; s) &= \frac{2\kappa(\omega)\tilde{S}_{yx}(\omega)}{\kappa(\omega)\tilde{S}_{xx}(\omega) - \tilde{S}_{yy}(\omega) + \sqrt{[\kappa(\omega)\tilde{S}_{xx}(\omega) - \tilde{S}_{yy}(\omega)]^2 + 4\kappa(\omega)|\tilde{S}_{xy}(\omega)|^2}} \\ &= \frac{2\tilde{S}_{xy}^*}{\tilde{S}_{xx} - \frac{1}{\kappa(\omega)}\tilde{S}_{yy} + \sqrt{[\tilde{S}_{xx} - \frac{1}{\kappa(\omega)}\tilde{S}_{yy}]^2 + \frac{4}{\kappa(\omega)}|\tilde{S}_{xy}|^2}}\end{aligned}$$

PROOF: We can transform \hat{H}_κ from that found in Definition 11.8 (page 99) to the forms in this proposition by the technique of “rationalizing the denominator”¹⁸

$$\begin{aligned}\hat{H}_\kappa &\triangleq \frac{\tilde{S}_{yy} - \kappa\tilde{S}_{xx} + \sqrt{[\tilde{S}_{yy} - \kappa\tilde{S}_{xx}]^2 + 4\kappa|\tilde{S}_{xy}|^2}}{2\tilde{S}_{xy}} \quad \text{by definition of } \hat{H}_\kappa \text{ (Definition 11.8 page 99)} \\ &= \frac{\left[\tilde{S}_{yy} - \kappa\tilde{S}_{xx} + \sqrt{[\tilde{S}_{yy} - \kappa\tilde{S}_{xx}]^2 + 4\kappa|\tilde{S}_{xy}|^2}\right] \overbrace{\left[\tilde{S}_{yy} - \kappa\tilde{S}_{xx} - \sqrt{[\tilde{S}_{yy} - \kappa\tilde{S}_{xx}]^2 + 4\kappa|\tilde{S}_{xy}|^2}\right]}^{\text{"rationalizing factor"}}}{2\tilde{S}_{xy} \underbrace{\left[\tilde{S}_{yy} - \kappa\tilde{S}_{xx} - \sqrt{[\tilde{S}_{yy} - \kappa\tilde{S}_{xx}]^2 + 4\kappa|\tilde{S}_{xy}|^2}\right]}_{\text{"rationalizing factor}}} \\ &= \frac{[\tilde{S}_{yy} - \kappa\tilde{S}_{xx}]^2 - [\tilde{S}_{yy} - \kappa\tilde{S}_{xx}]^2 - 4\kappa|\tilde{S}_{xy}|^2}{2\tilde{S}_{xy} \left[\tilde{S}_{yy} - \kappa\tilde{S}_{xx} - \sqrt{[\tilde{S}_{yy} - \kappa\tilde{S}_{xx}]^2 + 4\kappa|\tilde{S}_{xy}|^2}\right]} = \frac{-4\kappa|\tilde{S}_{xy}|^2}{2\tilde{S}_{xy} \left[\tilde{S}_{yy} - \kappa\tilde{S}_{xx} - \sqrt{[\tilde{S}_{yy} - \kappa\tilde{S}_{xx}]^2 + 4\kappa|\tilde{S}_{xy}|^2}\right]} \\ &= \frac{2\kappa\tilde{S}_{xy}^*}{\kappa\tilde{S}_{xx} - \tilde{S}_{yy} + \sqrt{[\kappa\tilde{S}_{xx} - \tilde{S}_{yy}]^2 + 4\kappa|\tilde{S}_{xy}|^2}} = \frac{2\frac{\kappa}{s}\tilde{S}_{xy}^*}{\frac{\kappa}{s}\tilde{S}_{xx} - \frac{1}{s}\tilde{S}_{yy} + \sqrt{\frac{1}{s^4}[\kappa\tilde{S}_{xx} - \tilde{S}_{yy}]^2 + \frac{4\kappa}{s^4}|\tilde{S}_{xy}|^2}} \\ &= \frac{2\tilde{S}_{xy}^*}{\tilde{S}_{xx} - \frac{1}{\kappa}\tilde{S}_{yy} + \sqrt{[\tilde{S}_{xx} - \frac{1}{\kappa}\tilde{S}_{yy}]^2 + \frac{4}{\kappa}|\tilde{S}_{xy}|^2}}\end{aligned}$$

Integrity check for $s = 0$ and $s \rightarrow \infty$ cases: Let $p \triangleq \kappa$.

$$\begin{aligned}\lim_{p \rightarrow \infty} \hat{H}_\kappa &= \lim_{p \rightarrow \infty} \frac{2\tilde{S}_{yx}}{\tilde{S}_{xx} - \frac{1}{p}\tilde{S}_{yy} + \sqrt{\left[\tilde{S}_{xx} - \frac{1}{p}\tilde{S}_{yy}\right]^2 + \frac{4}{p}|\tilde{S}_{xy}|^2}} = \frac{2\tilde{S}_{yx}}{\tilde{S}_{xx} + \sqrt{[\tilde{S}_{xx}]^2}} \\ &= \frac{\tilde{S}_{yx}}{\tilde{S}_{xx}} \triangleq \hat{H}_1 \quad \text{by def. of } \hat{H}_1 \text{ (Definition 11.2 page 95)}\end{aligned}$$

$$\begin{aligned}\lim_{p \rightarrow 0} \hat{H}_\kappa &= \lim_{p \rightarrow 0} \frac{2p\tilde{S}_{yx}}{p\tilde{S}_{xx} - \tilde{S}_{yy} + \sqrt{[p\tilde{S}_{xx} - \tilde{S}_{yy}]^2 + 4p|\tilde{S}_{xy}|^2}} \\ &= \lim_{p \rightarrow 0} \frac{\frac{d}{dp}(2p\tilde{S}_{yx})}{\frac{d}{dp}\left(p\tilde{S}_{xx} - \tilde{S}_{yy} + \sqrt{[p\tilde{S}_{xx} - \tilde{S}_{yy}]^2 + 4p|\tilde{S}_{xy}|^2}\right)} \quad \text{by l'Hôpital's rule}\end{aligned}$$

¹⁸ Slaught and Lennes (1915), page 274 (“197. Rationalizing the Denominator.”) <https://archive.org/details/elementaryalgebr00slaurich/page/274> Note that the operation in the proof of Proposition 11.6 is being performed essentially in reverse...or rather “rationalizing the numerator”.



$$\begin{aligned}
 &= \lim_{p \rightarrow 0} -\frac{2\tilde{S}_{yx}}{\tilde{S}_{xx}(p\tilde{S}_{xx} - \tilde{S}_{yy}) + 2|\tilde{S}_{xy}|^2 + \tilde{S}_{xx}\sqrt{(p\tilde{S}_{xx} - \tilde{S}_{yy})^2 + 4p|\tilde{S}_{xy}|^2}} \\
 &= \frac{2\tilde{S}_{yy}\tilde{S}_{yx}}{-\tilde{S}_{xx}\tilde{S}_{yy} + 2|\tilde{S}_{xy}|^2 + \tilde{S}_{xx}\tilde{S}_{yy}} = \frac{2\tilde{S}_{yy}\tilde{S}_{yx}}{2|\tilde{S}_{xy}|^2} = \frac{\tilde{S}_{yy}}{\tilde{S}_{xy}} \\
 &\triangleq \hat{H}_2
 \end{aligned}$$

by def. of \hat{H}_2 (Definition 11.3 page 96)

11.6 Least squares estimates of non-linear systems

“The legendary Hungarian mathematician John von Neumann once referred to the theory of nonequilibrium systems as the “theory of non-elephants,” ... Nevertheless, such a theory of non-elephants will be attempted here.”

Per Bak, in “how nature works...” ¹⁹





Figure 11.3: Least Square estimation (Theorem 11.3 page 108)

Let S be the system illustrated to the right. If there is no measurement noise on the input and output and if H is linear time invariant, then $\tilde{H} = \tilde{S}_{xy}/\tilde{S}_{xx}$ (Corollary 5.1 page 36). But what if there is output measurement noise? And what if H is not LTI? What is the best least-squares estimate of \tilde{H} ? The answer depends on how you define “the best”.



The definition of “best” or “optimal” is given by a cost function $C(\hat{H})$. There are several possible cost functions. Definition 11.11 provides some. Theorem 11.3 then demonstrate optimal solutions with respect to these definitions.

Definition 11.11. Let S be a system defined as in Figure 11.3 (page 107) (A) or (B). Define the following COST FUNCTIONS for spectral LEAST-SQUARES estimates:

DEF	$C_{rq}(\hat{G}) \triangleq \tilde{F} \ r(n) - q(n)\ ^2 \triangleq \tilde{F} \langle r(n) - q(n) r(0) - q(0) \rangle \triangleq \tilde{F} E \left([r(n) - q(n)] [r(0) - q(0)]^* \right)$ $C_{sy}(\hat{H}) \triangleq \tilde{F} \ s(n) - y(n)\ ^2 \triangleq \tilde{F} \langle s(n) - y(n) s(0) - y(0) \rangle \triangleq \tilde{F} E \left([s(n) - y(n)] [s(0) - y(0)]^* \right)$
-----	---

¹⁹ Bak (2013) page 29 (§ Systems in Balance Are Not Complex)

Lemma 11.3. Let $C_{rq}(\hat{G})$ and $C_{sy}(\hat{H})$ be defined as in Definition 11.11 (page 107).

L E M	$C_{rq}(\hat{G}) = \tilde{S}_{pp}(\omega) \hat{G}(\omega) ^2 - \tilde{S}_{py}(\omega) \hat{G}(\omega) - \tilde{S}_{py}^*(\omega) \hat{G}^*(\omega) + \tilde{S}_{qq}(\omega)$ $C_{sy}(\hat{H}) = \tilde{S}_{xx}(\omega) \hat{H}(\omega) ^2 - \tilde{S}_{xy}(\omega) \hat{H}(\omega) - \tilde{S}_{xy}^*(\omega) \hat{H}^*(\omega) + \tilde{S}_{yy}(\omega)$
----------------------	---

PROOF:

$$C_{rq}(\hat{G})$$

$$\begin{aligned} &\triangleq \tilde{\mathbf{F}}\mathbf{E}\left(\left[\mathbf{r}(n) - \mathbf{q}(n)\right]\left[\mathbf{r}(0) - \mathbf{q}(0)\right]^*\right) && \text{by definition of } C_{rq} \quad (\text{Definition 11.11 page 107}) \\ &= \tilde{\mathbf{F}}\left[\mathbf{E}[\mathbf{r}(n)\mathbf{r}^*(0)] - \mathbf{E}[\mathbf{r}(n)\mathbf{q}^*(0)] - \mathbf{E}[\mathbf{q}(n)\mathbf{r}^*(0)] + \mathbf{E}[\mathbf{q}(n)\mathbf{q}^*(0)]\right] && \text{by linearity of E} \quad (\text{Theorem 1.1 page 4}) \\ &\triangleq \tilde{\mathbf{F}}\left[R_{rr}(m) - R_{rq}(m) - R_{qr}(m) + R_{qq}(m)\right] && \text{by definition of } R_{xy} \quad (\text{Definition 2.4 page 12}) \\ &\triangleq [\tilde{S}_{rr}(\omega) - \tilde{S}_{rq}(\omega) - \tilde{S}_{qr}(\omega) + \tilde{S}_{qq}(\omega)] && \text{by definition of } \tilde{S}_{xy} \quad (\text{Definition 6.3 page 44}) \\ &= \boxed{\tilde{S}_{pp}(\omega) |\hat{G}(\omega)|^2 - \tilde{S}_{py}(\omega) \hat{G}(\omega) - \tilde{S}_{py}^*(\omega) \hat{G}^*(\omega) + \tilde{S}_{qq}(\omega)} && \text{by (A)-(D) and Corollary 7.8 page 59} \end{aligned}$$

$$C_{sy}(\hat{H})$$

$$\begin{aligned} &\triangleq \tilde{\mathbf{F}}\mathbf{E}\left(\left[\mathbf{s}(n) - \mathbf{y}(n)\right]\left[\mathbf{s}(0) - \mathbf{y}(0)\right]^*\right) && \text{by definition of } C_{sy} \quad (\text{Definition 11.11 page 107}) \\ &= \tilde{\mathbf{F}}\left[\mathbf{E}[\mathbf{s}(n)\mathbf{s}^*(0)] - \mathbf{E}[\mathbf{s}(n)\mathbf{y}^*(0)] - \mathbf{E}[\mathbf{y}(n)\mathbf{s}^*(0)] + \mathbf{E}[\mathbf{y}(n)\mathbf{y}^*(0)]\right] && \text{by linearity of E} \quad (\text{Theorem 1.1 page 4}) \\ &\triangleq \tilde{\mathbf{F}}\left[R_{ss}(m) - R_{sy}(m) - R_{ys}(m) + R_{yy}(m)\right] && \text{by definition of } R_{xy} \quad (\text{Definition 2.4 page 12}) \\ &\triangleq [\tilde{S}_{ss}(\omega) - \tilde{S}_{sy}(\omega) - \tilde{S}_{ys}(\omega) + \tilde{S}_{yy}(\omega)] && \text{by definition of } \tilde{S}_{xy} \quad (\text{Definition 6.3 page 44}) \\ &= \boxed{\tilde{S}_{xx}(\omega) |\hat{H}(\omega)|^2 - \tilde{S}_{xy}(\omega) \hat{H}(\omega) - \tilde{S}_{xy}^*(\omega) \hat{H}^*(\omega) + \tilde{S}_{yy}(\omega)} && \text{by (A)-(D) and Corollary 7.8 (page 59)} \end{aligned}$$

Theorem 11.3. Let \mathbf{S} be the system illustrated in Figure 11.3 page 107 (A) or (B).

T H M	$\left. \begin{array}{l} (A). \mathbf{x}, \mathbf{u}, \text{ and } \mathbf{v} \text{ are WSS} \\ (B). \mathbf{x}, \mathbf{u}, \text{ and } \mathbf{v} \text{ are UNCORRELATED} \\ (C). \mathbf{Eu} = \mathbf{Ev} = 0 \text{ (ZERO-MEAN)} \\ (D). \hat{G}_{ls} \text{ and } \hat{H}_{ls} \text{ are LTI} \end{array} \right\} \Rightarrow \left\{ \begin{array}{l} (1). \arg \min_{\hat{G}} C_{rq}(\hat{G}) = \frac{\tilde{S}_{yp}(\omega)}{\tilde{S}_{pp}(\omega)} \\ (2). \arg \min_{\hat{H}} C_{sy}(\hat{H}) = \frac{\tilde{S}_{yx}(\omega)}{\tilde{S}_{xx}(\omega)} \end{array} \right.$
----------------------	---

PROOF:

- Define shorthand function names $\hat{G} \triangleq \hat{G}_{ls}$ and $\hat{H} \triangleq \hat{H}_{ls}$.

2. lemma:

$$\begin{aligned} 0 &= \frac{\partial}{\partial \hat{G}_R} C_{rq}(\hat{G}) \\ &= \frac{\partial}{\partial \hat{G}_R} \left(\tilde{S}_{pp} |\hat{G}|^2 - \hat{G} \tilde{S}_{py} - \hat{G}^* \tilde{S}_{py}^* + \tilde{S}_{qq} \right) && \text{by Lemma 11.3 page 108} \\ &= \frac{\partial}{\partial \hat{G}_R} \left(\tilde{S}_{pp} [\hat{G}_R^2 + \hat{G}_I^2] - (\hat{G}_R + i\hat{G}_I) \tilde{S}_{py} - (\hat{G}_R + i\hat{G}_I)^* \tilde{S}_{py}^* + \tilde{S}_{qq} \right) \\ &= 2\hat{G}_R \tilde{S}_{pp} - \tilde{S}_{py} - \tilde{S}_{py}^* + \frac{\partial}{\partial \hat{G}_R} \tilde{S}_{qq} \xrightarrow{0} && \text{because } q \text{ does not vary with } \hat{G} \\ &= 2\hat{G}_R \tilde{S}_{pp} - 2\mathbf{R}_e \tilde{S}_{py} \\ &= 2\hat{G}_R \tilde{S}_{pp} - 2\mathbf{R}_e \tilde{S}_{yp} && \text{by Corollary 2.2 page 15} \\ &\Rightarrow \boxed{\hat{G}_R(\omega) = \frac{\mathbf{R}_e \tilde{S}_{yp}(\omega)}{\tilde{S}_{pp}(\omega)}} \end{aligned}$$

3. lemma:

$$\begin{aligned}
 0 &= \frac{\partial}{\partial \hat{G}_I} C_{rq}(\hat{G}) \\
 &= \frac{\partial}{\partial \hat{G}_I} \left(\tilde{S}_{pp} |\hat{G}|^2 - \hat{G} \tilde{S}_{py} - \hat{G}^* \tilde{S}_{py}^* + \tilde{S}_{qq} \right) && \text{by Lemma 11.3 page 108} \\
 &= \frac{\partial}{\partial \hat{G}_I} [\tilde{S}_{pp} [\hat{G}_R^2 + \hat{G}_I^2] - (\hat{G}_R + i\hat{G}_I) \tilde{S}_{py} - (\hat{G}_R - i\hat{G}_I) \tilde{S}_{py}^* + \tilde{S}_{qq}] \\
 &= 2\hat{G}_I \tilde{S}_{pp} - i\tilde{S}_{py} + i\tilde{S}_{py}^* + \frac{\partial}{\partial \hat{G}_I} \tilde{S}_{qq} \xrightarrow{0} && \text{because } q \text{ does not vary with } \hat{G} \\
 &= 2\hat{G}_I \tilde{S}_{pp} - 2i(i\mathbf{I}_m \tilde{S}_{py}) \\
 &= 2\hat{G}_I \tilde{S}_{pp} + 2i(i\mathbf{I}_m \tilde{S}_{yp}) && \text{by Corollary 2.2 page 15} \\
 \implies \hat{G}_I(\omega) &= \frac{\mathbf{I}_m \tilde{S}_{yp}(\omega)}{\tilde{S}_{pp}(\omega)}
 \end{aligned}$$

4. Proof for $\hat{G} \triangleq \hat{G}_{ls}$ expression:

$$\begin{aligned}
 \hat{G}(\omega) &= \hat{G}_R(\omega) + i\hat{G}_I(\omega) \\
 &= \frac{\mathbf{R}_e \tilde{S}_{yp}(\omega)}{\tilde{S}_{pp}(\omega)} + \frac{i\mathbf{I}_m \tilde{S}_{yp}(\omega)}{\tilde{S}_{pp}(\omega)} && \text{by (2) lemma and (3) lemma} \\
 &= \frac{\tilde{S}_{yp}(\omega)}{\tilde{S}_{pp}(\omega)}
 \end{aligned}$$

5. lemma:

$$\begin{aligned}
 0 &= \frac{\partial}{\partial \hat{H}_R} C_{sy}(\hat{H}) \\
 &= \frac{\partial}{\partial \hat{H}_R} \left(\tilde{S}_{xx} |\hat{H}|^2 - \hat{H} \tilde{S}_{xy} - \hat{H}^* \tilde{S}_{xy}^* + \tilde{S}_{yy} \right) && \text{by Lemma 11.3 page 108} \\
 &= \frac{\partial}{\partial \hat{H}_R} (\tilde{S}_{xx} [\hat{H}_R^2 + \hat{H}_I^2] - (\hat{H}_R + i\hat{H}_I) \tilde{S}_{xy} - (\hat{H}_R + i\hat{H}_I)^* \tilde{S}_{xy}^* + \tilde{S}_{yy}) \\
 &= 2\hat{H}_R \tilde{S}_{xx} - \tilde{S}_{xy} - \tilde{S}_{xy}^* + \frac{\partial}{\partial \hat{H}_R} \tilde{S}_{yy} \xrightarrow{0} && \text{because } y \text{ does not vary with } \hat{H} \\
 &= 2\hat{H}_R \tilde{S}_{xx} - 2\mathbf{R}_e \tilde{S}_{xy} \\
 &= 2\hat{H}_R \tilde{S}_{xx} - 2\mathbf{R}_e \tilde{S}_{yx} && \text{by Corollary 2.2 page 15} \\
 \implies \hat{H}_R(\omega) &= \frac{\mathbf{R}_e \tilde{S}_{yx}(\omega)}{\tilde{S}_{xx}(\omega)}
 \end{aligned}$$

6. lemma:

$$\begin{aligned}
 0 &= \frac{\partial}{\partial \hat{H}_I} C_{sy}(\hat{H}) \\
 &= \frac{\partial}{\partial \hat{H}_I} \left(\tilde{S}_{xx} |\hat{H}|^2 - \tilde{S}_{xy} \hat{H} - \tilde{S}_{xy}^* \hat{H}^* + \tilde{S}_{yy} \right) && \text{by Lemma 11.3 page 108} \\
 &= \frac{\partial}{\partial \hat{H}_I} [\tilde{S}_{xx} [\hat{H}_R^2 + \hat{H}_I^2] - \tilde{S}_{xy} (\hat{H}_R + i\hat{H}_I) - \tilde{S}_{xy}^* (\hat{H}_R - i\hat{H}_I) + \tilde{S}_{yy}] \\
 &= 2\hat{H}_I \tilde{S}_{xx} - i\tilde{S}_{xy} + i\tilde{S}_{xy}^* + \frac{\partial}{\partial \hat{H}_I} \tilde{S}_{yy} \xrightarrow{0} && \text{because } q \text{ does not vary with } \hat{H}
 \end{aligned}$$

$$\begin{aligned}
 &= 2\hat{H}_I \tilde{S}_{xx} - 2i(i\mathbf{I}_m \tilde{S}_{xy}) \\
 &= 2\hat{H}_I \tilde{S}_{xx} + 2i(i\mathbf{I}_m \tilde{S}_{yx}) \\
 &= 2\hat{H}_I \tilde{S}_{xx} - 2\mathbf{I}_m \tilde{S}_{yx} \\
 \implies \hat{H}_I(\omega) &= \frac{\mathbf{I}_m \tilde{S}_{yx}(\omega)}{\tilde{S}_{xx}(\omega)}
 \end{aligned}$$

by Corollary 2.2 page 15

7. Proof for $\hat{H} \triangleq \hat{H}_{ls}$ expression:

$$\begin{aligned}
 \hat{H}(\omega) &= \hat{H}_R(\omega) + i\hat{H}_I(\omega) \\
 &= \frac{\mathbf{R}_e \tilde{S}_{yp}(\omega)}{\tilde{S}_{xx}(\omega)} + \frac{i\mathbf{I}_m \tilde{S}_{yp}(\omega)}{\tilde{S}_{xx}(\omega)} \\
 &= \frac{\mathbf{R}_e \tilde{S}_{yx}(\omega)}{\tilde{S}_{xx}(\omega)} + \frac{i\mathbf{I}_m \tilde{S}_{yx}(\omega)}{\tilde{S}_{xx}(\omega)} \\
 &= \frac{\tilde{S}_{yx}(\omega)}{\tilde{S}_{xx}(\omega)}
 \end{aligned}$$

by (5) lemma and (6) lemma

by Theorem 7.4 page 55

Using Theorem 11.3 (previous) we can see that the optimal **least-squares** operators \hat{G}_{ls} and \hat{H}_{ls} for the **non-linear** operator \mathbf{T} in Figure 11.3 (page 107) (A) and (B) are (next two corollaries)

$$\begin{aligned}
 (1). \quad \hat{G}_{ls}(\omega) &= \frac{\tilde{S}_{yx}(\omega)}{\tilde{S}_{xx}(\omega) + \tilde{S}_{uu}(\omega)} \quad \text{for (A)—communication system} \\
 (2). \quad \hat{G}_{ls}(\omega) &= \frac{\tilde{S}_{yx}(\omega)}{\tilde{S}_{xx}(\omega) - \tilde{S}_{uu}(\omega)} \quad \text{for (B)—measurement system} \\
 (3). \quad \hat{H}_{ls}(\omega) &= \frac{\tilde{S}_{yx}(\omega)}{\tilde{S}_{xx}(\omega)} \quad \text{for either (A) or (B)}
 \end{aligned}$$


Corollary 11.1. Let \mathbf{S} be the system illustrated in Figure 11.3 page 107 (A).

THM


$$\left\{ \begin{array}{l} \text{hypotheses of Theorem 11.3} \\ \text{page 108} \end{array} \right\} \implies \left\{ \begin{array}{l} (1). \quad \arg \min_{\hat{G}_{ls}} C_{rq}(\hat{G}_{ls}) = \frac{\tilde{S}_{yx}(\omega)}{\tilde{S}_{xx}(\omega) + \tilde{S}_{uu}(\omega)} \\ (2). \quad \arg \min_{\hat{H}_{ls}} C_{sy}(\hat{H}_{ls}) = \frac{\tilde{S}_{yx}(\omega)}{\tilde{S}_{xx}(\omega)} \end{array} \right\}$$

PROOF:

$$\begin{aligned}
 \hat{G}_{ls} &= \frac{\tilde{S}_{yx}(\omega)}{\tilde{S}_{pp}(\omega)} && \text{by Theorem 11.3 page 108} \\
 &= \frac{\tilde{S}_{yx}(\omega)}{\tilde{S}_{xx}(\omega) + \tilde{S}_{uu}(\omega)} && \text{by Theorem 7.1 page 51} \\
 \hat{H}_{ls} &= \frac{\tilde{S}_{yx}(\omega)}{\tilde{S}_{xx}(\omega)} && \text{by Theorem 11.3 page 108}
 \end{aligned}$$



(A) communications LTI additive noise model



(B) measurement LTI additive noise model

Figure 11.4: Additive noise systems with LTI operator \mathbf{H}

Corollary 11.2. Let \mathbf{S} be the system illustrated in Figure 11.3 page 107 (B).

T H M	$\left\{ \begin{array}{l} \text{hypotheses of Theorem 11.3} \\ \text{page 108} \end{array} \right\} \Rightarrow \left\{ \begin{array}{l} (1). \quad \arg \min_{\hat{\mathbf{G}}_{ls}} C_{rq}(\hat{\mathbf{G}}_{ls}) = \frac{\tilde{S}_{yx}(\omega)}{\tilde{S}_{xx}(\omega) - \tilde{S}_{uu}(\omega)} \\ (2). \quad \arg \min_{\hat{\mathbf{H}}_{ls}} C_{sy}(\hat{\mathbf{H}}_{ls}) = \frac{\tilde{S}_{yx}(\omega)}{\tilde{S}_{xx}(\omega)} \end{array} \right\}$
----------------------	--

PROOF:

$$\begin{aligned} \hat{\mathbf{G}} &= \frac{\tilde{S}_{yx}(\omega)}{\tilde{S}_{pp}(\omega)} && \text{by Theorem 11.3 page 108} \\ &= \frac{\tilde{S}_{yx}(\omega)}{\tilde{S}_{xx}(\omega) - \tilde{S}_{uu}(\omega)} && \text{by Theorem 7.1 page 51} \\ \hat{\mathbf{H}} &= \frac{\tilde{S}_{yx}(\omega)}{\tilde{S}_{xx}(\omega)} && \text{by Theorem 11.3 page 108} \end{aligned}$$

It follows immediately from Corollary 11.1 (page 110) and Corollary 11.2 (page 111) that, in the special case of no input noise ($u(n) = 0$), $\hat{\mathbf{H}}_1$ is the optimal least-squares estimate of $\tilde{\mathbf{H}}$ (next corollary).

Corollary 11.3.²⁰ Let \mathbf{S} be the system illustrated in Figure 11.3 page 107 (A) or (B).

C O R	$\left\{ \begin{array}{l} (1). \quad \text{hypotheses of Theorem 11.3 and} \\ (2). \quad u(n) = 0 \end{array} \right\} \Rightarrow \left\{ \hat{\mathbf{G}}_{ls}(\omega) = \hat{\mathbf{H}}_{ls}(\omega) = \hat{\mathbf{H}}_1(\omega) \right\}$
----------------------	---

11.7 Least squares estimates of linear systems

The previous section did assume the estimates $\hat{\mathbf{H}}_1$ and $\hat{\mathbf{H}}_2$ to be *linear time invariant (LTI)*, but it did *not* assume that the system transfer function \mathbf{T} itself to be *LTI*. But making the LTI assumption on \mathbf{H} yields some interesting and insightful results, such as those in this section.

Theorem 11.4 (Estimating \mathbf{H} in communication additive noise system). Let \mathbf{S} be the system illustrated in Figure 11.4 page 111 (A).

²⁰ Bendat and Piersol (1980) pages 98–100 (5.1.1 Optimal Character of Calculations; note: proof minimizing \tilde{S}_{vv} but yields same result), Bendat and Piersol (1993) pages 106–109 (5.1.1 Optimality of Calculations), Bendat and Piersol (2010) pages 187–190 (6.1.4 Optimum Frequency Response Functions)

THM

$$\left\{ \begin{array}{l} (A). \quad \mathbf{H} \text{ is} \\ (B). \quad \mathbf{x}(n) \text{ is} \\ (C). \quad \mathbf{x}(n), \mathbf{u}(n), \text{ and } \mathbf{v}(n) \text{ are UNCORRELATED} \end{array} \right. \quad \left. \begin{array}{l} \text{LINEAR TIME INVARIANT (LTI) and} \\ \text{WIDE-SENSE STATIONARY (WSS) and} \\ \text{UNCORRELATED} \end{array} \right\}$$

$$\Rightarrow \left\{ \begin{array}{l} (1). \quad \hat{\mathbf{H}}_1(\omega) = \tilde{\mathbf{H}}(\omega) \\ (2). \quad \hat{\mathbf{H}}_2(\omega) = \frac{\tilde{\mathbf{S}}_{vv}(\omega)}{\tilde{\mathbf{H}}^*(\omega)\tilde{\mathbf{S}}_{xx}(\omega)} + \tilde{\mathbf{H}}(\omega) \left[1 + \frac{\tilde{\mathbf{S}}_{uu}(\omega)}{\tilde{\mathbf{S}}_{xx}(\omega)} \right] \end{array} \right. \quad \left. \begin{array}{l} \text{and} \end{array} \right\}$$

PROOF:

$$\begin{aligned} \hat{\mathbf{H}}_1(\omega) &\triangleq \frac{\tilde{\mathbf{S}}_{yx}(\omega)}{\tilde{\mathbf{S}}_{xx}(\omega)} \\ &= \frac{\tilde{\mathbf{H}}(\omega)\tilde{\mathbf{S}}_{xx}(\omega)}{\tilde{\mathbf{S}}_{xx}(\omega)} \\ &= \tilde{\mathbf{H}}(\omega) \end{aligned}$$

by definition of $\hat{\mathbf{H}}_1$ (Definition 11.2 page 95)

$$\begin{aligned} \hat{\mathbf{H}}_2(\omega) &\triangleq \frac{\tilde{\mathbf{S}}_{yy}(\omega)}{\tilde{\mathbf{S}}_{xy}(\omega)} \\ &= \frac{\tilde{\mathbf{S}}_{yy}(\omega)}{\tilde{\mathbf{H}}^*(\omega)\tilde{\mathbf{S}}_{xx}(\omega)} \\ &= \frac{\tilde{\mathbf{S}}_{vv}(\omega) + \tilde{\mathbf{S}}_{qq}(\omega)}{\tilde{\mathbf{H}}^*(\omega)\tilde{\mathbf{S}}_{xx}(\omega)} \\ &= \frac{\tilde{\mathbf{S}}_{vv}(\omega) + \tilde{\mathbf{H}}^*(\omega)\tilde{\mathbf{H}}(\omega)\tilde{\mathbf{S}}_{pp}(\omega)}{\tilde{\mathbf{H}}^*(\omega)\tilde{\mathbf{S}}_{xx}(\omega)} \\ &= \frac{\tilde{\mathbf{S}}_{vv}(\omega)}{\tilde{\mathbf{H}}^*(\omega)\tilde{\mathbf{S}}_{xx}(\omega)} + \frac{\tilde{\mathbf{H}}^*(\omega)\tilde{\mathbf{H}}(\omega)[\tilde{\mathbf{S}}_{xx}(\omega) + \tilde{\mathbf{S}}_{uu}(\omega)]}{\tilde{\mathbf{H}}^*(\omega)\tilde{\mathbf{S}}_{xx}(\omega)} \\ &= \frac{\tilde{\mathbf{S}}_{vv}(\omega)}{\tilde{\mathbf{H}}^*(\omega)\tilde{\mathbf{S}}_{xx}(\omega)} + \tilde{\mathbf{H}}(\omega) \left[1 + \frac{\tilde{\mathbf{S}}_{uu}(\omega)}{\tilde{\mathbf{S}}_{xx}(\omega)} \right] \end{aligned}$$

by Corollary 7.5 page 57

by definition of $\hat{\mathbf{H}}_2$ (Definition 11.3 page 96)

by Corollary 7.5 page 57

by Theorem 7.1 page 51

by Corollary 5.3 page 37

⇒

Theorem 11.5 (Estimating \mathbf{H} in measurement additive noise system). ²¹ Let \mathbf{S} be the system illustrated in Figure 11.4 page 111 (B).

THM

$$\left\{ \begin{array}{l} (A). \quad \mathbf{H} \text{ is} \\ (B). \quad \mathbf{x}(n) \text{ is} \\ (C). \quad \mathbf{x}(n), \mathbf{u}(n), \text{ and } \mathbf{v}(n) \text{ are UNCORRELATED} \end{array} \right. \quad \left. \begin{array}{l} \text{LINEAR TIME INVARIANT and} \\ \text{WIDE-SENSE STATIONARY and} \\ \text{UNCORRELATED} \end{array} \right\}$$

$$\Rightarrow \left\{ \begin{array}{l} (1). \quad \hat{\mathbf{H}}_1(\omega) = \tilde{\mathbf{H}}(\omega) \left[\frac{1}{1 + \frac{\tilde{\mathbf{S}}_{ww}(\omega)}{\tilde{\mathbf{S}}_{pp}(\omega)}} \right] \quad (\text{UNDER-ESTIMATED}) \text{ and} \\ (2). \quad \hat{\mathbf{H}}_2(\omega) = \tilde{\mathbf{H}}(\omega) \left[1 + \frac{\tilde{\mathbf{S}}_{vv}(\omega)}{\tilde{\mathbf{S}}_{qq}(\omega)} \right] \quad (\text{OVER-ESTIMATED}) \end{array} \right. \quad \left. \begin{array}{l} \text{and} \end{array} \right\}$$

²¹ Shin and Hammond (2008) page 294 ($H_1(f) = H(f)$ (9.70); $H_2(f) = H(f)(1 + S_{n_y n_y}(f)/S_{yy}(f))$ (9.71)), Shin and Hammond (2008) page 294 ($H_1(f) = H(f)/(1 + S_{n_x n_x}/S_{xx}(f))$ (9.72); $H_2(f) = H(f)$ (9.73)), Mitchell (1982) page 277 ($H_1(f) = H_0(f)/(1 + G_{nn}/G_{uu})$) Mitchell (1982) page 278 ($H_2(f) = H_0(f)(1 + G_{mm}/G_{vv})$)

PROOF:

$$\begin{aligned}
 \hat{H}_1(\omega) &\triangleq \frac{\tilde{S}_{yx}(\omega)}{\tilde{S}_{xx}(\omega)} && \text{by definition of } \hat{H}_1 && (\text{Definition 11.2 page 95}) \\
 &= \frac{\tilde{S}_{pp}(\omega)\tilde{H}(\omega)}{\tilde{S}_{xx}(\omega)} && \text{by Corollary 7.4 page 57} \\
 &= \frac{\tilde{S}_{pp}(\omega)\tilde{H}(\omega)}{\tilde{S}_{pp}(\omega) + \tilde{S}_{ww}(\omega)} && \text{by hypothesis (A)} && \text{and Corollary 5.3 page 37} \\
 &= \tilde{H}(\omega) \left[\frac{1}{1 + \frac{\tilde{S}_{ww}(\omega)}{\tilde{S}_{pp}(\omega)}} \right] \\
 \hat{H}_2(\omega) &\triangleq \frac{\tilde{S}_{yy}(\omega)}{\tilde{S}_{xy}(\omega)} && \text{by definition of } \hat{H}_2 && (\text{Definition 11.3 page 96}) \\
 &= \frac{\tilde{S}_{qq}(\omega) + \tilde{S}_{vv}(\omega)}{\tilde{S}_{xy}(\omega)} && \text{by hypothesis (C)} && \text{and Corollary 7.1 page 52} \\
 &= \frac{\tilde{S}_{qq}(\omega) + \tilde{S}_{vv}(\omega)}{\tilde{S}_{xq}(\omega)} && \text{by hypothesis (C)} && \text{and Theorem 7.4 page 55} \\
 &= \frac{\tilde{S}_{qq}(\omega) + \tilde{S}_{vv}(\omega)}{\tilde{S}_{pq}(\omega)} && \text{by hypothesis (C)} && \text{and Lemma 7.3 page 54} \\
 &= \frac{\tilde{S}_{qq}(\omega) + \tilde{S}_{vv}(\omega)}{\tilde{S}_{qq}(\omega)/\tilde{H}(\omega)} && \text{by LTI hypothesis (A)} && \text{and Corollary 5.3 page 37} \\
 &= \tilde{H}(\omega) \left[1 + \frac{\tilde{S}_{vv}(\omega)}{\tilde{S}_{qq}(\omega)} \right] && \text{by hypotheses (A) and (B)} && \text{and Corollary 5.3 page 37}
 \end{aligned}$$



Corollary 11.4. Let S be the system illustrated in Figure 11.4 (page 111).

COR	$\left\{ \begin{array}{l} (A). \text{ hypotheses of Theorem 11.5 and} \\ (B). \mathbf{u}(n) = \mathbf{u}(n) = 0 \text{ (NO INPUT NOISE)} \end{array} \right\} \Rightarrow \left\{ \begin{array}{l} \hat{H}_1(\omega) = \tilde{H}(\omega) \text{ (UNBIASED)} \end{array} \right\}$
	$\left\{ \begin{array}{l} (A). \text{ hypotheses of Theorem 11.5 and} \\ (B). \mathbf{v}(n) = 0 \text{ (NO OUTPUT NOISE)} \end{array} \right\} \Rightarrow \left\{ \begin{array}{l} \hat{H}_2(\omega) = \tilde{H}(\omega) \text{ (UNBIASED)} \end{array} \right\}$

Lemma 11.4. Let S be the system illustrated in Figure 11.4 (page 111).

LEM	$\left\{ \begin{array}{l} \text{There exists } \kappa(\omega) \text{ such that } \tilde{S}_{vv}(\omega) = \kappa(\omega)\tilde{S}_{uu}(\omega) \end{array} \right\}$ $\Rightarrow \left\{ \begin{array}{l} \tilde{S}_{uu}(\omega) = \frac{ \hat{H}(\omega) ^2 \tilde{S}_{xx}(\omega) - \hat{H}(\omega)\tilde{S}_{xy}(\omega) - \hat{H}^*(\omega)\tilde{S}_{xy}^*(\omega) + \tilde{S}_{yy}(\omega)}{\kappa(\omega) + \hat{H}(\omega) ^2} \end{array} \right\}$
-----	--

PROOF:

- Development based on results of previous chapters:

$$\begin{aligned}
 \tilde{S}_{vv} &= \tilde{S}_{yy} - \tilde{S}_{qq} && \text{by Corollary 7.1 page 52} \\
 &= \tilde{S}_{yy} - \tilde{S}_{pq}\hat{H} && \text{by Corollary 5.3 page 37} \\
 &= \tilde{S}_{yy} - \tilde{S}_{xy}\hat{H} && \text{by Theorem 7.4 page 55} \\
 \tilde{S}_{uu} &= \tilde{S}_{xx} - \tilde{S}_{pp} && \text{by Corollary 7.1 page 52}
 \end{aligned}$$

$$\begin{aligned}
&= \tilde{S}_{xx} - \frac{\tilde{S}_{qp}}{\hat{H}} && \text{by Corollary 5.3 page 37} \\
&= \tilde{S}_{xx} - \frac{\tilde{S}_{yx}}{\hat{H}} && \text{by Theorem 7.4 page 55} \\
\tilde{S}_{uu} \left[|\hat{H}|^2 + \kappa \right] &= |\hat{H}|^2 \tilde{S}_{uu} + \kappa \tilde{S}_{uu} \\
&\triangleq \tilde{S}_{uu} |\hat{H}|^2 + \tilde{S}_{vv} && \text{by definition of } \kappa(\omega) \\
&= |\hat{H}|^2 \left[\tilde{S}_{xx} - \frac{\tilde{S}_{yx}}{\hat{H}} \right] + [\tilde{S}_{yy} - \tilde{S}_{xy} \hat{H}] \\
&= |\hat{H}|^2 \tilde{S}_{xx} - \hat{H}^* \tilde{S}_{yx} - \tilde{S}_{xy} \hat{H} + \tilde{S}_{yy} \\
\Rightarrow \tilde{S}_{uu}(\omega) &= \frac{|\hat{H}(\omega)|^2 \tilde{S}_{xx}(\omega) - \hat{H}(\omega) \tilde{S}_{xy}(\omega) - \hat{H}^*(\omega) \tilde{S}_{xy}^*(\omega) + \tilde{S}_{yy}(\omega)}{\kappa(\omega) + |\hat{H}(\omega)|^2}
\end{aligned}$$

2. Development of Wicks and Vold ([Wicks and Vold \(1986\)](#)):

$$\begin{aligned}
\tilde{Y} - \tilde{V} &= \tilde{Q} = \hat{H}\tilde{P} = \hat{H}(\tilde{X} - \tilde{U}) && \text{by definition of } \hat{H} \\
\hat{H}\tilde{U} - \tilde{V} &= \hat{H}\tilde{X} - \tilde{Y} && \text{by left distributive prop.} \quad (\text{Theorem M.4 page 285}) \\
E([\hat{H}\tilde{U} - \tilde{V}] [\hat{H}\tilde{U} - \tilde{V}]^*) &= E([\hat{H}\tilde{X} - \tilde{Y}] [\hat{H}\tilde{X} - \tilde{Y}]^*) \\
|\hat{H}|^2 \tilde{S}_{uu} - \hat{H} \cancel{\tilde{S}_{uv}}^0 - \hat{H}^* \cancel{\tilde{S}_{vu}}^0 + \tilde{S}_{vv} &= |\hat{H}|^2 \tilde{S}_{xx} - \hat{H} \tilde{S}_{xy} - \hat{H}^* \tilde{S}_{xy}^* + \tilde{S}_{yy} && \text{because } u \text{ and } v \text{ are uncorrelated} \\
|\hat{H}|^2 \tilde{S}_{uu} + \kappa \tilde{S}_{uu} &= |\hat{H}|^2 \tilde{S}_{xx} - \hat{H} \tilde{S}_{xy} - \hat{H}^* \tilde{S}_{xy}^* + \tilde{S}_{yy} && \text{by hypothesis}
\end{aligned}$$

⇒

Theorem 11.6. [22](#) Let S be the system illustrated in Figure 11.4 (page 111). Let $\hat{H}_k(\omega)$ be the transfer function estimate defined in Definition 11.7 (page 99).

T H M	$\left\{ \begin{array}{l} (1). \text{ There exists } \kappa(\omega) \text{ such that} \\ (2). \tilde{S}_{vv}(\omega) = \kappa(\omega) \tilde{S}_{uu}(\omega) \end{array} \right. \text{ and } \Rightarrow \left\{ \begin{array}{l} \arg \min_{\hat{H}} C(\hat{H}) = \hat{H}_k(\omega) \\ (\hat{H}_k \text{ is the "optimal" estimator for minimizing system noise}) \end{array} \right.$
-------------	--

PROOF:

1. Let $F \triangleq |\hat{H}(\omega)|^2 \tilde{S}_{xx}(\omega) - \hat{H}(\omega) \tilde{S}_{xy}(\omega) - \hat{H}^*(\omega) \tilde{S}_{xy}^*(\omega) + \tilde{S}_{yy}(\omega)$ (numerator in Lemma 11.4) and $G \triangleq \kappa(\omega) + |\hat{H}(\omega)|^2$ (denominator in Lemma 11.4)

2. lemma $\left(\frac{\partial}{\partial \hat{H}_R} \tilde{S}_{uu} \right)$:

$$\begin{aligned}
\boxed{0} &= \frac{1}{2} G^2 0 = \frac{1}{2} G^2 \frac{\partial}{\partial \hat{H}_R} \tilde{S}_{uu} && \text{set } \frac{\partial}{\partial \hat{H}_R} \tilde{S}_{uu} = 0 \text{ to find optimum } \hat{H}_R \\
&= \frac{1}{2} G^2 \frac{\partial}{\partial \hat{H}_R} \frac{F}{G} && \text{by Lemma 11.4 page 113} \\
&= \frac{1}{2} G^2 \frac{(F'G - G'F)}{G^2} && \text{by Quotient Rule} \\
&= \frac{1}{2}(F'G - G'F)
\end{aligned}$$

²² [Wicks and Vold \(1986\)](#) page 898 (has additional s in denominator), [Shin and Hammond \(2008\)](#) page 293 ((9.67)), [White et al. \(2006\)](#) page 679 (6))

$$\begin{aligned}
 &= \frac{1}{2} [2\hat{H}_R \tilde{S}_{xx} - \tilde{S}_{xy} - \tilde{S}_{xy}^*] G - \frac{1}{2} 2\hat{H}_R F \quad \text{by definition of } F, G \\
 &= \boxed{\hat{H}_R \tilde{S}_{xx} G - G R_e \tilde{S}_{xy} - \hat{H}_R F}
 \end{aligned}$$

(item (1) page 114)

3. lemma $\left(\frac{\partial}{\partial \hat{H}_I} \tilde{S}_{uu}\right)$:

$$\begin{aligned}
 \boxed{0} &= \frac{1}{2} G^2 0 = \frac{1}{2} G^2 \frac{\partial}{\partial \hat{H}_I} \tilde{S}_{uu} && \text{set } \frac{\partial}{\partial \hat{H}_I} \tilde{S}_{uu} = 0 \text{ to find optimum } \hat{H}_I \\
 &= \frac{1}{2} G^2 0 = \frac{1}{2} G^2 \frac{\partial}{\partial \hat{H}_I} \frac{F}{G} && \text{by Lemma 11.4 page 113} \\
 &= \frac{1}{2} G^2 \frac{(F'G - G'F)}{G^2} && \text{by Quotient Rule} \\
 &= \frac{1}{2} (F'G - G'F) \\
 &= \frac{1}{2} [2\hat{H}_I \tilde{S}_{xx} - i\tilde{S}_{xy} + i\tilde{S}_{xy}^*] G - \frac{1}{2} 2\hat{H}_I F \quad \text{by definition of } F, G \\
 &= \boxed{\hat{H}_I \tilde{S}_{xx} G + G I_m \tilde{S}_{xy} - \hat{H}_I F}
 \end{aligned}$$

(item (1) page 114)

4. Solve for \hat{H} ...

$$\begin{aligned}
 0 = 0 + i0 &= \frac{1}{2} G^2 0 + \frac{1}{2} G^2 0 = \frac{1}{2} G^2 \frac{\partial}{\partial \hat{H}_R} \tilde{S}_{uu} + i \frac{1}{2} G^2 \frac{\partial}{\partial \hat{H}_I} \tilde{S}_{uu} \\
 &= [\hat{H}_R \tilde{S}_{xx} G - G R_e \tilde{S}_{xy} - \hat{H}_R F] + i[\hat{H}_I \tilde{S}_{xx} G + G I_m \tilde{S}_{xy} - \hat{H}_I F] \quad \text{by (2) lemma and (3) lemma} \\
 &= \hat{H} \tilde{S}_{xx} G - \tilde{S}_{xy}^* G - \hat{H} F \quad \text{because } R_e(z) + iI_m(z) = z \text{ and } R_e(z) - iI_m(z) = z^* \\
 &= \hat{H} \tilde{S}_{xx} G - \tilde{S}_{yx} G - \hat{H} F \quad \text{by Corollary 2.2 page 15} \\
 &= \hat{H} \tilde{S}_{xx} (\kappa + |\hat{H}|^2) - \tilde{S}_{yx} (\kappa + |\hat{H}|^2) - \hat{H} (|\hat{H}|^2 \tilde{S}_{xx} - \hat{H} \tilde{S}_{xy} - \hat{H}^* \tilde{S}_{xy}^* + \tilde{S}_{yy}) \quad \text{by } F, G \text{ defs.} \\
 &= \hat{H} \tilde{S}_{xx} \left(\kappa + |\hat{H}|^2 \right) - \tilde{S}_{yx} \left(\kappa + |\hat{H}|^2 \right) - \hat{H} \left(|\hat{H}|^2 \tilde{S}_{xx} - \hat{H} \tilde{S}_{xy} - \hat{H}^* \tilde{S}_{xy}^* + \tilde{S}_{yy} \right) \\
 &= \kappa \hat{H} \tilde{S}_{xx} - \tilde{S}_{yx} \left(\kappa + |\hat{H}|^2 \right) + \left(\hat{H}^2 \tilde{S}_{xy} + |\hat{H}|^2 \tilde{S}_{xy}^* - \hat{H} \tilde{S}_{yy} \right) \\
 &= \kappa \hat{H} \tilde{S}_{xx} - \kappa \tilde{S}_{yx} - \tilde{S}_{yx} |\hat{H}|^2 + \left(\hat{H}^2 \tilde{S}_{xy} + |\hat{H}|^2 \tilde{S}_{xy}^* - \hat{H} \tilde{S}_{yy} \right) \\
 &= \hat{H}^2 \tilde{S}_{xy} + \hat{H} [\kappa \tilde{S}_{xx} - \tilde{S}_{yy}] - \kappa \tilde{S}_{xy}^* \\
 \Rightarrow \hat{H} &= \frac{(\tilde{S}_{yy} - \kappa \tilde{S}_{xx}) \pm \sqrt{(\tilde{S}_{yy} - \kappa \tilde{S}_{xx})^2 + 4\kappa |\tilde{S}_{xy}|^2}}{2\tilde{S}_{xy}}
 \end{aligned}$$

by Quadratic Equation

11.8 Coherence

11.8.1 Application

Coherence has two basic purposes:

1. The *coherence* of x and y is a measure of how closely x and y are statistically related. That is, it is an indication of how much x and y "cohere" or "stick" together

2. The *coherence* of x and y is a measure of how reliable are the estimates \hat{H}_1 and \hat{H}_2 (Definition 11.2 page 95, Definition 11.3 page 96). If the coherence is 0.70 or above, then we can have high confidence that the estimates \hat{H}_1 and \hat{H}_2 are “good” estimates.²³

11.8.2 Definitions

Definition 11.12. ²⁴ Let S be a system with input $x(n)$ and output $y(n)$.

DEF

The **complex coherence function** is defined as $C_{xy}(\omega) \triangleq \frac{\tilde{S}_{xy}^*(\omega)}{\sqrt{\tilde{S}_{xx}(\omega)\tilde{S}_{yy}(\omega)}}$

The **ordinary coherence function** is defined as $\gamma_{xy}^2(\omega) \triangleq \frac{|\tilde{S}_{xy}(\omega)|^2}{\tilde{S}_{xx}(\omega)\tilde{S}_{yy}(\omega)}$

Proposition 11.7.

P R P $\gamma_{xy}^2(\omega) = \frac{\hat{H}_1(\omega)}{\hat{H}_2(\omega)}$

PROOF:

$$\gamma_{xy}^2(\omega) \triangleq \frac{|\tilde{S}_{xy}|^2}{\tilde{S}_{xx}\tilde{S}_{yy}} \quad \text{by definition of } \gamma_{xy}^2 \quad (\text{Definition 11.12 page 116})$$

$$= \frac{\tilde{S}_{xy}^*\tilde{S}_{xx}}{\tilde{S}_{yy}\tilde{S}_{xy}} \triangleq \frac{\hat{H}_1}{\hat{H}_2} \quad \text{by definitions of } \hat{H}_1 \text{ and } \hat{H}_2 \quad (\text{Definition 11.2 page 95, Definition 11.3 page 96})$$

Remark 11.2. Note that the *complex transmissibility* \tilde{T}'_{xy} of Remark 11.1 provides a nice mathematical symmetry (always a good sign of good direction) with *coherence* in the system identification family tree. In particular, note that the following:

R E M $C_{xy} \triangleq \sqrt{\frac{\hat{H}_1^*}{\hat{H}_2}}$ whereas $\tilde{T}'_{xy} \triangleq \sqrt{\hat{H}_1\hat{H}_2}$

PROOF:

$$\sqrt{\frac{\hat{H}_1^*(\omega)}{\hat{H}_2(\omega)}} \quad \text{by definition of } \hat{H}_{gm} \quad (\text{Definition 11.5 page 97})$$

11.8.3 A warning

Estimators yield, as the name implies, estimates. These estimates in general contain some error.

²³ Liang and Lee (2015) pages 363–365 (7.4.2 COHERENCE FUNCTION)

²⁴ Chen et al. (2012) page 4699(1), (2), Liang and Lee (2015) pages 363–365 (7.4.2 Coherence function), Ewins (1986) page 131 ($\gamma^2 = H_1(\omega)/H_2(\omega)$ (3.8))

Example 11.1 (The K=1 Welch estimate of coherence). Suppose we have two *uncorrelated* stationary sequences $x(n)$ and $y(n)$. Then, there CSD $S_{xy}(\omega)$ should be 0 because

$$\begin{aligned} S_{xy}(\omega) &\triangleq \check{\mathbf{F}}\mathbf{E}_{xy}(m) \\ &= \check{\mathbf{F}}\mathbf{E}[x(n)y[n+m]] \\ &= \check{\mathbf{F}}[\mathbf{E}_x(n)][\mathbf{E}_y[n+m]] \\ &= \check{\mathbf{F}}[0][0] \\ &= 0 \end{aligned}$$

This will give a coherence of 0 also:

$$C(\omega) = \frac{S_{xy}}{\sqrt{S_{xx}S_{yy}}} = 0$$

However, the Welch estimate with $K = 1$ will yield


$$\begin{aligned} |C(\omega)| &= \left| \frac{S_{xy}}{\sqrt{S_{xx}S_{yy}}} \right| \\ &= \left| \frac{(\tilde{\mathbf{F}}x)(\tilde{\mathbf{F}}y)^*}{\sqrt{|\tilde{\mathbf{F}}x|^2|\tilde{\mathbf{F}}y|^2}} \right| \\ &= 1 \end{aligned}$$

CHAPTER 12


ESTIMATING NOISE

Estimating noise in a system is difficult and many estimation methods are possible.

- Thong et al. (2001)
- Zheng et al. (2002)
- Kim and Kamel (2004)
- Kamel and Sim (2004)



(A) communications LTI additive noise model



(B) measurement LTI additive noise model

Figure 12.1: Additive noise systems with LTI operator \mathbf{H}

CHAPTER 13

COMMUNICATION CHANNELS

13.1 System model




Figure 13.1: Communication system model

A communication system is an operator \mathbf{S} over an information sequence u that generates an estimated information sequence $\hat{s}u$. The system operator factors into a receive operator \mathbf{R} , a channel operator \mathbf{C} , and a transmit operator \mathbf{T} such that

$$\mathbf{S} = \mathbf{RCT}.$$

The transmit operator operates on an information sequence u to generate a channel signal $s(t; u)$. The channel operator operates on the transmitted signal $s(t; u)$ to generate the received signal $r(t; u)$. The receive operator operates on the received signal $r(t; u)$ to generate the estimate $\hat{s}u$ (see Figure 13.1 (page 121)).

Definition 13.1. Let U be the set of all sequences u and let

DEF	$\mathbf{S} : U \rightarrow U$ (system operator)
	$\mathbf{T} : U \rightarrow \mathbb{R}^\infty$ (transmit operator)
	$\mathbf{C} : \mathbb{R}^\infty \rightarrow \mathbb{R}^\infty$ (channel operator)
	$\mathbf{R} : \mathbb{R}^\infty \rightarrow U$ (receive operator)

be operators. A **digital communication system** is the operation \mathbf{S} on the set of information sequences U such that $\mathbf{S} \stackrel{\triangle}{=} \mathbf{RCT}$.

Communication systems can be continuous or discrete valued in time and/or amplitude:

$s(t) = a(t)\psi(t)$	continuous time t	discrete time t
continuous amplitude $a(t)$	analog communications	discrete-time communications
discrete amplitude $a(t)$	—	digital communications

In this document, we normally take the approach that

1. \mathbf{C} is stochastic
2. There is no structural constraint on \mathbf{R} .
3. \mathbf{R} is optimum with respect to the ML-criterion.

These characteristics are explained more fully below.

13.1.1 Channel operator

Real-world physical channels perform a number of operations on a signal. Often these operations are closely modeled by a channel operator \mathbf{C} . Properties that characterize a particular channel operator associated with some physical channel include

- ☛ linear or non-linear
- ☛ time-invariant or time-variant
- ☛ memoryless or non-memoryless
- ☛ deterministic or stochastic.

Examples of physical channels include free space, air, water, soil, copper wire, and fiber optic cable. Information is carried through a channel using some physical process. These processes include:

Process	Example
☛ electromagnetic waves	free space, air
☛ acoustic waves	water, soil
☛ electric field potential (voltage)	wire
☛ light	fiber optic cable
☛ quantum mechanics	

13.1.2 Receive operator

Let \mathbf{I} be the *identity operator*.¹ Ideally, \mathbf{R} is selected such that $\mathbf{RCT} = \mathbf{I}$. In this case we say that \mathbf{R} is the *left inverse*² of \mathbf{CT} and denote this left inverse by \mathbf{C} . One example of a system where this inverse exists is the noiseless ISI system. While this is quite useful for mathematical analysis and system design, \mathbf{C} does not actually exist for any real-world system.

When \mathbf{C} does not exist, the “ideal” \mathbf{R} is one that is optimum

1. with respect to some *criterion* (or cost function)
2. and sometimes under some structural *constraint*.

¹ \mathbf{I} is the *identity operator* if for any operator \mathbf{X} , $\mathbf{XI} = \mathbf{IX} = \mathbf{X}$.

² $\mathbf{X}^{-1}\mathbf{X}$ is the *left inverse* of \mathbf{X} if $\mathbf{X}^{-1}\mathbf{XX} = \mathbf{I}$.

$\mathbf{X}^{-1}\mathbf{X}$ is the *right inverse* of \mathbf{X} if $\mathbf{XX}^{-1}\mathbf{X} = \mathbf{I}$.

$\mathbf{X}^{-1}\mathbf{X}$ is the *inverse* of \mathbf{X} if $\mathbf{X}^{-1}\mathbf{XX} = \mathbf{XX}^{-1}\mathbf{X} = \mathbf{I}$.



When a structural constraint is imposed on \mathbf{R} , the solution is called *structured*; otherwise, it is called *non-structured*.³ A common example of a structured approach is the use of a transversal filter (FIR filter in DSP) in which optimal coefficients are found for the filter. A structured \mathbf{R} is only optimal with respect to the imposed constraint. Even though \mathbf{R} may be optimal with respect to this structure, \mathbf{R} may not be optimal in general; that is, there may be another structure that would lead to a “better” solution. In a non-structured approach, \mathbf{R} is free to take any form whatsoever (practical or impractical) and therefore leads to the best of the best solutions.

The nature of \mathbf{R} depends heavily on the nature of \mathbf{C} . If \mathbf{C} does not exist, then the ideal \mathbf{R} is one that is optimal with respect to some criterion. If \mathbf{C} is deterministic, then appropriate optimization criterion may include

- least square error (LSE) criterion
- minimum absolute error criterion
- minimum peak distortion criterion.

If \mathbf{C} is stochastic then appropriate optimization criterion may include

- | | |
|---|--|
| • Bayes: | pdf known and cost function defined |
| • Maximum a posteriori probability (MAP): | pdf known and uniform cost function |
| • Maximum likelihood (ML): | pdf known and no prior probability information |
| • mini-max: | pdf not known but a cost function is defined |
| • Neyman-Pearson: | pdf not known and no cost function defined. |

Making \mathbf{R} optimum with respect to one of these criterion leads to an *estimate* $\hat{s}u = \mathbf{RCTu}$ that is also optimum with respect to the same criterion (Definition 8.1 page 64).

13.2 Optimization in the case of additional operations




Figure 13.2: Theorem of reversibility

Often in communication systems, an additional operator \mathbf{X} is inserted such that (see Figure 13.2 (page 123))

$$\mathbf{S} = \mathbf{RXCT}.$$

An example of such an operator \mathbf{X} is a receive filter. Is it still possible to find an \mathbf{R} that will perform as well as the case where \mathbf{X} is not inserted? In general, the answer is “no”. For example, if $\mathbf{Xr} = 0$, then all received information is lost and obviously there is no \mathbf{R} that can recover from this event. However, in the case where the right inverse $\mathbf{X}^{-1}\mathbf{X}$ of \mathbf{X} exists, then the answer to the question is “yes” and an optimum \mathbf{R} still exists. That is, it doesn't matter if an \mathbf{X} is inserted into system as long as \mathbf{X} is invertible. This is stated formally in the next theorem.

Theorem 13.1 (Theorem of Reversibility). ⁴ Let

³ Trees (2001) page 12

⁴ Trees (2001) pages 289–290

- 4) $\hat{\theta} = \mathbf{R}^{-1}\mathbf{C}\mathbf{T}\mathbf{u}$ be the optimum estimate of u
- 5) \mathbf{X} be an operator with right inverse $\mathbf{X}^{-1}\mathbf{X}$.

Then there exists some \mathbf{R}' such that

**T
H
M** $\hat{\theta} = \mathbf{R}'\mathbf{X}\mathbf{C}\mathbf{T}\mathbf{u}$.

PROOF: Let $\mathbf{R}' = \mathbf{R}\mathbf{X}^{-1}\mathbf{X}$. Then

$$\mathbf{R}'\mathbf{X}\mathbf{C}\mathbf{T}\mathbf{u} = \mathbf{R}\mathbf{X}^{-1}\mathbf{X}\mathbf{C}\mathbf{T}\mathbf{u} = \mathbf{R}\mathbf{C}\mathbf{T}\mathbf{u} = \hat{\theta}$$



13.3 Channel Statistics

The receiver needs to make a decision as to what sequence (u) the transmitter has sent. This decision should be optimal in some sense. Very often the optimization criterion is chosen to be the *maximal likelihood (ML)* criterion. The information that the receiver can use to make an optimal decision is the received signal $r(t)$.

If the symbols in $r(t)$ are statistically *independent*, then the optimal estimate of the current symbol depends only on the current symbol period of $r(t)$. Using other symbol periods of $r(t)$ has absolutely no additional benefit. Note that the AWGN channel is *memoryless*; that is, the way the channel treats the current symbol has nothing to do with the way it has treated any other symbol. Therefore, if the symbols sent by the transmitter into the channel are independent, the symbols coming out of the channel are also independent.

However, also note that the symbols sent by the transmitter are often very intentionally not independent; but rather a strong relationship between symbols is intentionally introduced. This relationship is called *channel coding*. With proper channel coding, it is theoretically possible to reduce the probability of communication error to any arbitrarily small value as long as the channel is operating below its *channel capacity*.

This chapter assumes that the received symbols are statistically independent; and therefore optimal decisions at the receiver for the current symbol are made only from the current symbol period of $r(t)$.

The received signal $r(t)$ over a single symbol period contains an uncountably infinite number of points. That is a lot. It would be nice if the receiver did not have to look at all those uncountably infinite number of points when making an optimal decision. And in fact the receiver does indeed not have to. As it turns out, a single finite set of *statistics* $\{r_1, r_2, \dots, r_N\}$ is sufficient for the receiver to make an optimal decision as to which value the transmitter sent.

Definition 13.2. Let C be an additive noise channel

CHAPTER 14

OPTIMAL SYMBOL DETECTION

14.1 ML Estimation

Theorem 14.1. In an AWGN channel with received signal $r(t) = s(t; \phi) + n(t)$ Let

- ➊ $r(t) = s(t; \phi) + n(t)$ be the received signal in an AWGN channel
- ➋ $n(t)$ a Gaussian white noise process
- ➌ $s(t; \phi)$ the transmitted signal such that

$$s(t; \phi) = \sum_{n \in \mathbb{Z}} a_n \lambda(t - nT) \cos(2\pi f_c t + \theta_n + \phi).$$

Then the optimal ML estimate of ϕ is either of the two equivalent expressions

THM

$$\begin{aligned} \hat{\phi}_{\text{ml}} &= -\text{atan} \left[\frac{\sum_{n \in \mathbb{Z}} a_n \int_{t \in \mathbb{R}} r(t) \lambda(t - nT) \sin(2\pi f_c t + \theta_n) dt}{\sum_{n \in \mathbb{Z}} a_n \int_{t \in \mathbb{R}} r(t) \lambda(t - nT) \cos(2\pi f_c t + \theta_n) dt} \right] \\ &= \arg_{\phi} \left(\sum_{n \in \mathbb{Z}} a_n \int_{t \in \mathbb{R}} r(t) [\lambda(t - nT) \sin(2\pi f_c t + \theta_n + \phi)] dt = 0 \right). \end{aligned}$$

PROOF:

$$\begin{aligned} \hat{\phi}_{\text{ml}} &= \arg_{\phi} \left(2 \int_{t \in \mathbb{R}} r(t) \left[\frac{\partial}{\partial \phi} s(t; \phi) \right] dt = \frac{\partial}{\partial \phi} \int_{t \in \mathbb{R}} s^2(t; \phi) dt \right) \quad \text{by Theorem 9.6 page 80} \\ &= \arg_{\phi} \left(2 \int_{t \in \mathbb{R}} r(t) \left[\frac{\partial}{\partial \phi} s(t; \phi) \right] dt = \frac{\partial}{\partial \phi} \|s(t; \phi)\|^2 dt \right) \\ &= \arg_{\phi} \left(2 \int_{t \in \mathbb{R}} r(t) \left[\frac{\partial}{\partial \phi} s(t; \phi) \right] dt = 0 \right) \\ &= \arg_{\phi} \left(\int_{t \in \mathbb{R}} r(t) \left[\frac{\partial}{\partial \phi} \sum_{n \in \mathbb{Z}} a_n \lambda(t - nT) \cos(2\pi f_c t + \theta_n + \phi) \right] dt = 0 \right) \end{aligned}$$

$$\begin{aligned}
&= \arg_{\phi} \left(- \sum_{n \in \mathbb{Z}} a_n \int_{t \in \mathbb{R}} r(t) [\lambda(t - nT) \sin(2\pi f_c t + \theta_n + \phi)] dt = 0 \right) \\
&= \arg_{\phi} \left(\sum_{n \in \mathbb{Z}} a_n \int_{t \in \mathbb{R}} r(t) \lambda(t - nT) [\sin(2\pi f_c t + \theta_n) \cos(\phi) + \sin(\phi) \cos(2\pi f_c t + \theta_n)] dt = 0 \right) \\
&= \arg_{\phi} \left(\sum_{n \in \mathbb{Z}} a_n \int_{t \in \mathbb{R}} r(t) \lambda(t - nT) \sin(\phi) \cos(2\pi f_c t + \theta_n) dt = - \sum_{n \in \mathbb{Z}} a_n \int_{t \in \mathbb{R}} r(t) \lambda(t - nT) \sin(2\pi f_c t + \theta_n) \cos(\phi) dt \right) \\
&= \arg_{\phi} \left(\sin(\phi) \sum_{n \in \mathbb{Z}} a_n \int_{t \in \mathbb{R}} r(t) \lambda(t - nT) \cos(2\pi f_c t + \theta_n) dt = -\cos(\phi) \sum_{n \in \mathbb{Z}} a_n \int_{t \in \mathbb{R}} r(t) \lambda(t - nT) \sin(2\pi f_c t + \theta_n) dt \right) \\
&= \arg_{\phi} \left(\frac{\sin(\phi)}{\cos(\phi)} = - \frac{\sum_{n \in \mathbb{Z}} a_n \int_{t \in \mathbb{R}} r(t) \lambda(t - nT) \sin(2\pi f_c t + \theta_n) dt}{\sum_{n \in \mathbb{Z}} a_n \int_{t \in \mathbb{R}} r(t) \lambda(t - nT) \cos(2\pi f_c t + \theta_n) dt} \right) \\
&= \arg_{\phi} \left(\tan(\phi) = - \frac{\sum_{n \in \mathbb{Z}} a_n \int_{t \in \mathbb{R}} r(t) \lambda(t - nT) \sin(2\pi f_c t + \theta_n) dt}{\sum_{n \in \mathbb{Z}} a_n \int_{t \in \mathbb{R}} r(t) \lambda(t - nT) \cos(2\pi f_c t + \theta_n) dt} \right) \\
&= \arg_{\phi} \left(\phi = -\text{atan} \left(\frac{\sum_{n \in \mathbb{Z}} a_n \int_{t \in \mathbb{R}} r(t) \lambda(t - nT) \sin(2\pi f_c t + \theta_n) dt}{\sum_{n \in \mathbb{Z}} a_n \int_{t \in \mathbb{R}} r(t) \lambda(t - nT) \cos(2\pi f_c t + \theta_n) dt} \right) \right) \\
&= -\text{atan} \left(\frac{\sum_{n \in \mathbb{Z}} a_n \int_{t \in \mathbb{R}} r(t) \lambda(t - nT) \sin(2\pi f_c t + \theta_n) dt}{\sum_{n \in \mathbb{Z}} a_n \int_{t \in \mathbb{R}} r(t) \lambda(t - nT) \cos(2\pi f_c t + \theta_n) dt} \right)
\end{aligned}$$

⇒

14.2 Generalized coherent modulation




Figure 14.1: Distributions of orthonormal components

Theorem 14.2. Let

- (V, ⟨· | ·⟩, S) be a modulation space
- Ψ ≜ {ψ_n(t) : n = 1, 2, …, N} be a set of orthonormal functions that span S
- r_n ≜ ⟨r(t) | ψ_n(t)⟩
- R ≜ {r_n : n = 1, 2, …, N}
- s_n(m) ≜ ⟨s(t; m) | ψ_n(t)⟩

and let V be partitioned into **decision regions**

$$\{D_m : m = 1, 2, \dots, |S|\}$$

such that

$$r(t) \in D_{\hat{m}} \iff \hat{m} = \arg \max_m P\{s(t; m) | r(t)\}.$$



Then the **probability of detection error** is

$$\text{T H M} \quad P\{\text{error}\} = 1 - \left(\frac{1}{\sqrt{2\pi\sigma^2}} \right)^N \sum_m P\{m \text{ sent}\} \int_{\mathbf{r} \in D_m} \exp \frac{-1}{2\sigma^2} \sum_{n=1}^N [\dot{r}_n - \dot{s}_n(m)]^2 d\mathbf{r}.$$

PROOF:

$$\begin{aligned} P\{\text{error}\} &= 1 - P\{\text{no error}\} \\ &= 1 - \sum_m P\{(m \text{ sent}) \wedge (\hat{m} = m \text{ detected})\} \\ &= 1 - \sum_m P\{(\hat{m} = m \text{ detected}) | (m \text{ sent})\} P\{m \text{ sent}\} \\ &= 1 - \sum_m P\{m \text{ sent}\} P\{\mathbf{r}|(m \text{ sent})\} \\ &= 1 - \sum_m P\{m \text{ sent}\} \int_{\mathbf{r} \in D_m} p[\mathbf{r}|(m \text{ sent})] d\mathbf{r} \\ &= 1 - \sum_m P\{m \text{ sent}\} \int_{\mathbf{r} \in D_m} \prod_n p[\dot{r}_n|m] d\mathbf{r} \\ &= 1 - \sum_m P\{m \text{ sent}\} \int_{\mathbf{r} \in D_m} \prod_{n=1}^N \frac{1}{\sqrt{2\pi\sigma^2}} \exp \frac{-[\dot{r}_n - E\dot{r}_n]^2}{2\sigma^2} d\mathbf{r} \\ &= 1 - \left(\frac{1}{\sqrt{2\pi\sigma^2}} \right)^N \sum_m P\{m \text{ sent}\} \int_{\mathbf{r} \in D_m} \exp \frac{-1}{2\sigma^2} \sum_{n=1}^N [\dot{r}_n - \dot{s}_n(m)]^2 d\mathbf{r} \end{aligned}$$

14.3 Frequency Shift Keying (FSK)

Theorem 14.3. In an FSK modulation space, the optimal ML estimator of m is

$$\text{T H M} \quad \hat{m} = \arg \max_m \dot{r}_m.$$

PROOF:

$$\begin{aligned} \hat{m} &= \arg \max_m P\{\mathbf{r}(t)|s(t; m)\} \\ &= \arg \min_m \sum_{n=1}^N [\dot{r}_n - \dot{s}_n(m)]^2 && \text{by Theorem 9.6 (page 80)} \\ &= \arg \min_m \sum_{n=1}^N [\dot{r}_n^2 - 2\dot{r}_n \dot{s}_n(m) + \dot{s}_n^2(m)] \\ &= \arg \min_m \sum_{n=1}^N [-2\dot{r}_n \dot{s}_n(m) + \dot{s}_n^2(m)] && \dot{r}_n^2 \text{ is independent of } m \\ &= \arg \min_m \sum_{n=1}^N [-2\dot{r}_n a \bar{\delta}_{mn} + a^2 \bar{\delta}_{mn}] \\ &= \arg \min_m [-2a \dot{r}_m + a^2] \end{aligned}$$

$$= \arg \min_m [-\dot{r}_m]$$

$$= \arg \max_m [\dot{r}_m]$$

a and 2 independent of m

⇒

Theorem 14.4. If an FSK modulation space let

$$\begin{array}{ll} z_2 & \triangleq \dot{r}_1(1) - \dot{r}_2(1) \\ z_3 & \triangleq \dot{r}_1(1) - \dot{r}_3(1) \\ \vdots & \\ z_M & \triangleq \dot{r}_1(1) - \dot{r}_M(1) \end{array} \quad \left| \begin{array}{l} z_2 > 0 \implies \dot{r}_1 > \dot{r}_2 \quad | \quad m = 1 \\ z_3 > 0 \implies \dot{r}_1 > \dot{r}_3 \quad | \quad m = 1 \\ \vdots \\ z_M > 0 \implies \dot{r}_1 > \dot{r}_M \quad | \quad m = 1 \end{array} \right.$$

Then the **probability of detection error** is

T H M $P\{\text{error}\} = 1 - \frac{M-1}{M} \int_0^\infty \int_0^\infty \cdots \int_0^\infty p(z_2, z_3, \dots, z_M) dz_2 dz_3 \cdots dz_M$ where

$$p(z_2, z_3, \dots, z_M) = \frac{1}{(2\pi)^{\frac{M-1}{2}} \sqrt{\det R}} \exp -\frac{1}{2} \begin{bmatrix} z_2 - \dot{s} \\ z_3 - \dot{s} \\ \vdots \\ z_M - \dot{s} \end{bmatrix}^T R^{-1} \begin{bmatrix} z_2 - \dot{s} \\ z_3 - \dot{s} \\ \vdots \\ z_M - \dot{s} \end{bmatrix}$$

and

$$R = \begin{bmatrix} \text{cov}[z_2, z_2] & \text{cov}[z_2, z_3] & \cdots & \text{cov}[z_2, z_M] \\ \text{cov}[z_3, z_2] & \text{cov}[z_3, z_3] & \cdots & \text{cov}[z_3, z_M] \\ \vdots & \vdots & \ddots & \vdots \\ \text{cov}[z_M, z_2] & \text{cov}[z_M, z_3] & \cdots & \text{cov}[z_M, z_M] \end{bmatrix} = N_o \begin{bmatrix} 2 & 1 & \cdots & 1 \\ 1 & 2 & \cdots & 1 \\ \vdots & \vdots & \ddots & \vdots \\ 1 & 1 & \cdots & 2 \end{bmatrix}$$

The inverse matrix R^{-1} is equivalent to (????)

$$R^{-1} \stackrel{?}{=} \frac{1}{MN_o} \begin{bmatrix} M-1 & -1 & \cdots & -1 \\ -1 & M-1 & \cdots & -1 \\ \vdots & \vdots & \ddots & \vdots \\ -1 & -1 & \cdots & M-1 \end{bmatrix}$$

PROOF:

$$\begin{aligned} E z_k &= E [\dot{r}_{11} - \dot{r}_{1k}] \\ &= E \dot{r}_{11} - E \dot{r}_{1k} \\ &= \dot{s} - 0 \\ &= \dot{s} \end{aligned}$$



$$\begin{aligned}
\text{cov}[z_m, z_n] &= \mathbb{E}[z_m z_n] - [\mathbb{E}z_m][\mathbb{E}z_n] \\
&= \mathbb{E}[(\dot{r}_{11} - \dot{r}_{1m})(\dot{r}_{11} - \dot{r}_{1n})] - \dot{s}^2 \\
&= \mathbb{E}[\dot{r}_{11}^2 - \dot{r}_{11}\dot{r}_{1n} - \dot{r}_{1m}\dot{r}_{11}\dot{r}_{1m}\dot{r}_{1n}] - \dot{s}^2 \\
&= [\text{var } \dot{r}_{11} + (\mathbb{E}\dot{r}_{11})^2] - \mathbb{E}[\dot{r}_{11}] \mathbb{E}[\dot{r}_{1n}] - \mathbb{E}[\dot{r}_{1m}] \mathbb{E}[\dot{r}_{11}] + [\text{cov}[\dot{r}_{1m}, \dot{r}_{1n}] + (\mathbb{E}\dot{r}_{1m})(\mathbb{E}\dot{r}_{1n})] - \dot{s}^2 \\
&= [\text{var } \dot{r}_{11} + \dot{s}^2] - a \cdot 0 - 0 \cdot a + [\text{cov}[\dot{r}_{1m}, \dot{r}_{1n}] + 0 \cdot 0] - \dot{s}^2 \\
&= \text{var } \dot{r}_{11} + \text{cov}[\dot{r}_{1m}, \dot{r}_{1n}] \\
&= N_o + \text{cov}[\dot{r}_{1m}, \dot{r}_{1n}] \\
&= \begin{cases} 2N_o & \text{for } m = n \\ N_o & \text{for } m \neq n. \end{cases}
\end{aligned}$$

$$P\{\text{error}\} = 1 - P\{\text{no error}\}$$

$$\begin{aligned}
&= 1 - \sum_{m=1}^M P\{\text{m transmitted}\} \wedge (\forall k \neq m, \dot{r}_m > \dot{r}_k) \\
&= 1 - (M-1)P\{1 \text{ transmitted}\} \wedge (\dot{r}_{11} > \dot{r}_{12}) \wedge (\dot{r}_{11} > \dot{r}_{13}) \wedge \dots \wedge (\dot{r}_{11} > \dot{r}_{1M}) \\
&= 1 - (M-1)P\{(\dot{r}_{11} - \dot{r}_{12} > 0) \wedge (\dot{r}_{11} - \dot{r}_{13} > 0) \wedge \dots \wedge (\dot{r}_{11} - \dot{r}_{1M} > 0) | 1 \text{ transmitted}\} P\{1 \text{ transmitted}\} \\
&= 1 - \frac{M-1}{M} P\{(z_2 > 0) \wedge (z_3 > 0) \wedge \dots \wedge (z_M > 0) | 1 \text{ transmitted}\} \\
&= 1 - \frac{M-1}{M} \int_0^\infty \int_0^\infty \dots \int_0^\infty p(z_2, z_3, \dots, z_M) dz_2 dz_3 \dots dz_M.
\end{aligned}$$



14.4 Quadrature Amplitude Modulation (QAM)

14.4.1 Receiver statistics




Figure 14.2: Distributions of QAM components

Theorem 14.5. Let $(V, \langle \cdot | \cdot \rangle)$ be a QAM modulation space such that

$$\begin{aligned}
\mathbf{r}(t) &= \mathbf{s}(t; m) + \mathbf{n}(t) \\
\dot{r}_c &\triangleq \langle \mathbf{r}(t) | \psi_c(t) \rangle \\
\dot{r}_s &\triangleq \langle \mathbf{r}(t) | \psi_s(t) \rangle.
\end{aligned}$$

Then $(\dot{r}_c|m)$ and $(\dot{r}_s|m)$ are **independent** and have **marginal distributions**

$$\begin{aligned} (\dot{r}_c|m) &\sim \mathcal{N}(a_m, \sigma^2) = \mathcal{N}(r_m \cos \theta_m, \sigma^2) \\ (\dot{r}_s|m) &\sim \mathcal{N}(b_m, \sigma^2) = \mathcal{N}(r_m \sin \theta_m, \sigma^2). \end{aligned}$$

PROOF: See Theorem 9.5 (page 79) page 79.

14.4.2 Detection

Theorem 14.6. Let $(V, \langle \cdot | \cdot \rangle, S)$ be a QAM modulation space with

$$\begin{aligned} \mathbf{r}(t) &= \mathbf{s}(t; m) + \mathbf{n}(t) \\ \dot{r}_c &\triangleq \langle \mathbf{r}(t) | \psi_c(t) \rangle \\ \dot{r}_s &\triangleq \langle \mathbf{r}(t) | \psi_s(t) \rangle. \end{aligned}$$

Then $\{\dot{r}_c, \dot{r}_s\}$ are sufficient statistics for optimal ML detection and the optimal ML estimate of m is

$$\hat{m}_{\text{ml}}[m] = \arg \min_m [(\dot{r}_c - a_m)^2 + (\dot{r}_s - b_m)^2].$$

PROOF:

$$\begin{aligned} \hat{m}_{\text{ml}}[m] &= \arg \max_m P\{\mathbf{r}(t)|\mathbf{s}(t; m)\} && \text{by Definition 8.1 (page 64)} \\ &= \arg \min_m \sum_{n=1}^N [\dot{r}_n - \dot{s}_n(m)]^2 && \text{by Theorem 9.6 (page 80)} \\ &= \arg \min_m [(\dot{r}_c - a_m)^2 + (\dot{r}_s - b_m)^2] \end{aligned}$$

14.4.3 Probability of error




Figure 14.3: QAM-16 constellation and decision regions

Theorem 14.7. In a QAM-16 constellation as shown in Figure 14.3 (page 130), the probability of error is

$$P\{\text{error}\} = \frac{9}{4}Q^2 \left(\frac{\dot{s}_{21} - \dot{s}_{11}}{2N_o} \right).$$

PROOF: Let

$$d \triangleq \dot{s}_{21} - \dot{s}_{11}.$$

Then

$$\begin{aligned} P\{\text{error}\} &= \sum_{m=1}^M P\{[s(t; m) \text{ transmitted}] \wedge [(\dot{r}_1, \dot{r}_2) \notin D_m]\} \\ &= \sum_{m=1}^M P\{[(\dot{r}_1, \dot{r}_2) \notin D_m] | [s(t; m) \text{ transmitted}]\} P\{[s(t; m) \text{ transmitted}]\} \\ &= \frac{1}{M} \sum_{m=1}^M P\{[(\dot{r}_1, \dot{r}_2) \notin D_m] | [s(t; m) \text{ transmitted}]\} \\ &= \frac{1}{M} [4P\{(\dot{r}_1, \dot{r}_2) \notin D_1 | s_1(t)\} + 8P\{(\dot{r}_1, \dot{r}_2) \notin D_2 | s_2(t)\} + 4P\{(\dot{r}_1, \dot{r}_2) \notin D_6 | s_6(t)\}] \\ &= \frac{1}{M} \left[4 \int \int_{(x,y) \notin D_1} p_{xy|1}(x, y) dx dy + 8 \int \int_{(x,y) \notin D_2} p_{xy|2}(x, y) dx dy + \right. \\ &\quad \left. 4 \int \int_{(x,y) \notin D_6} p_{xy|6}(x, y) dx dy \right] \\ &= \frac{1}{M} \left[4 \int \int_{(x,y) \notin D_1} p_{x|1}(x)p_{y|1}(y) dx dy + 8 \int \int_{(x,y) \notin D_2} p_{x|2}(x)p_{y|2}(y) dx dy + \right. \\ &\quad \left. 4 \int \int_{(x,y) \notin D_6} p_{x|6}(x)p_{y|6}(y) dx dy \right] \\ &= \frac{1}{M} \left[4Q\left(\frac{d}{2N_o}\right)Q\left(\frac{d}{2N_o}\right) + 8Q\left(\frac{d}{2N_o}\right)2Q\left(\frac{d}{2N_o}\right) + 4 \cdot 2Q\left(\frac{d}{2N_o}\right)2Q\left(\frac{d}{2N_o}\right) \right] \\ &= \frac{9}{4}Q^2\left(\frac{d}{2N_o}\right) \end{aligned}$$



14.5 Phase Shift Keying (PSK)

14.5.1 Receiver statistics




Figure 14.4: Distributions of PSK components

Theorem 14.8. Let

$$\begin{aligned}\dot{r}_c &\triangleq \langle \mathbf{r}(t) | \psi_c(t) \rangle \\ \dot{r}_s &\triangleq \langle \mathbf{r}(t) | \psi_s(t) \rangle \\ \theta_m &\triangleq \text{atan} \left[\frac{\dot{r}_s(m)}{\dot{r}_c(m)} \right].\end{aligned}$$

The statistics $(\dot{r}_c|m)$ and $(\dot{r}_s|m)$ are **independent** with marginal distributions

$$\begin{aligned}(\dot{r}_c|m) &\sim \mathcal{N}(r \cos \theta_m, \sigma^2) \\ (\dot{r}_s|m) &\sim \mathcal{N}(r \sin \theta_m, \sigma^2) \\ p_{\theta_m}(\theta|m) &= \int_0^\infty x p_{\dot{r}_c}(x|m) p_{\dot{r}_s}(x \tan \theta|m) dx.\end{aligned}$$

PROOF:

Independence and marginal distributions of $\dot{r}_1(m)$ and $\dot{r}_2(m)$ follow directly from Theorem 9.5 (page 79) (page 79).

Let $X \triangleq \dot{r}_1(m)$, $Y \triangleq \dot{r}_2(m)$ and $\Theta \triangleq \theta_m$. Then¹

$$\begin{aligned}p_\theta(\theta)d\theta &\triangleq P\{\theta < \Theta \leq \theta + d\theta\} \\ &= P\left\{\theta < \text{atan} \frac{Y}{X} \leq \theta + d\theta\right\} \\ &= P\left\{\tan(\theta) < \frac{Y}{X} \leq \tan(\theta + d\theta)\right\} \\ &= P\left\{\tan(\theta) < \frac{Y}{X} \leq \tan \theta + (1 + \tan^2 \theta) d\theta\right\} \\ &= \int_0^\infty P\left\{\left[\tan \theta < \frac{Y}{X} \leq \tan \theta + (1 + \tan^2 \theta) d\theta\right] \wedge \left[(x < X \leq x + dx)\right]\right\} \\ &= \int_0^\infty P\left\{\tan \theta < \frac{Y}{x} \leq \tan \theta + (1 + \tan^2 \theta) d\theta \mid x < X \leq x + dx\right\} P\{x < X \leq x + dx\} \\ &= \int_0^\infty P\left\{x \tan \theta < Y \leq x \tan \theta + x(1 + \tan^2 \theta) d\theta \mid X = x\right\} p_x(x) dx \\ &= \int_0^\infty [p_Y(x \tan \theta) x(1 + \tan^2 \theta)] p_x(x) dx d\theta \\ &= (1 + \tan^2 \theta) \int_0^\infty x p_Y(x \tan \theta) p_x(x) dx d\theta \\ &\implies p_\theta(\theta)d\theta = (1 + \tan^2 \theta) \int_0^\infty x p_Y(x \tan \theta) p_x(x) dx\end{aligned}$$

¹A similar example is in [Papoulis \(1991\)](#), page 138

14.5.2 Detection

Theorem 14.9. Let $(V, \langle \cdot | \cdot \rangle, S)$ be a PSK modulation space with

$$\begin{aligned} r(t) &= s(t; m) + n(t) \\ \dot{r}_c &\triangleq \langle r(t) | \psi_c(t) \rangle \\ \dot{r}_s &\triangleq \langle r(t) | \psi_s(t) \rangle. \end{aligned}$$

Then $\{\dot{r}_c, \dot{r}_s\}$ are sufficient statistics for optimal ML detection and the optimal ML estimate of m is

$$\hat{m}_{\text{ml}}[m] = \arg \min_m [(\dot{r}_1 - r \cos \theta_m)^2 + (\dot{r}_2 - r \sin \theta_m)^2].$$

PROOF:

$$\begin{aligned} \hat{m}_{\text{ml}}[m] &= \arg \max_m P\{r(t) | s(t; m)\} && \text{by Definition 8.1 (page 64)} \\ &= \arg \min_m \sum_{n=1}^N [\dot{r}_n - \dot{s}_n(m)]^2 && \text{by Theorem 9.6 (page 80)} \\ &= \arg \min_m [(\dot{r}_1 - r \cos \theta_m)^2 + (\dot{r}_2 - r \sin \theta_m)^2]. \end{aligned}$$

14.5.3 Probability of error




Figure 14.5: PSK-8 Decision regions

Theorem 14.10. The probability of error using PSK modulation is

$$P\{ \text{error} \} = M \left[1 - \int_{\frac{2\pi}{M} \left(m - \frac{3}{2} \right)}^{\frac{2\pi}{M} \left(m - \frac{1}{2} \right)} p_{\theta_1}(\theta) d\theta \right].$$

PROOF: See Figure 14.5 (page 133).

$$\begin{aligned}
 P\{\text{error}\} &= \sum_{m=1}^M P\{\text{error}|s(t; m) \text{ was transmitted}\} \\
 &= M P\{\text{error}|s_1(t) \text{ was transmitted}\} \\
 &= M \left[1 - \int_{\frac{2\pi}{M}(m-\frac{3}{2})}^{\frac{2\pi}{M}(m-\frac{1}{2})} p_{\theta_1}(\theta) d\theta \right].
 \end{aligned}$$

☞

14.6 Pulse Amplitude Modulation (PAM)

14.6.1 Receiver statistics




Figure 14.6: Distribution of PAM component

Theorem 14.11. Let $(V, \langle \cdot | \cdot \rangle)$ be a PAM modulation space such that

$$\begin{aligned}
 r(t) &= s(t; m) + n(t) \\
 \dot{r}_c &\triangleq \langle r(t) | \psi_c(t) \rangle \\
 \dot{r}_s &\triangleq \langle r(t) | \psi_s(t) \rangle.
 \end{aligned}$$

Then $(\dot{r}|m)$ has **distribution**

$$\dot{r}(m) \sim N(a_m, \sigma^2).$$

☞ PROOF: This follows directly from Theorem 9.5 (page 79) (page 79). ☞

14.6.2 Detection

Theorem 14.12. Let $(V, \langle \cdot | \cdot \rangle, S)$ be a PAM modulation space with

$$\begin{aligned}
 r(t) &= s(t; m) + n(t) \\
 \dot{r} &\triangleq \langle r(t) | \psi(t) \rangle.
 \end{aligned}$$

Then \dot{r} is a sufficient statistic for the optimal ML detection of m and the optimal ML estimate of m is

$$\hat{u}_{\text{ml}}[m] = \arg \min_m |\dot{r} - a_m|.$$

PROOF:

$$\begin{aligned}
 \hat{u}_{\text{ml}}[m] &= \arg \max_m P\{r(t)|a_m\} && \text{by Definition 8.1 (page 64)} \\
 &= \arg \min_m \sum_{n=1}^N [\dot{r}_n - \dot{s}_n(m)]^2 && \text{by Theorem 9.6 (page 80)} \\
 &= \arg \min_m |\dot{r} - \dot{s}(m)|^2 \\
 &= \arg \min_m |\dot{r} - \dot{s}(m)|
 \end{aligned}$$



14.6.3 Probability of error

Theorem 14.13. *The probability of detection error in a PAM modulation space is*

$$P\{\text{error}\} = 2 \frac{M-1}{M} Q\left[\frac{a_2 - a_1}{2\sqrt{N_o}}\right].$$

PROOF: Let $d \triangleq a_2 - a_1$ and $\sigma \triangleq \sqrt{\text{var } \dot{r}} = \sqrt{N_o}$. Also, let the decision regions D_m be as illustrated in Figure 14.7 (page 135). Then

$$\begin{aligned}
 P\{\text{error}\} &= \sum_{m=1}^M P\{s(t; m) \text{ sent} \wedge r \notin D_m\} \\
 &= \sum_{m=1}^M P\{\dot{r} \notin D_m | s(t; m) \text{ sent}\} P\{s(t; m) \text{ sent}\} \\
 &= \sum_{m=1}^M P\{\dot{r}_m \notin D_m\} \frac{1}{M} \\
 &= \frac{1}{M} \left(Q\left[\frac{d}{2\sigma}\right] + 2Q\left[\frac{d}{2\sigma}\right] + \dots + 2Q\left[\frac{d}{2\sigma}\right] + Q\left[\frac{d}{2\sigma}\right] \right) \\
 &= 2 \frac{M-1}{M} Q\left[\frac{d}{2\sigma}\right] \\
 &= 2 \frac{M-1}{M} Q\left[\frac{\dot{s}_2 - \dot{s}_1}{2\sqrt{N_o}}\right]
 \end{aligned}$$




Figure 14.7: 4-ary PAM in AWGN channel

CHAPTER 15

NETWORK DETECTION

15.1 Detection

For detection, we need

1. Cost function: for hard decisions, its range must be linearly ordered. For soft decisions, it can be a lattice.
2. system joint and marginal probabilities (for Bayesian detection)

15.2 Bayesian Estimation

Definition 15.1.

DEF	$H \triangleq \{h_1, h_2, h_3, \dots\}$	set of hypotheses
DEF	$D \triangleq \{D_1, D_2, D_3, \dots\}$	partition—decision regions
DEF	$X \triangleq \{X_1, X_2, X_3, \dots\}$	set of sensor inputs

$$\begin{aligned} cost(h; P) &= \min_D \sum_i P \{ [X \in D_i] \wedge [H \neq h_i] \} \\ &= \min_D \sum_i P \{ X \in D_i \mid H \neq h_i \} P \{ H \neq h_i \} \\ &= \min_D \sum_i \sum_{j \neq i} [1 - P \{ X \in D_i \mid H = h_j \}] \sum_{j \neq i} [1 - P \{ H = h_j \}] \end{aligned}$$

$$\hat{h} = \arg_h cost(h; P)$$

15.3 Joint Gaussian Model

Assume convexity ...

$$\begin{aligned}
 \mathbf{D} &= \arg_{\mathbf{D}} \min_{\mathbf{D}} \text{cost}(h; P) \\
 &= \arg_{\mathbf{D}} \left\{ \frac{\partial}{\partial \mathbf{D}} \sum_i \int_{D_i} p(\mathbf{x}|H \neq h_i) \underbrace{p(H \neq h_i)}_c d\mathbf{x} = 0 \right\} \\
 &= \arg_{\mathbf{D}} \left\{ \frac{\partial}{\partial \mathbf{D}} c \sum_i \int_{D_i} p(\mathbf{x}|H \neq h_i) d\mathbf{x} = 0 \right\} \\
 &= \arg_{\mathbf{D}} \left\{ \frac{\partial}{\partial \mathbf{D}} \sum_i \left[1 - \sum_{j \neq i} \int_{D_i} p(\mathbf{x}|H = h_i) d\mathbf{x} \right] = 0 \right\} \\
 &= \arg_{\mathbf{D}} \left\{ \frac{\partial}{\partial \mathbf{D}} \sum_i \left[1 - \sum_{j \neq i} \int_{D_i} \frac{1}{\sqrt{(2\pi)^n |\mathbf{M}|}} \exp -\frac{1}{2} (\mathbf{x} - \mathbb{E}\mathbf{x})^T \mathbf{M}^{-1} (\mathbf{x} - \mathbb{E}\mathbf{x}) d\mathbf{z} \right] = 0 \right\} \\
 &= \arg_{\mathbf{D}} \left\{ \sum_i \left[1 - \sum_{j \neq i} \frac{\partial}{\partial \mathbf{D}} \int_{D_i} \frac{1}{\sqrt{(2\pi)^n |\mathbf{M}|}} \exp -\frac{1}{2} (\mathbf{x} - \mathbb{E}\mathbf{x})^T \mathbf{M}^{-1} (\mathbf{x} - \mathbb{E}\mathbf{x}) d\mathbf{z} \right] = 0 \right\} \\
 &= \arg_{\mathbf{D}} \left\{ \sum_i \left[1 - \sum_{j \neq i} \left[\begin{array}{c} \frac{\partial}{\partial D_1} \\ \frac{\partial}{\partial D_2} \\ \vdots \\ \frac{\partial}{\partial D_n} \end{array} \right] \int_{D_i} \frac{1}{\sqrt{(2\pi)^n |\mathbf{M}|}} \exp -\frac{1}{2} (\mathbf{x} - \mathbb{E}\mathbf{x})^T \mathbf{M}^{-1} (\mathbf{x} - \mathbb{E}\mathbf{x}) d\mathbf{z} \right] = 0 \right\} \\
 &= \arg_{\mathbf{D}} \left\{ \sum_i \left[1 - \sum_{j \neq i} \left[\begin{array}{c} \frac{\partial y_1}{\partial x_1} & \frac{\partial y_2}{\partial x_1} & \dots & \frac{\partial y_m}{\partial x_1} \\ \frac{\partial y_1}{\partial x_2} & \frac{\partial y_2}{\partial x_2} & \dots & \frac{\partial y_m}{\partial x_2} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial y_1}{\partial x_n} & \frac{\partial y_2}{\partial x_n} & \dots & \frac{\partial y_m}{\partial x_n} \end{array} \right] \underbrace{\text{Jacobian matrix}}_{\text{J}} \right] \right\}
 \end{aligned}$$

For two variable Gaussian ...

$$\begin{aligned}
 \text{cost} &= \min_{\mathbf{D}} \sum_i \int_{D_i} p(\mathbf{x}|H \neq h_i) \underbrace{p(H \neq h_i)}_c d\mathbf{x} \\
 &= \min_{\mathbf{D}} c \sum_i \int_{D_i} p(\mathbf{x}|H \neq h_i) d\mathbf{x} \\
 &= \min_{\mathbf{D}} c \sum_i \left[1 - \sum_{j \neq i} \int_{D_i} p(\mathbf{x}|H = h_i) d\mathbf{x} \right]
 \end{aligned}$$

$$= \min_D c \sum_i \left[1 - \sum_{j \neq i} \int_{D_j} \frac{1}{2\pi\sqrt{|M|}} \exp\left(\frac{z_1^2 E[z_2 z_2] - 2z_1 z_2 E[z_1 z_2] + z_2^2 E[z_1 z_1]}{-2|M|}\right) dz \right]$$

15.4 2 hypothesis, 2 sensor detection


Theorem 15.1 (centralized case). Let $(\Omega, \mathbb{E}, \mathbb{P})$ be a probability space. Let $D \subseteq \mathbb{E}$ be the DECISION REGION indicating hypothesis $H = h_1$. Let $\pi_0 \triangleq \mathbb{P}\{H = h_0\}$ and $\pi_1 \triangleq \mathbb{P}\{H = h_1\}$.

THM	$D = \arg \min_D \left[\underbrace{\mathbb{P}\{(x, y) \in D H = h_0\} \pi_0}_{\text{error for } H = h_0} + \underbrace{\mathbb{P}\{(x, y) \in D^c H = h_1\} \pi_1}_{\text{error for } H = h_1} \right]$ $= \arg \min_D \left[\underbrace{\pi_0 \int_D p_0(x, y) dx dy}_{\text{error for } H = h_0} + \underbrace{\pi_1 \int_D p_1(x, y) dx dy}_{\text{error for } H = h_1} \right]$
-----	---

PROOF:

$$\begin{aligned} D &= \arg \min_D [\mathbb{P}\{\text{error}\}] && \text{by definition of decision region } D \\ &= \arg \min_D [\mathbb{P}\{\text{error} \wedge H = h_0\} + \mathbb{P}\{\text{error} \wedge H = h_1\}] \\ &= \arg \min_D [\mathbb{P}\{\text{error}|H = h_0\} \pi_0 + \mathbb{P}\{\text{error}|H = h_1\} \pi_1] \\ &= \arg \min_D [\mathbb{P}\{(x, y) \in D | H = h_0\} \pi_0 + \mathbb{P}\{(x, y) \in D^c | H = h_1\} \pi_1] \\ &= \arg \min_D \left[\pi_0 \int_D p_0(x, y) dx dy + \pi_1 \int_D p_1(x, y) dx dy \right] \end{aligned}$$

Example 15.1. In the centralized case, the decision regions D in the xy -plane can be any arbitrary shape, as illustrated to the right.



Definition 15.2.

DEF Let \mathbf{P}_x and \mathbf{P}_y be set projection operators such that $D_x \triangleq \mathbf{P}_x D$
 $D_y \triangleq \mathbf{P}_y D$

Proposition 15.1. Let $+$ represent MINKOWSKI ADDITION

PRP $D = D_x + D_y$

Theorem 15.2 (distributed AND case). Let $(\Omega, \mathbb{E}, \mathbb{P})$ be a probability space. Let $D \subsetneq \mathbb{E}$ be the DECISION REGION indicating hypothesis $H = h_1$. Let $\pi_0 \triangleq \mathbb{P}\{H = h_0\}$ and $\pi_1 \triangleq \mathbb{P}\{H = h_1\}$. Let $E \triangleq D^c$.

T H M


$$D = \arg \min_D \left(\begin{array}{l} \mathbb{P}\{x \in E, y \in E\} \{H = h_1\} \pi_1 + \\ \mathbb{P}\{x \in E, y \in D\} \{H = h_1\} \pi_1 + \\ \mathbb{P}\{x \in D, y \in E\} \{H = h_1\} \pi_1 + \\ \mathbb{P}\{x \in D, y \in D\} \{H = h_0\} \pi_0 \end{array} \right)$$

PROOF:

x	y	H	$x \wedge y$	
0	0	0	0	
0	1	0	0	
1	0	0	0	
1	1	0	1	error
0	0	1	0	error
0	1	1	0	error
1	0	1	0	error
1	1	1	1	

$$\begin{aligned} D &= \arg \min_D [\mathbb{P}\{\text{error}\}] && \text{by definition of decision region } D \\ &= \arg \min_D [\mathbb{P}\{\text{error} \wedge H = h_0\} + \mathbb{P}\{\text{error} \wedge H = h_1\}] \\ &= \arg \min_D [\mathbb{P}\{\text{error}|H = h_0\} \pi_0 + \mathbb{P}\{\text{error}|H = h_1\} \pi_1] \\ &= \arg \min_D \left(\begin{array}{l} \mathbb{P}\{x \in E_x, y \in E_y\} \{H = h_1\} \pi_1 + \\ \mathbb{P}\{x \in D_x, y \in E_y\} \{H = h_1\} \pi_1 + \\ \mathbb{P}\{x \in E_x, y \in D_y\} \{H = h_1\} \pi_1 + \\ \mathbb{P}\{x \in D_x, y \in D_y\} \{H = h_0\} \pi_0 \end{array} \right) \end{aligned}$$

Example 15.2. In the distributed AND case, the decision regions D in the xy -plane are only simple rectangular shapes, as illustrated to the right.



Proposition 15.2.

P R P In general, distributed AND detection is suboptimal.

PROOF: Because only rectangular decision regions are possible, detection is suboptimal.

Theorem 15.3.¹

T H M For the distributed AND detection

$$D_x = \left\{ x \mid \pi_0 \int_{D_y} p_0(x, y) dx dy \leq \pi_1 \int_{D_y} p_1(x, y) dx dy \right\}$$

¹ Willett et al. (2000), page 3268


PROOF:

$$\begin{aligned} D_x = \{x | y \in D_y\} &\implies P\{(x, y) | H = h_0\} \pi_0 \leq P\{(x, y) | H = h_1\} \pi_1 \\ &= \left\{ x | \pi_0 \int_{D_y} p_0(x, y) dx dy \leq \pi_1 \int_{D_y} p_1(x, y) dx dy \right\} \end{aligned}$$



Part IV

Appendices



APPENDIX A

PROBABILITY SPACE




Figure A.1: Lattice of mathematical spaces



“It is not certain that everything is certain.”
Blaise Pascal (1623–1662), mathematician ¹

¹ quote: http://en.wikiquote.org/wiki/Blaise_Pascal
image: http://en.wikipedia.org/wiki/Image:Blaise_pascal.jpg

A.1 Probability functions

Definition A.1. ² Let $(X, \vee, \wedge, \neg, 0, 1; \leq)$ be a LATTICE WITH NEGATION.

The function P is a **probability function** if

- | | | |
|-------------|--|-------------------------|
| D
E
F | (1). $P(1) = 1$ | (NORMALIZED) and |
| | (2). $P(x) \geq 0 \quad \forall x \in X$ | (NONNEGATIVE) and |
| | (3). $\bigwedge_{n=1}^{\infty} x_n = 0 \implies P\left(\bigvee_{n=1}^{\infty} x_n\right) = \sum_{n=1}^{\infty} P(x_n) \quad \forall x_n \in X$ | (σ -ADDITIVE) . |

Remark A.1. The advantage of this definition is that P is a *measure*, and hence all the power of measure theory is subsequently at one's disposal in using P . However, it has often been argued that the requirement of σ -additivity is unnecessary for a probability function. Even as early as 1930, de Finetti argued against it, in what became a kind of polite running debate with Fréchet.³ In fact, Kolmogorov himself provided some argument against σ -additivity when referring to the closely related *Axiom of Continuity* saying, "Since the new axiom is essential for infinite fields of probability only, it is almost impossible to elucidate its empirical meaning...For, in describing any observable random process we can obtain only finite fields of probability...." But in its support he added, "This limitation has been found expedient in researches of the most diverse sort."⁴

There are several other definitions of probability that only require *additivity* rather than σ -*additivity*. On a *Boolean lattice*, the **traditional probability** function is defined as⁵

- | | | |
|---|-------------------------------------|-------------------|
| (1). | $P(1) = 1$ | (normalized) and |
| (2). | $P(x) \geq 0 \quad \forall x \in X$ | (nonnegative) and |
| (3). $x \wedge y = 0 \implies P(x \vee y) = P(x) + P(y) \quad \forall x, y \in X$ | (additive) . | |

This definition implies (on a *Boolean lattice*) that

- | | | |
|--|--|---------------------|
| (a). | $P(0) = 0$ | (nondegenerate) and |
| (b). | $P(x) \leq 1 \quad \forall x \in X$ | (upper bounded) and |
| (c). | $P(x) = 1 - P(x^\perp) \quad \forall x \in X$ | and |
| (d). | $P(x \vee y) \leq P(x) + P(y) \quad \forall x, y \in X$ | (subadditive) and |
| (e). | $P(x \vee y) = P(x) + P(y) - P(x \wedge y) \quad \forall x, y \in X$ | and |
| (f). $x \leq y \implies P(x) \leq P(y) \quad \forall x, y \in X$ | (monotone) . | |

On a *distributive pseudocomplemented lattice*, the **generalized probability** function has been defined as⁶

- | | | |
|---|------------|---------------------|
| (1). | $P(0) = 0$ | (nondegenerate) and |
| (2). | $P(1) = 1$ | (normalized) and |
| (3). $0 \leq P(1) \leq 1$ | and | |
| (4). $P(x \vee y) = P(x) + P(y) - P(x \wedge y) \quad \forall x, y \in X$ | . . | |

On an *orthomodular lattice*, or a *finite modular lattice*, the **quantum probability** function is defined as⁷

- | | | |
|--|--------------|---------------------|
| (1). | $P(0) = 0$ | (nondegenerate) and |
| (2). | $P(1) = 1$ | (normalized) and |
| (3). $x \perp y \implies P(x \vee y) = P(x) + P(y) \quad \forall x, y \in X$ | (additive) . | |

However, for lattices that are not *distributive*, *modular*, or *orthomodular*, none of these definitions

² Billingsley (1995) pages 22–23 (Probability Measures), Kolmogorov (1933a), Kolmogorov (1933b), page 16 (*field of probability*), Pap (1995) pages 8–9 (Definition 2.3(13)), Kalmbach (1986) page 27

³ de Finetti (1930a), Fréchet (1930a), de Finetti (1930b), Fréchet (1930b), de Finetti (1930c), Cifarelli and Regazzini (1996) pages 258–260

⁴ Kolmogorov (1933b), page 15

⁵ Papoulis (1991) pages 21–22, Kolmogorov (1933b), page 2 (§1. Axioms I–V)

⁶ Narens (2014) page 118, Narens (2007)

⁷ Greechie (1971) page 126 (DEFINITIONS), Narens (2014) page 118

work out so well. Take for example the O_6 lattice with the “very reasonable” probability function given in Example ?? (page ??). This probability space (O_6, P) fails to be any of the 4 probability functions defined in this Remark. It fails to be a *measure-theoretic* or *traditional probability* function because

$$a \wedge b = 0 \quad \text{but} \quad P(a \vee b) = P(1) = 1 \neq \frac{1}{3} + \frac{1}{2} = P(a) + P(b).$$

It fails to be a *generalized probability* function because

$$P(a \vee b) = P(1) = 1 \neq \frac{1}{3} + \frac{1}{2} - 0 = P(a) + P(b) - P(0) = P(a) + P(b) - P(a \wedge b).$$

It fails to be an *quantum probability* function because

$$a \perp b = 0 \quad \text{but} \quad P(a \vee b) = P(1) = 1 \neq \frac{1}{3} + \frac{1}{2} = P(a) + P(b).$$

In each of these cases, the function P fails to be *additive*. The solution of Definition A.1 (page 146) is simply to “switch off” *additivity* when the lattice is not *distributive*. This method is a little “crude”, but at least it allows us to define probability on a very wide class of lattices, while retaining compatibility with the *Boolean* case.

A.2 Probability Space

In mathematics, a *space* is simply a set and in the most general definition, nothing else. However, normally for a space to actually be useful, some additional structure is added. One of the most general additional structures is a *topology*; and a space together with a topology is called a *topological space*. A topological space imposes additional structure on a space in the form of subsets and guarantees that these subsets are closed under such fundamental operations as set *union* and set *intersection*. With the additional structure available in a topological space, we are able to analyze such basic concepts as continuity, convergence, and connectivity.

However for a great number of mathematical applications, we need to *measure* mathematical objects—the most general measurement being measures on subsets of some set. Examples of measurement in mathematics include integration and probability. Before measurement can be effectively performed on a set, the set must be equipped with a subset structure. In analysis, arguably the most fundamental subset structure is the humble *topology* (Definition ?? page ??). However, a simple topology does not provide sufficient structure for effective measurement. For example, often we would not only like to measure some subset A , but also its complement A^c . A topology is not closed under the complement operation. So instead of a topology only, we equip the space with a more powerful (and thus less general) structure called a σ -*algebra* (*sigma-algebra*) (Definition ?? page ??). A σ -*algebra* is a subset structure that is closed under set complement. A set together with a σ -*algebra* is called a *measurable space*. And a set together with a σ -*algebra* and a *measure* on that σ -*algebra* is called a *measure space* (Definition ?? page ??).

The next definition presents a very important measure space—the *probability space*.

Definition A.2.

D E F The triple (Ω, \mathbb{E}, P) is a **probability space** if

- (1). Ω is a SET
- (2). \mathbb{E} is a σ -ALGEBRA on Ω (Definition ?? page ??) and
- (3). $P : \mathbb{E} \rightarrow [0, 1]$ is a MEASURE on \mathbb{E} (Definition ?? page ??) .

If $S \triangleq (\Omega, \mathbb{E}, P)$ is a PROBABILITY SPACE then x is an **outcome** in S if $x \in \Omega$, A is an **event** in S if $A \in \mathbb{E}$, and PA is the **probability** of A in S if A is an EVENT in S .

Definition A.3.⁸ Let (Ω, \mathbb{E}, P) be a PROBABILITY SPACE (Definition A.2 page 147).

⁸  Papoulis (1990) page 52 (Independent Events)

D E F Two EVENTS A and B in \mathbb{E} are **independent** if
 $P(A \cap B) = P(A)P(B)$

Definition A.4. ⁹ Let (Ω, \mathbb{E}, P) be a PROBABILITY SPACE (Definition A.2 page 147). Let x and y be EVENTS in \mathbb{E} .

D E F The **conditional probability** of x given y is defined as
 $P(A|B) \triangleq \frac{P(A \cap B)}{P(B)}$

A.3 Properties

Proposition A.1.

P R P (Ω, \mathbb{E}, P) is a PROBABILITY SPACE \implies (Ω, \mathbb{E}, P) is a MEASURE SPACE
(every probability space is a measure space)

Theorem A.1. ¹⁰ Let (Ω, \mathbb{E}, P) be a PROBABILITY SPACE (Definition A.2 page 147).

T H M

- (1). $0 \leq P(x) \leq 1 \quad \forall x \in X$ (BOUNDED) and
- (2). $P(x) = 1 - P(x^\perp) \quad \forall x \in X$ (PARTITION OF UNITY) and
- (3). $x \leq y \implies P(y^\perp) \leq P(x^\perp) \quad \forall x, y \in X$ (ANTITONE)

PROOF:

1. Proof for $0 \leq P(x) \leq 1$:

$$\begin{aligned} 0 &= P(0) && \text{by by nondegenerate property of } P \text{ (Definition A.2 page 147)} \\ &\leq P(x) && \text{because } 0 \leq x \text{ and monotone property of } P \\ &\leq P(1) && \text{because } x \leq 1 \text{ and monotone property of } P \\ &= 1 && \text{by normalized property of } P \end{aligned}$$

2. Proof for $P(x) = 1 - P(x^\perp)$:

(a) Proof that P is *additive* (Definition A.2 page 147) over $\{0, x, x^\perp\} \subseteq X$:

- i. $\{0, x, x^\perp\}$ is *distributive*.
- ii. $x \wedge x^\perp = 0$ for all $x \in X$ by the *non-contradiction* property of *orthocomplemented lattices*.
- iii. Therefore, by Definition A.2, P is *additive* over $\{0, x, x^\perp\}$.

(b) Then ...

$$\begin{aligned} 1 - P(x^\perp) &= P(1) - P(x^\perp) && \text{by normalized property of } P && \text{(Definition A.2 page 147)} \\ &= P(x \vee x^\perp) - P(x^\perp) && \text{by excluded middle property of ortho. lat.} \\ &= P(x) + P(x^\perp) - P(x^\perp) && \text{by additive property of } (\Omega, \mathbb{E}, P) && \text{(item (2a) page 148)} \\ &= P(x) && \text{by field property of } (\mathbb{R}, +, \cdot, 0, 1) \end{aligned}$$

3. Proof for $x \leq y \implies P(y^\perp) \leq P(x^\perp)$:

$$\begin{aligned} x \leq y &\implies y^\perp \leq x^\perp && \text{by antitone property of orthocomplemented lattices} \\ &\implies P(y^\perp) \leq P(x^\perp) && \text{by monotone property of } P \end{aligned}$$

(Definition A.2 page 147)

⁹  Papoulis (1990) page 45 (2-3 Conditional Probability and Independence)

¹⁰ property (1):  Papoulis (1991) page 21 ((2-11))

Theorem A.2. ¹¹ Let (Ω, \mathbb{E}, P) be a PROBABILITY SPACE (Definition A.2 page 147).

T H M	L is BOOLEAN (Definition ?? page ??)	$\left\{ \begin{array}{l} 1. \quad P(x \vee y) = P(x) + P(y) - P(x \wedge y) \quad \forall x, y \in X \quad \text{and} \\ 2. \quad P(x \vee y) \leq P(x) + P(y) \quad \forall x, y \in X \quad (\text{BOOLE'S INEQUALITY}) \end{array} \right\}$
-------------	---	--

PROOF:

1. lemma: Proof that $P((\neg x) \wedge y) = P(y) - P(x \wedge y)$:

$$\begin{aligned} P(y) - P(xy) &= P(1 \wedge y) - P(xy) && \text{by definition of } 1 \text{ and } \wedge \\ &= P[(x \vee x^\perp)y] - P(xy) && \text{by excluded middle property of Boolean lattices} \\ &= P(xy \vee x^\perp y) - P(xy) && \text{by distributive property of Boolean lattices} \\ &= P(xy) + P(x^\perp y) - P(xy) && \text{because } (xy)(x^\perp y) = 0 \text{ and by additive property} \\ &= P(x^\perp y) \end{aligned}$$

2. Proof that $P(x \vee y) = P(x) + P(y) - P(x \wedge y)$:

$$\begin{aligned} P(x \vee y) &= P(x \vee x^\perp y) && \text{by property of Boolean lattices} \\ &= P(x) + P(x^\perp y) && \text{because } (x)(x^\perp y) = 0 \text{ and by additive property} \\ &= P(x) + P(y) - P(x \wedge y) && \text{by item (1) (page 149)} \end{aligned}$$

Theorem A.3 (sum of products). Let $(X, \vee, \wedge, 0, 1 ; \leq)$ be a BOUNDED LATTICE, (Ω, \mathbb{E}, P) a PROBABILITY SPACE (Definition A.2 page 147), and $\{y, x_1, x_2, x_3, \dots\}$ a subset of X .

T H M	$\left\{ \begin{array}{l} 1. \quad L \text{ is DISTRIBUTIVE} \\ 2. \quad \{x_1, x_2, \dots\} \text{ is a PARTITION of } y \end{array} \right. \text{ and} \right\} \Rightarrow \left\{ \begin{array}{l} 1. \quad P(y) = \sum_n P(x_n) \quad \text{and} \\ 2. \quad P(y) = \sum_n P(y \wedge x_n) \quad \text{and} \\ 3. \quad P(z \wedge y) = \sum_n P(z \wedge x_n) \end{array} \right\}$
-------------	--

PROOF:

1. Proof that P is *additive* (Definition A.2 page 147) on (Ω, \mathbb{E}, P) :

(a) Proof that $(yx_n) \wedge (yx_m) = 0$ for $n \neq m$:

$$\begin{aligned} (yx_n) \wedge (yx_m) &= y(x_n x_m) && \text{by definition of } \wedge \\ &= y \wedge 0 && \text{by mutually exclusive property of partitions} \\ &= 0 && \text{by lower bounded property of bounded lattices} \end{aligned}$$

(b) Proof that L is *distributive*: by *distributive hypothesis*

2. Proof that $P(y) = \sum_n P(x_n)$

$$\begin{aligned} P(y) &= P(yx_1 \vee yx_2 \vee \dots \vee yx_n) && \text{by item (1) and additive property} \\ &= \sum_n P(yx_n) && \text{by item (1) and additive property} \quad (\text{Definition A.2 page 147}) \\ &= \sum_n P(y|x_n)P(x_n) && \text{by conditional probability} \quad (\text{Definition A.4 page 148}) \end{aligned}$$

¹¹  Papoulis (1991) page 21 ((2-13)),  Feller (1970) pages 22–23 ((7.4),(7.6))




Figure A.2: Double coin toss (Example A.2 page 150)

As described in Definition A.2 (page 147), every *probability space* (Ω, \mathbb{E}, P) contains a probability *measure* $P : \mathbb{E} \rightarrow [0, 1]$. This probability *measure* has some basic properties as described in Theorem A.4 (next).

Theorem A.4. Let (Ω, \mathbb{E}, P) be a PROBABILITY SPACE. Let B be a set and $\{B_n | n = 1, 2, \dots, N\}$ a set of sets.

T H M

$$\left\{ \begin{array}{l} \{B_n | n = 1, 2, \dots, N\} \text{ is a} \\ \text{PARTITION of } B. \end{array} \right\} \implies \left\{ \begin{array}{l} (1). \quad P(B) = \sum_{n=1}^N P(B_n) \quad \forall B \in \mathbb{E} \quad \text{and} \\ (2). \quad P(AB) = \sum_{n=1}^N P(AB_n) \quad \forall A, B \in \mathbb{E} \end{array} \right\}$$


PROOF: P is a *measure* and by Definition ?? (page ??).

Proposition A.2. Let (Ω, \mathbb{E}, P) be a probability space, and X a RANDOM VARIABLE with PROBABILITY DENSITY FUNCTION $p_x(x)$ and CUMULATIVE DISTRIBUTION FUNCTION $c_x(x)$.

- P R P**
- (1). $c_x(x)$ is MONOTONE and
 - (2). $p_x(x)$ is CONTINUOUS $\implies c_x(x)$ is STRICTLY MONOTONE and
 - (3). $p_x(x)$ is CONTINUOUS $\implies c_x(x)$ is INVERTIBLE

A.4 Examples

Example A.1 (single coin toss). Let \circledcirc represent “heads” and \bullet represent “tails” in a coin toss. Let $0 < p < 1$ be the probability of a head. A *probability space* (Ω, \mathbb{E}, P) for a single coin toss is as follows:



Example A.2 (Double coin toss). Let \circledcirc represent “heads” and \bullet represent “tails” in a double coin toss in which each toss is *independent* (Definition A.3 page 147) of the other. Let $0 < p < 1$ be the probability of a head. The *probability space* (Ω, \mathbb{E}, P) is illustrated in Figure A.2 (page 150).

PROOF:

$$\begin{aligned}
 P(\Omega) &= 1 && \text{by } \textit{normalized} \text{ property of } P && (\text{Definition A.1 page 146}) \\
 P(C) &= P\{\odot\odot\} && \text{by definition of } C \\
 &= P(\odot)P(\odot) && \text{by definition of } \textit{independence} && (\text{Definition A.3 page 147}) \\
 &= p^2 && \text{by definition of } p \\
 P(A) &= P\{\bullet\bullet\} && \text{by definition of } A \\
 &= P(\bullet)P(\bullet) && \text{by definition of } \textit{independence} && (\text{Definition A.3 page 147}) \\
 &= \{1 - P(\odot)\}\{1 - P(\odot)\} && \text{by } \textit{antitone} \text{ property of } P && (\text{Theorem A.1 page 148}) \\
 &= (1 - p)^2 && \text{by definition of } p \\
 P(B) &= P\{(\bullet\odot), (\odot\bullet)\} && \text{by definition of } B \\
 &= P\{\bullet\odot\} + P\{\odot\bullet\} && \text{by } \textit{additive} \text{ property of } P && (\text{Definition A.1 page 146}) \\
 &= P(\bullet)P(\odot) + P(\odot)P(\bullet) && \text{by definition of } \textit{independence} && (\text{Definition A.3 page 147}) \\
 &= (1 - p)p + p(1 - p) && \text{by } \textit{antitone} \text{ property of } P && (\text{Theorem A.1 page 148}) \text{ and definition of } p \\
 &= -2p^2 + p + 1 \\
 P(A \cup B) &= P(A) + P(B) - P(A \cap B) && \text{by Theorem A.2} \\
 &= P(A) + P(B) - P(\emptyset) \\
 &= (1 - p)^2 + (-2p^2 + p + 1) + 0 && \text{by previous results} \\
 &= -p^2 - p + 1 \\
 P(\emptyset) &= 0 && \text{by } \textit{nondegenerate} \text{ property of } P && (\text{Definition A.1 page 146})
 \end{aligned}$$

EX

$$\begin{aligned}
 \Omega &= \{\square, \square, \square, \square, \square, \square\} \\
 \Xi &= \left\{ \underbrace{\{\}}_{\emptyset}, \underbrace{\{\square, \square, \square\}}_{\text{odd}}, \underbrace{\{\square, \square, \square, \square\}}_{\text{even}}, \underbrace{\{\square, \square, \square, \square, \square, \square\}}_{\Omega} \right\} \\
 P(X) &= \left\{ \begin{array}{lll} 0 & \text{for } e = \{\} & (\emptyset) \\ 1 & \text{for } e = \{\square, \square, \square, \square, \square, \square\} & (\Omega) \\ p & \text{for } e = \{\square, \square, \square\} & (\text{odd}) \\ 1 - p & \text{for } e = \{\square, \square, \square, \square\} & (\text{even}) \end{array} \right\}
 \end{aligned}$$




Figure A.3: even/odd die probability space (Example A.3 page 151)

Example A.3 (even/odd die toss). The *probability space* for an **even/odd die toss**, with $0 < p < 1$ being the probability of the die toss being odd, is illustrated in Figure A.3 (page 151).

PROOF:

$$\begin{aligned}
 P(\Omega) &= 1 && \text{by } \textit{normalized} \text{ property of } P && (\text{Definition A.1 page 146}) \\
 P(C) &= P\{\odot\odot\} && \text{by definition of } C \\
 &= P(\odot)P(\odot) && \text{by definition of } \textit{independence} && (\text{Definition A.3 page 147}) \\
 &= p^2 && \text{by definition of } p \\
 P(A) &= P\{\square, \square, \square\} && \text{by definition of } A \\
 &= p && \text{by definition of } p \\
 P(B) &= P\{\square, \square, \square, \square\} && \text{by definition of } B \\
 &= P\{\square, \square, \square, \square\}^c && \text{by definition of set complement } c \\
 &= PA^c && \text{by definition of } A
 \end{aligned}$$

$= P(\neg A)$	by definition of \neg	
$= 1 - P(A)$	by Theorem A.1 page 148	
$= 1 - p$	by definition of p	
$P(\emptyset) = 0$	by <i>nondegenerate</i> property of P	(Definition A.1 page 146)



The two previous *even/odd die* example (Example A.5 page 152) is in essence the same as the *single coin toss* (Example A.1 page 150). The next offers a little more complexity.

E X		$P(\emptyset) = 0 \quad P(\Omega) = 1$ $P(A) = (1-p)^2 \quad P(A \cup B) = (1-p)^2 + 2p(1-p)$ $P(B) = 2p(1-p) \quad P(A \cup C) = 2p^2 - 2p + 1$ $P(C) = p^2 \quad P(B \cup C) = -p^2 + 2p$
--------	--	--

Figure A.4: 3-4-2 die example (Example A.4 page 152)

Example A.4. Suppose we have a “fair” die and we are primarily interested in the events of the first three $\{\square, \square, \square\}$, the next two, $\{\square, \square\}$ and the final one $\{\square\}$. The resulting *probability space* is illustrated in Figure A.4 (page 152).

The two previous examples (Example A.5 page 152, Example A.4 page 152) illustrate a *probability spaces* in which the events are *mutually exclusive*. The (next) illustrates one where events are *not*.

Example A.5. Suppose we have a “fair” die and we are primarily interested in the events of the first four ($\{\square, \square, \square, \square\}$) (that is, whether one roll of the die will produce a value in the set $\{\square, \square, \square, \square\}$) and the last three ($\{\square, \square, \square\}$). However, these events do not by themselves form a σ -algebra. Rather under the \cap and \cup operations, these two events generate a total of eight possible events that together form a σ -algebra. The resulting *probability space* is illustrated in Figure A.5 (page 153).

But why go through all the trouble of requiring a σ -algebra? Having a σ -algebra in place ensures that anything we might possibly want to measure *can* be measured. It makes sure all possible combinations are taken into account. And why go through the additional trouble of requiring a measure space? With a measure space available, expressing the measure over a complex set is often greatly simplified because the measure space provides nice algebraic properties (namely the σ -additive property). Example A.6 (next) illustrates how a rather complex σ -algebra (64 elements) can be compactly represented in a measure space.

Example A.6. Suppose we have a “fair” dice and we are interested in measuring over the power set of events (largest possible algebra— $2^6 = 64$ events). This leads to the probability space (Ω, \mathbb{E}, P) where

E X	$\Omega = \{\square, \square, \square, \square, \square, \square, \square, \square\}$ $\mathbb{E} = \mathcal{P}(\Omega)$ (the power-set of Ω) $P(e) = \frac{1}{6} e $ ($\frac{1}{6}$ times the number of possible outcomes in event e)
--------	--

Example A.7 (Gaussian distribution on \mathbb{R}). Let \mathcal{B} be the *Borel algebra* on \mathbb{R} . Let $\mathcal{L} \triangleq (\mathcal{B}, \subseteq)$ be the lattice formed by the elements of \mathcal{B} —this lattice is a *Boolean algebra*. Let

$$P(A) \triangleq \frac{1}{\sqrt{2\pi\sigma^2}} \int_A e^{-\frac{x^2}{2\sigma^2}} dx \text{ for } A \subseteq \mathcal{B}$$

and where \int is the *Lebesgue integral* (Definition ?? page ??). Then (\mathcal{L}, P) is a **probability space**.

E X

$$\Omega = \{\square, \square, \square, \square, \square, \square\}$$

$$\mathbb{E} = \left\{ \underbrace{\{\}}_{\emptyset}, \underbrace{\{\square, \square, \square, \square, \square, \square\}}_{\Omega}, \underbrace{\{\square, \square, \square, \square, \square\}}_{\text{first four}}, \underbrace{\{\square, \square, \square, \square\}}_{\text{last three}}, \right.$$

$$\left. \underbrace{\{\square\}}_{\{1234\} \cap \{456\}}, \underbrace{\{\square, \square, \square, \square, \square, \square\}}_{\{4\}^c}, \underbrace{\{\square, \square\}}_{\{4\}^c \cap \{456\}}, \underbrace{\{\square, \square, \square\}}_{\{1234\} \cap \{4\}^c} \right\}$$

$$\mathbb{P}(e) = \begin{cases} 0 & \text{for } e = \{\} \\ 1 & \text{for } e = \{\square, \square, \square, \square, \square, \square\} \\ \frac{1}{2} & \text{for } e = \{\square, \square, \square, \square, \square\} \\ \frac{1}{3} & \text{for } e = \{\square, \square, \square, \square\} \\ \frac{1}{6} & \text{for } e = \{\square, \square, \square\} \\ \frac{1}{6} & \text{for } e = \{\square, \square, \square, \square, \square\} \\ \frac{1}{3} & \text{for } e = \{\square, \square, \square\} \\ \frac{1}{2} & \text{for } e = \{\square, \square, \square, \square\} \end{cases}$$

Figure A.5: First 4 / last 3 die example (Example A.5 page 152)

Example A.8 (Gaussian noise). Let $X \sim N(0, \sigma^2)$ be a random variable with Gaussian distribution. We can construct the following probability space (Ω, \mathbb{E}, P) :

E X

$$\Omega = \mathbb{R}$$

$$\mathbb{E} = \{\emptyset, \Omega\} \cup \{(a, b) | a, b \in \mathbb{R}, a < b\}$$

$$\mathbb{P}_x = \begin{cases} 0 & \text{for } x = \emptyset \\ 1 & \text{for } x = \Omega \\ \frac{1}{\sqrt{2\pi\sigma^2}} \int_a^b e^{-\frac{x^2}{2\sigma^2}} dx & \text{otherwise} \end{cases}$$

Example A.9. The set of outcomes Ω can also be a set of waveforms:

E X

$$\Omega = \left\{ \begin{array}{c} \square \square \square \square \square \square \\ \square \square \square \square \square \square \\ \square \square \square \square \square \square \end{array} \right\}$$

$$\mathbb{E} = \mathcal{P}(\Omega)$$

$$\mathbb{P}(e) = \frac{1}{7}|e|$$

A.5 Probability subspaces

Example A.10. Suppose a random process is capable of producing three values $\Omega \triangleq \{x, y, z\}$. There are five *algebras of sets* on Ω and therefore five probability spaces $(\Omega, \mathbb{E}_n, P)$ on Ω with the five values of \mathbb{E}_n listed below:¹²

¹² algebra of sets: Definition ?? page ??





Figure A.6: Euclidean 3-dimensional space partitioned as a power lattice



$$\begin{aligned}
 E_1 &= \{ \emptyset, X \} \\
 E_2 &= \{ \emptyset, \{x\}, \{y, z\}, X \} \\
 E_3 &= \{ \emptyset, \{y\}, \{x, z\}, X \} \\
 E_4 &= \{ \emptyset, \{z\}, \{x, y\}, X \} \\
 E_5 &= \{ \emptyset, \{x\}, \{y\}, \{z\}, \{x, y\}, \{x, z\}, \{y, z\}, X \}
 \end{aligned}$$

Suppose the samples of Ω are generated by a physical process such that they are all equally likely to occur. Then by varying the σ -algebra E_n effectively varies the probability distribution of the probability space (Ω, E_n, P) . This is illustrated by the figure to the left.

APPENDIX B

PROBABILITY DENSITY FUNCTIONS



“While writing my book I had an argument with Feller. He asserted that everyone said “random variable” and I asserted that everyone said “chance variable.” We obviously had to use the same name in our books, so we decided the issue by a stochastic procedure. That is, we tossed for it and he won.”¹

Joseph Leonard Doob (1910–2004), pioneer of and key contributor to mathematical probability¹

B.1 Random variables

The concept of the *random variable* is widely used in probability and random processes. Before discussing what a *random variable* is, note two things that a *random variable* is *not* (next remark).

Remark B.1. ² As pointed out by others, the term “random variable” is a “misnomer”:

**R
E
M**

- A *random variable* is **not random**.
- A *random variable* is **not a variable**.

What is it then? It is a *function* (next definition). In particular, it is a function that maps from an underlying stochastic process into \mathbb{R} . Any “randomness” (whatever that means) it may appear to have comes from the stochastic process it is mapping *from*. But the function itself (the *random*

¹ quote: [Snell \(1997\)](#), page 307, [Snell \(2005\)](#), page 251

image: <http://www.dartmouth.edu/~chance/Doob/conversation.html>

² [Miller \(2006\)](#) page 130, [Feldman and Valdez-Flores \(2010\)](#) page 4 (“The name “random variable” is actually a misnomer, since it is not random and not a variable....the *random variable* simply maps each point (outcome) in the sample space to a number on the real line...Technically, the space into which the *random variable* maps the sample space may be more general than the real line...”), [Curry and Feldman \(2010\)](#) page 4, [Trivedi \(2016\)](#) page 2.1 (“The term “random variable” is actually a misnomer, since a *random variable* X is really a function whose domain is the sample space S , and whose range is the set of all real numbers, \mathbb{R} .”)

variable itself) is very deterministic and well-defined. What gives it the appearance of being random is that the outcome ω of the experiment appears to be random to the observer. So the *random variable* $X(\omega)$ is simply a function of an underlying mechanism that appears to be random.

Definition B.1. ³ Let (Ω, \mathbb{E}, P) be a PROBABILITY SPACE (Definition A.2 page 147).

DEF

A **random variable** X is any function in \mathbb{R}^{Ω} .

B.2 Probability distributions

The probability information about σ -algebra \mathbb{E} in a *probability space* (Definition A.2 page 147) is completely specified by *measure* P . However, sometimes it is more convenient to express this same *measure* information in terms of the *probability density function* or the *cummulative distribution function* of the *probability space*.

Definition B.2. ⁴ Let X be a RANDOM VARIABLE (Definition B.1 page 156) on a PROBABILITY SPACE (Ω, \mathbb{E}, P) .

DEF

X has **cummulative distribution function** (cdf) $c_X(x)$ if

$$c_X(x) \triangleq P\{x \in \mathbb{E} | X < x\}$$

X **probability density function** (pdf) $p_X(x)$ if

$$p_X(x) \triangleq \frac{d}{dx} c_X(x) \triangleq \frac{d}{dx} P\{x \in \mathbb{E} | X < x\}$$

Remark B.2. Suppose X be a *random variable* on a *probability space* (Ω, \mathbb{E}, P) . Note that

- Both X and \mathbb{E} are *functions*.
- But X is a function that maps from Ω to \mathbb{R} ,
- whereas P is a function that maps from \mathbb{E} to \mathbb{R} .

Definition B.3. Let (Ω, \mathbb{E}, P) be a PROBABILITY SPACE (Definition A.2 page 147) and X and $Y : \Omega \rightarrow \mathbb{R}$ random variables. Then a **joint probability density function** $p_{XY} : \mathbb{E} \times \Omega \rightarrow [0 : 1]$ and a **joint cumulative distribution function** $c_{XY} : \mathbb{E} \times \Omega \rightarrow [0 : 1]$ are defined as

DEF

$$c_{XY}(x, y) \triangleq P\{X \leq x | Y \leq y\} \quad (\text{JOINT CUMULATIVE DISTRIBUTION FUNCTION})$$

$$p_{XY}(x, y) \triangleq \frac{d}{dy} \frac{d}{dx} c_{XY}(x, y) \quad (\text{JOINT PROBABILITY DENSITY FUNCTION})$$

Definition B.4. Let (Ω, \mathbb{E}, P) be a PROBABILITY SPACE (Definition A.2 page 147) and X a random variable. Then a **conditional probability density function** $p_X : \mathbb{E} \times \Omega \rightarrow [0 : 1]$ and a **conditional cumulative distribution function** $c_X : \mathbb{E} \times \Omega \rightarrow [0 : 1]$ are defined as

DEF

$$c_X(x|y) \triangleq P\{X \leq x | Y = y\} \quad (\text{CONDITIONAL CUMULATIVE DISTRIBUTION FUNCTION—CDF})$$

$$p_X(x|y) \triangleq \frac{d}{dx} c_X(x|y) \quad (\text{CONDITIONAL PROBABILITY DENSITY FUNCTION—PDF})$$

B.3 Properties

Definition B.2 (page 156) defines the pdf and cdf of a *probability space* (Ω, \mathbb{E}, P) in terms of *measure* P . Conversely, the probability *measure* $P\{a \leq X < b\}$ of an event $\{a \leq X < b\}$ can be expressed in terms of either the pdf or cdf.

³  Papoulis (1991), page 63

⁴  von der Linden et al. (2014) page 93 (Definitions 7.1, 7.2)

Proposition B.1. Let X a RANDOM VARIABLE with PDF p_x and CDF c_x (Definition B.2 page 156) on the PROBABILITY SPACE (Ω, \mathbb{E}, P) (Definition A.2 page 147).

P R P

$$\left\{ \begin{array}{l} (1). c_x(x) \text{ and } c_y(y) \text{ are CONTINUOUS OR} \\ (2). p_x(x) \text{ and } p_y(y) \text{ are CONTINUOUS} \end{array} \right\} \implies \left\{ \begin{array}{l} p_x(x) = \lim_{\epsilon \rightarrow 0} \frac{1}{\epsilon} P\{x \leq X < x + \epsilon\} \\ p_{xy}(x, y) = \lim_{\epsilon \rightarrow 0} \frac{1}{\epsilon} P\{x \leq X < x + \epsilon \wedge y \leq Y < y + \epsilon\} \end{array} \right\}$$

PROOF:

$$\begin{aligned} p_x(x) &\triangleq \frac{d}{dx} c_x(x) && \text{by definition of } p_x && (\text{Definition B.2 page 156}) \\ &= \lim_{\epsilon \rightarrow 0} \frac{1}{\epsilon} P\{x \in \mathbb{R} | x \leq X < x + \epsilon\} && \text{by definition of } \frac{d}{dx} && (\text{Definition ?? page ??}) \end{aligned}$$



Theorem B.1. Let (Ω, \mathbb{E}, P) be a probability space, X be a random variable, and (a, b) a real interval.

T H M

$$\left\{ \begin{array}{l} (1). c_x(x) \text{ is CONTINUOUS OR} \\ (2). p_x(x) \text{ is CONTINUOUS} \end{array} \right\} \implies \left\{ P\{a < X \leq b\} = c_x(b) - c_x(a) = \int_a^b p_x(x) dx \right\}$$

PROOF:

$$\begin{aligned} P\{a < X \leq b\} &= P\{X \leq b\} - P\{X < a\} && \text{by sum of products} && (\text{Theorem A.3 page 149}) \\ &= P\{X \leq b\} - P\{X \leq a\} && \text{by continuity hypothesis} \\ &\triangleq c_x(b) - c_x(a) && \text{by definition of } c_x && (\text{Definition B.2 page 156}) \end{aligned}$$

$$\begin{aligned} \int_a^b p_x(x) dx &\triangleq \int_a^b \left[\frac{d}{dx} c_x(x) \right] dx && \text{by definition of } p_x && (\text{Definition B.2 page 156}) \\ &= c_x(x)|_{x=b} - c_x(x)|_{x=a} && \text{by Fundamental theorem of calculus} \\ &= c_x(b) - c_x(a) \end{aligned}$$



Theorem B.2. Let (Ω, \mathbb{E}, P) be a PROBABILITY SPACE, X be a RANDOM VARIABLE, and $(a : b)$ a REAL INTERVAL.

T H M

$$P\{a \leq X < b\} = \int_a^b p_x(x) dx = \int_{-\infty}^b c_x(x) dx - \int_{-\infty}^a c_x(x) dx$$

The properties of the pdf follow closely the properties of measure P .

Theorem B.3. ⁵

T H M

$$\left\{ \begin{array}{l} (A). c_x(x) \text{ is CONTINUOUS OR} \\ (B). p_x(x) \text{ is CONTINUOUS} \end{array} \right\} \implies \left\{ \begin{array}{l} (1). p_{x|y}(x|y) = \frac{p_{xy}(x,y)}{p_y(y)} \text{ and} \\ (2). p_x(x) = \int_{y \in \mathbb{R}} p_{xy}(x,y) dy \end{array} \right\}$$

⁵ Papoulis (1990) page 158 (Auxiliary Variable)

PROOF:

$$\begin{aligned}
 p_{X|Y}(x|y) &\triangleq \frac{d}{dx} c_{X|Y}(x|y) && \text{by definition of } c_X \quad (\text{Definition A.4 page 148}) \\
 &\triangleq \lim_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon} P\{x \leq X < x + \varepsilon | Y = y\} && \text{by definition of } \frac{d}{dx} \quad (\text{Definition ?? page ??}) \\
 &\triangleq \lim_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon} \frac{P\{(x \leq X < x + \varepsilon) \wedge (Y = y)\}}{P\{Y = y\}} && \text{by definition of } P\{A|B\} \quad (\text{Definition A.4 page 148}) \\
 &= \lim_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon} \frac{P\{(x \leq X < x + \varepsilon) \wedge (y \leq Y < y + \varepsilon)\}}{P\{y \leq Y < y + \varepsilon\}} && \text{by continuity hypothesis} \\
 &= \frac{\lim_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon} P\{(x \leq X < x + \varepsilon) \wedge (y \leq Y < y + \varepsilon)\}}{\lim_{\varepsilon \rightarrow 0} P\{y \leq Y < y + \varepsilon\}} && \text{by property of } \lim_{\varepsilon \rightarrow 0} \\
 &= \frac{p_{XY}(x, y)}{p_Y(y)} && \text{by Proposition B.1 page 157}
 \end{aligned}$$

$$\begin{aligned}
 \int_{y \in \mathbb{R}} p_{XY}(x, y) dy &\triangleq \int_{y \in \mathbb{R}} \left[\frac{d}{dy} \frac{d}{dx} c_{XY}(x, y) \right] dy && \text{by definition of } p_X \quad (\text{Definition B.2 page 156}) \\
 &= \frac{d}{dx} c_{XY}(x, y) && \\
 &\triangleq \lim_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon} \int_{y \in \mathbb{R}} P\{x \leq X < x + \varepsilon, y \leq Y < y + \varepsilon\} dy && \text{by definition of } \frac{d}{dx} \quad (\text{Definition ?? page ??}) \\
 &= \lim_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon} P\{x \leq X < x + \varepsilon\} && \\
 &= p_X(x) && \text{by Proposition B.1 page 157}
 \end{aligned}$$

⇒

Theorem B.4.

T	$c_X(\sup \mathbb{R}) = 1$
H	$c_X(\inf \mathbb{R}) = 0$

PROOF:

$$\begin{aligned}
 c_X(\sup \mathbb{R}) &\triangleq P\{X \leq \sup \mathbb{R}\} && \text{by definition of } c_X \quad (\text{Definition B.2 page 156}) \\
 &= 1 \\
 c_X(\inf \mathbb{R}) &\triangleq P\{X \leq \inf \mathbb{R}\} && \text{by definition of } c_X \quad (\text{Definition B.2 page 156}) \\
 &= 0
 \end{aligned}$$

⇒

The properties of the pdf follow closely the properties of measure P.

Theorem B.5.

T	$c_{X Y}(x y) = \frac{\frac{d}{dy} c_{XY}(x, y)}{p_Y(y)}$	$p_{X Y}(x y) = \frac{p_{XY}(x, y)}{p_Y(y)}$
---	---	--



PROOF:

$$\begin{aligned}
 c_{X|Y}(x|y) &\triangleq P\{X \leq x | Y = y\} && \text{by definition of } c_{X|Y} && (\text{Definition B.4 page 156}) \\
 &\triangleq \frac{P\{X \leq x | Y = y\}}{P\{Y = y\}} && \text{by definition of } P\{X|Y\} && (\text{Definition A.4 page 148}) \\
 &= \lim_{\epsilon \rightarrow 0} \frac{P\{X \leq x | y < Y \leq y + \epsilon\}}{P\{y < Y \leq y + \epsilon\}} \\
 &= \lim_{\epsilon \rightarrow 0} \frac{[P\{X \leq x | Y \leq y + \epsilon\} - P\{X \leq x | Y \leq y\}]/\epsilon}{[P\{Y \leq y + \epsilon\} - P\{Y \leq y\}]/\epsilon} \\
 &\triangleq \lim_{\epsilon \rightarrow 0} \frac{[c_{XY}(x, y + \epsilon) - c_{XY}(x, y)]/\epsilon}{[c_Y(y + \epsilon) - c_Y(y)]/\epsilon} && \text{by definition of } c_{XY} && (\text{Definition B.3 page 156}) \\
 &\triangleq \frac{\frac{d}{dy}c_{XY}(x, y)}{\frac{d}{dy}c_Y(y)} && \text{by definition of } \frac{d}{dy}f(y) \\
 &\triangleq \frac{\frac{d}{dy}c_{XY}(x, y)}{p_Y(y)} && \text{by definition of } p_Y && (\text{Definition B.2 page 156}) \\
 &= \frac{\frac{d}{dy}c_{XY}(x, y)}{p_Y(y)} && \text{because } y \text{ is fixed}
 \end{aligned}$$

$$\begin{aligned}
 p_{X|Y}(x|y) &\triangleq \frac{d}{dx}c_{X|Y}(x|y) && \text{by definition of } p_{X|Y} && (\text{Definition B.4 page 156}) \\
 &= \frac{d}{dx} \frac{\frac{d}{dy}c_{XY}(x, y)}{p_Y(y)} && \text{by previous result} \\
 &= \frac{\frac{d}{dx}\frac{d}{dy}c_{XY}(x, y)}{p_Y(y)} && \text{because } p_Y(y) \text{ is not a function of } x \\
 &\triangleq \frac{p_{XY}(x, y)}{p_Y(y)} && \text{by definition of } p_{XY}(x, y) && (\text{Definition B.3 page 156})
 \end{aligned}$$



Theorem B.6. Let (Ω, \mathbb{E}, P) be a probability space.

T H M	$\int_{x \in \mathbb{R}} p_X(x) dx = 1$	$\int_{x \in \mathbb{R}} p_{X Y}(x y) dx = 1$
	$\int_{y \in \mathbb{R}} p_{XY}(x, y) dy = p_X(x) \quad \forall x \in \Omega$	$\int_{x \in \mathbb{R}} \int_{y \in \mathbb{R}} p_{XY}(x, y) dy dx = 1$

PROOF:

$$\begin{aligned}
 \int_{\mathbb{R}} p_X(x) dx &= c_X(\sup \mathbb{R}) - c_X(\inf \mathbb{R}) && \text{by Theorem B.1 page 157} \\
 &= 1 - 0 \\
 &= 1 && \text{because 0 is the additive identity element in } (\mathbb{R}, +, \cdot, 0, 1) \\
 \int_{x \in \mathbb{R}} p_{X|Y}(x|y) dx &\triangleq \int_{x \in \mathbb{R}} \frac{d}{dx}c_{X|Y}(x|y) dx && \text{by definition of } p_{X|Y}(x|y) (\text{Definition B.4 page 156}) \\
 &= c_{X|Y}(\sup \mathbb{R}|y) - c_{X|Y}(\inf \mathbb{R}|y) && \text{by Fundamental theorem of calculus}
 \end{aligned}$$

$$= 1 - 0$$

$$= 1$$

because 0 is the additive identity element in $(\mathbb{R}, +, \cdot, 0, 1)$

$$\int_{y \in \mathbb{R}} p_{XY}(x, y) dy = \int_{y \in \mathbb{R}} p_{YX}(y, x) dy$$

$$= \int_{y \in \mathbb{R}} p_{Y|X}(y|x) p_X(x) dy$$

by Theorem B.5 page 158

$$= p_X(x) \int_{y \in \mathbb{R}} p_{Y|X}(y|x) dy$$

because $p_X(x)$ is not a function of y

$$= p_X(x) \cdot 1$$

by previous result

$$= p_X(x)$$

because 1 is the multiplicative identity element in $(\mathbb{R}, +, \cdot, 0, 1)$

$$\int_{x \in \mathbb{R}} \int_{y \in \mathbb{R}} p_{XY}(x, y) dy dx = \int_{x \in \mathbb{R}} p_X(x) dx$$

by previous result

$$= 1$$

by previous result



APPENDIX C

SOME PROBABILITY DENSITY FUNCTIONS

C.1 Discrete distributions

*Example C.1.*¹ Suppose we throw two “fair” dice and want to know the probabilities of their sum. Let X represent the sum of the face values of the two dice. The resulting probability distribution is illustrated in Figure C.1 (page 162) and has probability space as follows:

EX	$\Omega = \{\square\square, \square\bullet, \bullet\square, \dots, \bullet\bullet\}$
	$\mathbb{E} = \{2^{\{X=n n=2,3,\dots,10,11, \text{ or } 12\}}\}$
	$P(e) = \frac{1}{36} e $

C.2 Continuous distributions

C.2.1 Uniform distribution

Definition C.1. The **uniform distribution** $p_x(x)$ is defined as

DEF	$p_x(x) \triangleq \begin{cases} 1 & \text{for } 0 < x \leq 1 \\ 0 & \text{otherwise} \end{cases}$
-----	--

Note that although “simple” in form, in light of *Wold's Theorem*, the value of the *uniform distribution* should *not* be taken lightly.

¹  Osgood (2002)




Figure C.1: Probability distribution for two dice (see Example C.1 page 161)


C.2.2 Gaussian distribution

“Tout le monde y croit cependant, me disait un jour M. Lippmann, car les expérimentateurs s’irrégularisent que c’est un théorème de mathématiques, et les mathématiciens que c’est un fait expérimental.”



“Everyone believes in it [(the normal distribution)] however, said to me one day Mr. Lippmann, because the experimenters imagine that it is a theorem of mathematics, and mathematicians that it is an experimental fact.”²

Bernard A. Lippmann as told by Henri Poincaré ²

Figure C.2: Gaussian pdf with $\mu = 0$ and $\sigma \in [0.1, 2]$.

Definition C.2. The **Gaussian distribution** (or **normal distribution**) has pdf

D E F $p_x(x) \triangleq \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{(x-\mu)^2}{2\sigma^2}}$ A random variable X with this distribution is denoted

D E F $X \sim N(\mu, \sigma^2)$ The function $Q(x)$ is defined as the area under a Gaussian PDF with zero mean

² quote: Poincaré (1912), page 171
translation: assisted by Google Translate
image:

and variance equal to one from x to infinity such that

D E F

$$Q(x) \triangleq \frac{1}{\sqrt{2\pi}} \int_x^{\infty} e^{-\frac{u^2}{2}} du$$

C.2.3 Gamma distribution

Definition C.3. ³ Let $b \in \mathbb{R}$. The **gamma function** $\Gamma(b)$ is

D E F

$$\Gamma(b) \triangleq \int_0^{\infty} x^{b-1} e^{-x} dx$$

Proposition C.1. ⁴ Let $b \in \mathbb{R}$ and $n \in \mathbb{N}$.

P R P

$$\begin{aligned}\Gamma(b) &= (b-1)\Gamma(b-1) \\ \Gamma(n) &= (n-1)!\end{aligned}$$

PROOF: Let

$$\begin{aligned}u &= x^{b-1} & du &= (b-1)x^{b-2} dx \\ dv &= e^{-x} dx & v &= -e^{-x}\end{aligned}$$

$$\begin{aligned}\Gamma(b) &\triangleq \int_0^{\infty} x^{b-1} e^{-x} dx \\ &= \int_{x=0}^{\infty} u dv \\ &= uv \Big|_{x=0}^{\infty} - \int_{x=0}^{\infty} v du \\ &= -x^{b-1} e^{-x} \Big|_{x=0}^{\infty} + (b-1) \int_{x=0}^{\infty} e^{-x} x^{b-1} dx \\ &= (-0+0) + (b-1)\Gamma(b-1)\end{aligned}$$

Note that

$$\Gamma(1) = \int_0^{\infty} x^{1-1} e^{-x} dx = \int_0^{\infty} e^{-x} dx = -e^{-x} \Big|_0^{\infty} = -0+1 = 1$$

$$\begin{aligned}\Gamma(n) &= (n-1)\Gamma(n-1) \\ &= (n-1)(n-2)\Gamma(n-2) \\ &= (n-1)(n-2)(n-3)\Gamma(n-3) \\ &\vdots \\ &= (n-1)(n-2)(n-3) \cdots (1)\Gamma(1) \\ &= (n-1)(n-2)(n-3) \cdots (1) \\ &\triangleq (n-1)!\end{aligned}$$

Definition C.4. A **Gamma distribution** (b, λ) has pdf

D E F

$$p_x(x) \triangleq \frac{\lambda}{\Gamma(b)} e^{-\lambda x} (\lambda x)^{b-1}$$

³ Papoulis (1991), page 79, Ross (1998), page 222

⁴ Ross (1998), page 223

Theorem C.1. ⁵ Let X and Y be RANDOM VARIABLES on a PROBABILITY SPACE $(\Omega, \mathbb{E}, \mathbb{P})$.

T H M	$\left\{ \begin{array}{ll} (A). & X \text{ and } Y \text{ are INDEPENDENT} \\ (B). & X \text{ has GAMMA DISTRIBUTION } (a, \lambda) \quad \text{and} \\ (C). & Y \text{ has GAMMA DISTRIBUTION } (b, \lambda) \quad \text{and} \\ (D). & Z \triangleq X + Y \end{array} \right\} \Rightarrow \left\{ \begin{array}{l} Z \text{ has Gamma distribution} \\ (a + b, \lambda). \end{array} \right\}$
-------------	---

PROOF:

$$p_Z(z) = p_X(z) \star p_Y(z)$$

$$= \int_{u \in \mathbb{R}} p_X(u)p_Y(z-u) du \quad \text{by definition of convolution (Definition N.3 page 312)}$$

$$= \int_0^z \frac{1}{\Gamma(a)} \lambda e^{-\lambda u} (\lambda u)^{a-1} \frac{1}{\Gamma(b)} \lambda e^{-\lambda(z-u)} (\lambda(z-u))^{b-1} du$$

$$= \frac{1}{\Gamma(a)} \frac{1}{\Gamma(b)} e^{-\lambda z} \lambda^{1+1+a-1+b-1} \int_0^z u^{a-1} (z-u)^{b-1} du$$

$$= \frac{1}{\Gamma(a)} \frac{1}{\Gamma(b)} \lambda e^{-\lambda z} \lambda^{a+b-1} \int_0^1 (vz)^{a-1} (z-vz)^{b-1} z dv$$

$$= \frac{1}{\Gamma(a)} \frac{1}{\Gamma(b)} \lambda e^{-\lambda z} \lambda^{a+b-1} z^{a-1+b-1+1} \int_0^1 v^{a-1} (1-v)^{b-1} dv$$

$$= \left[\frac{1}{\Gamma(a)} \frac{1}{\Gamma(b)} \int_0^1 v^{a-1} (1-v)^{b-1} dv \right] \lambda e^{-\lambda z} (\lambda z)^{a+b-1}$$

$$= C \lambda e^{-\lambda z} (\lambda z)^{a+b-1}$$

$$= \frac{\lambda}{\Gamma(a+b)} e^{-\lambda z} (\lambda z)^{a+b-1}$$

$\Rightarrow p_Z(z)$ is a $(a+b, \lambda)$ Gamma distribution

where C is some constant

C must be the value that makes $\int_z p_Z(z) = 1$

C.2.4 Chi-squared distributions

Definition C.5. ⁶ Let $p(x)$ be a PROBABILITY DENSITY FUNCTION on a PROBABILITY SPACE $(\Omega, \mathbb{E}, \mathbb{P})$.

D E F $p(x)$ is a **chi-square distribution** if

$$p(x) \triangleq \left\{ \begin{array}{ll} 0 & \text{if } x < 0 \\ \frac{1}{\sqrt{2\pi\sigma^2 x}} \exp\left[-\frac{x}{2\sigma^2}\right] & \text{if } x \geq 0 \end{array} \right\} \quad \text{for } \sigma > 0$$

Theorem C.2. ⁷

The following distributions are equivalent:

- | | |
|-------------|---|
| T
H
M | <ol style="list-style-type: none"> (1). chi-squared distribution and (2). distribution of X^2 where $X \sim N(0, \sigma^2)$ and (3). Gamma distribution $\left(\frac{1}{2}, \frac{1}{2\sigma^2}\right)$ |
|-------------|---|

PROOF:

⁵ Ross (1998), page 266

⁶ Proakis (2001), page 41, Papoulis (1990) page 219 (7-4 Special Distributions of Statistics, (7-78))

⁷ Ross (1998), page 267

1. Proof that χ^2 has chi-squared distribution:

$$\begin{aligned}
 p_Y(y) &= \frac{1}{2\sqrt{y}} \left[p_X(-\sqrt{y}) + p_X(\sqrt{y}) \right] && \text{by Corollary 4.3 page 29} \\
 &= \frac{1}{2\sqrt{y}} \left[\frac{1}{\sqrt{2\pi\sigma^2}} \exp -\frac{(-\sqrt{y}-0)^2}{2\sigma^2} + \frac{1}{\sqrt{2\pi\sigma^2}} \exp -\frac{(+\sqrt{y}-0)^2}{2\sigma^2} \right] \\
 &= \frac{1}{2\sqrt{y}} \left[2 \frac{1}{\sqrt{2\pi\sigma^2}} \exp -\frac{y}{2\sigma^2} \right] \\
 &= \frac{1}{\sqrt{2\pi\sigma^2}y} \exp -\frac{y}{2\sigma^2}
 \end{aligned}$$

2. Proof that chi-distribution is a Gamma distribution (b, λ) :

$$\begin{aligned}
 b &\triangleq \frac{1}{2} \\
 \lambda &\triangleq \frac{1}{2\sigma^2}
 \end{aligned}$$

$$\begin{aligned}
 \frac{1}{\sqrt{2\pi\sigma^2}y} \exp -\frac{y}{2\sigma^2} &= \frac{1}{\sqrt{\pi}} \lambda^{1/2} \lambda^{1/2} (\lambda y)^{-1/2} e^{-\lambda y} \\
 &= \frac{\lambda}{\sqrt{\pi}} (\lambda y)^{b-1} e^{-\lambda y}
 \end{aligned}$$



Definition C.6. ⁸ The **Chi-squared distribution with n degrees of freedom** has pdf

D E F

$$p_Y(y) \triangleq \begin{cases} 0 & : y < 0 \\ \frac{1}{2\sigma^2\Gamma(n/2)} \left(\frac{y}{2\sigma^2} \right)^{\frac{n}{2}-1} \exp -\frac{y}{2\sigma^2} & : y \geq 0 \end{cases}$$

Theorem C.3. ⁹ The following distributions are equivalent:

1. chi-squared distribution with n degrees of freedom

2. the distribution of $\sum_{k=1}^n X_k^2$ where $\{X_k | X_k \sim N(0, \sigma^2), k = 1, 2, \dots, n\}$ are independent random variables.

3. Gamma distribution $\left(\frac{n}{2}, \frac{1}{2\sigma^2}\right)$.

PROOF:

1. Prove chi-squared distribution with n degrees of freedom is the Gamma distribution $\left(\frac{n}{2}, \frac{1}{2\sigma^2}\right)$:

$$\begin{aligned}
 \lambda &\triangleq \frac{1}{2\sigma^2} \\
 b &\triangleq \frac{1}{2} \\
 \frac{1}{2\sigma^2\Gamma(n/2)} \left(\frac{y}{2\sigma^2} \right)^{\frac{n}{2}-1} \exp -\frac{y}{2\sigma^2} &= \frac{\lambda}{\Gamma(nb)} (\lambda y)^{nb-1} \exp -\lambda y
 \end{aligned}$$

⁸ Proakis (2001), page 41

⁹ Ross (1998), page 267

2. Prove $\sum_{k=1}^n X^2$ is Gamma $\left(\frac{n}{2}, \frac{1}{2\sigma^2}\right)$:

(a) By Theorem C.2, X_k has Gamma distribution $\left(\frac{1}{2}, \frac{1}{2\sigma^2}\right)$.

(b) By Theorem C.1, $\sum_{k=1}^n X_k^2$ has distribution $\left(\frac{n}{2}, \frac{1}{2\sigma^2}\right)$.

Definition C.7. ¹⁰ A **noncentral chi-square distribution** (μ, σ^2) has pdf

$$\text{DEF} \quad p_Y(y) = \frac{1}{\sqrt{2\pi\sigma^2 y}} \exp \frac{y + \mu^2}{-2\sigma^2} \cosh \frac{\mu\sqrt{y}}{\sigma^2}$$

Theorem C.4.

T **H** **M** The following distributions are equivalent:

- (1). NON-CENTRAL CHI-SQUARED DISTRIBUTION (μ, σ^2)
- (2). distribution of X^2 where $X \sim N(\mu, \sigma^2)$

PROOF:

1. Proof that $Y = X^2$ has a non-central chi-squared distribution:

$$\begin{aligned} p_Y(y) &= \frac{1}{2\sqrt{y}} \left[p_X(-\sqrt{y}) + p_X(\sqrt{y}) \right] \quad \text{by Corollary 4.3 page 29} \\ &= \frac{1}{2\sqrt{y}} \left[\frac{1}{\sqrt{2\pi\sigma^2}} \exp -\frac{(-\sqrt{y} - \mu)^2}{2\sigma^2} + \frac{1}{\sqrt{2\pi\sigma^2}} \exp -\frac{(+\sqrt{y} - \mu)^2}{2\sigma^2} \right] \\ &= \frac{1}{2\sqrt{y}} \left[\frac{1}{\sqrt{2\pi\sigma^2}} \exp -\frac{y + \mu^2}{2\sigma^2} \exp \frac{-2\mu\sqrt{y}}{2\sigma^2} + \frac{1}{\sqrt{2\pi\sigma^2}} \exp -\frac{y + \mu^2}{2\sigma^2} \exp \frac{2\mu\sqrt{y}}{2\sigma^2} \right] \\ &= \frac{1}{\sqrt{2\pi\sigma^2 y}} \exp -\frac{y + \mu^2}{2\sigma^2} \frac{1}{2} \left[\exp \frac{2\mu\sqrt{y}}{2\sigma^2} + \exp \frac{-2\mu\sqrt{y}}{2\sigma^2} \right] \\ &= \frac{1}{\sqrt{2\pi\sigma^2 y}} \exp \frac{y + \mu^2}{-2\sigma^2} \cosh \frac{\mu\sqrt{y}}{\sigma^2} \end{aligned}$$

Definition C.8. ¹¹ The α th-order modified Bessel function of the first kind $I_\alpha(x)$ is

$$\text{DEF} \quad I_\alpha(x) = \sum_{k=0}^{\infty} \frac{1}{k! \Gamma(\alpha + k + 1)} \left(\frac{x}{2}\right)^{\alpha+2k}$$

Definition C.9. ¹² The **noncentral chi-square with n -degrees of freedom** distribution has pdf

$$\text{DEF} \quad p_Y(y) = \frac{1}{2\sigma^2} \left(\frac{y}{s^2}\right)^{\frac{n-2}{4}} \exp \frac{y + s^2}{-2\sigma^2} I_{n/2-1} \left(\sqrt{y} \frac{s}{\sigma^2}\right) \quad \text{where } s^2 \triangleq \sum_{k=1}^n \mu_k^2$$

¹⁰ Proakis (2001), page 42

¹¹ Proakis (2001), page 43

¹² Proakis (2001), page 43




Figure C.3: *Joint Gaussian distributions $p_{xy}(x, y)$ with varying correlations*

C.2.5 Radial distributions

Definition C.10. ¹³ *The Rayleigh distribution is the pdf*

$$\text{DEF } p_R(r) = \begin{cases} 0 & \text{for } r < 0 \\ \frac{r}{\sigma^2} \exp -\frac{r^2}{2\sigma^2} & \text{for } r \geq 0 \end{cases}$$

Note that by Proposition 4.3, this distribution is equivalent to the distribution of $R = \sqrt{X^2 + Y^2}$ where X and Y are independent random variables each with distribution $N(0, \sigma^2)$.

Definition C.11. ¹⁴ *The Rice distribution is the pdf*

$$\text{DEF } p_R(r) = \begin{cases} 0 & \text{for } r < 0 \\ \frac{r}{\sigma^2} \exp \frac{r^2+s^2}{-2\sigma^2} I_0 \left(\frac{rs}{\sigma^2} \right) & \text{for } r \geq 0 \end{cases}$$

C.3 Joint Gaussian distributions

Definition C.12 (Joint Gaussian pdf). ¹⁵

$$\text{DEF } p(x_1, x_2, \dots, x_n) \triangleq \frac{1}{\sqrt{(2\pi)^n |\mathbf{M}|}} \exp -\frac{1}{2}(\mathbf{x} - \mathbf{Ex})^T \mathbf{M}^{-1} (\mathbf{x} - \mathbf{Ex}) \quad (\text{Gaussian joint pdf})$$

$$\mathbf{x} \triangleq \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}$$

$$Z_k \triangleq X_k - \mathbf{E}X_k$$

(zero mean random variables)

$$\mathbf{M} \triangleq \begin{bmatrix} E[Z_1 Z_1] & E[Z_1 Z_2] & \cdots & E[Z_1 Z_n] \\ E[Z_2 Z_1] & E[Z_2 Z_2] & \cdots & E[Z_2 Z_n] \\ \vdots & \vdots & \ddots & \vdots \\ E[Z_n Z_1] & E[Z_n Z_2] & \cdots & E[Z_n Z_n] \end{bmatrix}$$


(correlation matrix)

Example C.2 (1 variable joint Gaussian pdf). The **Gaussian distribution** (or **normal distribution**) has pdf

¹³ Proakis (2001), page 44

¹⁴ Proakis (2001), page 46

¹⁵ Proakis (2001), page 49, Moon and Stirling (2000), page 34

Figure C.4: Gaussian pdf with $\mu = 0$ and $\sigma \in [0.1, 2]$.

E X

$$p_x(x) \triangleq \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{(x-\mu)^2}{2\sigma^2}}$$

$$\begin{aligned} t &= \arg_t \min_t \left[\frac{1}{2} \int_t^\infty \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{(x-\mu)^2}{2\sigma^2}} + \frac{1}{2} \int_{-\infty}^t \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{(x-\eta)^2}{2\sigma^2}} \right] \\ &= \arg_t \left\{ \frac{\partial}{\partial t} \left[\frac{1}{2} \int_t^\infty \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{(x-\mu)^2}{2\sigma^2}} + \frac{1}{2} \int_{-\infty}^t \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{(x-\eta)^2}{2\sigma^2}} \right] = 0 \right\} \\ &= \arg_t \left\{ \frac{1}{2\sqrt{2\pi\sigma^2}} \left[\frac{\partial}{\partial t} \int_t^\infty e^{-\frac{(x-\mu)^2}{2\sigma^2}} + \frac{\partial}{\partial t} \int_{-\infty}^t e^{-\frac{(x-\eta)^2}{2\sigma^2}} \right] = 0 \right\} \\ &= \arg_t \left\{ \left[\left(e^{-\frac{(\infty-\mu)^2}{2\sigma^2}} 0 - e^{-\frac{(t-\mu)^2}{2\sigma^2}} 1 \right) + \left(e^{-\frac{(t-\mu)^2}{2\sigma^2}} 1 - e^{-\frac{(\infty-\mu)^2}{2\sigma^2}} 0 \right) \right] = 0 \right\} \\ &= \arg_t \left\{ \left[e^{-\frac{(t-\eta)^2}{2\sigma^2}} - e^{-\frac{(t-\mu)^2}{2\sigma^2}} \right] = 0 \right\} \\ &= \arg_t \{ (t - \eta)^2 = (t - \mu)^2 \} \\ &= \frac{\mu + \eta}{2} \end{aligned}$$

Example C.3 (2 variable joint Gaussian pdf).

E X

$$\begin{aligned} z_1 &\triangleq x_1 - \mathbb{E}x_1 \\ z_2 &\triangleq x_2 - \mathbb{E}x_2 \\ |M| &\triangleq |\mathbb{E}[z_1z_1]\mathbb{E}[z_2z_2] - \mathbb{E}[z_1z_2]\mathbb{E}[z_1z_2]| \\ p(x_1, x_2) &\triangleq \frac{1}{2\pi\sqrt{|M|}} \exp\left(\frac{z_1^2\mathbb{E}[z_2z_2] - 2z_1z_2\mathbb{E}[z_1z_2] + z_2^2\mathbb{E}[z_1z_1]}{-2|M|}\right) \end{aligned}$$

APPENDIX D

SPECTRAL THEORY

D.1 Operator Spectrum

Definition D.1. ¹ Let $\mathbf{A} \in \mathcal{B}(X, Y)$ be an operator over the linear spaces $X = (X, F, \oplus, \otimes)$ and $Y \triangleq (Y, F, \oplus, \otimes)$. Let $\mathcal{N}(\mathbf{A})$ be the NULL SPACE of \mathbf{A} .

D E F An **eigenvalue** of \mathbf{A} is any value λ such that there exists \mathbf{x} such that $\mathbf{Ax} = \lambda\mathbf{x}$.

The **eigenspace** H_λ of \mathbf{A} at eigenvalue λ is $\mathcal{N}(\mathbf{A} - \lambda\mathbf{I})$.

An **eigenvector** of \mathbf{A} associated with eigenvalue λ is any element of $\mathcal{N}(\mathbf{A} - \lambda\mathbf{I})$.

Example D.1. ² Let \mathbf{D} be the differential operator.

The set $\{e^{\lambda x} | \lambda \in \mathbb{C}\}$ are the eigenvectors of \mathbf{D} .

E X	$\rho(\mathbf{D}) = \emptyset$ (D has no non-spectral points whatsoever)
	$\sigma_p(\mathbf{D}) = \sigma(\mathbf{D})$ (the spectrum of D is all eigenvalues)
	$\sigma_c(\mathbf{D}) = \emptyset$ (D has no continuous spectrum)
	$\sigma_r(\mathbf{D}) = \emptyset$ (D has no resolvent spectrum)

PROOF:

$$\begin{aligned} (\mathbf{D} - \lambda\mathbf{I})e^{\lambda x} &= \mathbf{D}e^{\lambda x} - \lambda\mathbf{I}e^{\lambda x} \\ &= \lambda e^{\lambda x} - \lambda e^{\lambda x} \\ &= 0 \end{aligned} \quad \forall \lambda \in \mathbb{C}$$

This theorem and proof needs more work and investigation to prove/disprove its claims.

Definition D.2. ³ Let $\mathbf{A} \in \mathcal{B}(X, Y)$ be an operator over the linear spaces $X = (X, F, \oplus, \otimes)$ and $Y \triangleq (Y, F, \oplus, \otimes)$.

¹ Bollobás (1999), page 168, Descartes (1637), Descartes (1954), Cayley (1858), Hilbert (1904), page 67, Hilbert (1912),

² Pedersen (2000), page 79

³ Michel and Herget (1993), page 439

quantity	$\mathcal{N}(\mathbf{A} - \lambda\mathbf{I}) = \{\mathbf{0}\}$ ($\mathbf{x} = \mathbf{0}$ is the only solution)	$\overline{\mathcal{R}(\mathbf{A} - \lambda\mathbf{I})} = \mathbf{X}$ (dense)	$(\mathbf{A} - \lambda\mathbf{I})^{-1} \in \mathcal{B}(\mathbf{X}, \mathbf{Y})$ (continuous/bounded)
$\rho(\mathbf{A})$ (resolvent set)	1	1	1
$\sigma_p(\mathbf{A})$ (point spectrum)	0		
$\sigma_r(\mathbf{A})$ (residual spectrum)	1	0	
$\sigma_c(\mathbf{A})$ (continuous spectrum)	1	1	0

Table D.1: Spectrum of an operator \mathbf{A}

The **resolvent set** $\rho(\mathbf{A})$ of operator \mathbf{A} is defined as

$$\text{DEF } \rho(\mathbf{A}) \triangleq \left\{ \lambda \in F \mid \begin{array}{l} 1. \quad \mathcal{N}(\mathbf{A} - \lambda\mathbf{I}) = \{\mathbf{0}\} \\ 2. \quad \overline{\mathcal{R}(\mathbf{A} - \lambda\mathbf{I})} = \mathbf{X} \\ 3. \quad (\mathbf{A} - \lambda\mathbf{I})^{-1} \in \mathcal{B}(\mathbf{X}, \mathbf{Y}) \end{array} \right. \begin{array}{l} \text{(no non-zero eigenvectors)} \\ \text{(the range is dense in } \mathbf{X} \text{).} \\ \text{(inverse is continuous/bounded).} \end{array} \right\}$$

The **spectrum** $\sigma(\mathbf{A})$ of operator \mathbf{A} is defined as

$$\sigma(\mathbf{A}) \triangleq F \setminus \rho(\mathbf{A}).$$

Definition D.3. ⁴ Let $\mathbf{A} \in \mathcal{B}(\mathbf{X}, \mathbf{Y})$ be an operator over the linear spaces $\mathbf{X} = (X, F, \oplus, \otimes)$ and $\mathbf{Y} \triangleq (Y, F, \oplus, \otimes)$.

The **point spectrum** $\sigma_p(\mathbf{A})$ of operator \mathbf{A} is defined as


$$\sigma_p(\mathbf{A}) \triangleq \left\{ \lambda \in F \mid \begin{array}{l} 1. \quad \mathcal{N}(\mathbf{A} - \lambda\mathbf{I}) \supsetneq \{\mathbf{0}\} \\ \text{(has non-zero eigenvector)} \end{array} \right\}$$

The **residual spectrum** $\sigma_r(\mathbf{A})$ of operator \mathbf{A} is defined as

$$\sigma_r(\mathbf{A}) \triangleq \left\{ \lambda \in F \mid \begin{array}{l} 1. \quad \mathcal{N}(\mathbf{A} - \lambda\mathbf{I}) = \{\mathbf{0}\} \\ 2. \quad \overline{\mathcal{R}(\mathbf{A} - \lambda\mathbf{I})} \neq \mathbf{X} \\ \text{(not dense in } \mathbf{X} \text{—has gaps).} \end{array} \right. \text{and} \right\}$$

The **continuous spectrum** $\sigma_c(\mathbf{A})$ of operator \mathbf{A} is defined as

$$\sigma_c(\mathbf{A}) \triangleq \left\{ \lambda \in F \mid \begin{array}{l} 1. \quad \mathcal{N}(\mathbf{A} - \lambda\mathbf{I}) = \{\mathbf{0}\} \\ 2. \quad \overline{\mathcal{R}(\mathbf{A} - \lambda\mathbf{I})} = \mathbf{X} \\ 3. \quad (\mathbf{A} - \lambda\mathbf{I})^{-1} \notin \mathcal{B}(\mathbf{X}, \mathbf{Y}) \end{array} \right. \begin{array}{l} \text{(no non-zero eigenvectors)} \\ \text{(dense in } \mathbf{X}.) \\ \text{(not continuous / not bounded)} \end{array} \right. \text{and} \right\}$$



The spectral components' definitions are illustrated in the figure to the left and summarized in Table D.1 (page 170). Let a family of operators $\mathbf{B}(\lambda)$ be defined with respect to an operator \mathbf{A} such that $\mathbf{B}(\lambda) \triangleq (\mathbf{A} - \lambda\mathbf{I})$. Normally, we might expect a “normal” or “regular” or even “mundane” operator $\mathbf{B}(\lambda)$ to have the properties

1. $\mathbf{B}(\lambda)\mathbf{x} = \mathbf{0}$ if and only if $\mathbf{x} = \mathbf{0}$
2. $\mathbf{B}(\lambda)\mathbf{x}$ spans virtually all of \mathbf{X} as we vary \mathbf{x}
3. $\mathbf{B}^{-1}(\lambda)$ is continuous.

After all, these are the properties that we would have if $\mathbf{B}(\lambda)$ were simply an affine operator in the

⁴ [Bollobás \(1999\)](#), page 168, [Hilbert \(1906\)](#) pages 169–172

field of real numbers— such as $[\mathbf{B}(\lambda)](x) \triangleq [\lambda](x) = \lambda x$ which is 0 if and only if $x = 0$, has range $\mathcal{R}(\lambda) = \mathbb{R}$, and its inverse $\lambda^{-1}x$ is continuous.

If for some λ the operator $\mathbf{B}(\lambda)$ does have all these “regular” properties, then that λ part of the *resolvent set* of \mathbf{A} and λ is called *regular*. However if for some λ the operator $\mathbf{B}(\lambda)$ fails any of these conditions, then that λ part of the *spectrum* of \mathbf{A} . And which conditions it fails determines which component of the spectrum it is in.

Theorem D.1. ⁵ Let $\mathbf{A} \in \mathcal{B}(X, Y)$ be an operator.

T
H
M

$$\sigma(\mathbf{A}) = \sigma_p(\mathbf{A}) \cup \sigma_c(\mathbf{A}) \cup \sigma_r(\mathbf{A})$$

Theorem D.2 (Spectral Theorem). ^{6 7} Let $\mathbf{N} \in Y^X$ be an operator.

T
H
M

$$\left. \begin{array}{l} (1). \underbrace{\mathbf{N}^* \mathbf{N} = \mathbf{N} \mathbf{N}^*}_{\mathbf{N} \text{ is NORMAL}} \\ (2). \mathbf{N} \text{ is COMPACT} \end{array} \right\} \Rightarrow$$

$$\left\{ \begin{array}{l} (1). \mathbf{N} = \sum_n \lambda_n \mathbf{P}_n \\ (2). \sum_n \mathbf{P}_n = \mathbf{I} \\ (3). \mathbf{P}_n \mathbf{P}_m = \delta_{n-m} \mathbf{P}_n \\ (4). \dim(\mathcal{H}_n) < \infty \\ (5). |\{\lambda_n | \lambda_n \neq 0\}| \text{ is COUNTABLY INFINITE} \end{array} \right.$$

where

$$\begin{aligned} (\lambda_n)_{n \in \mathbb{Z}} &\triangleq \sigma_p(\mathbf{N}) && \text{(eigenvalues of } \mathbf{N}) \\ \mathcal{H}_n &\triangleq \mathcal{N}(\mathbf{N} - \lambda_n \mathbf{I}) && \text{(\lambda}_n \text{ is the eigenspace of } \mathbf{N} \text{ at } \lambda_n \text{ in } Y) \\ \mathbf{H}_n &= \mathbf{P}_n Y && \text{(\mathbf{P}_n \text{ is the projection operator that generates } \mathcal{H}_n)} \end{aligned}$$

D.2 Fredholm kernels

Definition D.4. ⁸

D
E
F

A **Fredholm operator** \mathbf{K} is defined as

$$[\mathbf{K}\mathbf{f}](t) \triangleq \underbrace{\int_a^b \kappa(t, s)\mathbf{f}(s) ds}_{\text{kernel}} \quad \forall \mathbf{f} \in L_2([a, b])$$

*Fredholm integral equation of the first kind*⁹

Example D.2. Examples of Fredholm operators include

1. Fourier Transform $[\tilde{\mathbf{F}}\mathbf{x}](f) = \int_t x(t)e^{-i2\pi ft} dt$ $\kappa(t, f) = e^{-i2\pi ft}$
2. Inverse Fourier Transform $[\tilde{\mathbf{F}}^{-1}\tilde{\mathbf{x}}](t) = \int_f \tilde{x}(f)e^{i2\pi ft} df$ $\kappa(f, t) = e^{i2\pi ft}$
3. Laplace operator $[\mathbf{L}\mathbf{x}](s) = \int_t x(t)e^{-st} dt$ $\kappa(t, s) = e^{-st}$
4. autocorrelation operator $[\mathbf{R}\mathbf{x}](t) = \int_s R(t, s)x(s) ds$ $\kappa(t, s) = R(t, s)$

Theorem D.3. Let \mathbf{K} be a Fredholm operator with kernel $\kappa(t, s)$ and adjoint \mathbf{K}^* .

T
H
M

$$[\mathbf{K}\mathbf{f}](t) = \int_A \kappa(t, s)\mathbf{f}(s) ds \iff [\mathbf{K}^*\mathbf{f}](t) = \int_A \kappa^*(s, t)\mathbf{f}(s) ds$$

⁵ Michel and Herget (1993), page 440

⁶ Michel and Herget (1993), page 457, ⁷ Bollobás (1999), page 200

⁷ Hilbert (1906), ⁸ Hilbert (1912), ⁹ von Neumann (1929), ¹⁰ de Witt (1659)

⁸ Michel and Herget (1993), page 425

⁹ The equation $\int_u \kappa(t, s)\mathbf{f}(s) ds$ is a **Fredholm integral equation of the first kind** and $\kappa(t, u)$ is the **kernel** of the equation. References: ¹¹ Fredholm (1900), ¹² Fredholm (1903), page 365, ¹³ Michel and Herget (1993), page 97, ¹⁴ Keener (1988), page 101

PROOF:

$$\begin{aligned}
 [\mathbf{K}f](t) &= \int_A \kappa(t, s)f(s) ds \\
 \Leftrightarrow \langle [\mathbf{K}f](t) | g(t) \rangle &= \left\langle \int_s \kappa(t, s)f(s) ds | g(t) \right\rangle \text{ by left hypothesis} \\
 &= \int_s f(s) \langle \kappa(t, s) | g(t) \rangle ds \text{ by additivity property of } \langle \triangle | \nabla \rangle \text{ (Definition I.1 page 233)} \\
 &= \int_s f(s) \langle g(t) | \kappa(t, s) \rangle^* ds \text{ by conjugate symmetry property of } \langle \triangle | \nabla \rangle \text{ (Definition I.1 page 233)} \\
 &= \langle f(s) | \langle g(t) | \kappa(t, s) \rangle \rangle \text{ by local definition of } \langle \triangle | \nabla \rangle \\
 &= \left\langle f(s) | \underbrace{\int_t \kappa^*(t, s)g(t) dt}_{[\mathbf{K}^*g](s)} \right\rangle \text{ by local definition of } \langle \triangle | \nabla \rangle \\
 \Leftrightarrow [\mathbf{K}^*g](s) &= \int_A \kappa^*(t, s)g(t) dt \text{ by right hypothesis} \\
 \Leftrightarrow [\mathbf{K}^*g](\sigma) &= \int_A \kappa^*(\tau, \sigma)g(\tau) d\tau \text{ by change of variable: } \tau = t, \sigma = s \\
 \Leftrightarrow [\mathbf{K}^*f](t) &= \int_A \kappa^*(s, t)f(s) ds \text{ by change of variable: } t = \sigma, s = \tau, f = g
 \end{aligned}$$

⇒

Theorem D.4. ¹⁰ Let \mathbf{K} be an Fredholm operator with kernel $\kappa(t, s)$ and adjoint \mathbf{K}^* .

THM

$$\mathbf{K} = \mathbf{K}^* \iff \underbrace{\kappa(t, s)}_{\text{kernel is conjugate symmetric}} = \underbrace{\kappa^*(s, t)}_{\text{kernel is conjugate symmetric}}$$

PROOF:

$$\begin{aligned}
 \mathbf{K} = \mathbf{K}^* &\iff \int_A \kappa(t, s)f(s) ds = \int_A \kappa^*(s, t)f(s) ds \text{ by Theorem D.3 page 171} \\
 &\iff \kappa(t, s) = \kappa^*(s, t)
 \end{aligned}$$

⇒

Theorem D.5 (Mercer's Theorem). ¹¹ Let \mathbf{K} be an Fredholm operator with kernel $\kappa(t, s)$ and eigen-system $((\lambda_n, \phi_n(t)))_{n \in \mathbb{Z}}$.

THM

$$\left. \begin{array}{l} 1. \underbrace{\int_a^b \int_a^b \kappa(t, s)f(t)f^*(s) dt ds \geq 0}_{\text{positive}} \quad \text{and} \\ 2. \kappa(t, s) \text{ is continuous on } [a, b] \times [a, b] \end{array} \right\} \Rightarrow \left\{ \begin{array}{l} 1. \kappa(t, s) = \sum_n \lambda_n \phi_n(t) \phi_n^*(s) \quad \text{and} \\ 2. \kappa(t, s) \text{ converges absolutely and uniformly on } [a, b] \times [a, b] \end{array} \right.$$

PROOF:

⇒

¹⁰ Michel and Herget (1993), page 430

¹¹ Gohberg et al. (2003), page 198, Courant and Hilbert (1930), pages 138–140, Mercer (1909), page 439

APPENDIX E

MATRIX CALCULUS

Optimization problems often require finding the value of some parameter which results in some measure reaching a minimum or maximum value. Often this optimal parameter value can be found by solving the single equation generated by the partial derivative of the measure with respect to the parameter. When there are several parameters, optimization often requires several simultaneous equations generated by the partial derivatives of the measure with respect to each parameter. The need for several partial derivatives and several simultaneous equations leads to a natural union of two branches of mathematics—partial differential equations and linear algebra. In general, we would like to not only be able to take the partial derivative of a scalar with respect to another scalar, but to be able to take the partial derivative of a vector with respect to another vector. This generalization is the problem addressed in this section. Other references are also available.¹

E.1 First derivative of a vector with respect to a vector

Definition E.1.

\mathbf{x} is a vector with the following properties:

D E F

$$1. \quad \mathbf{x} \triangleq \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} \quad (\text{n element column vector})$$

$$2. \quad \frac{\partial}{\partial x_k} x_j = \delta_{kj} \quad ((x_1, x_2, \dots, x_n) \text{ are mutually independent})$$

Definition E.2 (Jacobian matrix).² The gradient of \mathbf{y} with respect to \mathbf{x} , as well as the gradient of \mathbf{y}^T with respect to \mathbf{x} , is defined as

¹ [Graham \(1981\)](#) (Chapter 4), [Haykin \(2001\)](#) (Appendix B), [Moon and Stirling \(2000\)](#) (Appendix E), [Scharf \(1991\)](#), pages 274–276, [Trees \(2002\)](#) (Section A.7), [Felippa \(1999\)](#)

² [Graham \(1981\)](#), page 52, [Scharf \(1991\)](#), page 274, [Trees \(2002\)](#), page 1398

D E F $\frac{\partial \mathbf{y}}{\partial \mathbf{x}} \triangleq \frac{\partial \mathbf{y}^T}{\partial \mathbf{x}} \triangleq \underbrace{\begin{bmatrix} \frac{\partial y_1}{\partial x_1} & \frac{\partial y_2}{\partial x_1} & \dots & \frac{\partial y_m}{\partial x_1} \\ \frac{\partial y_1}{\partial x_2} & \frac{\partial y_2}{\partial x_2} & \dots & \frac{\partial y_m}{\partial x_2} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial y_1}{\partial x_n} & \frac{\partial y_2}{\partial x_n} & \dots & \frac{\partial y_m}{\partial x_n} \end{bmatrix}}_{n \times m \text{ matrix}} \quad \forall \mathbf{y} \in \mathbb{C}^m$

Remark E.1. Depending on whether \mathbf{x} and \mathbf{y} are scalars or vectors, $\frac{\partial \mathbf{y}}{\partial \mathbf{x}}$ takes on the following forms:³

	y scalar	y vector
x scalar	$\frac{\partial y}{\partial x}$	$\frac{\partial \mathbf{y}}{\partial x} = \begin{bmatrix} \frac{\partial y_1}{\partial x} & \frac{\partial y_2}{\partial x} & \dots & \frac{\partial y_m}{\partial x} \end{bmatrix}$
x vector	$\frac{\partial y}{\partial \mathbf{x}} = \begin{bmatrix} \frac{\partial y}{\partial x_1} \\ \frac{\partial y}{\partial x_2} \\ \vdots \\ \frac{\partial y}{\partial x_n} \end{bmatrix}$	$\frac{\partial \mathbf{y}}{\partial \mathbf{x}} \triangleq \begin{bmatrix} \frac{\partial y_1}{\partial x_1} & \frac{\partial y_2}{\partial x_1} & \dots & \frac{\partial y_m}{\partial x_1} \\ \frac{\partial y_1}{\partial x_2} & \frac{\partial y_2}{\partial x_2} & \dots & \frac{\partial y_m}{\partial x_2} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial y_1}{\partial x_n} & \frac{\partial y_2}{\partial x_n} & \dots & \frac{\partial y_m}{\partial x_n} \end{bmatrix}$

Lemma E.1. Let $\mathbf{x} \in \mathbb{R}^n$ be a vector. Then

L E M $\frac{\partial}{\partial x_k} x_i x_j = \bar{\delta}_{ik} x_j + \bar{\delta}_{jk} x_i = \begin{cases} 2x_k & \text{for } i = j = k \\ x_j & \text{for } i = k \text{ and } j \neq k \\ x_i & \text{for } i \neq k \text{ and } j = k \\ 0 & \text{otherwise} \end{cases}$

Lemma E.2.

L E M $(\mathbf{x}^H \mathbf{A} \mathbf{x}) = \sum_{i=1}^n \sum_{j=1}^n a_{ij} x_i^* x_j$ $\underbrace{\mathbf{A} \in (\mathbb{C}^n \times \mathbb{C}^n)}_{n \times n \text{ array}} \text{ and } \underbrace{\mathbf{x} \in \mathbb{C}^n}_{n \text{ element column vector}}$

PROOF:

$$\begin{aligned}
 \mathbf{x}^H \mathbf{A} \mathbf{x} &\triangleq \begin{bmatrix} x_1 & x_2 & \dots & x_n \end{bmatrix}^* \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & \dots & a_{nn} \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} \quad \text{by definitions of } \mathbf{A} \text{ and } \mathbf{x} \\
 &= \begin{bmatrix} x_1 & x_2 & \dots & x_n \end{bmatrix}^* \sum_{i=1}^n x_i \begin{bmatrix} a_{1i} \\ a_{2i} \\ \vdots \\ a_{ni} \end{bmatrix} \\
 &= \sum_{i=1}^n x_i \begin{bmatrix} x_1 & x_2 & \dots & x_n \end{bmatrix}^* \begin{bmatrix} a_{1i} \\ a_{2i} \\ \vdots \\ a_{ni} \end{bmatrix} \\
 &= \sum_{i=1}^n x_i \sum_{j=1}^n a_{ji} x_j^*
 \end{aligned}$$

³For the generalization of the partial derivative of a matrix with respect to a matrix, see [Graham \(1981\)](#) (chapter 6). Graham uses *kronecker products* to handle the additional dimensions(?)

$$= \sum_{i=1}^n \sum_{j=1}^n a_{ij} x_i^* x_j$$

**Lemma E.3.****L
E
M**

$$\frac{\partial}{\partial \mathbf{x}} [a(\mathbf{x}) b(\mathbf{x})] = a(\mathbf{x}) \left[\frac{\partial}{\partial \mathbf{x}} b(\mathbf{x}) \right] + \left[\frac{\partial}{\partial \mathbf{x}} a(\mathbf{x}) \right] b(\mathbf{x})$$

$\underbrace{\forall a, b : \mathbb{R}^n \rightarrow \mathbb{R}}$

$a(\mathbf{x}), b(\mathbf{x})$ are functions from a vector \mathbf{x} to a scalar in \mathbb{R}

PROOF:

$$\begin{aligned} \frac{\partial}{\partial \mathbf{x}} [a(\mathbf{x}) b(\mathbf{x})] &= \begin{bmatrix} \frac{\partial}{\partial x_1} [a(\mathbf{x}) b(\mathbf{x})] \\ \frac{\partial}{\partial x_2} [a(\mathbf{x}) b(\mathbf{x})] \\ \vdots \\ \frac{\partial}{\partial x_n} [a(\mathbf{x}) b(\mathbf{x})] \end{bmatrix} \\ &= \begin{bmatrix} a(\mathbf{x}) \frac{\partial b(\mathbf{x})}{\partial x_1} + b(\mathbf{x}) \frac{\partial a(\mathbf{x})}{\partial x_1} \\ a(\mathbf{x}) \frac{\partial b(\mathbf{x})}{\partial x_2} + b(\mathbf{x}) \frac{\partial a(\mathbf{x})}{\partial x_2} \\ \vdots \\ a(\mathbf{x}) \frac{\partial b(\mathbf{x})}{\partial x_n} + b(\mathbf{x}) \frac{\partial a(\mathbf{x})}{\partial x_n} \end{bmatrix} \\ &= \begin{bmatrix} a(\mathbf{x}) \frac{\partial b(\mathbf{x})}{\partial x_1} \\ a(\mathbf{x}) \frac{\partial b(\mathbf{x})}{\partial x_2} \\ \vdots \\ a(\mathbf{x}) \frac{\partial b(\mathbf{x})}{\partial x_n} \end{bmatrix} + \begin{bmatrix} \frac{\partial a(\mathbf{x})}{\partial x_1} b(\mathbf{x}) \\ \frac{\partial a(\mathbf{x})}{\partial x_2} b(\mathbf{x}) \\ \vdots \\ \frac{\partial a(\mathbf{x})}{\partial x_n} b(\mathbf{x}) \end{bmatrix} \\ &= a(\mathbf{x}) \left[\frac{\partial b(\mathbf{x})}{\partial \mathbf{x}} \right] + \left[\frac{\partial a(\mathbf{x})}{\partial \mathbf{x}} \right] b(\mathbf{x}) \end{aligned}$$

**Theorem E.1.** ⁴**L
E
M**

$$\frac{\partial}{\partial \mathbf{x}} \mathbf{x} = \mathbf{I} \quad \forall \mathbf{x} \in \mathbb{R}^n$$

PROOF:

$$\begin{aligned} \frac{\partial}{\partial \mathbf{x}} \mathbf{x} &= \begin{bmatrix} \frac{\partial x_1}{\partial x_1} & \frac{\partial x_2}{\partial x_1} & \dots & \frac{\partial x_n}{\partial x_1} \\ \frac{\partial x_1}{\partial x_2} & \frac{\partial x_1}{\partial x_2} & \dots & \frac{\partial x_n}{\partial x_2} \\ \frac{\partial x_1}{\partial x_3} & \frac{\partial x_2}{\partial x_3} & \dots & \frac{\partial x_n}{\partial x_3} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial x_1}{\partial x_n} & \frac{\partial x_2}{\partial x_n} & \dots & \frac{\partial x_n}{\partial x_n} \end{bmatrix} \\ &= \begin{bmatrix} \bar{\delta}_{11} & \bar{\delta}_{21} & \dots & \bar{\delta}_{n1} \\ \bar{\delta}_{12} & \bar{\delta}_{22} & \dots & \bar{\delta}_{n2} \\ \vdots & \vdots & \ddots & \vdots \\ \bar{\delta}_{1n} & \bar{\delta}_{2n} & \dots & \bar{\delta}_{nn} \end{bmatrix} \end{aligned}$$

by Definition E.2 page 173

by Definition E.1 page 173 (mutual independence property)

⁴ Scharf (1991), page 274, Trees (2002), page 1398

$$\begin{aligned}
 &= \begin{bmatrix} 1 & 0 & \cdots & 0 \\ 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 1 \end{bmatrix} && \text{by definition of kronecker delta function } \delta \\
 &= \mathbf{I} && \text{by definition of identity operator } \mathbf{I}
 \end{aligned}$$

⇒

Theorem E.2.

T H M $\frac{\partial}{\partial \mathbf{x}}(\mathbf{A}\mathbf{x}) = \mathbf{A}^T + \sum_{i=1}^n \left(\frac{\partial}{\partial \mathbf{x}} [\begin{array}{cccc} a_{1i} & a_{2i} & \cdots & a_{mi} \end{array}] \right) \mathbf{x}_i \quad \forall \mathbf{x} \in \mathbb{R}^n, \mathbf{A} \in \mathbb{C}^m \times \mathbb{C}^n$

PROOF: Let

$$\mathbf{x} \triangleq \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} \quad \mathbf{A} \triangleq \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}$$

$$\frac{\partial}{\partial \mathbf{x}} (\mathbf{A}\mathbf{x}) = \frac{\partial}{\partial \mathbf{x}} \left(\begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} \right)$$

$$= \frac{\partial}{\partial \mathbf{x}} \begin{bmatrix} \sum_{i=1}^n a_{1i}x_i \\ \sum_{i=1}^n a_{2i}x_i \\ \vdots \\ \sum_{i=1}^n a_{mi}x_i \end{bmatrix} \quad \text{by matrix multiplication}$$

$$= \frac{\partial}{\partial \mathbf{x}} \sum_{i=1}^n \begin{bmatrix} a_{1i}x_i \\ a_{2i}x_i \\ \vdots \\ a_{mi}x_i \end{bmatrix}$$

$$= \sum_{i=1}^n \frac{\partial}{\partial \mathbf{x}} \begin{bmatrix} a_{1i}x_i \\ a_{2i}x_i \\ \vdots \\ a_{mi}x_i \end{bmatrix}$$

$$= \sum_{i=1}^n \begin{bmatrix} \frac{\partial a_{1i}x_i}{\partial x_1} & \frac{\partial a_{2i}x_i}{\partial x_1} & \cdots & \frac{\partial a_{mi}x_i}{\partial x_1} \\ \frac{\partial a_{1i}x_i}{\partial x_2} & \frac{\partial a_{2i}x_i}{\partial x_2} & \cdots & \frac{\partial a_{mi}x_i}{\partial x_2} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial a_{1i}x_i}{\partial x_n} & \frac{\partial a_{2i}x_i}{\partial x_n} & \cdots & \frac{\partial a_{mi}x_i}{\partial x_n} \end{bmatrix}$$

$$= \sum_{i=1}^n \begin{bmatrix} a_{1i} \frac{\partial x_i}{\partial x_1} + \frac{\partial a_{1i}}{\partial x_1} x_i & a_{2i} \frac{\partial x_i}{\partial x_1} + \frac{\partial a_{2i}}{\partial x_1} x_i & \cdots & a_{mi} \frac{\partial x_i}{\partial x_1} + \frac{\partial a_{mi}}{\partial x_1} x_i \\ a_{1i} \frac{\partial x_i}{\partial x_2} + \frac{\partial a_{1i}}{\partial x_2} x_i & a_{2i} \frac{\partial x_i}{\partial x_2} + \frac{\partial a_{2i}}{\partial x_2} x_i & \cdots & a_{mi} \frac{\partial x_i}{\partial x_2} + \frac{\partial a_{mi}}{\partial x_2} x_i \\ \vdots & \vdots & \ddots & \vdots \\ a_{1i} \frac{\partial x_i}{\partial x_n} + \frac{\partial a_{1i}}{\partial x_n} x_i & a_{2i} \frac{\partial x_i}{\partial x_n} + \frac{\partial a_{2i}}{\partial x_n} x_i & \cdots & a_{mi} \frac{\partial x_i}{\partial x_n} + \frac{\partial a_{mi}}{\partial x_n} x_i \end{bmatrix}$$

by Definition E.2 page 173

by Lemma E.3 page 175



$$= \sum_{i=1}^n \begin{bmatrix} a_{1i} \frac{\partial x_i}{\partial x_1} & a_{2i} \frac{\partial x_i}{\partial x_1} & \cdots & a_{mi} \frac{\partial x_i}{\partial x_1} \\ a_{1i} \frac{\partial x_i}{\partial x_2} & a_{2i} \frac{\partial x_i}{\partial x_2} & \cdots & a_{mi} \frac{\partial x_i}{\partial x_2} \\ \vdots & \vdots & \ddots & \vdots \\ a_{1i} \frac{\partial x_i}{\partial x_n} & a_{2i} \frac{\partial x_i}{\partial x_n} & \cdots & a_{mi} \frac{\partial x_i}{\partial x_n} \end{bmatrix} + \sum_{i=1}^n \begin{bmatrix} \frac{\partial a_{1i}}{\partial x_1} x_i & \frac{\partial a_{2i}}{\partial x_1} x_i & \cdots & \frac{\partial a_{mi}}{\partial x_1} x_i \\ \frac{\partial a_{1i}}{\partial x_2} x_i & \frac{\partial a_{2i}}{\partial x_2} x_i & \cdots & \frac{\partial a_{mi}}{\partial x_2} x_i \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial a_{1i}}{\partial x_n} x_i & \frac{\partial a_{2i}}{\partial x_n} x_i & \cdots & \frac{\partial a_{mi}}{\partial x_n} x_i \end{bmatrix}$$

$$= \sum_{i=1}^n \begin{bmatrix} a_{1i} \bar{\delta}_{i1} & a_{2i} \bar{\delta}_{i1} & \cdots & a_{mi} \bar{\delta}_{i1} \\ a_{1i} \bar{\delta}_{i2} & a_{2i} \bar{\delta}_{i2} & \cdots & a_{mi} \bar{\delta}_{i2} \\ \vdots & \vdots & \ddots & \vdots \\ a_{1i} \bar{\delta}_{in} & a_{2i} \bar{\delta}_{in} & \cdots & a_{mi} \bar{\delta}_{in} \end{bmatrix} + \sum_{i=1}^n \left(\frac{\partial}{\partial \mathbf{x}} [a_{1i} \ a_{2i} \ \cdots \ a_{mi}] \right) x_i \quad \text{by Lemma E.1}$$

$$= \begin{bmatrix} a_{11} & a_{21} & \cdots & a_{m1} \\ a_{12} & a_{22} & \cdots & a_{m2} \\ \vdots & \vdots & \ddots & \vdots \\ a_{1n} & a_{2n} & \cdots & a_{mn} \end{bmatrix} + \sum_{i=1}^n \left(\frac{\partial}{\partial \mathbf{x}} [a_{1i} \ a_{2i} \ \cdots \ a_{mi}] \right) x_i \quad \text{by definition of } \bar{\delta}$$

$$= \mathbf{A}^T + \sum_{i=1}^n \left(\frac{\partial}{\partial \mathbf{x}} [a_{1i} \ a_{2i} \ \cdots \ a_{mi}] \right) x_i$$

⇒

Theorem E.3 (Affine equations). ⁵

T H M	A and B are independent of x \implies $\begin{cases} \frac{\partial}{\partial \mathbf{x}}(\mathbf{Ax}) = \mathbf{A}^T & \forall \mathbf{x} \in \mathbb{R}^n, \mathbf{A} \in \mathbb{C}^m \times \mathbb{C}^n \\ \frac{\partial}{\partial \mathbf{x}}(\mathbf{x}^T \mathbf{B}) = \mathbf{B} & \forall \mathbf{x} \in \mathbb{R}^n, \mathbf{B} \in \mathbb{C}^n \times \mathbb{C}^m \end{cases}$
-------------	---

PROOF: Let $\mathbf{B} \triangleq \mathbf{A}^T$.1. Proof that $\frac{\partial}{\partial \mathbf{x}}(\mathbf{Ax}) = \mathbf{A}^T$:

$$\begin{aligned} \frac{\partial}{\partial \mathbf{x}}(\mathbf{Ax}) &= \mathbf{A}^T + \sum_{i=1}^n \left(\frac{\partial}{\partial \mathbf{x}} [a_{1i} \ a_{2i} \ \cdots \ a_{mi}] \right) x_i && \text{by Theorem E.2 page 176} \\ &= \mathbf{A}^T + \sum_{i=1}^n \left[\frac{\partial}{\partial \mathbf{x}} a_{1i} \ \frac{\partial}{\partial \mathbf{x}} a_{2i} \ \cdots \ \frac{\partial}{\partial \mathbf{x}} a_{mi} \right] x_i \\ &= \mathbf{A}^T + \sum_{i=1}^n \left[\begin{array}{cccc} 0 & 0 & \cdots & 0 \end{array} \right] x_i && \text{by left hypothesis} \\ &= \mathbf{A}^T \end{aligned}$$

2. Proof that $\frac{\partial}{\partial \mathbf{x}}(\mathbf{x}^T \mathbf{B}) = \mathbf{B}$:

$$\begin{aligned} \frac{\partial}{\partial \mathbf{x}}(\mathbf{x}^T \mathbf{B}) &= \frac{\partial}{\partial \mathbf{x}}(\mathbf{x}^T \mathbf{A}^T) && \text{by definition of } \mathbf{B} \\ &= \frac{\partial}{\partial \mathbf{x}}[(\mathbf{Ax})^T] && \\ &= \frac{\partial}{\partial \mathbf{x}}(\mathbf{Ax}) && \text{by Definition E.2 page 173} \\ &= \mathbf{A}^T && \text{by Theorem E.3 page 177} \\ &= \mathbf{B} && \text{by definition of } \mathbf{B} \end{aligned}$$

⇒

⁵  Graham (1981), page 54

Theorem E.4 (Product rule). ⁶ Let \mathbf{y} and \mathbf{z} be functions of \mathbf{x} and

T	H	M	$\frac{\partial}{\partial \mathbf{x}} \mathbf{z}^T \mathbf{y} = \frac{\partial \mathbf{z}}{\partial \mathbf{x}} \mathbf{y} + \frac{\partial \mathbf{y}}{\partial \mathbf{x}} \mathbf{z}$	forall $\mathbf{x} \in \mathbb{R}^n$, $\mathbf{y} \in \mathbb{R}^m$, $\mathbf{z} \in \mathbb{R}^m$
---	---	---	--	--

PROOF:

$$\begin{aligned}
 \frac{\partial}{\partial \mathbf{x}} \mathbf{z}^T \mathbf{y} &= \frac{\partial}{\partial \mathbf{x}} \sum_{k=1}^m z_k y_k \\
 &= \sum_{k=1}^m \frac{\partial}{\partial \mathbf{x}} z_k y_k \\
 &= \sum_{k=1}^m \frac{\partial z_k}{\partial \mathbf{x}} y_k + \sum_{k=1}^m \frac{\partial y_k}{\partial \mathbf{x}} z_k \quad \text{by Lemma E.3 page 175} \\
 &= \left[\begin{array}{cccc} \frac{\partial z_1}{\partial x_1} y_1 & + & \frac{\partial z_2}{\partial x_1} y_2 & + \cdots + \frac{\partial z_n}{\partial x_1} y_n \\ \frac{\partial z_1}{\partial x_1} y_1 & + & \frac{\partial z_2}{\partial x_1} y_2 & + \cdots + \frac{\partial z_n}{\partial x_1} y_n \\ \vdots & & \ddots & \vdots \\ \frac{\partial z_1}{\partial x_1} y_1 & + & \frac{\partial z_2}{\partial x_1} y_2 & + \cdots + \frac{\partial z_n}{\partial x_1} y_n \end{array} \right] + \left[\begin{array}{cccc} \frac{\partial y_1}{\partial x_1} z_1 & + & \frac{\partial y_2}{\partial x_1} z_2 & + \cdots + \frac{\partial y_n}{\partial x_1} z_n \\ \frac{\partial y_1}{\partial x_1} z_1 & + & \frac{\partial y_2}{\partial x_1} z_2 & + \cdots + \frac{\partial y_n}{\partial x_1} z_n \\ \vdots & & \ddots & \vdots \\ \frac{\partial y_1}{\partial x_1} z_1 & + & \frac{\partial y_2}{\partial x_1} z_2 & + \cdots + \frac{\partial y_n}{\partial x_1} z_n \end{array} \right] \\
 &= \left[\begin{array}{ccc} \frac{\partial z_1}{\partial x_1} & \frac{\partial z_2}{\partial x_1} & \cdots & \frac{\partial z_n}{\partial x_1} \\ \frac{\partial z_1}{\partial x_1} & \frac{\partial z_2}{\partial x_1} & \cdots & \frac{\partial z_n}{\partial x_1} \\ \vdots & \ddots & \vdots & \vdots \\ \frac{\partial z_1}{\partial x_1} & \frac{\partial z_2}{\partial x_1} & \cdots & \frac{\partial z_n}{\partial x_1} \end{array} \right] \left[\begin{array}{c} y_1 \\ y_2 \\ \vdots \\ y_n \end{array} \right] + \left[\begin{array}{ccc} \frac{\partial y_1}{\partial x_1} & \frac{\partial y_2}{\partial x_1} & \cdots & \frac{\partial y_n}{\partial x_1} \\ \frac{\partial y_1}{\partial x_1} & \frac{\partial y_2}{\partial x_1} & \cdots & \frac{\partial y_n}{\partial x_1} \\ \vdots & \ddots & \vdots & \vdots \\ \frac{\partial y_1}{\partial x_1} & \frac{\partial y_2}{\partial x_1} & \cdots & \frac{\partial y_n}{\partial x_1} \end{array} \right] \left[\begin{array}{c} z_1 \\ z_2 \\ \vdots \\ z_n \end{array} \right] \\
 &= \frac{\partial \mathbf{z}}{\partial \mathbf{x}} \mathbf{y} + \frac{\partial \mathbf{y}}{\partial \mathbf{x}} \mathbf{z}
 \end{aligned}$$

Theorem E.5.

T	H	M	$\frac{\partial}{\partial \mathbf{x}} (\mathbf{x}^T \mathbf{A} \mathbf{x}) = \mathbf{A} \mathbf{x} + \mathbf{A}^T \mathbf{x} + \left[\sum_{i=1}^n \left(\frac{\partial}{\partial \mathbf{x}} [a_{1i} \ a_{2i} \ \cdots \ a_{ni}] \right) x_i \right] \mathbf{x}$	forall $\mathbf{x} \in \mathbb{R}^n$, $\mathbf{A} \in \mathbb{R}^{n \times n}$
---	---	---	--	---

PROOF:

$$\begin{aligned}
 \frac{\partial}{\partial \mathbf{x}} (\mathbf{x}^T \mathbf{A} \mathbf{x}) &= \left[\frac{\partial}{\partial \mathbf{x}} \mathbf{x} \right] \mathbf{A} \mathbf{x} + \left[\frac{\partial}{\partial \mathbf{x}} \mathbf{A} \mathbf{x} \right] \mathbf{x} \quad \text{by Theorem E.4 page 178} \\
 &= \mathbf{I} \mathbf{A} \mathbf{x} + \left[\mathbf{A}^T + \sum_{i=1}^n \left(\frac{\partial}{\partial \mathbf{x}} [a_{1i} \ a_{2i} \ \cdots \ a_{ni}] \right) x_i \right] \mathbf{x} \quad \text{by Theorem E.1 and Theorem E.2} \\
 &= \mathbf{A} \mathbf{x} + \mathbf{A}^T \mathbf{x} + \left[\sum_{i=1}^n \left(\frac{\partial}{\partial \mathbf{x}} [a_{1i} \ a_{2i} \ \cdots \ a_{ni}] \right) x_i \right] \mathbf{x} \quad \text{by definition of identity operator } \mathbf{I}
 \end{aligned}$$

Theorem E.6 (Quadratic form). ⁷

T	H	M	\mathbf{A} is independent of \mathbf{x} $\Rightarrow \frac{\partial}{\partial \mathbf{x}} (\mathbf{x}^T \mathbf{A} \mathbf{x}) = \mathbf{A} \mathbf{x} + \mathbf{A}^T \mathbf{x}$	forall $\mathbf{x} \in \mathbb{R}^n$, $\mathbf{A} \in \mathbb{R}^{n \times n}$
---	---	---	---	---

⁶ Scharf (1991), page 274, Trees (2002), page 1398

⁷ Graham (1981), page 54

PROOF:

$$\begin{aligned}\frac{\partial}{\partial \mathbf{x}}(\mathbf{x}^T \mathbf{A} \mathbf{x}) &= \left[\frac{\partial}{\partial \mathbf{x}} \mathbf{x} \right] \mathbf{A} \mathbf{x} + \left[\frac{\partial}{\partial \mathbf{x}} \mathbf{A} \mathbf{x} \right] \mathbf{x} \\ &= \mathbf{I} \mathbf{A} \mathbf{x} + \mathbf{A}^T \mathbf{x}\end{aligned}$$

by Theorem E.4 page 178

by Theorem E.1 page 175 and Theorem E.3 page 177

Corollary E.1.⁸

COR $\frac{\partial}{\partial \mathbf{x}}(\mathbf{x}^T \mathbf{x}) = 2\mathbf{x} \quad \forall \mathbf{x} \in \mathbb{R}^n$

PROOF:

$$\begin{aligned}\frac{\partial}{\partial \mathbf{x}}(\mathbf{x}^T \mathbf{x}) &= \frac{\partial}{\partial \mathbf{x}}(\mathbf{x}^T \mathbf{I} \mathbf{x}) \\ &= \mathbf{I} \mathbf{x} + \mathbf{I}^T \mathbf{x} \\ &= \mathbf{x} + \mathbf{x} \\ &= 2\mathbf{x}\end{aligned}$$

by property of identity operator I

by previous result 3.

by property of identity operator I **Theorem E.7** (Chain rule).⁹ Let \mathbf{z} be a function of \mathbf{y} and \mathbf{y} a function of \mathbf{x} and

$$\mathbf{x} \triangleq \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} \quad \mathbf{y} \triangleq \begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_m \end{bmatrix} \quad \mathbf{z} \triangleq \begin{bmatrix} z_1 \\ z_2 \\ \vdots \\ z_k \end{bmatrix}$$

THM $\frac{\partial}{\partial \mathbf{x}} \mathbf{z} = \frac{\partial \mathbf{y}}{\partial \mathbf{x}} \frac{\partial \mathbf{z}}{\partial \mathbf{y}}$

PROOF:

$$\begin{aligned}\frac{\partial \mathbf{z}}{\partial \mathbf{x}} &= \begin{bmatrix} \frac{\partial z_1}{\partial x_1} & \frac{\partial z_1}{\partial x_2} & \cdots & \frac{\partial z_1}{\partial x_n} \\ \frac{\partial z_2}{\partial x_1} & \frac{\partial z_2}{\partial x_2} & \cdots & \frac{\partial z_2}{\partial x_n} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial z_k}{\partial x_1} & \frac{\partial z_k}{\partial x_2} & \cdots & \frac{\partial z_k}{\partial x_n} \end{bmatrix} \\ &= \begin{bmatrix} \sum_{j=0}^m \frac{\partial z_1}{\partial y_j} \frac{\partial y_j}{\partial x_1} & \sum_{j=0}^m \frac{\partial z_2}{\partial y_j} \frac{\partial y_j}{\partial x_1} & \cdots & \sum_{j=0}^m \frac{\partial z_k}{\partial y_j} \frac{\partial y_j}{\partial x_1} \\ \sum_{j=0}^m \frac{\partial z_1}{\partial y_j} \frac{\partial y_j}{\partial x_2} & \sum_{j=0}^m \frac{\partial z_2}{\partial y_j} \frac{\partial y_j}{\partial x_2} & \cdots & \sum_{j=0}^m \frac{\partial z_k}{\partial y_j} \frac{\partial y_j}{\partial x_2} \\ \vdots & \vdots & \ddots & \vdots \\ \sum_{j=0}^m \frac{\partial z_1}{\partial y_j} \frac{\partial y_j}{\partial x_n} & \sum_{j=0}^m \frac{\partial z_2}{\partial y_j} \frac{\partial y_j}{\partial x_n} & \cdots & \sum_{j=0}^m \frac{\partial z_k}{\partial y_j} \frac{\partial y_j}{\partial x_n} \end{bmatrix} \\ &= \begin{bmatrix} \frac{\partial y_1}{\partial x_1} & \frac{\partial y_2}{\partial x_1} & \cdots & \frac{\partial y_m}{\partial x_1} \\ \frac{\partial y_1}{\partial x_2} & \frac{\partial y_2}{\partial x_2} & \cdots & \frac{\partial y_m}{\partial x_2} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial y_1}{\partial x_n} & \frac{\partial y_2}{\partial x_n} & \cdots & \frac{\partial y_m}{\partial x_n} \end{bmatrix} \begin{bmatrix} \frac{\partial z_1}{\partial y_1} & \frac{\partial z_2}{\partial y_1} & \cdots & \frac{\partial z_k}{\partial y_1} \\ \frac{\partial z_1}{\partial y_2} & \frac{\partial z_2}{\partial y_2} & \cdots & \frac{\partial z_k}{\partial y_2} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial z_1}{\partial y_m} & \frac{\partial z_2}{\partial y_m} & \cdots & \frac{\partial z_k}{\partial y_m} \end{bmatrix} \\ &= \frac{\partial \mathbf{y}}{\partial \mathbf{x}} \frac{\partial \mathbf{z}}{\partial \mathbf{y}}\end{aligned}$$

⁸ Graham (1981), page 54⁹ Graham (1981), pages 54–55

E.2 First derivative of a matrix with respect to a scalar

Definition E.3. Let $x \in \mathbb{R}$, $\{y_{jk} \in \mathbb{C} | j = 1, 2, \dots, m; k = 1, 2, \dots, n\}$ and

$$Y = \underbrace{\begin{bmatrix} y_{11} & y_{12} & \cdots & y_{1n} \\ y_{21} & y_{22} & \cdots & y_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ y_{m1} & y_{m2} & \cdots & y_{mn} \end{bmatrix}}_{m \times n \text{ matrix}}$$

The derivative of Y with respect to x is

D E F

$$\frac{dY}{dx} \triangleq \underbrace{\begin{bmatrix} \frac{dy_{11}}{dx} & \frac{dy_{12}}{dx} & \cdots & \frac{dy_{1n}}{dx} \\ \frac{dy_{21}}{dx} & \frac{dy_{22}}{dx} & \cdots & \frac{dy_{2n}}{dx} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{dy_{m1}}{dx} & \frac{dy_{m2}}{dx} & \cdots & \frac{dy_{mn}}{dx} \end{bmatrix}}_{m \times n \text{ matrix}}$$

Theorem E.8.¹⁰ Let $x \in \mathbb{R}$, $\{y_{jp} \in \mathbb{C} | j = 1, 2, \dots, m; p = 1, 2, \dots, n\}$, $\{w_{jp} \in \mathbb{C} | j = 1, 2, \dots, n; p = 1, 2, \dots, k\}$, and

$$Y = \underbrace{\begin{bmatrix} y_{11} & y_{12} & \cdots & y_{1n} \\ y_{21} & y_{22} & \cdots & y_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ y_{m1} & y_{m2} & \cdots & y_{mn} \end{bmatrix}}_{m \times n \text{ matrix}} \quad W = \underbrace{\begin{bmatrix} w_{11} & w_{12} & \cdots & w_{1k} \\ w_{21} & w_{22} & \cdots & w_{2k} \\ \vdots & \vdots & \ddots & \vdots \\ w_{p1} & w_{p2} & \cdots & w_{pk} \end{bmatrix}}_{p \times k \text{ matrix}}$$

T H M

$\frac{d}{dx}(Y + W) = \frac{d}{dx}Y + \frac{d}{dx}W$	(for $p = m, k = n$)
$\frac{d}{dx}(YW) = \left(\frac{d}{dx}Y\right)W + Y\left(\frac{d}{dx}W\right)$	(for $p = n$)
$\frac{d}{dx}(Y^T) = \left(\frac{d}{dx}Y\right)^T$	
$\frac{d}{dx}(Y^{-1}) = -Y^{-1}\left(\frac{d}{dx}Y\right)Y^{-1}$	(for $m = n$ and Y invertible)

PROOF:

$$\begin{aligned} \frac{d}{dx}(Y + W) &= \frac{d}{dx} \left(\begin{bmatrix} y_{11} & y_{12} & \cdots & y_{1n} \\ y_{21} & y_{22} & \cdots & y_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ y_{m1} & y_{m2} & \cdots & y_{mn} \end{bmatrix} + \begin{bmatrix} w_{11} & w_{12} & \cdots & w_{1n} \\ w_{21} & w_{22} & \cdots & w_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ w_{m1} & w_{m2} & \cdots & w_{mn} \end{bmatrix} \right) \\ &= \frac{d}{dx} \begin{bmatrix} y_{11} + w_{11} & y_{12} + w_{12} & \cdots & y_{1n} + w_{1n} \\ y_{21} + w_{21} & y_{22} + w_{22} & \cdots & y_{2n} + w_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ y_{m1} + w_{m1} & y_{m2} + w_{m2} & \cdots & y_{mn} + w_{mn} \end{bmatrix} \end{aligned}$$

¹⁰ Gradshteyn and Ryzhik (1980), pages 1106–1107

$$\begin{aligned}
&= \begin{bmatrix} (y_{11} + w_{11})' & (y_{12} + w_{12})' & \cdots & (y_{1n} + w_{1n})' \\ (y_{21} + w_{21})' & (y_{22} + w_{22})' & \cdots & (y_{2n} + w_{2n})' \\ \vdots & \vdots & \ddots & \vdots \\ (y_{m1} + w_{m1})' & (y_{m2} + w_{m2})' & \cdots & (y_{mn} + w_{mn})' \end{bmatrix} \\
&= \begin{bmatrix} y'_{11} + w'_{11} & y'_{12} + w'_{12} & \cdots & y'_{1n} + w'_{1n} \\ y'_{21} + w'_{21} & y'_{22} + w'_{22} & \cdots & y'_{2n} + w'_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ y'_{m1} + w'_{m1} & y'_{m2} + w'_{m2} & \cdots & y'_{mn} + w'_{mn} \end{bmatrix} \\
&= \begin{bmatrix} y'_{11} & y'_{12} & \cdots & y'_{1n} \\ y'_{21} & y'_{22} & \cdots & y'_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ y'_{m1} & y'_{m2} & \cdots & y'_{mn} \end{bmatrix} + \begin{bmatrix} w'_{11} & w'_{12} & \cdots & w'_{1n} \\ w'_{21} & w'_{22} & \cdots & w'_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ w'_{m1} & w'_{m2} & \cdots & w'_{mn} \end{bmatrix} \\
&= \frac{d}{dx} \begin{bmatrix} y_{11} & y_{12} & \cdots & y_{1n} \\ y_{21} & y_{22} & \cdots & y_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ y_{m1} & y_{m2} & \cdots & y_{mn} \end{bmatrix} + \frac{d}{dx} \begin{bmatrix} w_{11} & w_{12} & \cdots & w_{1n} \\ w_{21} & w_{22} & \cdots & w_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ w_{m1} & w_{m2} & \cdots & w_{mn} \end{bmatrix} \\
&= \frac{d}{dx} Y + \frac{d}{dx} W
\end{aligned}$$

$$\begin{aligned}
\frac{d}{dx}(YW) &= \frac{d}{dx} \left(\begin{bmatrix} y_{11} & y_{12} & \cdots & y_{1n} \\ y_{21} & y_{22} & \cdots & y_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ y_{m1} & y_{m2} & \cdots & y_{mn} \end{bmatrix} \begin{bmatrix} w_{11} & w_{12} & \cdots & w_{1k} \\ w_{21} & w_{22} & \cdots & w_{2k} \\ \vdots & \vdots & \ddots & \vdots \\ w_{n1} & w_{n2} & \cdots & w_{nk} \end{bmatrix} \right) \\
&= \frac{d}{dx} \begin{bmatrix} \sum_{j=1}^n y_{1j} w_{j1} & \sum_{j=1}^n y_{1j} w_{j2} & \cdots & \sum_{j=1}^n y_{1j} w_{jk} \\ \sum_{j=1}^n y_{2j} w_{j1} & \sum_{j=1}^n y_{2j} w_{j2} & \cdots & \sum_{j=1}^n y_{2j} w_{jk} \\ \vdots & \vdots & \ddots & \vdots \\ \sum_{j=1}^n y_{mj} w_{j1} & \sum_{j=1}^n y_{mj} w_{j2} & \cdots & \sum_{j=1}^n y_{mj} w_{jk} \end{bmatrix} \\
&= \frac{d}{dx} \sum_{j=1}^n \begin{bmatrix} y_{1j} w_{j1} & y_{1j} w_{j2} & \cdots & y_{1j} w_{jk} \\ y_{2j} w_{j1} & y_{2j} w_{j2} & \cdots & y_{2j} w_{jk} \\ \vdots & \vdots & \ddots & \vdots \\ y_{mj} w_{j1} & y_{mj} w_{j2} & \cdots & y_{mj} w_{jk} \end{bmatrix} \\
&= \sum_{j=1}^n \frac{d}{dx} \begin{bmatrix} y_{1j} w_{j1} & y_{1j} w_{j2} & \cdots & y_{1j} w_{jk} \\ y_{2j} w_{j1} & y_{2j} w_{j2} & \cdots & y_{2j} w_{jk} \\ \vdots & \vdots & \ddots & \vdots \\ y_{mj} w_{j1} & y_{mj} w_{j2} & \cdots & y_{mj} w_{jk} \end{bmatrix} \\
&= \sum_{j=1}^n \begin{bmatrix} \frac{d}{dx}(y_{1j} w_{j1}) & \frac{d}{dx}(y_{1j} w_{j2}) & \cdots & \frac{d}{dx}(y_{1j} w_{jk}) \\ \frac{d}{dx}(y_{2j} w_{j1}) & \frac{d}{dx}(y_{2j} w_{j2}) & \cdots & \frac{d}{dx}(y_{2j} w_{jk}) \\ \vdots & \vdots & \ddots & \vdots \\ \frac{d}{dx}(y_{mj} w_{j1}) & \frac{d}{dx}(y_{mj} w_{j2}) & \cdots & \frac{d}{dx}(y_{mj} w_{jk}) \end{bmatrix} \\
&= \sum_{j=1}^n \begin{bmatrix} \vdots & \vdots & \ddots & \vdots \\ y'_{1j} w_{j1} + y_{1j} w'_{j1} & y'_{1j} w_{j2} + y_{1j} w'_{j2} & \cdots & y'_{1j} w_{jk} + y_{1j} w'_{jk} \\ y'_{2j} w_{j1} + y_{2j} w'_{j1} & y'_{2j} w_{j2} + y_{2j} w'_{j2} & \cdots & y'_{2j} w_{jk} + y_{2j} w'_{jk} \\ y'_{mj} w_{j1} + y_{mj} w'_{j1} & y'_{mj} w_{j2} + y_{mj} w'_{j2} & \cdots & y'_{mj} w_{jk} + y_{mj} w'_{jk} \end{bmatrix}
\end{aligned}$$

$$\begin{aligned}
 &= \sum_{j=1}^n \left(\begin{bmatrix} y'_{1j} w_{j1} & y'_{1j} w_{j2} & \cdots & y'_{1j} w_{jk} \\ y'_{2j} w_{j1} & y'_{2j} w_{j2} & \cdots & y'_{2j} w_{jk} \\ \vdots & \vdots & \ddots & \vdots \\ y'_{mj} w_{j1} & y'_{mj} w_{j2} & \cdots & y'_{mj} w_{jk} \end{bmatrix} + \begin{bmatrix} y_{1j} w'_{j1} & y_{1j} w'_{j2} & \cdots & y_{1j} w'_{jk} \\ y_{2j} w'_{j1} & y_{2j} w'_{j2} & \cdots & y_{2j} w'_{jk} \\ \vdots & \vdots & \ddots & \vdots \\ y_{mj} w'_{j1} & y_{mj} w'_{j2} & \cdots & y_{mj} w'_{jk} \end{bmatrix} \right) \\
 &= \left(\frac{d}{dx} Y \right) W + Y \left(\frac{d}{dx} W \right)
 \end{aligned}$$

$$\begin{aligned}
 \frac{d}{dx} (Y^T) &= \frac{d}{dx} \left(\begin{bmatrix} y_{11} & y_{12} & \cdots & y_{1n} \\ y_{21} & y_{22} & \cdots & y_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ y_{m1} & y_{m2} & \cdots & y_{mn} \end{bmatrix}^T \right) \\
 &= \frac{d}{dx} \begin{bmatrix} y_{11} & y_{21} & \cdots & y_{n1} \\ y_{12} & y_{22} & \cdots & y_{n2} \\ \vdots & \vdots & \ddots & \vdots \\ y_{1n} & y_{2n} & \cdots & y_{nn} \end{bmatrix} \\
 &= \begin{bmatrix} y'_{11} & y'_{21} & \cdots & y'_{n1} \\ y'_{12} & y'_{22} & \cdots & y'_{n2} \\ \vdots & \vdots & \ddots & \vdots \\ y'_{1n} & y'_{2n} & \cdots & y'_{nn} \end{bmatrix} \\
 &= \begin{bmatrix} y'_{11} & y'_{12} & \cdots & y'_{1n} \\ y'_{21} & y'_{22} & \cdots & y'_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ y'_{m1} & y'_{m2} & \cdots & y'_{mn} \end{bmatrix}^T \\
 &= \left(\frac{d}{dx} \begin{bmatrix} y_{11} & y_{12} & \cdots & y_{1n} \\ y_{21} & y_{22} & \cdots & y_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ y_{m1} & y_{m2} & \cdots & y_{mn} \end{bmatrix} \right)^T
 \end{aligned}$$

$$\begin{aligned}
 \frac{d}{dx} (Y^{-1}) &= \frac{d}{dx} \frac{\text{adj} Y}{|Y|} \\
 &\vdots \\
 &\text{no proof at this time} \\
 &\vdots \\
 &= -Y^{-1} \left(\frac{d}{dx} Y \right) Y^{-1}
 \end{aligned}$$



E.3 Second derivative of a scalar with respect to a vector

Definition E.4. ¹¹ Let

$$\mathbf{x} \triangleq \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}$$

¹¹ Lieb and Loss (2001), page 240, Horn and Johnson (1990), page 167

The **Hessian matrix** of a scalar y with respect to the vector \mathbf{x} is

DEF

$$\frac{\partial^2 y}{\partial \mathbf{x}^2} = \frac{\partial}{\partial \mathbf{x}} \left(\frac{\partial y}{\partial \mathbf{x}} \right) = \frac{\partial}{\partial \mathbf{x}} \begin{bmatrix} \frac{\partial y}{\partial x_1} \\ \frac{\partial y}{\partial x_2} \\ \vdots \\ \frac{\partial y}{\partial x_n} \end{bmatrix} = \underbrace{\begin{bmatrix} \frac{\partial}{\partial x_1} \frac{\partial y}{\partial x_1} & \frac{\partial}{\partial x_1} \frac{\partial y}{\partial x_2} & \cdots & \frac{\partial}{\partial x_1} \frac{\partial y}{\partial x_n} \\ \frac{\partial}{\partial x_2} \frac{\partial y}{\partial x_1} & \frac{\partial}{\partial x_2} \frac{\partial y}{\partial x_2} & \cdots & \frac{\partial}{\partial x_2} \frac{\partial y}{\partial x_n} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial}{\partial x_n} \frac{\partial y}{\partial x_1} & \frac{\partial}{\partial x_n} \frac{\partial y}{\partial x_2} & \cdots & \frac{\partial}{\partial x_n} \frac{\partial y}{\partial x_n} \end{bmatrix}}_{n \times n \text{ matrix}}$$

E.4 Multiple derivatives of a vector with respect to a scalar

Definition E.5. Let

$$\mathbf{y} \triangleq \begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_m \end{bmatrix}$$

The derivative of a vector \mathbf{y} with respect to the scalar x is

DEF

$$\begin{bmatrix} \mathbf{y} \\ \frac{d}{dx} \mathbf{y} \\ \frac{d^2}{dx^2} \mathbf{y} \\ \vdots \\ \frac{d^n}{dx^n} \mathbf{y} \end{bmatrix} = \underbrace{\begin{bmatrix} y_1 & y_2 & \cdots & y_m \\ \frac{d}{dx} y_1 & \frac{d}{dx} y_2 & \cdots & \frac{d}{dx} y_m \\ \frac{d^2}{dx^2} y_1 & \frac{d^2}{dx^2} y_2 & \cdots & \frac{d^2}{dx^2} y_m \\ \vdots & \vdots & \ddots & \vdots \\ \frac{d^n}{dx^n} y_1 & \frac{d^n}{dx^n} y_2 & \cdots & \frac{d^n}{dx^n} y_m \end{bmatrix}}_{(n+1) \times m \text{ matrix}}$$

APPENDIX F

LINEAR SPACES



“The geometric calculus, in general, consists in a system of operations on geometric entities, and their consequences, analogous to those that algebra has on the numbers. It permits the expression in formulas of the results of geometric constructions, the representation with equations of propositions of geometry, and the substitution of a transformation of equations for a verbal argument.”¹

Giuseppe Peano (1858–1932), Italian mathematician, credited with being one of the first to introduce the concept of the *linear space* (*vector space*).¹

F.1 Definition and basic results

A *metric space* is a set together with nothing else save a *metric* that gives the space a *topology* (Definition ?? page ??). A *linear space* (next definition) in general has no topology but does have some additional *algebraic* structure that is useful in generalizing a number of mathematical concepts. If one wishes to have both algebraic structure and a topology, then this can be accomplished by appending a *topology* to a *linear space* giving a *topological linear space* (Definition ?? page ??), a *metric* giving a *metric linear space*, an *inner product* giving an *inner product space* (Definition I.1 page 233), or a *norm* giving a *normed linear space* (Definition J.1 page 249).

Definition F.1. ² Let $(\mathbb{F}, +, \cdot, 0, 1)$ be a FIELD. Let X be a set, let $+$ be an OPERATOR (Definition M.1 page 281) in X^{X^2} , and let \otimes be an operator in $X^{\mathbb{F} \times X}$.

¹ quote: Peano (1888b), page ix

image http://en.wikipedia.org/wiki/File:Giuseppe_Peano.jpg, public domain

² Kubrusly (2001) pages 40–41 (Definition 2.1 and following remarks), Haaser and Sullivan (1991), page 41, Halmos (1948), pages 1–2, Peano (1888a) (Chapter IX), Peano (1888b), pages 119–120, Banach (1922) pages 134–135




Figure F.1: Lattice of mathematical spaces

The structure $\Omega \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}))$ is a **linear space** over $(\mathbb{F}, +, \cdot, 0, 1)$ if

- | | | |
|-----|---|-------------------------------|
| DEF | 1. $\exists 0 \in X$ such that $x + 0 = x \quad \forall x \in X$ | (+ IDENTITY) |
| | 2. $\exists y \in X$ such that $x + y = 0 \quad \forall x \in X$ | (+ INVERSE) |
| | 3. $(x + y) + z = x + (y + z) \quad \forall x, y, z \in X$ | (+ is ASSOCIATIVE) |
| | 4. $x + y = y + x \quad \forall x, y \in X$ | (+ is COMMUTATIVE) |
| | 5. $1 \cdot x = x \quad \forall x \in X$ | (· IDENTITY) |
| | 6. $\alpha \cdot (\beta \cdot x) = (\alpha \cdot \beta) \cdot x \quad \forall \alpha, \beta \in S \text{ and } x \in X$ | (· ASSOCIATES with ·) |
| | 7. $\alpha \cdot (x + y) = (\alpha \cdot x) + (\alpha \cdot y) \quad \forall \alpha \in S \text{ and } x, y \in X$ | (· DISTRIBUTES over +) |
| | 8. $(\alpha + \beta) \cdot x = (\alpha \cdot x) + (\beta \cdot x) \quad \forall \alpha, \beta \in S \text{ and } x \in X$ | (· PSEUDO-DISTRIBUTES over +) |

The set X is called the **underlying set**. The elements of X are called **vectors**. The elements of \mathbb{F} are called **scalars**. A LINEAR SPACE is also called a **vector space**. If $\mathbb{F} \triangleq \mathbb{R}$, then Ω is a **real linear space**. If $\mathbb{F} \triangleq \mathbb{C}$, then Ω is a **complex linear space**.

Definition F.2. Let $L_1 \triangleq (X_1, +, \cdot, (\mathbb{F}_1, \dot{+}, \dot{\times}))$ and $L_2 \triangleq (X_2, +, \cdot, (\mathbb{F}_2, \dot{+}, \dot{\times}))$.

Ω_2 is a **linear subspace** of Ω_1 if

- | | |
|-----|--|
| DEF | 1. L_1 is a LINEAR SPACE (Definition F.1 page 185) and |
| | 2. L_2 is a LINEAR SPACE (Definition F.1 page 185) and |
| | 3. $\mathbb{F}_2 \subseteq \mathbb{F}_1$ and |
| | 4. $X_2 \subseteq X_1$ and |

Remark F.1. ³ By the first four conditions (*) listed in Definition F.1, $(X, +)$ is a **commutative group** (or **abelian group**).

³ Akhiezer and Glazman (1993), page 1, Haaser and Sullivan (1991), page 41

Often when discussing a linear space, the operator \cdot is simply expressed with juxtaposition (e.g. αx is equivalent to $\alpha \cdot x$). In doing this, there is no risk of ambiguity between scalar-vector multiplication and scalar-scalar multiplication because the operands uniquely identify the precise operator.⁴

Example F.1 (tuples in \mathbb{F}^N).⁵ Let $(x_n)_1^N$ be an N -tuple (Definition P.1 page 329) over a field $(\mathbb{F}, +, \cdot, 0, 1)$.

E X	$X \triangleq \{(x_n)_1^N x_n \in \mathbb{F}\}$ and $(x_n)_1^N + (y_n)_1^N \triangleq (x_n + y_n)_1^N \quad \forall (x_n)_1^N \in X \quad \text{and}$ $\alpha \cdot (x_n)_1^N \triangleq (\alpha \times x_n)_1^N \quad \forall (x_n)_1^N \in X, \alpha \in \mathbb{F}.$
----------------	---

Then the structure $(X, +, \cdot, (\mathbb{F}, +, \times))$ is a *linear space*.

Example F.2 (real numbers).⁶ Let $(\mathbb{R}, +, \cdot, 0, 1)$ be the field of real numbers.

E X	The structure $(\mathbb{R}, +, \cdot, (\mathbb{R}, +, \cdot))$ is a <i>linear space</i> . That is, the field of real numbers forms a linear space over itself.
----------------	---

Example F.3 (functions).⁷ Let $(\mathbb{F}, +, \cdot, 0, 1)$ be a field. Let Y^X be the set of all functions with domain X and range Y .

E X	Let $[f + g](x) \triangleq f(x) + g(x) \quad \forall f, g \in Y^X \quad (\text{pointwise addition})$ and $[\alpha \cdot f](x) \triangleq \alpha \cdot [f(x)] \quad \forall f \in Y^X, \alpha \in \mathbb{F}.$
----------------	--

Then the structure $(Y^X, +, \cdot, (\mathbb{F}, +, \times))$ is a *linear space*.

Example F.4 (functions onto \mathbb{F}).⁸ Let $(\mathbb{F}, +, \cdot, 0, 1)$ be a field. Let \mathbb{F}^X be the set of all functions with domain X and range \mathbb{F} .

E X	Let $[f + g](x) \triangleq f(x) + g(x) \quad \forall f, g \in \mathbb{F}^X \quad (\text{pointwise addition})$ and $[\alpha \cdot f](x) \triangleq \alpha \cdot [f(x)] \quad \forall f \in \mathbb{F}^X, \alpha \in \mathbb{F}.$
----------------	--

Then the structure $(\mathbb{F}^X, +, \cdot, (\mathbb{F}, +, \cdot))$ is a *linear space*.

Theorem F.1 (Additive identity properties).⁹ Let $(X, +, \cdot, (\mathbb{F}, +, \times))$ be a linear space, 0 the ADDITIVE IDENTITY ELEMENT (Definition ?? page ??) with respect to $+$, and $\mathbb{0}$ the ADDITIVE IDENTITY ELEMENT with respect to \cdot .

T H M	1. $0x = \mathbb{0} \quad \forall x \in X$ 2. $\alpha\mathbb{0} = \mathbb{0} \quad \forall \alpha \in \mathbb{F}$ 3. $\alpha x = \mathbb{0} \implies \alpha = 0 \text{ or } x = \mathbb{0}$ 4. $x + x = x \implies x = \mathbb{0}$ 5. $\alpha \neq 0 \text{ and } x \neq \mathbb{0} \implies \alpha x \neq \mathbb{0}$
----------------------	--

PROOF:

⁴ *Operator overload* is a technique in which two fundamentally different operators or functions share the same symbol or label. It is inherent in the programming language C++ and is therein called *operator overload*. In C++, you can define two (or more) operators or functions that share the same symbol or name, but yet are completely different. Two such operators (or functions) are distinguished from each other by the type of their operands. So for example, in C++, you can define an $m \times n$ matrix *type* and use operator overload to define a $+$ operator that operates on this new matrix type. So if variables x and y are of floating point type and A and B are of the matrix type, you can then add either type using the same syntax style:

$x+y$ (add two floating point numbers)
 $A+B$ (add two matrices)

Even though both of these operations “look” the same, they are of course fundamentally different.

⁵  Kubrusly (2001) page 41 (Example 2D)

⁶  Kubrusly (2001) page 41 (Example 2D),  Hamel (1905)

⁷  Kubrusly (2001) page 42 (Example 2F)

⁸  Kubrusly (2001) page 41 (Example 2E)

⁹  Berberian (1961) page 6 (Theorem 1),  Michel and Herget (1993) page 77



1. Proof that $0x = \emptyset$:

$$\begin{aligned}
 0x &= 0x + 0\emptyset && \text{by definition of } + \text{ additive identity element} \\
 &= 0x + 0x + (-0x) && \text{by definition of } + \text{ additive inverse} \\
 &= (0 + 0)x + (-0 \cdot x) && \text{by definition of } + \text{ additive identity element} \\
 &= 0x + (-0x) && \text{by Definition F.1 property 4} \\
 &= \emptyset && \text{by definition of } + \text{ additive identity element}
 \end{aligned}$$

2. Proof that $\alpha\emptyset = \emptyset$:

$$\begin{aligned}
 \alpha\emptyset &= \alpha(0x) && \text{by item 1} \\
 &= (\alpha 0)x && \text{by Definition F.1 property 6} \\
 &= 0x \\
 &= \emptyset && \text{by item 1}
 \end{aligned}$$

3. Proof that $\alpha \neq 0$ and $x \neq \emptyset \implies \alpha x \neq \emptyset$: Suppose $\alpha x = \emptyset$. Then

$$\begin{aligned}
 x &= \left(\frac{1}{\alpha}\right)x \\
 &= \frac{1}{\alpha}(\alpha x) \\
 &= \frac{1}{\alpha}\emptyset \\
 &= \emptyset && \text{by item 2} \\
 \implies x &= \emptyset
 \end{aligned}$$

This is a *contradiction* and so $\alpha x \neq \emptyset$.

4. Proof that $\alpha x = \emptyset \implies \alpha = 0$ or $x = \emptyset$: contrapositive argument of item 3

5. Proof that $x + x = x \implies x = \emptyset$:

$$\begin{aligned}
 x &= x + \emptyset && \text{by } \textit{additive identity property} \text{ (Definition F.1 page 185)} \\
 &= x + [x + (-x)] && \text{by } \textit{additive inverse property} \text{ (Definition F.1 page 185)} \\
 &= [x + x] + (-x) && \text{by } \textit{associative property} \text{ (Definition F.1 page 185)} \\
 &= x + (-x) && \text{by left hypothesis} \\
 &= \emptyset && \text{by } \textit{additive inverse property} \text{ (Definition F.1 page 185)}
 \end{aligned}$$

Definition F.3. ¹⁰ Let $\Omega \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}))$ be a linear space with vectors $x, y \in X$. Let $-y$ be the additive inverse of y such that $y + (-y) = \emptyset$.

D E F The **difference** of x and y is $x + (-y)$ and is denoted $x - y$.

Theorem F.2 (Additive inverse properties). ¹¹ Let $\Omega \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}))$ be a linear space, \emptyset the ADDITIVE IDENTITY ELEMENT with respect to $+$, and $-x$ the ADDITIVE INVERSE (Definition ?? page ??) of x with respect to $+$.

T H M	1. $x + y = \emptyset \implies x = -y \quad \forall x, y \in X$ (additive inverse is UNIQUE) 2. $(-\alpha)x = -(\alpha x) = \alpha(-x) \quad \forall x \in X, \alpha \in \mathbb{F}$ 3. $\alpha(x - y) = \alpha x - \alpha y \quad \forall x, y \in X, \alpha \in \mathbb{F}$ (DISTRIBUTIVE) 4. $(\alpha - \beta)x = \alpha x - \beta x \quad \forall x \in X, \alpha, \beta \in \mathbb{F}$ (DISTRIBUTIVE)
-------	---

¹⁰ Berberian (1961) page 7 (Definition 1)

¹¹ Berberian (1961) page 7 (Corollary 1), Michel and Herget (1993) page 77, Peano (1888a) (Chapter IX), Peano (1888b) pages 119–120, Banach (1922) pages 134–135

PROOF:

1. Proof that $x + y = 0 \implies x = -y$:

$$\begin{aligned} x &= x - 0 \\ &= x - (x + y) && \text{by left hypothesis} \\ &= (x - x) - y \\ &= 0 - y \\ &= -y \end{aligned}$$

2. Proof that $(-\alpha)x = -(\alpha x)$:

$$\begin{aligned} 0 &= 0x && \text{by Theorem F.1 page 187} \\ &= (\alpha - \alpha)x \\ &= [\alpha + (-\alpha)]x && \text{by field property of } \mathbb{F} \\ &= \alpha x + (-\alpha)x && \text{by Definition F.1 page 185} \\ \implies -(\alpha x) &= (-\alpha)x && \text{by item (1) page 189} \end{aligned}$$

3. Proof that $\alpha(-x) = -(\alpha x)$:

$$\begin{aligned} 0 &= \alpha 0 && \text{by Theorem F.1 page 187} \\ &= \alpha[x + (-x)] \\ &= \alpha x + \alpha(-x) && \text{by definition of additive identity element } -x \\ &= \alpha x + \alpha(-x) && \text{by Definition F.1 page 185} \\ \implies -(\alpha x) &= \alpha(-x) && \text{by item (1) page 189} \end{aligned}$$

4. Proof that $\alpha(x - y) = \alpha x - \alpha y$:

$$\begin{aligned} \alpha(x - y) &= \alpha[x + (-y)] && \text{by Definition F.3 page 188} \\ &= \alpha x + \alpha(-y) \\ &= \alpha x + (-\alpha y) && \text{by item (3) page 189} \\ &= \alpha x - \alpha y && \text{by Definition F.3 page 188} \end{aligned}$$

5. Proof that $(\alpha - \beta)x = \alpha x - \beta x$:

$$\begin{aligned} (\alpha - \beta)x &= [\alpha + (-\beta)]x && \text{by field properties of } \mathbb{F} \\ &= \alpha x + (-\beta)x \\ &= \alpha x + [-(\beta x)] && \text{by item (2) page 189} \\ &= \alpha x - (\beta x) && \text{by Definition F.3 page 188} \end{aligned}$$

⇒

Theorem F.3. ¹² Let $(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}))$ be a linear space, 0 the additive identity element with respect to $+$, and $-x$ additive inverse of x with respect to $+$.

T
H
M

1. $\alpha x = \alpha y$ and $\alpha \neq 0 \implies x = y \quad \forall x, y \in X$
2. $\alpha x = \beta x$ and $x \neq 0 \implies \alpha = \beta \quad \forall x, y \in X, \alpha, \beta \in \mathbb{F}$
3. $z + x = z + y \implies x = y \quad \forall x, y, z \in X$

¹² Michel and Herget (1993) page 77, Peano (1888a) (Chapter IX), Peano (1888b) pages 119–120, Banach (1922) pages 134–135

PROOF:

1. Proof that $\alpha x = \alpha y$ and $\alpha \neq 0 \implies x = y$:

$$\begin{aligned} 0 &= \frac{1}{\alpha}(0) && \text{by left hypothesis } (\alpha \neq 0) \\ &= \frac{1}{\alpha}(\alpha x - \alpha y) && \text{by left hypothesis } (\alpha x = \alpha y) \\ &= \frac{1}{\alpha}\alpha(x - y) && \text{by Definition E1 page 185} \\ &= x - y \end{aligned}$$

2. Proof that $\alpha x = \beta x$ and $x \neq 0 \implies \alpha = \beta$:

$$\begin{aligned} 0 &= \alpha x + (-\alpha x) && \text{by definition of additive inverse} \\ &= \beta x + (-\alpha x) && \text{by left hypothesis} \\ &= \beta x + (-\alpha)x && \text{by Theorem F2 page 188} \\ &= [\beta + (-\alpha)]x && \text{by Definition E1 page 185} \\ \implies \beta - \alpha &= 0 && \text{by Theorem E1 page 187} \\ \implies \alpha &= \beta && \text{by field properties of } \mathbb{F} \end{aligned}$$

3. Proof that $z + x = z + y \implies x = y$:

$$\begin{aligned} 0 &= (z + x) - (z + y) && \text{by Definition E1 property 1} \\ &= (x + z) - (z + y) && \text{by Definition E1 property 3} \\ &= (x + z) + [(-1)z + (-1)y] && \text{by previous result 2.} \\ &= (x + z) + (-z - y) \\ &= x + (z - z) - y \\ &= x - y \end{aligned}$$

F.2 Order on Linear Spaces

Definition F.4. ¹³ Let $\Omega \triangleq (X, +, \cdot, (\mathbb{R}, \dot{+}, \dot{\times}))$ be a real linear space.

The pair (Ω, \leq) is an ordered linear space if

- | | | |
|----------------------|--|-----|
| D
E
F | <ol style="list-style-type: none"> 1. $x \leq y \implies x + z \leq y + z \quad \forall z \in X$ 2. $x \leq y \implies \alpha x \leq \alpha y \quad \forall \alpha \in \mathbb{F}$ | and |
|----------------------|--|-----|

A vector x is positive if $0 \leq x$.

The positive cone X^+ of (X, \leq) is the set $X^+ \triangleq \{x \in X | 0 \leq x\}$.

Definition F.5. ¹⁴ Let (X, \leq) be an ordered linear space.

The tuple $L \triangleq (X, \vee, \wedge; \leq)$ is a Riesz space if L is a lattice.

A RIESZ SPACE is also called a vector lattice.

Theorem F.4. ¹⁵ Let $(X, \vee, \wedge; \leq)$ be a Riesz space (Definition F.5 page 190).

T H M	$x \vee y = -[(-x) \wedge (-y)]$ $x + (y \vee z) = (x + y) \vee (x + z)$ $\alpha(x \vee y) = (\alpha x) \vee (\alpha y)$ $x + y = (x \wedge y) + (x \vee y)$	$x \wedge y = -[(-x) \vee (-y)]$ $x + (y \wedge z) = (x + y) \wedge (x + z)$ $\alpha(x \wedge y) = (\alpha x) \wedge (\alpha y)$	$\forall x, y \in X$ $\forall x, y, z \in X$ $\forall x, y \in X, \alpha \geq 0$ $\forall x, y \in X, \alpha \in \mathbb{F}$
----------------------	---	--	---

¹³ Aliprantis and Burkinshaw (2006) pages 1-2

¹⁴ Aliprantis and Burkinshaw (2006) page 2

¹⁵ Aliprantis and Burkinshaw (2006) page 3 (Theorem 1.2)

PROOF:

1. Proof that $x \vee y = -[(-x) \wedge (-y)]$:

$(-x) \wedge (-y) \leq -x$	$(-x) \wedge (-y) \leq -y$
$x \leq -[(-x) \wedge (-y)]$	$y \leq -[(-x) \wedge (-y)]$
$x \vee y \leq -[(-x) \wedge (-y)]$	
$x \leq x \vee y$	$y \leq x \vee y$
$-(x \vee y) \leq -x$	$-(x \vee y) \leq -y$
$-(x \vee y) \leq (-x) \wedge (-y)$	
$-[(-x) \wedge (-y)] \leq x \vee y$	

2. Proof that $x \wedge y = -[(-x) \vee (-y)]$:

$x \vee y = -[(-x) \wedge (-y)]$	by item (1)
$(-x) \vee (-y) = -[(-(-x)) \wedge (-(-y))]$	replace x with $-x$ and y with y
$(-x) \vee (-y) = -[x \wedge y]$	$-(-x) = x$
$-[x \wedge y] = (-x) \vee (-y)$	by symmetry of $=$ relation
$x \wedge y = -[(-x) \vee (-y)]$	multiply both sides by -1

3. Proof that $x + (y \vee z) = (x + y) \vee (x + z)$:

$x + y \leq x + (y \vee z)$	$x + z \leq x + (y \vee z)$
$(x + y) \vee (x + z) \leq x + (y \vee z)$	
$y = -x + (x + y)$	$z = -x + (x + z)$
$\leq -x + [(x + y) \vee (x + z)]$	$\leq -x + [(x + y) \vee (x + z)]$
$y \vee z \leq -x + [(x + y) \vee (x + z)]$	
$x + (y \vee z) \leq (x + y) \vee (x + z)$	

4. Proof that $x + (y \wedge z) = (x + y) \wedge (x + z)$:

$x + y \geq x + (y \wedge z)$	$x + z \geq x + (y \wedge z)$
$(x + y) \wedge (x + z) \geq x + (y \wedge z)$	
$y = -x + (x + y)$	$z = -x + (x + z)$
$\geq -x + [(x + y) \wedge (x + z)]$	$\geq -x + [(x + y) \wedge (x + z)]$
$y \wedge z \geq -x + [(x + y) \wedge (x + z)]$	
$x + (y \wedge z) \geq (x + y) \wedge (x + z)$	

5. Proof that $\alpha(x \vee y) = (\alpha x) \vee (\alpha y)$ for $\alpha \geq 0$:

$x \leq x \vee y$	$y \leq x \vee y$	
$\alpha x \leq \alpha(x \vee y)$	$\alpha y \leq \alpha(x \vee y)$	by Definition F.4 page 190
$(\alpha x) \vee (\alpha y) \leq \alpha(x \vee y)$		
$\alpha x \leq (\alpha x) \vee (\alpha y)$	$\alpha y \leq (\alpha x) \vee (\alpha y)$	
$x \leq \alpha^{-1}(\alpha x) \vee (\alpha y)$	$y \leq \alpha^{-1}(\alpha x) \vee (\alpha y)$	
$x \vee y \leq \alpha^{-1}(\alpha x) \vee (\alpha y)$		
$\alpha(x \vee y) \leq (\alpha x) \vee (\alpha y)$		

6. Proof that $\alpha(x \wedge y) = (\alpha x) \wedge (\alpha y)$ for $\alpha \geq 0$:

$x \geq x \wedge y$	$y \geq x \wedge y$	
$\alpha x \geq \alpha(x \wedge y)$	$\alpha y \geq \alpha(x \wedge y)$	
$(\alpha x) \wedge (\alpha y) \geq \alpha(x \wedge y)$		by Definition F.4 page 190

$\alpha x \geq (\alpha x) \wedge (\alpha y)$	$\alpha y \geq (\alpha x) \wedge (\alpha y)$	
$x \geq \alpha^{-1}(\alpha x) \wedge (\alpha y)$	$y \geq \alpha^{-1}(\alpha x) \wedge (\alpha y)$	
$x \wedge y \geq \alpha^{-1}(\alpha x) \wedge (\alpha y)$		
$\alpha(x \wedge y) \geq (\alpha x) \wedge (\alpha y)$		

7. Proof that $x + y = (x \wedge y) + (x \vee y)$:

$x \leq x \vee y$	$y \leq x \vee y$
$x + y \leq (x \vee y) + y$	$x + vy \leq x + (x \vee y)$
$x + y - (x \vee y) \leq y$	$x + vy - (x \vee y) \leq x$
$x + y - (x \vee y) \leq x \wedge y$	
$x + y \leq (x \vee y) + (x \wedge y)$	
$x \wedge y \leq x$	$x \wedge y \leq y$
$0 \leq x - (x \wedge y)$	$0 \leq y - (x \wedge y)$
$y \leq y + x - (x \wedge y)$	$x \leq x + y - (x \wedge y)$
$y \leq x + y - (x \wedge y)$	$x \leq x + y - (x \wedge y)$
$x \vee y \leq x + y - (x \wedge y)$	
$(x \wedge y) + (x \vee y) \leq x + y$	



Definition F.6. ¹⁶ Let $(X, \vee, \wedge; \leq)$ be a RIESZ SPACE (Definition F.5 page 190).

D E F x^+ is defined as $x^+ \triangleq x \vee \emptyset$ and is called the **positive part** of x .
 x^- is defined as $x^- \triangleq (-x) \vee \emptyset$ and is called the **negative part** of x .
 $|x|$ is defined as $|x| \triangleq x \vee (-x)$ and is called the **absolute value** of x .

Theorem F.5. ¹⁷ Let $(X, \vee, \wedge; \leq)$ be a RIESZ SPACE (Definition F.5 page 190).

T H M	$y - z = x$ and $y \wedge z = \emptyset$	\Leftrightarrow	$\left\{ \begin{array}{l} y = x^+ \text{ and} \\ z = x^- \end{array} \right.$
-------	---	-------------------	---

PROOF:

1. Proof that left hypothesis \implies right hypothesis:

$$\begin{aligned}
 x^+ &= x \vee \emptyset && \text{by definition of } x^+ \text{ Definition F.6 page 192} \\
 &= (y - z) \vee \emptyset && \text{by left hypothesis} \\
 &= (y - z) \vee (z - z) \\
 &= (y \vee z) - z && \text{by Theorem F.4 page 190} \\
 &= [y + z - (y \wedge z)] - z && \text{by Theorem F.4 page 190} \\
 &= y - (y \wedge z) \\
 &= y - \emptyset && \text{by left hypothesis} \\
 &= y \\
 x^- &= (-x) \vee \emptyset && \text{by definition of } x^- \text{ Definition F.6 page 192} \\
 &= (z - y) \vee \emptyset && \text{by left hypothesis} \\
 &= (z - y) \vee (y - y) \\
 &= (z \vee y) - y && \text{by Theorem F.4 page 190}
 \end{aligned}$$

¹⁶ Aliprantis and Burkinshaw (2006) page 4, Istrătescu (1987) page 129

¹⁷ Aliprantis and Burkinshaw (2006) page 4 (Theorem 1.3)

$$\begin{aligned}
 &= [z + y - (z \wedge y)] - z && \text{by Theorem F.4 page 190} \\
 &= z - (z \wedge y) \\
 &= z - \emptyset && \text{by left hypothesis} \\
 &= z
 \end{aligned}$$

2. Proof that left hypothesis \iff right hypothesis:

$$\begin{aligned}
 y - z &= x^+ - x^- && \text{by right hypothesis} \\
 &= [x \vee \emptyset] - [(-x) \vee \emptyset] && \text{by Definition F.6 page 192} \\
 &= (x \vee \emptyset) + (x \wedge \emptyset) && \text{by Theorem F.4 page 190} \\
 &= x && \text{by Theorem F.4 page 190} \\
 y \wedge z &= x^+ \wedge x^- && \text{by right hypothesis} \\
 &= [x^- + (x^+ - x^-)] \wedge [x^- + \emptyset] && \text{by Theorem F.4 page 190} \\
 &= x^- + [(x^+ - x^-) \wedge \emptyset] && \text{by right hypothesis} \\
 &= x^- + [(y - z) \wedge \emptyset] && \text{by previous result} \\
 &= x^- + (x \wedge \emptyset) && \text{by Theorem F.4 page 190} \\
 &= x^- - [-x \vee \emptyset] && \text{by definition of } x^- \text{ (Definition F.6 page 192)} \\
 &= x^- - x && \text{by definition of } x^- \text{ (Definition F.6 page 192)} \\
 &= \emptyset
 \end{aligned}$$



Theorem F.6. ¹⁸ Let $(X, \vee, \wedge; \leq)$ be a RIESZ SPACE (Definition F.5 page 190). Let x^+ the POSITIVE PART of $x \in X$, x^- the NEGATIVE PART of $x \in X$, and $|x|$ the ABSOLUTE VALUE (Definition F.6 page 192) of $x \in X$.

T H M	$ x = x^+ + x^-$ $x = (x - y)^+ + (x \wedge y)$ $\forall x \in X$
-------------	---



PROOF:

$$\begin{aligned}
 |x| &= x \vee (-x) && \text{by definition of } |x| \text{ (Definition F.6 page 192)} \\
 &= (2x - x) \vee (\emptyset - x) \\
 &= (-x + 2x) \vee (-x + \emptyset) && \text{by commutative property (Definition F.1 page 185)} \\
 &= (-x) + (2x \vee \emptyset) && \text{by Theorem F.4 page 190} \\
 &= (2x \vee \emptyset) - x && \text{by the commutative property (Definition F.1 page 185)} \\
 &= 2(x \vee \emptyset) - x && \text{by Theorem F.4 page 190} \\
 &= 2x^+ - x && \text{by definition of } x^+ \text{ (Definition F.6 page 192)} \\
 &= 2x^+ - (x^+ - x^-) && \text{by 1} \\
 &= x^+ + x^- \\
 x &= x + \emptyset && x + y - y \\
 &= (x \vee y) + (x \wedge y) - y && \text{by Theorem F.4 page 190} \\
 &= [(x - y) \vee (y - y)] + (x \wedge y) && \text{by Theorem F.4 page 190} \\
 &= [(x - y) \vee \emptyset] + (x \wedge y) \\
 &= (x - y)^+ + (x \wedge y) && \text{by definition of } x^+ \text{ (Definition F.6 page 192)}
 \end{aligned}$$



¹⁸ Aliprantis and Burkinshaw (2006) page 4

Theorem E.7. ¹⁹ Let $(X, \vee, \wedge; \leq)$ be a RIESZ SPACE (Definition F.5 page 190). Let x^+ the POSITIVE PART of $x \in X$, x^- the NEGATIVE PART of $x \in X$, and $|x|$ the ABSOLUTE VALUE (Definition F.6 page 192) of $x \in X$.

T
H
M

1. $x \vee y = \frac{1}{2}(x + y + |x - y|) \quad \forall x, y \in X$
2. $x \wedge y = \frac{1}{2}(x + y - |x - y|) \quad \forall x, y \in X$
3. $|x - y| = (x \vee y) - (x \wedge y) \quad \forall x, y \in X$
4. $|x| \vee |y| = \frac{1}{2}(|x + y| + |x - y|) \quad \forall x, y \in X$
5. $|x| \wedge |y| = \frac{1}{2}(|x + y| - |x - y|) \quad \forall x, y \in X$

PROOF:

$$(x + y + |x - y|) = (x + y) + [(x - y) \vee (y - x)] \quad \text{by Definition F.6 page 192}$$

$$= [(x + y) + (x - y)] \vee [(x + y) + (y - x)] \quad \text{by Theorem F.4 page 190}$$

$$= (2x) \vee (2y)$$

$$= 2(x \vee y) \quad \text{by Theorem F.4 page 190}$$

$$(x + y - |x - y|) = (x + y) - [(x - y) \vee (y - x)] \quad \text{by Definition F.6 page 192}$$

$$= (x + y) - [(-(y - x)) \vee (-(x - y))] \quad \text{by Theorem F.4 page 190}$$

$$= (x + y) + [(y - x) \wedge (x - y)] \quad \text{by Theorem F.4 page 190}$$

$$= [(x + y) + (y - x)] \wedge [(x + y) + (x - y)] \quad \text{by Theorem F.4 page 190}$$

$$= (2y) \wedge (2x)$$

$$= 2(y \wedge x) \quad \text{by Theorem F.4 page 190}$$

$$= 2(x \wedge y) \quad \text{by Theorem F.4 page 190}$$

$$|x - y| = \frac{1}{2}(x + y + |x - y|) - \frac{1}{2}(x + y - |x - y|)$$

$$= (x \vee y) - (x \wedge y) \quad \text{by 1 and 2}$$

$$|x + y| + |x - y| = \frac{1}{2}(\emptyset + |2x + 2y|) + |x - y|$$

$$= \frac{1}{2}[(x + y) + (-x - y) + |(x + y) - (-x - y)|] + |x - y|$$

$$= [(x + y) \vee (-x - y)] + |x - y| \quad \text{by 1}$$

$$= [(x + y) + |x - y|] \vee [(-x - y) + |x - y|] \quad \text{by Theorem F.4 page 190}$$

$$= 2(x \vee y) \vee 2[(-y) + (-x) + |(-y) - (-x)|] \quad \text{by 1}$$

$$= 2(x \vee y) \vee 2[(-y) \vee (-x)] \quad \text{by 1}$$

$$= 2([x \vee (-x)] \vee (y \vee (-y)))$$

$$= 2(|x| \vee |y|) \quad \text{by Definition F.6 page 192}$$

$$||x + y| - |x - y|| = 2(|x + y| \vee |x - y|) - (|x + y| + |x - y|)$$

$$= (|x + y + x - y| + |x + y - x + y|) - 2(|x| \vee |y|) \quad \text{by 1}$$

$$= 2(|x| + |y|) - 2(|x| \vee |y|) \quad \text{by 3}$$

$$= 2(|x| \vee |y|) \quad \text{by Theorem F.4 page 190}$$

Definition F.7. ²⁰ Let $(X, \vee, \wedge; \leq)$ be a RIESZ SPACE (Definition F.5 page 190). Let x^+ the POSITIVE PART of $x \in X$, x^- the NEGATIVE PART of $x \in X$, and $|x|$ the ABSOLUTE VALUE (Definition F.6 page 192) of $x \in X$.

D
E
F

x and y are disjoint, denoted by $x \perp y$, if

$$|x| \wedge |y| = \emptyset.$$

Two subsets U and V of X are disjoint, denoted by $U \perp V$ if

$$x \perp y \quad \forall x \in U \text{ and } y \in V$$

¹⁹ Aliprantis and Burkinshaw (2006) page 5 (Theorem 1.4)

²⁰ Aliprantis and Burkinshaw (2006) page 5

Definition F.8. ²¹ Let $(X, \vee, \wedge; \leq)$ be a RIESZ SPACE (Definition F.5 page 190). Let x^+ the POSITIVE PART of $x \in X$, x^- the NEGATIVE PART of $x \in X$, and $|x|$ the ABSOLUTE VALUE (Definition F.6 page 192) of $x \in X$. Let Y be a subset of X .

D E F Y^d is the **disjoint complement** of Y if $Y^d \triangleq \{x \in X | x \perp y \quad \forall y \in Y\}$.
The quantity Y^{dd} is defined as $(Y^d)^d$.

Definition F.9. ²² Let $(X, \vee, \wedge; \leq)$ be a RIESZ SPACE (Definition F.5 page 190). Let x^+ the POSITIVE PART of $x \in X$, x^- the NEGATIVE PART of $x \in X$, and $|x|$ the ABSOLUTE VALUE (Definition F.6 page 192) of $x \in X$.

D E F	$ A \triangleq \{ a a \in A\}$ $A^+ \triangleq \{a^+ a \in A\}$ $A^- \triangleq \{a^- a \in A\}$ $A \vee B \triangleq \{a \vee b a \in A \text{ and } b \in B\}$ $A \wedge B \triangleq \{a \wedge b a \in A \text{ and } b \in B\}$ $x \vee A \triangleq \{x \vee a a \in A\}$ $x \wedge A \triangleq \{x \wedge a a \in A\}$
-------	--

²¹  Aliprantis and Burkinshaw (2006) page 5

²²  Aliprantis and Burkinshaw (2006) page 7

APPENDIX G

LINEAR COMBINATIONS

G.1 Linear combinations in linear spaces

A *linear space* (Definition F.1 page 185) in general is not equipped with a *topology*. Without a topology, it is not possible to determine whether an *infinite sum* of vectors converges. Therefore in this section (dealing with linear spaces), all definitions related to sums of vectors will be valid for *finite* sums (Definition L.1 page 267) only (finite “ N ”).

Definition G.1. ¹ Let $\{x_n \in X \mid n=1,2,\dots,N\}$ be a set of vectors in a LINEAR SPACE $(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}))$.

D E F A vector $x \in X$ is a **linear combination** of the vectors in $\{x_n\}$ if

there exists $\{\alpha_n \in \mathbb{F} \mid n=1,2,\dots,N\}$ such that
$$x = \sum_{n=1}^N \alpha_n x_n.$$

Definition G.2. ² Let $(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}))$ be a linear space and Y be a subset of X .

D E F The **linear span** of Y is defined as $\text{span}Y \triangleq \left\{ \sum_{y \in Y} \alpha_y y \mid \alpha_y \in \mathbb{F}, y \in Y \right\}.$

The set Y spans a set A if $A \subseteq \text{span}Y$.

Proposition G.1. ³ Let $\{x_n \in X \mid n=1,2,\dots,N\}$ be a set of vectors in a LINEAR SPACE $L \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}))$.

- P R P**
1. $\text{span}\{x_n\}$ is a LINEAR SPACE (Definition F.1 page 185) and
 2. $\text{span}\{x_n\}$ is a LINEAR SUBSPACE of L (Definition F.2 page 186).

Definition G.3. ⁴ Let $L \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}))$ be a LINEAR SPACE.

D E F The set $Y \triangleq \{x_n \in X \mid n=1,2,\dots,N\}$ is **linearly independent** in L if
$$\left\{ \sum_{n=1}^N \alpha_n x_n = 0 \right\} \implies \{\alpha_1 = \alpha_2 = \dots = \alpha_N = 0\}.$$


The set Y is **linearly dependent** in L if Y is not linearly independent in L .

¹ Berberian (1961) page 11 (Definition I.4.1), Kubrusly (2001) page 46

² Michel and Herget (1993) page 86 (3.3.7 Definition), Kurdila and Zabarankin (2005) page 44, Searcoid (2002) page 71 (Definition 3.2.5—more general definition)

³ Kubrusly (2001) page 46

⁴ Bachman and Narici (1966) pages 3–4, Christensen (2003) page 2, Heil (2011) page 156 (Definition 5.7)

Figure G.1: Lattice of *linear combinations*

Definition G.4. ⁵ Let $\{x_n \in X | n=1,2,\dots,N\}$ be a set of vectors in a LINEAR SPACE $L \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}))$.

D E F The set $\{x_n\}$ is a **Hamel basis** for L if

1. $\{x_n\}$ SPANS L (Definition G.2 page 197) and
2. $\{x_n\}$ is LINEARLY INDEPENDENT in L (Definition G.1 page 197) .

A HAMEL BASIS is also called a **linear basis**.

Definition G.5. ⁶ Let $L \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}))$ be a LINEAR SPACE. Let x be a VECTOR in L and $Y \triangleq \{x_n \in X | n=1,2,\dots,N\}$ be a set of vectors in L .

D E F The expression $\sum_{n=1}^N \alpha_n x_n$ is the **expansion** of x on Y in L if $x = \sum_{n=1}^N \alpha_n x_n$.

In this case, the sequence $(\alpha_n)_{n=1}^N$ is the **coordinates** of x with respect to Y in L .
If $\alpha_N \neq 0$, then N is the **dimension** $\dim L$ of L .

Theorem G.1. ⁷ Let $\{x_n | n=1,2,\dots,N\}$ be a HAMEL BASIS (Definition G.4 page 199) for a LINEAR SPACE $(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}))$.

T H M $\left\{ x = \sum_{n=1}^N \alpha_n x_n = \sum_{n=1}^N \beta_n x_n \right\} \implies \underbrace{\alpha_n = \beta_n}_{\text{coordinates of } x \text{ are UNIQUE}} \quad \forall x \in X$

PROOF:

$$\begin{aligned} 0 &= x - x \\ &= \sum_{n=1}^N \alpha_n x_n - \sum_{n=1}^N \beta_n x_n \\ &= \sum_{n=1}^N (\alpha_n - \beta_n) x_n \\ \implies &\{x_n\} \text{ is linearly dependent if } (\alpha_n - \beta_n) \neq 0 \quad \forall n = 1, 2, \dots, N \\ \implies &(\alpha_n - \beta_n) = 0 \quad \forall n = 1, 2, \dots, N \quad (\text{because } \{x_n\} \text{ is a basis and therefore must be linearly independent}) \\ \implies &\alpha_n = \beta_n \text{ for } n = 1, 2, \dots, N \end{aligned}$$

Theorem G.2. ⁸ Let $L \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}))$ be a LINEAR SPACE.

T H M $\left\{ \begin{array}{l} 1. \{x_n \in X | n=1,2,\dots,N\} \text{ is a HAMEL BASIS for } L \\ 2. \{y_n \in X | n=1,2,\dots,M\} \text{ is a set of LINEARLY INDEPENDENT vectors in } L \end{array} \right\} \implies \left\{ \begin{array}{l} 1. M \leq N \\ 2. M = N \implies \{y_n | n=1,2,\dots,M\} \text{ is a BASIS for } L \\ 3. M \neq N \implies \{y_n | n=1,2,\dots,M\} \text{ is NOT a basis for } L \end{array} \right\}$

PROOF:

⁵ Hamel (1905), Bachman and Narici (1966) page 4, Kubrusly (2001) pages 48–49 (Section 2.4), Young (2001) page 1, Carothers (2005) page 25, Heil (2011) page 125 (Definition 4.1)

⁶ Hamel (1905), Bachman and Narici (1966) page 4, Kubrusly (2001) pages 48–49 (Section 2.4), Young (2001) page 1, Carothers (2005) page 25, Heil (2011) page 125 (Definition 4.1)

⁷ Michel and Herget (1993) pages 89–90 (Theorem 3.3.25)

⁸ Michel and Herget (1993) pages 90–91 (Theorem 3.3.26)

1. Proof that $\{y_1, x_1, \dots, x_{N-1}\}$ is a *basis* for L :

(a) Proof that $\{y_1, x_1, \dots, x_{N-1}\}$ spans L :

i. Because $\{x_n|_{n=1,2,\dots,N}\}$ is a *basis* for L , there exists $\beta \in \mathbb{F}$ and $\{\alpha_n \in \mathbb{F}|_{n=1,2,\dots,N}\}$ such that

$$\beta y_1 + \sum_{n=1}^N \alpha_n x_n = 0.$$

ii. Select an n such that $\alpha_n \neq 0$ and renumber (if necessary) the above indices such that

$$x_n = -\frac{\beta}{\alpha_n} y_1 + \sum_{n=1}^{N-1} \frac{\alpha_n}{\alpha_n} x_n.$$

iii. Then, for any $y \in X$, we can write

$$\begin{aligned} y &= \sum_{n=1}^N \gamma_{n \in \mathbb{Z}} x_n \\ &= \left(\sum_{n=1}^{N-1} \gamma_{n \in \mathbb{Z}} x_n \right) + \gamma_{n \in \mathbb{Z}} \left(-\frac{\beta}{\alpha_n} y_1 - \sum_{n=1}^{N-1} \frac{\alpha_n}{\alpha_n} x_n \right) \\ &= -\frac{\beta \gamma_n}{\alpha_n} y_1 + \sum_{n=1}^{N-1} \left(\gamma_n - \frac{\alpha_n \gamma_n}{\alpha_n} \right) x_n \\ &= \delta y_1 + \sum_{n=1}^{N-1} \delta_{n \in \mathbb{Z}} x_n \end{aligned}$$

iv. This implies that $\{y_1, x_1, \dots, x_{N-1}\}$ spans L :

(b) Proof that $\{y_1, x_1, \dots, x_{N-1}\}$ is *linearly independent*:

i. If $\{y_1, x_1, \dots, x_{N-1}\}$ is *linearly dependent*, then there exists $\{\epsilon, \epsilon_1, \dots, \epsilon_{N-1}\}$ such that

$$\epsilon y_1 + \left(\sum_{n=1}^{N-1} \epsilon_{n \in \mathbb{Z}} x_n \right) + 0 x_n = 0.$$

ii. item (1(b)i) implies that the coordinate of y_1 associated with x_n is 0.

$$y_1 = -\left(\sum_{n=1}^{N-1} \frac{\epsilon_n}{\epsilon} x_n \right) + 0 x_n = 0.$$

iii. item (1(a)i) implies that the coordinate of y_1 associated with x_n is *not* 0.

$$y_1 = -\sum_{n=1}^N \frac{\alpha_n}{\beta} x_n.$$

iv. This implies that item (1(b)i) (that the set is linearly dependent) is *false* because item (1(b)ii) and item (1(b)iii) contradict each other.

v. This implies $\{y_1, x_1, \dots, x_{N-1}\}$ is *linearly independent*.

2. Proof that $\{y_1, y_2, x_1, \dots, x_{N-2}\}$ is a *basis*: Repeat item (1).

3. Suppose $m = n$. Proof that $\{y_1, y_2, \dots, y_M\}$ is a *basis*: Repeat item (1) $M - 1$ times.

4. Proof that $M \not> N$:

(a) Suppose that $M = N + 1$.

(b) Then because $\{y_n|_{n=1,2,\dots,N}\}$ is a *basis*, there exists $\{\zeta_n|_{n=1,2,\dots,N+1}\}$ such that

$$\sum_{n=1}^{N+1} \zeta_{n \in \mathbb{Z}} y_{n \in \mathbb{Z}} = 0.$$

(c) This implies that $\{y_n|_{n=1,2,\dots,N+1}\}$ is *linearly dependent*.

(d) This implies that $\{y_n|_{n=1,2,\dots,N+1}\}$ is *not* a basis.

(e) This implies that $M > N$.

5. Proof that $M \neq N \implies \{y_n|_{n=1,2,\dots,M}\}$ is *not* a basis for L :

(a) Proof that $M > N \implies \{y_n|_{n=1,2,\dots,M}\}$ is *not* a basis for L : same as in item (4).

(b) Proof that $M < N \implies \{y_n|_{n=1,2,\dots,M}\}$ is *not* a basis for L :

i. Suppose $M = N - 1$.

ii. Then $\{y_n|_{n=1,2,\dots,N-1}\}$ is a *basis* and there exists λ such that

$$\sum_{n=1}^N \lambda_{n \in \mathbb{Z}} y_{n \in \mathbb{Z}} = 0.$$

iii. This implies that $\{y_n|_{n=1,2,\dots,N}\}$ is *linearly dependent* and is *not* a basis.

iv. But this contradicts item (3), therefore $M \neq N - 1$.

v. Because $M = N$ yields a basis but $M = N - 1$ does not, $M < N - 1$ also does not yield a basis.

Corollary G.1. ⁹ Let $L \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}))$ be a linear space.

COR $\left\{ \begin{array}{l} 1. \quad \{x_n \in X | n=1,2,\dots,N\} \text{ is a HAMEL BASIS for } L \text{ and} \\ 2. \quad \{y_n \in X | n=1,2,\dots,M\} \text{ is a HAMEL BASIS for } L \end{array} \right\} \implies \{N = M\}$

(all Hamel bases for L have the same number of vectors)

PROOF: This follows from Theorem G.2 (page 199).

G.2 Bases in topological linear spaces

A linear space supports the concept of the *span* of a set of vectors (Definition G.2 page 197). In a topological linear space $\Omega \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), T)$, a set A is said to be *total* in Ω if the span of A is *dense* in Ω . In this case, A is said to be a *total set* or a *complete set*. However, this use of “complete” in a “complete set” is not equivalent to the use of “complete” in a “complete metric space”. ¹⁰ In this text, except for these comments and Definition G.6, “complete” refers to the metric space definition only.

If a set is both *total* and *linearly independent* (Definition G.3 page 197) in Ω , then that set is a *Hamel basis* (Definition G.4 page 199) for Ω .

Definition G.6. ¹¹ Let A^- be the CLOSURE of A in a TOPOLOGICAL LINEAR SPACE $\Omega \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), T)$. Let $\text{span}A$ be the SPAN (Definition G.2 page 197) of a set A .

DEF A set of vectors A is **total** (or **complete** or **fundamental**) in Ω if
 $(\text{span}A)^- = \Omega$ (SPAN of A is DENSE in Ω).

⁹ Kubrusly (2001) page 52 (Theorem 2.7), Michel and Herget (1993) page 91 (Theorem 3.3.31)

¹⁰ Haaser and Sullivan (1991) pages 296–297 (6.Orthogonal Bases), Rynne and Youngson (2008) page 78 (Remark 3.50), Heil (2011) page 21 (Remark 1.26)

¹¹ Young (2001) page 19 (Definition 1.5.1), Sohrab (2003) page 362 (Definition 9.2.3), Gupta (1998) page 134 (Definition 2.4), Bachman and Narici (1966) pages 149–153 (Definition 9.3, Theorems 9.9 and 9.10)

G.3 Schauder bases in Banach spaces

Definition G.7. ¹² Let $\mathcal{B} \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \|\cdot\|)$ be a BANACH SPACE. Let \doteq represent STRONG CONVERGENCE in \mathcal{B} .

The countable set $\{x_n \in X \mid n \in \mathbb{N}\}$ is a **Schauder basis** for \mathcal{B} if for each $x \in X$

1. $\exists (\alpha_n \in \mathbb{F})_{n \in \mathbb{N}}$ such that $x \doteq \sum_{n=1}^{\infty} \alpha_n x_n$ (STRONG CONVERGENCE in \mathcal{B}) and

2. $\left\{ \sum_{n=1}^{\infty} \alpha_n x_n \doteq \sum_{n=1}^{\infty} \beta_n x_n \right\} \Rightarrow \{(\alpha_n) = (\beta_n)\}$ (COEFFICIENT FUNCTIONALS are UNIQUE)

DEF

In this case, $\sum_{n=1}^{\infty} \alpha_n x_n$ is the **expansion** of x on $\{x_n \mid n \in \mathbb{N}\}$ and

the elements of (α_n) are the **coefficient functionals** associated with the basis $\{x_n\}$. Coefficient functionals are also called **coordinate functionals**.

In a Banach space, the existence of a Schauder basis implies that the space is *separable* (Theorem G.3 ¹³ page 202). The question of whether the converse is also true was posed by Banach himself in 1932, and became known as “*The basis problem*”. This remained an open question for many years. The question was finally answered some 41 years later in 1973 by Per Enflo (University of California at Berkley), with the answer being “no”. Enflo constructed a counterexample in which a separable Banach space does *not* have a Schauder basis. ¹⁴ Life is simpler in Hilbert spaces where the converse is true: a Hilbert space has a Schauder basis *if and only if* it is separable (Theorem G.11 page 214).

Theorem G.3. ¹⁵ Let $\mathcal{B} \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \|\cdot\|)$ be a BANACH SPACE. Let \mathbb{Q} be the field of rational numbers.

T H M $\left\{ \begin{array}{l} 1. \mathcal{B} \text{ has a SCHAUDER BASIS and} \\ 2. \mathbb{Q} \text{ is DENSE in } \mathbb{F}. \end{array} \right\} \Rightarrow \{ \mathcal{B} \text{ is SEPARABLE} \}$

PROOF:

1. lemma:

$$\begin{aligned} \left| \left\{ x \mid \exists (\alpha_n \in \mathbb{Q})_{n \in \mathbb{N}} \text{ such that } \lim_{N \rightarrow \infty} \left\| x - \sum_{n=1}^N \alpha_n x_n \right\| = 0 \right\} \right| &= |\mathbb{Q} \times \mathbb{N}| \\ &= |\mathbb{Z} \times \mathbb{Z}| \\ &= |\mathbb{Z}| \\ &= \text{countably infinite} \end{aligned}$$

¹² Carothers (2005) pages 24–25, Christensen (2003) pages 46–49 (Definition 3.1.1 and page 49), Young (2001) page 19 (Section 6), Singer (1970), page 17, Schauder (1927), Schauder (1928)

¹³ Banach (1932a), page 111

¹⁴ Enflo (1973), Lindenstrauss and Tzafriri (1977) pages 84–95 (Section 2.d)

¹⁵ Bachman et al. (2000) page 112 (3.4.8), Giles (2000) page 17, Heil (2011) page 21 (Theorem 1.27)

2. remainder of proof:

\mathcal{B} has a Schauder basis $(\mathbf{x}_n)_{n \in \mathbb{N}}$

\implies for every $\mathbf{x} \in \mathcal{B}$, there exists $(\alpha_n \in \mathbb{F})_{n \in \mathbb{N}}$ such that $\mathbf{x} \doteq \sum_{n=1}^{\infty} \alpha_n \mathbf{x}_n$ by Definition G.7 page 202

\implies for every $\mathbf{x} \in \mathcal{B}$, there exists $(\alpha_n \in \mathbb{F})_{n \in \mathbb{N}}$ such that $\lim_{N \rightarrow \infty} \left\| \mathbf{x} - \sum_{n=1}^N \alpha_n \mathbf{x}_n \right\| = 0$

\implies for every $\mathbf{x} \in \mathcal{B}$, there exists $(\alpha_n \in \mathbb{Q})_{n \in \mathbb{N}}$ such that $\lim_{N \rightarrow \infty} \left\| \mathbf{x} - \sum_{n=1}^N \alpha_n \mathbf{x}_n \right\| = 0$ because $\mathbb{Q}^- = \mathbb{F}$

$\implies \mathcal{B} = \left\{ \mathbf{x} \mid \exists (\alpha_n \in \mathbb{Q})_{n \in \mathbb{N}} \text{ such that } \lim_{N \rightarrow \infty} \left\| \mathbf{x} - \sum_{n=1}^N \alpha_n \mathbf{x}_n \right\| = 0 \right\}$

$\implies \mathcal{B} = \left\{ \mathbf{x} \mid \exists (\alpha_n \in \mathbb{Q})_{n \in \mathbb{N}} \text{ such that } \mathbf{x} = \lim_{N \rightarrow \infty} \sum_{n=1}^N \alpha_n \mathbf{x}_n \right\}$

$\implies \mathcal{B}$ is separable by (1) lemma page 202



Definition G.8. ¹⁶ Let $\{\mathbf{x}_n | n \in \mathbb{N}\}$ and $\{\mathbf{y}_n | n \in \mathbb{N}\}$ be SCHAUDER BASES of a BANACH SPACE $(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \|\cdot\|)$.

D E F

$\{\mathbf{x}_n\}$ is equivalent to $\{\mathbf{y}_n\}$

if there exists a BOUNDED INVERTIBLE operator \mathbf{R} in X^X such that $\mathbf{R}\mathbf{x}_n = \mathbf{y}_n \quad \forall n \in \mathbb{Z}$

Theorem G.4. ¹⁷ Let $\{\mathbf{x}_n | n \in \mathbb{N}\}$ and $\{\mathbf{y}_n | n \in \mathbb{N}\}$ be SCHAUDER BASES of a BANACH SPACE $(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \|\cdot\|)$.

T H M

$\{\{\mathbf{x}_n\} \text{ is EQUIVALENT to } \{\mathbf{y}_n\}\}$

$\iff \left\{ \sum_{n=1}^{\infty} \alpha_n \mathbf{x}_n \text{ is CONVERGENT} \iff \sum_{n=1}^{\infty} \alpha_n \mathbf{y}_n \text{ is CONVERGENT} \right\}$

Lemma G.1. ¹⁸ Let $\Omega \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \mathbf{T})$ be a topological linear space. Let $\text{span}A$ be the SPAN of a set A (Definition G.2 page 197). Let $\tilde{\mathbf{f}}(\omega)$ and $\tilde{\mathbf{g}}(\omega)$ be the FOURIER TRANSFORMS (Definition N.2 page 309) of the functions $\mathbf{f}(x)$ and $\mathbf{g}(x)$, respectively, in $L^2_{\mathbb{R}}$ (Definition ?? page ??). Let $\check{\mathbf{a}}(\omega)$ be the DTFT (Definition O.1 page 319) of a sequence $(a_n)_{n \in \mathbb{Z}}$ in $\ell^2_{\mathbb{R}}$ (Definition P.2 page 329).

L E M

$\left\{ \begin{array}{l} (1). \left\{ \mathbf{T}^n \mathbf{f} | n \in \mathbb{Z} \right\} \text{ is a SCHAUDER BASIS for } \Omega \text{ and} \\ (2). \left\{ \mathbf{T}^n \mathbf{g} | n \in \mathbb{Z} \right\} \text{ is a SCHAUDER BASIS for } \Omega \end{array} \right\} \implies \left\{ \begin{array}{l} \exists (a_n)_{n \in \mathbb{Z}} \text{ such that} \\ \tilde{\mathbf{f}}(\omega) = \check{\mathbf{a}}(\omega) \tilde{\mathbf{g}}(\omega) \end{array} \right\}$

PROOF: Let \mathbf{V}'_0 be the space spanned by $\{\mathbf{T}^n \phi | n \in \mathbb{Z}\}$.

$$\begin{aligned} \tilde{\mathbf{f}}(\omega) &\triangleq \tilde{\mathbf{F}}\mathbf{f} && \text{by definition of } \tilde{\mathbf{F}} && \text{(Definition N.2 page 309)} \\ &= \tilde{\mathbf{F}} \sum_{n \in \mathbb{Z}} a_n \mathbf{T}\mathbf{g} && \text{by (2)} \\ &= \sum_{n \in \mathbb{Z}} a_n \tilde{\mathbf{F}}\mathbf{g} \end{aligned}$$

¹⁶ Young (2001) page 25 (Definition 1.8.1, Theorem 1.8.7)

¹⁷ Young (2001) page 25 (Definition 1.8.1, Theorem 1.8.7)

¹⁸ Daubechies (1992), page 140

$$\begin{aligned}
 &= \underbrace{\sum_{n \in \mathbb{Z}} a_n e^{-i\omega n} \tilde{\mathbf{F}} \mathbf{g}}_{\check{a}(\omega)} \quad \text{by Corollary ?? page ??} \\
 &= \check{a}(\omega) \tilde{\mathbf{g}}(\omega) \quad \text{by definition of } \check{\mathbf{F}} \text{ and } \tilde{\mathbf{F}} \quad \text{by (Definition O.1 page 319, Definition N.2 page 309)}
 \end{aligned}$$

$$\begin{aligned}
 V_0 &\triangleq \left\{ f(x) | f(x) = \sum_{n \in \mathbb{Z}} b_n T^n g(x) \right\} \\
 &= \left\{ f(x) | \tilde{\mathbf{F}} f(x) = \tilde{\mathbf{F}} \sum_{n \in \mathbb{Z}} b_n T^n g(x) \right\} \\
 &= \left\{ f(x) | \tilde{f}(\omega) = \tilde{b}(\omega) \tilde{\mathbf{g}}(\omega) \right\} \\
 &= \left\{ f(x) | \tilde{f}(\omega) = \tilde{b}(\omega) \check{a}(\omega) \tilde{f}(\omega) \right\} \\
 &= \left\{ f(x) | \tilde{f}(\omega) = \tilde{c}(\omega) \tilde{f}(\omega) \right\} \quad \text{where } \tilde{c}(\omega) \triangleq \tilde{b}(\omega) \check{a}(\omega) \\
 &= \left\{ f(x) | f(x) = \sum_{n \in \mathbb{Z}} c_n f(x - n) \right\} \\
 &\triangleq V'_0
 \end{aligned}$$

→

G.4 Linear combinations in inner product spaces

In an *inner product space*, *orthogonality* is a special case of *linear independence*; or alternatively, linear independence is a generalization of orthogonality (next theorem).

Theorem G.5. ¹⁹ Let $\{x_n \in X | n=1,2,\dots,N\}$ be a set of vectors in an INNER PRODUCT SPACE (Definition I.1 page 233) $(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), (\triangle | \nabla))$.

T H M	$\left\{ \begin{array}{l} \{x_n\} \text{ is ORTHOGONAL} \\ (\text{Definition I.4 page 245}) \end{array} \right\} \implies \left\{ \begin{array}{l} \{x_n\} \text{ is LINEARLY INDEPENDENT} \\ (\text{Definition G.1 page 197}) \end{array} \right\}$
----------------------	--

PROOF:

1. Proof using *Pythagorean theorem* (Theorem I.10 page 246):

Let $(\alpha_n)_{n \in \mathbb{N}}$ be a sequence with at least one nonzero element.

$$\begin{aligned}
 \left\| \sum_{n=1}^N \alpha_n x_n \right\|^2 &= \sum_{n=1}^N \|\alpha_n x_n\|^2 \quad \text{by left hypoth. and Pythagorean Theorem (Theorem I.10 page 246)} \\
 &= \sum_{n=1}^N |\alpha_n|^2 \|x_n\|^2 \quad \text{by definition of } \|\cdot\| \quad (\text{Definition J.1 page 249}) \\
 &> 0 \\
 \implies \sum_{n=1}^N \alpha_n x_n &\neq 0 \\
 \implies (\alpha_n)_{n \in \mathbb{N}} \text{ is linearly independent} &\quad \text{by definition of linear independence} \quad (\text{Definition G.3 page 197})
 \end{aligned}$$

¹⁹  Aliprantis and Burkinshaw (1998) page 283 (Corollary 32.8),  Kubrusly (2001) page 352 (Proposition 5.34)

2. Alternative proof:

$$\begin{aligned}
 \sum_{n=1}^N \alpha_n \mathbf{x}_n = \mathbf{0} &\implies \left\langle \sum_{n=1}^N \alpha_n \mathbf{x}_n \mid \mathbf{x}_m \right\rangle = \langle \mathbf{0} \mid \mathbf{x}_m \rangle \\
 &\implies \sum_{n=1}^N \alpha_n \langle \mathbf{x}_n \mid \mathbf{x}_m \rangle = 0 \\
 &\implies \sum_{n=1}^N \alpha_n \bar{\delta}(k-m) = 0 \\
 &\implies \alpha_m = 0 \quad \text{for } m = 1, 2, \dots, N
 \end{aligned}$$

⇒

Theorem G.6 (Bessel's Equality). ²⁰ Let $\{\mathbf{x}_n \in X \mid n=1, 2, \dots, N\}$ be a set of vectors in an INNER PRODUCT SPACE (Definition I.1 page 233) $(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle \mid \triangledown \rangle)$ and with $\|\mathbf{x}\| \triangleq \sqrt{\langle \mathbf{x} \mid \mathbf{x} \rangle}$ (Definition I.2 page 238).

THM

$$\left\{ \begin{array}{l} \{\mathbf{x}_n\} \text{ is ORTHONORMAL} \\ (\text{Definition I.4 page 245}) \end{array} \right\} \implies \left\{ \underbrace{\left\| \mathbf{x} - \sum_{n=1}^N \langle \mathbf{x} \mid \mathbf{x}_n \rangle \mathbf{x}_n \right\|^2}_{\text{approximation error}} = \|\mathbf{x}\|^2 - \sum_{n=1}^N |\langle \mathbf{x} \mid \mathbf{x}_n \rangle|^2 \quad \forall \mathbf{x} \in X \right\}$$

PROOF:

$$\begin{aligned}
 & \left\| \mathbf{x} - \sum_{n=1}^N \langle \mathbf{x} \mid \mathbf{x}_n \rangle \mathbf{x}_n \right\|^2 \\
 &= \|\mathbf{x}\|^2 + \left\| \sum_{n=1}^N \langle \mathbf{x} \mid \mathbf{x}_n \rangle \mathbf{x}_n \right\|^2 - 2\Re \left\langle \mathbf{x} \mid \sum_{n=1}^N \langle \mathbf{x} \mid \mathbf{x}_n \rangle \mathbf{x}_n \right\rangle && \text{by polar identity} && (\text{Lemma I.1 page 237}) \\
 &= \|\mathbf{x}\|^2 + \left\| \sum_{n=1}^N \langle \mathbf{x} \mid \mathbf{x}_n \rangle \mathbf{x}_n \right\|^2 - 2\Re \left[\left(\sum_{n=1}^N \langle \mathbf{x} \mid \mathbf{x}_n \rangle \right)^* \langle \mathbf{x} \mid \mathbf{x}_n \rangle \right] && \text{by property of } \langle \triangle \mid \triangledown \rangle && (\text{Definition I.1 page 233}) \\
 &= \|\mathbf{x}\|^2 + \sum_{n=1}^N \|\langle \mathbf{x} \mid \mathbf{x}_n \rangle \mathbf{x}_n\|^2 - 2\Re \left[\left(\sum_{n=1}^N \langle \mathbf{x} \mid \mathbf{x}_n \rangle \right)^* \langle \mathbf{x} \mid \mathbf{x}_n \rangle \right] && \text{by Pythagorean Theorem} && (\text{Theorem I.10 page 246}) \\
 &= \|\mathbf{x}\|^2 + \sum_{n=1}^N \|\langle \mathbf{x} \mid \mathbf{x}_n \rangle \mathbf{x}_n\|^2 - 2\Re \left(\sum_{n=1}^N \langle \mathbf{x} \mid \mathbf{x}_n \rangle^* \langle \mathbf{x} \mid \mathbf{x}_n \rangle \right) \\
 &= \|\mathbf{x}\|^2 + \sum_{n=1}^N |\langle \mathbf{x} \mid \mathbf{x}_n \rangle|^2 \underbrace{\|\mathbf{x}_n\|^2}_1 - 2\Re \left(\sum_{n=1}^N \langle \mathbf{x} \mid \mathbf{x}_n \rangle^* \langle \mathbf{x} \mid \mathbf{x}_n \rangle \right) && \text{by property of } \|\cdot\| && (\text{Definition J.1 page 249}) \\
 &= \|\mathbf{x}\|^2 + \sum_{n=1}^N |\langle \mathbf{x} \mid \mathbf{x}_n \rangle|^2 \cdot 1 - 2\Re \left(\sum_{n=1}^N \langle \mathbf{x} \mid \mathbf{x}_n \rangle^* \langle \mathbf{x} \mid \mathbf{x}_n \rangle \right) && \text{by def. of orthonormality} && (\text{Definition I.4 page 245}) \\
 &= \|\mathbf{x}\|^2 + \sum_{n=1}^N |\langle \mathbf{x} \mid \mathbf{x}_n \rangle|^2 - 2\Re \sum_{n=1}^N |\langle \mathbf{x} \mid \mathbf{x}_n \rangle|^2 \\
 &= \|\mathbf{x}\|^2 + \sum_{n=1}^N |\langle \mathbf{x} \mid \mathbf{x}_n \rangle|^2 - 2 \sum_{n=1}^N |\langle \mathbf{x} \mid \mathbf{x}_n \rangle|^2 && \text{because } |\cdot| \text{ is real}
 \end{aligned}$$

²⁰ Bachman et al. (2000) page 103, Pedersen (2000) pages 38–39

$$= \|x\|^2 - \sum_{n=1}^N |\langle x | x_n \rangle|^2$$

⇒

Theorem G.7 (Bessel's inequality). ²¹ Let $\{x_n \in X \mid n=1,2,\dots,N\}$ be a set of vectors in an INNER PRODUCT SPACE (Definition I.1 page 233) $(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle | \triangleright \rangle)$ and with $\|x\| \triangleq \sqrt{\langle x | x \rangle}$ (Definition I.2 page 238).

T H M	$\left\{ \begin{array}{l} \{x_n\} \text{ is ORTHONORMAL} \\ (\text{Definition I.4 page 245}) \end{array} \right\} \Rightarrow \left\{ \begin{array}{l} \sum_{n=1}^N \langle x x_n \rangle ^2 \leq \ x\ ^2 \quad \forall x \in X \end{array} \right\}$
-------------	--

PROOF:

$$\begin{aligned} 0 &\leq \left\| x - \sum_{n=1}^N \langle x | x_n \rangle x_n \right\|^2 && \text{by definition of } \|\cdot\| && (\text{Definition J.1 page 249}) \\ &= \|x\|^2 - \sum_{n=1}^N |\langle x | x_n \rangle|^2 && \text{by Bessel's Equality} && (\text{Theorem G.6 page 205}) \end{aligned}$$

⇒

The *Best Approximation Theorem* (next) shows that

- ➊ the best sequence for representing a vector is the sequence of projections of the vector onto the sequence of basis functions
- ➋ the error of the projection is orthogonal to the projection.

Theorem G.8 (Best Approximation Theorem). ²² Let $\{x_n \in X \mid n=1,2,\dots,N\}$ be a set of vectors in an INNER PRODUCT SPACE (Definition I.1 page 233) $(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle | \triangleright \rangle)$ and with $\|x\| \triangleq \sqrt{\langle x | x \rangle}$ (Definition I.2 page 238).

T H M	$\left\{ \begin{array}{l} \{x_n\} \text{ is ORTHONORMAL} \\ (\text{Definition I.4 page 245}) \end{array} \right\} \Rightarrow \left\{ \begin{array}{l} 1. \arg \min_{(\alpha_n)_{n=1}^N} \left\ x - \sum_{n=1}^N \alpha_n x_n \right\ = \underbrace{(\langle x x_n \rangle)_{n=1}^N}_{\text{best } \alpha_n = \langle x x_n \rangle} \quad \forall x \in X \quad \text{and} \\ 2. \underbrace{\left(\sum_{n=1}^N \langle x x_n \rangle x_n \right)}_{\text{approximation}} \perp \underbrace{\left(x - \sum_{n=1}^N \langle x x_n \rangle x_n \right)}_{\text{approximation error}} \quad \forall x \in X \end{array} \right\}$
-------------	---

PROOF:

²¹ Giles (2000) pages 54–55 (3.13 Bessel's inequality), Bollobás (1999) page 147, Aliprantis and Burkinshaw (1998) page 284

²² Walter and Shen (2001), pages 3–4, Pedersen (2000), page 39, Edwards (1995), pages 94–100, Weyl (1940)

1. Proof that $(\langle \mathbf{x} | \mathbf{x}_n \rangle)$ is the best sequence:

$$\begin{aligned}
 & \left\| \mathbf{x} - \sum_{n=1}^N \alpha_n \mathbf{x}_n \right\|^2 \\
 &= \| \mathbf{x} \|^2 - 2 \Re \left\langle \mathbf{x} \mid \sum_{n=1}^N \alpha_n \mathbf{x}_n \right\rangle + \left\| \sum_{n=1}^N \alpha_n \mathbf{x}_n \right\|^2 \\
 &= \| \mathbf{x} \|^2 - 2 \Re \left(\sum_{n=1}^N \alpha_n^* \langle \mathbf{x} | \mathbf{x}_n \rangle \right) + \sum_{n=1}^N \| \alpha_n \mathbf{x}_n \|^2 \quad \text{by Pythagorean Theorem} \quad (\text{Theorem I.10 page 246}) \\
 &= \| \mathbf{x} \|^2 - 2 \Re \left(\sum_{n=1}^N \alpha_n^* \langle \mathbf{x} | \mathbf{x}_n \rangle \right) + \sum_{n=1}^N | \alpha_n |^2 + \underbrace{\left[\sum_{n=1}^N | \langle \mathbf{x} | \mathbf{x}_n \rangle |^2 - \sum_{n=1}^N | \langle \mathbf{x} | \mathbf{x}_n \rangle |^2 \right]}_0 \\
 &= \left[\| \mathbf{x} \|^2 - \sum_{n=1}^N | \langle \mathbf{x} | \mathbf{x}_n \rangle |^2 \right] + \sum_{n=1}^N \left[| \langle \mathbf{x} | \mathbf{x}_n \rangle |^2 - 2 \Re_e [\alpha_n^* \langle \mathbf{x} | \mathbf{x}_n \rangle] + | \alpha_n |^2 \right] \\
 &= \left[\| \mathbf{x} \|^2 - \sum_{n=1}^N | \langle \mathbf{x} | \mathbf{x}_n \rangle |^2 \right] + \sum_{n=1}^N [| \langle \mathbf{x} | \mathbf{x}_n \rangle |^2 - \alpha_n^* \langle \mathbf{x} | \mathbf{x}_n \rangle - \alpha_n \langle \mathbf{x} | \mathbf{x}_n \rangle^* + | \alpha_n |^2] \\
 &= \left\| \mathbf{x} - \sum_{n=1}^N \langle \mathbf{x} | \mathbf{x}_n \rangle \mathbf{x}_n \right\|^2 + \sum_{n=1}^N | \langle \mathbf{x} | \mathbf{x}_n \rangle - \alpha_n |^2 \quad \text{by Bessel's Equality} \quad (\text{Theorem G.6 page 205}) \\
 &\geq \left\| \mathbf{x} - \sum_{n=1}^N \langle \mathbf{x} | \mathbf{x}_n \rangle \mathbf{x}_n \right\|^2
 \end{aligned}$$

2. Proof that the approximation and approximation error are orthogonal:

$$\begin{aligned}
 \left\langle \sum_{n=1}^N \langle \mathbf{x} | \mathbf{x}_n \rangle \mathbf{x}_n \mid \mathbf{x} - \sum_{n=1}^N \langle \mathbf{x} | \mathbf{x}_n \rangle \mathbf{x}_n \right\rangle &= \left\langle \sum_{n=1}^N \langle \mathbf{x} | \mathbf{x}_n \rangle \mathbf{x}_n \mid \mathbf{x} \right\rangle - \left\langle \sum_{n=1}^N \langle \mathbf{x} | \mathbf{x}_n \rangle \mathbf{x}_n \mid \sum_{n=1}^N \langle \mathbf{x} | \mathbf{x}_n \rangle \mathbf{x}_n \right\rangle \\
 &= \sum_{n=1}^N \langle \mathbf{x} | \mathbf{x}_n \rangle^* \langle \mathbf{x} | \mathbf{x}_n \rangle - \sum_{n=1}^N \sum_{m=1}^N \langle \mathbf{x} | \mathbf{x}_n \rangle \langle \mathbf{x} | \mathbf{x}_m \rangle^* \langle \mathbf{x}_n | \mathbf{x}_m \rangle \\
 &= \sum_{n=1}^N | \langle \mathbf{x} | \mathbf{x}_n \rangle |^2 - \sum_{n=1}^N \sum_{m=1}^N \langle \mathbf{x} | \mathbf{x}_n \rangle \langle \mathbf{x} | \mathbf{x}_m \rangle^* \bar{\delta}_{nm} \\
 &= \sum_{n=1}^N | \langle \mathbf{x} | \mathbf{x}_n \rangle |^2 - \sum_{n=1}^N | \langle \mathbf{x} | \mathbf{x}_n \rangle |^2 \\
 &= 0
 \end{aligned}$$



G.5 Orthonormal bases in Hilbert spaces

Definition G.9. Let $\{ \mathbf{x}_n \in X \mid n=1,2,\dots,N \}$ be a set of vectors in an INNER PRODUCT SPACE (Definition I.1 page 233) $\mathcal{Q} \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), (\Delta \mid \nabla))$.

D E F The set $\{x_n\}$ is an **orthogonal basis** for Ω if $\{x_n\}$ is ORTHOGONAL and is

a SCHAUDER BASIS for Ω .

The set $\{x_n\}$ is an **orthonormal basis** for Ω if $\{x_n\}$ is ORTHONORMAL and is a SCHAUDER BASIS for Ω .

Definition G.10. ²³ Let $H \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle | \nabla \rangle)$ be a Hilbert space.

D E F Suppose there exists a set $\{x_n \in X \mid n \in \mathbb{N}\}$ such that $x \doteq \sum_{n=1}^{\infty} \langle x | x_n \rangle x_n$.

Then the quantities $\langle x | x_n \rangle$ are called the **Fourier coefficients** of x and the sum

$\sum_{n=1}^{\infty} \langle x | x_n \rangle x_n$ is called the **Fourier expansion** of x or the **Fourier series** for x .

Lemma G.2 (Perfect reconstruction). Let $\{x_n \in X \mid n \in \mathbb{N}\}$ be a set of vectors in a HILBERT SPACE $H \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle | \nabla \rangle)$.

L E M $\left\{ \begin{array}{l} (1). \quad \{x_n\} \text{ is a BASIS for } H \\ (2). \quad \{x_n\} \text{ is ORTHONORMAL} \end{array} \right. \text{ and } \Rightarrow x \doteq \underbrace{\sum_{n=1}^{\infty} \underbrace{\langle x | x_n \rangle}_{\text{Fourier coefficient}} x_n}_{\text{Fourier expansion}} \quad \forall x \in X$

PROOF:

$$\begin{aligned} \langle x | x_n \rangle &= \left\langle \sum_{m \in \mathbb{Z}} \alpha_m x_m | x_n \right\rangle && \text{by left hypothesis (1)} \\ &= \sum_{m \in \mathbb{Z}} \alpha_m \langle x_m | x_n \rangle && \text{by homogeneous property of } \langle \triangle | \nabla \rangle \quad (\text{Definition I.1 page 233}) \\ &= \sum_{m \in \mathbb{Z}} \alpha_m \delta_{n-m} && \text{by left hypothesis (2)} \quad (\text{Definition I.4 page 245}) \\ &= \alpha_n \end{aligned}$$

Proposition G.2. ²⁴ Let $\{x_n \in X \mid n \in \mathbb{N}\}$ be a set of vectors in a HILBERT SPACE $H \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle | \nabla \rangle)$.

P R P $\|x\|^2 \doteq \underbrace{\sum_{n=1}^{\infty} |\langle x | x_n \rangle|^2}_{\text{PARSEVAL FRAME}} \iff x \doteq \underbrace{\sum_{n=1}^{\infty} \langle x | x_n \rangle x_n}_{\text{FOURIER EXPANSION (Definition G.10 page 208)}} \quad \forall x \in X$

PROOF:

²³ Fabian et al. (2010) page 27 (Theorem 1.55), Young (2001) page 6, Young (1980) page 6

²⁴ Han et al. (2007) pages 93–94 (Proposition 3.11)

1. Proof that *Parseval frame* \iff *Fourier expansion*

$$\begin{aligned}
 \|x\|^2 &\triangleq \langle x | x \rangle && \text{by definition of } \|\cdot\| && (\text{Definition J.1 page 249}) \\
 &= \left\langle \sum_{n=1}^{\infty} \langle x | x_n \rangle x | x_n \right\rangle && \text{by right hypothesis} \\
 &\stackrel{*}{=} \sum_{n=1}^{\infty} \langle x | x_n \rangle \langle x | x_n \rangle && \text{by property of } \langle \triangle | \triangledown \rangle && (\text{Definition I.1 page 233}) \\
 &\stackrel{*}{=} \sum_{n=1}^{\infty} \langle x | x_n \rangle \langle x | x_n \rangle^* && \text{by property of } \langle \triangle | \triangledown \rangle && (\text{Definition I.1 page 233}) \\
 &\stackrel{*}{=} \sum_{n=1}^{\infty} |\langle x | x_n \rangle|^2 && \text{by property of } \mathbb{C} && (\text{Definition H.7 page 231})
 \end{aligned}$$

2. Proof that *Parseval frame* \implies *Fourier expansion*

(a) Let $(e_n)_{n \in \mathbb{N}}$ be the *standard orthonormal basis* such that the n th element of e_n is 1 and all other elements are 0.

(b) Let M be an operator in H such that $Mx \triangleq \sum_{n=1}^{\infty} \langle x | x_n \rangle e_n$.

(c) lemma: M is *isometric*. Proof:

$$\begin{aligned}
 \|Mx\|^2 &= \left\| \sum_{n=1}^{\infty} \langle x | x_n \rangle e_n \right\|^2 && \text{by definition of } M && (\text{item (2b) page 209}) \\
 &= \sum_{n=1}^{\infty} \|\langle x | x_n \rangle e_n\|^2 && \text{by Pythagorean Theorem} && (\text{Theorem I.10 page 246}) \\
 &= \sum_{n=1}^{\infty} |\langle x | x_n \rangle|^2 \|e_n\|^2 && \text{by homogeneous property of } \|\cdot\| && (\text{Definition J.1 page 249}) \\
 &= \sum_{n=1}^{\infty} |\langle x | x_n \rangle|^2 && \text{by definition of orthonormal} && (\text{Definition I.4 page 245}) \\
 &= \|x\|^2 && \text{by Parseval frame hypothesis} \\
 \implies M &\text{ is isometric} && \text{by definition of isometric} && (\text{Definition M.10 page 300})
 \end{aligned}$$

(d) Let $(u_n)_{n \in \mathbb{N}}$ be an *orthonormal basis* for H .

(e) Proof for *Fourier expansion*:

$$\begin{aligned}
 x &= \sum_{n=1}^{\infty} \langle x | u_n \rangle u_n && \text{by Fourier expansion (Proposition G.3 page 212)} \\
 &= \sum_{n=1}^{\infty} \langle Mx | Mu_n \rangle u_n && \text{by (2c) lemma page 209 and Theorem M.21 page 301} \\
 &= \sum_{n=1}^{\infty} \left\langle \sum_{m=1}^{\infty} \langle x | x_m \rangle e_m | \sum_{k=1}^{\infty} \langle u_n | x_k \rangle e_k \right\rangle u_n && \text{by item (2b) page 209} \\
 &= \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \langle x | x_m \rangle \sum_{k=1}^{\infty} \langle u_n | x_k \rangle^* \langle e_m | e_k \rangle u_n && \text{by Definition I.1 page 233} \\
 &= \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \langle x | x_m \rangle \langle u_n | x_m \rangle^* u_n && \text{by item (2a) page 209 and Definition I.4 page 245}
 \end{aligned}$$

$$\begin{aligned}
 &= \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \langle \mathbf{x} | \mathbf{x}_m \rangle \langle \mathbf{x}_m | \mathbf{u}_n \rangle \mathbf{u}_n \\
 &= \sum_{m=1}^{\infty} \langle \mathbf{x} | \mathbf{x}_m \rangle \sum_{n=1}^{\infty} \langle \mathbf{x}_m | \mathbf{u}_n \rangle \mathbf{u}_n \\
 &= \sum_{m=1}^{\infty} \langle \mathbf{x} | \mathbf{x}_m \rangle \mathbf{x}_m
 \end{aligned}
 \quad \begin{array}{l} \text{by Definition I.1 page 233} \\ \text{by item (2d) page 209} \end{array}$$

☞

When is a set of orthonormal vectors in a Hilbert space \mathbf{H} *total*? Theorem G.9 (next) offers some help.

Theorem G.9 (The Fourier Series Theorem). ²⁵ Let $\{\mathbf{x}_n \in X\}_{n \in \mathbb{N}}$ be a set of vectors in a HILBERT SPACE $\mathbf{H} \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \Delta | \nabla \rangle)$ and let $\|\mathbf{x}\| \triangleq \sqrt{\langle \mathbf{x} | \mathbf{x} \rangle}$ (Definition I.2 page 238).

THM	$(A) \{\mathbf{x}_n\} \text{ is ORTHONORMAL in } \mathbf{H} \implies$ $\left\{ \begin{array}{ll} (1). \quad (\text{span}\{\mathbf{x}_n\})^\perp = \mathbf{H} & \{\{\mathbf{x}_n\} \text{ is TOTAL in } \mathbf{H}\} \\ \Leftrightarrow (2). \quad \langle \mathbf{x} \mathbf{y} \rangle \triangleq \sum_{n=1}^{\infty} \langle \mathbf{x} \mathbf{x}_n \rangle \langle \mathbf{y} \mathbf{x}_n \rangle^* \quad \forall \mathbf{x}, \mathbf{y} \in X & \text{(GENERALIZED PARSEVAL'S IDENTITY)} \\ \Leftrightarrow (3). \quad \ \mathbf{x}\ ^2 \triangleq \sum_{n=1}^{\infty} \langle \mathbf{x} \mathbf{x}_n \rangle ^2 \quad \forall \mathbf{x} \in X & \text{(PARSEVAL'S IDENTITY)} \\ \Leftrightarrow (4). \quad \mathbf{x} \triangleq \sum_{n=1}^{\infty} \langle \mathbf{x} \mathbf{x}_n \rangle \mathbf{x}_n \quad \forall \mathbf{x} \in X & \text{(FOURIER SERIES EXPANSION)} \end{array} \right. \end{array} \right\}$
-----	--

⇒ PROOF:

1. Proof that (1) \implies (2):

$$\begin{aligned}
 \langle \mathbf{x} | \mathbf{y} \rangle &\triangleq \left\langle \sum_{n=1}^{\infty} \langle \mathbf{x} | \mathbf{x}_n \rangle \mathbf{x}_n \mid \sum_{m=1}^{\infty} \langle \mathbf{y} | \mathbf{x}_m \rangle \mathbf{x}_m \right\rangle \quad \text{by (A) and (1)} \\
 &\triangleq \sum_{n=1}^{\infty} \langle \mathbf{x} | \mathbf{x}_n \rangle \left\langle \mathbf{x}_n \mid \sum_{m=1}^{\infty} \langle \mathbf{y} | \mathbf{x}_m \rangle \mathbf{x}_m \right\rangle \quad \text{by property of } \langle \cdot | \cdot \rangle \quad \text{(Definition I.1 page 233)} \\
 &\triangleq \sum_{n=1}^{\infty} \langle \mathbf{x} | \mathbf{x}_n \rangle \sum_{m=1}^{\infty} \langle \mathbf{y} | \mathbf{x}_m \rangle^* \langle \mathbf{x}_n | \mathbf{x}_m \rangle \quad \text{by property of } \langle \cdot | \cdot \rangle \quad \text{(Definition I.1 page 233)} \\
 &\triangleq \sum_{n=1}^{\infty} \langle \mathbf{x} | \mathbf{x}_n \rangle \sum_{m=1}^{\infty} \langle \mathbf{y} | \mathbf{x}_m \rangle^* \bar{\delta}_{mn} \quad \text{by (A)} \\
 &\triangleq \sum_{n=1}^{\infty} \langle \mathbf{x} | \mathbf{x}_n \rangle \langle \mathbf{y} | \mathbf{x}_n \rangle^* \quad \text{by definition of } \bar{\delta}_n \quad \text{(Definition I.3 page 245)}
 \end{aligned}$$

²⁵ Bachman and Narici (1966) pages 149–155 (Theorem 9.12), Kubrusly (2001) pages 360–363 (Theorem 5.48), Aliprantis and Burkinshaw (1998) pages 298–299 (Theorem 34.2), Christensen (2003) page 57 (Theorem 3.4.2), Berberian (1961) pages 52–53 (Theorem II§8.3), Heil (2011) pages 34–35 (Theorem 1.50), Bracewell (1978) page 112 (Rayleigh's theorem)

2. Proof that (2) \implies (3):

$$\begin{aligned} \|\mathbf{x}\|^2 &\triangleq \langle \mathbf{x} | \mathbf{x} \rangle && \text{by definition of } \textit{induced norm} && (\text{Theorem I.4 page 238}) \\ &= \sum_{n=1}^{\infty} \langle \mathbf{x} | \mathbf{x}_n \rangle \langle \mathbf{x} | \mathbf{x}_n \rangle^* && \text{by (2)} \\ &= \sum_{n=1}^{\infty} |\langle \mathbf{x} | \mathbf{x}_n \rangle|^2 \end{aligned}$$

3. Proof that (3) \iff (4) *not* using (A): by Proposition G.2 page 208

4. Proof that (3) \implies (1) (proof by contradiction):

(a) Suppose $\{\mathbf{x}_n\}$ is *not total*.

(b) Then there must exist a vector \mathbf{y} in H such that the set $B \triangleq \{\mathbf{x}_n\} \cup \mathbf{y}$ is *orthonormal*.

(c) Then $1 = \|\mathbf{y}\|^2 \neq \sum_{n=1}^{\infty} |\langle \mathbf{y} | \mathbf{x}_n \rangle|^2 = 0$.

(d) But this contradicts (3), and so $\{\mathbf{x}_n\}$ must be *total* and (3) \implies (1).

5. Extraneous proof that (3) \implies (4) (this proof is not really necessary here):

$$\begin{aligned} \left\| \mathbf{x} - \sum_{n=1}^{\infty} \langle \mathbf{x} | \mathbf{x}_n \rangle \mathbf{x}_n \right\|^2 &= \|\mathbf{x}\|^2 - \sum_{n=1}^{\infty} |\langle \mathbf{x} | \mathbf{x}_n \rangle|^2 && \text{by Bessel's Equality} && (\text{Theorem G.6 page 205}) \\ &= 0 && \text{by (3)} \\ \implies \mathbf{x} &\stackrel{*}{=} \sum_{n=1}^{\infty} \langle \mathbf{x} | \mathbf{x}_n \rangle \mathbf{x}_n && \text{by definition of } \stackrel{*}{=} \end{aligned}$$

6. Extraneous proof that (A) \implies (4) (this proof is not really necessary here)

(a) The sequence $\sum_{n=1}^N |\langle \mathbf{x} | \mathbf{x}_n \rangle|^2$ is *monotonically increasing* in n .

(b) By Bessel's inequality (page 206), the sequence is upper bounded by $\|\mathbf{x}\|^2$:

$$\sum_{n=1}^N |\langle \mathbf{x} | \mathbf{x}_n \rangle|^2 \leq \|\mathbf{x}\|^2$$

(c) Because this sequence is both monotonically increasing and bounded in n , it must equal its bound in the limit as n approaches infinity:

$$\lim_{N \rightarrow \infty} \sum_{n=1}^N |\langle \mathbf{x} | \mathbf{x}_n \rangle|^2 = \|\mathbf{x}\|^2 \tag{G.1}$$

(d) If we combine this result with *Bessel's Equality* (Theorem G.6 page 205) we have

$$\begin{aligned} \lim_{N \rightarrow \infty} \left\| \mathbf{x} - \sum_{n=1}^N \langle \mathbf{x} | \mathbf{x}_n \rangle \mathbf{x}_n \right\|^2 &= \|\mathbf{x}\|^2 - \lim_{N \rightarrow \infty} \sum_{n=1}^N |\langle \mathbf{x} | \mathbf{x}_n \rangle|^2 && \text{by Bessel's Equality (Theorem G.6 page 205)} \\ &= \|\mathbf{x}\|^2 - \|\mathbf{x}\|^2 && \text{by equation (G.1) page 211} \\ &= 0 \end{aligned}$$



Proposition G.3 (Fourier expansion). Let $\{x_n \in X | n \in \mathbb{N}\}$ be a set of vectors in a HILBERT SPACE $(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), (\triangle | \nabla))$.

P R P	$\underbrace{\{x_n\} \text{ is an ORTHONORMAL BASIS for } H}_{(A)} \quad \Rightarrow \quad \underbrace{\left\{ x \doteq \sum_{n=1}^{\infty} \alpha_n x_n \quad \Leftrightarrow \quad \underbrace{\alpha_n = \langle x x_n \rangle}_{(2)} \right\}}_{(1)}$
-------------	---

PROOF:

1. Proof that (1) \Rightarrow (2): by Lemma G.2 page 208

2. Proof that (1) \Leftarrow (2):

$$\begin{aligned}
 \left\| x - \sum_{n=1}^{\infty} \alpha_n x_n \right\|^2 &= \left\| x - \sum_{n=1}^{\infty} \langle x | x_n \rangle x_n \right\|^2 && \text{by right hypothesis} \\
 &= \|x\|^2 - \sum_{n=1}^{\infty} |\langle x | x_n \rangle|^2 && \text{by Bessel's equality} \quad (\text{Theorem G.6 page 205}) \\
 &= 0 && \text{by Parseval's Identity} \quad (\text{Theorem G.9 page 210}) \\
 \stackrel{\text{def}}{\Leftrightarrow} \quad x &\doteq \sum_{n=1}^{\infty} \langle x | x_n \rangle x_n && \text{by definition of strong convergence}
 \end{aligned}$$

⇒

Proposition G.4 (Riesz-Fischer Theorem). ²⁶ Let $\{x_n \in X | n \in \mathbb{N}\}$ be a set of vectors in a HILBERT SPACE $(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), (\triangle | \nabla))$.

P R P	$\underbrace{\{x_n\} \text{ is an ORTHONORMAL BASIS for } H}_{(A)} \quad \Rightarrow \quad \left\{ \underbrace{\sum_{n=1}^{\infty} \alpha_n ^2 < \infty}_{(1)} \quad \Leftrightarrow \quad \underbrace{\exists x \in H \text{ such that } \alpha_n = \langle x x_n \rangle}_{(2)} \right\}$
-------------	--

PROOF:

1. Proof that (1) \Rightarrow (2):

(a) If (1) is true, then let $x \doteq \sum_{n \in \mathbb{N}} \alpha_n x_n$.

(b) Then

$$\begin{aligned}
 \langle x | x_n \rangle &= \left\langle \sum_{m \in \mathbb{N}} \alpha_m x_m | x_n \right\rangle && \text{by definition of } x \\
 &= \sum_{m \in \mathbb{N}} \alpha_m \langle x_m | x_n \rangle && \text{by homogeneous property of } (\triangle | \nabla) \quad (\text{Definition I.1 page 233}) \\
 &= \sum_{m \in \mathbb{N}} \alpha_m \bar{\delta}_{mn} && \text{by (A)} \\
 &= \sum_{m \in \mathbb{N}} \alpha_m && \text{by definition of } \bar{\delta} \quad (\text{Definition I.3 page 245})
 \end{aligned}$$

²⁶ Young (2001) page 6

2. Proof that (1) \iff (2):

$$\begin{aligned} \sum_{n \in \mathbb{N}} |\alpha_n|^2 &= \sum_{n \in \mathbb{N}} |\langle x | x_n \rangle|^2 && \text{by (2)} \\ &\leq \|x\|^2 && \text{by Bessel's Inequality} && \text{(Theorem G.7 page 206)} \\ &\leq \infty \end{aligned}$$



Theorem G.10. ²⁷

**T
H
M**

All SEPARABLE HILBERT SPACES are ISOMORPHIC. That is,

$$\left\{ \begin{array}{l} \mathbf{X} \text{ is a separable} \\ \text{Hilbert space} \\ \mathbf{Y} \text{ is a separable} \\ \text{Hilbert space} \end{array} \right. \text{ and } \Rightarrow \left\{ \begin{array}{l} \text{there is a BIJECTIVE operator } \mathbf{M} \in \mathbf{Y}^{\mathbf{X}} \text{ such that} \\ (1). \quad \mathbf{y} = \mathbf{M}\mathbf{x} \quad \forall \mathbf{x} \in \mathbf{X}, \mathbf{y} \in \mathbf{Y} \quad \text{and} \\ (2). \quad \|\mathbf{M}\mathbf{x}\| = \|\mathbf{x}\| \quad \forall \mathbf{x} \in \mathbf{X} \quad \text{and} \\ (3). \quad \langle \mathbf{M}\mathbf{x} | \mathbf{M}\mathbf{y} \rangle = \langle \mathbf{x} | \mathbf{y} \rangle \quad \forall \mathbf{x} \in \mathbf{X}, \mathbf{y} \in \mathbf{Y} \end{array} \right\}$$

PROOF:

1. Let $\mathbf{X} \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \Delta | \nabla \rangle)$ be a *separable Hilbert space* with *orthonormal basis* $\{x_n\}_{n \in \mathbb{N}}$. Let $\mathbf{Y} \triangleq (Y, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \Delta | \nabla \rangle)$ be a *separable Hilbert space* with *orthonormal basis* $\{y_n\}_{n \in \mathbb{N}}$.
2. Proof that there exists *bijection* operator \mathbf{M} and its inverse \mathbf{M}^{-1} between $\{x_n\}$ and $\{y_n\}$:
 - (a) Let \mathbf{M} be defined such that $y_n \triangleq \mathbf{M}x_n$.
 - (b) Thus \mathbf{M} is a *bijection* between $\{x_n\}$ and $\{y_n\}$.
 - (c) Because \mathbf{M} is a *bijection* between $\{x_n\}$ and $\{y_n\}$, \mathbf{M} has an inverse operator \mathbf{M}^{-1} between $\{x_n\}$ and $\{y_n\}$ such that $x_n = \mathbf{M}^{-1}y_n$.
3. Proof that \mathbf{M} and \mathbf{M}^{-1} are *bijection* operators between \mathbf{X} and \mathbf{Y} :
 - (a) Proof that \mathbf{M} maps \mathbf{X} into \mathbf{Y} :

(a) Proof that \mathbf{M} maps \mathbf{X} into \mathbf{Y} :

$$\begin{aligned} \mathbf{x} \in \mathbf{X} &\iff \mathbf{x} \triangleq \sum_{n \in \mathbb{N}} \langle \mathbf{x} | \mathbf{x}_n \rangle \mathbf{x}_n && \text{by Fourier expansion} && \text{(Theorem G.9 page 210)} \\ &\implies \exists \mathbf{y} \in \mathbf{Y} \text{ such that } \langle \mathbf{y} | \mathbf{y}_n \rangle = \langle \mathbf{x} | \mathbf{x}_n \rangle && \text{by Riesz-Fischer Thm.} && \text{(Proposition G.4 page 212)} \\ &\implies \\ \mathbf{y} &= \sum_{n \in \mathbb{N}} \langle \mathbf{y} | \mathbf{y}_n \rangle \mathbf{y}_n && \text{by Fourier expansion} && \text{(Theorem G.9 page 210)} \\ &= \sum_{n \in \mathbb{N}} \langle \mathbf{x} | \mathbf{x}_n \rangle \mathbf{y}_n && \text{by Riesz-Fischer Thm.} && \text{(Proposition G.4 page 212)} \\ &= \sum_{n \in \mathbb{N}} \langle \mathbf{x} | \mathbf{x}_n \rangle \mathbf{M}\mathbf{x}_n && \text{by definition of } \mathbf{M} && \text{(item (2a) page 213)} \\ &= \mathbf{M} \sum_{n \in \mathbb{N}} \langle \mathbf{x} | \mathbf{x}_n \rangle \mathbf{x}_n && \text{by prop. of linear ops.} && \text{(Theorem M.1 page 282)} \\ &= \mathbf{M}\mathbf{x} && \text{by definition of } \mathbf{x} \end{aligned}$$

²⁷ Young (2001) page 6

(b) Proof that \mathbf{M}^{-1} maps \mathbf{Y} into \mathbf{X} :

$$\begin{aligned}
 y \in \mathbf{Y} &\iff y \doteq \sum_{n \in \mathbb{N}} \langle y | y_n \rangle y_n && \text{by Fourier expansion (Theorem G.9 page 210)} \\
 &\implies \exists x \in \mathbf{X} \text{ such that } \langle x | x_n \rangle = \langle y | y_n \rangle \text{ by Riesz-Fischer Thm.} && \text{(Proposition G.4 page 212)} \\
 &\implies \\
 x &= \sum_{n \in \mathbb{N}} \langle x | x_n \rangle x_n && \text{by Fourier expansion (Theorem G.9 page 210)} \\
 &= \sum_{n \in \mathbb{N}} \langle y | y_n \rangle x_n && \text{by Riesz-Fischer Thm. (Proposition G.4 page 212)} \\
 &= \sum_{n \in \mathbb{N}} \langle y | y_n \rangle \mathbf{M}^{-1} y_n && \text{by definition of } \mathbf{M}^{-1} \text{ (item (2c) page 213)} \\
 &= \mathbf{M}^{-1} \sum_{n \in \mathbb{N}} \langle y | y_n \rangle y_n && \text{by prop. of linear ops. (Theorem M.1 page 282)} \\
 &= \mathbf{M}^{-1} y && \text{by definition of } y
 \end{aligned}$$

4. Proof for (2):

$$\begin{aligned}
 \|\mathbf{M}x\|^2 &= \left\| \mathbf{M} \sum_{n \in \mathbb{N}} \langle x | x_n \rangle x_n \right\|^2 && \text{by Fourier expansion (Theorem G.9 page 210)} \\
 &= \left\| \sum_{n \in \mathbb{N}} \langle x | x_n \rangle \mathbf{M}x_n \right\|^2 && \text{by property of linear operators (Theorem M.1 page 282)} \\
 &= \left\| \sum_{n \in \mathbb{N}} \langle x | x_n \rangle y_n \right\|^2 && \text{by definition of } \mathbf{M} \text{ (item (2a) page 213)} \\
 &= \sum_{n \in \mathbb{N}} |\langle x | x_n \rangle|^2 && \text{by Parseval's Identity (Proposition G.4 page 212)} \\
 &= \left\| \sum_{n \in \mathbb{N}} \langle x | x_n \rangle x_n \right\|^2 && \text{by Parseval's Identity (Proposition G.4 page 212)} \\
 &= \|x\|^2 && \text{by Fourier expansion (Theorem G.9 page 210)}
 \end{aligned}$$

5. Proof for (3): by (2) and Theorem M.21 page 301



Theorem G.11. ²⁸ Let \mathbf{H} be a HILBERT SPACE.

T	H	M	\mathbf{H} has a SCHAUDER BASIS	\iff	\mathbf{H} is SEPARABLE
---	---	---	-----------------------------------	--------	---------------------------

Theorem G.12. ²⁹ Let \mathbf{H} be a HILBERT SPACE.

T	H	M	\mathbf{H} has an ORTHONORMAL BASIS	\iff	\mathbf{H} is SEPARABLE
---	---	---	---------------------------------------	--------	---------------------------

²⁸ Bachman et al. (2000) page 112 (3.4.8), Berberian (1961) page 53 (Theorem II§8.3)

²⁹ Kubrusly (2001) page 357 (Proposition 5.43)

G.6 Riesz bases in Hilbert spaces

Definition G.11. ³⁰ Let $\{x_n \in X | n \in \mathbb{N}\}$ be a set of vectors in a SEPARABLE HILBERT SPACE $H \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle | \nabla \rangle)$

DEF $\{x_n\}$ is a **Riesz basis** for H if $\{x_n\}$ is EQUIVALENT (Definition G.8 page 203) to some ORTHONORMAL BASIS (Definition G.9 page 207) in H .

Definition G.12. ³¹ Let $(x_n \in X)_{n \in \mathbb{N}}$ be a sequence of vectors in a SEPARABLE HILBERT SPACE $H \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle | \nabla \rangle)$.

The sequence (x_n) is a **Riesz sequence** for H if

DEF $\exists A, B \in \mathbb{R}^+$ such that $A \sum_{n=1}^{\infty} |\alpha_n|^2 \leq \left\| \sum_{n=1}^{\infty} \alpha_n x_n \right\|^2 \leq B \sum_{n=1}^{\infty} |\alpha_n|^2 \quad \forall (\alpha_n) \in \ell_{\mathbb{F}}^2$.

Definition G.13. Let $X \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle | \nabla \rangle)$ be an INNER PRODUCT SPACE (Definition I.1 page 233).

DEF The sequences $(x_n \in X)_{n \in \mathbb{Z}}$ and $(y_n \in X)_{n \in \mathbb{Z}}$ are **biorthogonal** with respect to each other in X if $\langle x_n | y_m \rangle = \delta_{nm}$

Lemma G.3. ³² Let $\{x_n | n \in \mathbb{N}\}$ be a sequence in a HILBERT SPACE $X \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle | \nabla \rangle)$. Let $\{y_n | n \in \mathbb{N}\}$ be a sequence in a HILBERT SPACE $Y \triangleq (Y, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle | \nabla \rangle)$. Let

LEM $\left\{ \begin{array}{l} (i). \quad \{x_n\} \text{ is TOTAL in } X \\ (ii). \quad \text{There exists } A > 0 \text{ such that } A \sum_{n \in C} |\alpha_n|^2 \leq \left\| \sum_{n \in C} \alpha_n x_n \right\|^2 \text{ for finite } C \\ (iii). \quad \text{There exists } B > 0 \text{ such that } \left\| \sum_{n=1}^{\infty} b_n y_n \right\|^2 \leq B \sum_{n=1}^{\infty} |b_n|^2 \quad \forall (b_n)_{n \in \mathbb{N}} \in \ell_{\mathbb{F}}^2 \end{array} \right\} \Rightarrow \left\{ \begin{array}{l} (1). \quad \mathbf{R}^\circ \text{ is a linear bounded operator that maps from } \text{span}\{x_n\} \text{ to } \text{span}\{y_n\} \\ \text{where } \mathbf{R}^\circ \sum_{n \in C} c_n x_n \triangleq \sum_{n \in C} c_n y_n, \text{ for some sequence } (c_n) \text{ and finite set } C \\ (2). \quad \mathbf{R} \text{ has a unique extension to a bounded operator } \mathbf{R} \text{ that maps from } X \text{ to } Y \\ (3). \quad \|\mathbf{R}^\circ\| \leq \frac{B}{A} \\ (4). \quad \|\mathbf{R}\| \leq \frac{B}{A} \end{array} \right\}$

Theorem G.13. ³³ Let $\{x_n \in X | n \in \mathbb{N}\}$ be a set of vectors in a SEPARABLE HILBERT SPACE $H \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle | \nabla \rangle)$.

THM $\left\{ \begin{array}{l} \{x_n\} \text{ is a RIESZ BASIS} \\ \text{for } H \end{array} \right\} \iff \left\{ \begin{array}{l} (1). \quad \{x_n\} \text{ is TOTAL in } H \\ (2). \quad \exists A, B \in \mathbb{R}^+ \text{ such that } A \sum_{n=1}^{\infty} |\alpha_n|^2 \leq \left\| \sum_{n=1}^{\infty} \alpha_n x_n \right\|^2 \leq B \sum_{n=1}^{\infty} |\alpha_n|^2 \end{array} \right\}$

PROOF:

³⁰ Young (2001) page 27 (Definition 1.8.2), Christensen (2003) page 63 (Definition 3.6.1), Heil (2011) page 196 (Definition 7.9)

³¹ Christensen (2003) pages 66–68 (page 68 and (3.24) on page 66), Wojtaszczyk (1997) page 20 (Definition 2.6)

³² Christensen (2003) pages 65–66 (Lemma 3.6.5)

³³ Young (2001) page 27 (Theorem 1.8.9), Christensen (2003) page 66 (Theorem 3.6.6), Heil (2011) pages 197–198 (Theorem 7.13), Christensen (2008) pages 61–62 (Theorem 3.3.7)

1. Proof for (\implies) case:(a) Proof that *Riesz basis* hypothesis \implies (1): all bases for H are *total* in H .(b) Proof that *Riesz basis* hypothesis \implies (2):i. Let $(\mathbf{u}_n)_{n \in \mathbb{N}}$ be an *orthonormal basis* for H .ii. Let \mathbf{R} be a *bounded bijective* operator such that $\mathbf{x}_n = \mathbf{R}\mathbf{u}_n$.iii. Proof for upper bound B :

$$\begin{aligned}
 \left\| \sum_{n=1}^{\infty} \alpha_n \mathbf{x}_n \right\|^2 &= \left\| \sum_{n=1}^{\infty} \alpha_n \mathbf{R}\mathbf{u}_n \right\|^2 && \text{by definition of } \mathbf{R} && (\text{item (1(b)ii)}) \\
 &= \left\| \mathbf{R} \sum_{n=1}^{\infty} \alpha_n \mathbf{u}_n \right\|^2 && \text{by Theorem M.1 page 282} \\
 &\leq \|\mathbf{R}\|^2 \left\| \sum_{n=1}^{\infty} \alpha_n \mathbf{u}_n \right\|^2 && \text{by Theorem M.6 page 288} \\
 &= \|\mathbf{R}\|^2 \sum_{n=1}^{\infty} \|\alpha_n \mathbf{u}_n\|^2 && \text{by Pythagorean Theorem} && (\text{Theorem I.10 page 246}) \\
 &= \|\mathbf{R}\|^2 \sum_{n=1}^{\infty} |\alpha_n|^2 \|\mathbf{u}_n\|^2 && \text{by homogeneous property of norms} && (\text{Definition J.1 page 249}) \\
 &= \underbrace{\|\mathbf{R}\|^2}_{B} \sum_{n=1}^{\infty} |\alpha_n|^2 && \text{by definition of orthonormality} && (\text{Definition I.4 page 245})
 \end{aligned}$$

iv. Proof for lower bound A :

$$\begin{aligned}
 \left\| \sum_{n=1}^{\infty} \alpha_n \mathbf{x}_n \right\|^2 &= \frac{\|\mathbf{R}^{-1}\|^2}{\|\mathbf{R}^{-1}\|^2} \left\| \sum_{n=1}^{\infty} \alpha_n \mathbf{x}_n \right\|^2 && \text{because } \|\mathbf{R}^{-1}\| > 0 && (\text{Proposition M.1 page 286}) \\
 &\geq \frac{1}{\|\mathbf{R}^{-1}\|^2} \left\| \mathbf{R}^{-1} \sum_{n=1}^{\infty} \alpha_n \mathbf{x}_n \right\|^2 && \text{by Theorem M.6 page 288} \\
 &= \frac{1}{\|\mathbf{R}^{-1}\|^2} \left\| \mathbf{R}^{-1} \sum_{n=1}^{\infty} \alpha_n \mathbf{R}\mathbf{u}_n \right\|^2 && \text{by definition of } \mathbf{R} && (\text{item (1(b)ii) page 216}) \\
 &= \frac{1}{\|\mathbf{R}^{-1}\|^2} \left\| \mathbf{R}^{-1} \mathbf{R} \sum_{n=1}^{\infty} \alpha_n \mathbf{u}_n \right\|^2 && \text{by property of linear operators} && (\text{Theorem M.1 page 282}) \\
 &= \frac{1}{\|\mathbf{R}^{-1}\|^2} \left\| \sum_{n=1}^{\infty} \alpha_n \mathbf{u}_n \right\|^2 && \text{by definition of inverse op.} && (\text{Definition M.2 page 281}) \\
 &= \frac{1}{\|\mathbf{R}^{-1}\|^2} \sum_{n=1}^{\infty} \|\alpha_n \mathbf{u}_n\|^2 && \text{by Pythagorean Theorem} && (\text{Theorem I.10 page 246}) \\
 &= \frac{1}{\|\mathbf{R}^{-1}\|^2} \sum_{n=1}^{\infty} |\alpha_n|^2 \|\mathbf{u}_n\|^2 && \text{by } \|\cdot\| \text{ homogeneous prop.} && (\text{Definition J.1 page 249}) \\
 &= \underbrace{\frac{1}{\|\mathbf{R}^{-1}\|^2} \sum_{n=1}^{\infty} |\alpha_n|^2}_{A} && \text{by def. of orthonormality} && (\text{Definition I.4 page 245})
 \end{aligned}$$

2. Proof for (\implies) case:

- (a) Let $\{u_n\}_{n \in \mathbb{N}}$ be an *orthonormal basis* for H .
- (b) Using (2) and Lemma G.3 (page 215), construct an bounded extension operator R such that $Ru_n = x_n$ for all $n \in \mathbb{N}$.
- (c) Using (2) and Lemma G.3 (page 215), construct an bounded extension operator S such that $Sx_n = u_n$ for all $n \in \mathbb{N}$.
- (d) Then, $RVx = VRx \implies V = R^{-1}$, and so R is a bounded invertible operator
- (e) and $\{x_n\}$ is a *Riesz sequence*.



Theorem G.14. ³⁴ Let $H \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \Delta | \nabla \rangle)$ be a SEPARABLE HILBERT SPACE.

T
H
M

$$\left\{ \begin{array}{l} (\mathbf{x}_n \in H)_{n \in \mathbb{Z}} \text{ is a} \\ \text{RIESZ BASIS for } H \end{array} \right\} \implies \left\{ \begin{array}{l} \text{There exists } (\mathbf{y}_n \in H)_{n \in \mathbb{Z}} \text{ such that} \\ (1). (\mathbf{x}_n) \text{ and } (\mathbf{y}_n) \text{ are BIORTHOGONAL and} \\ (2). (\mathbf{y}_n) \text{ is also a RIESZ BASIS for } H \text{ and} \\ (3). \exists B > A > 0 \text{ such that} \\ A \sum_{n=1}^{\infty} |a_n|^2 \leq \left\| \sum_{n=1}^{\infty} a_n \mathbf{x}_n \right\|^2 = \left\| \sum_{n=1}^{\infty} a_n \mathbf{y}_n \right\|^2 \leq B \sum_{n=1}^{\infty} |a_n|^2 \\ \forall (a_n)_{n \in \mathbb{N}} \in \ell_{\mathbb{F}}^2 \end{array} \right\}$$

PROOF:

1. Proof for (1):

- (a) Let e_n be the *unit vector* in H such that the n th element of e_n is 1 and all other elements are 0.
- (b) Let M be an operator on H such that $Me_n = x_n$.
- (c) Note that M is *isometric*, and as such $\|Mx\| = \|x\| \quad \forall x \in H$.
- (d) Let $y_n \triangleq (M^{-1})^*$.
- (e) Then,

$$\begin{aligned} \langle y_n | x_m \rangle &= \left\langle (M^{-1})^* e_n | M e_m \right\rangle \\ &= \langle e_n | M^{-1} M e_m \rangle \\ &= \langle e_n | e_m \rangle \\ &= \bar{\delta}_{nm} \\ \implies \{x_n\} \text{ and } \{y_n\} \text{ are biorthogonal} &\quad \text{by Definition I.4 page 245} \end{aligned}$$

³⁴ Wojtaszczyk (1997) page 20 (Lemma 2.7(a))

2. Proof for (3):

$$\begin{aligned}
 \left\| \sum_{n \in \mathbb{Z}} \alpha_n y_n \right\| &= \left\| \sum_{n \in \mathbb{Z}} \alpha_n (\mathbf{M}^{-1})^* e_n \right\| && \text{by definition of } y_n && \text{(Proposition 1d page 217)} \\
 &= \left\| (\mathbf{M}^{-1})^* \sum_{n \in \mathbb{Z}} \alpha_n e_n \right\| && \text{by property of linear ops.} \\
 &= \left\| \sum_{n \in \mathbb{Z}} \alpha_n e_n \right\| && \text{because } (\mathbf{M}^{-1})^* \text{ is isometric} && \text{(Definition M.10 page 300)} \\
 &= \left\| \mathbf{M} \sum_{n \in \mathbb{Z}} \alpha_n e_n \right\| && \text{because } \mathbf{M} \text{ is isometric} && \text{(Definition M.10 page 300)} \\
 &= \left\| \sum_{n \in \mathbb{Z}} \alpha_n \mathbf{M} e_n \right\| && \text{by property of linear operators} \\
 &= \left\| \sum_{n \in \mathbb{Z}} \alpha_n x_n \right\| && \text{by definition of } \mathbf{M} \\
 \implies \{y_n\} &\text{ is a Riesz basis} && \text{by left hypothesis}
 \end{aligned}$$

3. Proof for (2): by (3) and definition of *Riesz basis* (Definition G.11 page 215)

 **Proposition G.5.** ³⁵ Let $\{x_n | n \in \mathbb{N}\}$ be a set of vectors in a HILBERT SPACE $\mathbf{H} \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle | \nabla \rangle)$.

P R P	$ \left\{ \begin{array}{l} \{x_n\} \text{ is a RIESZ BASIS for } \mathbf{H} \text{ with} \\ A \sum_{n=1}^{\infty} a_n ^2 \leq \left\ \sum_{n=1}^{\infty} a_n x_n \right\ ^2 \leq B \sum_{n=1}^{\infty} a_n ^2 \\ \forall \{a_n\} \in \ell_{\mathbb{F}}^2 \end{array} \right\} \Rightarrow \left\{ \begin{array}{l} \{x_n\} \text{ is a FRAME for } \mathbf{H} \text{ with} \\ \underbrace{\frac{1}{B} \ x\ ^2 \leq \sum_{n=1}^{\infty} \langle x x_n \rangle ^2 \leq \frac{1}{A} \ x\ ^2}_{\text{STABILITY CONDITION}} \\ \forall x \in \mathbf{H} \end{array} \right\} $
-------------	---

 PROOF:

1. Let $\{y_n | n \in \mathbb{N}\}$ be a *Riesz basis* that is *biorthonormal* to $\{x_n | n \in \mathbb{N}\}$ (Theorem G.14 page 217).

2. Let $x \triangleq \sum_{n=1}^{\infty} a_n y_n$.

3. lemma:

$$\begin{aligned}
 \sum_{n=1}^{\infty} |\langle x | x_n \rangle|^2 &= \sum_{n=1}^{\infty} \left| \left\langle \sum_{m=1}^{\infty} a_m y_m | x_n \right\rangle \right|^2 && \text{by definition of } x && \text{(item (2) page 218)} \\
 &= \sum_{n=1}^{\infty} \left| \sum_{m=1}^{\infty} a_m \langle y_m | x_n \rangle \right|^2 && \text{by homogeneous property of } \langle \triangle | \nabla \rangle && \text{(Definition I.1 page 233)} \\
 &= \sum_{n=1}^{\infty} \left| \sum_{m=1}^{\infty} a_m \bar{\delta}_{mn} \right|^2 && \text{by definition of biorthonormal} && \text{(Definition G.13 page 215)} \\
 &= \sum_{n=1}^{\infty} |a_n|^2 && \text{by definition of } \bar{\delta} && \text{(Definition I.3 page 245)}
 \end{aligned}$$

³⁵  Igari (1996) page 220 (Lemma 9.8),  Wojtaszczyk (1997) pages 20–21 (Lemma 2.7(a))

4. Then

$$\begin{aligned}
 A \sum_{n=1}^{\infty} |a_n|^2 &\leq \left\| \sum_{n=1}^{\infty} a_n \mathbf{x}_n \right\|^2 \leq B \sum_{n=1}^{\infty} |a_n|^2 && \text{by definition of } \{\mathbf{y}_n\} \text{ (item (1) page 218)} \\
 \implies A \sum_{n=1}^{\infty} |a_n|^2 &\leq \left\| \sum_{n=1}^{\infty} a_n \mathbf{y}_n \right\|^2 \leq B \sum_{n=1}^{\infty} |a_n|^2 && \text{by definition of } \{\mathbf{y}_n\} \text{ (item (1) page 218)} \\
 \implies A \sum_{n=1}^{\infty} |a_n|^2 &\leq \|\mathbf{x}\|^2 \leq B \sum_{n=1}^{\infty} |a_n|^2 && \text{by definition of } \mathbf{x} \text{ (item (2) page 218)} \\
 \implies A \sum_{n=1}^{\infty} |\langle \mathbf{x} | \mathbf{x}_n \rangle|^2 &\leq \|\mathbf{x}\|^2 \leq B \sum_{n=1}^{\infty} |\langle \mathbf{x} | \mathbf{x}_n \rangle|^2 && \text{by (3) lemma} \\
 \implies \frac{1}{B} \|\mathbf{x}\|^2 &\leq \sum_{n=1}^{\infty} |\langle \mathbf{x} | \mathbf{x}_n \rangle|^2 \leq \frac{1}{A} \|\mathbf{x}\|^2
 \end{aligned}$$



Theorem G.15 (Battle-Lemarié orthogonalization). ³⁶ Let $\tilde{f}(\omega)$ be the FOURIER TRANSFORM (Definition N.2 page 309) of a function $f \in L^2_{\mathbb{R}}$.

THM	$ \left\{ \begin{array}{l} 1. \quad \left\{ \mathbf{T}^n f \mid n \in \mathbb{Z} \right\} \text{ is a RIESZ BASIS for } L^2_{\mathbb{R}} \quad \text{and} \\ 2. \quad \tilde{f}(\omega) \triangleq \frac{\tilde{g}(\omega)}{\sqrt{2\pi \sum_{n \in \mathbb{Z}} \tilde{g}(\omega + 2\pi n) ^2}} \end{array} \right\} \Rightarrow \left\{ \begin{array}{l} \left\{ \mathbf{T}^n f \mid n \in \mathbb{Z} \right\} \\ \text{is an ORTHONORMAL BASIS for } L^2_{\mathbb{R}} \end{array} \right\} $
-----	---

PROOF:

1. Proof that $\{\mathbf{T}^n f \mid n \in \mathbb{Z}\}$ is orthonormal:

$$\begin{aligned}
 \tilde{S}_{\phi\phi}(\omega) &= 2\pi \sum_{n \in \mathbb{Z}} |\tilde{f}(\omega + 2\pi n)|^2 && \text{by Theorem ?? page ??} \\
 &= 2\pi \sum_{n \in \mathbb{Z}} \left| \frac{\tilde{g}(\omega + 2\pi n)}{\sqrt{2\pi \sum_{m \in \mathbb{Z}} |\tilde{g}(\omega + 2\pi(m-n))|^2}} \right|^2 && \text{by left hypothesis} \\
 &= \sum_{n \in \mathbb{Z}} \left| \frac{\tilde{g}(\omega + 2\pi n)}{\sqrt{\sum_{m \in \mathbb{Z}} |\tilde{g}(\omega + 2\pi m)|^2}} \right|^2 \\
 &= \sum_{n \in \mathbb{Z}} \left| \frac{1}{\sqrt{\sum_{m \in \mathbb{Z}} |\tilde{g}(\omega + 2\pi m)|^2}} \right|^2 |\tilde{g}(\omega + 2\pi n)|^2 \\
 &= \frac{1}{\sum_{m \in \mathbb{Z}} |\tilde{g}(\omega + 2\pi m)|^2} \sum_{n \in \mathbb{Z}} |\tilde{g}(\omega + 2\pi n)|^2 \\
 &= 1 \\
 \implies \{f_n \mid n \in \mathbb{Z}\} &\text{ is orthonormal} && \text{by Theorem ?? page ??}
 \end{aligned}$$

³⁶ Wojtaszczyk (1997) page 25 (Remark 2.4), Vidakovic (1999), page 71, Mallat (1989), page 72, Mallat (1999), page 225, Daubechies (1992) page 140 ((5.3.3))

2. Proof that $\{\mathbf{T}^n f \mid n \in \mathbb{Z}\}$ is a basis for V_0 : by Lemma G.1 page 203.



G.7 Frames in Hilbert spaces

Definition G.14. ³⁷ Let $\{x_n \in X \mid n \in \mathbb{N}\}$ be a set of vectors in a HILBERT SPACE $H \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \Delta | \nabla \rangle)$.

The set $\{x_n\}$ is a **frame** for H if (STABILITY CONDITION)

$$\exists A, B \in \mathbb{R}^+ \text{ such that } A \|x\|^2 \leq \sum_{n=1}^{\infty} |\langle x | x_n \rangle|^2 \leq B \|x\|^2 \quad \forall x \in X.$$

The quantities A and B are **frame bounds**.

D E F The quantity A' is the **optimal lower frame bound** if

$$A' = \sup \{A \in \mathbb{R}^+ \mid A \text{ is a lower frame bound}\}.$$

The quantity B' is the **optimal upper frame bound** if

$$B' = \inf \{B \in \mathbb{R}^+ \mid B \text{ is an upper frame bound}\}.$$

A frame is a **tight frame** if $A = B$.

A frame is a **normalized tight frame** (or a **Parseval frame**) if $A = B = 1$.

A frame $\{x_n \mid n \in \mathbb{N}\}$ is an **exact frame** if for some $m \in \mathbb{Z}$, $\{x_n \mid n \in \mathbb{N}\} \setminus \{x_m\}$ is NOT a frame.

A frame is a *Parseval frame* (Definition G.14) if it satisfies *Parseval's Identity* (Theorem G.9 page 210). All orthonormal bases are Parseval frames (Theorem G.9 page 210); but not all Parseval frames are orthonormal bases.

Definition G.15. Let $\{x_n\}$ be a **frame** (Definition G.14 page 220) for the HILBERT SPACE $H \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \Delta | \nabla \rangle)$. Let S be an OPERATOR on H .

D E F S is a **frame operator** for $\{x_n\}$ if $Sf(x) = \sum_{n \in \mathbb{Z}} \langle f | x_n \rangle x_n \quad \forall f \in H$.

Theorem G.16. ³⁸ Let S be a FRAME OPERATOR (Definition G.15 page 220) of a FRAME $\{x_n\}$ (Definition G.14 page 220) for the HILBERT SPACE $H \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \Delta | \nabla \rangle)$.

T H M

- (1). S is INVERTIBLE.
- (2). $f(x) = \sum_{n \in \mathbb{Z}} \langle f | S^{-1} x_n \rangle x_n = \sum_{n \in \mathbb{Z}} \langle f | x_n \rangle S^{-1} x_n \quad \forall f \in H$

Theorem G.17. ³⁹ Let $\{x_n \in X \mid n=1,2,\dots,N\}$ be a set of vectors in a HILBERT SPACE $H \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \Delta | \nabla \rangle)$.

T H M $\{x_n\}$ is a FRAME for $\text{span}\{x_n\}$.

PROOF:

³⁷ Young (2001) pages 154–155, Christensen (2003) page 88 (Definitions 5.1.1, 5.1.2), Heil (2011) pages 204–205 (Definition 8.2), Jørgensen et al. (2008) page 267 (Definition 12.22), Duffin and Schaeffer (1952) page 343, Daubechies et al. (1986), page 1272

³⁸ Christensen (2008) pages 100–102 (Theorem 5.1.7)

³⁹ Christensen (2003) page 3

1. Upper bound: Proof that there exists B such that $\sum_{n=1}^N |\langle \mathbf{x} | \mathbf{x}_n \rangle|^2 \leq B \|\mathbf{x}\|^2 \quad \forall \mathbf{x} \in \mathcal{H}$:

$$\begin{aligned} \sum_{n=1}^N |\langle \mathbf{x} | \mathbf{x}_n \rangle|^2 &\leq \sum_{n=1}^N \langle \mathbf{x}_n | \mathbf{x}_n \rangle \langle \mathbf{x} | \mathbf{x} \rangle \quad \text{by Cauchy-Schwarz inequality (Theorem I.2 page 234)} \\ &= \underbrace{\left\{ \sum_{n=1}^N \|\mathbf{x}_n\|^2 \right\}}_B \|\mathbf{x}\|^2 \end{aligned}$$

2. Lower bound: Proof that there exists A such that $A \|\mathbf{x}\|^2 \leq \sum_{n=1}^N |\langle \mathbf{x} | \mathbf{x}_n \rangle|^2 \quad \forall \mathbf{x} \in \mathcal{H}$:

$$\begin{aligned} \sum_{n=1}^N |\langle \mathbf{x} | \mathbf{x}_n \rangle|^2 &= \sum_{n=1}^N \left| \left\langle \mathbf{x}_n | \frac{\mathbf{x}}{\|\mathbf{x}\|} \right\rangle \right|^2 \|\mathbf{x}\|^2 \\ &\geq \underbrace{\left(\inf_y \left\{ \sum_{n=1}^N |\langle \mathbf{x}_n | y \rangle|^2 | \|y\| = 1 \right\} \right)}_A \|\mathbf{x}\|^2 \end{aligned}$$


Example G.1. Let $(\mathbb{R}^2, +, \cdot, (\mathbb{R}, \dot{+}, \dot{\times}), \langle \triangle | \triangleright \rangle)$ be an inner product space with $\left\langle \begin{bmatrix} x_1 \\ y_1 \end{bmatrix} | \begin{bmatrix} x_2 \\ y_2 \end{bmatrix} \right\rangle \triangleq x_1 x_2 + y_1 y_2$. Let \mathbf{S} be the *frame operator* (Definition G.15 page 220) with *inverse* \mathbf{S}^{-1} .

EX

Let $N \in \{3, 4, 5, \dots\}$, $\theta \in \mathbb{R}$, and $r \in \mathbb{R}^+$ ($r > 0$).

Let $\mathbf{x}_n \triangleq r \begin{bmatrix} \cos(\theta + 2n\pi/N) \\ \sin(\theta + 2n\pi/N) \end{bmatrix} \quad \forall n \in \{0, 1, \dots, N-1\}$.

Then, $(\mathbf{x}_0, \mathbf{x}_1, \dots, \mathbf{x}_{N-1})$ is a **tight frame** for \mathbb{R}^2 with *frame bound* $A = \frac{Nr^2}{2}$.



Moreover, $\mathbf{S} = \frac{Nr^2}{2} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$, $\mathbf{S}^{-1} = \frac{2}{Nr^2} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$, and $\mathbf{v} = \frac{2}{Nr^2} \sum_{n=0}^{N-1} \langle \mathbf{v} | \mathbf{x}_n \rangle \mathbf{x}_n \quad \forall \mathbf{v} \in \mathbb{R}^2$.

PROOF:

1. Proof that $(\mathbf{x}_0, \mathbf{x}_1, \dots, \mathbf{x}_{N-1})$ is a *tight frame* with *frame bound* $A = \frac{Nr^2}{2}$: Let $\mathbf{v} \triangleq (x, y) \in \mathbb{R}^2$.

$$\begin{aligned} \sum_{n=0}^{N-1} |\langle \mathbf{v} | \mathbf{x}_n \rangle|^2 &\triangleq \sum_{n=0}^{N-1} \left| \mathbf{v}^H \begin{bmatrix} \cos\left(\theta + \frac{2n\pi}{N}\right) \\ \sin\left(\theta + \frac{2n\pi}{N}\right) \end{bmatrix} \right|^2 && \text{by definitions of } \mathbf{v} \text{ of } \langle \mathbf{y} | \mathbf{x} \rangle \\ &\triangleq \sum_{n=0}^{N-1} r^2 \left| x \cos\left(\theta + \frac{2n\pi}{N}\right) + y \sin\left(\theta + \frac{2n\pi}{N}\right) \right|^2 && \text{by definition of } \mathbf{y}^H \mathbf{x} \text{ operation} \\ &= r^2 x^2 \sum_{n=0}^{N-1} \cos^2\left(\theta + \frac{2n\pi}{N}\right) + r^2 y^2 \sum_{n=0}^{N-1} \sin^2\left(\theta + \frac{2n\pi}{N}\right) + r^2 xy \sum_{n=0}^{N-1} \cos\left(\theta + \frac{2n\pi}{N}\right) \sin\left(\theta + \frac{2n\pi}{N}\right) \\ &= r^2 x^2 \frac{N}{2} + r^2 y^2 \frac{N}{2} + r^2 xy 0 && \text{by Corollary ?? page ??} \\ &= (x^2 + y^2) \frac{Nr^2}{2} = \underbrace{\left(\frac{Nr^2}{2} \right)}_A \|\mathbf{v}\|^2 && \text{by definition of } \|\mathbf{v}\| \end{aligned}$$

2. Proof that $\mathbf{S} = \frac{Nr^2}{2} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$:

(a) Let $e_1 \triangleq \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ and $e_2 \triangleq \begin{bmatrix} 0 \\ 1 \end{bmatrix}$.

(b) lemma: $\mathbf{S}e_1 = \frac{Nr^2}{2}e_1$. Proof:

$$\begin{aligned}\mathbf{S}e_1 &= \sum_{n=0}^{N-1} \langle e_1 | \mathbf{x}_n \rangle \mathbf{x}_n \\ &= \sum_{n=0}^{N-1} r \cos\left(\theta + \frac{2n\pi}{N}\right) r \begin{bmatrix} \cos\left(\theta + \frac{2n\pi}{N}\right) \\ \sin\left(\theta + \frac{2n\pi}{N}\right) \end{bmatrix} = r^2 \sum_{n=0}^{N-1} \begin{bmatrix} \cos^2\left(\theta + \frac{2n\pi}{N}\right) \\ \cos\left(\theta + \frac{2n\pi}{N}\right) \sin\left(\theta + \frac{2n\pi}{N}\right) \end{bmatrix} \\ &= r^2 \begin{bmatrix} N/2 \\ 0 \end{bmatrix} = \frac{Nr^2}{2}e_1 \quad \text{by Summation around unit circle (Corollary ?? page ??)}$$

(c) lemma: $\mathbf{S}e_2 = \frac{Nr^2}{2}e_2$. Proof:

$$\begin{aligned}\mathbf{S}e_2 &= \sum_{n=0}^{N-1} \langle e_2 | \mathbf{x}_n \rangle \mathbf{x}_n \\ &= \sum_{n=0}^{N-1} r \sin\left(\theta + \frac{2n\pi}{N}\right) r \begin{bmatrix} \cos\left(\theta + \frac{2n\pi}{N}\right) \\ \sin\left(\theta + \frac{2n\pi}{N}\right) \end{bmatrix} = r^2 \sum_{n=0}^{N-1} \begin{bmatrix} \sin\left(\theta + \frac{2n\pi}{N}\right) \cos\left(\theta + \frac{2n\pi}{N}\right) \\ \sin^2\left(\theta + \frac{2n\pi}{N}\right) \end{bmatrix} \\ &= r^2 \begin{bmatrix} 0 \\ N/2 \end{bmatrix} = \frac{Nr^2}{2}e_2 \quad \text{by Summation around unit circle (Corollary ?? page ??)}$$

(d) Complete the proof of item (2) using Eigendecomposition $\mathbf{S} = \mathbf{Q}\Lambda\mathbf{Q}^{-1}$:

$$\mathbf{S}e_1 = \frac{Nr^2}{2}e_1 \quad \text{by (2c) lemma}$$

$\Rightarrow e_1$ is an eigenvector of \mathbf{S} with eigenvalue $\frac{Nr^2}{2}$

$$\mathbf{S}e_2 = \frac{Nr^2}{2}e_2 \quad \text{by (2c) lemma}$$

$\Rightarrow e_2$ is an eigenvector of \mathbf{S} with eigenvalue $\frac{Nr^2}{2}$

$$\overbrace{\mathbf{S} = \underbrace{\begin{bmatrix} 1 & 1 \\ e_1 & e_2 \\ 1 & 1 \end{bmatrix}}_{\mathbf{Q}} \underbrace{\begin{bmatrix} \frac{Nr^2}{2} & 0 \\ 0 & \frac{Nr^2}{2} \end{bmatrix}}_{\Lambda} \underbrace{\begin{bmatrix} 1 & 1 \\ e_1 & e_2 \\ 1 & 1 \end{bmatrix}}_{\mathbf{Q}^{-1}}}^{\text{Eigendecomposition of } \mathbf{S}} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} \frac{Nr^2}{2} & 0 \\ 0 & \frac{Nr^2}{2} \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}^{-1} = \frac{Nr^2}{2} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

3. Proof that $\mathbf{S}^{-1} = \frac{2}{Nr^2} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$:

$$\mathbf{S}\mathbf{S}^{-1} = \frac{Nr^2}{2} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \frac{2}{Nr^2} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \triangleq \mathbf{I}_2 \quad \text{by item (2)}$$

$$\mathbf{S}^{-1}\mathbf{S} = \frac{2}{Nr^2} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \frac{Nr^2}{2} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \triangleq \mathbf{I}_2 \quad \text{by item (2)}$$

4. Proof that $\mathbf{v} = \frac{2}{Nr^2} \sum_{n=0}^{N-1} (\mathbf{v}^H \mathbf{x}_n) \mathbf{x}_n$:

$$\mathbf{v} = \sum_{n=0}^{N-1} \langle \mathbf{v} | \mathbf{S}^{-1} \mathbf{x}_n \rangle \mathbf{x}_n = \sum_{n=0}^{N-1} \left\langle \mathbf{v} \mid \frac{2}{Nr^2} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \mathbf{x}_n \right\rangle \mathbf{x}_n \quad \text{by item (3)}$$

$$= \frac{2}{Nr^2} \sum_{n=0}^{N-1} \langle \mathbf{v} | \mathbf{x}_n \rangle \mathbf{x}_n = \frac{2}{Nr^2} \sum_{n=0}^{N-1} (\mathbf{v}^H \mathbf{x}_n) \mathbf{x}_n \quad \text{by definition of } \langle \mathbf{y} | \mathbf{x} \rangle$$


Example G.2 (Peace Frame/Mercedes Frame). ⁴⁰ Let $(\mathbb{R}^2, +, \cdot, (\mathbb{R}, \dot{+}, \dot{\times}), \langle \triangle | \nabla \rangle)$ be an inner product space with $\langle (x_1, y_1) | (x_2, y_2) \rangle \triangleq x_1y_1 + x_2y_2$. Let \mathbf{S} be the *frame operator* (Definition G.15 page 220) with inverse \mathbf{S}^{-1} .

Let $\mathbf{x}_1 \triangleq \frac{1}{2} \begin{bmatrix} 0 \\ 2 \end{bmatrix}$, $\mathbf{x}_2 \triangleq \frac{1}{2} \begin{bmatrix} -\sqrt{3} \\ -1 \end{bmatrix}$, and $\mathbf{x}_3 \triangleq \frac{1}{2} \begin{bmatrix} \sqrt{3} \\ -1 \end{bmatrix}$.

E X Then, $(\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3)$ is a **tight frame** for \mathbb{R}^2 with *frame bound* $A = \frac{3}{2}$.

Moreover, $\mathbf{S} = \frac{3}{2} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$, $\mathbf{S}^{-1} = \frac{2}{3} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$,

and $\mathbf{v} = \frac{2}{3} \sum_{n=1}^3 \langle \mathbf{v} | \mathbf{x}_n \rangle \mathbf{x}_n \triangleq \frac{2}{3} \sum_{n=1}^3 (\mathbf{v}^H \mathbf{x}_n) \mathbf{x}_n \quad \forall \mathbf{v} \in \mathbb{R}^2$.



PROOF:

1. This frame is simply a special case of the frame presented in Example G.1 (page 221) with $r = 1$, $N = 3$, and $\theta = \pi/2$.

2. Let's give it a try! Let $\mathbf{v} \triangleq \begin{bmatrix} 1 \\ 1 \end{bmatrix}$.

$$\begin{aligned} \sum_{n=1}^3 \langle \mathbf{v} | \mathbf{S}^{-1} \mathbf{x}_n \rangle \mathbf{x}_n &= \frac{2}{3} \sum_{n=1}^3 (\mathbf{v}^H \mathbf{x}_n) \mathbf{x}_n && \text{by Example G.1 page 221} \\ &= (\mathbf{v}^H \mathbf{x}_1) \mathbf{x}_1 + (\mathbf{v}^H \mathbf{x}_2) \mathbf{x}_2 + (\mathbf{v}^H \mathbf{x}_3) \mathbf{x}_3 \\ &= \frac{2}{3} \left(\left(\mathbf{v}^H \frac{1}{2} \begin{bmatrix} 0 \\ 2 \end{bmatrix} \right) \mathbf{x}_1 + \left(\mathbf{v}^H \frac{1}{2} \begin{bmatrix} -\sqrt{3} \\ -1 \end{bmatrix} \right) \mathbf{x}_2 + \left(\mathbf{v}^H \frac{1}{2} \begin{bmatrix} \sqrt{3} \\ -1 \end{bmatrix} \right) \mathbf{x}_3 \right) \\ &= \frac{2}{3} \cdot \frac{1}{2} \left(\left(\mathbf{v}^H \begin{bmatrix} 0 \\ 2 \end{bmatrix} \right) \mathbf{x}_1 + \left(\mathbf{v}^H \begin{bmatrix} -\sqrt{3} \\ -1 \end{bmatrix} \right) \mathbf{x}_2 + \left(\mathbf{v}^H \begin{bmatrix} \sqrt{3} \\ -1 \end{bmatrix} \right) \mathbf{x}_3 \right) \\ &= \frac{1}{3} \left((2) \frac{1}{2} \begin{bmatrix} 0 \\ 2 \end{bmatrix} + (-\sqrt{3}-1) \frac{1}{2} \begin{bmatrix} -\sqrt{3} \\ -1 \end{bmatrix} + (\sqrt{3}-1) \frac{1}{2} \begin{bmatrix} \sqrt{3} \\ -1 \end{bmatrix} \right) \\ &= \frac{1}{6} \left[\begin{array}{l} 2(0) + (-\sqrt{3}-1)(-\sqrt{3}) + (\sqrt{3}-1)(\sqrt{3}) \\ 2(2) + (-\sqrt{3}-1)(-1) + (\sqrt{3}-1)(-1) \end{array} \right] \\ &= \frac{1}{6} \left[\begin{array}{l} 0 + (3+\sqrt{3}) + (3-\sqrt{3}) \\ 4 + (1+\sqrt{3}) + (1-\sqrt{3}) \end{array} \right] = \begin{bmatrix} 1 \\ 1 \end{bmatrix} \triangleq \mathbf{v} \end{aligned}$$

In Example G.1 (page 221) and Example G.2 (page 223), the frame operator \mathbf{S} and its inverse \mathbf{S}^{-1} were computed. In general however, it is not always necessary or even possible to compute these, as illustrated in Example G.3 (next).


Example G.3. ⁴¹ Let $(\mathbb{R}^2, +, \cdot, (\mathbb{R}, \dot{+}, \dot{\times}), \langle \triangle | \nabla \rangle)$ be an inner product space with $\langle (x_1, y_1) | (x_2, y_2) \rangle \triangleq x_1y_1 + x_2y_2$. Let \mathbf{S} be the *frame operator* (Definition G.15 page 220) with inverse \mathbf{S}^{-1} .

⁴⁰ Heil (2011) pages 204–205 ($r = 1$ case), Byrne (2005) page 80 ($r = 1$ case), Han et al. (2007) page 91 (Example 3.9, $r = \sqrt{2}/3$ case)

⁴¹ Christensen (2003) pages 7–8 (?)

E

Let p and q be orthonormal vectors in $\mathbf{X} \triangleq \text{span}\{p, q\}$.
 Let $x_1 \triangleq p$, $x_2 \triangleq p + q$, and $x_3 \triangleq p - q$.
 Then, $\{x_1, x_2, x_3\}$ is a frame for \mathbf{X} with frame bounds $A = 0$ and $B = 5$.



Moreover,
 $S^{-1}x_1 = \frac{1}{3}p$ and
 $S^{-1}x_2 = \frac{1}{3}p + \frac{1}{2}q$ and
 $S^{-1}x_3 = \frac{1}{3}p - \frac{1}{2}q$.

PROOF:

1. Proof that (x_1, x_2, x_3) is a frame with frame bounds $A = 0$ and $B = 5$:

$$\begin{aligned} \sum_{n=1}^3 |\langle v | x_n \rangle|^2 &\triangleq |\langle v | p \rangle|^2 + |\langle v | p + q \rangle|^2 + |\langle v | p - q \rangle|^2 && \text{by definitions of } x_1, x_2, \text{ and } x_3 \\ &= |\langle v | p \rangle|^2 + |\langle v | p \rangle + \langle v | q \rangle|^2 + |\langle v | p \rangle - \langle v | q \rangle|^2 && \text{by additivity of } \langle \Delta | \nabla \rangle \text{ (Definition I.1 page 233)} \\ &= |\langle v | p \rangle|^2 + (|\langle v | p \rangle|^2 + |\langle v | q \rangle|^2 + \langle v | p \rangle \langle v | q \rangle^* + \langle v | q \rangle \langle v | p \rangle^*) \\ &\quad + (|\langle v | p \rangle|^2 + |\langle v | q \rangle|^2 - \langle v | p \rangle \langle v | q \rangle^* - \langle v | q \rangle \langle v | p \rangle^*) \\ &= 3|\langle v | p \rangle|^2 + 2|\langle v | q \rangle|^2 \\ &\leq 3\|v\| \|p\| + 2\|v\| \|q\| && \text{by CS Inequality (Theorem I.2 page 234)} \\ &= \|v\| (3\|p\| + 2\|q\|) \\ &= \boxed{5\|v\|} && \text{by orthonormality of } p \text{ and } q \end{aligned}$$

2. lemma: $Sp = 3p$, $Sq = 2q$, $S^{-1}p = \frac{1}{3}p$, and $S^{-1}q = \frac{1}{2}q$. Proof:

$$\begin{aligned} Sp &\triangleq \sum_{n=1}^3 \langle p | x_n \rangle x_n \\ &= \langle p | p \rangle p + \langle p | p + q \rangle (p + q) + \langle p | p - q \rangle (p - q) \\ &= (1)p + (1+0)(p+q) + (1-0)(p-q) \\ &= 3p \\ \implies S^{-1}p &= \frac{1}{3}p \\ Sq &\triangleq \sum_{n=1}^3 \langle q | x_n \rangle x_n \\ &= \langle q | p \rangle p + \langle q | p + q \rangle (p + q) + \langle q | p - q \rangle (p - q) \\ &= (0)q + (0+1)(p+q) + (0-1)(p-q) \\ &= 2q \\ \implies S^{-1}q &= \frac{1}{2}q \end{aligned}$$

3. Remark: Without knowing p and q , from (2) lemma it follows that it is not possible to compute S or S^{-1} explicitly.

4. Proof that $S^{-1}x_1 = \frac{1}{3}p$, $S^{-1}x_2 = \frac{1}{3}p + \frac{1}{2}q$ and $S^{-1}x_3 = \frac{1}{3}p - \frac{1}{2}q$:

$$\begin{aligned} S^{-1}x_1 &\triangleq S^{-1}p && \text{by definition of } x_1 \\ &= \frac{1}{3}p && \text{by (2) lemma} \\ S^{-1}x_2 &\triangleq S^{-1}(p + q) && \text{by definition of } x_2 \\ &= \frac{1}{3}p + \frac{1}{2}q && \text{by (2) lemma} \end{aligned}$$

$$\begin{aligned} \mathbf{S}^{-1}\mathbf{x}_3 &\triangleq \mathbf{S}^{-1}(\mathbf{p} - \mathbf{q}) \\ &= \frac{1}{3}\mathbf{p} - \frac{1}{2}\mathbf{q} \end{aligned}$$

by definition of \mathbf{x}_2
by (2) lemma

5. Check that $\mathbf{v} = \sum_n \langle \mathbf{v} | \mathbf{x}_n \rangle \mathbf{x}_n = \langle \mathbf{v} | \mathbf{p} \rangle \mathbf{p} + \langle \mathbf{v} | \mathbf{q} \rangle \mathbf{q}$:

$$\begin{aligned} \mathbf{v} &= \sum_{n=1}^3 \langle \mathbf{v} | \mathbf{S}^{-1}\mathbf{x}_n \rangle \mathbf{x}_n \\ &= \langle \mathbf{v} | \mathbf{S}^{-1}\mathbf{p} \rangle \mathbf{p} + \langle \mathbf{v} | \mathbf{S}^{-1}(\mathbf{p} + \mathbf{q}) \rangle (\mathbf{p} + \mathbf{q}) + \langle \mathbf{v} | \mathbf{S}^{-1}(\mathbf{p} - \mathbf{q}) \rangle (\mathbf{p} - \mathbf{q}) \\ &= \left\langle \mathbf{v} \mid \frac{1}{3}\mathbf{p} \right\rangle \mathbf{p} + \left\langle \mathbf{v} \mid \frac{1}{3}\mathbf{p} + \frac{1}{2}\mathbf{q} \right\rangle (\mathbf{p} + \mathbf{q}) + \left\langle \mathbf{v} \mid \frac{1}{3}\mathbf{p} - \frac{1}{2}\mathbf{q} \right\rangle (\mathbf{p} - \mathbf{q}) \\ &= \left(\frac{1}{3} + \frac{1}{3} + \frac{1}{3} \right) \langle \mathbf{v} | \mathbf{p} \rangle \mathbf{p} + \left(\frac{1}{3} - \frac{1}{3} \right) \langle \mathbf{v} | \mathbf{p} \rangle \mathbf{q} + \left(\frac{1}{2} - \frac{1}{2} \right) \langle \mathbf{v} | \mathbf{q} \rangle \mathbf{p} + \left(\frac{1}{2} + \frac{1}{2} \right) \langle \mathbf{v} | \mathbf{q} \rangle \mathbf{q} \\ &= \langle \mathbf{v} | \mathbf{p} \rangle \mathbf{p} + \langle \mathbf{v} | \mathbf{q} \rangle \mathbf{q} \end{aligned}$$



APPENDIX H

NORMED ALGEBRAS

H.1 Algebras

All *linear spaces* (Definition F.1 page 185) are equipped with an operation by which vectors in the spaces can be added together. Linear spaces also have an operation that allows a scalar and a vector to be “multiplied” together. But linear spaces in general have no operation that allows two vectors to be multiplied together. A linear space together with such an operator is an **algebra**.¹

There are many many possible algebras—many more than one can shake a stick at, as indicated by Michiel Hazewinkel in his book, *Handbook of Algebras*: “Algebra, as we know it today (2005), consists of many different ideas, concepts and results. A reasonable estimate of the number of these different items would be somewhere between 50,000 and 200,000. Many of these have been named and many more could (and perhaps should) have a “name” or other convenient designation.”²

Definition H.1. ³ Let \mathbf{A} be an ALGEBRA.

D E F An algebra \mathbf{A} is **unital** if $\exists u \in \mathbf{A}$ such that $ux = xu = x \quad \forall x \in \mathbf{A}$

Definition H.2. ⁴ Let \mathbf{A} be an UNITAL ALGEBRA (Definition H.1 page 227) with unit e .

D E F The **spectrum** of $x \in \mathbf{A}$ is $\sigma(x) \triangleq \{\lambda \in \mathbb{C} \mid \lambda e - x \text{ is not invertible}\}$.
The **resolvent** of $x \in \mathbf{A}$ is $\rho_x(\lambda) \triangleq (\lambda e - x)^{-1} \quad \forall \lambda \notin \sigma(x)$.
The **spectral radius** of $x \in \mathbf{A}$ is $r(x) \triangleq \sup \{|\lambda| \mid \lambda \in \sigma(x)\}$.

¹ Fuchs (1995) page 2

² Hazewinkel (2000) page v

³ Folland (1995) page 1

⁴ Folland (1995) pages 3–4

H.2 Star-Algebras

Definition H.3. ⁵ Let A be an ALGEBRA.

The pair $(A, *)$ is a ***-algebra**, or **star-algebra**, if

- DEF 1. $(x + y)^* = x^* + y^*$ $\forall x, y \in A$ (DISTRIBUTIVE) and
 2. $(\alpha x)^* = \bar{\alpha} x^*$ $\forall x \in A, \alpha \in \mathbb{C}$ (CONJUGATE LINEAR) and
 3. $(xy)^* = y^* x^*$ $\forall x, y \in A$ (ANTIAUTOMORPHIC) and
 4. $x^{**} = x$ $\forall x \in A$ (INVOLUTORY)

The operator $*$ is called an **involution** on the algebra A .

Proposition H.1. ⁶ Let $(A, *)$ be an UNITAL *-ALGEBRA.

PRP x is invertible $\Rightarrow \begin{cases} 1. x^* \text{ is INVERTIBLE } \forall x \in A \text{ and} \\ 2. (x^*)^{-1} = (x^{-1})^* \quad \forall x \in A \end{cases}$

PROOF: Let e be the unit element of $(A, *)$.

1. Proof that $e^* = e$:

$$\begin{aligned} x e^* &= (x e^*)^{**} && \text{by involutory property of } * && (\text{Definition H.3 page 228}) \\ &= (x^* e^{**})^* && \text{by antiautomorphic property of } * && (\text{Definition H.3 page 228}) \\ &= (x^* e)^* && \text{by involutory property of } * && (\text{Definition H.3 page 228}) \\ &= (x^*)^* && \text{by definition of } e && \\ &= x && \text{by involutory property of } * && (\text{Definition H.3 page 228}) \\ e^* x &= (e^* x)^{**} && \text{by involutory property of } * && (\text{Definition H.3 page 228}) \\ &= (e^{**} x^*)^* && \text{by antiautomorphic property of } * && (\text{Definition H.3 page 228}) \\ &= (e x^*)^* && \text{by involutory property of } * && (\text{Definition H.3 page 228}) \\ &= (x^*)^* && \text{by definition of } e && \\ &= x && \text{by involutory property of } * && (\text{Definition H.3 page 228}) \end{aligned}$$

2. Proof that $(x^*)^{-1} = (x^{-1})^*$:

$$\begin{aligned} (x^{-1})^* (x^*) &= [x (x^{-1})]^* && \text{by antiautomorphic and involution properties of } * && (\text{Definition H.3 page 228}) \\ &= e^* \\ &= e && \text{by item (1) page 228} && \\ (x^*) (x^{-1})^* &= [x^{-1} x]^* && \text{by antiautomorphic and involution properties of } * && (\text{Definition H.3 page 228}) \\ &= e^* \\ &= e && \text{by item (1) page 228} && \end{aligned}$$

Definition H.4. ⁷ Let $(A, \|\cdot\|)$ be a *-ALGEBRA (Definition H.3 page 228).

DEF  An element $x \in A$ is **hermitian** or **self-adjoint** if $x^* = x$.

 An element $x \in A$ is **normal** if $xx^* = x^*x$.

 An element $x \in A$ is a **projection** if $xx = x$ (INVOLUTORY) and $x^* = x$ (HERMITIAN).

⁵  Rickart (1960), page 178,  Gelfand and Naimark (1964), page 241

⁶  Folland (1995) page 5

⁷  Rickart (1960), page 178,  Gelfand and Naimark (1964), page 242

Theorem H.1. ⁸ Let $(A, \|\cdot\|)$ be a $*$ -ALGEBRA (Definition H.3 page 228).

T
H
M

$$\underbrace{x = x^* \text{ and } y = y^*}_{x \text{ and } y \text{ are hermitian}}$$

 \Rightarrow

$$\begin{cases} x + y = (x + y)^* & (x + y \text{ is selfadjoint}) \\ x^* = (x^*)^* & (x^* \text{ is selfadjoint}) \\ \underbrace{xy = (xy)^*}_{(xy) \text{ is hermitian}} \iff \underbrace{xy = yx}_{\text{commutative}} & \end{cases}$$

PROOF:

$$\begin{aligned} (x + y)^* &= x^* + y^* && \text{by distributive property of } * \\ &= x + y && \text{by left hypothesis} \end{aligned} \quad (\text{Definition H.3 page 228})$$

$$(x^*)^* = x \quad \text{by involutory property of } * \quad (\text{Definition H.3 page 228})$$

Proof that $xy = (xy)^* \implies xy = yx$

$$\begin{aligned} xy &= (xy)^* && \text{by left hypothesis} \\ &= y^* x^* && \text{by antiautomorphic property of } * \\ &= yx && \text{by left hypothesis} \end{aligned} \quad (\text{Definition H.3 page 228})$$

Proof that $xy = (xy)^* \iff xy = yx$

$$\begin{aligned} (xy)^* &= (yx)^* && \text{by left hypothesis} \\ &= x^* y^* && \text{by antiautomorphic property of } * \\ &= xy && \text{by left hypothesis} \end{aligned} \quad (\text{Definition H.3 page 228})$$

Definition H.5 (Hermitian components). ⁹ Let $(A, \|\cdot\|)$ be a $*$ -ALGEBRA (Definition H.3 page 228).

DEF

$$\begin{aligned} \text{The real part of } x \text{ is defined as } \mathbf{R}_e x &\triangleq \frac{1}{2}(x + x^*) \\ \text{The imaginary part of } x \text{ is defined as } \mathbf{I}_m x &\triangleq \frac{1}{2i}(x - x^*) \end{aligned}$$

Theorem H.2. ¹⁰ Let $(A, *)$ be a $*$ -ALGEBRA (Definition H.3 page 228).

T
H
M

$$\begin{aligned} \Re x &= (\Re x)^* && \forall x \in A && (\Re x \text{ is hermitian}) \\ \Im x &= (\Im x)^* && \forall x \in A && (\Im x \text{ is hermitian}) \end{aligned}$$

PROOF:

$$\begin{aligned} (\Re x)^* &= \left(\frac{1}{2}(x + x^*)\right)^* && \text{by definition of } \Re \\ &= \frac{1}{2}(x^* + x^{**}) && \text{by distributive property of } * \\ &= \frac{1}{2}(x^* + x) && \text{by involutory property of } * \\ &= \Re x && \text{by definition of } \Re \\ (\Im x)^* &= \left(\frac{1}{2i}(x - x^*)\right)^* && \text{by definition of } \Im \end{aligned} \quad (\text{Definition H.5 page 229})$$

⁸ Michel and Herget (1993) page 429

⁹ Michel and Herget (1993) page 430, Rickart (1960), page 179, Gelfand and Naimark (1964), page 242

¹⁰ Michel and Herget (1993) page 430, Halmos (1998) page 42

$$\begin{aligned}
 &= \frac{1}{2i}(x^* - x^{**}) && \text{by distributive property of } * && (\text{Definition H.3 page 228}) \\
 &= \frac{1}{2i}(x^* - x) && \text{by involutory property of } * && (\text{Definition H.3 page 228}) \\
 &= \Im x && \text{by definition of } \Im && (\text{Definition H.5 page 229})
 \end{aligned}$$

⇒

Theorem H.3 (Hermitian representation). ¹¹ Let $(A, *)$ be a $*$ -ALGEBRA (Definition H.3 page 228).

T	H	M	$a = x + iy \iff x = \Re a \text{ and } y = \Im a$
---	---	---	--

PROOF:

Proof that $a = x + iy \implies x = \Re a \text{ and } y = \Im a$:

$$\begin{aligned}
 &a = x + iy && \text{by left hypothesis} \\
 \implies &a^* = (x + iy)^* && \text{by definition of adjoint} && (\text{Definition H.4 page 228}) \\
 &= x^* - iy^* && \text{by distributive property of } * && (\text{Definition H.3 page 228}) \\
 &= x - iy && \text{by Theorem H.2 page 229} \\
 \implies &x = a - iy && \text{by solving for } x \text{ in } a = x + iy \text{ equation} \\
 &x = a^* + iy && \text{by solving for } x \text{ in } a^* = x - iy \text{ equation} \\
 \implies &x + x = a + a^* && \text{by adding previous 2 equations} \\
 \implies &2x = a + a^* && \text{by solving for } x \text{ in previous equation} \\
 \implies &x = \frac{1}{2}(a + a^*) && \\
 &= \Re a && \text{by definition of } \Re && (\text{Definition H.5 page 229}) \\
 \\
 &iy = a - x && \text{by solving for } iy \text{ in } a = x + iy \text{ equation} \\
 &iy = -a^* + x && \text{by solving for } iy \text{ in } a = x + iy \text{ equation} \\
 \implies &iy + iy = a - a^* && \text{by adding previous 2 equations} \\
 \implies &y = \frac{1}{2i}(a - a^*) && \text{by solving for } iy \text{ in previous equations} \\
 &= \Im a && \text{by definition of } \Im && (\text{Definition H.5 page 229})
 \end{aligned}$$

Proof that $a = x + iy \iff x = \Re a \text{ and } y = \Im a$:

$$\begin{aligned}
 x + iy &= \Re a + i\Im a && \text{by right hypothesis} \\
 &= \underbrace{\frac{1}{2}(a + a^*)}_{\Re a} + i\underbrace{\frac{1}{2i}(a - a^*)}_{\Im a} && \text{by definition of } \Re \text{ and } \Im && (\text{Definition H.5 page 229}) \\
 &= \left(\frac{1}{2}a + \frac{1}{2}a\right) + \left(\frac{1}{2}a^* - \frac{1}{2}a^*\right) \xrightarrow{0} 0 \\
 &= a
 \end{aligned}$$

⇒

¹¹ Michel and Herget (1993) page 430, Rickart (1960), page 179, Gelfand and Neumark (1943b), page 7

H.3 Normed Algebras

Definition H.6. ¹² Let \mathbf{A} be an algebra.

**D
E
F**

The pair $(\mathbf{A}, \|\cdot\|)$ is a **normed algebra** if

$$\|xy\| \leq \|x\| \|y\| \quad \forall x, y \in \mathbf{A} \quad (\text{multiplicative condition})$$

A normed algebra $(\mathbf{A}, \|\cdot\|)$ is a **Banach algebra** if $(\mathbf{A}, \|\cdot\|)$ is also a Banach space.

Proposition H.2.

**P
R
P**

$(\mathbf{A}, \|\cdot\|)$ is a normed algebra \implies multiplication is **continuous** in $(\mathbf{A}, \|\cdot\|)$

PROOF:

1. Define $f(x) \triangleq zx$. That is, the function f represents multiplication of x times some arbitrary value z .
2. Let $\delta \triangleq \|x - y\|$ and $\epsilon \triangleq \|f(x) - f(y)\|$.
3. To prove that multiplication (f) is *continuous* with respect to the metric generated by $\|\cdot\|$, we have to show that we can always make ϵ arbitrarily small for some $\delta > 0$.
4. And here is the proof that multiplication is indeed continuous in $(\mathbf{A}, \|\cdot\|)$:

$$\begin{aligned} \|f(x) - f(y)\| &\triangleq \|zx - zy\| && \text{by definition of } f && \text{(item (1) page 231)} \\ &= \|z(x - y)\| \\ &\leq \|z\| \|x - y\| && \text{by definition of normed algebra} && \text{(Definition H.6 page 231)} \\ &\triangleq \|z\| \delta && \text{by definition of } \delta && \text{(item (2) page 231)} \\ &\leq \epsilon && \text{for some value of } \delta > 0 \end{aligned}$$

Theorem H.4 (Gelfand-Mazur Theorem). ¹³ Let \mathbb{C} be the field of complex numbers.

**T
H
M**

$(\mathbf{A}, \|\cdot\|)$ is a Banach algebra
every nonzero $x \in \mathbf{A}$ is invertible } $\implies \mathbf{A} \cong \mathbb{C}$ (\mathbf{A} is isomorphic to \mathbb{C})

H.4 C* Algebras

Definition H.7. ¹⁴

**D
E
F**

The triple $(\mathbf{A}, \|\cdot\|, *)$ is a **C* algebra** if

1. $(\mathbf{A}, \|\cdot\|)$ is a Banach algebra and
2. $(\mathbf{A}, *)$ is a $*$ -algebra and
3. $\|x^* x\| = \|x\|^2 \quad \forall x \in \mathbf{A}$

A **C* algebra** $(\mathbf{A}, \|\cdot\|, *)$ is also called a **C star algebra**.

¹² Rickart (1960), page 2, Berberian (1961) page 103 (Theorem IV.9.2)

¹³ Folland (1995) page 4, Mazur (1938) ((statement)), Gelfand (1941) ((proof))

¹⁴ Folland (1995) page 1, Gelfand and Naimark (1964), page 241, Gelfand and Neumark (1943a), Gelfand and Neumark (1943b)

Theorem H.5. ¹⁵ Let A be an algebra.

T
H
M

$$(A, \|\cdot\|, *) \text{ is a } C^* \text{ algebra} \implies \|x^*\| = \|x\|$$

PROOF:

$$\begin{aligned}
 \|x\| &= \frac{1}{\|x\|} \|x\|^2 \\
 &= \frac{1}{\|x\|} \|x^* x\| && \text{by definition of } C^* \text{-algebra} && (\text{Definition H.7 page 231}) \\
 &\leq \frac{1}{\|x\|} \|x^*\| \|x\| && \text{by definition of normed algebra} && (\text{Definition H.6 page 231}) \\
 &= \|x^*\| \\
 \|x^*\| &\leq \|x^{**}\| && \text{by previous result} \\
 &= \|x\| && \text{by involution property of } * && (\text{Definition H.3 page 228})
 \end{aligned}$$

⇒

¹⁵ [Folland \(1995\) page 1](#), [Gelfand and Neumark \(1943b\), page 4](#), [Gelfand and Neumark \(1943a\)](#)

APPENDIX I

INNER PRODUCT SPACES

I.1 Definition and basic results

Definition I.1.¹ Let $\Omega \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}))$ be a LINEAR SPACE (Definition F.1 page 185).

A FUNCTIONAL $\langle \Delta | \nabla \rangle \in \mathbb{F}^{X \times X}$ is an **inner product** on Ω if

- | | | | | |
|-----|--|---|-----------------------|-----|
| DEF | 1. $\langle \alpha x y \rangle = \alpha \langle x y \rangle$ | $\forall x, y \in X, \forall \alpha \in \mathbb{C}$ | (HOMOGENEOUS) | and |
| | 2. $\langle x + y u \rangle = \langle x u \rangle + \langle y u \rangle$ | $\forall x, y, u \in X$ | (ADDITIVE) | and |
| | 3. $\langle x y \rangle = \langle y x \rangle^*$ | $\forall x, y \in X$ | (CONJUGATE SYMMETRIC) | and |
| | 4. $\langle x x \rangle \geq 0$ | $\forall x \in X$ | (NON-NEGATIVE) | and |
| | 5. $\langle x x \rangle = 0 \iff x = 0$ | $\forall x \in X$ | (NON-ISOTROPIC) | |

An inner product is also called a **scalar product**.

The tuple $(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \Delta | \nabla \rangle)$ is called an **inner product space**.

Theorem I.1.² Let $\Omega \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \Delta | \nabla \rangle)$ be a LINEAR SPACE (Definition F.1 page 185).

- | | | |
|-----|--|---|
| THM | 1. $\langle x y + z \rangle = \langle x y \rangle + \langle x z \rangle$ | $\forall x, y, z \in X$ |
| | 2. $\langle x \alpha y \rangle = \alpha^* \langle x y \rangle$ | $\forall x, y \in X, \alpha \in \mathbb{F}$ |
| | 3. $\langle x 0 \rangle = \langle 0 x \rangle = 0$ | $\forall x \in X$ |
| | 4. $\langle x - y z \rangle = \langle x z \rangle - \langle y z \rangle$ | $\forall x, y, z \in X$ |
| | 5. $\langle x y - z \rangle = \langle x y \rangle - \langle x z \rangle$ | $\forall x, y, z \in X$ |
| | 6. $\langle x z \rangle = \langle y z \rangle$ | $\forall z \in X \neq \{0\} \iff x = y$ |
| | 7. $\langle x y \rangle = 0$ | $\forall x \in X \iff y = 0$ |

PROOF:

$$\begin{aligned}
 \langle x | y + z \rangle &= \langle y + z | x \rangle^* && \text{by conjugate symmetric property of } \langle \Delta | \nabla \rangle && \text{(Definition I.1 page 233)} \\
 &= (\langle y | x \rangle + \langle z | x \rangle)^* && \text{by additive property of } \langle \Delta | \nabla \rangle && \text{(Definition I.1 page 233)} \\
 &= \langle y | x \rangle^* + \langle z | x \rangle^* && \text{by distributive property of } * && \text{(Definition H.3 page 228)} \\
 &= \langle x | y \rangle + \langle x | z \rangle && \text{by conjugate symmetric property of } \langle \Delta | \nabla \rangle && \text{(Definition I.1 page 233)} \\
 \langle x | \alpha y \rangle &= \langle \alpha y | x \rangle^* && \text{by conjugate symmetric property of } \langle \Delta | \nabla \rangle && \text{(Definition I.1 page 233)}
 \end{aligned}$$

¹ Istrătescu (1987) page 111 (Definition 4.1.1), Bollobás (1999) pages 130–131, Haaser and Sullivan (1991) page 277, Aliprantis and Burkinshaw (1998), page 276, Peano (1888b) page 72

² Berberian (1961) page 27, Haaser and Sullivan (1991) page 277

$= (\alpha \langle y x \rangle)^*$	by <i>homogeneous</i> property of $\langle \Delta \nabla \rangle$	(Definition I.1 page 233)
$= \alpha^* \langle y x \rangle^*$	by <i>antiautomorphic</i> property of $*$	(Definition H.3 page 228)
$= \alpha^* \langle x y \rangle$	by <i>conjugate symmetric</i> property of $\langle \Delta \nabla \rangle$	(Definition I.1 page 233)
$\langle x 0 \rangle = \langle 0 x \rangle^*$	by <i>conjugate symmetric</i> property of $\langle \Delta \nabla \rangle$	(Definition I.1 page 233)
$= \langle 0 \cdot y x \rangle^*$		
$= (0 \cdot \langle y x \rangle)^*$	by <i>homogeneous</i> property of $\langle \Delta \nabla \rangle$	(Definition I.1 page 233)
$= 0$		
$\langle 0 x \rangle = \langle 0 \cdot y x \rangle$		
$= (0 \cdot \langle y x \rangle)$	by <i>homogeneous</i> property of $\langle \Delta \nabla \rangle$	(Definition I.1 page 233)
$= 0$		
$\langle x - y z \rangle = \langle x + (-y) z \rangle$	by definition of $+$	
$= \langle x z \rangle + \langle -y z \rangle$	by <i>additive</i> property of $\langle \Delta \nabla \rangle$	(Definition I.1 page 233)
$= \langle x z \rangle - \langle y z \rangle$	by <i>homogeneous</i> property of $\langle \Delta \nabla \rangle$	(Definition I.1 page 233)
$\langle x y - z \rangle = \langle y - z x \rangle^*$	by <i>conjugate symmetric</i> property of $\langle \Delta \nabla \rangle$	(Definition I.1 page 233)
$= (\langle y x \rangle - \langle z x \rangle)^*$	by 4.	
$= \langle y x \rangle^* - \langle z x \rangle^*$	by <i>distributive</i> property of $*$	(Definition H.3 page 228)
$= \langle x y \rangle - \langle x z \rangle$	by <i>conjugate symmetric</i> property of $\langle \Delta \nabla \rangle$	(Definition I.1 page 233)
$\langle x z \rangle = \langle y z \rangle$	$\forall z$	
$\iff \langle x z \rangle - \langle y z \rangle = 0$	$\forall z$	by property of complex numbers
$\iff \langle x - y z \rangle = 0$	$\forall z$	by 4.
$\iff x - y = 0$	$\forall z$	by <i>non-isotropic</i> property of $\langle \Delta \nabla \rangle$ (Definition I.1 page 233)

Proof that $\langle x | y \rangle = 0 \implies y = 0$:

1. Suppose $y \neq 0$;
2. Then $\langle y | y \rangle \neq 0$ by the *non-isotropic* property of $\langle \Delta | \nabla \rangle$ (Definition I.1 page 233)
3. But because $y \in X$, the left hypothesis implies that $\langle y | y \rangle = 0$.
4. This is a *contradiction*.
5. Therefore $y \neq 0$ must be incorrect and $y = 0$ must be correct.

Proof that $\langle x | y \rangle = 0 \iff y = 0$:

$$\begin{aligned} \langle x | y \rangle &= \langle x | 0 \rangle && \text{by right hypothesis} \\ &= 0 && \text{by Theorem I.1 page 233} \end{aligned}$$

⇒

One of the most useful and widely used inequalities in analysis is the *Cauchy-Schwarz Inequality* (sometimes also called the *Cauchy-Bunyakovsky-Schwarz Inequality*). In fact, we will use this inequality shortly to prove that every inner product space *has* a norm and therefore every inner product space *is* a normed linear space.

Theorem I.2 (Cauchy-Schwarz Inequality). ³ Let $(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \Delta | \nabla \rangle)$ be an INNER PRODUCT SPACE and $|\cdot| \in \mathbb{R}^{\mathbb{C}}$ an ABSOLUTE VALUE function. Let $\|\cdot\|$ be a function in $\mathbb{R}^{\mathbb{F}}$ such that $\|x\| \triangleq$

³ Haaser and Sullivan (1991) page 278, Aliprantis and Burkinshaw (1998), page 278, Cauchy (1821) page 455, Bunyakovsky (1859) page 6, Schwarz (1885)

$\sqrt{\langle x | x \rangle}$.⁴

T H M	$ \langle x y \rangle ^2 \leq \langle x x \rangle \langle y y \rangle$	$\forall x, y \in X$
	$ \langle x y \rangle ^2 = \langle x x \rangle \langle y y \rangle \iff \exists \alpha \in \mathbb{F} \text{ such that } y = \alpha x \quad \forall x, y \in X$	
	$ \langle x y \rangle \leq \ x\ \ y\ \quad \forall x, y \in X$	
	$ \langle x y \rangle = \ x\ \ y\ \iff \exists \alpha \in \mathbb{F} \text{ such that } y = \alpha x \quad \forall x, y \in X$	

PROOF:

1. Proof that $|\langle x | y \rangle| \leq \|x\| \|y\|$:⁵

(a) $y = \emptyset$ case:

$$\begin{aligned}
 |\langle x | y \rangle|^2 &= |\langle x | \emptyset \rangle|^2 && \text{by } y = \emptyset \text{ hypothesis} \\
 &= |\langle \emptyset | x \rangle|^2 && \text{by Definition I.1 page 233} \\
 &= |\langle 00 | x \rangle|^2 && \text{by Definition F.1 page 185} \\
 &= |0 \langle 0 | x \rangle|^2 && \text{by Definition I.1 page 233} \\
 &= 0 \\
 &= \langle x | x \rangle \langle \emptyset | \emptyset \rangle \\
 &= \langle x | x \rangle \langle y | y \rangle && \text{by } y = \emptyset \text{ hypothesis}
 \end{aligned}$$

(b) $y \neq \emptyset$ case: Let $\lambda \triangleq \frac{\langle x | y \rangle}{\langle y | y \rangle}$.

$$\begin{aligned}
 0 &\leq \langle x - \lambda y | x - \lambda y \rangle && \text{by Definition I.1} \\
 &= \langle x | x - \lambda y \rangle + \langle -\lambda y | x - \lambda y \rangle && \text{by Definition I.1} \\
 &= \langle x - \lambda y | x \rangle^* - \lambda \langle y | x - \lambda y \rangle && \text{by Definition I.1} \\
 &= \langle x | x \rangle^* + \langle -\lambda y | x \rangle^* - \lambda \langle x - \lambda y | y \rangle^* && \text{by Definition I.1} \\
 &= \langle x | x \rangle^* - \lambda^* \langle y | x \rangle^* - \lambda \langle x | y \rangle^* - \lambda \langle -\lambda y | y \rangle^* && \text{by Definition I.1} \\
 &= \langle x | x \rangle - \lambda^* \langle x | y \rangle - \lambda \langle x | y \rangle^* + \lambda \lambda^* \langle y | y \rangle^* && \text{by Definition I.1} \\
 &= \langle x | x \rangle + \left[\frac{\langle x | y \rangle}{\langle y | y \rangle} \lambda^* \langle y | y \rangle - \lambda^* \langle x | y \rangle \right] - \frac{\langle x | y \rangle}{\langle y | y \rangle} \langle x | y \rangle^* && \text{by definition of } \lambda \\
 &= \langle x | x \rangle - \frac{1}{\langle y | y \rangle} |\langle x | y \rangle|^2 \\
 \implies |\langle x | y \rangle|^2 &\leq \langle x | x \rangle \langle y | y \rangle
 \end{aligned}$$

2. Proof that $|\langle x | y \rangle|^2 = \langle x | x \rangle \langle y | y \rangle \iff y = ax$:

Let $\frac{1}{a} \triangleq \lambda \triangleq \frac{\langle x | y \rangle}{\langle y | y \rangle}$. Then...

$$\begin{aligned}
 y &= ax \\
 \iff x &= \lambda y \\
 \iff x - \lambda y &= \emptyset \\
 \iff 0 &= \langle x - \lambda y | x - \lambda y \rangle && \text{by Definition I.1 page 233} \\
 &= \langle x | x - \lambda y \rangle + \langle -\lambda y | x - \lambda y \rangle && \text{by Definition I.1 page 233} \\
 &= \langle x - \lambda y | x \rangle^* - \lambda \langle y | x - \lambda y \rangle && \text{by Definition I.1 page 233} \\
 &\vdots && \text{(same steps as in 1(b))}
 \end{aligned}$$

⁴The function $\|\cdot\|$ is a *norm* (Theorem I.4 page 238) and is called the *norm induced by the inner product* $\langle \Delta | \nabla \rangle$ (Definition I.2 page 238).


⁵  Haaser and Sullivan (1991), page 278

$$\iff |\langle \mathbf{x} | \mathbf{y} \rangle|^2 = \langle \mathbf{x} | \mathbf{x} \rangle \langle \mathbf{y} | \mathbf{y} \rangle$$

$$= \langle \mathbf{x} | \mathbf{x} \rangle - \frac{1}{\langle \mathbf{y} | \mathbf{y} \rangle} |\langle \mathbf{x} | \mathbf{y} \rangle|^2$$

3. Alternate proof for $|\langle \mathbf{x} | \mathbf{y} \rangle| \leq \|\mathbf{x}\| \|\mathbf{y}\|$: (Note: This is essentially the same proof as used by Schwarz).⁶

(a) Proof that $(a\lambda^2 + b\lambda + c \geq 0 \quad \forall \lambda \in \mathbb{R}) \implies b^2 \leq 4ac$ (quadratic discriminant inequality):



Let $k \in (0, \infty)$, and $r_1, r_2 \in \mathbb{C}$ be the roots of $a\lambda^2 + b\lambda + c = 0$. Then

$$\begin{aligned} 0 &\leq a\lambda^2 + b\lambda + c && \text{by left hypothesis} \\ &= k(\lambda - r_1)(\lambda - r_2) && \text{by definition of } r_1 \text{ and } r_2 \\ &= k(\lambda^2 - r_1\lambda - r_2\lambda + r_1r_2) \\ \implies \lambda^2 - r_1\lambda - r_2\lambda + r_1r_2 &\geq 0 \\ \implies r_1 &= r_2^* && \text{because } r_1r_2 \geq 0 \text{ for } \lambda = 0 \end{aligned}$$

The *quadratic equation* places another constraint on r_1 and r_2 :

$$\begin{aligned} \frac{b^2 + \sqrt{b^2 - 4ac}}{2a} &= r_1 && \text{by quadratic equation} \\ &= r_2^* && \text{by previous result} \\ &= \left(\frac{b^2 - \sqrt{b^2 - 4ac}}{2a} \right)^* && \text{by quadratic equation} \end{aligned}$$

The only way for this to be true is if $b^2 \leq 4ac$ (the **discriminate** is non-positive).

(b) Proof that $\langle \mathbf{y} | \mathbf{y} \rangle \lambda^2 + 2|\langle \mathbf{x} | \mathbf{y} \rangle| \lambda + \langle \mathbf{x} | \mathbf{x} \rangle \geq 0 \quad \forall \lambda \in \mathbb{R}$:

$$\begin{aligned} 0 &\leq \langle \mathbf{x} + \alpha \mathbf{y} | \mathbf{x} + \alpha \mathbf{y} \rangle && \text{by Definition I.1 page 233} \\ &= \langle \mathbf{x} | \mathbf{x} + \alpha \mathbf{y} \rangle + \langle \alpha \mathbf{y} | \mathbf{x} + \alpha \mathbf{y} \rangle && \text{by Definition I.1 page 233} \\ &= \langle \mathbf{x} | \mathbf{x} + \alpha \mathbf{y} \rangle + \alpha \langle \mathbf{y} | \mathbf{x} + \alpha \mathbf{y} \rangle && \text{by Definition I.1 page 233} \\ &= \langle \mathbf{x} + \alpha \mathbf{y} | \mathbf{x} \rangle^* + \alpha \langle \mathbf{x} + \alpha \mathbf{y} | \mathbf{y} \rangle^* && \text{by Definition I.1 page 233} \\ &= \langle \mathbf{x} | \mathbf{x} \rangle^* + \langle \alpha \mathbf{y} | \mathbf{x} \rangle^* + \alpha \langle \mathbf{x} | \mathbf{y} \rangle^* + \alpha \langle \alpha \mathbf{y} | \mathbf{y} \rangle^* && \text{by Definition I.1 page 233} \\ &= \langle \mathbf{x} | \mathbf{x} \rangle^* + \alpha^* \langle \mathbf{y} | \mathbf{x} \rangle^* + \alpha \langle \mathbf{x} | \mathbf{y} \rangle^* + \alpha \alpha^* \langle \mathbf{y} | \mathbf{y} \rangle^* && \text{by Definition I.1 page 233} \\ &= \langle \mathbf{x} | \mathbf{x} \rangle + \alpha^* \langle \mathbf{x} | \mathbf{y} \rangle + (\alpha^* \langle \mathbf{x} | \mathbf{y} \rangle)^* + |\alpha|^2 \langle \mathbf{y} | \mathbf{y} \rangle && \text{by Definition I.1 page 233} \\ &= \langle \mathbf{x} | \mathbf{x} \rangle + 2\Re(\alpha^* \langle \mathbf{x} | \mathbf{y} \rangle) + |\alpha|^2 \langle \mathbf{y} | \mathbf{y} \rangle && \text{by Definition I.1 page 233} \\ &\leq \langle \mathbf{x} | \mathbf{x} \rangle + 2|\alpha^* \langle \mathbf{x} | \mathbf{y} \rangle| + |\alpha|^2 \langle \mathbf{y} | \mathbf{y} \rangle && \text{by Definition I.1 page 233} \\ &= \langle \mathbf{x} | \mathbf{x} \rangle + 2|\langle \mathbf{x} | \mathbf{y} \rangle||\alpha| + \langle \mathbf{y} | \mathbf{y} \rangle |\alpha|^2 && \text{by Definition I.1 page 233} \\ &= \langle \mathbf{y} | \mathbf{y} \rangle |\alpha|^2 + 2|\langle \mathbf{x} | \mathbf{y} \rangle| |\alpha| + \langle \mathbf{x} | \mathbf{x} \rangle && \text{by Definition I.1 page 233} \\ &= \underbrace{\langle \mathbf{y} | \mathbf{y} \rangle}_{a} \lambda^2 + \underbrace{2|\langle \mathbf{x} | \mathbf{y} \rangle|}_{b} \lambda + \underbrace{\langle \mathbf{x} | \mathbf{x} \rangle}_{c} && \text{because } \lambda \triangleq |\alpha| \in \mathbb{R} \end{aligned}$$

(c) The above equation is in the quadratic form used in the lemma of part (a).

$$\begin{aligned} \left(\underbrace{2|\langle \mathbf{x} | \mathbf{y} \rangle|}_{b} \right)^2 &\leq 4 \underbrace{\langle \mathbf{y} | \mathbf{y} \rangle}_{a} \underbrace{\langle \mathbf{x} | \mathbf{x} \rangle}_{c} && \text{by the results of parts (a) and (b)} \\ \implies |\langle \mathbf{x} | \mathbf{y} \rangle|^2 &\leq \langle \mathbf{x} | \mathbf{x} \rangle \langle \mathbf{y} | \mathbf{y} \rangle \end{aligned}$$

⁶ [Aliprantis and Burkinshaw \(1998\)](#), page 278, [Steele \(2004\)](#), page 11

4. Proof that $|\langle x | y \rangle| \leq \|x\| \|y\|$:

This follows directly from the definition $\|x\| \triangleq \sqrt{\langle x | x \rangle}$.

5. Proof that $|\langle x | y \rangle| = \|x\| \|y\| \iff \exists \alpha \in \mathbb{C} \text{ such that } y = \alpha x$:

This follows directly from the definition $\|x\| \triangleq \sqrt{\langle x | x \rangle}$.

Corollary I.1. ⁷ Let $(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle | \nabla \rangle)$ be an INNER PRODUCT SPACE.

COR

$\langle x | y \rangle$ is CONTINUOUS in both x and y .

PROOF: Let $\|x\| \triangleq \sqrt{\langle x | x \rangle}$.

$$\begin{aligned} |\langle x + \epsilon | y \rangle - \langle x | y \rangle|^2 &= |\langle x + \epsilon - x | y \rangle|^2 && \text{by additivity of } \langle \triangle | \nabla \rangle && \text{(Definition I.1 page 233)} \\ &= |\langle \epsilon | y \rangle|^2 \\ &\leq \|\epsilon\|^2 \|y\| && \text{by Cauchy-Schwarz Inequality} && \text{(Theorem I.2 page 234)} \end{aligned}$$

I.2 Relationship between norms and inner products

I.2.1 Norms induced by inner products

Lemma I.1 (Polar Identity). ⁸ Let $(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle | \nabla \rangle)$ be an INNER PRODUCT SPACE (Definition I.1 page 233). Let $\Re z$ represent the real part of $z \in \mathbb{C}$. Let $\|\cdot\|$ be a function in $\mathbb{R}^{\mathbb{F}}$ such that $\|x\| \triangleq \sqrt{\langle x | x \rangle}$.⁹

LEM

$$\|x + y\|^2 = \|x\|^2 + 2\Re_e [\langle x | y \rangle] + \|y\|^2 \quad \forall x, y \in X$$

PROOF:

$$\begin{aligned} \|x + y\|^2 &= \langle x + y | x + y \rangle && \text{by definition of induced norm} && \text{(Theorem I.4 page 238)} \\ &= \langle x | x + y \rangle + \langle y | x + y \rangle && \text{by Definition I.1 page 233} \\ &= \langle x + y | x \rangle^* + \langle x + y | y \rangle^* && \text{by Definition I.1 page 233} \\ &= \langle x | x \rangle^* + \langle y | x \rangle^* + \langle x | y \rangle^* + \langle y | y \rangle^* && \text{by Definition I.1 page 233} \\ &= \langle x | x \rangle + \langle x | y \rangle + \langle x | y \rangle^* + \langle y | y \rangle && \text{by definition of inner product} && \text{(Definition I.1 page 233)} \\ &= \|x\|^2 + 2\Re \langle x | y \rangle + \|y\|^2 && \text{by definition of induced norm} && \text{(Theorem I.4 page 238)} \end{aligned}$$

⁷ Bollobás (1999) page 132, Aliprantis and Burkinshaw (1998) page 279 (Lemma 32.4)

⁸ Conway (1990) page 4, Heil (2011) page 27 (Lemma 1.36(a))

⁹ The function $\|\cdot\|$ is a norm (Theorem I.4 page 238) and is called the norm induced by the inner product $\langle \triangle | \nabla \rangle$ (Definition I.2 page 238).

Theorem I.3 (Minkowski's inequality). ¹⁰ Let $(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle | \triangleright \rangle)$ be an INNER PRODUCT SPACE. Let $\|\cdot\|$ be a function in $\mathbb{R}^{\mathbb{F}}$ such that $\|x\| \triangleq \sqrt{\langle x | x \rangle}$.¹¹

T H M	$\ x + y\ \leq \ x\ + \ y\ \quad \forall x, y \in X$
-------------	---

PROOF:

$$\begin{aligned} \|x + y\|^2 &= \|x\|^2 + 2\Re\langle x | y \rangle + \|y\|^2 && \text{by Polar Identity (Lemma I.1 page 237)} \\ &\leq \|x\|^2 + 2|\langle x | y \rangle| + \|y\|^2 \\ &\leq \|x\|^2 + 2\sqrt{\langle x | x \rangle}\sqrt{\langle y | y \rangle} + \|y\|^2 && \text{by Cauchy-Schwarz Inequality (Theorem I.2 page 234)} \\ &= (\|x\| + \|y\|)^2 \end{aligned}$$



Theorem I.4 (induced norm). ¹² Let $\Omega \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle | \triangleright \rangle)$ be an INNER PRODUCT SPACE (Definition I.1 page 233).

T H M	$\ x\ \triangleq \sqrt{\langle x x \rangle} \implies \ \cdot\ \text{ is a NORM}$
-------------	--

PROOF: For a function to be a norm, it must satisfy the four properties listed in Definition J.1 (page 249).

1. Proof that $\|\cdot\|$ is a norm:

- (a) Proof that $\|x\| > 0$ for $x \neq 0$ (non-negative):
By Definition I.1 page 233, all inner products have this property.
- (b) Proof that $\|x\| = 0 \iff x = 0$ (non-isometric):
By Definition I.1, all inner products have this property.
- (c) Prove $\|ax\| = |a| \|x\|$ (homogeneous):

$$\|ax\| \triangleq \sqrt{\langle ax | ax \rangle} = \sqrt{aa^* \langle x | x \rangle} = \sqrt{|a|^2 \langle x | x \rangle} = |a| \|x\|$$

- (d) Proof that $\|x + y\| \leq \|x\| + \|y\|$ (subadditive): This is true by Minkowski's inequality page 238

2. Proof that every inner product space is a normed linear space:

Since every inner product induces a norm, so every inner product space has a norm (the norm induced by the inner product) and is therefore a normed linear space.



Theorem I.4 (previous theorem) demonstrates that in any inner product space $(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle | \triangleright \rangle)$, the function $\|x\| \triangleq \sqrt{\langle x | x \rangle}$ is a norm. That is, $\|x\|$ is the *norm induced by the inner product*. This norm is formally defined next.

Definition I.2. Let $(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle | \triangleright \rangle)$ be an INNER PRODUCT SPACE (Definition I.1 page 233).

D E F	The norm induced by the inner product $\langle \triangle \triangleright \rangle$ is defined as $\ x\ \triangleq \sqrt{\langle x x \rangle}$
-------------	---

¹⁰ Aliprantis and Burkinshaw (1998) pages 278–279 (Theorem 32.3), Maligranda (1995), Minkowski (1910) page 115

¹¹ The function $\|\cdot\|$ is a *norm* (Theorem I.4 page 238) and is called the *normed induced by the inner product $\langle \triangle | \triangleright \rangle$* (Definition I.2 page 238).

¹² Aliprantis and Burkinshaw (1998), pages 278–279, Haaser and Sullivan (1991) page 278

I.2.2 Inner products induced by norms

Theorem I.4 (page 238) demonstrates that if a *linear space* (Definition F.1 page 185) has an *inner product* (Definition I.1 page 233), then that inner product always induces a *norm* (Definition J.1 page 249), and the relationship between the two is simply $\|x\| = \sqrt{\langle x | x \rangle}$ (Definition I.2 page 238). But what about the converse? What if a linear space has a norm—can that norm also induce an inner product? The answer in general is “no”: Not all norms can induce an inner product. But a less harsh answer is “sometimes”: Some norms **can** induce inner products. This leads to some important and interesting questions:

1. How many different inner products can be induced from a single norm? The answer turns out to be **at most one**, but maybe none (Theorem I.5 page 239).
2. When a norm *can* induce an inner product, what is that (unique) inner product? The inner product expressed in terms of the norm is given by the *Polarization Identity* (Theorem I.6 page 240).
3. Which norms can induce an inner product and which ones cannot? The answer is that norms that satisfy the *parallelogram law* (Theorem I.7 page 241) **can** induce an inner product; and the ones that don't, cannot (Theorem I.7 page 241).

Theorem I.5. ¹³ Let $(X, +, \cdot, (\mathbb{F}, +, \dot{\times}), \|\cdot\|)$ be a NORMED LINEAR SPACE (Definition J.1 page 249).

T H M	$\left. \begin{array}{l} \exists \langle \Delta \nabla \rangle \text{ and } (\cdot \cdot) \text{ such that} \\ \ x\ ^2 = \langle x x \rangle = (x x) \quad \forall x \in X \end{array} \right\} \Rightarrow \underbrace{\langle x y \rangle = (x y)}_{\dots \text{then those two inner products are equivalent.}} \quad \forall x, y \in X$ <p>If a norm is induced by two inner products...</p>
-------------	--

PROOF:

$$\begin{aligned}
 2 \langle x | y \rangle &= [\langle x | y \rangle + \langle y | x \rangle] + [\langle x | y \rangle - \langle y | x \rangle] \\
 &= [\langle x | y \rangle + \langle y | x \rangle] + i [-i \langle x | y \rangle + i \langle y | x \rangle] \\
 &= [\langle x | y \rangle + \langle y | x \rangle] + i [-\langle ix | y \rangle - \langle y | ix \rangle] \\
 &= \left(\underbrace{[\langle x | y \rangle + \langle y | x \rangle + \langle x | x \rangle + \langle y | y \rangle]}_{\langle x+y | x+y \rangle} - \underbrace{[\langle x | x \rangle + \langle y | y \rangle]}_{\text{residue}} \right) \\
 &\quad - i \left(\underbrace{[\langle ix | y \rangle + \langle y | ix \rangle + \langle ix | ix \rangle + \langle y | y \rangle]}_{\langle ix+y | ix+y \rangle} - \underbrace{[\langle ix | ix \rangle + \langle y | y \rangle]}_{\text{residue}} \right) \\
 &= ([\langle x+y | x+y \rangle] - [\langle x | x \rangle + \langle y | y \rangle]) - i ([\langle ix+y | ix+y \rangle] - [\langle ix | ix \rangle + \langle y | y \rangle]) \\
 &= ([\langle x+y | x+y \rangle] - [\langle x | x \rangle + \langle y | y \rangle]) - i ([\langle ix+y | ix+y \rangle] - [\langle ix | ix \rangle + \langle y | y \rangle]) \\
 &= \left(\underbrace{[\langle x | y \rangle + \langle y | x \rangle + \langle x | x \rangle + \langle y | y \rangle]}_{\langle x+y | x+y \rangle} - \underbrace{[\langle x | x \rangle + \langle y | y \rangle]}_{\text{residue}} \right) \\
 &\quad - i \left(\underbrace{[\langle ix | y \rangle + \langle y | ix \rangle + \langle ix | ix \rangle + \langle y | y \rangle]}_{\langle ix+y | ix+y \rangle} - \underbrace{[\langle ix | ix \rangle + \langle y | y \rangle]}_{\text{residue}} \right)
 \end{aligned}$$

¹³ Aliprantis and Burkinshaw (1998), page 280, Bollobás (1999), page 132, Jordan and von Neumann (1935) page 721

$$\begin{aligned}
 &= [(\mathbf{x} | \mathbf{y}) + (\mathbf{y} | \mathbf{x})] + i [-(i\mathbf{x} | \mathbf{y}) - (\mathbf{y} | i\mathbf{x})] \\
 &= [(\mathbf{x} | \mathbf{y}) + (\mathbf{y} | \mathbf{x})] + i [-i(\mathbf{x} | \mathbf{y}) + i(\mathbf{y} | \mathbf{x})] \\
 &= [(\mathbf{x} | \mathbf{y}) + (\mathbf{y} | \mathbf{x})] + [(\mathbf{x} | \mathbf{y}) - (\mathbf{y} | \mathbf{x})] \\
 &= 2(\mathbf{x} | \mathbf{y})
 \end{aligned}$$

⇒

Theorem I.6 (Polarization Identities). ¹⁴ Let $\Omega \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}))$ be a linear space, $\langle \Delta | \nabla \rangle \in \mathbb{F}^{X \times X}$ a function, and $\|\mathbf{x}\| \triangleq \sqrt{\langle \mathbf{x} | \mathbf{x} \rangle}$.

THM

$(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \Delta | \nabla \rangle)$ is an inner product space \Rightarrow

$$4 \langle \mathbf{x} | \mathbf{y} \rangle = \underbrace{\begin{cases} \|\mathbf{x} + \mathbf{y}\|^2 - \|\mathbf{x} - \mathbf{y}\|^2 + i \|\mathbf{x} + i\mathbf{y}\|^2 - i \|\mathbf{x} - i\mathbf{y}\|^2 & \text{for } \mathbb{F} = \mathbb{C} \quad \forall \mathbf{x}, \mathbf{y} \in X \\ \|\mathbf{x} + \mathbf{y}\|^2 - \|\mathbf{x} - \mathbf{y}\|^2 & \text{for } \mathbb{F} = \mathbb{R} \quad \forall \mathbf{x}, \mathbf{y} \in X \end{cases}}_{\text{inner product induced by norm}}$$

⇒


PROOF:

1. These follow directly from properties of *bilinear functionals* (Theorem ?? page ??).

2. Alternative proof for $\mathbb{F} = \mathbb{C}$ case:

$$\begin{aligned}
 &\|\mathbf{x} + \mathbf{y}\|^2 - \|\mathbf{x} - \mathbf{y}\|^2 + i \|\mathbf{x} + i\mathbf{y}\|^2 - i \|\mathbf{x} - i\mathbf{y}\|^2 \\
 &= \underbrace{\|\mathbf{x}\|^2 + \|\mathbf{y}\|^2 + 2\Re \langle \mathbf{x} | \mathbf{y} \rangle}_{\langle \mathbf{x} + \mathbf{y} | \mathbf{x} + \mathbf{y} \rangle} - \underbrace{(\|\mathbf{x}\|^2 + \|\mathbf{y}\|^2 + 2\Re \langle \mathbf{x} | -\mathbf{y} \rangle)}_{\langle \mathbf{x} - \mathbf{y} | \mathbf{x} - \mathbf{y} \rangle} \\
 &\quad + i \underbrace{(\|\mathbf{x}\|^2 + \|\mathbf{y}\|^2 + 2\Re \langle \mathbf{x} | i\mathbf{y} \rangle)}_{i \langle \mathbf{x} + i\mathbf{y} | \mathbf{x} + i\mathbf{y} \rangle} - i \underbrace{(\|\mathbf{x}\|^2 + \|\mathbf{y}\|^2 + 2\Re \langle \mathbf{x} | -i\mathbf{y} \rangle)}_{i \langle \mathbf{x} - i\mathbf{y} | \mathbf{x} - i\mathbf{y} \rangle} \quad \text{by Lemma I.1 page 237} \\
 &= \underbrace{\|\mathbf{x}\|^2 + \|\mathbf{y}\|^2 + 2\Re \langle \mathbf{x} | \mathbf{y} \rangle}_{\langle \mathbf{x} + \mathbf{y} | \mathbf{x} + \mathbf{y} \rangle} - \underbrace{(\|\mathbf{x}\|^2 + \|\mathbf{y}\|^2 - 2\Re \langle \mathbf{x} | \mathbf{y} \rangle)}_{\langle \mathbf{x} - \mathbf{y} | \mathbf{x} - \mathbf{y} \rangle} \\
 &\quad + i \underbrace{(\|\mathbf{x}\|^2 + \|\mathbf{y}\|^2 + 2\Re \langle \mathbf{x} | i\mathbf{y} \rangle)}_{i \langle \mathbf{x} + i\mathbf{y} | \mathbf{x} + i\mathbf{y} \rangle} - i \underbrace{(\|\mathbf{x}\|^2 + \|\mathbf{y}\|^2 - 2\Re \langle \mathbf{x} | i\mathbf{y} \rangle)}_{i \langle \mathbf{x} - i\mathbf{y} | \mathbf{x} - i\mathbf{y} \rangle} \quad \text{by Definition I.1 page 233} \\
 &= 4\Re \langle \mathbf{x} | \mathbf{y} \rangle + 4i\Re \langle \mathbf{x} | i\mathbf{y} \rangle \\
 &= 2 \underbrace{(\langle \mathbf{x} | \mathbf{y} \rangle + \langle \mathbf{x} | \mathbf{y} \rangle^*)}_{4\Re \langle \mathbf{x} | \mathbf{y} \rangle} + 2i \underbrace{(\langle \mathbf{x} | i\mathbf{y} \rangle + \langle \mathbf{x} | i\mathbf{y} \rangle^*)}_{4i\Re \langle \mathbf{x} | i\mathbf{y} \rangle} \\
 &= 2(\langle \mathbf{x} | \mathbf{y} \rangle + \langle \mathbf{x} | \mathbf{y} \rangle^*) + 2i(i^* \langle \mathbf{x} | \mathbf{y} \rangle + (i^{**}) \langle \mathbf{x} | \mathbf{y} \rangle^*) \\
 &= 2(\langle \mathbf{x} | \mathbf{y} \rangle + \langle \mathbf{x} | \mathbf{y} \rangle^*) + 2i(-i \langle \mathbf{x} | \mathbf{y} \rangle + i \langle \mathbf{x} | \mathbf{y} \rangle^*) \quad \text{by Definition I.1 page 233} \\
 &= 2\langle \mathbf{x} | \mathbf{y} \rangle + 2\langle \mathbf{x} | \mathbf{y} \rangle^* + 2\langle \mathbf{x} | \mathbf{y} \rangle - 2\langle \mathbf{x} | \mathbf{y} \rangle^* \\
 &= 4\langle \mathbf{x} | \mathbf{y} \rangle
 \end{aligned}$$

⇒



In plane geometry (\mathbb{R}^2), the *parallelogram law* states that the sum of the squares of the lengths of the sides of a parallelogram is equal to the sum of the squares of the lengths of its diagonals. This is illustrated in the figure to the left.

¹⁴ Berberian (1961) pages 29–30 (Theorem II.3.3), Istrătescu (1987) page 110 (Proposition 4.1.5), Bollobás (1999), page 132, Jordan and von Neumann (1935) page 721

Acutally, the parallelogram law can be generalized to *any inner product space* (not just in the plane). And if the parallelogram law happens to hold true in a normed linear space, then that normed linear space is actually an *inner product space*. The parallelogram law and its relation to inner product spaces is stated in the next theorem.

Theorem I.7 (Parallelogram law). ¹⁵ Let $\Omega \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle | \triangleright \rangle)$ and $\|x\| \triangleq \sqrt{\langle x | x \rangle}$.

T H M	Ω is an inner product space $\iff \underbrace{2\ x\ ^2 + 2\ y\ ^2 = \ x+y\ ^2 + \ x-y\ ^2}_{\text{PARALLELOGRAM LAW / VON NEUMANN-JORDAN CONDITION}}$ $\forall x, y \in \Omega$
-------------	--

PROOF:

1. Proof that $\exists \langle x | y \rangle$ such that $\|x\|^2 = \langle x | x \rangle \implies$ [parallelogram law is true]:

$$\begin{aligned} \|x+y\|^2 + \|x-y\|^2 &= [\|x\|^2 + \|y\|^2 + 2R_e[2\langle x | y \rangle]] + [\|x\|^2 + \|y\|^2 + 2R_e[2\langle x | -y \rangle]] \\ &\quad \text{by Lemma I.1 page 237} \\ &= [\|x\|^2 + \|y\|^2 + 2R_e[2\langle x | y \rangle]] + [\|x\|^2 + \|y\|^2 - 2R_e[2\langle x | y \rangle]] \\ &= 2\|x\|^2 + 2\|y\|^2 \end{aligned}$$

2. Proof that $\exists \langle x | y \rangle$ such that $\|x\|^2 = \langle x | x \rangle \iff$ [parallelogram law is true]:

Note that if an inner product exists in the norm linear space $(\Omega, \|\cdot\|)$, then that norm linear space is actually an inner product space $(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle | \triangleright \rangle)$. And if it is an inner product space, then by Theorem I.6 page 240 that inner product must be given by the **Polarization Identity**

$$\langle x | y \rangle = \|ax + y\|^2 - \|ax - y\|^2 + i\|ax + iy\|^2 - i\|ax - iy\|^2.$$

Therefore, here we must use the parallelogram law to show that the bilinear function $f(x, y) \triangleq \langle x | y \rangle$ given on the left hand side of the “=” relation is indeed an inner product—that is, that it satisfies the requirements of Definition I.1 page 233.

(a) Proof that $\langle x | x \rangle \geq 0$ (non-negative):

$$\begin{aligned} 4\langle x | x \rangle &\triangleq \|x+x\|^2 - \|x-x\|^2 + 0 \quad \text{by Polarization Identity} \\ &= \|2x\|^2 - 0 + i(\|x+ix\|^2 - \|x-ix\|^2) \quad \text{by Definition J.1 page 249} \\ &= |2|^2\|x\|^2 + i(\|x+ix\|^2 - |i|\|x-ix\|^2) \\ &= 4\|x\|^2 + i(\|x+ix\|^2 - \|ix+x\|^2) \quad \text{by Definition J.1 page 249} \\ &= 4\|x\|^2 \quad \text{by Definition J.1 page 249} \\ &\geq 0 \end{aligned}$$

(b) Proof that $\langle x | x \rangle = 0 \iff x = \emptyset$ (non-isotropic):

$$\begin{aligned} 4\langle x | x \rangle &= 4\|x\|^2 && \text{by result of part (a)} \\ &= 0 &\iff x = \emptyset & \text{by Definition J.1 page 249} \end{aligned}$$

¹⁵ Amir (1986), page 8, Istrătescu (1987) page 110, Day (1973), page 151, Halmos (1998), page 14, Aliprantis and Burkinshaw (1998) pages 280–281 (Theorem 32.6), Riesz (1934) page 36?, Jordan and von Neumann (1935) pages 721–722

(c) Proof that $\langle \mathbf{x} + \mathbf{u} | \mathbf{y} \rangle = \langle \mathbf{x} | \mathbf{y} \rangle + \langle \mathbf{u} | \mathbf{y} \rangle$ (additive):¹⁶

$$\begin{aligned}
4 \langle \mathbf{x} + \mathbf{y} | \mathbf{z} \rangle &= 8 \left\langle \frac{\mathbf{x} + \mathbf{y}}{2} | \mathbf{z} \right\rangle && \text{by Definition I.1 page 233} \\
&= 2 \left\| \frac{\mathbf{x} + \mathbf{y}}{2} + \mathbf{z} \right\|^2 - 2 \left\| \frac{\mathbf{x} + \mathbf{y}}{2} - \mathbf{z} \right\|^2 \\
&\quad + 2i \left\| \frac{\mathbf{x} + \mathbf{y}}{2} + \mathbf{z} \right\|^2 - 2i \left\| \frac{\mathbf{x} + \mathbf{y}}{2} - i\mathbf{z} \right\|^2 && \text{by } \textit{Polarization Identity} \\
&= \left(2 \left\| \frac{\mathbf{x} + \mathbf{y}}{2} + \mathbf{z} \right\|^2 + 2 \left\| \frac{\mathbf{x} - \mathbf{y}}{2} \right\|^2 \right) \\
&\quad - \left(2 \left\| \frac{\mathbf{x} + \mathbf{y}}{2} - \mathbf{z} \right\|^2 + 2 \left\| \frac{\mathbf{x} - \mathbf{y}}{2} \right\|^2 \right) \\
&\quad + \left(2i \left\| \frac{\mathbf{x} + \mathbf{y}}{2} + \mathbf{z} \right\|^2 + 2i \left\| \frac{\mathbf{x} - \mathbf{y}}{2} \right\|^2 \right) \\
&\quad - \left(2i \left\| \frac{\mathbf{x} + \mathbf{y}}{2} - i\mathbf{z} \right\|^2 + 2i \left\| \frac{\mathbf{x} - \mathbf{y}}{2} \right\|^2 \right) \\
&= (\|\mathbf{x} + \mathbf{z}\|^2 + \|\mathbf{y} + \mathbf{z}\|^2) - (\|\mathbf{x} - \mathbf{z}\|^2 + \|\mathbf{y} - \mathbf{z}\|^2) \\
&\quad + (i\|\mathbf{x} + \mathbf{z}\|^2 + i\|\mathbf{y} + \mathbf{z}\|^2) - (i\|\mathbf{x} - i\mathbf{z}\|^2 + i\|\mathbf{y} - i\mathbf{z}\|^2) && \text{by parallelogram law} \\
&= (\|\mathbf{x} + \mathbf{z}\|^2 - \|\mathbf{x} - \mathbf{z}\|^2 + i\|\mathbf{x} + \mathbf{z}\|^2 - i\|\mathbf{x} - i\mathbf{z}\|^2) \\
&\quad + (\|\mathbf{y} + \mathbf{z}\|^2 - \|\mathbf{y} - \mathbf{z}\|^2 + i\|\mathbf{y} + \mathbf{z}\|^2 - i\|\mathbf{y} - i\mathbf{z}\|^2) \\
&= 4 \langle \mathbf{x} | \mathbf{z} \rangle + 4 \langle \mathbf{y} | \mathbf{z} \rangle && \text{by } \textit{Polarization Identity}
\end{aligned}$$

(d) Proof that $\langle \mathbf{x} | \mathbf{y} \rangle = \langle \mathbf{y} | \mathbf{y} \rangle^*$ (*conjugate symmetric*):

$$\begin{aligned}
4 \langle \mathbf{x} | \mathbf{y} \rangle &\triangleq \|\mathbf{x} + \mathbf{y}\|^2 - \|\mathbf{x} - \mathbf{y}\|^2 + i\|\mathbf{x} + i\mathbf{y}\|^2 - i\|\mathbf{x} - i\mathbf{y}\|^2 && \text{by } \textit{Polarization Identity} \\
&= \|\mathbf{y} + \mathbf{x}\|^2 - \|\mathbf{y} - \mathbf{x}\|^2 + i\|i(\mathbf{y} - i\mathbf{x})\|^2 - i\|-i(\mathbf{y} + i\mathbf{x})\|^2 && \text{by Definition F.1 page 185} \\
&= \|\mathbf{y} + \mathbf{x}\|^2 - \|\mathbf{y} - \mathbf{x}\|^2 + i\|\mathbf{y} - i\mathbf{x}\|^2 - i\|\mathbf{y} + i\mathbf{x}\|^2 && \text{by Definition J.1 page 249} \\
&= (\|\mathbf{y} + \mathbf{x}\|^2 - \|\mathbf{y} - \mathbf{x}\|^2 - i\|\mathbf{y} - i\mathbf{x}\|^2 + i\|\mathbf{y} + i\mathbf{x}\|^2)^* \\
&= (\|\mathbf{y} + \mathbf{x}\|^2 - \|\mathbf{y} - \mathbf{x}\|^2 + i\|\mathbf{y} + i\mathbf{x}\|^2 - i\|\mathbf{y} - i\mathbf{x}\|^2)^* \\
&\triangleq 4 \langle \mathbf{y} | \mathbf{x} \rangle^* && \text{by } \textit{Polarization Identity}
\end{aligned}$$

(e) Proof that $\langle \alpha\mathbf{x} | \mathbf{y} \rangle = \alpha \langle \mathbf{x} | \mathbf{y} \rangle$ (*homogeneous*):¹⁷

i. Proof that $\langle \alpha\mathbf{x} | \mathbf{y} \rangle$ is linear in α :

$$\begin{aligned}
0 &\leq \|\alpha\mathbf{x} + \mathbf{y}\| - \|\beta\mathbf{x} + \mathbf{y}\| \\
&\leq \|(\alpha\mathbf{x} + \mathbf{y}) - (\beta\mathbf{x} + \mathbf{y})\| && \text{by Theorem J.2 page 250} \\
&\leq \|(\alpha - \beta)\mathbf{x}\|
\end{aligned}$$

This implies that as $\alpha \rightarrow \beta$, $\|\alpha\mathbf{x} + \mathbf{y}\| \rightarrow \|\beta\mathbf{x} + \mathbf{y}\|$, which by definition implies that $\|\alpha\mathbf{x} + \mathbf{y}\|$ linear in α . And by the parallelogram law, $\langle \alpha\mathbf{x} | \mathbf{y} \rangle$ is also linear in α .

ii. Proof that $\langle n\mathbf{x} | \mathbf{y} \rangle = n \langle \mathbf{x} | \mathbf{y} \rangle$ for $n \in \mathbb{Z}$ (integer case):

A. Proof for $n = \pm 1$:

$$\begin{aligned}
\langle n\mathbf{x} | \mathbf{y} \rangle &= \langle \pm 1\mathbf{x} | \mathbf{y} \rangle && \text{by } n = \pm 1 \text{ hypothesis} \\
&= \pm 1 \langle \mathbf{x} | \mathbf{y} \rangle && \text{by definition of } \textit{inner product} && \text{(Definition I.1 page 233)} \\
&= n \langle \mathbf{x} | \mathbf{y} \rangle && \text{by } n = \pm 1 \text{ hypothesis}
\end{aligned}$$

¹⁶ Aliprantis and Burkinshaw (1998), page 281

¹⁷ Aliprantis and Burkinshaw (1998), page 138

B. Proof for $n = 0$:

$$\begin{aligned}
 \langle nx | y \rangle &= \langle 0x | y \rangle && \text{by } n = 0 \text{ hypothesis} \\
 &= \langle x - x | y \rangle \\
 &= \langle x | y \rangle + \langle -1x | y \rangle \\
 &= \langle x | y \rangle - 1 \langle x | y \rangle \\
 &= \langle x | y \rangle - \langle x | y \rangle \\
 &= 0 \langle x | y \rangle \\
 &= n \langle x | y \rangle && \text{by } n = 0 \text{ hypothesis}
 \end{aligned}$$

C. Proof for $n = \pm 2$:

$$\begin{aligned}
 \langle nx | y \rangle &= \langle \pm 2x | y \rangle && \text{by } n = \pm 1 \text{ hypothesis} \\
 &= \langle \pm(x + x) | y \rangle \\
 &= \pm \langle x + x | y \rangle && \text{by definition of inner product} \quad (\text{Definition I.1 page 233}) \\
 &= \pm (\langle x | y \rangle + \langle x | y \rangle) && \text{by additive property} \\
 &= \pm 2 \langle x | y \rangle \\
 &= n \langle x | y \rangle && \text{by } n = \pm 2 \text{ hypothesis}
 \end{aligned}$$

D. Proof that $[n \text{ case}] \implies [n \pm 1 \text{ case}]$:

$$\begin{aligned}
 \langle (n \pm 1)x | y \rangle &= \langle nx \pm 1x | y \rangle \\
 &= \langle nx | y \rangle + \langle \pm 1x | y \rangle && \text{by additive property} \\
 &= n \langle x | y \rangle \pm 1 \langle x | y \rangle && \text{by left hypothesis} \\
 &= (n \pm 1) \langle x | y \rangle
 \end{aligned}$$

iii. Proof that $\langle qx | y \rangle = q \langle x | y \rangle$ for $q \in \mathbb{Q}$ (rational number case):

$$\begin{aligned}
 \frac{n}{m} \langle x | y \rangle &= \frac{n}{m} \left\langle \frac{m}{m} x | y \right\rangle && \text{where } n, m \in \mathbb{Z} \text{ and } m \neq 0 \\
 &= \frac{nm}{m} \left\langle \frac{1}{m} x | y \right\rangle && \text{by previous result} \\
 &= \frac{m}{m} \left\langle \frac{n}{m} x | y \right\rangle && \text{by previous result} \\
 &= \left\langle \frac{n}{m} x | y \right\rangle
 \end{aligned}$$

iv. Proof that $\langle rx | y \rangle = r \langle x | y \rangle$ for all $r \in \mathbb{R}$ (real number case):

Because \mathbb{Q} is dense in \mathbb{R} and because $\|\alpha x + y\|$ is continuous in α , so $\langle \alpha x | y \rangle = \alpha \langle x | y \rangle$ for all $\alpha \in \mathbb{R}$.

v. Proof that $\langle cx | y \rangle = c \langle x | y \rangle$ for all $c \in \mathbb{C}$ (complex number case):

No proof at this time.



Remark I.1. ¹⁸ The inner product has already been defined in Definition I.1 (page 233) as a bilinear function that is *non-negative, non-isotropic, homogeneous, additive, and conjugate symmetric*. However, given a normed linear space, we could alternatively define the inner product using the *parallelogram law* (Theorem I.7 page 241) together with the *Polarization Identity* (Theorem I.6 page 240). Under this new definition, an inner product *exists* if the parallelogram law is satisfied, and is *specified*, in terms of the norm, by the Polarization Identity.

¹⁸ Loomis (1953), pages 23–24, Kubrusly (2001) page 317

Of the uncountably infinite number of $\ell_{\mathbb{F}}^p$ norms, only the norm for $p = 2$ induces an inner product (Proposition I.1, next).

Proposition I.1. ¹⁹ Let $\|(x_n)_{n \in \mathbb{Z}}\|_p$ be the $\ell_{\mathbb{F}}^p$ norm of the sequence (x_n) in the space $\ell_{\mathbb{F}}^p$.

P R P	$\ (x_n)\ _p$ induces an inner product $\iff p = 2$
-------------	---

PROOF:

1. Proof that $\|\cdot\|_p$ induces an inner product $\iff p = 2$ (using the *Parallelogram law* page 241):

$$\begin{aligned}
 & \|x + y\|_p^2 + \|x - y\|_p^2 \\
 &= \|x + y\|_2^2 + \|x - y\|_2^2 && \text{by right hypothesis} \\
 &= \left(\sum_{n \in \mathbb{Z}} |x_n + y_n|^2 \right)^{\frac{2}{p}} + \left(\sum_{n \in \mathbb{Z}} |x_n - y_n|^2 \right)^{\frac{2}{p}} && \text{by definition of } \|\cdot\|_p \\
 &= \sum_{n \in \mathbb{Z}} (x_n + y_n)(x_n + y_n)^* + \sum_{n \in \mathbb{Z}} (x_n - y_n)(x_n - y_n)^* \\
 &= \sum_{n \in \mathbb{Z}} \left(|x_n|^2 + |y_n|^2 + 2\Re(x_n y_n) \right) + \sum_{n \in \mathbb{Z}} \left(|x_n|^2 + |y_n|^2 - 2\Re(x_n y_n) \right) \\
 &= 2 \sum_{n \in \mathbb{Z}} |x_n|^2 + 2 \sum_{n \in \mathbb{Z}} |y_n|^2 \\
 &= 2 \|x\|_2^2 + 2 \|y\|_2^2 && \text{by definition of } \|\cdot\|_p \\
 &= 2 \|x\|_p^2 + 2 \|y\|_p^2 && \text{by right hypothesis} \\
 &\implies \|\cdot\|_2 \text{ induces an inner product} && \text{by Theorem I.7 page 241}
 \end{aligned}$$

2. Proof that $\|\cdot\|_p$ induces an inner product $\implies p = 2$:

(a) Let $x \triangleq (1, 0)$ and $y \triangleq (0, 1)$. Then ²⁰


$$\begin{aligned}
 \|x + y\|_p^2 + \|x - y\|_p^2 &= \left(\sum_{n \in \mathbb{Z}} |x_n + y_n|^p \right)^{\frac{2}{p}} + \left(\sum_{n \in \mathbb{Z}} |x_n - y_n|^p \right)^{\frac{2}{p}} && \text{by definition of } \|\cdot\|_p \\
 &= (|1+0|^p + |0+1|^p)^{\frac{2}{p}} + (|1-0|^p + |0-1|^p)^{\frac{2}{p}} && \text{by definitions of } x \text{ and } y \\
 &= 2^{\frac{2}{p}} + 2^{\frac{2}{p}} \\
 &= 2 \cdot 2^{\frac{2}{p}} \\
 2 \|x\|_p^2 + 2 \|y\|_p^2 &= 2 \left(\sum_{n \in \mathbb{Z}} |x_n|^p \right)^{\frac{2}{p}} + 2 \left(\sum_{n \in \mathbb{Z}} |y_n|^p \right)^{\frac{2}{p}} && \text{by definition of } \|\cdot\|_p \\
 &= 2(|1|^p + |0|^p)^{\frac{2}{p}} + 2(|1|^p + |0|^p)^{\frac{2}{p}} && \text{by definitions of } x \text{ and } y \\
 &= 2 + 2 \\
 &= 4 \\
 2 \cdot 2^{\frac{2}{p}} = 4 &= \implies 2^{\frac{2}{p}} = 2 \\
 &= \implies p = 2
 \end{aligned}$$

¹⁹  Kubrusly (2001) pages 318–319 (Example 5B)

²⁰ <http://groups.google.com/group/sci.math/msg/531b1173f08871e9>

(b) Proof that $2^{2/p}$ is monotonic decreasing in p (and so $p = 2$ is the only solution):

$$\begin{aligned}\frac{d}{dp} 2^{\frac{2}{p}} &= \frac{d}{dp} e^{\ln 2^{\frac{2}{p}}} \\ &= \left(e^{\ln 2^{\frac{2}{p}}} \right) \frac{d}{dp} \ln 2^{\frac{2}{p}} \\ &= \left(2^{\frac{2}{p}} \right) \frac{d}{dp} (2 \ln 2) \frac{1}{p} \\ &= \left(2^{\frac{2}{p}} \right) 2 \ln 2 \left(-\frac{1}{p^2} \right) \\ &< 0 \quad \forall p \in (0, \infty)\end{aligned}$$



I.3 Orthogonality

Definition I.3.

D E F The Kronecker delta function δ_n is defined as $\delta_n \triangleq \begin{cases} 1 & \text{for } n = 0 \quad \text{and} \\ 0 & \text{for } n \neq 0: \end{cases} \quad \forall n \in \mathbb{Z}$

Definition I.4. Let $(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \Delta | \nabla \rangle)$ be an INNER PRODUCT SPACE (Definition I.1 page 233).

Two vectors x and y in X are **orthogonal** if

$$\langle x | y \rangle = \begin{cases} 0 & \text{for } x \neq y \\ c \in \mathbb{F} \setminus 0 & \text{for } x = y \end{cases}$$

The notation $x \perp y$ implies x and y are **ORTHOGONAL**.

A set $Y \in \mathcal{P}^X$ is **orthogonal** if $x \perp y \quad \forall x, y \in Y$.

A set Y is **orthonormal** if it is ORTHOGONAL and $\langle y | y \rangle = 1 \quad \forall y \in Y$.

A sequence $(x_n \in X)_{n \in \mathbb{Z}}$ is **orthogonal** if $\langle x_n | x_m \rangle = c \delta_{nm}$ for some $c \in \mathbb{R} \setminus 0$.

A sequence $(x_n \in X)_{n \in \mathbb{Z}}$ is **orthonormal** if $\langle x_n | x_m \rangle = \delta_{nm}$.

The definition of the orthogonality relation \perp has several immediate consequences (next theorem):

Theorem I.8.²¹ Let $(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \Delta | \nabla \rangle)$ be an INNER PRODUCT SPACE.

- T H M**
1. $x \perp x \iff x = 0 \quad \forall x \in X$
 2. $x \perp y \implies \alpha x \perp y \quad \forall \alpha \in \mathbb{R}, x, y \in X \quad (\text{homogeneous})$
 3. $x \perp y \iff y \perp x \quad \forall x, y \in X \quad (\text{symmetry})$
 4. $x \perp y \text{ and } y \perp z \implies x \perp (y + z) \quad \forall x, y, z \in X \quad (\text{additive})$
 5. $\exists \beta \in \mathbb{R} \text{ such that } x \perp (\beta x + y) \quad \forall x \in X \setminus 0, y \in X$

Theorem I.9. Let $(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \Delta | \nabla \rangle)$ be an INNER PRODUCT SPACE.

- T H M**
1. $\langle x | y \rangle = 0 \quad \text{and}$
 2. $x + y = 0$
- $$\iff \begin{cases} 1. \quad x = 0 \quad \text{and} \\ 2. \quad y = 0 \end{cases} \quad \forall x, y \in X$$

PROOF:

²¹ James (1945), page 292, Drljević (1989) page 232

1. Proof that $x = y = \emptyset$:

$$\begin{aligned}
 0 &= \langle \emptyset | \emptyset \rangle && \text{by non-isotropic property of } \langle \triangle | \nabla \rangle \text{ (Definition I.1 page 233)} \\
 &= \langle x + y | x + y \rangle && \text{by left hypothesis 2} \\
 &= \langle x | x + y \rangle + \langle y | x + y \rangle && \text{by additive property of } \langle \triangle | \nabla \rangle \text{ (Definition I.1 page 233)} \\
 &= \langle x | x \rangle + \langle x | y \rangle + \langle x | y \rangle^* + \langle y | y \rangle && \text{by conjugate symmetric and additive properties of } \langle \triangle | \nabla \rangle \\
 &= \underbrace{\langle x | x \rangle}_{\geq 0} + 0 + 0 + \underbrace{\langle y | y \rangle}_{\geq 0} && \text{by left hypothesis 1} \\
 \implies x &= \emptyset \text{ and } y = \emptyset && \text{by non-negative and non-isotropic properties of } \langle \triangle | \nabla \rangle
 \end{aligned}$$

2. Proof that $\langle x | y \rangle = 0$:

$$\begin{aligned}
 \langle x | y \rangle &= \langle \emptyset | \emptyset \rangle && \text{by right hypotheses} \\
 &= 0 && \text{by non-isotropic property of } \langle \triangle | \nabla \rangle \text{ (Definition I.1 page 233)}
 \end{aligned}$$

3. Proof that $x + y = \emptyset$:

$$\begin{aligned}
 x + y &= \emptyset + \emptyset && \text{by right hypotheses} \\
 &= \emptyset
 \end{aligned}$$



The *triangle inequality* theorem for vectors in a *normed linear space* (Theorem J.1 page 249) demonstrates that

$\left\| \sum_{n=1}^N x_n \right\| \leq \sum_{n=1}^N \|x_n\|$. The *Pythagorean Theorem* (next) demonstrates that this *inequality* becomes *equality* when the set $\{x_n\}$ is orthogonal.

Theorem I.10 (Pythagorean Theorem). ²² Let $\{x_n \in X | n=1,2,\dots,N\}$ be a set of vectors in an INNER PRODUCT SPACE (Definition I.1 page 233) ($X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle | \nabla \rangle$) and let $\|x\| \triangleq \sqrt{\langle x | x \rangle}$ (Definition I.2 page 238).

THEM	$\{x_n\}$ is ORTHOGONAL	$\iff \left\ \sum_{n=1}^N x_n \right\ ^2 = \sum_{n=1}^N \ x_n\ ^2 \quad \forall N \in \mathbb{N}$
------	-------------------------	--

PROOF: 1. Proof for (\implies) case:

$$\begin{aligned}
 \left\| \sum_{n=1}^N x_n \right\|^2 &= \left\langle \sum_{n=1}^N x_n | \sum_{m=1}^N x_m \right\rangle && \text{by def. of } \|\cdot\| && \text{(Definition J.1 page 249)} \\
 &= \sum_{n=1}^N \sum_{m=1}^N \langle x_n | x_m \rangle && \text{by def. of } \langle \triangle | \nabla \rangle && \text{(Definition I.1 page 233)} \\
 &= \sum_{n=1}^N \sum_{m=1}^N \langle x_n | x_m \rangle \delta_{n-m} && \text{by left hypothesis} \\
 &= \sum_{n=1}^N \langle x_n | x_n \rangle && \text{by def. of } \bar{\delta} && \text{(Definition I.3 page 245)} \\
 &= \sum_{n=1}^N \|x_n\|^2 && \text{by def. of } \|\cdot\| && \text{(Definition J.1 page 249)}
 \end{aligned}$$

²² Aliprantis and Burkinshaw (1998) pages 282–283 (Theorem 32.7), Kubrusly (2001) page 324 (Proposition 5.8), Bollobás (1999) pages 132–133 (Theorem 3)

2. Proof for (\Leftarrow) case:

$$\begin{aligned} 4 \langle \mathbf{x} | \mathbf{y} \rangle &= \| \mathbf{x} + \mathbf{y} \|^2 - \| \mathbf{x} - \mathbf{y} \|^2 + i \| \mathbf{x} + i\mathbf{y} \|^2 - i \| \mathbf{x} - i\mathbf{y} \|^2 \quad \text{by polarization identity (Theorem I.6 page 240)} \\ &= (\| \mathbf{x} \|^2 + \| \mathbf{y} \|^2) - (\| \mathbf{x} \|^2 + \| -\mathbf{y} \|^2) + i (\| \mathbf{x} \|^2 + \| i\mathbf{y} \|^2) - i (\| \mathbf{x} \|^2 + \| -i\mathbf{y} \|^2) \quad \text{by right hypothesis} \\ &= (\| \mathbf{x} \|^2 + \| \mathbf{y} \|^2) - (\| \mathbf{x} \|^2 + |-1|^2 \| \mathbf{y} \|^2) + i (\| \mathbf{x} \|^2 + |i|^2 \| \mathbf{y} \|^2) - i (\| \mathbf{x} \|^2 + |-i|^2 \| \mathbf{y} \|^2) \quad \text{by definition of } \| \cdot \| \\ &= (\| \mathbf{x} \|^2 + \| \mathbf{y} \|^2) - (\| \mathbf{x} \|^2 + \| \mathbf{y} \|^2) + i (\| \mathbf{x} \|^2 + \| \mathbf{y} \|^2) - i (\| \mathbf{x} \|^2 + \| \mathbf{y} \|^2) \quad \text{by def. of } | \cdot | \\ &= 0 \end{aligned}$$



APPENDIX J

NORMED LINEAR SPACES

J.1 Definition and basic results

Definition J.1. ¹ Let $(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}))$ be a LINEAR SPACE (Definition F.1 page 185) and $|\cdot| \in \mathbb{R}^{\mathbb{F}}$ the ABSOLUTE VALUE function.

A functional $\|\cdot\|$ in \mathbb{R}^X is a **norm** if

- | | | | |
|-----|---|------------------------------------|-----|
| DEF | 1. $\ x\ \geq 0$ $\forall x \in X$ | (STRICTLY POSITIVE) | and |
| | 2. $\ x\ = 0 \iff x = 0$ $\forall x \in X$ | (NONDEGENERATE) | and |
| | 3. $\ \alpha x\ = \alpha \ x\ $ $\forall x \in X, \alpha \in \mathbb{C}$ | (HOMOGENEOUS) | and |
| | 4. $\ x + y\ \leq \ x\ + \ y\ $ $\forall x, y \in X$ | (SUBADDITIVE/TRIANGLE INEQUALITY). | |

A **normed linear space** is the tuple $(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \|\cdot\|)$.

The definition of the *norm* (Definition J.1 page 249) requires that any two vectors in a norm space be *subadditive* (they satisfy the *triangle inequality* property) such that $\|x_1 + x_2\| \leq \|x_1\| + \|x_2\|$. Actually, in **any** normed linear space, this property holds true for **any** finite number of vectors—not just two—such that $\|x_1 + x_2 + \dots + x_N\| \leq \|x_1\| + \|x_2\| + \dots + \|x_N\|$ (next theorem).

Theorem J.1 (triangle inequality). ² Let $(x_n \in X)_1^N$ be an N -TUPLE (Definition P.1 page 329) of vectors in a NORMED LINEAR SPACE $(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \|\cdot\|)$.

THM	$\left\ \sum_{n=1}^N x_n \right\ \leq \sum_{n=1}^N \ x_n\ \quad \forall N \in \mathbb{N}, x_n \in V$
-----	---

PROOF: Proof is by induction:

¹ Aliprantis and Burkinshaw (1998), pages 217–218, Banach (1932a), page 53, Banach (1932b), page 33, Banach (1922) page 135

² Michel and Herget (1993), page 344, Euclid (circa 300BC) (Book I Proposition 20)

1. Proof for the $N = 1$ case:

$$\begin{aligned} \left\| \sum_{n=1}^1 \mathbf{x}_n \right\| &= \|\mathbf{x}_1\| \\ &= \sum_{n=1}^1 \|\mathbf{x}_1\| \end{aligned}$$

2. Proof for the $N = 2$ case:

$$\begin{aligned} \left\| \sum_{n=1}^2 \mathbf{x}_n \right\| &= \left\| \sum_{n=1}^2 \mathbf{x}_n \right\| \\ &= \|\mathbf{x}_1 + \mathbf{x}_2\| \\ &\leq \|\mathbf{x}_1\| + \|\mathbf{x}_2\| && \text{by Definition J.1 page 249 (triangle inequality)} \\ &= \sum_{n=1}^2 \|\mathbf{x}_n\| \end{aligned}$$

3. Proof that [N case] \implies [$N + 1$ case]:

$$\begin{aligned} \left\| \sum_{n=1}^{N+1} \mathbf{x}_n \right\| &= \left\| \sum_{n=1}^N \mathbf{x}_n + \mathbf{x}_{N+1} \right\| \\ &\leq \left\| \sum_{n=1}^N \mathbf{x}_n \right\| + \|\mathbf{x}_{N+1}\| && \text{by Definition J.1 page 249 (triangle inequality)} \\ &\leq \sum_{n=1}^N \|\mathbf{x}_n\| + \|\mathbf{x}_{N+1}\| && \text{by left hypothesis} \\ &= \sum_{n=1}^{N+1} \|\mathbf{x}_n\| \end{aligned}$$



Theorem J.2 (Reverse Triangle Inequality). ³ Let $(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \|\cdot\|)$ be a NORMED LINEAR SPACE (Definition J.1 page 249).

T
H
M

$$\underbrace{\|\|\mathbf{x}\| - \|\mathbf{y}\|\|}_{\text{REVERSE TRIANGLE INEQUALITY}} \leq \|\mathbf{x} - \mathbf{y}\| \quad \forall \mathbf{x}, \mathbf{y} \in X$$


PROOF:

$$\begin{aligned} \|\|\mathbf{x}\| - \|\mathbf{y}\|\| &= \|(\mathbf{x} - \mathbf{y}) + \mathbf{y}\| - \|\mathbf{y}\| \\ &\leq \|\mathbf{x} - \mathbf{y}\| + \|\mathbf{y}\| - \|\mathbf{y}\| && \text{by Definition J.1 page 249} \\ &= \|\mathbf{x} - \mathbf{y}\| \\ &= \|\mathbf{x} - \mathbf{y}\| && \text{by Definition J.1 page 249} \end{aligned}$$

$$\begin{aligned} \|\mathbf{x} - \mathbf{y}\| &\leq \|\mathbf{x} - \mathbf{0}\| + \|\mathbf{0} - \mathbf{y}\| \\ &= \|\mathbf{x}\| + |-1| \|\mathbf{y}\| && \text{by previous result with } u = 0 \\ &= \|\mathbf{x}\| + \|\mathbf{y}\| && \text{by Definition J.1 page 249} \end{aligned}$$



³ Aliprantis and Burkinshaw (1998), page 218, Giles (2000) page 2, Banach (1922) page 136



The shortest distance between two vectors is always the difference of the vectors. This is proven in next and illustrated to the left in the Euclidean space \mathbb{R}^2 (the plane)

Proposition J.1. ⁴ Let $(X, +, \cdot, (\mathbb{F}, +, \dot{\times}), \|\cdot\|)$ be a NORMED LINEAR SPACE (Definition J.1 page 249).

P **R** **P**
$$\|x - y\| \leq \|x - u\| + \|u - y\| \quad \forall x, y, u \in X$$

PROOF:

$$\begin{aligned} \|x - y\| &= \|(x - u) + (u - y)\| \\ &\leq \|x - u\| + \|u - y\| \end{aligned} \quad \text{by Definition J.1 page 249}$$

Example J.1 (The usual norm). ⁵ Let $\mathbb{R}^\mathbb{R}$ be the set of all functions with domain and range the set of *real numbers* \mathbb{R} .

E **X** The absolute value (Definition ?? page ??) $|\cdot| \in \mathbb{R}^\mathbb{R}$ is a *norm*.

Example J.2 (l_p norms). Let $(x_n)_{n \in \mathbb{Z}}$ be a sequence (Definition P.1 page 329) of real numbers. An uncountably infinite number of norms is provided by the $\ell_p^{\mathbb{F}}$ norms $\|(x_n)\|_p$:

E **X**
$$\|(x_n)\|_p \triangleq \left(\sum_{n \in \mathbb{Z}} |x_n|^p \right)^{\frac{1}{p}}$$
 is a norm for $p \in [1 : \infty]$

J.2 Relationship between metrics and norms

J.2.1 Metrics generated by norms

The concept of *length* is very closely related to the concept of *distance*. Thus it is not surprising that a *norm* (a “length” function) can be used to define a *metric* (a “distance” function) on any *metric linear space* (Definition ?? page ??). Another way to say this is that the norm of a normed linear space *induces* a metric on this space. And so every normed linear space also has a metric. And because every normed linear space has a metric, **every normed linear space is also a metric space**. Actually this can be generalized one step further in that every metric space is also a *topological space*. And so **every normed linear space is also a topological space**. In symbols,

$$\text{normed linear space} \implies \text{metric space} \implies \text{topological space}.$$

⁴ Aliprantis and Burkinshaw (1998), page 218

⁵ Giles (1987), page 3

Theorem J.3. ⁶ Let $d \in \mathbb{R}^{X \times X}$ be a function on a REAL normed linear space $(X, +, \cdot, (\mathbb{R}, \dot{+}, \dot{\times}), \|\cdot\|)$. Let $B(x, r) \triangleq \{y \in X \mid \|y - x\| < r\}$ be the OPEN BALL of radius r centered at a point x .

T H M $d(x, y) \triangleq \|x - y\|$ is a metric on X

PROOF: The proof follows directly from the definition of a metric (not included in this text) the definition of *norm* (Definition J.1 page 249). \Rightarrow

The previous theorem defined a metric $d(x, y)$ induced by the norm $\|x\|$. The next definition defines this metric formally.

Definition J.2. ⁷ Let $(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \|\cdot\|)$ be a NORMED LINEAR SPACE (Definition J.1 page 249).

D E F The metric induced by the norm $\|\cdot\|$ is the function $d \in \mathbb{R}^X$ such that

$$d(x, y) \triangleq \|x - y\| \quad \forall x, y \in X$$

Due to its algebraic structure, every norm is *continuous* (Corollary J.1 page 252). This makes norm spaces very useful in analysis. For a function f to be *continuous*, for every $\epsilon > 0$ there must exist a $\delta > 0$ such that $|f(x + \delta) - f(x)| < \epsilon$. The *Reverse Triangle Inequality* (Theorem J.2 page 250) shows this to be true when $f(\cdot) \triangleq \|\cdot\|$.

Corollary J.1. ⁸ Let $\Omega \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \|\cdot\|)$ be a NORMED LINEAR SPACE (Definition J.1 page 249).

C O R The norm $\|\cdot\|$ is CONTINUOUS in Ω .

PROOF: This follows from these concepts:

1. The fact that $d(x, y) \triangleq \|x - y\|$ is a *metric* (Theorem J.3 page 252).
2. *Continuity* in a metric space.
3. The *Reverse Triangle Inequality* (Theorem J.2 page 250).

Theorem J.4 (next) demonstrates that **all open or closed balls in any normed linear space** are *convex*. However, the converse is not true—that is, a metric not generated by a norm may still produce a ball that is convex.

Theorem J.4. ⁹ Let $(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), d)$ be a METRIC LINEAR SPACE (Definition ?? page ??). Let B be the OPEN BALL $B(p, r) \triangleq \{x \in X \mid d(p, x) < r\}$ (open ball with respect to metric d centered at point p and with radius r).

T H M
$$\left. \begin{array}{l} \exists \|\cdot\| \in \mathbb{R}^X \text{ such that} \\ d(x, y) = \|y - x\| \\ \text{d is generated by a norm} \end{array} \right\} \Rightarrow \left\{ \begin{array}{l} 1. \quad B(x, r) = x + B(0, r) \\ 2. \quad B(0, r) = r B(0, 1) \\ 3. \quad B(x, r) \text{ is CONVEX} \\ 4. \quad x \in B(0, r) \iff -x \in B(0, r) \quad (\text{SYMMETRIC}) \end{array} \right.$$

⁶ Michel and Herget (1993), page 344, Banach (1932a) page 53

⁷ Giles (2000) page 1 (1.1 Definition)

⁸ Giles (2000) page 2

⁹ Giles (2000) page 2 (1.2 Remarks), Giles (1987) pages 22–26 (2.4 Theorem, 2.11 Theorem)

PROOF:

1. Proof that $d(x + z, y + vz) = d(x, y)$ (invariant):

$$\begin{aligned} d(x + z, y + vz) &= \|(y + vz) - (x + z)\| && \text{by left hypothesis} \\ &= \|y - x\| \\ &= d(x, y) && \text{by left hypothesis} \end{aligned}$$

2. Proof that $B(x, r) = x + B(0, r)$:

$$\begin{aligned} B(x, r) &= \{y \in X | d(x, y) < r\} && \text{by definition of open ball } B \\ &= \{y \in X | d(y - x, y - x) < r\} && \text{by right result 1.} \\ &= \{y \in X | d(0, y - x) < r\} \\ &= \{u + x \in X | d(0, u) < r\} && \text{let } u \triangleq y - x \\ &= x + \{u \in X | d(0, u) < r\} \\ &= x + B(0, r) && \text{by definition of open ball } B \end{aligned}$$

3. Proof that $B(0, r) = r B(0, 1)$:

$$\begin{aligned} B(0, r) &= \{y \in X | d(0, y) < r\} && \text{by definition of open ball } B \\ &= \left\{ y \in X | \frac{1}{r} d(0, y) < 1 \right\} \\ &= \left\{ y \in X | \frac{1}{r} \|y - 0\| < 1 \right\} && \text{by left hypothesis} \\ &= \left\{ y \in X | \left\| \frac{1}{r} y - \frac{1}{r} 0 \right\| < 1 \right\} && \text{by homogeneous property of } \|\cdot\| \text{ page 249} \\ &= \left\{ y \in X | d\left(\frac{1}{r} 0, \frac{1}{r} y\right) < 1 \right\} && \text{by left hypothesis} \\ &= \{ru \in X | d(0, u) < 1\} && \text{let } u \triangleq \frac{1}{r} y \\ &= r \{u \in X | d(0, u) < 1\} \\ &= r B(0, 1) && \text{by definition of open ball } B \end{aligned}$$

4. Proof that $B(p, r)$ is convex:

We must prove that for any pair of points x and y in the open ball $B(p, r)$, any point $\lambda x + (1 - \lambda)y$ is also in the open ball. That is, the distance from any point $\lambda x + (1 - \lambda)y$ to the ball's center p must be less than r .

$$\begin{aligned} d(p, \lambda x + (1 - \lambda)y) &= \|p - \lambda x - (1 - \lambda)y\| && \text{by left hypothesis} \\ &= \left\| \underbrace{\lambda p + (1 - \lambda)p - \lambda x - (1 - \lambda)y}_{p} \right\| \\ &= \|\lambda p - \lambda x + (1 - \lambda)p - (1 - \lambda)y\| \\ &\leq \|\lambda p - \lambda x\| + \|(1 - \lambda)p - (1 - \lambda)y\| && \text{by subadditivity property of } \|\cdot\| \text{ page 249} \\ &= |\lambda| \|p - x\| + |1 - \lambda| \|p - y\| && \text{by homogeneous property of } \|\cdot\| \text{ page 249} \\ &= \lambda \|p - x\| + (1 - \lambda) \|p - y\| && \text{because } 0 \leq \lambda \leq 1 \\ &\leq \lambda r + (1 - \lambda)r && \text{because } x, y \text{ are in the ball } B(p, r) \\ &= r \end{aligned}$$

5. Proof that $x \in B(\mathbf{0}, r) \iff -x \in B(\mathbf{0}, r)$ (symmetric):

$$\begin{aligned}
 x \in B(\mathbf{0}, r) &\iff x \in \{y \in X \mid d(\mathbf{0}, y) < r\} && \text{by definition of open ball } B \\
 &\iff x \in \{y \in X \mid \|y - \mathbf{0}\| < r\} && \text{by left hypothesis} \\
 &\iff x \in \{y \in X \mid \|y\| < r\} \\
 &\iff x \in \{y \in X \mid \|(-1)(-y)\| < r\} \\
 &\iff x \in \{y \in X \mid \| -1 \| \| -y \| < r\} && \text{by homogeneous property of } \|\cdot\| \text{ page 249} \\
 &\iff x \in \{y \in X \mid \| -y - \mathbf{0} \| < r\} \\
 &\iff x \in \{y \in X \mid d(\mathbf{0}, -y) < r\} && \text{by left hypothesis} \\
 &\iff x \in \{-u \in X \mid d(\mathbf{0}, u) < r\} && \text{let } u \triangleq -y \\
 &\iff x \in (-\{u \in X \mid d(\mathbf{0}, u) < r\}) \\
 &\iff x \in (-B(\mathbf{0}, r)) \\
 &\iff -x \in B(\mathbf{0}, r)
 \end{aligned}$$

⇒

Theorem J.4 (page 252) demonstrates that if a metric d in a metric space $(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), d)$ is generated by a norm, then the ball $B(x, r)$ in that metric linear space is *convex*. However, the converse is not true. That is, it is possible for the balls in a metric space (Y, p) to be *convex*, but yet the metric p not be generated by a norm.




Figure J.1: Open balls in (\mathbb{R}^0, d_n) , (\mathbb{R}, d_n) , (\mathbb{R}^2, d_n) , and (\mathbb{R}^3, d_n) .

J.2.2 Norms generated by metrics

Every normed linear space is also a metric linear space (Theorem J.3 page 252). That is, a metric linear space generates a *normed linear space*. However, the converse is not true—not every metric linear space is a *normed linear space*. A characterization of metric linear spaces that *are* normed linear spaces is given by Theorem J.5 page 255.

Lemma J.1. ¹⁰ Let $(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), d)$ be a METRIC LINEAR SPACE. Let $\|x\| \triangleq d(x, \mathbf{0}) \forall x \in X$.

¹⁰ Oikhberg and Rosenthal (2007), page 599

L E M

$$\underbrace{d(x+z, y+z) = d(x, y) \quad \forall x, y, z \in X}_{\text{TRANSLATION INVARIANT}} \implies \begin{cases} 1. \quad \|x\| = \|-x\| & \forall x \in X \quad \text{and} \\ 2. \quad \|x\| = 0 \iff x = 0 & \forall x \in X \quad \text{and} \\ 3. \quad \|x+y\| \leq \|x\| + \|y\| & \forall x, y \in X \end{cases}$$

PROOF:

1. Proof that $\|x\| = \|-x\|$:

$$\begin{aligned} \|x\| &= d(x, 0) && \text{by definition of } \|\cdot\| \\ &= d(x - x, 0 - x) && \text{by translation invariance hypothesis} \\ &= d(0, -x) \\ &= \|-x\| && \text{by definition of } \|\cdot\| \end{aligned}$$

2a. Proof that $\|x\| = 0 \implies x = 0$:

$$\begin{aligned} 0 &= \|x\| && \text{by left hypothesis} \\ &= d(x, 0) && \text{by definition of } \|\cdot\| \\ &= d(x, 0) && \text{by definition of } \|\cdot\| \\ &\implies x = 0 && \text{by property of metrics} \end{aligned}$$

2b. Proof that $\|x\| = 0 \iff x = 0$:

$$\begin{aligned} \|x\| &= d(x, 0) && \text{by definition of } \|\cdot\| \\ &= d(0, 0) && \text{by right hypothesis} \\ &= 0 && \text{by property of metrics} \end{aligned}$$

3. Proof that $\|x+y\| \leq \|x\| + \|y\|$:

$$\begin{aligned} \|x+y\| &= d(x+y, 0) && \text{by definition of } \|\cdot\| \\ &= d(x+y - y, 0 - y) && \text{by translation invariance hypothesis} \\ &= d(x, -y) \\ &\leq d(x, 0) + d(0, y) && \text{by property of metrics} \\ &= d(x, 0) + d(y, 0) && \text{by property of metrics} \\ &= \|x\| + \|y\| && \text{by definition of } \|\cdot\| \end{aligned}$$

Theorem J.5. ¹¹ Let $(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}))$ be a LINEAR SPACE. Let $d(x, y) \triangleq \|x - y\| \forall x, y \in X$.

T H M

$$\left. \begin{array}{l} 1. \quad d(x+z, y+z) = d(x, y) \quad \forall x, y, z \in X \quad (\text{TRANSLATION INVARIANT}) \quad \text{and} \\ 2. \quad d(\alpha x, \alpha y) = |\alpha| d(x, y) \quad \forall x, y \in X, \alpha \in \mathbb{F} \quad (\text{HOMOGENEOUS}) \end{array} \right\} \iff \|\cdot\| \text{ is a NORM}$$

PROOF:

1. Proof of \implies assertion:

- (a) Proof that $\|\cdot\|$ is *strictly positive*: This follows directly from the definition of d .
- (b) Proof that $\|\cdot\|$ is *nondegenerate*: This follows directly from Lemma J.1 (page 254).
- (c) Proof that $\|\cdot\|$ is *homogeneous*: This follows from the second left hypothesis.

¹¹  Bollobás (1999), page 21

(d) Proof that $\|\cdot\|$ satisfies the *triangle-inequality*: This follows directly from Lemma J.1 (page 254).

2. Proof of \Leftarrow assertion:


$$\begin{aligned}
 d(x+z, y+z) &= \|(x+z) - (y+z)\| && \text{by definition of } d \\
 &= \|x - y\| \\
 &= d(x, y) && \text{by definition of } d \\
 d(\alpha x, \alpha y) &= \|(\alpha x) - (\alpha y)\| && \text{by definition of } d \\
 &= \|\alpha(x - y)\| \\
 &= |\alpha| \|x - y\| && \text{by definition of } \|\cdot\| \text{ page 249} \\
 &= |\alpha| d(x, y) && \text{by definition of } d
 \end{aligned}$$



J.3 Orthogonality on normed linear spaces

Traditionally, *orthogonality* (Definition I.4 page 245) is a property defined in *inner product spaces* (Definition I.1 page 233). However, the concept of orthogonality can be extended to *normed linear spaces* (Definition J.1 page 249). Here are some examples:

- ① *Isosceles orthogonality*: Definition J.3 page 256
- ② *Pythagorean orthogonality*: Definition J.4 page 258
- ③ *Birkhoff orthogonality*: Definition J.5 page 258



Isosceles orthogonality (Definition J.3 page 256) can be illustrated using a *parallelogram*, as illustrated in the figure to the upper left. In this case, orthogonality implies that the parallelogram is a rectangle, which in turn implies that the lengths of the two diagonals are equal ($\|x + y\| = \|x - y\|$). Isosceles orthogonality can also be illustrated with a triangle where the sides are of lengths $\|x + y\|$ and $\|x - y\|$ and base of length $\|x + x\|$. In this case if x and y are orthogonal, then the triangle is *isosceles*. This is illustrated in figure to the lower left. Isosceles orthogonality is formally defined next.

Definition J.3. ¹² Let $(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \|\cdot\|)$ be a NORMED LINEAR SPACE (Definition J.1 page 249).

D E F Two vectors x and y are **orthogonal in the sense of James** if

$$\|x + y\| = \|x - y\|.$$

This property is also called **isosceles orthogonality** or **James orthogonality**.

Theorem J.6. Let $(X, +, \cdot, (\mathbb{R}, \dot{+}, \dot{\times}), \langle \Delta | \nabla \rangle)$ be an INNER-PRODUCT SPACE (Definition I.1 page 233) with induced norm $\|x\| \triangleq \sqrt{\langle x | x \rangle}$, ISOSCELES ORTHOGONALITY (Definition J.3 page 256) relation \oplus , and inner-product relation ORTHOGONALITY (Definition I.4 page 245) relation \perp .

T
H
M

$$\underbrace{x \oplus y}_{\text{orthogonal in the sense of James}}$$

\iff

$$\underbrace{x \perp y}_{\text{orthogonal in the sense of inner-product space}}$$

¹² James (1945) page 292 (DEFINITION 2.1), Amir (1986) page 24, Dunford and Schwartz (1957), page 93

PROOF:

1. Proof that $x \odot y \implies x \perp y$:

$$\begin{aligned}
 & 4 \langle x | y \rangle \\
 &= \underbrace{\|x + y\|^2 - \|x - y\|^2}_{0 \text{ by } x \odot y \text{ hypothesis}} + i \|x + iy\|^2 - i \|x - iy\|^2 \quad \text{by polarization identity (Theorem I.6 page 240)} \\
 &= 0 + i \|x + iy\|^2 - i \|x - iy\|^2 \quad \text{by } x \odot y \text{ hypothesis} \\
 &= i [\|x\|^2 + \|y\|^2 + 2\Re \langle x | iy \rangle] \\
 &\quad - i [\|x\|^2 + \|y\|^2 + 2\Re \langle x | -iy \rangle] \quad \text{by Polar Identity (Lemma I.1 page 237)} \\
 &= i [\|x\|^2 + \|y\|^2 + 2\Re \langle x | iy \rangle] \\
 &\quad - i [\|x\|^2 + \|y\|^2 + 2\Re \langle x | -iy \rangle] \quad \text{by Definition J.1 page 249 and Definition I.1 page 233} \\
 &= 4i\Re \langle x | iy \rangle \\
 &= 4i\Re [i^* \langle x | y \rangle] \\
 &= 0 \quad \text{because inner-product space is real } (\mathbb{F} = \mathbb{R})
 \end{aligned}$$

2. Proof that $x \odot y \iff x \perp y$:

$$\begin{aligned}
 \|x + y\|^2 &= \|x\|^2 + \|y\|^2 + 2\Re \langle x | y \rangle \quad \text{by Polar Identity (Lemma I.1 page 237)} \\
 &= \|x\|^2 + \|y\|^2 + 0 \quad \text{by } x \perp y \text{ hypothesis} \\
 &= \|x\|^2 + \|y\|^2 - 2\Re \langle x | y \rangle \quad \text{0 when } x \perp y \text{ by } x \perp y \text{ hypothesis} \\
 &= \|x\|^2 + \|y\|^2 + 2\Re \langle x | -y \rangle \\
 &= \|x - y\|^2 \quad \text{by Polar Identity (Lemma I.1 page 237)}
 \end{aligned}$$

Theorem J.7. ¹³ Let $(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \|\cdot\|)$ be a normed linear space and with ISOSCELES ORTHOGONALITY (Definition J.3 page 256) relation \odot .

T H M	$x \odot y \iff y \odot x \iff \alpha x \odot \alpha y \quad \forall \alpha \in \mathbb{F}$
-------------	---


PROOF:

$$\begin{aligned}
 x \odot y &\implies \|x + y\| = \|x - y\| \quad \text{by Definition J.3 page 256} \\
 &\implies \|x + y\| = |-1| \|x - y\| \\
 &\implies \|x + y\| = \|(x - y)\| \quad \text{by Definition J.1 page 249} \\
 &\implies \|y + x\| = \|y - x\| \quad \text{by Definition E.1 page 185} \\
 &\implies y \odot x \quad \text{by Definition J.3 page 256} \\
 \\
 y \odot x &\implies \|y + x\| = \|y - x\| \quad \text{by Definition J.3 page 256} \\
 &\implies |\alpha| \|y + x\| = |\alpha| \|y - x\| \\
 &\implies \|\alpha(y + x)\| = \|\alpha(y - x)\| \quad \text{by Definition J.1 page 249} \\
 &\implies \|\alpha y + \alpha x\| = \|\alpha y - \alpha x\| \\
 &\implies \|\alpha x + \alpha y\| = \|-(\alpha x - \alpha y)\| \quad \text{by Definition E.1 page 185}
 \end{aligned}$$

¹³ Amir (1986) page 24

$$\begin{aligned} \Rightarrow \| \alpha x + \alpha y \| &= | -1 | \| \alpha x - \alpha y \| && \text{by Definition J.1 page 249} \\ \Rightarrow \| \alpha x + \alpha y \| &= \| \alpha x - \alpha y \| \\ \Rightarrow \alpha x \odot \alpha y & && \text{by Definition J.3 page 256} \end{aligned}$$

$$\begin{aligned} \alpha x \odot \alpha y \Rightarrow \| \alpha x + \alpha y \| &= \| \alpha x - \alpha y \| && \text{by Definition J.3 page 256} \\ \Rightarrow \| \alpha(x + y) \| &= \| \alpha(x - y) \| && \text{by Definition E.1 page 185} \\ \Rightarrow |\alpha| \|x + y\| &= |\alpha| \|x - y\| && \text{by Definition J.1 page 249} \\ \Rightarrow \|x + y\| &= \|x - y\| && \text{by Definition J.1 page 249} \\ \Rightarrow x \odot y & && \text{by Definition J.3 page 256} \end{aligned}$$




If a triangle in a plane has two perpendicular sides of lengths a and b and a hypotenuse of length c , then by the *Pythagorean Theorem* (Theorem I.10 page 246), $a^2 + b^2 = c^2$. This concept of orthogonality can be generalized to normed linear spaces. Two vectors x and y (with lengths $\|x\|$ and $\|y\|$) are orthogonal when $\|x\|^2 + \|y\|^2 = \|x - y\|^2$ ($x - y$ is a kind of "hypotenuse"). This kind of orthogonality is defined next and illustrated in the figure to the left.

Definition J.4. ¹⁴ Let $(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \|\cdot\|)$ be a NORMED LINEAR SPACE (Definition J.1 page 249).

D E F Two vectors x and y are **orthogonal in the Pythagorean sense** if
 $\|x - y\|^2 = \|x\|^2 + \|y\|^2$.
This relationship is also called **Pythagorean orthogonality**.

Theorem J.8. ¹⁵ Let $(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle | \nabla \rangle)$ be an INNER-PRODUCT SPACE (Definition I.1 page 233) with induced norm $\|x\| \triangleq \sqrt{\langle x | x \rangle}$, PYTHAGOREAN ORTHOGONALITY (Definition J.4 page 258) relation \odot , and inner-product relation ORTHOGONALITY (Definition I.4 page 245) relation \perp .

T H M	$\underbrace{x \odot y}_{\text{orthogonal in the Pythagorean sense}}$	\iff	$\underbrace{x \perp y}_{\text{orthogonal in the sense of inner-product space}}$
-------	---	--------	--



Besides *isosceles orthogonality* (Definition J.3 page 256), orthogonality in normed linear spaces can be defined using *Birkhoff orthogonality*, as defined in Definition J.5 (next) and illustrated to the left.

Definition J.5. ¹⁶ Let $(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \|\cdot\|)$ be a NORMED LINEAR SPACE (Definition J.1 page 249).

D E F Two vectors x and y are **orthogonal in the sense of Birkhoff** if
 $\|x\| \leq \|x + \alpha y\| \quad \forall \alpha \in \mathbb{F}$.
This relationship is also called **Birkhoff orthogonality**.

Theorem J.9. Let $(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle | \nabla \rangle)$ be an INNER-PRODUCT SPACE (Definition I.1 page 233) with induced norm $\|x\| \triangleq \sqrt{\langle x | x \rangle}$, BIRKHOFF ORTHOGONALITY relation \odot (Definition J.5 page 258), and inner-product relation ORTHOGONALITY relation \perp (Definition I.4 page 245).

T H M	$\underbrace{x \odot y}_{\text{orthogonal in the sense of Birkhoff}}$	\iff	$\underbrace{x \perp y}_{\text{orthogonal in the sense of inner-product space}}$
-------	---	--------	--

¹⁴ James (1945) page 292 (DEFINITION 2.2), Amir (1986) page 57, Drljević (1989) page 232

¹⁵ Amir (1986) page 57

¹⁶ Amir (1986) page 33, Dunford and Schwartz (1957), page 93, James (1947) page 265

APPENDIX K

INTERVALS AND CONVEXITY

K.1 Intervals

In the real number system, for $a \leq b$, the *interval* $[a : b]$ is the set a and b and all the numbers inbetween, as in $[a : b] \triangleq \{x \in \mathbb{R} | a \leq x \leq b\}$. This concept can be easily generalized:

- In an **ordered set**, if two elements x and y are *comparable* and $x \leq y$, then we say that x and y and all the elements inbetween, as determined by the ordering relation \leq , are the interval $[a : b]$.
- In a **lattice**, the concept of the *interval* can be generalized even further. In an arbitrary ordered set, the interval $[x : y]$ of item (K.1) is restricted to the case in which x and y are *comparable*. This restriction can be lifted (Definition K.2 page 259) with the additional structure of upper and lower bounds provided by lattices.
- A **metric space** in general has no *order relation* \leq . But intervals can still be defined (Definition K.4 page 260) in a metric space in terms of the *triangle inequality*.
- A **linear space** (Definition F.1 page 185) over a real or complex field in general has no *order relation* that compares *vectors* in the space, but the standard order relation \leq for real numbers \mathbb{R} can still be used (Definition K.5 page 260) to define an interval in a linear space.

Definition K.1 (intervals on ordered sets). ¹ Let (X, \leq) be an ORDERED SET.

DEF	The set $[x : y] \triangleq \{z \in X x \leq z \leq y\}$ is called a closed interval and
DEF	The set $(x : y] \triangleq \{z \in X x < z \leq y\}$ is called a half-open interval and
DEF	The set $[x : y) \triangleq \{z \in X x \leq z < y\}$ is called a half-open interval and
DEF	The set $(x : y) \triangleq \{z \in X x < z < y\}$ is called an open interval .

Definition K.2 (intervals on lattices). ² Let $(X, \vee, \wedge; \leq)$ be a LATTICE.

DEF	The set $[x : y] \triangleq \{z \in X x \wedge y \leq z \leq x \vee y\}$ is called a closed interval .
DEF	The set $(x : y] \triangleq \{z \in X x \wedge y < z \leq x \vee y\}$ is called a half-open interval .
DEF	The set $[x : y) \triangleq \{z \in X x \wedge y \leq z < x \vee y\}$ is called a half-open interval .
DEF	The set $(x : y) \triangleq \{z \in X x \wedge y < z < x \vee y\}$ is called an open interval .

When x and y are comparable and $x \leq y$, then Definition K.2 (previous) simplifies to item (K.1)

¹ Apostol (1975) page 4, Ore (1935) page 409

² Duthie (1942) page 2, Ore (1935) page 425 (quotient structures)

(page 259).

Definition K.3. ³ Let $L \triangleq (X, \vee, \wedge; \leq)$ be a LATTICE with dual L^* . Let $[x : y]$ be a CLOSED INTERVAL (Definition K.2 page 259) on set X . The sublattices $L[x : y]$ and $L^*[x : y]$ are defined as follows:

DEF	$L[x : y] \triangleq \{z \in L z \in [x : y]\} \quad \forall x, y \in X$
DEF	$L^*[x : y] \triangleq \{z \in L^* z \in [x : y]\} \quad \forall x, y \in X$

Definition K.4. ⁴

DEF	In a METRIC SPACE (X, d) , the set $[a : b]$ is the closed interval from x to y and is defined as $[x : y] \triangleq \{z \in X d(x, z) + d(z, y) = d(x, y)\}$. An element $z \in X$ is geodesically between x and y if $z \in [x : y]$.
-----	---

Definition K.5. ⁵

DEF	In a LINEAR SPACE $(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}))$ (Definition F.1 page 185), $[x : y] \triangleq \{\lambda x + (1 - \lambda)y = z 0 \leq \lambda \leq 1\}$ is called a closed interval and $(x : y] \triangleq \{\lambda x + (1 - \lambda)y = z 0 < \lambda \leq 1\}$ is called a half-open interval and $[x : y) \triangleq \{\lambda x + (1 - \lambda)y = z 0 \leq \lambda < 1\}$ is called a half-open interval and $(x : y) \triangleq \{\lambda x + (1 - \lambda)y = z 0 < \lambda < 1\}$ is called an open interval .
-----	--

K.2 Convex sets


Using the concept of the *interval* (previous section), we can define the *convex set* (next definition).

Definition K.6. ⁶ Let X be a SET in an ORDERED SET (X, \leq) , a LATTICE $(X, \vee, \wedge; \leq)$, a METRIC SPACE (X, d) , or a LINEAR SPACE $(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}))$.

DEF	A subset $D \subseteq X$ is a convex set in X if $x, y \in D \implies [x : y] \subseteq D$.
-----	--


A set that is **not** convex is **concave**.

Example K.1. Consider the Euclidean space \mathbb{R}^2 (a special case of a *linear space*).



$\Leftarrow \begin{cases} \text{The figure to the left is a} \\ \text{convex set in } \mathbb{R}^2. \end{cases}$

$\Rightarrow \begin{cases} \text{The figure to the right is a} \\ \text{concave set in } \mathbb{R}^2. \end{cases}$



Example K.2. In a *metric space*, examples of *convex sets* are *convex balls*. Examples include those balls generated by the following metrics:

- Taxi-cab metric
- Euclidean metric
- Sup metric
- Tangential metric

³ Maeda and Maeda (1970), page 1

⁴ van de Vel (1993) page 8

⁵ Barvinok (2002) page 2

⁶ Barvinok (2002) page 5

Examples of metrics generating balls which are *not* convex include the following:

- ➊ Parabolic metric
- ➋ Exponential metric

K.3 Convex functions




Figure K.1: Convex and concave functions

Definition K.7. ⁷ Let $(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}))$ be a LINEAR SPACE (Definition F.1 page 185) and D a CONVEX SET (Definition K.6 page 260) in X .

A function $f \in F^D$ is **convex** if

$$f(\lambda x + [1 - \lambda]y) \leq \lambda f(x) + (1 - \lambda) f(y) \quad \forall x, y \in D \text{ and } \forall \lambda \in (0, 1)$$


A function $g \in F^D$ is **strictly convex** if

$$g(\lambda x + [1 - \lambda]y) = \lambda g(x) + (1 - \lambda) g(y) \quad \forall x, y \in D, x \neq y, \text{ and } \forall \lambda \in (0, 1)$$

A function $f \in F^D$ is **concave** if $-f$ is CONVEX.

A function $f \in F^D$ is **affine** iff is CONVEX and CONCAVE.

Example K.3. The function $f(x) = 2^x$ is a **convex function** (Definition K.7 page 261), as illustrated to the right.



Definition K.8. ⁸ Let $(X, +, \cdot, (\mathbb{R}, \dot{+}, \dot{\times}))$ be a LINEAR SPACE (Definition F.1 page 185).

The **epigraph** $\text{epi}(f)$ and **hypograph** $\text{hyp}(f)$ of a functional $f \in \mathbb{R}^X$ are defined as


$$\text{epi}(f) \triangleq \{(x, y) \in X \times \mathbb{R} | y \geq f(x)\}$$

$$\text{hyp}(f) \triangleq \{(x, y) \in X \times \mathbb{R} | y \leq f(x)\}$$

Example K.4.

⁷ Simon (2011) page 2, Barvinok (2002) page 2, Bollobás (1999), page 3, Jensen (1906), page 176, Clarkson (1936) (strictly convex)

⁸ Beer (1993) page 13 (§1.3), Aubin and Frankowska (2009) page 222, Aubin (2011) page 223



Proposition K.1.⁹ Let $(X, +, \cdot, (\mathbb{R}, \dot{+}, \dot{\times}))$ be a LINEAR SPACE (Definition F.1 page 185). Let f be a FUNCTIONAL in \mathbb{R}^X .

P R P	$\left\{ \begin{array}{l} f \text{ is a} \\ \text{CONVEX FUNCTION} \end{array} \right\} \Leftrightarrow \left\{ \begin{array}{l} \text{epi}(f) \text{ is a} \\ \text{CONVEX SET} \end{array} \right\}$
----------------------	--

Often a function can be proven to be *convex* or *concave*. *Convex* and *concave* functions are defined in Definition K.9 (page 262) (next) and illustrated in Figure K.1 (page 261).

Definition K.9. Let

$$y(x) = \frac{f(b) - f(a)}{b - a}(x - a) + f(a).$$

A function $f : \mathbb{R} \rightarrow \mathbb{R}$ is

- | | |
|----------------------|--|
| D
E
F | <ul style="list-style-type: none"> (1). convex <i>in $(a : b)$ if $f(x) \leq y(x)$ for $x \in (a : b)$</i> (2). concave <i>in $(a : b)$ if $f(x) \geq y(x)$ for $x \in (a : b)$</i> (3). strictly convex <i>in $(a : b)$ if $f(x) < y(x)$ for $x \in (a : b)$</i> (4). strictly concave <i>in $(a : b)$ if $f(x) > y(x)$ for $x \in (a : b)$</i> |
|----------------------|--|

Theorem K.1 (Jensen's Inequality).¹⁰ Let $(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}))$ be a LINEAR SPACE (Definition F.1 page 185), D a subset of X , and f a functional in \mathbb{F}^D . Let \sum be the SUMMATION OPERATOR (Definition L.1 page 267).

T H M	$\left\{ \begin{array}{ll} 1. & D \text{ is CONVEX} & \text{and} \\ 2. & f \text{ is CONVEX} & \text{and} \\ 3. & \sum_{n=1}^N \lambda_n = 1 & (\text{WEIGHTS}) \end{array} \right\} \implies f\left(\sum_{n=1}^N \lambda_n x_n\right) \leq \sum_{n=1}^N \lambda_n f(x_n) \quad \forall x_n \in D, N \in \mathbb{N}$
----------------------	--

PROOF: Proof is by induction:

1. Proof that statement is true for $N = 1$:

$$\begin{aligned} f\left(\sum_{n=1}^{N=1} \lambda_n x_n\right) &= f(\lambda_1 x_1) \\ &\leq f(\lambda_1 x_1) \\ &= \sum_{n=1}^{N=1} \lambda_n f(x_n) \end{aligned}$$

⁹ Udriste (1994) page 63, Kurdila and Zabarankin (2005) page 178 (Proposition 6.1.1), Rockafellar (1970) page 23 (Section 4 Convex Functions), Çinlar and Vanderbei (2013) page 86 (5.4 Theorem)

¹⁰ Mitrinović et al. (2010) page 6, Bollobás (1999) page 3, Lay (1982) page 7, Jensen (1906), pages 179–180

2. Proof that statement is true for $N = 2$:

$$\begin{aligned} f\left(\sum_{n=1}^{N=2} \lambda_n x_n\right) &= f(\lambda_1 x_1 + \lambda_2 x_2) \\ &\leq \lambda_1 f(x_1) + \lambda_2 f(x_2) && \text{by convexity hypothesis} \\ &= \sum_{n=1}^{N=2} \lambda_n f(x_n) \end{aligned}$$

3. Proof that if the statement is true for N , then it is also true for $N + 1$:

$$\begin{aligned} f\left(\sum_{n=1}^{N+1} \lambda_n x_n\right) &= f\left(\sum_{n=1}^N \lambda_n x_n + \lambda_{N+1} x_{N+1}\right) \\ &= f\left([1 - \lambda_{N+1}] \sum_{n=1}^N \frac{\lambda_n}{1 - \lambda_{N+1}} x_n + \lambda_{N+1} x_{N+1}\right) \\ &\leq [1 - \lambda_{N+1}] f\left(\sum_{n=1}^N \frac{\lambda_n}{1 - \lambda_{N+1}} x_n\right) + \lambda_{N+1} f(x_{N+1}) && \text{by convexity hypothesis} \\ &\leq [1 - \lambda_{N+1}] \sum_{n=1}^N \frac{\lambda_n}{1 - \lambda_{N+1}} f(x_n) + \lambda_{N+1} f(x_{N+1}) && \text{by "true for } N\text{" hypothesis} \\ &= \sum_{n=1}^N \lambda_n f(x_n) + \lambda_{N+1} f(x_{N+1}) \\ &= \sum_{n=1}^{N+1} \lambda_n f(x_n) \end{aligned}$$

4. Since the statement is true for $N = 1$, $N = 2$, and true for $N \implies$ true for $N + 1$, then it is true for $N = 1, 2, 3, 4, \dots$



The next theorem gives another form of convex functions that is a little less intuitive but provides powerful analytic results.

Theorem K.2. Let $f : \mathbb{R} \rightarrow \mathbb{R}$. For every $x_1, x_2 \in (a, b)$ and $\lambda \in [0, 1]$

T
H
M

f is convex in $(a, b) \iff f(\lambda x_1 + (1 - \lambda)x_2) \leq \lambda f(x_1) + (1 - \lambda)f(x_2)$

PROOF:

1. prove f is convex $\implies f(\lambda x_1 + (1 - \lambda)x_2) \leq \lambda f(x_1) + (1 - \lambda)f(x_2)$:

$$\begin{aligned} f(\lambda x_1 + (1 - \lambda)x_2) &\leq \frac{f(b) - f(a)}{b - a} [\lambda x_1 + (1 - \lambda)x_2 - a] + f(a) \\ &= \frac{f(x_2) - f(x_1)}{x_2 - x_1} [\lambda x_1 + (1 - \lambda)x_2 - x_1] + f(x_1) \\ &= \frac{f(x_2) - f(x_1)}{x_2 - x_1} [(x_2 - x_1)(1 - \lambda)] + f(x_1) \\ &= (1 - \lambda)f(x_2) - (1 - \lambda)f(x_1) + f(x_1) \\ &= \lambda f(x_1) + (1 - \lambda)f(x_2) \end{aligned}$$

2. prove f is convex $\iff f(\lambda x_1 + (1 - \lambda)x_2) \leq \lambda f(x_1) + (1 - \lambda)f(x_2)$:

Let $x = \lambda(b - a) + a$ Notice that as λ varies from 0 to 1, x varies from b to a . So free variable λ works as a change of variable for free variable x .

$$\begin{aligned}\lambda &= \frac{x - a}{b - a} \\ f(x) &= f(\lambda(b - a) + a) \\ &\leq \lambda f(b) + (1 - \lambda)f(a) \\ &= \lambda[f(b) - f(a)] + f(a) \\ &= \frac{f(b) - f(a)}{b - a}(x - a) + f(a)\end{aligned}$$



Taking the second derivative of a function provides a convenient test for whether that function is convex.

Theorem K.3. ¹¹

T
H
M

$f''(x) > 0 \implies f$ is convex

PROOF:

$$\begin{aligned}f(x) &= f(x_0) + f'(x_0)(x - x_0) + f''(c)(x - x_0)^2 \\ &\geq f(x_0) + f'(x_0)(x - x_0) \\ &= f(x_0) + f'(x_0)(x - \lambda x_1 - (1 - \lambda)x_2)\end{aligned}$$

$$\begin{aligned}f(x_1) &\geq f(x_0) + f'(x_0)(x_1 - \lambda x_1 - (1 - \lambda)x_2) \\ &= f(x_0) + f'(x_0)(1 - \lambda)(x_1 - x_2) \\ &= f(x_0) - f'(x_0)(1 - \lambda)(x_2 - x_1)\end{aligned}$$

$$\begin{aligned}f(x_2) &\geq f(x_0) + f'(x_0)(x_2 - \lambda x_1 - (1 - \lambda)x_2) \\ &= f(x_0) + f'(x_0)\lambda(x_2 - x_1)\end{aligned}$$

$$\begin{aligned}\lambda f(x_1) + (1 - \lambda)f(x_2) &\geq \lambda [f(x_0) - f'(x_0)(1 - \lambda)(x_2 - x_1)] + (1 - \lambda) [f(x_0) + f'(x_0)\lambda(x_2 - x_1)] \\ &= \lambda [f(x_0) - f'(x_0)(1 - \lambda)(x_2 - x_1)] + [f(x_0) + f'(x_0)\lambda(x_2 - x_1)] - \lambda [f(x_0) + f'(x_0)\lambda(x_2 - x_1)] \\ &= f(x_0) \\ &= f(\lambda x_1 + (1 - \lambda)x_2)\end{aligned}$$

By Theorem K.2 (page 263), $f(x)$ is convex.



K.4 Literature

Literature survey:

¹¹ Cover and Thomas (1991), pages 24–25

1. Abstract convexity:

- [Edelman and Jamison \(1985\)](#)
- [van de Vel \(1993\)](#)
- [Hörmander \(1994\)](#)

2. Order convexity (lattice theory):

- [Edelman \(1986\)](#)

3. Metric convexity:

- [Menger \(1928\)](#)
- [Blumenthal \(1970\) page 41 \(?\)](#)
- [Khamsi and Kirk \(2001\) pages 35–38](#)



APPENDIX L

FINITE SUMS



“I think that it was Harald Bohr who remarked to me that “all analysts spend half their time hunting through the literature for inequalities which they want to use and cannot prove.””¹

G.H. Hardy (1877–1947) in his “Presidential Address” to the London Mathematical Society on November 8, 1928, about a remark that he suggested was from Harald Bohr (1887–1951), Danish mathematician pictured to the left.¹

L.1 Summation

Definition L.1. ² Let $+$ be an addition operator on a tuple $(x_n)_m^N$.

The summation of (x_n) from index m to index N with respect to $+$ is

$$\sum_{n=m}^N x_n \triangleq \begin{cases} 0 & \text{for } N < m \\ \left(\sum_{n=m}^{N-1} x_n \right) + x_N & \text{for } N \geq m \end{cases}$$

Theorem L.1 (Generalized associative property). ³ Let $+$ be an addition operator on a tuple $(x_n)_m^N$.

+ is ASSOCIATIVE \implies

$$\sum_{n=m}^L x_n + \left(\sum_{n=L+1}^M x_n + \sum_{n=M+1}^N x_n \right) = \left(\sum_{n=m}^L x_n + \sum_{n=L+1}^M x_n \right) + \sum_{n=M+1}^N x_n \quad \text{for } m < L < M \leq N$$

$\overbrace{\hspace{10em}}$
 $\sum_{n=m}^N$ is ASSOCIATIVE

PROOF:

¹ quote: [Hardy \(1929\)](#), page 64

image: http://www-history.mcs.st-andrews.ac.uk/PictDisplay/Bohr_Harald.html

² reference: [Berberian \(1961\)](#) page 8 (Definition I.3.1)

“ Σ ” notation: [Fourier \(1820\)](#) page 280

³ [Berberian \(1961\)](#) pages 9–10 (Theorem I.3.1)

1. Proof for $N < m$ case: $\sum_{n=m}^N x_n = 0$.

2. Proof for $N = m$ case: $\sum_{n=m}^m x_n = \left(\sum_{n=m}^{m-1} x_n \right) + x_m = 0 + x_m = x_m$.

3. Proof for $N = m + 1$ case: $\sum_{n=m}^{m+1} x_n = \left(\sum_{n=m}^m x_n \right) + x_{m+1} = x_m + x_{m+1}$

4. Proof for $N = m + 2$ case:

$$\begin{aligned}\sum_{n=m}^{m+2} x_n &= \left(\sum_{n=m}^{m+1} x_n \right) + x_{m+2} \\ &= (x_m + x_{m+1}) + x_{m+2} \\ &= x_m + (x_{m+1} + x_{m+2})\end{aligned}$$

by Definition L.1 page 267

by item (3)

by left hypothesis

5. Proof that N case $\implies N + 1$ case:

$$\begin{aligned}\sum_{n=m}^{N+1} x_n &= \underbrace{\left(\sum_{n=m}^N x_n \right)}_{\text{associative}} + x_{N+1} \\ &= \left(\sum_{n=m}^L x_n + \left(\sum_{n=L+1}^M x_n + \sum_{n=M+1}^N x_n \right) \right) + x_{N+1} \\ &= \left(\left(\sum_{n=m}^L x_n + \sum_{n=L+1}^M x_n \right) + \sum_{n=M+1}^N x_n \right) + x_{N+1} \\ &= \left(\sum_{n=m}^L x_n + \sum_{n=L+1}^M x_n \right) + \left(\sum_{n=M+1}^N x_n + x_{N+1} \right) \\ &= \left(\sum_{n=m}^L x_n + \sum_{n=L+1}^M x_n \right) + \left(\sum_{n=M+1}^{N+1} x_n \right)\end{aligned}$$

by Definition L.1 page 267

L.2 Means

L.2.1 Weighted ϕ -means

Definition L.2. ⁴

The $(\lambda_n)_1^N$ weighted ϕ -mean of a tuple $(x_n)_1^N$ is defined as

$$M_\phi((x_n)) \triangleq \phi^{-1} \left(\sum_{n=1}^N \lambda_n \phi(x_n) \right)$$

where ϕ is a CONTINUOUS and STRICTLY MONOTONIC function in $\mathbb{R}^{\mathbb{R}^+}$

and $(\lambda_n)_{n=1}^N$ is a sequence of weights for which $\sum_{n=1}^N \lambda_n = 1$.

Lemma L.1. ⁵ Let $M_\phi((x_n))$ be the $(\lambda_n)_1^N$ weighted ϕ -mean of a tuple $(x_n)_1^N$. Let the property CONVEX be defined as in Definition K.7 (page 261).

DEF	$\phi\psi^{-1}$ is CONVEX and ϕ is INCREASING $\implies M_\phi((x_n)) \geq M_\psi((x_n))$
LEM	$\phi\psi^{-1}$ is CONVEX and ϕ is DECREASING $\implies M_\phi((x_n)) \leq M_\psi((x_n))$
LEM	$\phi\psi^{-1}$ is CONCAVE and ϕ is INCREASING $\implies M_\phi((x_n)) \leq M_\psi((x_n))$
LEM	$\phi\psi^{-1}$ is CONCAVE and ϕ is DECREASING $\implies M_\phi((x_n)) \geq M_\psi((x_n))$

PROOF:

1. Case where $\phi\psi^{-1}$ is convex and ϕ is increasing:

$$\begin{aligned} M_\phi((x_n)) &\triangleq \phi^{-1} \left(\sum_{n=1}^N \lambda_n \phi(x_n) \right) && \text{by definition of } M_\phi \text{ (Definition L.2 page 269)} \\ &= \phi^{-1} \left(\sum_{n=1}^N \lambda_n \phi\psi^{-1}\psi(x_n) \right) && \text{by definition of } \psi^{-1} \\ &\geq \phi^{-1} \left(\phi\psi^{-1} \sum_{n=1}^N \lambda_n \psi(x_n) \right) && \text{by Jensen's Inequality (Theorem K.1 page 262)} \\ &= \left(\psi^{-1} \sum_{n=1}^N \lambda_n \psi(x_n) \right) && \text{by definition of } \psi^{-1} \\ &\triangleq M_\psi((x_n)) && \text{by definition of } M_\psi \text{ (Definition L.2 page 269)} \end{aligned}$$

2. Case where $\phi\psi^{-1}$ is convex and ϕ is decreasing:

$$\begin{aligned} M_\phi((x_n)) &\triangleq \phi^{-1} \left(\sum_{n=1}^N \lambda_n \phi(x_n) \right) && \text{by definition of } M_\phi \text{ (Definition L.2 page 269)} \\ &= \phi^{-1} \left(\sum_{n=1}^N \lambda_n \phi\psi^{-1}\psi(x_n) \right) && \text{by definition of } \psi^{-1} \\ &\leq \phi^{-1} \left(\phi\psi^{-1} \sum_{n=1}^N \lambda_n \psi(x_n) \right) && \text{by Jensen's Inequality and because } \phi^{-1} \text{ is decreasing} \end{aligned}$$

⁴  Bollobás (1999) page 5

⁵  Pečarić et al. (1992) page 107,  Bollobás (1999) page 5,  Hardy et al. (1952) page 75

$$\begin{aligned}
 &= \left(\psi^{-1} \sum_{n=1}^N \lambda_n \psi(x_n) \right) && \text{by definition of } \psi^{-1} \\
 &\triangleq M_\psi(\langle x_n \rangle) && \text{by definition of } M_\psi \text{ (Definition L.2 page 269)}
 \end{aligned}$$



One of the most well known inequalities in mathematics is *Minkowski's Inequality* (1910, Theorem L.5 page 276). In 1946, H.P. Mulholland submitted a result⁶ that generalizes Minkowski's Inequality to an equal weighted ϕ -mean. And Milovanović and Milovanović (1979) generalized this even further to a *weighted* ϕ -mean (Theorem L.2, next).

Theorem L.2. ⁷

T H M	$ \left\{ \begin{array}{ll} \text{1. } \phi \text{ is CONVEX} & \text{and} \\ \text{2. } \phi \text{ is STRICTLY MONOTONIC} & \text{and} \\ \text{3. } \phi(0) = 0 & \text{and} \\ \text{4. } \log \circ \phi \circ \exp \text{ is CONVEX} & \end{array} \right\} \implies $ $ \phi^{-1} \left(\sum_{n=1}^N \lambda_n \phi(x_n + y_n) \right) \leq \phi^{-1} \left(\sum_{n=1}^N \lambda_n \phi(x_n) \right) + \phi^{-1} \left(\sum_{n=1}^N \lambda_n \phi(y_n) \right) $
----------------------	--

L.2.2 Power means

Definition L.3. ⁸ Let $M_{\phi(x;r)}(\langle x_n \rangle)$ be the $\langle \lambda_n \rangle_1^N$ weighted ϕ -mean of a NON-NEGATIVE tuple $\langle x_n \rangle_1^N$ (Definition L.2 page 269).

DEF A mean $M_{\phi(x;r)}(\langle x_n \rangle)$ is a **power mean** with parameter r if $\phi(x) \triangleq x^r$. That is,

$$M_{\phi(x;r)}(\langle x_n \rangle) = \left(\sum_{n=1}^N \lambda_n (x_n)^r \right)^{\frac{1}{r}}$$

Theorem L.3. ⁹ Let $M_{\phi(x;r)}(\langle x_n \rangle)$ be POWER MEAN with parameter r of an N -tuple $\langle x_n \rangle_1^N$. Let \mathbb{R}^* be the set of extended real numbers ($\mathbb{R}^* \triangleq \mathbb{R} \cup \{-\infty, \infty\}$).¹⁰

T H M	$ M_{\phi(x;r)}(\langle x_n \rangle) \triangleq \left(\sum_{n=1}^N \lambda_n (x_n)^r \right)^{\frac{1}{r}} \text{ is CONTINUOUS and STRICTLY INCREASING in } \mathbb{R}^*. $ $ M_{\phi(x;r)}(\langle x_n \rangle) = \begin{cases} \min_{n=1,2,\dots,N} \langle x_n \rangle & \text{for } r = -\infty \\ \prod_{n=1}^N x_n^{\lambda_n} & \text{for } r = 0 \\ \max_{n=1,2,\dots,N} \langle x_n \rangle & \text{for } r = +\infty \end{cases} $
----------------------	--

PROOF:

⁶ Mulholland (1950)

⁷ Milovanović and Milovanović (1979), Bullen (2003) page 306 (Theorem 9)

⁸ Bullen (2003) page 175, Bollobás (1999) page 6

⁹ Bullen (2003) pages 175–177 (see also page 203), Bollobás (1999) pages 6–8, Besso (1879), Bienaymé (1840) page 68

¹⁰ Rana (2002) pages 385–388 (Appendix A)

1. Proof that $M_{\phi(x;r)}$ is *strictly increasing* in r :

(a) Let r and s be such that $-\infty < r < s < \infty$.

(b) Let $\phi_r \triangleq x^r$ and $\phi_s \triangleq x^s$. Then $\phi_r \phi_s^{-1} = x^{\frac{r}{s}}$.

(c) The composite function $\phi_r \phi_s^{-1}$ is *convex* or *concave* depending on the values of r and s :

	$r < 0$ (ϕ_r decreasing)	$r > 0$ (ϕ_r increasing)
$s < 0$	convex	(not possible)
$s > 0$	convex	concave

(d) Therefore by Lemma L.1 (page 269),

$$-\infty < r < s < \infty \implies M_{\phi(x;r)}(\langle x_n \rangle) < M_{\phi(x;s)}(\langle x_n \rangle).$$

2. Proof that $M_{\phi(x;r)}$ is continuous in r for $r \in \mathbb{R} \setminus 0$: The sum of continuous functions is continuous. For the cases of $r \in \{-\infty, 0, \infty\}$, see the items that follow.

3. Lemma: $M_{\phi(x;-r)}(\langle x_n \rangle) = \{M_{\phi(x;r)}(\langle x_n^{-1} \rangle)\}^{-1}$. Proof:

$$\begin{aligned} \{M_{\phi(x;r)}(\langle x_n^{-1} \rangle)\}^{-1} &= \left\{ \left(\sum_{n=1}^N \lambda_n (x_n^{-1})^r \right)^{\frac{1}{r}} \right\}^{-1} && \text{by definition of } M_{\phi} \\ &= \left(\sum_{n=1}^N \lambda_n (x_n)^{-r} \right)^{\frac{1}{-r}} \\ &= M_{\phi(x;-r)}(\langle x_n \rangle) && \text{by definition of } M_{\phi} \end{aligned}$$

4. Proof that $\lim_{r \rightarrow \infty} M_{\phi}(\langle x_n \rangle) = \max_{n \in \mathbb{Z}} \langle x_n \rangle$:

(a) Let $x_m \triangleq \max_{n \in \mathbb{Z}} \langle x_n \rangle$

(b) Note that $\lim_{r \rightarrow \infty} M_{\phi} \leq \max_{n \in \mathbb{Z}} \langle x_n \rangle$ because

$$\begin{aligned} \lim_{r \rightarrow \infty} M_{\phi}(\langle x_n \rangle) &= \lim_{r \rightarrow \infty} \left(\sum_{n=1}^N \lambda_n x_n^r \right)^{\frac{1}{r}} && \text{by definition of } M_{\phi} \\ &\leq \lim_{r \rightarrow \infty} \left(\sum_{n=1}^N \lambda_n x_m^r \right)^{\frac{1}{r}} && \text{by definition of } x_m \text{ in item (4a) and because} \\ &&& \phi(x) \triangleq x^r \text{ and } \phi^{-1} \text{ are both increasing or both} \\ &&& \text{decreasing} \\ &= \lim_{r \rightarrow \infty} \left(x_m^r \underbrace{\sum_{n=1}^N \lambda_n}_{1} \right)^{\frac{1}{r}} && \text{because } x_m \text{ is a constant} \\ &= \lim_{r \rightarrow \infty} (x_m^r \cdot 1)^{\frac{1}{r}} \\ &= x_m \\ &= \max_{n \in \mathbb{Z}} \langle x_n \rangle && \text{by definition of } x_m \text{ in item (4a)} \end{aligned}$$

(c) But also note that $\lim_{r \rightarrow \infty} M_\phi(\{x_n\}) \geq \max_{n \in \mathbb{Z}} (\{x_n\})$ because

$$\begin{aligned} \lim_{r \rightarrow \infty} M_\phi(\{x_n\}) &= \lim_{r \rightarrow \infty} \left(\sum_{n=1}^N \lambda_n x_n^r \right)^{\frac{1}{r}} && \text{by definition of } M_\phi \\ &\geq \lim_{r \rightarrow \infty} (w_m x_m^r)^{\frac{1}{r}} && \text{by definition of } x_m \text{ in item (4a) and because} \\ &&& \phi(x) \triangleq x^r \text{ and } \phi^{-1} \text{ are both increasing or both} \\ &&& \text{decreasing} \\ &= \lim_{r \rightarrow \infty} w_m^{\frac{1}{r}} x_m^r \\ &= x_m \\ &= \max_{n \in \mathbb{Z}} (\{x_n\}) && \text{by definition of } x_m \text{ in item (4a)} \end{aligned}$$

(d) Combining items (b) and (c) we have $\lim_{r \rightarrow \infty} M_\phi = \max_{n \in \mathbb{Z}} (\{x_n\})$.

5. Proof that $\lim_{r \rightarrow -\infty} M_\phi(\{x_n\}) = \min_{n \in \mathbb{Z}} (\{x_n\})$:

$$\begin{aligned} \lim_{r \rightarrow -\infty} M_{\phi(x;r)}(\{x_n\}) &= \lim_{r \rightarrow \infty} M_{\phi(x;-r)}(\{x_n\}) && \text{by change of variable } r \\ &= \lim_{r \rightarrow \infty} \{M_{\phi(x;r)}(\{x_n^{-1}\})\}^{-1} && \text{by Lemma in item (3) page 271} \\ &= \lim_{r \rightarrow \infty} \frac{1}{M_{\phi(x;r)}(\{x_n^{-1}\})} \\ &= \frac{\lim_{r \rightarrow \infty} 1}{\lim_{r \rightarrow \infty} M_{\phi(x;r)}(\{x_n^{-1}\})} && \text{by property of lim } ^{11} \\ &= \frac{1}{\max_{n \in \mathbb{Z}} (\{x_n^{-1}\})} && \text{by item (4)} \\ &= \frac{1}{\left(\min_{n \in \mathbb{Z}} (\{x_n\}) \right)^{-1}} \\ &= \min_{n \in \mathbb{Z}} (\{x_n\}) \end{aligned}$$

6. Proof that $\lim_{r \rightarrow 0} M_\phi(\{x_n\}) = \prod_{n=1}^N x_n^{\lambda_n}$:

$$\begin{aligned} \lim_{r \rightarrow 0} M_\phi(\{x_n\}) &= \lim_{r \rightarrow 0} \exp \{ \ln \{ M_\phi(\{x_n\}) \} \} \\ &= \lim_{r \rightarrow 0} \exp \left\{ \ln \left\{ \left(\sum_{n=1}^N \lambda_n (x_n^r) \right)^{\frac{1}{r}} \right\} \right\} && \text{by definition of } M_\phi \\ &= \exp \left\{ \frac{\frac{\partial}{\partial r} \ln \left(\sum_{n=1}^N \lambda_n (x_n^r) \right)}{\frac{\partial}{\partial r} r} \right\}_{r=0} && \text{by l'Hôpital's rule}^{12} \end{aligned}$$

¹¹  Rudin (1976) page 85 (4.4 Theorem)

$$\begin{aligned}
&= \exp \left\{ \frac{\sum_{n=1}^N \lambda_n \frac{\partial}{\partial r} (x_n^r)}{\sum_{n=1}^N \lambda_n (x_n^r)} \right\}_{r=0} \\
&= \exp \left\{ \frac{\sum_{n=1}^N \lambda_n \frac{\partial}{\partial r} \exp (\ln (x_n^r))}{\sum_{n=1}^N \lambda_n} \right\}_{r=0} \\
&= \exp \left\{ \frac{\sum_{n=1}^N \lambda_n \frac{\partial}{\partial r} \exp (r \ln (x_n))}{1} \right\}_{r=0} \\
&= \exp \left\{ \sum_{n=1}^N \lambda_n \frac{\partial}{\partial r} \exp (r \ln (x_n)) \right\}_{r=0} \\
&= \exp \left\{ \sum_{n=1}^N \lambda_n \exp \{r \ln x_n\} \ln (x_n) \right\}_{r=0} \\
&= \exp \left\{ \sum_{n=1}^N \lambda_n \ln (x_n) \right\} \\
&= \exp \left\{ \sum_{n=1}^N \ln (x_n^{\lambda_n}) \right\} \\
&= \exp \left\{ \ln \prod_{n=1}^N x_n^{\lambda_n} \right\} \\
&= \prod_{n=1}^N x_n^{\lambda_n}
\end{aligned}$$



Definition L.4. Let $(x_n)_1^N$ be a tuple. Let $(\lambda_n)_1^N$ be a tuple of weighting values such that $\sum_{n=1}^N \lambda_n = 1$.

¹² Rudin (1976) page 109 (5.13 Theorem)

DEF

$$\text{The \textbf{harmonic mean} of } (\lambda_n) \text{ is defined as } \mu_h \triangleq \left(\sum_{n=1}^N \lambda_n \frac{1}{x_n} \right)^{-1}$$

$$\text{The \textbf{geometric mean} of } (\lambda_n) \text{ is defined as } \mu_g \triangleq \prod_{n=1}^N x_n^{\lambda_n}$$

$$\text{The \textbf{arithmetic mean} of } (\lambda_n) \text{ is defined as } \mu_a \triangleq \sum_{n=1}^N \lambda_n x_n$$

$$\text{The \textbf{average} of } (\lambda_n) \text{ is defined as } \mu_a \triangleq \frac{1}{N} \sum_{n=1}^N x_n$$

Corollary L.1. ¹³ Let $(\lambda_n)_1^N$ be a tuple. Let $(x_n)_1^N$ be a tuple of weighting values such that $\sum_{n=1}^N \lambda_n = 1$.

COR

$$\min(\lambda_n) \leq \underbrace{\left(\sum_{n=1}^N \lambda_n \frac{1}{x_n} \right)^{-1}}_{\text{harmonic mean}} \leq \underbrace{\prod_{n=1}^N x_n^{\lambda_n}}_{\text{geometric mean}} \leq \underbrace{\sum_{n=1}^N \lambda_n x_n}_{\text{arithmetic mean}} \leq \max(\lambda_n)$$

PROOF:

- These five means are all special cases of the *power mean* $M_{\phi(x:r)}$ (Definition L.3 page 270):

$r = \infty$:	$\max(\lambda_n)$
$r = 1$:	arithmetic mean
$r = 0$:	geometric mean
$r = -1$:	harmonic mean
$r = -\infty$:	$\min(\lambda_n)$
- The inequalities follow directly from Theorem L.3 (page 270).
- Generalized AM-GM inequality: If one is only concerned with the arithmetic mean and geometric mean, their relationship can be established directly using *Jensen's Inequality*:

$$\begin{aligned} \sum_{n=1}^N \lambda_n x_n &= b^{\log_b \left(\sum_{n=1}^N \lambda_n x_n \right)} \\ &\geq b^{\left(\sum_{n=1}^N \lambda_n \log_b x_n \right)} && \text{by Jensen's Inequality (Theorem K.1 page 262)} \\ &= \prod_{n=1}^N b^{\left(\lambda_n \log_b x_n \right)} \\ &= \prod_{n=1}^N b^{\left(\log_b x_n \right) \lambda_n} \\ &= \prod_{n=1}^N x_n^{\lambda_n} \end{aligned}$$

¹³ [Bullen \(2003\) page 71](#), [Bollobás \(1999\) page 5](#), [Cauchy \(1821\) pages 457–459](#) (Note II, theorem 17), [Jensen \(1906\) page 183](#)

Lemma L.2 (Young's Inequality). ¹⁴

LEM

$$\begin{aligned} xy &< \frac{x^p}{p} + \frac{y^q}{q} \quad \text{with } \frac{1}{p} + \frac{1}{q} = 1 \quad \forall 1 < p < \infty, x, y \geq 0, \quad \text{but } y \neq x^{p-1} \\ xy &= \frac{x^p}{p} + \frac{y^q}{q} \quad \text{with } \frac{1}{p} + \frac{1}{q} = 1 \quad \forall 1 < p < \infty, x, y \geq 0, \quad \text{and } y = x^{p-1} \end{aligned}$$

PROOF:

1. Proof that $\frac{1}{p-1} = q - 1$:

$$\begin{aligned} \frac{1}{p} + \frac{1}{q} = 1 &\iff \frac{q}{q} + \frac{q}{p} = q \\ &\iff q\left(1 - \frac{1}{p}\right) = 1 \\ &\iff q = \frac{1}{1 - \frac{1}{p}} \\ &\iff q = \frac{p}{p-1} \\ &\iff q - 1 = \frac{p}{p-1} - \frac{p-1}{p-1} \\ &\iff q - 1 = \frac{p - (p-1)}{p-1} \\ &\iff q - 1 = \frac{1}{p-1} \end{aligned}$$

2. Proof that $v = u^{p-1} \iff u = v^{q-1}$:

$$\begin{aligned} u &= v^{\frac{1}{p-1}} && \text{by left hypothesis} \\ &= v^{q-1} && \text{by item (1)} \end{aligned}$$


3. Proof that $v = u^{p-1}$ is monotonically increasing in u and $u = v^{q-1}$ is monotonically increasing in v :

$$\begin{aligned} \frac{dv}{du} &= \frac{d}{du} u^{p-1} \\ &= (p-1)u^{p-2} \\ &> 0 \\ \frac{du}{dv} &= \frac{d}{dv} v^{q-1} \\ &= (q-1)v^{q-2} \\ &> 0 \end{aligned}$$

4. Proof that $xy \leq \frac{x^p}{p} + \frac{y^q}{q}$:

¹⁴ Carothers (2000), page 43, Tolsted (1964), page 5, Maligranda (1995), page 257, Hardy et al. (1952) (Theorem 24), Young (1912) page 226

$$\begin{aligned} xy &\leq \int_0^x u^{p-1} du + \int_0^y v^{q-1} dv \\ &= \frac{u^p}{p} \Big|_0^x + \frac{v^q}{q} \Big|_0^y \\ &= \frac{x^p}{p} + \frac{y^q}{q} \end{aligned}$$



Theorem L.4 (Hölder's Inequality). ¹⁵ Let $(x_n \in \mathbb{C})_1^N$ and $(y_n \in \mathbb{C})_1^N$ be complex N -tuples.

T H M	$\underbrace{\sum_{n=1}^N x_n y_n }_{\ x \cdot y\ _1} \leq \underbrace{\left(\sum_{n=1}^N x_n ^p \right)^{\frac{1}{p}}}_{\ x\ _p} \underbrace{\left(\sum_{n=1}^N y_n ^q \right)^{\frac{1}{q}}}_{\ y\ _q} \quad \text{with} \quad \frac{1}{p} + \frac{1}{q} = 1 \quad \forall 1 < p < \infty$
-------------	---

PROOF: Let $\|(x_n)\|_p \triangleq \left(\sum_{n=1}^N |x_n|^p \right)^{\frac{1}{p}}$.

$$\begin{aligned} \sum_{n=1}^N |x_n y_n| &= \|(x_n)\|_p \|(y_n)\|_q \sum_{n=1}^N \frac{|x_n y_n|}{\|(x_n)\|_p \|(y_n)\|_q} \\ &= \|(x_n)\|_p \|(y_n)\|_q \sum_{n=1}^N \frac{|x_n|}{\|(x_n)\|_p} \frac{|y_n|}{\|(y_n)\|_q} \\ &\leq \|(x_n)\|_p \|(y_n)\|_q \sum_{n=1}^N \left(\frac{1}{p} \frac{|x_n|^p}{\|(x_n)\|_p^p} + \frac{1}{q} \frac{|y_n|^q}{\|(y_n)\|_q^q} \right) \quad \text{by Young's Inequality (Lemma L.2 page 275)} \\ &= \|(x_n)\|_p \|(y_n)\|_q \left(\frac{1}{p} \cdot \frac{\sum |x_n|^p}{\|(x_n)\|_p^p} + \frac{1}{q} \cdot \frac{\sum |y_n|^q}{\|(y_n)\|_q^q} \right) \\ &= \|(x_n)\|_p \|(y_n)\|_q \left(\frac{1}{p} \frac{\|(x_n)\|_p^p}{\|(x_n)\|_p^p} + \frac{1}{q} \frac{\|(y_n)\|_q^q}{\|(y_n)\|_q^q} \right) \\ &= \|(x_n)\|_p \|(y_n)\|_q \underbrace{\left(\frac{1}{p} + \frac{1}{q} \right)}_1 \end{aligned}$$

by definition of $\|\cdot\|$

by $\frac{1}{p} + \frac{1}{q} = 1$ constraint

Theorem L.5 (Minkowski's Inequality for sequences). ¹⁶ Let $(x_n \in \mathbb{C})_1^N$ and $(y_n \in \mathbb{C})_1^N$ be complex N -tuples.

¹⁵ [Bullen \(2003\)](#) page 178 (2.1), [Carothers \(2000\)](#), page 44, [Tolsted \(1964\)](#), page 6, [Maligranda \(1995\)](#), page 257, [Hardy et al. \(1952\)](#) (Theorem 11), [Hölder \(1889\)](#)

¹⁶ [Bullen \(2003\)](#) page 179, [Carothers \(2000\)](#), page 44, [Tolsted \(1964\)](#), page 7, [Maligranda \(1995\)](#), page 258, [Hardy et al. \(1952\)](#) (Theorem 24), [Minkowski \(1910\)](#) page 115

T H M

$$\left(\sum_{n=1}^N |x_n + y_n|^p \right)^{\frac{1}{p}} \leq \left(\sum_{n=1}^N |x_n|^p \right)^{\frac{1}{p}} + \left(\sum_{n=1}^N |y_n|^p \right)^{\frac{1}{p}} \quad \forall 1 < p < \infty$$

PROOF:

1. Define q in terms of p such that $\frac{1}{p} + \frac{1}{q} = 1$

2. Proof that $\frac{1}{q} = \frac{p-1}{p}$:

$$\begin{aligned} \frac{1}{p} + \frac{1}{q} = 1 &\iff \frac{1}{q} = 1 - \frac{1}{p} \\ &\iff \frac{1}{q} = \frac{p-1}{p} \end{aligned}$$

3. Define $\|\cdot\|$ as follows:

$$\|x\|_p \triangleq \left(\sum_{n=1}^N |x_n|^p \right)^{\frac{1}{p}}$$

4. Proof that $\|x_n + y_n\|_p \leq \|x_n\|_p + \|y_n\|_p$:

$$\begin{aligned} \|x_n + y_n\|_p^p &= \sum_{n=1}^N |x_n + y_n|^p && \text{by definition of } \|\cdot\|_p \\ &= \sum_{n=1}^N |x_n + y_n| |x_n + y_n|^{p-1} && \text{by homogeneous property of } |\cdot| \\ &\leq \sum_{n=1}^N |x_n| |x_n + y_n|^{p-1} + \sum_{n=1}^N |y_n| |x_n + y_n|^{p-1} && \text{by subadditive property of } |\cdot| \\ &= \sum_{n=1}^N |x_n(x_n + y_n)^{p-1}| + \sum_{n=1}^N |y_n(x_n + y_n)^{p-1}| && \text{by homogeneous property of } |\cdot| \\ &\leq \|x_n\|_p \|(x_n + y_n)^{p-1}\|_q + \|y_n\|_p \|(x_n + y_n)^{p-1}\|_q && \text{by Hölder's Inequality page 276} \\ &= (\|x_n\|_p + \|y_n\|_p) \|(x_n + y_n)^{p-1}\|_q \\ &= (\|x_n\|_p + \|y_n\|_p) \left(\sum_{n=1}^N |(x_n + y_n)^{p-1}|^q \right)^{\frac{1}{q}} && \text{by definition of } \|\cdot\|_p \\ &= (\|x_n\|_p + \|y_n\|_p) \left(\sum_{n=1}^N |(x_n + y_n)^{\frac{p}{p-1}}|^{\frac{p-1}{p}} \right)^{\frac{p-1}{p}} && \text{by item (2)} \\ &= (\|x_n\|_p + \|y_n\|_p) \left(\sum_{n=1}^N |(x_n + y_n)|^p \right)^{\frac{p-1}{p}} \\ &= (\|x_n\|_p + \|y_n\|_p) \|x_n + y_n\|^{p-1} \\ &\implies \|x_n + y_n\|_p \leq \|x_n\|_p + \|y_n\|_p \end{aligned}$$



“Cauchy is the only one occupied with pure mathematics: Poisson, Fourier, Ampere, etc., busy themselves exclusively with magnetism and other physical subjects.”¹⁷
 in an 1826 letter written by Niels Henrik Abel¹⁸

Theorem L.6 (Cauchy-Schwarz Inequality).¹⁸ Let $(x_n \in \mathbb{C})_1^N$ and $(y_n \in \mathbb{C})_1^N$ be complex N -tuples.

T H M	$\left \sum_{n=1}^N x_n y_n^* \right ^2 \leq \left(\sum_{n=1}^N x_n ^2 \right) \left(\sum_{n=1}^N y_n ^2 \right)$ $\left \sum_{n=1}^N x_n y_n^* \right ^2 = \left(\sum_{n=1}^N x_n ^2 \right) \left(\sum_{n=1}^N y_n ^2 \right) \iff \exists a \in \mathbb{C} \text{ such that } y = ax \quad \forall x, y \in X$
----------------------	---

PROOF:

1. The Cauchy-Schwarz inequality for sequences is a special case of the Hölder inequality (Theorem L.4) for $p = q = 2$.
2. Alternatively, the Cauchy-Schwarz inequality for sequences is a special case of the *Cauchy-Schwarz inequality* in inner-product spaces:
 - (a) $\langle x_n | y_n \rangle \triangleq \sum_{n=1}^N x_n y_n$ is an inner-product and $((\langle x_n | \rangle, \langle \cdot | \cdot \rangle))$ is an inner-product space.
 - (b) By the *Cauchy-Schwarz Inequality for inner-product spaces* (Theorem I.2 page 234),

$$|\langle x | y \rangle|^2 \leq \langle x | x \rangle \langle y | y \rangle$$

3. Not only does the Hölder inequality imply the Cauchy-Schwarz inequality, but somewhat surprisingly, the converse is also true: The Cauchy-Schwarz inequality implies the Hölder inequality.¹⁹

PROPOSITION L.1.²⁰

P R P	$(x + y)^p \leq 2^p(x^p + y^p) \quad \forall x, y \geq 0, 1 < p < \infty$
----------------------	---

PROOF:

$$\begin{aligned}
 (x + y)^p &\leq (2 \max \{x, y\})^p \\
 &= 2^p(\max \{x, y\})^p \\
 &= 2^p(\max \{x^p, y^p\}) \\
 &\leq 2^p(x^p + y^p)
 \end{aligned}$$

¹⁷ quote: [Boyer and Merzbach \(2011\) page 462](#)

image: http://en.wikipedia.org/wiki/File:Augustin-Louis_Cauchy_1901.jpg, public domain

¹⁸ [Aliprantis and Burkinshaw \(1998\)](#), page 278, [Scharz \(1885\)](#), [Bouniakowsky \(1859\)](#), [Hardy et al. \(1952\)](#) page 25 (Theorem 11), [Cauchy \(1821\)](#) page 455 (???)

¹⁹ [Bullen \(2003\)](#) pages 183–185 (Theorem 5)

²⁰ [Carothers \(2000\)](#), page 43

L.3 Power Sums

Theorem L.7 (Geometric Series). ²¹

T H M

$$\sum_{k=0}^{n-1} r^k = \frac{1 - r^n}{1 - r} \quad \forall r \in \mathbb{C} \setminus \{0\}$$

PROOF:

$$\begin{aligned} \sum_{k=0}^{n-1} r^k &= \left(\frac{1}{1-r} \right) \left[(1-r) \sum_{k=0}^{n-1} r^k \right] \\ &= \left(\frac{1}{1-r} \right) \left[\sum_{k=0}^{n-1} r^k - r \sum_{k=0}^{n-1} r^k \right] \\ &= \left(\frac{1}{1-r} \right) \left[\sum_{k=0}^{n-1} r^k - \left(\sum_{k=0}^{n-1} r^k - 1 + r^n \right) \right] \\ &= \left(\frac{1}{1-r} \right) [1 - r^n] \\ &= \frac{1 - r^n}{1 - r} \end{aligned}$$

Lemma L.3. Let $f(t)$ be a function.

L E M

$$S(x) \triangleq \sum_{n \in \mathbb{Z}} f(x + n\tau) = S(x + \tau) \quad (\text{S}(x) \text{ is PERIODIC with period } \tau)$$

PROOF:

$$\begin{aligned} S(x + \tau) &\triangleq \sum_{n \in \mathbb{Z}} f(x + \tau + n\tau) \\ &= \sum_{n \in \mathbb{Z}} f(x + (n+1)\tau) \\ &= \sum_{m \in \mathbb{Z}} f(x + m\tau) \quad \text{where } m \triangleq n + 1 \\ &\triangleq S(x) \end{aligned}$$

Proposition L.2 (Power Sums). ²²

P R P

$\sum_{k=1}^n k$	$= \frac{n(n+1)}{2}$	$\forall n \in \mathbb{N}$
$\sum_{k=1}^n k^2$	$= \frac{n(n+1)(2n+1)}{6}$	$\forall n \in \mathbb{N}$
$\sum_{k=1}^n k^3$	$= \frac{n^2(n+1)^2}{4}$	$\forall n \in \mathbb{N}$
$\sum_{k=1}^n k^4$	$= \frac{n(n+1)(2n+1)(3n^2+3n-1)}{30}$	$\forall n \in \mathbb{N}$

²¹ Hall and Knight (1894), page 39 (article 55)

²² Amann and Escher (2008) pages 51–57, Menini and Oystaeyen (2004) page 91 (Exercises 5.36–5.39)

PROOF:

1. Proof that $\sum_{k=1}^n k = \frac{n(n+1)}{2}$: (proof by induction)

$$\begin{aligned}\sum_{k=1}^{n=1} k &= 1 \\ &= \frac{1(1+1)}{2} \\ &= \left. \frac{n(n+1)}{2} \right|_{n=1}\end{aligned}$$

$$\begin{aligned}\sum_{k=1}^{n+1} k &= \left(\sum_{k=1}^n k \right) + (n+1) \\ &= \left(\frac{n(n+1)}{2} \right) + (n+1) \\ &= (n+1) \left(\frac{n}{2} + 1 \right) \\ &= (n+1) \left(\frac{n+2}{2} \right) \\ &= \frac{(n+1)(n+2)}{2}\end{aligned}$$

by left hypothesis

2. Proof that $\sum_{k=1}^n k^2 = \frac{n(n+1)(2n+1)}{6}$: (proof by induction)

$$\begin{aligned}\sum_{k=1}^{n=1} k^2 &= 1 \\ &= \frac{1(1+1)(2+1)}{6} \\ &= \left. \frac{n(n+1)(2n+1)}{6} \right|_{n=1}\end{aligned}$$

$$\begin{aligned}\sum_{k=1}^{n+1} k^2 &= \left(\sum_{k=1}^n k^2 \right) + (n+1)^2 \\ &= \left(\frac{n(n+1)(2n+1)}{6} \right) + (n+1)^2 \\ &= (n+1) \left(\frac{n(2n+1) + 6(n+1)}{6} \right) \\ &= (n+1) \left(\frac{2n^2 + 7n + 6}{6} \right) \\ &= (n+1) \left(\frac{(n+2)(2n+3)}{6} \right) \\ &= \frac{(n+1)[(n+1)+1][2(n+1)+1]}{6}\end{aligned}$$

by left hypothesis

APPENDIX M

OPERATORS ON LINEAR SPACES



“And I am not afraid to say that there is a way to advance algebra as far beyond what Vieta and Descartes have left us as Vieta and Descartes carried it beyond the ancients....we need still another analysis which is distinctly geometrical or linear, and which will express situation directly as algebra expresses magnitude directly.”

Gottfried Leibniz (1646–1716), German mathematician, in a September 8, 1679 letter to Christian Huygens.¹

M.1 Operators on linear spaces

M.1.1 Operator Algebra

An operator is simply a function that maps from a linear space to another linear space (or to the same linear space).

Definition M.1. ²

D E F A function A in Y^X is an **operator** in Y^X if
 X and Y are both LINEAR SPACES (Definition F.1 page 185).

Two operators A and B in Y^X are **equal** if $Ax = Bx$ for all $x \in X$. The inverse relation of an operator A in Y^X always exists as a *relation* in 2^{XY} , but may not always be a *function* (may not always be an operator) in Y^X .

The operator $I \in X^X$ is the *identity* operator if $Ix = I$ for all $x \in X$.

Definition M.2. ³ Let X^X be the set of all operators with from a LINEAR SPACE X to X . Let I be an operator in X^X . Let $\mathbb{I}(X)$ be the IDENTITY ELEMENT in X^X .

¹ quote: Leibniz (1679) pages 248–249

image: http://en.wikipedia.org/wiki/File:Gottfried_Wilhelm_von_Leibniz.jpg, public domain

² Heil (2011) page 42

³ Michel and Herget (1993) page 411




Figure M.1: Some operator types

DEF

I is the **identity operator** in $\mathbf{X}^{\mathbf{X}}$ if $\mathbf{I} = \mathbb{I}(\mathbf{X})$.

M.1.2 Linear operators

Definition M.3. ⁴ Let $\mathbf{X} \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}))$ and $\mathbf{Y} \triangleq (Y, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}))$ be linear spaces.

DEF

An operator $\mathbf{L} \in \mathbf{Y}^{\mathbf{X}}$ is **linear** if

1. $\mathbf{L}(x + y) = \mathbf{L}x + \mathbf{L}y \quad \forall x, y \in \mathbf{X}$ (ADDITIONAL) and
2. $\mathbf{L}(\alpha x) = \alpha \mathbf{L}x \quad \forall x \in \mathbf{X}, \alpha \in \mathbb{F}$ (HOMOGENEOUS).

The set of all linear operators from \mathbf{X} to \mathbf{Y} is denoted $\mathcal{L}(\mathbf{X}, \mathbf{Y})$ such that
 $\mathcal{L}(\mathbf{X}, \mathbf{Y}) \triangleq \{\mathbf{L} \in \mathbf{Y}^{\mathbf{X}} | \mathbf{L} \text{ is linear}\}$.

Theorem M.1. ⁵ Let \mathbf{L} be an operator from a linear space \mathbf{X} to a linear space \mathbf{Y} , both over a field \mathbb{F} .

THM

$$\mathbf{L} \text{ is LINEAR} \implies \begin{cases} 1. \mathbf{L}\emptyset = \emptyset & \text{and} \\ 2. \mathbf{L}(-x) = -(\mathbf{L}x) & \forall x \in \mathbf{X} \text{ and} \\ 3. \mathbf{L}(x - y) = \mathbf{L}x - \mathbf{L}y & \forall x, y \in \mathbf{X} \text{ and} \\ 4. \mathbf{L}\left(\sum_{n=1}^N \alpha_n x_n\right) = \sum_{n=1}^N \alpha_n (\mathbf{L}x_n) & x_n \in \mathbf{X}, \alpha_n \in \mathbb{F} \end{cases}$$

PROOF:

⁴ Kubrusly (2001) page 55, Aliprantis and Burkinshaw (1998) page 224, Hilbert et al. (1927) page 6, Stone (1932) page 33

⁵ Berberian (1961) page 79 (Theorem IV.1.1)

1. Proof that $\mathbf{L}\mathbf{0} = \mathbf{0}$:

$$\begin{aligned}\mathbf{L}\mathbf{0} &= \mathbf{L}(0 \cdot \mathbf{0}) && \text{by additive identity property (Theorem F.1 page 187)} \\ &= 0 \cdot (\mathbf{L}\mathbf{0}) && \text{by } \textit{homogeneous} \text{ property of } \mathbf{L} \text{ (Definition M.3 page 282)} \\ &= \mathbf{0} && \text{by additive identity property (Theorem F.1 page 187)}\end{aligned}$$

2. Proof that $\mathbf{L}(-\mathbf{x}) = -(\mathbf{Lx})$:

$$\begin{aligned}\mathbf{L}(-\mathbf{x}) &= \mathbf{L}(-1 \cdot \mathbf{x}) && \text{by additive inverse property (Theorem F.2 page 188)} \\ &= -1 \cdot (\mathbf{Lx}) && \text{by } \textit{homogeneous} \text{ property of } \mathbf{L} \text{ (Definition M.3 page 282)} \\ &= -(\mathbf{Lx}) && \text{by additive inverse property (Theorem F.2 page 188)}\end{aligned}$$

3. Proof that $\mathbf{L}(\mathbf{x} - \mathbf{y}) = \mathbf{Lx} - \mathbf{Ly}$:

$$\begin{aligned}\mathbf{L}(\mathbf{x} - \mathbf{y}) &= \mathbf{L}(\mathbf{x} + (-\mathbf{y})) && \text{by additive inverse property (Theorem F.2 page 188)} \\ &= \mathbf{L}(\mathbf{x}) + \mathbf{L}(-\mathbf{y}) && \text{by } \textit{linearity} \text{ property of } \mathbf{L} \text{ (Definition M.3 page 282)} \\ &= \mathbf{Lx} - \mathbf{Ly} && \text{by 2.}\end{aligned}$$

4. Proof that $\mathbf{L}\left(\sum_{n=1}^N \alpha_n \mathbf{x}_n\right) = \sum_{n=1}^N \alpha_n (\mathbf{Lx}_n)$:

(a) Proof for $N = 1$:

$$\begin{aligned}\mathbf{L}\left(\sum_{n=1}^N \alpha_n \mathbf{x}_n\right) &= \mathbf{L}(\alpha_1 \mathbf{x}_1) && \text{by } N = 1 \text{ hypothesis} \\ &= \alpha_1 (\mathbf{Lx}_1) && \text{by } \textit{homogeneous} \text{ property of Definition M.3 page 282}\end{aligned}$$

(b) Proof that N case $\implies N + 1$ case:

$$\begin{aligned}\mathbf{L}\left(\sum_{n=1}^{N+1} \alpha_n \mathbf{x}_n\right) &= \mathbf{L}\left(\alpha_{N+1} \mathbf{x}_{N+1} + \sum_{n=1}^N \alpha_n \mathbf{x}_n\right) \\ &= \mathbf{L}(\alpha_{N+1} \mathbf{x}_{N+1}) + \mathbf{L}\left(\sum_{n=1}^N \alpha_n \mathbf{x}_n\right) && \text{by } \textit{linearity} \text{ property of Definition M.3 page 282} \\ &= \alpha_{N+1} \mathbf{L}(\mathbf{x}_{N+1}) + \sum_{n=1}^N \mathbf{L}(\alpha_n \mathbf{x}_n) && \text{by left } N + 1 \text{ hypothesis} \\ &= \sum_{n=1}^{N+1} \mathbf{L}(\alpha_n \mathbf{x}_n)\end{aligned}$$



Theorem M.2. ⁶ Let $\mathcal{L}(\mathbf{X}, \mathbf{Y})$ be the set of all linear operators from a linear space \mathbf{X} to a linear space \mathbf{Y} . Let $\mathcal{N}(\mathbf{L})$ be the NULL SPACE of an operator \mathbf{L} in $\mathbf{Y}^{\mathbf{X}}$ and $\mathcal{I}(\mathbf{L})$ the IMAGE SET of \mathbf{L} in $\mathbf{Y}^{\mathbf{X}}$.

T H M	$\mathcal{L}(\mathbf{X}, \mathbf{Y})$ is a linear space $\mathcal{N}(\mathbf{L})$ is a linear subspace of \mathbf{X} $\mathcal{I}(\mathbf{L})$ is a linear subspace of \mathbf{Y}	(space of linear transforms) $\forall \mathbf{L} \in \mathbf{Y}^{\mathbf{X}}$ $\forall \mathbf{L} \in \mathbf{Y}^{\mathbf{X}}$
-------------	---	--

PROOF:

⁶ Michel and Herget (1993) pages 98–104, Berberian (1961) pages 80–85 (Theorem IV.1.4 and Theorem IV.3.1)

1. Proof that $\mathcal{N}(\mathbf{L})$ is a linear subspace of \mathbf{X} :

- (a) $0 \in \mathcal{N}(\mathbf{L}) \implies \mathcal{N}(\mathbf{L}) \neq \emptyset$
- (b) $\mathcal{N}(\mathbf{L}) \triangleq \{x \in \mathbf{X} | \mathbf{L}x = 0\} \subseteq \mathbf{X}$
- (c) $x + y \in \mathcal{N}(\mathbf{L}) \implies 0 = \mathbf{L}(x + y) = \mathbf{L}(y + x) \implies y + x \in \mathcal{N}(\mathbf{L})$
- (d) $\alpha \in \mathbb{F}, x \in \mathbf{X} \implies 0 = \mathbf{L}x \implies 0 = \alpha \mathbf{L}x \implies 0 = \mathbf{L}(\alpha x) \implies \alpha x \in \mathcal{N}(\mathbf{L})$

2. Proof that $\mathcal{I}(\mathbf{L})$ is a linear subspace of \mathbf{Y} :

- (a) $0 \in \mathcal{I}(\mathbf{L}) \implies \mathcal{I}(\mathbf{L}) \neq \emptyset$
- (b) $\mathcal{I}(\mathbf{L}) \triangleq \{y \in \mathbf{Y} | \exists x \in \mathbf{X} \text{ such that } y = \mathbf{L}x\} \subseteq \mathbf{Y}$
- (c) $x + y \in \mathcal{I}(\mathbf{L}) \implies \exists v \in \mathbf{X} \text{ such that } \mathbf{L}v = x + y = y + x \implies y + x \in \mathcal{I}(\mathbf{L})$
- (d) $\alpha \in \mathbb{F}, x \in \mathcal{I}(\mathbf{L}) \implies \exists x \in \mathbf{X} \text{ such that } y = \mathbf{L}x \implies \alpha y = \alpha \mathbf{L}x = \mathbf{L}(\alpha x) \implies \alpha x \in \mathcal{I}(\mathbf{L})$

⇒

Example M.1. ⁷ Let $C([a : b], \mathbb{R})$ be the set of all *continuous* functions from the closed real interval $[a : b]$ to \mathbb{R} .

E **X** $C([a : b], \mathbb{R})$ is a linear space.

Theorem M.3. ⁸ Let $\mathcal{L}(\mathbf{X}, \mathbf{Y})$ be the set of linear operators from a linear space \mathbf{X} to a linear space \mathbf{Y} . Let $\mathcal{N}(\mathbf{L})$ be the NULL SPACE of a linear operator $\mathbf{L} \in \mathcal{L}(\mathbf{X}, \mathbf{Y})$.

T	$\mathbf{L}x = \mathbf{L}y \iff x - y \in \mathcal{N}(\mathbf{L})$
H	\mathbf{L} is INJECTIVE $\iff \mathcal{N}(\mathbf{L}) = \{0\}$

PROOF:

1. Proof that $\mathbf{L}x = \mathbf{L}y \implies x - y \in \mathcal{N}(\mathbf{L})$:

$$\begin{aligned} \mathbf{L}(x - y) &= \mathbf{L}x - \mathbf{L}y && \text{by Theorem M.1 page 282} \\ &= 0 && \text{by left hypothesis} \\ &\implies x - y \in \mathcal{N}(\mathbf{L}) && \text{by definition of null space} \end{aligned}$$

2. Proof that $\mathbf{L}x = \mathbf{L}y \iff x - y \in \mathcal{N}(\mathbf{L})$:

$$\begin{aligned} \mathbf{L}y &= \mathbf{L}y + 0 && \text{by definition of linear space (Definition E.1 page 185)} \\ &= \mathbf{L}y + \mathbf{L}(x - y) && \text{by right hypothesis} \\ &= \mathbf{L}y + (\mathbf{L}x - \mathbf{L}y) && \text{by Theorem M.1 page 282} \\ &= (\mathbf{L}y - \mathbf{L}y) + \mathbf{L}x && \text{by associative and commutative properties (Definition E.1 page 185)} \\ &= \mathbf{L}x \end{aligned}$$

3. Proof that \mathbf{L} is *injective* $\iff \mathcal{N}(\mathbf{L}) = \{0\}$:

$$\begin{aligned} \mathbf{L} \text{ is injective} &\iff \{(\mathbf{L}x = \mathbf{L}y \iff x = y) \quad \forall x, y \in \mathbf{X}\} \\ &\iff \{[\mathbf{L}x - \mathbf{L}y = 0 \iff (x - y) = 0] \quad \forall x, y \in \mathbf{X}\} \\ &\iff \{[\mathbf{L}(x - y) = 0 \iff (x - y) = 0] \quad \forall x, y \in \mathbf{X}\} \\ &\iff \mathcal{N}(\mathbf{L}) = \{0\} \end{aligned}$$

⁷ Eidelman et al. (2004) page 3

⁸ Berberian (1961) page 88 (Theorem IV.1.4)

Theorem M.4.⁹ Let \mathcal{W} , \mathcal{X} , \mathcal{Y} , and \mathcal{Z} be linear spaces over a field \mathbb{F} .

T H M	1. $L(MN) = (LM)N$ 2. $L(M \dotplus N) = (LM) \dotplus (LN)$ 3. $(L \dotplus M)N = (LN) \dotplus (MN)$ 4. $\alpha(LM) = (\alpha L)M = L(\alpha M)$	$\forall L \in \mathcal{L}(\mathcal{Z}, \mathcal{W}), M \in \mathcal{L}(\mathcal{Y}, \mathcal{Z}), N \in \mathcal{L}(\mathcal{X}, \mathcal{Y})$ $\forall L \in \mathcal{L}(\mathcal{Y}, \mathcal{Z}), M \in \mathcal{L}(\mathcal{X}, \mathcal{Y}), N \in \mathcal{L}(\mathcal{X}, \mathcal{Y})$ $\forall L \in \mathcal{L}(\mathcal{Y}, \mathcal{Z}), M \in \mathcal{L}(\mathcal{Y}, \mathcal{Z}), N \in \mathcal{L}(\mathcal{X}, \mathcal{Y})$ $\forall L \in \mathcal{L}(\mathcal{Y}, \mathcal{Z}), M \in \mathcal{L}(\mathcal{X}, \mathcal{Y}), \alpha \in \mathbb{F}$	(ASSOCIATIVE) (LEFT DISTRIBUTIVE) (RIGHT DISTRIBUTIVE) (HOMOGENEOUS)
----------------------	---	--	---

PROOF:

1. Proof that $L(MN) = (LM)N$: Follows directly from property of *associative* operators.

2. Proof that $L(M \dotplus N) = (LM) \dotplus (LN)$:

$$\begin{aligned}
 [L(M \dotplus N)]x &= L[(M \dotplus N)x] \\
 &= L[(Mx) \dotplus (Nx)] \\
 &= [L(Mx)] \dotplus [L(Nx)] \quad \text{by } \textit{additive} \text{ property Definition M.3 page 282} \\
 &= [(LM)x] \dotplus [(LN)x]
 \end{aligned}$$

3. Proof that $(L \dotplus M)N = (LN) \dotplus (MN)$: Follows directly from property of *associative* operators.

4. Proof that $\alpha(LM) = (\alpha L)M$: Follows directly from *associative* property of linear operators.

5. Proof that $\alpha(LM) = L(\alpha M)$:

$$\begin{aligned}
 [\alpha(LM)]x &= \alpha[(LM)x] \\
 &= L[\alpha(Mx)] \quad \text{by } \textit{homogeneous} \text{ property Definition M.3 page 282} \\
 &= L[(\alpha M)x] \\
 &= [L(\alpha M)]x
 \end{aligned}$$

Theorem M.5 (Fundamental theorem of linear equations).  Michel and Herget (1993) page 99 Let $\mathcal{Y}^{\mathcal{X}}$ be the set of all operators from a linear space \mathcal{X} to a linear space \mathcal{Y} . Let $\mathcal{N}(L)$ be the NULL SPACE of an operator L in $\mathcal{Y}^{\mathcal{X}}$ and $\mathcal{I}(L)$ the IMAGE SET of L in $\mathcal{Y}^{\mathcal{X}}$ (Definition ?? page ??).

T H M	$\dim \mathcal{I}(L) + \dim \mathcal{N}(L) = \dim \mathcal{X}$ $\forall L \in \mathcal{Y}^{\mathcal{X}}$
----------------------	--

PROOF: Let $\{\psi_k | k = 1, 2, \dots, p\}$ be a basis for \mathcal{X} constructed such that $\{\psi_{p-n+1}, \psi_{p-n+2}, \dots, \psi_p\}$ is a basis for

⁹  Berberian (1961) page 88 (Theorem IV.5.1)

$\mathcal{N}(\mathbf{L})$.

Let $p \triangleq \dim \mathbf{X}$.

Let $n \triangleq \dim \mathcal{N}(\mathbf{L})$.

$$\begin{aligned}
 \dim \mathcal{I}(\mathbf{L}) &= \dim \{y \in \mathbf{Y} \mid \exists x \in \mathbf{X} \text{ such that } y = \mathbf{L}x\} \\
 &= \dim \left\{ y \in \mathbf{Y} \mid \exists (\alpha_1, \alpha_2, \dots, \alpha_p) \text{ such that } y = \mathbf{L} \sum_{k=1}^p \alpha_k \psi_k \right\} \\
 &= \dim \left\{ y \in \mathbf{Y} \mid \exists (\alpha_1, \alpha_2, \dots, \alpha_p) \text{ such that } y = \sum_{k=1}^p \alpha_k \mathbf{L}\psi_k \right\} \\
 &= \dim \left\{ y \in \mathbf{Y} \mid \exists (\alpha_1, \alpha_2, \dots, \alpha_p) \text{ such that } y = \sum_{k=1}^{p-n} \alpha_k \mathbf{L}\psi_k + \sum_{k=1}^n \alpha_k \mathbf{L}\psi_k \right\} \\
 &= \dim \left\{ y \in \mathbf{Y} \mid \exists (\alpha_1, \alpha_2, \dots, \alpha_p) \text{ such that } y = \sum_{k=1}^{p-n} \alpha_k \mathbf{L}\psi_k + \mathbb{0} \right\} \\
 &= p - n \\
 &= \dim \mathbf{X} - \dim \mathcal{N}(\mathbf{L})
 \end{aligned}$$

Note: This “proof” may be missing some necessary detail.



M.2 Operators on Normed linear spaces

M.2.1 Operator norm

Definition M.4. ¹⁰ Let $\mathcal{L}(\mathbf{X}, \mathbf{Y})$ be the space of linear operators over normed linear spaces \mathbf{X} and \mathbf{Y} .
¹¹

The **operator norm** $\|\cdot\|$ is defined as

$$\|\mathbf{A}\| \triangleq \sup_{x \in \mathbf{X}} \{\|\mathbf{Ax}\| \mid \|x\| \leq 1\} \quad \forall \mathbf{A} \in \mathcal{L}(\mathbf{X}, \mathbf{Y})$$

The pair $(\mathcal{L}(\mathbf{X}, \mathbf{Y}), \|\cdot\|)$ is the **normed space of linear operators** on (\mathbf{X}, \mathbf{Y}) .

Proposition M.1 (next) shows that the functional defined in Definition M.4 (previous) is a *norm*.¹²

Proposition M.1. ¹³ Let $(\mathcal{L}(\mathbf{X}, \mathbf{Y}), \|\cdot\|)$ be the normed space of linear operators over the normed linear spaces $\mathbf{X} \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \|\cdot\|)$ and $\mathbf{Y} \triangleq (Y, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \|\cdot\|)$.

The functional $\|\cdot\|$ is a **norm** on $\mathcal{L}(\mathbf{X}, \mathbf{Y})$. In particular,

- | | | |
|-------------|---|--|
| P
R
P | 1. $\ \mathbf{A}\ \geq 0$ (NON-NEGATIVE)
2. $\ \mathbf{A}\ = 0 \iff \mathbf{A} \stackrel{\circ}{=} \mathbb{0}$ (NONDEGENERATE)
3. $\ \alpha \mathbf{A}\ = \alpha \ \mathbf{A}\ $ (HOMOGENEOUS)
4. $\ \mathbf{A} \dot{+} \mathbf{B}\ \leq \ \mathbf{A}\ + \ \mathbf{B}\ $ (SUBADDITIVE). | and
and
and
and |
|-------------|---|--|

Moreover, $(\mathcal{L}(\mathbf{X}, \mathbf{Y}), \|\cdot\|)$ is a **normed linear space**.

¹⁰ Rudin (1991) page 92, Aliprantis and Burkinshaw (1998) page 225

¹¹ The operator norm notation $\|\cdot\|$ is introduced (as a Matrix norm) in

Horn and Johnson (1990), page 290

¹² norm $\|\cdot\|$: Definition J.1 (page 249)

¹³ Rudin (1991) page 93

PROOF:

1. Proof that $\|\mathbf{A}\| > 0$ for $\mathbf{A} \neq \mathbb{0}$:

$$\begin{aligned} \|\mathbf{A}\| &\triangleq \sup_{x \in X} \{\|\mathbf{Ax}\| \mid \|x\| \leq 1\} \\ &> 0 \end{aligned} \quad \text{by definition of } \|\cdot\| \text{ (Definition M.4 page 286)}$$

2. Proof that $\|\mathbf{A}\| = 0$ for $\mathbf{A} \stackrel{\circ}{=} \mathbb{0}$:

$$\begin{aligned} \|\mathbf{A}\| &\triangleq \sup_{x \in X} \{\|\mathbf{Ax}\| \mid \|x\| \leq 1\} \\ &= \sup_{x \in X} \{\|\mathbb{0}x\| \mid \|x\| \leq 1\} \\ &= 0 \end{aligned} \quad \text{by definition of } \|\cdot\| \text{ (Definition M.4 page 286)}$$

3. Proof that $\|\alpha\mathbf{A}\| = |\alpha| \|\mathbf{A}\|$:

$$\begin{aligned} \|\alpha\mathbf{A}\| &\triangleq \sup_{x \in X} \{\|\alpha\mathbf{Ax}\| \mid \|x\| \leq 1\} \\ &= \sup_{x \in X} \{|\alpha| \|\mathbf{Ax}\| \mid \|x\| \leq 1\} \\ &= |\alpha| \sup_{x \in X} \{\|\mathbf{Ax}\| \mid \|x\| \leq 1\} \\ &= |\alpha| \|\mathbf{A}\| \end{aligned} \quad \begin{array}{l} \text{by definition of } \|\cdot\| \text{ (Definition M.4 page 286)} \\ \text{by definition of } \|\cdot\| \text{ (Definition M.4 page 286)} \\ \text{by definition of sup} \\ \text{by definition of } \|\cdot\| \text{ (Definition M.4 page 286)} \end{array}$$

4. Proof that $\|\mathbf{A} + \mathbf{B}\| \leq \|\mathbf{A}\| + \|\mathbf{B}\|$:

$$\begin{aligned} \|\mathbf{A} + \mathbf{B}\| &\triangleq \sup_{x \in X} \{\|(A + B)x\| \mid \|x\| \leq 1\} \\ &= \sup_{x \in X} \{\|\mathbf{Ax} + \mathbf{Bx}\| \mid \|x\| \leq 1\} \\ &\leq \sup_{x \in X} \{\|\mathbf{Ax}\| + \|\mathbf{Bx}\| \mid \|x\| \leq 1\} \\ &\leq \sup_{x \in X} \{\|\mathbf{Ax}\| \mid \|x\| \leq 1\} + \sup_{x \in X} \{\|\mathbf{Bx}\| \mid \|x\| \leq 1\} \\ &\triangleq \|\mathbf{A}\| + \|\mathbf{B}\| \end{aligned} \quad \begin{array}{l} \text{by definition of } \|\cdot\| \text{ (Definition M.4 page 286)} \\ \text{by definition of } \|\cdot\| \text{ (Definition M.4 page 286)} \\ \text{by definition of } \|\cdot\| \text{ (Definition M.4 page 286)} \end{array}$$



Lemma M.1. Let $(\mathcal{L}(X, Y), \|\cdot\|)$ be the normed space of linear operators over normed linear spaces $X \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \|\cdot\|)$ and $Y \triangleq (Y, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \|\cdot\|)$.

L
E
M

$$\|\mathbf{L}\| = \sup_x \{\|\mathbf{Lx}\| \mid \|x\| = 1\} \quad \forall x \in \mathcal{L}(X, Y)$$

PROOF: ¹⁴

1. Proof that $\sup_x \{\|\mathbf{Lx}\| \mid \|x\| \leq 1\} \geq \sup_x \{\|\mathbf{Lx}\| \mid \|x\| = 1\}$:

$$\sup_x \{\|\mathbf{Lx}\| \mid \|x\| \leq 1\} \geq \sup_x \{\|\mathbf{Lx}\| \mid \|x\| = 1\} \quad \text{because } A \subseteq B \implies \sup A \leq \sup B$$

14



Many many thanks to former NCTU Ph.D. student Chien Yao (Chinese: 姚建; PinYin: Yáo Jiàn) for his brilliant help with this proof. (If you are viewing this text as a pdf file, zoom in on the figure to the left to see text from Chien Yao's 2007 April 16 email.)

2. Let the subset $Y \subsetneq X$ be defined as

$$Y \triangleq \left\{ y \in X \mid \begin{array}{l} 1. \quad \|Ly\| = \sup_{x \in X} \{\|Lx\| \mid \|x\| \leq 1\} \text{ and} \\ 2. \quad 0 < \|y\| \leq 1 \end{array} \right\}$$

3. Proof that $\sup_x \{\|Lx\| \mid \|x\| \leq 1\} \leq \sup_x \{\|Lx\| \mid \|x\| = 1\}$:

$$\begin{aligned} \sup_{x \in X} \{\|Lx\| \mid \|x\| \leq 1\} &= \|Ly\| && \text{by definition of set } Y \\ &= \frac{\|y\|}{\|y\|} \|Ly\| \\ &= \|y\| \left\| \frac{1}{\|y\|} Ly \right\| && \text{by homogeneous property (page 249)} \\ &= \|y\| \left\| L \frac{y}{\|y\|} \right\| && \text{by homogeneous property (page 282)} \\ &\leq \|y\| \sup_{y \in Y} \left\{ \left\| L \frac{y}{\|y\|} \right\| \right\} && \text{by definition of supremum} \\ &= \|y\| \sup_{y \in Y} \left\{ \left\| L \frac{y}{\|y\|} \right\| \mid \left\| \frac{y}{\|y\|} \right\| = 1 \right\} && \text{because } \left\| \frac{y}{\|y\|} \right\| = 1 \text{ for all } y \in Y \\ &\leq \sup_{y \in Y} \left\{ \left\| L \frac{y}{\|y\|} \right\| \mid \left\| \frac{y}{\|y\|} \right\| = 1 \right\} && \text{because } 0 < \|y\| \leq 1 \\ &\leq \sup_{x \in X} \{\|Lx\| \mid \|x\| = 1\} && \text{because } \frac{y}{\|y\|} \in X \quad \forall y \in Y \end{aligned}$$

4. By (1) and (3),

$$\sup_{x \in X} \{\|Lx\| \mid \|x\| \leq 1\} = \sup_{x \in X} \{\|Lx\| \mid \|x\| = 1\}$$



Proposition M.2. ¹⁵ Let \mathbf{I} be the identity operator in the normed space of linear operators $(\mathcal{L}(X, X), \|\cdot\|)$.

P R P $\|\mathbf{I}\| = 1$

PROOF:

$$\begin{aligned} \|\mathbf{I}\| &\triangleq \sup \{\|\mathbf{Ix}\| \mid \|x\| \leq 1\} && \text{by definition of } \|\cdot\| \text{ (Definition M.4 page 286)} \\ &= \sup \{\|x\| \mid \|x\| \leq 1\} && \text{by definition of } \mathbf{I} \text{ (Definition M.2 page 281)} \\ &= 1 \end{aligned}$$



Theorem M.6. ¹⁶ Let $(\mathcal{L}(X, Y), \|\cdot\|)$ be the normed space of linear operators over normed linear spaces X and Y .

T H M	$\ Lx\ \leq \ \mathbf{L}\ \ x\ \quad \forall L \in \mathcal{L}(X, Y), x \in X$
	$\ \mathbf{KL}\ \leq \ \mathbf{K}\ \ \mathbf{L}\ \quad \forall K, L \in \mathcal{L}(X, Y)$

¹⁵ Michel and Herget (1993) page 410

¹⁶ Rudin (1991) page 103, Aliprantis and Burkinshaw (1998) page 225

PROOF:

1. Proof that $\|\mathbf{L}x\| \leq \|\mathbf{L}\| \ \|x\|$:

$$\begin{aligned}
 \|\mathbf{L}x\| &= \frac{\|x\|}{\|x\|} \|\mathbf{L}x\| \\
 &= \|x\| \left\| \frac{1}{\|x\|} \mathbf{L}x \right\| \\
 &= \|x\| \left\| \mathbf{L} \frac{x}{\|x\|} \right\| \\
 &\triangleq \|x\| \|\mathbf{L}y\| \\
 &\leq \|x\| \sup_y \|\mathbf{L}y\| \\
 &= \|x\| \sup_y \{ \|\mathbf{L}y\| \mid \|y\| = 1 \} \\
 &\triangleq \|x\| \|\mathbf{L}\|
 \end{aligned}$$

by property of norms
by property of linear operators
where $y \triangleq \frac{x}{\|x\|}$
by definition of supremum
because $\|y\| = \left\| \frac{x}{\|x\|} \right\| = \frac{\|x\|}{\|x\|} = 1$
by definition of operator norm

2. Proof that $\|\mathbf{KL}\| \leq \|\mathbf{K}\| \ \|\mathbf{L}\|$:

$$\begin{aligned}
 \|\mathbf{KL}\| &\triangleq \sup_{x \in X} \{ \|(\mathbf{KL})x\| \mid \|x\| \leq 1 \} \\
 &= \sup_{x \in X} \{ \|\mathbf{K}(\mathbf{L}x)\| \mid \|x\| \leq 1 \} \\
 &\leq \sup_{x \in X} \{ \|\mathbf{K}\| \ \|\mathbf{L}x\| \mid \|x\| \leq 1 \} \\
 &\leq \sup_{x \in X} \{ \|\mathbf{K}\| \ \|\mathbf{L}\| \ \|x\| \mid \|x\| \leq 1 \} \\
 &= \sup_{x \in X} \{ \|\mathbf{K}\| \ \|\mathbf{L}\| \ 1 \mid \|x\| \leq 1 \} \\
 &= \|\mathbf{K}\| \ \|\mathbf{L}\|
 \end{aligned}$$

by Definition M.4 page 286 ($\|\cdot\|$)
by 1.
by 1.
by definition of sup
by definition of sup

M.2.2 Bounded linear operators

Definition M.5. ¹⁷ Let $(\mathcal{L}(X, Y), \|\cdot\|)$ be a normed space of linear operators.

D E F An operator \mathbf{B} is **bounded** if $\|\mathbf{B}\| < \infty$.

The quantity $\mathcal{B}(X, Y)$ is the set of all **bounded linear operators** on (X, Y) such that $\mathcal{B}(X, Y) \triangleq \{ \mathbf{L} \in \mathcal{L}(X, Y) \mid \|\mathbf{L}\| < \infty \}$.

Theorem M.7. ¹⁸ Let $(\mathcal{L}(X, Y), \|\cdot\|)$ be the set of linear operators over normed linear spaces $X \triangleq (X, +, \cdot, (\mathbb{F}, +, \times), \|\cdot\|)$ and $Y \triangleq (Y, +, \cdot, (\mathbb{F}, +, \times), \|\cdot\|)$.

The following conditions are all EQUIVALENT:

- | | |
|--------------|--|
| T H M | <ol style="list-style-type: none"> 1. \mathbf{L} is continuous at A SINGLE POINT $x_0 \in X$ $\forall \mathbf{L} \in \mathcal{L}(X, Y)$ 2. \mathbf{L} is CONTINUOUS (at every point $x \in X$) $\forall \mathbf{L} \in \mathcal{L}(X, Y)$ 3. $\ \mathbf{L}\ < \infty$ (\mathbf{L} is BOUNDED) $\forall \mathbf{L} \in \mathcal{L}(X, Y)$ 4. $\exists M \in \mathbb{R}$ such that $\ \mathbf{L}x\ \leq M \ x\ \quad \forall \mathbf{L} \in \mathcal{L}(X, Y), x \in X$ |
|--------------|--|

¹⁷ Rudin (1991) pages 92–93

¹⁸ Aliprantis and Burkinshaw (1998) page 227

PROOF:

1. Proof that 1 \implies 2:

$$\begin{aligned}
 \epsilon &> \|Lx - Lx_0\| && \text{by hypothesis 1} \\
 &= \|L(x - x_0)\| && \text{by linearity (Definition M.3 page 282)} \\
 &= \|L(x + y - x_0 - y)\| \\
 &= \|L(x + y) - L(x_0 + y)\| && \text{by linearity (Definition M.3 page 282)} \\
 \implies L &\text{ is continuous at point } x + y \\
 \implies L &\text{ is continuous at every point in } X && \text{(hypothesis 2)}
 \end{aligned}$$

2. Proof that 2 \implies 1: obvious.

3. Proof that 4 \implies 2:¹⁹

$$\begin{aligned}
 \|Lx\| \leq M \|x\| &\implies \|L(x - y)\| \leq M \|x - y\| && \text{by hypothesis 4} \\
 &\implies \|Lx - Ly\| \leq M \|x - y\| && \text{by linearity of } L \text{ (Definition M.3 page 282)} \\
 &\implies \|Lx - Ly\| \leq \epsilon \text{ whenever } M \|x - y\| < \epsilon \\
 &\implies \|Lx - Ly\| \leq \epsilon \text{ whenever } \|x - y\| < \frac{\epsilon}{M} && \text{(hypothesis 2)}
 \end{aligned}$$

4. Proof that 3 \implies 4:

$$\begin{aligned}
 \|Lx\| &\leq \underbrace{\|L\|}_M \|x\| && \text{by Theorem M.6 page 288} \\
 &= M \|x\| && \text{where } M \triangleq \|L\| < \infty \text{ (by hypothesis 1)}
 \end{aligned}$$

5. Proof that 1 \implies 3:²⁰

$$\begin{aligned}
 \|L\| = \infty &\implies \{\|Lx\| \mid \|x\| \leq 1\} = \infty \\
 &\implies \exists (x_n) \text{ such that } \|x_n\| = 1 \text{ and } \|L\| = \{\|Lx_n\| \mid \|x_n\| \leq 1\} = \infty \\
 &\implies \|x_n\| = 1 \text{ and } \infty = \|L\| = \|Lx_n\| \\
 &\implies \|x_n\| = 1 \text{ and } \|Lx_n\| \geq n \\
 &\implies \frac{1}{n} \|x_n\| = \frac{1}{n} \text{ and } \frac{1}{n} \|Lx_n\| \geq 1 \\
 &\implies \left\| \frac{x_n}{n} \right\| = \frac{1}{n} \text{ and } \left\| L \frac{x_n}{n} \right\| \geq 1 \\
 &\implies \lim_{n \rightarrow \infty} \left\| \frac{x_n}{n} \right\| = 0 \text{ and } \lim_{n \rightarrow \infty} \left\| L \frac{x_n}{n} \right\| \geq 1 \\
 \implies L &\text{ is not continuous at 0}
 \end{aligned}$$

But by hypothesis, L is continuous. So the statement $\|L\| = \infty$ must be *false* and thus $\|L\| < \infty$ (L is *bounded*).

¹⁹ Bollobás (1999), page 29

²⁰ Aliprantis and Burkinshaw (1998), page 227

M.2.3 Adjoint on normed linear spaces

Definition M.6. Let $\mathcal{B}(X, Y)$ be the space of bounded linear operators on normed linear spaces X and Y . Let X^* be the TOPOLOGICAL DUAL SPACE of X .

D E F B^* is the **adjoint** of an operator $B \in \mathcal{B}(X, Y)$ if
 $f(Bx) = [B^*f](x) \quad \forall f \in X^*, x \in X$

Theorem M.8. ²¹ Let $\mathcal{B}(X, Y)$ be the space of bounded linear operators on normed linear spaces X and Y .

T H M $(A + B)^* = A^* + B^* \quad \forall A, B \in \mathcal{B}(X, Y)$
 $(\lambda A)^* = \lambda A^* \quad \forall A, B \in \mathcal{B}(X, Y)$
 $(AB)^* = B^*A^* \quad \forall A, B \in \mathcal{B}(X, Y)$

PROOF:

$$[A + B]^*f(x) = f([A + B]x) \quad \text{by definition of adjoint} \quad (\text{Definition M.6 page 291})$$

$$[\lambda A]^*f(x) = f([\lambda A]x) \quad \text{by definition of adjoint} \quad (\text{Definition M.6 page 291})$$

$$[AB]^*f(x) = f([AB]x) \quad \text{by definition of adjoint} \quad (\text{Definition M.6 page 291})$$

Theorem M.9. ²² Let $\mathcal{B}(X, Y)$ be the space of bounded linear operators on normed linear spaces X and Y . Let B^* be the adjoint of an operator B .

T H M $\|B\| = \|B^*\| \quad \forall B \in \mathcal{B}(X, Y)$

PROOF:

$$\|B\| \triangleq \sup \{ \|Bx\| \mid \|x\| \leq 1 \} \quad \text{by Definition M.4 page 286}$$

$$\stackrel{?}{=} \sup \{ |g(Bx; y^*)| \mid \|x\| \leq 1, \|y^*\| \leq 1 \}$$

$$= \sup \{ |f(x; B^*y^*)| \mid \|x\| \leq 1, \|y^*\| \leq 1 \}$$

$$\triangleq \sup \{ \|B^*y^*\| \mid \|x\| \leq 1, \|y^*\| \leq 1 \}$$

$$= \sup \{ \|B^*y^*\| \mid \|y^*\| \leq 1 \}$$

$$\triangleq \|B^*\|$$

by Definition M.4 page 286

²¹ Bollobás (1999), page 156

²² Rudin (1991) page 98

M.2.4 More properties



“Beginning with the third year of studies, most of my mathematical work was really started in conversations with Mazur and Banach. And according to Banach some of my own contributions were characterized by a certain “strangeness” in the formulation of problems and in the outline of possible proofs. As he told me once some years later, he was surprised how often these “strange” approaches really worked.”²³

Stanislaus M. Ulam (1909–1984), Polish mathematician ²³

Theorem M.10 (Mazur-Ulam theorem). ²⁴ Let $\phi \in \mathcal{L}(\mathbf{X}, \mathbf{Y})$ be a function on normed linear spaces $(\mathbf{X}, \|\cdot\|_{\mathbf{X}})$ and $(\mathbf{Y}, \|\cdot\|_{\mathbf{Y}})$. Let $\mathbf{I} \in \mathcal{L}(\mathbf{X}, \mathbf{X})$ be the identity operator on $(\mathbf{X}, \|\cdot\|_{\mathbf{X}})$.

T
H
M

$$\left. \begin{array}{l} 1. \underbrace{\phi^{-1}\phi = \phi\phi^{-1} = \mathbf{I}}_{\text{bijective}} \\ 2. \underbrace{\|\phi\mathbf{x} - \phi\mathbf{y}\|_{\mathbf{Y}} = \|\mathbf{x} - \mathbf{y}\|_{\mathbf{X}}}_{\text{isometric}} \end{array} \right\} \text{and } \Rightarrow \underbrace{\phi([1 - \lambda]\mathbf{x} + \lambda\mathbf{y}) = [1 - \lambda]\phi\mathbf{x} + \lambda\phi\mathbf{y} \forall \lambda \in \mathbb{R}}_{\text{affine}}$$

PROOF: Proof not yet complete.

1. Let ψ be the reflection of \mathbf{z} in \mathbf{X} such that $\psi\mathbf{x} = 2\mathbf{z} - \mathbf{x}$

$$(a) \|\psi\mathbf{x} - \mathbf{z}\| = \|\mathbf{x} - \mathbf{z}\|$$

2. Let $\lambda \triangleq \sup_g \{\|g\mathbf{z} - \mathbf{z}\|\}$

3. Proof that $g \in W \implies g^{-1} \in W$:

Let $\hat{\mathbf{x}} \triangleq g^{-1}\mathbf{x}$ and $\hat{\mathbf{y}} \triangleq g^{-1}\mathbf{y}$.

$$\begin{aligned} \|g^{-1}\mathbf{x} - g^{-1}\mathbf{y}\| &= \|\hat{\mathbf{x}} - \hat{\mathbf{y}}\| && \text{by definition of } \hat{\mathbf{x}} \text{ and } \hat{\mathbf{y}} \\ &= \|g\hat{\mathbf{x}} - g\hat{\mathbf{y}}\| && \text{by left hypothesis} \\ &= \|gg^{-1}\mathbf{x} - gg^{-1}\mathbf{y}\| && \text{by definition of } \hat{\mathbf{x}} \text{ and } \hat{\mathbf{y}} \\ &= \|\mathbf{x} - \mathbf{y}\| && \text{by definition of } g^{-1} \end{aligned}$$

²³ quote: [Ulam \(1991\)](#), page 33

image: <http://www-history.mcs.st-andrews.ac.uk/Biographies/Ulam.html>

²⁴ [Oikhberg and Rosenthal \(2007\)](#), page 598, [Väisälä \(2003\)](#), page 634, [Giles \(2000\)](#), page 11, [Dunford and Schwartz \(1957\)](#), page 91, [Mazur and Ulam \(1932\)](#)

4. Proof that $gz = z$:

$$\begin{aligned}
 2\lambda &= 2 \sup \{ \|gz - z\| \} && \text{by definition of } \lambda \text{ item (2)} \\
 &\leq 2 \|gz - z\| && \text{by definition of sup} \\
 &= \|2z - 2gz\| \\
 &= \|\psi gz - gz\| && \text{by definition of } \psi \text{ item (1)} \\
 &= \|g^{-1}\psi gz - g^{-1}gz\| && \text{by item (3)} \\
 &= \|g^{-1}\psi gz - z\| && \text{by definition of } g^{-1} \\
 &= \|\psi g^{-1}\psi gz - z\| \\
 &= \|g^*z - z\| \\
 &\leq \lambda && \text{by definition of } \lambda \text{ item (2)} \\
 &\implies 2\lambda \leq \lambda \\
 &\implies \lambda = 0 \\
 &\implies gz = z
 \end{aligned}$$

5. Proof that $\phi\left(\frac{1}{2}\mathbf{x} + \frac{1}{2}\mathbf{y}\right) = \frac{1}{2}\phi\mathbf{x} + \frac{1}{2}\phi\mathbf{y}$:

$$\begin{aligned}
 \phi\left(\frac{1}{2}\mathbf{x} + \frac{1}{2}\mathbf{y}\right) &= \\
 &= \frac{1}{2}\phi\mathbf{x} + \frac{1}{2}\phi\mathbf{y}
 \end{aligned}$$

6. Proof that $\phi([1 - \lambda]\mathbf{x} + \lambda\mathbf{y}) = [1 - \lambda]\phi\mathbf{x} + \lambda\phi\mathbf{y}$:

$$\begin{aligned}
 \phi([1 - \lambda]\mathbf{x} + \lambda\mathbf{y}) &= \\
 &= [1 - \lambda]\phi\mathbf{x} + \lambda\phi\mathbf{y}
 \end{aligned}$$

Theorem M.11 (Neumann Expansion Theorem). ²⁵ Let $\mathbf{A} \in \mathbf{X}^\mathbf{X}$ be an operator on a linear space \mathbf{X} . Let $\mathbf{A}^0 \triangleq \mathbf{I}$.

T H M	$ \left. \begin{array}{l} 1. \quad \mathbf{A} \in \mathcal{B}(\mathbf{X}, \mathbf{X}) \quad (\mathbf{A} \text{ is bounded}) \\ 2. \quad \ \mathbf{A}\ < 1 \end{array} \right\} \implies \left\{ \begin{array}{ll} 1. & (\mathbf{I} - \mathbf{A})^{-1} \quad \text{exists} \\ 2. & \ (\mathbf{I} - \mathbf{A})^{-1}\ \leq \frac{1}{1 - \ \mathbf{A}\ } \\ 3. & (\mathbf{I} - \mathbf{A})^{-1} = \sum_{n=0}^{\infty} \mathbf{A}^n \\ & \text{with uniform convergence} \end{array} \right. $
-------------	--

M.3 Operators on Inner product spaces

M.3.1 General Results

Theorem M.12. ²⁶ Let $\mathbf{A}, \mathbf{B} \in \mathcal{B}(\mathbf{X}, \mathbf{X})$ be bounded linear operators on an inner product space $\mathbf{X} \triangleq (X, +, \cdot, (\mathbb{F}, +, \hat{\times}), \langle \triangle | \triangleright \rangle)$.

T H M	$ \begin{array}{lll} \langle \mathbf{Bx} x \rangle = 0 & \forall x \in X & \iff \mathbf{Bx} = \mathbf{0} \quad \forall x \in X \\ \langle \mathbf{Ax} x \rangle = \langle \mathbf{Bx} x \rangle & \forall x \in X & \iff \mathbf{A} = \mathbf{B} \end{array} $
-------------	---

²⁵ Michel and Herget (1993) page 415

²⁶ Rudin (1991) page 310 (Theorem 12.7, Corollary)

PROOF:

1. Proof that $\langle \mathbf{Bx} | x \rangle = 0 \implies \mathbf{Bx} = \mathbb{0}$:

$$\begin{aligned}
 0 &= \langle \mathbf{B}(x + \mathbf{Bx}) | (x + \mathbf{Bx}) \rangle + i \langle \mathbf{B}(x + i\mathbf{Bx}) | (x + i\mathbf{Bx}) \rangle && \text{by left hypothesis} \\
 &= \{\langle \mathbf{Bx} + \mathbf{B}^2 x | x + \mathbf{Bx} \rangle\} + i\{\langle \mathbf{Bx} + i\mathbf{B}^2 x | x + i\mathbf{Bx} \rangle\} && \text{by Definition M.3 page 282} \\
 &= \{\langle \mathbf{Bx} | x \rangle + \langle \mathbf{Bx} | \mathbf{Bx} \rangle + \langle \mathbf{B}^2 x | x \rangle + \langle \mathbf{B}^2 x | \mathbf{Bx} \rangle\} && \text{by Definition I.1 page 233} \\
 &\quad + i\{\langle \mathbf{Bx} | x \rangle - i\langle \mathbf{Bx} | \mathbf{Bx} \rangle + i\langle \mathbf{B}^2 x | x \rangle - i^2 \langle \mathbf{B}^2 x | \mathbf{Bx} \rangle\} \\
 &= \{0 + \langle \mathbf{Bx} | \mathbf{Bx} \rangle + \langle \mathbf{B}^2 x | x \rangle + 0\} + i\{0 - i\langle \mathbf{Bx} | \mathbf{Bx} \rangle + i\langle \mathbf{B}^2 x | x \rangle - i^2 0\} && \text{by left hypothesis} \\
 &= \{\langle \mathbf{Bx} | \mathbf{Bx} \rangle + \langle \mathbf{B}^2 x | x \rangle\} + \{\langle \mathbf{Bx} | \mathbf{Bx} \rangle - \langle \mathbf{B}^2 x | x \rangle\} \\
 &= 2\langle \mathbf{Bx} | \mathbf{Bx} \rangle \\
 &= 2\|\mathbf{Bx}\|^2 \\
 &\implies \mathbf{Bx} = \mathbb{0} && \text{by Definition J.1 page 249}
 \end{aligned}$$

2. Proof that $\langle \mathbf{Bx} | x \rangle = 0 \iff \mathbf{Bx} = \mathbb{0}$: by property of inner products (Theorem I.1 page 233).

3. Proof that $\langle \mathbf{Ax} | x \rangle = \langle \mathbf{Bx} | x \rangle \implies \mathbf{A} \doteq \mathbf{B}$:

$$\begin{aligned}
 0 &= \langle \mathbf{Ax} | x \rangle - \langle \mathbf{Bx} | x \rangle && \text{by left hypothesis} \\
 &= \langle \mathbf{Ax} - \mathbf{Bx} | x \rangle && \text{by } \textit{additivity} \text{ property of } \langle \triangle | \nabla \rangle \text{ (Definition I.1 page 233)} \\
 &= \langle (\mathbf{A} - \mathbf{B})x | x \rangle && \text{by definition of operator addition} \\
 \implies &(\mathbf{A} - \mathbf{B})x = \mathbb{0} && \text{by item 1} \\
 \implies &\mathbf{A} = \mathbf{B} && \text{by definition of operator subtraction}
 \end{aligned}$$

4. Proof that $\langle \mathbf{Ax} | x \rangle = \langle \mathbf{Bx} | x \rangle \iff \mathbf{A} \doteq \mathbf{B}$:

$$\langle \mathbf{Ax} | x \rangle = \langle \mathbf{Bx} | x \rangle \quad \text{by } \mathbf{A} \doteq \mathbf{B} \text{ hypothesis}$$

⇒

M.3.2 Operator adjoint

A fundamental concept of operators on inner product spaces is the *operator adjoint* (Proposition M.3 page 294). The adjoint of an operator is a kind of generalization of the conjugate of a complex number in that

- Both are *star-algebras* (Theorem M.13 page 295).
- Both support decomposition into “real” and “imaginary” parts (Theorem H.3 page 230).

Structurally, the operator adjoint provides a convenient symmetric relationship between the *range space* and *null space* of an operator (Theorem M.14 page 296).

Proposition M.3. ²⁷ Let $\mathcal{B}(\mathbf{H}, \mathbf{H})$ be the space of bounded linear operators on a Hilbert space \mathbf{H} .²⁸

P R P An operator \mathbf{B}^* is the ADJOINT of $\mathbf{B} \in \mathcal{B}(\mathbf{H}, \mathbf{H})$ if
 $\langle \mathbf{Bx} | y \rangle = \langle x | \mathbf{B}^* y \rangle \quad \forall x, y \in \mathbf{H}$.

²⁷ Michel and Herget (1993) page 220, Rudin (1991) page 311, Giles (2000), page 182, von Neumann (1929) page 49, Stone (1932) page 41

²⁸ bounded operator: Definition M.5 (page 289); adjoint: Definition M.6 (page 291)

PROOF:

1. For fixed y , $f(x) \triangleq \langle x | y \rangle$ is a *functional* in \mathbb{F}^X .

2. \mathbf{B}^* is the *adjoint* of \mathbf{B} because

$$\begin{aligned}\langle \mathbf{B}x | y \rangle &\triangleq f(\mathbf{B}x) \\ &\triangleq \mathbf{B}^*f(x) \\ &= \langle x | \mathbf{B}^*y \rangle\end{aligned}\quad \text{by Definition M.6 (page 291)}$$

Example M.2 (Matrix algebra: $A^* = A^H$). In matrix algebra,

E
X

- 4 The inner product operation $\langle x | y \rangle$ is represented by $y^H x$.
- 4 The linear operator is represented as a matrix A .
- 4 The operation of A on vector x is represented as Ax .
- 4 The adjoint of matrix A is the Hermitian matrix A^H .

PROOF:

$$\langle Ax | y \rangle \triangleq y^H Ax = [(Ax)^H y]^H = [x^H A^H y]^H = (A^H y)^H x \triangleq \langle x | A^H y \rangle$$

Structures that satisfy the four conditions of the next theorem are known as **-algebras* ("star-algebras", Definition H.3 page 228). Other structures which are *-algebras include the *field of complex numbers* \mathbb{C} and any *ring of complex square $n \times n$ matrices*.²⁹

Theorem M.13 (operator star-algebra). ³⁰ Let H be a Hilbert space with operators $\mathbf{A}, \mathbf{B} \in \mathcal{B}(H, H)$ and with adjoints $\mathbf{A}^*, \mathbf{B}^* \in \mathcal{B}(H, H)$. Let $\bar{\alpha}$ be the complex conjugate of some $\alpha \in \mathbb{C}$.

T
H
M

The pair $(H, *)$ is a **-ALGEBRA* (STAR-ALGEBRA). In particular,

1. $(\mathbf{A} + \mathbf{B})^* = \mathbf{A}^* + \mathbf{B}^* \quad \forall \mathbf{A}, \mathbf{B} \in H$ (DISTRIBUTIVE) and
2. $(\alpha \mathbf{A})^* = \bar{\alpha} \mathbf{A}^* \quad \forall \mathbf{A} \in H$ (CONJUGATE LINEAR) and
3. $(\mathbf{AB})^* = \mathbf{B}^* \mathbf{A}^* \quad \forall \mathbf{A}, \mathbf{B} \in H$ (ANTIAUTOMORPHIC) and
4. $\mathbf{A}^{**} = \mathbf{A} \quad \forall \mathbf{A} \in H$ (INVOLUTARY)

PROOF:

$$\begin{aligned}\langle x | (\mathbf{A} + \mathbf{B})^* y \rangle &= \langle (\mathbf{A} + \mathbf{B})x | y \rangle && \text{by definition of adjoint} && \text{(Proposition M.3 page 294)} \\ &= \langle \mathbf{Ax} | y \rangle + \langle \mathbf{Bx} | y \rangle && \text{by definition of inner product} && \text{(Definition I.1 page 233)} \\ &= \langle x | \mathbf{A}^* y \rangle + \langle x | \mathbf{B}^* y \rangle && \text{by definition of operator addition} && \\ &= \langle x | \mathbf{A}^* y + \mathbf{B}^* y \rangle && \text{by definition of inner product} && \text{(Definition I.1 page 233)} \\ &= \langle x | (\mathbf{A}^* + \mathbf{B}^*) y \rangle && \text{by definition of operator addition} && \end{aligned}$$

$$\begin{aligned}\langle x | (\alpha \mathbf{A})^* y \rangle &= \langle (\alpha \mathbf{A})x | y \rangle && \text{by definition of adjoint} && \text{(Proposition M.3 page 294)} \\ &= \langle \alpha (\mathbf{Ax}) | y \rangle && \text{by definition of scalar multiplication} && \\ &= \alpha \langle \mathbf{Ax} | y \rangle && \text{by definition of inner product} && \text{(Definition I.1 page 233)} \\ &= \alpha \langle x | \mathbf{A}^* y \rangle && \text{by definition of adjoint} && \text{(Proposition M.3 page 294)}\end{aligned}$$

²⁹ Sakai (1998) page 1

³⁰ Halmos (1998), pages 39–40, Rudin (1991) page 311

$$= \langle x | \alpha^* A^* y \rangle \quad \text{by definition of inner product} \quad (\text{Definition I.1 page 233})$$

$$\begin{aligned} \langle x | (AB)^* y \rangle &= \langle (AB)x | y \rangle && \text{by definition of adjoint} && (\text{Proposition M.3 page 294}) \\ &= \langle A(Bx) | y \rangle && \text{by definition of operator multiplication} && \\ &= \langle (Bx) | A^* y \rangle && \text{by definition of adjoint} && (\text{Proposition M.3 page 294}) \\ &= \langle x | B^* A^* y \rangle && \text{by definition of adjoint} && (\text{Proposition M.3 page 294}) \end{aligned}$$

$$\begin{aligned} \langle x | A^{**} y \rangle &= \langle A^* x | y \rangle && \text{by definition of adjoint} && (\text{Proposition M.3 page 294}) \\ &= \langle y | A^* x \rangle^* && \text{by definition of inner product} && (\text{Definition I.1 page 233}) \\ &= \langle Ay | x \rangle^* && \text{by definition of adjoint} && (\text{Proposition M.3 page 294}) \\ &= \langle x | Ay \rangle && \text{by definition of inner product} && (\text{Definition I.1 page 233}) \end{aligned}$$

⇒

Theorem M.14. ³¹ Let \mathcal{Y}^X be the set of all operators from a linear space X to a linear space Y . Let $\mathcal{N}(L)$ be the NULL SPACE of an operator L in \mathcal{Y}^X and $\mathcal{I}(L)$ the IMAGE SET of L in \mathcal{Y}^X .

T	$\mathcal{N}(A) = \mathcal{I}(A^*)^\perp$
H	
M	$\mathcal{N}(A^*) = \mathcal{I}(A)^\perp$

PROOF:

$$\begin{aligned} \mathcal{I}(A^*)^\perp &= \{y \in H \mid \langle y | u \rangle = 0 \quad \forall u \in \mathcal{I}(A^*)\} \\ &= \{y \in H \mid \langle y | A^* x \rangle = 0 \quad \forall x \in H\} \\ &= \{y \in H \mid \langle Ay | x \rangle = 0 \quad \forall x \in H\} \quad \text{by definition of } A^* && (\text{Proposition M.3 page 294}) \\ &= \{y \in H \mid Ay = 0\} \\ &= \mathcal{N}(A) \quad \text{by definition of } \mathcal{N}(A) \end{aligned}$$

$$\begin{aligned} \mathcal{I}(A)^\perp &= \{y \in H \mid \langle y | u \rangle = 0 \quad \forall u \in \mathcal{I}(A)\} \\ &= \{y \in H \mid \langle y | Ax \rangle = 0 \quad \forall x \in H\} \quad \text{by definition of } \mathcal{I} \\ &= \{y \in H \mid \langle A^* y | x \rangle = 0 \quad \forall x \in H\} \quad \text{by definition of } A^* && (\text{Proposition M.3 page 294}) \\ &= \{y \in H \mid A^* y = 0\} \\ &= \mathcal{N}(A^*) \quad \text{by definition of } \mathcal{N}(A) \end{aligned}$$

⇒

M.4 Special Classes of Operators

M.4.1 Projection operators

Definition M.7. ³² Let $\mathcal{B}(X, Y)$ be the space of bounded linear operators on normed linear spaces X and Y . Let P be a bounded linear operator in $\mathcal{B}(X, Y)$.

D	
E	P is a projection operator if $P^2 = P$.
F	

³¹ Rudin (1991) page 312

³² Rudin (1991) page 133 (5.15 Projections), Kubrusly (2001) page 70, Bachman and Narici (1966) page 6, Halmos (1958) page 73 (§41. Projections)

Theorem M.15. ³³ Let $\mathcal{B}(X, Y)$ be the space of bounded linear operators on normed linear spaces X and Y . Let P be a bounded linear operator in $\mathcal{B}(X, Y)$ with NULL SPACE $\mathcal{N}(P)$ and IMAGE SET $I(P)$.

T H M	1. $P^2 = P$ (P is a projection operator) and 2. $\Omega = X \hat{+} Y$ (Y complements X in Ω) and 3. $P\Omega = X$ (P projects onto X)	$\left\{ \begin{array}{l} 1. \quad I(P) = X \\ 2. \quad \mathcal{N}(P) = Y \\ 3. \quad \Omega = I(P) \hat{+} \mathcal{N}(P) \end{array} \right.$ and and
----------------------	---	---

PROOF:

$$\begin{aligned} I(P) &= P\Omega \\ &= P(\Omega_1 + \Omega_2) \\ &= P\Omega_1 + P\Omega_2 \\ &= \Omega_1 + \{0\} \\ &= \Omega_1 \end{aligned}$$

$$\begin{aligned} \mathcal{N}(P) &= \{x \in \Omega | Px = 0\} \\ &= \{x \in (\Omega_1 + \Omega_2) | Px = 0\} \\ &= \{x \in \Omega_1 | Px = 0\} + \{x \in \Omega_2 | Px = 0\} \\ &= \{0\} + \Omega_2 \\ &= \Omega_2 \end{aligned}$$



Theorem M.16. ³⁴ Let $\mathcal{B}(X, Y)$ be the space of bounded linear operators on normed linear spaces X and Y . Let P be a bounded linear operator in $\mathcal{B}(X, Y)$.

T H M	$\underbrace{P^2 = P}_{P \text{ is a projection operator}} \iff \underbrace{(I - P)^2 = (I - P)}_{(I - P) \text{ is a projection operator}}$
----------------------	--

PROOF:

Proof that $P^2 = P \implies (I - P)^2 = (I - P)$:

$$\begin{aligned} (I - P)^2 &= (I - P)(I - P) \\ &= I(I - P) + (-P)(I - P) \\ &= I - P - PI + P^2 \\ &= I - P - P + P \quad \text{by left hypothesis} \\ &= I - P \end{aligned}$$

Proof that $P^2 = P \iff (I - P)^2 = (I - P)$:

$$\begin{aligned} P^2 &= \underbrace{I - P - P + P^2}_{(I - P)^2} - (I - P - P) \\ &= (I - P)^2 - (I - P - P) \\ &= (I - P) - (I - P - P) \quad \text{by right hypothesis} \\ &= P \end{aligned}$$



³³ Michel and Herget (1993) pages 120–121

³⁴ Michel and Herget (1993) page 121

M.4.2 Self Adjoint Operators

Definition M.8. ³⁵ Let $\mathbf{B} \in \mathcal{B}(H, H)$ be a bounded operator with adjoint \mathbf{B}^* on a Hilbert space H .

D E F The operator \mathbf{B} is said to be **self-adjoint** or **hermitian** if $\mathbf{B} \doteq \mathbf{B}^*$.

Example M.3 (Autocorrelation operator). Let $x(t)$ be a random process with autocorrelation

$$R_{xx}(t, u) \triangleq \underbrace{E[x(t)x^*(u)]}_{\text{expectation}}$$

Let an autocorrelation operator \mathbf{R} be defined as $[\mathbf{R}f](t) \triangleq \int_{\mathbb{R}} R_{xx}(t, u)f(u) du$.

E X $\mathbf{R} = \mathbf{R}^*$ (The autocorrelation operator \mathbf{R} is *self-adjoint*)

PROOF:

1. First note that the *autocorrelation kernel* $R_{xx}(t, u)$ is *hermitian symmetric*:

$$\begin{aligned} R_{xx}(t, u) &\triangleq E[x(t)x^*(u)] = [Ex^*(t)x(u)]^* = [Ex(u)x^*(t)]^* \\ &= R_{xx}^*(u, t) \end{aligned}$$

2. Proof that the *autocorrelation operator* \mathbf{R} is *self-adjoint*:

$$\begin{aligned} \langle \mathbf{R}f | g \rangle &= \left\langle \int_{u \in \mathbb{R}} R_{xx}(t, u)f(u) du | g(t) \right\rangle && \text{by definition of } \mathbf{R} \\ &= \int_{u \in \mathbb{R}} f(u) \langle R_{xx}(t, u) | g(t) \rangle du \\ &= \int_{u \in \mathbb{R}} f(u) \int_t R_{xx}(t, u)g^*(t) dt du \\ &= \int_{u \in \mathbb{R}} f(u) \int_t R_{xx}^*(u, t)g^*(t) dt du && \text{by 1.} \\ &= \int_{u \in \mathbb{R}} f(u) \left[\int_t R_{xx}(u, t)g(t) dt \right]^* du \\ &= \int_{u \in \mathbb{R}} f(u) [\mathbf{R}g]^* du && \text{by definition of } \mathbf{R} \\ &= \langle f | \mathbf{R}g \rangle \end{aligned}$$

Theorem M.17. ³⁶ Let $\mathbf{S} : H \rightarrow H$ be an operator over a Hilbert space H with eigenvalues $\{\lambda_n\}$ and eigenfunctions $\{\psi_n\}$ such that $\mathbf{S}\psi_n = \lambda_n\psi_n$ and let $\|x\| \triangleq \sqrt{\langle x | x \rangle}$.

T H M $\mathbf{S} = \mathbf{S}^*$ \Rightarrow $\left\{ \begin{array}{ll} 1. \langle \mathbf{S}x | x \rangle \in \mathbb{R} & (\text{the hermitian quadratic form of } \mathbf{S} \text{ is real}) \\ 2. \lambda_n \in \mathbb{R} & (\text{eigenvalues of } \mathbf{S} \text{ are real}) \\ 3. \lambda_n \neq \lambda_m \implies \langle \psi_n | \psi_m \rangle = 0 & (\text{eigenfunctions associated with distinct eigenvalues are orthogonal}) \end{array} \right.$

PROOF:

³⁵Historical works regarding self-adjoint operators: von Neumann (1929) page 49, “linearer Operator R selbstadjungiert oder Hermitesch”, Stone (1932) page 50 (“self-adjoint transformations”)

³⁶Lax (2002), pages 315–316, Keener (1988), pages 114–119

1. Proof that $\mathbf{S} = \mathbf{S}^* \implies \langle \mathbf{S}\mathbf{x} | \mathbf{x} \rangle \in \mathbb{R}$:

$$\begin{aligned} \langle \mathbf{x} | \mathbf{S}\mathbf{x} \rangle &= \langle \mathbf{S}\mathbf{x} | \mathbf{x} \rangle \\ &= \langle \mathbf{x} | \mathbf{S}\mathbf{x} \rangle^* \end{aligned} \quad \begin{array}{l} \text{by left hypothesis} \\ \text{by definition of } \langle \triangle | \nabla \rangle \text{ Definition I.1 page 233} \end{array}$$

2. Proof that $\mathbf{S} = \mathbf{S}^* \implies \lambda_n \in \mathbb{R}$:

$$\begin{aligned} \lambda_n \|\psi_n\|^2 &= \lambda_n \langle \psi_n | \psi_n \rangle \\ &= \langle \lambda_n \psi_n | \psi_n \rangle \\ &= \langle \mathbf{S}\psi_n | \psi_n \rangle \\ &= \langle \psi_n | \mathbf{S}\psi_n \rangle \\ &= \langle \psi_n | \lambda_n \psi_n \rangle \\ &= \lambda_n^* \langle \psi_n | \psi_n \rangle \\ &= \lambda_n^* \|\psi_n\|^2 \end{aligned} \quad \begin{array}{l} \text{by definition} \\ \text{by definition of } \langle \triangle | \nabla \rangle \text{ Definition I.1 page 233} \\ \text{by definition of eigenpairs} \\ \text{by left hypothesis} \\ \text{by definition of eigenpairs} \\ \text{by definition of } \langle \triangle | \nabla \rangle \text{ Definition I.1 page 233} \\ \text{by definition} \end{array}$$

3. Proof that $\mathbf{S} = \mathbf{S}^* \implies [\lambda_n \neq \lambda_m \implies \langle \psi_n | \psi_m \rangle = 0]$:

$$\begin{aligned} \lambda_n \langle \psi_n | \psi_m \rangle &= \langle \lambda_n \psi_n | \psi_m \rangle \\ &= \langle \mathbf{S}\psi_n | \psi_m \rangle \\ &= \langle \psi_n | \mathbf{S}\psi_m \rangle \\ &= \langle \psi_n | \lambda_m \psi_m \rangle \\ &= \lambda_m^* \langle \psi_n | \psi_m \rangle \\ &= \lambda_m \langle \psi_n | \psi_m \rangle \end{aligned} \quad \begin{array}{l} \text{by definition of } \langle \triangle | \nabla \rangle \text{ Definition I.1 page 233} \\ \text{by definition of eigenpairs} \\ \text{by left hypothesis} \\ \text{by definition of eigenpairs} \\ \text{by definition of } \langle \triangle | \nabla \rangle \text{ Definition I.1 page 233} \\ \text{because } \lambda_m \text{ is real} \end{array}$$

This implies for $\lambda_n \neq \lambda_m \neq 0$, $\langle \psi_n | \psi_m \rangle = 0$.

M.4.3 Normal Operators

Definition M.9. ³⁷ Let $\mathcal{B}(X, Y)$ be the space of bounded linear operators on normed linear spaces X and Y . Let N^* be the adjoint of an operator $N \in \mathcal{B}(X, Y)$.

D E F N is **normal** if $N^*N = NN^*$.

Theorem M.18. ³⁸ Let $\mathcal{B}(H, H)$ be the space of bounded linear operators on a Hilbert space H . Let $\mathcal{N}(N)$ be the NULL SPACE of an operator N in $\mathcal{B}(H, H)$ and $\mathcal{I}(N)$ the IMAGE SET of N in $\mathcal{B}(H, H)$.

T H M $\underbrace{N^*N = NN^*}_{N \text{ is normal}} \iff \|N^*x\| = \|Nx\| \quad \forall x \in H$

PROOF:

³⁷ Rudin (1991) page 312, Michel and Herget (1993) page 431, Dieudonné (1969), page 167, Frobenius (1878), Frobenius (1968) page 391

³⁸ Rudin (1991) pages 312–313

1. Proof that $\mathbf{N}^*\mathbf{N} = \mathbf{N}\mathbf{N}^* \implies \|\mathbf{N}^*\mathbf{x}\| = \|\mathbf{N}\mathbf{x}\|$:

$$\begin{aligned} \|\mathbf{N}\mathbf{x}\|^2 &= \langle \mathbf{N}\mathbf{x} | \mathbf{N}\mathbf{x} \rangle \\ &= \langle \mathbf{x} | \mathbf{N}^*\mathbf{N}\mathbf{x} \rangle \\ &= \langle \mathbf{x} | \mathbf{N}\mathbf{N}^*\mathbf{x} \rangle \\ &= \langle \mathbf{N}\mathbf{x} | \mathbf{N}^*\mathbf{x} \rangle \\ &= \|\mathbf{N}^*\mathbf{x}\|^2 \end{aligned}$$

by definition
by Proposition M.3 page 294 (definition of \mathbf{N}^*)
by left hypothesis (\mathbf{N} is normal)
by Proposition M.3 page 294 (definition of \mathbf{N}^*)
by definition

2. Proof that $\mathbf{N}^*\mathbf{N} = \mathbf{N}\mathbf{N}^* \iff \|\mathbf{N}^*\mathbf{x}\| = \|\mathbf{N}\mathbf{x}\|$:

$$\begin{aligned} \langle \mathbf{N}^*\mathbf{N}\mathbf{x} | \mathbf{x} \rangle &= \langle \mathbf{N}\mathbf{x} | \mathbf{N}^*\mathbf{x} \rangle && \text{by Proposition M.3 page 294 (definition of } \mathbf{N}^*) \\ &= \langle \mathbf{N}\mathbf{x} | \mathbf{N}\mathbf{x} \rangle && \text{by Theorem M.13 page 295 (property of adjoint)} \\ &= \|\mathbf{N}\mathbf{x}\|^2 && \text{by definition} \\ &= \|\mathbf{N}^*\mathbf{x}\|^2 && \text{by right hypothesis } (\|\mathbf{N}^*\mathbf{x}\| = \|\mathbf{N}\mathbf{x}\|) \\ &= \langle \mathbf{N}^*\mathbf{x} | \mathbf{N}^*\mathbf{x} \rangle && \text{by definition} \\ &= \langle \mathbf{N}\mathbf{N}^*\mathbf{x} | \mathbf{x} \rangle && \text{by Proposition M.3 page 294 (definition of } \mathbf{N}^*) \end{aligned}$$



Theorem M.19. ³⁹ Let $\mathcal{B}(\mathbf{H}, \mathbf{H})$ be the space of bounded linear operators on a Hilbert space \mathbf{H} . Let $\mathcal{N}(\mathbf{N})$ be the NULL SPACE of an operator \mathbf{N} in $\mathcal{B}(\mathbf{H}, \mathbf{H})$ and $\mathcal{I}(\mathbf{N})$ the IMAGE SET of \mathbf{N} in $\mathcal{B}(\mathbf{H}, \mathbf{H})$.

T H M	$\underbrace{\mathbf{N}^*\mathbf{N} = \mathbf{N}\mathbf{N}^*}_{\mathbf{N} \text{ is normal}}$	\implies	$\underbrace{\mathcal{N}(\mathbf{N}^*) = \mathcal{N}(\mathbf{N})}_{\mathbf{N} \text{ and } \mathbf{N}^* \text{ have the same null space}}$
----------------------------------	---	------------	--



PROOF:

$$\begin{aligned} \mathcal{N}(\mathbf{N}^*) &= \{ \mathbf{x} | \mathbf{N}^*\mathbf{x} = 0 \quad \forall \mathbf{x} \in \mathbf{X} \} && \text{(definition of } \mathcal{N}^*) \\ &= \{ \mathbf{x} | \| \mathbf{N}^*\mathbf{x} \| = 0 \quad \forall \mathbf{x} \in \mathbf{X} \} && \text{by definition of } \| \cdot \| \text{ (Definition J.1 page 249)} \\ &= \{ \mathbf{x} | \| \mathbf{N}\mathbf{x} \| = 0 \quad \forall \mathbf{x} \in \mathbf{X} \} \\ &= \{ \mathbf{x} | \mathbf{N}\mathbf{x} = 0 \quad \forall \mathbf{x} \in \mathbf{X} \} && \text{by definition of } \| \cdot \| \text{ (Definition J.1 page 249)} \\ &= \mathcal{N}(\mathbf{N}) && \text{(definition of } \mathcal{N}^*) \end{aligned}$$



M.4.4 Isometric operators

An operator on a pair of normed linear spaces is *isometric* (next definition) if it is an *isometry*.

Definition M.10. Let $(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \|\cdot\|)$ and $(Y, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \|\cdot\|)$ be NORMED LINEAR SPACES (Definition J.1 page 249).

D E F	An operator $\mathbf{M} \in \mathcal{L}(X, Y)$ is isometric if $\ \mathbf{M}\mathbf{x}\ = \ \mathbf{x}\ \quad \forall \mathbf{x} \in X$
----------------------------------	---



Theorem M.20. ⁴⁰ Let $(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \|\cdot\|)$ and $(Y, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \|\cdot\|)$ be NORMED LINEAR SPACES. Let \mathbf{M} be a linear operator in $\mathcal{L}(X, Y)$.

T H M	$\underbrace{\ \mathbf{M}\mathbf{x}\ = \ \mathbf{x}\ \quad \forall \mathbf{x} \in X}_{\text{isometric in length}}$	\iff	$\underbrace{\ \mathbf{M}\mathbf{x} - \mathbf{M}\mathbf{y}\ = \ \mathbf{x} - \mathbf{y}\ \quad \forall \mathbf{x}, \mathbf{y} \in X}_{\text{isometric in distance}}$
----------------------------------	--	--------	--



³⁹ Rudin (1991) pages 312–313

⁴⁰ Kubrusly (2001) page 239 (Proposition 4.37), Berberian (1961) page 27 (Theorem IV.7.5)

PROOF:

1. Proof that $\|\mathbf{M}\mathbf{x}\| = \|\mathbf{x}\| \implies \|\mathbf{M}\mathbf{x} - \mathbf{M}\mathbf{y}\| = \|\mathbf{x} - \mathbf{y}\|$:

$$\begin{aligned} \|\mathbf{M}\mathbf{x} - \mathbf{M}\mathbf{y}\| &= \|\mathbf{M}(\mathbf{x} - \mathbf{y})\| && \text{by definition of linear operators (Definition M.3 page 282)} \\ &= \|\mathbf{M}\mathbf{u}\| && \text{let } \mathbf{u} \triangleq \mathbf{x} - \mathbf{y} \\ &= \|\mathbf{x} - \mathbf{y}\| && \text{by left hypothesis} \end{aligned}$$

2. Proof that $\|\mathbf{M}\mathbf{x}\| = \|\mathbf{x}\| \iff \|\mathbf{M}\mathbf{x} - \mathbf{M}\mathbf{y}\| = \|\mathbf{x} - \mathbf{y}\|$:

$$\begin{aligned} \|\mathbf{M}\mathbf{x}\| &= \|\mathbf{M}(\mathbf{x} - \mathbf{0})\| \\ &= \|\mathbf{M}\mathbf{x} - \mathbf{M}\mathbf{0}\| && \text{by definition of linear operators (Definition M.3 page 282)} \\ &= \|\mathbf{x} - \mathbf{0}\| && \text{by right hypothesis} \\ &= \|\mathbf{x}\| \end{aligned}$$



Isometric operators have already been defined (Definition M.10 page 300) in the more general normed linear spaces, while Theorem M.20 (page 300) demonstrated that in a normed linear space \mathbf{X} , $\|\mathbf{M}\mathbf{x}\| = \|\mathbf{x}\| \iff \|\mathbf{M}\mathbf{x} - \mathbf{M}\mathbf{y}\| = \|\mathbf{x} - \mathbf{y}\|$ for all $\mathbf{x}, \mathbf{y} \in \mathbf{X}$. Here in the more specialized inner product spaces, Theorem M.21 (next) demonstrates two additional equivalent properties.

Theorem M.21. ⁴¹ Let $\mathcal{B}(\mathbf{X}, \mathbf{X})$ be the space of bounded linear operators on a normed linear space $\mathbf{X} \triangleq (X, +, \cdot, (\mathbb{F}, +, \dot{\times}), \|\cdot\|)$. Let \mathbf{N} be a bounded linear operator in $\mathcal{L}(\mathbf{X}, \mathbf{X})$, and \mathbf{I} the identity operator in $\mathcal{L}(\mathbf{X}, \mathbf{X})$. Let $\|\mathbf{x}\| \triangleq \sqrt{\langle \mathbf{x} | \mathbf{x} \rangle}$.

The following conditions are all equivalent:

- | | |
|-------------|---|
| T
H
M | 1. $\mathbf{M}^* \mathbf{M} = \mathbf{I}$ \iff
2. $\langle \mathbf{M}\mathbf{x} \mathbf{M}\mathbf{y} \rangle = \langle \mathbf{x} \mathbf{y} \rangle \quad \forall \mathbf{x}, \mathbf{y} \in X \quad (\mathbf{M} \text{ is surjective}) \iff$
3. $\ \mathbf{M}\mathbf{x} - \mathbf{M}\mathbf{y}\ = \ \mathbf{x} - \mathbf{y}\ \quad \forall \mathbf{x}, \mathbf{y} \in X \quad (\text{isometric in distance}) \iff$
4. $\ \mathbf{M}\mathbf{x}\ = \ \mathbf{x}\ \quad \forall \mathbf{x} \in X \quad (\text{isometric in length}) \iff$ |
|-------------|---|

PROOF:

1. Proof that (1) \implies (2):

$$\begin{aligned} \langle \mathbf{M}\mathbf{x} | \mathbf{M}\mathbf{y} \rangle &= \langle \mathbf{x} | \mathbf{M}^* \mathbf{M}\mathbf{y} \rangle && \text{by Proposition M.3 page 294 (definition of adjoint)} \\ &= \langle \mathbf{x} | \mathbf{I}\mathbf{y} \rangle && \text{by (1)} \\ &= \langle \mathbf{x} | \mathbf{y} \rangle && \text{by Definition M.2 page 281 (definition of I)} \end{aligned}$$

2. Proof that (2) \implies (4):

$$\begin{aligned} \|\mathbf{M}\mathbf{x}\| &= \sqrt{\langle \mathbf{M}\mathbf{x} | \mathbf{M}\mathbf{x} \rangle} && \text{by definition of } \|\cdot\| \\ &= \sqrt{\langle \mathbf{x} | \mathbf{x} \rangle} && \text{by right hypothesis} \\ &= \|\mathbf{x}\| && \text{by definition of } \|\cdot\| \end{aligned}$$

3. Proof that (2) \iff (4):

$$\begin{aligned} 4 \langle \mathbf{M}\mathbf{x} | \mathbf{M}\mathbf{y} \rangle &= \|\mathbf{M}\mathbf{x} + \mathbf{M}\mathbf{y}\|^2 - \|\mathbf{M}\mathbf{x} - \mathbf{M}\mathbf{y}\|^2 + i \|\mathbf{M}\mathbf{x} + i\mathbf{M}\mathbf{y}\|^2 - i \|\mathbf{M}\mathbf{x} - i\mathbf{M}\mathbf{y}\|^2 && \text{by polarization id.} \\ &= \|\mathbf{M}(\mathbf{x} + \mathbf{y})\|^2 - \|\mathbf{M}(\mathbf{x} - \mathbf{y})\|^2 + i \|\mathbf{M}(\mathbf{x} + i\mathbf{y})\|^2 - i \|\mathbf{M}(\mathbf{x} - i\mathbf{y})\|^2 && \text{by Definition M.3} \\ &= \|\mathbf{x} + \mathbf{y}\|^2 - \|\mathbf{x} - \mathbf{y}\|^2 + i \|\mathbf{x} + i\mathbf{y}\|^2 - i \|\mathbf{x} - i\mathbf{y}\|^2 && \text{by left hypothesis} \end{aligned}$$

⁴¹ Michel and Herget (1993) page 432 (Theorem 7.5.8), Kubrusly (2001) page 391 (Proposition 5.72)

4. Proof that (3) \iff (4): by Theorem M.20 page 300

5. Proof that (4) \implies (1):

$$\begin{aligned}
 \langle \mathbf{M}^* \mathbf{M} \mathbf{x} | \mathbf{x} \rangle &= \langle \mathbf{M} \mathbf{x} | \mathbf{M}^{**} \mathbf{x} \rangle && \text{by Proposition M.3 page 294 (definition of adjoint)} \\
 &= \langle \mathbf{M} \mathbf{x} | \mathbf{M} \mathbf{x} \rangle && \text{by Theorem M.13 page 295 (property of adjoint)} \\
 &= \| \mathbf{M} \mathbf{x} \|^2 && \text{by definition} \\
 &= \| \mathbf{x} \|^2 && \text{by left hypothesis with } y = 0 \\
 &= \langle \mathbf{x} | \mathbf{x} \rangle && \text{by definition} \\
 &= \langle \mathbf{I} \mathbf{x} | \mathbf{x} \rangle && \text{by Definition M.2 page 281 (definition of } \mathbf{I} \text{)} \\
 \implies \mathbf{M}^* \mathbf{M} &= \mathbf{I} && \forall \mathbf{x} \in X
 \end{aligned}$$

⇒

Theorem M.22. ⁴² Let $\mathcal{B}(\mathbf{X}, \mathbf{Y})$ be the space of bounded linear operators on normed linear spaces \mathbf{X} and \mathbf{Y} . Let \mathbf{M} be a bounded linear operator in $\mathcal{B}(\mathbf{X}, \mathbf{Y})$, and \mathbf{I} the identity operator in $\mathcal{L}(\mathbf{X}, \mathbf{X})$. Let Λ be the set of eigenvalues of \mathbf{M} . Let $\| \mathbf{x} \| \triangleq \sqrt{\langle \mathbf{x} | \mathbf{x} \rangle}$.

T H M	$\underbrace{\mathbf{M}^* \mathbf{M} = \mathbf{I}}$ <small>\mathbf{M} is isometric</small>	\implies	$\left\{ \begin{array}{l} \ \mathbf{M} \ = 1 \quad (\text{UNIT LENGTH}) \quad \text{and} \\ \lambda = 1 \quad \forall \lambda \in \Lambda \end{array} \right.$
----------------------------------	--	------------	--

PROOF:

1. Proof that $\mathbf{M}^* \mathbf{M} = \mathbf{I} \implies \| \mathbf{M} \| = 1$:

$$\begin{aligned}
 \| \mathbf{M} \| &= \sup_{\mathbf{x} \in \mathbf{X}} \{ \| \mathbf{M} \mathbf{x} \| \mid \| \mathbf{x} \| = 1 \} && \text{by Definition M.4 page 286} \\
 &= \sup_{\mathbf{x} \in \mathbf{X}} \{ \| \mathbf{x} \| \mid \| \mathbf{x} \| = 1 \} && \text{by Theorem M.21 page 301} \\
 &= \sup_{\mathbf{x} \in \mathbf{X}} \{ 1 \} \\
 &= 1
 \end{aligned}$$

2. Proof that $|\lambda| = 1$: Let (\mathbf{x}, λ) be an eigenvector-eigenvalue pair.

$$\begin{aligned}
 1 &= \frac{1}{\| \mathbf{x} \|} \| \mathbf{x} \| \\
 &= \frac{1}{\| \mathbf{x} \|} \| \mathbf{M} \mathbf{x} \| && \text{by Theorem M.21 page 301} \\
 &= \frac{1}{\| \mathbf{x} \|} \| \lambda \mathbf{x} \| && \text{by definition of } \lambda \\
 &= \frac{1}{\| \mathbf{x} \|} |\lambda| \| \mathbf{x} \| && \text{by homogeneous property of } \|\cdot\| \\
 &= |\lambda|
 \end{aligned}$$

⇒

Example M.4 (One sided shift operator). ⁴³ Let \mathbf{X} be the set of all sequences with range $\mathbb{W} (0, 1, 2, \dots)$ and shift operators defined as

1. $\mathbf{S}_r(x_0, x_1, x_2, \dots) \triangleq (0, x_0, x_1, x_2, \dots)$ (right shift operator)
2. $\mathbf{S}_l(x_0, x_1, x_2, \dots) \triangleq (x_1, x_2, x_3, \dots)$ (left shift operator)

⁴² Michel and Herget (1993) page 432

⁴³ Michel and Herget (1993) page 441

- E X**
1. \mathbf{S}_r is an isometric operator.
 2. $\mathbf{S}_r^* = \mathbf{S}_l$

PROOF:

1. Proof that $\mathbf{S}_r^* = \mathbf{S}_l$:

$$\begin{aligned}
 \langle \mathbf{S}_r(x_0, x_1, x_2, \dots) | (y_0, y_1, y_2, \dots) \rangle &= \langle (0, x_0, x_1, x_2, \dots) | (y_0, y_1, y_2, \dots) \rangle \\
 &= \sum_{n=1}^{\infty} x_{n-1} y_n^* \\
 &= \sum_{n=0}^{\infty} x_n y_{n+1}^* \\
 &= \sum_{n=0}^{\infty} x_n y_{n+1}^* \\
 &= \langle (x_0, x_1, x_2, \dots) | (y_1, y_2, y_3, \dots) \rangle \\
 &= \left\langle (x_0, x_1, x_2, \dots) | \underbrace{\mathbf{S}_l(y_0, y_1, y_2, \dots)}_{\mathbf{S}_r^*} \right\rangle
 \end{aligned}$$

2. Proof that \mathbf{S}_r is isometric ($\mathbf{S}_r^* \mathbf{S}_r = \mathbf{I}$):

$$\begin{aligned}
 \mathbf{S}_r^* \mathbf{S}_r &= \mathbf{S}_l \mathbf{S}_r \\
 &= \mathbf{I}
 \end{aligned}
 \quad \text{by 1.}$$

M.4.5 Unitary operators

Definition M.11. ⁴⁴ Let $\mathcal{B}(\mathbf{X}, \mathbf{Y})$ be the space of bounded linear operators on normed linear spaces \mathbf{X} and \mathbf{Y} . Let \mathbf{U} be a bounded linear operator in $\mathcal{B}(\mathbf{X}, \mathbf{Y})$, and \mathbf{I} the identity operator in $\mathcal{B}(\mathbf{X}, \mathbf{X})$.

D E F The operator \mathbf{U} is **unitary** if $\mathbf{U}^* \mathbf{U} = \mathbf{U} \mathbf{U}^* = \mathbf{I}$.

Proposition M.4. Let $\mathcal{B}(\mathbf{X}, \mathbf{Y})$ be the space of bounded linear operators on normed linear spaces \mathbf{X} and \mathbf{Y} . Let \mathbf{U} and \mathbf{V} be bounded linear operators in $\mathcal{B}(\mathbf{X}, \mathbf{Y})$.

P R P $\left. \begin{array}{l} \mathbf{U} \text{ is UNITARY} \\ \mathbf{V} \text{ is UNITARY} \end{array} \right\} \implies (\mathbf{UV}) \text{ is UNITARY.}$

⁴⁴ [Rudin \(1991\)](#) page 312, [Michel and Herget \(1993\)](#) page 431, [Autonne \(1901\)](#) page 209, [Autonne \(1902\)](#), [Schur \(1909\)](#), [Steen \(1973\)](#)

PROOF:

$$\begin{aligned}
 (\mathbf{U}\mathbf{V})(\mathbf{U}\mathbf{V})^* &= (\mathbf{U}\mathbf{V})(\mathbf{V}^*\mathbf{U}^*) && \text{by Theorem M.8 page 291} \\
 &= \mathbf{U}(\mathbf{V}\mathbf{V}^*)\mathbf{U}^* && \text{by associative property} \\
 &= \mathbf{U}\mathbf{I}\mathbf{U}^* && \text{by definition of unitary operators—Definition M.11 page 303} \\
 &= \mathbf{I} && \text{by definition of unitary operators—Definition M.11 page 303}
 \end{aligned}$$

$$\begin{aligned}
 (\mathbf{U}\mathbf{V})^*(\mathbf{U}\mathbf{V}) &= (\mathbf{V}^*\mathbf{U}^*)(\mathbf{U}\mathbf{V}) && \text{by Theorem M.8 page 291} \\
 &= \mathbf{V}^*(\mathbf{U}^*\mathbf{U})\mathbf{V} && \text{by associative property} \\
 &= \mathbf{V}^*\mathbf{I}\mathbf{V} && \text{by definition of unitary operators—Definition M.11 page 303} \\
 &= \mathbf{I} && \text{by definition of unitary operators—Definition M.11 page 303}
 \end{aligned}$$



Theorem M.23. ⁴⁵ Let $\mathcal{B}(\mathbf{H}, \mathbf{H})$ be the space of bounded linear operators on a Hilbert space \mathbf{H} . Let \mathbf{U} be a bounded linear operator in $\mathcal{B}(\mathbf{H}, \mathbf{H})$, and $\mathcal{I}(\mathbf{U})$ the IMAGE SET of \mathbf{U} .

T
H
M

The following conditions are **equivalent**:

1. $\mathbf{U}\mathbf{U}^* = \mathbf{U}^*\mathbf{U} = \mathbf{I}$ (unitary) \iff
2. $\langle \mathbf{U}\mathbf{x} | \mathbf{U}\mathbf{y} \rangle = \langle \mathbf{U}^*\mathbf{x} | \mathbf{U}^*\mathbf{y} \rangle = \langle \mathbf{x} | \mathbf{y} \rangle$ and $\mathcal{I}(\mathbf{U}) = X$ (surjective) \iff
3. $\|\mathbf{U}\mathbf{x} - \mathbf{U}\mathbf{y}\| = \|\mathbf{U}^*\mathbf{x} - \mathbf{U}^*\mathbf{y}\| = \|\mathbf{x} - \mathbf{y}\|$ and $\mathcal{I}(\mathbf{U}) = X$ (isometric in distance) \iff
4. $\|\mathbf{U}\mathbf{x}\| = \|\mathbf{x}\|$ and $\mathcal{I}(\mathbf{U}) = X$ (isometric in length) \iff

PROOF:

1. Proof that (1) \implies (2):

(a) $\langle \mathbf{U}\mathbf{x} | \mathbf{U}\mathbf{y} \rangle = \langle \mathbf{U}^*\mathbf{x} | \mathbf{U}^*\mathbf{y} \rangle = \langle \mathbf{x} | \mathbf{y} \rangle$ by Theorem M.21 (page 301).

(b) Proof that $\mathcal{I}(\mathbf{U}) = X$:

$$\begin{aligned}
 X &\supseteq \mathcal{I}(\mathbf{U}) && \text{because } \mathbf{U} \in X^X \\
 &\supseteq \mathcal{I}(\mathbf{U}\mathbf{U}^*) \\
 &= \mathcal{I}(\mathbf{I}) && \text{by left hypothesis } (\mathbf{U}^*\mathbf{U} = \mathbf{U}\mathbf{U}^* = \mathbf{I}) \\
 &= X && \text{by Definition M.2 page 281 (definition of } \mathbf{I})
 \end{aligned}$$

2. Proof that (2) \iff (3) \iff (4): by Theorem M.21 page 301.

3. Proof that (3) \implies (1):

(a) Proof that $\|\mathbf{U}\mathbf{x} - \mathbf{U}\mathbf{y}\| = \|\mathbf{x} - \mathbf{y}\| \implies \mathbf{U}^*\mathbf{U} = \mathbf{I}$: by Theorem M.21 page 301

(b) Proof that $\|\mathbf{U}^*\mathbf{x} - \mathbf{U}^*\mathbf{y}\| = \|\mathbf{x} - \mathbf{y}\| \implies \mathbf{U}\mathbf{U}^* = \mathbf{I}$:

$$\begin{aligned}
 \|\mathbf{U}^*\mathbf{x} - \mathbf{U}^*\mathbf{y}\| = \|\mathbf{x} - \mathbf{y}\| &\implies \mathbf{U}^{**}\mathbf{U}^* = \mathbf{I} && \text{by Theorem M.21 page 301} \\
 &\implies \mathbf{U}\mathbf{U}^* = \mathbf{I} && \text{by Theorem M.13 page 295}
 \end{aligned}$$



Theorem M.24. Let $\mathcal{B}(\mathbf{H}, \mathbf{H})$ be the space of bounded linear operators on a Hilbert space \mathbf{H} . Let \mathbf{U} be a bounded linear operator in $\mathcal{B}(\mathbf{H}, \mathbf{H})$, $\mathcal{N}(\mathbf{U})$ the NULL SPACE of \mathbf{U} , and $\mathcal{I}(\mathbf{U})$ the IMAGE SET of \mathbf{U} .

T
H
M

$$\underbrace{\mathbf{U}\mathbf{U}^* = \mathbf{U}^*\mathbf{U} = \mathbf{I}}_{\mathbf{U} \text{ is unitary}} \implies \left\{ \begin{array}{lcl} \mathbf{U}^{-1} & = & \mathbf{U}^* \\ \mathcal{I}(\mathbf{U}) & = & \mathcal{I}(\mathbf{U}^*) & = & X \\ \mathcal{N}(\mathbf{U}) & = & \mathcal{N}(\mathbf{U}^*) & = & \{\mathbf{0}\} \\ \|\mathbf{U}\| & = & \|\mathbf{U}^*\| & = & 1 & \text{(UNIT LENGTH)} \end{array} \right.$$

⁴⁵ Rudin (1991) pages 313–314 (Theorem 12.13), Knapp (2005a) page 45 (Proposition 2.6)

PROOF:

1. Note that \mathbf{U} , \mathbf{U}^* , and \mathbf{U}^{-1} are all both **isometric** and **normal**:

$$\begin{aligned}\mathbf{U}^*\mathbf{U} &= \mathbf{I} \implies \mathbf{U} \text{ is isometric} \\ \mathbf{U}\mathbf{U}^* &= \mathbf{U}^*\mathbf{U} = \mathbf{I} \implies \mathbf{U}^* \text{ is isometric} \\ \mathbf{U}^{-1} &= \mathbf{U}^* \implies \mathbf{U}^{-1} \text{ is isometric}\end{aligned}$$

$$\begin{aligned}\mathbf{U}^*\mathbf{U} &= \mathbf{U}\mathbf{U}^* = \mathbf{I} \implies \mathbf{U} \text{ is normal} \\ \mathbf{U}\mathbf{U}^* &= \mathbf{U}^*\mathbf{U} = \mathbf{I} \implies \mathbf{U}^* \text{ is normal} \\ \mathbf{U}^{-1} &= \mathbf{U}^* \implies \mathbf{U}^{-1} \text{ is normal}\end{aligned}$$

2. Proof that $\mathbf{U}^*\mathbf{U} = \mathbf{U}\mathbf{U}^* = \mathbf{I} \implies \mathcal{I}(\mathbf{U}) = \mathcal{I}(\mathbf{U}^*) = \mathcal{H}$: by Theorem M.23 page 304.

3. Proof that $\mathbf{U}^*\mathbf{U} = \mathbf{U}\mathbf{U}^* = \mathbf{I} \implies \mathcal{N}(\mathbf{U}) = \mathcal{N}(\mathbf{U}^*) = \mathcal{N}(\mathbf{U}^{-1})$:

$$\begin{aligned}\mathcal{N}(\mathbf{U}^*) &= \mathcal{N}(\mathbf{U}) && \text{because } \mathbf{U} \text{ and } \mathbf{U}^* \text{ are both normal and by Theorem M.18 page 299} \\ &= \mathcal{I}(\mathbf{U})^\perp && \text{by Theorem M.14 page 296} \\ &= X^\perp && \text{by above result} \\ &= \{\emptyset\} && \text{by Proposition ?? page ??}\end{aligned}$$

4. Proof that $\mathbf{U}^*\mathbf{U} = \mathbf{U}\mathbf{U}^* = \mathbf{I} \implies \|\mathbf{U}^{-1}\| = \|\mathbf{U}^*\| = \|\mathbf{U}\| = 1$:

Because \mathbf{U} , \mathbf{U}^* , and \mathbf{U}^{-1} are all isometric and by Theorem M.22 page 302.



Example M.5. Examples of *Fredholm integral operators* include

1. Fourier Transform $[\tilde{\mathbf{F}}\mathbf{x}](f) = \int_t x(t)e^{-i2\pi ft} dt \quad \kappa(t, f) = e^{-i2\pi ft}$
2. Inverse Fourier Transform $[\tilde{\mathbf{F}}^{-1}\tilde{\mathbf{x}}](t) = \int_f \tilde{x}(f)e^{i2\pi ft} df \quad \kappa(f, t) = e^{i2\pi ft}$
3. Laplace operator $[\mathbf{L}\mathbf{x}](s) = \int_t x(t)e^{-st} dt \quad \kappa(t, s) = e^{-st}$

Example M.6 (Translation operator). Let $\mathbf{X} = L^2_{\mathbb{R}}$ and $\mathbf{T} \in \mathbf{X}^{\mathbf{X}}$ be defined as

$$\mathbf{T}\mathbf{f}(x) \triangleq \mathbf{f}(x - 1) \quad \forall \mathbf{f} \in L^2_{\mathbb{R}} \quad (\text{translation operator})$$

E X	<ol style="list-style-type: none"> 1. $\mathbf{T}^{-1}\mathbf{f}(x) = \mathbf{f}(x + 1) \quad \forall \mathbf{f} \in L^2_{\mathbb{R}}$ (inverse translation operator) 2. $\mathbf{T}^* = \mathbf{T}^{-1}$ (T is invertible) 3. $\mathbf{T}^*\mathbf{T} = \mathbf{T}\mathbf{T}^* = \mathbf{I}$ (T is unitary)
--------	--

PROOF:

1. Proof that $\mathbf{T}^{-1}\mathbf{f}(x) = \mathbf{f}(x + 1)$:

$$\begin{aligned}\mathbf{T}^{-1}\mathbf{T} &= \mathbf{I} \\ \mathbf{T}\mathbf{T}^{-1} &= \mathbf{I}\end{aligned}$$

2. Proof that \mathbf{T} is unitary:

$$\begin{aligned}\langle \mathbf{T}\mathbf{f}(x) | \mathbf{g}(x) \rangle &= \langle \mathbf{f}(x - 1) | \mathbf{g}(x) \rangle && \text{by definition of T} \\ &= \int_x \mathbf{f}(x - 1)\mathbf{g}^*(x) dx \\ &= \int_x \mathbf{f}(x)\mathbf{g}^*(x + 1) dx \\ &= \langle \mathbf{f}(x) | \mathbf{g}(x + 1) \rangle \\ &= \left\langle \mathbf{f}(x) | \underbrace{\mathbf{T}^{-1}\mathbf{g}(x)}_{\mathbf{T}^*} \right\rangle && \text{by 1.}\end{aligned}$$

Example M.7 (Dilation operator). Let $\mathbf{X} = L^2_{\mathbb{R}}$ and $\mathbf{T} \in \mathbf{X}^{\mathbf{X}}$ be defined as

$$\mathbf{D}\mathbf{f}(x) \triangleq \sqrt{2}\mathbf{f}(2x) \quad \forall \mathbf{f} \in L^2_{\mathbb{R}} \quad (\text{dilation operator})$$

- | | |
|--------|--|
| E
X | 1. $\mathbf{D}^{-1}\mathbf{f}(x) = \frac{1}{\sqrt{2}}\mathbf{f}\left(\frac{1}{2}x\right) \quad \forall \mathbf{f} \in L^2_{\mathbb{R}}$ (inverse dilation operator)
2. $\mathbf{D}^* = \mathbf{D}^{-1}$ (\mathbf{D} is invertible)
3. $\mathbf{D}^*\mathbf{D} = \mathbf{D}\mathbf{D}^* = \mathbf{I}$ (\mathbf{D} is unitary) |
|--------|--|

PROOF:

1. Proof that $\mathbf{D}^{-1}\mathbf{f}(x) = \frac{1}{\sqrt{2}}\mathbf{f}\left(\frac{1}{2}x\right)$:

$$\begin{aligned} \mathbf{D}^{-1}\mathbf{D} &= \mathbf{I} \\ \mathbf{D}\mathbf{D}^{-1} &= \mathbf{I} \end{aligned}$$

2. Proof that \mathbf{D} is unitary:

$$\begin{aligned} \langle \mathbf{D}\mathbf{f}(x) | \mathbf{g}(x) \rangle &= \left\langle \sqrt{2}\mathbf{f}(2x) | \mathbf{g}(x) \right\rangle && \text{by definition of } \mathbf{D} \\ &= \int_x \sqrt{2}\mathbf{f}(2x)\mathbf{g}^*(x) dx \\ &= \int_{u \in \mathbb{R}} \sqrt{2}\mathbf{f}(u)\mathbf{g}^*\left(\frac{1}{2}u\right) \frac{1}{2} du && \text{let } u \triangleq 2x \implies dx = \frac{1}{2} du \\ &= \int_{u \in \mathbb{R}} \mathbf{f}(u) \left[\frac{1}{\sqrt{2}}\mathbf{g}\left(\frac{1}{2}u\right) \right]^* du \\ &= \left\langle \mathbf{f}(x) | \frac{1}{\sqrt{2}}\mathbf{g}\left(\frac{1}{2}x\right) \right\rangle \\ &= \left\langle \mathbf{f}(x) | \underbrace{\mathbf{D}^{-1}\mathbf{g}(x)}_{\mathbf{D}^*} \right\rangle && \text{by 1.} \end{aligned}$$

Example M.8 (Delay operator). Let \mathbf{X} be the set of all sequences and $\mathbf{D} \in \mathbf{X}^{\mathbf{X}}$ be a delay operator.

- | | |
|--------|---|
| E
X | The delay operator $\mathbf{D}((x_n))_{n \in \mathbb{Z}} \triangleq ((x_{n-1}))_{n \in \mathbb{Z}}$ is unitary. |
|--------|---|

PROOF: The inverse \mathbf{D}^{-1} of the delay operator \mathbf{D} is

$$\mathbf{D}^{-1}((x_n))_{n \in \mathbb{Z}} \triangleq ((x_{n+1}))_{n \in \mathbb{Z}}.$$

$$\begin{aligned} \langle \mathbf{D}((x_n)) | (y_n) \rangle &= \langle ((x_{n-1})) | (y_n) \rangle && \text{by definition of } \mathbf{D} \\ &= \sum_n x_{n-1} y_n^* \\ &= \sum_n x_n y_{n+1}^* \\ &= \langle ((x_n)) | ((y_{n+1})) \rangle \\ &= \left\langle ((x_n)) | \underbrace{\mathbf{D}^{-1}((y_n))}_{\mathbf{D}^*} \right\rangle \end{aligned}$$

Therefore, $\mathbf{D}^* = \mathbf{D}^{-1}$. This implies that $\mathbf{D}\mathbf{D}^* = \mathbf{D}^*\mathbf{D} = \mathbf{I}$ which implies that \mathbf{D} is unitary.

Example M.9 (Fourier transform). Let $\tilde{\mathbf{F}}$ be the *Fourier Transform* and $\tilde{\mathbf{F}}^{-1}$ the *inverse Fourier Transform* operator

$$[\tilde{\mathbf{F}}\mathbf{x}](f) \triangleq \int_t \mathbf{x}(t) e^{\underbrace{-i2\pi f t}_{\kappa(t, f)}} dt \quad [\tilde{\mathbf{F}}^{-1}\tilde{\mathbf{x}}](t) \triangleq \int_f \tilde{\mathbf{x}}(f) e^{\underbrace{i2\pi f t}_{\kappa^*(t, f)}} df.$$

E X $\tilde{\mathbf{F}}^* = \tilde{\mathbf{F}}^{-1}$ (the Fourier Transform operator $\tilde{\mathbf{F}}$ is unitary)

PROOF:

$$\begin{aligned} \langle \tilde{\mathbf{F}}\mathbf{x} | \tilde{\mathbf{y}} \rangle &= \left\langle \int_t \mathbf{x}(t) e^{-i2\pi f t} dt | \tilde{\mathbf{y}}(f) \right\rangle \\ &= \int_t \mathbf{x}(t) \left\langle e^{-i2\pi f t} | \tilde{\mathbf{y}}(f) \right\rangle dt \\ &= \int_t \mathbf{x}(t) \int_f e^{-i2\pi f t} \tilde{\mathbf{y}}^*(f) df dt \\ &= \int_t \mathbf{x}(t) \left[\int_f e^{i2\pi f t} \tilde{\mathbf{y}}(f) df \right]^* dt \\ &= \left\langle \mathbf{x}(t) | \int_f \tilde{\mathbf{y}}(f) e^{i2\pi f t} df \right\rangle \\ &= \left\langle \mathbf{x} | \underbrace{\tilde{\mathbf{F}}^{-1}\tilde{\mathbf{y}}}_{\tilde{\mathbf{F}}^*} \right\rangle \end{aligned}$$

This implies that $\tilde{\mathbf{F}}$ is unitary ($\tilde{\mathbf{F}}^* = \tilde{\mathbf{F}}^{-1}$). ➡

Example M.10 (Rotation matrix). ⁴⁶ Let the rotation matrix $\mathbf{R}_\theta : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be defined as

$$\mathbf{R}_\theta \triangleq \begin{bmatrix} \cos\theta & -\sin\theta \\ \sin\theta & \cos\theta \end{bmatrix}$$

E X

1.	$\mathbf{R}_\theta^{-1} = \mathbf{R}_{-\theta}$
2.	$\mathbf{R}_\theta^* = \mathbf{R}_{-\theta}$ (\mathbf{R} is unitary)

PROOF:

$$\begin{aligned} \mathbf{R}^* &= \mathbf{R}^H \\ &= \begin{bmatrix} \cos\theta & -\sin\theta \\ \sin\theta & \cos\theta \end{bmatrix}^H && \text{by definition of } \mathbf{R} \\ &= \begin{bmatrix} \cos\theta & \sin\theta \\ -\sin\theta & \cos\theta \end{bmatrix} && \text{by definition of Hermetian transpose operator } H \\ &= \begin{bmatrix} \cos(-\theta) & -\sin(-\theta) \\ \sin(-\theta) & \cos(-\theta) \end{bmatrix} && \text{by Theorem ?? page ??} \\ &= \mathbf{R}_{-\theta} && \text{by definition of } \mathbf{R} \\ &= \mathbf{R}^{-1} && \text{by 1.} \end{aligned}$$

⁴⁶  Noble and Daniel (1988), page 311

M.5 Operator order

Definition M.12. ⁴⁷ Let $P \in Y^X$ be an operator.

D E F P is **positive** if $\langle Px | x \rangle \geq 0 \forall x \in X$.
This condition is denoted $P \geq 0$.

Theorem M.25. ⁴⁸

T H M $\underbrace{P \geq 0 \text{ and } Q \geq 0}_{P \text{ and } Q \text{ are both positive}}$	\implies	$\begin{cases} (P + Q) \geq 0 & ((P + Q) \text{ is positive}) \\ A^*PA \geq 0 & \forall A \in \mathcal{B}(X, X) \quad (A^*PA \text{ is positive}) \\ A^*A \geq 0 & \forall A \in \mathcal{B}(X, X) \quad (A^*A \text{ is positive}) \end{cases}$
--	------------	--

PROOF:

$$\begin{aligned}
 \langle (P + Q)x | x \rangle &= \langle Px | x \rangle + \langle Qx | x \rangle && \text{by additive property of } \langle \cdot | \cdot \rangle \text{ (Definition I.1 page 233)} \\
 &\geq \langle Px | x \rangle && \text{by left hypothesis} \\
 &\geq 0 && \text{by left hypothesis} \\
 \langle A^*PAx | x \rangle &= \langle PAx | Ax \rangle && \text{by definition of adjoint (Proposition M.3 page 294)} \\
 &= \langle Py | y \rangle && \text{where } y \triangleq Ax \\
 &\geq 0 && \text{by left hypothesis} \\
 \langle Ix | x \rangle &= \langle x | x \rangle && \text{by definition of } I \text{ (Definition M.2 page 281)} \\
 &\geq 0 && \text{by non-negative property of } \langle \cdot | \cdot \rangle \text{ (Definition I.1 page 233)} \\
 &\implies I \text{ is positive} && \\
 \langle A^*Ax | x \rangle &= \langle A^*Ix | x \rangle && \text{by definition of } I \text{ (Definition M.2 page 281)} \\
 &\geq 0 && \text{by two previous results}
 \end{aligned}$$

Definition M.13. ⁴⁹ Let $A, B \in \mathcal{B}(X, Y)$ be bounded operators.

D E F $\underbrace{A \geq B}_{\text{"A is greater than or equal to B"}}$ \iff $\underbrace{(A - B) \geq 0}_{\text{"(A - B) is positive"}}$

⁴⁷ Michel and Herget (1993) page 429 (Definition 7.4.12)

⁴⁸ Michel and Herget (1993) page 429

⁴⁹ Michel and Herget (1993) page 429

APPENDIX N

FOURIER TRANSFORM



“The analytical equations ... extend to all general phenomena. There cannot be a language more universal and more simple, more free from errors and from obscurities, ... mathematical analysis is as extensive as nature itself; ...it preserves every principle which it has once acquired; it grows and strengthens itself incessantly in the midst of the many variations and errors of the human mind. Its chief attribute is clearness; it has no marks to express confused notions. It brings together phenomena the most diverse, and discovers the hidden analogies which unite them.”

Joseph Fourier (1768–1830)¹

N.1 Definitions

This chapter deals with the *Fourier Transform* in the space of *Lebesgue square-integrable functions* $L^2_{(\mathbb{R}, \mathcal{B}, \mu)}$, where \mathbb{R} is the set of real numbers, \mathcal{B} is the set of *Borel sets* on \mathbb{R} , μ is the standard *Borel measure* on \mathbb{R} , and

$$L^2_{(\mathbb{R}, \mathcal{B}, \mu)} \triangleq \left\{ f \in \mathbb{R}^{\mathbb{R}} \mid \int_{\mathbb{R}} |f|^2 d\mu < \infty \right\}.$$

Furthermore, $\langle \Delta | \nabla \rangle$ is the *inner product* induced by the operator $\int_{\mathbb{R}} d\mu$ such that

$$\langle f | g \rangle \triangleq \int_{\mathbb{R}} f(x)g^*(x) dx,$$

and $(L^2_{(\mathbb{R}, \mathcal{B}, \mu)}, \langle \Delta | \nabla \rangle)$ is a *Hilbert space*.

Definition N.1. Let κ be a FUNCTION in $\mathbb{C}^{\mathbb{R}^2}$.

D E F The function κ is the **Fourier kernel** if $\kappa(x, \omega) \triangleq e^{i\omega x} \quad \forall x, \omega \in \mathbb{R}$

Definition N.2. ² Let $L^2_{(\mathbb{R}, \mathcal{B}, \mu)}$ be the space of all LEBESGUE SQUARE-INTEGRABLE FUNCTIONS.

¹ quote: [Fourier \(1878\)](#), pages 7–8 (Preliminary Discourse)

image: <http://en.wikipedia.org/wiki/File:Fourier2.jpg>, public domain

² [Bachman et al. \(2000\)](#) page 363, [Chorin and Hald \(2009\)](#) page 13, [Loomis and Bolker \(1965\)](#), page 144, [Knapp \(2005b\)](#) pages 374–375, [Fourier \(1822\)](#), [Fourier \(1878\)](#) page 336?

DEF

The Fourier Transform operator $\tilde{\mathbf{F}}$ is defined as

$$[\tilde{\mathbf{F}}f](\omega) \triangleq \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} f(x) e^{-i\omega x} dx \quad \forall f \in L^2_{(\mathbb{R}, \mathcal{B}, \mu)}$$

This definition of the Fourier Transform is also called the **unitary Fourier Transform**.

Remark N.1 (Fourier transform scaling factor).³ If the Fourier transform operator $\tilde{\mathbf{F}}$ and inverse Fourier transform operator $\tilde{\mathbf{F}}^{-1}$ are defined as

$$\tilde{\mathbf{F}}f(x) \triangleq F(\omega) \triangleq A \int_{\mathbb{R}} f(x) e^{-i\omega x} dx \quad \text{and} \quad \tilde{\mathbf{F}}^{-1}\tilde{f}(\omega) \triangleq B \int_{\mathbb{R}} F(\omega) e^{i\omega x} dx$$

then A and B can be any constants as long as $AB = \frac{1}{2\pi}$. The Fourier transform is often defined with the scaling factor A set equal to 1 such that $[\tilde{\mathbf{F}}f(x)](\omega) \triangleq \int_{\mathbb{R}} f(x) e^{-i\omega x} dx$. In this case, the inverse Fourier transform operator $\tilde{\mathbf{F}}^{-1}$ is either defined as

- $[\tilde{\mathbf{F}}^{-1}f(x)](f) \triangleq \int_{\mathbb{R}} f(x) e^{i2\pi f x} dx$ (using oscillatory frequency free variable f) or
- $[\tilde{\mathbf{F}}^{-1}f(x)](\omega) \triangleq \frac{1}{2\pi} \int_{\mathbb{R}} f(x) e^{i\omega x} dx$ (using angular frequency free variable ω).

In short, the 2π has to show up somewhere, either in the argument of the exponential ($e^{-i2\pi f t}$) or in front of the integral ($\frac{1}{2\pi} \int \dots$). One could argue that it is unnecessary to burden the exponential argument with the 2π factor ($e^{-i2\pi f t}$), and thus could further argue in favor of using the angular frequency variable ω thus giving the inverse operator definition $[\tilde{\mathbf{F}}^{-1}f(x)](\omega) \triangleq \frac{1}{2\pi} \int_{\mathbb{R}} f(x) e^{-i\omega x} dx$. But this causes a new problem. In this case, the Fourier operator $\tilde{\mathbf{F}}$ is not *unitary* (see Theorem N.2 page 310)—in particular, $\tilde{\mathbf{F}}\tilde{\mathbf{F}}^* \neq \mathbf{I}$, where $\tilde{\mathbf{F}}^*$ is the *adjoint* of $\tilde{\mathbf{F}}$; but rather, $\tilde{\mathbf{F}} \left(\frac{1}{2\pi} \tilde{\mathbf{F}}^* \right) = \left(\frac{1}{2\pi} \tilde{\mathbf{F}}^* \right) \tilde{\mathbf{F}} = \mathbf{I}$. But if we define the operators $\tilde{\mathbf{F}}$ and $\tilde{\mathbf{F}}^{-1}$ to both have the scaling factor $\frac{1}{\sqrt{2\pi}}$, then $\tilde{\mathbf{F}}$ and $\tilde{\mathbf{F}}^{-1}$ are inverses and $\tilde{\mathbf{F}}$ is *unitary*—that is, $\tilde{\mathbf{F}}\tilde{\mathbf{F}}^* = \tilde{\mathbf{F}}^*\tilde{\mathbf{F}} = \mathbf{I}$.

N.2 Operator properties

Theorem N.1 (Inverse Fourier transform).⁴ Let $\tilde{\mathbf{F}}$ be the Fourier Transform operator (Definition N.2 page 309). The inverse $\tilde{\mathbf{F}}^{-1}$ of $\tilde{\mathbf{F}}$ is

THM

$$[\tilde{\mathbf{F}}^{-1}\tilde{f}](x) = \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} \tilde{f}(\omega) e^{i\omega x} d\omega \quad \forall \tilde{f} \in L^2_{(\mathbb{R}, \mathcal{B}, \mu)}$$

Theorem N.2. Let $\tilde{\mathbf{F}}$ be the Fourier Transform operator with inverse $\tilde{\mathbf{F}}^{-1}$ and adjoint $\tilde{\mathbf{F}}^*$.

THM

$$\tilde{\mathbf{F}}^* = \tilde{\mathbf{F}}^{-1}$$

PROOF:

$$\begin{aligned} \langle \tilde{\mathbf{F}}f | g \rangle &= \left\langle \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} f(x) e^{-i\omega x} dx | g(\omega) \right\rangle && \text{by definition of } \tilde{\mathbf{F}} \text{ page 309} \\ &= \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} f(x) \langle e^{-i\omega x} | g(\omega) \rangle dx && \text{by additive property of } \langle \cdot | \cdot \rangle \text{ page 233} \\ &= \int_{\mathbb{R}} f(x) \frac{1}{\sqrt{2\pi}} \langle g(\omega) | e^{-i\omega x} \rangle^* dx && \text{by conjugate symmetric property of } \langle \cdot | \cdot \rangle \text{ page 233} \end{aligned}$$

³ Chorin and Hald (2009) page 13, Jeffrey and Dai (2008) pages xxxi–xxxii, Knapp (2005b) pages 374–375

⁴ Chorin and Hald (2009) page 13

$$\begin{aligned}
 &= \left\langle f(x) \mid \frac{1}{\sqrt{2\pi}} \langle g(\omega) \mid e^{-i\omega x} \rangle \right\rangle && \text{by definition of } \langle \triangle \mid \nabla \rangle \\
 &= \left\langle f \mid \underbrace{\tilde{F}^{-1}g}_{\tilde{F}^*} \right\rangle && \text{by Theorem N.1 page 310}
 \end{aligned}$$



The Fourier Transform operator has several nice properties:

- \tilde{F} is *unitary*⁵ (Corollary N.1—next corollary).
- Because \tilde{F} is unitary, it automatically has several other nice properties (Theorem N.3 page 311).

Corollary N.1. Let I be the identity operator and let \tilde{F} be the Fourier Transform operator with adjoint \tilde{F}^* and inverse \tilde{F}^{-1} .

C O R	$\tilde{F}\tilde{F}^* = \tilde{F}^*\tilde{F} = I$ (\tilde{F} is unitary)
	$\tilde{F}^* = \tilde{F}^{-1}$



PROOF: This follows directly from the fact that $\tilde{F}^* = \tilde{F}^{-1}$ (Theorem N.2 page 310).

Theorem N.3. Let \tilde{F} be the Fourier transform operator with adjoint \tilde{F}^* and inverse \tilde{F} . Let $\|\cdot\|$ be the operator norm with respect to the vector norm $\|\cdot\|$ with respect to the Hilbert space $(\mathbb{C}^{\mathbb{R}}, \langle \triangle \mid \nabla \rangle)$. Let $\mathcal{R}(A)$ be the range of an operator A .

T H M	$\mathcal{R}(F\tau) = \mathcal{R}(\tilde{F}^{-1}) = L^2_{\mathbb{R}}$ $\ \tilde{F}\ = \ \tilde{F}^{-1}\ = 1$ (UNITARY) $\langle \tilde{F}f \mid \tilde{F}g \rangle = \langle \tilde{F}^{-1}f \mid \tilde{F}^{-1}g \rangle = \langle f \mid g \rangle$ (PARSEVAL'S EQUATION) $\ \tilde{F}f\ = \ \tilde{F}^{-1}f\ = \ f\ $ (PLANCHEREL'S FORMULA) $\ \tilde{F}f - \tilde{F}g\ = \ \tilde{F}^{-1}f - \tilde{F}^{-1}g\ = \ f - g\ $ (ISOMETRIC)
-------------	---



PROOF: These results follow directly from the fact that \tilde{F} is unitary (Corollary N.1 page 311) and from the properties of unitary operators (Theorem M.24 page 304).



Theorem N.4 (Shift relations). Let \tilde{F} be the Fourier transform operator.

T H M	$\tilde{F}[f(x-u)](\omega) = e^{-i\omega u} [\tilde{F}f(x)](\omega)$ $[\tilde{F}(e^{ivx}g(x))](\omega) = [\tilde{F}g(x)](\omega - v)$
-------------	--



PROOF:

$$\begin{aligned}
 \tilde{F}[f(x-u)](\omega) &= \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} f(x-u)e^{-i\omega x} dx && \text{by definition of } \tilde{F} && (\text{Definition N.2 page 309}) \\
 &= \frac{1}{\sqrt{2\pi}} \int_{v \in \mathbb{R}} f(v)e^{-i\omega(u+v)} dv && \text{where } v \triangleq x - u \implies t = u + v \\
 &= e^{-i\omega u} \frac{1}{\sqrt{2\pi}} \int_{v \in \mathbb{R}} f(v)e^{-i\omega v} dv && \\
 &= e^{-i\omega u} \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} f(x)e^{-i\omega x} dx && \text{by change of variable } t = v \\
 &= e^{-i\omega u} [\tilde{F}f(x)](\omega) && \text{by definition of } \tilde{F} && (\text{Definition N.2 page 309}) \\
 [\tilde{F}(e^{ivx}g(x))](\omega) &= \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} e^{ivx}g(x)e^{-i\omega x} dx && \text{by definition of } \tilde{F} && (\text{Definition N.2 page 309})
 \end{aligned}$$

⁵ unitary operators: Definition M.11 page 303

$$= \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} g(x) e^{-i(\omega-v)x} dx$$

$$= [\tilde{\mathbf{F}}g(x)](\omega - v) \quad \text{by definition of } \tilde{\mathbf{F}} \quad (\text{Definition N.2 page 309})$$

⇒

Theorem N.5 (Complex conjugate). *Let $\tilde{\mathbf{F}}$ be the Fourier Transform operator and $*$ represent the complex conjugate operation on the set of complex numbers.*

T H M	$\tilde{\mathbf{F}}f^*(-x) = -[\tilde{\mathbf{F}}f(x)]^* \quad \forall f \in L^2_{(\mathbb{R}, \mathcal{B}, \mu)}$
	$f \text{ is real} \implies \tilde{f}(-\omega) = [\tilde{f}(\omega)]^* \quad \forall \omega \in \mathbb{R}$ REALITY CONDITION

PROOF:

$$[\tilde{\mathbf{F}}f^*(-x)](\omega) \triangleq \frac{1}{\sqrt{2\pi}} \int f^*(-x) e^{-i\omega x} dx \quad \text{by definition of } \tilde{\mathbf{F}} \quad (\text{Definition N.2 page 309})$$

$$= \frac{1}{\sqrt{2\pi}} \int f^*(u) e^{i\omega u} (-1) du \quad \text{where } u \triangleq -x \implies dx = -du$$

$$= - \left[\frac{1}{\sqrt{2\pi}} \int f(u) e^{-i\omega u} du \right]^*$$

$$\triangleq -[\tilde{\mathbf{F}}f(x)]^* \quad \text{by definition of } \tilde{\mathbf{F}} \quad (\text{Definition N.2 page 309})$$

$$\tilde{f}(-\omega) \triangleq \frac{1}{\sqrt{2\pi}} \int f(x) e^{-i(-\omega)x} dx \quad \text{by definition of } \tilde{\mathbf{F}} \quad (\text{Definition N.2 page 309})$$

$$= \left[\frac{1}{\sqrt{2\pi}} \int f^*(x) e^{-i\omega x} dx \right]^*$$

$$= \left[\frac{1}{\sqrt{2\pi}} \int f(x) e^{-i\omega x} dx \right]^* \quad \text{by } f \text{ is real hypothesis}$$

$$\triangleq \tilde{f}^*(\omega) \quad \text{by definition of } \tilde{\mathbf{F}} \quad (\text{Definition N.2 page 309})$$

⇒

N.3 Convolution

Definition N.3. ⁶

D E F The convolution operation is defined as

$$[f \star g](x) \triangleq f(x) \star g(x) \triangleq \int_{u \in \mathbb{R}} f(u) g(x-u) du \quad \forall f, g \in L^2_{(\mathbb{R}, \mathcal{B}, \mu)}$$

Theorem P.2 (next) demonstrates that multiplication in the “time domain” is equivalent to convolution in the “frequency domain” and vice-versa.

Theorem N.6 (convolution theorem). ⁷ Let $\tilde{\mathbf{F}}$ be the Fourier Transform operator and \star the convolution operator.

⁶ Bachman (1964), page 6, Bracewell (1978) page 108 (Convolution theorem)

⁷ Bracewell (1978) page 110

T H M

$\underbrace{\tilde{F}[f(x) \star g(x)](\omega)}_{\text{convolution in "time domain"}},$	$= \underbrace{\sqrt{2\pi} [\tilde{F}f](\omega) [\tilde{F}g](\omega)}_{\text{multiplication in "frequency domain"}}, \quad \forall f, g \in L^2_{(\mathbb{R}, \mathcal{B}, \mu)}$
$\underbrace{\tilde{F}[f(x)g(x)](\omega)}_{\text{multiplication in "time domain"}},$	$= \underbrace{\frac{1}{\sqrt{2\pi}} [\tilde{F}f](\omega) \star [\tilde{F}g](\omega)}_{\text{convolution in "frequency domain"}}, \quad \forall f, g \in L^2_{(\mathbb{R}, \mathcal{B}, \mu)}.$

PROOF:

$$\begin{aligned}
 \tilde{F}[f(x) \star g(x)](\omega) &= \tilde{F}\left[\int_{u \in \mathbb{R}} f(u)g(x-u) du\right](\omega) && \text{by definition of } \star \text{ (Definition N.3 page 312)} \\
 &= \int_{u \in \mathbb{R}} f(u)[\tilde{F}g(x-u)](\omega) du \\
 &= \int_{u \in \mathbb{R}} f(u)e^{-i\omega u} [\tilde{F}g(x)](\omega) du && \text{by Theorem N.4 page 311} \\
 &= \sqrt{2\pi} \left(\underbrace{\frac{1}{\sqrt{2\pi}} \int_{u \in \mathbb{R}} f(u)e^{-i\omega u} du}_{[\tilde{F}f](\omega)} \right) [\tilde{F}g](\omega) \\
 &= \sqrt{2\pi} [\tilde{F}f](\omega) [\tilde{F}g](\omega) && \text{by definition of } \tilde{F} \text{ (Definition N.2 page 309)} \\
 \tilde{F}[f(x)g(x)](\omega) &= \tilde{F}[(\tilde{F}^{-1}\tilde{F}f(x))g(x)](\omega) && \text{by definition of operator inverse (page 281)} \\
 &= \tilde{F}\left[\left(\frac{1}{\sqrt{2\pi}} \int_{v \in \mathbb{R}} [\tilde{F}f(x)](v)e^{ivx} dv\right) g(x)\right](\omega) && \text{by Theorem N.1 page 310} \\
 &= \frac{1}{\sqrt{2\pi}} \int_{v \in \mathbb{R}} [\tilde{F}f(x)](v) [\tilde{F}(e^{ivx} g(x))](\omega, v) dv \\
 &= \frac{1}{\sqrt{2\pi}} \int_{v \in \mathbb{R}} [\tilde{F}f(x)](v) [\tilde{F}g(x)](\omega - v) dv && \text{by Theorem N.4 page 311} \\
 &= \frac{1}{\sqrt{2\pi}} [\tilde{F}f](\omega) \star [\tilde{F}g](\omega) && \text{by definition of } \star \text{ (Definition N.3 page 312)}
 \end{aligned}$$

N.4 Real valued functions




Figure N.1: Fourier transform components of real-valued signal

Theorem N.7. Let $f(x)$ be a function in $L^2_{\mathbb{R}}$ and $\tilde{f}(\omega)$ the FOURIER TRANSFORM of $f(x)$.

T H M

$\left\{ \begin{array}{l} f(x) \text{ is REAL-VALUED} \\ (f \in \mathbb{R}^{\mathbb{R}}) \end{array} \right\}$	$\Rightarrow \left\{ \begin{array}{ll} \tilde{f}(\omega) &= \tilde{f}^*(-\omega) & (\text{HERMITIAN SYMMETRIC}) \\ \mathbf{R}_e[\tilde{f}(\omega)] &= \mathbf{R}_e[\tilde{f}(-\omega)] & (\text{SYMMETRIC}) \\ \mathbf{I}_m[\tilde{f}(\omega)] &= -\mathbf{I}_m[\tilde{f}(-\omega)] & (\text{ANTI-SYMMETRIC}) \\ \tilde{f}(\omega) &= \tilde{f}(-\omega) & (\text{SYMMETRIC}) \\ \angle[\tilde{f}(\omega)] &= \angle[\tilde{f}(-\omega)] & (\text{ANTI-SYMMETRIC}). \end{array} \right\}$
--	---

PROOF:

$$\begin{aligned}
 \tilde{f}(\omega) &\triangleq [\tilde{\mathbf{F}}f(x)](\omega) \triangleq \langle f(x) | e^{i\omega x} \rangle = \langle f(x) | e^{i(-\omega)x} \rangle^* \triangleq \tilde{f}^*(-\omega) \\
 \mathbf{R}_e[\tilde{f}(\omega)] &= \mathbf{R}_e[\tilde{f}^*(-\omega)] = \mathbf{R}_e[\tilde{f}(-\omega)] \\
 \mathbf{I}_m[\tilde{f}(\omega)] &= \mathbf{I}_m[\tilde{f}^*(-\omega)] = -\mathbf{I}_m[\tilde{f}(-\omega)] \\
 |\tilde{f}(\omega)| &= |\tilde{f}^*(-\omega)| = |\tilde{f}(-\omega)| \\
 \angle\tilde{f}(\omega) &= \angle\tilde{f}^*(-\omega) = -\angle\tilde{f}(-\omega)
 \end{aligned}$$

⇒

N.5 Moment properties

Definition N.4.⁸

DEF

The quantity M_n is the ***n*th moment** of a function $f(x) \in L^2_{\mathbb{R}}$ if

$$M_n \triangleq \int_{\mathbb{R}} x^n f(x) dx \quad \text{for } n \in \mathbb{W}.$$

Lemma N.1.⁹ Let M_n be the ***n*th moment** (Definition N.4 page 314) and $\tilde{f}(\omega) \triangleq [\tilde{\mathbf{F}}f](\omega)$ the **FOURIER TRANSFORM** (Definition N.2 page 309) of a function $f(x)$ in $L^2_{\mathbb{R}}$ (Definition ?? page ??).

LEM

$$M_n = \sqrt{2\pi}(i)^n \left[\frac{d}{d\omega} \right]^n \tilde{f}(\omega) \Big|_{\omega=0} \quad \forall n \in \mathbb{W}, f \in L^2_{\mathbb{R}}$$

$$\left[\frac{d}{d\omega} \right]^n \tilde{f}(\omega) \Big|_{\omega=0} = \frac{1}{\sqrt{2\pi}} (-i)^n M_n \quad \forall n \in \mathbb{W}, f \in L^2_{\mathbb{R}}$$

PROOF:

$$\begin{aligned}
 \sqrt{2\pi}(i)^n \left[\left[\frac{d}{d\omega} \right]^n \tilde{f}(\omega) \right]_{\omega=0} &= \sqrt{2\pi}(i)^n \left[\left[\frac{d}{d\omega} \right]^n \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} f(x) e^{-i\omega x} dx \right]_{\omega=0} \quad \text{by definition of } \tilde{\mathbf{F}} \text{ (Definition N.2 page 309)} \\
 &= (i)^n \int_{\mathbb{R}} f(x) \left[\left[\frac{d}{d\omega} \right]^n e^{-i\omega x} \right] dx \Big|_{\omega=0} \\
 &= (i)^n \int_{\mathbb{R}} f(x) [(-i)^n x^n e^{-i\omega x}] dx \Big|_{\omega=0} \\
 &= (-i^2)^n \int_{\mathbb{R}} f(x) x^n dx \\
 &= \int_{\mathbb{R}} f(x) x^n dx \\
 &\triangleq M_n \quad \text{by definition of } M_n \text{ (Definition N.4 page 314)}
 \end{aligned}$$

⇒

Lemma N.2.¹⁰ Let M_n be the ***n*th moment** (Definition N.4 page 314) and $\tilde{f}(\omega) \triangleq [\tilde{\mathbf{F}}f](\omega)$ the **FOURIER TRANSFORM** (Definition N.2 page 309) of a function $f(x)$ in $L^2_{\mathbb{R}}$ (Definition ?? page ??).

LEM

$$M_n = 0 \iff \left[\frac{d}{d\omega} \right]^n \tilde{f}(\omega) \Big|_{\omega=0} = 0 \quad \forall n \in \mathbb{W}$$

PROOF:

⁸ Jawerth and Sweldens (1994), pages 16–17, Sweldens and Piessens (1993), page 2, Vidakovic (1999), page 83

⁹ Goswami and Chan (1999), pages 38–39

¹⁰ Vidakovic (1999), pages 82–83, Mallat (1999), pages 241–242

1. Proof for (\implies) case:

$$\begin{aligned} 0 &= \langle f(x) | x^n \rangle && \text{by left hypothesis} \\ &= \sqrt{2\pi}(-i)^{-n} \left[\frac{d}{d\omega} \right]^n \tilde{f}(\omega) \Big|_{\omega=0} && \text{by Lemma N.1 page 314} \\ &\implies \left[\frac{d}{d\omega} \right]^n \tilde{f}(\omega) \Big|_{\omega=0} = 0 \end{aligned}$$

2. Proof for (\Leftarrow) case:

$$\begin{aligned} 0 &= \left[\frac{d}{d\omega} \right]^n \tilde{f}(\omega) \Big|_{\omega=0} && \text{by right hypothesis} \\ &= \left[\frac{d}{d\omega} \right]^n \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} f(x) e^{-i\omega x} dx \Big|_{\omega=0} && \text{by definition of } \tilde{f}(\omega) \\ &= \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} f(x) \left[\frac{d}{d\omega} \right]^n e^{-i\omega x} dx \Big|_{\omega=0} \\ &= \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} f(x) [(-i)^n x^n e^{-i\omega x}] dx \Big|_{\omega=0} \\ &= (-i)^n \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} f(x) x^n dx \\ &= (-i)^n \frac{1}{\sqrt{2\pi}} \langle f(x) | x^n \rangle && \text{by definition of } \langle \cdot | \cdot \rangle \text{ in } L^2_{\mathbb{R}} \text{ (Definition ?? page ??)} \end{aligned}$$



Lemma N.3 (Strang-Fix condition). ¹¹ Let $f(x)$ be a function in $L^2_{\mathbb{R}}$ and M_n the n TH MOMENT (Definition N.4 page 314) off $f(x)$. Let T be the TRANSLATION OPERATOR (Definition ?? page ??).

L E M	$\sum_{k \in \mathbb{Z}} \underbrace{T^k x^n f(x)}_{\text{STRANG-FIX CONDITION in "time"}} = M_n \iff \underbrace{\left[\frac{d}{d\omega} \right]^n \tilde{f}(\omega) \Big _{\omega=2\pi k} = \frac{1}{\sqrt{2\pi}} (-i)^n \bar{\delta}_k M_n}_{\text{STRANG-FIX CONDITION in "frequency"}}$
----------------------	---

PROOF:

1. Proof for (\implies) case:

$$\begin{aligned} \left[\left[\frac{d}{d\omega} \right]^n \tilde{f}(\omega) \right]_{\omega=2\pi k} &= \sum_{k \in \mathbb{Z}} \left[\left[\frac{d}{d\omega} \right]^n \tilde{f}(\omega) \right]_{\omega=2\pi k} e^{i2\pi kx} \bar{\delta}_k \\ &= \frac{1}{\sqrt{2\pi}} \sum_{k \in \mathbb{Z}} \left[\left[\frac{d}{d\omega} \right]^n \int_{\mathbb{R}} f(x) e^{-i\omega x} dx \right]_{\omega=2\pi k} e^{i2\pi kx} \bar{\delta}_k && \text{by Definition N.2 page 309} \\ &= \frac{1}{\sqrt{2\pi}} \sum_{k \in \mathbb{Z}} \left[\int_{\mathbb{R}} f(x) (-ix)^n e^{-i\omega x} dx \right]_{\omega=2\pi k} e^{i2\pi kx} \bar{\delta}_k \\ &= (-i)^n \frac{1}{\sqrt{2\pi}} \sum_{k \in \mathbb{Z}} \left[\int_{\mathbb{R}} x^n f(x) e^{-i\omega x} dx \right]_{\omega=2\pi k} e^{i2\pi kx} \bar{\delta}_k \\ &= (-i)^n \frac{1}{\sqrt{2\pi}} \sum_{k \in \mathbb{Z}} (x - k)^n f(x - k) \bar{\delta}_k && \text{by PSF (Theorem ?? page ??)} \\ &= \frac{1}{\sqrt{2\pi}} (-i)^n \bar{\delta}_k M_n && \text{by left hypothesis} \end{aligned}$$

¹¹ Jawerth and Sweldens (1994), pages 16–17, Sweldens and Piessens (1993), page 2, Vidakovic (1999), page 83, Mallat (1999), pages 241–243, Fix and Strang (1969)

2. Proof for (\Leftarrow) case:

$$\begin{aligned}
 \frac{1}{\sqrt{2\pi}}(-i)^n M_n &= \frac{1}{\sqrt{2\pi}} \sum_{k \in \mathbb{Z}} [(-i)^n \delta_k M_n] e^{-i2\pi kx} && \text{by definition of } \delta \\
 &= \sum_{k \in \mathbb{Z}} \left[\left[\frac{d}{d\omega} \right]^n \tilde{f}(\omega) \right] \Big|_{\omega=2\pi k} e^{-i2\pi kx} && \text{by right hypothesis} \\
 &= \sum_{k \in \mathbb{Z}} \left[\left[\frac{d}{d\omega} \right]^n \int_{\mathbb{R}} f(x) e^{-i\omega x} dx \right] \Big|_{\omega=2\pi k} e^{-i2\pi kx} \\
 &= \sum_{k \in \mathbb{Z}} \left[\int_{\mathbb{R}} f(x) (-ix)^n e^{-i\omega x} dx \right] \Big|_{\omega=2\pi k} e^{-i2\pi kx} \\
 &= (-i)^n \sum_{k \in \mathbb{Z}} \left[\int_{\mathbb{R}} x^n f(x) e^{-i\omega x} dx \right] \Big|_{\omega=2\pi k} e^{-i2\pi kx} \\
 &= (-i)^n \sum_{k \in \mathbb{Z}} (x - k)^n f(x - k) && \text{by PSF} \\
 &&& \text{(Theorem ?? page ??)}
 \end{aligned}$$





N.6 Examples

Example N.1 (rectangular pulse). Let $\tilde{f}(\omega)$ be the Fourier transform of a function $f(x) \in L^2_{\mathbb{R}}$.

$$f(x) = \begin{cases} c & \text{for } x \in [a : b] \\ 0 & \text{otherwise} \end{cases}$$

$$\tilde{f}(\omega) = \frac{c(b-a)}{\sqrt{2\pi}} e^{-i\left(\frac{a+b}{2}\omega\right)} \left[\frac{\sin\left(\frac{b-a}{2}\omega\right)}{\left(\frac{b-a}{2}\omega\right)} \right]$$

E
X



PROOF:

$$\begin{aligned}
 \tilde{f}(\omega) &= \tilde{F}[f(x)](\omega) && \text{by definition of } \tilde{f}(\omega) \\
 &= e^{-i\left(\frac{a+b}{2}\omega\right)} \tilde{F}\left[f\left(x - \frac{a+b}{2}\right)\right](\omega) && \text{by shift relation} \\
 &= e^{-i\left(\frac{a+b}{2}\omega\right)} \tilde{F}\left[c \mathbb{1}_{[a:b]} \left(x - \frac{a+b}{2}\right)\right](\omega) && \text{by definition of } f(x) \\
 &= e^{-i\left(\frac{a+b}{2}\omega\right)} \tilde{F}\left[c \mathbb{1}_{\left[-\frac{b-a}{2} : \frac{b-a}{2}\right]}(x)\right](\omega) && \text{by definition of } \mathbb{1} \\
 &= \frac{1}{\sqrt{2\pi}} e^{-i\left(\frac{a+b}{2}\omega\right)} \int_{\mathbb{R}} c \mathbb{1}_{\left[-\frac{b-a}{2} : \frac{b-a}{2}\right]}(x) e^{-i\omega x} dx && \text{by definition of } \tilde{F} \\
 &= \frac{1}{\sqrt{2\pi}} e^{-i\left(\frac{a+b}{2}\omega\right)} \int_{-\frac{b-a}{2}}^{\frac{b-a}{2}} c e^{-i\omega x} dx && \text{by definition of } \mathbb{1} \\
 &= \frac{c}{\sqrt{2\pi}} e^{-i\left(\frac{a+b}{2}\omega\right)} \frac{1}{-i\omega} e^{-i\omega x} \Big|_{-\frac{b-a}{2}}^{\frac{b-a}{2}} && \\
 &= \frac{2c}{\sqrt{2\pi}\omega} e^{-i\left(\frac{a+b}{2}\omega\right)} \left[\frac{e^{i\left(\frac{b-a}{2}\omega\right)} - e^{-i\left(\frac{b-a}{2}\omega\right)}}{2i} \right] &&
 \end{aligned}$$





$$= \frac{c(b-a)}{\sqrt{2\pi}} e^{-i\left(\frac{a+b}{2}\omega\right)} \left[\frac{\sin\left(\frac{b-a}{2}\omega\right)}{\left(\frac{b-a}{2}\omega\right)} \right]$$

by Euler formulas

(Corollary ?? page ??)

Example N.2 (triangle). Let $\tilde{f}(\omega)$ be the Fourier transform of a function $f(x) \in L^2_{\mathbb{R}}$.

E X	$f(x) = \begin{cases} c \left[1 - \frac{ 2x-b-a }{b-a} \right] & \text{for } x \in [a : b) \\ 0 & \text{otherwise} \end{cases}$	$\tilde{f}(\omega) = \frac{c(b-a)^2}{4\sqrt{2\pi}} e^{-i\left(\frac{a+b}{2}\omega\right)} \left[\frac{\sin\left(\frac{b-a}{4}\omega\right)}{\left(\frac{b-a}{4}\omega\right)} \right]^2$
----------------------	--	---





PROOF:

$$\begin{aligned}
 \tilde{f}(\omega) &= \tilde{F}[f(x)](\omega) && \text{by definition of } \tilde{f}(\omega) \\
 &= e^{-i\left(\frac{a+b}{2}\omega\right)} \tilde{F}\left[f\left(x - \frac{a+b}{2}\right)\right](\omega) && \text{by shift relation} \quad (\text{Theorem N.4 page 311}) \\
 &= \tilde{F}\left[c\left(1 - \frac{|2x-b-a|}{b-a}\right) \mathbb{1}_{[a:b)}(x)\right](\omega) && \text{by definition of } f(x) \\
 &= c \tilde{F}\left[\mathbb{1}_{\left[\frac{a}{2}:\frac{b}{2}\right)}(x) \star \mathbb{1}_{\left[\frac{a}{2}:\frac{b}{2}\right)}(x)\right](\omega) \\
 &= c \sqrt{2\pi} \tilde{F}\left[\mathbb{1}_{\left[\frac{a}{2}:\frac{b}{2}\right)}\right] \tilde{F}\left[\mathbb{1}_{\left[\frac{a}{2}:\frac{b}{2}\right)}\right] && \text{by convolution theorem} \quad (\text{Theorem P.2 page 332}) \\
 &= c \sqrt{2\pi} \left(\tilde{F}\left[\mathbb{1}_{\left[\frac{a}{2}:\frac{b}{2}\right)}\right] \right)^2 \\
 &= c \sqrt{2\pi} \left(\frac{\left(\frac{b}{2} - \frac{a}{2}\right)}{\sqrt{2\pi}} e^{-i\left(\frac{a+b}{4}\omega\right)} \left[\frac{\sin\left(\frac{b-a}{4}\omega\right)}{\left(\frac{b-a}{4}\omega\right)} \right] \right)^2 && \text{by Rectangular pulse ex.} \quad \text{Example N.1 page 316} \\
 &= \frac{c(b-a)^2}{4\sqrt{2\pi}} e^{-i\left(\frac{a+b}{2}\omega\right)} \left[\frac{\sin\left(\frac{b-a}{4}\omega\right)}{\left(\frac{b-a}{4}\omega\right)} \right]^2
 \end{aligned}$$

Example N.3. Let a function f be defined in terms of the cosine function (Definition ?? page ??) as follows:

E X	$f(x) \triangleq \begin{cases} \cos^2\left(\frac{\pi}{2}x\right) & \text{for } x \leq 1 \\ 0 & \text{otherwise} \end{cases}$	$\tilde{f}(\omega) = \frac{1}{2\sqrt{2\pi}} \left[\underbrace{\frac{2\sin\omega}{\omega}}_{2\operatorname{sinc}(\omega)} + \underbrace{\frac{\sin(\omega+\pi)}{(\omega+\pi)}}_{\operatorname{sinc}(\omega+\pi)} + \underbrace{\frac{\sin(\omega-\pi)}{(\omega-\pi)}}_{\operatorname{sinc}(\omega-\pi)} \right]$
----------------------	--	--



PROOF: Let $\mathbb{1}_A(x)$ be the *set indicator function* (Definition ?? page ??) on a set A .

$$\begin{aligned}
 \tilde{f}(\omega) &\triangleq \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} f(x)e^{-i\omega x} dx && \text{by definition of } \tilde{f}(\omega) \text{ (Definition N.2)} \\
 &= \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} \cos^2\left(\frac{\pi}{2}x\right) \mathbb{1}_{[-1:1]}(x) e^{-i\omega x} dx && \text{by definition of } f(x) \\
 &= \frac{1}{\sqrt{2\pi}} \int_{-1}^1 \cos^2\left(\frac{\pi}{2}x\right) e^{-i\omega x} dx && \text{by definition of } \mathbb{1} \text{ (Definition ??)} \\
 &= \frac{1}{\sqrt{2\pi}} \int_{-1}^1 \left[\frac{e^{i\frac{\pi}{2}x} + e^{-i\frac{\pi}{2}x}}{2} \right]^2 e^{-i\omega x} dx && \text{by Corollary ?? page ??} \\
 &= \frac{1}{4\sqrt{2\pi}} \int_{-1}^1 [2 + e^{i\pi x} + e^{-i\pi x}] e^{-i\omega x} dx \\
 &= \frac{1}{4\sqrt{2\pi}} \int_{-1}^1 2e^{-i\omega x} + e^{-i(\omega+\pi)x} + e^{-i(\omega-\pi)x} dx \\
 &= \frac{1}{4\sqrt{2\pi}} \left[2 \frac{e^{-i\omega x}}{-i\omega} + \frac{e^{-i(\omega+\pi)x}}{-i(\omega+\pi)} + \frac{e^{-i(\omega-\pi)x}}{-i(\omega-\pi)} \right]_{-1}^1 \\
 &= \frac{1}{2\sqrt{2\pi}} \left[2 \frac{e^{-i\omega} - e^{+i\omega}}{-2i\omega} + \frac{e^{-i(\omega+\pi)} - e^{+i(\omega+\pi)}}{-2i(\omega+\pi)} + \frac{e^{-i(\omega-\pi)} - e^{+i(\omega-\pi)}}{-2i(\omega-\pi)} \right]_{-1}^1 \\
 &= \frac{1}{2\sqrt{2\pi}} \left[\underbrace{\frac{2\sin\omega}{\omega}}_{2 \operatorname{sinc}(\omega)} + \underbrace{\frac{\sin(\omega+\pi)}{(\omega+\pi)}}_{\operatorname{sinc}(\omega+\pi)} + \underbrace{\frac{\sin(\omega-\pi)}{(\omega-\pi)}}_{\operatorname{sinc}(\omega-\pi)} \right]
 \end{aligned}$$



APPENDIX O

DISCRETE TIME FOURIER TRANSFORM


O.1 Definition

Definition O.1.

D E F The discrete-time Fourier transform \check{F} of $(x_n)_{n \in \mathbb{Z}}$ is defined as

$$[\check{F}(x_n)](\omega) \triangleq \sum_{n \in \mathbb{Z}} x_n e^{-i\omega n} \quad \forall (x_n)_{n \in \mathbb{Z}} \in \ell^2_{\mathbb{R}}$$

If we compare the definition of the *Discrete Time Fourier Transform* (Definition O.1 page 319) to the definition of the Z-transform (Definition P.4 page 330), we see that the DTFT is just a special case of the more general Z-Transform, with $z = e^{i\omega}$. If we imagine $z \in \mathbb{C}$ as a complex plane, then $e^{i\omega}$ is a unit circle in this plane. The “frequency” ω in the DTFT is the unit circle in the much larger z-plane, as illustrated to the right.



O.2 Properties

Proposition O.1 (DTFT periodicity). Let $\check{x}(\omega) \triangleq \check{F}[(x_n)](\omega)$ be the DISCRETE-TIME FOURIER TRANSFORM (Definition O.1 page 319) of a sequence $(x_n)_{n \in \mathbb{Z}}$ in $\ell^2_{\mathbb{R}}$.

P R P

$\check{x}(\omega) = \underbrace{\check{x}(\omega + 2\pi n)}_{\text{PERIODIC with period } 2\pi} \quad \forall n \in \mathbb{Z}$
--

PROOF:

$$\begin{aligned} \check{x}(\omega + 2\pi n) &= \sum_{m \in \mathbb{Z}} x_m e^{-i(\omega+2\pi n)m} \\ &= \sum_{m \in \mathbb{Z}} x_m e^{-i\omega m} e^{-i2\pi nm} \\ &= \check{x}(\omega) \end{aligned}$$



Theorem O.1. Let $\tilde{x}(\omega) \triangleq \check{F}[(x[n])](\omega)$ be the DISCRETE-TIME FOURIER TRANSFORM (Definition O.1 page 319) of a sequence $(x_n)_{n \in \mathbb{Z}}$ in $\ell^2_{\mathbb{R}}$.

T H M	$\left\{ \begin{array}{l} \tilde{x}(\omega) \triangleq \check{F}(x[n]) \\ \end{array} \right\} \implies \left\{ \begin{array}{l} (1). \quad \check{F}(x[-n]) = \tilde{x}(-\omega) \quad \text{and} \\ (2). \quad \check{F}(x^*[n]) = \tilde{x}^*(-\omega) \quad \text{and} \\ (3). \quad \check{F}(x^*[-n]) = \tilde{x}^*(\omega) \end{array} \right\}$
-------------	---

PROOF:

$$\begin{aligned} \check{F}(x[-n]) &\triangleq \sum_{n \in \mathbb{Z}} x[-n]e^{-i\omega n} && \text{by definition of DTFT} && (\text{Definition O.1 page 319}) \\ &= \sum_{m \in \mathbb{Z}} x[m]e^{i\omega m} && \text{where } m \triangleq -n \implies n = -m \\ &= \sum_{m \in \mathbb{Z}} x[m]e^{-i(-\omega)m} \\ &\triangleq \tilde{x}(-\omega) && \text{by left hypothesis} \end{aligned}$$

$$\begin{aligned} \check{F}(x^*[n]) &\triangleq \sum_{n \in \mathbb{Z}} x^*[n]e^{-i\omega n} && \text{by definition of DTFT} && (\text{Definition O.1 page 319}) \\ &= \left(\sum_{n \in \mathbb{Z}} x[n]e^{i\omega n} \right)^* && \text{by distributive property of } *-\text{algebras} && (\text{Definition H.3 page 228}) \\ &= \left(\sum_{n \in \mathbb{Z}} x[n]e^{-i(-\omega)n} \right)^* \\ &\triangleq \tilde{x}^*(-\omega) && \text{by left hypothesis} \end{aligned}$$

$$\begin{aligned} \check{F}(x^*[-n]) &\triangleq \sum_{n \in \mathbb{Z}} x^*[-n]e^{-i\omega n} && \text{by definition of DTFT} && (\text{Definition O.1 page 319}) \\ &= \left(\sum_{n \in \mathbb{Z}} x[-n]e^{i\omega n} \right)^* && \text{by distributive property of } *-\text{algebras} && (\text{Definition H.3 page 228}) \\ &= \left(\sum_{m \in \mathbb{Z}} x[m]e^{-i\omega m} \right)^* && \text{where } m \triangleq -n \implies n = -m \\ &\triangleq \tilde{x}^*(\omega) && \text{by left hypothesis} \end{aligned}$$

Theorem O.2. Let $\tilde{x}(\omega) \triangleq \check{F}[(x[n])](\omega)$ be the DISCRETE-TIME FOURIER TRANSFORM (Definition O.1 page 319) of a sequence $(x[n])_{n \in \mathbb{Z}}$ in $\ell^2_{\mathbb{R}}$.

T H M	$\left\{ \begin{array}{l} (1). \quad \tilde{x}(\omega) \triangleq \check{F}(x[n]) \\ (2). \quad (x[n]) \text{ is REAL-VALUED} \end{array} \right\} \implies \left\{ \begin{array}{l} (1). \quad \check{F}(x[-n]) = \tilde{x}(-\omega) \quad \text{and} \\ (2). \quad \check{F}(x^*[n]) = \tilde{x}^*(-\omega) = \tilde{x}(\omega) \quad \text{and} \\ (3). \quad \check{F}(x^*[-n]) = \tilde{x}^*(\omega) = \tilde{x}(-\omega) \end{array} \right\}$
-------------	--


PROOF:

$$\begin{aligned} \check{F}(x[-n]) &\triangleq \sum_{n \in \mathbb{Z}} x[-n]e^{-i\omega n} && \text{by definition of DTFT} && (\text{Definition O.1 page 319}) \\ &= \sum_{m \in \mathbb{Z}} x[m]e^{i\omega m} && \text{where } m \triangleq -n \implies n = -m \\ &= \sum_{m \in \mathbb{Z}} x[m]e^{-i(-\omega)m} \end{aligned}$$

$$\triangleq \check{x}(-\omega) \quad \text{by left hypothesis}$$

$$\begin{aligned} \check{x}^*(-\omega) &= \check{\mathbf{F}}(\mathbf{x}^*[n]) && \text{by Theorem O.1 page 320} \\ &= \check{\mathbf{F}}(\mathbf{x}[n]) && \text{by real-valued hypothesis} \\ &= [\check{x}(\omega)] && \text{by definition of } \check{x}(\omega) \quad (\text{Definition O.1 page 319}) \end{aligned}$$

$$\begin{aligned} \check{x}^*(\omega) &= \check{\mathbf{F}}(\mathbf{x}^*[-n]) && \text{by Theorem O.1 page 320} \\ &= \check{\mathbf{F}}(\mathbf{x}[-n]) && \text{by real-valued hypothesis} \\ &= [\check{x}(-\omega)] && \text{by result (1)} \end{aligned}$$



Proposition O.2. Let $\check{x}(z)$ be the Z-TRANSFORM (Definition P.4 page 330) and $\check{x}(\omega)$ the DISCRETE-TIME FOURIER TRANSFORM (Definition O.1 page 319) of (x_n) .

P	$\left\{ \sum_{n \in \mathbb{Z}} x_n = c \right\}$	\Leftrightarrow	$\underbrace{\left\{ \check{x}(z) \Big _{z=1} = c \right\}}_{(2) z \text{ domain}}$	\Leftrightarrow	$\underbrace{\left\{ \check{x}(\omega) \Big _{\omega=0} = c \right\}}_{(3) \text{ frequency domain}}$	$\forall (x_n)_{n \in \mathbb{Z}} \in \ell^2_{\mathbb{R}}, c \in \mathbb{R}$
P R P	$\underbrace{(1) \text{ time domain}}$					

PROOF:

1. Proof that (1) \Rightarrow (2):

$$\begin{aligned} \check{x}(z) \Big|_{z=1} &= \sum_{n \in \mathbb{Z}} x_n z^{-n} \Big|_{z=1} && \text{by definition of } \check{x}(z) \text{ (Definition P.4 page 330)} \\ &= \sum_{n \in \mathbb{Z}} x_n && \text{because } z^n = 1 \text{ for all } n \in \mathbb{Z} \\ &= c && \text{by hypothesis (1)} \end{aligned}$$

2. Proof that (2) \Rightarrow (3):

$$\begin{aligned} \check{x}(\omega) \Big|_{\omega=0} &= \sum_{n \in \mathbb{Z}} x_n e^{-i\omega n} \Big|_{\omega=0} && \text{by definition of } \check{x}(\omega) \quad (\text{Definition O.1 page 319}) \\ &= \sum_{n \in \mathbb{Z}} x_n z^{-n} \Big|_{z=1} && \\ &= \check{x}(z) \Big|_{z=1} && \text{by definition of } \check{x}(z) \quad (\text{Definition P.4 page 330}) \\ &= c && \text{by hypothesis (2)} \end{aligned}$$

3. Proof that (3) \implies (1):

$$\begin{aligned} \sum_{n \in \mathbb{Z}} x_n &= \sum_{n \in \mathbb{Z}} x_n e^{-i\omega n} \Big|_{\omega=0} \\ &= \check{x}(\omega) && \text{by definition of } \check{x}(\omega) && (\text{Definition O.1 page 319}) \\ &= c && \text{by hypothesis (3)} \end{aligned}$$



Proposition O.3. *If the coefficients are real, then the magnitude response (MR) is symmetric.*

PROOF:

$$\begin{aligned} |\tilde{h}(-\omega)| &\triangleq |\check{h}(z)|_{z=e^{-i\omega}} \\ &= \left| \sum_{m \in \mathbb{Z}} x[m] e^{i\omega m} \right| \\ &= \underbrace{\left| \left(\sum_{m \in \mathbb{Z}} x[m] e^{-i\omega m} \right)^* \right|}_{\text{if } x[m] \text{ is real}} \\ &\triangleq |\check{h}(z)|_{z=e^{-i\omega}} && \triangleq |\tilde{h}(\omega)| \end{aligned}$$



Proposition O.4.¹

P R P	$\underbrace{\sum_{n \in \mathbb{Z}} (-1)^n x_n = c}_{(1) \text{ in "time"}} \iff \underbrace{\check{x}(z) _{z=-1} = c}_{(2) \text{ in "z domain"}} \iff \underbrace{\check{x}(\omega) _{\omega=\pi} = c}_{(3) \text{ in "frequency"}}$ $\iff \underbrace{\left(\sum_{n \in \mathbb{Z}} h_{2n}, \sum_{n \in \mathbb{Z}} h_{2n+1} \right)}_{(4) \text{ sum of even, sum of odd}} = \left(\frac{1}{2} \left(\sum_{n \in \mathbb{Z}} h_n + c \right), \frac{1}{2} \left(\sum_{n \in \mathbb{Z}} h_n - c \right) \right)$ $\forall c \in \mathbb{R}, (x_n)_{n \in \mathbb{Z}}, (y_n)_{n \in \mathbb{Z}} \in \ell^2_{\mathbb{R}}$
-------------	--

PROOF:

1. Proof that (1) \implies (2):

$$\begin{aligned} \check{x}(z)|_{z=-1} &= \sum_{n \in \mathbb{Z}} x_n z^{-n} \Big|_{z=-1} \\ &= \sum_{n \in \mathbb{Z}} (-1)^n x_n \\ &= c && \text{by (1)} \end{aligned}$$

¹ Chui (1992) page 123

2. Proof that (2) \implies (3):

$$\begin{aligned} \sum_{n \in \mathbb{Z}} x_n e^{-i\omega n} \Big|_{\omega=\pi} &= \sum_{n \in \mathbb{Z}} (-1)^n x_n \\ &= \sum_{n \in \mathbb{Z}} (-1)^{-n} x_n && = \sum_{n \in \mathbb{Z}} z^{-n} x_n \Big|_{z=-1} \\ &= c && \text{by (2)} \end{aligned}$$

3. Proof that (3) \implies (1):

$$\begin{aligned} \sum_{n \in \mathbb{Z}} (-1)^n x_n &= \sum_{n \in \mathbb{Z}} (-1)^{-n} x_n \\ &= \sum_{n \in \mathbb{Z}} e^{-i\omega n} x_n \Big|_{\omega=\pi} \\ &= c && \text{by (3)} \end{aligned}$$

4. Proof that (2) \implies (4):

$$(a) \text{ Define } A \triangleq \sum_{n \in \mathbb{Z}} h_{2n} \quad B \triangleq \sum_{n \in \mathbb{Z}} h_{2n+1}.$$

(b) Proof that $A - B = c$:

$$\begin{aligned} c &= \sum_{n \in \mathbb{Z}} (-1)^n x_n && \text{by (2)} \\ &= \underbrace{\sum_{n \in \mathbb{Z}_e} (-1)^n x_n}_{\text{even terms}} + \underbrace{\sum_{n \in \mathbb{Z}_o} (-1)^n x_n}_{\text{odd terms}} \\ &= \sum_{n \in \mathbb{Z}} (-1)^{2n} x_{2n} + \sum_{n \in \mathbb{Z}} (-1)^{2n+1} x_{2n+1} \\ &= \underbrace{\sum_{n \in \mathbb{Z}} x_{2n}}_A - \underbrace{\sum_{n \in \mathbb{Z}} x_{2n+1}}_B \\ &\triangleq A - B && \text{by definitions of } A \text{ and } B \end{aligned}$$

$$(c) \text{ Proof that } A + B = \sum_{n \in \mathbb{Z}} x_n:$$

$$\begin{aligned} \sum_{n \in \mathbb{Z}} x_n &= \sum_{n \text{ even}} x_n + \sum_{n \text{ odd}} x_n \\ &= \underbrace{\sum_{n \in \mathbb{Z}} x_{2n}}_A + \underbrace{\sum_{n \in \mathbb{Z}} x_{2n+1}}_B \\ &= A + B && \text{by definitions of } A \text{ and } B \end{aligned}$$

(d) This gives two simultaneous equations:

$$A - B = c$$

$$A + B = \sum_{n \in \mathbb{Z}} x_n$$

(e) Solutions to these equations give

$$\begin{aligned}\sum_{n \in \mathbb{Z}} x_{2n} &\triangleq A \\ \sum_{n \in \mathbb{Z}} x_{2n+1} &\triangleq B\end{aligned}\begin{aligned}&= \frac{1}{2} \left(\sum_{n \in \mathbb{Z}} x_n + c \right) \\ &= \frac{1}{2} \left(\sum_{n \in \mathbb{Z}} x_n - c \right)\end{aligned}$$

5. Proof that (2) \Leftarrow (4):

$$\begin{aligned}\sum_{n \in \mathbb{Z}} (-1)^n x_n &= \underbrace{\sum_{n \in \mathbb{Z}_e} (-1)^n x_n}_{\text{even terms}} + \underbrace{\sum_{n \in \mathbb{Z}_o} (-1)^n x_n}_{\text{odd terms}} \\ &= \sum_{n \in \mathbb{Z}} (-1)^{2n} x_{2n} + \sum_{n \in \mathbb{Z}} (-1)^{2n+1} x_{2n+1} \\ &= \sum_{n \in \mathbb{Z}} x_{2n} - \sum_{n \in \mathbb{Z}} x_{2n+1} \\ &= \frac{1}{2} \left(\sum_{n \in \mathbb{Z}} x_n + c \right) - \frac{1}{2} \left(\sum_{n \in \mathbb{Z}} x_n - c \right) \quad \text{by (3)} \\ &= c\end{aligned}$$



Lemma O.1. Let $\tilde{f}(\omega)$ be the DTFT (Definition O.1 page 319) of a sequence $(x_n)_{n \in \mathbb{Z}}$.

L E M	$\underbrace{(x_n \in \mathbb{R})_{n \in \mathbb{Z}}}_{\text{REAL-VALUED sequence}}$	\Rightarrow	$\underbrace{ \check{x}(\omega) ^2 = \check{x}(-\omega) ^2}_{\text{EVEN}}$	$\forall (x_n)_{n \in \mathbb{Z}} \in \ell^2_{\mathbb{R}}$
----------------------------------	--	---------------	---	--

PROOF:

$$\begin{aligned}|\check{x}(\omega)|^2 &= |\check{x}(z)|^2 \Big|_{z=e^{i\omega}} \\ &= \check{x}(z)\check{x}^*(z) \Big|_{z=e^{i\omega}} \\ &= \left[\sum_{n \in \mathbb{Z}} x_n z^{-n} \right] \left[\sum_{m \in \mathbb{Z}} x_m z^{-m} \right]^* \Big|_{z=e^{i\omega}} \\ &= \left[\sum_{n \in \mathbb{Z}} x_n z^{-n} \right] \left[\sum_{m \in \mathbb{Z}} x_m^* (z^*)^{-m} \right] \Big|_{z=e^{i\omega}} \\ &= \sum_{n \in \mathbb{Z}} \sum_{m \in \mathbb{Z}} x_n x_m^* z^{-n} (z^*)^{-m} \Big|_{z=e^{i\omega}} \\ &= \sum_{n \in \mathbb{Z}} \left[|x_n|^2 + \sum_{m > n} x_n x_m^* z^{-n} (z^*)^{-m} + \sum_{m < n} x_n x_m^* z^{-n} (z^*)^{-m} \right] \Big|_{z=e^{i\omega}} \\ &= \sum_{n \in \mathbb{Z}} \left[|x_n|^2 + \sum_{m > n} x_n x_m e^{i\omega(m-n)} + \sum_{m < n} x_n x_m^* e^{i\omega(m-n)} \right] \\ &= \sum_{n \in \mathbb{Z}} \left[|x_n|^2 + \sum_{m > n} x_n x_m e^{i\omega(m-n)} + \sum_{m > n} x_n x_m e^{-i\omega(m-n)} \right] \\ &= \sum_{n \in \mathbb{Z}} \left[|x_n|^2 + \sum_{m > n} x_n x_m (e^{i\omega(m-n)} + e^{-i\omega(m-n)}) \right]\end{aligned}$$

$$\begin{aligned}
 &= \sum_{n \in \mathbb{Z}} \left[|x_n|^2 + \sum_{m>n} x_n x_m 2\cos[\omega(m-n)] \right] \\
 &= \sum_{n \in \mathbb{Z}} |x_n|^2 + 2 \sum_{n \in \mathbb{Z}} \sum_{m>n} x_n x_m \cos[\omega(m-n)]
 \end{aligned}$$

Since \cos is real and even, then $|\check{x}(\omega)|^2$ must also be real and even. \Rightarrow

Theorem O.3 (inverse DTFT). ² Let $\check{x}(\omega)$ be the DISCRETE-TIME FOURIER TRANSFORM (Definition O.1 page 319) of a sequence $(x_n)_{n \in \mathbb{Z}} \in \ell^2_{\mathbb{R}}$. Let \check{x}^{-1} be the inverse of \check{x} .

T H M	$\underbrace{\left\{ \check{x}(\omega) \triangleq \sum_{n \in \mathbb{Z}} x_n e^{-i\omega n} \right\}}_{\check{x}(\omega) \triangleq \check{F}(x_n)} \quad \Rightarrow \quad \underbrace{\left\{ x_n = \frac{1}{2\pi} \int_{-\pi}^{\pi} \check{x}(\omega) e^{i\omega n} d\omega \quad \forall \omega \in \mathbb{R} \right\}}_{(x_n) = \check{F}^{-1}(\check{x}(\omega))} \quad \forall (x_n)_{n \in \mathbb{Z}} \in \ell^2_{\mathbb{R}}$
----------------------	---

\Leftarrow PROOF:

$$\begin{aligned}
 \frac{1}{2\pi} \int_{-\pi}^{\pi} \check{x}(\omega) e^{i\omega n} d\omega &= \frac{1}{2\pi} \int_{-\pi}^{\pi} \underbrace{\left[\sum_{m \in \mathbb{Z}} x_m e^{-i\omega m} \right]}_{\check{x}(\omega)} e^{i\omega n} d\omega && \text{by definition of } \check{x}(\omega) \\
 &= \frac{1}{2\pi} \int_{-\pi}^{\pi} \sum_{m \in \mathbb{Z}} x_m e^{-i\omega(m-n)} d\omega \\
 &= \frac{1}{2\pi} \sum_{m \in \mathbb{Z}} x_m \int_{-\pi}^{\pi} e^{-i\omega(m-n)} d\omega \\
 &= \frac{1}{2\pi} \sum_{m \in \mathbb{Z}} x_m [2\pi \delta_{m-n}] \\
 &= x_n
 \end{aligned}$$

Theorem O.4 (orthonormal quadrature conditions). ³ Let $\check{x}(\omega)$ be the DISCRETE-TIME FOURIER TRANSFORM (Definition O.1 page 319) of a sequence $(x_n)_{n \in \mathbb{Z}} \in \ell^2_{\mathbb{R}}$. Let $\bar{\delta}_n$ be the KRONECKER DELTA FUNCTION at n (Definition I.3 page 245).

T H M	$ \begin{aligned} \sum_{m \in \mathbb{Z}} x_m y_{m-2n}^* &= 0 \iff \check{x}(\omega) \check{y}^*(\omega) + \check{x}(\omega + \pi) \check{y}^*(\omega + \pi) = 0 && \forall n \in \mathbb{Z}, \forall (x_n), (y_n) \in \ell^2_{\mathbb{R}} \\ \sum_{m \in \mathbb{Z}} x_m x_{m-2n}^* &= \bar{\delta}_n \iff \check{x}(\omega) ^2 + \check{x}(\omega + \pi) ^2 = 2 && \forall n \in \mathbb{Z}, \forall (x_n), (y_n) \in \ell^2_{\mathbb{R}} \end{aligned} $
----------------------	--

\Leftarrow PROOF: Let $z \triangleq e^{i\omega}$.

²  J.S.Chitode (2009) page 3-95 ((3.6.2))

³  Daubechies (1992) pages 132-137 ((5.1.20),(5.1.39))

1. Proof that $2 \sum_{n \in \mathbb{Z}} \left[\sum_{k \in \mathbb{Z}} x_k y_{k-2n}^* \right] e^{-i2\omega n} = \check{x}(\omega) \check{y}^*(\omega) + \check{x}(\omega + \pi) \check{y}^*(\omega + \pi)$:

$$\begin{aligned}
 & 2 \sum_{n \in \mathbb{Z}} \left[\sum_{k \in \mathbb{Z}} x_k y_{k-2n}^* \right] e^{-i2\omega n} \\
 &= 2 \sum_{k \in \mathbb{Z}} x_k \sum_{n \in \mathbb{Z}} y_{k-2n}^* z^{-2n} \\
 &= 2 \sum_{k \in \mathbb{Z}} x_k \sum_{n \text{ even}} y_{k-n}^* z^{-n} \\
 &= \sum_{k \in \mathbb{Z}} x_k \sum_{n \in \mathbb{Z}} y_{k-n}^* z^{-n} (1 + e^{i\pi n}) \\
 &= \sum_{k \in \mathbb{Z}} x_k \sum_{n \in \mathbb{Z}} y_{k-n}^* z^{-n} + \sum_{k \in \mathbb{Z}} x_k \sum_{n \in \mathbb{Z}} y_{k-n}^* z^{-n} e^{i\pi n} \\
 &= \sum_{k \in \mathbb{Z}} x_k \sum_{m \in \mathbb{Z}} y_m^* z^{-(k-m)} + \sum_{k \in \mathbb{Z}} x_k \sum_{m \in \mathbb{Z}} y_m^* e^{-i(\omega+\pi)(k-m)} \quad \text{where } m \triangleq k - n \\
 &= \sum_{k \in \mathbb{Z}} x_k z^{-k} \sum_{m \in \mathbb{Z}} y_m^* z^m + \sum_{k \in \mathbb{Z}} x_k e^{-i(\omega+\pi)k} \sum_{m \in \mathbb{Z}} y_m^* e^{+i(\omega+\pi)m} \\
 &= \sum_{k \in \mathbb{Z}} x_k e^{-i\omega k} \left[\sum_{m \in \mathbb{Z}} y_m^* e^{-i\omega m} \right]^* + \sum_{k \in \mathbb{Z}} x_k e^{-i(\omega+\pi)k} \left[\sum_{m \in \mathbb{Z}} y_m^* e^{-i(\omega+\pi)m} \right]^* \\
 &\triangleq \check{x}(\omega) \check{y}^*(\omega) + \check{x}(\omega + \pi) \check{y}^*(\omega + \pi)
 \end{aligned}$$

2. Proof that $\sum_{m \in \mathbb{Z}} x_m y_{m-2n}^* = 0 \implies \check{x}(\omega) \check{y}^*(\omega) + \check{x}(\omega + \pi) \check{y}^*(\omega + \pi) = 0$:

$$\begin{aligned}
 0 &= 2 \sum_{n \in \mathbb{Z}} \left[\sum_{k \in \mathbb{Z}} x_k y_{k-2n}^* \right] e^{-i2\omega n} && \text{by left hypothesis} \\
 &= \check{x}(\omega) \check{y}^*(\omega) + \check{x}(\omega + \pi) \check{y}^*(\omega + \pi) && \text{by item (1)}
 \end{aligned}$$

3. Proof that $\sum_{m \in \mathbb{Z}} x_m y_{m-2n}^* = 0 \iff \check{x}(\omega) \check{y}^*(\omega) + \check{x}(\omega + \pi) \check{y}^*(\omega + \pi) = 0$:

$$\begin{aligned}
 2 \sum_{n \in \mathbb{Z}} \left[\sum_{k \in \mathbb{Z}} x_k y_{k-2n}^* \right] e^{-i2\omega n} &= \check{x}(\omega) \check{y}^*(\omega) + \check{x}(\omega + \pi) \check{y}^*(\omega + \pi) && \text{by item (1)} \\
 &= 0 && \text{by right hypothesis}
 \end{aligned}$$

Thus by the above equation, $\sum_{n \in \mathbb{Z}} \left[\sum_{k \in \mathbb{Z}} x_k y_{k-2n}^* \right] e^{-i2\omega n} = 0$. The only way for this to be true is if $\sum_{k \in \mathbb{Z}} x_k y_{k-2n}^* = 0$.

4. Proof that $\sum_{m \in \mathbb{Z}} x_m x_{m-2n}^* = \bar{\delta}_n \implies |\check{x}(\omega)|^2 + |\check{x}(\omega + \pi)|^2 = 2$:

Let $g_n \triangleq x_n$.

$$\begin{aligned}
 2 &= 2 \sum_{n \in \mathbb{Z}} \bar{\delta}_n e^{-i2\omega n} \\
 &= 2 \sum_{n \in \mathbb{Z}} \left[\sum_{k \in \mathbb{Z}} x_k y_{k-2n}^* \right] e^{-i2\omega n} && \text{by left hypothesis} \\
 &= \check{x}(\omega) \check{y}^*(\omega) + \check{x}(\omega + \pi) \check{y}^*(\omega + \pi) && \text{by item (1)}
 \end{aligned}$$

5. Proof that $\sum_{m \in \mathbb{Z}} x_m x_{m-2n}^* = \bar{\delta}_n \iff |\check{x}(\omega)|^2 + |\check{x}(\omega + \pi)|^2 = 2$:

Let $g_n \triangleq x_n$.

$$\begin{aligned}
 2 \sum_{n \in \mathbb{Z}} \left[\sum_{k \in \mathbb{Z}} x_k y_{k-2n}^* \right] e^{-i2\omega n} &= \check{x}(\omega) \check{y}^*(\omega) + \check{x}(\omega + \pi) \check{y}^*(\omega + \pi) && \text{by item (1)} \\
 &= 2 && \text{by right hypothesis}
 \end{aligned}$$

Thus by the above equation, $\sum_{n \in \mathbb{Z}} [\sum_{k \in \mathbb{Z}} x_k y_{k-2n}^*] e^{-i2\omega n} = 1$. The only way for this to be true is if $\sum_{k \in \mathbb{Z}} x_k y_{k-2n}^* = \delta_n$.



O.3 Derivatives

Theorem O.5. ⁴ Let $\check{x}(\omega)$ be the DTFT (Definition O.1 page 319) of a sequence $(x_n)_{n \in \mathbb{Z}}$.

T H M	$(A) \quad \left[\frac{d}{d\omega} \right]^n \check{x}(\omega) \Big _{\omega=0} = 0 \iff \sum_{k \in \mathbb{Z}} k^n x_k = 0 \quad (B) \quad \forall n \in \mathbb{W}$
	$(C) \quad \left[\frac{d}{d\omega} \right]^n \check{x}(\omega) \Big _{\omega=\pi} = 0 \iff \sum_{k \in \mathbb{Z}} (-1)^k k^n x_k = 0 \quad (D) \quad \forall n \in \mathbb{W}$

PROOF:

1. Proof that (A) \implies (B):

$$\begin{aligned}
 0 &= \left[\frac{d}{d\omega} \right]^n \check{x}(\omega) \Big|_{\omega=0} && \text{by hypothesis (A)} \\
 &= \left[\frac{d}{d\omega} \right]^n \sum_{k \in \mathbb{Z}} x_k e^{-i\omega k} \Big|_{\omega=0} && \text{by definition of } \check{x}(\omega) \text{ (Definition O.1 page 319)} \\
 &= \sum_{k \in \mathbb{Z}} x_k \left[\frac{d}{d\omega} \right]^n e^{-i\omega k} \Big|_{\omega=0} \\
 &= \sum_{k \in \mathbb{Z}} x_k \left[(-i)^n k^n e^{-i\omega k} \right] \Big|_{\omega=0} \\
 &= (-i)^n \sum_{k \in \mathbb{Z}} k^n x_k
 \end{aligned}$$

2. Proof that (A) \iff (B):

$$\begin{aligned}
 \left[\frac{d}{d\omega} \right]^n \check{x}(\omega) \Big|_{\omega=0} &= \left[\frac{d}{d\omega} \right]^n \sum_{k \in \mathbb{Z}} x_k e^{-i\omega k} \Big|_{\omega=0} && \text{by definition of } \check{x}(\omega) \\
 &= \sum_{k \in \mathbb{Z}} x_k \left[\left[\frac{d}{d\omega} \right]^n e^{-i\omega k} \right] \Big|_{\omega=0} \\
 &= \sum_{k \in \mathbb{Z}} x_k \left[(-i)^n k^n e^{-i\omega k} \right] \Big|_{\omega=0} \\
 &= (-i)^n \sum_{k \in \mathbb{Z}} k^n x_k \\
 &= 0 && \text{by hypothesis (B)}
 \end{aligned}$$

⁴ Vidakovic (1999), pages 82–83, Mallat (1999), pages 241–242

3. Proof that (C) \implies (D):

$$\begin{aligned}
 0 &= \left[\frac{d}{d\omega} \right]^n \check{x}(\omega) \Big|_{\omega=\pi} && \text{by hypothesis (C)} \\
 &= \left[\frac{d}{d\omega} \right]^n \sum_{k \in \mathbb{Z}} x_k e^{-i\omega k} \Big|_{\omega=\pi} && \text{by definition of } \check{x} \text{ (Definition O.1 page 319)} \\
 &= \sum_{k \in \mathbb{Z}} x_k \left[\frac{d}{d\omega} \right]^n e^{-i\omega k} \Big|_{\omega=\pi} \\
 &= \sum_{k \in \mathbb{Z}} x_k \left[(-i)^n k^n e^{-i\omega k} \right] \Big|_{\omega=\pi} \\
 &= \sum_{k \in \mathbb{Z}} x_k \left[(-i)^n k^n (-1)^k \right] \\
 &= (-i)^n \sum_{k \in \mathbb{Z}} (-1)^k k^n x_k
 \end{aligned}$$

4. Proof that (C) \iff (D):

$$\begin{aligned}
 \left[\frac{d}{d\omega} \right]^n \check{x}(\omega) \Big|_{\omega=\pi} &= \left[\frac{d}{d\omega} \right]^n \sum_{k \in \mathbb{Z}} x_k e^{-i\omega k} \Big|_{\omega=\pi} && \text{by definition of } \check{x} \text{ (Definition O.1 page 319)} \\
 &= \sum_{k \in \mathbb{Z}} x_k \left[\frac{d}{d\omega} \right]^n e^{-i\omega k} \Big|_{\omega=\pi} \\
 &= \sum_{k \in \mathbb{Z}} x_k \left[(-i)^n k^n e^{-i\omega k} \right] \Big|_{\omega=\pi} \\
 &= \sum_{k \in \mathbb{Z}} x_k \left[(-i)^n k^n (-1)^k \right] \\
 &= (-i)^n \sum_{k \in \mathbb{Z}} (-1)^k k^n x_k \\
 &= 0 && \text{by hypothesis (D)}
 \end{aligned}$$



APPENDIX P

OPERATIONS ON SEQUENCES

P.1 Convolution operator

Definition P.1. ¹ Let X^Y be the set of all functions from a set Y to a set X . Let \mathbb{Z} be the set of integers.

D E F A function f in X^Y is a **sequence** over X if $Y = \mathbb{Z}$.

A sequence may be denoted in the form $(x_n)_{n \in \mathbb{Z}}$ or simply as (x_n) .

Definition P.2. ² Let $(\mathbb{F}, +, \cdot, 0, 1)$ be a FIELD (Definition ?? page ??).

D E F The space of all absolutely square summable sequences $\ell_{\mathbb{F}}^2$ over \mathbb{F} is defined as

$$\ell_{\mathbb{F}}^2 \triangleq \left\{ (x_n)_{n \in \mathbb{Z}} \mid \sum_{n \in \mathbb{Z}} |x_n|^2 < \infty \right\}$$

The space $\ell_{\mathbb{R}}^2$ is an example of a *separable Hilbert space*. In fact, $\ell_{\mathbb{R}}^2$ is the *only* separable Hilbert space in the sense that all separable Hilbert spaces are isomorphically equivalent. For example, $\ell_{\mathbb{R}}^2$ is isomorphic to $L_{\mathbb{R}}^2$, the space of all absolutely square Lebesgue integrable functions.

Definition P.3.

D E F The **convolution operation \star** is defined as

$$(x_n) \star (y_n) \triangleq \left(\sum_{m \in \mathbb{Z}} x_m y_{n-m} \right)_{n \in \mathbb{Z}} \quad \forall (x_n)_{n \in \mathbb{Z}}, (y_n)_{n \in \mathbb{Z}} \in \ell_{\mathbb{R}}^2$$

Proposition P.1. Let \star be the CONVOLUTION OPERATOR (Definition P.3 page 329).

P R P $(x_n) \star (y_n) = (y_n) \star (x_n) \quad \forall (x_n)_{n \in \mathbb{Z}}, (y_n)_{n \in \mathbb{Z}} \in \ell_{\mathbb{R}}^2 \quad (\star \text{ is COMMUTATIVE})$

¹ Bromwich (1908), page 1, Thomson et al. (2008) page 23 (Definition 2.1), Joshi (1997) page 31

² Kubrusly (2011) page 347 (Example 5.K)

PROOF:

$$\begin{aligned}
 [x \star y](n) &\triangleq \sum_{m \in \mathbb{Z}} x_m y_{n-m} && \text{by Definition P.3 page 329} \\
 &= \sum_{k \in \mathbb{Z}} x_{n-k} y(k) && \text{where } k = n - m \iff m = n - k \\
 &= \sum_{k \in \mathbb{Z}} x_{n-k} y(k) && \text{by change commutivity of addition} \\
 &= \sum_{m \in \mathbb{Z}} x_{n-m} y_m && \text{by change of variables} \\
 &= \sum_{m \in \mathbb{Z}} y_m x_{n-m} && \text{by commutative property of the field over } \mathbb{C} \\
 &\triangleq (y \star x)_n && \text{by Definition P.3 page 329}
 \end{aligned}$$



Proposition P.2. Let \star be the CONVOLUTION OPERATOR (Definition P.3 page 329). Let $\ell^2_{\mathbb{R}}$ be the set of ABSOLUTELY SUMMABLE sequences (Definition P.2 page 329).

$$\boxed{\begin{array}{l} \text{P} \\ \text{R} \\ \text{P} \end{array} \left\{ \begin{array}{l} (A). \quad x(n) \in \ell^2_{\mathbb{R}} \text{ and} \\ (B). \quad y(n) \in \ell^2_{\mathbb{R}} \end{array} \right\} \Rightarrow \left\{ \sum_{k \in \mathbb{Z}} x[k] y[n+k] = x[-n] \star y(n) \right\}}$$

PROOF:

$$\begin{aligned}
 \sum_{k \in \mathbb{Z}} x[k] y[n+k] &= \sum_{-p \in \mathbb{Z}} x[-p] y[n-p] && \text{where } p \triangleq -k && \Rightarrow k = -p \\
 &= \sum_{p \in \mathbb{Z}} x[-p] y[n-p] && \text{by absolutely summable hypothesis} && \text{(Definition P.2 page 329)} \\
 &= \sum_{p \in \mathbb{Z}} x'[p] y[n-p] && \text{where } x'[n] \triangleq x[-n] && \Rightarrow x[-n] = x'[n] \\
 &\triangleq x'[n] \star y[n] && \text{by definition of convolution } \star && \text{(Definition P.3 page 329)} \\
 &\triangleq x[-n] \star y[n] && \text{by definition of } x'[n]
 \end{aligned}$$



P.2 Z-transform

Definition P.4. ³

$$\boxed{\begin{array}{l} \text{D} \\ \text{E} \\ \text{F} \end{array} \text{The z-transform } \mathbf{Z} \text{ of } (x_n)_{n \in \mathbb{Z}} \text{ is defined as} \\
 [\mathbf{Z}(x_n)](z) \triangleq \underbrace{\sum_{n \in \mathbb{Z}} x_n z^{-n}}_{\text{Laurent series}} \quad \forall (x_n) \in \ell^2_{\mathbb{R}}}$$

Theorem P.1. Let $X(z) \triangleq \mathbf{Z}x[n]$ be the Z-TRANSFORM of $x[n]$.

$$\boxed{\begin{array}{l} \text{T} \\ \text{H} \\ \text{M} \end{array} \left\{ \check{x}(z) \triangleq \mathbf{Z}(x[n]) \right\} \Rightarrow \left\{ \begin{array}{l} (1). \quad \mathbf{Z}(\alpha x[n]) = \alpha \check{x}(z) \quad \forall (x_n) \in \ell^2_{\mathbb{R}} \text{ and} \\ (2). \quad \mathbf{Z}(x[n-k]) = z^{-k} \check{x}(z) \quad \forall (x_n) \in \ell^2_{\mathbb{R}} \text{ and} \\ (3). \quad \mathbf{Z}(x[-n]) = \check{x}\left(\frac{1}{z}\right) \quad \forall (x_n) \in \ell^2_{\mathbb{R}} \text{ and} \\ (4). \quad \mathbf{Z}(x^*[n]) = \check{x}^*\left(z^*\right) \quad \forall (x_n) \in \ell^2_{\mathbb{R}} \text{ and} \\ (5). \quad \mathbf{Z}(x^*[-n]) = \check{x}^*\left(\frac{1}{z^*}\right) \quad \forall (x_n) \in \ell^2_{\mathbb{R}} \end{array} \right\}}$$

³ Laurent series: [Abramovich and Aliprantis \(2002\) page 49](#)

PROOF:

$$\begin{aligned}
 \alpha \mathbb{Z} \check{x}(z) &\triangleq \alpha \mathbb{Z}(\check{x}[n]) && \text{by definition of } \check{x}(z) \\
 &\triangleq \alpha \sum_{n \in \mathbb{Z}} x[n] z^{-n} && \text{by definition of } \mathbb{Z} \text{ operator} \\
 &\triangleq \sum_{n \in \mathbb{Z}} (\alpha x[n]) z^{-n} && \text{by distributive property} \\
 &\triangleq \mathbb{Z}(\alpha x[n]) && \text{by definition of } \mathbb{Z} \text{ operator} \\
 z^{-k} \check{x}(z) &= z^{-k} \mathbb{Z}(x[n]) && \text{by definition of } \check{x}(z) \quad (\text{left hypothesis}) \\
 &\triangleq z^{-k} \sum_{n=-\infty}^{n=+\infty} x[n] z^{-n} && \text{by definition of } \mathbb{Z} \quad (\text{Definition P.4 page 330}) \\
 &= \sum_{n=-\infty}^{n=+\infty} x[n] z^{-n-k} \\
 &= \sum_{m=k=-\infty}^{m=k=+\infty} x[m-k] z^{-m} && \text{where } m \triangleq n+k \implies n = m - k \\
 &= \sum_{m=-\infty}^{m=+\infty} x[m-k] z^{-m} \\
 &= \sum_{n=-\infty}^{n=+\infty} x[n-k] z^{-n} && \text{where } n \triangleq m \\
 &\triangleq \mathbb{Z}(x[n-k]) && \text{by definition of } \mathbb{Z} \quad (\text{Definition P.4 page 330}) \\
 \mathbb{Z}(x^*[n]) &\triangleq \sum_{n \in \mathbb{Z}} x^*[n] z^{-n} && \text{by definition of } \mathbb{Z} \quad (\text{Definition P.4 page 330}) \\
 &\triangleq \left(\sum_{n \in \mathbb{Z}} x[n] (z^*)^{-n} \right)^* && \text{by definition of } \mathbb{Z} \quad (\text{Definition P.4 page 330}) \\
 &\triangleq \check{x}^*(z^*) && \text{by definition of } \mathbb{Z} \quad (\text{Definition P.4 page 330}) \\
 \mathbb{Z}(x[-n]) &\triangleq \sum_{n \in \mathbb{Z}} x[-n] z^{-n} && \text{by definition of } \mathbb{Z} \quad (\text{Definition P.4 page 330}) \\
 &= \sum_{-m \in \mathbb{Z}} x[m] z^m && \text{where } m \triangleq -n \implies n = -m \\
 &= \sum_{m \in \mathbb{Z}} x[m] z^m && \text{by absolutely summable property} \quad (\text{Definition P.2 page 329}) \\
 &= \sum_{m \in \mathbb{Z}} x[m] \left(\frac{1}{z} \right)^{-m} && \text{by absolutely summable property} \quad (\text{Definition P.2 page 329}) \\
 &\triangleq \check{x}\left(\frac{1}{z}\right) && \text{by definition of } \mathbb{Z} \quad (\text{Definition P.4 page 330}) \\
 \mathbb{Z}(x^*[-n]) &\triangleq \sum_{n \in \mathbb{Z}} x^*[-n] z^{-n} && \text{by definition of } \mathbb{Z} \quad (\text{Definition P.4 page 330}) \\
 &= \sum_{-m \in \mathbb{Z}} x^*[m] z^m && \text{where } m \triangleq -n \implies n = -m \\
 &= \sum_{m \in \mathbb{Z}} x^*[m] z^m && \text{by absolutely summable property} \quad (\text{Definition P.2 page 329}) \\
 &= \sum_{m \in \mathbb{Z}} x^*[m] \left(\frac{1}{z} \right)^{-m} && \text{by absolutely summable property} \quad (\text{Definition P.2 page 329}) \\
 &= \left(\sum_{m \in \mathbb{Z}} x[m] \left(\frac{1}{z^*} \right)^{-m} \right)^* && \text{by absolutely summable property} \quad (\text{Definition P.2 page 329})
 \end{aligned}$$




Figure P.1: Direct form 1 order 2 IIR filters

$$\triangleq \check{x}^* \left(\frac{1}{z^*} \right) \quad \text{by definition of } \mathbf{Z} \quad (\text{Definition P.4 page 330})$$

Theorem P.2 (convolution theorem). *Let \star be the convolution operator (Definition P.3 page 329).*

T H M	$\mathbf{Z} \underbrace{\left(\langle x_n \rangle \star \langle y_n \rangle \right)}_{\text{sequence convolution}} = \underbrace{\left(\mathbf{Z} \langle x_n \rangle \right) \left(\mathbf{Z} \langle y_n \rangle \right)}_{\text{series multiplication}}$	$\forall \langle x_n \rangle_{n \in \mathbb{Z}}, \langle y_n \rangle_{n \in \mathbb{Z}} \in \ell^2_{\mathbb{R}}$
-------------	--	--

PROOF:

$$\begin{aligned}
 [\mathbf{Z}(x \star y)](z) &\triangleq \mathbf{Z} \left(\sum_{m \in \mathbb{Z}} x_m y_{n-m} \right) && \text{by Definition P.3 page 329} \\
 &\triangleq \sum_{n \in \mathbb{Z}} \sum_{m \in \mathbb{Z}} x_m y_{n-m} z^{-n} && \text{by Definition P.4 page 330} \\
 &= \sum_{n \in \mathbb{Z}} \sum_{m \in \mathbb{Z}} x_m y_{n-m} z^{-n} \\
 &= \sum_{m \in \mathbb{Z}} \sum_{n \in \mathbb{Z}} x_m y_{n-m} z^{-n} \\
 &= \sum_{m \in \mathbb{Z}} \sum_{k \in \mathbb{Z}} x_m y_k z^{-(m+k)} && \text{where } k = n - m \iff n = m + k \\
 &= \left[\sum_{m \in \mathbb{Z}} x_m z^{-m} \right] \left[\sum_{k \in \mathbb{Z}} y_k z^{-k} \right] \\
 &\triangleq (\mathbf{Z} \langle x_n \rangle) (\mathbf{Z} \langle y_n \rangle) && \text{by Definition P.4 page 330}
 \end{aligned}$$

P.3 From z-domain back to time-domain

$$\check{y}(z) = b_0 X(z) + b_1 z^{-1} X(z) + b_2 z^{-2} X(z) - a_1 z^{-1} Y(z) + a_2 z^{-2} Y(z)$$

$$y[n] = b_0 x[n] + b_1 x[n-1] + b_2 x[n-2] - a_1 y[n-1] - a_2 y[n-2]$$

Example P.1. See Figure P.1 (page 332)

$$\frac{3z^2 + 5z + 7}{2z^2 + 10z + 12} = \frac{3z^2 + 5z + 7}{2(z^2 + 5z + 6)} = \frac{(3z^2 + 5z + 7)z^{-2}}{z^2 + 5z + 6} = \frac{(3z^2 + 5z + 7)z^{-2}}{1 + 5z^{-1} + 6z^{-2}}$$

P.4 Zero locations

The system property of *minimum phase* is defined in Definition P.5 (next) and illustrated in Figure P.2 (page 333).




Figure P.2: Minimum Phase filter

Definition P.5. ⁴ Let $\check{x}(z) \triangleq \mathbf{Z}(x_n)$ be the Z TRANSFORM (Definition P.4 page 330) of a sequence $(x_n)_{n \in \mathbb{Z}}$ in $\ell_{\mathbb{R}}^2$. Let $(z_n)_{n \in \mathbb{Z}}$ be the ZEROS of $\check{x}(z)$.

DEF The sequence (x_n) is **minimum phase** if

$$\underbrace{|z_n| < 1}_{\check{x}(z) \text{ has all its ZEROS inside the unit circle}} \quad \forall n \in \mathbb{Z}$$

The impulse response of a minimum phase filter has most of its energy concentrated near the beginning of its support, as demonstrated next.

Theorem P.3 (Robinson's Energy Delay Theorem). ⁵ Let $p(z) \triangleq \sum_{n=0}^N a_n z^{-n}$ and $q(z) \triangleq \sum_{n=0}^N b_n z^{-n}$ be polynomials.

THM $\left\{ \begin{array}{l} p \text{ is MINIMUM PHASE} \\ q \text{ is NOT minimum phase} \end{array} \right. \text{ and } \Rightarrow \sum_{n=0}^{m-1} |a_n|^2 \geq \sum_{n=0}^{m-1} |b_n|^2 \quad \forall 0 \leq m \leq N$

$$\underbrace{\text{"energy" of the first } m \text{ coefficients of } p(z)}_{\text{of the first } m \text{ coefficients of } p(z)} \geq \underbrace{\text{"energy" of the first } m \text{ coefficients of } q(z)}_{\text{of the first } m \text{ coefficients of } q(z)}$$

But for more *symmetry*, put some zeros inside and some outside the unit circle.

Example P.2. An example of a minimum phase polynomial is the Daubechies-4 scaling function. The minimum phase polynomial causes most of the energy to be concentrated near the origin, making it very *asymmetric*. In contrast, the Symlet-4 has a design very similar to that of Daubechies-4, but the selected zeros are not all within the unit circle in the complex z plane. This results in a

⁴ Farina and Rinaldi (2000) page 91, Dumitrescu (2007) page 36

⁵ Dumitrescu (2007) page 36, Robinson (1962), Robinson (1966) (???), Claerbout (1976), pages 52–53




Figure P.3: Daubechies-4 and Symlet-4 scaling functions pole-zero plots

scaling function that is more symmetric and less contracted near the origin. Both scaling functions are illustrated in Figure P.3 (page 334).

P.5 Pole locations

Definition P.6.

D E F A filter (or system or operator) \mathbf{H} is **causal** if its current output does not depend on future inputs.

Definition P.7.

D E F A filter (or system or operator) \mathbf{H} is **time-invariant** if the mapping it performs does not change with time.

Definition P.8.

D E F An operation \mathbf{H} is **linear** if any output y_n can be described as a linear combination of inputs x_n as in

$$y_n = \sum_{m \in \mathbb{Z}} h(m)x(n-m).$$

For a filter to be *stable*, place all the poles *inside* the unit circle.

Theorem P.4. A causal LTI filter is **stable** if all of its poles are **inside** the unit circle.

Example P.3. Stable/unstable filters are illustrated in Figure P.4 (page 335).

True or False? This filter has no poles:








Figure P.4: Pole-zero plot stable/unstable causal LTI filters (Example P.3 page 334)

$$H(z) = b_0 + b_1 z^{-1} + b_2 z^{-2}$$



$$H(z) = b_0 + b_1 z^{-1} + b_2 z^{-2} = \frac{z^2}{z^2} \times \frac{b_0 + b_1 z^{-1} + b_2 z^{-2}}{1} = \frac{b_0 z^2 + b_1 z^1 + b_2}{z^2}$$



P.6 Mirroring for real coefficients

If you want real coefficients, choose poles and zeros in conjugate pairs (next).

Proposition P.3.

P R P	$\left\{ \begin{array}{l} \text{ZEROS and POLES} \\ \text{occur in CONJUGATE PAIRS} \end{array} \right\} \Rightarrow \left\{ \begin{array}{l} \text{COEFFICIENTS} \\ \text{are REAL.} \end{array} \right\}$
-------------	---




Figure P.5: Conjugate pair structure yielding real coefficients

PROOF:

$$\begin{aligned}(z - p_1)(z - p_1^*) &= [z - (a + ib)][z - (a - ib)] \\&= z^2 + [-a + ib - ib - a]z - [ib]^2 \\&= z^2 - 2az + b^2\end{aligned}$$

Example P.4. See Figure P.5 (page 336).

$$\begin{aligned}H(z) &= G \frac{[z - z_1][z - z_2]}{[z - p_1][z - p_2]} = G \frac{[z - (1+i)][z - (1-i)]}{[z - (-^{2/3} + i^{1/2})][z - (-^{2/3} - i^{1/2})]} \\&= G \frac{z^2 - z[(1-i) + (1+i)] + (1-i)(1+i)}{z^2 - z[(-^{2/3} + i^{1/2}) + (-^{2/3} - i^{1/2})] + (-^{2/3} + i^{1/2})(-^{2/3} - i^{1/2})} \\&= G \frac{z^2 - 2z + 2}{z^2 - \frac{4}{3}z + (\frac{1}{3} + \frac{1}{4})} = G \frac{z^2 - 2z + 2}{z^2 - \frac{4}{3}z + \frac{19}{12}}\end{aligned}$$

P.7 Rational polynomial operators

A digital filter is simply an operator on $\ell_{\mathbb{R}}^2$. If the digital filter is a causal LTI system, then it can be expressed as a rational polynomial in z as shown next.

Lemma P.1. A causal LTI operator H can be expressed as a rational expression $\check{h}(z)$.

$$\begin{aligned}\check{h}(z) &= \frac{b_0 + b_1 z^{-1} + b_2 z^{-2} + \dots + b_N z^{-N}}{1 + a_1 z^{-1} + a_2 z^{-2} + \dots + a_N z^{-N}} \\&= \frac{\sum_{n=0}^N b_n z^{-n}}{1 + \sum_{n=1}^N a_n z^{-n}}\end{aligned}$$

A filter operation $\check{h}(z)$ can be expressed as a product of its roots (poles and zeros).

$$\begin{aligned}\check{h}(z) &= \frac{b_0 + b_1 z^{-1} + b_2 z^{-2} + \dots + b_N z^{-N}}{1 + a_1 z^{-1} + a_2 z^{-2} + \dots + a_N z^{-N}} \\&= \alpha \frac{(z - z_1)(z - z_2) \dots (z - z_N)}{(z - p_1)(z - p_2) \dots (z - p_N)}\end{aligned}$$

where α is a constant, z_i are the zeros, and p_i are the poles. The poles and zeros of such a rational expression are often plotted in the z-plane with a unit circle about the origin (representing $z = e^{i\omega}$). Poles are marked with \times and zeros with \circ . An example is shown in Figure P.6 page 337. Notice that in this figure the zeros and poles are either real or occur in complex conjugate pairs.




Figure P.6: Pole-zero plot for rational expression with real coefficients

P.8 Filter Banks

Conjugate quadrature filters (next definition) are used in *filter banks*. If $\check{x}(z)$ is a *low-pass filter*, then the conjugate quadrature filter of $\check{y}(z)$ is a *high-pass filter*.

Definition P.9.⁶ Let $(x_n)_{n \in \mathbb{Z}}$ and $(y_n)_{n \in \mathbb{Z}}$ be SEQUENCES (Definition P.1 page 329) in $\ell^2_{\mathbb{R}}$ (Definition P.2 page 329).

D E F The sequence (y_n) is a **conjugate quadrature filter** with shift N with respect to (x_n) if
 $y_n = \pm(-1)^n x_{N-n}^*$

A CONJUGATE QUADRATURE FILTER is also called a **CQF** or a **Smith-Barnwell filter**.

Any triple $((x_n), (y_n), N)$ in this form is said to satisfy the
conjugate quadrature filter condition or the **CQF condition**.

Theorem P.5 (CQF theorem).⁷ Let $\check{y}(\omega)$ and $\check{x}(\omega)$ be the DTFTs (Definition O.1 page 319) of the sequences $(y_n)_{n \in \mathbb{Z}}$ and $(x_n)_{n \in \mathbb{Z}}$, respectively, in $\ell^2_{\mathbb{R}}$ (Definition P.2 page 329).

T H M	$\underbrace{y_n = \pm(-1)^n x_{N-n}^*}_{(1) \text{ CQF in "time"} } \iff \check{y}(z) = \pm(-1)^N z^{-N} \check{x}^*\left(\frac{-1}{z^*}\right) \quad (2) \text{ CQF in "z-domain"}$ $\iff \check{y}(\omega) = \pm(-1)^N e^{-i\omega N} \check{x}^*(\omega + \pi) \quad (3) \text{ CQF in "frequency"}$ $\iff x_n = \pm(-1)^N (-1)^n y_{N-n}^* \quad (4) \text{ "reversed" CQF in "time"}$ $\iff \check{x}(z) = \pm z^{-N} \check{y}^*\left(\frac{-1}{z^*}\right) \quad (5) \text{ "reversed" CQF in "z-domain"}$ $\iff \check{x}(\omega) = \pm e^{-i\omega N} \check{y}^*(\omega + \pi) \quad (6) \text{ "reversed" CQF in "frequency"}$
-------	--

$\forall N \in \mathbb{Z}$

PROOF:

⁶ Strang and Nguyen (1996) page 109, Haddad and Akansu (1992) pages 256–259 (section 4.5), Vaidyanathan (1993) page 342 ((7.2.7), (7.2.8)), Smith and Barnwell (1984a), Smith and Barnwell (1984b), Mintzer (1985)

⁷ Strang and Nguyen (1996) page 109, Mallat (1999) pages 236–238 ((7.58),(7.73)), Haddad and Akansu (1992) pages 256–259 (section 4.5), Vaidyanathan (1993) page 342 ((7.2.7), (7.2.8))

1. Proof that (1) \Rightarrow (2):

$$\begin{aligned}
 \check{y}(z) &= \sum_{n \in \mathbb{Z}} y_n z^{-n} && \text{by definition of } z\text{-transform} && (\text{Definition P.4 page 330}) \\
 &= \sum_{n \in \mathbb{Z}} (\pm)(-1)^n x_{N-n}^* z^{-n} && \text{by (1)} \\
 &= \pm \sum_{m \in \mathbb{Z}} (-1)^{N-m} x_m^* z^{-(N-m)} && \text{where } m \triangleq N - n \Rightarrow && n = N - m \\
 &= \pm(-1)^N z^{-N} \sum_{m \in \mathbb{Z}} (-1)^{-m} x_m^* (z^{-1})^{-m} \\
 &= \pm(-1)^N z^{-N} \sum_{m \in \mathbb{Z}} x_m^* \left(-\frac{1}{z}\right)^{-m} \\
 &= \pm(-1)^N z^{-N} \left[\sum_{m \in \mathbb{Z}} x_m \left(-\frac{1}{z^*}\right)^{-m} \right]^* \\
 &= \pm(-1)^N z^{-N} \check{x}^* \left(\frac{-1}{z^*}\right) && \text{by definition of } z\text{-transform} && (\text{Definition P.4 page 330})
 \end{aligned}$$

2. Proof that (1) \Leftarrow (2):

$$\begin{aligned}
 \check{y}(z) &= \pm(-1)^N z^{-N} \check{x}^* \left(\frac{-1}{z^*}\right) && \text{by (2)} \\
 &= \pm(-1)^N z^{-N} \left[\sum_{m \in \mathbb{Z}} x_m \left(\frac{-1}{z^*}\right)^{-m} \right]^* && \text{by definition of } z\text{-transform} && (\text{Definition P.4 page 330}) \\
 &= \pm(-1)^N z^{-N} \left[\sum_{m \in \mathbb{Z}} x_m^* (-z^{-1})^{-m} \right] && \text{by definition of } z\text{-transform} && (\text{Definition P.4 page 330}) \\
 &= \sum_{m \in \mathbb{Z}} (\pm)(-1)^{N-m} x_m^* z^{-(N-m)} \\
 &= \sum_{m \in \mathbb{Z}} (\pm)(-1)^n x_{N-n}^* z^{-n} && \text{where } n = N - m \Rightarrow && m \triangleq N - n \\
 &\Rightarrow x_n = \pm(-1)^n x_{N-n}^*
 \end{aligned}$$

3. Proof that (1) \Rightarrow (3):

$$\begin{aligned}
 \check{y}(\omega) &\triangleq \check{x}(z) \Big|_{z=e^{i\omega}} && \text{by definition of DTFT (Definition O.1 page 319)} \\
 &= \left[\pm(-1)^N z^{-N} \check{x} \left(\frac{-1}{z^*}\right) \right]_{z=e^{i\omega}} && \text{by (2)} \\
 &= \pm(-1)^N e^{-i\omega N} \check{x} (e^{i\pi} e^{i\omega}) \\
 &= \pm(-1)^N e^{-i\omega N} \check{x} (e^{i(\omega+\pi)}) \\
 &= \pm(-1)^N e^{-i\omega N} \check{x}(\omega + \pi) && \text{by definition of DTFT (Definition O.1 page 319)}
 \end{aligned}$$

4. Proof that (1) \Rightarrow (6):

$$\begin{aligned}
 \check{x}(\omega) &= \sum_{n \in \mathbb{Z}} y_n e^{-i\omega n} && \text{by definition of DTFT} && (\text{Definition O.1 page 319}) \\
 &= \sum_{n \in \mathbb{Z}} \underbrace{\pm(-1)^n x_{N-n}^*}_{CQF} e^{-i\omega n} && \text{by (1)} \\
 &= \sum_{m \in \mathbb{Z}} \pm(-1)^{N-m} x_m^* e^{-i\omega(N-m)} && \text{where } m \triangleq N - n \Rightarrow && n = N - m \\
 &= \pm(-1)^N e^{-i\omega N} \sum_{m \in \mathbb{Z}} (-1)^m x_m^* e^{i\omega m}
 \end{aligned}$$



$$\begin{aligned}
&= \pm(-1)^N e^{-i\omega N} \sum_{m \in \mathbb{Z}} e^{i\pi m} x_m^* e^{i\omega m} \\
&= \pm(-1)^N e^{-i\omega N} \sum_{m \in \mathbb{Z}} x_m^* e^{i(\omega+\pi)m} \\
&= \pm(-1)^N e^{-i\omega N} \left[\sum_{m \in \mathbb{Z}} x_m e^{-i(\omega+\pi)m} \right]^* \\
&= \pm(-1)^N e^{-i\omega N} \check{x}^*(\omega + \pi) \quad \text{by definition of DTFT} \quad (\text{Definition O.1 page 319})
\end{aligned}$$

5. Proof that (1) \iff (3):

$$\begin{aligned}
y_n &= \frac{1}{2\pi} \int_{-\pi}^{+\pi} \check{y}(\omega) e^{i\omega n} d\omega \quad \text{by inverse DTFT} \quad (\text{Theorem O.3 page 325}) \\
&= \frac{1}{2\pi} \int_{-\pi}^{+\pi} \underbrace{\pm(-1)^N e^{-iN\omega}}_{\text{right hypothesis}} \check{x}^*(\omega + \pi) e^{i\omega n} d\omega \quad \text{by right hypothesis} \\
&= \pm(-1)^N \frac{1}{2\pi} \int_{-\pi}^{+\pi} \check{x}^*(\omega + \pi) e^{i\omega(n-N)} d\omega \quad \text{by right hypothesis} \\
&= \pm(-1)^N \frac{1}{2\pi} \int_0^{2\pi} \check{x}^*(v) e^{i(v-\pi)(n-N)} dv \quad \text{where } v \triangleq \omega + \pi \implies \omega = v - \pi \\
&= \pm(-1)^N e^{-i\pi(n-N)} \frac{1}{2\pi} \int_0^{2\pi} \check{x}^*(v) e^{iv(n-N)} dv \\
&= \pm(-1)^N \underbrace{(-1)^N}_{e^{i\pi N}} \underbrace{(-1)^n}_{e^{-i\pi n}} \left[\frac{1}{2\pi} \int_0^{2\pi} \check{x}(v) e^{iv(N-n)} dv \right]^* \\
&= \pm(-1)^n x_{N-n}^* \quad \text{by inverse DTFT} \quad (\text{Theorem O.3 page 325})
\end{aligned}$$

6. Proof that (1) \iff (4):

$$\begin{aligned}
y_n = \pm(-1)^n x_{N-n}^* &\iff (\pm)(-1)^n y_n = (\pm)(\pm)(-1)^n (-1)^n x_{N-n}^* \\
&\iff \pm(-1)^n y_n = x_{N-n}^* \\
&\iff (\pm(-1)^n y_n)^* = (x_{N-n}^*)^* \\
&\iff \pm(-1)^n y_n^* = x_{N-n} \\
&\iff x_{N-n} = \pm(-1)^n y_n^* \\
&\iff x_m = \pm(-1)^{N-m} y_{N-m}^* \quad \text{where } m \triangleq N - n \implies n = N - m \\
&\iff x_m = \pm(-1)^{N-m} y_{N-m}^* \\
&\iff x_m = \pm(-1)^N (-1)^m y_{N-m}^* \\
&\iff x_n = \pm(-1)^N (-1)^n y_{N-n}^* \quad \text{by change of free variables}
\end{aligned}$$

7. Proofs for (5) and (6): not included. See proofs for (2) and (3).



Theorem P.6. ⁸ Let $\check{y}(\omega)$ and $\check{x}(\omega)$ be the DTFTs (Definition O.1 page 319) of the sequences $(y_n)_{n \in \mathbb{Z}}$ and $(x_n)_{n \in \mathbb{Z}}$, respectively, in $\ell^2_{\mathbb{R}}$ (Definition P.2 page 329).

T H M

Let $y_n = \pm(-1)^n x_{N-n}^*$ (CQF CONDITION, Definition P.9 page 337). Then

$$\left\{
\begin{aligned}
(A) \quad \left[\frac{d}{d\omega} \right]^n \check{y}(\omega) \Big|_{\omega=0} &= 0 \iff \left[\frac{d}{d\omega} \right]^n \check{x}(\omega) \Big|_{\omega=\pi} = 0 && (B) \\
&\iff \sum_{k \in \mathbb{Z}} (-1)^k k^n x_k &= 0 && (C) \\
&\iff \sum_{k \in \mathbb{Z}} k^n y_k &= 0 && (D)
\end{aligned}
\right\} \forall n \in \mathbb{W}$$

⁸ Vidakovic (1999), pages 82–83, Mallat (1999), pages 241–242

PROOF:

1. Proof that (A) \implies (B):


$$\begin{aligned}
 0 &= \left[\frac{d}{d\omega} \right]^n \check{y}(\omega) \Big|_{\omega=0} && \text{by (A)} \\
 &= \left[\frac{d}{d\omega} \right]^n (\pm)(-1)^N e^{-i\omega N} \check{x}^*(\omega + \pi) \Big|_{\omega=0} && \text{by CQF theorem (Theorem P.5 page 337)} \\
 &= (\pm)(-1)^N \sum_{\ell=0}^n \binom{n}{\ell} \left[\frac{d}{d\omega} \right]^\ell [e^{-i\omega N}] \cdot \left[\frac{d}{d\omega} \right]^{n-\ell} [\check{x}^*(\omega + \pi)] \Big|_{\omega=0} && \text{by Leibnitz GPR (Lemma ?? page ??)} \\
 &= (\pm)(-1)^N \sum_{\ell=0}^n \binom{n}{\ell} -iN^\ell e^{-i\omega N} \left[\frac{d}{d\omega} \right]^{n-\ell} [\check{x}^*(\omega + \pi)] \Big|_{\omega=0} \\
 &= (\pm)(-1)^N e^{-i0N} \sum_{\ell=0}^n \binom{n}{\ell} -iN^\ell \left[\frac{d}{d\omega} \right]^{n-\ell} [\check{x}^*(\omega + \pi)] \Big|_{\omega=0} \\
 &\implies \check{x}^{(0)}(\pi) = 0 \\
 &\implies \check{x}^{(1)}(\pi) = 0 \\
 &\implies \check{x}^{(2)}(\pi) = 0 \\
 &\implies \check{x}^{(3)}(\pi) = 0 \\
 &\implies \check{x}^{(4)}(\pi) = 0 \\
 &\vdots \quad \vdots \\
 &\implies \check{x}^{(n)}(\pi) = 0 \quad \text{for } n = 0, 1, 2, \dots
 \end{aligned}$$

2. Proof that (A) \iff (B):

$$\begin{aligned}
 0 &= \left[\frac{d}{d\omega} \right]^n \check{x}(\omega) \Big|_{\omega=\pi} && \text{by (B)} \\
 &= \left[\frac{d}{d\omega} \right]^n (\pm) e^{-i\omega N} \check{y}^*(\omega + \pi) \Big|_{\omega=\pi} && \text{by CQF theorem (Theorem P.5 page 337)} \\
 &= (\pm) \sum_{\ell=0}^n \binom{n}{\ell} \left[\frac{d}{d\omega} \right]^\ell [e^{-i\omega N}] \cdot \left[\frac{d}{d\omega} \right]^{n-\ell} [\check{y}^*(\omega + \pi)] \Big|_{\omega=\pi} && \text{by Leibnitz GPR (Lemma ?? page ??)} \\
 &= (\pm) \sum_{\ell=0}^n \binom{n}{\ell} (-iN)^\ell e^{-i\omega N} \left[\frac{d}{d\omega} \right]^{n-\ell} [\check{y}^*(\omega + \pi)] \Big|_{\omega=\pi} \\
 &= (\pm) e^{-i\pi N} \sum_{\ell=0}^n \binom{n}{\ell} -iN^\ell \left[\frac{d}{d\omega} \right]^{n-\ell} [\check{y}^*(\omega + \pi)] \Big|_{\omega=\pi} \\
 &= (\pm)(-1)^N \sum_{\ell=0}^n \binom{n}{\ell} -iN^\ell \left[\frac{d}{d\omega} \right]^{n-\ell} [\check{y}^*(\omega + \pi)] \Big|_{\omega=\pi} \\
 &\implies \check{y}^{(0)}(0) = 0 \\
 &\implies \check{y}^{(1)}(0) = 0 \\
 &\implies \check{y}^{(2)}(0) = 0 \\
 &\implies \check{y}^{(3)}(0) = 0 \\
 &\implies \check{y}^{(4)}(0) = 0 \\
 &\vdots \quad \vdots \\
 &\implies \check{y}^{(n)}(0) = 0 \quad \text{for } n = 0, 1, 2, \dots
 \end{aligned}$$

3. Proof that (B) \iff (C): by Theorem O.5 page 327


4. Proof that (A) \iff (D): by Theorem O.5 page 327



5. Proof that (CQF) $\not\Leftarrow$ (A): Here is a counterexample: $\check{y}(\omega) = 0$.

P.9 Inverting non-minimum phase filters

Minimum phase filters are easy to invert: each *zero* becomes a *pole* and each *pole* becomes a *zero*.



$$\begin{aligned}
|A(z)|_{z=e^{i\omega}} &= \frac{1}{r} \left| \frac{z - re^{i\phi}}{z - \frac{1}{r}e^{i\phi}} \right|_{z=e^{i\omega}} \\
&= \left| e^{i\phi} \left(\frac{e^{-i\phi}z - r}{rz - e^{i\phi}} \right) \right|_{z=e^{i\omega}} \\
&= \left| -z \left(\frac{rz^{-1} - e^{-i\phi}}{rz - e^{i\phi}} \right) \right|_{z=e^{i\omega}} \\
&= \left| \frac{1}{e^{-iv}} \left(\frac{re^{-i\omega} - e^{-i\phi}}{(re^{i\omega} - e^{i\phi})^*} \right) \right| \\
&\equiv 1
\end{aligned}$$

$$\begin{aligned}
&= \left| \frac{z - re^{i\phi}}{rz - e^{i\phi}} \right|_{z=e^{i\omega}} \\
&= \left| z \left(\frac{e^{-i\phi} - rz^{-1}}{rz - e^{i\phi}} \right) \right|_{z=e^{i\omega}} \\
&= \left| \widetilde{e^{-i\pi}} e^{i\omega} \left(\frac{re^{-i\omega} - e^{-i\phi}}{re^{i\omega} - e^{i\phi}} \right) \right| \\
&= \left| \frac{re^{-i\omega} - e^{-i\phi}}{re^{-i\omega} - e^{i\phi}} \right|
\end{aligned}$$

APPENDIX Q


B-SPLINES

Q.1 Definitions

Definition Q.1. Let X be a set.

D E F The **step function** $\sigma \in \mathbb{R}^{\mathbb{R}}$ is defined as

$$\sigma(x) \triangleq \mathbb{1}_{[0:\infty)}(x) \quad \forall x \in \mathbb{R}.$$



Lemma Q.1. Let $\sigma(x)$ be the STEP FUNCTION (Definition Q.1 page 343).

L E M $\{g(x) > 0\} \implies \{\sigma[g(x)f(x)] = \sigma[f(x)]\} \quad \forall f, g \in \mathbb{R}^{\mathbb{R}}$

PROOF:

$$\begin{aligned}
 \sigma[g(x)f(x)] &\triangleq \mathbb{1}_{[0:\infty)}[g(x)f(x)] && \text{by definition of } \sigma(x) && (\text{Definition Q.1 page 343}) \\
 &\triangleq \begin{cases} 1 & \text{for } g(x)f(x) \geq 0 \\ 0 & \text{otherwise} \end{cases} && \text{by definition of } \mathbb{1}_A(x) && (\text{Definition ?? page ??}) \\
 &= \begin{cases} 1 & \text{for } f(x) \geq 0 \\ 0 & \text{otherwise} \end{cases} && \text{by } g(x) > 0 \text{ hypothesis} \\
 &\triangleq \mathbb{1}_{[0:\infty)}[f(x)] && \text{by definition of } \mathbb{1}_A(x) && (\text{Definition ?? page ??}) \\
 &\triangleq \sigma[f(x)] && \text{by definition of } \sigma(x) && (\text{Definition Q.1 page 343})
 \end{aligned}$$

Definition Q.2.¹ Let $\mathbb{1}$ be the SET INDICATOR function (Definition ?? page ??). Let $f(x) \star g(x)$ represent the CONVOLUTION operation (Definition N.3 page 312).

D E F The **n th order cardinal B-spline** $N_n(x)$ for $n \in \mathbb{W}$ is defined as

$$N_n(x) \triangleq \begin{cases} \mathbb{1}_{[0:1)}(x) & \text{for } n = 0 \\ N_{n-1}(x) \star N_0(x) & \text{for } n \in \mathbb{W} \setminus 0 \end{cases} \quad \forall x \in \mathbb{R}$$

Lemma Q.2.²

¹ Chui (1992) page 85 ((4.2.1)), Christensen (2008) page 140, Chui (1988) page 1

² Christensen (2008) page 140, Chui (1992) page 85 ((4.2.1)), Chui (1988) page 1, Prasad and Iyengar (1997) page 145

L E M $N_n(x) = \int_{\tau=0}^{\tau=1} N_{n-1}(x - \tau) d\tau \quad \forall n \in \{1, 2, 3, \dots\}$

PROOF:

$$\begin{aligned}
 N_n(x) &\triangleq N_{n-1}(x) \star N_0(x) && \text{by definition of } N_n(x) && (\text{Definition Q.2 page 343}) \\
 &\triangleq \int_{\mathbb{R}} N_{n-1}(x - \tau) N_0(\tau) d\tau && \text{by definition of convolution operation } \star && (\text{Definition N.3 page 312}) \\
 &\triangleq \int_{\mathbb{R}} N_{n-1}(x - \tau) \mathbb{1}_{[0:1]}(\tau) d\tau && \text{by definition of } N_0(x) && (\text{Definition Q.2 page 343}) \\
 &= \int_{[0:1]} N_{n-1}(x - \tau) d\tau && \text{by definition of } \mathbb{1}_A(x) && (\text{Definition ?? page ??}) \\
 &= \int_{[0:1]} N_{n-1}(x - \tau) d\tau \\
 &\triangleq \int_0^1 N_{n-1}(x - \tau) d\tau
 \end{aligned}$$



Lemma Q.3. Let $f(x)$ be a FUNCTION in $\mathbb{R}^{\mathbb{R}}$. Let $F(x)$ be the ANTI-DERIVATIVE of $f(x)$.

Let $\sigma(x)$ be the STEP FUNCTION (Definition Q.1 page 343).

L E M

$$\begin{aligned}
 &\int_{y=a}^{y=b} f(x - y) \sigma(x - y) dy \\
 &= \left\{ \begin{array}{ll} - \int_{y=x-a}^{y=x-b} f(y) dy & \text{for } x \geq b \\ - \int_{y=x-a}^{y=0} f(y) dy & \text{for } a \leq x \leq b \\ 0 & \text{for } x \leq a \end{array} \right\} = \left\{ \begin{array}{ll} F(x - a) - F(x - b) & \text{for } x \geq b \\ F(x - a) - F(0) & \text{for } a \leq x \leq b \\ 0 & \text{for } x \leq a \end{array} \right\} \\
 &= [F(x - a) - F(0)]\sigma(x - a) + [F(0) - F(x - b)]\sigma(x - b)
 \end{aligned}$$

PROOF:

$$\begin{aligned}
 \int_{y=a}^{y=b} f(x - y) \sigma(x - y) dy &= \left\{ \begin{array}{ll} \int_{y=a}^{y=b} f(x - y) dy & \text{for } x \geq b \\ \int_{y=a}^{y=x} f(x - y) dy & \text{for } a \leq x \leq b \\ 0 & \text{for } x \leq a \end{array} \right\} && \text{by definition of } \sigma \text{ (Definition Q.1 page 343)} \\
 &= \left\{ \begin{array}{ll} - \int_{u=x-a}^{u=x-b} f(u) du & \text{for } x \geq b \\ - \int_{u=x-a}^{u=0} f(u) du & \text{for } a \leq x \leq b \\ 0 & \text{for } x \leq a \end{array} \right\} && \text{where } u \triangleq x - y \implies y = x - u \\
 &= \left\{ \begin{array}{ll} - \int_{y=x-a}^{y=x-b} f(y) dy & \text{for } x \geq b \\ - \int_{y=x-a}^{y=0} f(y) dy & \text{for } a \leq x \leq b \\ 0 & \text{for } x \leq a \end{array} \right\} && \text{by change of dummy variable } (u \rightarrow y) \\
 &= \left\{ \begin{array}{ll} F(x - a) - F(x - b) & \text{for } x \geq b \\ F(x - a) - F(0) & \text{for } a \leq x \leq b \\ 0 & \text{for } x \leq a \end{array} \right\} && \text{by Fundamental Theorem of Calculus} \\
 &= [F(x - a) - F(x - b)]\sigma(x - b) + [F(x - a) - F(0)][\sigma(x - a) - \sigma(x - b)] \\
 &= [F(x - a) - F(0)]\sigma(x - a) + [F(x - a) - F(x - b) - F(x - a) + F(0)]\sigma(x - b) \\
 &= [F(x - a) - F(0)]\sigma(x - a) + [F(0) - F(x - b)]\sigma(x - b)
 \end{aligned}$$






Figure Q.1: some low order B-splines (Example Q.1 page 345)

Lemma Q.4. Let $\sigma(x)$ be the STEP FUNCTION (Definition Q.1 page 343).

$$\text{LEM} \quad \int_{\tau=0}^{\tau=1} (x - \tau - k)^n \sigma(x - \tau - k) d\tau = \frac{1}{n+1} [(x - k)^{n+1} \sigma(x - k) - (x - k - 1)^{n+1} \sigma(x - k - 1)]$$

PROOF:

$$\begin{aligned}
 & \int_{\tau=0}^{\tau=1} (x - \tau - k)^n \sigma(x - \tau - k) d\tau \\
 &= \int_{y=k}^{y=k+1} (x - y)^n \sigma(x - y) dy && \text{where } y \triangleq \tau + k \implies \tau = y - k \\
 &= [F(x - k) - F(0)] \sigma(x - k) + [F(0) - F(x - k - 1)] \sigma(x - k - 1) && \text{by Lemma Q.3 (page 344), where } f(x) \triangleq x^n \\
 &= \frac{[(x - k)^{n+1} - 0] \sigma(x - k) + [0 - (x - k - 1)^{n+1}] \sigma(x - k - 1)}{n+1} && \text{because } F(x) \triangleq \int f(x) dx = \frac{x^{n+1}}{n+1} + c \\
 &= \frac{1}{n+1} [(x - k)^{n+1} \sigma(x - k) - (x - k - 1)^{n+1} \sigma(x - k - 1)]
 \end{aligned}$$

*Example Q.1.*³ Let $\sigma(x)$ be the step function (Definition Q.1 page 343). Let $\binom{n}{k}$ be the binomial coefficient (Definition ?? page ??). The 0th order B-spline (Definition Q.2 page 343) $N_0(x)$ can be expressed as follows:

$$\text{EX} \quad N_0(x) = \left\{ \begin{array}{ll} 1 & \text{for } x \in [0 : 1) \\ 0 & \text{otherwise} \end{array} \right\} = \left\{ \sum_{k=0}^1 (-1)^k \binom{1}{k} (x - k)^0 \sigma(x - k) \quad \forall x \in \mathbb{R} \right\}$$

The B-spline $N_0(x)$ is illustrated in Figure Q.1 (page 345).

³ Schumaker (2007) page 136 (Table 1)

PROOF:

$$\begin{aligned}
 N_0(x) &= \mathbb{1}_{[0:1]}(x) && \text{by definition of } N_0(x) \\
 &= \sigma(x) - \sigma(x-1) && \text{by definition of } \sigma(x) \\
 &= \left[\binom{1}{0} \sigma(x) - \binom{1}{1} \sigma(x-1) \right] && \text{by definition of binomial coefficient } \binom{n}{k} \\
 &= \sum_{k=0}^1 (-1)^k \binom{1}{k} (x-k)^0 \sigma(x-k) && \text{by definition of } \sum \text{ operator}
 \end{aligned}$$

Example Q.2.⁴ Let $\sigma(x)$ be the step function. Let $\binom{n}{k}$ be the binomial coefficient.

The 1st order B-spline $N_1(x)$ can be expressed as follows:

E X

$$N_1(x) = \begin{cases} x & \text{for } x \in [0 : 1] \\ -x+2 & \text{for } x \in [1 : 2] \\ 0 & \text{otherwise} \end{cases} = \left\{ \sum_{k=0}^2 (-1)^k \binom{2}{k} (x-k) \sigma(x-k) \quad \forall x \in \mathbb{R} \right\}$$

The B-spline $N_1(x)$ is illustrated in Figure Q.1 (page 345).

PROOF:

$$\begin{aligned}
 N_1(x) &= \int_{\tau=0}^{\tau=1} N_0(x-\tau) d\tau && \text{by Lemma Q.2 page 343} \\
 &= \int_{\tau=0}^{\tau=1} \sum_{k=0}^1 (-1)^k \binom{1}{k} (x-\tau-k)^0 \sigma(x-\tau-k) d\tau && \text{by Example Q.1 page 345} \\
 &= \sum_{k=0}^1 (-1)^k \binom{1}{k} \int_{\tau=0}^{\tau=1} (x-\tau-k)^0 \sigma(x-\tau-k) d\tau && \text{by linearity of } \int d\tau \text{ operator} \\
 &= \sum_{k=0}^1 (-1)^k \binom{1}{k} \frac{1}{0+1} [(x-k)^{0+1} \sigma(x-k) - (x-k-1)^{0+1} \sigma(x-k-1)] && \text{by Lemma Q.4 page 345} \\
 &= \begin{pmatrix} 1\{(x-0)\sigma(x-0) - (x-1)\sigma(x-1)\} \\ -1\{(x-1)\sigma(x-1) - (x-2)\sigma(x-2)\} \end{pmatrix} \\
 &= x\sigma(x) - 2(x-1)\sigma(x-1) + (x-2)\sigma(x-2) \\
 &= \sum_{k=0}^2 (-1)^k \binom{2}{k} (x-k) \sigma(x-k) && \text{by def. of } \binom{n}{k} \text{ (Definition ?? page ??)} \\
 &= \begin{cases} x & \text{for } x \in [0 : 1] \\ -x+2 & \text{for } x \in [1 : 2] \\ 0 & \text{otherwise} \end{cases} && \text{by def. of } \sigma(x) \text{ (Definition Q.1 page 343)}
 \end{aligned}$$

Example Q.3.⁵ Let $\sigma(x)$ be the step function. Let $\binom{n}{k}$ be the binomial coefficient.

The 2nd order B-spline $N_2(x)$ can be expressed as follows:

E X

$$N_2(x) = \frac{1}{2} \begin{cases} x^2 & \text{for } x \in [0 : 1] \\ -2x^2 + 6x - 3 & \text{for } x \in [1 : 2] \\ x^2 - 6x + 9 & \text{for } x \in [2 : 3] \\ 0 & \text{otherwise} \end{cases} = \left\{ \frac{1}{2} \sum_{k=0}^3 (-1)^k \binom{3}{k} (x-k)^2 \sigma(x-k) \quad \forall x \in \mathbb{R} \right\}$$

The B-spline $N_2(x)$ is illustrated in Figure Q.1 (page 345).

⁴ Christensen (2008) page 148 (Exercise 6.2), Christensen (2010) page 212 (Exercise 10.2), Heil (2011) pages 142–143 (Definition 4.22 (The Schauder System)), Schumaker (2007) page 136 (Table 1), Stoer and Bulirsch (2002) page 124

⁵ Christensen (2008) page 148 (Exercise 6.2), Christensen (2010) page 212 (Exercise 10.2), Schumaker (2007) page 136 (Table 1), Stoer and Bulirsch (2002) page 124

PROOF:

$$\begin{aligned}
 N_2(x) &= \int_{\tau=0}^{\tau=1} N_1(x - \tau) d\tau && \text{by Lemma Q.2 page 343} \\
 &= \int_{\tau=0}^{\tau=1} \sum_{k=0}^2 (-1)^k \binom{2}{k} (x - \tau - k) \sigma(x - \tau - k) d\tau && \text{by Example Q.2 page 346} \\
 &= \sum_{k=0}^2 (-1)^k \binom{2}{k} \int_{\tau=0}^{\tau=1} (x - \tau - k) \sigma(x - \tau - k) d\tau && \text{by linearity of } \int d\tau \text{ operator} \\
 &= \sum_{k=0}^2 (-1)^k \binom{2}{k} \frac{1}{1+1} [(x - k)^{1+1} \sigma(x - k) - (x - k - 1)^{1+1} \sigma(x - k - 1)] && \text{by Lemma Q.4 page 345} \\
 &= \frac{1}{2} \left(\begin{array}{c} 1 \quad \{(x-0)^2 \sigma(x-0) - (x-1)^2 \sigma(x-1)\} \\ -2 \quad \{(x-1)^2 \sigma(x-1) - (x-2)^2 \sigma(x-2)\} \\ +1 \quad \{(x-2)^2 \sigma(x-2) - (x-3)^2 \sigma(x-3)\} \end{array} \right) \\
 &= \frac{1}{2} [x^2 \sigma(x) - 3(x-1)^2 \sigma(x-1) + 3(x-2)^2 \sigma(x-2) - (x-3)^2 \sigma(x-3)] \\
 &= \frac{1}{2} \sum_{k=0}^3 (-1)^k \binom{3}{k} (x-k)^2 \sigma(x-k) && \text{by def. of } \binom{n}{k} \text{ (Definition ?? page ??)} \\
 &= \frac{1}{2} \left\{ \begin{array}{ll} x^2 & \text{for } x \in [0 : 1] \\ -2x^2 + 6x - 3 & \text{for } x \in [1 : 2] \\ x^2 - 6x + 9 & \text{for } x \in [2 : 3] \\ 0 & \text{otherwise} \end{array} \right\} && \text{by def. of } \sigma(x) \text{ (Definition Q.1 page 343)}
 \end{aligned}$$

The final steps of this proof can be calculated “by hand” or by using the free and open source software package *Maxima* along with the script file listed in Section ?? (page ??). \Rightarrow

Q.2 Algebraic properties

Theorem Q.1 (next) presents a closed form expression for an *n*th order B-spline $N_n(x)$ based on the definition of $N_n(x)$ given in Definition Q.2 (page 343). Alternatively, Theorem Q.1 could serve as the definition and Definition Q.2 as a property.

Theorem Q.1.⁶ Let $N_n(x)$ be the *n*th ORDER B-SPLINE (Definition Q.2 page 343). Let $\sigma(x)$ be the STEP FUNCTION (Definition Q.1 page 343).

T H M	$N_n(x) = \frac{1}{n!} \sum_{k=0}^{n+1} (-1)^k \binom{n+1}{k} (x-k)^n \sigma(x-k) \quad \forall n \in \{0, 1, 2, \dots\} = \mathbb{W}$
-------------	--

PROOF: Proof follows by induction:

1. base case (choose one):
 - Proof for $n = 0$ case: by Example Q.1 (page 345).
 - Proof for $n = 1$ case: by Example Q.2 (page 346).
 - Proof for $n = 2$ case: by Example Q.3 (page 346).

⁶ Christensen (2008) page 142 (Theorem 6.1.3), Chui (1992) page 84 ((4.1.12))

2. inductive step—proof that n case $\implies n + 1$ case:

$$\begin{aligned}
 N_{n+1}(x) &= \int_0^1 N_n(x - \tau) d\tau && \text{by Lemma Q.2 page 343} \\
 &= \int_0^1 \frac{1}{n!} \sum_{k=0}^{k=n+1} (-1)^k \binom{n+1}{k} (x - \tau - k)^n \sigma(x - \tau - k) d\tau && \text{by induction hypothesis} \\
 &= \frac{1}{n!} \sum_{k=0}^{k=n+1} (-1)^k \binom{n+1}{k} \int_0^1 (x - \tau - k)^n \sigma(x - \tau - k) d\tau && \text{by linearity of } \int d\tau \text{ operator} \\
 &= \frac{1}{n!} \sum_{k=0}^{k=n+1} (-1)^k \binom{n+1}{k} \frac{1}{n+1} [(x - k)^{n+1} \sigma(x - k) - (x - k - 1)^{n+1} \sigma(x - k - 1)] && \text{by Lemma Q.4 page 345} \\
 &= \frac{1}{(n+1)!} \sum_{k=0}^{k=n+1} (-1)^k \binom{n+1}{k} [(x - k)^{n+1} \sigma(x - k) - (x - k - 1)^{n+1} \sigma(x - k - 1)] \\
 &= \frac{1}{(n+1)!} \left[\sum_{k=0}^{k=n+1} (-1)^k \binom{n+1}{k} (x - k)^{n+1} \sigma(x - k) - \sum_{k=0}^{k=n+1} (-1)^k \binom{n+1}{k} (x - k - 1)^{n+1} \sigma(x - k - 1) \right] \\
 &= \frac{1}{(n+1)!} \left[\sum_{k=0}^{k=n+1} (-1)^k \binom{n+1}{k} (x - k)^{n+1} \sigma(x - k) - \sum_{m=1}^{m=n+2} (-1)^{m-1} \binom{n+1}{m-1} (x - m)^{n+1} \sigma(x - m) \right]
 \end{aligned}$$


where $m \triangleq k + 1 \implies k = m - 1$

$$\begin{aligned}
 &= \frac{1}{(n+1)!} \left(\sum_{k=0}^{k=n+1} (-1)^k \binom{n+1}{k} (x - k)^{n+1} \sigma(x - k) - \sum_{m=1}^{m=n+2} (-1)^{m-1} \left[\binom{n+2}{m} - \binom{n+1}{m} \right] (x - m)^{n+1} \sigma(x - m) \right) && \text{by Pascal's identity / Stifel formula (Theorem ?? page ??)} \\
 &= \frac{1}{(n+1)!} \left(\sum_{m=1}^{m=n+2} (-1)^m \binom{n+2}{m} (x - m)^{n+1} \sigma(x - m) - \sum_{m=1}^{m=n+2} (-1)^m \binom{n+1}{m} (x - m)^{n+1} \sigma(x - m) + \sum_{k=0}^{k=n+1} (-1)^k \binom{n+1}{k} (x - k)^{n+1} \sigma(x - k) \right) && \text{note } (-1)^{m-1} = -(-1)^m \\
 &= \frac{1}{(n+1)!} \left(\sum_{m=0}^{m=n+2} (-1)^m \binom{n+2}{m} (x - m)^{n+1} \sigma(x - m) - (-1)^0 \binom{n+2}{0} (x - 0)^{n+1} \sigma(x - 0) - \sum_{m=1}^{m=n+1} (-1)^m \binom{n+1}{m} (x - m)^{n+1} \sigma(x - m) - (-1)^{n+2} \binom{n+1}{n+2} (x - n - 2)^{n+1} \sigma(x - n - 2) + \sum_{k=1}^{k=n+1} (-1)^k \binom{n+1}{k} (x - k)^{n+1} \sigma(x - k) + (-1)^0 \binom{n+1}{0} (x - 0)^{n+1} \sigma(x - 0) \right) && \begin{array}{ll} \text{(A)} & \text{desired } n + 1 \text{ case} \\ \text{(B)} & \text{cancelled by (F)} \\ \text{(C)} & \text{cancelled by (E)} \\ \text{(D)} & \binom{n+1}{n+2} = 0 \text{ by Proposition ?? page ??} \\ \text{(E)} & \text{cancelled by (C)} \\ \text{(F)} & \binom{n+2}{0} = \binom{n+1}{0} = 1, \text{ so (F) is cancelled by (B)} \end{array} \\
 &= \frac{1}{(n+1)!} \sum_{m=0}^{m=n+2} (-1)^m \binom{n+2}{m} (x - m)^{n+1} \sigma(x - m) && (n + 1 \text{ case})
 \end{aligned}$$

*Example Q.4.*⁷ Let $N_3(x)$ be the 3rd order B-spline (Definition Q.2 page 343).⁸

E X

$$N_3(x) = \frac{1}{6} \begin{cases} x^3 & \text{for } 0 \leq x \leq 1 \\ -3x^3 + 12x^2 - 12x + 4 & \text{for } 1 \leq x \leq 2 \\ 3x^3 - 24x^2 + 60x - 44 & \text{for } 2 \leq x \leq 3 \\ -x^3 + 12x^2 - 48x + 64 & \text{for } 3 \leq x \leq 4 \\ 0 & \text{otherwise} \end{cases}$$



PROOF: This expression can be calculated “by hand” using Theorem Q.1 (page 347) or by using the free and open source software package *Maxima* along with the script file listed in Section ?? (page ??). \Rightarrow

Example Q.5. Let $N_4(x)$ be the 4th order B-spline (Definition Q.2 page 343).

E X

$$N_4(x) = \frac{1}{24} \begin{cases} x^4 & \text{for } 0 \leq x \leq 1 \\ -4x^4 + 20x^3 - 30x^2 + 20x - 5 & \text{for } 1 \leq x \leq 2 \\ 6x^4 - 60x^3 + 210x^2 - 300x + 155 & \text{for } 2 \leq x \leq 3 \\ -4x^4 + 60x^3 - 330x^2 + 780x - 655 & \text{for } 3 \leq x \leq 4 \\ x^4 - 20x^3 + 150x^2 - 500x + 625 & \text{for } 4 \leq x \leq 5 \\ 0 & \text{otherwise} \end{cases}$$

PROOF: This expression can be calculated “by hand” using Theorem Q.1 (page 347) or by using the free and open source software package *Maxima* along with the script file listed in Section ?? (page ??). \Rightarrow

Example Q.6. Let $N_5(x)$ be the 5th order B-spline (Definition Q.2 page 343).

E X

$$N_5(x) = \frac{1}{120} \begin{cases} x^5 & \text{for } 0 \leq x \leq 1 \\ -5x^5 + 30x^4 - 60x^3 + 60x^2 - 30x + 6 & \text{for } 1 \leq x \leq 2 \\ 10x^5 - 120x^4 + 540x^3 - 1140x^2 + 1170x - 474 & \text{for } 2 \leq x \leq 3 \\ -10x^5 + 180x^4 - 1260x^3 + 4260x^2 - 6930x + 4386 & \text{for } 3 \leq x \leq 4 \\ 5x^5 - 120x^4 + 1140x^3 - 5340x^2 + 12270x - 10974 & \text{for } 4 \leq x \leq 5 \\ -x^5 + 30x^4 - 360x^3 + 2160x^2 - 6480x + 7776 & \text{for } 5 \leq x \leq 6 \\ 0 & \text{otherwise} \end{cases}$$

The 5th order B-spline $N_5(x)$ is illustrated in Figure Q.1 (page 345).

PROOF: This expression can be calculated “by hand” using Theorem Q.1 (page 347) or by using the free and open source software package *Maxima* along with the script file listed in Section ?? (page ??). \Rightarrow

Example Q.7. Let $N_6(x)$ be the 6th order B-spline (Definition Q.2 page 343).

E X

$$N_6(x) = \frac{1}{720} \begin{cases} x^6 & \text{for } 0 \leq x \leq 1 \\ -6x^6 + 42x^5 - 105x^4 + 140x^3 - 105x^2 + 42x - 7 & \text{for } 1 \leq x \leq 2 \\ 15x^6 - 210x^5 + 1155x^4 - 3220x^3 + 4935x^2 - 3990x + 1337 & \text{for } 2 \leq x \leq 3 \\ -20x^6 + 420x^5 - 3570x^4 + 15680x^3 - 37590x^2 + 47040x - 24178 & \text{for } 3 \leq x \leq 4 \\ 15x^6 - 420x^5 + 4830x^4 - 29120x^3 + 96810x^2 - 168000x + 119182 & \text{for } 4 \leq x \leq 5 \\ -6x^6 + 210x^5 - 3045x^4 + 23380x^3 - 100065x^2 + 225750x - 208943 & \text{for } 5 \leq x \leq 6 \\ x^6 - 42x^5 + 735x^4 - 6860x^3 + 36015x^2 - 100842x + 117649 & \text{for } 6 \leq x \leq 7 \\ 0 & \text{otherwise} \end{cases}$$

The 6th order B-spline $N_6(x)$ is illustrated in Figure Q.1 (page 345).

PROOF: This expression can be calculated “by hand” using Theorem Q.1 (page 347) or by using the free and open source software package *Maxima* along with the script file listed in Section ?? (page ??). \Rightarrow

⁷ Schumaker (2007) page 136 (Table 1), Shizgal (2015) page 92 ((2.199)), Szabó and Horváth (2004) page 146 ((4)), Wei and Billings (2006) page 578 (Table 1), Maleknejad et al. (2013) ((9))

⁸ For help with plotting B-splines, see APPENDIX R (page 375).

Example Q.8. Let $N_7(x)$ be the 7th order B-spline (Definition Q.2 page 343).

$$\begin{aligned} \text{E} \\ \text{x} \end{aligned} \quad \begin{aligned} 7!N_7(x) &= 5040N_7(x) = \\ &\left\{ \begin{array}{ll} x^7 & \text{for } 0 \leq x \leq 1 \\ -7x^7 + 56x^6 - 168x^5 + 280x^4 - 280x^3 + 168x^2 - 56x + 8 & \text{for } 1 \leq x \leq 2 \\ 21x^7 - 336x^6 + 2184x^5 - 7560x^4 + 15400x^3 - 18648x^2 + 12488x - 3576 & \text{for } 2 \leq x \leq 3 \\ -35x^7 + 840x^6 - 8400x^5 + 45360x^4 - 143360x^3 + 267120x^2 - 273280x + 118896 & \text{for } 3 \leq x \leq 4 \\ 35x^7 - 1120x^6 + 15120x^5 - 111440x^4 + 483840x^3 - 1238160x^2 + 1733760x - 1027984 & \text{for } 4 \leq x \leq 5 \\ -21x^7 + 840x^6 - 14280x^5 + 133560x^4 - 741160x^3 + 2436840x^2 - 4391240x + 3347016 & \text{for } 5 \leq x \leq 6 \\ 7x^7 - 336x^6 + 6888x^5 - 78120x^4 + 528920x^3 - 2135448x^2 + 4753336x - 4491192 & \text{for } 6 \leq x \leq 7 \\ -x^7 + 56x^6 - 1344x^5 + 17920x^4 - 143360x^3 + 688128x^2 - 1835008x + 2097152 & \text{for } 7 \leq x \leq 8 \\ 0 & \text{otherwise} \end{array} \right\} \end{aligned}$$

The 7th order B-spline $N_7(x)$ is illustrated in Figure Q.1 (page 345).

PROOF: This expression can be calculated “by hand” using Theorem Q.1 (page 347) or by using the free and open source software package *Maxima* along with the script file listed in Section ?? (page ??). \Rightarrow

Example Q.9. ⁹ The $(n+1)^2$ coefficients of the order $n, n-1, \dots, 0$ monomials of each B-spline $N_n(x)$ multiplied by $n!$ induce an *integer sequence*

$x \triangleq (1, 1, 0, -1, 2, 1, 0, 0, -2, 6, -3, 1, -6, 9, 1, 0, 0, 0, -3, 12, -12, 4, 3, -24, 60, -44, -1, 12, -48, 64, \dots)$ as more fully listed in Table Q.1 (page 374). In this sequence $x \triangleq (x_0, x_1, x_2, \dots)$, the coefficients for the *order n* B-spline $N_n(x)$ begin at the sequence index value

$$p \triangleq \sum_{k=0}^n k^2 = \frac{1}{6}n(n+1)(2n+1) \quad \text{and end at index value } p + (n+1)^2 - 1.$$

For example, the coefficients for $N_3(x)$ begin at index value $p \triangleq 0 + 1 + 4 + 9 = 14$ and end at index value $p + 4^2 - 1 = 29$. Using these coefficients gives the following expression for $N_3(x)$:

$$N_3(x) = \left[\begin{array}{cccc} 1 & 0 & 0 & 0 \\ -3 & 12 & -12 & 4 \\ 3 & -24 & 60 & -44 \\ -1 & 12 & -48 & 64 \end{array} \right] \left[\begin{array}{c} x^3 \\ x^2 \\ x \\ 1 \end{array} \right] = \left\{ \begin{array}{ll} x^3 & \text{for } 0 \leq x < 1 \\ -3x^3 + 12x^2 - 12x + 4 & \text{for } 1 \leq x < 2 \\ 3x^3 - 24x^2 + 60x - 44 & \text{for } 2 \leq x < 3 \\ -x^3 + 12x^2 - 48x + 64 & \text{for } 3 \leq x < 4 \\ 0 & \text{otherwise} \end{array} \right\}$$

...which agrees with the result presented in Example Q.4 (page 349).

PROOF:

1. The coefficients for the sequence x may be computed with assistance from *Maxima* together with the script file listed in Section ?? (page ??).
2. Proof that $\sum_{k=0}^n k^2 = \frac{1}{6}n(n+1)(2n+1)$: The summation is a *power sum*. The relation may be proved using *induction*.¹⁰
 - (a) Base case: $n=0$ case ...

$$\begin{aligned} \sum_{k=0}^{n=0} k^2 &= 0 \\ &= \frac{0(0+1)(2 \cdot 0 + 1)}{6} \\ &= \frac{n(n+1)(2n+1)}{6} \Big|_{n=0} \end{aligned}$$

⁹ Greenhoe (2017b)

¹⁰ Greenhoe (2017a), pages 186–187 (Proposition 11.2 (Power Sums))

(b) Base case: $n=1$ case ...

$$\begin{aligned}\sum_{k=0}^{k=1} k^2 &= 0 + 1 \\ &= \frac{1(1+1)(2 \cdot 1 + 1)}{6} \\ &= \frac{n(n+1)(2n+1)}{6} \Big|_{n=1}\end{aligned}$$

(c) inductive step—proof that n case $\implies n+1$ case:

$$\begin{aligned}\sum_{k=0}^{n+1} k^2 &= \left(\sum_{k=0}^n k^2 \right) + (n+1)^2 \\ &= \left(\frac{n(n+1)(2n+1)}{6} \right) + (n+1)^2 && \text{by } n \text{ case hypothesis} \\ &= (n+1) \left(\frac{n(2n+1) + 6(n+1)}{6} \right) \\ &= (n+1) \left(\frac{2n^2 + 7n + 6}{6} \right) \\ &= (n+1) \left(\frac{(n+2)(2n+3)}{6} \right) \\ &= \frac{(n+1)[(n+1)+1][2(n+1)+1]}{6}\end{aligned}$$



Theorem Q.2. ¹¹

T H M	$\frac{d}{dx} N_n(x) = N_{n-1}(x) - N_{n-1}(x-1) \quad \forall n \in \mathbb{N}, \forall x \in \mathbb{R}$
-------------	--

PROOF:

1. Proof using Lemma Q.2 (page 343) and the *Fundamental Theorem of Calculus*:

$$\begin{aligned}\frac{d}{dx} N_n(x) &= \frac{d}{dx} \int_0^1 N_{n-1}(x-\tau) d\tau && \text{by Lemma Q.2 page 343} \\ &= \frac{d}{dx} \int_{x-u=0}^{x-u=1} N_{n-1}(u)(-1) du && \text{where } u \triangleq x - \tau \implies \tau = x - u \\ &= \frac{d}{dx} \int_{u=x-1}^{u=x} N_{n-1}(u) du \\ &= \left\{ \frac{d}{dx} \int N_{n-1}(u) du \Big|_{u=x} \right\} - \left\{ \frac{d}{dx} \int N_{n-1}(u) du \Big|_{u=x-1} \right\} && \text{by Fundamental Theorem of Calculus}^{12} \\ &= \left\{ N_{n-1}(x) \frac{d}{dx}(x) \right\} - \left\{ N_{n-1}(x-1) \frac{d}{dx}(x-1) \right\} && \text{by Chain Rule}^{13} \\ &= N_{n-1}(x) - N_{n-1}(x-1)\end{aligned}$$

¹¹ Höllig (2003) page 25 (3.2), Schumaker (2007) page 121 (Theorem 4.16)

¹² Hijab (2011) page 163 (Theorem 4.4.3)

¹³ Hijab (2011) pages 73–74 (Theorem 3.1.2)

2. Proof using Lemma Q.2 (page 343) and induction:

(a) Base case ...proof for $n = 1$ case:

$$\begin{aligned}
 N_0(x) - N_0(x-1) &= \underbrace{\sigma(x) - \sigma(x-1)}_{N_0(x)} - \underbrace{[\sigma(x-1) - \sigma(x-2)]}_{N_0(x-1)} \quad \text{by Example Q.1 page 345} \\
 &= \sigma(x) - 2\sigma(x-1) + \sigma(x-2) \\
 &= \sum_{k=0}^2 (-1)^k \binom{2}{k} \sigma(x-k) \\
 &= \frac{d}{dx} \sum_{k=0}^2 (-1)^k \binom{2}{k} (x-k) \sigma(x-k) \\
 &= \frac{d}{dx} N_1(x) \quad \text{by Example Q.2 page 346}
 \end{aligned}$$

(b) Base case ...proof for $n = 2$ case:

$$\begin{aligned}
 N_1(x) - N_1(x-1) &= \underbrace{x\sigma(x) - 2(x-1)\sigma(x-1) + (x-2)\sigma(x-2)}_{N_1(x)} \\
 &\quad - \underbrace{[(x-1)\sigma(x-1) - 2(x-2)\sigma(x-2) + (x-3)\sigma(x-3)]}_{N_1(x-1)} \quad \text{by Example Q.2 page 346} \\
 &= x\sigma(x) + [-2x + 2 - x + 1]\sigma(x-1) + [x - 2 + 2x - 4]\sigma(x-2) + [-x + 3]\sigma(x-3) \\
 &= x\sigma(x) + [-3x + 3]\sigma(x-1) + [3x - 6]\sigma(x-2) + [-x + 3]\sigma(x-3) \\
 &= \frac{d}{dx} \left\{ \begin{array}{l} \frac{1}{2}x^2\sigma(x) + \left[-\frac{3}{2}x^2 + 3x - \frac{1}{2} \right] \sigma(x-1) + \left[\frac{3}{2}x^2 - 6x + 3 \right] \sigma(x-2) \\ \quad + \left[-\frac{1}{2}x^2 + 3x - \frac{5}{2} \right] \sigma(x-3) \end{array} \right\} \\
 &= \frac{d}{dx} N_2(x) \quad \text{by Example Q.3 page 346}
 \end{aligned}$$

(c) Proof that n case $\implies n+1$ case:

$$\begin{aligned}
 \frac{d}{dx} N_{n+1}(x) &= \frac{d}{dx} \int_0^1 N_n(x-\tau) d\tau \quad \text{by Lemma Q.2 page 343} \\
 &= \int_0^1 \frac{d}{d\tau} N_n(x-\tau) d\tau \quad \text{by Leibniz Integration Rule (Theorem ?? page ??)} \\
 &= \int_0^1 [N_{n-1}(x-\tau) - N_{n-1}(x-1-\tau)] d\tau \quad \text{by left hypothesis} \\
 &= \int_0^1 N_{n-1}(x-\tau) d\tau - \int_0^1 N_{n-1}(x-1-\tau) d\tau \\
 &= N_n(x) - N_n(x-1) \quad \text{by Lemma Q.2 page 343}
 \end{aligned}$$

Theorem Q.3 (B-spline recursion). ¹⁴ Let $N_n(x)$ be the n TH ORDER B-SPLINE (Definition Q.2 page 343).

T H M	$N_n(x) = \frac{x}{n} N_{n-1}(x) + \frac{n+1-x}{n} N_{n-1}(x-1) \quad \forall n \in \{1, 2, 3, \dots\}, \forall x \in \mathbb{R}$
-------------	---

¹⁴ Christensen (2008) page 140 (Theorem 6.1.1), Höllig (2003) page 27 (3.4), Schumaker (2007) page 120 (Theorem 4.15), de Boor (2001) page 90 (B-Spline Property (i)), Chui (1988) page 2 (Theorem 1.1), Wojtaszczyk (1997) page 53 (Theorem 3.7), Cox (1972), de Boor (1972)

PROOF:

1. Base case ...proof for $n = 1$ case:

$$\begin{aligned} \frac{x}{1} N_0(x) + \frac{1+1-x}{1} N_0(x-1) &= \underbrace{\frac{x}{1} [\sigma(x) - \sigma(x-1)]}_{N_0(x)} + \underbrace{\frac{1+1-x}{1} [\sigma(x-1) - \sigma(x-2)]}_{N_0(x-1)} \\ &= x\sigma(x) + [-x - x + 2]\sigma(x-1) + [x - 2]\sigma(x-2) \\ &= N_1(x) \quad \text{by Example Q.2 page 346} \end{aligned}$$

2. Induction step ...proof that n case $\Rightarrow n+1$ case:

$$\begin{aligned} &\frac{x}{n+1} N_n(x) + \frac{n+2-x}{n+1} N_n(x-1) + c_1 \\ &= \int \frac{d}{dx} \left\{ \frac{x}{n+1} N_n(x) + \frac{n+2-x}{n+1} N_n(x-1) \right\} dx \\ &= \int \underbrace{\frac{1}{n+1} N_n(x) + \frac{x}{n+1} \frac{d}{dx} N_n(x)}_{\frac{d}{dx} \frac{x}{n+1} N_n(x)} + \underbrace{\frac{-1}{n+1} N_n(x-1) + \frac{n+2-x}{n} \frac{d}{dx} N_n(x-1)}_{\frac{d}{dx} \frac{n+2-x}{n+1} N_n(x-1)} dx \\ &\quad \text{by product rule} \\ &= \int \frac{1}{n+1} \left[\underbrace{\frac{x}{n} N_{n-1}(x) + \frac{n+1-x}{n} N_{n-1}(x-1)}_{\text{by } n \text{ hypothesis}} \right] + \frac{x}{n+1} \underbrace{[N_{n-1}(x) - N_{n-1}(x-1)]}_{\text{by Theorem Q.2 page 351}} \\ &\quad - \underbrace{\left[\frac{x-1}{n^2+n} N_{n-1}(x-1) + \frac{n-x+2}{n(n+1)} N_{n-1}(x-2) \right]}_{\text{by induction hypothesis}} \\ &\quad + \frac{n+2-x}{n+1} \underbrace{[N_{n-1}(x-1) - N_{n-1}(x-2)]}_{\text{by Theorem Q.2 page 351}} dx \\ &= \int \left[\frac{x}{n(n+1)} + \frac{x}{n+1} \right] N_{n-1}(x) + \left[\frac{n-x+1}{n(n+1)} - \frac{x-1}{n(n+1)} + \frac{n+2-2x}{n+1} \right] N_{n-1}(x-1) \\ &\quad + \left[\frac{-n-2+x}{n(n+1)} + \frac{-n-2+x}{n+1} \right] N_{n-1}(x-2) dx \\ &= \int \left[\frac{x+nx}{n(n+1)} \right] N_{n-1}(x) + \left[\frac{n+2-2x+n(n+2-2x)}{n(n+1)} \right] N_{n-1}(x-1) \\ &\quad + \left[\frac{-n-2+x+n(-n-2+x)}{n(n+1)} \right] N_{n-1}(x-2) dx \\ &= \int \left[\frac{x}{n} \right] N_{n-1}(x) + \left[\frac{n+2-2x}{n} \right] N_{n-1}(x-1) + \left[\frac{-n-2+x}{n} \right] N_{n-1}(x-2) dx \\ &= \int \underbrace{\left[\frac{x}{n} \right] N_{n-1}(x)}_{N_n(x)} + \underbrace{\left[\frac{n+1-x}{n} \right] N_{n-1}(x-1)}_{N_{n-1}(x-1)} \\ &\quad - \underbrace{\left[\frac{x-1}{n} \right] N_{n-1}(x-1) - \left[\frac{n+2-x}{n} \right] N_{n-1}(x-2)}_{N_{n-1}(x-1)} dx \\ &= \int N_n(x) - N_n(x-1) dx \quad \text{by } n \text{ hypothesis} \\ &= \int \frac{d}{dx} N_{n+1}(x) dx \quad \text{by Theorem Q.2 page 351} \\ &= N_{n+1}(x) + c_2 \end{aligned}$$

Proof that $c_1 = c_2$: By item (2) (page 354), $N_n(x) = 0$ for $x < 0$. Therefore, $c_1 = c_2$.



Theorem Q.4 (B-spline general form). ¹⁵ Let $N_n(x)$ be the n TH ORDER B-SPLINE (Definition Q.2 page 343). Let $\text{supp } f$ be the SUPPORT of a function $f \in \mathbb{R}^{\mathbb{R}}$.

T
H
M

1. $N_n(x) \geq 0 \quad \forall n \in \mathbb{W}, \quad \forall x \in \mathbb{R}$ (NON-NEGATIVE)
2. $\text{supp } N_n(x) = [0 : n + 1] \quad \forall n \in \mathbb{W}$ (CLOSED SUPPORT)
3. $\int_{\mathbb{R}} N_n(x) dx = 1 \quad \forall n \in \mathbb{W}$ (UNIT AREA)
4. $N_n\left(\frac{n+1}{2} - x\right) = N_n\left(\frac{n+1}{2} + x\right) \quad \forall n \in \mathbb{N} \quad \forall x \in \mathbb{R}$ (SYMMETRIC about $x = \frac{n+1}{2}$)

PROOF:

1. Proof that $N_n(x) \geq 0$ (proof by induction):

(a) base case...proof that $N_0(x) \geq 0$:

$$\begin{aligned} N_0(x) &\triangleq \mathbb{1}_{[0:1]}(x) && \text{by definition of } N_0(x) && (\text{Definition Q.2 page 343}) \\ &\geq 0 && \text{by definition of } \mathbb{1}_A(x) && (\text{Definition ?? page ??}) \end{aligned}$$

(b) inductive step—proof that $\{N_n(x) \geq 0\} \implies \{N_{n+1}(x) \geq 0\}$:

$$\begin{aligned} N_{n+1}(x) &= \int_{\tau=0}^{\tau=1} N_n(x - \tau) d\tau && \text{by Lemma Q.2 page 343} \\ &\geq 0 && \text{by induction hypothesis } (N_n(x) \geq 0) \end{aligned}$$

2. Proof that $\text{supp } N_n(x) = [0 : n + 1]$ (proof by induction):

(a) Base case ...proof that $\text{supp } N_0 = [0 : 1]$:

$$\begin{aligned} \text{supp } N_0 &\triangleq \text{supp } \mathbb{1}_{[0:1]} && \text{by definition of } N_0(x) && (\text{Definition Q.2 page 343}) \\ &= \{[0 : 1]\}^- && \text{by definition of support operator} \\ &= [0 : 1] && \text{by definition of closure operator} \end{aligned}$$

(b) Induction step ...proof that $\{\text{supp } N_n = [0 : n + 1]\} \implies \{\text{supp } N_{n+1} = [0 : n + 2]\}$:

$$\begin{aligned} \text{supp } N_{n+1}(x) &= \text{supp } \int_{\tau=0}^{\tau=1} N_n(x - \tau) d\tau && \text{by Lemma Q.2 page 343} \\ &= \text{supp } \int_{[0:1]} N_n(x - \tau) d\tau && \text{by def. of Lebesgue integration} \\ &= \{x \in \mathbb{R} | (x - \tau) \in [0 : n + 1] \text{ for some } \tau \in [0 : 1]\}^- && \text{by induction hypothesis} \\ &= [0 : n + 1] \cup [0 + 1 : n + 1 + 1]^- \\ &= [0 : n + 2]^- \\ &= [0 : n + 2] && \text{by property of closure operator} \end{aligned}$$

3. Proof that $\int_{\mathbb{R}} N_n(x) dx = 1$ (proof by induction):

¹⁵ Christensen (2008) page 140 (Theorem 6.1.1), Höllig (2003) page 27 (3.4), Schumaker (2007) page 120 (Theorem 4.15), de Boor (2001) page 90 (B-Spline Property (i)), Chui (1988) page 2 (Theorem 1.1), Wojtaszczyk (1997) page 53 (Theorem 3.7), Cox (1972), de Boor (1972), Prasad and Iyengar (1997) pages 145–146 (Theorem 6.2)



(a) Base case ...proof that $\int_{\mathbb{R}} N_0(x) dx = 1$:

$$\begin{aligned} \int_{\mathbb{R}} N_0(x) dx &= \int_{\mathbb{R}} \mathbb{1}_{[0:1]} dx && \text{by definition of } N_0(x) && (\text{Definition Q.2 page 343}) \\ &= \int_{[0:1)} 1 dx && \text{by definition of } \mathbb{1}_A(x) && (\text{Definition ?? page ??}) \\ &= \int_{[0:1]} 1 dx && \text{by property of Lebesgue integration} \\ &= 1 \end{aligned}$$

(b) Induction step ...proof that $\{\int_{\mathbb{R}} N_n(x) dx = 1\} \implies \{\int_{\mathbb{R}} N_{n+1} dx = 1\}$:

$$\begin{aligned} \int_{\mathbb{R}} N_{n+1}(x) dx &= \int_{\mathbb{R}} \int_0^1 N_n(x - \tau) d\tau dx && \text{by Lemma Q.2 page 343} \\ &= \int_0^1 \int_{\mathbb{R}} N_n(x - \tau) dx d\tau \\ &= \int_0^1 \int_{\mathbb{R}} N_n(u) du d\tau && \text{where } u \triangleq x - \tau \implies \tau = x - u \\ &= \int_0^1 1 d\tau && \text{by induction hypothesis} \\ &= 1 \end{aligned}$$

4. Proof that $N_n(x)$ is *symmetric* for $n \in \{1, 2, 3, \dots\}$:

(a) Note that $N_0(x)$ ($n = 0$) is *not symmetric* (in particular it fails at $x = 1/2$) because

$$N_0\left(\frac{0+1}{2} - \frac{1}{2}\right) = N_0(0) = 1 \neq 0 = N_1(1) = N_0\left(\frac{0+1}{2} + \frac{1}{2}\right)$$

(b) Base case ...proof for $n = 1$ case:

$$\begin{aligned} N_1\left(\frac{1+1}{2} - x\right) &= N_1(1-x) \\ &= \begin{cases} (1-x) & \text{for } 1-x \in [0 : 1] \\ -(1-x)+2 & \text{for } 1-x \in [1 : 2] \\ 0 & \text{otherwise} \end{cases} && \text{by Example Q.2 page 346} \\ &= \begin{cases} -x+1 & \text{for } -x \in [-1 : 0] \\ x+1 & \text{for } -x \in [0 : 1] \\ 0 & \text{otherwise} \end{cases} \\ &= \begin{cases} x+1 & \text{for } x \in [-1 : 0] \\ -x+1 & \text{for } x \in [0 : 1] \\ 0 & \text{otherwise} \end{cases} \\ &= \begin{cases} (1+x) & \text{for } 1+x \in [0 : 1] \\ -(1+x)+2 & \text{for } 1+x \in [1 : 2] \\ 0 & \text{otherwise} \end{cases} \\ &= N_1(1+x) && \text{by Example Q.2 page 346} \\ &= N_1\left(\frac{1+1}{2} + x\right) \end{aligned}$$

(c) Induction step ...proof that $n - 1$ case $\implies n$ case:

$$\begin{aligned}
 & N_n\left(\frac{n+1}{2} + x\right) \\
 &= \frac{\frac{n+1}{2} + x}{n} N_{n-1}\left(\frac{n+1}{2} + x\right) + \frac{n+1 - \left(\frac{n+1}{2} + x\right)}{n} N_{n-1}\left(\frac{n+1}{2} + x - 1\right) \quad \text{by Theorem Q.3 page 352} \\
 &= \frac{n+1 - \left(\frac{n+1}{2} - x\right)}{n} N_{n-1}\left(\frac{n}{2} + \left[x + \frac{1}{2}\right]\right) + \frac{\frac{n+1}{2} - x}{n} N_{n-1}\left(\frac{n}{2} + \left[x - \frac{1}{2}\right]\right) \\
 &= \frac{n+1 - \left(\frac{n+1}{2} - x\right)}{n} N_{n-1}\left(\frac{n}{2} - \left[x + \frac{1}{2}\right]\right) + \frac{\frac{n+1}{2} - x}{n} N_{n-1}\left(\frac{n}{2} - \left[x - \frac{1}{2}\right]\right) \quad \text{by induction hypothesis} \\
 &= \frac{n+1 - \left(\frac{n+1}{2} - x\right)}{n} N_{n-1}\left(\left[\frac{n+1}{2} - x\right] - 1\right) + \frac{\frac{n+1}{2} - x}{n} N_{n-1}\left(\frac{n+1}{2} - x\right) \\
 &= N_n\left(\frac{n+1}{2} - x\right) \quad \text{by Theorem Q.3 page 352}
 \end{aligned}$$



Q.3 Projection properties

In the case where $(N_n(x - k))_{k \in \mathbb{Z}}$ is to be used as a basis in some subspace of $L^2_{\mathbb{R}}$, one may want to *project* a function $f(x)$ onto a basis function $N_n(x - k)$. This is especially true when $(N_n(x - k))$ is *orthogonal*; but in the case of *B-splines* this is only true when $n = 0$ (Theorem Q.8 page 366). Nevertheless, projection of a function onto $N_n(x - k)$, or the projection of $N_n(x)$ onto another basis function (such as the complex exponential in the case of *Fourier analysis* as in Lemma Q.5 page 358), is still useful. Projection in an *inner product space* is typically performed using the *inner product* $\langle f(x) | N_n(x - k) \rangle$; and in the space $L^2_{\mathbb{R}}$, this inner product is typically defined as an *integral* such that

$$\langle f(x) | N_n(x - k) \rangle \triangleq \int_{\mathbb{R}} f(x) N_n(x - k) dx.$$

As it turns out, there is a way to compute this inner product that only involves the function $f(x)$ and the order parameter n (next theorem).

Theorem Q.5. ¹⁶ Let f be a continuous function in $L^2_{\mathbb{R}}$ and $f^{(n)}$ the n th derivative of $f(x)$.

THEOREM

$$\begin{aligned}
 (1). \quad \int_{\mathbb{R}} f(x) N_n(x) dx &= \int_{[0:1]^{n+1}} f(x_1 + x_2 + \dots + x_{n+1}) dx_1 dx_2 \dots dx_{n+1} \\
 (2). \quad \int_{\mathbb{R}} f^{(n)}(x) N_n(x) dx &= \sum_{k=0}^{n+1} (-1)^{n-k+1} \binom{n+1}{k} f(k)
 \end{aligned}$$

PROOF:

1. Proof for (1) (proof by induction):

(a) Base case ...proof for $n = 0$ case:

$$\int_{\mathbb{R}} f(x) N_0(x) dx = \int_{[0:1]} f(x) dx \quad \text{by definition of } N_0(x) \quad (\text{Definition Q.2 page 343})$$

¹⁶ Chui (1992) page 85 ((4.2.2), (4.2.3)), Christensen (2008) page 140 (Theorem 6.1.1)

(b) Inductive step—proof that n case $\Rightarrow n + 1$ case:

$$\begin{aligned}
 & \int_{\mathbb{R}} f(x) N_{n+1}(x) dx \\
 &= \int_{\mathbb{R}} \left[\int_0^1 N_n(x - \tau) d\tau \right] f(x) dx && \text{by Lemma Q.2 page 343} \\
 &= \int_{[0:1]} \int_{\mathbb{R}} N_n(x - \tau) f(x) dx d\tau \\
 &= \int_{[0:1]} \int_{\mathbb{R}} N_n(u) f(u + \tau) du d\tau && \text{where } u \triangleq x - \tau \Rightarrow x = u + \tau \\
 &= \int_{[0:1]} \int_{[0:1]^{n+1}} f(u_1 + u_2 + \dots + u_{n+1} + \tau) du_1 du_2 \dots du_{n+1} d\tau && \text{by induction hypothesis} \\
 &= \int_{[0:1]^{n+2}} f(u_1 + u_2 + \dots + u_{n+1} + u_{n+2}) du_1 du_2 \dots du_{n+2} d\tau \\
 &= \int_{[0:1]^{n+2}} f(x_1 + x_2 + \dots + x_{n+1} + x_{n+2}) dx_1 dx_2 \dots dx_{n+2} && \text{by change of variables } u_k \rightarrow x_k
 \end{aligned}$$

2. Proof for (2):

$$\begin{aligned}
 \int_{\mathbb{R}} f^{(n)}(x) N_n(x) dx &= \int_{[0:1]^{n+1}} f^{(n)} \left(\sum_{k=1}^{n+1} x_k \right) dx_1 dx_2 \dots dx_{n+1} && \text{by (1)} \\
 &= \sum_{k=0}^{n+1} (-1)^{n-k+1} \binom{n+1}{k} f(k) && \text{by Theorem ?? page ??}
 \end{aligned}$$




Figure Q.2: Zero locations for B-cardinal spline $N_n(x)$ scaling coefficients

Q.4 Fourier analysis

Simply put, no matter what new and fancy basis sequences are discovered, the *Fourier transform* never goes out of style. This is largely because the *kernel* of the Fourier transform—the *complex exponential* function—has two properties that makes it extremely special:

- ➊ The complex exponential is an eigenvalue of any *linear time invariant* (LTI) operator (Theorem ?? page ??).
- ➋ The complex exponential generates a *continuous point spectrum* for the *differential operator*.

Thus, we might expect the projection of the *B-spline* function $N_n(x)$ onto the complex exponential (essentially the *Fourier transform* of $N_n(x)$,...next lemma) to be useful. Such a hunch would be confirmed because it is useful for proving that

- ☞ the sequence $(N_n(x - k))_{k \in \mathbb{Z}}$ is a *Riesz basis* (Lemma Q.6 page 361, Theorem Q.8 page 366) and
- ☞ the sequence $(N_n(x - k))_{k \in \mathbb{Z}}$ is a *multiresolution analysis* (Theorem Q.10 page 369).

Lemma Q.5. ¹⁷ Let $\tilde{\mathbf{F}}$ be the FOURIER TRANSFORM operator (Definition N.2 page 309).

L E M	$\tilde{\mathbf{F}}N_n(\omega) = \frac{1}{\sqrt{2\pi}} \left(\frac{1 - e^{-i\omega}}{i\omega} \right)^{n+1} = \frac{1}{\sqrt{2\pi}} e^{-i\frac{(n+1)\omega}{2}} \left(\frac{\sin(\omega_2)}{\omega_2} \right)^{n+1} \triangleq \frac{1}{\sqrt{2\pi}} e^{-i\frac{(n+1)\omega}{2}} \left(\text{sinc} \frac{\omega}{2} \right)^{n+1}$
-------------	---

☞ PROOF:


1. Proof using Theorem Q.5 page 356:

$$\begin{aligned}
 \tilde{\mathbf{F}}N_n(\omega) &= \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} N_n(x) e^{-i\omega x} dx && \text{by definition of } \tilde{\mathbf{F}} \quad (\text{Definition N.2 page 309}) \\
 &= \frac{1}{\sqrt{2\pi}} \int_{[0:1]^{n+1}} e^{-i\omega(x_1+x_2+\dots+x_{n+1})} dx_1 dx_2 \dots dx_{n+1} && \text{by Theorem Q.5} \\
 &= \frac{1}{\sqrt{2\pi}} \prod_{k=1}^{n+1} \left(\int_{[0:1]} e^{-i\omega x_k} dx_k \right) && \text{because } e^{x+y} = e^x e^y \\
 &= \frac{1}{\sqrt{2\pi}} \left(\int_0^1 e^{-i\omega x} dx \right)^{n+1} = \frac{1}{\sqrt{2\pi}} \left(\frac{e^{-i\omega x} \Big|_0^1}{-i\omega} \right)^{n+1} \\
 &= \frac{1}{\sqrt{2\pi}} \left(\frac{1 - e^{-i\omega}}{i\omega} \right)^{n+1} = \frac{1}{\sqrt{2\pi}} \left[e^{-i\frac{\omega}{2}} \left(\frac{e^{i\frac{\omega}{2}} - e^{-i\frac{\omega}{2}}}{i\omega} \right) \right]^{n+1} \\
 &= \frac{1}{\sqrt{2\pi}} \left[e^{-i\frac{\omega}{2}} \left(\frac{2i \sin\left(\frac{\omega}{2}\right)}{\frac{2i\omega}{2}} \right) \right]^{n+1} && \text{by Euler formulas} \quad (\text{Corollary ?? page ??}) \\
 &= \frac{1}{\sqrt{2\pi}} e^{-i\frac{(n+1)\omega}{2}} \left(\frac{\sin(\omega_2)}{\omega_2} \right)^{n+1}
 \end{aligned}$$

2. Proof using *rectangular pulse* example (Example N.1 page 316) and *Convolution Theorem* (Theorem P.2 page 332):


$$\begin{aligned}
 \tilde{\mathbf{F}}N_n(\omega) &= \left[\sqrt{2\pi} \right]^n [\tilde{\mathbf{F}}N_0]^{n+1} && \text{by Convolution Theorem} \quad (\text{Theorem P.2 page 332}) \\
 &= \frac{1}{\sqrt{2\pi}} \left[\sqrt{2\pi} \frac{c(b-a)}{\sqrt{2\pi}} e^{-i\left(\frac{a+b}{2}\omega\right)} \left(\frac{\sin\left(\frac{b-a}{2}\omega\right)}{\left(\frac{b-a}{2}\omega\right)} \right) \right]^{n+1} && \text{by rectangular pulse example} \\
 &= \frac{1}{\sqrt{2\pi}} \left[\sqrt{2\pi} \frac{1}{\sqrt{2\pi}} e^{-i\left(\frac{\omega}{2}\right)} \left(\frac{\sin\left(\frac{\omega}{2}\right)}{(\omega_2)} \right) \right]^{n+1} && \text{with } a = 0, b = c = 1 \quad (\text{Example N.1 page 316}) \\
 &= \frac{1}{\sqrt{2\pi}} e^{-i\left(\frac{(n+1)\omega}{2}\right)} \left(\frac{\sin\left(\frac{\omega}{2}\right)}{(\omega_2)} \right)^{n+1}
 \end{aligned}$$

Example Q.10. The Fourier transform magnitude $|\tilde{\mathbf{F}}N_0](\omega)|$ of the 0 order B-spline $N_0(x)$ is illustrated to the right.




¹⁷ [Christensen \(2008\) page 142](#) (Corollary 6.1.2)

Example Q.11. The Fourier transform magnitude $|[\tilde{F}N_1](\omega)|$ of the 1st order B-spline $N_1(x)$ is illustrated to the right.



Example Q.12. The Fourier transform magnitude $|[\tilde{F}N_2](\omega)|$ of the 2nd order B-spline $N_2(x)$ is illustrated to the right.



Q.5 Basis properties

Q.5.1 Uniqueness properties

Coefficients of a *basis sequence* are not always *unique*. Take for example a very trivial sequence (α_1, α_2) in which the coefficients are summed. If $f(x) \triangleq \alpha_1 + \alpha_2$ and $g(x) \triangleq \beta_1 + \beta_2$,

$$\begin{aligned} \text{then } \{(\alpha_1, \alpha_2) = (\beta_1, \beta_2)\} &\implies f(x) = g(x) \\ \text{but } f(x) = g(x) &\implies \{(\alpha_1, \alpha_2) = (\beta_1, \beta_2)\}, \end{aligned}$$

because for example if $(\alpha_1, \alpha_2) = (1, 2)$ and $(\beta_1, \beta_2) = (-6, 9)$, then $f(x) = g(x)$, but $(\alpha_1, \alpha_2) \neq (\beta_1, \beta_2)$. This example demonstrates that the “if and only if” condition \iff does not hold and coefficients are not unique in all *basis sequences*. But arguably a minimal requirement for any practical basis sequence is that the coefficients are *unique* (the “if and only if” condition \iff holds). And indeed, in a *B-spline* basis sequence $(N_n(x - k))_{k \in \mathbb{Z}}$, the coefficients $(\alpha_k)_{k \in \mathbb{Z}}$ are *unique*, as demonstrated by Theorem Q.6 (next).

Theorem Q.6. ¹⁸ Let $N_n(x)$ be the n TH-ORDER B-SPLINE (Definition Q.2 page 343). Let

$$f(x) \triangleq \sum_{k \in \mathbb{Z}} \alpha_k N_n(x - k) \quad \text{and} \quad g(x) \triangleq \sum_{k \in \mathbb{Z}} \beta_k N_n(x - k).$$

T
H
M

$$\left\{ f(x) = g(x) \quad \forall x \in \mathbb{R} \right\} \iff \underbrace{\{(\alpha_k)_{k \in \mathbb{Z}} = (\beta_k)_{k \in \mathbb{Z}}\}}_{\text{coefficients are UNIQUE}}$$

PROOF:

1. Proof that \iff condition holds:

$$\begin{aligned} f(x) &\triangleq \sum_{k \in \mathbb{Z}} \alpha_k N_n(x - k) && \text{by definition of } f(x) \\ &= \sum_{k \in \mathbb{Z}} \beta_k N_n(x - k) && \text{by right hypothesis} \\ &\triangleq g(x) && \text{by definition of } g(x) \end{aligned}$$

2. Proof that \implies condition holds (proof by contradiction):

(a) Suppose it does *not* hold.

¹⁸ Wojtaszczyk (1997) page 55 (Theorem 3.11)

(b) Then there exists sequences $(\alpha_k)_{k \in \mathbb{Z}}$ and $(\beta_k)_{k \in \mathbb{Z}}$ such that
 $(\alpha_k) - (\beta_k) \triangleq (\text{alpha}_k - \beta_k) \neq (0, 0, 0, \dots)$
but also such that $f(x) - g(x) = 0 \forall x \in \mathbb{R}$.

(c) If this were possible, then

$$\begin{aligned} 0 &= f(x) - g(x) \\ &= \sum_{m \in \mathbb{Z}} \alpha_m N_n(x - m) - \sum_{m \in \mathbb{Z}} \beta_m N_n(x - m) \\ &= \sum_{m \in \mathbb{Z}} (\alpha_m - \beta_m) N_n(x - m) \\ &= \sum_{m=0}^{m=n} (\alpha_m - \beta_m) \frac{1}{n!} \left[\sum_{k=0}^{n+1} (-1)^k \binom{n+1}{k} (x - k)^n \sigma(x - k) \right] \end{aligned} \quad \text{by Theorem Q.1 page 347}$$

(d) But this is *impossible* because $N(x)$ is *non-negative* (Theorem Q.4 page 354).

(e) Therefore, there is a contradiction, and the \Rightarrow condition *does* hold.



Q.5.2 Partition of unity properties

In the case in which a sequence of *B-splines* $(N_n(x - k))_{k \in \mathbb{Z}}$ is to be used as a *basis* for some subspace of $L^2_{\mathbb{R}}$, arguably one of the most important properties for the sequence to have is the *partition of unity* property such that $\sum_{k \in \mathbb{Z}} N_n(x - k) = 1$. This allows for convenient representation of the most basic functions, such as constants.¹⁹ As it turns out, B-splines *do* have this property (next theorem).

Theorem Q.7 (B-spline partition of unity). ²⁰ Let $N_n(x)$ be the *n*TH ORDER B-SPLINE (Definition Q.2 page 343).

T	$\sum_{k \in \mathbb{Z}} N_n(x - k) = 1 \quad \forall n \in \mathbb{W}$	(PARTITION OF UNITY)
---	---	----------------------

PROOF:

1. lemma: $\sum_{k \in \mathbb{Z}} N_0(x - k) = 1$. Proof:

$$\begin{aligned} \sum_{k \in \mathbb{Z}} N_0(x - k) &= \sum_{k \in \mathbb{Z}} \mathbb{1}_{[0:1]}(x - k) && \text{by definition of } N_0(x) && \text{(Definition Q.2 page 343)} \\ &= 1 && \text{by definition of } \mathbb{1}_A(x) && \text{(Definition ?? page ??)} \end{aligned}$$

2. Proof for this theorem follows from the $n = 0$ case ((1) lemma page 360), the definition of $N_n(x)$ (Definition Q.2 page 343), and Corollary ?? (page ??).

3. Alternatively, this theorem can be proved by *induction*:

(a) Base case ($n = 0$ case): by (1) lemma.

¹⁹ Jawerth and Sweldens (1994) page 8

²⁰ Christensen (2008) page 140 (Theorem 6.1.1), Höllig (2003) page 27 (3.4), Schumaker (2007) page 120 (Theorem 4.15), de Boor (2001) page 90 (B-Spline Property (i)), Chui (1988) page 2 (Theorem 1.1), Wojtaszczyk (1997) page 53 (Theorem 3.7), Cox (1972), de Boor (1972)





Figure Q.3: *auto-power spectrum* $\tilde{S}_n(\omega)$ plots of *B-splines* $N_n(x)$ (Lemma Q.6 page 361) For C and L^AT_EX source code to generate such a plot, see Section ?? (page ??).


(b) Inductive step—proof that $\sum_{k \in \mathbb{Z}} N_n(x - k) = 1 \implies \sum_{k \in \mathbb{Z}} N_{n+1}(x - k) = 1$:

$$\begin{aligned}
 \sum_{k \in \mathbb{Z}} N_{n+1}(x - k) &= \sum_{k \in \mathbb{Z}} \int_{\tau=0}^{\tau=1} N_n(x - k - \tau) d\tau && \text{by Lemma Q.2 page 343} \\
 &= \sum_{k \in \mathbb{Z}} \int_{x-u=0}^{x-u=1} N_n(u - k)(-1) du && \text{where } u \triangleq x - \tau \implies \tau = x - u \\
 &= \sum_{k \in \mathbb{Z}} \int_{u=x-1}^{u=x} N_n(u - k) du \\
 &= \int_{u=x-1}^{u=x} \left(\sum_{k \in \mathbb{Z}} N_n(u - k) \right) du \\
 &= \int_{u=x-1}^{u=x} 1 du && \text{by induction hypothesis} \\
 &= 1
 \end{aligned}$$


Example Q.13. The *partition of unity* property for the 0 *order* B-spline $N_0(x)$ (Example Q.1 page 345) is illustrated to the right.



Example Q.14. The *partition of unity* property for the 1st order B-spline $N_1(x)$ (Example Q.2 page 346) is illustrated to the right.



Example Q.15. The *partition of unity* property for the 2nd order B-spline $N_2(x)$ (Example Q.3 page 346) is illustrated to the right.



Q.5.3 Riesz basis properties

Lemma Q.6. Let $N_n(x)$ be the n th ORDER B-SPLINE (Definition Q.2 page 343).

Let $\tilde{S}_n(\omega) \triangleq 2\pi \sum_{k \in \mathbb{Z}} |\tilde{F}N_n(\omega - 2\pi k)|^2$ be the AUTO-POWER SPECTRUM (Definition ?? page ??) of $N_n(x)$.

LEM	(1). $0 < \tilde{S}_n(\omega) \leq 1 \quad \forall \omega \in \mathbb{R} \quad , \quad \forall n \in \mathbb{W}$ (2). $\tilde{S}_n(\omega) = 1 \quad \forall \omega \in \mathbb{R} \quad , \quad \text{for } n = 0$	(3). $\tilde{S}_n(0) = 1 \quad \forall n \in \mathbb{W}$ (4). $\tilde{S}_n(\pi) \leq \frac{1}{3} \quad \forall n \in \mathbb{W} \setminus \{0\}$	(Note: see illustration in Figure Q.3 page 361.)
-----	--	---	---

PROOF:

1. lemma: $\tilde{S}_n(\omega) = \sum_{k \in \mathbb{Z}} \left[\frac{\sin\left(\frac{\omega}{2}\right)}{k\pi - \frac{\omega}{2}} \right]^{2(n+1)}$. Proof:

$$\tilde{S}_n(\omega) \triangleq 2\pi \sum_{k \in \mathbb{Z}} |\tilde{\mathbf{F}}\mathbf{N}_n(\omega - 2\pi k)|^2 \quad \text{by Definition ?? page ??}$$

$$= 2\pi \sum_{k \in \mathbb{Z}} \left| \frac{1}{\sqrt{2\pi}} e^{-i \frac{(n+1)(\omega - 2\pi k)}{2}} \left(\frac{\sin\left(\frac{\omega - 2\pi k}{2}\right)}{\frac{\omega - 2\pi k}{2}} \right)^{n+1} \right|^2 \quad \text{by Lemma Q.5 page 358}$$

$$= \sum_{k \in \mathbb{Z}} \left[\frac{\sin\left(\frac{\omega - 2\pi k}{2}\right)}{\frac{\omega - 2\pi k}{2}} \right]^{2(n+1)}$$

$$= \sum_{k \in \mathbb{Z}} \left[\frac{\sin\left(\frac{\omega}{2} - k\pi\right)}{k\pi - \frac{\omega}{2}} \right]^{2(n+1)}$$

$$= \sum_{k \in \mathbb{Z}} \left[\frac{(-1)^k \sin\left(\frac{\omega}{2}\right)}{k\pi - \frac{\omega}{2}} \right]^{2(n+1)}$$

$$= \sum_{k \in \mathbb{Z}} \left[\frac{\sin\left(\frac{\omega}{2}\right)}{k\pi - \frac{\omega}{2}} \right]^{2(n+1)}$$

2. lemma (one sided series form):

$$\begin{aligned} \tilde{S}_n(\omega) &= \sum_{k \in \mathbb{Z}} \left[\frac{\sin\left(\frac{\omega}{2}\right)}{k\pi - \frac{\omega}{2}} \right]^{2(n+1)} \\ &= \left[\frac{\sin(\omega/2)}{(\omega/2)} \right]^{2(n+1)} + \left[\frac{\sin(\omega/2)}{(\pi/2)} \right]^{2(n+1)} \left(\sum_{k=1}^{\infty} \left[\frac{1}{2k - \frac{\omega}{\pi}} \right]^{2(n+1)} + \sum_{k=1}^{\infty} \left[\frac{1}{2k + \frac{\omega}{\pi}} \right]^{2(n+1)} \right) \end{aligned} \quad \text{by (1) lemma}$$

3. lemma: $\tilde{S}_n(\omega)$ is *continuous* for all $\omega \in \mathbb{R}$.

Proof: $\sin(\omega/2)$ and $\omega/2$ are *continuous*, so $\tilde{S}_n(\omega)$ is *continuous* as well.

4. lemma: $\tilde{S}_n(\omega)$ is *periodic* with period 2π (and so we only need to examine $\tilde{S}_n(\omega)$ for $\omega \in [0 : 2\pi]$). Proof of *periodicity*: This follows directly from Proposition ?? (page ??).

5. lemma: $\tilde{S}_n(-\omega) = \tilde{S}_n(\omega)$ (*symmetric* about 0) and $\tilde{S}_n(\pi - \omega) = \tilde{S}_n(\pi + \omega)$ (*symmetric* about π). Proof: This follows directly from Proposition ?? (page ??).



6. Proof that $\tilde{S}_n(0) = 1$:

$$\begin{aligned}
 \tilde{S}_n(0) &= \lim_{\omega \rightarrow 0} \tilde{S}_n(\omega) && \text{by (3) lemma} \\
 &= \lim_{\omega \rightarrow 0} \left[\left[\frac{\sin(\omega/2)}{(\omega/2)} \right]^{2(n+1)} + \left[\frac{\sin(\omega/2)}{(\pi/2)} \right]^{2(n+1)} \left(\sum_{k=1}^{\infty} \left[\frac{1}{2k - \frac{\omega}{\pi}} \right]^{2(n+1)} + \sum_{k=1}^{\infty} \left[\frac{1}{2k + \frac{\omega}{\pi}} \right]^{2(n+1)} \right) \right] && \text{by (2) lemma} \\
 &= \lim_{\omega \rightarrow 0} \left[\frac{\cos\left(\frac{\omega}{2}\right)}{-\frac{1}{2}} \right]^{2(n+1)} + 0 && \text{by l'Hôpital's rule} \\
 &= (-1)^{2(n+1)} = 1
 \end{aligned}$$

7. Proof that $\tilde{S}_n(\pi)$ converges to some value > 0 :

(a) Proof that $\tilde{S}_n(\pi) > 0$:

$$\begin{aligned}
 \tilde{S}_n(\pi) &= \left[\frac{\sin(\pi/2)}{(\pi/2)} \right]^{2(n+1)} + \left[\frac{\sin(\pi/2)}{(\pi/2)} \right]^{2(n+1)} \left(\sum_{k=1}^{\infty} \left[\frac{1}{2k - \frac{\pi}{\pi}} \right]^{2(n+1)} + \sum_{k=1}^{\infty} \left[\frac{1}{2k + \frac{\pi}{\pi}} \right]^{2(n+1)} \right) && \text{by (2) lemma} \\
 &= \left(\frac{2}{\pi} \right)^{2(n+1)} \left[1 + \left(\frac{1}{1} \right)^{2(n+1)} + \left(\frac{1}{3} \right)^{2(n+1)} + \left(\frac{1}{3} \right)^{2(n+1)} + \left(\frac{1}{5} \right)^{2(n+1)} + \left(\frac{1}{5} \right)^{2(n+1)} + \dots \right] \\
 &= 2 \left(\frac{2}{\pi} \right)^{2(n+1)} \underbrace{\sum_{k=1}^{\infty} \left[\frac{1}{2k-1} \right]^{2(n+1)}}_{\text{Dirichlet Lambda function } \lambda(2n+2)} \\
 &> 0 && \text{because } x^2 > 0 \text{ for all } x \in \mathbb{R} \setminus \{0\}
 \end{aligned}$$

(b) Proof that $\tilde{S}_n(\pi)$ converges:

$$\begin{aligned}
 \tilde{S}_n(\pi) &= 2 \left(\frac{2}{\pi} \right)^{2(n+1)} \sum_{k=1}^{\infty} \left(\frac{1}{2k-1} \right)^{2(n+1)} && \text{by item (7a)} \\
 &\leq 2 \left(\frac{2}{\pi} \right)^{2(n+1)} \sum_{k=1}^{\infty} \left(\frac{1}{k} \right)^{2(n+1)} \\
 &\leq 2 \left(\frac{2}{\pi} \right)^{2(n+1)} \sum_{k=1}^{\infty} \left(\frac{1}{k} \right)^2 \\
 &\implies \text{convergence} && \text{by comparison test}
 \end{aligned}$$

(c) Tighter bounds for $\tilde{S}_n(\pi)$ for certain values of $n \in \{0, 1, 2, 3, 4\}$:

$$\begin{aligned}
 \tilde{S}_n(\pi) &= 2 \left(\frac{2}{\pi} \right)^{2(n+1)} \sum_{k=1}^{\infty} \left(\frac{1}{2k-1} \right)^{2(n+1)} && \text{by item (7a)} \\
 &= 2 \left(\frac{2}{\pi} \right)^{2(n+1)} U_{2(n+1)} && \text{by } \text{Jolley (1961), pages 56–57 ((307))} \\
 &= 2 \left(\frac{2}{\pi} \right)^{2(n+1)} \left[\frac{\pi^{2(n+1)} \alpha_{n+1}}{(4)[(2n+2)!]} \right] && \text{by } \text{Jolley (1961), pages 56–57 ((307))} \\
 &= \frac{2^{2n+1} \alpha_{n+1}}{(2n+2)!}
 \end{aligned}$$

$$\begin{aligned}
 &= \left\{ \begin{array}{ll} \frac{2^1(1)}{2!} & \text{for } n = 0 \quad (\alpha_1 = 1) \\ \frac{2^2(1)}{4!} & \text{for } n = 1 \quad (\alpha_2 = 1) \\ \frac{2^3(3)}{6!} & \text{for } n = 2 \quad (\alpha_3 = 3) \\ \frac{2^7(17)}{8!} & \text{for } n = 3 \quad (\alpha_4 = 17) \\ \frac{2^9(155)}{10!} & \text{for } n = 4 \quad (\alpha_5 = 155) \end{array} \right\} \quad \text{by } \text{Jolley (1961), page 234 (1130)} \\
 &= \left\{ \begin{array}{ll} 1 & \text{for } n = 0 \\ \frac{1}{3} & \text{for } n = 1 \\ \frac{2}{15} & \text{for } n = 2 \\ \frac{15}{315} & \text{for } n = 3 \\ \frac{62}{2835} & \text{for } n = 4 \end{array} \right\} = \left\{ \begin{array}{ll} 1 & \text{for } n = 0 \\ 0.3333333333333333 \dots & \text{for } n = 1 \\ 0.1333333333333333 \dots & \text{for } n = 2 \\ 0.0539682539682 \dots & \text{for } n = 3 \\ 0.0218694885361 \dots & \text{for } n = 4 \end{array} \right\}
 \end{aligned}$$

(d) Being important for the $n = 0$ case, note that²¹

$$\underbrace{\sum_{k=1}^{\infty} \left(\frac{1}{2k-1} \right)^2}_{\text{Dirichlet Lambda function } \lambda(2)} = 1 + \frac{1}{3^2} + \frac{1}{5^2} + \frac{1}{7^2} + \dots = \frac{\pi^2}{8}$$

(e) Proof that $\tilde{S}_n(\pi) \leq \frac{1}{3}$: because $\tilde{S}_n(\pi) = \frac{1}{3}$ for $n = 1$ (item (7c) page 363) and because $\tilde{S}_n(\pi)$ is decreasing for increasing n .

8. lemma: $\tilde{S}_n(\omega)$ converges to some value $> 0 \forall \omega \in \mathbb{R}$. Proof:

(a) For $\omega = 0$, $\tilde{S}_n(\omega) = 1$ by item (6).

(b) Proof that $\tilde{S}_n(\omega) > 0$ for $\omega \in (0 : 2\pi)$:

$$\begin{aligned}
 \tilde{S}_n(\omega) &= \left[\frac{\sin(\omega/2)}{(\omega/2)} \right]^{2(n+1)} + \left[\frac{\sin(\omega/2)}{(\pi/2)} \right]^{2(n+1)} \left(\sum_{k=1}^{\infty} \left[\frac{1}{2k - \frac{\omega}{\pi}} \right]^{2(n+1)} + \sum_{k=1}^{\infty} \left[\frac{1}{2k + \frac{\omega}{\pi}} \right]^{2(n+1)} \right) \quad \text{by (2) lemma} \\
 &> 0
 \end{aligned}$$

(c) Proof that $\tilde{S}_n(\omega)$ converges:

$$\begin{aligned}
 \text{i. lemma: } \sum_{k=1}^{\infty} \left[\frac{1}{2k \pm \frac{\omega}{\pi}} \right]^{2(n+1)} \text{ converges. Proof:} \\
 \lim_{b \rightarrow \infty} \int_1^b \left[\frac{1}{2y \pm \frac{\omega}{\pi}} \right]^{2(n+1)} dy &= \lim_{b \rightarrow \infty} \int_1^b \left[2y \pm \frac{\omega}{\pi} \right]^{-2n-2} dy \\
 &= \lim_{b \rightarrow \infty} \frac{\left[2y \pm \frac{\omega}{\pi} \right]^{-2n-1}}{2(-2n-1)} \Big|_1^b \\
 &= \lim_{b \rightarrow \infty} \left(\frac{-1}{2(2n+1)} \right) \left[\frac{1}{\left[2b \pm \frac{\omega}{\pi} \right]^{2n+1}} - \frac{1}{\left[2 \pm \frac{\omega}{\pi} \right]^{2n+1}} \right] \\
 &= 0 + \frac{1}{2(2n+1) \left[2 \pm \frac{\omega}{\pi} \right]^{2n+1}} \\
 &< \infty \quad \forall \omega \in [0 : 2\pi] \\
 \Rightarrow \sum_{k=1}^{\infty} \left[\frac{1}{2k \pm \frac{\omega}{\pi}} \right]^{2(n+1)} \text{ converges} &\quad \text{by integral test}
 \end{aligned}$$

²¹ [Nahin \(2011\) page 153](#), [Bailey et al. \(2013\) page 334 \(Catalan's Constant\)](#), [Bailey et al. \(2011\) \(15\)](#), [Wells \(1987\) page 36 \(Dictionary entry for \$\pi\$: pages 31–37\)](#), [Heinbockel \(2010\) page 94 \(2.27 Dirichlet Lambda function\)](#)

ii. completion of proof using (8(c)i) lemma ...

$$\begin{aligned}\tilde{S}_n(\omega) &= \left[\frac{\sin(\omega/2)}{(\omega/2)} \right]^{2(n+1)} + \left[\frac{\sin(\omega/2)}{(\pi/2)} \right]^{2(n+1)} \left(\sum_{k=1}^{\infty} \left[\frac{1}{2k - \frac{\omega}{\pi}} \right]^{2(n+1)} + \sum_{k=1}^{\infty} \left[\frac{1}{2k + \frac{\omega}{\pi}} \right]^{2(n+1)} \right) \text{ by (2) lemma} \\ &\implies \tilde{S}_n(\omega) \text{ converges } \forall \omega \in (0 : 2\pi) \quad \text{by (8(c)i) lemma}\end{aligned}$$

9. lemma (an expression for $\tilde{S}'_n(\omega)$):

$$\begin{aligned}\tilde{S}'_n(\omega) &\triangleq \frac{d}{d\omega} \tilde{S}_n(\omega) \\ &= \frac{d}{d\omega} \sum_{k \in \mathbb{Z}} \left[\frac{\sin\left(\frac{\omega}{2}\right)}{k\pi - \frac{\omega}{2}} \right]^{2(n+1)} \quad \text{by (1) lemma page 362} \\ &= \sum_{k \in \mathbb{Z}} \frac{d}{d\omega} \left[\frac{\sin\left(\frac{\omega}{2}\right)}{k\pi - \frac{\omega}{2}} \right]^{2(n+1)} \quad \text{by linearity of } \frac{d}{d\omega} \text{ operator} \\ &= \sum_{k \in \mathbb{Z}} 2(n+1) \left[\frac{\sin\left(\frac{\omega}{2}\right)}{k\pi - \frac{\omega}{2}} \right]^{2n+1} \frac{d}{d\omega} \left[\frac{\sin\left(\frac{\omega}{2}\right)}{k\pi - \frac{\omega}{2}} \right] \quad \text{by power rule} \\ &= 2(n+1) \sum_{k \in \mathbb{Z}} \left[\frac{\sin\left(\frac{\omega}{2}\right)}{k\pi - \frac{\omega}{2}} \right]^{2n+1} \left[\frac{\frac{1}{2} \cos\left(\frac{\omega}{2}\right) \left(k\pi - \frac{\omega}{2}\right) - \sin\left(\frac{\omega}{2}\right) \left(-\frac{1}{2}\right)}{\left(k\pi - \frac{\omega}{2}\right)^2} \right] \quad \text{by quotient rule} \\ &= (n+1) \sum_{k \in \mathbb{Z}} \left[\frac{\sin\left(\frac{\omega}{2}\right)}{k\pi - \frac{\omega}{2}} \right]^{2n+1} \left[\frac{\cos\left(\frac{\omega}{2}\right) \left(k\pi - \frac{\omega}{2}\right) + \sin\left(\frac{\omega}{2}\right)}{\left(k\pi - \frac{\omega}{2}\right)^2} \right]\end{aligned}$$

10. lemma: $\tilde{S}'_n(0) = \tilde{S}'_n(\pi) = 0$. Proof: This follows from Proposition ?? (page ??). Here is alternate proof:

$$\begin{aligned}\tilde{S}'_n(0) &= \lim_{\omega \rightarrow 0} \tilde{S}'_n(\omega) \\ &= \lim_{\omega \rightarrow 0} (n+1) \sum_{k \in \mathbb{Z}} \left[\frac{\sin\left(\frac{\omega}{2}\right)}{k\pi - \frac{\omega}{2}} \right]^{2n+1} \left[\frac{\cos\left(\frac{\omega}{2}\right) \left(k\pi - \frac{\omega}{2}\right) + \sin\left(\frac{\omega}{2}\right)}{\left(k\pi - \frac{\omega}{2}\right)^2} \right] \quad \text{by (9) lemma} \\ &= \lim_{\omega \rightarrow 0} (n+1) \left[\frac{\sin\left(\frac{\omega}{2}\right)}{-\frac{\omega}{2}} \right]^{2n+1} \left[\frac{\cos\left(\frac{\omega}{2}\right) \left(-\frac{\omega}{2}\right) + \sin\left(\frac{\omega}{2}\right)}{\left(-\frac{\omega}{2}\right)^2} \right] \\ &= (n+1) \lim_{\omega \rightarrow 0} \left[\frac{\sin\left(\frac{\omega}{2}\right)}{-\frac{\omega}{2}} \right]^{2n+1} \lim_{\omega \rightarrow 0} \left[\frac{\cos\left(\frac{\omega}{2}\right) \left(-\frac{\omega}{2}\right) + \sin\left(\frac{\omega}{2}\right)}{\left(-\frac{\omega}{2}\right)^2} \right] \\ &= (n+1) [-1]^{2n+1} \lim_{\omega \rightarrow 0} \left[\frac{-\frac{1}{2} \sin\left(\frac{\omega}{2}\right) \left(-\frac{\omega}{2}\right) + \cos\left(\frac{\omega}{2}\right) \left(-\frac{1}{2}\right) + \cos\left(\frac{\omega}{2}\right) \left(\frac{1}{2}\right)}{-\frac{2}{2} \left(-\frac{\omega}{2}\right)} \right] \quad \text{by l'Hôpital's rule} \\ &= (1)(0) \\ &= 0\end{aligned}$$

$$\begin{aligned}
\tilde{S}'_n(\pi) &= (n+1) \sum_{k \in \mathbb{Z}} \left[\frac{\sin\left(\frac{\pi}{2}\right)}{k\pi - \frac{\pi}{2}} \right]^{2n+1} \left[\frac{\cos\left(\frac{\pi}{2}\right)\left(k\pi - \frac{\pi}{2}\right) + \sin\left(\frac{\pi}{2}\right)}{\left(k\pi - \frac{\pi}{2}\right)^2} \right] \\
&= (n+1) \sum_{k \in \mathbb{Z}} \left[\frac{1}{k\pi - \frac{\pi}{2}} \right]^{2n+1} \left[\frac{0\left(k\pi - \frac{\pi}{2}\right) + 1}{\left(k\pi - \frac{\pi}{2}\right)^2} \right] \\
&= (n+1) \left(\frac{2}{\pi} \right)^{2n+3} \sum_{k \in \mathbb{Z}} \left[\frac{1}{2k-1} \right]^{2n+3} \\
&= (n+1) \left(\frac{2}{\pi} \right)^{2n+3} \left[\left(\frac{1}{1} \right)^{2n+3} + \left(\frac{1}{-1} \right)^{2n+3} + \left(\frac{1}{3} \right)^{2n+3} + \left(\frac{1}{-3} \right)^{2n+3} + \dots \right] \\
&= (n+1) \left(\frac{2}{\pi} \right)^{2n+3} \sum_{k=1}^{\infty} (-1)^{k+1} \alpha_k \quad \text{where } \alpha_k \triangleq \begin{cases} \left(\frac{1}{k} \right)^{2n+3} & \text{for } k \text{ odd} \\ \left(\frac{1}{k-1} \right)^{2n+3} & \text{for } k \text{ even} \end{cases} \\
&= 0 \quad \text{because } \lim_{k \rightarrow \infty} \alpha_k = 0 \text{ and by Alternating Series Test}
\end{aligned}$$

11. lemma: $\tilde{S}_n(\omega)$ is *decreasing* with respect to $\omega \in [0 : \pi]$. Proof:

$$\begin{aligned}
\tilde{S}'_n(\omega) &= (n+1) \sum_{k \in \mathbb{Z}} \left[\frac{\sin\left(\frac{\omega}{2}\right)}{k\pi - \frac{\omega}{2}} \right]^{2n+1} \left[\frac{\cos\left(\frac{\omega}{2}\right)\left(k\pi - \frac{\omega}{2}\right) + \sin\left(\frac{\omega}{2}\right)}{\left(k\pi - \frac{\omega}{2}\right)^2} \right] \quad \text{by (9) lemma page 365} \\
&= (n+1) \underbrace{\left(\sin \frac{\omega}{2} \right)^{2n+1}}_{\geq 0 \text{ for } \omega \in [0 : 2\pi]} \sum_{k \in \mathbb{Z}} \left[\frac{1}{k\pi - \frac{\omega}{2}} \right]^{2n+2} \left[\underbrace{\left(\cos \frac{\omega}{2} \right)}_{\text{sign change at } \omega = \pi} + \underbrace{\frac{\sin \frac{\omega}{2}}{k\pi - \frac{\omega}{2}}}_{\substack{\text{decreasing w.r.t. } \omega \in \mathbb{R}}} \right]
\end{aligned}$$

12. lemma: $\tilde{S}_n(\omega)$ is *increasing* with respect to $\omega \in [\pi : 2\pi]$. Proof: This is true because $\tilde{S}_n(\omega)$ is *decreasing* in $[0 : \pi]$ ((11) lemma) and because $\tilde{S}_n(\omega)$ is *symmetric* about $\omega = \pi$ ((5) lemma).

13. Proof that $0 < \tilde{S}_n(\omega) \leq 1$:

- (a) $\tilde{S}_n(\omega) > 0$ by (8) lemma and
- (b) $\tilde{S}_n(0) = 1$ by item (6) and
- (c) $\tilde{S}_n(\omega)$ is *decreasing* from $\omega = 0$ to $\omega = \pi$ by (11) lemma and
- (d) $\tilde{S}_n(\omega)$ is *increasing* from $\omega = \pi$ to $\omega = 2\pi$ by (12) lemma and
- (e) $\tilde{S}_n(2\pi) = 1$ because $\tilde{S}_n(2\pi) = \tilde{S}_n(0)$ by (4) lemma.

Theorem Q.8. ²²

T H M	1. $(N_n(x-k))_{k \in \mathbb{Z}}$ is a RIESZ BASIS <i>for</i> $\text{span}(N_n(x-k))_{k \in \mathbb{Z}}$ $\iff n = 0$ 2. $(N_n(x-k))_{k \in \mathbb{Z}}$ is an ORTHONORMAL BASIS <i>for</i> $\text{span}(N_n(x-k))_{k \in \mathbb{Z}}$
-------------	---

PROOF:

²² Wojtaszczyk (1997) page 56 (Proposition 3.12), Prasad and Iyengar (1997) page 148 (Theorem 6.3), Forster and Massopust (2009) page 66 (Theorem 2.25)

1. Proof that $(N_n(x - k))_{k \in \mathbb{Z}}$ is a *Riesz basis* for $\text{span}(N_n(x - k))_{k \in \mathbb{Z}}$:

$$\begin{aligned} 0 < \tilde{S}_n(\omega) &\leq 1 && \text{by Lemma Q.6 page 361 (1)} \\ \implies (N_n(x - k))_{k \in \mathbb{Z}} &\text{ is a } Riesz \text{ basis for } \text{span}(N_n(x - k))_{k \in \mathbb{Z}} && \text{by Theorem ?? page ??} \end{aligned}$$

2. Proof that $\{n = 0\} \iff (N_n(x - k))_{k \in \mathbb{Z}}$ is an *orthonormal basis* for $\text{span}(N_n(x - k))_{k \in \mathbb{Z}}$:

$$\begin{aligned} n = 0 \iff \tilde{S}_n(\omega) &= 1 && \text{by Lemma Q.6 page 361 (2), (4)} \\ \iff (N_n(x - k))_{k \in \mathbb{Z}} &\text{ is an orthonormal basis for } \text{span}(N_n(x - k))_{k \in \mathbb{Z}} && \text{by Theorem ?? page ??} \end{aligned}$$



Q.6 Mutiresolution properties

Q.6.1 Introduction

In 1989, Stéphane G. Mallat introduced the *Mutiresolution Analysis* (MRA) structure (Definition ?? page ??) An MRA is very powerful because it can be used to approximate functions at incrementally increasing “scales” of resolution, and furthermore induces a *wavelet*. In fact, the MRA has become the dominate wavelet construction method. Moreover, P.G. Lemarié has proved that all wavelets with *compact support* are generated by an MRA. ²³

Q.6.2 B-spline dyadic decomposition

One key feature of an MRA is *dyadic decomposition* such that $N_n(x) = \sum_k \alpha_n N_n(2x - k)$ for some sequence (α_n) . As it turns out, *B-splines* also have this property (next theorem).

Theorem Q.9 (*B-spline dyadic decomposition*). ²⁴ Let $N_n(x)$ be the n TH ORDER B-SPLINE.

T H M	$N_n(x) = \frac{1}{2^n} \sum_{k=0}^{n+1} \binom{n+1}{k} N_n(2x - k) \quad \forall n \in \mathbb{W}, \forall x \in \mathbb{R}$
-------------	---

PROOF:

1. Base case ...proof for $n = 0$ case:

$$\begin{aligned} N_0(x) &= \mathbb{1}_{[0:1]}(x) && \text{by definition of } \mathbb{1}_A(x) \quad (\text{Definition ?? page ??}) \\ &= \mathbb{1}_{[0:\frac{1}{2}]}(x) + \mathbb{1}_{[\frac{1}{2}:1]}(x) \\ &= \mathbb{1}_{[2x0:2x\frac{1}{2}]}(2x) + \mathbb{1}_{[2x\frac{1}{2}-1:2x1-1]}(2x - 1) \\ &= \mathbb{1}_{[0:1]}(2x) + \mathbb{1}_{[0:1]}(2x - 1) \\ &= \frac{1}{2^0} \sum_{k=0}^{0+1} \binom{0+1}{k} N_0(2x - k) \end{aligned}$$

²³ Mallat (1999) page 240, Definition ?? (page ??)

²⁴ Prasad and Iyengar (1997) pages 151–152 (proof using Fourier transform)


2. Induction step...proof that n case $\implies n + 1$ case:

$$\begin{aligned}
 N_{n+1}(x) &= \int_0^1 N_n(x - \tau) d\tau && \text{by Lemma Q.2 page 343} \\
 &= \int_0^1 \frac{1}{2^n} \sum_{k=0}^{n+1} \binom{n+1}{k} N_n(2x - 2\tau - k) d\tau && \text{by induction hypothesis} \\
 &= \frac{1}{2^n} \sum_{k=0}^{n+1} \binom{n+1}{k} \int_{\tau=0}^{\tau=1} N_n(2x - 2\tau - k) d\tau && \text{by linearity of } \sum \text{ operator} \\
 &= \frac{1}{2^n} \sum_{k=0}^{n+1} \binom{n+1}{k} \int_{u=0}^{u=2} N_n(2x - u - k) \frac{1}{2} du && \text{where } u \triangleq 2\tau \implies \tau = \frac{1}{2}u \\
 &= \frac{1}{2^{n+1}} \sum_{k=0}^{n+1} \binom{n+1}{k} \left[\int_{u=0}^{u=1} N_n(2x - k - u) du + \int_{u=1}^{u=2} N_n(2x - k - u) du \right] \\
 &= \frac{1}{2^{n+1}} \sum_{k=0}^{n+1} \binom{n+1}{k} \left[\int_{u=0}^{u=1} N_n(2x - k - u) du + \int_{v=0}^{v=1} N_n(2x - k - v - 1) dv \right] \text{ where } v \triangleq u - 1 \implies u = v + 1 \\
 &= \frac{1}{2^{n+1}} \sum_{k=0}^{n+1} \binom{n+1}{k} [N_n(2x - k) + N_n(2x - k - 1)] && \text{by Lemma Q.2 page 343} \\
 &= \frac{1}{2^{n+1}} \left[\sum_{k=0}^{n+1} \binom{n+1}{k} N_n(2x - k) + \sum_{m=1}^{n+2} \binom{n+1}{m-1} N_n(2x - m) \right] \text{ where } m \triangleq k + 1 \implies k = m - 1 \\
 &= \frac{1}{2^{n+1}} \left[\underbrace{\sum_{k=1}^{n+1} \left[\binom{n+1}{k} + \binom{n+1}{k-1} \right] N_n(2x - k)}_{\text{common indices of above two summations}} + \underbrace{\binom{n+1}{0} N_n(2x - 0)}_{k=0 \text{ term}} + \underbrace{\binom{n+2}{n+2} N_n(2x - n - 2)}_{m=n+2 \text{ term}} \right] \\
 &= \frac{1}{2^{n+1}} \left[\underbrace{\sum_{k=1}^{n+1} \binom{n+2}{k} N_n(2x - k)}_{\text{by Stifel formula (Theorem ?? page ??)}} + \underbrace{\binom{n+2}{0} N_n(2x - 0)}_{\text{because } \binom{n+1}{0} = 1 = \binom{n+2}{0}} + \binom{n+2}{n+2} N_n(2x - n - 2) \right] \\
 &= \frac{1}{2^{n+1}} \sum_{k=0}^{n+2} \binom{n+2}{k} N_n(2x - k)
 \end{aligned}$$

Example Q.16.²⁵ The 0 order B-spline dyadic decomposition

$$N_0(x) = \frac{1}{1} \sum_{k=0}^{k=1} \binom{1}{k} N_0(2x - k)$$

is illustrated to the right.




²⁵ Strang (1989) page 615 (Box function), Strang and Nguyen (1996) page 441 (Box function)

Example Q.17. ²⁶The 1st order B-spline dyadic decomposition

$$N_1(x) = \frac{1}{2} \sum_{k=0}^{k=2} \binom{2}{k} N_1(2x - k)$$


is illustrated to the right.



Example Q.18. The 2nd order B-spline dyadic decomposition

$$N_2(x) = \frac{1}{4} \sum_{k=0}^{k=3} \binom{3}{k} N_2(2x - k)$$

is illustrated to the right.



Q.6.3 B-spline MRA scaling functions

Theorem Q.10. Let $f N_n(x)$ be the n TH ORDER B-SPLINE (Definition Q.2 page 343).

Let $V_j \triangleq \text{span}(N_n(2^j x - k))_{k \in \mathbb{Z}}$.

T H M $(V_j)_{j \in \mathbb{Z}}$ is a MULTIRESOLUTION ANALYSIS on $L^2_{\mathbb{R}}$ with SCALING FUNCTION $\phi(x) \triangleq N_n(x)$

PROOF:

1. lemma: $(N_n(x - k))_{k \in \mathbb{Z}}$ is a *Riesz sequence*. Proof: by Theorem Q.8 (page 366).

2. lemma: $\exists (h_k)$ such that $N_n(x) = \sum_{k \in \mathbb{Z}} h_k N_n(2x - k)$. Proof: by Theorem Q.9 (page 367). In fact, note that $h_k = \frac{1}{2^n \sqrt{2}} \binom{n+1}{k}$

3. lemma: $\tilde{F}N_n(\omega)$ is *continuous* at 0. Proof:

$$\tilde{F}N_n(\omega) = \frac{1}{\sqrt{2\pi}} e^{-i\frac{(n+1)\omega}{2}} \left(\text{sinc} \frac{\omega}{2} \right)^{n+1} \quad \text{by Lemma Q.5 page 358}$$

\implies continuous at 0 by known property of sinc function

4. lemma: $\tilde{\phi}(0) \neq 0$. Proof:

$$\begin{aligned} \tilde{F}N_n(0) &= \frac{1}{\sqrt{2\pi}} e^{-i\frac{(n+1)\omega}{2}} \left(\text{sinc} \frac{\omega}{2} \right)^{n+1} \Big|_{\omega=0} && \text{by Lemma Q.5 page 358} \\ &= 1 \cdot \frac{1}{1/2} = 2 && \text{by } l'Hôpital's \text{ rule} \\ &\neq 0 \end{aligned}$$

5. The completion of this proof follows directly from (1) lemma, (2) lemma, (3) lemma, (4) lemma, and Theorem ?? (page ??).

²⁶ Strang (1989) page 615 (Hat function), Strang and Nguyen (1996) page 442 (Hat function), Heil (2011) page 380 (Fig. 12.10)




Figure Q.4: *dilation equation* demonstrations for selected B-splines (Example Q.19 page 370)

Q.6.4 B-spline MRA coefficient sequences

Because each *B-spline* $N_n(x)$ is the *scaling function* for an *MRA* (Theorem Q.10 page 369), each *B-spline* also satisfies the *dilation equation* (Theorem ?? page ??) such that

$$N_n(x) = \sqrt{2} \sum_{k \in \mathbb{Z}} h_k N(2x - k) \quad \text{where} \quad h_k = \begin{cases} \frac{\sqrt{2}}{2^{n+1}} \binom{n+1}{k} & \text{for } n = 0, 1, \dots, n \\ 0 & \text{otherwise} \end{cases}$$

The resulting sequence $(h_k)_{k \in \mathbb{Z}}$ is the *ordern B-spline MRA coefficient sequence* induced by the *order n B-spline MRA scaling sequence* $\phi(x) \triangleq N_n(x)$.²⁷

Example Q.19. See Figure Q.4 (page 370) for some *dilation equation* demonstrations of selected B-splines.

Theorem Q.11 (*B-spline scaling coefficients*). *Let $(L^2_{\mathbb{R}}, (\mathcal{V}_j), \phi, (h_n))$ be an MRA system (Definition ?? page ??). Let $N_n(x)$ be a nth ORDER B-SPLINE (Definition Q.2 page 343).*

T H M	$\underbrace{\phi(x) \triangleq N_n(x)}_{(1) \text{ B-spline scaling function}} \implies \underbrace{(h_k)}_{\text{scaling sequence in "time"} \atop \text{in "time" domain}} = \begin{cases} \frac{\sqrt{2}}{2^{n+1}} \binom{n}{k} & \text{for } k = 0, 1, \dots, n \\ 0 & \text{otherwise} \end{cases}$ $\iff \check{h}(z) \Big _{z \triangleq e^{i\omega}} = \frac{\sqrt{2}}{2^n} (1 + z^{-1})^{n+1} \Big _{z \triangleq e^{i\omega}} \quad (3) \text{ scaling sequence in "z domain"}$ $\iff \check{h}(\omega) = 2\sqrt{2} e^{-i\frac{n+1}{2}\omega} \left[\cos\left(\frac{\omega}{2}\right) \right] \quad (4) \text{ scaling sequence in "frequency"}$
----------------------	--

PROOF:

1. Proof that (1) \implies (3): By Theorem Q.10 page 369 we know that $N_n(x)$ is a *scaling function* (Definition ?? page ??). So then we know that we can use Lemma ?? page ??.

$$\begin{aligned}
 \check{h}(\omega) &= \sqrt{2} \frac{\tilde{\phi}(2\omega)}{\tilde{\phi}(\omega)} && \text{by Lemma ?? page ??} \\
 &= \sqrt{2} \frac{\tilde{N}_n(2\omega)}{\tilde{N}_n(\omega)} && \text{by (1)} \\
 &= \sqrt{2} \frac{\frac{1}{\sqrt{2\pi}} \left(\frac{1-e^{-i2\omega}}{2i\omega} \right)^{n+1}}{\frac{1}{\sqrt{2\pi}} \left(\frac{1-e^{-i\omega}}{i\omega} \right)^{n+1}} && \text{by Lemma Q.5 page 358}
 \end{aligned}$$

²⁷For Octave/ MatLab code useful for plotting a function given a sequence of coefficients (h_k) , see Section ?? (page ??).

$$\begin{aligned}
&= \frac{\sqrt{2}}{2^{n+1}} \left(\frac{1 - z^{-2}}{1 - z^{-1}} \right)^{n+1} \Big|_{z=e^{i\omega}} \\
&= \frac{\sqrt{2}}{2^{n+1}} \left[\left(\frac{1 - z^{-2}}{1 - z^{-1}} \right) \left(\frac{1 + z^{-1}}{1 + z^{-1}} \right) \right]^{n+1} \Big|_{z=e^{i\omega}} \\
&= \frac{\sqrt{2}}{2^{n+1}} \left(\frac{(1 - z^{-2})(1 + z^{-1})}{1 - z^{-2}} \right)^{n+1} \Big|_{z=e^{i\omega}} \\
&= \frac{\sqrt{2}}{2^n} (1 + z^{-1})^{n+1} \Big|_{z \triangleq e^{i\omega}}
\end{aligned}$$

2. Proof that (3) \iff (2):

$$\begin{aligned}
\check{h}(z) \Big|_{z \triangleq e^{i\omega}} &= \frac{\sqrt{2}}{2^n} (1 + z^{-1})^{n+1} \Big|_{z \triangleq e^{i\omega}} && \text{by (3)} \\
&= \frac{\sqrt{2}}{2^n} \left(\sum_{k=0}^{n+1} \binom{n}{k} z^{-k} \right) \Big|_{z \triangleq e^{i\omega}} && \text{by binomial theorem} \\
\iff h_k &= \frac{\sqrt{2}}{2^{n+1}} \binom{n}{k} && \text{by definition of } Z \text{ transform (Definition P.4 page 330)}
\end{aligned}$$

3. Proof that (3) \implies (4):

$$\begin{aligned}
\tilde{h}(\omega) &= \check{h}(z) \Big|_{z \triangleq e^{i\omega}} && \text{by definition of DTFT (Definition O.1 page 319)} \\
&= \frac{\sqrt{2}}{2^n} (1 + z^{-1})^{n+1} \Big|_{z \triangleq e^{i\omega}} && \text{by (3)} \\
&= \frac{\sqrt{2}}{2^n} (1 + e^{-i\omega})^{n+1} && \text{by definition of } z \\
&= \frac{\sqrt{2}}{2^n} \left[e^{-i\frac{1}{2}\omega} \left(e^{i\frac{\omega}{2}} + e^{-i\frac{\omega}{2}} \right) \right]^{n+1} \\
&= \frac{\sqrt{2}}{2^n} e^{-i\frac{n+1}{2}\omega} \left[2\cos\left(\frac{\omega}{2}\right) \right]^{n+1} \\
&= 2\sqrt{2} e^{-i\frac{n+1}{2}\omega} \left[\cos\left(\frac{\omega}{2}\right) \right]^{n+1}
\end{aligned}$$

4. Proof that (3) \iff (4):

$$\begin{aligned}
\check{h}(z) \Big|_{z \triangleq e^{i\omega}} &= \check{h}(e^{i\omega}) \\
&= \tilde{h}(\omega) \\
&= 2\sqrt{2} e^{-i\frac{n+1}{2}\omega} \left[\cos\left(\frac{\omega}{2}\right) \right]^{n+1} && \text{by (4)} \\
&= \frac{\sqrt{2}}{2^n} e^{-i\frac{n+1}{2}\omega} \left[2\cos\left(\frac{\omega}{2}\right) \right]^{n+1} \\
&= \frac{\sqrt{2}}{2^n} \left[e^{-i\frac{1}{2}\omega} \left(e^{i\frac{\omega}{2}} + e^{-i\frac{\omega}{2}} \right) \right]^{n+1} \\
&= \frac{\sqrt{2}}{2^n} (1 + e^{-i\omega})^{n+1} \\
&= \frac{\sqrt{2}}{2^n} (1 + z^{-1})^{n+1} \Big|_{z \triangleq e^{i\omega}}
\end{aligned}$$

Example Q.20 (2 coefficient case). ²⁸ Let $(L_{\mathbb{R}}^2, (\mathcal{V}_j), \phi, (h_n))$ be an MRA system (Definition ?? page ??).

E X

$$\left\{ \begin{array}{l} 1. \text{ supp}\phi(x) = [0 : 1] \quad \text{and} \\ 2. (\phi(x - k)) \text{ forms a} \\ \text{partition of unity} \end{array} \right\} \xleftrightarrow{(A)} h_n = \left\{ \begin{array}{ll} \frac{\sqrt{2}}{2} & \text{for } n = 0 \\ \frac{\sqrt{2}}{2} & \text{for } n = 1 \\ 0 & \text{otherwise} \end{array} \right\} \xleftrightarrow{(B)} \underbrace{\{\phi(x) = N_0(x)\}}_{(C)}$$

PROOF:

1. Proof that (A) \implies (B):

- (a) lemma: Only h_0 and h_1 are *non-zero*; All other coefficients h_k are 0. Proof: This follows from $\text{supp}\phi(x) = [0 : 1]$ (Definition ?? page ??) and by Theorem ?? page ??.
- (b) lemma (equations for (h_k)): Because (h_k) is a *scaling coefficient sequence* (Definition ?? page ??), it must satisfy the *admissibility equation* (Theorem ?? page ??). And because $(\phi(x - k))$ forms a *partition of unity*, it must satisfy the equations given by Theorem ?? (page ??). (1a) lemma and these two constraints yield two simultaneous equations and two unknowns:

$$\begin{aligned} h_0 + h_1 &= \sqrt{2} && \text{(admissibility condition)} \\ h_0 - h_1 &= 0 && \text{(partition of unity/zero at } -1 \text{ / vanishing 0th moment)} \end{aligned}$$

- (c) lemma: The equations provided by (1b) lemma can be expressed in matrix algebra form as follows...

$$\underbrace{\begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}}_A \begin{bmatrix} h_0 \\ h_1 \end{bmatrix} = \begin{bmatrix} \sqrt{2} \\ 0 \end{bmatrix}$$

- (d) lemma: The *inverse A*⁻¹ of A can be expressed as demonstrated below...

$$\begin{aligned} \left[\begin{array}{cc|cc} 1 & 1 & 1 & 0 \\ 1 & -1 & 0 & 1 \end{array} \right] &\rightarrow \left[\begin{array}{cc|cc} 2 & 0 & 1 & 1 \\ 1 & -1 & 0 & 1 \end{array} \right] \rightarrow \left[\begin{array}{cc|cc} 2 & 0 & 1 & 1 \\ 0 & -1 & -\frac{1}{2} & \frac{1}{2} \end{array} \right] \rightarrow \left[\begin{array}{cc|cc} 1 & 0 & \frac{1}{2} & \frac{1}{2} \\ 0 & 1 & \frac{1}{2} & -\frac{1}{2} \end{array} \right] \\ \implies A^{-1} &= \frac{1}{2} \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} \end{aligned}$$

- (e) Proof for the values of (h_k) (B):

$$\begin{aligned} \begin{bmatrix} h_0 \\ h_1 \end{bmatrix} &= A^{-1} A \begin{bmatrix} h_0 \\ h_1 \end{bmatrix} = A^{-1} \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} \begin{bmatrix} h_0 \\ h_1 \end{bmatrix} && \text{by (1c) lemma} \\ &= A^{-1} \begin{bmatrix} \sqrt{2} \\ 0 \end{bmatrix} && \text{by (1c) lemma} \\ &= \frac{1}{2} \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} \begin{bmatrix} \sqrt{2} \\ 0 \end{bmatrix} && \text{by (1d) lemma} \\ &= \frac{\sqrt{2}}{2} \begin{bmatrix} 1 \\ 1 \end{bmatrix} \end{aligned}$$

²⁸ [Haar \(1910\)](#), [Wojtaszczyk \(1997\)](#) pages 14–15 (“Sources and comments”)

2. Proof that (B) \implies (C):

$$\begin{aligned}
 (B) \implies \phi(x) &= \sum_{k \in \mathbb{Z}} h_k \sqrt{2} \phi(2x - k) && \text{dilation equation} \\
 &= \sum_{k=0}^{k=1} \left(\frac{\sqrt{2}}{2} \right) \sqrt{2} \phi(2x - k) && \text{by item (1e) page 372} \\
 &= \sum_{k=0}^{k=1} \phi(2x - k) \\
 &= \sum_{k=0}^{k=1} \binom{1}{k} \phi(2x - k) && \text{by definition of } \binom{n}{k} \\
 \implies (D) &&& \text{by } B\text{-spline dyadic decomposition} \quad (\text{Theorem Q.9 page 367})
 \end{aligned}$$

3. Proof that (B) \Leftarrow (C):

$$\begin{aligned}
 (C) \implies N_0(x) &= \sum_{k=0}^{k=1} \binom{1}{k} N_0(2x - k) && \text{by } B\text{-spline dyadic decomposition} \quad (\text{Theorem Q.9 page 367}) \\
 &= \sum_{k=0}^{k=1} \left(\frac{\sqrt{2}}{2} \right) \sqrt{2} N_0(2x - k) && \text{by definition of } \binom{n}{k} \\
 &= \sum_{k \in \mathbb{Z}} h_k \sqrt{2} N_0(2x - k) && \text{by definition of } \binom{n}{k} \\
 \implies (B) &&&
 \end{aligned}$$

4. Proof that (A) \Leftarrow (C):

1. Proof that (C) $\implies \text{supp } \phi(x) = [0 : 1]$: by Theorem Q.4 (page 354)
2. Proof that (C) $\implies (\phi(x - k))$ forms a *partition of unity*: by Theorem Q.7 (page 360)



E X	n=0,	(÷0!)	1;						
	n=1,	(÷1!)	1, 0; -1, 2;						
	n=2,	(÷2!)	1, 0, 0; -2, 6, -3; 1, -6, 9;						
	n=3,	(÷3!)	1, 0, 0, 0; -3, 12, -12, 4; 3, -24, 60, -44; -1, 12, -48, 64;						
	n=4,	(÷4!)	1, 0, 0, 0, 0; -4, 20, -30, 20, -5; 6, -60, 210, -300, 155; -4, 60, -330, 780, -655; 1, -20, 150, -500, 625;						
	n=5,	(÷5!)	1, 0, 0, 0, 0, 0; -5, 30, -60, 60, -30, 6; 10, -120, 540, -1140, 1170, -474; -10, 180, -1260, 4260, -6930, 4386; 5, -120, 1140, -5340, 12270, -10974; -1, 30, -360, 2160, -6480, 7776;						
	n=6,	(÷6!)	1, 0, 0, 0, 0, 0, 0; -6, 42, -105, 140, -105, 42, -7; 15, -210, 1155, -3220, 4935, -3990, 1337; -20, 420, -3570, 15680, -37590, 47040, -24178; 15, -420, 4830, -29120, 96810, -168000, 119182; -6, 210, -3045, 23380, -100065, 225750, -208943; 1, -42, 735, -6860, 36015, -100842, 117649;						
	n=7,	(÷7!)	1, 0, 0, 0, 0, 0, 0, 0; -7, 56, -168, 280, -280, 168, -56, 8; 21, -336, 2184, -7560, 15400, -18648, 12488, -3576; -35, 840, -8400, 45360, -143360, 267120, -273280, 118896; 35, -1120, 15120, -111440, 483840, -1238160, 1733760, -1027984; -21, 840, -14280, 133560, -741160, 2436840, -4391240, 3347016; 7, -336, 6888, -78120, 528920, -2135448, 4753336, -4491192; -1, 56, -1344, 17920, -143360, 688128, -1835008, 2097152;						
	n=8,	(÷8!)	1, 0, 0, 0, 0, 0, 0, 0, 0; -8, 72, -252, 504, -630, 504, -252, 72, -9; 28, -504, 3780, -15624, 39690, -64008, 64260, -36792, 9207; -56, 1512, -17388, 111384, -436590, 1079064, -1650348, 1432872, -541917; 70, -2520, 39060, -340200, 1821330, -6146280, 12800340, -15082200, 7715619; -56, 2520, -49140, 541800, -3691170, 15903720, -42324660, 63667800, -41503131; 28, -1512, 35532, -474264, 3929310, -20674584, 67410252, -124449192, 99584613; -8, 504, -13860, 217224, -2121210, 13208328, -51179940, 112731192, -107948223; 1, -72, 2268, -40824, 459270, -3306744, 14880348, -38263752, 43046721						

Table Q.1: Coefficients of the *B-splines* $N_n(x)$ multiplied by $n!$ (Example Q.9 page 350)

APPENDIX R

SOURCE CODE

The free and open source software package Maxima has been used to compute some of the algebraic expressions for *B-splines* used in APPENDIX Q (page 343):

```
1 /*=====
2 * Daniel J. Greenhoe
3 * Maxima script file
4 * To execute this script, start Maxima in a command window
5 * in the subdirectory containing this file (e.g. c:\math\maxima\
6 * and then after the (%i...) prompt enter
7 * batchload("bspline.max")$
8 * Data produced will be written to the file "bsplineout.txt".
9 * reference: http://maxima.sourceforge.net/documentation.html
10 */
11 /*
12 * initialize script
13 */
14 reset();
15 kill(all);
16 load(orthopoly);
17 display2d:false; /* 2-dimensional display */
18 writefile ("bsplineout.txt");
19 /*
20 * n = B-spline order parameter
21 * may be set to any value in {1,2,3,...}
22 */
23 n:2;
24 print("=====");
25 print("Daniel J. Greenhoe");
26 print("Output file for nth order B-spline Nn(x) calculation, n=",n," .");
27 print("Output produced using Maxima running the script file bspline.max");
28 print("=====");
29 Nnx:(1/n!)*sum((-1)^k*binomial(n+1,k)*(x-k)^n*unit_step(x-k),k,0,n+1);
30 print("=====");
31 print("      n+1      k (n+1)      n      ");
32 print("      n! Nn(x) = SUM (-1) ( ) (x-k)  step(x-k) ,n=",n," ");
33 print("      k=0      ( k )      ");
34 print("      ,n+1,      k ( ,n+1,)      ,n);
35 print(n,"! Nn(x) = SUM (-1) ( ) (x-k)  step(x-k)");
36 print("      k=0      ( k )");
37 print("      = ",expand(n!*Nnx));
38 print("=====");
39 assume(x<=0);   print(n!,"N(x)= ",expand(n!*Nnx)," for x<=0");   forget(x<=0);
40 for i:0 thru n step 1 do(
41   assume(x>i,x<(i+1)),
42   print(n!,"N(x)= ",expand(n!*Nnx)," for ",i,"<x<",i+1),
43   tex(expand(n!*Nnx),"djh.tex"),/*write output in TeX format to file "djh.tex"*/
44   forget(x>i,x<(i+1))
45 );
46 assume(x>(n+1)); print(n!,"N(x)= ",expand(n!*Nnx)," for x>",n+1); forget(x>(n+1));
```

```

47 print("-----");
48 print(" values at some specific points x:           ");
49 print("-----");
50 y:Nnx,x=(n+1)/2;print("N(",(n+1)/2,")= ",y," (center value)");
51 y:Nnx,x=(n+2)/2;print("N(",(n+2)/2,")= ",y);
52 y:Nnx,x=(n+3)/2;print("N(",(n+3)/2,")= ",y);
53 /*-----*/
54 * close output file
55 *-----*/
56 closefile();

```

Once the polynomial expressions for a *B-spline* have been calculated, they can be plotted within a \LaTeX environment using the [pst-plot package](#) along with a \LaTeX source file such as the following:¹

```

1 %=====
2 % Daniel J. Greenhoe
3 % LaTeX file
4 % N_3(x) B-spline
5 % nominal unit = 10mm
6 %=====
7 \begin{pspicture}(-1,-0.5)(5,1)
8 %
9 % parameters
10 %
11 \psset{plotpoints=64,labelsep=1pt}
12 %
13 % axes
14 %
15 \psaxes[linewidth=0.75pt, linecolor=axis ,yAxis=false ,ticks=x, labels=x]{<->}(0,0)(-1,0)(5,1)% x axis
16 \psaxes[linewidth=0.75pt, linecolor=axis ,xAxis=false ,ticks=x, labels=x]{->}(0,0)(-1,0)(5,1)% y axis
17 %
18 % annotation
19 %
20 \psline [linestyle=dashed, linewidth=0.75pt, linecolor=red](2,0)(2,0.667)%  

21 \psline [linestyle=dashed, linewidth=0.75pt, linecolor=red](0,0.667)(2,0.667)%  

22 \psline [linestyle=dashed, linewidth=0.75pt, linecolor=red](1,0)(1,0.1667)%  


23 \psline [linestyle=dashed, linewidth=0.75pt, linecolor=red](3,0)(3,0.1667)%  

24 \psline [linestyle=dashed, linewidth=0.75pt, linecolor=red](0,0.1667)(3,0.1667)%  


25 \uput[180](0,0.667){$\frac{2}{3}$}  

26 \uput[180](0,0.1667){$\frac{1}{6}$}
27 %
28 % function plot
29 %
30 \psplot{0}{1}{+1 x 3 exp mul}                                6 div% for 0<=x<=1
31 \psplot{1}{2}{-3 x 3 exp mul +12 x 2 exp mul add -12 x mul add +4 add 6 div}% for 1<=x<=2
32 \psplot{2}{3}{+3 x 3 exp mul -24 x 2 exp mul add +60 x mul add -44 add 6 div}% for 2<=x<=3
33 \psplot{3}{4}{-1 x 3 exp mul +12 x 2 exp mul add -48 x mul add +64 add 6 div}% for 3<=x<=4
34 \psline(0,0)(-0.5,0)\psline[linestyle=dotted](-0.5,0)(-0.75,0)%          % for x<=0
35 \psline(4,0)(4.5,0)\psline[linestyle=dotted](4.5,0)(4.75,0)%          % for x>=4
36 \end{pspicture}

```



Alternatively, one can plot $N_3(x)$ more or less directly from the equation given in Theorem Q.1 (page 347) without first calculating the polynomial expressions:



```

1 %=====
2 % Daniel J. Greenhoe
3 % LaTeX file
4 % N_3(x) B-spline
5 % nominal unit = 10mm
6 %=====
7 \begin{pspicture}(-1,-0.5)(5,1)
8 %
9 % parameters
10 %
11 \psset{plotpoints=64,labelsep=1pt}

```

¹For help with PostScript®math operators, see [Adobe \(1999\)](#), pages 508–509 (Arithmetic and Math Operators).

```

12 %
13 % axes
14 %
15 \psaxes[linewidth=0.75pt, linecolor=axis, yAxis=false, ticks=x, labels=x]{<->}(0,0)(-1,0)(5,1)% x axis
16 \psaxes[linewidth=0.75pt, linecolor=axis, xAxis=false, ticks=x, labels=x]{->}(0,0)(-1,0)(5,1)% y axis
17 %
18 % annotation
19 %
20 \psline[linestyle=dashed, linewidth=0.75pt, linecolor=red](2,0)(2,0.667)%
21 \psline[linestyle=dashed, linewidth=0.75pt, linecolor=red](0,0.667)(2,0.667)%
22 \psline[linestyle=dashed, linewidth=0.75pt, linecolor=red](1,0)(1,0.1667)%
23 \psline[linestyle=dashed, linewidth=0.75pt, linecolor=red](3,0)(3,0.1667)%
24 \psline[linestyle=dashed, linewidth=0.75pt, linecolor=red](0,0.1667)(3,0.1667)%
25 \uput[180](0,0.667){$\frac{2}{3}$}%
26 \uput[180](0,0.1667){$\frac{1}{6}$}%
27 %
28 % for n=3
29 % 
$$\frac{1}{n!} \sum_{k=1}^{n+1} (-1)^k \binom{n+1}{k} (x-k)^n s(x-k) = \frac{1}{3!} \sum_{k=1}^4 (-1)^k \binom{4}{k} (x-k)^3 s(x-k)$$

30 % where  $s(x) = 0$  for  $x < 0$  and  $1$  for  $x \geq 0$  (step function)
31 %
32 %
33 \psplot{0}{1}{1 x 0 sub 3 exp mul 6 div}%
34 \psplot{1}{2}{1 x 0 sub 3 exp mul 4 x 1 sub 3 exp mul sub 6 div}%
35 \psplot{2}{3}{1 x 0 sub 3 exp mul 4 x 1 sub 3 exp mul sub 6 x 2 sub 3 exp mul add 6 div}%
36 \psplot{3}{4}{1 x 0 sub 3 exp mul 4 x 1 sub 3 exp mul sub 6 x 2 sub 3 exp mul add 4 x 3 sub
   3 exp mul sub 6 div}%
37 \psplot{4}{4.5}{1 x 0 sub 3 exp mul 4 x 1 sub 3 exp mul sub 6 x 2 sub 3 exp mul add 4 x 3 sub
   3 exp mul sub 1 x 4 sub 3 exp mul add 6 div}%
38 %
39 % 
$$N_3(x) = \frac{[(4\text{choose}0)(x-0)^3 - (4\text{choose}1)(x-1)^3 + (4\text{choose}2)(x-2)^3 - (4\text{choose}3)(x-3)^3 + (4\text{choose}4)(x-4)^3]/3!}{6}$$

40 % 
$$= \frac{1}{6} [(x-0)^3 - 4(x-1)^3 + 6(x-2)^3 - (x-3)^3 + (x-4)^3]$$

41 %
42 \psline(0,0)(-0.5,0)%
43 \psline[linestyle=dotted](-0.5,0)(-0.75,0)%
44 \psline[linestyle=dotted](4.5,0)(4.75,0)%
45 \end{pspicture}%

```


Back Matter




“It appears to me that if one wants to make progress in mathematics, one should study the masters and not the pupils.”

Niels Henrik Abel (1802–1829), Norwegian mathematician ²




“When evening comes, I return home and go to my study. On the threshold I strip naked, taking off my muddy, sweaty workaday clothes, and put on the robes of court and palace, and in this graver dress I enter the courts of the ancients and am welcomed by them, and there I taste the food that alone is mine, and for which I was born. And there I make bold to speak to them and ask the motives of their actions, and they, in their humanity reply to me. And for the space of four hours I forget the world, remember no vexation, fear poverty no more, tremble no more at death; I pass indeed into their world.”

Niccolò Machiavelli (1469–1527), Italian political philosopher, in a 1513 letter to friend Francesco Vettori. ³



ancient library of Alexandria



The Book Worm by Carl Spitzweg, circa 1850



“To sit alone in the lamplight with a book spread out before you, and hold intimate converse with men of unseen generations—such is a pleasure beyond compare.”

Yoshida Kenko (Urabe Kaneyoshi) (1283? – 1350?), Japanese author and Buddhist monk ⁵

² quote: [Simmons \(2007\)](#), page 187.

image: http://en.wikipedia.org/wiki/Image:Niels_Henrik_Abel.jpg, public domain

³ quote: [Machiavelli \(1961\)](#), page 139?.

image: http://commons.wikimedia.org/wiki/File:Santi_di_Tito_-_Niccolo_Machiavelli%27s_portrait_headcrop.jpg, public domain

⁴ <http://en.wikipedia.org/wiki/File:Ancientlibraryalex.jpg>, public domain http://en.wikipedia.org/wiki/File:Carl_Spitzweg_021.jpg

⁵ quote: [Kenko \(circa 1330\)](#)

image: http://en.wikipedia.org/wiki/Yoshida_Kenko



BIBLIOGRAPHY

- M. Abom. A note on random errors in frequency response estimators. *Journal of Sound and Vibration*, 107(2):355–358, 1986. doi: 10.1016/0022-460X(86)90244-0. URL <http://adsabs.harvard.edu/abs/1986JSV...107..355A>.
- Yuri A. Abramovich and Charalambos D. Aliprantis. *An Invitation to Operator Theory*. American Mathematical Society, Providence, Rhode Island, 2002. ISBN 0-8218-2146-6. URL <http://books.google.com/books?vid=ISBN0821821466>.
- Adobe. *PostScript Language Reference*. Addison-Wesley Publishing Company, Reading, Massachusetts, 1999. ISBN 0-201-37922-8. URL <http://wwwimages.adobe.com/content/dam/Adobe/en/devnet/postscript/pdfs/PLRM.pdf>.
- N. I. Akhiezer and I. M. Glazman. *Theory of Linear Operators in Hilbert Spaces*, volume 1. Dover, New York, 1993. URL <http://books.google.com/books?vid=ISBN0486677486>. Translated from the original Russian text *Teoriia lineinykh operatorov v Gil'bertovom prostranstve*.
- Charalambos D. Aliprantis and Owen Burkinshaw. *Principles of Real Analysis*. Academic Press, London, 3 edition, 1998. ISBN 9780120502578. URL <http://www.amazon.com/dp/0120502577>.
- Charalambos D. Aliprantis and Owen Burkinshaw. *Positive Operators*. Springer, Dordrecht, 2006. ISBN 9781402050077. URL <http://books.google.com/books?vid=ISBN1402050070>. reprint of Academic Press 1985 edition.
- Randall J. Allemand, R. Zimmerman, and David L. Brown. Techniques of reducing noise in frequency response measurements. *ASME Winter Annual Meeting*, 1979. session on Application of System Identification Techniques.
- Randall J. Allemand, David L. Brown, and Robert W. Rost. *Experimental Modal Analysis And Dynamic Component Synthesis*, 2(AFWAL-TR-87-3069), December 1987. URL <https://apps.dtic.mil/dtic/tr/fulltext/u2/a195145.pdf>.
- Theodore T. Allen. *Introduction to Engineering Statistics and Lean Six Sigma: Statistical Quality Control and Design of Experiments and Systems*. Springer, 3 edition, 2018. ISBN 9781447174202.
- Herbert Amann and Joachim Escher. *Analysis II*. Birkhäuser Verlag AG, Basel–Boston–Berlin, 2008. ISBN 978-3-7643-7472-3. URL <http://books.google.com/books?vid=ISBN3764374721>.
- Dan Amir. *Characterizations of Inner Product Spaces*, volume 20 of *Operator Theory: Advances and Applications*. Birkhäuser Verlag, Basel, 1986. ISBN 3-7643-1774-4. URL <http://www.worldcat.org/isbn/3764317744>.

- John E. Angus. The probability integral transform and related results. *SIAM Review*, 36(4):652–654, February 25 1994. doi: 10.1137/1036146. URL <http://dx.doi.org/10.1137/1036146>.
- Tom M. Apostol. *Mathematical Analysis*. Addison-Wesley series in mathematics. Addison-Wesley, Reading, 2 edition, 1975. ISBN 986-154-103-9. URL <http://books.google.com/books?vid=ISBN0201002884>.
- Jean-Pierre Aubin. *Applied Functional Analysis*, volume 47 of *Pure and Applied Mathematics: A Wiley Series of Texts, Monographs and Tracts*. John Wiley & Sons, 2 edition, September 30 2011. ISBN 9781118030974. URL <http://books.google.com/books?vid=ISBN1118030974>.
- Jean-Pierre Aubin and Hélène Frankowska. *Set-Valued Analysis*. Modern Birkhäuser Classics. Springer, March 2 2009. ISBN 9780817648480. URL <http://books.google.com/books?vid=ISBN0817648488>.
- Léon Autonne. Sur l'hermitien (on the hermitian). In *Comptes Rendus Des Séances De L'Académie Des Sciences*, volume 133, pages 209–268. De L'Académie des sciences (Academy of Sciences), Paris, 1901. URL <http://visualiseur.bnf.fr/Visualiseur?O=NUMM-3089>. Comptes Rendus Des Séances De L'Académie Des Sciences (Reports Of the Meetings Of the Academy of Science).
- Léon Autonne. Sur l'hermitien (on the hermitian). *Rendiconti del Circolo Matematico di Palermo*, 16:104–128, 1902. Rendiconti del Circolo Matematico di Palermo (Statements of the Mathematical Circle of Palermo).
- George Bachman. *Elements of Abstract Harmonic Analysis*. Academic paperbacks. Academic Press, New York, 1964. URL <http://books.google.com/books?id=ZP8-AAAAIAAJ>.
- George Bachman and Lawrence Narici. *Functional Analysis*. Academic Press textbooks in mathematics; Pure and Applied Mathematics Series. Academic Press, 1 edition, 1966. ISBN 9780486402512. URL <http://books.google.com/books?vid=ISBN0486402517>. “unabridged republication” available from Dover (isbn 0486402517).
- George Bachman, Lawrence Narici, and Edward Beckenstein. *Fourier and Wavelet Analysis*. Universitext Series. Springer, 2000. ISBN 9780387988993. URL <http://books.google.com/books?vid=ISBN0387988998>.
- David H. Bailey, Jonathan M. Borwein, and Andrew Mattingly. The computation of previously inaccessible digits of π^2 and catalan's constant (2013). page 23, March 24 2011. URL <https://pdfs.semanticscholar.org/0b15/22a1b2f76729e19be2be1bca9f57031a4db9.pdf>.
- David H. Bailey, Jonathan M. Borwein, and Andrew Mattingly. 20. the computation of previously inaccessible digits of π^2 and catalan's constant (2013). In *Pi: The Next Generation: A Sourcebook on the Recent History of Pi and Its Computation*, pages 324–339. Springer, 2013. ISBN 9783319323770. URL <http://books.google.com/books?vid=ISBN3319323776>.
- Per Bak. *How Nature Works: the science of self-organized criticality*. Springer Science & Business Media, 2013. ISBN 9781475754261. URL <http://books.google.com/books?vid=ISBN1475754264>.
- N Balakrishnan and Chin-Diew Lai. *Continuous Bivariate Distributions*. Springer Science & Business Media, 2 edition, May 31 2009. ISBN 9780387096148. URL <http://books.google.com/books?vid=ISBN0387096140>.
- Stefan Banach. Sur les opérations dans les ensembles abstraits et leur applications aux équations intégrales (on abstract operations and their applications to the integral equations). *Fundamenta Mathematicae*, 3:133–181, 1922. URL <http://matwbn.icm.edu.pl/ksiazki/fm/fm3/fm3120.pdf>.

- Stefan Banach. *Théorie des opérations linéaires*. Monografje Matematyczne, Warsaw, Poland, 1932a. URL <http://matwbn.icm.edu.pl/kstrecs.php?tom=1&wyd=10>. (Theory of linear operations).
- Stefan Banach. *Theory of Linear Operations*, volume 38 of *North-Holland mathematical library*. North-Holland, Amsterdam, 1932b. ISBN 0444701842. URL <http://www.amazon.com/dp/0444701842/>. English translation of 1932 French edition, published in 1987.
- Alexander Barvinok. *A Course in Convexity*, volume 54 of *Graduate studies in mathematics*. American Mathematical Society, 2002. ISBN 9780821872314. URL <http://books.google.com/books?vid=ISBN0821872311>.
- Gerald Beer. *Topologies on Closed and Closed Convex Sets*, volume 268 of *Mathematics and Its Applications*. October 31 1993. ISBN 9780792325314. URL <http://books.google.com/books?vid=ISBN0792325311>.
- J. S. Bendat. Statistical errors in measurement of coherence functions and input/output quantities. *Journal of Sound and Vibration*, 59(3):405–421, August 08 1978. doi: 10.1016/S0022-460X(78)80007-8. URL [https://doi.org/10.1016/S0022-460X\(78\)80007-8](https://doi.org/10.1016/S0022-460X(78)80007-8).
- Julius S. Bendat and Allan G. Piersol. *Engineering Applications of Correlation and Spectral Analysis*. John Wiley & Sons, 1980. ISBN 9780471058878. URL <http://www.amazon.com/dp/0471058874>.
- Julius S. Bendat and Allan G. Piersol. *Engineering Applications of Correlation and Spectral Analysis*. Wiley-Interscience, 2 edition, 1993. ISBN 9780471570554. URL <http://www.amazon.com/dp/0471570559>.
- Julius S. Bendat and Allan G. Piersol. *Random Data: Analysis and Measurement Procedures*, volume 729 of *Wiley Series in Probability and Statistics*. John Wiley & Sons, 4 edition, 2010. ISBN 9781118210826. URL <http://books.google.com/books?vid=ISBN1118210824>.
- Sterling Khazag Berberian. *Introduction to Hilbert Space*. Oxford University Press, New York, 1961. URL <http://books.google.com/books?vid=ISBN0821819127>.
- D. Besso. Teoremi elementari sui massimi i minimi. *Annuari Ist. Tech. Roma*, pages 7–24, 1879. see Bullen(2003) pages 453, 203.
- M. Jales Bienaymé. Société philomatique de paris—extraits des procès-verbaux. *Scéance*, pages 67–68, June 13 1840. URL <http://www.archive.org/details/extratsdesproc46183941soci>. see Bullen(2003) pages 453, 203.
- Patrick Billingsley. *Probability And Measure*. Wiley series in probability and mathematical statistics. Wiley, 3 edition, 1995. ISBN 978-0471007104. URL <http://books.google.com/books?vid=ISBN0471007102>.
- Leonard Mascot Blumenthal. *Theory and Applications of Distance Geometry*. Chelsea Publishing Company, Bronx, New York, USA, 2 edition, 1970. ISBN 0-8284-0242-6. URL <http://books.google.com/books?vid=ISBN0828402426>.
- Béla Bollobás. *Linear Analysis; an introductory course*. Cambridge mathematical textbooks. Cambridge University Press, Cambridge, 2 edition, March 1 1999. ISBN 978-0521655774. URL <http://books.google.com/books?vid=ISBN0521655773>.
- V. Bouniakowsky. Sur quelques inégalités concernant les intégrales ordinaires et les intégrales aux différences finies. *Mémoires de l'Acad. de St.-Pétersbourg*, 1(9), 1859. URL http://www-stat.wharton.upenn.edu/~steele/Publications/Books/CSMC/CSMC_index.html.

- Carl Benjamin Boyer and Uta C. Merzbach. *A History of Mathematics*. Wiley, New York, 3 edition, January 11 2011. ISBN 0470525487. URL <http://books.google.com/books?vid=ISBN0470525487>.
- Ronald Newbold Bracewell. *The Fourier transform and its applications*. McGraw-Hill electrical and electronic engineering series. McGraw-Hill, 2, illustrated, international student edition edition, 1978. ISBN 9780070070134. URL <http://books.google.com/books?vid=ISBN007007013X>.
- Thomas John I'Anson Bromwich. *An Introduction to the Theory of Infinite Series*. Macmillan and Company, 1 edition, 1908. ISBN 9780821839768. URL <http://www.archive.org/details/anintroduction00bromgoog>.
- P. S. Bullen. *Handbook of Means and Their Inequalities*, volume 560 of *Mathematics and Its Applications*. Kluwer Academic Publishers, Dordrecht, Boston, 2 edition, 2003. ISBN 9781402015229. URL <http://books.google.com/books?vid=ISBN1402015224>.
- Viktor Yakovlevich Bunyakovsky. Sur quelques inégalités concernant les intégrales ordinaires et les intégrales aux différences finies. *Mémoires de L'Académie Impériale des Sciences de St.-Pétersbourg*, 1(9):1–18, 1859. URL <http://www-stat.wharton.upenn.edu/~steele/Publications/Books/CSMC/bunyakovsky.pdf>. (Mémoires of the Imperial Academy of Sciences of Saint Petersburg); (On Some Inequalities concerning the Ordinary Integrals and Integrals with the Finite Differences).
- Charles L. Byrne. *Signal Processing: A Mathematical Approach*. AK Peters Series. A K Peters, 2005. ISBN 1568812426.
- James A. Cadzow. *Foundations of Digital Signal Processing and Data Analysis*. Macmillan Publishing Company, New York, 1987. ISBN 0023180102. URL http://frontweb.vuse.vanderbilt.edu/vuse_web/directory/facultybio.asp?FacultyID=9.
- Peter E. Caines. *Linear Stochastic Systems*, volume 100 of *Wiley Series in Probability and Statistics*. J. Wiley, 1988. ISBN 9780471081012. URL <http://books.google.com/books?vid=ISBN0471081019>.
- Peter E. Caines. *Linear Stochastic Systems*, volume 77 of *Classics in Applied Mathematics*. SIAM, 2018. ISBN 9781611974713. URL <http://books.google.com/books?vid=ISBN1611974712>.
- Florian Cajori. A history of mathematical notations; notations mainly in higher mathematics. In *A History of Mathematical Notations; Two Volumes Bound as One*, volume 2. Dover, Mineola, New York, USA, 1993. ISBN 0-486-67766-4. URL <http://books.google.com/books?vid=ISBN0486677664>. reprint of 1929 edition by *The Open Court Publishing Company*.
- James J. Callahan. *Advanced Calculus: A Geometric View*. Springer Science & Business Media, September 9 2010. ISBN 9781441973320. URL <http://books.google.com/books?vid=ISBN144197332X>.
- Thomas G. Carne and C. R. Dohrmann. Improving experimental frequency response function matrices for admittance modeling. In *Proceedings of the XXIV International Modal Analysis Conference (IMAC)*, St. Louis, Missouri, USA, 2006. URL <https://www.researchgate.net/publication/265790751>.
- N. L. Carothers. *A Short Course on Banach Space Theory*. Number 64 in London Mathematical Society Student Texts. Cambridge University Press, Cambridge, 2005. ISBN 0521842832. URL <http://www.comm.musashi-tech.ac.jp/~arimoto/Drarimoto/preprints/calothers/Assemble.ps>.

- N.L. Carothers. *Real Analysis*. Cambridge University Press, Cambridge, 2000. ISBN 978-0521497565. URL <http://books.google.com/books?vid=ISBN0521497566>.
- Augustin-Louis Cauchy. *Part 1: Analyse Algebrique*. Cours D'Analyse de L'école Royale Polytechnique. Debure frères, Paris, 1821. ISBN 2-87647-053-5. URL <http://www.archive.org/details/coursanalyse00caucgoog>. Systems design course of the Polytechnic Royal School; 1st Part: Algebraic analysis).
- Arthur Cayley. A memoir on the theory of matrices. *Philosophical Transactions of the Royal Society of London*, 148:17–37, 1858. ISSN 1364-503X. URL <http://www.jstor.org/view/02610523/ap000059/00a00020/0>.
- Si-Wei Chen, Xue-Song Wang, and Motoyuki Sato. Polinsar complex coherence estimation based on covariance matrix similarity test. *IEEE Transactions on Geoscience and Remote Sensing*, 50(11):4699–4710, November 2012. doi: 10.1109/TGRS.2012.2192937. URL https://www.researchgate.net/profile/Si_Wei_Chen/publication/281271737.
- Alexandre J. Chorin and Ole H. Hald. *Stochastic Tools in Mathematics and Science*, volume 1 of *Surveys and Tutorials in the Applied Mathematical Sciences*. Springer, New York, 2 edition, 2009. ISBN 978-1-4419-1001-1. URL <http://books.google.com/books?vid=ISBN9781441910011>.
- Ole Christensen. *An Introduction to Frames and Riesz Bases*. Applied and Numerical Harmonic Analysis. Birkhäuser, Boston/Basel/Berlin, 2003. ISBN 0-8176-4295-1. URL <http://books.google.com/books?vid=ISBN0817642951>.
- Ole Christensen. *Frames and bases: An Introductory Course*. Applied and Numerical Harmonic Analysis. Birkhäuser, Boston/Basel/Berlin, 2008. ISBN 9780817646776. URL <http://books.google.com/books?vid=ISBN0817646779>.
- Ole Christensen. *Functions, Spaces, and Expansions: Mathematical Tools in Physics and Engineering*. Applied and Numerical Harmonic Analysis. Birkhäuser, Boston/Basel/Berlin, 2010. ISBN 978-0-8176-4979-1. doi: 10.1007/978-0-8176-4980-7. URL <http://books.google.com/books?vid=ISBN0817649794>.
- Charles K. Chui. *Multivariate Splines*, volume 54 of *CBMS-NSF regional conference series in applied mathematics*. SIAM, 1988. ISBN 1611970172. URL <http://books.google.com/books?vid=ISBN1611970172>.
- Charles K. Chui. *An Introduction to Wavelets*. Academic Press, San Diego, California, USA, January 3 1992. ISBN 9780121745844. URL <http://books.google.com/books?vid=ISBN0121745848>.
- Donato Michele Cifarelli and Eugenio Regazzini. De finetti's contribution to probability and statistics. *Statistical Science*, 11(4):253–282, 1996. URL <http://www.jstor.org/stable/2246020>.
- Erhan Çinlar and Robert J Vanderbei. *Real and Convex Analysis*. Undergraduate Texts in Mathematics. Springer, January 4 2013. ISBN 1461452570. URL <http://books.google.com/books?vid=ISBN1461452570>.
- Jon F. Claerbout. *Fundamentals of Geophysical Data Processing with Applications to Petroleum Prospecting*. International series in the earth and planetary sciences, Tab Mastering Electronics Series. McGraw-Hill, New York, 1976. ISBN 9780070111172. URL <http://sep.stanford.edu/sep/prof/>.
- James A. Clarkson. Uniformly convex spaces. *Transactions of the American Mathematical Society*, 40(3):396–414, December 1936. URL <http://www.jstor.org/stable/1989630>.

- R. E. Cobb and L. D. Mitchell. Estimation of uncorrelated content in experimentally measured frequency response functions using three measurement channels. *Mechanical Systems and Signal Processing*, 4(6):449–461, November 1990. URL [https://doi.org/10.1016/0888-3270\(90\)90045-M](https://doi.org/10.1016/0888-3270(90)90045-M).
- Richard E. Cobb. *Confidence Bands, Measurement Noise, and Multiple Input - Multiple Output Measurements Using the Three-Channel Frequency Response Function Estimator*. Ph.d. dissertation, May 1988. URL <http://hdl.handle.net/10919/53675>. “Dissertation submitted to the Faculty of the Virginia Polytechnic Institute and State University in partial fulfillment of the requirements for the degree of Doctor of Philosophy in Mechanical Engineering”.
- John B. Conway. *A Course in Functional Analysis*, volume 96 of *Graduate Texts in Mathematics*. Springer, 2 edition, 1990. ISBN 0-387-97245-5. URL <http://books.google.com/books?vid=ISBN0387972455>.
- R. Courant and D. Hilbert. *Methods of Mathematical Physics*, volume 1. Interscience Publishers, New York, 1930. URL <http://www.worldcat.org/isbn/0471504475>.
- T.M. Cover and Joy A. Thomas. *Elements of Information Theory*. John Wiley & Sons, Inc., New York, 1991. ISBN 0-471-06259-6. URL <http://www.amazon.com/dp/0471062596>.
- M. G. Cox. The numerical evaluation of b-splines. *IMA Journal of Applied Mathematics*, 10(2):134–149, 1972. doi: 10.1093/imamat/10.2.134. URL <http://imamat.oxfordjournals.org/content/10/2/134.abstract>.
- Guy L. Curry and Richard M. Feldman. *Manufacturing Systems Modeling and Analysis*. Technology & Engineering. Springer Science & Business Media, 2 edition, 2010. ISBN 9783642166181. URL <http://books.google.com/books?vid=ISBN3642166180>.
- Ingrid Daubechies. *Ten Lectures on Wavelets*. Society for Industrial and Applied Mathematics, Philadelphia, 1992. ISBN 0-89871-274-2. URL <http://www.amazon.com/dp/0898712742>.
- Ingrid Daubechies, A. Grossman, and Y. Meyer. Painless nonorthogonal expansions. *Journal of Mathematical Physics*, 27(5):1271–1283, May 1986. ISSN 0022-2488. URL <link.aip.org/link/?jmp/27/1271>.
- Mahlon Marsh Day. *Normed Linear Spaces*, volume 21 of *Ergebnisse der Mathematik und ihrer Grenzgebiete*. Springer-Verlag, Berlin, Heidelberg, New York, 3 edition, 1973. ISBN 0387061487. URL <http://books.google.com/books?id=UZ3vSAAACAAJ>.
- Bruno de Finetti. Sui passaggi al limite nel calcolo delle probabilità. *Rendiconti del Reale Istituto Lombardo di Scienze e Lettere*, 63:155–166, 1930a.
- Bruno de Finetti. A proposito dell'estensione del teorema delle probabilità totali alle classi numerabili. *Rendiconti del Reale Istituto Lombardo di Scienze e Lettere*, 63:901–905, 1930b.
- Bruno de Finetti. Ancora sull'estensione alle classi numerabili del teorema delle probabilità totali. *Rendiconti del Reale Istituto Lombardo di Scienze e Lettere*, 63:1063–1069, 1930c.
- Johan de Witt. *Elementa curvarum linearum (Elements of Curves)*. 1659. Elementa curvarum linearum was an appendix to de Witt's translation of of Descartes' *La Géométrie*.
- Carl R. de Boor. On calculating with b-splines. *Journal of Approximation Theory*, pages 50–62, 1972.

- Carl R. de Boor. *A Practical Guide to Splines*, volume 27 of *Applied Mathematical Sciences*. Springer, revised edition, 2001. ISBN 0387953663. URL <http://books.google.com/books?vid=ISBN0387953663>.
- F.M. Dekking, C. Kraaikamp, H.P. Lopuhaa, and L.E. Meester. *A Modern Introduction to Probability and Statistics: Understanding Why and How*. Springer Texts in Statistics. Springer Science & Business Media, 2006. ISBN 9781846281686. URL <http://books.google.com/books?vid=ISBN1846281687>.
- René Descartes. *La géométrie*. 1637. URL http://historical.library.cornell.edu/math/math_D.html.
- René Descartes. *Regulae ad directionem ingenii*. 1684a. URL http://www.fh-augsburg.de/~harsch/Chronologia/Lspost17/Descartes/des_re00.html.
- René Descartes. *Rules for Direction of the Mind*. 1684b. URL http://en.wikisource.org/wiki/Rules_for_the_Direction_of_the_Mind.
- René Descartes. *The Geometry of Rene Descartes*. Courier Dover Publications, June 1 1954. ISBN 0486600688. URL <http://books.google.com/books?vid=isbn0486600688>. originally published by Open Court Publishing, Chicago, 1925; translation of La géométrie.
- Luc Devroye. *Non-Uniform Random Variate Generation*. Springer-Verlag, New York, 1986. ISBN 0387963057. URL <http://www.eirene.de/Devroye.pdf>.
- Jean Alexandre Dieudonné. *Foundations of Modern Analysis*. Academic Press, New York, 1969. ISBN 1406727911. URL <http://books.google.com/books?vid=ISBN1406727911>.
- Hamid Drljević. On the representation of functionals and the stability of mappings in hilbert and banach spaces. In Themistocles M Rassias, editor, *Topics in mathematical analysis: a volume dedicated to the memory of A.L. Cauchy*, volume II of *Series in Pure Mathematics*, pages 231–245. World Scientific Publishing Company, 1989. ISBN 9971506661. URL <http://books.google.com/books?vid=ISBN9971506661>.
- R. J. Duffin and A. C. Schaeffer. A class of nonharmonic fourier series. *Transactions of the American Mathematical Society*, 72(2):341–366, March 1952. ISSN 1088-6850. URL <http://www.jstor.org/stable/1990760>.
- Bogdan Dumitrescu. *Positive Trigonometric Polynomials and Signal Processing Applications*. Signals and Communication Technology. Springer, 2007. ISBN 978-1-4020-5124-1. URL <http://gen.lib.rus.ec/get?md5=5346e169091b2d928d8333cd053300f9>.
- Nelson Dunford and Jacob T. Schwartz. *Linear operators. Part 1, General Theory*, volume 7 of *Pure and applied mathematics*. Interscience Publishers, New York, 1957. ISBN 0471226394. URL <http://www.amazon.com/dp/0471608483>. with the assistance of William G. Bade and Robert G. Bartle.
- W.D. Duthie. Segments of ordered sets. *Transactions of the American Mathematical Society*, 51(1): 1–14, January 1942. doi: 10.2307/1989978. URL <http://www.jstor.org/stable/1989978>.
- Paul H. Edelman. Abstract convexity and meet-distributive lattices. In Ivan Rival, editor, *Combinatorics and ordered sets: Proceedings of the AMS-IMS-SIAM joint summer research conference, held August 11–17, 1985*, volume 57 of *Contemporary Mathematics*, pages 127–150, Providence RI, 1986. American Mathematical Society. ISBN 0821850512. URL <http://books.google.com/books?vid=ISBN0821850512>. conference held in Arcata California.

- Paul H. Edelman and Robert E. Jamison. The theory of convex geometries. *Geometriae Dedicata*, 19(3):247–270, December 1985. ISSN 0046-5755. doi: 10.1007/BF00149365. URL <http://www.springerlink.com/content/n4344856887387gw/>.
- Robert E. Edwards. *Functional Analysis: Theory and Applications*. Dover books on mathematics. Dover, New York, 1995. ISBN 0-486-68143-2. URL <http://books.google.com/books?vid=ISBN0486681432>.
- Yuli Eidelman, Vitali D. Milman, and Antonis Tsolomitis. *Functional Analysis: An Introduction*, volume 66 of *Graduate Studies in Mathematics*. American Mathematical Society, 2004. ISBN 0821836463. URL <http://books.google.com/books?vid=ISBN0821836463>.
- Per Enflo. A counterexample to the approximation problem in banach spaces. *Acta Mathematica*, 130:309–317, 1973. URL <http://link.springer.com/content/pdf/10.1007/BF02392270>.
- Euclid. *Elements*. circa 300BC. URL <http://farside.ph.utexas.edu/euclid.html>.
- David Ewen. *The Book of Modern Composers*. Alfred A. Knopf, New York, 1950. URL <http://books.google.com/books?id=yHw4AAAAIAAJ>.
- David Ewen. *The New Book of Modern Composers*. Alfred A. Knopf, New York, 3 edition, 1961. URL <http://books.google.com/books?id=bZIaAAAAMAAJ>.
- D. J. Ewins. *Modal Testing: Theory and Practice*. Research Studies Press, “reprinted with amendments april 1986” edition, April 1986. ISBN 0863800173. URL <http://www.amazon.com/dp/086380036X>.
- Marián Fabian, Petr Habala, Petr Hájek, Vicente Montesinos, Václav Zizler, and Václav Zizler. *Banach Space Theory: The Basis for Linear and Nonlinear Analysis*. CMS Books in Mathematics. Springer, 2010. ISBN 1441975144. URL <http://books.google.com/books?vid=isbn1441975144>.
- Lorenzo Farina and Sergio Rinaldi. *Positive Linear Systems: Theory and Applications*. Pure and applied mathematics. John Wiley & Sons, 1 edition, July 3 2000. ISBN 9780471384564. URL <http://books.google.com/books?vid=ISBN0471384569>.
- Richard M. Feldman and Ciriaco Valdez-Flores. *Applied Probability and Stochastic Processes*. Technology & Engineering. Springer Science & Business Media, 2 edition, 2010. ISBN 9783642051586. URL <http://books.google.com/books?vid=ISBN3642051588>.
- Carlos A. Felippa. *Matrix Calculus*. University of Colorado at Boulder, August 18 1999. URL <http://caswww.colorado.edu/courses.d/IFEM.d/>.
- William Feller. *An Introduction to Probability Theory and its Applications Volume I*. Wiley series in probability and mathematical statistics. John Wiley & Sons, 3, revised edition, 1970. ISBN 9780471257080. URL <http://www.amazon.com/dp/0471257087>.
- R. A. Fisher. On the mathematical foundations of theoretical statistics. *Philosophical Transactions of the Royal Society*, January 1922. URL <https://doi.org/10.1098/rsta.1922.0009>.
- G.L. Fix and G. Strang. Fourier analysis of the finite element method in ritz-galerkin theory. *Studies in Applied Mathematics*, 48:265–273, 1969.
- Gerald B. Folland. *A Course in Abstract Harmonic Analysis*. Studies in Advanced Mathematics. CRC Press, Boca Raton, 1995. ISBN 0-8493-8490-7. URL <http://books.google.com/books?vid=ISBN0849384907>.

- Brigitte Forster and Peter Massopust, editors. *Four Short Courses on Harmonic Analysis: Wavelets, Frames, Time-Frequency Methods, and Applications to Signal and Image Analysis*. Applied and Numerical Harmonic Analysis. Springer, November 19 2009. ISBN 9780817648909. URL <http://books.google.com/books?vid=ISBN0817648909>.
- Jean-Baptiste-Joseph Fourier. Refroidissement séculaire du globe terrestre". In M. Gaston Darboux, editor, *Œuvres De Fourier*, volume 2, pages 271–288. Ministère de L'instruction Publique, Paris, France, April 1820. URL <http://gallica.bnf.fr/ark:/12148/bpt6k33707/f276.image>. original paper at pages 58–70.
- Jean-Baptiste-Joseph Fourier. *Théorie Analytique de la Chaleur (The Analytical Theory of Heat)*. Chez Firmin Didot, pere et fils, Paris, 1822. URL <http://books.google.com/books?vid=04X2vlqZx7hydlQUWEq&id=TDQJAAAAIAAJ>.
- Jean-Baptiste-Joseph Fourier. *The Analytical Theory of Heat (Théorie Analytique de la Chaleur)*. Cambridge University Press, Cambridge, February 20 1878. URL <http://www.archive.org/details/analyticaltheory00fourrich>. 1878 English translation of the original 1822 French edition. A 2003 Dover edition is also available: isbn 0486495310.
- Maurice Fréchet. Sur l'extension du théorème des probabilités totales au cas d'une suite infinie d'événements. *Ren- diconti del Reale Istituto Lombardo di Scienze e Lettere*, 63:899–900, 1930a.
- Maurice Fréchet. Sur l'extension du théorème des probabilités totales au cas d'une suite infinie d'événements. *Ren- diconti del Reale Istituto Lombardo di Scienze e Lettere*, 63:1059–1062, 1930b.
- Erik Ivar Fredholm. Sur une nouvelle méthode pour la résolution du problème de dirichlet (on a new method for the resolution of the problem of dirichlet). *Oefversigt af Kongl. Sv. Vetenskaps-Academiens Förfhandlingar*, 57:39–66, 1900.
- Erik Ivar Fredholm. Sur une classe d'équations fonctionnelles (on a class of functional equations). *Acta Mathematica*, 27(1):365–390, December 1903. ISSN 0001-5962. doi: 10.1007/BF02421317. URL <http://www.springerlink.com/content/c41371137837p252/>.
- Ferdinand Georg Frobenius. Über lineare substitutionen und bilineare formen. *Journal für die reine und angewandte Mathematik (Crelle's Journal)*, 84:1–63, 1878. ISSN 0075-4102. URL <http://www.digizeitschriften.de/home/services/pdfterms/?ID=509796>.
- Ferdinand Georg Frobenius. Über lineare substitutionen und bilineare formen. In Jean Pierre Serre, editor, *Gesammelte Abhandlungen (Collected Papers)*, volume I, pages 343–405. Springer, Berlin, 1968. URL <http://www.worldcat.org/oclc/253015>. reprint of Frobenius' 1878 paper.
- Jürgen Fuchs. *Affine Lie Algebras and Quantum Groups: An Introduction, With Applications in Conformal Field Theory*. Cambridge Monographs on Mathematical Physics. Cambridge University Press, 1995. ISBN 052148412X. URL <http://books.google.com/books?vid=ISBN052148412X>.
- Israel M. Gelfand. Normierte ringe. *Mat. Sbornik*, 9(51):3–24, 1941.
- Israel M. Gelfand and Mark A. Naimark. *Normed Rings with an Involution and their Representations*, pages 240–274. Chelsea Publishing Company, Bronx, 1964. ISBN 0821820222. URL <http://books.google.com/books?vid=ISBN0821820222>.
- Israel M. Gelfand and Mark A. Neumark. On the imbedding of normed rings into the ring of operators in hilbert space. *Mat. Sbornik*, 12(54:2):197–217, 1943a.

- Israel M. Gelfand and Mark A. Neumark. *On the imbedding of normed rings into the ring of operators in Hilbert Space*, pages 3–19. 1943b. ISBN 0821851756. URL <http://books.google.com/books?vid=ISBN0821851756>.
- John Robilliard Giles. *Introduction to the Analysis of Metric Spaces*. Number 3 in Australian Mathematical Society lecture series. Cambridge University Press, Cambridge, 1987. ISBN 978-0521359283. URL <http://books.google.com/books?vid=ISBN0521359287>.
- John Robilliard Giles. *Introduction to the Analysis of Normed Linear Spaces*. Number 13 in Australian Mathematical Society lecture series. Cambridge University Press, Cambridge, 2000. ISBN 0-521-65375-4. URL <http://books.google.com/books?vid=ISBN0521653754>.
- Israel Gohberg, Seymour Goldberg, and Marinus A. Kaashoek. *Basic Classes of Linear Operators*. Birkhäuser, Basel, 1 edition, 2003. ISBN 3764369302. URL <http://books.google.com/books?vid=ISBN3764369302>.
- Steve Goldman. *Vibration Spectrum Analysis: A Practical Approach*. Industrial Press Inc., 1999. ISBN 9780831130886. URL <http://books.google.com/books?vid=ISBN0831130881>.
- Jaideva C. Goswami and Andrew K. Chan. *Fundamentals of Wavelets; Theory, Algorithms, and Applications*. John Wiley & Sons, Inc., 1999. ISBN 0-471-19748-3. URL http://vadkudr.boom.ru/Collection/fundwave_contents.html.
- H. G. D. Goyder. Foolproof methods for frequency response measurements. Technical Report ADP003677, United Kingdom Atomic Energy Authority London (England), 1984. URL <https://apps.dtic.mil/dtic/tr/fulltext/u2/a143301.pdf#page=118>.
- I.S. Gradshteyn and I.M. Ryzhik. *Table of Integrals, Series, and Products*. 1980. ISBN 0-12-294760-6. URL <http://www.worldcat.org/isbn/0122947606>.
- Alexander Graham. *Kronecker Products and Matrix Calculus: With Applications*. Ellis Horwood Series; Mathematics and its Applications. Ellis Horwood Limited, Chichester, 1981. ISBN 0-85312-391-8. URL <http://books.google.com/books?vid=ISBN0853123918>.
- R.J Greechie. Orthomodular lattices admitting no states. *Journal of Combinatorial Theory, Series A*, 10(2):119–132, March 1971. URL [http://dx.doi.org/10.1016/0097-3165\(71\)90015-X](http://dx.doi.org/10.1016/0097-3165(71)90015-X).
- Daniel J. Greenhoe. *Structure and Analysis of Mathematical Spaces*, volume 2 of *Mathematical Structure and Design series*. version 0.30 edition, January 2017a. URL <https://www.researchgate.net/publication/312201065>.
- Daniel J. Greenhoe. The sequence $a(n,m)$ of the m polynomial coefficients of the n -th order b-spline scaled by $n!$, read by rows, with n in $0,1,2,\dots$ and m in $1,2,3,\dots,(n+1)2$. *Online Encyclopedia of Integer Sequences*, July 2017b. URL <https://oeis.org/A289358>.
- Amritava Gupta. *Real & Abstract Analysis*. Academic Publishers Calcutta, 1998. ISBN 9788186358443. URL <http://books.google.com/books?vid=ISBN8186358447>. first published in 1998, reprinted in 2000 and 2006.
- Alfréd Haar. Zur theorie der orthogonalen funktionensysteme. *Mathematische Annalen*, 69:331–371, September 1910. ISSN 1432-1807. doi: 10.1007/BF01456326.
- Norman B. Haaser and Joseph A. Sullivan. *Real Analysis*. Dover Publications, New York, 1991. ISBN 0-486-66509-7. URL <http://books.google.com/books?vid=ISBN0486665097>.

- Paul R. Haddad and Ali N. Akansu. *Multiresolution Signal Decomposition: Transforms, Subbands, and Wavelets*. Academic Press, October 1 1992. ISBN 0323138365. URL <http://books.google.com/books?vid=ISBN0323138365>.
- Henry Sinclair Hall and Samuel Ratcliffe Knight. *Higher algebra, a sequel to elementary algebra for schools*. Macmillan, London, 1894. URL <http://www.archive.org/details/higheralgebraas00kniggoog>.
- Paul R. Halmos. *Finite Dimensional Vector Spaces*. Princeton University Press, Princeton, 1 edition, 1948. ISBN 0691090955. URL <http://books.google.com/books?vid=isbn0691090955>.
- Paul R. Halmos. *Finite Dimensional Vector Spaces*. Springer-Verlag, New York, 2 edition, 1958. ISBN 0-387-90093-4. URL <http://books.google.com/books?vid=isbn0387900934>.
- Paul R. Halmos. *Introduction to Hilbert Space and the Theory of Spectral Multiplicity*. Chelsea Publishing Company, New York, 2 edition, 1998. ISBN 0821813781. URL <http://books.google.com/books?vid=ISBN0821813781>.
- Georg Hamel. Eine basis aller zahlen und die unstetigen lösungen der funktionalgleichung $f(x + y) = f(x) + f(y)$. *Mathematische Annalen*, 60(3):459–462, 1905. URL <http://gdz.sub.uni-goettingen.de/dms/load/img/?PPN=GDZPPN002260395&IDDOC=28580>.
- Deguang Han, Keri Kornelson, David Larson, and Eric Weber. *Frames for Undergraduates*, volume 40 of *Student Mathematical Library*. American Mathematical Society, 2007. ISBN 0821842129. URL <http://books.google.com/books?vid=ISBN0821842129>. Deguang Han = ???
- G.H. Hardy. Prolegomena to a chapter on inequalities. *Journal of the London Mathematical Society*, 1–4:61–78, November 8 1929. URL http://jlms.oxfordjournals.org/content/vols1-4/issue13/index.dtl#PRESIDENTIAL_ADDRESS. “Presidential Address” to the London Mathematical Society.
- Godfrey H. Hardy. *A Mathematician's Apology*. Cambridge University Press, Cambridge, 1940. URL <http://www.math.ualberta.ca/~mss/misc/A%20Mathematician's%20Apology.pdf>.
- Godfrey Harold Hardy, John Edensor Littlewood, and George Pólya. *Inequalities*. Cambridge Mathematical Library. Cambridge University Press, Cambridge, 2 edition, 1952. URL <http://books.google.com/books?vid=ISBN0521358809>.
- Simon Haykin. *Adaptive Filter Theory*. Prentice Hall, Upper Saddle River, 4 edition, September 24 2001. ISBN 978-0130901262. URL <http://books.google.com/books?vid=isbn0130901261>.
- Michiel Hazewinkel, editor. *Handbook of Algebras*, volume 2. North-Holland, Amsterdam, 1 edition, 2000. ISBN 044450396X. URL <http://books.google.com/books?vid=ISBN044450396X>.
- Jean Van Heijenoort. *From Frege to Gödel : A Source Book in Mathematical Logic, 1879-1931*. Harvard University Press, Cambridge, Massachusetts, 1967. URL <http://www.hup.harvard.edu/catalog/VANFGX.html>.
- Christopher Heil. *A Basis Theory Primer. Applied and Numerical Harmonic Analysis*. Birkhäuser, Boston, expanded edition edition, 2011. ISBN 9780817646868. URL <http://books.google.com/books?vid=ISBN9780817646868>.
- J. H. Heinbockel. *Introduction to Finite and Infinite Series and Related Topics*. Trafford Publishing, 2010. ISBN 9781426949548. URL <http://books.google.com/books?vid=ISBN9781426949545>.

- Carl W. Helstrom. *Probability and Stochastic Processes for Engineers*. Maxwell Macmillan international editions in engineering. Macmillan, 2 edition, 1991. ISBN 9780023535710. URL <http://books.google.com/books?vid=ISBN0023535717>.
- Omar Hijab. *Introduction to Calculus and Classical Analysis*. Undergraduate Texts in Mathematics. Springer, 3 edition, 2011. ISBN 9781441994875. URL <http://books.google.com/books?vid=ISBN1441994874>.
- David Hilbert. Grundzüge einer allgemeinen theorie der linearen integralgleichungen (fundamentals of a general theory of the linear integral equations). *Mathematisch-Physikalische Klasse (Mathematical-Physical Class)*, pages 49–91, March 1904. URL <http://dz-srv1.sub.uni-goettingen.de/sub/digbib/loader?did=D57552>. Report 1 of 6.
- David Hilbert. Grundzüge einer allgemeinen theorie der linearen integralgleichungen (fundamentals of a general theory of the linear integral equations). *Mathematisch-Physikalische Klasse (Mathematical-Physical Class)*, pages 157–228, March 1906. URL <http://dz-srv1.sub.uni-goettingen.de/sub/digbib/loader?did=D58133>. Report 4 of 6.
- David Hilbert. *Grundzüge einer allgemeinen Theorie der linearen Integralgleichungen (Fundamentals of a general theory of the linear integral equations)*, volume 26 of *Fortschritte der mathematischen Wissenschaften in Monographien (Progress of the mathematical sciences in Monographs)*. B.G. Teubner, Leipzig und Berlin, 1912. URL <http://www.worldcat.org/oclc/13468199>.
- David Hilbert, Lothar Nordheim, and John von Neumann. über die grundlagen der quantenmechanik (on the bases of quantum mechanics). *Mathematische Annalen*, 98:1–30, 1927. ISSN 0025-5831 (print) 1432-1807 (online). URL <http://dz-srv1.sub.uni-goettingen.de/cache/toc/D27776.html>.
- Otto Hölder. üeber einen mittelwerthssatz. *Göttingen Nachrichten*, pages 38–47, 1889. URL <http://www.digizeitschriften.de/dms/img/?PPN=GDZPPN00252421X>.
- Klaus Höllig. *Finite Element Methods With B-Splines*, volume 26 of *Frontiers in Applied Mathematics*. SIAM, 2003. ISBN 0-89871-533-4. URL <http://books.google.com/books?vid=ISBN0898715334>.
- Lars Hörmander. *Notions of Convexity*. Birkhäuser, Boston, 1994. ISBN 0817637990. URL <http://books.google.com/books?vid=ISBN0817637990>.
- Roger A. Horn and Charles R. Johnson. *Matrix Analysis*. Cambridge University Press, Cambridge, 1990. ISBN 0-521-30586-1. URL <http://books.google.com/books?vid=isbn0521305861>. Library: QA188H66 1985.
- Alfred Edward Housman. *More Poems*. Alfred A. Knopf, 1936. URL <http://books.google.com/books?id=rTMiAAAAMAAJ>.
- Emmanuel C. Ifeachor and Barrie W. Jervis. *Digital Signal Processing: A Practical Approach*. Electronic systems engineering series. Prentice Hall, 1993. ISBN 020154413X. URL <http://www.amazon.com/dp/020154413X>.
- Emmanuel C. Ifeachor and Barrie W. Jervis. *Digital Signal Processing: A Practical Approach*. Electronic systems engineering series. Prentice Hall, 2002. ISBN 9780201596199. URL <http://www.amazon.com/dp/0201596199>.
- Satoru Igari. *Real Analysis—With an Introduction to Wavelet Theory*, volume 177 of *Translations of mathematical monographs*. American Mathematical Society, 1996. ISBN 9780821821046. URL <http://books.google.com/books?vid=ISBN0821821040>.

- Vasile I. Istrătescu. *Inner Product Structures: Theory and Applications*. Mathematics and Its Applications. D. Reidel Publishing Company, 1987. ISBN 9789027721822. URL <http://books.google.com/books?vid=ISBN9027721823>.
- Robert C. James. Orthogonality in normed linear spaces. *Duke Mathematical Journal*, 12(2):291–302, 1945. ISSN 0012-7094. doi: 10.1215/S0012-7094-45-01223-3. URL <http://projecteuclid.org/euclid.dmj/1077473105>.
- Robert C. James. Orthogonality and linear functionals in normed linear spaces. *Transactions of the American Mathematical Society*, 61(2):265–292, March 1947. ISSN 1088-6850. doi: 10.2307/1990220. URL <http://www.jstor.org/stable/1990220>.
- Bjorn Jawerth and Wim Sweldens. An overview of wavelet based multiresolutional analysis. *SIAM Review*, 36:377–412, September 1994. URL <http://cm.bell-labs.com/who/wim/papers/papers.html#overview>.
- Alan Jeffrey and Hui Hui Dai. *Handbook of Mathematical Formulas and Integrals*. Handbook of Mathematical Formulas and Integrals Series. Academic Press, 4 edition, January 18 2008. ISBN 9780080556840. URL <http://books.google.com/books?vid=ISBN0080556841>.
- J. L. W. V. Jensen. Sur les fonctions convexes et les inégalités entre les valeurs moyennes (on the convex functions and the inequalities between the average values). *Acta Mathematica*, 30(1):175–193, December 1906. ISSN 0001-5962. doi: 10.1007/BF02418571. URL <http://www.springerlink.com/content/r55q1411g840j446/>.
- Leonard Benjamin William Jolley. *Summation of Series*. Dover, New York, second revised edition, 1961. URL <http://plouffe.fr/simon/math/SummationofSeries.pdf>.
- P. Jordan and J. von Neumann. On inner products in linear, metric spaces. *Annals of Mathematics*, 36(3):719–723, July 1935. doi: 10.2307/1968653. URL <http://www.jstor.org/stable/1968653>.
- Palle E. T. Jørgensen, Kathy Merrill, Judith Packer, and Lawrence W. Baggett. *Representations, Wavelets and Frames: A Celebration of the Mathematical Work of Lawrence Baggett*. Applied and Numerical Harmonic Analysis. Birkhäuser, 2008. ISBN 9780817646820. URL <http://books.google.com/books?vid=ISBN0817646825>.
- K. D. Joshi. *Applied Discrete Structures*. New Age International, New Delhi, 1997. ISBN 8122408265. URL <http://books.google.com/books?vid=ISBN8122408265>.
- J.S. Chitode. *Signals And Systems*. Technical Publications, 2009. ISBN 9788184316780. URL <http://books.google.com/books?vid=ISBN818431678X>.
- Gudrun Kalmbach. *Measures and Hilbert Lattices*. World Scientific, Singapore, 1986. ISBN 9789971500092. URL <http://books.google.com/books?vid=ISBN9971500094>.
- Nidal S. Kamel and K. S. Sim. Image signal-to-noise ratio and noise variance estimation using autoregressive model. *The Journal of Scanning Microscopes*, 26:277–281, September 15 2004. URL <https://onlinelibrary.wiley.com/doi/pdf/10.1002/sca.4950260605>.
- Steven M. Kay. *Modern Spectral Estimation: Theory and Application*. Prentice-Hall signal processing series. Prentice Hall, 1988. ISBN 9788131733561. URL <http://books.google.com/books?vid=ISBN8131733564>.
- James P. Keener. *Principles of Applied Mathematics; Transformation and Approximation*. Addison-Wesley Publishing Company, Reading, Massachusetts, 1988. ISBN 0-201-15674-1. URL <http://www.worldcat.org/isbn/0201156741>.

- Yoshida Kenko. *The Tsurezure Gusa of Yoshida No Kaneyoshi. Being the meditations of a recluse in the 14th Century (Essays in Idleness)*. circa 1330. URL <http://www.humanistictexts.org/kenko.htm>. 1911 translation of circa 1330 text.
- Mohamed A. Khamsi and W.A. Kirk. *An Introduction to Metric Spaces and Fixed Point Theory*. John Wiley, New York, 2001. ISBN 978-0471418252. URL <http://books.google.com/books?vid=isbn0471418250>.
- K.S. Kim and Nidal S. Kamel. Image signal-to-noise ratio and noise variance estimation using autoregressive model. *The Journal of Scanning Microscopes*, 26(6):135–139, September 24 2004. URL <https://doi.org/10.1002/sca.4950260605>.
- Anthony W Knapp. *Advanced Real Analysis*. Cornerstones. Birkhäuser, Boston, Massachusetts, USA, 1 edition, July 29 2005a. ISBN 0817643826. URL <http://books.google.com/books?vid=ISBN0817643826>.
- Anthony W Knapp. *Basic Real Analysis*. Cornerstones. Birkhäuser, Boston, Massachusetts, USA, 1 edition, July 29 2005b. ISBN 0817632506. URL <http://books.google.com/books?vid=ISBN0817632506>.
- Andrei N. Kolmogorov. *Grundbegriffe der Wahrscheinlichkeitsrechnung*. Springer, Berlin, 1933a.
- Andrei Nikolaevich Kolmogorov. *Foundations of the theory of probability*. Chelsea Publishing Company, New Yourk, 2 edition, 1933b. ISBN B0006AUOGO. URL <http://statweb.stanford.edu/~cgates/PERSI/Courses/Phil166-266/Kolmogorov-Foundations.pdf>. 1956 2nd edition English translation of A. N. Kolmogorov's 1933 "Grundbegriffe der Wahrscheinlichkeitsrechnung".
- Lambert Herman Koopmans. volume 22 of *Probability and mathematical statistics*. Academic Press, 2 edition, 1995. ISBN 9780124192515.
- Carlos S. Kubrusly. *The Elements of Operator Theory*. Springer, 1 edition, 2001. ISBN 9780817641740. URL <http://books.google.com/books?vid=ISBN0817641742>.
- Carlos S. Kubrusly. *The Elements of Operator Theory*. Springer, 2 edition, 2011. ISBN 9780817649975. URL <http://books.google.com/books?vid=ISBN0817649972>.
- Andrew Kurdila and Michael Zabarankin. *Convex Functional Analysis. Systems & Control: Foundations & Applications*. Birkhäuser, Boston, 2005. ISBN 9783764321987. URL <http://books.google.com/books?vid=ISBN3764321989>.
- Peter D. Lax. *Functional Analysis*. John Wiley & Sons Inc., USA, 2002. ISBN 0-471-55604-1. URL <http://www.worldcat.org/isbn/0471556041>. QA320.L345 2002.
- Steven R. Lay. *Convex Sets and their Applications*. John Wiley & Sons, New York, 1982. ISBN 0-471-09584-2. URL <http://books.google.com/books?vid=isbn0471095842>.
- Q. Leclere, Bert Roozen, and Celine Sandier. On the use of the hs estimator for the experimental assessment of transmissibility matrices. *Mechanical Systems and Signal Processing*, 43(1-2):237–245, 2014. URL <https://hal.archives-ouvertes.fr/hal-00911916/document>. <hal-00911916>.
- Gottfried Wilhelm Leibniz. Letter to christian huygens, 1679. In Leroy E. Loemker, editor, *Philosophical Papers and Letters*, volume 2 of *The New Synthese Historical Library*, chapter 27, pages 248–249. Kluwer Academic Press, Dordrecht, 2 edition, September 8 1679. ISBN 902770693X. URL <http://books.google.com/books?vid=ISBN902770693X>.

- J. Leuridan, J. De Vis, and Van der Auweraer H. F. Lembregts D. "a comparison of some frequency response function measurement techniques. pages 908–918, 1986. URL <https://www.researchgate.net/publication/283994172>.
- Zach Liang and George C. Lee. *Random Vibration: Mechanical, Structural, and Earthquake Engineering Applications*. Advances in Earthquake Engineering. Taylor & Francis, illustrated edition, 2015. ISBN 9781498702348. URL <http://www.amazon.com/dp/1498702341>.
- Elliott H. Lieb and Michael Loss. *Analysis*, volume 14 of *Graduate studies in mathematics*. American Mathematical Society, Providence, Rhode Island, USA, 2 edition, 2001. ISBN 0821827839. URL <http://books.google.com/books?vid=ISBN0821827839>.
- Joram Lindenstrauss and Lior Tzafriri. *Classical Banach Spaces I: Sequence Spaces*, volume 92 of *Ergebnisse der Mathematik und ihrer Grenzgebiete*. Springer-Verlag, 1977. ISBN 978-3642665592. URL <http://www.amazon.com/dp/3642665594>.
- Anders Lindquist and Giorgio Picci. *Linear Stochastic Systems: A Geometric Approach to Modeling, Estimation and Identification*, volume 1 of *Series in Contemporary Mathematics*. Springer, 2015. ISBN 9783662457504. URL <http://books.google.com/books?vid=ISBN3662457504>.
- Lynn H. Loomis. *An Introduction to Abstract Harmonic Analysis*. The University Series in Higher Mathematics. D. Van Nostrand Company, Toronto, 1953. URL <http://books.google.com/books?id=aNg-AAAAIAAJ>.
- Lynn H. Loomis and Ethan D. Bolker. *Harmonic analysis*. Mathematical Association of America, 1965. URL <http://books.google.com/books?id=MEfvAAAAMAAJ>.
- Niccolò Machiavelli. *The Literary Works of Machiavelli: Mandragola, Clizia, A Dialogue on Language, and Belfagor, with Selections from the Private Correspondence*. Oxford University Press, 1961. ISBN 0313212481. URL <http://www.worldcat.org/isbn/0313212481>.
- Fumitomo Maeda and Shûichirô Maeda. *Theory of Symmetric lattices*, volume 173 of *Die Grundlehren der mathematischen Wissenschaften in Einzeldarstellungen*. Springer-Verlag, Berlin/New York, 1970. URL <http://books.google.com/books?id=4oeBAAAIAAJ>.
- Khosrow Maleknejad, Monireh Nosrati Sahlan, and Azadeh Ostad. Numerical solution of fractional integro-differential equation by using cubic b-spline wavelets. In *Proceedings of the World Congress on Engineering*, volume I, London, July 3–5 2013. URL <https://pdfs.semanticscholar.org/bbe0/55f40768bbefe9e6476bdce701effb9243f7.pdf>.
- Lech Maligranda. A simple proof of the hölder and the minkowski inequality. *The American Mathematical Monthly*, 102(3):256–259, March 1995. URL <http://www.jstor.org/stable/2975013>.
- Stéphane G. Mallat. Multiresolution approximations and wavelet orthonormal bases of $l^2(\mathbb{R})$. *Transactions of the American Mathematical Society*, 315(1):69–87, September 1989. URL http://blanche.polytechnique.fr/~mallat/papiers/math_multiresolution.pdf.
- Stéphane G. Mallat. *A Wavelet Tour of Signal Processing*. Elsevier, 2 edition, September 15 1999. ISBN 0-12-466606-X. URL <http://books.google.com/books?vid=ISBN012466606X>.
- R. Viswanathan Mandyam D. Srinath, PK. Rajasekaran. *Introduction to Statistical Signal Processing with Applications*. Prentice Hall Inc, Upper Saddle River, 1996. ISBN 013125295X. URL <http://engr.smu.edu/ee/mds/>.
- Dimitris G. Manolakis, Vinay K. Ingle, and Stephen M. Kogon. *Statistical and Adaptive Signal Processing*. Mc-Graw Hill, Boston, 2000. ISBN 0-07-040051-2.

- Dimitris G. Manolakis, Vinay K. Ingle, and Stephen M. Kogon. *Statistical and Adaptive Signal Processing: Spectral Estimation, Signal Modeling, Adaptive Filtering, and Array Processing*. Artech House signal processing library. Artech House, illustrated, reprint edition, 2005. ISBN 9781580536103. URL <http://www.amazon.com/dp/1580536107>.
- MatLab. cpsd: Cross power spectral density. *MatLab*, accessed 2018 November 26 2018a. URL <https://www.mathworks.com/help/signal/ref/cpsd.html>.
- MatLab. xcorr: Cross-correlation. *MatLab*, accessed 2018 November 26 2018b. URL <https://www.mathworks.com/help/signal/ref/xcorr.html>.
- Stefan Mazur. Sur les anneaux linéaires. *Comptes rendus de l'Académie des sciences*, 207:1025–1027, 1938.
- Stefan Mazur and Stanislaus M. Ulam. Sur les transformations isométriques d'espaces vectoriels normées. *Comptes rendus de l'Académie des sciences*, 194:946–948, 1932.
- Karl Menger. Untersuchungen über allgemeine metrik. *Mathematische Annalen*, 100:75–163, 1928. ISSN 0025-5831. URL <http://link.springer.com/article/10.1007/BF01455705>. (Investigations on general metric).
- Claudia Menini and Freddy Van Oystaeyen. *Abstract Algebra; A Comprehensive Treatment*. Marcel Dekker Inc, New York, April 2004. ISBN 0-8247-0985-3. URL <http://books.google.com/books?vid=isbn0824709853>.
- James Mercer. Functions of positive and negative type and their connection with the theory of integral equations. *Philosophical Transactions of the Royal Society of London*, 209:415–446, 1909. ISSN 02643952. URL <http://www.jstor.org/stable/91043>.
- Anthony N. Michel and Charles J. Herget. *Applied Algebra and Functional Analysis*. Dover Publications, Inc., 1993. ISBN 0-486-67598-X. URL <http://books.google.com/books?vid=ISBN048667598X>. original version published by Prentice-Hall in 1981.
- Gregory K. Miller. *Probability: Modeling and Applications to Random Processes*. Wiley, August 25 2006. URL <http://www.amazon.com/dp/0471458929>.
- Gradimir V. Milovanović and Igorž. Milovanović. On a generalization of certain results of a. osztowski and a. lupaš. *Publikacije Elektrotehničkog Fakulteta (Publications Electrical Engineering)*, (643):62–69, 1979. URL <http://www.mi.sanu.ac.rs/~gvm/radovi/643.pdf>.
- Hermann Minkowski. *Geometrie der Zahlen*. Druck und Verlag von B.G. Teubner, Leipzig, 1910. URL <http://www.archive.org/details/geometriederzah100minkrich>. Geometry of Numbers.
- Fred Mintzer. Filters for distortion-free two-band multi-rate filter banks. *IEEE Transactions on Acoustics, Speech and Signal Processing*, 32, 1985.
- Larry D. Mitchell. An improved method for the fast fourier transform (fft) calculation of the frequency response function. In M. Kunt and F. De Coulon, editors, *Proceedings of EUSIPCO-80, First European Signal Processing Conference*, Signal processing, theories and applications, pages 159–160, Lausanne, Switzerland, September 16–19 1980. Elsevier. URL <http://www.worldcat.org/oclc/63416846>. citation from Cobb(1988).
- Larry D. Mitchell. Improved methods for the fast fourier transform (fft) calculation of the frequency response function. *Journal of Mechanical Design*, 104(2):277–279, April 01 1982. doi: 10.1115/1.3256336. URL <http://mechanicaldesign.asmedigitalcollection.asme.org/article.aspx?articleid=1451670>.

- Dragoslav S. Mitrinović, J. E. Pečarić, and Arlington M. Fink. *Classical and New Inequalities in Analysis*, volume 61 of *Mathematics and its Applications (East European Series)*. Kluwer Academic Publishers, Dordrecht, Boston, London, 2010. ISBN 978-90-481-4225-5. URL <http://www.amazon.com/dp/0792320646>.
- Todd K. Moon and Wynn C. Stirling. *Mathematical Methods and Algorithms for Signal Processing*. Prentice Hall, Upper Saddle River, 2000. ISBN 0-201-36186-8. URL <http://books.google.com/books?vid=isbn0201361868>.
- H. P. Mulholland. On generalizations of minkowski's inequality in the form of a triangle inequality. *Proceedings of the London Mathematical Society*, s2-51:294–307, 1950. URL <http://plms.oxfordjournals.org/content/s2-51/1/294.extract>. received 1946 October 10, read 1947 June 19.
- Paul J. Nahin. *Dr. Euler's Fabulous Formula: Cures Many Mathematical Ills*. Princeton Science Library. Princeton University Press, reprint, revised edition, April 25 2011. ISBN 9781400838479. URL <http://books.google.com/books?vid=ISBN1400838479>.
- Louis Narens. *Theories of Probability: An examination of logical and qualitative foundations*, volume 2 of *Advanced series on mathematical psychology*. World Scientific, January 2007. ISBN 9789812708014. URL <http://www.amazon.com/dp/9812708014>.
- Louis Narens. Alternative probability theories for cognitive psychology. *Topics in Cognitive Science*, 6(1):114–120, January 2014. ISSN 1756-8757. doi: 10.1111/tops.12071. URL <http://onlinelibrary.wiley.com/doi/10.1111/tops.12071/full>.
- Oliver Nelles. *Nonlinear System Identification*. Springer, New York, 2001. ISBN 9783540673699.
- Ben Noble and James W. Daniel. *Applied Linear Algebra*. Prentice-Hall, Englewood Cliffs, NJ, USA, 3 edition, 1988. ISBN 0-13-041260-0. URL <http://www.worldcat.org/isbn/0130412600>. Library QA184.N6 1988 512.5 87-11511.
- Timur Oikhberg and Haskell Rosenthal. A metric characterization of normed linear spaces. *Rocky Mountain Journal Of Mathematics*, 37(2):597–608, 2007. URL <http://www.ma.utexas.edu/users/rosenth1/pdf-papers/95-oikh.pdf>.
- Oystein Ore. On the foundation of abstract algebra. i. *The Annals of Mathematics*, 36(2):406–437, April 1935. URL <http://www.jstor.org/stable/1968580>.
- Brad Osgood. The fourier transform and its applications; ee261: Standford university course. university course notes, 2002. URL <http://www.stanford.edu/class/ee261/>.
- Lincoln P. Paine. *Warships of the World to 1900*. Ships of the World Series. Houghton Mifflin Harcourt, 2000. ISBN 9780395984147. URL <http://books.google.com/books?vid=ISBN9780395984149>.
- Endre Pap. *Null-Additive Set Functions*, volume 337 of *Mathematics and Its Applications*. Kluwer Academic Publishers, 1995. ISBN 0792336585. URL <http://www.amazon.com/dp/0792336585>.
- Athanasios Papoulis. *Probability, Random Variables, and Stochastic Processes*. McGraw-Hill Series in Electrical Engineering. McGraw-Hill Book Company, New York, 2 edition, 1984. ISBN 0070484686. URL <http://books.google.com/books?vid=ISBN0070484686>.
- Athanasios Papoulis. *Probability, Random Variables, and Stochastic Processes*. McGraw-Hill, New York, 3 edition, 1991. ISBN 0070484775. URL <http://books.google.com/books?vid=ISBN0070484775>.

- Anthanasios Papoulis. *Probability, Random Variables, and Stochastic Processes*. McGraw-Hill series in electrical and computer engineering. Tata McGraw-Hill, New York, 4 edition, 2002. ISBN 9780071226615. URL <http://books.google.com/books?vid=ISBN0071226613>.
- Athanasiros Papoulis. *Probability & Statistics*. Prentice-Hall international editions. Prentice Hall, illustrated edition, 1990. ISBN 9780137116980. URL <http://books.google.com/books?vid=ISBN0137116985>.
- Giuseppe Peano. *Calcolo geometrico secondo l'Ausdehnungslehre di H. Grassmann preceduto dalle operazioni della logica deduttiva*. Fratelli Bocca Editori, Torino, 1888a. Geometric Calculus: According to the Ausdehnungslehre of H. Grassmann.
- Giuseppe Peano. *Geometric Calculus: According to the Ausdehnungslehre of H. Grassmann*. Springer (2000), 1888b. ISBN 0817641262. URL <http://books.google.com/books?vid=isbn0817641262>. originally published in 1888 in Italian.
- Michael Pedersen. *Functional Analysis in Applied Mathematics and Engineering*. Chapman & Hall/CRC, New York, 2000. ISBN 9780849371691. URL <http://books.google.com/books?vid=ISBN0849371694>. Library QA320.P394 1999.
- J. E. Pečarić, Frank Proschan, and Yung Liang Tong. *Convex Functions, Partial Orderings, and Statistical Applications*, volume 187 of *Mathematics in Science and Engineering*. Academic Press, San Diego, California, 1992. ISBN 978-0125492508. URL <http://books.google.com/books?vid=ISBN0125492502>.
- Rik Pintelon and Johan Schoukens. *System Identification: A Frequency Domain Approach*. John Wiley & Sons, 2 edition, 2012. ISBN 9781118287392. URL <http://books.google.com/books?vid=ISBN1118287398>.
- Henri Poincaré. *Calcul Des Probabilités*. De L'ecole Polytechnique, Paris, 1912. URL <http://www.archive.org/details/calculdeprobabil00poinrich>.
- Karl Raimund Popper. *Conjectures and refutations: the growth of scientific knowledge*. Basic Books, 1962. URL <https://books.google.com/books?id=OBgNAQAAIAAJ>.
- Karl Raimund Popper. *Conjectures and refutations: the growth of scientific knowledge*. Routledge, 2 edition, 1963. ISBN 9781135971304.
- Lakshman Prasad and Sundararaja S. Iyengar. *Wavelet Analysis with Applications to Image Processing*. CRC Press LLC, Boca Raton, 1997. ISBN 978-0849331695. URL <http://books.google.com/books?vid=ISBN0849331692>. Library TA1637.P7 1997.
- John G. Proakis. *Digital Communications*. McGraw Hill, 4 edition, 2001. ISBN 0-07-232111-3. URL <http://www.mhhe.com/>.
- Inder K. Rana. *An Introduction to Measure and Integration*, volume 45 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, R.I., 2 edition, 2002. ISBN 978-0821829745. URL <http://books.google.com/books?vid=ISBN0821829742>.
- Charles Earl Rickart. *General Theory of Banach Algebras*. University series in higher mathematics. D. Van Nostrand Company, Yale University, 1960. URL <http://books.google.com/books?id=PVrvAAAAMAAJ>.
- Frigyes Riesz. Zur theorie des hilbertschen raumes. *Acta Scientiarum Mathematicarum*, 7:34–38, 1934. URL <http://www.math.u-szeged.hu/acta/>. The Theory of Hilbert Space.

- Enders A. Robinson. *Random Wavelets and Cybernetic Systems*, volume 9 of *Griffins Statistical Monographs & Courses*. Lubrecht & Cramer Limited, London, June 1962. ISBN 0852640757. URL <http://books.google.com/books?vid=ISBN0852640757>.
- Enders A. Robinson. Multichannel z-transforms and minimum delay. *Geophysics*, 31(3):482–500, June 1966. doi: 10.1190/1.1439788. URL <http://dx.doi.org/10.1190/1.1439788>.
- R. Tyrrell Rockafellar. *Convex Analysis*. Princeton landmarks in mathematics. Princeton University Press, 1970. ISBN 9780691015866. URL <http://books.google.com/books?vid=ISBN0691015864>.
- Sheldon Ross. *A First Course in Probability*. Prentice Hall, Upper Saddle River, New Jersey, USA 07458, 5 edition, 1998. ISBN 0-13-746314-6. URL <http://www.amazon.com/dp/0137463146>.
- George G. Roussas. *An Introduction to Probability and Statistical Inference*. Academic Press, 2 edition, October 21 2014. ISBN 9780128004371. URL <http://books.google.com/books?vid=ISBN0128004371>.
- Walter Rudin. *Principles of Mathematical Analysis*. McGraw-Hill, New York, 3 edition, 1976. ISBN 007054235X. URL <http://books.google.com/books?vid=ISBN007054235X>. Library QA300.R8 1976.
- Walter Rudin. *Functional Analysis*. McGraw-Hill, New York, 2 edition, 1991. ISBN 0-07-118845-2. URL <http://www.worldcat.org/isbn/0070542252>. Library QA320.R83 1991.
- Bryan P. Rynne and Martin A. Youngson. *Linear Functional Analysis*. Springer undergraduate mathematics series. Springer, 2 edition, January 1 2008. ISBN 9781848000056. URL <http://books.google.com/books?vid=ISBN1848000057>.
- Shôichirô Sakai. *C*-Algebras and W*-Algebras*. Springer-Verlag, Berlin, 1 edition, 1998. ISBN 9783540636335. URL <http://books.google.com/books?vid=ISBN3540636331>. reprint of 1971 edition.
- Louis L. Scharf. *Statistical Signal Processing*. Addison-Wesley Publishing Company, Reading, MA, 1991. ISBN 0-201-19038-9.
- H. A. Scharz. über ein die flächen kleinsten flächeninhalts betreffendes problem der variation-srechnung. *Acta Soc. Scient. Fen.*, 15:315–362, 1885. URL http://www-stat.wharton.upenn.edu/~steele/Publications/Books/CSMC/CSMC_index.html.
- Juljusz Schauder. Zur theorie stetiger abbildungen in funktionalräumen. *Mathematische Zeitschrift*, 26:47–65, 1927. URL <http://eudml.org/doc/167911;jsessionid=156A34EBAB6C0E2DDAAC8C1232D23E8F>.
- Juljusz Schauder. Eine eigenschaft des haarschen orthogonalsystems. *Mathematische Zeitschrift*, 28:317–320, 1928.
- Larry L. Schumaker. *Spline Functions: Basic Theory*. Cambridge Mathematical Library. Cambridge University Press, 3 edition, August 2007. ISBN 978-0-521-70512-7. URL <http://books.google.com/books?vid=ISBN0521705126>.
- Isaac Schur. Über die charakterischen wurzeln einer linearen substitution mit enier anwendung auf die theorie der integralgleichungen (over the characteristic roots of one linear substitution with an application to the theory of the integral). *Mathematische Annalen*, 66:488–510, 1909. URL <http://dz-srv1.sub.uni-goettingen.de/cache/toc/D38231.html>.

- Karl Hermann Amandus Schwarz. *Ueber Ein Die Flachen Kleinsten Flächeninhalts; Betreffendes Problem Der Variationsrechnung* (Over flattening smallest flat contents;. October 31 1885. URL <http://www-stat.wharton.upenn.edu/~steele/Publications/Books/CSMC/Schwarz.pdf>. Festschrift Zum Jubelgeburtstage des Herrn Karl Weierstrass (anniversary publication to comemerate the birthdays of Mr. Karl Weierstrass).
- Mícheál Ó Searcoid. *Elements of Abstract Analysis*. Springer Undergraduate Mathematics Series. Springer, 2002. ISBN 9781852334246. URL <http://books.google.com/books?vid=ISBN185233424X>.
- Jun Shao. *Mathematical Statistics*. Springer Texts in Statistics. Springer Science & Business Media, 2003. ISBN 9780387953823. URL <http://books.google.com/books?vid=ISBN0387953825>.
- Kihong Shin and Joseph Hammond. *Fundamentals of Signal Processing for Sound and Vibration Engineers*. John Wiley & Sons, 2008. ISBN 9780470725641.
- Bernard Shizgal. *Spectral Methods in Chemistry and Physics: Applications to Kinetic Theory and Quantum Mechanics*. Scientific Computation. Springer, January 7 2015. ISBN 9789401794541. URL <http://books.google.com/books?vid=ISBN9401794545>.
- George Finlay Simmons. *Calculus Gems: Brief Lives and Memorable Mathematicians*. Mathematical Association of America, Washington DC, 2007. ISBN 0883855615. URL <http://books.google.com/books?vid=ISBN0883855615>.
- Barry Simon. *Convexity: An Analytic Viewpoint*, volume 187 of *Cambridge Tracts in Mathematics*. Cambridge University Press, May 19 2011. ISBN 9781107007314. URL <http://books.google.com/books?vid=ISBN1107007313>.
- Iván Singer. *Bases in Banach Spaces I*, volume 154 of *Grundlehren der mathematischen Wissenschaften in Einzeldarstellungen mit besonderer Berücksichtigung der Anwendungsgebiete*. Springer-Verlag New York, New York, 1970. ISBN 9780387048338.
- Herbert Ellsworth Slaught and Nels Johann Lennes. *Elementary Algebra*. Allyn and Bacon, 1915. URL <https://archive.org/details/elementaryalgebr00slaurich/>.
- M.J.T. Smith and T.P. Barnwell. A procedure for designing exact reconstruction filter banks for tree-structured subband coders. *IEEE International Conference on Acoustics, Speech and Signal Processing*, 9:421–424, 1984a. T.P. Barnwell is T.P. Barnwell III.
- M.J.T. Smith and T.P. Barnwell. The design of digital filters for exact reconstruction in subband coding. *IEEE Transactions on Acoustics, Speech and Signal Processing*, 34(3):434–441, June 1984b. ISSN 0096-3518. doi: 10.1109/TASSP.1986.1164832. T.P. Barnwell is T.P. Barnwell III.
- J. Laurie Snell. A conversation with joe doob. *Statistical Science*, 12(4):301–311, 1997. URL <http://www.jstor.org/stable/2246220>. similar article on web at <http://www.dartmouth.edu/~chance/Doob/conversation.html>.
- J. Laurie Snell. Obituary: Joseph leonard doob. *Journal of Applied Probability*, 42(1):247–256, 2005. URL <http://projecteuclid.org/euclid.jap/1110381384>.
- Houshang H. Sohrab. *Basic Real Analysis*. Birkhäuser, Boston, 1 edition, 2003. ISBN 978-0817642112. URL <http://books.google.com/books?vid=ISBN0817642110>.
- J. Michael Steele. *The Cauchy-Schwarz Master Class: An Introduction to the Art of Mathematical Inequalities*. Cambridge University Press, Cambridge, April 26 2004. ISBN 052154677X. URL http://www-stat.wharton.upenn.edu/~steele/Publications/Books/CSMC/CSMC_index.htm.

- Lynn Arthur Steen. Highlights in the history of spectral theory. *The American Mathematical Monthly*, 80(4):359–381, April 1973. ISSN 00029890. URL <http://www.jstor.org/stable/2319079>.
- Josef Stoer and R. Bulirsch. *Introduction to Numerical Analysis*, volume 12 of *Texts in Applied Mathematics*. Springer Science & Business Media, 3 edition, August 21 2002. ISBN 9780387954523. URL <http://books.google.com/books?vid=ISBN038795452X>.
- Marshall Harvey Stone. *Linear transformations in Hilbert space and their applications to analysis*, volume 15 of *American Mathematical Society. Colloquium publications*. American Mathematical Society, New York, 1932. URL <http://books.google.com/books?vid=ISBN0821810154>. 1990 reprint of the original 1932 edition.
- Gilbert Strang. Wavelets and dilation equations: A brief introduction. *SIAM Review*, 31(4):614–627, December 1989. ISSN 0036-1445. URL <https://pdfs.semanticscholar.org/d118/a25a91488b4b37457cdc506472819de90523.pdf>.
- Gilbert Strang and Truong Nguyen. *Wavelets and Filter Banks*. Wellesley-Cambridge Press, Wellesley, MA, 1996. ISBN 9780961408879. URL <http://books.google.com/books?vid=ISBN0961408871>.
- Yuri M. Suhov, Michael Kelbert, and Mark Kelbert. *Probability and Statistics by Example: Volume 1, Basic Probability and Statistics*, volume 1 of *Probability and Statistics by Example*. Cambridge University Press, illustrated, reprint edition, October 13 2005. ISBN 9780521847667.
- Wim Sweldens and Robert Piessens. Wavelet sampling techniques. In *1993 Proceedings of the Statistical Computing Section*, pages 20–29. American Statistical Association, August 1993. URL <http://citeseer.ist.psu.edu/18531.html>.
- Tamás Szabó and Gábor Horváth. An efficient hardware implementation of feed-forward neural networks. *Applied Intelligence*, 1:143–158, 2004. URL https://www.researchgate.net/profile/Gabor_Horvath/publication/263107833.
- Brian S. Thomson, Andrew M. Bruckner, and Judith B. Bruckner. *Elementary Real Analysis*. www.classicalrealanalysis.com, 2 edition, 2008. ISBN 9781434843678. URL <http://classicalrealanalysis.info/com/Elementary-Real-Analysis.php>.
- J. T. L. Thong, K. S. Sim, and J. C. H. Phang. Single-image signal-to-noise ratio estimation. *The Journal of Scanning Microscopes*, 23:328–336, December 07 2001. URL <https://doi.org/10.1002/sca.4950230506>.
- Elmer Tolsted. An elementary derivation of cauchy, hölder, and minkowski inequalities from young's inequality. *Mathematics Magazine*, 37:2–12, 1964. URL <http://mathdl.maa.org/mathDL/22/?pa=content&sa=viewDocument&nodeId=3036>.
- Harry L. Van Trees. *Detection, Estimation, and Modulation Theory, Part I*. Wiley-Interscience, reprint edition edition, September 27 2001. ISBN 0471095176. URL http://ece.gmu.edu/faculty_info/van.html.
- Harry L. Van Trees. *Otimum Array Processing; Part IV of Detection, Estimation, and Modulation Theory*. Wiley-Interscience, New York, 2002. ISBN 0-471-09390-4. URL http://ece.gmu.edu/faculty_info/van.html.
- Kishor S. Trivedi. *Probability and Statistics with Reliability, Queuing, and Computer Science Applications*. John Wiley & Sons, 2 edition, 2016. ISBN 9781119314202.

- Constantin Udriste. *Convex Functions and Optimization Methods on Riemannian Manifolds*, volume 297 of *Mathematics and Its Applications*. Springer, July 31 1994. ISBN 0792330021. URL <http://books.google.com/books?vid=ISBN0792330021>.
- Stanislaw Marcin Ulam. *Adventures of a Mathematician*. University of California Press, Berkeley, 1991. ISBN 0520071549. URL <http://books.google.com/books?vid=ISBN0520071549>.
- P.P. Vaidyanathan. *Multirate Systems and Filter Banks*. Prentice Hall Signal Processing Series. Prentice Hall, 1993. ISBN 0136057187. URL <http://books.google.com/books?vid=ISBN0136057187>.
- Jussi Väisälä. A proof of the mazur-ulam theorem. *The American Mathematical Monthly*, 110(7): 633–635, August–September 2003. URL <http://www.helsinki.fi/~jvaisala/mazurulam.pdf>.
- M.L.J. van de Vel. *Theory of Convex Structures*, volume 50 of *North-Holland Mathematical Library*. North-Holland, Amsterdam, 1993. ISBN 978-0444815057. URL <http://books.google.com/books?vid=ISBN0444815058>.
- Brani Vidakovic. *Statistical Modeling by Wavelets*. John Wiley & Sons, Inc, New York, 1999. ISBN 0-471-29365-2. URL <http://www.amazon.com/dp/0471293652>.
- Wolfgang von der Linden, Volker Dose, and Udo Von Toussaint. *Bayesian Probability Theory: Applications in the Physical Sciences*. Cambridge University Press, 2014. ISBN 9781107035904. URL <http://books.google.com/books?vid=ISBN1107035902>.
- John von Neumann. Allgemeine eigenwerttheorie hermitescher funktionaloperatoren. *Mathematische Annalen*, 102(1):49–131, 1929. ISSN 0025-5831 (print) 1432-1807 (online). URL <http://resolver.sub.uni-goettingen.de/purl?GDZPPN002273535>. General eigenvalue theory of Hermitian functional operators.
- Gilbert G. Walter and XiaoPing Shen. *Wavelets and Other Orthogonal Systems*. Chapman and Hall/CRC, New York, 2 edition, 2001. ISBN 1-58488-227-1. URL <http://books.google.com/books?vid=ISBN1584882271>. Library QA403.3.W34 2000.
- H. L. Wei and S. A. Billings. An efficient nonlinear cardinal b-spline model for high tide forecasts at the venice lagoon. *Nonlinear Processes in Geophysics*, 13:577–584, 2006. URL <https://www.nonlin-processes-geophys.net/13/577/2006/>.
- David Wells. *The Penguin Dictionary of Curious and Interesting Numbers*. revised edition, April 7 1987. ISBN 9780140080292. URL <http://www.amazon.com/dp/0140080295>.
- Hermann Weyl. The method of orthogonal projection in potential theory. *Duke Mathematical Journal*, 7(1):411–444, 1940. URL <http://projecteuclid.org/euclid.dmj/1077492266>.
- P. R. White, M. H. Tan, and J. K. Hammond. Analysis of the maximum likelihood, total least squares and principal component approaches for frequency response function estimation. *Journal of Sound and Vibration*, 290(3–5):676–689, March 2006. URL <https://doi.org/10.1016/j.jsv.2005.04.029>.
- Alfred L. Wicks and Harvard Vold. The h_s frequency response function estimator. In *Proceedings of the 4th International Modal Analysis Conference*, pages 897–899, Los Angeles Airport Marriott Hotel, California, USA, February 3–6 1986. URL <https://www.tib.eu/en/search/id/TIBKAT%3A246489537/Proceedings-of-the-4th-International-Modal-Analysis/>.
- Samuel Stanley Wilks. Mathematical statistics. page 644, 1963.

- Peter Willett, Peter F. Swaszek, and Rick S. Blum. The good, bad, and ugly: Distributed detection of a known signal in dependent gaussian noise. *IEEE Transactions on Signal Processing*, 48(12):3266–3279, December 2000.
- P. Wojtaszczyk. *A Mathematical Introduction to Wavelets*, volume 37 of *London Mathematical Society student texts*. Cambridge University Press, February 13 1997. ISBN 9780521578943. URL <http://books.google.com/books?vid=ISBN0521578949>.
- W. J. Yan and W. X. Ren. Operational modal parameter identification from power spectrum density transmissibility. *Computer-Aided Civil and Infrastructure Engineering*, 27(3):202–217, 2012. URL <https://doi.org/10.1111/j.1467-8667.2011.00735.x>.
- Frank Yates and Kenneth Mather. Ronald aylmer fisher, 1890–1962. *Biographical Memoirs of Fellows of the Royal Society of London*, 9:90–120, November 1963. URL <https://doi.org/10.1098/rsbm.1963.0006>. print ISSN:0080-4606, online ISSN:1748-8494, <https://pdfs.semanticscholar.org/6ccd/4303f68f3543fc794c376ce00395947a79b6.pdf>.
- Robert M. Young. *An introduction to nonharmonic Fourier series*, volume 93 of *Pure and applied mathematics*. Academic Press, 1 edition, 1980. ISBN 0127728503. URL <http://books.google.com/books?vid=ISBN0127728503>.
- Robert M. Young. *An introduction to nonharmonic Fourier series*, volume 93 of *Pure and applied mathematics*. Academic Press, revised first edition, May 16 2001. ISBN 0127729550. URL <http://books.google.com/books?vid=ISBN0127729550>.
- William Henry Young. On classes of summable functions and their fourier series. *Proceedings of the Royal Society of London*, 87(594):225–229, August 1912. URL <http://www.archive.org/details/philtrans02496252>.
- Yu Zhang, Tong Zhu, and Jing Zhou. Understanding power spectrum density transmissibility. *Shock and Vibration*, 2016(8479721):9, 2016. URL <http://dx.doi.org/10.1155/2016/8479721>.
- Mu-Qiao Zheng, She-Ping Hu, and Li Li. Improved methods for the estimate of frequency response functions. pages 917–921, 2002. URL <https://pdfs.semanticscholar.org/edfc/0572a727633e729204bc9a8cbce4eb3211e5.pdf>.
- Yun-Lai Zhou and Magd Abdel Wahab. Republished paper. rapid early damage detection using transmissibility with distance measure analysis under unknown excitation in long-term health monitoring. *Journal of Vibroengineering*, 20(1):823–831, 2018. doi: 10.21595/jve.2016.19718. URL <https://doi.org/10.21595/jve.2016.19718>.



REFERENCE INDEX

- Abramovich and Aliprantis (2002), 330
Aliprantis and Burkinshaw (2006), 190, 192–195
Abom (1986), 96
Aliprantis and Burkinshaw (1998), 204, 206, 210, 233, 234, 236–239, 241, 242, 246, 249–251, 278, 282, 286, 288–290
Adobe (1999), 376
Akhiezer and Glazman (1993), 186
Allemand et al. (1979), 96
Allemand et al. (1987), 96
Allen (2018), 6
Amann and Escher (2008), 279
Amir (1986), 241, 256–258
Angus (1994), 25
Apostol (1975), 259
Aubin and Frankowska (2009), 261
Aubin (2011), 261
Autonne (1901), 303
Autonne (1902), 303
Bachman (1964), 312
Bachman and Narici (1966), 197, 199, 201, 210, 296
Bachman et al. (2000), 202, 205, 214, 309
Bailey et al. (2011), 364
Bailey et al. (2013), 364
Bak (2013), 107
Balakrishnan and Lai (2009), 25
Banach (1922), 185, 188, 189, 249, 250
Banach (1932b), 249
Banach (1932a), 202, 249, 252
Barvinok (2002), 260, 261
Beer (1993), 261
Bendat (1978), 96
Bendat and Piersol (1980), 111
Bendat and Piersol (1993), 95, 111
Bendat and Piersol (2010), 95, 96, 111
?, 44
Berberian (1961), 187, 188, 197, 210, 214, 231, 233, 240, 267, 282–285, 300
Besso (1879), 270
Bienaymé (1840), 270
Billingsley (1995), 146
Blumenthal (1970), 265
Bollobás (1999), 169–171, 206, 233, 237, 239, 240, 246, 255, 261, 262, 269, 270, 274, 290, 291
Bouniakowsky (1859), 278
Boyer and Merzbach (2011), 278
Bracewell (1978), 210, 312
Bromwich (1908), 329
Bullen (2003), 270, 274, 276, 278
Bunyakovsky (1859), 234
Byrne (2005), 223
Cadzow (1987), 11, 12, 44
Caines (1988), 8
Caines (2018), 8
Callahan (2010), 26
Carne and Dohrmann (2006), 98
Carothers (2000), 275, 276, 278
Carothers (2005), 199, 202
Cauchy (1821), 234, 274, 278
Cayley (1858), 169
Chen et al. (2012), 116
J.S.Chitode (2009), 325
Chorin and Hald (2009), 309, 310
Christensen (2003), 197, 202, 210, 215, 220, 223
Christensen (2008), 215, 220, 343, 346, 347, 352, 354, 356, 358, 360
Christensen (2010), 346
Chui (1988), 343, 352, 354, 360
Chui (1992), 322, 343, 347, 356
Cifarelli and Regazzini (1996), 146
Çinlar and Vanderbei (2013), 262
Claerbout (1976), 333
Cobb (1988), 93, 96, 100
Cobb and Mitchell (1990), 100
Conway (1990), 237
Courant and Hilbert (1930), 172
Cover and Thomas (1991), 6, 264
Cox (1972), 352, 354, 360
Curry and Feldman (2010), 155
Daubechies (1992), 203, 219, 325
Day (1973), 241
de Witt (1659), 171
de Boor (1972), 352, 354, 360
de Boor (2001), 352, 354, 360
de Finetti (1930a), 146
de Finetti (1930b), 146
de Finetti (1930c), 146
Dekking et al. (2006), 6
Descartes (1954), 169
Descartes (1637), 169
Devroye (1986), 25, 29
Daubechies et al. (1986), 220
Dieudonné (1969), 299

- Drljević (1989), 245, 258
 Duffin and Schaeffer (1952), 220
 Dumitrescu (2007), 333
 Dunford and Schwartz (1957), 256, 258, 292
 Duthie (1942), 259
 Edelman and Jamison (1985), 265
 Edelman (1986), 265
 Edwards (1995), 206
 Eidelman et al. (2004), 284
 Enflo (1973), 202
 Ewen (1950), viii
 Ewen (1961), viii
 Ewins (1986), 94, 116
 Fabian et al. (2010), 208
 Farina and Rinaldi (2000), 333
 Feldman and Valdez-Flores (2010), 155
 Felippa (1999), 173
 Feller (1970), 149
 Fisher (1922), 75
 Fix and Strang (1969), 315
 Folland (1995), 227, 228, 231, 232
 Forster and Massopust (2009), 366
 Fourier (1820), 267
 Fourier (1878), 309
 Fourier (1822), 309
 Fréchet (1930a), 146
 Fréchet (1930b), 146
 Fredholm (1900), 18, 171
 Fredholm (1903), 18, 171
 Frobenius (1968), 299
 Frobenius (1878), 299
 Fuchs (1995), 227
 Gelfand (1941), 231
 Gelfand and Neumark (1943b), 230–232
 Gelfand and Neumark (1943a), 231, 232
 Gelfand and Naimark (1964), 228, 229, 231
 Giles (1987), 251, 252
 Giles (2000), 202, 206, 250, 252, 292, 294
 Gohberg et al. (2003), 172
 Goldman (1999), 95
 Goswami and Chan (1999), 314
 Goyder (1984), 96, 100
 Gradshteyn and Ryzhik (1980), 180
 Graham (1981), 173, 177–179
 Greechie (1971), 146
 Greenhoe (2017a), 350
 Gupta (1998), 201
 Haar (1910), 372
- Haaser and Sullivan (1991), 185, 186, 201, 233–235, 238
 Haddad and Akansu (1992), 337
 Hall and Knight (1894), 279
 Halmos (1948), 185
 Halmos (1958), 296
 Halmos (1998), 229, 241, 295
 Hamel (1905), 187, 199
 Han et al. (2007), 208, 223
 Hardy (1929), 267
 Hardy et al. (1952), 269, 275, 276, 278
 Hazewinkel (2000), 227
 Heijenoort (1967), viii
 Heil (2011), 197, 199, 201, 202, 210, 215, 220, 223, 237, 281, 346, 369
 Heinbockel (2010), 364
 Helstrom (1991), 44
 Hijab (2011), 351
 Hilbert (1904), 169
 Hilbert (1906), 170, 171
 Hilbert (1912), 169, 171
 Hilbert et al. (1927), 282
 Hölder (1889), 276
 Höllig (2003), 351, 352, 354, 360
 Hörmander (1994), 265
 Horn and Johnson (1990), 71, 182, 286
 Housman (1936), viii
 Ifeachor and Jervis (1993), 68
 Ifeachor and Jervis (2002), 68
 Igari (1996), 218
 Istrătescu (1987), 192, 233, 240, 241
 James (1945), 245, 256, 258
 James (1947), 258
 Jawerth and Sweldens (1994), 314, 315, 360
 Jeffrey and Dai (2008), 310
 Jensen (1906), 6, 261, 262, 274
 Jolley (1961), 363, 364
 Jordan and von Neumann (1935), 239–241
 Jørgensen et al. (2008), 220
 Joshi (1997), 329
 Kalmbach (1986), 146
 Kamel and Sim (2004), 119
 Kay (1988), 44, 65, 92
 Keener (1988), 18–20, 171, 298
 Kenko (circa 1330), 380
 Khamsi and Kirk (2001), 265
 Kim and Kamel (2004), 119
 Knapp (2005a), 304
 Knapp (2005b), 309, 310
 Kolmogorov (1933b), 146
 Kolmogorov (1933a), 146
- Koopmans (1995), 44
 Kubrusly (2001), 185, 187, 197, 199, 201, 204, 210, 214, 243, 244, 246, 282, 296, 300, 301
 Kubrusly (2011), 329
 Kurdila and Zabarankin (2005), 197, 262
 Lax (2002), 298
 Lay (1982), 262
 Leclere et al. (2014), 99, 105
 Leibniz (1679), 281
 Leuridan et al. (1986), 93, 96
 Liang and Lee (2015), 116
 Lieb and Loss (2001), 182
 von der Linden et al. (2014), 156
 Lindenstrauss and Tzafriri (1977), 202
 Lindquist and Picci (2015), 8
 Loomis (1953), 243
 Loomis and Bolker (1965), 309
 Machiavelli (1961), 379
 Maeda and Maeda (1970), 260
 Maleknejad et al. (2013), 349
 Maligranda (1995), 238, 275, 276
 Mallat (1989), 219
 Mallat (1999), 219, 314, 315, 327, 337, 339, 367
 Manolakis et al. (2005), 44
 MatLab (2018a), 11
 MatLab (2018b), 11
 Mazur and Ulam (1932), 292
 Mazur (1938), 231
 Menger (1928), 265
 Menini and Oystaeyen (2004), 279
 Mercer (1909), 172
 Michel and Herget (1993), 18, 169, 171, 172, 187–189, 197, 199, 201, 229, 230, 249, 252, 281, 283, 285, 288, 293, 294, 297, 299, 301–303, 308
 Manolakis et al. (2000), 72
 Miller (2006), 155
 Milovanović and Milovanović (1979), 270
 Minkowski (1910), 238, 276
 Mintzer (1985), 337
 Mitchell (1980), 96
 Mitchell (1982), 96, 97, 112
 Mitrinović et al. (2010), 262
 Moon and Stirling (2000), 8, 167
 Mulholland (1950), 270
 Nahin (2011), 364
 Narens (2014), 146
 Nelles (2001), 63, 71, 73

- Noble and Daniel (1988), 307
Greenhoe (2017b), 350
Oikhberg and Rosenthal (2007), 254, 292
Ore (1935), 259
Osgood (2002), 161
Paine (2000), vi
Pap (1995), 146
Papoulis (1984), 11, 12, 18, 26, 28, 29, 44
Papoulis (1990), 6, 26, 31, 147, 148, 157, 164
Papoulis (2002), 44
Papoulis (1991), 17, 26, 28–30, 35–37, 39, 41, 45, 50, 132, 146, 148, 149, 156, 163
Peano (1888b), 185, 188, 189, 233
Peano (1888a), 188, 189
Pečarić et al. (1992), 269
Pedersen (2000), 169, 205, 206
Pintelon and Schoukens (2012), 96, 100
Poincaré (1912), 162
Popper (1962), 93
Popper (1963), 93
Prasad and Iyengar (1997), 343, 354, 366, 367
Proakis (2001), 26, 28, 29, 164–167
Rana (2002), 270
Rickart (1960), 228–231
Riesz (1934), 241
Robinson (1962), 333
Robinson (1966), 333
Rockafellar (1970), 262
Ross (1998), 6, 7, 163–165
Roussas (2014), 25
Rudin (1991), 286, 288, 289, 291, 293–296, 299, 300, 303, 304
Rudin (1976), 272, 273
Rynne and Youngson (2008), 201
Sakai (1998), 295
Scharf (1991), 173, 175, 178
Schauder (1927), 202
Schauder (1928), 202
Schumaker (2007), 345, 346, 349, 351, 352, 354, 360
Schur (1909), 303
Scharz (1885), 278
Schwarz (1885), 234
Searcoid (2002), 197
Shao (2003), 6
Shin and Hammond (2008), 93, 96, 99, 100, 105, 112, 114
Shizgal (2015), 349
Simmons (2007), 379
Simon (2011), 261
Singer (1970), 202
Slaughter and Lennes (1915), 106
Smith and Barnwell (1984a), 337
Smith and Barnwell (1984b), 337
Snell (1997), 155
Snell (2005), 155
Sohrab (2003), 201
Mandyam D. Srinath (1996), 63, 81, 84, 85
Steele (2004), 236
Steen (1973), 169, 303
Stoer and Bulirsch (2002), 346
Stone (1932), 282, 294, 298
Strang and Nguyen (1996), 337, 368, 369
Strang (1989), 368, 369
Suhov et al. (2005), 6
Sweldens and Piessens (1993), 314, 315
Szabó and Horváth (2004), 349
Thomson et al. (2008), 329
Thong et al. (2001), 119
Tolsted (1964), 275, 276
Trivedi (2016), 155
Udriste (1994), 262
Ulam (1991), 292
Vaidyanathan (1993), 337
Väistälä (2003), 292
Trees (2001), 123
Trees (2002), 173, 175, 178
van de Vel (1993), 260, 265
Vidakovic (1999), 219, 314, 315, 327, 339
von Neumann (1929), 171, 294, 298
Walter and Shen (2001), 206
Wei and Billings (2006), 349
Wells (1987), 364
Weyl (1940), 206
White et al. (2006), 96, 99, 100, 114
Wicks and Vold (1986), 99, 114
Wilks (1963), 4, 7, 11
Willett et al. (2000), 140
Wojtaszczyk (1997), 215, 217–219, 352, 354, 359, 360, 366, 372
Yan and Ren (2012), 95
Yates and Mather (1963), 11
Young (1912), 275
Young (1980), 208
Young (2001), 199, 201–203, 208, 212, 213, 215, 220
Zhang et al. (2016), 95
Zheng et al. (2002), 97, 99, 119
Zhou and Wahab (2018), 95

SUBJECT INDEX

0 order B-spline, 361
0 order B-spline dyadic decomposition, 368
0th order B-spline, 345
1st order B-spline, 346, 361
1st order B-spline dyadic decomposition, 369
2nd order B-spline, 346, 361
2nd order B-spline dyadic decomposition, 369
3rd order B-spline, 349
4th order B-spline, 349
5th order B-spline, 349
6th order B-spline, 349
7th order B-spline, 350
 C^* -algebra, 231, 232
 C^* -algebra, 232
 $\Gamma(b)$, 163
*-algebra, 12, 15, 35–37, 228, 228–230, 295, 320
 $\langle \Delta | \nabla \rangle$, 233
 α th-order modified Bessel function of the first kind, 166
 n th derivative, 356
 n th moment, 314, 314, 315
 n th order B-spline, 347, 352, 354, 360, 361, 367, 369, 370
 n th order cardinal B-spline, 343
 n th-order B-spline, 359
 N -tuple, 187, 249
 ϕ -mean, 269
(ESD), 44
*-algebras, 295
 \LaTeX , vi
 \TeX-Gyre Project , vi
 \Xe\LaTeX , vi
attention markers, 128, 132, 243, 292
 problem, 169, 286, 291
 σ -additive, 146, 152
 σ -additivity, 146

σ -algebra, 147, 152, 154
0 order B-spline, 358
1st order B-spline, 359
2 coefficient case, 372
2nd order B-spline, 359
inverse, 122
Abel, 278
Abel, Niels Henrik, 379
abelian group, 186
absolute value, x, 192, 193–195, 234, 249, 251
absolutely summable, 15, 330, 331
ADC, 95
additive, 19, 146–149, 151, 233, 234, 243, 245, 246, 282, 285
additive identity, 12, 187–189
additive identity element, 187, 188
additive inverse, 188, 189
additive property, 310
additivity, 19, 146, 147, 224, 237, 294
adjoint, 230, 291, 291, 294, 295, 310
admissibility condition, 372
admissibility equation, 372
Adobe Systems Incorporated, vi
affine, 170, 261, 292
Affine equations, 177
algebra, 185, 227, 227, 228
algebra of sets, xi, 153
algebraic, 185
algebras
 C^* -algebra, 231
 *-algebra, 228
algebras of sets, 153
Alternating Series Test, 366
AM-GM inequality, 274
analog-to-digital converter, 95
AND, xi
anti-derivative, 344
anti-symmetric, 13, 48, 313
anti-symmetric, 48
antiautomorphic, 14, 228, 229, 234, 295
antilinear, 229
antitone, 148, 151
Aristotle, 11, 17
arithmetic mean, 91, 97, 274
arithmetic mean geometric mean inequality, 274
Arithmetic Mean transfer function estimate, 96
associates, 186
associative, 5, 186, 188, 267, 268, 284, 285, 304
asymmetric, 333
auto-correlation, 12, 13, 48
auto-power spectrum, 361
auto-spectral density, 14, 16
autocorrelation, 298
autocorrelation function, 18
autocorrelation kernel, 298
autocorrelation operator, 18, 20, 298
Avant-Garde, vi
average, 92, 274
average power, 44
AWGN, 75
Axiom of Continuity, 146
B-spline, 34, 349, 350, 356, 357, 359, 360, 367, 370, 374–376
B-spline dyadic decomposition, 367, 373
B-spline general form, 354
B-spline partition of unity,

<p>360 B-spline recursion, 352 B-spline scaling coefficients, 370 B-splines, 361 Bak, Per, 107 ball, 260 Banach algebra, 231 Banach space, 202, 203 basis, 20, 199–201, 208, 360 frame, 220 orthogonal, 207 orthonormal, 207 Riesz, 215 tight frame, 220 basis sequence, 359 Battle-Lemarié orthogonalization, 219 Bayesian, 63 Bayesian estimate, 64 Bessel's Equality, 205, 206, 207, 211 Bessel's Inequality, 213 Bessel's inequality, 206 Best Approximation Theorem, 206, 206 biased, 96 bijection, 213 bijective, xi, 213, 292 bilinear functionals, 240 binomial coefficient, 345, 346 Binomial Theorem, 5 binomial theorem, 371 biorthogonal, 215, 217, 218 Birkhoff orthogonality, 256, 258, 258 Bohr, Harald, 267 Boole's inequality, 149 Boolean, 147, 149 Boolean algebra, 152 Boolean lattice, 146, 149 Borel algebra on \mathbb{R}, 152 Borel measure, 309 Borel sets, 309 bounded, xi, 148, 203, 289 bounded bijective, 216 bounded lattice, 149 bounded linear operators, 289 bounded operator, 289, 294 C star algebra, 231 calculus, 185 Carl Spitzweg, 379 Cartesian product, x Cauchy Schwartz inequality, 103, 104 Cauchy-Bunyakovsky-Schwarz Inequality, 234 Cauchy-Schwartz inequality, 9</p>	<p>Cauchy-Schwarz Inequality, 89, 234, 234, 237, 238, 278 Cauchy-Schwarz inequality, 221, 278 Cauchy-Schwarz Inequality for inner-product spaces, 278 causal, 334, 334 cdf, 25, 156, 156, 157 Chain Rule, 351 chain rule, 27, 179 channel capacity, 124 channel coding, 124 channel statistics, 124 characteristic function, x Chebyshev's inequality, 7 chi-square distribution, 164 Chi-squared distribution, 164 chi-squared distribution, 164 Chi-squared distribution with n degrees of freedom, 165, 165 chi-squared distribution with n degrees of freedom, 165 closed interval, 259, 260, 260 closed support, 354 closure, 201, 354 coefficient functionals, 202, 202 coefficients, 335 coherence, 115, 116 communication system, 110, 121 communications additive noise cross-correlation, 55 communications additive noise model, 94, 107 Communications additive noise system, 55 communications LTI additive noise model, 111, 119 commutative, 12, 186, 193, 284, 329, 330 commutative group, 186 compact, 171 compact support, 367 comparable, 259 comparison test, 363 complement, x complete, 201 complete metric space, 201 complete set, 201 complex, 104 complex coherence, 116 complex exponential, 357 complex linear space, 186 complex transmissibility, 98, 98, 116</p>	<p>complex-valued, 98 concave, 260, 260, 261, 261, 262, 262, 269, 271 conditional cumulative distribution function, 156, 156 conditional probability, 149 conditional probability density function, 156, 156 conditional probability of x given y, 148 conjuage symmetric, 18 conjugate linear, 228, 229, 295 conjugate pairs, 335 conjugate quadrature filter, 337, 337 conjugate quadrature filter condition, 337 Conjugate quadrature filters, 337 conjugate symmetric, 13, 15, 18, 48, 172, 233, 234, 242, 243, 246 conjugate symmetric property, 310 conjugate symmetry, 19 conjugate-symmetric, 15 consistent, 92 constant, 12, 43, 64 constraint, 122 Continuity, 252 continuity, 157, 158 continuous, xi, 25, 150, 157, 231, 237, 252, 269, 270, 284, 356, 362, 369 Continuous data whitening, 87 continuous point spectrum, 357 continuous spectrum, 170 contradiction, 188 convergence, 363 convergent, 44, 203 converges, 363–365 Convex, 262 convex, 5, 6, 252, 254, 260, 261, 261, 262, 262, 269–271 functional, 261 strictly, 261 convex function, 261, 262 convex set, 260, 260–262 convolution, 31, 32, 35, 36, 38, 45, 57, 164, 312, 329, 329, 330, 343 convolution operation, 312 convolution operator, 329, 330 Convolution Theorem, 38, 358 convolution theorem, 312, 317, 332</p>
---	---	--

coordinate functionals, 202	eigenvector, 169	bound, 220
coordinates, 199	epigraph, 261	optimal upper frame
correlated, 52, 94, 95	equal, 281	bound, 220
cost function, 63, 107	equivalent, 203	ordered linear space,
countably infinite, 171, 202	even/odd die toss, 151	190
counting measure, xi	event, 147	orthogonal basis, 208
CQF, 337, 337, 338	exact frame, 220	orthogonal in the
CQF condition, 337, 339	expansion, 199, 202	Pythagorean sense, 258
CQF theorem, 337, 340	Fourier coefficients, 208	orthogonal in the sense
Cramér-Rao bound, 86	Fourier expansion, 208	of Birkhoff, 258
Cramér-Rao lower bound, 84	Fourier series, 208	orthogonal in the sense
criterion, 122	frame, 220	of James, 256
critical point, 26, 26, 28, 29	frame bounds, 220	orthonormal basis, 208
cross spectral density, 14	Fredholm operator, 171	outcome, 147
cross-correlation, 12, 13, 18, 18	fundamental, 201	Parseval frame, 220
CS Inequality, 224	Gaussian distribution,	point spectrum, 170
CSD, 14	162, 167	positive, 190, 308
cummulative distribution function, 156, 156	gradient of y with respect to x , 173	positive cone, 190
cumulative distribution function, 25, 150	gradient of y^T with respect to x , 173	positive part, 192
decision region, 139, 140	half-open interval, 259,	power mean, 270
decreasing, 269, 366	260	probability space, 147,
definitions	Hamel basis, 199	152
C^* algebra, 231, 232	hermitian, 228	projection, 228
*-algebra, 12, 15, 35–37,	Hessian matrix, 183	Pythagorean orthogonality, 258
228, 320	hypograph, 261	real linear space, 186
n th order cardinal B-spline, 343	inner product space,	residual spectrum, 170
absolute value, 192	233	resolvent, 227
algebra, 227	intervals on lattices, 259	resolvent set, 170
Banach algebra, 231	intervals on ordered sets, 259	Riesz basis, 215
biorthogonal, 215	isometric, 300	Riesz space, 190
Birkhoff orthogonality, 258	isosceles orthogonality, 256	scalars, 186
bounded, 289	James orthogonality, 256	Schauder basis, 202
bounded linear operators, 289	linear basis, 199	self-adjoint, 228
C star algebra, 231	linear combination, 197	sequence, 329
closed interval, 259, 260	linear space, 186	set projection operators, 139
coefficient functionals, 202	linear subspace, 186	Smith-Barnwell filter, 337
complete, 201	minimum phase, 333	space of all absolutely square summable sequences, 329
complex linear space, 186	multiplicative condition, 231	spans, 197
conditional probability of x given y , 148	negative part, 192	spectral radius, 227
continuous spectrum, 170	norm, 249	spectrum, 170, 227
convex set, 260	norm induced by the inner product, 238	star-algebra, 228
coordinate functionals, 202	normal, 228, 299	summation, 267
coordinates, 199	normal distribution, 162, 167	tight frame, 220
CQF, 337	normalized tight frame, 220	total, 201
difference, 188	normed algebra, 231	underlying set, 186
disjoint, 194	normed linear space, 286	unital, 227
disjoint complement, 195	normed space of linear operators, 286	unitary, 303
eigenspace, 169	open interval, 259, 260	vector lattice, 190
eigenvalue, 169	operator norm, 286	vector space, 186
	optimal lower frame	vectors, 186

differentiable, 26
 differentiable function, 26, 28, 29
 differential operator, 357
 dilation, 306
 dilation equation, 370, 373
 dimension, 199
 direction, 72
 Dirichlet Lambda function, 363, 364
 Discrete data whitening, 87
 Discrete Time Fourier Series, xii
 Discrete Time Fourier Transform, xii, 14, 319
 discrete-time Fourier transform, 319, 319–321, 325
 discriminate, 236
 disjoint, 194
 disjoint complement, 195
 distance, 251
 distributes, 186
 distributive, 12, 35–37, 146–149, 188, 228–230, 233, 234, 295, 320, 331
 distributive pseudocomplemented lattice, 146
 domain, x
 Double coin toss, 150, 150
 DTFT, 14, 15, 93, 203, 320, 324, 327, 337–339, 371
 DTFT periodicity, 319
 dyadic decomposition, 367
 efficient, 81, 84, 85, 87
 Eigendecomposition, 222
 eigenspace, 169
 eigenvalue, 169, 222
 eigenvector, 169, 222
 empty set, xi
 energy, 333
 energy spectral density, 44
 ensemble average, 44
 epigraph, 261
 equal, 281
 equalities
 Bessel's, 205
 equality
 triangle, 250
 equality by definition, x
 equality relation, x
 equations
 parallelogram law, 241
 equivalent, 203, 203, 215
 ergodic in the mean, 44, 44, 45
 estimate, 64, 93, 123
 estimation theory, 63
 Euclidean metric, 254
 Euclidean space, 260
 Euler formulas, 317, 358
 even, 324
 even/odd die, 152
 even/odd die probability space, 151
 even/odd die toss, 151
 event, 147, 147, 148
 events, 147
 exact frame, 220
 examples
 2 coefficient case, 372
 Double coin toss, 150
 even/odd die, 152
 even/odd die toss, 151
 Mercedes Frame, 223
 Peace Frame, 223
 Rectangular pulse, 317
 rectangular pulse, 316, 358
 single coin toss, 150, 152
 The usual norm, 251
 triangle, 317
 tuples in \mathbb{F}^N , 187
 excluded middle, 148, 149
 exclusive OR, xi
 existential quantifier, xi
 exists, 44
 expansion, 199, 202
 expectation, 5, 94
 expectation operator, 3, 3, 4, 6
 false, x
 field, 185, 187, 329
 field of complex numbers, 295
 field of probability, 146
 filter banks, 337
 finite modular lattice, 146
 FontLab Studio, vi
 for each, xi
 Fourier analysis, 356
 fourier analysis, 309
 Fourier coefficients, 208
 Fourier expansion, 208, 208, 209, 212, 213, 214
 Fourier kernel, 309
 Fourier Series, xi
 Fourier series, 208
 Fourier series expansion, 210
 Fourier Transform, xi, xii, 45, 219, 307, 309, 310, 313, 358
 adjoint, 310
 Fourier transform, 203, 307, 314, 316, 317, 357
 inverse, 310
 Fourier transform magnitude, 358, 359
 Fourier transform scaling factor, 310
 Fourier, Joseph, 309
 frame, 218, 220, 220, 224
 frame bound, 221, 223
 frame bounds, 220, 224
 frame operator, 220, 220, 221, 223
 frames, 198
 Fredholm integral equation of the first kind, 18, 171
 Fredholm integral operators, 305
 Fredholm operator, 171, 171, 172
 Fredholm operators, 171
 Free Software Foundation, vi
 Frequency Response Function, 93
 Frequency Response Identification, 93
 Frequency Shift Keying
 coherent, 127
 FRI, 93
 FSK
 coherent, 127
 function, 26, 155, 156, 281, 309, 344
 even, 324
 step, 343
 function overload, 13
 functional, 233, 262, 295
 functions, xi
 0 order B-spline, 361
 0th order B-spline, 345
 1st order B-spline, 346, 361
 2nd order B-spline, 346, 361
 3rd order B-spline, 349
 4th order B-spline, 349
 5th order B-spline, 349
 6th order B-spline, 349
 7th order B-spline, 350
 n th-order modified Bessel function of the first kind, 166
 nth derivative, 356
 nth moment, 314
 nth order B-spline, 347, 352, 354, 360, 361, 367, 369, 370
 nth-order B-spline, 359
 ϕ -mean, 269
 (ESD), 44
 0 order B-spline, 358
 1st order B-spline, 359
 2nd order B-spline, 359
 absolute value, 193, 234, 249
 adjoint, 230
 anti-derivative, 344
 arithmetic mean, 91, 97, 274
 Arithmetic Mean transfer function estimate, 96
 auto-correlation, 12, 13,

<p>48 auto-power spectrum, 361 auto-spectral density, 16 autocorrelation function, 18 autocorrelation operator, 18 average, 92, 274 average power, 44 B-spline, 34, 349, 350, 356, 357, 359, 360, 367, 370, 374–376 B-splines, 361 Bayesian estimate, 64 binomial coefficient, 345, 346 Borel measure, 309 cdf, 25, 156, 156, 157 chi-square distribution, 164 Chi-squared distribution with n degrees of freedom, 165 coherence, 115, 116 complex coherence, 116 complex exponential, 357 complex transmissibility, 98, 98, 116 conditional cumulative distribution function, 156, 156 conditional probability density function, 156, 156 conjugate quadrature filter, 337 continuous point spectrum, 357 cost function, 63, 107 cross-correlation, 12, 13, 18 cummulative distribution function, 156, 156 cumulative distribution function, 25, 150 dilation equation, 370, 373 Dirichlet Lambda function, 363, 364 Discrete Time Fourier Transform, 319 discrete-time Fourier transform, 319–321 DTFT, 324, 327, 338, 339 eigenvector, 222 energy spectral density, 44 estimate, 64 Fourier kernel, 309</p>	<p>Fourier transform, 314, 316, 317 Fredholm integral equation of the first kind, 171 function, 156 Gamma distribution, 163 gamma function, 163 generalized probability, 146, 147 Geometric mean, 97 geometric mean, 97, 274 geometric mean estimator, 97 Geometric mean transfer function estimate, 97 Harmonic mean, 98, 99 harmonic mean, 274 Harmonic mean transfer function estimate, 98 impulse response, 14, 15, 35, 37, 38, 56, 93 induced norm, 211, 237 inner product, 8, 9, 233, 237, 242, 243, 309, 356 input, 14, 15, 35, 37 integer sequence, 350 Inverse Method transfer function estimate, 96 Jensen's inequality, 5 joint cumulative distribution function, 156, 156 Joint Gaussian distribution, 167 joint probability density function, 156, 156 kernel, 357 Kronecker delta function, 245 kronecker delta function, 325 Least Squares Technique, 96 Least Squares transfer function estimate, 96 linear functional, 291 magnitude, 98 MAP estimate, 64, 75 maximum a-posteriori probability estimate, 64 maximum likelihood estimate, 64 mean, 3, 12, 12, 18, 35 mean square error, 65, 65 Mean square estimate, 64 measure, 3, 146, 147, 150, 156, 157 measure-theoretic, 147 mini-max estimate, 64</p>	<p>ML estimate, 64, 75, 77, 80, 82, 85 MM estimate, 64 MS estimate, 64 non-central chi-squared distribution, 166 noncentral chi-square distribution, 166 noncentral chi-square with n-degrees of freedom, 166 norm, 8, 9 order n B-spline, 350 order n B-spline MRA coefficient sequence, 370 order n B-spline MRA scaling sequence, 370 ordinary coherence, 116 ordinary transmissibility, 98 output, 14, 15, 35, 37 pdf, 156, 156, 157 phase, 98, 104 power spectral density, 16, 44 power sum, 350 probability, 147 probability density function, 3, 25, 34, 150, 156, 156, 164 probability function, 146 quantization noise, 95 quantum probability, 146, 147 random process, 17, 20, 44 random sequence, 11, 11–14, 35, 39, 94 random variable, 3–8, 17, 25, 26, 28, 29, 31, 33, 150, 155, 156, 156, 157, 164 Rayleigh distribution, 167 Rice distribution, 167 Riesz sequence, 215 scalar product, 233 scaling coefficient sequence, 372 scaling function, 369, 370 Scaling transfer function estimate, 99, 99 sequence, 11, 34, 370 set indicator function, 318 signal-to-noise ratio, 95 sine sweep, 93 SNR, 95 spectral power, 16, 44, 95</p>
---	---	--

step function, 343, 343–
 347
 Total Least Squares
 transfer function estimate,
100
 Total least squares trans-
 fer function estimate, 99
 traditional probability,
146, 147
 transfer function esti-
 mate $\hat{H}_k(\omega; \kappa)$, 99
 transfer function esti-
 mate $\hat{H}_c(\omega)$, 100
 translation operator,
315
 Transmissibility, 98
 transmissibility, 95, 97
 transmissibility $\tilde{T}_{xy}(\omega)$,
95
 uniform distribution,
161, 161
 variance, 3, 12, 12, 91
 wavelet, 367
 Z transform, 371
 z transform, 333
 Z-transform, 321
 z-transform, 338
 fundamental, 201
 Fundamental Theorem of
 Calculus, 344, 351
 Fundamental theorem of
 calculus, 157, 159
 Fundamental theorem of lin-
 ear equations, 285
 Gamma distribution, 163,
 164, 165
 gamma function, 163, 163
 Gaussian, 33
 Gaussian distribution, 162,
 162, 167
 Gelfand-Mazur Theorem,
 231
 General ML estimation, 80
 Generalized AM-GM in-
 equality, 274
 generalized arithmetic mean
 geometric mean inequality,
 274
 Generalized associative
 property, 267
 Generalized Parseval's Iden-
 tity, 210
 generalized probability, 146,
 147
 Generalized triangle in-
 equality, 9
 geodesically between, 260
 Geometric mean, 97
 geometric mean, 97, 274
 geometric mean estimator,
 97
 Geometric mean transfer
 function estimate, 97
 Geometric Series, 279
 geometry, 185
 GNU Octave
 for, 375
 step, 375
 Golden Hind, vi
 gradient, 72
 gradient of y with respect to
 x , 173
 gradient of y^T with respect to
 x , 173
 greatest lower bound, xi
 Gutenberg Press, vi
 Hölder's Inequality, 276
 half-open interval, 259, 260
 Hamel bases, 198
 Hamel basis, 199, 199, 201
 Handbook of Algebras, 227
 Hardy, G.H., 267
 harmonic analysis, 309
 Harmonic mean, 98, 99
 harmonic mean, 274
 Harmonic mean transfer
 function estimate, 98
 hermitian, 228, 228, 298
 hermitian components, 230
 Hermitian representation,
230
 Hermitian symmetric, 313
 hermitian symmetric, 298
 Hessian, 73
 Hessian matrix, 182, 183
 Heuristic, vi
 high-pass filter, 337
 Hilbert space, 208, 210, 212,
 214, 215, 218, 220, 309
 homogeneous, 5, 208, 209,
 212, 216, 218, 233, 234, 238,
 242, 243, 245, 249, 255, 282,
 283, 285, 286
 Housman, Alfred Edward, vii
 hypograph, 261
 hypotenuse, 258
 identities
 Fourier expansion, 208,
 209
 Parseval frame, 208, 209
 identity, 186, 281
 identity element, 281
 identity operator, 122, 176,
 281, 282
 if, xi
 if and only if, xi
 image, x
 image set, 283, 285, 296, 297,
 299, 300, 304
 imaginary part, xi, 229
 implied by, xi

implies, xi
 implies and is implied by, xi
 impossible, 360
 impulse response, 14, 15, 35,
 37, 38, 56, 93
 inclusive OR, xi
 increasing, 269, 366
 independence, 77, 80, 151
 independent, 7, 31, 34, 81,
 124, 148, 150, 164
 indicator function, x
 induced norm, 211, 237, 238
 induces, 251
 induction, 350, 352, 360
 inequalities
 AM-GM, 274
 Cauchy-Bunyakovsky-
 Schwarz, 234, 278
 Cauchy-Schwarz, 234,
 278
 Hölder, 276
 Jensen's, 262
 Minkowski, 238
 Minkowski (sequences),
 276
 quadratic discriminant,
 236
 Young, 275
 inequality
 Bessel's, 206
 triangle, 246, 249, 286
 infinite sum, 197
 injective, xi, 284
 inner product, 8, 9, 185, 233,
 237, 239, 242, 243, 309, 356
 Polarization Identity,
 240
 uniqueness, 239
 inner product space, 8, 9,
 185, 204–207, 215, 233, 234,
 237, 238, 241, 245, 246, 256,
 356
 inner-product, xi
 inner-product space, 256,
 258
 input, 14, 15, 35, 37
 inside, 334
 instantaneous error, 72
 integer sequence, 350
 integral, 356
 integral test, 364
 intersection, x, 147
 interval, 259, 260
 intervals on lattices, 259
 intervals on ordered sets, 259
 into, 213, 214
 inverse, 186, 221, 223, 372
 inverse DTFT, 325, 339
 Inverse Fourier transform,
310
 inverse Fourier Transform,

<p>307 Inverse Method transfer function estimate, 96 Inverse probability integral transform, 25 invertible, 26, 150, 203, 220, 228 involuntary, 295 involution, 228, 228, 232 involutory, 15, 228–230 irreflexive ordering relation, xi isometric, 209, 217, 218, 292, 300, 300, 311 isometric operator, 299, 301–303 isometry, 300 isomorphic, 213 isosceles, 256 Isosceles orthogonality, 256 isosceles orthogonality, 256, 256–258</p> <p>Jacobian matrix, 173 James orthogonality, 256 Jensen's Inequality, 6, 262, 269, 274 Jensen's inequality, 5, 6, 6 join, xi joint cumulative distribution function, 156, 156 Joint Gaussian distribution, 167 joint probability density function, 156, 156</p> <p>Kaneyoshi, Urabe, 379 Karhunen-Loève Expansion, 20 Kenko, Yoshida, 379 kernel, 18, 357 Kolmogorov's inequality, 7 Kronecker delta function, 245 kronecker delta function, 176, 325 kronecker product, 174 kronecker products, 174</p> <p>l'Hôpital's rule, 102, 105, 106, 272, 363, 365, 369 lattice, 190, 259, 260 lattice with negation, 146 Laurent series, 330 Law of the Unconscious Statistician, 6 Least squares, 69 least squares, 69, 71 Least squares estimation, 107 least squares estimation, 72</p>	<p>Least squares estimations, 107 Least Squares Technique, 96 Least Squares transfer function estimate, 96 least upper bound, xi least-squares, 107, 110 Lebesgue integral, 152 Lebesgue integration, 354, 355 Lebesgue square-integrable functions, 309 left distributive, 5, 114, 285 left inverse, 122 Leibnitz GPR, 340 Leibniz Integration Rule, 352 Leibniz, Gottfried, ix, 281 length, 251 linear, 4, 5, 35, 36, 54, 55, 57, 94, 104, 282, 282, 334 linear basis, 199 linear bounded, xi linear combination, 197 linear combinations, 198 linear functional, 291 linear independence, 204 linear operator, 4, 5, 52, 54, 55, 216, 218 linear operators, 214, 282, 291 linear ops., 214, 218 linear space, 185, 186, 186, 187, 197, 199, 227, 233, 239, 249, 255, 259–262, 281 linear spaces, 281 linear span, 197 linear subspace, 186, 197 linear time invariant, 37, 41, 95, 107, 111, 112, 357 linear time invariant systems, 45 linear time-invariant, 57 linearity, 5, 19, 27, 30, 31, 45, 51, 55–57, 91, 108, 283, 346–348, 365, 368 Linearity of E, 4 linearly dependent, 197, 199–201 linearly independent, 197, 199–201, 204 Lippmann, Bernard A., 162 Liquid Crystal, vi low-pass filter, 337 lower bounded, 149 LTI, 13, 15, 35–38, 45, 56–59, 95, 96, 107, 108, 111–113, 119, 357 operations on non-stationary random processes, 41 operations on WSS ran-</p> <p>dom processes, 43 Machiavelli, Niccolò, 379 magnitude, 98 MAP, 64, 75 MAP estimate, 64, 75, 77 maps to, x Markov's inequality, 6, 7 matched filter, 88 MatLab, 370 matrix rotation, 307 matrix calculus, 173 matrix:quadratic form, 178, 179 Maxima, 347, 349, 350 maximal likelihood (ML), 124 maximum a-posteriori, 64 maximum a-posteriori probability estimate, 64 maximum a-posteriori probability estimation, 75 maximum likelihood, 64 maximum likelihood estimate, 64 maximum likelihood estimation, 75, 80 amplitude, 81 general, 80 phase, 84 Mazur-Ulam theorem, 292 mean, 3, 3, 12, 12, 18, 35 mean square error, 65, 65 Mean square estimate, 64 measurable space, 147 measure, 3, 146, 147, 150, 156, 157 measure space, 147, 148 measure-theoretic, 147 measurement additive noise cross-correlation, 55 measurement additive noise model, 94, 107 Measurement additive noise system, 55 measurement LTI additive noise model, 111, 119 measurement system, 110 meet, xi memoryless, 124 Mercedes Frame, 223 Mercer's Theorem, 22, 172 metric, xi, 185, 251, 252 Euclidean, 254 generated by norm, 252 induced by norm, 252 parabolic, 254 sup, 254 taxi-cab, 254 metric induced by the norm, 252</p>
--	--

metric linear space, 185, 251, 252, 254
 metric space, 185, 252, 259, 260
 mini-max estimate, 64
 Minimum mean square estimation, 67
 Minimum phase, 341
 minimum phase, 38, 49, 333, 333, 334
 energy, 333
 Minkowski addition, 139
 Minkowski's Inequality, 270
 Minkowski's inequality, 238
 Minkowski's Inequality for sequences, 276
 ML, 64, 75
 ML amplitude estimation, 81
 ML estimate, 64, 75, 77, 80, 82, 85
 ML estimation of a function of a parameter, 85
 ML phase estimation, 84
 MM estimate, 64
 modular, 146
 monotone, 146, 148, 150
 monotonically decreasing, 101
 MRA, 370
 MRA system, 370, 372
 MS estimate, 64
 multiplicative condition, 231
 Multiresolution Analysis, 367
 multiresolution analysis, 358, 369
 mutually exclusive, 149, 152
 negative part, 192, 193–195
 Neumann Expansion Theorem, 293
 Newton's method, 72
 no input noise, 113
 no output noise, 113
 noise
 colored, 87
 non-central chi-squared distribution, 166
 non-contradiction, 148
 non-isometric, 238
 non-isotropic, 233, 234, 241, 243, 246
 non-linear, 54, 55, 94, 110
 non-negative, 19, 20, 233, 238, 241, 243, 246, 270, 286, 354, 360
 non-positive, 236
 non-stationary, 44
 non-structured, 123
 non-zero, 372
 Noncentral chi-square distribution, 166

noncentral chi-square distribution, 166
 noncentral chi-square with n -degrees of freedom, 166
 nondegenerate, 146, 148, 151, 152, 249, 255, 286
 nonnegative, 146
 nonparametric, 63
 norm, 8, 9, 185, 235, 237–239, 249, 249, 251, 252, 255, 286
 induced by inner product, 238
 Polarization Identity, 240
 usual, 251
 norm induced by the inner product, 238, 238
 normal, 171, 228, 299, 299, 300
 normal distribution, 162, 162, 167
 normal operator, 299, 303
 normalized, 146, 148, 151
 normalized tight frame, 220
 normed algebra, 231, 231, 232
 normed linear space, 185, 239, 246, 249–252, 254, 256, 258, 286
 normed linear spaces, 300
 normed space of linear operators, 286
 NOT, x
 not ergodic in the mean, 44
 not included
 metric, 252
 not invertible, 26
 not necessarily linear, 37, 51
 not symmetric, 355
 not total, 211
 null space, x, 169, 283–285, 294, 296, 297, 299, 300, 304
 O₆ lattice, 147
 Octave, 370
 one sided shift operator, 302
 only if, xi
 open ball, 252
 open interval, 259, 260
 operations
 0 order B-spline dyadic decomposition, 368
 1st order B-spline dyadic decomposition, 369
 2nd order B-spline dyadic decomposition, 369
 adjoint, 291, 291
 auto-spectral density, 14
 autocorrelation operator, 20
 closure, 354

communications additive noise model, 94
 communications LTI additive noise model, 111, 119
 Continuous data whitening, 87
 convolution, 31, 32, 35, 36, 38, 57, 164, 329, 330, 343
 convolution operation, 312
 cross spectral density, 14
 CSD, 14
 detection, 63
 differential operator, 357
 Discrete data whitening, 87
 Discrete Time Fourier Series, xii
 Discrete Time Fourier Transform, xii, 14
 discrete-time Fourier transform, 319
 DTFT, 14, 15, 93, 203, 320, 337, 339, 371
 Eigendecomposition, 222
 ensemble average, 44
 estimate, 93
 expectation, 5, 94
 expectation operator, 3, 3, 4, 6
 Fourier analysis, 356
 Fourier Series, xi
 Fourier Transform, xi, xii, 45, 219, 307, 310, 313
 Fourier transform, 203, 357
 Fourier transform magnitude, 358, 359
 frame operator, 220, 220, 221, 223
 Fredholm integral equation of the first kind, 18
 Frequency Response Function, 93
 Frequency Response Identification, 93
 FRF, 93
 identity operator, 282
 imaginary part, 229
 integral, 356
 inverse, 221, 223, 372
 inverse Fourier Transform, 307
 involution, 228
 kernel, 18
 Least squares estimation, 107

<p>Least squares estimations, 107 Lebesgue integral, 152 Lebesgue integration, 354, 355 linear operator, 4, 5, 52, 54, 55 linear operators, 291 linear span, 197 measure, 147 measurement additive noise model, 94 measurement LTI additive noise model, 111, 119 operator, 220, 281 power spectral density, 14 project, 356 projection, 296 PSD, 14 rationalizing the denominator, 106 real part, 229 sampling, 4 Sum, 32 summation operator, 262 support, 354 system identification, 93 time average, 44 unitary Fourier Transform, 310 variance, 44 z-domain cross spectral density, 14 z-domain power spectral density, 14 Z-Transform, xii z-transform, 330, 330 Z-transform operator, 14 operator, 185, 220, 281 adjoint, 229 autocorrelation, 298 bounded, 289 definition, 281 delay, 306 dilation, 306 identity, 281 isometric, 299, 301–303 linear, 282 norm, 286 normal, 299, 300, 303 null space, 296 positive, 308 projection, 296 range, 296 self-adjoint, 298 shift, 302 translation, 305 unbounded, 289</p>	<p>unitary, 299, 303, 304, 311 operator adjoint, 294 operator norm, xi, 286 operator star-algebra, 295 optimal lower frame bound, 220 optimal receiver, 75 optimal upper frame bound, 220 order, x, xi order n B-spline, 350 order n B-spline MRA coefficient sequence, 370 order n B-spline MRA scaling sequence, 370 order relation, 259 ordered linear space, 190 ordered pair, x ordered set, 259, 260 ordinary coherence, 116 ordinary transmissibility, 98 orthocomplemented lattice, 148 orthogonal, 19, 204, 207, 208, 245, 245, 246, 356 orthogonal basis, 208 orthogonal in the Pythagorean sense, 258 orthogonal in the sense of Birkhoff, 258 orthogonal in the sense of James, 256 orthogonality, 204, 256, 258 Birkhoff, 258 inner product space, 245 isosceles, 256 James, 256 Pythagorean, 258 orthomodular, 146 orthomodular lattice, 146 orthonomal, 245 orthonormal, 205–211, 224, 245 orthonormal bases, 198 orthonormal basis, 126, 208, 212–217, 219, 366, 367 Orthonormal decomposition, 88 orthonormal expansion, 208, 212 orthonormal quadrature conditions, 325 orthonormality, 205, 216, 224 orthonormal basis, 209 outcome, 147 outcomes, 147 output, 14, 15, 35, 37 over estimate, 96 over-estimated, 112 PAM, 134</p>	<p>parabolic metric, 254 parallelogram, 240, 256 Parallelogram Law, 9 Parallelogram law, 241, 244 parallelogram law, 239–241, 243 parametric, 63 Parseval frame, 208, 209, 220, 220 Parseval frames, 198 Parseval's equation, 311 Parseval's Identity, 210, 212, 214, 220 Parseval's Theorem, 16 partition, 149, 150 partition of unity, 34, 148, 360, 361, 372, 373 Pascal's identity, 348 Pascal, Blaise, 145 pdf, 156, 156, 157 Peace Frame, 223 Per Enflo, 202 Perfect reconstruction, 208 periodic, 279, 319, 362 periodicity, 362 phase, 98, 104 Phase Shift Keying, 131 Plancheral's formula, 45, 81 Plancherel's formula, 311 Poincaré, Henri, 162 point spectrum, 170 pointwise addition, 187 Polar Identity, 237, 238, 257 polar identity, 205 Polarization Identities, 240 Polarization Identity, 239, 241–243 polarization identity, 247, 257 pole, 341 poles, 49, 335 polynomial approximation, 71 Popper, Karl, 93 positive, 19, 20, 190, 308 positive cone, 190 positive definite, 19, 20 positive part, 192, 193–195 power mean, 270, 270, 274 power rule, 365 power set, xi power spectral density, 14, 16, 44 power sum, 350 Power Sums, 279 probability, 147 probability density function, 3, 25, 34, 150, 156, 156, 164 probability function, 146 Probability integral transform, 25</p>
--	--	--

<p>probability measure, 147 probability space, 3–5, 8, 9, 11, 17, 33, 147, 147–151, 152, 152, 156, 157, 164 probability space, 152 product rule, 175, 178, 353 project, 356 projection, 228, 296 projection operator, 296 projection statistics Additive Gaussian noise channel, 78 Additive noise channel, 77 Additive white Gaussian noise channel, 79 Additive white noise channel, 79 projections, 80 proper subset, x proper superset, x properties σ-additive, 146, 152 σ-additivity, 146 absolute value, x absolutely summable, 15, 330, 331 additive, 19, 146–149, 151, 233, 234, 243, 245, 246, 282, 285 additive identity, 12, 188 additive inverse, 188 additivity, 19, 146, 147, 224, 237, 294 admissibility condition, 372 admissibility equation, 372 affine, 170, 261, 292 algebra of sets, xi AND, xi anti-symmetric, 13, 48, 313 anti-symmetric, 48 antiautomorphic, 14, 228, 229, 234, 295 antitone, 148, 151 associates, 186 associative, 5, 186, 188, 267, 268, 284, 285, 304 AWGN, 75 Bayesian, 63 biased, 96 bijection, 213 bijective, 213, 292 biorthogonal, 217, 218 Birkhoff orthogonality, 258 Boole's inequality, 149 Boolean, 147, 149 Boolean lattice, 149</p>	<p>bounded, 148, 203 bounded bijective, 216 Cartesian product, x causal, 334 characteristic function, x closed support, 354 closure, 201 commutative, 12, 186, 193, 284, 329, 330 compact support, 367 comparable, 259 complement, x complex, 104 complex-valued, 98 concave, 260, 260, 261, 261, 262, 269, 271 conjugate linear, 228, 295 conjugate quadrature filter condition, 337 conjugate symmetric, 13, 15, 18, 48, 172, 233, 234, 242, 243, 246 conjugate symmetry, 19 conjugate-symmetric, 15 consistent, 92 constant, 12, 43, 64 Continuity, 252 continuity, 157, 158 continuous, 25, 150, 157, 237, 252, 269, 270, 284, 356, 362, 369 convergence, 363 convergent, 44, 203 converges, 363–365 convex, 5, 6, 252, 254, 260, 261, 261, 262, 262, 269–271 correlated, 52, 94, 95 countably infinite, 171, 202 counting measure, xi CQF condition, 337, 339 critical point, 26, 26, 28, 29 decreasing, 269, 366 dense, 201, 202 difference, x differentiable, 26 differentiable function, 28, 29 distributes, 186 distributive, 12, 35–37, 146–149, 188, 228–230, 233, 234, 295, 320, 331 domain, x dyadic decomposition, 367 efficient, 81, 84, 85, 87</p>	empty set, xi equality by definition, x equality relation, x equivalent, 203, 215 ergodic in the mean, 44, 45 even, 324 excluded middle, 148, 149 existential quantifier, xi exists, 44 false, x for each, xi Fredholm operator, 171, 172 Fredholm operators, 171 Gamma distribution, 164 Gaussian, 33 geodesically between, 260 greatest lower bound, xi hermitian, 228, 298 Hermitian symmetric, 313 hermitian symmetric, 298 homogeneous, 5, 208, 209, 212, 216, 218, 233, 234, 238, 242, 243, 245, 249, 255, 282, 283, 285, 286 hypotenuse, 258 identity, 186 identity operator, 176 if, xi if and only if, xi image, x imaginary part, xi implied by, xi implies, xi implies and is implied by, xi impossible, 360 inclusive OR, xi increasing, 269, 366 independence, 77, 80, 151 independent, 7, 31, 34, 81, 148, 150, 164 indicator function, x injective, 284 inner-product, xi inside, 334 intersection, x into, 213, 214 invertible, 26, 150, 203, 220, 228 involuntary, 295 involution, 228, 232 involutory, 15, 228–230
---	--	---

irreflexive ordering relation, xi	nondegenerate, 146, 148, 151, 152, 249, 286	reality condition, 312
isometric, 209, 217, 218, 292, 300, 311	nonnegative, 146	reflexive ordering relation, xi
isomorphic, 213	nonparametric, 63	relation, x
Isosceles orthogonality, 256	normalized, 146, 148, 151	relational and, x
isosceles orthogonality, 256–258	NOT, x	reverse triangle inequality, 250
join, xi	not ergodic in the mean, 44	right distributive, 5, 285
kronecker delta function, 176	not invertible, 26	ring of sets, xi
least upper bound, xi	not necessarily linear, 37, 51	self adjoint, 19, 298
least-squares, 107, 110	not symmetric, 355	self-adjoint, 19, 298, 298
left distributive, 5, 114, 285	not total, 211	separable, 202, 203, 214
linear, 4, 5, 35, 36, 54, 55, 57, 94, 104, 282, 282	null space, x	set of algebras of sets, xi
linear independence, 204	only if, xi	set of rings of sets, xi
linear space, 187	operator norm, xi	set of topologies, xi
linear time invariant, 37, 95, 107, 111, 112, 357	order, x, xi	space of linear transforms, 283
linear time-invariant, 57	ordered pair, x	span, xi
linearity, 5, 19, 27, 30, 31, 45, 51, 55–57, 91, 108, 283, 346–348, 365, 368	orthogonal, 19, 204, 208, 245, 245, 246, 356	spans, 199, 200
linearly dependent, 197, 199–201	orthogonality, 204, 256, 258	stability condition, 218,
linearly independent, 197, 199–201, 204	orthomodular, 146	220 stable, 334
lower bounded, 149	orthonomal, 245	stationary in correlation, 43
LTI, 13, 15, 35–38, 56– 59, 95, 96, 107, 108, 111–113, 119, 357	orthonormal, 205, 206, 208–211, 224, 245	stationary in the 1st moment, 12
maps to, x	orthonormality, 205, 216, 224	stationary in the 2nd moment, 12
meet, xi	over estimate, 96	stationary in the mean, 43, 44
metric, xi	over-estimated, 112	Strang-Fix condition, 315
Minimum phase, 341	parallelogram, 240, 256	strictly concave, 262
minimum phase, 38, 333, 334	parallelogram law, 243	strictly convex, 261, 262
modular, 146	parametric, 63	strictly increasing, 270,
monotone, 146, 148, 150	Parseval's Theorem, 16	271 strictly monotone, 150
monotonically decreasing, 101	partition of unity, 34, 148, 360, 361, 372, 373	strictly monotonic, 269,
mutually exclusive, 149, 152	periodic, 279, 319, 362	270 strictly monotonic in-
no input noise, 113	periodicity, 362	creasing, 103
no output noise, 113	pointwise addition, 187	strictly positive, 249
non-contradiction, 148	Polarization Identity, 243	strong convergence, 202, 212
non-isometric, 238	positive, 19, 20	subadditive, 146, 238,
non-isotropic, 233, 234, 241, 243, 246	positive definite, 19, 20	249, 286
non-linear, 54, 55, 94, 110	power set, xi	subset, x
non-negative, 19, 20, 233, 238, 241, 243, 246, 270, 286, 354, 360	proper subset, x	super set, x
non-positive, 236	proper superset, x	symmetric, 13, 48, 252,
non-stationary, 44	pseudo-distributes, 186	313, 322, 354, 355, 362, 366
non-zero, 372	Pythagorean orthogonality, 258	symmetric , 48
	Pythagorean Theorem, 258	symmetric difference, x
	quadratic discriminant inequality, 236	symmetry, 245, 333
	randomness, 155	there exists, xi
	range, x	tight frame, 221
	real, 19, 252, 257, 322, 335	time-invariant, 334
	real part, xi	topology of sets, xi
	real-valued, 15, 20, 98, 313, 320, 321, 324	total, 201, 210, 211, 215,
		216 translation invariant,
		255

triangle inequality, 259 triangle inequality, 249 true, x unbiased, 81, 82, 91, 92, 113 uncorrelated, 20, 51–59, 76, 91, 92, 94–96, 108, 112, 114, 117 under estimate, 96 under estimates, 96 under-estimated, 112 uniformly distributed, 25, 34 union, x unique, 26, 188, 199, 202, 359 unit area, 354 unit length, 302, 304 unitary, 303, 304, 311 universal quantifier, xi upper bounded, 146 vanishing 0th moment, 372 vector norm, xi von Neumann-Jordan condition, 241 wide sense stationary, 12, 12–14, 37, 43, 43, 44, 52, 91, 92, 95 wide-sense stationary, 112 WSS, 12, 13, 15, 36–38, 43, 43, 51, 53–59, 91, 95, 96, 108, 112 zero at -1, 372 zero measurement error, 95 zero measurement noise, 95 zero-mean, 7, 51–59, 108 PSD, 14 pseudo-distributes, 186 PSF, 315, 316 PSK, 131 pstricks, vi Pulse Amplitude Modulation, 134 Pythagorean orthogonality, 256, 258, 258 Pythagorean Theorem, 204, 205, 207, 209, 216, 246, 246, 258 Pythagorean theorem, 204 QAM, 129 quadratic discriminate inequality, 236 Quadratic Equation, 115 quadratic equation, 236 quadratic form, 178, 179 Quadrature Amplitude Modulation, 129	quantization noise, 95 quantum probability, 146, 147 quotes Abel, 278 Abel, Niels Henrik, 379 Aristotle, 11, 17 Bak, Per, 107 Bohr, Harald, 267 Descartes, René, ix Fourier, Joseph, 309 Hardy, G.H., 267 Housman, Alfred Edward, vii Kaneyoshi, Urabe, 379 Kenko, Yoshida, 379 Leibniz, Gottfried, ix, 281 Lippmann, Bernard A., 162 Machiavelli, Niccolò, 379 Pascal, Blaise, 145 Poincaré, Henri, 162 Popper, Karl, 93 Russell, Bertrand, vii Stravinsky, Igor, vii Ulam, Stanislaus M., 292 Quotient Rule, 114, 115 quotient rule, 365 quotient structures, 259 random process, 17, 17, 20, 44 random sequence, 11, 11–14, 35, 39, 94 random variable, 3–8, 17, 17, 25, 26, 28, 29, 31, 33, 150, 155, 156, 156, 157, 164 randomness, 155 range, x range space, 294 rational expression, 38, 49 rationalizing factor, 106 rationalizing the denominator, 106 Rayleigh distribution, 167, 167 real, 19, 252, 257, 322, 335 real interval, 157 real linear space, 186 real numbers, 251 real part, xi, 229 real-valued, 15, 20, 98, 313, 320, 321, 324 reality condition, 312 Rectangular pulse, 317 rectangular pulse, 316, 358 recursively, 71 reflection, 292 reflexive ordering relation, xi regular, 171	relation, x, 26, 281 relational and, x relations, xi order relation, 259 residual spectrum, 170 resolvent, 227 resolvent set, 170, 171 Reverse Triangle Inequality, 250, 252 reverse triangle inequality, 250 Rice distribution, 167, 167 Riesz bases, 198 Riesz basis, 215, 215–219, 358, 366, 367 Riesz sequence, 215, 217, 369 Riesz space, 190, 190, 192–195 Riesz-Fischer Theorem, 212 Riesz-Fischer Thm., 213, 214 right distributive, 5, 285 right inverse, 122 ring of complex square $n \times n$ matrices, 295 ring of sets, xi Robinson's Energy Delay Theorem, 333 rotation matrix, 307 Russell, Bertrand, vii sampling, 4 scalar product, 233 scalars, 186 scaling coefficient sequence, 372 scaling function, 99, 369, 370 scaling parameter, 99 Scaling transfer function estimate, 99, 99 Schauder bases, 198, 203 Schauder basis, 202, 202, 203, 208, 214 self adjoint, 19, 298 self-adjoint, 19, 228, 298, 298 semilinear, 229 separable, 202, 203, 214 separable Hilbert space, 213, 215, 217, 329 separable Hilbert spaces, 213 sequence, 11, 34, 251, 329, 370 sequences, 337 set, 147, 185, 260 set indicator, 343 set indicator function, 318 set of algebras of sets, xi set of rings of sets, xi set of topologies, xi set projection operators, 139 sets open ball, 252 real numbers, 251
---	--	--

shift operator, 302	strictly monotonic, 269, 270	of x given y , 148
shift relation, 316, 317	strictly monotonic increasing, 103	conjugate pairs, 335
sigma-algebra, 147	strictly positive, 249, 255	conjugate quadrature
Signal matching, 88	strong convergence, 202, 212	filter, 337
signal to noise ratio, 88	structured, 123	Conjugate quadrature
signal-to-noise ratio, 95	structures	filters, 337
sinc, 316, 317	C^* algebra, 231	convex function, 261,
sine sweep, 93	C^* -algebra, 232	262
single coin toss, 150, 152	$*$ -algebra, 12, 15, 35–37,	convex set, 260, 260–262
Smith-Barnwell filter, 337	228, 228–230, 295, 320	convolution operator,
SNR, 95	N -tuple, 187, 249	329, 330
source code, 375	$*$ -algebras, 295	coordinates, 199
space, 147	σ -algebra, 147, 152, 154	CQF, 337, 338
inner product, 233	absolute value, 194, 195	critical point, 26
linear, 185	additive identity ele-	differentiable function,
metric, 252	ment, 187, 188	26
normed vector, 249	additive inverse, 188	discrete-time Fourier
vector, 185	adjoint, 294, 295	transform, 325
space of all absolutely square	algebra, 185, 227, 227,	distributive pseudo-
Lebesgue integrable functions, 329	228	complemented lattice, 146
space of all absolutely square	algebraic, 185	Double coin toss, 150
summable sequences, 329	autocorrelation kernel,	epigraph, 261
space of linear transforms, 283	298	Euclidean space, 260
span, xi, 201, 203	autocorrelation opera-	even/odd die probabili-
spans, 197, 199, 200	tor, 298	ity space, 151
spectral power, 16, 44, 95	ball, 260	even/odd die toss, 151
spectral radius, 227	Banach space, 202, 203	event, 147, 147, 148
Spectral Theorem, 171	basis, 20, 199–201, 208,	expansion, 199
spectral theorem, 171	360	field, 185, 187, 329
spectrum, 170, 171, 227	basis sequence, 359	field of complex num-
stability, 334	bijection, 213	bers, 295
stability condition, 218, 220	bijective, 213	field of probability, 146
stable, 334	bilinear functionals, 240	filter banks, 337
standard orthonormal basis, 209	Boolean algebra, 152	finite modular lattice,
star-algebra, 228, 228, 295	Boolean lattice, 146, 149	146
star-algebras, 294, 295	Borel algebra on \mathbb{R} , 152	Fourier Transform, 309
state, 72	Borel sets, 309	frame, 218, 220, 220, 224
stationary in correlation, 43	bounded lattice, 149	frames, 198
stationary in the 1st moment, 12	bounded operator, 294	function, 26, 155, 309,
stationary in the 2nd moment, 12	C star algebra, 231	344
stationary in the mean, 43, 44	calculus, 185	functional, 233, 262, 295
statistical average, 72	closed interval, 259,	geometry, 185
statistics, 124	260, 260	half-open interval, 259,
steepest descent, 72	coefficient functionals,	260
step function, 343, 343–347	202	Hamel bases, 198
step size, 72	coefficients, 335	Hamel basis, 199, 199,
Stifel formula, 368	communication system,	201
Stifel formula, 348	110	high-pass filter, 337
stochastic, 66	communications additive noise model, 107	Hilbert space, 208, 210,
Strang-Fix condition, 315, 315	Communications additive noise system, 55	212, 214, 215, 218, 220, 309
Stravinsky, Igor, vii	complete metric space,	hypograph, 261
strictly concave, 262	201	identity, 186
strictly convex, 261, 261, 262	complete set, 201	identity element, 281
strictly increasing, 270, 271	complex linear space,	image set, 283, 304
strictly monotone, 150	186	infinite sum, 197
	conditional probability,	inner product, 185, 239
	149	inner product space, 8,
	conditional probability	9, 185, 204–207, 215, 233,
		234, 237, 238, 241, 245, 246,
		256, 356
		inner-product space,

- 256, 258
 interval, 259, 260
 intervals on lattices, 259
 intervals on ordered sets, 259
 inverse, 186
 isometry, 300
 lattice, 259, 260
 lattice with negation, 146
 Laurent series, 330
 Lebesgue square-integrable functions, 309
 linear basis, 199
 linear combination, 197
 linear combinations, 198
 linear operator, 216, 218
 linear operators, 214
 linear ops., 214, 218
 linear space, 185, 186, 186, 197, 199, 227, 233, 239, 249, 255, 259–262, 281
 linear spaces, 281
 linear subspace, 186, 197
 low-pass filter, 337
 measurable space, 147
 measure, 147
 measure space, 147, 148
 measurement additive noise model, 107
 Measurement additive noise system, 55
 measurement system, 110
 metric, 185, 252
 metric linear space, 185, 251, 252, 254
 metric space, 185, 252, 259, 260
 MRA, 370
 MRA system, 370, 372
 Multiresolution Analysis, 367
 multiresolution analysis, 358, 369
 negative part, 193–195
 norm, 185, 235, 237–239, 249, 251, 252, 255
 normalized tight frame, 220
 normed algebra, 231, 232
 normed linear space, 185, 239, 246, 249–252, 254, 256, 258
 normed linear spaces, 300
 null space, 284, 304
 O_6 lattice, 147
 open ball, 252
 open interval, 259, 260
 operator, 185
 ordered set, 259, 260
 orthocomplemented lattice, 148
 orthogonal basis, 208
 orthomodular lattice, 146
 orthonormal bases, 198
 orthonormal basis, 208, 212–217, 219, 366, 367
 orthonormal basis, 209
 outcome, 147
 Parseval frame, 220
 Parseval frames, 198
 Parseval's equation, 311
 partition, 149, 150
 Plancherel's formula, 311
 pole, 341
 poles, 335
 positive part, 193–195
 probability space, 3–5, 8, 9, 11, 17, 33, 147, 147–151, 152, 152, 156, 157, 164
 probability space, 152
 quotient structures, 259
 random variable, 156
 real interval, 157
 real linear space, 186
 relation, 26
 resolvent, 227
 Riesz bases, 198
 Riesz basis, 215–219, 358, 366, 367
 Riesz sequence, 217, 369
 Riesz space, 190, 192–195
 ring of complex square $n \times n$ matrices, 295
 scalars, 186
 Schauder bases, 198, 203
 Schauder basis, 202, 202, 203, 208, 214
 separable Hilbert space, 213, 215, 217, 329
 separable Hilbert spaces, 213
 sequence, 251
 sequences, 337
 set, 147, 185, 260
 sigma-algebra, 147
 space, 147
 space of all absolutely square Lebesgue integrable functions, 329
 space of all absolutely square summable sequences, 329
 span, 201, 203
 spectral radius, 227
 spectrum, 227
 standard orthonormal basis, 209
 star-algebra, 228, 295
 support, 354
 system, 13, 37, 51–59, 93, 95–100, 103, 107, 108, 110–114, 116
 tight frame, 220, 221, 223
 tight frames, 198
 topological dual space, 291
 topological linear space, 185, 201
 topology, 147, 185, 197
 total set, 201
 underlying set, 186
 unit vector, 217
 unital *-algebra, 228
 unital algebra, 227
 vector, 199
 vector space, 8, 185, 186
 vector space of random variables, 4
 vectors, 186, 259
 weights, 262
 zero, 341
 zeros, 333, 335
 subadditive, 146, 238, 249, 286
 subset, x
 sufficient statistic, 75
 Sufficient statistic theorem, 75
 Sum, 32
 sum of products, 27, 31, 149, 157
 summation, 267
 Summation around unit circle, 222
 summation operator, 262
 sup metric, 254
 super set, x
 support, 354
 surjective, xi
 symmetric, 13, 48, 252, 313, 322, 354, 355, 362, 366
 conjugate, 18
 symmetric, 48
 symmetric difference, x
 symmetry, 245, 333
 system, 13, 37, 51–59, 93, 95–100, 103, 107, 108, 110–114, 116
 system identification, 93
 taxi-cab metric, 254
 The basis problem, 202
 The Book Worm, 379

The Fourier Series Theorem, 210 The usual norm, 251 Theorem of Reversibility, 123 theorems Affine equations, 177 Alternating Series Test, 366 B-spline dyadic decomposition, 367, 373 B-spline general form, 354 B-spline partition of unity, 360 B-spline recursion, 352 B-spline scaling coefficients, 370 Battle-Lemarié orthogonalization, 219 Bessel's Equality, 205, 206, 207, 211 Bessel's Inequality, 213 Bessel's inequality, 206 Best Approximation Theorem, 206, 206 Binomial Theorem, 5 binomial theorem, 371 Cauchy Schwartz inequality, 103, 104 Cauchy-Schwartz inequality, 9 Cauchy-Schwarz Inequality, 89, 234, 237, 238, 278 Cauchy-Schwarz inequality, 221 chain rule, 27 Chebyshev's inequality, 7 communications additive noise cross-correlation, 55 comparison test, 363 Convolution Theorem, 38, 358 convolution theorem, 312, 317, 332 CQF theorem, 337, 340 CS Inequality, 224 dilation equation, 370 DTFT periodicity, 319 Euler formulas, 317, 358 Fourier expansion, 209, 212, 213, 214 Fourier series expansion, 210 Fundamental Theorem of Calculus, 344, 351 Fundamental theorem of calculus, 157, 159	Fundamental theorem of linear equations, 285 Gelfand-Mazur, 231 General ML estimation, 80 Generalized AM-GM inequality, 274 Generalized associative property, 267 Generalized Parseval's Identity, 210 Generalized triangle inequality, 9 Geometric Series, 279 Hölder's Inequality, 276 Hermitian representation, 230 induced norm, 238 induction, 350, 352, 360 integral test, 364 inverse DTFT, 325, 339 Inverse Fourier transform, 310 Inverse probability integral transform, 25 Jensen's Inequality, 6, 262, 269, 274 Jensen's inequality, 6, 6 Karhunen-Loève Expansion, 20 Kolmogorov's inequality, 7 l'Hôpital's rule, 102, 105, 106, 272, 363, 365, 369 Law of the Unconscious Statistician, 6 Least squares, 69 Leibnitz GPR, 340 Leibniz Integration Rule, 352 Linearity of E, 4 Markov's inequality, 6, 7 Mazur-Ulam theorem, 292 measurement additive noise cross-correlation, 55 Mercer's Theorem, 22, 172 Minimum mean square estimation, 67 Minkowski's Inequality, 270 Minkowski's inequality, 238 Minkowski's Inequality for sequences, 276 ML amplitude estimation, 81 ML estimation of a function of a parameter, 85 ML phase estimation, 84	Neumann Expansion Theorem, 293 operator star-algebra, 295 orthonormal quadrature conditions, 325 Parallelogram Law, 9 Parallelogram law, 241 parallelogram law, 241 Parseval's Identity, 210, 212, 214, 220 Pascal's identity, 348 Perfect reconstruction, 208 Plancheral's formula, 45, 81 Polar Identity, 237, 238, 257 polar identity, 205 Polarization Identities, 240 Polarization Identity, 241–243 polarization identity, 247, 257 power rule, 365 Power Sums, 279 Probability integral transform, 25 PSF, 315, 316 Pythagorean Theorem, 204, 205, 207, 209, 216, 246, 246 Pythagorean theorem, 204 Quadratic Equation, 115 Quotient Rule, 114, 115 quotient rule, 365 Reverse Triangle Inequality, 250, 252 Riesz-Fischer Theorem, 212 Riesz-Fischer Thm., 213, 214 Robinson's Energy Delay Theorem, 333 shift relation, 316, 317 Spectral Theorem, 171 spectral theorem, 171 Stifel formal, 368 Stifel formula, 348 Strang-Fix condition, 315 Sufficient statistic theorem, 75 sum of products, 27, 31, 149, 157 Summation around unit circle, 222 The Fourier Series Theorem, 210
--	---	--

Theorem of Reversibility, 123 triangle inequality, 246 , 249 Wiener-Hopf equations, 68 , 68 Wold's Theorem, 161 Young's Inequality, 275 , 276 there exists, xi tight frame, 220 , 221 , 223 tight frames, 198 time average, 44 time-invariant, 334 , 334 topological dual space, 291 topological linear space, 185 , 201 topological space, 147 , 251 topology, 147 , 185 , 197 topology of sets, xi total, 201 , 201 , 210 , 211 , 215 , 216 Total Least Squares transfer function estimate, 100 Total least squares transfer function estimate, 99 total set, 201 traditional probability, 146 , 147 transfer function estimate $\hat{H}_k(\omega; \kappa)$, 99 transfer function estimate $\hat{H}_c(\omega)$, 100 transform inverse Fourier, 310 translation, 305 translation invariant, 255 translation operator, 315 Transmissibility, 98 transmissibility, 95 , 97 transmissibility $\tilde{T}_{xy}(\omega)$, 95 triangle, 317 triangle inequality, 246 , 249 , 249 , 259 , 286 triangle inequality, 249 triangle-inequality, 256 true, x	tuples in \mathbb{F}^N , 187 Ulam, Stanislaus M., 292 unbiased, 81 , 82 , 91 , 92 , 113 uncorrelated, 20 , 51–59 , 76 , 91 , 92 , 94–96 , 108 , 112 , 114 , 117 under estimate, 96 under estimates, 96 under-estimated, 112 underlying set, 186 uniform distribution, 161 , 161 uniformly distributed, 25 , 34 union, x , 147 unique, 26 , 188 , 199 , 202 , 359 unit area, 354 unit length, 302 , 304 unit vector, 217 unital, 227 unital $*$ -algebra, 228 unital algebra, 227 unitary, 303 , 303 , 304 , 307 , 310 , 311 unitary Fourier Transform, 310 unitary operator, 299 , 303 universal quantifier, xi upper bounded, 146 usual norm, 251 Utopia, vi	vanishing 0th moment, 372 vanishing moments, 314 , 327 variance, 3 , 3 , 12 , 12 , 44 , 91 vector, 199 vector lattice, 190 vector norm, xi vector space, 8 , 185 , 186 vector space of random variables, 4 vectors, 186 , 259 von Neumann-Jordan condition, 241
		wavelet, 367 weighted, 270 weights, 262 whitening filter, 38 , 48 wide sense stationary, 12 , 12–14 , 37 , 43 , 43 , 44 , 52 , 91 , 92 , 95 wide-sense stationary, 112 Wiener-Hopf equations, 68 , 68 Wold's Theorem, 161 Wronskian, 183 WSS, 12 , 12 , 13 , 15 , 36–38 , 43 , 43 , 51 , 53–59 , 91 , 95 , 96 , 108 , 112

Young's Inequality, [275](#), [276](#)

values nth moment, 314 Cramér-Rao bound, 86 Cramér-Rao lower bound, 84 dimension, 199 eigenvalue, 222 frame bound, 221 , 223 frame bounds, 224 MAP estimate, 77 ML estimate, 77 rationalizing factor, 106 scaling function, 99 scaling parameter, 99 sufficient statistic, 75	Z transform, 371 z transform, 333 z-domain cross spectral density, 14 z-domain power spectral density, 14 Z-Transform, xii Z-transform, 321 z-transform, 330 , 330 , 338 Z-transform operator, 14 zero, 341 zero at -1 , 322 , 372 zero measurement error, 95 zero measurement noise, 95 zero-mean, 7 , 51–59 , 108 zeros, 49 , 333 , 335
--	--

License

This document is provided under the terms of the Creative Commons license CC BY-NC-ND 4.0. For an exact statement of the license, see

<https://creativecommons.org/licenses/by-nc-nd/4.0/legalcode>

The icon  appearing throughout this document is based on one that was once at

<https://creativecommons.org/>

where it was stated, "Except where otherwise noted, content on this site is licensed under a Creative

Commons Attribution 4.0 International license."





...last page ...please stop reading ...

