
Frames and Bases Structure and Design

VERSION 0.21







TITLE PAGE Daniel J. Greenhoe page v

title: Frames and Bases Structure and Design

document type: book

series: Mathematical Structure and Design

volume:

author: Daniel J. Greenhoe

VERSION 0.21 version:

2021 March 16 (Tuesday) 08:55am UTC time stamp: Copyright © 2021 Daniel J. Greenhoe copyright:

license: Creative Commons license CC BY-NC-ND 4.0

typesetting engine: X₇IM_FX

document url: https://github.com/dgreenhoe/pdfs/blob/master/msdframes.pdf

https://www.researchgate.net/project/Mathematical-Structure-and-Design

This text was typeset using $X \equiv LAT_E X$, which is part of the $T_E X$ family of typesetting engines, which is arguably the greatest development since the Gutenberg Press. Graphics were rendered using the *pstricks* and related packages, and $LAT_E X$ graphics support.

The main roman, *italic*, and **bold** font typefaces used are all from the *Heuristica* family of typefaces (based on the *Utopia* typeface, released by *Adobe Systems Incorporated*). The math font is XITS from the XITS font project. The font used in quotation boxes is adapted from *Zapf Chancery Medium Italic*, originally from URW++ Design and Development Incorporated. The font used for the text in the title is Adventor (similar to *Avant-Garde*) from the *TEX-Gyre Project*. The font used for the ISBN in the footer of individual pages is LOUID CRYSTAL (*Liquid Crystal*) from *FontLab Studio*. The Latin handwriting font is *Lavi* from the *Free Software Foundation*.

The ship appearing throughout this text is loosely based on the *Golden Hind*, a sixteenth century English galleon famous for circumnavigating the globe.¹



¹ Paine (2000) page 63 ⟨Golden Hind⟩




Tell me of runes to graveThat hold the bursting wave,Or bastions to designFor longer date than mine. ♥

Alfred Edward Housman, English poet (1859–1936) ²



The uninitiated imagine that one must await inspiration in order to create. That is a mistake. I am far from saying that there is no such thing as inspiration; quite the opposite. It is found as a driving force in every kind of human activity, and is in no wise peculiar to artists. But that force is brought into action by an effort, and that effort is work. Just as appetite comes by eating so work brings inspiration, if inspiration is not discernible at the beginning. ♥

Igor Fyodorovich Stravinsky (1882–1971), Russian-born composer ³



As I think about acts of integrity and grace, I realise that there is nothing in my knowledge to compare with Frege's dedication to truth. His entire life's work was on the verge of completion, much of his work had been ignored to the benefit of men infinitely less capable, his second volume was about to be published, and upon finding that his fundamental assumption was in error, he responded with intellectual pleasure clearly submerging any feelings of personal disappointment. It was almost superhuman and a telling indication of that of which men are capable if their dedication is to creative work and knowledge instead of cruder efforts to dominate and be known.

Bertrand Russell (1872–1970), British mathematician, in a 1962 November 23 letter to Dr. van Heijenoort.



page viii Daniel J. Greenhoe Title page

image: http://en.wikipedia.org/wiki/Image:Housman.jpg

image: http://en.wikipedia.org/wiki/Image:Igor_Stravinsky.jpg

4 quote: ## Heijenoort (1967) page 127

image: http://www-history.mcs.st-andrews.ac.uk/PictDisplay/Russell.html



SYMBOLS

rugula XVI. Quae vero praesentem mentis attentionem non requirunt, etiamsi ad conclusionem necessaria sint, illa melius est per brevissimas notas designare quam per integras figuras: ita enim memoria non poterit falli, nec tamen interim cogitatio distrahetur ad haec retinenda, dum aliis deducendis incumbit.



"Rule XVI. As for things which do not require the immediate attention of the mind, however necessary they may be for the conclusion, it is better to represent them by very concise symbols rather than by complete figures. It will thus be impossible for our memory to go wrong, and our mind will not be distracted by having to retain these while it is taken up with deducing other matters."

René Descartes (1596–1650), French philosopher and mathematician ⁵



Gottfried Leibniz (1646–1716), German mathematician, ⁶

Symbol list

symbol	description	
numbers:		
\mathbb{Z}	integers	$\dots, -3, -2, -1, 0, 1, 2, 3, \dots$
W	whole numbers	0, 1, 2, 3,

...continued on next page...

⁵quote: Descartes (1684a) (rugula XVI), translation: Descartes (1684b) (rule XVI), image: Frans Hals (circa 1650), http://en.wikipedia.org/wiki/Descartes, public domain

⁶quote: ☐ Cajori (1993) ⟨paragraph 540⟩, image: http://en.wikipedia.org/wiki/File:Gottfried_Wilhelm_von_Leibniz.jpg, public domain

symbol	description		
N	natural numbers	1, 2, 3,	
\mathbb{Z}^{\dashv}	non-positive integers	$\dots, -3, -2, -1, 0$	
\mathbb{Z}^-	negative integers	\dots , 3, 2, 1, 0 \dots , -3, -2, -1	
\mathbb{Z}_{o}	odd integers	\dots , -3 , -1 , 1 , 3 , \dots	
\mathbb{Z}_{e}	even integers	\dots , -4 , -2 , 0 , 2 , 4 , \dots	
—e ℚ	rational numbers	$\frac{m}{n}$ with $m \in \mathbb{Z}$ and $n \in \mathbb{Z} \setminus 0$	
\mathbb{R}	real numbers	completion of \mathbb{Q}	
R ⊢		_	
\mathbb{R}^{\dashv}	non-negative real numbers	$[0,\infty)$	
	non-positive real numbers	$(-\infty,0]$	
R ⁺	positive real numbers	$(0, \infty)$	
R ⁻	negative real numbers	$(-\infty,0)$	
\mathbb{R}^*	extended real numbers	$\mathbb{R}^* \triangleq \mathbb{R} \cup \{-\infty, \infty\}$	
C	complex numbers	(-ft	
F	arbitrary field	(often either $\mathbb R$ or $\mathbb C$)	
∞	positive infinity		
$-\infty$	negative infinity	0.14150065	
π	pi	3.14159265	
relations:	1		
R	relation		
O	relational and		
$X \times Y$	Cartesian product of X and Y		
(\triangle, ∇)	ordered pair	1	
	absolute value of a complex nu	imber z	
=	equality relation		
≜	equality by definition		
→	maps to		
€	is an element of		
∉	is not an element of		
9 (®)	domain of a relation ®		
$\mathcal{J}(\mathbb{R})$	image of a relation ®		
R (®)	range of a relation ®		
$\mathscr{N}(\mathbb{R})$	null space of a relation ®		
set relations:	ou boot		
<u>_</u>	subset		
-	proper subset		
⊆ ♀ ⊄	super set		
<i>⊋</i> <i>⋆</i>	proper superset		
¥ *	is not a subset of		
•			
_	operations on sets:		
$A \cup B$			
$A \cap B$	set intersection		
$A \triangle B$	set symmetric difference		
$egin{array}{c} A ackslash B \ A^{c} \end{array}$	set difference		
	set complement		
	· set order		
$\mathbb{1}_A(x)$ set indicator function or characteristic function			
logic:	"true" condition		
1	true contaition		



SYMBOL LIST Daniel J. Greenhoe page xi

symbol	description		
0	"false" condition		
¬	logical NOT operation		
\wedge	logical AND operation		
V	logical inclusive OR operation		
\oplus	logical exclusive OR operation		
\Longrightarrow	"implies";	"only if"	
$\begin{array}{c} \Longrightarrow \\ \Longleftrightarrow \\ \Longleftrightarrow \end{array}$	"implied by";	"if"	
\iff	"if and only if";	"implies and is implied by"	
\forall	universal quantifier:	"for each"	
3	existential quantifier:	"there exists"	
order on sets:			
V	join or least upper bound		
\wedge	meet or greatest lower bound		
≤	reflexive ordering relation	"less than or equal to"	
≤ ≥ <	reflexive ordering relation	"greater than or equal to"	
	irreflexive ordering relation	"less than"	
>	irreflexive ordering relation	"greater than"	
measures on s	sets:		
X	order or counting measure of a	set X	
distance spac			
d	metric or distance function		
linear spaces:			
$\ \cdot\ $	vector norm		
•	operator norm		
$\langle \triangle \nabla \rangle$			
$span(m{V})$	span of a linear space V		
algebras:			
\Re	real part of an element in a *-al		
$\mathfrak F$	imaginary part of an element in	ı a *-algebra	
set structures			
T	a topology of sets		
\boldsymbol{R}	a ring of sets		
\boldsymbol{A}	an algebra of sets		
Ø	empty set		
_	2^X power set on a set X		
sets of set stru			
$\mathcal{T}(X)$	set of topologies on a set X		
$\mathscr{R}(X)$	S .		
, ,	$\mathcal{A}(X)$ set of algebras of sets on a set X		
	classes of relations/functions/operators:		
_	2^{XY} set of <i>relations</i> from X to Y		
Y^X	set of $functions$ from X to Y		
$\mathscr{S}_{j}(X,Y)$	set of <i>surjective</i> functions from X to Y		
$\mathscr{F}_{j}(X,Y)$			
	$\mathscr{B}_{j}(X,Y)$ set of <i>bijective</i> functions from X to Y		
$\mathscr{B}(\boldsymbol{X},\boldsymbol{Y})$	$\mathscr{B}(X,Y)$ set of bounded functions/operators from X to Y		
$\mathscr{L}(\pmb{X}, \pmb{Y})$	$\mathscr{L}(X, Y)$ set of <i>linear bounded</i> functions/operators from X to Y		
$\mathscr{C}(\pmb{X},\pmb{Y})$	$\mathscr{C}(\boldsymbol{X},\boldsymbol{Y})$ set of <i>continuous</i> functions/operators from \boldsymbol{X} to \boldsymbol{Y}		
specific transforms/operators:			

...continued on next page...





page xii Daniel J. Greenhoe Symbol List

symbol	description	
F	Fourier Transform operator (Definition I.2 page 196)	
$\mathbf{\hat{F}}$	Fourier Series operator (Definition N.1 page 247)	
Ĕ	F Discrete Time Fourier Series operator (Definition M.1 page 237)	
${f Z}$	Z-Transform operator (Definition J.4 page 208)	
$ ilde{f}(\omega)$	$\tilde{f}(\omega)$ Fourier Transform of a function $f(x) \in L^2_{\mathbb{D}}$	
$reve{x}(\omega)$	Discrete Time Fourier Transform of a sequence $(x_n \in \mathbb{C})_{n \in \mathbb{Z}}$	
$\check{x}(z)$	<i>Z-Transform</i> of a sequence $(x_n \in \mathbb{C})_{n \in \mathbb{Z}}$	

SYMBOL INDEX

P, 193
$T_n(x)$, 178
C, 39
Q, 44
\mathbb{R} , 39
1, 40
D_n , 184
J _n , 193
K _n , 191
V _n , 193
$\bar{\delta}_n^n$, 20
$S_{ff}(z), 255$
$\check{S}_{fg}(z)$, 255
$(((V_i), \subseteq), 59$
$(\mathbf{A}, \ \cdot\ , *), 151$
$\left(\boldsymbol{L}_{\mathbb{R}}^{2},\left(\!\left(\boldsymbol{V}_{j}\right)\!\right),54$
$(\kappa_n)_{n\in\mathbb{Z}}, 191$

$\ \cdot\ $, 116 $L^2_{(\mathbb{R},\mathcal{B},\mu)}$, 141
$L^2_{\mathbb{R}}$, 141
$PW_{\sigma}^{2}, 50$
$\frac{d}{dx}$, 222
$\exp(ix)$, 160
tan, 165
★ , 207
$\mathscr{L}(\mathbb{C},\mathbb{C})$, 50
cos, 165
$\cos(x)$, 155
sin, 165
$\sin(x)$, 155
$\tilde{S}_{ff}(\omega)$, 255
$\tilde{S}_{fg}(\omega)$, 255
F , 196
∨, <mark>70</mark>

\wedge , 70
w_N , 185
<i>X</i> , 39
<i>Y</i> , 39
ℂ ^ℂ , 39
$\mathbb{R}^{\mathbb{R}}$, 39
$D^*, 42$
\mathbf{D}_{α} , 40
I _m , 149
I, 112
L, 223
$R_{\rm e}$, 149
$T^*, 42$
T, 40
T_{τ} , 40
Z , 208
ϕ , 54
span, 9
' '

Y^{X} , 39
[x], 70
$\lfloor x \rfloor$, 70
T_n , 178
*, 148
$\hat{\mathbf{F}}^{-1}$, 248
$\hat{\mathbf{F}}^*$, 249
$\hat{\mathbf{F}}$, 247
 · , 117
★ , 145
$\mathcal{B}(\boldsymbol{X},\boldsymbol{Y}),\boldsymbol{120}$
Y^X , 113
ρ , 147, 266
$\sigma_{\rm c}$, 266
$\sigma_{\rm p}$, 266
$\sigma_{\rm r}^{\rm r}$, 266
σ , 147, 266
r, 147
•

page xiv	Daniel J. Greenhoe	




_CONTENTS

	Title page	
	Typesetting	
	Quotes	vii
	Symbol list	ix
	Symbol index	xiii
	Contents	XV
	Analysis and Transforms	_
1	Analyses and Transforms	1 1
	1.1 Abstract spaces	
	·	
	1.6 Operator inducing analyses	
	1.7 Wavelet analyses	6
2	Linear Combinations	9
	2.1 Linear combinations in linear spaces	9
	2.2 Bases in topological linear spaces	
	2.3 Schauder bases in Banach spaces	
	2.4 Linear combinations in inner product spaces	
	2.5 Orthonormal bases in Hilbert spaces	
	2.6 Riesz bases in Hilbert spaces	27
	2.7 Frames in Hilbert spaces	
_		
3	Transversal Operators	39
	3.1 Families of Functions	
	3.2 Definitions and algebraic properties	
	3.3 Linear space properties	
	3.4 Inner product space properties	
	Normed linear space properties	
	3.6 Fourier transform properties	
	Basis theory properties	50
4	MRA Structures	53
	4.1 Introduction	
	4.2 Definition	
	4.3 Order structure	
	4.4 Dilation equation	
	4.5 Necessary Conditions	
	4.6 Sufficient conditions	
	4.7 Support size	
	4.8 Scaling functions with partition of unity	
5	Wavelet Structures	79
	5.1 Introduction	79

oage xvi	Daniel J. Greenhoe	CONTENTS

	5.2	5.1.2 Analyses	79 80 81 81
	5.3 5.4 5.5	Order structure	83 83
	5.6 5.7	Necessary conditions	85 88
	5.8 5.9	Support size	89 90
Αŗ	pend	lices	93
A	Alge	ebraic structures	95
В	Line		97
	B.1	Subspaces of a linear space	
		Subspaces of an inner product space	
		Subspace Metrics	
		Literature	
С		· · · · · · · · · · · · · · · · · · ·	11
	C.1	Operators on linear spaces	
		C.1.1 Operator Algebra 1 C.1.2 Linear operators 1	
	C.2	Operators on Normed linear spaces	
		C.2.1 Operator norm	
		C.2.2 Bounded linear operators	
		C.2.3 Adjoints on normed linear spaces	
	C_{2}	C.2.4 More properties	
	0.3	C.3.1 General Results	
		C.3.2 Operator adjoint	
	C.4	Special Classes of Operators	
		C.4.1 Projection operators	
		C.4.2 Self Adjoint Operators	
		C.4.3 Normal Operators 1 C.4.4 Isometric operators 1	
		C.4.5 Unitary operators	
	C.5	Operator order	
D	Calc	eulus 1	41
E	Con	volution 1	45
		Definition	45
	E.2	Properties	45
F	Nor	med Algebras	47
	F.1	Algebras	
	F.2	Star-Algebras	
	F.3	Normed Algebras	
	F.4	C* Algebras	51
G			53
		Definition Candidates 1 Definitions 1	
		Basic properties	
		The complex exponential	
	G.5	Trigonometric Identities	62
		Planar Geometry	
	G.7	Trigonometric functions of complex numbers	68





CONTENTS Daniel J. Greenhoe page xvii

	G.8	The power of the exponential	170
н	Tria	onometric Polynomials	173
		Trigonometric reduction	
	H.3		
	H.4		
	H.5	Trigonometric summations	
	H.6		
L	Fou	rier Transform	195
	1.1	Introduction	
	1.2	Definitions	196
	1.3	Operator properties	197
	1.4	Transversal properties	198
	1.5	Convolution relations	198
	1.6	Calculus relations	199
	1.7	Real valued functions	200
	1.8	Moment properties	200
	1.9	Examples	203
J		ransform Communication of the	207
	J.1	Convolution operator	
	J.2	Z-transform	
	J.3	From z-domain back to time-domain	
	J.4	Zero locations	
	J.5	Pole locations	
	J.6	Mirroring for real coefficients	
	J.7	Rational polynomial operators	
	J.8	Filter Banks	
	J.9	Inverting non-minimum phase filters	219
	T	lau Funanciana (Tuancfausa)	001
K		lor Expansions (Transforms)	221
	K.1		
	K.2	Taylor Expansion	222
ı.	Lan	lace Transform	223
-	Lup I 1	Operator Definition	
	L.2	Operator Inverse	
	L.3	Transversal properties	
	L.4	Linear properties	
			7.7.3
	15		
	L.5	Modulation properties	226
	L.6	Modulation properties	226 228
	L.6 L.7	Modulation properties	226 228 233
	L.6 L.7 L.8	Modulation properties	226 228 233 235
	L.6 L.7	Modulation properties	226 228 233 235
M	L.6 L.7 L.8 L.9	Modulation properties	226 228 233 235
M	L.6 L.7 L.8 L.9	Modulation properties	226 228 233 235 236 237
M	L.6 L.7 L.8 L.9	Modulation properties Causality properties Exponential decay properties Product properties Calculus properties Crete Time Fourier Transform Definition	226 228 233 235 236 237 237
M	L.6 L.7 L.8 L.9 Disc M.1 M.2	Modulation properties	226 228 233 235 236 237 237 237
M	L.6 L.7 L.8 L.9 Disc M.1 M.2	Modulation properties Causality properties Exponential decay properties Product properties Calculus properties Calculus properties Definition Properties	226 228 233 235 236 237 237 237
	L.6 L.7 L.8 L.9 Disc M.1 M.2 M.3	Modulation properties Causality properties Exponential decay properties Product properties Calculus properties Crete Time Fourier Transform Definition Properties Derivatives	226 228 233 235 236 237 237 245 247
	L.6 L.7 L.8 L.9 Disc M.1 M.2 M.3	Modulation properties Causality properties Exponential decay properties Product properties Calculus properties Crete Time Fourier Transform Definition Properties Derivatives	226 228 233 235 236 237 237 245 247
	L.6 L.7 L.8 L.9 Disc M.1 M.2 M.3	Modulation properties Causality properties Exponential decay properties Product properties Calculus properties Crete Time Fourier Transform Definition Properties Derivatives	226 228 233 235 236 237 237 245 247
	L.6 L.7 L.8 L.9 Disc M.1 M.2 M.3	Modulation properties Causality properties Exponential decay properties Product properties Calculus properties Calculus properties Crete Time Fourier Transform Definition Properties Derivatives prier Series Definition	226 228 233 235 236 237 237 245 247 247 248
N	L.6 L.7 L.8 L.9 Disc M.1 M.2 M.3 Fou N.1 N.2 N.3	Modulation properties Causality properties Exponential decay properties Product properties Calculus properties Calculus properties Crete Time Fourier Transform Definition Properties Derivatives rier Series Definition Inverse Fourier Series operator Fourier series for compactly supported functions	226 228 233 235 236 237 237 245 247 247 248 250
N	L.6 L.7 L.8 L.9 Disc M.1 M.2 M.3 Fou N.1 N.2 N.3	Modulation properties Causality properties Exponential decay properties Product properties Calculus properties Calculus properties Crete Time Fourier Transform Definition Properties Derivatives rier Series Definition Inverse Fourier Series operator	226 228 233 235 236 237 237 245 247 247 248
N O	L.6 L.7 L.8 L.9 Disc M.1 M.2 M.3 Fou N.1 N.2 N.3	Modulation properties Causality properties Exponential decay properties Product properties Calculus properties Calculus properties Crete Time Fourier Transform Definition Properties Derivatives rier Series Definition Inverse Fourier Series operator Fourier series for compactly supported functions	226 228 233 235 236 237 237 245 247 247 248 250
N O	L.6 L.7 L.8 L.9 Disc M.1 M.2 M.3 Fou N.1 N.2 N.3	Modulation properties Causality properties Exponential decay properties Product properties Calculus properties Calculus properties Crete Time Fourier Transform Definition Properties Derivatives rier Series Definition Inverse Fourier Series operator Fourier series for compactly supported functions t Wavelet Transform (FWT)	226 228 233 235 236 237 237 245 247 248 250 251



page xviii Daniel J. Greenhoe CONTENTS

Q	Continuous Random Processes				
	Q.1 Definitions	263			
	Q.2 Properties	264			
R	Spectral Theory	265			
	Spectral Theory R.1 Operator Spectrum	265			
	R.2 Fredholm kernels	267			
Ва	ack Matter	269			
	ack Matter References	270			
	Reference Index	291			
	Subject Index	295			
	License	307			
	End of document				

CHAPTER 1			



The analytical equations, unknown to the ancient geometers, which Descartes was the first to introduce into the study of curves and surfaces, ... they extend to all general phenomena. There cannot be a language more universal and more simple, more free from errors and from obscurities, ... mathematical analysis is as extensive as nature itself; it defines all perceptible relations, measures times, spaces, forces, temperatures; this difficult science is formed slowly, but it preserves every principle which it has once acquired; it grows and strengthens itself incessantly in the midst of the many variations and errors of the human mind. Its chief attribute is clearness; it has no marks to express confused notions. It brings together phenomena the most diverse, and discovers the hidden analogies which unite them.

ANALYSES AND TRANSFORMS

Joseph Fourier (1768–1830) ¹

1.1 Abstract spaces

The **abstract space** was introduced by Maurice Fréchet in his 1906 Ph.D. thesis.² An *abstract space* in mathematics does not really have a rigorous definition; but in general it is a set together with some other unifying structure. Examples of spaces include *topological spaces*, *metric spaces*, and *linear spaces* (*vector spaces*).

² Fréchet (1906),

Fréchet (1928). "A collection of these abstract elements will be called an abstract set. If to this set there is added some rule of association of these elements, or some relation between them, the set will be called an abstract space."—Maurice Fréchet

1.2 Lattice of subspaces

An abstract space can be decomposed into one or more *subspaces*. Roughly speaking, a subspace of an abstract space is simply a subset the abstract space that has the same properties of that abstract space. The subspaces can be ordered under the ordering relation \subseteq (subset or equal to relation) to form a *lattice*.





Figure 1.1: lattice of topologies on $X \triangleq \{x, y, z\}$ (Example 1.1 page 2)

Example 1.1. ³The power set 2^X is a *topology* on the set X. But there are also 28 other topologies on $\{x, y, z\}$, and these are all *subspaces* of $2^{\{x, y, z\}}$. Let a given topology in $\mathcal{T}(\{x, y, z\})$ be represented by a Hasse diagram as illustrated to the right, where a circle present means the indicated set is in the topology, and a circle absent means the indicated set is not in the topology. The lattice of the 29 topologies $(\mathcal{T}(\{x, y, z\}), \cup, \cap; \subseteq)$ is illustrated in Figure 1.1 (page 2). The lattice of these 29 topologies is *non-distributive* (it contains the *N5 lattice*). The five topologies illustrated by red shaded nodes are also *algebras of sets*.



Example 1.2. The power set 2^X is an *algebra of sets* on the set X. But there are also 14 other algebras of sets on $\{w, x, y, z\}$, and these are all *subspaces* of $2^{\{w, x, y, z\}}$. The *lattice of algebras of sets* on $\{w, x, y, z\}$ is illustrated in Figure 1.2 (page 3).

A *linear subspace* is a subspace of a *linear space* (*vector space*). Linear subspaces have some special properties: Every linear subspace contains the additive identity zero vector, and every linear subspace is *convex*.

⁴ Isham (1999) page 44, Isham (1989) page 1516, I Steiner (1966) page 386



1.3. ANALYSES Daniel J. Greenhoe page 3





Figure 1.2: lattice of *algebras of sets* on $\{w, x, y, z\}$ (Example 1.2 page 2)

Example 1.3. The 3-dimensional Euclidean space \mathbb{R}^3 contains the 2-dimensional xy-plane and xz-plane subspaces, which in turn both contain the 1-dimensional x-axis subspace. These subspaces are illustrated in the figure to the right and in Figure B.1 (page 97).



1.3 Analyses

An **analysis** of a space X is any lattice of subspaces of X. The partial or complete reconstruction of X from this set is a **synthesis**.

Example 1.4. The lattices of subspaces illustrated in Figure 1.4 (page 4) are all *analyses* of \mathbb{R}^3 .

1.4 Transform

Definition 1.1. A transform on a space **X** is a sequence of projection operators that induces an ANAL-YSIS on **X**.

Section 1.3 defined an **analysis** of a space X as is any lattice of subspaces of X. In like manner, an **analysis** of a function f(x) with respect to a transform T is simply the transform T of f (Tf). Such

⁵The word *analysis* comes from the Greek word ἀνάλυσις, meaning "dissolution" (Perschbacher (1990) page 23 (entry 359)), which in turn means "the resolution or separation into component parts" (Black et al. (2009), http://dictionary.reference.com/browse/dissolution)








Figure 1.3: lattice of subspaces of \mathbb{R}^3 (Example B.1 page 97)

linearly ordered analysis of \mathbb{R}^3 M-3 analysis of \mathbb{R}^3 wavelet-like analysis of \mathbb{R}^3

Figure 1.4: some analyses of \mathbb{R}^3 (Example 1.4 page 3)

an analysis or transform is often represented as the sequence of coefficients (λ_n) multiplying the

basis vectors
$$(\psi_n(x))$$
 such that $f(x) = \mathbf{T}f(x) = \sum_{n \in \mathbb{Z}} \lambda_n \psi_n(x)$

Example 1.5. A Fourier analysis is a sequence of subspaces with sinusoidal bases. Examples of subspaces in a Fourier analysis include $V_1 = \text{span}\{e^{ix}\}$, $V_{2.3} = \text{span}\{e^{i2.3x}\}$, $V_{\sqrt{2}} = \text{span}\{e^{i\sqrt{2}x}\}$, etc. A **transform** is a set of *projection operators* that maps a family of functions (e.g. $L^2_{\mathbb{R}}$) into an analysis. The Fourier transform" for Fourier Analysis is (Definition I.2 page 196)

$$[\tilde{\mathbf{F}}\mathbf{f}](\omega) \triangleq \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} \mathbf{f}(x)e^{-i\omega x} dx$$

Properties of subspace order structures 1.5

The ordered set of all linear subspaces of a *Hilbert space* is an *orthomodular lattice*. Orthomodular lattices (and hence Hilbert subspaces) have some special properties (next theorem). One is that they satisfy de Morgan's law.

(DE MORGAN) and (DE MORGAN) and and and

⁶ ∄ Beran (1985) pages 30–33, ∄ Birkhoff and Neumann (1936) page 830 ⟨L74⟩, ∄ Beran (1976) pages 251–252






Figure 1.5: some common transforms

Most transforms have a very simple M-n order structure, as illustrated to the right and in Figure 1.5 page 5. The M-n lattices for $n \ge 3$ are *modular* but not *distributive*. Analyses typically have one subspace that is a *scaling* subspace; and this subspace is often simply a family of constants (as is the case with *Fourier Analysis*). There is one noteable exception to this—MRA induced *wavelet analysis* (Definition 5.1 page 81).




1.6 Operator inducing analyses



An *analysis* is often defined in terms of

a small number (e.g. 2) operators. Two such operators are the *translation operator* and the *dilation operator* (Definition 3.3 page 40).



Example 1.6. In Fourier analysis, continuous dilations (Definition 3.3 page 40) of the complex exponential form a basis (Definition 2.7 page 14) for the space of square integrable functions $L^2_{\mathbb{R}}$ (Definition D.1 page 141) such that $L^2_{\mathbb{R}} = \operatorname{span} \left\{ \mathbf{D}_{\omega} e^{ix} |_{\omega \in \mathbb{R}} \right\}$.

Example 1.7. In Fourier series analysis (Theorem N.1 page 248), discrete dilations of the complex exponential form a basis for $\mathcal{L}^2_{\mathbb{R}}(0:2\pi)$ such that $\mathcal{L}^2_{\mathbb{R}}(0:2\pi)=\operatorname{span}\left\{\mathbf{D}_je^{ix}\big|j\in\mathbb{Z}\right\}$.





Figure 1.6: some wavelet transforms

1.7 Wavelet analyses

The term "wavelet" comes from the French word "ondelette", meaning "small wave". And in essence, wavelets are "small waves" (as opposed to the "long waves" of Fourier analysis) that form a basis for the Hilbert space $\boldsymbol{L}_{\mathbb{R}}^2$.

A special characteristic of wavelet analysis is that there is not just one scaling subspace, (as is with the case of Fourier and several other analyses), but an entire sequence of scaling subspaces (Figure 1.6 page 6). These scaling subspaces are *linearly ordered* with respect to the ordering relation ⊆. In wavelet theory, this structure is called a *multireso-lution analysis*, or *MRA* (Definition 4.1 page 54). The MRA was introduced by Stéphane G. Mallat in 1989. The concept of a scaling space was perhaps first introduced by Taizo Iijima in 1959 in Japan, and later as the *Gaussian Pyramid* by Burt and Adelson in the 1980s in the West. 9



The MRA has become the dominate wavelet construction method. Moreover, P.G. Lemarié has proved that all wavelets with *compact support* are generated by an MRA.¹¹


¹¹ Lemarié (1990), Mallat (1999) page 240




¹⁰ Mallat (1989) page 70, ☐ Iijima (1959), ☐ Burt and Adelson (1983), ☐ Adelson and Burt (1981), ☐ Lindeberg (1993), ☐ Alvarez et al. (1993), ☐ Guichard et al. (2012) pages 23–24 (§3.2.1 Scale-Space Extrema), ☐ Guichard et al. (20xx) pages 77–78 (§5.2.1 Scale-Space Extrema), ☐ Weickert (1999) (historical survey)

1.7. WAVELET ANALYSES Daniel J. Greenhoe page 7

A second special characteristic of wavelet analysis is that it's order structure with respect to the \subseteq relation is not a simple M-n lattice (as is with the case of Fourier and several other analyses). Rather, it is a lattice of the form illustrated to the right and in Figure 1.6 (page 6). This lattice is *non-complemented*, *non-distributive*, *non-modular*, and *non-Boolean* (Proposition 5.1 page 83). 12



In the world of mathematical structures, the order structure of wavelet analyses is quite rare, but not completely unique. One example of a system with similar structure is the set of $Primorial^{14}$ numbers together with the | ("divides") ordering relation 13 as illustrated to the right.



The basis sequence of most transform are fixed with no design freedom For example, the Fourier Transform uses the complex exponential, Taylor Expansion uses monomials of the form $(x - a)^n$. However, there are an infinite number of wavelet basis sequences—lots and lots of design freedom. For information regarding designing wavelet basis sequences, see α Greenhoe (2013).

However, one arguable disadvantage is that wavelets do not support a **convolution theorem**—a theorem enjoyed by the Fourier transforms, Laplace Transform, and Z Transform. These other transforms induce a convolution theorem because they are defined in terms of an exponential (e.g. $e^{-i\omega t}$, $e^{-i\omega n}$, e^{-st} , z^{-n}), and exponentials sport the property $a^{x+y} = a^x a^y$.

¹⁴

Sloane (2014) (http://oeis.org/A002110),
☐ Greenhoe (2013) page 30





Linear combinations in linear spaces 2.1

A *linear space* (Definition C.1 page 111) in general is not equipped with a *topology*. Without a topology, it is not possible to determine whether an *infinite sum* of vectors converges. Therefore in this section (dealing with linear spaces), all definitions related to sums of vectors will be valid for finite sums only (finite "N").

Definition 2.1. Let $\{x_n \in X | n=1,2,...,N\}$ be a set of vectors in a Linear space $(X, +, \cdot, (\mathbb{F}, \dotplus, \dot{\times}))$.

A vector $x \in X$ is a **linear combination** of the vectors in $\{x_n\}$ if

D E

there exists $\{\alpha_n \in \mathbb{F} | n=1,2,...,N\}$ such that $x = \sum_{n=1}^{\infty} \alpha_n x_n$.

Definition 2.2. Let $(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}))$ be a linear space and Y be a subset of X.

D E F

The **linear span** of Y is defined as $\operatorname{span} Y \triangleq \left\{ \sum_{\gamma \in \Gamma} \alpha_{\gamma} \mathbf{y}_{\gamma} \middle| \alpha_{\gamma} \in \mathbb{F}, \mathbf{y}_{\gamma} \in Y \right\}.$

The set Y **spans** a set A if $A \subseteq \operatorname{span} Y$.

Proposition 2.1. ³ Let $\{x_n \in X \mid n=1,2,...,N\}$ be a set of vectors in a Linear space $\mathbf{L} \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}))$.

```
1. span\{x_n\} is a LINEAR SPACE (Definition C.1 page 111) and 2. span\{x_n\} is a LINEAR SUBSPACE of {\bf L} .
```

Definition 2.3. 4 Let $\mathbf{L} \triangleq \left(X, +, \cdot, (\mathbb{F}, \dotplus, \dot{\times})\right)$ be a Linear space.

The set $Y \triangleq \left\{x_n \in X \middle| n=1,2,...,N\right\}$ is **linearly independent** in \mathbf{L} if $\left\{\sum_{n=1}^N \alpha_n x_n = 0\right\} \implies \left\{\alpha_1 = \alpha_2 = \cdots = \alpha_N = 0\right\}.$ The set Y is **linearly dependent** in L if Y is not linearly independent in L.

¹ ■ Berberian (1961) page 11 (Definition I.4.1), ■ Kubrusly (2001) page 46

⁽²⁰⁰²⁾ page 71 (Definition 3.2.5—more general definition)

³ Kubrusly (2001) page 46

⁴ Bachman and Narici (1966) pages 3–4,
☐ Christensen (2003) page 2, ☐ Heil (2011) page 156 (Definition 5.7)




Figure 2.1: Lattice of linear combinations

Definition 2.4. ⁵ Let $\{x_n \in X \mid n=1,2,...,N\}$ be a set of vectors in a LINEAR SPACE $\mathbf{L} \triangleq (X, +, \cdot, (\mathbb{F}, +, \dot{\mathbf{x}}))$.

D E F

```
The set \{x_n\} is a Hamel basis for L if

1. \{x_n\} SPANS L
```

1. $\{x_n\}$ SPANS L

2. $\{x_n\}$ is LINEARLY INDEPENDENT in L (Definition 2.1 page 9)

A HAMEL BASIS is also called a **linear basis**.

Definition 2.5. ⁶ Let $L \triangleq (X, +, \cdot, (\mathbb{F}, +, \dot{x}))$ be a linear space. Let x be a vector in L and $Y \triangleq$ $\{x_n \in X \mid n=1,2,...,N\}$ be a set of vectors in L.



The expression $\sum_{n=1}^{N} \alpha_n \mathbf{x}_n$ is the **expansion** of \mathbf{x} on Y in \mathbf{L} if $\mathbf{x} = \sum_{n=1}^{N} \alpha_n \mathbf{x}_n$.

In this case, the sequence $(\alpha_n)_{n=1}^N$ is the **coordinates** of x with respect to Y in L. If $\alpha_N \neq 0$, then N is the **dimension** dim**L** of **L**.

Theorem 2.1. ⁷ Let $\{x_n | n=1,2,...,N\}$ be a Hamel basis (Definition 2.4 page 11) for a Linear space

[♠]Proof:

$$\begin{split} & = \mathbf{x} - \mathbf{x} \\ & = \sum_{n=1}^{N} \alpha_n \mathbf{x}_n - \sum_{n=1}^{N} \beta_n \mathbf{x}_n \\ & = \sum_{n=1}^{N} \left(\alpha_n - \beta_n \right) \mathbf{x}_n \\ & \Longrightarrow \left\{ \mathbf{x}_n \right\} \text{ is } \textit{linearly dependent if } \left(\alpha_n - \beta_n \right) \neq 0 \qquad \forall n = 1, 2, \dots, N \\ & \Longrightarrow \left(\alpha_n - \beta_n \right) = 0 \qquad \forall n = 1, 2, \dots, N \qquad \text{(because } \left\{ \mathbf{x}_n \right\} \text{ is a } \textit{basis } \text{and therefore must be } \textit{linearly independent)} \\ & \Longrightarrow \alpha_n = \beta_n \text{ for } n = 1, 2, \dots, N \end{split}$$

```
Theorem 2.2. <sup>8</sup> Let L \triangleq (X, +, \cdot, (\mathbb{F}, +, \times)) be a Linear space.

1. \{x_n \in X | n=1,2,...,N\} is a Hamel basis for L
2. \{y_n \in X | n=1,2,...,M\} is a set of Linearly independent vectors in L
    T
H
M

\begin{cases}
1. & M \leq N \\
2. & M = N \implies \{y_n | n=1,2,...,M\} \text{ is a BASIS for } \mathbf{L} \\
3. & M \neq N \implies \{y_n | n=1,2,...,M\} \text{ is NOT a basis for } \mathbf{L}
\end{cases}
```

[♠]Proof:

1. Proof that $\{y_1, x_1, \dots, x_{N-1}\}$ is a *basis* for *L*:



⁵ Hamel (1905), ∂ Bachman and Narici (1966) page 4, ∂ Kubrusly (2001) pages 48–49 (Section 2.4), ∂ Young (2001) page 1, ☐ Carothers (2005) page 25, ☐ Heil (2011) page 125 ⟨Definition 4.1⟩

⁶🗒 Hamel (1905), *᠗* Bachman and Narici (1966) page 4, *᠗* Kubrusly (2001) pages 48–49 (Section 2.4), *᠗* Young

⁷ Michel and Herget (1993) pages 89–90 (Theorem 3.3.25)

⁸ Michel and Herget (1993) pages 90–91 (Theorem 3.3.26)

- (a) Proof that $\{y_1, x_1, ..., x_{N-1}\}$ spans L:
 - i. Because $\{x_n \mid n=1,2,...,N\}$ is a *basis* for L, there exists $\beta \in \mathbb{F}$ and $\{\alpha_n \in \mathbb{F} \mid n=1,2,...,N\}$ such that $\beta y_1 + \sum_{i=1}^{N} \alpha_n x_n = 0$.
 - ii. Select an n such that $\alpha_n \neq 0$ and renumber (if necessary) the above indices such that

$$x_n = -\frac{\beta}{\alpha_n} y_1 + \sum_{n=1}^{N-1} \frac{\alpha_n}{\alpha_n} x_n.$$

iii. Then, for any $y \in X$, we can write

$$y = \sum_{n=1}^{N} \gamma_{n \in \mathbb{Z}} x_n$$

$$= \left(\sum_{n=1}^{N-1} \gamma_{n \in \mathbb{Z}} x_n\right) + \gamma_{n \in \mathbb{Z}} \left(-\frac{\beta}{\alpha_n} y_1 - \sum_{n=1}^{N-1} \frac{\alpha_n}{\alpha_n} x_n\right)$$

$$= -\frac{\beta \gamma_n}{\alpha_n} y_1 + \sum_{n=1}^{N-1} \left(\gamma_n - \frac{\alpha_n \gamma_n}{\alpha_n}\right) x_n$$

$$= \delta y_1 + \sum_{n=1}^{N-1} \delta_{n \in \mathbb{Z}} x_n$$

- iv. This implies that $\{y_1, x_1, ..., x_{N-1}\}$ spans L:
- (b) Proof that $\{y_1, x_1, ..., x_{N-1}\}$ is linearly independent:
 - i. If $\{y_1, x_1, \dots, x_{N-1}\}$ is *linearly dependent*, then there exists $\{\epsilon, \epsilon_1, \dots, \epsilon_{N-1}\}$ such that $\epsilon y_1 + \left(\sum_{n=1}^{N-1} \epsilon_{n \in \mathbb{Z}} x_n\right) + 0x_n = 0.$
 - ii. item (1(b)i) implies that the coordinate of y_1 associated with x_n is 0.

$$\mathbf{y}_1 = -\left(\sum_{n=1}^{N-1} \frac{\epsilon_n}{\epsilon} \mathbf{x}_n\right) + 0 \mathbf{x}_n = 0.$$

iii. item (1(a)i) implies that the coordinate of y_1 associated with x_n is not 0.

$$\mathbf{y}_1 = -\sum_{n=1}^N \frac{\alpha_n}{\beta} \mathbf{x}_n.$$

- iv. This implies that item (1(b)i) (that the set is linearly dependent) is *false* because item (1(b)ii) and item (1(b)iii) *contradict* each other.
- v. This implies $\{y_1, x_1, ..., x_{N-1}\}$ is linearly independent.
- 2. Proof that $\{y_1, y_2, x_1, ..., x_{N-2}\}$ is a *basis*: Repeat item (1).
- 3. Suppose m = n. Proof that $\{y_1, y_2, ..., y_M\}$ is a *basis*: Repeat item (1) M 1 times.
- 4. Proof that M > N:
 - (a) Suppose that M = N + 1.
 - (b) Then because $\{ \mathbf{y}_n | n=1,2,...,N \}$ is a *basis*, there exists $\{ \zeta_n | n=1,2,...,N+1 \}$ such that $\sum_{n=1}^{N+1} \zeta_{n \in \mathbb{Z}} \mathbf{y}_{n \in \mathbb{Z}} = 0.$
 - (c) This implies that $\{y_n|_{n=1,2,...,N+1}\}$ is *linearly dependent*.
 - (d) This implies that $\{y_n|_{n=1,2,...,N+1}\}$ is *not* a basis.



- (e) This implies that M > N.
- 5. Proof that $M \neq N \implies \{y_n|_{n=1,2,...,M}\}$ is *not* a basis for L:
 - (a) Proof that $M > N \implies \{y_n | n=1,2,...,M\}$ is *not* a basis for L: same as in item (4).
 - (b) Proof that $M < N \implies \{y_n|_{n=1,2,...,M}\}$ is *not* a basis for L:
 - i. Suppose m = N 1.
 - ii. Then $\{y_n|_{n=1,2,...,N-1}\}$ is a *basis* and there exists λ such that

$$\sum_{n=1}^{N} \lambda_{n \in \mathbb{Z}} \mathbf{y}_{n \in \mathbb{Z}} = 0.$$

- iii. This implies that $\{y_n | n=1,2,...,N\}$ is *linearly dependent* and is *not* a basis.
- iv. But this contradicts item (3), therefore $M \neq N 1$.
- v. Because M = N yields a basis but M = N 1 does not, M < N 1 also does not yield a basis.

Corollary 2.1. ⁹ Let $L \triangleq (X, +, \cdot, (\mathbb{F}, +, \dot{\times}))$ be a linear space.

 $\left\{ x_n \in X \mid n=1,2,...,N \right\}$ is a Hamel basis for L and $\left\{ y_n \in X \mid n=1,2,...,M \right\}$ is a Hamel basis for L $\{N = M\}$ (all Hamel bases for **L** have the same number of vectors)

 $^{\circ}$ Proof: This follows from Theorem 2.2 (page 11).

Bases in topological linear spaces 2.2

A linear space supports the concept of the *span* of a set of vectors (Definition 2.2 page 9). In a topological linear space $\Omega \triangleq (X, +, \cdot, (\mathbb{F}, +, \times), T)$, a set A is said to be total in Ω if the span of A is dense in Ω . In this case, A is said to be a *total set* or a *complete set*. However, this use of "complete" in a "complete set" is not equivalent to the use of "complete" in a "complete metric space". 10 In this text, except for these comments and Definition 2.6, "complete" refers to the metric space definition only.

If a set is both total and linearly independent (Definition 2.3 page 9) in Ω , then that set is a Hamel basis (Definition 2.4 page 11) for Ω .

Definition 2.6. ¹¹ Let A^- be the closure of a A in a topological linear space $\mathbf{\Omega} \triangleq (X, +, \cdot, (\mathbb{F}, +, \dot{\times}), T)$ Let span A be the SPAN (Definition 2.2 page 9) of a set A.

A set of vectors A is **total** (or **complete** or **fundamental**) in Ω if $(\operatorname{span} A)^- = \Omega$ (SPAN of A is dense in Ω).

¹¹ ♥ Young (2001) page 19 〈Definition 1.5.1〉, ● Sohrab (2003) page 362 〈Definition 9.2.3〉, ● Gupta (1998) page 134 (Definition 2.4), Bachman and Narici (1966) pages 149–153 (Definition 9.3, Theorems 9.9 and 9.10)



⁹ Kubrusly (2001) page 52 (Theorem 2.7), \mathcal{A} Michel and Herget (1993) page 91 (Theorem 3.3.31)

¹⁰ Haaser and Sullivan (1991) pages 296–297 (6·Orthogonal Bases), ₽ Rynne and Youngson (2008) page 78 〈Remark 3.50〉, **/** Heil (2011) page 21 〈Remark 1.26〉

DEF

2.3 Schauder bases in Banach spaces

Definition 2.7. Let $\mathbf{B} \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \|\cdot\|)$ be a Banach space. Let $\stackrel{\star}{=}$ represent strong convergence in \mathbf{B} .

CONVERGENCE in **B**.

The countable set $\{x_n \in X \mid n \in \mathbb{N}\}$ is a **Schauder basis** for **B** if for each $x \in X$

1.
$$\exists (\alpha_n \in \mathbb{F})_{n \in \mathbb{N}}$$
 such that $x \stackrel{\star}{=} \sum_{n=1}^{\infty} \alpha_n x_n$ (Strong convergence in B) and

2.
$$\left\{\sum_{n=1}^{\infty}\alpha_{n}x_{n} \stackrel{\star}{=} \sum_{n=1}^{\infty}\beta_{n}x_{n}\right\} \implies \left\{\left(\left(\alpha_{n}\right)\right) = \left(\left(\beta_{n}\right)\right)\right\} \quad \text{(coefficient functionals are unique)}$$

In this case, $\sum_{n=1}^{\infty} \alpha_n \mathbf{x}_n$ is the **expansion** of \mathbf{x} on $\{\mathbf{x}_n | n \in \mathbb{N}\}$ and

the elements of (α_n) are the **coefficient functionals** associated with the basis $\{x_n\}$. Coefficient functionals are also called **coordinate functionals**.

In a Banach space, the existence of a Schauder basis implies that the space is *separable* (Theorem 2.3 page 14). The question of whether the converse is also true was posed by Banach himself in 1932, ¹³ and became know as "*The basis problem*". This remained an open question for many years. The question was finally answered some 41 years later in 1973 by Per Enflo (University of California at Berkley), with the answer being "no". Enflo constructed a counterexample in which a separable Banach space does *not* have a Schauder basis. ¹⁴ Life is simpler in Hilbert spaces where the converse *is* true: a Hilbert space has a Schauder basis *if and only if* it is separable (Theorem 2.11 page 27).

Theorem 2.3. ¹⁵ Let $\mathbf{B} \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \|\cdot\|)$ be a Banach space. Let \mathbb{Q} be the field of rational numbers.

$$\begin{bmatrix}
\mathsf{T} \\
\mathsf{H} \\
\mathsf{M}
\end{bmatrix}
\begin{bmatrix}
1. & \mathbf{B} \text{ has a Schauder basis} & and \\
2. & \mathbb{Q} \text{ is dense } in \mathbb{F}.
\end{bmatrix}
\implies \left\{ \mathbf{B} \text{ is separable } \right\}$$

[♠]Proof:

1. lemma:

$$\left\| \left\{ x | \exists \left(\alpha_n \mathbb{Q} \right)_{n \in \mathbb{N}} \text{ such that } \lim_{N \to \infty} \left\| x - \sum_{n=1}^{N} \alpha_n x_n \right\| = 0 \right\} \right\| = |\mathbb{Q} \times \mathbb{N}|$$

$$= |\mathbb{Z} \times \mathbb{Z}|$$

$$= |\mathbb{Z}|$$

$$= countably infinite$$



¹² ☐ Carothers (2005) pages 24–25, ☐ Christensen (2003) pages 46–49 〈Definition 3.1.1 and page 49〉, ☐ Young (2001) page 19 〈Section 6〉, ☐ Singer (1970) page 17, ☐ Schauder (1927), ☐ Schauder (1928)

¹³ 🛮 Banach (1932a) page 111

¹⁴ ■ Enflo (1973), ■ Lindenstrauss and Tzafriri (1977) pages 84–95 (Section 2.d)

2. remainder of proof:

B has a Schauder basis $(x_n)_{n\in\mathbb{N}}$

$$\implies$$
 for every $\mathbf{x} \in \mathbf{B}$, there exists $(\alpha_n \in \mathbb{F})_{n \in \mathbb{N}}$ such that $\mathbf{x} \stackrel{*}{=} \sum_{n=1}^{\infty} \alpha_n \mathbf{x}_n$ by Definition 2.7 page 14

$$\implies$$
 for every $\mathbf{x} \in \mathbf{B}$, there exists $(\alpha_n \in \mathbb{F})_{n \in \mathbb{N}}$ such that $\lim_{N \to \infty} \left\| \mathbf{x} - \sum_{n=1}^N \alpha_n \mathbf{x}_n \right\| = 0$

$$\implies$$
 for every $\mathbf{x} \in \mathbf{B}$, there exists $(\alpha_n \in \mathbb{Q})_{n \in \mathbb{N}}$ such that $\lim_{N \to \infty} \left\| \mathbf{x} - \sum_{n=1}^N \alpha_n \mathbf{x}_n \right\| = 0$ because $\mathbb{Q}^- = \mathbb{F}$

$$\implies \boldsymbol{B} = \left\{ \boldsymbol{x} | \exists \left(\alpha_n \mathbb{Q} \right)_{n \in \mathbb{N}} \text{ such that } \lim_{N \to \infty} \left\| \boldsymbol{x} - \sum_{n=1}^N \alpha_n \boldsymbol{x}_n \right\| = 0 \right\}$$

$$\implies \mathbf{\textit{B}} = \left\{ \left. x | \exists \left((\alpha_n \mathbb{Q}) \right)_{n \in \mathbb{N}} \text{ such that } \mathbf{\textit{x}} = \lim_{N \to \infty} \sum_{n=1}^N \alpha_n \mathbf{\textit{x}}_n \right. \right\}$$

 \implies **B** is separable by (1) lemma page 14

Definition 2.8. ¹⁶ Let $\{x_n | n \in \mathbb{N}\}$ and $\{y_n | n \in \mathbb{N}\}$ be Schauder bases of a Banach space $(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \|\cdot\|).$

 $\{x_n\}$ is **equivalent** to $\{y_n\}$ if there exists a BOUNDED INVERTIBLE operator **R** in X^X such that $\mathbf{R}x_n = y_n$

Theorem 2.4. ¹⁷ Let $\{x_n \mid n \in \mathbb{N}\}$ and $\{y_n \mid n \in \mathbb{N}\}$ be Schauder bases of a Banach space $(X, +, \cdot, (\mathbb{F}, \dotplus, \dot{\times}), ||\cdot||).$

$$\left\{ \left\{ \boldsymbol{x}_{n} \right\} \text{ is equivalent to } \left\{ \boldsymbol{y}_{n} \right\} \right\} \\ \iff \left\{ \sum_{n=1}^{\infty} \alpha_{n} \boldsymbol{x}_{n} \text{ is convergent } \iff \sum_{n=1}^{\infty} \alpha_{n} \boldsymbol{y}_{n} \text{ is convergent} \right\}$$

Lemma 2.1. 18 Let $\Omega \triangleq (X, +, \cdot, (\mathbb{F}, +, \dot{\times}), T)$ be a topological linear space. Let span A be the SPAN of a set A (Definition 2.2 page 9). Let $\tilde{\mathsf{f}}(\omega)$ and $\tilde{\mathsf{g}}(\omega)$ be the FOURIER TRANSFORMs (Definition 1.2 page 196) of the functions f(x) and g(x), respectively, in $L^2_{\mathbb{R}}$ (Definition D.1 page 141). Let $\check{a}(\omega)$ be the DTFT (Definition M.1 page 237) of a sequence $(a_n)_{n\in\mathbb{Z}}$ in $\mathscr{C}^2_{\mathbb{R}}$ (Definition J.2 page 207).

 $^{\otimes}$ Proof: Let V'_0 be the space spanned by $\{\mathbf{T}^n \phi | n \in \mathbb{Z}\}$.

$$\begin{split} \tilde{\mathbf{f}}(\omega) &\triangleq \tilde{\mathbf{F}}\mathbf{f} & \text{by definition of } \tilde{\mathbf{F}} \\ &= \tilde{\mathbf{F}} \sum_{n \in \mathbb{Z}} a_n \mathbf{T} \mathbf{g} \\ &= \sum_{n \in \mathbb{Z}} a_n \tilde{\mathbf{F}} \mathbf{T} \mathbf{g} \end{split}$$

 $^{^{16}}$ Young (2001) page 25 (Definition 1.8.1, Theorem 1.8.7)

¹⁷ Young (2001) page 25 (Definition 1.8.1, Theorem 1.8.7)

¹⁸ Daubechies (1992) page 140

$$=\underbrace{\sum_{n\in\mathbb{Z}}a_ne^{-i\omega n}\tilde{\mathbf{F}}\mathbf{g}}_{\check{\mathbf{g}}(\omega)}$$

(Corollary 3.1 page 47)

 $= \breve{a}(\omega)\tilde{g}(\omega)$

by definition of $\check{\mathbf{F}}$ and $\tilde{\mathbf{F}}$

(Definition M.1 page 237) (Definition I.2 page 196)

$$\begin{split} & \boldsymbol{V}_0 \triangleq \left\{ \mathbf{f}(x) | \mathbf{f}(x) = \sum_{n \in \mathbb{Z}} b_n \mathbf{T}^n \mathbf{g}(x) \right\} \\ & = \left\{ \mathbf{f}(x) | \tilde{\mathbf{F}} \mathbf{f}(x) = \tilde{\mathbf{F}} \sum_{n \in \mathbb{Z}} b_n \mathbf{T}^n \mathbf{g}(x) \right\} \\ & = \left\{ \mathbf{f}(x) | \tilde{\mathbf{f}}(\omega) = \tilde{\mathbf{b}}(\omega) \tilde{\mathbf{g}}(\omega) \right\} \\ & = \left\{ \mathbf{f}(x) | \tilde{\mathbf{f}}(\omega) = \tilde{\mathbf{b}}(\omega) \tilde{\mathbf{a}}(\omega) \tilde{\mathbf{f}}(\omega) \right\} \\ & = \left\{ \mathbf{f}(x) | \tilde{\mathbf{f}}(\omega) = \tilde{\mathbf{c}}(\omega) \tilde{\mathbf{f}}(\omega) \right\} \qquad \text{where } \tilde{\mathbf{c}}(\omega) \triangleq \tilde{\mathbf{b}}(\omega) \tilde{\mathbf{a}}(\omega) \\ & = \left\{ \mathbf{f}(x) | \mathbf{f}(x) = \sum_{n \in \mathbb{Z}} \mathbf{c}_n \mathbf{f}(x - n) \right\} \\ & \triangleq \boldsymbol{V}_0' \end{split}$$

 \blacksquare

2.4 Linear combinations in inner product spaces

Definition 2.9. Let $(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle | \nabla \rangle)$ be an INNER PRODUCT SPACE (Definition C.9 page 124).

Two vectors \mathbf{x} and \mathbf{y} in X are **orthogonal** if $\langle \mathbf{x} \mid \mathbf{y} \rangle = \begin{cases} 0 & \text{for } \mathbf{x} \neq \mathbf{y} \\ c \in \mathbb{F} \setminus 0 & \text{for } \mathbf{x} = \mathbf{y} \end{cases}$

In an *inner product space*, *orthogonality* is a special case of *linear independence*; or alternatively, linear independence is a generalization of orthogonality (next theorem).

Theorem 2.5. ¹⁹ Let $\{x_n \in X \mid n=1,2,...,N\}$ be a set of vectors in an INNER PRODUCT SPACE (Definition C.9 page 124) $(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle \mid \nabla \rangle)$.

NPROOF:

1. Proof using *Pythagorean theorem*: Let $(\alpha_n)_{n\in\mathbb{N}}$ be a sequence with at least one nonzero element.

¹⁹ ⚠ Aliprantis and Burkinshaw (1998) page 283 ⟨Corollary 32.8⟩, Д Kubrusly (2001) page 352 ⟨Proposition 5.34⟩



$$\left\| \sum_{n=1}^{N} \alpha_n \mathbf{x}_n \right\|^2 = \sum_{n=1}^{N} \|\alpha_n \mathbf{x}_n\|^2$$
 by left hypoth. and *Pythagorean Theorem*
$$= \sum_{n=1}^{N} |\alpha_n|^2 \|\mathbf{x}_n\|^2$$
 by definition of $\|\cdot\|$ (Definition C.5 page 116) > 0

$$\implies \sum_{n=1}^{N} \alpha_n x_n \neq 0$$

 $\implies (x_n)_{n\in\mathbb{N}}$ is linearly independent—by definition of linear independence

(Definition 2.3 page 9)

2. Alternative proof:

$$\sum_{n=1}^{N} \alpha_{n} \mathbf{x}_{n} = 0 \implies \left\langle \sum_{n=1}^{N} \alpha_{n} \mathbf{x}_{n} \mid \mathbf{x}_{m} \right\rangle = \langle 0 \mid \mathbf{x}_{m} \rangle$$

$$\implies \sum_{n=1}^{N} \alpha_{n} \langle \mathbf{x}_{n} \mid \mathbf{x}_{m} \rangle = 0$$

$$\implies \sum_{n=1}^{N} \alpha_{n} \bar{\delta}(k - m) = 0$$

$$\implies \alpha_{m} = 0 \quad \text{for } m = 1, 2, ..., N$$

Theorem 2.6 (Bessel's Equality). ²⁰ Let $\{x_n \in X \mid n=1,2,...,N\}$ be a set of vectors in an INNER PRODUCT SPACE (Definition C.9 page 124) $(X, +, \cdot, (\mathbb{F}, \dotplus, \dot{\times}), \langle \triangle \mid \nabla \rangle)$ and with $\|x\| \triangleq \sqrt{\langle x \mid x \rangle}$.

 $\left\{ \begin{array}{c} \left\{ \boldsymbol{x}_{n} \right\} \text{ is ORTHONORMAL} \\ \text{(Definition 2.9 page 16)} \end{array} \right\} \quad \Longrightarrow \quad \left\{ \underbrace{\left\| \boldsymbol{x} - \sum_{n=1}^{N} \left\langle \boldsymbol{x} \mid \boldsymbol{x}_{n} \right\rangle \boldsymbol{x}_{n} \right\|^{2}}_{approximation \ error} = \|\boldsymbol{x}\|^{2} - \sum_{n=1}^{N} |\left\langle \boldsymbol{x} \mid \boldsymbol{x}_{n} \right\rangle|^{2} \quad \forall \boldsymbol{x} \in X \right\}$

№PROOF:

$$\left\| \mathbf{x} - \sum_{n=1}^{N} \langle \mathbf{x} \mid \mathbf{x}_{n} \rangle \mathbf{x}_{n} \right\|^{2}$$

$$= \|\mathbf{x}\|^{2} + \left\| \sum_{n=1}^{N} \langle \mathbf{x} \mid \mathbf{x}_{n} \rangle \mathbf{x}_{n} \right\|^{2} - 2\Re \left\langle \mathbf{x} \mid \sum_{n=1}^{N} \langle \mathbf{x} \mid \mathbf{x}_{n} \rangle \mathbf{x}_{n} \right\rangle \quad \text{by polar identity}$$

$$= \|\mathbf{x}\|^{2} + \left\| \sum_{n=1}^{N} \langle \mathbf{x} \mid \mathbf{x}_{n} \rangle \mathbf{x}_{n} \right\|^{2} - 2\Re \left[\left(\sum_{n=1}^{N} \langle \mathbf{x} \mid \mathbf{x}_{n} \rangle \right)^{*} \langle \mathbf{x} \mid \mathbf{x}_{n} \rangle \right] \quad \text{by property of } \langle \triangle \mid \nabla \rangle \quad \text{(Definition C.9 page 124)}$$

$$= \|\mathbf{x}\|^{2} + \sum_{n=1}^{N} \|\langle \mathbf{x} \mid \mathbf{x}_{n} \rangle \mathbf{x}_{n} \|^{2} - 2\Re \left[\left(\sum_{n=1}^{N} \langle \mathbf{x} \mid \mathbf{x}_{n} \rangle \right)^{*} \langle \mathbf{x} \mid \mathbf{x}_{n} \rangle \right] \quad \text{by Pythagorean Theorem}$$

²⁰ Bachman et al. (2002) page 103, Pedersen (2000) pages 38–39

$$= \|\mathbf{x}\|^2 + \sum_{n=1}^{N} \|\langle \mathbf{x} \mid \mathbf{x}_n \rangle \mathbf{x}_n\|^2 - 2\Re \left(\sum_{n=1}^{N} \langle \mathbf{x} \mid \mathbf{x}_n \rangle^* \langle \mathbf{x} \mid \mathbf{x}_n \rangle \right)$$

$$= \|\mathbf{x}\|^2 + \sum_{n=1}^{N} |\langle \mathbf{x} \mid \mathbf{x}_n \rangle|^2 \underbrace{\|\mathbf{x}_n\|^2}_{1} - 2\Re \left(\sum_{n=1}^{N} \langle \mathbf{x} \mid \mathbf{x}_n \rangle^* \langle \mathbf{x} \mid \mathbf{x}_n \rangle \right)$$
 by property of $\|\cdot\|$ (Definition C.5 page 116)
$$= \|\mathbf{x}\|^2 + \sum_{n=1}^{N} |\langle \mathbf{x} \mid \mathbf{x}_n \rangle|^2 \cdot 1 - 2\Re \left(\sum_{n=1}^{N} \langle \mathbf{x} \mid \mathbf{x}_n \rangle^* \langle \mathbf{x} \mid \mathbf{x}_n \rangle \right)$$
 by def. of *orthonormality* (Definition 2.9 page 16)
$$= \|\mathbf{x}\|^2 + \sum_{n=1}^{N} |\langle \mathbf{x} \mid \mathbf{x}_n \rangle|^2 - 2\Re \sum_{n=1}^{N} |\langle \mathbf{x} \mid \mathbf{x}_n \rangle|^2$$
 because $|\cdot|$ is real
$$= \|\mathbf{x}\|^2 - \sum_{n=1}^{N} |\langle \mathbf{x} \mid \mathbf{x}_n \rangle|^2$$

Theorem 2.7 (Bessel's inequality). ²¹ Let $\{x_n \in X \mid n=1,2,...,N\}$ be a set of vectors in an INNER PRODUCT SPACE (Definition C.9 page 124) $(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle | \nabla \rangle)$ and with $||x|| \triangleq \sqrt{\langle x | x \rangle}$.

$$\left\{ \begin{array}{c} \left\{ \boldsymbol{x}_{n} \right\} \text{ is ORTHONORMAL} \\ \text{(Definition 2.9 page 16)} \end{array} \right\} \quad \Longrightarrow \quad \left\{ \begin{array}{c} \sum_{n=1}^{N} \left| \langle \boldsymbol{x} \mid \boldsymbol{x}_{n} \rangle \right|^{2} & \leq & \|\boldsymbol{x}\|^{2} & \forall \boldsymbol{x} \in \boldsymbol{X} \end{array} \right\}$$

№ Proof:

$$0 \le \left\| \mathbf{x} - \sum_{n=1}^{N} \left\langle \mathbf{x} \mid \mathbf{x}_{n} \right\rangle \mathbf{x}_{n} \right\|^{2}$$
 by definition of $\| \cdot \|$ (Definition C.5 page 116)
$$= \left\| \mathbf{x} \right\|^{2} - \sum_{n=1}^{N} \left| \left\langle \mathbf{x} \mid \mathbf{x}_{n} \right\rangle \right|^{2}$$
 by Bessel's Equality (Theorem 2.6 page 17)

The Best Approximation Theorem (next) shows that

- the best sequence for representing a vector is the sequence of projections of the vector onto the sequence of basis functions
- the error of the projection is orthogonal to the projection.

Theorem 2.8 (Best Approximation Theorem). ²² Let $\{x_n \in X \mid n=1,2,...,N\}$ be a set of vectors in an INNER PRODUCT SPACE (Definition C.9 page 124) $(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle \mid \nabla \rangle)$ and with $\|x\| \triangleq \sqrt{\langle x \mid x \rangle}$.

²² Walter and Shen (2001) pages 3–4, Pedersen (2000) page 39, Edwards (1995) pages 94–100, Weyl (1940)



²¹ ☐ Giles (2000) pages 54–55 (3.13 Bessel's inequality), ☐ Bollobás (1999) page 147, ☐ Aliprantis and Burkinshaw (1998) page 284

$$\left\{
\begin{cases}
\left\{x_{n}\right\} is \\
\text{ORTHONORMAL} \\
\left(\text{Definition 2.9 page 16}\right)
\end{cases}
\Longrightarrow$$

1.
$$\arg\min_{\left(\alpha_{n}\right)_{n=1}^{N}}\left\|\mathbf{x}-\sum_{n=1}^{N}\alpha_{n}\mathbf{x}_{n}\right\| = \underbrace{\left(\left\langle\mathbf{x}\mid\mathbf{x}_{n}\right\rangle\right)_{n=1}^{N}}_{best\ \alpha_{n}=\left\langle\mathbf{x}\mid\mathbf{x}_{n}\right\rangle} \quad \forall \mathbf{x} \in X \quad and$$

2. $\underbrace{\left(\sum_{n=1}^{N}\left\langle\mathbf{x}\mid\mathbf{x}_{n}\right\rangle\mathbf{x}_{n}\right)}_{approximation} \perp \underbrace{\left(\mathbf{x}-\sum_{n=1}^{N}\left\langle\mathbf{x}\mid\mathbf{x}_{n}\right\rangle\mathbf{x}_{n}\right)}_{approximation} \quad \forall \mathbf{x} \in X$

^ℚProof:

1. Proof that $(\langle x \mid x_n \rangle)$ is the best sequence:

$$\begin{split} & \left\| \boldsymbol{x} - \sum_{n=1}^{N} \alpha_{n} \boldsymbol{x}_{n} \right\|^{2} \\ &= \|\boldsymbol{x}\|^{2} - 2\Re\left\langle \boldsymbol{x} \mid \sum_{n=1}^{N} \alpha_{n} \boldsymbol{x}_{n} \right\rangle + \left\| \sum_{n=1}^{N} \alpha_{n} \boldsymbol{x}_{n} \right\|^{2} \\ &= \|\boldsymbol{x}\|^{2} - 2\Re\left(\sum_{n=1}^{N} \alpha_{n}^{*} \langle \boldsymbol{x} \mid \boldsymbol{x}_{n} \rangle \right) + \sum_{n=1}^{N} \|\alpha_{n} \boldsymbol{x}_{n}\|^{2} \quad \text{by Pythagorean Theorem} \\ &= \|\boldsymbol{x}\|^{2} - 2\Re\left(\sum_{n=1}^{N} \alpha_{n}^{*} \langle \boldsymbol{x} \mid \boldsymbol{x}_{n} \rangle \right) + \sum_{n=1}^{N} \|\alpha_{n}\|^{2} + \underbrace{\left[\sum_{n=1}^{N} \left| \langle \boldsymbol{x} \mid \boldsymbol{x}_{n} \rangle \right|^{2} - \sum_{n=1}^{N} \left| \langle \boldsymbol{x} \mid \boldsymbol{x}_{n} \rangle \right|^{2} \right]}_{0} \\ &= \left[\|\boldsymbol{x}\|^{2} - \sum_{n=1}^{N} \left| \langle \boldsymbol{x} \mid \boldsymbol{x}_{n} \rangle \right|^{2} \right] + \sum_{n=1}^{N} \left[\left| \langle \boldsymbol{x} \mid \boldsymbol{x}_{n} \rangle \right|^{2} - 2\Re_{\mathbf{e}} \left[\alpha_{n}^{*} \langle \boldsymbol{x} \mid \boldsymbol{x}_{n} \rangle \right] + \left| \alpha_{n} \right|^{2} \right] \\ &= \left[\|\boldsymbol{x}\|^{2} - \sum_{n=1}^{N} \left| \langle \boldsymbol{x} \mid \boldsymbol{x}_{n} \rangle \right|^{2} \right] + \sum_{n=1}^{N} \left[\left| \langle \boldsymbol{x} \mid \boldsymbol{x}_{n} \rangle \right|^{2} - \alpha_{n}^{*} \langle \boldsymbol{x} \mid \boldsymbol{x}_{n} \rangle - \alpha_{n} \langle \boldsymbol{x} \mid \boldsymbol{x}_{n} \rangle^{*} + |\alpha_{n}|^{2} \right] \\ &= \left\| \boldsymbol{x} - \sum_{n=1}^{N} \langle \boldsymbol{x} \mid \boldsymbol{x}_{n} \rangle \boldsymbol{x}_{n} \right\|^{2} + \sum_{n=1}^{N} \left| \langle \boldsymbol{x} \mid \boldsymbol{x}_{n} \rangle - \alpha_{n} \right|^{2} \quad \text{by Bessel's Equality} \end{aligned} \tag{Theorem 2.6 page 17)} \\ &\geq \left\| \boldsymbol{x} - \sum_{n=1}^{N} \langle \boldsymbol{x} \mid \boldsymbol{x}_{n} \rangle \boldsymbol{x}_{n} \right\|^{2} \end{aligned}$$

2. Proof that the approximation and approximation error are orthogonal:

$$\left\langle \sum_{n=1}^{N} \langle \mathbf{x} \mid \mathbf{x}_{n} \rangle \mathbf{x}_{n} \mid \mathbf{x} - \sum_{n=1}^{N} \langle \mathbf{x} \mid \mathbf{x}_{n} \rangle \mathbf{x}_{n} \right\rangle = \left\langle \sum_{n=1}^{N} \langle \mathbf{x} \mid \mathbf{x}_{n} \rangle \mathbf{x}_{n} \mid \mathbf{x} \right\rangle - \left\langle \sum_{n=1}^{N} \langle \mathbf{x} \mid \mathbf{x}_{n} \rangle \mathbf{x}_{n} \mid \sum_{n=1}^{N} \langle \mathbf{x} \mid \mathbf{x}_{n} \rangle \mathbf{x}_{n} \right\rangle$$

$$= \sum_{n=1}^{N} \langle \mathbf{x} \mid \mathbf{x}_{n} \rangle^{*} \langle \mathbf{x} \mid \mathbf{x}_{n} \rangle - \sum_{n=1}^{N} \sum_{m=1}^{N} \langle \mathbf{x} \mid \mathbf{x}_{n} \rangle \langle \mathbf{x} \mid \mathbf{x}_{m} \rangle^{*} \langle \mathbf{x}_{n} \mid \mathbf{x}_{m} \rangle$$

$$= \sum_{n=1}^{N} |\langle \mathbf{x} \mid \mathbf{x}_{n} \rangle|^{2} - \sum_{n=1}^{N} \sum_{m=1}^{N} \langle \mathbf{x} \mid \mathbf{x}_{n} \rangle \langle \mathbf{x} \mid \mathbf{x}_{m} \rangle^{*} \bar{\delta}_{nm}$$

$$= \sum_{n=1}^{N} |\langle \mathbf{x} \mid \mathbf{x}_{n} \rangle|^{2} - \sum_{n=1}^{N} |\langle \mathbf{x} \mid \mathbf{x}_{n} \rangle|^{2}$$

Orthonormal bases in Hilbert spaces 2.5

Definition 2.10. Let $\{x_n \in X | n=1,2,...,N\}$ be a set of vectors in an INNER PRODUCT SPACE (Definition C.9) $\textit{page 124)} \ \pmb{\varOmega} \triangleq \big(X, \, +, \, \cdot, \, (\mathbb{F}, \, \dot{+}, \, \dot{\times}), \, \big\langle \triangle \, | \, \nabla \big\rangle \big).$

D E F

The set $\{x_n\}$ is an **orthogonal basis** for Ω if $\{x_n\}$ is ORTHOGONAL and is *a* Schauder basis for Ω .

The set $\{x_n\}$ is an **orthonormal basis** for Ω if $\{x_n\}$ is orthonormal and is *a* Schauder basis for Ω .

Definition 2.11. ²³ Let $\mathbf{H} \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle | \nabla \rangle)$ be a Hilbert space.

Suppose there exists a set $\{x_n \in X \mid n \in \mathbb{N}\}$ such that $x \stackrel{*}{=} \sum_{i=1}^n \langle x \mid x_n \rangle x_n$.

Then the quantities $\langle x \mid x_n \rangle$ are called the **Fourier coefficients** of x and the sum $\sum_{n=1}^{\infty} \langle x \mid x_n \rangle x_n \text{ is called the Fourier expansion of } x \text{ or the Fourier series for } x.$

Definition 2.12.



The **Kronecker delta function** $\bar{\delta}_n$ is defined as $\bar{\delta}_n \triangleq \begin{cases} 1 & \text{for } n = 0 \\ 0 & \text{otherwise} \end{cases}$

$$\bar{\delta}_n \triangleq \left\{ \begin{array}{ll} 1 & \textit{for } n = 0 & \textit{and} \\ 0 & \textit{otherwise} \end{array} \right. \quad \forall n \in \mathbb{Z}$$

Lemma 2.2 (Perfect reconstruction). Let $\{x_n \in X \mid n \in \mathbb{N}\}$ be a set of vectors in a Hilbert space $\mathbf{H} \triangleq$ $(X, +, \cdot, (\mathbb{F}, +, \dot{\times}), \langle \triangle \mid \nabla \rangle).$

$$\left\{ \begin{array}{ll} \text{(1).} & (\mathbf{x}_n) \text{ is a BASIS for } \mathbf{H} \\ \text{(2).} & (\mathbf{x}_n) \text{ is ORTHONORMAL} \end{array} \right\} \qquad \Longrightarrow \qquad \mathbf{x} \stackrel{\star}{=} \sum_{n=1}^{\infty} \underbrace{\langle \mathbf{x} \mid \mathbf{x}_n \rangle}_{Fourier coefficient} \mathbf{x}_n \qquad \forall \mathbf{x} \in X$$

^ℚProof:

$$\langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle = \left\langle \sum_{m \in \mathbb{Z}} \alpha_m \boldsymbol{x}_m \mid \boldsymbol{x}_n \right\rangle \qquad \text{by left hypothesis (1)}$$

$$= \sum_{m \in \mathbb{Z}} \alpha_m \left\langle \boldsymbol{x}_m \mid \boldsymbol{x}_n \right\rangle \qquad \text{by } homogeneous \text{ property of } \langle \triangle \mid \nabla \rangle \qquad \text{(Definition C.9 page 124)}$$


$$= \sum_{m \in \mathbb{Z}} \alpha_m \bar{\delta}_{n-m} \qquad \text{by left hypothesis (2)} \qquad \text{(Definition 2.9 page 16)}$$

$$= \alpha_n$$

Proposition 2.2. ²⁴ Let $\{x_n \in X \mid n \in \mathbb{N}\}$ be a set of vectors in a Hilbert space $\mathbf{H} \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle \mid \nabla \rangle).$

²³ Fabian et al. (2010) page 27 ⟨Theorem 1.55⟩, Young (2001) page 6, Young (1980) page 6 ²⁴ Han et al. (2007) pages 93–94 (Proposition 3.11)





♥Proof:

1. Proof that Parseval frame \leftarrow Fourier expansion

$$\|x\|^{2} \triangleq \langle x \mid x \rangle \qquad \text{by definition of } \|\cdot\|$$

$$= \left\langle \sum_{n=1}^{\infty} \langle x \mid x_{n} \rangle x \mid x_{n} \right\rangle \qquad \text{by right hypothesis}$$

$$\stackrel{+}{=} \sum_{n=1}^{\infty} \langle x \mid x_{n} \rangle \langle x \mid x_{n} \rangle \qquad \text{by property of } \langle \triangle \mid \nabla \rangle$$

$$\stackrel{+}{=} \sum_{n=1}^{\infty} \langle x \mid x_{n} \rangle \langle x \mid x_{n} \rangle^{*} \qquad \text{by property of } \langle \triangle \mid \nabla \rangle$$

$$\stackrel{+}{=} \sum_{n=1}^{\infty} \left| \langle x \mid x_{n} \rangle \right|^{2} \qquad \text{by property of } \mathbb{C} \qquad \text{(Definition F.7 page 151)}$$

2. Proof that Parseval frame \implies Fourier expansion

- (a) Let $(e_n)_{n\in\mathbb{N}}$ be the *standard othornormal basis* such that the *n*th element of e_n is 1 and all other elements are 0.
- (b) Let **M** be an operator in **H** such that $\mathbf{M} \mathbf{x} \triangleq \sum_{n=1}^{\infty} \langle \mathbf{x} \mid \mathbf{x}_n \rangle e_n$.
- (c) lemma: **M** is *isometric*. Proof:

$$\|\mathbf{M}\boldsymbol{x}\|^{2} = \left\|\sum_{n=1}^{\infty} \langle \boldsymbol{x} \mid \boldsymbol{x}_{n} \rangle e_{n}\right\|^{2} \qquad \text{by definition of } \mathbf{M} \qquad \text{(item (2b) page 21)}$$

$$= \sum_{n=1}^{\infty} \left\|\langle \boldsymbol{x} \mid \boldsymbol{x}_{n} \rangle e_{n}\right\|^{2} \qquad \text{by } Pythagorean \ Theorem}$$

$$= \sum_{n=1}^{\infty} \left|\langle \boldsymbol{x} \mid \boldsymbol{x}_{n} \rangle\right|^{2} \left\|e_{n}\right\|^{2} \qquad \text{by } homogeneous \ property of } \|\cdot\| \qquad \text{(Definition C.5 page 116)}$$

$$= \sum_{n=1}^{\infty} \left|\langle \boldsymbol{x} \mid \boldsymbol{x}_{n} \rangle\right|^{2} \qquad \text{by definition of } orthonormal \qquad \text{(Definition 2.9 page 16)}$$

$$= \|\boldsymbol{x}\|^{2} \qquad \text{by Parseval frame hypothesis}$$

$$\implies \mathbf{M} \text{ is } isometric \qquad \text{by definition of } isometric \qquad \text{(Definition C.13 page 132)}$$

(d) Let $(u_n)_{n\in\mathbb{N}}$ be an *orthornormal basis* for H.





(e) Proof for Fourier expansion:

$$x = \sum_{n=1}^{\infty} \langle x | u_n \rangle u_n \qquad \text{by Fourier expansion (Proposition 2.3 page 24)}$$

$$= \sum_{n=1}^{\infty} \langle \mathbf{M} x | \mathbf{M} u_n \rangle u_n \qquad \text{by (2c) lemma page 21 and Theorem C.23 page 133}$$

$$= \sum_{n=1}^{\infty} \left\langle \sum_{m=1}^{\infty} \langle x | x_m \rangle e_m | \sum_{k=1}^{\infty} \langle u_n | x_k \rangle e_k \right\rangle u_n \quad \text{by item (2b) page 21}$$

$$= \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \langle x | x_m \rangle \sum_{k=1}^{\infty} \langle u_n | x_k \rangle^* \langle e_m | e_k \rangle u_n \quad \text{by Definition C.9 page 124}$$

$$= \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \langle x | x_m \rangle \langle u_n | x_m \rangle^* u_n \quad \text{by item (2a) page 21 and Definition 2.9 page 16}$$

$$= \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \langle x | x_m \rangle \langle x_m | u_n \rangle u_n \quad \text{by Definition C.9 page 124}$$

$$= \sum_{m=1}^{\infty} \langle x | x_m \rangle \sum_{n=1}^{\infty} \langle x_m | u_n \rangle u_n \quad \text{by Definition C.9 page 124}$$

$$= \sum_{m=1}^{\infty} \langle x | x_m \rangle \sum_{n=1}^{\infty} \langle x_m | u_n \rangle u_n \quad \text{by item (2d) page 21}$$

When is a set of orthonormal vectors in a Hilbert space *H total*? Theorem 2.9 (next) offers some help.

Theorem 2.9 (The Fourier Series Theorem). ²⁵ Let $\{x_n \in X \mid n \in \mathbb{N}\}$ be a set of vectors in a Hilbert

Theorem 2.9 (The Fourier Series Theorem). Let
$$\{x_n \in X \mid n \in \mathbb{N}\}\$$
 be a set of vectors in a function of the series and the series are considered by the series of vectors in a function of the series are considered by the series are considered by the series expansion of the series expansion

^ℚProof:

²⁵ ■ Bachman and Narici (1966) pages 149–155 (Theorem 9.12), ■ Kubrusly (2001) pages 360–363 (Theorem 5.48), Aliprantis and Burkinshaw (1998) pages 298–299 (Theorem 34.2), Christensen (2003) page 57 (Theorem 3.4.2), Berberian (1961) pages 52–53 ⟨Theorem II§8.3⟩, Heil (2011) pages 34–35 ⟨Theorem 1.50⟩, Bracewell (1978) page 112 (Rayleigh's theorem)



₽

1. Proof that $(1) \Longrightarrow (2)$:

2. Proof that $(2) \Longrightarrow (3)$:

$$||x||^{2} \triangleq \langle x | x \rangle$$
 by definition of *induced norm*

$$= \sum_{n=1}^{\infty} \langle x | x_{n} \rangle \langle x | x_{n} \rangle^{*}$$
 by (2)
$$= \sum_{n=1}^{\infty} |\langle x | x_{n} \rangle|^{2}$$

- 3. Proof that (3) \iff (4) *not* using (A): by Proposition 2.2 page 20
- 4. Proof that (3) \Longrightarrow (1) (proof by contradiction):
 - (a) Suppose $\{x_n\}$ is *not total*.
 - (b) Then there must exist a vector y in H such that the set $B \triangleq \{x_n\} \cup y$ is *orthonormal*.

(c) Then
$$1 = ||y||^2 \neq \sum_{n=1}^{\infty} |\langle y | x_n \rangle|^2 = 0$$
.

- (d) But this contradicts (3), and so $\{x_n\}$ must be *total* and (3) \Longrightarrow (1).
- 5. Extraneous proof that $(3) \Longrightarrow (4)$ (this proof is not really necessary here):

$$\left\| \mathbf{x} - \sum_{n=1}^{\infty} \langle \mathbf{x} \mid \mathbf{x}_n \rangle \mathbf{x}_n \right\|^2 = \|\mathbf{x}\|^2 - \sum_{n=1}^{\infty} |\langle \mathbf{x} \mid \mathbf{x}_n \rangle|^2 \qquad \text{by } Bessel's Equality} \qquad \text{(Theorem 2.6 page 17)}$$

$$= 0 \qquad \qquad \text{by (3)}$$

$$\implies \mathbf{x} \stackrel{\star}{=} \sum_{n=1}^{\infty} \langle \mathbf{x} \mid \mathbf{x}_n \rangle \mathbf{x}_n \qquad \text{by definition of } \stackrel{\star}{=}$$

- 6. Extraneous proof that (A) \Longrightarrow (4) (this proof is not really necessary here)
 - (a) The sequence $\sum_{n=1}^{N} |\langle x | x_n \rangle|^2$ is *monotonically increasing* in *n*.
 - (b) By Bessel's inequality (page 18), the sequence is upper bounded by $||x||^2$:

$$\sum_{n=1}^{N} \left| \langle \mathbf{x} \mid \mathbf{x}_n \rangle \right|^2 \le \|\mathbf{x}\|^2$$





—>

(c) Because this sequence is both monotonically increasing and bounded in n, it must equal its bound in the limit as n approaches infinity:

$$\lim_{N \to \infty} \sum_{n=1}^{N} \left| \langle \mathbf{x} \mid \mathbf{x}_n \rangle \right|^2 = \|\mathbf{x}\|^2 \tag{2.1}$$

(d) If we combine this result with Bessel's Equality (Theorem 2.6 page 17) we have

$$\lim_{N \to \infty} \left\| \mathbf{x} - \sum_{n=1}^{N} \langle \mathbf{x} \mid \mathbf{x}_n \rangle \mathbf{x}_n \right\|^2 = \|\mathbf{x}\|^2 - \lim_{N \to \infty} \sum_{n=1}^{N} |\langle \mathbf{x} \mid \mathbf{x}_n \rangle|^2 \quad \text{by Bessel's Equality (Theorem 2.6 page 17)}$$

$$= \|\mathbf{x}\|^2 - \|\mathbf{x}\|^2 \quad \text{by equation (2.1) page 24}$$

$$= 0$$

Proposition 2.3 (Fourier expansion). Let $\{x_n \in X \mid n \in \mathbb{N}\}$ be a set of vectors in a Hilbert space $(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle \mid \nabla \rangle)$.

$$\begin{cases}
x_n \text{ is an Orthonormal Basis } for \mathbf{H} \\
(A)
\end{cases}
\implies
\begin{cases}
x = \sum_{n=1}^{\infty} \alpha_n x_n \\
(1)
\end{cases}
\iff
\underbrace{\alpha_n = \langle x \mid x_n \rangle}_{(2)}$$

№ Proof:

- 1. Proof that (1) \Longrightarrow (2): by Lemma 2.2 page 20
- 2. Proof that $(1) \Leftarrow (2)$:


$$\left\| \mathbf{x} - \sum_{n=1}^{\infty} \alpha_n \mathbf{x}_{n \in \mathbb{Z}} \right\|^2 = \left\| \mathbf{x} - \sum_{n=1}^{\infty} \langle \mathbf{x} \mid \mathbf{x}_n \rangle \mathbf{x}_{n \in \mathbb{Z}} \right\|^2 \quad \text{by right hypothesis}$$

$$= \|\mathbf{x}\|^2 - \sum_{n=1}^{\infty} |\langle \mathbf{x} \mid \mathbf{x}_n \rangle|^2 \quad \text{by Bessel's equality} \qquad \text{(Theorem 2.6 page 17)}$$

$$= 0 \quad \qquad \text{by $Parseval's Identity} \qquad \text{(Theorem 2.9 page 22)}$$

$$\stackrel{\text{def}}{\iff} \mathbf{x} \stackrel{*}{=} \sum_{n=1}^{\infty} \langle \mathbf{x} \mid \mathbf{x}_n \rangle \mathbf{x}_n \quad \text{by definition of $strong convergence}$$

Proposition 2.4 (Riesz-Fischer Theorem). ²⁶ Let $\{x_n \in X \mid n \in \mathbb{N}\}$ be a set of vectors in a Hilbert space $(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle \mid \nabla \rangle)$.



[♠]Proof:

²⁶ Young (2001) page 6



- 1. Proof that $(1) \Longrightarrow (2)$:
 - (a) If (1) is true, then let $\mathbf{x} \triangleq \sum_{n \in \mathbb{N}} \alpha_n \mathbf{x}_n$.
 - (b) Then

$$\begin{split} \langle \boldsymbol{x} \, | \, \boldsymbol{x}_n \rangle &= \left\langle \sum_{m \in \mathbb{N}} \alpha_m \boldsymbol{x}_m \, | \, \boldsymbol{x}_n \right\rangle & \text{by definition of } \boldsymbol{x} \\ &= \sum_{m \in \mathbb{N}} \alpha_m \left\langle \boldsymbol{x}_m \, | \, \boldsymbol{x}_n \right\rangle & \text{by } homogeneous \text{ property of } \langle \triangle \, | \, \nabla \rangle & \text{(Definition C.9 page 124)} \\ &= \sum_{m \in \mathbb{N}} \alpha_m \bar{\delta}_{mn} & \text{by (A)} \\ &= \sum_{m \in \mathbb{N}} \alpha_n & \text{by definition of } \bar{\delta} & \text{(Definition 2.12 page 20)} \end{split}$$

2. Proof that $(1) \Leftarrow (2)$:

$$\sum_{n \in \mathbb{N}} |\alpha_n|^2 = \sum_{n \in \mathbb{N}} |\langle \mathbf{x} \mid \mathbf{x}_n \rangle|^2$$
 by (2)
$$\leq ||\mathbf{x}||^2$$
 by Bessel's Inequality (Theorem 2.7 page 18)
$$\leq \infty$$

Theorem 2.10. 27

.

T H M All separable Hilbert spaces are isomorphic. That is, $\begin{cases}
X \text{ is a separable} \\
Hilbert space & and} \\
Y \text{ is a separable} \\
Hilbert space
\end{cases}
\Rightarrow
\begin{cases}
there is a bijective operator <math>\mathbf{M} \in Y^X$ such that $\mathbf{M} = \mathbf{M} = \mathbf{M$

[♠]Proof:

- 1. Let $\mathbf{X} \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle \mid \nabla \rangle)$ be a *separable Hilbert space* with *orthonormal basis* $\{\mathbf{x}_n \mid n \in \mathbb{N}\}$. Let $\mathbf{Y} \triangleq (Y, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle \mid \nabla \rangle)$ be a *separable Hilbert space* with *orthonormal basis* $\{\mathbf{y}_n \mid n \in \mathbb{N}\}$.
- 2. Proof that there exists *bijective* operator **M** and its inverse \mathbf{M}^{-1} between $\{x_n\}$ and $\{y_n\}$:
 - (a) Let **M** be defined such that $y_n \triangleq \mathbf{M}x_n$.
 - (b) Thus **M** is a *bijection* between $\{x_n\}$ and $\{y_n\}$.
 - (c) Because **M** is a *bijection* between $\{x_n\}$ and $\{y_n\}$, **M** has an inverse operator \mathbf{M}^{-1} between $\{x_n\}$ and $\{y_n\}$ such that $x_n = \mathbf{M}^{-1}y_n$.
- 3. Proof that M and M^{-1} are *bijective* operators between X and Y:

(a) Proof that **M** maps **X** into **Y**:

$$x \in X \iff x \stackrel{*}{=} \sum_{n \in \mathbb{N}} \langle x \mid x_n \rangle x_n \qquad \qquad \text{by } Fourier \ expansion} \qquad \text{(Theorem 2.9 page 22)}$$

$$\implies \exists y \in Y \quad \text{such that} \quad \langle y \mid y_n \rangle = \langle x \mid x_n \rangle \quad \text{by } Riesz\text{-}Fischer \ Thm. \qquad \text{(Proposition 2.4 page 24)}$$

$$\implies y = \sum_{n \in \mathbb{N}} \langle y \mid y_n \rangle y_n \qquad \qquad \text{by } Fourier \ expansion} \qquad \text{(Theorem 2.9 page 22)}$$

$$= \sum_{n \in \mathbb{N}} \langle x \mid x_n \rangle y_n \qquad \qquad \text{by } Riesz\text{-}Fischer \ Thm. \qquad \text{(Proposition 2.4 page 24)}$$

$$= \sum_{n \in \mathbb{N}} \langle x \mid x_n \rangle M x_n \qquad \qquad \text{by definition of } M \qquad \text{(item (2a) page 25)}$$

$$= M \sum_{n \in \mathbb{N}} \langle x \mid x_n \rangle x_n \qquad \qquad \text{by prop. of linear ops.} \qquad \text{(Theorem C.1 page 113)}$$

$$= Mx \qquad \qquad \text{by definition of } x$$

(b) Proof that M^{-1} maps Y into X:

4. Proof for (2):

$$\|\mathbf{M}\mathbf{x}\|^{2} = \left\|\mathbf{M}\sum_{n\in\mathbb{N}}\langle\mathbf{x}\,|\,\mathbf{x}_{n}\rangle\,\mathbf{x}_{n}\right\|^{2} \qquad \text{by } Fourier \, expansion} \qquad \text{(Theorem 2.9 page 22)}$$

$$= \left\|\sum_{n\in\mathbb{N}}\langle\mathbf{x}\,|\,\mathbf{x}_{n}\rangle\,\mathbf{M}\mathbf{x}_{n}\right\|^{2} \qquad \text{by property of } linear \, operators \qquad \text{(Theorem C.1 page 113)}$$

$$= \left\|\sum_{n\in\mathbb{N}}\langle\mathbf{x}\,|\,\mathbf{x}_{n}\rangle\,\mathbf{y}_{n}\right\|^{2} \qquad \text{by definition of } \mathbf{M} \qquad \text{(item (2a) page 25)}$$

$$= \sum_{n\in\mathbb{N}}\left|\langle\mathbf{x}\,|\,\mathbf{x}_{n}\rangle\right|^{2} \qquad \text{by } Parseval's \, Identity \qquad \text{(Proposition 2.4 page 24)}$$

$$= \left\|\sum_{n\in\mathbb{N}}\langle\mathbf{x}\,|\,\mathbf{x}_{n}\rangle\,\mathbf{x}_{n}\right\|^{2} \qquad \text{by } Parseval's \, Identity \qquad \text{(Proposition 2.4 page 24)}$$

$$= \|\mathbf{x}\|^{2} \qquad \text{by } Fourier \, expansion \qquad \text{(Theorem 2.9 page 22)}$$

5. Proof for (3): by (2) and Theorem C.23 page 133

Theorem 2.11. ²⁸ *Let H be a* HILBERT SPACE.

H is SEPARABLE *H has a* Schauder basis

Theorem 2.12. 29 *Let* H *be* a Hilbert space

H has an Orthonormal basis *H* is separable

Riesz bases in Hilbert spaces 2.6

Definition 2.13. 30 Let $\{x_n \in X \mid n \in \mathbb{N}\}$ be a set of vectors in a SEPARABLE HILBERT SPACE $\mathbf{H} \triangleq (X, +, \cdot, (\mathbb{F}, +, \cdot, \mathbb{F}))$

 $\{m{x}_n\}$ is a **Riesz basis** for **H** if $\{m{x}_n\}$ is EQUIVALENT (Definition 2.8 page 15) to some ORTHONORMAL BASIS (Definition 2.10 page 20) in H.

Definition 2.14. ³¹ Let $(x_n \in X)_{n \in \mathbb{N}}$ be a sequence of vectors in a Separable Hilbert space $\mathbf{H} \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle \mid \nabla \rangle).$

The sequence (x_n) is a **Riesz sequence** for **H** if $A\sum_{n=1}^{\infty} |\alpha_n|^2 \le \left\| \sum_{n=1}^{\infty} \alpha_n \mathbf{x}_n \right\|^2 \le B\sum_{n=1}^{\infty} |\alpha_n|^2$ E $\exists A, B \in \mathbb{R}^+$ such that

Definition 2.15. Let $X \triangleq (X, +, \cdot, (\mathbb{F}, +, \dot{x}), \langle \triangle | \nabla \rangle)$ be an INNER PRODUCT SPACE (Definition C.9 page 124).

The sequences $(\mathbf{x}_n \in X)_{n \in \mathbb{Z}}$ and $(\mathbf{y}_n \in X)_{n \in \mathbb{Z}}$ are **biorthogonal** with respect to each other in **X** if

Lemma 2.3. ³² Let $\{x_n \mid n \in \mathbb{N}\}$ be a sequence in a Hilbert space $\mathbf{X} \triangleq (X, +, \cdot, (\mathbb{F}, +, \dot{\times}), \langle \triangle \mid \nabla \rangle)$. Let $\{ y_n | n \in \mathbb{N} \}$ be a sequence in a Hilbert space $Y \triangleq (Y, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle | \nabla \rangle)$. Let

(i). $\{x_n\}$ is total in X(ii). There exists A > 0 such that $A \sum_{n \in C} |a_n|^2 \le \left\| \sum_{n \in C} a_n \mathbf{x}_n \right\|^2$ for finite C and (iii). There exists B > 0 such that $\left\| \sum_{n=1}^{\infty} b_n y_n \right\|^2 \le B \sum_{n=1}^{\infty} \left| b_n \right|^2 \quad \forall (b_n)_{n \in \mathbb{N}} \in \mathscr{C}_{\mathbb{F}}^2$ (1). \mathbf{R}° is a linear bounded operator that maps from $\operatorname{span}\{x_n\}$ to $\operatorname{span}\{y_n\}$ where $\mathbf{R}^{\circ}\sum_{n\in C}c_nx_n\triangleq\sum_{n\in C}c_ny_n$, for some sequence (c_n) and finite set C(2). **R** has a unique extension to a bounded operator **R** that maps from **X** to **Y** and and

²⁸ ■ Bachman et al. (2002) page 112 ⟨3.4.8⟩, ■ Berberian (1961) page 53 ⟨Theorem II§8.3⟩

²⁹ Kubrusly (2001) page 357 (Proposition 5.43)

³¹ Christensen (2003) pages 66–68 ⟨page 68 and (3.24) on page 66⟩, Wojtaszczyk (1997) page 20 ⟨Definition 2.6⟩

³² Christensen (2003) pages 65–66 (Lemma 3.6.5)

³⁰ ✓ Young (2001) page 27 〈Definition 1.8.2〉, <a> Christensen (2003) page 63 〈Definition 3.6.1〉, <a> Heil (2011) page 196 (Definition 7.9)

Theorem 2.13. 33 Let $\{x_n \in X \mid n \in \mathbb{N}\}$ be a set of vectors in a SEPARABLE HILBERT SPACE $\mathcal{H} \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle \mid \nabla \rangle).$

$$\begin{array}{l}
\mathbf{H} = (X, +, \cdot, (\mathbb{F}, +, \times), (\triangle | \vee \rangle). \\
\mathbf{H} \\$$

№PROOF:

- 1. Proof for (\Longrightarrow) case:
 - (a) Proof that *Riesz basis* hypothesis \implies (1): all bases for H are *total* in H.
 - (b) Proof that *Riesz basis* hypothesis \implies (2):
 - i. Let $(u_n)_{n\in\mathbb{N}}$ be an *orthonormal basis* for H.
 - ii. Let **R** be a *bounded bijective* operator such that $x_n = \mathbf{R} u_n$.
 - iii. Proof for upper bound *B*:

$$\left\|\sum_{n=1}^{\infty} \alpha_{n} \mathbf{x}_{n}\right\|^{2} = \left\|\sum_{n=1}^{\infty} \alpha_{n} \mathbf{R} \mathbf{u}_{n}\right\|^{2} \quad \text{by definition of } \mathbf{R} \qquad \text{(item (1(b)ii))}$$

$$= \left\|\mathbf{R} \sum_{n=1}^{\infty} \alpha_{n} \mathbf{u}_{n}\right\|^{2} \quad \text{by Theorem C.1 page 113}$$

$$\leq \|\mathbf{R}\|^{2} \left\|\sum_{n=1}^{\infty} \alpha_{n} \mathbf{u}_{n}\right\|^{2} \quad \text{by Theorem C.6 page 119}$$

$$= \|\mathbf{R}\|^{2} \sum_{n=1}^{\infty} \|\alpha_{n} \mathbf{u}_{n}\|^{2} \quad \text{by } Pythagorean \ Theorem$$

$$= \|\mathbf{R}\|^{2} \sum_{n=1}^{\infty} |\alpha|^{2} \|\mathbf{u}_{n}\|^{2} \quad \text{by } homogeneous \ property of norms} \quad \text{(Definition C.5 page 116)}$$

$$= \|\mathbf{R}\|^{2} \sum_{n=1}^{\infty} |\alpha|^{2} \quad \text{by definition of } orthonormality \quad \text{(Definition 2.9 page 16)}$$

iv. Proof for lower bound *A*:

$$\left\|\sum_{n=1}^{\infty} \alpha_{n} \boldsymbol{x}_{n}\right\|^{2} = \frac{\left\|\mathbf{R}^{-1}\right\|^{2}}{\left\|\mathbf{R}^{-1}\right\|^{2}} \left\|\sum_{n=1}^{\infty} \alpha_{n} \boldsymbol{x}_{n}\right\|^{2} \quad \text{because } \left\|\mathbf{R}^{-1}\right\| > 0 \quad \text{(Proposition C.1 page 117)}$$

$$\geq \frac{1}{\left\|\mathbf{R}^{-1}\right\|^{2}} \left\|\mathbf{R}^{-1} \sum_{n=1}^{\infty} \alpha_{n} \boldsymbol{x}_{n}\right\|^{2} \quad \text{by Theorem C.6 page 119}$$

$$= \frac{1}{\left\|\mathbf{R}^{-1}\right\|^{2}} \left\|\mathbf{R}^{-1} \sum_{n=1}^{\infty} \alpha_{n} \mathbf{R} \boldsymbol{u}_{n}\right\|^{2} \quad \text{by definition of } \mathbf{R} \quad \text{(item (1(b)ii) page 28)}$$

$$= \frac{1}{\left\|\mathbf{R}^{-1}\right\|^{2}} \left\|\mathbf{R}^{-1} \mathbf{R} \sum_{n=1}^{\infty} \alpha_{n} \boldsymbol{u}_{n}\right\|^{2} \quad \text{by property of } linear operators \quad \text{(Theorem C.1 page 113)}$$



$$= \frac{1}{\|\|\mathbf{R}^{-1}\|\|^2} \left\| \sum_{n=1}^{\infty} \alpha_n \mathbf{u}_n \right\|^2 \qquad \text{by definition of inverse op.} \qquad \text{(Definition C.3 page 112)}$$

$$= \frac{1}{\|\|\mathbf{R}^{-1}\|\|^2} \sum_{n=1}^{\infty} \|\alpha_n \mathbf{u}_n\|^2 \qquad \text{by } Pythagorean \ Theorem}$$

$$= \frac{1}{\|\|\mathbf{R}^{-1}\|\|^2} \sum_{n=1}^{\infty} |\alpha_n|^2 \|\mathbf{u}_n\|^2 \qquad \text{by } \|\cdot\| \ homogeneous \ \text{prop.} \qquad \text{(Definition C.5 page 116)}$$

$$= \frac{1}{\|\|\mathbf{R}^{-1}\|\|^2} \sum_{n=1}^{\infty} |\alpha_n|^2 \qquad \text{by def. of } orthonormality \qquad \text{(Definition 2.9 page 16)}$$

2. Proof for (\Longrightarrow) case:

- (a) Let $\{u_n | n \in \mathbb{N}\}$ be an *orthonormal basis* for H.
- (b) Using (2) and Lemma 2.3 (page 27), construct an bounded extension operator **R** such that $\mathbf{R}\mathbf{u}_n = \mathbf{x}_n \text{ for all } n \in \mathbb{N}.$
- (c) Using (2) and Lemma 2.3 (page 27), construct an bounded extension operator S such that $\mathbf{S}\mathbf{x}_n = \mathbf{u}_n \text{ for all } n \in \mathbb{N}.$
- (d) Then, $\mathbf{R}\mathbf{V}\mathbf{x} = \mathbf{V}\mathbf{R}\mathbf{x} \implies \mathbf{V} = \mathbf{R}^{-1}$, and so **R** is a bounded invertible operator
- (e) and $\{x_n\}$ is a *Riesz sequence*.

Theorem 2.14. 34 Let $H \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle \mid \nabla \rangle)$ be a SEPARABLE HILBERT SPACE.

There exists $(y_n \in H)_{n \in \mathbb{Z}}$ such that

(1). (x_n) and (y_n) are BIORTHOGONAL and

(2). (y_n) is also a RIESZ BASIS for H and

(3). $\exists B > A > 0$ such that $\sum_{n=1}^{\infty} |a_n|^2 = |a_n|$ $A\sum_{n=1}^{\infty} |a_n|^2 \le \left\| \sum_{n=1}^{\infty} a_n x_n \right\|^2 = \left\| \sum_{n=1}^{\infty} a_n y_n \right\|^2 \le B\sum_{n=1}^{\infty} |a_n|^2$ $\forall (a_n)_{n\in\mathbb{N}}\in\mathscr{C}^2_{\mathbb{F}}$

^ℚProof:

1. Proof for (1):

- (a) Let e_n be the *unit vector* in H such that the *n*th element of e_n is 1 and all other elements are 0.
- (b) Let **M** be an operator on **H** such that $Me_n = x_n$.
- (c) Note that **M** is *isometric*, and as such $||\mathbf{M}\mathbf{x}|| = ||\mathbf{x}|| \quad \forall \mathbf{x} \in \mathbf{H}$.
- (d) Let $\mathbf{y}_n \triangleq (\mathbf{M}^{-1})^*$.
- (e) Then,

$$\langle \mathbf{y}_{n} | \mathbf{x}_{m} \rangle = \left\langle \left(\mathbf{M}^{-1} \right)^{*} e_{n} | \mathbf{M} e_{m} \right\rangle$$

$$= \left\langle e_{n} | \mathbf{M}^{-1} \mathbf{M} e_{m} \right\rangle$$

$$= \left\langle e_{n} | e_{m} \right\rangle$$

$$= \bar{\delta}_{nm}$$

$$\Longrightarrow \left\{ \mathbf{x}_{n} \right\} \text{ and } \left\{ \mathbf{y}_{n} \right\} \text{ are biorthogonal}$$

by Definition 2.9 page 16



³⁴ Wojtaszczyk (1997) page 20 (Lemma 2.7(a))

2. Proof for (3):

$$\left\| \sum_{n \in \mathbb{Z}} \alpha_{n} \mathbf{y}_{n} \right\| = \left\| \sum_{n \in \mathbb{Z}} \alpha_{n} (\mathbf{M}^{-1})^{*} e_{n} \right\| \qquad \text{by definition of } \mathbf{y}_{n} \qquad \text{(Proposition 1d page 29)}$$

$$= \left\| (\mathbf{M}^{-1})^{*} \sum_{n \in \mathbb{Z}} \alpha_{n} e_{n} \right\| \qquad \text{by property of } linear \ ops.$$

$$= \left\| \sum_{n \in \mathbb{Z}} \alpha_{n} e_{n} \right\| \qquad \text{because } (\mathbf{M}^{-1})^{*} \text{ is } isometric \qquad \text{(Definition C.13 page 132)}$$

$$= \left\| \mathbf{M} \sum_{n \in \mathbb{Z}} \alpha_{n} e_{n} \right\| \qquad \text{because } \mathbf{M} \text{ is } isometric \qquad \text{(Definition C.13 page 132)}$$

$$= \left\| \sum_{n \in \mathbb{Z}} \alpha_{n} \mathbf{M} e_{n} \right\| \qquad \text{by property of } linear \ operators$$

$$= \left\| \sum_{n \in \mathbb{Z}} \alpha_{n} \mathbf{x}_{n} \right\| \qquad \text{by definition of } \mathbf{M}$$

$$\implies \{ \mathbf{y}_{n} \} \text{ is a } Riesz \ basis \qquad \text{by left hypothesis}$$

3. Proof for (2): by (3) and definition of Riesz basis (Definition 2.13 page 27)

Proposition 2.5. ³⁵ Let $\{x_n \mid n \in \mathbb{N}\}$ be a set of vectors in a Hilbert space $\mathbf{H} \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle \mid \nabla \rangle)$.

$$\left\{ \begin{cases} \left\{ \mathbf{x}_{n} \right\} \text{ is } a \text{ Riesz basis for } \mathbf{H} \text{ with} \\ A \sum_{n=1}^{\infty} \left| a_{n} \right|^{2} \leq \left\| \sum_{n=1}^{\infty} a_{n} \mathbf{x}_{n} \right\|^{2} \leq B \sum_{n=1}^{\infty} \left| a_{n} \right|^{2} \right\} \implies \left\{ \underbrace{\begin{cases} \left\{ \mathbf{x}_{n} \right\} \text{ is } a \text{ frame for } \mathbf{H} \text{ with} \\ \frac{1}{B} \left\| \mathbf{x} \right\|^{2} \leq \sum_{n=1}^{\infty} \left| \left\langle \mathbf{x} \mid \mathbf{x}_{n} \right\rangle \right|^{2} \leq \frac{1}{A} \left\| \mathbf{x} \right\|^{2} \right\} \\ \forall \mathbf{x} \in \mathbf{H} \end{cases}$$

[♠]Proof:

- 1. Let $\{y_n | n \in \mathbb{N}\}$ be a *Riesz basis* that is *biorthogonal* to $\{x_n | n \in \mathbb{N}\}$ (Theorem 2.14 page 29).
- 2. Let $\mathbf{x} \triangleq \sum_{n=1}^{\infty} a_n \mathbf{y}_n$.
- 3. lemma:

$$\sum_{n=1}^{\infty} \left| \langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle \right|^2 = \sum_{n=1}^{\infty} \left| \left\langle \sum_{m=1}^{\infty} a_n \boldsymbol{y}_m \mid \boldsymbol{x}_n \right\rangle \right|^2 \quad \text{by definition of } \boldsymbol{x} \qquad \text{(item (2) page 30)}$$

$$= \sum_{n=1}^{\infty} \left| \sum_{m=1}^{\infty} a_n \left\langle \boldsymbol{y}_m \mid \boldsymbol{x}_n \right\rangle \right|^2 \quad \text{by } homogeneous \text{ property of } \langle \triangle \mid \nabla \rangle \quad \text{(Definition C.9 page 124)}$$

$$= \sum_{n=1}^{\infty} \left| \sum_{m=1}^{\infty} a_n \bar{\delta}_{mn} \right|^2 \quad \text{by definition of } biorthogonal \quad \text{(Definition 2.15 page 27)}$$

$$= \sum_{n=1}^{\infty} \left| a_n \right|^2 \quad \text{by definition of } \bar{\delta} \quad \text{(Definition 2.12 page 20)}$$

³⁵ **☐** Igari (1996) page 220 ⟨Lemma 9.8⟩, **☐** Wojtaszczyk (1997) pages 20–21 ⟨Lemma 2.7(a)⟩



$$A \sum_{n=1}^{\infty} |a_n|^2 \leq \left\| \sum_{n=1}^{\infty} a_n x_n \right\|^2 \leq B \sum_{n=1}^{\infty} |a_n|^2 \quad \text{by definition of } \{y_n\} \text{ (item (1) page 30)}$$

$$\Rightarrow A \sum_{n=1}^{\infty} |a_n|^2 \leq \left\| \sum_{n=1}^{\infty} a_n y_n \right\|^2 \leq B \sum_{n=1}^{\infty} |a_n|^2 \quad \text{by definition of } \{y_n\} \text{ (item (1) page 30)}$$

$$\Rightarrow A \sum_{n=1}^{\infty} |a_n|^2 \leq \|x\|^2 \leq B \sum_{n=1}^{\infty} |a_n|^2 \quad \text{by definition of } x \text{ (item (2) page 30)}$$

$$\Rightarrow A \sum_{n=1}^{\infty} |\langle x | x_n \rangle|^2 \leq \|x\|^2 \leq B \sum_{n=1}^{\infty} |\langle x | x_n \rangle|^2 \quad \text{by (3) lemma}$$

$$\Rightarrow \frac{1}{B} \|x\|^2 \leq \sum_{n=1}^{\infty} |\langle x | x_n \rangle|^2 \leq \frac{1}{A} \|x\|^2$$

Theorem 2.15 (Battle-Lemarié orthogonalization). 36 Let $\tilde{f}(\omega)$ be the Fourier Transform (Definition

1.2 page 196) of a function
$$f \in \mathcal{L}_{\mathbb{R}}^{2}$$
.

$$\begin{bmatrix}
1. & \left\{ \mathbf{T}^{n} \mathbf{g} \middle| n \in \mathbb{Z} \right\} \text{ is a Riesz basis for } \mathcal{L}_{\mathbb{R}}^{2} & \text{and} \\
2. & \tilde{f}(\omega) \triangleq \frac{\tilde{\mathbf{g}}(\omega)}{\sqrt{2\pi \sum_{n \in \mathbb{Z}} |\tilde{\mathbf{g}}(\omega + 2\pi n)|^{2}}}
\end{bmatrix} \implies \left\{ \begin{array}{c} \left\{ \mathbf{T}^{n} \mathbf{f} \middle| n \in \mathbb{Z} \right\} \\ \text{is an Orthonormal basis for } \mathcal{L}_{\mathbb{R}}^{2} \end{array} \right\}$$

^ℚProof:

1. Proof that $\{ \mathbf{T}^n \mathbf{f} | n \in \mathbb{Z} \}$ is orthonormal:

$$\tilde{S}_{\phi\phi}(\omega) = 2\pi \sum_{n \in \mathbb{Z}} \left| \tilde{f}(\omega + 2\pi n) \right|^{2}$$
 by Theorem P.1 page 255
$$= 2\pi \sum_{n \in \mathbb{Z}} \left| \frac{\tilde{g}(\omega + 2\pi n)}{\sqrt{2\pi \sum_{m \in \mathbb{Z}} |\tilde{g}(\omega + 2\pi (n - m))|^{2}}} \right|^{2}$$
 by left hypothesis
$$= \sum_{n \in \mathbb{Z}} \left| \frac{\tilde{g}(\omega + 2\pi n)}{\sqrt{\sum_{m \in \mathbb{Z}} |\tilde{g}(\omega + 2\pi m)|^{2}}} \right|^{2}$$

$$= \sum_{n \in \mathbb{Z}} \left| \frac{1}{\sqrt{\sum_{m \in \mathbb{Z}} |\tilde{g}(\omega + 2\pi m)|^{2}}} \right|^{2} |\tilde{g}(\omega + 2\pi n)|^{2}$$

$$= \frac{1}{\sum_{m \in \mathbb{Z}} |\tilde{g}(\omega + 2\pi m)|^{2}} \sum_{n \in \mathbb{Z}} |\tilde{g}(\omega + 2\pi n)|^{2}$$

$$= 1$$

$$\Rightarrow \{ f_{n} |_{n \in \mathbb{Z}} \} \text{ is orthonormal}$$
 by Theorem P.3 page 261

³⁶ Wojtaszczyk (1997) page 25 (Remark 2.4), 🏿 Vidakovic (1999) page 71, 🖨 Mallat (1989) page 72, 🗐 Mallat (1999) page 225, **■** Daubechies (1992) page 140 ((5.3.3))





2. Proof that $\{\mathbf{T}^n \mathbf{f} | n \in \mathbb{Z}\}$ is a basis for V_0 : by Lemma 2.1 page 15.

Frames in Hilbert spaces 2.7

Definition 2.16. ³⁷ Let $\{x_n \in X \mid n \in \mathbb{N}\}$ be a set of vectors in a HILBERT SPACE $H \triangleq (X, +, \cdot, (\mathbb{F}, +, \dot{\times}), \langle \triangle \mid \nabla \rangle).$ The set $\{x_n\}$ is a **frame** for H if (STABILITY CONDITION)

$$\exists A, B \in \mathbb{R}^+$$
 such that $A \|x\|^2 \le \sum_{n=1}^{\infty} |\langle x | x_n \rangle|^2 \le B \|x\|^2$ $\forall x \in X$.

The quantities A and B are frame bounds.

The quantity A' is the **optimal lower frame bound** if

 $A' = \sup \{ A \in \mathbb{R}^+ | A \text{ is a lower frame bound} \}.$

The quantity B' is the **optimal upper frame bound** if

 $B' = \inf \{ B \in \mathbb{R}^+ | B \text{ is an upper frame bound} \}.$

A frame is a **tight frame** if A = B.

A frame is a normalized tight frame (or a Parseval frame) if A = B = 1.

A frame $\{x_n | n \in \mathbb{N}\}$ is an **exact frame** if for some $m \in \mathbb{Z}$, $\{x_n | n \in \mathbb{N}\} \setminus \{x_m\}$ is not a frame.

A frame is a Parseval frame (Definition 2.16) if it satisfies Parseval's Identity (Theorem 2.9 page 22). All orthonormal bases are Parseval frames (Theorem 2.9 page 22); but not all Parseval frames are orthonormal bases.

Definition 2.17. Let $\{x_n\}$ be a **frame** (Definition 2.16 page 32) for the HILBERT SPACE $H \triangleq (X, +, \cdot, (\mathbb{F}, +, \dot{\times}), \langle \triangle | \nabla \rangle)$. Let **S** be an OPERATOR on **H**.

$$\begin{array}{c} \textbf{D} \\ \textbf{E} \\ \textbf{F} \end{array} \textbf{S} \ \textit{is a frame operator } for \left\{ \boldsymbol{x}_n \right\} \ \textit{if} \qquad \textbf{Sf}(\boldsymbol{x}) = \sum_{n \in \mathbb{Z}} \left\langle \mathbf{f} \mid \boldsymbol{x}_n \right\rangle \boldsymbol{x}_n \qquad \forall \mathbf{f} \in \mathcal{H}.$$

Theorem 2.16. ³⁸ Let S be a Frame Operator (Definition 2.17 page 32) of a Frame $\{x_n\}$ (Definition 2.16 page 32) for the Hilbert space $\mathbf{H} \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle \mid \nabla \rangle).$

(1). **S** *is* INVERTIBLE. and
(2).
$$f(x) = \sum_{n \in \mathbb{Z}} \langle f | S^{-1}x_n \rangle x_n = \sum_{n \in \mathbb{Z}} \langle f | x_n \rangle S^{-1}x_n \quad \forall f \in H$$

Theorem 2.17. ³⁹ Let $\{x_n \in X \mid n=1,2,...,N\}$ be a set of vectors in a Hilbert space $\mathbf{H} \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle \mid \nabla \rangle).$

$$\{x_n\}$$
 is a frame for $\{x_n\}$.

[♠]Proof:

³⁹ Christensen (2003) page 3



³⁷ ☑ Young (2001) pages 154–155, ② Christensen (2003) page 88 (Definitions 5.1.1, 5.1.2), ② Heil (2011) pages 204– 205 (Definition 8.2), Jørgensen et al. (2008) page 267 (Definition 12.22), Duffin and Schaeffer (1952) page 343, Daubechies et al. (1986) page 1272

 $^{^{38}}$ Christensen (2008) pages 100–102 (Theorem 5.1.7)

1. Upper bound: Proof that there exists B such that $\sum_{n=1}^{N} |\langle x | x_n \rangle|^2 \le B ||x||^2 \quad \forall x \in H$:

$$\sum_{n=1}^{N} \left| \langle \mathbf{x} \mid \mathbf{x}_{n} \rangle \right|^{2} \leq \sum_{n=1}^{N} \langle \mathbf{x}_{n} \mid \mathbf{x}_{n} \rangle \langle \mathbf{x} \mid \mathbf{x} \rangle$$
 by Cauchy-Schwarz inequality
$$= \underbrace{\left\{ \sum_{n=1}^{N} \left\| \mathbf{x}_{n} \right\|^{2} \right\}}_{B} \|\mathbf{x}\|^{2}$$


2. Lower bound: Proof that there exists *A* such that $A \|x\|^2 \le \sum_{n=1}^N |\langle x | x_n \rangle|^2 \quad \forall x \in H$:


$$\sum_{n=1}^{N} \left| \langle \mathbf{x} \mid \mathbf{x}_n \rangle \right|^2 = \sum_{n=1}^{N} \left| \left\langle \mathbf{x}_n \mid \frac{\mathbf{x}}{\|\mathbf{x}\|} \right\rangle \right|^2 \|\mathbf{x}\|^2$$

$$\geq \underbrace{\left(\inf_{\mathbf{y}} \left\{ \sum_{n=1}^{N} \left| \left\langle \mathbf{x}_n \mid \mathbf{y} \right\rangle \right|^2 | \|\mathbf{y}\| = 1 \right\} \right)}_{A} \|\mathbf{x}\|^2$$

Example 2.1. Let $(\mathbb{R}^2, +, \cdot, (\mathbb{R}, \dot{+}, \dot{\times}), \langle \triangle | \nabla \rangle)$ be an inner product space with $\left(\begin{bmatrix} x_1 \\ y_1 \end{bmatrix} | \begin{bmatrix} x_2 \\ y_2 \end{bmatrix}\right) \triangleq x_1x_2 + y_1y_2$. Let **S** be the *frame operator* (Definition 2.17 page 32) with *inverse* \mathbf{S}^{-1} .

Let $N \in \{3, 4, 5, ...\}$, $\theta \in \mathbb{R}$, and $r \in \mathbb{R}^+$ (r > 0). Let $\mathbf{x}_n \triangleq r \begin{bmatrix} \cos(\theta + 2n\pi/N) \\ \sin(\theta + 2n\pi/N) \end{bmatrix} \quad \forall_{n \in \{0, 1, ..., N-1\}}$.





Moreover,
$$\mathbf{S} = \frac{Nr^2}{2} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$
, $\mathbf{S}^{-1} = \frac{2}{Nr^2} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$, and $\mathbf{v} = \frac{2}{Nr^2} \sum_{n=0}^{N-1} \langle \mathbf{v} \mid \mathbf{x}_n \rangle \mathbf{x}_n \quad \forall \mathbf{v} \in \mathbb{R}^2$.

№PROOF:

1. Proof that $(x_0, x_1, \dots, x_{N-1})$ is a *tight frame* with *frame bound* $A = \frac{Nr^2}{2}$: Let $\mathbf{v} \triangleq (x, y) \in \mathbb{R}^2$.

$$\sum_{n=0}^{N-1} \left| \langle \boldsymbol{v} \mid \boldsymbol{x}_{n} \rangle \right|^{2} \triangleq \sum_{n=0}^{N-1} \left| \boldsymbol{v}^{\mathbf{H}} \boldsymbol{r} \begin{bmatrix} \cos \left(\theta + \frac{2n\pi}{N} \right) \\ \sin \left(\theta + \frac{2n\pi}{N} \right) \end{bmatrix} \right|^{2}$$
 by definitions of \boldsymbol{v} of $\langle \boldsymbol{y} \mid \boldsymbol{x} \rangle$

$$\triangleq \sum_{n=0}^{N-1} r^{2} \left| \operatorname{xcos} \left(\theta + \frac{2n\pi}{N} \right) + \operatorname{ysin} \left(\theta + \frac{2n\pi}{N} \right) \right|^{2}$$
 by definition of $\boldsymbol{y}^{\mathbf{H}} \boldsymbol{x}$ operation
$$= r^{2} x^{2} \sum_{n=0}^{N-1} \cos^{2} \left(\theta + \frac{2n\pi}{N} \right) + r^{2} y^{2} \sum_{n=0}^{N-1} \sin^{2} \left(\theta + \frac{2n\pi}{N} \right) + r^{2} x y \sum_{n=0}^{N-1} \cos \left(\theta + \frac{2n\pi}{N} \right) \sin \left(\theta + \frac{2n\pi}{N} \right)$$

$$= r^{2} x^{2} \frac{N}{2} + r^{2} y^{2} \frac{N}{2} + r^{2} x y 0$$
 by Corollary H.1 page 189
$$= \left(x^{2} + y^{2} \right) \frac{Nr^{2}}{2} = \left(\frac{Nr^{2}}{2} \right) \boldsymbol{v}^{\mathbf{H}} \boldsymbol{v} \triangleq \left(\frac{Nr^{2}}{2} \right) \|\boldsymbol{v}\|^{2}$$
 by definition of $\|\boldsymbol{v}\|$

2. Proof that $\mathbf{S} = \frac{Nr^2}{2} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$:



(a) Let
$$e_1 \triangleq \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$
 and $e_2 \triangleq \begin{bmatrix} 0 \\ 1 \end{bmatrix}$.

(b) lemma: $\mathbf{S}\mathbf{e}_1 = \frac{Nr^2}{2}\mathbf{e}_1$. Proof:

$$\begin{split} \mathbf{S}e_1 &= \sum_{n=0}^{N-1} \left\langle e_1 \mid \mathbf{x}_n \right\rangle \mathbf{x}_n \\ &= \sum_{n=0}^{N-1} r \mathrm{cos} \Big(\theta + \frac{2n\pi}{N} \Big) r \begin{bmatrix} \mathrm{cos} \Big(\theta + \frac{2n\pi}{N} \Big) \\ \mathrm{sin} \Big(\theta + \frac{2n\pi}{N} \Big) \end{bmatrix} = r^2 \sum_{n=0}^{N-1} \begin{bmatrix} \mathrm{cos}^2 \Big(\theta + \frac{2n\pi}{N} \Big) \\ \mathrm{cos} \Big(\theta + \frac{2n\pi}{N} \Big) \mathrm{sin} \Big(\theta + \frac{2n\pi}{N} \Big) \end{bmatrix} \\ &= r^2 \begin{bmatrix} N/2 \\ 0 \end{bmatrix} = \frac{Nr^2}{2} e_1 \quad \text{by Summation around unit circle (Corollary H.1 page 189)} \end{split}$$

(c) lemma: $\mathbf{S}\mathbf{e}_2 = \frac{Nr^2}{2}\mathbf{e}_2$. Proof:

$$\begin{split} \mathbf{S}\boldsymbol{e}_2 &= \sum_{n=0}^{N-1} \left\langle \boldsymbol{e}_2 \mid \boldsymbol{x}_n \right\rangle \boldsymbol{x}_n \\ &= \sum_{n=0}^{N-1} r \mathrm{sin} \Big(\theta + \frac{2n\pi}{N} \Big) r \Bigg[\begin{array}{c} \cos \Big(\theta + \frac{2n\pi}{N} \Big) \\ \sin \Big(\theta + \frac{2n\pi}{N} \Big) \end{array} \Bigg] = r^2 \sum_{n=0}^{N-1} \Bigg[\begin{array}{c} \sin \Big(\theta + \frac{2n\pi}{N} \Big) \cos \Big(\theta + \frac{2n\pi}{N} \Big) \\ \sin^2 \Big(\theta + \frac{2n\pi}{N} \Big) \end{array} \Bigg] \\ &= r^2 \Bigg[\begin{array}{c} 0 \\ N/2 \end{array} \Bigg] = \frac{Nr^2}{2} \boldsymbol{e}_2 \qquad \text{by Summation around unit circle (Corollary H.1 page 189)} \end{split}$$

(d) Complete the proof of item (2) using *Eigendecomposition* $S = QAQ^{-1}$:

$$\mathbf{S}\mathbf{e}_1 = \frac{Nr^2}{2}\mathbf{e}_1$$
 by (2c) lemma

 \implies e_1 is an eigenvector of **S** with eigenvalue $\frac{Nr^2}{2}$

$$\mathbf{S}\mathbf{e}_2 = \frac{Nr^2}{2}\mathbf{e}_2$$
 by (2c) lemma

 \implies e_2 is an eigenvector of **S** with eigenvalue $\frac{Nr^2}{2}$

Eigendecomposition of S

$$\mathbf{S} = \underbrace{\begin{bmatrix} \mid & \mid \\ e_1 & e_2 \\ \mid & \mid \end{bmatrix}}_{\mathbf{Q}} \underbrace{\begin{bmatrix} \frac{Nr^2}{2} & 0 \\ 0 & \frac{Nr^2}{2} \end{bmatrix}}_{\mathbf{A}} \underbrace{\begin{bmatrix} \mid & \mid \\ e_1 & e_2 \\ \mid & \mid \end{bmatrix}}_{\mathbf{Q}^{-1}}^{-1} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} \frac{Nr^2}{2} & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & \frac{Nr^2}{2} \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}^{-1} = \frac{Nr^2}{2} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

3. Proof that $S^{-1} = \frac{2}{Nr^2} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$:

$$\mathbf{S}\mathbf{S}^{-1} = \frac{Nr^2}{2} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \frac{2}{Nr^2} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \triangleq \mathbf{I}_2$$
 by item (2)

$$\mathbf{S}^{-1}\mathbf{S} = \frac{2}{Nr^2} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \frac{Nr^2}{2} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \triangleq \mathbf{I}_2$$
 by item (2)

4. Proof that $\mathbf{v} = \frac{2}{Nr^2} \sum_{n=0}^{N-1} (\mathbf{v}^{\mathbf{H}} \mathbf{x}_n) \mathbf{x}_n$:

$$\boldsymbol{v} = \sum_{n=0}^{N-1} \left\langle \boldsymbol{v} \mid \mathbf{S}^{-1} \boldsymbol{x}_n \right\rangle \boldsymbol{x}_n = \sum_{n=0}^{N-1} \left\langle \boldsymbol{v} \mid \frac{2}{Nr^2} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \boldsymbol{x}_n \right\rangle \boldsymbol{x}_n$$
 by item (3)

$$= \frac{2}{Nr^2} \sum_{n=0}^{N-1} \langle \boldsymbol{v} | \boldsymbol{x}_n \rangle \boldsymbol{x}_n = \frac{2}{Nr^2} \sum_{n=0}^{N-1} (\boldsymbol{v}^{\mathbf{H}} \boldsymbol{x}_n) \boldsymbol{x}_n$$
 by definition of $\langle \boldsymbol{y} | \boldsymbol{x} \rangle$




Example 2.2 (Peace Frame/Mercedes Frame). ⁴⁰ Let $(\mathbb{R}^2, +, \cdot, (\mathbb{R}, \dot{+}, \dot{\times}), \langle \triangle | \nabla \rangle)$ be an inner product space with $\langle (x_1, y_1) | (x_2, y_2) \rangle \triangleq x_1 y_1 + x_2 y_2$. Let **S** be the *frame operator* (Definition 2.17 page 32) with inverse S^{-1} .

Let
$$\mathbf{x}_1 \triangleq \frac{1}{2} \begin{bmatrix} 0 \\ 2 \end{bmatrix}$$
, $\mathbf{x}_2 \triangleq \frac{1}{2} \begin{bmatrix} -\sqrt{3} \\ -1 \end{bmatrix}$, and $\mathbf{x}_3 \triangleq \frac{1}{2} \begin{bmatrix} \sqrt{3} \\ -1 \end{bmatrix}$.

Then, (x_1, x_2, x_3) is a **tight frame** for \mathbb{R}^2 with

frame bound
$$A = \frac{3}{2}$$
.
Moreover, $\mathbf{S} = \frac{3}{2} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$, $\mathbf{S}^{-1} = \frac{2}{3} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$,

and
$$\mathbf{v} = \frac{2}{3} \sum_{n=1}^{3} \langle \mathbf{v} | \mathbf{x}_n \rangle \mathbf{x}_n \triangleq \frac{2}{3} \sum_{n=1}^{3} (\mathbf{v}^{\mathbf{H}} \mathbf{x}_n) \mathbf{x}_n \quad \forall \mathbf{v} \in \mathbb{R}^2.$$



NPROOF:

- 1. This frame is simply a special case of the frame presented in Example 2.1 (page 33) with r = 1, N = 3, and $\theta = \pi h$.
- 2. Let's give it a try! Let $\mathbf{v} \triangleq \begin{bmatrix} 1 \\ 1 \end{bmatrix}$.

$$\sum_{n=1}^{3} \left\langle v \mid \mathbf{S}^{-1} \mathbf{x}_{n} \right\rangle \mathbf{x}_{n} = \frac{2}{3} \sum_{n=1}^{3} \left(v^{\mathbf{H}} \mathbf{x}_{n} \right) \mathbf{x}_{n}$$
 by Example 2.1 page 33
$$= \left(v^{\mathbf{H}} \mathbf{x}_{1} \right) \mathbf{x}_{1} + \left(v^{\mathbf{H}} \mathbf{x}_{2} \right) \mathbf{x}_{2} + \left(v^{\mathbf{H}} \mathbf{x}_{3} \right) \mathbf{x}_{3}$$

$$= \frac{2}{3} \left(\left(v^{\mathbf{H}} \frac{1}{2} \begin{bmatrix} 0 \\ 2 \end{bmatrix} \right) \mathbf{x}_{1} + \left(v^{\mathbf{H}} \frac{1}{2} \begin{bmatrix} -\sqrt{3} \\ -1 \end{bmatrix} \right) \mathbf{x}_{2} + \left(v^{\mathbf{H}} \frac{1}{2} \begin{bmatrix} \sqrt{3} \\ -1 \end{bmatrix} \right) \mathbf{x}_{3} \right)$$

$$= \frac{2}{3} \cdot \frac{1}{2} \left(\left(v^{\mathbf{H}} \begin{bmatrix} 0 \\ 2 \end{bmatrix} \right) \mathbf{x}_{1} + \left(v^{\mathbf{H}} \begin{bmatrix} -\sqrt{3} \\ -1 \end{bmatrix} \right) \mathbf{x}_{2} + \left(v^{\mathbf{H}} \begin{bmatrix} \sqrt{3} \\ -1 \end{bmatrix} \right) \mathbf{x}_{3} \right)$$

$$= \frac{1}{3} \left((2) \frac{1}{2} \begin{bmatrix} 0 \\ 2 \end{bmatrix} + \left(-\sqrt{3} - 1 \right) \frac{1}{2} \begin{bmatrix} -\sqrt{3} \\ -1 \end{bmatrix} + \left(\sqrt{3} - 1 \right) \frac{1}{2} \begin{bmatrix} \sqrt{3} \\ -1 \end{bmatrix} \right)$$

$$= \frac{1}{6} \begin{bmatrix} 2(0) + (-\sqrt{3} - 1)(-\sqrt{3}) + (\sqrt{3} - 1)(\sqrt{3}) \\ 2(2) + (-\sqrt{3} - 1)(-1) + (\sqrt{3} - 1)(-1) \end{bmatrix}$$

$$= \frac{1}{6} \begin{bmatrix} 0 + (3 + \sqrt{3}) + (3 - \sqrt{3}) \\ 4 + (1 + \sqrt{3}) + (1 - \sqrt{3}) \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \end{bmatrix} \triangleq \mathbf{v}$$

In Example 2.1 (page 33) and Example 2.2 (page 35), the frame operator S and its inverse S^{-1} were computed. In general however, it is not always necessary or even possible to compute these, as illustrated in Example 2.3 (next).


Example 2.3. ⁴¹ Let $(\mathbb{R}^2, +, \cdot, (\mathbb{R}, \dotplus, \dot{x}), \langle \triangle | \nabla \rangle)$ be an inner product space with $\langle (x_1, y_1) | (x_2, y_2) \rangle \triangleq x_1 y_1 + x_2 y_2$. Let **S** be the *frame operator* (Definition 2.17 page 32) with *inverse* \mathbf{S}^{-1} .

⁴⁰ ← Heil (2011) pages 204–205 ⟨r = 1 case⟩, ← Byrne (2005) page 80 ⟨r = 1 case⟩, ← Han et al. (2007) page 91 $\langle \text{Example 3.9}, r = \sqrt{2/3 \text{ case}} \rangle$

☐ Christensen (2003) pages 7–8 ⟨?⟩

Let p and q be *orthonormal* vectors in $X \triangleq \text{span}\{p, q\}$.

Let $x_1 \triangleq p$, $x_2 \triangleq p + q$, and $x_3 \triangleq p - q$. Then, $\{x_1, x_2, x_3\}$ is a **frame** for X with *frame bounds* A = 0 and B = 5.



Moreover,

$$\mathbf{S}^{-1} \mathbf{x}_1 = \frac{1}{3} \mathbf{p}$$
 and $\mathbf{S}^{-1} \mathbf{x}_2 = \frac{1}{3} \mathbf{p} + \frac{1}{2} \mathbf{q}$ and $\mathbf{S}^{-1} \mathbf{x}_3 = \frac{1}{3} \mathbf{p} - \frac{1}{2} \mathbf{q}$.

№ Proof:

1. Proof that (x_1, x_2, x_3) is a *frame* with *frame bounds* A = 0 and B = 5:

$$\sum_{n=1}^{3} \left| \langle \boldsymbol{v} \, | \, \boldsymbol{x}_{n} \rangle \right|^{2} \triangleq \left| \langle \boldsymbol{v} \, | \, \boldsymbol{p} \rangle \right|^{2} + \left| \langle \boldsymbol{v} \, | \, \boldsymbol{p} + \boldsymbol{q} \rangle \right|^{2} + \left| \langle \boldsymbol{v} \, | \, \boldsymbol{p} - \boldsymbol{q} \rangle \right|^{2}$$
 by definitions of \boldsymbol{x}_{1} , \boldsymbol{x}_{2} , and \boldsymbol{x}_{3}

$$= \left| \langle \boldsymbol{v} \, | \, \boldsymbol{p} \rangle \right|^{2} + \left| \langle \boldsymbol{v} \, | \, \boldsymbol{p} \rangle + \langle \boldsymbol{v} \, | \, \boldsymbol{q} \rangle \right|^{2} + \left| \langle \boldsymbol{v} \, | \, \boldsymbol{p} \rangle - \langle \boldsymbol{v} \, | \, \boldsymbol{q} \rangle \right|^{2}$$
 by $\boldsymbol{additivity}$ of $\langle \triangle \, | \, \nabla \rangle$ (Definition C.9 page 124)
$$= \left| \langle \boldsymbol{v} \, | \, \boldsymbol{p} \rangle \right|^{2} + \left| \langle \boldsymbol{v} \, | \, \boldsymbol{p} \rangle \right|^{2} + \left| \langle \boldsymbol{v} \, | \, \boldsymbol{q} \rangle \right|^{2} + \langle \boldsymbol{v} \, | \, \boldsymbol{p} \rangle \langle \boldsymbol{v} \, | \, \boldsymbol{q} \rangle^{*} + \langle \boldsymbol{v} \, | \, \boldsymbol{q} \rangle \langle \boldsymbol{v} \, | \, \boldsymbol{p} \rangle^{*})$$

$$+ \left(\left| \langle \boldsymbol{v} \, | \, \boldsymbol{p} \rangle \right|^{2} + \left| \langle \boldsymbol{v} \, | \, \boldsymbol{q} \rangle \right|^{2} - \langle \boldsymbol{v} \, | \, \boldsymbol{p} \rangle \langle \boldsymbol{v} \, | \, \boldsymbol{q} \rangle^{*} - \langle \boldsymbol{v} \, | \, \boldsymbol{q} \rangle \langle \boldsymbol{v} \, | \, \boldsymbol{p} \rangle^{*} \right)$$

$$= 3 \left| \langle \boldsymbol{v} \, | \, \boldsymbol{p} \rangle \right|^{2} + 2 \left| \langle \boldsymbol{v} \, | \, \boldsymbol{q} \rangle \right|^{2}$$

$$\leq 3 \left\| \boldsymbol{v} \, \| \, \| \boldsymbol{p} \| + 2 \left\| \boldsymbol{v} \, \| \, \| \boldsymbol{q} \right\|$$
 by \boldsymbol{CS} $\boldsymbol{Inequality}$

$$= \left\| \boldsymbol{v} \, \| \, (3 \left\| \boldsymbol{p} \right\| + 2 \left\| \boldsymbol{q} \right\| \right)$$

$$= 5 \left\| \boldsymbol{v} \, \|$$
 by $\boldsymbol{orthonormality}$ of \boldsymbol{p} and \boldsymbol{q}

2. lemma: $\mathbf{S}p = 3p$, $\mathbf{S}q = 2q$, $\mathbf{S}^{-1}p = \frac{1}{3}p$, and $\mathbf{S}^{-1}q = \frac{1}{2}q$. Proof:

$$\mathbf{S}\boldsymbol{p} \triangleq \sum_{n=1}^{3} \langle \boldsymbol{p} \,|\, \boldsymbol{x}_{n} \rangle \,\boldsymbol{x}_{n}$$

$$= \langle \boldsymbol{p} \,|\, \boldsymbol{p} \rangle \,\boldsymbol{p} + \langle \boldsymbol{p} \,|\, \boldsymbol{p} + \boldsymbol{q} \rangle \,(\boldsymbol{p} + \boldsymbol{q}) + \langle \boldsymbol{p} \,|\, \boldsymbol{p} - \boldsymbol{q} \rangle \,(\boldsymbol{p} - \boldsymbol{q})$$

$$= (1)\boldsymbol{p} + (1 + 0)(\boldsymbol{p} + \boldsymbol{q}) + (1 - 0)(\boldsymbol{p} - \boldsymbol{q})$$

$$= 3\boldsymbol{p}$$

$$\Rightarrow \mathbf{S}^{-1}\boldsymbol{p} = \frac{1}{3}\boldsymbol{p}$$

$$\mathbf{S}\boldsymbol{q} \triangleq \sum_{n=1}^{3} \langle \boldsymbol{q} \,|\, \boldsymbol{x}_{n} \rangle \,\boldsymbol{x}_{n}$$

$$= \langle \boldsymbol{q} \,|\, \boldsymbol{p} \rangle \,\boldsymbol{p} + \langle \boldsymbol{q} \,|\, \boldsymbol{p} + \boldsymbol{q} \rangle \,(\boldsymbol{p} + \boldsymbol{q}) + \langle \boldsymbol{q} \,|\, \boldsymbol{p} - \boldsymbol{q} \rangle \,(\boldsymbol{p} - \boldsymbol{q})$$

$$= (0)\boldsymbol{q} + (0 + 1)(\boldsymbol{p} + \boldsymbol{q}) + (0 - 1)(\boldsymbol{p} - \boldsymbol{q})$$

$$= 2\boldsymbol{q}$$

$$\Rightarrow \mathbf{S}^{-1}\boldsymbol{q} = \frac{1}{2}\boldsymbol{q}$$

- 3. Remark: Without knowing p and q, from (2) lemma it follows that it is not possible to compute S or S^{-1} explicitly.
- 4. Proof that $S^{-1}x_1 = \frac{1}{2}p$, $S^{-1}x_2 = \frac{1}{2}p + \frac{1}{2}q$ and $S^{-1}x_3 = \frac{1}{2}p \frac{1}{2}q$:

$$\mathbf{S}^{-1}\mathbf{x}_{1} \triangleq \mathbf{S}^{-1}\mathbf{p}$$
 by definition of \mathbf{x}_{1}

$$= \frac{1}{3}\mathbf{p}$$
 by (2) lemma

$$\mathbf{S}^{-1}\mathbf{x}_{2} \triangleq \mathbf{S}^{-1}(\mathbf{p} + \mathbf{q})$$
 by definition of \mathbf{x}_{2}

$$= \frac{1}{3}\mathbf{p} + \frac{1}{2}\mathbf{q}$$
 by (2) lemma

$$\mathbf{S}^{-1} \mathbf{x}_3 \triangleq \mathbf{S}^{-1} (\mathbf{p} - \mathbf{q})$$
$$= \frac{1}{3} \mathbf{p} - \frac{1}{2} \mathbf{q}$$

by definition of x_2

by (2) lemma

5. Check that $v = \sum_{n} \langle v \mid x_n \rangle x_n = \langle v \mid p \rangle p + \langle v \mid q \rangle q$:

$$v = \sum_{n=1}^{3} \left\langle v \mid \mathbf{S}^{-1} \mathbf{x}_{n} \right\rangle \mathbf{x}_{n}$$

$$= \left\langle v \mid \mathbf{S}^{-1} \mathbf{p} \right\rangle \mathbf{p} + \left\langle v \mid \mathbf{S}^{-1} (\mathbf{p} + \mathbf{q}) \right\rangle (\mathbf{p} + \mathbf{q}) + \left\langle v \mid \mathbf{S}^{-1} (\mathbf{p} - \mathbf{q}) \right\rangle (\mathbf{p} - \mathbf{q})$$

$$= \left\langle v \mid \frac{1}{3} \mathbf{p} \right\rangle \mathbf{p} + \left\langle v \mid \frac{1}{3} \mathbf{p} + \frac{1}{2} \mathbf{q} \right\rangle (\mathbf{p} + \mathbf{q}) + \left\langle v \mid \frac{1}{3} \mathbf{p} - \frac{1}{2} \mathbf{q} \right\rangle (\mathbf{p} - \mathbf{q})$$

$$= \left(\frac{1}{3} + \frac{1}{3} + \frac{1}{3} \right) \left\langle v \mid \mathbf{p} \right\rangle \mathbf{p} + \left(\frac{1}{3} - \frac{1}{3} \right) \left\langle v \mid \mathbf{p} \right\rangle \mathbf{q} + \left(\frac{1}{2} - \frac{1}{2} \right) \left\langle v \mid \mathbf{q} \right\rangle \mathbf{p} + \left(\frac{1}{2} + \frac{1}{2} \right) \left\langle v \mid \mathbf{q} \right\rangle \mathbf{q}$$

$$= \left\langle v \mid \mathbf{p} \right\rangle \mathbf{p} + \left\langle v \mid \mathbf{q} \right\rangle \mathbf{q}$$

 \blacksquare

TRANSVERSAL OPERATORS

"Je me plaisois surtout aux mathématiques, à cause de la certitude et de l'évidence de leurs raisons: mais je ne remarquois point encore leur vrai usage; et, pensant qu'elles ne servoient qu'aux arts mécaniques, je m'étonnois de ce que leurs fondements étant si fermes et si solides, on n'avoit rien bâti dessus de plus relevé:



"I was especially delighted with the mathematics, on account of the certitude and evidence of their reasonings; but I had not as yet a precise knowledge of their true use; and thinking that they but contributed to the advancement of the mechanical arts, I was astonished that foundations, so strong and solid, should have had no loftier superstructure reared on them."

René Descartes, philosopher and mathematician (1596–1650)

3.1 Families of Functions

This text is largely set in the space of $Lebesgue\ square-integrable\ functions\ L^2_{\mathbb{R}}$ (Definition D.1 page 141). The space $L^2_{\mathbb{R}}$ is a subspace of the space $\mathbb{R}^{\mathbb{R}}$, the set of all functions with $domain\ \mathbb{R}$ (the set of real numbers) and $range\ \mathbb{R}$. The space $\mathbb{R}^{\mathbb{R}}$ is a subspace of the space $\mathbb{C}^{\mathbb{C}}$, the set of all functions with $domain\ \mathbb{C}$ (the set of complex numbers) and $range\ \mathbb{C}$. That is, $L^2_{\mathbb{R}}\subseteq\mathbb{R}^{\mathbb{R}}\subseteq\mathbb{C}^{\mathbb{C}}$. In general, the notation Y^X represents the set of all functions with domain X and range Y (Definition 3.1 page 39). Although this notation may seem curious, note that for finite X and finite Y, the number of functions (elements) in Y^X is $|Y^X| = |Y|^{|X|}$.

Definition 3.1. *Let X and Y be sets.*

The space Y^X represents the set of all functions with DOMAIN X and RANGE Y such that $Y^X \triangleq \{f(x)|f(x): X \to Y\}$

translation: Descartes (1637c) (part I, paragraph 10)

nage: http://en.wikipedia.org/wiki/File:Frans_Hals_-_Portret_van_Ren%C3%A9_Descartes.jpg, public domain

¹ quote: Descartes (1637b)

Definition 3.2. 2 Let X be a set.


D E

The indicator function
$$\mathbb{1} \in \{0,1\}^{2^X}$$
 is defined as
$$\mathbb{1}_A(x) = \begin{cases} 1 & \text{for } x \in A & \forall x \in X, A \in 2^X \\ 0 & \text{for } x \notin A & \forall x \in X, A \in 2^X \end{cases}$$
 The indicator function $\mathbb{1}$ is also called the **characteristic function**.


3.2 **Definitions and algebraic properties**

Much of the wavelet theory developed in this text is constructed using the translation operator T and the **dilation operator D** (next).

Definition 3.3. ³



Example 3.1. Let **T** and **D** be defined as in Definition 3.3 (page 40).



Proposition 3.1. Let T_{τ} be a TRANSLATION OPERATOR (Definition 3.3 page 40).

$$\begin{array}{c|c}
\mathbf{P} \\
\mathbf{R} \\
\mathbf{P}
\end{array}
\qquad
\begin{array}{c}
\mathbf{T}_{\tau}^{n} \mathbf{f}(x) = \sum_{n \in \mathbb{Z}} \mathbf{T}_{\tau}^{n} \mathbf{f}(x + \tau) & \forall \mathbf{f} \in \mathbb{R}^{\mathbb{R}}
\end{array}
\qquad
\left(\sum_{n \in \mathbb{Z}} \mathbf{T}_{\tau}^{n} \mathbf{f}(x) \text{ is Periodic with period } \tau\right)$$

^ℚProof:

$$\sum_{n\in\mathbb{Z}}\mathbf{T}_{\tau}^{n}\mathsf{f}(x+\tau) = \sum_{n\in\mathbb{Z}}\mathsf{f}(x-n\tau+\tau) \qquad \text{by definition of } \mathbf{T}_{\tau} \qquad \text{(Definition 3.3 page 40)}$$

$$= \sum_{m\in\mathbb{Z}}\mathsf{f}(x-m\tau) \qquad \text{where } m\triangleq n-1 \qquad \Longrightarrow n=m+1$$

$$= \sum_{m\in\mathbb{Z}}\mathbf{T}_{\tau}^{m}\mathsf{f}(x) \qquad \text{by definition of } \mathbf{T}_{\tau} \qquad \text{(Definition 3.3 page 40)}$$

In a linear space, every operator has an *inverse*. Although the inverse always exists as a *relation*, it may not exist as a function or as an operator. But in some cases the inverse of an operator is itself an operator. The inverses of the operators **T** and **D** both exist as operators, as demonstrated next.

^{18 (}Definitions 2.3,2.4), Mammler (2008) page A-21, Machman et al. (2002) page 473, Macker (2004) page 260, Zayed (2004) page ,

☐ Heil (2011) page 250 (Notation 9.4),
☐ Casazza and Lammers (1998) page 74,
☐ Goodman et al. (1993a) page 639, Heil and Walnut (1989) page 633 (Definition 1.3.1), Dai and Lu (1996) page 81, Dai and Larson (1998) page 2



⁴⁴⁰

Proposition 3.2 (transversal operator inverses). *Let* **T** *and* **D** *be as defined in Definition 3.3 page 40.*

T has an inverse \mathbf{T}^{-1} in $\mathbb{C}^{\mathbb{C}}$ expressed by the relation $\mathbf{T}^{-1}\mathbf{f}(x) = \mathbf{f}(x+1) \quad \forall \mathbf{f} \in \mathbb{C}^{\mathbb{C}}$ (translation operator inverse). **D** has an inverse \mathbf{D}^{-1} in $\mathbb{C}^{\mathbb{C}}$ expressed by the relation $\mathbf{D}^{-1}\mathsf{f}(x) = \frac{\sqrt{2}}{2}\,\mathsf{f}\left(\frac{1}{2}x\right) \quad \forall \mathsf{f} \in \mathbb{C}^{\mathbb{C}}$ (dilation operator inverse).

[♠]Proof:

Р

1. Proof that T^{-1} is the inverse of T:

$$\mathbf{T}^{-1}\mathbf{T}f(x) = \mathbf{T}^{-1}f(x-1) \qquad \text{by defintion of } \mathbf{T}$$

$$= f([x+1]-1)$$

$$= f(x)$$

$$= f([x-1]+1)$$

$$= \mathbf{T}f(x+1) \qquad \text{by defintion of } \mathbf{T}$$

$$= \mathbf{T}\mathbf{T}^{-1}f(x)$$

$$\Rightarrow \mathbf{T}^{-1}\mathbf{T} = \mathbf{I} = \mathbf{T}\mathbf{T}^{-1}$$

2. Proof that \mathbf{D}^{-1} is the inverse of \mathbf{D} :

$$\mathbf{D}^{-1}\mathbf{D}\mathbf{f}(x) = \mathbf{D}^{-1}\sqrt{2}\mathbf{f}(2x) \qquad \text{by defintion of } \mathbf{D}$$

$$= \left(\frac{\sqrt{2}}{2}\right)\sqrt{2}\mathbf{f}\left(2\left[\frac{1}{2}x\right]\right)$$

$$= \mathbf{f}(x)$$

$$= \sqrt{2}\left[\frac{\sqrt{2}}{2}\mathbf{f}\left(\frac{1}{2}[2x]\right)\right]$$

$$= \mathbf{D}\left[\frac{\sqrt{2}}{2}\mathbf{f}\left(\frac{1}{2}x\right)\right] \qquad \text{by defintion of } \mathbf{D}$$

$$= \mathbf{D}\mathbf{D}^{-1}\mathbf{f}(x)$$

$$\Rightarrow \mathbf{D}^{-1}\mathbf{D} = \mathbf{I} = \mathbf{D}\mathbf{D}^{-1}$$

Proposition 3.3. Let T and D be as defined in Definition 3.3 page 40.

Let $\mathbf{D}^0 = \mathbf{T}^0 \triangleq \mathbf{I}$ be the identity operator.

$$\mathbf{p} \quad \mathbf{D}^{j} \mathbf{T}^{n} \mathbf{f}(x) = 2^{j/2} \mathbf{f}(2^{j} x - n) \qquad \forall j, n \in \mathbb{Z}, \mathbf{f} \in \mathbb{C}^{\mathbb{C}}$$

Linear space properties 3.3

Proposition 3.4. Let T and D be as in Definition 3.3 page 40.

$$\mathbf{P}_{\mathbf{R}} \mathbf{D}^{j} \mathbf{T}^{n} [\mathsf{f} \mathsf{g}] = 2^{-j/2} [\mathbf{D}^{j} \mathbf{T}^{n} \mathsf{f}] [\mathbf{D}^{j} \mathbf{T}^{n} \mathsf{g}] \qquad \forall j,n \in \mathbb{Z}, \mathsf{f} \in \mathbb{C}^{\mathbb{C}}$$

[♠]Proof:

$$\mathbf{D}^{j}\mathbf{T}^{n}[f(x)g(x)] = 2^{j/2}f(2^{j}x - n)g(2^{j}x - n)$$
 by Proposition 3.3 page 41

$$= 2^{-j/2}[2^{j/2}f(2^{j}x - n)][2^{j/2}g(2^{j}x - n)]$$
 by Proposition 3.3 page 41

$$= 2^{-j/2}[\mathbf{D}^{j}\mathbf{T}^{n}f(x)][\mathbf{D}^{j}\mathbf{T}^{n}g(x)]$$
 by Proposition 3.3 page 41




In general the operators **T** and **D** are *noncommutative* (**TD** \neq **DT**), as demonstrated by Counterexample 3.1 (next) and Proposition 3.5 (page 42).

Counterexample 3.1.



As illustrated to the right, it is **not** always true that **TD** = **DT**:



Proposition 3.5 (commutator relation). ⁴ Let T and D be as in Definition 3.3 page 40.

New Proof:

$$\mathbf{D}^{j}\mathbf{T}^{2^{j}n}\mathsf{f}(x) = 2^{j/2}\,\mathsf{f}(2^{j}x-2^{j}n) \qquad \text{by Proposition 3.4 page 41}$$

$$= 2^{j/2}\,\mathsf{f}\left(2^{j}[x-n]\right) \qquad \text{by } distributivity \text{ of the field } (\mathbb{R},+,\cdot,0,1) \qquad \text{(Definition A.6 page 96)}$$

$$= \mathbf{T}^{n}2^{j/2}\,\mathsf{f}\left(2^{j}x\right) \qquad \text{by definition of } \mathbf{T} \qquad \text{(Definition 3.3 page 40)}$$

$$= \mathbf{T}^{n}\mathbf{D}^{j}\mathsf{f}(x) \qquad \text{by definition of } \mathbf{D} \qquad \text{(Definition 3.3 page 40)}$$

$$\mathbf{D}^{j}\mathbf{T}^{n}\mathsf{f}(x) = 2^{j/2}\,\mathsf{f}(2^{j}x-n) \qquad \text{by Proposition 3.4 page 41}$$

$$= 2^{j/2}\,\mathsf{f}\left(2^{j}\left[x-2^{-j/2}n\right]\right) \qquad \text{by } distributivity \text{ of the field } (\mathbb{R},+,\cdot,0,1) \qquad \text{(Definition A.6 page 96)}$$


$$= \mathbf{T}^{2^{-j/2}n}2^{j/2}\,\mathsf{f}\left(2^{j}x\right) \qquad \text{by definition of } \mathbf{T} \qquad \text{(Definition 3.3 page 40)}$$

$$= \mathbf{T}^{2^{-j/2}n}\mathbf{D}^{j}\mathsf{f}(x) \qquad \text{by definition of } \mathbf{D} \qquad \text{(Definition 3.3 page 40)}$$

3.4 Inner product space properties

In an inner product space, every operator has an *adjoint* (Proposition C.3 page 125) and this adjoint is always itself an operator. In the case where the adjoint and inverse of an operator U coincide, then U is said to be *unitary* (Definition C.14 page 135). And in this case, U has several nice properties (see Proposition 3.9 and Theorem 3.1 page 45). Proposition 3.6 (next) gives the adjoints of **D** and **T**, and Proposition 3.7 (page 43) demonstrates that both **D** and **T** are unitary. Other examples of unitary operators include the *Fourier Transform operator* $\tilde{\mathbf{F}}$ (Corollary I.1 page 197) and the *rotation matrix operator* (Example C.5 page 137).

Proposition 3.6. Let **T** be the Translation operator (Definition 3.3 page 40) with adjoint \mathbf{T}^* and \mathbf{D} the dilation operator with adjoint \mathbf{D}^* (Definition C.8 page 121).



⁴ Christensen (2003) page 42 ⟨equation (2.9)⟩, Dai and Larson (1998) page 21, Goodman et al. (1993a) page 641, Goodman et al. (1993b) page 110



№ Proof:

1. Proof that $T^*f(x) = f(x + 1)$:

$$\langle \mathsf{g}(x) \, | \, \mathbf{T}^*\mathsf{f}(x) \rangle = \langle \mathsf{g}(u) \, | \, \mathbf{T}^*\mathsf{f}(u) \rangle \qquad \qquad \text{by change of variable } x \to u$$

$$= \langle \mathbf{T}\mathsf{g}(u) \, | \, \mathsf{f}(u) \rangle \qquad \qquad \text{by definition of adjoint } \mathbf{T}^* \qquad \text{(Definition C.8 page 121)}$$

$$= \langle \mathsf{g}(u-1) \, | \, \mathsf{f}(u) \rangle \qquad \qquad \text{by definition of } \mathbf{T} \qquad \qquad \text{(Definition 3.3 page 40)}$$

$$= \langle \mathsf{g}(x) \, | \, \mathsf{f}(x+1) \rangle \qquad \qquad \text{where } x \triangleq u-1 \implies u=x+1$$

$$\Longrightarrow \mathbf{T}^*\mathsf{f}(x) = \mathsf{f}(x+1)$$

2. Proof that $\mathbf{D}^* f(x) = \frac{\sqrt{2}}{2} f\left(\frac{1}{2}x\right)$:

$$\langle \mathbf{g}(x) \, | \, \mathbf{D}^* \mathbf{f}(x) \rangle = \langle \mathbf{g}(u) \, | \, \mathbf{D}^* \mathbf{f}(u) \rangle \qquad \qquad \text{by change of variable } x \to u \\ = \langle \mathbf{D} \mathbf{g}(u) \, | \, \mathbf{f}(u) \rangle \qquad \qquad \text{by definition of } \mathbf{D}^* \qquad \qquad \text{(Definition C.8 page 121)} \\ = \langle \sqrt{2} \mathbf{g}(2u) \, | \, \mathbf{f}(u) \rangle \qquad \qquad \text{by definition of } \mathbf{D} \qquad \qquad \text{(Definition 3.3 page 40)} \\ = \int_{u \in \mathbb{R}} \sqrt{2} \mathbf{g}(2u) \mathbf{f}^*(u) \, \mathrm{du} \qquad \qquad \text{by definition of } \langle \triangle \, | \, \nabla \rangle \\ = \int_{x \in \mathbb{R}} \mathbf{g}(x) \left[\sqrt{2} \mathbf{f}\left(\frac{x}{2}\right) \frac{1}{2} \right]^* \, \mathrm{dx} \qquad \text{where } x = 2u \\ = \langle \mathbf{g}(x) \, | \, \frac{\sqrt{2}}{2} \mathbf{f}\left(\frac{x}{2}\right) \rangle \qquad \qquad \text{by definition of } \langle \triangle \, | \, \nabla \rangle \\ \implies \mathbf{D}^* \mathbf{f}(x) = \frac{\sqrt{2}}{2} \, \mathbf{f}\left(\frac{x}{2}\right)$$

Proposition 3.7. ⁵ Let \mathbf{T} and \mathbf{D} be as in Definition 3.3 (page 40). Let \mathbf{T}^{-1} and \mathbf{D}^{-1} be as in Proposition 3.2 (page 41).

P T is UNITARY in $L_{\mathbb{R}}^2$ ($\mathbf{T}^{-1} = \mathbf{T}^*$ in $L_{\mathbb{R}}^2$).
P D is UNITARY in $L_{\mathbb{R}}^2$ ($\mathbf{D}^{-1} = \mathbf{D}^*$ in $L_{\mathbb{R}}^2$).

♥Proof:

 $\mathbf{T}^{-1} = \mathbf{T}^*$ by Proposition 3.2 page 41 and Proposition 3.6 page 42 by the definition of *unitary* operators (Definition C.14 page 135) $\mathbf{D}^{-1} = \mathbf{D}^*$ by Proposition 3.2 page 41 and Proposition 3.6 page 42 by the definition of *unitary* operators (Definition C.14 page 135)

3.5 Normed linear space properties

Proposition 3.8. *Let* **D** *be the* DILATION OPERATOR (Definition 3.3 page 40).



⁵ Christensen (2003) page 41 ⟨Lemma 2.5.1⟩, Wojtaszczyk (1997) page 18 ⟨Lemma 2.5⟩



^ℚProof:

1. Proof that (1) \leftarrow *constant* property:

$$\mathbf{D}f(x) \triangleq \sqrt{2}f(2x)$$
 by definition of \mathbf{D} (Definition 3.3 page 40)
= $\sqrt{2}f(x)$ by *constant* hypothesis

2. Proof that (2) \leftarrow *constant* property:

$$\|f(x) - f(x+h)\| = \|f(x) - f(x)\| \quad \text{by } constant \text{ hypothesis}$$

$$= \|0\|$$

$$= 0 \quad \text{by } nondegenerate \text{ property of } \|\cdot\|$$

$$\leq \varepsilon$$

$$\implies \forall h > 0, \ \exists \varepsilon \quad \text{such that} \quad \|f(x) - f(x+h)\| < \varepsilon$$

$$\stackrel{\text{def}}{\iff} f(x) \text{ is } continuous$$

- 3. Proof that $(1,2) \implies constant$ property:
 - (a) Suppose there exists $x, y \in \mathbb{R}$ such that $f(x) \neq f(y)$.
 - (b) Let $(x_n)_{n\in\mathbb{N}}$ be a sequence with limit x and $(y_n)_{n\in\mathbb{N}}$ a sequence with limit y
 - (c) Then

$$0 < \|f(x) - f(y)\|$$
 by assumption in item (3a) page 44
$$= \lim_{n \to \infty} \|f(x_n) - f(y_n)\|$$
 by (2) and definition of (x_n) and (y_n) in item (3b) page 44
$$= \lim_{n \to \infty} \|f(2^m x_n) - f(2^\ell y_n)\| \quad \forall m, \ell \in \mathbb{Z}$$
 by (1)
$$= 0$$

(d) But this is a *contradiction*, so f(x) = f(y) for all $x, y \in \mathbb{R}$, and f(x) is *constant*.

Remark 3.1.


In Proposition 3.8 page 43, it is not possible to remove the continuous constraint outright, as demonstrated by the next two counterexamples.

Counterexample 3.2. Let f(x) be a function in $\mathbb{R}^{\mathbb{R}}$.

CNT

Let
$$f(x) \triangleq \begin{cases} 0 & \text{for } x = 0 \\ 1 & \text{otherwise.} \end{cases}$$

Then $\mathbf{D}f(x) \triangleq \sqrt{2}f(2x) = \sqrt{2}f(x)$, but f(x) is not constant.



—>


Counterexample 3.3. Let f(x) be a function in $\mathbb{R}^{\mathbb{R}}$.

Let \mathbb{Q} be the set of *rational numbers* and $\mathbb{R} \setminus \mathbb{Q}$ the set of *irrational numbers*.

CNT

Let
$$f(x) \triangleq \begin{cases} 1 & \text{for } x \in \mathbb{Q} \\ -1 & \text{for } x \in \mathbb{R} \setminus \mathbb{Q}. \end{cases}$$

Then $\mathbf{D}f(x) \triangleq \sqrt{2}f(2x) = \sqrt{2}f(x)$, but $f(x)$ is *not constant*.



Proposition 3.9 (Operator norm). Let **T** and **D** be as in Definition 3.3 page 40. Let \mathbf{T}^{-1} and \mathbf{D}^{-1} be as in Proposition 3.2 page 41. Let \mathbf{T}^* and \mathbf{D}^* be as in Proposition 3.6 page 42. Let $\|\cdot\|$ and $\langle \triangle \mid \nabla \rangle$ be as in Definition D.1 page 141. Let $\|\cdot\|$ be the operator norm (Definition C.6 page 117) induced by $\|\cdot\|$.

$$\|\mathbf{T}\| = \|\mathbf{D}\| = \|\mathbf{T}^*\| = \|\mathbf{D}^*\| = \|\mathbf{T}^{-1}\| = \|\mathbf{D}^{-1}\| = 1$$

PROOF: These results follow directly from the fact that T and D are *unitary* (Proposition 3.7 page 43) and from Theorem C.25 page 136 and Theorem C.26 page 136.

Theorem 3.1. Let **T** and **D** be as in Definition 3.3 page 40.

Let \mathbf{T}^{-1} and \mathbf{D}^{-1} be as in Proposition 3.2 page 41. Let $\|\cdot\|$ and $\langle \triangle \mid \nabla \rangle$ be as in Definition D.1 page 141.

T H M	1.	T f	=	D f	=	f	$\forall f \in \mathcal{L}^2_{\mathbb{R}}$	(ISOMETRIC IN LENGTH)
	2.	$\ \mathbf{T}f-\mathbf{T}g\ $		$\ \mathbf{D}f - \mathbf{D}g\ $		$\ f - g\ $	$\forall f,g{\in} oldsymbol{L}_{\mathbb{R}}^2$	(ISOMETRIC IN DISTANCE)
	3.	$\ \mathbf{T}^{-1}f - \mathbf{T}^{-1}g\ $	=	$\left\ \mathbf{D}^{-1}f-\mathbf{D}^{-1}g\right\ $	=	$\ f - g\ $	$\forall f,g \in \mathcal{L}_{\mathbb{R}}^2$	(ISOMETRIC IN DISTANCE)
	4.	$\langle \mathbf{Tf} \mid \mathbf{Tg} \rangle$	=	$\langle \mathbf{D} f \mid \mathbf{D} g \rangle$	=	$\langle f \mid g \rangle$	$\forall f,g{\in} oldsymbol{L}_{\mathbb{R}}^2$	(SURJECTIVE)
	5.	$\langle \mathbf{T}^{-1} \mathbf{f} \mathbf{T}^{-1} \mathbf{g} \rangle$	=	$\langle \mathbf{D}^{-1} f \mid \mathbf{D}^{-1} g \rangle$	=	$\langle f \mid g \rangle$	$\forall f,\!g{\in} oldsymbol{L}_{\mathbb{R}}^2$	(SURJECTIVE)

 \P Proof: These results follow directly from the fact that **T** and **D** are *unitary* (Proposition 3.7 page 43) and from Theorem C.25 page 136 and Theorem C.26 page 136.

Proposition 3.10. Let **T** be as in Definition 3.3 page 40. Let **A*** be the ADJOINT (Definition C.8 page 121) of an operator **A**. Let the property "SELF ADJOINT" be defined as in Definition C.11 (page 129).

$$\left(\sum_{n\in\mathbb{Z}}\mathbf{T}^{n}\right) = \left(\sum_{n\in\mathbb{Z}}\mathbf{T}^{n}\right)^{*} \qquad \left(The\ operator\left[\sum_{n\in\mathbb{Z}}\mathbf{T}^{n}\right]\ is\ \text{Self-Adjoint}\right)$$

№PROOF:

$$\left\langle \left(\sum_{n\in\mathbb{Z}}\mathbf{T}^n\right)\mathbf{f}(x)\,|\,\mathbf{g}(x)\right\rangle = \left\langle \sum_{n\in\mathbb{Z}}\mathbf{f}(x-n)\,|\,\mathbf{g}(x)\right\rangle \qquad \text{by definition of }\mathbf{T} \qquad \text{(Definition 3.3 page 40)}$$

$$= \left\langle \sum_{n\in\mathbb{Z}}\mathbf{f}(x+n)\,|\,\mathbf{g}(x)\right\rangle \qquad \text{by }commutative \text{ property} \qquad \text{(Definition A.5 page 96)}$$

$$= \sum_{n\in\mathbb{Z}}\left\langle \mathbf{f}(x+n)\,|\,\mathbf{g}(x)\right\rangle \qquad \text{by }additive \text{ property of }\left\langle \triangle\mid \nabla\right\rangle$$

$$= \sum_{n\in\mathbb{Z}}\left\langle \mathbf{f}(u)\,|\,\mathbf{g}(u-n)\right\rangle \qquad \text{where }u\triangleq x+n$$

$$= \left\langle \mathbf{f}(u)\,\left|\,\sum_{n\in\mathbb{Z}}\mathbf{g}(u-n)\right\rangle \qquad \text{by }additive \text{ property of }\left\langle \triangle\mid \nabla\right\rangle$$

$$= \left\langle \mathbf{f}(x)\,\left|\,\sum_{n\in\mathbb{Z}}\mathbf{g}(x-n)\right\rangle \qquad \text{by change of variable: }u\to x$$


$$= \left\langle \mathbf{f}(x)\,\left|\,\sum_{n\in\mathbb{Z}}\mathbf{T}^n\mathbf{g}(x)\right\rangle \qquad \text{by definition of }\mathbf{T} \qquad \text{(Definition 3.3 page 40)}$$

$$\Leftrightarrow \left(\sum_{n\in\mathbb{Z}}\mathbf{T}^n\right) = \left(\sum_{n\in\mathbb{Z}}\mathbf{T}^n\right)^* \qquad \text{by definition of }adjoint \qquad \text{(Proposition C.3 page 125)}$$

$$\Leftrightarrow \left(\sum_{n\in\mathbb{Z}}\mathbf{T}^n\right) \text{ is }self-adjoint \qquad \text{by definition of }self-adjoint \qquad \text{(Definition C.11 page 129)}$$

 \blacksquare







3.6 Fourier transform properties

Proposition 3.11. Let T and D be as in Definition 3.3 page 40.

Let **B** be the Two-Sided Laplace transform defined as [**B**f](s) $\triangleq \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} f(x)e^{-sx} dx$.

№ Proof:

$$\mathbf{BT}^{n} \mathbf{f}(x) = \mathbf{Bf}(x - n) \qquad \text{by definition of } \mathbf{T}$$

$$= \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} \mathbf{f}(x - n)e^{-sx} \, dx \qquad \text{by definition of } \mathbf{B}$$

$$= \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} \mathbf{f}(u)e^{-s(u+n)} \, du \qquad \text{where } u \triangleq x - n$$

$$= e^{-sn} \left[\frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} \mathbf{f}(u)e^{-su} \, du \right]$$

$$= e^{-sn} \mathbf{Bf}(x) \qquad \text{by definition of } \mathbf{B}$$

$$\begin{aligned} \mathbf{B}\mathbf{D}^{j}\mathbf{f}(x) &= \mathbf{B}\big[2^{j/2}\,\mathbf{f}\big(2^{j}x\big)\big] & \text{by definition of }\mathbf{D} \\ &= \frac{1}{\sqrt{2\pi}}\,\int_{\mathbb{R}}\big[2^{j/2}\,\mathbf{f}\big(2^{j}x\big)\big]e^{-sx}\,\,\mathrm{d}x & \text{by definition of }\mathbf{B} \\ &= \frac{1}{\sqrt{2\pi}}\,\int_{\mathbb{R}}\big[2^{j/2}\,\mathbf{f}(u)\big]e^{-s2^{-j}}2^{-j}\,\,\mathrm{d}u & \text{let }u\triangleq 2^{j}x \implies x=2^{-j}u \\ &= \frac{\sqrt{2}}{2}\,\frac{1}{\sqrt{2\pi}}\,\int_{\mathbb{R}}\mathbf{f}(u)e^{-s2^{-j}u}\,\,\mathrm{d}u \\ &= \mathbf{D}^{-1}\,\bigg[\frac{1}{\sqrt{2\pi}}\,\int_{\mathbb{R}}\mathbf{f}(u)e^{-su}\,\,\mathrm{d}u\bigg] & \text{by Proposition 3.6 page 42 and} & \text{Proposition 3.7 page 43} \\ &= \mathbf{D}^{-j}\,\mathbf{B}\,\mathbf{f}(x) & \text{by definition of }\mathbf{B} \\ &\mathbf{D}\mathbf{B}\,\mathbf{f}(x) = \mathbf{D}\bigg[\frac{1}{\sqrt{2\pi}}\,\int_{\mathbb{R}}\mathbf{f}(x)e^{-sx}\,\,\mathrm{d}x\bigg] & \text{by definition of }\mathbf{B} \\ &= \frac{\sqrt{2}}{\sqrt{2\pi}}\,\int_{\mathbb{R}}\mathbf{f}(x)e^{-2sx}\,\,\mathrm{d}x & \text{by definition of }\mathbf{D} \end{aligned} \tag{Definition 3.3 page 40}$$

$$= \frac{\sqrt{2}}{\sqrt{2\pi}} \int_{\mathbb{R}} f\left(\frac{u}{2}\right) e^{-su} \frac{1}{2} du \qquad \text{let } u \triangleq 2x \implies x = \frac{1}{2}u$$

$$= \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} \left[\frac{\sqrt{2}}{2} f\left(\frac{u}{2}\right)\right] e^{-su} du$$

$$= \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} \left[\mathbf{D}^{-1} f\right](u) e^{-su} du \qquad \text{by Proposition 3.6 page 42 and} \qquad \text{Proposition 3.7 page 43}$$

$$= \mathbf{B} \mathbf{D}^{-1} f(x) \qquad \text{by definition of } \mathbf{B}$$

$$\mathbf{B}^{-1}\mathbf{D}^{-1}\mathbf{B} = \mathbf{B}^{-1}\mathbf{B}\mathbf{D}$$
 by previous result
$$= \mathbf{D}$$
 by definition of operator inverse (Definition C.3 page 112)
$$\mathbf{B}\mathbf{D}^{-1}\mathbf{B}^{-1} = \mathbf{D}\mathbf{B}\mathbf{B}^{-1}$$
 by previous result

= **D** by definition of operator inverse (Definition C.3 page 112)

$$\begin{aligned} \textbf{DBD} &= \textbf{DD}^{-1}\textbf{B} & \text{by previous result} \\ &= \textbf{B} & \text{by definition of operator inverse} & \text{(Definition C.3 page 112)} \\ \textbf{D}^{-1}\textbf{BD}^{-1} &= \textbf{D}^{-1}\textbf{DB} & \text{by previous result} \\ &= \textbf{B} & \text{by definition of operator inverse} & \text{(Definition C.3 page 112)} \end{aligned}$$

Corollary 3.1. Let \mathbf{T} and \mathbf{D} be as in Definition 3.3 page 40. Let $\tilde{\mathbf{f}}(\omega) \triangleq \tilde{\mathbf{F}}\mathbf{f}(x)$ be the Fourier Transform (Definition 1.2 page 196) of some function $\mathbf{f} \in \mathcal{L}^2_{\mathbb{R}}$ (Definition D.1 page 141).

1.
$$\tilde{\mathbf{F}}\mathbf{T}^{n} = e^{-i\omega n}\tilde{\mathbf{F}}$$

2. $\tilde{\mathbf{F}}\mathbf{D}^{j} = \mathbf{D}^{-j}\tilde{\mathbf{F}}$
3. $\mathbf{D}\tilde{\mathbf{F}} = \tilde{\mathbf{F}}\mathbf{D}^{-1}$
4. $\mathbf{D} = \tilde{\mathbf{F}}\mathbf{D}^{-1}\tilde{\mathbf{F}}^{-1} = \tilde{\mathbf{F}}^{-1}\mathbf{D}^{-1}\tilde{\mathbf{F}}$
5. $\tilde{\mathbf{F}} = \mathbf{D}\tilde{\mathbf{F}}\mathbf{D} = \mathbf{D}^{-1}\tilde{\mathbf{F}}\mathbf{D}^{-1}$

 $^{\text{N}}$ Proof: These results follow directly from Proposition 3.11 page 46 with $\tilde{\mathbf{F}} = \mathbf{B}|_{s=i\omega}$.

Proposition 3.12. Let **T** and **D** be as in Definition 3.3 page 40. Let $\tilde{f}(\omega) \triangleq \tilde{\mathbf{F}} f(x)$ be the Fourier Transform (Definition 1.2 page 196) of some function $f \in L^2_{\mathbb{R}}$ (Definition D.1 page 141).

$$\mathbf{\tilde{F}}\mathbf{D}^{j}\mathbf{T}^{n}\mathbf{f}(x) = \frac{1}{2^{j/2}}e^{-i\frac{\omega}{2^{j}}n}\tilde{\mathbf{f}}\left(\frac{\omega}{2^{j}}\right)$$

№Proof:

$$\mathbf{\tilde{F}}\mathbf{D}^{j}\mathbf{T}^{n}f(x) = \mathbf{D}^{-j}\mathbf{\tilde{F}}\mathbf{T}^{n}f(x) \qquad \text{by Corollary 3.1 page 47 (3)}
= \mathbf{D}^{-j}e^{-i\omega n}\mathbf{\tilde{F}}f(x) \qquad \text{by Corollary 3.1 page 47 (3)}
= \mathbf{D}^{-j}e^{-i\omega n}\mathbf{\tilde{f}}(\omega) \qquad \text{by Proposition 3.2 page 41}$$

Proposition 3.13. Let **T** be the translation operator (Definition 3.3 page 40). Let $\tilde{\mathbf{f}}(\omega) \triangleq \tilde{\mathbf{F}}\mathbf{f}(x)$ be the FOURIER TRANSFORM (Definition 1.2 page 196) of a function $\mathbf{f} \in \mathbf{L}^2_{\mathbb{R}}$. Let $\check{\mathbf{a}}(\omega)$ be the DTFT (Definition M.1 page 237) of a sequence $(a_n)_{n\in\mathbb{Z}} \in \boldsymbol{\ell}^2_{\mathbb{R}}$ (Definition J.2 page 207).


$$\begin{array}{c} sequence \, (\!(a_n)\!)_{n \in \mathbb{Z}} \in \mathscr{C}^2_{\mathbb{R}} \, (\text{Definition J.2 page 207}). \\ \\ \overset{\mathsf{P}}{\underset{\mathsf{P}}{\mathsf{R}}} \, \tilde{\mathbf{F}} \sum_{n \in \mathbb{Z}} a_n \mathbf{T}^n \phi(x) = \check{\mathbf{a}}(\omega) \tilde{\phi}(\omega) \qquad \forall (a_n) \in \mathscr{C}^2_{\mathbb{R}}, \phi(x) \in \mathscr{L}^2_{\mathbb{R}} \\ \end{array}$$

♥Proof:

$$\begin{split} \tilde{\mathbf{F}} \sum_{n \in \mathbb{Z}} a_n \mathbf{T}^n \phi(x) &= \sum_{n \in \mathbb{Z}} a_n \tilde{\mathbf{F}} \mathbf{T}^n \phi(x) \\ &= \sum_{n \in \mathbb{Z}} a_n e^{-i\omega n} \tilde{\mathbf{F}} \phi(x) & \text{by Corollary 3.1 page 47} \\ &= \left[\sum_{n \in \mathbb{Z}} a_n e^{-i\omega n} \right] \tilde{\phi}(\omega) & \text{by definition of } \tilde{\phi}(\omega) \\ &= \check{\mathbf{a}}(\omega) \tilde{\phi}(\omega) & \text{by definition of } DTFT \text{ (Definition M.1 page 237)} \end{split}$$

 \Rightarrow





Definition 3.4. Let $L^2_{(\mathbb{R},\mathcal{B},\mu)}$ be the space of Lebesgue square-integrable functions (Definition D.1 page 141). Let $\ell^2_{\mathbb{R}}$ be the space of all absolutely square summable sequences over \mathbb{R} (Definition D.1 page 141).

DE

S is the **sampling operator** in $\mathscr{C}^{2}_{\mathbb{R}}^{\mathcal{L}^{2}_{\mathbb{R}}}$ if $[\mathbf{Sf}(x)](n) \triangleq f\left(\frac{2\pi}{\tau}n\right)$ $\forall f \in \mathcal{L}^{2}_{(\mathbb{R},\mathcal{B},\mu)}, \tau \in \mathbb{R}^{+}$

Theorem 3.2 (Poisson Summation Formula—PSF). ⁶ Let $\tilde{f}(\omega)$ be the Fourier transform (Definition 1.2 page 196) of a function $f(x) \in L^2_{\mathbb{R}}$. Let S be the SAMPLING OPERATOR (Definition 3.4 page 48).

$$\underbrace{\sum_{n \in \mathbb{Z}} \mathbf{T}_{\tau}^{n} \mathbf{f}(x) = \sum_{n \in \mathbb{Z}} \mathbf{f}(x + n\tau)}_{summation \ in "time"} = \underbrace{\sqrt{\frac{2\pi}{\tau}} \hat{\mathbf{F}}^{-1} \mathbf{S} \tilde{\mathbf{F}}[\mathbf{f}(x)]}_{operator \ notation} = \underbrace{\frac{\sqrt{2\pi}}{\tau} \sum_{n \in \mathbb{Z}} \tilde{\mathbf{f}}\left(\frac{2\pi}{\tau}n\right) e^{i\frac{2\pi}{\tau}nx}}_{summation \ in "frequency"}$$

№ Proof:

1. lemma: If $h(x) \triangleq \sum_{n \in \mathbb{Z}} f(x + n\tau)$ then $h \equiv \hat{\mathbf{F}}^{-1}\hat{\mathbf{F}}h$. Proof:

Note that h(x) is *periodic* with period τ . Because h is periodic, it is in the domain of $\hat{\mathbf{F}}$ and thus $h \equiv \hat{\mathbf{F}}^{-1}\hat{\mathbf{F}}h$.

2. Proof of PSF (this theorem—Theorem 3.2):

$$\begin{split} \sum_{n\in\mathbb{Z}} \mathsf{f}(x+n\tau) &= \hat{\mathbf{F}}^{-1} \hat{\mathbf{F}} \sum_{n\in\mathbb{Z}} \mathsf{f}(x+n\tau) & \text{by (1) lemma page 48} \\ &= \hat{\mathbf{F}}^{-1} \left[\frac{1}{\sqrt{\tau}} \int_{0}^{\tau} \left(\sum_{n\in\mathbb{Z}} \mathsf{f}(x+n\tau) \right) e^{-i\frac{2\pi}{\tau}kx} \, \mathrm{d}x \right] & \text{by definition of } \hat{\mathbf{F}} & \text{(Definition N.1 page 247)} \\ &= \hat{\mathbf{F}}^{-1} \left[\frac{1}{\sqrt{\tau}} \sum_{n\in\mathbb{Z}} \int_{0}^{\tau} \mathsf{f}(x+n\tau) e^{-i\frac{2\pi}{\tau}kx} \, \mathrm{d}x \right] \\ &= \hat{\mathbf{F}}^{-1} \left[\frac{1}{\sqrt{\tau}} \sum_{n\in\mathbb{Z}} \int_{u=n\tau}^{u=(n+1)\tau} \mathsf{f}(u) e^{-i\frac{2\pi}{\tau}k(u-n\tau)} \, \mathrm{d}u \right] & \text{where } u \triangleq x+n\tau \implies x = u-n\tau \\ &= \hat{\mathbf{F}}^{-1} \left[\frac{1}{\sqrt{\tau}} \sum_{n\in\mathbb{Z}} e^{i2\pi kn^{\bullet}} \int_{u=n\tau}^{u=(n+1)\tau} \mathsf{f}(u) e^{-i\frac{2\pi}{\tau}ku} \, \mathrm{d}u \right] \\ &= \sqrt{\frac{2\pi}{\tau}} \hat{\mathbf{F}}^{-1} \left[\frac{1}{\sqrt{2\pi}} \int_{u\in\mathbb{R}} \mathsf{f}(u) e^{-i\left(\frac{2\pi}{\tau}k\right)u} \, \mathrm{d}u \right] & \text{by evaluation of } \hat{\mathbf{F}}^{-1} & \text{(Theorem N.1 page 248)} \\ &= \sqrt{\frac{2\pi}{\tau}} \, \hat{\mathbf{F}}^{-1} \left[\left[\tilde{\mathbf{F}}\mathsf{f}(x) \right] \left(\frac{2\pi}{\tau}k \right) \right] & \text{by definition of } \hat{\mathbf{S}} & \text{(Definition 1.2 page 196)} \\ &= \sqrt{\frac{2\pi}{\tau}} \, \hat{\mathbf{F}}^{-1} \mathbf{S} \tilde{\mathbf{F}} \mathsf{f} & \text{by definition of } \hat{\mathbf{S}} & \text{(Definition 3.4 page 48)} \\ &= \frac{\sqrt{2\pi}}{\tau} \, \sum_{i=1}^{\infty} \tilde{\mathbf{f}} \left(\frac{2\pi}{\tau}n \right) e^{i\frac{2\pi}{\tau}nx} & \text{by evaluation of } \hat{\mathbf{F}}^{-1} & \text{(Theorem N.1 page 248)} \end{aligned}$$




🤲 Frames and Bases Structure and Design [VERSION 021] 🥙

₽

Theorem 3.3 (Inverse Poisson Summation Formula—IPSF).

Let $\tilde{f}(\omega)$ be the Fourier transform (Definition 1.2 page 196) of a function $f(x) \in L^2_{\mathbb{R}}$.



NPROOF:

1. lemma: If $h(\omega) \triangleq \sum_{n \in \mathbb{Z}} \tilde{f}\left(\omega + \frac{2\pi}{\tau}n\right)$, then $h \equiv \hat{\mathbf{F}}^{-1}\hat{\mathbf{F}}h$. Proof: Note that $h(\omega)$ is periodic with period $2\pi/T$:

$$\mathsf{h}\left(\omega + \frac{2\pi}{\tau}\right) \triangleq \sum_{n \in \mathbb{Z}} \tilde{\mathsf{f}}\left(\omega + \frac{2\pi}{\tau} + \frac{2\pi}{\tau}n\right) = \sum_{n \in \mathbb{Z}} \tilde{\mathsf{f}}\left(\omega + (n+1)\frac{2\pi}{\tau}\right) = \sum_{n \in \mathbb{Z}} \tilde{\mathsf{f}}\left(\omega + \frac{2\pi}{\tau}n\right) \triangleq \mathsf{h}(\omega)$$

Because h is periodic, it is in the domain of $\hat{\mathbf{F}}$ and is equivalent to $\hat{\mathbf{F}}^{-1}\hat{\mathbf{F}}$ h.

2. Proof of IPSF (this theorem—Theorem 3.3):

$$\begin{split} &\sum_{n\in\mathbb{Z}}\tilde{\mathbf{f}}\left(\omega+\frac{2\pi}{\tau}n\right)\\ &=\hat{\mathbf{F}}^{-1}\hat{\mathbf{f}}\sum_{n\in\mathbb{Z}}\tilde{\mathbf{f}}\left(\omega+\frac{2\pi}{\tau}n\right) & \text{by (1) lemma page 49} \\ &=\hat{\mathbf{F}}^{-1}\left[\sqrt{\frac{\tau}{2\pi}}\int_{0}^{\frac{2\pi}{\tau}}\sum_{n\in\mathbb{Z}}\tilde{\mathbf{f}}\left(\omega+\frac{2\pi}{\tau}n\right)e^{-i\omega\frac{2\pi}{2\pi i\tau}k}\,\mathrm{d}\omega\right] & \text{by definition of }\hat{\mathbf{F}} & \text{(Definition N.1 page 247)} \\ &=\hat{\mathbf{F}}^{-1}\left[\sqrt{\frac{\tau}{2\pi}}\sum_{n\in\mathbb{Z}}\int_{0}^{\frac{2\pi}{\tau}}\tilde{\mathbf{f}}\left(\omega+\frac{2\pi}{\tau}n\right)e^{-i\omega Tk}\,\mathrm{d}\omega\right] \\ &=\hat{\mathbf{F}}^{-1}\left[\sqrt{\frac{\tau}{2\pi}}\sum_{n\in\mathbb{Z}}\int_{u=\frac{2\pi}{\tau}}^{u=\frac{2\pi}{\tau}(n+1)}\tilde{\mathbf{f}}\left(u\right)e^{-i(u-\frac{2\pi}{\tau}n)Tk}\,\mathrm{d}u\right] & \text{where } u\triangleq\omega+\frac{2\pi}{\tau}n \implies \omega=u-\frac{2\pi}{\tau}n \\ &=\hat{\mathbf{F}}^{-1}\left[\sqrt{\frac{\tau}{2\pi}}\sum_{n\in\mathbb{Z}}e^{i2\pi nk^{\star}}\int_{\frac{2\pi}{\tau}n}^{\frac{2\pi}{\tau}(n+1)}\tilde{\mathbf{f}}\left(u\right)e^{-iu\tau k}\,\mathrm{d}u\right] \\ &=\hat{\mathbf{F}}^{-1}\left[\sqrt{\frac{\tau}{2\pi}}\int_{\mathbb{R}}\tilde{\mathbf{f}}\left(u\right)e^{-iu\tau k}\,\mathrm{d}u\right] \\ &=\hat{\mathbf{F}}^{-1}\left[\sqrt{\frac{\tau}{2\pi}}\int_{\mathbb{R}}\tilde{\mathbf{f}}\left(u\right)e^{iu(-\tau k)}\,\mathrm{d}u\right] \\ &=\sqrt{\tau}\,\hat{\mathbf{F}}^{-1}\left[\left[\tilde{\mathbf{F}}^{-1}\tilde{\mathbf{f}}\left(-k\tau\right)\right]\right] & \text{by value of }\tilde{\mathbf{F}}^{-1} \\ &=\sqrt{\tau}\,\hat{\mathbf{F}}^{-1}\mathbf{S}\tilde{\mathbf{F}}^{-1}\,\tilde{\mathbf{f}} & \text{by definition of } \mathbf{S} & \text{(Definition 1.2 page 196)} \\ &=\sqrt{\tau}\,\hat{\mathbf{F}}^{-1}\mathbf{S}\mathbf{f}\left(x\right) & \text{by definition of } \tilde{\mathbf{F}} & \text{(Definition 1.2 page 196)} \end{split}$$



(Definition 3.4 page 48)

(Theorem N.1 page 248)

 $= \sqrt{\tau} \, \hat{\mathbf{F}}^{-1} \mathsf{f}(-k\tau)$

 $= \sqrt{\tau} \frac{1}{\sqrt{\frac{2\pi}{\tau}}} \sum_{k \in \mathbb{Z}} f(-k\tau) e^{i2\pi \frac{i\pi}{\tau} k\omega}$

by definition of S

by definition of $\hat{\mathbf{F}}^{-1}$

⁷ Gauss (1900) page 88

$$= \frac{\tau}{\sqrt{\frac{2\pi}{\tau}}} \sum_{k \in \mathbb{Z}} f(-k\tau)e^{ik\tau\omega}$$
 by definition of $\hat{\mathbf{F}}^{-1}$ (Theorem N.1 page 248)
$$= \frac{\tau}{\sqrt{2\pi}} \sum_{m \in \mathbb{Z}} f(m\tau)e^{-i\omega m\tau}$$
 let $m \triangleq -k$

Remark 3.2. The left hand side of the *Poisson Summation Formula* (Theorem 3.2 page 48) is very similar to the *Zak Transform* \mathbf{Z} : ⁸

to the Zak Transform **Z**: ⁸

$$(\mathbf{Z}f)(t,\omega) \triangleq \sum_{n \in \mathbb{Z}} f(x+n\tau)e^{i2\pi n\omega}$$

Remark 3.3. A generalization of the *Poisson Summation Formula* (Theorem 3.2 page 48) is the **Selberg Trace Formula**. ⁹

3.7 Basis theory properties

Example 3.2 (linear functions). ¹⁰ Let **T** be the *translation operator* (Definition 3.3 page 40). Let $\mathcal{L}(\mathbb{C},\mathbb{C})$ be the set of all *linear* functions in $L^2_{\mathbb{R}}$.

1.
$$\{x, \mathbf{T}x\}$$
 is a *basis* for $\mathcal{L}(\mathbb{C}, \mathbb{C})$ and 2. $f(x) = f(1)x - f(0)\mathbf{T}x$ $\forall f \in \mathcal{L}(\mathbb{C}, \mathbb{C})$

PROOF: By left hypothesis, f is *linear*; so let $f(x) \triangleq ax + b$

$$f(1)x - f(0)\mathbf{T}x = f(1)x - f(0)(x - 1)$$
 by Definition 3.3 page 40

$$= (ax + b)|_{x=1} x - (ax + b)|_{x=0} (x - 1)$$
 by left hypothesis and definition of f

$$= (a + b)x - b(x - 1)$$
 by left hypothesis and definition of f

$$= ax + bx - bx + b$$
 by left hypothesis and definition of f

Example 3.3 (Cardinal Series). Let **T** be the *translation operator* (Definition 3.3 page 40). The *Paley-Wiener* class of functions PW_{σ}^2 are those functions which are "bandlimited" with respect to their Fourier transform (Definition 1.2 page 196). The cardinal series forms an orthogonal basis for such a space. The *Fourier coefficients* (Definition 2.11 page 20) for a projection of a function f onto the Cardinal series basis elements is particularly simple—these coefficients are samples of f(x) taken at regular intervals. In fact, one could represent the coefficients using inner product notation with the *Dirac delta distribution* δ as follows:

$$\langle f(x) | \mathbf{T}^{n} \delta(x) \rangle \triangleq \int_{\mathbb{R}} f(x) \delta(x - n) \, dx \triangleq f(n)$$
1.
$$\left\{ \mathbf{T}^{n} \frac{\sin(\pi x)}{\pi x} \middle| n \in \mathbb{N} \right\} \text{ is a } basis \text{ for } \mathbf{PW}_{\sigma}^{2} \text{ and}$$
2.
$$f(x) = \sum_{n=1}^{\infty} f(n) \mathbf{T}^{n} \frac{\sin(\pi x)}{\pi x} \qquad \forall f \in \mathbf{PW}_{\sigma}^{2}, \sigma \leq \frac{1}{2}$$
Cardinal series

⁸ Janssen (1988) page 24, Zayed (1996) page 482

¹⁰ Higgins (1996) page 2 (1.1 General introduction)



⁹ ■ Lax (2002) page 349, ■ Selberg (1956), ■ Terras (1999)

Example 3.4 (Fourier Series).

(1). $\left\{ \mathbf{D}_{n} e^{ix} \mid n \in \mathbb{Z} \right\}$ is a *basis* for $\mathbf{L}(0:2\pi)$ (2). $\mathbf{f}(x) = \frac{1}{\sqrt{2\pi}} \sum_{n \in \mathbb{Z}} \alpha_{n} \mathbf{D}_{n} e^{ix} \quad \forall x \in (0:2\pi), \mathbf{f} \in \mathbf{L}(0:2\pi)$ where E X $\alpha_n \triangleq \frac{1}{\sqrt{2\pi}} \int_0^{2\pi} f(x) \mathbf{D}_n e^{-ix} dx \quad \forall f \in L(0:2\pi)$

[♠]Proof: See Theorem N.1 page 248.

Example 3.5 (Fourier Transform). 11

(1).
$$\left\{ \mathbf{D}_{\omega} e^{ix} | \omega \in \mathbb{R} \right\}$$
 is a *basis* for $\mathbf{L}_{\mathbb{R}}^{2}$ and (2). $f(x) = \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} \tilde{f}(\omega) \mathbf{D}_{x} e^{i\omega} d\omega \quad \forall f \in \mathbf{L}_{\mathbb{R}}^{2}$ where $\tilde{f}(\omega) \triangleq \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} f(x) \mathbf{D}_{\omega} e^{-ix} dx \quad \forall f \in \mathbf{L}_{\mathbb{R}}^{2}$

Example 3.6 (Gabor Transform). 12

(1).
$$\left\{ \left(\mathbf{T}_{\tau} e^{-\pi x^{2}} \right) \left(\mathbf{D}_{\omega} e^{ix} \right) \middle| \tau, \omega \in \mathbb{R} \right\}$$
 is a basis for $\mathbf{L}_{\mathbb{R}}^{2}$ and

(2). $\mathbf{f}(x) = \int_{\mathbb{R}} \mathbf{G}(\tau, \omega) \mathbf{D}_{x} e^{i\omega} d\omega \qquad \forall x \in \mathbb{R}, \mathbf{f} \in \mathbf{L}_{\mathbb{R}}^{2}$ where $\mathbf{G}(\tau, \omega) \triangleq \int_{\mathbb{R}} \mathbf{f}(x) \left(\mathbf{T}_{\tau} e^{-\pi x^{2}} \right) \left(\mathbf{D}_{\omega} e^{-ix} \right) dx \quad \forall x \in \mathbb{R}, \mathbf{f} \in \mathbf{L}_{\mathbb{R}}^{2}$

Example 3.7 (wavelets). Let $\psi(x)$ be a *wavelet*.

(1).
$$\left\{ \mathbf{D}^{k} \mathbf{T}^{n} \psi(x) \middle| k, n \in \mathbb{Z} \right\}$$
 is a *basis* for $\mathcal{L}_{\mathbb{R}}^{2}$ and
(2). $f(x) = \sum_{k \in \mathbb{Z}} \sum_{n \in \mathbb{Z}} \alpha_{k,n} \mathbf{D}^{k} \mathbf{T}^{n} \psi(x) \quad \forall f \in \mathcal{L}_{\mathbb{R}}^{2}$ where
$$\alpha_{n} \triangleq \int_{\mathbb{R}} f(x) \mathbf{D}^{k} \mathbf{T}^{n} \psi^{*}(x) \, dx \quad \forall f \in \mathcal{L}_{\mathbb{R}}^{2}$$

¹¹cross reference: Definition I.2 page 196

¹² Gabor (1946),

Qian and Chen (1996) ⟨Chapter 3⟩,
Forster and Massopust (2009) page 32 ⟨Definition 1.69⟩

...on fait la science avec des faits comme une maison avec des pierres; mais une accumulation de faits n'est pas plus une science qu'un tas de pierres n'est une maison.




Science is built up of facts, as a house is built of stones; but an accumulation of facts is no more a science than a heap of stones is a house.

Jules Henri Poincaré (1854-1912), physicist and mathematician ¹

4.1 Introduction

In 1989, Stéphane G. Mallat introduced the *Multiresolution* Analysis (MRA, Definition 4.1 page 54) method for wavelet construction. The MRA has become the dominate wavelet construction method. This text uses the MRA method extensively, and combines the MRA "scaling subspaces" (Definition 4.1 page 54) with "wavelet subspaces" (Definition 5.1 page 81) to form a subspace structure as represented by the Hasse diagram to the right. The Fast Wavelet Transform combines both sets of subspaces as well, providing the results of projections onto both wavelet and MRA subspaces. Moreover, P.G. Lemarié has proved that all wavelets with compact support are generated by an MRA.²



The MRA is an **analysis** of the linear space $L^2_{\mathbb{R}}$. An analysis of a linear space \boldsymbol{X} is any sequence $(\boldsymbol{V}_j)_{j\in\mathbb{Z}}$ of linear subspaces of \boldsymbol{X} . The partial or complete reconstruction of \boldsymbol{X} from $(\boldsymbol{V}_j)_{j\in\mathbb{Z}}$ is a synthesis.³ An analysis is completely *characterized* by a *transform*. For example, a Fourier analysis is a sequence of subspaces with sinusoidal bases. Examples of subspaces in a Fourier analysis

¹ quote: Poincaré (1902a) (Chapter IX, paragraph 7) translation: Poincaré (1902b) page 141 http://www-groups.dcs.st-and.ac.uk/~history/PictDisplay/Poincare.html

² ⚠ Lemarié (1990), ⋒ Mallat (1999) page 240

³The word *analysis* comes from the Greek word ἀνάλυσις, meaning "dissolution" (
Perschbacher (1990) page 23

include $V_1 = \text{span}\{e^{ix}\}$, $V_{2.3} = \text{span}\{e^{i2.3x}\}$, $V_{\sqrt{2}} = \text{span}\{e^{i\sqrt{2}x}\}$, etc. A **transform** is loosely defined as a function that maps a family of functions into an analysis. A very useful transform (a "Fourier transform") for Fourier Analysis is (Definition I.2 page 196)

$$[\tilde{\mathbf{F}}\mathbf{f}](\omega) \triangleq \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} \mathbf{f}(x)e^{-i\omega x} dx$$

4.2 **Definition**

A multiresolution analysis provides "coarse" approximations of a function in a linear space $L^2_{\mathbb{R}}$ at multiple "scales" or "resolutions". Key to this process is a sequence of *scaling functions*. Most traditional transforms feature a single scaling function $\phi(x)$ set equal to one $(\phi(x) = 1)$. This allows for convenient representation of the most basic functions, such as constants.⁴ A multiresolution system, on the other hand, uses a generalized form of the scaling concept:

- 1. Instead of the scaling function simply being set *equal to unity* ($\phi(x) = 1$), a multiresolution system (Definition 4.3 page 63) is often constructed in such a way that the scaling function $\phi(x)$ forms a partition of unity such that $\sum_{n \in \mathbb{Z}} \mathbf{T}^n \phi(x) = 1$.
- 2. Instead of there being *just one* scaling function, there is an entire sequence of scaling functions $(\mathbf{D}^j \phi(x))_{i \in \mathbb{Z}}$, each corresponding to a different "resolution".

Definition 4.1. ⁵ Let $(V_j)_{j \in \mathbb{Z}}$ be a sequence of subspaces on $L^2_{\mathbb{R}}$ (Definition D.1 page 141). Let A^- be the CLOSURE of a set A.

The sequence $(V_j)_{j\in\mathbb{Z}}$ is a multiresolution analysis on $L^2_{\mathbb{R}}$ if

1. $V_j = V_j^ \forall j\in\mathbb{Z}$ (CLOSED) and

2. $V_j \subset V_{j+1}$ $\forall j\in\mathbb{Z}$ (LINEARLY ORDERED) and

3. $\left(\bigcup_{j\in\mathbb{Z}}V_j\right)^- = L^2_{\mathbb{R}}$ (DENSE in $L^2_{\mathbb{R}}$) and

4. $\dot{\mathbf{f}} \in \mathbf{V}_{j} \stackrel{'}{\Longleftrightarrow} \mathbf{D} \mathbf{f} \in \mathbf{V}_{j+1} \quad \forall j \in \mathbb{Z}, \mathbf{f} \in \mathbf{L}_{\mathbb{R}}^{2} \quad (\text{SELF-SIMILAR})$ 5. $\exists \phi \quad \text{such that} \quad \left\{ \mathbf{T}^{n} \phi | n \in \mathbb{Z} \right\} \quad is \ a \ \text{RIESZ BASIS} \quad for \ \mathbf{V}_{0}.$

A MULTIRESOLUTION ANALYSIS is also called an MRA.

An element V_j of $(V_j)_{j\in\mathbb{Z}}$ is a scaling subspace of the space $L^2_{\mathbb{R}}$. The pair $(L^2_{\mathbb{R}}, (V_j))$ is a multiresolution analysis space, or MRA space.

The function ϕ is the scaling function of the MRA space.

The traditional definition of the MRA also includes the following:

1.
$$f \in V_j \iff T^n f \in V_j \quad \forall n, j \in \mathbb{Z}, f \in L^2_{\mathbb{R}}$$
 (translation invariant)

1.
$$f \in V_j \iff T^n f \in V_j \quad \forall n, j \in \mathbb{Z}, f \in L^2_{\mathbb{R}}$$
 (translation invariant)
2. $\bigcap_{j \in \mathbb{Z}} V_j = \{\emptyset\}$ (greatest lower bound is **0**)


⟨entry 359⟩), which in turn means "the resolution or separation into component parts" (Black et al. (2009), http://original.com/parts | Black et al. (2009), http://original.com/parts //dictionary.reference.com/browse/dissolution)

⁴ Jawerth and Sweldens (1994) page 8

⁵ Hernández and Weiss (1996) page 44, Mallat (1999) page 221 ⟨Definition 7.1⟩, Mallat (1989) page 70, Meyer (1992) page 21 (Definition 2.2.1), <a> Christensen (2003) page 284 (Definition 13.1.1), <a> Bachman et al. (2002) 140 (Riesz basis: page 139)

However, Proposition 4.1 (next) demonstrates that both of these follow from the *MRA* as defined in Definition 4.1.

Proposition 4.1. ⁶



♦Proof: Proof for (1):

$$\mathbf{T}^n \mathbf{f} \in \mathbf{V}_j \\ \iff \mathbf{T}^n \mathbf{f} \in \operatorname{span} \left\{ \mathbf{D}^j \mathbf{T}^m \phi |_{m \in \mathbb{Z}} \right\} \\ \iff \exists \left((\alpha_n)_{n \in \mathbb{Z}} \right) \quad \text{such that} \quad \mathbf{T}^n \mathbf{f}(x) = \sum_{k \in \mathbb{Z}} \alpha_k \mathbf{D}^j \mathbf{T}^k \phi(x) \quad \text{by definition of } \{\phi\} \quad \text{(Definition 4.1 page 54)} \\ \iff \exists \left((\alpha_n)_{n \in \mathbb{Z}} \right) \quad \text{such that} \quad \mathbf{f}(x) = \mathbf{T}^{-n} \sum_{k \in \mathbb{Z}} \alpha_k \mathbf{D}^j \mathbf{T}^k \phi(x) \quad \text{by definition of } \mathbf{T} \quad \text{(Definition 3.3 page 40)} \\ = \sum_{k \in \mathbb{Z}} \alpha_k \mathbf{T}^{-n} \mathbf{D}^j \mathbf{T}^k \phi(x) \\ = \sum_{k \in \mathbb{Z}} \alpha_k \mathbf{D}^j \mathbf{T}^{k-2n} \phi(x) \quad \text{by } commutator \ relation} \quad \text{(Proposition 3.5 page 42)} \\ = \sum_{\ell \in \mathbb{Z}} \alpha_{\ell+2n} \mathbf{D}^j \mathbf{T}^\ell \phi(x) \quad \text{where } \ell \triangleq k-2n \implies k = \ell+2n \\ = \sum_{\ell \in \mathbb{Z}} \beta_\ell \mathbf{D}^j \mathbf{T}^\ell \phi(x) \quad \text{where } \beta_\ell \triangleq \alpha_{\ell+2n} \\ \iff \mathbf{f} \in \mathbf{V}_i \quad \text{by def. of } \{\mathbf{T}^n \phi\} \quad \text{(Definition 4.1 page 54)} \\ \end{cases}$$

Proof for (2):

- 1. Let \mathbf{P}_j be the *projection operator* that generates the scaling subspace \mathbf{V}_j such that $\mathbf{V}_j = \{\mathbf{P}_j \mathbf{f} | \mathbf{f} \in \mathbf{L}_{\mathbb{R}}^2\}$
- 2. lemma: Functions with *compact support* are *dense* in $L^2_{\mathbb{R}}$. Therefore, we only need to prove that the proposition is true for functions with support in [-R:R], for all R>0.
- 3. For some function $f \in \mathcal{L}^2_{\mathbb{R}}$, let $(f_n)_{n \in \mathbb{Z}}$ be a sequence of functions in $\mathcal{L}^2_{\mathbb{R}}$ with *compact support* such that $\operatorname{sppf}_n \subseteq [-R:R]$ for some R > 0 and $f(x) = \lim_{n \to \infty} (f_n(x))$.
- 4. lemma: $\bigcap V_j = \{0\}$ \iff $\lim_{j \to -\infty} ||P_j f|| = 0$ $\forall f \in \mathcal{L}^2_{\mathbb{R}}$. Proof:

$$\bigcap_{j\in\mathbb{Z}} \mathbf{V}_j = \bigcap_{j\in\mathbb{Z}} \left\{ \mathbf{P}_j \mathbf{f} | \mathbf{f} \in \mathbf{L}_{\mathbb{R}}^2 \right\} \qquad \text{by definition of } \mathbf{V}_j \qquad \text{(definition 1 page 55)}$$

$$= \lim_{j\to -\infty} \left\{ \mathbf{P}_j \mathbf{f} | \mathbf{f} \in \mathbf{L}_{\mathbb{R}}^2 \right\} \qquad \text{by definition of } \cap$$

$$= \mathbb{O} \iff \lim_{j\to -\infty} \left\| \mathbf{P}_j \mathbf{f} \right\| = 0 \qquad \text{by } \textit{nondegenerate} \text{ property of } \|\cdot\| \qquad \text{(Definition C.5 page 116)}$$

5. lemma: $\lim_{j \to -\infty} \|\mathbf{P}_j \mathbf{f}\| = 0 \quad \forall \mathbf{f} \in L_{\mathbb{R}}^2$. Proof:

⁶ ☐ Hernández and Weiss (1996) page 45 〈Theorem 1.6〉, ☐ Wojtaszczyk (1997) pages 19–28 〈Proposition 2.14〉, ☐ Pinsky (2002) pages 313–314 〈Lemma 6.4.28〉





Let $\mathbb{1}_{A(x)}$ be the *set indicator function* (Definition 3.2 page 40)

$$\lim_{j\to-\infty} \left\| \mathbf{P}_j \mathsf{f} \right\|^2$$

$$=\lim_{j\to-\infty}\left\|\mathbf{P}_{j}\lim_{n\to\infty}\left(\mathbf{f}_{n}\right)\right\|^{2}$$

$$\leq \lim_{j \to -\infty} B \sum_{n \in \mathbb{Z}} \left| \left\langle \mathbf{P}_{j} \lim_{n \to \infty} \left(\mathbf{f}_{n} \right) \mid \mathbf{D}^{j} \mathbf{T}^{n} \boldsymbol{\phi} \right\rangle \right|^{2}$$

$$= \lim_{j \to -\infty} B \sum_{n \in \mathbb{Z}} \left| \left\langle \lim_{n \to \infty} (f_n) \mid \mathbf{D}^j \mathbf{T}^n \phi \right\rangle \right|^2$$

$$= \lim_{j \to -\infty} B \sum_{n \in \mathbb{Z}} \left| \left\langle \mathbb{1}_{[-R:R]}(x) \lim_{n \to \infty} (f_n) \mid \mathbf{D}^j \mathbf{T}^n \phi(x) \right\rangle \right|^2$$

$$= \lim_{j \to -\infty} B \sum_{n \in \mathbb{Z}} \left| \left\langle \lim_{n \to \infty} \left(\mathsf{f}_n \right) \mid \mathbb{1}_{[-R:R]}(x) \mathbf{D}^j \mathbf{T}^n \phi(x) \right\rangle \right|^2$$

$$\leq \lim_{j \to -\infty} B \sum_{n \in \mathbb{Z}} \left\| \lim_{n \to \infty} (f_n) \right\|^2 \left\| \mathbb{1}_{[-R:R]}(x) \mathbf{D}^j \mathbf{T}^n \phi(x) \right\|^2$$

$$= \lim_{j \to -\infty} B \sum_{n \in \mathbb{Z}} \|\mathbf{f}\|^2 \|\mathbb{1}_{[-R:R]}(x) \mathbf{D}^j \mathbf{T}^n \phi(x)\|^2$$

$$= \lim_{j \to -\infty} B \sum_{n \in \mathbb{Z}} \|\mathbf{f}\|^2 \left\| \left[\underbrace{\mathbf{D}^j \mathbf{D}^{-j}}_{\mathbf{I}} \mathbb{1}_{[-R:R]}(x) \right] \left[\mathbf{D}^j \mathbf{T}^n \phi(x) \right] \right\|^2$$

$$= \lim_{j \to -\infty} B \sum_{n \in \mathbb{Z}} \|\mathbf{f}\|^2 \|2^{j/2} \mathbf{D}^j \{ \left[\mathbf{D}^{-j} \mathbb{1}_{[-R:R]}(x) \right] \left[\mathbf{T}^n \phi(x) \right] \} \|^2$$

$$= \lim_{j \to -\infty} B \sum_{n \in \mathbb{Z}} \|\mathbf{f}\|^2 \|\mathbf{D}^j \left\{ 2^{j/2} 2^{-j/2} \mathbb{1}_{[-R:R]} (2^{-j} x) \left[\mathbf{T}^n \phi(x) \right] \right\} \|^2$$

$$=\lim_{j\to-\infty}B\sum_{n\in\mathbb{Z}}\|\mathsf{f}\|^2\left\|\mathbf{D}^j\left\{\left[\underbrace{\mathbf{T}^n\mathbf{T}^{-n}}_{\mathsf{I}_{[-R:R]}}(2^{-j}x)\right]\left[\mathbf{T}^n\phi(x)\right]\right\}\right\|^2$$

$$= \lim_{j \to -\infty} B \sum_{n \in \mathbb{Z}} \|\mathbf{f}\|^2 \|\mathbf{D}^j \left\{ \left[\mathbf{T}^n \mathbb{1}_{[-R:R]} (2^{-j} x + n) \right] \left[\mathbf{T}^n \phi(x) \right] \right\} \|^2$$

$$= \lim_{j \to -\infty} B \sum_{n \in \mathbb{Z}} \|\mathbf{f}\|^2 \|\mathbf{D}^j \mathbf{T}^n \{ \mathbb{1}_{[-R:R]} (2^{-j} x + n) \phi(x) \} \|^2$$

$$= \lim_{j \to -\infty} B \sum_{n \in \mathbb{Z}} \|f\|^2 \|\mathbb{1}_{[-R:R]} (2^{-j} x + n) \phi(x)\|^2$$

$$= B \|f\|^2 \sum_{n \in \mathbb{Z}} \lim_{j \to -\infty} \|\mathbb{1}_{[-2^j R + n: 2^j R + n]}(u) \phi(2^{-j} (u - n))\|^2$$

$$= B \|f\|^2 \sum_{j \to -\infty} \lim_{j \to -\infty} \int_{-2j}^{2^j R + n} |\phi(2^{-j}(u - n))|^2 du$$

$$= B \|\mathbf{f}\|^2 \sum_{n \in \mathbb{Z}} \int_n^n |\phi(0)|^2 \ \mathrm{d}\mathbf{u}$$

=0

by definition 3 page 55

by frame property (Proposition 2.5 page 30)

by definition of P_j (definition 1 page 55)

by definition of (f_n) (definition 3 page 55)

prop. of $\langle \triangle \mid \bigtriangledown \rangle$ in $\mathcal{L}^2_{\mathbb{R}}$ (Definition D.1 page 141)

by CS Inequality

by definition of (f_n) (definition 3 page 55)

by property of D (Proposition 3.2 page 41)

by Proposition 3.4 page 41

by property of **D** (Proposition 3.2 page 41)

by property of **T** (Proposition 3.2 page 41)

by property of T (Proposition 3.2 page 41)

by property of ${\bf D}$ (Proposition 3.2 page 41)

by *unitary* prop. (Theorem 3.1 page 45)

 $u \triangleq 2^{j}x + n \implies x = 2^{-j}(u - n)$

6. Final step in proof that $\bigcap V_j = \{0\}$: by (4) lemma page 55 and (5) lemma page 56

4.2. DEFINITION Daniel J. Greenhoe page 57

Proposition 4.2. ⁷

$$\left\{ \begin{array}{l} \text{(1).} \quad \left(\mathbf{T}^n \phi \right) \text{ is a Riesz sequence} \quad \text{and} \\ \text{(2).} \quad \tilde{\phi}(\omega) \text{ is continuous at } 0 \quad \text{ and} \\ \text{(3).} \quad \tilde{\phi}(0) \neq 0 \end{array} \right\} \Longrightarrow \left\{ \left(\bigcup_{j \in \mathbb{Z}} \mathbf{V}_j \right)^- = \mathbf{L}_{\mathbb{R}}^2 \quad (\text{dense in } \mathbf{L}_{\mathbb{R}}^2) \right\}$$

♥Proof:

- 1. Let \mathbf{P}_j be the *projection operator* that generates the scaling subspace \mathbf{V}_j such that $\mathbf{V}_i = \{\mathbf{P}_j \mathbf{f} | \mathbf{f} \in \mathbf{H}\}$
- 2. definition: Choose $f \in \mathcal{L}^2_{\mathbb{R}}$ such that $f \perp \bigcup_{j \in \mathbb{Z}} V_j$. Let $\tilde{f}(\omega)$ be the *Fourier Transform* (Definition I.2 page 196) of f(x).
- 3. lemma: The function f (definition 2 page 57) *exists* because the set of functions that can be chosen to be f at least contains 0 (it is not the emptyset). Proof:

$$f(x) = 0 \implies \left\langle f \mid \left\{ h \in L_{\mathbb{R}}^{2} \mid h \in \bigcup_{j \in \mathbb{Z}} V_{j} \right\} \right\rangle$$

$$= \left\langle 0 \mid \left\{ h \in L_{\mathbb{R}}^{2} \mid h \in \bigcup_{j \in \mathbb{Z}} V_{j} \right\} \right\rangle$$

$$= 0$$

$$\implies f \perp \bigcup_{j \in \mathbb{Z}} V_{j}$$

$$\implies f \text{ exists}$$

4. lemma: $\|\mathbf{P}_{j}\mathbf{f}\| = 0 \quad \forall j \in \mathbb{Z}$. Proof:

$$\|\mathbf{P}_{f}\| = \|0\|$$
 by definition of f (definition 2 page 57)
= 0 by *nondegenerate* property of $\|\cdot\|$

- 5. definition: Choose some function $g \in L^2_{\mathbb{R}}$ such that $\tilde{g}(\omega) = \tilde{f}(\omega)\mathbb{1}_{[-R:R]}$ (Definition 3.2 page 40) for some R > 0 and such that $\|f g\| < \varepsilon$. Let $\tilde{g}(\omega)$ be the *Fourier Transform* (Definition 1.2 page 196) of g(x).
- 6. lemma: The function g (definition 5 page 57) exists. Proof: For some (possibly very large) R,

$$\varepsilon > \|\tilde{\mathbf{f}}(\omega) - \tilde{\mathbf{g}}(\omega)\| \qquad \text{by definition of g} \qquad \text{(definition 5 page 57)}$$

$$= \|\tilde{\mathbf{F}}\mathbf{f}(x) - \tilde{\mathbf{F}}\mathbf{g}(x)\| \qquad \text{by definition of } \tilde{\mathbf{f}} \text{ and } \tilde{\mathbf{g}} \qquad \text{(definition 2 page 57), (definition 5 page 57)}$$

$$= \|\tilde{\mathbf{F}}\big[\mathbf{f}(x) - \mathbf{g}(x)\big]\| \qquad \text{by } \text{linearity of } \tilde{\mathbf{F}} \qquad \text{(Definition C.4 page 113)}$$

$$= \|\mathbf{f}(x) - \mathbf{g}(x)\| \qquad \text{by } \text{unitary property of } \tilde{\mathbf{F}} \qquad \text{(Theorem I.2 page 197)}$$

$$\implies \mathbf{g} \text{ exists} \qquad \text{because it's possible to satisfy definition 5 page 57}$$

7. lemma: $\|\mathbf{P}_{j}\mathbf{g}\| < \varepsilon \quad \forall j \in \mathbb{Z}$ for sufficiently large R. Proof:

$\varepsilon > \ f - g\ $	by definition of g	(definition 5 page 57)
$\geq \ \mathbf{P}_{j}[f-g]\ $	by property of projection operators	(Definition C.10 page 127)
$= \left\ \mathbf{P}_{j} f - \mathbf{P}_{j} g \right\ $	by <i>additive</i> property of \mathbf{P}_{j}	(Definition C.4 page 113)
$\geq \left \left\ \mathbf{P}_{j} f \right\ - \left\ \mathbf{P}_{j} g \right\ \right $	by Reverse Triangle Inequality	
$= \left 0 - \left\ \mathbf{P}_{j} \mathbf{g} \right\ \right $	by ((4) lemma page 57)	
$= \ \mathbf{P}_j \mathbf{g}\ $	by <i>strictly positive</i> property of $\ \cdot\ $	(Definition C.5 page 116)

Wojtaszczyk (1997) pages 28–31 (Proposition 2.15)



8. lemma: g = 0. Proof:

$$\begin{aligned} 0 &= \lim_{j \to \infty} \left\| \mathbf{P}_j \mathbf{g} \right\|^2 & \text{by (7) lemma page 57} \\ &\geq \lim_{j \to \infty} A \sum_{n \in \mathbb{Z}} \left| \left\langle \mathbf{P}_j \mathbf{g} \mid \mathbf{D}^j \mathbf{T}^n \phi \right\rangle \right|^2 & \text{by frame property} & \text{(Proposition 2.5 page 30)} \\ &= \lim_{j \to \infty} A \sum_{n \in \mathbb{Z}} \left| \left\langle \mathbf{g} \mid \mathbf{D}^j \mathbf{T}^n \phi \right\rangle \right|^2 & \text{by definition of } \mathbf{P}_j & \text{(item (1) page 57)} \\ &= \lim_{j \to \infty} A \sum_{n \in \mathbb{Z}} \left| \left\langle \mathbf{\tilde{F}} \mathbf{g} \mid \mathbf{\tilde{F}} \mathbf{D}^j \mathbf{T}^n \phi \right\rangle \right|^2 & \text{by unitary property of } \mathbf{\tilde{F}} & \text{(Theorem I.2 page 197)} \\ &= \lim_{j \to \infty} A \sum_{n \in \mathbb{Z}} \left| \left\langle \mathbf{\tilde{g}}(\omega) \mid 2^{-j/2} e^{-i2^{-j}\omega n} \tilde{\phi}(2^{-j}\omega) \right\rangle \right|^2 & \text{by Proposition 3.12 page 47} \\ &= \lim_{j \to \infty} A \sum_{n \in \mathbb{Z}} \left| \left\langle \mathbf{\tilde{g}}(\omega) \tilde{\phi}^* (2^{-j}\omega) \mid 2^{-j/2} e^{-i2^{-j}\omega n} \right\rangle \right|^2 & \text{by property of } \left\langle \triangle \mid \nabla \right\rangle \text{ in } \boldsymbol{L}_{\mathbb{R}}^2 \\ &= \lim_{j \to \infty} A \left\| \mathbf{\tilde{g}}(\omega) \tilde{\phi}^* (2^{-j}\omega) \right\|^2 & \text{by Parseval's Identity} & \text{(Theorem 2.9 page 22)} \\ &= A \left\| \mathbf{\tilde{g}}(\omega) \tilde{\phi}^* (0) \right\|^2 & \text{by left hypothesis (2)} \\ &= A \left\| \tilde{\phi}^* (0) \right\|^2 \left\| \mathbf{\tilde{g}}(\omega) \right\|^2 & \text{by unitary property of } \mathbf{\tilde{F}} & \text{(Theorem I.2 page 197)} \\ &\Rightarrow \|\mathbf{g}\| = 0 & \text{by unitary property of } \mathbf{\tilde{F}} & \text{(Theorem I.2 page 197)} \\ &\Rightarrow \|\mathbf{g}\| = 0 & \text{by left hypothesis (3)} \\ &\Leftrightarrow \mathbf{g} = 0 & \text{by nondegenerate property of } \| \cdot \| \end{aligned}$$

9. Final step in proof that $\left(\bigcup_{i\in\mathbb{Z}}V_i\right)^-=L_{\mathbb{R}}^2$:

$$g = 0$$

$$\Rightarrow f = 0$$

$$\Rightarrow \left(\bigcup_{j \in \mathbb{Z}} V_j\right)^- = L_{\mathbb{R}}^2$$
by (8) lemma page 58
by definition of g
(definition 5 page 57)

Definition 4.1 defines an MRA on the space $L_{\mathbb{R}}^2$, which is a special case of a *separable Hilbert space*. A Hilbert space is a *linear space* that is equipped with an *inner product*, is *complete* with respect to the *metric* induced by the inner product, and contains a subset that is *dense* in $L_{\mathbb{R}}^2$.

An *inner product* on a linear space endows the linear space with a *topology*. The sum such as $\sum_{n=1}^{N} \alpha_n f_n$ is finite and thus suitable for a finite linear space only. An infinite space requires an infinite sum $\sum_{n=1}^{\infty} \alpha_n \phi_n$, and an infinite sum is defined in terms of a limit. The limit, in turn, is defined in terms of a *topology*. The *inner product* induces a *norm* (Definition C.5 page 116) which induces a *metric* which induces a topology.

Definition 4.1 defines each subspace V_j to be *closed* ($V_j = V_j^-$) in $L^2_{\mathbb{R}}$. As one might imagine, the properties of *completeness* and *closure* are closely related. Moreover, Every *complete* sequence is also *bounded*, and so each subspace V_j is *bounded* as well.

Proposition 4.3. Let $(L^2_{\mathbb{R}}, (V_j))$ be an MRA space.

Each subspace V_j is COMPLETE.




- 1. By definition Definition 4.1, $\boldsymbol{L}_{\mathbb{R}}^2$ is complete.
- 2. In any metric space, (which includes all inner product spaces such as $L_{\mathbb{R}}^2$), a *closed* subspace of a *complete* metric space is itself also *complete*.
- 3. In any *complete* metric space X (which includes all Hilbert spaces such as $L_{\mathbb{R}}^2$), the two properties coincide—that is, a subspace is complete *if and only if* it is closed in the space X.
- 4. So because $L_{\mathbb{R}}^2$ is *complete* and each V_j is *closed*, then each V_j is also *complete*.

₽

4.3 Order structure

A multiresolution analysis (Definition 4.1 page 54) together with the set inclusion relation \subseteq forms the *linearly ordered set* $((V_j), \subseteq)$, illustrated to the right by a *Hasse diagram*. Subspaces V_j increase in "size" with increasing j. That is, they contain more and more vectors (functions) for larger and larger j—with the upper limit of this sequence being $L^2_{\mathbb{R}}$. Alternatively, we can say that approximation within a subspace V_j yields greater "resolution" for increasing j.



The $least\ upper\ bound\ (l.u.b.)$ of the linearly ordered set $\left(\left(\begin{pmatrix} \textit{\textbf{V}}_j \end{pmatrix}\right), \subseteq\right)$ is $\textit{\textbf{L}}^2_{\mathbb{R}}$ (Definition 4.1 page 54):

$$\left(\bigcup_{j\in\mathbb{Z}} \mathbf{V}_j\right)^{-} = \mathbf{L}_{\mathbb{R}}^2$$

The greatest lower bound (g.l.b.) of the linearly ordered set $((V_j), \subseteq)$ is $\mathbf{0}$ (Proposition 4.1 page 55): $\bigcap_{i=1}^n V_j = \mathbf{0}$.

All linear subspaces contain the zero vector (Proposition B.3 page 99). So the intersection of any two subspaces must at least contain \mathbb{O} . If the intersection of any two linear subspaces X and Y is exactly $\{\mathbb{O}\}$, then for any vector in the sum of those subspaces $\{u \in X + Y\}$ there are **unique** vectors $f \in X$ and $g \in Y$ such that u = f + g. This is *not* necessarily true if the intersection contains more than just $\{\mathbb{O}\}$ (Theorem B.1 page 101).

4.4 Dilation equation

Several functions in mathematics exhibit a kind of self-similar or recursive property:

- If a function f(x) is *linear*, then (Example 3.2 page 50) f(x) = f(1)x f(0)Tx.



$$f(x) = \sum_{n=1}^{\infty} f(n) \mathbf{T}^n \frac{\sin[\pi(x)]}{\pi(x)}.$$

B-splines are another example:
$$\mathsf{N}_n(x) = \frac{1}{n} x \mathsf{N}_{n-1}(x) - \frac{1}{n} x \mathbf{T} \mathsf{N}_{n-1}(x) + \frac{n+1}{n} \mathbf{T} \mathsf{N}_{n-1}(x) \qquad \forall n \in \mathbb{N} \setminus \{1\}, \forall x \in \mathbb{R}.$$

The scaling function $\phi(x)$ (Definition 4.1 page 54) also exhibits a kind of *self-similar* property. By Definition 4.1 page 54, the dilation **D**f of each vector f in V_0 is in V_1 . If $\{\mathbf{T}^n \phi | n \in \mathbb{Z}\}$ is a basis for V_0 , then $\{\mathbf{DT}^n\phi|_{n\in\mathbb{Z}}\}\$ is a basis for V_1 , $\{\mathbf{D}^2\mathbf{T}^n\phi|_{n\in\mathbb{Z}}\}$ is a basis for V_2 , ...; and in general $\{\mathbf{D}^j\mathbf{T}^m\phi|_{j\in\mathbb{Z}}\}$ is a basis for V_j . Also, if ϕ is in V_0 , then it is also in V_1 (because $V_0\subset V_1$). And because ϕ is in V_1 and because $\{\mathbf{DT}^n \phi | n \in \mathbb{Z}\}$ is a basis for V_1 , ϕ is a linear combination of the elements in $\{\mathbf{DT}^n \phi | n \in \mathbb{Z}\}$. That is, ϕ can be represented as a linear combination of translated and dilated versions of itself. The resulting equation is called the *dilation equation* (Definition 4.2, next).⁸

Definition 4.2. 9 Let $(L^2_{\mathbb{R}}, (V_i))$ be a multiresolution analysis space with scaling function ϕ (Def-

D E F

inition 4.1 page 54). Let $(h_n)_{n\in\mathbb{Z}}$ be a SEQUENCE (Definition J.1 page 207) in $\mathscr{C}^2_{\mathbb{R}}$ (Definition J.2 page 207).

The EQUATION $\left\{\phi(x) = \sum_{n\in\mathbb{Z}} h_n \mathbf{DT}^n \phi(x) \quad \forall x \in \mathbb{R}\right\}$ is called the **dilation equation**.

It is also called the **refinement equation**, **two-scale difference equation**, and two-scale relation.

Remark 4.1.

The dilation equation under the defintions of T and D evaluates to R E $\phi(x) = \sum_{n \in \mathbb{Z}} \mathsf{h}_n \phi(2x - n).$

^ℚProof:

$$\phi(x) = \sum_{n \in \mathbb{Z}} \mathsf{h}_n \mathbf{D} \mathbf{T}^n \phi(x)$$

$$= \sum_{n \in \mathbb{Z}} \mathsf{h}_n \mathbf{D} \phi(x - n) \qquad \text{by definition of } \mathbf{T} \qquad \text{(Definition 3.3 page 40)}$$

$$= \sum_{n \in \mathbb{Z}} \mathsf{h}_n \phi(2x - n) \qquad \text{by definition of } \mathbf{D} \qquad \text{(Definition 3.3 page 40)}$$

Theorem 4.1 (dilation equation). Let an MRA SPACE and SCALING FUNCTION be as defined in Definition 4.1 page 54.

 $\exists (h_n)_{n \in \mathbb{Z}} \text{ such that}$ $\phi(x) = \sum_{n \in \mathbb{Z}} h_n \mathbf{DT}^n \phi(x)$ $(L_{\mathbb{R}}^2, (V_j))$ is an MRA space $\{uith \text{ scaling function } \phi\}$ DILATION EQUATION IN "TIME"

Jawerth and Sweldens (1994) page 7



⁸The property of *translation invariance* is of particular significance in the theory of *normed linear spaces* (a Hilbert space is a complete normed linear space equipped with an inner product).

№PROOF:

$$\phi \in V_0 \qquad \qquad \text{by definition of MRA} \qquad \text{(Definition 4.1 page 54)}$$

$$\subseteq V_1 \qquad \qquad \text{by definition of MRA} \qquad \text{(Definition 4.1 page 54)}$$

$$\triangleq \operatorname{span} \left\{ \left. \mathbf{DT}^n \phi(x) \right|_{n \in \mathbb{Z}} \right\} \qquad \qquad \operatorname{by definition of } V_j \qquad \text{(Definition 4.1 page 54)}$$

$$\Longrightarrow \exists \left. \left(\mathbf{h}_n \right)_{n \in \mathbb{Z}} \quad \text{such that} \quad \phi(x) = \sum_{n \in \mathbb{Z}} \mathbf{h}_n \mathbf{DT}^n \phi(x) \qquad \text{by definition of span} \qquad \text{(Definition 2.2 page 9)}$$

Lemma 4.1. ¹⁰ Let $\phi(x)$ be a function in $L^2_{\mathbb{R}}$ (Definition D.1 page 141). Let $\tilde{\phi}(\omega)$ be the Fourier transform (Definition I.2 page 196) of $\phi(x)$. Let $\check{\mathsf{h}}(\omega)$ be the DISCRETE TIME FOURIER TRANSFORM (Definition M.1 page 237) of a sequence $(\mathsf{h}_n)_{n\in\mathbb{Z}}$.

$$\frac{dente \left(\prod_{n} J_{n \in \mathbb{Z}} \right)}{(A)} \quad \phi(x) = \sum_{n \in \mathbb{Z}} h_{n} \mathbf{D} \mathbf{T}^{n} \phi(x) \quad \forall x \in \mathbb{R} \quad \Longleftrightarrow \quad \tilde{\phi}(\omega) = \frac{\sqrt{2}}{2} \, \check{\mathsf{h}} \left(\frac{\omega}{2} \right) \, \tilde{\phi} \left(\frac{\omega}{2} \right) \qquad \forall \omega \in \mathbb{R} \qquad (1)$$

$$\Leftrightarrow \quad \tilde{\phi}(\omega) = \tilde{\phi} \left(\frac{\omega}{2^{N}} \right) \prod_{n=1}^{N} \frac{\sqrt{2}}{2} \, \check{\mathsf{h}} \left(\frac{\omega}{2^{n}} \right) \quad \forall n \in \mathbb{N}, \omega \in \mathbb{R} \quad (2)$$

♥Proof:

1. Proof that (A) \Longrightarrow (1):

$$\begin{split} \tilde{\phi}(\omega) &\triangleq \tilde{\mathbf{F}} \phi \\ &= \tilde{\mathbf{F}} \sum_{n \in \mathbb{Z}} \mathsf{h}_n \mathbf{D} \mathbf{T}^n \phi(x) \\ &= \sum_{n \in \mathbb{Z}} \mathsf{h}_n \tilde{\mathbf{F}} \mathbf{D} \mathbf{T}^n \phi(x) \\ &= \sum_{n \in \mathbb{Z}} \mathsf{h}_n \frac{\sqrt{2}}{2} e^{-i\frac{\omega}{2}n} \phi\left(\frac{\omega}{2}\right) \\ &= \sum_{n \in \mathbb{Z}} \mathsf{h}_n \frac{\sqrt{2}}{2} e^{-i\frac{\omega}{2}n} \phi\left(\frac{\omega}{2}\right) \\ &= \frac{\sqrt{2}}{2} \left[\sum_{n \in \mathbb{Z}} \mathsf{h}_n e^{-i\frac{\omega}{2}n} \right] \tilde{\phi}\left(\frac{\omega}{2}\right) \\ &= \frac{\sqrt{2}}{2} \, \check{\mathsf{h}}\left(\frac{\omega}{2}\right) \, \tilde{\phi}\left(\frac{\omega}{2}\right) \end{split}$$
 by definition of *DTFT* (Definition M.1 page 237)

2. Proof that (A) \Leftarrow (1):

$$\begin{aligned} \phi(x) &= \tilde{\mathbf{F}}^{-1} \tilde{\phi}(\omega) & \text{by definition of } \tilde{\phi}(\omega) \\ &= \tilde{\mathbf{F}}^{-1} \frac{\sqrt{2}}{2} \, \check{\mathsf{h}} \Big(\frac{\omega}{2} \Big) \, \tilde{\phi} \Big(\frac{\omega}{2} \Big) & \text{by (1)} \\ &= \tilde{\mathbf{F}}^{-1} \frac{\sqrt{2}}{2} \, \sum_{n \in \mathbb{Z}} h_n e^{-i\frac{\omega}{2}n} \, \tilde{\phi} \Big(\frac{\omega}{2} \Big) & \text{by definition of } DTFT \\ &= \frac{\sqrt{2}}{2} \, \sum_{n \in \mathbb{Z}} h_n \tilde{\mathbf{F}}^{-1} e^{-i\frac{\omega}{2}n} \, \tilde{\phi} \Big(\frac{\omega}{2} \Big) & \text{by property of linear operators} \\ &= \frac{\sqrt{2}}{2} \, \sum_{n \in \mathbb{Z}} h_n \tilde{\mathbf{F}}^{-1} \tilde{\mathbf{F}} \mathbf{D} \mathbf{T}^n \phi & \text{by Proposition 3.12 page 47} \\ &= \sum_{n \in \mathbb{Z}} \mathsf{h}_n \mathbf{D} \mathbf{T}^n \phi(x) & \text{by definition of } operator inverse \end{aligned}$$



¹⁰ Mallat (1999) page 228

- 3. Proof that $(1) \Longrightarrow (2)$:
 - (a) Proof for N = 1 case:

$$\begin{split} \tilde{\phi}\left(\frac{\omega}{2^{N}}\right) \left. \prod_{n=1}^{N} \frac{\sqrt{2}}{2} \check{\mathsf{h}}\left(\frac{\omega}{2^{n}}\right) \right|_{N=1} &= \frac{\sqrt{2}}{2} \, \check{\mathsf{h}}\left(\frac{\omega}{2}\right) \tilde{\phi}\left(\frac{\omega}{2}\right) \\ &= \tilde{\phi}(\omega) \end{split} \qquad \qquad \text{by (1)}$$

(b) Proof that $[N \text{ case}] \Longrightarrow [N+1 \text{ case}]$:

$$\begin{split} \tilde{\phi}\Big(\frac{\omega}{2^{N+1}}\Big) & \prod_{n=1}^{N+1} \frac{\sqrt{2}}{2} \check{\mathsf{h}}\Big(\frac{\omega}{2^n}\Big) = \left[\prod_{n=1}^N \frac{\sqrt{2}}{2} \check{\mathsf{h}}\Big(\frac{\omega}{2^n}\Big)\right] \underbrace{\frac{\sqrt{2}}{2} \check{\mathsf{h}}\Big(\frac{\omega}{2^{N+1}}\Big)}_{\tilde{\phi}(\omega/2^N)} \\ & = \tilde{\phi}(\omega/2^N) \prod_{n=1}^N \frac{\sqrt{2}}{2} \check{\mathsf{h}}\Big(\frac{\omega}{2^n}\Big) \\ & = \tilde{\phi}(\omega) \end{split} \qquad \qquad \text{by [N case] hypothesis} \end{split}$$

4. Proof that $(1) \Leftarrow (2)$:

$$\tilde{\phi}(\omega) = \tilde{\phi}\left(\frac{\omega}{2^N}\right) \prod_{n=1}^N \frac{\sqrt{2}}{2} \check{h}\left(\frac{\omega}{2^n}\right) \bigg|_{N=1}$$

$$= \tilde{\phi}\left(\frac{\omega}{2}\right) \frac{\sqrt{2}}{2} \check{h}\left(\frac{\omega}{2}\right)$$

$$= \frac{\sqrt{2}}{2} \check{h}\left(\frac{\omega}{2}\right) \tilde{\phi}\left(\frac{\omega}{2}\right)$$
by (2)

Lemma 4.2. Let $\phi(x)$ be a function in $\mathbf{L}^2_{\mathbb{R}}$ (Definition D.1 page 141). Let $\tilde{\phi}(\omega)$ be the Fourier transform (Definition 1.2 page 196) of $\phi(x)$. Let $\check{\mathsf{h}}(\omega)$ be the DISCRETE TIME FOURIER TRANSFORM (Definition M.1 page 237) of (h_n) . Let $\prod_{n=1}^\infty x_n \triangleq \lim_{N\to\infty} \prod_{n=1}^N x_n$, with respect to the standard norm in $\mathbf{L}^2_{\mathbb{R}}$.

$$\left\{
\begin{array}{l}
\tilde{\phi}(\omega) = C \prod_{n=1}^{\infty} \frac{\sqrt{2}}{2} \check{h}\left(\frac{\omega}{2^{n}}\right) \\
\forall C > 0, \omega \in \mathbb{R}
\end{array}
\right\} \implies \phi(x) = \sum_{n \in \mathbb{Z}} h_{n} \mathbf{D} \mathbf{T}^{n} \phi(x) \qquad \forall x \in \mathbb{R}$$

$$\Leftrightarrow \tilde{\phi}(\omega) = \frac{\sqrt{2}}{2} \check{h}\left(\frac{\omega}{2}\right) \tilde{\phi}\left(\frac{\omega}{2}\right) \qquad \forall \omega \in \mathbb{R}$$

$$\Leftrightarrow \tilde{\phi}(\omega) = \tilde{\phi}\left(\frac{\omega}{2^{N}}\right) \prod_{n=1}^{N} \frac{\sqrt{2}}{2} \check{h}\left(\frac{\omega}{2^{n}}\right) \qquad \forall n \in \mathbb{N}, \omega \in \mathbb{R}$$

$$(1)$$

♥Proof:

1. Proof that (1) \iff (2) \iff (3): by Lemma 4.1 page 61

₽

2. Proof that (A) \Longrightarrow (2):

$$\begin{split} \tilde{\phi}(\omega) &= C \prod_{n=1}^{\infty} \frac{\sqrt{2}}{2} \check{\mathsf{h}} \left(\frac{\omega}{2^n} \right) \\ &= C \frac{\sqrt{2}}{2} \check{\mathsf{h}} \left(\frac{\omega}{2} \right) \prod_{n=1}^{\infty} \frac{\sqrt{2}}{2} \check{\mathsf{h}} \left(\frac{\omega}{2^{n+1}} \right) \\ &= C \frac{\sqrt{2}}{2} \check{\mathsf{h}} \left(\frac{\omega}{2} \right) \prod_{n=1}^{\infty} \frac{\sqrt{2}}{2} \check{\mathsf{h}} \left(\frac{\omega/2}{2^n} \right) \\ &= \frac{\sqrt{2}}{2} \check{\mathsf{h}} \left(\frac{\omega}{2} \right) \left[C \prod_{n=1}^{\infty} \frac{\sqrt{2}}{2} \check{\mathsf{h}} \left(\frac{\omega/2}{2^n} \right) \right] \\ &= \frac{\sqrt{2}}{2} \check{\mathsf{h}} \left(\frac{\omega}{2} \right) \tilde{\phi} \left(\frac{\omega}{2} \right) \end{split}$$

by left hypothesis

by left hypothesis

Proposition 4.4. Let $\phi(x)$ be a function in $L^2_{\mathbb{R}}$ (Definition D.1 page 141). Let $\tilde{\phi}(\omega)$ be the Fourier transform (Definition I.2 page 196) of $\phi(x)$. Let $\check{h}(\omega)$ be the DISCRETE TIME FOURIER TRANSFORM (Definition M.1 page 237) of $((\mathsf{h}_n)).\ Let \prod_{n=1} x_n \triangleq \lim_{N \to \infty} \prod_{n=1} x_n, \ with \ respect \ to \ the \ standard \ norm \ in \ \boldsymbol{L}^2_{\mathbb{R}}.$

$$\begin{cases} \tilde{\phi}(\omega) \text{ is } \\ \text{continuous } \\ \text{at } \omega = 0 \end{cases} \Rightarrow \begin{cases} \tilde{\phi}(\omega) = \sum_{n \in \mathbb{Z}} h_n DT^n \phi(x) & \forall x \in \mathbb{R} \\ \Leftrightarrow \tilde{\phi}(\omega) = \frac{\sqrt{2}}{2} \check{h} \left(\frac{\omega}{2}\right) \tilde{\phi} \left(\frac{\omega}{2}\right) & \forall \omega \in \mathbb{R} \\ \Leftrightarrow \tilde{\phi}(\omega) = \tilde{\phi} \left(\frac{\omega}{2^N}\right) \prod_{n=1}^N \frac{\sqrt{2}}{2} \check{h} \left(\frac{\omega}{2^n}\right) & \forall n \in \mathbb{N}, \omega \in \mathbb{R} \end{cases}$$

$$\Leftrightarrow \tilde{\phi}(\omega) = \tilde{\phi}(0) \prod_{n=1}^\infty \frac{\sqrt{2}}{2} \check{h} \left(\frac{\omega}{2^n}\right) & \omega \in \mathbb{R} \end{cases}$$

$$\Leftrightarrow \tilde{\phi}(\omega) = \tilde{\phi}(0) \prod_{n=1}^\infty \frac{\sqrt{2}}{2} \check{h} \left(\frac{\omega}{2^n}\right) & \omega \in \mathbb{R} \end{cases}$$

$$(1)$$

[♠]Proof:

- 1. Proof that (1) \iff (2) \iff (3): by Lemma 4.1 page 61
- 2. Proof that $(3) \Longrightarrow (4)$:

$$\tilde{\phi}(0) \prod_{n=1}^{\infty} \frac{\sqrt{2}}{2} \check{h} \left(\frac{\omega}{2^n}\right) = \lim_{N \to \infty} \tilde{\phi}\left(\frac{\omega}{2^N}\right) \prod_{n=1}^{N} \frac{\sqrt{2}}{2} \check{h} \left(\frac{\omega}{2^n}\right)$$
 by *continuity* and definition of $\prod_{n=1}^{\infty} x_n$ by (3) and Lemma 4.1 page 61

3. Proof that (2) \Leftarrow (4): by Lemma 4.2 page 62


Definition 4.3 (next) formally defines the coefficients that appear in Theorem 4.1 (page 60).

Definition 4.3. Let $(L^2_{\mathbb{R}}, (V_j))$ be a multiresolution analysis space with scaling function ϕ . Let $(h_n)_{n \in \mathbb{Z}}$

be a sequence of coefficients such that $\phi = \sum_{n \in \mathbb{Z}} h_n \mathbf{D} \mathbf{T}^n \phi$.

A multiresolution system is the tuple $(L_{\mathbb{R}}^2, (V_j), \phi, (h_n))$. The sequence $(h_n)_{n \in \mathbb{Z}}$ is the scaling coefficient sequence. A multiresolution system is also called an MRA system. An MRA system is an **orthonormal MRA system** if $\{ \mathbf{T}^n \phi | n \in \mathbb{Z} \}$ is Orthonormal.

Theorem 4.2. Let $(L_{\mathbb{R}}^2, (V_j), \phi, (h_n))$ be an MRA SYSTEM (Definition 4.3 page 63). Let $\operatorname{span} A$ be the LINEAR SPAN (Definition 2.2 page 9) of a set A.



[♠]Proof: Proof is by induction: 11

1. induction basis (proof for j = 0 case):


2. induction step (proof that j case $\implies j + 1$ case):

$$\begin{split} & \operatorname{sgn}\left\{\mathbf{D}^{j+1}\mathbf{T}^{n}\phi \Big| \ n\in\mathbb{Z}\right\} \\ & = \left\{f\in L_{\mathbb{R}}^{2}|\exists \left(\alpha_{n}\right) \quad \text{such that} \quad f(x) = \sum_{n\in\mathbb{Z}}\alpha_{n}\mathbf{D}^{j+1}\mathbf{T}^{n}\phi\right\} \quad \text{by definition of span} \qquad \text{(Definition 2.2 page 9)} \\ & = \left\{f\in L_{\mathbb{R}}^{2}|\exists \left(\alpha_{n}\right) \quad \text{such that} \quad f(x) = \mathbf{D}\sum_{n\in\mathbb{Z}}\alpha_{n}\mathbf{D}^{j}\mathbf{T}^{n}\phi\right\} \\ & = \left\{f\in L_{\mathbb{R}}^{2}|\exists \left(\alpha_{n}\right) \quad \text{such that} \quad \mathbf{D}^{-1}[\mathbf{D}f(x)] = \sum_{n\in\mathbb{Z}}\alpha_{n}\mathbf{D}^{j}\mathbf{T}^{n}\phi\right\} \\ & = \mathbf{D}\left\{f\in L_{\mathbb{R}}^{2}|\exists \left(\alpha_{n}\right) \quad \text{such that} \quad f(x) = \sum_{n\in\mathbb{Z}}\alpha_{n}\mathbf{D}^{j}\mathbf{T}^{n}\phi\right\} \\ & = \mathbf{D}\left\{f\in L_{\mathbb{R}}^{2}|\exists \left(\alpha_{n}\right) \quad \text{such that} \quad f(x) = \sum_{n\in\mathbb{Z}}\alpha_{n}\mathbf{D}^{j}\mathbf{T}^{n}\phi\right\} \\ & = \mathbf{D}\operatorname{span}\left\{\mathbf{D}^{j}\mathbf{T}^{n}\phi\right|_{n\in\mathbb{Z}}\right\} \qquad \qquad \text{by definition of span} \qquad \text{(Definition 2.2 page 9)} \\ & = \mathbf{D}V_{j} \qquad \qquad \text{by induction hypothesis} \\ & = V_{j+1} \qquad \qquad \text{(Definition 4.1 page 54)} \end{split}$$

Example 4.1.


In the *Haar* MRA, the scaling function $\phi(x)$ is the *pulse function*

$$\phi(x) = \begin{cases} 1 & \text{for } x \in [0:1) \\ 0 & \text{otherwise.} \end{cases}$$



In the subspace V_j $(j \in \mathbb{Z})$ the scaling functions are

$$\mathbf{D}^{j}\phi(x) = \begin{cases} (2)^{j/2} & \text{for } x \in [0:(2^{-j})) \\ 0 & \text{otherwise.} \end{cases}$$




¹¹ Smith (2011) page 4

The scaling subspace V_0 is the span $V_0 \triangleq \text{span} \{ \mathbf{T}^n \phi | n \in \mathbb{Z} \}$. The scaling subspace V_j is the span $V_j \triangleq \text{span} \{ \mathbf{D}^j \mathbf{T}^n \phi | n \in \mathbb{Z} \}$. Note that $\| \mathbf{D}^j \mathbf{T}^n \phi \|$ for each resolution j and shift n is unity:


$$\|\mathbf{D}^{j}\mathbf{T}^{n}\phi\|^{2} = \|\phi\|^{2} \qquad \text{by } unitary \text{ properties of } \mathbf{T} \text{ and } \mathbf{D} \qquad \text{(Theorem 3.1 page 45)}$$

$$= \int_{0}^{1} |1|^{2} \, \mathrm{dx} \qquad \text{by definition of } \|\cdot\| \text{ on } \mathbf{L}_{\mathbb{R}}^{2} \qquad \text{(Definition D.1 page 141)}$$

Let $f(x) = \sin(\pi x)$. Suppose we want to project f(x) onto the subspaces V_0 , V_1 , V_2 ,



The values of the transform coefficients for the subspace V_j are given by



$$\begin{split} \left[\mathbf{R}_{j} \mathbf{f}(x) \right] &(n) = \frac{1}{\left\| \mathbf{D}^{j} \mathbf{T}^{n} \phi \right\|^{2}} \left\langle \mathbf{f}(x) \left| \mathbf{D}^{j} \mathbf{T}^{n} \phi \right\rangle \\ &= \frac{1}{\left\| \phi \right\|^{2}} \left\langle \mathbf{f}(x) \left| 2^{j/2} \phi (2^{j} x - n) \right\rangle \\ &= 2^{j/2} \left\langle \mathbf{f}(x) \left| \phi (2^{j} x - n) \right\rangle \\ &= 2^{j/2} \int_{2^{-j} n}^{2^{-j} (n+1)} \mathbf{f}(x) \, \mathrm{d}x \\ &= 2^{j/2} \int_{2^{-j} n}^{2^{-j} (n+1)} \sin(\pi x) \, \mathrm{d}x \\ &= 2^{j/2} \left(-\frac{1}{\pi} \right) \cos(\pi x) \Big|_{2^{-j} n}^{2^{-j} (n+1)} \\ &= \frac{2^{j/2}}{\pi} \left[\cos(2^{-j} n\pi) - \cos(2^{-j} (n+1)\pi) \right] \end{split}$$

by Proposition 3.3 page 41

And the projection $\mathbf{A}_n f(x)$ of the function f(x) onto the subspace \mathbf{V}_j is

$$\begin{aligned} \mathbf{A}_{j}\mathbf{f}(x) &= \sum_{n \in \mathbb{Z}} \left\langle \mathbf{f}(x) \mid \mathbf{D}^{j} \mathbf{T}^{n} \phi \right\rangle \mathbf{D}^{j} \mathbf{T}^{n} \phi \\ &= \frac{2^{j/2}}{\pi} \sum_{n \in \mathbb{Z}} \left[\cos\left(2^{-j} n \pi\right) - \cos\left(2^{-j} (n+1) \pi\right) \right] 2^{j/2} \phi \left(2^{j} x - n\right) \\ &= \frac{2^{j}}{\pi} \sum_{n \in \mathbb{Z}} \left[\cos\left(2^{-j} n \pi\right) - \cos\left(2^{-j} (n+1) \pi\right) \right] \phi \left(2^{j} x - n\right) \end{aligned}$$

The transforms of $sin(\pi x)$ into the subspaces V_0 , V_1 , and V_2 , as well as the approximations in those subspaces are as illustrated in Figure 4.1 (page 66).








Figure 4.1: Projections of $sin(\pi x)$ on Haar subspaces (Example 4.1 page 64)

4.5 Necessary Conditions

Theorem 4.3 (admissibility condition). Let $\check{\mathsf{h}}(z)$ be the Z-transform (Definition J.4 page 208) and $\check{\mathsf{h}}(\omega)$ the DISCRETE-TIME FOURIER TRANSFORM (Definition M.1 page 237) of a sequence $(\mathsf{h}_n)_{n\in\mathbb{Z}}$.

$$\left\{\left(\boldsymbol{L}_{\mathbb{R}}^{2},\,\left(\!\left(\boldsymbol{V}_{j}\right)\!\right),\,\phi,\,\left(\!\left(h_{n}\right)\!\right)\,is\,an\,\text{MRA SYSTEM (Definition 4.3 page 63)}\right\}$$

$$\Longrightarrow \left\{\sum_{n\in\mathbb{Z}}\mathsf{h}_{n}=\sqrt{2}\right\} \iff \left\{\check{\mathsf{h}}(z)\Big|_{z=1}=\sqrt{2}\right\} \iff \left\{\check{\mathsf{h}}(\omega)\Big|_{\omega=0}=\sqrt{2}\right\}$$

$$(1)\,\text{ADMISSIBILITY in "time"}$$

$$(2)\,\text{ADMISSIBILITY in "z domain"}$$

$$(3)\,\text{ADMISSIBILITY in "frequency"}$$

[♠]Proof:

1. Proof that MRA system \implies (1):

$$\begin{split} \sum_{n \in \mathbb{Z}} \mathsf{h}_n &= \frac{\int_{\mathbb{R}} \phi(x) \, \mathrm{d} \mathsf{x}}{\int_{\mathbb{R}} \phi(x) \, \mathrm{d} \mathsf{x}} \sum_{n \in \mathbb{Z}} \mathsf{h}_n \\ &= \frac{1}{\int_{\mathbb{R}} \phi(x) \, \mathrm{d} \mathsf{x}} \int_{\mathbb{R}} \sum_{n \in \mathbb{Z}} \mathsf{h}_n \phi(x) \, \mathrm{d} \mathsf{x} \\ &= \frac{1}{\int_{\mathbb{R}} \phi(x) \, \mathrm{d} \mathsf{x}} \int_{\mathbb{R}} \sum_{n \in \mathbb{Z}} \mathsf{h}_n \frac{\sqrt{2}}{\sqrt{2}} \phi(2y - n) 2 \, \mathrm{d} \mathsf{y} \qquad \text{let } y \triangleq \frac{x + n}{2} \implies x = 2y - n \implies \mathrm{d} \mathsf{x} = 2 \, \mathrm{d} \mathsf{y} \\ &= \frac{2}{\sqrt{2}} \frac{1}{\int_{\mathbb{R}} \phi(x) \, \mathrm{d} \mathsf{x}} \int_{\mathbb{R}} \sum_{n \in \mathbb{Z}} \mathsf{h}_n \mathbf{D} \mathbf{T}^n \phi(y) \, \mathrm{d} \mathsf{y} \qquad \text{by definitions of } \mathbf{T} \text{ and } \mathbf{D} \text{ (Definition 3.3 page 40)} \\ &= \sqrt{2} \frac{1}{\int_{\mathbb{R}} \phi(x) \, \mathrm{d} \mathsf{x}} \int_{\mathbb{R}} \phi(y) \, \mathrm{d} \mathsf{y} \qquad \text{by } dilation \ equation \text{ (Theorem 4.1 page 60)} \\ &= \sqrt{2} \end{split}$$

2. Alternate proof that MRA system \implies (1):



Let $f(x) \triangleq 1 \quad \forall x \in \mathbb{R}$.

$$\begin{split} \langle \phi \, | \, \mathsf{f} \, \rangle &= \left\langle \sum_{n \in \mathbb{Z}} \mathsf{h}_n \mathbf{D} \mathbf{T}^n \phi \, | \, \mathsf{f} \, \right\rangle & \text{by } \textit{dilation equation} & \text{(Theorem 4.1 page 60)} \\ &= \sum_{n \in \mathbb{Z}} \mathsf{h}_n \, \langle \mathbf{D} \mathbf{T}^n \phi \, | \, \mathsf{f} \, \rangle & \text{by linearity of } \langle \triangle \, | \, \nabla \rangle \\ &= \sum_{n \in \mathbb{Z}} \mathsf{h}_n \, \langle \phi \, | \, (\mathbf{D} \mathbf{T}^n)^* \mathsf{f} \, \rangle & \text{by definition of operator adjoint} & \text{(Theorem C.13 page 126)} \\ &= \sum_{n \in \mathbb{Z}} \mathsf{h}_n \, \langle \phi \, | \, (\mathbf{T}^*)^n \mathbf{D}^* \mathsf{f} \, \rangle & \text{by property of operator adjoint} & \text{(Theorem C.13 page 126)} \\ &= \sum_{n \in \mathbb{Z}} \mathsf{h}_n \, \langle \phi \, | \, (\mathbf{T}^{-1})^n \mathbf{D}^{-1} \mathsf{f} \, \rangle & \text{by unitary property of } \mathbf{T} \text{ and } \mathbf{D} & \text{(Proposition 3.7 page 43)} \\ &= \sum_{n \in \mathbb{Z}} \mathsf{h}_n \, \left\langle \phi \, | \, (\mathbf{T}^{-1})^n \frac{\sqrt{2}}{2} \mathsf{f} \, \right\rangle & \text{because } \mathsf{f} \text{ is a constant hypothesis} & \text{and by Proposition 3.2 page 41} \\ &= \sum_{n \in \mathbb{Z}} \mathsf{h}_n \, \left\langle \phi \, | \, \frac{\sqrt{2}}{2} \mathsf{f} \, \right\rangle & \text{by } \mathsf{f}(x) = 1 \text{ definition} \\ &= \sum_{n \in \mathbb{Z}} \mathsf{h}_n \, \frac{\sqrt{2}}{2} \, \langle \phi \, | \, \mathsf{f} \, \rangle & \text{by property of } \langle \triangle \, | \, \nabla \rangle \\ &= \frac{\sqrt{2}}{2} \, \left\langle \phi \, | \, \mathsf{f} \, \right\rangle & \sum_{n \in \mathbb{Z}} \mathsf{h}_n \\ &\Longrightarrow \sum_{n \in \mathbb{Z}} \mathsf{h}_n = \sqrt{2} \end{split}$$

- 3. Proof that (1) \iff (2) \iff (3): by Proposition M.2 page 239.
- 4. Proof for \Leftarrow part: by Counterexample 4.1 page 67.

$$\begin{array}{c} \text{Counterexample 4.1. Let } \left(\mathcal{L}_{\mathbb{R}}^2, \, \left(\!\! \left(\mathcal{V}_j \!\! \right), \, \phi, \, \left(h_n \!\! \right) \!\! \right) \text{ be an } \textit{MRA system } \text{(Definition 4.3 page 63).} \\ \\ \left\{ \left(\!\!\! \left(\!\!\! h_n \!\! \right) \right) \triangleq \sqrt{2} \bar{\delta}_{n-1} \triangleq \left\{ \begin{array}{c} \sqrt{2} & \text{for } n=1 \\ 0 & \text{otherwise.} \end{array} \right. & \longrightarrow \left\{ \phi(x) = 0 \right\} \\ \text{which means} \\ \left\{ \sum_{n \in \mathbb{Z}} \mathsf{h}_n = \sqrt{2} \right\} & \Longrightarrow \left\{ \left(\mathcal{L}_{\mathbb{R}}^2, \, \left(\!\! \left(\mathcal{V}_j \!\! \right), \, \phi, \, \left(h_n \!\! \right) \!\! \right) \text{ is an MRA system for } \mathcal{L}_{\mathbb{R}}^2. \right\} \\ \end{array}$$

[♠]Proof:

$$\phi(x) = \sum_{n \in \mathbb{Z}} \mathsf{h}_n \mathbf{D} \mathbf{T}^n \phi(x) \qquad \text{by } \textit{dilation equation} \qquad \text{(Theorem 4.1 page 60)}$$

$$= \sum_{n \in \mathbb{Z}} \mathsf{h}_n \phi(2x - n) \qquad \text{by definitions of } \mathbf{D} \text{ and } \mathbf{T} \qquad \text{(Definition 3.3 page 40)}$$

$$= \sum_{n \in \mathbb{Z}} \sqrt{2} \bar{\delta}_{n-1} \phi(2x - n) \qquad \text{by definitions of } (\mathsf{h}_n)$$

$$= \sqrt{2} \phi(2x - 1) \qquad \text{by definition of } \phi(x)$$

$$\Rightarrow \phi(x) = 0$$

—>

This implies $\phi(x)=0$, which implies that $\left(\boldsymbol{L}_{\mathbb{R}}^{2},\,\left(\boldsymbol{V}_{j}\right),\,\phi,\,\left(h_{n}\right)\right)$ is *not* an *MRA system* for $\boldsymbol{L}_{\mathbb{R}}^{2}$ because

$$\left(\bigcup_{j\in\mathbb{Z}} \mathbf{\textit{V}}_j\right)^- = \left(\bigcup_{j\in\mathbb{Z}} \operatorname{span}\left\{\mathbf{D}^j\mathbf{T}^n\phi|_{n\in\mathbb{Z}}\right\}\right)^- \neq \mathbf{\textit{L}}_{\mathbb{R}}^2$$

(the *least upper bound* is *not* $L^2_{\mathbb{R}}$).

Theorem 4.4 (Quadrature condition in "time"). Let $(L_{\mathbb{R}}^2, (V_j), \phi, (h_n))$ be an MRA SYSTEM (Definition 4.3 page 63).

$$\begin{array}{c|c} \mathbf{T} & \mathbf{H} & \sum_{m \in \mathbb{Z}} \mathsf{h}_m \sum_{k \in \mathbb{Z}} \mathsf{h}_k^* \left\langle \phi \mid \mathbf{T}^{2n-m+k} \phi \right\rangle = \left\langle \phi \mid \mathbf{T}^n \phi \right\rangle \qquad \forall n \in \mathbb{Z} \end{array}$$

№ Proof:

$$\langle \phi \, | \, \mathbf{T}^n \phi \rangle = \left\langle \sum_{m \in \mathbb{Z}} \mathsf{h}_m \mathbf{D} \mathbf{T}^m \phi \, | \, \mathbf{T}^n \sum_{k \in \mathbb{Z}} \mathsf{h}_k \mathbf{D} \mathbf{T}^k \phi \right\rangle \qquad \text{by } dilation \ equation}$$
 (Theorem 4.1 page 60)
$$= \sum_{m \in \mathbb{Z}} \mathsf{h}_m \sum_{k \in \mathbb{Z}} \mathsf{h}_k^* \left\langle \mathbf{D} \mathbf{T}^m \phi \, | \, \mathbf{T}^n \mathbf{D} \mathbf{T}^k \phi \right\rangle \qquad \text{by properties of } \langle \triangle \, | \, \nabla \rangle$$

$$= \sum_{m \in \mathbb{Z}} \mathsf{h}_m \sum_{k \in \mathbb{Z}} \mathsf{h}_k^* \left\langle \phi \, | \, (\mathbf{D} \mathbf{T}^m)^* \, \mathbf{T}^n \mathbf{D} \mathbf{T}^k \phi \right\rangle \qquad \text{by definition of operator adjoint} \qquad \text{(Proposition C.3 page 125)}$$

$$= \sum_{m \in \mathbb{Z}} \mathsf{h}_m \sum_{k \in \mathbb{Z}} \mathsf{h}_k^* \left\langle \phi \, | \, (\mathbf{D} \mathbf{T}^m)^* \, \mathbf{D} \mathbf{T}^{2n} \mathbf{T}^k \phi \right\rangle \qquad \text{by Proposition 3.5 page 42}$$

$$= \sum_{m \in \mathbb{Z}} \mathsf{h}_m \sum_{k \in \mathbb{Z}} \mathsf{h}_k^* \left\langle \phi \, | \, \mathbf{T}^{*m} \mathbf{D}^* \mathbf{D} \mathbf{T}^{2n} \mathbf{T}^k \phi \right\rangle \qquad \text{by operator star-algebra properties} \qquad \text{(Theorem C.13 page 126)}$$

$$= \sum_{m \in \mathbb{Z}} \mathsf{h}_m \sum_{k \in \mathbb{Z}} \mathsf{h}_k^* \left\langle \phi \, | \, \mathbf{T}^{-m} \mathbf{D}^{-1} \mathbf{D} \mathbf{T}^{2n} \mathbf{T}^k \phi \right\rangle \qquad \text{by Proposition 3.7 page 43}$$

$$= \sum_{m \in \mathbb{Z}} \mathsf{h}_m \sum_{k \in \mathbb{Z}} \mathsf{h}_k^* \left\langle \phi \, | \, \mathbf{T}^{2n-m+k} \phi \right\rangle$$

Theorem 4.5 (next) presents the *quadrature necessary conditions* of a *wavelet system*. These relations simplify dramatically in the special case of an *orthonormal wavelet system* (Theorem M.4 page 243).

Theorem 4.5 (Quadrature condition in "frequency"). ¹² Let $(L_{\mathbb{R}}^2, (V_j), \phi, (h_n))$ be an MRA system (Definition 4.3 page 63). Let $\tilde{\mathbf{x}}(\omega)$ be the DISCRETE TIME FOURIER TRANSFORM (Definition M.1 page 237) for a sequence $(x_n)_{n\in\mathbb{Z}}$ in $\boldsymbol{\mathcal{E}}_{\mathbb{R}}^2$. Let $\tilde{\mathbf{x}}(\omega)$ be the Auto-Power spectrum (Definition P.3 page 255) of ϕ .

$$|\check{\mathbf{h}}(\omega)|^2 \tilde{\mathbf{S}}_{\phi\phi}(\omega) + |\check{\mathbf{h}}(\omega + \pi)|^2 \tilde{\mathbf{S}}_{\phi\phi}(\omega + \pi) = 2\tilde{\mathbf{S}}_{\phi\phi}(2\omega)$$

$$Note: \dot{S}_{\phi\phi}(\omega) = 1$$
 $for orthonormal MRA$

♥Proof:

$$\begin{split} &2\tilde{\mathbf{S}}_{\phi\phi}(2\omega) \\ &= 2(2\pi) \sum_{n \in \mathbb{Z}} \left| \tilde{\phi}(2\omega + 2\pi n) \right|^2 \qquad \qquad \text{by Theorem P.1 page 255} \\ &= 2(2\pi) \sum_{n \in \mathbb{Z}} \left| \frac{\sqrt{2}}{2} \check{\mathbf{h}} \left(\frac{2\omega + 2\pi n}{2} \right) \tilde{\phi} \left(\frac{2\omega + 2\pi n}{2} \right) \right|^2 \qquad \qquad \qquad \text{by Lemma 4.1 page 61} \\ &= 2\pi \sum_{n \in \mathbb{Z}_e} \left| \check{\mathbf{h}} \left(\frac{2\omega + 2\pi n}{2} \right) \right|^2 \left| \tilde{\phi} \left(\frac{2\omega + 2\pi n}{2} \right) \right|^2 + 2\pi \sum_{n \in \mathbb{Z}_e} \left| \check{\mathbf{h}} \left(\frac{2\omega + 2\pi n}{2} \right) \right|^2 \left| \tilde{\phi} \left(\frac{2\omega + 2\pi n}{2} \right) \right|^2 \end{split}$$



$$= 2\pi \sum_{n \in \mathbb{Z}} \left| \check{\mathbf{h}} \left(\omega + 2\pi n \right) \right|^{2} \left| \tilde{\phi} \left(\omega + 2\pi n \right) \right|^{2} + 2\pi \sum_{n \in \mathbb{Z}} \left| \check{\mathbf{h}} \left(\omega + 2\pi n + \pi \right) \right|^{2} \left| \tilde{\phi} \left(\omega + 2\pi n + \pi \right) \right|^{2}$$

$$= 2\pi \sum_{n \in \mathbb{Z}} \left| \check{\mathbf{h}} \left(\omega \right) \right|^{2} \left| \tilde{\phi} \left(\omega + 2\pi n \right) \right|^{2} + 2\pi \sum_{n \in \mathbb{Z}} \left| \check{\mathbf{h}} \left(\omega + \pi \right) \right|^{2} \left| \tilde{\phi} \left(\omega + 2\pi n + \pi \right) \right|^{2}$$
 by Proposition M.1 page 237
$$= \left| \check{\mathbf{h}} \left(\omega \right) \right|^{2} \left(2\pi \sum_{n \in \mathbb{Z}} \left| \tilde{\phi} \left(\omega + 2\pi n \right) \right|^{2} \right) + \left| \check{\mathbf{h}} \left(\omega + \pi \right) \right|^{2} \left(2\pi \sum_{n \in \mathbb{Z}} \left| \tilde{\phi} \left(\omega + \pi + 2\pi n \right) \right|^{2} \right)$$

$$= \left| \check{\mathbf{h}} \left(\omega \right) \right|^{2} \check{\mathbf{S}}_{\phi\phi}(\omega) + \left| \check{\mathbf{h}} \left(\omega + \pi \right) \right|^{2} \check{\mathbf{S}}_{\phi\phi}(\omega + \pi)$$
 by Theorem P.1 page 255

Sufficient conditions 4.6

Theorem 4.6 (next) gives a set of *sufficient* conditions on the *scaling function* (Definition 4.1 page 54) ϕ to generate an MRA.

$$\begin{array}{l} \textbf{Theorem 4.6.} \quad \overset{13}{\text{Let }} V_j \triangleq \text{span } \{ \mathbf{T} \phi(x) | n \in \mathbb{Z} \} \text{ (Definition 2.2 page 9).} \\ \\ \begin{pmatrix} (1). & ((\mathbf{T}^n \phi)) \text{ is a RIESZ SEQUENCE (Definition 2.14 page 27)} & and \\ (2). & \exists ((\mathbf{h}_n)) \text{ such that } \phi(x) = \sum_{n \in \mathbb{Z}} h_n \mathbf{D} \mathbf{T}^n \phi(x) & and \\ (3). & \tilde{\phi}(\omega) \text{ is CONTINUOUS at 0} & and \\ (4). & \tilde{\phi}(0) \neq 0 & & & \\ \end{pmatrix} \\ \Rightarrow \begin{cases} \begin{pmatrix} (\mathbf{V}_j) \\ j \in \mathbb{Z} \text{ is an MRA} \\ \text{(Definition 4.1 page 54)} \end{pmatrix} \\ \\ & \text{(Definition 4.1 page 54)} \\ \end{pmatrix}$$

 $^{\circ}$ Proof: For this to be true, each of the conditions in the definition of an MRA (Definition 4.1 page 54) must be satisfied:

- 1. Proof that each V_i is *closed*: by definition of span
- 2. Proof that (V_i) is linearly ordered:

$$V_i \subseteq V_{i+1} \iff \operatorname{span}\{\mathbf{D}^j \mathbf{T}^n \phi\} \subseteq \operatorname{span}\{\mathbf{D}^{j+1} \mathbf{T}^n \phi\} \iff (2)$$

- 3. Proof that $\bigcup_{j\in\mathbb{Z}} V_j$ is *dense* in $L^2_{\mathbb{R}}$: by Proposition 4.2 page 57
- 4. Proof of *self-similar* property:

$$\left\{\mathsf{f} \in \mathbf{V}_j \iff \mathbf{D}\mathsf{f} \in \mathbf{V}_{j+1}\right\} \iff \mathsf{f} \in \mathsf{span}\{\mathbf{T}^n \phi\} \iff \mathbf{D}\mathsf{f} \in \mathsf{span}\{\mathbf{D}\mathbf{T}^n \phi\} \iff (2)$$

5. Proof for *Riesz basis*: by (1) and Proposition 4.2 page 57.

Wojtaszczyk (1997) page 28 ⟨Theorem 2.13⟩, ₱ Pinsky (2002) page 313 ⟨Theorem 6.4.27⟩



₿

Support size

The *support* of a function is what it's non-zero part "sits" on. If the support of the scaling coefficients (h_n) goes from say [0,3] in \mathbb{Z} , what is the support of the scaling function $\phi(x)$? The answer is [0, 3] in \mathbb{R} —essentially the same as the support of (h_n) except that the two functions have different domains (\mathbb{Z} versus \mathbb{R}). This concept is defined in Definition 4.4 (next definition), and proven in Theorem 4.7 (next theorem).

Definition 4.4. Let $(L_{\mathbb{R}}^2, (V_j), \phi, (h_n))$ be an MRA system (Definition 4.3 page 63). Let X^- represent the Closure of a set X in $L_{\mathbb{R}}^2$, $\vee X$ the least upper bound of an ordered set (X, \leq) , $\wedge X$ the Greatest LOWER BOUND $of(X, \leq)$, and



Theorem 4.7 (support size). ¹⁴ Let $(L_{\mathbb{R}}^2, (V_j), \phi, (h_n))$ be an MRA SYSTEM (Definition 4.3 page 63). Let supp f be the support of a function f (Definition 4.4 page 70).

$$\frac{\mathsf{T}}{\mathsf{H}}$$
 $\mathrm{sm}\phi=\mathrm{sm}h$

[♠]Proof:

- 1. Definitions: $supp \phi \triangleq [a, b]$ $supph \triangleq [k, m].$
- 2. lemma: $supp \phi(x) = [a, b] \iff supp \phi(2x) = \left| \frac{a}{2}, \frac{b}{2} \right|$
- 3. lemma: $sup[\lambda \phi(x)] = sup[\phi(x)] \quad \forall \lambda \in \mathbb{R} \setminus 0$
- 4. Proof that k = a:

$$a = \bigwedge \operatorname{supp} \phi(x)$$

$$\triangleq \bigwedge \operatorname{supp} \left[\sum_{n \in \mathbb{Z}} h_n \mathbf{D} \mathbf{T}^n \phi(x) \right]$$

$$= \bigwedge \operatorname{supp} \left[\sqrt{2} \sum_{n \in \mathbb{Z}} h_n \phi(2x - n) \right]$$

$$= \bigwedge \operatorname{supp} \left[\sum_{n \in \mathbb{Z}} h_n \phi(2x - n) \right]$$

$$= \bigwedge \operatorname{supp} \left[h_k \phi(2x - k) \right]$$

$$= \bigwedge \operatorname{supp} \left[\phi(2x - k) \right]$$

$$= \bigwedge \operatorname{supp} \left[\phi\left(2\left[x - \frac{k}{2}\right]\right) \right]$$

$$= \bigwedge \left\{ t | \phi\left(2\left[x - \frac{k}{2}\right]\right) \neq 0 \right\}$$

by definition of a (item (1) page 70)

by dilation equation (Theorem 4.1 page 60)

by definition of T and D (Definition 3.3 page 40)

by (3) lemma

because n = k is the *least value* of n for which $h_n \neq 0$

by (3) lemma

by definition of supp (Definition 4.4 page 70)

Mallat (1999) pages 243–244



$$= x \quad \text{such that} \quad x - \frac{k}{2} = \frac{a}{2}$$

$$= \frac{k}{2} + \frac{a}{2}$$

$$\Longrightarrow$$

$$\iff$$

by (2) lemma

$$\frac{k}{2} = a - \frac{a}{2}$$

5. Proof that m = b:

$$b = \bigvee \operatorname{spp}\phi(x)$$

$$\triangleq \bigvee \operatorname{spp} \left[\sum_{n \in \mathbb{Z}} h_n \mathbf{D} \mathbf{T}^n \phi(x) \right]$$

$$= \bigvee \operatorname{spp} \left[\sqrt{2} \sum_{n \in \mathbb{Z}} h_n \phi(2x - n) \right]$$

$$= \bigvee \operatorname{spp} \left[\sum_{n \in \mathbb{Z}} h_n \phi(2x - n) \right]$$

$$= \bigvee \operatorname{spp} \left[h_m \phi(2x - m) \right]$$

$$= \bigvee \operatorname{spp} \left[\phi(2x - m) \right]$$

$$= \bigvee \left\{ t | \phi\left(2\left[x - \frac{m}{2}\right]\right) \right\}$$

$$= x \quad \text{such that} \quad x - \frac{m}{2} = \frac{b}{2}$$

$$= \frac{m}{2} + \frac{b}{2}$$

$$\implies \frac{m}{2} = b - \frac{b}{2}$$

$$\iff m = b$$

by definition of b

(item (1) page 70)

by dilation equation

(Theorem 4.1 page 60)

by definition of T and D

(Definition 3.3 page 40)

by (3) lemma

because n = m is the greatest value of n for which $h_n \neq 0$


by (3) lemma

by definition of supp

(Definition 4.4 page 70)


by (2) lemma

Example 4.2.




Scaling functions with partition of unity 4.8

The Z transform (Definition J.4 page 208) of a sequence (h_n) with sum $\sum_{n\in\mathbb{Z}}(-1)^nh_n=0$ has a zero at z = -1. Somewhat surprisingly, the partition of unity and zero at z = -1 properties are actually equivalent (next theorem).



Theorem 4.8. ¹⁵ $Let(L_{\mathbb{R}}^2, (V_j), \phi, (h_n))$ be a multiresolution system (Definition 4.3 page 63). $Let \tilde{\mathbf{F}} f(\omega)$ be the Fourier transform (Definition 1.2 page 196) of a function $f \in L_{\mathbb{R}}^2$. Let $\bar{\delta}_n$ be the Kronecker Delta Function (Definition 2.12 page 20). Let c be some contant in $\mathbb{R} \setminus 0$.



 $^{\mathbb{Q}}$ Proof: Let \mathbb{Z}_{e} be the set of even integers and \mathbb{Z}_{o} the set of odd integers.

1. Proof that $(1) \Leftarrow (2)$:

$$\begin{split} \sum_{n\in\mathbb{Z}}\mathbf{T}^n\phi &= \sum_{n\in\mathbb{Z}}\mathbf{T}^n\left[\sum_{m\in\mathbb{Z}}\mathsf{h}_m\mathbf{D}\mathbf{T}^m\phi\right] & \text{by } \textit{dilation equation} \qquad \text{(Theorem 4.1 page 60)} \\ &= \sum_{m\in\mathbb{Z}}\mathsf{h}_m\sum_{n\in\mathbb{Z}}\mathbf{D}\mathbf{T}^n\mathbf{D}\mathbf{T}^m\phi \\ &= \sum_{m\in\mathbb{Z}}\mathsf{h}_m\sum_{n\in\mathbb{Z}}\mathbf{D}\mathbf{T}^2\mathbf{T}^m\phi \qquad \qquad \text{by } \textit{commutator relation} \qquad \text{(Proposition 3.5 page 42)} \\ &= \mathbf{D}\sum_{m\in\mathbb{Z}}\mathsf{h}_m\sum_{n\in\mathbb{Z}}\mathbf{T}^{2n}\mathbf{T}^m\phi \qquad \qquad \text{by } \textit{PSF} \qquad \qquad \text{(Theorem 3.2 page 48)} \\ &= \mathbf{D}\sum_{m\in\mathbb{Z}}\mathsf{h}_m\left[\sqrt{\frac{2\pi}{2}}\hat{\mathbf{F}}^{-1}\mathbf{S}_2\tilde{\mathbf{F}}(\mathbf{T}^m\phi)\right] \qquad \text{by } \textit{PSF} \qquad \qquad \text{(Theorem 3.2 page 48)} \\ &= \sqrt{\pi}\mathbf{D}\sum_{m\in\mathbb{Z}}\mathsf{h}_m\hat{\mathbf{F}}^{-1}\mathbf{S}_2e^{-i\omega m}\tilde{\mathbf{F}}\phi \qquad \qquad \text{by Corollary 3.1 page 47} \\ &= \sqrt{\pi}\mathbf{D}\sum_{m\in\mathbb{Z}}\mathsf{h}_m\hat{\mathbf{F}}^{-1}e^{-i\frac{2\pi}{2}km}\mathbf{S}_2\tilde{\mathbf{F}}\phi \qquad \qquad \text{by definition of } \mathbf{S} \qquad \text{(Definition 3.4 page 48)} \\ &= \sqrt{\pi}\mathbf{D}\sum_{m\in\mathbb{Z}}\mathsf{h}_m\hat{\mathbf{F}}^{-1}(-1)^{km}\mathbf{S}_2\tilde{\mathbf{F}}\phi \qquad \qquad \text{by definition of } \mathbf{S} \qquad \text{(Definition 3.4 page 48)} \\ &= \sqrt{\pi}\mathbf{D}\sum_{m\in\mathbb{Z}}\mathsf{h}_m\hat{\mathbf{F}}^{-1}(-1)^{km}\mathbf{S}_2\tilde{\mathbf{F}}\phi \qquad \qquad \text{by definition of } \hat{\mathbf{F}}^{-1} \qquad \text{(Theorem N.1 page 248)} \\ &= \frac{\sqrt{2\pi}}{2}\mathbf{D}\sum_{k\in\mathbb{Z}}\left(\mathbf{S}_2\tilde{\mathbf{F}}\phi\right)e^{i\pi kx}\sum_{m\in\mathbb{Z}}(-1)^{km}\mathsf{h}_m \\ &= \frac{\sqrt{2\pi}}{2}\mathbf{D}\sum_{k\in\mathbb{Z}_a}\left(\mathbf{S}_2\tilde{\mathbf{F}}\phi\right)e^{i\pi kx}\sum_{m\in\mathbb{Z}}(-1)^{km}\mathsf{h}_m + \frac{\sqrt{2\pi}}{2}\mathbf{D}\sum_{k\in\mathbb{Z}_a}\left(\mathbf{S}_2\tilde{\mathbf{F}}\phi\right)e^{i\pi kx}\sum_{m\in\mathbb{Z}}(-1)^{km}\mathsf{h}_m \\ &= \frac{\sqrt{2\pi}}{2}\mathbf{D}\sum_{k\in\mathbb{Z}_a}\left(\mathbf{S}_2\tilde{\mathbf{F}}\phi\right)e^{i\pi kx}\sum_{m\in\mathbb{Z}}\mathsf{h}_m + \frac{\sqrt{2\pi}}{2}\mathbf{D}\sum_{k\in\mathbb{Z}_a}\left(\mathbf{S}_2\tilde{\mathbf{F}}\phi\right)e^{i\pi kx}\sum_{m\in\mathbb{Z}}(-1)^{km}\mathsf{h}_m \\ &= \frac{\sqrt{2\pi}}{2}\mathbf{D}\sum_{k\in\mathbb{Z}_a}\left(\mathbf{S}_2\tilde{\mathbf{F}}\phi\right)e^{i\pi kx}\sum_{m\in\mathbb{Z}}\mathsf{h}_m\mathbf{D}^{-1}\mathbf{D}^{-1}\mathbf{D}^{-1}\mathbf{D}^{-1}\mathbf{D}^{-1}\mathbf{D}^{-1}\mathbf{D}^{-1}\mathbf{D}^{-1}\mathbf{D}^{-1}\mathbf{D}^{-$$

¹⁵ Jawerth and Sweldens (1994) page 8, *■* Chui (1992) page 123



$$= \sqrt{\pi} \mathbf{D} \sum_{k \in \mathbb{Z}_{e}} (\mathbf{S}_{2} \tilde{\mathbf{F}} \phi) e^{i\pi kx} \qquad \text{by Theorem 4.3 (page 66)} \quad \text{and right hypothesis}$$

$$= \sqrt{\pi} \mathbf{D} \sum_{k \in \mathbb{Z}_{e}} \tilde{\phi} \left(\frac{2\pi}{2}k\right) e^{i\pi kx} \qquad \text{by definitions of } \tilde{\mathbf{F}} \text{ and } \mathbf{S}_{2}$$

$$= \sqrt{\pi} \mathbf{D} \sum_{k \in \mathbb{Z}} \tilde{\phi}(2\pi k) e^{i2\pi kx} \qquad \text{by definition of } \mathbb{Z}_{e}$$

$$= \frac{1}{\sqrt{2}} \mathbf{D} \left\{ \sqrt{2\pi} \sum_{k \in \mathbb{Z}} \tilde{\phi}(2\pi k) e^{i2\pi kx} \right\}$$

$$= \frac{1}{\sqrt{2}} \mathbf{D} \sum_{n \in \mathbb{Z}} \phi(x+n) \qquad \text{by } PSF \qquad \text{(Theorem 3.2 page 48))}$$

$$= \frac{1}{\sqrt{2}} \mathbf{D} \sum_{n} \mathbf{T}^{n} \phi \qquad \text{by definition of } \mathbf{T} \qquad \text{(Definition 3.3 page 40)}$$

The above equation sequence demonstrates that

$$\mathbf{D}\sum_{n}\mathbf{T}^{n}\phi=\sqrt{2}\sum_{n}\mathbf{T}^{n}\phi$$

(essentially that $\sum_n \mathbf{T}^n \phi$ is equal to it's own dilation). This implies that $\sum_n \mathbf{T}^n \phi$ is a constant (Proposition

2. Proof that $(1) \Longrightarrow (2)$:

$$c = \sum_{n \in \mathbb{Z}} \mathbf{T}^n \phi \qquad \qquad \text{by left hypothesis}$$

$$= \sqrt{2\pi} \ \hat{\mathbf{F}}^{-1} \mathbf{S} \hat{\mathbf{F}} \phi \qquad \qquad \text{by Lemma 4.1 page 61}$$

$$= \sqrt{2\pi} \ \hat{\mathbf{F}}^{-1} \mathbf{S} \sqrt{2} \left(\mathbf{D}^{-1} \sum_{n \in \mathbb{Z}} \mathbf{h}_n e^{-i\omega n} \right) \left(\mathbf{D}^{-1} \hat{\mathbf{F}} \phi \right) \qquad \qquad \text{by Lemma 4.1 page 61}$$

$$= 2\sqrt{\pi} \ \hat{\mathbf{F}}^{-1} \left(\mathbf{S} \mathbf{D}^{-1} \sum_{n \in \mathbb{Z}} \mathbf{h}_n e^{-i\omega n} \right) \left(\mathbf{S} \hat{\mathbf{F}} \mathbf{D} \phi \right) \qquad \qquad \text{by Corollary 3.1 page 47}$$

$$= 2\sqrt{\pi} \ \hat{\mathbf{F}}^{-1} \left(\mathbf{S} \frac{1}{\sqrt{2}} \sum_{n \in \mathbb{Z}} \mathbf{h}_n e^{-i\frac{\omega}{2}n} \right) \left(\mathbf{S} \hat{\mathbf{F}} \mathbf{D} \phi \right) \qquad \qquad \text{by evaluation of } \mathbf{D}^{-1} \qquad \text{(Proposition 3.2 page 41)}$$

$$= \sqrt{2\pi} \ \hat{\mathbf{F}}^{-1} \left(\sum_{n \in \mathbb{Z}} \mathbf{h}_n e^{-i\frac{2\pi i}{2}n} \right) \left(\mathbf{S} \hat{\mathbf{F}} \mathbf{D} \phi \right) \qquad \qquad \text{by definition of } \mathbf{S} \qquad \text{(Definition 3.4 page 48)}$$

$$= \sqrt{2\pi} \ \hat{\mathbf{F}}^{-1} \left(\sum_{n \in \mathbb{Z}} \mathbf{h}_n (-1)^{kn} \right) \left(\mathbf{S} \mathbf{D}^{-1} \mathbf{F} \phi \right) \qquad \qquad \text{by definition of } \mathbf{S} \qquad \text{(Definition 3.4 page 48)}$$

$$= \sqrt{2\pi} \ \hat{\mathbf{F}}^{-1} \left(\sum_{n \in \mathbb{Z}} \mathbf{h}_n (-1)^{kn} \right) \left(\mathbf{S} \frac{1}{\sqrt{2}} \tilde{\phi} \left(\frac{\omega}{2} \right) \right) \qquad \qquad \text{by definition of } \mathbf{S} \qquad \text{(Definition 3.4 page 48)}$$

$$= \sqrt{\pi} \sum_{k \in \mathbb{Z}} \sum_{n \in \mathbb{Z}} \mathbf{h}_n (-1)^{kn} \ \tilde{\phi}(\pi k) e^{i2\pi kx} + \sqrt{\pi} \sum_{k \text{ odd}} \sum_{n \in \mathbb{Z}} \mathbf{h}_n (-1)^{kn} \ \tilde{\phi}(\pi k) e^{i2\pi kx}$$

$$= \sqrt{\pi} \sum_{k \text{ even}} \sum_{n \in \mathbb{Z}} \mathbf{h}_n (-1)^{kn} \tilde{\phi}(\pi k) e^{i2\pi kx} + \sqrt{\pi} \sum_{k \text{ odd}} \sum_{n \in \mathbb{Z}} \mathbf{h}_n (-1)^{kn} \ \tilde{\phi}(\pi k) e^{i2\pi kx}$$

$$= \sqrt{\pi} \sum_{k \text{ even}} \sum_{n \in \mathbb{Z}} \mathbf{h}_n \left(\sum_{n \in \mathbb{Z}} \mathbf{h}_n \right) \left(\sum_{n \in \mathbb{Z}} \mathbf{h}_n (-1)^{kn} \right) \left(\sum_{n \in \mathbb{Z}} \mathbf{h}_n (-1)^{kn} \right) \tilde{\phi}(\pi k) e^{i2\pi kx}$$

<u>©</u> **0⊗**⊜

$$=\sqrt{\pi}\sum_{k\in\mathbb{Z}}\sqrt{2}\,\tilde{\phi}(\pi 2k)e^{i2\pi 2kx}+\sqrt{\pi}\sum_{k\in\mathbb{Z}}\left(\sum_{n\in\mathbb{Z}}\mathsf{h}_n(-1)^n\right)\tilde{\phi}(\pi[2k+1])e^{i2\pi[2k+1]x}\quad\text{by Theorem 4.3 page 66}$$

$$=\frac{\sqrt{2\pi}}{\sqrt{2\pi}}\tilde{\phi}(0)+\sqrt{\pi}e^{i2\pi x}\sum_{n\in\mathbb{Z}}\mathsf{h}_n(-1)^n\sum_{k\in\mathbb{Z}}\tilde{\phi}(\pi[2k+1])e^{i4\pi kx}\quad\text{by left hypothesis and Theorem \ref{eq:page \ref{eq:page final}}}\text{ because the right side must equal }c$$

3. Proof that $(2) \Longrightarrow (3)$:

$$\sum_{n \in \mathbb{Z}_e} \mathsf{h}_n = \sum_{n \in \mathbb{Z}_o} \mathsf{h}_n = \frac{1}{2} \sum_{n \in \mathbb{Z}} \mathsf{h}_n \qquad \qquad \text{by (2) and Proposition M.4 page 240}$$

$$= \frac{\sqrt{2}}{2} \qquad \qquad \text{by } admissibility \ condition \ (\text{Theorem 4.3 page 66})$$

4. Proof that $(2) \Leftarrow (3)$:

$$\frac{\sqrt{2}}{2} = \underbrace{\sum_{n \in \mathbb{Z}_{e}} (-1)^{n} \mathsf{h}_{n}}_{\text{even terms}} + \underbrace{\sum_{n \in \mathbb{Z}_{o}} (-1)^{n} \mathsf{h}_{n}}_{\text{odd terms}}$$
by (3)
$$\implies \underbrace{\sum_{n \in \mathbb{Z}} (-1)^{n} \mathsf{h}_{n}}_{\text{n} = 0}$$
by Proposition M.4 page 240

Not every function that forms a *partition of unity* is a *basis* for an *MRA*, as formerly stated next and demonstrated by Counterexample 4.2 (page 74) and Counterexample 4.3 (page 76).

Proposition 4.5.

 $\phi(x) \text{ generates a PARTITION OF UNITY} \implies \phi(x) \text{ generates an MRA system.}$

№ Proof: By Counterexample 4.2 (page 74) and Counterexample 4.3 (page 76).

Counterexample 4.2. Let a function ϕ be defined in terms of the sine function (Definition G.2 page 155) as follows:

 $\phi(x) \triangleq \begin{cases} \sin^2(\frac{\pi}{2}x) & \text{for } x \in [0:2] \\ 0 & \text{otherwise} \end{cases}$

Then $\int_{\mathbb{R}} \phi(x) dx = 1$ and ϕ induces a *partition of unity*

-1 0 1 2 3 -5 -4 -3 -2 -1 0 1 2 3 4 5

but $\{ \mathbf{T}^n \phi | n \in \mathbb{Z} \}$ does **not** generate an *MRA*.


 \mathbb{Q} PROOF: Let $\mathbb{1}_A(x)$ be the *set indicator function* (Definition 3.2 page 40) on a set A.

- 1. Proof that $\int_{\mathbb{R}} \phi(x) dx = 1$: by Example **??** (page **??**)
- 2. Proof that $\phi(x)$ forms a *partition of unity*: by Example **??** (page **??**)
- 3. Proof that $\phi(x) \notin \text{span} \{ \mathbf{DT}^n \phi(x) | n \in \mathbb{Z} \}$ (and so does not generate an *MRA*):



CNT

- (a) Note that the *support* (Definition 4.4 page 70) of ϕ is $supp \phi = [0:2]$.
- (b) Therefore, the *support* of (h_n) is $supp(h_n) = \{0, 1, 2\}$ (Theorem 4.7 page 70).
- (c) So if $\phi(x)$ is an MRA, we only need to compute $\{h_0, h_1, h_2\}$ (the rest would be 0).



Here would be the values of $\{h_1, h_2, h_3\}$:

$$\phi(x) = \sum_{n \in \mathbb{Z}} \mathsf{h}_n \mathbf{D} \mathbf{T}^n \phi(x)$$

$$= \sum_{n \in \mathbb{Z}} \mathsf{h}_n \mathbf{D} \mathbf{T}^n \sin^2 \left(\frac{\pi}{2} x\right) \mathbb{1}_{[0:2]}(x)$$

$$= \sum_{n \in \mathbb{Z}} \mathsf{h}_n \sin^2 \left(\frac{\pi}{2} (2x - n)\right) \mathbb{1}_{[0:2]}(2x - n)$$

$$= \sum_{n = 0}^2 \mathsf{h}_n \sin^2 \left(\frac{\pi}{2} (2x - n)\right) \mathbb{1}_{[0:2]}(2x - n)$$
 by Theorem 4.7 page 70

(d) The values of (h_0, h_1, h_2) can be conveniently calculated at the knot locations $x = \frac{1}{2}$, x = 1, and $x = \frac{3}{2}$ (see the diagram in item (3c) page 75):

$$\begin{split} \frac{\sqrt{2}}{2} \cdot \frac{1}{2} &= \frac{\sqrt{2}}{2} \left(\frac{1}{\sqrt{2}} \right)^2 \\ &= \frac{\sqrt{2}}{2} \sin^2 \left(\frac{\pi}{4} \right) \\ &\triangleq \frac{\sqrt{2}}{2} \phi \left(\frac{1}{2} \right) \\ &= \frac{\sqrt{2}}{2} \sqrt{2} \sum_{n \in \mathbb{Z}} h_n \sin^2 \left(\frac{\pi}{2} (1-n) \right) \mathbb{I}_{[0:2]} (1-n) \\ &= h_0 \sin^2 \left(\frac{\pi}{2} (1-0) \right) \mathbb{I}_{[0:2]} (1-0) + h_1 \sin^2 \left(\frac{\pi}{2} (1-1) \right) \mathbb{I}_{[0:2]} (1-1) \\ &+ h_2 \sin^2 \left(\frac{\pi}{2} (1-2) \right) \mathbb{I}_{[0:2]} (1-2) \\ &= h_0 \cdot 1 \cdot 1 + h_1 \cdot 0 \cdot 1 + h_2 (-1) \cdot 0 \\ &= h_0 \end{split}$$

$$\begin{split} \frac{\sqrt{2}}{2} \cdot 1 &= \frac{\sqrt{2}}{2}(1)^2 \\ &= \frac{\sqrt{2}}{2} \sin^2 \left(\frac{\pi}{2}\right) \\ &\triangleq \frac{\sqrt{2}}{2} \phi(1) \\ &= \frac{\sqrt{2}}{2} \sqrt{2} \sum_{n \in \mathbb{Z}} \mathsf{h}_n \sin^2 \left(\frac{\pi}{2}(2-n)\right) \mathbb{1}_{[0:2]}(2-n) \\ &= \mathsf{h}_0 \sin^2 \left(\frac{\pi}{2}(2-0)\right) \mathbb{1}_{[0:2]}(2-0) + \mathsf{h}_1 \sin^2 \left(\frac{\pi}{2}(2-1)\right) \mathbb{1}_{[0:2]}(2-1) \\ &+ \mathsf{h}_2 \sin^2 \left(\frac{\pi}{2}(2-2)\right) \mathbb{1}_{[0:2]}(2-2) \end{split}$$

ⓒ ⓑ ⑤

=
$$h_0 \cdot 0 \cdot 1 + h_1 \cdot 1 \cdot 1 + h_2 \cdot 0 \cdot 1$$

= h_1


$$\begin{split} \frac{\sqrt{2}}{2} \cdot \frac{1}{2} &= \frac{\sqrt{2}}{2} \left(\frac{1}{-\sqrt{2}} \right)^2 \\ &= \frac{\sqrt{2}}{2} \sin^2 \left(\frac{3\pi}{4} \right) \\ &\triangleq \frac{\sqrt{2}}{2} \phi \left(\frac{3}{2} \right) \\ &= \frac{\sqrt{2}}{2} \sqrt{2} \sum_{n \in \mathbb{Z}} h_n \sin^2 \left(\frac{\pi}{2} (3-n) \right) \mathbb{I}_{[0:2]} (3-n) \\ &= h_0 \sin^2 \left(\frac{\pi}{2} (3-0) \right) \mathbb{I}_{[0:2]} (3-0) + h_1 \sin^2 \left(\frac{\pi}{2} (3-1) \right) \mathbb{I}_{[0:2]} (3-1) \\ &+ h_2 \sin^2 \left(\frac{\pi}{2} (3-2) \right) \mathbb{I}_{[0:2]} (3-2) \\ &= h_0 \cdot (-1) \cdot 0 + h_1 \cdot 0 \cdot 1 + h_2 1 \cdot 1 \\ &= h_2 \end{split}$$

(e) These values for (h_0, h_1, h_2) are valid for the knot locations $x = \frac{1}{2}$, x = 1, and $x = \frac{3}{2}$, **but** they don't satisfy the *dilation equation* (Theorem 4.1 page 60). In particular, $\phi(x) \neq \sum_{n} h_n \mathbf{DT}^n \phi(x)$

$$\phi(x) \neq \sum_{n \in \mathbb{Z}} \mathsf{h}_n \mathbf{D} \mathbf{T}^n \phi(x)$$

(see the diagram in item (3c) page 75)

Counterexample 4.3 (raised sine). ¹⁶ Let a function f be defined in terms of a shifted cosine function (Definition G.1 page 155) as follows:



 $^{\circ}$ Proof: Let $\mathbb{1}_A(x)$ be the *set indicator function* (Definition 3.2 page 40) on a set A.

1. Proof that $\phi(x)$ forms a partition of unity:

$$\sum_{n \in \mathbb{Z}} \mathbf{T}^n \phi(x) = \sum_{n \in \mathbb{Z}} \mathbf{T}^n \phi(x+1)$$
$$= \sum_{n \in \mathbb{Z}} \phi(x+1-n)$$

by Proposition 3.1 page 40

by Definition 3.3 page 40

¹⁶ Proakis (2001) pages 560–561

$$= \sum_{n \in \mathbb{Z}} \frac{1}{2} \{1 + \cos[\pi(|x - 1 + 1 - n|)]\} \mathbb{1}_{[0:2)}(x + 1 - n) \qquad \text{by definition of } \phi(x)$$

$$= \sum_{n \in \mathbb{Z}} \frac{1}{2} \{1 + \cos[\pi(|x - n|)]\} \mathbb{1}_{[-1:1)}(x - n) \qquad \text{by Definition 3.2 page 40}$$

$$= \sum_{n \in \mathbb{Z}} \frac{1}{2} \left\{1 + \cos\left[\frac{\pi}{\beta}\left(|x - n| - \frac{1 - \beta}{2}\right)\right]\right\} \mathbb{1}_{[-1:1)}(x - n) \Big|_{\beta = 1}$$

$$= 1 \qquad \text{by Example $\ref{eq: page ?\ref{eq: pag$$

- 2. Proof that $\phi(x) \notin \text{span} \{ \mathbf{DT}^n \phi(x) | n \in \mathbb{Z} \}$ (and so does not generate an *MRA*):
 - (a) Note that the *support* (Definition 4.4 page 70) of ϕ is $supp \phi = [0:2]$.
 - (b) Therefore, the *support* of (h_n) is $supp(h_n) = \{0, 1, 2\}$ (Theorem 4.7 page 70).
 - (c) So if $\phi(x)$ is an *MRA*, we only need to compute $\{h_0, h_1, h_2\}$ (the rest would be 0). Here would be the values of $\{h_1, h_2, h_3\}$:

$$\begin{aligned} \phi(x) &= \sum_{n \in \mathbb{Z}} \mathsf{h}_n \mathbf{D} \mathbf{T}^n \phi(x) \\ &= \sum_{n \in \mathbb{Z}} \mathsf{h}_n \mathbf{D} \mathbf{T}^n \; \frac{1}{2} \bigg\{ \; 1 + \cos[\pi(|x-1|)] \bigg\} \, \mathbb{1}_{[0:2]}(x) \qquad \qquad \text{by definition of } \phi(x) \\ &= \sum_{n \in \mathbb{Z}} \mathsf{h}_n \; \frac{\sqrt{2}}{2} \bigg\{ \; 1 + \cos[\pi(|2x-1-n|)] \bigg\} \, \mathbb{1}_{[0:2]}(2x-n) \qquad \qquad \text{by Definition 3.3 page 40} \\ &= \sum_{n=0}^2 \mathsf{h}_n \; \frac{\sqrt{2}}{2} \bigg\{ \; 1 + \cos[\pi(|2x-1-n|)] \bigg\} \, \mathbb{1}_{[0:2]}(2x-n) \qquad \qquad \text{by Theorem 4.7 page 70} \end{aligned}$$

(d) The values of (h_0, h_1, h_2) can be conveniently calculated at the knot locations $x = \frac{1}{2}$, x = 1, and $x = \frac{3}{2}$ (see the diagram in item (3c) page 75):

$$\frac{1}{2} = \sum_{n=0}^{2} h_n \frac{\sqrt{2}}{2} \left\{ 1 + \cos[\pi(|2x - 1 - n|)] \right\} \mathbb{1}_{[0:2]}(2x - n) \Big|_{x = \frac{1}{2}}$$

$$= h_0 \frac{\sqrt{2}}{2} \left\{ 1 + \cos[1 - 1 - 0] \right\}$$

$$= h_0 \sqrt{2}$$

$$\Rightarrow h_0 = \frac{\sqrt{2}}{4}$$

$$1 = \sum_{n=0}^{2} h_{n} \frac{\sqrt{2}}{2} \left\{ 1 + \cos[\pi(|2x - 1 - n|)] \right\} \mathbb{1}_{[0:2]}(2x - n) \Big|_{x=1}$$

$$= h_{1} \frac{\sqrt{2}}{2} \left\{ 1 + \cos[2 - 1 - 1] \right\}$$

$$= h_{1} \sqrt{2}$$

$$\implies h_{1} = \frac{\sqrt{2}}{2}$$

$$\frac{1}{2} = \sum_{n=0}^{2} \mathsf{h}_{n} \frac{\sqrt{2}}{2} \left\{ 1 + \cos[\pi(|2x - 1 - n|)] \right\} \mathbb{1}_{[0:2]} (2x - n) \bigg|_{x=0}$$



$$= h_2 \frac{\sqrt{2}}{2} \left\{ 1 + \cos[1 - 1 - 0] \right\}$$

$$= h_2 \sqrt{2}$$

$$\implies h_2 = \frac{\sqrt{2}}{4}$$

(e) These values for (h_0, h_1, h_2) are valid for the knot locations $x = \frac{1}{2}$, x = 1, and $x = \frac{3}{2}$, **but** they don't satisfy the *dilation equation* (Theorem 4.1 page 60). In particular (see diagram), $\phi(x) \neq \sum_{n \in \mathbb{Z}} \mathsf{h}_n \mathbf{DT}^n \phi(x)$.

Example 4.3 (2 coefficient case/Haar wavelet system/order 0 B-spline wavelet system). ¹⁷ Let $(L_{\mathbb{R}}^2, (V_j), (W_j), \phi, \psi, (h_n), (g_n))$ be an *wavelet system*.

E X

$$\begin{cases} 1. & \text{supp}\phi(x) = [0:1] \\ 2. & admissibility condition \\ 3. & partition of unity \end{cases} & \text{(Theorem 4.7 page 70)} \quad \text{and} \\ 3. & partition of unity} \end{cases} \Longrightarrow \begin{cases} \frac{n + h_n}{0} \\ \frac{\sqrt{2}}{2} \\ 1 + \frac{\sqrt{2}}{2} \\ \text{other} = 0 \end{cases}$$

NPROOF:

- 1. Proof that (1) \implies that only h_0 and h_1 are non-zero: by Theorem 4.7 page 70.
- 2. Proof for values of h_0 and h_1 :
 - (a) Method 1: Under the constraint of two non-zero scaling coefficients, a scaling function design is fully constrained using the *admissibility equation* (Theorem 4.3 page 66) and the *partition of unity* constraint. The partition of unity formed by $\phi(x)$ is illustrated in Example **??** (page **??**). Here are the equations:

$$h_0 + h_1 = \sqrt{2}$$
 (admissibility equation Theorem 4.3 page 66)
 $h_0 - h_1 = 0$ (partition of unity/zero at -1 Theorem 4.8 page 72)
Here are the calculations for the coefficients:

$$(h_0+h_1)+(h_0-h_1)=2h_0 \hspace{1cm} =\sqrt{2} \hspace{1cm} \text{(add two equations together)}$$

$$(h_0+h_1)-(h_0-h_1)=2h_1 \hspace{1cm} =\sqrt{2} \hspace{1cm} \text{(subtract second from first)}$$

₽

 \Rightarrow

¹⁷ Haar (1910), Wojtaszczyk (1997) pages 14–15 ("Sources and comments")



The bottom line for mathematicians is that the architecture has to be right. In all the mathematics that I did, the essential point was to find the right architecture. It's like building a bridge. Once the main lines of the structure are right, then the details miraculously fit. The problem is the overall design. ♥

Freeman Dyson (1923–2020), physicist and mathematician ¹

5.1 Introduction

5.1.1 What are wavelets?

In Fourier analysis, continuous dilations (Definition 3.3 page 40) of the complex exponential (Definition G.4 page 160) form a basis (Definition 2.7 page 14) for the space of square integrable functions $\boldsymbol{L}^2_{\mathbb{R}}$ (Definition D.1 page 141) such that

$$\mathcal{L}_{\mathbb{R}}^{2}=\operatorname{span}\left\{ \mathbf{D}_{\omega}e^{ix}|_{\omega\in\mathbb{R}}
ight\} .$$

In Fourier series analysis (Theorem N.1 page 248), discrete dilations of the complex exponential form a basis for $L^2_{\mathbb{R}}(0:2\pi)$ such that

$$\mathbf{L}_{\mathbb{R}}^{2}(0:2\pi) = \operatorname{span}\left\{\left.\mathbf{D}_{i}e^{ix}\right|_{j\in\mathbb{Z}}\right\}.$$

In Wavelet analysis, for some *mother wavelet* (Definition 5.1 page 81) $\psi(x)$,

$$\boldsymbol{L}_{\mathbb{R}}^{2} = \operatorname{span} \left\{ \mathbf{D}_{\omega} \mathbf{T}_{\tau} \psi(x) | \omega, \tau \in \mathbb{R} \right\}.$$

However, the ranges of parameters ω and τ can be much reduced to the countable set \mathbb{Z} resulting in a *dyadic* wavelet basis such that for some mother wavelet $\psi(x)$,

$$\tilde{\mathbf{L}}_{\mathbb{R}}^{2} = \operatorname{span}\left\{\mathbf{D}^{j}\mathbf{T}^{n}\psi(x)|j,n\in\mathbb{Z}\right\}.$$

This text deals almost exclusively with dyadic wavelets. Wavelets that are both dyadic and com-


image: http://www.isepp.org/Media/Speaker%20Images/95-96%20Images/dyson.jpg

¹ quote: Albers and Dyson (1994) page 20

pactly supported have the attractive feature that they can be easily implemented in hardware or software by use of the *Fast Wavelet Transform* (Figure O.1 page 253).


5.1.2 Analyses

An analysis can be partially characterized by its order structure with respect to an order relation such as the set inclusion relation \subseteq . Most transforms have a very simple M-n order structure, as illustrated to the right. The M-n lattices for $n \ge 3$ are modular but not distributive. Analyses typically have one subspace that is a scaling subspace; and this subspace is often simply a family of constants (as is the case with Fourier Analysis).




A special characteristic of wavelet analysis is that there is not just one scaling subspace, but an entire sequence of scaling subspaces. These scaling subspaces are *linearly ordered* with respect to the ordering relation \subseteq . In wavelet theory, this structure is called a *multiresolution analysis*, or MRA (Definition 4.1 page 54).


The MRA was introduced by Stéphane G. Mallat in 1989. The concept of a scaling space was perhaps first introduced by Taizo Iijima in 1959 in Japan, and later as the *Gaussian Pyramid* by Burt and Adelson in the 1980s in the West.²



A second special characteristic of wavelet analysis is that it's order structure with respect to the \subseteq relation is not a simple M-n lattice (as is with the case of Fourier and other analyses). Rather, it is a lattice of the form illustrated to the right. This lattice is non-complemented, non-distributive, non-modular, and non-Boolean (Proposition 5.1 page 83).



The wavelet subspace structure is similar in form to that of the *Primorial numbers*, illustrated to the right by a *Hasse diagram*.



An analysis can be represented using three different structures:

- ① sequence of subspaces
- ② sequence of basis coefficients
- 3 sequence of basis vectors

 $^3 \subseteq Sloane(2014) \langle http://oeis.org/A002110 \rangle$



 $^{^2}$ Mallat (1989) page 70, ☐ Iijima (1959), ☐ Burt and Adelson (1983), ☐ Adelson and Burt (1981), ☐ Lindeberg (1993), ☐ Alvarez et al. (1993), ☐ Guichard et al. (2012) pages 23–24 (\$3.2.1 Scale-Space Extrema), ☐ Guichard et al. (20xx) pages 77–78 (\$5.2.1 Scale-Space Extrema), ☐ Weickert (1999) (historical survey)

5.2. DEFINITION Daniel J. Greenhoe page 81

These structures are isomorphic to each other, and can therefore be used interchangeably.




Figure 5.1 (page 91) illustrate the order structures of some analyses, including two wavelet analyses:

Definition 5.2

The term "wavelet" comes from the French word "ondelette", meaning "small wave". And in essence, wavelets are "small waves" (as opposed to the "long waves" of Fourier analysis) that form a basis for the Hilbert space L_{\square}^2 .

Definition 5.1. ⁶ Let **T** and **D** be as defined in Definition 3.3 page 40. A function $\psi(x)$ in $L^2_{\mathbb{R}}$ is a wavelet function for $L^2_{\mathbb{R}}$ if $\{\mathbf{D}^j\mathbf{T}^n\psi|_{j,n\in\mathbb{Z}}\}$ is a RIESZ BASIS for $L^2_{\mathbb{R}}$.

In this case, ψ is also called the **mother wavelet** of the basis $\{\mathbf{D}^{j}\mathbf{T}^{n}\psi|_{j,n\in\mathbb{Z}}\}$. The sequence of subspaces $(W_j)_{j\in\mathbb{Z}}$ is the wavelet analysis induced by ψ , where each subspace W_j is defined

 $\mathbf{W}_{j} riangleq ext{span} \left\{ \left. \mathbf{D}^{j} \mathbf{T}^{n} \psi \right| n \in \mathbb{Z}
ight\} .$

A wavelet analysis ((\boldsymbol{W}_j)) is often constructed from a multiresolution analysis (Definition 4.1 page 54) ((\boldsymbol{V}_j)) under the relationship

where $\hat{+}$ is subspace addition (*Minkowski addition*). $V_{j+1} = V_j + W_j,$

By this relationship alone, (W_i) is in no way uniquely defined in terms of a multiresolution analysis (V_i) . In general there are many possible complements of a subspace V_i . To uniquely define such a wavelet subspace, one or more additional constraints are required. One of the most common additional constraints is *orthogonality*, such that V_i and W_i are orthogonal to each other.

Dilation equation 5.3

Suppose $(\mathbf{T}^n \psi)_{n \in \mathbb{Z}}$ is a basis for \mathbf{W}_0 . By Definition 5.1 page 81, the wavelet subspace \mathbf{W}_0 is contained in the scaling subspace V_1 . By Definition 4.1 page 54, the sequence $(\mathbf{DT}^n \phi)_{n \in \mathbb{Z}}$ is a basis for V_1 . Because W_0 is contained in V_1 , the sequence $(\mathbf{DT}^n \phi)_{n \in \mathbb{Z}}$ is also a basis for W_0 .

Theorem 5.1 (wavelet dilation equation). Let $\left(\mathbf{L}_{\mathbb{R}}^{2},\,\left(\!\!\left(\mathbf{\textit{V}}_{j}\right)\!\!\right),\,\phi,\,\left(\!\!\left(h_{n}\right)\!\!\right)\right)$ be a multiresolution system (Definition 4.3 page 63) and $(\mathbf{W}_j)_{j\in\mathbb{Z}}$ be a WAVELET ANALYSIS (Definition 5.1 page 81) with respect to

[■] Wojtaszczyk (1997) page 17 (Definition 2.1)





Strang and Nguyen (1996) page ix,
 Atkinson and Han (2009) page 191

 $\left(m{L}_{\mathbb{R}}^{2},\,\left(m{V}_{j}
ight) ,\,\phi,\,\left(h_{n}
ight)
ight) \,and\,with\,$ WAVELET FUNCTION ψ (Definition 5.1 page 81).

$$\exists (g_n)_{n \in \mathbb{Z}} \text{ such that } \psi = \sum_{n \in \mathbb{Z}} g_n \mathbf{D} \mathbf{T}^n \phi$$

[♠]Proof:

$$\begin{split} & \psi \in \textit{\textbf{W}}_0 \\ & \subseteq \textit{\textbf{V}}_1 \\ & = \text{span} \, (\mathbf{D}\mathbf{T}^n \phi(x))_{n \in \mathbb{Z}} \\ & \Longrightarrow \, \exists \, (g_n)_{n \in \mathbb{Z}} \quad \text{such that} \quad \psi = \sum_{n \in \mathbb{Z}} g_n \mathbf{D}\mathbf{T}^n \phi \end{split}$$

by Definition 5.1 page 81 by Definition 5.1 page 81 by Definition 4.1 page 54 (MRA)

₽

A wavelet system (next definition) consists of two subspace sequences:

- 4.1 page 54) provides "coarse" approximations of a function in $L^2_{\mathbb{R}}$ at different "scales" or resolutions.
- $\overset{\circ}{\text{A}}$ A wavelet analysis (W_j) provides the "detail" of the function missing from the approximation provided by a given scaling subspace (Definition 5.1 page 81).

Definition 5.2. $Let\left(\mathbf{L}_{\mathbb{R}}^{2},\,\left(\mathbf{V}_{j}\right),\,\phi,\,\left(h_{n}\right)\right)$ be a multiresolution system (Definition 4.1 page 54) and $\left(\mathbf{W}_{j}\right)_{j\in\mathbb{Z}}$ a wavelet analysis (Definition 5.1 page 81) with respect to $\left(\mathbf{V}_{j}\right)_{j\in\mathbb{Z}}$. $Let\left(g_{n}\right)_{n\in\mathbb{Z}}$ be a sequence of coefficients.

D E F A wavelet system is the tuple $(L_{\mathbb{R}}^2, (V_j), (W_j), \phi, \psi, (h_n), (g_n))$ and the sequence $(g_n)_{n \in \mathbb{Z}}$ that satisfies the equation $\psi = \sum_{n \in \mathbb{Z}} g_n \mathbf{D} \mathbf{T}^n \phi$

is the wavelet coefficient sequence.


Remark 5.1. The pair of coefficient sequences $((h_n), (g_n))$ generates the scaling function $\phi(x)$ (Definition 4.1 page 54) and the wavelet function $\psi(x)$ (Definition 5.1 page 81). These functions in turn generate the multiresolution analysis (V_j) (Definition 4.1 page 54) and the wavelet analysis (W_j) (Definition 5.1 page 81). Therefore, the coefficient sequence pair $((h_n), (g_n))$ totally defines a wavelet system $(L_{\mathbb{R}}^2, (V_j), (W_j), \phi, \psi, (h_n), (g_n))$ (Definition 5.2 page 82).

Furthermore, especially in the case of orthonormal wavelets, the wavelet coefficient sequence $(g_n)_{n\in\mathbb{Z}}$ is often defined in terms of the scaling coefficient sequence $(h_n)_{n\in\mathbb{Z}}$ in a very simple and straightforward manner. Therefore, in the case of an orthonormal wavelet system, the coefficient scaling sequence $(h_n)_{n\in\mathbb{Z}}$ often totally defines the entire wavelet system. And in this case, designing a wavelet system is only a matter of finding a handful of scaling coefficients (h_1, h_2, \ldots, h_n) ...because once you have these, you can generate everything else.



Order structure 5.4

The wavelet system $\left(L_{\mathbb{R}}^2,\,\left(\!\!\left(\boldsymbol{V}_{j}^{}\right)\!\!\right),\,\left(\!\!\left(\boldsymbol{W}_{j}^{}\right)\!\!\right),\,\phi,\,\psi,\,\left(\!\!\left(h_{n}^{}\right)\!\!\right),\,\left(\!\!\left(g_{n}^{}\right)\!\!\right)$ (Definition 5.2 page 82) together with the set inclusion relation \subseteq forms an ordered set, illustrated to the right by a Hasse diagram.



Proposition 5.1. Let $(L_{\mathbb{R}}^2, (V_j), (W_j), \phi, \psi, (h_n), (g_n))$ be a wavelet system with order relation \subseteq . The lattice $\mathbf{L} \triangleq ((\mathbf{V}_i), (\mathbf{W}_i), \vee, \wedge; \subseteq)$ has the following properties:

- 1. L is nondistributive.
- 2. L is NONMODULAR. R
 - 3. L is noncomplemented.
 - 4. **L** is nonBoolean.



- 1. Proof that *L* is *nondistributive*:
 - (a) L contains the N5 lattice.
 - (b) Because *L* contains the *N5* lattice, *L* is *nondistributive*.
- 2. Proof that *L* is *nonmodular* and *nondistributive*:
 - (a) L contains the N5 lattice.

3. Proof that *L* is *noncomplemented*:

(b) Because L contains the N5 lattice, L is nonmodular.

$$x' = y' = v' = z$$


$$z' = \{x, y, v\}$$

$$x'' = (x')'$$

$$= z'$$

$$= \{x, y, v\}$$

 $\neq x$



- 4. Proof that *L* is *nonBoolean*:
 - (a) L is nondistributive (item (1)).
 - (b) Because *L* is *nondistributive*, it is *nonBoolean*.

Subspace algebraic structure 5.5

Theorem 5.2. Let $\left(\mathbf{L}_{\mathbb{R}}^{2},\,\left(\!\!\left(\mathbf{V}_{j}^{}\right)\!\!\right),\,\left(\!\!\left(\mathbf{W}_{j}^{}\right)\!\!\right),\,\phi,\,\psi,\,\left(\!\!\left(h_{n}^{}\right)\!\!\right),\,\left(\!\!\left(g_{n}^{}\right)\!\!\right)\right)$ be a WAVELET SYSTEM (Definition 5.2 page 82). Let $V_1 + V_2$ represent Minkowski addition of two subspaces V_1 and V_2 of a Hilbert space H.

$$\begin{array}{lll} \boldsymbol{L}_{\mathbb{R}}^{2} & = & \lim_{j \to \infty} \boldsymbol{V}_{j} & (\boldsymbol{L}_{\mathbb{R}}^{2} \text{ is equivalent to one very large scaling subspace}) \\ & = & \boldsymbol{V}_{j} + \boldsymbol{W}_{j} + \boldsymbol{W}_{j+1} + \boldsymbol{W}_{j+2} + \cdots & \left(\begin{array}{c} \boldsymbol{L}_{\mathbb{R}}^{2} \text{ is equivalent to one scaling space} \\ \text{and a sequence of wavelet subspaces} \end{array} \right) \\ & = & \cdots + \boldsymbol{W}_{-2} + \boldsymbol{W}_{-1} + \boldsymbol{W}_{0} + \boldsymbol{W}_{1} + \boldsymbol{W}_{2} + \cdots & (\boldsymbol{L}_{\mathbb{R}}^{2} \text{ is equivalent to a sequence of wavelet subspaces}) \end{array}$$

♥Proof:

1. Proof for (1):

$$L_{\mathbb{R}}^2 = \lim_{i \to \infty} V_i$$
 by Definition 4.1 page 54

2. Proof for (2):

$$\underbrace{V_{j} + W_{j}}_{V_{j+1}} + W_{j+1} + W_{j+2} + \cdots = \underbrace{V_{j+1} + W_{j+1}}_{V_{j+2}} + W_{j+2} + W_{j+3} + \cdots \\
= \underbrace{V_{j+2} + W_{j+2}}_{V_{j+3}} + W_{j+3} + W_{j+4} + \cdots \\
= \underbrace{V_{j+3} + W_{j+3}}_{V_{j+3}} + W_{j+4} + W_{j+5} + \cdots \\
= \underbrace{V_{j+3} + W_{j+3}}_{V_{j+4}} + W_{j+5} + W_{j+6} + \cdots \\
= \underbrace{V_{j+5} + W_{j+5}}_{V_{j+5}} + W_{j+6} + W_{j+6} + \cdots \\
= \lim_{j \to \infty} V_{j+5} + W_{j+5} + W_{j+6} + W_{j+6} + \cdots \\
= L_{\mathbb{R}}^{2}$$

3. Proof for (3):

$$L_{\mathbb{R}}^{2} = \underbrace{V_{0}}_{V_{-1} \hat{+} W_{-1}} + W_{0} \hat{+} W_{1} \hat{+} W_{2} \hat{+} W_{3} \hat{+} \cdots$$

$$= \underbrace{V_{-1}}_{V_{-2} \hat{+} W_{-2}} + W_{-1} \hat{+} W_{0} \hat{+} W_{1} \hat{+} W_{2} \hat{+} W_{3} \hat{+} \cdots$$

$$= \underbrace{V_{-2}}_{V_{-3} \hat{+} W_{-2}} + W_{-1} \hat{+} W_{0} \hat{+} W_{1} \hat{+} W_{2} \hat{+} W_{3} \hat{+} \cdots$$

$$= \underbrace{V_{-3}}_{V_{-4} \hat{+} W_{-3}} + W_{-2} \hat{+} W_{-1} \hat{+} W_{0} \hat{+} W_{1} \hat{+} W_{2} \hat{+} W_{3} \hat{+} \cdots$$

$$\vdots$$

$$= \cdots \hat{+} W_{-3} \hat{+} W_{-2} \hat{+} W_{-1} \hat{+} W_{0} \hat{+} W_{1} \hat{+} W_{2} \hat{+} W_{3} \hat{+} \cdots$$

₽

Remark 5.2. In the special case that two subspaces W_1 and W_2 are *orthogonal* to each other, then the *subspace addition* operation $W_1 + W_2$ is frequently expressed as $W_1 + W_2$. In the case of an *orthonormal wavelet system*, the expressions in Theorem 5.2 (page 83) could be expressed as


$$L_{\mathbb{R}}^{2} = \lim_{j \to \infty} V_{j}$$

$$= V_{j} \oplus W_{j} \oplus W_{j+1} \oplus W_{j+2} \oplus \cdots$$

$$= \cdots \oplus W_{-2} \oplus W_{-1} \oplus W_{0} \oplus W_{1} \oplus W_{2} \oplus \cdots$$

5.6 Necessary conditions

Theorem 5.3 (quadrature conditions in "time"). Let $(L_{\mathbb{R}}^2, (V_j), (W_j), \phi, \psi, (h_n), (g_n))$ be a wavelet system (Definition 5.2 page 82).



[♠]Proof:

- 1. Proof for (1): by Theorem 4.4 page 68.
- 2. Proof for (2):

$$\langle \psi \mid \mathbf{T}^n \psi \rangle = \left\langle \sum_{m \in \mathbb{Z}} g_m \mathbf{D} \mathbf{T}^m \phi \mid \mathbf{T}^n \sum_{k \in \mathbb{Z}} g_k \mathbf{D} \mathbf{T}^k \phi \right\rangle \qquad \text{by } wavelet \, dilation \, equation} \qquad \text{(Theorem 5.1 page 81)}$$

$$= \sum_{m \in \mathbb{Z}} g_m \sum_{k \in \mathbb{Z}} g_k^* \left\langle \mathbf{D} \mathbf{T}^m \phi \mid \mathbf{T}^n \mathbf{D} \mathbf{T}^k \phi \right\rangle \qquad \text{by properties of } \langle \triangle \mid \nabla \rangle \qquad \text{(Definition C.9 page 124)}$$

$$= \sum_{m \in \mathbb{Z}} g_m \sum_{k \in \mathbb{Z}} g_k^* \left\langle \phi \mid (\mathbf{D} \mathbf{T}^m)^* \mathbf{T}^n \mathbf{D} \mathbf{T}^k \phi \right\rangle \qquad \text{by def. of operator adjoint} \qquad \text{(Proposition C.3 page 125)}$$

$$= \sum_{m \in \mathbb{Z}} g_m \sum_{k \in \mathbb{Z}} g_k^* \left\langle \phi \mid (\mathbf{D} \mathbf{T}^m)^* \mathbf{D} \mathbf{T}^{2n} \mathbf{T}^k \phi \right\rangle \qquad \text{by Proposition 3.5 page 42}$$

$$= \sum_{m \in \mathbb{Z}} g_m \sum_{k \in \mathbb{Z}} g_k^* \left\langle \phi \mid \mathbf{T}^{*m} \mathbf{D}^* \mathbf{D} \mathbf{T}^{2n} \mathbf{T}^k \phi \right\rangle \qquad \text{by operator star-algebra prop.} \qquad \text{(Theorem C.13 page 126)}$$

$$= \sum_{m \in \mathbb{Z}} g_m \sum_{k \in \mathbb{Z}} g_k^* \left\langle \phi \mid \mathbf{T}^{-m} \mathbf{D}^{-1} \mathbf{D} \mathbf{T}^{2n} \mathbf{T}^k \phi \right\rangle \qquad \text{by Proposition 3.7 page 43}$$

$$= \sum_{m \in \mathbb{Z}} g_m \sum_{k \in \mathbb{Z}} g_k^* \left\langle \phi \mid \mathbf{T}^{2n-m+k} \phi \right\rangle$$

3. Proof for (3):

$$\begin{split} &\langle \phi \,|\, \mathbf{T}^n \psi \rangle \\ &= \left\langle \sum_{m \in \mathbb{Z}} h_m \mathbf{D} \mathbf{T}^m \phi \,|\, \mathbf{T}^n \sum_{k \in \mathbb{Z}} g_k \mathbf{D} \mathbf{T}^k \phi \right\rangle \qquad \text{by Theorem 4.1 page 60} \qquad \text{and Theorem 5.1 page 81} \\ &= \sum_{m \in \mathbb{Z}} h_m \sum_{k \in \mathbb{Z}} g_k^* \left\langle \mathbf{D} \mathbf{T}^m \phi \,|\, \mathbf{T}^n \mathbf{D} \mathbf{T}^k \phi \right\rangle \qquad \text{by properties of } \langle \triangle \,|\, \nabla \rangle \qquad \text{(Definition C.9 page 124)} \\ &= \sum_{m \in \mathbb{Z}} h_m \sum_{k \in \mathbb{Z}} g_k^* \left\langle \phi \,|\, (\mathbf{D} \mathbf{T}^m)^* \,\mathbf{T}^n \mathbf{D} \mathbf{T}^k \phi \right\rangle \qquad \text{by definition of operator adjoint} \qquad \text{(Proposition C.3 page 125)} \\ &= \sum_{m \in \mathbb{Z}} h_m \sum_{k \in \mathbb{Z}} g_k^* \left\langle \phi \,|\, (\mathbf{D} \mathbf{T}^m)^* \,\mathbf{D} \mathbf{T}^{2n} \mathbf{T}^k \phi \right\rangle \qquad \text{by Proposition 3.5 page 42} \\ &= \sum_{m \in \mathbb{Z}} h_m \sum_{k \in \mathbb{Z}} g_k^* \left\langle \phi \,|\, \mathbf{T}^{*m} \mathbf{D}^* \mathbf{D} \mathbf{T}^{2n} \mathbf{T}^k \phi \right\rangle \qquad \text{by operator star-algebra properties} \qquad \text{(Theorem C.13 page 126)} \\ &= \sum_{m \in \mathbb{Z}} h_m \sum_{k \in \mathbb{Z}} g_k^* \left\langle \phi \,|\, \mathbf{T}^{-m} \mathbf{D}^{-1} \mathbf{D} \mathbf{T}^{2n} \mathbf{T}^k \phi \right\rangle \qquad \text{by Proposition 3.7 page 43} \\ &= \sum_{m \in \mathbb{Z}} h_m \sum_{k \in \mathbb{Z}} g_k^* \left\langle \phi \,|\, \mathbf{T}^{2n-m+k} \phi \right\rangle \end{aligned}$$



 \Box

Proposition 5.2. Let $(L_{\mathbb{R}}^2, (V_j), (W_j), \phi, \psi, (h_n), (g_n))$ be a wavelet system. Let $\tilde{\phi}(\omega)$ and $\tilde{\psi}(\omega)$ be the Fourier transforms (Definition 1.2 page 196) of $\phi(x)$ and $\psi(x)$, respectively. Let $\check{g}(\omega)$ be the Discrete time Fourier transform (Definition M.1 page 237) of (g_n) .

$$\tilde{\psi}(\omega) = \frac{\sqrt{2}}{2} \, \tilde{\mathsf{g}}\!\left(\frac{\omega}{2}\right) \tilde{\phi}\!\left(\frac{\omega}{2}\right)$$

№PROOF:

$$\begin{split} \tilde{\psi}(\omega) &\triangleq \tilde{\mathbf{F}} \psi \\ &= \tilde{\mathbf{F}} \sum_{n \in \mathbb{Z}} g_n \mathbf{D} \mathbf{T}^n \phi & \text{by } wavelet \, dilation \, equation} \\ &= \sum_{n \in \mathbb{Z}} g_n \tilde{\mathbf{F}} \mathbf{D} \mathbf{T}^n \phi & \text{by Corollary 3.1 page 47} \\ &= \sum_{n \in \mathbb{Z}} g_n \mathbf{D}^{-1} \tilde{\mathbf{F}} \mathbf{T}^n \phi & \text{by Corollary 3.1 page 47} \\ &= \sum_{n \in \mathbb{Z}} g_n \sqrt{2} (\mathbf{D}^{-1} e^{-i\omega n}) (\mathbf{D}^{-1} \tilde{\mathbf{F}} \phi) & \text{by Proposition 3.4 page 41} \\ &= \sqrt{2} \left(\mathbf{D}^{-1} \sum_{n \in \mathbb{Z}} g_n e^{-i\omega n} \right) (\mathbf{D}^{-1} \tilde{\mathbf{F}} \phi) & \text{by definition of } \tilde{\mathbf{F}} & \text{(Definition M.1 page 237)} \\ &= \sqrt{2} \sqrt{2} \tilde{\mathbf{g}} \left(\frac{\omega}{2} \right) \frac{\sqrt{2}}{2} \tilde{\phi} \left(\frac{\omega}{2} \right) & \text{by property of } \mathbf{D} & \text{(Proposition 3.2 page 41)} \\ &= \frac{\sqrt{2}}{2} \, \tilde{\mathbf{g}} \left(\frac{\omega}{2} \right) \tilde{\phi} \left(\frac{\omega}{2} \right) & \text{by property of } \mathbf{D} & \text{(Proposition 3.2 page 41)} \end{split}$$

Theorem 5.4 (next) presents the *quadrature* necessary conditions of a wavelet system. These relations simplify dramatically in the special case of an *orthonormal wavelet system* (Theorem M.4 page 243).

Theorem 5.4 (Quadrature conditions in "frequency"). ⁷ Let $(L_{\mathbb{R}}^2, (V_j), (W_j), \phi, \psi, (h_n), (g_n))$ be a wavelet system. Let $\tilde{\mathbf{x}}(\omega)$ be the discrete time Fourier transform (Definition M.1 page 237) for a sequence $(x_n)_{n\in\mathbb{Z}}$ in $\boldsymbol{\mathcal{C}}_{\mathbb{R}}^2$. Let $\tilde{\mathbf{S}}_{\phi\phi}(\omega)$ be the auto-power spectrum (Definition P.3 page 255) of ϕ , $\tilde{\mathbf{S}}_{\psi\psi}(\omega)$ be the auto-power spectrum of ψ , and $\tilde{\mathbf{S}}_{\phi\psi}(\omega)$ be the cross-power spectrum of ϕ and ψ .

PROOF:

1. Proof for (1): by Theorem 4.5 page 68.

⁷ Chui (1992) page 135, Goswami and Chan (1999) page 110



2. Proof for (2):

$$\begin{split} &2\tilde{\mathbf{S}}_{\psi\psi}(2\omega)\triangleq 2(2\pi)\sum_{n\in\mathbb{Z}}|\tilde{\psi}(2\omega+2\pi n)|^2\\ &=2(2\pi)\sum_{n\in\mathbb{Z}}\left|\frac{\sqrt{2}}{2}\check{\mathbf{g}}\left(\frac{2\omega+2\pi n}{2}\right)\tilde{\phi}\left(\frac{2\omega+2\pi n}{2}\right)\right|^2 \quad \text{by Lemma 4.1 page 61}\\ &=2\pi\sum_{n\in\mathbb{Z}_e}\left|\check{\mathbf{g}}\left(\frac{2\omega+2\pi n}{2}\right)\right|^2\left|\tilde{\phi}\left(\frac{2\omega+2\pi n}{2}\right)\right|^2+\\ &2\pi\sum_{n\in\mathbb{Z}_e}\left|\check{\mathbf{g}}\left(\frac{2\omega+2\pi n}{2}\right)\right|^2\left|\tilde{\phi}\left(\frac{2\omega+2\pi n}{2}\right)\right|^2\\ &=2\pi\sum_{n\in\mathbb{Z}}\left|\check{\mathbf{g}}(\omega+2\pi n)|^2\left|\tilde{\phi}(\omega+2\pi n)\right|^2+2\pi\sum_{n\in\mathbb{Z}}\left|\check{\mathbf{g}}(\omega+2\pi n+\pi)|^2\left|\tilde{\phi}(\omega+2\pi n+\pi)\right|^2\\ &=2\pi\sum_{n\in\mathbb{Z}}\left|\check{\mathbf{g}}(\omega)\right|^2\left|\tilde{\phi}(\omega+2\pi n)\right|^2+2\pi\sum_{n\in\mathbb{Z}}\left|\check{\mathbf{g}}(\omega+\pi)\right|^2\left|\tilde{\phi}(\omega+2\pi n+\pi)\right|^2\\ &=|\check{\mathbf{g}}(\omega)|^2\left(2\pi\sum_{n\in\mathbb{Z}}\left|\tilde{\phi}(\omega+2\pi n)\right|^2+\right)|\check{\mathbf{g}}(\omega+\pi)|^2\left(2\pi\sum_{n\in\mathbb{Z}}\left|\tilde{\phi}(\omega+\pi+2\pi n)\right|^2\right)\\ &=|\check{\mathbf{g}}(\omega)|^2\tilde{\mathbf{S}}_{\phi\phi}(\omega)+|\check{\mathbf{g}}(\omega+\pi)|^2\tilde{\mathbf{S}}_{\phi\phi}(\omega+\pi) \quad \text{by Theorem P.1 page 255} \end{split}$$

3. Proof for (3):

$$\begin{split} 2\tilde{\mathbf{S}}_{\phi\psi}(2\omega) &= 2(2\pi) \sum_{n \in \mathbb{Z}} \tilde{\phi}(2\omega + 2\pi n) \tilde{\psi}^*(2\omega + 2\pi n) \\ &= 2(2\pi) \sum_{n \in \mathbb{Z}} \frac{\sqrt{2}}{2} \check{\mathbf{h}} (\omega + \pi n) \, \tilde{\phi} (\omega + \pi n) \, \frac{\sqrt{2}}{2} \, \check{\mathbf{g}}^* (\omega + \pi n) \, \tilde{\phi}^* (\omega + \pi n) \quad \text{by Lemma 4.1 page 61} \\ &= 2\pi \sum_{n \in \mathbb{Z}_0} \check{\mathbf{h}} (\omega + \pi n) \, \check{\mathbf{g}}^* (\omega + \pi n) \, \big| \tilde{\phi} (\omega + \pi n) \big|^2 \\ &= 2\pi \sum_{n \in \mathbb{Z}_0} \check{\mathbf{h}} (\omega + \pi n) \, \check{\mathbf{g}}^* (\omega + \pi n) \, \big| \tilde{\phi} (\omega + \pi n) \big|^2 \\ &+ 2\pi \sum_{n \in \mathbb{Z}_0} \check{\mathbf{h}} (\omega + \pi n) \, \check{\mathbf{g}}^* (\omega + \pi n) \, \big| \tilde{\phi} (\omega + \pi n) \big|^2 \\ &= 2\pi \sum_{n \in \mathbb{Z}} \check{\mathbf{h}} (\omega + 2\pi n + \pi) \, \check{\mathbf{g}}^* (\omega + 2\pi n + \pi) \, \big| \tilde{\phi} (\omega + 2\pi n + \pi) \big|^2 \\ &+ 2\pi \sum_{n \in \mathbb{Z}} \check{\mathbf{h}} (\omega + 2\pi n) \, \check{\mathbf{g}}^* (\omega + 2\pi n) \, \big| \tilde{\phi} (\omega + 2\pi n) \big|^2 \\ &= 2\pi \sum_{n \in \mathbb{Z}} \check{\mathbf{h}} (\omega + \pi) \, \check{\mathbf{g}}^* (\omega + \pi) \, \big| \tilde{\phi} (\omega + 2\pi n + \pi) \big|^2 + 2\pi \sum_{n \in \mathbb{Z}} \check{\mathbf{h}} (\omega) \, \check{\mathbf{g}}^* (\omega) \, \big| \tilde{\phi} (\omega + 2\pi n) \big|^2 \\ &= \check{\mathbf{h}} (\omega) \, \check{\mathbf{g}}^* (\omega) \, \bigg(2\pi \sum_{n \in \mathbb{Z}} \big| \tilde{\phi} (\omega + 2\pi n) \big|^2 \bigg) \\ &+ \check{\mathbf{h}} (\omega + \pi) \, \check{\mathbf{g}}^* (\omega + \pi) \, \bigg(2\pi \sum_{n \in \mathbb{Z}} \big| \tilde{\phi} (\omega + \pi + 2\pi n) \big|^2 \bigg) \\ &= \check{\mathbf{h}} (\omega) \check{\mathbf{g}}^* (\omega) \, \bigg(2\pi \sum_{n \in \mathbb{Z}} \big| \tilde{\phi} (\omega + 2\pi n) \big|^2 \bigg) + \check{\mathbf{h}} (\omega + \pi) \, \check{\mathbf{g}}^* (\omega + \pi) \, \bigg(2\pi \sum_{n \in \mathbb{Z}} \big| \tilde{\phi} (\omega + \pi + 2\pi n) \big|^2 \bigg) \\ &= \check{\mathbf{h}} (\omega) \check{\mathbf{g}}^* (\omega) \, \bigg(2\pi \sum_{n \in \mathbb{Z}} \big| \tilde{\phi} (\omega + 2\pi n) \big|^2 \bigg) + \check{\mathbf{h}} (\omega + \pi) \, \check{\mathbf{g}}^* (\omega + \pi) \, \bigg(2\pi \sum_{n \in \mathbb{Z}} \big| \tilde{\phi} (\omega + \pi + 2\pi n) \big|^2 \bigg) \\ &= \check{\mathbf{h}} (\omega) \check{\mathbf{g}}^* (\omega) \, \bigg(2\pi \sum_{n \in \mathbb{Z}} \big| \tilde{\phi} (\omega + 2\pi n) \big|^2 \bigg) + \check{\mathbf{h}} (\omega + \pi) \, \check{\mathbf{g}}^* (\omega + \pi) \, \bigg(2\pi \sum_{n \in \mathbb{Z}} \big| \tilde{\phi} (\omega + \pi + 2\pi n) \big|^2 \bigg) \\ &= \check{\mathbf{h}} (\omega) \check{\mathbf{g}}^* (\omega) \, \bigg(2\pi \sum_{n \in \mathbb{Z}} \big| \tilde{\phi} (\omega + 2\pi n) \big|^2 \bigg) + \check{\mathbf{h}} (\omega + \pi) \, \check{\mathbf{g}}^* (\omega + \pi) \, \bigg(2\pi \sum_{n \in \mathbb{Z}} \big| \tilde{\phi} (\omega + \pi + 2\pi n) \big|^2 \bigg) \\ &= \check{\mathbf{h}} (\omega) \check{\mathbf{g}}^* (\omega) \, \bigg(2\pi \sum_{n \in \mathbb{Z}} \big| \tilde{\phi} (\omega + 2\pi n) \big|^2 \bigg) + \check{\mathbf{h}} (\omega + \pi) \, \bigg(2\pi \sum_{n \in \mathbb{Z}} \big| \tilde{\phi} (\omega + \pi + \pi) \big|^2 \bigg) + \check{\mathbf{h}} (\omega + \pi) \, \bigg(2\pi \sum_{n \in \mathbb{Z}} \big| \tilde{\phi} (\omega + \pi + \pi) \big|^2 \bigg) + \check{\mathbf{h}} (\omega + \pi) \, \bigg(2\pi \sum_{n \in \mathbb{Z}} \big| \tilde{\phi} (\omega + \pi) \big|^2 \bigg) + \check{\mathbf{h}} (\omega + \pi) \, \bigg(2\pi \sum_{n \in \mathbb{Z}} \big| \tilde{\phi} (\omega + \pi) \big|^2 \bigg) + \check{\mathbf{h}} (\omega + \pi$$

 \Box

ⓒ ⓑ ⑤



5.7 Sufficient condition

In this text, an often used sufficient condition for designing the *wavelet coefficient sequence* (g_n) (Definition 5.2 page 82) is the *conjugate quadrature filter condition* (Definition J.9 page 215). It expresses the sequence (g_n) in terms of the *scaling coefficient sequence* (Definition 4.3 page 63) and a "shift" integer N as $g_n = \pm (-1)^n h_{N-n}^*$. The CQF condition has the following "nice" properties:

- 1. Given a scaling coefficient sequence (h_n) (Definition 4.3 page 63), it is extremely simple to compute the wavelet coefficient sequence (g_n) (Definition 5.2 page 82).
- 2. If $\{\mathbf{T}\phi\}$ of a *wavelet system* $(\mathbf{L}_{\mathbb{R}}^2, (V_j), (W_j), \phi, \psi, (h_n), (g_n))$ (Definition 5.2 page 82) is *orthonormal* and $((g_n), (h_n), N)$ satisfies the *CQF condition*, then $\{\mathbf{T}^n\psi\}$ is also *orthonormal*.
- 3. If $\{\mathbf{T}\phi\}$ of a wavelet system $\left(\boldsymbol{L}_{\mathbb{R}}^2, \left(\!\!\left(\boldsymbol{V}_j\!\!\right)\!\!\right), \left(\!\!\left(\boldsymbol{W}_j\!\!\right)\!\!\right), \phi, \psi, \left(\!\!\left(h_n\!\!\right)\!\!\right), \left(\!\!\left(g_n\!\!\right)\!\!\right)$ (Definition 5.2 page 82) is orthonormal and $\left(\left(g_n\!\!\right), \left(h_n\!\!\right), N\!\!\right)$ satisfies the CQF condition, then the wavelet subspace \boldsymbol{W}_0 is orthonormal to the scaling subspace \boldsymbol{V}_0 ($\boldsymbol{W}_0 \perp \boldsymbol{V}_0$).

Theorem 5.5. Let $(L_{\mathbb{R}}^2, (V_j), (W_j), \phi, \psi, (h_n), (g_n))$ be a WAVELET SYSTEM (Definition 5.2 page 82). Let $\check{\mathbf{g}}(\omega)$ be the DTFT (Definition M.1 page 237) and $\check{\mathbf{g}}(z)$ the Z-TRANSFORM (Definition J.4 page 208) of (g_n) .

$$\underbrace{g_n = \pm (-1)^n h_{N-n}^*, \, N \in \mathbb{Z}}_{\text{CONJUGATE QUADRATURE FILTER}} \iff \underbrace{\breve{g}(\omega)}_{\text{CONJUGATE QUADRATURE FILTER}} \iff \underbrace{\breve{g}(\omega)}_{n \in \mathbb{Z}} = \pm (-1)^N e^{-i\omega N} \check{\mathsf{h}}^*(\omega + \pi) \Big|_{\omega = \pi}}_{\omega = \pi} \qquad (1)$$

$$\iff \underbrace{\breve{g}(z)}_{z=-1} = \sqrt{2} \qquad (2)$$

$$\iff \check{\mathsf{g}}(z)\Big|_{z=-1} = \sqrt{2} \qquad (3)$$

$$\iff \check{\mathsf{g}}(\omega)\Big|_{\omega = \pi} = \sqrt{2} \qquad (4)$$

♥Proof:

- 1. Proof that CQF \iff (1): by Theorem J.5 page 215
- 2. Proof that $CQF \Longrightarrow (4)$:

$$\begin{split} \breve{\mathbf{g}}(\pi) &= \breve{\mathbf{g}}(\omega) \Big|_{\omega = \pi} \\ &= \pm (-1)^N e^{-i\omega N} \breve{\mathbf{h}}^*(\omega + \pi) \Big|_{\omega = \pi} \qquad \text{by } CQF \, theorem \qquad \text{(Theorem J.5 page 215)} \\ &= \pm (-1)^N e^{-i\pi N} \breve{\mathbf{h}}^*(2\pi) \\ &= \pm (-1)^N (-1)^N \breve{\mathbf{h}}^*(0) \qquad \text{by } DTFT \, periodicity} \qquad \text{(Proposition M.1 page 237)} \\ &= \sqrt{2} \qquad \qquad \text{by } admissibility \, condition} \end{split}$$

3. Proof that (2) \iff (3) \iff (4): by Proposition M.4 page 240

5.8. SUPPORT SIZE Daniel J. Greenhoe page 89

5.8 Support size

Theorem 5.6 (support size). ⁸ Let $(L_{\mathbb{R}}^2, (V_j), (W_j), \phi, \psi, (h_n), (g_n))$ be a WAVELET SYSTEM (Definition 5.2 page 82) induced by the CQF CONDITIONS (Theorem 5.5 page 88). Let supple be the support of a function f (Definition 4.4 page 70).


$$\sup_{\mathbf{M}} \phi = \sup_{\mathbf{M}} \mathbf{M}$$

$$\sup_{\mathbf{M}} \psi = \left[\frac{N - (n_2 - n_1)}{2} : \frac{N + (n_2 - n_1)}{2} \right]$$

NPROOF:

- 1. Proof that $spp \phi = sph$: by Theorem 4.7 (page 70)
- 2. Proof that $spp \psi = \left[\frac{N (n_2 n_1)}{2} : \frac{N + (n_2 n_1)}{2} \right]$:

Example 5.1. Here are some examples using Daubechies wavelet functions.



⁸ 🍠 Mallat (1999) pages 243–244

page 90

Daniel J. Greenhoe

CHAPTER 5. WAVELET STRUCTURES

5.9 Examples

No further examples of wavelets are presented in this section. Examples begin in the next chapter which is about a property called the *partition of unity*. Other design constraints leading to wavelets with more "powerful" properties include *vanishing moments* (Chapter ?? page ??), *orthonormality*, *compact support*, and *minimum phase* (Definition J.5 page 211).

5.9. EXAMPLES Daniel J. Greenhoe page 91




Figure 5.1: examples of the order structures of some analyses

5.9. EXAMPLES Daniel J. Greenhoe page 93





ALGEBRAIC STRUCTURES



In this book, learned reader, you have the rules of algebra... It unties the knot not only where one term is equal to another or two to one but also where two are equal to two or three to one.... this most abstruse and unsurpassed treasury of the entire arithmetic being brought to light and, as in a theater, exposed to the sight of all...

Gerolamo Cardano (1501–1576), Italian mathematician, physician, and astrologer ¹

A set together with one or more operations forms several standard mathematical structures:

 $group \supseteq ring \supseteq commutative ring \supseteq integral domain \supseteq field$

Definition A.1. ² Let X be a set and \diamondsuit : $X \times X \to X$ be an operation on X.

```
The pair (X, \diamondsuit) is a group if

1. \exists e \in X such that e \diamondsuit x = x \diamondsuit e = x \forall x \in X (IDENTITY element) and

2. \exists (-x) \in X such that (-x) \diamondsuit x = x \diamondsuit (-x) = e \forall x \in X (INVERSE element) and

3. x \diamondsuit (y \diamondsuit z) = (x \diamondsuit y) \diamondsuit z \forall x, y, z \in X (ASSOCIATIVE)
```

Definition A.2. 3 Let $+: X \times X \to X$ and $\cdot: X \times X \to X$ be operations on a set X. Furthermore, let the operation \cdot also be represented by juxtapostion as in $a \cdot b \equiv ab$.

```
The triple (X, +, \cdot) is a ring if
             1. (X, +) is a group.
                                                                         (additive group)
                                                                                                              and
D
E
             2. \quad x(yz)
                               = (xy)z
                                                         \forall x, y, z \in X
                                                                         (ASSOCIATIVE with respect to ·)
                                                                                                              and
             3. x(y+z) = (xy) + (xz)
                                                        \forall x, y, z \in X
                                                                         (· is left distributive over +)
                                                                                                              and
             4. (x + y)z = (xz) + (yz)
                                                        \forall x,y,z \in X
                                                                         (· is right distributive over +).
```

Definition A.3. ⁴

```
1 quote: Cardano (1545) page 1
image: http://en.wikipedia.org/wiki/Image:Cardano.jpg
2 Durbin (2000) page 29
3 Durbin (2000) pages 114-115
4 Durbin (2000) page 118
```

and and

and

D

E

```
A triple (X, +, \cdot) is a commutative ring if
```

1. $(X, +, \cdot)$ is a RING

and

 $2. \quad xy = yx$

 $\forall x,y \in X$ (COMMUTATIVE).

Definition A.4. ⁵ *Let R be a* COMMUTATIVE RING (*Definition A.3 page 95*).

	A function	$ \cdot $ in	$\mathbb{R}^{\mathbb{R}}$	is an absolute	value (or	modulus) if
D	1.	x	\geq	0	$x \in \mathbb{R}$	(NON-NEGATIVE)
E	2.	x	=	$0 \iff x = 0$	$x \in \mathbb{R}$	(NONDEGENERATE)

3. $|xy| = |x| \cdot |y|$ $x,y \in \mathbb{R}$ (Homogeneous / Submultiplicative) 4. $|x+y| \le |x| + |y|$ $x,y \in \mathbb{R}$ (Subadditive / Triangle inequality)

Definition A.5. 6

The structure $F \triangleq (X, +, \cdot, 0, 1)$ is a **field** if

1. $(X,+,\cdot)$ is a ring

ng)

and

 $2. \quad xy = yx$

 $\forall x,y \in X$ (commutative with respect to ·) and

3. $(X \setminus \{0\}, \cdot)$ is a group

(group with respect to ·).

Definition A.6. ⁷ *Let* $V = (F, +, \cdot)$ *be a* VECTOR SPACE *and* $\otimes : V \times V \rightarrow V$ *be a vector-vector multiplication operator.*

An **algebra** is any pair (V, \otimes) that satisfies $(\otimes \text{ is represented by juxtaposition})$

	1.	(ux)y	=	u(xy)	$\forall u, x, y \in V$	(ASSOCIATIVE)	and
D E F	2.	u(x + y)	=	(ux) + (uy)	$\forall u, x, y \in V$	(LEFT DISTRIBUTIVE)	and
F	3.	(u+x)y	=	(uy) + (xy)	$\forall u, x, y \in V$	(RIGHT DISTRIBUTIVE)	and
	4.	$\alpha(xy)$	=	$(\alpha \mathbf{x})\mathbf{y} = \mathbf{x}(\alpha \mathbf{y})$	$\forall x,y \in V \ and \ \alpha \in F$	(SCALAR COMMUTATIVE)	

⁷ Abramovich and Aliprantis (2002) page 3, Michel and Herget (1993) page 56



⁵ Cohn (2002) page 312

APPENDIX B	
	l
	I INFAR SUBSPACES

B.1 Subspaces of a linear space

Linear spaces (Definition C.1 page 111) can be decomposed into a collection of *linear subspaces* (Definition B.1 page 98). Often such a collection along with an *order relation* forms a *lattice*.





Figure B.1: lattice of subspaces of \mathbb{R}^3 (Example B.1 page 97)

Example B.1. The 3-dimensional Euclidean space \mathbb{R}^3 contains the 2-dimensional xy-plane and xz-plane subspaces, which in turn both contain the 1-dimensional x-axis subspace. These subspaces are illustrated in the figure to the right and in Figure B.1 (page 97).



E X D E F

Definition B.1. Let $\Omega \triangleq (X, +, \cdot, (\mathbb{F}, +, \dot{\times}))$ be a LINEAR SPACE (Definition C.1 page 111).

Every *linear space* (Definition C.1 page 111) **X** has at least two *linear subspaces*—itself and **0** (Proposition B.1 page 98), called the *trivial linear space*. The *linear span* (Definition 2.2 page 9) of every subset of a linear linear space is a subspace (Proposition B.2 page 99). Every *linear subspace* contains the "zero" vector 0, and is *convex*.

Proposition B.1.
2
 Let $X \triangleq (X, +, \cdot, (\mathbb{F}, \dotplus, \dot{\times}))$ and $0 \triangleq (\{0\}, +, \cdot, (\mathbb{F}, \dotplus, \dot{\times}))$.

Proposition B.1. 2 Let $X \triangleq (X, +, \cdot, (\mathbb{F}, \dotplus, \dot{\times}))$ and $0 \triangleq (\{0\}, +, \cdot, (\mathbb{F}, \dotplus, \dot{\times}))$.

Proposition B.1. 2 Let $X \triangleq (X, +, \cdot, (\mathbb{F}, \dotplus, \dot{\times}))$ and $0 \triangleq (\{0\}, +, \cdot, (\mathbb{F}, \dotplus, \dot{\times}))$.

Proposition B.1. 2 Let $X \triangleq (X, +, \cdot, (\mathbb{F}, \dotplus, \dot{\times}))$ and $0 \triangleq (\{0\}, +, \cdot, (\mathbb{F}, \dotplus, \dot{\times}))$.

 $\$ Proof: For a structure to be a linear subspace of X, it must satisfy the requirements of Definition B.1 (page 98).

- 1. Proof that $\{0\}$ is a linear subspace:
 - (a) Note that $\{0\} \neq \emptyset$.
 - (b) Proof that $x, y \in \{0\} \implies x + y \in \{0\}$:

$$x + y = 0 + 0$$
 by $x, y \in \{0\}$ hypothesis $= 0$ $\in \{0\}$

(c) Proof that $x \in \{0\}, \alpha \in \mathbb{F} \implies \alpha x \in \{0\}$:

$$\alpha x = \alpha \mathbb{O}$$
 by $x \in \{0\}$ hypothesis by definition of $0 \in \{0\}$

- 2. Proof that Ω is a linear subspace of itself:
 - (a) Proof that $X \neq \emptyset$:

$$X \neq \emptyset$$

(b) Proof that $x, y \in X \implies x + y \in X$:

$$x + y \in \{0\}$$
 because $+: X \times X \to X$ (X is closed under vector addition)

(c) Proof that $x \in X$, $\alpha \in \mathbb{F} \implies \alpha x \in X$:

$$\alpha x \in X$$
 because $\cdot : \mathbb{F} \times X \to X$ (X is closed under scalar-vector multiplication)

² Michel and Herget (1993) pages 81–83, A Haaser and Sullivan (1991) page 43



¹ Michel and Herget (1993) page 81 ⟨Definition 3.2.1⟩, Berberian (1961) page 13 ⟨Definition I.5.1⟩, Halmos (1958) page 16

Proposition B.2. 3 Let $X \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{x}))$ be a LINEAR SPACE (Definition C.1 page 111). Let span be the LINEAR SPAN of a set Y in X.

$$\left\{ \begin{array}{l} Y \text{ is } a \text{ subset of the set } X \\ (Y \subseteq X) \end{array} \right\} \implies \left\{ \begin{array}{l} \text{span} Y \text{ is } a \text{ linear subspace of } \textbf{\textit{X}}. \end{array} \right\}$$

Proposition B.3. 4 Let $X \triangleq (X, +, \cdot, (\mathbb{F}, +, \dot{x}))$ be a LINEAR SPACE and \emptyset the zero vector of X.

[♠]Proof:

$$Y \text{ is a } subspace \implies \exists (\alpha y) \in Y \quad \forall \alpha \in \mathbb{F}$$
 by Definition B.1 page 98
$$\implies \exists 0 \in Y \qquad \qquad \text{because } \alpha = 0 \in \mathbb{F}$$

$$Y$$
 is a linear subspace $\implies x + y \in Y \ \forall x, y \in Y$
 $\implies \lambda x + (1 - \lambda)y \in Y \ \forall x, y \in Y$
 $\implies Y$ is $convex$


Definition B.2. ⁵ Let $\mathbf{X} \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}))$ and $\mathbf{Y} \triangleq (Y, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}))$ be LINEAR SUBSPACES (Definition B.1 page 98) of a LINEAR SPACE (Definition C.1 page 111) $\Omega \triangleq (\Omega, +, \cdot, (\mathbb{F}, +, \dot{x}))$.

Example B.2. Some examples of operations on subspaces in \mathbb{R}^3 are illustrated next: Remark B.1.

Notice the similarities between the properties of linear subspaces in a linear space (Proposition B.4 page 100) and the properties of closed sets in a topological space:

linear subspaces	closed sets
0	Ø
Ω	$egin{array}{c} arnothing \ oldsymbol{\Omega} \end{array}$
X + Y	$X \cup Y$
N	_
$\bigcap_{n=1} X_n$	$\bigcap_{\gamma \in \Gamma} X_{\gamma}$
n=1	$\gamma \in \Gamma$

One key difference is that the union of two linear subspaces is not in general a linear subspace. For example, if x is the vector [10] in the x direction linear subspace of \mathbb{R}^2 and y is the vector [01] in the y direction linear subspace, then x + y is not in the union of the two linear subspaces (it is not on the x axis or y axis but rather at (1,1).




³ Michel and Herget (1993) page 86

⁴ Michel and Herget (1993) page 81


⁵ Wedderburn (1907) page 79

Michel and Herget (1993) page 82

In general, the set of all linear subspaces of a linear space Ω is *not* closed under the subspace union (v) operation; that is, the union of of two linear subspaces is *not* necessarily a linear subspace. However the set is closed under Minkowski sum (+) and subspace intersection (A). Proposition B.4 (next) shows four useful objects are always subspaces. Some of these in Euclidean space \mathbb{R}^3 are illustrated to the right.



Proposition B.4. 7 Let \boldsymbol{X} be a LINEAR SPACE (Definition C.1 page 111).



 $^{\circ}$ Proof: For a structure to be a linear subspace of X, it must satisfy the requirements of Definition B.1 (page 98).

- 1. Proof that $X_1 + X_2 + \cdots + X_N$ is a *linear subspace* (proof by induction):
 - (a) proof for N = 1 case: by left hypothesis.
 - (b) proof for N = 2 case:
 - i. proof that $X_1 + X_2 \neq \emptyset$:

$$\begin{aligned} \textbf{\textit{X}}_1 \, \hat{+} \, \textbf{\textit{X}}_2 &= \big\{ \textbf{\textit{v}} + \textbf{\textit{w}} | \textbf{\textit{v}} \in \textbf{\textit{X}}_1 \text{ and } \textbf{\textit{w}} \in \textbf{\textit{Y}} \big\} \\ &\supseteq \big\{ \textbf{\textit{v}} + \textbf{\textit{w}} | \textbf{\textit{v}} \in \{ \mathbb{0} \} \subseteq \textbf{\textit{X}}_1 \text{ and } \textbf{\textit{w}} \in \{ \mathbb{0} \} \subseteq \textbf{\textit{X}}_2 \big\} \\ &= \{ \mathbb{0} + \mathbb{0} \} \\ &= \{ \mathbb{0} \} \\ &\neq \varnothing \end{aligned}$$
by Definition B.2 page 99

ii. proof that $x, y \in X_1 + X_2 \implies x + y \in X_1 + X_2$:

$$x + y = (v_1 + w_1) + (v_2 + w_2)$$
 by $x, y \in X_1 + X_2$ hypothesis
$$= \underbrace{(v_1 + v_2)}_{\text{in } X_1} + \underbrace{(w_1 + w_2)}_{\text{in } X_2 \text{ because } X_2 \text{ is a linear subspace}}$$

$$\in \{v + w | v \in X_1 \text{ and } w \in Y\}$$
 by Definition B.2 page 99

iii. proof that $v \in X_1 + X_2$, $\alpha \in F \implies \alpha v \in X_1 + X_2$:

$$\alpha \mathbf{x} = \alpha(\mathbf{v}_1 + \mathbf{w}_1)$$
 by $\mathbf{x} \in \mathbf{X}_1 + \mathbf{X}_2$ hypothesis
$$= \underbrace{\alpha \mathbf{v}_1}_{\text{in } \mathbf{X}_1} + \underbrace{\alpha \mathbf{w}_1}_{\text{in } \mathbf{X}_2 \text{ because } \mathbf{X}_2 \text{ is a linear subspace}}$$

$$\in \left\{ \mathbf{v} + \mathbf{w} | \mathbf{v} \in \mathbf{X}_1 \text{ and } \mathbf{w} \in \mathbf{Y} \right\}$$

$$= \mathbf{X}_1 + \mathbf{X}_2$$
 by Definition B.2 page 99

(c) Proof that $[N \text{ case}] \implies [N+1 \text{ case}]$:

$$\mathbf{X}_1 \; \hat{+} \; \mathbf{X}_2 \; \hat{+} \; \cdots \; \hat{+} \; \mathbf{X}_{N+1} = \underbrace{\left(\mathbf{X}_1 \; \hat{+} \; \mathbf{X}_2 \; \hat{+} \; \cdots \; \hat{+} \; \mathbf{X}_N\right)}_{\text{linear subspace by } N \text{ case hypothsis}} \; \hat{+} \; \mathbf{X}_{N+1}$$

 \implies linear subspace by N = 2 case (item (1b) page 100)

⁷ Michel and Herget (1993) pages 81–83



- 2. Proof that $X_1 \cap X_2 \cap \cdots \cap X_N$ is a *linear subspace* (proof by induction):
 - (a) proof for N = 1 case: X_1 is a linear subspace by left hypothesis.
 - (b) Proof for N = 2 case:
 - i. proof that $X \cap Y \neq \emptyset$:

$$X \cap Y = \{x \in X | x \in X \text{ and } w \in Y\}$$

$$\supseteq \{x \in X | x \in \{0\} \subseteq X \text{ and } x \in \{0\} \subseteq Y\}$$

$$= \{0 + 0\}$$

$$= \{0\}$$

$$\neq \emptyset$$

ii. proof that $x, y \in X \cap Y \implies x + y \in X \cap Y$:

$$x, y \in X \cap Y \implies x, y \in X \text{ and } x, y \in Y$$

 $\implies x + y \in X \text{ and } x + y \in Y \text{ because } X \text{ and } Y \text{ are linear subspaces}$
 $\implies x + y \in X \cap Y$

iii. proof that $v \in X \cap Y$, $\alpha \in F \implies \alpha v \in X \cap Y$:

$$x \in X \cap Y \implies x \in X \text{ and } x \in Y$$

$$\implies \alpha x \in X \text{ and } \alpha x \in Y \qquad \text{because } X \text{ and } Y \text{ are linear subspaces}$$

$$\implies \alpha x \in X \cap Y$$

(c) Proof that $[N \text{ case}] \implies [N+1 \text{ case}]$:

$$X_1 \cap X_2 \cap \cdots \cap X_{N+1} = \underbrace{\left(X_1 \cap X_2 \cap \cdots \cap X_N\right)}_{\text{linear subspace by } N \text{ case hypothsis}} \cap X_{N+1}$$

$$\implies \text{linear subspace by } N = 2 \text{ case (item (2b) page 101)}$$

Every linear subspace contains the zero vector $\mathbb O$ (Proposition B.3 page 99). But if a pair of linear subspaces of a linear space **X** only have 0 in commmon, then any vector in **X** can be uniquely represented by a single vector from each of the two subspaces (next).

Theorem B.1. 8 Let $\mathbf{X} \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}))$ and $\mathbf{Y} \triangleq (Y, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}))$ be Linear subspaces (Definition B.1 page 98) of a LINEAR SPACE (Definition C.1 page 111) $\mathbf{\Omega} \triangleq (\Omega, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}))$.

$$X \cap Y = \{\emptyset\} \iff \begin{cases} \text{for every } \mathbf{u} \in \mathbf{X} + \mathbf{Y} \text{ there exist } \mathbf{x} \in X \text{ and } \mathbf{y} \in Y \text{ such that } \\ 1. \quad \mathbf{u} = \mathbf{x} + \mathbf{y} \qquad \text{and} \\ 2. \quad \mathbf{x} \text{ and } \mathbf{y} \text{ are UNIQUE.} \end{cases}$$

^ℚProof:

1. Proof that $X \cap Y = \{0\} \implies unique x, y$: Suppose that x and y are not unique, but rather $u = x_1 + y_1 = x_2 + y_2$ where $x_1, x_2 \in X$ and $y_1, y_2 \in Y$.

$$u = x_1 + y_1 = x_2 + y_2 \implies \underbrace{x_1 - x_2}_{\in X} = \underbrace{y_2 - y_1}_{\in Y}$$

$$\implies x_1 - x_2, y_2 - y_1 \in X \cap Y$$

$$\implies x_1 - x_2 = y_2 - y_1 = 0$$
 by left hypothesis
$$\implies x_1 = x_2 \quad \text{and} \quad y_2 = y_1$$

$$\implies x \text{ and } y \text{ are } unique$$

⁸ Michel and Herget (1993) page 83 (Theorem 3.2.12),

Kubrusly (2001) page 67 (Theorem 2.14)

Kubrusly (2001) page 67 (Theorem 2.14)



2. Proof that $X \cap Y = \{0\} \iff unique x, y$:

$$u = x + y$$

$$= x + y + y - y$$

$$= (x + y) + (y - y)$$

$$\Rightarrow x \text{ and } y \text{ are } not \text{ unique if } y \neq \emptyset$$

$$\Rightarrow y = \emptyset$$

$$\Rightarrow X \cap Y = \{\emptyset\}$$
borsome vector $y \in X \cap Y$
because $x \in X$ and $y \in X \cap Y$...

Theorem B.2. ⁹Let Ω be a linear subspace and 2^{Ω} the set of closed linear subspaces of Ω .

$$(2^{\Omega}, \, \hat{+}, \, \Omega, \, \mathbf{0}, \, \mathbf{\Omega}; \, \subseteq) \text{ is a LATTICE. In particular}$$

$$X \, \hat{+} \, X = X \qquad X \, \Omega \, X = X \qquad \forall X \in 2^{\Omega}$$

$$X \, \hat{+} \, Y = Y \, \hat{+} \, X \qquad X \, \Omega \, Y = Y \, \Omega \, X \qquad \forall X, Y \in 2^{\Omega}$$

$$(X \, \hat{+} \, Y) \, \hat{+} \, Z = X \, \hat{+} \, (Y \, \hat{+} \, Z) \qquad (X \, \Omega \, Y) \, \Omega \, Z = X \, \Omega \, (Y \, \Omega \, Z) \qquad \forall X, Y, Z \in 2^{\Omega}$$

$$X \, \hat{+} \, (X \, \Omega \, Y) = X \qquad X \, \Omega \, (X \, \hat{+} \, Y) = X \qquad \forall X, Y \in 2^{\Omega}$$

PROOF: These results follow directly from the properties of lattices.

B.2 Subspaces of an inner product space

Definition B.3. ¹⁰ Let $\Omega \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle | \nabla \rangle)$ be an INNER PRODUCT SPACE (Definition C.9 page 124).

The **orthogonal complement** A^{\perp} in Ω of a set $A \subseteq X$ is $A^{\perp} \triangleq \{x \in X | \langle x | y \rangle = 0 \quad \forall y \in A\}.$ The expression $A^{\perp \perp}$ is defined as $(A^{\perp})^{\perp}$.

Proposition B.5. 11 Let $(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{x}), \langle \triangle | \nabla \rangle)$ be an INNER PRODUCT SPACE (Definition C.9 page 124).

$$\begin{array}{c} \mathsf{P} \\ \mathsf{R} \\ \mathsf{P} \end{array} \quad A \subseteq B \quad \Longrightarrow \quad B^{\perp} \subseteq A^{\perp} \quad \forall A, B \in 2^{X} \quad \text{(antitone)} \end{array}$$

№PROOF:

$$B^{\perp} \triangleq \{x \in X \mid \langle x \mid y \rangle = 0 \quad \forall y \in B\}$$
 by definition of B^{\perp} (Definition B.3 page 102)
$$\subseteq \{x \in X \mid \langle x \mid y \rangle = 0 \quad \forall y \in A\}$$
 by definition of A^{\perp} (Definition B.3 page 102)

Every *linear space* **X** contains **0** and **X** as *linear subspaces* (Proposition B.1 page 98). If **X** is also an *inner product space*, then **0** and **X** are *orthogonal complements* of each other (next proposition).

¹¹ ■ Berberian (1961) page 60 (Theorem III.2.2), ■ Kubrusly (2011) page 326




⁹ Iturrioz (1985) pages 56–57

¹⁰ Berberian (1961) page 59 ⟨Definition III.2.1⟩, Michel and Herget (1993) page 382, Kubrusly (2001) page 328

₽

Proposition B.6. ¹² Let $\Omega \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle | \nabla \rangle)$ be an inner product space (Definition C.9 page 124) and \emptyset the vector additive identity element (Definition C.1 page 111) in Ω .



№ Proof:

$$\{ \mathbb{O} \}^{\perp} = \{ \boldsymbol{x} \in \boldsymbol{X} | \langle \boldsymbol{x} | \boldsymbol{y} \rangle = 0 \quad \forall \boldsymbol{y} \in \{ \mathbb{O} \} \}$$
 by definition of \bot (Definition B.3 page 102)
$$= \{ \boldsymbol{x} \in \boldsymbol{X} | \langle \boldsymbol{x} | \mathbb{O} \rangle = 0 \}$$

$$= \boldsymbol{X}$$

$$\boldsymbol{X}^{\perp} = \{ \boldsymbol{x} \in \boldsymbol{X} | \langle \boldsymbol{x} | \boldsymbol{y} \rangle = 0 \quad \forall \boldsymbol{y} \in \boldsymbol{X} \}$$
 by definition of \bot Definition B.3 page 102
$$= \{ \boldsymbol{x} \in \boldsymbol{X} | \langle \boldsymbol{x} | \boldsymbol{x} \rangle = 0 \}$$

$$= \{ \mathbb{O} \}$$

For any set A contained in a linear space X, $A^{\perp \perp}$ is a *linear subspace*, and it is the smallest linear subspace containing the set A ($A^{\perp \perp} = \operatorname{span} A$, next theorem). In the case that A is a *linear subspace* rather than just a subset, results simplify significantly (next corollary).

Theorem B.3. ¹³ Let $\Omega \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle | \nabla \rangle)$ be an inner product space (Definition C.9 page 124). Let $\operatorname{span} A$ be the span of a set A (Definition 2.2 page 9).

$$\left\{
\begin{array}{l}
A \text{ is a subset of } X \\
(A \subseteq X)
\end{array}
\right\} \implies
\left\{
\begin{array}{l}
1. \quad A \cap A^{\perp} = \begin{cases}
\{\emptyset\} & \text{if } \emptyset \in A \\
\emptyset & \text{if } \emptyset \notin A
\end{cases}
\right.$$

$$\left\{
\begin{array}{l}
A \text{ is a subset of } X \\
A \subseteq A^{\perp \perp} = \text{span } A
\end{cases}
\right.$$

$$A^{\perp} = A^{\perp \perp} = A^{\perp} = A^{-\perp} = (\text{span } A)^{\perp} \quad \text{and}$$

$$\left\{
\begin{array}{l}
A^{\perp} \text{ is a subspace of } \Omega
\end{array}
\right.$$

№PROOF:

1. Proof that $A \cap A^{\perp} = \cdots$:

$$A \cap A^{\perp} = \{ x \in X | x \in A \} \cap \{ x \in X | \langle x | y \rangle \quad \forall y \in A \}$$
 by definition of A^{\perp}

$$= \{ x \in X | x \in A \text{ and } \langle x | y \rangle \quad \forall y \in A \}$$

$$= \{ \begin{cases} \{0\} & \text{if } 0 \in A \\ \emptyset & \text{if } 0 \notin A \end{cases}$$

2. Proof that $A \subseteq A^{\perp \perp} = \operatorname{span} A$:

$$x \in A \implies \{x\}^{\perp \perp} \subseteq A^{\perp \perp}$$
$$\implies x \in \{x\}^{\perp \perp} \subseteq A^{\perp \perp}$$
$$\implies x \in A^{\perp \perp}$$

but

$$x \in A^{\perp \perp} \implies x \in A$$

Here is an example for the \implies part using the linear space \mathbb{R}^3 :

¹² ■ Kubrusly (2011) page 326, ■ Michel and Herget (1993) page 383

¹³ Michel and Herget (1993) page 383, Kubrusly (2011) page 326

- (a) Let $A \triangleq \{i\}$, where *i* is the unit vector on the x-axis.
- (b) Then $A^{\perp} = \{x \in X | x \in yz \text{ plane}\}.$
- (c) Then $A^{\perp \perp} = \{ x \in X | x \in x \text{ axis} \}.$
- (d) Therefore, $A \subseteq A^{\perp \perp}$
- 3. Proof for A^{\perp} equivalent expressions:
 - (a) Proof that $A^{\perp} = A^{\perp \perp \perp}$:

$$A^{\perp} \subseteq (A^{\perp})^{\perp \perp}$$
 by item (2)
 $= (A^{\perp \perp})^{\perp}$
 $= A^{\perp \perp \perp}$ by Definition B.3 page 102
 $A^{\perp \perp \perp} = (A^{\perp \perp})^{\perp}$ by Definition B.3 page 102
 $\subseteq A^{\perp}$ by item (2) and Proposition B.5 (page 102)

- (b) Proof that $A^{\perp \perp \perp} = (\operatorname{span} A)^{\perp}$: follows directly from item (2) $(A^{\perp \perp} = \operatorname{span} A)$.
- (c) Proof that $A^{\perp} = A^{\perp -}$:
 - i. Let (x_n) be an A^{\perp} -valued sequence that converges to the limit x in X.
 - ii. The limit point x must be in A^{\perp} because for all $y \in A$

$$\langle x \mid y \rangle = \langle \lim x_n \mid y \rangle$$
 by definition of the sequence (x_n)
= $\lim \langle x_n \mid y \rangle$
= 0 because (x_n) is A^{\perp} -valued

- iii. Because $\langle x | y \rangle = 0 \quad \forall y \in A, x \text{ is in } A^{\perp}.$
- iv. Because A^{\perp} contains all its limit points, and by the *Closed Set Theorem* (Theorem ?? page ??), it must be *closed* $(A^{\perp} = A^{\perp^{-}})$
- (d) Proof that $A^{\perp} = A^{-\perp}$:
 - i. Let $x \in A^{\perp}$ and $y \in A^{-}$.
 - ii. Let (y_n) be an A^{\perp} -valued sequence that converges in X to y.
 - iii. Thus $A^{\perp} \perp A^{-}$ because

$$\langle y \mid x \rangle = \langle \lim y_n \mid x \rangle$$
 by definition of (y_n)
= $\lim \langle y_n \mid x \rangle$
= 0 because (y_n) is A^{\perp} -valued

- iv. Because $A^{\perp} \perp A^{-}$, so $A^{\perp} \subseteq A^{\perp -}$.
- v. But $A^{\perp^-} \subseteq A^{\perp}$ because

$$A \subseteq A^- \implies A^{\perp^-} \subseteq A^{\perp}$$
 by *antitone* property (Proposition B.5 page 102)

- vi. And so $A^{\perp} = A^{\perp^{-}}$.
- 4. Proof that A^{\perp} is a **subspace** of Ω (must satisfy the conditions of Definition B.1 page 98):
 - (a) Proof that $A^{\perp} \neq \emptyset$: A^{\perp} has at least one element, the element 0...

$$\langle 0 \mid y \rangle = 0 \quad \forall y \in A$$
 by definition of 0 by definition of A^{\perp} (Definition B.3 page 102)

(b) Proof that $A^{\perp} \subseteq X$:

$$\mathbf{u} \in A^{\perp} \implies \mathbf{u} \in \{\mathbf{x} \in X \mid \langle \mathbf{x} \mid \mathbf{y} \rangle = 0 \quad \forall \mathbf{y} \in A\}$$
 by definition of A^{\perp} (Definition B.3 page 102)
$$\implies \mathbf{u} \in X$$
 by definition of sets

(c) Proof that $u, v \in A^{\perp} \implies (u + v) \in A^{\perp}$:

$$u, v \in A^{\perp} \implies \langle u \mid y \rangle = \langle v \mid y \rangle = 0 \quad \forall y \in A \quad \text{by definition of } A^{\perp} \text{ (Definition B.3 page 102)}$$

$$\implies \langle u \mid y \rangle + \langle v \mid y \rangle = 0 \quad \forall y \in A$$

$$\implies \langle u + v \mid y \rangle = 0 \quad \forall y \in A \quad \text{by } additive \text{ property of } \langle \triangle \mid \nabla \rangle \text{ (Definition C.9 page 124)}$$

$$\implies u + v \in A^{\perp} \quad \text{by definition of } A^{\perp} \text{ (Definition B.3 page 102)}$$

(d) Proof that $v \in \Omega \implies \alpha v \in A^{\perp}$:

$$oldsymbol{v} \in A^\perp \implies \langle oldsymbol{v} \,|\, oldsymbol{y} \rangle = 0 \qquad \forall oldsymbol{y} \in A \qquad \text{by definition of } A^\perp \text{ (Definition B.3 page 102)} \ \implies \alpha \,\langle oldsymbol{v} \,|\, oldsymbol{y} \rangle = \alpha \cdot 0 \qquad \forall oldsymbol{y} \in A \ \implies \langle \alpha oldsymbol{v} \,|\, oldsymbol{y} \rangle = 0 \qquad \forall oldsymbol{y} \in A \qquad \text{by } homogeneous \text{ property of } \langle \triangle \,|\, \nabla \rangle \text{ (Definition C.9 page 124)} \ \implies \alpha oldsymbol{v} \in A^\perp \qquad \qquad \text{by definition of } A^\perp \text{ (Definition B.3 page 102)} \$$


Corollary B.1. Let $\mathbf{X} \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle \mid \nabla \rangle)$ and $\mathbf{Y} \triangleq (Y, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle \mid \nabla \rangle)$ be inner product spaces. Let $\operatorname{span} Y$ be the span of the set Y (Definition 2.2 page 9).

 $\left\{ \begin{array}{l} \textbf{Y} \text{ is a linear subspace of } \textbf{X} \end{array} \right\} \qquad \Longrightarrow \qquad \left\{ \begin{array}{l} 1. \quad \textbf{Y} \cap \textbf{Y}^{\perp} &= \{0\} & \text{and} \\ 2. \quad \textbf{Y} &= \textbf{Y}^{\perp \perp} = \text{span} \textbf{Y} & \text{and} \\ 3. \quad \textbf{Y}^{\perp} &= \textbf{Y}^{\perp \perp \perp} & \text{and} \\ 4. \quad \textbf{Y}^{\perp} \text{ is a subspace of } \textbf{X} \end{array} \right\}$

♥Proof:

- 1. Proof that $Y \cap Y^{\perp} = \{0\}$: This follows from Theorem B.3 (page 103) and the fact that all subspaces contain the zero vector 0 (Proposition B.3 page 99).
- 2. Proof that $Y = Y^{\perp \perp} = \operatorname{span} Y$: This follows directly from Theorem B.3 (page 103).
- 3. Proof that $\mathbf{Y}^{\perp} = \mathbf{Y}^{\perp \perp \perp}$: This follows directly from Theorem B.3 (page 103).
- 4. Proof that Y^{\perp} is a **subspace** of X: This follows directly from Theorem B.3 (page 103).

Theorem B.4. ¹⁴ Let $\mathbf{Y} \triangleq (Y, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle \mid \nabla \rangle)$ and $\mathbf{Z} \triangleq (Z, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle \mid \nabla \rangle)$ be linear subspaces of an inner product space $\mathbf{\Omega} \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle \mid \nabla \rangle)$.



NPROOF:

$$x \in Y \cap Z \implies x \in Y \text{ and } x \in Z$$
 by definition of \cap
$$\implies \langle x \mid x \rangle = 0$$
 by hypothesis $Y \perp Z$
$$\implies x = \emptyset$$
 by $non\text{-isotropic}$ property of $\langle \triangle \mid \nabla \rangle$ (Definition C.9 page 124)

¹⁴ Kubrusly (2001) page 324

Frames and Bases Structure and Design [VERSION 021] thtps://github.com/dgreenhoe/pdfs/blob/master/msdframes.pdf

₽

Theorem B.5. ¹⁵ Let $\mathbf{Y} \triangleq (Y, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle \mid \nabla \rangle)$ and $\mathbf{Z} \triangleq (Z, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle \mid \nabla \rangle)$ be linear subspaces of an inner product space $\mathbf{\Omega} \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle \mid \nabla \rangle)$.

$$\left\{ \begin{array}{l} 1. \quad \textbf{Y} \perp \textbf{Z} \text{ and} \\ 2. \quad \textbf{x} \in \textbf{Y} + \textbf{Z} \end{array} \right\} \implies \left\{ \begin{array}{l} 1. \quad \textit{There exists } \textbf{y} \in \textbf{Y} \text{ and } \textbf{z} \in \textbf{Z} \text{ such that } \textbf{x} = \textbf{y} + \textbf{z} \text{ and} \\ 2. \quad \textbf{y} \text{ and } \textbf{z} \text{ are UNIQUE.} \end{array} \right\}$$

- **№** Proof:
 - 1. Proof that y and z exist: by definition of Minkowski addition operator $\hat{+}$ (Definition B.2 page 99).
 - 2. Proof that *y* and *z* are *unique*:
 - (a) Suppose $x = y_1 + z_1 = y_1 + z_2$ for $y_1, y_2 \in Y$ and $z_1, z_2 \in Z$.
 - (b) This implies

$$0 = x - x$$

$$= (y_1 + z_1) - (y_1 + z_2)$$

$$= (y_1 - y_2) + (z_1 - z_2)$$
in Z

- (c) Because $y_1 y_2 \in Y$, $z_1 z_2 \in Z$, $(y_1 y_2) + (z_1 + z_2) = 0$, and $(y_1 y_2 \mid z_1 z_2) = 0$, then by Theorem **??** (page **??**), $y_1 y_2 = 0$ and $z_1 z_2 = 0$.
- (d) This implies $y_1 = y_2$ and $z_1 = z_2$.
- (e) This implies y and z are unique.

₽

B.3 Subspaces of a Hilbert Space

Theorem B.6. ¹⁶ Let $\mathbf{H} \triangleq (X, +, \cdot, (\mathbb{F}, +, \dot{\times}), \langle \triangle \mid \nabla \rangle)$ be a Hilbert space (Definition **??** page **??**). Let Y be a subset of X, and let $d(x, Y) \triangleq \inf_{\mathbf{y} \in Y} \|\mathbf{x} - \mathbf{y}\|$.

- **№** Proof:
 - 1. Let $\delta \triangleq \inf\{x y | y \in Y\}$.
 - 2. Let $(y_n)_{n\in\mathbb{Z}}$ be a sequence such that $||x-y_n|| \to \delta$.

¹⁶ ■ Kubrusly (2001) page 330 〈Theorem 5.13〉, Aliprantis and Burkinshaw (1998) page 290 〈Theorem 33.6〉, Berberian (1961) page 68 〈Theorem III.5.1〉



¹⁵ Berberian (1961) page 61 (Theorem III.2.3)

3. Proof that (y_n) is *Cauchy*:

$$\lim_{m,n\to\infty} \|\mathbf{y}_{n} - \mathbf{y}_{m}\|^{2}$$

$$= \lim_{m,n\to\infty} \|(\mathbf{y}_{n} - x) + (x - \mathbf{y}_{m})\|^{2}$$

$$= \lim_{m,n\to\infty} \left\{ -\|(\mathbf{y}_{n} - x) - (x - \mathbf{y}_{m})\|^{2} + 2\|\mathbf{y}_{n} - x\|^{2} + 2\|x - \mathbf{y}_{m}\|^{2} \right\} \quad \text{by parallelogram law (page ??)}$$

$$= \lim_{m,n\to\infty} \left\{ -4 \left\| \frac{1}{2} \mathbf{y}_{n} + \frac{1}{2} \mathbf{y}_{m} - x \right\|^{2} + 2\|\mathbf{y}_{n} - x\|^{2} + 2\|x - \mathbf{y}_{m}\|^{2} \right\}$$

$$\leq \lim_{m,n\to\infty} \left\{ -4\delta^{2} + 2\|\mathbf{y}_{n} - x\|^{2} + 2\|x - \mathbf{y}_{m}\|^{2} \right\} \quad \text{by definition of } \delta \text{ (item (1))}$$

$$= -4\delta^{2} + \lim_{m,n\to\infty} \left\{ 2\|\mathbf{y}_{n} - x\|^{2} \right\} + \lim_{m,n\to\infty} \left\{ 2\|x - \mathbf{y}_{m}\|^{2} \right\}$$

$$= -4\delta^{2} + 2\delta^{2} + 2\delta^{2}$$

$$= 0$$

- 4. Proof that d(x, Y) = ||x y||: because (y_n) is *Cauchy* (item (1)) and by the *closed* hypothesis.
- 5. Proof that y is *unique*: Because in a metric space, the limit of a convergent sequence is *unique*.

Theorem B.7. Let $H \triangleq (X, +, \cdot, (\mathbb{F}, +, \dot{\times}), \langle \triangle | \nabla \rangle)$ be a HILBERT SPACE (Definition ?? page ??). Let $\mathsf{d}(x,Y) \triangleq \inf_{\mathbf{y} \in Y} \|\mathbf{x} - \mathbf{y}\|. \ Let \ \mathbf{Y} \triangleq \left(Y, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \left\langle \triangle \mid \nabla \right\rangle \right) \ and \ Y^{\perp} \ the \ \mathrm{ORTHOGONAL \ COMPLEMENT} \ of$

There exists $p \in Y$ such that 1. d(x, Y) = ||x - p|| and 2. p is UNIQUE and $\{ Y \text{ is } a \text{ SUBSPACE } of H \}$ 3. $x - p \in Y^{\perp}$.

Theorem B.8 (Projection Theorem). ¹⁸ Let
$$\mathbf{H} \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle | \nabla \rangle)$$
 be a Hilbert space.

THE H A SUBSPACE of \mathbf{H} \Rightarrow $Y + Y^{\perp} = \mathbf{H}$

^ℚProof:

$$\mathbf{Y} \,\hat{+} \, \mathbf{Y}^{\perp} = \begin{bmatrix} \mathbf{Y} \,\hat{+} \, \mathbf{Y}^{\perp} \end{bmatrix}^{\perp \perp}$$
 by Corollary B.1 page 105
 $= \begin{bmatrix} \mathbf{Y}^{\perp} \, \boldsymbol{\cap} \, \mathbf{Y}^{\perp \perp} \end{bmatrix}^{\perp}$ by Proposition B.5 (page 102)
 $= \{0\}^{\perp}$ by Corollary B.1 page 105
 $= \mathbf{H}$ by Proposition B.6 page 103

The inclusion relation \subseteq is an order relation on the set of subspaces of a linear space Ω .



¹⁷ Kubrusly (2001) page 330 (Theorem 5.13)

¹⁸ ■ Bachman and Narici (1966) page 172 (Theorem 10.8),
 Kubrusly (2001) page 339 (Theorem 5.20)

Proposition B.7. Let S be the set of subspaces of a linear space Ω . Let \subseteq be the inclusion relation.

P R P

 (S, \subseteq) is an **ordered set**

 \mathbb{Q} Proof: (S, \subseteq) is an *ordered set* and because

1.	$X \subseteq X$	∀ X ∈ S	(reflexive)	and	preorder	
2.	$X \subseteq Y$ and $Y \subseteq Z \implies X \subseteq Z$	$\forall X,Y,Z \in S$	(transitive)	and		\Rightarrow
3.	$X \subseteq Y$ and $Y \subseteq X \implies X = Y$	$\forall X,Y \in S$	(anti-symmetric)	_	,	

Theorem B.9. ¹⁹Let H be a Hilbert space and 2^H the set of closed linear subspaces of H.


	$(2^{\mathbf{H}}, \hat{+}, \hat{n}, 0, \mathbf{H}; \subseteq)$ is an orthomodular lattice. In particular				
	$1. \mathbf{X} \hat{+} \mathbf{X}^{\perp} = \mathbf{A}$		∀ X ∈ H	(COMPLEMENTED)	
Ţ	$2. \boldsymbol{X} \cap \boldsymbol{X}^{\perp} = 0$		∀ X ∈ H	(COMPLEMENTED)	
T H M	3. $(\boldsymbol{X}^{\perp})^{\perp} = \boldsymbol{\lambda}$		∀ X ∈ H	(INVOLUTORY)	
	4. $\mathbf{X} \leq \mathbf{Y} \implies \mathbf{Y}$		∀ X , Y ∈ H	(ANTITONE)	
	5. $X \leq Y \implies X$	$\mathbf{X} + (\mathbf{X}^{\perp} \cap \mathbf{Y}) = \mathbf{Y}$	$\forall \boldsymbol{X}, \boldsymbol{Y} \in X$	(ORTHOMODULAR IDENTITY)	

№ Proof:

- 1. Proof for *complemented* (1) property: by *Projection Theorem* (Theorem B.8 page 107).
- 2. Proof for *complemented* (2) property: by Corollary B.1 (page 105).
- 3. Proof for *involutory* property: by Corollary B.1 (page 105).
- 4. Proof for *antitone* property: by Proposition B.5 (page 102).
- 5. Proof for *orthomodular identity* property:
- 6. Proof that lattice is *orthomodular*: by 5 properties and definition of *orthomodular lattice*.

This concept is illustrated to the right where $X, Y \in 2^H$ are linear subspaces of the linear space H and

$$X \subseteq Y \implies Y = X + (X^{\perp} \cap Y).$$



Corollary B.2. Let H be a Hilbert space with orthogonality operation \bot . Let $(2^H, \uparrow, \alpha, 0, H; \subseteq)$ be the lattice of subspaces of H.

$$\begin{array}{c} \mathbf{C} \\ \mathbf{O} \\ \mathbf{R} \end{array} \begin{array}{c} (\boldsymbol{X} \, \hat{+} \, \boldsymbol{Y})^{\perp} & = \quad \boldsymbol{X}^{\perp} \, \boldsymbol{\Omega} \, \boldsymbol{Y}^{\perp} \quad \forall \boldsymbol{X}, \boldsymbol{Y} \in 2^{H} \quad (\text{De Morgan}) \quad and \\ (\boldsymbol{X} \, \boldsymbol{\Omega} \, \boldsymbol{Y})^{\perp} & = \quad \boldsymbol{X}^{\perp} \, \hat{+} \, \boldsymbol{Y}^{\perp} \quad \forall \boldsymbol{X}, \boldsymbol{Y} \in 2^{H} \quad (\text{De Morgan}) \end{array}$$

[♠]Proof: By properties of *orthocomplemented lattices* .

 \Rightarrow

¹⁹ Iturrioz (1985) pages 56–57



B.4. SUBSPACE METRICS Daniel J. Greenhoe page 109

B.4 Subspace Metrics

Definition B.4 (Hilbert space gap metric). ²⁰ Let X be a **Hilbert space** and S the set of subspaces of X. Then we define the following metric between subspaces of X.

```
 d(V, W) \triangleq \|P - Q\| \quad \forall V, W \in S  (the distance between subspaces V and W is the size of the difference of their projection operators)  where \ V \triangleq PX  (P is the projection operator that generates the subspace V)  and \ W \triangleq QX  (Q is the projection operator that generates the subspace W).
```

Definition B.5 (Banach space gap metric). Let X be a **Banach space** and S the set of subspaces of X. Then we define the following metric between subspaces of X.

```
d(V, W) \triangleq \max \left\{ \sup_{v \in V, \|v\| = 1} p(v, W), \sup_{w \in W, \|w\| = 1} p(w, V) \right\} \forall V, w \in S
where p(v, W) \triangleq \inf_{w \in W} \|v - w\| \qquad (metric from the point v to the subspace W)
```

Definition B.6 (Schäffer's metric). ²²

```
d(V, W) = \log(1 + \max\{r(V, W), r(W, V)\}) \quad where
r(V, W) \triangleq \begin{cases} \inf\{\|\mathbf{A} - \mathbf{I}\| | \mathbf{A}\mathbf{V} = \mathbf{W}\} & \text{if } \mathbf{A} \text{ and } \mathbf{A}^{-1} \text{ both exist} \\ 1 & \text{otherwise} \end{cases}
```

B.5 Literature

LITERATURE SURVEY:

1. Lattice of subspaces

Husimi (1937)

Sasaki (1954)

Loomis (1955)

von Neumann (1960)

Holland (1970)

Amemiya and Araki (1966)

■ Gudder (1979)

Gudder (2005)

2. Characterizations of lattice of Hilbert subspaces (cf @ Iturrioz (1985) page 60):

Piron (1964a) (using pre-Hilbert spaces)

Piron (1964b) (using pre-Hilbert spaces)

Amemiya and Araki (1966) (using pre-Hilbert spaces)

3. Metrics on subspaces:

Burago et al. (2001)



²⁰ ☐ Deza and Deza (2006) page 235, ☐ Akhiezer and Glazman (1993) page 69, ☐ Berkson (1963) page 8, ☐ Krein and Krasnoselski (1947)

²² Massera and Schäffer (1958) pages 562–563, Berkson (1963) pages 7–8





²¹ Akhiezer and Glazman (1993) page 70, Berkson (1963) page 8, Krein et al. (1948)

APPENDIX C	
I	
	OPERATORS ON LINEAR SPACES



■ And I am not afraid to say that there is a way to advance algebra as far beyond what Vieta and Descartes have left us as Vieta and Descartes carried it beyond the ancients.... we need still another analysis which is distinctly geometrical or linear, and which will express situation directly as algebra expresses magnitude directly. Gottfried Leibniz (1646–1716), German mathematician, in a September 8, 1679 letter to Christian Huygens.

C.1 Operators on linear spaces

C.1.1 Operator Algebra

An operator is simply a function that maps from a linear space to another linear space (or to the same linear space).

Definition C.1. ² Let $(\mathbb{F}, +, \cdot, 0, 1)$ be a FIELD (Definition A.5 page 96). Let X be a set, let + be an OPERATOR (Definition C.2 page 112) in X^{X^2} , and let \otimes be an operator in $X^{\mathbb{F} \times X}$.

² Kubrusly (2001) pages 40–41 ⟨Definition 2.1 and following remarks⟩, ■ Haaser and Sullivan (1991) page 41, ■ Halmos (1948) pages 1–2, ■ Peano (1888a) ⟨Chapter IX⟩, ■ Peano (1888b) pages 119–120, ■ Banach (1922) pages 134–135




Figure C.1: Some operator types

```
The structure \Omega \triangleq (X, +, \cdot, (\mathbb{F}, +, \times)) is a linear space over (\mathbb{F}, +, \cdot, 0, 1) if
         1. \exists 0 \in X such that x + 0 = x
                                                                                                                             (+ IDENTITY)
             \exists y \in X
                               such that x + y = 0
                                                                                                 \forall x \in X
                                                                                                                            (+ INVERSE)
                                        (x+y)+z = x+(y+z)
                                                                                                \forall x, y, z \in X
                                                                                                                            (+ is associative)
                                                 x + y = y + x
         4.
                                                                                                \forall x,y \in X
                                                                                                                            (+ is COMMUTATIVE)
                                                   1 \cdot x = x
                                                                                                \forall x \in X
                                                                                                                            (· IDENTITY)
                                           \alpha \cdot (\beta \cdot \mathbf{x}) = (\alpha \cdot \beta) \cdot \mathbf{x} \forall \alpha, \beta \in S \text{ and } \mathbf{x} \in X \text{ (Associates with )}
                                         \alpha \cdot (\mathbf{x} + \mathbf{y}) = (\alpha \cdot \mathbf{x}) + (\alpha \cdot \mathbf{y}) \quad \forall \alpha \in S \text{ and } \mathbf{x}, \mathbf{y} \in X
         7.
                                                                                                                            (· DISTRIBUTES over +)
                                         (\alpha + \beta) \cdot \mathbf{x} = (\alpha \cdot \mathbf{x}) + (\beta \cdot \mathbf{x}) \quad \forall \alpha, \beta \in S \text{ and } \mathbf{x} \in X
                                                                                                                            (· PSEUDO-DISTRIBUTES over +)
The set X is called the underlying set. The elements of X are called vectors. The elements of \mathbb{F}
are called scalars. A linear space is also called a vector space. If \mathbb{F} \triangleq \mathbb{R}, then \Omega is a real linear
space. If \mathbb{F} \triangleq \mathbb{C}, then \Omega is a complex linear space.
```

Definition C.2. ³


E

A function A in Y^X is an **operator** in Y^X if X and Y are both LINEAR SPACES (Definition C.1 page 111).

Two operators **A** and **B** in Y^X are **equal** if Ax = Bx for all $x \in X$. The inverse relation of an operator **A** in Y^X always exists as a *relation* in 2^{XY} , but may not always be a *function* (may not always be an operator) in Y^X .

The operator $\mathbf{I} \in \mathbf{X}^{\mathbf{X}}$ is the *identity* operator if $\mathbf{I}\mathbf{x} = \mathbf{I}$ for all $\mathbf{x} \in \mathbf{X}$.

Definition C.3. ⁴ Let X^X be the set of all operators with from a linear space X to X. Let I be an operator in X^X . Let $\mathbb{Q}(X)$ be the identity element in X^X .



³ Heil (2011) page 42

⁴ Michel and Herget (1993) page 411



C.1.2 Linear operators

Definition C.4. ⁵ Let $\mathbf{X} \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}))$ and $\mathbf{Y} \triangleq (Y, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}))$ be linear spaces.

D E F An operator $L \in Y^X$ is **linear** if

1.
$$L(x + y) = Lx + Ly \quad \forall x, y \in X$$
 (ADDITIVE) and

2.
$$\mathbf{L}(\alpha \mathbf{x}) = \alpha \mathbf{L} \mathbf{x}$$
 $\forall \mathbf{x} \in \mathbf{X}, \forall \alpha \in \mathbb{F}$ (homogeneous).

The set of all linear operators from X to Y is denoted $\mathcal{L}(X, Y)$ such that $\mathcal{L}(X, Y) \triangleq \{L \in Y^X | L \text{ is linear}\}$.

Theorem C.1. ⁶ *Let* \mathbf{L} *be an operator from a linear space* \mathbf{X} *to a linear space* \mathbf{Y} *, both over a field* \mathbb{F} .

$$\left\{ \mathbf{L} \text{ is LINEAR} \right\} \qquad \Longrightarrow \qquad \left\{ \begin{array}{lll}
1. & \mathbf{L} \mathbb{O} & = & \mathbb{O} & & \text{and} \\
2. & \mathbf{L} (-\mathbf{x}) & = & -(\mathbf{L}\mathbf{x}) & \forall \mathbf{x} \in \mathbf{X} & \text{and} \\
3. & \mathbf{L} (\mathbf{x} - \mathbf{y}) & = & \mathbf{L}\mathbf{x} - \mathbf{L}\mathbf{y} & \forall \mathbf{x}, \mathbf{y} \in \mathbf{X} & \text{and} \\
4. & \mathbf{L} \left(\sum_{n=1}^{N} \alpha_n \mathbf{x}_n \right) & = & \sum_{n=1}^{N} \alpha_n \left(\mathbf{L}\mathbf{x}_n \right) & \mathbf{x}_n \in \mathbf{X}, \alpha_n \in \mathbb{F} \end{array} \right\}$$

№PROOF:

1. Proof that L0 = 0:

2. Proof that L(-x) = -(Lx):

$$\mathbf{L}(-\mathbf{x}) = \mathbf{L}(-1 \cdot \mathbf{x})$$
 by *additive inverse* property $= -1 \cdot (\mathbf{L}\mathbf{x})$ by *homogeneous* property of \mathbf{L} (Definition C.4 page 113) $= -(\mathbf{L}\mathbf{x})$ by *additive inverse* property

3. Proof that L(x - y) = Lx - Ly:

$$\mathbf{L}(x-y) = \mathbf{L}(x+(-y))$$
 by additive inverse property $= \mathbf{L}(x) + \mathbf{L}(-y)$ by linearity property of \mathbf{L} (Definition C.4 page 113) $= \mathbf{L}x - \mathbf{L}y$ by item (2)

- 4. Proof that $\mathbf{L}\left(\sum_{n=1}^{N} \alpha_n \mathbf{x}_n\right) = \sum_{n=1}^{N} \alpha_n (\mathbf{L} \mathbf{x}_n)$:
 - (a) Proof for N = 1:

$$\mathbf{L}\left(\sum_{n=1}^{N} \alpha_{n} \mathbf{x}_{n}\right) = \mathbf{L}(\alpha_{1} \mathbf{x}_{1}) \qquad \text{by } N = 1 \text{ hypothesis}$$

$$= \alpha_{1}(\mathbf{L} \mathbf{x}_{1}) \qquad \text{by } homogeneous \text{ property of } \mathbf{L} \qquad \text{(Definition C.4 page 113)}$$

⁶ Berberian (1961) page 79 (Theorem IV.1.1)





 $^{^5}$ % Kubrusly (2001) page 55, Aliprantis and Burkinshaw (1998) page 224, Hilbert et al. (1927) page 6, Stone (1932) page 33

Daniel J. Greenhoe

(b) Proof that N case $\implies N+1$ case:

$$\begin{split} \mathbf{L} \left(\sum_{n=1}^{N+1} \alpha_n \mathbf{x}_n \right) &= \mathbf{L} \left(\alpha_{N+1} \mathbf{x}_{N+1} + \sum_{n=1}^{N} \alpha_n \mathbf{x}_n \right) \\ &= \mathbf{L} \left(\alpha_{N+1} \mathbf{x}_{N+1} \right) + \mathbf{L} \left(\sum_{n=1}^{N} \alpha_n \mathbf{x}_n \right) \quad \text{by } \textit{linearity } \textit{property of } \mathbf{L} \quad \textit{(Definition C.4 page 113)} \\ &= \alpha_{N+1} \mathbf{L} \left(\mathbf{x}_{N+1} \right) + \sum_{n=1}^{N} \mathbf{L} \left(\alpha_n \mathbf{x}_n \right) \quad \quad \textit{by } \textit{left } N+1 \textit{ hypothesis} \\ &= \sum_{n=1}^{N+1} \mathbf{L} \left(\alpha_n \mathbf{x}_n \right) \end{split}$$

Theorem C.2. ⁷ Let $\mathcal{L}(X, Y)$ be the set of all linear operators from a linear space X to a linear space Y. Let $\mathcal{N}(L)$ be the NULL SPACE of an operator L in Y^X and $\mathcal{L}(L)$ the IMAGE SET of L in Y^X .

т	$\mathscr{L}(\pmb{X},\pmb{Y})$	is a linear space		(space of linear transforms)
Ĥ	$\mathcal{N}(\mathbf{L})$	is a linear subspace of X	$\forall \mathbf{L} \in \mathbf{Y}^{\mathbf{X}}$	
M	$\mathcal{F}(\mathbf{L})$	is a linear subspace of Y	$\forall \mathbf{L} \in \mathbf{Y}^{\mathbf{X}}$	

№ Proof:

- 1. Proof that $\mathcal{N}(\mathbf{L})$ is a linear subspace of X:
 - (a) $0 \in \mathcal{N}(L) \implies \mathcal{N}(L) \neq \emptyset$
 - (b) $\mathcal{N}(\mathbf{L}) \triangleq \{ x \in X | \mathbf{L}x = 0 \} \subseteq X$
 - (c) $x + y \in \mathcal{N}(L) \implies 0 = L(x + y) = L(y + x) \implies y + x \in \mathcal{N}(L)$
 - (d) $\alpha \in \mathbb{F}$, $x \in X \implies 0 = Lx \implies 0 = \alpha Lx \implies 0 = L(\alpha x) \implies \alpha x \in \mathcal{N}(L)$
- 2. Proof that $\mathcal{J}(\mathbf{L})$ is a linear subspace of \mathbf{Y} :
 - (a) $0 \in \mathcal{I}(L) \implies \mathcal{I}(L) \neq \emptyset$
 - (b) $\mathcal{J}(L) \triangleq \{y \in Y | \exists x \in X \text{ such that } y = Lx\} \subseteq Y$
 - (c) $x + y \in \mathcal{J}(L) \implies \exists v \in X$ such that $Lv = x + y = y + x \implies y + x \in \mathcal{J}(L)$
 - (d) $\alpha \in \mathbb{F}$, $x \in \mathcal{J}(L) \implies \exists x \in X$ such that $y = Lx \implies \alpha y = \alpha Lx = L(\alpha x) \implies \alpha x \in \mathcal{J}(L)$

Example C.1. ⁸ Let $\mathcal{C}([a:b], \mathbb{R})$ be the set of all *continuous* functions from the closed real interval [a:b] to \mathbb{R} .

 $\mathscr{C}([a:b],\mathbb{R})$ is a linear space.

Theorem C.3. ⁹ Let $\mathcal{L}(X, Y)$ be the set of linear operators from a linear space X to a linear space Y. Let $\mathcal{N}(L)$ be the NULL SPACE of a linear operator $L \in \mathcal{L}(X, Y)$.

Ţ	Lx = Ly	\iff	$x - y \in \mathcal{N}(L)$
M	$\mathbf{L}\mathbf{x} = \mathbf{L}\mathbf{y}$ \mathbf{L} is injective	\iff	$\mathcal{N}(\mathbf{L}) = \{0\}$

⁷ ■ Michel and Herget (1993) pages 98–104, ■ Berberian (1961) pages 80–85 (Theorem IV.1.4 and Theorem IV.3.1)

 $^{^9}$ Berberian (1961) page 88 (Theorem IV.1.4)



⁸ Eidelman et al. (2004) page 3

 \blacksquare

^ℚProof:

1. Proof that $Lx = Ly \implies x - y \in \mathcal{N}(L)$:

$$L(x - y) = Lx - Ly$$
 by Theorem C.1 page 113
 by left hypothesis

$$\implies x - y \in \mathcal{N}(L)$$
 by definition of *Null Space*

2. Proof that $Lx = Ly \iff x - y \in \mathcal{N}(L)$:

$$\mathbf{L}y = \mathbf{L}y + \mathbf{0}$$
 by definition of linear space (Definition C.1 page 111)

 $= \mathbf{L}y + \mathbf{L}(x - y)$ by right hypothesis

 $= \mathbf{L}y + (\mathbf{L}x - \mathbf{L}y)$ by Theorem C.1 page 113

 $= (\mathbf{L}y - \mathbf{L}y) + \mathbf{L}x$ by associative and commutative properties (Definition C.1 page 111)

 $= \mathbf{L}x$

3. Proof that **L** is *injective* $\iff \mathcal{N}(\mathbf{L}) = \{0\}$:

L is injective
$$\iff \{(\mathbf{L}\mathbf{x} = \mathbf{L}\mathbf{y} \iff \mathbf{x} = \mathbf{y}) \ \forall \mathbf{x}, \mathbf{y} \in X\}$$

$$\iff \{[\mathbf{L}\mathbf{x} - \mathbf{L}\mathbf{y} = 0 \iff (\mathbf{x} - \mathbf{y}) = 0] \ \forall \mathbf{x}, \mathbf{y} \in X\}$$

$$\iff \{[\mathbf{L}(\mathbf{x} - \mathbf{y}) = 0 \iff (\mathbf{x} - \mathbf{y}) = 0] \ \forall \mathbf{x}, \mathbf{y} \in X\}$$

$$\iff \mathcal{N}(\mathbf{L}) = \{0\}$$

Theorem C.4. 10 Let W, X, Y, and Z be linear spaces over a field \mathbb{F} .

		, , ,	, 3	
	1. L(MN) =	(LM)N	$\forall L {\in} \mathscr{L}(\boldsymbol{Z}, \boldsymbol{W}), M {\in} \mathscr{L}(\boldsymbol{Y}, \boldsymbol{Z}), N {\in} \mathscr{L}(\boldsymbol{X}, \boldsymbol{Y})$	(ASSOCIATIVE)
I		$(LM) \stackrel{\circ}{+} (LN)$	$\forall L \in \mathcal{L}(Y, Z), M \in \mathcal{L}(X, Y), N \in \mathcal{L}(X, Y)$	(LEFT DISTRIBUTIVE)
М	3. $(\mathbf{L} + \mathbf{M}) \hat{\mathbf{N}} =$	$(LN) \stackrel{\circ}{+} (MN)$	$\forall L {\in} \mathscr{L}(\textbf{\textit{Y}},\textbf{\textit{Z}}), M {\in} \mathscr{L}(\textbf{\textit{Y}},\textbf{\textit{Z}}), N {\in} \mathscr{L}(\textbf{\textit{X}},\textbf{\textit{Y}})$	(RIGHT DISTRIBUTIVE)
	4. $\alpha(LM)$ =	$(\alpha \mathbf{L})\mathbf{M} = \mathbf{L}(\alpha \mathbf{M})$	$\forall \mathbf{L} {\in} \mathcal{L}(\mathbf{\textit{Y}},\mathbf{\textit{Z}}), \mathbf{M} {\in} \mathcal{L}(\mathbf{\textit{X}},\mathbf{\textit{Y}}), \alpha {\in} \mathbb{F}$	(HOMOGENEOUS)

№PROOF:

- 1. Proof that L(MN) = (LM)N: Follows directly from property of associative operators.
- 2. Proof that L(M + N) = (LM) + (LN):

$$\begin{aligned} \left[\mathbf{L} \big(\mathbf{M} + \mathbf{N} \big) \right] \mathbf{x} &= \mathbf{L} \left[\big(\mathbf{M} + \mathbf{N} \big) \mathbf{x} \right] \\ &= \mathbf{L} \left[(\mathbf{M} \mathbf{x}) + (\mathbf{N} \mathbf{x}) \right] \\ &= \left[\mathbf{L} (\mathbf{M} \mathbf{x}) \right] + \left[\mathbf{L} (\mathbf{N} \mathbf{x}) \right] \end{aligned} \quad \text{by additive property Definition C.4 page 113} \\ &= \left[(\mathbf{L} \mathbf{M}) \mathbf{x} \right] + \left[(\mathbf{L} \mathbf{N}) \mathbf{x} \right] \end{aligned}$$

- 3. Proof that (L + M)N = (LN) + (MN): Follows directly from property of *associative* operators.
- 4. Proof that $\alpha(LM) = (\alpha L)M$: Follows directly from *associative* property of linear operators.
- 5. Proof that $\alpha(LM) = L(\alpha M)$:

$$\begin{split} & [\alpha(\mathbf{L}\mathbf{M})] \boldsymbol{x} = \alpha[(\mathbf{L}\mathbf{M})\boldsymbol{x}] \\ & = \mathbf{L}[\alpha(\mathbf{M}\boldsymbol{x})] \qquad \qquad \text{by $homogeneous$ property Definition C.4 page 113} \\ & = \mathbf{L}[(\alpha\mathbf{M})\boldsymbol{x}] \\ & = [\mathbf{L}(\alpha\mathbf{M})]\boldsymbol{x} \end{split}$$



¹⁰ Berberian (1961) page 88 (Theorem IV.5.1)

Theorem C.5 (Fundamental theorem of linear equations). ¹¹ Let Y^X be the set of all operators from a linear space X to a linear space Y. Let $\mathcal{N}(L)$ be the NULL SPACE of an operator L in Y^X and $\mathcal{F}(L)$ the IMAGE SET of L in Y^X .

$$\dim \mathcal{J}(\mathbf{L}) + \dim \mathcal{N}(\mathbf{L}) = \dim \mathbf{X} \qquad \forall \mathbf{L} \in \mathbf{Y}^{\mathbf{X}}$$

PROOF: Let $\{\psi_k | k = 1, 2, ..., p\}$ be a basis for \boldsymbol{X} constructed such that $\{\psi_{p-n+1}, \psi_{p-n+2}, ..., \psi_p\}$ is a basis for $\boldsymbol{\mathcal{N}}(\mathbf{L})$.

Let
$$p \triangleq \dim X$$
.
Let $n \triangleq \dim \mathcal{N}(\mathbf{L})$.

$$\begin{split} \dim \mathscr{F}(\mathbf{L}) &= \dim \left\{ \mathbf{y} \in \mathbf{Y} | \exists \mathbf{x} \in \mathbf{X} \quad \text{such that} \quad \mathbf{y} = \mathbf{L} \mathbf{x} \right\} \\ &= \dim \left\{ \begin{array}{l} \mathbf{y} \in \mathbf{Y} | \exists (\alpha_1, \alpha_2, \dots, \alpha_p) \quad \text{such that} \quad \mathbf{y} = \mathbf{L} \sum_{k=1}^p \alpha_k \mathbf{L} \psi_k \right\} \\ &= \dim \left\{ \begin{array}{l} \mathbf{y} \in \mathbf{Y} | \exists (\alpha_1, \alpha_2, \dots, \alpha_p) \quad \text{such that} \quad \mathbf{y} = \sum_{k=1}^p \alpha_k \mathbf{L} \psi_k \right\} \\ &= \dim \left\{ \begin{array}{l} \mathbf{y} \in \mathbf{Y} | \exists (\alpha_1, \alpha_2, \dots, \alpha_p) \quad \text{such that} \quad \mathbf{y} = \sum_{k=1}^{p-n} \alpha_k \mathbf{L} \psi_k + \sum_{k=1}^n \alpha_k \mathbf{L} \psi_k \right\} \\ &= \dim \left\{ \begin{array}{l} \mathbf{y} \in \mathbf{Y} | \exists (\alpha_1, \alpha_2, \dots, \alpha_p) \quad \text{such that} \quad \mathbf{y} = \sum_{k=1}^{p-n} \alpha_k \mathbf{L} \psi_k + \mathbf{0} \right\} \\ &= p-n \\ &= \dim \mathbf{X} - \dim \mathscr{N}(\mathbf{L}) \end{split}$$

Note: This "proof" may be missing some necessary detail.

C.2 Operators on Normed linear spaces

C.2.1 Operator norm

Definition C.5. ¹² Let $V = (X, \mathbb{F}, \hat{+}, \cdot)$ be a linear space and \mathbb{F} be a field with absolute value function $|\cdot| \in \mathbb{R}^{\mathbb{F}}$ (Definition A.4 page 96).

A norm is any functional $\|\cdot\|$ in \mathbb{R}^X that satisfies

1. $\|x\| \ge 0$ $\forall x \in X$ (Strictly positive) and

2. $\|x\| = 0 \iff x = \emptyset$ $\forall x \in X$ (Nondegenerate) and

3. $\|ax\| = |a| \|x\|$ $\forall x \in X, a \in \mathbb{C}$ (Homogeneous) and

4. $\|x + y\| \le \|x\| + \|y\|$ $\forall x, y \in X$ (Subadditive/triangle inquality).

A normed linear space is the pair $(V, \|\cdot\|)$.

¹² ■ Aliprantis and Burkinshaw (1998) pages 217–218, ■ Banach (1932a) page 53, ■ Banach (1932b) page 33, ■ Banach (1922) page 135



¹¹ Michel and Herget (1993) page 99

Definition C.6. ¹³ Let $\mathcal{L}(X, Y)$ be the space of linear operators over normed linear spaces X and Y.

D E F

```
The operator norm \|\cdot\| is defined as \|\|\mathbf{A}\|\| \triangleq \sup_{\mathbf{x} \in \mathbf{X}} \{\|\mathbf{A}\mathbf{x}\| \mid \|\mathbf{x}\| \le 1\} \qquad \forall \mathbf{A} \in \mathcal{L}(\mathbf{X}, \mathbf{Y})
The pair (\mathcal{L}(\mathbf{X}, \mathbf{Y}), \|\|\cdot\|) is the normed space of linear operators on (\mathbf{X}, \mathbf{Y}).
```

Proposition C.1 (next) shows that the functional defined in Definition C.6 (previous) is a *norm* (Definition C.5 page 116).

Proposition C.1. ¹⁵ Let $(\mathcal{L}(X, Y), |||\cdot|||)$ be the normed space of linear operators over the normed linear spaces $X \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), ||\cdot||)$ and $Y \triangleq (Y, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), ||\cdot||)$.

	,	,	*	,		
	The functional $\ \cdot\ $ is a $oldsymbol{I}$	norm on $\mathcal{L}(X,$	Y). In particular,			
	$1. \mathbf{A} \geq 0$		$\forall \mathbf{A} \in \mathcal{L}(\mathbf{X}, \mathbf{Y})$	(NON-NEGATIVE)	and	
P R	$2. \mathbf{A} = 0$	$\iff \mathbf{A} \stackrel{\circ}{=} \mathbb{0}$	$\forall \mathbf{A} \in \mathcal{L}(\mathbf{X}, \mathbf{Y})$	(NONDEGENERATE)	and	
P	3. $\ \alpha \mathbf{A} \ = \alpha \ $		$\forall \mathbf{A} {\in} \mathcal{L}(\mathbf{X}, \mathbf{Y}), \alpha {\in} \mathbb{F}$	(HOMOGENEOUS)	and	
	$4. \mathbf{A} + \mathbf{B} \leq \mathbf{A} $	$\ \mathbf{H}\ $	$\forall \mathbf{A} \in \mathcal{L}(\mathbf{X}, \mathbf{Y})$	(SUBADDITIVE).		
	Moreover, $(\mathscr{L}(X, Y), \ \cdot\)$ is a normed linear space .					

[♥]Proof:

1. Proof that $\|\mathbf{A}\| > 0$ for $\mathbf{A} \neq 0$:

$$\|\mathbf{A}\| \triangleq \sup_{\mathbf{x} \in \mathbf{X}} \{ \|\mathbf{A}\mathbf{x}\| \mid \|\mathbf{x}\| \le 1 \}$$
$$> 0$$

by definition of |||·||| (Definition C.6 page 117)

2. Proof that $\|\mathbf{A}\| = 0$ for $\mathbf{A} \stackrel{\circ}{=} 0$:

$$|||\mathbf{A}||| \triangleq \sup_{x \in X} \{||\mathbf{A}x|| \mid ||x|| \le 1\}$$
$$= \sup_{x \in X} \{||0x|| \mid ||x|| \le 1\}$$
$$= 0$$

by definition of |||·||| (Definition C.6 page 117)

3. Proof that $\|\alpha A\| = |\alpha| \|A\|$:

$$\begin{aligned} \| \alpha \mathbf{A} \| & \triangleq \sup_{\boldsymbol{x} \in \boldsymbol{X}} \left\{ \| \alpha \mathbf{A} \boldsymbol{x} \| \mid \| \boldsymbol{x} \| \leq 1 \right\} \\ &= \sup_{\boldsymbol{x} \in \boldsymbol{X}} \left\{ |\alpha| \| \mathbf{A} \boldsymbol{x} \| \mid \| \boldsymbol{x} \| \leq 1 \right\} \\ &= |\alpha| \sup_{\boldsymbol{x} \in \boldsymbol{X}} \left\{ \| \mathbf{A} \boldsymbol{x} \| \mid \| \boldsymbol{x} \| \leq 1 \right\} \end{aligned} \qquad \text{by definition of } \| \cdot \| \text{ (Definition C.6 page 117)}$$

$$= |\alpha| \sup_{\boldsymbol{x} \in \boldsymbol{X}} \left\{ \| \mathbf{A} \boldsymbol{x} \| \mid \| \boldsymbol{x} \| \leq 1 \right\}$$

$$= |\alpha| \| \| \mathbf{A} \|$$

$$= |\alpha| \| \| \mathbf{A} \|$$

$$\text{by definition of } \| \cdot \| \text{ (Definition C.6 page 117)}$$





¹³ ■ Rudin (1991) page 92, ■ Aliprantis and Burkinshaw (1998) page 225

¹⁴The operator norm notation |||·||| is introduced (as a Matrix norm) in


Horn and Johnson (1990) page 290

¹⁵ Rudin (1991) page 93

$$\| \mathbf{A} + \mathbf{B} \| \triangleq \sup_{x \in X} \left\{ \| (\mathbf{A} + \mathbf{B})x \| \mid \|x\| \le 1 \right\}$$
 by definition of $\| \cdot \|$ (Definition C.6 page 117)
$$= \sup_{x \in X} \left\{ \| \mathbf{A}x + \mathbf{B}x \| \mid \|x\| \le 1 \right\}$$
 by definition of $\| \cdot \|$ (Definition C.6 page 117)
$$\le \sup_{x \in X} \left\{ \| \mathbf{A}x \| \mid \|x\| \le 1 \right\} + \sup_{x \in X} \left\{ \| \mathbf{B}x \| \mid \|x\| \le 1 \right\}$$
 by definition of $\| \cdot \|$ (Definition C.6 page 117)
$$\triangleq \| \| \mathbf{A} \| + \| \| \mathbf{B} \|$$
 by definition of $\| \cdot \|$ (Definition C.6 page 117)

₽

Lemma C.1. Let $(\mathcal{L}(X, Y), |||\cdot|||)$ be the normed space of linear operators over normed linear spaces $X \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), ||\cdot||)$ and $Y \triangleq (Y, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), ||\cdot||)$.



Daniel J. Greenhoe

[♠]Proof: ¹⁶

1. Proof that $\sup_{x} \{ \|\mathbf{L}x\| \mid \|x\| \le 1 \} \ge \sup_{x} \{ \|\mathbf{L}x\| \mid \|x\| = 1 \}$:

$$\sup_{\mathbf{x}} \{ \|\mathbf{L}\mathbf{x}\| \mid \|\mathbf{x}\| \le 1 \} \ge \sup_{\mathbf{x}} \{ \|\mathbf{L}\mathbf{x}\| \mid \|\mathbf{x}\| = 1 \} \qquad \text{because } A \subseteq B \implies \sup_{\mathbf{x}} A \le \sup_{\mathbf{x}} B$$

2. Let the subset $Y \subseteq X$ be defined as

$$Y \triangleq \left\{ \begin{array}{ll} 1. & \|\mathbf{L}\mathbf{y}\| = \sup_{\mathbf{x} \in X} \{\|\mathbf{L}\mathbf{x}\| \mid \|\mathbf{x}\| \le 1\} \text{ and } \\ 2. & 0 < \|\mathbf{y}\| \le 1 \end{array} \right\}$$

3. Proof that $\sup_{\mathbf{x}} \{ \|\mathbf{L}\mathbf{x}\| \mid \|\mathbf{x}\| \le 1 \} \le \sup_{\mathbf{x}} \{ \|\mathbf{L}\mathbf{x}\| \mid \|\mathbf{x}\| = 1 \}$:

$$\sup_{x \in X} \{ \|\mathbf{L}x\| \mid \|x\| \le 1 \} = \|\mathbf{L}y\|$$
 by definition of set Y

$$= \frac{\|y\|}{\|y\|} \|\mathbf{L}y\|$$
 by homogeneous property (page 116)
$$= \|y\| \left\| \mathbf{L} \frac{y}{\|y\|} \right\|$$
 by homogeneous property (page 113)
$$\le \|y\| \sup_{y \in Y} \left\{ \left\| \mathbf{L} \frac{y}{\|y\|} \right\| \right\}$$
 by definition of supremum
$$= \|y\| \sup_{y \in Y} \left\{ \left\| \mathbf{L} \frac{y}{\|y\|} \right\| \mid \left\| \frac{y}{\|y\|} \right\| = 1 \right\}$$
 because $\left\| \frac{y}{\|y\|} \right\| = 1$ for all $y \in Y$

$$\le \sup_{y \in Y} \left\{ \left\| \mathbf{L} \frac{y}{\|y\|} \right\| \mid \left\| \frac{y}{\|y\|} \right\| = 1 \right\}$$
 because $0 < \|y\| \le 1$

$$\le \sup_{x \in X} \left\{ \|\mathbf{L}x\| \mid \|x\| = 1 \right\}$$
 because $\frac{y}{\|y\|} \in X$ $\forall y \in Y$



Many many thanks to former NCTU Ph.D. student Chien Yao (Chinese: 姚建; PinYin: Yáo Jiàn) for his brilliant help with this proof. (If you are viewing this text as a pdf file, zoom in on the figure to the left to see text from Chien Yao's 2007 April 16 email.)



4. By (1) and (3),

$$\sup_{x \in X} \{ \|\mathbf{L}x\| \mid \|x\| \le 1 \} = \sup_{x \in X} \{ \|\mathbf{L}x\| \mid \|x\| = 1 \}$$

 \blacksquare

Proposition C.2. ¹⁷ Let **I** be the identity operator in the normed space of linear operators $(\mathcal{L}(X, X), \|\|\cdot\|\|)$.



♥Proof:

$$\|\|\mathbf{I}\|\| \triangleq \sup \{ \|\mathbf{I}x\| \mid \|x\| \le 1 \}$$

= $\sup \{ \|x\| \mid \|x\| \le 1 \}$
= 1

by definition of |||·||| (Definition C.6 page 117) by definition of I (Definition C.3 page 112)

 \blacksquare

Theorem C.6. ¹⁸ Let $(\mathcal{L}(X, Y), \|\|\cdot\|\|)$ be the normed space of linear operators over normed linear spaces X and Y.



^ℚProof:

1. Proof that $||Lx|| \le |||L||| ||x||$:

$$\|\mathbf{L}\mathbf{x}\| = \frac{\|\mathbf{x}\|}{\|\mathbf{x}\|} \|\mathbf{L}\mathbf{x}\|$$

$$= \|\mathbf{x}\| \left\| \frac{1}{\|\mathbf{x}\|} \mathbf{L}\mathbf{x} \right\|$$

$$= \|\mathbf{x}\| \left\| \mathbf{L} \frac{\mathbf{x}}{\|\mathbf{x}\|} \right\|$$

$$\triangleq \|\mathbf{x}\| \|\mathbf{L}\mathbf{y}\|$$

$$\leq \|\mathbf{x}\| \sup_{\mathbf{y}} \|\mathbf{L}\mathbf{y}\|$$

$$= \|\mathbf{x}\| \sup_{\mathbf{y}} \{ \|\mathbf{L}\mathbf{y}\| \mid \|\mathbf{y}\| = 1 \}$$

$$\triangleq \|\mathbf{x}\| \|\mathbf{L}\|$$

by property of norms

by property of linear operators

where
$$y \triangleq \frac{x}{\|x\|}$$

by definition of supremum

because
$$||y|| = \left\| \frac{x}{||x||} \right\| = \frac{||x||}{||x||} = 1$$

by definition of operator norm



¹⁷ Michel and Herget (1993) page 410

¹⁸ Rudin (1991) page 103, Aliprantis and Burkinshaw (1998) page 225

2. Proof that $|||KL||| \le |||K||| |||L|||$:

Daniel J. Greenhoe

C.2.2 Bounded linear operators

Definition C.7. 19 Let $(\mathcal{L}(X, Y), \|\|\cdot\|)$ be a normed space of linear operators.

Ε

An operator **B** is **bounded** if $|||\mathbf{B}||| < \infty$.

The quantity $\mathcal{B}(X, Y)$ is the set of all **bounded linear operators** on (X, Y) such that $\mathscr{B}(X, Y) \triangleq \{ \mathbf{L} \in \mathscr{L}(X, Y) | |||\mathbf{L}||| < \infty \}.$

Theorem C.7. 20 Let $(\mathcal{L}(X, Y), \|\|\cdot\|\|)$ be the set of linear operators over normed linear spaces $\mathbf{X} \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \|\cdot\|) \text{ and } \mathbf{Y} \triangleq (Y, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \|\cdot\|).$

Т

The following conditions are all EQUIVALENT:

- 1. L is continuous at a single point $x_0 \in X \quad \forall L \in \mathcal{L}(X,Y)$ 2. L is continuous (at every point $x \in X$) $\forall \mathbf{L} \in \mathcal{L}(\mathbf{X}, \mathbf{Y})$
- 3. $|||\mathbf{L}||| < \infty$ (L is bounded) $\forall L \in \mathcal{L}(X,Y)$
- 4. $\exists M \in \mathbb{R}$ such that $\|\mathbf{L}\mathbf{x}\| \leq M \|\mathbf{x}\|$ $\forall \mathbf{L} \in \mathcal{L}(\mathbf{X}, \mathbf{Y}), \mathbf{x} \in X$

^ℚProof:

1. Proof that $1 \implies 2$:

$$\begin{aligned} \epsilon &> \left\| \mathbf{L} \mathbf{x} - \mathbf{L} \mathbf{x}_0 \right\| & \text{by hypothesis 1} \\ &= \left\| \mathbf{L} (\mathbf{x} - \mathbf{x}_0) \right\| & \text{by linearity (Definition C.4 page 113)} \\ &= \left\| \mathbf{L} (\mathbf{x} + \mathbf{y} - \mathbf{x}_0 - \mathbf{y}) \right\| & \text{by linearity (Definition C.4 page 113)} \\ &\Rightarrow \mathbf{L} \text{ is continuous at point } \mathbf{x} + \mathbf{y} \\ &\Rightarrow \mathbf{L} \text{ is continuous at every point in } X & \text{(hypothesis 2)} \end{aligned}$$

2. Proof that $2 \implies 1$: obvious:

²⁰ Aliprantis and Burkinshaw (1998) page 227



¹⁹ Rudin (1991) pages 92–93

3. Proof that $4 \implies 2^{21}$

$$\begin{aligned} \|\|\mathbf{L}x\|\| &\leq M \, \|x\| \implies \|\|\mathbf{L}(x-y)\|\| \leq M \, \|x-y\| & \text{by hypothesis 4} \\ &\implies \|\|\mathbf{L}x-\mathbf{L}y\|\| \leq M \, \|x-y\| & \text{by linearity of } \mathbf{L} \text{ (Definition C.4 page 113)} \\ &\implies \|\|\mathbf{L}x-\mathbf{L}y\|\| \leq \epsilon \text{ whenever } M \, \|x-y\| < \epsilon \\ &\implies \|\|\mathbf{L}x-\mathbf{L}y\|\| \leq \epsilon \text{ whenever } \|x-y\| < \frac{\epsilon}{M} & \text{(hypothesis 2)} \end{aligned}$$

4. Proof that $3 \implies 4$:

$$\|\mathbf{L}x\| \le \|\mathbf{L}\| \|x\|$$
 by Theorem C.6 page 119
$$= M \|x\|$$
 where $M \triangleq \|\mathbf{L}\| < \infty$ (by hypothesis 1)

5. Proof that $1 \implies 3^{22}$

$$\|\|\mathbf{L}\|\| = \infty \implies \{\|\mathbf{L}\mathbf{x}\| \mid \|\mathbf{x}\| \le 1\} = \infty$$

$$\implies \exists (\mathbf{x}_n) \quad \text{such that} \quad \|\mathbf{x}_n\| = 1 \text{ and } \|\|\mathbf{L}\|\| = \{\|\mathbf{L}\mathbf{x}_n\| \mid \|\mathbf{x}_n\| \le 1\} = \infty$$

$$\implies \|\mathbf{x}_n\| = 1 \text{ and } \infty = \|\|\mathbf{L}\|\| = \|\mathbf{L}\mathbf{x}_n\|$$

$$\implies \|\mathbf{x}_n\| = 1 \text{ and } \|\mathbf{L}\mathbf{x}_n\| \ge n$$

$$\implies \frac{1}{n} \|\mathbf{x}_n\| = \frac{1}{n} \text{ and } \frac{1}{n} \|\mathbf{L}\mathbf{x}_n\| \ge 1$$

$$\implies \|\frac{\mathbf{x}_n}{n}\| = \frac{1}{n} \text{ and } \|\mathbf{L}\frac{\mathbf{x}_n}{n}\| \ge 1$$

$$\implies \lim_{n \to \infty} \|\frac{\mathbf{x}_n}{n}\| = 0 \text{ and } \lim_{n \to \infty} \|\mathbf{L}\frac{\mathbf{x}_n}{n}\| \ge 1$$

$$\implies \mathbf{L} \text{ is not continuous at } 0$$

But by hypothesis, L is continuous. So the statement $\|\|L\|\| = \infty$ must be false and thus $\|\|L\|\| < \infty$ (L is bounded).

Adjoints on normed linear spaces **C.2.3**

Definition C.8. Let $\mathcal{B}(X, Y)$ be the space of BOUNDED LINEAR OPERATORS on normed linear spaces **X** and **Y**. Let X^* be the TOPOLOGICAL DUAL SPACE of **X**.

 \mathbf{B}^* is the **adjoint** of an operator $\mathbf{B} \in \mathcal{B}(\mathbf{X}, \mathbf{Y})$ if $f(\mathbf{B}\mathbf{x}) = [\mathbf{B}^*\mathbf{f}](\mathbf{x})$ $\forall f \in X^*, x \in X$

Theorem C.8. 23 Let $\mathcal{B}(X,Y)$ be the space of bounded linear operators on normed linear SPACES X and Y

Т	$(\mathbf{A} \stackrel{\circ}{+} \mathbf{B})^*$	=	$\mathbf{A}^* \stackrel{\circ}{+} \mathbf{B}^*$	$\forall \mathbf{A}, \mathbf{B} \in \mathcal{B}(\mathbf{X}, \mathbf{Y})$
н	$(\lambda \mathbf{A})^*$	=	$\lambda \mathbf{A}^*$	$\forall A,B \in \mathcal{B}(X,Y)$
M	$(\mathbf{AB})^*$	=	$\mathbf{B}^*\mathbf{A}^*$	$\forall A,B \in \mathcal{B}(X,Y)$

²¹ Bollobás (1999) page 29





²² Aliprantis and Burkinshaw (1998) page 227

²³ Bollobás (1999) page 156

№ Proof:

$$\left[\mathbf{A} \stackrel{\circ}{+} \mathbf{B} \right]^* \mathbf{f}(\mathbf{x}) = \mathbf{f} \left(\left[\mathbf{A} \stackrel{\circ}{+} \mathbf{B} \right] \mathbf{x} \right) \qquad \text{by definition of adjoint} \qquad \text{(Definition C.8 page 121)}$$

$$= \mathbf{f}(\mathbf{A}\mathbf{x} + \mathbf{B}\mathbf{x}) \qquad \text{by definition of linear operators} \qquad \text{(Definition C.4 page 113)}$$

$$= \mathbf{f}(\mathbf{A}\mathbf{x}) + \mathbf{f}(\mathbf{B}\mathbf{x}) \qquad \text{by definition of } \mathbf{adjoint} \qquad \text{(Definition C.8 page 121)}$$

$$= \mathbf{A}^* \mathbf{f}(\mathbf{x}) + \mathbf{B}^* \mathbf{f}(\mathbf{x}) \qquad \text{by definition of } \mathbf{adjoint} \qquad \text{(Definition C.8 page 121)}$$

$$= \left[\mathbf{A}^* + \mathbf{B}^* \right] \mathbf{f}(\mathbf{x}) \qquad \text{by definition of } \mathbf{adjoint} \qquad \text{(Definition C.8 page 121)}$$

$$= \lambda \mathbf{f}(\mathbf{A}\mathbf{x}) \qquad \text{by definition of } \mathbf{adjoint} \qquad \text{(Definition C.8 page 121)}$$

$$= \left[\lambda \mathbf{A}^* \right] \mathbf{f}(\mathbf{x}) \qquad \text{by definition of } \mathbf{adjoint} \qquad \text{(Definition C.8 page 121)}$$

$$= \mathbf{f}(\mathbf{A}\mathbf{B}\mathbf{x}) \qquad \text{by definition of } \mathbf{adjoint} \qquad \text{(Definition C.8 page 121)}$$

$$= \mathbf{f}(\mathbf{A}\mathbf{B}\mathbf{x}) \qquad \text{by definition of } \mathbf{adjoint} \qquad \text{(Definition C.8 page 121)}$$

$$= \mathbf{B}^* \mathbf{f}(\mathbf{A}^* \mathbf{f}) \mathbf{f}(\mathbf{x}) \qquad \text{by definition of } \mathbf{adjoint} \qquad \text{(Definition C.8 page 121)}$$

$$= \mathbf{B}^* \mathbf{f}(\mathbf{A}^* \mathbf{f}) \mathbf{f}(\mathbf{x}) \qquad \text{by definition of } \mathbf{adjoint} \qquad \text{(Definition C.8 page 121)}$$

$$= \mathbf{B}^* \mathbf{f}(\mathbf{A}^* \mathbf{f}) \mathbf{f}(\mathbf{x}) \qquad \text{by definition of } \mathbf{adjoint} \qquad \text{(Definition C.8 page 121)}$$

$$= \mathbf{B}^* \mathbf{f}(\mathbf{A}^* \mathbf{f}) \mathbf{f}(\mathbf{x}) \qquad \text{by definition of } \mathbf{adjoint} \qquad \text{(Definition C.8 page 121)}$$

$$= \mathbf{B}^* \mathbf{f}(\mathbf{A}^* \mathbf{f}) \mathbf{f}(\mathbf{x}) \qquad \text{by definition of } \mathbf{adjoint} \qquad \text{(Definition C.8 page 121)}$$

$$= \mathbf{B}^* \mathbf{f}(\mathbf{A}^* \mathbf{f}) \mathbf{f}(\mathbf{x}) \qquad \text{by definition of } \mathbf{adjoint} \qquad \text{(Definition C.8 page 121)}$$

$$= \mathbf{B}^* \mathbf{f}(\mathbf{A}^* \mathbf{f}) \mathbf{f}(\mathbf{x}) \qquad \text{by definition of } \mathbf{adjoint} \qquad \text{(Definition C.8 page 121)}$$

$$= \mathbf{B}^* \mathbf{f}(\mathbf{A}^* \mathbf{f}) \mathbf{f}(\mathbf{A}) \qquad \text{(Definition C.8 page 121)}$$

$$= \mathbf{B}^* \mathbf{f}(\mathbf{A}^* \mathbf{f}) \mathbf{f}(\mathbf{A}) \qquad \text{(Definition C.8 page 121)}$$

$$= \mathbf{B}^* \mathbf{f}(\mathbf{A}^* \mathbf{f}) \mathbf{f}(\mathbf{A}) \qquad \text{(Definition C.8 page 121)}$$

$$= \mathbf{B}^* \mathbf{f}(\mathbf{A}^* \mathbf{f}) \mathbf{f}(\mathbf{A}) \qquad \text{(Definition C.8 page 121)}$$

$$= \mathbf{A}^* \mathbf{f}(\mathbf{A}) \qquad \mathbf{A}^* \mathbf{f}(\mathbf{A}) \qquad \mathbf{A}^* \mathbf{f}(\mathbf{A}) \qquad \mathbf{A}^* \mathbf{f}(\mathbf$$

Theorem C.9. ²⁴ Let $\mathcal{B}(X, Y)$ be the space of BOUNDED LINEAR OPERATORS on normed linear spaces X and Y. Let \mathbf{B}^* be the adjoint of an operator \mathbf{B} .

$$\begin{array}{c} \mathsf{T} \\ \mathsf{H} \\ \mathsf{M} \end{array} | || \mathsf{B} ||| = \left| \left| \left| \mathsf{B}^* \right| \right| \qquad \forall \mathsf{B} \in \mathscr{B}(\mathsf{X}, \mathsf{Y})$$

№ Proof:

|||**B**|||
$$\triangleq \sup \{ ||\mathbf{B}x|| \mid ||x|| \le 1 \}$$
 by Definition C.6 page 117
 $\stackrel{?}{=} \sup \{ ||\mathbf{g}(\mathbf{B}x; \mathbf{y}^*)|| ||x|| \le 1, ||\mathbf{y}^*|| \le 1 \}$
 $= \sup \{ ||\mathbf{f}(\mathbf{x}; \mathbf{B}^* \mathbf{y}^*)|| ||x|| \le 1, ||\mathbf{y}^*|| \le 1 \}$
 $\triangleq \sup \{ ||\mathbf{B}^* \mathbf{y}^*|| \mid ||\mathbf{y}^*|| \le 1 \}$
 $\triangleq |||\mathbf{B}^*|||$ by Definition C.6 page 117

C.2.4 More properties



*Beginning with the third year of studies, most of my mathematical work was really started in conversations with Mazur and Banach. And according to Banach some of my own contributions were characterized by a certain "strangeness" in the formulation of problems and in the outline of possible proofs. As he told me once some years later, he was surprised how often these "strange" approaches really worked. Stanislaus M. Ulam (1909–1984), Polish mathematician ²⁵

²⁴ Rudin (1991) page 98



—>

Theorem C.10 (Mazur-Ulam theorem). ²⁶ Let $\phi \in \mathcal{L}(X, Y)$ be a function on normed linear spaces $(X, \|\cdot\|_X)$ and $(Y, \|\cdot\|_Y)$. Let $I \in \mathcal{L}(X, X)$ be the identity operator on $(X, \|\cdot\|_X)$.

1.
$$\frac{\phi^{-1}\phi = \phi\phi^{-1} = \mathbf{I}}{\text{bijective}}$$
2.
$$\|\phi \mathbf{x} - \phi \mathbf{y}\|_{Y} = \|\mathbf{x} - \mathbf{y}\|_{X} \quad \forall \mathbf{x}, \mathbf{y} \in X$$

$$\underset{isometric}{\text{isometric}}$$

$$\Rightarrow \underbrace{\phi([1 - \lambda]\mathbf{x} + \lambda \mathbf{y}) = [1 - \lambda]\phi\mathbf{x} + \lambda \phi\mathbf{y} \forall \lambda \in \mathbb{R}}_{affine}$$

♠PROOF: Proof not yet complete.

1. Let ψ be the reflection of z in X such that $\psi x = 2z - x$

(a)
$$\|\psi x - z\| = \|x - z\|$$

2. Let
$$\lambda \triangleq \sup_{g} \{ \|gz - z\| \}$$

3. Proof that $g \in W \implies g^{-1} \in W$:

Let
$$\hat{\mathbf{x}} \triangleq \mathbf{g}^{-1}\mathbf{x}$$
 and $\hat{\mathbf{y}} \triangleq \mathbf{g}^{-1}\mathbf{y}$.

$$||g^{-1}x - g^{-1}y|| = ||\hat{x} - \hat{y}||$$

$$= ||g\hat{x} - g\hat{y}||$$

$$= ||gg^{-1}x - gg^{-1}y||$$

$$= ||x - y||$$

by definition of \hat{x} and \hat{y} by left hypothesis by definition of \hat{x} and \hat{y} by definition of g^{-1}

4. Proof that gz = z:

$$2\lambda = 2 \sup \{ \|gz - z\| \}$$

$$\leq 2 \|gz - z\|$$

$$= \|2z - 2gz\|$$

$$= \|\varphi gz - gz\|$$

$$= \|g^{-1}\psi gz - g^{-1}gz\|$$

$$= \|g^{-1}\psi gz - z\|$$

$$= \|\varphi g^{-1}\psi gz - z\|$$

$$= \|g^*z - z\|$$

$$\leq \lambda$$

$$\implies 2\lambda \leq \lambda$$

$$\implies \lambda = 0$$

$$\implies gz = z$$

by definition of λ item (2) by definition of sup

by definition of ψ item (1) by item (3) by definition of g^{-1}

by definition of λ item (2)

5. Proof that $\phi\left(\frac{1}{2}\mathbf{x} + \frac{1}{2}\mathbf{y}\right) = \frac{1}{2}\phi\mathbf{x} + \frac{1}{2}\phi\mathbf{y}$:

$$\phi\left(\frac{1}{2}x + \frac{1}{2}y\right) =$$

$$= \frac{1}{2}\phi x + \frac{1}{2}\phi y$$

Ulam (1991) page 33

image: http://www-history.mcs.st-andrews.ac.uk/Biographies/Ulam.html

²⁶ ☑ Oikhberg and Rosenthal (2007) page 598, ② Väisälä (2003) page 634, ② Giles (2000) page 11, ② Dunford and Schwartz (1957) page 91, Mazur and Ulam (1932)

6. Proof that $\phi([1-\lambda]x + \lambda y) = [1-\lambda]\phi x + \lambda \phi y$:


Daniel J. Greenhoe

$$\phi([1 - \lambda]x + \lambda y) =$$

$$= [1 - \lambda]\phi x + \lambda \phi y$$

₽

Theorem C.11 (Neumann Expansion Theorem). ²⁷ Let $A \in X^X$ be an operator on a linear space X.



C.3 Operators on Inner product spaces

C.3.1 General Results

Definition C.9. ²⁸ Let $\Omega \triangleq (X, +, \cdot, (\mathbb{F}, +, \dot{\times}))$ be a linear space.

```
A function \langle \triangle \mid \nabla \rangle \in \mathbb{F}^{X \times X} is an inner product on \Omega if
                               \langle x \mid x \rangle \geq 0
                                                                                                                              (non-negative)
                                                                                                                                                                       and
                                 \langle \boldsymbol{x} \mid \boldsymbol{x} \rangle = 0 \iff \boldsymbol{x} = 0
                                                                                              \forall x \in X
                                                                                                                              (nondegenerate)
                                                                                                                                                                       and
                               \langle \alpha x \mid y \rangle = \alpha \langle x \mid y \rangle
DEF
                                                                                              \forall x,y \in X, \forall \alpha \in \mathbb{C}
                                                                                                                              (homogeneous)
                                                                                                                                                                       and
                    4. \langle x + y | u \rangle = \langle x | u \rangle + \langle y | u \rangle
                                                                                              \forall x, y, u \in X
                                                                                                                              (additive)
                                                                                                                                                                       and
                                  \langle x | y \rangle = \langle y | x \rangle^*
                                                                                              \forall x, y \in X
                                                                                                                              (conjugate symmetric).
        An inner product is also called a scalar product.
        An inner product space is the pair (\Omega, \langle \triangle \mid \nabla \rangle).
```

Theorem C.12. ²⁹ *Let* \mathbf{A} , $\mathbf{B} \in \mathcal{B}(\mathbf{X}, \mathbf{X})$ *be* Bounded linear operators *on an inner product space* $\mathbf{X} \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle | \nabla \rangle).$

Ţ.	$\langle \mathbf{B} \mathbf{x} \mid \mathbf{x} \rangle$	=	0	$\forall x \in X$	$\iff \\ \Leftrightarrow \\$	$\mathbf{B}\mathbf{x}$	=	0	∀ <i>x</i> ∈ <i>X</i>
M	$\langle \mathbf{A} \mathbf{x} \mid \mathbf{x} \rangle$	=	$\langle \mathbf{B} \mathbf{x} \mid \mathbf{x} \rangle$	$\forall x \in X$	\iff	A	=	B	

№ Proof:

²⁹ Rudin (1991) page 310 (Theorem 12.7, Corollary)



²⁷ Michel and Herget (1993) page 415

²⁸ Haaser and Sullivan (1991) page 277, Aliprantis and Burkinshaw (1998) page 276, Peano (1888b) page 72

1. Proof that $\langle \mathbf{B} \mathbf{x} \mid \mathbf{x} \rangle = 0 \implies \mathbf{B} \mathbf{x} = 0$:

$$0 = \langle \mathbf{B}(\mathbf{x} + \mathbf{B}\mathbf{x}) \mid (\mathbf{x} + \mathbf{B}\mathbf{x}) \rangle + i \langle \mathbf{B}(\mathbf{x} + i\mathbf{B}\mathbf{x}) \mid (\mathbf{x} + i\mathbf{B}\mathbf{x}) \rangle$$
 by left hypothesis
$$= \left\{ \langle \mathbf{B}\mathbf{x} + \mathbf{B}^2\mathbf{x}) \mid \mathbf{x} + \mathbf{B}\mathbf{x} \rangle \right\} + i \left\{ \langle \mathbf{B}\mathbf{x} + i\mathbf{B}^2\mathbf{x}) \mid \mathbf{x} + i\mathbf{B}\mathbf{x} \rangle \right\}$$
 by Definition C.4 page 113 by Definition C.9 page 124
$$+ i \left\{ \langle \mathbf{B}\mathbf{x} \mid \mathbf{x} \rangle + i \langle \mathbf{B}\mathbf{x} \mid \mathbf{B}\mathbf{x} \rangle + i \langle \mathbf{B}^2\mathbf{x} \mid \mathbf{x} \rangle - i^2 \langle \mathbf{B}^2\mathbf{x} \mid \mathbf{B}\mathbf{x} \rangle \right\}$$
 by Definition C.9 page 124
$$+ i \left\{ \langle \mathbf{B}\mathbf{x} \mid \mathbf{x} \rangle - i \langle \mathbf{B}\mathbf{x} \mid \mathbf{B}\mathbf{x} \rangle + i \langle \mathbf{B}^2\mathbf{x} \mid \mathbf{x} \rangle - i^2 \langle \mathbf{B}^2\mathbf{x} \mid \mathbf{B}\mathbf{x} \rangle \right\}$$
 by left hypothesis
$$= \left\{ \langle \mathbf{B}\mathbf{x} \mid \mathbf{B}\mathbf{x} \rangle + \langle \mathbf{B}^2\mathbf{x} \mid \mathbf{x} \rangle \right\} + \left\{ \langle \mathbf{B}\mathbf{x} \mid \mathbf{B}\mathbf{x} \rangle - \langle \mathbf{B}^2\mathbf{x} \mid \mathbf{x} \rangle \right\}$$
 by left hypothesis
$$= \left\{ \langle \mathbf{B}\mathbf{x} \mid \mathbf{B}\mathbf{x} \rangle + \langle \mathbf{B}^2\mathbf{x} \mid \mathbf{x} \rangle \right\} + \left\{ \langle \mathbf{B}\mathbf{x} \mid \mathbf{B}\mathbf{x} \rangle - \langle \mathbf{B}^2\mathbf{x} \mid \mathbf{x} \rangle \right\}$$
 by Definition C.5 page 116

- 2. Proof that $\langle \mathbf{B} \mathbf{x} \mid \mathbf{x} \rangle = 0 \iff \mathbf{B} \mathbf{x} = 0$: by property of inner products.
- 3. Proof that $\langle \mathbf{A} \mathbf{x} \mid \mathbf{x} \rangle = \langle \mathbf{B} \mathbf{x} \mid \mathbf{x} \rangle \implies \mathbf{A} \stackrel{\circ}{=} \mathbf{B}$:

$$0 = \langle \mathbf{A} x \mid \mathbf{x} \rangle - \langle \mathbf{B} \mathbf{x} \mid \mathbf{x} \rangle \qquad \text{by left hypothesis}$$

$$= \langle \mathbf{A} \mathbf{x} - \mathbf{B} \mathbf{x} \mid \mathbf{x} \rangle \qquad \text{by additivity property of } \langle \triangle \mid \nabla \rangle \text{ (Definition C.9 page 124)}$$

$$= \langle (\mathbf{A} - \mathbf{B}) \mathbf{x} \mid \mathbf{x} \rangle \qquad \text{by definition of operator addition}$$

$$\implies (\mathbf{A} - \mathbf{B}) \mathbf{x} = 0 \qquad \text{by item 1}$$

$$\implies \mathbf{A} = \mathbf{B} \qquad \text{by definition of operator subtraction}$$

4. Proof that $\langle \mathbf{A} \mathbf{x} \mid \mathbf{x} \rangle = \langle \mathbf{B} \mathbf{x} \mid \mathbf{x} \rangle \iff \mathbf{A} \stackrel{\circ}{=} \mathbf{B}$:

$$\langle \mathbf{A} \mathbf{x} \mid \mathbf{x} \rangle = \langle \mathbf{B} \mathbf{x} \mid \mathbf{x} \rangle$$

by $\mathbf{A} \stackrel{\circ}{=} \mathbf{B}$ hypothesis

C.3.2 Operator adjoint

A fundamental concept of operators on inner product spaces is the *operator adjoint* (Proposition C.3 page 125). The adjoint of an operator is a kind of generalization of the conjugate of a complex number in that

- Both are *star-algebras* (Theorem C.13 page 126).
- Both support decomposition into "real" and "imaginary" parts (Theorem F.3 page 150).

Structurally, the operator adjoint provides a convenient symmetric relationship between the *range space* and *Null Space* of an operator (Theorem C.14 page 127).

Proposition C.3. ³⁰ Let $\mathcal{B}(H, H)$ be the space of Bounded Linear Operators (Definition C.7 page 120) on a Hilbert space H.

An operator
$$\mathbf{B}^*$$
 is the **adjoint** of $\mathbf{B} \in \mathcal{B}(H, H)$ if $\langle \mathbf{B} x | y \rangle = \langle x | \mathbf{B}^* y \rangle$ $\forall x, y \in H$.

^ℚProof:

Frames and Bases Structure and Design [VERSION 021] thtps://github.com/dgreenhoe/pdfs/blob/master/msdframes.pdf



³⁰ Michel and Herget (1993) page 220, Rudin (1991) page 311, Giles (2000) page 182, von Neumann (1929) page 49, Stone (1932) page 41

- 1. For fixed y, $f(x) \triangleq \langle x | y \rangle$ is a functional in \mathbb{F}^X .
- 2. \mathbf{B}^* is the *adjoint* of \mathbf{B} because

$$\langle \mathbf{B}x \mid y \rangle \triangleq \mathsf{f}(\mathbf{B}x)$$

$$\triangleq \mathbf{B}^*\mathsf{f}(x) \qquad \text{by definition of } operator \, adjoint \qquad \text{(Definition C.8 page 121)}$$

$$= \langle x \mid \mathbf{B}^*y \rangle$$

Example C.2.

In matrix algebra ("linear algebra")

The inner product operation $\langle x \mid y \rangle$ is represented by $y^H x$.

The linear operator is represented as a matrix A.

The operation of A on a vector x is represented as

The adjoint of matrix A is the Hermitian matrix A^H .

♥Proof:

E X

$$\langle Ax \mid y \rangle \triangleq y^H Ax = [(Ax)^H y]^H = [x^H A^H y]^H = (A^H y)^H x \triangleq \langle x \mid A^H y \rangle$$

Structures that satisfy the four conditions of the next theorem are known as *-algebras ("star-algebras" (Definition F.3 page 148). Other structures which are *-algebras include the *field of complex numbers* $\mathbb C$ and any *ring of complex square* $n \times n$ *matrices*. 31

Theorem C.13 (operator star-algebra). ³² *Let* H *be* a HILBERT SPACE *with operators* A, $B \in \mathcal{B}(H, H)$ *and with adjoints* A^* , $B^* \in \mathcal{B}(H, H)$. *Let* $\bar{\alpha}$ *be the complex conjugate of some* $\alpha \in \mathbb{C}$.

-	The pair $(H, *)$ is a *-algebra (star-algebra). In particular,									
Т	1.	$(\mathbf{A} \stackrel{\circ}{+} \mathbf{B})^*$	=	$A^* + B^*$	∀ A , B ∈ <i>H</i>	(DISTRIBUTIVE)	and			
н	2.	$(\alpha \mathbf{A})^*$	=	$\bar{\alpha}\mathbf{A}^*$	∀ A , B ∈ <i>H</i>	(CONJUGATE LINEAR)	and			
M	3.	$(AB)^*$	=	$\mathbf{B}^*\mathbf{A}^*$	∀ A , B ∈ <i>H</i>	(ANTIAUTOMORPHIC)	and			
	4.	\mathbf{A}^{**}	=	A	$\forall A,B \in H$	(INVOLUTARY)				

[♠]Proof:

³² Halmos (1998a) pages 39–40, Rudin (1991) page 311



³¹ Sakai (1998) page 1

$\langle x \mid (\mathbf{A}\mathbf{B})^* y \rangle = \langle (\mathbf{A}\mathbf{B})x \mid y \rangle$	by definition of adjoint	(Proposition C.3 page 125)
$= \langle \mathbf{A}(\mathbf{B}\mathbf{x}) \mid \mathbf{y} \rangle$	by definition of operator multiplication	
$= \langle (\mathbf{B}\mathbf{x}) \mid \mathbf{A}^* \mathbf{y} \rangle$	by definition of adjoint	(Proposition C.3 page 125)
$= \langle x \mid \mathbf{B}^* \mathbf{A}^* y \rangle$	by definition of adjoint	(Proposition C.3 page 125)
$\langle x \mid \mathbf{A}^{**} y \rangle = \langle \mathbf{A}^* x \mid y \rangle$	by definition of adjoint	(Proposition C.3 page 125)
$=\langle y \mid \mathbf{A}^* \mathbf{x} \rangle^*$	by definition of inner product	(Definition C.9 page 124)
$= \langle \mathbf{A} \mathbf{y} \mathbf{x} \rangle^*$	by definition of adjoint	(Proposition C.3 page 125)
$=\langle x \mid \mathbf{A}y \rangle$	by definition of inner product	(Definition C.9 page 124)

Theorem C.14. 33 Let Y^X be the set of all operators from a linear space X to a linear space Y. Let $\mathcal{N}(\mathbf{L})$ be the Null Space of an operator \mathbf{L} in $\mathbf{Y}^{\mathbf{X}}$ and $\mathbf{\mathcal{F}}(\mathbf{L})$ the image set of \mathbf{L} in $\mathbf{Y}^{\mathbf{X}}$.

$$\begin{array}{c|c} \mathbf{T} & \mathcal{N}(\mathbf{A}) = \boldsymbol{\mathscr{I}}(\mathbf{A}^*)^{\perp} \\ \mathbf{M} & \mathcal{N}(\mathbf{A}^*) = \boldsymbol{\mathscr{I}}(\mathbf{A})^{\perp} \\ \end{array}$$

[♠]Proof:

$$\begin{split} \mathscr{F}(\mathbf{A}^*)^\perp &= \big\{ y \in H | \, \langle y \, | \, u \rangle = 0 \quad \forall u \in \mathscr{F}(\mathbf{A}^*) \big\} \\ &= \big\{ y \in H | \, \langle y \, | \, x \rangle = 0 \quad \forall x \in H \big\} \\ &= \{ y \in H | \, \langle \mathbf{A} y \, | \, x \rangle = 0 \quad \forall x \in H \} \\ &= \{ y \in H | \mathbf{A} y = 0 \} \\ &= \mathscr{N}(\mathbf{A}) \end{split} \qquad \text{by definition of } \mathscr{N}(\mathbf{A}) \end{split}$$

$$\mathscr{F}(\mathbf{A})^\perp &= \{ y \in H | \, \langle y \, | \, u \rangle = 0 \quad \forall u \in \mathscr{F}(\mathbf{A}) \} \\ &= \{ y \in H | \, \langle y \, | \, u \rangle = 0 \quad \forall u \in \mathscr{F}(\mathbf{A}) \} \\ &= \{ y \in H | \, \langle y \, | \, x \rangle = 0 \quad \forall x \in H \} \\ &= \{ y \in H | \, \langle \mathbf{A}^* y \, | \, x \rangle = 0 \quad \forall x \in H \} \\ &= \{ y \in H | \mathbf{A}^* y = 0 \} \\ &= \mathscr{N}(\mathbf{A}^*) \qquad \text{by definition of } \mathscr{N}(\mathbf{A}) \end{split}$$

$$(\text{Proposition C.3 page 125})$$

Special Classes of Operators C.4

Projection operators C.4.1

Definition C.10. ³⁴ Let $\mathcal{B}(X, Y)$ be the space of BOUNDED LINEAR OPERATORS on normed linear spaces X and Y. Let P be a bounded linear operator in $\mathcal{B}(X, Y)$.



³³ Rudin (1991) page 312

³⁴ ■ Rudin (1991) page 126 (5.15 Projections), ■ Kubrusly (2001) page 70, ■ Bachman and Narici (1966) page 26, Halmos (1958) page 73 (§41. Projections)



Theorem C.15. ³⁵ Let $\mathcal{B}(X, Y)$ be the space of BOUNDED LINEAR OPERATORS on normed linear spaces X and Y. Let P be a bounded linear operator in $\mathcal{B}(X, Y)$ with NULL SPACE $\mathcal{N}(P)$ and IMAGE SET $\mathcal{F}(P)$.

$$\begin{bmatrix}
1. & \mathbf{P}^2 &= \mathbf{P} & (\mathbf{P} \text{ is a projection operator}) & and \\
2. & \mathbf{\Omega} &= \mathbf{X} + \mathbf{Y} & (\mathbf{Y} \text{ compliments } \mathbf{X} \text{ in } \mathbf{\Omega}) & and \\
3. & \mathbf{P}\mathbf{\Omega} &= \mathbf{X} & (\mathbf{P} \text{ projects onto } \mathbf{X})
\end{bmatrix}$$

$$\begin{vmatrix}
1. & \mathbf{F}(\mathbf{P}) &= \mathbf{X} & and \\
2. & \mathcal{N}(\mathbf{P}) &= \mathbf{Y} & and \\
3. & \mathbf{\Omega} &= \mathbf{\mathcal{F}}(\mathbf{P}) + \mathcal{N}(\mathbf{P})$$

№PROOF:

$$\begin{split} \boldsymbol{\mathcal{J}}(\mathbf{P}) &= \mathbf{P}\boldsymbol{\mathcal{Q}} \\ &= \mathbf{P}(\boldsymbol{\mathcal{Q}}_1 + \boldsymbol{\mathcal{Q}}_2) \\ &= \mathbf{P}\boldsymbol{\mathcal{Q}}_1 + \mathbf{P}\boldsymbol{\mathcal{Q}}_2 \\ &= \boldsymbol{\mathcal{Q}}_1 + \{0\} \\ &= \boldsymbol{\mathcal{Q}}_1 \end{split}$$

$$\mathcal{N}(\mathbf{P}) = \{ \mathbf{x} \in \mathbf{\Omega} | \mathbf{P} \mathbf{x} = \mathbf{0} \}$$

$$= \{ \mathbf{x} \in (\mathbf{\Omega}_1 + \mathbf{\Omega}_2) | \mathbf{P} \mathbf{x} = \mathbf{0} \}$$

$$= \{ \mathbf{x} \in \mathbf{\Omega}_1 | \mathbf{P} \mathbf{x} = \mathbf{0} \} + \{ \mathbf{x} \in \mathbf{\Omega}_2 | \mathbf{P} \mathbf{x} = \mathbf{0} \}$$

$$= \{ \mathbf{0} \} + \mathbf{\Omega}_2$$

$$= \mathbf{\Omega}_2$$

Theorem C.16. ³⁶ Let $\mathcal{B}(X, Y)$ be the space of BOUNDED LINEAR OPERATORS on normed linear spaces X and Y. Let P be a bounded linear operator in $\mathcal{B}(X, Y)$.



♥PROOF:

Proof that
$$\mathbf{P}^2 = \mathbf{P} \implies (\mathbf{I} - \mathbf{P})^2 = (\mathbf{I} - \mathbf{P})$$
:

$$(I - P)^{2} = (I - P)(I - P)$$

$$= I(I - P) + (-P)(I - P)$$

$$= I - P - PI + P^{2}$$

$$= I - P - P + P$$

$$= I - P$$

by left hypothesis

4 Proof that
$$\mathbf{P}^2 = \mathbf{P} \iff (\mathbf{I} - \mathbf{P})^2 = (\mathbf{I} - \mathbf{P})$$
:

$$\mathbf{P}^{2} = \underbrace{\mathbf{I} - \mathbf{P} - \mathbf{P} + \mathbf{P}^{2}}_{(\mathbf{I} - \mathbf{P})^{2}} - (\mathbf{I} - \mathbf{P} - \mathbf{P})$$

$$= (\mathbf{I} - \mathbf{P})^{2} - (\mathbf{I} - \mathbf{P} - \mathbf{P})$$

$$= (\mathbf{I} - \mathbf{P}) - (\mathbf{I} - \mathbf{P} - \mathbf{P})$$

$$= \mathbf{P}$$

by right hypothesis

³⁶ Michel and Herget (1993) page 121



₽

³⁵ Michel and Herget (1993) pages 120–121

Theorem C.17. ³⁷ *Let* H *be a* Hilbert space *and* P *an operator in* H^H *with adjoint* P^* , Null Space $\mathcal{N}(\mathbf{P})$, and image set $\mathcal{J}(\mathbf{P})$.

If **P** is a projection operator, then the following are equivalent: (P is self-adjoint) T H M $\mathbf{P}^*\mathbf{P} = \mathbf{P}\mathbf{P}^*$ (P is NORMAL) $\mathscr{J}(\mathbf{P}) = \mathscr{N}(\mathbf{P})^{\perp}$ 4. $\langle \mathbf{P} \mathbf{x} \mid \mathbf{x} \rangle = \|\mathbf{P} \mathbf{x}\|^2$

PROOF: This proof is incomplete at this time.

Proof that $(1) \Longrightarrow (2)$:

$$\mathbf{P}^*\mathbf{P} = \mathbf{P}^{**}\mathbf{P}^*$$
by (1)
$$= \mathbf{PP}^*$$
by Theorem C.13 page 126

Proof that $(1) \Longrightarrow (3)$:

$$\mathcal{J}(\mathbf{P}) = \mathcal{N}(\mathbf{P}^*)^{\perp}$$
 by Theorem C.14 page 127
= $\mathcal{N}(\mathbf{P})^{\perp}$ by (1)

Proof that $(3) \Longrightarrow (4)$:

Proof that $(4) \Longrightarrow (1)$:

Self Adjoint Operators C.4.2

Definition C.11. ³⁸ Let $\mathbf{B} \in \mathcal{B}(\mathbf{H}, \mathbf{H})$ be a BOUNDED operator with adjoint \mathbf{B}^* on a Hilbert space Н.

The operator **B** is said to be **self-adjoint** or **hermitian** if $\mathbf{B} \stackrel{\circ}{=} \mathbf{B}^*$.

Example C.3 (Autocorrelation operator). Let x(t) be a random process with autocorrelation $R_{xx}(t, u) \triangleq \underbrace{E[x(t)x^*(u)]}_{\text{expectation}}$

Let an autocorrelation operator **R** be defined as [**R**f](t) $\triangleq \int_{\mathbb{R}} R_{\underline{x}\underline{x}}(t,u) f(u) du$.

 $\mathbf{R} = \mathbf{R}^*$ (The auto-correlation operator **R** is *self-adjoint*)

Theorem C.18. ³⁹ Let $S: H \to H$ be an operator over a Hilbert space H with eigenvalues $\{\lambda_n\}$ and eigenfunctions $\{\psi_n\}$ such that $\mathbf{S}\psi_n = \lambda_n \psi_n$ and let $\|\mathbf{x}\| \triangleq \sqrt{\langle \mathbf{x} \mid \mathbf{x} \rangle}$.

$$\left\{ \begin{array}{l} \mathbf{T} \\ \mathbf{H} \\ \mathbf{N} \end{array} \right\} \left\{ \begin{array}{l} \mathbf{S} = \mathbf{S}^* \\ \mathbf{S} \text{ is } \textit{self adjoint} \end{array} \right\} \implies \left\{ \begin{array}{l} 1. & \langle \mathbf{S} \mathbf{x} \mid \mathbf{x} \rangle \in \mathbb{R} \\ 2. & \lambda_n \in \mathbb{R} \\ 3. & \lambda_n \neq \lambda_m \implies \langle \psi_n \mid \psi_m \rangle = 0 \end{array} \right. \text{ (the hermitian quadratic form of \mathbf{S} is real-valued)} \\ \left\{ \begin{array}{l} \mathbf{S} = \mathbf{S}^* \\ \mathbf{S} \text{ is } \textit{self adjoint} \end{array} \right\} \implies \left\{ \begin{array}{l} 1. & \langle \mathbf{S} \mathbf{x} \mid \mathbf{x} \rangle \in \mathbb{R} \\ 2. & \lambda_n \in \mathbb{R} \\ 3. & \lambda_n \neq \lambda_m \implies \langle \psi_n \mid \psi_m \rangle = 0 \end{array} \right. \text{ (eigenvectors are ORTHOGONAL)}$$

³⁹ 🛮 Lax (2002) pages 315–316, 📳 Keener (1988) pages 114–119, 📳 Bachman and Narici (1966) page 24 (Theorem 2.1), Bertero and Boccacci (1998) page 225 (\$"9.2 SVD of a matrix ... If all eigenvectors are normalized...")



³⁷ Rudin (1991) page 314

³⁸Historical works regarding self-adjoint operators: a von Neumann (1929) page 49, "linearer Operator R selbstadjungiert oder Hermitesch",

Stone (1932) page 50 ⟨"self-adjoint transformations"⟩

1. Proof that $S = S^* \implies \langle Sx \mid x \rangle \in \mathbb{R}$:

Daniel J. Greenhoe

$$\langle x \mid Sx \rangle = \langle Sx \mid x \rangle$$
 by left hypothesis
= $\langle x \mid Sx \rangle^*$ by definition of $\langle \triangle \mid \nabla \rangle$ Definition C.9 page 124

2. Proof that $S = S^* \implies \lambda_n \in \mathbb{R}$:

$$\lambda_{n} \|\psi_{n}\|^{2} = \lambda_{n} \langle \psi_{n} | \psi_{n} \rangle \qquad \text{by definition}$$

$$= \langle \lambda_{n} \psi_{n} | \psi_{n} \rangle \qquad \text{by definition of } \langle \triangle | \nabla \rangle \text{ Definition C.9 page 124}$$

$$= \langle \mathbf{S} \psi_{n} | \psi_{n} \rangle \qquad \text{by definition of eigenpairs}$$

$$= \langle \psi_{n} | \mathbf{S} \psi_{n} \rangle \qquad \text{by left hypothesis}$$

$$= \langle \psi_{n} | \lambda_{n} \psi_{n} \rangle \qquad \text{by definition of eigenpairs}$$

$$= \lambda_{n}^{*} \langle \psi_{n} | \psi_{n} \rangle \qquad \text{by definition of } \langle \triangle | \nabla \rangle \text{ Definition C.9 page 124}$$

$$= \lambda_{n}^{*} \|\psi_{n}\|^{2} \qquad \text{by definition}$$

3. Proof that $S = S^* \implies [\lambda_n \neq \lambda_m \implies \langle \psi_n | \psi_m \rangle = 0]$:

$$\lambda_{n} \langle \psi_{n} | \psi_{m} \rangle = \langle \lambda_{n} \psi_{n} | \psi_{m} \rangle \qquad \text{by definition of } \langle \triangle | \nabla \rangle \text{ Definition C.9 page 124}$$

$$= \langle \mathbf{S} \psi_{n} | \psi_{m} \rangle \qquad \text{by definition of eigenpairs}$$

$$= \langle \psi_{n} | \mathbf{S} \psi_{m} \rangle \qquad \text{by left hypothesis}$$

$$= \langle \psi_{n} | \lambda_{m} \psi_{m} \rangle \qquad \text{by definition of eigenpairs}$$

$$= \lambda_{m}^{*} \langle \psi_{n} | \psi_{m} \rangle \qquad \text{by definition of } \langle \triangle | \nabla \rangle \text{ Definition C.9 page 124}$$

$$= \lambda_{m} \langle \psi_{n} | \psi_{m} \rangle \qquad \text{because } \lambda_{m} \text{ is real}$$

This implies for $\lambda_n \neq \lambda_m \neq 0$, $\langle \psi_n | \psi_m \rangle = 0$.


₽

C.4.3 Normal Operators

Definition C.12. 40 Let $\mathcal{B}(X, Y)$ be the space of BOUNDED LINEAR OPERATORS on normed linear spaces X and Y. Let \mathbb{N}^* be the adjoint of an operator $\mathbb{N} \in \mathcal{B}(X, Y)$.



Theorem C.19. ⁴¹ Let $\mathcal{B}(H, H)$ be the space of BOUNDED LINEAR OPERATORS on a HILBERT SPACE H. Let $\mathcal{N}(N)$ be the NULL SPACE of an operator N in $\mathcal{B}(H, H)$ and $\mathcal{F}(N)$ the image set of N in $\mathcal{B}(H, H)$.



⁴¹ Rudin (1991) pages 312–313



№PROOF:

1. Proof that $\mathbf{N}^*\mathbf{N} = \mathbf{N}\mathbf{N}^* \implies \|\mathbf{N}^*x\| = \|\mathbf{N}x\|$:

$$||\mathbf{N}x||^2 = \langle \mathbf{N}x \mid \mathbf{N}x \rangle$$
 by definition

$$= \langle x \mid \mathbf{N}^* \mathbf{N}x \rangle$$
 by Proposition C.3 page 125 (definition of \mathbf{N}^*)

$$= \langle x \mid \mathbf{N}\mathbf{N}^* x \rangle$$
 by left hypothesis (\mathbf{N} is normal)

$$= \langle \mathbf{N}x \mid \mathbf{N}^* x \rangle$$
 by Proposition C.3 page 125 (definition of \mathbf{N}^*)

$$= ||\mathbf{N}^* x||^2$$
 by definition

2. Proof that $N^*N = NN^* \iff ||N^*x|| = ||Nx||$:

$$\langle \mathbf{N}^* \mathbf{N} \mathbf{x} \mid \mathbf{x} \rangle = \langle \mathbf{N} \mathbf{x} \mid \mathbf{N}^{**} \mathbf{x} \rangle \qquad \text{by Proposition C.3 page 125 (definition of } \mathbf{N}^*)$$

$$= \langle \mathbf{N} \mathbf{x} \mid \mathbf{N} \mathbf{x} \rangle \qquad \text{by Theorem C.13 page 126 (property of adjoint)}$$


$$= \|\mathbf{N} \mathbf{x}\|^2 \qquad \text{by definition}$$

$$= \|\mathbf{N}^* \mathbf{x}\|^2 \qquad \text{by right hypothesis } (\|\mathbf{N}^* \mathbf{x}\| = \|\mathbf{N} \mathbf{x}\|)$$

$$= \langle \mathbf{N}^* \mathbf{x} \mid \mathbf{N}^* \mathbf{x} \rangle \qquad \text{by definition}$$

$$= \langle \mathbf{N} \mathbf{N}^* \mathbf{x} \mid \mathbf{x} \rangle \qquad \text{by Proposition C.3 page 125 (definition of } \mathbf{N}^*)$$


Theorem C.20. ⁴² Let $\mathcal{B}(H, H)$ be the space of BOUNDED LINEAR OPERATORS on a HILBERT SPACE H. Let $\mathcal{N}(N)$ be the NULL SPACE of an operator N in $\mathcal{B}(H, H)$ and $\mathcal{F}(N)$ the image set of N in $\mathcal{B}(H, H)$.



№PROOF:

$$\mathcal{N}(\mathbf{N}^*) = \left\{ x | \mathbf{N}^* x = 0 \quad \forall x \in \mathbf{X} \right\}$$
 by definition of *Null Space*
$$= \left\{ x | \| \mathbf{N}^* x \| = 0 \quad \forall x \in \mathbf{X} \right\}$$
 by definition of $\| \cdot \|$ (Definition C.5 page 116)
$$= \left\{ x | \| \mathbf{N} x \| = 0 \quad \forall x \in \mathbf{X} \right\}$$
 by definition of $\| \cdot \|$ (Definition C.5 page 116)
$$= \mathcal{N}(\mathbf{N})$$
 by definition of *Null Space* \mathcal{N}

Theorem C.21. ⁴³ Let $\mathcal{B}(H, H)$ be the space of bounded linear operators on a Hilbert space H. Let $\mathcal{N}(N)$ be the Null Space of an operator N in $\mathcal{B}(H, H)$ and $\mathcal{F}(N)$ the image set of N in $\mathcal{B}(H, H)$.



[♠]PROOF: The proof in (1) is flawed. This implies that (2) is also flawed. *■* Rudin (1991) page 313 claims both to be true.

Frames and Bases Structure and Design [VERSION 021] thtps://github.com/dgreenhoe/pdfs/blob/master/msdframes.pdf



⁴² Rudin (1991) pages 312–313

⁴³ Rudin (1991) pages 312–313

1. Proof that $N^*N = NN^* \implies N^*\psi = \lambda^*\psi$:

Daniel J. Greenhoe

$$\mathbf{N}\psi = \lambda\psi$$

$$\Longleftrightarrow$$

$$0 = \mathcal{N}(\mathbf{N} - \lambda \mathbf{I})$$

$$= \mathcal{N}([\mathbf{N} - \lambda \mathbf{I}]^*)$$

$$= \mathcal{N}(\mathbf{N}^* - [\lambda \mathbf{I}]^*)$$
by $\mathcal{N}(\mathbf{N}) = \mathcal{N}(\mathbf{N}^*)$

$$= \mathcal{N}(\mathbf{N}^* - \lambda^* \mathbf{I}^*)$$
by Theorem C.13 page 126
$$= \mathcal{N}(\mathbf{N}^* - \lambda^* \mathbf{I})$$

$$\Longrightarrow$$

$$(\mathbf{N}^* - \lambda^* \mathbf{I})\psi = 0$$

$$\Longleftrightarrow \mathbf{N}^*\psi = \lambda^*\psi$$

2. Proof that $\mathbf{N}^*\mathbf{N} = \mathbf{N}\mathbf{N}^* \implies [\lambda_n \neq \lambda_m \implies \langle \psi_n | \psi_m \rangle = 0]$:

$$\begin{array}{ll} \lambda_{n} \left\langle \psi_{n} \mid \psi_{m} \right\rangle = \left\langle \lambda_{n} \psi_{n} \mid \psi_{m} \right\rangle & \text{by definition of } \left\langle \triangle \mid \bigtriangledown \right\rangle \text{ Definition C.9 page 124} \\ = \left\langle \mathbf{N} \psi_{n} \mid \psi_{m} \right\rangle & \text{by definition of eigenpairs} \\ = \left\langle \psi_{n} \mid \mathbf{N}^{*} \psi_{m} \right\rangle & \text{by Proposition C.3 page 125 (definition of adjoint)} \\ = \left\langle \psi_{n} \mid \lambda_{m}^{*} \psi_{m} \right\rangle & \text{by (4.)} \\ = \lambda_{m} \left\langle \psi_{n} \mid \psi_{m} \right\rangle & \text{by definition of } \left\langle \triangle \mid \bigtriangledown \right\rangle \text{ Definition C.9 page 124} \end{array}$$

This implies for $\lambda_n \neq \lambda_m \neq 0$, $\langle \psi_n | \psi_m \rangle = 0$.

Isometric operators C.4.4

An operator on a pair of normed linear spaces is *isometric* (next definition) if it is an *isometry*.

Definition C.13. Let $(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \|\cdot\|)$ and $(Y, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \|\cdot\|)$ be NORMED LINEAR SPACES (Definition C.5 page 116).

An operator
$$\mathbf{M} \in \mathcal{L}(\mathbf{X}, \mathbf{Y})$$
 is **isometric** if $\|\mathbf{M}\mathbf{x}\| = \|\mathbf{x}\| \quad \forall \mathbf{x} \in X$.

 $\textbf{Theorem C.22.} \overset{44}{\overset{}{}} Let \big(X,\,+,\,\cdot,\,(\mathbb{F},\,\dot{+},\,\dot{\times}),\,\|\cdot\|\big) \, and \, \big(Y,\,+,\,\cdot,\,(\mathbb{F},\,\dot{+},\,\dot{\times}),\,\|\cdot\|\big) \, be \, \text{normed linear spaces.}$ Let **M** be a linear operator in $\mathcal{L}(\mathbf{X}, \mathbf{Y})$.

$$\|\mathbf{M}x\| = \|x\| \quad \forall x \in X \qquad \Longleftrightarrow \qquad \|\mathbf{M}x - \mathbf{M}y\| = \|x - y\| \quad \forall x, y \in X$$
isometric in length isometric in distance

^ℚProof:

1. Proof that $||Mx|| = ||x|| \implies ||Mx - My|| = ||x - y||$:

$$\|\mathbf{M}x - \mathbf{M}y\| = \|\mathbf{M}(x - y)\|$$
 by definition of linear operators (Definition C.4 page 113)
 $= \|\mathbf{M}u\|$ let $u \triangleq x - y$
 $= \|x - y\|$ by left hypothesis

⁴⁴ Kubrusly (2001) page 239 (Proposition 4.37), Berberian (1961) page 27 (Theorem IV.7.5)




2. Proof that $||Mx|| = ||x|| \iff ||Mx - My|| = ||x - y||$:

$$\|\mathbf{M}x\| = \|\mathbf{M}(x - 0)\|$$

= $\|\mathbf{M}x - \mathbf{M}0\|$ by definition of linear operators (Definition C.4 page 113)
= $\|x - 0\|$ by right hypothesis
= $\|x\|$

Isometric operators have already been defined (Definition C.13 page 132) in the more general normed linear spaces, while Theorem C.22 (page 132) demonstrated that in a normed linear space X, $||Mx|| = ||x|| \iff ||Mx - My|| = ||x - y||$ for all $x, y \in X$. Here in the more specialized inner product spaces, Theorem C.23 (next) demonstrates two additional equivalent properties.

Theorem C.23. ⁴⁵ Let $\mathcal{B}(\mathbf{X}, \mathbf{X})$ be the space of BOUNDED LINEAR OPERATORS on a normed linear space $\mathbf{X} \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \|\cdot\|)$. Let \mathbf{N} be a bounded linear operator in $\mathcal{L}(\mathbf{X}, \mathbf{X})$, and \mathbf{I} the identity operator in $\mathcal{L}(\mathbf{X}, \mathbf{X})$. Let $\|\mathbf{x}\| \triangleq \sqrt{\langle \mathbf{x} | \mathbf{x} \rangle}$.



№PROOF:

1. Proof that $(1) \Longrightarrow (2)$:

$$\langle \mathbf{M} \mathbf{x} \mid \mathbf{M} \mathbf{y} \rangle = \langle \mathbf{x} \mid \mathbf{M}^* \mathbf{M} \mathbf{y} \rangle$$
 by Proposition C.3 page 125 (definition of adjoint)

$$= \langle \mathbf{x} \mid \mathbf{I} \mathbf{y} \rangle$$
 by (1)

$$= \langle \mathbf{x} \mid \mathbf{y} \rangle$$
 by Definition C.3 page 112 (definition of **I**)

2. Proof that $(2) \Longrightarrow (4)$:

$$\|\mathbf{M}x\| = \sqrt{\langle \mathbf{M}x \, | \, \mathbf{M}x \rangle}$$
 by definition of $\|\cdot\|$ by right hypothesis $\|\mathbf{M}x\| = \|\mathbf{M}x\|$ by definition of $\|\cdot\|$

3. Proof that $(2) \Leftarrow (4)$:

$$4 \langle \mathbf{M} \mathbf{x} | \mathbf{M} \mathbf{y} \rangle = \|\mathbf{M} \mathbf{x} + \mathbf{M} \mathbf{y}\|^{2} - \|\mathbf{M} \mathbf{x} - \mathbf{M} \mathbf{y}\|^{2} + i \|\mathbf{M} \mathbf{x} + i \mathbf{M} \mathbf{y}\|^{2} - i \|\mathbf{M} \mathbf{x} - i \mathbf{M} \mathbf{y}\|^{2}$$
by polarization id.

$$= \|\mathbf{M} (\mathbf{x} + \mathbf{y})\|^{2} - \|\mathbf{M} (\mathbf{x} - \mathbf{y})\|^{2} + i \|\mathbf{M} (\mathbf{x} + i \mathbf{y})\|^{2} - i \|\mathbf{M} (\mathbf{x} - i \mathbf{y})\|^{2}$$
by Definition C.4

$$= \|\mathbf{x} + \mathbf{y}\|^{2} - \|\mathbf{x} - \mathbf{y}\|^{2} + i \|\mathbf{x} + i \mathbf{y}\|^{2} - i \|\mathbf{x} - i \mathbf{y}\|^{2}$$
by left hypothesis

4. Proof that (3) \iff (4): by Theorem C.22 page 132



5. Proof that $(4) \Longrightarrow (1)$:

$$\langle \mathbf{M}^* \mathbf{M} \boldsymbol{x} \mid \boldsymbol{x} \rangle = \langle \mathbf{M} \boldsymbol{x} \mid \mathbf{M}^{**} \boldsymbol{x} \rangle \qquad \text{by Proposition C.3 page 125 (definition of adjoint)}$$

$$= \langle \mathbf{M} \boldsymbol{x} \mid \mathbf{M} \boldsymbol{x} \rangle \qquad \text{by Theorem C.13 page 126 (property of adjoint)}$$

$$= \|\mathbf{M} \boldsymbol{x}\|^2 \qquad \text{by definition}$$


$$= \|\boldsymbol{x}\|^2 \qquad \text{by left hypothesis with } \boldsymbol{y} = 0$$

$$= \langle \boldsymbol{x} \mid \boldsymbol{x} \rangle \qquad \text{by definition}$$

$$= \langle \mathbf{I} \boldsymbol{x} \mid \boldsymbol{x} \rangle \qquad \text{by Definition C.3 page 112 (definition of I)}$$

$$\Rightarrow \quad \mathbf{M}^* \mathbf{M} = \mathbf{I} \qquad \forall \boldsymbol{x} \in X$$

Theorem C.24. ⁴⁶ Let $\mathcal{B}(X, Y)$ be the space of BOUNDED LINEAR OPERATORS on normed linear spaces X and Y. Let M be a bounded linear operator in $\mathcal{B}(X, Y)$, and I the identity operator in $\mathcal{L}(X, X)$. Let Λ be the set of eigenvalues of M. Let $||x|| \triangleq \sqrt{\langle x | x \rangle}$.



[♥]PROOF:

1. Proof that $\mathbf{M}^*\mathbf{M} = \mathbf{I} \implies |||\mathbf{M}||| = 1$:

$$\|\mathbf{M}\| = \sup_{\mathbf{x} \in X} \{ \|\mathbf{M}\mathbf{x}\| \mid \|\mathbf{x}\| = 1 \}$$
 by Definition C.6 page 117
$$= \sup_{\mathbf{x} \in X} \{ \|\mathbf{x}\| \mid \|\mathbf{x}\| = 1 \}$$
 by Theorem C.23 page 133
$$= \sup_{\mathbf{x} \in X} \{ 1 \}$$

$$= 1$$

2. Proof that $|\lambda| = 1$: Let (x, λ) be an eigenvector-eigenvalue pair.

$$1 = \frac{1}{\|x\|} \|x\|$$

$$= \frac{1}{\|x\|} \|Mx\|$$
 by Theorem C.23 page 133
$$= \frac{1}{\|x\|} \|\lambda x\|$$
 by definition of λ

$$= \frac{1}{\|x\|} |\lambda| \|x\|$$
 by homogeneous property of $\|\cdot\|$

$$= |\lambda|$$

Example C.4 (One sided shift operator). ⁴⁷ Let \boldsymbol{X} be the set of all sequences with range \mathbb{W} (0, 1, 2, ...) and shift operators defined as

1.
$$\mathbf{S}_r(x_0, x_1, x_2, \dots) \triangleq (0, x_0, x_1, x_2, \dots)$$
 (right shift operator)
2. $\mathbf{S}_l(x_0, x_1, x_2, \dots) \triangleq (x_1, x_2, x_3, \dots)$ (left shift operator)

1. \mathbf{S}_r is an isometric operator. 2. $\mathbf{S}_r^* = \mathbf{S}_I$

⁴⁶ Michel and Herget (1993) page 432

⁴⁷ Michel and Herget (1993) page 441



^ℚProof:

1. Proof that $S_r^* = S_l$:

$$\langle \mathbf{S}_{r} (x_{0}, x_{1}, x_{2}, \dots) | (y_{0}, \mathbf{y}_{1}, \mathbf{y}_{2}, \dots) \rangle = \langle (0, x_{0}, x_{1}, x_{2}, \dots) | (y_{0}, \mathbf{y}_{1}, \mathbf{y}_{2}, \dots) \rangle$$

$$= \sum_{n=1}^{\infty} \mathbf{x}_{n-1} \mathbf{y}_{n}^{*}$$

$$= \sum_{n=0}^{\infty} \mathbf{x}_{n} \mathbf{y}_{n+1}^{*}$$

$$= \sum_{n=0}^{\infty} \mathbf{x}_{n} \mathbf{y}_{n+1}^{*}$$

$$= \langle (x_{0}, x_{1}, x_{2}, \dots) | (y_{1}, \mathbf{y}_{2}, \mathbf{y}_{3}, \dots) \rangle \rangle$$

$$= \langle (x_{0}, x_{1}, x_{2}, \dots) | (\mathbf{y}_{1}, \mathbf{y}_{2}, \mathbf{y}_{3}, \dots) \rangle \rangle$$

2. Proof that S_r is isometric ($S_r^*S_r = I$):

$$\mathbf{S}_r^* \mathbf{S}_r = \mathbf{S}_l \mathbf{S}_r$$

$$= \mathbf{I}$$
by 1.

C.4.5Unitary operators

Let $\mathcal{B}(X,Y)$ be the space of BOUNDED LINEAR OPERATORS on normed linear spaces X and Y. Let U be a bounded linear operator in $\mathcal{B}(X,Y)$, and I the identity operator in $\mathcal{B}(\boldsymbol{X},\boldsymbol{X}).$

The operator U is unitary if $U^*U = UU^* = I$.

Proposition C.4. Let $\mathcal{B}(X, Y)$ be the space of BOUNDED LINEAR OPERATORS on normed linear spaces **X** and **Y**. Let **U** and **V** be bounded linear operators in $\mathscr{B}(X, Y)$.

$$\left.\begin{array}{ccc}
P & U \text{ is UNITARY} & and \\
P & V \text{ is UNITARY}
\end{array}\right\} \qquad \Longrightarrow \qquad (UV) \text{ is UNITARY}.$$

[♠]Proof:

$$(\mathbf{UV})(\mathbf{UV})^* = (\mathbf{UV})(\mathbf{V}^*\mathbf{U}^*) \qquad \text{by Theorem C.8 page 121}$$

$$= \mathbf{U}(\mathbf{VV}^*)\mathbf{U}^* \qquad \text{by associative property}$$

$$= \mathbf{UIU}^* \qquad \text{by definition of } \underbrace{unitary} \text{ operators} \qquad \text{(Definition C.14 page 135)}$$

$$= \mathbf{I} \qquad \text{by definition of } \underbrace{unitary} \text{ operators} \qquad \text{(Definition C.14 page 135)}$$

$$(\mathbf{UV})^*(\mathbf{UV}) = (\mathbf{V}^*\mathbf{U}^*)(\mathbf{UV}) \qquad \text{by Theorem C.8 page 121}$$

$$= \mathbf{V}^*(\mathbf{U}^*\mathbf{U})\mathbf{V} \qquad \text{by associative property}$$

$$= \mathbf{V}^*\mathbf{IV} \qquad \text{by definition of } \underbrace{unitary} \text{ operators} \qquad \text{(Definition C.14 page 135)}$$

$$= \mathbf{I} \qquad \text{by definition of } \underbrace{unitary} \text{ operators} \qquad \text{(Definition C.14 page 135)}$$

⁴⁸ Rudin (1991) page 312, 🏿 Michel and Herget (1993) page 431, 🖨 Autonne (1901) page 209, 🖨 Autonne (1902),





 \Rightarrow

Theorem C.25. ⁴⁹ Let $\mathcal{B}(H, H)$ be the space of BOUNDED LINEAR OPERATORS on a HILBERT SPACE H. Let $\mathcal{J}(\mathbf{U})$ be the image set of \mathbf{U} .

If U is a bounded linear operator ($U \in \mathcal{B}(H, H)$), then the following conditions are equivalent:

T

- 1. $\mathbf{U}\mathbf{U}^* = \mathbf{U}^*\mathbf{U} = \mathbf{I}$
- (UNITARY)

- 2. $\langle \mathbf{U} \mathbf{x} | \mathbf{U} \mathbf{y} \rangle = \langle \mathbf{U}^* \mathbf{x} | \mathbf{U}^* \mathbf{y} \rangle = \langle \mathbf{x} | \mathbf{y} \rangle$ and $\mathbf{\mathscr{I}}(\mathbf{U}) = X$
- (SURJECTIVE)
- 3. $\|\mathbf{U}x \mathbf{U}y\| = \|\mathbf{U}^*x \mathbf{U}^*y\| = \|x y\|$ and $\mathscr{F}(\mathbf{U}) = X$ (isometric in distance)

 $4. \quad \|\mathbf{U}\mathbf{x}\| = \|\mathbf{x}\|$

- and $\mathcal{J}(\mathbf{U}) = X$ (ISOMETRIC IN LENGTH)

^ℚProof:

- 1. Proof that $(1) \implies (2)$:
 - (a) $\langle \mathbf{U} \mathbf{x} | \mathbf{U} \mathbf{y} \rangle = \langle \mathbf{U}^* \mathbf{x} | \mathbf{U}^* \mathbf{y} \rangle = \langle \mathbf{x} | \mathbf{y} \rangle$ by Theorem C.23 (page 133).
 - (b) Proof that $\mathcal{J}(\mathbf{U}) = X$:

$$X \supseteq \mathcal{F}(\mathbf{U})$$
 because $\mathbf{U} \in X^X$
 $\supseteq \mathcal{F}(\mathbf{U}\mathbf{U}^*)$
 $= \mathcal{F}(\mathbf{I})$ by left hypothesis $(\mathbf{U}^*\mathbf{U} = \mathbf{U}\mathbf{U}^* = \mathbf{I})$
 $= X$ by Definition C.3 page 112 (definition of \mathbf{I})

- 2. Proof that (2) \iff (3) \iff (4): by Theorem C.23 page 133.
- 3. Proof that (3) \implies (1):
 - (a) Proof that $||\mathbf{U}x \mathbf{U}y|| = ||x y|| \implies \mathbf{U}^*\mathbf{U} = \mathbf{I}$: by Theorem C.23 page 133
 - (b) Proof that $\|\mathbf{U}^*x \mathbf{U}^*y\| = \|x y\| \implies \mathbf{U}\mathbf{U}^* = \mathbf{I}$:

$$\|\mathbf{U}^* \mathbf{x} - \mathbf{U}^* \mathbf{y}\| = \|\mathbf{x} - \mathbf{y}\| \implies \qquad \mathbf{U}^{**} \mathbf{U}^* = \mathbf{I}$$
 by Theorem C.23 page 133 by Theorem C.13 page 126

Theorem C.26. Let $\mathcal{B}(H, H)$ be the space of BOUNDED LINEAR OPERATORS on a HILBERT SPACE H. Let U be a bounded linear operator in $\mathcal{B}(H,H)$, $\mathcal{N}(U)$ the Null Space of U, and $\mathcal{F}(U)$ the image set of U.

$$\underbrace{\mathbf{U}\mathbf{U}^* = \mathbf{U}^*\mathbf{U} = \mathbf{I}}_{\mathbf{U} \text{ is unitary}} \Longrightarrow \left\{ \begin{array}{ll} \mathbf{U}^{-1} &=& \mathbf{U}^* & \text{and} \\ \boldsymbol{\mathcal{I}}(\mathbf{U}) &=& \boldsymbol{\mathcal{I}}(\mathbf{U}^*) &=& \boldsymbol{X} & \text{and} \\ \boldsymbol{\mathcal{N}}(\mathbf{U}) &=& \boldsymbol{\mathcal{N}}(\mathbf{U}^*) &=& \{0\} & \text{and} \\ \|\mathbf{U}\| &=& \|\mathbf{U}^*\| &=& 1 & \text{(unit length)} \end{array} \right\}$$

[♠]Proof:

1. Note that U, U^* , and U^{-1} are all both *isometric* and *normal*:

⁴⁹ ■ Rudin (1991) pages 313–314 (Theorem 12.13), ■ Knapp (2005a) page 45 (Proposition 2.6)



- 2. Proof that $U^*U = UU^* = I \implies \mathcal{J}(U) = \mathcal{J}(U^*) = H$: by Theorem C.25 page 136.
- 3. Proof that $\mathbf{U}^*\mathbf{U} = \mathbf{U}\mathbf{U}^* = \mathbf{I} \implies \mathcal{N}(\mathbf{U}) = \mathcal{N}(\mathbf{U}^*) = \mathcal{N}(\mathbf{U}^{-1})$:

$$\mathcal{N}(\mathbf{U}^*) = \mathcal{N}(\mathbf{U})$$
 because \mathbf{U} and \mathbf{U}^* are both *normal* and by Theorem C.20 page 131 by Theorem C.14 page 127 by above result $= \{0\}$


4. Proof that $\mathbf{U}^*\mathbf{U} = \mathbf{U}\mathbf{U}^* = \mathbf{I} \implies \|\mathbf{U}^{-1}\| = \|\mathbf{U}^*\| = \|\mathbf{U}\| = 1$: Because U, U^* , and U^{-1} are all isometric and by Theorem C.24 page 134.


Example C.5 (Rotation matrix). ⁵⁰

$$\underbrace{\left\{ \mathbf{R}_{\theta} \triangleq \begin{bmatrix} \cos\theta & -\sin\theta \\ \sin\theta & \cos\theta \end{bmatrix} \right\}}_{\mathbf{rotation \ matrix} \ \mathbf{R}_{\theta} : \ \mathbb{R}^{2} \to \mathbb{R}^{2}} \qquad \Longrightarrow \qquad \left\{ \begin{array}{ccc} \text{(1).} & \mathbf{R}^{-1}{}_{\theta} & = & \mathbf{R}_{-\theta} & \text{ and } \\ \text{(2).} & \mathbf{R}^{*}{}_{\theta} & = & \mathbf{R}^{-1}{}_{\theta} & \text{ (R is unitary)} \end{array} \right\}$$

^ℚProof:

$$\begin{split} \mathbf{R}^* &= \mathbf{R}^H \\ &= \begin{bmatrix} \cos\theta & -\sin\theta \\ \sin\theta & \cos\theta \end{bmatrix}^H & \text{by definition of } \mathbf{R} \\ &= \begin{bmatrix} \cos\theta & \sin\theta \\ -\sin\theta & \cos\theta \end{bmatrix} & \text{by definition of } Hermetian \ transpose \ operator \ H \\ &= \begin{bmatrix} \cos(-\theta) & -\sin(-\theta) \\ \sin(-\theta) & \cos(-\theta) \end{bmatrix} & \text{(Theorem G.2 page 157)} \\ &= \mathbf{R}_{-\theta} & \text{by definition of } \mathbf{R} \\ &= \mathbf{R}^{-1} & \text{by 1}. \end{split}$$





Example C.8 (Translation operator). Let $X = L_{\mathbb{R}}^2$ and $T \in X^X$ be defined as

$$\mathbf{Tf}(x) \triangleq \mathbf{f}(x-1) \quad \forall \mathbf{f} \in L_{\mathbb{R}}^2$$
 (translation operator)

⁵¹ ☐ Gel'fand (1963) page 4, ☐ Gelfand et al. (2018) page 4



⁵⁰ ■ Noble and Daniel (1988) page 311

1.
$$\mathbf{T}^{-1} f(x) = f(x+1)$$
 $\forall f \in L^2_{\mathbb{R}}$ (inverse translation operator)

2. 3. $\mathbf{T}^*\mathbf{T} = \mathbf{T}\mathbf{T}^* = \mathbf{I}$

(T is invertible) (T is unitary)

[♠]Proof:

1. Proof that $T^{-1}f(x) = f(x + 1)$:

$$\mathbf{T}^{-1}\mathbf{T} = \mathbf{I}$$
$$\mathbf{T}\mathbf{T}^{-1} = \mathbf{I}$$

2. Proof that **T** is unitary:

$$\langle \mathbf{T}f(x) | g(x) \rangle = \langle f(x-1) | g(x) \rangle$$
 by definition of \mathbf{T}

$$= \int_{x} f(x-1)g^{*}(x) dx$$

$$= \int_{x} f(x)g^{*}(x+1) dx$$

$$= \langle f(x) | g(x+1) \rangle$$

$$= \left\langle f(x) | \underbrace{\mathbf{T}^{-1}}_{\mathbf{T}^{*}} g(x) \right\rangle$$
 by 1.

Example C.9 (Dilation operator). Let $\pmb{X} = \pmb{L}_{\mathbb{R}}^2$ and $\mathbf{T} \in \pmb{X}^{\pmb{X}}$ be defined as

$$\mathbf{Df}(x) \triangleq \sqrt{2}\mathbf{f}(2x) \quad \forall \mathbf{f} \in L_{\mathbb{R}}^2 \quad \text{(dilation operator)}$$

1.
$$\mathbf{D}^{-1} f(x) = \frac{1}{\sqrt{2}} f\left(\frac{1}{2}x\right) \quad \forall f \in L_{\mathbb{R}}^2$$
 (inverse dilation operator)
2. $\mathbf{D}^* = \mathbf{D}^{-1}$ (D is invertible)

(D is invertible)

 $\mathbf{D}^*\mathbf{D} = \mathbf{D}\mathbf{D}^* = \mathbf{I}$

(D is unitary)

1. Proof that $\mathbf{D}^{-1} f(x) = \frac{1}{\sqrt{2}} f\left(\frac{1}{2}x\right)$:

$$\mathbf{D}^{-1}\mathbf{D} = \mathbf{I}$$
$$\mathbf{D}\mathbf{D}^{-1} = \mathbf{I}$$

2. Proof that **D** is unitary:

$$\begin{split} \langle \mathbf{D} \mathbf{f}(x) \, | \, \mathbf{g}(x) \rangle &= \left\langle \sqrt{2} \mathbf{f}(2x) \, | \, \mathbf{g}(x) \right\rangle & \text{by definition of } \mathbf{D} \\ &= \int_x \sqrt{2} \mathbf{f}(2x) \mathbf{g}^*(x) \, \mathrm{d} \mathbf{x} \\ &= \int_{u \in \mathbb{R}} \sqrt{2} \mathbf{f}(u) \mathbf{g}^* \left(\frac{1}{2}u\right) \frac{1}{2} \, \mathrm{d} \mathbf{u} & \text{let } u \triangleq 2x \implies \mathrm{d} \mathbf{x} = \frac{1}{2} \, \mathrm{d} \mathbf{u} \\ &= \int_{u \in \mathbb{R}} \mathbf{f}(u) \left[\frac{1}{\sqrt{2}} \mathbf{g} \left(\frac{1}{2}u\right) \right]^* \, \mathrm{d} \mathbf{u} \\ &= \left\langle \mathbf{f}(x) \, | \, \frac{1}{\sqrt{2}} \mathbf{g} \left(\frac{1}{2}x\right) \right\rangle \\ &= \left\langle \mathbf{f}(x) \, | \, \frac{\mathbf{D}^{-1}}{\mathbf{D}^*} \mathbf{g}(x) \right\rangle & \text{by } 1. \end{split}$$

[♠]Proof:

₽

Example C.10 (Delay operator). Let X be the set of all sequences and $D \in X^X$ be a delay operator.

The delay operator $\mathbf{D}(x_n)_{n\in\mathbb{Z}} \triangleq (x_{n-1})_{n\in\mathbb{Z}}$ is unitary.

 \triangle Proof: The inverse \mathbf{D}^{-1} of the delay operator \mathbf{D} is

$$\mathbf{D}^{-1} \left(x_n \right)_{n \in \mathbb{Z}} \triangleq \left(x_{n+1} \right)_{n \in \mathbb{Z}}.$$

$$\langle \mathbf{D} ((x_n)) | ((y_n)) \rangle = \langle ((x_{n-1})) | ((y_n)) \rangle$$
 by definition of \mathbf{D}

$$= \sum_{n} x_{n-1} y_n^*$$

$$= \sum_{n} x_n y_{n+1}^*$$

$$= \langle ((x_n)) | ((y_{n+1})) \rangle$$

$$= \langle ((x_n)) | ((y_n)) \rangle$$

Therefore, $\mathbf{D}^* = \mathbf{D}^{-1}$. This implies that $\mathbf{D}\mathbf{D}^* = \mathbf{D}^*\mathbf{D} = \mathbf{I}$ which implies that \mathbf{D} is unitary.

Example C.11 (Fourier transform). Let $\tilde{\mathbf{F}}$ be the *Fourier Transform* and $\tilde{\mathbf{F}}^{-1}$ the *inverse Fourier Transform* operator (Theorem N.1 page 248)

$$[\tilde{\mathbf{F}}\mathbf{x}](f) \triangleq \int_{t} \mathbf{x}(t) \underbrace{e^{-i2\pi f t}}_{\kappa(t,f)} d\mathbf{t} \qquad \qquad \left[\tilde{\mathbf{F}}^{-1}\tilde{\mathbf{x}}\right](t) \triangleq \int_{f} \tilde{\mathbf{x}}(f) \underbrace{e^{i2\pi f t}}_{\kappa^{*}(t,f)} d\mathbf{f}.$$

 $\tilde{\mathbf{F}}^* = \tilde{\mathbf{F}}^{-1}$ (the Fourier Transform operator $\tilde{\mathbf{F}}$ is unitary)

№PROOF:

$$\begin{split} \left\langle \tilde{\mathbf{F}} \mathbf{x} \mid \tilde{\mathbf{y}} \right\rangle &= \left\langle \int_{t} \mathbf{x}(t) e^{-i2\pi f t} \, \mathrm{d} \mathbf{t} \mid \tilde{\mathbf{y}}(f) \right\rangle \\ &= \int_{t} \mathbf{x}(t) \left\langle e^{-i2\pi f t} \mid \tilde{\mathbf{y}}(f) \right\rangle \, \mathrm{d} \mathbf{t} \\ &= \int_{t} \mathbf{x}(t) \int_{f} e^{-i2\pi f t} \tilde{\mathbf{y}}^{*}(f) \, \mathrm{d} \mathbf{f} \, \mathrm{d} \mathbf{t} \\ &= \int_{t} \mathbf{x}(t) \left[\int_{f} e^{i2\pi f t} \tilde{\mathbf{y}}(f) \, \mathrm{d} \mathbf{f} \right]^{*} \, \mathrm{d} \mathbf{t} \\ &= \left\langle \mathbf{x}(t) \mid \int_{f} \tilde{\mathbf{y}}(f) e^{i2\pi f t} \, \mathrm{d} \mathbf{f} \right\rangle \\ &= \left\langle \mathbf{x} \mid \underbrace{\tilde{\mathbf{F}}^{-1}}_{\tilde{\mathbf{F}}^{*}} \tilde{\mathbf{y}} \right\rangle \end{split}$$

This implies that $\tilde{\mathbf{F}}$ is unitary ($\tilde{\mathbf{F}}^* = \tilde{\mathbf{F}}^{-1}$).



C.5 Operator order

Definition C.15. ⁵² Let $P \in Y^X$ be an operator.

Daniel J. Greenhoe

P is positive if $\langle \mathbf{P}x \mid x \rangle \geq 0 \ \forall x \in \mathbf{X}$. This condition is denoted $\mathbf{P} \geq 0$.

Theorem C.27. 53

01EIII 6.27.						
		(P+Q)	≥	0		$((\mathbf{P} + \mathbf{Q}) \text{ is positive})$
$\mathbf{P} \ge 0$ and $\mathbf{Q} \ge 0$	\Longrightarrow	A^*PA	\geq	0	$\forall \mathbf{A} \in \mathcal{B}(\mathbf{X}, \mathbf{X})$	(A*PA is positive)
P and Q are both positive		A^*A	\geq	0	$\forall \mathbf{A} \in \mathcal{B}(\mathbf{X}, \mathbf{X})$	(A*A is positive)
	$\mathbf{P} \ge 0 \text{ and } \mathbf{Q} \ge 0$	$\underbrace{\mathbf{P} \ge 0 \ and \ \mathbf{Q} \ge 0} \qquad \Longrightarrow \qquad$	$\underbrace{\mathbf{P} \ge 0 \text{ and } \mathbf{Q} \ge 0}_{\mathbf{A}^* \mathbf{P} \mathbf{A}} \implies \begin{cases} (\mathbf{P} + \mathbf{Q}) \\ \mathbf{A}^* \mathbf{P} \mathbf{A} \\ \mathbf{A}^* \mathbf{A} \end{cases}$	$\underbrace{\mathbf{P} \geq 0 \text{ and } \mathbf{Q} \geq 0}_{\mathbf{A}^* \mathbf{A}} \implies \begin{cases} (\mathbf{P} + \mathbf{Q}) \geq \\ \mathbf{A}^* \mathbf{P} \mathbf{A} \geq \\ \mathbf{A}^* \mathbf{A} > \end{cases}$	$\underbrace{\mathbf{P} \ge 0 \text{ and } \mathbf{Q} \ge 0}_{\mathbf{A}^* \mathbf{A}} \implies \begin{cases} (\mathbf{P} + \mathbf{Q}) \ge 0 \\ \mathbf{A}^* \mathbf{P} \mathbf{A} \ge 0 \\ \mathbf{A}^* \mathbf{A} > 0 \end{cases}$	$ \underbrace{\mathbf{P} \ge 0 \text{ and } \mathbf{Q} \ge 0}_{\mathbf{A}^* \mathbf{P} \mathbf{A}} \implies \begin{cases} (\mathbf{P} + \mathbf{Q}) \ge 0 \\ \mathbf{A}^* \mathbf{P} \mathbf{A} \ge 0 \\ \mathbf{A}^* \mathbf{A} > 0 \end{cases} \forall \mathbf{A} \in \mathcal{B}(\mathbf{X}, \mathbf{X})$

№ Proof:

$$\langle (\mathbf{P} + \mathbf{Q}) \boldsymbol{x} \mid \boldsymbol{x} \rangle = \langle \mathbf{P} \boldsymbol{x} \mid \boldsymbol{x} \rangle + \langle \mathbf{Q} \boldsymbol{x} \mid \boldsymbol{x} \rangle \qquad \text{by additive property of } \langle \triangle \mid \nabla \rangle \text{ (Definition C.9 page 124)}$$

$$\geq \langle \mathbf{P} \boldsymbol{x} \mid \boldsymbol{x} \rangle \qquad \text{by left hypothesis}$$

$$\geq 0 \qquad \text{by left hypothesis}$$

$$\langle \mathbf{A}^* \mathbf{P} \mathbf{A} \boldsymbol{x} \mid \boldsymbol{x} \rangle = \langle \mathbf{P} \mathbf{A} \boldsymbol{x} \mid \mathbf{A} \boldsymbol{x} \rangle \qquad \text{by definition of adjoint (Proposition C.3 page 125)}$$

$$= \langle \mathbf{P} \boldsymbol{y} \mid \boldsymbol{y} \rangle \qquad \text{where } \boldsymbol{y} \triangleq \mathbf{A} \boldsymbol{x}$$

$$\geq 0 \qquad \text{by left hypothesis}$$

$$\langle \mathbf{I} \boldsymbol{x} \mid \boldsymbol{x} \rangle = \langle \boldsymbol{x} \mid \boldsymbol{x} \rangle \qquad \text{by definition of I (Definition C.3 page 112)}$$


$$\geq 0 \qquad \text{by non-negative property of } \langle \triangle \mid \nabla \rangle \text{ (Definition C.9 page 124)}$$

$$\Rightarrow \mathbf{I} \text{ is positive}$$

$$\langle \mathbf{A}^* \mathbf{A} \boldsymbol{x} \mid \boldsymbol{x} \rangle = \langle \mathbf{A}^* \mathbf{I} \mathbf{A} \boldsymbol{x} \mid \boldsymbol{x} \rangle \qquad \text{by definition of I (Definition C.3 page 112)}$$

$$\geq 0 \qquad \text{by two previous results}$$

Definition C.16. ⁵⁴ *Let* $A, B \in \mathcal{B}(X, Y)$ *be* BOUNDED *operators*.



⁵⁴ Michel and Herget (1993) page 429



⁵² Michel and Herget (1993) page 429 (Definition 7.4.12)

⁵³ Michel and Herget (1993) page 429

Definition D.1. Let \mathbb{R} be the set of real numbers, \mathscr{B} the set of Borel sets on \mathbb{R} , and μ the standard Borel measure on \mathscr{B} . Let $\mathbb{R}^{\mathbb{R}}$ be as in Definition 3.1 page 39.

The space of Lebesgue square-integrable functions $L^2_{(\mathbb{R},\mathscr{B},\mu)}$ (or $L^2_{\mathbb{R}}$) is defined as

$$\mathbf{L}_{\mathbb{R}}^{2} \triangleq \mathbf{L}_{(\mathbb{R},\mathcal{B},\mu)}^{2} \triangleq \left\{ f \in \mathbb{R}^{\mathbb{R}} | \left(\int_{\mathbb{R}} |f|^{2} \right)^{\frac{1}{2}} d\mu < \infty \right\}.$$

The **standard inner product** $\langle \triangle \mid \nabla \rangle$ on $L^2_{\mathbb{R}}$ is defined as

$$\langle f(x) | g(x) \rangle \triangleq \int_{\mathbb{D}} f(x)g^*(x) dx.$$

The **standard norm** $\|\cdot\|$ on $L^2_{\mathbb{R}}$ is defined as $\|f(x)\| \triangleq \langle f(x) | f(x) \rangle^{\frac{1}{2}}$

Definition D.2. *Let* f(x) *be a* FUNCTION *in* $\mathbb{R}^{\mathbb{R}}$.

$$\frac{\mathsf{d}}{\mathsf{d}\mathsf{x}}\mathsf{f}(x) \triangleq \mathsf{f}'(x) \triangleq \lim_{\varepsilon \to 0} \frac{\mathsf{f}(x+\varepsilon) - \mathsf{f}(x)}{\varepsilon}$$

Proposition D.1.

$$\begin{bmatrix}
(1). & f(x) \text{ is Continuous} & and \\
(2). & f(a+x) = f(a-x) \\
\text{SYMMETRIC about a point a}
\end{bmatrix}
\implies
\begin{cases}
(1). & f'(a+x) = -f'(a-x) \\
(2). & f'(a) = 0
\end{cases}$$
(ANTI-SYMMETRIC about a)

♥Proof:

DEF

$$f'(a+x) = \lim_{\varepsilon \to 0} \frac{1}{\varepsilon} [f(a+x+\varepsilon) - f(a+x-\varepsilon)]$$

$$= \lim_{\varepsilon \to 0} \frac{1}{\varepsilon} [f(a-x-\varepsilon) - f(a-x+\varepsilon)]$$
by hpothesis (2)
$$= -\lim_{\varepsilon \to 0} \frac{1}{\varepsilon} [f(a-x+\varepsilon) - f(a-x-\varepsilon)]$$

$$= -f(a-x)$$

$$f'(a) = \frac{1}{\varepsilon} f'(a+0) + \frac{1}{\varepsilon} f'(a-0)$$

$$f'(a) = \frac{1}{2}f'(a+0) + \frac{1}{2}f'(a-0)$$
$$= \frac{1}{2}[f'(a+0) - f'(a+0)]$$

by previous result

₽

= 0

Lemma D.1.

$$f(x)$$
 is invertible $\Longrightarrow \left\{ \frac{d}{dy} f^{-1}(y) = \frac{1}{\frac{d}{dx} f[f^{-1}(y)]} \right\}$

^ℚProof:

$$\frac{d}{dy} f^{-1}(y) \triangleq \lim_{\varepsilon \to 0} \frac{f^{-1}(y + \varepsilon) - f^{-1}(y)}{\varepsilon} \qquad \text{by definition of } \frac{d}{dy}$$

$$= \lim_{\delta \to 0} \frac{1}{\left[\frac{f(x + \delta) - f(x)}{\delta}\right]} \Big|_{x \triangleq f^{-1}(y)} \qquad \text{because in the limit, } \frac{\Delta y}{\Delta x} = \left(\frac{\Delta x}{\Delta y}\right)^{-1}$$

$$\triangleq \frac{1}{\frac{d}{dx} f(x)} \Big|_{x \triangleq f^{-1}(y)} \qquad \text{by definition of } \frac{d}{dx}$$

$$= \frac{1}{\frac{d}{dx} f\left[f^{-1}(y)\right]} \qquad \text{because } x \triangleq f^{-1}(y)$$

by definition of $\frac{d}{dv}$

by definition of $\frac{d}{dx}$

(Definition D.2 page 141)

(Definition D.2 page 141)

because $x \triangleq f^{-1}(v)$

Theorem D.1. Let f be a continuous function in
$$L^2_{\mathbb{R}}$$
 and $f^{(n)}$ the nth derivative of f.
$$\int_{[0:1)^n} f^{(n)} \left(\sum_{k=1}^n x_k \right) \mathrm{d} \mathsf{x}_1 \ \mathrm{d} \mathsf{x}_2 \cdots \ \mathrm{d} \mathsf{x}_n = \sum_{k=0}^n (-1)^{n-k} \binom{n}{k} f(k) \qquad \forall n \in \mathbb{N}$$

[♠]Proof: Proof by induction:

1. Base case ...proof for n = 1 case:

$$\int_{[0:1)} f^{(1)}(x) dx = f(1) - f(0)$$

$$= (-1)^{1+1} {1 \choose 1} f(1) + (-1)^{1+0} {1 \choose 0} f(0)$$

$$= \sum_{k=0}^{1} (-1)^{n-k} {n \choose k} f(k)$$

by Fundamental theorem of calculus

¹ Chui (1992) page 86 ⟨item (ii)⟩, Prasad and Iyengar (1997) pages 145–146 ⟨Theorem 6.2 (b)⟩

2. Induction step ...proof that n case $\implies n+1$ case:

$$\begin{split} &\int_{[0:1)^{n+1}} \mathsf{f}^{(n+1)} \Biggl(\sum_{k=1}^{n+1} x_k \Biggr) \, \mathrm{d} \mathbf{x}_1 \, \mathrm{d} \mathbf{x}_2 \cdots \, \mathrm{d} \mathbf{x}_{n+1} \\ &= \int_{[0:1)^n} \Biggl[\int_0^1 \mathsf{f}^{(n+1)} \Biggl(x_{n+1} + \sum_{k=1}^n x_k \Biggr) \, \mathrm{d} \mathbf{x}_{n+1} \Biggr] \, \mathrm{d} \mathbf{x}_1 \, \mathrm{d} \mathbf{x}_2 \cdots \, \mathrm{d} \mathbf{x}_n \\ &= \int_{[0:1)^n} \Biggl[\mathsf{f}^{(n)} \Biggl(x_{n+1} + \sum_{k=1}^n x_k \Biggr) \Biggr|_{x_{n+1}=0}^{|x_{n+1}=0|} \mathrm{d} \mathbf{x}_1 \, \mathrm{d} \mathbf{x}_2 \cdots \, \mathrm{d} \mathbf{x}_n \quad \text{by } \textit{Fundamental theorem of calculus} \\ &= \int_{[0:1)^n} \Biggl[\mathsf{f}^{(n)} \Biggl(1 + \sum_{k=1}^n x_k \Biggr) - \mathsf{f}^{(n)} \Biggl(0 + \sum_{k=1}^n x_k \Biggr) \Biggr] \, \mathrm{d} \mathbf{x}_1 \, \mathrm{d} \mathbf{x}_2 \cdots \, \mathrm{d} \mathbf{x}_n \\ &= \sum_{k=0}^n (-1)^{n-k} \binom{n}{k} \mathsf{f}(k+1) - \sum_{k=0}^n (-1)^{n-k} \binom{n}{k} \mathsf{f}(k) \quad \text{by induction hypothesis} \\ &= \sum_{m=1}^{m=n+1} (-1)^{n-m+1} \binom{n}{m-1} \mathsf{f}(m) + \sum_{k=0}^n (-1)^{n-k} \binom{n}{k} \mathsf{f}(k) \quad \text{where } m \triangleq k+1 \implies k = m-1 \\ &= \Biggl[\mathsf{f}(n+1) + \sum_{k=1}^n (-1)^{n-k+1} \binom{n}{k-1} \mathsf{f}(k) \Biggr] + \Biggl[(-1)^{n+1} \mathsf{f}(0) + \sum_{k=1}^n (-1)^{n-k+1} \binom{n}{k} \mathsf{f}(k) \Biggr] \\ &= \mathsf{f}(n+1) + (-1)^{n+1} \mathsf{f}(0) + \sum_{k=1}^n (-1)^{n-k+1} \Biggl[\binom{n}{k-1} + \binom{n}{k} \Biggr] \mathsf{f}(k) \\ &= (-1)^0 \binom{n+1}{n+1} \mathsf{f}(n+1) + (-1)^{n+1} \binom{n+1}{0} \mathsf{f}(0) + \sum_{k=1}^n (-1)^{n-k+1} \binom{n+1}{k} \mathsf{f}(k) \end{aligned} \quad \text{by } \textit{Stifel formula} \\ &= \sum_{k=0}^{n+1} (-1)^{n-k+1} \binom{n+1}{k} \mathsf{f}(k)$$

Some proofs invoke differentiation multiple times. This is simplified thanks to the *Leibniz rule*, also called the *generalized product rule* (*GPR*, next lemma). The Leibniz rule is remarkably similar in form to the *binomial theorem*.

Lemma D.2 (Leibniz rule / generalized product rule). 2 Let f(x), $g(x) \in L^2_{\mathbb{R}}$ with derivatives $f^{(n)}(x) \triangleq \frac{d^n}{dx^n} f(x)$ and $g^{(n)}(x) \triangleq \frac{d^n}{dx^n} g(x)$ for n = 0, 1, 2, ..., and $\binom{n}{k} \triangleq \frac{n!}{(n-k)!k!}$ (binomial coefficient). Then

$$\frac{\mathsf{L}}{\mathsf{M}} \quad \frac{\mathrm{d}^n}{\mathrm{d}x^n} [\mathsf{f}(x)\mathsf{g}(x)] = \sum_{k=0}^n \binom{n}{k} \mathsf{f}^{(k)}(x) \mathsf{g}^{(n-k)}(x)$$

Example D.1.

$$\frac{\mathsf{E}}{\mathsf{X}} \frac{\mathrm{d}^3}{\mathrm{d}x^3} \big[\mathsf{f}(x)\mathsf{g}(x) \big] = \mathsf{f}'''(x)\mathsf{g}(x) + 3\mathsf{f}''(x)\mathsf{g}'(x) + 3\mathsf{f}'(x)\mathsf{g}''(x) + \mathsf{f}(x)\mathsf{g}'''(x)$$

Theorem D.2 (Leibniz integration rule). ³

³ Flanders (1973) page 615 ⟨(1.1)⟩ Talvila (2001), Knapp (2005b) page 389 ⟨Chapter VII⟩, Protter and Morrey (2012) page 422 ⟨Leibniz Rule. Theorem 1.⟩, http://planetmath.org/encyclopedia/DifferentiationUnderIntegralSign.html





² ■ Ben-Israel and Gilbert (2002) page 154, ■ Leibniz (1710)



 $\frac{d}{dx} \int_{a(x)}^{b(x)} g(t) dt = g[b(x)]b'(x) - g[a(x)]a'(x)$

APPENDIX E	
I	
	CONVOLUTION

E.1 Definition

```
Definition E.1.

The convolution operation is defined as
                    [f \star g](x) \triangleq f(x) \star g(x) \triangleq \int_{u \in \mathbb{R}} f(u)g(x - u) du \qquad \forall f, g \in L^2_{(\mathbb{R}, \mathcal{B}, \mu)}
```

Properties E.2

```
Theorem E.1. ^2

f \star g = g \star f \qquad \text{(COMMUTATIVE)}
f \star (g \star h) = (g \star g) \star h
(\alpha f) \star g = \alpha (f \star g) = f \star (\alpha g) \quad \forall \alpha \in \mathbb{C}
                                                                                                                               (ASSOCIATEVE)
             f \star (g + h) = (f \star g) + (f \star h)
                                                                                                                               (DISTRIBUTIVE)
```

¹ Bachman et al. (2002) page 268 ⟨Definition 5.2.1, but with $1/2\pi$ scaling factor⟩,
Bachman (1964) page 6, Bracewell (1978) page 224 ⟨Table 11.1 Theorems for the Laplace Transform⟩
 Bachman et al. (2002) pages 268–270



APPENDIX F	
I	
	NORMED ALGEBRAS

F.1 Algebras

All *linear space*s are equipped with an operation by which vectors in the spaces can be added together. Linear spaces also have an operation that allows a scalar and a vector to be "multiplied" together. But linear spaces in general have no operation that allows two vectors to be multiplied together. A linear space together with such an operator is an **algebra**. ¹

There are many many possible algebras—many more than one can shake a stick at, as indicated by Michiel Hazewinkel in his book, *Handbook of Algebras*: "Algebra, as we know it today (2005), consists of many different ideas, concepts and results. A reasonable estimate of the number of these different items would be somewhere between 50,000 and 200,000. Many of these have been named and many more could (and perhaps should) have a "name" or other convenient designation."²

```
Definition F.1. <sup>3</sup> Let \mathbf{A} be an ALGEBRA.

An algebra \mathbf{A} is unital if \exists u \in \mathbf{A} such that ux = xu = x \forall x \in \mathbf{A}
```

Definition F.2. ⁴ Let **A** be an UNITAL ALGEBRA (Definition F.1 page 147) with unit e.

```
The spectrum of x \in \mathbf{A} is \sigma(x) \triangleq \left\{\lambda \in \mathbb{C} | \lambda e - x \text{ is not invertible} \right\}.

The resolvent of x \in \mathbf{A} is \rho_x(\lambda) \triangleq (\lambda e - x)^{-1} \forall \lambda \notin \sigma(x).

The spectral radius of x \in \mathbf{A} is r(x) \triangleq \sup\{|\lambda| | \lambda \in \sigma(x)\}.
```

¹ Fuchs (1995) page 2

² Hazewinkel (2000) page v

³ Folland (1995) page 1

⁴ Folland (1995) pages 3–4

D E F

Star-Algebras **F.2**

Definition F.3. 5 Let A be an ALGEBRA.

The pair(A, *) is a *-algebra, or star-algebra, if 1. $(x + y)^* = x^* + y^* \quad \forall x, y \in A$ (DISTRIBUTIVE) and $(\alpha x)^*$ $= \bar{\alpha}x^*$ $\forall x \in A, \alpha \in \mathbb{C}$ (CONJUGATE LINEAR) and $= v^*x^*$ 3. $\forall x, y \in \mathbf{A}$ (ANTIAUTOMORPHIC) and $\forall x \in A$ (INVOLUTORY) The operator * is called an **involution** on the algebra A.

Proposition F.1. 6 *Let* (A, *) *be an* UNITAL *-ALGEBRA.

	F	(,)				
P	• •		(1. x	* <i>is</i> invertible	∀ <i>x</i> ∈ A	and
R P	x is invertible	\Rightarrow	2. ($(x^*)^{-1} = (x^{-1})^*$	∀ <i>x</i> ∈ A	

 $^{\lozenge}$ Proof: Let *e* be the unit element of (A, *).

1. Proof that $e^* = e$:

$$x e^* = (x e^*)^{**}$$
 by $involutory$ property of * (Definition F.3 page 148)
 $= (x^* e^{**})^*$ by $antiautomorphic$ property of * (Definition F.3 page 148)
 $= (x^* e)^*$ by $involutory$ property of * (Definition F.3 page 148)
 $= (x^*)^*$ by definition of e
 $= x$ by $involutory$ property of * (Definition F.3 page 148)
 $e^* x = (e^* x)^{**}$ by $involutory$ property of * (Definition F.3 page 148)
 $= (e^{**} x^*)^*$ by $antiautomorphic$ property of * (Definition F.3 page 148)
 $= (e^* x^*)^*$ by $involutory$ property of * (Definition F.3 page 148)
 $= (x^*)^*$ by definition of e
 $= x$ by $involutory$ property of * (Definition F.3 page 148)

2. Proof that $(x^*)^{-1} = (x^{-1})^*$:

```
(x^{-1})^*(x^*) = [x(x^{-1})]^*
                            by antiautomorphic and involution properties of * (Definition F.3 page 148)
                            by item (1) page 148
(x^*)(x^{-1})^* = [x^{-1}x]^*
                            by antiautomorphic and involution properties of * (Definition F.3 page 148)
                            by item (1) page 148
             = e
```

- **Definition F.4.** ⁷ Let $(A, \|\cdot\|)$ be a *-ALGEBRA (Definition F.3 page 148). An element $x \in A$ is **hermitian** or **self-adjoint** if $x^* = x$.
 - \blacktriangleleft An element $x \in \mathbf{A}$ is **normal** if $xx^* = x^*x$.
 - \blacktriangleleft An element $x \in \mathbf{A}$ is a **projection** if xx = x (involutory) and $x^* = x$ (hermitian).
 - ⁵ Rickart (1960) page 178, 🗈 Gelfand and Naimark (1964), page 241
 - ⁶ Folland (1995) page 5
 - ⁷ Rickart (1960) page 178, 🖰 Gelfand and Naimark (1964), page 242



D E

Theorem F.1. 8 Let $(A, \|\cdot\|)$ be a *-ALGEBRA (Definition F.3 page 148).

 $\begin{array}{c}
\mathsf{T} \\
\mathsf{H} \\
\mathsf{M}
\end{array}
\qquad x = x^* \ and \ y = y^* \\
x \ and \ y \ are \ \mathsf{HERMITIAN}
\end{array}
\implies \begin{cases}
x + y = (x + y)^* & (x + y \ is \ self \ adjoint) \\
x^* = (x^*)^* & (x^* \ is \ self \ adjoint) \\
xy = (xy)^* \iff xy = yx \\
(xy) \ is \ \mathsf{HERMITIAN}
\end{cases}$

№ Proof:

$$(x + y)^* = x^* + y^*$$
 by *distributive* property of * (Definition F.3 page 148)
= $x + y$ by left hypothesis

$$(x^*)^* = x$$
 by *involutory* property of * (Definition F.3 page 148)

Proof that $xy = (xy)^* \implies xy = yx$

$$xy = (xy)^*$$
 by left hypothesis
 $= y^*x^*$ by *antiautomorphic* property of * (Definition F.3 page 148)
 $= yx$ by left hypothesis

Proof that $xy = (xy)^* \iff xy = yx$

$$(xy)^* = (yx)^*$$
 by left hypothesis
 $= x^*y^*$ by antiautomorphic property of * (Definition F.3 page 148)
 $= xy$ by left hypothesis

Definition F.5 (Hermitian components). 9 Let $(A, \|\cdot\|)$ be a *-ALGEBRA (Definition F.3 page 148).

The **real part** of x is defined as $\mathbf{R}_{e}x \triangleq \frac{1}{2}(x+x^{*})$ The **imaginary part** of x is defined as $\mathbf{I}_{m}x \triangleq \frac{1}{2i}(x-x^{*})$

Theorem F.2. 10 Let (A, *) be a *-ALGEBRA (Definition F.3 page 148).

I	$\mathbf{R}_{e}x$	=	$(\mathbf{R}_{e}x)^*$	∀ <i>x</i> ∈ A	$(\mathbf{R}_{e}x\ is\ HERMITIAN)$
M	$\mathbf{I}_{m}x$	=	$(\mathbf{I}_{m}x)^*$	$\forall x \in A$	$(\mathbf{I}_{m}x\ is\ HERMITIAN)$

NPROOF:

D

$$\begin{split} \left(\mathbf{R}_{\mathrm{e}}x\right)^{*} &= \left(\frac{1}{2}\left(x+x^{*}\right)\right)^{*} & \text{by definition of } \mathfrak{R} \\ &= \frac{1}{2}\left(x^{*}+x^{**}\right) & \text{by } \textit{distributive} \text{ property of } * & \text{(Definition F.3 page 148)} \\ &= \frac{1}{2}\left(x^{*}+x\right) & \text{by } \textit{involutory} \text{ property of } * & \text{(Definition F.3 page 148)} \\ &= \mathbf{R}_{\mathrm{e}}x & \text{by definition of } \mathfrak{R} & \text{(Definition F.5 page 149)} \\ &\left(\mathbf{I}_{\mathrm{m}}x\right)^{*} &= \left(\frac{1}{2i}\left(x-x^{*}\right)\right)^{*} & \text{by definition of } \mathfrak{F}. \end{split}$$



⁸ Michel and Herget (1993) page 429

⁹ Michel and Herget (1993) page 430, Rickart (1960) page 179, Gelfand and Naimark (1964), page 242

¹⁰ ■ Michel and Herget (1993) page 430, ■ Halmos (1998a) page 42

$$= \frac{1}{2i}(x^* - x^{**})$$
 by *distributive* property of * (Definition F.3 page 148)
$$= \frac{1}{2i}(x^* - x)$$
 by *involutory* property of * (Definition F.3 page 148)
$$= \mathbf{I}_m x$$
 by definition of \mathfrak{F} (Definition F.5 page 149)

Theorem F.3 (Hermitian representation). 11 Let (A, *) be a *-ALGEBRA (Definition F.3 page 148).



$$a = x + iy$$

$$\iff$$

$$x = \mathbf{R}_{\mathbf{e}}a$$

$$a = x + iy$$
 \iff $x = \mathbf{R}_{e}a$ and $y = \mathbf{I}_{m}a$

[♠]Proof:

! Proof that $a = x + iy \implies x = \mathbf{R}_e a$ and $y = \mathbf{I}_m a$:

! Proof that $a = x + iy \iff x = \mathbf{R}_e a$ and $y = \mathbf{I}_m a$:

$$x + iy = \mathbf{R}_{e}a + i\mathbf{I}_{m}a$$
 by right hypothesis
$$= \underbrace{\frac{1}{2}(a + a^{*})}_{\mathbf{R}_{e}a} + i\underbrace{\frac{1}{2i}(a - a^{*})}_{\mathbf{I}_{m}a}$$
 by definition of \Re and \Im (Definition F.5 page 149)
$$= \left(\frac{1}{2}a + \frac{1}{2}a\right) + \left(\frac{1}{2}a^{*} - \frac{1}{2}a^{*}\right)^{-0}$$

$$= a$$

¹¹ ■ Michel and Herget (1993) page 430,
Rickart (1960) page 179, Gelfand and Neumark (1943b) page 7



F.3. NORMED ALGEBRAS Daniel J. Greenhoe page 151

Normed Algebras F.3

Definition F.6. 12 Let A be an algebra.

The pair $(\mathbf{A}, \|\cdot\|)$ is a **normed algebra** if E $\forall x, y \in \mathbf{A}$ $||xy|| \le ||x|| \, ||y||$

(multiplicative condition)

A normed algebra $(\mathbf{A}, \|\cdot\|)$ is a **Banach algebra** if $(\mathbf{A}, \|\cdot\|)$ is also a Banach space.

Proposition F.2.



 $(A, \|\cdot\|)$ is a normed algebra

multiplication is continuous $in(A, ||\cdot||)$

^ℚProof:

- 1. Define $f(x) \triangleq zx$. That is, the function f represents multiplication of x times some arbitrary value z.
- 2. Let $\delta \triangleq ||x y||$ and $\epsilon \triangleq ||f(x) f(y)||$.
- 3. To prove that multiplication (f) is *continuous* with respect to the metric generated by $\|\cdot\|$, we have to show that we can always make ϵ arbitrarily small for some $\delta > 0$.
- 4. And here is the proof that multiplication is indeed continuous in $(A, \|\cdot\|)$:

```
\|f(x) - f(y)\| \triangleq \|zx - zy\|
                                             by definition of f
                                                                                                   (item (1) page 151)
                 = \|z(x - y)\|
                 \leq ||z|| ||x - y||
                                             by definition of normed algebra
                                                                                                   (Definition F.6 page 151)
                 \triangleq ||z|| \delta
                                             by definition of \delta
                                                                                                   (item (2) page 151)
                                             for some value of \delta > 0
                  <\epsilon
```

Theorem F.4 (Gelfand-Mazur Theorem). ¹³ Let \mathbb{C} be the field of complex numbers.

 $(\mathbf{A}, \|\cdot\|)$ is a Banach algebra every nonzero $x \in \mathbf{A}$ is invertible

 $A \equiv \mathbb{C}$ (A is isomorphic to \mathbb{C})

C* Algebras **F.4**

Definition F.7. ¹⁴

```
The triple (\mathbf{A}, \|\cdot\|, *) is a C^* algebra if
                                     is a Banach algebra and
             1. (A, \|\cdot\|)
D
E
             2. (A,*)
                                     is a *-algebra
                                                                   and
             3. ||x^*x|| = ||x||^2 \quad \forall x \in \mathbf{A}
     AC^* algebra (A, \|\cdot\|, *) is also called a C star algebra.
```

¹² ■ Rickart (1960) page 2, ■ Berberian (1961) page 103 (Theorem IV.9.2)

¹⁴ Folland (1995) page 1, 🖰 Gelfand and Naimark (1964), page 241, 📃 Gelfand and Neumark (1943a), 🖰 Gelfand and Neumark (1943b)





¹³ Folland (1995) page 4, 🗓 Mazur (1938) ((statement)), 🖫 Gelfand (1941) ((proof))

Theorem F.5. 15 Let **A** be an algebra.



 $(A, \|\cdot\|, *)$ is $a C^*$ algebra

 \Longrightarrow

$$||x^*|| = ||x||$$

№ Proof:

$$\|x\| = \frac{1}{\|x\|} \|x\|^2$$

$$= \frac{1}{\|x\|} \|x^*x\| \qquad \text{by definition of } C^* \text{-} algebra \qquad \text{(Definition F.7 page 151)}$$

$$\leq \frac{1}{\|x\|} \|x^*\| \|x\| \qquad \text{by definition of } normed \ algebra \qquad \text{(Definition F.6 page 151)}$$

$$= \|x^*\|$$

$$\|x^*\| \leq \|x^{**}\| \qquad \text{by previous result}$$

$$= \|x\| \qquad \text{by } involution \text{ property of } * \qquad \text{(Definition F.3 page 148)}$$

 \Rightarrow

 $^{^{15}}$ $\slash\hspace{-0.6em}$ Folland (1995) page 1, $\slash\hspace{-0.6em}$ Gelfand and Neumark (1943b) page 4, $\slash\hspace{-0.6em}$ Gelfand and Neumark (1943a)

Definition Candidates G.1

There are several ways of defining the sine and cosine functions, including the following: ¹

1. **Planar geometry:** Trigonometric functions are traditionally introduced as they have come to us historically—that is, as related to the parameters of triangles.²

$$\cos x \triangleq \frac{x}{r}$$
$$\sin x \triangleq \frac{y}{r}$$

2. Complex exponential: The cosine and sine functions are the real and imaginary parts of the complex exponential such that³

$$\cos x \triangleq \mathbf{R}_{e} e^{ix} \qquad \sin x \triangleq \mathbf{I}_{m} (e^{ix})$$

3. **Polynomial:** Let $\sum_{n=0}^{\infty} x_n \triangleq \lim_{N \to \infty} \sum_{n=0}^{N} x_n$ in some topological space. The sine and cosine functions

can be defined in terms of Taylor expansions such that⁴

$$\cos(x) \triangleq \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n)!} x^{2n} = 1 - \frac{x^2}{2} + \frac{x^4}{4!} - \frac{x^6}{6!} + \cdots$$

$$\sin(x) \triangleq \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n)!} x^{2n+1} - x - \frac{x^3}{4!} + \frac{x^5}{4!} - \frac{x^7}{4!} + \cdots$$

 $[\]sin(x) \triangleq \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n+1)!} x^{2n+1} = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + \cdots$

¹The term *sine* originally came from the Hindu word *jiva* and later adapted to the Arabic word *jiba*. Abrabic-Latin translator Robert of Chester apparently confused this word with the Arabic word jaib, which means "bay" or "inlet" thus resulting in the Latin translation sinus, which also means "bay" or "inlet". Reference: Boyer and Merzbach (1991) page 252

² Abramowitz and Stegun (1972) page 78

⁴ Rosenlicht (1968) page 157, Abramowitz and Stegun (1972) page 74

4. **Product of factors:** Let $\prod_{n=0}^{\infty} x_n \triangleq \lim_{N \to \infty} \prod_{n=0}^{N} x_n$ in some topological space. The sine and cosine functions can be defined in terms of a product of factors such that⁵

$$\cos(x) \triangleq \prod_{n=1}^{\infty} \left[1 - \left(\frac{x}{(2n-1)\frac{\pi}{2}} \right)^2 \right] \qquad \sin(x) \triangleq x \prod_{n=1}^{\infty} \left[1 - \left(\frac{x}{n\pi} \right)^2 \right]$$

5. **Partial fraction expansion:** The sine function can be defined in terms of a partial fraction expansion such that⁶

$$\sin(x) \triangleq \frac{1}{\frac{1}{x} + 2x \sum_{n=1}^{\infty} \frac{(-1)^n}{x^2 - (n\pi)^2}} \qquad \cos(x) \triangleq \underbrace{\left(\frac{1}{x} + 2x \sum_{n=1}^{\infty} \frac{1}{x^2 - (n\pi)^2}\right)}_{\cot(x)} \sin(x)$$

6. **Differential operator:** The sine and cosine functions can be defined as solutions to differential equations expressed in terms of the differential operator $\frac{d}{dx}$ such that

$$cos(x) \triangleq f(x)$$
 where $\frac{d^2}{dx^2}f + f = 0$ $f(0) = 1$ $\frac{d}{dx}f(0) = 0$ $f(0) = 0$

7. **Integral operator:** The sine and cosine functions can be defined as inverses of integrals of square roots of rational functions such that⁷

$$cos(x) \triangleq f^{-1}(x) \text{ where } f(x) \triangleq \underbrace{\int_{x}^{1} \sqrt{\frac{1}{1 - y^{2}}} dy}_{arccos(x)}$$

 $sin(x) \triangleq g^{-1}(x) \text{ where } g(x) \triangleq \underbrace{\int_{0}^{x} \sqrt{\frac{1}{1 - y^{2}}} dy}_{arcsin(x)}$

For purposes of analysis, it can be argued that the more natural approach for defining harmonic functions is in terms of the differentiation operator $\frac{d}{dx}$ (Definition G.1 page 155). Support for such an approach includes the following:

- Both sine and cosine are very easily represented analytically as polynomials with coefficients involving the operator $\frac{d}{dx}$ (Theorem G.1 page 156).
- All solutions of homogeneous second order differential equations are linear combinations of sine and cosine (Theorem G.3 page 158).
- Sine and cosine themselves are related to each other in terms of the differentiation operator (Theorem G.4 page 159).

⁷ Abramowitz and Stegun (1972) page 79



⁵ Abramowitz and Stegun (1972) page 75

⁶ Abramowitz and Stegun (1972) page 75

G.2. DEFINITIONS page 155 Daniel J. Greenhoe

The complex exponential function is a solution of a second order homogeneous differential equation (Definition G.4 page 160).

Sine and cosine are orthogonal with respect to an innerproduct generated by an integral operator—which is a kind of inverse differential operator (Section G.6 page 168).

G.2 Definitions

Definition G.1. ⁸ *Let C be the* SPACE OF ALL CONTINUOUSLY DIFFERENTIABLE REAL FUNCTIONS *and* $\frac{d}{dx} \in C^C$ the differentiation operator.

D E F

The function $f \in \mathbb{C}^{\mathbb{C}}$ is the **cosine** function $\cos(x) \triangleq f(x)$ if

- 1. $\frac{d^2}{dx^2}f + f = 0 \quad (second \ order \ homogeneous \ differential \ equation)$ 2. $f(0) = 1 \quad (first \ initial \ condition)$ 3. $\left[\frac{d}{dx}f\right](0) = 0 \quad (second \ initial \ condition).$

Definition G.2. ⁹ Let C and $\frac{d}{dx} \in C^C$ be defined as in definition of $\cos(x)$ (Definition G.1 page 155). The function $f \in \mathbb{C}^{\mathbb{C}}$ is the **sine** function $\sin(x) \triangleq f(x)$ if

D Ε

- 1. $\frac{d^2}{dx^2}f + f = 0$ (second order homogeneous differential equation) 2. f(0) = 0 (first initial condition)
- and
- $\left[\frac{d}{dt}f\right](0) = 1$ (second initial condition).

Definition G.3. 10

D E F

Let π ("pi") be defined as the element in $\mathbb R$ such that

- (1). $\cos\left(\frac{\pi}{2}\right) = 0$ and
- > 0 and (2).
- (3). π is the **smallest** of all elements in \mathbb{R} that satisfies (1) and (2).

G.3 Basic properties

Lemma G.1. 11 Let **C** be the space of all continuously differentiable real functions and $\frac{d}{dx} \in C^C$ the differentiation operator.

$$\begin{cases} \frac{d^{2}}{dx^{2}}f + f = 0 \end{cases} \iff \begin{cases} f(x) = [f](0) \sum_{n=0}^{\infty} (-1)^{n} \frac{x^{2n}}{(2n)!} + \left[\frac{d}{dx}f\right](0) \sum_{n=0}^{\infty} (-1)^{n} \frac{x^{2n+1}}{(2n+1)!} \\ = \left(f(0) + \left[\frac{d}{dx}f\right](0)x\right) - \left(\frac{f(0)}{2!}x^{2} + \frac{\left[\frac{d}{dx}f\right](0)}{3!}x^{3}\right) + \left(\frac{f(0)}{4!}x^{4} + \frac{\left[\frac{d}{dx}f\right](0)}{5!}x^{5}\right) \dots \end{cases}$$

¹¹ Rosenlicht (1968) page 156, Liouville (1839)





⁸ Rosenlicht (1968) page 157, Flanigan (1983) pages 228–229

⁹ Rosenlicht (1968) page 157, **₽** Flanigan (1983) pages 228–229

¹⁰ Rosenlicht (1968) page 158

 $^{\mathbb{Q}}$ Proof: Let $f'(x) \triangleq \frac{d}{dx} f(x)$.

$$f'''(x) = -\left[\frac{d}{dx}f\right](x)$$

$$f^{(4)}(x) = -\left[\frac{d}{dx}f\right](x)$$

$$= -\left[\frac{d^2}{dx^2}f\right](x) = f(x)$$

1. Proof that
$$\left[\frac{d^2}{dx^2}f\right](x) + f(x) = 0 \implies f(x) = \sum_{n=0}^{\infty} (-1)^n \left[\frac{f(0)}{(2n)!}x^{2n} + \frac{\left[\frac{d}{dx}f\right](0)}{(2n+1)!}x^{2n+1}\right]$$
:

$$f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(0)}{n!} x^n \quad \text{by Taylor expansion}$$

$$= f(0) + \left[\frac{d}{dx}f\right](0)x - \frac{\left[\frac{d^2}{dx^2}f\right](0)}{2!} x^2 - \frac{f^3(0)}{3!} x^3 + \frac{f^4(0)}{4!} x^4 + \frac{f^5(0)}{5!} x^5 - \cdots$$

$$= f(0) + \left[\frac{d}{dx}f\right](0)x - \frac{f(0)}{2!} x^2 - \frac{\left[\frac{d}{dx}f\right](0)}{3!} x^3 + \frac{f(0)}{4!} x^4 + \frac{\left[\frac{d}{dx}f\right](0)}{5!} x^5 - \cdots$$

$$= f(x) = \sum_{n=0}^{\infty} (-1)^n \left[\frac{f(0)}{(2n)!} x^{2n} + \frac{\left[\frac{d}{dx}f\right](0)}{(2n+1)!} x^{2n+1}\right]$$

2. Proof that
$$\left[\frac{d^2}{dx^2}f\right](x) + f(x) = 0 \iff f(x) = \sum_{n=0}^{\infty} (-1)^n \left[\frac{f(0)}{(2n)!}x^{2n} + \frac{\left[\frac{d}{dx}f\right](0)}{(2n+1)!}x^{2n+1}\right]$$
:

$$\begin{split} \left[\frac{d^2}{dx^2} f \right](x) &= \frac{d}{dx} \frac{d}{dx} \left[f(x) \right] \\ &= \frac{d}{dx} \frac{d}{dx} \sum_{n=0}^{\infty} (-1)^n \left[\frac{f(0)}{(2n)!} x^{2n} + \frac{\left[\frac{d}{dx} f \right](0)}{(2n+1)!} x^{2n+1} \right] \\ &= \sum_{n=1}^{\infty} (-1)^n \left[\frac{(2n)(2n-1)f(0)}{(2n)!} x^{2n-2} + \frac{(2n+1)(2n) \left[\frac{d}{dx} f \right](0)}{(2n+1)!} x^{2n-1} \right] \\ &= \sum_{n=1}^{\infty} (-1)^n \left[\frac{f(0)}{(2n-2)!} x^{2n-2} + \frac{\left[\frac{d}{dx} f \right](0)}{(2n-1)!} x^{2n-1} \right] \\ &= \sum_{n=0}^{\infty} (-1)^{n+1} \left[\frac{f(0)}{(2n)!} x^{2n} + \frac{\left[\frac{d}{dx} f \right](0)}{(2n+1)!} x^{2n+1} \right] \\ &= -f(x) \end{split}$$

by right hypothesis

by right hypothesis

Theorem G.1 (Taylor series for cosine/sine). 12

 $\cos(x) = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n}}{(2n)!} = 1 - \frac{x^2}{2} + \frac{x^4}{4!} - \frac{x^6}{6!} + \cdots$ $\forall x \in \mathbb{R}$ $\sin(x) = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{(2n+1)!} = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + \cdots$


¹² Rosenlicht (1968) page 157



^ℚProof:

$$\cos(x) = \underbrace{f(0) \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n}}{(2n)!}}_{\text{even terms}} + \underbrace{\left[\frac{d}{dx} f\right](0) \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{(2n+1)!}}_{\text{odd terms}}$$
by Lemma G.1 page 155
$$= 1 \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n}}{(2n)!} + 0 \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{(2n+1)!}$$
by cos initial conditions (Definition G.1 page 155)
$$= \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n}}{(2n)!} + \underbrace{\left[\frac{d}{dx} f\right](0) \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{(2n+1)!}}_{\text{odd terms}}$$
by Lemma G.1 page 155
$$= 0 \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n}}{(2n)!} + 1 \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{(2n+1)!}$$
by sin initial conditions (Definition G.2 page 155)
$$= \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{(2n+1)!}$$

Theorem G.2. ¹³




[♠]Proof:

$$\cos(0) = 1 - \frac{x^2}{2} + \frac{x^4}{4!} - \frac{x^6}{6!} + \cdots \bigg|_{x=0}$$
 by Taylor series for cosine (Theorem G.1 page 156)
$$= 1$$

$$\sin(0) = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + \cdots \bigg|_{x=0}$$
 by Taylor series for sine (Theorem G.1 page 156)
$$= 0$$

$$\cos(-x) = 1 - \frac{(-x)^2}{2} + \frac{(-x)^4}{4!} - \frac{(-x)^6}{6!} + \cdots$$
 by Taylor series for cosine (Theorem G.1 page 156)
$$= 1 - \frac{x^2}{2} + \frac{x^4}{4!} - \frac{x^6}{6!} + \cdots$$
 by Taylor series for cosine (Theorem G.1 page 156)
$$= \cos(x)$$
 by Taylor series for sine (Theorem G.1 page 156)
$$\sin(-x) = (-x) - \frac{(-x)^3}{3!} + \frac{(-x)^5}{5!} - \frac{(-x)^7}{7!} + \cdots$$
 by Taylor series for sine (Theorem G.1 page 156)
$$= -\left[x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + \cdots\right]$$
 by Taylor series for sine (Theorem G.1 page 156)

Lemma G.2. ¹⁴



¹³ Rosenlicht (1968) page 157

¹⁴ Rosenlicht (1968) page 158



♥Proof:

$$\cos(1) = 1 - \frac{x^2}{2} + \frac{x^4}{4!} - \frac{x^6}{6!} + \cdots \Big|_{x=1}$$
$$= 1 - \frac{1}{2} + \frac{1}{4!} - \frac{1}{6!} + \cdots$$

$$\cos(2) = 1 - \frac{x^2}{2} + \frac{x^4}{4!} - \frac{x^6}{6!} + \cdots \Big|_{x=2}$$
$$= 1 - \frac{4}{2} + \frac{16}{24} - \frac{64}{720} + \cdots$$

by Taylor series for cosine

(Theorem G.1 page 156)

by Taylor series for cosine

(Theorem G.1 page 156)

$$x \in (0:2)$$
 \implies each term in the sequence $\left(\left(x - \frac{x^3}{3!}\right), \left(\frac{x^5}{5!} - \frac{x^7}{7!}\right), \left(\frac{x^9}{9!} - \frac{x^{11}}{11!}\right), \dots\right)$ is > 0 \implies $\sin(x) > 0$

Proposition G.1. Let π be defined as in Definition G.3 (page 155).



The value π *exists in* \mathbb{R} .

 $2 < \pi < 4$. *(B)*.

< 0

^ℚProof:

$$\cos(1) > 0$$
$$\cos(2) < 0$$

$$\implies 1 < \frac{\pi}{2} < 2$$

$$\implies 2 < \pi < 4$$

by Lemma G.2 page 157

by Lemma G.2 page 157

Theorem G.3. 15 Let C be the space of all continuously differentiable real functions and $\frac{d}{dx} \in C^C$ the differentiation operator. Let $f'(0) \triangleq \left[\frac{d}{dx}f\right](0)$. $\begin{cases} \frac{d^2}{dx^2}f + f = 0 \end{cases} \iff \left\{ f(x) = f(0)\cos(x) + f'(0)\sin(x) \right\}$



$$\left\{ \frac{d^2}{dx^2} f + f = 0 \right\}$$

$$\iff$$

$$\left\{ f(x) = f(0)\cos(x) + f'(0)\sin(x) \right\}$$

[♠]Proof:

1. Proof that $\left[\frac{d^2}{dx^2}f\right](x) = -f(x) \implies f(x) = f(0)\cos(x) + \left[\frac{d}{dx}f\right](0)\sin(x)$:

$$f(x) = f(0) \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n}}{(2n)!} + \left[\frac{d}{dx} f\right](0) \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{(2n+1)!}$$
 by left hypothesis and Lemma G.1 page 15
$$\int_{\sin(x)}^{\cos(x)} (-1)^n \frac{x^{2n}}{(2n+1)!}$$
 by definitions of cos and sin (Definition G.1 page 155, Definition G.2 page 155)

by left hypothesis and Lemma G.1 page 155

¹⁵ Rosenlicht (1968) page 157.



2. Proof that $\frac{d^2}{dx^2}f = -f \iff f(x) = f(0)\cos(x) + \left[\frac{d}{dx}f\right](0)\sin(x)$:

$$f(x) = f(0)\cos x + \left[\frac{d}{dx}f\right](0)\sin x$$
 by right hypothesis
$$= f(0)\sum_{n=0}^{\infty} (-1)^n \frac{x^{2n}}{(2n)!} + \left[\frac{d}{dx}f\right](0)\sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{(2n+1)!}$$

$$\implies \frac{d^2}{dx^2}f + f = 0$$
 by Lemma G.1 page 155

Remark G.1. The general solution for the *non-homogeneous* equation $\frac{d^2}{dx^2}f(x) + f(x) = g(x)$ with initial conditions f(a) = 1 and $f'(a) = \rho$ is $f(x) = \cos(x) + \rho \sin(x) + \int_a^x g(y) \sin(x - y) dy$. This type of equation is called a *Volterra integral equation of the second type*. References: Folland (1992) page 371, Liouville (1839). Volterra equation references: Pedersen (2000) page 99, Lalescu (1908), **Lalescu** (1911)

Theorem G.4. ¹⁶ Let $\frac{d}{dx} \in C^C$ be the differentiation operator.

$$\frac{\mathbf{d}}{\mathbf{d}\mathbf{d}}\cos(x) = -\sin(x) \quad \forall x \in \mathbb{R} \quad \left| \begin{array}{c} \frac{\mathbf{d}}{\mathbf{d}\mathbf{k}}\sin(x) \\ \end{array} \right| = \cos(x) \quad \forall x \in \mathbb{R} \quad \left| \begin{array}{c} \cos^2(x) + \sin^2(x) \\ \end{array} \right| = 1 \quad \forall x \in \mathbb{R}$$

[♠]Proof:


$$\frac{d}{dx}\cos(x) = \frac{d}{dx}\sum_{n=0}^{\infty} (-1)^n \frac{x^{2n}}{(2n)!}$$
 by $Taylor series$ (Theorem G.1 page 156)
$$= \sum_{n=1}^{\infty} (-1)^n \frac{2nx^{2n-1}}{(2n)!} = \sum_{n=1}^{\infty} (-1)^n \frac{x^{2n-1}}{(2n-1)!} = \sum_{n=0}^{\infty} (-1)^{n+1} \frac{x^{2n}}{(2n)!}$$
 by $Taylor series$ (Theorem G.1 page 156)
$$\frac{d}{dx}\sin(x) = \frac{d}{dx}\sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{(2n+1)!}$$
 by $Taylor series$ (Theorem G.1 page 156)
$$= \sum_{n=0}^{\infty} (-1)^n \frac{(2n+1)x^{2n}}{(2n+1)!} = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n}}{(2n)!}$$
 by $Taylor series$ (Theorem G.1 page 156)
$$\frac{d}{dx}\left[\cos^2(x) + \sin^2(x)\right] = -2\cos(x)\sin(x) + 2\sin(x)\cos(x)$$
 by $Taylor series$ (Theorem G.1 page 156)
$$\frac{d}{dx}\left[\cos^2(x) + \sin^2(x)\right] = -2\cos(x)\sin(x) + 2\sin(x)\cos(x)$$
 by $Taylor series$ (Theorem G.1 page 156)
$$\Rightarrow \cos^2(x) + \sin^2(x)$$
 is $constant$

$$\Rightarrow \cos^2(x) + \sin^2(x)$$

$$= \cos^2(0) + \sin^2(0)$$

$$= 1 + 0 = 1$$
 by Theorem G.2 page 157

Proposition G.2.



¹⁶ Rosenlicht (1968) page 157

Frames and Bases Structure and Design [VERSION 021] https://github.com/dgreenhoe/pdfs/blob/master/msdframes.pdf





№PROOF:

$$\sin(\pi h) = \pm \sqrt{\sin^2(\pi h) + 0}$$

$$= \pm \sqrt{\sin^2(\pi h) + \cos^2(\pi h)}$$
 by definition of π (Definition G.3 page 155)
$$= \pm \sqrt{1}$$
 by Theorem G.4 page 159
$$= \pm 1$$

$$= 1$$
 by Lemma G.2 page 157

\blacksquare

G.4 The complex exponential

Definition G.4.


The function $f \in \mathbb{C}^{\mathbb{C}}$ is the **exponential function** $\exp(ix) \triangleq f(x)$ if

1. $\frac{d^2}{dx^2}f + f = 0$ (second order homogeneous differential equation) and

2. f(0) = 1 (first initial condition) and

3. $\left[\frac{d}{dx}f\right](0) = i$ (second initial condition).

Theorem G.5 (Euler's Identity). 17



♥PROOF:

D E F

$$\exp(ix) = f(0)\cos(x) + \left[\frac{d}{dx}f\right](0)\sin(x)$$
 by Theorem G.3 page 158
= $\cos(x) + i\sin(x)$ by Definition G.4 page 160



Proposition G.3.


$$e^{-i\pi h} = -i \mid e^{i\pi h} = i$$

♥Proof:

$$e^{i\pi h} = \cos(\pi h) + i\sin(\pi h)$$
 by $Euler's Identity$ (Theorem G.5 page 160)
$$= 0 + i$$
 by Theorem G.2 (page 157) and Proposition G.2 (page 159)
$$e^{-i\pi h} = \cos(\pi h) + i\sin(\pi h)$$
 by $Euler's Identity$ (Theorem G.5 page 160)
$$= \cos(\pi h) - i\sin(\pi h)$$
 by Theorem G.2 page 157
$$= 0 - i$$
 by Theorem G.2 (page 157) and Proposition G.2 (page 159)

₽

Corollary G.1.



¹⁷ Euler (1748), Bottazzini (1986) page 12



♥Proof:

$$|e^{ix}| = \cos(x) + i\sin(x)$$
 by Euler's Identity (Theorem G.5 page 160)
$$= \sum_{n \in \mathbb{W}} \frac{(-1)^n x^{2n}}{(2n)!} + i \sum_{n \in \mathbb{W}} \frac{(-1)^n x^{2n+1}}{(2n+1)!}$$
 by Taylor series (Theorem G.1 page 156)
$$= \sum_{n \in \mathbb{W}} \frac{(i^2)^n x^{2n}}{(2n)!} + \sum_{n \in \mathbb{W}} \frac{i(i^2)^n x^{2n+1}}{(2n+1)!}$$


$$= \sum_{n \in \mathbb{W} \cap \mathbb{Z}_e} \frac{(ix)^n}{n!} + \sum_{n \in \mathbb{W} \cap \mathbb{Z}_o} \frac{(ix)^n}{n!}$$

$$= \sum_{n \in \mathbb{W}} \frac{(ix)^n}{(2n)!} + \sum_{n \in \mathbb{W}} \frac{(ix)^n}{(2n+1)!}$$

$$= \sum_{n \in \mathbb{W}} \frac{(ix)^n}{n!} + \sum_{n \in \mathbb{W} \cap \mathbb{Z}_o} \frac{(ix)^n}{n!}$$

$$= \sum_{n \in \mathbb{W}} \frac{(ix)^n}{n!}$$


Corollary G.2 (Euler formulas). 18



[♠]Proof:

$$\begin{split} \boxed{\mathbf{R}_{\mathrm{e}}\Big(e^{ix}\Big)} &\triangleq \frac{e^{ix} + \left(e^{ix}\right)^*}{2} = \frac{e^{ix} + e^{-ix}}{2} & \text{by definition of } \mathfrak{R} & \text{(Definition F.5 page 149)} \\ &= \frac{\cos(x) + i\sin(x)}{2} + \frac{\cos(-x) + i\sin(-x)}{2} & \text{by } Euler's \ Identity & \text{(Theorem G.5 page 160)} \\ &= \frac{\cos(x) + i\sin(x)}{2} + \frac{\cos(x) - i\sin(x)}{2} & = \frac{\cos(x)}{2} + \frac{\cos(x)}{2} & = \cos(x) \\ \boxed{\mathbf{I}_{\mathrm{m}}\Big(e^{ix}\Big)} &\triangleq \frac{e^{ix} - \left(e^{ix}\right)^*}{2i} = \frac{e^{ix} - e^{-ix}}{2i} & \text{by definition of } \mathfrak{T} & \text{(Definition F.5 page 149)} \\ &= \frac{\cos(x) + i\sin(x)}{2i} - \frac{\cos(-x) + i\sin(-x)}{2i} & \text{by } Euler's \ Identity & \text{(Theorem G.5 page 160)} \\ &= \frac{\cos(x) + i\sin(x)}{2i} - \frac{\cos(x) - i\sin(x)}{2i} & = \frac{i\sin(x)}{2i} + \frac{i\sin(x)}{2i} & = \sin(x) \\ \end{bmatrix}$$

Theorem G.6. 19



№Proof:

$$e^{\alpha} e^{\beta} = \left(\sum_{n \in \mathbb{W}} \frac{\alpha^{n}}{n!}\right) \left(\sum_{m \in \mathbb{W}} \frac{\beta^{m}}{m!}\right)$$
 by Corollary G.1 page 160
$$= \sum_{n \in \mathbb{W}} \sum_{k=0}^{n} \frac{\alpha^{k}}{k!} \frac{\beta^{n-k}}{(n-k)!}$$

$$= \sum_{n \in \mathbb{W}} \sum_{k=0}^{n} \frac{n!}{n!} \frac{\alpha^{k}}{k!} \frac{\beta^{n-k}}{(n-k)!}$$

¹⁹ Rudin (1987) page 1

$$= \sum_{n \in \mathbb{W}} \frac{1}{n!} \sum_{k=0}^{n} \frac{n!}{k! (n-k)!} \alpha^{k} \beta^{n-k}$$

$$= \sum_{n \in \mathbb{W}} \frac{1}{n!} \sum_{k=0}^{n} \binom{n}{k} \alpha^{k} \beta^{n-k}$$

$$= \sum_{n \in \mathbb{W}} \frac{(\alpha + \beta)^{n}}{n!}$$

$$= e^{\alpha + \beta}$$

Daniel J. Greenhoe

by the Binomial Theorem

by Corollary G.1 page 160

₽

G.5 Trigonometric Identities

Theorem G.7 (shift identities).

Theorem 6 (offite facilities).								
I	$\cos\left(x+\frac{\pi}{2}\right)$	=	$-\sin x$	$\forall x \in \mathbb{R}$	$\sin\left(x+\frac{\pi}{2}\right)$	=	cosx	$\forall x \in \mathbb{R}$
H M	$\cos\left(x-\frac{\pi}{2}\right)$	=	$\sin x$	$\forall x \in \mathbb{R}$	$\sin\left(x-\frac{\pi}{2}\right)$	=	$-\cos x$	$\forall x \in \mathbb{R}$

NPROOF:

$$\cos\left(x+\frac{\pi}{2}\right) = \frac{e^{i\left(x+\frac{\pi}{2}\right)} + e^{-i\left(x+\frac{\pi}{2}\right)}}{2} \qquad \text{by $Euler formulas} \qquad \text{(Corollary G.2 page 161)}$$

$$= \frac{e^{ix}e^{i\frac{\pi}{2}} + e^{-ix}e^{-i\frac{\pi}{2}}}{2} \qquad \text{by $e^{a\beta} = e^a e^{\beta}$ result} \qquad \text{(Theorem G.6 page 161)}$$

$$= \frac{e^{ix}(i) + e^{-ix}(-i)}{2} \qquad \text{by Proposition G.3 page 160}$$

$$= \frac{e^{ix} - e^{-ix}}{-2i} \qquad \text{by $Euler formulas} \qquad \text{(Corollary G.2 page 161)}$$

$$\cos\left(x-\frac{\pi}{2}\right) = \frac{e^{i\left(x-\frac{\pi}{2}\right)} + e^{-i\left(x-\frac{\pi}{2}\right)}}{2} \qquad \text{by $Euler formulas} \qquad \text{(Corollary G.2 page 161)}$$

$$= \frac{e^{ix}e^{-i\frac{\pi}{2}} + e^{-ix}e^{+i\frac{\pi}{2}}}{2} \qquad \text{by $e^{a\beta} = e^a e^{\beta}$ result} \qquad \text{(Theorem G.6 page 161)}$$

$$= \frac{e^{ix}(-i) + e^{-ix}(i)}{2} \qquad \text{by Proposition G.3 page 160}$$

$$= \frac{e^{ix} - e^{-ix}}{2i} \qquad \text{by $Euler formulas} \qquad \text{(Corollary G.2 page 161)}$$

$$\sin\left(x+\frac{\pi}{2}\right) = \cos\left(\left[x+\frac{\pi}{2}\right] - \frac{\pi}{2}\right) \qquad \text{by previous result}$$

$$= \cos(x)$$

$$\sin\left(x-\frac{\pi}{2}\right) = -\cos\left(\left[x-\frac{\pi}{2}\right] + \frac{\pi}{2}\right) \qquad \text{by previous result}$$

$$= -\cos(x)$$

Theorem G.8 (product identities).

	(A).	cosxcosy	=	$\frac{1}{2}\cos(x-y)$	+	$^{1}h\cos(x+y)$	$\forall x,y \in \mathbb{R}$
H	(B).	$\cos x \sin y$	=	$-1/2\sin(x-y)$	+	$^{1}h\sin(x+y)$	$\forall x,y \in \mathbb{R}$
М	(C).	$\sin x \cos y$	=	$\frac{1}{2}\sin(x-y)$	+	$^{1}h\sin(x+y)$	$\forall x,y \in \mathbb{R}$
	(D).	$\sin x \sin y$	=	$^{1}h\cos(x-y)$	_	$^{1}/\cos(x+y)$	$\forall x,y \in \mathbb{R}$

^ℚProof:

1. Proof for (A) using Euler formulas (Corollary G.2 page 161) (algebraic method requiring *complex number system* \mathbb{C}):

$$\cos x \cos y = \left(\frac{e^{ix} + e^{-ix}}{2}\right) \left(\frac{e^{iy} + e^{-iy}}{2}\right)$$
 by *Euler formulas* (Corollary G.2 page 161)
$$= \frac{e^{i(x+y)} + e^{i(x-y)} + e^{i(-x+y)} + e^{i(-x-y)}}{4}$$

$$= \frac{e^{i(x+y)} + e^{-i(x+y)}}{4} + \frac{e^{i(x-y)} + e^{-i(x-y)}}{4}$$
 by *Euler formulas* (Corollary G.2 page 161)
$$= \frac{1}{2}\cos(x+y) + \frac{1}{2}\cos(x-y)$$

2. Proof for (A) using Volterra integral equation (Theorem G.3 page 158) (differential equation method requiring only *real number system* \mathbb{R}):

$$f(x) \triangleq \frac{1}{2} cos(x - y) + \frac{1}{2} cos(x + y)$$

$$\Rightarrow \frac{d}{dx} f(x) = -\frac{1}{2} cos(x - y) - \frac{1}{2} cos(x + y)$$
by Theorem G.4 page 159
$$\Rightarrow \frac{d^2}{dx^2} f(x) = -\frac{1}{2} cos(x - y) - \frac{1}{2} cos(x + y)$$
by Theorem G.4 page 159
$$\Rightarrow \frac{d^2}{dx^2} f(x) + f(x) = 0$$
by additive inverse property
$$\Rightarrow \frac{1}{2} cos(x - y) + \frac{1}{2} cos(x + y) = \frac{\left[\frac{1}{2} cos(0 - y) + \frac{1}{2} cos(0 + y)\right] cos(x)}{f''(0)} + \frac{\left[-\frac{1}{2} cos(0 - y) - \frac{1}{2} cos(0 + y)\right] cos(x)}{f''(0)}$$

$$\Rightarrow \frac{1}{2} cos(x - y) + \frac{1}{2} cos(x + y) = cosycosx + 0sin(x)$$

$$\Rightarrow cosxcosy = \frac{1}{2} cos(x - y) + \frac{1}{2} cos(x - y) + \frac{1}{2} cos(x + y)$$

3. Proof for (B) using Euler formulas (Corollary G.2 page 161):

$$sinxsiny = \left(\frac{e^{ix} - e^{-ix}}{2i}\right) \left(\frac{e^{iy} - e^{-iy}}{2i}\right)$$

$$= \frac{e^{i(x+y)} - e^{i(x-y)} - e^{i(-x+y)} + e^{i(-x-y)}}{-4}$$

$$= \frac{e^{i(x+y)} + e^{-i(x+y)} - e^{i(x-y)} - e^{-i(x-y)}}{-4}$$

$$= \frac{e^{i(x+y)} + e^{-i(x+y)}}{-4} - \frac{e^{i(x-y)} + e^{-i(x-y)}}{-4}$$

$$= \frac{2\cos(x-y)}{4} - \frac{2\cos(x+y)}{4}$$
by Corollary G.2 page 161
$$= \frac{1}{2}\cos(x-y) - \frac{1}{2}\cos(x+y)$$

4. Proofs for (C) and (D) using (A) and (B):

$$\begin{aligned} \cos x \sin y &= \cos(x) \cos\left(y - \frac{\pi}{2}\right) & \text{by } \textit{shift identities} \\ &= \frac{1}{2} \cos\left(x + y - \frac{\pi}{2}\right) + \frac{1}{2} \cos\left(x - y + \frac{\pi}{2}\right) & \text{by (A)} \\ &= \frac{1}{2} \sin(x + y) - \frac{1}{2} \sin(x - y) & \text{by } \textit{shift identities} \end{aligned} \qquad \text{(Theorem G.7 page 162)} \\ \sin x \cos y &= \cos y \sin x \\ &= \frac{1}{2} \sin(y + x) - \frac{1}{2} \sin(y - x) & \text{by (B)} \\ &= \frac{1}{2} \sin(x + y) + \frac{1}{2} \sin(x - y) & \text{by Theorem G.2 page 157} \end{aligned}$$

Proposition G.4.

P	(A).	$\cos(\pi)$	=	-1	(C).	$\cos(2\pi)$ $\sin(2\pi)$	=	1	(E).	$e^{i\pi}$	=	-1
P	(B).	$sin(\pi)$	=	0	(D).	$\sin(2\pi)$	=	0	(F).	$e^{i2\pi}$	=	0

№ PROOF:

$$\begin{aligned} \cos(\pi) &= -1 + 1 + \cos(\pi) \\ &= -1 + 2[\frac{1}{2}\cos(\pi h - \pi h) + \frac{1}{2}\cos(\pi h + \pi h)] \\ &= -1 + 2\cos(\pi h - \pi h) + \frac{1}{2}\cos(\pi h + \pi h)] \\ &= -1 + 2\cos(\pi h - \pi h) + \frac{1}{2}\cos(\pi h + \pi h)] \\ &= -1 + 2(0)(0) \\ &= -1 + 2(0)(0) \\ &= -1 \end{aligned}$$
 by definition of π (Definition G.3 page 155)
$$= -1$$
 sin(π) = $0 + \sin(\pi)$ by $\sin(0) = 0$ result (Theorem G.2 page 157)
$$= 2\cos(\pi h)\sin(\pi h)$$
 by $\sin(0) = 0$ result (Theorem G.2 page 157)
$$= 2\cos(\pi h)\sin(\pi h)$$
 by πh definition of π (Definition G.3 page 155)
$$= 0$$
 (Theorem G.8 page 162)
$$= 2(0)\sin(\pi h)$$
 by definition of π (Definition G.3 page 155)
$$= 0$$
 (Definition G.3 page 155)
$$= 0$$
 (Theorem G.8 page 162)
$$= 2(\frac{1}{2}\cos(\pi - \pi) + \frac{1}{2}\cos(\pi + \pi)] - 1$$
 by $\cos(0) = 1$ result (Theorem G.2 page 157)
$$= 2\cos(\pi h)\cos(\pi) - 1$$
 by πh definition of πh (Theorem G.8 page 162)
$$= 2(-1)(-1) - 1$$
 by πh definition of πh (Theorem G.8 page 162)
$$= 2(-1)(-1) - 1$$
 by πh definition of πh (Theorem G.8 page 162)
$$= 2(-1)(-1) - 1$$
 by πh definition of πh (Theorem G.8 page 162)
$$= 2(-1)(-1) - 1$$
 by πh definition of πh (Theorem G.8 page 162)
$$= 2(-1)(-1) - 1$$
 by πh definition of πh (Definition G.3 page 157)
$$= 2\sin(\pi h)\cos(\pi h) + \frac{1}{2}\sin(\pi h) + \frac{1}{2}\sin$$

Theorem G.9 (double angle formulas). ²⁰

		(A).	$\cos(x+y)$	=	$\cos x \cos y - \sin x \sin y$	$\forall x,y \in \mathbb{R}$
	Ā	(B).	$\sin(x+y)$	=	$\sin x \cos y + \cos x \sin y$	$\forall x,y \in \mathbb{R}$
M	(C)	tan(x + y)	=	$\tan x + \tan y$	$\forall x,y \in \mathbb{R}$	
	(C).	tan(x + y)		$1 - \tan x \tan y$	v <i>x,y</i> ∈™	

№PROOF:

1. Proof for (A) using product identities (Theorem G.8 page 162).

$$\cos(x+y) = \underbrace{\frac{1}{2}\cos(x+y) + \frac{1}{2}\cos(x+y)}_{\cos(x+y)} + \underbrace{\frac{1}{2}\cos(x-y) - \frac{1}{2}\cos(x-y)}_{0}$$

$$= \left[\frac{1}{2}\cos(x-y) + \frac{1}{2}\cos(x+y)\right] - \left[\frac{1}{2}\cos(x-y) - \frac{1}{2}\cos(x+y)\right]$$

$$= \cos x \cos y - \sin x \sin y$$

by Theorem G.8 page 162

2. Proof for (A) using Volterra integral equation (Theorem G.3 page 158):

$$f(x) \triangleq \cos(x+y) \implies \frac{d}{dx}f(x) = -\sin(x+y) \qquad \text{by Theorem G.4 page 159}$$

$$\implies \frac{d^2}{dx^2}f(x) = -\cos(x+y) \qquad \text{by Theorem G.4 page 159}$$

$$\implies \frac{d^2}{dx^2}f(x) + f(x) = 0 \qquad \text{by additive inverse property}$$

$$\implies \cos(x+y) = \cos y \cos x - \sin y \sin x \qquad \text{by Theorem G.3 page 158}$$

$$\implies \cos(x+y) = \cos x \cos y - \sin x \sin y \qquad \text{by commutative property}$$

3. Proof for (B) and (C) using (A):

$$\sin(x+y) = \cos\left(x - \frac{\pi}{2} + y\right)$$
 by shift identities (Theorem G.7 page 162)
$$= \cos\left(x - \frac{\pi}{2}\right)\cos(y) - \sin\left(x - \frac{\pi}{2}\right)\sin(y)$$
 by (A)
$$= \sin(x)\cos(y) + \cos(x)\sin(y)$$
 by shift identities (Theorem G.7 page 162)

$$\tan(x+y) = \frac{\sin(x+y)}{\cos(x+y)}$$

$$= \frac{\sin x \cos y + \cos x \sin y}{\cos x \cos y - \sin x \sin y} \qquad \text{by (A)}$$


$$= \left(\frac{\sin x \cos y + \cos x \sin y}{\cos x \cos y - \sin x \sin y}\right) \left(\frac{\cos x \cos y}{\cos x \cos y}\right)$$

$$= \frac{\frac{\sin x \cos y}{\cos x \cos y} + \frac{\cos x \sin y}{\cos x \cos y}}{\frac{\cos x \cos y}{\cos x \cos y}} = \frac{\frac{\sin x}{\cos x} + \frac{\sin y}{\cos y}}{1 - \frac{\sin x}{\cos x} \frac{\sin y}{\cos y}} = \frac{\tan x + \tan y}{1 - \tan x \tan y}$$

Theorem G.10 (trigonometric periodicity).

		` 0		1	J ,							
ηl	(A).	$\cos(x + M\pi)$	=	$(-1)^M \cos(x)$	$\forall x \in \mathbb{R},$	$M \in \mathbb{Z}$	(D).	$\cos(x + 2M\pi)$	=	cos(x)	$\forall x \in \mathbb{R},$	$M \in \mathbb{Z}$
Ĥ	(B).	$\sin(x + M\pi)$	=	$(-1)^M \sin(x)$	$\forall x \in \mathbb{R},$	$M \in \mathbb{Z}$	(E).	$\sin(x + 2M\pi)$ $\sin(x + 2M\pi)$ $\sin(x + 2M\pi)$	=	sin(x)	$\forall x \in \mathbb{R},$	$M \in \mathbb{Z}$
M	(C).	$e^{i(x+M\pi)}$	=	$(-1)^{M}e^{ix}$	$\forall x \in \mathbb{R},$	$M \in \mathbb{Z}$	(F).	$e^{i(x+2M\pi)}$	=	e^{ix}	$\forall x \in \mathbb{R},$	$M \in \mathbb{Z}$

²⁰Expressions for $\cos(\alpha + \beta)$, $\sin(\alpha + \beta)$, and $\sin^2 x$ appear in works as early as Ptolemy (circa 100AD). Reference: http://en.wikipedia.org/wiki/History_of_trigonometric_functions





- 1. Proof for (A):
 - (a) M = 0 case: $\cos(x + 0\pi) = \cos(x) = (-1)^0 \cos(x)$
 - (b) Proof for M > 0 cases (by induction):
 - i. Base case M = 1:

$$\cos(x+\pi) = \cos x \cos \pi - \sin x \sin \pi$$
 by double angle formulas (Theorem G.9 page 165)
 $= \cos x(-1) - \sin x(0)$ by $\cos \pi = -1$ result (Proposition G.4 page 164)
 $= (-1)^1 \cos x$

ii. Inductive step...Proof that M case $\implies M + 1$ case:

$$\cos(x + [M+1]\pi) = \cos([x+\pi] + M\pi)$$

$$= (-1)^{M} \cos(x + \pi)$$
 by induction hypothesis (*M* case)
$$= (-1)^{M} (-1) \cos(x)$$
 by base case (item (1(b)i) page 166)
$$= (-1)^{M+1} \cos(x)$$

$$\implies M+1 \text{ case}$$

(c) Proof for M < 0 cases: Let $N \triangleq -M ... \implies N > 0$.

$$\cos(x + M\pi) \triangleq \cos(x - N\pi) \qquad \text{by definition of } N$$

$$= \cos(x)\cos(-N\pi) - \sin(x)\sin(-N\pi) \qquad \text{by double angle formulas} \qquad \text{(Theorem G.9 page 165)}$$

$$= \cos(x)\cos(N\pi) + \sin(x)\sin(N\pi) \qquad \text{by Theorem G.2 page 157}$$

$$= \cos(x)\cos(0 + N\pi) + \sin(x)\sin(0 + N\pi)$$

$$= \cos(x)(-1)^N\cos(0) + \sin(x)(-1)^N\sin(0) \qquad \text{by } M \geq 0 \text{ results} \qquad \text{(item (1b) page 166)}$$

$$= (-1)^N\cos(x) \qquad \text{by } \cos(0) = 1, \sin(0) = 0 \text{ results} \qquad \text{(Theorem G.2 page 157)}$$

$$\triangleq (-1)^{-M}\cos(x) \qquad \text{by definition of } N$$

$$= (-1)^M\cos(x) \qquad \text{by definition of } N$$

(d) Proof using complex exponential:

$$\cos(x + M\pi) = \frac{e^{i(x + M\pi)} + e^{-i(x + M\pi)}}{2}$$
 by Euler formulas (Corollary G.2 page 161)

$$= e^{iM\pi} \left[\frac{e^{ix} + e^{-ix}}{2} \right]$$
 by $e^{\alpha\beta} = e^{\alpha}e^{\beta}$ result (Theorem G.6 page 161)

$$= \left(e^{i\pi} \right)^{M} \cos x$$
 by Euler formulas (Corollary G.2 page 161)

$$= (-1)^{M} \cos x$$
 by $e^{i\pi} = -1$ result (Proposition G.4 page 164)

- 2. Proof for (B):
 - (a) M = 0 case: $\sin(x + 0\pi) = \sin(x) = (-1)^0 \sin(x)$
 - (b) Proof for M > 0 cases (by induction):
 - i. Base case M = 1:

$$\sin(x + \pi) = \sin x \cos \pi + \cos x \sin \pi$$
 by double angle formulas (Theorem G.9 page 165)
 $= \sin x (-1) - \cos x (0)$ by $\sin \pi = 0$ results (Proposition G.4 page 164)
 $= (-1)^1 \sin x$

ii. Inductive step...Proof that M case $\implies M + 1$ case:

$$\sin(x + [M+1]\pi) = \sin([x+\pi] + M\pi)$$

$$= (-1)^M \sin(x + \pi)$$

$$= (-1)^M (-1)\sin(x)$$
by induction hypothesis (*M* case)
by base case (item (2(b)i) page 166)
$$= (-1)^{M+1} \sin(x)$$

$$\implies M+1 \text{ case}$$

(c) Proof for M < 0 cases: Let $N \triangleq -M ... \implies N > 0$.

$$\sin(x + M\pi) \triangleq \sin(x - N\pi) \qquad \text{by definition of } N$$

$$= \sin(x)\sin(-N\pi) - \sin(x)\sin(-N\pi) \qquad \text{by double angle formulas} \qquad \text{(Theorem G.9 page 165)}$$

$$= \sin(x)\sin(N\pi) + \sin(x)\sin(N\pi) \qquad \text{by Theorem G.2 page 157}$$

$$= \sin(x)\sin(0 + N\pi) + \sin(x)\sin(0 + N\pi)$$

$$= \sin(x)(-1)^N\sin(0) + \sin(x)(-1)^N\sin(0) \qquad \text{by } M \ge 0 \text{ results} \qquad \text{(item (2b) page 166)}$$

$$= (-1)^N\sin(x) \qquad \text{by } \sin(0) = 1, \sin(0) = 0 \text{ results} \qquad \text{(Theorem G.2 page 157)}$$

$$\triangleq (-1)^{-M}\sin(x) \qquad \text{by definition of } N$$

$$= (-1)^{M}\sin(x) \qquad \text{by definition of } N$$

(d) Proof using complex exponential:

$$\sin(x + M\pi) = \frac{e^{i(x + M\pi)} - e^{-i(x + M\pi)}}{2i} \qquad \text{by } \textit{Euler formulas} \qquad \text{(Corollary G.2 page 161)}$$

$$= e^{iM\pi} \left[\frac{e^{ix} - e^{-ix}}{2i} \right] \qquad \text{by } e^{\alpha\beta} = e^{\alpha}e^{\beta} \text{ result} \qquad \text{(Theorem G.6 page 161)}$$

$$= \left(e^{i\pi} \right)^{M} \sin x \qquad \text{by } \textit{Euler formulas} \qquad \text{(Corollary G.2 page 161)}$$

$$= (-1)^{M} \sin x \qquad \text{by } e^{i\pi} = -1 \text{ result} \qquad \text{(Proposition G.4 page 164)}$$


3. Proof for (C):

$$e^{i(x+M\pi)}=e^{iM\pi}e^{ix}$$
 by $e^{\alpha\beta}=e^{\alpha}e^{\beta}$ result (Theorem G.6 page 161)
$$=\left(e^{i\pi}\right)^{M}\left(e^{ix}\right)$$

$$=\left(-1\right)^{M}e^{ix}$$
 by $e^{i\pi}=-1$ result (Proposition G.4 page 164)

4. Proofs for (D), (E), and (F): $\cos(i(x + 2M\pi)) = (-1)^{2M}\cos(ix) = \cos(ix)$ by (A) $\sin(i(x + 2M\pi)) = (-1)^{2M}\sin(ix) = \sin(ix)$ by (B) $e^{i(x+2M\pi)} = (-1)^{2M}e^{ix} = e^{ix}$ by (C)

Theorem G.11 (half-angle formulas/squared identities).



№ Proof:

$$\cos^2 x \triangleq (\cos x)(\cos x) = \frac{1}{2}\cos(x-x) + \frac{1}{2}\cos(x+x) \qquad \text{by } product \, identities} \qquad \text{(Theorem G.8 page 162)}$$

$$= \frac{1}{2}[1+\cos(2x)] \qquad \qquad \text{by } \cos(0) = 1 \, \text{result} \qquad \text{(Theorem G.2 page 157)}$$

$$\sin^2 x = (\sin x)(\sin x) = \frac{1}{2}\cos(x-x) - \frac{1}{2}\cos(x+x) \qquad \text{by } product \, identities} \qquad \text{(Theorem G.8 page 162)}$$

$$= \frac{1}{2}[1-\cos(2x)] \qquad \qquad \text{by } \cos(0) = 1 \, \text{result} \qquad \text{(Theorem G.2 page 157)}$$

$$\cos^2 x + \sin^2 x = \frac{1}{2}[1+\cos(2x)] + \frac{1}{2}[1-\cos(2x)] = 1 \qquad \text{by } (A) \, \text{and } (B)$$

$$\text{note: see also} \qquad \text{Theorem G.4 page 159}$$

□>

G.6 Planar Geometry

The harmonic functions cos(x) and sin(x) are *orthogonal* to each other in the sense

$$\langle \cos(x) | \sin(x) \rangle = \int_{-\pi}^{+\pi} \cos(x) \sin(x) \, dx$$

$$= \frac{1}{2} \int_{-\pi}^{+\pi} \sin(x - x) \, dx + \frac{1}{2} \int_{-\pi}^{+\pi} \sin(x + x) \, dx \qquad \text{by Theorem G.8 page 162}$$

$$= \frac{1}{2} \int_{-\pi}^{+\pi} \sin(0) \, dx + \frac{1}{2} \int_{-\pi}^{+\pi} \sin(2x) \, dx$$

$$= -\frac{1}{2} \cdot \frac{1}{2} \cos(2x) \Big|_{-\pi}^{+\pi} \cos(2x)$$

$$= -\frac{1}{4} [\cos(2\pi) - \cos(-2\pi)]$$


$$= 0$$

Because cos(x) are sin(x) are orthogonal, they can be conveniently represented by the x and y axes in a plane—because perpendicular axes in a plane are also orthogonal. Vectors in the plane can be represented by linear combinations of cosx and sinx. Let tan x be defined as

$$\tan x \triangleq \frac{\sin x}{\cos x}.$$

We can also define a value θ to represent the angle between such a vector and the x-axis such that

$$\theta = \tan^{-1}\left(\frac{\sin\theta}{\cos\theta}\right)$$



$$\begin{array}{cccc}
\cos\theta & \triangleq & \frac{x}{r} & \sec\theta & \triangleq & \frac{r}{x} \\
\sin\theta & \triangleq & \frac{y}{r} & \csc\theta & \triangleq & \frac{x}{y} \\
\tan\theta & \triangleq & \frac{y}{x} & \cot\theta & \triangleq & \frac{x}{y}
\end{array}$$

G.7 Trigonometric functions of complex numbers

Definition G.5. 21 $\begin{array}{cccc}
\text{D} & \cosh(z) & \triangleq & \frac{e^z + e^{-z}}{2} & \forall z \in \mathbb{C} \\
\text{sinh}(z) & \triangleq & \frac{e^z - e^{-z}}{2} & \forall z \in \mathbb{C}
\end{array}$

²¹ Saxelby (1920) page 225



Theorem G.12. 22

	cosh(ix)	=	cos(x)	$\forall x$	∈ℝ
	sinh(ix)	=	$i\sin(x)$	$\forall x$	∈ℝ
I	cos(ix)	=	cosh(x)	$\forall x$	∈ℝ
H	sin(ix)	=	$i \sinh(x)$	$\forall x$	∈ℝ
	$\cos(x+iy)$	=	$\cos(x)\cosh(y) - i\sin(x)\sinh(y)$	∀ <i>x</i> , j	∕∈R
	$\sin(x+iy)$	=	$\sin(x)\cosh(y) + i\cos(x)\sinh(y)$	∀ <i>x</i> ,,	√∈R

№PROOF:

$$\cosh(ix) \triangleq \frac{e^{ix} + e^{-ix}}{2} \qquad \text{by definition of } \cosh(x) \qquad \text{(Definition G.5 page 168)}$$

$$= \cos(x) \qquad \text{by } Euler's \, Identity \qquad \text{(Theorem G.5 page 160)}$$

$$\sinh(ix) \triangleq \frac{e^{ix} - e^{-ix}}{2} \qquad \text{by definition of } \sinh(x) \qquad \text{(Definition G.5 page 168)}$$

$$\triangleq i \left[\frac{e^{ix} - e^{-ix}}{2i} \right] \qquad \text{by definition of } \sinh(x) \qquad \text{(Definition G.5 page 168)}$$

$$= i \sin(x) \qquad \text{by } Euler's \, Identity \qquad \text{(Theorem G.5 page 160)}$$

$$\cos(ix) \triangleq \frac{e^{iix} + e^{-iix}}{2} \qquad \text{by } Euler's \, Identity \qquad \text{(Theorem G.5 page 160)}$$

$$= \frac{e^{-x} + e^{-x}}{2}$$

$$= \frac{e^{x} + e^{-x}}{2}$$

$$\triangleq \cosh(x) \qquad \text{by definition of } \cosh(x) \qquad \text{(Definition G.5 page 160)}$$

$$\sin(ix) \triangleq \frac{e^{iix} - e^{-iix}}{2i} \qquad \text{by } Euler's \, Identity \qquad \text{(Theorem G.5 page 160)}$$

$$\sin(ix) \triangleq \frac{e^{iix} - e^{-iix}}{2i} \qquad \text{by } Euler's \, Identity \qquad \text{(Theorem G.5 page 160)}$$

$$= \frac{e^{-x} - e^{x}}{2i}$$

$$= -(-i^{2}) \left[\frac{e^{x} - e^{-x}}{2i} \right]$$

$$= i \left[\frac{e^{x} - e^{-x}}{2i} \right]$$

$$\triangleq i \sinh(x) \qquad \text{by } definition of \cosh(x) \qquad \text{(Definition G.5 page 168)}$$

$$\cos(x + iy) = \cos(x) \cos(iy) - \sin(x) \sin(iy) \qquad \text{by } double \, angle \, formulas}$$

$$= \cos(x) \cosh(y) - i \sin(x) \sinh(y) \qquad \text{by } previous \, results}$$

$$\sin(x + iy) = \sin(x) \cos(iy) + \cos(x) \sin(iy) \qquad \text{by } double \, angle \, formulas}$$

$$= \sin(x) \cosh(y) + i \cos(x) \sin(iy) \qquad \text{by } previous \, results}$$

²²https://proofwiki.org/wiki/Cosine_of_Complex_Number, https://proofwiki.org/wiki/Sine_of_Complex_Number,

Saxelby (1920) pages 416-417





₽

G.8 The power of the exponential



Gentlemen, that is surely true, it is absolutely paradoxical; we cannot understand it, and we don't know what it means. But we have proved it, and therefore we know it must be the truth.

→

Benjamin Peirce (1809–1880), American Harvard University mathematician after proving $e^{i\pi} = -1$ in a lecture. ²³



♣ Young man, in mathematics you don't understand things. You just get used to
them.
♠

John von Neumann (1903–1957), Hungarian-American mathematician, as allegedly told to Gary Zukav by Felix T. Smith, Head of Molecular Physics at Stanford Research Institute, about a "physicist friend". 24

The following corollary presents one of the most amazing relations in all of mathematics. It shows a simple and compact relationship between the transcendental numbers π and e, the imaginary number i, and the additive and multiplicative identity elements 0 and 1. The fact that there is any relationship at all is somewhat amazing; but for there to be such an elegant one is truly one of the worders of the world of numbers.

Corollary G.3. ²⁵

$$\begin{array}{c} \mathbf{C} \\ \mathbf{O} \\ \mathbf{R} \end{array} e^{i\pi} + 1 = 0$$

[♠]Proof:

$$e^{ix}\big|_{x=\pi} = [\cos x + i \sin x]_{x=\pi}$$
 by Euler's Identity (Theorem G.5 page 160)
 $= -1 + i \cdot 0$ by Proposition G.4 page 164
 $= -1$

There are many transforms available, several of them integral transforms $[\mathbf{A}\mathbf{f}](s) \triangleq \int_t \mathbf{f}(s)\kappa(t,s) \, ds$ using different kernels $\kappa(t,s)$. But of all of them, two of the most often used themselves use an exponential kernel:

- ① The *Laplace Transform* with kernel $\kappa(t, s) \triangleq e^{st}$
- ② The Fourier Transform with kernel $\kappa(t, \omega) \triangleq e^{i\omega t}$.

quote: A Kasner and Newman (1940) page 104

image: http://www-history.mcs.st-andrews.ac.uk/history/PictDisplay/Peirce_Benjamin.html

24 quote: **Zukav** (1980) page 208

image: http://en.wikipedia.org/wiki/John_von_Neumann

The quote appears in a footnote in Zukav (1980) that reads like this: Dr. Felix Smith, Head of Molecular Physics, Stanford Research Institute, once related to me the true story of a physicist friend who worked at Los Alamos after World War II. Seeking help on a difficult problem, he went to the great Hungarian mathematician, John von Neumann, who was at Los Alamos as a consultant. "Simple," said von Neumann. "This can be solved by using the method of characteristics." After the explanation the physicist said, "I'm afraid I don't understand the method of characteristics." "Young man," said von Neumann, "in mathematics you don't understand things, you just get used to them."

²⁵ ■ Euler (1748), ■ Euler (1988) ⟨chapter 8?⟩, http://www.daviddarling.info/encyclopedia/E/Eulers_formula.



Of course, the Fourier kernel is just a special case of the Laplace kernel with $s = i\omega$ ($i\omega$ is a unit circle in s if s is depicted as a plane with real and imaginary axes). What is so special about exponential kernels? Is it just that they were discovered sooner than other kernels with other transforms? The answer in general is "no". The exponential has two properties that makes it extremely special:

- Mark The exponential is an eigenvalue of any linear time invariant (LTI) operator (Theorem G.13)
- **4** The exponential generates a *continuous point spectrum* for the *differential operator*.

Theorem G.13. ²⁶ Let L be an operator with kernel $h(t, \omega)$ and $\check{\mathsf{h}}(s) \triangleq \left\langle \left. \mathsf{h}(t, \omega) \right| e^{st} \right\rangle$ (LAPLACE TRANSFORM).

$$\left\{ \begin{array}{l} \text{L } is \text{ Linear } and \\ \text{2. } L is \text{ TIME-INVARIANT} \end{array} \right\} \qquad \Longrightarrow \qquad \left\{ \begin{array}{l} \text{Le}^{st} = \underbrace{\check{\mathsf{h}}^*(-s)}_{eigenvalue} \underbrace{e^{st}}_{eigenvector} \end{array} \right\}$$

№Proof:

$$[\mathbf{L}e^{st}](s) = \langle e^{su} \mid \mathsf{h}((t;u),s) \rangle \qquad \text{by linear hypothesis}$$

$$= \langle e^{su} \mid \mathsf{h}((t-u),s) \rangle \qquad \text{by time-invariance hypothesis}$$

$$= \langle e^{s(t-v)} \mid \mathsf{h}(v,s) \rangle \qquad \text{let } v = t - u \implies u = t - v$$

$$= e^{st} \langle e^{-sv} \mid \mathsf{h}(v,s) \rangle \qquad \text{by additivity of } \langle \triangle \mid \nabla \rangle$$

$$= \langle \mathsf{h}(v,s) \mid e^{-sv} \rangle^* e^{st} \qquad \text{by conjugate symmetry of } \langle \triangle \mid \nabla \rangle$$

$$= \langle \mathsf{h}(v,s) \mid e^{(-s)v} \rangle^* e^{st}$$

$$= \check{\mathsf{h}}^*(-s) e^{st} \qquad \text{by definition of } \check{\mathsf{h}}(s)$$

²⁶ Mallat (1999) page 2, ...page 2 online: http://www.cmap.polytechnique.fr/~mallat/WTintro.pdf





TRIGONOMETRIC POLYNOMIALS



■ I turn aside with a shudder of horror from this lamentable plague of functions which have no derivatives.

■

Charles Hermite (1822 – 1901), French mathematician, in an 1893 letter to Stieltjes, in response to the "pathological" everywhere continuous but nowhere differentiable *Weierstrass functions* $f(x) = \sum_{n=0}^{\infty} a^n \cos(b^n \pi x)$.

H.1 Trigonometric expansion

Theorem H.1 (DeMoivre's Theorem).

$$\begin{array}{c} \mathsf{T} \\ \mathsf{H} \\ \mathsf{M} \end{array} \left(re^{ix} \right)^n = r^n (\cos nx + i \sin nx) \qquad \forall r, x \in \mathbb{R}$$

♥Proof:

$$(re^{ix})^n = r^n e^{inx}$$

= $r^n (\cos nx + i\sin nx)$ by Euler's identity (Theorem G.5 page 160)

The cosine with argument nx can be expanded as a polynomial in cos(x) (next).

Theorem H.2 (trigonometric expansion). ²

1 quote: ☐ Hermite (1893) translation: ☐ Lakatos (1976) page 19 image: http://www-groups.dcs.sx-and.ac.uk/~history/PictDisplay/Hermite.html 2 ☐ Rivlin (1974) page 3 ⟨(1.8)⟩

$$\cos(nx) = \sum_{k=0}^{\left\lfloor \frac{n}{2} \right\rfloor} \sum_{m=0}^{k} (-1)^{k+m} {n \choose 2k} {k \choose m} (\cos x)^{n-2(k-m)} \qquad \forall n \in \mathbb{W} \text{ and } x \in \mathbb{R}$$

$$\sin(nx) = \sum_{k=0}^{\left\lfloor \frac{n}{2} \right\rfloor} \sum_{m=0}^{k} (-1)^{k+m} {n \choose 2k} {k \choose m} (\sin x)^{n-2(k-m)} \qquad \forall n \in \mathbb{W} \text{ and } x \in \mathbb{R}$$

♥Proof:

$$\begin{aligned} \cos(nx) &= \Re \left(\cos nx + i \sin nx \right) \\ &= \Re \left(e^{inx} \right) \\ &= \Re \left[\left(e^{ix} \right)^n \right] \\ &= \Re \left[\left(\cos x + i \sin x \right)^n \right] \\ &= \Re \left[\left(\cos x + i \sin x \right)^n \right] \\ &= \Re \left[\sum_{k \in \mathbb{Z}} \binom{n}{k} (\cos x)^{n-k} x \sin^k x \right] \\ &= \Re \left[\sum_{k \in \mathbb{Z}} i^k \binom{n}{k} \cos^{n-k} x \sin^k x + i \sum_{k \in \{1,3,\dots,n\}} \binom{n}{k} \cos^{n-k} x \sin^k x \right] \\ &= \Re \left[\sum_{k \in \{0,4,\dots,n\}} \binom{n}{k} \cos^{n-k} x \sin^k x + i \sum_{k \in \{3,7,\dots,n\}} \binom{n}{k} \cos^{n-k} x \sin^k x \right] \\ &= \sum_{k \in \{0,4,\dots,n\}} \binom{n}{k} \cos^{n-k} x \sin^k x + -i \sum_{k \in \{3,7,\dots,n\}} \binom{n}{k} \cos^{n-k} x \sin^k x \right] \\ &= \sum_{k \in \{0,2,\dots,n\}} \binom{n}{k} (-1)^{\frac{k}{2}} \cos^{n-k} x \sin^k x \\ &= \sum_{k \in \{0,2,\dots,n\}} \binom{n}{k} (-1)^{\frac{k}{2}} \cos^{n-k} x \sin^k x \\ &= \sum_{k \in \{0,2,\dots,n\}} \binom{n}{k} (-1)^{k} \cos^{n-2k} x \sin^{2k} x \\ &= \sum_{k \in \{0,2,\dots,n\}} \binom{n}{k} (-1)^{k} \cos^{n-2k} x \sin^{2k} x \\ &= \sum_{k \in \{0,2,\dots,n\}} \binom{n}{2k} (-1)^{k} \cos^{n-2k} x \sin^{2k} x \\ &= \sum_{k \in \{0,2,\dots,n\}} \binom{n}{2k} (-1)^{k} \cos^{n-2k} x (1 - \cos^2 x)^{k} \\ &= \sum_{k \in \{0,2,\dots,n\}} \binom{n}{2k} (-1)^{k} \cos^{n-2k} x \left[\sum_{m = 0}^{k} \binom{k}{m} (-1)^{m} \cos^{2m} x \right] \\ &= \sum_{k = 0}^{\left\lfloor \frac{n}{2} \right\rfloor} \sum_{m = 0}^{k} (-1)^{k+m} \binom{n}{2k} \binom{k}{m} \cos^{n-2(k-m)} x \\ \sin(nx) &= \cos\left(nx - \frac{\pi}{2}\right) \\ &= \cos\left(n\left[x - \frac{\pi}{2n}\right]\right) \\ &= \sum_{k = 0}^{\left\lfloor \frac{n}{2} \right\rfloor} \sum_{m = 0}^{k} (-1)^{k+m} \binom{n}{2k} \binom{k}{m} \cos^{n-2(k-m)} \left(n\left[x - \frac{\pi}{2n}\right]\right) \end{aligned}$$

₽

$$= \sum_{k=0}^{\left\lfloor \frac{n}{2} \right\rfloor} \sum_{m=0}^{k} (-1)^{k+m} \binom{n}{2k} \binom{k}{m} \cos^{n-2(k-m)} \left(nx - \frac{\pi}{2} \right)$$

$$= \sum_{k=0}^{\left\lfloor \frac{n}{2} \right\rfloor} \sum_{m=0}^{k} (-1)^{k+m} \binom{n}{2k} \binom{k}{m} \sin^{n-2(k-m)} (nx)$$

Example H.1.



$$\cos 5x = 16\cos^5 x - 20\cos^3 x + 5\cos x$$

 $\sin 5x = 16\sin^5 x - 20\sin^3 x + 5\sin x$.

[♠]Proof:

1. Proof using *DeMoivre's Theorem* (Theorem H.1 page 173):

$$\begin{aligned} \cos 5x + i \sin 5x \\ &= e^{i5x} \\ &= (e^{ix})^5 \\ &= (\cos x + i \sin x)^5 \\ &= \sum_{k=0}^{5} {5 \choose k} [\cos x]^{5-k} [i \sin x]^k \\ &= {5 \choose 0} [\cos x]^{5-0} [i \sin x]^0 + {5 \choose 1} [\cos x]^{5-1} [i \sin x]^1 + {5 \choose 2} [\cos x]^{5-2} [i \sin x]^2 + \\ {5 \choose 3} [\cos x]^{5-3} [i \sin x]^3 + {5 \choose 4} [\cos x]^{5-4} [i \sin x]^4 + {5 \choose 5} [\cos x]^{5-5} [i \sin x]^5 \\ &= 1 \cos^5 x + i 5 \cos^4 x \sin x - 10 \cos^3 x \sin^2 x - i 10 \cos^2 x \sin^3 x + 5 \cos x \sin^4 x + i 1 \sin^5 x \\ &= [\cos^5 x - 10 \cos^3 x \sin^2 x + 5 \cos x \sin^4 x] + i [5 \cos^4 x \sin x - 10 \cos^2 x \sin^3 x + \sin^5 x] \\ &= [\cos^5 x - 10 \cos^3 x (1 - \cos^2 x) + 5 \cos x (1 - \cos^2 x) (1 - \cos^2 x)] + \\ i [5(1 - \sin^2 x)(1 - \sin^2 x) \sin x - 10(1 - \sin^2 x) \sin^3 x + \sin^5 x] \\ &= [\cos^5 x - 10 (\cos^3 x - \cos^5 x) + 5 \cos x (1 - 2\cos^2 x + \cos^4 x)] + \\ i [5(1 - 2\sin^2 x + \sin^4 x) \sin x - 10 (\sin^3 x - \sin^5 x) + \sin^5 x] \\ &= [\cos^5 x - 10 (\cos^3 x - \cos^5 x) + 5 (\cos x - 2\cos^3 x + \cos^5 x)] + \\ i [5(\sin x - 2\sin^3 x + \sin^5 x) - 10 (\sin^3 x - \sin^5 x) + \sin^5 x] \\ &= [16 \cos^5 x - 20 \cos^3 x + 5 \cos x] + i [16 \sin^5 x - 20 \sin^3 x + 5 \sin x] \\ \cos 5x &= \frac{16 \cos^5 x - 20 \cos^3 x + 5 \cos x}{\sin^5 x} + i [16 \sin^5 x - 20 \sin^3 x + 5 \sin x] \end{aligned}$$

2. Proof using trigonometric expansion (Theorem H.2 page 173):

$$\cos 5x = \sum_{k=0}^{\left\lfloor \frac{5}{2} \right\rfloor} \sum_{m=0}^{k} (-1)^{k+m} \binom{n}{2k} \binom{k}{m} (\cos x)^{n-2(k-m)}$$

$$= \sum_{k=0}^{2} \sum_{m=0}^{k} (-1)^{k+m} \binom{n}{2k} \binom{k}{m} (\cos x)^{5-2(k-m)}$$

$$= (-1)^{0} \binom{5}{0} \binom{0}{0} \cos^{5} x + (-1)^{1} \binom{5}{2} \binom{1}{0} \cos^{3} x + (-1)^{2} \binom{5}{2} \binom{1}{1} \cos^{5} x + (-1)^{2} \binom{5}{4} \binom{2}{0} \cos^{1} x + (-1)^{3} \binom{5}{4} \binom{2}{1} \cos^{3} x + (-1)^{4} \binom{5}{4} \binom{2}{2} \cos^{5} x$$

Frames and Bases Structure and Design [VERSION 021]






Figure H.1: Lattice of harmonic cosines $\{\cos(nx)|n=0,1,2,...\}$

$$= +(1)(1)\cos^5 x - (10)(1)\cos^3 x + (10)(1)\cos^5 x + (5)(1)\cos x - (5)(2)\cos^3 x + (5)(1)\cos^5 x$$

$$= +(1+10+5)\cos^5 x + (-10-10)\cos^3 x + 5\cos x$$

$$= 16\cos^5 x - 20\cos^3 x + 5\cos x$$

Example H.2. 3

			$\cos x \mid n$	cosnx	polynomial in cosx
		$\cos 0x = 1$		1	$8\cos^4 x - 8\cos^2 x + 1$
E X		$\cos 1x = \cos^1 x$	5	$\cos 5x =$	$= 16\cos^5 x - 20\cos^3 x + 5\cos x$
	2	$\cos 2x = 2\cos^2 x - 1$	6	$\cos 6x =$	$= 32\cos^6 x - 48\cos^4 x + 18\cos^2 x - 1$
		$\cos 3x = 4\cos^3 x - 3\cos x$	7	$\cos 7x =$	$= 64\cos^7 x - 112\cos^5 x + 56\cos^3 x - 7\cos x$

NPROOF:

$$\cos 2x = \sum_{k=0}^{\left\lfloor \frac{2}{2} \right\rfloor} \sum_{m=0}^{k} (-1)^{k+m} {3 \choose 2k} {k \choose m} (\cos x)^{2-2(k-m)}$$

$$= (-1)^{0} {3 \choose 0} {0 \choose 0} \cos^{2}x + (-1)^{1} {3 \choose 2} {1 \choose 0} \cos^{0}x + (-1)^{2} {3 \choose 2} {1 \choose 1} \cos^{2}x$$

$$= +(1)(1)\cos^{2}x - (1)(1) + (1)(1)\cos^{2}x$$

$$= 2\cos^{2}x - 1$$

$$\cos 3x = \sum_{k=0}^{\left\lfloor \frac{3}{2} \right\rfloor} \sum_{m=0}^{k} (-1)^{k+m} {3 \choose 2k} {k \choose m} (\cos x)^{3-2(k-m)}$$

$$= (-1)^0 {3 \choose 0} {0 \choose 0} \cos^3 x + (-1)^1 {3 \choose 2} {1 \choose 0} \cos^1 x + (-1)^2 {3 \choose 2} {1 \choose 1} \cos^3 x$$

$$= + {3 \choose 0} {0 \choose 0} \cos^3 x - {3 \choose 2} {1 \choose 0} \cos^1 x + {3 \choose 2} {1 \choose 1} \cos^3 x$$

$$= +(1)(1)\cos^3 x - (3)(1)\cos^1 x + (3)(1)\cos^3 x$$

$$= 4\cos^3 x - 3\cos x$$

$$\cos 4x = \sum_{k=0}^{\left\lfloor \frac{4}{2} \right\rfloor} \sum_{m=0}^{k} (-1)^{k+m} {4 \choose 2k} {k \choose m} (\cos x)^{4-2(k-m)}$$

³ Abramowitz and Stegun (1972) page 795, Guillemin (1957) page 593 \langle (21) \rangle , Sloane (2014) \langle http://oeis.org/A039991 \rangle , Sloane (2014) \langle http://oeis.org/A028297 \rangle



$$\begin{split} &= \sum_{k=0}^2 \sum_{m=0}^k (-1)^{k+m} \binom{4}{2k} \binom{k}{m} (\cos x)^{4-2(k-m)} \\ &= (-1)^{0+0} \binom{4}{2 \cdot 0} \binom{0}{0} (\cos x)^{4-2(0-0)} + (-1)^{1+0} \binom{4}{2 \cdot 1} \binom{1}{0} (\cos x)^{4-2(1-0)} \\ &\quad + (-1)^{1+1} \binom{4}{2 \cdot 1} \binom{1}{1} (\cos x)^{4-2(1-1)} + (-1)^{2+0} \binom{4}{2 \cdot 2} \binom{2}{0} (\cos x)^{4-2(2-0)} \\ &\quad + (-1)^{2+1} \binom{4}{2 \cdot 2} \binom{2}{1} (\cos x)^{4-2(2-1)} + (-1)^{2+2} \binom{4}{2 \cdot 2} \binom{2}{2} (\cos x)^{4-2(2-2)} \\ &= (1)(1) \cos^4 x - (6)(1) \cos^2 x + (6)(1) \cos^4 x + (1)(1) \cos^0 x - (1)(2) \cos^2 x + (1)(1) \cos^4 x \\ &= 8 \cos^4 x - 8 \cos^2 x + 1 \end{split}$$

 $\cos 5x = 16\cos^5 x - 20\cos^3 x + 5\cos x$ see Example H.1 page 175

$$\begin{aligned} \cos 6x &= \sum_{k=0}^{\left \lfloor \frac{6}{2} \right \rfloor} \sum_{m=0}^{k} (-1)^{k+m} \binom{6}{2k} \binom{k}{m} (\cos x)^{6-2(k-m)} \\ &= (-1)^0 \binom{6}{0} \binom{0}{0} \cos^6 x + (-1)^1 \binom{6}{2} \binom{1}{0} \cos^4 x + (-1)^2 \binom{6}{2} \binom{1}{1} \cos^6 x + (-1)^2 \binom{6}{4} \binom{2}{0} \cos^2 x + \\ &\quad (-1)^3 \binom{6}{4} \binom{2}{1} \cos^4 x + (-1)^4 \binom{6}{4} \binom{2}{2} \cos^6 x + (-1)^3 \binom{6}{6} \binom{3}{0} \cos^0 x + (-1)^4 \binom{6}{6} \binom{3}{1} \cos^2 x + \\ &\quad (-1)^5 \binom{6}{6} \binom{3}{2} \cos^4 x + (-1)^6 \binom{6}{6} \binom{3}{3} \cos^6 x \\ &= +(1)(1) \cos^6 x - (15)(1) \cos^4 x + (15)(1) \cos^6 x + (15)(1) \cos^2 x - (15)(2) \cos^4 x + (15)(1) \cos^6 x \\ &\quad - (1)(1) \cos^0 x + (1)(3) \cos^2 x - (1)(3) \cos^4 x + (1)(1) \cos^6 x \\ &= 32 \cos^6 x - 48 \cos^4 x + 18 \cos^2 x - 1 \end{aligned}$$

$$\cos 7x = \sum_{k=0}^{\left[\frac{7}{2}\right]} \sum_{m=0}^{k} (-1)^{k+m} {n \choose 2k} {k \choose m} (\cos x)^{n-2(k-m)}$$

$$= \sum_{k=0}^{3} \sum_{m=0}^{k} (-1)^{k+m} {n \choose 2k} {k \choose m} (\cos x)^{7-2(k-m)}$$

$$= (-1)^{0} {n \choose 0} {0 \choose 0} \cos^{7}x + (-1)^{1} {n \choose 2} {1 \choose 0} \cos^{5}x + (-1)^{2} {n \choose 2} {1 \choose 1} \cos^{7}x + (-1)^{2} {n \choose 4} {2 \choose 0} \cos^{3}x$$

$$+ (-1)^{3} {n \choose 4} {2 \choose 1} \cos^{5}x + (-1)^{4} {n \choose 4} {2 \choose 2} \cos^{7}x + (-1)^{3} {n \choose 6} {3 \choose 0} \cos^{1}x + (-1)^{4} {n \choose 6} {3 \choose 1} \cos^{3}x$$

$$+ (-1)^{5} {n \choose 6} {3 \choose 2} \cos^{5}x + (-1)^{6} {n \choose 6} {3 \choose 3} \cos^{7}x$$

$$= (1)(1)\cos^{7}x - (21)(1)\cos^{5}x + (21)(1)\cos^{7}x + (35)(1)\cos^{3}x$$

$$- (35)(2)\cos^{5}x + (35)(1)\cos^{7}x - (7)(1)\cos^{1}x + (7)(3)\cos^{3}x$$

$$- (7)(3)\cos^{5}x + (7)(1)\cos^{7}x$$

$$= (1 + 21 + 35 + 7)\cos^{7}x - (21 + 70 + 21)\cos^{5}x + (35 + 21)\cos^{3}x - (7)\cos^{1}x$$

$$= 64\cos^{7}x - 112\cos^{5}x + 56\cos^{3}x - 7\cos x$$




Figure H.2: Lattice of Chebyshev polynomials $\{T_n(x)|n=0,1,2,3\}$

Note: Trigonometric expansion of cos(nx) for particular values of n can also be performed with the free software package $Maxima^{TM}$ using the syntax illustrated to the right:

```
trigexpand(cos(2*x));
trigexpand(cos(3*x));
trigexpand(cos(4*x));
trigexpand(cos(5*x));
trigexpand(cos(6*x));
trigexpand(cos(7*x));
```

Definition H.1.

D E

The nth Chebyshev polynomial of the first kind is defined as $T_n(x) \triangleq \cos nx$ where $\cos x \triangleq x$

Theorem H.3. ⁵ Let $T_n(x)$ be a Chebyshev polynomial with $n \in \mathbb{W}$.

$$\begin{array}{ccc} T & n \text{ is even} & \Longrightarrow & T_n(x) \text{ is even.} \\ M & n \text{ is odd} & \Longrightarrow & T_n(x) \text{ is odd.} \end{array}$$

Example H.3. Let $T_n(x)$ be a *Chebyshev polynomial* with $n \in \mathbb{W}$.

$$T_0(x) = 1
T_1(x) = x
T_2(x) = 2x^2 - 1
T_3(x) = 4x^3 - 3x$$

$$T_0(x) = 1
T_2(x) = 8x^4 - 8x^2 + 1
T_5(x) = 16x^5 - 20x^3 + 5x
T_6(x) = 32x^6 - 48x^4 + 18x^2 - 1$$

№ Proof: Proof of these equations follows directly from Example H.2 (page 176).

H.2 Trigonometric reduction

Theorem H.2 (page 173) showed that $\cos nx$ can be expressed as a polynomial in $\cos x$. Conversely, Theorem H.4 (next) shows that a polynomial in $\cos x$ can be expressed as a linear combination of $(\cos nx)_{n\in\mathbb{Z}}$.

Theorem H.4 (trigonometric reduction).

⁵ ☐ Rivlin (1974) page 5 ⟨(1.13)⟩, ☐ Süli and Mayers (2003) page 242 ⟨Lemma 8.2⟩, ☐ Davidson and Donsig (2010) page 222 ⟨exercise 10.7.A(a)⟩



⁴ maxima pages 157–158 (10.5 Trigonometric Functions)

$$X = \operatorname{span} \left\{ \cos^n(2\pi x) | n = 0, 1, 2, 3 \right\}$$
scaling subspace

Figure H.3: Lattice of exponential cosines $\{\cos^n x | n = 0, 1, 2, 3\}$

$$\cos^{n} x = \frac{1}{2^{n}} \sum_{k=0}^{n} \binom{n}{k} \cos[(n-2k)x]$$

$$= \begin{cases} \frac{1}{2^{n}} \binom{n}{\frac{n}{2}} + \frac{1}{2^{n-1}} \sum_{k=0}^{\frac{n}{2}-1} \binom{n}{k} \cos[(n-2k)x] & \text{for } n \text{ even} \\ \frac{1}{2^{n-1}} \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} \binom{n}{k} \cos[(n-2k)x] & \text{for } n \text{ odd} \end{cases}$$

[♠]Proof:

$$\cos^{n} x = \left(\frac{e^{ix} + e^{-ix}}{2}\right)^{n}$$

$$= \mathbf{R}_{e} \left[\left(\frac{e^{ix} + e^{-ix}}{2}\right)^{n} \right]$$

$$= \mathbf{R}_{e} \left[\frac{1}{2^{n}} \sum_{k=0}^{n} \binom{n}{k} e^{i(n-k)x} e^{-ikx} \right]$$

$$= \mathbf{R}_{e} \left[\frac{1}{2^{n}} \sum_{k=0}^{n} \binom{n}{k} e^{i(n-2k)x} \right]$$

$$= \mathbf{R}_{e} \left[\frac{1}{2^{n}} \sum_{k=0}^{n} \binom{n}{k} (\cos[(n-2k)x] + i\sin[(n-2k)x]) \right]$$

$$= \mathbf{R}_{e} \left[\frac{1}{2^{n}} \sum_{k=0}^{n} \binom{n}{k} \cos[(n-2k)x] + i\frac{1}{2^{n}} \sum_{k=0}^{n} \binom{n}{k} \sin[(n-2k)x] \right]$$

$$= \frac{1}{2^{n}} \sum_{k=0}^{n} \binom{n}{k} \cos[(n-2k)x]$$

$$= \begin{cases} \frac{1}{2^{n}} \binom{n}{\frac{n}{2}} + \frac{1}{2^{n-1}} \sum_{k=0}^{\frac{n}{2}-1} \binom{n}{k} \cos[(n-2k)x] & : n \text{ even} \\ \frac{1}{2^{n-1}} \sum_{k=0}^{n} \binom{n}{k} \cos[(n-2k)x] & : n \text{ odd} \end{cases}$$

Example H.4. 6



 $^{^6}$ ⚠ Abramowitz and Stegun (1972) page 795, \trianglerighteq Sloane (2014) $\langle \texttt{http://oeis.org/A100257} \rangle$, \trianglerighteq Sloane (2014) $\langle \texttt{http://oeis.org/A008314} \rangle$

	n	$\cos^n x$	trigonometric reduction	n	$\cos^n x$		trigonometric reduction
	0	$\cos^0 x =$	1	4	$\cos^4 x$	=	$\frac{\cos 4x + 4\cos 2x + 3}{2^3}$
E X	1	$\cos^1 x =$	cosx	5	$\cos^5 x$	=	$\frac{2^3}{\cos 5x + 5\cos 3x + 10\cos x}$
	2	$\cos^2 x =$	$\frac{\cos 2x + 1}{2}$		l		$\frac{2^4}{\cos 6x + 6\cos 4x + 15\cos 2x + 10}$
	3	$\cos^3 x =$	$\frac{\cos 3x + 3\cos x}{2^2}$	7	$\cos^7 x$	=	$\frac{\cos 7x + 7\cos 5x + 21\cos 3x + 35\cos x}{2^6}$

♥Proof:

$$\cos^{0}x = \frac{1}{2^{n}} \sum_{k=0}^{n} {n \choose k} \cos([n-2k]x) \bigg|_{n=0}$$

$$= \frac{1}{2^{0}} \sum_{k=0}^{0} {n \choose k} \cos[(0-2k)x]$$

$$= {n \choose 0} \cos[(0-2\cdot 0)x]$$

$$= 1$$

$$\cos^{1}x = \frac{1}{2^{n}} \sum_{k=0}^{n} {n \choose k} \cos([n-2k]x) \bigg|_{n=1}$$

$$= \frac{1}{2^{1}} \sum_{k=0}^{1} {n \choose k} \cos[(1-2k)x]$$

$$= \frac{1}{2} \left[{n \choose 0} \cos[(1-2\cdot 0)x] + {n \choose 1} \cos[(1-2\cdot 1)x] \right]$$

$$= \frac{1}{2} \left[1\cos x + 1\cos(-x) \right]$$

$$= \frac{1}{2} \left[\cos x + \cos x \right]$$

$$= \cos x$$

$$\cos^{2}x = \frac{1}{2^{n}} \sum_{k=0}^{n} {n \choose k} \cos([n-2k]x) \bigg|_{n=2}$$

$$= \frac{1}{2^{2}} \sum_{k=0}^{2} {n \choose k} \cos([2-2k]x)$$

$$= \frac{1}{2^{2}} \left[{n \choose 0} \cos([2-2\cdot 0]x) + {n \choose 1} \cos([2-2\cdot 1]x) + {n \choose 2} \cos([2-2\cdot 2]x) + \right]$$

$$= \frac{1}{2^{2}} \left[1\cos(2x) + 2\cos(0x) + 1\cos(-2x) \right]$$

$$= \frac{1}{2^{2}} \left[\cos(2x) + 2 + \cos(2x) \right]$$

$$= \frac{1}{2} \left[\cos(2x) + 1 \right]$$

$$\cos^{3}x = \frac{1}{2^{n}} \sum_{k=0}^{n} {n \choose k} \cos([n-2k]x) \bigg|_{n=3}$$

$$= \frac{1}{2^{3}} \sum_{k=0}^{3} {n \choose k} \cos([3-2k]x)$$

$$\begin{aligned}
&= \frac{1}{2^3} \left[\log(3x) + 3\cos(1x) + 3\cos(-1x) + 1\cos(-3x) \right] \\
&= \frac{1}{2^3} \left[\cos(3x) + 3\cos(x) + 3\cos(x) + \cos(3x) \right] \\
&= \frac{1}{2^2} \left[\cos(3x) + 3\cos(x) \right] \\
&= \frac{1}{2^2} \left[\cos(3x) + 3\cos(x) \right] \\
&= \frac{1}{2^2} \sum_{k=0}^{n} \binom{n}{k} \cos\left(\left[n - 2k \right] x \right) \right|_{n=4} \\
&= \frac{1}{2^4} \sum_{k=0}^{4} \binom{4}{k} \cos\left(\left[4 - 2k \right] x \right) \\
&= \frac{1}{2^4} \left[1\cos(4x) + 4\cos(2x) + 6\cos(0x) + 4\cos(-2x) + 1\cos(-4x) \right] \\
&= \frac{1}{2^3} \left[\cos(4x) + 4\cos(2x) + 3 \right] \\
&\cos^5 x = \frac{1}{16} \sum_{k=0}^{1} \binom{5}{k} \cos\left(\left[5 - 2k \right) x \right] \\
&= \frac{1}{16} \left[\binom{5}{0} \cos 5x + \binom{5}{1} \cos 3x + \binom{5}{2} \cos x \right] \\
&= \frac{1}{16} \left[\cos 5x + 5\cos 3x + 10\cos x \right] \\
&\cos^6 x = \frac{1}{2^6} \binom{6}{6} + \frac{1}{2^{6-1}} \sum_{k=0}^{\frac{5}{2}-1} \binom{6}{k} \cos\left(\left[6 - 2k \right) x \right] \\
&= \frac{1}{6^4} 20 + \frac{1}{3^2} \left[\binom{6}{0} \cos 6x + \binom{6}{1} \cos 4x \binom{6}{2} \cos 2x \right] \\
&= \frac{1}{3^2} \left[\cos 6x + 6\cos 4x + 15\cos 2x + 10 \right] \\
&\cos^7 x = \frac{1}{2^{7-1}} \sum_{k=0}^{\frac{7}{2}} \binom{7}{k} \cos\left(\left[7 - 2k \right) x \right] \\
&= \frac{1}{6^4} \left[\binom{7}{0} \cos 7x + \binom{7}{1} \cos 5x + \binom{7}{2} \cos 3x + \binom{7}{3} \cos x \right] \\
&= \frac{1}{6^4} \left[\cos 7x + 7\cos 5x + 21\cos 3x + 35\cos x \right] \end{aligned}$$

Note: Trigonometric reduction of $\cos^n(x)$ for particular values of n can also be performed with the free software package $Maxima^{TM}$ using the syntax illustrated to the right:⁷

```
trigreduce((cos(x))^2);
trigreduce((cos(x))^3);
trigreduce((cos(x))^4);
trigreduce((cos(x))^5);
trigreduce((cos(x))^6);
trigreduce((cos(x))^7);
```



⁷ http://maxima.sourceforge.net/docs/manual/en/maxima_15.html maxima page 158 (10.5 Trigonometric Functions)

H.3 Spectral Factorization

Theorem H.5 (Fejér-Riesz spectral factorization). 8 Let $[0, \infty) \subsetneq \mathbb{R}$ and

$$p(e^{ix}) \triangleq \sum_{n=-N}^{N} a_n e^{inx}$$
 (Laurent trigonometric polynomial order 2N)
$$q(e^{ix}) \triangleq \sum_{n=1}^{N} b_n e^{inx}$$
 (standard trigonometric polynomial order N)

$$\begin{array}{c} \mathbf{T} \\ \mathbf{H} \\ \mathbf{M} \end{array} \mathbf{p} \left(e^{ix} \right) \in [0, \infty) \quad \forall x \in [0, 2\pi] \qquad \Longrightarrow \qquad \left\{ \begin{array}{c} \exists \, (b_n)_{n \in \mathbb{Z}} \quad \text{such that} \\ \mathbf{p} \left(e^{ix} \right) = \mathbf{q} \left(e^{ix} \right) \, \mathbf{q}^* \left(e^{ix} \right) & \forall x \in \mathbb{R} \end{array} \right.$$

♥Proof:

1. Proof that $a_n = a_{-n}^* \left((a_n)_{n \in \mathbb{Z}} \text{ is } Hermitian \ symmetric} \right)$: Let $a_n \triangleq r_n e^{i\phi_n}$, $r_n, \phi_n \in \mathbb{R}$. Then

$$\begin{split} & p\left(e^{inx}\right) \triangleq \sum_{n=-N}^{N} a_n e^{inx} \\ & = \sum_{n=-N}^{N} r_n e^{i\phi_n} e^{inx} \\ & = \sum_{n=-N}^{N} r_n e^{inx + \phi_n} \\ & = \sum_{n=-N}^{N} r_n \cos(nx + \phi_n) + i \sum_{n=-N}^{N} r_n \sin(nx + \phi_n) \\ & = \sum_{n=-N}^{N} r_n \cos(nx + \phi_n) + i \left[r_0 \sin(0x + \phi_0) + \sum_{n=1}^{N} r_n \sin(nx + \phi_n) + \sum_{n=1}^{N} r_{-n} \sin(-nx + \phi_{-n}) \right] \\ & = \sum_{n=-N}^{N} r_n \cos(nx + \phi_n) + i \left[r_0 \sin(\phi_0) + \sum_{n=1}^{N} r_n \sin(nx + \phi_n) - \sum_{n=1}^{N} r_{-n} \sin(nx - \phi_{-n}) \right] \\ & \Rightarrow r_n = r_{-n}, \ \phi_n = -\phi_{-n} \ \Rightarrow a_n = a_{-n}^*, \ a_0 \in \mathbb{R} \end{split}$$


2. Because the coefficients $(c_n)_{n\in\mathbb{Z}}$ are $Hermitian\ symmetric$, the zeros of P(z) occur in $conjugate\ reciprical pairs$. This means that if $\sigma\in\mathbb{C}$ is a zero of P(z) ($P(\sigma)=0$), then $\frac{1}{\sigma^*}$ is also a zero of P(z) (P(z)=0). In the complex z plane, this relationship means zeros are reflected across the unit circle such that

$$\frac{1}{\sigma^*} = \frac{1}{(re^{i\phi})^*} = \frac{1}{r} \frac{1}{e^{-i\phi}} = \frac{1}{r} e^{i\phi}$$

⁸ Pinsky (2002) pages 330–331



H.4. DIRICHLET KERNEL Daniel J. Greenhoe page 183



3. Because the zeros of p(z) occur in conjugate recipricol pairs, $p\left(e^{ix}\right)$ can be factored:

$$\begin{split} &\mathsf{p}\left(e^{ix}\right) = \mathsf{p}(z)|_{z=e^{ix}} \\ &= z^{-N} C \prod_{n=1}^{N} (z - \sigma_n) \prod_{n=1}^{N} \left(z - \frac{1}{\sigma_n^*}\right) \bigg|_{z=e^{ix}} \\ &= C \prod_{n=1}^{N} (z - \sigma_n) \prod_{n=1}^{N} z^{-1} \left(z - \frac{1}{\sigma_n^*}\right) \bigg|_{z=e^{ix}} \\ &= C \prod_{n=1}^{N} (z - \sigma_n) \prod_{n=1}^{N} \left(1 - \frac{1}{\sigma_n^*} z^{-1}\right) \bigg|_{z=e^{ix}} \\ &= C \prod_{n=1}^{N} (z - \sigma_n) \prod_{n=1}^{N} \left(z^{-1} - \sigma_n^*\right) \left(-\frac{1}{\sigma_n^*}\right) \bigg|_{z=e^{ix}} \\ &= \left[C \prod_{n=1}^{N} \left(-\frac{1}{\sigma_n^*}\right)\right] \left[\prod_{n=1}^{N} (z - \sigma_n)\right] \left[\prod_{n=1}^{N} \left(\frac{1}{z^*} - \sigma_n\right)\right]^* \bigg|_{z=e^{ix}} \\ &= \left[C_2 \prod_{n=1}^{N} (z - \sigma_n)\right] \left[C_2 \prod_{n=1}^{N} \left(\frac{1}{z^*} - \sigma_n\right)\right]^* \bigg|_{z=e^{ix}} \\ &= \mathsf{q}(z) \mathsf{q}^* \left(\frac{1}{z^*}\right) \bigg|_{z=e^{ix}} \\ &= \mathsf{q}\left(e^{ix}\right) \mathsf{q}^* \left(e^{ix}\right) \end{split}$$

H.4 Dirichlet Kernel



Dirichlet alone, not I, nor Cauchy, nor Gauss knows what a completely rigorous proof is. Rather we learn it first from him. When Gauss says he has proved something it is clear; when Cauchy says it, one can wager as much pro as con; when Dirichlet says it, it is certain.

Carl Gustav Jacob Jacobi (1804–1851), Jewish-German mathematician ⁹

image: http://en.wikipedia.org/wiki/File:Carl_Jacobi.jpg, public domain





The Dirichlet Kernel is critical in proving what is not immediately obvious in examining the Fourier Series—that for a broad class of periodic functions, a function can be recovered from (with uniform convergence) its Fourier Series analysis.

Definition H.2. ¹⁰

E

The **Dirichlet Kernel**
$$D_n \in \mathbb{R}^{\mathbb{W}}$$
 with period τ is defined as

Daniel J. Greenhoe

$$\mathsf{D}_n(x) \triangleq \frac{1}{\tau} \sum_{k=-n}^n e^{i\frac{2\pi}{\tau}kx}$$

Proposition H.1. 11 Let D_n be the DIRICHLET KERNEL with period τ (Definition H.2 page 184).

$$D_n(x) = \frac{1}{\tau} \frac{\sin\left(\frac{\pi}{\tau}[2n+1]x\right)}{\sin\left(\frac{\pi}{\tau}x\right)}$$

^ℚProof:

$$\begin{split} \mathsf{D}_{n}(x) &\triangleq \frac{1}{\tau} \sum_{k=-n}^{n} e^{i\frac{2\pi}{\tau}nx} & \text{by definition of } \mathsf{D}_{n} \\ &= \frac{1}{\tau} \sum_{k=0}^{2n} e^{i\frac{2\pi}{\tau}(k-n)x} = \frac{1}{\tau} e^{-i\frac{2\pi n}{\tau}x} \sum_{k=0}^{2n} e^{i\frac{2\pi}{\tau}kx} = \frac{1}{\tau} e^{-i\frac{2\pi}{\tau}nx} \sum_{k=0}^{2n} \left(e^{i\frac{2\pi}{\tau}x} \right)^{k} \\ &= \frac{1}{\tau} e^{-i\frac{2\pi n}{\tau}x} \frac{1 - \left(e^{i\frac{2\pi}{\tau}x} \right)^{2n+1}}{1 - e^{i\frac{2\pi}{\tau}x}} & \text{by geometric series} \\ &= \frac{1}{\tau} e^{-i\frac{2\pi}{\tau}nx} \frac{1 - e^{i\frac{2\pi}{\tau}(2n+1)x}}{1 - e^{i\frac{2\pi}{\tau}(2n+1)x}} = \frac{1}{\tau} e^{-i\frac{2\pi n}{\tau}x} \left(\frac{e^{i\frac{\pi}{\tau}(2n+1)x}}{e^{i\frac{\pi}{\tau}x}} \right) \frac{e^{-i\frac{\pi}{\tau}(2n+1)x} - e^{i\frac{\pi}{\tau}(2n+1)x}}{e^{-i\frac{\pi}{\tau}x} - e^{i\frac{\pi}{\tau}x}} \\ &= \frac{1}{\tau} e^{-i\frac{2\pi n}{\tau}x} \left(e^{i\frac{2\pi n}{\tau}x} \right) \frac{-2i\sin\left[\frac{\pi}{\tau}(2n+1)x\right]}{-2i\sin\left[\frac{\pi}{\tau}x\right]} = \frac{1}{\tau} \frac{\sin\left[\frac{\pi}{\tau}(2n+1)x\right]}{\sin\left[\frac{\pi}{\tau}x\right]} \end{split}$$

Proposition H.2. 12 Let D_n be the DIRICHLET KERNEL with period τ (Definition H.2 page 184).

$$\int_{0}^{\tau} \mathsf{D}_{n}(x) \, \mathsf{dx} = 1$$

^ℚProof:

$$\begin{split} \int_0^\tau \mathsf{D}_n(x) \, \mathrm{d} \mathbf{x} &\triangleq \int_0^\tau \frac{1}{\tau} \sum_{k=-n}^n e^{i\frac{2\pi}{\tau}nx} \, \mathrm{d} \mathbf{x} \\ &= \frac{1}{\tau} \sum_{k=-n}^n \int_{-\frac{\tau}{2}}^{\frac{\tau}{2}} e^{i\frac{2\pi}{\tau}nx} \, \mathrm{d} \mathbf{x} \\ &= \frac{1}{\tau} \sum_{k=-n}^n \int_{-\frac{\tau}{2}}^{\frac{\tau}{2}} \cos\left(\frac{2\pi}{\tau}nx\right) + i \sin\left(\frac{2\pi}{\tau}nx\right) \, \mathrm{d} \mathbf{x} \end{split}$$

¹² ■ Bruckner et al. (1997) pages 620–621






Figure H.4: D_n function for N = 1, 5, 10, 20. $D_n \rightarrow \text{comb.}$ (See Proposition H.1 page 184).

$$= \frac{1}{\tau} \sum_{k=-n}^{n} \int_{-\frac{\tau}{2}}^{\frac{\tau}{2}} \cos\left(\frac{2\pi}{\tau} n x\right) dx$$

$$= \frac{1}{\tau} \sum_{k=-n}^{n} \frac{\sin\left(\frac{2\pi}{\tau} n x\right)}{\frac{2\pi}{\tau} n} \Big|_{-\frac{\tau}{2}}^{\frac{\tau}{2}}$$

$$= \frac{1}{\tau} \sum_{k=-n}^{n} \left[\frac{\sin\left(\frac{2\pi}{\tau} n \frac{\tau}{2}\right)}{\frac{2\pi}{\tau} n} - \frac{\sin\left(-\frac{2\pi}{\tau} n \frac{\tau}{2}\right)}{\frac{2\pi}{\tau} n} \right]$$

$$= \frac{1}{\tau} \frac{\tau}{2} \sum_{k=-n}^{n} \left[\frac{\sin(\pi n)}{\pi n} + \frac{\sin(\pi n)}{\pi n} \right]$$

$$= \frac{1}{2} \left[2 \frac{\sin(\pi n)}{\pi n} \right]_{k=0}$$

$$= 1$$

Proposition H.3. Let D_n be the DIRICHLET KERNEL with period τ (Definition H.2 page 184). Let w_N (the "WIDTH" of $D_n(x)$) be the distance between the two points where the center pulse of $D_n(x)$ intersects the x axis.

$$\begin{array}{ccc}
 & D_n(0) & = \frac{1}{\tau}(2n+1) \\
 & w_n & = \frac{2\tau}{2n+1}
\end{array}$$

Frames and Bases Structure and Design [VERSON 021] https://github.com/dgreenhoe/pdfs/blob/master/msdframes.pdf



♥Proof:

$$\begin{split} \mathsf{D}_n(0) &= \left. \mathsf{D}_n(x) \right|_{t=0} \\ &= \frac{1}{\tau} \frac{\sin \left[\frac{\pi}{\tau} (2n+1)x \right]}{\sin \left[\frac{\pi}{\tau} t \right]} \bigg|_{t=0} \\ &= \frac{1}{\tau} \frac{\frac{\mathsf{d}}{\mathsf{d} \mathsf{x}} \sin \left[\frac{\pi}{\tau} (2n+1)x \right]}{\frac{\mathsf{d}}{\mathsf{d} \mathsf{x}} \sin \left[\frac{\pi}{\tau} t \right]} \bigg|_{t=0} \\ &= \frac{1}{\tau} \frac{\frac{\pi}{\tau} (2n+1)}{\frac{\pi}{\tau}} \frac{\cos \left[\frac{\pi}{\tau} (2n+1)x \right]}{\cos \left[\frac{\pi}{\tau} t \right]} \bigg|_{t=0} \\ &= \frac{1}{\tau} \frac{\frac{\pi}{\tau} (2n+1)}{\frac{\pi}{\tau}} \frac{1}{1} \\ &= \frac{1}{\tau} (2n+1) \end{split}$$

by Proposition H.1 page 184

by l'Hôpital's rule

The center pulse of kernel $D_n(x)$ intersects the x axis at

$$t = \pm \frac{\tau}{(2n+1)}$$

which implies

$$w_n = \frac{\tau}{2n+1} + \frac{\tau}{2n+1} = \frac{2\tau}{(2n+1)}.$$

Proposition H.4. ¹³ Let D_n be the DIRICHLET KERNEL with period τ (Definition H.2 page 184).

$$\mathsf{D}_n(x) = \mathsf{D}_n(-x)$$

 $D_n(x) = D_n(-x)$ (D_n is an even function)

[♠]Proof:

$$D_n(x) = \frac{1}{\tau} \frac{\sin\left[\frac{\pi}{\tau}(2n+1)x\right]}{\sin\left[\frac{\pi}{\tau}t\right]}$$

$$= \frac{1}{\tau} \frac{-\sin\left[-\frac{\pi}{\tau}(2n+1)x\right]}{-\sin\left[-\frac{\pi}{\tau}t\right]}$$

$$= \frac{1}{\tau} \frac{\sin\left[\frac{\pi}{\tau}(2n+1)(-x)\right]}{\sin\left[\frac{\pi}{\tau}(-x)\right]}$$

$$= D_n(-x)$$

by Proposition H.1 page 184

because sinx is an *odd* function

by Proposition H.1 page 184

¹³ Bruckner et al. (1997) pages 620–621



Trigonometric summations H.5

 $\sum_{n=0}^{N-1} \cos(nx) = \frac{1}{2} + \frac{\sin(\left[N - \frac{1}{2}\right]x)}{2\sin(\frac{1}{2}x)} = \frac{\sin(\left[N - \frac{1}{2}\right]x) + \sin(\frac{1}{2}x)}{2\sin(\frac{1}{2}x)}$ $\sum_{n=0}^{N-1} \sin(nx) = \frac{1}{2}\cot(\frac{1}{2}x) + \frac{\cos(\left[N - \frac{1}{2}\right]x)}{2\sin(\frac{1}{2}x)} = \frac{\cos(\left[N - \frac{1}{2}\right]x) + \cos(\frac{1}{2}x)}{2\sin(\frac{1}{2}x)}$ $\forall x \in \mathbb{R}$

[♠]Proof:

$$\sum_{n=0}^{N-1} \cos(nx) = \sum_{n=0}^{N-1} \Re e^{inx} = \Re \sum_{n=0}^{N-1} e^{inx} = \Re \sum_{n=0}^{N-1} \left(e^{ix} \right)^n$$

$$= \Re \left[\frac{1 - e^{iNx}}{1 - e^{ix}} \right] \qquad \text{by geometric series}$$

$$= \Re \left[\left(\frac{e^{i\frac{1}{2}Nx}}{e^{i\frac{1}{2}x}} \right) \left(\frac{e^{-i\frac{1}{2}Nx} - e^{i\frac{1}{2}Nx}}{e^{-i\frac{1}{2}x} - e^{i\frac{1}{2}x}} \right) \right]$$

$$= \Re \left[\left(e^{i\frac{1}{2}(N-1)x} \right) \left(\frac{-i\frac{1}{2}\sin\left(\frac{1}{2}Nx\right)}{-i\frac{1}{2}\sin\left(\frac{1}{2}x\right)} \right) \right]$$

$$= \cos\left(\frac{1}{2}(N-1)x \right) \left(\frac{\sin\left(\frac{1}{2}Nx\right)}{\sin\left(\frac{1}{2}x\right)} \right)$$

$$= \frac{-\frac{1}{2}\sin\left(-\frac{1}{2}x\right) + \frac{1}{2}\sin\left(\left[N-\frac{1}{2}x\right]x\right)}{\sin\left(\frac{1}{2}x\right)}$$
by product identities

(Theorem G.8 page 162)
$$= \frac{1}{2} + \frac{\sin\left(\left[N-\frac{1}{2}\right]x\right)}{2\sin\left(\frac{1}{2}x\right)}$$

$$\sum_{n=0}^{N-1} \sin(nx) = \sum_{n=0}^{N-1} \mathfrak{F}e^{inx} = \mathfrak{F}\sum_{n=0}^{N-1} e^{inx} = \mathfrak{F}\sum_{n=0}^{N-1} \left(e^{ix}\right)^n$$


$$= \mathfrak{F}\left[\frac{1 - e^{iNx}}{1 - e^{ix}}\right] \qquad \text{by geometric series}$$

$$= \mathfrak{F}\left[\left(\frac{e^{i\frac{1}{2}Nx}}{e^{i\frac{1}{2}x}}\right)\left(\frac{e^{-i\frac{1}{2}Nx} - e^{i\frac{1}{2}Nx}}{e^{-ix/2} - e^{i\frac{1}{2}x}}\right)\right]$$

$$= \mathfrak{F}\left[\left(e^{i(N-1)x/2}\right)\left(\frac{-\frac{1}{2}i\sin\left(\frac{1}{2}Nx\right)}{-\frac{1}{2}i\sin\left(\frac{1}{2}x\right)}\right)\right]$$

¹⁴ Muniz (1953) page 140 ⟨"Lagrange's Trigonometric Identities"⟩,

Jeffrey and Dai (2008) pages 128–130 ⟨2.4.1.6 Sines, Cosines, and Tagents of Multiple Angles; (14), (13)





$$= \sin\left(\frac{(N-1)x}{2}\right) \left(\frac{\sin\left(\frac{1}{2}Nx\right)}{\sin\left(\frac{1}{2}x\right)}\right)$$

$$= \frac{\frac{1}{2}\cos\left(-\frac{1}{2}x\right) - \frac{1}{2}\cos\left(\left[N-\frac{1}{2}\right]x\right)}{\sin\left(\frac{1}{2}x\right)}$$
by product identities (Theorem G.8 page 162)
$$= \frac{1}{2}\cot\left(\frac{1}{2}x\right) + \frac{\cos\left(\left[N-\frac{1}{2}\right]x\right)}{2\sin\left(\frac{1}{2}x\right)}$$

Note that these results (summed with indices from n = 0 to n = N - 1) are compatible with $\underline{\mathbb{R}}$ Muniz (1953) page 140 (summed with indices from n = 1 to n = N) as demonstrated next:

$$\sum_{n=0}^{N-1} \cos(nx) = \sum_{n=1}^{N} \cos(nx) + [\cos(0x) - \cos(Nx)]$$

$$= \left[-\frac{1}{2} + \frac{\sin(\left[N + \frac{1}{2}x\right]x)}{2\sin(\frac{1}{2}x)} \right] + [\cos(0x) - \cos(Nx)] \qquad \text{by } \text{ Muniz (1953) page 140}$$

$$= \left(1 - \frac{1}{2} \right) + \frac{\sin(\left[N + \frac{1}{2}x\right]x) - 2\sin(\frac{1}{2}x)\cos(Nx)}{2\sin(\frac{1}{2}x)}$$

$$= \frac{1}{2} + \frac{\sin(\left[N + \frac{1}{2}x\right]x) - 2\left[\sin(\left[\frac{1}{2} - N\right]x\right) + \sin\left[\left(\frac{1}{2} + N\right)x\right]\right]}{2\sin(\frac{1}{2}x)} \qquad \text{by Theorem G.8 page 162}$$

$$= \frac{1}{2} + \frac{\sin(\frac{1}{2}(2N - 1)x)}{2\sin(\frac{1}{2}x)} \qquad \implies \text{above result}$$

$$\sum_{n=0}^{N-1} \sin(nx) = \sum_{n=1}^{N} \sin(nx) + [\sin(0x) - \sin(Nx)]$$

$$= \frac{1}{2} \cot(\frac{1}{2}x) - \frac{\cos(\left[N + \frac{1}{2}x\right]x)}{2\sin(\frac{1}{2}x)} + [0 - \sin(Nx)] \qquad \text{by } \text{ Muniz (1953) page 140}$$

$$= \frac{1}{2} \cot(\frac{1}{2}x) - \frac{\cos(\left[N + \frac{1}{2}x\right]x) - 2\sin(\frac{1}{2}x)\sin(Nx)}{2\sin(\frac{1}{2}x)}$$

$$= \frac{1}{2} \cot(\frac{1}{2}x) - \frac{\cos(\left[N + \frac{1}{2}x\right]x) - \left[\cos(\left[\frac{1}{2} - N\right]x) - \cos(\left[\frac{1}{2} + N\right]x)\right]}{2\sin(\frac{1}{2}x)}$$

$$= \frac{1}{2} \cot(\frac{1}{2}x) + \frac{\cos(\left[N - \frac{1}{2}x\right]x)}{2\sin(\frac{1}{2}x)} \qquad \implies \text{above result}$$

Theorem H.7. ¹⁵

¹⁵ Jeffrey and Dai (2008) pages $128-130 \langle 2.4.1.6 \rangle$ Sines, Cosines, and Tagents of Multiple Angles; (16) and (17)



[♠]Proof:

$$\sum_{n=0}^{N-1} \cos(nx + y) = \sum_{n=0}^{N-1} \left[\cos(nx)\cos(y) - \sin(nx)\sin(y) \right]$$
 by double angle formulas (Theorem G.9 page 165)
$$= \cos(y) \sum_{n=0}^{N-1} \cos(nx) - \sin(y) \sum_{n=0}^{N-1} \sin(nx)$$

$$\sum_{n=0}^{N-1} \sin(nx + y) = \sum_{n=0}^{N-1} \left[\cos(nx)\cos(y) + \sin(nx)\sin(y) \right]$$
 by double angle formulas (Theorem G.9 page 165)
$$= \cos(y) \sum_{n=0}^{N-1} \cos(nx) + \sin(y) \sum_{n=0}^{N-1} \sin(nx)$$

Corollary H.1 (Summation around unit circle).

$$\sum_{n=0}^{N-1} \cos\left(\theta + \frac{2nM\pi}{N}\right) = \sum_{n=0}^{N-1} \sin\left(\theta + \frac{2nM\pi}{N}\right) = \sum_{n=0}^{N-1} \cos\left(\theta + \frac{2nM\pi}{N}\right) \sin\left(\theta + \frac{2nM\pi}{N}\right) = 0 \quad \forall \theta \in \mathbb{R}$$

$$\forall \theta \in \mathbb{R}$$

[♠]Proof:

$$\begin{split} &\sum_{n=0}^{N-1} \cos\left(\theta + \frac{2nM\pi}{N}\right) \\ &= \cos(\theta) \sum_{n=0}^{N-1} \cos\left(\frac{2nM\pi}{N}\right) - \sin(\theta) \sum_{n=0}^{N-1} \sin\left(\frac{2nM\pi}{N}\right) \\ &= \cos(\theta) \left[\frac{1}{2} + \frac{\sin\left(\left[N - \frac{1}{2}\right]\frac{2M\pi}{N}\right)}{2\sin\left(\frac{1}{2}\frac{2M\pi}{N}\right)}\right] - \sin(\theta) \left[\frac{1}{2}\cot\left(\frac{1}{2}\frac{2M\pi}{N}\right) + \frac{\cos\left(\left[N - \frac{1}{2}\right]\frac{2M\pi}{N}\right)}{2\sin\left(\frac{1}{2}\frac{2M\pi}{N}\right)}\right] \quad \text{by Theorem H.6 page 187} \\ &= \cos(\theta) \left[\frac{1}{2} - \frac{\sin\left(\frac{M\pi}{N} - 2M\pi\right)}{2\sin\left(\frac{M\pi}{N}\right)}\right] - \sin(\theta) \left[\frac{1}{2}\cot\left(\frac{M\pi}{N}\right) - \frac{\cos\left(\frac{M\pi}{N} - 2M\pi\right)}{2\sin\left(\frac{M\pi}{N}\right)}\right] \\ &= \cos(\theta) \left[\frac{1}{2} - \frac{1}{2}\frac{\sin\left(\frac{M\pi}{N}\right)}{\sin\left(\frac{M\pi}{N}\right)}\right] - \sin(\theta) \left[\frac{1}{2}\cot\left(\frac{M\pi}{N}\right) - \frac{1}{2}\cot\left(\frac{M\pi}{N}\right)\right] \quad \text{by trigonometric periodicity} \\ &= \cos(\theta)[0] - \sin(\theta)[0] \\ &= 0 \end{split}$$

$$\sum_{n=0}^{N-1} \sin\left(\theta + \frac{2nM\pi}{N}\right) = \sum_{n=0}^{N-1} \cos\left(\theta - \frac{\pi}{2} + \frac{2nM\pi}{N}\right) \qquad \text{by shift identities} \qquad \text{(Theorem G.7 page 162)}$$

$$= \sum_{n=0}^{N-1} \cos\left(\phi + \frac{2nM\pi}{N}\right) \qquad \text{where } \phi \triangleq \theta - \frac{\pi}{2}$$

$$= 0 \qquad \qquad \text{by previous result}$$

$$\begin{split} &\sum_{n=0}^{N-1} \cos\left(\theta + \frac{2nM\pi}{N}\right) \sin\left(\theta + \frac{2nM\pi}{N}\right) \\ &= -\frac{1}{2} \sum_{n=0}^{N-1} \sin\left(\left[\theta + \frac{2nM\pi}{N}\right] - \left[\theta + \frac{2nM\pi}{N}\right]\right) + \frac{1}{2} \sum_{n=0}^{N-1} \sin\left(\left[\theta + \frac{2nM\pi}{N}\right] + \left[\theta + \frac{2nM\pi}{N}\right]\right) \quad \text{by Theorem G.8 page 162} \\ &= -\frac{1}{2} \sum_{n=0}^{N-1} \sin(\theta)^n + \frac{1}{2} \sum_{n=0}^{N-1} \sin\left(2\theta + \frac{4nM\pi}{N}\right) \\ &= \frac{1}{2} \sin(2\theta) \sum_{n=0}^{N-1} \sin\left(\frac{4nM\pi}{N}\right) + \frac{1}{2} \cos(2\theta) \sum_{n=0}^{N-1} \sin\left(\frac{4nM\pi}{N}\right) \\ &= \cos(2\theta) \left[\frac{1}{2} + \frac{\sin\left(\left[N - \frac{1}{2}\right] \frac{4M\pi}{N}\right)}{2\sin\left(\frac{1}{2} \frac{2M\pi}{N}\right)}\right] - \sin(2\theta) \left[\frac{1}{2} \cot\left(\frac{1}{2} \frac{4M\pi}{N}\right) + \frac{\cos\left(\left[N - \frac{1}{2}\right] \frac{4M\pi}{N}\right)}{2\sin\left(\frac{1}{2} \frac{4M\pi}{N}\right)}\right] \quad \text{by Theorem H.6 page 187} \\ &= \cos(2\theta) \left[\frac{1}{2} - \frac{\sin\left(\frac{2M\pi}{N} - 4M\pi\right)}{2\sin\left(\frac{2M\pi}{N}\right)}\right] - \sin(2\theta) \left[\frac{1}{2} \cot\left(\frac{2M\pi}{N}\right) - \frac{\cos\left(\frac{2M\pi}{N} - 4M\pi\right)}{2\sin\left(\frac{2M\pi}{N}\right)}\right] \\ &= \cos(\theta) \left[\frac{1}{2} - \frac{1}{2} \frac{\sin\left(\frac{2M\pi}{N}\right)}{\sin\left(\frac{2M\pi}{N}\right)}\right] - \sin(\theta) \left[\frac{1}{2} \cot\left(\frac{2M\pi}{N}\right) - \frac{1}{2} \cot\left(\frac{2M\pi}{N}\right)\right] \quad \text{by trigonometric periodicity} \\ &= \cos(\theta) [0] - \sin(\theta) [0] \\ &= 0 \end{split}$$

$$\sum_{n=0}^{N-1} \cos^2\left(\theta + \frac{2nM\pi}{N}\right) = \frac{1}{2} \sum_{n=0}^{N-1} \left[1 + \cos\left(2\theta + \frac{4nM\pi}{N}\right)\right]$$
by Theorem G.11 page 167
$$= \frac{1}{2} \sum_{n=0}^{N-1} \left[1 + \cos(2\theta)\cos\left(\frac{4nM\pi}{N}\right) - \sin(2\theta)\sin\left(\frac{4nM\pi}{N}\right)\right]$$
by Theorem G.9 page 165
$$= \frac{1}{2} \sum_{n=0}^{N-1} 1 + \frac{1}{2}\cos(2\theta) \sum_{n=0}^{N-1} \cos\left(\frac{4nM\pi}{N}\right) - \frac{1}{2}\sin(2\theta) \sum_{n=0}^{N-1} \sin\left(\frac{4nM\pi}{N}\right)$$

$$= \left[\frac{1}{2} \sum_{n=0}^{N-1} 1\right] + \frac{1}{2}\cos(2\theta)0 - \frac{1}{2}\sin(2\theta)0$$
by previous results
$$= \frac{N}{2}$$

$$\sum_{n=0}^{N-1} \sin^2\left(\theta + \frac{2nM\pi}{N}\right) = \sum_{n=0}^{N-1} \cos^2\left(\theta - \frac{\pi}{2} + \frac{2nM\pi}{N}\right)$$
 by shift identities (Theorem G.7 page 162)
$$= \sum_{n=0}^{N-1} \cos^2\left(\phi + \frac{2nM\pi}{N}\right)$$
 where $\phi \triangleq \theta - \frac{\pi}{2}$

$$= \frac{N}{2}$$
 by previous result

Summability Kernels H.6

Definition H.3. ¹⁶ Let $(\kappa_n)_{n\in\mathbb{Z}}$ be a sequence of CONTINUOUS 2π PERIODIC functions. The sequence $(\kappa_n)_{n\in\mathbb{Z}}$ is a **summability kernel** if 1. $\frac{1}{2\pi} \int_{0}^{2\pi} \kappa_{n}(x) dx = 1 \quad \forall n \in \mathbb{Z} \quad and$ 2. $\frac{1}{2\pi} \int_{0}^{2\pi} \left| \kappa_{n}(x) \right| dx \in \mathbb{R} \quad \forall n \in \mathbb{Z} \quad and$ 3. $\lim_{n \to \infty} \int_{\delta}^{2\pi - \delta} \left| \kappa_{n}(x) \right| dx = 0 \quad \forall n \in \mathbb{Z}, 0 < \delta < \pi$

Theorem H.8. ¹⁷ Let $(\kappa_n)_{n\in\mathbb{Z}}$ be a sequence. Let \mathbb{T} be the quotient $\mathbb{R}/2\pi\mathbb{Z}$.

1. $f \in L^1(\mathbb{T})$ \Rightarrow $f(x) = \lim_{n \to \infty} \frac{1}{2\pi} \int_{\mathbb{T}} \kappa_n(x) f(x-x) dx$


The *Dirichlet kernel* (Definition H.2 page 184) is *not* a *summability kernel*. Examples of kernels that *are* summability kernels include

1. Fejér's kernel (Definition H.4 page 191) de la Vallée Poussin kernel (Definition H.5 page 193) Jackson kernel (Definition H.6 page 193) 4. Poisson kernel (Definition H.7 page 193.)

Definition H.4. 18

Fejér's kernel K_n is defined as DEF $K_n(x) \triangleq \sum_{k=-\infty}^{k=n} \left(1 - \frac{|k|}{n+1}\right) e^{ikx}$

Proposition H.5. ¹⁹ Let K_n be Fejér's kernel (Definition H.4 page 191).



¹⁶ ☑ Cerdà (2010) page 56, ❷ Katznelson (2004) page 10, ❷ de Reyna (2002) page 21, ❷ Walnut (2002) pages 40–41, Heil (2011) page 440,
Istrăţescu (1987) page 309



¹⁷ Katznelson (2004) page 11

¹⁸ Katznelson (2004) page 12

¹⁹ ■ Katznelson (2004) page 12, Heil (2011) page 448

№PROOF:

1. Lemma: Proof that $\sin^2 \frac{x}{2} \equiv \frac{-1}{4} (e^{-ix} - 2 + e^{ix})$:

$$\sin^{2} \frac{x}{2} \equiv \left(\frac{e^{-i\frac{x}{2}} - e^{+i\frac{x}{2}}}{2i}\right)^{2}$$
 by Euler Formulas (Corollary G.2 page 161)

$$\equiv \frac{-1}{4} \left(e^{-2i\frac{x}{2}} - 2e^{-i\frac{x}{2}}e^{i\frac{x}{2}} + e^{2i\frac{x}{2}}\right)$$

$$\equiv \frac{-1}{4} \left(e^{-ix} - 2 + e^{ix}\right) :$$

2. Lemma:

$$2|k|-|k+1|-|k-1| = \begin{cases} -2 & \text{for } k=0\\ 0 & \text{for } k \in \mathbb{Z} \backslash 0 \end{cases}$$

3. Proof that
$$K_n(x) = \frac{1}{n+1} \left(\frac{\sin\frac{n+1}{2}x}{\sin\frac{1}{2}x}\right)^2$$
:
$$-4(n+1)\left(\sin\frac{1}{2}x\right)^2 K_n(x)$$

$$= -4(n+1)\left(\frac{-1}{4}\right)\left(e^{-ix} - 2 + e^{ix}\right)K_n(x) \quad \text{by item (1)}$$

$$= (n+1)\left(e^{-ix} - 2 + e^{ix}\right) \sum_{k=-n}^{k=n} \left(1 - \frac{|k|}{n+1}\right)e^{ikx} \quad \text{by Definition II.4}$$

$$= (n+1)\frac{1}{n+1}\left(e^{-ix} - 2 + e^{ix}\right) \sum_{k=-n}^{k=n} (n+1-|k|)e^{ikx}$$

$$= e^{-ix} \sum_{k=-n}^{k=n} (n+1-|k|)e^{ikx} - 2 \sum_{k=-n}^{k=n} (n+1-|k|)e^{ikx}e^{ix} \sum_{k=-n}^{k=n} (n+1-|k|)e^{ikx}$$

$$= \sum_{k=-n}^{k=n} (n+1-|k|)e^{i(k-1)x} - 2 \sum_{k=-n}^{k=n} (n+1-|k|)e^{ikx} \sum_{k=-n}^{k=n} (n+1-|k|)e^{i(k+1)x}$$

$$= \sum_{k=-n-1}^{k=n-1} (n+1-|k|+1)e^{ikx} - 2 \sum_{k=-n}^{k=n} (n+1-|k|)e^{ikx} \sum_{k=-n+1}^{k=n+1} (n+1-|k-1|)e^{ikx}$$

$$= e^{-i(n+1)x} + 2e^{-inx} + \sum_{k=-n+1}^{k=n-1} (n+1-|k-1|)e^{ikx} + 2e^{-inx} + \sum_{k=-n+1}^{k=n-1} (n+1-|k-1|)e^{ikx}$$

$$= e^{-i(n+1)x} + \sum_{k=-n+1}^{k=n-1} (n+1-|k-1|)e^{ikx} + 2e^{-inx} + \sum_{k=-n+1}^{k=n-1} (n+1-|k-1|)e^{ikx} + 2e^{-inx} + \sum_{k=-n+1}^{k=n-1} (n+1-|k-1|)e^{ikx} + 2e^{-inx} + \sum_{k=-n+1}^{k=n-1} (n+1-|k-1|)e^{-ikx} + 2e^{-inx} + 2e^{-i$$

$$= e^{-i(n+1)x} + e^{i(n+1)x} + \sum_{k=-n+1}^{k=n-1} [(n+1-|k+1|) - 2(n+1-|k|) + (n+1-|k-1|)] e^{ikx}$$

$$= e^{-i(n+1)x} + e^{i(n+1)x} + \sum_{k=-n+1}^{k=n-1} (2|k| - |k+1| - |k-1|) e^{ikx}$$

$$= e^{-i(n+1)x} + e^{i(n+1)x} - 2 \quad \text{by item (2)}$$

$$= -4 \left(\sin\frac{n+1}{2}x\right)^2 \quad \text{by item (1)}$$

Definition H.5. 20 Let K_n be FEJÉR'S KERNEL (Definition H.4 page 191).

The **de la Vallée Poussin kernel** \vee_n is defined as E $V_n(x) \triangleq 2K_{2n+1}(x) - K_n(x)$

Definition H.6. ²¹ Let K_n be Fejér's Kernel (Definition H.4 page 191). The **Jackson kernel** J_n is defined as

 $J_n(x) \triangleq \left\| K_n \right\|^{-2} K_n^2(x)$

Definition H.7. 22

The **Poisson kernel** P is defined as D E F $P(r,x) \triangleq \sum_{k \in \mathbb{Z}} r^{|k|} e^{ikx}$

 $^{^{20}}$ Katznelson (2004) page 16

²¹ Katznelson (2004) page 17

²² Katznelson (2004) page 16



APPENDIX I	
1	
	FOURIER TRANSFORM



□ Up to this point we have supposed that the function whose development is required in a series of sines of multiple arcs can be developed in a series arranged according to powers of the variable χ, ... We can extend the same results to any functions, even to those which are discontinuous and entirely arbitrary. ... even entirely arbitrary functions may be developed in series of sines of multiple arcs. ♥

Joseph Fourier (1768–1830)

I.1 Introduction

Historically, before the Fourier Transform was the Taylor Expansion (transform). The Taylor Expansion demonstrates that for **analytic** functions, knowledge of the derivatives of a function at a location x = a allows you to determine (predict) arbitrarily closely all the points f(x) in the vicinity of x = a (Chapter K page 221). But analytic functions are by definition functions for which all their derivatives exist. Thus, if a function is *discontinuous*, it is simply not a candidate for the Taylor Expansion. And some 300 years ago, mathematician giants of the day were fairly content with this.

But then in came an engineer named Joseph Fourier whose day job was working as a governor of lower Egypt under Napolean. He claimed that, rather than expansion based on derivatives, one could expand based on integrals over sinusoids, and that this would work not just for analytic functions, but for **discontinuous** ones as well!²

Needless to say, this did not go over too well initially in the mathematical community. But over time (on the order of 200 or so years), the Fourier Transform has in many ways won the day.



¹ quote: *☐* Fourier (1878) page 184,186 ⟨\$219,220⟩

image: http://en.wikipedia.org/wiki/File:Fourier2.jpg, public domain

² Robinson (1982) page 886

³Caricature of Legendre (left) and Fourier (right), 1820, by Julien-Léopold Boilly (1796–1874). "Album de 73

Definitions

This chapter deals with the Fourier Transform in the space of Lebesgue square-integrable functions $L^2_{(\mathbb{R},\mathcal{B},\mu)}$, where \mathbb{R} is the set of real numbers, \mathcal{B} is the set of *Borel sets* on \mathbb{R} , μ is the standard *Borel measure* on \mathcal{B} , and

$$L^2_{(\mathbb{R},\mathscr{B},\mu)} \triangleq \left\{ f \in \mathbb{R}^{\mathbb{R}} | \int_{\mathbb{R}} |f|^2 d\mu < \infty \right\}.$$

Daniel J. Greenhoe

 $\mathbf{L}^{2}_{(\mathbb{R},\mathcal{B},\mu)} \triangleq \bigg\{ \mathsf{f} \in \mathbb{R}^{\mathbb{R}} | \int_{\mathbb{R}} |\mathsf{f}|^{2} \, \mathsf{d}\mu < \infty \bigg\}.$ Furthermore, $\langle \triangle | \nabla \rangle$ is the *inner product* induced by the operator $\int_{\mathbb{R}} \, \mathsf{d}\mu$ such that

$$\langle f | g \rangle \triangleq \int_{\mathbb{R}} f(x) g^*(x) dx,$$

and $\left(L^2_{(\mathbb{R},\mathscr{B},\mu)},\langle\triangle\mid\nabla\rangle\right)$ is a *Hilbert space*.

Definition I.1. *Let* κ *be a* Function $in \mathbb{C}^{\mathbb{R}^2}$.



The function κ is the **Fourier kernel** if $\kappa(x,\omega) \triangleq e^{i\omega x}$

$$\kappa(x,\omega) \triangleq e^{i\omega x}$$

 $\forall x,\omega \in \mathbb{R}$

Definition I.2. 4 Let $L^2_{(\mathbb{R},\mathcal{B},\mu)}$ be the space of all Lebesgue square-integrable functions.

E

The **Fourier Transform** operator $ilde{\mathbf{F}}$ is defined as

$$\left[\tilde{\mathbf{F}}\mathbf{f}\right](\omega) \triangleq \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} \mathbf{f}(x) e^{-i\omega x} dx \quad \forall \mathbf{f} \in L^{2}_{(\mathbb{R}, \mathcal{B}, \mu)}$$

This definition of the Fourier Transform is also called the unitary Fourier Transform.

Remark I.1 (Fourier transform scaling factor). 5 If the Fourier transform operator $\tilde{\mathbf{F}}$ and inverse Fourier transform operator $\tilde{\mathbf{F}}^{-1}$ are defined as

$$\tilde{\mathbf{F}} \mathbf{f}(x) \triangleq \mathbf{F}(\omega) \triangleq A \int_{\mathbb{R}} \mathbf{f}(x) e^{-i\omega x} dx$$
 and $\tilde{\mathbf{F}}^{-1} \tilde{\mathbf{f}}(\omega) \triangleq B \int_{\mathbb{R}} \mathbf{F}(\omega) e^{i\omega x} d\omega$

then A and B can be any constants as long as $AB = \frac{1}{2\pi}$. The Fourier transform is often defined with the scaling factor A set equal to 1 such that $\left[\tilde{\mathbf{F}}f(x)\right](\omega) \triangleq \int_{\mathbb{R}} f(x) e^{-i\omega x} dx$. In this case, the inverse Fourier transform operator $\tilde{\mathbf{F}}^{-1}$ is either defined as

(using oscillatory frequency free variable
$$f$$
) or $\left[\tilde{\mathbf{F}}^{-1}\mathsf{f}(x)\right](\omega) \triangleq \frac{1}{2\pi} \int_{\mathbb{R}} \mathsf{f}(x) \, e^{i\omega x} \, \mathrm{dx}$ (using angular frequency free variable ω).

$$\left[\tilde{\mathbf{F}}^{-1}\mathsf{f}(x)\right](\omega) \triangleq \frac{1}{2\pi} \int_{\mathbb{R}} \mathsf{f}(x) \, e^{i\omega x} \, dx \quad \text{(using angular frequency free variable } \omega\text{)}$$

In short, the 2π has to show up somewhere, either in the argument of the exponential $(e^{-i2\pi ft})$ or in front of the integral $(\frac{1}{2\pi} \int \cdots)$. One could argue that it is unnecessary to burden the exponential argument with the 2π factor $(e^{-i2\pi ft})$, and thus could further argue in favor of using the angular frequency variable ω thus giving the inverse operator definition $\left[\tilde{\mathbf{F}}^{-1}\mathsf{f}(x)\right](\omega) \triangleq \frac{1}{2\pi} \int_{\mathbb{R}} \mathsf{f}(x) \, e^{-i\omega x} \, dx$. But this causes a new problem. In this case, the Fourier operator $\tilde{\mathbf{F}}$ is not *unitary* (see Theorem I.2 page 197)—in particular, $\tilde{\mathbf{F}}\tilde{\mathbf{F}}^* \neq \mathbf{I}$, where $\tilde{\mathbf{F}}^*$ is the *adjoint* of $\tilde{\mathbf{F}}$; but rather, $\tilde{\mathbf{F}}\left(\frac{1}{2\pi}\tilde{\mathbf{F}}^*\right) = \left(\frac{1}{2\pi}\tilde{\mathbf{F}}^*\right)\tilde{\mathbf{F}} = \mathbf{I}$. But if we define the operators $\tilde{\mathbf{F}}$ and $\tilde{\mathbf{F}}^{-1}$ to both have the scaling factor $\frac{1}{\sqrt{2\pi}}$, then $\tilde{\mathbf{F}}$ and $\tilde{\mathbf{F}}^{-1}$ are inverses and $\tilde{\mathbf{F}}$ is unitary—that is, $\tilde{\mathbf{F}}\tilde{\mathbf{F}}^* = \tilde{\mathbf{F}}^*\tilde{\mathbf{F}} = \mathbf{I}$.

Portraits-Charge Aquarelle's des Membres de l'Institute (watercolor portrait #29). Biliotheque de l'Institut de France." Public domain. https://en.wikipedia.org/wiki/File:Legendre_and_Fourier_(1820).jpg

 $^{^5}$ Chorin and Hald (2009) page 13, 🏿 Jeffrey and Dai (2008) pages xxxi–xxxii, 🗷 Knapp (2005b) pages 374–375



⁴ Bachman et al. (2002) page 363, 🛭 Chorin and Hald (2009) page 13, 🗐 Loomis and Bolker (1965) page 144, Knapp (2005b) pages 374–375, Fourier (1822), Fourier (1878) page 336?

Operator properties I.3

Theorem I.1 (Inverse Fourier transform). 6 Let $\tilde{\mathbf{F}}$ be the Fourier Transform operator (Definition 1.2 page 196). The inverse $\tilde{\mathbf{F}}^{-1}$ of $\tilde{\mathbf{F}}$ is

$$\begin{bmatrix} \tilde{\mathbf{F}}^{-1}\tilde{\mathbf{f}} \end{bmatrix}(x) = \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} \tilde{\mathbf{f}}(\omega) e^{i\omega x} \, d\omega \qquad \forall \tilde{\mathbf{f}} \in L^2_{(\mathbb{R}, \mathcal{B}, \mu)}$$

Theorem I.2. Let $\tilde{\mathbf{F}}$ be the Fourier Transform operator with inverse $\tilde{\mathbf{F}}^{-1}$ and adjoint $\tilde{\mathbf{F}}^*$.

$$\mathbf{\tilde{F}}^* = \mathbf{\tilde{F}}^{-1}$$

[♠]Proof:

$$\begin{split} \left\langle \tilde{\mathbf{F}} \mathsf{f} \mid \mathsf{g} \right\rangle &= \left\langle \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} \mathsf{f}(x) \, e^{-i\omega x} \, \, \mathsf{dx} \mid \mathsf{g}(\omega) \right\rangle & \text{by definition of } \tilde{\mathbf{F}} \text{ page } 196 \\ &= \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} \mathsf{f}(x) \, \left\langle e^{-i\omega x} \mid \mathsf{g}(\omega) \right\rangle \, \, \mathsf{dx} & \text{by } \textit{additive property of } \left\langle \triangle \mid \nabla \right\rangle \\ &= \int_{\mathbb{R}} \mathsf{f}(x) \, \frac{1}{\sqrt{2\pi}} \, \left\langle \mathsf{g}(\omega) \mid e^{-i\omega x} \right\rangle^* \, \, \mathsf{dx} & \text{by } \textit{conjugate symmetric property of } \left\langle \triangle \mid \nabla \right\rangle \\ &= \left\langle \mathsf{f}(x) \mid \frac{1}{\sqrt{2\pi}} \, \left\langle \mathsf{g}(\omega) \mid e^{-i\omega x} \right\rangle \right\rangle & \text{by definition of } \left\langle \triangle \mid \nabla \right\rangle \\ &= \left\langle \mathsf{f} \mid \tilde{\mathbf{F}}^{-1} \mathsf{g} \right\rangle & \text{by Theorem I.1 page } 197 \end{split}$$

The Fourier Transform operator has several nice properties:

- $\stackrel{\text{def}}{=}$ F is unitary ⁷ (Corollary I.1—next corollary).
- Because $\tilde{\mathbf{F}}$ is unitary, it automatically has several other nice properties (Theorem 1.3 page 197).

Corollary I.1. Let I be the identity operator and let $\tilde{\mathbf{F}}$ be the Fourier Transform operator with adjoint $\tilde{\mathbf{F}}^*$ and inverse $\tilde{\mathbf{F}}^{-1}$.

$$\tilde{\mathbf{F}}^* = \tilde{\mathbf{F}}^* = \mathbf{F}^{-1}$$

$$\tilde{\mathbf{F}}^* = \tilde{\mathbf{F}}^{-1}$$
(\tilde{\mathbf{F}} is unitary)

 $^{\circ}$ Proof: This follows directly from the fact that $\tilde{\mathbf{F}}^* = \tilde{\mathbf{F}}^{-1}$ (Theorem I.2 page 197).

Theorem I.3. Let $\tilde{\mathbf{F}}$ be the Fourier transform operator with adjoint $\tilde{\mathbf{F}}^*$ and inverse $\tilde{\mathbf{F}}$. Let $\|\cdot\|$ be the operator norm with respect to the vector norm $\|\cdot\|$ with respect to the Hilbert space $(\mathbb{C}^{\mathbb{R}}, \langle \triangle \mid \nabla \rangle)$. Let $\mathcal{R}(\mathbf{A})$ be the range of an operator \mathbf{A} .

 $^{\circ}$ Proof: These results follow directly from the fact that $\tilde{\mathbf{F}}$ is unitary (Corollary I.1 page 197) and from the properties of unitary operators (Theorem C.26 page 136).

⁷ unitary operators: Definition C.14 page 135





—>

⁶ Chorin and Hald (2009) page 13

I.4 Transversal properties


Theorem I.4 (Shift relations). Let $\tilde{\mathbf{F}}$ be the Fourier Transform operator (Definition 1.2 page 196).

$$\begin{array}{lll} \mathbf{\tilde{F}} & \mathbf{\tilde{F}}[\mathbf{f}(x-y)](\omega) & = & e^{-i\omega y} \left[\mathbf{\tilde{F}}\mathbf{f}(x)\right](\omega) \\ \mathbf{M} & \left[\mathbf{\tilde{F}}\left(e^{irx}\mathbf{g}(x)\right)\right](\omega) & = & \left[\mathbf{\tilde{F}}\mathbf{g}(x)\right](\omega-r) \end{array}$$

PROOF: Let L be the *Laplace Transform* operator (Definition L.1 page 223).

$$\begin{split} \tilde{\mathbf{F}}[\mathbf{f}(x-y)](\omega) &= \mathbf{L}[\mathbf{f}(x-y)](s)|_{s=i\omega} & \text{by definition of } \mathbf{L} & \text{(Definition L.1 page 223)} \\ &= e^{-sy} \left[\mathbf{L}\mathbf{f}(x)](s)|_{s=i\omega} & \text{by Laplace } translation \text{ property} & \text{(Theorem L.2 page 224)} \\ &= e^{-i\omega y} \left[\tilde{\mathbf{F}}\mathbf{f}(x) \right](\omega) & \text{by definition of } \tilde{\mathbf{F}} & \text{(Definition I.2 page 196)} \\ \left[\tilde{\mathbf{F}}\left(e^{irx}\mathbf{g}(x)\right) \right](\omega) &= \left[\mathbf{L}\left(e^{irx}\mathbf{g}(x)\right) \right](s)|_{s=i\omega} & \text{by definition of } \mathbf{L} & \text{(Definition L.1 page 223)} \\ &= \left[\left[\mathbf{L}\mathbf{g}(x) \right](s-r) \right]|_{s=i\omega} & \text{by Laplace } dilation \text{ property} & \text{(Theorem L.2 page 224)} \\ &= \left[\tilde{\mathbf{F}}\mathbf{g}(x) \right](\omega-r) & \text{by definition of } \tilde{\mathbf{F}} & \text{(Definition I.2 page 196)} \end{split}$$

Theorem I.5 (Complex conjugate). Let $\tilde{\mathbf{F}}$ be the Fourier Transform operator and * represent the complex conjugate operation on the set of complex numbers.



[♠]Proof:

$$\begin{split} \left[\tilde{\mathbf{F}}\mathsf{f}^*(-x)\right](\omega) &\triangleq \frac{1}{\sqrt{2\pi}} \int \mathsf{f}^*(-x)e^{-i\omega x} \, \, \mathsf{d} \mathsf{x} \qquad \text{by definition of } \tilde{\mathbf{F}} \qquad \text{(Definition I.2 page 196)} \\ &= \frac{1}{\sqrt{2\pi}} \int \mathsf{f}^*(u)e^{i\omega u}(-1) \, \, \mathsf{d} \mathsf{u} \qquad \text{where } u \triangleq -x \implies \mathsf{d} \mathsf{x} = - \, \mathsf{d} \mathsf{u} \\ &= -\left[\frac{1}{\sqrt{2\pi}} \int \mathsf{f}(u)e^{-i\omega u} \, \, \mathsf{d} \mathsf{u}\right]^* \\ &\triangleq -\left[\tilde{\mathbf{F}}\mathsf{f}(x)\right]^* \qquad \text{by definition of } \tilde{\mathbf{F}} \qquad \text{(Definition I.2 page 196)} \end{split}$$


I.5 Convolution relations

Theorem J.2 (next) demonstrates that multiplication in the "time domain" is equivalent to convolution in the "frequency domain" and vice-versa.

Theorem I.6 (convolution theorem). ⁸ Let $\tilde{\mathbf{F}}$ be the Fourier Transform operator (Definition 1.2 page 196) and \star the convolution operator (Definition E.1 page 145).

⁸ ■ Bachman et al. (2002) pages 269–270 (5.2.3 Convolutions to Products), ■ Bachman (1964) page 8, ■ Bracewell (1978) page 110





 $^{\circ}$ Proof: Let **L** be the *Laplace Transform* operator (Definition L.1 page 223).

$$\begin{split} \tilde{\mathbf{F}}\big[\mathbf{f}(x)\star\mathbf{g}(x)\big](\omega) &= \mathbf{L}\big[\mathbf{f}(x)\star\mathbf{g}(x)\big](s)\big|_{s=i\omega} & \text{by definition of } \mathbf{L} & \text{(Definition L.1 page 223)} \\ &= \sqrt{2\pi}\big[\mathbf{L}\mathbf{f}\big](s)\, \big[\mathbf{L}\mathbf{g}\big](s)\big|_{s=i\omega} & \text{by } Laplace \ convolution \ \text{result} & \text{(Theorem L.6 page 235)} \\ &= \sqrt{2\pi}\big[\tilde{\mathbf{F}}\mathbf{f}\big](\omega)\, \big[\tilde{\mathbf{F}}\mathbf{g}\big](\omega) & \\ &= \sqrt{2\pi}\big[\tilde{\mathbf{F}}\mathbf{f}\big](\omega)\, \big[\tilde{\mathbf{F}}\mathbf{g}\big](\omega) & \\ &= \frac{1}{\sqrt{2\pi}}\big[\mathbf{L}\mathbf{f}\big](s)\star \big[\mathbf{L}\mathbf{g}\big](s)\big|_{s=i\omega} & \\ &= \frac{1}{\sqrt{2\pi}}\big[\tilde{\mathbf{F}}\mathbf{f}\big](\omega)\star \big[\tilde{\mathbf{F}}\mathbf{g}\big](\omega) & \end{split}$$

Calculus relations I.6

Theorem I.7. Let $\tilde{\mathbf{F}}$ be the Fourier Transform operator (Definition 1.2 page 196).

$$\begin{cases} \lim_{t \to -\infty} \mathbf{x}(t) = 0 \end{cases} \implies \left\{ \tilde{\mathbf{F}} \left[\frac{\mathsf{d}}{\mathsf{dt}} \mathbf{x}(t) \right] = i\omega \left[\tilde{\mathbf{F}} \mathbf{x} \right](\omega) \right\}$$

 $^{\circ}$ Proof: Let L be the *Laplace Transform* operator (Definition L.1 page 223).

$$\tilde{\mathbf{F}} \left[\frac{\mathsf{d}}{\mathsf{dt}} \mathbf{x}(t) \right] \triangleq \mathbf{L} \left[\frac{\mathsf{d}}{\mathsf{dt}} \mathbf{x}(t) \right] (s) \Big|_{s=i\omega}$$
 by definitions of **L** and $\tilde{\mathbf{F}}$ (Definition L.1 page 223)
$$= s[\mathbf{L}\mathbf{x}(t)](s)|_{s=i\omega}$$
 by Theorem L.7 page 236
$$= i\omega [\tilde{\mathbf{F}}\mathbf{x}](\omega)$$

Theorem I.8. Let $\tilde{\mathbf{F}}$ be the Fourier Transform operator (Definition 1.2 page 196).

$$\mathbf{\tilde{F}} \int_{u=-\infty}^{u=t} \mathsf{x}(u) \, \mathsf{du} = \frac{1}{i\omega} \big[\mathbf{\tilde{F}} \mathsf{x} \big](\omega)$$

Let L be the *Laplace Transform* operator (Definition L.1 page 223).

$$\tilde{\mathbf{F}} \int_{u=-\infty}^{u=t} \mathsf{x}(u) \, \mathsf{du} \triangleq \mathbf{L} \int_{u=-\infty}^{u=t} \mathsf{x}(u) \, \mathsf{du} \bigg|_{s=i\omega}$$

$$= \frac{1}{s} [\mathbf{L}\mathsf{x}(t)](s) \bigg|_{s=i\omega} \qquad \text{by Theorem L.7 page 236}$$

$$= \frac{1}{i\omega} [\tilde{\mathbf{F}}\mathsf{x}(t)](\omega)$$

Real valued functions I.7




Figure I.1: Fourier transform components of real-valued signal

Theorem I.9. Let f(x) be a function in $L^2_{\mathbb{R}}$ and $\tilde{f}(\omega)$ the Fourier Transform of f(x).

Theorem I.9. Let
$$f(x)$$
 be a function in $L^2_{\mathbb{R}}$ and $\tilde{f}(\omega)$ the Fourier Transform of $f(x)$.

$$\begin{cases}
f(x) \text{ is real-valued} \\
(f \in \mathbb{R}^{\mathbb{R}})
\end{cases}
\Rightarrow
\begin{cases}
\tilde{f}(\omega) = \tilde{f}^*(-\omega) & (\text{Hermitian Symmetric}) \\
R_e[\tilde{f}(\omega)] = R_e[\tilde{f}(-\omega)] & (\text{Symmetric}) \\
I_m[\tilde{f}(\omega)] = -I_m[\tilde{f}(-\omega)] & (\text{Symmetric}) \\
|\tilde{f}(\omega)| = |\tilde{f}(-\omega)| & (\text{Symmetric}) \\
|\tilde{f}(\omega)| = |\tilde{f}(-\omega)| & (\text{Symmetric}).
\end{cases}$$

[♠]Proof:

$$\begin{array}{llll} \tilde{\mathbf{f}}(\omega) & \triangleq & [\tilde{\mathbf{F}}\mathbf{f}(x)](\omega) & \triangleq & \left\langle \mathbf{f}(x) \,|\, e^{i\omega x} \right\rangle & = & \left\langle \mathbf{f}(x) \,|\, e^{i(-\omega)x} \right\rangle^* & \triangleq & \tilde{\mathbf{f}}^*(-\omega) \\ \mathbf{R}_{\mathrm{e}} \left[\tilde{\mathbf{f}}(\omega) \right] & = & \mathbf{R}_{\mathrm{e}} \left[\tilde{\mathbf{f}}^*(-\omega) \right] & = & \mathbf{R}_{\mathrm{e}} \left[\tilde{\mathbf{f}}(-\omega) \right] \\ \mathbf{I}_{\mathrm{m}} \left[\tilde{\mathbf{f}}(\omega) \right] & = & \mathbf{I}_{\mathrm{m}} \left[\tilde{\mathbf{f}}^*(-\omega) \right] & = & -\mathbf{I}_{\mathrm{m}} \left[\tilde{\mathbf{f}}(-\omega) \right] \\ |\tilde{\mathbf{f}}(\omega)| & = & |\tilde{\mathbf{f}}^*(-\omega)| & = & |\tilde{\mathbf{f}}(-\omega)| \\ \mathcal{L}\tilde{\mathbf{f}}(\omega) & = & \mathcal{L}\tilde{\mathbf{f}}^*(-\omega) & = & -\mathcal{L}\tilde{\mathbf{f}}(-\omega) \end{array}$$

Moment properties I.8

Definition I.3. 9

The quantity M_n is the n**th moment** of a function $f(x) \in L_{\mathbb{R}}^2$ if $M_n \triangleq \int_{\mathbb{R}} x^n f(x) dx$ for $n \in \mathbb{W}$.

Lemma I.1. ¹⁰ Let M_n be the nth moment (Definition 1.3 page 200) and $\tilde{f}(\omega) \triangleq \left[\tilde{\mathbf{F}}f\right](\omega)$ the Fourier trans-


⁹ Jawerth and Sweldens (1994) pages 16–17, E Sweldens and Piessens (1993) page 2, € Vidakovic (1999) page 83 ¹⁰ Goswami and Chan (1999) pages 38–39



♥Proof:

$$\begin{split} \sqrt{2\pi}(i)^n \Big[\Big[\frac{\mathrm{d}}{\mathrm{d}\omega} \Big]^n \tilde{\mathsf{f}}(\omega) \Big]_{\omega=0} &= \sqrt{2\pi}(i)^n \Big[\Big[\frac{\mathrm{d}}{\mathrm{d}\omega} \Big]^n \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} \mathsf{f}(x) e^{-i\omega x} \, \mathrm{d}x \Big]_{\omega=0} \quad \text{by definition of } \tilde{\mathbf{F}} \quad \text{(Definition I.2 page 196)} \\ &= (i)^n \int_{\mathbb{R}} \mathsf{f}(x) \Big[\Big[\frac{\mathrm{d}}{\mathrm{d}\omega} \Big]^n e^{-i\omega x} \Big] \, \mathrm{d}x \Big|_{\omega=0} \\ &= (i)^n \int_{\mathbb{R}} \mathsf{f}(x) \Big[(-i)^n x^n e^{-i\omega x} \Big] \, \mathrm{d}x \Big|_{\omega=0} \\ &= (-i^2)^n \int_{\mathbb{R}} \mathsf{f}(x) x^n \, \mathrm{d}x \\ &= \int_{\mathbb{R}} x^n \mathsf{f}(x) \, \mathrm{d}x \\ &\triangleq \mathsf{M}_n \quad \text{by definition of } \mathsf{M}_n \quad \text{(Definition I.3 page 200)} \end{split}$$

Lemma I.2. 11 Let M_n be the nth moment (Definition 1.3 page 200) and $\tilde{f}(\omega) \triangleq [\tilde{\mathbf{F}}f](\omega)$ the Fourier trans-FORM (Definition I.2 page 196) of a function f(x) in $\mathcal{L}^2_{\mathbb{R}}$ (Definition D.1 page 141).



^ℚProof:

1. Proof for (\Longrightarrow) case:

$$0 = \langle \mathbf{f}(x) \mid x^n \rangle$$

$$= \sqrt{2\pi} (-i)^{-n} \left[\frac{\mathbf{d}}{\mathbf{d}\omega} \right]^n \tilde{\mathbf{f}}(\omega) \Big|_{\omega=0}$$

$$\implies \left[\frac{\mathbf{d}}{\mathbf{d}\omega} \right]^n \tilde{\mathbf{f}}(\omega) \Big|_{\omega=0} = 0$$

by left hypothesis

by Lemma I.1 page 200

2. Proof for (\Leftarrow) case:

$$0 = \left[\frac{d}{d\omega} \right]^n \tilde{f}(\omega) \Big|_{\omega=0}$$

$$= \left[\frac{d}{d\omega} \right]^n \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} f(x) e^{-i\omega x} dx \Big|_{\omega=0}$$

$$= \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} f(x) \left[\frac{d}{d\omega} \right]^n e^{-i\omega x} dx \Big|_{\omega=0}$$

$$= \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} f(x) \left[(-i)^n x^n e^{-i\omega x} \right] dx \Big|_{\omega=0}$$

$$= (-i)^n \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} f(x) x^n dx$$

$$= (-i)^n \frac{1}{\sqrt{2\pi}} \langle f(x) | x^n \rangle$$

by right hypothesis

by definition of $\tilde{f}(\omega)$

by definition of $\langle \cdot | \cdot \rangle$ in $\mathcal{L}^2_{\mathbb{R}}$ (Definition D.1 page 141)

¹¹ Vidakovic (1999) pages 82–83, Mallat (1999) pages 241–242

🚧 Frames and Bases Structure and Design [VERSION 0.21] 🚧 https://github.com/dgreenhoe/pdfs/blob/master/msdframes.pdf





Lemma I.3 (Strang-Fix condition). ¹² Let f(x) be a function in $L^2_{\mathbb{R}}$ and M_n the nth moment (Definition 1.3 page 200) of f(x). Let T be the translation operator (Definition 3.3 page 40).

 $\sum_{k \in \mathbb{Z}} \mathbf{T}^k x^n \mathbf{f}(x) = \mathbf{M}_n \qquad \Longleftrightarrow \qquad \underbrace{\left[\frac{\mathrm{d}}{\mathrm{d}\omega}\right]^n \tilde{\mathbf{f}}(\omega)}_{\text{Strang-Fix condition } in \text{ "time"}} = \frac{1}{\sqrt{2\pi}} (-i)^n \bar{\delta}_k \mathbf{M}_n$

♥Proof:

1. Proof for (\Longrightarrow) case:

$$\begin{split} \left[\left[\frac{\mathrm{d}}{\mathrm{d}\omega} \right]^n & \tilde{\mathbf{f}}(\omega) \right]_{\omega = 2\pi k} &= \sum_{k \in \mathbb{Z}} \left[\left[\frac{\mathrm{d}}{\mathrm{d}\omega} \right]^n \tilde{\mathbf{f}}(\omega) \right]_{\omega = 2\pi k} e^{i2\pi kx} \bar{\delta}_k \\ &= \frac{1}{\sqrt{2\pi}} \sum_{k \in \mathbb{Z}} \left[\left[\frac{\mathrm{d}}{\mathrm{d}\omega} \right]^n \int_{\mathbb{R}} \mathbf{f}(x) e^{-i\omega x} \, \mathrm{d}x \right]_{\omega = 2\pi k} e^{i2\pi kx} \bar{\delta}_k \quad \text{by definition of } \tilde{\mathbf{f}}(\omega) \quad \text{(Definition I.2 page 196)} \\ &= \frac{1}{\sqrt{2\pi}} \sum_{k \in \mathbb{Z}} \left[\int_{\mathbb{R}} \mathbf{f}(x) (-ix)^n e^{-i\omega x} \, \mathrm{d}x \right]_{\omega = 2\pi k} e^{i2\pi kx} \bar{\delta}_k \\ &= (-i)^n \frac{1}{\sqrt{2\pi}} \sum_{k \in \mathbb{Z}} \left[\int_{\mathbb{R}} x^n \mathbf{f}(x) e^{-i\omega x} \, \mathrm{d}x \right]_{\omega = 2\pi k} e^{i2\pi kx} \bar{\delta}_k \\ &= (-i)^n \frac{1}{\sqrt{2\pi}} \sum_{k \in \mathbb{Z}} (x - k)^n \mathbf{f}(x - k) \bar{\delta}_k \qquad \text{by PSF} \quad \text{(Theorem 3.2 page 48)} \\ &= \frac{1}{\sqrt{2\pi}} (-i)^n \bar{\delta}_k \mathsf{M}_n \qquad \text{by left hypothesis} \end{split}$$

2. Proof for (\Leftarrow) case:


$$\begin{split} \frac{1}{\sqrt{2\pi}}(-i)^n \mathsf{M}_n &= \frac{1}{\sqrt{2\pi}} \sum_{k \in \mathbb{Z}} \left[(-i)^n \bar{\delta}_k \mathsf{M}_n \right] e^{-i2\pi kx} & \text{by definition of } \bar{\delta} \\ &= \sum_{k \in \mathbb{Z}} \left[\left[\frac{\mathrm{d}}{\mathrm{d}\omega} \right]^n \tilde{\mathsf{f}}(\omega) \right] \Big|_{\omega = 2\pi k} e^{-i2\pi kx} & \text{by right hypothesis} \\ &= \sum_{k \in \mathbb{Z}} \left[\left[\frac{\mathrm{d}}{\mathrm{d}\omega} \right]^n \int_{\mathbb{R}} \mathsf{f}(x) e^{-i\omega x} \, \mathrm{d}x \right] \Big|_{\omega = 2\pi k} e^{-i2\pi kx} \\ &= \sum_{k \in \mathbb{Z}} \left[\int_{\mathbb{R}} \mathsf{f}(x) (-ix)^n e^{-i\omega x} \, \mathrm{d}x \right] \Big|_{\omega = 2\pi k} e^{-i2\pi kx} \\ &= (-i)^n \sum_{k \in \mathbb{Z}} \left[\int_{\mathbb{R}} x^n \mathsf{f}(x) e^{-i\omega x} \, \mathrm{d}x \right] \Big|_{\omega = 2\pi k} e^{-i2\pi kx} \\ &= (-i)^n \sum_{k \in \mathbb{Z}} \left[x^n \mathsf{f}(x) e^{-i\omega x} \, \mathrm{d}x \right] \Big|_{\omega = 2\pi k} e^{-i2\pi kx} \end{split}$$
Theorem 3.2 page 48)

¹² ■ Jawerth and Sweldens (1994) pages 16–17, ■ Sweldens and Piessens (1993) page 2, ■ Vidakovic (1999) page 83, ■ Mallat (1999) pages 241–243, ■ Fix and Strang (1969)



Examples I.9

Example I.1 (rectangular pulse). Let $\tilde{f}(\omega)$ be the *Fourier transform* of a function $f(x) \in L^2_{\mathbb{R}}$.




^ℚProof:

E X

$$\begin{split} &\tilde{\mathbf{f}}(\omega) = \tilde{\mathbf{F}}[\mathbf{f}(x)](\omega) & \text{by definition of } \tilde{\mathbf{f}}(\omega) \\ &= e^{-i\left(\frac{a+b}{2}\right)\omega} \tilde{\mathbf{F}}\Big[\mathbf{f}\left(x-\frac{a+b}{2}\right)\Big](\omega) & \text{by shift relation} & \text{(Theorem I.4 page 198)} \\ &= e^{-i\left(\frac{a+b}{2}\right)\omega} \tilde{\mathbf{F}}\Big[c\,\mathbb{I}_{[a:b)}\Big(x-\frac{a+b}{2}\Big)\Big](\omega) & \text{by definition of } \mathbf{f}(x) \\ &= e^{-i\left(\frac{a+b}{2}\right)\omega} \tilde{\mathbf{F}}\Big[c\,\mathbb{I}_{\left[-\frac{b-a}{2}:\frac{b-a}{2}\right)}(x)\Big](\omega) & \text{by definition of } \mathbb{I} & \text{(Definition 3.2 page 40)} \\ &= \frac{1}{\sqrt{2}\pi}e^{-i\left(\frac{a+b}{2}\right)\omega} \int_{\mathbb{R}} c\,\mathbb{I}_{\left[-\frac{b-a}{2}:\frac{b-a}{2}\right)}(x)e^{-i\omega x} \,\mathrm{d}x & \text{by definition of } \mathbb{I} & \text{(Definition 1.2 page 196)} \\ &= \frac{1}{\sqrt{2}\pi}e^{-i\left(\frac{a+b}{2}\right)\omega} \int_{-\frac{b-a}{2}}^{\frac{b-a}{2}} ce^{-i\omega x} \,\mathrm{d}x & \text{by definition of } \mathbb{I} & \text{(Definition 3.2 page 40)} \\ &= \frac{c}{\sqrt{2}\pi}e^{-i\left(\frac{a+b}{2}\right)\omega} \frac{1}{-i\omega}e^{-i\omega x} \Big|_{-\frac{b-a}{2}}^{\frac{b-a}{2}} \\ &= \frac{2c}{\sqrt{2}\pi\omega}e^{-i\left(\frac{a+b}{2}\right)\omega} \left[\frac{e^{i\left(\frac{b-a}{2}\omega\right)}-e^{-i\left(\frac{b-a}{2}\omega\right)}}{2i}\right] \\ &= \frac{c(b-a)}{\sqrt{2\pi}}e^{-i\left(\frac{a+b}{2}\omega\right)} \left[\frac{\sin\left(\frac{b-a}{2}\omega\right)}{\left(\frac{b-a}{2}\omega\right)}\right] & \text{by $Euler formulas} \end{aligned} \tag{Corollary G.2 page 161} \end{split}$$

Example I.2 (triangle). Let $\tilde{f}(\omega)$ be the *Fourier transform* of a function $f(x) \in L^2_{\mathbb{R}}$.



[♠]Proof:

EX

 $\tilde{\mathbf{f}}(\omega) = \tilde{\mathbf{F}}[\mathbf{f}(x)](\omega)$ by definition of $\tilde{f}(\omega)$

$$= e^{-i\left(\frac{a+b}{2}\right)\omega}\tilde{\mathbf{F}}\left[\mathbf{f}\left(x-\frac{a+b}{2}\right)\right](\omega) \qquad \text{by shift relation} \qquad \text{(Theorem I.4 page 198)}$$

$$= \tilde{\mathbf{F}}\left[c\left(1-\frac{|2x-b-a|}{b-a}\right)\mathbb{I}_{[a:b)}(x)\right](\omega) \qquad \text{by definition of } \mathbf{f}(x)$$

$$= c\tilde{\mathbf{F}}\left[\mathbb{I}_{\left[\frac{a}{2}:\frac{b}{2}\right)}(x)\star\mathbb{I}_{\left[\frac{a}{2}:\frac{b}{2}\right)}(x)\right](\omega)$$


$$= c\sqrt{2\pi}\tilde{\mathbf{F}}\left[\mathbb{I}_{\left[\frac{a}{2}:\frac{b}{2}\right)}\right]\tilde{\mathbf{F}}\left[\mathbb{I}_{\left[\frac{a}{2}:\frac{b}{2}\right)}\right] \qquad \text{by convolution theorem} \qquad \text{(Theorem J.2 page 210)}$$

$$= c\sqrt{2\pi}\left(\tilde{\mathbf{F}}\left[\mathbb{I}_{\left[\frac{a}{2}:\frac{b}{2}\right)}\right]\right)^2$$

$$= c\sqrt{2\pi}\left(\frac{\left(\frac{b}{2}-\frac{a}{2}\right)}{\sqrt{2\pi}}e^{-i\left(\frac{a+b}{4}\omega\right)}\left[\frac{\sin\left(\frac{b-a}{4}\omega\right)}{\left(\frac{b-a}{4}\omega\right)}\right]^2 \qquad \text{by Rectangular pulse ex.} \qquad \text{Example I.1 page 203}$$

$$= \frac{c(b-a)^2}{4\sqrt{2\pi}}e^{-i\left(\frac{a+b}{2}\omega\right)}\left[\frac{\sin\left(\frac{b-a}{4}\omega\right)}{\left(\frac{b-a}{4}\omega\right)}\right]^2$$

Example I.3. Let a function f be defined in terms of the cosine function (Definition G.1 page 155) as follows:



 $^{\circ}$ Proof: Let $\mathbb{1}_A(x)$ be the *set indicator function* (Definition 3.2 page 40) on a set A.

$$\begin{split} &\tilde{\mathsf{f}}(\omega) \triangleq \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} \mathsf{f}(x) e^{-i\omega x} \, \mathrm{d}x & \text{by definition of } \tilde{\mathsf{f}}(\omega) \text{ (Definition 1.2)} \\ &= \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} \cos^2 \left(\frac{\pi}{2}x\right) \mathbbm{1}_{[-1:1]}(x) e^{-i\omega x} \, \mathrm{d}x & \text{by definition of } \mathbbm{f}(x) \\ &= \frac{1}{\sqrt{2\pi}} \int_{-1}^1 \cos^2 \left(\frac{\pi}{2}x\right) e^{-i\omega x} \, \mathrm{d}x & \text{by definition of } \mathbbm{1} \text{ (Definition 3.2)} \\ &= \frac{1}{\sqrt{2\pi}} \int_{-1}^1 \left[e^{i\frac{\pi}{2}x} + e^{-i\frac{\pi}{2}x} \right]^2 e^{-i\omega x} \, \mathrm{d}x & \text{by Corollary G.2 page 161} \\ &= \frac{1}{4\sqrt{2\pi}} \int_{-1}^1 \left[2 + e^{i\pi x} + e^{-i\pi x} \right] e^{-i\omega x} \, \mathrm{d}x \\ &= \frac{1}{4\sqrt{2\pi}} \int_{-1}^1 2 e^{-i\omega x} + e^{-i(\omega + \pi)x} + e^{-i(\omega - \pi)x} \, \mathrm{d}x \\ &= \frac{1}{4\sqrt{2\pi}} \left[2 \frac{e^{-i\omega x}}{-i\omega} + \frac{e^{-i(\omega + \pi)x}}{-i(\omega + \pi)} + \frac{e^{-i(\omega - \pi)x}}{-i(\omega - \pi)} \right]_{-1}^1 \\ &= \frac{1}{2\sqrt{2\pi}} \left[2 \frac{e^{-i\omega}}{-2i\omega} + \frac{e^{-i(\omega + \pi)}}{-2i(\omega + \pi)} + \frac{e^{-i(\omega - \pi)}}{-2i(\omega - \pi)} \right]_{-1}^1 \end{split}$$

I.9. EXAMPLES Daniel J. Greenhoe page 205

$$=\frac{1}{2\sqrt{2\pi}}\left[\underbrace{\frac{2\mathrm{sin}\omega}{\omega}}_{2\,\mathrm{sinc}(\omega)} + \underbrace{\frac{\mathrm{sin}(\omega+\pi)}{(\omega+\pi)}}_{\mathrm{sinc}(\omega+\pi)} + \underbrace{\frac{\mathrm{sin}(\omega-\pi)}{(\omega-\pi)}}_{\mathrm{sinc}(\omega-\pi)}\right]$$



Example I.4. ¹³

$$\tilde{\mathbf{F}}\left[e^{-\alpha|x|}\right] = \frac{1}{\sqrt{2\pi}} \left[\frac{2\alpha}{\alpha^2 + \omega^2}\right]$$

^ℚProof:

1. Proof using *Laplace Transform*:

$$\begin{split} \sqrt{2\pi}\tilde{\mathbf{F}}\big[e^{-\alpha|x|}\big] &\triangleq \left[\sqrt{2\pi}\right] \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} e^{-\alpha|x|} e^{-i\omega x} \, \mathrm{d}x \quad \text{by definition of } \tilde{\mathbf{F}} \\ &= \left[\int_{\mathbb{R}} e^{-\alpha|x|} e^{-sx} \, \mathrm{d}x\right]_{s=i\omega} \\ &= \left[\frac{2\alpha}{\alpha^2 - s^2}\right]_{s=i\omega} \quad \forall \mathbf{R}_{\mathrm{e}}(s) \in (-\alpha:\alpha) \quad \text{by Corollary L.9 page 233} \\ &= \frac{2\alpha}{\alpha^2 + \omega^2} \quad \text{because } s = i\omega \text{ is in } (-\alpha:\alpha) \end{split}$$

2. Alternate proof:

$$\sqrt{2\pi}\tilde{\mathbf{F}}\left[e^{-\alpha|x|}\right] \triangleq \left[\sqrt{2\pi}\right] \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{-\alpha|x|} e^{-i\omega x} \, \mathrm{d}x \qquad \text{by definition}$$

$$= \int_{-\infty}^{0} e^{-\alpha(-x)} e^{-i\omega x} \, \mathrm{d}x + \int_{0}^{\infty} e^{-\alpha(x)} e^{-i\omega x} \, \mathrm{d}x$$

$$= \int_{-\infty}^{0} e^{x(\alpha-i\omega)} \, \mathrm{d}x + \int_{0}^{\infty} e^{x(-\alpha-i\omega)} \, \mathrm{d}x$$

$$= \frac{e^{x(\alpha-i\omega)}}{\alpha-i\omega} \Big|_{-\infty}^{0} + \frac{e^{x(-\alpha-i\omega)}}{-\alpha-i\omega} \Big|_{0}^{\infty} \qquad \text{by Fundame}$$

$$= \left[\frac{1}{\alpha-i\omega} - 0\right] + \left[0 - \frac{1}{-\alpha-i\omega}\right]$$

$$= \left[\frac{1}{\alpha-i\omega}\right] \left[\frac{\alpha-i\omega}{\alpha-i\omega}\right] + \left[\frac{1}{\alpha+i\omega}\right] \left[\frac{\alpha+i\omega}{\alpha+i\omega}\right]$$

$$= \frac{\alpha-i\omega}{\alpha^2+\omega^2} + \frac{\alpha+i\omega}{\alpha^2+\omega^2}$$

$$= \left[\frac{2\alpha}{\alpha^2+\omega^2}\right]$$

by definition of $\tilde{\mathbf{F}}$ (Definition I.2 page 196)

by Fundamental Theorem of Calculus

¹³https://math.stackexchange.com/questions/4015842/





Convolution operator I.1

Definition J.1. Let X^Y be the set of all functions from a set Y to a set X. Let \mathbb{Z} be the set of integers.

A function f in X^Y is a **sequence** over X if $Y = \mathbb{Z}$.

A sequence may be denoted in the form $(x_n)_{n\in\mathbb{Z}}$ or simply as (x_n) .

Definition J.2. 2 Let $(\mathbb{F}, +, \cdot, 0, 1)$ be a FIELD (Definition A.5 page 96).

The space of all absolutely square summable sequences $\mathscr{C}^2_{\mathbb{F}}$ over \mathbb{F} is defined as

$$\mathscr{C}_{\mathbb{F}}^2 \triangleq \left\{ \left(\left(x_n \right)_{n \in \mathbb{Z}} \mid \sum_{n \in \mathbb{Z}} \left| x_n \right|^2 < \infty \right\}$$

The space $\mathscr{C}^2_{\mathbb{R}}$ is an example of a *separable Hilbert space*. In fact, $\mathscr{C}^2_{\mathbb{R}}$ is the *only* separable Hilbert space in the sense that all separable Hilbert spaces are isomorphically equivalent. For example, $\mathscr{C}^2_{\mathbb{R}}$ is isomorphic to $L^2_{\mathbb{R}}$, the space of all absolutely square Lebesgue integrable functions.

Definition J.3.

D E F

D E The **convolution** operation \star is defined as

$$(x_n) \star (y_n) \triangleq \left(\sum_{m \in \mathbb{Z}} x_m y_{n-m} \right)_{n \in \mathbb{Z}} \forall (x_n)_{n \in \mathbb{Z}}, (y_n)_{n \in \mathbb{Z}} \in \mathscr{C}^2_{\mathbb{R}}$$

Proposition J.1. Let \star be the CONVOLUTION OPERATOR (Definition J.3 page 207).

¹ Bromwich (1908) page 1, Thomson et al. (2008) page 23 ⟨Definition 2.1⟩, Joshi (1997) page 31
² Kubrusly (2011) page 347 ⟨Example 5.K⟩

^ℚProof:

$$[x \star y](n) \triangleq \sum_{m \in \mathbb{Z}} x_m y_{n-m} \qquad \text{by Definition J.3 page 207}$$

$$= \sum_{k \in \mathbb{Z}} x_{n-k} y(k) \qquad \text{where } k \triangleq n-m \implies m=n-k$$

$$= \sum_{k \in \mathbb{Z}} x_{n-k} y(k) \qquad \text{by } commutativity \text{ of addition}$$

$$= \sum_{m \in \mathbb{Z}} x_{n-m} y_m \qquad \text{by change of variables}$$

$$= \sum_{m \in \mathbb{Z}} y_m x_{n-m} \qquad \text{by commutative property of the field over } \mathbb{C}$$

$$\triangleq (y \star x)_n \qquad \text{by Definition J.3 page 207}$$

Proposition J.2. Let \star be the Convolution operator (Definition J.3 page 207). Let $\mathscr{C}^2_{\mathbb{R}}$ be the set of Abso-LUTELY SUMMABLE Sequences (Definition J.2 page 207).

$$\begin{array}{c} \mathbf{P} \\ \mathbf{R} \\ \mathbf{P} \end{array} \left\{ \begin{array}{ccc} (A). & \mathbf{x}(n) & \in & \boldsymbol{\mathcal{C}}_{\mathbb{R}}^2 & and \\ (B). & \mathbf{y}(n) & \in & \boldsymbol{\mathcal{C}}_{\mathbb{R}}^2 \end{array} \right\} \implies \left\{ \begin{array}{ccc} \sum_{k \in \mathbb{Z}} \mathbf{x}[k] \mathbf{y}[n+k] & = & \mathbf{x}[-n] \star \mathbf{y}(n) \\ k \in \mathbb{Z} \end{array} \right\}$$

[♠]Proof:

$$\sum_{k \in \mathbb{Z}} \mathsf{x}[k]\mathsf{y}[n+k] = \sum_{-p \in \mathbb{Z}} \mathsf{x}[-p]\mathsf{y}[n-p] \qquad \text{where } p \triangleq -k \qquad \Longrightarrow k = -p$$

$$= \sum_{p \in \mathbb{Z}} \mathsf{x}[-p]\mathsf{y}[n-p] \qquad \text{by } absolutely \, summable \, \text{hypothesis} \qquad \text{(Definition J.2 page 207)}$$

$$= \sum_{p \in \mathbb{Z}} \mathsf{x}'[p]\mathsf{y}[n-p] \qquad \text{where } \mathsf{x}'[n] \triangleq \mathsf{x}[-n] \qquad \Longrightarrow \mathsf{x}[-n] = \mathsf{x}'[n]$$

$$\triangleq \mathsf{x}'[n] \star \mathsf{y}[n] \qquad \text{by definition of } convolution \star \qquad \text{(Definition J.3 page 207)}$$

$$\triangleq \mathsf{x}[-n] \star \mathsf{y}[n] \qquad \text{by definition of } \mathsf{x}'[n]$$

₽


Z-transform

Definition J.4. ³

D E F

The z-transform
$$\mathbf{Z}$$
 of $(x_n)_{n \in \mathbb{Z}}$ is defined as
$$\left[\mathbf{Z}(x_n)\right](z) \triangleq \sum_{n \in \mathbb{Z}} x_n z^{-n} \quad \forall (x_n) \in \mathscr{C}_{\mathbb{R}}^2$$
Laurent series

Theorem J.1. Let $X(z) \triangleq \mathbf{Z} \times [n]$ be the z-transform of $\times [n]$.



³Laurent series: Abramovich and Aliprantis (2002) page 49



J.2. Z-TRANSFORM Daniel J. Greenhoe page 209

[♠]Proof:

$\alpha \mathbb{Z}\check{\mathbf{x}}(z) \triangleq \alpha \mathbf{Z} (\mathbf{x}[n])$	by definition of $\check{x}(z)$	
$\triangleq \alpha \sum_{n} x[n] z^{-n}$	by definition of Z operator	
$\triangleq \sum_{n\in\mathbb{Z}}^{n\in\mathbb{Z}} (\alpha x[n]) z^{-n}$	by distributive property	
$\triangleq \mathbf{Z} \left(\alpha x[n] \right)$	by definition of ${f Z}$ operator	
$z^{-k}\check{x}(z) = z^{-k}\mathbf{Z}\left(x[n]\right)$	by definition of $\check{\mathbf{x}}(z)$	(left hypothesis)
$\triangleq z^{-k} \sum_{n=-\infty}^{n=+\infty} x[n] z^{-n}$	by definition of ${f Z}$	(Definition J.4 page 208)
$= \sum_{n=-\infty}^{n=+\infty} x[n]z^{-n-k}$ $= \sum_{n=-\infty}^{m-k=+\infty} x[n]z^{-n-k}$		
$=\sum_{m-k=-\infty}x[m-k]z^{-m}$	where $m \triangleq n + k$	$\implies n = m - k$
$=\sum_{m=-\infty}^{m=+\infty} x[m-k]z^{-m}$		
$= \sum_{n=-\infty}^{n=+\infty} x[n-k]z^{-n}$	where $n \triangleq m$	
$\triangleq \mathbf{Z} \left(\left(\mathbf{x}[n-k] \right) \right)$	by definition of ${f Z}$	(Definition J.4 page 208)
$\mathbf{Z}(\mathbf{x}^*[n]) \triangleq \sum_{\mathbf{z} \in \mathbb{Z}^n} \mathbf{x}^*[n] \mathbf{z}^{-n}$	by definition of ${f Z}$	(Definition J.4 page 208)
$\triangleq \left(\sum_{n\in\mathbb{Z}} x[n](z^*)^{-n}\right)^*$	by definition of ${f Z}$	(Definition J.4 page 208)
$\triangleq \check{X}^*(z^*)$	by definition of ${f Z}$	(Definition J.4 page 208)
$\mathbf{Z}((x[-n])) \triangleq \sum_{n \in \mathbb{Z}} x[-n]z^{-n}$	by definition of ${f Z}$	(Definition J.4 page 208)
$=\sum_{-m\in\mathbb{Z}}x[m]z^m$	where $m \triangleq -n$	$\implies n = -m$
$=\sum_{m\in\mathbb{Z}}x[m]z^m$	by absolutely summable property	(Definition J.2 page 207)
$= \sum_{m \in \mathbb{Z}} x[m] \left(\frac{1}{z}\right)^{-m}$	by absolutely summable property	(Definition J.2 page 207)
$\triangleq \check{x}\left(\frac{1}{7}\right)$	by definition of ${f Z}$	(Definition J.4 page 208)
$\mathbf{Z}\left(\mathbf{x}^*[-n]\right) \triangleq \sum_{n \in \mathbb{Z}} \mathbf{x}^*[-n] z^{-n}$	by definition of ${f Z}$	(Definition J.4 page 208)
$=\sum_{-m\in\mathbb{Z}}x^*[m]z^m$	where $m \triangleq -n$	$\implies n = -m$
$=\sum_{m\in\mathbb{Z}}x^*[m]z^m$	by absolutely summable property	(Definition J.2 page 207)
$= \sum_{m \in \mathbb{Z}} x^*[m] \left(\frac{1}{z}\right)^{-m}$	by absolutely summable property	(Definition J.2 page 207)
$= \left(\sum_{m \in \mathbb{Z}} x[m] \left(\frac{1}{z^*}\right)^{-m}\right)^*$	by absolutely summable property	(Definition J.2 page 207)






Figure J.1: Direct form 1 order 2 IIR filters

$$\triangleq \check{\mathsf{x}}^* \left(\frac{1}{z^*} \right) \qquad \qquad \mathsf{by definition of } \mathbf{Z} \qquad \qquad \mathsf{(Definition J.4 page 208)}$$

Theorem J.2 (convolution theorem). Let \star be the convolution operator (Definition J.3 page 207).

$$\mathbf{Z} \underbrace{\left(\left((x_n) \star (y_n) \right) \right)}_{sequence\ convolution} = \underbrace{\left(\mathbf{Z} \left((x_n) \right) \left(\mathbf{Z} \left((y_n) \right) \right)}_{series\ multiplication} \qquad \forall (x_n)_{n \in \mathbb{Z}}, (y_n)_{n \in \mathbb{Z}} \in \mathcal{C}^2_{\mathbb{R}}$$

[♠]Proof:

$$[\mathbf{Z}(x \star y)](z) \triangleq \mathbf{Z} \left(\sum_{m \in \mathbb{Z}} x_m y_{n-m} \right) \qquad \text{by definition of } \star \qquad \text{(Definition J.3 page 207)}$$

$$\triangleq \sum_{n \in \mathbb{Z}} \sum_{m \in \mathbb{Z}} x_m y_{n-m} z^{-n} \qquad \text{by definition of } \mathbf{Z} \qquad \text{(Definition J.4 page 208)}$$

$$= \sum_{n \in \mathbb{Z}} \sum_{m \in \mathbb{Z}} x_m y_{n-m} z^{-n} \qquad = \sum_{m \in \mathbb{Z}} \sum_{n \in \mathbb{Z}} x_m y_{n-m} z^{-n}$$

$$= \sum_{m \in \mathbb{Z}} \sum_{k \in \mathbb{Z}} x_m y_k z^{-(m+k)} \qquad \text{where } k \triangleq n-m \qquad \Longleftrightarrow n = m+k$$

$$= \left[\sum_{m \in \mathbb{Z}} x_m z^{-m} \right] \left[\sum_{k \in \mathbb{Z}} y_k z^{-k} \right]$$

$$\triangleq \left[\mathbf{Z} \left(x_n \right) \right] \left[\mathbf{Z} \left(y_n \right) \right] \qquad \text{by definition of } \mathbf{Z} \qquad \text{(Definition J.4 page 208)}$$

J.3 From z-domain back to time-domain

$$\check{\mathbf{y}}(z) = b_0 X(z) + b_1 z^{-1} X(z) + b_2 z^{-2} X(z) - a_1 z^{-1} Y(z) + a_2 z^{-2} Y(z)$$

$$y[n] = b_0x[n] + b_1x[n-1] + b_2x[n-2] - a_1y[n-1] - a_2y[n-2]$$

Example J.1. See Figure J.1 (page 210)

$$\frac{3z^2 + 5z + 7}{2z^2 + 10z + 12} = \frac{3z^2 + 5z + 7}{2(z^2 + 5z + 6)} = \frac{\left(3hz^2 + 5hz + 7h\right)}{z^2 + 5z + 6} = \frac{\left(3hz^2 + 5hz + 7hz\right)}{1 + 5z^{-1} + 6z^{-2}}$$



J.4. ZERO LOCATIONS Daniel J. Greenhoe page 211

J.4 Zero locations

The system property of *minimum phase* is defined in Definition J.5 (next) and illustrated in Figure J.2 (page 211).




Figure J.2: Minimum Phase filter

Definition J.5. ⁴ Let $\check{\mathbf{x}}(z) \triangleq \mathbf{Z}(x_n)$ be the Z TRANSFORM (Definition J.4 page 208) of a sequence $(x_n)_{n \in \mathbb{Z}}$ in $\mathscr{C}^2_{\mathbb{R}}$. Let $(z_n)_{n \in \mathbb{Z}}$ be the ZEROS of $\check{\mathbf{x}}(z)$.

```
The sequence (x_n) is minimum phase if  |z_n| < 1 \quad \forall n \in \mathbb{Z} 
 |x(z)| \text{ has all its zeros inside the unit circle}
```

The impulse response of a minimum phase filter has most of its energy concentrated near the beginning of its support, as demonstrated next.

Theorem J.3 (Robinson's Energy Delay Theorem). ⁵ Let $p(z) \triangleq \sum_{n=0}^{N} a_n z^{-n}$ and $q(z) \triangleq \sum_{n=0}^{N} b_n z^{-n}$ be polynomials.

$$\left\{ \begin{array}{l} \mathsf{p} \quad \text{is minimum phase} \\ \mathsf{q} \quad \text{is not } minimum \ phase \\ \end{array} \right\} \implies \sum_{n=0}^{m-1} \left| a_n \right|^2 \geq \sum_{n=0}^{m-1} \left| b_n \right|^2 \qquad \forall 0 \leq m \leq N$$

But for more *symmetry*, put some zeros inside and some outside the unit circle (Figure J.3 page 212).

Example J.2. An example of a minimum phase polynomial is the Daubechies-4 scaling function. The minimum phase polynomial causes most of the energy to be concentrated near the origin, making it very *asymmetric*. In contrast, the Symlet-4 has a design very similar to that of Daubechies-4, but the selected zeros are not all within the unit circle in the complex z plane. This results in a scaling function that is more symmetric and less contrated near the origin. Both scaling functions are illustrated in Figure J.3 (page 212).



⁴ Farina and Rinaldi (2000) page 91, Dumitrescu (2007) page 36

⁵ ☐ Dumitrescu (2007) page 36, ☐ Robinson (1962), ☐ Robinson (1966) ⟨???⟩, ☐ Claerbout (1976) pages 52–53




Figure J.3: Daubechies-4 and Symlet-4 scaling functions pole-zero plots

J.5 Pole locations

Definition J.6.

A filter (or system or operator) **H** is **causal** if its current output does not depend on future inputs.

Definition J.7.

A filter (or system or operator) **H** is **time-invariant** if the mapping it performs does not change with time.

Definition J.8.

An operation **H** is **linear** if any output y_n can be described as a linear combination of inputs x_n as in $y_n = \sum_{m \in \mathbb{Z}} h(m) x(n-m) .$

For a filter to be *stable*, place all the poles *inside* the unit circle.

Theorem J.4. A causal LTI filter is **stable** if all of its poles are **inside** the unit circle.

Example J.3. Stable/unstable filters are illustrated in Figure J.4 (page 213).

True or False? This filter has no poles:

$$H(z) = b_0 + b_1 z^{-1} + b_2 z^{-2}$$

$$\mathsf{H}(z) = b_0 + b_1 z^{-1} + b_2 z^{-2} = \frac{z^2}{z^2} \times \frac{b_0 + b_1 z^{-1} + b_2 z^{-2}}{1} = \frac{b_0 z^2 + b_1 z^1 + b_2 z^{-2}}{z^2}$$






Figure J.4: Pole-zero plot stable/unstable causal LTI filters (Example J.3 page 212)




Figure J.5: FIR filters

J.6 Mirroring for real coefficients

If you want real coefficients, choose poles and zeros in conjugate pairs (next).

Proposition J.3.



[♠]Proof:

$$(z - p_1)(z - p_1^*) = [z - (a + ib)][z - (a - ib)]$$
$$= z^2 + [-a + ib - ib - a]z - [ib]^2$$




Figure J.6: Conjugate pair structure yielding real coefficients

$$= z^2 - 2az + b^2$$

Example J.4. See Figure J.6 (page 213).

$$\begin{split} H(z) &= G\frac{\left[z-z_1\right]\left[z-z_2\right]}{\left[z-p_1\right]\left[z-p_2\right]} = G\frac{\left[z-(1+i)\right]\left[z-(1-i)\right]}{\left[z-(-2/3+i^1/2)\right]\left[z-(-2/3-i^1/2)\right]} \\ &= G\frac{z^2-z\left[(1-i)+(1+i)\right]+(1-i)(1+i)}{z^2-z\left[(-2/3+i^1/2)+(-2/3+i^1/2)\right]+(-2/3+i^1/2)} \\ &= G\frac{z^2-2z+2}{z^2-4/3z+(4/3+1/4)} = G\frac{z^2-2z+2}{z^2-4/3z+1^9/12} \end{split}$$

J.7 Rational polynomial operators

A digital filter is simply an operator on $\mathscr{E}^2_{\mathbb{R}}$. If the digital filter is a causal LTI system, then it can be expressed as a rational polynomial in z as shown next.

Lemma J.1. A causal LTI operator **H** can be expressed as a rational expression $\check{h}(z)$.

$$\begin{split} \check{\mathbf{h}}(z) &= \frac{b_0 + b_1 z^{-1} + b_2 z^{-2} + \dots + b_N z^{-N}}{1 + a_1 z^{-1} + a_2 z^{-2} + \dots + a_N z^{-N}} \\ &= \frac{\sum\limits_{n=0}^{N} b_n z^{-n}}{1 + \sum\limits_{n=1}^{N} a_n z^{-n}} \end{split}$$

A filter operation $\check{h}(z)$ can be expressed as a product of its roots (poles and zeros).

$$\begin{split} \check{\mathbf{h}}(z) &= \frac{b_0 + b_1 z^{-1} + b_2 z^{-2} + \dots + b_N z^{-N}}{1 + a_1 z^{-1} + a_2 z^{-2} + \dots + a_N z^{-N}} \\ &= \alpha \frac{(z - z_1)(z - z_2) \dots (z - z_N)}{(z - p_1)(z - p_2) \dots (z - p_N)} \end{split}$$

where α is a constant, z_i are the zeros, and p_i are the poles. The poles and zeros of such a rational expression are often plotted in the z-plane with a unit circle about the origin (representing $z = e^{i\omega}$). Poles are marked with \times and zeros with \bigcirc . An example is shown in Figure J.7 page 215. Notice that in this figure the zeros and poles are either real or occur in complex conjugate pairs.

J.8 Filter Banks

Conjugate quadrature filters (next definition) are used in filter banks. If $\check{x}(z)$ is a low-pass filter, then the conjugate quadrature filter of $\check{y}(z)$ is a high-pass filter.

page 215 J.8. FILTER BANKS Daniel J. Greenhoe




Figure J.7: Pole-zero plot for rational expression with real coefficients

Definition J.9. ⁶ Let $(x_n)_{n\in\mathbb{Z}}$ and $(y_n)_{n\in\mathbb{Z}}$ be SEQUENCES (Definition J.1 page 207) in $\mathscr{C}^2_{\mathbb{R}}$ (Definition J.2 page 207). The sequence (y_n) is a **conjugate quadrature filter** with shift N with respect to (x_n) if

 $y_n = \pm (-1)^n x_{N-n}^*$

A CONJUGATE QUADRATURE FILTER is also called a **CQF** or a **Smith-Barnwell filter**. Any triple $((x_n), (y_n), N)$ in this form is said to satisfy the

conjugate quadrature filter condition or the CQF condition.

Theorem J.5 (CQF theorem). ⁷ Let $\check{y}(\omega)$ and $\check{x}(\omega)$ be the DTFTs (Definition M.1 page 237) of the sequences $(y_n)_{n\in\mathbb{Z}}$ and $(x_n)_{n\in\mathbb{Z}}$, respectively, in $\ell_{\mathbb{R}}^2$ (Definition J.2 page 207).

$$(y_n)_{n\in\mathbb{Z}} \ and \ (x_n)_{n\in\mathbb{Z}}, \ respectively, \ in \ \mathscr{E}_{\mathbb{R}}^{} \ (Definition J.2 \ page 207).$$

$$y_n = \pm (-1)^n x_{N-n}^* \qquad \Longleftrightarrow \qquad \check{y}(z) = \pm (-1)^N z^{-N} \check{x}^* \left(\frac{-1}{z^*}\right) \qquad (2) \quad \text{CQF in "z-domain"}$$

$$\iff \check{y}(\omega) = \pm (-1)^N e^{-i\omega N} \check{x}^* (\omega + \pi) \qquad (3) \quad \text{CQF in "frequency"}$$

$$\iff x_n = \pm (-1)^N (-1)^n y_{N-n}^* \qquad (4) \quad \text{"reversed" CQF in "time"}$$

$$\iff \check{x}(z) = \pm z^{-N} \check{y}^* \left(\frac{-1}{z^*}\right) \qquad (5) \quad \text{"reversed" CQF in "z-domain"}$$

$$\iff \check{x}(\omega) = \pm e^{-i\omega N} \check{y}^* (\omega + \pi) \qquad (6) \quad \text{"reversed" CQF in "frequency"}$$

$$\forall N \in \mathbb{Z}$$

^ℚProof:

D E F

1. Proof that $(1) \implies (2)$:

$$\begin{split} \check{\mathbf{y}}(z) &= \sum_{n \in \mathbb{Z}} y_n z^{-n} & \text{by definition of } z\text{-}transform \\ &= \sum_{n \in \mathbb{Z}} (\pm)(-1)^n x_{N-n}^* z^{-n} & \text{by (1)} \\ &= \pm \sum_{m \in \mathbb{Z}} (-1)^{N-m} x_m^* z^{-(N-m)} & \text{where } m \triangleq N-n \implies n = N-m \\ &= \pm (-1)^N z^{-N} \sum_{m \in \mathbb{Z}} (-1)^{-m} x_m^* \left(z^{-1}\right)^{-m} \\ &= \pm (-1)^N z^{-N} \sum_{m \in \mathbb{Z}} x_m^* \left(-\frac{1}{z}\right)^{-m} \\ &= \pm (-1)^N z^{-N} \left[\sum_{m \in \mathbb{Z}} x_m \left(-\frac{1}{z^*}\right)^{-m}\right]^* \end{split}$$

⁶ Strang and Nguyen (1996) page 109, Haddad and Akansu (1992) pages 256–259 ⟨section 4.5⟩, Vaidyanathan (1993) page 342 ⟨(7.2.7), (7.2.8)⟩, Smith and Barnwell (1984a), Smith and Barnwell (1984b), Mintzer (1985) ⁷

■ Strang and Nguyen (1996) page 109,
■ Mallat (1999) pages 236–238 ⟨(7.58),(7.73)⟩,
■ Haddad and Akansu (1992) pages 256–259 (section 4.5), ■ Vaidyanathan (1993) page 342 ((7.2.7), (7.2.8))





$$= \pm (-1)^{N} z^{-N} \check{\mathbf{x}}^* \left(\frac{-1}{z^*} \right)$$

by definition of *z-transform*

(Definition J.4 page 208)

2. Proof that $(1) \iff (2)$:

$$\dot{\mathbf{y}}(z) = \pm (-1)^N z^{-N} \dot{\mathbf{x}}^* \left(\frac{-1}{z^*}\right) \qquad \text{by (2)}$$

$$= \pm (-1)^N z^{-N} \left[\sum_{m \in \mathbb{Z}} x_m \left(\frac{-1}{z^*}\right)^{-m} \right]^* \qquad \text{by definition of } z\text{-}transform \qquad \text{(Definition J.4 page 208)}$$

$$= \pm (-1)^N z^{-N} \left[\sum_{m \in \mathbb{Z}} x_m^* \left(-z^{-1}\right)^{-m} \right] \qquad \text{by definition of } z\text{-}transform \qquad \text{(Definition J.4 page 208)}$$

$$= \sum_{m \in \mathbb{Z}} (\pm) (-1)^{N-m} x_m^* z^{-(N-m)}$$

$$= \sum_{m \in \mathbb{Z}} (\pm) (-1)^n x_{N-n}^* z^{-n} \qquad \text{where } n = N - m \implies m \triangleq N - n$$

$$\Rightarrow x_n = \pm (-1)^n x_{N-n}^*$$

3. Proof that $(1) \implies (3)$:

$$\begin{split} \breve{\mathbf{y}}(\omega) &\triangleq \breve{\mathbf{x}}(z) \Big|_{z=e^{i\omega}} & \text{by definition of } DTFT \text{ (Definition M.1 page 237)} \\ &= \left[\pm (-1)^N z^{-N} \breve{\mathbf{x}} \left(\frac{-1}{z^*} \right) \right]_{z=e^{i\omega}} & \text{by (2)} \\ &= \pm (-1)^N e^{-i\omega N} \breve{\mathbf{x}} \left(e^{i\pi} e^{i\omega} \right) \\ &= \pm (-1)^N e^{-i\omega N} \breve{\mathbf{x}} \left(e^{i(\omega+\pi)} \right) \\ &= \pm (-1)^N e^{-i\omega N} \breve{\mathbf{x}} (\omega+\pi) & \text{by definition of } DTFT \text{ (Definition M.1 page 237)} \end{split}$$

4. Proof that $(1) \implies (6)$:

$$\begin{split} &\check{\mathbf{x}}(\omega) = \sum_{n \in \mathbb{Z}} y_n e^{-i\omega n} & \text{by definition of } DTFT & \text{(Definition M.1 page 237)} \\ &= \sum_{n \in \mathbb{Z}} \pm (-1)^n x_{N-n}^* e^{-i\omega n} & \text{by (1)} \\ &= \sum_{m \in \mathbb{Z}} \pm (-1)^{N-m} x_m^* e^{-i\omega (N-m)} & \text{where } m \triangleq N-n \implies n = N-m \\ &= \pm (-1)^N e^{-i\omega N} \sum_{m \in \mathbb{Z}} (-1)^m x_m^* e^{i\omega m} \\ &= \pm (-1)^N e^{-i\omega N} \sum_{m \in \mathbb{Z}} e^{i\pi m} x_m^* e^{i\omega m} \\ &= \pm (-1)^N e^{-i\omega N} \sum_{m \in \mathbb{Z}} x_m^* e^{i(\omega + \pi)m} \\ &= \pm (-1)^N e^{-i\omega N} \left[\sum_{m \in \mathbb{Z}} x_m e^{-i(\omega + \pi)m} \right]^* \\ &= \pm (-1)^N e^{-i\omega N} \check{\mathbf{x}}^* (\omega + \pi) & \text{by definition of } DTFT & \text{(Definition M.1 page 237)} \end{split}$$

5. Proof that $(1) \Leftarrow (3)$:

$$y_n = \frac{1}{2\pi} \int_{-\pi}^{+\pi} \check{y}(\omega) e^{i\omega n} \, d\omega \qquad \qquad \text{by } inverse \, DTFT \qquad \text{(Theorem M.3 page 243)}$$

$$= \frac{1}{2\pi} \int_{-\pi}^{+\pi} \underbrace{\pm (-1)^N e^{-iN\omega} \check{x}^*(\omega + \pi) e^{i\omega n}}_{\text{right hypothesis}} \, d\omega \qquad \qquad \text{by right hypothesis}$$

$$= \pm (-1)^N \frac{1}{2\pi} \int_{-\pi}^{+\pi} \check{x}^*(\omega + \pi) e^{i\omega(n-N)} \, d\omega \qquad \qquad \text{by right hypothesis}$$

$$= \pm (-1)^N \frac{1}{2\pi} \int_0^{2\pi} \check{x}^*(v) e^{i(v-\pi)(n-N)} \, dv \qquad \qquad \text{where } v \triangleq \omega + \pi \implies \omega = v - \pi$$

$$= \pm (-1)^N e^{-i\pi(n-N)} \frac{1}{2\pi} \int_0^{2\pi} \check{x}^*(v) e^{iv(n-N)} \, dv$$

$$= \pm (-1)^N \underbrace{(-1)^N (-1)^N (-1)^n}_{e^{-i\pi n}} \left[\frac{1}{2\pi} \int_0^{2\pi} \check{x}(v) e^{iv(N-n)} \, dv \right]^*$$

$$= \pm (-1)^n x_{N-n}^* \qquad \qquad \text{by } inverse \, DTFT \qquad \text{(Theorem M.3 page 243)}$$

6. Proof that $(1) \iff (4)$:

$$y_{n} = \pm (-1)^{n} x_{N-n}^{*} \iff (\pm)(-1)^{n} y_{n} = (\pm)(\pm)(-1)^{n} (-1)^{n} x_{N-n}^{*}$$

$$\iff \pm (-1)^{n} y_{n} = x_{N-n}^{*}$$

$$\iff (\pm (-1)^{n} y_{n})^{*} = (x_{N-n}^{*})^{*}$$

$$\iff \pm (-1)^{n} y_{n}^{*} = x_{N-n}$$

$$\iff x_{N-n} = \pm (-1)^{n} y_{n}^{*}$$

$$\iff x_{m} = \pm (-1)^{N-m} y_{N-m}^{*}$$

$$\iff x_{m} = \pm (-1)^{N-m} y_{N-m}^{*}$$

$$\iff x_{m} = \pm (-1)^{N} (-1)^{m} y_{N-m}^{*}$$

$$\iff x_{n} = \pm (-1)^{N} (-1)^{m} y_{N-n}^{*}$$

$$\iff x_{n} = \pm (-1)^{N} (-1)^{n} y_{N-n}^{*}$$

7. Proofs for (5) and (6): not included. See proofs for (2) and (3).

Theorem J.6. ⁸ Let $\check{y}(\omega)$ and $\check{x}(\omega)$ be the DTFTs (Definition M.1 page 237) of the sequences $(y_n)_{n\in\mathbb{Z}}$ and

Theorem J.6. ⁸ Let
$$\S(\omega)$$
 and $\S(\omega)$ be the DTFTs (Definition M.1 page 237) of the set $(x_n)_{n\in\mathbb{Z}}$, respectively, in $\mathscr{C}^2_{\mathbb{R}}$ (Definition J.2 page 207).

Let $y_n = \pm (-1)^n x_{N-n}^*$ (CQF CONDITION, Definition J.9 page 215). Then
$$\begin{cases}
(A) & \left[\frac{\mathsf{d}}{\mathsf{d}\omega}\right]^n \S(\omega)\right|_{\omega=0} = 0 \iff \left[\frac{\mathsf{d}}{\mathsf{d}\omega}\right]^n \S(\omega)\right|_{\omega=\pi} = 0 & \text{(B)} \\
\Leftrightarrow \sum_{k\in\mathbb{Z}} (-1)^k k^n x_k = 0 & \text{(C)} \\
\Leftrightarrow \sum_{k\in\mathbb{Z}} k^n y_k = 0 & \text{(D)}
\end{cases}$$

^ℚProof:

⁸ Vidakovic (1999) pages 82–83, A Mallat (1999) pages 241–242



1. Proof that (A) \Longrightarrow (B):


$$\begin{array}{lll} 0 = \left[\frac{\mathrm{d}}{\mathrm{d}\omega} \right]^n \check{\mathbf{y}}(\omega) \Big|_{\omega=0} & \text{by (A)} \\ = \left[\frac{\mathrm{d}}{\mathrm{d}\omega} \right]^n (\pm) (-1)^N e^{-i\omega N} \check{\mathbf{x}}^* (\omega + \pi) \Big|_{\omega=0} & \text{by CQF theorem} & \text{(Theorem J.5 page 215)} \\ = (\pm) (-1)^N \sum_{\ell=0}^n \binom{n}{\ell} \left[\frac{\mathrm{d}}{\mathrm{d}\omega} \right]^\ell \left[e^{-i\omega N} \right] \cdot \left[\frac{\mathrm{d}}{\mathrm{d}\omega} \right]^{n-\ell} \left[\check{\mathbf{x}}^* (\omega + \pi) \right] \Big|_{\omega=0} & \text{by $Leibnitz$ GPR} & \text{(Lemma D.2 page 143)} \\ = (\pm) (-1)^N \sum_{\ell=0}^n \binom{n}{\ell} -i N^\ell e^{-i\omega N} \left[\frac{\mathrm{d}}{\mathrm{d}\omega} \right]^{n-\ell} \left[\check{\mathbf{x}}^* (\omega + \pi) \right] \Big|_{\omega=0} & \\ = (\pm) (-1)^N e^{-i0N} \sum_{\ell=0}^n \binom{n}{\ell} -i N^\ell \left[\frac{\mathrm{d}}{\mathrm{d}\omega} \right]^{n-\ell} \left[\check{\mathbf{x}}^* (\omega + \pi) \right] \Big|_{\omega=0} & \\ & \Longrightarrow \check{\mathbf{x}}^{(0)}(\pi) = 0 & \\ & \Longrightarrow \check{\mathbf{x}}^{(1)}(\pi) = 0 & \\ & \Longrightarrow \check{\mathbf{x}}^{(2)}(\pi) = 0 & \\ & \Longrightarrow \check{\mathbf{x}}^{(3)}(\pi) = 0 & \\ & \Longrightarrow \check{\mathbf{x}}^{(4)}(\pi) = 0 & \\ & \vdots & \vdots & \\ & \Longrightarrow \check{\mathbf{x}}^{(n)}(\pi) = 0 & \text{for $n=0,1,2,\dots$} \end{array}$$

2. Proof that (A) \Leftarrow (B):

$$\begin{array}{lll} 0 = \left[\frac{\mathrm{d}}{\mathrm{d} \mathrm{o}}\right]^n \check{\mathbf{x}}(\omega) \Big|_{\omega = \pi} & \text{by (B)} \\ = \left[\frac{\mathrm{d}}{\mathrm{d} \mathrm{o}}\right]^n (\pm) e^{-i\omega N} \check{\mathbf{y}}^* (\omega + \pi) \Big|_{\omega = \pi} & \text{by } CQF \ theorem & \text{(Theorem J.5 page 215)} \\ = (\pm) \sum_{\ell = 0}^n \binom{n}{\ell} \left[\frac{\mathrm{d}}{\mathrm{d} \mathrm{o}}\right]^\ell \left[e^{-i\omega N}\right] \cdot \left[\frac{\mathrm{d}}{\mathrm{d} \mathrm{o}}\right]^{n-\ell} \left[\check{\mathbf{y}}^* (\omega + \pi)\right] \Big|_{\omega = \pi} & \text{by } Leibnitz \ GPR & \text{(Lemma D.2 page 143)} \\ = (\pm) \sum_{\ell = 0}^n \binom{n}{\ell} (-iN)^\ell e^{-i\omega N} \left[\frac{\mathrm{d}}{\mathrm{d} \mathrm{o}}\right]^{n-\ell} \left[\check{\mathbf{y}}^* (\omega + \pi)\right] \Big|_{\omega = \pi} & \\ = (\pm) e^{-i\pi N} \sum_{\ell = 0}^n \binom{n}{\ell} -iN^\ell \left[\frac{\mathrm{d}}{\mathrm{d} \mathrm{o}}\right]^{n-\ell} \left[\check{\mathbf{y}}^* (\omega + \pi)\right] \Big|_{\omega = \pi} & \\ = (\pm) (-1)^N \sum_{\ell = 0}^n \binom{n}{\ell} -iN^\ell \left[\frac{\mathrm{d}}{\mathrm{d} \mathrm{o}}\right]^{n-\ell} \left[\check{\mathbf{y}}^* (\omega + \pi)\right] \Big|_{\omega = \pi} & \\ & \Longrightarrow \ \check{\mathbf{y}}^{(0)}(0) = 0 & \\ & \Longrightarrow \ \check{\mathbf{y}}^{(1)}(0) = 0 & \\ & \Longrightarrow \ \check{\mathbf{y}}^{(3)}(0) = 0 & \\ & \Longrightarrow \ \check{\mathbf{y}}^{(4)}(0) = 0 & \\ & \Longrightarrow \ \check{\mathbf{y}}^{(n)}(0) = 0 & \text{for } n = 0, 1, 2, \dots \end{array}$$


- 3. Proof that (B) \iff (C): by Theorem M.5 page 245
- 4. Proof that (A) \iff (D): by Theorem M.5 page 245
- 5. Proof that (CQF) \Leftarrow (A): Here is a counterexample: $\check{y}(\omega) = 0$.





J.9 Inverting non-minimum phase filters

Minimum phase filters are easy to invert: each zero becomes a pole and each pole becomes a zero.



$$\begin{split} |A(z)|_{z=e^{i\omega}} &= \frac{1}{r} \left| \frac{z - re^{i\phi}}{z - \frac{1}{r}e^{i\phi}} \right|_{z=e^{i\omega}} \\ &= \left| e^{i\phi} \left(\frac{e^{-i\phi}z - r}{rz - e^{i\phi}} \right) \right|_{z=e^{i\omega}} \\ &= \left| -z \left(\frac{rz^{-1} - e^{-i\phi}}{rz - e^{i\phi}} \right) \right|_{z=e^{i\omega}} \\ &= \left| \frac{1}{e^{-iv}} \left(\frac{re^{-i\omega} - e^{-i\phi}}{(re^{i\omega} - e^{i\phi})^*} \right) \right| \\ &= \frac{1}{e^{-iv}} \left(\frac{re^{-i\omega} - e^{-i\phi}}{(re^{i\omega} - e^{i\phi})^*} \right) \\ &= 1 \end{split}$$

page 220 Daniel J. Greenhoe APPENDIX J. Z TRANSFORM







TAYLOR EXPANSIONS (TRANSFORMS)

K.1 Introduction

For modeling real-world processes above the quantum level, measurements are *continuous* in time—that is, the first derivative of a function over time representing the measurement *exists*.

But even for "simple" physical systems, it is not just the first derivative that matters. For example, the classical "vibrating string" vertical displacement u(x,t) wave equation can be described as

$$\frac{\partial^2 u}{\partial x^2} - \frac{1}{c^2} \frac{\partial^2 u}{\partial t^2} = 0$$

Not only do physical systems demonstrate heavy dependence on the derivatives of their measurement functions, but also commonly exhibit *oscillation*, as demonstrated by sunspot activity over the last 300 years or earthquake activity (Figure K.1 page 222).

In fact, derivatives and oscillations are fundamentally linked as demonstrated by the fact that all solutions of homogeneous second order differential equations are linear combinations of sine and cosine functions (Theorem G.3 page 158):

cosine functions (Theorem G.3 page 158):
$$\left\{\frac{\mathrm{d}^2}{\mathrm{d}x^2}\mathsf{f} + \mathsf{f} = 0\right\} \iff \left\{\mathsf{f}(x) = \mathsf{f}(0)\cos(x) + \mathsf{f}'(0)\sin(x)\right\} \qquad \forall \mathsf{f} \in \mathcal{C}, \forall x \in \mathbb{R}$$

Derivatives are calculated *locally* about a point. Oscillations are observed *globally* over a range, and analyzed (decomposed) by projecting the function onto a sequence of basis functions—sinusoids in the case of Fourier Transform family. Projection is accomplished using inner products, and often these are calculated using *integration*. Note that derivatives and integrals are also fundamentally linked as demonstrated by the *Fundamental Theorem of Calculus*…which shows that integration can be calculated using anti-differentiation:

$$\int_{a}^{b} f(x) dx = F(b) - F(a) \quad \text{where } F(x) \text{ is the } antiderivative \text{ of } f(x).$$

Brook Taylor showed that for *analytic* functions, knowledge of the derivatives of a function at a location x = a allows you to determine (predict) arbitrarily closely all the points f(x) in the vicinity

¹ analytic functions: Functions for which all their derivatives exist.

Daniel J. Greenhoe




Figure K.1: Sunspot and earthquake measurements

of
$$x = a$$
:²

$$f(x) = f(a) + \frac{1}{1!}f'(a)[x - a] + \frac{1}{2!}f''(x)[x - a]^2 + \frac{1}{3!}f'''(x)[x - a]^3 + \cdots$$

On the other hand, the Fourier Transform is a kind of counter-part of the Taylor expansion:³

On	On the other name, the rounter transjorm is a kind of counter part of the rayior expe					
	Taylor coefficients	Fourier coefficients				
	Depend on derivatives $\frac{d^n}{dx^n} f(x)$	Depend on integrals $\int_{-\infty}^{\infty} f(x)e^{-i\omega x} dx$				
	Behavior in the vicinity of a point.	$J_{x \in \mathbb{R}}$ Behavior over the entire function.				
44	Demonstrate trends locally.	Demonstrate trends globally, such as oscillations.				
44	Admits <i>analytic</i> functions only.	Admits <i>non-analytic</i> functions as well.				
44	Function must be <i>continuous</i> .	Function can be <i>discontinuous</i> .				

K.2 Taylor Expansion

Theorem K.1 (Taylor Series). ⁴ Let C be the space of all ANALYTIC functions and $\frac{d}{dt}$ in C^C the DIF-FERENTIATION OPERATOR.

A Taylor Series about the point
$$x = a$$
 of a function $f(x) \in C^C$ is
$$f(x) = \sum_{n=0}^{\infty} \frac{\left[\frac{d}{dx}^n f\right](a)}{n!} \underbrace{(x-a)^n}_{basis function} \forall a \in \mathbb{R}, f \in C$$
A Maclaurin Series is a Taylor Series about the point $a = 0$.

² Robinson (1982) page 886

³ Robinson (1982) page 886

⁴ Flanigan (1983) page 221 ⟨Theorem 15⟩, Strichartz (1995) page 281, Sohrab (2003) page 317 ⟨Theorem 8.4.9), Taylor (1715), Taylor (1717), Maclaurin (1742)

La langue de l'analyse, la plus parfaite de toutes les langues, tant par elle-même un puissant instrument de découvertes; ses notations, lorsqu'elles sont nécessaires et heureusement imaginées, sont des germes de nouveaux calculs.

"The language of analysis, most perfect of all, being in itself a powerful instrument of discoveries, its notations, especially when they are necessary and happily imagined, are the seeds of new calculi."

Pierre-Simon Laplace¹

L.1 Operator Definition

Definition L.1. ² Let $L^2_{(\mathbb{R},\mathcal{B},\mu)}$ be the space of all Lebesgue square-integrable functions.

D E F The **Laplace Transform** operator **L** is here defined as $[\mathbf{Lf}](s) \triangleq \int_{x \in \mathbb{R}} \mathsf{f}(x)e^{-sx} \, \mathsf{dx} \qquad \forall \mathsf{f} \in L^2_{(\mathbb{R}, \mathcal{B}, \mu)}$

Such integrals may *converge* for certain values of *s* and *diverge* for others.

Definition L.2. Let L[g(x)] be the LAPLACE TRANSFORM (Definition L.1 page 223) of a function g(x).

The set $\mathbf{R} \propto \mathbf{L}[\mathbf{g}(x)]$ of all s for which $\mathbf{L}[\mathbf{g}(x)]$ CONVERGES is the **Region of Convergence** of $\mathbf{L}[\mathbf{g}(x)]$.

In this text, the region of convergence may in places be specified using the *open interval* (A : B) and *closed interval* [A : B].

¹ ■ Laplace (1814) page xxxi ⟨Introduction⟩, ■ Laplace (1812), ■ Laplace (1902) pages 48–49, ■ Moritz (1914) page 200 ⟨Quote 1222., but "conceived" not "imagined", and "are so many germs" not "are the seeds"⟩, https://todayinsci.com/L/Laplace_Pierre-LaplacePierre-Analysis-Quotations.htm, https://translate.google.com/,

² Bracewell (1978) page 219 ⟨Chapter 11 The Laplace transform⟩, ② van der Pol and Bremmer (1959) page 13 ⟨5. Strip of convergence of the Laplace integral⟩, ② Levy (1958) page 2 ⟨"two-sided transformation"⟩, ② Betten (2008b) page 295 ⟨(B.1)⟩

Remark L.1. A scaling factor $\frac{1}{\sqrt{2\pi}}$ in front of $\int_{\mathbb{R}}$ in Definition L.1 is not typically found in references offering definitions of the Laplace Transform, and is not included here either. That is not to say, however, that it's not a good idea. Including it would make the operator \mathbf{L} more directly compatible with the *Unitary Fourier Transform* operator $\tilde{\mathbf{F}}$ (Definition I.2 page 196). Note also that a $\frac{1}{2\pi}$ scaling factor is included in [a Bachman et al. (2002) page 268] in their definition of *convolution* (Definition E.1 page 145, Section L.8 page 235).

L.2 Operator Inverse

Theorem L.1. ³

$$\mathsf{g}(x) = \mathbf{L}^{-1}[\mathsf{G}(s)] \triangleq \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \mathsf{G}(s) e^{sx} ds \quad \textit{for some } c \in \mathbb{R}^+$$

L.3 Transversal properties

Theorem L.2. ⁴ Let $\mathbf{L}[\mathbf{g}(x)]$ be the Laplace Transform (Definition L.1 page 223) of a function $\mathbf{g}(x)$. Let the Region of Convergence of $\mathbf{L}[\mathbf{g}(x)](s)$ be $A \leq \mathbf{R}_{\mathbf{e}}(s) \leq B$ with $(A, B) \in \mathbb{R}^2$.

			Мар	pping	_	Region of (Convergence	Domain	Property
		$\mathbf{L}\left[\mathbf{g}(x-\alpha)\right]$	=	$e^{-\alpha s} \mathbf{L} [\mathbf{g}(x)](s)$	for	$\mathbf{R}_{\mathbf{e}}(s)$	$\in [A:B]$	$\forall x, \alpha \in \mathbb{C}$	(TRANSLATION)
Ň	Å	$\mathbf{L}\big[g(\alpha x)\big]$	=	$\frac{1}{ \alpha } \mathbf{L} \big[g(x) \big] \left(\frac{s}{\alpha} \right)$	for	$\mathbf{R}_{e}\left(\frac{s}{\alpha}\right)$	$\in [A:B]$	$\forall x, \alpha \in \mathbb{C}$	(DILATION)

[♠]Proof:

⁴ Bracewell (1978) page 224 (Table 11.1: "Shift" and "Similarity" entries), ■ Levy (1958) page 15 (Equation 0.8)



 $^{^3}$ Bracewell (1978) page 220 (Chapter 11 The Laplace transform)

$$= \begin{cases} \frac{1}{\alpha} \int_{u=-\infty}^{u=\infty} \mathsf{g}(u) e^{-(s/\alpha)u} \, \, \mathrm{d}u & \text{if } \alpha \geq 0 \qquad \forall A \leq \mathbf{R}_{\mathsf{e}} \left(\frac{s}{\alpha}\right) \leq B \\ \frac{1}{\alpha} \int_{u=-\infty}^{u=-\infty} \mathsf{g}(u) e^{-(s/\alpha)u} \, \, \mathrm{d}u & \text{otherwise} \quad \forall A \leq \mathbf{R}_{\mathsf{e}} \left(\frac{s}{\alpha}\right) \leq B \end{cases}$$

$$= \begin{cases} \frac{1}{\alpha} \int_{u=-\infty}^{u=\infty} \mathsf{g}(u) e^{-(s/\alpha)u} \, \, \mathrm{d}u & \text{if } \alpha \geq 0 \qquad \forall A \leq \mathbf{R}_{\mathsf{e}} \left(\frac{s}{\alpha}\right) \leq B \\ -\frac{1}{\alpha} \int_{u=-\infty}^{u=\infty} \mathsf{g}(u) e^{-(s/\alpha)u} \, \, \mathrm{d}u & \text{otherwise} \quad \forall A \leq \mathbf{R}_{\mathsf{e}} \left(\frac{s}{\alpha}\right) \leq B \end{cases}$$

$$= \frac{1}{|\alpha|} \int_{x \in \mathbb{R}} \mathsf{g}(x) e^{-(s/\alpha)x} \, \, \mathrm{d}x \qquad \forall A \leq \mathbf{R}_{\mathsf{e}} \left(\frac{s}{\alpha}\right) \leq B \quad \text{by change of variable} \qquad u \to x$$

$$\triangleq \frac{1}{|\alpha|} \left[\mathbf{L} \mathsf{g}(x) \right] \left(\frac{s}{\alpha}\right) \qquad \forall A \leq \mathbf{R}_{\mathsf{e}} \left(\frac{s}{\alpha}\right) \leq B \quad \text{by definition of } \mathbf{L} \qquad \text{(Definition L.1 page 223)}$$

Corollary L.1. ⁵ *Let* L, G(s), *A*, and *B* be defined as in Theorem L.2 (page 224).

C	Mapping	Region of Convergence	Domain	Property
R	$\mathbf{L}\big[g(-x)\big] = G(-s)$	for $\mathbf{R}_{e}(s) \in [-B:-A]$	$\forall x, \alpha \in \mathbb{C}$	(REVERSAL)

№ Proof:

$$\begin{split} \mathbf{L}\big[\mathsf{g}(-x)\big] &= \mathbf{L}\big[\mathsf{g}([-1]x)\big] & \mathbf{R}_\mathsf{e}(s) \in [A:B] & \text{by definition of unary operator } - \\ &= \mathbf{L}\bigg[\frac{1}{|-1|}\mathsf{g}\bigg(\frac{x}{-1}\bigg)\bigg] & \mathbf{R}_\mathsf{e}\bigg(\frac{s}{-1}\bigg) \in [A:B] & \text{by } \textit{dilation property (Theorem L.2 page 224)} \\ &= \mathsf{G}(-s) & \mathbf{R}_\mathsf{e}(s) \in [-B:-A] \end{split}$$

L.4 Linear properties

Theorem L.3. ⁶ Let L be the Laplace Transform operator (Definition L.1 page 223). Let $G(s) \triangleq [Lg(x)]$ and $F(s) \triangleq [Lf(x)]$. Let the Region of Convergence of G(s) be $A \leq \mathbf{R}_{e}(s) \leq B$ and the Region of Convergence of G(s) be $G(s) \neq B$ and the Region of Convergence of G(s) be $G(s) \neq B$.

т	Mapping				Region of Convergence	Domain	Property
H	$\mathbf{L}\left[f(x) + g(x)\right]$	=	F(s) + G(s)	for	$\mathbf{R}_{\mathbf{e}}(s) \in [A : B] \cap [C : D]$	$\forall x, \alpha \in \mathbb{C}$	(ADDITIVE)
M	$\mathbf{L}[\alpha g(x)]$	=	$\alpha G(s)$	for	$\mathbf{R}_{e}(s) \in [A : B]$	$\forall x, \alpha \in \mathbb{C}$	(HOMOGENEOUS)

Corollary L.2 (Linear Properties). Let L be the Laplace Transform operator (Definition L.1 page 223). Let A and B be real numbers such that [A:B] is the Region of Convergence of $\mathbf{L}[g(x)]$. Let C and D be real numbers such that [C:D] is the Region of Convergence of $\mathbf{L}[f(x)]$. Let A_n and B_n be real numbers such that $[A_n:B_n]$ is the Region of Convergence of $\mathbf{L}[g_n(x)]$.



page 225

 $^{^5}$ Bracewell (1978) page 224 (Table 11.1 Theorems for the Laplace Transform)

⁶ Bracewell (1978) page 224 ⟨Table 11.1 Theorems for the Laplace Transform⟩, ☐ Betten (2008a) page 296 ⟨(B.6)⟩, ☐ Levy (1958) page 13 ⟨Equation 0.2⟩, ☐ van der Pol and Bremmer (1959) page 22 ⟨Introduction⟩, ☐ Shafii-Mousavi (2015) page 7 ⟨Theorem 1.4⟩

		Мар	ping		Region of Convergence	Domain
	L [0]	=	0	for	$\mathbf{R}_{\mathbf{e}}(s) \in [A : B]$	$\forall x \in \mathbb{C}$
С	$\mathbf{L}[-g(x)]$	=	$-\mathbf{L}[\mathbf{g}(x)]$	for	$\mathbf{R}_{\mathbf{e}}(s) \in [A : B]$	$\forall x \in \mathbb{C}$
O R	$\mathbf{L}\left[f(x)-g(x)\right]$	=	$\mathbf{L}[g(x)] - \mathbf{L}[f(x)]$	for	$\mathbf{R}_{e}(s) \in [A : B] \cap [C : D]$	$\forall x \in \mathbb{C}$
n	<u>[</u>		<u>N</u>		N	
	$\mathbf{L} \sum \alpha_n g_n(x) \Big $	=	$\sum \alpha_n \mathbf{L} \big[g_n(x) \big]$	for	$\mathbf{R}_{\mathbf{e}}(s) \in \bigcap \left[A_n : B_n \right]$	$\forall x, \alpha_n {\in} \mathbb{C}$
	<u>n=1</u>		n=1		n=1	

[№]PROOF:

- 1. By Theorem L.3 (page 225), the operator Laplace Transform operator L is additive and homogeneous.
- 2. By item (1) and Definition C.4 (page 113), L is *linear*.
- 3. By item (2) and Theorem C.1 (page 113), the four properties listed follow.

L.5 Modulation properties

Theorem L.4. ⁷ Let L be the Laplace Transform operator (Definition L.1 page 223). Let $G(s) \triangleq [Lg(x)]$. Let the Region of Convergence of G(s) be $A \leq \mathbf{R}_{e}(s) \leq B$.

Т	Mapping		Region of Convergence	Domain	Property
H	$\frac{\mathbf{L}\left[e^{-\alpha x}\mathbf{g}(x)\right]}{\mathbf{L}\left[e^{-\alpha x}\mathbf{g}(x)\right]} =$	$G(s + \alpha)$	for $A - \mathbf{R}_{e}(\alpha) \le \mathbf{R}_{e}(s) \le B - \mathbf{R}_{e}(\alpha)$		(MODULATION)

№ Proof:

$$\begin{split} \mathbf{L} \big[e^{-\alpha x} \mathbf{g}(x) \big] &\triangleq \int_{x \in \mathbb{R}} e^{-\alpha x} \mathbf{g}(x) e^{-sx} \, \mathrm{d}x & \text{by definition of } \mathbf{L} & \text{(Definition L.1 page 223)} \\ &= \int_{x \in \mathbb{R}} \mathbf{g}(x) e^{-(s+\alpha)x} \, \mathrm{d}x & A \leq \mathbf{R}_{\mathrm{e}}(s+\alpha) \leq B & b^{x+y} = b^x b^y \\ &\triangleq \big[\mathbf{L} \mathbf{g}(x) \big](s+\alpha) & A - \mathbf{R}_{\mathrm{e}}(\alpha) \leq \mathbf{R}_{\mathrm{e}}(s) \leq B - \mathbf{R}_{\mathrm{e}}(\alpha) & \text{(Definition L.1 page 223)} \\ &\triangleq \mathbf{G}(s+\alpha) & A - \mathbf{R}_{\mathrm{e}}(\alpha) \leq \mathbf{R}_{\mathrm{e}}(s) \leq B - \mathbf{R}_{\mathrm{e}}(\alpha) & \text{by definition of } \mathbf{G}(s) \end{split}$$

Corollary L.3. ⁸ Let **L** be the Laplace Transform operator (Definition L.1 page 223). Let $G(s) \triangleq [Lg(x)]$ Let the Region of Convergence of G(s) be $A \leq R_e(s) \leq B$.

	Mapping		Region of Convergence	Domain
	$\mathbf{L}\left[\cos(\omega_o x)\mathbf{g}(x)\right] = \frac{1}{2}\mathbf{G}(s - i\omega_o) + \frac{1}{2}\mathbf{G}(s + i\omega_o)$	for	$\mathbf{R}_{e}(s) \in [A : B]$	$\forall x \omega_o \in \mathbb{C}$
CO	$\mathbf{L}\left[\sin(\omega_o x)\mathbf{g}(x)\right] = -\frac{i}{2}\mathbf{G}(s - i\omega_o) + \frac{i}{2}\mathbf{G}(s + i\omega_o)$	for	$\mathbf{R}_{e}(s) \in [A : B]$	$\forall x\omega_o{\in}\mathbb{C}$
R	$\mathbf{L}\left[\cosh\left(\omega_{o}x\right)\mathbf{g}(x)\right] = \frac{1}{2}\mathbf{G}(s-\omega_{o}) + \frac{1}{2}\mathbf{G}(s+\omega_{o})$	for	$\mathbf{R}_{e}(s) \in \left[A + \left \mathbf{R}_{e}(\omega_o) \right : B - \left \mathbf{R}_{e}(\omega_o) \right \right]$	$\forall x\omega_o{\in}\mathbb{C}$
	$\mathbf{L}\big[\sinh\big(\omega_o x\big)g(x)\big] = \frac{1}{2}G(s-\omega_o) - \frac{1}{2}G(s+\omega_o)$	for	$\mathbf{R}_{e}(s) \in \left[A + \left \mathbf{R}_{e}(\omega_o) \right : B - \left \mathbf{R}_{e}(\omega_o) \right \right]$	$\forall x\omega_o{\in}\mathbb{C}$

⁷ ■ Bracewell (1978) page 224 ⟨Table 11.1: "Modulation" entry⟩, ■ Levy (1958) page 19 ⟨Equation 1.2⟩

⁸ Bracewell (1978) page 224 (Table 11.1 Theorems for the Laplace Transform)



^ℚProof:

1. Mappings:

$$\begin{split} \mathbf{L} \big[\cosh \big(\omega_o x \big) \mathbf{g}(x) \big] &= \mathbf{L} \bigg[\bigg(\frac{e^{\omega_o x} + e^{-\omega_o x}}{2} \bigg) \mathbf{g}(x) \bigg] & \text{by definition of } \cosh(x) & \text{(Definition G.5 page 168)} \\ &= \frac{1}{2} \mathbf{L} \big[e^{\omega_o x} \mathbf{g}(x) \big](s) + \frac{1}{2} \mathbf{L} \big[e^{-\omega_o x} \mathbf{g}(x) \big](s) & \text{by } additive \text{ property} \\ &= \frac{1}{2} \mathbf{L} \big[\mathbf{g}(x) \big](s - \omega) + \frac{1}{2} \mathbf{L} \big[\mathbf{g}(x) \big](s + \omega) & \text{by } modulation \text{ prop.} \\ &= \frac{1}{2} \mathbf{G}(s - \omega_o) + \frac{1}{2} \mathbf{G}(s + \omega_o) & \text{by definition of } \mathbf{G}(s) \end{split}$$
 (Theorem L.4 page 226)

$$\begin{split} \mathbf{L} \big[\sinh \big(\omega_o x \big) \mathbf{g}(x) \big] &= \mathbf{L} \Bigg[\bigg(\frac{e^{\omega_o x} - e^{-\omega_o x}}{2} \bigg) \mathbf{g}(x) \bigg] & \text{by definition of sinh}(x) & \text{(Definition G.5 page 168)} \\ &= \frac{1}{2} \mathbf{L} \big[e^{\omega_o x} \mathbf{g}(x) \big](s) - \frac{1}{2} \mathbf{L} \big[e^{-\omega_o x} \mathbf{g}(x) \big](s) & \text{by $additive$ property} \\ &= \frac{1}{2} \mathbf{L} \big[\mathbf{g}(x) \big](s - \omega) - \frac{1}{2} \mathbf{L} \big[\mathbf{g}(x) \big](s + \omega) & \text{by $modulation$ prop.} \\ &= \frac{1}{2} \mathbf{G}(s - \omega_o) - \frac{1}{2} \mathbf{G}(s + \omega_o) & \text{by definition of $\mathbf{G}(s)} \end{split}$$
 (Theorem L.4 page 226)

$$\begin{split} \mathbf{L} \big[\cos \big(\omega_o x \big) \mathbf{g}(x) \big] &= \mathbf{L} \big[\cosh \big(i \omega_o x \big) \mathbf{g}(x) \big] & \text{by Theorem G.12 page 169} \\ &= \frac{1}{2} \mathbf{G}(s - i \omega_o) + \frac{1}{2} \mathbf{G}(s + i \omega_o) & \text{by } \mathbf{L} \big[\cos (\omega_o x) \mathbf{g}(x) \big] \text{ result} \end{split}$$

$$\begin{split} \mathbf{L} \big[\sin \big(\omega_o x \big) \mathbf{g}(x) \big] &= \mathbf{L} \big[-i^2 \sin \big(\omega_o x \big) \mathbf{g}(x) \big] \\ &= -i \mathbf{L} \big[i \sin \big(\omega_o x \big) \mathbf{g}(x) \big] & \text{by $homogeneous$ property} \quad \text{(Theorem L.3 page 225)} \\ &= -i \mathbf{L} \big[\sinh \big(i \omega_o x \big) \mathbf{g}(x) \big] & \text{by Theorem G.12 page 169} \\ &= -\frac{i}{2} \mathbf{G}(s - i \omega_o) + \frac{i}{2} \mathbf{G}(s + i \omega_o) & \text{by } \mathbf{L} \big[\sin (\omega_o x) \mathbf{g}(x) \big] \text{ result} \end{split}$$

2. Region of Convergence of $L[\cos(\omega_o x)g(x)]$ and $L[\sin(\omega_o x)g(x)]$:

$$\begin{aligned} & \mathbf{RocL} \Big[^{\cos /\!\!\!\!\!/}_{\sin} \left(\omega_o x \right) \mathbf{g}(x) \Big] \\ & = \mathbf{RocL} \Big[\left(\frac{e^{i \omega_o x} \pm e^{-i \omega_o x}}{2} \right) \mathbf{g}(x) \Big] & \text{by } Euler's \, Identity & \text{(Theorem G.5 page 160)} \\ & = \mathbf{Roc} \Big(\mathbf{L} \Big[\frac{e^{i \omega_o x}}{2} \mathbf{g}(x) \Big] \pm \mathbf{L} \Big[\frac{e^{-i \omega_o x}}{2} \mathbf{g}(x) \Big] \Big) & \text{by } additive \, \mathbf{property} & \text{(Theorem L.3 page 225)} \\ & = \mathbf{RocL} \Big[\left(\frac{e^{-i \omega_o x}}{2} \right) \mathbf{g}(x) \Big] \cap \mathbf{RocL} \Big[\left(\frac{e^{i \omega_o x}}{2} \right) \mathbf{g}(x) \Big] \\ & = \left[A - \mathbf{R}_{\mathrm{e}}(i \omega) : B - \mathbf{R}_{\mathrm{e}}(i \omega) \right] \cap \left[A - \mathbf{R}_{\mathrm{e}}(-i \omega) : B - \mathbf{R}_{\mathrm{e}}(-i \omega) \right] \\ & = \left[A - 0 : B - 0 \right] \cap \left[A - 0 : B - 0 \right] \\ & = \left[A : B \right] \end{aligned}$$

3. Region of Convergence of L[$\cosh(\omega_o x)g(x)$] and L[$\sinh(\omega_o x)g(x)$]:


$$\mathbf{RocL}\left[\frac{\cos h/\sin \left(\omega_{o} x\right) g(x)}{2}\right]$$

$$= \mathbf{RocL}\left[\left(\frac{e^{\omega_{o} x} \pm e^{-\omega_{o} x}}{2}\right) g(x)\right]$$
by def. $\cosh(x)$, $\sinh(x)$ (Definition G.5 page 168)

$$\begin{split} &= \mathbf{Roc} \bigg(\mathbf{L} \bigg[\frac{e^{\omega_o x}}{2} \mathbf{g}(x) \bigg] \pm \mathbf{L} \bigg[\frac{e^{-\omega_o x}}{2} \mathbf{g}(x) \bigg] \bigg) \qquad \text{by $additive$ property} \qquad \text{(Theorem L.3 page 225)} \\ &= \mathbf{Roc} \mathbf{L} \bigg[\bigg(\frac{e^{-\omega_o x}}{2} \bigg) \mathbf{g}(x) \bigg] \cap \mathbf{Roc} \mathbf{L} \bigg[\bigg(\frac{e^{\omega_o x}}{2} \bigg) \mathbf{g}(x) \bigg] \\ &= \big[A - \mathbf{R}_{\mathrm{e}}(\omega_o) : B - \mathbf{R}_{\mathrm{e}}(\omega_o) \big] \cap \big[A - \mathbf{R}_{\mathrm{e}}(-\omega_o) : B - \mathbf{R}_{\mathrm{e}}(-\omega_o) \big] \\ &= \begin{cases} \big[A + \mathbf{R}_{\mathrm{e}}(\omega_o) : B - \mathbf{R}_{\mathrm{e}}(\omega_o) \big] & \text{for $\omega \geq 0$} \\ \big[A - \mathbf{R}_{\mathrm{e}}(\omega_o) : B + \mathbf{R}_{\mathrm{e}}(\omega_o) \big] & \text{otherwise} \end{cases} \\ &= \big[A + \big| \mathbf{R}_{\mathrm{e}}(\omega_o) \big| : B - \big| \mathbf{R}_{\mathrm{e}}(\omega_o) \big| \big] \qquad \text{by definition of $|x|$} \end{split}$$

L.6 Causality properties

Definition L.3. 9 The Heaviside step function $\mu(x)$ or unit step function is defined as



Theorem L.5. 10 Let L be the LAPLACE TRANSFORM operator (Definition L.1 page 223) and $\mu(x)$ the UNIT STEP function (Definition L.3 page 228).

		Марр	ing		Reg	ion of Convergence	Domain
H	(1).	$\mathbf{L}[\mu(x)]$	=	$\frac{1}{s}$	for	$\mathbf{R}_{e}(s) > 0$	$\forall x \in \mathbb{R}$
M	(2).	$\mathbf{L}[\mu(-x)]$	=	$-\frac{1}{s}$	for	$\mathbf{R}_{e}(s) < 0$	$\forall x \in \mathbb{R}$

[♠]Proof:

$$\begin{aligned} \mathbf{L}[\mu(x)] &\triangleq \int_{\mathbb{R}} \mu(x) e^{-sx} \, \mathrm{d}x & \text{by definition of } \mathbf{L} & \text{(Definition L.1 page 223)} \\ &= \int_{0}^{\infty} e^{-sx} \, \mathrm{d}x & \text{by definition of } \mu(x) & \text{(Definition L.3 page 228)} \\ &= \frac{e^{-sx}}{-s} \Big|_{0}^{\infty} & \text{by } Fundamental Theorem of Calculus} \\ &= \lim_{x \to \infty} \left[\frac{e^{-sx}}{-s} \right] - \left(\frac{e^{0}}{-s} \right) \\ &= 0 + \frac{1}{s} & \forall \mathbf{R}_{\mathbf{e}}(s) > 0 \\ &= \frac{1}{s} & \forall \mathbf{R}_{\mathbf{e}}(s) > 0 \end{aligned}$$

$$\mathbf{L}[\mu(-x)] = \mathbf{L}[\mu(x)](-s) & \mathbf{R}_{\mathbf{e}}(s) < 0 & \text{by } reversal \text{ property} \\ &= \frac{-1}{s} & \mathbf{R}_{\mathbf{e}}(s) < 0 & \text{by } (1) \end{aligned}$$

by (1)

¹⁰ Bracewell (1978) page 227 (Table 11.2 Some Laplace transforms)



—>

⁹ Betten (2008a) page 285

Corollary L.4. ¹¹ Let L be the LAPLACE TRANSFORM operator (Definition L.1 page 223) and $\mu(x)$ the UNIT STEP function.

№PROOF:

$$\begin{split} \mathbf{L} \big[e^{-\alpha x} \mu(x) \big](s) &= \mathbf{L} [\mu(x)](s+\alpha) & \text{by modulation} \\ &= \frac{1}{s+\alpha} & \forall \mathbf{R}_{\mathrm{e}}(s) \in \left(0 - \mathbf{R}_{\mathrm{e}}(\alpha) : \infty - \mathbf{R}_{\mathrm{e}}(\alpha) \right) \text{ by Theorem L.5 page 228} \\ &= \frac{1}{s+\alpha} & \forall \mathbf{R}_{\mathrm{e}}(s) > -\mathbf{R}_{\mathrm{e}}(\alpha) \end{split}$$

$$\begin{split} \mathbf{L} \big[e^{-\alpha x} \mu(-x) \big](s) &= \mathbf{L} [\mu(-x)](s+\alpha) & \text{by modulation} & \text{(Theorem L.4 page 226)} \\ &= \frac{-1}{s+\alpha} & \forall \mathbf{R}_{\mathrm{e}}(s) \in \left(-\infty - \mathbf{R}_{\mathrm{e}}(\alpha) : 0 - (-\mathbf{R}_{\mathrm{e}}(\alpha))\right) \text{ by Theorem L.5 page 228} \\ &= \frac{-1}{s+\alpha} & \forall \mathbf{R}_{\mathrm{e}}(s) < \mathbf{R}_{\mathrm{e}}(\alpha) & \text{by anti-causality} & \text{(Theorem L.5 page 228)} \end{split}$$

Corollary L.5. ¹² Let L be the Laplace Transform operator (Definition L.1 page 223) and $\mu(x)$ the Unit STEP function.

	J	Mapping			Reg	ion of Convergence	Domain
	(1).	$\mathbf{L}\big[\cos(\omega_o x)\mu(x)\big]$	=	$\frac{s}{s^2 + \omega_o^2}$		$\mathbf{R}_{e}(s) > 0$	$x,\omega_o\in\mathbb{R}$
CO	(2).	$\mathbf{L}\big[\sin(\omega_o x)\mu(x)\big]$	=	$\frac{\omega_o}{s^2 + \omega_o^2}$	for	$\mathbf{R}_{e}(s) > 0$	$x,\omega_o\in\mathbb{R}$
R	(3).	$\mathbf{L}\big[\cos(\omega_o x)\mu(-x)\big]$	=	$\frac{-s}{s^2 + \omega_o^2}$		$\mathbf{R}_{e}(s) < 0$	$x,\omega_o\in\mathbb{R}$
	(4).	$\mathbf{L}\big[\sin(\omega_o x)\mu(-x)\big]$	=	$\frac{-\omega_o}{s^2 + \omega_o^2}$	for	$\mathbf{R}_{e}(s) < 0$	$x,\omega_o\in\mathbb{R}$

NPROOF:

$$\begin{split} \mathbf{L} \big[\cos(\omega_o x) \mu(x) \big](s) &= \frac{1}{2} \mathbf{L} [\mu(x)] (s - i\omega_o) + \frac{1}{2} \mathbf{L} [\mu(x)] (s + i\omega_o) \\ &= \frac{1}{2} \left[\frac{1}{s - i\omega_o} \right] + \frac{1}{2} \left[\frac{1}{s + i\omega_o} \right] \\ &= \frac{1}{2} \left[\frac{1}{s - i\omega_o} \right] \left[\frac{s + i\omega_o}{s + i\omega_o} \right] + \frac{1}{2} \left[\frac{1}{s + i\omega_o} \right] \left[\frac{s - i\omega_o}{s - i\omega_o} \right] \\ &= \frac{1}{2} \left[\frac{(s + i\omega_o) + (s - i\omega_o)}{s^2 + \omega_o^2} \right] \\ &= \frac{s}{s^2 + \omega_o^2} \end{split} \qquad \qquad \mathbf{R}_{\mathrm{e}}(s) > 0 \end{split}$$

¹² ■ Bracewell (1978) page 227 〈Table 11.2 Some Laplace transforms〉, ■ Shafii-Mousavi (2015) page 3 〈Table 1, using One-Sided Laplace Transform〉





¹¹ and and Bremmer (1959) page 22 ⟨Introduction⟩, shafii-Mousavi (2015) page 3 ⟨Table 1, using One-Sided Laplace Transform⟩, van der Pol and Bremmer (1959) page 26 ⟨(8) seems to have an error: $\frac{s}{s+\alpha}$ ⟩

$$\begin{split} \mathbf{L} \big[\sin(\omega_o x) \mu(x) \big](s) &= -\frac{i}{2} \mathbf{L} [\mu(x)] (s - i \omega_o) + \frac{i}{2} \mathbf{L} [\mu(x)] (s + i \omega_o) & \text{by } modulation \quad \text{(Corollary L.3 page 226)} \\ &= -\frac{i}{2} \left[\frac{1}{s - i \omega_o} \right] + \frac{i}{2} \left[\frac{1}{s + i \omega_o} \right] & \mathbf{R}_{\mathrm{e}}(s) > 0 \quad \text{by } causal \text{ prop.} \quad \text{(Theorem L.5 page 228)} \\ &= -\frac{i}{2} \left[\frac{1}{s - i \omega_o} \right] \left[\frac{s + i \omega_o}{s + i \omega_o} \right] + \frac{i}{2} \left[\frac{1}{s + i \omega_o} \right] \left[\frac{s - i \omega_o}{s - i \omega_o} \right] & \text{(Rationalizing the Denominator)} \\ &= \frac{i}{2} \left[\frac{-(s + i \omega_o) + (s - i \omega_o)}{s^2 + \omega_o^2} \right] & \mathbf{R}_{\mathrm{e}}(s) > 0 \\ &= \frac{\omega_o}{s^2 + \omega_o^2} & \mathbf{R}_{\mathrm{e}}(s) > 0 \end{split}$$

$$\begin{split} \mathbf{L} \big[\mu(-x) \mathrm{cos}(\omega_o x) \big](s) &= \mathbf{L} \big[\mu(-x) \mathrm{cos}(\omega_o(-x)) \big](s) \\ &= \mathbf{L} \big[\mu(x) \mathrm{cos}(\omega_o x) \big](-s) \\ &= \frac{(-s)}{(-s)^2 + \omega_o^2} \\ &= \frac{-s}{s^2 + \omega_o^2} \\ \mathbf{R}_{\mathbf{e}}(s) < 0 \\ \mathbf{E} \big[\mathrm{sin}(\omega_o x) \mu(-x) \big](s) \\ &= -\mathbf{L} \big[\mathrm{sin}(\omega_o(-x)) \mu(-x) \big](s) \\ &= -\mathbf{L} \big[\mathrm{sin}(\omega_o(-x)) \mu(-x) \big](s) \\ &= -\mathbf{L} \big[\mathrm{sin}(\omega_o x) \mu(x) \big](-s) \\ &= -\mathbf{L} \big[\mathrm{$$

Corollary L.6. ¹³ *Let* L *be the* LAPLACE TRANSFORM *operator* (Definition L.1 page 223) and $\mu(x)$ the UNIT STEP function.

	Ма	Mapping						
	(1). $\mathbf{L}\left[\cosh(\omega_o x)\mu(x)\right]$	(x) =	$\frac{s}{s^2 - \omega_o^2}$	for	$\mathbf{R}_{e}(s) > \left \omega_o \right $	$x,\omega_o \in \mathbb{R}$		
CO	(2). $\mathbf{L}\left[\sinh(\omega_o x)\mu(x)\right]$	<u>(</u>)] =	$\frac{\omega_o}{s^2 - \omega_o^2}$	for	$\mathbf{R}_{e}(s) > \left \omega_o \right $	$x,\omega_o\in\mathbb{R}$		
R	(3). $\mathbf{L}\left[\sinh(\omega_o x)\mu(-\omega_o x)\right]$		$\frac{-s}{s^2 - \omega_o^2}$	for	$\mathbf{R}_{e}(s) < \left \omega_o \right $	$x,\omega_o\in\mathbb{R}$		
	(4). $\mathbf{L}[\sinh(\omega_o x)\mu(-$	-x)] =	$\frac{-\omega_o}{s^2 - \omega_o^2}$	for	$\mathbf{R}_{e}(s) < \left \omega_o \right $	$x,\omega_o\in\mathbb{R}$		

№PROOF:

1. Mappings for $\mathbf{L}[\cosh(\omega_o x)\mu(x)]$ and $\mathbf{L}[\sinh(\omega_o x)\mu(x)]$:

$$\mathbf{L}\left[\cosh(\omega_{o}x)\mu(x)\right](s) = \frac{1}{2}\mathbf{L}[\mu(x)](s - \omega_{o}) + \frac{1}{2}\mathbf{L}[\mu(x)](s + \omega_{o}) \quad \text{by } modulation \qquad \text{(Corollary L.3 page 226)}$$

$$= \frac{1}{2}\left[\frac{1}{s - \omega_{o}}\right] + \frac{1}{2}\left[\frac{1}{s + \omega_{o}}\right] \quad \text{by } causal \text{ property} \quad \text{(Theorem L.5 page 228)}$$

¹³ Shafii-Mousavi (2015) page 3 (Table 1, using One-Sided Laplace Transform)



$$\begin{split} &=\frac{1}{2}\left[\frac{1}{s-\omega_o}\right]\left[\frac{s+\omega_o}{s+\omega_o}\right] + \frac{1}{2}\left[\frac{1}{s+\omega_o}\right]\left[\frac{s-\omega_o}{s-\omega_o}\right] \\ &=\frac{1}{2}\left[\frac{(s+\omega_o)+(s-\omega_o)}{s^2-\omega_o^2}\right] \\ &=\frac{s}{s^2-\omega_o^2} \end{split}$$

$$\begin{split} \mathbf{L} \big[& \sinh(\omega_o x) \mu(x) \big](s) = \frac{1}{2} \mathbf{L} [\mu(x)] (s - \omega_o) - \frac{1}{2} \mathbf{L} [\mu(x)] (s + \omega_o) \quad \text{by } modulation \\ &= \frac{1}{2} \left[\frac{1}{s - \omega_o} \right] - \frac{1}{2} \left[\frac{1}{s + \omega_o} \right] \quad \text{by } causal \text{ property} \quad \text{(Theorem L.5 page 228)} \\ &= \frac{1}{2} \left[\frac{1}{s - \omega_o} \right] \left[\frac{s + \omega_o}{s + \omega_o} \right] - \frac{1}{2} \left[\frac{1}{s + \omega_o} \right] \left[\frac{s - \omega_o}{s - \omega_o} \right] \\ &= \frac{1}{2} \left[\frac{(s + \omega_o) - (s - \omega_o)}{s^2 - \omega_o^2} \right] \\ &= \frac{\omega_o}{s^2 - \omega_o^2} \end{split}$$

2. Region of Convergence of $L[\cosh(\omega_o x)\mu(x)]$ and $L[\sinh(\omega_o x)\mu(x)]$:

$$\begin{aligned} \mathbf{RocL} \big[\cosh(\omega_o x) \mu(x) \big] &= \big[A + \big| \mathbf{R}_{\mathsf{e}}(\omega_o) \big| \ : \ B - \big| \mathbf{R}_{\mathsf{e}}(\omega_o) \big| \big] & \text{by Corollary L.3 page 226} \\ &= \big(0 + \big| \mathbf{R}_{\mathsf{e}}(\omega_o) \big| \ : \ \infty - \big| \mathbf{R}_{\mathsf{e}}(\omega_o) \big| \big) & \text{by Theorem L.5 page 228} \\ &= \big(\big| \mathbf{R}_{\mathsf{e}}(\omega_o) \big| \ : \ \infty \big) \\ &\implies \mathbf{RocL} \big[\cosh(\omega_o x) \mu(x) \big] > \big| \mathbf{R}_{\mathsf{e}}(\omega_o) \big| \end{aligned}$$

3. Mappings for $\mathbf{L}[\cosh(\omega_o x)\mu(-x)]$ and $\mathbf{L}[\sinh(\omega_o x)\mu(-x)]$:

$$\begin{split} \mathbf{L} \big[\cosh(\omega_o x) \mu(-x) \big](s) &= \mathbf{L} \big[\cosh(\omega_o (-x)) \mu(-x) \big](s) \\ &= \mathbf{L} \big[\cosh(\omega_o x) \mu(x) \big](-s) \qquad \text{by } reversal \text{ property} \\ &= \frac{(-s)}{(-s)^2 - \omega_o^2} \qquad \text{by previous result} \\ &= \frac{-s}{s^2 - \omega_o^2} \end{split}$$

$$\begin{split} \mathbf{L} \big[\sinh(\omega_o x) \mu(-x) \big](s) &= \mathbf{L} \big[-\sinh(\omega_o (-x)) \mu(-x) \big](s) \\ &= -\mathbf{L} \big[\sinh(\omega_o (-x)) \mu(-x) \big](s) \quad \text{by } homogeneous \text{ property} \qquad \text{(Theorem L.3 page 225)} \\ &= -\mathbf{L} \big[\sinh(\omega_o x) \mu(x) \big](-s) \quad \text{by } reversal \text{ property} \qquad \text{(Corollary L.1 page 225)} \\ &= \frac{-\omega_o}{(-s)^2 - \omega_o^2} \qquad \text{by previous result} \\ &= \frac{-\omega_o}{s^2 - \omega_o^2} \end{split}$$

4. Region of Convergence of $\mathbf{L} [\cosh(\omega_o x)\mu(-x)]$ and $\mathbf{L} [\sinh(\omega_o x)\mu(-x)]$:

$$\begin{aligned} \mathbf{RocL} \big[\cosh(\omega_o x) \mu(-x) \big] &= \big[A + \big| \mathbf{R}_{\mathsf{e}}(\omega_o) \big| : B - \big| \mathbf{R}_{\mathsf{e}}(\omega_o) \big| \big] & \text{by Corollary L.3 page 226} \\ &= \big(-\infty + \big| \mathbf{R}_{\mathsf{e}}(\omega_o) \big| : 0 - \big| \mathbf{R}_{\mathsf{e}}(\omega_o) \big| \big) & \text{by Theorem L.5 page 228} \\ &= \big(-\infty : \big| \mathbf{R}_{\mathsf{e}}(\omega_o) \big| \big) \\ &\Longrightarrow \mathbf{RocL} \big[\cosh(\omega_o x) \mu(-x) \big] < \big| \mathbf{R}_{\mathsf{e}}(\omega_o) \big| \end{aligned}$$

□>

Corollary L.7. ¹⁴ *Let* **L** *be the* Laplace Transform *operator* (Definition L.1 page 223) and $\mu(x)$ the unit step function.

	Mapping		Region of Convergence	Domain
	(1). $\mathbf{L}\left[\cos(\omega_o x)e^{-\alpha x}\mu(x)\right]$	=	$\frac{s+\alpha}{(s+\alpha)^2 + \omega_o^2} for$	$\mathbf{R}_{e}(s) > -\mathbf{R}_{e}(\alpha) x, \omega_o \in \mathbb{R}; \alpha \in \mathbb{C}$
CO	(2). $\mathbf{L}\left[\sin(\omega_o x)e^{-\alpha x}\mu(x)\right]$	=	$\frac{\omega_o}{(s+\alpha)^2 + \omega_o^2} for$	$\mathbf{R}_{e}(s) > -\mathbf{R}_{e}(\alpha) x, \omega_o \in \mathbb{R}; \alpha \in \mathbb{C}$
R	(3). $\mathbf{L}\left[\cos(\omega_o x)e^{\alpha x}\mu(-x)\right]$	=	$(s-\alpha)^2 + \omega_o^2$	$\mathbf{R}_{e}(s) > -\mathbf{R}_{e}(\alpha) x, \omega_o \in \mathbb{R}; \alpha \in \mathbb{C}$
	(4). $\mathbf{L}\left[\sin(\omega_o x)e^{\alpha x}\mu(-x)\right]$	=	$\frac{-\omega_o}{(s-\alpha)^2 + \omega_o^2} for$	$\mathbf{R}_{e}(s) > -\mathbf{R}_{e}(\alpha) x, \omega_o \in \mathbb{R}; \alpha \in \mathbb{C}$

№ Proof:

$$\begin{split} \mathbf{L} \big[\cos(\omega_o x) e^{-ax} \mu(x) \big] (s) &= \mathbf{L} \big[\mu(x) \cos(\omega_o x) \big] (s+\alpha) & \text{by } modulation \text{ property} \\ &= \frac{s+\alpha}{(s+\alpha)^2 + \omega_o^2} & \forall \mathbf{R}_\mathbf{e}(s) \in \left(0 - \mathbf{R}_\mathbf{e}(\alpha) : \infty - \mathbf{R}_\mathbf{e}(\alpha) \right) & \text{by Corollary L.5} \\ &= \frac{s+\alpha}{(s+\alpha)^2 + \omega_o^2} & \forall \mathbf{R}_\mathbf{e}(s) > -\mathbf{R}_\mathbf{e}(\alpha) \\ &= \frac{s+\alpha}{(s+\alpha)^2 + \omega_o^2} & \forall \mathbf{R}_\mathbf{e}(s) > -\mathbf{R}_\mathbf{e}(\alpha) \\ &= \frac{\omega_o}{(s+\alpha)^2 + \omega_o^2} & \forall \mathbf{R}_\mathbf{e}(s) \in \left(0 - \mathbf{R}_\mathbf{e}(\alpha) : \infty - \mathbf{R}_\mathbf{e}(\alpha) \right) & \text{by Corollary L.5} \\ &= \frac{\omega_o}{(s+\alpha)^2 + \omega_o^2} & \forall \mathbf{R}_\mathbf{e}(s) \in \left(0 - \mathbf{R}_\mathbf{e}(\alpha) : \infty - \mathbf{R}_\mathbf{e}(\alpha) \right) & \text{by Corollary L.5} \\ &= \frac{\omega_o}{(s+\alpha)^2 + \omega_o^2} & \forall \mathbf{R}_\mathbf{e}(s) > -\mathbf{R}_\mathbf{e}(\alpha) \\ &= \frac{-(s-\alpha)}{(s-\alpha)^2 + \omega_o^2} & \forall \mathbf{R}_\mathbf{e}(s) \in \left(0 - \mathbf{R}_\mathbf{e}(\alpha) : \infty - \mathbf{R}_\mathbf{e}(\alpha) \right) & \text{by Corollary L.5} \\ &= \frac{-(s-\alpha)}{(s-\alpha)^2 + \omega_o^2} & \forall \mathbf{R}_\mathbf{e}(s) \in \left(0 - \mathbf{R}_\mathbf{e}(\alpha) : \infty - \mathbf{R}_\mathbf{e}(\alpha) \right) & \text{by Corollary L.5} \\ &= \frac{-s+\alpha}{(s-\alpha)^2 + \omega_o^2} & \forall \mathbf{R}_\mathbf{e}(s) > -\mathbf{R}_\mathbf{e}(\alpha) \\ &= \frac{-s+\alpha}{(s-\alpha)^2 + \omega_o^2} & \forall \mathbf{R}_\mathbf{e}(s) > -\mathbf{R}_\mathbf{e}(\alpha) \\ &= \frac{-\omega_o}{(s-\alpha)^2 + \omega_o^2} & \forall \mathbf{R}_\mathbf{e}(s) \in \left(0 - \mathbf{R}_\mathbf{e}(\alpha) : \infty - \mathbf{R}_\mathbf{e}(\alpha) \right) & \text{by Corollary L.5} \\ &= \frac{-\omega_o}{(s-\alpha)^2 + \omega_o^2} & \forall \mathbf{R}_\mathbf{e}(s) \in \left(0 - \mathbf{R}_\mathbf{e}(\alpha) : \infty - \mathbf{R}_\mathbf{e}(\alpha) \right) & \text{by Corollary L.5} \\ &= \frac{-\omega_o}{(s-\alpha)^2 + \omega_o^2} & \forall \mathbf{R}_\mathbf{e}(s) \in \left(0 - \mathbf{R}_\mathbf{e}(\alpha) : \infty - \mathbf{R}_\mathbf{e}(\alpha) \right) & \text{by Corollary L.5} \\ &= \frac{-\omega_o}{(s-\alpha)^2 + \omega_o^2} & \forall \mathbf{R}_\mathbf{e}(s) \in \left(0 - \mathbf{R}_\mathbf{e}(\alpha) : \infty - \mathbf{R}_\mathbf{e}(\alpha) \right) & \text{by Corollary L.5} \\ &= \frac{-\omega_o}{(s-\alpha)^2 + \omega_o^2} & \forall \mathbf{R}_\mathbf{e}(s) \in \left(0 - \mathbf{R}_\mathbf{e}(\alpha) : \infty - \mathbf{R}_\mathbf{e}(\alpha) \right) & \text{by Corollary L.5} \\ &= \frac{-\omega_o}{(s-\alpha)^2 + \omega_o^2} & \forall \mathbf{R}_\mathbf{e}(s) \in \left(0 - \mathbf{R}_\mathbf{e}(\alpha) : \infty - \mathbf{R}_\mathbf{e}(\alpha) \right) & \text{by Corollary L.5} \\ &= \frac{-\omega_o}{(s-\alpha)^2 + \omega_o^2} & \forall \mathbf{R}_\mathbf{e}(s) \in \left(0 - \mathbf{R}_\mathbf{e}(\alpha) : \infty - \mathbf{R}_\mathbf{e}(\alpha) \right) & \text{by Corollary L.5} \\ &= \frac{-\omega_o}{(s-\alpha)^2 + \omega_o^2} & \forall \mathbf{R}_\mathbf{e}(s) \in \left(0 - \mathbf{R}_\mathbf{e}(\alpha) : \infty - \mathbf{R}_\mathbf{e}(\alpha) \right) & \text{by Corollary L.5} \\ &= \frac{-\omega_o}{(s-\alpha)^2 + \omega_o^2} & \forall \mathbf{R}_\mathbf{e}(s) \in \left(0 - \mathbf{R}_\mathbf{e}(\alpha) : \infty - \mathbf{R}_\mathbf{e}(\alpha) \right) & \text{by Corollary L.5} \\ &= \frac{-\omega_o}{(s-\alpha)^2 + \omega_o^2} & \forall \mathbf{R}_\mathbf{e}$$

Corollary L.8. Let L be the LAPLACE TRANSFORM operator (Definition L.1 page 223).

C	$\mathbf{L}\left[\cos(\omega_o x)\right]$	is divergent is divergent	$\forall s \in \mathbb{C}$	$\forall x, \omega_o \in \mathbb{R}$
R	$\mathbf{L}\left[\sin(\omega_o x)\right]$	is divergent	$\forall s \in \mathbb{C}$	$\forall x, \omega_o \in \mathbb{R} \setminus \{0\}$

№ Proof:

$$\mathbf{L}\left[\cos(\omega_{o}x)\right] = \underbrace{\mathbf{L}\left[\mu(x)\cos(\omega_{o}x)\right]}_{\forall \mathbf{R}_{e}(s) > 0} + \underbrace{\mathbf{L}\left[\mu(-x)\cos(\omega_{o}x)\right]}_{\forall \mathbf{R}_{e}(s) < 0}$$
by Corollary L.5 page 229
$$= \underbrace{\frac{s}{s^{2} + \omega_{o}^{2}}}_{\forall \mathbf{R}_{e}(s) > 0} + \underbrace{\frac{-s}{s^{2} + \omega_{o}^{2}}}_{\forall \mathbf{R}_{e}(s) < 0}$$
by Corollary L.5 page 229

¹⁴ Bracewell (1978) page 227 (Table 11.2 Some Laplace transforms)



$$= \begin{cases} 0 & \forall \mathbf{R}_{e}(s) \in (-\infty:0) \cap (0:\infty) = \emptyset \\ \infty & \forall s \in \mathbb{C} \end{cases}$$

$$\Rightarrow \mathbf{L}[\cos(\omega_{o}x)] \text{ is } \mathbf{divergent} \ \forall s \in \mathbb{C}$$

$$\mathbf{L}[\sin(\omega_{o}x)] = \mathbf{L}[\mu(x)\sin(\omega_{o}x)] + \mathbf{L}[\mu(-x)\sin(\omega_{o}x)] \qquad \text{by Corollary L.5 page 229}$$

$$= \frac{\omega_{o}}{s^{2} + \omega_{o}^{2}} + \frac{-\omega_{o}}{s^{2} + \omega_{o}^{2}} \qquad \text{by Corollary L.5 page 229}$$

$$= \begin{cases} 0 & \forall \mathbf{R}_{e}(s) \in (-\infty:0) \cap (0:\infty) = \emptyset \\ \infty & \forall s \in \mathbb{C} \end{cases}$$

$$\Rightarrow \mathbf{L}[\sin(\omega_{o}x)] \text{ is } \mathbf{divergent} \ \forall s \in \mathbb{C}$$

L.7 Exponential decay properties

Corollary L.9. ¹⁵ Let L be the Laplace Transform operator (Definition L.1 page 223) and $\mu(x)$ the unit step function. Let $A \triangleq \mathbf{R}_{\mathbf{e}}(\alpha)$.

	J	C \	<i>'</i>	
С	Mapping		Region of Convergence	Domain
O R	$\mathbf{L}\big[e^{-\alpha x }\big] =$	$=\frac{2\alpha}{\alpha^2-s^2}$	for $\mathbf{R}_{e}(s) \in (-A : A)$	$x,\alpha\in\mathbb{C}$

№ Proof:

$$\begin{split} \mathbf{L} \big[e^{-\alpha|x|} \big] &= \mathbf{L} \big[e^{-\alpha|x|} \mu(x) + e^{-\alpha|x|} \mu(-x) \big] & \text{by definition of } \mu(x) & \text{(Definition L.3 page 228)} \\ &= \mathbf{L} \big[e^{-\alpha|x|} \mu(x) \big] + \mathbf{L} \big[e^{-\alpha|x|} \mu(-x) \big] & \text{by homogeneous property} & \text{(Theorem L.3 page 225)} \\ &= \mathbf{L} \big[e^{-\alpha x} \mu(x) \big] + \mathbf{L} \big[e^{\alpha x} \mu(-x) \big] & \text{by Definition L.3 page 228} & \text{and Corollary L.4 page 229} \\ &= \left[\frac{1}{s+\alpha} \right] + \left[\frac{-1}{s-\alpha} \right] & \forall \mathbf{R_e}(s) \in \left(-\mathbf{R_e}(\alpha) : \mathbf{R_e}(\alpha) \right) & \text{by Corollary L.4 page 229} \\ &= \frac{(s-\alpha)-(s+\alpha)}{(s+\alpha)(s-\alpha)} & \forall \mathbf{R_e}(s) \in \left(-\mathbf{R_e}(\alpha) : \mathbf{R_e}(\alpha) \right) \\ &= \frac{2\alpha}{\alpha^2-s^2} & \forall \mathbf{R_e}(s) \in \left(-\mathbf{R_e}(\alpha) : \mathbf{R_e}(\alpha) \right) \end{split}$$

Corollary L.10. ¹⁶ *Let* **L** *be the* Laplace Transform *operator* (Definition L.1 page 223) and $\mu(x)$ the Unit Step function. Let $A \triangleq \mathbf{R}_{\mathbf{e}}(\alpha)$.

	oral juntotto, to the large with					
		Mapping		Region of Convergence	Domain	
С	c	(1). $\mathbf{L}\left[\cos(\omega_o x)e^{-\alpha x }\mu(x)\right]$	$= \frac{s+\alpha}{(s+\alpha)^2 + \omega_o^2}$	for $\mathbf{R}_{e}(s) \in (-A:A)$	$x,\alpha\in\mathbb{R}$	
	O R	(2). $\mathbf{L}\left[\cos(\omega_o x)e^{-\alpha x }\mu(-x)\right]$	$=\frac{-s+\alpha}{s}$	for $\mathbf{R}_{e}(s) \in (-A : A)$	$x,\alpha\in\mathbb{R}$	
		(3). $\mathbf{L}\left[\cos(\omega_o x)e^{-\alpha x }\right]$	$= \frac{(s-\alpha)^2 + \omega_o^2}{s+\alpha} + \frac{-s+\alpha}{(s-\alpha)^2 + \omega_o^2}$	for $\mathbf{R}_{e}(s) \in (-A : A)$	$x,\alpha\in\mathbb{R}$	

¹⁵ Bracewell (1978) page 227 (Table 11.2 Some Laplace transforms),

🤲 Frames and Bases Structure and Design [VERSIDN 021] 🤲



Bracewell (1978) page 227 (Table 11.2 Some Laplace transforms), \triangle Levy (1958) page 19 (with $\psi = 0$, $\alpha_0 = \alpha$, and $\alpha_1 = 1$), http://ece-research.unm.edu/bsanthan/ece541/table ME.pdf

^ℚProof:

1. Proof for (1):

$$\begin{split} &\mathbf{L}\big[\cos(\omega_o x)e^{-\alpha|x|}\mu(x)\big](s) \\ &= \mathbf{L}\big[\cos(\omega_o x)e^{-\alpha x}\mu(x)\big](s) \\ &= \mathbf{L}\big[\cos(\omega_o x)\mu(x)\big](s+\alpha) \quad \forall \mathbf{R}_{\mathrm{e}}(s) \in \left(0-\mathbf{R}_{\mathrm{e}}(\alpha): \infty-\mathbf{R}_{\mathrm{e}}(\alpha)\right) \quad \text{by } \textit{modulation } \textit{prop.} \quad \textit{(Theorem L.4 page 226)} \\ &= \frac{s+\alpha}{(s+\alpha)^2+\omega_o^2} \quad \forall \mathbf{R}_{\mathrm{e}}(s) \in \left(-\mathbf{R}_{\mathrm{e}}(\alpha): \mathbf{R}_{\mathrm{e}}(\alpha)\right) \quad \text{by } \textit{Corollary L.5 page 229} \end{split}$$

2. Proof for (2):

$$\begin{split} &\mathbf{L} \big[\cos(\omega_o x) e^{-\alpha|x|} \, \mu(-x) \big] \\ &= \mathbf{L} \big[\cos(\omega_o x) e^{\alpha x} \, \mu(-x) \big] \qquad \text{by definition of } \mu(x) \qquad \text{(Definition L.3 page 228)} \\ &= \mathbf{L} \big[\cos(-\omega_o x) e^{\alpha x} \, \mu(-x) \big] \qquad \text{by } even \, \text{property of } \cos(x) \qquad \text{(Theorem G.2 page 157)} \\ &= \mathbf{L} \big[e^{\alpha x} \cos(\omega_o(-x)) \mu(-x) \big] \\ &= \mathbf{L} \cos(\omega_o(-x)) \mu(-x) (s-\alpha) \qquad \text{by } modulation \, \text{property} \qquad \text{(Theorem L.4 page 226)} \\ &= \mathbf{L} \cos(\omega_o(-x)) \mu(-x) (s-\alpha) \qquad \text{by } modulation \, \text{property} \qquad \text{(Theorem L.4 page 226)} \\ &= \frac{-s+\alpha}{(s-\alpha)^2 + \omega_o^2} \quad \forall \mathbf{R_e}(s) \in \left(-\mathbf{R_e}(\alpha) : \mathbf{R_e}(\alpha)\right) \quad \text{by Corollary L.5 and Theorem L.4 page 226} \end{split}$$

3. Proof for (3):

$$\begin{split} \mathbf{L} \big[\cos(\omega_o x) e^{-\alpha|x|} \big] &= \mathbf{L} \big[\cos(\omega_o x) e^{-\alpha|x|} \mu(x) \big] + \mathbf{L} \big[\cos(\omega_o x) e^{-\alpha|x|} \mu(-x) \big] \\ &= \mathbf{L} \big[\cos(\omega_o x) e^{-\alpha x} \mu(x) \big] + \mathbf{L} \big[\cos(-\omega_o x) e^{\alpha x} \mu(-x) \big] \\ &= \frac{s + \alpha}{(s + \alpha)^2 + \omega_o^2} + \frac{-s + \alpha}{(s - \alpha)^2 + \omega_o^2} \qquad \qquad \forall \mathbf{R}_{\mathrm{e}}(s) \in \left(-\mathbf{R}_{\mathrm{e}}(\alpha) : \mathbf{R}_{\mathrm{e}}(\alpha) \right) \end{split}$$

Corollary L.11. 17 Let L be the LAPLACE TRANSFORM operator (Definition L.1 page 223) and $\mu(x)$ the UNIT STEP function. Let $A \triangleq \mathbf{R}_{\bullet}(\alpha)$.

	J					
	Mapping		Region of Convergence	Domain		
С	(1). $\mathbf{L}\left[\sin(\omega_o x)e^{-\alpha x }\mu(x)\right]$	$= \frac{\omega_o}{(s+\alpha)^2 + \omega_o^2}$	for $\mathbf{R}_{e}(s) \in (-A : A)$	$x,\alpha\in\mathbb{R}$		
O R	(2). $\mathbf{L}\left[\sin(\omega_o x)e^{-\alpha x }\mu(-x)\right]$	$ = \frac{-\omega_o}{(s-\alpha)^2 + \omega_o^2} $	for $\mathbf{R}_{\mathbf{e}}(s) \in (-A : A)$	$x,\alpha\in\mathbb{R}$		
	(3). $\mathbf{L}\left[\sin(\omega_o x)e^{-\alpha x }\right]$	$= \frac{\omega_o + \omega_o}{(s+\alpha)^2 + \omega_o^2} + \frac{-\omega_o}{(s-\alpha)^2 + \omega_o^2}$	for $\mathbf{R}_{\mathbf{e}}(s) \in (-A : A)$	$x,\alpha\in\mathbb{R}$		

[♠]Proof:

1. Proof for (1):

$$\begin{split} &\mathbf{L} \big[\sin(\omega_o x) e^{-\alpha |x|} \mu(x) \big] \\ &= \mathbf{L} \big[\sin(\omega_o x) e^{-\alpha x} \mu(x) \big] \quad \text{by definition of } \mu(x) \qquad \text{(Definition L.3 page 228)} \\ &= \frac{s + \alpha}{(\omega_o)^2 + \omega_o^2} \qquad \forall \mathbf{R}_{\mathrm{e}}(s) \in \left(-\mathbf{R}_{\mathrm{e}}(\alpha) : \mathbf{R}_{\mathrm{e}}(\alpha) \right) \quad \text{by Corollary L.5 page 229 and Theorem L.4 page 226} \end{split}$$

Bracewell (1978) page 227 (Table 11.2 Some Laplace transforms), \triangle Levy (1958) page 19 (with $\psi = 0$, $\alpha_0 = \alpha$, and $\alpha_1 = 1$, http://ece-research.unm.edu/bsanthan/ece541/table ME.pdf



2. Proof for (2):

$$\begin{split} &\mathbf{L}\big[\sin(\omega_o x)e^{-\alpha|x|}\mu(-x)\big]\\ &=\mathbf{L}\big[\sin(-\omega_o x)e^{\alpha x}\mu(-x)\big] \quad \text{by definition of }\mu(x) \qquad \text{(Definition L.3 page 228)}\\ &=\mathbf{L}\big[-\sin(\omega_o x)e^{\alpha x}\mu(-x)\big] \quad \text{by } odd \text{ property of }\sin(x) \qquad \text{(Theorem G.2 page 157)}\\ &=-\mathbf{L}\big[\sin(\omega_o x)e^{\alpha x}\mu(-x)\big] \quad \text{by } homogeneous \text{ property} \qquad \text{(Theorem L.3 page 225)}\\ &=\frac{-\omega_o}{(s-\alpha)^2+\omega_o^2} \qquad \forall \mathbf{R_e}(s) \in \left(-\mathbf{R_e}(\alpha):\mathbf{R_e}(\alpha)\right) \quad \text{by Theorem L.4 page 226 and Corollary L.5} \end{split}$$

3. Proof for (3):

$$\begin{split} \mathbf{L} \big[\sin(\omega_o x) e^{-\alpha |x|} \big] &= \mathbf{L} \big[\sin(\omega_o x) e^{-\alpha |x|} \mu(x) \big] + \mathbf{L} \big[\sin(\omega_o x) e^{-\alpha |x|} \mu(-x) \big] \\ &= \mathbf{L} \big[\sin(\omega_o x) e^{-\alpha x} \mu(x) \big] + \mathbf{L} \big[\sin(-\omega_o x) e^{\alpha x} \mu(-x) \big] \\ &= \frac{\omega_o}{(s+\alpha)^2 + \omega_o^2} + \frac{-\omega_o}{(s-\alpha)^2 + \omega_o^2} \end{split} \qquad \forall \mathbf{R}_{\mathbf{e}}(s) \in \left(-\mathbf{R}_{\mathbf{e}}(\alpha) : \mathbf{R}_{\mathbf{e}}(\alpha) \right) \end{split}$$

Product properties L.8

Theorem L.6 (next) demonstrates that multiplication in the "time domain" is equivalent to convolution in the "s domain" and vice-versa.

Theorem L.6 (convolution theorem). 18 Let L be the LAPLACE TRANSFORM operator (Definition L.1 page 223) and \star the convolution operator (Definition E.1 page 145). Let A, B, C, and D be defined as in Corollary L.2 (page 225).

$$\begin{array}{lll} \mathbf{T} & \mathbf{H} \\ \mathbf{H} & \mathbf{L} \big[\mathbf{f}(x) \star \mathbf{g}(x) \big](s) &= & [\mathbf{L} \mathbf{f}](s) \big[\mathbf{L} \mathbf{g} \big](s) & \forall \mathbf{R}_{\mathbf{e}}(s) \in [A:B] \cap [C:D] & \forall \mathbf{f}, \mathbf{g} \in \mathcal{L}^2_{(\mathbb{R},\mathcal{B},\mu)} \\ & \mathbf{L} \big[\mathbf{f}(x) \mathbf{g}(x) \big](s) &= & [\mathbf{L} \mathbf{f}](s) \star \big[\mathbf{L} \mathbf{g} \big](s) & \forall \mathbf{R}_{\mathbf{e}}(s) \in [A+C:B+D], \ c \in (A:B) & \forall \mathbf{f}, \mathbf{g} \in \mathcal{L}^2_{(\mathbb{R},\mathcal{B},\mu)}. \end{array}$$

^ℚProof:

$$\mathbf{L}\big[\mathsf{f}(x)\star\mathsf{g}(x)\big](s) = \mathbf{L}\left[\int_{u\in\mathbb{R}}\mathsf{f}(u)\mathsf{g}(x-u)\,\mathsf{d}\mathsf{u}\right](s) \qquad \text{by definition of }\star \qquad \text{(Definition E.1 page 145)}$$

$$= \int_{u\in\mathbb{R}}\mathsf{f}(u)\big[\mathbf{L}\mathsf{g}(x-u)\big](s)\,\mathsf{d}\mathsf{u}$$

$$= \int_{u\in\mathbb{R}}\mathsf{f}(u)e^{-su}\,\big[\mathbf{L}\mathsf{g}(x)\big](s)\,\mathsf{d}\mathsf{u} \qquad \text{by } translation \text{ property} \qquad \text{(Theorem L.2 page 224)}$$

$$= \underbrace{\left(\int_{u\in\mathbb{R}}\mathsf{f}(u)e^{-su}\,\mathsf{d}\mathsf{u}\right)}_{[\mathbf{L}\mathsf{f}](s)} \big[\mathbf{L}\mathsf{g}\big](s)$$

$$= [\mathbf{L}\mathsf{f}](s)\,\big[\mathbf{L}\mathsf{g}\big](s) \qquad \qquad \mathbf{R}_{\mathsf{e}}(s) \in [A:B] \cap [C:D] \qquad \text{by definition of }\mathbf{L}$$

$$\mathbf{L}[\mathbf{f}(x)\mathbf{g}(x)](s) = \mathbf{L}\left[\left(\mathbf{L}^{-1}\mathbf{L}\mathbf{f}(x)\right)\mathbf{g}(x)\right](s) \qquad \text{by def. of operator inverse} \qquad \text{(Definition C.3 page 112)}$$

$$= \mathbf{L}\left[\left(\int_{v \in \mathbb{R}} \left[\mathbf{L}\mathbf{f}(x)\right](v)e^{sxv} \, dv\right)\mathbf{g}(x)\right](s) \qquad \text{by Theorem L.1 page 224}$$

¹⁸ 🗐 Bracewell (1978) page 224, 🥥 Bachman et al. (2002) pages 268–270, 🥥 Bachman (1964) page 8



$$= \int_{v \in \mathbb{R}} [\mathbf{Lf}(x)](v) [\mathbf{L}(e^{sxv} \mathbf{g}(x))](s,v) \, dv$$

$$= \int_{v \in \mathbb{R}} [\mathbf{Lf}(x)](v) [\mathbf{Lg}(x)](s-v) \, dv \qquad \text{by Theorem L.2 page 224}$$

$$= [\mathbf{Lf}](s) \star [\mathbf{Lg}](s) \qquad \text{by definition of } \star \qquad \text{(Definition E.1 page 145)}$$

Calculus properties **L.9**

Theorem L.7. 19 Let L be the LAPLACE TRANSFORM operator (Definition L.1 page 223).

$$\begin{cases}
\lim_{x \to -\infty} g(x) = 0
\end{cases} \implies \left\{ L \left[\frac{d}{dt} g(x) \right] \right\} = s \left[Lg \right](s)$$

$$L \int_{u=-\infty}^{u=x} g(u) du = \frac{1}{s} \left[Lg \right](s)$$

^ℚProof:

$$\mathbf{L}\left[\frac{\mathsf{d}}{\mathsf{d}\mathsf{x}}\mathsf{g}(x)\right] \triangleq \int_{x \in \mathbb{R}} \underbrace{\left[\frac{\mathsf{d}}{\mathsf{d}\mathsf{x}}\mathsf{g}(x)\right]}_{\mathsf{d}\mathsf{v}} \underbrace{e^{-sx}}_{\mathsf{u}} \, \mathsf{d}\mathsf{x} \qquad \text{by definition of } \mathbf{L}$$

$$= \underbrace{e^{-sx}}_{\mathsf{u}} \underbrace{\mathsf{g}(x)}_{\mathsf{v}} \Big|_{x=-\infty}^{x=+\infty} - \int_{x \in \mathbb{R}} \underbrace{\mathsf{g}(x)(-s)e^{-sx}}_{\mathsf{d}\mathsf{u}} \, \mathsf{d}\mathsf{x} \qquad \text{by Integration by Parts}$$

$$= e^{-s\infty} \underbrace{\mathsf{g}(\infty)}_{\mathsf{v}} - e^{s\infty} \underbrace{\mathsf{g}(-\infty)^{-0}}_{\mathsf{v}}(-s) \int_{x \in \mathbb{R}} \mathsf{g}(x)e^{-sx} \, \mathsf{d}\mathsf{x} \qquad \text{by left hypothesis}$$

$$\triangleq s \Big[\mathbf{L} \mathbf{g} \Big](s) \qquad \text{by definition of } \mathbf{L} \qquad \text{(Definition L.1 page 223)}$$

$$\mathbf{L} \int_{u=-\infty}^{u=x} \mathbf{g}(u) \, d\mathbf{u} \triangleq \int_{x=-\infty}^{x=+\infty} \left[\int_{u=-\infty}^{u=x} \mathbf{g}(u) \, d\mathbf{u} \right] e^{-sx} \, d\mathbf{x} \qquad \text{by definition of } \mathbf{L}$$

$$= \int_{x=-\infty}^{x=+\infty} \left[\int_{u=-\infty}^{u=+\infty} \mathbf{g}(u) \mu(x-u) \, d\mathbf{u} \right] e^{-sx} \, d\mathbf{x}$$

$$= \int_{v=-\infty}^{v=+\infty} \int_{u=-\infty}^{u=+\infty} \mathbf{g}(u) h(v) e^{-s(u+v)} \, d\mathbf{u} \, d\mathbf{v}$$

$$= \left[\int_{v=-\infty}^{v=+\infty} \mu(v) e^{-sv} \, d\mathbf{v} \right] \left[\int_{u=-\infty}^{u=+\infty} \mathbf{g}(u) e^{-su} \, d\mathbf{u} \right]$$

$$= \left[\int_{v=0}^{v=\infty} e^{-sv} \, d\mathbf{v} \right] \left[\mathbf{L}\mathbf{g} \right] (s)$$

$$= \frac{1}{-s} e^{-sv} \Big|_{v=0}^{v=\infty} \left[\mathbf{L}\mathbf{g} \right] (s) \qquad \text{by } Fundamental Theorem of Calculus}$$

$$= \frac{1}{s} \left[\mathbf{L}\mathbf{g} \right] (s) \qquad \text{by definition of } \mathbf{L} \qquad \text{(Definition } \mathbf{L}.1 \text{ page 223)}$$

¹⁹ ■ Betten (2008b) page 301 ((B.27)), ■ Levy (1958) page 15 (Equation 0.7)



DISCRETE TIME FOURIER TRANSFORM


M.1 Definition

Definition M.1.

D E F

The discrete-time Fourier transform
$$\check{\mathbf{F}}$$
 of $(x_n)_{n\in\mathbb{Z}}$ is defined as $[\check{\mathbf{F}}(x_n)](\omega) \triangleq \sum_{n\in\mathbb{Z}} x_n e^{-i\omega n} \quad \forall (x_n)_{n\in\mathbb{Z}} \in \mathscr{C}^2_{\mathbb{R}}$

If we compare the definition of the *Discrete Time Fourier Transform* (Definition M.1 page 237) to the definition of the Z-transform (Definition J.4 page 208), we see that the DTFT is just a special case of the more general Z-Transform, with $z=e^{i\omega}$. If we imagine $z\in\mathbb{C}$ as a complex plane, then $e^{i\omega}$ is a unit circle in this plane. The "frequency" ω in the DTFT is the unit circle in the much larger z-plane, as illustrated to the right.



M.2 Properties

Proposition M.1 (DTFT periodicity). Let $\check{\mathbf{x}}(\omega) \triangleq \check{\mathbf{F}}\big[(x_n)\big](\omega)$ be the discrete-time Fourier transform (Definition M.1 page 237) of a sequence $(x_n)_{n\in\mathbb{Z}}$ in $\boldsymbol{\ell}^2_{\mathbb{R}}$.

$$\stackrel{\mathsf{P}}{\underset{\mathsf{P}}{\mathsf{R}}} \underbrace{\check{\mathsf{X}}(\omega) = \check{\mathsf{X}}(\omega + 2\pi n)}_{\text{PERIODIC with period } 2\pi} \quad \forall n \in \mathbb{Z}$$

NPROOF:

$$\check{\mathbf{x}}(\omega + 2\pi n) = \sum_{m \in \mathbb{Z}} x_m e^{-i(\omega + 2\pi n)m} = \sum_{m \in \mathbb{Z}} x_m e^{-i\omega m} e^{-i2\pi nm}$$

$$= \sum_{m \in \mathbb{Z}} x_m e^{-i\omega m} = \check{\mathbf{x}}(\omega)$$

₽

Theorem M.1. Let $\check{\mathbf{x}}(\omega) \triangleq \check{\mathbf{F}}[(\mathbf{x}[n])](\omega)$ be the discrete-time Fourier transform (Definition M.1 page 237) of a sequence $(x_n)_{n\in\mathbb{Z}}$ in $\mathscr{C}^2_{\mathbb{R}}$.

$$\left\{\begin{array}{ccc} \tilde{\mathbf{x}}(\omega) & \triangleq & \check{\mathbf{F}}(\mathbf{x}[n]) \end{array}\right\} & \Longrightarrow & \left\{\begin{array}{ccc} (1). & \check{\mathbf{F}}(\mathbf{x}[-n]) & = & \tilde{\mathbf{x}}(-\omega) & and \\ (2). & \check{\mathbf{F}}(\mathbf{x}^*[n]) & = & \tilde{\mathbf{x}}^*(-\omega) & and \\ (3). & \check{\mathbf{F}}(\mathbf{x}^*[-n]) & = & \tilde{\mathbf{x}}^*(\omega) \end{array}\right\}$$

№PROOF:

$$\mathbf{\check{F}}(\mathbf{x}[-n]) \triangleq \sum_{n \in \mathbb{Z}} \mathbf{x}[-n]e^{-i\omega n} \qquad \text{by definition of } DTFT \qquad \text{(Definition M.1 page 237)}$$

$$= \sum_{m \in \mathbb{Z}} \mathbf{x}[m]e^{i\omega m} \qquad \text{where } m \triangleq -n \implies n = -m$$

$$= \sum_{m \in \mathbb{Z}} \mathbf{x}[m]e^{-i(-\omega)m}$$

$$\triangleq \tilde{\mathbf{x}}(-\omega) \qquad \text{by left hypothesis}$$

$$\mathbf{\check{F}}(\mathbf{x}^*[n]) \triangleq \sum_{n \in \mathbb{Z}} \mathbf{x}^*[n]e^{-i\omega n} \qquad \text{by definition of } DTFT \qquad \text{(Definition M.1 page 237)}$$

$$\mathbf{F}(\mathbf{x}^*[n]) \triangleq \sum_{n \in \mathbb{Z}} \mathbf{x}^*[n]e^{-i\omega n} \qquad \text{by definition of } DTFT \qquad \text{(Definition M.1 page 237)}$$

$$= \left(\sum_{n \in \mathbb{Z}} \mathbf{x}[n]e^{i\omega n}\right)^* \qquad \text{by } distributive \text{ property of } *-\mathbf{algebras} \qquad \text{(Definition F.3 page 148)}$$

$$= \left(\sum_{n \in \mathbb{Z}} \mathbf{x}[n]e^{-i(-\omega)n}\right)^*$$

$$\triangleq \tilde{\mathbf{x}}^*(-\omega) \qquad \text{by left hypothesis}$$

$$\check{\mathbf{F}} (\mathbf{x}^*[-n]) \triangleq \sum_{n \in \mathbb{Z}} \mathbf{x}^*[-n]e^{-i\omega n} \qquad \text{by definition of } DTFT \qquad \text{(Definition M.1 page 237)}$$

$$= \left(\sum_{n \in \mathbb{Z}} \mathbf{x}[-n]e^{i\omega n}\right)^* \qquad \text{by } distributive \text{ property of } *-\mathbf{algebras} \qquad \text{(Definition F.3 page 148)}$$

$$= \left(\sum_{m \in \mathbb{Z}} \mathbf{x}[m]e^{-i\omega m}\right)^* \qquad \text{where } m \triangleq -n \implies n = -m$$

$$\triangleq \tilde{\mathbf{x}}^*(\omega) \qquad \text{by left hypothesis}$$

Theorem M.2. Let $\check{\mathbf{x}}(\omega) \triangleq \check{\mathbf{F}}[(\mathbf{x}[n])](\omega)$ be the DISCRETE-TIME FOURIER TRANSFORM (Definition M.1 page 237) of a sequence $(\mathbf{x}[n])_{n\in\mathbb{Z}}$ in $\mathscr{C}^2_{\mathbb{R}}$.

$$\begin{bmatrix}
\mathsf{T} \\
\mathsf{H} \\
\mathsf{M}
\end{bmatrix}
\begin{cases}
(1). \quad \tilde{\mathsf{X}}(\omega) \triangleq \tilde{\mathsf{F}}(\mathsf{X}[n]) & \text{and} \\
(2). \quad (\mathsf{X}[n]) \text{ is REAL-VALUED}
\end{cases}
\Rightarrow
\begin{bmatrix}
(1). \quad \tilde{\mathsf{F}}(\mathsf{X}[-n]) = \tilde{\mathsf{X}}(-\omega) & \text{and} \\
(2). \quad \tilde{\mathsf{F}}(\mathsf{X}^*[n]) = \tilde{\mathsf{X}}^*(-\omega) = \tilde{\mathsf{X}}(\omega) & \text{and} \\
(3). \quad \tilde{\mathsf{F}}(\mathsf{X}^*[-n]) = \tilde{\mathsf{X}}^*(\omega) = \tilde{\mathsf{X}}(-\omega)
\end{bmatrix}$$

№ Proof:

$$\mathbf{\breve{F}}\left(\mathbf{x}[-n]\right) \triangleq \sum_{n \in \mathbb{Z}} \mathbf{x}[-n]e^{-i\omega n} \qquad \text{by definition of } DTFT \qquad \text{(Definition M.1 page 237)}$$

$$= \sum_{m \in \mathbb{Z}} \mathbf{x}[m]e^{i\omega m} \qquad \text{where } m \triangleq -n \implies n = -m$$

$$= \sum_{m \in \mathbb{Z}} \mathbf{x}[m]e^{-i(-\omega)m}$$

$$\triangleq \tilde{\mathbf{x}}(-\omega)$$

by left hypothesis

$$\begin{bmatrix} \tilde{\mathbf{x}}^*(-\omega) \end{bmatrix} = \begin{bmatrix} \tilde{\mathbf{F}} (\mathbf{x}^*[n]) \end{bmatrix}$$
$$= \tilde{\mathbf{F}} (\mathbf{x}[n])$$
$$= \tilde{\mathbf{x}}(\omega) \end{bmatrix}$$

by Theorem M.1 page 238 by real-valued hypothesis


by definition of $\tilde{x}(\omega)$

(Definition M.1 page 237)


$$\begin{bmatrix} \tilde{\mathbf{x}}^*(\omega) \end{bmatrix} = \begin{bmatrix} \check{\mathbf{F}} (\mathbf{x}^*[-n]) \\ = \check{\mathbf{F}} (\mathbf{x}[-n]) \\ = \begin{bmatrix} \tilde{\mathbf{x}}(-\omega) \end{bmatrix}$$

by Theorem M.1 page 238 by real-valued hypothesis

by result (1)



Proposition M.2. Let $\check{x}(z)$ be the Z-transform (Definition J.4 page 208) and $\check{x}(\omega)$ the discrete-time Fourier TRANSFORM (Definition M.1 page 237) $of(x_n)$.



[♠]Proof:

1. Proof that (1) \implies (2):

$$\begin{aligned}
\check{\mathsf{x}}(z)\Big|_{z=1} &= \sum_{n\in\mathbb{Z}} x_n z^{-n} \\
&= \sum_{n\in\mathbb{Z}} x_n \\
&= c
\end{aligned}$$
by definition of $\check{\mathsf{x}}(z)$ (Definition J.4 page 208)
$$\begin{aligned}
\mathsf{because}\ z^n &= 1 \text{ for all } n \in \mathbb{Z} \\
\mathsf{by hypothesis}\ (1)
\end{aligned}$$

2. Proof that (2) \Longrightarrow (3):

$$\begin{split} \breve{\mathbf{x}}(\omega)\Big|_{\omega=0} &= \sum_{n\in\mathbb{Z}} x_n e^{-i\omega n} \Bigg|_{\omega=0} & \text{by definition of } \breve{\mathbf{x}}(\omega) & \text{(Definition M.1 page 237)} \\ &= \sum_{n\in\mathbb{Z}} x_n z^{-n} \Bigg|_{z=1} & \text{by definition of } \breve{\mathbf{x}}(z) & \text{(Definition J.4 page 208)} \\ &= c & \text{by hypothesis (2)} \end{split}$$

3. Proof that (3) \implies (1):

$$\begin{split} \sum_{n \in \mathbb{Z}} x_n &= \sum_{n \in \mathbb{Z}} x_n e^{-i\omega n} \bigg|_{\omega = 0} \\ &= \breve{\mathsf{x}}(\omega) & \text{by definition of } \breve{\mathsf{x}}(\omega) & \text{(Definition M.1 page 237)} \\ &= c & \text{by hypothesis (3)} \end{split}$$

Proposition M.3. If the coefficients are **real**, then the magnitude response (MR) is **symmetric**.

[♠]Proof:

$$\begin{aligned} \left| \tilde{\mathsf{h}}(-\omega) \right| &\triangleq \left| \check{\mathsf{h}}(z) \right|_{z=e^{-i\omega}} \\ &= \left| \sum_{m \in \mathbb{Z}} \mathsf{x}[m] e^{i\omega m} \right| \\ &= \left| \left(\sum_{m \in \mathbb{Z}} \mathsf{x}[m] e^{-i\omega m} \right)^* \right| \\ &= \left| \left(\sum_{m \in \mathbb{Z}} \mathsf{x}[m] e^{-i\omega m} \right)^* \right| \\ &\triangleq \left| \check{\mathsf{h}}(z) \right|_{z=e^{-i\omega}} \end{aligned}$$

$$\triangleq \left| \check{\mathsf{h}}(\omega) \right|$$

Proposition M.4.
$$\underbrace{\sum_{n \in \mathbb{Z}} (-1)^n x_n = c}_{(1) \text{ in "time"}} \iff \underbrace{\check{\mathsf{X}}(z)|_{z=-1} = c}_{(2) \text{ in "z domain"}} \iff \check{\check{\mathsf{X}}}(\omega)|_{\omega=\pi} = c}_{(3) \text{ in "frequency"}}$$

$$\iff \underbrace{\left(\sum_{n \in \mathbb{Z}} h_{2n}, \sum_{n \in \mathbb{Z}} h_{2n+1}\right) = \left(\frac{1}{2} \left(\sum_{n \in \mathbb{Z}} h_n + c\right), \frac{1}{2} \left(\sum_{n \in \mathbb{Z}} h_n - c\right)\right)}_{(4) \text{ sum of even, sum of odd}}$$

$$\forall c \in \mathbb{R}, (x_n)_{n \in \mathbb{Z}}, (y_n)_{n \in \mathbb{Z}} \in \ell_{\mathbb{R}}^2$$

[♠]Proof:

1. Proof that $(1) \Longrightarrow (2)$:

$$|\check{\mathbf{x}}(z)|_{z=-1} = \sum_{n \in \mathbb{Z}} x_n z^{-n} \Big|_{z=-1}$$

$$= \sum_{n \in \mathbb{Z}} (-1)^n x_n$$

$$= c$$

¹ Chui (1992) page 123



by (1)

2. Proof that $(2) \Longrightarrow (3)$:

$$\sum_{n \in \mathbb{Z}} x_n e^{-i\omega n} \bigg|_{\omega = \pi} = \sum_{n \in \mathbb{Z}} (-1)^n x_n$$

$$= \sum_{n \in \mathbb{Z}} (-1)^{-n} x_n \qquad = \sum_{n \in \mathbb{Z}} z^{-n} x_n \bigg|_{z = -1}$$

$$= c \qquad \qquad \text{by (2)}$$

3. Proof that (3) \Longrightarrow (1):

$$\sum_{n \in \mathbb{Z}} (-1)^n x_n = \sum_{n \in \mathbb{Z}} (-1)^{-n} x_n$$

$$= \sum_{n \in \mathbb{Z}} e^{-i\omega n} x_n \Big|_{\omega = \pi}$$

$$= c \qquad \text{by (3)}$$

- 4. Proof that $(2) \Longrightarrow (4)$:
 - (a) Define $A \triangleq \sum_{n \in \mathbb{Z}} h_{2n}$ $B \triangleq \sum_{n \in \mathbb{Z}} h_{2n+1}$.
 - (b) Proof that A B = c:

$$c = \sum_{n \in \mathbb{Z}} (-1)^n x_n$$

$$= \sum_{n \in \mathbb{Z}_e} (-1)^n x_n + \sum_{n \in \mathbb{Z}_o} (-1)^n x_n$$

$$= \sum_{n \in \mathbb{Z}} (-1)^{2n} x_{2n} + \sum_{n \in \mathbb{Z}} (-1)^{2n+1} x_{2n+1}$$

$$= \sum_{n \in \mathbb{Z}} x_{2n} - \sum_{n \in \mathbb{Z}} x_{2n+1}$$

by (2)

 $\triangleq A - B$ by definitions of A and B (c) Proof that $A + B = \sum_{n \in \mathbb{Z}} x_n$:

$$\sum_{n \in \mathbb{Z}} x_n = \sum_{n \text{ even}} x_n + \sum_{n \text{ odd}} x_n$$

$$= \sum_{n \in \mathbb{Z}} x_{2n} + \sum_{n \in \mathbb{Z}} x_{2n+1}$$

$$= A + B$$

by definitions of A and B

(d) This gives two simultaneous equations:

$$A - B = c$$

$$A + B = \sum_{n \in \mathbb{Z}} x_n$$

(e) Solutions to these equations give

$$\sum_{n \in \mathbb{Z}} x_{2n} \triangleq A \qquad \qquad = \frac{1}{2} \left(\sum_{n \in \mathbb{Z}} x_n + c \right)$$

$$\sum_{n \in \mathbb{Z}} x_{2n+1} \triangleq B \qquad \qquad = \frac{1}{2} \left(\sum_{n \in \mathbb{Z}} x_n - c \right)$$

5. Proof that $(2) \Leftarrow (4)$:

$$\sum_{n \in \mathbb{Z}} (-1)^n x_n = \sum_{n \in \mathbb{Z}_e} (-1)^n x_n + \sum_{n \in \mathbb{Z}_o} (-1)^n x_n$$

$$= \sum_{n \in \mathbb{Z}} (-1)^{2n} x_{2n} + \sum_{n \in \mathbb{Z}} (-1)^{2n+1} x_{2n+1}$$

$$= \sum_{n \in \mathbb{Z}} x_{2n} - \sum_{n \in \mathbb{Z}} x_{2n+1}$$

$$= \frac{1}{2} \left(\sum_{n \in \mathbb{Z}} x_n + c \right) - \frac{1}{2} \left(\sum_{n \in \mathbb{Z}} x_n - c \right)$$
by (3)
$$= c$$

Lemma M.1. Let $\tilde{f}(\omega)$ be the DTFT (Definition M.1 page 237) of a sequence $(x_n)_{n\in\mathbb{Z}}$.

 $\underbrace{(x_n \in \mathbb{R})_{n \in \mathbb{Z}}}_{\text{REAL-VALUED sequence}} \implies \underbrace{|\breve{\mathbf{x}}(\omega)|^2 = |\breve{\mathbf{x}}(-\omega)|^2}_{\text{EVEN}} \qquad \forall (x_n)_{n \in \mathbb{Z}} \in \mathscr{E}_{\mathbb{R}}^2$

♥Proof:

$$\begin{split} |\check{\mathsf{x}}(\omega)|^2 &= |\check{\mathsf{x}}(z)|^2\big|_{z=e^{i\omega}} \\ &= \check{\mathsf{x}}(z)\check{\mathsf{x}}^*(z)\big|_{z=e^{i\omega}} \\ &= \left[\sum_{n\in\mathbb{Z}} x_n z^{-n}\right] \left[\sum_{m\in\mathbb{Z}} x_m z^{-n}\right]^*\big|_{z=e^{i\omega}} \\ &= \left[\sum_{n\in\mathbb{Z}} x_n z^{-n}\right] \left[\sum_{m\in\mathbb{Z}} x_m^* (z^*)^{-m}\right]_{z=e^{i\omega}} \\ &= \sum_{n\in\mathbb{Z}} \sum_{m\in\mathbb{Z}} x_n x_m^* z^{-n} (z^*)^{-m}\big|_{z=e^{i\omega}} \\ &= \sum_{n\in\mathbb{Z}} \left[|x_n|^2 + \sum_{m>n} x_n x_m^* z^{-n} (z^*)^{-m} + \sum_{m< n} x_n x_m^* z^{-n} (z^*)^{-m}\right]_{z=e^{i\omega}} \\ &= \sum_{n\in\mathbb{Z}} \left[|x_n|^2 + \sum_{m>n} x_n x_m e^{i\omega(m-n)} + \sum_{m< n} x_n x_m e^{i\omega(m-n)}\right] \\ &= \sum_{n\in\mathbb{Z}} \left[|x_n|^2 + \sum_{m>n} x_n x_m e^{i\omega(m-n)} + \sum_{m>n} x_n x_m e^{-i\omega(m-n)}\right] \\ &= \sum_{n\in\mathbb{Z}} \left[|x_n|^2 + \sum_{m>n} x_n x_m (e^{i\omega(m-n)} + e^{-i\omega(m-n)})\right] \end{split}$$

$$= \sum_{n \in \mathbb{Z}} \left[|x_n|^2 + \sum_{m > n} x_n x_m 2 \cos[\omega(m-n)] \right]$$
$$= \sum_{n \in \mathbb{Z}} |x_n|^2 + 2 \sum_{n \in \mathbb{Z}} \sum_{m > n} x_n x_m \cos[\omega(m-n)]$$

Since cos is real and even, then $|\check{\mathbf{x}}(\omega)|^2$ must also be real and even.

Theorem M.3 (inverse DTFT). ² Let $\check{\mathbf{x}}(\omega)$ be the DISCRETE-TIME FOURIER TRANSFORM (Definition M.1 page 237) of a sequence $(x_n)_{n\in\mathbb{Z}}\in\mathscr{C}^2_{\mathbb{R}}$. Let $\check{\mathbf{x}}^{-1}$ be the inverse of $\check{\mathbf{x}}$.

$$\underbrace{\left\{ \begin{tabular}{l} \begin$$

№ Proof:

$$\frac{1}{2\pi} \int_{\alpha-\pi}^{\alpha+\pi} \check{\mathsf{x}}(\omega) e^{i\omega n} \, \mathrm{d}\omega = \frac{1}{2\pi} \int_{\alpha-\pi}^{\alpha+\pi} \left[\sum_{m \in \mathbb{Z}} x_m e^{-i\omega m} \right] e^{i\omega n} \, \mathrm{d}\omega \qquad \text{by definition of } \check{\mathsf{x}}(\omega)$$

$$= \frac{1}{2\pi} \int_{\alpha-\pi}^{\alpha+\pi} \sum_{m \in \mathbb{Z}} x_m e^{-i\omega(m-n)} \, \mathrm{d}\omega$$

$$= \frac{1}{2\pi} \sum_{m \in \mathbb{Z}} x_m \int_{\alpha-\pi}^{\alpha+\pi} e^{-i\omega(m-n)} \, \mathrm{d}\omega$$

$$= \frac{1}{2\pi} \sum_{m \in \mathbb{Z}} x_m \left[2\pi \bar{\delta}_{m-n} \right]$$

$$= x_n$$

Theorem M.4 (orthonormal quadrature conditions). 3 Let $\check{\mathbf{x}}(\omega)$ be the DISCRETE-TIME FOURIER TRANSFORM (Definition M.1 page 237) of a sequence $(x_n)_{n\in\mathbb{Z}}\in\mathscr{C}^2_{\mathbb{R}}$. Let $\bar{\delta}_n$ be the KRONECKER DELTA FUNCTION at n (Definition 2.12 page 20).

 \bigcirc Proof: Let $z \triangleq e^{i\omega}$.

³ Daubechies (1992) pages $132-137 \langle (5.1.20), (5.1.39) \rangle$





² J.S.Chitode (2009) page 3-95 ((3.6.2))

1. Proof that
$$2\sum_{n\in\mathbb{Z}}\left[\sum_{k\in\mathbb{Z}}x_ky_{k-2n}^*\right]e^{-i2\omega n}=\breve{\mathbf{x}}(\omega)\breve{\mathbf{y}}^*(\omega)+\breve{\mathbf{x}}(\omega+\pi)\breve{\mathbf{y}}^*(\omega+\pi)$$
:

$$\begin{split} &2\sum_{n\in\mathbb{Z}}\left[\sum_{k\in\mathbb{Z}}x_{k}y_{k-2n}^{*}\right]e^{-i2\omega n}\\ &=2\sum_{k\in\mathbb{Z}}x_{k}\sum_{n\in\mathbb{Z}}y_{k-2n}^{*}z^{-2n}\\ &=2\sum_{k\in\mathbb{Z}}x_{k}\sum_{n\,\text{even}}y_{k-n}^{*}z^{-n}\\ &=\sum_{k\in\mathbb{Z}}x_{k}\sum_{n\in\mathbb{Z}}y_{k-n}^{*}z^{-n}\left(1+e^{i\pi n}\right)\\ &=\sum_{k\in\mathbb{Z}}x_{k}\sum_{n\in\mathbb{Z}}y_{k-n}^{*}z^{-n}+\sum_{k\in\mathbb{Z}}x_{k}\sum_{n\in\mathbb{Z}}y_{k-n}^{*}z^{-n}e^{i\pi n}\\ &=\sum_{k\in\mathbb{Z}}x_{k}\sum_{m\in\mathbb{Z}}y_{m}^{*}z^{-(k-m)}+\sum_{k\in\mathbb{Z}}x_{k}\sum_{m\in\mathbb{Z}}y_{m}^{*}e^{-i(\omega+\pi)(k-m)} \qquad \text{where } m\triangleq k-n\\ &=\sum_{k\in\mathbb{Z}}x_{k}z^{-k}\sum_{m\in\mathbb{Z}}y_{m}^{*}z^{m}+\sum_{k\in\mathbb{Z}}x_{k}e^{-i(\omega+\pi)k}\sum_{m\in\mathbb{Z}}y_{m}^{*}e^{+i(\omega+\pi)m}\\ &=\sum_{k\in\mathbb{Z}}x_{k}e^{-i\omega k}\left[\sum_{m\in\mathbb{Z}}y_{m}e^{-i\omega m}\right]^{*}+\sum_{k\in\mathbb{Z}}x_{k}e^{-i(\omega+\pi)k}\left[\sum_{m\in\mathbb{Z}}y_{m}e^{-i(\omega+\pi)m}\right]^{*}\\ &\triangleq \check{\mathsf{X}}(\omega)\check{\mathsf{y}}^{*}(\omega)+\check{\mathsf{X}}(\omega+\pi)\check{\mathsf{y}}^{*}(\omega+\pi) \end{split}$$

2. Proof that $\sum_{m \in \mathbb{Z}} x_m y_{m-2n}^* = 0 \implies \check{\mathbf{x}}(\omega) \check{\mathbf{y}}^*(\omega) + \check{\mathbf{x}}(\omega + \pi) \check{\mathbf{y}}^*(\omega + \pi) = 0$:

$$0 = 2 \sum_{n \in \mathbb{Z}} \left[\sum_{k \in \mathbb{Z}} x_k y_{k-2n}^* \right] e^{-i2\omega n}$$
 by left hypothesis
= $\check{\mathbf{x}}(\omega) \check{\mathbf{y}}^*(\omega) + \check{\mathbf{x}}(\omega + \pi) \check{\mathbf{y}}^*(\omega + \pi)$ by item (1)

3. Proof that $\sum_{m \in \mathbb{Z}} x_m y_{m-2n}^* = 0 \iff \check{\mathbf{x}}(\omega) \check{\mathbf{y}}^*(\omega) + \check{\mathbf{x}}(\omega + \pi) \check{\mathbf{y}}^*(\omega + \pi) = 0$:

$$2\sum_{n\in\mathbb{Z}} \left[\sum_{k\in\mathbb{Z}} x_k y_{k-2n}^* \right] e^{-i2\omega n} = \breve{\mathsf{x}}(\omega) \breve{\mathsf{y}}^*(\omega) + \breve{\mathsf{x}}(\omega + \pi) \breve{\mathsf{y}}^*(\omega + \pi) \qquad \text{by item (1)}$$
$$= 0 \qquad \qquad \text{by right hypothesis}$$

Thus by the above equation, $\sum_{n\in\mathbb{Z}}\left[\sum_{k\in\mathbb{Z}}x_ky_{k-2n}^*\right]e^{-i2\omega n}=0$. The only way for this to be true is if $\sum_{k\in\mathbb{Z}}x_ky_{k-2n}^*=0$.

4. Proof that $\sum_{m\in\mathbb{Z}} x_m x_{m-2n}^* = \bar{\delta}_n \implies |\breve{\mathsf{x}}(\omega)|^2 + |\breve{\mathsf{x}}(\omega' + \pi)|^2 = 2$: Let $g_n \triangleq x_n$.

$$2 = 2 \sum_{n \in \mathbb{Z}} \bar{\delta}_{n \in \mathbb{Z}} e^{-i2\omega n}$$

$$= 2 \sum_{n \in \mathbb{Z}} \left[\sum_{k \in \mathbb{Z}} x_k y_{k-2n}^* \right] e^{-i2\omega n}$$
 by left hypothesis
$$= \check{\mathsf{x}}(\omega) \check{\mathsf{y}}^*(\omega) + \check{\mathsf{x}}(\omega + \pi) \check{\mathsf{y}}^*(\omega + \pi)$$
 by item (1)


5. Proof that $\sum_{m \in \mathbb{Z}} x_m x_{m-2n}^* = \bar{\delta}_n \iff |\check{\mathbf{x}}(\omega)|^2 + |\check{\mathbf{x}}(\omega' + \pi)|^2 = 2$: Let $g_n \triangleq x_n$.

$$2\sum_{n\in\mathbb{Z}}\left[\sum_{k\in\mathbb{Z}}x_ky_{k-2n}^*\right]e^{-i2\omega n} = \breve{\mathsf{x}}(\omega)\breve{\mathsf{y}}^*(\omega) + \breve{\mathsf{x}}(\omega+\pi)\breve{\mathsf{y}}^*(\omega+\pi) \qquad \text{by item (1)}$$

$$= 2 \qquad \qquad \text{by right hypothesis}$$

Thus by the above equation, $\sum_{n\in\mathbb{Z}} \left[\sum_{k\in\mathbb{Z}} x_k y_{k-2n}^*\right] e^{-i2\omega n} = 1$. The only way for this to be true is if $\sum_{k\in\mathbb{Z}} x_k y_{k-2n}^* = \bar{\delta}_n.$

Derivatives M.3



[♠]Proof:

1. Proof that $(A) \implies (B)$:

$$\begin{split} 0 &= \left[\frac{\mathrm{d}}{\mathrm{d}\omega}\right]^n \check{\mathbf{x}}(\omega)\Big|_{\omega=0} \\ &= \left[\frac{\mathrm{d}}{\mathrm{d}\omega}\right]^n \sum_{k \in \mathbb{Z}} x_k e^{-i\omega k}\Big|_{\omega=0} \\ &= \sum_{k \in \mathbb{Z}} x_k \left[\frac{\mathrm{d}}{\mathrm{d}\omega}\right]^n e^{-i\omega k}\Big|_{\omega=0} \\ &= \sum_{k \in \mathbb{Z}} x_k \left[(-i)^n k^n e^{-i\omega k}\right]\Big|_{\omega=0} \\ &= (-i)^n \sum_{k \in \mathbb{Z}} k^n x_k \end{split}$$


by hypothesis (A)

by definition of $\check{x}(\omega)$ (Definition M.1 page 237)

2. Proof that $(A) \leftarrow (B)$:

$$\begin{split} \left[\frac{\mathrm{d}}{\mathrm{d}\omega} \right]^n \check{\mathbf{x}}(\omega) \Big|_{\omega=0} &= \left[\frac{\mathrm{d}}{\mathrm{d}\omega} \right]^n \sum_{k \in \mathbb{Z}} x_k e^{-i\omega k} \Big|_{\omega=0} \\ &= \sum_{k \in \mathbb{Z}} x_k \left[\left[\frac{\mathrm{d}}{\mathrm{d}\omega} \right]^n e^{-i\omega k} \right] \Big|_{\omega=0} \\ &= \sum_{k \in \mathbb{Z}} x_k \left[(-i)^n k^n e^{-i\omega k} \right] \Big|_{\omega=0} \\ &= (-i)^n \sum_{k \in \mathbb{Z}} k^n x_k \\ &= 0 \end{split} \qquad \text{by hypothesis (B)}$$

⁴ ■ Vidakovic (1999) pages 82–83, ■ Mallat (1999) pages 241–242





3. Proof that $(C) \implies (D)$:

$$\begin{split} 0 &= \left[\frac{\mathrm{d}}{\mathrm{d}\omega}\right]^n \check{\mathsf{X}}(\omega) \Big|_{\omega = \pi} \\ &= \left[\frac{\mathrm{d}}{\mathrm{d}\omega}\right]^n \sum_{k \in \mathbb{Z}} x_k e^{-i\omega k} \Big|_{\omega = \pi} \\ &= \sum_{k \in \mathbb{Z}} x_k \left[\frac{\mathrm{d}}{\mathrm{d}\omega}\right]^n e^{-i\omega k} \Big|_{\omega = \pi} \\ &= \sum_{k \in \mathbb{Z}} x_k \left[(-i)^n k^n e^{-i\omega k}\right] \Big|_{\omega = \pi} \\ &= \sum_{k \in \mathbb{Z}} x_k \left[(-i)^n k^n (-1)^k\right] \\ &= (-i)^n \sum_{k \in \mathbb{Z}} (-1)^k k^n x_k \end{split}$$

by hypothesis (C)

by definition of \check{x} (Definition M.1 page 237)

4. Proof that $(C) \iff (D)$:

$$\begin{split} \left[\frac{\mathrm{d}}{\mathrm{d}\omega}\right]^n &\check{\mathsf{x}}(\omega) \Big|_{\omega=\pi} = \left[\frac{\mathrm{d}}{\mathrm{d}\omega}\right]^n \sum_{k \in \mathbb{Z}} x_k e^{-i\omega k} \Big|_{\omega=\pi} \\ &= \sum_{k \in \mathbb{Z}} x_k \left[\frac{\mathrm{d}}{\mathrm{d}\omega}\right]^n e^{-i\omega k} \Big|_{\omega=\pi} \\ &= \sum_{k \in \mathbb{Z}} x_k \left[(-i)^n k^n e^{-i\omega k}\right] \Big|_{\omega=\pi} \\ &= \sum_{k \in \mathbb{Z}} x_k \left[(-i)^n k^n (-1)^k\right] \\ &= (-i)^n \sum_{k \in \mathbb{Z}} (-1)^k k^n x_k \\ &= 0 \end{split}$$

by definition of \breve{x} (Definition M.1 page 237)

by hypothesis (D)

₽

• ...et la nouveauté de l'objet, jointe à son importance, a déterminé la classe à couronner cet ouvrage, en observant cependant que la manière dont l'auteur parvient à ses équations n'est pas exempte de difficultés, et que son analyse, pour les intégrer, laisse encore quelque chose à désirer, soit relativement à la généralité, soit même du coté de la rigueur.



• ... and the innovation of the subject, together with its importance, convinced the committee to crown this work. By observing however that the way in which the author arrives at his equations is not free from difficulties, and the analysis of which, to integrate them, still leaves something to be desired, either relative to generality, or even on the side of rigour.

A competition awards committee consisting of the mathematical giants Lagrange, Laplace, Legendre, and others, commenting on Fourier's 1807 landmark paper *Dissertation on the propagation of heat in solid bodies* that introduced the *Fourier Series*. ¹

N.1 Definition

The *Fourier Series* expansion of a periodic function is simply a complex trigonometric polynomial. In the special case that the periodic function is even, then the Fourier Series expansion is a cosine polynomial.

Definition N.1. ²

The Fourier Series operator $\hat{\mathbf{F}}: \mathbf{L}_{\mathbb{R}}^2 \to \mathcal{E}_{\mathbb{R}}^2$ is defined as $[\hat{\mathbf{F}}f](n) \triangleq \frac{1}{\sqrt{\tau}} \int_0^{\tau} f(x)e^{-i\frac{2\pi}{\tau}nx} \, \mathrm{dx} \qquad \forall f \in \left\{ f \in \mathbf{L}_{\mathbb{R}}^2 | f \text{ is periodic with period } \tau \right\}$

² ■ Katznelson (2004) page 3

Inverse Fourier Series operator N.2

Theorem N.1. Let $\hat{\mathbf{F}}$ be the Fourier Series operator.



The inverse Fourier Series operator
$$\hat{\mathbf{F}}^{-1}$$
 is given by
$$\left[\hat{\mathbf{F}}^{-1}\left(\tilde{\mathbf{x}}_{n}\right)_{n\in\mathbb{Z}}\right](x)\triangleq\frac{1}{\sqrt{\tau}}\sum_{n\in\mathbb{Z}}\tilde{\mathbf{x}}_{n}e^{i\frac{2\pi}{\tau}nx}\qquad\forall(\tilde{\mathbf{x}}_{n})\in\mathscr{C}_{\mathbb{R}}^{2}$$

№ Proof: The proof of the pointwise convergence of the Fourier Series is notoriously difficult. It was conjectured in 1913 by Nokolai Luzin that the Fourier Series for all square summable periodic functions are pointwise convergent: Luzin (1913)

Fifty-three years later (1966) at a conference in Moscow, Lennart Axel Edvard Carleson presented one of the most spectacular results ever in mathematics; he demonstrated that the Luzin conjecture is indeed correct. Carleson formally published his result that same year:

Carleson (1966)

Carleson's proof is expounded upon in Reyna's (2002) 175 page book:

de Reyna (2002)

Interestingly enough, Carleson started out trying to disprove Luzin's conjecture. Carleson said this in an interview published in 2001:³ "Well, the problem of course presents itself already when you are a student and I was thinking of the problem on and off, but the situation was more interesting than that. The great authority in those days was Zygmund and he was completely convinced that what one should produce was not a proof but a counter-example. When I was a young student in the United States, I met Zygmund and I had an idea how to produce some very complicated functions for a counter-example and Zygmund encouraged me very much to do so. I was thinking about it for about 15 years on and off, on how to make these counter-examples work and the interesting thing that happened was that I suddenly realized why there should be a counter-example and how you should produce it. I thought I really understood what was the back ground and then to my amazement I could prove that this "correct" counter-example couldn't exist and therefore I suddenly realized that what you should try to do was the opposite, you should try to prove what was not fashionable, namely to prove convergence. The most important aspect in solving a mathematical problem is the conviction of what is the true result! Then it took like 2 or 3 years using the technique that had been developed during the past 20 years or so. It is actually a problem related to analytic functions basically even though it doesn't look that way.

For now, if you just want some intuitive justification for the Fourier Series, and you can somehow imagine that the Dirichlet kernel generates a comb function of Dirac delta functions, then perhaps what follows may help (or not). It is certainly not mathematically rigorous and is by no means a real proof (but at least it is less than 175 pages).

$$\begin{aligned} \left[\hat{\mathbf{F}}^{-1}\hat{\mathbf{F}}\mathbf{x}\right](x) &= \hat{\mathbf{F}}^{-1}\underbrace{\left[\frac{1}{\sqrt{\tau}}\int_{0}^{\tau}\mathbf{x}(x)e^{-i\frac{2\pi}{\tau}nx}\,\mathrm{d}\mathbf{x}\right]}_{\hat{\mathbf{F}}\mathbf{x}} \qquad \text{by definition of } \hat{\mathbf{F}} \end{aligned} \qquad \text{Definition N.1 page 247}$$

$$&= \frac{1}{\sqrt{\tau}}\sum_{n\in\mathbb{Z}}\left[\frac{1}{\sqrt{\tau}}\int_{0}^{\tau}\mathbf{x}(u)e^{-i\frac{2\pi}{\tau}nu}\,\mathrm{d}\mathbf{u}\right]e^{i\frac{2\pi}{\tau}nx} \qquad \text{by definition of } \hat{\mathbf{F}}^{-1}$$

$$&= \frac{1}{\sqrt{\tau}}\sum_{n\in\mathbb{Z}}\frac{1}{\sqrt{\tau}}\int_{0}^{\tau}\mathbf{x}(u)e^{-i\frac{2\pi}{\tau}nu}e^{i\frac{2\pi}{\tau}nx}\,\mathrm{d}\mathbf{u}$$

$$&= \frac{1}{\sqrt{\tau}}\sum_{n\in\mathbb{Z}}\frac{1}{\sqrt{\tau}}\int_{0}^{\tau}\mathbf{x}(u)e^{i\frac{2\pi}{\tau}n(x-u)}\,\mathrm{d}\mathbf{u}$$

³ Carleson and Engquist (2001)



$$=\int_{0}^{\tau} \mathbf{x}(u) \frac{1}{\tau} \sum_{n \in \mathbb{Z}} e^{i\frac{2\tau}{\tau} (\mathbf{x} - \mathbf{u})} \, \mathrm{d}\mathbf{u}$$

$$=\int_{0}^{\tau} \mathbf{x}(u) \left[\sum_{n \in \mathbb{Z}} \delta(\mathbf{x} - \mathbf{u} - n\tau) \right] \, \mathrm{d}\mathbf{u}$$

$$=\sum_{n \in \mathbb{Z}} \int_{u=0}^{u=\tau} \mathbf{x}(u) \delta(\mathbf{x} - \mathbf{u} - n\tau) \, \mathrm{d}\mathbf{u}$$

$$=\sum_{n \in \mathbb{Z}} \int_{v=n\tau=0}^{v=\tau} \mathbf{x}(v - n\tau) \delta(\mathbf{x} - v) \, \mathrm{d}\mathbf{v} \qquad \text{where } v \triangleq u + n\tau$$

$$=\sum_{n \in \mathbb{Z}} \int_{v=n\tau}^{v=(n+1)\tau} \mathbf{x}(v - n\tau) \delta(\mathbf{x} - v) \, \mathrm{d}\mathbf{v} \qquad \text{where } v \triangleq u + n\tau$$

$$=\sum_{n \in \mathbb{Z}} \int_{v=n\tau}^{v=(n+1)\tau} \mathbf{x}(v) \delta(\mathbf{x} - v) \, \mathrm{d}\mathbf{v} \qquad \text{because } \mathbf{x} \text{ is periodic with period } \tau$$

$$=\int_{\mathbb{R}} \mathbf{x}(v) \delta(\mathbf{x} - v) \, \mathrm{d}\mathbf{v}$$

$$=\mathbf{x}(\mathbf{x})$$

$$=\mathbf{I} \tilde{\mathbf{x}}(n) \qquad \text{by definition of } \mathbf{I} \qquad \text{(Definition C.3 page 112)}$$

$$[\hat{\mathbf{F}} \hat{\mathbf{F}}^{-1} \tilde{\mathbf{x}}](n) = \hat{\mathbf{F}} \left[\frac{1}{\sqrt{\tau}} \sum_{k \in \mathbb{Z}} \tilde{\mathbf{x}}(k) e^{i\frac{2\tau}{\tau}k\mathbf{x}} \right] \qquad \text{by definition of } \hat{\mathbf{F}}^{-1}$$

$$= \frac{1}{\tau} \int_{0}^{\tau} \left[\sum_{k \in \mathbb{Z}} \tilde{\mathbf{x}}(k) e^{i\frac{2\tau}{\tau}(k-n)\mathbf{x}} \right] d\mathbf{x}$$

$$= \sum_{k \in \mathbb{Z}} \tilde{\mathbf{x}}(k) \frac{1}{\tau} \int_{0}^{\tau} e^{i\frac{2\tau}{\tau}(k-n)\mathbf{x}} \, \mathrm{d}\mathbf{x} \right]$$

$$= \sum_{k \in \mathbb{Z}} \tilde{\mathbf{x}}(k) \frac{1}{\tau} \left[\frac{1}{\tau^2 \pi}(k-n) e^{i\frac{2\tau}{\tau}(k-n)\mathbf{x}} \right]_{0}^{\tau}$$

$$= \sum_{k \in \mathbb{Z}} \tilde{\mathbf{x}}(k) \frac{1}{\tau^2 \pi}(k-n) \left[e^{i2\pi(k-n)} - 1 \right]$$

 $= \sum_{k=1}^{\infty} \tilde{\mathbf{x}}(k) \, \bar{\delta}(k-n) \lim_{x \to 0} \left| \frac{e^{i2\pi x} - 1}{i2\pi x} \right|$ $= \tilde{\mathbf{x}}(n) \left. \frac{\frac{\mathrm{d}}{\mathrm{d}\mathbf{x}} \left(e^{i2\pi x} - 1 \right)}{\frac{\mathrm{d}}{\mathrm{d}\mathbf{x}} (i2\pi x)} \right|_{\mathbf{x} = 0}$

by l'Hôpital's rule

 $= \tilde{\mathsf{x}}(n) \left. \frac{i2\pi e^{i2\pi x}}{i2\pi} \right|_{x=0}$

 $= \tilde{x}(n)$

 $= \mathbf{I}\tilde{\mathbf{x}}(n)$

by definition of I

(Definition C.3 page 112)

The **Fourier Series adjoint** operator $\hat{\mathbf{F}}^*$ is given by



🚧 Frames and Bases Structure and Design [VERSION 021] 🌉



№PROOF:

$$\begin{split} \left\langle \hat{\mathbf{F}} \mathbf{x}(x) \,|\, \tilde{\mathbf{y}}(n) \right\rangle_{\mathbb{Z}} &= \left\langle \frac{1}{\sqrt{\tau}} \int_{0}^{\tau} \mathbf{x}(x) e^{-i\frac{2\pi}{\tau}nx} \,\, \mathrm{d}\mathbf{x} \,|\, \tilde{\mathbf{y}}(n) \right\rangle_{\mathbb{Z}} \quad \text{by definition of } \hat{\mathbf{F}} \\ &= \frac{1}{\sqrt{\tau}} \int_{0}^{\tau} \mathbf{x}(x) \left\langle e^{-i\frac{2\pi}{\tau}nx} \,|\, \tilde{\mathbf{y}}(n) \right\rangle_{\mathbb{Z}} \,\, \mathrm{d}\mathbf{x} \quad \text{by additivity property of } \left\langle \triangle \,|\, \nabla \right\rangle \\ &= \int_{0}^{\tau} \mathbf{x}(x) \frac{1}{\sqrt{\tau}} \left\langle \tilde{\mathbf{y}}(n) \,|\, e^{-i\frac{2\pi}{\tau}nx} \right\rangle_{\mathbb{Z}}^{*} \,\, \mathrm{d}\mathbf{x} \quad \text{by property of } \left\langle \triangle \,|\, \nabla \right\rangle \\ &= \int_{0}^{\tau} \mathbf{x}(x) \big[\hat{\mathbf{F}}^{-1} \tilde{\mathbf{y}}(n) \big]^{*} \,\, \mathrm{d}\mathbf{x} \qquad \text{by definition of } \hat{\mathbf{F}}^{-1} \end{split} \tag{Theorem N.1 page 248} \\ &= \left\langle \mathbf{x}(x) \,|\, \hat{\underline{\mathbf{F}}}^{-1} \tilde{\mathbf{y}}(n) \right\rangle_{\mathbb{R}} \end{split}$$

The Fourier Series operator has several nice properties:

- \clubsuit $\hat{\mathbf{F}}$ is unitary 4 (Corollary N.1 page 250).
- Because $\hat{\mathbf{F}}$ is unitary, it automatically has several other nice properties such as being *isometric*, and satisfying *Parseval's equation*, satisfying *Plancheral's formula*, and more (Corollary N.2 page 250).

Corollary N.1. Let **I** be the identity operator and let $\hat{\mathbf{F}}$ be the Fourier Series operator with adjoint $\hat{\mathbf{F}}^*$.

$$\{\hat{\mathbf{F}}\hat{\mathbf{F}}^* = \hat{\mathbf{F}}^*\hat{\mathbf{F}} = \mathbf{I} \}$$
 ($\hat{\mathbf{F}}$ is **unitary** ... and thus also normal and isometric)

 igthedarmoonup Proof: This follows directly from the fact that $\hat{\mathbf{F}}^* = \hat{\mathbf{F}}^{-1}$ (Theorem N.2 page 249).

Corollary N.2. Let $\hat{\mathbf{F}}$ be the Fourier series operator with adjoint $\hat{\mathbf{F}}^*$ and inverse $\hat{\mathbf{F}}^{-1}$.

 $\ ^{\ }\ ^{\ }$ PROOF: These results follow directly from the fact that $\hat{\mathbf{F}}$ is unitary (Corollary N.1 page 250) and from the properties of unitary operators (Theorem C.26 page 136).

N.3 Fourier series for compactly supported functions

Theorem N.3.

T H M

The set
$$\left\{ \left. \frac{1}{\sqrt{\tau}} e^{i\frac{2\pi}{\tau}nx} \right| n \in \mathbb{Z} \right. \right\}$$

is an Orthonormal basis for all functions f(x) with support in $[0:\tau]$.

⁴unitary operators: Definition C.14 page 135



APPENDIX O___

FAST WAVELET TRANSFORM (FWT)

The Fast Wavelet Transform can be computed using simple discrete filter operations (as a conjugate mirror filter).

Definition 0.1 (Wavelet Transform). Let the wavelet transform $W: \{f: \mathbb{R} \to \mathbb{C}\} \to \{w: \mathbb{Z}^2 \to \mathbb{C}\}$ be defined as ¹

$$[\mathbf{W}\mathsf{f}](j,n) \triangleq \left\langle \mathsf{f}(x) \, | \, \psi_{k,n}(x) \right\rangle$$

Definition 0.2. The following relations are defined as described below:

-	2 chilitation 3.2. The following the defined the december 2 court								
		scaling coefficients					$\langle f(x) \phi_{j,n}(x) \rangle$		
	D E F	wavelet coefficients	$w_j: \mathbb{Z} \to \mathbb{C}$	such that	$w_j(n)$	≜	$\langle f(x) \psi_{j,n}(x) \rangle$		
١	F	scaling filter coefficients	$\bar{h} : \mathbb{Z} o \mathbb{C}$	such that	h(n)	≜	h(-n)		
		wavelet filter coefficients	$\bar{g}:\mathbb{Z}\to\mathbb{C}$	such that	$\bar{g}(n)$	≜	g(-n)		

The scaling and wavelet filter coefficients at scale j are equal to the filtered and downsampled (Theorem $\ref{theorem}$) scaling filter coefficients at scale j+1:²

- \clubsuit The convolution (Definition J.3 page 207) of $v_{j+1}(n)$ with $\bar{\mathsf{h}}(n)$ and then downsampling by 2 produces $v_j(n)$.
- $\stackrel{\checkmark}{=}$ The convolution of $v_{j+1}(n)$ with $\bar{\mathfrak{g}}(n)$ and then downsampling by 2 produces $w_j(n)$.

This is formally stated and proved in the next theorem.

Laplace Transform
$$\mathcal{L}f(s) \triangleq \langle f(x) | e^{sx} \rangle \triangleq \int_x f(x)e^{-sx} dx$$
Continuous Fourier Transform $\mathcal{F}f(\omega) \triangleq \langle f(x) | e^{i\omega x} \rangle \triangleq \int_x f(x)e^{-i\omega x} dx$
Fourier Series Transform $\mathcal{F}_s f(k) \triangleq \langle f(x) | e^{i\frac{2\pi}{T}kx} \rangle \triangleq \int_x f(x)e^{-i\frac{2\pi}{T}kx} dx$
Laplace Transform $\mathcal{F}_s f(\omega) \triangleq \langle f(x) | e^{i\omega x} \rangle \triangleq \int_x f(x)e^{-i\omega x} dx$
Laplace Transform $\mathcal{F}_s f(\omega) \triangleq \langle f(x) | e^{i\omega x} \rangle \triangleq \int_x f(x)e^{-i\omega x} dx$
Laplace Transform $\mathcal{F}_s f(\omega) \triangleq \langle f(x) | e^{i\omega x} \rangle \triangleq \int_x f(x)e^{-i\omega x} dx$
Laplace Transform $\mathcal{F}_s f(\omega) \triangleq \langle f(x) | e^{i\omega x} \rangle \triangleq \int_x f(x)e^{-i\omega x} dx$
Laplace Transform $\mathcal{F}_s f(\omega) \triangleq \langle f(x) | e^{i\omega x} \rangle \triangleq \int_x f(x)e^{-i\omega x} dx$
Laplace Transform $\mathcal{F}_s f(\omega) \triangleq \langle f(x) | e^{i\omega x} \rangle \triangleq \int_x f(\omega)e^{-i\omega x} dx$
Laplace Transform $\mathcal{F}_s f(\omega) \triangleq \langle f(x) | e^{i\omega x} \rangle \triangleq \int_x f(\omega)e^{-i\omega x} dx$
Laplace Transform $\mathcal{F}_s f(\omega) \triangleq \langle f(x) | e^{i\omega x} \rangle \triangleq \int_x f(\omega)e^{-i\omega x} dx$
Laplace Transform $\mathcal{F}_s f(\omega) \triangleq \langle f(\omega) | e^{i\omega x} \rangle \triangleq \int_x f(\omega)e^{-i\omega x} dx$
Laplace Transform $\mathcal{F}_s f(\omega) \triangleq \langle f(\omega) | e^{i\omega x} \rangle \triangleq \int_x f(\omega)e^{-i\omega x} dx$
Laplace Transform $\mathcal{F}_s f(\omega) \triangleq \langle f(\omega) | e^{i\omega x} \rangle \triangleq \int_x f(\omega)e^{-i\omega x} dx$
Laplace Transform $\mathcal{F}_s f(\omega) \triangleq \langle f(\omega) | e^{i\omega x} \rangle \triangleq \int_x f(\omega)e^{-i\omega x} dx$
Laplace Transform $\mathcal{F}_s f(\omega) \triangleq \langle f(\omega) | e^{i\omega x} \rangle \triangleq \int_x f(\omega)e^{-i\omega x} dx$
Laplace Transform $\mathcal{F}_s f(\omega) \triangleq \langle f(\omega) | e^{i\omega x} \rangle \triangleq \int_x f(\omega)e^{-i\omega x} dx$
Laplace Transform $\mathcal{F}_s f(\omega) \triangleq \langle f(\omega) | e^{i\omega x} \rangle \triangleq \int_x f(\omega)e^{-i\omega x} dx$
Laplace Transform $\mathcal{F}_s f(\omega) \triangleq \langle f(\omega) | e^{i\omega x} \rangle \triangleq \int_x f(\omega)e^{-i\omega x} dx$
Laplace Transform $\mathcal{F}_s f(\omega) \triangleq \langle f(\omega) | e^{i\omega x} \rangle \triangleq \int_x f(\omega)e^{-i\omega x} dx$
Laplace Transform $\mathcal{F}_s f(\omega) \triangleq \langle f(\omega) | e^{i\omega x} \rangle \triangleq \int_x f(\omega)e^{-i\omega x} dx$
Laplace Transform $\mathcal{F}_s f(\omega) \triangleq \langle f(\omega) | e^{i\omega x} \rangle \triangleq \int_x f(\omega)e^{-i\omega x} dx$
Laplace Transform $\mathcal{F}_s f(\omega) \triangleq \langle f(\omega) | e^{i\omega x} \rangle \triangleq \int_x f(\omega)e^{-i\omega x} dx$
Laplace Transform $\mathcal{F}_s f(\omega) \triangleq \langle f(\omega) | e^{i\omega x} \rangle \triangleq$

¹Notice that this definition is similar to the definition of transforms of other analysis systems:

² Mallat (1999) page 257, Burrus et al. (1998) page 35

 $v_j(n) = [\bar{h} \star v_{j+1}](2n)$ $w_i(n) = [\bar{g} \star v_{i+1}](2n)$

[♠]Proof:

$$\begin{split} v_{j}(n) &= \left\langle f(x) \mid \phi_{j,n}(x) \right\rangle \\ &= \left\langle f(x) \mid \sqrt{2^{j}} \phi \left(2^{j} x - n \right) \right\rangle \\ &= \left\langle f(x) \mid \sqrt{2^{j}} \sqrt{2} \sum_{m} h(m) \phi \left(2(2^{j} x - n) - m \right) \right\rangle \\ &= \left\langle f(x) \mid \sum_{m} h(j) \sqrt{2^{j+1}} \phi \left(2^{j+1} x - 2n - m \right) \right\rangle \\ &= \sum_{m} h(m) \left\langle f(x) \mid \sqrt{2^{j+1}} \phi \left(2^{j+1} x - 2n - m \right) \right\rangle \\ &= \sum_{m} h(m) \left\langle f(x) \mid \phi_{j+1,2n+m}(x) \right\rangle \\ &= \sum_{m} h(m) v_{j+1}(2n + m) \\ &= \sum_{p} h(p - 2n) v_{j+1}(p) \\ &= \left[h + v_{j+1} \right] (2n) \\ w_{j}(n) &= \left\langle f(x) \mid \psi_{j,n}(x) \right\rangle \\ &= \left\langle f(x) \mid \sqrt{2^{j}} \psi \left(2^{j} x - n \right) \right\rangle \\ &= \left\langle f(x) \mid \sqrt{2^{j}} \psi \left(2^{j} x - n \right) \right\rangle \\ &= \sum_{m} g(m) \left\langle f(x) \mid \sqrt{2^{j+1}} \phi \left(2^{j+1} x - 2n - m \right) \right\rangle \\ &= \sum_{m} g(m) \left\langle f(x) \mid \psi_{j+1,2n+m}(x) \right\rangle \\ &= \sum_{m} g(m) \left\langle f(x) \mid \psi_{j+1,2n+m}(x) \right\rangle \\ &= \sum_{m} g(m) v_{j+1}(2n + m) \\ &= \sum_{p} g(p - 2n) v_{j+1}(p) \\ &= \sum_{m} g(2n - p) v_{j+1}(p) \end{split}$$
 let $p = 2n + m \iff m = p - 2n$

These filtering and downsampling operations are equivalent to the operations performed by a filter bank. Therefore, a filter bank can be used to implement a Fast Wavelet Transform (FWT), as illustrated in Figure 0.1 (page 253).



 $= [\bar{g} \star v_{i+1}](2n)$

₽




Figure O.1: *k*-Stage Fast Wavelet Transform



POWER SPECTRUM FUNCTIONS

P.1 Correlation

Definition P.1 and Definition P.2 define four quantities. In this document, the quantities' notation and terminology are similar to those used in the study of random processes.

```
Definition P.1. <sup>1</sup> Let \langle \triangle | \nabla \rangle be the STANDARD INNER PRODUCT in L^2_{\mathbb{R}} (Definition D.1 page 141).

Property Rfg(n) \triangleq \langle f(x) | T^n g(x) \rangle, n \in \mathbb{Z}; f, g \in L^2_{\mathbb{F}}, is the cross-correlation function of f and g. R_{ff}(n) \triangleq \langle f(x) | T^n f(x) \rangle, n \in \mathbb{Z}; f \in L^2_{\mathbb{F}}, is the autocorrelation function of f.
```

Definition P.2. Let $R_{fg}(n)$ and $R_{ff}(n)$ be the sequences defined in Definition P.1 page 255. Let $\mathbf{Z}(x_n)$ be the Z-TRANSFORM (Definition J.4 page 208) of a sequence $(x_n)_{n\in\mathbb{Z}}$.

```
\check{S}_{fg}(z) \triangleq \mathbf{Z}[\mathsf{R}_{fg}(n)], \quad f,g \in \mathcal{L}_{\mathbb{F}}^2, \quad is \ the \ complex \ cross-power \ spectrum \ of \ f \ and \ g.

\check{S}_{ff}(z) \triangleq \mathbf{Z}[\mathsf{R}_{fg}(n)], \quad f,g \in \mathcal{L}_{\mathbb{F}}^2, \quad is \ the \ complex \ auto-power \ spectrum \ of \ f.
```

Power Spectrum P.2

```
Definition P.3. ^3 Let \check{S}_{fg}(z) and \check{S}_{ff}(z) be the functions defined in Definition P.2 page 255. \check{S}_{fg}(\omega) \triangleq \check{S}_{fg}(e^{i\omega}), \ \forall f,g \in L_{\mathbb{F}}^2, \ is the \ {\it cross-power spectrum of f} \ and \ g. \check{S}_{ff}(\omega) \triangleq \check{S}_{ff}(e^{i\omega}), \ \forall f \in L_{\mathbb{F}}^2, \ is the \ {\it auto-power spectrum of f}.
```

Theorem P.1. ⁴ Let $\tilde{S}_{fg}(\omega)$ and $\tilde{S}_{ff}(\omega)$ be defined as in Definition P.3 (page 255).

Let
$$\tilde{\mathbf{f}}(\omega)$$
 be the Fourier transform (Definition I.2 page 196) of a function $\mathbf{f}(x) \in \mathcal{L}_{\mathbb{F}}^2$.

$$\tilde{\mathbf{S}}_{\mathrm{fg}}(\omega) = 2\pi \sum_{n \in \mathbb{Z}} \tilde{\mathbf{f}}(\omega + 2\pi n) \tilde{\mathbf{g}}^*(\omega + 2\pi n) \quad \forall \mathbf{f}, \mathbf{g} \in \mathcal{L}_{\mathbb{F}}^2$$

$$\tilde{\mathbf{S}}_{\mathrm{ff}}(\omega) = 2\pi \sum_{n \in \mathbb{Z}} \left| \tilde{\mathbf{f}}(\omega + 2\pi n) \right|^2 \quad \forall \mathbf{f} \in \mathcal{L}_{\mathbb{F}}^2$$

¹ Chui (1992) page 134, Papoulis (1991) pages 294–332 ⟨(10-29), (10-169)⟩

³ Chui (1992) page 134, Papoulis (1991) page 333 ⟨(10-179)⟩

⁴ Chui (1992) page 135

 \bigcirc Proof: Let $z \triangleq e^{i\omega}$.

$$\begin{split} \tilde{\mathbf{S}}_{\mathsf{f}\mathsf{g}}(\omega) &\triangleq \check{\mathbf{S}}_{\mathsf{f}\mathsf{g}}(z) \\ &= \sum_{n \in \mathbb{Z}} \mathsf{R}_{\mathsf{f}\mathsf{g}}(n) z^{-n} \\ &= \sum_{n \in \mathbb{Z}} \left\langle \mathsf{f}(x) \, | \, \mathsf{g}(x-n) \right\rangle z^{-n} \\ &= \sum_{n \in \mathbb{Z}} \left\langle \tilde{\mathbf{F}}[\mathsf{f}(x)] \, | \, \tilde{\mathbf{F}}\big[\mathsf{g}(x-n)\big] \right\rangle z^{-n} \\ &= \sum_{n \in \mathbb{Z}} \left\langle \tilde{\mathsf{f}}(v) \, | \, e^{-ivn} \tilde{\mathsf{g}}(v) \right\rangle z^{-n} \\ &= \sum_{n \in \mathbb{Z}} \sqrt{2\pi} \left[\frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} \tilde{\mathsf{f}}(v) \tilde{\mathsf{g}}^*(v) e^{ivu} \, \mathrm{dv} \right]_{u=n} z^{-n} \\ &= \sqrt{2\pi} \sum_{n \in \mathbb{Z}} \left[\tilde{\mathbf{F}}^{-1} \Big(\sqrt{2\pi} \tilde{\mathsf{f}}(v) \tilde{\mathsf{g}}^*(v) \Big) \right]_{u=n} e^{-i\omega n} \\ &= 2\pi \sum_{n \in \mathbb{Z}} \tilde{\mathsf{f}}(\omega + 2\pi n) \tilde{\mathsf{g}}^*(\omega + 2\pi n) \end{split}$$

by definition of
$$\tilde{S}_{fg}$$
 (Definition P.3 page 255)
by definition of \tilde{S}_{fg} (Definition P.2 page 255)
by definition of \tilde{S}_{fg} (Definition P.3 page 255)
by unitary property of \tilde{F} (Theorem I.3 page 197)
by shift relation (Theorem I.4 page 198)
by definition of $\boldsymbol{L}_{\mathbb{R}}^2$ (Definition D.1 page 141)
by Theorem I.1 page 197
by $IPSF$ with $\tau=1$ (Theorem 3.3 page 49)

(Definition P.3 page 255)

$$\begin{split} \tilde{S}_{ff}(\omega) &= \left. \tilde{S}_{fg}(\omega) \right|_{g=f} & \text{by definition of } \tilde{S}_{fg}(\omega) \\ &= 2\pi \sum_{n \in \mathbb{Z}} \tilde{f}(\omega + 2\pi n) \tilde{g}^*(\omega + 2\pi n) \right|_{g \triangleq f} & \text{by previous result} \\ &= 2\pi \sum_{n \in \mathbb{Z}} \tilde{f}(\omega + 2\pi n) \tilde{f}^*(\omega + 2\pi n) \\ &= 2\pi \sum_{n \in \mathbb{Z}} \left| \tilde{f}(\omega + 2\pi n) \right|^2 & \text{because } |z|^2 \triangleq zz^* \quad \forall z \in \mathbb{C} \end{split}$$

Proposition P.1. Let $\tilde{S}_{ff}(\omega)$ be defined as in Definition P.3 (page 255).

 $\tilde{S}_{ff}(\omega) \geq 0 \quad \text{(NON-NEGATIVE)}$

[♠]Proof:

$$\tilde{S}_{ff}(\omega) = 2\pi \sum_{n \in \mathbb{Z}} |\tilde{f}(\omega + 2\pi n)|^2$$
 by Theorem P.1 page 255
$$\geq 0$$
 because $|z| \geq 0 \quad \forall z \in \mathbb{C}$

Proposition P.2. Let $\tilde{S}_{fg}(\omega)$ and $\tilde{S}_{ff}(\omega)$ be defined as in Definition P.3 (page 255).

 $\tilde{S}_{fg}(\omega + 2\pi) = \tilde{S}_{fg}(\omega) \quad (\text{PERIODIC with period } 2\pi)$ $\tilde{S}_{ff}(\omega + 2\pi) = \tilde{S}_{ff}(\omega) \quad (\text{PERIODIC with period } 2\pi)$



№PROOF:

$$\tilde{S}_{fg}(\omega + 2\pi) = 2\pi \sum_{n \in \mathbb{Z}} \tilde{f}(\omega + 2\pi + 2\pi n) \tilde{g}^*(\omega + 2\pi + 2\pi n)$$
by Theorem P.1 page 255
$$= 2\pi \sum_{n \in \mathbb{Z}} \tilde{f}[\omega + 2\pi (n+1)] \tilde{g}^*[\omega + 2\pi (n+1)]$$

$$= 2\pi \sum_{m \in \mathbb{Z}} \tilde{f}[\omega + 2\pi m] \tilde{g}^*[\omega + 2\pi m]$$
where $m \triangleq n+1$

$$= \tilde{S}_{fg}(\omega)$$
by Theorem P.1 page 255
$$\tilde{S}_{ff}(\omega + 2\pi) = \tilde{S}_{fg}(\omega + 2\pi)|_{g=f}$$

$$= \tilde{S}_{fg}(\omega)|_{g=f}$$
by previous result
$$= \tilde{S}_{ff}(\omega)$$

Proposition P.3. Let $\tilde{S}_{fg}(\omega)$ and $\tilde{S}_{ff}(\omega)$ be defined as in Definition P.3 (page 255).

♥Proof:

$$\begin{split} \tilde{S}_{fg}(-\omega) &= 2\pi \sum_{n \in \mathbb{Z}} \tilde{f}(-\omega + 2\pi n) \tilde{g}^*(-\omega + 2\pi n) \\ &= 2\pi \sum_{n \in \mathbb{Z}} \tilde{f}^*(\omega - 2\pi n) \tilde{g}(\omega - 2\pi n) \\ &= 2\pi \sum_{m \in \mathbb{Z}} \tilde{g}(\omega + 2\pi m) \tilde{f}^*(\omega + 2\pi m) \\ &= 2\pi \sum_{m \in \mathbb{Z}} \tilde{g}(\omega + 2\pi m) \tilde{f}^*(\omega + 2\pi m) \\ &= \tilde{S}_{gf}(\omega) \end{split} \qquad \text{by Theorem P.1 page 255} \\ \tilde{S}_{fg}(\pi - \omega) &= 2\pi \sum_{n \in \mathbb{Z}} \tilde{f}(\pi - \omega + 2\pi n) \tilde{g}^*(\pi - \omega + 2\pi n) \\ &= 2\pi \sum_{n \in \mathbb{Z}} \tilde{f}^*(-\pi + \omega - 2\pi n) \tilde{g}(-\pi + \omega - 2\pi n) \\ &= 2\pi \sum_{n \in \mathbb{Z}} \tilde{f}^*(\pi + \omega - 2\pi - 2\pi n) \tilde{g}(\pi + \omega - 2\pi - 2\pi n) \\ &= 2\pi \sum_{n \in \mathbb{Z}} \tilde{f}^*(\pi + \omega + 2\pi n) \tilde{g}(\pi + \omega - 2\pi - 2\pi n) \\ &= 2\pi \sum_{n \in \mathbb{Z}} \tilde{f}^*(\pi + \omega + 2\pi n) \tilde{f}^*(\pi + \omega + 2\pi n) \\ &= 2\pi \sum_{n \in \mathbb{Z}} \tilde{g}(\pi + \omega + 2\pi m) \tilde{f}^*(\pi + \omega + 2\pi m) \\ &= \tilde{S}_{gf}(\pi + \omega) \end{split} \qquad \text{where } m \triangleq -n - 1 \\ &= \tilde{S}_{gf}(\pi + \omega) \end{aligned}$$

$$\tilde{S}_{ff}(\pi - \omega) = \tilde{S}_{fg}(\pi - \omega)|_{g \triangleq f}$$

by definition of $g(g \triangleq f)$

 $=\tilde{S}_{ff}(+\omega)$

$$= \tilde{S}_{gf}(\pi + \omega) \Big|_{g \triangleq f}$$
$$= \tilde{S}_{ff}(\pi + \omega)$$

by previous result by definition of g ($g \triangleq f$)

Proposition P.4. Let $\tilde{S}_{ff}(\omega)$ be the AUTO-POWER SPECTRUM (Definition P.3 page 255) of a function $f(x) \in \mathcal{L}^2_{\mathbb{R}}$ and $\tilde{S}'_{ff}(\omega) \triangleq \frac{d}{d\omega} \tilde{S}_{ff}(\omega)$ (Definition D.2 page 141).

$$\begin{cases} \text{and } S'_{\mathsf{ff}}(\omega) \stackrel{\triangle}{=} \frac{\mathsf{d}}{\mathsf{d}\omega} S_{\mathsf{ff}}(\omega) \text{ (Definition D.2 page 141)}. \\ \\ \left\{ \begin{array}{l} \text{(a).} & \mathsf{f} \text{ is REAL} \quad and \\ \text{(b).} & \tilde{S}_{\mathsf{ff}}(\omega) \text{ is CONTINUOUS } at \omega = 0 \end{array} \right\} \qquad \Longrightarrow \begin{cases} \text{(1).} & \tilde{S}'_{\mathsf{ff}}(0) = 0 \quad and \\ \text{(2).} & \tilde{S}'_{\mathsf{ff}}(\omega) = -\tilde{S}'_{\mathsf{ff}}(-\omega) \quad \forall \omega \in \mathbb{R} \\ \\ \text{ANTI-SYMMETRIC about 0} \end{cases} \\ \\ \left\{ \begin{array}{l} \text{(c).} & \mathsf{f} \text{ is REAL} \quad and \\ \text{(d).} & \tilde{S}'_{\mathsf{ff}}(\omega) \text{ is CONTINUOUS } at \omega = \pi \end{array} \right\} \qquad \Longrightarrow \begin{cases} \text{(3).} & \tilde{S}'_{\mathsf{ff}}(\pi) = 0 \quad and \\ \text{(4).} & \tilde{S}'_{\mathsf{ff}}(\pi + \omega) = -\tilde{S}'_{\mathsf{ff}}(\pi - \omega) \quad \forall \omega \in \mathbb{R} \\ \\ \text{ANTI-SYMMETRIC about } \pi \end{cases} \end{cases}$$

№ Proof: This follows from Proposition P.3 (page 257) and Proposition D.1 (page 141).

Theorem P.2 (next) is a major result and provides strong motivation for bothering with *power spectrum* functions in the first place. In particular, the *auto-power spectrum* being *bounded* provides a necessary and sufficient condition for a sequence of functions $(\phi(x-n))_{n\in\mathbb{Z}}$ to be a *Riesz basis* (Definition 2.13 page 27) for the *span* span $(\phi(x-n))$ of the sequence.

Theorem P.2. ⁵ Let $\tilde{S}_{ff}(\omega)$ be defined as in Definition P.3 (page 255). Let $\|\cdot\|$ be defined as in Definition D.1 (page 141). Let 0 < A < B.

$$\underbrace{\left\{A\sum_{n\in\mathbb{N}}\left|a_{n}\right|^{2}\leq\left\|\sum_{n\in\mathbb{Z}}a_{n}\phi(x-n)\right\|^{2}\leq B\sum_{n\in\mathbb{N}}\left|\alpha_{n}\right|^{2}\quad\forall(a_{n})\in\mathscr{C}_{\mathbb{F}}^{2}\right\}}_{\left(\left(\phi(x-n)\right)\text{ is a Riesz basis for span }\left(\left(\phi(x-n)\right)\right)\text{ (Theorem 2.13 page 28)}}\Longleftrightarrow\left\{A\leq\widetilde{S}_{\phi\phi}(\omega)\leq B\right\}$$

[♠]Proof:

1. lemma:

$$\left\| \sum_{n \in \mathbb{Z}} a_n \phi(x - n) \right\|^2 = \left\| \tilde{\mathbf{F}} \sum_{n \in \mathbb{Z}} a_n \phi(x - n) \right\|^2 \qquad \text{because } \tilde{\mathbf{F}} \text{ is } unitary \text{ (Theorem I.2 page 197)}$$

$$= \left\| \breve{\mathbf{a}}(\omega) \tilde{\phi}(\omega) \right\|^2 \qquad \text{by Proposition 3.13 page 47}$$

$$= \int_{\mathbb{R}} \left| \breve{\mathbf{a}}(\omega) \tilde{\phi}(\omega) \right|^2 d\omega \qquad \text{by definition of } \| \cdot \|$$

$$= \sum_{n \in \mathbb{Z}} \int_0^{2\pi} \left| \breve{\mathbf{a}}(\omega + 2\pi n) \tilde{\phi}(\omega + 2\pi n) \right|^2 d\omega$$

$$= \int_0^{2\pi} \sum_{n \in \mathbb{Z}} \left| \breve{\mathbf{a}}(\omega + 2\pi n) \right|^2 \left| \tilde{\phi}(\omega + 2\pi n) \right|^2 d\omega$$

$$= \int_0^{2\pi} \sum_{n \in \mathbb{Z}} \left| \breve{\mathbf{a}}(\omega) \right|^2 \left| \tilde{\phi}(\omega + 2\pi n) \right|^2 d\omega \qquad \text{by Proposition M.1 page 237}$$



$$= \int_0^{2\pi} |\check{\mathbf{a}}(\omega)|^2 \frac{1}{2\pi} 2\pi \sum_{n \in \mathbb{Z}} |\tilde{\phi}(\omega + 2\pi n)|^2 d\omega$$
$$= \frac{1}{2\pi} \int_0^{2\pi} |\check{\mathbf{a}}(\omega)|^2 \tilde{\mathbf{S}}_{\phi\phi}(\omega) d\omega$$

by definition of $\tilde{\mathsf{S}}_{\phi\phi}(\omega)$ (Theorem P.1 page 255)

2. lemma:

$$\begin{split} \frac{1}{2\pi} \int_0^{2\pi} |\check{\mathbf{a}}(\omega)|^2 \, \mathrm{d}\omega &= \frac{1}{2\pi} \int_0^{2\pi} \left| \sum_{n \in \mathbb{Z}} a_n e^{-i\omega n} \right|^2 \, \mathrm{d}\omega \qquad \qquad \text{by def. of } DTFT \text{ (Definition M.1 page 237)} \\ &= \frac{1}{2\pi} \int_0^{2\pi} \left[\sum_{n \in \mathbb{Z}} a_n e^{-i\omega n} \right] \left[\sum_{m \in \mathbb{Z}} a_m e^{-i\omega m} \right]^* \, \mathrm{d}\omega \\ &= \frac{1}{2\pi} \int_0^{2\pi} \left[\sum_{n \in \mathbb{Z}} a_n e^{-i\omega n} \right] \left[\sum_{m \in \mathbb{Z}} a_m^* e^{i\omega m} \right] \, \mathrm{d}\omega \\ &= \frac{1}{2\pi} \sum_{n \in \mathbb{Z}} \sum_{m \in \mathbb{Z}} a_n a_m^* \int_0^{2\pi} e^{-i\omega(n-m)} \, \mathrm{d}\omega \\ &= \frac{1}{2\pi} \sum_{n \in \mathbb{Z}} \sum_{m \in \mathbb{Z}} a_n a_m^* 2\pi \bar{\delta}_{nm} \\ &= \sum_{n \in \mathbb{Z}} \left| a_n \right|^2 \qquad \qquad \text{by definition of } \bar{\delta} \text{ (Definition 2.12 page 20)} \end{split}$$

3. Proof for (\Leftarrow) case:

$$A \sum_{n \in \mathbb{Z}} |a_n|^2 = \frac{A}{2\pi} \int_0^{2\pi} |\check{\mathbf{a}}(\omega)|^2 d\omega \qquad \text{by (2) lemma page 259}$$

$$= \frac{1}{2\pi} \int_0^{2\pi} |\check{\mathbf{a}}(\omega)|^2 A d\omega$$

$$\leq \frac{1}{2\pi} \int_0^{2\pi} |\check{\mathbf{a}}(\omega)|^2 \tilde{\mathbf{S}}_{\phi\phi}(\omega) d\omega \qquad \text{by right hypothesis}$$

$$= \left\| \sum_{n \in \mathbb{Z}} a_n \phi(x - n) \right\|^2 \qquad \text{by (1) lemma page 258}$$

$$= \frac{1}{2\pi} \int_0^{2\pi} |\check{\mathbf{a}}(\omega)|^2 \tilde{\mathbf{S}}_{\phi\phi}(\omega) d\omega \qquad \text{by (1) lemma page 258}$$

$$\leq \frac{1}{2\pi} \int_0^{2\pi} |\check{\mathbf{a}}(\omega)|^2 B d\omega \qquad \text{by right hypothesis}$$

$$= \frac{B}{2\pi} \int_0^{2\pi} |\check{\mathbf{a}}(\omega)|^2 d\omega$$

$$= B \sum_{n \in \mathbb{Z}} |a_n|^2 \qquad \text{by (2) lemma page 259}$$

4. Proof for (\Longrightarrow) case:

- (a) Let $Y \triangleq \left\{ \omega \in [0:2\pi] | \tilde{S}_{\phi\phi}(\omega) > \alpha \right\}$ and $X \triangleq \left\{ \omega \in [0:2\pi] | \tilde{S}_{\phi\phi}(\omega) < \alpha \right\}$
- (b) Let $\mathbb{1}_{A(x)}$ be the $set\ indicator$ (Definition 3.2 page 40) of a set A. Let $(b_n)_{n\in\mathbb{Z}}$ be the $inverse\ DTFT$ (Theorem M.3 page 243) of $\mathbb{1}_Y(\omega)$ such that $\mathbb{1}_Y(\omega)\triangleq\sum b_ne^{-i\omega n}\triangleq \tilde{\mathbf{b}}(\omega)$.





Let $(a_n)_{n\in\mathbb{Z}}$ be the *inverse DTFT* (Theorem M.3 page 243) of $\mathbb{1}_X(\omega)$ such that $\mathbb{1}_X(\omega) \triangleq \sum_{n\in\mathbb{N}} a_n e^{-i\omega n} \triangleq \check{\mathbf{a}}(\omega)$.

(c) Proof that $\alpha \leq B$:

Let $\mu(A)$ be the *measure* of a set A.

$$\begin{split} \boxed{B} \sum_{n \in \mathbb{Z}} \left| b_n \right|^2 &\geq \left\| \sum_{n \in \mathbb{Z}} b_n \phi(x - n) \right\|^2 \qquad \text{by left hypothesis} \\ &= \frac{1}{2\pi} \int_0^{2\pi} \left| \tilde{\mathbf{b}}(\omega) \right|^2 \tilde{\mathbf{S}}_{\phi\phi}(\omega) \, \mathrm{d}\omega \qquad \text{by (1) lemma page 258} \\ &= \frac{1}{2\pi} \int_0^{2\pi} \left| \mathbb{1}_Y(\omega) \right|^2 \tilde{\mathbf{S}}_{\phi\phi}(\omega) \, \mathrm{d}\omega \qquad \text{by definition of } \mathbb{1}_Y(\omega) \qquad \text{(item (4b) page 259)} \\ &= \frac{1}{2\pi} \int_Y \left| \mathbb{1} \right|^2 \tilde{\mathbf{S}}_{\phi\phi}(\omega) \, \mathrm{d}\omega \qquad \text{by definition of } \mathbb{1}_Y(\omega) \qquad \text{(item (4b) page 259)} \\ &\geq \frac{\alpha}{2\pi} \mu(Y) \qquad \text{by definition of } Y \qquad \text{(item (4a) page 259)} \\ &= \int_0^{2\pi} \left| \mathbb{1}_Y(\omega) \right|^2 \, \mathrm{d}\omega \qquad \text{by definition of } \mathbb{1}_Y(\omega) \qquad \text{(item (4b) page 259)} \\ &= \int_0^{2\pi} \left| \sum_{n \in \mathbb{Z}} b_n e^{-i\omega n} \right|^2 \, \mathrm{d}\omega \qquad \text{by definition of } \mathbb{6}(\omega) \qquad \text{(item (4b) page 259)} \\ &= \int_0^{2\pi} \left| \tilde{\mathbf{b}}(\omega) \right|^2 \, \mathrm{d}\omega \qquad \text{by definition of } \tilde{\mathbf{b}}(\omega) \qquad \text{(item (4b) page 259)} \\ &= \frac{2\pi}{2\pi} \sum_{n \in \mathbb{Z}} \left| b_n \right|^2 \qquad \text{by (2) lemma page 259} \end{split}$$

(d) Proof that $\tilde{S}_{\phi\phi}(\omega) \leq B$:

- (i). $\tilde{S}_{\phi\phi}(\omega) > \alpha$ whenever $\omega \in Y$ (item (4a) page 259).
- (ii). But even then, $\alpha \leq B$ (item (4c) page 260).
- (iii). So, $\tilde{S}_{\phi\phi}(\omega) \leq B$.

(e) Proof that $A \leq \alpha$:

Let $\mu(A)$ be the *measure* of a set *A*.

$$\begin{split} \boxed{A} \sum_{n \in \mathbb{Z}} \left| a_n \right|^2 & \leq \left\| \sum_{n \in \mathbb{Z}} a_n \phi(x - n) \right\|^2 \qquad \text{by left hypothesis} \\ & = \frac{1}{2\pi} \int_0^{2\pi} \left| \mathbf{a}(\omega) \right|^2 \tilde{\mathbf{S}}_{\phi\phi}(\omega) \, \mathrm{d}\omega \qquad \text{by (1) lemma page 258} \\ & = \frac{1}{2\pi} \int_0^{2\pi} \left| \mathbb{1}_X(\omega) \right|^2 \tilde{\mathbf{S}}_{\phi\phi}(\omega) \, \mathrm{d}\omega \qquad \text{by definition of } \mathbb{1}_X(\omega) \qquad \text{(Definition 3.2 page 40)} \\ & = \frac{1}{2\pi} \int_X \left| 1 \right|^2 \tilde{\mathbf{S}}_{\phi\phi}(\omega) \, \mathrm{d}\omega \qquad \text{by definition of } \mathbb{1}_X(\omega) \qquad \text{(Definition 3.2 page 40)} \\ & \leq \frac{\alpha}{2\pi} \mu(X) \qquad \text{by definition of } X \qquad \text{(item (4a) page 259)} \\ & = \int_0^{2\pi} \left| \mathbb{1}_X(\omega) \right|^2 \, \mathrm{d}\omega \qquad \text{by definition of } \mathbb{1}_X(\omega) \qquad \text{(Definition 3.2 page 40)} \\ & = \int_0^{2\pi} \left| \sum_{n \in \mathbb{Z}} a_n e^{-i\omega n} \right|^2 \, \mathrm{d}\omega \qquad \text{by definition of } (a_n) \qquad \text{((2) lemma page 259)} \\ & = \int_0^{2\pi} \left| \mathbf{a}(\omega) \right|^2 \, \mathrm{d}\omega \qquad \text{by definition of } \mathbf{a}(\omega) \qquad \text{((2) lemma page 259)} \end{split}$$

$$= \left[\alpha\right] \sum_{n \in \mathbb{Z}} \left| a_n \right|^2$$

by (2) lemma page 259

- (f) Proof that $A \leq \tilde{S}_{\phi\phi}(\omega)$:
 - $\tilde{S}_{\phi\phi}(\omega) < \alpha \text{ whenever } \omega \in X$ (item (4a) page 259).
 - (ii). But even then, $A \leq \alpha$ (item (4e) page 260).
 - (iii). So, $A \leq \tilde{S}_{\phi\phi}(\omega)$.

In the case that f and g are *orthonormal*, the spectral density relations simplify considerably (next).

Theorem P.3. ⁶ Let \tilde{S}_{ff} and \tilde{S}_{fg} be the SPECTRAL DENSITY FUNCTIONS (Definition P.3 page 255).

I	$\langle f(x) f(x-n) \rangle$	=	$\bar{\delta}_n$	(($f(x-n)$) is orthonormal) ($f(x)$ is orthogonal to ($g(x-n)$))	\Leftrightarrow	$\tilde{S}_{ff}(\omega)$	=	1	$\forall f \in \mathcal{L}_{\mathbb{F}}^2$
M	$\langle f(x) g(x-n) \rangle$	=	0	(f(x) is orthogonal to $(g(x - n))$)	\iff	$\tilde{S}_{fg}(\omega)$	=	0	$\forall f,g \in \mathcal{L}_{\mathbb{F}}^2$

[♠]Proof:

- 1. Proof that $\langle f(x) | f(x-n) \rangle = \bar{\delta}_n \iff \tilde{S}_{ff}(\omega) = 1$: This follows directly from Theorem P.2 (page 258) with A = B = 1 (by Parseval's Identity Theorem 2.9 page 22 since $\{T^n f\}$ is orthonormal)
- 2. Alternate proof that $\langle f(x) | f(x-n) \rangle = \bar{\delta}_n \implies \tilde{S}_{ff}(\omega) = 1$:

$$\begin{split} \tilde{\mathsf{S}}_{\mathsf{ff}}(\omega) &= \sum_{n \in \mathbb{Z}} \mathsf{R}_{\mathsf{ff}}(n) e^{-i\omega n} & \text{by definition of } \tilde{\mathsf{S}}_{\mathsf{fg}} & \text{(Definition P.3 page 255)} \\ &= \sum_{n \in \mathbb{Z}} \left\langle \mathsf{f}(x) \, | \, \mathsf{f}(x-n) \right\rangle e^{-i\omega n} & \text{by definition of } \mathsf{R}_{\mathsf{ff}} & \text{(Definition P.1 page 255)} \\ &= \sum_{n \in \mathbb{Z}} \bar{\delta}_n e^{-i\omega n} & \text{by left hypothesis} \\ &= 1 & \text{by definition of } \bar{\delta} & \text{(Definition 2.12 page 20)} \end{split}$$

3. Alternate proof that $\langle f(x) | f(x-n) \rangle = \bar{\delta}_n \iff \tilde{S}_{ff}(\omega) = 1$:

$$\begin{split} &\langle \mathsf{f}(x) \, | \, \mathsf{f}(x-n) \rangle \\ &= \langle \tilde{\mathbf{F}}\mathsf{f}(x) \, | \, \tilde{\mathbf{F}}\mathsf{f}(x-n) \rangle \\ &= \langle \tilde{\mathbf{f}}(\omega) \, | \, e^{-i\omega n} \tilde{\mathbf{f}}(\omega) \rangle \\ &= \int_{\mathbb{R}} \tilde{\mathbf{f}}(\omega) e^{i\omega n} \tilde{\mathbf{f}}^*(\omega) \, \mathrm{d}\omega \\ &= \int_{\mathbb{R}} |\tilde{\mathbf{f}}(\omega)|^2 e^{i\omega n} \, \mathrm{d}\omega \\ &= \sum_{n \in \mathbb{Z}} \int_{2\pi n}^{2\pi (n+1)} |\tilde{\mathbf{f}}(\omega)|^2 e^{i\omega n} \, \mathrm{d}\omega \\ &= \sum_{n \in \mathbb{Z}} \int_{0}^{2\pi} |\tilde{\mathbf{f}}(u+2\pi n)|^2 e^{i(u+2\pi n)n} \, \mathrm{d}u \\ &= \frac{1}{2\pi} \int_{0}^{2\pi} \left[2\pi \sum_{n \in \mathbb{Z}} |\tilde{\mathbf{f}}(u+2\pi n)|^2 \right] e^{iun} e^{i2\pi n n} \, \mathrm{d}u \\ &= \frac{1}{2\pi} \int_{0}^{2\pi} \tilde{\mathbf{S}}_{\mathsf{ff}}(\omega) e^{iun} \, \mathrm{d}u \\ &= \frac{1}{2\pi} \int_{0}^{2\pi} \tilde{\mathbf{S}}_{\mathsf{ff}}(\omega) e^{iun} \, \mathrm{d}u \end{split} \qquad \text{by Theorem P.1 page 255}$$

 6 Hernández and Weiss (1996) page 50 ⟨Proposition 2.1.11⟩, \bigcirc Wojtaszczyk (1997) page 23 ⟨Corollary 2.9⟩, ☐ IGARI (1996) PAGES 214-215 (LEMMA 9.2), ☐ PINSKY (2002) PAGE 306 (COROLLARY 6.4.9)



$$=\frac{1}{2\pi}\int_0^{2\pi}e^{iun}\,\mathrm{d}u$$
$$=\bar{\delta}_n$$

by right hypothesis

by definition of $\bar{\delta}$

(Definition 2.12 page 20)

4. Proof that $\langle f(x) | g(x - n) \rangle = 0 \implies \tilde{S}_{fg}(\omega) = 0$:

$$\tilde{S}_{fg}(\omega) = \sum_{n \in \mathbb{Z}} R_{fg}(n) e^{-i\omega n}$$

$$= \sum_{n \in \mathbb{Z}} \langle f(x) | g(x - n) \rangle e^{-i\omega n}$$

$$= \sum_{n \in \mathbb{Z}} 0 e^{-i\omega n}$$

$$= 0$$

by definition of \tilde{S}_{fg}

(Definition P.3 page 255)

by definition of R_{fg}

(Definition P.1 page 255)

by left hypothesis

5. Proof that $\langle f(x) | g(x - n) \rangle = 0 \iff \tilde{S}_{fg}(\omega) = 0$:

$$\langle f(x) | g(x - n) \rangle$$

$$= \langle \tilde{F}f(x) | \tilde{F}g(x - n) \rangle$$

$$= \langle \tilde{f}(\omega) | e^{-i\omega n} \tilde{g}(\omega) \rangle$$

$$= \int_{\mathbb{R}} \tilde{f}(\omega) e^{i\omega n} \tilde{g}^*(\omega) d\omega$$

$$= \int_{\mathbb{R}} \tilde{f}(\omega) \tilde{g}^*(\omega) e^{i\omega n} d\omega$$

$$= \sum_{n \in \mathbb{Z}} \int_{2\pi n}^{2\pi (n+1)} \tilde{f}(\omega) \tilde{g}^*(\omega) e^{i\omega n} d\omega$$

$$= \sum_{n \in \mathbb{Z}} \int_{0}^{2\pi} \tilde{f}(u + 2\pi n) \tilde{g}^*(u + 2\pi n) e^{i(u + 2\pi n)n} du$$

$$= \frac{1}{2\pi} \int_{0}^{2\pi} \left[2\pi \sum_{n \in \mathbb{Z}} \tilde{f}(u + 2\pi n) \tilde{g}^*(u + 2\pi n) \right] e^{iun} e^{i2\pi nn} du$$

$$= \frac{1}{2\pi} \int_{0}^{2\pi} \tilde{S}_{fg}(u) e^{iun} du$$

 $= \frac{1}{2\pi} \int_{0}^{2\pi} 0 \cdot e^{iun} \, du$

by unitary property of $ilde{\mathbf{F}}$ (Theorem I.3 page 197)

by $shift\ property\ of\ ilde{\mathbf{F}}$ (Theorem I.4 page 198)

by definition of $\langle \triangle \mid \nabla \rangle$ (Definition D.1 page 141)

where $u \triangleq \omega - 2\pi n \implies \omega = u + 2\pi n$

by Theorem P.1 page 255

by right hypothesis

₽



Q.1 Definitions

Definition Q.1. ¹ *Let* (Ω, \mathbb{E}, P) *be a* PROBABILITY SPACE.

```
The function x: \Omega \to \mathbb{R} is a random variable.
The function y: \mathbb{R} \times \Omega \to \mathbb{R} is a random process.
```

The random process $x(t, \omega)$, where t commonly represents time and $\omega \in \Omega$ is an outcome of an experiment, can take on more specialized forms depending on whether t and ω are fixed or allowed to vary. These forms are illustrated in Figure Q.1 page 263^2 and Figure Q.2 page 264.

$\mathbf{x}(t,\omega)$	fixed t	variable t				
fixed ω	number	time function				
variable ω	random variable	random process				

Figure Q.1: Specialized forms of a random process $x(t, \omega)$

Definition Q.2. 3 *Let* x(t) *and* y(t) *be random processes.*

n	The mean	$\mu_{X}(t)$	of $x(t)$ is	$\mu_{X}(t)$	≜	E[x(t)]
Ē	The cross-correlation	$R_{xy}(t)$	of $x(t)$ and $y(t)$ is	$R_{xy}(t, u)$	≜	$E[x(t)y^*(u)]$
F	The auto-correlation function					$E\left[x(t)x^*(u)\right]$

Remark Q.1. ⁴ The equation $\int_{u \in \mathbb{R}} \mathsf{R}_{\mathsf{x}\mathsf{x}}(t,u) \mathsf{f}(u)$ du is a *Fredholm integral equation of the first kind* and $\mathsf{R}_{\mathsf{x}\mathsf{x}}(t,u)$ is the *kernel* of the equation.

² Papoulis (1991) pages 285–286

³ Papoulis (1984) page 216 $\langle R_{xy}(t_1, t_2) = E\{x(t_1)y^*(t_2)\}$ (9-35) \rangle

⁴ Fredholm (1900), Fredholm (1903) page 365, Michel and Herget (1993) page 97, Keener (1988) page 101




Figure Q.2: Example of a random process $x(t, \omega)$

Q.2 Properties

Theorem Q.1. Let x(t) and y(t) be random processes with cross-correlation $R_{xy}(t, u)$ and let $R_{xx}(t, u)$ be the auto-correlation of x(t).

$$\begin{array}{c} \mathsf{T} \\ \mathsf{H} \\ \mathsf{M} \end{array} \begin{array}{c} \mathsf{R}_{\mathsf{X}\mathsf{X}}(t,u) &= \; \mathsf{R}_{\mathsf{X}\mathsf{X}}^*(u,t) \quad \text{(CONJUGATE SYMMETRIC)} \\ \mathsf{R}_{\mathsf{X}\mathsf{Y}}(t,u) &= \; \mathsf{R}_{\mathsf{Y}\mathsf{X}}^*(u,t) \end{array}$$

№ Proof:

$$R_{xx}(t,u) \triangleq E[x(t)x^*(u)] \qquad = E[x^*(u)x(t)] = (E[x(u)x^*(t)])^* \qquad \triangleq R_{xx}^*(u,t)$$

$$R_{xy}(t,u) \triangleq E[x(t)y^*(u)] \qquad = E[y^*(u)x(t)] = (E[y(u)x^*(t)])^* \qquad \triangleq R_{yx}^*(u,t)$$

₽

APPENDIX	\exists
----------	-----------

SPECTRAL THEORY

R.1 Operator Spectrum

Definition R.1. Let $\mathbf{A} \in \mathcal{B}(\mathbf{X}, \mathbf{Y})$ be an operator over the linear spaces $\mathbf{X} = (X, F, \oplus, \otimes)$ and $\mathbf{Y} \triangleq (Y, F, \oplus, \otimes)$. Let $\mathcal{N}(\mathbf{A})$ be the NULL SPACE of \mathbf{A} .

An **eigenvalue** of **A** is any value λ such that there exists **x** such that $\mathbf{A}\mathbf{x} = \lambda \mathbf{x}$.

The **eigenspace** H_{λ} of A at eigenvalue λ is $\mathcal{N}(A - \lambda I)$.

An **eigenvector** of \mathbf{A} associated with eigenvalue λ is any element of $\mathcal{N}(\mathbf{A} - \lambda \mathbf{I})$.

Example R.1. ² Let **D** be the differntial operator.

	The set $\{\epsilon$	$e^{\lambda x} \lambda \in \mathbb{C}$	are the eigenvectors of D .			
	$\rho(\mathbf{D}) =$	Ø	(D has no non-spectral points whatsoever)			
E X	$\sigma_{p}(\mathbf{D}) = \sigma(\mathbf{D})$		(the spectrum of ${\bf D}$ is all eigenvalues)			
	$\sigma_{c}(\mathbf{D}) =$	Ø	(D has no continuous spectrum)			
	$\sigma_{r}(\mathbf{D}) =$	Ø	(D has no resolvent spectrum)			

№ Proof:

$$(\mathbf{D} - \lambda \mathbf{I})e^{\lambda x} = \mathbf{D}e^{\lambda x} - \lambda \mathbf{I}e^{\lambda x}$$
$$= \lambda e^{\lambda x} - \lambda e^{\lambda x}$$
$$= 0$$

This theorem and proof needs more work and investigation to prove/disprove its claims.

Definition R.2. ³ Let $\mathbf{A} \in \mathcal{B}(\mathbf{X}, \mathbf{Y})$ be an operator over the linear spaces $\mathbf{X} = (X, F, \oplus, \otimes)$ and $\mathbf{Y} \triangleq (Y, F, \oplus, \otimes)$.

 $\forall \lambda \in \mathbb{C}$

¹ ■ Bollobás (1999) page 168, ■ Descartes (1637a), ■ Descartes (1954), ■ Cayley (1858), ■ Hilbert (1904) page 67, ■ Hilbert (1912),

² Pedersen (2000) page 79

³ Michel and Herget (1993) page 439

DEF

Table R.1: Spectrum of an operator A

The **resolvent set** $\rho(A)$ of operator A is defined as

$$\rho(\mathbf{A}) \triangleq \begin{cases} 1. & \mathcal{N}(\mathbf{A} - \lambda \mathbf{I}) = \{0\} \\ \lambda \in F \mid 2. & \mathcal{R}(\mathbf{A} - \lambda \mathbf{I}) = \mathbf{X} \\ 3. & (\mathbf{A} - \lambda \mathbf{I})^{-1} \in \mathcal{B}(\mathbf{X}, \mathbf{Y}) \end{cases} & \text{(inverse is continuous/bounded).} \qquad and$$

The **spectrum** $\sigma(\mathbf{A})$ of operator **A** is defined as


$$\sigma(\mathbf{A}) \triangleq F \setminus \rho(\mathbf{A}).$$

Definition R.3. ⁴ Let $A \in \mathcal{B}(X, Y)$ be an operator over the linear spaces $X = (X, F, \oplus, \otimes)$ and $Y \triangleq$ (Y, F, \oplus, \otimes) . The **point spectrum** $\sigma_{o}(\mathbf{A})$ of operator **A** is defined as

 $\sigma_{\mathsf{D}}(\mathbf{A}) \triangleq \{ \lambda \in F | 1. \ \mathcal{N}(\mathbf{A} - \lambda \mathbf{I}) \supseteq \{ 0 \}$ (has non-zero eigenvector) The **residual spectrum** $\sigma_r(\mathbf{A})$ of operator **A** is defined as (no non-zero eigenvectors)

$$\sigma_{\mathsf{r}}(\mathbf{A}) \triangleq \left\{ \lambda \in F \middle| \begin{array}{l} 1. & \mathcal{N}(\mathbf{A} - \lambda \mathbf{I}) = \{\emptyset\} \\ 2. & \overline{\mathcal{R}}(\mathbf{A} - \lambda \mathbf{I}) \neq \mathbf{X} \end{array} \right. \text{ (not dense in } \mathbf{X} - \text{has gaps).}$$

The **continuous spectrum** $\sigma_c(\mathbf{A})$ of operator **A** is defined as



The spectral components' definitions are illustrated in the figure to the left and summarized in Table R.1 (page 266). Let a family of operators $\mathbf{B}(\lambda)$ be defined with respect to an operator **A** such that $\mathbf{B}(\lambda) \triangleq$ $(\mathbf{A} - \lambda \mathbf{I})$. Normally, we might expect a "normal" or "regular" or even "mundane" operator $\mathbf{B}(\lambda)$ to have the properties

- 1. $\mathbf{B}(\lambda)\mathbf{x} = 0$ if and only if $\mathbf{x} = 0$
- 2. $\mathbf{B}(\lambda)\mathbf{x}$ spans virtually all of \mathbf{X} as we vary x
- 3. $\mathbf{B}^{-1}(\lambda)$ is continuous.

After all, these are the properties that we would have if $\mathbf{B}(\lambda)$ were simply an affine operator in the

⁴ Bollobás (1999) page 168, Hilbert (1906) pages 169–172



field of real numbers—such as $[\mathbf{B}(\lambda)](x) \triangleq [\lambda](x) = \lambda x$ which is 0 if and only if x = 0, has range $\Re(\lambda) = \mathbb{R}$, and its inverse $\lambda^{-1}x$ is continuous.

If for some λ the operator $\mathbf{B}(\lambda)$ does have all these "regular" properties, then that λ part of the resolvent set of **A** and λ is called *regular*. However if for some λ the operator **B**(λ) fails any of these conditions, then that λ part of the *spectrum* of **A**. And which conditions it fails determines which component of the spectrum it is in.

Theorem R.1. ⁵ Let $A \in \mathcal{B}(X, Y)$ be an operator.

$$\begin{array}{c} \mathbf{T} \\ \mathbf{H} \\ \mathbf{M} \end{array} \sigma(\mathbf{A}) = \sigma_{\mathsf{p}}(\mathbf{A}) \cup \sigma_{\mathsf{c}}(\mathbf{A}) \cup \sigma_{\mathsf{r}}(\mathbf{A})$$

Theorem R.2 (Spectral Theorem). 6 Let $N \in Y^X$ be an operator

Theorem R.2 (Spectral Theorem).
6
 Let $\mathbf{N} \in Y^{X}$ be an operator.

(A). $\mathbf{N}^{*}\mathbf{N} = \mathbf{N}\mathbf{N}^{*}$
(B). \mathbf{N} is COMPACT

$$\begin{cases} (1). & \mathbf{N} = \sum_{n} \lambda_{n} \mathbf{P}_{n} \\ (2). & \sum_{n} \mathbf{P}_{n} = \mathbf{I} \\ (3). & \mathbf{P}_{n} \mathbf{P}_{m} = \bar{\delta}_{n-m} \mathbf{P}_{n} \\ (4). & \dim(H_{n}) < \infty \\ (5). & \left| \left\{ \lambda_{n} \middle| \lambda_{n} \neq 0 \right\} \middle| \text{is COUNTABLY INFINITE} \right. \end{cases}$$

where
$$(\lambda_{n})_{n \in \mathbb{Z}} \triangleq \sigma_{p}(\mathbf{N}) \qquad \text{(eigenvalues of } \mathbf{N}) \\ H_{n} \triangleq \mathcal{N}(\mathbf{N} - \lambda_{n}\mathbf{I}) \qquad \text{(} \lambda_{n} \text{ is the eigenspace of } \mathbf{N} \text{ at } \lambda_{n} \text{ in } \mathbf{Y}) \\ H_{n} = \mathbf{P}_{n} \mathbf{Y} \qquad (\mathbf{P}_{n} \text{ is the projection operator that generates } \mathbf{H}_{n})$$

R.2 Fredholm kernels

Definition R.4. 7

D E F

A **Fredholm operator K** is defined as Fredholm integral equation of the first kind ⁸

Example R.2. Examples of Fredholm operators include

1. Fourier Transform
$$[\tilde{\mathbf{F}}\mathbf{x}](f) = \int_t \mathbf{x}(t)e^{-i2\pi ft} \, dt \quad \kappa(t,f) = e^{-i2\pi ft}$$

2. Inverse Fourier Transform $[\tilde{\mathbf{F}}^{-1}\tilde{\mathbf{x}}](t) = \int_f \tilde{\mathbf{x}}(f)e^{i2\pi ft} \, df \quad \kappa(f,t) = e^{i2\pi ft}$
3. Laplace operator $[\mathbf{L}\mathbf{x}](s) = \int_t \mathbf{x}(t)e^{-st} \, dt \quad \kappa(t,s) = e^{-st}$
4. autocorrelation operator $[\mathbf{R}\mathbf{x}](t) = \int_s R(t,s)\mathbf{x}(s) \, ds \quad \kappa(t,s) = R(t,s)$

Theorem R.3. Let **K** be a Fredholm operator with kernel $\kappa(t, s)$ and adjoint **K***.

$$[\mathbf{K}f](t) = \int_{A} \kappa(t, s) f(s) ds \qquad \Longleftrightarrow \qquad [\mathbf{K}^*f](t) = \int_{A} \kappa^*(s, t) f(s) ds$$

⁵ Michel and Herget (1993) page 440

⁶ ☑ Michel and Herget (1993) page 457, ② Bollobás (1999) page 200, 및 Hilbert (1906), ② Hilbert (1912), ② von Neumann (1929), a de Witt (1659)

⁷ ■ Michel and Herget (1993) page 425

⁸The equation $\int_{u} \kappa(t, s) f(s) ds$ is a **Fredholm integral equation of the first kind** and $\kappa(t, u)$ is the **kernel** of the equation. References: Fredholm (1900), Fredholm (1903) page 365, Michel and Herget (1993) page 97, Keener (1988) page 101

№ Proof:

$$\begin{aligned} & \langle \mathbf{K}f \rangle(t) = \int_{A} \kappa(t,s)f(s) \, \mathrm{d}s \\ \Leftrightarrow & \langle \mathbf{K}f \rangle(t) \, | \, \mathsf{g}(t) \rangle = \left\langle \int_{s} \kappa(t,s)f(s) \, \mathrm{d}s \, | \, \mathsf{g}(t) \right\rangle \qquad \text{by left hypothesis} \\ & = \int_{s} f(s) \, \langle \kappa(t,s) \, | \, \mathsf{g}(t) \rangle \, \, \mathrm{d}s \qquad \text{by additivity property of } \langle \triangle \, | \, \nabla \rangle \\ & = \int_{s} f(s) \, \langle \, \mathsf{g}(t) \, | \, \kappa(t,s) \rangle^{*} \, \, \mathrm{d}s \qquad \text{by conjugate symmetry property of } \langle \triangle \, | \, \nabla \rangle \\ & = \langle f(s) \, | \, \langle \, \mathsf{g}(t) \, | \, \kappa(t,s) \rangle \rangle \qquad \text{by local definition of } \langle \triangle \, | \, \nabla \rangle \\ & = \left\langle f(s) \, | \, \int_{t} \kappa^{*}(t,s) \, \mathsf{g}(t) \, \, \mathrm{d}t \right\rangle \qquad \text{by local definition of } \langle \triangle \, | \, \nabla \rangle \\ & \Leftrightarrow \left[\mathbf{K}^{*}\mathbf{g} \right](s) = \int_{A} \kappa^{*}(t,s) \, \mathsf{g}(t) \, \, \mathrm{d}t \qquad \text{by right hypothesis} \\ & \Leftrightarrow \left[\mathbf{K}^{*}\mathbf{g} \right](s) = \int_{A} \kappa^{*}(\tau,\sigma) \, \mathsf{g}(\tau) \, \, \mathrm{d}\tau \qquad \text{by change of variable: } \tau = t, \, \sigma = s \\ & \Leftrightarrow \left[\mathbf{K}^{*}\mathbf{f} \right](t) = \int_{A} \kappa^{*}(s,t) \, \mathsf{f}(s) \, \, \mathrm{d}s \qquad \text{by change of variable: } t = \sigma, \, s = \tau, \, f = \mathbf{g} \end{aligned}$$

Corollary R.1. ⁹ *Let* **K** *be an Fredholm operator with kernel* $\kappa(t, s)$ *and adjoint* **K***.

 $\begin{array}{c}
\mathbf{K} = \mathbf{K}^* \\
\mathbf{K} \text{ is self-adjoint}
\end{array}$ $\longleftrightarrow \qquad \underbrace{\kappa(t, s) = \kappa^*(s, t)}_{\text{kernel is conjugate symmetric}}$

♥Proof:

$$\mathbf{K} = \mathbf{K}^* \iff \int_A \kappa(t, s) \mathsf{f}(s) \, ds = \int_A \kappa^*(s, t) \mathsf{f}(s) \, ds \qquad \text{by Theorem R.3 page 267}$$

$$\iff \kappa(t, s) = \kappa^*(s, t)$$

Theorem R.4 (Mercer's Theorem). ¹⁰ Let **K** be an Fredholm operator with kernel $\kappa(t, s)$ and eigensystem $\{(\lambda_n, \phi_n(t))\}_{n \in \mathbb{Z}}$.

 $\left\{ \begin{array}{l} \text{(A).} \quad \int_{a}^{b} \int_{a}^{b} \kappa(t,s) f(t) f^{*}(s) \ \mathrm{d}t \geq 0 \quad and \\ \\ \text{positive} \\ \text{(B).} \quad \kappa(t,s) \ is \ \text{CONTINUOUS} \ on \\ \\ [a:b] \times [a:b] \end{array} \right\} \Longrightarrow \left\{ \begin{array}{l} \text{(1).} \quad \kappa(t,s) = \sum_{n} \lambda_{n} \phi_{n}(t) \phi_{n}^{*}(s) \quad and \\ \\ \text{(2).} \quad \kappa(t,s) \ \text{CONVERGES ABSOLUTELY} \\ \\ and \ \text{UNIFORMLY} \ on \\ \\ [a:b] \times [a:b] \end{array} \right\}$

¹⁰ Gohberg et al. (2003) page 198, Courant and Hilbert (1930) pages 138−140, Mercer (1909) page 439



⁹ Michel and Herget (1993) page 430

Back Matter




It appears to me that if one wants to make progress in mathematics, one should study the masters and not the pupils.


Niels Henrik Abel (1802–1829), Norwegian mathematician ¹¹



When evening comes, I return home and go to my study. On the threshold I strip naked, taking off my muddy, sweaty workaday clothes, and put on the robes of court and palace, and in this graver dress I enter the courts of the ancients and am welcomed by them, and there I taste the food that alone is mine, and for which I was born. And there I make bold to speak to them and ask the motives of their actions, and they, in their humanity reply to me. And for the space of four hours I forget the world, remember no vexation, fear poverty no more, tremble no more at death; I pass indeed into their world.

Niccolò Machiavelli (1469–1527), Italian political philosopher, in a 1513 letter to friend Francesco Vettori. 12







The Book Worm by Carl Spitzweg, circa 1850

13

¹¹ quote: **Simmons** (2007) page 187.

¹² quote: Machiavelli (1961) page 139?.

http://en.wikipedia.org/wiki/File:Ancientlibraryalex.jpg, public domain http://en.wikipedia.org/wiki/File:Carl_Spitzweg_021.jpg,



★ To sit alone in the lamplight with a book spread out before you, and hold intimate converse with men of unseen generations—such is a pleasure beyond compare.

 Yoshida Kenko (Urabe Kaneyoshi) (1283? – 1350?), Japanese author and Buddhist monk

quote: Kenko (circa 1330)

image: http://en.wikipedia.org/wiki/Yoshida_Kenko



I I	
RIRI IOCRADHV	

- Yuri A. Abramovich and Charalambos D. Aliprantis. *An Invitation to Operator Theory*. American Mathematical Society, Providence, Rhode Island, 2002. ISBN 0-8218-2146-6. URLhttp://books.google.com/books?vid=ISBN0821821466.
- Milton Abramowitz and Irene A. Stegun, editors. *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables.* National Bureau of Standards, 1972. URL http://www.cs.bham.ac.uk/~aps/research/projects/as/book.php.
- Edward H. Adelson and Peter J. Burt. Image data compression with the laplacian pyramid. In *Proceedings of the Pattern Recognition and Information Processing Conference*, pages 218–223, Dallas Texas, 1981. IEEE Computer Society Press. URL citeseerx.ist.psu.edu/viewdoc/summary?doi=10.1.1.50.791.
- N. I. Akhiezer and I. M. Glazman. *Theory of Linear Operators in Hilbert Spaces*, volume 1. Dover, New York, 1993. URL http://books.google.com/books?vid=ISBN0486677486. Translated from the original Russian text *Teoriia lineinykh operatorov v Gil'bertovom prostranstve*.
- Donald J. Albers and Freeman Dyson. Freeman dyson: Mathematician, physicist, and writer. *The College Mathematics Journal*, 25(1):3–21, January 1994. doi: 10.2307/2687079. URL http://www.jstor.org/stable/2687079.
- Charalambos D. Aliprantis and Owen Burkinshaw. *Principles of Real Analysis*. Acedemic Press, London, 3 edition, 1998. ISBN 9780120502578. URL http://www.amazon.com/dp/0120502577.
- Luis Alvarez, Frédéric Guichard, Pierre-Louis Lions, and Jean Michel Morel. Axioms and fundamental equations of image processing. *Archive for Rational Mechanics and Analysis*, 123(3):199–257, 1993. URL http://link.springer.com/article/10.1007/BF00375127.
- Ichiro Amemiya and Huzihiro Araki. A remark on piron's paper. *Publications of the Research Institute for Mathematical Sciences, Kyoto University*, 2(3):423–427, 1966. ISSN 0034-5318. doi: 10.2977/prims/1195195769. URL http://projecteuclid.org/euclid.prims/1195195769.
- George E. Andrews, Richard Askey, and Ranjan Roy. *Special Functions*, volume 71 of *Encyclopedia of mathematics and its applications*. Cambridge University Press, Cambridge, U.K., new edition, February 15 2001. ISBN 0521789885. URL http://books.google.com/books?vid=ISBN0521789885.

Kendall E. Atkinson and Weimin Han. *Theoretical Numerical Analysis: A Functional Analysis Framework*, volume 39 of *Texts in Applied Mathematics*. Springer, 3 edition, 2009. ISBN 9781441904584. URL http://books.google.com/books?vid=ISBN1441904581.

- Léon Autonne. Sur l'hermitien (on the hermitian). In *Comptes Rendus Des SéAnces De L'AcadéMie Des Sciences*, volume 133, pages 209–268. De L'Académie des sciences (Academy of Sciences), Paris, 1901. URL http://visualiseur.bnf.fr/Visualiseur?0=NUMM-3089. Comptes Rendus Des SéAnces De L'AcadéMie Des Sciences (Reports Of the Meetings Of the Academy of Science).
- Léon Autonne. Sur l'hermitien (on the hermitian). *Rendiconti del Circolo Matematico di Palermo*, 16:104–128, 1902. Rendiconti del Circolo Matematico di Palermo (Statements of the Mathematical Circle of Palermo).
- George Bachman. *Elements of Abstract Harmonic Analysis*. Academic paperbacks. Academic Press, New York, 1964. ISBN 9781483267562. URL http://books.google.com/books?vid=ISBN9781483267562.
- George Bachman and Lawrence Narici. *Functional Analysis*. Academic Press textbooks in mathematics; Pure and Applied Mathematics Series. Academic Press, 1 edition, 1966. ISBN 9780486402512. URL http://books.google.com/books?vid=ISBN0486402517. "unabridged republication" available from Dover (isbn 0486402517).
- George Bachman, Lawrence Narici, and Edward Beckenstein. *Fourier and Wavelet Analysis*. Universitext Series. Springer, 2002. ISBN 9780387988993. URL http://books.google.com/books?vid=ISBN9780387988993. "Corrected second printing, 2002".
- Stefan Banach. Sur les opérations dans les ensembles abstraits et leur applications aux équations intégrales (on abstract operations and their applications to the integral equations). *Fundamenta Mathematicae*, 3:133–181, 1922. URL http://matwbn.icm.edu.pl/ksiazki/fm/fm3/fm3120.pdf.
- Stefan Banach. *Théorie des opérations linéaires*. Monografje Matematyczne, Warsaw, Poland, 1932a. URL http://matwbn.icm.edu.pl/kstresc.php?tom=1&wyd=10. (Theory of linear operations).
- Stefan Banach. *Theory of Linear Operations*, volume 38 of *North-Holland mathematical library*. North-Holland, Amsterdam, 1932b. ISBN 0444701842. URL http://www.amazon.com/dp/0444701842/. English translation of 1932 French edition, published in 1987.
- Adi Ben-Israel and Robert P. Gilbert. *Computer-supported calculus*. Texts and monographs in symbolic computation. Springer, 2002. ISBN 3-211-82924-5. URL http://books.google.com/books?vid=ISBN3211829245.
- Ladislav Beran. Three identities for ortholattices. *Notre Dame Journal of Formal Logic*, 17(2):251–252, 1976. doi: 10.1305/ndjfl/1093887530. URL http://projecteuclid.org/euclid.ndjfl/1093887530.
- Ladislav Beran. *Orthomodular Lattices: Algebraic Approach*. Mathematics and Its Applications (East European Series). D. Reidel Publishing Company, Dordrecht, 1985. ISBN 90-277-1715-X. URL http://books.google.com/books?vid=ISBN902771715X.
- Sterling Khazag Berberian. *Introduction to Hilbert Space*. Oxford University Press, New York, 1961. URL http://books.google.com/books?vid=ISBN0821819127.
- Earl Berkson. Some metrics on the subspaces of a banach space. *Pacific Journal of Mathematics*, 13(1):7–22, 1963. URL http://projecteuclid.org/euclid.pjm/1103035953.



M. Bertero and P. Boccacci. *Introduction to Inverse Problems in Imaging*. CRC Press, 1998. ISBN 9781439822067. URL http://books.google.com/books?vid=ISBN9781439822067.

- Josef Betten. *Creep Mechanics*, chapter A. The Heaviside and Dirac Functions, pages 285–294. Springer Science & Business Media, 3 edition, 2008a. ISBN 9783540850519. URL https://link.springer.com/content/pdf/bbm:978-3-540-85051-9/1.pdf. https://www.springer.com/gp/book/9783540850502.
- Josef Betten. Creep Mechanics, chapter B. The LAPLACE Transformation, pages 295–343. Springer Science & Business Media, 3 edition, 2008b. ISBN 9783540850519. URL https://link.springer.com/content/pdf/bbm:978-3-540-85051-9/1.pdf. https://www.springer.com/gp/book/9783540850502.
- Garrett Birkhoff and John Von Neumann. The logic of quantum mechanics. *The Annals of Mathematics*, 37(4):823–843, October 1936. URL http://www.jstor.org/stable/1968621.
- Duncan Black, Ian Brooks, and et al. Robert Groves, editors. *Collins English Dictionary—Complete and Unabridged*. HarperCollins Publishers, 10 edition, 2009. ISBN 978-0-00-732119-3. URL http://books.google.com/books?vid=ISBN9780007321193.
- Béla Bollobás. *Linear Analysis; an introductory course*. Cambridge mathematical textbooks. Cambridge University Press, Cambridge, 2 edition, March 1 1999. ISBN 978-0521655774. URL http://books.google.com/books?vid=ISBN0521655773.
- Umberto Bottazzini. *The Higher Calculus: A History of Real and Complex Analysis from Euler to Weierstrass*. Springer-Verlag, New York, 1986. ISBN 0-387-96302-2. URL http://books.google.com/books?vid=ISBN0387963022.
- Carl Benjamin Boyer and Uta C. Merzbach. *A History of Mathematics*. Wiley, New York, 2 edition, 1991. ISBN 0471543977. URL http://books.google.com/books?vid=ISBN0471543977.
- Ronald Newbold Bracewell. *The Fourier transform and its applications*. McGraw-Hill electrical and electronic engineering series. McGraw-Hill, 2, illustrated, international student edition edition, 1978. ISBN 9780070070134. URL http://books.google.com/books?vid=ISBN007007013X.
- Thomas John I'Anson Bromwich. *An Introduction to the Theory of Infinite Series*. Macmillan and Company, 1 edition, 1908. ISBN 9780821839768. URL http://www.archive.org/details/anintroductiont00bromgoog.
- Andrew M. Bruckner, Judith B. Bruckner, and Brian S. Thomson. *Real Analysis*. Prentice-Hall, Upper Saddle River, N.J., 1997. ISBN 9780134588865. URL http://books.google.com/books?vid=ISBN013458886X.
- Dmitri Burago, Yuri Burago, and Sergei Ivanov. *A Course in Metric Geometry*, volume 33 of *Graduate Studies in Mathematics*. American Mathematical Society, 2001. ISBN 978-0821821299. URL http://books.google.com/books?vid=ISBN0821821296.
- C. Sidney Burrus, Ramesh A. Gopinath, and Haitao Guo. *Introduction to Wavelets and Wavelet Transforms; A Primer*. Prentice Hall, Upper Saddle River, New Jersey, USA, 1998. ISBN 0-13-489600-9. URL http://www.dsp.rice.edu/software/wavebook.shtml.
- Peter J. Burt and Edward H. Adelson. The laplacian pyramid as a compact image code. *IEEE Transactions On Communications*, COM-3L(4):532-540, April 1983. URL http://citeseer.ist.psu.edu/burt831aplacian.html.





Charles L. Byrne. *Signal Processing: A Mathematical Approach*. AK Peters Series. A K Peters, 2005. ISBN 1568812426.

- Florian Cajori. A history of mathematical notations; notations mainly in higher mathematics. In *A History of Mathematical Notations; Two Volumes Bound as One*, volume 2. Dover, Mineola, New York, USA, 1993. ISBN 0-486-67766-4. URL http://books.google.com/books?vid=ISBN0486677664. reprint of 1929 edition by *The Open Court Publishing Company*.
- Gerolamo Cardano. *Ars Magna or the Rules of Algebra*. Dover Publications, Mineola, New York, 1545. ISBN 0486458733. URL http://www.amazon.com/dp/0486458733. English translation of the Latin *Ars Magna* edition, published in 2007.
- Lennart Axel Edvard Carleson. Convergence and growth of partial sums of fourier series. *Acta Mathematica*, 116(1):135–157, 1966. doi: 10.1007/BF02392815. URL http://dx.doi.org/10.1007/BF02392815.
- Lennart Axel Edvard Carleson and Björn Engquist. After the 'golden age': what next? lennart carleson interviewed by björn engquist. In Björn Engquist and Wilfried Schmid, editors, *Mathematics Unlimited—2001 and beyond*, pages 455–461. Springer, Berlin, 2001. ISBN 978-3-540-66913-5. URL https://link.springer.com/chapter/10.1007/978-3-642-56478-9_22.
- N. L. Carothers. A Short Course on Banach Space Theory. Number 64 in London Mathematical Society Student Texts. Cambridge University Press, Cambridge, 2005. ISBN 0521842832. URL http://www.comm.musashi-tech.ac.jp/~arimoto/Drarimoto/preprints/calothers/Assemble.ps.
- Peter G. Casazza and Mark C. Lammers. Bracket products for weyl-heisenberg frames. In Hans G. Feichtinger and Thomas Strohmer, editors, *Gabor Analysis and Algorithms: Theory and Applications*, Applied and Numerical Harmonic Analysis, pages 71–98. Birkhäuser, 1998. ISBN 9780817639594.
- Clémente Ibarra Castanedo. *Quantitative subsurface defect evaluation by pulsed phase thermography: depth retrieval with the phase.* PhD thesis, Université Laval, October 2005. URL http://archimede.bibl.ulaval.ca/archimede/fichiers/23016/23016.html. Faculte' Des Sciences Et De Génie.
- Arthur Cayley. A memoir on the theory of matrices. *Philosophical Transactions of the Royal Society of London*, 148:17–37, 1858. ISSN 1364-503X. URL http://www.jstor.org/view/02610523/ap000059/00a00020/0.
- Joan Cerdà. *Linear functional analysis*, volume 116 of *Graduate studies in mathematics*. American Mathematical Society, July 16 2010. ISBN 0821851152. URL http://books.google.com/books?vid=ISBN0821851152.
- Alexandre J. Chorin and Ole H. Hald. *Stochastic Tools in Mathematics and Science*, volume 1 of *Surveys and Tutorials in the Applied Mathematical Sciences*. Springer, New York, 2 edition, 2009. ISBN 978-1-4419-1001-1. URL http://books.google.com/books?vid=ISBN9781441910011.
- Ole Christensen. *An Introduction to Frames and Riesz Bases*. Applied and Numerical Harmonic Analysis. Birkhäuser, Boston/Basel/Berlin, 2003. ISBN 0-8176-4295-1. URL http://books.google.com/books?vid=ISBN0817642951.
- Ole Christensen. *Frames and bases: An Introductory Course.* Applied and Numerical Harmonic Analysis. Birkhäuser, Boston/Basel/Berlin, 2008. ISBN 9780817646776. URL http://books.google.com/books?vid=ISBN0817646779.



Charles K. Chui. *An Introduction to Wavelets*. Academic Press, San Diego, California, USA, January 3 1992. ISBN 9780121745844. URL http://books.google.com/books?vid=ISBN0121745848.

- Jon F. Claerbout. Fundamentals of Geophysical Data Processing with Applications to Petroleum Prospecting. International series in the earth and planetary sciences, Tab Mastering Electronics Series. McGraw-Hill, New York, 1976. ISBN 9780070111172. URL http://sep.stanford.edu/sep/prof/.
- Paul M. Cohn. *Basic Algebra; Groups, Rings and Fields*. Springer, December 6 2002. ISBN 1852335874. URL http://books.google.com/books?vid=isbn1852335874.
- R. Courant and D. Hilbert. *Methods of Mathematical Physics*, volume 1. Interscience Publishers, New York, 1930. URL http://www.worldcat.org/isbn/0471504475.
- Xingde Dai and David R. Larson. *Wandering vectors for unitary systems and orthogonal wavelets*. Number 640 in Memoirs of the American Mathematical Society. American Mathematical Society, Providence R.I., July 1998. ISBN 0821808001. URL http://books.google.com/books?vid=ISBN0821808001.
- Xingde Dai and Shijie Lu. Wavelets in subspaces. *Michigan Math. J.*, 43(1):81–98, 1996. doi: 10. 1307/mmj/1029005391. URL http://projecteuclid.org/euclid.mmj/1029005391.
- Ingrid Daubechies. *Ten Lectures on Wavelets*. Society for Industrial and Applied Mathematics, Philadelphia, 1992. ISBN 0-89871-274-2. URL http://www.amazon.com/dp/0898712742.
- Ingrid Daubechies, A. Grossman, and Y. Meyer. Painless nonorthogonal expansions. *Journal of Mathematical Physics*, 27(5):1271–1283, May 1986. ISSN 0022-2488. URL link.aip.org/link/?jmp/27/1271.
- Kenneth R. Davidson and Allan P. Donsig. *Real Analysis and Applications*. Springer, 2010. ISBN 9781441900050. URL http://books.google.com/books?vid=ISBN1441900055.
- Charles Jean de la Vallée-Poussin. Sur l'intégrale de lebesgue. *Transactions of the American Mathematical Society*, 16(4):435–501, October 1915. URL http://www.jstor.org/stable/1988879.
- Juan-Arias de Reyna. *Pointwise Convergence of Fourier Series*, volume 1785 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin/Heidelberg/New York, 2002. ISBN 3540432701. URL http://books.google.com/books?vid=ISBN3540432701.
- Johan de Witt. Elementa curvarum linearum (elements of curves). In *La Géométrie*. 1659. Elementa curvarum linearum was an appendix to de Witt's translation of Descartes' *La Géométrie*.
- René Descartes. *La géométrie*. 1637a. URLhttp://historical.library.cornell.edu/math/math_D.html.
- René Descartes. *Discours de la méthode pour bien conduire sa raison, et chercher la verite' dans les sciences*. Jan Maire, Leiden, 1637b. URL http://www.gutenberg.org/etext/13846.
- René Descartes. Discourse on the Method of Rightly Conducting the Reason in the Search for Truth in the Sciences. 1637c. URL http://www.gutenberg.org/etext/59.
- René Descartes. Regulae ad directionem ingenii. 1684a. URL http://www.fh-augsburg.de/~harsch/Chronologia/Lspost17/Descartes/des_re00.html.
- René Descartes. Rules for Direction of the Mind. 1684b. URL http://en.wikisource.org/wiki/Rules_for_the_Direction_of_the_Mind.





René Descartes. *The Geometry of Rene Descartes*. Courier Dover Publications, June 1 1954. ISBN 0486600688. URL http://books.google.com/books?vid=isbn0486600688. orginally published by Open Court Publishing, Chicago, 1925; translation of La géométrie.

- Elena Deza and Michel-Marie Deza. *Dictionary of Distances*. Elsevier Science, Amsterdam, 2006. ISBN 0444520872. URL http://books.google.com/books?vid=ISBN0444520872.
- Jean Alexandre Dieudonné. *Foundations of Modern Analysis*. Academic Press, New York, 1969. ISBN 1406727911. URL http://books.google.com/books?vid=ISBN1406727911.
- R. J. Duffin and A. C. Schaeffer. A class of nonharmonic fourier series. *Transactions of the American Mathematical Society*, 72(2):341–366, March 1952. ISSN 1088-6850. URL http://www.jstor.org/stable/1990760.
- Bogdan Dumitrescu. *Positive Trigonometric Polynomials and Signal Processing Applications*. Signals and Communication Technology. Springer, 2007. ISBN 978-1-4020-5124-1. URL gen.lib.rus.ec/get?md5=5346e169091b2d928d8333cd053300f9.
- Nelson Dunford and Jacob T. Schwartz. *Linear operators. Part 1, General Theory*, volume 7 of *Pure and applied mathematics*. Interscience Publishers, New York, 1957. ISBN 0471226394. URL http://www.amazon.com/dp/0471608483. with the assistance of William G. Bade and Robert G. Bartle.
- John R. Durbin. *Modern Algebra; An Introduction*. John Wiley & Sons, Inc., 4 edition, 2000. ISBN 0-471-32147-8. URL http://www.worldcat.org/isbn/0471321478.
- Robert E. Edwards. *Functional Analysis: Theory and Applications*. Dover books on mathematics. Dover, New York, 1995. ISBN 0-486-68143-2. URL http://books.google.com/books?vid=ISBN0486681432.
- Yuli Eidelman, Vitali D. Milman, and Antonis Tsolomitis. *Functional Analysis: An Introduction*, volume 66 of *Graduate Studies in Mathematics*. American Mathematical Society, 2004. ISBN 0821836463. URL http://books.google.com/books?vid=ISBN0821836463.
- Per Enflo. A counterexample to the approximation problem in banach spaces. *Acta Mathematica*, 130:309–317, 1973. URL http://link.springer.com/content/pdf/10.1007/BF02392270.
- Leonhard Euler. *Introductio in analysin infinitorum*, volume 1. Marcum-Michaelem Bousquet & Socios, Lausannæ, 1748. URL http://www.math.dartmouth.edu/~euler/pages/E101.html. Introduction to the Analysis of the Infinite.
- Leonhard Euler. *Introduction to the Analysis of the Infinite*. Springer, 1988. ISBN 0387968245. URL http://books.google.com/books?vid=ISBN0387968245. translation of 1748 Introductio in analysin infinitorum.
- David Ewen. *The Book of Modern Composers*. Alfred A. Knopf, New York, 1950. URLhttp://books.google.com/books?id=yHw4AAAAIAAJ.
- David Ewen. *The New Book of Modern Composers*. Alfred A. Knopf, New York, 3 edition, 1961. URL http://books.google.com/books?id=bZIaAAAAMAAJ.
- Marián Fabian, Petr Habala, Petr Hájek, Vicente Montesinos, Vaclav Zizler, and Václav Zizler. *Banach Space Theory: The Basis for Linear and Nonlinear Analysis.* CMS Books in Mathematics. Springer, 2010. ISBN 1441975144. URL http://books.google.com/books?vid=isbn1441975144.



Lorenzo Farina and Sergio Rinaldi. Positive Linear Systems: Theory and Applications. Pure and applied mathematics. John Wiley & Sons, 1 edition, July 3 2000. ISBN 9780471384564. URL http: //books.google.com/books?vid=ISBN0471384569.

- G.L. Fix and G. Strang. Fourier analysis of the finite element method in ritz-galerkin theory. *Studies* in Applied Mathematics, 48:265–273, 1969.
- Harley Flanders. Differentiation under the integral sign. The American Mathematical Monthly, 80(6):615-627, June-July 1973. URL http://sgpwe.izt.uam.mx/files/users/uami/ jdf/proyectos/Derivar_inetegral.pdf. http://www.jstor.org/pss/2319163.
- Francis J. Flanigan. Complex Variables; Harmonic and Analytic Functions. Dover, New York, 1983. ISBN 9780486613888. URL http://books.google.com/books?vid=ISBN0486613887.
- Gerald B. Folland. Fourier Analysis and its Applications. Wadsworth & Brooks / Cole Advanced Books & Software, Pacific Grove, California, USA, 1992. ISBN 0-534-17094-3. URL http://www. worldcat.org/isbn/0534170943.
- Gerald B. Folland. A Course in Abstract Harmonic Analysis. Studies in Advanced Mathematics. CRC Press, Boca Raton, 1995. ISBN 0-8493-8490-7. URL http://books.google.com/books?vid= ISBN0849384907.
- Brigitte Forster and Peter Massopust, editors. Four Short Courses on Harmonic Analysis: Wavelets, Frames, Time-Frequency Methods, and Applications to Signal and Image Analysis. Applied and Numerical Harmonic Analysis. Springer, November 19 2009. ISBN 9780817648909. URL http: //books.google.com/books?vid=ISBN0817648909.
- Jean-Baptiste-Joseph Fourier. Mémoire sur la propagation de la chaleur dans les corps solides (dissertation on the propagation of heat in solid bodies). In M. Gaston Darboux, editor, Œuvres De Fourier, pages 215-221. Ministère de L'instruction Publique, Paris, France, 2 edition, December 21 1807. URL http://gallica.bnf.fr/ark:/12148/bpt6k33707/f220n7.
- Jean-Baptiste-Joseph Fourier. Théorie Analytique de la Chaleur (The Analytical Theory of Heat). Chez Firmin Didot, pere et fils, Paris, 1822. URL http://books.google.com/books?vid= 04X2vlqZx7hydlQUWEq&id=TDQJAAAAIAAJ.
- Jean-Baptiste-Joseph Fourier. The Analytical Theory of Heat (Théorie Analytique de la Chaleur). Cambridge University Press, Cambridge, February 20 1878. URL http://www.archive.org/ details/analyticaltheory00fourrich. 1878 English translation of the original 1822 French edition. A 2003 Dover edition is also available: isbn 0486495310.
- Maurice René Fréchet. Sur quelques points du calcul fonctionnel (on some points of functional calculation). Rendiconti del Circolo Matematico di Palermo, 22:1-74, April 22 1906. URL https: //www.lpsm.paris/pageperso/mazliak/Frechet 1906.pdf. Rendicontidel Circolo Matematico di Palermo (Statements of the Mathematical Circle of Palermo).
- Maurice René Fréchet. Les Espaces abstraits et leur théorie considérée comme introduction a l'analyse générale. Borel series. Gauthier-Villars, Paris, 1928. URL http://books.google.com/ books?id=9czoHQAACAAJ. Abstract spaces and their theory regarded as an introduction to general analysis.
- Erik Ivar Fredholm. Sur une nouvelle méthode pour la résolution du problème de dirichlet (on a new method for the resolution of the problem of dirichlet). /:Oefversigt af Kongl. Sv. Vetenskaps-Academiens Förhandlingar, 57:39–66, 1900.





Erik Ivar Fredholm. Sur une classe d'equations fonctionnelles (on a class of functional equations). *Acta Mathematica*, 27(1):365–390, December 1903. ISSN 0001-5962. doi: 10.1007/BF02421317. URL http://www.springerlink.com/content/c41371137837p252/.

- Ferdinand Georg Frobenius. Uber lineare substitutionen und bilineare formen. *Journal für die reine und angewandte Mathematik (Crelle's Journal)*, 84:1–63, 1878. ISSN 0075-4102. URL http://www.digizeitschriften.de/home/services/pdfterms/?ID=509796.
- Ferdinand Georg Frobenius. Uber lineare substitutionen und bilineare formen. In Jean Pierre Serre, editor, *Gesammelte Abhandlungen (Collected Papers)*, volume I, pages 343–405. Springer, Berlin, 1968. URL http://www.worldcat.org/oclc/253015. reprint of Frobenius' 1878 paper.
- Jürgen Fuchs. Affine Lie Algebras and Quantum Groups: An Introduction, With Applications in Conformal Field Theory. Cambridge Monographs on Mathematical Physics. Cambridge University Press, 1995. ISBN 052148412X. URL http://books.google.com/books?vid=ISBN052148412X.
- Dennis Gabor. Theory of communication. *Journal of the Institution of Electrical Engineers*, 93(26): 429–457, November 1946. URL http://bigwww.epfl.ch/chaudhury/gabor.pdf.
- Carl Friedrich Gauss. *Carl Friedrich Gauss Werke*, volume 8. Königlichen Gesellschaft der Wissenschaften, B.G. Teubneur In Leipzig, Göttingen, 1900. URL http://gdz.sub.uni-goettingen.de/dms/load/img/?PPN=PPN236010751.
- Israel M. Gelfand. Normierte ringe. *Mat. Sbornik*, 9(51)(1):3-24, 1941. URL http://www.mathnet.ru/links/d19894c5545d2f75b3e1037162f86d58/sm6046.pdf.
- Israel M. Gelfand and Mark A. Naimark. Normed rings with an involution and their representations. In *Commutative Normed Rings*, number 170 in AMS Chelsea Publishing Series, pages 240–274. Chelsea Publishing Company, Bronx, 1964. ISBN 9780821820223. URL http://books.google.com/books?vid=ISBN0821820222.
- Israel M. Gelfand and Mark A. Neumark. On the imbedding of normed rings into the ring of operators in hilbert space. *Mat. Sbornik*, 12(54:2):197–217, 1943a.
- Israel M. Gelfand and Mark A. Neumark. On the imbedding of normed rings into the ring of operators in hilbert space. In Robert S. Doran, editor, *C*-algebras: 1943–1993: a Fifty Year Celebration: Ams Special Session Commenorating the First Fifty Years of C*-Algebra Theory January 13–14, 1993*, pages 3–19. American Mathematical Society, 1943b. ISBN 978-0821851753. URL http://books.google.com/books?vid=ISBN0821851756.
- Israel M. Gelfand, R. A. Minlos, and Z. .Ya. Shapiro. *Representations of the rotation and Lorentz groups and their applications*. Courier Dover Publications, reprint edition, 2018. ISBN 9780486823850. Izrail' Moiseevich Gel'fand.
- Izrail' Moiseevich Gel'fand. *Representations of the rotation and Lorentz groups and their applications.* Pergamon Press book, 1963. 2018 Dover edition available.
- John Robilliard Giles. *Introduction to the Analysis of Normed Linear Spaces*. Number 13 in Australian Mathematical Society lecture series. Cambridge University Press, Cambridge, 2000. ISBN 0-521-65375-4. URL http://books.google.com/books?vid=ISBN0521653754.
- Israel Gohberg, Seymour Goldberg, and Marinus A. Kaashoek. *Basic Classes of Linear Operators*. Birkhäuser, Basel, 1 edition, 2003. ISBN 3764369302. URL http://books.google.com/books?vid=ISBN3764369302.



T. N. T. Goodman, S. L. Lee, and W. S. Tang. Wavelets in wandering subspaces. *Transactions of the A.M.S.*, 338(2):639–654, August 1993a. URL http://www.jstor.org/stable/2154421. Transactions of the American Mathematical Society.

- T. N. T. Goodman, S. L. Lee, and W. S. Tang. Wavelets in wandering subspaces. *Advances in Computational Mathematics 1*, pages 109–126, February 1993b.
- Jaideva C. Goswami and Andrew K. Chan. Fundamentals of Wavelets; Theory, Algorithms, and Applications. John Wiley & Sons, Inc., 1999. ISBN 0-471-19748-3. URL http://vadkudr.boom.ru/Collection/fundwave_contents.html.
- Daniel J. Greenhoe. Wavelet Structure and Design, volume 3 of Mathematical Structure and Design series. Abstract Space Publishing, August 2013. ISBN 9780983801139. URL http://books.google.com/books?vid=ISBN0983801134. revised online version available at https://www.researchgate.net/publication/312529555.
- Stanley Gudder. *Stochastic Methods in Quantum Mechanics*. North Holland, 1979. ISBN 0444002995. URL http://books.google.com/books?vid=ISBN0444002995.
- Stanley Gudder. *Stochastic Methods in Quantum Mechanics*. Dover, Mineola NY, 2005. ISBN 0486445321. URL http://books.google.com/books?vid=ISBN0486445321.
- Frédéric Guichard, Jean-Michel Morel, and Robert Ryan. *Contrast invariant image analysis and PDE's.* 2012. URL http://dev.ipol.im/~morel/JMMBook2012.pdf.
- Frédéric Guichard, Jean-Michel Morel, and Robert Ryan. *Contrast invariant image analysis and PDE's.* 20xx. URL http://dev.ipol.im/~morel/JMMBookOct04_complete.pdf.
- Ernst Adolph Guillemin. *Synthesis of Passive Networks: Theory and Methods Appropriate to the Realization and Approximation Problems*. John Wiley & Sons, 1957. ISBN 9780882754819. URL http://books.google.com.tw/books?id=JQ4nAAAAMAAJ.
- Amritava Gupta. *Real & Abstract Analysis*. Academic Publishers Calcutta, 1998. ISBN 9788186358443. URL http://books.google.com/books?vid=ISBN8186358447. first published in 1998, reprinted in 2000 and 2006.
- Alfréd Haar. Zur theorie der orthogonalen funktionensysteme. *Mathematische Annalen*, 69:331–371, September 1910. ISSN 1432-1807. doi: 10.1007/BF01456326.
- Norman B. Haaser and Joseph A. Sullivan. *Real Analysis*. Dover Publications, New York, 1991. ISBN 0-486-66509-7. URL http://books.google.com/books?vid=ISBN0486665097.
- Paul R. Haddad and Ali N. Akansu. *Multiresolution Signal Decomposition: Transforms, Subbands, and Wavelets.* Acedemic Press, October 1 1992. ISBN 0323138365. URL http://books.google.com/books?vid=ISBN0323138365.
- Paul R. Halmos. *Finite Dimensional Vector Spaces*. Princeton University Press, Princeton, 1 edition, 1948. ISBN 0691090955. URL http://books.google.com/books?vid=isbn0691090955.
- Paul R. Halmos. *Finite Dimensional Vector Spaces*. Springer Science & Business Media, New York, 1 edition, 1958. ISBN 9781461263876. URL http://books.google.com/books?vid=isbn9781461263876. 2012 "Softcover reprint of the hardcover 1st edition 1987"; *Undergraduate Texts in Mathematics*.





Paul R. Halmos. *Intoduction to Hilbert Space and the Theory of Spectral Multiplicity*. Chelsea Publishing Company, New York, 2 edition, 1998a. ISBN 0821813781. URL http://books.google.com/books?vid=ISBN0821813781.

- Paul R. Halmos. *Introduction to Hilbert Space and the Theory of Spectral Multiplicity*. AMS Chelsea, Providence RI, 2 edition, 1998b. ISBN 0821813781. URL http://books.google.com/books?vid=ISBN 0821813781.
- Georg Hamel. Eine basis aller zahlen und die unstetigen lösungen der funktionalgleichung f(x + y) = f(x) + f(y). Mathematische Annalen, 60(3):459–462, 1905. URL http://gdz.sub.uni-goettingen.de/dms/load/img/?PPN=GDZPPN002260395&IDDOC=28580.
- Deguang Han, Keri Kornelson, David Larson, and Eric Weber. *Frames for Undergraduates*, volume 40 of *Student Mathematical Library*. American Mathematical Society, 2007. ISBN 0821842129. URL http://books.google.com/books?vid=ISBN0821842129. Deguang Han = ???
- Godfrey H. Hardy. *A Mathematician's Apology*. Cambridge University Press, Cambridge, 1940. URL http://www.math.ualberta.ca/~mss/misc/A%20Mathematician's%20Apology.pdf.
- Felix Hausdorff. *Set Theory*. Chelsea Publishing Company, New York, 3 edition, 1937. ISBN 0828401195. URL http://books.google.com/books?vid=ISBN0828401195. 1957 translation of the 1937 German *Grundzüge der Mengenlehre*.
- Michiel Hazewinkel, editor. *Handbook of Algebras*, volume 2. North-Holland, Amsterdam, 1 edition, 2000. ISBN 044450396X. URL http://books.google.com/books?vid=ISBN044450396X.
- Jean Van Heijenoort. From Frege to Gödel: A Source Book in Mathematical Logic, 1879-1931. Harvard University Press, Cambridge, Massachusetts, 1967. URL http://www.hup.harvard.edu/catalog/VANFGX.html.
- Christopher Heil. *A Basis Theory Primer*. Applied and Numerical Harmonic Analysis. Birkhäuser, Boston, expanded edition edition, 2011. ISBN 9780817646868. URL http://books.google.com/books?vid=ISBN9780817646868.
- Christopher E. Heil and David F. Walnut. Continuous and discrete wavelet transforms. *Society for Industrial and Applied Mathematics*, 31(4), December 1989. URL http://citeseer.ist.psu.edu/viewdoc/download?doi=10.1.1.132.1241&rep=rep1&type=pdf.
- Charles Hermite. Lettre à stieltjes. In B. Baillaud and H. Bourget, editors, *Correspondance d'Hermite et de Stieltjes*, volume 2, pages 317–319. Gauthier-Villars, Paris, May 20 1893. published in 1905.
- Eugenio Hernández and Guido Weiss. *A First Course on Wavelets*. CRC Press, New York, 1996. ISBN 0849382742. URL http://books.google.com/books?vid=ISBN0849382742.
- John Rowland Higgins. Sampling Theory in Fourier and Signal Analysis: Foundations. Oxford Science Publications. Oxford University Press, August 1 1996. ISBN 9780198596998. URL http://books.google.com/books?vid=ISBN0198596995.
- David Hilbert. Grundzüge einer allgemeinen theorie der linearen integralgleichungen (fundamentals of a general theory of the linear integral equations). *Mathematisch-Physikalische Klasse (Mathematical-Physical Class)*, pages 49–91, March 1904. URL http://dz-srv1.sub.uni-goettingen.de/sub/digbib/loader?did=D57552. Report 1 of 6.



David Hilbert. Grundzüge einer allgemeinen theorie der linearen integralgleichungen (fundamentals of a general theory of the linear integral equations). *Mathematisch-Physikalische Klasse (Mathematical-Physical Class)*, pages 157–228, March 1906. URL http://dz-srv1.sub.uni-goettingen.de/sub/digbib/loader?did=D58133. Report 4 of 6.

- David Hilbert. *Grundzüge einer allgemeinen Theorie der linearen Integralgleichungen (Fundamentals of a general theory of the linear integral equations)*, volume 26 of *Fortschritte der mathematischen Wissenschaften in Monographien (Progress of the mathematical sciences in Monographs)*. B.G. Teubner, Leipzig und Berlin, 1912. URL http://www.worldcat.org/oclc/13468199.
- David Hilbert, Lothar Nordheim, and John von Neumann. über die grundlagen der quantenmechanik (on the bases of quantum mechanics). *Mathematische Annalen*, 98:1–30, 1927. ISSN 0025-5831 (print) 1432-1807 (online). URL http://dz-srv1.sub.uni-goettingen.de/cache/toc/D27776.html.
- Samuel S. Holland, Jr. The current interest in orthomodular lattices. In James C. Abbott, editor, *Trends in Lattice Theory*, pages 41–126. Van Nostrand-Reinhold, New York, 1970. URL http://books.google.com/books?id=ZfA-AAAAIAAJ. from Preface: "The present volume contains written versions of four talks on lattice theory delivered to a symposium on Trends in Lattice Theory held at the United States Naval Academy in May of 1966.".
- Roger A. Horn and Charles R. Johnson. *Matrix Analysis*. Cambridge University Press, Cambridge, 1990. ISBN 0-521-30586-1. URL http://books.google.com/books?vid=isbn0521305861. Library: QA188H66 1985.
- Alfred Edward Housman. *More Poems*. Alfred A. Knopf, 1936. URL http://books.google.com/books?id=rTMiAAAAMAAJ.
- K Husimi. Studies on the foundations of quantum mechanics i. *Proceedings of the Physico-Mathematical Society of Japan*, 19:766–789, 1937.
- Satoru Igari. *Real Analysis*—*With an Introduction to Wavelet Theory*, volume 177 of *Translations of mathematical monographs*. American Mathematical Society, 1996. ISBN 9780821821046. URL http://books.google.com/books?vid=ISBN0821821040.
- Taizo Iijima. Basic theory of pattern observation. *Papers of Technical Group on Automata and Automatic Control*, December 1959. see Weickert 1999 for historical information.
- Chris J. Isham. *Modern Differential Geometry for Physicists*. World Scientific Publishing, New Jersey, 2 edition, 1999. ISBN 9810235623. URL http://books.google.com/books?vid=ISBN9810235623.
- C.J. Isham. Quantum topology and quantisation on the lattice of topologies. *Classical and Quantum Gravity*, 6:1509–1534, November 1989. doi: 10.1088/0264-9381/6/11/007. URL http://www.iop.org/EJ/abstract/0264-9381/6/11/007.
- Vasile I. Istrățescu. *Inner Product Structures: Theory and Applications*. Mathematics and Its Applications. D. Reidel Publishing Company, 1987. ISBN 9789027721822. URL http://books.google.com/books?vid=ISBN9027721823.
- Luisa Iturrioz. Ordered structures in the description of quantum systems: mathematical progress. In *Methods and applications of mathematical logic: proceedings of the VII Latin American Symposium on Mathematical Logic held July 29-August 2, 1985*, volume 69, pages 55–75, Providence Rhode Island, July 29–August 2 1985. Sociedade Brasileira de Lógica, Sociedade Brasileira de Matemática, and the Association for Symbolic Logic, AMS Bookstore (1988). ISBN 0821850768.





A. J. E. M. Janssen. The zak transform: A signal transform for sampled time-continuous signals. *Philips Journal of Research*, 43(1):23–69, 1988.


- Bjorn Jawerth and Wim Sweldens. An overview of wavelet based multiresolution analysis. SIAM Review, 36:377–412, September 1994. URL http://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.45.5864&rep=rep1&type=pdf.
- Alan Jeffrey and Hui Hui Dai. *Handbook of Mathematical Formulas and Integrals*. Handbook of Mathematical Formulas and Integrals Series. Academic Press, 4 edition, January 18 2008. ISBN 9780080556840. URL http://books.google.com/books?vid=ISBN0080556841.
- Palle E. T. Jørgensen, Kathy Merrill, Judith Packer, and Lawrence W. Baggett. *Representations, Wavelets and Frames: A Celebration of the Mathematical Work of Lawrence Baggett.* Applied and Numerical Harmonic Analysis. Birkhäuser, 2008. ISBN 9780817646820. URL http://books.google.com/books?vid=ISBN0817646825.
- K. D. Joshi. *Applied Discrete Structures*. New Age International, New Delhi, 1997. ISBN 8122408265. URL http://books.google.com/books?vid=ISBN8122408265.
- J.S.Chitode. *Signals And Systems*. Technical Publications, 2009. ISBN 9788184316780. URL http://books.google.com/books?vid=ISBN818431678X.
- Jean-Pierre Kahane. Partial differential equations, trigonometric series, and the concept of function around 1800: a brief story about lagrange and fourier. In Dorina Mitrea and V. G. Mazía, editors, *Perspectives In Partial Differential Equations, Harmonic Analysis And Applications: a Volume in Honor of Vladimir G. Maz'ya's 70th birthday*, volume 79 of *Proceedings of Symposia in Pure Mathematics*, pages 187–206. American Mathematical Society, 2008. ISBN 0821844245. URL http://books.google.com/books?vid=ISBN0821844245.
- Shizuo Kakutani and George W. Mackey. Ring and lattice characterizations of complex hilbert space. *Bulletin of the American Mathematical Society*, 52:727–733, 1946. doi: 10.1090/S0002-9904-1946-08644-9. URL http://www.ams.org/bull/1946-52-08/S0002-9904-1946-08644-9/.
- David W. Kammler. *A First Course in Fourier Analysis*. Cambridge University Press, 2 edition, 2008. ISBN 9780521883405. URL http://books.google.com/books?vid=ISBN0521883407.
- Edward Kasner and James Roy Newman. *Mathematics and the Imagination*. Simon and Schuster, 1940. ISBN 0486417034. URL http://books.google.com/books?vid=ISBN0486417034. "unabridged and unaltered republication" available from Dover.
- Yitzhak Katznelson. *An Introduction to Harmonic Analysis*. Cambridge mathematical library. Cambridge University Press, 3 edition, 2004. ISBN 0521543592. URL http://books.google.com/books?vid=ISBN0521543592.
- James P. Keener. *Principles of Applied Mathematics; Transformation and Approximation*. Addison-Wesley Publishing Company, Reading, Massachusets, 1988. ISBN 0201156741. URL http://www.worldcat.org/isbn/0201156741.
- Yoshida Kenko. The Tsuredzure Gusa of Yoshida No Kaneyoshi. Being the meditations of a recluse in the 14th Century (Essays in Idleness). circa 1330. URL http://www.humanistictexts.org/kenko.htm. 1911 translation of circa 1330 text.
- Anthony W Knapp. *Advanced Real Analysis*. Cornerstones. Birkhäuser, Boston, Massachusetts, USA, 1 edition, July 29 2005a. ISBN 0817643826. URL http://books.google.com/books?vid=ISBN0817643826.



Anthony W Knapp. Basic Real Analysis. Cornerstones. Birkhäuser, Boston, Massachusetts, USA, 1 edition, July 29 2005b. ISBN 0817632506. URL http://books.google.com/books?vid= ISBN0817632506.

- M. G. Krein and M. A. Krasnoselski. Fundamental theorems concerning the extension of hermitian operators and some of their applications to the theory of orthogonal polynomials and the moment problem. Uspekhi Matematicheskikh Nauk, 2(3), 1947. URL http://www.turpion.org/ php/homes/pa.phtml?jrnid=rm. (Russian Mathematical Surveys).
- M. G. Krein, M. A. Krasnoselski, and D. P. Milman. Concerning the deficiency numbers of linear operators in banach space and some geometric questions. Sbornik Trudov Inst. Matem. AN Ukrainian SSR, 11:97-112, 1948.
- Carlos S. Kubrusly. The Elements of Operator Theory. Springer, 1 edition, 2001. **ISBN** 9780817641740. URL http://books.google.com/books?vid=ISBN0817641742.
- The Elements of Operator Theory. Carlos S. Kubrusly. **ISBN** Springer, 2 edition, 2011. 9780817649975. URL http://books.google.com/books?vid=ISBN0817649972.
- Andrew Kurdila and Michael Zabarankin. Convex Functional Analysis. Systems & Control: Foundations & Applications. Birkhäuser, Boston, 2005. ISBN 9783764321987. URL http://books. google.com/books?vid=ISBN3764321989.
- Joseph-Louis Lagrange, Pierre-Simon Laplace, Étienne Louis Malus, René Just Haüy, and Adrien-Marie Legendre. Proclamation des prix décernés dans la séance publique de 6 janvier 1812. Esprit des Journaux, Français et étrangers par Une Societe de Gens Delettres, 2:111-112, January 6 1812a. URL http://books.google.com/books?id=QpUUAAAAQAAJ.
- Joseph-Louis Lagrange, Pierre-Simon Laplace, Étienne Louis Malus, René Just Haüy, and Adrien-Marie Legendre. Proclamation des prix décernés dans la séance publique de 6 janvier 1812. Mercure De France, Journal Littéraire et Politique, 50:374–375, January 6 1812b. URL http: //books.google.com/books?id=8HxBAAAAAAJ.
- Imre Lakatos. Proofs and Refutations: The Logic of Mathematical Discovery. Cambridge University Press, Cambridge, 1976. ISBN 0521290384. URL http://books.google.com/books?vid= ISBN0521290384.
- Traian Lalescu. Sur les équations de Volterra. PhD thesis, University of Paris, 1908. advisor was Émile Picard.
- Traian Lalescu. Introduction à la théorie des équations intégrales (Introduction to the Theory of Integral Equations). Librairie Scientifique A. Hermann, Paris, 1911. URL http://www.worldcat. org/oclc/1278521. first book about integral equations ever published.
- Pierre-Simon Laplace. Theorie Analytique des Probabilités. 1812.
- Pierre-Simon Laplace. Theorie Analytique des Probabilités. Paris, 2 edition, 1814. URL https: //archive.org/details/bub_gb_6MRLAAAAMAAJ/.
- Pierre-Simon Laplace. A Philosophical Essay on Probabilities. Wiley, 1902. URL https://books. google.com/books?id=DfI7AQAAMAAJ.
- Rupert Lasser. Introduction to Fourier Series, volume 199 of Monographs and textbooks in pure and applied mathematics. Marcel Dekker, New York, New York, USA, February 8 1996. ISBN 978-0824796105. URL http://books.google.com/books?vid=ISBN0824796101. QA404.L33 1996.







Peter D. Lax. Functional Analysis. John Wiley & Sons Inc., USA, 2002. ISBN 0-471-55604-1. URL http://www.worldcat.org/isbn/0471556041. QA320.L345 2002.


- Gottfried W. Leibniz. Symbolismus memorabilis calculi algebraici et infinitesimalis, in comparatione potentiarum et differentiarum; et de lege homogeneorum transcendentali. *Miscellanea Berolinensia ad incrementum scientiarum, ex scriptis Societati Regiae scientarum,* pages 160–165, 1710. URL http://bibliothek.bbaw.de/bibliothek-digital/digitalequellen/schriften/anzeige/index_html?band=01-misc/1& seite:int=184.
- Gottfried Wilhelm Leibniz. Letter to christian huygens, 1679. In Leroy E. Loemker, editor, *Philosophical Papers and Letters*, volume 2 of *The New Synthese Historical Library*, chapter 27, pages 248–249. Kluwer Academic Press, Dordrecht, 2 edition, September 8 1679. ISBN 902770693X. URL http://books.google.com/books?vid=ISBN902770693X.
- Pierre Gilles Lemarié, editor. Les Ondelettes en 1989, volume 1438 of Lecture Notes in Mathematics. Springer-Verlag, Berlin, 1990. ISBN 978-3-540-52932-3. URL http://link.springer.com/book/10.1007/BFb0083510/.
- Ezra C. Levy. A New Table of Laplace Transformation Pairs. Space Technology Laboratories, Inc., P.O. Box 95001 Los Angeles 45, California, November 1958. "This Laplace transform table contains 1024 possible transformation pairs in nine sections....In this table we shall deal with the one-sided Laplace transformation, as contrasted to the two-sided transformation...The reason for this choice is that for the systems that are most commonly considered the solution of interest is that corresponding to real values of time, and which assumes knowledge of the initial conditions, i. e., the dynamic characteristics of the system at t=0."
- Tony Lindeberg. *Scale-Space Theory in Computer Vision*. The Springer International Series in Engineering and Computer Science. Springer, 1993. ISBN 9780792394181. URL http://books.google.com/books?vid=ISBN0792394186.
- Joram Lindenstrauss and Lior Tzafriri. *Classical Banach Spaces I: Sequence Spaces*, volume 92 of *Ergebnisse der Mathematik und ihrer Grenzgebiete*. Springer-Verlag, 1977. ISBN 978-3642665592. URL http://www.amazon.com/dp/3642665594.
- J. Liouville. Sur l'integration d'une classe d'équations différentielles du second ordre en quantités finies explicites. *Journal De Mathematiques Pures Et Appliquees*, 4:423–456, 1839. URL http://gallica.bnf.fr/ark:/12148/bpt6k16383z.
- Lynn H. Loomis. *The Lattice Theoretic Background of the Dimension Theory of Operator Algebras*, volume 18 of *Memoirs of the American Mathematical Society*. American Mathematical Society, Providence RI, 1955. ISBN 0821812181. URL http://books.google.com/books?id=P3V1_1XCFRkC.
- Lynn H. Loomis and Ethan D. Bolker. *Harmonic analysis*. Mathematical Association of America, 1965. URL http://books.google.com/books?id=MEfvAAAAMAAJ.
- Nikolai Luzin. Sur la convergence des séries trigonom etriers de fourier. *C. R. Acad. Sci.*, 156:1655–1658, 1913.
- Niccolò Machiavelli. The Literary Works of Machiavelli: Mandragola, Clizia, A Dialogue on Language, and Belfagor, with Selections from the Private Correspondence. Oxford University Press, 1961. ISBN 0313212481. URL http://www.worldcat.org/isbn/0313212481.
- Colin Maclaurin. Treatise of Fluxions. W. Baynes, 1742. URL http://www.amazon.com/dp/B000863E7M.



Stéphane G. Mallat. Multiresolution approximations and wavelet orthonormal bases of $l^2(\mathbb{R})$. *Transactions of the American Mathematical Society*, 315(1):69–87, September 1989. URL http://blanche.polytechnique.fr/~mallat/papiers/math_multiresolution.pdf.

- Stéphane G. Mallat. *A Wavelet Tour of Signal Processing*. Elsevier, 2 edition, September 15 1999. ISBN 9780124666061. URL http://books.google.com/books?vid=ISBN012466606X.
- J. L. Massera and J. J. Schäffer. Linear differential equations and functional analysis, i. *The Annals of Mathematics, 2nd Series*, 67(3):517–573, May 1958. URL http://www.jstor.org/stable/1969871.
- maxima. Maxima Manual version 5.28.0. 5.28.0 edition. URL http://maxima.sourceforge.net/documentation.html.
- Stefan Mazur. Sur les anneaux linéaires. *Comptes rendus de l'Académie des sciences*, 207:1025–1027, 1938.
- Stefan Mazur and Stanislaus M. Ulam. Sur les transformations isométriques d'espaces vectoriels normées. *Comptes rendus de l'Académie des sciences*, 194:946–948, 1932.
- James Mercer. Functions of positive and negative type and their connection with the theory of integral equations. *Philosophical Transactions of the Royal Society of London*, 209:415–446, 1909. ISSN 02643952. URL http://www.jstor.org/stable/91043.
- Yves Meyer. Wavelets and Operators, volume 37 of Cambridge Studies in Advanced Mathematics. Cambridge University Press, January 12 1992. ISBN 0521458692. URL http://books.google.com/books?vid=ISBN0521458692.
- Anthony N. Michel and Charles J. Herget. *Applied Algebra and Functional Analysis*. Dover Publications, Inc., 1993. ISBN 048667598X. URL http://books.google.com/books?vid=ISBN048667598X. original version published by Prentice-Hall in 1981.
- Fred Mintzer. Filters for distortion-free two-band multi-rate filter banks. *IEEE Transactions on Acoustics, Speech and Signal Processing*, 32, 1985.
- Robert Édouard Moritz. *Memorabilia Mathematica; Or, The Philomath's Quotation-book.* Macmillan, 1914. URL https://archive.org/details/memorabiliamath02morigoog.
- Eddie Ortiz Muniz. A method for deriving various formulas in electrostatics and electromagnetism using lagrange's trigonometric identities. *American Journal of Physics*, 21(140), 1953. doi: 10. 1119/1.1933371. URL http://dx.doi.org/10.1119/1.1933371.
- Ben Noble and James W. Daniel. *Applied Linear Algebra*. Prentice-Hall, Englewood Cliffs, NJ, USA, 3 edition, 1988. ISBN 0130412600. URL http://www.worldcat.org/isbn/0130412600.
- Timur Oikhberg and Haskell Rosenthal. A metric characterization of normed linear spaces. *Rocky Mountain Journal Of Mathematics*, 37(2):597–608, 2007. URL http://www.ma.utexas.edu/users/rosenthl/pdf-papers/95-oikh.pdf.
- Judith Packer. Applications of the work of stone and von neumann to wavelets. In Robert S. Doran and Richard V. Kadison, editors, *Operator Algebras, Quantization, and Noncommutative Geometry: A Centennial Celebration Honoring John Von Neumann and Marshall H. Stone: AMS Special Session on Operator Algebras, Quantization, and Noncommutative Geometry, a Centennial Celebration Honoring John Von Neumann and Marshall H. Stone, January 15-16, 2003, Baltimore, Maryland, volume 365 of Contemporary mathematics—American Mathematical Society, pages 253–280, Baltimore, Maryland, 2004. American Mathematical Society. ISBN 9780821834022. URL http://books.google.com/books?vid=isbn0821834029.*







Lincoln P. Paine. Warships of the World to 1900. Ships of the World Series. Houghton Mifflin Harcourt, 2000. ISBN 9780395984147. URL http://books.google.com/books?vid=ISBN9780395984149.

- Anthanasios Papoulis. *Probability, Random Variables, and Stochastic Processes*. McGraw-Hill Series in Electrical Engineering. McGraw-Hill Book Company, New York, 2 edition, 1984. ISBN 9780070484689. URL http://books.google.com/books?vid=ISBN0070484686.
- Anthanasios Papoulis. *Probability, Random Variables, and Stochastic Processes*. McGraw-Hill, New York, 3 edition, 1991. ISBN 0070484775. URL http://books.google.com/books?vid=ISBN0070484775.
- Giuseppe Peano. *Calcolo geometrico secondo l'Ausdehnungslehre di H. Grassmann preceduto dalle operazioni della logica deduttiva*. Fratelli Bocca Editori, Torino, 1888a. Geometric Calculus: According to the *Ausdehnungslehre* of H. Grassmann.
- Giuseppe Peano. Geometric Calculus: According to the Ausdehnungslehre of H. Grassmann. Springer (2000), 1888b. ISBN 0817641262. URL http://books.google.com/books?vid=isbn0817641262. originally published in 1888 in Italian.
- Michael Pedersen. Functional Analysis in Applied Mathematics and Engineering. Chapman & Hall/CRC, New York, 2000. ISBN 9780849371691. URL http://books.google.com/books?vid=ISBN0849371694. Library QA320.P394 1999.
- Wesley J. Perschbacher, editor. *The New Analytical Greek Lexicon*. Hendrickson Publishers, Peabody, Mass., 1990. ISBN 978-0-943575-33-9. URL http://www.amazon.com/dp/0943575338.
- Mark A. Pinsky. *Introduction to Fourier Analysis and Wavelets*. Brooks/Cole, Pacific Grove, 2002. ISBN 9780534376604. URL http://www.amazon.com/dp/0534376606.
- C Piron. Axiomatique quantique. *Helvetica Physica Acta*, 37:439–468, 1964a. ISSN 0018-0238. English translation completed by M. Cole.
- C Piron. Qunatum axiomatics. *Helvetica Physica Acta*, 1964b. English translation of *Axiomatique quantique*, RB4 Technical memo 107/106/104, GPO Engineering Department (London).
- Henri Poincaré. La Science et l'hypothèse. 1902a. URL http://fr.wikisource.org/wiki/La_Science_et_1%27hypoth%C3%A8se. (Science and Hypothesis).
- Henri Poincaré. *Science and Hypothesis*. Dover Publications (1952), New York, 1902b. ISBN 0486602214. URL http://books.google.com/books?vid=isbn0486602214. translation of La Science et l'hypothèse.
- Lakshman Prasad and Sundararaja S. Iyengar. Wavelet Analysis with Applications to Image Processing. CRC Press LLC, Boca Raton, 1997. ISBN 978-0849331695. URL http://books.google.com/books?vid=ISBN0849331692. Library TA1637.P7 1997.
- John G. Proakis. *Digital Communications*. McGraw Hill, 4 edition, 2001. ISBN 0-07-232111-3. URL http://www.mhhe.com/.
- Murray H. Protter and Charles B. Jr. Morrey. *Intermediate Calculus*. Undergraduate Texts in Mathematics. Springer Science & Business Media, 2, illustrated edition, 2012. ISBN 9781461210863. URL http://books.google.com/books?vid=ISBN1461210860. "Softcover reprint of the hard-cover 2nd edition 1985" (page iv).



Ptolemy. *Ptolemy's Almagest*. Springer-Verlag (1984), New York, circa 100AD. ISBN 0387912207. URL http://gallica.bnf.fr/ark:/12148/bpt6k3974x.

- Shie Qian and Dapang Chen. *Joint time-frequency analysis: methods and applications*. PTR Prentice Hall, 1996. ISBN 9780132543842. URL http://books.google.com/books?vid=ISBN0132543842.
- Charles Earl Rickart. *General Theory of Banach Algebras*. University series in higher mathematics. D. Van Nostrand Company, Yale University, 1960. URL http://books.google.com/books?id=PVrvAAAAMAAJ.
- Theodore J. Rivlin. *The Chebyshev Polynomials*. Pure and Applied Mathematics: A Wiley-Interscience Series of Texts, Monographs and Tracts. John Wiley & Sons, New York, 1974. ISBN 0-471-72470-X. URL http://books.google.com/books?vid=ISBN047172470X.
- Enders A. Robinson. *Random Wavelets and Cybernetic Systems*, volume 9 of *Griffins Statistical Monographs & Courses*. Lubrecht & Cramer Limited, London, June 1962. ISBN 0852640757. URL http://books.google.com/books?vid=ISBN0852640757.
- Enders A. Robinson. Multichannel z-transforms and minimum delay. *Geophyics*, 31(3):482–500, June 1966. doi: 10.1190/1.1439788. URL http://dx.doi.org/10.1190/1.1439788.
- Enders A. Robinson. A historical perspective of spectrum estimation. *Proceedings of the IEEE*, 70 (9):885–907, September 1982. doi: 10.1109/PROC.1982.12423. URL http://www.archive.ece.cmu.edu/~ece792/handouts/Robinson82.pdf.
- Maxwell Rosenlicht. *Introduction to Analysis*. Dover Publications, New York, 1968. ISBN 0-486-65038-3. URL http://books.google.com/books?vid=ISBN0486650383.
- Walter Rudin. Real and Complex Analysis. McGraw-Hill Book Company, New York, New York, USA, 3 edition, 1987. ISBN 9780070542341. URL http://www.amazon.com/dp/0070542341. Library QA300.R8 1976.
- Walter Rudin. Functional Analysis. McGraw-Hill, New York, 2 edition, 1991. ISBN 0-07-118845-2. URL http://www.worldcat.org/isbn/0070542252. Library QA320.R83 1991.
- Bryan P. Rynne and Martin A. Youngson. *Linear Functional Analysis*. Springer undergraduate mathematics series. Springer, 2 edition, January 1 2008. ISBN 9781848000056. URL http://books.google.com/books?vid=ISBN1848000057.
- Shôichirô Sakai. *C*-Algebras and W*-Algebras*. Springer-Verlag, Berlin, 1 edition, 1998. ISBN 9783540636335. URL http://books.google.com/books?vid=ISBN3540636331. reprint of 1971 edition.
- Usa Sasaki. Orthocomplemented lattices satisfying the exchange axiom. *Journal of Science of the Hiroshima University*, 17:293–302, 1954. ISSN 0386-3034. URL http://journalseek.net/cgi-bin/journalseek/journalsearch.cgi?field=issn&query=0386-3034.
- Frank Moulton Saxelby. *A course in practical mathematics*. Longmans, Green, and Co., 39 Paternoster Row, London, 6 edition, 1920. URL https://archive.org/details/courseinpractica00saxerich/.
- Juljusz Schauder. Zur theorie stetiger abbildungen in funktionalräumen. *Mathematische Zeitschrift*, 26:47–65, 1927. URL http://eudml.org/doc/167911;jsessionid=156A34EBAB6C0E2DDAAC8C1232D23E8F.





Juljusz Schauder. Eine eigenschaft des haarschen orthogonalsystems. *Mathematische Zeitschrift*, 28:317–320, 1928.

- Gert Schubring. *Conflicts Between Generalization, Rigor, and Intuition: Number Concepts Underlying the Development of Analysis in 17th–19th Century France and Germany.* Sources and studies in the history of mathematics and physical sciences. Springer, New York, 1 edition, June 2005. ISBN 0387228365. URL http://books.google.com/books?vid=ISBN0387228365.
- Isaac Schur. Uber die charakterischen wurzeln einer linearen substitution mit enier anwendung auf die theorie der integralgleichungen (over the characteristic roots of one linear substitution with an application to the theory of the integral). *Mathematische Annalen*, 66:488–510, 1909. URL http://dz-srv1.sub.uni-goettingen.de/cache/toc/D38231.html.
- Mícheál Ó Searcóid. *Elements of Abstract Analysis*. Springer Undergraduate Mathematics Series. Springer, 2002. ISBN 9781852334246. URL http://books.google.com/books?vid=ISBN185233424X.
- Atle Selberg. Harmonic analysis and discontinuous groups in weakly symmetric riemannian spaces with applications to dirichlet series. *Journal of the Indian Mathematical Society*, 20:47–87, 1956.
- Morteza Shafii-Mousavi. *Laplace Transforms Essentials*. Essentials. Research & Education Association, Piscataway, New Jersey, USA, 2015. ISBN 9780738672458. URL http://books.google.com/books?vid=ISBN9780738672458.
- George Finlay Simmons. *Calculus Gems: Brief Lives and Memorable Mathematicians*. Mathematical Association of America, Washington DC, 2007. ISBN 0883855615. URL http://books.google.com/books?vid=ISBN0883855615.
- Iván Singer. Bases in Banach Spaces I, volume 154 of Grundlehren der mathematischen Wissenschaften in Einzeldarstellungen mit besonderer Berücksichtigung der Anwendungsgebiete. Springer-Verlag New York, New York, 1970. ISBN 9780387048338.
- Neil J. A. Sloane. On-line encyclopedia of integer sequences. World Wide Web, 2014. URL http://oeis.org/.
- Jonathan D. H. Smith. *Introduction to Abstract Algebra*. CRC Press, March 23 2011. ISBN 1420063723. URL http://books.google.com/books?vid=ISBN1420063723.
- M.J.T. Smith and T.P. Barnwell. A procedure for designing exact reconstruction filter banks for tree-structured subband coders. *IEEE International Conference on Acoustics, Speech and Signal Processing*, 9:421–424, 1984a. T.P. Barnwell is T.P. Barnwell III.
- M.J.T. Smith and T.P. Barnwell. The design of digital filters for exact reconstruction in subband coding. *IEEE Transactions on Acoustics, Speech and Signal Processing*, 34(3):434–441, June 1984b. ISSN 0096-3518. doi: 10.1109/TASSP.1986.1164832. T.P. Barnwell is T.P. Barnwell III.
- Houshang H. Sohrab. *Basic Real Analysis*. Birkhäuser, Boston, 1 edition, 2003. ISBN 978-0817642112. URL http://books.google.com/books?vid=ISBN0817642110.
- Lynn Arthur Steen. Highlights in the history of spectral theory. *The American Mathematical Monthly*, 80(4):359–381, April 1973. ISSN 00029890. URL http://www.jstor.org/stable/2319079.
- Anne K. Steiner. The lattice of topologies: Structure and complementation. *Transactions of the American Mathematical Society*, 122(2):379–398, April 1966. URL http://www.jstor.org/stable/1994555.



Marshall Harvey Stone. *Linear transformations in Hilbert space and their applications to analysis*, volume 15 of *American Mathematical Society. Colloquium publications*. American Mathematical Society, New York, 1932. URL http://books.google.com/books?vid=ISBN0821810154. 1990 reprint of the original 1932 edition.

- Gilbert Strang and Truong Nguyen. *Wavelets and Filter Banks*. Wellesley-Cambridge Press, Wellesley, MA, 1996. ISBN 9780961408879. URL http://books.google.com/books?vid=ISBN0961408871.
- Robert S. Strichartz. *The Way of Analysis*. Jones and Bartlett Publishers, Boston-London, 1995. ISBN 978-0867204711. URL http://books.google.com/books?vid=ISBN0867204710.
- Endre Süli and David F. Mayers. *An Introduction to Numerical Analysis*. Cambridge University Press, August 28 2003. ISBN 9780521007948. URL http://books.google.com/books?vid=ISBN0521007941.
- Wim Sweldens and Robert Piessens. Wavelet sampling techniques. In 1993 Proceedings of the Statistical Computing Section, pages 20–29. American Statistical Association, August 1993. URL http://citeseer.ist.psu.edu/18531.html.
- Erik Talvila. Necessary and sufficient conditions for differentiating under the integral sign. *The American Mathematical Monthly*, 108(6):544–548, June–July 2001. URL http://arxiv.org/abs/math/0101012.
- Brook Taylor. *Methodus Incrementorum Directa et Inversa*. London, 1715. URL https://books.google.com/books?id=oA5tpwAACAAJ.
- Brook Taylor. *Methodus Incrementorum Directa et Inversa*. London, 1717. URL https://books.google.com/books?id=zbgYn4lq4-EC.
- Audrey Terras. *Fourier Analysis on Finite Groups and Applications*. Number 43 in London Mathematical Society Student Texts. Cambridge University Press, Cambridge, 1999. ISBN 0-521-45718-1. URL http://books.google.com/books?vid=ISBN0521457181.
- Brian S. Thomson, Andrew M. Bruckner, and Judith B. Bruckner. *Elementary Real Analysis*. www.classicalrealanalysis.com, 2 edition, 2008. ISBN 9781434843678. URL http://classicalrealanalysis.info/com/Elementary-Real-Analysis.php.
- Stanislaw Marcin Ulam. *Adventures of a Mathematician*. University of California Press, Berkeley, 1991. ISBN 0520071549. URL http://books.google.com/books?vid=ISBN0520071549.
- P.P. Vaidyanathan. *Multirate Systems and Filter Banks*. Prentice Hall Signal Processing Series. Prentice Hall, 1993. ISBN 0136057187. URL http://books.google.com/books?vid=ISBN0136057187.
- Jussi Väisälä. A proof of the mazur-ulam theorem. *The American Mathematical Monthly*, 110(7): 633–635, August–September 2003. URL http://www.helsinki.fi/~jvaisala/mazurulam.pdf.
- Balthasar van der Pol and Hendricus Bremmer. Operational Calculus based on the Two-Sided Laplace Integral. Cambridge at the University Press, 2nd, reprinted edition, 1959. URL https://archive.org/details/in.ernet.dli.2015.141269/.
- Brani Vidakovic. *Statistical Modeling by Wavelets*. John Wiley & Sons, Inc, New York, 1999. ISBN 9780471293651. URL http://www.amazon.com/dp/0471293652.





John von Neumann. Allgemeine eigenwerttheorie hermitescher funktionaloperatoren. *Mathematische Annalen*, 102(1):49–131, 1929. ISSN 0025-5831 (print) 1432-1807 (online). URL http://resolver.sub.uni-goettingen.de/purl?GDZPPN002273535. General eigenvalue theory of Hermitian functional operators.

- John von Neumann. *Continuous Geometry*. Princeton mathematical series. Princeton University Press, Princeton, 1960. URL http://books.google.com/books?id=3bjq0gAACAAJ.
- David F. Walnut. *An Introduction to Wavelet Analysis*. Applied and numerical harmonic analysis. Springer, 2002. ISBN 0817639624. URL http://books.google.com/books?vid=ISBN0817639624.
- Gilbert G. Walter and XiaoPing Shen. *Wavelets and Other Orthogonal Systems*. Chapman and Hall/CRC, New York, 2 edition, 2001. ISBN 9781584882275. URL http://books.google.com/books?vid=ISBN1584882271.
- Heinrich Martin Weber. Die allgemeinen grundlagen der galois'schen gleichungstheorie. *Mathematische Annalen*, 43(4):521–549, December 1893. URL http://resolver.sub.uni-goettingen.de/purl?GDZPPN002254670. The general foundation of Galois' equation theory.
- Joseph H. Maclagan Wedderburn. On hypercomplex numbers. *Proceedings of the London Mathematical Society*, 6:77–118, 1907. URL http://plms.oxfordjournals.org/cgi/reprint/s2-6/1/77.
- Joachim Weickert. Linear scale-space has first been proposed in japan. *Journal of Mathematical Imaging and Vision*, 10:237–252, May 1999. URL http://dl.acm.org/citation.cfm?id=607668.
- Hermann Weyl. The method of orthogonal projection in potential theory. *Duke Mathematical Journal*, 7(1):411–444, 1940. URL http://projecteuclid.org/euclid.dmj/1077492266.
- W. John Wilbur. Quantum logic and the locally convex spaces. *Transactions of the American Mathematical Society*, 207:343–360, June 1975. URL http://www.jstor.org/stable/1997181.
- P. Wojtaszczyk. *A Mathematical Introduction to Wavelets*, volume 37 of *London Mathematical Society student texts*. Cambridge University Press, February 13 1997. ISBN 9780521578943. URL http://books.google.com/books?vid=ISBN0521578949.
- Robert M. Young. *An introduction to nonharmonic Fourier series*, volume 93 of *Pure and applied mathematics*. Academic Press, 1 edition, 1980. ISBN 0127728503. URL http://books.google.com/books?vid=ISBN0127728503.
- Robert M. Young. *An introduction to nonharmonic Fourier series*, volume 93 of *Pure and applied mathematics*. Academic Press, revised first edition, May 16 2001. ISBN 0127729550. URL http://books.google.com/books?vid=ISBN0127729550.
- Ahmed I. Zayed. *Handbook of Function and Generalized Function Transformations*. Mathematical Sciences Reference Series. CRC Press, Boca Raton, 1996. ISBN 0849378516. URL http://books.google.com/books?vid=ISBN0849378516.
- Ahmed I. Zayed. A prelude to sampling, wavelets, and tomography. In John Benedetto and Ahmed I. Zayed, editors, *Sampling, Wavelets, and Tomography*, Applied and Numerical Harmonic Analysis, pages 1–32. Springer, 2004. ISBN 9780817643041. URL http://books.google.com/books?vid=ISBN0817643044.
- Gary Zukav. *The Dancing Wu Li Masters : An Overview of the New Physics*. Bantam Books, New York, 1980. ISBN 055326382X. URL http://books.google.com/books?vid=ISBN055326382X.



REFERENCE INDEX

54, 243

Abramovich and Aliprantis (2002), 96, 208Aliprantis and Burkinshaw (1998), 16, 18, 22, 40, 106, 113, 116, 117, 119–121, 124 Adelson and Burt (1981), 6, 80 Akhiezer and Glazman (1993), 109Ptolemy (circa 100AD), 165 Alvarez et al. (1993), 6, 80 Amemiya and Araki (1966), Andrews et al. (2001), 48 Abramowitz and Stegun (1972), 153, 154, 176, 179 Atkinson and Han (2009), 6, Autonne (1901), 135 Autonne (1902), 135 Bachman (1964), 145, 198, Bachman and Narici (1966), 9, 11, 13, 22, 107, 127, 129 Bachman et al. (2002), 14, 17, 27, 40, 54, 145, 196, 198, 224, 235 Banach (1922), 111, 116 Banach (1932b), 116 Banach (1932a), 14, 116 Gilbert Ben-Israel and (2002), 143Beran (1976), 4 Beran (1985), 4 Berberian (1961), 9, 22, 27, 98, 102, 106, 113-115, 132, Berkson (1963), 109 Bertero and Boccacci (1998), Betten (2008a), 225, 228 Betten (2008b), 223, 236

Birkhoff and Neumann (1936), 4, 109Bollobás (1999), 18, 121, 265-Bottazzini (1986), 160, 161 Boyer and Merzbach (1991), 153 Bracewell (1978), 22, 145, 198, 223-226, 228, 229, 232-235 Bromwich (1908), 207 Bruckner et al. (1997), 184, Burago et al. (2001), 109 Burrus et al. (1998), 251 Burt and Adelson (1983), 6, Byrne (2005), 35 Cardano (1545), 95 Carleson and **Engquist** (2001), 248Carothers (2005), 11, 14 Casazza and Lammers (1998), 40Cayley (1858), 265 Cerdà (2010), 191 J.S.Chitode (2009), 243 Chorin and Hald (2009), 196, Christensen (2003), 9, 14, 22, 27, 28, 32, 35, 40, 42, 43, 54 Christensen (2008), 28, 32 Chui (1992), 68, 72, 86, 142, 240, 255 Claerbout (1976), 211 Cohn (2002), 96 Black et al. (2009), 3, 54 Courant and Hilbert (1930), 268 Dai and Lu (1996), 40 Dai and Larson (1998), 40, 42

Daubechies (1992), 15, 31,

Davidson and Donsig (2010), de Witt (1659), 267 Descartes (1954), 265 Descartes (1637a), 265 Descartes (1637b), 39 Deza and Deza (2006), 109 Daubechies et al. (1986), 32 Dieudonné (1969), 130 Duffin and Schaeffer (1952), 32 Dumitrescu (2007), 211 Dunford and Schwartz (1957), 123Durbin (2000), 95, 96 Albers and Dyson (1994), 79 Lagrange et al. (1812a), 247 Edwards (1995), 18 Eidelman et al. (2004), 114 Enflo (1973), 14 Euler (1748), 153, 160, 161, 170 Ewen (1950), viii Ewen (1961), viii Fabian et al. (2010), 20 Farina and Rinaldi (2000), Fix and Strang (1969), 202 Flanders (1973), 143 Flanigan (1983), 155, 222 Folland (1995), 147, 148, 151, Folland (1992), 48, 159, 184 Forster and Massopust (2009), 51Fourier (1807), 247 Fourier (1878), 1, 195, 196 Fourier (1822), 196 Fréchet (1906), 1 Fréchet (1928), 1

Fredholm (1900), 263, 267

Fredholm (1903), 263, 267 Iijima (1959), 6, 80 Mazur and Ulam (1932), 123 Frobenius (1968), 130 Isham (1989), 2 Mazur (1938), 151 Frobenius (1878), 130 Isham (1999), 2 Lagrange et al. (1812b), 247 Fuchs (1995), 147 Istrățescu (1987), 191 Mercer (1909), 268 Gabor (1946), 51 Iturrioz (1985), 102, 108, 109 Meyer (1992), 54 Gauss (1900), 49 Janssen (1988), 50 Michel and Herget (1993), 9, and Gelfand (1941), 151 **Jawerth Sweldens** 11, 13, 96, 98–103, 112, 114, Gelfand Neumark (1994), 54, 60, 72, 200, 202and 116, 119, 124, 125, 128, 130, (1943b), 150–152 Jeffrey and Dai (2008), 187, 133–135, 140, 149, 150, 263, Gelfand and Neumark 188, 196 265, 267, 268 (1943a), 151, 152 Jørgensen et al. (2008), 32 Mintzer (1985), 215 Moritz (1914), 223 Gel'fand (1963), 137 Joshi (1997), 207 Gelfand and Naimark (1964), Kahane (2008), 247 Muniz (1953), 187, 188 148, 149, 151 Kammler (2008), 40 Noble and Daniel (1988), 137 Gelfand et al. (2018), 137 Kasner and Newman (1940), Oikhberg and Rosenthal Giles (2000), 14, 18, 123, 125 (2007), 123Gohberg et al. (2003), 268 Katznelson (2004), 184, 191, Packer (2004), 40 Goodman et al. (1993b), 42 193, 247 Paine (2000), vi Goodman et al. (1993a), 40, Keener (1988), 129, 263, 267 Papoulis (1984), 263 42 Kenko (circa 1330), 270 Papoulis (1991), 255, 263 Goswami and Chan (1999), Knapp (2005a), 136 Peano (1888b), 111, 124 68, 86, 200 Pedersen (2000), 17, 18, 159, Knapp (2005b), 48, 143, 196 Greenhoe (2013), 7 Krein and Krasnoselski Gudder (1979), 109 (1947), 109Perschbacher (1990), 3, 54 Gudder (2005), 109 Krein et al. (1948), 109 Pinsky (2002), 55, 69, 182, Guichard et al. (2012), 6, 80 Kubrusly (2001), 9, 11, 13, 258, 261 Guichard et al. (20xx), 6, 80 16, 22, 27, 101, 102, 105–107, Poincaré (1902b), 53 de la Vallée-Poussin (1915), Guillemin (1957), 176 111, 113, 127, 132, 133 Gupta (1998), 13 Kubrusly (2011), 102, 103, Haar (1910), 78 207 Prasad and Iyengar (1997), Kurdila Haaser and Sullivan (1991), and Zabarankin 142 (2005), 9Proakis (2001), 76 13, 98, 111, 124 Haddad and Akansu (1992), Lakatos (1976), 173 Protter and Morrey (2012), 215 Lalescu (1908), 159 Halmos (1948), 111 Lalescu (1911), 159 Qian and Chen (1996), 51 Laplace (1812), 223 de Reyna (2002), 191 Halmos (1958), 98, 127 Halmos (1998b), 109 Laplace (1814), 223 Rickart (1960), 148-151 Halmos (1998a), 126, 149 Laplace (1902), 223 Rivlin (1974), 173, 178 Hamel (1905), 11 Robinson (1962), 211 Lasser (1996), 48 Han et al. (2007), 20, 35 Lax (2002), 50, 129 Robinson (1966), 211 Hausdorff (1937), 40 Leibniz (1710), 143 Robinson (1982), 195, 222 Leibniz (1679), 111 Rosenlicht (1968), 153, 155-Hazewinkel (2000), 147 Heijenoort (1967), viii Lemarié (1990), 6, 53 Heil and Walnut (1989), 40 Levy (1958), 223-226, 233, Rudin (1991), 117, 119, 120, Heil (2011), 9, 11, 13, 14, 22, 234, 236 122, 124-127, 129-131, 135, 27, 28, 32, 35, 40, 112, 184, Lindeberg (1993), 6, 80 136 Lindenstrauss and Tzafriri Rudin (1987), 48, 161 Hermite (1893), 173 (1977), 14Rynne and Youngson (2008), Liouville (1839), 155, 159 Hernández and Weiss (1996), Loomis (1955), 109 54, 55, 261 Sakai (1998), 126 Higgins (1996), 50 Loomis and Bolker (1965), Sasaki (1954), 109 Hilbert (1904), 265 196 Saxelby (1920), 168, 169 Hilbert (1906), 266, 267 Machiavelli (1961), 269 Schauder (1927), 14 Hilbert (1912), 265, 267 Maclaurin (1742), 222 Schauder (1928), 14 Hilbert et al. (1927), 113 Mallat (1989), 6, 31, 54, 80 Schubring (2005), 183 Holland (1970), 109 Mallat (1999), 6, 31, 53, 54, Schur (1909), 135 Horn and Johnson (1990), 61, 70, 89, 171, 201, 202, 215, Searcóid (2002), 9 217, 245, 251 Selberg (1956), 50 Housman (1936), viii Massera and Schäffer (1958), Shafii-Mousavi (2015), 225, Husimi (1937), 109 229, 230 Igari (1996), 30, 258, 261 maxima, 178, 181 Simmons (2007), 269



REFERENCE INDEX Daniel J. Greenhoe page 293

Singer (1970), 14 Smith and Barnwell (1984a), Smith and Barnwell (1984b), 215 Smith (2011), 64 Sohrab (2003), 13, 222 Steen (1973), 135, 265 Steiner (1966), 2 Stone (1932), 113, 125, 129 Strang and Nguyen (1996), 6, 81, 215 Strichartz (1995), 222 Süli and Mayers (2003), 178 Sweldens and Piessens

(1993), 200, 202 Talvila (2001), 143 Taylor (1717), 222 Taylor (1715), 222 Terras (1999), 50 Thomson et al. (2008), 207 Ulam (1991), 123 Vaidyanathan (1993), 215 Väisälä (2003), 123 van der Pol and Bremmer (1959), 223, 225, 229 Vidakovic (1999), 31, 200-202, 217, 245 von Neumann (1929), 125, 129, 267

von Neumann (1960), 109 Walnut (2002), 40, 54, 191 Walter and Shen (2001), 18 Weber (1893), 96 Wedderburn (1907), 99 Weyl (1940), 18 Wojtaszczyk (1997), 27, 29-31, 40, 43, 55, 57, 69, 78, 81, 258, 261 Young (1980), 20 Young (2001), 11, 13–15, 20, 24, 25, 27, 28, 32 Zayed (2004), 40 Zayed (1996), 50 Zukav (1980), 170



page 294 Daniel J. Greenhoe REFERENCE INDEX





SUBJECT INDEX

C* algebra, 151, 152	analysis, 3 , 3, 5, 53	23, 24
C^* -algebra, 152	analytic, 195, 221, 222	Bessel's Inequality, 25
*-algebra, 126, 148 , 148–150,	AND, xi	Bessel's inequality, 18
238	anti-causality, 229	Best Approximation Theo-
<i>n</i> th moment, 200 , 200–202	anti-symmetric, 108, 141,	rem, 18 , 18
*-algebras, 126	200, 258	bijection, 25
ŁTĘX, vi	antiautomorphic, 126, 148,	bijective, xi, 25, 123
TEX-Gyre Project, vi	149	Binomial Theorem, 162
XHETEX, vi	antiderivative, 221	binomial theorem, 143
attention markers, 123	antilinear, 149	biorthogonal, 27 , 29, 30
problem, 116, 122, 129,	antitone, 102, 104, 108	Borel measure, 141, 196
131, 265	aperature, 109	Borel sets, 141, 196
2 coefficient case, 78	associates, 112	bounded, xi, 15, 58, 120 , 129,
,	associateve, 145	140, 258
Abel, Niels Henrik, 269	associative, 95, 96, 112, 115,	bounded bijective, 28
absolute value, x, 96	135	bounded linear operator,
absolutely summable, 208,	asymmetric, 211	136
209	auto-correlation function,	bounded linear operators,
abstract space, 1, 1	263	120 , 121, 122, 124, 125, 127,
additive, 45, 57, 105, 113, 115,	auto-power spectrum, 68,	128, 130, 131, 133–136
124, 225–228	86, 255 , 258	bounded operator, 120
additive identity, 113	autocorrelation, 129	•
additive inverse, 113, 163,	autocorrelation function,	C star algebra, 151
165	255	Cardano, Gerolamo, 95
additive property, 197	Avant-Garde, vi	Cardinal Series, 50
additivity, 36, 125		Cardinal series, 50, 59
adjoint, 42, 45, 121 , 122, 125 ,	B-spline $N_1(x)$, 71	Carl Spitzweg, 269
126, 150, 196	B-spline $N_2(x)$, 71	Cartesian product, x
admissibility, 66	B-splines, 60	Cauchy, 107
admissibility condition, 66,	Banach algebra, 151	Cauchy-Schwarz inequality,
74, 78, 88	Banach space, 14, 15	33
admissibility equation, 78	bandlimited, 50, 59	causal, 212 , 212, 229–231
Adobe Systems Incorpo-	basis, 5, 11–13, 20, 50, 51, 64,	ceiling, 70
rated, vi	74, 79	characteristic function, x, 40
affine, 123, 266	frame, 32	characterized, 53
algebra, 95, 96 , 147 , 147, 148	orthogonal, 20	Chebyshev polynomial, 178
algebra of sets, xi, 2	orthonormal, 20	Chebyshev polynomial of the
algebras	Riesz, 27, 28	first kind, 178
C*-algebra, 151	tight frame, 32	Chebyshev polynomials, 178
*-algebra, 148	Battle-Lemarié orthogonal-	closed, 54, 58, 59, 69, 104,
algebras of sets, 2, 3	ization, 31	106, 107
analyses, 3	Bessel's Equality, 17 , 18, 19,	closed interval, 223
• '	± J' ' ' ' '	,

Closed Set Theorem, 104	convolution, 198, 207 , 207,	eigenspace, <mark>265</mark>
closed sets, 99	208, 224	equivalent, 15
closure, 13, 54, 58, 70	convolution operation, 145	exact frame, 32
coefficient functionals, 14,	convolution operator, 207,	expansion, 11 , 14
14	208	exponential function,
coefficients, 213	• • • •	160
comb function, 248	204, 210 , 235	field, 96
commutative, 45, 96, 112,	coordinate functionals, 14	Fourier coefficients, 20
115, 145, 165, 207, 208	coordinates, 11	Fourier expansion, 20
commutative ring, 95, 96 , 96	cosine, 155	Fourier series, 20
commutativity, 208	countably infinite, 14, 267	frame, 32
commutator relation, 42 , 55,	counting measure, xi	frame bounds, 32
72	CQF, 215 , 215, 216	fundamental, 13
compact, 267	CQF condition, 88, 215 , 217	group, 95
compact support, 6, 53, 55,	CQF conditions, 89	Hamel basis, 11
90 compactly supported 80	CQF theorem, 88, 215 , 218 cross-correlation, 263 , 264	hermitian, 148
compactly supported, 80 complement, x		inner product space, 124
complemented, 108	255	linear basis, 11
complete, 13 , 58, 59	cross-power spectrum, 86,	linear combination, 9
complete metric space, 13	255	linear manifold, 98
complete set, 13	CS Inequality, 36, 56	linear space, 112
completeness, 58	Co mequancy, 50, 50	linear subspace, 98
complex auto-power spec-	Daubechies wavelet func-	MRA, 54
trum, 255	tion, 89	MRA space, 54
complex cross-power spec-	Daubechies-1, 89	MRA system, 63
trum, 255	Daubechies-2, 89	multiplicative condi-
complex exponential, 5, 79	Daubechies-3, 89	tion, 151
complex linear space, 112	Daubechies-3 scaling func-	multiresolution analy-
complex number system,	tion, 71	sis, 54
163	de la Vallée Poussin kernel,	multiresolution analysis
conjuage symmetric, 264	191, 193	space, 54
conjugate linear, 126, 148,	de Morgan, 4, 108	multiresolution system,
149	de Morgan's law, 4	63
conjugate pairs, 213	definitions	normal, <mark>148</mark>
conjugate quadrature filter,	C* algebra, 151 , 152	normalized tight frame,
88, 215 , 215	*-algebra, 148 , 238	32
conjugate quadrature filter	abstract space, 1	normed algebra, 151
condition, 88, 215	algebra, 96 , 147	normed linear space,
Conjugate quadrature filters,	Banach algebra, 151 biorthogonal, 27	116, 117
214	bounded linear opera-	normed space of linear
conjugate recipricol pairs,	tors, 120	operators, 117
182	Cetar algebra 151	optimal lower frame
conjugate symmetric, 124,	coefficient functionals,	bound, 32
264, 268	1.4	optimal upper frame
conjugate symmetric prop-	commutative ring, 96	bound, 32
erty, 197	complete, 13	orthogonal basis, 20
constant, 43, 44, 159	complex auto newer	orthogonal comple-
continuity, 63	spectrum, 255	ment, 102
continuous, xi, 5, 43, 44, 57,	complex cross-power	orthonormal basis, 20
63, 69, 79, 114, 141, 151, 191,	cnoctrum 255	orthonormal MRA sys-
221, 222, 258, 268	complex linear space,	tem, 63
continuous point spectrum,	112	Parseval frame, 32
171	continuous spectrum,	point spectrum, 266
continuous spectrum, 266	266	projection, 148 real linear space, 112
converge, 223 convergent, 15	coordinate functionals,	Region of Convergence,
converges, 223	1.4	223
converges absolutely, 268	coordinates, 11	residual spectrum, 266
convex, 2, 98, 99, 106	CQF, 215	resolvent, 147
convexity, 107	dilation operator in-	resolvent set, 266
Convenies, 101	verse, 41	10001v011t 30t, 200





page 297 SUBJECT INDEX Daniel J. Greenhoe

Riesz basis, 27	Dirac delta distribution, 50	Fourier Series, 51
ring, <mark>95</mark>	Dirichlet Kernel, 184–186	Fourier Transform, 51
scalars, 112	Dirichlet kernel, 191	Gabor Transform, 51
scaling function, 54	discontinuous, 195, 222	Haar scaling function,
scaling subspace, 54	discrete, 5, 79	64
Schauder basis, 14	Discrete Time Fourier Series,	Haar wavelet system, 78
Selberg Trace Formula,	xii	linear functions, 50
50	Discrete Time Fourier Trans-	Mercedes Frame, 35
self-adjoint, 148	form, xii, 237	order 0 B-spline wavelet
Smith-Barnwell filter,	Discrete time Fourier trans-	system, 78
215	form, 61–63, 86	Peace Frame, 35
space of all abso-	discrete time Fourier trans-	raised cosine, 77
lutely square summable se-	form, 68, 86	raised sine, 76
quences, 207	discrete-time Fourier trans-	Rectangular pulse, 204
space of Lebesgue	form, 66, 237 , 237–239, 243	rectangular pulse, 203
square-integrable functions,	Dissertation on the propaga-	triangle, 203
141	tion of heat in solid bodies,	wavelets, 51
spans, 9	247	exclusive OR, xi
spectral radius, 147	distributes, 112	existential quantifier, xi
spectrum, 147 , 266	distributive, 5, 80, 126, 145,	exists, 221
standard inner product,	148–150, 209, 238	expansion, 11, 14
141	distributivity, 42	exponential function, 160
standard norm, 141	diverge, 223	false, xi
star-algebra, 148	divergent, 232, 233	Fast Wavelet Transform, 53,
subspace intersection,	domain, x, 39	80, 252
99	double angle formulas, 165 ,	fast wavelet transform, 251
subspace union, 99	166, 167, 169, 189	Fejér's kernel, 191 , 191, 193
support, 70	DTFT, 15, 47, 61, 88, 215–217,	Fejér-Riesz spectral factor-
synthesis, 3	238, 242, 245, 259	ization, 182 , 182
tight frame, 32	DTFT periodicity, 88, 237	field, 95, 96 , 111, 207
total, 13	dyadic, 79	field of complex numbers,
translation operator in-	Dyson, Freeman, 79	126
verse, 41 underlying set, 112	Eigendecomposition, 34	filter banks, 214
unital, 147	eigenspace, 265	floor, 70
vector space, 112	eigenvalue, 34, 265	FontLab Studio, vi
vector space, 112 vectors, 112	eigenvector, 34, 265	for each, <mark>xi</mark>
wavelet analysis, 81	empty set, <mark>xi</mark>	Fourier Analysis, 5, 80
wavelet coefficient se-	equal, 112	Fourier analysis, 4, 5
quence, 82	equalities	fourier analysis, 195
wavelet system, 82	Bessel's, 17	Fourier coefficients, 20 , 50
delay, 139	equality by definition, x	Fourier expansion, 20, 20–
DeMoivre's Theorem, 173,	equality relation, x	22, 24 , 26
173, 175	equation, 60	Fourier kernel, 196
dense, 13, 14, 54, 55, 57, 58,	equations	Fourier Series, xii, 51, 247
69	dilation equation, 60 , 60	Fourier series, 20
Descartes, René, ix, 39	refinement equation, 60	Fourier Series adjoint, 249
difference, x	two-scale difference	Fourier series analysis, 5
differential operator, 171	equation, 60	Fourier series expansion, 22
differentiation operator, 222	two-scale relation, 60	Fourier Series operator, 247
dilation, 138, 198, 224, 225	equivalent, 15 , 15, 27, 136	Fourier Transform, xii, 31, 47,
dilation equation, 60 , 60, 66–	Euler Formulas, 192	51 , 57, 137, 139, 170, 196 ,
68, 70–72, 76, 78	Euler formulas, 161 , 162,	196, 199, 200, 222
dilation equation in	163, 166, 167, 203	adjoint, 197
"time hyperpage, 60	Euler's Identity, 160 , 160,	Fourier transform, 4, 15, 48,
dilation operator, 5, 40, 40,	161, 164, 169, 227	49, 54, 61–63, 72, 86, 139, 200,
42, 43	even, 157, 178, 186, 230, 234,	201, 203, 255
dilation operator adjoint, 42	242	inverse, 197
dilation operator inverse, 41	exact frame, 32	Fourier Transform operator,
dimension, 11	examples	42
Dirac delta, 248	2 coefficient case, 78	Fourier transform scaling
	Cardinal Series, 50	factor, 196



Fourier, Joseph, 1, 195	de la Vallée Poussin ker-	sequence, 207
frame, 30, 32 , 32, 36	nel, 191, 193	set indicator, 259
frame bound, 33, 35	dilation equation, 76, 78	set indicator function,
frame bounds, 32, 36	dilation operator, 43	56, 74, 76, 204
frame operator, 32 , 32, 33, 35	Dirac delta, 248	sine, 155
frame property, 56, 58	Dirichlet Kernel, 184,	spectral density func-
frames, 10	186	tion, 261
Fredholm integral equation	Dirichlet kernel, 191	standard inner product,
of the first kind, 263, 267	Discrete Time Fourier	255
Fredholm integral operators,	Transform, 237	subspace addition, 84
137	Discrete time Fourier	summability kernel, 191
Fredholm operator, 267 , 267,	transform, 61–63, 86	Taylor expansion, 153
268	discrete time Fourier	translation operator, 40,
Fredholm operators, 267	transform, 68, 86	202
Free Software Foundation, vi	discrete-time Fourier	unit step, 228–230, 232–
function, 40, 112, 141, 196	transform, 66, 237–239	234
characteristic, 40	DTFT, 216, 242, 245	unit step function, 228
even, 242	eigenvector, 34	Volterra integral equa-
indicator, 40	Fejér's kernel, 191 , 191,	tion, 163, 165
functional, 126	193	Volterra integral equa-
functions, xi	Fourier coefficients, 50	tion of the second type, 159
<i>n</i> th moment, 200, 201	Fourier kernel, 196	wavelet, 51
absolute value, 96	Fourier transform, 48,	wavelet function, 81 , 82
adjoint, 150	49, 61–63, 72, 86, 200, 201,	z transform, 211
antiderivative, 221	203, 255	Z-transform, 66, 239
auto-correlation func-	Fredholm integral equa-	z-transform, 215, 216,
tion, 263	tion of the first kind, 267	255
auto-power spectrum,	Heaviside step function,	Zak Transform, 50
68, 86, 255 , 258	228	fundamental, 13
autocorrelation func-	indicator function, 40	Fundamental Theorem of
tion, 255	induced norm, 23	Calculus, 221, 228, 236
11011, 200		
B-spline N _{x} (x) 71	inner product 124 196	Fundamental theorem of
B-spline $N_1(x)$, 71	inner product, 124 , 196 Jackson kernel 191 193	Fundamental theorem of calculus 142 143
B-spline $N_2(x)$, 71	Jackson kernel, 191, 193	calculus, 142, 143
B-spline $N_2(x)$, 71 B-splines, 60	Jackson kernel, 191, 193 Kronecker delta func-	calculus, 142, 143 Fundamental theorem of lin-
B-spline N ₂ (x), 71 B-splines, 60 Borel measure, 141, 196	Jackson kernel, 191, 193 Kronecker delta function, 20 , 72	calculus, 142, 143 Fundamental theorem of linear equations, 116 , 116
B-spline $N_2(x)$, 71 B-splines, 60 Borel measure, 141, 196 characteristic function,	Jackson kernel, 191, 193 Kronecker delta function, 20 , 72 kronecker delta func-	calculus, 142, 143 Fundamental theorem of lin-
B-spline N ₂ (x), 71 B-splines, 60 Borel measure, 141, 196 characteristic function,	Jackson kernel, 191, 193 Kronecker delta function, 20, 72 kronecker delta function, 243	calculus, 142, 143 Fundamental theorem of linear equations, 116, 116 FWT, 251, 252
B-spline N ₂ (x), 71 B-splines, 60 Borel measure, 141, 196 characteristic function, 40 Chebyshev polynomial,	Jackson kernel, 191, 193 Kronecker delta function, 20, 72 kronecker delta function, 243 linear functional, 122	calculus, 142, 143 Fundamental theorem of linear equations, 116, 116 FWT, 251, 252 g.l.b., 59
B-spline N ₂ (x), 71 B-splines, 60 Borel measure, 141, 196 characteristic function, 40 Chebyshev polynomial, 178	Jackson kernel, 191, 193 Kronecker delta function, 20, 72 kronecker delta function, 243 linear functional, 122 mean, 263	calculus, 142, 143 Fundamental theorem of linear equations, 116, 116 FWT, 251, 252 g.l.b., 59 Gabor Transform, 51
B-spline N ₂ (x), 71 B-splines, 60 Borel measure, 141, 196 characteristic function, 40 Chebyshev polynomial, 178 Chebyshev polynomial	Jackson kernel, 191, 193 Kronecker delta function, 20, 72 kronecker delta function, 243 linear functional, 122 mean, 263 measure, 260	calculus, 142, 143 Fundamental theorem of linear equations, 116, 116 FWT, 251, 252 g.l.b., 59 Gabor Transform, 51 gap metric, 109
B-spline N ₂ (x), 71 B-splines, 60 Borel measure, 141, 196 characteristic function, 40 Chebyshev polynomial, 178 Chebyshev polynomial of the first kind, 178	Jackson kernel, 191, 193 Kronecker delta function, 20, 72 kronecker delta function, 243 linear functional, 122 mean, 263 measure, 260 Minkowski addition, 83	calculus, 142, 143 Fundamental theorem of linear equations, 116, 116 FWT, 251, 252 g.l.b., 59 Gabor Transform, 51 gap metric, 109 Gaussian Pyramid, 6, 80
B-spline N ₂ (x), 71 B-splines, 60 Borel measure, 141, 196 characteristic function, 40 Chebyshev polynomial, 178 Chebyshev polynomial of the first kind, 178 comb function, 248	Jackson kernel, 191, 193 Kronecker delta function, 20, 72 kronecker delta function, 243 linear functional, 122 mean, 263 measure, 260 Minkowski addition, 83 modulus, 96	calculus, 142, 143 Fundamental theorem of linear equations, 116, 116 FWT, 251, 252 g.l.b., 59 Gabor Transform, 51 gap metric, 109 Gaussian Pyramid, 6, 80 Gelfand-Mazur Theorem,
B-spline N ₂ (x), 71 B-splines, 60 Borel measure, 141, 196 characteristic function, 40 Chebyshev polynomial, 178 Chebyshev polynomial of the first kind, 178 comb function, 248 complex exponential, 5,	Jackson kernel, 191, 193 Kronecker delta function, 20, 72 kronecker delta function, 243 linear functional, 122 mean, 263 measure, 260 Minkowski addition, 83 modulus, 96 mother wavelet, 79, 81	calculus, 142, 143 Fundamental theorem of linear equations, 116, 116 FWT, 251, 252 g.l.b., 59 Gabor Transform, 51 gap metric, 109 Gaussian Pyramid, 6, 80 Gelfand-Mazur Theorem, 151
B-spline N ₂ (x), 71 B-splines, 60 Borel measure, 141, 196 characteristic function, 40 Chebyshev polynomial, 178 Chebyshev polynomial of the first kind, 178 comb function, 248 complex exponential, 5,	Jackson kernel, 191, 193 Kronecker delta function, 20, 72 kronecker delta function, 243 linear functional, 122 mean, 263 measure, 260 Minkowski addition, 83 modulus, 96 mother wavelet, 79, 81 norm, 116, 117	calculus, 142, 143 Fundamental theorem of linear equations, 116, 116 FWT, 251, 252 g.l.b., 59 Gabor Transform, 51 gap metric, 109 Gaussian Pyramid, 6, 80 Gelfand-Mazur Theorem, 151 Generalized Parseval's Iden-
B-spline N ₂ (x), 71 B-splines, 60 Borel measure, 141, 196 characteristic function, 40 Chebyshev polynomial, 178 Chebyshev polynomial of the first kind, 178 comb function, 248 complex exponential, 5, 79 conjugate quadrature	Jackson kernel, 191, 193 Kronecker delta function, 20, 72 kronecker delta function, 243 linear functional, 122 mean, 263 measure, 260 Minkowski addition, 83 modulus, 96 mother wavelet, 79, 81 norm, 116, 117 operator norm, 117	calculus, 142, 143 Fundamental theorem of linear equations, 116, 116 FWT, 251, 252 g.l.b., 59 Gabor Transform, 51 gap metric, 109 Gaussian Pyramid, 6, 80 Gelfand-Mazur Theorem, 151 Generalized Parseval's Identity, 22
B-spline N ₂ (x), 71 B-splines, 60 Borel measure, 141, 196 characteristic function, 40 Chebyshev polynomial, 178 Chebyshev polynomial of the first kind, 178 comb function, 248 complex exponential, 5, 79 conjugate quadrature filter, 215	Jackson kernel, 191, 193 Kronecker delta function, 20, 72 kronecker delta function, 243 linear functional, 122 mean, 263 measure, 260 Minkowski addition, 83 modulus, 96 mother wavelet, 79, 81 norm, 116, 117 operator norm, 117 Parseval's equation, 250	calculus, 142, 143 Fundamental theorem of linear equations, 116, 116 FWT, 251, 252 g.l.b., 59 Gabor Transform, 51 gap metric, 109 Gaussian Pyramid, 6, 80 Gelfand-Mazur Theorem, 151 Generalized Parseval's Identity, 22 generalized product rule,
B-spline N ₂ (x), 71 B-splines, 60 Borel measure, 141, 196 characteristic function, 40 Chebyshev polynomial, 178 Chebyshev polynomial of the first kind, 178 comb function, 248 complex exponential, 5, 79 conjugate quadrature filter, 215 continuous point spec-	Jackson kernel, 191, 193 Kronecker delta function, 20, 72 kronecker delta function, 243 linear functional, 122 mean, 263 measure, 260 Minkowski addition, 83 modulus, 96 mother wavelet, 79, 81 norm, 116, 117 operator norm, 117 Parseval's equation, 250 Plancheral's formula,	calculus, 142, 143 Fundamental theorem of linear equations, 116, 116 FWT, 251, 252 g.l.b., 59 Gabor Transform, 51 gap metric, 109 Gaussian Pyramid, 6, 80 Gelfand-Mazur Theorem, 151 Generalized Parseval's Identity, 22 generalized product rule, 143, 143
B-spline N ₂ (x), 71 B-splines, 60 Borel measure, 141, 196 characteristic function, 40 Chebyshev polynomial, 178 Chebyshev polynomial of the first kind, 178 comb function, 248 complex exponential, 5, 79 conjugate quadrature filter, 215 continuous point spectrum, 171	Jackson kernel, 191, 193 Kronecker delta function, 20, 72 kronecker delta function, 243 linear functional, 122 mean, 263 measure, 260 Minkowski addition, 83 modulus, 96 mother wavelet, 79, 81 norm, 116, 117 operator norm, 117 Parseval's equation, 250 Plancheral's formula,	calculus, 142, 143 Fundamental theorem of linear equations, 116, 116 FWT, 251, 252 g.l.b., 59 Gabor Transform, 51 gap metric, 109 Gaussian Pyramid, 6, 80 Gelfand-Mazur Theorem, 151 Generalized Parseval's Identity, 22 generalized product rule, 143, 143 geometric series, 184, 187
B-spline N ₂ (x), 71 B-splines, 60 Borel measure, 141, 196 characteristic function, 40 Chebyshev polynomial, 178 Chebyshev polynomial of the first kind, 178 comb function, 248 complex exponential, 5, 79 conjugate quadrature filter, 215 continuous point spectrum, 171 cosine, 155	Jackson kernel, 191, 193 Kronecker delta function, 20, 72 kronecker delta function, 243 linear functional, 122 mean, 263 measure, 260 Minkowski addition, 83 modulus, 96 mother wavelet, 79, 81 norm, 116, 117 operator norm, 117 Parseval's equation, 250 Plancheral's formula,	calculus, 142, 143 Fundamental theorem of linear equations, 116, 116 FWT, 251, 252 g.l.b., 59 Gabor Transform, 51 gap metric, 109 Gaussian Pyramid, 6, 80 Gelfand-Mazur Theorem, 151 Generalized Parseval's Identity, 22 generalized product rule, 143, 143 geometric series, 184, 187 globally, 221
B-spline N ₂ (x), 71 B-splines, 60 Borel measure, 141, 196 characteristic function, 40 Chebyshev polynomial, 178 Chebyshev polynomial of the first kind, 178 comb function, 248 complex exponential, 5, 79 conjugate quadrature filter, 215 continuous point spectrum, 171 cosine, 155 cross-correlation, 263	Jackson kernel, 191, 193 Kronecker delta function, 20, 72 kronecker delta function, 243 linear functional, 122 mean, 263 measure, 260 Minkowski addition, 83 modulus, 96 mother wavelet, 79, 81 norm, 116, 117 operator norm, 117 Parseval's equation, 250 Plancheral's formula, 250 Plancherel's formula,	calculus, 142, 143 Fundamental theorem of linear equations, 116, 116 FWT, 251, 252 g.l.b., 59 Gabor Transform, 51 gap metric, 109 Gaussian Pyramid, 6, 80 Gelfand-Mazur Theorem, 151 Generalized Parseval's Identity, 22 generalized product rule, 143, 143 geometric series, 184, 187 globally, 221 GNU Octave
B-spline N ₂ (x), 71 B-splines, 60 Borel measure, 141, 196 characteristic function, 40 Chebyshev polynomial, 178 Chebyshev polynomial of the first kind, 178 comb function, 248 complex exponential, 5, 79 conjugate quadrature filter, 215 continuous point spectrum, 171 cosine, 155 cross-correlation, 263 cross-correlation func-	Jackson kernel, 191, 193 Kronecker delta function, 20, 72 kronecker delta function, 243 linear functional, 122 mean, 263 measure, 260 Minkowski addition, 83 modulus, 96 mother wavelet, 79, 81 norm, 116, 117 operator norm, 117 Parseval's equation, 250 Plancheral's formula, 250 Plancherel's formula, 250 Poisson kernel, 191, 193	calculus, 142, 143 Fundamental theorem of linear equations, 116, 116 FWT, 251, 252 g.l.b., 59 Gabor Transform, 51 gap metric, 109 Gaussian Pyramid, 6, 80 Gelfand-Mazur Theorem, 151 Generalized Parseval's Identity, 22 generalized product rule, 143, 143 geometric series, 184, 187 globally, 221 GNU Octave cos, 178, 181
B-spline N ₂ (x), 71 B-splines, 60 Borel measure, 141, 196 characteristic function, 40 Chebyshev polynomial, 178 Chebyshev polynomial of the first kind, 178 comb function, 248 complex exponential, 5, 79 conjugate quadrature filter, 215 continuous point spectrum, 171 cosine, 155 cross-correlation, 263 cross-correlation function, 255	Jackson kernel, 191, 193 Kronecker delta function, 20, 72 kronecker delta function, 243 linear functional, 122 mean, 263 measure, 260 Minkowski addition, 83 modulus, 96 mother wavelet, 79, 81 norm, 116, 117 operator norm, 117 Parseval's equation, 250 Plancheral's formula, 250 Plancherel's formula, 250 Poisson kernel, 191, 193 Poisson Summation	calculus, 142, 143 Fundamental theorem of linear equations, 116, 116 FWT, 251, 252 g.l.b., 59 Gabor Transform, 51 gap metric, 109 Gaussian Pyramid, 6, 80 Gelfand-Mazur Theorem, 151 Generalized Parseval's Identity, 22 generalized product rule, 143, 143 geometric series, 184, 187 globally, 221 GNU Octave cos, 178, 181 Golden Hind, vi
B-spline N ₂ (x), 71 B-splines, 60 Borel measure, 141, 196 characteristic function, 40 Chebyshev polynomial, 178 Chebyshev polynomial of the first kind, 178 comb function, 248 complex exponential, 5, 79 conjugate quadrature filter, 215 continuous point spectrum, 171 cosine, 155 cross-correlation, 263 cross-correlation function, 255 cross-power spectrum,	Jackson kernel, 191, 193 Kronecker delta function, 20, 72 kronecker delta function, 243 linear functional, 122 mean, 263 measure, 260 Minkowski addition, 83 modulus, 96 mother wavelet, 79, 81 norm, 116, 117 operator norm, 117 Parseval's equation, 250 Plancheral's formula, 250 Plancherel's formula, 250 Poisson kernel, 191, 193 Poisson Summation Formula, 50	calculus, 142, 143 Fundamental theorem of linear equations, 116, 116 FWT, 251, 252 g.l.b., 59 Gabor Transform, 51 gap metric, 109 Gaussian Pyramid, 6, 80 Gelfand-Mazur Theorem, 151 Generalized Parseval's Identity, 22 generalized product rule, 143, 143 geometric series, 184, 187 globally, 221 GNU Octave cos, 178, 181 Golden Hind, vi GPR, 143
B-spline N ₂ (x), 71 B-splines, 60 Borel measure, 141, 196 characteristic function, 40 Chebyshev polynomial, 178 Chebyshev polynomial of the first kind, 178 comb function, 248 complex exponential, 5, 79 conjugate quadrature filter, 215 continuous point spectrum, 171 cosine, 155 cross-correlation, 263 cross-correlation function, 255 cross-power spectrum, 86, 255	Jackson kernel, 191, 193 Kronecker delta function, 20, 72 kronecker delta function, 243 linear functional, 122 mean, 263 measure, 260 Minkowski addition, 83 modulus, 96 mother wavelet, 79, 81 norm, 116, 117 operator norm, 117 Parseval's equation, 250 Plancheral's formula, 250 Plancherel's formula, 250 Poisson kernel, 191, 193 Poisson Summation Formula, 50 power spectrum, 258	calculus, 142, 143 Fundamental theorem of linear equations, 116, 116 FWT, 251, 252 g.l.b., 59 Gabor Transform, 51 gap metric, 109 Gaussian Pyramid, 6, 80 Gelfand-Mazur Theorem, 151 Generalized Parseval's Identity, 22 generalized product rule, 143, 143 geometric series, 184, 187 globally, 221 GNU Octave cos, 178, 181 Golden Hind, vi GPR, 143 greatest lower bound, xi, 54,
B-spline N ₂ (x), 71 B-splines, 60 Borel measure, 141, 196 characteristic function, 40 Chebyshev polynomial, 178 Chebyshev polynomial of the first kind, 178 comb function, 248 complex exponential, 5, 79 conjugate quadrature filter, 215 continuous point spectrum, 171 cosine, 155 cross-correlation, 263 cross-correlation function, 255 cross-power spectrum, 86, 255 Daubechies wavelet	Jackson kernel, 191, 193 Kronecker delta function, 20, 72 kronecker delta function, 243 linear functional, 122 mean, 263 measure, 260 Minkowski addition, 83 modulus, 96 mother wavelet, 79, 81 norm, 116, 117 operator norm, 117 Parseval's equation, 250 Plancheral's formula, 250 Plancherel's formula, 250 Poisson kernel, 191, 193 Poisson Summation Formula, 50 power spectrum, 258 random process, 263	calculus, 142, 143 Fundamental theorem of linear equations, 116, 116 FWT, 251, 252 g.l.b., 59 Gabor Transform, 51 gap metric, 109 Gaussian Pyramid, 6, 80 Gelfand-Mazur Theorem, 151 Generalized Parseval's Identity, 22 generalized product rule, 143, 143 geometric series, 184, 187 globally, 221 GNU Octave cos, 178, 181 Golden Hind, vi GPR, 143 greatest lower bound, xi, 54, 55, 59, 70
B-spline N ₂ (x), 71 B-splines, 60 Borel measure, 141, 196 characteristic function, 40 Chebyshev polynomial, 178 Chebyshev polynomial of the first kind, 178 comb function, 248 complex exponential, 5, 79 conjugate quadrature filter, 215 continuous point spectrum, 171 cosine, 155 cross-correlation, 263 cross-correlation function, 255 cross-power spectrum, 86, 255 Daubechies wavelet function, 89	Jackson kernel, 191, 193 Kronecker delta function, 20, 72 kronecker delta function, 243 linear functional, 122 mean, 263 measure, 260 Minkowski addition, 83 modulus, 96 mother wavelet, 79, 81 norm, 116, 117 operator norm, 117 Parseval's equation, 250 Plancheral's formula, 250 Plancherel's formula, 250 Poisson kernel, 191, 193 Poisson Summation Formula, 50 power spectrum, 258 random process, 263 random variable, 263	calculus, 142, 143 Fundamental theorem of linear equations, 116, 116 FWT, 251, 252 g.l.b., 59 Gabor Transform, 51 gap metric, 109 Gaussian Pyramid, 6, 80 Gelfand-Mazur Theorem, 151 Generalized Parseval's Identity, 22 generalized product rule, 143, 143 geometric series, 184, 187 globally, 221 GNU Octave cos, 178, 181 Golden Hind, vi GPR, 143 greatest lower bound, xi, 54, 55, 59, 70 greatest value, 71
B-spline N ₂ (x), 71 B-splines, 60 Borel measure, 141, 196 characteristic function, 40 Chebyshev polynomial, 178 Chebyshev polynomial of the first kind, 178 comb function, 248 complex exponential, 5, 79 conjugate quadrature filter, 215 continuous point spectrum, 171 cosine, 155 cross-correlation, 263 cross-correlation function, 255 cross-power spectrum, 86, 255 Daubechies wavelet function, 89 Daubechies-1, 89	Jackson kernel, 191, 193 Kronecker delta function, 20, 72 kronecker delta function, 243 linear functional, 122 mean, 263 measure, 260 Minkowski addition, 83 modulus, 96 mother wavelet, 79, 81 norm, 116, 117 operator norm, 117 Parseval's equation, 250 Plancheral's formula, 250 Plancherel's formula, 250 Poisson kernel, 191, 193 Poisson Summation Formula, 50 power spectrum, 258 random process, 263 random variable, 263 Riesz sequence, 27	calculus, 142, 143 Fundamental theorem of linear equations, 116, 116 FWT, 251, 252 g.l.b., 59 Gabor Transform, 51 gap metric, 109 Gaussian Pyramid, 6, 80 Gelfand-Mazur Theorem, 151 Generalized Parseval's Identity, 22 generalized product rule, 143, 143 geometric series, 184, 187 globally, 221 GNU Octave cos, 178, 181 Golden Hind, vi GPR, 143 greatest lower bound, xi, 54, 55, 59, 70 greatest value, 71 group, 95, 95
B-spline N ₂ (x), 71 B-splines, 60 Borel measure, 141, 196 characteristic function, 40 Chebyshev polynomial, 178 Chebyshev polynomial of the first kind, 178 comb function, 248 complex exponential, 5, 79 conjugate quadrature filter, 215 continuous point spectrum, 171 cosine, 155 cross-correlation, 263 cross-correlation function, 255 cross-power spectrum, 86, 255 Daubechies wavelet function, 89 Daubechies-1, 89 Daubechies-2, 89	Jackson kernel, 191, 193 Kronecker delta function, 20, 72 kronecker delta function, 243 linear functional, 122 mean, 263 measure, 260 Minkowski addition, 83 modulus, 96 mother wavelet, 79, 81 norm, 116, 117 operator norm, 117 Parseval's equation, 250 Plancheral's formula, 250 Plancherel's formula, 250 Poisson kernel, 191, 193 Poisson Summation Formula, 50 power spectrum, 258 random process, 263 random variable, 263 Riesz sequence, 27 scalar product, 124	calculus, 142, 143 Fundamental theorem of linear equations, 116, 116 FWT, 251, 252 g.l.b., 59 Gabor Transform, 51 gap metric, 109 Gaussian Pyramid, 6, 80 Gelfand-Mazur Theorem, 151 Generalized Parseval's Identity, 22 generalized product rule, 143, 143 geometric series, 184, 187 globally, 221 GNU Octave cos, 178, 181 Golden Hind, vi GPR, 143 greatest lower bound, xi, 54, 55, 59, 70 greatest value, 71
B-spline N ₂ (x), 71 B-splines, 60 Borel measure, 141, 196 characteristic function, 40 Chebyshev polynomial, 178 Chebyshev polynomial of the first kind, 178 comb function, 248 complex exponential, 5, 79 conjugate quadrature filter, 215 continuous point spectrum, 171 cosine, 155 cross-correlation, 263 cross-correlation function, 255 cross-power spectrum, 86, 255 Daubechies wavelet function, 89 Daubechies-1, 89 Daubechies-2, 89 Daubechies-3, 89	Jackson kernel, 191, 193 Kronecker delta function, 20, 72 kronecker delta function, 243 linear functional, 122 mean, 263 measure, 260 Minkowski addition, 83 modulus, 96 mother wavelet, 79, 81 norm, 116, 117 operator norm, 117 Parseval's equation, 250 Plancheral's formula, 250 Plancherel's formula, 250 Poisson kernel, 191, 193 Poisson Summation Formula, 50 power spectrum, 258 random process, 263 random variable, 263 Riesz sequence, 27 scalar product, 124 scaling coefficient se-	calculus, 142, 143 Fundamental theorem of linear equations, 116, 116 FWT, 251, 252 g.l.b., 59 Gabor Transform, 51 gap metric, 109 Gaussian Pyramid, 6, 80 Gelfand-Mazur Theorem, 151 Generalized Parseval's Identity, 22 generalized product rule, 143, 143 geometric series, 184, 187 globally, 221 GNU Octave cos, 178, 181 Golden Hind, vi GPR, 143 greatest lower bound, xi, 54, 55, 59, 70 greatest value, 71 group, 95, 95
B-spline N ₂ (x), 71 B-splines, 60 Borel measure, 141, 196 characteristic function, 40 Chebyshev polynomial, 178 Chebyshev polynomial of the first kind, 178 comb function, 248 complex exponential, 5, 79 conjugate quadrature filter, 215 continuous point spectrum, 171 cosine, 155 cross-correlation, 263 cross-correlation function, 255 cross-power spectrum, 86, 255 Daubechies wavelet function, 89 Daubechies-1, 89 Daubechies-2, 89	Jackson kernel, 191, 193 Kronecker delta function, 20, 72 kronecker delta function, 243 linear functional, 122 mean, 263 measure, 260 Minkowski addition, 83 modulus, 96 mother wavelet, 79, 81 norm, 116, 117 operator norm, 117 Parseval's equation, 250 Plancheral's formula, 250 Plancherel's formula, 250 Poisson kernel, 191, 193 Poisson Summation Formula, 50 power spectrum, 258 random process, 263 random variable, 263 Riesz sequence, 27 scalar product, 124	calculus, 142, 143 Fundamental theorem of linear equations, 116, 116 FWT, 251, 252 g.l.b., 59 Gabor Transform, 51 gap metric, 109 Gaussian Pyramid, 6, 80 Gelfand-Mazur Theorem, 151 Generalized Parseval's Identity, 22 generalized product rule, 143, 143 geometric series, 184, 187 globally, 221 GNU Octave cos, 178, 181 Golden Hind, vi GPR, 143 greatest lower bound, xi, 54, 55, 59, 70 greatest value, 71 group, 95, 95 Gutenberg Press, vi





Subject Index Daniel J. Greenhoe page 299

half-angle formulas, 167	into, 26	Lebesgue square-integrable
Hamel bases, 10	inverse, 33, 35, 40, 41, 95, 112	functions, 39, 196, 223
Hamel basis, 11, 11, 13	inverse DTFT, 217, 243, 259,	left distributive, 95, 96, 115
Handbook of Algebras, 147	260	Leibnitz GPR, 218
harmonic analysis, 195	inverse Fourier Series, 248	Leibniz integration rule, 143
Harmonic shifted orthonor-	Inverse Fourier Transform,	Leibniz rule, 143 , 143
mality requirement, 261	137	Leibniz, Gottfried, ix, 111
Hasse diagram, 53, 59, 80, 83	Inverse Fourier transform,	linear, 50, 59, 113 , 113, 171
Heaviside step function, 228	197	212, 226
Hermetian transpose, 137	inverse Fourier Transform,	linear basis, 11
Hermite, Charles, 173	139	linear bounded, xi
hermitian, 129 , 148 , 148, 149	Inverse Poisson Summation	linear combination, 9
hermitian components, 150	Formula, 49 , 49	linear combinations, 10
Hermitian representation,	invertible, 15, 32, 142, 148	linear functional, 122
150	involutary, 126	linear functions, 50
Hermitian symmetric, 182,	involution, 148 , 148, 152	linear independence, 16, 17
200	involutory, 108, 148–150	linear manifold, 98
Heuristica, vi	IPSF, 49 , 49, 256	linear operator, 28, 30
high-pass filter, 214	irrational numbers, 44	linear operators, 26, 113, 122
Hilbert space, 4, 20, 22, 24,	irreflexive ordering relation,	linear ops., 26, 30
27, 30, 32, 106, 107, 125, 126,	xi	Linear space, 97
129–131, 136, 196	isometric, 21, 29, 30, 123,	linear space, 1, 2, 9, 11, 58
homogeneous, 20, 21, 25, 28–	132 , 132, 136, 197, 250	98–102, 112 , 112, 147
30, 58, 96, 105, 113, 115–117,	isometric in distance, 45, 136	linear spaces, 112
124, 225–227, 230, 231, 233,	isometric in length, 45, 136	linear span, 9 , 64, 98, 99
235	isometric operator, 133–135	linear subspace, 2, 9, 97, 98
Housman, Alfred Edward, vii	isometry, 132	98–103, 105
	isomorphic, 25	linear subspaces, 99
identities	isomorphic, 20	linear time invariant, 171
Fourier expansion, 21,	Jackson kernel, 191, 193	linearity, 57, 67, 113, 114
22	Jacobi, Carl Gustav Jacob,	linearly dependent, 9 , 11–13
Parseval frame, 21	183	linearly independent, 9 , 11-
identity, 95, 112	jaib, 153	13, 16, 17
identity element, 112		linearly ordered, 6, 54, 69, 80
identity operator, 41, 112,	jiba, 153 jiva, 153	linearly ordered set, 59
112	join, xi	Liquid Crystal, vi
if, xi	join, xi	locally, 221
if and only if, xi	11.11.1.050	low-pass filter, 214
image, x	Kaneyoshi, Urabe, 270	•
image set, 114, 116, 127–131,	Kenko, Yoshida, 270	Machiavelli, Niccolò, 269
136	kernel, 263	Maclaurin Series, 222
imaginary part, xi, 149	Kronecker delta function, 20 ,	maps to, x
implied by, xi	72	matrix
implies, xi	kronecker delta function,	rotation, 137
implies and is implied by, xi	243	Maxima, 178, 181
inclusive OR, xi		Mazur-Ulam theorem, 123
indicator function, x, 40	l'Hôpital's rule, 186, 249	mean, <mark>263</mark>
induced norm, 23	l.u.b., 59	measure, 260
inequality	Lagrange trigonometric	meet, <mark>xi</mark>
Bessel's, 18	identities, 187	Mercedes Frame, 35
triangle, 116, 117	Laplace convolution, 199	Mercer's Theorem, 268 , 268
infinite sum, 9	Laplace operator, 137	metric, xi, 58
injective, xi, 114, 115	Laplace Transform, 170, 198,	metric space, 1
inner product, 58, 124 , 196	199, 205, 223 , 223–226, 228–	metrics
inner product space, 16–18,	230, 232–236	gap, 109
20, 27, 102, 103, 105, 106, 124	Laplace transform, 171	Schäffer, 109
inner-product, xi	lattice, 2, 97, 102	Minimum phase, 219
inside, 212	lattice of algebras of sets, 2	minimum phase, 90, 211
integral domain, 95	Laurent series, 208	211, 212
integration, 221	least upper bound, xi, 59, 68,	Minkowski addition, 81, 83
Integration by Parts, 236	70	99
intersection, x	least value, 70	modular, 5, 80





page 300 Daniel J. Greenhoe Subject Index

modulation, 226, 227, 229–	one sided snift operator, 134	inverse Fourier Trans-
232, 234	only if, <mark>xi</mark>	form, 139
modulus, 96	open interval, 223	involution, 148
mother wavelet, 79, 81	opening, 109	kernel, 263
MRA, 6, 54 , 54, 55, 61, 69, 74–	operations	Laplace operator, 137
77, 80	adjoint, 42, 45, 121 , 122,	Laplace Transform, 198,
MRA space, 54 , 54, 58, 60	125	199, 205, 223 , 223–226, 228–
-		
MRA system, 63 , 63, 64, 66–	analysis, 3 , 53	230, 232–236
68, 70	closure, 70	Laplace transform, 171
multiplicative condition,	convolution, 207, 208,	linear operators, 122
151	224	linear span, 9
Multiresolution Analysis, 53	convolution operation,	Maclaurin Series, 222
multiresolution analysis, 6,	145	Minkowski addition, 81
54 , 54, 59, 80, 82	differential operator,	modulation, 226
multiresolution analysis	171	operator, 32, 112
J T		-
space, 54 , 60	differentiation operator,	operator adjoint, 126
multiresolution anaysis, 81	222	operator inverse, 61
multiresolution system, 63,	dilation, 224	projection, 127
72, 81, 82	dilation operator, 5, 40,	projection operator, 55,
12, 01, 02	-	
NET W. O	40, 42	57
N5 lattice, 2	dilation operator ad-	Rationalizing the De-
Neumann Expansion Theo-	joint, 42	nominator, 229, 230
rem, 124	Discrete Time Fourier	real part, 149
non-analytic, 222		
	Series, xii	reflection operator, 137
non-Boolean, 7, 80	Discrete Time Fourier	reversal, 225
non-complemented, 7, 80	Transform, xii	rotation matrix, 137
non-distributive, 2, 7, 80	discrete-time Fourier	rotation operator, 137
non-homogeneous, 159		
	transform, 237	sampling operator, 48,
non-isotropic, 105	DTFT, 15, 47, 61, 88, 215,	48
non-modular, 7, 80	217, 238, 259	Taylor Series, 222, 222
non-negative, 96, 117, 124,	Eigendecomposition, 34	transform, 3 , 54
256		
	Fast Wavelet Transform,	translation, 224
nonBoolean, 83	252	translation operator, 5,
noncommutative, 42	Fourier Series, xii, 247	40 , 40, 42
noncomplemented, 83		
nondegenerate, 44, 55, 57,	Fourier Series adjoint,	translation operator ad-
	249	joint, 42
58, 96, 116, 117, 124	Fourier Series operator,	Unitary Fourier Trans-
nondistributive, 83	247	form, 224
nonmodular, 83	Fourier Transform, xii,	unitary Fourier Trans-
norm, 58, 116 , 117		•
	31, 47, 57, 137, 139, 196 , 196,	form, 196
normal, 129, 130 , 130, 131,	199, 200, 222	Z-Transform, <mark>xii</mark>
136, 137, 148 , 250, 267	Fourier transform, 4, 15,	Z-transform, 88
normal operator, 130, 135	54	z-transform, 208 , 208
normalized tight frame, 32		
0	frame operator, 32 , 32,	operator, 32, 40, 111, 112
normed algebra, 151, 151,	33, 35	adjoint, 149
152	Fredholm integral equa-	autocorrelation, 129
normed linear space, 116,	tion of the first kind, 263	bounded, 120
117		
	Fredholm operator, 267	definition, 112
normed linear spaces, 60,	FWT, 252	delay, 139
121, 132	Hermetian transpose,	dilation, 138
normed space of linear oper-	-	
	137	identity, 112
ators, 117	identity operator, 41,	isometric, 133–135
NOT, xi	112	linear, 113
not constant, 44	imaginary part, 149	norm, 117
not total, 23		
	integration, 221	normal, 130, 131, 135
not unique, 102	inverse, 33, 35, 41	Null Space, 127
Null Space, 114–116, 125,	inverse DTFT, 259, 260	positive, 140
127–131, 136	inverse Fourier Series,	projection, 127
null space, x, 265		- ,
11a11 space, A, 200	248	range, 127
odd 157 170 100 000 005	Inverse Fourier Trans-	self-adjoint, 129
odd, 157, 178, 186, 230, 235	form, 137	shift, 134
andelette 6 81	•	· · · · · · · · · · · · · · · · · · ·





translation, 137	periodic, 40, 48, 191, 237, 256	129, 140, 258
unbounded, 120	Plancheral's formula, 250	bounded bijective, 28
unitary, 135, 136, 197, 250	Plancherel's formula, 197, 250	Cartesian product, x Cauchy, 107
operator adjoint, 125, 126	Poincaré, Jules Henri, 53	causal, 212 , 229–231
operator inverse, 61	point spectrum, 266	characteristic function,
operator norm, xi, 45, 117	Poisson kernel, 191, 193	X
operator star-algebra, 126	Poisson Summation For-	closed, 54, 58, 59, 69,
optimal lower frame bound,	mula, 48 , 50	104, 106, 107
32	polar identity, 17	closure, 13, 58
optimal upper frame bound,	pole, 219	commutative, 45, 96,
32	poles, 213	112, 115, 145, 165, 207, 208
order, x, xi	polynomial	commutativity, 208
order 0 B-spline wavelet sys-	trigonometric, 173	compact, 267
tem, 78 order relation, 97	positive, 140 power set, xi	compact support, 6, 53, 55, 90
ordered pair, x	power spectrum, 258	compactly supported,
ordered set, 70, 83, 108	Primorial numbers, 7, 80	80
orthnormal, 88	probability space, 263	complement, x
orthocomplemented lattice,	product identities, 163 , 164,	complemented, 108
108	165, 167, 187, 188	complete, 58, 59
orthogonal, 16, 16, 20, 84,	projection, 127 , 148	completeness, 58
129, 168, 261	projection operator, 55, 57,	conjugate linear, 126,
orthogonal basis, 20	127, 129	148
orthogonal complement,	projection operators, 4	conjugate quadrature
102 , 102, 107 orthogonality, 16, 81	Projection Theorem, 107 , 108	filter condition, 88, 215 conjugate symmetric,
inner product space, 16	proper subset, x	124, 264, 268
orthomodular, 108	proper superset, x	constant, 43, 44, 159
orthomodular identity, 108	properties	continuity, 63
orthomodular lattice, 4, 108	absolute value, x	continuous, 5, 43, 44, 57,
orthonormal, 17–23, 36, 63,	absolutely summable,	63, 69, 79, 114, 141, 191, 221,
68, 88, 261	208, 209	222, 258, 268
orthonormal bases, 10	additive, 45, 57, 105, 113,	converge, 223
orthonormal basis, 20, 24,	115, 124, 225–228	convergent, 15
25, 27–29, 31, 250	additive identity, 113	converges, 223
orthonormal expansion, 20,	additive inverse, 113,	converges absolutely,
24 orthonormal MRA system,	163, 165 additivity, 36, 125	268 convex, 2, 98, 99, 106
63	admissibility, 66	convexity, 107
orthonormal quadrature	admissibility condition,	countably infinite, 14,
conditions, 243	78	267
orthonormal wavelet sys-	affine, 123, 266	counting measure, xi
tem, 68, 84, 86	algebra of sets, xi	CQF condition, 88, 215,
orthonormality, 18, 28, 29,	analytic, 195, 221, 222	217
36, 90	AND, xi	de Morgan, 4, 108
orthornormal basis, 21	anti-causality, 229	de Morgan's law, 4
oscillation, 221	anti-symmetric, 108, 141, 200, 258	dense, 13, 14, 54, 55, 57,
Paley-Wiener, 50	antiautomorphic, 126,	58, 69 difference, x
parallelogram law, 107	148, 149	dilation, 198, 225
Parseval frame, 21, 32 , 32	antitone, 102, 104, 108	dilation equation in
Parseval frames, 10	associates, 112	"time hyperpage, 60
Parseval's equation, 197, 250	associateve, 145	discontinuous, 195, 222
Parseval's Identity, 22, 24, 26,	associative, 95, 96, 112,	discrete, 5, 79
32, 58, 261 partition of unity, 54, 71, 72,	115, 135	distributes, 112
74, 76, 78, 90	bandlimited, 59	distributive, 5, 80, 126,
Peace Frame, 35	bijection, 25	145, 148–150, 209, 238
Peirce, Benjamin, 170	bijective, 25, 123	distributivity, 42
Per Enflo, 14	biorthogonal, 29, 30	divergent 222 222
Perfect reconstruction, 20	bounded, 15, 58, 120 ,	divergent, 232, 233





domain, x	115	orthonormal, 17–23, 36,
dyadic, 79	linear, 50, 59, 113 , 113,	63, 68, 88, 261
empty set, <mark>xi</mark>	171, 226	orthonormality, 18, 28,
equal, 112	linear independence,	29, 36, 90
equality by definition, x	16, 17	oscillation, 221
equality relation, x	linear time invariant,	Paley-Wiener, 50
equivalent, 15, 27, 136	171	partition of unity, 71, 72,
even, 157, 178, 186, 230,	linearity, 57, 67, 113, 114	74, 76, 78, 90
234, 242	linearly dependent, 9,	periodic, 40, 48, 191,
exclusive OR, <mark>xi</mark>	11–13	237, 256
existential quantifier, xi	linearly independent, 9,	positive, 140
exists, 221	11–13, 16, 17	power set, xi
false, <mark>xi</mark>	linearly ordered, 6, 54,	proper subset, x
for each, xi	69, 80	proper superset, x
frame property, 56, 58	locally, 221	pseudo-distributes, 112
Fredholm operator, 267,	maps to, x	range, x
268	meet, xi	real, 213, 240, 258
Fredholm operators, 267	metric, xi	real part, xi
globally, 221	Minimum phase, 219	real-valued, 129, 200,
greatest lower bound, xi	minimum phase, 90,	238, 239, 242
hermitian, 129 , 148, 149	211 , 211, 212	recursive, 59
Hermitian symmetric,	modular, 5, 80	reflexive, 108
182, 200	modulation, 227, 229–	reflexive ordering rela-
homogeneous, 20, 21,	232, 234	tion, xi
25, 28–30, 58, 96, 105, 113,	non-analytic, 222	regular, 267
115–117, 124, 225–227, 230,	non-Boolean, 7, 80	relation, x
231, 233, 235	non-complemented, 7,	relational and, x
identity, 95, 112	80	reversal, 228, 230, 231
if, xi	non-distributive, 2, 7, 80	right distributive, 95, 96,
if and only if, xi	non-homogeneous, 159	115
image, x	non-isotropic, 105	ring of sets, xi
imaginary part, xi	non-modular, 7, 80	scalar commutative, 96
implied by, xi	non-negative, 96, 117,	self adjoint, 45, 129
implies, xi	124, 256	self-adjoint, 45, 129 , 129
implies and is implied	nonBoolean, 83	self-similar, 54, 59, 60,
by, xi	noncommutative, 42	64, 69
inclusive OR, xi	noncomplemented, 83	separable, 14, 15, 27
indicator function, x	nondegenerate, 44, 55,	set of algebras of sets, xi
injective, 114, 115	57, 58, 96, 116, 117, 124	set of tinge of sets, xi
inner-product, xi	nondistributive, 83	set of topologies, xi
inside, 212	nonmodular, 83	shift property, 261, 262
intersection, x	normal, 129, 130 , 136,	similar, 46
into, 26	137, 250, 267	space of linear trans-
inverse, 95	NOT, xi	forms, 114
inversible, 15, 32, 142,	not constant, 44	span, xi
148	not total, 23	spans, 11, 12
involutary, 126	not unique, 102	stability condition, 30,
•	-	32
involution, 148, 152	null space, x	
involutory, 108, 148–150	odd, 157, 178, 186, 230,	stable, 212
irreflexive ordering rela-	235	Strang-Fix condition,
tion, xi	only if, xi	202
isometric, 21, 29, 30,	operator norm, xi	strictly positive, 57, 116
123, 132 , 132, 136, 197, 250	order, x, xi	strong convergence, 14,
isometric in distance,	ordered pair, x	24
45, 136	orthogonal 16 16 20	subadditive, 96, 116, 117
isometric in length, 45,	orthogonal, 16 , 16, 20,	submultiplicative, 96
136	84, 129, 261	subset, x
isomorphic, 25	orthogonality, 16	summability kernel, 191
join, xi	orthomodular, 108	super set, x
least upper bound, xi	orthomodular identity,	support, 75, 77
left distributive, 95, 96,	108	surjective, 45, 136





SUBJECT INDEX page 303 Daniel J. Greenhoe

symmetric, 141, 200,	Poincaré, Jules Henri, 53	Theorem, 211
240, 257	Russull, Bertrand, vii	rotation matrix, 137
symmetric difference, x	Stravinsky, Igor, <mark>vii</mark>	rotation matrix operator, 42
symmetry, 211	Ulam, Stanislaus M., 122	rotation operator, 137
there exists, xi	von Neumann, John,	Russull, Bertrand, vii
tight frame, 33	170	sampling operator, 48, 48
time-invariant, 171, 212	raised cosine, 77	scalar commutative, 96
topology of sets, xi	raised cosine, 77	scalar product, 124
total, 13, 22, 23, 27, 28	random process, 263 , 263	scalars, 112
transitive, 108	random processes, 255	scaling, 5, 80
translation, 198, 235	random variable, 263 , 263	scaling coefficient sequence,
translation invariance,	range, x, 39	63 , 88
60	range space, 125	scaling coefficients, 251
translation invariant, 54,	rational numbers, 44	scaling filter coefficients, 251
triangle inequality 06	Rationalizing the Denomi-	scaling filters, 252
triangle inequality, 96 triangle inquality, 116	nator, 229, 230	scaling function, 54 , 54, 60,
true, x	real, 213, 240, 258	69
uniformly, 268	real linear space, 112	scaling functions, 54
union, x	real number system, 163	scaling subspace, 54 , 88
unique, 11, 14, 59, 101,	real part, xi, 149	Schauder bases, 10, 15
102, 106, 107	real-valued, 129, 200, 238,	Schauder basis, 14 , 14, 15,
unit length, 134, 136	239, 242	20, 27
unitary, 42, 43, 45, 56–	Rectangular pulse, 204	Schäffer's metric, 109
58, 65, 135 , 135–137, 197,	rectangular pulse, 203	Selberg Trace Formula, 50
250, 256, 258, 261, 262	recursive, 59	self adjoint, 45, 129
universal quantifier, xi	refinement equation, 60	self-adjoint, 45, 129 , 129, 148
vanishing moments, 90	reflection, 123	self-similar, 54, 59, 60, 64, 69
vector norm, xi	reflection operator, 137	semilinear, 149
zero at $z = -1, 71, 72$	reflexive, 108	separable, 14, 15, 27
pseudo-distributes, 112	reflexive ordering relation, xi	separable Hilbert space, 25,
PSF, 48 , 72, 73, 202	Region of Convergence, 223 ,	27–29, 58, 207
pstricks, vi	224–226	separable Hilbert spaces, 25
pulse function, 64	regular, 267	sequence, 60, 207
Pythagorean Theorem, 17,	relation, x, 40, 112	sequences, 215 set indicator, 259
19, 21, 28, 29	relational and, x relations, xi	set indicator, 255 set indicator function, 56, 74,
Pythagorean theorem, 16	function, 40	76, 204
quadrature, 86	operator, 40	set of algebras of sets, xi
Quadrature condition, 68	relation, 40	set of tings of sets, xi
quadrature condition, 85	residual spectrum, 266	set of topologies, xi
Quadrature conditions, 86	resolution, 54, 59	shift identities, 162 , 164, 165,
quadrature necessary condi-	resolvent, 147	190, 191
tions, 68	resolvent set, 266 , 267	shift operator, 134
quotes	reversal, 225, 228, 230, 231	shift property, 261, 262
Abel, Niels Henrik, 269	Reverse Triangle Inequality,	shift relation, 203, 204, 256
Cardano, Gerolamo, 95	57	similar, 46
Descartes, René, ix, 39	Riesz bases, 10	sinc, 203
Dyson, Freeman, 79	Riesz basis, 27 , 28–31, 54, 69,	sine, 153, 155
Fourier, Joseph, 1, 195	81, 258	sinus, 153
Hermite, Charles, 173	Riesz sequence, 27 , 29, 57, 69	Smith-Barnwell filter, 215
Housman, Alfred Ed-	Riesz-Fischer Theorem, 24	space
ward, vii	Riesz-Fischer Thm., 26	inner product, 124
Jacobi, Carl Gustav Ja-	right distributive, 95, 96, 115	linear, 111
cob, 183	ring, 95 , 95, 96	linear subspace, 98
Kaneyoshi, Urabe, 270	absolute value, 96	Minkowski addition, 99
Kenko, Yoshida, 270	commutative, 95	normed vector, 116
Leibniz, Gottfried, ix,	modulus, 96	orthogonal, 102
111	ring of complex square $n \times n$	vector, 111
Machiavelli, Niccolò,	matrices, 126	space of all absolutely square
269 D	ring of sets, xi	Lebesgue integrable func-
Peirce, Benjamin, 170	Robinson's Energy Delay	tions, 207



space of all absolutely square	bounded linear opera-	Hasse diagram, 53, 59
summable sequences, 207	tors, 120 , 121, 122, 124, 125,	high-pass filter, 214
space of all absolutely square	127, 128, 130, 131, 133–136	Hilbert space, 4, 20, 22,
summable sequences over ℝ,	C star algebra, 151	24, 27, 30, 32, 106, 107, 125,
48	Cardinal series, 50, 59	126, 129–131, 136, 196
space of all continuously dif-	closed interval, 223	identity, 112
ferentiable real functions,	closure, 54	identity element, 112
155	coefficient functionals,	image set, 114, 116, 127–
space of Lebesgue square-	14	131, 136
integrable functions, 48, 141	coefficients, 213	infinite sum, 9
space of linear transforms,	commutative ring, 95,	inner product, 58
114	96 , 96	inner product space, 16-
space of square integrable	complete metric space,	18, 20, 27, 102, 103, 105, 106,
functions, 5, 79	13	124
span, xi, 13, 15, 258	complete set, 13	integral domain, 95
spans, 9 , 11, 12	complex linear space,	inverse, 40, 112
spectral density function,	112	irrational numbers, 44
261	complex number sys-	isometry, 132
spectral factorization, 182	tem, 163	l.u.b., 59
spectral radius, 147	conjugate pairs, 213	lattice, 2, 97, 102
Spectral Theorem, 267	conjugate quadrature	lattice of algebras of sets,
spectrum, 147 , 266 , 267	filter, 88, 215	2
squared identities, 167	Conjugate quadrature	Laurent series, 208
stability, 212	filters, 214	least upper bound, 59,
stability condition, 30, 32	conjugate recipricol	68
stable, 212	pairs, 182	Lebesgue square-
standard inner product, 141,	continuous spectrum,	integrable functions, 39, 196,
255	266	223
standard norm, 141	convolution operator,	linear basis, 11
standard othornormal basis,	207, 208	linear combination, 9
21	coordinates, 11	linear combinations, 10
star-algebra, 126, 148 , 148	CQF, 215 , 215, 216	linear operator, 28, 30
star-algebras, 125, 126	CQF condition, 88	linear operators, 26
Stifel formula, 143	Dirac delta distribution,	linear ops., 26, 30
Strang-Fix condition, 202,	50	Linear space, 97
202	discrete-time Fourier	linear space, 1, 2, 9, 11,
Stravinsky, Igor, vii	transform, 243	58, 98–102, 112 , 112, 147
strictly positive, 57, 116	domain, 39	linear spaces, 112
strong convergence, 14, 24	eigenspace, 265	linear span, 64, 98, 99
structures	equation, 60	linear subspace, 2, 9, 97,
C* algebra, 151	expansion, 11	98 , 98–103, 105
C*-algebra, 152	Fast Wavelet Transform,	low-pass filter, 214
*-algebra, 126, 148 , 148–	53, 80	metric, 58
150, 238	field, 95, 96 , 111, 207	metric space, 1
*-algebras, 126	field of complex num-	MRA, 6, 54, 55, 61, 69,
abstract space, 1, 1	bers, 126	74–77, 80
adjoint, 126	filter banks, 214	MRA space, 54, 58, 60
algebra, 96 , 147 , 147,	Fourier Analysis, 5	MRA system, 63 , 63, 64,
148	Fourier analysis, 4, 5	66–68, 70
algebra of sets, 2	Fourier series analysis, 5	Multiresolution Analy-
algebras of sets, 3	frame, 30, 32 , 32, 36	sis, 53
analyses, 3	frames, 10	multiresolution analy-
analysis, 3, 5	function, 141, 196	sis, 6, 54 , 54, 59, 80, 82
Banach space, 14, 15	functional, 126	multiresolution analysis
basis, 5, 11–13, 20, 50,	g.l.b., 59	space, 60
51, 64, 74, 79	Gaussian Pyramid, 6, 80	multiresolution anaysis,
bijection, 25	greatest lower bound,	81
bijective, 25	54, 55, 59	multiresolution system,
Borel sets, 141, 196	group, 95 , 95	63 , 72, 81, 82
bounded linear opera-	Hamel bases, 10	N5 lattice, 2
tor, 136	Hamel basis, 11 , 11, 13	norm, <mark>58</mark>





SUBJECT INDEX Daniel J. Greenhoe page 305

normalized tight frame,	Riesz sequence, 29, 57,	underlying set, 112 unit vector, 29
normed algebra, 151,	ring, 95 , 95, 96	unital *-algebra, 148
152	ring of complex square $n \times n$ matrices, 126	unital algebra, 147 vector, 11
normed linear space, 116, 117	scalars, 112	vector, 11 vector additive identity
normed linear spaces,	scaling coefficient se-	element, 103
60, 121, 132	quence, 88	vector space, 1, 2, 96,
normed space of linear	scaling subspace 88	112
operators, 117 Null Space, 114, 115,	scaling subspace, 88 Schauder bases, 10, 15	vectors, 112 wavelet analysis, 5, 81,
129–131, 136	Schauder basis, 14, 14,	81, 82
null space, 265	15, 20, 27	wavelet coefficient se-
open interval, 223	separable Hilbert space,	quence, 82, 88
operator, 111 order relation, 97	25, 27–29, 58, 207 separable Hilbert	wavelet subspace, 88 wavelet system, 68, 78,
ordered set, 70, 83, 108	spaces, 25	82 , 82, 83, 88, 89
orthocomplemented	sequence, 60	zero, 219
lattice, 108	sequences, 215	zeros, 211, 213
orthogonal basis, 20 orthogonal comple-	Smith-Barnwell filter, 215	subadditive, 96, 116, 117 submultiplicative, 96
ment, 102, 107	space of all absolutely	subset, x, 99, 103, 106
orthomodular lattice, 4,	square Lebesgue integrable	subspace, 2, 99, 103–105, 107
108	functions, 207	subspace addition, 84
orthonormal basis, 10	space of all abso-	subspace intersection, 99
orthonormal basis, 20 , 24, 25, 27–29, 31, 250	lutely square summable sequences, 207	subspace union, 99 subspaces, 2
orthonormal MRA sys-	space of all absolutely	sum of even, 72
tem, 63	square summable sequences	sum of odd, 72
orthonormal wavelet	over R, 48	summability kernel, 191 , 191
system, 68, 84, 86 orthornormal basis, 21	space of all contin- uously differentiable real	Summation around unit circle, 34, 189
Parseval frame, 32	functions, 155	super set, x
Parseval frames, 10	space of Lebesgue	support, 70 , 70, 75, 77
Parseval's equation, 197	square-integrable functions,	support size, 70 , 70, 89
partition of unity, 74 Plancherel's formula,	48, 141 space of square inte-	surjective, xi, 45, 136 symmetric, 141, 200, 240, 257
197	grable functions, 5, 79	conjugate, 264
point spectrum, 266	span, 13, 15, 258	symmetric difference, x
pole, 219	spectral radius, 147	symmetry, 211
poles, 213	spectrum, 147 , 266 , 267	synthesis, 3 , 53
Primorial numbers, 7, 80 probability space, 263	standard othornormal basis, 21	Taylor expansion, 153
projection operator, 129	star-algebra, 126, 148	Taylor Series, 222, 222
projection operators, 4	star-algebras, 125	Taylor series, 159, 161 Taylor series for cosine, 157,
quadrature, 86	subset, 99, 103, 106	158
quadrature necessary conditions, 68	subspace, 2, 99, 103– 105, 107	Taylor series for cosine/sine,
random processes, 255	subspaces, 2	156
range, 39	support, 70	Taylor series for sine, 157 The basis problem, 14
rational numbers, 44	tight frame, 32 , 33, 35	The Book Worm, 269
real linear space, 112 real number system, 163	tight frames, 10 topological dual space,	The Fourier Series Theorem,
Region of Convergence,	121	22
223 , 224–226	topological linear space,	theorems
residual spectrum, 266	13	admissibility condition, 66 , 74, 88
resolvent set 266, 267	topological space, 1	Battle-Lemarié orthogo-
resolvent set, 266 , 267 Riesz bases, 10	topology, 2, 9, 58 total set, 13	nalization, 31
Riesz basis, 28–31, 54,	10141 001, 10	
10C3Z Dasis, Z0-31, 34,	translation operator, 50	Bessel's Equality, 17 , 18, 19, 23, 24





Bessel's inequality, 18	Leibnitz GPR, 218	sine, 156
Best Approximation	Leibniz integration rule,	Taylor series for sine,
Theorem, 18 , 18	143	157
Binomial Theorem, 162	Leibniz rule, 143 , 143	The Fourier Series Theo-
binomial theorem, 143	Mazur-Ulam theorem,	rem, 22
Cauchy-Schwarz in-	123	transversal operator in-
equality, 33	Mercer's Theorem, 268,	verses, 41
Closed Set Theorem, 104	268	trigonometric expan-
commutator relation,	Neumann Expansion	sion, 173
42 , 55, 72	Theorem, 124	trigonometric periodic-
convolution theorem, 7,	operator star-algebra,	ity, 165 , 189, 190
198 , 204, 210 , 235	126	trigonometric reduc-
CQF conditions, 89	orthonormal quadra-	tion, 178
CQF theorem, 88, 215 ,	ture conditions, 243	wavelet dilation equa-
218	parallelogram law, 107	tion, 81 , 85, 86, 89
CS Inequality, 36, 56	Parseval's Identity, 22,	there exists, xi
DeMoivre's Theorem,	24, 26, 32, 58, 261	tight frame, 32 , 33, 35
173 , 173, 175	Perfect reconstruction,	tight frames, 10
dilation equation, 60 ,	20	time-invariant, 171, 212 , 212
66–68, 70–72	Poisson Summation	topological dual space, 121
double angle formulas,	Formula, 48	topological linear space, 121
165 , 166, 167, 169, 189	polar identity, 17	topological space, 1
DTFT periodicity, 88,	product identities, 163 ,	topological space, 1 topology, 2, 9, 58
237	164, 165, 167, 187, 188	topology, 2, 3, 36 topology of sets, xi
Euler formulas, 161,		1 0.
		total, 13, 13, 22, 23, 27, 28
162, 163, 166, 167, 203	107, 108	total set, 13
Euler's Identity, 160 ,	PSF, 48 , 72, 73, 202	transform, 3 , 4, 53, 54
160, 161, 164, 169, 227	Pythagorean Theorem,	inverse Fourier, 197
Fejér-Riesz spectral fac-	17, 19, 21, 28, 29	transitive, 108
torization, 182 , 182	Pythagorean theorem,	translation, 137, 198, 224,
Fourier expansion, 22,	16	235
24 , 26	Quadrature condition,	translation invariance, 60
Fourier series expan-	68	translation invariant, 54, 55
sion, 22	quadrature condition,	translation operator, 5, 40,
Fundamental Theorem	85	40, 42, 50, 202
of Calculus, 221, 228, 236	Quadrature conditions,	translation operator adjoint,
Fundamental theorem	86	42
of calculus, 142, 143	Reverse Triangle In-	translation operator inverse,
Fundamental theorem	equality, <mark>57</mark>	41
of linear equations, 116 , 116	Riesz-Fischer Theorem,	transversal operator in-
Gelfand-Mazur Theo-	24	verses, 41
rem, 151	Riesz-Fischer Thm., 26	triangle, <mark>203</mark>
Generalized Parseval's	Robinson's Energy De-	triangle inequality, 96, 117
Identity, 22	lay Theorem, 211	triangle inquality, 116
generalized product	shift identities, 162 , 164,	trigonometric expansion,
rule, 143	165, 190, 191	173
half-angle formulas, 167	shift relation, 203, 204,	trigonometric periodicity,
Hermitian representa-	256	165 , 189, 190
tion, 150	Spectral Theorem, 267	trigonometric reduction,
Integration by Parts, 236	squared identities, 167	178
inverse DTFT, 217, 243	Stifel formula, 143	trivial linear space, 98
Inverse Fourier trans-	Strang-Fix condition,	true, x
form, 197	202	two-scale difference equa-
Inverse Poisson Sum-	Summation around unit	tion, 60
mation Formula, 49, 49	circle, 34, 189	two-scale relation, 60
IPSF, 49 , 256	support size, 70 , 70, 89	two-sided Laplace trans-
	Taylor Series, 222	form, 46
l'Hôpital's rule, 186, 249		101111, 40
Lagrange trigonometric	Taylor series, 159, 161	Ulam, Stanislaus M., 122
identities, 187	Taylor series for cosine,	underlying set, 112
Laplace convolution,	157, 158	uniformly, 268
199	Taylor series for cosine/-	union v





LICENSE Daniel J. Greenhoe page 307

floor, 70	wavelet dilation equation,
frame bound, 33, 35	81 , 85, 86, 89
frame bounds, 36	wavelet filter coefficients,
greatest lower bound, 70	251
greatest value, 71	wavelet filters, 252
least upper bound, 70	wavelet function, 81, 82
least value, 70	wavelet subspace, 88
vanishing moments, 90, 201,	wavelet system, 68, 78, 82,
245	82, 83, 85, 86, 88, 89
vector, 11	wavelet transform, 251
vector additive identity ele-	wavelets, 51
ment, 103	Weierstrass functions, 173
vector norm, <mark>xi</mark>	width, 185
vector space, 1, 2, 96, 112	
vectors, 112	
Volterra integral equation,	z transform, 211
163, 165	
Volterra integral equation of	Z-Transform, xii
the second type, 159	Z-transform, 66, 88, 239
von Neumann, John, 170	z-transform, 208 , 208, 215,
wavelet E1	216, 255
	Zak Transform, 50
	zero, 219
	zero at -1, 240
	zero at $z = -1, 71, 72$
wavelet coefficients, 251	zeros, 211, 213
	frame bound, 33, 35 frame bounds, 36 greatest lower bound, 70 greatest value, 71 least upper bound, 70 least value, 70 vanishing moments, 90, 201, 245 vector, 11 vector additive identity element, 103 vector norm, xi vector space, 1, 2, 96, 112 vectors, 112 Volterra integral equation, 163, 165 Volterra integral equation of the second type, 159

License

This document is provided under the terms of the Creative Commons license CC BY-NC-ND 4.0. For an exact statement of the license, see

```
https://creativecommons.org/licenses/by-nc-nd/4.0/legalcode
```

The icon expearing throughout this document is based on one that was once at https://creativecommons.org/

where it was stated, "Except where otherwise noted, content on this site is licensed under a Creative Commons Attribution 4.0 International license."





page 308 Daniel J. Greenhoe LICENSE





Daniel J. Greenhoe page 309 LAST PAGE



