A Book Concerning Statistical Signal Processing

VERSION 0.14X









Daniel J. Greenhoe





***Signal Processing ABCs* series**
volume 2



title:	*A Book Concerning Statistical Signal Processing*
document type:	`book`
series:	*Signal Processing ABCs*
volume:	2
author:	Daniel J. Greenhoe
version:	VERSION 0.14X
time stamp:	2019 August 16 (Friday) 11:07am UTC
copyright:	Copyright © 2019 Daniel J. Greenhoe
license:	Creative Commons license CC BY-NC-ND 4.0
typesetting engine:	XƎLaTeX
document url:	https://www.researchgate.net/project/Signal-Processing-ABCs

This text was typeset using XƎLaTeX, which is part of the TeXfamily of typesetting engines, which is arguably the greatest development since the Gutenberg Press. Graphics were rendered using the *pstricks* and related packages, and LaTeX graphics support.

The main roman, *italic*, and **bold** font typefaces used are all from the *Heuristica* family of typefaces (based on the *Utopia* typeface, released by *Adobe Systems Incorporated*). The math font is XITS from the XITS font project. The font used in quotation boxes is adapted from *Zapf Chancery Medium Italic*, originally from URW++ Design and Development Incorporated. The font used for the text in the title is Adventor (similar to *Avant-Garde*) from the *TeX-Gyre Project*. The font used for the ISBN in the footer of individual pages is LIQUID CRYSTAL (*Liquid Crystal*) from *FontLab Studio*. The Latin handwriting font is *Lavi* from the *Free Software Foundation*.

The ship appearing throughout this text is loosely based on the *Golden Hind*, a sixteenth century English galleon famous for circumnavigating the globe.[1]

[1] Paine (2000) page 63 ⟨Golden Hind⟩

❝*Here, on the level sand,*
Between the sea and land,
What shall I build or write
Against the fall of night? ❞

❝*Tell me of runes to grave*
That hold the bursting wave,
Or bastions to design
For longer date than mine. ❞

Alfred Edward Housman, English poet (1859–1936) [2]

❝*The uninitiated imagine that one must await inspiration in order to create. That is a mistake. I am far from saying that there is no such thing as inspiration; quite the opposite. It is found as a driving force in every kind of human activity, and is in no wise peculiar to artists. But that force is brought into action by an effort, and that effort is work. Just as appetite comes by eating so work brings inspiration, if inspiration is not discernible at the beginning.* ❞

Igor Fyodorovich Stravinsky (1882–1971), Russian-born composer [3]



❝*As I think about acts of integrity and grace, I realise that there is nothing in my knowledge to compare with Frege's dedication to truth. His entire life's work was on the verge of completion, much of his work had been ignored to the benefit of men infinitely less capable, his second volume was about to be published, and upon finding that his fundamental assumption was in error, he responded with intellectual pleasure clearly submerging any feelings of personal disappointment. It was almost superhuman and a telling indication of that of which men are capable if their dedication is to creative work and knowledge instead of cruder efforts to dominate and be known.* ❞

Bertrand Russell (1872–1970), British mathematician, in a 1962 November 23 letter to Dr. van Heijenoort. [4]

[2] quote: Housman (1936), page 64 ⟨"Smooth Between Sea and Land"⟩, Hardy (1940) ⟨section 7⟩
image: http://en.wikipedia.org/wiki/Image:Housman.jpg

[3] quote: Ewen (1961), page 408, Ewen (1950)
image: http://en.wikipedia.org/wiki/Image:Igor_Stravinsky.jpg

[4] quote: Heijenoort (1967), page 127
image: http://www-history.mcs.st-andrews.ac.uk/PictDisplay/Russell.html

BY-NC-ND
COPYRIGHT © 2019 DANIEL J. GREENHOE

SYMBOLS

❝ *rugula XVI. Quae vero praesentem mentis attentionem non requirunt, etiamsi ad conclusionem necessaria sint, illa melius est per brevissimas notas designare quam per integras figuras: ita enim memoria non poterit falli, nec tamen interim cogitatio distrahetur ad haec retinenda, dum aliis deducendis incumbit.* ❞

❝ *Rule XVI. As for things which do not require the immediate attention of the mind, however necessary they may be for the conclusion, it is better to represent them by very concise symbols rather than by complete figures. It will thus be impossible for our memory to go wrong, and our mind will not be distracted by having to retain these while it is taken up with deducing other matters.* ❞

René Descartes (1596–1650), French philosopher and mathematician [5]

❝ *In signs one observes an advantage in discovery which is greatest when they express the exact nature of a thing briefly and, as it were, picture it; then indeed the labor of thought is wonderfully diminished.* ❞

Gottfried Leibniz (1646–1716), German mathematician, [6]

Symbol list

symbol	description	
numbers:		
$\mathbb{Z}$	integers	$\ldots, -3, -2, -1, 0, 1, 2, 3, \ldots$
$\mathbb{W}$	whole numbers	$0, 1, 2, 3, \ldots$
$\mathbb{N}$	natural numbers	$1, 2, 3, \ldots$
$\mathbb{Z}^{\dashv}$	non-positive integers	$\ldots, -3, -2, -1, 0$

…continued on next page…

[5] quote: Descartes (1684a) ⟨rugula XVI⟩, translation: Descartes (1684b) ⟨rule XVI⟩, image: Frans Hals (circa 1650), http://en.wikipedia.org/wiki/Descartes, public domain

[6] quote: Cajori (1993) ⟨paragraph 540⟩, image: http://en.wikipedia.org/wiki/File:Gottfried_Wilhelm_von_Leibniz.jpg, public domain

symbol	description	
$\mathbb{Z}^-$	negative integers	$\ldots, -3, -2, -1$
$\mathbb{Z}_o$	odd integers	$\ldots, -3, -1, 1, 3, \ldots$
$\mathbb{Z}_e$	even integers	$\ldots, -4, -2, 0, 2, 4, \ldots$
$\mathbb{Q}$	rational numbers	$\frac{m}{n}$ with $m \in \mathbb{Z}$ and $n \in \mathbb{Z}\backslash 0$
$\mathbb{R}$	real numbers	completion of $\mathbb{Q}$
$\mathbb{R}^{\vdash}$	non-negative real numbers	$[0, \infty)$
$\mathbb{R}^{\dashv}$	non-positive real numbers	$(-\infty, 0]$
$\mathbb{R}^+$	positive real numbers	$(0, \infty)$
$\mathbb{R}^-$	negative real numbers	$(-\infty, 0)$
$\mathbb{R}^*$	extended real numbers	$\mathbb{R}^* \triangleq \mathbb{R} \cup \{-\infty, \infty\}$
$\mathbb{C}$	complex numbers	
$\mathbb{F}$	arbitrary field	(often either $\mathbb{R}$ or $\mathbb{C}$)
∞	positive infinity	
$-\infty$	negative infinity	
π	pi	$3.14159265\ldots$
relations:		
®	relation	
$\circledwedge$	relational and	
$X \times Y$	Cartesian product of X and Y	
$(\triangle, \triangledown)$	ordered pair	
$\lvert z \rvert$	absolute value of a complex number z	
$=$	equality relation	
$\triangleq$	equality by definition	
$\rightarrow$	maps to	
$\in$	is an element of	
$\notin$	is not an element of	
$\mathcal{D}(®)$	domain of a relation ®	
$\mathcal{I}(®)$	image of a relation ®	
$\mathcal{R}(®)$	range of a relation ®	
$\mathcal{N}(®)$	null space of a relation ®	
set relations:		
$\subseteq$	subset	
$\subsetneq$	proper subset	
$\supseteq$	super set	
$\supsetneq$	proper superset	
$\nsubseteq$	is not a subset of	
$\not\subset$	is not a proper subset of	
operations on sets:		
$A \cup B$	set union	
$A \cap B$	set intersection	
$A \triangle B$	set symmetric difference	
$A \backslash B$	set difference	
A^c	set complement	
$\lvert \cdot \rvert$	set order	
$\mathbb{1}_A(x)$	set indicator function or characteristic function	
logic:		
1	"true" condition	
0	"false" condition	
$\neg$	logical NOT operation	

...continued on next page...

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

symbol	description	
$\wedge$	logical AND operation	
$\vee$	logical inclusive OR operation	
$\oplus$	logical exclusive OR operation	
$\Longrightarrow$	"implies";	"only if"
$\Longleftarrow$	"implied by";	"if"
$\Longleftrightarrow$	"if and only if";	"implies and is implied by"
$\forall$	universal quantifier:	"for each"
$\exists$	existential quantifier:	"there exists"
order on sets:		
$\vee$	join or least upper bound	
$\wedge$	meet or greatest lower bound	
$\leq$	reflexive ordering relation	"less than or equal to"
$\geq$	reflexive ordering relation	"greater than or equal to"
$<$	irreflexive ordering relation	"less than"
$>$	irreflexive ordering relation	"greater than"
measures on sets:		
$\lvert X \rvert$	order or counting measure of a set X	
distance spaces:		
d	metric or distance function	
linear spaces:		
$\lVert\cdot\rVert$	vector norm	
$\lvert\lvert\lvert\cdot\rvert\rvert\rvert$	operator norm	
$\langle\triangle \mid \triangledown\rangle$	inner-product	
$\mathsf{span}(\boldsymbol{V})$	span of a linear space $\boldsymbol{V}$	
algebras:		
$\Re$	real part of an element in a $*$-algebra	
$\Im$	imaginary part of an element in a $*$-algebra	
set structures:		
$\boldsymbol{T}$	a topology of sets	
$\boldsymbol{R}$	a ring of sets	
$\boldsymbol{A}$	an algebra of sets	
$\varnothing$	empty set	
2^X	power set on a set X	
sets of set structures:		
$\mathcal{T}(X)$	set of topologies on a set X	
$\mathcal{R}(X)$	set of rings of sets on a set X	
$\mathcal{A}(X)$	set of algebras of sets on a set X	
classes of relations/functions/operators:		
2^{XY}	set of *relations* from X to Y	
Y^X	set of *functions* from X to Y	
$\mathcal{S}_{\mathsf{j}}(X,Y)$	set of *surjective* functions from X to Y	
$\mathcal{I}_{\mathsf{j}}(X,Y)$	set of *injective* functions from X to Y	
$\mathcal{B}_{\mathsf{j}}(X,Y)$	set of *bijective* functions from X to Y	
$\mathcal{B}(\boldsymbol{X},\boldsymbol{Y})$	set of *bounded* functions/operators from $\boldsymbol{X}$ to $\boldsymbol{Y}$	
$\mathcal{L}(\boldsymbol{X},\boldsymbol{Y})$	set of *linear bounded* functions/operators from $\boldsymbol{X}$ to $\boldsymbol{Y}$	
$\mathcal{C}(\boldsymbol{X},\boldsymbol{Y})$	set of *continuous* functions/operators from $\boldsymbol{X}$ to $\boldsymbol{Y}$	
specific transforms/operators:		
$\tilde{\mathbf{F}}$	*Fourier Transform* operator (Definition O.2 page 333)	
$\hat{\mathbf{F}}$	*Fourier Series* operator	

...continued on next page...

COPYRIGHT © 2019 DANIEL J. GREENHOE

symbol	description
$\check{\mathbf{F}}$	*Discrete Time Fourier Series* operator (Definition P.1 page 343)
$\mathbf{Z}$	*Z-Transform* operator (Definition Q.4 page 354)
$\tilde{\mathsf{f}}(\omega)$	*Fourier Transform* of a function $\mathsf{f}(x) \in \boldsymbol{L}^2_{\mathbb{R}}$
$\check{\mathsf{x}}(\omega)$	*Discrete Time Fourier Transform* of a sequence $(\!(x_n \in \mathbb{C})\!)_{n\in\mathbb{Z}}$
$\check{\mathsf{x}}(z)$	*Z-Transform* of a sequence $(\!(x_n \in \mathbb{C})\!)_{n\in\mathbb{Z}}$

BY-NC-ND
COPYRIGHT © 2019 DANIEL J. GREENHOE

SYMBOL INDEX

$\bar{\delta}_n$, 267
$(\boldsymbol{A}, \|\cdot\|, *)$, 253
$\|\cdot\|$, 271
$\perp$, 216
$\star$, 353
σ, 367
$\tilde{\mathbf{F}}$, 334
$|\boldsymbol{x}|$, 214
$\boldsymbol{x}^-$, 214
$\boldsymbol{x}^+$, 214
d, 217
$\mathbf{I}_{\mathsf{m}}$, 251
$\mathbf{I}$, 304
$\mathbf{R}_{\mathsf{e}}$, 251
$\mathbf{Z}$, 354
$\mathsf{N}_n(x)$, 367
$\mathsf{epi}(\mathsf{f})$, 283
$\mathsf{hyp}(\mathsf{f})$, 283
span, 219
$*$, 250
$|||\cdot|||$, 308
$\textcircled{\perp}$, 278, 280
$\star$, 336
$\mathcal{B}(\boldsymbol{X}, \boldsymbol{Y})$, 311
$(\cdot : \cdot)$, 281, 282
$(\cdot : \cdot]$, 281, 282
$[\cdot : \cdot)$, 281, 282
$[\cdot : \cdot]$, 281, 282
$\boldsymbol{Y}^{\boldsymbol{X}}$, 304
ρ, 190, 249
σ_{c}, 190
σ_{p}, 190
σ_{r}, 190
σ, 190, 249
r, 249

Copyright © 2019 Daniel J. Greenhoe

CONTENTS

Title page . . . v
Typesetting . . . vi
Quotes . . . vii
Symbol list . . . ix
Symbol index . . . xiii
Contents . . . xv

I Statistical Analysis **1**

1 Expectation operator **3**
1.1 Definitions . . . 3
1.2 Expectation as a linear operator . . . 4
1.3 Expectation inequalities . . . 6
1.4 Joint and conditional probability spaces . . . 7
1.5 Expectation inner product space . . . 8

2 Random Sequences **11**
2.1 Definitions . . . 11
2.2 Properties . . . 12
2.3 Wide Sense Stationary processes . . . 12
2.4 Spectral density . . . 14
2.5 Spectral Power . . . 16

3 Continuous Random Processes **17**
3.1 Definitions . . . 17
3.2 Eigen-analysis of random processes . . . 18
3.3 Properties . . . 20

II Statistical Processing **23**

4 Operations on Random Variables **25**
4.1 Functions of one random variable . . . 25
4.2 Functions of two random variables . . . 31

5 Operators on Discrete Random Sequences **35**
5.1 LTI operators on random sequences . . . 35
5.2 LTI operators on WSS random sequences . . . 36
5.3 Parallel operators on WSS random sequences . . . 37
5.4 Whitening discrete random sequences . . . 38

6 Operators on Continuous Random Sequences **41**
6.1 LTI Operations on non-stationary random processes . . . 41
6.2 LTI Operations on WSS random processes . . . 43
6.3 Whitening continuous random sequences . . . 48

7 Additive noise on random sequences 51
7.1 Additive noise and correlation . . . 51
7.2 Additive noise and operators . . . 53
7.3 Additive noise and LTI operators . . . 56
7.4 Additive noise and dual operators . . . 58

III Statistical Estimation 61

8 Estimation Overview 63
8.1 Estimation types . . . 63
8.2 Estimation criterion . . . 64
8.3 Measures of estimator quality . . . 65
8.4 Estimation techniques . . . 66
8.5 Sequential decoding . . . 67

9 Norm Minimization 69
9.1 Minimum mean square estimation . . . 70
9.2 Least squares . . . 72

10 Gradient Search Techniques 75
10.1 Gradient search techniques . . . 75
10.2 Direct search . . . 77

11 KL-expansion application 79
11.1 Sufficient statistics . . . 79
11.2 Optimal symbol estimation . . . 84
11.3 Colored noise . . . 91
11.4 Signal matching . . . 92

12 Moment Estimation 95
12.1 Mean Estimation . . . 95
12.2 Variance Estimation . . . 98
12.3 Estimates in terms of moment estimates . . . 101

13 Correlation Estimation 103

14 Spectral Estimation 105

15 Density Estimation 107

16 System Identification 109
16.1 Estimation techniques . . . 109
16.2 Additive noise system models . . . 110
16.3 Transfer function estimate definitions and interpretation . . . 111
16.4 Estimator relationships . . . 116
16.5 Alternate forms . . . 121
16.6 Least squares estimates of non-linear systems . . . 123
16.7 Least squares estimates of linear systems . . . 127
16.8 Coherence . . . 131
16.8.1 Application . . . 131
16.8.2 Definitions . . . 132
16.8.3 A warning . . . 132

17 Estimating Noise 135

IV Statistical Detection 137

18 Communication channels 139
18.1 System model . . . 139
18.1.1 Channel operator . . . 140
18.1.2 Receive operator . . . 140

BY-NC-ND Copyright © 2019 Daniel J. Greenhoe

18.2 Optimization in the case of additional operations . . . 141
18.3 Channel Statistics . . . 142

19 Optimal Symbol Detection **143**
19.1 ML Estimation . . . 143
19.2 Generalized coherent modulation . . . 144
19.3 Frequency Shift Keying (FSK) . . . 145
19.4 Quadrature Amplitude Modulation (QAM) . . . 147
19.4.1 Receiver statistics . . . 147
19.4.2 Detection . . . 148
19.4.3 Probability of error . . . 148
19.5 Phase Shift Keying (PSK) . . . 149
19.5.1 Receiver statistics . . . 149
19.5.2 Detection . . . 151
19.5.3 Probability of error . . . 151
19.6 Pulse Amplitude Modulation (PAM) . . . 152
19.6.1 Receiver statistics . . . 152
19.6.2 Detection . . . 152
19.6.3 Probability of error . . . 153

20 Network Detection **155**
20.1 Detection . . . 155
20.2 Bayesian Estimation . . . 155
20.3 Joint Gaussian Model . . . 156
20.4 2 hypothesis, 2 sensor detection . . . 157

V Appendices **161**

A Probability Space **163**
A.1 Probability functions . . . 164
A.2 Probability Space . . . 165
A.3 Properties . . . 166
A.4 Examples . . . 168
A.5 Probability subspaces . . . 171

B Probability Density Functions **173**
B.1 Random variables . . . 173
B.2 Probability distributions . . . 174
B.3 Properties . . . 174

C Some Probability Density Functions **179**
C.1 Discrete distributions . . . 179
C.2 Continuous distributions . . . 179
C.2.1 Uniform distribution . . . 179
C.2.2 Gaussian distribution . . . 180
C.2.3 Gamma distribution . . . 181
C.2.4 Chi-squared distributions . . . 182
C.2.5 Radial distributions . . . 185
C.3 Joint Gaussian distributions . . . 186

D Spectral Theory **189**
D.1 Operator Spectrum . . . 189
D.2 Fredholm kernels . . . 191

E Matrix Calculus **193**
E.1 First derivative of a vector with respect to a vector . . . 193
E.2 First derivative of a matrix with respect to a scalar . . . 200
E.3 Second derivative of a scalar with respect to a vector . . . 202
E.4 Multiple derivatives of a vector with respect to a scalar . . . 203

F Algebraic structures **205**

Copyright © 2019 Daniel J. Greenhoe
BY-NC-ND

G Linear spaces 207
G.1 Definition and basic results . . . 207
G.2 Order on Linear Spaces . . . 212

H Linear Combinations 219
H.1 Linear combinations in linear spaces . . . 219
H.2 Bases in topological linear spaces . . . 223
H.3 Schauder bases in Banach spaces . . . 224
H.4 Linear combinations in inner product spaces . . . 226
H.5 Orthonormal bases in Hilbert spaces . . . 229
H.6 Riesz bases in Hilbert spaces . . . 237
H.7 Frames in Hilbert spaces . . . 242

I Normed Algebras 249
I.1 Algebras . . . 249
I.2 Star-Algebras . . . 250
I.3 Normed Algebras . . . 253
I.4 C* Algebras . . . 253

J Inner Product Spaces 255
J.1 Definition and basic results . . . 255
J.2 Relationship between norms and inner products . . . 259
J.2.1 Norms induced by inner products . . . 259
J.2.2 Inner products induced by norms . . . 261
J.3 Orthogonality . . . 267

K Normed Linear Spaces 271
K.1 Definition and basic results . . . 271
K.2 Relationship between metrics and norms . . . 273
K.2.1 Metrics generated by norms . . . 273
K.2.2 Norms generated by metrics . . . 276
K.3 Orthogonality on normed linear spaces . . . 278

L Intervals and Convexity 281
L.1 Intervals . . . 281
L.2 Convex sets . . . 282
L.3 Convex functions . . . 283
L.4 Literature . . . 286

M Finite Sums 289
M.1 Summation . . . 289
M.2 Means . . . 290
M.2.1 Weighted ϕ-means . . . 290
M.2.2 Power means . . . 292
M.3 Inequalities on power means . . . 295
M.4 Power Sums . . . 300

N Operators on Linear Spaces 303
N.1 Operators on linear spaces . . . 303
N.1.1 Operator Algebra . . . 303
N.1.2 Linear operators . . . 304
N.2 Operators on Normed linear spaces . . . 308
N.2.1 Operator norm . . . 308
N.2.2 Bounded linear operators . . . 311
N.2.3 Adjoints on normed linear spaces . . . 313
N.2.4 More properties . . . 314
N.3 Operators on Inner product spaces . . . 315
N.3.1 General Results . . . 315
N.3.2 Operator adjoint . . . 316
N.4 Special Classes of Operators . . . 318
N.4.1 Projection operators . . . 318
N.4.2 Self Adjoint Operators . . . 320

BY-NC-ND
Copyright © 2019 Daniel J. Greenhoe

N.4.3 Normal Operators . . . 321
N.4.4 Isometric operators . . . 323
N.4.5 Unitary operators . . . 326
N.5 Operator order . . . 331

O Fourier Transform **333**
O.1 Definitions . . . 333
O.2 Operator properties . . . 334
O.3 Convolution . . . 336
O.4 Real valued functions . . . 337
O.5 Moment properties . . . 338
O.6 Examples . . . 340

P Discrete Time Fourier Transform **343**
P.1 Definition . . . 343
P.2 Properties . . . 343
P.3 Derivatives . . . 351

Q Operations on Sequences **353**
Q.1 Convolution operator . . . 353
Q.2 Z-transform . . . 354
Q.3 From z-domain back to time-domain . . . 356
Q.4 Zero locations . . . 357
Q.5 Pole locations . . . 358
Q.6 Mirroring for real coefficients . . . 359
Q.7 Rational polynomial operators . . . 360
Q.8 Filter Banks . . . 361
Q.9 Inverting non-minimum phase filters . . . 365

R B-Splines **367**
R.1 Definitions . . . 367
R.2 Algebraic properties . . . 371
R.3 Projection properties . . . 380
R.4 Fourier analysis . . . 381
R.5 Basis properties . . . 383
R.5.1 Uniqueness properties . . . 383
R.5.2 Partition of unity properties . . . 384
R.5.3 Riesz basis properties . . . 385
R.6 Mutiresolution properties . . . 391
R.6.1 Introduction . . . 391
R.6.2 B-spline dyadic decomposition . . . 391
R.6.3 B-spline MRA scaling functions . . . 393
R.6.4 B-spline MRA coefficient sequences . . . 394

S Source Code **399**

Back Matter **403**
References . . . 404
Reference Index . . . 431
Subject Index . . . 435
License . . . 452
End of document . . . 453

Copyright © 2019 Daniel J. Greenhoe
BY-NC-ND

Copyright © 2019 Daniel J. Greenhoe

Part I

Statistical Analysis

CHAPTER 1

EXPECTATION OPERATOR

1.1 Definitions

In a *probability space* $(\Omega, \mathbb{E}, \mathsf{P})$ (Definition A.2 page 165), all probability information is contained in the *measure* P. Often times this information is overwhelming and a simpler statistic, which does not offer so much information, is sufficient. Some of the most common statistics can be conveniently expressed in terms of the *expectation operator* E.

Definition 1.1. *Let* $(\Omega, \mathbb{E}, \mathsf{P})$ *be a* PROBABILITY SPACE *(Definition A.2 page 165) and* X *a* RANDOM VARIABLE *(Definition ?? page ??) on* $(\Omega, \mathbb{E}, \mathsf{P})$ *with* PROBABILITY DENSITY FUNCTION p_{X}.

DEF *The* ***expectation operator*** E_{X} *on* X *is defined as*

$$\mathsf{E}_{\mathsf{X}}\mathsf{X} \triangleq \int_{x \in \mathbb{F}} x \mathsf{p}_{\mathsf{X}}(x)\, \mathrm{d}x.$$

We already said that a *random variable* X is neither random nor a variable, but is rather a function of an underlying process that does appear to be random. However, because it is a function of a process that does appear random, the *random variable* X also appears to be random. That is, if we don't know the outcome of of the underlying experimental process, then we also don't know for sure what X is, and so X does indeed appear to be random. However, eventhough X appears to be random, the expected value $\mathsf{E}_{\mathsf{X}}\mathsf{X}$ of X is **not random**. Rather it is a fixed value (like 0 or 7.9 or −2.6).

On the other hand, eventhough $\mathsf{E}\mathsf{X}$ is **not random**, note that $\mathsf{E}(\mathsf{X}|\mathsf{Y})$ **is random**. This is because $\mathsf{E}(\mathsf{X}|\mathsf{Y})$ is a function of Y. That is, once we know that Y equals some fixed value y (like 0 or 2.7 or −5.1) then $\mathsf{E}(\mathsf{X}|\mathsf{Y} = y)$ is also fixed. However, if we don't know the value of Y, then Y is still a *random variable* and the expression $\mathsf{E}(\mathsf{X}|\mathsf{Y})$ is also random (a function of *random variable* Y).

Two common statistics that are conveniently expressed in terms of the expectation operator are the *mean* and *variance*. The mean is an indicator of the "middle" of a probability distribution and the variance is an indicator of the "spread".

Definition 1.2. *Let* X *be a* RANDOM VARIABLE *on the* PROBABILITY SPACE $(\Omega, \mathbb{E}, \mathsf{P})$.

DEF
(1). *The* ***mean*** μ_{X} *of* X *is* $\quad \mu_{\mathsf{X}} \triangleq \mathsf{E}_{\mathsf{X}}\mathsf{X}$

(2). *The* ***variance*** $\mathsf{Var}(\mathsf{X})$ *or* σ^2_{X} *of* X *is* $\quad \mathsf{Var}(\mathsf{X}) \triangleq \mathsf{E}_{\mathsf{X}}\left[(\mathsf{X} - \mathsf{E}_{\mathsf{X}}\mathsf{X})^2\right]$

1.2 Expectation as a linear operator

The next theorem demonstrates that the operator E is a *linear operator* (Definition N.3 page 304)—which in turn makes E part of a distinguished club of operators along with fellow member operators differentiation $\frac{\mathsf{d}}{\mathsf{d}x}$, integration $\int \mathsf{d}x$, Laplace $\mathbf{L}$, Fourier $\tilde{\mathbf{F}}$, z-transform $\mathbf{Z}$, etc. Because E is a linear operator, it immediately inherits all the properties that it's linear operator birthright grants it (Corollary 1.1 page 4).

Theorem 1.1 (Linearity of E). [1] *Let* X *be a* RANDOM VARIABLE *on a* PROBABILITY SPACE $(\Omega, \mathbb{E}, \mathsf{P})$.

THM

$$\mathsf{E}_{\mathsf{x}}(a\mathsf{X} + b\mathsf{Y} + c) = \left(a\mathsf{E}_{\mathsf{x}}\mathsf{X}\right) + \left(b\mathsf{E}_{\mathsf{x}}\mathsf{Y}\right) + c \quad \forall_{a,b,c\in\mathbb{R}} \quad \text{(LINEAR)}$$

✎PROOF:

$$\begin{aligned}
\mathsf{E}_{\mathsf{xy}}(a\mathsf{X} + b\mathsf{Y} + c) &\triangleq \int_{x\in\mathbb{R}}\int_{y\in\mathbb{R}} [ax + by + c]\mathsf{p}_{\mathsf{xy}}(x,y)\,\mathsf{d}y\,\mathsf{d}x \qquad \text{by definition of } \mathsf{E} \text{ (Definition 1.1 page 3)} \\
&= \int_{x\in\mathbb{R}}\int_{y\in\mathbb{R}} ax\mathsf{p}_{\mathsf{xy}}(x,y)\,\mathsf{d}y\,\mathsf{d}x + \int_{x\in\mathbb{R}}\int_{y\in\mathbb{R}} by\mathsf{p}_{\mathsf{xy}}(x,y)\,\mathsf{d}y\,\mathsf{d}x + \int_{x\in\mathbb{R}}\int_{y\in\mathbb{R}} c\mathsf{p}_{\mathsf{xy}}(x,y)\,\mathsf{d}y\,\mathsf{d}x \\
&= \int_{x\in\mathbb{R}} ax \underbrace{\int_{y\in\mathbb{R}} \mathsf{p}_{\mathsf{xy}}(x,y)\,\mathsf{d}y}_{\mathsf{p}_{\mathsf{x}}(x)}\,\mathsf{d}x + \int_{y\in\mathbb{R}} by \underbrace{\int_{x\in\mathbb{R}} \mathsf{p}_{\mathsf{xy}}(x,y)\,\mathsf{d}x}_{\mathsf{p}_{\mathsf{y}}(y)}\,\mathsf{d}y + c\underbrace{\int_{y\in\mathbb{R}}\int_{x\in\mathbb{R}} \mathsf{p}_{\mathsf{xy}}(x,y)\,\mathsf{d}x\,\mathsf{d}y}_{1} \\
&= a\underbrace{\int_{x\in\mathbb{R}} x\mathsf{p}_{\mathsf{x}}(x)\,\mathsf{d}x}_{\mathsf{EX}} + b\underbrace{\int_{y\in\mathbb{R}} y\mathsf{p}_{\mathsf{y}}(y)\,\mathsf{d}y}_{\mathsf{EY}} + c \\
&= \left(a\mathsf{E}_{\mathsf{x}}\mathsf{X}\right) + \left(b\mathsf{E}_{\mathsf{y}}\mathsf{Y}\right) + c
\end{aligned}$$

✏

Corollary 1.1. *Let* E *be the* EXPECTATION OPERATOR *over a* PROBABILITY SPACE $(\Omega, \mathbb{E}, \mathsf{P})$. *Let* $spLLF$ *be a* VECTOR SPACE OF RANDOM VARIABLES *over* $(\Omega, \mathbb{E}, \mathsf{P})$.

COR

(1). $\mathsf{E}\mathbb{0} = \mathbb{0}$ *and*

(2). $\mathsf{E}(-\mathsf{X}) = -(\mathsf{EX}) \quad \forall_{\mathsf{X}\in L^2_{\mathbb{F}}}$ *and*

(3). $\mathsf{E}(\mathsf{X} - \mathsf{Y}) = \mathsf{EX} - \mathsf{EY} \quad \forall_{\mathsf{X},\mathsf{Y}\in L^2_{\mathbb{F}}}$ *and*

(4). $\mathsf{E}\left(\sum_{n=1}^{N} \alpha_n \mathsf{X}_n\right) = \sum_{n=1}^{N} \alpha_n\left(\mathsf{EX}_n\right) \quad \forall_{\alpha_n\in\mathbb{F}}, \quad \forall_{\mathsf{X}\in L^2_{\mathbb{F}}}$

✎PROOF: These all follow immediately from the fact that E is a *linear operator* and from Theorem N.1 (page 304). ✏

Remark 1.1. Projecting a stochastic process onto a basis often yields valuable insights into the nature of the underlying data. Typical projection operators include the Fourier operator $\tilde{\mathbf{F}}$, Laplace $\mathbf{L}$, and z-transform $\mathbf{Z}$ ...not to mention wavelet operators. But note that any such projection on a random sequence simply produces another random sequence. For example, the Fourier transform $\tilde{\mathbf{F}}\mathsf{x}(n)$ of a random sequence $\mathsf{x}(n)$ is another random sequence.

One way to overcome this difficulty is to simply invoke the *sampling* operator $\mathbf{S}\mathsf{x}(n)$ (CHAPTER **??** page **??**), yielding a deterministic sequence, and then take the Fourier transform of the resulting deterministic sequence. The problem here is that every time you resample the sequence, you will very likely get a different Fourier transform.

[1] Wilks (1963b), page 73 ⟨§3.2 "Mean value of a random variable"⟩

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

Arguably a better approach (and the standard one at that) is to first invoke the expectation operator $\mathsf{Ex}(n)$, also yielding a deterministic sequence.

The good news here is that because E and all the above mentioned operators are *linear*, we can do all the standard arithmetic acrobatics associated with linear algebra operators (next corollary).

Corollary 1.2. *Let* $\mathbf{M}$ *and* $\mathbf{N}$ *be* LINEAR OPERATORS *(Definition N.3 page 304)*.

COR					
1.	$\mathsf{E}(\mathbf{MN})$	$=$	$(\mathsf{E}\mathbf{M})\mathbf{N}$	$\forall \mathsf{E}\in\mathcal{L}(\boldsymbol{Z},\boldsymbol{W}), \mathbf{M}\in\mathcal{L}(\boldsymbol{Y},\boldsymbol{Z}), \mathbf{N}\in\mathcal{L}(\boldsymbol{X},\boldsymbol{Y})$	(ASSOCIATIVE)
2.	$\mathsf{E}(\mathbf{M}\mathbin{\mathring{+}}\mathbf{N})$	$=$	$(\mathsf{E}\mathbf{M})\mathbin{\mathring{+}}(\mathsf{E}\mathbf{N})$	$\forall \mathsf{E}\in\mathcal{L}(\boldsymbol{Y},\boldsymbol{Z}), \mathbf{M}\in\mathcal{L}(\boldsymbol{X},\boldsymbol{Y}), \mathbf{N}\in\mathcal{L}(\boldsymbol{X},\boldsymbol{Y})$	(LEFT DISTRIBUTIVE)
3.	$(\mathsf{E}\mathbin{\mathring{+}}\mathbf{M})\mathbf{N}$	$=$	$(\mathsf{E}\mathbf{N})\mathbin{\mathring{+}}(\mathbf{MN})$	$\forall \mathsf{E}\in\mathcal{L}(\boldsymbol{Y},\boldsymbol{Z}), \mathbf{M}\in\mathcal{L}(\boldsymbol{Y},\boldsymbol{Z}), \mathbf{N}\in\mathcal{L}(\boldsymbol{X},\boldsymbol{Y})$	(RIGHT DISTRIBUTIVE)
4.	$\alpha(\mathsf{E}\mathbf{M})$	$=$	$(\alpha\mathsf{E})\mathbf{M} = \mathsf{E}(\alpha\mathbf{M})$	$\forall \mathsf{E}\in\mathcal{L}(\boldsymbol{Y},\boldsymbol{Z}), \mathbf{M}\in\mathcal{L}(\boldsymbol{X},\boldsymbol{Y}), \alpha\in\mathbb{F}$	(HOMOGENEOUS)

✎PROOF: These all follow immediately from the fact that E is a *linear operator* and from Theorem N.4 (page 307). ✏

Corollary 1.3. *Let* X *be a* RANDOM VARIABLE *on a* PROBABILITY SPACE $(\Omega, \mathbb{E}, \mathsf{P})$.

COR				
	$\mathsf{Var}(a\mathsf{X}+b)$	$=$	$a^2\mathsf{Var}(\mathsf{X})$	$\forall a,b\in\mathbb{R}$
	$\mathsf{Var}(\mathsf{X})$	$=$	$\mathsf{E}_{\mathsf{x}}(\mathsf{X}^2) - (\mathsf{E}_{\mathsf{x}}\mathsf{X})^2$	

✎PROOF:

$$
\begin{aligned}
\mathsf{Var}(\mathsf{X}) &\triangleq \mathsf{E}_{\mathsf{x}}\left[(\mathsf{X}-\mathsf{E}_{\mathsf{x}}\mathsf{X})^2\right] && \text{by definition of } \mathsf{Var} && \text{(Definition 1.2 page 3)}\\
&= \mathsf{E}_{\mathsf{x}}\left[\mathsf{X}^2 - 2\mathsf{X}\mathsf{E}_{\mathsf{x}}\mathsf{X} + (\mathsf{E}_{\mathsf{x}}\mathsf{X})^2\right] && \text{by } \textit{Binomial Theorem} && \\
&= \mathsf{E}_{\mathsf{x}}\mathsf{X}^2 - \mathsf{E}_{\mathsf{x}}\left[2\mathsf{X}\mathsf{E}_{\mathsf{x}}\mathsf{X}\right] + \mathsf{E}_{\mathsf{x}}(\mathsf{E}_{\mathsf{x}}\mathsf{X})^2 && \text{by } \textit{linearity} \text{ of } \mathsf{E} && \text{(Theorem 1.1 page 4)}\\
&= \mathsf{E}_{\mathsf{x}}\mathsf{X}^2 - 2(\mathsf{E}_{\mathsf{x}}\mathsf{X})[\mathsf{E}_{\mathsf{x}}\mathsf{X}] + (\mathsf{E}_{\mathsf{x}}\mathsf{X})^2 && && \\
&= \mathsf{E}_{\mathsf{x}}(\mathsf{X}^2) - (\mathsf{E}_{\mathsf{x}}\mathsf{X})^2 && && \\
\mathsf{Var}(a\mathsf{X}+b) &= \mathsf{E}_{\mathsf{x}}(a\mathsf{X}+b)^2 - [\mathsf{E}_{\mathsf{x}}(a\mathsf{X}+b)]^2 && && \\
&= \mathsf{E}_{\mathsf{x}}(a^2\mathsf{X}^2 + 2ab\mathsf{X} + b^2) - [a(\mathsf{E}_{\mathsf{x}}\mathsf{X}) + b]^2 && && \\
&= a^2\mathsf{E}_{\mathsf{x}}\mathsf{X}^2 + 2ab\mathsf{E}_{\mathsf{x}}\mathsf{X} + b^2 - \left[a^2[\mathsf{E}_{\mathsf{x}}\mathsf{X}]^2 + 2ab\mathsf{E}_{\mathsf{x}}\mathsf{X} + b^2\right] && \text{by } \textit{linearity} \text{ of } \mathsf{E} && \text{(Theorem 1.1 page 4)}\\
&= a^2\left[\mathsf{E}_{\mathsf{x}}\mathsf{X}^2 - (\mathsf{E}_{\mathsf{x}}\mathsf{X})^2\right] && && \\
&\triangleq a^2\mathsf{Var}(\mathsf{X}) && \text{by previous result} &&
\end{aligned}
$$

✏



Figure 1.1: Jensen's inequality

Jensen's inequality is an extremely useful application of *convex*ity (Definition L.9 page 284) to the *expectation* operator. Jensen's inequality is stated in Corollary 1.4 (next) and illustrated in Figure 1.1 (page 5).

COPYRIGHT © 2019 DANIEL J. GREENHOE

Corollary 1.4 (Jensen's inequality). [2] *Let* f *be a function in* $\mathbb{R}^{\mathbb{R}}$ *and* X *be a* RANDOM VARIABLE *on* $(\Omega, \mathbb{E}, \mathsf{P})$.

COR $\{\mathsf{f}$ *is* CONVEX$\} \quad \Longrightarrow \quad \{\mathsf{f}(\mathsf{EX}) \leq \mathsf{Ef}(\mathsf{X})\}$

✎PROOF:

1. Proof 1: Let $mx + c$ be a "support line" under $\mathsf{f}(x)$ (Figure 1.1 page 5) such that

$$\begin{aligned} mx + c &< \mathsf{f}(x) \quad \text{for } x \neq \mathsf{EX} \\ mx + c &= \mathsf{f}(x) \quad \text{for } x = \mathsf{EX}. \end{aligned}$$

   Then

$$\begin{aligned} \mathsf{f}(\mathsf{EX}) &= m[\mathsf{EX}] + c \\ &= \mathsf{E}[mX + c] \\ &\leq \mathsf{Ef}(\mathsf{X}) \end{aligned}$$

2. Proof 2 (alternate proof):

$$\begin{aligned} \mathsf{f}(\mathsf{EX}) &\triangleq \mathsf{f}\left(\sum_{x\in\mathbb{E}} x\mathsf{P}(x)\right) \\ &\leq \sum_{x\in\mathbb{E}} \mathsf{f}(x)\mathsf{P}(x) \qquad \text{by } \textit{Jensen's inequality} \text{ for convex sets} \qquad \text{(Theorem L.1 page 284)} \end{aligned}$$

✏

Example 1.1. [3] Some examples of *Jensen's Inequality* (Corollary 1.4 page 6) applied to the *expectation operator* are the following:

EX
$$\begin{aligned} (\mathsf{EX})^{-1} &< \mathsf{E}\left(\mathsf{X}^{-1}\right) \\ \mathsf{E}\left(\log \mathsf{X}\right) &< \log(\mathsf{EX}) \\ e^{-\mathsf{EX}} &\leq \mathsf{E}\left[e^{-\mathsf{X}}\right] \end{aligned}$$

Theorem 1.2 (Law of the Unconscious Statistician). [4]

THM $\mathsf{E}\left[\mathsf{g}(\mathsf{X})\right] = \int_{x\in\mathbb{R}} \mathsf{g}(x)\mathsf{p}_{\mathsf{x}}(x)\, \mathrm{d}x$

1.3 Expectation inequalities

Theorem 1.3 (Markov's inequality). [5] *Let* $\mathsf{X} : \Omega \to [0, \infty)$ *be a non-negative valued* RANDOM VARIABLE *and* $a \in (0, \infty)$. *Then*

THM $\mathsf{P}\{\mathsf{X} \geq a\} \leq \frac{1}{a}\mathsf{EX}$

[2] Shao (2003) page 31 ⟨"1.3 Distributions and Their Characteristics"⟩, Cover and Thomas (1991), page 25, Jensen (1906), pages 179–180

[3] Shao (2003) pages 31–32 ⟨"Example 1.18"⟩, Dekking et al. (2006) page 110 ⟨"8.5 Solutions to the quick exercises"⟩

[4] Suhov et al. (2005) page 145 ⟨(2.69)⟩, Allen (2018) page 490 ⟨18.3.4 The Law of the Unconscious Statistician⟩, Papoulis (1990) page 124 ⟨Fundamental Theorem⟩

[5] Ross (1998), page 395

BY-NC-ND | COPYRIGHT © 2019 DANIEL J. GREENHOE

✎PROOF:

$$
\begin{aligned}
I &\triangleq \begin{cases} 1 & \text{for} \quad \mathsf{X} \geq a \\ 0 & \text{for} \quad \mathsf{X} < a \end{cases} \\
aI &\leq \mathsf{X} \\
I &\leq \frac{1}{a}\mathsf{X} \\
\mathsf{E}I &\leq \mathsf{E}\left(\frac{1}{a}\mathsf{X}\right)
\end{aligned}
$$

$$
\begin{aligned}
\mathsf{P}\{\mathsf{X} \geq a\} &= 1 \cdot \mathsf{P}\{\mathsf{X} \geq a\} + 0 \cdot \mathsf{P}\{\mathsf{X} < a\} \\
&= \mathsf{E}I \\
&\leq \mathsf{E}\left(\frac{1}{a}\mathsf{X}\right) \\
&= \frac{1}{a}\mathsf{E}\mathsf{X}
\end{aligned}
$$

✏

Theorem 1.4 (Chebyshev's inequality)**.** [6] *Let* X *be a* RANDOM VARIABLE *with mean* μ *and variance* σ^2.

THM $\mathsf{P}\{|\mathsf{X} - \mu| \geq a\} \leq \dfrac{\sigma^2}{a^2}$

✎PROOF:

$$
\begin{aligned}
\mathsf{P}\{|\mathsf{X} - \mu| \geq a\} &= \mathsf{P}\left\{(\mathsf{X} - \mu)^2 \geq a^2\right\} \\
&\leq \frac{1}{a^2}\mathsf{E}(\mathsf{X} - \mu)^2 && \text{by } \textit{Markov's inequality} && \text{(Theorem 1.3 page 6)} \\
&= \frac{\sigma^2}{a^2}
\end{aligned}
$$

✏

Theorem 1.5 (Kolmogorov's inequality)**.** [7] *Let* X *be a* RANDOM VARIABLE *with mean* μ *and variance* σ^2.

THM $\left\{\begin{array}{lll} (A). & (\mathsf{x}_n) \textit{ are } \text{INDEPENDENT} & \textit{and} \\ (B). & \textit{Each } \mathsf{x}_n \textit{ has } \text{ZERO-MEAN} & \textit{and} \\ (C). & \textit{Each } \mathsf{x}_n \textit{ has variance } \sigma^2 & \end{array}\right\} \implies \mathsf{P}\left[\left|\sum_{n=1}^{N} \mathsf{x}_n\right| < \lambda \sum_{n=1}^{N} \mathsf{x}_n^2\right] \geq 1 - \frac{1}{\lambda^2}$

1.4 Joint and conditional probability spaces

Sometimes the problem of finding the expected value of a *random variable* X can be simplified by "conditioning X on Y".

Theorem 1.6. *Let* X *and* Y *be* RANDOM VARIABLES*. Then*

THM $\mathsf{E}_{\mathsf{x}}\mathsf{X} = \mathsf{E}_{\mathsf{y}}\mathsf{E}_{\mathsf{x}|\mathsf{y}}(\mathsf{X}|\mathsf{Y})$

[6] Ross (1998), page 396
[7] Wilks (1963b), page 107 ⟨§4.5 "Kolmogorov's inequality"⟩

COPYRIGHT © 2019 DANIEL J. GREENHOE | CC BY-NC-ND

✎PROOF:

$$\begin{aligned}
\mathsf{E}_{\mathsf{y}}\mathsf{E}_{\mathsf{x}|\mathsf{y}}(\mathsf{X}|\mathsf{Y}) &\triangleq \mathsf{E}_{\mathsf{y}}\left[\int_{x\in\mathbb{R}} x\mathsf{p}(\mathsf{X}=x|\mathsf{Y})\,\mathrm{d}x\right] \\
&\triangleq \int_{y\in\mathbb{R}}\left[\int_{x\in\mathbb{R}} x\mathsf{p}(x|\mathsf{Y}=y)\,\mathrm{d}x\right]\mathsf{p}(y)\,\mathrm{d}y \\
&= \int_{y\in\mathbb{R}}\int_{x\in\mathbb{R}} x\mathsf{p}(x|y)\mathsf{p}(y)\,\mathrm{d}x\,\mathrm{d}y \\
&= \int_{x\in\mathbb{R}} x\int_{y\in\mathbb{R}} \mathsf{p}(x,y)\,\mathrm{d}y\,\mathrm{d}x \\
&= \int_{x\in\mathbb{R}} x\mathsf{p}(x)\,\mathrm{d}x \\
&\triangleq \mathsf{E}_{\mathsf{x}}\mathsf{X}
\end{aligned}$$

✏

1.5 Expectation inner product space

When possible, we like to generalize any given mathematical structure to a more general mathematical structure and then take advantage of the properties of that more general structure. Such a generalization can be done with *random variables*. Random variables can be viewed as vectors in a vector space. Furthermore, the expectation of the product of two *random variables* (e.g. $\mathsf{E}(\mathsf{XY})$) can be viewed as an *inner product* in an *inner product space*. Since we have an *inner product* space, we can then immediately use all the properties of *inner product spaces*, *norm*ed spaces, vector spaces, metric spaces, and topological spaces.

Theorem 1.7. [8] *Let R be a ring, $(\Omega,\mathbb{E},\mathsf{P})$ be a* PROBABILITY SPACE, E *the expectation operator, and $\boldsymbol{V}=\{\mathsf{X}|\mathsf{X}:\Omega\to R\}$ be the set of all random vectors in* PROBABILITY SPACE $(\Omega,\mathbb{E},\mathsf{P})$.

THM

(1).	$\boldsymbol{V}\triangleq\{\mathsf{X}\|\mathsf{X}:\Omega\to R\}$	*is a* VECTOR SPACE.
(2).	$\langle\mathsf{X}\,\|\,\mathsf{Y}\rangle\triangleq\mathsf{E}(\mathsf{XY}^*)$	*is an* INNER PRODUCT.
(3).	$\|\mathsf{X}\|\triangleq\sqrt{\mathsf{E}(\mathsf{XX}^*)}$	*is a* NORM.
(4).	$(\boldsymbol{V},\langle\triangle\,\|\,\triangledown\rangle)$	*is an* INNER PRODUCT SPACE.

✎PROOF:

1. Proof that $\boldsymbol{V}$ is a vector space:

1)	$\forall\mathsf{X},\mathsf{Y},\mathsf{Z}\in\boldsymbol{V}$	$(\mathsf{X}+\mathsf{Y})+\mathsf{Z}=\mathsf{X}+(\mathsf{Y}+\mathsf{Z})$	(+ is associative)
2)	$\forall\mathsf{X},\mathsf{Y}\in\boldsymbol{V}$	$\mathsf{X}+\mathsf{Y}=\mathsf{Y}+\mathsf{X}$	(+ is commutative)
3)	$\exists 0\in\boldsymbol{V}$ such that $\forall\mathsf{X}\in\boldsymbol{V}$	$\mathsf{X}+0=\mathsf{X}$	(+ identity)
4)	$\forall\mathsf{X}\in\boldsymbol{V}\exists\mathsf{Y}\in\boldsymbol{V}$ such that	$\mathsf{X}+\mathsf{Y}=0$	(+ inverse)
5)	$\forall\alpha\in S$ and $\mathsf{X},\mathsf{Y}\in\boldsymbol{V}$	$\alpha\cdot(\mathsf{X}+\mathsf{Y})=(\alpha\cdot\mathsf{X})+(\alpha\cdot\mathsf{Y})$	($\cdot$ distributes over +)
6)	$\forall\alpha,\beta\in S$ and $\mathsf{X}\in\boldsymbol{V}$	$(\alpha+\beta)\cdot\mathsf{X}=(\alpha\cdot\mathsf{X})+(\beta\cdot\mathsf{X})$	($\cdot$ pseudo-distributes over +)
7)	$\forall\alpha,\beta\in S$ and $\mathsf{X}\in\boldsymbol{V}$	$\alpha(\beta\cdot\mathsf{X})=(\alpha\cdot\beta)\cdot\mathsf{X}$	($\cdot$ associates with $\cdot$)
8)	$\forall\mathsf{X}\in\boldsymbol{V}$	$1\cdot\mathsf{X}=\mathsf{X}$	($\cdot$ identity)

[8] Lindquist and Picci (2015) pages 25–26 ⟨2.1 Hilbert Space of Second-Order Random Variables. $\langle\xi\,|\,\eta\rangle=\mathsf{E}\{\xi\bar{\eta}\}$⟩, Caines (1988) page 21 ⟨$Exy=\int_\Omega x(\omega)y(\omega)dP(\omega)$⟩, Caines (2018) page 21 ⟨$Exy=\int_\Omega x(\omega)y(\omega)dP(\omega)$⟩, Moon and Stirling (2000), pages 105–106

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

2. Proof that $\langle \mathsf{X} \mid \mathsf{Y} \rangle \triangleq \mathsf{E}(\mathsf{X}\mathsf{Y}^*)$ is an *inner product*.

1)	$\mathsf{E}(\mathsf{X}\mathsf{X}^*)$	≥ 0	$\forall \mathsf{X} \in \boldsymbol{V}$	(non-negative)
2)	$\mathsf{E}(\mathsf{X}\mathsf{X}^*)$	$= 0 \iff \mathsf{X} = 0$	$\forall \mathsf{X} \in \boldsymbol{V}$	(non-degenerate)
3)	$\mathsf{E}(\alpha \mathsf{X}\mathsf{Y}^*)$	$= \alpha \mathsf{E}(\mathsf{X}\mathsf{Y}^*)$	$\forall \mathsf{X}, \mathsf{Y} \in \boldsymbol{V},\ \forall \alpha \in \mathbb{C}$	(homogeneous)
4)	$\mathsf{E}[(\mathsf{X}+\mathsf{Y})\mathsf{Z}^*]$	$= \mathsf{E}(\mathsf{X}\mathsf{Z}^*) + \mathsf{E}(Y\mathsf{Z}^*)$	$\forall \mathsf{X}, \mathsf{Y}, \mathsf{Z} \in \boldsymbol{V}$	(additive)
5)	$\mathsf{E}(\mathsf{X}\mathsf{Y}^*)$	$= \mathsf{E}(\mathsf{Y}\mathsf{X}^*)$	$\forall \mathsf{X}, \mathsf{Y} \in \boldsymbol{V}$	(conjugate symmetric).

3. Proof that $\|\mathsf{X}\| \triangleq \sqrt{\mathsf{E}(\mathsf{X}\mathsf{X}^*)}$ is a *norm*: This *norm* is simply induced by the above *inner product*.

4. Proof that $(\boldsymbol{V}, \langle \triangle \mid \triangledown \rangle)$ is an *inner product space*: Because $\boldsymbol{V}$ is a vector space and $\langle \triangle \mid \triangledown \rangle$ is an *inner product*, $(\boldsymbol{V}, \langle \triangle \mid \triangledown \rangle)$ is an *inner product space*.

The next theorem gives some results that follow directly from vector space properties:

Theorem 1.8. *Let* $(\Omega, \mathbb{E}, \mathsf{P})$ *be a* PROBABILITY SPACE *with expectation functional* E.

THM

1.	$\sqrt{\mathsf{E}\left(\sum_{n=1}^{N} \mathsf{X}_n\right)} \leq \sum_{n=1}^{N} \mathsf{E}(\mathsf{X}_n X_n^*)$		(GENERALIZED TRIANGLE INEQUALITY)
2.	$\left\|\mathsf{E}(\mathsf{X}\mathsf{Y}^*)\right\|^2 \leq \mathsf{E}(\mathsf{X}\mathsf{X}^*)\,\mathsf{E}(Y Y^*)$		(CAUCHY-SCHWARTZ INEQUALITY)
3.	$2\mathsf{E}(\mathsf{X}\mathsf{X}^*) + 2\mathsf{E}(Y Y^*) = \mathsf{E}[(\mathsf{X}+\mathsf{Y})(\mathsf{X}+\mathsf{Y})^*] + \mathsf{E}[(\mathsf{X}-Y)(\mathsf{X}-Y)^*]$		(PARALLELOGRAM LAW)

✎PROOF:

1. $\left(\mathbb{R}^{\Omega}, \mathsf{E}(\mathsf{x},\mathsf{y})\right)$ is an *inner product space*. Proof: Theorem 1.7 (page 8).

2. Because it is an *inner product space*, the other properties follow:
   1. Generalized triangle inequality: Theorem K.1 page 271
   2. Cauchy-Schwartz inequality: Theorem J.2 page 256
   3. Parallelogram Law: Theorem J.7 page 263

COPYRIGHT © 2019 DANIEL J. GREENHOE
BY-NC-ND

Copyright © 2019 Daniel J. Greenhoe

CHAPTER 2

RANDOM SEQUENCES

❝ *A likely impossibility is always preferable to an unconvincing possibility.*❞
Aristotle (384 BC – 322 BC) [1]

❝ *We are quite in danger of sending highly trained and highly intelligent young men out into the world with tables of erroneous numbers under their arms, and with a dense fog in the place where their brains ought to be. In this century, of course, they will be working on guided missiles and advising the medical profession on the control of disease, and there is no limit to the extent to which they could impede every sort of national effort.*❞

Ronald A. Fisher, (1890–1962), Statistician, at a lecture in 1958 at Michigan State University [2]

2.1 Definitions

Definition 2.1.

DEF

A ***random sequence*** *is a* SEQUENCE *over a* PROBABILITY SPACE *(Definition A.2 page 165)*.

Definition 2.2. [3] *Let* $\mathsf{x}(n)$ *and* $\mathsf{y}(n)$ *be* RANDOM SEQUENCES.

[1] quote: http://en.wikiquote.org/wiki/Aristotle
image: http://en.wikipedia.org/wiki/Aristotle

[2] quote: Yates and Mather (1963) page 107. image: http://www.genetics.org/content/154/4/1419

[3] Papoulis (1984) page 263 $\langle R_{xy}(m) = E\{\mathsf{x}(m)\mathsf{y}^*(0)\}\rangle$, Wilks (1963b), page 77 ⟨§3.4 "Moments of two-dimensional random variables"⟩, Cadzow (1987) page 341 $\langle r_{xy}(m) = E[\mathsf{x}(m)\mathsf{y}^*(0)]\rangle$, MatLab (2018b) $\langle R_{xy}(m) = E\{x_{n+m}y_n^*\}\rangle$, MatLab (2018a) $\langle R_{xy}(m) = E\{x_{n+m}y_n^*\}\rangle$

DEF

The ***mean***	$\mu_{\mathsf{x}}(n)$	*of* $\mathsf{x}(n)$ *is*	$\mu_{\mathsf{x}}(n)$	$\triangleq$	$\mathsf{E}[\mathsf{x}(n)]$
The ***variance***	$\sigma^2_{\mathsf{x}}(n)$	*of* $\mathsf{x}(n)$ *is*	$\sigma^2_{\mathsf{x}}(n)$	$\triangleq$	$\mathsf{E}\Big(\big[\mathsf{x}(n)-\mu_{\mathsf{x}}(n)\big]^2\Big)$
The ***cross-correlation***	$\mathsf{R}_{\mathsf{xy}}(n,m)$	*of* $\mathsf{x}(n)$ *and* $\mathsf{y}(n)$ *is*	$\mathsf{R}_{\mathsf{xy}}(n,m)$	$\triangleq$	$\mathsf{E}\big[\mathsf{x}(n+m)\mathsf{y}^*(n)\big]$
The ***auto-correlation***	$\mathsf{R}_{\mathsf{xx}}(n,m)$	*of* $\mathsf{x}(n)$ *is*	$\mathsf{R}_{\mathsf{xx}}(n,m)$	$\triangleq$	$\mathsf{R}_{\mathsf{xy}}(n,m)\big\vert_{\mathsf{y}=\mathsf{x}}$

2.2 Properties

Theorem 2.1.

THM

$$\mathsf{R}_{\mathsf{xx}}(n,m) = \mathsf{R}^*_{\mathsf{xx}}(n+m,-m)$$
$$\mathsf{R}_{\mathsf{xy}}(n,m) = \mathsf{R}^*_{\mathsf{yx}}(n+m,-m)$$

✎PROOF:

$\mathsf{R}_{\mathsf{xy}}(n,m) \triangleq \mathsf{E}\big[\mathsf{x}(n+m)\mathsf{y}^*(n)\big]$	by definition of $\mathsf{R}_{\mathsf{xy}}(n,m)$	(Definition 2.2 page 11)
$= \mathsf{E}\big[\mathsf{y}^*(n)\mathsf{x}(n+m)\big]$	by *commutative* property of $(\mathbb{C},+,\cdot,0,1)$	(Definition F.5 page 206)
$= \big(\mathsf{E}\big[\mathsf{y}(n)\mathsf{x}^*(n+m)\big]\big)^*$	by *distributive* property of $*$-**algebras**	(Definition I.3 page 250)
$= \big(\mathsf{E}\big[\mathsf{y}(n+m-m)\mathsf{x}^*(n+m)\big]\big)^*$	by *additive identity* property of $(\mathbb{R},+,\cdot,0,1)$	(Definition F.5 page 206)
$\triangleq \mathsf{R}^*_{\mathsf{yx}}(n+m,-m)$	by definition of $\mathsf{R}_{\mathsf{xy}}(n,m)$	(Definition 2.2 page 11)

$\mathsf{R}_{\mathsf{xx}}(n,m) = \mathsf{R}_{\mathsf{xy}}(n,m)\big\vert_{\mathsf{y}=\mathsf{x}}$	by $\mathsf{y}=\mathsf{x}$ constraint
$= \mathsf{R}^*_{\mathsf{xy}}(n+m,-m)\big\vert_{\mathsf{y}=\mathsf{x}}$	by previous result
$= \mathsf{R}^*_{\mathsf{xx}}(n+m,-m)$	by $\mathsf{y}=\mathsf{x}$ constraint

✏

2.3 Wide Sense Stationary processes

Definition 2.3. *Let* $\mathsf{x}(n)$ *be a* RANDOM SEQUENCE *with* MEAN $\mu_{\mathsf{x}}(n)$ *and* VARIANCE $\sigma^2(n)$ *(Definition 2.2 page 11)*.

DEF

$\mathsf{x}(n)$ *is* ***wide sense stationary (WSS)*** *if*

1. $\mu_{\mathsf{x}}(n)$ *is* CONSTANT *with respect to* n (STATIONARY IN THE 1ST MOMENT) *and*
2. $\sigma^2_{\mathsf{x}}(n)$ *is* CONSTANT *with respect to* n (STATIONARY IN THE 2ND MOMENT)

Definition 2.4. [4] *Let* $\mathsf{x}(n)$ *be a* RANDOM SEQUENCE *with statistics* $\mu_{\mathsf{x}}(n)$, $\sigma^2_{\mathsf{x}}(n)$, $\mathsf{R}_{\mathsf{xx}}(n,m)$, *and* $\mathsf{R}_{\mathsf{xy}}(n,m)$ *(Definition 2.2 page 11)*.

DEF

$\{$ x *and* y *are* WIDE SENSE STATIONARY $\}$ $\Longrightarrow$

(1).	*The* ***mean***	μ_{x}	*of* $\mathsf{x}(n)$ *is*	μ_{x}	$\triangleq$	$\mathsf{E}[\mathsf{x}(0)]$
(2).	*The* ***variance***	σ^2_{x}	*of* $\mathsf{x}(n)$ *is*	σ^2_{x}	$\triangleq$	$\mathsf{E}\Big(\big[\mathsf{x}(0)-\mu_{\mathsf{x}}\big]^2\Big)$
(4).	*The* ***cross-correlation***	$\mathsf{R}_{\mathsf{xy}}(m)$	*of* $\mathsf{x}(n)$ *and* $\mathsf{y}(n)$ *is*	$\mathsf{R}_{\mathsf{xy}}(m)$	$\triangleq$	$\mathsf{E}\big[\mathsf{x}(m)\mathsf{y}^*(0)\big]$
(3).	*The* ***auto-correlation***	$\mathsf{R}_{\mathsf{xx}}(m)$	*of* $\mathsf{x}(n)$ *is*	$\mathsf{R}_{\mathsf{xx}}(m)$	$\triangleq$	$\mathsf{R}_{\mathsf{xy}}(m)\big\vert_{\mathsf{y}=\mathsf{x}}$

[4] Papoulis (1984) page 263 ($\langle$"$R_{xy}(\tau) = E\big\{\mathsf{x}(t+\tau)\mathsf{y}^*(t)\big\}$"$\rangle$, Cadzow (1987) page 341 $\langle r_{xy}(n) = E\big[\mathsf{x}(k+n)\mathsf{y}^*(k)\big]$ (10.41)$\rangle$

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

Remark 2.1. The $\mathsf{R}_{\mathsf{xy}}(n,m)$ of Definition 2.2 (page 11) and the $\mathsf{R}_{\mathsf{xy}}(m)$ of Definition 2.4 (page 12) (etc.) are examples of *function overload*—that is, functions that use the same mnemonic but are distinguished by different domains. Perhaps a more common example of function overload is the "+" mnemonic. Traditionally it is used with domain of the natural numbers $\mathbb{N}$ as in $3+2$. Later it was extended for domain real numbers $\mathbb{R}$ as in $\sqrt{3}+\sqrt{2}$, or even complex numbers $\mathbb{C}$ as in $\left(\sqrt{3}+i\sqrt{2}\right)+(e+i\pi)$. And it was even more dramatically extended for use with domain $\mathbb{R}^N \times \mathbb{R}^M$ in "linear algebra" as in

$$\begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} + \begin{bmatrix} 5 & 6 \\ 7 & 8 \end{bmatrix} = \begin{bmatrix} 6 & 8 \\ 10 & 12 \end{bmatrix}$$

Remark 2.2. The definition for $\mathsf{R}_{\mathsf{xy}}(m)$ can be defined with the conjugate $*$ on either x or y, or on neither or both; and moreover x may either lead or lag y. In total, there are $2\times 2\times 2 = 8$ different ways to define $\mathsf{R}_{\mathsf{xy}}(m)$.[5] and $\mathsf{R}_{\mathsf{xx}}(m)$ involve complex numbers. This may seem curious when typical ADCs provide real-valued sequences. Note however that complex-valued sequences often come up in signal processing due to two common situations:[6]

1. The presence of an *FFT* operator in the signal processing path
2. communications channel processing involving phase discrimination (e.g. PSK and QAM).

Proposition 2.1. *Let* $\mathsf{y}(n)$ *be a* RANDOM SEQUENCE, $\mathsf{x}(n)$ *a* RANDOM SEQUENCE *with* AUTO-CORRELATION $\mathsf{R}_{\mathsf{xx}}(n,m)$, *and* R_{xy} *the* CROSS-CORRELATION *of* x *and* y.

PRP

$$\left\{\begin{array}{l} \mathsf{x} \text{ and } \mathsf{y} \text{ are} \\ \text{WIDE SENSE STATIONARY} \\ \text{(WSS) (Definition 6.1 page 43)} \end{array}\right\} \Longrightarrow \left\{\begin{array}{lcll} \mathsf{R}_{\mathsf{xx}}(n,m) & = & \mathsf{R}_{\mathsf{xx}}(m) & \forall n\in\mathbb{Z} \\ \mathsf{R}_{\mathsf{xy}}(n,m) & = & \mathsf{R}_{\mathsf{xy}}(m) & \forall n\in\mathbb{Z} \\ \text{(Definition 2.2 page 11)} & & \text{(Definition 2.4 page 12)} & \end{array}\right\}$$

✎PROOF:

$$\begin{aligned}
\mathsf{R}_{\mathsf{xy}}(n,m) &\triangleq \mathsf{E}\left[\mathsf{x}[n+m]\mathsf{y}^*[n]\right] && \text{by definition of } \mathsf{R}_{\mathsf{xy}}(n,m) && \text{(Definition 2.2 page 11)} \\
&= \mathsf{E}\left[\mathsf{x}[n-n+m]\mathsf{y}^*[n-n]\right] && \text{by } \textit{wide sense stationary} \text{ hypothesis} \\
&= \mathsf{E}\left[\mathsf{x}[m]\mathsf{y}^*[0]\right] \\
&\triangleq \mathsf{R}_{\mathsf{xy}}(m) && \text{by definition of } \mathsf{R}_{\mathsf{xy}}(m) && \text{(Definition 2.4 page 12)} \\
\mathsf{R}_{\mathsf{xx}}(n,m) &= \mathsf{R}_{\mathsf{xy}}(n,m)\big|_{\mathsf{y}=\mathsf{x}} \\
&= \mathsf{R}_{\mathsf{xy}}(m)\big|_{\mathsf{y}=\mathsf{x}} && \text{by previous result} \\
&= \mathsf{R}_{\mathsf{xx}}(m)
\end{aligned}$$

✎



Figure 2.1: *auto-correlation* $\mathsf{R}_{\mathsf{xx}}(m)$

[5] Greenhoe (2019)

[6] S. Lawrence Marple (1987) page ?, S. Lawrence Marple (2019) pages 48–50 ⟨§"2.12 Extra: Source of Complex-Valued Signals"⟩

COPYRIGHT © 2019 DANIEL J. GREENHOE

Corollary 2.1. *Let* $\mathsf{x}(n)$ *be a* RANDOM SEQUENCE *with* AUTO-CORRELATION $\mathsf{R}_{\mathsf{xx}}(n,m)$, $\mathsf{y}(n)$ *a* RANDOM SEQUENCE *with* AUTO-CORRELATION $\mathsf{R}_{\mathsf{yy}}(n,m)$, *and* $\mathsf{R}_{\mathsf{xy}}(n,m)$ *the* CROSS-CORRELATION *of* x *and* y. *Let* $\mathbf{S}$ *be a* SYSTEM *with input* $\mathsf{x}(n)$ *and output* $\mathsf{y}(n)$.

COR

$$\left\{\begin{array}{lll} (A). & \mathsf{x} \text{ is WSS} & \text{and} \\ (B). & \mathsf{y} \text{ is WSS} & \text{and} \\ (C). & \mathbf{S} \text{ is LTI} & \end{array}\right\} \Longrightarrow \left\{\begin{array}{llllll} (1). & \mathsf{R}_{\mathsf{xy}}(m) & = & \mathsf{R}^*_{\mathsf{yx}}(-m) & & \text{and} \\ (2). & \mathsf{R}_{\mathsf{xx}}(m) & = & \mathsf{R}^*_{\mathsf{xx}}(-m) & \text{(CONJUGATE SYMMETRIC)} & \text{and} \\ (3). & \mathbf{R}_{\mathsf{e}}\mathsf{R}_{\mathsf{xx}}(m) & = & \mathbf{R}_{\mathsf{e}}\mathsf{R}_{\mathsf{xx}}(-m) & \text{(SYMMETRIC)} & \text{and} \\ (4). & \mathbf{I}_{\mathsf{m}}\mathsf{R}_{\mathsf{xx}}(m) & = & -\mathbf{I}_{\mathsf{m}}\mathsf{R}_{\mathsf{xx}}(-m) & \text{(ANTI-SYMMETRIC)} & \text{and} \\ (5). & \left|\mathsf{R}_{\mathsf{xx}}(m)\right| & = & \left|\mathsf{R}_{\mathsf{xx}}(-m)\right| & \text{(SYMMETRIC)} & \text{and} \\ (6). & \angle\mathsf{R}_{\mathsf{xx}}(m) & = & -\angle\mathsf{R}_{\mathsf{xx}}(-m) & \text{(ANTI-SYMMETRIC)} & \end{array}\right\}$$

✎PROOF:

$$\begin{array}{rlll} \mathsf{R}_{\mathsf{xy}}(m) & = \mathsf{R}_{\mathsf{xy}}(n,m) & \text{by Proposition 2.1 page 13} & \text{and hypotheses (A),(B)} \\ & = \mathsf{R}^*_{\mathsf{yx}}(n+m,-m) & \text{by Theorem 2.1 page 12} & \text{and hypothesis (B)} \\ & = \mathsf{R}^*_{\mathsf{yx}}(-m) & \text{by Proposition 2.1 page 13} & \text{and hypothesis (A)} \\ \mathsf{R}_{\mathsf{xx}}(m) & = \mathsf{R}_{\mathsf{xx}}(n,m) & \text{by Proposition 2.1 page 13} & \text{and hypothesis (A)} \\ & = \mathsf{R}^*_{\mathsf{xx}}(n+m,-m) & \text{by Theorem 2.1 page 12} & \text{and hypothesis (B)} \\ & = \mathsf{R}^*_{\mathsf{xx}}(-m) & \text{by Proposition 2.1 page 13} & \text{and hypothesis (A)} \end{array}$$

✏

2.4 Spectral density

Definition 2.5. *Let* $\mathsf{x}(n)$ *and* $\mathsf{y}(n)$ *be* WIDE SENSE STATIONARY RANDOM SEQUENCES *with auto-correlation* $\mathsf{R}_{\mathsf{xx}}(m)$ *and cross-correlation* $\mathsf{R}_{\mathsf{xy}}(m)$. *Let* $\mathbf{Z}$ *be the* Z-TRANSFORM OPERATOR *(Definition Q.4 page 354)*.

DEF

The ***z-domain cross spectral density*** ***(CSD)*** $\check{\mathsf{S}}_{\mathsf{xy}}(z)$ *of* $\mathbf{x}$ *and* $\mathbf{y}$ *is*

$$\check{\mathsf{S}}_{\mathsf{xy}}(z) \triangleq \mathbf{Z}\mathsf{R}_{\mathsf{xy}}(m) \triangleq \sum_{m\in\mathbb{Z}} \mathsf{R}_{\mathsf{xy}}(m)z^{-m}$$

The ***z-domain power spectral density*** ***(PSD)*** $\check{\mathsf{S}}_{\mathsf{xx}}(z)$ *of* $\mathbf{x}$ *is* $\quad \check{\mathsf{S}}_{\mathsf{xx}}(z) \triangleq \check{\mathsf{S}}_{\mathsf{xy}}(z)\big|_{\mathsf{y}(n)=\mathsf{x}(n)}$

Definition 2.6. *Let* $\mathsf{x}(n)$ *and* $\mathsf{y}(n)$ *be* WIDE SENSE STATIONARY RANDOM SEQUENCES *with auto-correlation* $\mathsf{R}_{\mathsf{xx}}(m)$ *and cross-correlation* $\mathsf{R}_{\mathsf{xy}}(m)$. *Let* $\check{\mathbf{F}}$ *be the* DISCRETE TIME FOURIER TRANSFORM (DTFT) *operator (Definition P.1 page 343)*.

DEF

The ***auto-spectral density*** $\check{\mathsf{S}}_{\mathsf{xx}}(z)$ *of* $\mathbf{x}$ *is*

$$\tilde{\mathsf{S}}_{\mathsf{xx}}(z) \triangleq \check{\mathbf{F}}\mathsf{R}_{\mathsf{xx}}(m) \triangleq \sum_{m\in\mathbb{Z}} \mathsf{R}_{\mathsf{xx}}(m)e^{-i\omega m}$$

The ***cross spectral density*** ***(CSD)*** $\check{\mathsf{S}}_{\mathsf{xy}}(z)$ *of* $\mathbf{x}$ *and* $\mathbf{y}$ *is*

$$\tilde{\mathsf{S}}_{\mathsf{xy}}(z) \triangleq \check{\mathbf{F}}\mathsf{R}_{\mathsf{xy}}(m) \triangleq \sum_{m\in\mathbb{Z}} \mathsf{R}_{\mathsf{xy}}(m)e^{-i\omega m}$$

The ***auto-spectral density*** *is also called* ***power spectral density (PSD)***.

Theorem 2.2. *Let* $\mathbf{S}$ *be a system with* IMPULSE RESPONSE $\mathsf{h}(n)$, INPUT $\mathsf{x}(n)$, *and* OUTPUT $\mathsf{y}(n)$.

THM

$$\left\{\ \mathsf{x} \text{ and } \mathsf{y} \text{ are WIDE SENSE STATIONARY}\ \right\} \Longrightarrow \left\{\begin{array}{llll} (1). & \check{\mathsf{S}}_{\mathsf{xx}}(z) & = & \check{\mathsf{S}}^*_{\mathsf{xx}}\left(\frac{1}{z^*}\right) \quad \text{and} \\ (2). & \check{\mathsf{S}}_{\mathsf{yx}}(z) & = & \check{\mathsf{S}}^*_{\mathsf{xy}}\left(\frac{1}{z^*}\right) \end{array}\right\}$$

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

✎PROOF:

$$\begin{aligned}
\check{\mathsf{S}}_{\mathsf{yx}}(z) &\triangleq \mathbf{Z}\mathsf{R}_{\mathsf{yx}}(m) && \text{by definition of } \check{\mathsf{S}}_{\mathsf{xy}}(z) && \text{(Definition 2.6 page 14)}\\
&\triangleq \sum_{m\in\mathbb{Z}} \mathsf{R}_{\mathsf{yx}}(m)z^{-m} && \text{by definition of } \mathbf{Z} && \text{(Definition Q.4 page 354)}\\
&\triangleq \sum_{m\in\mathbb{Z}} \mathsf{R}^*_{\mathsf{xy}}(-m)z^{-m} && \text{by Corollary 2.1 page 14} && \\
&= \left[\sum_{m\in\mathbb{Z}} \mathsf{R}_{\mathsf{xy}}(-m)(z^*)^{-m}\right]^* && \text{by } \textit{antiautomorphic} \text{ property of *-algebras} && \text{(Definition I.3 page 250)}\\
&= \left[\sum_{-p\in\mathbb{Z}} \mathsf{R}_{\mathsf{xy}}(p)(z^*)^{p}\right]^* && \text{where } p \triangleq -m && \implies m = -p\\
&= \left[\sum_{p\in\mathbb{Z}} \mathsf{R}_{\mathsf{xy}}(p)(z^*)^{p}\right]^* && \text{by } \textit{absolutely summable} \text{ property} && \text{(Definition Q.2 page 353)}\\
&= \left[\sum_{p\in\mathbb{Z}} \mathsf{R}_{\mathsf{xy}}(p)\left(\frac{1}{z^*}\right)^{-p}\right]^* && && \\
&= \check{\mathsf{S}}^*_{\mathsf{xy}}\left(\frac{1}{z^*}\right) && \text{by definition of } \mathbf{Z} && \text{(Definition Q.4 page 354)}\\
\check{\mathsf{S}}_{\mathsf{xx}}(z) &= \check{\mathsf{S}}_{\mathsf{xy}}(z)\big|_{\mathsf{y}=\mathsf{x}} && && \\
&= \check{\mathsf{S}}^*_{\mathsf{yx}}(z)\big|_{\mathsf{y}=\mathsf{x}} && && \\
&= \check{\mathsf{S}}^*_{\mathsf{xy}}\left(\frac{1}{z^*}\right)\Big|_{\mathsf{y}=\mathsf{x}} && \text{by (2)—previous result} && \\
&= \check{\mathsf{S}}^*_{\mathsf{xx}}\left(\frac{1}{z^*}\right) && &&
\end{aligned}$$

✏

Corollary 2.2. *Let* $\mathbf{S}$ *be a system with* IMPULSE RESPONSE $\mathsf{h}(n)$, INPUT $\mathsf{x}(n)$, *and* OUTPUT $\mathsf{y}(n)$.

COR

$$\left\{\begin{array}{ll} (A). & \mathsf{h} \textit{ is } \text{LTI} \quad \textit{and} \\ (B). & \mathsf{x} \textit{ and } \mathsf{y} \textit{ are } \text{WSS} \end{array}\right\} \implies \left\{\begin{array}{llll} (1). & \tilde{\mathsf{S}}^*_{\mathsf{xy}}(\omega) = \tilde{\mathsf{S}}_{\mathsf{yx}}(\omega) & \text{(CONJUGATE-SYMMETRIC)} & \textit{and} \\ (2). & \tilde{\mathsf{S}}^*_{\mathsf{xx}}(\omega) = \tilde{\mathsf{S}}_{\mathsf{xx}}(\omega) & \text{(CONJUGATE SYMMETRIC)} & \textit{and} \\ (3). & \tilde{\mathsf{S}}_{\mathsf{xx}}(\omega) \in \mathbb{R} & \text{(REAL-VALUED)} & \end{array}\right\}$$

✎PROOF:

$$\begin{aligned}
\tilde{\mathsf{S}}^*_{\mathsf{xy}}(\omega) &= \check{\mathsf{S}}^*_{\mathsf{xy}}(z)\big|_{z=e^{i\omega}} && \text{by definition of } \textit{DTFT} && \text{(Definition P.1 page 343)}\\
&= \check{\mathsf{S}}^{**}_{\mathsf{yx}}\left(\frac{1}{z^*}\right)\Big|_{z=e^{i\omega}} && \text{by Theorem 2.2 page 14} && \\
&= \check{\mathsf{S}}_{\mathsf{yx}}\left(\frac{1}{z^*}\right)\Big|_{z=e^{i\omega}} && \text{by } \textit{involutory} \text{ property of } *\textbf{-algebras} && \text{(Definition I.3 page 250)}\\
&= \check{\mathsf{S}}_{\mathsf{yx}}\left(\frac{1}{e^{i\omega *}}\right) && && \\
&= \check{\mathsf{S}}_{\mathsf{yx}}\left(e^{i\omega}\right) && && \\
&= \tilde{\mathsf{S}}_{\mathsf{yx}}(\omega) && \text{by definition of } \textit{DTFT} && \text{(Definition P.1 page 343)}\\
\tilde{\mathsf{S}}^*_{\mathsf{xx}}(\omega) &= \check{\mathsf{S}}^*_{\mathsf{xx}}(z)\big|_{z=e^{i\omega}} && \text{by definition of } \textit{DTFT} && \text{(Definition P.1 page 343)}\\
&= \check{\mathsf{S}}^{**}_{\mathsf{xx}}\left(\frac{1}{z^*}\right)\Big|_{z=e^{i\omega}} && \text{by Theorem 2.2 page 14} && \\
&= \check{\mathsf{S}}_{\mathsf{xx}}\left(\frac{1}{z^*}\right)\Big|_{z=e^{i\omega}} && \text{by } \textit{involutory} \text{ property of } *\textbf{-algebras} && \text{(Definition I.3 page 250)}\\
&= \check{\mathsf{S}}_{\mathsf{xx}}\left(\frac{1}{e^{i\omega *}}\right) && &&
\end{aligned}$$

COPYRIGHT © 2019 DANIEL J. GREENHOE

$$
\begin{aligned}
&= \check{S}_{xx}\left(e^{i\omega}\right) \\
&= \tilde{S}_{xx}(\omega) && \text{by definition of } \textit{DTFT} && \text{(Definition P.1 page 343)} \\
&\implies \tilde{S}_{xx}(\omega) \text{ is } \textit{real-valued} \\
\tilde{S}^*_{xx}(\omega) &= \tilde{S}^*_{xy}(\omega)\big|_{y=x} \\
&= \tilde{S}_{yx}(\omega)\big|_{y=x} && \text{by previous result} \\
&= \tilde{S}_{xx}(\omega)
\end{aligned}
$$

✏

2.5 Spectral Power

The term "*spectral power*" is a bit of an oxymoron because "spectral" deals with leaving the time-domain for the frequency-domain, howbeit the concept of power is solidly founded on the concept of time in that power = energy per time.

However, *Parseval's Theorem* (Proposition H.2 page 230) demonstrates that power in time can also be calculated in frequency. So, it makes some sense to speak of the term "spectral power". Moreover, one way to estimate this power is to average the Fourier Transforms of the product $|x(n)|^2 = x(n)x^*(n)$…that is, to use an estimate of the auto-spectral density $\tilde{S}_{xx}(\omega)$. Thus, an alternate name for *auto-spectral density* is **power spectral density** (PSD).

BY-NC-ND Copyright © 2019 Daniel J. Greenhoe

CHAPTER 3

CONTINUOUS RANDOM PROCESSES



❝ *A likely impossibility is always preferable to an unconvincing possibility.* ❞
Aristotle (384 BC – 322 BC) [1]

3.1 Definitions

Definition 3.1. [2] *Let* $(\Omega, \mathbb{E}, \mathsf{P})$ *be a* PROBABILITY SPACE.

DEF
The function $\mathsf{x} : \Omega \to \mathbb{R}$ *is a* ***random variable****.*
The function $\mathsf{y} : \mathbb{R} \times \Omega \to \mathbb{R}$ *is a* ***random process****.*

The random process $\mathsf{x}(t, \omega)$, where t commonly represents time and $\omega \in \Omega$ is an outcome of an experiment, can take on more specialized forms depending on whether t and ω are fixed or allowed to vary. These forms are illustrated in Figure 3.1 page 17[3] and Figure 3.2 page 18.

$\mathsf{x}(t, \omega)$	fixed t	variable t
fixed ω	number	time function
variable ω	random variable	random process

Figure 3.1: Specialized forms of a random process $\mathsf{x}(t, \omega)$

[1] quote: http://en.wikiquote.org/wiki/Aristotle
image: http://en.wikipedia.org/wiki/Aristotle
[2] Papoulis (1991), page 63, Papoulis (1991), page 285
[3] Papoulis (1991), pages 285–286



Figure 3.2: Example of a random process $\mathsf{x}(t,\omega)$

Definition 3.2. [4] *Let* $\mathsf{x}(t)$ *and* $\mathsf{y}(t)$ *be random processes.*

DEF						
	The ***mean***	$\mu_{\mathsf{X}}(t)$	*of* $\mathsf{x}(t)$ *is*	$\mu_{\mathsf{X}}(t)$	$\triangleq$	$\mathsf{E}[\mathsf{x}(t)]$
	The ***cross-correlation***	$\mathsf{R}_{\mathsf{xy}}(t)$	*of* $\mathsf{x}(t)$ *and* $\mathsf{y}(t)$ *is*	$\mathsf{R}_{\mathsf{xy}}(t,u)$	$\triangleq$	$\mathsf{E}\left[\mathsf{x}(t)\mathsf{y}^*(u)\right]$
	The ***auto-correlation function***	$\mathsf{R}_{\mathsf{xx}}(t)$	*of* $\mathsf{x}(t)$ *is*	$\mathsf{R}_{\mathsf{xx}}(t,u)$	$\triangleq$	$\mathsf{E}\left[\mathsf{x}(t)\mathsf{x}^*(u)\right]$

Remark 3.1. [5] The equation $\int_{u\in\mathbb{R}} \mathsf{R}_{\mathsf{xx}}(t,u)\mathsf{f}(u)\,\mathrm{d}u$ is a *Fredholm integral equation of the first kind* and $\mathsf{R}_{\mathsf{xx}}(t,u)$ is the *kernel* of the equation.

Theorem 3.1. *Let* $\mathsf{x}(t)$ *and* $\mathsf{y}(t)$ *be random processes with cross-correlation* $\mathsf{R}_{\mathsf{xy}}(t,u)$ *and let* $\mathsf{R}_{\mathsf{xx}}(t,u)$ *be the auto-correlation of* $\mathsf{x}(t)$.

THM				
	$\mathsf{R}_{\mathsf{xx}}(t,u)$	$=$	$\mathsf{R}^*_{\mathsf{xx}}(u,t)$	(CONJUGATE SYMMETRIC)
	$\mathsf{R}_{\mathsf{xy}}(t,u)$	$=$	$\mathsf{R}^*_{\mathsf{yx}}(u,t)$	

✎PROOF:

$$\begin{aligned}
\mathsf{R}_{\mathsf{xx}}(t,u) &\triangleq \mathsf{E}\left[\mathsf{x}(t)\mathsf{x}^*(u)\right] &&= \mathsf{E}\left[\mathsf{x}^*(u)\mathsf{x}(t)\right] = \left(\mathsf{E}\left[\mathsf{x}(u)\mathsf{x}^*(t)\right]\right)^* &&\triangleq \mathsf{R}^*_{\mathsf{xx}}(u,t)\\
\mathsf{R}_{\mathsf{xy}}(t,u) &\triangleq \mathsf{E}\left[\mathsf{x}(t)\mathsf{y}^*(u)\right] &&= \mathsf{E}\left[\mathsf{y}^*(u)\mathsf{x}(t)\right] = \left(\mathsf{E}\left[\mathsf{y}(u)\mathsf{x}^*(t)\right]\right)^* &&\triangleq \mathsf{R}^*_{\mathsf{yx}}(u,t)
\end{aligned}$$

✎

3.2 Eigen-analysis of random processes

Definition 3.3. *Let* $\mathsf{x}(t)$ *be random processes with* AUTO-CORRELATION *function* *(Definition 3.2 page 18)* $\mathsf{R}_{\mathsf{xx}}(t,u)$.

DEF	
	The ***auto-correlation operator*** $\mathbf{R}$ *of* $\mathsf{x}(t)$ *is defined as* $\mathbf{R}\mathsf{f} \triangleq \int_{u\in\mathbb{R}} \mathsf{R}_{\mathsf{xx}}(t,u)\mathsf{f}(u)\,\mathrm{d}u$

[4] Papoulis (1984) page 216 $\langle R_{xy}(t_1,t_2) = E\left\{\mathsf{x}(t_1)\mathsf{y}^*(t_2)\right\}$ (9-35)$\rangle$,

[5] Fredholm (1900), Fredholm (1903), page 365, Michel and Herget (1993), page 97, Keener (1988), page 101

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

Definition 3.4. *Let* $\mathsf{x}(t)$ *be a* RANDOM PROCESS *with* AUTO-CORRELATION $\mathsf{R}_{\mathsf{xx}}(\tau)$ *(Definition 3.2 page 18)*.

DEF

A RANDOM PROCESS $\mathsf{x}(t)$ *is* ***white*** *if* $\mathsf{R}_{\mathsf{xx}}(\tau) = \delta(\tau)$

If a random process $\mathsf{x}(t)$ is *white* (Definition 3.4 page 19) and the set $\Psi = \{\psi_1(t), \psi_2(t), \ldots, \psi_N(t)\}\}$ is **any** set of orthonormal basis functions, then the innerproducts $\langle n(t) \mid \psi_n(t)\rangle$ and $\langle n(t) \mid \psi_m(t)\rangle$ are *uncorrelated* for $m \neq n$. However, if $\mathsf{x}(t)$ is **colored** (not white), then the innerproducts are not in general uncorrelated. But if the elements of Ψ are chosen to be the eigenfunctions of $\mathbf{R}$ such that $\mathbf{R}\psi_n = \lambda_n\psi_n$, then by Theorem 3.1 (page 18), the set $\{\psi_n(t)\}$ are *orthogonal* and the innerproducts **are** *uncorrelated* even though $\mathsf{x}(t)$ is not white. This criterion is called the Karhunen-Loève criterion for $\mathsf{x}(t)$.

Theorem 3.2. *Let* $\mathbf{R}$ *be an* AUTO-CORRELATION *operator.*

THM

$$\langle \mathbf{R}\mathsf{x} \mid \mathsf{x}\rangle \geq 0 \qquad \forall \mathsf{x} \in \boldsymbol{X} \qquad \text{(NON-NEGATIVE)}$$
$$\langle \mathbf{R}\mathsf{x} \mid \mathsf{y}\rangle = \langle \mathsf{x} \mid \mathbf{R}\mathsf{y}\rangle \qquad \forall \mathsf{x}, \mathsf{y} \in \boldsymbol{X} \qquad \text{(SELF-ADJOINT)}$$

✎PROOF:

1. Proof that $\mathbf{R}$ is *non-negative*:

$$\begin{aligned}
\langle \mathbf{R}\mathsf{y} \mid \mathsf{y}\rangle &= \left\langle \int_{u\in\mathbb{R}} \mathsf{R}_{\mathsf{xx}}(t,u)\mathsf{y}(u)\,\mathrm{d}u \mid \mathsf{y}(t)\right\rangle && \text{by definition of } \mathbf{R} && \text{(Definition 3.3 page 18)}\\
&= \left\langle \int_{u\in\mathbb{R}} \mathsf{E}\left[\mathsf{x}(t)\mathsf{x}^*(u)\right]\mathsf{y}(u)\,\mathrm{d}u \mid \mathsf{y}(t)\right\rangle && \text{by definition of } \mathsf{R}_{\mathsf{xx}}(t,u) && \text{(Definition 3.2 page 18)}\\
&= \mathsf{E}\left[\left\langle \int_{u\in\mathbb{R}} \mathsf{x}(t)\mathsf{x}^*(u)\mathsf{y}(u)\,\mathrm{d}u \mid \mathsf{y}(t)\right\rangle\right] && \text{by } \textit{linearity} \text{ of } \langle\triangle \mid \triangledown\rangle \text{ and } \int && \text{(Definition J.1 page 255)}\\
&= \mathsf{E}\left[\int_{u\in\mathbb{R}} \mathsf{x}^*(u)\mathsf{y}(u)\,\mathrm{d}u\,\langle \mathsf{x}(t) \mid \mathsf{y}(t)\rangle\right] && \text{by } \textit{additivity} \text{ property of } \langle\triangle \mid \triangledown\rangle && \text{(Definition J.1 page 255)}\\
&= \mathsf{E}\left[\langle \mathsf{y}(u) \mid \mathsf{x}(u)\rangle\,\langle \mathsf{x}(t) \mid \mathsf{y}(t)\rangle\right] && \text{by local definition of } \langle\triangle \mid \triangledown\rangle && \text{(Definition J.1 page 255)}\\
&= \mathsf{E}\left[\langle \mathsf{x}(u) \mid \mathsf{y}(u)\rangle^*\,\langle \mathsf{x}(t) \mid \mathsf{y}(t)\rangle\right] && \text{by } \textit{conjugate symmetry} \text{ prop.} && \text{(Definition J.1 page 255)}\\
&= \mathsf{E}|\langle \mathsf{x}(t) \mid \mathsf{y}(t)\rangle|^2 && \text{by definition of } |\cdot| && \text{(Definition F.4 page 206)}\\
&\geq 0
\end{aligned}$$

2. Proof that $\mathbf{R}$ is self-adjoint:

$$\begin{aligned}
\langle [\mathbf{R}\mathsf{x}](t) \mid \mathsf{y}\rangle &= \left\langle \int_{u\in\mathbb{R}} \mathsf{R}_{\mathsf{xx}}(t,u)\mathsf{x}(u)\,\mathrm{d}u \mid \mathsf{y}(t)\right\rangle && \text{by definition of } \mathbf{R} && \text{(Definition 3.3 page 18)}\\
&= \int_{u\in\mathbb{R}} \mathsf{x}(u)\,\langle \mathsf{R}_{\mathsf{xx}}(t,u) \mid \mathsf{y}(t)\rangle\,\mathrm{d}u && \text{by } \textit{additive} \text{ property of } \langle\triangle \mid \triangledown\rangle && \text{(Definition J.1 page 255)}\\
&= \int_{u\in\mathbb{R}} \mathsf{x}(u)\,\langle \mathsf{y}(t) \mid \mathsf{R}_{\mathsf{xx}}(t,u)\rangle^*\,\mathrm{d}u && \text{by } \textit{conjugate symmetry} \text{ prop.} && \text{(Definition J.1 page 255)}\\
&= \langle \mathsf{x}(u) \mid \langle \mathsf{y}(t) \mid \mathsf{R}_{\mathsf{xx}}(t,u)\rangle\rangle && \text{by local definition of } \langle\triangle \mid \triangledown\rangle && \text{(Definition J.1 page 255)}\\
&= \left\langle \mathsf{x}(u) \mid \int_{t\in\mathbb{R}} \mathsf{y}(t)\mathsf{R}^*_{\mathsf{xx}}(t,u)\,\mathrm{d}t\right\rangle && \text{by local definition of } \langle\triangle \mid \triangledown\rangle && \text{(Definition J.1 page 255)}\\
&= \left\langle \mathsf{x}(u) \mid \int_{t\in\mathbb{R}} \mathsf{y}(t)\mathsf{R}_{\mathsf{xx}}(u,t)\,\mathrm{d}t\right\rangle && \text{by property of } \mathsf{R}_{\mathsf{xx}} && \text{(Theorem 3.1 page 18)}\\
&= \left\langle \mathsf{x}(u) \mid \underbrace{\mathbf{R}}_{\mathbf{R}^*}\mathsf{y}\right\rangle && \text{by definition of } \mathbf{R} && \text{(Definition 3.3 page 18)}
\end{aligned}$$

$$\Longrightarrow \qquad \mathbf{R} = \mathbf{R}^* \qquad \Longrightarrow \mathbf{R} \text{ is } \textit{self adjoint}$$

COPYRIGHT © 2019 DANIEL J. GREENHOE

3.3 Properties

Theorem 3.3. [6] *Let* $(\!(\lambda_n)\!)_{n\in\mathbb{Z}}$ *be the eigenvalues and* $(\!(\psi_n)\!)_{n\in\mathbb{Z}}$ *be the eigenfunctions of operator* $\mathbf{R}$ *such that* $\mathbf{R}\psi_n = \lambda_n\psi_n$.

THM

1. $\lambda_n \in \mathbb{R}$ *(eigenvalues of* $\mathbf{R}$ *are* REAL*)*
2. $\lambda_n \neq \lambda_m \implies \langle \psi_n \,|\, \psi_m \rangle = 0$ *(eigenfunctions associated with distinct eigenvalues are* ORTHOGONAL*)*
3. $\left\|\psi_n(t)\right\|^2 > 0 \implies \lambda_n \geq 0$ *(eigenvalues are* NON-NEGATIVE*)*
4. $\left\|\psi_n(t)\right\|^2 > 0, \langle \mathbf{R}\mathsf{f} \,|\, \mathsf{f} \rangle > 0 \implies \lambda_n > 0$ *(if* $\mathbf{R}$ *is* POSITIVE DEFINITE*, then eigenvalues are* POSITIVE*)*

✎PROOF:

1. Proof that eigenvalues are *real-valued*: Because $\mathbf{R}$ is self-adjoint, its eigenvalues are real (Theorem N.18 page 321).
2. eigenfunctions associated with distinct eigenvalues are orthogonal: Because $\mathbf{R}$ is self-adjoint, this property follows (Theorem N.18 page 321).
3. Proof that eigenvalues are *non-negative*:

$$\begin{aligned}
0 &\geq \langle \mathbf{R}\psi_n \,|\, \psi_n \rangle && \text{by definition of non-negative definite}\\
&= \langle \lambda_n\psi_n \,|\, \psi_n \rangle && \text{by hypothesis}\\
&= \lambda_n \langle \psi_n \,|\, \psi_n \rangle && \text{by definition of inner-products}\\
&= \lambda_n \left\|\psi_n\right\|^2 && \text{by definition of norm induced by inner-product}
\end{aligned}$$

4. Eigenvalues are *positive* if $\mathbf{R}$ is *positive definite*:

$$\begin{aligned}
0 &> \langle \mathbf{R}\psi_n \,|\, \psi_n \rangle && \text{by definition of } \textit{positive definite}\\
&= \langle \lambda_n\psi_n \,|\, \psi_n \rangle && \text{by hypothesis}\\
&= \lambda_n \langle \psi_n \,|\, \psi_n \rangle && \text{by definition of inner-products}\\
&= \lambda_n \left\|\psi_n\right\|^2 && \text{by definition of norm induced by inner-product}
\end{aligned}$$

✏

Theorem 3.4 (Karhunen-Loève Expansion)**.** [7] *Let* $\mathbf{R}$ *be the* AUTO-CORRELATION OPERATOR *(Definition 3.3 page 18) of a* RANDOM PROCESS $\mathsf{x}(t)$. *Let* $(\!(\lambda_n)\!)_{n\in\mathbb{Z}}$ *be the eigenvalues of* $\mathbf{R}$ *and* $(\!(\psi_n)\!)_{n\in\mathbb{Z}}$ *are the eigenfunctions of* $\mathbf{R}$ *such that* $\mathbf{R}\psi_n = \lambda_n\psi_n$.

THM

$$\underbrace{\left\|\psi_n(t)\right\| = 1}_{\{\psi_n(t)\}\ \textit{are}\ \text{NORMALIZED}} \implies \underbrace{\mathsf{E}\left(\left|\mathsf{x}(t) - \sum_{n\in\mathbb{Z}} \langle \mathsf{x}(t) \,|\, \psi_n(t)\rangle\, \psi_n(t)\right|^2\right) = 0}_{\text{CONVERGENCE IN PROBABILITY}} \qquad (\{\psi_n(t)\}\ \textit{is a}\ \text{BASIS}\ \textit{for}\ \mathsf{x}(t))$$

✎PROOF:

1. Define $\dot{x}_n \triangleq \langle \mathsf{x}(t) \,|\, \psi_n(t) \rangle$

[6] Keener (1988), pages 114–119
[7] Keener (1988), pages 114–119

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

2. Define $\mathbf{R}\mathsf{x}(t) \triangleq \int_{u\in\mathbb{R}} \mathsf{R}_{\mathsf{xx}}(t,u)\mathsf{x}(u)\,\mathrm{d}u$

3. lemma: $\mathsf{E}[\mathsf{x}(t)\mathsf{x}(t)] = \sum_{n\in\mathbb{Z}} \lambda_n \left|\psi_n(t)\right|^2$. Proof:

$$\mathsf{E}[\mathsf{x}(t)\mathsf{x}(t)] = \sum_{n\in\mathbb{Z}} \lambda_n \left|\psi_n(t)\right|^2 \qquad \begin{array}{ll} \text{by} & \textit{non-negative} \text{ property} \\ \text{and} & \textit{Mercer's Theorem} \end{array} \qquad \begin{array}{l} \text{(Theorem 3.2 page 19)} \\ \text{(Theorem D.4 page 192)} \end{array}$$

4. lemma:

$$\begin{aligned}
&\mathsf{E}\left[\mathsf{x}(t)\left(\sum_{n\in\mathbb{Z}} \dot{x}_n\psi_n(t)\right)^*\right] \\
&\triangleq \mathsf{E}\left[\mathsf{x}(t)\left(\sum_{n\in\mathbb{Z}} \int_{u\in\mathbb{R}} \mathsf{x}(u)\psi_n^*(u)\,\mathrm{d}u\,\psi_n(t)\right)^*\right] && \text{by definition of } \dot{x} && \text{(definition 1 page 20)} \\
&= \sum_{n\in\mathbb{Z}} \left(\int_{u\in\mathbb{R}} \mathsf{E}\left[\mathsf{x}(t)\mathsf{x}^*(u)\right]\psi_n(u)\,\mathrm{d}u\right)\psi_n^*(t) && \text{by } \textit{linearity} && \text{(Theorem 1.1 page 4)} \\
&\triangleq \sum_{n\in\mathbb{Z}} \left(\int_{u\in\mathbb{R}} \mathsf{R}_{\mathsf{xx}}(t,u)\psi_n(u)\,\mathrm{d}u\right)\psi_n^*(t) && \text{by definition of } \mathsf{R}_{\mathsf{xx}}(t,u) && \text{(Definition 3.2 page 18)} \\
&\triangleq \sum_{n\in\mathbb{Z}} \left(\mathbf{R}\psi_n(t)\psi_n^*(t)\right) && \text{by definition of } \mathbf{R} && \text{(definition 2 page 21)} \\
&= \sum_{n\in\mathbb{Z}} \lambda_n\psi_n(t)\psi_n^*(t) && \text{by property of } \textit{eigen-system} \\
&= \sum_{n\in\mathbb{Z}} \lambda_n \left|\psi_n(t)\right|^2
\end{aligned}$$

5. lemma:

$$\begin{aligned}
&\mathsf{E}\left[\sum_{n\in\mathbb{Z}} \dot{x}_n\psi_n(t)\left(\sum_{m\in\mathbb{Z}} \dot{x}_m\psi_m(t)\right)^*\right] \\
&\triangleq \mathsf{E}\left[\sum_{n\in\mathbb{Z}} \int_{u\in\mathbb{R}} \mathsf{x}(u)\psi_n^*(u)\,\mathrm{d}u\,\psi_n(t)\left(\sum_{m\in\mathbb{Z}} \int_{v} \mathsf{x}(v)\psi_m^*(v)\,\mathrm{d}v\,\psi_m(t)\right)^*\right] && \text{by definition of } \dot{x} && \text{(definition 1 pa} \\
&= \sum_{n\in\mathbb{Z}}\sum_{m\in\mathbb{Z}} \int_u \left(\int_v \mathsf{E}\left[\mathsf{x}(u)\mathsf{x}^*(v)\right]\psi_m(v)\,\mathrm{d}v\right)\psi_n^*(u)\,\mathrm{d}u\,\psi_n(t)\psi_m^*(t) && \text{by } \textit{linearity} && \text{(Theorem 1.1 p} \\
&= \sum_{n\in\mathbb{Z}}\sum_{m\in\mathbb{Z}} \int_u \left(\int_v \mathsf{R}_{\mathsf{xx}}(u,v)\psi_m(v)\,\mathrm{d}v\right)\psi_n^*(u)\,\mathrm{d}u\,\psi_n(t)\psi_m^*(t) && \text{by definition of } \mathsf{R}_{\mathsf{xx}}(t,u) && \text{(Definition 3.2 p} \\
&= \sum_{n\in\mathbb{Z}}\sum_{m\in\mathbb{Z}} \int_u \left(\mathbf{R}\psi_m(u)\right)\psi_n^*(u)\,\mathrm{d}u\,\psi_n(t)\psi_m^*(t) && \text{by definition of } \mathbf{R} && \text{(definition 2 pa} \\
&= \sum_{n\in\mathbb{Z}}\sum_{m\in\mathbb{Z}} \int_u \left(\lambda_m\psi_m(u)\right)\psi_n^*(u)\,\mathrm{d}u\,\psi_n(t)\psi_m^*(t) && \text{by property of } \textit{eigen-system} \\
&= \sum_{n\in\mathbb{Z}}\sum_{m\in\mathbb{Z}} \lambda_m\left(\int_{u\in\mathbb{R}} \psi_m(u)\psi_n^*(u)\,\mathrm{d}u\right)\psi_n(t)\psi_m^*(t) && \text{by } \textit{linearity} \\
&= \sum_{n\in\mathbb{Z}}\sum_{m\in\mathbb{Z}} \lambda_m \left\|\psi(t)\right\|^2 \bar{\delta}_{mn}\psi_n(t)\psi_m^*(t) && \text{by } \textit{orthogonal} \text{ property} && \text{(Theorem 3.3 p} \\
&= \sum_{n\in\mathbb{Z}}\sum_{m\in\mathbb{Z}} \lambda_m\bar{\delta}_{mn}\psi_n(t)\psi_m^*(t) && \text{by } \textit{normalized} \text{ hypothesis} \\
&= \sum_{n\in\mathbb{Z}} \lambda_n\psi_n(t)\psi_n^*(t) && \text{by definition of } \textit{Kronecker delta } \bar{\delta} && \text{(Definition J.3 p} \\
&= \sum_{n\in\mathbb{Z}} \lambda_n \left|\psi_n(t)\right|^2
\end{aligned}$$

Copyright © 2019 Daniel J. Greenhoe

6. Proof that $\{\psi_n(t)\}$ is a *basis* for $\mathsf{x}(t)$:

$$
\begin{aligned}
&\mathsf{E}\left(\left|\mathsf{x}(t) - \sum_{n\in\mathbb{Z}} \dot{x}_n\psi_n(t)\right|^2\right)\\
&= \mathsf{E}\left(\left[\mathsf{x}(t) - \sum_{n\in\mathbb{Z}} \dot{x}_n\psi_n(t)\right]\left[\mathsf{x}(t) - \sum_{m\in\mathbb{Z}} \dot{x}_m\psi_m(t)\right]^*\right)\\
&= \mathsf{E}\left(\mathsf{x}(t)\mathsf{x}^*(t) - \mathsf{x}(t)\left[\sum_{n\in\mathbb{Z}} \dot{x}_n\psi_n(t)\right]^* - \mathsf{x}^*(t)\sum_{n\in\mathbb{Z}} \dot{x}_n\psi_n(t) + \left[\sum_{n\in\mathbb{Z}} \dot{x}_n\psi_n(t)\right]\left[\sum_{m\in\mathbb{Z}} \dot{x}_m\psi_m(t)\right]^*\right)\\
&= \mathsf{E}\big(\mathsf{x}(t)\mathsf{x}^*(t)\big) - \mathsf{E}\left[\mathsf{x}(t)\left[\sum_{n\in\mathbb{Z}} \dot{x}_n\psi_n(t)\right]^*\right] - \mathsf{E}\left[\mathsf{x}^*(t)\sum_{n\in\mathbb{Z}} \dot{x}_n\psi_n(t)\right] + \mathsf{E}\left[\sum_{n\in\mathbb{Z}} \dot{x}_n\psi_n(t)\left[\sum_{m\in\mathbb{Z}} \dot{x}_m\psi_m(t)\right]^*\right]\\
&\qquad \text{by } \textit{linearity} \text{ of } \mathsf{E} \text{ (Theorem 1.1 page 4)}\\
&= \underbrace{\sum_{n\in\mathbb{Z}} \lambda_n\left|\psi_n(t)\right|^2}_{\text{by (3) lemma}} - \underbrace{\sum_{n\in\mathbb{Z}} \lambda_n\left|\psi_n(t)\right|^2}_{\text{by (4) lemma}} - \underbrace{\left[\sum_{n\in\mathbb{Z}} \lambda_n\left|\psi_n(t)\right|^2\right]^*}_{\text{by (4) lemma}} + \underbrace{\sum_{n\in\mathbb{Z}} \lambda_n\left|\psi_n(t)\right|^2}_{\text{by (5) lemma}}\\
&= 0
\end{aligned}
$$

✏

BY-NC-ND Copyright © 2019 Daniel J. Greenhoe

Part II

Statistical Processing

CHAPTER 4

OPERATIONS ON RANDOM VARIABLES

4.1 Functions of one random variable

Proposition 4.1. *Let* $(\Omega, \mathbb{E}, \mathsf{P})$ *be a probability space, and* X *a* RANDOM VARIABLE *with* CUMULATIVE DISTRIBUTION FUNCTION $\mathsf{c}_{\mathsf{X}}(x)$.

PRP

$$\left\{ \begin{array}{l} \mathsf{X} \textit{ is } \text{UNIFORMLY DISTRIBUTED} \\ \textit{(Definition C.1 page 179)} \end{array} \right\} \iff \mathsf{c}_{\mathsf{X}}(x) = \left\{ \begin{array}{ll} 0 & \textit{for } x < 0 \\ x & \textit{for } 0 < x \leq 1 \\ 1 & \textit{otherwise} \end{array} \right\}$$

Theorem 4.1 (Probability integral transform)**.** [1] *Let* $(\Omega, \mathbb{E}, \mathsf{P})$ *be a probability space. Let* X *be a* RANDOM VARIABLE *with* PROBABILITY DENSITY FUNCTION $\mathsf{p}_{\mathsf{X}}(x)$ *and* CUMULATIVE DISTRIBUTION FUNCTION $\mathsf{c}_{\mathsf{X}}(x)$. *Let* Y *be a* RANDOM VARIABLE CUMULATIVE DISTRIBUTION FUNCTION $\mathsf{c}_{\mathsf{Y}}(y)$.

THM

$$\left\{ \begin{array}{ll} \textit{(1).} & \mathsf{Y} = \mathsf{c}_{\mathsf{X}}(\mathsf{X}) \quad \textit{and} \\ \textit{(2).} & \mathsf{p}_{\mathsf{X}}(x) \textit{ is } \text{CONTINUOUS} \end{array} \right\} \implies \left\{ \begin{array}{l} \mathsf{Y} \textit{ is } \text{UNIFORMLY DISTRIBUTED} \\ \textit{(Definition C.1 page 179)} \end{array} \right\}$$

✎PROOF:

$\mathsf{c}_{\mathsf{Y}}(y)$	$\triangleq \mathsf{P}\{\mathsf{Y} \leq y\}$	by definition of *cdf*	(Definition B.2 page 174)
	$= \mathsf{P}\left\{\mathsf{c}_{\mathsf{X}}(\mathsf{X}) \leq y\right\}$	by hypothesis (1)	
	$= \mathsf{P}\left\{\mathsf{X} \leq \mathsf{c}_{\mathsf{X}}^{-1}(y)\right\}$	by hypothesis (2) and	Proposition A.2 page 168
	$\triangleq \mathsf{c}_{\mathsf{X}}\left[\mathsf{c}_{\mathsf{X}}^{-1}(y)\right]$	by definition of *cdf*	(Definition B.2 page 174)
	$= y$		
$\implies$	Y is *uniformly distributed*	by	Proposition 4.1 page 25

✏

Theorem 4.2 (Inverse probability integral transform)**.** [2] *Let* $(\Omega, \mathbb{E}, \mathsf{P})$ *be a probability space. Let* X *be a* RANDOM VARIABLE *with* PROBABILITY DENSITY FUNCTION $\mathsf{p}_{\mathsf{X}}(x)$ *and* CUMULATIVE DISTRIBUTION FUNCTION $\mathsf{c}_{\mathsf{X}}(x)$. *Let* Y *be a* RANDOM VARIABLE CUMULATIVE DISTRIBUTION FUNCTION $\mathsf{c}_{\mathsf{Y}}(y)$.

THM

$$\left\{ \begin{array}{ll} \textit{(1).} & \mathsf{Y} = \mathsf{c}_{\mathsf{Z}}^{-1}(\mathsf{X}) \quad \textit{and} \\ \textit{(2).} & \mathsf{Y} \textit{ is } \text{UNIFORMLY DISTRIBUTED} \quad \textit{and} \\ \textit{(3).} & \mathsf{p}_{\mathsf{Z}}(z) \textit{ is } \text{CONTINUOUS} \end{array} \right\} \implies \left\{ \begin{array}{l} \mathsf{p}_{\mathsf{Y}}(y) = \mathsf{p}_{\mathsf{Z}}(y) \\ (\mathsf{Y} \textit{ has distribution } \mathsf{p}_{\mathsf{Z}}(y)) \end{array} \right\}$$

[1] Angus (1994), Roussas (2014) page 232 ⟨Theorem 10⟩, Devroye (1986) page 28 ⟨Theorem 2.1⟩
[2] Devroye (1986) page 28 ⟨Theorem 2.1⟩, Balakrishnan and Lai (2009) page 624 ⟨14.2.1 Introduction⟩



Figure 4.1: $Y = f(X)$

✎PROOF:

$c_Y(y) \triangleq P\{Y \le y\}$	by definition of c_Y	(Definition B.2 page 174)
$= P\{c_Z^{-1}(X) \le y\}$	by hypothesis (1)	
$= P\{X \le c_Z(y)\}$	by hypothesis (3) and	Proposition A.2 page 168
$\triangleq c_X[c_Z(y)]$	by definition of c_X	(Definition B.2 page 174)
$= c_Z(y)$	because $0 \le c_Z(y) \le 1$ and by	Proposition 4.1 page 25
$\Longrightarrow \quad p_Y(y) = p_Z(y)$	(Y has the distribution of Z)	

✏

Definition 4.1. [3] *Let $f(x)$ be a* DIFFERENTIABLE FUNCTION *in $\mathbb{R}^{\mathbb{R}}$.*

DEF *A point $p \in \mathbb{R}$ is a **critical point** of $f(x)$ if*
$$f'(p) = 0.$$

Theorem 4.3. [4] *Let X and Y be* RANDOM VARIABLES *in $\mathbb{R}^{\mathbb{R}}$. Let f be a* DIFFERENTIABLE FUNCTION *in $\mathbb{R}^{\mathbb{R}}$ with N* CRITICAL POINTS *(Definition 4.1 page 26). Let the range of X be partitioned into $N+1$ partitions $\{A_n | n = 1, 2, \ldots, N+1\}$ with partition boundaries set at the N* CRITICAL POINTS *of $f(x)$—as illustrated in Figure 4.1 (page 26). Let $g_n(x) \triangleq f(x)$ but with domain restricted to $x \in A_n$.*

THM
$$\left\{\begin{array}{ll} (1). & Y = f(X) \quad \textit{and} \\ (2). & f \textit{ is } \text{DIFFERENTIABLE} \end{array}\right\} \Longrightarrow \left\{ p_Y(y) = \sum_{n=1}^{N+1} \frac{p_X\left(g_n^{-1}(y)\right)}{\left|f'\left(g_n^{-1}(y)\right)\right|} \right\}$$

✎PROOF:

1. The problem with a function $f(x)$ with at least $N = 1$ critical point is that $f^{-1}(y)$ is *not invertible*. That is, $f^{-1}(y)$ has more than one solution (and thus the *relation* $f^{-1}(y)$ is not a *function*). However, note that in each partition A_n, $f(x)$ is *invertible* and thus $f^{-1}(y)$ in that partition has a *unique* solution. Thus, each $g_n(x)$ *is* invertible in it's domain (and each $g_n^{-1}(y)$ exists as a function).

[3] Callahan (2010) page 189 ⟨Definition 6.1⟩

[4] Papoulis (1984) pages 95–96 ⟨"Fundamental Theorem"⟩, Papoulis (1990) page 157 ⟨"Fundamental Theorem"⟩, Papoulis (1991), page 93, Haykin (1994) page 235 ⟨0471571768⟩§"4.5 TRANSFORMATIONS OF RANDOM VARIABLES", Proakis (2001), page 30,

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

2. Using item (1), the remainder of the proof follows …

$$
\begin{aligned}
\mathsf{p}_{\mathsf{Y}}(y) &\triangleq \frac{\mathsf{d}}{\mathsf{d}y}\mathsf{P}\{\mathsf{Y} \le y\} && \text{by definition of } \mathsf{p}_{\mathsf{Y}} \text{ (Definition B.2 page 174)}\\
&= \frac{\mathsf{d}}{\mathsf{d}y}\mathsf{P}\{\mathsf{f}(\mathsf{X}) \le y\} && \text{by hypothesis (1)}\\
&= \frac{\mathsf{d}}{\mathsf{d}y}\sum_{n=1}^{N+1}\mathsf{P}\left\{\mathsf{f}(\mathsf{X}) \le y | \mathsf{X} \in A_n\right\} && \text{by } \textit{sum of products} \text{ (Theorem A.3 page 167)}\\
&= \frac{\mathsf{d}}{\mathsf{d}y}\sum_{n=1}^{N+1}\mathsf{P}\left\{\mathsf{f}(\mathsf{X}) \le y \mid \mathsf{X} \in A_n\right\}\mathsf{P}\left\{\mathsf{X} \in A_n\right\} && \text{by definition of } \mathsf{P}\{X|Y\} \text{ (Definition A.4 page 166)}\\
&= \frac{\mathsf{d}}{\mathsf{d}y}\sum_{n=1}^{N+1}\mathsf{P}\left\{\mathsf{g}_n(\mathsf{X}) \le y \,\middle|\, \mathsf{X} \in A_n\right\}\mathsf{P}\left\{\mathsf{X} \in A_n\right\} && \text{by definition of } \mathsf{g}_n(x)\\
&= \frac{\mathsf{d}}{\mathsf{d}y}\left\{\begin{array}{ll}\sum_{n=1}^{N+1}\mathsf{P}\left\{\mathsf{X} \le \mathsf{g}_n^{-1}(y) \,\middle|\, \mathsf{X} \in A_n\right\}\mathsf{P}\left\{\mathsf{X} \in A_n\right\} & \text{for } \mathsf{f}'(x) \ge 0\\ \sum_{n=1}^{N+1}\mathsf{P}\left\{\mathsf{X} \ge \mathsf{g}_n^{-1}(y) \,\middle|\, \mathsf{X} \in A_n\right\}\mathsf{P}\left\{\mathsf{X} \in A_n\right\} & \text{otherwise}\end{array}\right\} && \text{by item (1)}\\
&= \frac{\mathsf{d}}{\mathsf{d}y}\left\{\begin{array}{ll}\sum_{n=1}^{N+1}\mathsf{P}\left\{\mathsf{X} \le \mathsf{g}_n^{-1}(y) \,\middle|\, \mathsf{X} \in A_n\right\}\mathsf{P}\left\{\mathsf{X} \in A_n\right\} & \text{for } \mathsf{f}'(x) \ge 0\\ \sum_{n=1}^{N+1}\left[1-\mathsf{P}\left\{\mathsf{X} \le \mathsf{g}_n^{-1}(y) \,\middle|\, \mathsf{X} \in A_n\right\}\mathsf{P}\left\{\mathsf{X} \in A_n\right\}\right] & \text{otherwise}\end{array}\right\}\\
&= \frac{\mathsf{d}}{\mathsf{d}y}\left\{\begin{array}{ll}\sum_{n=1}^{N+1}\mathsf{P}\left\{\mathsf{X} \le \mathsf{g}_n^{-1}(y) | \mathsf{X} \in A_n\right\} & \text{for } \mathsf{f}'(x) \ge 0\\ \sum_{n=1}^{N+1}\left[1-\mathsf{P}\left\{\mathsf{X} \le \mathsf{g}_n^{-1}(y) | \mathsf{X} \in A_n\right\}\right] & \text{otherwise}\end{array}\right\} && \text{by definition of } \mathsf{P}\{X|Y\}\\
&= \frac{\mathsf{d}}{\mathsf{d}y}\left\{\begin{array}{ll}\sum_{n=1}^{N+1}\left(\mathsf{P}\left\{\mathsf{X} \le \mathsf{g}_n^{-1}(y)\right\}-\mathsf{P}\left\{\mathsf{X} < \min A_{n-1}\right\}\right) & \text{for } \mathsf{f}'(x) \ge 0\\ \sum_{n=1}^{N+1}\left[1-\left(\mathsf{P}\left\{\mathsf{X} \le \mathsf{g}_n^{-1}(y)\right\}-\mathsf{P}\left\{\mathsf{X} < \min A_{n-1}\right\}\right)\right] & \text{otherwise}\end{array}\right\}\\
&= \frac{\mathsf{d}}{\mathsf{d}y}\left\{\begin{array}{ll}\sum_{n=1}^{N+1}\mathsf{P}\left\{\mathsf{X} \le \mathsf{g}_n^{-1}(y)\right\} & \text{for } \mathsf{f}'(x) \ge 0\\ \sum_{n=1}^{N+1}\left[1-\mathsf{P}\left\{\mathsf{X} \le \mathsf{g}_n^{-1}(y)\right\}\right] & \text{otherwise}\end{array}\right\} && \text{because } \frac{\mathsf{d}}{\mathsf{d}y}\mathsf{P}\{\mathsf{X} < \text{a constant}\} = 0\\
&= \left\{\begin{array}{ll}\sum_{n=1}^{N+1}\frac{\mathsf{d}}{\mathsf{d}y}\mathsf{c}_{\mathsf{X}}\left[\mathsf{g}_n^{-1}(y)\right] & \text{for } \mathsf{f}'(x) \ge 0\\ \sum_{n=1}^{N+1}\frac{\mathsf{d}}{\mathsf{d}y}\left[1-\mathsf{c}_{\mathsf{X}}\left(\mathsf{g}_n^{-1}(y)\right)\right] & \text{otherwise}\end{array}\right\} && \text{by } \textit{linearity} \text{ of } \frac{\mathsf{d}}{\mathsf{d}y} \text{ operator}\\
&= \left\{\begin{array}{ll}\sum_{n=1}^{N+1}\mathsf{p}_{\mathsf{X}}\left[\mathsf{g}_n^{-1}(y)\right]\frac{\mathsf{d}}{\mathsf{d}y}\left[\mathsf{g}_n^{-1}(y)\right] & \text{for } \mathsf{f}'(x) \ge 0\\ \sum_{n=1}^{N+1}\left[-\mathsf{p}_{\mathsf{X}}\left[\mathsf{g}_n^{-1}(y)\right]\frac{\mathsf{d}}{\mathsf{d}y}\left[\mathsf{g}_n^{-1}(y)\right]\right] & \text{otherwise}\end{array}\right\} && \text{by definition of } \mathsf{p}_{\mathsf{X}} \text{ (Definition B.2 page 174)}\\
& && \text{and the } \textit{chain rule}
\end{aligned}
$$

Copyright © 2019 Daniel J. Greenhoe



Figure 4.2: $\mathsf{Y} = \frac{1}{\mathsf{X}}$

$$= \sum_{n=1}^{N+1} \mathsf{p}_{\mathsf{X}}\left(\mathsf{g}_n^{-1}(y)\right)\left|\frac{\mathsf{d}}{\mathsf{d}y}\left[\mathsf{g}_n^{-1}(y)\right]\right|$$

$$= \sum_{n=1}^{N+1} \frac{\mathsf{p}_{\mathsf{X}}\left(\mathsf{g}_n^{-1}(y)\right)}{\left|\mathsf{f}'\left(\mathsf{g}_n^{-1}(y)\right)\right|} \qquad \text{by Lemma ?? page ??}$$

✏

Corollary 4.1. [5] *Let* X *and* Y *be* RANDOM VARIABLES *in* $\mathbb{R}^{\mathbb{R}}$. *Let* $a, b \in \mathbb{R}$.

COR
$$\left\{\begin{array}{ll} (1). & \mathsf{Y} = a\mathsf{X} + b \quad \textit{and} \\ (2). & a \neq 0 \end{array}\right\} \Longrightarrow \left\{\mathsf{p}_{\mathsf{Y}}(y) = \frac{1}{|a|}\mathsf{p}_{\mathsf{X}}\left(\frac{y-b}{a}\right)\right\}$$

✎PROOF:

1. Note that $\mathsf{f}(x) = ax + b$ is a *differentiable function* with $N = 0$ *critical points* and $\mathsf{f}'(x) = a$.
2. The inverse of $\mathsf{f}(x)$ is $\mathsf{g}_1(y) = \mathsf{f}^{-1}(y) = \frac{y-b}{a}$.
3. It follows that

$$\mathsf{p}_{\mathsf{Y}}(y) = \sum_{n=1}^{N+1} \frac{\mathsf{p}_{\mathsf{X}}\left(\mathsf{g}_n^{-1}(y)\right)}{\left|\mathsf{f}'\left(\mathsf{g}_n^{-1}(y)\right)\right|} \qquad \text{by Theorem 4.3 (page 26)}$$

$$= \frac{\mathsf{p}_{\mathsf{X}}\left(\mathsf{f}^{-1}(y)\right)}{\left|\mathsf{f}'\left(\mathsf{f}^{-1}(y)\right)\right|} \qquad \text{because } N = 0$$

$$= \frac{\mathsf{p}_{\mathsf{X}}\left(\mathsf{f}^{-1}(y)\right)}{|a|} \qquad \text{by item (1)}$$

$$= \frac{1}{|a|}\mathsf{p}_{\mathsf{X}}\left(\frac{y-b}{a}\right) \qquad \text{by item (2)}$$

✏

[5] Papoulis (1984) page 96 ⟨"Illustrations" 1⟩, Papoulis (1991), page 95, Proakis (2001), page 29

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE



Figure 4.3: $\mathsf{Y} = rvX^2$

Corollary 4.2. [6]

COR $\left\{\mathsf{Y} = \frac{1}{\mathsf{X}}\right\} \implies \left\{\mathsf{p}_{\mathsf{Y}}(y) = \frac{1}{y^2}\mathsf{p}_{\mathsf{X}}\left(\frac{1}{y}\right) \quad \textit{for } y > 0\right\}$

✎PROOF:

1. Note that $\mathsf{f}(x) = 1/x$ is a *differentiable function* in $x > 0$ with $N = 0$ *critical points* and $\mathsf{f}'(x) = -1/x^2$.
2. The inverse of $\mathsf{f}(x)$ is $\mathsf{g}_1(y) = \mathsf{f}^{-1}(y) = \frac{1}{y}$.
3. It follows that

$$\begin{aligned}
\mathsf{p}_{\mathsf{Y}}(y) &= \sum_{n=1}^{N+1} \frac{\mathsf{p}_{\mathsf{X}}\left(\mathsf{g}_n^{-1}(y)\right)}{\left|\mathsf{f}'\left(\mathsf{g}_n^{-1}(y)\right)\right|} && \text{by Theorem 4.3 (page 26)} \\
&= \frac{\mathsf{p}_{\mathsf{X}}\left(\mathsf{f}^{-1}(y)\right)}{\left|\mathsf{f}'\left(\mathsf{f}^{-1}(y)\right)\right|} && \text{because } N = 0 \\
&= \frac{1}{\left|-1/(1/y)^2\right|}\mathsf{p}_{\mathsf{X}}\left(\frac{1}{y}\right) \\
&= \frac{1}{y^2}\mathsf{p}_{\mathsf{X}}\left(\frac{1}{y}\right)
\end{aligned}$$

✏

Corollary 4.3. [7] *Let* X *and* Y *be* RANDOM VARIABLES.

COR $\left\{\mathsf{Y} = \mathsf{X}^2\right\} \implies \left\{\mathsf{p}_{\mathsf{Y}}(y) = \frac{1}{2\sqrt{y}}\left[\mathsf{p}_{\mathsf{X}}(-\sqrt{y}) + \mathsf{p}_{\mathsf{X}}(\sqrt{y})\right]\right\}$

✎PROOF:

1. The roots of $y = x^2$ are $x_1 = -\sqrt{y}$ and $x_2 = +\sqrt{y}$.
2. The derivative of $\mathsf{f}(x) \triangleq y = x^2$ is $\mathsf{f}'(x) = 2x$.

[6] Papoulis (1984) page 97 ⟨Example 5-10⟩, Papoulis (1991), page 94
[7] Papoulis (1984) page 95 ⟨Example 5-9⟩, Devroye (1986) page 27 ⟨Example 4.4⟩, Papoulis (1991), page 95, Proakis (2001), page 29

COPYRIGHT © 2019 DANIEL J. GREENHOE



Figure 4.4: $Z = \tan \Theta$

3. And so it follows that …

$$
\begin{aligned}
\mathsf{p}_Y(y) &= \sum_{n=1}^{N} \frac{\mathsf{p}_X(x_n)}{|\mathsf{f}'(x_n)|} && \text{by Theorem 4.3 page 26} \\
&= \frac{\mathsf{p}_X(x_1)}{|\mathsf{f}'(x_1)|} + \frac{\mathsf{p}_X(x_2)}{|\mathsf{f}'(x_2)|} && \text{by definition of } \sum \\
&= \frac{\mathsf{p}_X(-\sqrt{y})}{|\mathsf{f}'(-\sqrt{y})|} + \frac{\mathsf{p}_X(\sqrt{y})}{|\mathsf{f}'(\sqrt{y})|} && \text{by item (1)} \\
&= \frac{\mathsf{p}_X(-\sqrt{y})}{2\sqrt{y}} + \frac{\mathsf{p}_X(\sqrt{y})}{2\sqrt{y}} && \text{by item (2)} \\
&= \frac{1}{2\sqrt{y}}\left[\mathsf{p}_X(-\sqrt{y}) + \mathsf{p}_X(\sqrt{y})\right] && \text{by } \textit{linearity} \text{ of } + \text{ operation}
\end{aligned}
$$

✏

Corollary 4.4. [8] *Let* $Z = \tan \Theta$. *Then*

COR $\{Z = \tan \Theta\} \quad \Longrightarrow \quad \left\{ \mathsf{p}_Z(z) = \frac{1}{1+z^2} \sum_{n \in Z} \mathsf{p}_\theta(\operatorname{atan}(z) + n\pi) \right\}$

✎PROOF:

1. The roots of $z = \tan\theta$ are $\{\theta_n = \operatorname{atan} z + n\pi | n \in \mathbb{Z}\}$.
2. The derivative of $z = \tan\theta$ is $\mathsf{f}'(\theta) = \sec^2\theta$.
3. It follows that

$$
\begin{aligned}
\mathsf{p}_Z(z) &= \sum_{n=1}^{N} \frac{\mathsf{p}_\theta(\theta_n)}{|\mathsf{f}'(\theta_n)|} \\
&= \sum_n \frac{\mathsf{p}_\theta(\operatorname{atan} z + n\pi)}{|\mathsf{f}'(\operatorname{atan} z + n\pi)|} \\
&= \sum_n \frac{\mathsf{p}_\theta(\operatorname{atan} z + n\pi)}{|\sec^2(\operatorname{atan} z + n\pi)|} \\
&= \sum_n \cos^2(\operatorname{atan} z + n\pi)\mathsf{p}_\theta(\operatorname{atan} z + n\pi)
\end{aligned}
$$

[8] Papoulis (1991), pages 99–100

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

$$= \cos^2(\operatorname{atan} z) \sum_n \mathsf{p}_\theta(\operatorname{atan} z + n\pi)$$

$$= \frac{1}{1+z^2} \sum_n \mathsf{p}_\theta(\operatorname{atan} z + n\pi)$$

✏

4.2 Functions of two random variables

Theorem 4.4. [9] *Let* X, Y, *and* Z *be* RANDOM VARIABLES. *Let* $\star$ *be the* CONVOLUTION *operator* *(Definition O.3 page 336)*.

THM

$$\left\{ \begin{array}{lll} (1). & \mathsf{Z} \triangleq \mathsf{X} + \mathsf{Y} & \textit{and} \\ (2). & \mathsf{X} \textit{ and } \mathsf{Y} \textit{ are } \text{INDEPENDENT} & \textit{(Definition A.3 page 165)} \end{array} \right\} \implies \left\{ \mathsf{p}_\mathsf{Z}(z) = \mathsf{p}_\mathsf{X}(z) \star \mathsf{p}_\mathsf{Y}(z) \right\}$$

✎PROOF:

$$\begin{aligned}
\mathsf{p}_\mathsf{Z}(z) &\triangleq \frac{\mathrm{d}}{\mathrm{d}z}\mathsf{c}_\mathsf{Z}(z) && \text{by definition of } \mathsf{p}_\mathsf{Z} && \text{(Definition B.2 page 174)} \\
&\triangleq \frac{\mathrm{d}}{\mathrm{d}z}\mathsf{P}\{\mathsf{Z} \le z\} && \text{by definition of } \mathsf{c}_\mathsf{Z} && \text{(Definition B.2 page 174)} \\
&= \frac{\mathrm{d}}{\mathrm{d}z}\mathsf{P}\{\mathsf{X} + \mathsf{Y} \le z\} && \text{by hypothesis (1)} \\
&= \frac{\mathrm{d}}{\mathrm{d}z}\lim_{\varepsilon\to 0}\sum_{n\in\mathbb{Z}} \mathsf{P}\left\{\mathsf{X}+\mathsf{Y} \le z \,|\, y + n\varepsilon < \mathsf{Y} \le y + (n+1)\varepsilon\right\} && \text{by } \textit{sum of products} && \text{(Theorem A.3 page 167)} \\
&= \frac{\mathrm{d}}{\mathrm{d}z}\int_{y\in\mathbb{R}} \mathsf{P}\left\{\mathsf{X}+\mathsf{Y} \le z \mid \mathsf{Y} = y\right\} \mathsf{p}_\mathsf{Y}(y)\,\mathrm{d}y && \text{by definiton of } \mathsf{P}\{\mathsf{X} \mid \mathsf{Y}\} && \text{(Definition A.4 page 166)} \\
&= \frac{\mathrm{d}}{\mathrm{d}z}\int_{y\in\mathbb{R}} \mathsf{P}\left\{\mathsf{X} \le z - y \mid \mathsf{Y} = y\right\} \mathsf{p}_\mathsf{Y}(y)\,\mathrm{d}y \\
&= \frac{\mathrm{d}}{\mathrm{d}z}\int_{y\in\mathbb{R}} \mathsf{P}\{\mathsf{X} \le z - y\}\mathsf{p}_\mathsf{Y}(y)\,\mathrm{d}y && \text{by hypothesis (2)} \\
&\triangleq \frac{\mathrm{d}}{\mathrm{d}z}\int_{y\in\mathbb{R}} \mathsf{c}_\mathsf{X}(z-y)\mathsf{p}_\mathsf{Y}(y)\,\mathrm{d}y && \text{by definition of } \mathsf{c}_\mathsf{X} && \text{(Definition B.2 page 174)} \\
&= \int_{y\in\mathbb{R}} \frac{\mathrm{d}}{\mathrm{d}z}\left[\mathsf{c}_\mathsf{X}(z-y)\mathsf{p}_\mathsf{Y}(y)\right]\mathrm{d}y && \text{by } \textit{linearity} \text{ of } \frac{\mathrm{d}}{\mathrm{d}z} \\
&= \int_{y\in\mathbb{R}} \left[\frac{\mathrm{d}}{\mathrm{d}z}\mathsf{c}_\mathsf{X}(z-y)\right]\mathsf{p}_\mathsf{Y}(y)\,\mathrm{d}y && \text{because } y \text{ is fixed inside the integral} \\
&\triangleq \int_{y} \mathsf{p}_\mathsf{X}(z-y)\mathsf{p}_\mathsf{Y}(y)\,\mathrm{d}y && \text{by definition of } \mathsf{p}_\mathsf{X} && \text{(Definition B.2 page 174)} \\
&= \mathsf{p}_\mathsf{X}(z) \star \mathsf{p}_\mathsf{Y}(z) && \text{by definition of } \star && \text{(Definition O.3 page 336)}
\end{aligned}$$

✏

Theorem 4.5. *Let*

- X_1 *and* X_2 *be random variables with joint distribution* $\mathsf{p}_{\mathsf{X}_1,\mathsf{X}_2}(x_1, x_2)$
- $\mathsf{Y}_1 = \mathsf{f}_1(x_1, x_2)$ *and* $\mathsf{Y}_2 = \mathsf{f}_2(x_1, x_2)$

[9] Papoulis (1990) page 160 ⟨Example 5.16⟩

COPYRIGHT © 2019 DANIEL J. GREENHOE



Figure 4.5: **Sum** of random variables yields **convolution** of pdfs (Theorem 4.4 page 31)

Then the joint distribution of Y_1 *and* Y_2 *is*

THM

$$\mathsf{p}_{\mathsf{Y}_1,\mathsf{Y}_2}(y_1,y_2) = \frac{\mathsf{p}_{\mathsf{X}_1,\mathsf{X}_2}(x_1,x_2)}{|J(x_1,x_2)|} = \frac{\mathsf{p}_{\mathsf{X}_1,\mathsf{X}_2}(x_1,x_2)}{\begin{vmatrix} \frac{\partial \mathsf{f}_1}{\partial x_1} & \frac{\partial \mathsf{f}_1}{\partial x_2} \\ \frac{\partial \mathsf{f}_2}{\partial x_1} & \frac{\partial \mathsf{f}_2}{\partial x_2} \end{vmatrix}} = \frac{\mathsf{p}_{\mathsf{X}_1,\mathsf{X}_2}(x_1,x_2)}{\frac{\partial \mathsf{f}_1}{\partial x_1}\frac{\partial \mathsf{f}_2}{\partial x_2} - \frac{\partial \mathsf{f}_1}{\partial x_2}\frac{\partial \mathsf{f}_2}{\partial x_1}}$$

Proposition 4.2. *Let* X *and* Y *be random variables with joint distribution* $\mathsf{p}_{\mathsf{XY}}(x,y)$ *and*

$$R^2 \triangleq X^2 + Y^2 \qquad\qquad \Theta \triangleq \operatorname{atan}\frac{\mathsf{Y}}{\mathsf{X}}.$$

Then

PRP

$$\mathsf{p}_{\mathsf{R},\Theta}(r,\theta) = r\ \mathsf{p}_{\mathsf{XY}}(r\cos\theta, r\sin\theta)$$

✎PROOF:

$$\begin{aligned}
\mathsf{p}_{\mathsf{R},\Theta}(r,\theta) &= \frac{\mathsf{p}_{\mathsf{XY}}(x,y)}{|J(x,y)|} = \frac{\mathsf{p}_{\mathsf{XY}}(x,y)}{\begin{vmatrix} \frac{\partial R}{\partial x} & \frac{\partial R}{\partial y} \\ \frac{\partial \theta}{\partial x} & \frac{\partial \theta}{\partial y} \end{vmatrix}} = \frac{\mathsf{p}_{\mathsf{XY}}(x,y)}{\begin{vmatrix} \frac{x}{\sqrt{x^2+y^2}} & \frac{y}{\sqrt{x^2+y^2}} \\ \frac{-y}{x^2+y^2} & \frac{x}{x^2+y^2} \end{vmatrix}} \\
&= \frac{\mathsf{p}_{\mathsf{XY}}(x,y)}{\frac{x}{\sqrt{x^2+y^2}}\frac{x}{x^2+y^2} - \frac{y}{\sqrt{x^2+y^2}}\frac{-y}{x^2+y^2}} \\
&= \frac{\mathsf{p}_{\mathsf{XY}}(x,y)}{\frac{x^2+y^2}{(x^2+y^2)^{3/2}}} \\
&= \mathsf{p}_{\mathsf{XY}}(x,y)\frac{(x^2+y^2)^{3/2}}{x^2+y^2} \\
&= \mathsf{p}_{\mathsf{XY}}(r\cos\theta, r\sin\theta)\frac{r^3}{r^2} \\
&= r\ \mathsf{p}_{\mathsf{XY}}(r\cos\theta, r\sin\theta)
\end{aligned}$$

✏

Proposition 4.3. *Let* $\mathsf{X} \sim \mathsf{N}\left(0,\sigma^2\right)$ *and* $\mathsf{Y} \sim \mathsf{N}\left(0,\sigma^2\right)$ *be independent random variables and*

$$R^2 \triangleq X^2 + Y^2 \qquad\qquad \Theta \triangleq \operatorname{atan}\frac{\mathsf{Y}}{\mathsf{X}}.$$

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

Then

PRP

1. *R and Θ are independent with joint distribution* $\mathsf{p}_{\mathsf{R},\Theta}(r,\theta) = \mathsf{p}_{\mathsf{R}}(r)\mathsf{p}_{\theta}(\theta)$
2. *R has Rayleigh distribution* $\mathsf{p}_{\mathsf{R}}(r) = \frac{r}{\sigma^2}\exp\frac{r^2}{-2\sigma^2}$
3. *Θ has uniform distribution* $\mathsf{p}_{\theta}(\theta) = \frac{1}{2\pi}$

✎PROOF:

$$\begin{aligned}
\mathsf{p}_{\mathsf{R},\Theta}(r,\theta) &= r\ \mathsf{p}_{\mathsf{XY}}(r\cos\theta, r\sin\theta) && \text{by Proposition 4.2 (page 32)}\\
&= r\ \mathsf{p}_{\mathsf{X}}(r\cos\theta)\ \mathsf{p}_{\mathsf{Y}}(r\sin\theta) && \text{by independence hypothesis}\\
&= r\ \frac{1}{\sqrt{2\pi\sigma^2}}\exp\frac{(r\cos\theta-0)^2}{-2\sigma^2}\frac{1}{\sqrt{2\pi\sigma^2}}\exp\frac{(r\sin\theta-0)^2}{-2\sigma^2}\\
&= \frac{1}{2\pi\sigma^2}\ r\ \exp\frac{r^2(\cos^2\theta+\sin^2\theta)}{-2\sigma^2}\\
&= \frac{1}{2\pi\sigma^2}\ r\ \exp\frac{r^2}{-2\sigma^2}\\
&= \left[\frac{1}{2\pi}\right]\left[\frac{r}{\sigma^2}\exp\frac{r^2}{-2\sigma^2}\right]
\end{aligned}$$

✏

Proposition 4.4. *Let* X *and* Y *be* RANDOM VARIABLES *with covariance* σ_{xy} *on a* PROBABILITY SPACE $(\Omega, \mathbb{E}, \mathsf{P})$.

PRP

$$\left\{\begin{array}{ll}
(A). & \mathsf{X}\ \textit{is}\ \text{GAUSSIAN}\ \textit{with}\ \mathsf{N}\left(\mu_{\mathsf{X}},\sigma_{\mathsf{X}}^2\right)\ \textit{and}\\
(B). & \mathsf{X}\ \textit{is}\ \text{GAUSSIAN}\ \textit{with}\ \mathsf{N}\left(\mu_{\mathsf{Y}},\sigma_{\mathsf{Y}}^2\right)\ \textit{and}\\
(C). & \sigma_{\mathsf{xy}} = \mathrm{cov}\,[\mathsf{X},\mathsf{Y}]
\end{array}\right\} \Longrightarrow \left\{\mathsf{P}\{\mathsf{X}>\mathsf{Y}\} = \mathsf{Q}\left(\frac{-\mu_{\mathsf{X}}+\mu_{\mathsf{Y}}}{\sigma_{\mathsf{X}}^2+\sigma_{\mathsf{Y}}^2-2\sigma_{\mathsf{xy}}}\right)\right\}$$

✎PROOF: Because X and Y are jointly Gaussian, their linear combination $\mathsf{Z} = rvX - \mathsf{Y}$ is also Gaussian. A Gaussian distribution is completely defined by its mean and variance. So, to determine the distribution of Z, we just have to determine the mean and variance of Z.

$$\begin{aligned}
\mathsf{E}Z &= \mathsf{E}\mathsf{X} - \mathsf{E}Y\\
&= \mu_{\mathsf{X}} - \mu_{\mathsf{Y}}
\end{aligned}$$

$$\begin{aligned}
\mathrm{var}\,Z &= \mathsf{E}Z^2 - (\mathsf{E}Z)^2\\
&= \mathsf{E}(\mathsf{X}-\mathsf{Y})^2 - (\mathsf{E}\mathsf{X}-\mathsf{E}Y)^2\\
&= \mathsf{E}(\mathsf{X}^2 - 2XY + Y^2) - [(\mathsf{E}\mathsf{X})^2 - 2\mathsf{E}\mathsf{X}\mathsf{E}Y + (\mathsf{E}Y)^2]\\
&= [\mathsf{E}X^2 - (\mathsf{E}\mathsf{X})^2] + [Y^2 - (\mathsf{E}Y)^2] - 2[\mathsf{E}XY - \mathsf{E}\mathsf{X}\mathsf{E}Y]\\
&= \mathrm{var}\,\mathsf{X} + \mathrm{var}\,Y - 2\mathrm{cov}\,[\mathsf{X},\mathsf{Y}]\\
&\triangleq \sigma_{\mathsf{X}}^2 + \sigma_{\mathsf{Y}}^2 - 2\sigma_{\mathsf{xy}}
\end{aligned}$$

$$\begin{aligned}
\mathsf{P}\{\mathsf{X}>\mathsf{Y}\} &= \mathsf{P}\{\mathsf{X}-\mathsf{Y}>0\}\\
&= \mathsf{P}\{Z>0\}\\
&= \mathsf{Q}\left(\frac{z-\mathsf{E}Z}{\mathrm{var}\,Z}\right)\Bigg|_{z=0}\\
&= \mathsf{Q}\left(\frac{0-\mu_{\mathsf{X}}+\mu_{\mathsf{Y}}}{\sigma_{\mathsf{X}}^2+\sigma_{\mathsf{Y}}^2-2\sigma_{\mathsf{xy}}}\right)
\end{aligned}$$

✏

COPYRIGHT © 2019 DANIEL J. GREENHOE
BY-NC-ND



Figure 4.6: The distributions of sums of independent uniformly distributed random variables (Example 4.1 page 34)

Example 4.1. Let $(\!(X_1, X_2, X_3, \ldots)\!)$ be a *sequence* of *independent* (Definition A.3 page 165) *uniformly distributed* random variables. Let $p_N(x)$ be the *probability density function* of $Y \triangleq \sum_{n=1}^{N} X_n$. Some of these distributions are illustrated in Figure 4.6 (page 34). Note that the distributions of the sequence $(\!(p_1, p_2, p_3, \ldots)\!)$ are all *B-splines* (Definition R.2 page 367) and all form a *partition of unity*.

BY-NC-ND Copyright © 2019 Daniel J. Greenhoe

CHAPTER 5

OPERATORS ON DISCRETE RANDOM SEQUENCES

5.1 LTI operators on random sequences

Theorem 5.1. [1] *Let* $\mathsf{x}(n)$ *be a* RANDOM SEQUENCE *with* MEAN μ_{X} *and* $\mathsf{y}(n)$ *a* RANDOM SEQUENCE *with* MEAN μ_{Y}. *Let* **S** *be a system with* IMPULSE RESPONSE $\mathsf{h}(n)$, INPUT $\mathsf{x}(n)$, *and* OUTPUT $\mathsf{y}(n)$.



THM

$$\left\{ \mathbf{S} \text{ is } (LTI) \right\} \Longrightarrow \left\{ \begin{array}{llll} (1). & \mu_{\mathsf{Y}}(n) & = \displaystyle\sum_{k\in\mathbb{Z}} \mathsf{h}(k)\mu_{\mathsf{X}}(n-k) & \triangleq \mathsf{h}(n) \star \mu_{\mathsf{X}}(n) \quad \text{and} \\ (2). & \mathsf{R}_{\mathsf{xy}}(n,m) & = \displaystyle\sum_{k\in\mathbb{Z}} \mathsf{h}^*(k)\mathsf{R}_{\mathsf{xx}}(n-k,m+k) & \quad \text{and} \\ (3). & \mathsf{R}_{\mathsf{yy}}(n,m) & = \displaystyle\sum_{k\in\mathbb{Z}} \mathsf{h}^*(k)\mathsf{R}_{\mathsf{yx}}(n-k,m+k) & \end{array} \right\}$$

✎PROOF:

$$\begin{aligned}
\mu_{\mathsf{Y}}(n) &\triangleq \mathsf{E}\left[\mathsf{y}(n)\right] && \text{by definition of } \mu_{\mathsf{Y}} && \text{(Definition 2.2 page 11)} \\
&= \mathsf{E}\left[\sum_{k\in\mathbb{Z}} \mathsf{h}(k)\mathsf{x}(n-k)\right] && \text{by } LTI \text{ hypothesis} \\
&= \sum_{k\in\mathbb{Z}} \mathsf{h}(k)\mathsf{E}[\mathsf{x}(n-k)] && \text{by } \textit{linear} \text{ property} \\
&= \sum_{k\in\mathbb{Z}} \mathsf{h}(k)\mu_{\mathsf{X}}(n-k) && \text{by definition of } \mu_{\mathsf{X}} && \text{(Definition 2.2 page 11)} \\
&\triangleq \mathsf{h}(n) \star \mu_{\mathsf{X}}(n) && \text{by definition of } \textit{convolution} && \text{(Definition Q.3 page 353)}
\end{aligned}$$

$$\begin{aligned}
\mathsf{R}_{\mathsf{xy}}(n,m) &\triangleq \mathsf{E}\left[\mathsf{x}(n+m)\mathsf{y}^*(n)\right] && \text{by definition of } \mathsf{R}_{\mathsf{xy}}(n,m) && \text{(Definition 2.2 page 11)} \\
&= \mathsf{E}\left[\mathsf{x}(n+m)(\mathsf{h}(n) \star \mathsf{x}(n))^*\right] && \text{by } LTI \text{ hypothesis} \\
&\triangleq \mathsf{E}\left[\mathsf{x}(n+m)\left(\sum_{k\in\mathbb{Z}} h(k)\mathsf{x}(n-k)\right)^*\right] && \text{by definition of } \textit{convolution} \star && \text{(Definition Q.3 page 353)} \\
&= \mathsf{E}\left[\mathsf{x}(n+m)\sum_{k\in\mathbb{Z}} \mathsf{h}^*(k)\mathsf{x}^*(n-k)\right] && \text{by } \textit{distributive} \text{ property of } *\textbf{-algebras} && \text{(Definition I.3 page 250)}
\end{aligned}$$

[1] Papoulis (1991), page 310

$$= \mathsf{E}\left[\sum_{k\in\mathbb{Z}} \mathsf{h}^*(k)\mathsf{x}(n+m)\mathsf{x}^*(n-k)\right]$$ by *distributive* property of $(\mathbb{C},+,\cdot,0,1)$ (Definition F.5 page 206)

$$= \sum_{k\in\mathbb{Z}} \mathsf{h}^*(k)\mathsf{E}\left[\mathsf{x}(n-k+k+m)\mathsf{x}^*(n-k)\right]$$ by *linear* property of E (Theorem 1.1 page 4)

$$\triangleq \sum_{k\in\mathbb{Z}} \mathsf{h}^*(k)\mathsf{R}_{\mathsf{xx}}(n-k,m+k)$$ by definition of $\mathsf{R}_{\mathsf{xx}}(n,m)$ (Definition 2.2 page 11)

$$\mathsf{R}_{\mathsf{yy}}(n,m) \triangleq \mathsf{E}\left[\mathsf{y}(n+m)\mathsf{y}^*(n)\right]$$ by definition of $\mathsf{R}_{\mathsf{xy}}(n,m)$ (Definition 2.2 page 11)

$$= \mathsf{E}\left[\mathsf{y}(n+m)(\mathsf{h}(n)\star\mathsf{x}(n))^*\right]$$ by *LTI* hypothesis

$$\triangleq \mathsf{E}\left[\mathsf{y}(n+m)\left(\sum_{k\in\mathbb{Z}} \mathsf{h}(k)\mathsf{x}(n-k)\right)^*\right]$$ by definition of *convolution* (Definition Q.3 page 353)

$$= \mathsf{E}\left[\mathsf{y}(n+m)\sum_{k\in\mathbb{Z}} \mathsf{h}^*(k)\mathsf{x}^*(n-k)\right]$$ by *distributive* property of $*$-**algebras** (Definition I.3 page 250)

$$= \mathsf{E}\left[\sum_{k\in\mathbb{Z}} \mathsf{h}^*(k)\mathsf{y}(n+m)\mathsf{x}^*(n-k)\right]$$ by *distributive* property of $(\mathbb{C},+,\cdot,0,1)$ (Definition F.5 page 206)

$$= \sum_{k\in\mathbb{Z}} \mathsf{h}^*(k)\mathsf{E}\left[\mathsf{y}(n-k+k+m)\mathsf{x}^*(n-k)\right]$$ by *linear* property of E (Theorem 1.1 page 4)

$$\triangleq \sum_{k\in\mathbb{Z}} \mathsf{h}^*(k)\mathsf{R}_{\mathsf{yx}}(n-k,m+k)$$ by definition of $\mathsf{R}_{\mathsf{xy}}(n,m)$ (Definition 2.2 page 11)

✏

5.2 LTI operators on WSS random sequences

Corollary 5.1. *Let* $\mathbf{S}$ *be the system defined in Theorem 5.1 (page 35).*

COR

$$\left.\begin{array}{ll}(A). & \mathbf{S}\ \textit{is}\ \text{LTI} \quad \textit{and}\\ (B). & \mathsf{x}(n)\ \textit{is}\ \text{WSS}\end{array}\right\} \Longrightarrow \left\{\begin{array}{lllll}(1). & \mu_{\mathsf{Y}} & = & \mu_{\mathsf{X}}\sum_{n\in\mathbb{Z}}\mathsf{h}(k) & \textit{and}\\ (2). & \mathsf{R}_{\mathsf{xy}}(m) & = & \mathsf{R}_{\mathsf{xx}}(m)\star\mathsf{h}^*(-m) \triangleq \sum_{k\in\mathbb{Z}}\mathsf{h}^*(k)\mathsf{R}_{\mathsf{xx}}(m+k) & \textit{and}\\ (3). & \mathsf{R}_{\mathsf{yy}}(m) & = & \mathsf{R}_{\mathsf{yx}}(m)\star\mathsf{h}^*(-m) \triangleq \sum_{k\in\mathbb{Z}}\mathsf{h}^*(k)\mathsf{R}_{\mathsf{xy}}(m+k) & \textit{and}\\ (4). & \mathsf{R}_{\mathsf{yy}}(m) & = & \mathsf{R}^*_{\mathsf{xx}}(m)\star\mathsf{h}(-m)\star\mathsf{h}^*(-m) & \end{array}\right.$$

✎PROOF:

$$\mu_{\mathsf{Y}} = \mu_{\mathsf{Y}}(n)$$ by Proposition 2.1 page 13 and hypothesis (A)

$$= \sum_{n\in\mathbb{Z}} \mathsf{h}(k)\mu_{\mathsf{X}}(n-k)$$ by Theorem 2.1 page 12 and hypothesis (B)

$$= \sum_{n\in\mathbb{Z}} \mathsf{h}(k)\mu_{\mathsf{X}}(0)$$ by Definition 6.1 page 43 and hypothesis (B)

$$= \mu_{\mathsf{X}}(0)\sum_{n\in\mathbb{Z}} \mathsf{h}(k)$$ by *linear* property of $\sum$

$$= \mu_{\mathsf{X}}\sum_{n\in\mathbb{Z}} \mathsf{h}(k)$$ by Proposition 2.1 page 13

[1] Papoulis (1991), page 323

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

$$
\begin{aligned}
\mathsf{R}_{\mathsf{xy}}(m) &\triangleq \mathsf{R}_{\mathsf{xy}}(0,m) && \text{by Proposition 2.1 page 13} && \text{and hypothesis (A)}\\
&= \sum_{k\in\mathbb{Z}} \mathsf{h}^*(k)\mathsf{R}_{\mathsf{xx}}(0-k,m+k) && \text{by Theorem 5.1 page 35} && \text{and hypothesis (B)}\\
&= \sum_{k\in\mathbb{Z}} \mathsf{h}^*(k)\mathsf{R}_{\mathsf{xx}}(m+k) && \text{by Proposition 2.1 page 13} && \text{and hypothesis (A)}\\
&= \mathsf{h}^*(-m) \star \mathsf{R}_{\mathsf{xx}}(m) && \text{by Proposition Q.2 page 354} \\
\mathsf{R}_{\mathsf{yy}}(m) &\triangleq \mathsf{R}_{\mathsf{yy}}(0,m) && \text{by Proposition 2.1 page 13} && \text{and hypothesis (A)}\\
&= \sum_{k\in\mathbb{Z}} \mathsf{h}^*(k)\mathsf{R}_{\mathsf{yx}}(n-k,m+k) && \text{by Theorem 5.1 page 35} && \text{and hypothesis (B)}\\
&= \sum_{k\in\mathbb{Z}} \mathsf{h}^*(k)\mathsf{R}_{\mathsf{yx}}(m+k) && \text{by Proposition 2.1 page 13} && \text{and hypothesis (A)}\\
&= \mathsf{h}^*(-m) \star \mathsf{R}_{\mathsf{yx}}(m) && \text{by Proposition Q.2 page 354} \\
\mathsf{R}_{\mathsf{yy}}(m) &= \mathsf{h}^*(-m) \star \mathsf{R}_{\mathsf{yx}}(m) && \text{by result (2)} \\
&= \mathsf{h}^*(-m) \star \mathsf{R}^*_{\mathsf{xy}}(m) && \text{by Corollary 2.1 page 14} \\
&= \mathsf{h}^*(-m) \star \left[\mathsf{h}^*(-m) \star \mathsf{R}_{\mathsf{xx}}(m)\right]^* && \text{by result (1)} \\
&= \mathsf{h}^*(-m) \star \mathsf{h}(-m) \star \mathsf{R}^*_{\mathsf{xx}}(m) && \text{by } \textit{distributive} \text{ property of } *\textbf{-algebras} && \text{(Definition I.3 page 250)}
\end{aligned}
$$

Corollary 5.2. [2] *Let* **S** *be a system with* IMPULSE RESPONSE $\mathsf{h}(n)$, INPUT $\mathsf{x}(n)$, *and* OUTPUT $\mathsf{y}(n)$.

COR

$$
\left\{\begin{array}{ll} (A). & \mathsf{h} \textit{ is } \text{LINEAR TIME INVARIANT} \quad \textit{and} \\ (B). & \mathsf{x} \textit{ and } \mathsf{y} \textit{ are } \text{WIDE SENSE STATIONARY} \end{array}\right\} \Longrightarrow \left\{\begin{array}{lll} (1). & \check{\mathsf{S}}_{\mathsf{xy}}(z) = \check{\mathsf{S}}_{\mathsf{xx}}(z)\check{\mathsf{H}}^*\left(\frac{1}{z^*}\right) & \textit{and} \\ (2). & \check{\mathsf{S}}_{\mathsf{yy}}(z) = \check{\mathsf{S}}_{\mathsf{yx}}(z)\check{\mathsf{H}}^*\left(\frac{1}{z^*}\right) & \textit{and} \\ (3). & \check{\mathsf{S}}_{\mathsf{yy}}(z) = \check{\mathsf{S}}_{\mathsf{xx}}(z)\check{\mathsf{H}}(z)\check{\mathsf{H}}^*\left(\frac{1}{z^*}\right) & \end{array}\right\}
$$

✎PROOF: The proof is given in Proposition **??** (page **??**) (1).

Corollary 5.3. *Let* **S** *be a system with* IMPULSE RESPONSE $\mathsf{h}(n)$, INPUT $\mathsf{x}(n)$, *and* OUTPUT $\mathsf{y}(n)$.

COR

$$
\left\{\begin{array}{lll} (A). & \mathsf{h} \textit{ is} & \text{LTI} \quad \textit{and} \\ (B). & \mathsf{x} \textit{ and } \mathsf{y} \textit{ are} & \text{WSS} \end{array}\right\} \Longrightarrow \left\{\begin{array}{lll} (1). & \tilde{\mathsf{S}}_{\mathsf{xy}}(\omega) = \tilde{\mathsf{S}}_{\mathsf{xx}}(\omega)\tilde{\mathsf{H}}^*(\omega) & \textit{and} \\ (2). & \tilde{\mathsf{S}}_{\mathsf{yy}}(\omega) = \tilde{\mathsf{S}}_{\mathsf{xy}}(\omega)\tilde{\mathsf{H}}(\omega) & \textit{and} \\ (3). & \tilde{\mathsf{S}}_{\mathsf{yy}}(\omega) = \tilde{\mathsf{S}}_{\mathsf{xx}}(\omega)\left|\tilde{\mathsf{H}}(\omega)\right|^2 & \end{array}\right\}
$$

✎PROOF: The proof is given in Proposition **??** (page **??**) (1).

5.3 Parallel operators on WSS random sequences

Theorem 5.2. *Let* **S** *be the* SYSTEM *illustrated to the right, where* **T** *is* NOT NECESSARILY LINEAR. *Let*

$$(\!(\mathsf{h}(n))\!) \triangleq \mathbf{H}\bar{\delta}(n) \triangleq \sum_{m\in\mathbb{Z}} \mathsf{h}(m)\bar{\delta}(n-m)$$

be the IMPULSE RESPONSE *of* **H**.



[2] Papoulis (1991), page 323

COPYRIGHT © 2019 DANIEL J. GREENHOE

THM

$$\left\{\begin{array}{lll}\text{(A).} & \mathsf{x}(n) \text{ is} & \textit{WSS} \quad \text{and} \\ \text{(B).} & \mathbf{H} \text{ is} & \textit{LTI}\end{array}\right\} \Longrightarrow \left\{\begin{array}{llll}\text{(1).} & \mathsf{R}_{\mathsf{wy}}(m) & = & \sum_{n\in\mathbb{Z}} \mathsf{h}(n)\mathsf{R}_{\mathsf{xy}}(m-n) \quad (\textit{convolution}) \\ & & \triangleq & \mathsf{h}(m) \star \mathsf{R}_{\mathsf{xy}}(m) \quad \text{and} \\ \text{(2).} & \check{\mathsf{S}}_{\mathsf{wy}}(z) & = & \check{\mathsf{H}}(z)\check{\mathsf{S}}_{\mathsf{xy}}(z) \quad \text{and} \\ \text{(3).} & \tilde{\mathsf{S}}_{\mathsf{wy}}(\omega) & = & \tilde{\mathsf{H}}(\omega)\tilde{\mathsf{S}}_{\mathsf{xy}}(\omega)\end{array}\right\}$$

✎PROOF:

$$\begin{aligned}
\mathsf{R}_{\mathsf{wy}}(m) &\triangleq \mathsf{E}\left[\mathsf{w}(m)\mathsf{y}^*(0)\right] && \text{by (A) and definition of } \mathsf{R}_{\mathsf{wy}} && \text{(Definition 2.4 page 12)}\\
&\triangleq \mathsf{E}\left([\mathbf{H}\mathsf{x}](m)\mathsf{y}^*(0)\right) && \text{by definition of } \mathbf{S} && \\
&= \mathbf{H}\mathsf{E}\left(\mathsf{x}(m)\mathsf{y}^*(0)\right) && \text{by } \textit{LTI} \text{ hypothesis} && \text{(B)}\\
&\triangleq \mathbf{H}\mathsf{R}_{\mathsf{xy}}(m) && \text{by definition of } \mathsf{R}_{\mathsf{xy}} && \text{(Definition 2.4 page 12)}\\
&= \sum_{n\in\mathbb{Z}} \mathsf{h}(n)\mathsf{R}_{\mathsf{xy}}(m-n) && \text{by definition of } \mathbf{H} \textit{ impulse response } (\!(\mathsf{h}(n))\!) && \\
&= \left[\mathsf{h}(m) \star \mathsf{R}_{\mathsf{xy}}(m)\right] && \text{by definition of } \textit{convolution} && \text{(Definition Q.3 page 353)}\\
\check{\mathsf{S}}_{\mathsf{wy}}(z) &\triangleq \mathbf{Z}\mathsf{R}_{\mathsf{wy}}(m) && \text{by definition of } \check{\mathsf{S}}_{\mathsf{wy}} && \text{(Definition 2.5 page 14)}\\
&= \left[\mathsf{h}(m) \star \mathsf{R}_{\mathsf{xy}}(m)\right] && \text{by previous result} && \\
&= \check{\mathsf{H}}(z)\check{\mathsf{S}}_{\mathsf{xy}}(z) && \text{by } \textit{Convolution Theorem} && \text{(Theorem Q.2 page 356)}\\
\tilde{\mathsf{S}}_{\mathsf{wy}}(\omega) &\triangleq \check{\mathbf{F}}\mathsf{R}_{\mathsf{wy}}(m) && \text{by definition of } \tilde{\mathsf{S}}_{\mathsf{wy}} && \text{(Definition 6.3 page 44)}\\
&= \left[\mathsf{h}(m) \star \mathsf{R}_{\mathsf{xy}}(m)\right] && \text{by previous result} && \\
&= \tilde{\mathsf{H}}(\omega)\tilde{\mathsf{S}}_{\mathsf{xy}}(\omega) && \text{by } \textit{Convolution Theorem} && \text{(Theorem Q.2 page 356)}
\end{aligned}$$

✏

5.4 Whitening discrete random sequences



Figure 5.1: Poles (×) and zeros (o) of a *minimum phase* filter

Definition 5.1. *Let $\check{\mathsf{H}}(z)$ be the z-transform of the impulse response of a filter. If $\check{\mathsf{H}}(z)$ can be expressed as a rational expression with poles and zeros $r_n e^{i\theta_n}$, then the filter is* ***minimum phase*** *if each $r_n < 1$ (all roots lie inside the unit circle in the complex z-plane).*

See Figure 5.1 page 38.

Note that if $\mathsf{L}(z)$ has a root at $z = re^{i\theta}$, then $\mathsf{L}^*(1/z^*)$ has a root at

$$\frac{1}{z^*} = \frac{1}{\left(re^{i\theta}\right)^*} = \frac{1}{re^{-i\theta}} = \frac{1}{r}e^{i\theta}.$$

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

That is, if $\mathsf{L}(z)$ has a root inside the unit circle, then $\mathsf{L}^*(1/z^*)$ has a root directly opposite across the unit circle boundary (see Figure 5.2 page 39). A causal stable filter $\check{\mathsf{H}}(z)$ must have all of its poles inside the unit circle. A minimum phase filter is a filter with both its poles and zeros inside the unit circle. One advantage of a minimum phase filter is that its recipricol (zeros become poles and poles become zeros) is also causal and stable.



Figure 5.2: Mirrored roots in complex-z plane



Figure 5.3: Innovations and whitening filters

The next theorem demonstrates a method for "whitening" a *random sequence* $\mathsf{x}(n)$ with a filter constructed from a decomposition of $\mathsf{R}_{\mathsf{xx}}(m)$. The technique is stated precisely in Theorem 5.3 page 39 and illustrated in Figure 5.3 page 39. Both imply two filters with impulse responses $l(n)$ and $\gamma(n)$. Filter $l(n)$ is referred to as the **innovations filter** (because it generates or "innovates" $\mathsf{x}(n)$ from a white noise process $\mathsf{w}(n)$) and $\gamma(n)$ is referred to as the **whitening filter** because it produces a white noise sequence when the input sequence is $\mathsf{x}(n)$.[3]

Theorem 5.3. *Let* $\mathsf{x}(n)$ *be a WSS* RANDOM SEQUENCE *with auto-correlation* $\mathsf{R}_{\mathsf{xx}}(m)$ *and spectral density* $\check{\mathsf{S}}_{\mathsf{xx}}(z)$. ***If*** $\check{\mathsf{S}}_{\mathsf{xx}}(z)$ *has a* ***rational expression****, then the following are true:*

1. *There exists a rational expression* $\mathsf{L}(z)$ *with minimum phase such that*
$$\check{\mathsf{S}}_{\mathsf{xx}}(z) = \mathsf{L}(z)\mathsf{L}^*\left(\frac{1}{z^*}\right).$$
2. *An LTI filter for which the Laplace transform of the impulse response* $\gamma(n)$ *is*
$$\Gamma(z) = \frac{1}{\mathsf{L}(z)}$$
is both causal and stable.
3. *If* $\mathsf{x}(n)$ *is the input to the filter* $\gamma(n)$*, the output* $\mathsf{y}(n)$ *is a* ***white noise sequence*** *such that*
$$\mathsf{S}_{\mathsf{yy}}(z) = 1 \qquad \mathsf{R}_{\mathsf{yy}}(m) = \bar{\delta}(m).$$

[3] Papoulis (1991), pages 401–402

COPYRIGHT © 2019 DANIEL J. GREENHOE

✎PROOF:

$$
\begin{aligned}
\mathsf{S}_{\mathsf{ww}}(z) &= \Gamma(z)\Gamma^*\left(\frac{1}{z^*}\right)\check{\mathsf{S}}_{\mathsf{xx}}(z)\\
&= \frac{1}{\mathsf{L}(z)}\frac{1}{\mathsf{L}^*\left(\frac{1}{z^*}\right)}\check{\mathsf{S}}_{\mathsf{xx}}(z)\\
&= \frac{1}{\mathsf{L}(z)}\frac{1}{\mathsf{L}^*\left(\frac{1}{z^*}\right)}\mathsf{L}(z)\mathsf{L}^*\left(\frac{1}{z^*}\right)\\
&= 1
\end{aligned}
$$

✏

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

CHAPTER 6

OPERATORS ON CONTINUOUS RANDOM SEQUENCES

6.1 LTI Operations on non-stationary random processes



Figure 6.1: Linear system with random process input and output

Theorem 6.1. [1] *Let* $\mathsf{h} : \mathbb{R} \to \mathbb{C}$ *be the impulse response of a linear time-invariant system and Let* $\mathsf{y}(t) = h(t) \star \mathsf{x}(t) \triangleq \int_{u\in\mathbb{R}} h(u)\mathsf{x}(t-u)\,\mathsf{d}u$ *as illustrated in Figure 6.1 page 41. Then*

THM

Correlation functions

$$\begin{aligned}
\mathsf{R}_{\mathsf{xy}}(t,u) &= \mathsf{R}_{\mathsf{xx}}(t,u) \star h^*(u) &&\triangleq \int_v \mathsf{h}^*(v)\mathsf{R}_{\mathsf{xx}}(t,u-v)\,\mathsf{d}v \\
\mathsf{R}_{\mathsf{yy}}(t,u) &= \mathsf{R}_{\mathsf{xy}}(t,u) \star \mathsf{h}(t) &&\triangleq \int_v \mathsf{h}(v)\mathsf{R}_{\mathsf{xy}}(t-v,u)\,\mathsf{d}v \\
\mathsf{R}_{\mathsf{yy}}(t,u) &= \mathsf{R}_{\mathsf{xx}}(t,u) \star \mathsf{h}(t) \star \mathsf{h}^*(u) &&\triangleq \int_w \mathsf{h}^*(w) \int_v \mathsf{h}(v)\mathsf{R}_{\mathsf{xx}}(t-v,u-w)\,\mathsf{d}v\,\mathsf{d}\omega
\end{aligned}$$

Laplace power spectral density functions

$$\begin{aligned}
\check{S}_{\mathsf{xy}}(s,r) &= \check{S}_{\mathsf{xx}}(s,r)\check{\mathsf{h}}^*(r^*) \\
\check{S}_{\mathsf{yy}}(s,r) &= \check{S}_{\mathsf{xy}}(s,r)\check{\mathsf{h}}(s) \\
\check{S}_{\mathsf{yy}}(s,r) &= \check{S}_{\mathsf{xx}}(s,r)\check{\mathsf{h}}(s)\check{\mathsf{h}}^*(r^*)
\end{aligned}$$

Power spectral density functions

$$\begin{aligned}
\mathsf{S}_{\mathsf{xy}}(f,g) &= \mathsf{S}_{\mathsf{xx}}(f,g)\tilde{\mathsf{h}}^*(-g) \\
\mathsf{S}_{\mathsf{yy}}(f,g) &= \mathsf{S}_{\mathsf{xy}}(f,g)\tilde{\mathsf{h}}(\omega) \\
\mathsf{S}_{\mathsf{yy}}(f,g) &= \mathsf{S}_{\mathsf{xx}}(f,g)\tilde{\mathsf{h}}(\omega)\tilde{\mathsf{h}}^*(-g)
\end{aligned}$$

[1] Papoulis (1991), page 312

✎PROOF:

$$
\begin{aligned}
\mathsf{R}_{\mathsf{xy}}(t,u) &\triangleq \mathsf{E}\big[\mathsf{x}(t)\mathsf{y}^*(u)\big] \\
&= \mathsf{E}\left[\mathsf{x}(t)\left(\int_v \mathsf{h}(v)\mathsf{x}(u-v)\,\mathrm{d}v\right)^*\right] \\
&= \mathsf{E}\left[\mathsf{x}(t)\int_v \mathsf{h}^*(v)\mathsf{x}^*(u-v)\,\mathrm{d}v\right] \\
&= \int_v \mathsf{h}^*(v)\mathsf{E}\big[\mathsf{x}(t)\mathsf{x}^*(u-v)\big]\,\mathrm{d}v \\
&= \int_v \mathsf{h}^*(v)\mathsf{R}_{\mathsf{xx}}(t,u-v)\,\mathrm{d}v \\
&\triangleq \mathsf{R}_{\mathsf{xx}}(t,u) \star h^*(u)
\end{aligned}
$$

$$
\begin{aligned}
\mathsf{R}_{\mathsf{yy}}(t,u) &\triangleq \mathsf{E}\big[\mathsf{y}(t)\mathsf{y}^*(u)\big] \\
&= \mathsf{E}\left[\left(\int_v \mathsf{h}(v)\mathsf{x}(t-v)\,\mathrm{d}v\right)\mathsf{y}^*(u)\right] \\
&= \int_v \mathsf{h}(v)\mathsf{E}\big[\mathsf{x}(t-v)\mathsf{y}^*(u)\big]\,\mathrm{d}v \\
&= \int_v \mathsf{h}(v)\mathsf{R}_{\mathsf{xy}}(t-v,u)\,\mathrm{d}v \\
&\triangleq \mathsf{R}_{\mathsf{xy}}(t,u) \star \mathsf{h}(t)
\end{aligned}
$$

$$
\begin{aligned}
\mathsf{R}_{\mathsf{yy}}(t,u) &\triangleq \mathsf{E}\big[\mathsf{y}(t)\mathsf{y}^*(u)\big] \\
&= \mathsf{E}\left[\left(\int_v \mathsf{h}(v)\mathsf{x}(t-v)\,\mathrm{d}v\right)\left(\int_w \mathsf{h}(w)\mathsf{x}(u-w)\,\mathrm{d}\omega\right)^*\right] \\
&= \int_w \mathsf{h}^*(w)\int_v \mathsf{h}(v)\mathsf{E}\big[\mathsf{x}(t-v)\mathsf{x}^*(u-w)\big]\,\mathrm{d}v\,\mathrm{d}\omega \\
&= \int_w \mathsf{h}^*(w)\int_v \mathsf{h}(v)\mathsf{R}_{\mathsf{xx}}(t-v,u-w)\,\mathrm{d}v\,\mathrm{d}\omega \\
&= \int_w \mathsf{h}^*(w)\big[\mathsf{R}_{\mathsf{xx}}(t,u-w) \star \mathsf{h}(t)\big]\,\mathrm{d}\omega \\
&\triangleq \mathsf{R}_{\mathsf{xx}}(t,u) \star \mathsf{h}(t) \star \mathsf{h}^*(u)
\end{aligned}
$$

$$
\begin{aligned}
\check{S}_{\mathsf{xy}}(s,r) &\triangleq \mathbf{L}\mathsf{R}_{\mathsf{xy}}(t,u) \\
&= \mathbf{L}[\mathsf{R}_{\mathsf{xx}}(t,u) \star \mathsf{h}^*(u)] \\
&= \mathbf{L}[\mathsf{R}_{\mathsf{xx}}(t,u)]\mathbf{L}[\mathsf{h}^*(u)] \\
&= \check{S}_{\mathsf{xx}}(s,r)\int_{u\in\mathbb{R}} \mathsf{h}^*(u)e^{-ru}\,\mathrm{d}u \\
&= \check{S}_{\mathsf{xx}}(s,r)\left[\int_{u\in\mathbb{R}} \mathsf{h}(u)e^{-r^*u}\,\mathrm{d}u\right]^* \\
&= \check{S}_{\mathsf{xx}}(s,r)\check{\mathsf{h}}^*(r^*)
\end{aligned}
$$

$$
\begin{aligned}
\check{S}_{\mathsf{yy}}(s,r) &\triangleq \mathbf{L}\mathsf{R}_{\mathsf{yy}}(t,u) \\
&= \mathbf{L}[\mathsf{R}_{\mathsf{xy}}(t,u) \star \mathsf{h}(t)] \\
&= \mathbf{L}[\mathsf{R}_{\mathsf{xy}}(t,u)]\mathbf{L}[\mathsf{h}(t)] \\
&= \check{S}_{\mathsf{xy}}(s,r)\check{\mathsf{h}}(s)
\end{aligned}
$$

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

$$
\begin{aligned}
&= \check{S}_{xy}(s,r)\check{h}(s)\\
&= \check{S}_{xx}(s,r)\check{h}^*(r^*)\check{h}(s)\\
&= \check{S}_{xx}(s,r)\check{h}(s)\check{h}^*(r^*)
\end{aligned}
$$

$$
\begin{aligned}
S_{xy}(f,g) &\triangleq \tilde{\mathbf{F}}R_{xy}(t,u)\\
&= \tilde{\mathbf{F}}[R_{xx}(t,u) \star h^*(u)]\\
&= \tilde{\mathbf{F}}[R_{xx}(t,u)]\tilde{\mathbf{F}}[h^*(u)]\\
&= S_{xx}(f,g)\int_{u\in\mathbb{R}} h^*(u)e^{-i2\pi gu}\,du\\
&= S_{xx}(f,g)\left[\int_{u\in\mathbb{R}} h(u)e^{i2\pi gu}\,du\right]^*\\
&= S_{xx}(f,g)\left[\int_{u\in\mathbb{R}} h(u)e^{-i2\pi(-g)u}\,du\right]^*\\
&= S_{xx}(f,g)\tilde{h}^*(-g)
\end{aligned}
$$

$$
\begin{aligned}
S_{yy}(f,g) &\triangleq \tilde{\mathbf{F}}R_{yy}(t,u)\\
&= \tilde{\mathbf{F}}[R_{xy}(t,u) \star h(t)]\\
&= \tilde{\mathbf{F}}[R_{xy}(t,u)]\tilde{\mathbf{F}}[h(t)]\\
&= S_{xy}(f,g)\tilde{h}(\omega)
\end{aligned}
$$

$$
\begin{aligned}
&= S_{xy}(f,g)\tilde{h}(\omega)\\
&= S_{xx}(f,g)\tilde{h}^*(-g)\tilde{h}(\omega)
\end{aligned}
$$

6.2 LTI Operations on WSS random processes

Definition 6.1.

DEF

A random process $x(t)$ *is* ***wide sense stationary*** *(WSS) if*

(1). $\mu_X(t)$ *is* CONSTANT *with respect to* t (STATIONARY IN THE MEAN) *and*

(2). $R_{xx}(t+\tau,t)$ *is* CONSTANT *with respect to* t (STATIONARY IN CORRELATION)

If a process $x(t)$ is *wide sense stationary*, mean and correlation are often written μ_X and $R_{xx}(\tau)$, respectively. If a pair of processes $x(t)$ and $y(t)$ are *WSS*, then their cross-correlation is commonly written $R_{xy}(\tau)$.

Definition 6.2. *Let* $x(t)$ *and* $y(t)$ *be WSS random processes.*

DEF

$$
\begin{aligned}
\check{S}_{xx}(s) &\triangleq \mathbf{L}R_{xx}(\tau) \triangleq \int_{\tau\in\mathbb{R}} R_{xx}(\tau)e^{-s\tau}\,d\tau\\
\check{S}_{yy}(s) &\triangleq \mathbf{L}R_{yy}(\tau) \triangleq \int_{\tau\in\mathbb{R}} R_{yy}(\tau)e^{-s\tau}\,d\tau\\
\check{S}_{xy}(s) &\triangleq \mathbf{L}R_{xy}(\tau) \triangleq \int_{\tau\in\mathbb{R}} R_{xy}(\tau)e^{-s\tau}\,d\tau\\
\check{S}_{yx}(s) &\triangleq \mathbf{L}R_{yx}(\tau) \triangleq \int_{\tau\in\mathbb{R}} R_{yx}(\tau)e^{-s\tau}\,d\tau
\end{aligned}
$$

COPYRIGHT © 2019 DANIEL J. GREENHOE

Definition 6.3. *Let* $\mathsf{x}(t)$ *and* $\mathsf{y}(t)$ *be WSS random processes.*

DEF

$$\tilde{\mathsf{S}}_{\mathsf{xx}}(\omega) \triangleq [\tilde{\mathbf{F}}\mathsf{R}_{\mathsf{xx}}(\tau)](\omega) \triangleq \int_{\tau\in\mathbb{R}} \mathsf{R}_{\mathsf{xx}}(\tau)e^{-i\omega\tau}\,\mathrm{d}\tau$$
$$\tilde{\mathsf{S}}_{\mathsf{yy}}(\omega) \triangleq [\tilde{\mathbf{F}}\mathsf{R}_{\mathsf{yy}}(\tau)](\omega) \triangleq \int_{\tau\in\mathbb{R}} \mathsf{R}_{\mathsf{yy}}(\tau)e^{-i\omega\tau}\,\mathrm{d}\tau$$
$$\tilde{\mathsf{S}}_{\mathsf{xy}}(\omega) \triangleq [\tilde{\mathbf{F}}\mathsf{R}_{\mathsf{xy}}(\tau)](\omega) \triangleq \int_{\tau\in\mathbb{R}} \mathsf{R}_{\mathsf{xy}}(\tau)e^{-i\omega\tau}\,\mathrm{d}\tau$$
$$\tilde{\mathsf{S}}_{\mathsf{yx}}(\omega) \triangleq [\tilde{\mathbf{F}}\mathsf{R}_{\mathsf{yx}}(\tau)](\omega) \triangleq \int_{\tau\in\mathbb{R}} \mathsf{R}_{\mathsf{yx}}(\tau)e^{-i\omega\tau}\,\mathrm{d}\tau$$

Definition 6.4. [2] *Let* $\mathsf{x}(t)$ *be a random variable that is* STATIONARY IN THE MEAN *such that* $\mathsf{E}[\mathsf{x}(t)]$ *is constant with respect to* t.

DEF

$\mathsf{x}(t)$ *is* ***ergodic in the mean*** *if*

$$\underbrace{\mathsf{E}[\mathsf{x}(t)]}_{\text{ENSEMBLE AVERAGE}} = \lim_{\tau\to\infty} \underbrace{\frac{1}{2\tau}\int_{-\tau}^{+\tau} \mathsf{x}(t)\,\mathrm{d}t}_{\text{TIME AVERAGE}}$$

Proposition 6.1.

PRP

$$\{\ \mathsf{x}(t) \text{ is NON-STATIONARY}\ \} \implies \{\ \mathsf{x}(t) \text{ is NOT ERGODIC IN THE MEAN}\ \}$$

✎PROOF: If $\mathsf{x}(t)$ is *non-stationary*, then $\mathsf{E}[\mathsf{x}(t)]$ is not constant with time. But $\lim_{\tau\to\infty}\frac{1}{2\tau}\int_{-\tau}^{+\tau}\mathsf{x}(t)\,\mathrm{d}t$ must be a constant (if it is *convergent*). ✏

Definition 6.5. [3] *Let* $\mathsf{x}(t)$ *be a* WIDE SENSE STATIONARY *random process.*

DEF

(1). *The* ***average power*** $\mathsf{P}_{\mathsf{avg}}$ *of* $\mathsf{x}(t)$ *is* $\quad \mathsf{P}_{\mathsf{avg}}\mathsf{x}(t) \triangleq \lim_{\tau\to\infty}\frac{1}{2\tau}\int_{t\in\mathbb{R}} |\mathsf{x}(t)|^2\,\mathrm{d}t$

(2). *The* ***energy spectral density*** $|\tilde{\mathsf{x}}(\omega)|^2$ *of* $\mathsf{x}(t)$ *is* $\quad |\tilde{\mathsf{x}}(\omega)|^2 \triangleq \left|\tilde{\mathbf{F}}\mathsf{x}(t)\right|^2$

Remark 6.1 (**spectral power**). Why does $\tilde{\mathsf{S}}_{\mathsf{xx}}(\omega)$ deserve the name *power spectral density*? This is answered by Theorem 6.2 (next). But to elaborate further, note that $\tilde{\mathsf{S}}_{\mathsf{xx}}$ is the spectral representation of the statistical relationship (the *variance*) between samples of $\mathsf{x}(t)$. For example, if there is no relationship, then $\tilde{\mathsf{S}}_{\mathsf{xx}}(\omega) = 1$. But in the case that $\mathsf{x}(t)$ is *ergodic in the mean*, then $\tilde{\mathsf{S}}_{\mathsf{xx}}$ takes on an additional meaning—it describes the "spectral power" present in $\mathsf{x}(t)$. This is demonstrated by the next theorem.

Theorem 6.2. *Let* $\mathsf{x}(t)$ *be a* RANDOM PROCESS.

THM

$$\left\{\begin{array}{ll} (A). & \mathsf{x}(t) \text{ is ERGODIC IN THE MEAN} \quad \textit{and} \\ (B). & \tilde{\mathsf{x}}(\omega) \text{ EXISTS} \end{array}\right\} \implies \left\{\begin{array}{lll} (1). & \tilde{\mathsf{S}}_{\mathsf{xx}}(\omega) = \lim_{\tau\to\infty}\frac{1}{2\tau}\underbrace{|\tilde{\mathsf{x}}(\omega)|^2}_{\text{(ESD)}} & \textit{and} \\ (2). & \mathsf{P}_{\mathsf{avg}}[\mathsf{x}(t)] = \int_{\omega\in\mathbb{R}} \tilde{\mathsf{S}}_{\mathsf{xx}}(\omega)\,\mathrm{d}\omega & \end{array}\right\}$$

[2] Papoulis (1984) page 246 ⟨Mean-Ergodic processes⟩, Papoulis (2002) page 523 ⟨12-1 ERGODICITY⟩, KAY (1988) PAGE 58 ⟨3.6 ERGODICITY OF THE AUTOCORRELATION FUNCTION⟩, MANOLAKIS ET AL. (2005) PAGE 106 ⟨ERGODIC RANDOM PROCESSES⟩, KOOPMANS (1995) PAGES 53–61, CADZOW (1987) PAGE 378 ⟨11.13 ERGODIC TIME SERIES⟩, HELSTROM (1991) PAGE 336

[3] ? page 177

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

✎PROOF:

$$
\begin{aligned}
\tilde{S}_{xx}(\omega) &\triangleq \int_{\tau\in\mathbb{R}} R_{xx}(\tau)e^{-i\omega\tau}\,d\tau && \text{by definition of } \tilde{S}_{xx}(\omega) && \text{(Definition 6.3 page 44)}\\
&= \int_{\tau\in\mathbb{R}} E\left[x(t+\tau)x^*(t)\right]e^{-i\omega\tau}\,d\tau && \text{by definition of } R_{xx}(t) && \text{(Definition 3.2 page 18)}\\
&= E\left[x^*(t)\int_{\tau\in\mathbb{R}} x(t+\tau)e^{-i\omega\tau}\,d\tau\right] && \text{by } \textit{linearity} \text{ of } E \text{ operator}\\
&= E\left[x^*(t)\int_{u\in\mathbb{R}} x(u)e^{-i\omega(u-t)}\,du\right] && \text{where } u \triangleq t+\tau \implies \tau = u-t\\
&= E\left[x^*(t)e^{i\omega t}\int_{u\in\mathbb{R}} x(u)e^{-i\omega u}\,du\right]\\
&= E\left[x^*(t)e^{i\omega t}\tilde{x}(\omega)\right] && \text{by definition of } \textit{Fourier Transform} && \text{(Definition O.2 page 333)}\\
&= E\left[x^*(t)e^{i\omega t}\right]\tilde{x}(\omega) && \text{by hypothesis (B)}\\
&= \left[\lim_{\tau\to\infty}\frac{1}{2\tau}\int_{-\tau}^{+\tau} x^*(t)e^{i\omega t}\,dt\right]\tilde{x}(\omega) && \text{by } \textit{ergodic in the mean} \text{ hypothesis} && \text{(Definition 6.4 page 44)}\\
&= \lim_{\tau\to\infty}\frac{1}{2\tau}\left[\int_{t\in\mathbb{R}} x(t)e^{-i\omega t}\,dt\right]^*\tilde{x}(\omega)\\
&= \lim_{\tau\to\infty}\frac{1}{2\tau}\tilde{x}^*(\omega)\tilde{x}(\omega) && \text{by hypothesis (B)}\\
&= \lim_{\tau\to\infty}\frac{1}{2\tau}|\tilde{x}(\omega)|^2
\end{aligned}
$$

$$
\begin{aligned}
\int_{\omega\in\mathbb{R}} \tilde{S}_{xx}(\omega)\,d\omega &= \int_{\omega\in\mathbb{R}} \lim_{\tau\to\infty}\frac{1}{2\tau}|\tilde{x}(\omega)|^2\,d\omega\\
&= \lim_{\tau\to\infty}\frac{1}{2\tau}\int_{\omega\in\mathbb{R}} |\tilde{x}(\omega)|^2\,d\omega\\
&= \lim_{\tau\to\infty}\frac{1}{2\tau}\int_{t\in\mathbb{R}} |x(t)|^2\,dt && \text{by } \textit{Plancheral's formula} && \text{(Theorem O.3 page 335, Theorem H.9 page 232)}\\
&= P_{avg} && \text{by definition of } P_{avg} && \text{(Definition 6.5 page 44)}
\end{aligned}
$$

Thus, $\tilde{S}_{xx}(\omega)$ is the power density of $x(t)$ in the frequency domain. ✏



Figure 6.2: Linear system with WSS random process input and output

Theorem 6.3. [4] *Let* $h : \mathbb{R} \to \mathbb{C}$ *be the impulse response of a linear time-invariant system and let* $y(t) = h(t) \star x(t) \triangleq \int_{u\in\mathbb{R}} h(u)x(t-u)\,du$ *as illustrated in Figure 6.1 page 41. Then*

[4] Papoulis (1991), pages 323–324

COPYRIGHT © 2019 DANIEL J. GREENHOE
BY-NC-ND

THM

$$
\begin{aligned}
\mathsf{R}_{\mathsf{xy}}(\tau) &= \mathsf{R}_{\mathsf{xx}}(\tau) \star h^*(-\tau) &&\triangleq \int_{u\in\mathbb{R}} h^*(-u)\mathsf{R}_{\mathsf{xx}}(\tau-u)\,\mathrm{d}u\\
\mathsf{R}_{\mathsf{yy}}(\tau) &= \mathsf{R}_{\mathsf{xy}}(\tau) \star h(\tau) &&\triangleq \int_{u\in\mathbb{R}} h(u)\mathsf{R}_{\mathsf{xy}}(\tau-u)\,\mathrm{d}u\\
\mathsf{R}_{\mathsf{yy}}(\tau) &= \mathsf{R}_{\mathsf{xx}}(\tau) \star h(\tau) \star h^*(-\tau) &&\triangleq \int_{v}\int_{u\in\mathbb{R}} h(u-v)h^*(-v)\mathsf{R}_{\mathsf{xx}}(\tau-u-v)\,\mathrm{d}u\,\mathrm{d}v\\
\\
\mathsf{S}_{\mathsf{xy}}(s) &= \mathsf{S}_{\mathsf{xx}}(s)\hat{h}^*(-s^*)\\
\mathsf{S}_{\mathsf{yy}}(s) &= \mathsf{S}_{\mathsf{xy}}(s)\hat{h}(s)\\
\mathsf{S}_{\mathsf{yy}}(s) &= \mathsf{S}_{\mathsf{xx}}(s)\hat{h}(s)\hat{h}^*(-s^*)\\
\\
\tilde{\mathsf{S}}_{\mathsf{xy}}(\omega) &= \tilde{\mathsf{S}}_{\mathsf{xx}}(\omega)\tilde{\mathsf{h}}^*(\omega)\\
\tilde{\mathsf{S}}_{\mathsf{yy}}(\omega) &= \tilde{\mathsf{S}}_{\mathsf{xy}}(\omega)\tilde{\mathsf{h}}(\omega)\\
\tilde{\mathsf{S}}_{\mathsf{yy}}(\omega) &= \tilde{\mathsf{S}}_{\mathsf{xx}}(\omega)|\tilde{\mathsf{h}}(\omega)|^2
\end{aligned}
$$

✎ PROOF:

$$
\begin{aligned}
\mathsf{R}_{\mathsf{xx}}(\tau) \star h^*(-\tau) &\triangleq \int_{u\in\mathbb{R}} h^*(-u)\mathsf{R}_{\mathsf{xx}}(\tau-u)\,\mathrm{d}u\\
&= \int_{u\in\mathbb{R}} h^*(-u)\mathsf{E}\left[\mathsf{x}(t)\mathsf{x}^*(t-\tau+u)\right]\,\mathrm{d}u\\
&= \mathsf{E}\left[\mathsf{x}(t)\int_{u\in\mathbb{R}} h^*(-u)\mathsf{x}^*(t-\tau+u)\,\mathrm{d}u\right]\\
&= \mathsf{E}\left[\mathsf{x}(t)\int_{u\in\mathbb{R}} h^*(u')\mathsf{x}^*(t-\tau-u')\,\mathrm{d}u'\right]\\
&= \mathsf{E}\left[\mathsf{x}(t)\mathsf{y}^*(t-\tau)\right]\\
&\triangleq \mathsf{R}_{\mathsf{xy}}(\tau)
\end{aligned}
$$

$$
\begin{aligned}
\mathsf{R}_{\mathsf{xy}}(\tau) \star h(\tau) &\triangleq \int_{u\in\mathbb{R}} h(u)\mathsf{R}_{\mathsf{xy}}(\tau-u)\,\mathrm{d}u\\
&= \int_{u\in\mathbb{R}} h(u)\mathsf{E}\left[\mathsf{x}(t+\tau-u)\mathsf{y}^*(t)\right]\,\mathrm{d}u\\
&= \mathsf{E}\left[\mathsf{y}^*(t)\int_{u\in\mathbb{R}} h(u)\mathsf{x}(t+\tau-u)\,\mathrm{d}u\right]\\
&= \mathsf{E}\left[\mathsf{y}^*(t)\mathsf{y}(t+\tau)\right]\\
&= \mathsf{E}\left[\mathsf{y}(t+\tau)\mathsf{y}^*(t)\right]\\
&\triangleq \mathsf{R}_{\mathsf{yy}}(\tau)
\end{aligned}
$$

$$
\begin{aligned}
\mathsf{R}_{\mathsf{yy}}(\tau) &= \mathsf{R}_{\mathsf{xy}}(\tau) \star h(\tau)\\
&= \mathsf{R}_{\mathsf{xx}}(\tau) \star h^*(-\tau) \star h(\tau)\\
&= \mathsf{R}_{\mathsf{xx}}(\tau) \star h(\tau) \star h^*(-\tau)
\end{aligned}
$$

$$
\begin{aligned}
\mathsf{S}_{\mathsf{xy}}(s) &\triangleq \mathbf{L}\mathsf{R}_{\mathsf{xy}}(\tau)\\
&\triangleq \int_{\tau\in\mathbb{R}} \mathsf{R}_{\mathsf{xy}}(\tau)e^{-s\tau}\,d\tau\\
&= \int_{\tau\in\mathbb{R}} \left[\mathsf{R}_{\mathsf{xx}}(\tau) \star h^*(-\tau)\right]e^{-s\tau}\,d\tau\\
&= \int_{\tau\in\mathbb{R}} \left[\int_{u\in\mathbb{R}} h^*(-u)\mathsf{R}_{\mathsf{xx}}(\tau-u)\,\mathrm{d}u\right]e^{-s\tau}\,d\tau\\
&= \int_{u\in\mathbb{R}} h^*(-u)\int_{\tau\in\mathbb{R}} \mathsf{R}_{\mathsf{xx}}(\tau-u)e^{-s\tau}\,d\tau\,\mathrm{d}u
\end{aligned}
$$

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

$$
\begin{aligned}
&= \int_{u\in\mathbb{R}} h^*(-u) \int_v \mathsf{R}_{\mathsf{xx}}(v) e^{-s(v+u)}\,\mathrm{d}v\,\mathrm{d}u && \text{where } v = \tau - u \iff \tau = v + u \\
&= \int_{u\in\mathbb{R}} h^*(-u) e^{-su}\,\mathrm{d}u \int_v \mathsf{R}_{\mathsf{xx}}(v) e^{-sv}\,\mathrm{d}v \\
&= \int_{u\in\mathbb{R}} h^*(u) e^{-s(-u)}\,\mathrm{d}u \int_v \mathsf{R}_{\mathsf{xx}}(v) e^{-sv}\,\mathrm{d}v \\
&= \left(\int_{u\in\mathbb{R}} h(u) e^{-(-s^*)u}\,\mathrm{d}u\right)^* \int_v \mathsf{R}_{\mathsf{xx}}(v) e^{-sv}\,\mathrm{d}v \\
&\triangleq \hat{h}^*(-s^*)\mathsf{S}_{\mathsf{xx}}(s)
\end{aligned}
$$

$$
\begin{aligned}
\mathsf{S}_{\mathsf{yy}}(s) &\triangleq \mathbf{L}\mathsf{R}_{\mathsf{yy}}(\tau) \\
&\triangleq \int_{\tau\in\mathbb{R}} \mathsf{R}_{\mathsf{yy}}(\tau) e^{-s\tau}\,d\tau \\
&= \int_{\tau\in\mathbb{R}} \left[\mathsf{R}_{\mathsf{xy}}(\tau) \star h(\tau)\right] e^{-s\tau}\,d\tau \\
&= \int_{\tau\in\mathbb{R}} \left[\int_{u\in\mathbb{R}} h(u)\mathsf{R}_{\mathsf{xy}}(\tau - u)\,\mathrm{d}u\right] e^{-s\tau}\,d\tau \\
&= \int_{u\in\mathbb{R}} h(u) \int_{\tau\in\mathbb{R}} \mathsf{R}_{\mathsf{xy}}(\tau - u) e^{-s\tau}\,d\tau\,\mathrm{d}u \\
&= \int_{u\in\mathbb{R}} h(u) \int_{\tau\in\mathbb{R}} \mathsf{R}_{\mathsf{xy}}(v) e^{-s(v+u)}\,d\tau\,\mathrm{d}u && \text{where } v = \tau - u \iff \tau = v + u \\
&= \int_{u\in\mathbb{R}} h(u) e^{-su}\,du \int_{\tau\in\mathbb{R}} \mathsf{R}_{\mathsf{xy}}(v) e^{-sv}\,d\tau \\
&\triangleq \hat{h}(s)\mathsf{S}_{\mathsf{xy}}(s)
\end{aligned}
$$

$$
\begin{aligned}
\mathsf{S}_{\mathsf{yy}}(s) &= \hat{h}(s)\mathsf{S}_{\mathsf{xy}}(s) \\
&= \hat{h}(s)\hat{h}^*(-s^*)\mathsf{S}_{\mathsf{xx}}(s)
\end{aligned}
$$

$$
\begin{aligned}
\tilde{\mathsf{S}}_{\mathsf{xy}}(\omega) &= \mathsf{S}_{\mathsf{xy}}(s)\big|_{s=j\omega} \\
&= \hat{h}^*(-s^*)\mathsf{S}_{\mathsf{xx}}(s)\big|_{s=j\omega} \\
&= \left(\int_{u\in\mathbb{R}} h(u) e^{-(-s^*)u}\,\mathrm{d}u\right)^* \int_v \mathsf{R}_{\mathsf{xx}}(v) e^{-sv}\,\mathrm{d}v\Big|_{s=j\omega} \\
&= \left(\int_{u\in\mathbb{R}} h(u) e^{-(-j\omega)^* u}\,\mathrm{d}u\right)^* \int_v \mathsf{R}_{\mathsf{xx}}(v) e^{-j\omega v}\,\mathrm{d}v \\
&= \left(\int_{u\in\mathbb{R}} h(u) e^{-j\omega u}\,\mathrm{d}u\right)^* \int_v \mathsf{R}_{\mathsf{xx}}(v) e^{-j\omega v}\,\mathrm{d}v \\
&\triangleq \tilde{\mathsf{h}}^*(\omega)\tilde{\mathsf{S}}_{\mathsf{xx}}(\omega)
\end{aligned}
$$

$$
\begin{aligned}
\tilde{\mathsf{S}}_{\mathsf{yy}}(\omega) &= \mathsf{S}_{\mathsf{yy}}(s)\big|_{s=j\omega} \\
&= \hat{h}(s)\mathsf{S}_{\mathsf{xy}}(s)\big|_{s=j\omega} \\
&= \int_{u\in\mathbb{R}} h(u) e^{-su}\,du \int_{\tau\in\mathbb{R}} \mathsf{R}_{\mathsf{xy}}(v) e^{-sv}\,d\tau\Big|_{s=j\omega} \\
&= \int_{u\in\mathbb{R}} h(u) e^{-j\omega u}\,du \int_{\tau\in\mathbb{R}} \mathsf{R}_{\mathsf{xy}}(v) e^{-j\omega v}\,d\tau \\
&= \tilde{\mathsf{h}}(\omega)\tilde{\mathsf{S}}_{\mathsf{xy}}(\omega)
\end{aligned}
$$

Copyright © 2019 Daniel J. Greenhoe

$$\begin{aligned}\tilde{S}_{yy}(\omega) &= \tilde{h}(\omega)\tilde{S}_{xy}(\omega)\\ &= \tilde{h}(\omega)\tilde{h}^*(\omega)\tilde{S}_{xx}(\omega)\\ &= |\tilde{h}(\omega)|^2\tilde{S}_{xx}(\omega)\end{aligned}$$

✏



Figure 6.3: *auto-correlation* $R_{xx}(\tau)$

Theorem 6.4. *Let* $x : \mathbb{R} \to \mathbb{C}$ *be a WSS random process with auto-correlation* $R_{xx}(\tau)$. *Then* $R_{xx}(\tau)$ *is* ***conjugate symmetric*** *such that (see Figure 6.3 page 48)*

THM

$$\begin{aligned}R_{xx}(\tau) &= R^*_{xx}(-\tau) && \text{(CONJUGATE SYMMETRIC)}\\ \mathbf{R}_e\left[R_{xx}(\tau)\right] &= \mathbf{R}_e\left[R_{xx}(-\tau)\right] && \text{(SYMMETRIC)}\\ \mathbf{I}_m\left[R_{xx}(\tau)\right] &= -\mathbf{I}_m\left[R_{xx}(-\tau)\right] && \text{(ANTI-SYMMETRIC)}\\ |R_{xx}(\tau)| &= |R_{xx}(-\tau)| && \text{(SYMMETRIC)}\\ \angle R_{xx}(\tau) &= \angle R_{xx}(-\tau) && \text{(ANTI-SYMMETRIC).}\end{aligned}$$

✎PROOF:

$$\begin{aligned}R^*_{xx}(\tau) &\triangleq \left(E\left[x(t-\tau)x^*(t)\right]\right)^*\\ &= E\left[x^*(t-\tau)x(t)\right]\\ &= E\left[x(t)x^*(t-\tau)\right]\\ &= E\left[x(u+\tau)x^*(u)\right] && \text{where } u \triangleq t-\tau \iff t = u+\tau\\ &\triangleq R_{xx}(\tau)\end{aligned}$$

$$\begin{aligned}\mathbf{R}_e\left[R_{xx}(\tau)\right] &= \mathbf{R}_e\left[R^*_{xx}(-\tau)\right] &&= \mathbf{R}_e\left[R_{xx}(-\tau)\right]\\ \mathbf{I}_m\left[R_{xx}(\tau)\right] &= \mathbf{I}_m\left[R^*_{xx}(-\tau)\right] &&= -\mathbf{I}_m\left[R_{xx}(-\tau)\right]\\ abs R_{xx}(\tau) &= \left|R^*_{xx}(-\tau)\right| &&= \left|R_{xx}(-\tau)\right|\\ \angle R_{xx}(\tau) &= \angle R^*_{xx}(-\tau) &&= -\angle R_{xx}(-\tau)\end{aligned}$$

✏

6.3 Whitening continuous random sequences

Simple algebraic operations on white noise processes (processes with autocorrelation $R_{xx}(\tau) = \delta(\tau)$) often produce *colored* noise (processes with autocorrelation $R_{xx}(\tau) \neq \delta(\tau)$). Sometimes we would like to process a colored noise process to produce a white noise process. This operation is known as *whitening*. Reasons for why we may want to whiten a noise process include

1. Samples from a white noise process are uncorrelated. If the noise process is Gaussian, then these samples are also independent which often greatly simplifies analysis.

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

2. Any orthonormal basis can be used to decompose a white noise process. This is not true of a colored noise process. Karhunen–Loève expansion can be used to decompose colored noise. [5]

Definition 6.6. *A **rational expression** $\mathsf{p}(s)$ is a polynomial divided by a polynomial such that*

DEF

$$\mathsf{p}(s) = \frac{\sum_{n=0}^{N} b_n s^n}{\sum_{n=0}^{M} a_n s^n}.$$

*The **zeros** of a rational expression are the roots of its numerator polynomial. The **poles** of a rational expression are the roots of its denominator polynomial.*

Definition 6.7. *Let $\check{\mathsf{h}}(s)$ be the Laplace transform of the impulse response of a filter. If $\check{\mathsf{h}}(s)$ can be expressed as a rational expression with poles and zeros at $a_n + ib_n$, then the filter is **minimum phase** if each $a_n < 0$ (all roots lie in the left hand side of the complex s-plane).*

Note that if $L(s)$ has a root at $s = -a + ib$, then $L^*(-s^*)$ has a root at

$$-s^* = -(-a + ib)^* = -(-a - ib) = a + ib.$$

That is, if $L(s)$ has a root in the left hand plane, then $L^*(-s^*)$ has a root directly opposite across the imaginary axis in the right hand plane (see Figure 6.4 page 49). A causal stable filter $\hat{h}(s)$ must have all of its poles in the left hand plane. A minimum phase filter is a filter with both its poles and zeros in the left hand plane. One advantage of a minimum phase filter is that its recipricol (zeros become poles and poles become zeros) is also causal and stable.



Figure 6.4: Mirrored roots in complex-s plane



Figure 6.5: Innovations and whitening filters

The next theorem demonstrates a method for "whitening" a random process $\mathsf{x}(t)$ with a filter constructed from a decomposition of $\mathsf{R}_{\mathsf{xx}}(\tau)$. The technique is stated precisely in Theorem 6.5 page 50

[5] *Karhunen–Loève expansion*: Section 3.2 page 18

Copyright © 2019 Daniel J. Greenhoe
CC BY-NC-ND

and illustrated in Figure 6.5 page 49. Both imply two filters with impulse responses $l(t)$ and $\gamma(t)$. Filter $l(t)$ is referred to as the **innovations filter** (because it generates or "innovates" $\mathsf{x}(t)$ from a white noise process $\mathsf{w}(t)$) and $\gamma(t)$ is referred to as the **whitening filter** because it produces a white noise sequence when the input sequence is $\mathsf{x}(t)$. [6]

Theorem 6.5. *Let $\mathsf{x}(t)$ be a WSS random process with autocorrelation $\mathsf{R}_{\mathsf{xx}}(\tau)$ and spectral density $\mathsf{S}_{\mathsf{xx}}(s)$. **If** $\mathsf{S}_{\mathsf{xx}}(s)$ has a **rational expression**, then the following are true:*

1. *There exists a rational expression $L(s)$ with minimum phase such that*

$$\mathsf{S}_{\mathsf{xx}}(s) = L(s)L^*(-s^*).$$

2. *An LTI filter for which the Laplace transform of the impulse response $\gamma(t)$ is*

$$\Gamma(s) = \frac{1}{L(s)}$$

   is both causal and stable.
3. *If $\mathsf{x}(t)$ is the input to the filter $\gamma(t)$, the output $\mathsf{y}(t)$ is a **white noise sequence** such that*

$$\mathsf{S}_{\mathsf{yy}}(s) = 1 \qquad\qquad \mathsf{R}_{\mathsf{yy}}(\tau) = \delta(\tau).$$

✎PROOF:

$$\begin{aligned}
\mathsf{S}_{\mathsf{ww}}(s) &= \Gamma(s)\Gamma^*(-s^*)\mathsf{S}_{\mathsf{xx}}(s) \\
&= \frac{1}{L(s)}\frac{1}{L^*(-s^*)}\mathsf{S}_{\mathsf{xx}}(s) \\
&= \frac{1}{L(s)}\frac{1}{L^*(-s^*)}L(s)L^*(-s^*) \\
&= 1
\end{aligned}$$

✏

[6] Papoulis (1991), pages 401–402

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

CHAPTER 7

ADDITIVE NOISE ON RANDOM SEQUENCES

7.1 Additive noise and correlation

Theorem 7.1. *Let* **S** *be the* SYSTEM *illustrated to the right, where* **T** *is* NOT NECESSARILY LINEAR.



THM

$$\left.\begin{array}{lll} \text{(A).} & \mathsf{x}(n) \text{ is} & \textit{WSS} \quad \text{and} \\ \text{(B).} & \mathsf{x}(n) \text{ and } \mathsf{v}(n) \text{ are} & \textit{uncorrelated} \quad \text{and} \\ \text{(C).} & \mathsf{v}(n) \text{ is} & \textit{zero-mean} \end{array}\right\} \Longrightarrow \left\{\begin{array}{llll} \text{(1).} & \mathsf{R}_{\mathsf{yv}}(m) & = & \mathsf{R}_{\mathsf{vv}}(m) \quad \text{and} \\ \text{(2).} & \mathsf{R}_{\mathsf{xy}}(m) & = & \mathsf{R}_{\mathsf{xx}}(m) \quad \text{and} \\ \text{(3).} & \mathsf{R}_{\mathsf{yy}}(m) & = & \mathsf{R}_{\mathsf{xx}}(m) + \mathsf{R}_{\mathsf{vv}}(m) \quad \text{and} \\ \text{(4).} & \mathsf{R}_{\mathsf{xx}}(m) & = & \mathsf{R}_{\mathsf{yy}}(m) + \mathsf{R}_{\mathsf{vv}}(m) - 2\mathbf{R}_{\mathsf{e}}\mathsf{R}_{\mathsf{yv}}(m) \end{array}\right.$$

✎PROOF:

$$\begin{aligned}
\mathsf{R}_{\mathsf{yv}}(m) &\triangleq \mathsf{E}\left[\mathsf{y}(m)\mathsf{v}^*(0)\right] && \text{by (A) and definition of } \mathsf{R}_{\mathsf{yv}} && \text{(Definition 2.4 page 12)} \\
&\triangleq \mathsf{E}\left[(\mathsf{x}(m)+\mathsf{v}(m))\mathsf{v}^*(0)\right] && \text{by definition of } \mathsf{y} \\
&= \mathsf{E}\left[\mathsf{x}(m)\mathsf{v}^*(0)\right] + \mathsf{E}\left[\mathsf{v}(m)\mathsf{v}^*(0)\right] && \text{by } \textit{linearity} \text{ of } \mathsf{E} && \text{(Theorem 1.1 page 4)} \\
&= \mathsf{E}\mathsf{x}(m)\mathsf{E}\mathsf{v}^*(0) + \mathsf{E}\left[\mathsf{v}(m)\mathsf{v}^*(0)\right] && \text{by } \textit{uncorrelated} \text{ hypothesis} && \text{(B)} \\
&= \mathsf{E}\mathsf{x}(m)\cancelto{0}{\mathsf{E}\mathsf{v}^*(0)} + \mathsf{E}\left[\mathsf{v}(m)\mathsf{v}^*(0)\right] && \text{by } \textit{zero-mean} \text{ hypothesis} && \text{(C)} \\
&= \mathsf{R}_{\mathsf{vv}}(m) && \text{by definition of } \mathsf{R}_{\mathsf{vv}} && \text{(Definition 2.4 page 12)} \\
\mathsf{R}_{\mathsf{xy}}(m) &\triangleq \mathsf{E}\left[\mathsf{x}(m)\mathsf{y}^*(0)\right] && \text{by (A) and definition of } \mathsf{R}_{\mathsf{xy}} && \text{(Definition 2.4 page 12)} \\
&\triangleq \mathsf{E}\left(\mathsf{x}(m)[\mathsf{x}(0)+\mathsf{v}(0)]^*\right) && \text{by definition of } \mathsf{y} \\
&= \mathsf{E}\left[\mathsf{x}(m)\mathsf{x}^*(0)\right] + \mathsf{E}\left[\mathsf{x}(m)\mathsf{v}^*(0)\right] && \text{by } \textit{linearity} \text{ of } \mathsf{E} && \text{(Theorem 1.1 page 4)} \\
&= \mathsf{E}\left[\mathsf{x}(m)\mathsf{x}^*(0)\right] + \mathsf{E}[\mathsf{x}(m)]\mathsf{E}\left[\mathsf{v}^*(0)\right] && \text{by } \textit{uncorrelated} \text{ hypothesis} && \text{(B)} \\
&= \mathsf{E}\left[\mathsf{x}(m)\mathsf{x}^*(0)\right] + \mathsf{E}[\mathsf{x}(m)]\cancelto{0}{\mathsf{E}\left[\mathsf{v}^*(0)\right]} && \text{by } \textit{zero-mean} \text{ hypothesis} && \text{(C)} \\
&= \mathsf{R}_{\mathsf{xx}}(m) && \text{by definition of } \mathsf{R}_{\mathsf{xx}} && \text{(Definition 2.4 page 12)}
\end{aligned}$$

$$\begin{aligned}
\mathsf{R}_{\mathsf{yy}}(m) &\triangleq \mathsf{E}\left[\mathsf{y}(m)\mathsf{y}^*(0)\right] && \text{by (A) and definition of } \mathsf{R}_{\mathsf{yy}} \\
&\triangleq \mathsf{E}\left[(\mathsf{x}(m)+\mathsf{v}(m))(\mathsf{x}(0)+\mathsf{v}(0))^*\right] && \text{by definition of } \mathsf{y} \\
&= \mathsf{E}\left[\mathsf{x}(m)\mathsf{x}^*(0)\right] + \mathsf{E}\left[\mathsf{x}(m)\mathsf{v}^*(0)\right] + \mathsf{E}\left[\mathsf{v}(m)\mathsf{x}^*(0)\right] + \mathsf{E}\left[\mathsf{v}(m)\mathsf{v}^*(0)\right] \\
&= \mathsf{E}\left[\mathsf{x}(m)\mathsf{x}^*(0)\right] + \mathsf{E}\mathsf{x}(m)\mathsf{E}\mathsf{v}^*(0) + \mathsf{E}\mathsf{v}(m)\mathsf{E}\mathsf{x}^*(0) + \mathsf{E}\left[\mathsf{v}(m)\mathsf{v}^*(0)\right] && \text{by } \textit{uncorrelated} \text{ hypothesis (B)} \\
&= \mathsf{E}\left[\mathsf{x}(m)\mathsf{x}^*(0)\right] + \mathsf{E}\mathsf{x}(m)\cancelto{0}{\mathsf{E}\mathsf{v}^*(0)} + \cancelto{0}{\mathsf{E}\mathsf{v}(m)}\mathsf{E}\mathsf{x}^*(0) + \mathsf{E}\left[\mathsf{v}(m)\mathsf{v}^*(0)\right] && \text{by } \textit{zero-mean} \text{ hypothesis (C)}
\end{aligned}$$

$$
\begin{aligned}
&= \mathsf{R}_{\mathsf{xx}}(m) + \mathsf{R}_{\mathsf{vv}}(m) && \text{by definition of } \mathsf{R}_{\mathsf{xx}} \\
\mathsf{R}_{\mathsf{xx}}(m) &\triangleq \mathsf{E}\big[\mathsf{x}(m)\mathsf{x}^*(0)\big] \\
&\triangleq \mathsf{E}\Big(\big[\mathsf{y}(m) - \mathsf{v}(m)\big]\big[\mathsf{y}(0) - \mathsf{v}(0)\big]^*\Big) \\
&= \mathsf{E}\big[\mathsf{y}(m)\mathsf{y}^*(0)\big] - \mathsf{E}\big[\mathsf{y}(m)\mathsf{v}^*(0)\big] - \mathsf{E}\big[\mathsf{v}(m)\mathsf{y}^*(0)\big] + \mathsf{E}\big[\mathsf{v}(m)\mathsf{v}^*(0)\big] \\
&\triangleq \mathsf{R}_{\mathsf{yy}}(m) - \mathsf{R}_{\mathsf{yv}}(m) - \mathsf{R}_{\mathsf{vy}}(m) + \mathsf{R}_{\mathsf{vv}}(m) \\
&= \mathsf{R}_{\mathsf{yy}}(m) + \mathsf{R}_{\mathsf{vv}}(m) - 2\mathbf{R}_{\mathsf{e}}\mathsf{R}_{\mathsf{yv}}(m)
\end{aligned}
$$

✏

Remark 7.1. Because in Theorem 7.1 $\mathsf{y} = \mathsf{x} + \mathsf{v}$ and $\mathsf{R}_{\mathsf{yy}} = \mathsf{R}_{\mathsf{xx}} + \mathsf{R}_{\mathsf{vv}}$, one might assume that R is a kind of *linear operator* (Definition N.3 page 304) and further assume that because $\mathsf{x} = \mathsf{y} - \mathsf{v}$ and $\mathsf{R}_{(-\mathsf{v})(-\mathsf{v})} = \mathsf{R}_{\mathsf{vv}}$, that $\mathsf{R}_{\mathsf{xx}} = \mathsf{R}_{\mathsf{yy}} + \mathsf{R}_{\mathsf{vv}}$. As Theorem 7.1 demonstrates, this is simply **not the case**. The problem here is that y and v are very much *correlated*—in fact y is obviously a *function* of v.

Corollary 7.1. *Let* $\mathbf{S}$ *be the* SYSTEM *illustrated in Theorem 7.1 (page 51).*

COR

$$
\left.\begin{array}{l}\textit{hypotheses of}\\ \textit{Theorem 7.1 (page 51)}\end{array}\right\} \Longrightarrow \left\{\begin{array}{lrcll}
(1). & \check{\mathsf{S}}_{\mathsf{yy}}(z) & = & \check{\mathsf{S}}_{\mathsf{xx}}(z) + \check{\mathsf{S}}_{\mathsf{vv}}(z) & \textit{and} \\
(2). & \check{\mathsf{S}}_{\mathsf{yv}}(z) & = & \check{\mathsf{S}}_{\mathsf{vv}}(z) & \textit{and} \\
(3). & \check{\mathsf{S}}_{\mathsf{yv}}(z) & = & \check{\mathsf{S}}_{\mathsf{yy}}(z) + \check{\mathsf{S}}_{\mathsf{vv}}(z) + \check{\mathsf{S}}_{\mathsf{yv}}(z) + \check{\mathsf{S}}^*_{\mathsf{yv}}(z^*) & \textit{and} \\
(4). & \tilde{\mathsf{S}}_{\mathsf{yy}}(\omega) & = & \tilde{\mathsf{S}}_{\mathsf{xx}}(\omega) + \tilde{\mathsf{S}}_{\mathsf{vv}}(\omega) & \textit{and} \\
(5). & \tilde{\mathsf{S}}_{\mathsf{yv}}(\omega) & = & \tilde{\mathsf{S}}_{\mathsf{vv}}(\omega) & \textit{and} \\
(6). & \tilde{\mathsf{S}}_{\mathsf{yv}}(\omega) & = & \tilde{\mathsf{S}}_{\mathsf{yy}}(\omega) + \tilde{\mathsf{S}}_{\mathsf{vv}}(\omega) + \tilde{\mathsf{S}}_{\mathsf{yv}}(\omega) + \tilde{\mathsf{S}}^*_{\mathsf{yv}}(-\omega) &
\end{array}\right.
$$

✎PROOF:

$$
\begin{aligned}
\check{\mathsf{S}}_{\mathsf{yy}}(z) &\triangleq \mathbf{Z}\mathsf{R}_{\mathsf{yy}}(m) && \text{by definition of } \check{\mathsf{S}}_{\mathsf{xx}} && \text{(Definition 2.5 page 14)} \\
&= \mathbf{Z}\mathsf{R}_{\mathsf{qq}}(m) + \mathbf{Z}\mathsf{R}_{\mathsf{vv}}(m) && \text{by previous result} && (1) \\
&= \check{\mathsf{S}}_{\mathsf{qq}}(z) + \check{\mathsf{S}}_{\mathsf{vv}}(z) && \text{by definition of } \check{\mathsf{S}}_{\mathsf{yy}} && \text{(Definition 6.3 page 44)} \\
\tilde{\mathsf{S}}_{\mathsf{yy}}(\omega) &\triangleq \check{\mathbf{F}}\mathsf{R}_{\mathsf{yy}}(m) && \text{by definition of } \tilde{\mathsf{S}}_{\mathsf{yy}} && \text{(Definition 6.3 page 44)} \\
&= \check{\mathbf{F}}\mathsf{R}_{\mathsf{qq}}(m) + \check{\mathbf{F}}\mathsf{R}_{\mathsf{vv}}(m) && \text{by previous result} && (1) \\
&= \tilde{\mathsf{S}}_{\mathsf{qq}}(\omega) + \tilde{\mathsf{S}}_{\mathsf{vv}}(\omega) && \text{by definition of } \tilde{\mathsf{S}}_{\mathsf{yy}} && \text{(Definition 6.3 page 44)}
\end{aligned}
$$

✏

Theorem 7.2. *Let* $\mathbf{S}$ *be the* SYSTEM *illustrated to the right:*



THM

$$
\left\{\begin{array}{lrcll}
(A). & \mathsf{x}(n) \text{ and } \mathsf{r}(n) \text{ are } \textit{wide sense stationary} & & & \text{and} \\
(B). & \mathsf{E}[\mathsf{x}(n)\mathsf{w}(n)] & = & \mathsf{E}\mathsf{x}(n)\mathsf{E}\mathsf{w}(n) \quad (\textit{uncorrelated}) & \text{and} \\
(C). & \mathsf{E}[\mathsf{r}(n)\mathsf{v}(n)] & = & \mathsf{E}\mathsf{r}(n)\mathsf{E}\mathsf{v}(n) \quad (\textit{uncorrelated}) & \text{and} \\
(D). & \mathsf{E}[\mathsf{w}(n)\mathsf{v}(n)] & = & \mathsf{E}\mathsf{w}(n)\mathsf{E}\mathsf{v}(n) \quad (\textit{uncorrelated}) & \text{and} \\
(E). & \mathsf{E}\mathsf{v}(n) = \mathsf{E}\mathsf{w}(n) & = & 0 \quad (\textit{zero-mean}) &
\end{array}\right\} \Longrightarrow \left\{\begin{array}{rcl}
\mathsf{R}_{\mathsf{sy}}(m) & = & \mathsf{R}_{\mathsf{sx}}(m) \\
 & = & \mathsf{R}_{\mathsf{ry}}(m) \\
 & = & \mathsf{R}_{\mathsf{rx}}(m)
\end{array}\right\}
$$

COPYRIGHT © 2019 DANIEL J. GREENHOE

✎PROOF:

$$
\begin{aligned}
R_{sy}(m) &\triangleq E\big[s(m)y^*(0)\big] && \text{by definition of } R_{xy} && \text{(Definition 2.4 page 12)}\\
&\triangleq E\big([r(m)+w(m)][x(0)+v(0)]^*\big) && \text{by definition of } \mathbf{S} && \\
&= E\big[r(m)x^*(0)\big] + E\big[r(m)v^*(0)\big] + E\big[w(m)x^*(0)\big] + E\big[w(m)v^*(0)\big] && && \\
&= E\big[r(m)x^*(0)\big] + Er(m)Ev^*(0) && && \\
&\qquad + Ew(m)Ex^*(0) + Ew(m)Ev^*(0) && \text{by } \textit{uncorrelated} \text{ hypotheses} && \text{(B), (C), and (D)}\\
&= E\big[r(m)x^*(0)\big] + Er(m)\cancelto{0}{Ev^*(0)} && && \\
&\qquad + \cancelto{0}{Ew(m)}Ex^*(0) + \cancelto{0}{Ew(m)}Ev^*(0) && \text{by } \textit{zero-mean} \text{ hypothesis} && \text{(E)}\\
&= R_{rx}(m) && \text{by definition of } R_{rx} && \text{(Definition 2.4 page 12)}\\
R_{sx}(m) &\triangleq E\big[s(m)x^*(0)\big] && \text{by definition of } R_{xy} && \text{(Definition 2.4 page 12)}\\
&\triangleq E\big([r(m)+w(m)]x^*(0)\big) && \text{by definition of } \mathbf{S} && \\
&= E\big[r(m)x^*(0)\big] + Ew(m)Ex^*(0) && \text{by } \textit{uncorrelated} \text{ hypothesis} && \text{(B)}\\
&= E\big[r(m)x^*(0)\big] + \cancelto{0}{Ew(m)}Ex^*(0) && \text{by } \textit{zero-mean} \text{ hypothesis} && \text{(E)}\\
&= R_{rx}(m) && \text{by definition of } R_{rx} && \text{(Definition 2.4 page 12)}\\
R_{ry}(m) &\triangleq E\big[r(m)y^*(0)\big] && \text{by definition of } R_{xy} && \text{(Definition 2.4 page 12)}\\
&\triangleq E\big(r(m)[x(0)+v(0)]^*\big) && \text{by definition of } \mathbf{S} && \\
&= E\big[r(m)x^*(0)\big] + Er(m)Ev^*(0) && \text{by } \textit{uncorrelated} \text{ hypothesis} && \text{(C)}\\
&= E\big[r(m)x^*(0)\big] + Er(m)\cancelto{0}{Ev^*(0)} && \text{by } \textit{zero-mean} \text{ hypothesis} && \text{(E)}\\
&= R_{rx}(m) && \text{by definition of } R_{rx} && \text{(Definition 2.4 page 12)}
\end{aligned}
$$

✎

Corollary 7.2. *Let* $\mathbf{S}$ *be the* SYSTEM *illustrated in Theorem 7.2 (page 52).*

COR
$$\left\{\begin{array}{l}\textit{hypotheses of}\\ \textit{Theorem 7.2 (page 52)}\end{array}\right\} \Longrightarrow \left\{\begin{array}{lllllllll} (1). & \check{S}_{sy}(z) & = & \check{S}_{sx}(z) & = & \check{S}_{ry}(z) & = & \check{S}_{rx}(z) & \textit{and}\\ (2). & \tilde{S}_{sy}(\omega) & = & \tilde{S}_{sx}(\omega) & = & \tilde{S}_{ry}(\omega) & = & \tilde{S}_{rx}(\omega) & \end{array}\right\}$$

✎PROOF:

$$
\begin{aligned}
\check{S}_{sy}(\omega) &\triangleq \mathbf{Z}R_{sy}(m) && \text{by definition of } \check{S}_{xy} && \text{(Definition 2.5 page 14)}\\
&= \mathbf{Z}R_{rx}(m) && \text{by previous result} && (1)\\
&= \check{S}_{rx}(z) && \text{by definition of } \check{S}_{xy} && \text{(Definition 2.5 page 14)}\\
\tilde{S}_{sy}(\omega) &\triangleq \check{\mathbf{F}}R_{sy}(m) && \text{by definition of } \tilde{S}_{xy} && \text{(Definition 6.3 page 44)}\\
&= \check{\mathbf{F}}R_{rx}(m) && \text{by previous result} && (1)\\
&= \tilde{S}_{rx}(\omega) && \text{by definition of } \tilde{S}_{xy} && \text{(Definition 6.3 page 44)}
\end{aligned}
$$

✎

7.2 Additive noise and operators

Lemma 7.1. *Let* $\mathbf{S}$ *be the* SYSTEM *illustrated in Figure 7.2 (page 55) (A).*

LEM
$$\left\{\begin{array}{llllll} (A). & R_{xx}(n_1,m) & = & R_{xx}(n_2,m) & \text{(WSS)} & \textit{and}\\ (B). & E[x(n)v(n)] & = & Ex(n)Ev(n) & \text{(UNCORRELATED)} & \textit{and}\\ (E). & Ev(n) & = & 0 & \text{(ZERO-MEAN)} & \end{array}\right\} \Longrightarrow \left\{\begin{array}{llll} (1). & R_{xy}(m) & = & R_{xq}(m) & \textit{and}\\ (2). & \check{S}_{xy}(z) & = & \check{S}_{xq}(z) & \textit{and}\\ (3). & \tilde{S}_{xy}(\omega) & = & \tilde{S}_{xq}(\omega) & \end{array}\right\}$$

COPYRIGHT © 2019 DANIEL J. GREENHOE | CC BY-NC-ND



(A) $\mathsf{R}_{\mathsf{xy}}(m) = \mathsf{R}_{\mathsf{xq}}(m)$ (Lemma 7.1 page 53)

(B) $\mathsf{R}_{\mathsf{py}}(m) = \mathsf{R}_{\mathsf{xy}}(m) + \mathsf{R}_{\mathsf{uy}}(m)$ (Lemma 7.2 page 54)

(C) $\mathsf{R}_{\mathsf{xy}}(m) = \mathsf{R}_{\mathsf{py}}(m)$ (Lemma 7.3 page 54)

Figure 7.1: Additive noise with *linear*/**non-linear** operator **T**

✎PROOF:

$\mathsf{R}_{\mathsf{xy}}(m) \triangleq \mathsf{E}\big[\mathsf{x}(m)\mathsf{y}^*(0)\big]$	by definition of R_{xy}	(Definition 2.4 page 12)
$\triangleq \mathsf{E}\big[\mathsf{x}(m)(\mathsf{q}(0)+\mathsf{v}(0))^*\big]$	by definition of **S**	(Figure 7.2 page 55) (A)
$= \mathsf{E}\big[\mathsf{x}(m)\mathsf{q}^*(0)+\mathsf{p}(m)\mathsf{v}^*(0)\big]$		
$= \mathsf{E}\big[\mathsf{x}(m)\mathsf{q}^*(0)\big]+\mathsf{E}\big[\mathsf{x}(m)\mathsf{v}^*(0)\big]$		
$= \mathsf{E}\big[\mathsf{x}(m)\mathsf{q}^*(0)\big]+[\mathsf{E}\mathsf{x}(m)]\big[\mathsf{E}\mathsf{v}^*(0)\big]$	by *uncorrelated* hypothesis	(B)
$= \mathsf{E}\big[\mathsf{x}(m)\mathsf{q}^*(0)\big]+\big[\mathsf{E}\mathsf{p}(m)\big]\cancelto{0}{\big[\mathsf{E}\mathsf{v}^*(0)\big]}$	by *zero-mean* hypothesis	(E)
$= \mathsf{R}_{\mathsf{xq}}(m)$	by definition of R_{xq}	(Definition 2.4 page 12)
$\check{\mathsf{S}}_{\mathsf{xy}}(z) \triangleq \mathbf{Z}\mathsf{R}_{\mathsf{xy}}(m)$	by definition of $\check{\mathsf{S}}_{\mathsf{xy}}$	(Definition 2.5 page 14)
$= \mathbf{Z}\mathsf{R}_{\mathsf{xq}}(m)$	by previous result	(1)
$= \check{\mathsf{S}}_{\mathsf{xq}}(z)$	by definition of $\check{\mathsf{S}}_{\mathsf{xq}}$	(Definition 2.5 page 14)
$\tilde{\mathsf{S}}_{\mathsf{xy}}(\omega) \triangleq \check{\mathbf{F}}\mathsf{R}_{\mathsf{xy}}(m)$	by definition of $\tilde{\mathsf{S}}_{\mathsf{xy}}$	(Definition 6.3 page 44)
$= \check{\mathbf{F}}\mathsf{R}_{\mathsf{xq}}(m)$	by previous result	(1)
$= \tilde{\mathsf{S}}_{\mathsf{xq}}(\omega)$	by definition of $\tilde{\mathsf{S}}_{\mathsf{xq}}$	(Definition 6.3 page 44)

✏

Lemma 7.2. *Let* **S** *be the* SYSTEM *illustrated in Figure 7.2 (page 55) (B).*

LEM
$$\left\{\begin{array}{llll} (A). & \mathsf{x}(n) \text{ is} & (\text{WSS}) & \text{and} \\ (B). & \mathsf{u}(n) \text{ is} & (\text{ZERO-MEAN}) & \text{and} \\ (C). & \mathsf{x}(n) \text{ and } \mathsf{u}(n) \text{ are} & (\text{UNCORRELATED}) & \end{array}\right\} \Longrightarrow \left\{\begin{array}{llll} (1). & \mathsf{R}_{\mathsf{pq}}(m) = \mathsf{R}_{\mathsf{xy}}(m)+\mathsf{R}_{\mathsf{uy}}(m) & & \text{and} \\ (2). & \check{\mathsf{S}}_{\mathsf{pq}}(z) = \check{\mathsf{S}}_{\mathsf{xy}}(z)+\check{\mathsf{S}}_{\mathsf{uy}}(z) & & \text{and} \\ (3). & \tilde{\mathsf{S}}_{\mathsf{pq}}(\omega) = \tilde{\mathsf{S}}_{\mathsf{xy}}(\omega)+\tilde{\mathsf{S}}_{\mathsf{uy}}(\omega) & & \end{array}\right\}$$

✎PROOF:

$\mathsf{R}_{\mathsf{xy}}(m) \triangleq \mathsf{E}\big[\mathsf{x}(m)\mathsf{y}^*(0)\big]$	by definition of R_{xy}	(Definition 2.4 page 12)
$\triangleq \mathsf{E}\big(\big[\mathsf{p}(m)-\mathsf{u}(m)\big]\mathsf{y}^*(0)\big)$	by definition of **S**	
$= \mathsf{E}\big[\mathsf{p}(m)\mathsf{y}^*(0)-\mathsf{u}(m)\mathsf{y}^*(0)\big]$		
$= \mathsf{E}\big[\mathsf{p}(m)\mathsf{y}^*(0)\big]-\mathsf{E}\big[\mathsf{u}(m)\mathsf{y}^*(0)\big]$	because E is a *linear operator*	(Theorem 1.1 page 4)
$\triangleq \mathsf{R}_{\mathsf{py}}(m)-\mathsf{R}_{\mathsf{uy}}(m)$	by definition of R_{xy}	(Definition 2.4 page 12)

✏

Lemma 7.3. *Let* **S** *be the* SYSTEM *illustrated in Figure 7.2 (page 55) (C).*

LEM
$$\left\{\begin{array}{llll} (A). & \mathsf{x}(n) \text{ is} & (\text{WSS}) & \text{and} \\ (B). & \mathsf{u}(n) \text{ is} & (\text{ZERO-MEAN}) & \text{and} \\ (C). & \mathsf{x}(n) \text{ and } \mathsf{u}(n) \text{ are} & (\text{UNCORRELATED}) & \end{array}\right\} \Longrightarrow \left\{\begin{array}{llll} (1). & \mathsf{R}_{\mathsf{xy}}(m) = \mathsf{R}_{\mathsf{py}}(m) & & \text{and} \\ (2). & \check{\mathsf{S}}_{\mathsf{xy}}(z) = \check{\mathsf{S}}_{\mathsf{py}}(z) & & \text{and} \\ (3). & \tilde{\mathsf{S}}_{\mathsf{xy}}(\omega) = \tilde{\mathsf{S}}_{\mathsf{py}}(\omega) & & \end{array}\right\}$$

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

✎PROOF:

$$
\begin{aligned}
\mathsf{R}_{\mathsf{xy}}(m) &\triangleq \mathsf{E}\big[\mathsf{x}(m)\mathsf{y}^*(0)\big] && \text{by definition of } \mathsf{R}_{\mathsf{py}} && \text{(Definition 2.4 page 12)}\\
&\triangleq \mathsf{E}\big(\big[\mathsf{p}(m)+\mathsf{u}(m)\big]\mathsf{y}^*(0)\big) && \text{by definition of } \mathbf{S} && \\
&= \mathsf{E}\big[\mathsf{p}(m)\mathsf{y}^*(0)+\mathsf{u}(m)\mathsf{y}^*(0)\big] && \text{by field properties of } (\mathbb{R},+,\cdot,0,1) && \\
&= \mathsf{E}\big[\mathsf{p}(m)\mathsf{y}^*(0)\big]+\mathsf{E}\big[\mathsf{u}(m)\mathsf{y}^*(0)\big] && \text{because } \mathsf{E} \text{ is a } \textit{linear operator} && \text{(Theorem 1.1 page 4)}\\
&= \mathsf{E}\big[\mathsf{p}(m)\mathsf{y}^*(0)\big]+\mathsf{E}[\mathsf{u}(m)]\mathsf{E}\big[\mathsf{y}^*(0)\big] && \text{by } \textit{uncorrelated} \text{ hypothesis} && \text{(C)}\\
&= \mathsf{E}\big[\mathsf{p}(m)\mathsf{y}^*(0)\big]+\mathsf{E}[\mathsf{u}(m)]\cancelto{0}{\mathsf{E}\big[\mathsf{y}^*(0)\big]} && \text{by } \textit{zero-mean} \text{ hypothesis} && \text{(B)}\\
&\triangleq \mathsf{R}_{\mathsf{py}}(m) && \text{by definition of } \mathsf{R}_{\mathsf{xy}} && \text{(Definition 2.4 page 12)}
\end{aligned}
$$

✏



(A) *Communications additive noise system*
$\tilde{\mathsf{S}}_{\mathsf{pq}}(\omega)=\tilde{\mathsf{S}}_{\mathsf{py}}(\omega)=\tilde{\mathsf{S}}_{\mathsf{xy}}(\omega)+\tilde{\mathsf{S}}_{\mathsf{uy}}(\omega)$
(Theorem 7.3 page 55)

(B) *Measurement additive noise system*
$\tilde{\mathsf{S}}_{\mathsf{pq}}(\omega)=\tilde{\mathsf{S}}_{\mathsf{py}}(\omega)=\tilde{\mathsf{S}}_{\mathsf{xq}}(\omega)=\tilde{\mathsf{S}}_{\mathsf{xy}}(\omega)$
(Theorem 7.4 page 55)

Figure 7.2: *linear* / **non-linear** additive noise systems

Theorem 7.3 (*communications additive noise cross-correlation*).
Let **S** *be the* SYSTEM *illustrated in Figure 7.2 page 55 (A).*

THM

$$
\left.\begin{array}{llll}
(A). & \mathsf{x}(n) \textit{ is} & \text{WSS} & \textit{and}\\
(B). & \mathsf{u}(n) \textit{ is} & \text{ZERO-MEAN} & \textit{and}\\
(C). & \mathsf{v}(n) \textit{ is} & \text{ZERO-MEAN} & \textit{and}\\
(D). & \mathsf{x}(n),\mathsf{u}(n),\mathsf{v}(n) \textit{ are} & \text{UNCORRELATED} &
\end{array}\right\}
\Longrightarrow
\left\{\begin{array}{lll}
(1). & \mathsf{R}_{\mathsf{pq}}(m)=\mathsf{R}_{\mathsf{py}}(m)=\mathsf{R}_{\mathsf{xy}}(m)+\mathsf{R}_{\mathsf{uy}}(m) & \textit{and}\\
(2). & \check{\mathsf{S}}_{\mathsf{pq}}(z)=\check{\mathsf{S}}_{\mathsf{py}}(z)=\check{\mathsf{S}}_{\mathsf{xy}}(z)+\check{\mathsf{S}}_{\mathsf{uy}}(z) & \textit{and}\\
(3). & \tilde{\mathsf{S}}_{\mathsf{pq}}(\omega)=\tilde{\mathsf{S}}_{\mathsf{py}}(\omega)=\tilde{\mathsf{S}}_{\mathsf{xy}}(\omega)+\tilde{\mathsf{S}}_{\mathsf{uy}}(\omega) &
\end{array}\right.
$$

✎PROOF:

$$
\begin{aligned}
\mathsf{R}_{\mathsf{pq}}(m) &= \mathsf{R}_{\mathsf{py}}(m) && \text{by Lemma 7.1 page 53} && \\
\mathsf{R}_{\mathsf{pq}}(m) &= \mathsf{R}_{\mathsf{xq}}(m)+\mathsf{R}_{\mathsf{uq}}(m) && \text{by Lemma 7.2 page 54} && \\
\mathsf{R}_{\mathsf{py}}(m) &\triangleq \mathsf{E}\big[\mathsf{p}(m)\mathsf{y}^*(0)\big] && \text{by definition } \mathsf{R}_{\mathsf{py}} && \text{(Definition 2.4 page 12)}\\
&\triangleq \mathsf{E}\big([\mathsf{x}(m)+\mathsf{u}(m)]\mathsf{y}^*(0)\big) && \text{by definition } \mathbf{S} && \text{(Figure 7.2 page 55) (A)}\\
&= \mathsf{E}\big[\mathsf{x}(m)\mathsf{y}^*(0)+\mathsf{u}(m)\mathsf{y}^*(0)\big] && && \\
&= \mathsf{E}\big[\mathsf{x}(m)\mathsf{y}^*(0)\big]+\mathsf{E}\big[\mathsf{u}(m)\mathsf{y}^*(0)\big] && \text{by } \textit{linearity} \text{ of } \mathsf{E} && \text{(Theorem 1.1 page 4)}\\
&= \mathsf{R}_{\mathsf{xy}}(m)+\mathsf{R}_{\mathsf{uy}}(m) && \text{by definitions } \mathsf{R}_{\mathsf{xy}} \text{ and } \mathsf{R}_{\mathsf{uy}} && \text{(Definition 2.4 page 12)}
\end{aligned}
$$

✏

Theorem 7.4 (*measurement additive noise cross-correlation*).
Let **S** *be the* SYSTEM *illustrated in Figure 7.2 page 55 (B).*

THM

$$
\left.\begin{array}{llll}
(A). & \mathsf{x}(n) \textit{ is} & \text{WSS} & \textit{and}\\
(B). & \mathsf{u}(n) \textit{ is} & \text{ZERO-MEAN} & \textit{and}\\
(C). & \mathsf{v}(n) \textit{ is} & \text{ZERO-MEAN} & \textit{and}\\
(D). & \mathsf{x}(n),\mathsf{u}(n),\mathsf{v}(n) \textit{ are} & \text{UNCORRELATED} &
\end{array}\right\}
\Longrightarrow
\left\{\begin{array}{lll}
(1). & \mathsf{R}_{\mathsf{pq}}(m)=\mathsf{R}_{\mathsf{py}}(m)=\mathsf{R}_{\mathsf{xq}}(m)=\mathsf{R}_{\mathsf{xy}}(m) & \textit{and}\\
(2). & \check{\mathsf{S}}_{\mathsf{pq}}(z)=\check{\mathsf{S}}_{\mathsf{py}}(z)=\check{\mathsf{S}}_{\mathsf{xq}}(z)=\check{\mathsf{S}}_{\mathsf{xy}}(z) & \textit{and}\\
(3). & \tilde{\mathsf{S}}_{\mathsf{pq}}(\omega)=\tilde{\mathsf{S}}_{\mathsf{py}}(\omega)=\tilde{\mathsf{S}}_{\mathsf{xq}}(\omega)=\tilde{\mathsf{S}}_{\mathsf{xy}}(\omega) &
\end{array}\right.
$$

COPYRIGHT © 2019 DANIEL J. GREENHOE | CC BY-NC-ND

✎PROOF:

$$
\begin{aligned}
\mathsf{R}_{\mathsf{pq}}(m) &= \mathsf{R}_{\mathsf{py}}(m) && \text{by Lemma 7.1 page 53} \\
\mathsf{R}_{\mathsf{pq}}(m) &= \mathsf{R}_{\mathsf{xq}}(m) && \text{by Lemma 7.3 page 54} \\
\mathsf{R}_{\mathsf{xy}}(m) &\triangleq \mathsf{E}\big[\mathsf{x}(m)\mathsf{y}^*(0)\big] && \text{by definition } \mathsf{R}_{\mathsf{xy}} \qquad \text{(Definition 2.4 page 12)} \\
&\triangleq \mathsf{E}\big(\big[\mathsf{p}(m)+\mathsf{u}(m)\big]\mathsf{y}^*(0)\big) && \text{by definition } \mathbf{S} \qquad \text{(Figure 7.2 page 55) (B)} \\
&= \mathsf{E}\big[\mathsf{p}(m)\mathsf{y}^*(0)+\mathsf{u}(m)\mathsf{y}^*(0)\big] \\
&= \mathsf{E}\big[\mathsf{p}(m)\mathsf{y}^*(0)\big]+\mathsf{E}\big[\mathsf{u}(m)\mathsf{y}^*(0)\big] && \text{by } \textit{linearity} \text{ of } \mathsf{E} \qquad \text{(Theorem 1.1 page 4)} \\
&= \mathsf{E}\big[\mathsf{p}(m)\mathsf{y}^*(0)\big]+\cancelto{0}{\mathsf{E}\big[\mathsf{u}(m)\mathsf{y}^*(0)\big]} && \text{by } \textit{uncorrelated} \text{ hypothesis} \qquad \text{(D)} \\
&= \mathsf{R}_{\mathsf{py}}(m) && \text{by definition of } \mathsf{R}_{\mathsf{py}} \qquad \text{(Definition 2.4 page 12)}
\end{aligned}
$$

✎

7.3 Additive noise and LTI operators



Figure 7.3: Additive noise with **LTI** operator **H**

Corollary 7.3. *Let* **S** *be the* SYSTEM *illustrated in Figure 7.3 (page 56) (A).*

COR

$$
\left\{
\begin{array}{llll}
(A). & \mathsf{x}(n) \textit{ is} & (\text{WSS}) & \textit{and} \\
(B). & \mathsf{u}(n) \textit{ is} & (\text{ZERO-MEAN}) & \textit{and} \\
(C). & \mathsf{x}(n) \textit{ and } \mathsf{u}(n) \textit{ are} & (\text{UNCORRELATED}) & \textit{and} \\
(D). & \mathbf{H} \textit{ is} & (\text{LTI}) &
\end{array}
\right\}
\Longrightarrow
\left\{ \tilde{\mathsf{S}}_{\mathsf{xy}}(\omega) = \tilde{\mathsf{H}}^*(\omega)\tilde{\mathsf{S}}_{\mathsf{xx}}(\omega) \right\}
$$

✎PROOF:

$$
\begin{aligned}
\tilde{\mathsf{S}}_{\mathsf{xy}}(\omega) &= \tilde{\mathsf{S}}_{\mathsf{xq}}(\omega) && \text{by Lemma 7.1 page 53} \\
&= \tilde{\mathsf{H}}^*(\omega)\tilde{\mathsf{S}}_{\mathsf{xx}}(\omega) && \text{by Corollary 5.3 page 37}
\end{aligned}
$$

✎

Theorem 7.5. *Let* **S** *be the* SYSTEM *illustrated in Figure 7.3 (page 56) (B).*

THM

$$
\left\{
\begin{array}{llll}
(A). & \mathsf{x}(n) \textit{ is} & (\text{WSS}) & \textit{and} \\
(B). & \mathsf{u}(n) \textit{ is} & (\text{ZERO-MEAN}) & \textit{and} \\
(C). & \mathsf{x}(n) \textit{ and } \mathsf{u}(n) \textit{ are} & (\text{UNCORRELATED}) & \textit{and} \\
(D). & \mathbf{H} \textit{ is} & (\text{LTI}) &
\end{array}
\right\}
\Longrightarrow
\left\{
\begin{array}{llll}
(1). & \mathsf{R}_{\mathsf{yx}}(m) = \mathsf{h}(m) \star \mathsf{R}_{\mathsf{xx}}(m) & \textit{and} \\
(2). & \check{\mathsf{S}}_{\mathsf{yx}}(z) = \check{\mathsf{H}}(z)\check{\mathsf{S}}_{\mathsf{xx}}(z) & \textit{and} \\
(3). & \tilde{\mathsf{S}}_{\mathsf{xy}}(\omega) = \tilde{\mathsf{H}}^*(\omega)\tilde{\mathsf{S}}_{\mathsf{xx}}(\omega) &
\end{array}
\right\}
$$

✎PROOF:

1. definition: Let $(\mathsf{h}(n))$ be the *impulse response* of operator **H** such that
$$\mathbf{H}\bar{\delta}(n) \triangleq \sum_{m\in\mathbb{Z}} \mathsf{h}(m)\bar{\delta}(n-m)$$

COPYRIGHT © 2019 DANIEL J. GREENHOE

2. lemma: $\mathbf{H}\boldsymbol{x}(n) = \sum_{m\in\mathbb{Z}} \mathsf{h}(n)\boldsymbol{x}(m-n) = \mathsf{h}(n) \star \mathsf{R}_{\mathsf{xx}}(n)$.

   Proof: by the *linear time-invariant* hypotheses (D) and definition of *convolution* operator $\star$ (Definition O.3 page 336)

3. Proof that $\mathsf{R}_{\mathsf{yx}}(m) = \mathsf{h}(n) \star \mathsf{R}_{\mathsf{xx}}(m)$:

$$\begin{aligned}
\mathsf{R}_{\mathsf{yx}}(m) &\triangleq \mathsf{E}\big[\mathsf{y}(m)\mathsf{x}^*(0)\big] && \text{by definition of } \mathsf{R}_{\mathsf{py}} && \text{(Definition 2.4 page 12)}\\
&= \mathsf{E}\big([\mathbf{H}\mathsf{x}(m) + \mathbf{H}\mathsf{u}(m)]\mathsf{x}^*(0)\big) && \text{by } \textit{linear} \text{ hypothesis} && \text{(D)}\\
&= \mathsf{E}\big(\big[\mathbf{H}\mathsf{x}^*(m)\big]\mathsf{x}^*(0) + [\mathbf{H}\mathsf{u}(0)]\mathsf{x}^*(0)\big) && && \\
&= \mathsf{E}\big(\big[\mathbf{H}\mathsf{x}^*(m)\big]\mathsf{x}^*(0)\big) + \mathsf{E}\big([\mathbf{H}\mathsf{u}(0)]\mathsf{x}^*(0)\big) && \text{by } \textit{linearity} \text{ of } \mathbf{E} && \text{(Theorem ?? page ??)}\\
&= \mathbf{H}\mathsf{E}\big[\mathsf{x}(m)\mathsf{x}^*(0)\big] + \mathbf{H}\mathsf{E}\big[\mathsf{u}(m)\mathsf{x}^*(0)\big] && \text{by } \textit{LTI} \text{ hypotheses} && \text{(D)}\\
&= \mathbf{H}\mathsf{E}\big[\mathsf{x}(m)\mathsf{x}^*(0)\big] + \mathbf{H}\mathsf{E}\mathsf{u}(m)\mathsf{E}\mathsf{x}^*(0) && \text{by } \textit{uncorrelated} \text{ hypothesis} && \text{(C)}\\
&= \mathbf{H}\mathsf{E}\big[\mathsf{x}(m)\mathsf{x}^*(0)\big] + \mathbf{H}\cancelto{0}{\mathsf{E}\mathsf{u}(m)}\mathsf{E}\mathsf{x}^*(0) && \text{by } \textit{zero-mean} \text{ hypothesis} && \text{(B)}\\
&= \mathbf{H}\mathsf{R}_{\mathsf{xx}}(m) && \text{by definition of } \mathsf{R}_{\mathsf{xx}} && \text{(Definition 2.4 page 12)}\\
&= \mathsf{h}(m) \star \mathsf{R}_{\mathsf{xx}}(m) && \text{by (2) lemma} &&
\end{aligned}$$

✏

When $\mathbf{H}$ is *LTI*, what effect does the additive uncorrelated noise sources have on the cross-statistical properties of x and y? Corollary 7.5 (next) demonstrates that, amazingly, under very general conditions, the noise sources have **no effect**.

Corollary 7.4. *Let* $\mathbf{S}$ *be the* SYSTEM *illustrated in Figure 7.3 (page 56) (C).*

COR

$$\left\{\begin{array}{lll}
(A). & \mathsf{x}(n) \textit{ is} & (\text{WSS}) \quad \textit{and}\\
(B). & \mathsf{u}(n) \textit{ is} & (\text{ZERO-MEAN}) \quad \textit{and}\\
(C). & \mathsf{x}(n) \textit{ and } \mathsf{w}(n) \textit{ are} & (\text{UNCORRELATED}) \quad \textit{and}\\
(D). & \mathbf{H} \textit{ is} & (\text{LTI})
\end{array}\right\} \Longrightarrow \left\{ \tilde{\mathsf{S}}_{\mathsf{xy}}(\omega) = \tilde{\mathsf{H}}^*(\omega)\tilde{\mathsf{S}}_{\mathsf{pp}}(\omega) \right\}$$

✎PROOF:

$$\begin{aligned}
\tilde{\mathsf{S}}_{\mathsf{xy}}(\omega) &= \tilde{\mathsf{S}}_{\mathsf{py}}(\omega) && \text{by Lemma 7.3 page 54}\\
&= \tilde{\mathsf{H}}^*(\omega)\tilde{\mathsf{S}}_{\mathsf{pp}}(\omega) && \text{by Corollary 5.3 page 37}
\end{aligned}$$

✏



(A) $\tilde{\mathsf{S}}_{\mathsf{xy}}(\omega) = \tilde{\mathsf{H}}^*(\omega)\tilde{\mathsf{S}}_{\mathsf{xx}}(\omega)$ (Corollary 7.5 page 57)

(B) $\tilde{\mathsf{S}}_{\mathsf{xy}}(\omega) = \tilde{\mathsf{H}}^*(\omega)\tilde{\mathsf{S}}_{\mathsf{pp}}(\omega)$ (Corollary 7.6 page 58)

Figure 7.4:

Corollary 7.5. *Let* $\mathbf{S}$ *be the* SYSTEM *illustrated in Figure 7.4 page 57 (A).*

COR

$$\left\{\begin{array}{lll}
(A). & \mathsf{x}(n) \textit{ is} & (\text{WSS}) \quad \textit{and}\\
(B). & \mathsf{u}(n) \textit{ is} & (\text{ZERO-MEAN}) \quad \textit{and}\\
(C). & \mathsf{v}(n) \textit{ is} & (\text{ZERO-MEAN}) \quad \textit{and}\\
(D). & \mathsf{x}(n), \mathsf{u}(n), \mathsf{v}(n) \textit{ are} & (\text{UNCORRELATED}) \quad \textit{and}\\
(E). & \mathbf{H} \textit{ is} & (\text{LTI})
\end{array}\right\} \Longrightarrow \left\{ \tilde{\mathsf{S}}_{\mathsf{xy}}(\omega) = \tilde{\mathsf{H}}^*(\omega)\tilde{\mathsf{S}}_{\mathsf{xx}}(\omega) \right\}$$

COPYRIGHT © 2019 DANIEL J. GREENHOE

✎PROOF:

$$\begin{aligned}\tilde{S}_{yx}(\omega) &= \tilde{S}_{qx}(\omega) && \text{by Lemma 7.1 page 53}\\ &= \tilde{H}(\omega)\tilde{S}_{xx}(\omega) && \text{by Corollary 5.3 page 37}\end{aligned}$$

✏

Corollary 7.6. *Let* **S** *be the* SYSTEM *illustrated in Figure 7.4 page 57 (B).*

COR

$$\left\{\begin{array}{llll} (A). & \mathsf{x}(n) \text{ is} & \text{WSS} & \text{and}\\ (B). & \mathsf{w}(n) \text{ is} & \text{ZERO-MEAN} & \text{and}\\ (C). & \mathsf{v}(n) \text{ is} & \text{ZERO-MEAN} & \text{and}\\ (D). & \mathsf{x}(n), \mathsf{w}(n), \mathsf{v}(n) \text{ are} & \text{UNCORRELATED} & \text{and}\\ (E). & \mathbf{H} \text{ is} & \text{LTI} & \end{array}\right\} \Longrightarrow \left\{ \tilde{S}_{xy}(\omega) = \tilde{H}^*(\omega)\tilde{S}_{pp}(\omega) \right\}$$

✎PROOF:

$$\begin{aligned}\tilde{S}_{yx}(\omega) &= \tilde{S}_{qx}(\omega) && \text{by Lemma 7.1 page 53}\\ &= \tilde{S}_{qp}(\omega) && \text{by Lemma 7.1 page 53}\\ &= \tilde{H}(\omega)\tilde{S}_{pp}(\omega) && \text{by Corollary 5.3 page 37}\end{aligned}$$

✏

7.4 Additive noise and dual operators



(A) dual communications additive noise system (Corollary 7.7 page 58)

(B) dual measurement additive noise system (Corollary 7.8 page 59)

Figure 7.5: Dual Additive Noise Systems

Corollary 7.7. *Let* **S** *be the* SYSTEM *illustrated in Figure 7.5 (page 58) (A).*

COR

$$\left\{\begin{array}{llll} (A). & \mathbf{H} \text{ is} & \text{LTI} & \text{and}\\ (B). & \mathsf{x}(n) \text{ is} & \text{WSS} & \text{and}\\ (C). & \mathsf{u} \text{ and } \mathsf{v} \text{ are} & \text{ZERO-MEAN} & \text{and}\\ (D). & \mathsf{x}, \mathsf{u}, \mathsf{v} \text{ are} & \text{UNCORRELATED} & \end{array}\right\} \Longrightarrow \left\{\begin{array}{llll} (1). & \check{S}_{sy}(z) = \check{H}(z)\left[\check{S}_{xy}(z) + \check{S}_{uy}(z)\right] & \text{and}\\ (2). & \tilde{S}_{sy}(\omega) = \tilde{H}(\omega)\left[\tilde{S}_{xy}(\omega) + \tilde{S}_{uy}(\omega)\right] & \end{array}\right\}$$

✎PROOF:

$$\begin{aligned}\check{S}_{sy}(z) &= \check{S}_{rq}(z) && \text{by Corollary 7.2 page 53} && \text{and (B), (C) and (D)}\\ &= \check{H}(z)\check{S}_{pq}(z) && \text{by Theorem 5.2 page 37} && \text{and (A)}\\ &= \check{H}(z)\left[\check{S}_{xq}(z) + \check{S}_{uq}(z)\right] && \text{by Lemma 7.2 page 54}\\ &= \check{H}(z)\left[\check{S}_{xy}(z) + \check{S}_{uy}(z)\right] && \text{by Lemma 7.1 page 53}\\ \tilde{S}_{sy}(\omega) &= \check{S}_{sy}(z)\big|_{z=e^{i\omega}}\\ &= \check{H}(z)\left[\check{S}_{xy}(z) + \check{S}_{uy}(z)\right]\big|_{z=e^{i\omega}} && \text{by previous result} && (1)\\ &= \tilde{H}(\omega)\left[\tilde{S}_{xy}(\omega) + \tilde{S}_{uy}(\omega)\right]\end{aligned}$$

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

Corollary 7.8. *Let* **S** *be the* SYSTEM *illustrated in Figure 7.5 (page 58) (B).*

COR

$$\left\{\begin{array}{llll} (A). & \mathbf{H}\ \textit{is} & \text{LTI} & \textit{and} \\ (B). & \mathsf{x}(n)\ \textit{is} & \text{WSS} & \textit{and} \\ (C). & \mathsf{u}\ \textit{and}\ \mathsf{v}\ \textit{are} & \text{ZERO-MEAN} & \textit{and} \\ (D). & \mathsf{p}, \mathsf{u}, \mathsf{v}\ \textit{are} & \text{UNCORRELATED} & \end{array}\right\} \Longrightarrow \left\{\begin{array}{llll} (1). & \check{\mathsf{S}}_{\mathsf{sy}}(z) & = & \check{\mathsf{H}}(z)\check{\mathsf{S}}_{\mathsf{xy}}(z) \quad \textit{and} \\ (2). & \tilde{\mathsf{S}}_{\mathsf{sy}}(\omega) & = & \tilde{\mathsf{H}}(\omega)\tilde{\mathsf{S}}_{\mathsf{xy}}(\omega) \end{array}\right\}$$

✎PROOF:

$$\begin{aligned}
\check{\mathsf{S}}_{\mathsf{sy}}(z) &= \check{\mathsf{S}}_{\mathsf{rq}}(z) && \text{by Corollary 7.2 page 53} && \text{and (B), (C) and (D)} \\
&= \check{\mathsf{H}}(z)\check{\mathsf{S}}_{\mathsf{pq}}(z) && \text{by Theorem 5.2 page 37} && \text{and (A)} \\
&= \check{\mathsf{H}}(z)\check{\mathsf{S}}_{\mathsf{xq}}(z) && \text{by Lemma 7.3 page 54} \\
&= \check{\mathsf{H}}(z)\check{\mathsf{S}}_{\mathsf{xy}}(z) && \text{by Lemma 7.1 page 53} \\
\tilde{\mathsf{S}}_{\mathsf{sy}}(\omega) &= \check{\mathsf{S}}_{\mathsf{sy}}(z)\big|_{z=e^{i\omega}} && \text{by definition of } \mathbf{Z} && \text{(Definition Q.4 page 354)} \\
&= \check{\mathsf{H}}(z)\check{\mathsf{S}}_{\mathsf{xy}}(z)\big|_{z=e^{i\omega}} && \text{by previous result} && (1) \\
&= \tilde{\mathsf{H}}(\omega)\tilde{\mathsf{S}}_{\mathsf{xy}}(\omega)
\end{aligned}$$

COPYRIGHT © 2019 DANIEL J. GREENHOE

Copyright © 2019 Daniel J. Greenhoe

Part III

Statistical Estimation

CHAPTER 8

ESTIMATION OVERVIEW

8.1 Estimation types

Estimation types. Let $\mathsf{x}(t;\theta)$ be a waveform with parameter θ. There are three basic types of estimation of x:

1. *detection*:
   - The waveform $\mathsf{x}(t;\theta_n)$ is known except for the value of parameter θ_n.
   - The parameter θ_n is one of a finite set of values.
   - Estimate θ_n and thereby also estimate $\mathsf{x}(t;\theta)$.
2. *parametric* estimation:
   - The waveform $\mathsf{x}(t;\theta)$ is known except for the value of parameter θ.
   - The parameter θ is one of an infinite set of values.
   - Estimate θ and thereby also estimate $\mathsf{x}(t;\theta)$.
3. *nonparametric* estimation:
   - The waveform $\mathsf{x}(t)$ is unknown and assumed without any parameter θ.
   - Estimate $\mathsf{x}(t)$.

Estimation criterion. Optimization requires a criterion against which the quality of an estimate is measured.[1] The most demanding and general criterion is the *Bayesian* criterion. The Bayesian criterion requires knowledge of the probability distribution functions and the definition of a *cost function*. Other criterion are special cases of the Bayesian criterion such that the cost function is defined in a special way, no cost function is defined, and/or the distribution is not known (Figure 8.2 page 66).

Estimation techniques. Estimation techniques can be classified into five groups (Figure 8.2 page 66):[2]

[1] Mandyam D. Srinath (1996) ⟨013125295X⟩.

[2] Nelles (2001) page 26 ⟨"Fig 2.2 Overview of linear and nonlinear optimization techniques"⟩, Nelles (2001) page 33 ⟨"Fig 2.5 The Bayes method is the most general approach but…"⟩, Nelles (2001) page 63 ⟨"Table 3.3 Relationship between linear recursive and nonlinear optimization techniques"⟩, Nelles (2001) page 66

1. sequential decoding
2. norm minimization
3. gradient search
4. inner product analysis
5. direct search

Sequential decoding is a non-linear estimation family. Perhaps the most famous of these is the Veterbi algorithm which uses a trellis to calculate the estimate. The Verterbi algorithm has been shown to yield an optimal estimate in the maximal likelihood (ML) sense. Norm minimization and gradient search algorithms are all linear algorithms. While this restriction to linear operations often simplifies calculations, it often yields an estimate that is not optimal in the ML sense.

8.2 Estimation criterion



Figure 8.1: Estimation criterion

Definition 8.1. *Let*

(A). $\mathsf{x}(t;\theta)$ *be a random process with unknown parameter* θ

(B). $\mathsf{y}(t)$ *an observed random process which is statistically dependent on* $\mathsf{x}(t;\theta)$

(C). $\mathsf{C}(\theta, \mathsf{p}(x, y))$ *be a cost function.*

Then the following ***estimates*** *are defined as described here:*

DEF

(1). ***Bayesian estimate*** $\hat{\theta}_{\mathsf{B}} \triangleq \arg\min_{\theta} C(\theta, \mathsf{p}(x, y))$

(2). ***Mean square estimate*** ***("MS estimate")*** $\hat{\theta}_{\mathsf{mms}} \triangleq \arg\min_{\theta} \mathsf{E}\, \|C(\theta, \mathsf{p}(x, y))\|^2$

(3). ***mini-max estimate*** ***("MM estimate")*** $\hat{\theta}_{\mathsf{mm}} \triangleq \arg\min_{\theta} \max_{\mathsf{p}} C(\theta, \mathsf{p}(x, y))$

(4). ***maximum a-posteriori probability estimate*** ***("MAP estimate")*** $\hat{\theta}_{\mathsf{map}} \triangleq \arg\max_{\theta} \mathsf{P}\{\mathsf{x}(t;\theta)|\mathsf{y}(t)\}$

(5). ***maximum likelihood estimate*** ***("ML estimate")*** $\hat{\theta}_{\mathsf{ml}} \triangleq \arg\max_{\theta} \mathsf{P}\{\mathsf{y}(t)|\mathsf{x}(t;\theta)\}$

BY-NC-ND Copyright © 2019 Daniel J. Greenhoe

Theorem 8.1. *Let* $\mathsf{x}(t;\theta)$ *be a random process with unknown parameter* θ.

THM $\{\mathsf{P}\{\theta\} = \text{CONSTANT}\} \quad \Longrightarrow \quad \{\hat{\theta}_{\mathsf{map}} = \hat{\theta}_{\mathsf{ml}}\}$

✎PROOF:

$$\begin{aligned}
\hat{\theta}_{\mathsf{map}} &\triangleq \arg\max_{\theta} \mathsf{P}\{\mathsf{x}(t;\theta)|\mathsf{y}(t)\} && \text{by definition of } \hat{\theta}_{\mathsf{map}} && \text{(Definition 8.1 page 64)}\\
&\triangleq \arg\max_{\theta} \frac{\mathsf{P}\{\mathsf{x}(t;\theta)\wedge \mathsf{y}(t)\}}{\mathsf{P}\{r(t)\}} && \text{by definition of } \textit{conditional probability} && \text{(Definition A.4 page 166)}\\
&\triangleq \arg\max_{\theta} \frac{\mathsf{P}\{r(t)|\mathsf{x}(t;\theta)\}\mathsf{P}\{\mathsf{x}(t;\theta)\}}{\mathsf{P}\{\mathsf{y}(t)\}} && \text{by definition of } \textit{conditional probability} && \text{(Definition A.4 page 166)}\\
&= \arg\max_{\theta} \mathsf{P}\{\mathsf{y}(t)|\mathsf{x}(t;\theta)\}\mathsf{P}\{\mathsf{x}(t;\theta)\} && \text{because } \mathsf{y}(t) \text{ is independent of } \theta\\
&= \arg\max_{\theta} \mathsf{P}\{\mathsf{y}(t)|\mathsf{x}(t;\theta)\}\\
&\triangleq \hat{\theta}_{\mathsf{ml}} && \text{by definition of } \hat{\theta}_{\mathsf{ml}} && \text{(Definition 8.1 page 64)}
\end{aligned}$$

✎

8.3 Measures of estimator quality

Definition 8.2. [3]

DEF *The* ***mean square error*** $\mathsf{mse}(\hat{\theta})$ *of an estimate* $\hat{\theta}$ *of a parameter* θ *is defined as*

$$\mathsf{mse}(\hat{\theta}) \triangleq \mathsf{E}\left[\left(\hat{\theta}-\theta\right)^2\right]$$

Definition 8.3. [4]

DEF *The* ***normalized rms error*** $\epsilon(\hat{\theta})$ *of an estimate* $\hat{\theta}$ *of a parameter* θ *is defined as*

$$\epsilon(\hat{\theta}) \triangleq \frac{\sqrt{\mathsf{mse}(\hat{\theta})}}{\theta} \triangleq \frac{\sqrt{\mathsf{E}\left[\left(\hat{\theta}-\theta\right)^2\right]}}{\theta}$$

Definition 8.4. [5]

DEF *The* ***mean integrated square error*** $\mathsf{mse}(\hat{\theta})$ *of an estimate* $\hat{\theta}$ *of a parameter* θ *is defined as*

$$\mathsf{mse}(\hat{\theta}) \triangleq \mathsf{E}\int_{\theta\in\mathbb{R}}\left[\left(\hat{\theta}-\theta\right)^2\right]$$

The *mean square error* of $\hat{\theta}$ can be expressed as the sum of two components: the variance of $\hat{\theta}$ and the bias of $\hat{\theta}$ squared (next Theorem). For an example of Theorem 8.2 in action, see the proof for the $\mathsf{mse}(\hat{\mu})$ of the *arithmetic mean estimate* as provided in Theorem 12.1 (page 95).

Theorem 8.2. [6] *Let* $\mathsf{mse}(\hat{\theta})$ *be the* MEAN SQUARE ERROR *(Definition 8.2 page 65) and* $\epsilon(\hat{\theta})$ *the* NORMALIZED

[3] Silverman (1986) page 35 ⟨§"1.3.2 Measures of discrepancy…"⟩, Bendat and Piersol (2010) ⟨§"1.4.3 Error Analysis Criteria"⟩, Bendat and Piersol (1966), page 183§"5.3 Statistical Errors for Parameter Estimates"

[4] Bendat and Piersol (2010) ⟨§"1.4.3 Error Analysis Criteria"⟩

[5] Silverman (1986) page 35 ⟨§"1.3.2 Measures of discrepancy…"⟩, Rosenblatt (1956) page 835 ⟨"integrated mean square error"⟩

[6] Choi (1978) page 76, Kay (1988) page 45 ⟨§"3.3 Estimation Theory"⟩, Stuart and Ord (1991) page 629 ⟨"Minium mean-square-error estimation"⟩, Clarkson (1993) page 51 ⟨§"2.6 Estimation of Moments"⟩, Bendat and Piersol (2010) ⟨§"1.4.3 Error Analysis Criteria"⟩, Bendat and Piersol (1966), page 183§"5.3 Statistical Errors for Parameter Estimates", Bendat and Piersol (1980) page 39 ⟨§"2.4.1 Bias versus Random Errors"⟩

Copyright © 2019 Daniel J. Greenhoe

RMS ERROR *(Definition 8.3 page 65) of an estimator* $\hat{\theta}$.

THM

$$\mathsf{mse}(\hat{\theta}) = \underbrace{\mathsf{E}\left[\left(\hat{\theta}-\mathsf{E}\hat{\theta}\right)^2\right]}_{\textit{variance of }\hat{\theta}} + \underbrace{\left[\mathsf{E}\hat{\theta}-\theta\right]^2}_{\textit{bias of }\hat{\theta}\textit{ squared}} \qquad \epsilon(\hat{\theta}) = \frac{\sqrt{\mathsf{E}\left[\left(\hat{\theta}-\mathsf{E}\hat{\theta}\right)^2\right]+\left[\mathsf{E}\hat{\theta}-\theta\right]^2}}{\theta}$$

✎PROOF:

$$\begin{aligned}
\mathsf{mse}(\hat{\theta}) &\triangleq \mathsf{E}\left[\left(\hat{\theta}-\theta\right)^2\right] && \text{by definition of mse} && \text{(Definition 8.2 page 65)}\\
&= \mathsf{E}\left[\left(\hat{\theta}\underbrace{-\mathsf{E}\hat{\theta}+\mathsf{E}\hat{\theta}}_{0}-\theta\right)^2\right] && \text{by } \textit{additive identity} \text{ property of } (\mathbb{C},+,\cdot,0,1)\\
&= \mathsf{E}\left[\left(\hat{\theta}-\mathsf{E}\hat{\theta}\right)^2+\underbrace{\left(\mathsf{E}\hat{\theta}-\theta\right)^2}_{\text{constant}}-2\left(\hat{\theta}-\mathsf{E}\hat{\theta}\right)\left(\mathsf{E}\hat{\theta}-\theta\right)\right] && \text{by } \textit{Binomial Theorem}\\
&= \mathsf{E}\left(\hat{\theta}-\mathsf{E}\hat{\theta}\right)^2+\left(\mathsf{E}\hat{\theta}-\theta\right)^2-2\mathsf{E}\left[\hat{\theta}\mathsf{E}\hat{\theta}-\hat{\theta}\theta-\mathsf{E}\hat{\theta}\hat{\theta}+\mathsf{E}\hat{\theta}\theta\right] && \text{by } \textit{linearity} \text{ of } \mathsf{E} && \text{(Theorem 1.1 page 4)}\\
&= \mathsf{E}\left(\hat{\theta}-\mathsf{E}\hat{\theta}\right)^2+\left(\mathsf{E}\hat{\theta}-\theta\right)^2-2\underbrace{\left[\mathsf{E}\hat{\theta}\mathsf{E}\hat{\theta}-\mathsf{E}\hat{\theta}\mathsf{E}\theta-\mathsf{E}\hat{\theta}\mathsf{E}\hat{\theta}+\mathsf{E}\hat{\theta}\mathsf{E}\theta\right]}_{0} && \text{by } \textit{linearity} \text{ of } \mathsf{E} && \text{(Theorem 1.1 page 4)}\\
&= \mathsf{E}\left(\hat{\theta}-\mathsf{E}\hat{\theta}\right)^2+\left(\mathsf{E}\hat{\theta}-\theta\right)^2
\end{aligned}$$

✏

Definition 8.5. [7]

DEF

An estimate $\hat{\theta}$ *of a parameter* θ *is a* ***minimum variance unbiased estimator*** *(MVUE) if*

(1). $\mathsf{E}\hat{\theta} = 0$ (UNBIASED) *and*

(2). *no other unbiased estimator* $\hat{\phi}$ *has smaller variance* $\mathrm{var}(\hat{\phi})$

8.4 Estimation techniques



Figure 8.2: Estimation techniques

[7] Choi (1978) page 76, Shao (2003) page 161 ⟨§"The UMVUE"⟩, Bolstad (2007) page 164 ⟨§"Minimum Variance Unbiased Estimator"⟩,

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

8.5 Sequential decoding



Figure 8.3: Viterbi algorithm trellis

It has been shown that the Viterbi algorithm (trellis) produces an optimal estimate in the maximal likelihood (ML) sense. A Verterbi trellis is shown in Figure 8.3 (page 67).

Copyright © 2019 Daniel J. Greenhoe
BY-NC-ND

Copyright © 2019 Daniel J. Greenhoe

CHAPTER 9

NORM MINIMIZATION



Figure 9.1: Estimation using gradient of cost function

Norm minimization techniques are very powerful in that an optimum solution can be computed in one step without iteration or recursion. In this section we present two types of norm minimization:[1]

1. minimum mean square estimation (MMSE):
   The MMS estimate is a *stochastic* estimate. To compute the MMS estimate, the we do not need to know the actual data values, but we must know certain system statistics which are the input data autocorrelation and input/output crosscorrelation. The cost function is the expected value of the norm squared error.
2. least square estimation (LSE):
   The LS estimate is a *deterministic* estimate. To compute the LS estimate, we must know the actual data values (although these may be "noisy" measurements). The cost function is the norm squared error.

Solutions to both are given in terms of two matrices:

[1] The Least Squares algorithm is nothing new to mathematics. It was first published by Legendre in 1805, but there is a credible claim by Gauss that he had it as far back as 1795. Gauss, by the way, was also the first to discover the FFT. References: Sorenson (1970) page 63, Plackett (1972), Stigler (1981), Dutka (1995)

Y: Autocorrelation matrix
W: Crosscorrelation matrix.

9.1 Minimum mean square estimation

Definition 9.1. *Let the following vectors, matrices, and functions be defined as follows:*

DEF

$\boldsymbol{x}$	$\in$	$\mathbb{C}^m$	*data vector*
$\boldsymbol{y}$	$\in$	$\mathbb{C}^n$	*processed data vector*
$\hat{\boldsymbol{y}}$	$\in$	$\mathbb{C}^n$	*processed data estimate vector*
$\boldsymbol{e}$	$\in$	$\mathbb{C}^n$	*error vector*
$\boldsymbol{p}$	$\in$	$\mathbb{R}^m$	*parameter vector*
U	$\in$	C_{mn}	*regression matrix*
R	$\in$	C_{mm}	*autocorrelation matrix*
W	$\in$	$\mathbb{C}^m$	*cross-correlation vector*
C	$:$	$\mathbb{R}^m \to \mathbb{R}^{\vdash}$	*cost function*

Theorem 9.1 (Minimum mean square estimation)**.** *Let*

$$\begin{aligned}
\hat{\boldsymbol{y}}(\boldsymbol{p}) &\triangleq U^H\boldsymbol{p}\\
\boldsymbol{e}(\boldsymbol{p}) &\triangleq \hat{\boldsymbol{y}} - \boldsymbol{y}\\
\mathsf{C}(\boldsymbol{p}) &\triangleq \mathsf{E}\,\|\boldsymbol{e}\|^2 \triangleq \mathsf{E}\left[\boldsymbol{e}^H\boldsymbol{e}\right]\\
\hat{\theta}_{\mathsf{mms}} &\triangleq \arg\min_{\boldsymbol{p}} \mathsf{C}(\boldsymbol{p})\\
R &\triangleq \mathsf{E}\left[UU^H\right]\\
W &\triangleq \mathsf{E}\left[U\boldsymbol{y}\right].
\end{aligned}$$

Then

THM

$$\begin{aligned}
\hat{\theta}_{\mathsf{mms}} &= (\mathbf{R}_{\mathsf{e}}Y)^{-1}(\mathbf{R}_{\mathsf{e}}W)\\
\mathsf{C}(\boldsymbol{p}) &= \boldsymbol{p}^H R\boldsymbol{p} - (W^H\boldsymbol{p})^* - W^H\boldsymbol{p} + \mathsf{E}\boldsymbol{y}^H\boldsymbol{y}\\
\nabla_{\boldsymbol{p}}\mathsf{C}(\boldsymbol{p}) &= 2\mathbf{R}_{\mathsf{e}}[Y]\boldsymbol{p} - 2\mathbf{R}_{\mathsf{e}}W\\
\mathsf{C}(\hat{\theta}_{\mathsf{mms}}) &= \begin{cases} \mathsf{E}\boldsymbol{y}^H\boldsymbol{y} + (\mathbf{R}_{\mathsf{e}}W^H)(\mathbf{R}_{\mathsf{e}}Y)^{-1}R(\mathbf{R}_{\mathsf{e}}Y)^{-1}(\mathbf{R}_{\mathsf{e}}W) - 2(\mathbf{R}_{\mathsf{e}}W^H)(\mathbf{R}_{\mathsf{e}}Y)^{-1}(\mathbf{R}_{\mathsf{e}}W) \\ \mathsf{E}\boldsymbol{y}^H\boldsymbol{y} - (\mathbf{R}_{\mathsf{e}}W^H)R^{-1}(\mathbf{R}_{\mathsf{e}}W) \quad \textit{if } R \textit{ is } \textbf{\textit{real-valued}} \end{cases}
\end{aligned}$$

✎PROOF: See APPENDIX E (page 193) for a Matrix Calculus reference.

1. Proof that cost $\mathsf{C}(\boldsymbol{p}) = \boldsymbol{p}^H R\boldsymbol{p} - (W^H)^*\boldsymbol{p} - W^H\boldsymbol{p} + \mathsf{E}\boldsymbol{y}^H\boldsymbol{y}$:

$$\begin{aligned}
\boxed{\mathsf{C}(\boldsymbol{p})} &\triangleq \mathsf{E}\,\|\boldsymbol{e}\|^2 && \text{by definition of } \textit{cost function}\ \mathsf{C}\\
&\triangleq \mathsf{E}\left[\boldsymbol{e}^H\boldsymbol{e}\right] && \text{by definition of } \textit{norm}\ \|\cdot\|\\
&\triangleq \mathsf{E}\left[(\hat{\boldsymbol{y}}-\boldsymbol{y})^H(\hat{\boldsymbol{y}}-\boldsymbol{y})\right] && \text{by definition of } \textit{error vector}\ \boldsymbol{e}\\
&\triangleq \mathsf{E}\left[\left(U^H\boldsymbol{p}-\boldsymbol{y}\right)^H\left(U^H\boldsymbol{p}-\boldsymbol{y}\right)\right] && \text{by definition of estimate } \hat{\boldsymbol{y}}\\
&= \mathsf{E}\left[\left(\boldsymbol{p}^HU-\boldsymbol{y}^H\right)\left(U^H\boldsymbol{p}-\boldsymbol{y}\right)\right] && \text{by } \textit{distributive} \text{ prop. of } \textit{*-algebras} \quad \text{(Definition I.3 page 250)}\\
&= \mathsf{E}\left[\boldsymbol{p}^HUU^H\boldsymbol{p} - \boldsymbol{p}^HU\boldsymbol{y} - \boldsymbol{y}^HU^H\boldsymbol{p} + \boldsymbol{y}^H\boldsymbol{y}\right] && \text{by } \textit{matrix algebra} \text{ ring property}
\end{aligned}$$

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

$$
\begin{aligned}
&= p^H \mathsf{E}\left[UU^H\right]p - p^H \mathsf{E}\left[U\boldsymbol{y}\right] - \mathsf{E}\left[\boldsymbol{y}^H U^H\right]p + \mathsf{E}\boldsymbol{y}^H\boldsymbol{y} && \text{by } \textit{linearity } \mathsf{E} \\
&= p^H \mathsf{E}\left[UU^H\right]p - (\mathsf{E}\left[U\boldsymbol{y}\right]^H p)^H - \mathsf{E}\left[U\boldsymbol{y}\right]^H p + \mathsf{E}\boldsymbol{y}^H\boldsymbol{y} \\
&= p^H Rp - (W^H p)^H - W^H p + \mathsf{E}\boldsymbol{y}^H\boldsymbol{y} \\
&= p^H Rp - (W^H p)^* - W^H p + \mathsf{E}\boldsymbol{y}^H\boldsymbol{y} \\
&= \boxed{p^H Rp - (W^H)^* p - W^H p + \mathsf{E}\boldsymbol{y}^H\boldsymbol{y}} \\
&= p^H Rp - 2\mathbf{R}_{\mathsf{e}}\left[W^H\right]p + \mathsf{E}\boldsymbol{y}^H\boldsymbol{y}
\end{aligned}
$$

(Theorem 1.1 page 4

2. Proof that optimal $p_{\text{opt}} = (\mathbf{R}_{\mathsf{e}}Y)^{-1}(\mathbf{R}_{\mathsf{e}}W)$:

$$
\begin{aligned}
\nabla_p \mathsf{C}(p) &= \nabla_p\left[p^H Rp - (W^H)^* p - W^H p + \mathsf{E}\boldsymbol{y}^H\boldsymbol{y}\right] && \text{by previous resut} \\
&= Rp + R^T p - \nabla_p\left[(W^H)^* p + W^H p\right] + 0 && \text{by } \textit{quadratic form} \text{ result (Theorem E.6 page 198)} \\
&= Rp + R^T p - [(W^H)^*]^T - [W^H]^T + 0 && \text{by } \textit{affine equations} \text{ result (Theorem E.3 page 197)} \\
&= Rp + (R^H)^* p - W - W^* && \text{by definition of } \textit{Hermitian Transpose}^H \\
&= Rp + R^* p - W - W^* && \text{because } R \text{ is } \textit{Hermitian symmetric} \\
&= (R + R^*)p - (W + W^*) && \text{by } \textit{ring} \text{ property} \\
&= 2(\mathbf{R}_{\mathsf{e}}Y)p - 2\mathbf{R}_{\mathsf{e}}W && \text{by definition of } \mathbf{R}_{\mathsf{e}} \text{ (Definition I.5 page 251)} \\
\Longrightarrow\ p_{\text{opt}} &= (\mathbf{R}_{\mathsf{e}}Y)^{-1}(\mathbf{R}_{\mathsf{e}}W) && \text{by setting } \nabla_p \mathsf{C}(p) = 0
\end{aligned}
$$

3. Cost of optimal p_{opt}:

$$
\begin{aligned}
\mathsf{C}(p_{\text{opt}}) &= p_{\text{opt}}^H R p_{\text{opt}} - 2\mathbf{R}_{\mathsf{e}}\left[W^H\right]p_{\text{opt}} + \mathsf{E}\boldsymbol{y}^H\boldsymbol{y} && \text{by item (1)} \\
&= [(\mathbf{R}_{\mathsf{e}}Y)^{-1}(\mathbf{R}_{\mathsf{e}}W)]^H R[(\mathbf{R}_{\mathsf{e}}Y)^{-1}(\mathbf{R}_{\mathsf{e}}W)] - 2\mathbf{R}_{\mathsf{e}}\left[W^H\right][(\mathbf{R}_{\mathsf{e}}Y)^{-1}(\mathbf{R}_{\mathsf{e}}W)] + \mathsf{E}\boldsymbol{y}^H\boldsymbol{y} && \text{by item (2)} \\
&= (\mathbf{R}_{\mathsf{e}}W^H)(\mathbf{R}_{\mathsf{e}}Y)^{-H} R(\mathbf{R}_{\mathsf{e}}Y)^{-1}(\mathbf{R}_{\mathsf{e}}W) - 2\mathbf{R}_{\mathsf{e}}\left[W^H\right](\mathbf{R}_{\mathsf{e}}Y)^{-1}(\mathbf{R}_{\mathsf{e}}W) + \mathsf{E}\boldsymbol{y}^H\boldsymbol{y} \\
&= (\mathbf{R}_{\mathsf{e}}W^H)(\mathbf{R}_{\mathsf{e}}R^H)^{-1} R(\mathbf{R}_{\mathsf{e}}Y)^{-1}(\mathbf{R}_{\mathsf{e}}W) - 2\mathbf{R}_{\mathsf{e}}\left[W^H\right](\mathbf{R}_{\mathsf{e}}Y)^{-1}(\mathbf{R}_{\mathsf{e}}W) + \mathsf{E}\boldsymbol{y}^H\boldsymbol{y} \\
&= (\mathbf{R}_{\mathsf{e}}W^H)(\mathbf{R}_{\mathsf{e}}Y)^{-1} R(\mathbf{R}_{\mathsf{e}}Y)^{-1}(\mathbf{R}_{\mathsf{e}}W) - 2(\mathbf{R}_{\mathsf{e}}W^H)(\mathbf{R}_{\mathsf{e}}Y)^{-1}(\mathbf{R}_{\mathsf{e}}W) + \mathsf{E}\boldsymbol{y}^H\boldsymbol{y}
\end{aligned}
$$

$$
\begin{aligned}
\mathsf{C}(p_{\text{opt}})|_{R\text{ real}} &= (\mathbf{R}_{\mathsf{e}}W^H)(\mathbf{R}_{\mathsf{e}}Y)^{-1} R(\mathbf{R}_{\mathsf{e}}Y)^{-1}(\mathbf{R}_{\mathsf{e}}W) - 2(\mathbf{R}_{\mathsf{e}}W^H)(\mathbf{R}_{\mathsf{e}}Y)^{-1}(\mathbf{R}_{\mathsf{e}}W) + \mathsf{E}\boldsymbol{y}^H\boldsymbol{y} \\
&= (\mathbf{R}_{\mathsf{e}}W^H)R^{-1}RR^{-1}(\mathbf{R}_{\mathsf{e}}W) - 2(\mathbf{R}_{\mathsf{e}}W^H)R^{-1}(\mathbf{R}_{\mathsf{e}}W) + \mathsf{E}\boldsymbol{y}^H\boldsymbol{y} \\
&= (\mathbf{R}_{\mathsf{e}}W^H)R^{-1}(\mathbf{R}_{\mathsf{e}}W) - 2(\mathbf{R}_{\mathsf{e}}W^H)R^{-1}(\mathbf{R}_{\mathsf{e}}W) + \mathsf{E}\boldsymbol{y}^H\boldsymbol{y} \\
&= \mathsf{E}\boldsymbol{y}^H\boldsymbol{y} - (\mathbf{R}_{\mathsf{e}}W^H)R^{-1}(\mathbf{R}_{\mathsf{e}}W)
\end{aligned}
$$

In many adaptive filter and equalization applications, the autocorrelation matrix U is simply the m-element random data vector $\boldsymbol{x}(k)$ at time k, as in the *Wiener-Hopf equations* (next).

Corollary 9.1 (Wiener-Hopf equations). [2]

COR

$$
\left\{ U \triangleq \boldsymbol{x}(k) \triangleq \begin{bmatrix} x(k) \\ x(k-1) \\ x(k-2) \\ \vdots \\ x(k-m+1) \end{bmatrix} \right\} \Longrightarrow \left\{ \begin{array}{rcl} \hat{\theta}_{\text{mms}} & = & R^{-1}W \\ \mathsf{C}(\hat{\theta}_{\text{mms}}) & = & W^T R^{-1} R R^{-1} W - 2W^T R^{-1} W + \mathsf{E}\boldsymbol{y}^T\boldsymbol{y} \end{array} \right\}
$$

[2] Ifeachor and Jervis (1993) pages 547–549 ⟨§"9.3 Basic Wiener filter theory"⟩, Ifeachor and Jervis (2002) pages 651–654 ⟨§"10.3 Basic Wiener filter theory"⟩, Kay (1988) page 51 ⟨§"3.3.3 Random Parameters"⟩

Copyright © 2019 Daniel J. Greenhoe



Figure 9.2: Adative filter example

✎PROOF: This is a special case of the more general case discussed in Theorem 9.1 (page 70). Here, the dimension of U is $m \times 1$ (n=1). As a result, $\boldsymbol{y}$, $\hat{\boldsymbol{y}}$, and $\boldsymbol{e}$ are simply scalar quantities (not vectors). In this special case, we have the following results (Figure 9.2 page 72):

$$\begin{aligned}
\hat{y}(\boldsymbol{p}) &\triangleq \boldsymbol{x}^T \boldsymbol{p} \\
e(\boldsymbol{p}) &\triangleq \hat{y} - y \\
\mathsf{C}(\boldsymbol{p}) &\triangleq \mathsf{E}\,\|\boldsymbol{e}\|^2 \triangleq \mathsf{E}\left[e^2\right] \\
\hat{\theta}_{\mathsf{mms}} &\triangleq \arg\min_{\boldsymbol{p}} \mathsf{C}(\boldsymbol{p}) \\
R &\triangleq \mathsf{E}\left[\boldsymbol{x}\boldsymbol{x}^T\right] \\
W &\triangleq \mathsf{E}[\boldsymbol{x} y] \\
\mathsf{C}(\boldsymbol{p}) &= \boldsymbol{p}^T R \boldsymbol{p} - 2W^T \boldsymbol{p} + \mathsf{E}\left[\boldsymbol{y}^T \boldsymbol{y}\right] \\
\nabla_{\boldsymbol{p}} \mathsf{C}(\boldsymbol{p}) &= 2R\boldsymbol{p} - 2W \\
\mathsf{C}(\hat{\theta}_{\mathsf{mms}})|_{R\ \text{real}} &= \mathsf{E}\boldsymbol{y}^T \boldsymbol{y} - W^T R^{-1} W.
\end{aligned}$$

✏

9.2 Least squares

Theorem 9.2 (Least squares)**.** *Let*

$$\begin{aligned}
\hat{\boldsymbol{y}}(\boldsymbol{p}) &\triangleq U^H \boldsymbol{p} \\
\boldsymbol{e}(\boldsymbol{p}) &\triangleq \hat{\boldsymbol{y}} - \boldsymbol{y} \\
\mathsf{C}(\boldsymbol{p}) &\triangleq \|\boldsymbol{e}\|^2 \triangleq \boldsymbol{e}^H \boldsymbol{e} \\
\hat{\theta}_{\mathsf{ls}} &\triangleq \arg\min_{\boldsymbol{p}} \mathsf{C}(\boldsymbol{p}) \\
R &\triangleq U U^H \\
W &\triangleq U \boldsymbol{y}.
\end{aligned}$$

COPYRIGHT © 2019 DANIEL J. GREENHOE

Then

THM

$$\begin{aligned}
\hat{\theta}_{\mathsf{ls}} &= (\mathbf{R}_{\mathsf{e}}Y)^{-1}(\mathbf{R}_{\mathsf{e}}W)\\
\mathsf{C}(\boldsymbol{p}) &= \boldsymbol{p}^H R\boldsymbol{p} - (W^H\boldsymbol{p})^* - W^H\boldsymbol{p} + \mathsf{E}\boldsymbol{y}^H\boldsymbol{y}\\
\nabla_{\boldsymbol{p}}\mathsf{C}(\boldsymbol{p}) &= 2\mathbf{R}_{\mathsf{e}}[Y]\boldsymbol{p} - 2\mathbf{R}_{\mathsf{e}}W\\
\mathsf{C}(\hat{\theta}_{\mathsf{ls}}) &= (\mathbf{R}_{\mathsf{e}}W^H)(\mathbf{R}_{\mathsf{e}}Y)^{-1}R(\mathbf{R}_{\mathsf{e}}Y)^{-1}(\mathbf{R}_{\mathsf{e}}W) - 2(\mathbf{R}_{\mathsf{e}}W^H)(\mathbf{R}_{\mathsf{e}}Y)^{-1}(\mathbf{R}_{\mathsf{e}}W) + \mathsf{E}\boldsymbol{y}^H\boldsymbol{y}\\
\mathsf{C}(\hat{\theta}_{\mathsf{ls}})|_{R\ real} &= \mathsf{E}\boldsymbol{y}^H\boldsymbol{y} - (\mathbf{R}_{\mathsf{e}}W^H)R^{-1}(\mathbf{R}_{\mathsf{e}}W).
\end{aligned}$$

✎PROOF: See APPENDIX E (page 193) for a Matrix Calculus reference.

$$\begin{aligned}
\mathsf{C}(\boldsymbol{p}) &\triangleq \|\boldsymbol{e}\|^2\\
&= \boldsymbol{e}^H\boldsymbol{e}\\
&= (\hat{\boldsymbol{y}} - \boldsymbol{y})^H(\hat{\boldsymbol{y}} - \boldsymbol{y})\\
&= \left(U^H\boldsymbol{p} - \boldsymbol{y}\right)^H\left(U^H\boldsymbol{p} - \boldsymbol{y}\right)\\
&= \left(\boldsymbol{p}^HU - \boldsymbol{y}^H\right)\left(U^H\boldsymbol{p} - \boldsymbol{y}\right)\\
&= \boldsymbol{p}^HUU^H\boldsymbol{p} - \boldsymbol{p}^HU\boldsymbol{y} - \boldsymbol{y}^HU^H\boldsymbol{p} + \boldsymbol{y}^H\boldsymbol{y}\\
&= \boldsymbol{p}^HR\boldsymbol{p} - (W^H\boldsymbol{p})^H - W^H\boldsymbol{p} + \boldsymbol{y}^H\boldsymbol{y}\\
&= \boldsymbol{p}^HR\boldsymbol{p} - (W^H\boldsymbol{p})^* - W^H\boldsymbol{p} + \boldsymbol{y}^H\boldsymbol{y}\\
&= \boldsymbol{p}^HR\boldsymbol{p} - (W^H)^*\boldsymbol{p} - W^H\boldsymbol{p} + \boldsymbol{y}^H\boldsymbol{y}\\
&= \boldsymbol{p}^HR\boldsymbol{p} - 2\mathbf{R}_{\mathsf{e}}\left[W^H\right]\boldsymbol{p} + \boldsymbol{y}^H\boldsymbol{y}
\end{aligned}$$

$$\begin{aligned}
\nabla_{\boldsymbol{p}}\mathsf{C}(\boldsymbol{p}) &= \nabla_{\boldsymbol{p}}\left[\boldsymbol{p}^HR\boldsymbol{p} - (W^H)^*\boldsymbol{p} - W^H\boldsymbol{p} + \boldsymbol{y}^H\boldsymbol{y}\right]\\
&= R\boldsymbol{p} + R^T\boldsymbol{p} - [(W^H)^*]^T - [W^H]^T + 0\\
&= R\boldsymbol{p} + (R^H)^*\boldsymbol{p} - W - W^*\\
&= R\boldsymbol{p} + R^*\boldsymbol{p} - W - W^*\\
&= (R + R^*)\boldsymbol{p} - (W + W^*)\\
&= 2(\mathbf{R}_{\mathsf{e}}Y)\boldsymbol{p} - 2\mathbf{R}_{\mathsf{e}}W
\end{aligned}$$

$$\boldsymbol{p}_{\mathsf{opt}} = (\mathbf{R}_{\mathsf{e}}Y)^{-1}(\mathbf{R}_{\mathsf{e}}W)$$

$$\begin{aligned}
\mathsf{C}(\boldsymbol{p}_{\mathsf{opt}}) &= \boldsymbol{p}_{\mathsf{opt}}^HR\boldsymbol{p}_{\mathsf{opt}} - 2\mathbf{R}_{\mathsf{e}}\left[W^H\right]\boldsymbol{p}_{\mathsf{opt}} + \boldsymbol{y}^H\boldsymbol{y}\\
&= [(\mathbf{R}_{\mathsf{e}}Y)^{-1}(\mathbf{R}_{\mathsf{e}}W)]^HR[(\mathbf{R}_{\mathsf{e}}Y)^{-1}(\mathbf{R}_{\mathsf{e}}W)] - 2\mathbf{R}_{\mathsf{e}}\left[W^H\right][(\mathbf{R}_{\mathsf{e}}Y)^{-1}(\mathbf{R}_{\mathsf{e}}W)] + \boldsymbol{y}^H\boldsymbol{y}\\
&= (\mathbf{R}_{\mathsf{e}}W^H)(\mathbf{R}_{\mathsf{e}}Y)^{-H}R(\mathbf{R}_{\mathsf{e}}Y)^{-1}(\mathbf{R}_{\mathsf{e}}W) - 2\mathbf{R}_{\mathsf{e}}\left[W^H\right](\mathbf{R}_{\mathsf{e}}Y)^{-1}(\mathbf{R}_{\mathsf{e}}W) + \boldsymbol{y}^H\boldsymbol{y}\\
&= (\mathbf{R}_{\mathsf{e}}W^H)(\mathbf{R}_{\mathsf{e}}R^H)^{-1}R(\mathbf{R}_{\mathsf{e}}Y)^{-1}(\mathbf{R}_{\mathsf{e}}W) - 2\mathbf{R}_{\mathsf{e}}\left[W^H\right](\mathbf{R}_{\mathsf{e}}Y)^{-1}(\mathbf{R}_{\mathsf{e}}W) + \boldsymbol{y}^H\boldsymbol{y}\\
&= (\mathbf{R}_{\mathsf{e}}W^H)(\mathbf{R}_{\mathsf{e}}Y)^{-1}R(\mathbf{R}_{\mathsf{e}}Y)^{-1}(\mathbf{R}_{\mathsf{e}}W) - 2(\mathbf{R}_{\mathsf{e}}W^H)(\mathbf{R}_{\mathsf{e}}Y)^{-1}(\mathbf{R}_{\mathsf{e}}W) + \boldsymbol{y}^H\boldsymbol{y}
\end{aligned}$$

$$\begin{aligned}
\mathsf{C}(\boldsymbol{p}_{\mathsf{opt}})|_{R\ \mathrm{real}} &= (\mathbf{R}_{\mathsf{e}}W^H)(\mathbf{R}_{\mathsf{e}}Y)^{-1}R(\mathbf{R}_{\mathsf{e}}Y)^{-1}(\mathbf{R}_{\mathsf{e}}W) - 2(\mathbf{R}_{\mathsf{e}}W^H)(\mathbf{R}_{\mathsf{e}}Y)^{-1}(\mathbf{R}_{\mathsf{e}}W) + \boldsymbol{y}^H\boldsymbol{y}\\
&= (\mathbf{R}_{\mathsf{e}}W^H)R^{-1}RR^{-1}(\mathbf{R}_{\mathsf{e}}W) - 2(\mathbf{R}_{\mathsf{e}}W^H)R^{-1}(\mathbf{R}_{\mathsf{e}}W) + \boldsymbol{y}^H\boldsymbol{y}\\
&= (\mathbf{R}_{\mathsf{e}}W^H)R^{-1}(\mathbf{R}_{\mathsf{e}}W) - 2(\mathbf{R}_{\mathsf{e}}W^H)R^{-1}(\mathbf{R}_{\mathsf{e}}W) + \boldsymbol{y}^H\boldsymbol{y}\\
&= \boldsymbol{y}^H\boldsymbol{y} - (\mathbf{R}_{\mathsf{e}}W^H)R^{-1}(\mathbf{R}_{\mathsf{e}}W)
\end{aligned}$$

✏

Example 9.1 (Polynomial approximation).
Suppose we **know** the locations $\{(x_n, y_n)|n = 1,2,3,4,5\}$ of 5 data points. Let $\boldsymbol{x}$ and $\boldsymbol{y}$ represent the

COPYRIGHT © 2019 DANIEL J. GREENHOE
CC BY-NC-ND

locations of these points such that

$$\boldsymbol{x} \triangleq \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \end{bmatrix} \qquad \boldsymbol{y} \triangleq \begin{bmatrix} y_1 \\ y_2 \\ y_3 \\ y_4 \\ y_5 \end{bmatrix}$$

Suppose we want to find a second order polynomial

$$cx^2 + bx + a$$

that best approximates these 5 points in the least squares sense. We define the matrix U (known) and vector $\hat{\theta}$ (to be computed) as follows:

$$U^H \triangleq \underbrace{\begin{bmatrix} 1 & x_1 & x_1^2 \\ 1 & x_2 & x_2^2 \\ 1 & x_3 & x_3^2 \\ 1 & x_4 & x_4^2 \\ 1 & x_5 & x_5^2 \end{bmatrix}}_{\text{Vandermonde matrix}}{}^{3} \qquad \hat{\theta} \triangleq \begin{bmatrix} a \\ b \\ c \end{bmatrix}$$

Then, using Theorem 9.2 (page 72), the best coefficients $\hat{\theta}$ for the polynomial are

$$\begin{aligned}
\hat{\theta} &= \begin{bmatrix} a \\ b \\ c \end{bmatrix} \\
&= R^{-1}W \\
&= (UU^H)^{-1}\,(U\boldsymbol{y}) \\
&= \left(\begin{bmatrix} 1 & x_1 & x_1^2 \\ 1 & x_2 & x_2^2 \\ 1 & x_3 & x_3^2 \\ 1 & x_4 & x_4^2 \\ 1 & x_5 & x_5^2 \end{bmatrix}^H \begin{bmatrix} 1 & x_1 & x_1^2 \\ 1 & x_2 & x_2^2 \\ 1 & x_3 & x_3^2 \\ 1 & x_4 & x_4^2 \\ 1 & x_5 & x_5^2 \end{bmatrix}\right)^{-1} \left(\begin{bmatrix} 1 & x_1 & x_1^2 \\ 1 & x_2 & x_2^2 \\ 1 & x_3 & x_3^2 \\ 1 & x_4 & x_4^2 \\ 1 & x_5 & x_5^2 \end{bmatrix}^H \begin{bmatrix} y_1 \\ y_2 \\ y_3 \\ y_4 \\ y_5 \end{bmatrix}\right)
\end{aligned}$$

[3] Horn and Johnson (1990)29

BY-NC-ND Copyright © 2019 Daniel J. Greenhoe

CHAPTER 10

GRADIENT SEARCH TECHNIQUES

10.1 Gradient search techniques

One of the biggest advantages of using a gradient search technique is that they can be implemented *recursively* as shown in the next equation. The general form of the gradient search parameter estimation techniques is[1]

THM $\boldsymbol{p}_n = \boldsymbol{p}_{n-1} - \eta_{n-1} R \left[\nabla_{\boldsymbol{p}} \mathsf{C}(\boldsymbol{p}_n)\right]$ where at time n

$\boldsymbol{p}_n$	is the *state*	(vector)
η_n	is the *step size*	(scalar)
Y	is the *direction*	(matrix)
$\nabla_{\boldsymbol{p}} \mathsf{C}(\boldsymbol{p}_n)$	is the *gradient* of the cost function $\mathsf{C}(\boldsymbol{p}_n)$	(vector)

Two major categories of gradient search techniques are

- steepest descent (includes LMS)
- Newton's method (includes RLS and Kalman filters).

The key difference between the two is that ***steepest descent* uses only first derivative information**, while ***Newton's method* uses both first and second derivative information** making it converge much faster but with significantly higher complexity.

First derivative techniques

Steepest descent. In this algorithm, $R = I$ (identity matrix). First derivative information is contained in $\nabla \mathsf{C}$. Second derivative information, if present, is contained in Y. Thus, steepest descent algorithms do not use second derivative information.

THM $\boldsymbol{p}_n = \boldsymbol{p}_{n-1} - \eta_{n-1} \left[\nabla_{\boldsymbol{p}} \mathsf{C}(\boldsymbol{p}_n)\right]$

[1] Nelles (2001), page 90

Least Mean Squares (LMS). [2] This is a special case of *steepest descent*. In minimum mean square estimation (Section 9.1 page 70), the cost function $\mathsf{C}(\boldsymbol{p})$ is defined as a *statistical average* of the error vector such that $\mathsf{C}(\boldsymbol{p}) = \mathsf{E}\left[\boldsymbol{e}^H \boldsymbol{e}\right]$. In this case the gradient $\nabla\mathsf{C}$ is difficult to compute. However, the LMS algorithm greatly simplifies the problem by instead defining the cost function as a function of the *instantaneous error* such that

$$\begin{aligned}
\boldsymbol{y} &= y(n)\\
\hat{\boldsymbol{y}} &= \hat{y}(n)\\
\mathsf{C}(\boldsymbol{p}) &= \|e(n)\|^2\\
&= e^2(n)\\
&= (\hat{y}(n) - y(n))^2
\end{aligned}$$

Computing the gradient of this cost function is then just a special case of *least squares estimation* (Section 9.2 page 72). Using LS, we let $U = \boldsymbol{x}^T$ and hence

$$\begin{aligned}
\nabla_{\boldsymbol{p}}\mathsf{C}(\boldsymbol{p}) &= 2U^T U\boldsymbol{p} - 2U^T\boldsymbol{y} && \text{by Theorem 9.2 page 72}\\
&= 2\boldsymbol{x}\boldsymbol{x}^T\boldsymbol{p} - 2\boldsymbol{x}y && \text{by above definitions}\\
&= 2\boldsymbol{x}\hat{y} - 2\boldsymbol{x}y\\
&= 2\boldsymbol{x}(\hat{y} - y)\\
&= 2\boldsymbol{x}e(n)
\end{aligned}$$

The LMS algorithm uses this instantaneous gradient for $\nabla\mathsf{C}$, lets $R = I$, and uses a constant step size η to give

THM $\boldsymbol{p}_n = \boldsymbol{p}_{n-1} - 2\eta\boldsymbol{x}_n e(n)$

Second derivative techniques

Newton's Method. This algorithm uses the *Hessian* matrix H, which is the second derivative of the cost function $\mathsf{C}(\boldsymbol{p})$, and lets $R = H^{-1}$.

$$\begin{aligned}
H_n &\triangleq && \nabla_{\boldsymbol{p}}\nabla_{\boldsymbol{p}}\mathsf{C}(\boldsymbol{p}_n)\\
\boldsymbol{p}_n &= \boldsymbol{p}_{n-1} - \eta_{n-1}H_n^{-1}\left[\nabla_{\boldsymbol{p}}\mathsf{C}(\boldsymbol{p}_n)\right]
\end{aligned}$$

Kalman filtering [3]

$$\begin{aligned}
\gamma(k) &= \frac{1}{x^T(k)P(k-1)\boldsymbol{x}(k) + 1}P(k-1)x(k)\\
P(k) &= (I - \gamma(k)x^T(k))P(k-1) + V\\
e(k) &= y(k) - x^T(k)\hat{\boldsymbol{p}}(k-1)\\
\hat{\boldsymbol{p}}(k) &= \hat{\boldsymbol{p}}(k-1) + \gamma(k)e(k)
\end{aligned}$$

[2] Manolakis et al. (2000), page 526
[3] Nelles (2001), page 66

BY-NC-ND Copyright © 2019 Daniel J. Greenhoe

Recursive Least Squares (RLS) [4] This algorithm is a special case of either the RLS with forgetting or the Kalman filter.

$$\begin{aligned}
\gamma(k) &= \frac{1}{x^T(k)P(k-1)\boldsymbol{x}(k)+1}P(k-1)x(k)\\
P(k) &= (I-\gamma(k)x^T(k))P(k-1)\\
e(k) &= y(k)-x^T(k)\hat{\boldsymbol{p}}(k-1)\\
\hat{\boldsymbol{p}}(k) &= \hat{\boldsymbol{p}}(k-1)+\gamma(k)e(k)
\end{aligned}$$

10.2 Direct search

A direct search algorithm may be used in cases where the cost function over $\boldsymbol{p}$ has several local minimums, making convergence difficult. Furthermore, direct search algorithms can be very computationally demanding.

[4] Nelles (2001), page 66

Copyright © 2019 Daniel J. Greenhoe
BY-NC-ND

Copyright © 2019 Daniel J. Greenhoe

CHAPTER 11

KL-EXPANSION APPLICATION

11.1 Sufficient statistics

Theorem 11.1 (page 79) (next) shows that the finite set $Y \triangleq \{\dot{y}_n | n = 1, 2, \ldots, N\}$ provides just as much information as having the entire $\mathsf{y}(t)$ waveform (an uncountably infinite number of values) with respect to the following cases:

1. the conditional probability of $\mathsf{x}(t;\hat{\theta})$ given $\mathsf{y}(t)$
2. the *MAP estimate* of the information sequence
3. the *ML estimate* of the information sequence.

That is, even with a drastic reduction in the amount of information from uncountably infinite to finite N, no information is lost with respect to the quantities listed above.

This amazing result is very useful in practical system implementation and also for proving other theoretical results (notably estimation and detection theorems which come later in this chapter).

Theorem 11.1 (Sufficient statistic theorem)**.** [1] *Let* **S** *be an additive White Gaussian noise system and* Ψ *an orthonormal basis for* $\mathsf{x}(t;\hat{\theta})$ *such that*

$$\begin{aligned} \mathsf{y}(t) &= \mathsf{x}(t;\hat{\theta}) + \mathsf{v}(t) \\ \Psi &= \{\psi_n | n = 1, 2, \ldots, N\} \end{aligned}$$

Then $Y \triangleq \{\dot{y}_n | n = 1, 2, \ldots, N\}$ *is a* ***sufficient statistic*** *for* $\mathsf{y}(t)$ *such that…*

THM

$$\left\{ \mathsf{v}(t) \text{ is AWGN} \right\} \Longrightarrow \left\{ \begin{array}{llll} (1). & \mathsf{P}\left\{\mathsf{x}(t;\hat{\theta})|\mathsf{y}(t)\right\} & = & \mathsf{P}\left\{\mathsf{x}(t;\hat{\theta})|Y\right\} \\ (2). & \hat{\theta}_{\mathsf{map}} \triangleq \arg\max_{\hat{\theta}} \mathsf{P}\left\{\mathsf{x}(t;\hat{\theta})|\mathsf{y}(t)\right\} & = & \arg\max_{\hat{\theta}} \mathsf{P}\left\{\mathsf{x}(t;\hat{\theta})|Y\right\} \\ (3). & \hat{\theta}_{\mathsf{ml}} \triangleq \arg\max_{\hat{\theta}} \mathsf{P}\left\{\mathsf{y}(t)|\mathsf{x}(t;\hat{\theta})\right\} & = & \arg\max_{\hat{\theta}} \mathsf{P}\left\{Y|\mathsf{x}(t;\hat{\theta})\right\} \end{array} \right\}$$

✎PROOF: Let $\mathsf{v}'(t) \triangleq \mathsf{v}(t) - \sum_{n=1}^{N} \dot{v}_n \psi_n(t)$.

[1] Fisher (1922) page 316 ⟨"Criterion of Sufficiency"⟩

1. The relationship between Y and $\mathsf{v}'(t)$ is given by

$$\begin{aligned}
\mathsf{y}(t) &= \sum_{n=1}^{N} \langle \mathsf{y}(t) \mid \psi_n(t) \rangle \psi_n(t) + \left[\mathsf{y}(t) - \sum_{n=1}^{N} \langle \mathsf{y}(t) \mid \psi_n(t) \rangle \psi_n(t) \right] \\
&= \sum_{n=1}^{N} \langle \mathsf{y}(t) \mid \psi_n(t) \rangle \psi_n(t) + \left[\mathsf{y}(t) - \sum_{n=1}^{N} \langle \mathsf{x}(t) + \mathsf{v}(t) \mid \psi_n(t) \rangle \psi_n(t) \right] \\
&= \sum_{n=1}^{N} \dot{y}_n \psi_n(t) + \left[\mathsf{x}(t) + \mathsf{v}(t) - \sum_{n=1}^{N} \langle \mathsf{x}(t) \mid \psi_n(t) \rangle \psi_n(t) - \sum_{n=1}^{N} \langle \mathsf{v}(t) \mid \psi_n(t) \rangle \psi_n(t) \right] \\
&= \sum_{n=1}^{N} \dot{y}_n \psi_n(t) + \mathsf{x}(t) + \mathsf{v}(t) - \mathsf{x}(t) - \left[\mathsf{v}(t) - \mathsf{v}'(t) \right] \\
&= \sum_{n=1}^{N} \dot{y}_n \psi_n(t) + \mathsf{v}'(t).
\end{aligned}$$

2. Proof that the set of statistics Y and the random process $\mathsf{v}'(t)$ are *uncorrelated*:

$$\begin{aligned}
\mathsf{E}\left[\dot{y}_n \mathsf{v}'(t)\right] &= \mathsf{E}\left[\langle \mathsf{y}(t) \mid \psi_n(t) \rangle \left(\mathsf{v}(t) - \sum_{n=1}^{N} \langle \mathsf{v}(t) \mid \psi_n(t) \rangle \psi_n(t) \right) \right] \\
&= \mathsf{E}\left[\langle \mathsf{x}(t) + \mathsf{v}(t) \mid \psi_n(t) \rangle \left(\mathsf{v}(t) - \sum_{n=1}^{N} \langle \mathsf{v}(t) \mid \psi_n(t) \rangle \psi_n(t) \right) \right] \\
&= \mathsf{E}\left[\left(\langle \mathsf{x}(t) \mid \psi_n(t) \rangle + \langle \mathsf{v}(t) \mid \psi_n(t) \rangle \right) \left(\mathsf{v}(t) - \sum_{n=1}^{N} \langle \mathsf{v}(t) \mid \psi_n(t) \rangle \psi_n(t) \right) \right] \\
&= \mathsf{E}\left[\left(\dot{x}_n + \dot{\upsilon}_n \right) \left(\mathsf{v}(t) - \sum_{n=1}^{N} \dot{\upsilon}_n \psi_n(t) \right) \right] \\
&= \mathsf{E}\left[\dot{x}_n \mathsf{v}(t) - \dot{x}_n \sum_{n=1}^{N} \dot{\upsilon}_n \psi_n(t) + \dot{\upsilon}_n \mathsf{v}(t) - \dot{\upsilon}_n \sum_{n=1}^{N} \dot{\upsilon}_n \psi_n(t) \right] \\
&= \mathsf{E}\left[\dot{x}_n \mathsf{v}(t)\right] - \mathsf{E}\left[\dot{x}_n \sum_{n=1}^{N} \dot{\upsilon}_n \psi_n(t) \right] + \mathsf{E}\left[\langle \mathsf{v}(t) \mid \psi_n(t) \rangle \mathsf{v}(t) \right] - \mathsf{E}\left[\sum_{m=1}^{N} \dot{\upsilon}_n \dot{\upsilon}_m \psi_m(t) \right] \\
&= \dot{x}_n \cancelto{0}{\mathsf{E}\mathsf{v}(t)} - \dot{x}_n \sum_{n=1}^{N} \cancelto{0}{\mathsf{E}\left[\dot{\upsilon}_n\right]} \psi_n(t) + \mathsf{E}\left[\left\langle \mathsf{v}(t)\mathsf{v}(\hat{\theta}) \mid \psi_n(\hat{\theta}) \right\rangle \right] - \sum_{m=1}^{N} \mathsf{E}\left[\dot{\upsilon}_n \dot{\upsilon}_m\right] \psi_m(t) \\
&= 0 - 0 + \left\langle \mathsf{E}\left[\mathsf{v}(t) n(\hat{\theta})\right] \mid \psi_n(\hat{\theta}) \right\rangle - \sum_{m=1}^{N} N_o \bar{\delta}_{mn} \psi_m(t) \qquad \text{(because } \dot{\upsilon}_n \text{ is white)} \\
&= \left\langle N_o \delta(t - \hat{\theta}) \mid \psi_n(\hat{\theta}) \right\rangle - N_o \psi_n(t) \\
&= N_o \psi_n(t) - N_o \psi_n(t) \\
&= 0
\end{aligned}$$

3. This implies $\dot{y}_n$ and $\mathsf{v}'(t)$ are uncorrelated. Since they are Gaussian processes (due to channel operator hypothesis), they are also independent.

4. Proof that $P\left\{\mathsf{x}(t;\hat{\theta})|\mathsf{y}(t)\right\} = P\left\{\mathsf{x}(t;\hat{\theta})|\dot{y}_1,\ \dot{y}_2, \ldots, \dot{y}_N\right\}$:

$$\mathsf{P}\left\{\mathsf{x}(t;\hat{\theta})|\mathsf{y}(t)\right\} = \mathsf{P}\left\{ \mathsf{x}(t;\hat{\theta})| \sum_{n=1}^{N} \dot{y}_n \psi_n(t) + \mathsf{v}'(t) \right\}$$

BY-NC-ND Copyright © 2019 Daniel J. Greenhoe

$$
\begin{aligned}
&= \mathsf{P}\left\{\mathsf{x}(t;\hat{\theta})|R,\mathsf{v}'(t)\right\} && \text{because } Y \text{ and } \mathsf{v}'(t) \text{ can be extracted by } \langle\cdots\,|\,\psi_n(t)\rangle\\
&= \frac{\mathsf{P}\left\{R,\mathsf{v}'(t)|\mathsf{x}(t;\hat{\theta})\right\}\,P\left\{\mathsf{x}(t;\hat{\theta})\right\}}{\mathsf{P}\{R,\mathsf{v}'(t)\}}\\
&= \frac{\mathsf{P}\left\{R|\mathsf{x}(t;\hat{\theta})\right\}\mathsf{P}\left\{\mathsf{v}'(t)|\mathsf{x}(t;\hat{\theta})\right\}\mathsf{P}\left\{\mathsf{x}(t;\hat{\theta})\right\}}{\mathsf{P}\{Y\}\mathsf{P}\{\mathsf{v}'(t)\}} && \text{by } \textit{independence} \text{ of } Y \text{ and } \mathsf{v}'(t)\\
&= \frac{\mathsf{P}\left\{R|\mathsf{x}(t;\hat{\theta})\right\}\mathsf{P}\left\{\mathsf{v}'(t)\right\}\mathsf{P}\left\{\mathsf{x}(t;\hat{\theta})\right\}}{\mathsf{P}\{Y\}\mathsf{P}\{\mathsf{v}'(t)\}}\\
&= \frac{\mathsf{P}\left\{R|\mathsf{x}(t;\hat{\theta})\right\}\mathsf{P}\left\{\mathsf{x}(t;\hat{\theta})\right\}}{\mathsf{P}\{Y\}}\\
&= \frac{\mathsf{P}\left\{R,\mathsf{x}(t;\hat{\theta})\right\}}{\mathsf{P}\{Y\}}\\
&= \mathsf{P}\left\{\mathsf{x}(t;\hat{\theta})|Y\right\}
\end{aligned}
$$

5. Proof that Y is a sufficient statistic for the *MAP estimate*:

$$
\begin{aligned}
\hat{\theta}_{\mathsf{map}} &\triangleq \arg\max_{\hat{\theta}} \mathsf{P}\left\{\mathsf{x}(t;\hat{\theta})|\mathsf{y}(t)\right\} && \text{by definition of } \textit{MAP estimate}\\
&= \arg\max_{\hat{\theta}} \mathsf{P}\left\{\mathsf{x}(t;\hat{\theta})|R\right\} && \text{by result 4.}
\end{aligned}
$$

6. Proof that Y is a sufficient statistic for the *ML estimate*:

$$
\begin{aligned}
\hat{\theta}_{\mathsf{ml}} &\triangleq \arg\max_{\hat{\theta}} \mathsf{P}\left\{\mathsf{y}(t)|\mathsf{x}(t;\hat{\theta})\right\} && \text{by definition of } \textit{ML estimate}\\
&= \arg\max_{\hat{\theta}} \mathsf{P}\left\{\sum_{n=1}^{N}\dot{y}_n\psi_n(t)+\mathsf{v}'(t)|\mathsf{x}(t;\hat{\theta})\right\}\\
&= \arg\max_{\hat{\theta}} \mathsf{P}\left\{R,\mathsf{v}'(t)|\mathsf{x}(t;\hat{\theta})\right\} && \text{because } Y \text{ and } \mathsf{v}'(t) \text{ can be extracted by } \langle\cdots\,|\,\psi_n(t)\rangle\\
&= \arg\max_{\hat{\theta}} \mathsf{P}\left\{R|\mathsf{x}(t;\hat{\theta})\right\}\mathsf{P}\left\{\mathsf{v}'(t)|\mathsf{x}(t;\hat{\theta})\right\} && \text{by } \textit{independence} \text{ of } Y \text{ and } \mathsf{v}'(t)\\
&= \arg\max_{\hat{\theta}} \mathsf{P}\left\{R|\mathsf{x}(t;\hat{\theta})\right\}\mathsf{P}\left\{\mathsf{v}'(t)\right\} && \text{by } \textit{independence} \text{ of } \mathsf{x}(t) \text{ and } \mathsf{v}'(t)\\
&= \arg\max_{\hat{\theta}} \mathsf{P}\left\{R|\mathsf{x}(t;\hat{\theta})\right\} && \text{by } \textit{independence} \text{ of } \mathsf{v}'(t) \text{ and } \hat{\theta}
\end{aligned}
$$

✏

Theorem 11.2. *Let* $\mathbf{C}=\mathbf{C}_{\mathsf{a}}$ *be an additive noise channel.*

THM

$$\underbrace{\mathbf{C}=\mathbf{C}_{\mathsf{a}}}_{\textit{additive noise channel}} \implies \left\{\ \mathsf{E}(\dot{y}_n|\theta) = \dot{x}_n(\theta)+\mathsf{E}\dot{\upsilon}_n\right.$$

✎PROOF:

$$
\begin{aligned}
\mathsf{E}(\dot{y}_n|\theta) &\triangleq \left(\langle\mathsf{y}(t)\,|\,\psi_n(t)\rangle\,|\theta\right)\\
&= \langle\mathsf{x}(t;\theta)+\mathsf{n}(t)\,|\,\psi_n(t)\rangle\\
&= \langle\mathsf{x}(t;\theta)\,|\,\psi_n(t)\rangle+\langle\mathsf{n}(t)\,|\,\psi_n(t)\rangle\\
&= \left\langle\sum_{k=1}^{N}\dot{x}_k(\theta)\psi_k(t)\,|\,\psi_n(t)\right\rangle+\dot{\upsilon}_n
\end{aligned}
$$

COPYRIGHT © 2019 DANIEL J. GREENHOE | CC BY-NC-ND

$$= \sum_{k=1}^{N} \dot{x}_k(\theta) \langle \psi_k(t) \mid \psi_n(t) \rangle + \dot{\upsilon}_n$$
$$= \dot{x}_n(\theta) + \dot{\upsilon}_n$$

$$\begin{aligned} \mathsf{E}(\dot{y}_n|\theta) &= \mathsf{E}\left[\dot{x}_n(\theta) + \dot{\upsilon}_n\right] \\ &= \mathsf{E}\dot{x}_n(\theta) + \mathsf{E}\dot{\upsilon}_n \\ &= \dot{x}_n(\theta) \end{aligned}$$



Figure 11.1: Additive Gaussian noise channel Statistics

Theorem 11.3. *Let* $\mathbf{C} = \mathbf{C}_{\mathsf{agn}}$ *be an additive gaussian noise channel with distribution* $\mathsf{n}(t) \sim \mathsf{N}\left(0, \sigma^2\right)$ *for all* t.

THM

$$\underbrace{\mathbf{C} = \mathbf{C}_{\mathsf{agn}}}_{\textit{additive Gaussian channel}} \implies \begin{cases} \mathsf{E}\dot{\upsilon}_n &= 0 \\ \mathsf{E}(\dot{y}_n|\theta) &= \dot{x}_n(\theta) \\ \dot{\upsilon}_n &\sim \mathsf{N}\left(0, \sigma^2\right) & \textit{(noise projections are Gaussian)} \\ \dot{y}_n|\theta &\sim \mathsf{N}\left(\dot{x}_n(\theta), \sigma^2\right) & \textit{(receiver projections are Gaussian)} \end{cases}$$

✎PROOF:

$$\begin{aligned} \mathsf{E}\dot{\upsilon}_n &= \mathsf{E}\langle \mathsf{n}(t) \mid \psi_n(t) \rangle \\ &= \langle \mathsf{E}\mathsf{n}(t) \mid \psi_n(t) \rangle \\ &= \langle 0 \mid \psi_n(t) \rangle \\ &= 0 \end{aligned}$$

$$\begin{aligned} (\dot{y}_n|\theta) &\triangleq \langle \mathsf{y}(t) \mid \psi_n(t) \rangle|\theta \\ &= \langle \mathsf{x}(t;\theta) + \mathsf{n}(t) \mid \psi_n(t) \rangle \\ &= \langle \mathsf{x}(t;\theta) \mid \psi_n(t) \rangle + \langle \mathsf{n}(t) \mid \psi_n(t) \rangle \\ &= \left\langle \sum_{k=1}^{N} \dot{x}_k(\theta)\psi_k(t) \mid \psi_n(t) \right\rangle + \dot{\upsilon}_n \\ &= \sum_{k=1}^{N} \dot{x}_k(\theta) \langle \psi_k(t) \mid \psi_n(t) \rangle + \dot{\upsilon}_n \\ &= \dot{x}_n(\theta) + \dot{\upsilon}_n \end{aligned}$$

$$\begin{aligned} \mathsf{E}(\dot{y}_n|\theta) &= \mathsf{E}\left[\dot{x}_n(\theta) + \dot{\upsilon}_n\right] \\ &= \mathsf{E}\dot{x}_n(\theta) + \mathsf{E}\dot{\upsilon}_n \\ &= \dot{x}_n(\theta) \end{aligned}$$

The distributions follow because they are linear operations on Gaussian processes.

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

Theorem 11.4. *Let* $\mathbf{C} = \mathbf{C}_{\text{awn}}$ *be an additive white noise channel.*

THM

$$\underbrace{\mathbf{C} = \mathbf{C}_{\text{awn}}}_{\text{additive white channel}} \implies \begin{cases} \mathsf{E}\dot{\upsilon}_n & = 0 & \text{(noise projection is zero-mean)} \\ \mathsf{E}(\dot{y}_n|\theta) & = \dot{x}_n(\theta) & \text{(expected receiver projection = transmitted projection)} \\ \text{cov}\left[\dot{\upsilon}_n, \dot{\upsilon}_m\right] & = \sigma^2\bar{\delta}_{nm} & \text{(noise projections are uncorrelated)} \\ \text{cov}\left[\dot{y}_n|\theta, \dot{y}_m|\theta\right] & = \sigma^2\bar{\delta}_{nm} & \text{(receiver projections are uncorrelated)} \end{cases}$$

✎PROOF: Because the noise is additive (see Theorem 11.2 (page 81))

$$\begin{aligned} \mathsf{E}\dot{\upsilon}_n &= 0 \\ (\dot{y}_n|\theta) &= \dot{x}_n(\theta) + \dot{\upsilon}_n \\ \mathsf{E}(\dot{y}_n|\theta) &= \dot{x}_n(\theta). \end{aligned}$$

Because the noise is also white,

$$\begin{aligned} \text{cov}\left[\dot{\upsilon}_m, \dot{\upsilon}_n\right] &= \text{cov}\left[\langle \mathsf{n}(t) \,|\, \psi_m(t)\rangle, \langle \mathsf{n}(t) \,|\, \psi_n(t)\rangle\right] \\ &= \mathsf{E}\left[\langle \mathsf{n}(t) \,|\, \psi_m(t)\rangle \langle \mathsf{n}(t) \,|\, \psi_n(t)\rangle\right] \\ &= \mathsf{E}\left[\langle \mathsf{n}(t) \,|\, \psi_m(t)\rangle \left\langle n(\hat{\theta}) \,|\, \psi_n(\hat{\theta})\right\rangle\right] \\ &= \mathsf{E}\left[\left\langle n(\hat{\theta}) \langle \mathsf{n}(t) \,|\, \psi_m(t)\rangle \,|\, \psi_n(\hat{\theta})\right\rangle\right] \\ &= \mathsf{E}\left[\left\langle \left\langle n(\hat{\theta})\mathsf{n}(t) \,|\, \psi_m(t)\right\rangle \,|\, \psi_n(\hat{\theta})\right\rangle\right] \\ &= \left\langle \left\langle \mathsf{E}\left[n(\hat{\theta})\mathsf{n}(t)\right] \,|\, \psi_m(t)\right\rangle \,|\, \psi_n(\hat{\theta})\right\rangle \\ &= \left\langle \left\langle \sigma^2\delta(t-\hat{\theta}) \,|\, \psi_m(t)\right\rangle \,|\, \psi_n(\hat{\theta})\right\rangle \\ &= \sigma^2 \langle \psi_n(t) \,|\, \psi_m(t)\rangle \\ &= \begin{cases} \sigma^2 & \text{for } n = m \\ 0 & \text{for } n \neq m. \end{cases} \end{aligned}$$

$$\begin{aligned} \text{cov}\left[\dot{y}_n|\theta, \dot{y}_m|\theta\right] &= \mathsf{E}\left[\dot{y}_n\dot{y}_m|\theta\right] - [\mathsf{E}\dot{y}_n|\theta][\mathsf{E}\dot{y}_m|\theta] \\ &= \mathsf{E}\left[(\dot{x}_n(\theta) + \dot{\upsilon}_n)(\dot{x}_m(\theta) + \dot{\upsilon}_m)\right] - \dot{x}_n(\theta)\dot{x}_m(\theta) \\ &= \mathsf{E}\left[(\dot{x}_n(\theta) + \dot{\upsilon}_n)(\dot{x}_m(\theta) + \dot{\upsilon}_m)\right] - \dot{x}_n(\theta)\dot{x}_m(\theta) \\ &= \mathsf{E}\left[\dot{x}_n(\theta)\dot{x}_m(\theta) + \dot{x}_n(\theta)\dot{\upsilon}_m + \dot{\upsilon}_n\dot{x}_m(\theta) + \dot{\upsilon}_n\dot{\upsilon}_m\right] - \dot{x}_n(\theta)\dot{x}_m(\theta) \\ &= \dot{x}_n(\theta)\dot{x}_m(\theta) + \dot{x}_n(\theta)\mathsf{E}\left[\dot{\upsilon}_m\right] + \mathsf{E}\left[\dot{\upsilon}_n\right]\dot{x}_m(\theta) + \mathsf{E}\left[\dot{\upsilon}_n\dot{\upsilon}_m\right] - \dot{x}_n(\theta)\dot{x}_m(\theta) \\ &= 0 + \dot{x}_n(\theta)\cdot 0 + 0\cdot\dot{x}_m(\theta) + \text{cov}\left[\dot{\upsilon}_n, \dot{\upsilon}_m\right] + [\mathsf{E}\dot{\upsilon}_n][\mathsf{E}\dot{\upsilon}_m] \\ &= \sigma^2\bar{\delta}_{nm} + 0\cdot 0 \\ &= \begin{cases} \sigma^2 & \text{for } n = m \\ 0 & \text{for } n \neq m. \end{cases} \end{aligned}$$

✏



Figure 11.2: Additive white Gaussian noise channel statistics

Theorem 11.5. *Let* $\mathbf{C} = \mathbf{C}_{\text{awgn}}$ *be an additive gaussian noise channel with distribution* $\mathsf{n}(t) \sim \mathsf{N}\left(0, \sigma^2\right)$ *for all* t.

COPYRIGHT © 2019 DANIEL J. GREENHOE

THM

$$\underbrace{\mathbf{C} = \mathbf{C}_{\mathsf{awgn}}}_{AWGN} \Longrightarrow \begin{cases} \dot{\upsilon}_n \sim \mathsf{N}\left(0, \sigma^2\right) & \textit{(noise projections are Gaussian)} \\ \dot{y}_n|\theta \sim \mathsf{N}\left(\dot{x}_n(\theta), \sigma^2\right) & \textit{(receiver projections are Gaussian)} \\ \operatorname{cov}\left[\dot{\upsilon}_n, \dot{\upsilon}_m\right] = \sigma^2\bar{\delta}_{nm} & \textit{(noise projections are uncorrelated)} \\ \operatorname{cov}\left[\dot{y}_n, \dot{y}_m\right] = \sigma^2\bar{\delta}_{nm} & \textit{(receiver projections are uncorrelated)} \\ \mathsf{P}\{\dot{\upsilon}_n = a \wedge \dot{\upsilon}_m = b\} = \mathsf{P}\{\dot{\upsilon}_n = a\}\mathsf{P}\{\dot{\upsilon}_m = b\} & \textit{(noise projections are independent)} \\ \mathsf{P}\{\dot{y}_n = a \wedge \dot{y}_m = b\} = \mathsf{P}\{\dot{y}_n = a\}\mathsf{P}\{\dot{y}_m = b\} & \textit{(receiver projections are independent)} \end{cases}$$

✎PROOF: The distributions follow because they are linear operations on Gaussian processes.

By Theorem 11.4 (page 83) (for AWN channel)

$$\begin{aligned} \mathsf{E}\dot{\upsilon}_n &= 0 \\ \operatorname{cov}\left[\dot{\upsilon}_m, \dot{\upsilon}_n\right] &= \sigma^2\bar{\delta}_{mn} \\ \dot{y}_n &= \dot{x}_n + \dot{\upsilon}_n \\ \mathsf{E}\dot{y}_n &= \dot{x}_n \\ \operatorname{cov}\left[\dot{y}_n, \dot{y}_m\right] &= \sigma^2\bar{\delta}_{mn} \end{aligned}$$

Because the processes are Gaussian, uncorrelatedness implies *independence*. ✏

11.2 Optimal symbol estimation

The AWGN projection statistics provided by Theorem 11.5 (page 83) help generate the optimal ML-estimates for a number of communication systems. These ML-estimates can be expressed in either of two standard forms:

- **Spectral decompostion**: The optimal estimate is expressed in terms of *projections* of signals onto orthonormal basis functions.
- **Matched signal**: The optimal estimate is expressed in terms of the (noisy) received signal correlated with ("matched" with) the (noiseless) transmitted signal.

Theorem 11.6 (page 84) (next) expresses the general optimal *ML estimate* in both of these forms.

Parameter detection is a special case of parameter estimation. In parameter detection, the estimate is a member of an finite set. In parameter estimation, the estimate is a member of an infinite set (Section 11.2 page 84).

Theorem 11.6 (General ML estimation)**.** *Let Ψ be an orthonormal set spanning $\mathsf{x}(t)$ such that*

$$\begin{aligned} \Psi &\triangleq (\!(\psi_1(t), \psi_2(t), \ldots, \psi_n(t))\!) \\ \dot{y}_n &\triangleq \langle \mathsf{y}(t) \,|\, \psi_n(t) \rangle \\ \dot{x}_n &\triangleq \langle \mathsf{x}(t) \,|\, \psi_n(t) \rangle \\ \mathsf{y}(t) &= \mathsf{x}(t;\hat{\theta}) + \mathsf{n}(t). \end{aligned}$$

Then the optimal ML-estimate $\hat{\theta}_{\mathsf{ml}}$ of parameter θ is

THM

$$\begin{aligned} \hat{\theta}_{\mathsf{ml}} &= \arg\min_{\hat{\theta}} \left[\sum_{n=1}^{N} [\dot{y}_n - \dot{x}_n(\hat{\theta})]^2\right] && \textit{(spectral decomposition)} \\ &= \arg\max_{\hat{\theta}} \left[2\left\langle \mathsf{y}(t) \,|\, \mathsf{x}(t;\hat{\theta})\right\rangle - \left\|\mathsf{x}(t;\hat{\theta})\right\|^2\right] && \textit{(matched signal)} \end{aligned}$$

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

✎PROOF:

$$
\begin{aligned}
\hat{\theta}_{\mathsf{ml}} &= \arg\max_{\hat{\theta}} \mathsf{P}\left\{\mathsf{y}(t)|\mathsf{x}(t;\hat{\theta})\right\} \\
&= \arg\max_{\hat{\theta}} \mathsf{P}\left\{\dot{y}_1, \dot{y}_2, \ldots, \dot{y}_n|\mathsf{x}(t;\hat{\theta})\right\} && \text{by Theorem 11.1 (page 79)} \\
&= \arg\max_{\hat{\theta}} \prod_{n=1}^{N} \mathsf{P}\left\{\dot{y}_n|\mathsf{x}(t;\hat{\theta})\right\} \\
&= \arg\max_{\hat{\theta}} \prod_{n=1}^{N} \mathsf{p}\left[\dot{\mathsf{y}}_{\mathsf{n}}|\mathsf{x}(\mathsf{t};\hat{\theta})\right] \\
&= \arg\max_{\hat{\theta}} \prod_{n=1}^{N} \frac{1}{\sqrt{2\pi\sigma^2}} \exp\frac{[\dot{y}_n - \dot{x}_n(\hat{\theta})]^2}{-2\sigma^2} && \text{by Theorem 11.5 (page 83)} \\
&= \arg\max_{\hat{\theta}} \left(\frac{1}{\sqrt{2\pi\sigma^2}}\right)^N \exp\frac{-1}{2\sigma^2}\sum_{n=1}^{N}[\dot{y}_n - \dot{x}_n(\hat{\theta})]^2 \\
&= \arg\max_{\hat{\theta}} \left[-\sum_{n=1}^{N}[\dot{y}_n - \dot{x}_n(\hat{\theta})]^2\right] \\
&= \arg\max_{\hat{\theta}} \left[-\lim_{N\to\infty}\sum_{n=1}^{N}[\dot{y}_n - \dot{x}_n(\hat{\theta})]^2\right] && \text{by Theorem 11.1 (page 79)} \\
&= \arg\max_{\hat{\theta}} \left[-\left\|\mathsf{y}(t) - \mathsf{x}(t;\hat{\theta})\right\|^2\right] && \text{by } \textit{Plancheral's formula} \quad \text{(Theorem H.9 page 232)} \\
&= \arg\max_{\hat{\theta}} \left[-\|\mathsf{y}(t)\|^2 + 2\mathbf{R}_{\mathsf{e}}\left\langle \mathsf{y}(t) \mid \mathsf{x}(t;\hat{\theta})\right\rangle - \left\|\mathsf{x}(t;\hat{\theta})\right\|^2\right] \\
&= \arg\max_{\hat{\theta}} \left[2\left\langle \mathsf{y}(t) \mid \mathsf{x}(t;\hat{\theta})\right\rangle - \left\|\mathsf{x}(t;\hat{\theta})\right\|^2\right] && \text{because } \mathsf{y}(t) \textit{ independent} \text{ of } \hat{\theta}
\end{aligned}
$$

✏

Theorem 11.7 (ML amplitude estimation)**.** [2] *Let* **S** *be an additive white gaussian noise system such that*

$$
\begin{aligned}
\mathsf{y}(t) &= [\mathbf{C}_{\mathsf{awgn}}s](t) = \mathsf{x}(t;a) + \mathsf{n}(t) \\
\mathsf{x}(t;a) &\triangleq a\lambda(t).
\end{aligned}
$$

Then

THM

$$
\begin{aligned}
\hat{a}_{\mathsf{ml}} &= \frac{1}{\|\lambda(t)\|^2}\left\langle \mathsf{y}(t) \mid \lambda(t)\right\rangle && \textit{(optimal ML-estimate of } a\textit{)} \\
&= \frac{1}{\|\lambda(t)\|^2}\sum_{n=1}^{N}\dot{y}_n\dot{\lambda}_n \\
\mathsf{E}\hat{a}_{\mathsf{ml}} &= a && (\hat{a}_{\mathsf{ml}} \textit{ is } \textbf{\textit{unbiased}}) \\
\operatorname{var}\hat{a}_{\mathsf{ml}} &= \frac{\sigma^2}{\|\lambda(t)\|^2} && \textit{(variance of estimate } \hat{a}_{\mathsf{ml}}\textit{)} \\
\operatorname{var}\hat{a}_{\mathsf{ml}} &= \textit{CR lower bound} && (\hat{a}_{\mathsf{ml}} \textit{ is an } \textbf{efficient} \textbf{\textit{estimate}})
\end{aligned}
$$

✎PROOF:

[2] Mandyam D. Srinath (1996) pages 158–159

COPYRIGHT © 2019 DANIEL J. GREENHOE
CC BY-NC-ND

1. *ML estimate* in "matched signal" form:

$$\begin{aligned}
\hat{a}_{\mathsf{ml}} &= \arg\max_a \left[2\left\langle \mathsf{y}(t) \mid \mathsf{x}(t;\hat{\theta})\right\rangle - \|\mathsf{x}(t;\phi)\|^2\right] && \text{by Theorem 11.6 (page 84)}\\
&= \arg\max_a \left[2\left\langle \mathsf{y}(t) \mid a\lambda(t)\right\rangle - \|a\lambda(t)\|^2\right] && \text{by hypothesis}\\
&= \arg_a \left[\frac{\partial}{\partial a}2a\left\langle \mathsf{y}(t) \mid \lambda(t)\right\rangle - \frac{\partial}{\partial a}a^2\|\lambda(t)\|^2 = 0\right]\\
&= \arg_a \left[2\left\langle \mathsf{y}(t) \mid \lambda(t)\right\rangle - 2a\|\lambda(t)\|^2 = 0\right]\\
&= \arg_a \left[\left\langle \mathsf{y}(t) \mid \lambda(t)\right\rangle = a\|\lambda(t)\|^2\right]\\
&= \frac{1}{\|\lambda(t)\|^2}\left\langle \mathsf{y}(t) \mid \lambda(t)\right\rangle
\end{aligned}$$

2. *ML estimate* in "spectral decomposition" form:

$$\begin{aligned}
\hat{a}_{\mathsf{ml}} &= \arg\min_a \left(\sum_{n=1}^{N}\left[\dot{y}_n - \dot{x}_n(a)\right]^2\right) && \text{by Theorem 11.6 (page 84)}\\
&= \arg_a \left(\frac{\partial}{\partial a}\sum_{n=1}^{N}\left[\dot{y}_n - \dot{x}_n(a)\right]^2 = 0\right)\\
&= \arg_a \left(2\sum_{n=1}^{N}\left[\dot{y}_n - \dot{x}_n(a)\right]\frac{\partial}{\partial a}\dot{x}_n(a) = 0\right)\\
&= \arg_a \left(\sum_{n=1}^{N}\left[\dot{y}_n - \left\langle a\lambda(t) \mid \psi_n(t)\right\rangle\right]\frac{\partial}{\partial a}\left\langle a\lambda(t) \mid \psi_n(t)\right\rangle = 0\right)\\
&= \arg_a \left(\sum_{n=1}^{N}\left[\dot{y}_n - a\left\langle \lambda(t) \mid \psi_n(t)\right\rangle\right]\frac{\partial}{\partial a}\left(a\left\langle \lambda(t) \mid \psi_n(t)\right\rangle\right) = 0\right)\\
&= \arg_a \left(\sum_{n=1}^{N}\left[\dot{y}_n - a\dot{\lambda}_n\right]\left\langle \lambda(t) \mid \psi_n(t)\right\rangle = 0\right)\\
&= \arg_a \left(\sum_{n=1}^{N}\left[\dot{y}_n - a\dot{\lambda}_n\right]\dot{\lambda}_n = 0\right)\\
&= \arg_a \left(\sum_{n=1}^{N}\dot{y}_n\dot{\lambda}_n = \sum_{n=1}^{N}a\dot{\lambda}_n^2\right)\\
&= \left(\frac{1}{\sum_{n=1}^{N}\dot{\lambda}_n^2}\right)\sum_{n=1}^{N}\dot{y}_n\dot{\lambda}_n\\
&= \frac{1}{\|\lambda(t)\|^2}\sum_{n=1}^{N}\dot{y}_n\dot{\lambda}_n
\end{aligned}$$

3. Prove that the estimate $\hat{a}_{\mathsf{ml}}$ is **unbiased**:

$$\begin{aligned}
\mathsf{E}\hat{a}_{\mathsf{ml}} &= \mathsf{E}\frac{1}{\|\lambda(t)\|^2}\int_{t\in\mathbb{R}}\mathsf{y}(t)\lambda(t)\,\mathrm{d}t && \text{by previous result}\\
&= \mathsf{E}\frac{1}{\|\lambda(t)\|^2}\int_{t\in\mathbb{R}}[a\lambda(t)+\mathsf{n}(t)]\lambda(t)\,\mathrm{d}t && \text{by hypothesis}\\
&= \frac{1}{\|\lambda(t)\|^2}\int_{t\in\mathbb{R}}\mathsf{E}[a\lambda(t)+\mathsf{n}(t)]\lambda(t)\,\mathrm{d}t && \text{by linearity of } \int\cdot\,\mathrm{d}t \text{ and } \mathsf{E}
\end{aligned}$$

BY-NC-ND Copyright © 2019 Daniel J. Greenhoe

$$
\begin{aligned}
&= \frac{1}{\|\lambda(t)\|^2} a \int_{t\in\mathbb{R}} \lambda^2(t)\,\mathrm{d}t && \text{by E operation}\\
&= \frac{1}{\|\lambda(t)\|^2} a\,\|\lambda(t)\|^2 && \text{by definition of } \|\cdot\|^2\\
&= a
\end{aligned}
$$

4. Compute the variance of $\hat{a}_{\mathsf{ml}}$:

$$
\begin{aligned}
\mathsf{E}\hat{a}_{\mathsf{ml}}^2 &= \mathsf{E}\left[\frac{1}{\|\lambda(t)\|^2}\int_{t\in\mathbb{R}} \mathsf{y}(t)\lambda(t)\,\mathrm{d}t\right]^2\\
&= \mathsf{E}\left[\frac{1}{\|\lambda(t)\|^4}\int_{t\in\mathbb{R}} \mathsf{y}(t)\lambda(t)\,\mathrm{d}t \int_{v} \mathsf{y}(v)\lambda(v)\,\mathrm{d}v\right]\\
&= \mathsf{E}\left[\frac{1}{\|\lambda(t)\|^4}\int_{t\in\mathbb{R}}\int_{v} [a\lambda(t)+\mathsf{n}(t)][a\lambda(v)+\mathsf{n}(v)]\lambda(t)\lambda(v)\,\mathrm{d}v\,\mathrm{d}t\right]\\
&= \mathsf{E}\left[\frac{1}{\|\lambda(t)\|^4}\int_{t\in\mathbb{R}}\int_{v} [a^2\lambda(t)\lambda(v)+a\lambda(t)\mathsf{n}(v)+a\lambda(v)\mathsf{n}(t)+\mathsf{n}(t)\mathsf{n}(v)]\lambda(t)\lambda(v)\,\mathrm{d}v\,\mathrm{d}t\right]\\
&= \left[\frac{1}{\|\lambda(t)\|^4}\int_{t\in\mathbb{R}}\int_{v} [a^2\lambda(t)\lambda(v)+0+0+\sigma^2\delta(t-v)]\lambda(t)\lambda(v)\,\mathrm{d}v\,\mathrm{d}t\right]\\
&= \frac{1}{\|\lambda(t)\|^4}\int_{t\in\mathbb{R}}\int_{v} a^2\lambda^2(t)\lambda^2(v)\,\mathrm{d}v\,\mathrm{d}t + \frac{1}{\|\lambda(t)\|^4}\int_{t\in\mathbb{R}}\int_{v} \sigma^2\delta(t-v)\lambda(t)\lambda(v)\,\mathrm{d}v\,\mathrm{d}t\\
&= \frac{1}{\|\lambda(t)\|^4} a^2 \int_{t\in\mathbb{R}} \lambda^2(t)\,\mathrm{d}t \int_{v} \lambda^2(v)\,\mathrm{d}v + \frac{1}{\|\lambda(t)\|^4}\sigma^2 \int_{t\in\mathbb{R}} \lambda^2(t)\,\mathrm{d}t\\
&= a^2 \frac{1}{\|\lambda(t)\|^4}\,\|\lambda(t)\|^2\,\|\lambda(v)\|^2 + \frac{1}{\|\lambda(t)\|^4}\sigma^2\,\|\lambda(t)\|^2\\
&= a^2 + \frac{\sigma^2}{\|\lambda(t)\|^2}
\end{aligned}
$$

$$
\begin{aligned}
\operatorname{var}\hat{a}_{\mathsf{ml}} &= \mathsf{E}\hat{a}_{\mathsf{ml}}^2 - (\mathsf{E}\hat{a}_{\mathsf{ml}})^2\\
&= \left(a^2 + \frac{\sigma^2}{\|\lambda(t)\|^2}\right) - \left(a^2\right)\\
&= \frac{\sigma^2}{\|\lambda(t)\|^2}
\end{aligned}
$$

5. Compute the Cramér-Rao Bound:

$$
\begin{aligned}
\mathsf{p}\left[\mathsf{y}(t)|\mathsf{s}(t;\mathsf{a})\right] &= \mathsf{p}\left[\dot{\mathsf{y}}_1, \dot{\mathsf{y}}_2, \ldots, \dot{\mathsf{y}}_N|\mathsf{s}(t;\mathsf{a})\right]\\
&= \prod_{n=1}^{N} \frac{1}{\sqrt{2\pi\sigma^2}} \exp\frac{(\dot{y}_n - a\dot{\lambda}_n)^2}{-2\sigma^2}\\
&= \left(\frac{1}{\sqrt{2\pi\sigma^2}}\right)^N \exp\frac{1}{-2\sigma^2}\sum_{n=1}^{N}(\dot{y}_n - a\dot{\lambda}_n)^2
\end{aligned}
$$

$$
\begin{aligned}
\frac{\partial}{\partial a}\ln \mathsf{p}\left[\mathsf{y}(t)|\mathsf{s}(t;\mathsf{a})\right] &= \frac{\partial}{\partial a}\ln\left(\frac{1}{\sqrt{2\pi\sigma^2}}\right)^N \exp\frac{1}{-2\sigma^2}\sum_{n=1}^{N}(\dot{y}_n - a\dot{\lambda}_n)^2\\
&= \frac{\partial}{\partial a}\ln\left(\frac{1}{\sqrt{2\pi\sigma^2}}\right)^N + \frac{\partial}{\partial a}\ln\exp\frac{1}{-2\sigma^2}\sum_{n=1}^{N}(\dot{y}_n - a\dot{\lambda}_n)^2\\
&= \frac{\partial}{\partial a}\frac{1}{-2\sigma^2}\sum_{n=1}^{N}(\dot{y}_n - a\dot{\lambda}_n)^2
\end{aligned}
$$

Copyright © 2019 Daniel J. Greenhoe

$$= \frac{1}{-2\sigma^2}\sum_{n=1}^{N} 2(\dot{y}_n - a\dot{\lambda}_n)(-\dot{\lambda}_n)$$

$$= \frac{1}{\sigma^2}\sum_{n=1}^{N} \dot{\lambda}_n(\dot{y}_n - a\dot{\lambda}_n)$$

$$\begin{aligned}
\frac{\partial^2}{\partial a^2}\ln \mathsf{p}\left[\mathsf{y(t)|s(t;a)}\right] &= \frac{\partial}{\partial a}\frac{\partial}{\partial a}\ln \mathsf{p}\left[\mathsf{y(t)|s(t;a)}\right] \\
&= \frac{\partial}{\partial a}\frac{1}{\sigma^2}\sum_{n=1}^{N} \dot{\lambda}_n(\dot{y}_n - a\dot{\lambda}_n) \\
&= \frac{1}{\sigma^2}\sum_{n=1}^{N} \dot{\lambda}_n(-\dot{\lambda}_n) \\
&= \frac{-1}{\sigma^2}\sum_{n=1}^{N} \dot{\lambda}_n^2 \\
&= \frac{-\|\lambda(t)\|^2}{\sigma^2}
\end{aligned}$$

$$\begin{aligned}
\operatorname{var}\hat{a}_{\mathsf{ml}} &\triangleq \mathsf{E}\left[\hat{a}_{\mathsf{ml}} - \mathsf{E}\hat{a}_{\mathsf{ml}}\right]^2 \\
&= \mathsf{E}\left[\hat{a}_{\mathsf{ml}} - a\right]^2 \\
&\geq \frac{-1}{\mathsf{E}\left(\frac{\partial^2}{\partial a^2}\ln \mathsf{p}\left[\mathsf{y(t)|s(t;a)}\right]\right)} \\
&= \frac{-1}{\mathsf{E}\left(\frac{-\|\lambda(t)\|^2}{\sigma^2}\right)} \\
&= \frac{\sigma^2}{\|\lambda(t)\|^2} \qquad \text{(Cramér-Rao lower bound of the variance)}
\end{aligned}$$

6. Prove that $\hat{a}_{\mathsf{ml}}$ is an ***efficient* estimate**:

   A estimate is *efficient* if $\operatorname{var}\hat{a}_{\mathsf{ml}} = \text{CR lower bound}$. We have already proven this, so $\hat{a}_{\mathsf{ml}}$ is an *efficient* estimate.

   Also, even without explicitly computing the variance of $\hat{a}_{\mathsf{ml}}$, the variance equals the *Cramér-Rao lower bound* (and hence $\hat{a}_{\mathsf{ml}}$ is an *efficient* estimate) if and only if

$$\hat{a}_{\mathsf{ml}} - a = \left(\frac{-1}{\mathsf{E}\left[\frac{\partial^2}{\partial a^2}\ln \mathsf{p}\left[\mathsf{y(t)|s(t;a)}\right]\right]}\right)\left(\frac{\partial}{\partial a}\ln \mathsf{p}\left[\mathsf{y(t)|s(t;a)}\right]\right)$$

$$\begin{aligned}
\left(\frac{-1}{\mathsf{E}\left(\frac{\partial^2}{\partial a^2}\ln \mathsf{p}\left[\mathsf{y(t)|s(t;a)}\right]\right)}\right)\left(\frac{\partial}{\partial a}\ln \mathsf{p}\left[\mathsf{y(t)|s(t;a)}\right]\right) &= \left(\frac{\sigma^2}{\|\lambda(t)\|^2}\right)\left(\frac{1}{\sigma^2}\sum_{n=1}^{N}\dot{\lambda}(\dot{y} - a\dot{\lambda})\right) \\
&= \frac{1}{\|\lambda(t)\|^2}\sum_{n=1}^{N}\dot{\lambda}\dot{y} - \frac{1}{\|\lambda(t)\|^2}\sum_{n=1}^{N}\dot{\lambda}^2 \\
&= \hat{a}_{\mathsf{ml}} - a
\end{aligned}$$

✎

Theorem 11.8 (ML phase estimation)**.** [3] *Let* **S** *be an additive white gaussian noise system such that*

[3] Mandyam D. Srinath (1996) pages 159–160

BY-NC-ND Copyright © 2019 Daniel J. Greenhoe

$$\begin{aligned} \mathsf{y}(t) &= [\mathbf{C}_{\mathsf{awgn}}s](t) = \mathsf{x}(t;\phi) + \mathsf{n}(t) \\ \mathsf{x}(t;\phi) &= A\cos(2\pi f_c t + \phi). \end{aligned}$$

Then the optimal ML-estimate of parameter ϕ is

THM

$$\hat{\phi}_{\mathsf{ml}} = -\operatorname{atan}\left(\frac{\langle \mathsf{y}(t) \mid \sin(2\pi f_c t)\rangle}{\langle \mathsf{y}(t) \mid \cos(2\pi f_c t)\rangle}\right)$$

✎PROOF:

$$\begin{aligned}
\hat{\phi}_{\mathsf{ml}} &= \arg\max_{\phi}\left[2\left\langle \mathsf{y}(t) \mid \mathsf{x}(t;\hat{\theta})\right\rangle - \|\mathsf{x}(t;\phi)\|^2\right] && \text{by Theorem 11.6 (page 84)} \\
&= \arg\max_{\phi}\left[2\left\langle \mathsf{y}(t) \mid \mathsf{x}(t;\phi)\right\rangle\right] && \text{because } \|\mathsf{x}(t;\phi)\| \text{ does not depend on } \phi \\
&= \arg_{\phi}\left[\frac{\partial}{\partial\phi}\left\langle \mathsf{y}(t) \mid \mathsf{x}(t;\phi)\right\rangle = 0\right] \\
&= \arg_{\phi}\left[\left\langle \mathsf{y}(t) \mid \frac{\partial}{\partial\phi}\mathsf{x}(t;\phi)\right\rangle = 0\right] && \text{because } \langle\cdot\mid\cdot\rangle \text{ is a linear operator} \\
&= \arg_{\phi}\left[\left\langle \mathsf{y}(t) \mid \frac{\partial}{\partial\phi}A\cos(2\pi f_c t + \phi)\right\rangle = 0\right] \\
&= \arg_{\phi}\left[\left\langle \mathsf{y}(t) \mid -A\sin(2\pi f_c t + \phi)\right\rangle = 0\right] \\
&= \arg_{\phi}\left[-A\left\langle \mathsf{y}(t) \mid \cos(2\pi f_c t)\sin\phi + \sin(2\pi f_c t)\cos\phi\right\rangle = 0\right] \\
&= \arg_{\phi}\left[\sin\phi\left\langle \mathsf{y}(t) \mid \cos(2\pi f_c t)\right\rangle = -\cos\phi\left\langle \mathsf{y}(t) \mid \sin(2\pi f_c t)\right\rangle\right] \\
&= \arg_{\phi}\left[\frac{\sin\phi}{\cos\phi} = -\frac{\langle \mathsf{y}(t) \mid \sin(2\pi f_c t)\rangle}{\langle \mathsf{y}(t) \mid \cos(2\pi f_c t)\rangle}\right] \\
&= \arg_{\phi}\left[\tan\phi = -\frac{\langle \mathsf{y}(t) \mid \sin(2\pi f_c t)\rangle}{\langle \mathsf{y}(t) \mid \cos(2\pi f_c t)\rangle}\right] \\
&= -\operatorname{atan}\left(\frac{\langle \mathsf{y}(t) \mid \sin(2\pi f_c t)\rangle}{\langle \mathsf{y}(t) \mid \cos(2\pi f_c t)\rangle}\right)
\end{aligned}$$

✏

Theorem 11.9 (ML estimation of a function of a parameter). [4] *Let* $\mathbf{S}$ *be an additive white gaussian noise system such that*

$$\begin{aligned} \mathsf{y}(t) &= [\mathbf{C}_{\mathsf{awgn}}s](t) = \mathsf{x}(t;\hat{\theta}) + \mathsf{n}(t) \\ \mathsf{x}(t;\hat{\theta}) &= \mathsf{g}(\hat{\theta}) \end{aligned}$$

where g *is one-to-one and onto (invertible).*

THM

Then the optimal ML-estimate of parameter u *is*
$$\hat{\theta}_{\mathsf{ml}} = \mathsf{g}^{-1}\left(\frac{1}{N}\sum_{n=1}^{N}\dot{y}_n\right).$$

If an ML ESTIMATE $\hat{\theta}_{\mathsf{ml}}$ *is unbiased* ($\mathsf{E}\hat{\theta}_{\mathsf{ml}} = \theta$) *then*
$$\operatorname{var}\hat{\theta}_{\mathsf{ml}} \geq \frac{\sigma^2}{N}\frac{1}{\left[\frac{\partial \mathsf{g}(\theta)}{\partial\theta}\right]^2}.$$

If $\mathsf{g}(\theta) = \theta$ *then* $\hat{\theta}_{\mathsf{ml}}$ *is an* ***efficient*** *estimate such that*
$$\operatorname{var}\hat{\theta}_{\mathsf{ml}} = \frac{\sigma^2}{N}.$$

✎PROOF:

$$\hat{\theta}_{\mathsf{ml}} = \arg\min_{\hat{\theta}}\left[\sum_{n=1}^{N}[\dot{y}_n - \mathsf{g}(\hat{\theta})]^2\right]$$

[4] Mandyam D. Srinath (1996) pages 142–143

COPYRIGHT © 2019 DANIEL J. GREENHOE
BY-NC-ND

$$
\begin{aligned}
&= \arg_{\hat\theta}\left[\frac{\partial}{\partial\hat\theta}\sum_{n=1}^{N}[\dot y_n - g(\hat\theta)]^2 = 0\right]\\
&= \arg_{\hat\theta}\left[2\sum_{n=1}^{N}[\dot y_n - g(\hat\theta)]\frac{\partial}{\partial\hat\theta}g(\hat\theta) = 0\right]\\
&= \arg_{\hat\theta}\left[2\sum_{n=1}^{N}[\dot y_n - g(\hat\theta)] = 0\right]\\
&= \arg_{\hat\theta}\left[\sum_{n=1}^{N}\dot y_n = Ng(\hat\theta)\right]\\
&= \arg_{\hat\theta}\left[g(\hat\theta) = \frac{1}{N}\sum_{n=1}^{N}\dot y_n\right]\\
&= \arg_{\hat\theta}\left[u = g^{-1}\left(\frac{1}{N}\sum_{n=1}^{N}\dot y_n\right)\right]\\
&= g^{-1}\left(\frac{1}{N}\sum_{n=1}^{N}\dot y_n\right)
\end{aligned}
$$

If $\hat\theta_{\mathsf{ml}}$ is unbiased ($\mathsf{E}\hat\theta_{\mathsf{ml}} = \theta$), we can use the *Cramér-Rao bound* to find a lower bound on the variance:

$$
\begin{aligned}
\operatorname{var}\hat\theta_{\mathsf{ml}} &\triangleq \mathsf{E}\left[\hat\theta_{\mathsf{ml}} - \mathsf{E}\hat\theta_{\mathsf{ml}}\right]^2\\
&= \mathsf{E}\left[\hat\theta_{\mathsf{ml}} - \theta\right]^2\\
&\geq \frac{-1}{\mathsf{E}\left(\frac{\partial^2}{\partial\theta^2}\ln p\left[y(t)|s(t;\theta)\right]\right)}\\
&= \frac{-1}{\mathsf{E}\left(\frac{\partial^2}{\partial\theta^2}\ln p\left[\dot y_1, \dot y_2, \ldots, \dot y_N|s(t;\theta)\right]\right)}\\
&= \frac{-1}{\mathsf{E}\left(\frac{\partial^2}{\partial\theta^2}\ln\left[\left(\frac{1}{\sqrt{2\pi\sigma^2}}\right)^N \exp\frac{-1}{2\sigma^2}\sum_{n=1}^{N}[\dot y_n - g(\theta)]^2\right]\right)}\\
&= \frac{-1}{\mathsf{E}\left(\frac{\partial^2}{\partial\theta^2}\ln\left[\left(\frac{1}{\sqrt{2\pi\sigma^2}}\right)^N\right] + \frac{\partial^2}{\partial\theta^2}\ln\left[\exp\frac{-1}{2\sigma^2}\sum_{n=1}^{N}[\dot y_n - g(\theta)]^2\right]\right)}\\
&= \frac{-1}{\mathsf{E}\left(\frac{\partial^2}{\partial\theta^2}\left(\frac{-1}{2\sigma^2}\sum_{n=1}^{N}[\dot y_n - g(\theta)]^2\right)\right)}\\
&= \frac{2\sigma^2}{\mathsf{E}\left(\frac{\partial}{\partial\theta}\frac{\partial}{\partial\theta}\sum_{n=1}^{N}[\dot y_n - g(\theta)]^2\right)}\\
&= \frac{2\sigma^2}{\mathsf{E}\left(-2\frac{\partial}{\partial\theta}\frac{\partial g(\theta)}{\partial\theta}\sum_{n=1}^{N}[\dot y_n - g(\theta)]\right)}\\
&= \frac{-\sigma^2}{\mathsf{E}\left(\frac{\partial g^2(\theta)}{\partial\theta^2}\sum_{n=1}^{N}[\dot y_n - g(\theta)] + \frac{\partial g(\theta)}{\partial\theta}\frac{\partial}{\partial\theta}\sum_{n=1}^{N}[\dot y_n - g(\theta)]\right)}
\end{aligned}
$$

BY-NC-ND Copyright © 2019 Daniel J. Greenhoe

$$
\begin{aligned}
&= \frac{-\sigma^2}{\mathsf{E}\left(\frac{\partial \mathsf{g}^2(\theta)}{\partial\theta^2}\sum_{n=1}^{N}[\dot{y}_n - \mathsf{g}(\theta)] - N\frac{\partial \mathsf{g}(\theta)}{\partial\theta}\frac{\partial \mathsf{g}(\theta)}{\partial\theta}\right)}\\
&= \frac{-\sigma^2}{\frac{\partial \mathsf{g}^2(\theta)}{\partial\theta^2}\sum_{n=1}^{N}\mathsf{E}[\dot{y}_n - \mathsf{g}(\theta)] - N\frac{\partial \mathsf{g}(\theta)}{\partial\theta}\frac{\partial \mathsf{g}(\theta)}{\partial\theta}}\\
&= \frac{-\sigma^2}{-N\frac{\partial \mathsf{g}(\theta)}{\partial\theta}\frac{\partial \mathsf{g}(\theta)}{\partial\theta}}\\
&= \frac{\sigma^2}{N}\frac{1}{\left[\frac{\partial \mathsf{g}(\theta)}{\partial\theta}\right]^2}
\end{aligned}
$$

The inequality becomes equality (an *efficient* estimate) if and only if

$$
\hat{\theta}_{\mathsf{ml}} - \theta = \left(\frac{-1}{\mathsf{E}\left(\frac{\partial^2}{\partial\theta^2}\ln \mathsf{p}\left[\mathsf{y(t)}|\mathsf{s(t;}\theta)\right]\right)}\right)\left(\frac{\partial}{\partial\theta}\ln \mathsf{p}\left[\mathsf{y(t)}|\mathsf{s(t;}\theta)\right]\right).
$$

$$
\begin{aligned}
\left(\frac{-1}{\mathsf{E}\left(\frac{\partial^2}{\partial\theta^2}\ln \mathsf{p}\left[\mathsf{y(t)}|\mathsf{s(t;}\theta)\right]\right)}\right)\left(\frac{\partial}{\partial\theta}\ln \mathsf{p}\left[\mathsf{y(t)}|\mathsf{s(t;}\theta)\right]\right) &= \left(\frac{\sigma^2}{N}\frac{1}{\left[\frac{\partial \mathsf{g}(\theta)}{\partial\theta}\right]^2}\right)\left(\frac{-1}{2\sigma^2}(2)\frac{\partial \mathsf{g}(\theta)}{\partial\theta}\sum_{n=1}^{N}[\dot{y}_n - \mathsf{g}(\theta)]\right)\\
&= -\frac{1}{N}\frac{1}{\frac{\partial \mathsf{g}(\theta)}{\partial\theta}}\left(\sum_{n=1}^{N}[\dot{y}_n - \mathsf{g}(\theta)]\right)\\
&= -\frac{1}{\frac{\partial \mathsf{g}(\theta)}{\partial\theta}}\left(\frac{1}{N}\sum_{n=1}^{N}\dot{y}_n - \mathsf{g}(\theta)\right)\\
&= -\frac{1}{\frac{\partial \mathsf{g}(\theta)}{\partial\theta}}\left(\hat{\theta}_{\mathsf{ml}} - \mathsf{g}(\theta)\right)\\
&= -(\hat{\theta}_{\mathsf{ml}} - \theta)
\end{aligned}
$$

✏

11.3 Colored noise

This chapter presented several theorems whose results depended on the noise being white. However if the noise is **colored**, then these results are invalid. But there is still hope for colored noise. Processing colored signals can be accomplished using two techniques:

1. Karhunen-Loève basis functions [5]
2. whitening filter [6]

Karhunen-Loève. If the noise is white, the set $\{\langle \mathsf{y}(t) \,|\, \psi_n(t)\rangle\}$ is a sufficient statistic regardless of which set $\{\psi_n(t)\}$ of orthonormal basis functions are used. If the noise is colored, and if $\{\psi_n(t)\}$

[5] **Karhunen-Loève**: Section 3.2 (page 18)
[6] *Continuous data whitening*: Section 6.3 page 48
Discrete data whitening: Section 5.4 page 38

Copyright © 2019 Daniel J. Greenhoe

satisfy the Karhunen-Loève criterion

$$\int_{t_2} \mathsf{R}_{\mathsf{xx}}(t_1, t_2)\psi_n(t_2)\, \mathrm{d}t_2 = \lambda_n \psi_n(t_1)$$

then $\{\langle \mathsf{y}(t) \mid \psi_n(t)\rangle\}$ is still a sufficient statistic.

Whitening filter. The whitening filter makes the received signal $\mathsf{y}(t)$ statistically white (uncorrelated in time). In this case, any orthonormal basis set can be used to generate sufficient statistics.

11.4 Signal matching

Detection methods. There are basically two types of detection methods:

1. signal matching
2. orthonormal decomposition.

Let S be the set of transmitted waveforms and Y be a set of orthonormal basis functions that span S. *Signal matching* computes the innerproducts of a received signal $\mathsf{y}(t)$ with each signal from S. *Orthonormal decomposition* computes the innerproducts of $\mathsf{y}(t)$ with each signal from the set Y.

In the case where $|S|$ is large, often $|R| << |S|$ making orthonormal decomposition much easier to implement. For example, in a QAM-64 modulation system, signal matching requires $|S| = 64$ innerproduct calculations, while orthonormal decomposition only requires $|R| = 2$ innerproduct calculations because all 64 signals in S can be spanned by just 2 orthonormal basis functions.

Maximizing SNR. Theorem 11.1 (page 79) shows that the innerproducts of $\mathsf{y}(t)$ with basis functions of Y is sufficient for optimal detection. Theorem 11.10 (page 92) (next) shows that a receiver can maximize the SNR of a received signal when signal matching is used.

Theorem 11.10. *Let* $\mathsf{x}(t)$ *be a transmitted signal,* $\mathsf{n}(t)$ *noise, and* $\mathsf{y}(t)$ *the received signal in an AWGN channel. Let the* SIGNAL TO NOISE RATIO *SNR be defined as*

$$\mathrm{SNR}[\mathsf{y}(t)] \triangleq \frac{|\langle \mathsf{x}(t) \mid \mathsf{x}(t)\rangle|^2}{\mathsf{E}\left[|\langle \mathsf{n}(t) \mid \mathsf{x}(t)\rangle|^2\right]}.$$

THM $\mathrm{SNR}[\mathsf{y}(t)] \leq \dfrac{2\,\|\mathsf{x}(t)\|^2}{N_o}$ *and is maximized (equality) when* $\mathsf{x}(t) = a\mathsf{x}(t)$, *where* $a \in \mathbb{R}$.

✎PROOF:

$$\begin{aligned}
\mathrm{SNR}[\mathsf{y}(t)] &\triangleq \frac{|\langle \mathsf{x}(t) \mid \mathsf{x}(t)\rangle|^2}{\mathsf{E}\left[|\langle \mathsf{n}(t) \mid \mathsf{x}(t)\rangle|^2\right]} \\
&= \frac{|\langle \mathsf{x}(t) \mid f(t)\rangle|^2}{\mathsf{E}\left[\left[\int_{t\in\mathbb{R}} \mathsf{n}(t)\mathsf{x}^*(t)\, dt\right]\left[\int_{\hat\theta} n(\hat\theta) f^*(\hat\theta)\, du\right]^*\right]} \\
&= \frac{|\langle \mathsf{x}(t) \mid \mathsf{x}(t)\rangle|^2}{\mathsf{E}\left[\int_{t\in\mathbb{R}}\int_{\hat\theta} \mathsf{n}(t) n^*(\hat\theta)\mathsf{x}^*(t)\mathsf{x}(\hat\theta)\, dt du\right]}
\end{aligned}$$

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

$$
\begin{aligned}
&= \frac{|\langle \mathsf{x}(t) \mid f(t)\rangle|^2}{\int_{t\in\mathbb{R}}\int_{\hat\theta} \mathsf{E}\left[\mathsf{n}(t)n^*(\hat\theta)\right]\mathsf{x}^*(t)\mathsf{x}(\hat\theta)\,dtdu}\\
&= \frac{|\langle \mathsf{x}(t) \mid \mathsf{x}(t)\rangle|^2}{\int_{t\in\mathbb{R}}\int_{\hat\theta} \frac{1}{2}N_o\delta(t-\hat\theta)\mathsf{x}^*(t)\mathsf{x}(\hat\theta)\,dtdu}\\
&= \frac{|\langle \mathsf{x}(t) \mid \mathsf{x}(t)\rangle|^2}{\frac{1}{2}N_o\int_{t\in\mathbb{R}} \mathsf{x}^*(t)\mathsf{x}(t)\,\mathrm{d}t}\\
&= \frac{|\langle \mathsf{x}(t) \mid \mathsf{x}(t)\rangle|^2}{\frac{1}{2}N_o\,\|\mathsf{x}(t)\|^2}\\
&\le \frac{|\,\|\mathsf{x}(t)\|\;\|\mathsf{x}(t)\|\,|^2}{\frac{1}{2}N_o\,\|\mathsf{x}(t)\|^2} && \text{by } \textit{Cauchy-Schwarz Inequality} \quad \text{(Theorem J.2 page 256)}\\
&= \frac{2\,\|\mathsf{x}(t)\|^2}{N_o}
\end{aligned}
$$

The Cauchy-Schwarz Inequality becomes an equality (SNR is maximized) when $\mathsf{x}(t) = a\mathsf{x}(t)$. ✏

Implementation. The innerproduct operations can be implemented using either

1. a correlator or
2. a matched filter.

A correlator is simply an integrator of the form $\langle \mathsf{y}(t) \mid f(t)\rangle = \int_0^T \mathsf{y}(t)f(t)\,\mathrm{d}t$.

A matched filter introduces a function $\mathsf{h}(t)$ such that $\mathsf{h}(t) = \mathsf{x}(T-t)$ (which implies $\mathsf{x}(t) = h(T-t)$) giving

$$
\underbrace{\langle \mathsf{y}(t) \mid \mathsf{x}(t)\rangle = \int_0^T \mathsf{y}(t)\mathsf{x}(t)\,\mathrm{d}t}_{\text{correlator}} = \underbrace{\int_0^\infty \mathsf{x}(\tau)h(t-\tau)\,\mathrm{d}\tau\Big|_{t=T} = \mathsf{x}(t)\star\mathsf{h}(t)|_{t=T}}_{\text{matched filter}}.
$$

This shows that $\mathsf{h}(t)$ is the impulse response of a filter operation sampled at time T. By Theorem 11.10 (page 92), the optimal impulse response is $\mathsf{h}(T-t) = f(t) = \mathsf{x}(t)$. That is, the optimal $\mathsf{h}(t)$ is just a "flipped" and shifted version of $\mathsf{x}(t)$.

Copyright © 2019 Daniel J. Greenhoe

Copyright © 2019 Daniel J. Greenhoe

CHAPTER 12

MOMENT ESTIMATION

12.1 Mean Estimation

Theorem 12.1. *Let* $\hat{\mu} \triangleq \sum_{n=1}^{N} \lambda_n \mathsf{x}_n$ *with* $\sum_{n=1}^{N} \lambda_n = 1$ *be the* ARITHMETIC MEAN *(Definition M.4 page 295)*.

THM

$$\left\{\begin{array}{lll} (A). & (\!(\mathsf{x}_n)\!) \text{ is WIDE SENSE STATIONARY} & \text{and} \\ (B). & \mu \triangleq \mathsf{E}\mathsf{x}_n & \text{and} \\ (C). & (\!(\mathsf{x}_n)\!) \text{ is UNCORRELATED} & \text{and} \\ (D). & \hat{\mu} \triangleq \sum_{n=1}^{N} \lambda_n \mathsf{x}_n \quad \text{(ARITHMETIC MEAN)} & \end{array}\right\} \Longrightarrow \left\{\begin{array}{lll} (1). & \mathsf{E}\hat{\mu} = \mu \quad \text{(UNBIASED)} & \text{and} \\ (2). & \mathrm{var}(\hat{\mu}) = \sigma^2 \sum_{n=1}^{N} \lambda_n^2 & \text{and} \\ (3). & \mathrm{mse}(\hat{\mu}) = \sigma^2 \sum_{n=1}^{N} \lambda_n^2 & \end{array}\right\}$$

✎PROOF:

$\mathsf{E}\hat{\mu} \triangleq \mathsf{E}\sum_{n\in\mathbb{Z}} \lambda_n \mathsf{x}_n$	by definition of *arithmetic mean*	(Definition M.4 page 295)
$= \sum_{n\in\mathbb{Z}} \lambda_n \mathsf{E}\mathsf{x}_n$	by *linearity* of E	(Theorem 1.1 page 4)
$= \mu \sum_{n\in\mathbb{Z}} \lambda_n$	by *WSS* hypothesis	(A)
$= \mu$	by $\sum \lambda_n = 1$ hypothesis	(Definition M.4 page 295)
$\mathrm{var}(\hat{\mu}) \triangleq \mathsf{E}(\hat{\mu} - \mathsf{E}\hat{\mu})^2$	by definition of *variance*	
$= \mathsf{E}(\hat{\mu} - \mu)^2$	by previous result	
$= \mathsf{E}\left(\sum_{n=1}^{N} \lambda_n \mathsf{x}_n - \mu\right)^2$	by definition of $\hat{\mu}$	
$= \mathsf{E}\left[\sum_{n=1}^{N} \lambda_n \mathsf{x}_n - \mu \underbrace{\sum_{n=1}^{N} \lambda_n}_{1}\right]^2$	by $\sum \lambda_n = 1$ hypothesis	(Definition M.4 page 295)

$$= \mathsf{E}\left[\sum_{n=1}^{N} \lambda_n(\mathsf{x}_n - \mu)\right]^2$$

$$= \mathsf{E}\left[\sum_{n=1}^{N} \lambda_n(\mathsf{x}_n - \mu) \sum_{m=1}^{N} \lambda_m(\mathsf{x}_m - \mu)\right]$$

$$= \sum_{n=1}^{N}\sum_{m=1}^{N} \lambda_n\lambda_m\big(\mathsf{E}\big[(\mathsf{x}_n - \mu)(\mathsf{x}_m - \mu)\big]\big)$$

$$= \sum_{n=1}^{N}\sum_{m=1}^{N} \lambda_n\lambda_m\big(\mathsf{E}\big[\mathsf{x}_n\mathsf{x}_m\big] - \mu\mathsf{E}\big[\mathsf{x}_n\big] - \mu\mathsf{E}\big[\mathsf{x}_m\big] + \mu^2\big)$$

$$= \sum_{n=1}^{N}\sum_{m=1}^{N} \lambda_n\lambda_m\big(\mathsf{E}\big[\mathsf{x}_n\mathsf{x}_m\big] - \mu^2 - \mu^2 + \mu^2\big)$$

$$= \sum_{n=1}^{N}\sum_{m=1}^{N} \lambda_n\lambda_m\big(\mathsf{E}\big[\mathsf{x}_n\mathsf{x}_m\big] - \mu^2\big)$$

$$= \sum_{n=1}^{N} \lambda_n^2\big(\mathsf{E}\big[\mathsf{x}_n^2\big] - \mu^2\big) + \sum_{n=1}^{N}\sum_{m\neq n} \lambda_n\lambda_m\big(\mathsf{E}\big[\mathsf{x}_n\mathsf{x}_m\big] - \mu^2\big)$$

$$= \sum_{n=1}^{N} \lambda_n^2\big(\mathsf{E}\big[\mathsf{x}_n^2\big] - \mu^2\big) + \sum_{n=1}^{N}\sum_{m\neq n} \lambda_n\lambda_m\big(\mathsf{E}\mathsf{x}_n\mathsf{E}\mathsf{x}_m - \mu^2\big)$$

$$= \sum_{n=1}^{N} \lambda_n^2\sigma^2 + \cancelto{0}{\sum_{n=1}^{N}\sum_{m\neq n} \lambda_n\lambda_m\big(\mu\mu - \mu^2\big)} \qquad \text{by } \textit{WSS} \text{ hypothesis} \qquad \text{(A)}$$

$$= \sigma^2 \sum_{n=1}^{N} \lambda_n^2$$

$$\mathsf{mse}(\hat{\mu}) = \mathsf{E}(\hat{\mu} - \mathsf{E}\hat{\mu})^2 + (\mathsf{E}\hat{\mu} - \mu)^2 \qquad \text{by Theorem 8.2 page 65}$$

$$= \sigma^2 \sum_{n=1}^{N} \lambda_n^2 + (\mu - \mu)^2 \qquad \text{by previous results}$$

$$= \sigma^2 \sum_{n=1}^{N} \lambda_n^2$$

✏

Definition 12.1.

DEF *The **average** $\hat{\mu}$ of a length N sequence $(\!|\mathsf{x}_n|\!)_1^N$ is defined as* $\hat{\mu} \triangleq \frac{1}{N}\sum_{n=1}^{N} \mathsf{x}_n$

Corollary 12.1. [1]

COR

$$\left\{\begin{array}{lll} \text{(A).} & (\!|\mathsf{x}_n|\!) \textit{ is } \text{WIDE SENSE STATIONARY} & \textit{and} \\ \text{(B).} & \mu \triangleq \mathsf{E}\mathsf{x}_n & \textit{and} \\ \text{(C).} & (\!|\mathsf{x}_n|\!) \textit{ is } \text{UNCORRELATED} & \textit{and} \\ \text{(D).} & \hat{\mu} \triangleq \frac{1}{N}\sum_{n=1}^{N} \mathsf{x}_n \quad \text{(AVERAGE)} & \end{array}\right\} \Longrightarrow \left\{\begin{array}{llll} \text{(1).} & \mathsf{E}\hat{\mu} = \mu & \text{(UNBIASED)} & \textit{and} \\ \text{(2).} & \mathsf{var}(\hat{\mu}) = \frac{\sigma^2}{N} & & \textit{and} \\ \text{(3).} & \mathsf{mse}(\hat{\mu}) = \frac{\sigma^2}{N} & \text{(CONSISTENT)} & \end{array}\right\}$$

✎PROOF: These results follow from Theorem 12.1 (page 95) with $\lambda_n = \frac{1}{N}$. ✏

[1] Kay (1988) page 45 ⟨§"3.3 ESTIMATION THEORY"⟩

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

The *arithmetic mean* estimator $\hat{\mu} \triangleq \sum \lambda_n \mathsf{x}_n$ is *unbiased* and *consistent* for any $\sum \lambda_n = 1$ and yields *mean square error* $\mathsf{mse}(\hat{\mu}) = \sigma^2 \sum \lambda_n^2$ (Theorem 12.1 page 95). But…

1. Said qualitatively: "What is the 'best' sequence $(\!(\lambda_n)\!)$ to use?"
2. Said quantitatively: "What sequence $(\!(\lambda_n)\!)$ yields the smallest $\mathsf{mse}(\hat{\mu})$?"

For example, would fashioning $(\!(\lambda_n)\!)$ to be a scaled version of a standard window function, like the *Hanning window*[2] illustrated below, yield the best $\mathsf{mse}(\hat{\mu})$?



Proposition 12.1 (page 98) answers question (2) stating that the best sequence in terms of minimal mse is the sequence $(\!(\lambda_n)\!) \triangleq \frac{1}{N}(\!(\ldots,1,1,1,\ldots)\!)$, which is the *average* estimator, which yields $\mathsf{mse}(\hat{\mu}) = \frac{\sigma^2}{N}$ (Corollary 12.1 page 96).



That is, it turns out that $\frac{1}{N} \leq \sum \lambda_n^2$ for all possible sequences $(\!(\lambda_n)\!)$. This fact is demonstrated by Lemma 12.1 (next), which in turn follows more or less directly from the ubiquitous *Cauchy-Schwarz Inequality* (Theorem M.6 page 298, Theorem J.2 page 256).

Lemma 12.1.

LEM
$$\left\{\sum_{n=1}^{N} \lambda_n = 1\right\} \quad \Longrightarrow \quad \left\{\frac{1}{N} \leq \sum_{n=1}^{N} \lambda_n^2\right\}$$

✎PROOF:

1. Let the sequence $(\!(a_n)\!)$ be defined as $(\!(a_n)\!) \triangleq (\!(\ldots,1,1,1,\ldots)\!)$
2. Let *inner product* $\langle a_n \mid b_n \rangle$ be defined as $\langle a_n \mid b_n \rangle \triangleq \sum_{n=1}^{N} a_n b_n$
3. Let *norm* $\|a_n\|$ be defined as $\|a_n\| \triangleq \sum_{n=1}^{N} a_n^2$
4. Proof of lemma:

$$\begin{aligned}
\boxed{\frac{1}{N}} &= \frac{1}{N}\left(\sum_{n=1}^{N} \lambda_n\right)^2 && \text{by } \sum_{n=1}^{N} \lambda_n = 1 \text{ hypothesis} \\
&= \frac{1}{N}\left(\sum_{n=1}^{N} a_n \lambda_n\right)^2 && \text{by } (\!(a_n)\!) \triangleq (\!(\ldots,1,1,1,\ldots)\!) \text{ definition} && \text{(definition 1 page 97)} \\
&\boxed{\leq} \frac{1}{N}\left(\sum_{n=1}^{N} a_n^2\right)\left(\sum_{n=1}^{N} \lambda_n^2\right) && \text{by } \textit{Cauchy-Schwartz inequality} && \text{(Theorem M.6 page 298)} \\
&\triangleq \frac{1}{N}\left(\sum_{n=1}^{N} 1^2\right)\left(\sum_{n=1}^{N} \lambda_n^2\right) && \text{by definition of } (\!(a_n)\!) && \text{(definition 1 page 97)} \\
&= \boxed{\sum_{n=1}^{N} \lambda_n^2}
\end{aligned}$$

[2] Abdaheer (2009), page 130

COPYRIGHT © 2019 DANIEL J. GREENHOE | CC BY-NC-ND

Proposition 12.1. *Let* mse(*average mean*) *be the mean square error of the* AVERAGE *estimator* *(Corollary 12.1 page 96)* *and* mse(*arithmetic mean*) *be the mean square error of the* ARITHMETIC *estimator* *(Theorem 12.1 page 95)*.

PRP

$$\text{mse}(\textit{average mean}) \leq \text{mse}(\textit{arithmetic mean})$$

PROOF:

$$\begin{aligned}
\text{mse}(\text{average mean}) &= \sigma^2 \frac{1}{N} && \text{by Corollary 12.1 page 96}\\
&\leq \sigma^2 \sum_{n=1}^{N} \lambda_n^2 && \text{by Lemma 12.1 page 97}\\
&= \text{mse}(\text{arithmetic mean}) && \text{by Theorem 12.1 page 95}
\end{aligned}$$

12.2 Variance Estimation

If we know the true *mean* μ of a stationary random process $(\!(\mathsf{x}_n)\!)$, then a reasonable estimate of the variance might be $\frac{1}{N}\sum_{n=1}^{N}(\mathsf{x}_n-\mu)^2$. This estimate has the highly touted property of being *unbiased*:

$$\mathsf{E}\left[\frac{1}{N}\sum_{n=1}^{N}(\mathsf{x}_n-\mu)^2\right] = \frac{1}{N}\sum_{n=1}^{N}\mathsf{E}\left[(\mathsf{x}_n-\mu)^2\right] = \frac{1}{N}\sum_{n=1}^{N}\sigma^2 = \sigma^2$$

Very good. However, in many cases we don't know the **true mean** μ, but rather only have an **estimated mean** $\hat{\mu} \triangleq \frac{1}{N} = \sum_{n=1}^{N}\mathsf{x}_n$. In this case, substituting in the estimated mean for the true mean as in $\hat{\text{var}}_B\big((\!(\mathsf{x}_n)\!)\big)$ (next definition) yields a *biased* variance estimate (next theorem).

Definition 12.2. [3] *Let* $\hat{\mu}$ *be an estimate of the mean of a random sequence* $(\!(\mathsf{x}_n)\!)$.

DEF

The ***sample variance*** $\hat{\text{var}}\big((\!(\mathsf{x}_n)\!)\big)$ *is defined as*

$$\hat{\text{var}}\big((\!(\mathsf{x}_n)\!)\big) \triangleq \frac{1}{N-1}\sum_{n=1}^{N}(\mathsf{x}_n-\hat{\mu})^2$$

The ***biased sample variance*** $\hat{\text{var}}_B\big((\!(\mathsf{x}_n)\!)\big)$ *is here defined as*

$$\hat{\text{var}}_B\big((\!(\mathsf{x}_n)\!)\big) \triangleq \frac{1}{N}\sum_{n=1}^{N}(\mathsf{x}_n-\hat{\mu})^2$$

The factor $\frac{N}{N-1}$ such that $\hat{\text{var}}\big((\!(\mathsf{x}_n)\!)\big) = \frac{N}{N-1}\hat{\text{var}}_B\big((\!(\mathsf{x}_n)\!)\big)$ is known as "*Bessel's correction*". Why such "correction" would be useful at all is demonstrated by Theorem 12.2 (next). Theorem 12.2 demonstrates that the *biased sample variance* $\hat{\text{var}}_B\big((\!(\mathsf{x}_n)\!)\big)$ is *biased*, and multiplication by $\frac{N}{N-1}$ makes it *unbiased*.

[3] Wilks (1963a), page 199 ⟨§"8.2 MEANS AND VARIANCES OF MEAN, VARIANCE,..."⟩, Wilks (1963b), page 199 ⟨§"(b) MEAN AND VARIANCE OF SAMPLE VARIANCE"⟩, Kenney (1947), page 125 ⟨"BESSEL'S CORRECTION"⟩, Bajpai (1967), page 509 ⟨???⟩

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

Theorem 12.2. [4] *Let $\hat{\mu}$ be the* AVERAGE *(Definition M.4 page 295) of a sequence $(\!(\mathsf{x}_n)\!)$. Let $\mu^4 \triangleq \mathsf{E}\left[\left(\mathsf{x}_n - \mu\right)^4\right]$ be the* 4TH CENTRAL MOMENT *of x_n.*

THM

$$\left.\begin{array}{lll} (A). & (\!(\mathsf{x}_n)\!) \text{ is WSS} & \text{and} \\ (B). & \mu \triangleq \mathsf{E}\mathsf{x}_n & \text{and} \\ (C). & (\!(\mathsf{x}_n)\!) \text{ is UNCORRELATED} & \end{array}\right\} \Longrightarrow \left\{\begin{array}{llcll} (1). & \mathsf{E}\hat{\mathsf{var}}_B(\!(\mathsf{x}_n)\!) & = & \frac{N-1}{N}\sigma^2 & \text{(BIASED)} \quad \text{and} \\ (2). & \mathsf{E}\hat{\mathsf{var}}(\!(\mathsf{x}_n)\!) & = & \sigma^2 & \text{(UNBIASED)} \quad \text{and} \\ (3). & \mathsf{var}\left[\hat{\mathsf{var}}(\!(\mathsf{x}_n)\!)\right] & = & \frac{1}{N}\left[\mu^4 - \left(\frac{N-3}{N-1}\right)\sigma^4\right] & \text{(CONSISTENT)} \end{array}\right.$$

✎PROOF:

1. lemma: $\mathsf{E}(\mathsf{x}_n\hat{\mu}) = \frac{1}{N}\sigma^2 + \mu^2$. Proof:

$$\begin{aligned}
\mathsf{E}\left(\mathsf{x}_n\hat{\mu}\right) &\triangleq \mathsf{E}\left(\mathsf{x}_n\frac{1}{N}\sum_{m=1}^{N}\mathsf{x}_m\right) && \text{by definition of } \textit{average} && \text{(Definition M.4 page 295)} \\
&= \mathsf{E}\left(\frac{1}{N}\sum_{m=1}^{N}\mathsf{x}_n\mathsf{x}_m\right) \\
&= \frac{1}{N}\sum_{m=1}^{N}\mathsf{E}\left(\mathsf{x}_n\mathsf{x}_m\right) && \text{by } \textit{linearity} \text{ of } \mathsf{E} && \text{(Theorem 1.1 page 4)} \\
&= \frac{1}{N}\left[\mathsf{E}\mathsf{x}_n^2 + \sum_{m\neq n}\mathsf{E}\left(\mathsf{x}_n\mathsf{x}_m\right)\right] \\
&= \frac{1}{N}\left[\mathsf{E}\mathsf{x}_n^2 + \sum_{m\neq n}\left(\mathsf{E}\mathsf{x}_n\right)\left(\mathsf{E}\mathsf{x}_m\right)\right] && \text{by } \textit{uncorrelated} \text{ hypothesis} && \text{(C)} \\
&= \frac{1}{N}\left[\left(\sigma^2+\mu^2\right) + (N-1)\mu^2\right] && \text{by Corollary 1.3 (page 5)} \\
&= \frac{1}{N}\sigma^2 + \mu^2
\end{aligned}$$

2. Proof for (1):

$$\begin{aligned}
\mathsf{E}\hat{\mathsf{var}}_B\left((\!(\mathsf{x}_n)\!)\right) &\triangleq \mathsf{E}\left[\frac{1}{N}\sum_{n=1}^{N}\left(\mathsf{x}_n - \hat{\mu}\right)^2\right] && \text{by definition of } \hat{\mathsf{var}}_B \text{ (Definition 12.2 page 98)} \\
&= \frac{1}{N}\mathsf{E}\left[\sum_{n=1}^{N}\left(\mathsf{x}_n \underbrace{-\mu+\mu}_{0} - \hat{\mu}\right)^2\right] \\
&= \frac{1}{N}\sum_{n=1}^{N}\left[\underbrace{\mathsf{E}\left(\mathsf{x}_n-\mu\right)^2}_{\sigma^2} + 2\mathsf{E}\left[(\mathsf{x}_n-\mu)(\mu-\hat{\mu})\right] + \mathsf{E}\left(\mu - \underbrace{\hat{\mu}}_{\mathsf{E}\mu}\right)^2\right] \\
&= \frac{1}{N}\sum_{n=1}^{N}\left[\sigma^2 + 2\mathsf{E}\left[\mathsf{x}_n\mu - \mathsf{x}_n\hat{\mu} - \mu^2 + \mu\hat{\mu}\right] + \frac{1}{N}\sigma^2\right] \\
&= \frac{1}{N}\sum_{n=1}^{N}\left[\sigma^2 + 2\left[\mu^2 - \mathsf{E}(\mathsf{x}_n\hat{\mu}) - \mu^2 + \mu^2\right] + \frac{1}{N}\sigma^2\right] && \text{by Corollary 12.1 page 96}
\end{aligned}$$

[4] Wilks (1963a), page 199 ⟨§"8.2 MEANS AND VARIANCES OF MEAN, VARIANCE,…"⟩, TUCKER (1965) PAGE 111 ⟨§"8.2 UNBIASED AND CONSISTENT ESTIMATES"⟩, STUART AND ORD (1991) PAGE 609 ⟨§"UNBIASED ESTIMATORS"⟩

COPYRIGHT © 2019 DANIEL J. GREENHOE

$$= \frac{1}{N}\sum_{n=1}^{N}\left[\sigma^2 + 2\left[\mu^2 - \left(\mu^2 + \frac{1}{N}\sigma^2\right)\right] + \frac{1}{N}\sigma^2\right] \qquad \text{by } \textit{unbiased} \text{ property of } \hat{\mu}$$

((1) lemma pa

$$= \frac{1}{N}\sum_{n=1}^{N}\left[\sigma^2 - \frac{1}{N}\sigma^2\right]$$

$$= \frac{N-1}{N}\sigma^2$$

3. Proof for (2):

$$\begin{aligned}
\mathsf{E}\hat{\text{var}}\big((\mathsf{x}_n)\big) &\triangleq \mathsf{E}\left[\frac{1}{N-1}\sum_{n=1}^{N}\left(\mathsf{x}_n - \hat{\mu}\right)^2\right] && \text{by definition of } \hat{\text{var}} && \text{(Definition 12.2 page 98)}\\
&= \frac{N}{N-1}\mathsf{E}\left[\frac{1}{N}\sum_{n=1}^{N-1}\left(\mathsf{x}_n - \hat{\mu}\right)^2\right] && \text{by } \textit{linearity} \text{ of } \mathsf{E} && \text{(Theorem 1.1 page 4)}\\
&= \frac{N}{N-1}\left[\frac{N-1}{N}\sigma^2\right] && \text{by } \hat{\text{var}}_B \text{ result} && \text{(item (2) page 99)}\\
&= \sigma^2
\end{aligned}$$

4. lemma: $\mathsf{E}\left[\left(\hat{\text{var}}\right)^2\right] = \frac{\mu^4}{N} + \frac{(N-1)^2+2}{N(N-1)}\sigma^4$. Proof: No proof here at this time. The assertion is made by Wilks (1963a), page 199 who also without there supplying a proof says, "Carrying out similar mean value operations we find after some reduction that" the result follows.

$$\begin{aligned}
\mathsf{E}\left[\left(\hat{\text{var}}\right)^2\right] &\triangleq \mathsf{E}\left[\left(\frac{1}{N-1}\sum_{n=1}^{N}\left(\mathsf{x}_n - \hat{\mu}\right)^2\right)^2\right] && \text{by definition of } \hat{\text{var}} \text{ (Definition 12.2 page 98)}\\
&= \mathsf{E}\left[\left(\frac{1}{N-1}\sum_{n=1}^{N}\left(\mathsf{x}_n - \hat{\mu}\right)^2\right)\left(\frac{1}{N-1}\sum_{m=1}^{N}\left(\mathsf{x}_m - \hat{\mu}\right)^2\right)\right]\\
&= \left(\frac{1}{N-1}\right)^2\mathsf{E}\left[\sum_{n=1}^{N}\sum_{m=1}^{N}\left(\mathsf{x}_n - \hat{\mu}\right)^2\left(\mathsf{x}_m - \hat{\mu}\right)^2\right]\\
&= \left(\frac{1}{N-1}\right)^2\mathsf{E}\left[\sum_{n=1}^{N}\sum_{m=1}^{N}\left(\mathsf{x}_n^2 - 2\mathsf{x}_n\hat{\mu} + \hat{\mu}^2\right)\left(\mathsf{x}_m^2 - 2\mathsf{x}_m\hat{\mu} + \hat{\mu}^2\right)\right]\\
&= \left(\frac{1}{N-1}\right)^2\mathsf{E}\left[\sum_{n=1}^{N}\sum_{m=1}^{N}\left[\begin{array}{c}\left(\mathsf{x}_n^2\mathsf{x}_m^2 - 2\hat{\mu}\mathsf{x}_n^2\mathsf{x}_m + \hat{\mu}^2\mathsf{x}_n^2\right) + \left(-2\hat{\mu}\mathsf{x}_n\mathsf{x}_m^2 + 4\hat{\mu}^2\mathsf{x}_n\mathsf{x}_m - 2\mathsf{x}_n\hat{\mu}^3\right)\\ + \left(\hat{\mu}^2\mathsf{x}_m^2 - 2\hat{\mu}^3\mathsf{x}_m + \hat{\mu}^4\right)\end{array}\right]\right]\\
&= \left(\frac{1}{N-1}\right)^2\sum_{n=1}^{N}\sum_{m=1}^{N}\left[\begin{array}{c}\mathsf{E}\left[\mathsf{x}_n^2\mathsf{x}_m^2\right] - 2\mathsf{E}\left[\hat{\mu}\mathsf{x}_n^2\mathsf{x}_m\right] - 2\mathsf{E}\left[\hat{\mu}\mathsf{x}_n\mathsf{x}_m^2\right] + \mathsf{E}\left[\hat{\mu}^2\mathsf{x}_n^2\right] + \mathsf{E}\left[\hat{\mu}^2\mathsf{x}_m^2\right]\\ +4\,\mathsf{E}\left[\hat{\mu}^2\mathsf{x}_n\mathsf{x}_m\right] - 2\mathsf{E}\left[\hat{\mu}^3\mathsf{x}_n\right] - 2\mathsf{E}\left[\hat{\mu}^3\mathsf{x}_m\right] + \mathsf{E}\left[\hat{\mu}^4\right]\end{array}\right]\\
&= \left(\frac{1}{N-1}\right)^2\sum_{n=1}^{N}\sum_{m=1}^{N}\left[\begin{array}{c}\mathsf{E}\left[\mathsf{x}_n^2\mathsf{x}_m^2\right] - 4\mathsf{E}\left[\hat{\mu}\mathsf{x}_n^2\mathsf{x}_m\right] + 2\mathsf{E}\left[\hat{\mu}^2\mathsf{x}_n^2\right]\\ +4\,\mathsf{E}\left[\hat{\mu}^2\mathsf{x}_n\mathsf{x}_m\right] - 4\mathsf{E}\left[\hat{\mu}^3\mathsf{x}_n\right] + \mathsf{E}\left[\hat{\mu}^4\right]\end{array}\right]\\
&\stackrel{?}{=} \frac{\mu^4}{N} + \frac{(N-1)^2+2}{N(N-1)}\sigma^4
\end{aligned}$$

BY-NC-ND Copyright © 2019 Daniel J. Greenhoe

5. Proof for (3):

$$\begin{aligned}
\operatorname{var}\left[\hat{\operatorname{var}}(\!(x_n)\!)\right] &\triangleq \mathsf{E}\left[\left(\hat{\operatorname{var}} - \mathsf{E}\hat{\operatorname{var}}\right)^2\right] && \text{by definition of } \hat{\operatorname{var}} && \text{(Definition 12.2 page 98)}\\
&= \mathsf{E}\left[\left(\hat{\operatorname{var}} - \sigma^2\right)^2\right] && \text{by (2)} && \text{(item (3) page 100)}\\
&= \mathsf{E}\left[\left(\hat{\operatorname{var}}\right)^2 - 2\sigma^2\hat{\operatorname{var}} + \left(\sigma^2\right)^2\right] && \text{by } \textit{Binomial Theorem} && \\
&= \mathsf{E}\left[\left(\hat{\operatorname{var}}\right)^2\right] - 2\sigma^2\mathsf{E}\left[\hat{\operatorname{var}}\right] + \mathsf{E}\left[\left(\sigma^2\right)^2\right] && \text{by } \textit{linearity} \text{ of } \mathsf{E} && \text{(Theorem 1.1 page 4)}\\
&= \mathsf{E}\left[\left(\hat{\operatorname{var}}\right)^2\right] - 2\left(\sigma^2\right)^2 + \left(\sigma^2\right)^2 && \text{by (2)} && \text{(item (3) page 100)}\\
&= \left[\frac{\mu^4}{N} + \left(\frac{(N-1)^2+2}{N(N-1)}\right)\sigma^4\right] - \sigma^4 && \text{by (4) lemma} && \\
&= \frac{1}{N}\left[\mu^4 + \left(\frac{(N^2-N+1)+2-N(N-1)}{N-1}\right)\sigma^4\right] && && \\
&= \frac{1}{N}\left[\mu^4 - \left(\frac{N-3}{N-1}\right)\sigma^4\right] && &&
\end{aligned}$$

✏

12.3 Estimates in terms of moment estimates

Definition 12.3.

DEF

*The **order-k moment estimate** is here defined as*

$$\hat{\mathsf{M}}_k(\!(x_n)\!) \triangleq \frac{1}{N}\sum_{n=1}^{N} x_n^k$$

Proposition 12.2.

PRP

$$\begin{aligned}
\hat{\mu}(\!(x_n)\!) &= \hat{\mathsf{M}}_1\\
\hat{\operatorname{var}}(\!(x_n)\!) &= \frac{N}{N-1}\hat{\mathsf{M}}_2 - \frac{N}{N-1}\hat{\mathsf{M}}_1^2
\end{aligned}$$

✎PROOF:

$$\begin{aligned}
\hat{\mu}(\!(x_n)\!) &\triangleq \frac{1}{N}\sum_{n=1}^{N} x_n && \text{by definition of } \hat{\mu} && \text{(Definition M.4 page 295)}\\
&\triangleq \hat{\mathsf{M}}_1 && \text{by definition of } \hat{\mathsf{M}}_1 && \text{(Definition 12.3 page 101)}
\end{aligned}$$

$$\begin{aligned}
\hat{\operatorname{var}}(\!(x_n)\!) &\triangleq \frac{1}{N-1}\sum_{n=1}^{N}\left(x_n - \hat{\mu}\right)^2 && \text{by definition of } \hat{\operatorname{var}} && \text{(Definition 12.2 page 98)}\\
&= \frac{1}{N-1}\sum_{n=1}^{N}\left(x_n^2 - 2x_n\hat{\mu} + (\hat{\mu})^2\right) && && \\
&= \frac{1}{N-1}\sum_{n=1}^{N} x_n^2 - 2\frac{1}{N-1}\sum_{n=1}^{N} x_n\hat{\mu} + \frac{1}{N-1}\sum_{n=1}^{N}(\hat{\mu})^2 && && \\
&= \frac{N}{N-1}\underbrace{\frac{1}{N}\sum_{n=1}^{N} x_n^2}_{\hat{\mathsf{M}}_2} - 2\underbrace{\hat{\mu}}_{\hat{\mathsf{M}}_1}\frac{N}{N-1}\underbrace{\frac{1}{N}\sum_{n=1}^{N} x_n}_{\hat{\mathsf{M}}_1} + \frac{N}{N-1}\underbrace{(\hat{\mu})^2}_{\hat{\mathsf{M}}_1^2} && &&
\end{aligned}$$

COPYRIGHT © 2019 DANIEL J. GREENHOE | CC BY-NC-ND

$$= \frac{N}{N-1}\hat{\mathrm{M}}_2 - \frac{N}{N-1}\hat{\mathrm{M}}_1^2$$

✏

Copyright © 2019 Daniel J. Greenhoe

CHAPTER 13

CORRELATION ESTIMATION

Definition 13.1. [1]

DEF

*The **windowed auto-correlation estimate** $\hat{\mathsf{R}}_{\mathsf{xx}}(m)$ is defined as*

$$\hat{\mathsf{R}}_{\mathsf{xx}}(m) \triangleq \frac{1}{N} \sum_{n=0}^{N-|m|} \mathsf{x}(n)\mathsf{x}(n+m)$$

Theorem 13.1. [2]

THM

$$\begin{aligned} \mathsf{E}\left[\hat{\mathsf{R}}_{\mathsf{xx}}(m)\right] &= \left(1 - \frac{|m|}{N}\right)\mathsf{R}_{\mathsf{xx}}(m) && \text{(ASYMPTOTICALLY UNBIASED)} \\ \operatorname{var}\left[\hat{\mathsf{R}}_{\mathsf{xx}}(m)\right] &= \frac{1}{N}\sum_{n\in\mathbb{Z}}\left[\mathsf{R}_{\mathsf{xx}}^2(n) + \mathsf{R}_{\mathsf{xx}}(n-m)\mathsf{R}_{\mathsf{xx}}(n+m)\right] && \text{(CONSISTENT)} \end{aligned}$$

[1] Vaseghi (2000) page 271 ⟨§"9.3.3 Energy-Spectral Density and Power-Spectral Density"⟩

[2] Jenkins and Watts (1968), Vaseghi (2000) page 272 ⟨§"9.3.3 Energy-Spectral Density and Power-Spectral Density"⟩

Copyright © 2019 Daniel J. Greenhoe

CHAPTER 14

SPECTRAL ESTIMATION

Quality of spectral estimators[1]

THM

Periodogram:	Q	$=$	1
Welch Method 0% overlap:	Q	$=$	$0.78N\Delta f$
Welch Method 50% overlap:	Q	$=$	$1.39N\Delta f$
Bartlett Method:	Q	$=$	$1.11N\Delta f$
Blackman-Tukey Method:	Q	$=$	$2.34N\Delta f$

[1] Proakis (2002) pages 452–457 ⟨§"8.2.4 Performance Characteristics of Nonparametric Power Spectrum Estimators"⟩, Proakis and Manolakis (1996) pages 916–919 ⟨§"12.2.4 Performance Characteristics of Nonparametric Power Spectrum Estimators"⟩, Rao and Swamy (2018) page 731 ⟨"Table 12.1 Comparison of performance of classical methods"⟩, Salivahanan and Vallavaraj (2001) page 606 ⟨§"12.5 Power Spectrum Estimation: Non-Parametric Methods"⟩, Ifeachor and Jervis (2002) pages 706–707 ⟨§"11.3.7 Comparison of the power spectral density estimation methods"⟩, J.S.Chitode (2009b) page P-100, Abdaheer (2009), page 204

Copyright © 2019 Daniel J. Greenhoe

CHAPTER 15

DENSITY ESTIMATION

References:

- Silverman (1986)

Copyright © 2019 Daniel J. Greenhoe

CHAPTER 16

SYSTEM IDENTIFICATION



❝ *I can therefore gladly admit that falsificationists like myself much prefer an attempt to solve an interesting problem by a bold conjecture, even (and especially) if it so turns out to be false, to any recital of a sequence of irrelevant truisms. We prefer this because we believe that this is the way in which we can learn from our mistakes and that in finding that our conjecture was false we shall have learned much about the truth, and shall have got nearer to the truth.* ❞

Karl R. Popper (1902–1994) [1]

16.1 Estimation techniques

Let **S** be a *system* with *impulse response* $\mathsf{h}(n)$ with with *DTFT* $\tilde{\mathsf{H}}(\omega)$, input $\mathsf{x}(n)$, and output $\mathsf{y}(n)$. Often in the field of "digital signal processing" (DSP), **S** is a "filter" with known $\mathsf{h}(n)$ and $\tilde{\mathsf{H}}(\omega)$ because the filter **S** was designed by a designer who had direct control over $\mathsf{h}(n)$.

However in many other practical situations, **S** is some other system for which $\mathsf{h}(n)$ and $\tilde{\mathsf{H}}(\omega)$ are *not* known…but which we may want to *estimate*. Examples of such **S** is a device on an industrial shaker table, a communication channel, or the entire earth.

Determining $\mathsf{h}(n)$ and/or $\tilde{\mathsf{H}}(\omega)$ is part of an operation called "*system identication*". Determining $\tilde{\mathsf{H}}(\omega)$ in particular is referred to as "*Frequency Response Identification*"[2] or as "*Frequency Response Function*" ("*FRF*") estimation.[3] *FRF* estimation is a challenging problem and one that many have devoted much effort to. This chapter describes some of that effort.

In the early days, people used a rather obvious technique for determining $\tilde{\mathsf{H}}(\omega)$—the humble *sine sweep*. That is, they drove the input with a sine wave with slowly increasing (or decreasing) frequency while measuring the resulting output. This technique, although effective, was "very slow".[4]

[1] quote: Popper (1962), page 231, Popper (1963) page 313
image: https://en.wikipedia.org/wiki/File:Karl_Popper.jpg, "no known copyright restrictions"

[2] Shin and Hammond (2008) page 292

[3] Cobb (1988) page 1 ⟨FRF "measurement"⟩

[4] Leuridan et al. (1986)911"Stepped Sine Testing", Cobb (1988) page 1 ⟨Chapter 1—Introduction⟩, Ewins



(A) **communications additive noise model**
The "input signal" is $x(n)$.

(B) **measurement additive noise model**
The "input signal" is $p(n)$.

In each model, $x(n)$ and $y(n)$ are "known", and $u(n)$, $v(n)$, and $w(n)$ are *not*.
In definition, the two models are **equivalent** under the relation $u(n) = -w(n)$.
In practice, they are **different**:

- in (A), x and u would be typically *uncorrelated*;
- in (B), x and $w = -u$ are very much *correlated* (x is a function of u).

Figure 16.1: Additive noise systems with *linear*/**non-linear** operator **T**

And there is another problem here—we don't always have control over the input signal. Examples of this include earthquake and volcanic activity analysis.

An alternative to the sine-sweep input is *random sequence* input. All the techniques that follow in this chapter are of this type. A problem with using random sequences directly for estimating $\tilde{H}(\omega)$ is that the estimate $\hat{H}(\omega)$ is itself also random. This is not what we want. We want an estimate that we can actually write down on paper or at least plot on paper.

A solution to this is to not use the random sequences directly to estimate $\tilde{H}(\omega)$, but instead to first use the *expectation* operator E (Definition 1.1 page 3). The expectation operator takes a quantity X that is inherently "random" (with some probability distribution $p(x)$) and turns it into a deterministic "constant" EX.

The operator E is also used by the spectral density functions $\tilde{S}_{xx}(\omega)$ and $\tilde{S}_{xy}(\omega)$ (Definition 6.3 page 44). And $\tilde{S}_{xx}(\omega)$ and $\tilde{S}_{xy}(\omega)$ are what are typically used to calculate an estimate $\hat{H}(\omega)$.

16.2 Additive noise system models

Consider the additive noise systems illustrated in Figure 17.1 (page 135).

- The illustration on the left is suitable for modeling a communications system where x is the transmitted signal, y is the received signal, u and v are thermal noise, and the "transfer function" **H** is the communications channel (air, water, wires, etc.) that one wishes to estimate.
- The illustration on the right is suitable for modeling a testing system where p is an input test signal (from an industrial shaker or from a naturally occurring signal originating from geophysical activity), w is measurement noise, x is the measured input contaminated by noise, and **H** is the device under test (a piece of equipment, a building, or the entire earth).

Note that the two models are an equivalent system **S** under the relation $u = -w$. But although one might expect such a sign difference to wreak mathematical havoc in resulting equations, this is

(1986) pages 125–140 ⟨3.7 Use of different excitation types⟩

BY-NC-ND
Copyright © 2019 Daniel J. Greenhoe

simply not the case here because

$$\tilde{S}_{ww} = \tilde{\mathbf{F}}E[w(m)w*(0)] = \tilde{\mathbf{F}}E\big[(-u(m))\big(-u^*(0)\big)\big] = \tilde{\mathbf{F}}E\big[(u(m))\big(u^*(0)\big)\big] = \tilde{S}_{uu}$$

So the sign difference is not that big of a difference after all. But there are some key differences in practice:

- In the communications model (on the left), the "input signal" is $x(n)$ and the frequency-domain input *signal-to-noise ratio* (*SNR*) is $\tilde{S}_{xx}(\omega)/\tilde{S}_{uu}(\omega)$. In the measurement model (on the right), the "input signal" is $p(n)$ and the frequency-domain input *signal-to-noise ratio* (*SNR*) is $\tilde{S}_{pp}(\omega)/\tilde{S}_{ww}(\omega) = \tilde{S}_{pp}(\omega)/\tilde{S}_{uu}(\omega)$.
- On the left, x and u would be typically *uncorrelated*; on the right, x and $w = -u$ are very much *correlated* (x is a function of u).

16.3 Transfer function estimate definitions and interpretation

As a first attempt at estimating the transfer function $\mathbf{H}$ of $\mathbf{S}$, or at least the magnitude squared of $\mathbf{H}$, we might assume $\mathbf{H}$ to be *LTI*, take a cue from the relation $\tilde{S}_{yy} = \tilde{S}_{xx}|\tilde{H}|^2$ of Corollary 5.3 (page 37), and arrive at a function called "*transmissibility*" (next definition).

Definition 16.1. [5] *Let* $\mathbf{S}$ *be a* SYSTEM *with input* $x(n)$ *and output* $y(n)$.

DEF ***transmissibility*** $\tilde{T}_{xy}(\omega)$ *is defined as* $$\tilde{T}_{xy}(\omega) \triangleq \sqrt{\frac{\tilde{S}_{yy}(\omega)}{\tilde{S}_{xx}(\omega)}}$$

Transmissibility is in essence the ratio of "*spectral power*" (Remark 6.1 page 44) output to *spectral power* input. Note that it is a real-valued function (because $\tilde{S}_{xx}$ and $\tilde{S}_{yy}$ are real-valued). We might suspect that we could attain better estimates of $\mathbf{H}$ by allowing the estimates to be complex-valued. And in fact, all the remaining estimates in this section are in general complex-valued.

And so to start (again), and in the very special (a.k.a unrealistic) case of $\mathbf{S}$ having *zero measurement noise* (*zero measurement error*) ($v = u = w = 0$), $h(n)$ being *linear time invariant* (*LTI*), and input $x(n)$ being *wide sense stationary*…then we can determine (a.k.a "identify") $h(n)$ or $\tilde{H}(\omega)$ exactly by $\tilde{H}(\omega) = \tilde{S}_{yx}(\omega)/\tilde{S}_{xx}(\omega)$ (Corollary 5.3 page 37).

However, in practical situations, there is measurement noise/error. Examples may include "road noise" from a test being performed in a moving vehicle or *quantization noise* from an *analog-to-digital converter* (*ADC*).

If the measurement error is at the output only (and under the assumptions of *LTI* and *WSS*) then $\hat{H}_1$ (next definition) is the ideal estimator in the sense that $\hat{H}_1 = \tilde{H}$ (Corollary 16.4 page 129).



Definition 16.2. [6] *Let* $\mathbf{S}$ *be a* SYSTEM *with input* $x(n)$ *and output* $y(n)$.

[5] Bendat and Piersol (2010) page 469 $\langle |H(f)| = \big[G_{yy}(f)/G_{xx}(f)\big]^{1/2}\rangle$, Yan and Ren (2012) page 204 $\langle (1)\ \big[G_{YY}(s)\big] = [H(s)]\big[G_{FF}(s)\big]\big[H^*(s)\big]^T\rangle$, Goldman (1999) page 179 ⟨Transmissibility …$H'_{ab} = G_{bb}/G_{aa}$ (note: differs by $\sqrt{\cdot}$ from Bendat and Piersol⟩, Zhang et al. (2016), Zhou and Wahab (2018) page 824, https://link.springer.com/chapter/10.1007/978-3-319-54109-9_4

[6] Bendat and Piersol (1993) pages 106–109 ⟨5.1.1 Optimality of Calculations⟩, Bendat and Piersol (2010) page 185 $\langle H_1(f) = G_{xy}(f)/G_{xx}(f)$ (6.37)⟩, Shin and Hammond (2008) page 293 $\langle H_1(f) = \tilde{S}_{xy}(f)/\tilde{S}_{xx}(f)$ (9.63); which dif-

COPYRIGHT © 2019 DANIEL J. GREENHOE | CC BY-NC-ND

DEF *The* ***Least Squares transfer function estimate*** $\hat{\mathsf{H}}_1(\omega)$ *of* **S** *is defined as* $\hat{\mathsf{H}}_1(\omega) \triangleq \dfrac{\tilde{\mathsf{S}}_{\mathsf{yx}}(\omega)}{\tilde{\mathsf{S}}_{\mathsf{xx}}(\omega)}$

The estimator $\hat{\mathsf{H}}_1$ is a good start. However in the early 1980s, L. D. Mitchell pointed out that in the presence of input noise, $\hat{\mathsf{H}}_1$ is far from ideal in that it is *biased* with respect to $\tilde{\mathsf{H}}$; in fact, $\hat{\mathsf{H}}_1$ *under estimates* $\tilde{\mathsf{H}}$ (Corollary 16.4 page 129). Mitchell proposed a new estimator $\hat{\mathsf{H}}_2$ (next definition).

This estimator has the special property that when there is input noise but no output noise (and under *LTI*, *WSS*, and *uncorrelated* assumptions), then it is ideal in the sense that $\hat{\mathsf{H}}_2(\omega) = \tilde{\mathsf{H}}(\omega)$ (Corollary 16.4 page 129).



Note also that in the case of both no input and no output noise, then $\hat{\mathsf{H}}_1 = \hat{\mathsf{H}}_2$ (Corollary 5.3 page 37).

Definition 16.3. [7] *Let* **S** *be a* SYSTEM *with input* $\mathsf{x}(n)$ *and output* $\mathsf{y}(n)$.

DEF *The* ***Inverse Method transfer function estimate*** $\hat{\mathsf{H}}_2(\omega)$ *of* **S** *is defined as* $\hat{\mathsf{H}}_2(\omega) \triangleq \dfrac{\tilde{\mathsf{S}}_{\mathsf{yy}}(\omega)}{\tilde{\mathsf{S}}_{\mathsf{xy}}(\omega)}$

Mitchell's $\hat{\mathsf{H}}_2$ contribution "generated a flurry of activity"[8] and soon more $\tilde{\mathsf{H}}$ estimators appeared. So far we have

- $\hat{\mathsf{H}}_1$ which is ideal when there is no input noise but *under estimates* $\tilde{\mathsf{H}}$ when there is (Corollary 16.4 page 129)
- $\hat{\mathsf{H}}_2$ which is ideal when there is no output noise but *over estimates* $\tilde{\mathsf{H}}$ when there is (Corollary 16.4 page 129).

But what about estimators for when there is noise on both input and output? Armed with two estimators that between them account for both input and output noise, an "ad hoc" solution might be to somehow take mean values of $\hat{\mathsf{H}}_1$ and $\hat{\mathsf{H}}_2$ to induce new estimators—this approach summarizes the next three definitions. An arguably more mature approach is to find estimators that are optimal with respect to least squares measures—and this approach summarizes Definition 16.9 – Definition 16.7 (page 115).

Definition 16.4. *Let* **S** *be a* SYSTEM *with input* $\mathsf{x}(n)$ *and output* $\mathsf{y}(n)$.

DEF *The* ***Arithmetic Mean transfer function estimate*** $\hat{\mathsf{H}}_{\mathsf{am}}(\omega)$ *of* **S** *is defined as*

$$\hat{\mathsf{H}}_{\mathsf{am}}(\omega) \triangleq \frac{\left|\tilde{\mathsf{S}}_{\mathsf{xy}}(\omega)\right|^2 + \tilde{\mathsf{S}}_{\mathsf{xx}}(\omega)\tilde{\mathsf{S}}_{\mathsf{yy}}(\omega)}{2\tilde{\mathsf{S}}_{\mathsf{xx}}(\omega)\tilde{\mathsf{S}}_{\mathsf{xy}}(\omega)}$$

fers from Definition 16.2, but see APPENDIX ?? page ?? ⟩, Bendat (1978)cited by Cobb(1988)—variance estimate for $\hat{\mathsf{H}}_1$, Allemang et al. (1979) ⟨cited by Shin(2008)⟩, Leuridan et al. (1986) page 910 ⟨*Least Squares Technique*; (8) $[G_{xx}](H) = [G_{xy}]$⟩, Abom (1986)cited by Cobb(1988)—variance estimate for $\hat{\mathsf{H}}_1$, Allemang et al. (1987) pages 54–55 ⟨5.3.1 H_1 Technique; $[H] = [G_{XF}][G_{FF}]^{-1}$ (11)⟩, Cobb (1988) page 2 ⟨[1] $\hat{H}(f) = \hat{G}_{yx}(f)/\hat{G}_{xx}(f)$ (1)⟩, Goyder (1984) page 438 ⟨$H(i\omega) = S_{qp}/S_{pp}$ (3)⟩, Pintelon and Schoukens (2012) page 233 ⟨$\hat{G}(\Omega_k) = S_{yu}(j\omega_k)S_{uu}^{-1}(j\omega_k)$ (7-30)⟩, White et al. (2006) page 678 ⟨$H_1(f) = \hat{S}_{x_m y_m}(f)/\hat{S}_{x_m x_m}(f)$ (1) which differs by conjugate, references Bendat and Piersol⟩,

[7] Shin and Hammond (2008) page 293 ⟨$H_2(f) = \tilde{\mathsf{S}}_{\mathsf{yy}}(f)/\tilde{\mathsf{S}}_{\mathsf{yx}}(f)$ (9.65); which differs from Definition 16.3, but see APPENDIX ?? page ?? ⟩, Bendat and Piersol (2010) page 186 ⟨$H_2(f) = G_{yy}(f)/G_{yx}(f)$ (6.42)⟩, Mitchell (1980) ⟨cited by Cobb(1988)⟩, Mitchell (1982) page 278 ⟨"Define what will be called an inverse method for calculation of a FRF as…"; $H_2(f) = G_{yy}/G_{yx}$ (6); Note this differs with Definition 16.3 by a conjugate, but note that Mitchell seems to follow Bendat (see his [3] and [4]), which would explain this difference (APPENDIX ?? page ??)⟩, Cobb (1988) page 3 ⟨[2] $\hat{H}(f) = \hat{G}_{yy}(f)/\hat{G}_{xy}(f)$ (1)⟩, White et al. (2006) page 678 ⟨$H_2(f) = \hat{S}_{y_m y_m}(f)/\hat{S}_{y_m x_m}(f)$ (2) which differs by conjugate, references Bendat and Piersol⟩

[8] Cobb (1988) page 3

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

Proposition 16.1. [9] *Let* $\mathbf{S}$ *be a* SYSTEM *with input* $\mathsf{x}(n)$ *and output* $\mathsf{y}(n)$.

PRP

$$\hat{\mathsf{H}}_{\mathsf{am}}(\omega) = \frac{\hat{\mathsf{H}}_1(\omega) + \hat{\mathsf{H}}_2(\omega)}{2} \quad \textit{(\textbf{arithmetic mean} of } \hat{\mathsf{H}}_1 \textit{ and } \hat{\mathsf{H}}_2\textit{)}$$

✎PROOF:

$$\hat{\mathsf{H}}_{\mathsf{am}}(\omega) \triangleq \frac{\left|\tilde{\mathsf{S}}_{\mathsf{xy}}(\omega)\right|^2 + \tilde{\mathsf{S}}_{\mathsf{xx}}(\omega)\tilde{\mathsf{S}}_{\mathsf{yy}}(\omega)}{2\tilde{\mathsf{S}}_{\mathsf{xx}}(\omega)\tilde{\mathsf{S}}_{\mathsf{xy}}(\omega)} \quad \text{by definition of } \hat{\mathsf{H}}_{\mathsf{am}} \quad \text{(Definition 16.4 page 112)}$$

$$= \frac{\tilde{\mathsf{S}}_{\mathsf{xy}}(\omega)\tilde{\mathsf{S}}^*_{\mathsf{xy}}(\omega)}{2\tilde{\mathsf{S}}_{\mathsf{xx}}(\omega)\tilde{\mathsf{S}}_{\mathsf{xy}}(\omega)} + \frac{\tilde{\mathsf{S}}_{\mathsf{xx}}(\omega)\tilde{\mathsf{S}}_{\mathsf{yy}}(\omega)}{2\tilde{\mathsf{S}}_{\mathsf{xx}}(\omega)\tilde{\mathsf{S}}_{\mathsf{xy}}(\omega)} = \frac{\frac{\tilde{\mathsf{S}}^*_{\mathsf{xy}}(\omega)}{\tilde{\mathsf{S}}_{\mathsf{xx}}(\omega)} + \frac{\tilde{\mathsf{S}}_{\mathsf{yy}}(\omega)}{\tilde{\mathsf{S}}_{\mathsf{xy}}(\omega)}}{2}$$

$$= \frac{\hat{\mathsf{H}}_1(\omega) + \hat{\mathsf{H}}_2(\omega)}{2} \quad \text{by definitions of } \hat{\mathsf{H}}_1 \text{ and } \hat{\mathsf{H}}_2 \quad \text{(Definition 16.2 page 111, Definition 16.3 page 112)}$$

✏

Definition 16.5. *Let* $\mathbf{S}$ *be a* SYSTEM *with input* $\mathsf{x}(n)$ *and output* $\mathsf{y}(n)$.

DEF

The ***Geometric mean transfer function estimate*** $\hat{\mathsf{H}}_{\mathsf{gm}}(\omega)$ *of* $\mathbf{S}$ *is defined as*

$$\hat{\mathsf{H}}_{\mathsf{gm}}(\omega) \triangleq \frac{\tilde{\mathsf{S}}^*_{\mathsf{xy}}(\omega)}{\left|\tilde{\mathsf{S}}_{\mathsf{xy}}(\omega)\right|}\sqrt{\frac{\tilde{\mathsf{S}}_{\mathsf{yy}}(\omega)}{\tilde{\mathsf{S}}_{\mathsf{xx}}(\omega)}}$$

Proposition 16.2. [10] *Let* $\mathbf{S}$ *be a* SYSTEM *with input* $\mathsf{x}(n)$ *and output* $\mathsf{y}(n)$.

PRP

$$\pm\hat{\mathsf{H}}_{\mathsf{gm}}(\omega) = \sqrt{\hat{\mathsf{H}}_1(\omega)\hat{\mathsf{H}}_2(\omega)} \quad \textit{(\textbf{geometric mean} of } \hat{\mathsf{H}}_1 \textit{ and } \hat{\mathsf{H}}_2\textit{)}$$

✎PROOF:

$$\pm\hat{\mathsf{H}}_{\mathsf{gm}}(\omega) \triangleq \pm\frac{\tilde{\mathsf{S}}^*_{\mathsf{xy}}(\omega)}{\left|\tilde{\mathsf{S}}_{\mathsf{xy}}(\omega)\right|}\sqrt{\frac{\tilde{\mathsf{S}}_{\mathsf{yy}}(\omega)}{\tilde{\mathsf{S}}_{\mathsf{xx}}(\omega)}} \quad \text{by definition of } \hat{\mathsf{H}}_{\mathsf{gm}} \quad \text{(Definition 16.5 page 113)}$$

$$= \sqrt{\frac{\left[\tilde{\mathsf{S}}^*_{\mathsf{xy}}(\omega)\right]^2}{\left|\tilde{\mathsf{S}}_{\mathsf{xy}}(\omega)\right|^2}\frac{\tilde{\mathsf{S}}_{\mathsf{yy}}(\omega)}{\tilde{\mathsf{S}}_{\mathsf{xx}}(\omega)}} = \sqrt{\frac{\tilde{\mathsf{S}}^*_{\mathsf{xy}}(\omega)}{\tilde{\mathsf{S}}_{\mathsf{xy}}(\omega)}\frac{\tilde{\mathsf{S}}_{\mathsf{yy}}(\omega)}{\tilde{\mathsf{S}}_{\mathsf{xx}}(\omega)}} = \sqrt{\frac{\tilde{\mathsf{S}}^*_{\mathsf{xy}}(\omega)}{\tilde{\mathsf{S}}_{\mathsf{xx}}(\omega)}\frac{\tilde{\mathsf{S}}_{\mathsf{yy}}(\omega)}{\tilde{\mathsf{S}}_{\mathsf{xy}}(\omega)}}$$

$$= \sqrt{\hat{\mathsf{H}}_1(\omega)\hat{\mathsf{H}}_2(\omega)} \quad \text{by definitions of } \hat{\mathsf{H}}_1 \text{ and } \hat{\mathsf{H}}_2 \quad \text{(Definition 16.2 page 111, Definition 16.3 page 112)}$$

$$= \textbf{Geometric mean} \text{ of } \hat{\mathsf{H}}_1(\omega) \text{ and } \hat{\mathsf{H}}_2(\omega)$$

Note that for a complex number $z \triangleq |z|e^{i\phi}$, $\sqrt{z}$ has two solutions:[11]

$$\sqrt{z} = \sqrt{|z|e^{i\phi}} = \{z_1, z_2\} = \left\{\sqrt{|z|}e^{i(\phi/2)}, \sqrt{|z|}e^{i(\phi/2+\pi)}\right\} = \pm\sqrt{|z|}e^{i(\phi/2)}$$

because $z_1^2 = z$ and $z_2^2 = z$. ✏

Note that the *geometric mean estimator* (Definition 16.5 page 113) and *transmissibility* (Definition 16.1 page 111) are closely related (next).

[9] Mitchell (1982) page 279 ⟨"Frequency Response Calculation: The Average Method"⟩, Zheng et al. (2002) page 918 ⟨"1.3 Arithmetic Mean Estimator H_3"⟩

[10] Zheng et al. (2002) page 918 ⟨"1.4 Geometric Mean Estimator H_4"⟩

[11] Many many thanks to Ben Cleveland for his help with this!!!

COPYRIGHT © 2019 DANIEL J. GREENHOE | CC BY-NC-ND

Proposition 16.3. *Let $\phi(\omega)$ be the* PHASE *of $\tilde{S}_{xy}(\omega)$ such that $\tilde{S}_{xy}(\omega) \triangleq |\tilde{S}_{xy}(\omega)|e^{-i\phi(\omega)}$*

PRP

$$\hat{H}_{gm}(\omega) = \tilde{T}_{xy}(\omega)\, e^{-i\phi(\omega)} \quad \left(\begin{array}{lcll} |\hat{H}_{gm}(\omega)| & = & \tilde{T}_{xy}(\omega) & \textit{is the } \textsc{magnitude} \textit{ of } \hat{H}_{gm}(\omega) \quad \textit{and} \\ \angle\hat{H}_{gm}(\omega) & = & -\angle\tilde{S}_{xy}(\omega) & \textit{is the } \textsc{phase} \textit{ of } \hat{H}_{gm}(\omega) \end{array} \right)$$

✎PROOF: Let $\phi(\omega)$ be the *phase* of

$$\begin{aligned}
\hat{H}_{gm}(\omega) &\triangleq \sqrt{\hat{H}_1(\omega)\hat{H}_2(\omega)} && \text{by definition of } \hat{H}_{gm} && \text{(Definition 16.5 page 113)} \\
&\triangleq \sqrt{\frac{\tilde{S}^*_{xy}(\omega)}{\tilde{S}_{xx}(\omega)}\frac{\tilde{S}_{yy}(\omega)}{\tilde{S}_{xy}(\omega)}} && \text{by definitions of } \hat{H}_1 \text{ and } \hat{H}_2 && \text{(Definition 16.2 page 111, Definition 16.3 page 112)} \\
&= \sqrt{\frac{\tilde{S}_{yy}(\omega)}{\tilde{S}_{xx}(\omega)}\frac{\tilde{S}^*_{xy}(\omega)}{\tilde{S}_{xy}(\omega)}} \\
&= \tilde{T}_{xy}(\omega)\sqrt{\frac{\tilde{S}^*_{xy}(\omega)}{\tilde{S}_{xy}(\omega)}} && \text{by definition of } \tilde{T}_{xy} && \text{(Definition 16.1 page 111)} \\
&= \tilde{T}_{xy}(\omega)\sqrt{\frac{|\tilde{S}_{xy}(\omega)|e^{-i\phi(\omega)}}{|\tilde{S}_{xy}(\omega)|e^{i\phi(\omega)}}} && \text{where } \tilde{S}_{xy}(\omega) \triangleq |\tilde{S}_{xy}(\omega)|e^{-i\phi(\omega)} \\
&= \tilde{T}_{xy}(\omega)\sqrt{e^{-i2\phi(\omega)}} \\
&= \tilde{T}_{xy}(\omega)\, e^{-i\phi(\omega)}
\end{aligned}$$

✏

Remark 16.1. *Transmissibility* $\tilde{T}_{xy}$ is a kind of "black sheep" of the system identification function family. All the other members of this family ($\hat{H}_1$, $\hat{H}_2$, $\hat{H}_v$, $\hat{H}_s$) are *complex-valued*, but $\tilde{T}_{xy}$ is only *real-valued*—a seemingly ordinary Joe born into a super-hero family. But Proposition 16.3 suggests that $\tilde{T}_{xy}$ is not simply a "black sheep", but rather a "dark horse" with abilities that can easily be unleashed by slight redefinition. In particular, Proposition 16.3 demonstrates that $\tilde{T}_{xy}$ is the *magnitude* of the geometric mean of $\hat{H}_1$ and $\hat{H}_2$. We can thus justifiably define a **complex transmissibility** function as $\hat{H}_{gm}$...and the magnitude of this *complex transmissibility* function is the *ordinary transmissibility* function of Definition 16.1 (page 111).

REM

complex transmissibility $\tilde{T}'_{xy}(\omega) \triangleq \hat{H}_{gm}(\omega)$

Definition 16.6. *Let* **S** *be a* SYSTEM *with input* $x(n)$ *and output* $y(n)$.

DEF

The ***Harmonic mean transfer function estimate*** $\hat{H}_{hm}(\omega)$ *of* **S** *is defined as*

$$\hat{H}_{hm}(\omega) \triangleq \frac{2\tilde{S}_{yy}(\omega)\tilde{S}^*_{xy}(\omega)}{\tilde{S}_{xx}(\omega)\tilde{S}_{yy}(\omega) + |\tilde{S}_{xy}(\omega)|^2}$$

Proposition 16.4. [12] *Let* **S** *be a* SYSTEM *with input* $x(n)$ *and output* $y(n)$.

PRP

$$\hat{H}_{hm}(\omega) = \frac{2}{\frac{1}{\hat{H}_1(\omega)} + \frac{1}{\hat{H}_2(\omega)}} \qquad (\textit{Harmonic mean of } \hat{H}_1 \textit{ and } \hat{H}_2)$$

[12] Carne and Dohrmann (2006) $\langle H_C = [H_A^{-1} + H_B^{-1}]^{-1} \rangle$

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

✎PROOF:

$$\hat{H}_{hm}(\omega) \triangleq \frac{2\tilde{S}_{yy}(\omega)\tilde{S}^*_{xy}(\omega)}{\tilde{S}_{xx}(\omega)\tilde{S}_{yy}(\omega) + \left|\tilde{S}_{xy}(\omega)\right|^2} \quad \text{by definition of } \hat{H}_{hm} \quad \text{(Definition 16.6 page 114)}$$

$$= \frac{2}{\dfrac{\tilde{S}_{xx}(\omega)\tilde{S}_{yy}(\omega) + \left|\tilde{S}_{xy}(\omega)\right|^2}{\tilde{S}_{yy}(\omega)\tilde{S}^*_{xy}(\omega)}} = \frac{2}{\dfrac{\tilde{S}_{xx}(\omega)}{\tilde{S}^*_{xy}(\omega)} + \dfrac{\tilde{S}_{xy}(\omega)}{\tilde{S}_{yy}(\omega)}}$$

$$= \frac{2}{\dfrac{1}{\hat{H}_1(\omega)} + \dfrac{1}{\hat{H}_2(\omega)}} \quad \text{by definitions of } \hat{H}_1 \text{ and } \hat{H}_2 \quad \text{(Definition 16.2 page 111, Definition 16.3 page 112)}$$

$$= \textbf{Harmonic mean} \text{ of } \hat{H}_1(\omega) \text{ and } \hat{H}_2(\omega)$$

✏

A bit of review reveals $\hat{H}_1$ at the low end of the estimation problem, $\hat{H}_2$ at the high end, and $\hat{H}_{hm}$, $\hat{H}_{gm}$, and $\hat{H}_{am}$ somewhere between. But these three "between" estimates are not shown to be optimal in any sense—they are just conceptually interesting. What we might really like is a family of estimators that

- include $\hat{H}_1$ and $\hat{H}_2$ as limiting cases
- include the between cases
- are optimal in some sense

The estimator $\hat{H}_\kappa(\omega;\kappa)$ is one such estimator (next definition) that

- has $\hat{H}_1$ and $\hat{H}_2$ as limiting cases (Theorem 16.1 page 117),
- is optimal in the least squares sense (Theorem 16.6 page 130), and
- allows for a system designer to specify an output-input spectral noise ratio $\kappa(\omega)$ that can vary with frequency ω.

Moreover, $\hat{H}_\kappa(\omega)$ includes some special cases:

- In the case of constant κ, $\hat{H}_\kappa$ simplifies to the *Scaling transfer function estimate* $\hat{H}_s$ (Definition 16.8 page 115).
- In the case of $\kappa = 1$, $\hat{H}_\kappa$ and $\hat{H}_s$ simplify to the *Total least squares transfer function estimate* $\hat{H}_v$ (Definition 16.9 page 116).

Definition 16.7. [13] *Let* **S** *be a* SYSTEM *with input* $x(n)$ *and output* $y(n)$.

DEF *The* ***transfer function estimate*** $\hat{H}_\kappa(\omega;\kappa)$ *with* ***scaling function*** $\kappa(\omega)$ *is defined as*

$$\hat{H}_\kappa(\omega;\kappa) \triangleq \frac{\tilde{S}_{yy}(\omega) - \kappa(\omega)\tilde{S}_{xx}(\omega) + \sqrt{\left[\tilde{S}_{yy}(\omega) - \kappa(\omega)\tilde{S}_{xx}(\omega)\right]^2 + 4\kappa(\omega)\left|\tilde{S}_{xy}(\omega)\right|^2}}{2\tilde{S}_{xy}(\omega)}$$

Definition 16.8. [14] *Let* **S** *be a* SYSTEM *with input* $x(n)$ *and output* $y(n)$.

DEF *The* ***Scaling transfer function estimate*** $\hat{H}_s(\omega;s)$ *of* **S** *with* ***scaling parameter*** $s \in [0:\infty)$ *is defined as* $\hat{H}_s(\omega;s) \triangleq \hat{H}_\kappa(\omega;\kappa)$ *with* $\kappa(\omega) \triangleq s^2$

[13] White et al. (2006) page 679 ⟨(6)⟩, Shin and Hammond (2008) page 293 ⟨(9.67)⟩

[14] Shin and Hammond (2008) page 293 ⟨(9.67) with $\kappa(\omega) = s^2$⟩, White et al. (2006) page 679 ⟨(6) with $\kappa(\omega) = s^2$⟩, Leclere et al. (2014) ⟨(10) $\kappa(f) = 1/s^2$ and x and y swapped⟩, Wicks and Vold (1986) page 898 ⟨has additional s in denominator⟩, Zheng et al. (2002) page 918 ⟨(10), seems to differ⟩

COPYRIGHT © 2019 DANIEL J. GREENHOE

Definition 16.9. [15] *Let* $\mathbf{S}$ *be a* SYSTEM *with input* $\mathsf{x}(n)$ *and output* $\mathsf{y}(n)$.

DEF *The* ***Total Least Squares transfer function estimate*** $\hat{\mathsf{H}}_{\mathsf{v}}(\omega)$ *of* $\mathbf{S}$ *is defined as*

$$\hat{\mathsf{H}}_{\mathsf{v}}(\omega) \triangleq \hat{\mathsf{H}}_{\kappa}(\omega;\kappa) \qquad \text{with } \kappa(\omega) = 1$$

The previous estimators all assumed two signals: an input $\mathsf{x}(n)$ and an output $\mathsf{y}(n)$. However, in many practical systems, there is a third signal that is "driving" the system. In 1984 Goyder proposed an estimator (next definition) that is based on three signals.

Definition 16.10 (Three channel estimate)**.** [16] *Let* $\mathbf{S}$ *be a system with input* $\mathsf{x}(n)$*, output* $\mathsf{y}(n)$*, and a driver* $\mathsf{p}(n)$.

DEF *The* ***transfer function estimate*** $\hat{\mathsf{H}}_{\mathsf{c}}(\omega)$ *is defined as*

$$\hat{\mathsf{H}}_{\mathsf{c}}(\omega) \triangleq \frac{\tilde{\mathsf{S}}_{\mathsf{py}}(\omega)}{\tilde{\mathsf{S}}_{\mathsf{px}}(\omega)}$$

16.4 Estimator relationships

Lemma 16.1.

LEM

$$\frac{\mathrm{d}}{\mathrm{d}p}\left[\tilde{\mathsf{S}}_{\mathsf{yy}} - p\tilde{\mathsf{S}}_{\mathsf{xx}} + \sqrt{\left(\tilde{\mathsf{S}}_{\mathsf{yy}} - p\tilde{\mathsf{S}}_{\mathsf{xx}}\right)^2 + 4p\left|\tilde{\mathsf{S}}_{\mathsf{xy}}\right|^2}\right] = \frac{\tilde{\mathsf{S}}_{\mathsf{xx}}\left(p\tilde{\mathsf{S}}_{\mathsf{xx}}-\tilde{\mathsf{S}}_{\mathsf{yy}}\right)+2\left|\tilde{\mathsf{S}}_{\mathsf{xy}}\right|^2-\tilde{\mathsf{S}}_{\mathsf{xx}}\sqrt{\left(p\tilde{\mathsf{S}}_{\mathsf{xx}}-\tilde{\mathsf{S}}_{\mathsf{yy}}\right)^2+4p\left|\tilde{\mathsf{S}}_{\mathsf{xy}}\right|^2}}{\sqrt{\left(p\tilde{\mathsf{S}}_{\mathsf{xx}}-\tilde{\mathsf{S}}_{\mathsf{yy}}\right)^2+4p\left|\tilde{\mathsf{S}}_{\mathsf{xy}}\right|^2}}$$

$$\frac{\mathrm{d}}{\mathrm{d}p}\left[p\tilde{\mathsf{S}}_{\mathsf{yy}} - \tilde{\mathsf{S}}_{\mathsf{xx}} + \sqrt{\left(p\tilde{\mathsf{S}}_{\mathsf{yy}} - \tilde{\mathsf{S}}_{\mathsf{xx}}\right)^2 + 4p\left|\tilde{\mathsf{S}}_{\mathsf{xy}}\right|^2}\right] = \frac{\tilde{\mathsf{S}}_{\mathsf{yy}}\left(p\tilde{\mathsf{S}}_{\mathsf{yy}}-\tilde{\mathsf{S}}_{\mathsf{xx}}\right)+2\left|\tilde{\mathsf{S}}_{\mathsf{xy}}\right|^2+\tilde{\mathsf{S}}_{\mathsf{yy}}\sqrt{\left(p\tilde{\mathsf{S}}_{\mathsf{yy}}-\tilde{\mathsf{S}}_{\mathsf{xx}}\right)^2+4p\left|\tilde{\mathsf{S}}_{\mathsf{xy}}\right|^2}}{\sqrt{\left(p\tilde{\mathsf{S}}_{\mathsf{yy}}-\tilde{\mathsf{S}}_{\mathsf{xx}}\right)^2+4p\left|\tilde{\mathsf{S}}_{\mathsf{xy}}\right|^2}}$$

✎PROOF:

$$\frac{\mathrm{d}}{\mathrm{d}p}\left[\tilde{\mathsf{S}}_{\mathsf{yy}} - p\tilde{\mathsf{S}}_{\mathsf{xx}} + \sqrt{\left(\tilde{\mathsf{S}}_{\mathsf{yy}} - p\tilde{\mathsf{S}}_{\mathsf{xx}}\right)^2 + 4p\left|\tilde{\mathsf{S}}_{\mathsf{xy}}\right|^2}\right] = -\tilde{\mathsf{S}}_{\mathsf{xx}} + \frac{-2\tilde{\mathsf{S}}_{\mathsf{xx}}\left(\tilde{\mathsf{S}}_{\mathsf{yy}} - p\tilde{\mathsf{S}}_{\mathsf{xx}}\right) + 4\left|\tilde{\mathsf{S}}_{\mathsf{xy}}\right|^2}{2\sqrt{\left(\tilde{\mathsf{S}}_{\mathsf{yy}} - p\tilde{\mathsf{S}}_{\mathsf{xx}}\right)^2 + 4p\left|\tilde{\mathsf{S}}_{\mathsf{xy}}\right|^2}}$$

$$= \frac{4\left|\tilde{\mathsf{S}}_{\mathsf{xy}}\right|^2 - 2\tilde{\mathsf{S}}_{\mathsf{xx}}\left(\tilde{\mathsf{S}}_{\mathsf{yy}} - p\tilde{\mathsf{S}}_{\mathsf{xx}}\right) - 2\tilde{\mathsf{S}}_{\mathsf{xx}}\sqrt{\left(\tilde{\mathsf{S}}_{\mathsf{yy}} - p\tilde{\mathsf{S}}_{\mathsf{xx}}\right)^2 + 4p\left|\tilde{\mathsf{S}}_{\mathsf{xy}}\right|^2}}{2\sqrt{\left(\tilde{\mathsf{S}}_{\mathsf{yy}} - p\tilde{\mathsf{S}}_{\mathsf{xx}}\right)^2 + 4p\left|\tilde{\mathsf{S}}_{\mathsf{xy}}\right|^2}}$$

$$= \frac{\tilde{\mathsf{S}}_{\mathsf{xx}}\left(p\tilde{\mathsf{S}}_{\mathsf{xx}} - \tilde{\mathsf{S}}_{\mathsf{yy}}\right) + 2\left|\tilde{\mathsf{S}}_{\mathsf{xy}}\right|^2 - \tilde{\mathsf{S}}_{\mathsf{xx}}\sqrt{\left(p\tilde{\mathsf{S}}_{\mathsf{xx}} - \tilde{\mathsf{S}}_{\mathsf{yy}}\right)^2 + 4p\left|\tilde{\mathsf{S}}_{\mathsf{xy}}\right|^2}}{\sqrt{\left(p\tilde{\mathsf{S}}_{\mathsf{xx}} - \tilde{\mathsf{S}}_{\mathsf{yy}}\right)^2 + 4p\left|\tilde{\mathsf{S}}_{\mathsf{xy}}\right|^2}}$$

$$\frac{\mathrm{d}}{\mathrm{d}p}\left[p\tilde{\mathsf{S}}_{\mathsf{yy}} - \tilde{\mathsf{S}}_{\mathsf{xx}} + \sqrt{\left(p\tilde{\mathsf{S}}_{\mathsf{yy}} - \tilde{\mathsf{S}}_{\mathsf{xx}}\right)^2 + 4p\left|\tilde{\mathsf{S}}_{\mathsf{xy}}\right|^2}\right] = \tilde{\mathsf{S}}_{\mathsf{yy}} + \frac{2\tilde{\mathsf{S}}_{\mathsf{yy}}\left(p\tilde{\mathsf{S}}_{\mathsf{yy}} - \tilde{\mathsf{S}}_{\mathsf{xx}}\right) + 4\left|\tilde{\mathsf{S}}_{\mathsf{xy}}\right|^2}{2\sqrt{\left(p\tilde{\mathsf{S}}_{\mathsf{yy}} - \tilde{\mathsf{S}}_{\mathsf{xx}}\right)^2 + 4p\left|\tilde{\mathsf{S}}_{\mathsf{xy}}\right|^2}}$$

$$= \frac{4\left|\tilde{\mathsf{S}}_{\mathsf{xy}}\right|^2 + 2\tilde{\mathsf{S}}_{\mathsf{yy}}\left(p\tilde{\mathsf{S}}_{\mathsf{yy}} - \tilde{\mathsf{S}}_{\mathsf{xx}}\right) + 2\tilde{\mathsf{S}}_{\mathsf{yy}}\sqrt{\left(p\tilde{\mathsf{S}}_{\mathsf{yy}} - \tilde{\mathsf{S}}_{\mathsf{xx}}\right)^2 + 4p\left|\tilde{\mathsf{S}}_{\mathsf{xy}}\right|^2}}{2\sqrt{\left(p\tilde{\mathsf{S}}_{\mathsf{yy}} - \tilde{\mathsf{S}}_{\mathsf{xx}}\right)^2 + 4p\left|\tilde{\mathsf{S}}_{\mathsf{xy}}\right|^2}}$$

[15] White et al. (2006) page 679 ⟨(6)⟩, Shin and Hammond (2008) page 294 ⟨(9.69)⟩

[16] Shin and Hammond (2008) page 297 ⟨$H_3(f) = S_{ry}(f)/S_{rx}(f)$ (9.78)⟩, Cobb (1988) page 4 ⟨${}^c\hat{H}(f) = \hat{G}_{ys}(f)/\hat{G}_{xs}(f)$ (1.4)⟩, Goyder (1984) page 440 ⟨$H(i\omega) = S_{qz}/S_{pz}$ (5)⟩, Cobb and Mitchell (1990) page 450 ⟨(1)⟩, Pintelon and Schoukens (2012) page 241 ⟨$\hat{G}(\Omega_k) = \hat{G}_{ry}(\Omega_k)\hat{G}_{ru}^{-1}(\Omega_k)$ (7-49)⟩

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

$$= \frac{\tilde{S}_{yy}(p\tilde{S}_{yy} - \tilde{S}_{xx}) + 2|\tilde{S}_{xy}|^2 + \tilde{S}_{yy}\sqrt{(p\tilde{S}_{yy} - \tilde{S}_{xx})^2 + 4p|\tilde{S}_{xy}|^2}}{\sqrt{(p\tilde{S}_{yy} - \tilde{S}_{xx})^2 + 4p|\tilde{S}_{xy}|^2}}$$

Lemma 16.2.

LEM

$$\begin{aligned}
\tilde{S}_{yy} - p\tilde{S}_{xx} + \sqrt{(\tilde{S}_{yy} - p\tilde{S}_{xx})^2 + 4p|\tilde{S}_{xy}|^2} &\geq 0\\
p\tilde{S}_{yy} - \tilde{S}_{xx} + \sqrt{(p\tilde{S}_{yy} - \tilde{S}_{xx})^2 + 4p|\tilde{S}_{xy}|^2} &\geq 0
\end{aligned}$$

✎PROOF:

$$\begin{aligned}
& \tilde{S}_{yy} - p\tilde{S}_{xx} + \sqrt{(\tilde{S}_{yy} - p\tilde{S}_{xx})^2 + 4p|\tilde{S}_{xy}|^2} && \geq 0\\
\iff & \sqrt{(p\tilde{S}_{xx} - \tilde{S}_{yy})^2 + 4p|\tilde{S}_{xy}|^2} && \geq p\tilde{S}_{xx} - \tilde{S}_{yy}\\
\iff & (p\tilde{S}_{xx} - \tilde{S}_{yy})^2 + 4p|\tilde{S}_{xy}|^2 && \geq (p\tilde{S}_{xx} - \tilde{S}_{yy})^2\\
\iff & 4p|\tilde{S}_{xy}|^2 && \geq 0\\
\iff & |\tilde{S}_{xy}| && \geq 0
\end{aligned}$$

$$\begin{aligned}
& p\tilde{S}_{yy} - \tilde{S}_{xx} + \sqrt{(p\tilde{S}_{yy} - \tilde{S}_{xx})^2 + 4p|\tilde{S}_{xy}|^2} && \geq 0\\
\iff & \sqrt{(\tilde{S}_{xx} - p\tilde{S}_{yy})^2 + 4p|\tilde{S}_{xy}|^2} && \geq \tilde{S}_{xx} - p\tilde{S}_{yy}\\
\iff & (\tilde{S}_{xx} - p\tilde{S}_{yy})^2 + 4p|\tilde{S}_{xy}|^2 && \geq (\tilde{S}_{xx} - p\tilde{S}_{yy})^2\\
\iff & 4p|\tilde{S}_{xy}|^2 && \geq 0\\
\iff & |\tilde{S}_{xy}| && \geq 0
\end{aligned}$$



Figure 16.2: $\hat{H}_s$ with $\tilde{S}_{xx} = \tilde{S}_{yy} = 1$ and $\tilde{S}_{xy} = \frac{1}{2}$

Theorem 16.1. *Let $\hat{H}_s$ be defined as in Definition 16.8 (page 115).*

THM

$$\begin{aligned}
\{s_1 < s_2\} \implies & |\hat{H}_s(\omega; s_2)| \leq \hat{H}_s(\omega; s_1) && (\hat{H}_s \textit{ is } \textbf{\textit{monotonically decreasing}} \textit{ in } s)\\
|\hat{H}_1(\omega)| \leq & |\hat{H}_s(\omega; s)| \leq && |\hat{H}_2(\omega)|\\
\hat{H}_s(\omega; s)\big|_{s=0} = & && \hat{H}_2(\omega)\\
\hat{H}_s(\omega; s)\big|_{s=1} = & && \hat{H}_v(\omega)\\
\lim_{s\to\infty} \hat{H}_s(\omega; s) = & && \hat{H}_1(\omega)
\end{aligned}$$

COPYRIGHT © 2019 DANIEL J. GREENHOE

✎PROOF: I. Proofs for equalities:

$$\hat{\mathsf{H}}_{\mathsf{s}}(\omega;s)\big|_{s=0} \triangleq \left.\frac{\tilde{\mathsf{S}}_{\mathsf{yy}} - s^2\tilde{\mathsf{S}}_{\mathsf{xx}} + \sqrt{\left[\tilde{\mathsf{S}}_{\mathsf{yy}} - s^2\tilde{\mathsf{S}}_{\mathsf{xx}}\right]^2 + 4s^2\left|\tilde{\mathsf{S}}_{\mathsf{xy}}\right|^2}}{2\tilde{\mathsf{S}}_{\mathsf{xy}}}\right|_{s=0} \quad \text{by def. of } \hat{\mathsf{H}}_{\mathsf{s}} \quad \text{(Definition 16.8 page 115)}$$

$$= \frac{\tilde{\mathsf{S}}_{\mathsf{yy}} - 0 + \sqrt{\left[\tilde{\mathsf{S}}_{\mathsf{yy}} - 0\right]^2 + 0}}{2\tilde{\mathsf{S}}_{\mathsf{xy}}} = \frac{\tilde{\mathsf{S}}_{\mathsf{yy}}}{\tilde{\mathsf{S}}_{\mathsf{xy}}}$$

$$\triangleq \hat{\mathsf{H}}_2 \quad \text{by def. of } \hat{\mathsf{H}}_2 \quad \text{(Definition 16.3 page 112)}$$

$$\hat{\mathsf{H}}_{\mathsf{s}}(\omega;s)\big|_{s=1} \triangleq \left.\frac{\tilde{\mathsf{S}}_{\mathsf{yy}} - s^2\tilde{\mathsf{S}}_{\mathsf{xx}} + \sqrt{\left[\tilde{\mathsf{S}}_{\mathsf{yy}} - s^2\tilde{\mathsf{S}}_{\mathsf{xx}}\right]^2 + 4s^2\left|\tilde{\mathsf{S}}_{\mathsf{xy}}\right|^2}}{2\tilde{\mathsf{S}}_{\mathsf{xy}}}\right|_{s=1} \quad \text{by def. of } \hat{\mathsf{H}}_{\mathsf{s}} \quad \text{(Definition 16.8 page 115)}$$

$$= \frac{\tilde{\mathsf{S}}_{\mathsf{yy}} - \tilde{\mathsf{S}}_{\mathsf{xx}} + \sqrt{\left[\tilde{\mathsf{S}}_{\mathsf{yy}} - \tilde{\mathsf{S}}_{\mathsf{xx}}\right]^2 + 4\left|\tilde{\mathsf{S}}_{\mathsf{xy}}\right|^2}}{2\tilde{\mathsf{S}}_{\mathsf{xy}}}$$

$$\triangleq \hat{\mathsf{H}}_{\mathsf{v}} \quad \text{by def. of } \hat{\mathsf{H}}_{\mathsf{v}} \quad \text{(Definition 16.9 page 116)}$$

$$\lim_{s\to\infty} \hat{\mathsf{H}}_{\mathsf{s}}(\omega;s) \triangleq \lim_{s\to\infty} \frac{\tilde{\mathsf{S}}_{\mathsf{yy}} - s^2\tilde{\mathsf{S}}_{\mathsf{xx}} + \sqrt{\left[\tilde{\mathsf{S}}_{\mathsf{yy}} - s^2\tilde{\mathsf{S}}_{\mathsf{xx}}\right]^2 + 4s^2\left|\tilde{\mathsf{S}}_{\mathsf{xy}}\right|^2}}{2\tilde{\mathsf{S}}_{\mathsf{xy}}} \quad \text{by def. of } \hat{\mathsf{H}}_{\mathsf{s}} \quad \text{(Definition 16.8 page 115)}$$

$$\triangleq \lim_{p\to 0} \frac{\tilde{\mathsf{S}}_{\mathsf{yy}} - {}^1\!/_p\tilde{\mathsf{S}}_{\mathsf{xx}} + \sqrt{\left[\tilde{\mathsf{S}}_{\mathsf{yy}} - {}^1\!/_p\tilde{\mathsf{S}}_{\mathsf{xx}}\right]^2 + 4{}^1\!/_p\left|\tilde{\mathsf{S}}_{\mathsf{xy}}\right|^2}}{2\tilde{\mathsf{S}}_{\mathsf{xy}}} \quad \text{where } p \triangleq \frac{1}{s^2}$$

$$= \lim_{p\to 0} \frac{p\tilde{\mathsf{S}}_{\mathsf{yy}} - \tilde{\mathsf{S}}_{\mathsf{xx}} + \sqrt{\left[p\tilde{\mathsf{S}}_{\mathsf{yy}} - \tilde{\mathsf{S}}_{\mathsf{xx}}\right]^2 + 4p\left|\tilde{\mathsf{S}}_{\mathsf{xy}}\right|^2}}{2p\tilde{\mathsf{S}}_{\mathsf{xy}}} \quad \text{by mult. by } 1 = \frac{p}{p}$$

$$= \lim_{p\to 0} \frac{\frac{\mathrm{d}}{\mathrm{d}p}\left[p\tilde{\mathsf{S}}_{\mathsf{yy}} - \tilde{\mathsf{S}}_{\mathsf{xx}} + \sqrt{\left[p\tilde{\mathsf{S}}_{\mathsf{yy}} - \tilde{\mathsf{S}}_{\mathsf{xx}}\right]^2 + 4p\left|\tilde{\mathsf{S}}_{\mathsf{xy}}\right|^2}\right]}{\frac{\mathrm{d}}{\mathrm{d}p}\left[2p\tilde{\mathsf{S}}_{\mathsf{xy}}\right]} \quad \text{by } \textit{l'Hôpital's rule}$$

$$= \lim_{p\to 0} \frac{\dfrac{\tilde{\mathsf{S}}_{\mathsf{yy}}\left(p\tilde{\mathsf{S}}_{\mathsf{yy}} - \tilde{\mathsf{S}}_{\mathsf{xx}}\right) + 2\left|\tilde{\mathsf{S}}_{\mathsf{xy}}\right|^2 + \tilde{\mathsf{S}}_{\mathsf{yy}}\sqrt{\left(p\tilde{\mathsf{S}}_{\mathsf{yy}} - \tilde{\mathsf{S}}_{\mathsf{xx}}\right)^2 + 4p\left|\tilde{\mathsf{S}}_{\mathsf{xy}}\right|^2}}{\sqrt{\left(p\tilde{\mathsf{S}}_{\mathsf{yy}} - \tilde{\mathsf{S}}_{\mathsf{xx}}\right)^2 + 4p\left|\tilde{\mathsf{S}}_{\mathsf{xy}}\right|^2}}}{2\tilde{\mathsf{S}}_{\mathsf{xy}}} \quad \text{by Lemma 16.1 page 116}$$

$$= \frac{\tilde{\mathsf{S}}_{\mathsf{yy}}\left(-\tilde{\mathsf{S}}_{\mathsf{xx}}\right) + 2\left|\tilde{\mathsf{S}}_{\mathsf{xy}}\right|^2 + \tilde{\mathsf{S}}_{\mathsf{yy}}\sqrt{\left(-\tilde{\mathsf{S}}_{\mathsf{xx}}\right)^2}}{2\tilde{\mathsf{S}}_{\mathsf{xy}}\sqrt{\left(-\tilde{\mathsf{S}}_{\mathsf{xx}}\right)^2}}$$

$$= \frac{2\left|\tilde{\mathsf{S}}_{\mathsf{xy}}\right|^2}{2\tilde{\mathsf{S}}_{\mathsf{xx}}\tilde{\mathsf{S}}_{\mathsf{xy}}} = \frac{\tilde{\mathsf{S}}^*_{\mathsf{xy}}}{\tilde{\mathsf{S}}_{\mathsf{xx}}} \triangleq \hat{\mathsf{H}}_1 \quad \text{by def. of } \hat{\mathsf{H}}_1 \quad \text{(Definition 16.2 page 111)}$$

II. Proof for monoticity:

1. Let $p \triangleq s^2$
2. lemma:

$$\boxed{\left[2\left|\tilde{\mathsf{S}}_{\mathsf{xy}}\right|^2 + \tilde{\mathsf{S}}_{\mathsf{xx}}\left(p\tilde{\mathsf{S}}_{\mathsf{xx}} - \tilde{\mathsf{S}}_{\mathsf{yy}}\right)\right]^2}$$
$$= 4\left|\tilde{\mathsf{S}}_{\mathsf{xy}}\right|^4 + 4\tilde{\mathsf{S}}_{\mathsf{xx}}\left|\tilde{\mathsf{S}}_{\mathsf{xy}}\right|^2\left(p\tilde{\mathsf{S}}_{\mathsf{xx}} - \tilde{\mathsf{S}}_{\mathsf{yy}}\right) + \tilde{\mathsf{S}}^2_{\mathsf{xx}}\left(p\tilde{\mathsf{S}}_{\mathsf{xx}} - \tilde{\mathsf{S}}_{\mathsf{yy}}\right)^2$$

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

$$\boxed{\le} 4|\tilde{S}_{xy}|^2\tilde{S}_{xx}\tilde{S}_{yy} + 4p\tilde{S}_{xx}|\tilde{S}_{xy}|^2\tilde{S}_{xx} - 4\tilde{S}_{xx}|\tilde{S}_{xy}|^2\tilde{S}_{yy} + \tilde{S}_{xx}^2(p\tilde{S}_{xx} - \tilde{S}_{yy})^2 \quad \left(\begin{array}{l}\text{by } \textit{Cauchy Schwartz inequality} \\ \text{(Theorem J.2 page 256)}\end{array}\right)$$

$$= \cancelto{0}{4\tilde{S}_{xx}\tilde{S}_{yy}|\tilde{S}_{xy}|^2} + 4p\tilde{S}_{xx}|\tilde{S}_{xy}|^2\tilde{S}_{xx} - \cancelto{0}{4\tilde{S}_{xx}\tilde{S}_{yy}|\tilde{S}_{xy}|^2} + \tilde{S}_{xx}^2(p\tilde{S}_{xx} - \tilde{S}_{yy})^2$$

$$= \tilde{S}_{xx}^2\left[(p\tilde{S}_{xx} - \tilde{S}_{yy})^2 + 4p|\tilde{S}_{xy}|^2\right]$$

$$= \boxed{\left[\tilde{S}_{xx}\sqrt{(p\tilde{S}_{xx} - \tilde{S}_{yy})^2 + 4p|\tilde{S}_{xy}|^2}\right]^2}$$

3. lemma: $2|\tilde{S}_{xy}|^2 + \tilde{S}_{xx}(p\tilde{S}_{xx} - \tilde{S}_{yy}) \le \tilde{S}_{xx}\sqrt{(p\tilde{S}_{xx} - \tilde{S}_{yy})^2 + 4p|\tilde{S}_{xy}|^2}$. Proof:

$$2|\tilde{S}_{xy}|^2 + \tilde{S}_{xx}(p\tilde{S}_{xx} - \tilde{S}_{yy}) \le \tilde{S}_{xx}\sqrt{(p\tilde{S}_{xx} - \tilde{S}_{yy})^2 + 4p|\tilde{S}_{xy}|^2}$$

$$\iff \left[2|\tilde{S}_{xy}|^2 + \tilde{S}_{xx}(p\tilde{S}_{xx} - \tilde{S}_{yy})\right]^2 \le \left[\tilde{S}_{xx}\sqrt{(p\tilde{S}_{xx} - \tilde{S}_{yy})^2 + 4p|\tilde{S}_{xy}|^2}\right]^2 \quad \left(\begin{array}{l}\text{because } \mathsf{f}(x) \triangleq x^2 \text{ is} \\ \textit{strictly monotonic increasing}\end{array}\right)$$

The previous inequality is true by (2) lemma, so (3) lemma also true.

4. Proof that $\frac{\mathsf{d}}{\mathsf{d}p}|\hat{H}_s| \le 0$:

$$\frac{\mathsf{d}}{\mathsf{d}p}|\hat{H}_s| \triangleq \frac{\mathsf{d}}{\mathsf{d}p}\left|\frac{\tilde{S}_{yy} - s^2\tilde{S}_{xx} + \sqrt{(\tilde{S}_{yy} - s^2\tilde{S}_{xx})^2 + 4s^2|\tilde{S}_{xy}|^2}}{2\tilde{S}_{xy}}\right| \quad \text{by def. of } \hat{H}_s \text{ (Definition 16.8 page 115)}$$

$$\triangleq \frac{\mathsf{d}}{\mathsf{d}p}\left[\frac{\tilde{S}_{yy} - p\tilde{S}_{xx} + \sqrt{(\tilde{S}_{yy} - p\tilde{S}_{xx})^2 + 4p|\tilde{S}_{xy}|^2}}{2\tilde{S}_{xy}}\right] \quad \text{by definition of } p \text{ (item (1) page 118)}$$

$$= \frac{\mathsf{d}}{\mathsf{d}p}\left(\frac{\left|\tilde{S}_{yy} - p\tilde{S}_{xx} + \sqrt{(\tilde{S}_{yy} - p\tilde{S}_{xx})^2 + 4p|\tilde{S}_{xy}|^2}\right|}{|2\tilde{S}_{xy}|}\right)$$

$$= \frac{\mathsf{d}}{\mathsf{d}p}\left(\frac{\tilde{S}_{yy} - p\tilde{S}_{xx} + \sqrt{(\tilde{S}_{yy} - p\tilde{S}_{xx})^2 + 4p|\tilde{S}_{xy}|^2}}{2|\tilde{S}_{xy}|}\right) \quad \text{by Lemma 16.2 page 117}$$

$$= \frac{\dfrac{\tilde{S}_{xx}(p\tilde{S}_{xx} - \tilde{S}_{yy}) + 2|\tilde{S}_{xy}|^2 - \tilde{S}_{xx}\sqrt{(p\tilde{S}_{xx} - \tilde{S}_{yy})^2 + 4p|\tilde{S}_{xy}|^2}}{\sqrt{(\tilde{S}_{yy} - p\tilde{S}_{xx})^2 + 4p|\tilde{S}_{xy}|^2}}}{2|\tilde{S}_{xy}|} \quad \text{by Lemma 16.1 page 116}$$

$$= \frac{\tilde{S}_{xx}(p\tilde{S}_{xx} - \tilde{S}_{yy}) + 2|\tilde{S}_{xy}|^2 - \tilde{S}_{xx}\sqrt{(p\tilde{S}_{xx} - \tilde{S}_{yy})^2 + 4p|\tilde{S}_{xy}|^2}}{2|\tilde{S}_{xy}|\sqrt{(\tilde{S}_{yy} - p\tilde{S}_{xx})^2 + 4p|\tilde{S}_{xy}|^2}}$$

$$\le 0 \quad \text{by (3) lemma}$$

✏

Theorem 16.2. *Let* $\mathbf{S}$ *be a* SYSTEM *with input* $\mathsf{x}(n)$ *and output* $\mathsf{y}(n)$.

THM

$$|\hat{H}_1(\omega)| \le |\hat{H}_{hm}(\omega)| \le |\hat{H}_{gm}(\omega)| \le |\hat{H}_{am}(\omega)| \le |\hat{H}_2(\omega)|$$

✎PROOF:

COPYRIGHT © 2019 DANIEL J. GREENHOE
CC BY-NC-ND

1. lemma: $\hat{H}_1(\omega) \le \hat{H}_2(\omega)$. Proof:

$$
\begin{aligned}
\left|\hat{H}_1\right| &\triangleq \left|\frac{\tilde{S}_{yx}(\omega)}{\tilde{S}_{xx}(\omega)}\right| && \text{by definition of } \hat{H}_1 && \text{(Definition 16.2 page 111)}\\
&= \left|\frac{\langle y \mid x\rangle}{\|x\|^2}\right| = \frac{|\langle y \mid x\rangle|}{\|x\|^2} \\
&\le \frac{|\langle y \mid x\rangle|}{\|x\|^2}\left|\frac{\|x\|\,\|y\|}{\langle y \mid x\rangle}\right|^2 && \text{by } \textit{Cauchy Schwartz inequality} && \text{Theorem J.2 page 256}\\
&= \frac{\|y\|^2}{|\langle y \mid x\rangle|} = \left|\frac{\|y\|^2}{\langle x \mid y\rangle}\right| = \left|\frac{\tilde{S}_{yy}(\omega)}{\tilde{S}_{xy}(\omega)}\right| \\
&= \left|\hat{H}_2\right| && \text{by definition of } \hat{H}_2 && \text{(Definition 16.3 page 112)}
\end{aligned}
$$

2. remainder of the proof:

$$
\begin{aligned}
\left|\hat{H}_1(\omega)\right| &= \min\left\{\hat{H}_1(\omega), \hat{H}_2(\omega)\right\} && \text{by (1) lemma} && \\
&\le \left|\hat{H}_{hm}(\omega)\right| && \text{by Corollary M.1 page 295} && \text{with } \lambda_1 = \lambda_2 = ½\\
&\le \left|\hat{H}_{gm}(\omega)\right| && \text{by Corollary M.1 page 295} && \text{with } \lambda_1 = \lambda_2 = ½\\
&\le \left|\hat{H}_{am}(\omega)\right| && \text{by Corollary M.1 page 295} && \text{with } \lambda_1 = \lambda_2 = ½\\
&\le \max\left\{\hat{H}_1(\omega), \hat{H}_2(\omega)\right\} && \text{by Corollary M.1 page 295} && \text{with } \lambda_1 = \lambda_2 = ½\\
&= \left|\hat{H}_2(\omega)\right| && \text{by (1) lemma} &&
\end{aligned}
$$

✏

Theorem 16.2 (page 119) compared the magnitudes of several transfer function estimates and demonstrated a simple *linear* relationship. What about phase? The phase of those estimates is even simpler than the magnitude, as demonstrated next.

Proposition 16.5 (Estimator phase)**.** *Let $z \triangleq |z|e^{i\phi}$ be a* COMPLEX *number in the set of complex numbers $\mathbb{C}$. Let $\angle z \triangleq \phi$ be the* PHASE *of z.*

PRP

$$
\begin{aligned}
\angle\hat{H}_1(\omega) &= \angle\hat{H}_{hm}(\omega) = \angle\hat{H}_{gm}(\omega) = \angle\hat{H}_{am}(\omega) = \angle\hat{H}_2(\omega) = \angle\hat{H}_s(\omega) = \angle\hat{H}_v(\omega) = \angle\hat{H}_\kappa(\omega)\\
&= \angle C_{xy}(\omega)\\
&= -\angle\tilde{S}_{xy}(\omega)
\end{aligned}
$$

✎ PROOF:

COPYRIGHT © 2019 DANIEL J. GREENHOE

$$\angle\hat{H}_1 \triangleq \angle\frac{\tilde{S}_{yx}}{\tilde{S}_{xx}} \qquad = \qquad = \angle\tilde{S}^*_{xy} = -\angle\tilde{S}_{xy}$$

$\uparrow$ (Definition 16.2 page 111)

$$\angle\hat{H}_{hm} \triangleq \angle\frac{2\tilde{S}_{yy}\tilde{S}^*_{xy}}{\tilde{S}_{xx}\tilde{S}_{yy} + |\tilde{S}_{xy}|^2} \qquad = \qquad = \angle\tilde{S}^*_{xy} = -\angle\tilde{S}_{xy}$$

$\uparrow$ (Definition 16.6 page 114)

$$\angle\hat{H}_{gm} \triangleq \angle\frac{\tilde{S}^*_{xy}}{|\tilde{S}_{xy}|}\sqrt{\frac{\tilde{S}_{yy}}{\tilde{S}_{xx}}} \qquad = \qquad = \angle\tilde{S}^*_{xy} = -\angle\tilde{S}_{xy}$$

$\uparrow$ (Definition 16.5 page 113)

$$\angle\hat{H}_{am} \triangleq \angle\frac{|\tilde{S}_{xy}|^2 + \tilde{S}_{xx}\tilde{S}_{yy}}{2\tilde{S}_{xx}\tilde{S}_{xy}} \qquad = \angle\frac{1}{\tilde{S}_{xy}} = \angle\tilde{S}^*_{xy} = -\angle\tilde{S}_{xy}$$

$\uparrow$ (Definition 16.4 page 112)

$$\angle\hat{H}_2 \triangleq \angle\frac{\tilde{S}_{yy}}{\tilde{S}_{xy}} \qquad = \angle\frac{1}{\tilde{S}_{xy}} = \angle\tilde{S}^*_{xy} = -\angle\tilde{S}_{xy}$$

$\uparrow$ (Definition 16.3 page 112)

$$\angle\hat{H}_s \triangleq \angle\frac{\tilde{S}_{yy} - s^2\tilde{S}_{xx} + \sqrt{[\tilde{S}_{yy} - s^2\tilde{S}_{xx}]^2 + 4s^2|\tilde{S}_{xy}|^2}}{2\tilde{S}_{xy}} \qquad = \angle\frac{1}{\tilde{S}_{xy}} = \angle\tilde{S}^*_{xy} = -\angle\tilde{S}_{xy}$$

$\uparrow$ (Definition 16.8 page 115)

$$\angle\hat{H}_v \triangleq \angle\frac{\tilde{S}_{yy} - \tilde{S}_{xx} + \sqrt{[\tilde{S}_{yy} - \tilde{S}_{xx}]^2 + 4|\tilde{S}_{xy}|^2}}{2\tilde{S}_{xy}} \qquad = \angle\frac{1}{\tilde{S}_{xy}} = \angle\tilde{S}^*_{xy} = -\angle\tilde{S}_{xy}$$

$\uparrow$ (Definition 16.9 page 116)

$$\angle\hat{H}_\kappa \triangleq \angle\frac{\tilde{S}_{yy} - \kappa\tilde{S}_{xx} + \sqrt{[\tilde{S}_{yy} - \kappa\tilde{S}_{xx}]^2 + 4\kappa|\tilde{S}_{xy}|^2}}{2\tilde{S}_{xy}} \qquad = \angle\frac{1}{\tilde{S}_{xy}} = \angle\tilde{S}^*_{xy} = -\angle\tilde{S}_{xy}$$

$\uparrow$ (Definition 16.7 page 115)

$$\angle C_{xy} \triangleq \angle\frac{\tilde{S}^*_{xy}}{\sqrt{\tilde{S}_{xx}\tilde{S}_{yy}}} \qquad = \qquad = \angle\tilde{S}^*_{xy} = -\angle\tilde{S}_{xy}$$

$\uparrow$ (Definition 16.12 page 132)

16.5 Alternate forms

Any standard kit of algebraic tricks should arguably always include the ability to swap the location of a square root between numerator and denominator. If you are of this persuasion, after traveling from the definition of $\hat{H}_s$ on page 115, you won't be dissappointed when arriving at the next proposition (Proposition 16.6 page 121). But it has more use than just allowing you to entertain friends at social occasions. It also makes it very easy to see (using only algebra) what previously employed *l'Hôpital's rule* (using calculus) in the proof of Theorem 16.1—that $\lim_{s\to\infty}\hat{H}_s = \hat{H}_1$.

Proposition 16.6. [17] *Let $\hat{H}_\kappa(\omega;\kappa)$ be defined as in Definition 16.7 (page 115).*

[17] Shin and Hammond (2008) page 293 ⟨(9.67)⟩, Leclere et al. (2014) ⟨(10) $\kappa(f) = 1/s^2$ and x and y swapped⟩

Copyright © 2019 Daniel J. Greenhoe

PRP

$$\hat{H}_\kappa(\omega;s) = \frac{2\kappa(\omega)\tilde{S}_{yx}(\omega)}{\kappa(\omega)\tilde{S}_{xx}(\omega)-\tilde{S}_{yy}(\omega)+\sqrt{\left[\kappa(\omega)\tilde{S}_{xx}(\omega)-\tilde{S}_{yy}(\omega)\right]^2+4\kappa(\omega)\left|\tilde{S}_{xy}(\omega)\right|^2}}$$

$$= \frac{2\tilde{S}^*_{xy}}{\tilde{S}_{xx}-\frac{1}{\kappa(\omega)}\tilde{S}_{yy}+\sqrt{\left[\tilde{S}_{xx}-\frac{1}{\kappa(\omega)}\tilde{S}_{yy}\right]^2+\frac{4}{\kappa(\omega)}\left|\tilde{S}_{xy}\right|^2}}$$

✎PROOF: We can transform $\hat{H}_\kappa$ from that found in Definition 16.8 (page 115) to the forms in this proposition by the technique of "*rationalizing the denominator*"[18]

$$\hat{H}_\kappa \triangleq \frac{\tilde{S}_{yy}-\kappa\tilde{S}_{xx}+\sqrt{\left[\tilde{S}_{yy}-\kappa\tilde{S}_{xx}\right]^2+4\kappa\left|\tilde{S}_{xy}\right|^2}}{2\tilde{S}_{xy}} \qquad \text{by definition of } \hat{H}_\kappa \text{ (Definition 16.8 page 115)}$$

$$= \frac{\left[\tilde{S}_{yy}-\kappa\tilde{S}_{xx}+\sqrt{\left[\tilde{S}_{yy}-\kappa\tilde{S}_{xx}\right]^2+4\kappa\left|\tilde{S}_{xy}\right|^2}\right]\overbrace{\left[\tilde{S}_{yy}-\kappa\tilde{S}_{xx}-\sqrt{\left[\tilde{S}_{yy}-\kappa\tilde{S}_{xx}\right]^2+4\kappa\left|\tilde{S}_{xy}\right|^2}\right]}^{\text{"rationalizing factor"}}}{2\tilde{S}_{xy}\underbrace{\left[\tilde{S}_{yy}-\kappa\tilde{S}_{xx}-\sqrt{\left[\tilde{S}_{yy}-\kappa\tilde{S}_{xx}\right]^2+4\kappa\left|\tilde{S}_{xy}\right|^2}\right]}_{\text{"rationalizing factor"}}}$$

$$= \frac{\left[\tilde{S}_{yy}-\kappa\tilde{S}_{xx}\right]^2-\left[\tilde{S}_{yy}-\kappa\tilde{S}_{xx}\right]^2-4\kappa\left|\tilde{S}_{xy}\right|^2}{2\tilde{S}_{xy}\left[\tilde{S}_{yy}-\kappa\tilde{S}_{xx}-\sqrt{\left[\tilde{S}_{yy}-\kappa\tilde{S}_{xx}\right]^2+4\kappa\left|\tilde{S}_{xy}\right|^2}\right]} = \frac{-4\kappa\left|\tilde{S}_{xy}\right|^2}{2\tilde{S}_{xy}\left[\tilde{S}_{yy}-\kappa\tilde{S}_{xx}-\sqrt{\left[\tilde{S}_{yy}-\kappa\tilde{S}_{xx}\right]^2+4\kappa\left|\tilde{S}_{xy}\right|^2}\right]}$$

$$= \boxed{\frac{2\kappa\tilde{S}^*_{xy}}{\kappa\tilde{S}_{xx}-\tilde{S}_{yy}+\sqrt{\left[\kappa\tilde{S}_{xx}-\tilde{S}_{yy}\right]^2+4\kappa\left|\tilde{S}_{xy}\right|^2}}} = \frac{2\frac{\kappa}{\kappa}\tilde{S}^*_{xy}}{\frac{\kappa}{\kappa}\tilde{S}_{xx}-\frac{1}{\kappa}\tilde{S}_{yy}+\sqrt{\frac{1}{s^4}\left[\kappa\tilde{S}_{xx}-\tilde{S}_{yy}\right]^2+\frac{4\kappa}{s^4}\left|\tilde{S}_{xy}\right|^2}}$$

$$= \frac{2\tilde{S}^*_{xy}}{\tilde{S}_{xx}-\frac{1}{\kappa}\tilde{S}_{yy}+\sqrt{\left[\frac{\kappa}{\kappa}\tilde{S}_{xx}-\frac{1}{\kappa}\tilde{S}_{yy}\right]^2+\frac{4}{\kappa}\left|\tilde{S}_{xy}\right|^2}} = \boxed{\frac{2\tilde{S}^*_{xy}}{\tilde{S}_{xx}-\frac{1}{\kappa}\tilde{S}_{yy}+\sqrt{\left[\tilde{S}_{xx}-\frac{1}{\kappa}\tilde{S}_{yy}\right]^2+\frac{4}{\kappa}\left|\tilde{S}_{xy}\right|^2}}}$$

Integrity check for $s=0$ and $s\to\infty$ cases: Let $p\triangleq\kappa$.

$$\lim_{p\to\infty}\hat{H}_\kappa = \lim_{p\to\infty}\frac{2\tilde{S}_{yx}}{\tilde{S}_{xx}-\frac{1}{p}\tilde{S}_{yy}+\sqrt{\left[\tilde{S}_{xx}-\frac{1}{p}\tilde{S}_{yy}\right]^2+\frac{4}{p}\left|\tilde{S}_{xy}\right|^2}} = \frac{2\tilde{S}_{yx}}{\tilde{S}_{xx}+\sqrt{\left[\tilde{S}_{xx}\right]^2}}$$

$$= \frac{\tilde{S}_{yx}}{\tilde{S}_{xx}} \triangleq \hat{H}_1 \qquad \text{by def. of } \hat{H}_1 \text{ (Definition 16.2 page 111)}$$

$$\lim_{p\to 0}\hat{H}_\kappa = \lim_{p\to 0}\frac{2p\tilde{S}_{yx}}{p\tilde{S}_{xx}-\tilde{S}_{yy}+\sqrt{\left[p\tilde{S}_{xx}-\tilde{S}_{yy}\right]^2+4p\left|\tilde{S}_{xy}\right|^2}}$$

$$= \lim_{p\to 0}\frac{\frac{d}{dp}\left(2p\tilde{S}_{yx}\right)}{\frac{d}{dp}\left(p\tilde{S}_{xx}-\tilde{S}_{yy}+\sqrt{\left[p\tilde{S}_{xx}-\tilde{S}_{yy}\right]^2+4p\left|\tilde{S}_{xy}\right|^2}\right)} \qquad \text{by } \textit{l'Hôpital's rule}$$

[18] Slaught and Lennes (1915), page 274 ⟨"197. Rationalizing the Denominator."⟩ https://archive.org/details/elementaryalgebr00slaurich/page/274 Note that the operation in the proof of Proposition 16.6 is being performed essentially in reverse…or rather "rationalizing the numerator".

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

$$= \lim_{p\to 0} \frac{2\tilde{S}_{yx}}{\dfrac{\tilde{S}_{xx}\left(p\tilde{S}_{xx}-\tilde{S}_{yy}\right)+2\left|\tilde{S}_{xy}\right|^2+\tilde{S}_{xx}\sqrt{\left(p\tilde{S}_{xx}-\tilde{S}_{yy}\right)^2+4p\left|\tilde{S}_{xy}\right|^2}}{\sqrt{\left(p\tilde{S}_{xx}-\tilde{S}_{yy}\right)^2+4p\left|\tilde{S}_{xy}\right|^2}}} \qquad \text{by Lemma 16.1 page 116}$$

$$= \frac{2\tilde{S}_{yy}\tilde{S}_{yx}}{-\tilde{S}_{xx}\tilde{S}_{yy}+2\left|\tilde{S}_{xy}\right|^2+\tilde{S}_{xx}\tilde{S}_{yy}} = \frac{2\tilde{S}_{yy}\tilde{S}_{yx}}{2\left|\tilde{S}_{xy}\right|^2} = \frac{\tilde{S}_{yy}}{\tilde{S}_{xy}}$$

$$\triangleq \hat{H}_2 \qquad \text{by def. of } \hat{H}_2 \text{ (Definition 16.3 page 112)}$$

16.6 Least squares estimates of non-linear systems

> *"The legendary Hungarian mathematician John von Neumann once referred to the the theory of nonequilibrium systems as the "theory of non-elephants," ...Nevertheless, such a theory of non-elephants will be attempted here."*
>
> Per Bak, in *"how nature works..."* [19]



(A) *Least squares estimation* for *communications additive noise model* (Corollary 16.1 page 126)

(B) *Least squares estimations* for *measurement additive noise model* (Corollary 16.2 page 127)

Figure 16.3: Least Square estimation (Theorem 16.3 page 124)

Let **S** be the *system* illustrated to the right. **If** there is no measurement noise on the input and output and **if H** is *linear time invariant*, then $\tilde{H} = \tilde{S}_{xy}/\tilde{S}_{xx}$ (Corollary 5.1 page 36). But what if there is output measurement noise? And what if **H** is not *LTI*? What is the best least-squares estimate of $\tilde{H}$? The answer depends on how you define "the best".



The defintion of "best" or "optimal" is given by a *cost function* $C(\hat{H})$. There are several possible cost functions. Definition 16.11 provides some. Theorem 16.3 then demonstrate optimal solutions with respect to these definitions.

Definition 16.11. *Let* **S** *be a system defined as in Figure 16.3 (page 123) (A) or (B). Define the following* COST FUNCTIONS *for spectral* LEAST-SQUARES *estimates:*

DEF

$$C_{rq}(\hat{G}) \triangleq \tilde{F}\,\|r(n)-q(n)\|^2 \triangleq \tilde{F}\,\langle r(n)-q(n)\,|\,r(0)-q(0)\rangle \triangleq \tilde{F}E\left(\left[r(n)-q(n)\right]\left[r(0)-q(0)\right]^*\right)$$

$$C_{sy}(\hat{H}) \triangleq \tilde{F}\,\|s(n)-y(n)\|^2 \triangleq \tilde{F}\,\langle s(n)-y(n)\,|\,s(0)-y(0)\rangle \triangleq \tilde{F}E\left(\left[s(n)-y(n)\right]\left[s(0)-y(0)\right]^*\right)$$

[19] Bak (2013) page 29 ⟨§ Systems in Balance Are Not Complex⟩

COPYRIGHT © 2019 DANIEL J. GREENHOE

Lemma 16.3. *Let* $C_{rq}(\hat{G})$ *and* $C_{sy}(\hat{H})$ *be defined as in Definition 16.11 (page 123).*

LEM

$$C_{rq}(\hat{G}) = \tilde{S}_{pp}(\omega)\left|\hat{G}(\omega)\right|^2 - \tilde{S}_{py}(\omega)\hat{G}(\omega) - \tilde{S}^*_{py}(\omega)\hat{G}^*(\omega) + \tilde{S}_{qq}(\omega)$$
$$C_{sy}(\hat{H}) = \tilde{S}_{xx}(\omega)\left|\hat{H}(\omega)\right|^2 - \tilde{S}_{xy}(\omega)\hat{H}(\omega) - \tilde{S}^*_{xy}(\omega)\hat{H}^*(\omega) + \tilde{S}_{yy}(\omega)$$

✎PROOF:

$\boxed{C_{rq}(\hat{G})}$

$$\begin{aligned}
&\triangleq \breve{\mathbf{F}}\mathsf{E}\Big(\big[r(n)-q(n)\big]\big[r(0)-q(0)\big]^*\Big) && \text{by definition of } C_{rq} && \text{(Definition 16.11 page 123)}\\
&= \tilde{\mathbf{F}}\big[\mathsf{E}\big[r(n)r^*(0)\big] - \mathsf{E}\big[r(n)q^*(0)\big] - \mathsf{E}\big[q(n)r^*(0)\big] + \mathsf{E}\big[q(n)q^*(0)\big]\big] && \text{by } \textit{linearity} \text{ of } \mathsf{E} && \text{(Theorem 1.1 page 4)}\\
&\triangleq \tilde{\mathbf{F}}\big[R_{rr}(m) - R_{rq}(m) - R_{qr}(m) + R_{qq}(m)\big] && \text{by definition of } R_{xy} && \text{(Definition 2.4 page 12)}\\
&\triangleq \big[\tilde{S}_{rr}(\omega) - \tilde{S}_{rq}(\omega) - \tilde{S}_{qr}(\omega) + \tilde{S}_{qq}(\omega)\big] && \text{by definition of } \tilde{S}_{xy} && \text{(Definition 6.3 page 44)}\\
&= \boxed{\tilde{S}_{pp}(\omega)\left|\hat{G}(\omega)\right|^2 - \tilde{S}_{py}(\omega)\hat{G}(\omega) - \tilde{S}^*_{py}(\omega)\hat{G}^*(\omega) + \tilde{S}_{qq}(\omega)} && \text{by (A)–(D) and} && \text{Corollary 7.8 page 59}
\end{aligned}$$

$\boxed{C_{sy}(\hat{H})}$

$$\begin{aligned}
&\triangleq \breve{\mathbf{F}}\mathsf{E}\Big(\big[s(n)-y(n)\big]\big[s(0)-y(0)\big]^*\Big) && \text{by definition of } C_{sy} && \text{(Definition 16.11 page 123)}\\
&= \tilde{\mathbf{F}}\big[\mathsf{E}\big[s(n)s^*(0)\big] - \mathsf{E}\big[s(n)y^*(0)\big] - \mathsf{E}\big[y(n)s^*(0)\big] + \mathsf{E}\big[y(n)y^*(0)\big]\big] && \text{by } \textit{linearity} \text{ of } \mathsf{E} && \text{(Theorem 1.1 page 4)}\\
&\triangleq \tilde{\mathbf{F}}\big[R_{ss}(m) - R_{sy}(m) - R_{ys}(m) + R_{yy}(m)\big] && \text{by definition of } R_{xy} && \text{(Definition 2.4 page 12)}\\
&\triangleq \big[\tilde{S}_{ss}(\omega) - \tilde{S}_{sy}(\omega) - \tilde{S}_{ys}(\omega) + \tilde{S}_{yy}(\omega)\big] && \text{by definition of } \tilde{S}_{xy} && \text{(Definition 6.3 page 44)}\\
&= \boxed{\tilde{S}_{xx}(\omega)\left|\hat{H}(\omega)\right|^2 - \tilde{S}_{xy}(\omega)\hat{H}(\omega) - \tilde{S}^*_{xy}(\omega)\hat{H}^*(\omega) + \tilde{S}_{yy}(\omega)} && \text{by (A)–(D) and} && \text{Corollary 7.8 (page 59)}
\end{aligned}$$

✏

Theorem 16.3. *Let* **S** *be the* SYSTEM *illustrated in Figure 16.3 page 123 (A) or (B).*

THM

$$\left.\begin{array}{lll}
(A). & x, u, \textit{and } v \textit{ are} & \text{WSS} \quad \textit{and}\\
(B). & x, u, \textit{and } v \textit{ are} & \text{UNCORRELATED} \quad \textit{and}\\
(C). & Eu = Ev = 0 & (\text{ZERO-MEAN}) \quad \textit{and}\\
(D). & \hat{G}_{ls} \textit{ and } \hat{H}_{ls} \textit{ are} & \text{LTI}
\end{array}\right\} \Longrightarrow \left\{\begin{array}{lll}
(1). & \arg\min\limits_{\hat{G}} C_{rq}(\hat{G}) & = \dfrac{\tilde{S}_{yp}(\omega)}{\tilde{S}_{pp}(\omega)}\\[2ex]
(2). & \arg\min\limits_{\hat{H}} C_{sy}(\hat{H}) & = \dfrac{\tilde{S}_{yx}(\omega)}{\tilde{S}_{xx}(\omega)}
\end{array}\right.$$

✎PROOF:

1. Define shorthand function names $\hat{G} \triangleq \hat{G}_{ls}$ and $\hat{H} \triangleq \hat{H}_{ls}$.
2. lemma:

$$\begin{aligned}
0 &= \frac{\partial}{\partial \hat{G}_R} C_{rq}(\hat{G})\\
&= \frac{\partial}{\partial \hat{G}_R}\Big(\tilde{S}_{pp}\left|\hat{G}\right|^2 - \hat{G}\tilde{S}_{py} - \hat{G}^*\tilde{S}^*_{py} + \tilde{S}_{qq}\Big) && \text{by Lemma 16.3 page 124}\\
&= \frac{\partial}{\partial \hat{G}_R}\big(\tilde{S}_{pp}\big[\hat{G}_R^2 + \hat{G}_I^2\big] - (\hat{G}_R + i\hat{G}_I)\tilde{S}_{py} - (\hat{G}_R + i\hat{G}_I)^*\tilde{S}^*_{py} + \tilde{S}_{qq}\big)\\
&= 2\hat{G}_R\tilde{S}_{pp} - \tilde{S}_{py} - \tilde{S}^*_{py} + \cancelto{0}{\frac{\partial}{\partial \hat{G}_R}\tilde{S}_{qq}} && \text{because q does not vary with } \hat{G}\\
&= 2\hat{G}_R\tilde{S}_{pp} - 2\mathbf{R}_e\tilde{S}_{py}\\
&= 2\hat{G}_R\tilde{S}_{pp} - 2\mathbf{R}_e\tilde{S}_{yp} && \text{by Corollary 2.2 page 15}\\
&\Longrightarrow \boxed{\hat{G}_R(\omega) = \frac{\mathbf{R}_e\tilde{S}_{yp}(\omega)}{\tilde{S}_{pp}(\omega)}}
\end{aligned}$$

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

3. lemma:

$$
\begin{aligned}
0 &= \frac{\partial}{\partial \hat{G}_I} C_{rq}(\hat{G}) \\
&= \frac{\partial}{\partial \hat{G}_I}\left(\tilde{S}_{pp}|\hat{G}|^2 - \hat{G}\tilde{S}_{py} - \hat{G}^*\tilde{S}_{py}^* + \tilde{S}_{qq}\right) && \text{by Lemma 16.3 page 124} \\
&= \frac{\partial}{\partial \hat{G}_I}\left[\tilde{S}_{pp}\left[\hat{G}_R^2 + \hat{G}_I^2\right] - (\hat{G}_R + i\hat{G}_I)\tilde{S}_{py} - (\hat{G}_R - i\hat{G}_I)\tilde{S}_{py}^* + \tilde{S}_{qq}\right] \\
&= 2\hat{G}_I\tilde{S}_{pp} - i\tilde{S}_{py} + i\tilde{S}_{py}^* + \cancelto{0}{\frac{\partial}{\partial \hat{G}_I}\tilde{S}_{qq}} && \text{because q does not vary with } \hat{G} \\
&= 2\hat{G}_I\tilde{S}_{pp} - 2i(i\mathbf{I}_m\tilde{S}_{py}) \\
&= 2\hat{G}_I\tilde{S}_{pp} + 2i(i\mathbf{I}_m\tilde{S}_{yp}) && \text{by Corollary 2.2 page 15} \\
&\Longrightarrow \boxed{\hat{G}_I(\omega) = \frac{\mathbf{I}_m\tilde{S}_{yp}(\omega)}{\tilde{S}_{pp}(\omega)}}
\end{aligned}
$$

4. Proof for $\hat{G} \triangleq \hat{G}_{ls}$ expression:

$$
\begin{aligned}
\boxed{\hat{G}(\omega)} &= \hat{G}_R(\omega) + i\hat{G}_I(\omega) \\
&= \frac{\mathbf{R}_e\tilde{S}_{yp}(\omega)}{\tilde{S}_{pp}(\omega)} + \frac{i\mathbf{I}_m\tilde{S}_{yp}(\omega)}{\tilde{S}_{pp}(\omega)} && \text{by (2) lemma and (3) lemma} \\
&= \boxed{\frac{\tilde{S}_{yp}(\omega)}{\tilde{S}_{pp}(\omega)}}
\end{aligned}
$$

5. lemma:

$$
\begin{aligned}
0 &= \frac{\partial}{\partial \hat{H}_R} C_{sy}(\hat{H}) \\
&= \frac{\partial}{\partial \hat{H}_R}\left(\tilde{S}_{xx}|\hat{H}|^2 - \hat{H}\tilde{S}_{xy} - \hat{H}^*\tilde{S}_{xy}^* + \tilde{S}_{yy}\right) && \text{by Lemma 16.3 page 124} \\
&= \frac{\partial}{\partial \hat{H}_R}\left(\tilde{S}_{xx}\left[\hat{H}_R^2 + \hat{H}_I^2\right] - (\hat{H}_R + i\hat{H}_I)\tilde{S}_{xy} - (\hat{H}_R + i\hat{H}_I)^*\tilde{S}_{xy}^* + \tilde{S}_{yy}\right) \\
&= 2\hat{H}_R\tilde{S}_{xx} - \tilde{S}_{xy} - \tilde{S}_{xy}^* + \cancelto{0}{\frac{\partial}{\partial \hat{H}_R}\tilde{S}_{yy}} && \text{because y does not vary with } \hat{H} \\
&= 2\hat{H}_R\tilde{S}_{xx} - 2\mathbf{R}_e\tilde{S}_{xy} \\
&= 2\hat{H}_R\tilde{S}_{xx} - 2\mathbf{R}_e\tilde{S}_{yx} && \text{by Corollary 2.2 page 15} \\
&\Longrightarrow \boxed{\hat{H}_R(\omega) = \frac{\mathbf{R}_e\tilde{S}_{yx}(\omega)}{\tilde{S}_{xx}(\omega)}}
\end{aligned}
$$

6. lemma:

$$
\begin{aligned}
0 &= \frac{\partial}{\partial \hat{H}_I} C_{sy}(\hat{H}) \\
&= \frac{\partial}{\partial \hat{H}_I}\left(\tilde{S}_{xx}|\hat{H}|^2 - \tilde{S}_{xy}\hat{H} - \tilde{S}_{xy}^*\hat{H}^* + \tilde{S}_{yy}\right) && \text{by Lemma 16.3 page 124} \\
&= \frac{\partial}{\partial \hat{H}_I}\left[\tilde{S}_{xx}\left[\hat{H}_R^2 + \hat{H}_I^2\right] - \tilde{S}_{xy}(\hat{H}_R + i\hat{H}_I) - \tilde{S}_{xy}^*(\hat{H}_R - i\hat{H}_I) + \tilde{S}_{yy}\right] \\
&= 2\hat{H}_I\tilde{S}_{xx} - i\tilde{S}_{xy} + i\tilde{S}_{xy}^* + \cancelto{0}{\frac{\partial}{\partial \hat{H}_R}\tilde{S}_{yy}} && \text{because q does not vary with } \hat{H}
\end{aligned}
$$

Copyright © 2019 Daniel J. Greenhoe
CC BY-NC-ND

$$= 2\hat{H}_I\tilde{S}_{xx} - 2i(i\mathbf{I}_m\tilde{S}_{xy})$$
$$= 2\hat{H}_I\tilde{S}_{xx} + 2i(i\mathbf{I}_m\tilde{S}_{yx}) \qquad \text{by Corollary 2.2 page 15}$$
$$= 2\hat{H}_I\tilde{S}_{xx} - 2\mathbf{I}_m\tilde{S}_{yx}$$
$$\Longrightarrow \boxed{\hat{H}_I(\omega) = \frac{\mathbf{I}_m\tilde{S}_{yx}(\omega)}{\tilde{S}_{xx}(\omega)}}$$

7. Proof for $\hat{H} \triangleq \hat{H}_{ls}$ expression:

$$\boxed{\hat{H}(\omega)} = \hat{H}_R(\omega) + i\hat{H}_I(\omega)$$
$$= \frac{\mathbf{R}_e\tilde{S}_{yp}(\omega)}{\tilde{S}_{xx}(\omega)} + \frac{i\mathbf{I}_m\tilde{S}_{yp}(\omega)}{\tilde{S}_{xx}(\omega)} \qquad \text{by (5) lemma and (6) lemma}$$
$$= \frac{\mathbf{R}_e\tilde{S}_{yx}(\omega)}{\tilde{S}_{xx}(\omega)} + \frac{i\mathbf{I}_m\tilde{S}_{yx}(\omega)}{\tilde{S}_{xx}(\omega)} \qquad \text{by Theorem 7.4 page 55}$$
$$= \boxed{\frac{\tilde{S}_{yx}(\omega)}{\tilde{S}_{xx}(\omega)}}$$

✏

Using Theorem 16.3 (previous) we can see that the optimal **least-squares** operators $\hat{G}_{ls}$ and $\hat{H}_{ls}$ for the **non-linear** operator $\mathbf{T}$ in Figure 16.3 (page 123) (A) and (B) are (next two corollaries)

(1).	$\hat{G}_{ls}(\omega)$	$= \dfrac{\tilde{S}_{yx}(\omega)}{\tilde{S}_{xx}(\omega) + \tilde{S}_{uu}(\omega)}$	for (A)—*communication system*
(2).	$\hat{G}_{ls}(\omega)$	$= \dfrac{\tilde{S}_{yx}(\omega)}{\tilde{S}_{xx}(\omega) - \tilde{S}_{uu}(\omega)}$	for (B)—*measurement system*
(3).	$\hat{H}_{ls}(\omega)$	$= \dfrac{\tilde{S}_{yx}(\omega)}{\tilde{S}_{xx}(\omega)}$	for either (A) or (B)

Corollary 16.1. *Let* $\mathbf{S}$ *be the* SYSTEM *illustrated in Figure 16.3 page 123 (A).*

THM

$$\left\{ \begin{array}{r} \textit{hypotheses of Theorem 16.3} \\ \textit{page 124} \end{array} \right\} \Longrightarrow \left\{ \begin{array}{lll} (1). & \arg\min\limits_{\hat{G}_{ls}} C_{rq}(\hat{G}_{ls}) & = \dfrac{\tilde{S}_{yx}(\omega)}{\tilde{S}_{xx}(\omega) + \tilde{S}_{uu}(\omega)} \\ (2). & \arg\min\limits_{\hat{H}_{ls}} C_{sy}(\hat{H}_{ls}) & = \dfrac{\tilde{S}_{yx}(\omega)}{\tilde{S}_{xx}(\omega)} \end{array} \right\}$$

✎ PROOF:

$$\hat{G}_{ls} = \frac{\tilde{S}_{yx}(\omega)}{\tilde{S}_{pp}(\omega)} \qquad \text{by Theorem 16.3 page 124}$$
$$= \frac{\tilde{S}_{yx}(\omega)}{\tilde{S}_{xx}(\omega) + \tilde{S}_{uu}(\omega)} \qquad \text{by Theorem 7.1 page 51}$$
$$\hat{H}_{ls} = \frac{\tilde{S}_{yx}(\omega)}{\tilde{S}_{xx}(\omega)} \qquad \text{by Theorem 16.3 page 124}$$

✏

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE



(A) **communications LTI additive noise model** (B) **measurement LTI additive noise model**

Figure 16.4: Additive noise systems with **LTI** operator **H**

Corollary 16.2. *Let* **S** *be the* SYSTEM *illustrated in Figure 16.3 page 123 (B).*

THM

$$\left\{ \begin{array}{r} \textit{hypotheses of Theorem 16.3} \\ \textit{page 124} \end{array} \right\} \Longrightarrow \left\{ \begin{array}{lll} (1). & \arg\min\limits_{\hat{G}_{ls}} C_{rq}(\hat{G}_{ls}) & = \dfrac{\tilde{S}_{yx}(\omega)}{\tilde{S}_{xx}(\omega) - \tilde{S}_{uu}(\omega)} \\ (2). & \arg\min\limits_{\hat{H}_{ls}} C_{sy}(\hat{H}_{ls}) & = \dfrac{\tilde{S}_{yx}(\omega)}{\tilde{S}_{xx}(\omega)} \end{array} \right\}$$

✎PROOF:

$$\begin{aligned} \hat{G} &= \frac{\tilde{S}_{yx}(\omega)}{\tilde{S}_{pp}(\omega)} && \text{by Theorem 16.3 page 124} \\ &= \frac{\tilde{S}_{yx}(\omega)}{\tilde{S}_{xx}(\omega) - \tilde{S}_{uu}(\omega)} && \text{by Theorem 7.1 page 51} \\ \hat{H} &= \frac{\tilde{S}_{yx}(\omega)}{\tilde{S}_{xx}(\omega)} && \text{by Theorem 16.3 page 124} \end{aligned}$$

✎

It follows immediately from Corollary 16.1 (page 126) and Corollary 16.2 (page 127) that, in the special case of no input noise ($u(n) = 0$), $\hat{H}_1$ is the optimal least-squares estimate of $\tilde{H}$ (next corollary).

Corollary 16.3. [20] *Let* **S** *be the* SYSTEM *illustrated in Figure 16.3 page 123 (A) or (B).*

COR

$$\left\{ \begin{array}{ll} (1). & \textit{hypotheses of Theorem 16.3} \quad \textit{and} \\ (2). & u(n) = 0 \end{array} \right\} \Longrightarrow \left\{ \hat{G}_{ls}(\omega) = \hat{H}_{ls}(\omega) = \hat{H}_1(\omega) \right\}$$

16.7 Least squares estimates of linear systems

The previous section did assume the estimates $\hat{H}_1$ and $\hat{H}_2$ to be *linear time invariant* (*LTI*), but it did *not* assume that the system transfer function **T** itself to be *LTI*. But making the LTI assumption on **H** yields some interesting and insightful results, such as those in this section.

Theorem 16.4 (Estimating H in communication additive noise system)**.** *Let* **S** *be the* SYSTEM *illustrated in Figure 16.4 page 127 (A).*

[20] Bendat and Piersol (1980) pages 98–100 ⟨5.1.1 Optimal Character of Calculations; note: proof minimizing $\tilde{S}_{vv}$ but yields same result⟩, Bendat and Piersol (1993) pages 106–109 ⟨5.1.1 Optimality of Calculations⟩, Bendat and Piersol (2010) pages 187–190 ⟨6.1.4 Optimum Frequency Response Functions⟩

COPYRIGHT © 2019 DANIEL J. GREENHOE | https://github.com/dgreenhoe/pdfs/blob/master/abcstat.pdf | CC BY-NC-ND

THM

$$\left\{\begin{array}{lll} (A). & \mathbf{H} \textit{ is} & \text{LINEAR TIME INVARIANT } (\text{LTI}) \textit{ and} \\ (B). & \mathsf{x}(n) \textit{ is} & \text{WIDE-SENSE STATIONARY } (\text{WSS}) \textit{ and} \\ (C). & \mathsf{x}(n), \mathsf{u}(n), \textit{and } \mathsf{v}(n) \textit{ are} & \text{UNCORRELATED} \end{array}\right\}$$

$$\Longrightarrow \left\{\begin{array}{lll} (1). & \hat{\mathsf{H}}_1(\omega) = \tilde{\mathsf{H}}(\omega) & \textit{and} \\ (2). & \hat{\mathsf{H}}_2(\omega) = \dfrac{\tilde{\mathsf{S}}_{\mathsf{vv}}(\omega)}{\tilde{\mathsf{H}}^*(\omega)\tilde{\mathsf{S}}_{\mathsf{xx}}(\omega)} + \tilde{\mathsf{H}}(\omega)\left[1 + \dfrac{\tilde{\mathsf{S}}_{\mathsf{uu}}(\omega)}{\tilde{\mathsf{S}}_{\mathsf{xx}}(\omega)}\right] & \end{array}\right\}$$

✎PROOF:

$$\begin{aligned}
\hat{\mathsf{H}}_1(\omega) &\triangleq \frac{\tilde{\mathsf{S}}_{\mathsf{yx}}(\omega)}{\tilde{\mathsf{S}}_{\mathsf{xx}}(\omega)} && \text{by definition of } \hat{\mathsf{H}}_1 && \text{(Definition 16.2 page 111)} \\
&= \frac{\tilde{\mathsf{H}}(\omega)\tilde{\mathsf{S}}_{\mathsf{xx}}(\omega)}{\tilde{\mathsf{S}}_{\mathsf{xx}}(\omega)} && \text{by Corollary 7.5 page 57} \\
&= \tilde{\mathsf{H}}(\omega) \\
\hat{\mathsf{H}}_2(\omega) &\triangleq \frac{\tilde{\mathsf{S}}_{\mathsf{yy}}(\omega)}{\tilde{\mathsf{S}}_{\mathsf{xy}}(\omega)} && \text{by definition of } \hat{\mathsf{H}}_2 && \text{(Definition 16.3 page 112)} \\
&= \frac{\tilde{\mathsf{S}}_{\mathsf{yy}}(\omega)}{\tilde{\mathsf{H}}^*(\omega)\tilde{\mathsf{S}}_{\mathsf{xx}}(\omega)} && \text{by Corollary 7.5 page 57} \\
&= \frac{\tilde{\mathsf{S}}_{\mathsf{vv}}(\omega) + \tilde{\mathsf{S}}_{\mathsf{qq}}(\omega)}{\tilde{\mathsf{H}}^*(\omega)\tilde{\mathsf{S}}_{\mathsf{xx}}(\omega)} && \text{by Theorem 7.1 page 51} \\
&= \frac{\tilde{\mathsf{S}}_{\mathsf{vv}}(\omega) + \tilde{\mathsf{H}}^*(\omega)\tilde{\mathsf{H}}(\omega)\tilde{\mathsf{S}}_{\mathsf{pp}}(\omega)}{\tilde{\mathsf{H}}^*(\omega)\tilde{\mathsf{S}}_{\mathsf{xx}}(\omega)} && \text{by Corollary 5.3 page 37} \\
&= \frac{\tilde{\mathsf{S}}_{\mathsf{vv}}(\omega)}{\tilde{\mathsf{H}}^*(\omega)\tilde{\mathsf{S}}_{\mathsf{xx}}(\omega)} + \frac{\tilde{\mathsf{H}}^*(\omega)\tilde{\mathsf{H}}(\omega)\left[\tilde{\mathsf{S}}_{\mathsf{xx}}(\omega) + \tilde{\mathsf{S}}_{\mathsf{uu}}(\omega)\right]}{\tilde{\mathsf{H}}^*(\omega)\tilde{\mathsf{S}}_{\mathsf{xx}}(\omega)} \\
&= \frac{\tilde{\mathsf{S}}_{\mathsf{vv}}(\omega)}{\tilde{\mathsf{H}}^*(\omega)\tilde{\mathsf{S}}_{\mathsf{xx}}(\omega)} + \tilde{\mathsf{H}}(\omega)\left[1 + \frac{\tilde{\mathsf{S}}_{\mathsf{uu}}(\omega)}{\tilde{\mathsf{S}}_{\mathsf{xx}}(\omega)}\right]
\end{aligned}$$

✏

Theorem 16.5 (Estimating H in measurement additive noise system). [21] *Let* **S** *be the* SYSTEM *illustrated in Figure 16.4 page 127 (B).*

THM

$$\left\{\begin{array}{lll} (A). & \mathbf{H} \textit{ is} & \text{LINEAR TIME INVARIANT } \textit{and} \\ (B). & \mathsf{x}(n) \textit{ is} & \text{WIDE-SENSE STATIONARY } \textit{and} \\ (C). & \mathsf{x}(n), \mathsf{u}(n), \textit{and } \mathsf{v}(n) \textit{ are} & \text{UNCORRELATED} \end{array}\right\}$$

$$\Longrightarrow \left\{\begin{array}{lll} (1). & \hat{\mathsf{H}}_1(\omega) = \tilde{\mathsf{H}}(\omega)\left[\dfrac{1}{1 + \dfrac{\tilde{\mathsf{S}}_{\mathsf{ww}}(\omega)}{\tilde{\mathsf{S}}_{\mathsf{pp}}(\omega)}}\right] & (\text{UNDER-ESTIMATED}) \textit{ and} \\ (2). & \hat{\mathsf{H}}_2(\omega) = \tilde{\mathsf{H}}(\omega)\left[1 + \dfrac{\tilde{\mathsf{S}}_{\mathsf{vv}}(\omega)}{\tilde{\mathsf{S}}_{\mathsf{qq}}(\omega)}\right] & (\text{OVER-ESTIMATED}) \end{array}\right\}$$

[21] Shin and Hammond (2008) page 294 $\langle H_1(f) = H(f)$ (9.70); $H_2(f) = H(f)\left(1 + S_{n_y n_y}(f)/S_{yy}(f)\right)$ (9.71)$\rangle$, Shin and Hammond (2008) page 294 $\langle H_1(f) = H(f)/\left(1 + S_{n_x n_x}/S_{xx}(f)\right)$ (9.72); $H_2(f) = H(f)$ (9.73)$\rangle$, Mitchell (1982) page 277 $\langle H_1(f) = H_0(f)/\left(1 + G_{nn}/G_{uu}\right)\rangle$ Mitchell (1982) page 278 $\langle H_2(f) = H_0(f)\left(1 + G_{mm}/G_{vv}\right)\rangle$

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

✎PROOF:

$$\begin{aligned}
\hat{H}_1(\omega) &\triangleq \frac{\tilde{S}_{yx}(\omega)}{\tilde{S}_{xx}(\omega)} && \text{by definition of } \hat{H}_1 && \text{(Definition 16.2 page 111)}\\
&= \frac{\tilde{S}_{pp}(\omega)\tilde{H}(\omega)}{\tilde{S}_{xx}(\omega)} && \text{by Corollary 7.4 page 57} \\
&= \frac{\tilde{S}_{pp}(\omega)\tilde{H}(\omega)}{\tilde{S}_{pp}(\omega)+\tilde{S}_{ww}(\omega)} && \text{by hypothesis (A)} && \text{and Corollary 5.3 page 37}\\
&= \tilde{H}(\omega)\left[\frac{1}{1+\frac{\tilde{S}_{ww}(\omega)}{\tilde{S}_{pp}(\omega)}}\right]
\end{aligned}$$

$$\begin{aligned}
\hat{H}_2(\omega) &\triangleq \frac{\tilde{S}_{yy}(\omega)}{\tilde{S}_{xy}(\omega)} && \text{by definition of } \hat{H}_2 && \text{(Definition 16.3 page 112)}\\
&= \frac{\tilde{S}_{qq}(\omega)+\tilde{S}_{vv}(\omega)}{\tilde{S}_{xy}(\omega)} && \text{by hypothesis (C)} && \text{and Corollary 7.1 page 52}\\
&= \frac{\tilde{S}_{qq}(\omega)+\tilde{S}_{vv}(\omega)}{\tilde{S}_{xq}(\omega)} && \text{by hypothesis (C)} && \text{and Theorem 7.4 page 55}\\
&= \frac{\tilde{S}_{qq}(\omega)+\tilde{S}_{vv}(\omega)}{\tilde{S}_{pq}(\omega)} && \text{by hypothesis (C)} && \text{and Lemma 7.3 page 54}\\
&= \frac{\tilde{S}_{qq}(\omega)+\tilde{S}_{vv}(\omega)}{\tilde{S}_{qq}(\omega)/\tilde{H}(\omega)} && \text{by } \textit{LTI} \text{ hypothesis (A)} && \text{and Corollary 5.3 page 37}\\
&= \tilde{H}(\omega)\left[1+\frac{\tilde{S}_{vv}(\omega)}{\tilde{S}_{qq}(\omega)}\right] && \text{by hypotheses (A) and (B)} && \text{and Corollary 5.3 page 37}
\end{aligned}$$

✏

Corollary 16.4. *Let* **S** *be the* SYSTEM *illustrated in Figure 16.4 (page 127).*

COR

$$\left\{\begin{array}{ll}(A). & \textit{hypotheses of Theorem 16.5 and}\\ (B). & \mathsf{u}(n)=\mathsf{u}(n)=0 \quad (\text{NO INPUT NOISE})\end{array}\right\} \Longrightarrow \left\{\hat{H}_1(\omega) = \tilde{H}(\omega) \quad (\text{UNBIASED})\right\}$$

$$\left\{\begin{array}{ll}(A). & \textit{hypotheses of Theorem 16.5 and}\\ (B). & \mathsf{v}(n)=0 \quad (\text{NO OUTPUT NOISE})\end{array}\right\} \Longrightarrow \left\{\hat{H}_2(\omega) = \tilde{H}(\omega) \quad (\text{UNBIASED})\right\}$$

Lemma 16.4. *Let* **S** *be the* SYSTEM *illustrated in Figure 16.4 (page 127).*

LEM

$$\left\{\textit{There exists } \kappa(\omega) \textit{ such that } \tilde{S}_{vv}(\omega)=\kappa(\omega)\tilde{S}_{uu}(\omega)\right\}$$
$$\Longrightarrow \left\{\tilde{S}_{uu}(\omega)=\frac{\left|\hat{H}(\omega)\right|^2\tilde{S}_{xx}(\omega)-\hat{H}(\omega)\tilde{S}_{xy}(\omega)-\hat{H}^*(\omega)\tilde{S}^*_{xy}(\omega)+\tilde{S}_{yy}(\omega)}{\kappa(\omega)+\left|\hat{H}(\omega)\right|^2}\right\}$$

✎PROOF:

1. Development based on results of previous chapters:

$$\begin{aligned}
\tilde{S}_{vv} &= \tilde{S}_{yy}-\tilde{S}_{qq} && \text{by Corollary 7.1 page 52}\\
&= \tilde{S}_{yy}-\tilde{S}_{pq}\hat{H} && \text{by Corollary 5.3 page 37}\\
&= \tilde{S}_{yy}-\tilde{S}_{xy}\hat{H} && \text{by Theorem 7.4 page 55}\\
\tilde{S}_{uu} &= \tilde{S}_{xx}-\tilde{S}_{pp} && \text{by Corollary 7.1 page 52}
\end{aligned}$$

COPYRIGHT © 2019 DANIEL J. GREENHOE

$$= \tilde{S}_{xx} - \frac{\tilde{S}_{qp}}{\hat{H}} \qquad \text{by Corollary 5.3 page 37}$$

$$= \tilde{S}_{xx} - \frac{\tilde{S}_{yx}}{\hat{H}} \qquad \text{by Theorem 7.4 page 55}$$

$$\tilde{S}_{uu}\left[|\hat{H}|^2 + \kappa\right] = |\hat{H}|^2\tilde{S}_{uu} + \kappa\tilde{S}_{uu}$$

$$\triangleq \tilde{S}_{uu}|\hat{H}|^2 + \tilde{S}_{vv} \qquad \text{by definition of } \kappa(\omega)$$

$$= |\hat{H}|^2\left[\tilde{S}_{xx} - \frac{\tilde{S}_{yx}}{\hat{H}}\right] + \left[\tilde{S}_{yy} - \tilde{S}_{xy}\hat{H}\right]$$

$$= |\hat{H}|^2\tilde{S}_{xx} - \hat{H}^*\tilde{S}_{yx} - \tilde{S}_{xy}\hat{H} + \tilde{S}_{yy}$$

$$\Longrightarrow \boxed{\tilde{S}_{uu}(\omega) = \frac{|\hat{H}(\omega)|^2\tilde{S}_{xx}(\omega) - \hat{H}(\omega)\tilde{S}_{xy}(\omega) - \hat{H}^*(\omega)\tilde{S}^*_{xy}(\omega) + \tilde{S}_{yy}(\omega)}{\kappa(\omega) + |\hat{H}(\omega)|^2}}$$

2. Development of Wicks and Vold (Wicks and Vold (1986)):

$$\tilde{Y} - \tilde{V} = \tilde{Q} = \hat{H}\tilde{P} = \hat{H}(\tilde{X} - \tilde{U}) \qquad \text{by definition of } \hat{H}$$

$$\hat{H}\tilde{U} - \tilde{V} = \hat{H}\tilde{X} - \tilde{Y} \qquad \text{by } \textit{left distributive} \text{ prop.} \qquad \text{(Theorem N.4 page 307}$$

$$E\left(\left[\hat{H}\tilde{U} - \tilde{V}\right]\left[\hat{H}\tilde{U} - \tilde{V}\right]^*\right) = E\left(\left[\hat{H}\tilde{X} - \tilde{Y}\right]\left[\hat{H}\tilde{X} - \tilde{Y}\right]^*\right)$$

$$|\hat{H}|^2\tilde{S}_{uu} - \hat{H}\cancelto{0}{\tilde{S}_{uv}} - \hat{H}^*\cancelto{0}{\tilde{S}_{vu}} + \tilde{S}_{vv} = |\hat{H}|^2\tilde{S}_{xx} - \hat{H}\tilde{S}_{xy} - \hat{H}^*\tilde{S}^*_{xy} + \tilde{S}_{yy} \qquad \text{because } u \text{ and } v \text{ are } \textit{uncorrelated}$$

$$|\hat{H}|^2\tilde{S}_{uu} + \kappa\tilde{S}_{uu} = |\hat{H}|^2\tilde{S}_{xx} - \hat{H}\tilde{S}_{xy} - \hat{H}^*\tilde{S}^*_{xy} + \tilde{S}_{yy} \qquad \text{by hypothesis}$$

Theorem 16.6. [22] *Let* **S** *be the* SYSTEM *illustrated in Figure 16.4 (page 127). Let* $\hat{H}_\kappa(\omega)$ *be the transfer function estimate defined in Definition 16.7 (page 115).*

THM

$$\left\{\begin{array}{ll} (1). & \textit{There exists } \kappa(\omega) \textit{ such that} \\ & \tilde{S}_{vv}(\omega) = \kappa(\omega)\tilde{S}_{uu}(\omega) \qquad \textit{and} \\ (2). & C(\hat{H}_{ls}) = \tilde{S}_{uu}(\omega) \end{array}\right\} \Longrightarrow \left\{\begin{array}{l} \arg\min_{\hat{H}} C(\hat{H}) = \hat{H}_\kappa(\omega) \\ (\hat{H}_\kappa \textit{ is the "optimal" estimator for minimizing system noise}) \end{array}\right\}$$

PROOF:

1. Let $F \triangleq |\hat{H}(\omega)|^2\tilde{S}_{xx}(\omega) - \hat{H}(\omega)\tilde{S}_{xy}(\omega) - \hat{H}^*(\omega)\tilde{S}^*_{xy}(\omega) + \tilde{S}_{yy}(\omega)$ (numerator in Lemma 16.4) and
   $G \triangleq \kappa(\omega) + |\hat{H}(\omega)|^2$ (denominator in Lemma 16.4)

2. lemma $\left(\frac{\partial}{\partial\hat{H}_R}\tilde{S}_{uu}\right)$:

$$\boxed{0} = \frac{1}{2}G^2 0 = \frac{1}{2}G^2\frac{\partial}{\partial\hat{H}_R}\tilde{S}_{uu} \qquad \text{set } \frac{\partial}{\partial\hat{H}_R}\tilde{S}_{uu} = 0 \text{ to find optimum } \hat{H}_R$$

$$= \frac{1}{2}G^2\frac{\partial}{\partial\hat{H}_R}\frac{F}{G} \qquad \text{by Lemma 16.4 page 129}$$

$$= \frac{1}{2}G^2\frac{(F'G - G'F)}{G^2} \qquad \text{by } \textit{Quotient Rule}$$

$$= \frac{1}{2}(F'G - G'F)$$

[22] Wicks and Vold (1986) page 898 ⟨has additional s in denominator⟩, Shin and Hammond (2008) page 293 ⟨(9.67)⟩, White et al. (2006) page 679 ⟨(6)⟩

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

$$= \frac{1}{2}\left[2\hat{H}_R\tilde{S}_{xx} - \tilde{S}_{xy} - \tilde{S}_{xy}^*\right]G - \frac{1}{2}2\hat{H}_R F \qquad \text{by definition of } F, G$$ (item (1) page 130)

$$= \boxed{\hat{H}_R\tilde{S}_{xx}G - G\mathbf{R}_e\tilde{S}_{xy} - \hat{H}_R F}$$

3. lemma $\left(\frac{\partial}{\partial\hat{H}_I}\tilde{S}_{uu}\right)$:

$$\boxed{0} = \frac{1}{2}G^2 0 = \frac{1}{2}G^2\frac{\partial}{\partial\hat{H}_I}\tilde{S}_{uu} \qquad \text{set } \frac{\partial}{\partial\hat{H}_I}\tilde{S}_{uu} = 0 \text{ to find optimum } \hat{H}_I$$

$$= \frac{1}{2}G^2 0 = \frac{1}{2}G^2\frac{\partial}{\partial\hat{H}_I}\frac{F}{G} \qquad \text{by Lemma 16.4 page 129}$$

$$= \frac{1}{2}G^2\frac{(F'G - G'F)}{G^2} \qquad \text{by } \textit{Quotient Rule}$$

$$= \frac{1}{2}(F'G - G'F)$$

$$= \frac{1}{2}\left[2\hat{H}_I\tilde{S}_{xx} - i\tilde{S}_{xy} + i\tilde{S}_{xy}^*\right]G - \frac{1}{2}2\hat{H}_I F \qquad \text{by definition of } F, G$$ (item (1) page 130)

$$= \boxed{\hat{H}_I\tilde{S}_{xx}G + G\mathbf{I}_m\tilde{S}_{xy} - \hat{H}_I F}$$

4. Solve for $\hat{H}$ …

$$0 = 0 + i0 = \frac{1}{2}G^2 0 + \frac{1}{2}G^2 0 = \frac{1}{2}G^2\frac{\partial}{\partial\hat{H}_R}\tilde{S}_{uu} + i\frac{1}{2}G^2\frac{\partial}{\partial\hat{H}_I}\tilde{S}_{uu}$$

$$= \left[\hat{H}_R\tilde{S}_{xx}G - G\mathbf{R}_e\tilde{S}_{xy} - \hat{H}_R F\right] + i\left[\hat{H}_I\tilde{S}_{xx}G + G\mathbf{I}_m\tilde{S}_{xy} - \hat{H}_I F\right] \qquad \text{by (2) lemma and (3) lemma}$$

$$= \hat{H}\tilde{S}_{xx}G - \tilde{S}_{xy}^*G - \hat{H}F \quad \text{because } \mathbf{R}_e(z) + i\mathbf{I}_m(z) = z \text{ and } \mathbf{R}_e(z) - i\mathbf{I}_m(z) = z^*$$

$$= \hat{H}\tilde{S}_{xx}G - \tilde{S}_{yx}G - \hat{H}F \qquad \text{by Corollary 2.2 page 15}$$

$$= \hat{H}\tilde{S}_{xx}\left(\kappa + |\hat{H}|^2\right) - \tilde{S}_{yx}\left(\kappa + |\hat{H}|^2\right) - \hat{H}\left(|\hat{H}|^2\tilde{S}_{xx} - \hat{H}\tilde{S}_{xy} - \hat{H}^*\tilde{S}_{xy}^* + \tilde{S}_{yy}\right) \qquad \text{by } F, G \text{ defs.}$$

$$= \hat{H}\tilde{S}_{xx}\left(\kappa + \cancel{|\hat{H}|^2}\right) - \tilde{S}_{yx}\left(\kappa + |\hat{H}|^2\right) - \hat{H}\left(\cancel{|\hat{H}|^2\tilde{S}_{xx}} - \hat{H}\tilde{S}_{xy} - \hat{H}^*\tilde{S}_{xy}^* + \tilde{S}_{yy}\right)$$

$$= \kappa\hat{H}\tilde{S}_{xx} - \tilde{S}_{yx}\left(\kappa + |\hat{H}|^2\right) + \left(\hat{H}^2\tilde{S}_{xy} + |\hat{H}|^2\tilde{S}_{xy}^* - \hat{H}\tilde{S}_{yy}\right)$$

$$= \kappa\hat{H}\tilde{S}_{xx} - \kappa\tilde{S}_{yx} - \cancel{\tilde{S}_{yx}|\hat{H}|^2} + \left(\hat{H}^2\tilde{S}_{xy} + \cancel{|\hat{H}|^2\tilde{S}_{xy}^*} - \hat{H}\tilde{S}_{yy}\right)$$

$$= \hat{H}^2\tilde{S}_{xy} + \hat{H}\left[\kappa\tilde{S}_{xx} - \tilde{S}_{yy}\right] - \kappa\tilde{S}_{xy}^*$$

$$\Longrightarrow \boxed{\hat{H} = \frac{\left(\tilde{S}_{yy} - \kappa\tilde{S}_{xx}\right) \pm \sqrt{\left(\tilde{S}_{yy} - \kappa\tilde{S}_{xx}\right)^2 + 4\kappa\left|\tilde{S}_{xy}\right|^2}}{2\tilde{S}_{xy}}} \qquad \text{by } \textit{Quadratic Equation}$$

16.8 Coherence

16.8.1 Application

Coherence has two basic purposes:

1. The *coherence* of x and y is a measure of how closely x and y are statistically related. That is, it is an indication of how much x and y "cohere" or "stick" together

Copyright © 2019 Daniel J. Greenhoe

2. The *coherence* of x and y is a measure of how reliable are the estimates $\hat{H}_1$ and $\hat{H}_2$ (Definition 16.2 page 111, Definition 16.3 page 112). If the coherence is 0.70 or above, then we can have high confidence that the estimates $\hat{H}_1$ and $\hat{H}_2$ are "good" estimates.[23]

16.8.2 Definitions

Definition 16.12. [24] *Let* S *be a* SYSTEM *with input* $x(n)$ *and output* $y(n)$.

DEF

The ***complex coherence*** *function is defined as* $C_{xy}(\omega) \triangleq \dfrac{\tilde{S}^*_{xy}(\omega)}{\sqrt{\tilde{S}_{xx}(\omega)\tilde{S}_{yy}(\omega)}}$

The ***ordinary coherence*** *function is defined as* $\gamma^2_{xy}(\omega) \triangleq \dfrac{\left|\tilde{S}_{xy}(\omega)\right|^2}{\tilde{S}_{xx}(\omega)\tilde{S}_{yy}(\omega)}$

Proposition 16.7.

PRP

$$\gamma^2_{xy}(\omega) = \frac{\hat{H}_1(\omega)}{\hat{H}_2(\omega)}$$

✎PROOF:

$$\boxed{\gamma^2_{xy}(\omega)} \triangleq \frac{\left|\tilde{S}_{xy}\right|^2}{\tilde{S}_{xx}\tilde{S}_{yy}} \qquad \text{by definition of } \gamma^2_{xy} \qquad \text{(Definition 16.12 page 132)}$$

$$= \frac{\tilde{S}^*_{xy}/\tilde{S}_{xx}}{\tilde{S}_{yy}/\tilde{S}_{xy}} \triangleq \boxed{\frac{\hat{H}_1}{\hat{H}_2}} \qquad \text{by definitions of } \hat{H}_1 \text{ and } \hat{H}_2 \qquad \text{(Definition 16.2 page 111, Definition 16.3 page 112)}$$

✏

Remark 16.2. Note that the *complex transmissibility* $\tilde{T}'_{xy}$ of Remark 16.1 provides a nice mathematical symmetry (always a good sign of good direction) with *coherence* in the system identification family tree. In particular, note that the following:

REM

$$C_{xy} \triangleq \sqrt{\frac{\hat{H}^*_1}{\hat{H}_2}} \text{ whereas } \tilde{T}'_{xy} \triangleq \sqrt{\hat{H}_1\hat{H}_2}$$

✎PROOF:

$$\sqrt{\frac{\hat{H}^*_1(\omega)}{\hat{H}_2(\omega)}} \qquad \text{by definition of } \hat{H}_{gm} \qquad \text{(Definition 16.5 page 113)}$$

✏

16.8.3 A warning

Estimators yield, as the name implies, estimates. These estimates in general contain some error.

[23] Liang and Lee (2015) pages 363–365 ⟨7.4.2COHERECE FUNCTION⟩

[24] Chen et al. (2012) page 4699(1), (2), Liang and Lee (2015) pages 363–365 ⟨7.4.2 Coherence function⟩, Ewins (1986) page 131 ⟨$\gamma^2 = H_1(\omega)/H_2(\omega)$ (3.8)⟩

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

Example 16.1 (The K=1 Welch estimate of coherence). Suppose we have two *uncorrelated* stationary sequences $\mathsf{x}(n)$ and $\mathsf{y}(n)$. Then, there CSD $\mathsf{S}_{\mathsf{xy}}(\omega)$ should be 0 because

$$\begin{aligned}
\mathsf{S}_{\mathsf{xy}}(\omega) &\triangleq \breve{\mathbf{F}}\mathsf{E}\mathsf{R}_{\mathsf{xy}}(m)\\
&= \breve{\mathbf{F}}\mathsf{E}[x(n)y[n+m]]\\
&= \breve{\mathbf{F}}\left[\mathsf{E}_{\mathsf{x}}(n)\right]\left[\mathsf{E}_{\mathsf{y}}[n+m]\right]\\
&= \breve{\mathbf{F}}[0][0]\\
&= 0
\end{aligned}$$

This will give a coherence of 0 also:

$$C(\omega) = \frac{\mathsf{S}_{\mathsf{xy}}}{\sqrt{\mathsf{S}_{\mathsf{xx}}\mathsf{S}_{\mathsf{yy}}}} = 0$$

However, the Welch estimate with $K = 1$ will yield

$$\begin{aligned}
|C(\omega)| &= \left|\frac{\mathsf{S}_{\mathsf{xy}}}{\sqrt{\mathsf{S}_{\mathsf{xx}}\mathsf{S}_{\mathsf{yy}}}}\right|\\
&= \left|\frac{(\tilde{\mathbf{F}}x)(\tilde{\mathbf{F}}y)^*}{\sqrt{\left|\tilde{\mathbf{F}}x\right|^2\left|\tilde{\mathbf{F}}y\right|^2}}\right|\\
&= 1
\end{aligned}$$

Copyright © 2019 Daniel J. Greenhoe
CC BY-NC-ND

Copyright © 2019 Daniel J. Greenhoe

CHAPTER 17

ESTIMATING NOISE

Estimating noise in a system is difficult and many estimation methods are possible.

- Thong et al. (2001)
- Zheng et al. (2002)
- Kim and Kamel (2004)
- Kamel and Sim (2004)



(A) **communications LTI additive noise model** (B) **measurement LTI additive noise model**

Figure 17.1: Additive noise systems with **LTI** operator **H**

Copyright © 2019 Daniel J. Greenhoe

Part IV

Statistical Detection

CHAPTER 18

COMMUNICATION CHANNELS

18.1 System model



Figure 18.1: Communication system model

A communication system is an operator. $\mathbf{S}$ over an information sequence u that generates an estimated information sequence $\hat{s}u$. The system operator factors into a receive operator $\mathbf{R}$, a channel operator $\mathbf{C}$, and a transmit operator $\mathbf{T}$ such that

$$\mathbf{S} = \mathbf{RCT}.$$

The transmit operator operates on an information sequence u to generate a channel signal $\mathsf{s}(t;\boldsymbol{u})$. The channel operator operates on the transmitted signal $\mathsf{s}(t;\boldsymbol{u})$ to generate the received signal $\mathsf{r}(t;\boldsymbol{u})$. The receive operator operates on the received signal $\mathsf{r}(t;\boldsymbol{u})$ to generate the estimate $\hat{s}u$ (see Figure 18.1 (page 139)).

Definition 18.1. *Let U be the set of all sequences u and let*

DEF

$$\begin{array}{llll} \mathbf{S} & : U & \to U & \textit{(system operator)} \\ \mathbf{T} & : U & \to \mathbb{R}^\infty & \textit{(transmit operator)} \\ \mathbf{C} & : \mathbb{R}^\infty & \to \mathbb{R}^\infty & \textit{(channel operator)} \\ \mathbf{R} & : \mathbb{R}^\infty & \to U & \textit{(receive operator)} \end{array}$$

*be operators. A **digital communication system** is the operation $\mathbf{S}$ on the set of information sequences U such that $\mathbf{S} \triangleq \mathbf{RCT}$.*

Communication systems can be continuous or discrete valued in time and/or amplitude:

$s(t) = a(t)\psi(t)$	continuous time t	discrete time t
continuous amplitude $a(t)$	analog communications	discrete-time communications
discrete amplitude $a(t)$	—	digital communications

In this document, we normally take the approach that

1. **C** is stochastic
2. There is no structural constraint on **R**.
3. **R** is optimum with respect to the ML-criterion.

These characteristics are explained more fully below.

18.1.1 Channel operator

Real-world physical channels perform a number of operations on a signal. Often these operations are closely modeled by a channel operator **C**. Properties that characterize a particular channel operator associated with some physical channel include

- linear or non-linear
- time-invariant or time-variant
- memoryless or non-memoryless
- deterministic or stochastic.

Examples of physical channels include free space, air, water, soil, copper wire, and fiber optic cable. Information is carried through a channel using some physical process. These processes include:

Process	Example
electromagnetic waves	free space, air
acoustic waves	water, soil
electric field potential (voltage)	wire
light	fiber optic cable
quantum mechanics	

18.1.2 Receive operator

Let **I** be the *identity operator*.[1] Ideally, **R** is selected such that $\mathbf{RCT} = \mathbf{I}$. In this case we say that **R** is the *left inverse*[2] of **CT** and denote this left inverse by **C**. One example of a system where this inverse exists is the noiseless ISI system. While this is quite useful for mathematical analysis and system design, **C** does not actually exist for any real-world system.

When **C** does not exist, the "ideal" **R** is one that is optimum

1. with respect to some *criterion* (or cost function)
2. and sometimes under some structural *constraint*.

[1] **I** is the *identity operator* if for any operator **X**, $\mathbf{XI} = \mathbf{IX} = \mathbf{X}$.

[2] $\mathbf{X}^{-1}X$ is the *left inverse* of **X** if $\mathbf{X}^{-1}X\mathbf{X} = \mathbf{I}$.
$\mathbf{X}^{-1}X$ is the *right inverse* of **X** if $\mathbf{X}\mathbf{X}^{-1}X = \mathbf{I}$.
$\mathbf{X}^{-1}X$ is the *inverse* of **X** if $\mathbf{X}^{-1}X\mathbf{X} = \mathbf{X}\mathbf{X}^{-1}X = \mathbf{I}$.

BY-NC-ND
Copyright © 2019 Daniel J. Greenhoe

When a structural constraint is imposed on $\mathbf{R}$, the solution is called *structured*; otherwise, it is called *non-structured*.[3] A common example of a structured approach is the use of a transversal filter (FIR filter in DSP) in which optimal coefficients are found for the filter. A structured $\mathbf{R}$ is only optimal with respect to the imposed constraint. Even though $\mathbf{R}$ may be optimal with respect to this structure, $\mathbf{R}$ may not be optimal in general; that is, there may be another structure that would lead to a "better" solution. In a non-structured approach, $\mathbf{R}$ is free to take any form whatsoever (practical or impractical) and therefore leads to the best of the best solutions.

The nature of $\mathbf{R}$ depends heavily on the nature of $\mathbf{C}$. If $\mathbf{C}$ does not exist, then the ideal $\mathbf{R}$ is one that is optimal with respect to some criterion. If $\mathbf{C}$ is deterministic, then appropriate optimization criterion may include

- least square error (LSE) criterion
- minimum absolute error criterion
- minimum peak distortion criterion.

If $\mathbf{C}$ is stochastic then appropriate optimization criterion may include

Bayes:	pdf known and cost function defined
Maximum aposteriori probability (MAP):	pdf known and uniform cost function
Maximum likelihood (ML):	pdf known and no prior probability information
mini-max:	pdf not known but a cost function is defined
Neyman-Pearson:	pdf not known and no cost function defined.

Making $\mathbf{R}$ optimum with respect to one of these criterion leads to an *estimate* $\hat{s}u = \mathbf{RCT}\boldsymbol{u}$ that is also optimum with respect to the same criterion (Definition 8.1 page 64).

18.2 Optimization in the case of additional operations



Figure 18.2: Theorem of reversibility

Often in communication systems, an additional operator $\mathbf{X}$ is inserted such that (see Figure 18.2 (page 141))

$$\mathbf{S} = \mathbf{RXCT}.$$

An example of such an operator $\mathbf{X}$ is a receive filter. Is it still possible to find an $\mathbf{R}$ that will perform as well as the case where $\mathbf{X}$ is not inserted? In general, the answer is "no". For example, if $\mathbf{X}r = 0$, then all received information is lost and obviously there is no $\mathbf{R}$ that can recover from this event. However, in the case where the right inverse $\mathbf{X}^{-1}X$ of $\mathbf{X}$ exists, then the answer to the question is "yes" and an optimum $\mathbf{R}$ still exists. That is, it doesn't matter if an $\mathbf{X}$ is inserted into system as long as $\mathbf{X}$ is invertible. This is stated formally in the next theorem.

Theorem 18.1 (Theorem of Reversibility). [4] *Let*

[3] Trees (2001) page 12
[4] Trees (2001) pages 289–290

Copyright © 2019 Daniel J. Greenhoe

- $\hat{\theta}_{=}\mathbf{RCT}\boldsymbol{u}$ *be the optimum estimate of* $\boldsymbol{u}$
- $\mathbf{X}$ *be an operator with right inverse* $\mathbf{X}^{-1}X$.

Then there exists some $\mathbf{R}'$ *such that*

THM $\hat{\theta}_{=}\mathbf{R}'\mathbf{XCT}\boldsymbol{u}$.

✎PROOF: Let $\mathbf{R}' = \mathbf{RX}^{-1}X$. Then

$$\mathbf{R}'\mathbf{XCT}\boldsymbol{u} = \mathbf{RX}^{-1}X\mathbf{CT}\boldsymbol{u} = \mathbf{RCT}\boldsymbol{u} = \hat{\theta}.$$

✏

18.3 Channel Statistics

The receiver needs to make a decision as to what sequence $(\!(u)\!)$ the transmitter has sent. This decision should be optimal in some sense. Very often the optimization criterion is chosen to be the *maximal likelihood (ML)* criterion. The information that the receiver can use to make an optimal decision is the received signal $\mathsf{r}(t)$.

If the symbols in $\mathsf{r}(t)$ are statistically *independent*, then the optimal estimate of the current symbol depends only on the current symbol period of $\mathsf{r}(t)$. Using other symbol periods of $\mathsf{r}(t)$ has absolutely no additional benefit. Note that the AWGN channel is *memoryless*; that is, the way the channel treats the current symbol has nothing to do with the way it has treated any other symbol. Therefore, if the symbols sent by the transmitter into the channel are independent, the symbols coming out of the channel are also independent.

However, also note that the symbols sent by the transmitter are often very intentionally not independent; but rather a strong relationship between symbols is intentionally introduced. This relationship is called *channel coding*. With proper channel coding, it is theoretically possible to reduce the probability of communication error to any arbitrarily small value as long as the channel is operating below its *channel capacity*.

This chapter assumes that the received symbols are statistically independent; and therefore optimal decisions at the receiver for the current symbol are made only from the current symbol period of $\mathsf{r}(t)$.

The received signal $\mathsf{r}(t)$ over a single symbol period contains an uncountably infinite number of points. That is a lot. It would be nice if the receiver did not have to look at all those uncountably infinite number of points when making an optimal decision. And in fact the receiver does indeed not have to. As it turns out, a single finite set of *statistics* $\{\dot{r}_1, \dot{r}_2, \ldots, \dot{r}_N\}$ is sufficient for the receiver to make an optimal decision as to which value the transmitter sent.

Definition 18.2. *Let* $\mathbf{C}$ *be an additive noise channel*

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

CHAPTER 19

OPTIMAL SYMBOL DETECTION

19.1 ML Estimation

Theorem 19.1. *In an AWGN channel with received signal* $\mathsf{r}(t) = s(t;\phi) + n(t)$ *Let*

- $\mathsf{r}(t) = s(t;\phi) + n(t)$ *be the received signal in an AWGN channel*
- $n(t)$ *a Gaussian white noise process*
- $\mathsf{s}(t;\phi)$ *the transmitted signal such that*

$$s(t;\phi) = \sum_{n\in\mathbb{Z}} a_n \lambda(t-nT)\cos(2\pi f_c t + \theta_n + \phi).$$

Then the optimal ML estimate of ϕ *is either of the two equivalent expressions*

THM

$$\begin{aligned}\hat{\phi}_{\mathsf{ml}} &= -\operatorname{atan}\left[\frac{\sum_{n\in\mathbb{Z}} a_n \int_{t\in\mathbb{R}} \mathsf{r}(t)\lambda(t-nT)\sin(2\pi f_c t + \theta_n)\,\mathsf{d}t}{\sum_{n\in\mathbb{Z}} a_n \int_{t\in\mathbb{R}} \mathsf{r}(t)\lambda(t-nT)\cos(2\pi f_c t + \theta_n)\,\mathsf{d}t}\right] \\ &= \arg_\phi\left(\sum_{n\in\mathbb{Z}} a_n \int_{t\in\mathbb{R}} \mathsf{r}(t)\left[\lambda(t-nT)\sin(2\pi f_c t + \theta_n + \phi)\right]\,\mathsf{d}t = 0\right).\end{aligned}$$

✎PROOF:

$$\begin{aligned}\hat{\phi}_{\mathsf{ml}} &= \arg_\phi\left(2\int_{t\in\mathbb{R}} \mathsf{r}(t)\left[\frac{\partial}{\partial\phi}s(t;\phi)\right]\,\mathsf{d}t = \frac{\partial}{\partial\phi}\int_{t\in\mathbb{R}} s^2(t;\phi)\,\mathsf{d}t\right) \qquad \text{by Theorem 11.6 page 84} \\ &= \arg_\phi\left(2\int_{t\in\mathbb{R}} \mathsf{r}(t)\left[\frac{\partial}{\partial\phi}s(t;\phi)\right]\,\mathsf{d}t = \frac{\partial}{\partial\phi}\|s(t;\phi)\|^2\,\mathsf{d}t\right) \\ &= \arg_\phi\left(2\int_{t\in\mathbb{R}} \mathsf{r}(t)\left[\frac{\partial}{\partial\phi}s(t;\phi)\right]\,\mathsf{d}t = 0\right) \\ &= \arg_\phi\left(\int_{t\in\mathbb{R}} \mathsf{r}(t)\left[\frac{\partial}{\partial\phi}\sum_{n\in\mathbb{Z}} a_n\lambda(t-nT)\cos(2\pi f_c t + \theta_n + \phi)\right]\,\mathsf{d}t = 0\right)\end{aligned}$$

$$= \arg_\phi\left(-\sum_{n\in\mathbb{Z}} a_n \int_{t\in\mathbb{R}} \mathsf{r}(t)\left[\lambda(t-nT)\sin(2\pi f_c t+\theta_n+\phi)\right]\,\mathrm{d}t = 0\right)$$

$$= \arg_\phi\left(\sum_{n\in\mathbb{Z}} a_n \int_{t\in\mathbb{R}} \mathsf{r}(t)\lambda(t-nT)\left[\sin(2\pi f_c t+\theta_n)\cos(\phi)+\sin(\phi)\cos(2\pi f_c t+\theta_n)\right]\,\mathrm{d}t = 0\right)$$

$$= \arg_\phi\left(\sum_{n\in\mathbb{Z}} a_n \int_{t\in\mathbb{R}} \mathsf{r}(t)\lambda(t-nT)\sin(\phi)\cos(2\pi f_c t+\theta_n)\,\mathrm{d}t = -\sum_{n\in\mathbb{Z}} a_n \int_{t\in\mathbb{R}} \mathsf{r}(t)\lambda(t-nT)\sin(2\pi f_c t+\theta_n)\cos(\phi)\,\mathrm{d}t\right)$$

$$= \arg_\phi\left(\sin(\phi)\sum_{n\in\mathbb{Z}} a_n \int_{t\in\mathbb{R}} \mathsf{r}(t)\lambda(t-nT)\cos(2\pi f_c t+\theta_n)\,\mathrm{d}t = -\cos(\phi)\sum_{n\in\mathbb{Z}} a_n \int_{t\in\mathbb{R}} \mathsf{r}(t)\lambda(t-nT)\sin(2\pi f_c t+\theta_n)\,\mathrm{d}t\right)$$

$$= \arg_\phi\left(\frac{\sin(\phi)}{\cos(\phi)} = -\frac{\sum_{n\in\mathbb{Z}} a_n \int_{t\in\mathbb{R}} \mathsf{r}(t)\lambda(t-nT)\sin(2\pi f_c t+\theta_n)\,\mathrm{d}t}{\sum_{n\in\mathbb{Z}} a_n \int_{t\in\mathbb{R}} \mathsf{r}(t)\lambda(t-nT)\cos(2\pi f_c t+\theta_n)\,\mathrm{d}t}\right)$$

$$= \arg_\phi\left(\tan(\phi) = -\frac{\sum_{n\in\mathbb{Z}} a_n \int_{t\in\mathbb{R}} \mathsf{r}(t)\lambda(t-nT)\sin(2\pi f_c t+\theta_n)\,\mathrm{d}t}{\sum_{n\in\mathbb{Z}} a_n \int_{t\in\mathbb{R}} \mathsf{r}(t)\lambda(t-nT)\cos(2\pi f_c t+\theta_n)\,\mathrm{d}t}\right)$$

$$= \arg_\phi\left(\phi = -\operatorname{atan}\left(\frac{\sum_{n\in\mathbb{Z}} a_n \int_{t\in\mathbb{R}} \mathsf{r}(t)\lambda(t-nT)\sin(2\pi f_c t+\theta_n)\,\mathrm{d}t}{\sum_{n\in\mathbb{Z}} a_n \int_{t\in\mathbb{R}} \mathsf{r}(t)\lambda(t-nT)\cos(2\pi f_c t+\theta_n)\,\mathrm{d}t}\right)\right)$$

$$= -\operatorname{atan}\left(\frac{\sum_{n\in\mathbb{Z}} a_n \int_{t\in\mathbb{R}} \mathsf{r}(t)\lambda(t-nT)\sin(2\pi f_c t+\theta_n)\,\mathrm{d}t}{\sum_{n\in\mathbb{Z}} a_n \int_{t\in\mathbb{R}} \mathsf{r}(t)\lambda(t-nT)\cos(2\pi f_c t+\theta_n)\,\mathrm{d}t}\right)$$

19.2 Generalized coherent modulation



Figure 19.1: Distributions of orthonormal components

Theorem 19.2. *Let*

- $(V, \langle\cdot|\cdot\rangle, S)$ *be a modulation space*
- $\Psi \triangleq \{\psi_n(t) : n = 1,2,\dots,N\}$ *be a set of orthonormal functions that span* S
- $\dot r_n \triangleq \langle \mathsf{r}(t) \,|\, \psi_n(t)\rangle$
- $R \triangleq \{\dot r_n : n = 1,2,\dots,N\}$
- $\dot s_n(m) \triangleq \langle s(t;m) \,|\, \psi_n(t)\rangle$

and let V *be partitioned into* ***decision regions***

$$\{D_m : m = 1,2,\dots,|S|\}$$

such that

$$\mathsf{r}(t) \in D_{\hat m} \iff \hat m = \arg\max_m \mathsf{P}\{s(t;m)|\mathsf{r}(t)\}\,.$$

BY-NC-ND Copyright © 2019 Daniel J. Greenhoe

*Then the **probability of detection error** is*

THM

$$\mathsf{P}\{error\} = 1 - \left(\frac{1}{\sqrt{2\pi\sigma^2}}\right)^N \sum_m \mathsf{P}\{m\ sent\} \int_{\boldsymbol{r}\in D_m} \exp\frac{-1}{2\sigma^2}\sum_{n=1}^N [\dot{r}_n - \dot{s}_n(m)]^2\ d\boldsymbol{r}.$$

✎PROOF:

$$\begin{aligned}
\mathsf{P}\{\text{error}\} &= 1 - \mathsf{P}\{\text{no error}\}\\
&= 1 - \sum_m \mathsf{P}\left\{(m \text{ sent}) \wedge (\hat{m} = m \text{ detected})\right\}\\
&= 1 - \sum_m \mathsf{P}\left\{(\hat{m} = m \text{ detected})|(m \text{ sent})\right\} \mathsf{P}\{m \text{ sent}\}\\
&= 1 - \sum_m \mathsf{P}\{m \text{ sent}\} \mathsf{P}\{\boldsymbol{r}|(m \text{ sent})\}\\
&= 1 - \sum_m \mathsf{P}\{m \text{ sent}\} \int_{\boldsymbol{r}\in D_m} \mathsf{p}\left[\boldsymbol{r}|(\mathsf{m} \text{ sent})\right] d\boldsymbol{r}\\
&= 1 - \sum_m \mathsf{P}\{m \text{ sent}\} \int_{\boldsymbol{r}\in D_m} \prod_n \mathsf{p}\left[\dot{\mathsf{r}}_\mathsf{n}|\mathsf{m}\right]\ d\boldsymbol{r}\\
&= 1 - \sum_m \mathsf{P}\{m \text{ sent}\} \int_{\boldsymbol{r}\in D_m} \prod_{n=1}^N \frac{1}{\sqrt{2\pi\sigma^2}} \exp\frac{-[\dot{r}_n - \mathsf{E}\dot{r}_n]^2}{2\sigma^2}\ d\boldsymbol{r}\\
&= 1 - \left(\frac{1}{\sqrt{2\pi\sigma^2}}\right)^N \sum_m \mathsf{P}\{m \text{ sent}\} \int_{\boldsymbol{r}\in D_m} \exp\frac{-1}{2\sigma^2}\sum_{n=1}^N [\dot{r}_n - \dot{s}_n(m)]^2\ d\boldsymbol{r}
\end{aligned}$$

✎

19.3 Frequency Shift Keying (FSK)

Theorem 19.3. *In an FSK modulation space, the optimal ML estimator of m is*

THM

$$\hat{m} = \arg\max_m \dot{r}_m.$$

✎PROOF:

$$\begin{aligned}
\hat{m} &= \arg\max_m \mathsf{P}\{\mathsf{r}(t)|s(t;m)\}\\
&= \arg\min_m \sum_{n=1}^N [\dot{r}_n - \dot{s}_n(m)]^2 && \text{by Theorem 11.6 (page 84)}\\
&= \arg\min_m \sum_{n=1}^N [\dot{r}_n^2 - 2\dot{r}_n\dot{s}_n(m) + \dot{s}_n^2(m)]\\
&= \arg\min_m \sum_{n=1}^N [-2\dot{r}_n\dot{s}_n(m) + \dot{s}_n^2(m)] && \dot{r}_n^2 \text{ is independent of } m\\
&= \arg\min_m \sum_{n=1}^N [-2\dot{r}_n a\bar{\delta}_{mn} + a^2\bar{\delta}_{mn}]\\
&= \arg\min_m [-2a\dot{r}_m + a^2]
\end{aligned}$$

COPYRIGHT © 2019 DANIEL J. GREENHOE
CC BY-NC-ND

$$= \arg\min_m[-\dot{r}_m] \qquad a \text{ and } 2 \text{ independent of } m$$
$$= \arg\max_m[\dot{r}_m]$$

✏

Theorem 19.4. *If an FSK modulation space let*

$$\begin{array}{rcl|lcl|l} z_2 & \triangleq & \dot{r}_1(1) - \dot{r}_2(1) & z_2 > 0 & \Longrightarrow & \dot{r}_1 > \dot{r}_2 & m = 1 \\ z_3 & \triangleq & \dot{r}_1(1) - \dot{r}_3(1) & z_3 > 0 & \Longrightarrow & \dot{r}_1 > \dot{r}_3 & m = 1 \\ & \vdots & & & & & \\ z_M & \triangleq & \dot{r}_1(1) - \dot{r}_M(1) & z_M > 0 & \Longrightarrow & \dot{r}_1 > \dot{r}_M & m = 1 \end{array}$$

Then the ***probability of detection error*** *is*

THM $$\mathsf{P}\{error\} = 1 - \frac{M-1}{M}\int_0^\infty\int_0^\infty\cdots\int_0^\infty \mathsf{p}(z_2, z_3, \ldots, z_M)\, dz_2 dz_3 \cdots dz_M$$ *where*

$$\mathsf{p}(z_2, z_3, \ldots, z_M) = \frac{1}{(2\pi)^{\frac{M-1}{2}}\sqrt{\det R}} \exp -\frac{1}{2}\begin{bmatrix} z_2 - \dot{s} \\ z_3 - \dot{s} \\ \vdots \\ z_M - \dot{s} \end{bmatrix}^T R^{-1} \begin{bmatrix} z_2 - \dot{s} \\ z_3 - \dot{s} \\ \vdots \\ z_M - \dot{s} \end{bmatrix}$$

and

$$R = \begin{bmatrix} \operatorname{cov}[z_2, z_2] & \operatorname{cov}[z_2, z_3] & \cdots & \operatorname{cov}[z_2, z_M] \\ \operatorname{cov}[z_3, z_2] & \operatorname{cov}[z_3, z_3] & \cdots & \operatorname{cov}[z_3, z_M] \\ \vdots & \vdots & \ddots & \vdots \\ \operatorname{cov}[z_M, z_2] & \operatorname{cov}[z_M, z_3] & \cdots & \operatorname{cov}[z_M, z_M] \end{bmatrix} = N_o \begin{bmatrix} 2 & 1 & \cdots & 1 \\ 1 & 2 & \cdots & 1 \\ \vdots & \vdots & \ddots & \vdots \\ 1 & 1 & \cdots & 2 \end{bmatrix}$$

The inverse matrix R^{-1} is equivalent to (????)

$$R^{-1} \stackrel{?}{=} \frac{1}{MN_o}\begin{bmatrix} M-1 & -1 & \cdots & -1 \\ -1 & M-1 & \cdots & -1 \\ \vdots & \vdots & \ddots & \vdots \\ -1 & -1 & \cdots & M-1 \end{bmatrix}$$

✎PROOF:

$$\begin{aligned} \mathsf{E}z_k &= \mathsf{E}\left[\dot{r}_{11} - \dot{r}_{1k}\right] \\ &= \mathsf{E}\dot{r}_{11} - \mathsf{E}\dot{r}_{1k} \\ &= \dot{s} - 0 \\ &= \dot{s} \end{aligned}$$

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

$$
\begin{aligned}
\operatorname{cov}\left[z_m, z_n\right] &= \mathsf{E}\left[z_m z_n\right] - [\mathsf{E} z_m][\mathsf{E} z_n] \\
&= \mathsf{E}\left[(\dot r_{11} - \dot r_{1m})(\dot r_{11} - \dot r_{1n})\right] - \dot s^2 \\
&= \mathsf{E}\left[\dot r_{11}^2 - \dot r_{11}\dot r_{1n} - \dot r_{1m}\dot r_{11}\dot r_{1m}\dot r_{1n}\right] - \dot s^2 \\
&= [\operatorname{var}\dot r_{11} + (\mathsf{E}\dot r_{11})^2] - \mathsf{E}\left[\dot r_{11}\right]\mathsf{E}\left[\dot r_{1n}\right] - \mathsf{E}\left[\dot r_{1m}\right]\mathsf{E}\left[\dot r_{11}\right] + [\operatorname{cov}\left[\dot r_{1m}, \dot r_{1n}\right] + (\mathsf{E}\dot r_{1m})(\mathsf{E}\dot r_{1n})] - \dot s^2 \\
&= [\operatorname{var}\dot r_{11} + \dot s^2] - a\cdot 0 - 0\cdot a + [\operatorname{cov}\left[\dot r_{1m}, \dot r_{1n}\right] + 0\cdot 0] - \dot s^2 \\
&= \operatorname{var}\dot r_{11} + \operatorname{cov}\left[\dot r_{1m}, \dot r_{1n}\right] \\
&= N_o + \operatorname{cov}\left[\dot r_{1m}, \dot r_{1n}\right] \\
&= \begin{cases} 2N_o & \text{for } m = n \\ N_o & \text{for } m \neq n. \end{cases}
\end{aligned}
$$

$$
\begin{aligned}
P\{\text{error}\} &= 1 - P\{\text{no error}\} \\
&= 1 - \sum_{m=1}^{M} P\{\text{m transmitted}) \wedge (\forall k \neq m, \dot r_m > \dot r_k\} \\
&= 1 - (M-1)P\{1 \text{ transmitted}) \wedge (\dot r_{11} > \dot r_{12}) \wedge (\dot r_{11} > \dot r_{13}) \wedge \cdots \wedge (\dot r_{11} > \dot r_{1M})\} \\
&= 1 - (M-1)P\{(\dot r_{11} - \dot r_{12} > 0) \wedge (\dot r_{11} - \dot r_{13} > 0) \wedge \cdots \wedge (\dot r_{11} - \dot r_{1M} > 0)|1 \text{ transmitted})\}P\{1 \text{ transmitted})\} \\
&= 1 - \frac{M-1}{M} P\{(z_2 > 0) \wedge (z_3 > 0) \wedge \cdots \wedge (z_M > 0)|1 \text{ transmitted})\} \\
&= 1 - \frac{M-1}{M} \int_0^\infty \int_0^\infty \cdots \int_0^\infty \mathsf{p}(z_2, z_3, \ldots, z_M)\, dz_2 dz_3 \cdots dz_M.
\end{aligned}
$$

✏

19.4 Quadrature Amplitude Modulation (QAM)

19.4.1 Receiver statistics



Figure 19.2: Distributions of QAM components

Theorem 19.5. *Let $(V, \langle\cdot\,|\,\cdot\rangle$ be a QAM modulation space such that*

$$
\begin{aligned}
\mathsf{r}(t) &= \mathsf{s}(t; m) + \mathsf{n}(t) \\
\dot r_c &\triangleq \langle \mathsf{r}(t)\,|\,\psi_c(t)\rangle \\
\dot r_s &\triangleq \langle \mathsf{r}(t)\,|\,\psi_s(t)\rangle\,.
\end{aligned}
$$

Copyright © 2019 Daniel J. Greenhoe

Then $(\dot{r}_c|m)$ *and* $(\dot{r}_s|m)$ *are* ***independent*** *and have* ***marginal distributions***

$$\begin{aligned}(\dot{r}_c|m) &\sim & \mathsf{N}\left(a_m,\sigma^2\right) &= \mathsf{N}\left(r_m\cos\theta_m,\sigma^2\right)\\ (\dot{r}_s|m) &\sim & \mathsf{N}\left(b_m,\sigma^2\right) &= \mathsf{N}\left(r_m\sin\theta_m,\sigma^2\right).\end{aligned}$$

✎PROOF: See Theorem 11.5 (page 83) page 83. ✏

19.4.2 Detection

Theorem 19.6. *Let* $(V,\langle\cdot\,|\,\cdot\rangle,S)$ *be a QAM modulation space with*

$$\begin{aligned}\mathsf{r}(t) &= \mathsf{s}(t;m)+\mathsf{n}(t)\\ \dot{r}_c &\triangleq & \langle\mathsf{r}(t)\,|\,\psi_c(t)\rangle\\ \dot{r}_s &\triangleq & \langle\mathsf{r}(t)\,|\,\psi_s(t)\rangle\,.\end{aligned}$$

Then $\{\dot{r}_c,\dot{r}_s\}$ *are sufficient statistics for optimal ML detection and the optimal ML estimate of* m *is*

$$\hat{u}_{\mathsf{ml}}[m] = \arg\min_m\left[(\dot{r}_c-a_m)^2+(\dot{r}_s-b_m)^2\right].$$

✎PROOF:

$$\begin{aligned}\hat{u}_{\mathsf{ml}}[m] &= \arg\max_m \mathsf{P}\left\{\mathsf{r}(t)|s(t;m)\right\} && \text{by Definition 8.1 (page 64)}\\ &= \arg\min_m \sum_{n=1}^{N}[\dot{r}_n-\dot{s}_n(m)]^2 && \text{by Theorem 11.6 (page 84)}\\ &= \arg\min_m\left[(\dot{r}_c-a_m)^2+(\dot{r}_s-b_m)^2\right]\end{aligned}$$

✏

19.4.3 Probability of error



Figure 19.3: QAM-16 cosstellation and decision regions

Theorem 19.7. *In a QAM-16 constellation as shown in Figure 19.3 (page 148), the probability of error is*

$$\mathsf{P}\left\{error\right\} = \frac{9}{4}Q^2\left(\frac{\dot{s}_{21}-\dot{s}_{11}}{2N_o}\right).$$

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

✎PROOF: Let

$$d \triangleq \dot{s}_{21} - \dot{s}_{11}.$$

Then

$$\begin{aligned}
\mathsf{P}\{\text{error}\} &= \sum_{m=1}^{M} \mathsf{P}\left\{[s(t;m) \text{ transmitted }] \wedge [(\dot{r}_1, \dot{r}_2) \notin D_m]\right\} \\
&= \sum_{m=1}^{M} \mathsf{P}\left\{[(\dot{r}_1, \dot{r}_2) \notin D_m] | [s(t;m) \text{ transmitted }]\right\} \mathsf{P}\left\{[s(t;m) \text{ transmitted }]\right\} \\
&= \frac{1}{M} \sum_{m=1}^{M} \mathsf{P}\left\{[(\dot{r}_1, \dot{r}_2) \notin D_m] | [s(t;m) \text{ transmitted }]\right\} \\
&= \frac{1}{M}\left[4\mathsf{P}\left\{(\dot{r}_1, \dot{r}_2) \notin D_1 | s_1(t)\right\} + 8\mathsf{P}\left\{(\dot{r}_1, \dot{r}_2) \notin D_2 | s_2(t)\right\} + 4\mathsf{P}\left\{(\dot{r}_1, \dot{r}_2) \notin D_6 | s_6(t)\right\}\right] \\
&= \frac{1}{M}\left[4\int\int_{(x,y)\notin D_1} \mathrm{p}_{xy|1}(x,y)\,\mathrm{d}x\,\mathrm{d}y + 8\int\int_{(x,y)\notin D_2} \mathrm{p}_{xy|2}(x,y)\,\mathrm{d}x\,\mathrm{d}y + \right. \\
&\quad \left. 4\int\int_{(x,y)\notin D_6} \mathrm{p}_{xy|6}(x,y)\,\mathrm{d}x\,\mathrm{d}y\right] \\
&= \frac{1}{M}\left[4\int\int_{(x,y)\notin D_1} \mathrm{p}_{x|1}(x)\mathrm{p}_{y|1}(y)\,\mathrm{d}x\,\mathrm{d}y + 8\int\int_{(x,y)\notin D_2} \mathrm{p}_{x|2}(x)\mathrm{p}_{y|2}(y)\,\mathrm{d}x\,\mathrm{d}y + \right. \\
&\quad \left. 4\int\int_{(x,y)\notin D_6} \mathrm{p}_{x|6}(x)\mathrm{p}_{y|6}(y)\,\mathrm{d}x\,\mathrm{d}y\right] \\
&= \frac{1}{M}\left[4Q\left(\frac{d}{2N_o}\right)Q\left(\frac{d}{2N_o}\right) + 8Q\left(\frac{d}{2N_o}\right)2Q\left(\frac{d}{2N_o}\right) + 4\cdot 2Q\left(\frac{d}{2N_o}\right)2Q\left(\frac{d}{2N_o}\right)\right] \\
&= \frac{9}{4}Q^2\left(\frac{d}{2N_o}\right)
\end{aligned}$$

✏

19.5 Phase Shift Keying (PSK)

19.5.1 Receiver statistics



Figure 19.4: Distributions of PSK components

COPYRIGHT © 2019 DANIEL J. GREENHOE
BY-NC-ND

Theorem 19.8. *Let*

$$\begin{aligned}
\dot{r}_c &\triangleq && \langle \mathsf{r}(t) \mid \psi_c(t)\rangle \\
\dot{r}_s &\triangleq && \langle \mathsf{r}(t) \mid \psi_s(t)\rangle \\
\theta_m &\triangleq && \operatorname{atan}\left[\frac{\dot{r}_s(m)}{\dot{r}_c(m)}\right].
\end{aligned}$$

The statistics $(\dot{r}_c|m)$ *and* $(\dot{r}_s|m)$ *are* ***independent*** *with marginal distributions*

$$\begin{aligned}
(\dot{r}_c|m) &\sim && \mathsf{N}\left(r\cos\theta_m, \sigma^2\right) \\
(\dot{r}_s|m) &\sim && \mathsf{N}\left(r\sin\theta_m, \sigma^2\right) \\
\mathrm{p}_{\theta_m}(\theta|m) &= \int_0^\infty x\mathrm{p}_{\dot{r}_c}(x|m)\mathrm{p}_{\dot{r}_s}(x\tan\theta|m)dx.
\end{aligned}$$

✎PROOF:

Indepence and marginal distributions of $\dot{r}_1(m)$ and $\dot{r}_2(m)$ follow directly from Theorem 11.5 (page 83) (page 83).

Let $X \triangleq \dot{r}_1(m)$, $Y \triangleq \dot{r}_2(m)$ and $\Theta \triangleq \theta_m$. Then[1]

$$\begin{aligned}
\mathsf{p}_\theta(\theta)d\theta &\triangleq \mathsf{P}\{\theta < \Theta \le \theta + d\theta\} \\
&= \mathsf{P}\left\{\theta < \operatorname{atan}\frac{Y}{X} \le \theta + d\theta\right\} \\
&= \mathsf{P}\left\{\tan(\theta) < \frac{Y}{X} \le \tan(\theta + d\theta)\right\} \\
&= \mathsf{P}\left\{\tan(\theta) < \frac{Y}{X} \le \tan\theta + (1+\tan^2\theta)\,\mathsf{d}\theta\right\} \\
&= \int_0^\infty \mathsf{P}\left\{\left[\tan\theta < \frac{Y}{X} \le \tan\theta + (1+\tan^2\theta)\,\mathsf{d}\theta\right] \wedge \left[(x < X \le x + \mathsf{d}x)\right]\right\} \\
&= \int_0^\infty \mathsf{P}\left\{\tan\theta < \frac{Y}{x} \le \tan\theta + (1+\tan^2\theta)\,\mathsf{d}\theta \;\middle|\; x < X \le x + dx\right\} \mathsf{P}\{x < X \le x + \mathsf{d}x\} \\
&= \int_0^\infty \mathsf{P}\left\{x\tan\theta < Y \le x\tan\theta + x(1+\tan^2\theta)\,\mathsf{d}\theta) \;\middle|\; X = x\right\} \mathsf{p}_\mathsf{X}(x)\,\mathsf{d}x \\
&= \int_0^\infty [\mathsf{p}_\mathsf{Y}(x\tan\theta)x(1+\tan^2\theta)]\mathsf{p}_\mathsf{X}(x)\,\mathsf{d}x\,\mathsf{d}\theta \\
&= (1+\tan^2\theta)\int_0^\infty x\mathsf{p}_\mathsf{Y}(x\tan\theta)\mathsf{p}_\mathsf{X}(x)\,\mathsf{d}x\,\mathsf{d}\theta \\
\Longrightarrow \\
\mathsf{p}_\theta(\theta)d\theta &= (1+\tan^2\theta)\int_0^\infty x\mathsf{p}_\mathsf{Y}(x\tan\theta)\mathsf{p}_\mathsf{X}(x)\,\mathsf{d}x
\end{aligned}$$

✏

[1] A similar example is in Papoulis (1991), page 138

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

19.5.2 Detection

Theorem 19.9. *Let $(V, \langle\cdot\,|\,\cdot\rangle, S)$ be a PSK modulation space with*

$$\begin{aligned} \mathsf{r}(t) &= \mathsf{s}(t;m) + \mathsf{n}(t) \\ \dot{r}_c &\triangleq \langle \mathsf{r}(t)\,|\,\psi_c(t)\rangle \\ \dot{r}_s &\triangleq \langle \mathsf{r}(t)\,|\,\psi_s(t)\rangle\,. \end{aligned}$$

Then $\{\dot{r}_c, \dot{r}_s\}$ are sufficient statistics for optimal ML detection and the optimal ML estimate of m is

$$\hat{u}_{\mathsf{ml}}[m] = \arg\min_m \left[(\dot{r}_1 - r\cos\theta_m)^2 + (\dot{r}_2 - r\sin\theta_m)^2\right].$$

✎PROOF:

$$\begin{aligned} \hat{u}_{\mathsf{ml}}[m] &= \arg\max_m \mathsf{P}\{\mathsf{r}(t)|s(t;m)\} && \text{by Definition 8.1 (page 64)} \\ &= \arg\min_m \sum_{n=1}^{N} [\dot{r}_n - \dot{s}_n(m)]^2 && \text{by Theorem 11.6 (page 84)} \\ &= \arg\min_m \left[(\dot{r}_1 - r\cos\theta_m)^2 + (\dot{r}_2 - r\sin\theta_m)^2\right]. \end{aligned}$$

✎

19.5.3 Probability of error



Figure 19.5: PSK-8 Decision regions

Theorem 19.10. *The probability of error using PSK modulation is*

$$\mathsf{P}\{error\} = M\left[1 - \int_{\frac{2\pi}{M}\left(m-\frac{3}{2}\right)}^{\frac{2\pi}{M}\left(m-\frac{1}{2}\right)} \mathrm{p}_{\theta_1}(\theta)\,d\theta\right].$$

✎PROOF: See Figure 19.5 (page 151).

COPYRIGHT © 2019 DANIEL J. GREENHOE

$$\begin{aligned}
\mathsf{P}\{\text{error}\} &= \sum_{m=1}^{M} \mathsf{P}\left\{\text{error}|s(t;m) \text{ was transmitted}\right\} \\
&= M\mathsf{P}\left\{\text{error}|s_1(t) \text{ was transmitted}\right\} \\
&= M\left[1 - \int_{\frac{2\pi}{M}\left(m-\frac{3}{2}\right)}^{\frac{2\pi}{M}\left(m-\frac{1}{2}\right)} \mathrm{p}_{\theta_1}(\theta)\, d\theta\right].
\end{aligned}$$

✏

19.6 Pulse Amplitude Modulation (PAM)

19.6.1 Receiver statistics



Figure 19.6: Distribution of PAM component

Theorem 19.11. *Let* $(V, \langle\cdot\,|\,\cdot\rangle$ *be a PAM modulation space such that*

$$\begin{aligned}
\mathsf{r}(t) &= \mathsf{s}(t;m) + \mathsf{n}(t) \\
\dot{r}_c &\triangleq \langle \mathsf{r}(t)\,|\,\psi_c(t)\rangle \\
\dot{r}_s &\triangleq \langle \mathsf{r}(t)\,|\,\psi_{\mathsf{s}}(t)\rangle\,.
\end{aligned}$$

Then $(\dot{r}|m)$ *has* ***distribution***

$$\dot{r}(m) \sim \mathsf{N}\left(a_m, \sigma^2\right).$$

✎PROOF: This follows directly from Theorem 11.5 (page 83) (page 83). ✏

19.6.2 Detection

Theorem 19.12. *Let* $(V, \langle\cdot\,|\,\cdot\rangle, S)$ *be a PAM modulation space with*

$$\begin{aligned}
\mathsf{r}(t) &= \mathsf{s}(t;m) + \mathsf{n}(t) \\
\dot{r} &\triangleq \langle \mathsf{r}(t)\,|\,\psi(t)\rangle\,.
\end{aligned}$$

Then $\dot{r}$ *is a sufficient statistic for the optimal ML detection of* m *and the optimal ML estimate of* m *is*

$$\hat{u}_{\mathsf{ml}}[m] = \arg\min_m |\dot{r} - a_m|.$$

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

✎PROOF:

$$\begin{aligned}
\hat{u}_{\mathsf{ml}}[m] &= \arg\max_m \mathsf{P}\left\{\mathsf{r}(t)|a_m\right\} && \text{by Definition 8.1 (page 64)}\\
&= \arg\min_m \sum_{n=1}^{N}[\dot{r}_n - \dot{s}_n(m)]^2 && \text{by Theorem 11.6 (page 84)}\\
&= \arg\min_m [\dot{r} - \dot{s}(m)]^2\\
&= \arg\min_m |\dot{r} - \dot{s}(m)|
\end{aligned}$$

✏

19.6.3 Probability of error

Theorem 19.13. *The probability of detection error in a PAM modulation space is*

$$\mathsf{P}\{error\} = 2\frac{M-1}{M}\mathsf{Q}\left[\frac{a_2 - a_1}{2\sqrt{N_o}}\right].$$

✎PROOF: Let $d \triangleq a_2 - a_1$ and $\sigma \triangleq \sqrt{\operatorname{var}\dot{r}} = \sqrt{N_o}$. Also, let the decision regions D_m be as illustrated in Figure 19.7 (page 153). Then

$$\begin{aligned}
\mathsf{P}\{error\} &= \sum_{m=1}^{M}\mathsf{P}\left\{s(t;m)\text{ sent }\wedge r \notin D_m\right\}\\
&= \sum_{m=1}^{M}\mathsf{P}\left\{\dot{r} \notin D_m|\mathsf{s}(t;m)\text{ sent }\right\}\mathsf{P}\left\{s(t;m)\text{ sent }\right\}\\
&= \sum_{m=1}^{M}\mathsf{P}\left\{\dot{r}_m \notin D_m\right\}\frac{1}{M}\\
&= \frac{1}{M}\left(\mathsf{Q}\left[\frac{d}{2\sigma}\right] + 2\mathsf{Q}\left[\frac{d}{2\sigma}\right] + \ldots 2\mathsf{Q}\left[\frac{d}{2\sigma}\right] + \mathsf{Q}\left[\frac{d}{2\sigma}\right]\right)\\
&= 2\frac{M-1}{M}\mathsf{Q}\left[\frac{d}{2\sigma}\right]\\
&= 2\frac{M-1}{M}\mathsf{Q}\left[\frac{\dot{s}_2 - \dot{s}_1}{2\sqrt{N_o}}\right]
\end{aligned}$$

✏



Figure 19.7: 4-ary PAM in AWGN channel

COPYRIGHT © 2019 DANIEL J. GREENHOE

Copyright © 2019 Daniel J. Greenhoe

CHAPTER 20

NETWORK DETECTION

20.1 Detection

For detection, we need

1. Cost function: for hard decisions, its range must be linearly ordered. For soft decisions, it can be a lattice.

2. system joint and marginal probabilities (for Bayesian detection)

20.2 Bayesian Estimation

Definition 20.1.

DEF				
	H	$\triangleq$	$\{h_1, h_2, h_3, \ldots\}$	*set of hypotheses*
	$\boldsymbol{D}$	$\triangleq$	$\{D_1, D_2, D_3, \ldots\}$	*partition—decision regions*
	X	$\triangleq$	$\{X_1, X_2, X_3, \ldots\}$	*set of sensor inputs*

$$
\begin{aligned}
cost(h; P) &= \min_{\boldsymbol{D}} \sum_i \mathsf{P}\left\{\left[X \in D_i\right] \wedge \left[H \neq h_i\right]\right\} \\
&= \min_{\boldsymbol{D}} \sum_i \mathsf{P}\left\{X \in D_i \;\middle|\; H \neq h_i\right\} \mathsf{P}\left\{H \neq h_i\right\} \\
&= \min_{\boldsymbol{D}} \sum_i \sum_{j \neq i} \left[1 - \mathsf{P}\left\{X \in D_i \;\middle|\; H = h_i\right\}\right] \sum_{j \neq i} \left[1 - \mathsf{P}\left\{H = h_i\right\}\right]
\end{aligned}
$$

$$\hat{h} = \arg_h cost(h; P)$$

20.3 Joint Gaussian Model

Assume convexity …

$$
\begin{aligned}
\boldsymbol{D} &= \arg_{\boldsymbol{D}} \min_{\boldsymbol{D}} cost(h;P) \\
&= \arg_{\boldsymbol{D}} \left\{ \frac{\partial}{\partial \boldsymbol{D}} \sum_i \int_{D_i} \mathsf{p}(\boldsymbol{x}|H \neq h_i) \underbrace{\mathsf{p}(H \neq h_i)}_{c} d\boldsymbol{x} = 0 \right\} \\
&= \arg_{\boldsymbol{D}} \left\{ \frac{\partial}{\partial \boldsymbol{D}} c \sum_i \int_{D_i} \mathsf{p}(\boldsymbol{x}|H \neq h_i) d\boldsymbol{x} = 0 \right\} \\
&= \arg_{\boldsymbol{D}} \left\{ \frac{\partial}{\partial \boldsymbol{D}} \sum_i \left[1 - \sum_{j \neq i} \int_{D_i} \mathsf{p}(\boldsymbol{x}|H = h_i) d\boldsymbol{x} \right] = 0 \right\} \\
&= \arg_{\boldsymbol{D}} \left\{ \frac{\partial}{\partial \boldsymbol{D}} \sum_i \left[1 - \sum_{j \neq i} \int_{D_i} \frac{1}{\sqrt{(2\pi)^n |\boldsymbol{M}|}} \exp -\frac{1}{2} (\boldsymbol{x} - \mathsf{E}\boldsymbol{x})^T \boldsymbol{M}^{-1} (\boldsymbol{x} - \mathsf{E}\boldsymbol{x}) d\boldsymbol{z} \right] = 0 \right\} \\
&= \arg_{\boldsymbol{D}} \left\{ \sum_i \left[1 - \sum_{j \neq i} \frac{\partial}{\partial \boldsymbol{D}} \int_{D_i} \frac{1}{\sqrt{(2\pi)^n |\boldsymbol{M}|}} \exp -\frac{1}{2} (\boldsymbol{x} - \mathsf{E}\boldsymbol{x})^T \boldsymbol{M}^{-1} (\boldsymbol{x} - \mathsf{E}\boldsymbol{x}) d\boldsymbol{z} \right] = 0 \right\} \\
&= \arg_{\boldsymbol{D}} \left\{ \sum_i \left[1 - \sum_{j \neq i} \begin{bmatrix} \frac{\partial}{\partial D_1} \\ \frac{\partial}{\partial D_2} \\ \vdots \ \frac{\partial}{\partial D_n} \end{bmatrix} \int_{D_i} \frac{1}{\sqrt{(2\pi)^n |\boldsymbol{M}|}} \exp -\frac{1}{2} (\boldsymbol{x} - \mathsf{E}\boldsymbol{x})^T \boldsymbol{M}^{-1} (\boldsymbol{x} - \mathsf{E}\boldsymbol{x}) d\boldsymbol{z} \right] = 0 \right\} \\
&= \arg_{\boldsymbol{D}} \left\{ \sum_i \left[1 - \sum_{j \neq i} \underbrace{\begin{bmatrix} \frac{\partial y_1}{\partial x_1} & \frac{\partial y_2}{\partial x_1} & \cdots & \frac{\partial y_m}{\partial x_1} \\ \frac{\partial y_1}{\partial x_2} & \frac{\partial y_2}{\partial x_2} & \cdots & \frac{\partial y_m}{\partial x_2} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial y_1}{\partial x_n} & \frac{\partial y_2}{\partial x_n} & \cdots & \frac{\partial y_m}{\partial x_n} \end{bmatrix}}_{\text{Jacobian matrix}} \right] \right\}
\end{aligned}
$$

For two variable Gaussian …

$$
\begin{aligned}
cost &= \min_{\boldsymbol{D}} \sum_i \int_{D_i} \mathsf{p}(\boldsymbol{x}|H \neq h_i) \underbrace{\mathsf{p}(H \neq h_i)}_{c} d\boldsymbol{x} \\
&= \min_{\boldsymbol{D}} c \sum_i \int_{D_i} \mathsf{p}(\boldsymbol{x}|H \neq h_i) d\boldsymbol{x} \\
&= \min_{\boldsymbol{D}} c \sum_i \left[1 - \sum_{j \neq i} \int_{D_i} \mathsf{p}(\boldsymbol{x}|H = h_i) d\boldsymbol{x} \right]
\end{aligned}
$$

BY-NC-ND Copyright © 2019 Daniel J. Greenhoe

$$= \min_{\boldsymbol{D}} c \sum_i \left[1 - \sum_{j \neq i} \int_{D_i} \frac{1}{2\pi\sqrt{|M|}} \exp\left(\frac{z_1^2 \mathsf{E}[z_2 z_2] - 2 z_1 z_2 \mathsf{E}[z_1 z_2] + z_2^2 \mathsf{E}[z_1 z_1]}{-2|M|} \right) d\boldsymbol{z} \right]$$

20.4 2 hypothesis, 2 sensor detection

Theorem 20.1 (centralized case)**.** *Let* $(\Omega, \mathbb{E}, \mathsf{P})$ *be a probability space. Let* $D \subsetneq \mathbb{E}$ *be the* DECISION REGION *indicating hypothesis* $H = h_1$*. Let* $\pi_0 \triangleq \mathsf{P}\{H = h_0\}$ *and* $\pi_1 \triangleq \mathsf{P}\{H = h_1\}$*.*

THM

$$\begin{aligned} D &= \arg\min_D \left[\underbrace{\mathsf{P}\left\{(x,y) \in D | H = h_0\right\} \pi_0}_{\text{error for } H = h_0} + \underbrace{\mathsf{P}\left\{(x,y) \in D^{\mathsf{c}} | H = h_1\right\} \pi_1}_{\text{error for } H = h_1} \right] \\ &= \arg\min_D \left[\underbrace{\pi_0 \int_D \mathrm{p}_0(x,y)\ \mathrm{d}x\ \mathrm{d}y}_{\text{error for } H = h_0} + \underbrace{\pi_1 \int_D \mathrm{p}_1(x,y)\ \mathrm{d}x\ \mathrm{d}y}_{\text{error for } H = h_1} \right] \end{aligned}$$

✎PROOF:

$$\begin{aligned} D &= \arg\min_D \left[\mathsf{P}\{\text{error}\}\right] && \text{by definition of decision region } D \\ &= \arg\min_D \left[\mathsf{P}\{\text{error} \wedge H = h_0\} + \mathsf{P}\{\text{error} \wedge H = h_1\}\right] \\ &= \arg\min_D \left[\mathsf{P}\left\{\text{error} | H = h_0\right\} \pi_0 + \mathsf{P}\left\{\text{error} | H = h_1\right\} \pi_1\right] \\ &= \arg\min_D \left[\mathsf{P}\left\{(x,y) \in D | H = h_0\right\} \pi_0 + \mathsf{P}\left\{(x,y) \in D^{\mathsf{c}} | H = h_1\right\} \pi_1\right] \\ &= \arg\min_D \left[\pi_0 \int_D \mathrm{p}_0(x,y)\ \mathrm{d}x\ \mathrm{d}y + \pi_1 \int_D \mathrm{p}_1(x,y)\ \mathrm{d}x\ \mathrm{d}y\right] \end{aligned}$$

✎▷



Example 20.1. In the centralized case, the decision regions D in the xy-plane can be any arbitrary shape, as illustrated to the right.

Definition 20.2.

DEF

Let $\mathbf{P}_x$ *and* $\mathbf{P}_y$ *be* ***set projection operators*** *such that*

$$\begin{aligned} D_x &\triangleq \mathbf{P}_x D \\ D_y &\triangleq \mathbf{P}_y D \end{aligned}$$

Proposition 20.1. *Let* $+$ *represent* MINKOWSKI ADDITION

PRP

$$\mathbf{D} = \mathbf{D}_x + \mathbf{D}_y$$

COPYRIGHT © 2019 DANIEL J. GREENHOE | CC BY-NC-ND

Theorem 20.2 (distributed AND case)**.** *Let* $(\Omega, \mathbb{E}, \mathsf{P})$ *be a probability space. Let* $D \subsetneq \mathbb{E}$ *be the* DECISION REGION *indicating hypothesis* $H = h_1$*. Let* $\pi_0 \triangleq \mathsf{P}\{H = h_0\}$ *and* $\pi_1 \triangleq \mathsf{P}\{H = h_1\}$*. Let* $E \triangleq D^c$*.*

THM

$$D = \arg\min_D \begin{pmatrix} \mathsf{P}\{ & x \in E, & y \in E & \}\{H = h_1\}\pi_1 & + \\ \mathsf{P}\{ & x \in E, & y \in D & \}\{H = h_1\}\pi_1 & + \\ \mathsf{P}\{ & x \in D, & y \in E & \}\{H = h_1\}\pi_1 & + \\ \mathsf{P}\{ & x \in D, & y \in D & \}\{H = h_0\}\pi_0 & \end{pmatrix}$$

✎PROOF:

x	y	H	$x \wedge y$	
0	0	0	0	
0	1	0	0	
1	0	0	0	
1	1	0	1	error
0	0	1	0	error
0	1	1	0	error
1	0	1	0	error
1	1	1	1	

$$\begin{aligned} D &= \arg\min_D [\mathsf{P}\{\text{error}\}] && \text{by definition of decision region } D \\ &= \arg\min_D \left[\mathsf{P}\{\text{error} \wedge H = h_0\} + \mathsf{P}\{\text{error} \wedge H = h_1\}\right] \\ &= \arg\min_D \left[\mathsf{P}\left\{\text{error}|H = h_0\right\}\pi_0 + \mathsf{P}\left\{\text{error}|H = h_1\right\}\pi_1\right] \\ &= \arg\min_D \begin{pmatrix} \mathsf{P}\{ & x \in E_x, & y \in E_y & \}\{H = h_1\}\pi_1 & + \\ \mathsf{P}\{ & x \in D_x, & y \in E_y & \}\{H = h_1\}\pi_1 & + \\ \mathsf{P}\{ & x \in E_x, & y \in D_y & \}\{H = h_1\}\pi_1 & + \\ \mathsf{P}\{ & x \in D_x, & y \in D_y & \}\{H = h_0\}\pi_0 & \end{pmatrix} \end{aligned}$$

✏

Example 20.2. In the distributed AND case, the decision regions D in the xy-plane are only simple rectangular shapes, as illustrated to the right.



Proposition 20.2.

PRP

In general, distributed AND *detection is suboptimal.*

✎PROOF: Because only rectangular decision regions are possible, detection is suboptimal. ✏

Theorem 20.3. [1]

THM

For the distributed AND *detection*

$$D_x = \left\{x|\pi_0 \int_{D_y} \mathrm{p}_0(x,y)\ \mathrm{d}x\ \mathrm{d}y \le \pi_1 \int_{D_y} \mathrm{p}_1(x,y)\ \mathrm{d}x\ \mathrm{d}y\right\}$$

[1] Willett et al. (2000), page 3268

COPYRIGHT © 2019 DANIEL J. GREENHOE
BY-NC-ND

✎PROOF:

$$D_x = \left\{x | y \in D_y \quad \Longrightarrow \quad \mathsf{P}\left\{(x,y)\,|H = h_0\right\}\pi_0 \le \mathsf{P}\left\{(x,y)\,|H = h_1\right\}\pi_1\right\}$$
$$= \left\{x | \pi_0 \int_{D_y} \mathrm{p}_0(x,y)\ \mathrm{d}x\ \mathrm{d}y \le \pi_1 \int_{D_y} \mathrm{p}_1(x,y)\ \mathrm{d}x\ \mathrm{d}y\right\}$$

✏

COPYRIGHT © 2019 DANIEL J. GREENHOE
BY-NC-ND

Copyright © 2019 Daniel J. Greenhoe

Part V

Appendices

APPENDIX A

PROBABILITY SPACE



Figure A.1: Lattice of mathematical spaces

❝ *It is not certain that everything is certain.*❞
Blaise Pascal (1623–1662), mathematician [1]

[1] quote: http://en.wikiquote.org/wiki/Blaise_Pascal
image: http://en.wikipedia.org/wiki/Image:Blaise_pascal.jpg

A.1 Probability functions

Definition A.1. [2] *Let* $(X, \vee, \wedge, \neg, 0, 1; \leq)$ *be a* LATTICE WITH NEGATION.

DEF

The function P *is a* ***probability function*** *if*

(1).	$\mathsf{P}(1)$	$=$	1		(NORMALIZED)	*and*
(2).	$\mathsf{P}(x)$	$\geq$	0	$\forall x \in X$	(NONNEGATIVE)	*and*
(3).	$\bigwedge_{n=1}^{\infty} x_n = 0 \implies \mathsf{P}\left(\bigvee_{n=1}^{\infty} x_n\right)$	$=$	$\sum_{n=1}^{\infty} \mathsf{P}(x_n)$	$\forall x_n \in X$	(σ-ADDITIVE)	.

Remark A.1. The advantage of this definition is that P is a *measure*, and hence all the power of measure theory is subsequently at one's disposal in using P. However, it has often been argued that the requirement of *σ-additivity* is unnecessary for a probability function. Even as early as 1930, de Finetti argued against it, in what became a kind of polite running debate with Fréchet.[3] In fact, Kolmogorov himself provided some argument against *σ-additivity* when referring to the closely related *Axiom of Continuity* saying, "Since the new axiom is essential for infinite fields of probability only, it is almost impossible to elucidate its empirical meaning…For, in describing any observable random process we can obtain only finite fields of probability.…" But in its support he added, "This limitation has been found expedient in researches of the most diverse sort."[4]

There are several other definitions of probability that only require *additivity* rather than *σ-additivity*. On a *Boolean lattice*, the **traditional probability** function is defined as[5]

(1).	$\mathsf{P}(1)$	$=$	1		(*normalized*)	and
(2).	$\mathsf{P}(x)$	$\geq$	0	$\forall x \in X$	(*nonnegative*)	and
(3).	$x \wedge y = 0 \implies \mathsf{P}(x \vee y)$	$=$	$\mathsf{P}(x) + \mathsf{P}(y)$	$\forall x,y \in X$	(*additive*)	.

This definition implies (on a *Boolean lattice*) that

(a).	$\mathsf{P}(0)$	$=$	0		(*nondegenerate*)	and
(b).	$\mathsf{P}(x)$	$\leq$	1	$\forall x \in X$	(*upper bounded*)	and
(c).	$\mathsf{P}(x)$	$=$	$1 - \mathsf{P}(x^{\perp})$	$\forall x \in X$		and
(d).	$\mathsf{P}(x \vee y)$	$\leq$	$\mathsf{P}(x) + \mathsf{P}(y)$	$\forall x,y \in X$	(*subadditive*)	and
(e).	$\mathsf{P}(x \vee y)$	$=$	$\mathsf{P}(x) + \mathsf{P}(y) - \mathsf{P}(x \wedge y)$	$\forall x,y \in X$		and
(f).	$x \leq y \implies \mathsf{P}(x)$	$\leq$	$\mathsf{P}(y)$	$\forall x,y \in X$	(*monotone*)	.

On a *distributive pseudocomplemented lattice*, the **generalized probability** function has been defined as[6]

(1).	$\mathsf{P}(0)$	$=$	0		(*nondegenerate*)	and
(2).	$\mathsf{P}(1)$	$=$	1		(*normalized*)	and
(3).	$0 \leq \mathsf{P}(1)$	$\leq$	1			and
(4).	$\mathsf{P}(x \vee y)$	$=$	$\mathsf{P}(x) + \mathsf{P}(y) - \mathsf{P}(x \wedge y)$	$\forall x,y \in X$		.

On an *orthomodular lattice*, or a *finite modular lattice*, the **quantum probability** function is defined as[7]

(1).	$\mathsf{P}(0)$	$=$	0		(*nondegenerate*)	and
(2).	$\mathsf{P}(1)$	$=$	1		(*normalized*)	and
(3).	$x \perp y \implies \mathsf{P}(x \vee y)$	$=$	$\mathsf{P}(x) + \mathsf{P}(y)$	$\forall x,y \in X$	(*additive*)	.

However, for lattices that are not *distributive*, *modular*, or *orthomodular*, none of these definitions

[2] Billingsley (1995) pages 22–23 ⟨Probability Measures⟩, Kolmogorov (1933a), Kolmogorov (1933b), page 16 ⟨*field of probability*⟩, Pap (1995) pages 8–9 ⟨Definition 2.3(13)⟩, Kalmbach (1986) page 27

[3] de Finetti (1930a), Fréchet (1930a), de Finetti (1930b), Fréchet (1930b), de Finetti (1930c), Cifarelli and Regazzini (1996) pages 258–260

[4] Kolmogorov (1933b), page 15

[5] Papoulis (1991) pages 21–22, Kolmogorov (1933b), page 2 ⟨§1. Axioms I–V⟩

[6] Narens (2014) page 118, Narens (2007)

[7] Greechie (1971) page 126 ⟨DEFINITIONS⟩, Narens (2014) page 118

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

work out so well. Take for example the O_6 *lattice* with the "very reasonable" probability function given in Example ?? (page ??). This probability space $(\mathrm{O}_6, \mathsf{P})$ fails to be any of the 4 probability functions defined in this Remark. It fails to be a *measure-theoretic* or *traditional probability* function because

$$a \wedge b = 0 \qquad \text{but} \qquad \mathsf{P}(a \vee b) = \mathsf{P}(1) = 1 \neq \tfrac{1}{3} + \tfrac{1}{2} = \mathsf{P}(a) + \mathsf{P}(b)\,.$$

It fails to be a *generalized probability* function because

$$\mathsf{P}(a \vee b) = \mathsf{P}(1) = 1 \neq \tfrac{1}{3} + \tfrac{1}{2} - 0 = \mathsf{P}(a) + \mathsf{P}(b) - \mathsf{P}(0) = \mathsf{P}(a) + \mathsf{P}(b) - \mathsf{P}(a \wedge b)\,.$$

It fails to be an *quantum probability* function because

$$a \perp b = 0 \qquad \text{but} \qquad \mathsf{P}(a \vee b) = \mathsf{P}(1) = 1 \neq \tfrac{1}{3} + \tfrac{1}{2} = \mathsf{P}(a) + \mathsf{P}(b)\,.$$

In each of these cases, the function P fails to be *additive*. The solution of Definition A.1 (page 164) is simply to "switch off" *additivity* when the lattice is not *distributive*. This method is a little "crude", but at least it allows us to define probability on a very wide class of lattices, while retaining compatibility with the *Boolean* case.

A.2 Probability Space

In mathematics, a *space* is simply a set and in the most general definition, nothing else. However, normally for a space to actually be useful, some additional structure is added. One of the most general additional structures is a *topology*; and a space together with a topology is called a *topological space*. A topological space imposes additional structure on a space in the form of subsets and guarantees that these subsets are closed under such fundamental operations as set *union* and set *intersection*. With the additional structure available in a topological space, we are able to analyze such basic concepts as continuity, convergence, and connectivity.

However for a great number of mathematical applications, we need to *measure* mathematical objects—the most general measurement being measures on subsets of some set. Examples of measurement in mathematics include integration and probability. Before measurement can be effectively performed on a set, the set must be equipped with a subset structure. In analysis, arguably the most fundamental subset structure is the humble *topology* (Definition ?? page ??). However, a simple topology does not provide sufficient structure for effective measurement. For example, often we would not only like to measure some subset A, but also its complement A^{c}. A topology is not closed under the complement operation. So instead of a topology only, we equip the space with a more powerful (and thus less general) structure called a *σ-algebra* (*sigma-algebra*) (Definition ?? page ??). A *σ-algebra* is a subset structure that is closed under set complement. A set together with a *σ-algebra* is called a *measurable space*. And a set together with a *σ-algebra* and a *measure* on that *σ-algebra* is called a *measure space* (Definition ?? page ??).

The next definition presents a very important measure space—the *probability space*.

Definition A.2.

DEF

The triple $(\Omega, \mathbb{E}, \mathsf{P})$ *is a* ***probability space*** *if*

(1). Ω *is a* SET *and*

(2). $\mathbb{E}$ *is a* σ-ALGEBRA *on* Ω *(Definition ?? page ??) and*

(3). $\mathsf{P} : \mathbb{E} \to [0,1]$ *is a* MEASURE *on* $\mathbb{E}$ *(Definition ?? page ??)* .

If $\boldsymbol{S} \triangleq (\Omega, \mathbb{E}, \mathsf{P})$ *is a* PROBABILITY SPACE *then* x *is an* ***outcome*** *in* $\boldsymbol{S}$ *if* $x \in \Omega$, A *is an* ***event*** *in* $\boldsymbol{S}$ *if* $A \in \mathbb{E}$, *and* $\mathsf{P}A$ *is the* ***probability*** *of* A *in* $\boldsymbol{S}$ *if* A *is an* EVENT *in* $\boldsymbol{S}$.

Definition A.3. [8] *Let* $(\Omega, \mathbb{E}, \mathsf{P})$ *be a* PROBABILITY SPACE *(Definition A.2 page 165)*.

[8] Papoulis (1990) page 52 ⟨Independent Events⟩

COPYRIGHT © 2019 DANIEL J. GREENHOE

DEF *Two* EVENTS *A and B in $\mathbb{E}$ are* ***independent*** *if*

$$\mathsf{P}(A \cap B) = \mathsf{P}(A)\mathsf{P}(B)$$

Definition A.4. [9] *Let $(\Omega, \mathbb{E}, \mathsf{P})$ be a* PROBABILITY SPACE *(Definition A.2 page 165). Let x and y be* EVENTS *in L.*

DEF *The* ***conditional probability of*** *x* ***given*** *y is defined as*

$$\mathsf{P}(A|B) \triangleq \frac{\mathsf{P}(A \cap B)}{\mathsf{P}(B)}$$

A.3 Properties

Proposition A.1.

PRP

$$\underbrace{(\Omega, \mathbb{E}, \mathsf{P}) \text{ is a PROBABILITY SPACE} \quad \Longrightarrow \quad (\Omega, \mathbb{E}, \mathsf{P}) \text{ is a MEASURE SPACE}}_{\text{(every probability space is a measure space)}}$$

Theorem A.1. [10] *Let $(\Omega, \mathbb{E}, \mathsf{P})$ be a* PROBABILITY SPACE *(Definition A.2 page 165).*

THM

(1).	0	$\leq$	$\mathsf{P}(x) \leq 1$	$\forall x \in X$	(BOUNDED)	*and*
(2).	$\mathsf{P}(x)$	$=$	$1 - \mathsf{P}(x^\perp)$	$\forall x \in X$	(PARTITION OF UNITY)	*and*
(3).	$x \leq y$	$\Longrightarrow$	$\mathsf{P}(y^\perp) \leq \mathsf{P}(x^\perp)$	$\forall x,y \in X$	(ANTITONE)	

✎PROOF:

1. Proof for $0 \leq \mathsf{P}(x) \leq 1$:

$$\begin{aligned}
\boxed{0} &= \mathsf{P}(0) && \text{by by } \textit{nondegenerate} \text{ property of } \mathsf{P} \text{ (Definition A.2 page 165)} \\
&\leq \boxed{\mathsf{P}(x)} && \text{because } 0 \leq x \text{ and } \textit{monotone} \text{ property of } \mathsf{P} \\
&\leq \mathsf{P}(1) && \text{because } x \leq 1 \text{ and } \textit{monotone} \text{ property of } \mathsf{P} \\
&= \boxed{1} && \text{by } \textit{normalized} \text{ property of } \mathsf{P}
\end{aligned}$$

2. Proof for $\mathsf{P}(x) = 1 - \mathsf{P}(x^\perp)$:

   (a) Proof that P is *additive* (Definition A.2 page 165) over $\{0, x, x^\perp\} \subseteq X$:

      i. $\{0, x, x^\perp\}$ is *distributive*.

      ii. $x \wedge x^\perp = 0$ for all $x \in X$ by the *non-contradiction* property of *orthocomplemented lattices*.

      iii. Therefore, by Definition A.2, P is *additive* over $\{0, x, x^\perp\}$.

   (b) Then …

$$\begin{aligned}
1 - \mathsf{P}(x^\perp) &= \mathsf{P}(1) - \mathsf{P}(x^\perp) && \text{by } \textit{normalized} \text{ property of } \mathsf{P} && \text{(Definition A.2 page 165)} \\
&= \mathsf{P}(x \vee x^\perp) - \mathsf{P}(x^\perp) && \text{by } \textit{excluded middle} \text{ property of ortho. lat.} \\
&= \mathsf{P}(x) + \mathsf{P}(x^\perp) - \mathsf{P}(x^\perp) && \text{by } \textit{additive} \text{ property of } (\Omega, \mathbb{E}, \mathsf{P}) && \text{(item (2a) page 166)} \\
&= \mathsf{P}(x) && \text{by field property of } (\mathbb{R}, +, \cdot, 0, 1)
\end{aligned}$$

3. Proof for $x \leq y \implies \mathsf{P}(y^\perp) \leq \mathsf{P}(x^\perp)$:

$$\begin{aligned}
x \leq y \implies y^\perp \leq x^\perp && \text{by } \textit{antitone} \text{ property of } \textit{orthocomplemented lattices} \\
\implies \mathsf{P}(y^\perp) \leq \mathsf{P}(x^\perp) && \text{by } \textit{monotone} \text{ property of } \mathsf{P} && \text{(Definition A.2 page 165)}
\end{aligned}$$

[9] Papoulis (1990) page 45 ⟨2-3 Conditional Probability and Indenpendence⟩

[10] property (1): Papoulis (1991) page 21 ⟨(2-11)⟩

BY-NC-ND | COPYRIGHT © 2019 DANIEL J. GREENHOE

Theorem A.2. [11] *Let* $(\Omega, \mathbb{E}, \mathsf{P})$ *be a* PROBABILITY SPACE *(Definition A.2 page 165)*.

THM

$$\left.\begin{array}{l} \boldsymbol{L} \text{ is BOOLEAN} \\ \text{(Definition ?? page ??)} \end{array}\right\} \Longrightarrow \left\{\begin{array}{lllll} 1. & \mathsf{P}(x \vee y) & = & \mathsf{P}(x) + \mathsf{P}(y) - \mathsf{P}(x \wedge y) & \forall x,y \in X \quad \text{and} \\ 2. & \mathsf{P}(x \vee y) & \leq & \mathsf{P}(x) + \mathsf{P}(y) & \forall x,y \in X \quad \text{(BOOLE'S INEQUALITY)} \end{array}\right.$$

✎PROOF:

1. lemma: Proof that $\mathsf{P}((\neg x) \wedge y) = \mathsf{P}(y) - \mathsf{P}(x \wedge y)$:

$$\begin{aligned} \mathsf{P}(y) - \mathsf{P}(xy) &= \mathsf{P}(1 \wedge y) - \mathsf{P}(xy) && \text{by definition of } 1 \text{ and } \wedge \\ &= \mathsf{P}\big[(x \vee x^{\perp})y\big] - \mathsf{P}(xy) && \text{by } \textit{excluded middle} \text{ property of } \textit{Boolean lattices} \\ &= \mathsf{P}(xy \vee x^{\perp}y) - \mathsf{P}(xy) && \text{by } \textit{distributive} \text{ property of } \textit{Boolean lattices} \\ &= \mathsf{P}(xy) + \mathsf{P}(x^{\perp}y) - \mathsf{P}(xy) && \text{because } (xy)(x^{\perp}y) = 0 \text{ and by } \textit{additive} \text{ property} \\ &= \mathsf{P}(x^{\perp}y) \end{aligned}$$

2. Proof that $\mathsf{P}(x \vee y) = \mathsf{P}(x) + \mathsf{P}(y) - \mathsf{P}(x \wedge y)$:

$$\begin{aligned} \mathsf{P}(x \vee y) &= \mathsf{P}(x \vee x^{\perp}y) && \text{by property of } \textit{Boolean lattices} \\ &= \mathsf{P}(x) + \mathsf{P}(x^{\perp}y) && \text{because } (x)(x^{\perp}y) = 0 \text{ and by } \textit{additive} \text{ property} \\ &= \mathsf{P}(x) + \mathsf{P}(y) - \mathsf{P}(x \wedge y) && \text{by item (1) (page 167)} \end{aligned}$$

Theorem A.3 (sum of products)**.** *Let* $(X, \vee, \wedge, 0, 1; \leq)$ *be a* BOUNDED LATTICE, $(\Omega, \mathbb{E}, \mathsf{P})$ *a* PROBABILITY SPACE *(Definition A.2 page 165)*, *and* $\{y, x_1, x_2, x_3, \ldots\}$ *a subset of* X.

THM

$$\left\{\begin{array}{lll} 1. & \boldsymbol{L} \text{ is DISTRIBUTIVE} & \text{and} \\ 2. & \{x_1, x_2, \ldots\} \text{ is a PARTITION of } y & \end{array}\right\} \Longrightarrow \left\{\begin{array}{llllll} 1. & \mathsf{P}(y) & = & \sum_n \mathsf{P}(x_n) & \text{and} \\ 2. & \mathsf{P}(y) & = & \sum_n \mathsf{P}(y \wedge x_n) & \text{and} \\ 3. & \mathsf{P}(z \wedge y) & = & \sum_n \mathsf{P}(z \wedge x_n) & \end{array}\right\}$$

✎PROOF:

1. Proof that P is *additive* (Definition A.2 page 165) on $(\Omega, \mathbb{E}, \mathsf{P})$:

   (a) Proof that $(yx_n) \wedge (yx_m) = 0$ for $n \neq m$:

$$\begin{aligned} (yx_n) \wedge (yx_m) &= y(x_n x_m) && \text{by definition of } \wedge \\ &= y \wedge 0 && \text{by } \textit{mutually exclusive} \text{ property of } \textit{partitions} \\ &= 0 && \text{by } \textit{lower bounded} \text{ property of } \textit{bounded lattices} \end{aligned}$$

   (b) Proof that $\boldsymbol{L}$ is *distributive*: by *distributive* hypothesis

2. Proof that $\mathsf{P}(y) = \sum_n \mathsf{P}(x_n)$

$$\begin{aligned} \mathsf{P}(y) &= \mathsf{P}\big(yx_1 \vee yx_2 \vee \cdots \vee yx_n\big) \\ &= \sum_n \mathsf{P}\big(yx_n\big) && \text{by item (1) and } \textit{additive} \text{ property} && \text{(Definition A.2 page 165)} \\ &= \sum_n \mathsf{P}\big(y|x_n\big)\mathsf{P}(x_n) && \text{by } \textit{conditional probability} && \text{(Definition A.4 page 166)} \end{aligned}$$

[11] Papoulis (1991) page 21 ⟨(2-13)⟩, Feller (1970) pages 22–23 ⟨(7.4),(7.6)⟩

COPYRIGHT © 2019 DANIEL J. GREENHOE

EX



$$\begin{aligned} \mathsf{P}(\varnothing) &= 0 \\ \mathsf{P}(A) &= (1-p)^2 \\ \mathsf{P}(B) &= 2p(1-p) \\ \mathsf{P}(C) &= p^2 \end{aligned} \qquad \begin{aligned} \mathsf{P}(\Omega) &= 1 \\ \mathsf{P}(A \cup B) &= (1-p)^2 + 2p(1-p) \\ \mathsf{P}(A \cup C) &= 2p^2 - 2p + 1 \\ \mathsf{P}(B \cup C) &= -p^2 + 2p \end{aligned}$$

Figure A.2: *Double coin toss* (Example A.2 page 168)

✏

As described in Definition A.2 (page 165), every *probability space* $(\Omega, \mathbb{E}, \mathsf{P})$ contains a probability *measure* $\mathsf{P} : \mathbb{E} \to [0,1]$. This probability *measure* has some basic properties as described in Theorem A.4 (next).

Theorem A.4. *Let* $(\Omega, \mathbb{E}, \mathsf{P})$ *be a* PROBABILITY SPACE. *Let* B *be a set and* $\{B_n | n = 1, 2, \ldots, N\}$ *a set of sets.*

THM

$$\left\{ \begin{array}{l} \{B_n | n = 1, 2, \ldots, N\} \text{ is a} \\ \text{PARTITION of } B. \end{array} \right\} \implies \left\{ \begin{array}{lll} (1). & \mathsf{P}(B) = \sum_{n=1}^{N} \mathsf{P}(B_n) & \forall B \in \mathbb{E} \quad \text{and} \\ (2). & \mathsf{P}(AB) = \sum_{n=1}^{N} \mathsf{P}(AB_n) & \forall A, B \in \mathbb{E} \end{array} \right\}$$

✎PROOF: P is a *measure* and by Definition **??** (page **??**). ✏

Proposition A.2. *Let* $(\Omega, \mathbb{E}, \mathsf{P})$ *be a probability space, and* X *a* RANDOM VARIABLE *with* PROBABILITY DENSITY FUNCTION $\mathsf{p}_{\mathsf{X}}(x)$ *and* CUMULATIVE DISTRIBUTION FUNCTION $\mathsf{c}_{\mathsf{X}}(x)$.

PRP

(1). $\mathsf{c}_{\mathsf{X}}(x)$ *is* MONOTONE *and*
(2). $\mathsf{p}_{\mathsf{X}}(x)$ *is* CONTINUOUS $\implies$ $\mathsf{c}_{\mathsf{X}}(x)$ *is* STRICTLY MONOTONE *and*
(3). $\mathsf{p}_{\mathsf{X}}(x)$ *is* CONTINUOUS $\implies$ $\mathsf{c}_{\mathsf{X}}(x)$ *is* INVERTIBLE

A.4 Examples

Example A.1 (single coin toss). Let ☺ represent "heads" and ● represent "tails" in a coin toss. Let $0 < p < 1$ be the probability of a head. A *probability space* $(\Omega, \mathbb{E}, \mathsf{P})$ for a single coin toss is as follows:

EX

$$\Omega = \{☺, \bullet\}$$

$$\mathbb{E} = \left\{ \underbrace{\varnothing}_{\text{neither}}, \underbrace{\{☺\}}_{\text{heads}}, \underbrace{\{\bullet\}}_{\text{tails}}, \underbrace{\{☺, \bullet\}}_{\text{heads or tails}} \right\}$$

$$\mathsf{P}(X) = \left\{ \begin{array}{lll} 0 & \text{for } X = \varnothing & \text{(neither heads nor tails)} \\ p & \text{for } X = \{☺\} & \text{(heads)} \\ 1-p & \text{for } X = \{\bullet\} & \text{(tails)} \\ 1 & \text{for } X = \{☺, \bullet\} & \text{(either heads or tails)} \end{array} \right\}$$



$$\begin{aligned} \mathsf{P}(\{☺, \bullet\}) &= 1 \\ \mathsf{P}(\{☺\}) &= p \\ \mathsf{P}(\{\bullet\}) &= 1-p \\ \mathsf{P}(\varnothing) &= 0 \end{aligned}$$

Example A.2 (Double coin toss). Let ☺ represent "heads" and ● represent "tails" in a double coin toss in which each toss is *independent* (Definition A.3 page 165) of the other. Let $0 < p < 1$ be the probability of a head. The *probability space* $(\Omega, \mathbb{E}, \mathsf{P})$ is illustrated in Figure A.2 (page 168).

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

✎PROOF:

$\mathsf{P}(\Omega) = 1$	by *normalized* property of P	(Definition A.1 page 164)
$\mathsf{P}(C) = \mathsf{P}\{☺☺\}$	by definition of C	
$= \mathsf{P}(☺)\mathsf{P}(☺)$	by definition of *independence*	(Definition A.3 page 165)
$= p^2$	by definition of p	
$\mathsf{P}(A) = \mathsf{P}\{●●\}$	by definition of A	
$= \mathsf{P}(●)\mathsf{P}(●)$	by definition of *independence*	(Definition A.3 page 165)
$= \{1 - \mathsf{P}(☺)\}\{1 - \mathsf{P}(☺)\}$	by *antitone* property of P	(Theorem A.1 page 166)
$= (1-p)^2$	by definition of p	
$\mathsf{P}(B) = \mathsf{P}\{(●☺), (☺●)\}$	by definition of B	
$= \mathsf{P}\{●☺\} + \mathsf{P}\{☺●\}$	by *additive* property of P	(Definition A.1 page 164)
$= \mathsf{P}\{●\}\mathsf{P}\{☺\} + \mathsf{P}\{☺\}\mathsf{P}\{●\}$	by definition of *independence*	(Definition A.3 page 165)
$= (1-p)p + p(1-p)$	by *antitone* property of P	(Theorem A.1 page 166) and definition of p
$= -2p^2 + p + 1$		
$\mathsf{P}(A \cup B) = \mathsf{P}(A) + \mathsf{P}(B) - \mathsf{P}(A \cap B)$	by Theorem A.2	
$= \mathsf{P}(A) + \mathsf{P}(B) - \mathsf{P}(\varnothing)$		
$= (1-p)^2 + (-2p^2 + p + 1) + 0$	by previous results	
$= -p^2 - p + 1$		
$\mathsf{P}(\varnothing) = 0$	by *nondegenerate* property of P	(Definition A.1 page 164)

✏



Figure A.3: *even/odd die probability space* (Example A.3 page 169)

Example A.3 (even/odd die toss). The *probability space* for an **even/odd die toss**, with $0 < p < 1$ being the probability of the die toss being odd, is illustrated in Figure A.3 (page 169).

✎PROOF:

$\mathsf{P}(\Omega) = 1$	by *normalized* property of P	(Definition A.1 page 164)
$\mathsf{P}(C) = \mathsf{P}\{☺☺\}$	by definition of C	
$= \mathsf{P}(☺)\mathsf{P}(☺)$	by definition of *independence*	(Definition A.3 page 165)
$= p^2$	by definition of p	
$\mathsf{P}(A) = \mathsf{P}\{⚀, ⚂, ⚄\}$	by definition of A	
$= p$	by definition of p	
$\mathsf{P}(B) = \mathsf{P}\{⚁, ⚃, ⚅\}$	by definition of B	
$= \mathsf{P}\{⚀, ⚂, ⚄\}^{\mathsf{c}}$	by definition of set complement c	
$= \mathsf{P}A^{\mathsf{c}}$	by definition of A	

COPYRIGHT © 2019 DANIEL J. GREENHOE | CC BY-NC-ND

$= \mathsf{P}(\neg A)$	by definition of $\neg$	
$= 1 - \mathsf{P}(A)$	by Theorem A.1 page 166	
$= 1 - p$	by definition of p	
$\mathsf{P}(\varnothing) = 0$	by *nondegenerate* property of P	(Definition A.1 page 164)

✏

The two previous *even/odd die* example (Example A.5 page 170) is in essence the same as the *single coin toss* (Example A.1 page 168). The next offers a little more complexity.

EX



$$\begin{aligned} \mathsf{P}(\varnothing) &= 0 & \mathsf{P}(\Omega) &= 1 \\ \mathsf{P}(A) &= (1-p)^2 & \mathsf{P}(A \cup B) &= (1-p)^2 + 2p(1-p) \\ \mathsf{P}(B) &= 2p(1-p) & \mathsf{P}(A \cup C) &= 2p^2 - 2p + 1 \\ \mathsf{P}(C) &= p^2 & \mathsf{P}(B \cup C) &= -p^2 + 2p \end{aligned}$$

Figure A.4: 3-4-2 die example (Example A.4 page 170)

Example A.4. Suppose we have a "fair" die and we are primarily interested in the events of the first three { ⚀, ⚁, ⚂ }, the next two, { ⚃, ⚄ } and the final one { ⚅ }. The resulting *probability space* is illustrated in Figure A.4 (page 170).

The two previous examples (Example A.5 page 170, Example A.4 page 170) illustrate a *probabilty spaces* in which the events are *mutually exclusive*. The (next) illustrates one where events are *not*.

Example A.5. Suppose we have a "fair" die and we are primarily interested in the events of the first four ({⚀,⚁,⚂,⚃}) (that is, whether one roll of the die will produce a value in the set { ⚀, ⚁, ⚂, ⚃ }) and the last three ({⚃,⚄,⚅}) However, these events do not by themselves form a σ-*algebra*. Rather under the $\cap$ and $\cup$ operations, these two events generate a total of eight possible events that together form a σ-*algebra*. The resulting *probability space* is illustrated in Figure A.5 (page 171).

But why go through all the trouble of requiring a σ-algebra? Having a σ-algebra in place ensures that anything we might possibly want to measure *can* be measured. It makes sure all possible combinations are taken into account. And why go through the additional trouble of requiring a measure space? With a measure space available, expressing the measure over a complex set is often greatly simplified because the measure space provides nice algebraic properties (namely the σ-*additive* property). Example A.6 (next) illustrates how a rather complex σ-algebra (64 elements) can be compactly represented in a measure space.

Example A.6. Suppose we have a "fair" dice and we are interested in measuring over the power set of events (largest possible algebra—$2^6 = 64$ events). This leads to the probability space $(\Omega, \mathbb{E}, \mathsf{P})$ where

EX

Ω	$= \{⚀,⚁,⚂,⚃,⚄,⚅\}$	
$\mathbb{E}$	$= \mathcal{P}(\Omega)$	(the power-set of Ω)
$\mathsf{P}(e)$	$= \frac{1}{6}\lvert e\rvert$	($\frac{1}{6}$ times the number of possible outcomes in event e)

Example A.7 (Gaussian distribution on $\mathbb{R}$). Let $\boldsymbol{B}$ be the *Borel algebra on* $\mathbb{R}$. Let $\boldsymbol{L} \triangleq (\boldsymbol{B}, \subseteq)$ be the lattice formed by the elements of $\boldsymbol{B}$—this lattice is a *Boolean algebra*. Let

$$\mathsf{P}(A) \triangleq \frac{1}{\sqrt{2\pi\sigma^2}} \int_A e^{\frac{x^2}{2\sigma^2}}\, dx \text{ for } A \subseteq \boldsymbol{B}$$

and where $\int$ is the *Lebesgue integral* (Definition **??** page **??**). Then $(\boldsymbol{L}, \mathsf{P})$ is a **probability space**.

BY-NC-ND Copyright © 2019 Daniel J. Greenhoe



Figure A.5: First 4 / last 3 die example (Example A.5 page 170)

Example A.8 (Gaussian noise). Let $X \sim \mathsf{N}\left(0, \sigma^2\right)$ be a random variable with Gaussian distribution. We can construct the following probability space $(\Omega, \mathbb{E}, \mathsf{P})$:

EX

$$\begin{aligned}
\Omega &= \mathbb{R} \\
\mathbb{E} &= \{\varnothing, \Omega\} \cup \{(a,b) | a, b \in \mathbb{R}, a < b\} \\
\mathsf{P}x &= \begin{cases} 0 & \text{for } x = \varnothing \\ 1 & \text{for } x = \Omega \\ \frac{1}{\sqrt{2\pi\sigma^2}} \int_a^b e^{\frac{x^2}{2\sigma^2}}\, \mathrm{d}x & \text{otherwise} \end{cases}
\end{aligned}$$

Example A.9. The set of outcomes Ω can also be a set of waveforms:

EX

$\Omega =$ 

$$\begin{aligned}
\mathbb{E} &= \mathcal{P}(\Omega) \\
\mathsf{P}(e) &= \tfrac{1}{7}|e|
\end{aligned}$$

A.5 Probability subspaces

Example A.10. Suppose a random process is capable of producing three values $\Omega \triangleq \{x, y, z\}$. There are five *algebras of sets* on Ω and therefore five probability spaces $(\Omega, \mathbb{E}_n, \mathsf{P})$ on Ω with the five values of $\mathbb{E}_n$ listed below: [12]

[12] *algebra of sets*: Definition ?? page ??

Copyright © 2019 Daniel J. Greenhoe



Figure A.6: Euclidean 3-dimensional space partitioned as a power lattice



$$
\begin{aligned}
\mathbb{E}_1 &= \{ \quad \varnothing, & & & & & & & X \quad \} \\
\mathbb{E}_2 &= \{ \quad \varnothing, & \{x\}, & & & & & \{y,z\}, & X \quad \} \\
\mathbb{E}_3 &= \{ \quad \varnothing, & & \{y\}, & & & \{x,z\}, & & X \quad \} \\
\mathbb{E}_4 &= \{ \quad \varnothing, & & & \{z\}, & \{x,y\}, & & & X \quad \} \\
\mathbb{E}_5 &= \{ \quad \varnothing, & \{x\}, & \{y\}, & \{z\}, & \{x,y\}, & \{x,z\}, & \{y,z\}, & X \quad \}
\end{aligned}
$$

Suppose the samples of Ω are generated by a physical process such that they are all equally likely to occur. Then by varying the *σ-algebra* $\mathbb{E}_n$ effectively varies the probability distribution of the probability space $(\Omega, \mathbb{E}_n, \mathsf{P})$. This is illustrated by the figure to the left.

BY-NC-ND Copyright © 2019 Daniel J. Greenhoe

APPENDIX B

PROBABILITY DENSITY FUNCTIONS

❝While writing my book I had an argument with Feller. He asserted that everyone said "random variable" and I asserted that everyone said "chance variable." We obviously had to use the same name in our books, so we decided the issue by a stochastic procedure. That is, we tossed for it and he won.❞

Joseph Leonard Doob (1910–2004), pioneer of and key contributor to mathematical probability[1]

B.1 Random variables

The concept of the *random variable* is widely used in probability and random processes. Before discussing what a *random variable is*, note two things that a *random variable* is *not* (next remark).

Remark B.1. [2] As pointed out by others, the term "random variable" is a "misnomer":

REM

- A *random variable* is **not random**.
- A *random variable* is **not a variable**.

What is it then? It is a *function* (next definition). In particular, it is a function that maps from an underlying stochastic process into $\mathbb{R}$. Any "*randomness*" (whatever that means) it may *appear* to have comes from the stochastic process it is mapping *from*. But the function itself (the *random*

[1] quote: Snell (1997), page 307, Snell (2005), page 251
image: http://www.dartmouth.edu/~chance/Doob/conversation.html

[2] Miller (2006) page 130, Feldman and Valdez-Flores (2010) page 4 ⟨"The name "random variable" is actually a misnomer, since it is not random and not a variable.…the *random variable* simply maps each point (outcome) in the sample space to a number on the real line…Technically, the space into which the *random variable* maps the sample space may be more general than the real line…"⟩, Curry and Feldman (2010) page 4, Trivedi (2016) page 2.1 ⟨"The term "random variable" is actually a misnomer, since a *random variable* X is really a function whose domain is the sample space S, and whose range is the set of all real numbers, $\mathbb{R}$."⟩

variable itself) is very deterministic and well-defined. What gives it the appearance of being random is that the outcome ω of the experiment appears to be random to the observer. So the *random variable* $\mathsf{X}(\omega)$ is simply a function of an underlying mechanism that appears to be random.

Definition B.1. [3] *Let* $(\Omega, \mathbb{E}, \mathsf{P})$ *be a* PROBABILITY SPACE *(Definition A.2 page 165)*.

DEF *A* ***random variable*** X *is any function in* $\mathbb{R}^{\Omega}$.

B.2 Probability distributions

The probability information about σ-algebra $\mathbb{E}$ in a *probability space* (Definition A.2 page 165) is completely specified by *measure* P. However, sometimes it is more convenient to express this same *measure* information in terms of the *probability density function* or the *cummulative distribution function* of the *probability space*.

Definition B.2. [4] *Let* X *be a* RANDOM VARIABLE *(Definition B.1 page 174) on a* PROBABILITY SPACE $(\Omega, \mathbb{E}, \mathsf{P})$.

DEF X *has* ***cummulative distribution function*** ***(cdf)*** $\mathsf{c}_{\mathsf{X}}(x)$ *if*

$$\mathsf{c}_{\mathsf{X}}(x) \triangleq \mathsf{P}\{x \in \mathbb{E} | \mathsf{X} < x\}$$

X ***probability density function*** ***(pdf)*** $\mathsf{p}_{\mathsf{X}}(x)$ *if*

$$\mathsf{p}_{\mathsf{X}}(x) \triangleq \frac{\mathsf{d}}{\mathsf{d}x}\mathsf{c}_{\mathsf{X}}(x) \triangleq \frac{\mathsf{d}}{\mathsf{d}x}\mathsf{P}\{x \in \mathbb{E} | \mathsf{X} < x\}$$

Remark B.2. Suppose X be a *random variable* on a *probability space* $(\Omega, \mathbb{E}, \mathsf{P})$. Note that

- Both X and $\mathbb{E}$ are *functions*.
- But X is a function that maps from Ω to $\mathbb{R}$,
- whereas P is a function that maps from $\mathbb{E}$ to $\mathbb{R}$.

Definition B.3. *Let* $(\Omega, \mathbb{E}, \mathsf{P})$ *be a* PROBABILITY SPACE *(Definition A.2 page 165) and* X *and* $\mathsf{Y} : \Omega \to \mathbb{R}$ *random variables. Then a* ***joint probability density function*** $\mathsf{p}_{\mathsf{X}} : \mathbb{E} \times \Omega \to [0 : 1]$ *and a* ***joint cumulative distribution function*** $\mathsf{c}_{\mathsf{X}} : \mathbb{E} \times \Omega \to [0 : 1]$ *are defined as*

DEF

$$\mathsf{c}_{\mathsf{XY}}(x, y) \triangleq \mathsf{P}\{\mathsf{X} \le x | \mathsf{Y} \le y\} \quad \text{(JOINT CUMULATIVE DISTRIBUTION FUNCTION)}$$

$$\mathsf{p}_{\mathsf{XY}}(x, y) \triangleq \frac{\mathsf{d}}{\mathsf{d}y}\frac{\mathsf{d}}{\mathsf{d}x}\mathsf{c}_{\mathsf{XY}}(x, y) \quad \text{(JOINT PROBABILITY DENSITY FUNCTION)}$$

Definition B.4. *Let* $(\Omega, \mathbb{E}, \mathsf{P})$ *be a* PROBABILITY SPACE *(Definition A.2 page 165) and* X *a random variable. Then a* ***conditional probability density function*** $\mathsf{p}_{\mathsf{X}} : \mathbb{E} \times \Omega \to [0 : 1]$ *and a* ***conditional cumulative distribution function*** $\mathsf{c}_{\mathsf{X}} : \mathbb{E} \times \Omega \to [0 : 1]$ *are defined as*

DEF

$$\mathsf{c}_{\mathsf{X}}(x|y) \triangleq \mathsf{P}\{\mathsf{X} \le x | \mathsf{Y} = y\} \quad \text{(CONDITIONAL CUMULATIVE DISTRIBUTION FUNCTION—CDF)}$$

$$\mathsf{p}_{\mathsf{X}}(x|y) \triangleq \frac{\mathsf{d}}{\mathsf{d}x}\mathsf{c}_{\mathsf{X}}(x|y) \quad \text{(CONDITIONAL PROBABILITY DENSITY FUNCTION—PDF)}$$

B.3 Properties

Definition B.2 (page 174) defines the pdf and cdf of a *probability space* $(\Omega, \mathbb{E}, \mathsf{P})$ in terms of *measure* P. Conversely, the probability *measure* $\mathsf{P}\{a \le \mathsf{X} < b\}$ of an event $\{a \le \mathsf{X} < b\}$ can be expressed in terms of either the pdf or cdf.

[3] Papoulis (1991), page 63
[4] von der Linden et al. (2014) page 93 ⟨Definitions 7.1, 7.2⟩

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

Proposition B.1. *Let* X *a* RANDOM VARIABLE *with* PDF p_{X} *and* CDF c_{X} *(Definition B.2 page 174) on the* PROBABILITY SPACE $(\Omega, \mathbb{E}, \mathsf{P})$ *(Definition A.2 page 165)*.

PRP

$$\left\{\begin{array}{ll} (1). & \mathsf{c}_{\mathsf{X}}(x) \text{ and } \mathsf{c}_{\mathsf{Y}}(y) \text{ are CONTINUOUS} \quad \text{OR} \\ (2). & \mathsf{p}_{\mathsf{X}}(x) \text{ and } \mathsf{p}_{\mathsf{Y}}(y) \text{ are CONTINUOUS} \end{array}\right\}$$

$$\Longrightarrow \left\{\begin{array}{rcl} \mathsf{p}_{\mathsf{X}}(x) & = & \lim_{\varepsilon\to 0}\frac{1}{\varepsilon}\mathsf{P}\{x \le \mathsf{X} < x+\varepsilon\} \\ \mathsf{p}_{\mathsf{XY}}(x,y) & = & \lim_{\varepsilon\to 0}\frac{1}{\varepsilon}\mathsf{P}\{x \le \mathsf{X} < x+\varepsilon \wedge y \le \mathsf{Y} < y+\varepsilon\} \end{array}\right\}$$

✎PROOF:

$$\begin{aligned} \mathsf{p}_{\mathsf{X}}(x) &\triangleq \frac{\mathsf{d}}{\mathsf{d}x}\mathsf{c}_{\mathsf{X}}(x) && \text{by definition of } \mathsf{p}_{\mathsf{X}} && \text{(Definition B.2 page 174)} \\ &= \lim_{\varepsilon\to 0}\frac{1}{\varepsilon}\mathsf{P}\{x \in \mathbb{R} | x \le \mathsf{X} < x+\varepsilon\} && \text{by definition of } \frac{\mathsf{d}}{\mathsf{d}x} && \text{(Definition ?? page ??)} \end{aligned}$$

✏

Theorem B.1. *Let* $(\Omega, \mathbb{E}, \mathsf{P})$ *be a probability space,* X *be a random variable, and* (a, b) *a real interval.*

THM

$$\left\{\begin{array}{ll} (1). & \mathsf{c}_{\mathsf{X}}(x) \text{ is CONTINUOUS} \quad \text{OR} \\ (2). & \mathsf{p}_{\mathsf{X}}(x) \text{ is CONTINUOUS} \end{array}\right\} \Longrightarrow \left\{ \mathsf{P}\{a < \mathsf{X} \le b\} = \mathsf{c}_{\mathsf{X}}(b) - \mathsf{c}_{\mathsf{X}}(a) = \int_a^b \mathsf{p}_{\mathsf{X}}(x)\,\mathsf{d}x \right\}$$

✎PROOF:

$$\begin{aligned} \mathsf{P}\{a < \mathsf{X} \le b\} &= \mathsf{P}\{\mathsf{X} \le b\} - \mathsf{P}\{\mathsf{X} < a\} && \text{by } \textit{sum of products} && \text{(Theorem A.3 page 167)} \\ &= \mathsf{P}\{\mathsf{X} \le b\} - \mathsf{P}\{\mathsf{X} \le a\} && \text{by } \textit{continuity} \text{ hypothesis} \\ &\triangleq \mathsf{c}_{\mathsf{X}}(b) - \mathsf{c}_{\mathsf{X}}(a) && \text{by definition of } \mathsf{c}_{\mathsf{X}} && \text{(Definition B.2 page 174)} \end{aligned}$$

$$\begin{aligned} \int_a^b \mathsf{p}_{\mathsf{X}}(x)\,\mathsf{d}x &\triangleq \int_a^b \left[\frac{\mathsf{d}}{\mathsf{d}x}\mathsf{c}_{\mathsf{X}}(x)\right]\mathsf{d}x && \text{by definition of } \mathsf{p}_{\mathsf{X}} && \text{(Definition B.2 page 174)} \\ &= \mathsf{c}_{\mathsf{X}}(x)\big|_{x=b} - \mathsf{c}_{\mathsf{X}}(x)\big|_{x=a} && \text{by } \textit{Fundamental theorem of calculus} \\ &= \mathsf{c}_{\mathsf{X}}(b) - \mathsf{c}_{\mathsf{X}}(a) \end{aligned}$$

✏

Theorem B.2. *Let* $(\Omega, \mathbb{E}, \mathsf{P})$ *be a* PROBABILITY SPACE, X *be a* RANDOM VARIABLE, *and* $(a : b)$ *a* REAL INTERVAL.

THM

$$\mathsf{P}\{a \le \mathsf{X} < b\} = \int_a^b \mathsf{p}_{\mathsf{X}}(x)\,\mathsf{d}x = \int_{-\infty}^b \mathsf{c}_{\mathsf{X}}(x)\,\mathsf{d}x - \int_{-\infty}^a \mathsf{c}_{\mathsf{X}}(x)\,\mathsf{d}x$$

The properties of the pdf follow closely the properties of *measure* P.

Theorem B.3. [5]

THM

$$\left\{\begin{array}{ll} (A). & \mathsf{c}_{\mathsf{X}}(x) \text{ is CONTINUOUS} \quad \text{OR} \\ (B). & \mathsf{p}_{\mathsf{X}}(x) \text{ is CONTINUOUS} \end{array}\right\} \Longrightarrow \left\{\begin{array}{lrcl} (1). & \mathsf{p}_{\mathsf{X}|\mathsf{Y}}(x|y) & = & \dfrac{\mathsf{p}_{\mathsf{XY}}(x,y)}{\mathsf{p}_{\mathsf{Y}}(y)} \quad \text{and} \\ (2). & \mathsf{p}_{\mathsf{X}}(x) & = & \displaystyle\int_{y\in\mathbb{R}} \mathsf{p}_{\mathsf{XY}}(x,y)\,\mathsf{d}y \end{array}\right\}$$

[5] Papoulis (1990) page 158 ⟨Auxiliary Variable⟩

COPYRIGHT © 2019 DANIEL J. GREENHOE

✎PROOF:

$$
\begin{aligned}
\mathsf{p}_{\mathsf{X}|\mathsf{Y}}(x|y) &\triangleq \frac{\mathsf{d}}{\mathsf{d}x}\mathsf{c}_{\mathsf{X}|\mathsf{Y}}(x|y) && \text{by definition of } \mathsf{c}_{\mathsf{X}} && \text{(Definition A.4 page 166)}\\
&\triangleq \lim_{\varepsilon\to 0}\frac{1}{\varepsilon}\mathsf{P}\{x \le \mathsf{X} < x+\varepsilon|\mathsf{Y}=y\} && \text{by definition of } \frac{\mathsf{d}}{\mathsf{d}x} && \text{(Definition ?? page ??)}\\
&\triangleq \lim_{\varepsilon\to 0}\frac{1}{\varepsilon}\frac{\mathsf{P}\{(x \le \mathsf{X} < x+\varepsilon)\wedge(\mathsf{Y}=y)\}}{\mathsf{P}\{\mathsf{Y}=y\}} && \text{by definition of } \mathsf{P}\{A|B\} && \text{(Definition A.4 page 166)}\\
&= \lim_{\varepsilon\to 0}\frac{1}{\varepsilon}\frac{\mathsf{P}\{(x \le \mathsf{X} < x+\varepsilon)\wedge(y \le \mathsf{Y} < y+\varepsilon)\}}{\mathsf{P}\{y \le \mathsf{Y} < y+\varepsilon\}} && \text{by } \textit{continuity} \text{ hypothesis}\\
&= \frac{\lim_{\varepsilon\to 0}\frac{1}{\varepsilon}\mathsf{P}\{(x \le \mathsf{X} < x+\varepsilon)\wedge(y \le \mathsf{Y} < y+\varepsilon)\}}{\lim_{\varepsilon\to 0}\mathsf{P}\{y \le \mathsf{Y} < y+\varepsilon\}} && \text{by property of } \lim_{\varepsilon\to 0}\\
&= \frac{\mathsf{p}_{\mathsf{XY}}(x,y)}{\mathsf{p}_{\mathsf{Y}}(y)} && \text{by Proposition B.1 page 175}
\end{aligned}
$$

$$
\begin{aligned}
\int_{y\in\mathbb{R}}\mathsf{p}_{\mathsf{XY}}(x,y)\,\mathsf{d}y &\triangleq \int_{y\in\mathbb{R}}\left[\frac{\mathsf{d}}{\mathsf{d}y}\frac{\mathsf{d}}{\mathsf{d}x}\mathsf{c}_{\mathsf{XY}}(x,y)\right]\mathsf{d}y && \text{by definition of } \mathsf{p}_{\mathsf{X}} && \text{(Definition B.2 page 174)}\\
&= \frac{\mathsf{d}}{\mathsf{d}x}\mathsf{c}_{\mathsf{XY}}(x,y)\\
&\triangleq \lim_{\varepsilon\to 0}\frac{1}{\varepsilon}\int_{y\in\mathbb{R}}\mathsf{P}\{x \le \mathsf{X} < x+\varepsilon, y \le \mathsf{Y} < y+\varepsilon\}\,\mathsf{d}y && \text{by definition of } \frac{\mathsf{d}}{\mathsf{d}x} && \text{(Definition ?? page ??)}\\
&= \lim_{\varepsilon\to 0}\frac{1}{\varepsilon}\mathsf{P}\{x \le \mathsf{X} < x+\varepsilon\}\\
&= \mathsf{p}_{\mathsf{X}}(x) && \text{by Proposition B.1 page 175}
\end{aligned}
$$

✏

Theorem B.4.

THM

$$
\begin{aligned}
\mathsf{c}_{\mathsf{X}}(\sup\mathbb{R}) &= 1\\
\mathsf{c}_{\mathsf{X}}(\inf\mathbb{R}) &= 0
\end{aligned}
$$

✎PROOF:

$$
\begin{aligned}
\mathsf{c}_{\mathsf{X}}(\sup\mathbb{R}) &\triangleq \mathsf{P}\{\mathsf{X} \le \sup\mathbb{R}\} && \text{by definition of } \mathsf{c}_{\mathsf{X}} && \text{(Definition B.2 page 174)}\\
&= 1\\
\mathsf{c}_{\mathsf{X}}(\inf\mathbb{R}) &\triangleq \mathsf{P}\{\mathsf{X} \le \inf\mathbb{R}\} && \text{by definition of } \mathsf{c}_{\mathsf{X}} && \text{(Definition B.2 page 174)}\\
&= 0
\end{aligned}
$$

✏

The properties of the pdf follow closely the properties of measure P.

Theorem B.5.

THM

$$
\mathsf{c}_{\mathsf{X}|\mathsf{Y}}(x|y) = \frac{\frac{\mathsf{d}}{\mathsf{d}y}\mathsf{c}_{\mathsf{XY}}(x,y)}{\mathsf{p}_{\mathsf{Y}}(y)} \qquad \mathsf{p}_{\mathsf{X}|\mathsf{Y}}(x|y) = \frac{\mathsf{p}_{\mathsf{XY}}(x,y)}{\mathsf{p}_{\mathsf{Y}}(y)}
$$

COPYRIGHT © 2019 DANIEL J. GREENHOE

✎PROOF:

$$
\begin{aligned}
\mathsf{c}_{\mathsf{X}|\mathsf{Y}}(x|y) &\triangleq \mathsf{P}\{\mathsf{X} \le x \mid \mathsf{Y} = y\} && \text{by definition of } \mathsf{c}_{\mathsf{X}|\mathsf{Y}} && \text{(Definition B.4 page 174)}\\
&\triangleq \frac{\mathsf{P}\{\mathsf{X} \le x|\mathsf{Y} = y\}}{\mathsf{P}\{\mathsf{Y} = y\}} && \text{by definition of } \mathsf{P}\{\mathsf{X}|\mathsf{Y}\} && \text{(Definition A.4 page 166)}\\
&= \lim_{\varepsilon\to 0} \frac{\mathsf{P}\{\mathsf{X} \le x|y < \mathsf{Y} \le y+\varepsilon\}}{\mathsf{P}\{y < \mathsf{Y} \le y+\varepsilon\}}\\
&= \lim_{\varepsilon\to 0} \frac{[\mathsf{P}\{\mathsf{X} \le x|\mathsf{Y} \le y+\varepsilon\} - \mathsf{P}\{\mathsf{X} \le x|\mathsf{Y} \le y\}]/\varepsilon}{[\mathsf{P}\{\mathsf{Y} \le y+\varepsilon\} - \mathsf{P}\{\mathsf{Y} \le y\}]/\varepsilon}\\
&\triangleq \lim_{\varepsilon\to 0} \frac{\left[\mathsf{c}_{\mathsf{XY}}(x, y+\varepsilon) - \mathsf{c}_{\mathsf{XY}}(x, y)\right]/\varepsilon}{\left[\mathsf{c}_{\mathsf{Y}}(y+\varepsilon) - \mathsf{c}_{\mathsf{Y}}(y)\right]/\varepsilon} && \text{by definition of } \mathsf{c}_{\mathsf{XY}} && \text{(Definition B.3 page 174)}\\
&\triangleq \frac{\frac{\mathsf{d}}{\mathsf{d}y}\mathsf{c}_{\mathsf{XY}}(x, y)}{\frac{\mathsf{d}}{\mathsf{d}y}\mathsf{c}_{\mathsf{Y}}(y)} && \text{by definition of } \frac{\mathsf{d}}{\mathsf{d}y}\mathsf{f}(y)\\
&\triangleq \frac{\frac{\mathsf{d}}{\mathsf{d}y}\mathsf{c}_{\mathsf{XY}}(x, y)}{\mathsf{p}_{\mathsf{Y}}(y)} && \text{by definition of } \mathsf{p}_{\mathsf{Y}} && \text{(Definition B.2 page 174)}\\
&= \frac{\frac{\mathsf{d}}{\mathsf{d}y}\mathsf{c}_{\mathsf{XY}}(x, y)}{\mathsf{p}_{\mathsf{Y}}(y)} && \text{because } y \text{ is fixed}
\end{aligned}
$$

$$
\begin{aligned}
\mathsf{p}_{\mathsf{X}|\mathsf{Y}}(x|y) &\triangleq \frac{\mathsf{d}}{\mathsf{d}x}\mathsf{c}_{\mathsf{X}|\mathsf{Y}}(x|y) && \text{by definition of } \mathsf{p}_{\mathsf{X}|\mathsf{Y}} && \text{(Definition B.4 page 174)}\\
&= \frac{\mathsf{d}}{\mathsf{d}x}\frac{\frac{\mathsf{d}}{\mathsf{d}y}\mathsf{c}_{\mathsf{XY}}(x, y)}{\mathsf{p}_{\mathsf{Y}}(y)} && \text{by previous result}\\
&= \frac{\frac{\mathsf{d}}{\mathsf{d}x}\frac{\mathsf{d}}{\mathsf{d}y}\mathsf{c}_{\mathsf{XY}}(x, y)}{\mathsf{p}_{\mathsf{Y}}(y)} && \text{because } \mathsf{p}_{\mathsf{Y}}(y) \text{ is not a function of } x\\
&\triangleq \frac{\mathsf{p}_{\mathsf{XY}}(x, y)}{\mathsf{p}_{\mathsf{Y}}(y)} && \text{by definition of } \mathsf{p}_{\mathsf{XY}}(x, y) && \text{(Definition B.3 page 174)}
\end{aligned}
$$

✏

Theorem B.6. *Let $(\Omega, \mathbb{E}, \mathsf{P})$ be a probability space.*

THM

$$
\begin{aligned}
\int_{x\in\mathbb{R}} \mathsf{p}_{\mathsf{X}}(x)\,\mathsf{d}x &= 1 \qquad\qquad \int_{x\in\mathbb{R}} \mathsf{p}_{\mathsf{X}|\mathsf{Y}}(x|y)\,\mathsf{d}x = 1\\
\int_{y\in\mathbb{R}} \mathsf{p}_{\mathsf{XY}}(x, y)\,\mathsf{d}y &= \mathsf{p}_{\mathsf{X}}(x) \quad \forall x\in\Omega \qquad \int_{x\in\mathbb{R}}\int_{y\in\mathbb{R}} \mathsf{p}_{\mathsf{XY}}(x, y)\,\mathsf{d}y\,\mathsf{d}x = 1
\end{aligned}
$$

✎PROOF:

$$
\begin{aligned}
\int_{\mathbb{R}} \mathsf{p}_{\mathsf{X}}(x)\,\mathsf{d}x &= \mathsf{c}_{\mathsf{X}}(\sup\mathbb{R}) - \mathsf{c}_{\mathsf{X}}(\inf\mathbb{R}) && \text{by Theorem B.1 page 175}\\
&= 1 - 0\\
&= 1 && \text{because } 0 \text{ is the additive identity element in } (\mathbb{R}, +, \cdot, 0, 1)\\
\int_{x\in\mathbb{R}} \mathsf{p}_{\mathsf{X}|\mathsf{Y}}(x|y)\,\mathsf{d}x &\triangleq \int_{x\in\mathbb{R}} \frac{\mathsf{d}}{\mathsf{d}x}\mathsf{c}_{\mathsf{X}|\mathsf{Y}}(x|y)\,\mathsf{d}x && \text{by definition of } \mathsf{p}_{\mathsf{X}|\mathsf{Y}}(x|y) \text{ (Definition B.4 page 174)}\\
&= \mathsf{c}_{\mathsf{X}|\mathsf{Y}}(\sup\mathbb{R}|y) - \mathsf{c}_{\mathsf{X}|\mathsf{Y}}(\inf\mathbb{R}|y) && \text{by } \textit{Fundamental theorem of calculus}
\end{aligned}
$$

COPYRIGHT © 2019 DANIEL J. GREENHOE

$$
\begin{aligned}
&= 1 - 0 \\
&= 1 && \text{because } 0 \text{ is the additive identity element in } (\mathbb{R}, +, \cdot, 0, 1) \\
\int_{y\in\mathbb{R}} \mathsf{p}_{\mathsf{XY}}(x,y)\,\mathsf{d}y &= \int_{y\in\mathbb{R}} \mathsf{p}_{\mathsf{YX}}(y,x)\,\mathsf{d}y \\
&= \int_{y\in\mathbb{R}} \mathsf{p}_{\mathsf{Y|X}}(y|x)\mathsf{p}_{\mathsf{X}}(x)\,\mathsf{d}y && \text{by Theorem B.5 page 176} \\
&= \mathsf{p}_{\mathsf{X}}(x)\int_{y\in\mathbb{R}} \mathsf{p}_{\mathsf{Y|X}}(y|x)\,\mathsf{d}y && \text{because } \mathsf{p}_{\mathsf{X}}(x) \text{ is not a function of } y \\
&= \mathsf{p}_{\mathsf{X}}(x)\cdot 1 && \text{by previous result} \\
&= \mathsf{p}_{\mathsf{X}}(x) && \text{because } 1 \text{ is the multiplicative identity element in } (\mathbb{R}, +, \cdot, 0, 1)
\end{aligned}
$$

$$
\begin{aligned}
\int_{x\in\mathbb{R}}\int_{y\in\mathbb{R}} \mathsf{p}_{\mathsf{XY}}(x,y)\,\mathsf{d}y\,\mathsf{d}x &= \int_{x\in\mathbb{R}} \mathsf{p}_{\mathsf{X}}(x)\,\mathsf{d}x && \text{by previous result} \\
&= 1 && \text{by previous result}
\end{aligned}
$$

✏

Copyright © 2019 Daniel J. Greenhoe

APPENDIX C

SOME PROBABILITY DENSITY FUNCTIONS

C.1 Discrete distributions

Example C.1. [1] Suppose we throw two "fair" dice and want to know the probabilities of their sum. Let X represent the sum of the face values of the two dice. The resulting probability distribution is illustrated in Figure C.1 (page 180) and has probability space as follows:

EX

$$\begin{aligned} \Omega &= \{\boxed{\cdot}\boxed{\cdot},\ \boxed{\cdot}\boxed{\cdot},\ \boxed{\cdot}\boxed{\cdot},\ \ldots,\ \boxed{\cdot}\boxed{\cdot}\} \\ \mathbb{E} &= \left\{2^{\{X=n \mid n=2,3,\ldots,10,11,\text{ or } 12\}}\right\} \\ \mathsf{P}(e) &= \frac{1}{36}\,|\,e\,| \end{aligned}$$

C.2 Continuous distributions

C.2.1 Uniform distribution

Definition C.1. *The* ***uniform distribution*** $\mathsf{p}_{\mathsf{X}}(x)$ *is defined as*

DEF

$$\mathsf{p}_{\mathsf{X}}(x) \triangleq \left\{ \begin{array}{ll} 1 & \textit{for}\, 0 < x \leq 1 \\ 0 & \textit{otherwise} \end{array} \right\}$$

Note that although "simple" in form, in light of *Wold's Theorem*, the value of the *uniform distribution* should *not* be taken lightly.

[1] Osgood (2002)



Figure C.1: Probability distribution for two dice (see Example C.1 page 179)

C.2.2 Gaussian distribution

❝Tout le monde y croit cependant, me disait un jour M. Lippmann, car les expérimentateurs s'irnaginent que c'est un théorème de mathématiques, et les mathématiciens que c'est un fait expérimental.❞

❝Everyone believes in it [(the normal distribution)] however, said to me one day Mr. Lippmann, because the experimenters imagine that it is a theorem of mathematics, and mathematicians that it is an experimental fact.❞

Bernard A. Lippmann as told by Henri Poincaré [2]



Figure C.2: Gaussian pdf with $\mu = 0$ and $\sigma \in [0.1, 2]$.

Definition C.2.

[2] quote: Poincaré (1912), page 171
translation: assisted by Google Translate
image:

BY-NC-ND Copyright © 2019 Daniel J. Greenhoe

DEF

*The **Gaussian distribution** (or **normal distribution**) has pdf*

$$\mathsf{p}_{\mathsf{X}}(x) \triangleq \frac{1}{\sqrt{2\pi\sigma^2}} e^{\frac{-(x-\mu)^2}{2\sigma^2}}$$

A random variable X *with this distribution is denoted*

$$\mathsf{X} \sim \mathsf{N}\left(\mu, \sigma^2\right)$$

The function $\mathsf{Q}(x)$ *is defined as the area under a Gaussian PDF with zero mean and variance equal to one from* x *to infinity such that*

$$\mathsf{Q}(x) \triangleq \frac{1}{\sqrt{2\pi}} \int_x^\infty e^{\frac{-u^2}{2}} \, \mathsf{d}u$$

C.2.3 Gamma distribution

Definition C.3. [3] *Let* $b \in \mathbb{R}$. *The **gamma function*** $\Gamma(b)$ *is*

DEF

$$\Gamma(b) \triangleq \int_0^\infty x^{b-1} e^{-x} \, \mathsf{d}x$$

Proposition C.1. [4] *Let* $b \in \mathbb{R}$ *and* $n \in \mathbb{N}$.

PRP

$$\begin{aligned} \Gamma(b) &= (b-1)\Gamma(b-1) \\ \Gamma(n) &= (n-1)! \end{aligned}$$

✎PROOF: Let

$$\begin{aligned} u &= x^{b-1} & \mathsf{d}u &= (b-1)x^{b-1} \, \mathsf{d}x \\ \mathsf{d}v &= e^{-x} \, \mathsf{d}x & v &= -e^{-x} \end{aligned}$$

$$\begin{aligned} \Gamma(b) &\triangleq \int_0^\infty x^{b-1} e^{-x} \, \mathsf{d}x \\ &= \int_{x=0}^\infty u \, \mathsf{d}v \\ &= uv|_{x=0}^\infty - \int_{x=0}^\infty v \, \mathsf{d}u \\ &= -x^{b-1}e^{-x}|_{x=0}^\infty + (b-1)\int_{x=0}^\infty e^{-x}x^{b-1} \, \mathsf{d}x \\ &= (-0+0) + (b-1)\Gamma(b-1) \end{aligned}$$

Note that

$$\Gamma(1) = \int_0^\infty x^{1-1}e^{-x} \, \mathsf{d}x = \int_0^\infty e^{-x} \, \mathsf{d}x = -e^{-x}|_0^\infty = -0+1 = 1$$

$$\begin{aligned} \Gamma(n) &= (n-1)\Gamma(n-1) \\ &= (n-1)(n-2)\Gamma(n-2) \\ &= (n-1)(n-2)(n-3)\Gamma(n-3) \\ &\vdots \\ &= (n-1)(n-2)(n-3)\cdots(1)\Gamma(1) \\ &= (n-1)(n-2)(n-3)\cdots(1) \\ &\triangleq (n-1)! \end{aligned}$$

[3] Papoulis (1991), page 79, Ross (1998), page 222
[4] Ross (1998), page 223

COPYRIGHT © 2019 DANIEL J. GREENHOE
CC BY-NC-ND

Definition C.4. *A **Gamma distribution** (b, λ) has pdf*

DEF
$$\mathsf{p}_{\mathsf{X}}(x) \triangleq \frac{\lambda}{\Gamma(b)} e^{-\lambda x}(\lambda x)^{b-1}$$

Theorem C.1. [5] *Let* X *and* Y *be* RANDOM VARIABLES *on a* PROBABILITY SPACE $(\Omega, \mathbb{E}, \mathsf{P})$.

THM
$$\left\{\begin{array}{lll} (A). & \mathsf{X} \text{ and } \mathsf{Y} \text{ are INDEPENDENT} & \text{and} \\ (B). & \mathsf{X} \text{ has GAMMA DISTRIBUTION } (a, \lambda) & \text{and} \\ (C). & \mathsf{Y} \text{ has GAMMA DISTRIBUTION } (b, \lambda) & \text{and} \\ (D). & \mathsf{Z} \triangleq \mathsf{X} + \mathsf{Y} & \end{array}\right\} \Longrightarrow \left\{\begin{array}{l} \mathsf{Z} \text{ has Gamma distribution} \\ (a+b, \lambda). \end{array}\right\}$$

✎PROOF:

$$\begin{aligned}
\mathsf{p}_{\mathsf{Z}}(z) &= \mathsf{p}_{\mathsf{X}}(z) \star \mathsf{p}_{\mathsf{Y}}(z) \\
&= \int_{u\in\mathbb{R}} \mathsf{p}_{\mathsf{X}}(u)\mathsf{p}_{\mathsf{Y}}(z-u)\,\mathrm{d}u && \text{by definition of } \textit{convolution} \quad \text{(Definition O.3 page 336)} \\
&= \int_0^z \frac{1}{\Gamma(a)}\lambda e^{-\lambda u}(\lambda u)^{a-1}\frac{1}{\Gamma(b)}\lambda e^{-\lambda(z-u)}(\lambda(z-u))^{b-1}\,\mathrm{d}u \\
&= \frac{1}{\Gamma(a)}\frac{1}{\Gamma(b)}e^{-\lambda z}\lambda^{1+1+a-1+b-1}\int_0^z u^{a-1}(z-u)^{b-1}\,\mathrm{d}u \\
&= \frac{1}{\Gamma(a)}\frac{1}{\Gamma(b)}\lambda e^{-\lambda z}\lambda^{a+b-1}\int_0^1 (vz)^{a-1}(z-vz)^{b-1}z\,\mathrm{d}v \\
&= \frac{1}{\Gamma(a)}\frac{1}{\Gamma(b)}\lambda e^{-\lambda z}\lambda^{a+b-1}z^{a-1+b-1+1}\int_0^1 v^{a-1}(1-v)^{b-1}\,\mathrm{d}v \\
&= \left[\frac{1}{\Gamma(a)}\frac{1}{\Gamma(b)}\int_0^1 v^{a-1}(1-v)^{b-1}\,\mathrm{d}v\right]\lambda e^{-\lambda z}(\lambda z)^{a+b-1} \\
&= C\lambda e^{-\lambda z}(\lambda z)^{a+b-1} && \text{where } C \text{ is some constant} \\
&= \frac{\lambda}{\Gamma(a+b)}e^{-\lambda z}(\lambda z)^{a+b-1} && C \text{ must be the value that makes } \int_z \mathsf{p}_{\mathsf{Z}}(z) = 1
\end{aligned}$$

$\Longrightarrow \mathsf{p}_{\mathsf{Z}}(z)$ is a $(a+b, \lambda)$ Gamma distribution

C.2.4 Chi-squared distributions

Definition C.5. [6] *Let* $\mathsf{p}(x)$ *be a* PROBABILITY DENSITY FUNCTION *on a* PROBABILITY SPACE $(\Omega, \mathbb{E}, \mathsf{P})$.

DEF
$\mathsf{p}(x)$ *is a **chi-square distribution** if*
$$\mathsf{p}(x) \triangleq \left\{\begin{array}{ll} 0 & \text{if } x < 0 \\ \dfrac{1}{\sqrt{2\pi\sigma^2 x}}\exp\left[-\dfrac{x}{2\sigma^2}\right] & \text{if } x \geq 0 \end{array}\right\} \qquad \text{for } \sigma > 0$$

Theorem C.2. [7]

[5] Ross (1998), page 266
[6] Proakis (2001), page 41, Papoulis (1990) page 219 ⟨7-4 Special Distributions of Statistics, (7-78)⟩
[7] Ross (1998), page 267

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

THM *The following distributions are equivalent:*

(1). *chi-squared distribution* *and*

(2). *distribution of* X^2 *where* $\mathsf{X} \sim \mathsf{N}\left(0, \sigma^2\right)$ *and*

(3). *Gamma distribution* $\left(\frac{1}{2}, \frac{1}{2\sigma^2}\right)$

✎PROOF:

1. Proof that X^2 has chi-squared distribution:

$$\begin{aligned}
\mathsf{p}_{\mathsf{Y}}(y) &= \frac{1}{2\sqrt{y}}\left[\mathsf{p}_{\mathsf{X}}(-\sqrt{y}) + \mathsf{p}_{\mathsf{X}}(\sqrt{y})\right] && \text{by Corollary 4.3 page 29} \\
&= \frac{1}{2\sqrt{y}}\left[\frac{1}{\sqrt{2\pi\sigma^2}}\exp-\frac{(-\sqrt{y}-0)^2}{2\sigma^2} + \frac{1}{\sqrt{2\pi\sigma^2}}\exp-\frac{(+\sqrt{y}-0)^2}{2\sigma^2}\right] \\
&= \frac{1}{2\sqrt{y}}\left[2\frac{1}{\sqrt{2\pi\sigma^2}}\exp-\frac{y}{2\sigma^2}+\right] \\
&= \frac{1}{\sqrt{2\pi\sigma^2 y}}\exp-\frac{y}{2\sigma^2}
\end{aligned}$$

2. Proof that chi-distribution is a Gamma distribution (b, λ):

$$\begin{aligned}
b &\triangleq \frac{1}{2} \\
\lambda &\triangleq \frac{1}{2\sigma^2}
\end{aligned}$$

$$\begin{aligned}
\frac{1}{\sqrt{2\pi\sigma^2 y}}\exp-\frac{y}{2\sigma^2} &= \frac{1}{\sqrt{\pi}}\lambda^{1/2}\lambda^{1/2}(\lambda y)^{-1/2}e^{-\lambda y} \\
&= \frac{\lambda}{\sqrt{\pi}}(\lambda y)^{b-1}e^{-\lambda y}
\end{aligned}$$

✏

Definition C.6. [8] *The* ***Chi-squared distribution with*** n ***degrees of freedom*** *has pdf*

DEF
$$\mathsf{p}_{\mathsf{Y}}(y) \triangleq \begin{cases} 0 & : y < 0 \\ \dfrac{1}{2\sigma^2\Gamma(n/2)}\left(\dfrac{y}{2\sigma^2}\right)^{\frac{n}{2}-1}\exp-\dfrac{y}{2\sigma^2} & : y \geq 0 \end{cases}$$

Theorem C.3. [9] *The following distributions are equivalent:*

1. *chi-squared distribution with* n *degrees of freedom*
2. *the distribution of* $\sum_{k=1}^{n} X_k^2$ *where* $\left\{X_k | X_k \sim \mathsf{N}\left(0, \sigma^2\right), k = 1, 2, \ldots, n\right\}$ *are independent random variables.*
3. *Gamma distribution* $\left(\frac{n}{2}, \frac{1}{2\sigma^2}\right)$.

[8] Proakis (2001), page 41
[9] Ross (1998), page 267

COPYRIGHT © 2019 DANIEL J. GREENHOE

✎PROOF:

1. Prove chi-squared distribution with n degrees of freedom is the Gamma distribution $\left(\frac{n}{2}, \frac{1}{2\sigma^2}\right)$:

$$\begin{aligned}
\lambda &\triangleq \frac{1}{2\sigma^2} \\
b &\triangleq \frac{1}{2} \\
\frac{1}{2\sigma^2\Gamma(n/2)}\left(\frac{y}{2\sigma^2}\right)^{\frac{n}{2}-1}\exp-\frac{y}{2\sigma^2} &= \frac{\lambda}{\Gamma(nb)}(\lambda y)^{nb-1}\exp-\lambda y
\end{aligned}$$

2. Prove $\sum_{k=1}^{n} X^2$ is Gamma $\left(\frac{n}{2}, \frac{1}{2\sigma^2}\right)$:

   (a) By Theorem C.2, X_k has Gamma distribution $\left(\frac{1}{2}, \frac{1}{2\sigma^2}\right)$.

   (b) By Theorem C.1, $\sum_{k=1}^{n} X_k^2$ has distribution $\left(\frac{n}{2}, \frac{1}{2\sigma^2}\right)$.

✏

Definition C.7. [10] *A **noncentral chi-square distribution** (μ, σ^2) has pdf*

DEF
$$\mathsf{p}_{\mathsf{Y}}(y) = \frac{1}{\sqrt{2\pi\sigma^2 y}}\exp\frac{y+\mu^2}{-2\sigma^2}\cosh\frac{\mu\sqrt{y}}{\sigma^2}$$

Theorem C.4.

THM
The following distributions are equivalent:

(1). NON-CENTRAL CHI-SQUARED DISTRIBUTION (μ, σ^2)

(2). *distribution of* X^2 *where* $\mathsf{X} \sim \mathsf{N}(\mu, \sigma^2)$

✎PROOF:

1. Proof that $\mathsf{Y} = X^2$ has a non-central chi-squared distribution:

$$\begin{aligned}
\mathsf{p}_{\mathsf{Y}}(y) &= \frac{1}{2\sqrt{y}}\left[\mathsf{p}_{\mathsf{X}}(-\sqrt{y}) + \mathsf{p}_{\mathsf{X}}(\sqrt{y})\right] \qquad \text{by Corollary 4.3 page 29} \\
&= \frac{1}{2\sqrt{y}}\left[\frac{1}{\sqrt{2\pi\sigma^2}}\exp-\frac{(-\sqrt{y}-\mu)^2}{2\sigma^2} + \frac{1}{\sqrt{2\pi\sigma^2}}\exp-\frac{(+\sqrt{y}-\mu)^2}{2\sigma^2}\right] \\
&= \frac{1}{2\sqrt{y}}\left[\frac{1}{\sqrt{2\pi\sigma^2}}\exp-\frac{y+\mu^2}{2\sigma^2}\exp\frac{-2\mu\sqrt{y}}{2\sigma^2} + \frac{1}{\sqrt{2\pi\sigma^2}}\exp-\frac{y+\mu^2}{2\sigma^2}\exp\frac{2\mu\sqrt{y}}{2\sigma^2}\right] \\
&= \frac{1}{\sqrt{2\pi\sigma^2 y}}\exp-\frac{y+\mu^2}{2\sigma^2}\frac{1}{2}\left[\exp\frac{2\mu\sqrt{y}}{2\sigma^2} + \exp\frac{-2\mu\sqrt{y}}{2\sigma^2}\right] \\
&= \frac{1}{\sqrt{2\pi\sigma^2 y}}\exp\frac{y+\mu^2}{-2\sigma^2}\cosh\frac{\mu\sqrt{y}}{\sigma^2}
\end{aligned}$$

✏

[10] Proakis (2001), page 42

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE



Figure C.3: *Joint Gaussian distributions* $\mathsf{p}_{\mathsf{XY}}(x, y)$ with varying correlations

Definition C.8. [11] *The* α***th-order modified Bessel function of the first kind*** $I_\alpha(x)$ *is*

DEF
$$I_\alpha(x) = \sum_{k=0}^{\infty} \frac{1}{k!\Gamma(\alpha+k+1)} \left(\frac{x}{2}\right)^{\alpha+2k}$$

Definition C.9. [12] *The* ***noncentral chi-square with*** n***-degrees of freedom*** *distribution has pdf*

DEF
$$\mathsf{p}_{\mathsf{Y}}(y) = \frac{1}{2\sigma^2}\left(\frac{y}{s^2}\right)^{\frac{n-2}{4}} \exp\frac{y+s^2}{-2\sigma^2} I_{n/2-1}\left(\sqrt{y}\frac{s}{\sigma^2}\right) \qquad \text{where } s^2 \triangleq \sum_{k=1}^{n} \mu_k^2$$

C.2.5 Radial distributions

Definition C.10. [13] *The* ***Rayleigh distribution*** *is the pdf*

DEF
$$\mathsf{p}_{\mathsf{R}}(r) = \begin{cases} 0 & \text{for} \quad r < 0 \\ \frac{r}{\sigma^2}\exp -\frac{r^2}{2\sigma^2} & \text{for} \quad r \geq 0 \end{cases}$$

Note that by Proposition 4.3, this distribution is equivalent to the distribution of $R = \sqrt{X^2+Y^2}$ where X and Y are independent random variables each with distribution $\mathsf{N}\left(0, \sigma^2\right)$.

Definition C.11. [14] *The* ***Rice distribution*** *is the pdf*

DEF
$$\mathsf{p}_{\mathsf{R}}(r) = \begin{cases} 0 & \text{for} \quad r < 0 \\ \frac{r}{\sigma^2}\exp\frac{r^2+s^2}{-2\sigma^2} I_o\left(\frac{rs}{\sigma^2}\right) & \text{for} \quad r \geq 0 \end{cases}$$

Copyright © 2019 Daniel J. Greenhoe

C.3 Joint Gaussian distributions

Definition C.12 (Joint Gaussian pdf). [15]

DEF

$$\mathsf{p}(x_1, x_2, \ldots, x_n) \triangleq \frac{1}{\sqrt{(2\pi)^n |\boldsymbol{M}|}} \exp -\frac{1}{2}(\boldsymbol{x} - \mathsf{E}\boldsymbol{x})^T \boldsymbol{M}^{-1}(\boldsymbol{x} - \mathsf{E}\boldsymbol{x}) \qquad \textit{(Gaussian joint pdf)}$$

$$\boldsymbol{x} \triangleq \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}$$

$$\mathsf{Z}_k \triangleq X_k - \mathsf{E}X_k \qquad \textit{(zero mean random variables)}$$

$$\boldsymbol{M} \triangleq \begin{bmatrix} \mathsf{E}[\mathsf{Z}_1\mathsf{Z}_1] & \mathsf{E}[\mathsf{Z}_1\mathsf{Z}_2] & \cdots & \mathsf{E}[\mathsf{Z}_1\mathsf{Z}_n] \\ \mathsf{E}[\mathsf{Z}_2\mathsf{Z}_1] & \mathsf{E}[\mathsf{Z}_2\mathsf{Z}_2] & & \mathsf{E}[\mathsf{Z}_2\mathsf{Z}_n] \\ \vdots & \vdots & \ddots & \vdots \\ \mathsf{E}[\mathsf{Z}_n\mathsf{Z}_1] & \mathsf{E}[\mathsf{Z}_n\mathsf{Z}_2] & \cdots & \mathsf{E}[\mathsf{Z}_n\mathsf{Z}_n] \end{bmatrix} \qquad \textit{(correlation matrix)}$$



Figure C.4: Gaussian pdf with $\mu = 0$ and $\sigma \in [0.1, 2]$.

Example C.2 (1 variable joint Gaussian pdf). The **Gaussian distribution** (or **normal distribution** has pdf

EX

$$\mathsf{p}_{\mathsf{x}}(x) \triangleq \frac{1}{\sqrt{2\pi\sigma^2}} e^{\frac{-(x-\mu)^2}{2\sigma^2}}$$

[11] Proakis (2001), page 43
[12] Proakis (2001), page 43
[13] Proakis (2001), page 44
[14] Proakis (2001), page 46
[15] Anderson (1984) page 21 ⟨Theorem 2.3.1⟩, Anderson (1958), page 14 ⟨§"2.3 The Multivariate Normal Distribution"⟩, Proakis (2001), page 49, Moon and Stirling (2000), page 34

BY-NC-ND Copyright © 2019 Daniel J. Greenhoe

$$
\begin{aligned}
t &= \arg_t \min_t \left[\frac{1}{2}\int_t^{\infty} \frac{1}{\sqrt{2\pi\sigma^2}} e^{\frac{-(x-\mu)^2}{2\sigma^2}} + \frac{1}{2}\int_{-\infty}^{t} \frac{1}{\sqrt{2\pi\sigma^2}} e^{\frac{-(x-\eta)^2}{2\sigma^2}}\right] \\
&= \arg_t \left\{\frac{\partial}{\partial t}\left[\frac{1}{2}\int_t^{\infty} \frac{1}{\sqrt{2\pi\sigma^2}} e^{\frac{-(x-\mu)^2}{2\sigma^2}} + \frac{1}{2}\int_{-\infty}^{t} \frac{1}{\sqrt{2\pi\sigma^2}} e^{\frac{-(x-\eta)^2}{2\sigma^2}}\right] = 0\right\} \\
&= \arg_t \left\{\frac{1}{2\sqrt{2\pi\sigma^2}}\left[\frac{\partial}{\partial t}\int_t^{\infty} e^{\frac{-(x-\mu)^2}{2\sigma^2}} + \frac{\partial}{\partial t}\int_{-\infty}^{t} e^{\frac{-(x-\eta)^2}{2\sigma^2}}\right] = 0\right\} \\
&= \arg_t \left\{\left[\left(e^{\frac{-(\infty-\mu)^2}{2\sigma^2}} 0 - e^{\frac{-(t-\mu)^2}{2\sigma^2}} 1\right) + \left(e^{\frac{-(t-\mu)^2}{2\sigma^2}} 1 - e^{\frac{-(\infty-\mu)^2}{2\sigma^2}} 0\right)\right] = 0\right\} \\
&= \arg_t \left\{\left[e^{\frac{-(t-\eta)^2}{2\sigma^2}} - e^{\frac{-(t-\mu)^2}{2\sigma^2}}\right] = 0\right\} \\
&= \arg_t \left\{(t-\eta)^2 = (t-\mu)^2\right\} \\
&= \frac{\mu+\eta}{2}
\end{aligned}
$$

Example C.3 (2 variable joint Gaussian pdf).

EX

$$
\begin{aligned}
z_1 &\triangleq x_1 - \mathsf{E}x_1 \\
z_2 &\triangleq x_2 - \mathsf{E}x_2 \\
|M| &\triangleq \left|\mathsf{E}[z_1 z_1]\mathsf{E}[z_2 z_2] - \mathsf{E}[z_1 z_2]\mathsf{E}[z_1 z_2]\right| \\
\mathsf{p}(x_1, x_2) &\triangleq \frac{1}{2\pi\sqrt{|M|}} \exp\left(\frac{z_1^2\mathsf{E}[z_2 z_2] - 2z_1 z_2 \mathsf{E}[z_1 z_2] + z_2^2 \mathsf{E}[z_1 z_1]}{-2|M|}\right)
\end{aligned}
$$

Copyright © 2019 Daniel J. Greenhoe

Copyright © 2019 Daniel J. Greenhoe

APPENDIX D

SPECTRAL THEORY

D.1 Operator Spectrum

Definition D.1. [1] *Let* $\mathbf{A} \in \mathcal{B}(\boldsymbol{X}, \boldsymbol{Y})$ *be an operator over the linear spaces* $\boldsymbol{X} = (X, F, \oplus, \otimes)$ *and* $\boldsymbol{Y} \triangleq (Y, F, \oplus, \otimes)$. *Let* $\mathcal{N}(\mathbf{A})$ *be the* NULL SPACE *of* $\mathbf{A}$.

DEF

An ***eigenvalue*** *of* $\mathbf{A}$ *is any value* λ *such that there exists* $\boldsymbol{x}$ *such that* $\mathbf{A}\boldsymbol{x} = \lambda\boldsymbol{x}$.
The ***eigenspace*** $\boldsymbol{H}_\lambda$ *of* $\mathbf{A}$ *at eigenvalue* λ *is* $\mathcal{N}(\mathbf{A} - \lambda\mathbf{I})$.
An ***eigenvector*** *of* $\mathbf{A}$ *associated with eigenvalue* λ *is any element of* $\mathcal{N}(\mathbf{A} - \lambda\mathbf{I})$.

Example D.1. [2] Let $\mathbf{D}$ be the differntial operator.

EX

The set $\left\{e^{\lambda x} | \lambda \in \mathbb{C}\right\}$ are the eigenvectors of $\mathbf{D}$.

$$\begin{aligned} \rho(\mathbf{D}) &= \varnothing && (\mathbf{D} \text{ has no non-spectral points whatsoever}) \\ \sigma_{\mathsf{p}}(\mathbf{D}) &= \sigma(\mathbf{D}) && (\text{the spectrum of } \mathbf{D} \text{ is all eigenvalues}) \\ \sigma_{\mathsf{c}}(\mathbf{D}) &= \varnothing && (\mathbf{D} \text{ has no continuous spectrum}) \\ \sigma_{\mathsf{r}}(\mathbf{D}) &= \varnothing && (\mathbf{D} \text{ has no resolvent spectrum}) \end{aligned}$$

✎PROOF:

$$\begin{aligned} (\mathbf{D} - \lambda\mathbf{I})e^{\lambda x} &= \mathbf{D}e^{\lambda x} - \lambda\mathbf{I}e^{\lambda x} \\ &= \lambda e^{\lambda x} - \lambda e^{\lambda x} \\ &= 0 && \forall \lambda \in \mathbb{C} \end{aligned}$$

This theorem and proof needs more work and investigation to prove/disprove its claims. ✏

Definition D.2. [3] *Let* $\mathbf{A} \in \mathcal{B}(\boldsymbol{X}, \boldsymbol{Y})$ *be an operator over the linear spaces* $\boldsymbol{X} = (X, F, \oplus, \otimes)$ *and* $\boldsymbol{Y} \triangleq (Y, F, \oplus, \otimes)$.

[1] Bollobás (1999), page 168, Descartes (1637), Descartes (1954), Cayley (1858), Hilbert (1904), page 67, Hilbert (1912),
[2] Pedersen (2000), page 79
[3] Michel and Herget (1993), page 439

quantity		$\mathcal{N}(\mathbf{A}-\lambda\mathbf{I}) = \{\mathbb{0}\}$ ($\boldsymbol{x} = \mathbb{0}$ is the only solution)	$\overline{\mathcal{R}(\mathbf{A}-\lambda\mathbf{I})} = \boldsymbol{X}$ (dense)	$(\mathbf{A}-\lambda\mathbf{I})^{-1} \in \mathcal{B}(\boldsymbol{X},\boldsymbol{Y})$ (continuous/bounded)
$\rho(\mathbf{A})$	(resolvent set)	1	1	1
$\sigma_{\mathsf{p}}(\mathbf{A})$	(point spectrum)	0		
$\sigma_{\mathsf{r}}(\mathbf{A})$	(residual spectrum)	1	0	
$\sigma_{\mathsf{c}}(\mathbf{A})$	(continuous spectrum)	1	1	0

Table D.1: Spectrum of an operator $\mathbf{A}$

DEF

*The **resolvent set** $\rho(\mathbf{A})$ of operator $\mathbf{A}$ is defined as*

$$\rho(\mathbf{A}) \triangleq \left\{ \lambda \in F \,\middle|\, \begin{array}{lll} 1. & \mathcal{N}(\mathbf{A}-\lambda\mathbf{I}) = \{\mathbb{0}\} & \textit{(no non-zero eigenvectors)} \quad \textit{and} \\ 2. & \overline{\mathcal{R}(\mathbf{A}-\lambda\mathbf{I})} = \boldsymbol{X} & \textit{(the range is dense in } \boldsymbol{X}\textit{).} \quad \textit{and} \\ 3. & (\mathbf{A}-\lambda\mathbf{I})^{-1} \in \mathcal{B}(\boldsymbol{X},\boldsymbol{Y}) & \textit{(inverse is continuous/bounded).} \end{array} \right\}$$

*The **spectrum** $\sigma(\mathbf{A})$ of operator $\mathbf{A}$ is defined as*

$$\sigma(\mathbf{A}) \triangleq F \setminus \rho(\mathbf{A}).$$

Definition D.3. [4] *Let $\mathbf{A} \in \mathcal{B}(\boldsymbol{X},\boldsymbol{Y})$ be an operator over the linear spaces $\boldsymbol{X} = (X, F, \oplus, \otimes)$ and $\boldsymbol{Y} \triangleq (Y, F, \oplus, \otimes)$.*

DEF

*The **point spectrum** $\sigma_{\mathsf{p}}(\mathbf{A})$ of operator $\mathbf{A}$ is defined as*

$$\sigma_{\mathsf{p}}(\mathbf{A}) \triangleq \left\{ \lambda \in F \,\middle|\, 1. \quad \mathcal{N}(\mathbf{A}-\lambda\mathbf{I}) \supseteq \{\mathbb{0}\} \quad \textit{(has non-zero eigenvector)} \right\}$$

*The **residual spectrum** $\sigma_{\mathsf{r}}(\mathbf{A})$ of operator $\mathbf{A}$ is defined as*

$$\sigma_{\mathsf{r}}(\mathbf{A}) \triangleq \left\{ \lambda \in F \,\middle|\, \begin{array}{lll} 1. & \mathcal{N}(\mathbf{A}-\lambda\mathbf{I}) = \{\mathbb{0}\} & \textit{(no non-zero eigenvectors)} \quad \textit{and} \\ 2. & \overline{\mathcal{R}(\mathbf{A}-\lambda\mathbf{I})} \neq \boldsymbol{X} & \textit{(not dense in } \boldsymbol{X}\textit{—has gaps).} \end{array} \right\}$$

*The **continuous spectrum** $\sigma_{\mathsf{c}}(\mathbf{A})$ of operator $\mathbf{A}$ is defined as*

$$\sigma_{\mathsf{c}}(\mathbf{A}) \triangleq \left\{ \lambda \in F \,\middle|\, \begin{array}{lll} 1. & \mathcal{N}(\mathbf{A}-\lambda\mathbf{I}) = \{\mathbb{0}\} & \textit{(no non-zero eigenvectors)} \quad \textit{and} \\ 2. & \overline{\mathcal{R}(\mathbf{A}-\lambda\mathbf{I})} = \boldsymbol{X} & \textit{(dense in } \boldsymbol{X}\textit{).} \quad \textit{and} \\ 3. & (\mathbf{A}-\lambda\mathbf{I})^{-1} \notin \mathcal{B}(\boldsymbol{X},\boldsymbol{Y}) & \textit{(not continuous / not bounded)} \end{array} \right\}$$



The spectral components' definitions are illustrated in the figure to the left and summarized in Table D.1 (page 190). Let a family of operators $\mathbf{B}(\lambda)$ be defined with respect to an operator $\mathbf{A}$ such that $\mathbf{B}(\lambda) \triangleq (\mathbf{A} - \lambda\mathbf{I})$. Normally, we might expect a "normal" or "regular" or even "mundane" operator $\mathbf{B}(\lambda)$ to have the properties

1. $\mathbf{B}(\lambda)\boldsymbol{x} = 0$ if and only if $\boldsymbol{x} = \mathbb{0}$
2. $\mathbf{B}(\lambda)\boldsymbol{x}$ spans virtually all of $\boldsymbol{X}$ as we vary $\boldsymbol{x}$
3. $\mathbf{B}^{-1}(\lambda)$ is continuous.

After all, these are the properties that we would have if $\mathbf{B}(\lambda)$ were simply an affine operator in the

[4] Bollobás (1999), page 168, Hilbert (1906) pages 169–172

BY-NC-ND Copyright © 2019 Daniel J. Greenhoe

field of real numbers— such as $[\mathbf{B}(\lambda)](x) \triangleq [\lambda](x) = \lambda x$ which is 0 if and only if $x = 0$, has range $\boldsymbol{\mathcal{R}}(\lambda) = \mathbb{R}$, and its inverse $\lambda^{-1}x$ is continuous.

If for some λ the operator $\mathbf{B}(\lambda)$ does have all these "regular" properties, then that λ part of the *resolvent set* of $\mathbf{A}$ and λ is called *regular*. However if for some λ the operator $\mathbf{B}(\lambda)$ fails any of these conditions, then that λ part of the *spectrum* of $\mathbf{A}$. And which conditions it fails determines which component of the spectrum it is in.

Theorem D.1. [5] *Let* $\mathbf{A} \in \mathcal{B}(\boldsymbol{X}, \boldsymbol{Y})$ *be an operator.*

THM

$$\sigma(\mathbf{A}) = \sigma_{\mathsf{p}}(\mathbf{A}) \cup \sigma_{\mathsf{c}}(\mathbf{A}) \cup \sigma_{\mathsf{r}}(\mathbf{A})$$

Theorem D.2 (Spectral Theorem)**.** [6] *Let* $\mathbf{N} \in Y^X$ *be an operator.*

THM

$$\left.\begin{array}{ll} (A). & \underbrace{\mathbf{N}^*\mathbf{N} = \mathbf{N}\mathbf{N}^*}_{\mathbf{N}\ \textit{is}\ \text{NORMAL}} \\ (B). & \mathbf{N}\ \textit{is}\ \text{COMPACT} \end{array}\right\} \quad \Longrightarrow \quad \left\{\begin{array}{lrcl} (1). & \mathbf{N} & = & \sum_n \lambda_n \mathbf{P}_n \\ (2). & \sum_n \mathbf{P}_n & = & \mathbf{I} \\ (3). & \mathbf{P}_n\mathbf{P}_m & = & \bar{\delta}_{n-m}\mathbf{P}_n \\ (4). & dim(\boldsymbol{H}_n) & < & \infty \\ (5). & \left|\left\{\lambda_n | \lambda_n \neq 0\right\}\right| & \textit{is} & \text{COUNTABLY INFINITE} \end{array}\right.$$

where

$$\begin{array}{rcll} (\!(\lambda_n)\!)_{n\in\mathbb{Z}} & \triangleq & \sigma_{\mathsf{p}}(\mathbf{N}) & \textit{(eigenvalues of}\ \mathbf{N}\textit{)} \\ \boldsymbol{H}_n & \triangleq & \mathcal{N}(\mathbf{N} - \lambda_n\mathbf{I}) & \textit{(}\lambda_n\ \textit{is the eigenspace of}\ \mathbf{N}\ \textit{at}\ \lambda_n\ \textit{in}\ \boldsymbol{Y}\textit{)} \\ \boldsymbol{H}_n & = & \mathbf{P}_n\boldsymbol{Y} & \textit{(}\mathbf{P}_n\ \textit{is the projection operator that generates}\ \boldsymbol{H}_n\textit{)} \end{array}$$

D.2 Fredholm kernels

Definition D.4. [7]

DEF

A ***Fredholm operator*** $\mathbf{K}$ *is defined as*

$$[\mathbf{K}\mathsf{f}](t) \triangleq \underbrace{\int_a^b \underbrace{\kappa(t,s)}_{\textit{kernel}}\mathsf{f}(s)\,\mathrm{d}s}_{\textit{Fredholm integral equation of the first kind}^{8}} \qquad \forall \mathsf{f} \in \boldsymbol{L}_2([a,b])$$

Example D.2. Examples of Fredholm operators include

1.	Fourier Transform	$[\tilde{\mathbf{F}}\mathsf{x}](f)$	$=$	$\int_t \mathsf{x}(t)e^{-i2\pi ft}\,\mathrm{d}t$	$\kappa(t,f)$	$=$	$e^{-i2\pi ft}$
2.	Inverse Fourier Transform	$[\tilde{\mathbf{F}}^{-1}\tilde{\mathsf{x}}](t)$	$=$	$\int_f \tilde{\mathsf{x}}(f)e^{i2\pi ft}\,\mathrm{d}f$	$\kappa(f,t)$	$=$	$e^{i2\pi ft}$
3.	Laplace operator	$[\mathbf{L}\mathsf{x}](s)$	$=$	$\int_t \mathsf{x}(t)e^{-st}\,\mathrm{d}t$	$\kappa(t,s)$	$=$	e^{-st}
4.	autocorrelation operator	$[\mathbf{R}\mathsf{x}](t)$	$=$	$\int_s R(t,s)\mathsf{x}(s)\,\mathrm{d}s$	$\kappa(t,s)$	$=$	$R(t,s)$

Theorem D.3. *Let* $\mathbf{K}$ *be a Fredholm operator with kernel* $\kappa(t,s)$ *and adjoint* $\mathbf{K}^*$.

THM

$$[\mathbf{K}\mathsf{f}](t) = \int_A \kappa(t,s)\mathsf{f}(s)\,\mathrm{d}s \quad \Longleftrightarrow \quad \left[\mathbf{K}^*\mathsf{f}\right](t) = \int_A \kappa^*(s,t)\mathsf{f}(s)\,\mathrm{d}s$$

[5] Michel and Herget (1993), page 440

[6] Michel and Herget (1993), page 457, Bollobás (1999), page 200, Hilbert (1906), Hilbert (1912), von Neumann (1929), de Witt (1659)

[7] Michel and Herget (1993), page 425

[8] The equation $\int_u \kappa(t,s)\mathsf{f}(s)\,\mathrm{d}s$ is a **Fredholm integral equation of the first kind** and $\kappa(t,u)$ is the **kernel** of the equation. References: Fredholm (1900), Fredholm (1903), page 365, Michel and Herget (1993), page 97, Keener (1988), page 101

Copyright © 2019 Daniel J. Greenhoe

✎PROOF:

$$
\begin{aligned}
& [\mathbf{K}\mathsf{f}](t) = \int_A \kappa(t,s)\mathsf{f}(s)\,\mathrm{d}s \\
\iff \langle [\mathbf{K}\mathsf{f}](t) \mid \mathsf{g}(t)\rangle &= \left\langle \int_s \kappa(t,s)\mathsf{f}(s)\,\mathrm{d}s \mid \mathsf{g}(t)\right\rangle && \text{by left hypothesis} \\
&= \int_s \mathsf{f}(s)\,\langle \kappa(t,s) \mid \mathsf{g}(t)\rangle\,\mathrm{d}s && \text{by additivity property of } \langle\triangle \mid \triangledown\rangle \text{ (Definition J.1 page 255)} \\
&= \int_s \mathsf{f}(s)\,\langle \mathsf{g}(t) \mid \kappa(t,s)\rangle^*\,\mathrm{d}s && \text{by conjugate symmetry property of } \langle\triangle \mid \triangledown\rangle \text{ (Definition J.1 page 255)} \\
&= \langle \mathsf{f}(s) \mid \langle \mathsf{g}(t) \mid \kappa(t,s)\rangle\rangle && \text{by local definition of } \langle\triangle \mid \triangledown\rangle \\
&= \left\langle \mathsf{f}(s) \mid \underbrace{\int_t \kappa^*(t,s)\mathsf{g}(t)\,\mathrm{d}t}_{[\mathbf{K}^*\mathsf{g}](s)}\right\rangle && \text{by local definition of } \langle\triangle \mid \triangledown\rangle \\
\iff [\mathbf{K}^*\mathsf{g}](s) &= \int_A \kappa^*(t,s)\mathsf{g}(t)\,\mathrm{d}t && \text{by right hypothesis} \\
\iff [\mathbf{K}^*\mathsf{g}](\sigma) &= \int_A \kappa^*(\tau,\sigma)\mathsf{g}(\tau)\,\mathrm{d}\tau && \text{by change of variable: } \tau = t,\ \sigma = s \\
\iff [\mathbf{K}^*\mathsf{f}](t) &= \int_A \kappa^*(s,t)\mathsf{f}(s)\,\mathrm{d}s && \text{by change of variable: } t = \sigma,\ s = \tau,\ \mathsf{f} = \mathsf{g}
\end{aligned}
$$

✏

Corollary D.1. [9] *Let* $\mathbf{K}$ *be an Fredholm operator with kernel* $\kappa(t,s)$ *and adjoint* $\mathbf{K}^*$.

COR

$$\underbrace{\mathbf{K} = \mathbf{K}^*}_{\mathbf{K}\text{ is self-adjoint}} \iff \underbrace{\kappa(t,s) = \kappa^*(s,t)}_{\text{kernel is conjugate symmetric}}$$

✎PROOF:

$$
\begin{aligned}
\mathbf{K} = \mathbf{K}^* &\iff \int_A \kappa(t,s)\mathsf{f}(s)\,\mathrm{d}s = \int_A \kappa^*(s,t)\mathsf{f}(s)\,\mathrm{d}s && \text{by Theorem D.3 page 191} \\
&\iff \kappa(t,s) = \kappa^*(s,t)
\end{aligned}
$$

✏

Theorem D.4 (Mercer's Theorem). [10] *Let* $\mathbf{K}$ *be an Fredholm operator with kernel* $\kappa(t,s)$ *and eigensystem* $(\!(\lambda_n, \phi_n(t))\!)_{n\in\mathbb{Z}}$.

THM

$$
\left\{
\begin{array}{ll}
(A). & \underbrace{\int_a^b \int_a^b \kappa(t,s)\mathsf{f}(t)\mathsf{f}^*(s)\,\mathrm{d}t \geq 0}_{\text{positive}} \quad \textit{and} \\
(B). & \kappa(t,s) \textit{ is } \text{CONTINUOUS } \textit{on} \\
 & [a:b]\times[a:b]
\end{array}
\right\}
\implies
\left\{
\begin{array}{ll}
(1). & \kappa(t,s) = \sum_n \lambda_n \phi_n(t)\phi_n^*(s) \quad \textit{and} \\
(2). & \kappa(t,s) \text{ CONVERGES ABSOLUTELY} \\
 & \textit{and } \text{UNIFORMLY } \textit{on} \\
 & [a:b]\times[a:b]
\end{array}
\right\}
$$

[9] Michel and Herget (1993), page 430

[10] Gohberg et al. (2003), page 198, Courant and Hilbert (1930), pages 138–140, Mercer (1909), page 439

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

APPENDIX E

MATRIX CALCULUS

Optimization problems often require finding the value of some parameter which results in some measure reaching a minimum or maximum value. Often this optimal parameter value can be found by solving the single equation generated by the partial derivative of the measure with respect to the parameter. When there are several parameters, optimization often requires several simultaneous equations generated by the partial derivatives of the measure with respect to each parameter. The need for several partial derivatives and several simultaneous equations leads to a natual union of two branches of mathematics— partial differential equations and linear algebra. In general, we would like to not only be able to take the partial derivative of a scalar with respect to another scalar, but to be able to take the partial derivative of a vector with respect to another vector. This generalization is the problem addressed in this section. Other references are also available.[1]

E.1 First derivative of a vector with respect to a vector

Definition E.1.

DEF

$\boldsymbol{x}$ is a vector with the following properties:

1. $$\boldsymbol{x} \triangleq \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} \qquad \textit{(n element column vector)}$$

2. $$\frac{\partial}{\partial x_k} x_j = \bar{\delta}_{kj} \qquad \textit{((}x_1, x_2, \ldots, x_n\textit{) are mutually independent)}$$

Definition E.2 (Jacobian matrix). [2] *The **gradient of $\boldsymbol{y}$ with respect to $\boldsymbol{x}$**, as well as the **gradient of $\boldsymbol{y}^T$ with respect to $\boldsymbol{x}$**, is defined as*

[1] Graham (1981) ⟨Chapter 4⟩, Haykin (2001) ⟨Appendix B⟩, Moon and Stirling (2000) ⟨Appendix E⟩, Scharf (1991), pages 274–276, Trees (2002) ⟨Section A.7⟩, Felippa (1999)

[2] Graham (1981), page 52, Graham (2018), page 5297804868241 78§"4.2 The Derivatives of Vectors", Scharf (1991), page 274, Trees (2002), page 1398, Anderson (1984) page 13 ⟨§"2.2.5 Transformation of Variables"⟩, Anderson (1958), page 11 ⟨§"2.2.5 Transformation of Variables"⟩

DEF

$$\frac{\partial \boldsymbol{y}}{\partial \boldsymbol{x}} \triangleq \frac{\partial \boldsymbol{y}^T}{\partial \boldsymbol{x}} \triangleq \underbrace{\begin{bmatrix} \frac{\partial y_1}{\partial x_1} & \frac{\partial y_2}{\partial x_1} & \cdots & \frac{\partial y_m}{\partial x_1} \\ \frac{\partial y_1}{\partial x_2} & \frac{\partial y_2}{\partial x_2} & \cdots & \frac{\partial y_m}{\partial x_2} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial y_1}{\partial x_n} & \frac{\partial y_2}{\partial x_n} & \cdots & \frac{\partial y_m}{\partial x_n} \end{bmatrix}}_{n \times m \text{ matrix}} \qquad \forall \boldsymbol{y} \in \mathbb{C}^m$$

Remark E.1. Depending on whether $\boldsymbol{x}$ and $\boldsymbol{y}$ are scalars or vectors, $\frac{\partial \boldsymbol{y}}{\partial \boldsymbol{x}}$ takes on the following forms:[3]

	y scalar	y vector
x scalar	$\frac{\partial y}{\partial x}$	$\frac{\partial \boldsymbol{y}}{\partial x} = \begin{bmatrix} \frac{\partial y_1}{\partial x} & \frac{\partial y_2}{\partial x} & \cdots & \frac{\partial y_m}{\partial x} \end{bmatrix}$
x vector	$\frac{\partial y}{\partial \boldsymbol{x}} = \begin{bmatrix} \frac{\partial y}{\partial x_1} \\ \frac{\partial y}{\partial x_2} \\ \vdots \\ \frac{\partial y}{\partial x_n} \end{bmatrix}$	$\frac{\partial \boldsymbol{y}}{\partial \boldsymbol{x}} \triangleq \begin{bmatrix} \frac{\partial y_1}{\partial x_1} & \frac{\partial y_2}{\partial x_1} & \cdots & \frac{\partial y_m}{\partial x_1} \\ \frac{\partial y_1}{\partial x_2} & \frac{\partial y_2}{\partial x_2} & \cdots & \frac{\partial y_m}{\partial x_2} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial y_1}{\partial x_n} & \frac{\partial y_2}{\partial x_n} & \cdots & \frac{\partial y_m}{\partial x_n} \end{bmatrix}$

Lemma E.1. *Let $\boldsymbol{x} \in \mathbb{R}^n$ be a vector. Then*

LEM

$$\frac{\partial}{\partial x_k} x_i x_j = \bar{\delta}_{ik} x_j + \bar{\delta}_{jk} x_i = \begin{cases} 2x_k & \text{for } i = j = k \\ x_j & \text{for } i = k \text{ and } j \neq k \\ x_i & \text{for } i \neq k \text{ and } j = k \\ 0 & \text{otherwise} \end{cases}$$

Lemma E.2.

LEM

$$\left(\boldsymbol{x}^H \mathbf{A} \boldsymbol{x}\right) = \sum_{i=1}^{n} \sum_{j=1}^{n} a_{ij} x_i^* x_j \qquad \forall \quad \begin{matrix} \mathbf{A} \in (\mathbb{C}^n \times \mathbb{C}^n) & \text{(}n \times n\text{ array)} & \text{and} \\ \boldsymbol{x} \in \mathbb{C}^n & \text{(}n\text{ element column vector)} & \end{matrix}$$

✎PROOF:

$$\boldsymbol{x}^H \mathbf{A} \boldsymbol{x} \triangleq \begin{bmatrix} x_1 & x_2 & \cdots & x_n \end{bmatrix}^* \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & \cdots & a_{nn} \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} \qquad \text{by definitions of } \mathbf{A} \text{ and } \boldsymbol{x}$$

$$= \begin{bmatrix} x_1 & x_2 & \cdots & x_n \end{bmatrix}^* \sum_{i=1}^{n} x_i \begin{bmatrix} a_{1i} \\ a_{2i} \\ \vdots \\ a_{ni} \end{bmatrix}$$

$$= \sum_{i=1}^{n} x_i \begin{bmatrix} x_1 & x_2 & \cdots & x_n \end{bmatrix}^* \begin{bmatrix} a_{1i} \\ a_{2i} \\ \vdots \\ a_{ni} \end{bmatrix}$$

$$= \sum_{i=1}^{n} x_i \sum_{j=1}^{n} a_{ji} x_j^*$$

[3] For the generalization of the partial derivative of a matrix with respect to a matrix, see Graham (1981) ⟨chapter 6⟩. Graham uses *kronecker products* to handle the additional dimensions(?)

BY-NC-ND | COPYRIGHT © 2019 DANIEL J. GREENHOE

$$= \sum_{i=1}^{n}\sum_{j=1}^{n} a_{ij} x_i^* x_j$$

Lemma E.3.

LEM

$$\frac{\partial}{\partial \boldsymbol{x}}[\mathsf{a}(\boldsymbol{x})\,\mathsf{b}(\boldsymbol{x})] = \mathsf{a}(\boldsymbol{x})\left[\frac{\partial}{\partial \boldsymbol{x}}\mathsf{b}(\boldsymbol{x})\right] + \left[\frac{\partial}{\partial \boldsymbol{x}}\mathsf{a}(\boldsymbol{x})\right]\mathsf{b}(\boldsymbol{x}) \qquad \underbrace{\forall \mathsf{a},\mathsf{b} : \mathbb{R}^n \to \mathbb{R}}_{\mathsf{a}(\boldsymbol{x}),\,\mathsf{b}(\boldsymbol{x})\ \textit{are functions from a vector } \boldsymbol{x} \textit{ to a scalar in } \mathbb{R}}$$

✎PROOF:

$$\frac{\partial}{\partial \boldsymbol{x}}[\mathsf{a}(\boldsymbol{x})\,\mathsf{b}(\boldsymbol{x})] = \begin{bmatrix} \frac{\partial}{\partial x_1}[\mathsf{a}(\boldsymbol{x})\,\mathsf{b}(\boldsymbol{x})] \\ \frac{\partial}{\partial x_2}[\mathsf{a}(\boldsymbol{x})\,\mathsf{b}(\boldsymbol{x})] \\ \vdots \\ \frac{\partial}{\partial x_n}[\mathsf{a}(\boldsymbol{x})\,\mathsf{b}(\boldsymbol{x})] \end{bmatrix} \qquad \text{by definition of } \frac{\partial}{\partial \boldsymbol{x}} \qquad \text{(Definition E.2 page 193)}$$

$$= \begin{bmatrix} \mathsf{a}(\boldsymbol{x})\frac{\partial \mathsf{b}(\boldsymbol{x})}{\partial x_1} + \mathsf{b}(\boldsymbol{x})\frac{\partial \mathsf{a}(\boldsymbol{x})}{\partial x_1} \\ \mathsf{a}(\boldsymbol{x})\frac{\partial \mathsf{b}(\boldsymbol{x})}{\partial x_2} + \mathsf{b}(\boldsymbol{x})\frac{\partial \mathsf{a}(\boldsymbol{x})}{\partial x_2} \\ \vdots \\ \mathsf{a}(\boldsymbol{x})\frac{\partial \mathsf{b}(\boldsymbol{x})}{\partial x_n} + \mathsf{b}(\boldsymbol{x})\frac{\partial \mathsf{a}(\boldsymbol{x})}{\partial x_n} \end{bmatrix} \qquad \text{by } \textit{linearity} \text{ of } \frac{\partial}{\partial \boldsymbol{x}}$$

$$= \begin{bmatrix} \mathsf{a}(\boldsymbol{x})\frac{\partial \mathsf{b}(\boldsymbol{x})}{\partial x_1} \\ \mathsf{a}(\boldsymbol{x})\frac{\partial \mathsf{b}(\boldsymbol{x})}{\partial x_2} \\ \\ \mathsf{a}(\boldsymbol{x})\frac{\partial \mathsf{b}(\boldsymbol{x})}{\partial x_n} \end{bmatrix} + \begin{bmatrix} \frac{\partial \mathsf{a}(\boldsymbol{x})}{\partial x_1}\mathsf{b}(\boldsymbol{x}) \\ \frac{\partial \mathsf{a}(\boldsymbol{x})}{\partial x_2}\mathsf{b}(\boldsymbol{x}) \\ \vdots \\ \frac{\partial \mathsf{a}(\boldsymbol{x})}{\partial x_n}\mathsf{b}(\boldsymbol{x}) \end{bmatrix} \qquad \text{by } \textit{linearity} \text{ of } \textit{vector addition}$$

$$= \mathsf{a}(\boldsymbol{x})\left[\frac{\partial \mathsf{b}(\boldsymbol{x})}{\partial \boldsymbol{x}}\right] + \left[\frac{\partial \mathsf{a}(\boldsymbol{x})}{\partial \boldsymbol{x}}\right]\mathsf{b}(\boldsymbol{x})$$

Theorem E.1. [4]

LEM

$$\frac{\partial}{\partial \boldsymbol{x}}\boldsymbol{x} = \mathbf{I} \qquad \forall \boldsymbol{x} \in \mathbb{R}^n$$

✎PROOF:

$$\frac{\partial}{\partial \boldsymbol{x}}\boldsymbol{x} = \begin{bmatrix} \frac{\partial x_1}{\partial x_1} & \frac{\partial x_2}{\partial x_1} & \cdots & \frac{\partial x_n}{\partial x_1} \\ \frac{\partial x_1}{\partial x_2} & \frac{\partial x_2}{\partial x_2} & \cdots & \frac{\partial x_n}{\partial x_2} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial x_1}{\partial x_n} & \frac{\partial x_2}{\partial x_n} & \cdots & \frac{\partial x_n}{\partial x_n} \end{bmatrix} \qquad \text{by Definition E.2 page 193}$$

$$= \begin{bmatrix} \bar{\delta}_{11} & \bar{\delta}_{21} & \cdots & \bar{\delta}_{n1} \\ \bar{\delta}_{12} & \bar{\delta}_{22} & \cdots & \bar{\delta}_{n2} \\ \vdots & \vdots & \ddots & \vdots \\ \bar{\delta}_{1n} & \bar{\delta}_{2n} & \cdots & \bar{\delta}_{nn} \end{bmatrix} \qquad \text{by Definition E.1 page 193 (mutual independence property)}$$

[4] Scharf (1991), page 274, Trees (2002), page 1398

COPYRIGHT © 2019 DANIEL J. GREENHOE

$$= \begin{bmatrix} 1 & 0 & \cdots & 0 \\ 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 1 \end{bmatrix} \qquad \text{by definition of kronecker delta function } \bar{\delta}$$

$$= \mathbf{I} \qquad \text{by definition of identity operator } \mathbf{I}$$

✏

Theorem E.2.

THM $$\frac{\partial}{\partial \boldsymbol{x}}(\mathbf{A}\boldsymbol{x}) = \mathbf{A}^T + \sum_{i=1}^{n} \left(\frac{\partial}{\partial \boldsymbol{x}} \begin{bmatrix} a_{1i} & a_{2i} & \cdots & a_{mi} \end{bmatrix} \right) x_i \qquad \forall \boldsymbol{x} \in \mathbb{R}^n,\ \mathbf{A} \in \mathbb{C}^m \times \mathbb{C}^n$$

✎PROOF: Let

$$\boldsymbol{x} \triangleq \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} \qquad \mathbf{A} \triangleq \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}$$

$$\frac{\partial}{\partial \boldsymbol{x}}(\mathbf{A}\boldsymbol{x}) = \frac{\partial}{\partial \boldsymbol{x}} \left(\begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} \right) \qquad \text{by definition of } A \text{ and } \boldsymbol{x}$$

$$= \frac{\partial}{\partial \boldsymbol{x}} \begin{bmatrix} \sum_{i=1}^{n} a_{1i}x_i \\ \sum_{i=1}^{n} a_{2i}x_i \\ \vdots \\ \sum_{i=1}^{n} a_{mi}x_i \end{bmatrix} \qquad \text{by matrix multiplication}$$

$$= \frac{\partial}{\partial \boldsymbol{x}} \sum_{i=1}^{n} \begin{bmatrix} a_{1i}x_i \\ a_{2i}x_i \\ \vdots \\ a_{mi}x_i \end{bmatrix}$$

$$= \sum_{i=1}^{n} \frac{\partial}{\partial \boldsymbol{x}} \begin{bmatrix} a_{1i}x_i \\ a_{2i}x_i \\ \vdots \\ a_{mi}x_i \end{bmatrix}$$

$$= \sum_{i=1}^{n} \begin{bmatrix} \frac{\partial a_{1i}x_i}{\partial x_1} & \frac{\partial a_{2i}x_i}{\partial x_1} & \cdots & \frac{\partial a_{mi}x_i}{\partial x_1} \\ \frac{\partial a_{1i}x_i}{\partial x_2} & \frac{\partial a_{2i}x_i}{\partial x_2} & \cdots & \frac{\partial a_{mi}x_i}{\partial x_2} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial a_{1i}x_i}{\partial x_n} & \frac{\partial a_{2i}x_i}{\partial x_n} & \cdots & \frac{\partial a_{mi}x_i}{\partial x_n} \end{bmatrix} \qquad \text{by Definition E.2 page 193}$$

$$= \sum_{i=1}^{n} \begin{bmatrix} a_{1i}\frac{\partial x_i}{\partial x_1} + \frac{\partial a_{1i}}{\partial x_1}x_i & a_{2i}\frac{\partial x_i}{\partial x_1} + \frac{\partial a_{2i}}{\partial x_1}x_i & \cdots & a_{mi}\frac{\partial x_i}{\partial x_1} + \frac{\partial a_{mi}}{\partial x_1}x_i \\ a_{1i}\frac{\partial x_i}{\partial x_2} + \frac{\partial a_{1i}}{\partial x_2}x_i & a_{2i}\frac{\partial x_i}{\partial x_2} + \frac{\partial a_{2i}}{\partial x_2}x_i & \cdots & a_{mi}\frac{\partial x_i}{\partial x_2} + \frac{\partial a_{mi}}{\partial x_2}x_i \\ \vdots & \vdots & \ddots & \vdots \\ a_{1i}\frac{\partial x_i}{\partial x_n} + \frac{\partial a_{1i}}{\partial x_n}x_i & a_{2i}\frac{\partial x_i}{\partial x_n} + \frac{\partial a_{2i}}{\partial x_n}x_i & \cdots & a_{mi}\frac{\partial x_i}{\partial x_n} + \frac{\partial a_{mi}}{\partial x_n}x_i \end{bmatrix} \qquad \text{by Lemma E.3 page 195}$$

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

$$= \sum_{i=1}^{n} \begin{bmatrix} a_{1i}\frac{\partial x_i}{\partial x_1} & a_{2i}\frac{\partial x_i}{\partial x_1} & \cdots & a_{mi}\frac{\partial x_i}{\partial x_1} \\ a_{1i}\frac{\partial x_i}{\partial x_2} & a_{2i}\frac{\partial x_i}{\partial x_2} & \cdots & a_{mi}\frac{\partial x_i}{\partial x_2} \\ \vdots & \vdots & \ddots & \vdots \\ a_{1i}\frac{\partial x_i}{\partial x_n} & a_{2i}\frac{\partial x_i}{\partial x_n} & \cdots & a_{mi}\frac{\partial x_i}{\partial x_n} \end{bmatrix} + \sum_{i=1}^{n} \begin{bmatrix} \frac{\partial a_{1i}}{\partial x_1}x_i & \frac{\partial a_{2i}}{\partial x_1}x_i & \cdots & \frac{\partial a_{mi}}{\partial x_1}x_i \\ \frac{\partial a_{1i}}{\partial x_2}x_i & \frac{\partial a_{2i}}{\partial x_2}x_i & \cdots & \frac{\partial a_{mi}}{\partial x_2}x_i \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial a_{1i}}{\partial x_n}x_i & \frac{\partial a_{2i}}{\partial x_n}x_i & \cdots & \frac{\partial a_{mi}}{\partial x_n}x_i \end{bmatrix}$$

$$= \sum_{i=1}^{n} \begin{bmatrix} a_{1i}\bar{\delta}_{i1} & a_{2i}\bar{\delta}_{i1} & \cdots & a_{mi}\bar{\delta}_{i1} \\ a_{1i}\bar{\delta}_{i2} & a_{2i}\bar{\delta}_{i2} & \cdots & a_{mi}\bar{\delta}_{i2} \\ \vdots & \vdots & \ddots & \vdots \\ a_{1i}\bar{\delta}_{in} & a_{2i}\bar{\delta}_{in} & \cdots & a_{mi}\bar{\delta}_{in} \end{bmatrix} + \sum_{i=1}^{n} \left(\frac{\partial}{\partial \boldsymbol{x}} \begin{bmatrix} a_{1i} & a_{2i} & \cdots & a_{mi} \end{bmatrix}\right) x_i \qquad \text{by Lemma E.1}$$

$$= \begin{bmatrix} a_{11} & a_{21} & \cdots & a_{m1} \\ a_{12} & a_{22} & \cdots & a_{m2} \\ \vdots & \vdots & \ddots & \vdots \\ a_{1n} & a_{2n} & \cdots & a_{mn} \end{bmatrix} + \sum_{i=1}^{n} \left(\frac{\partial}{\partial \boldsymbol{x}} \begin{bmatrix} a_{1i} & a_{2i} & \cdots & a_{mi} \end{bmatrix}\right) x_i \qquad \text{by definition of } \bar{\delta}$$

$$= \mathbf{A}^T + \sum_{i=1}^{n} \left(\frac{\partial}{\partial \boldsymbol{x}} \begin{bmatrix} a_{1i} & a_{2i} & \cdots & a_{mi} \end{bmatrix}\right) x_i$$

Theorem E.3 (Affine equations). [5]

THM *$\mathbf{A}$ and $\mathbf{B}$ are independent of $\boldsymbol{x}$* $\Longrightarrow$

$$\begin{cases} \frac{\partial}{\partial \boldsymbol{x}}(\mathbf{A}\boldsymbol{x}) = \mathbf{A}^T & \forall \boldsymbol{x}\in\mathbb{R}^n,\ \mathbf{A}\in\mathbb{C}^m\times\mathbb{C}^n \\ \frac{\partial}{\partial \boldsymbol{x}}(\boldsymbol{x}^T\mathbf{B}) = \mathbf{B} & \forall \boldsymbol{x}\in\mathbb{R}^n,\ \mathbf{B}\in\mathbb{C}^n\times\mathbb{C}^m \end{cases}$$

✎PROOF: Let $\mathbf{B} \triangleq \mathbf{A}^T$.

1. Proof that $\frac{\partial}{\partial \boldsymbol{x}}(A\boldsymbol{x}) = \mathbf{A}^T$:

$$\frac{\partial}{\partial \boldsymbol{x}}(\mathbf{A}\boldsymbol{x}) = \mathbf{A}^T + \sum_{i=1}^{n} \left(\frac{\partial}{\partial \boldsymbol{x}} \begin{bmatrix} a_{1i} & a_{2i} & \cdots & a_{mi} \end{bmatrix}\right) x_i \qquad \text{by Theorem E.2 page 196}$$

$$= \mathbf{A}^T + \sum_{i=1}^{n} \begin{bmatrix} \frac{\partial}{\partial \boldsymbol{x}}a_{1i} & \frac{\partial}{\partial \boldsymbol{x}}a_{2i} & \cdots & \frac{\partial}{\partial \boldsymbol{x}}a_{mi} \end{bmatrix} x_i$$

$$= \mathbf{A}^T + \sum_{i=1}^{n} \begin{bmatrix} 0 & 0 & \cdots & 0 \end{bmatrix} x_i \qquad \text{by left hypothesis}$$

$$= \mathbf{A}^T$$

2. Proof that $\frac{\partial}{\partial \boldsymbol{x}}(\boldsymbol{x}^T\mathbf{B}) = \mathbf{B}$:

$$\frac{\partial}{\partial \boldsymbol{x}}(\boldsymbol{x}^T\mathbf{B}) = \frac{\partial}{\partial \boldsymbol{x}}(\boldsymbol{x}^T\mathbf{A}^T) \qquad \text{by definition of } \mathbf{B}$$

$$= \frac{\partial}{\partial \boldsymbol{x}}\left[(\mathbf{A}\boldsymbol{x})^T\right]$$

$$= \frac{\partial}{\partial \boldsymbol{x}}(\mathbf{A}\boldsymbol{x}) \qquad \text{by Definition E.2 page 193}$$

$$= \mathbf{A}^T \qquad \text{by Theorem E.3 page 197}$$

$$= \mathbf{B} \qquad \text{by definition of } \mathbf{B}$$

[5] Graham (1981), page 54, Graham (2018), page 549780486824178§"4.2 The Derivatives of Vectors"

COPYRIGHT © 2019 DANIEL J. GREENHOE

Theorem E.4 (Product rule). [6] *Let $\boldsymbol{y}$ and $\boldsymbol{z}$ be functions of $\boldsymbol{x}$ and*

THM $$\frac{\partial}{\partial \boldsymbol{x}}\boldsymbol{z}^T\boldsymbol{y} = \frac{\partial \boldsymbol{z}}{\partial \boldsymbol{x}}\boldsymbol{y} + \frac{\partial \boldsymbol{y}}{\partial \boldsymbol{x}}\boldsymbol{z} \qquad \forall \boldsymbol{x}\in\mathbb{R}^n,\ \boldsymbol{y}\in\mathbb{R}^m,\ \boldsymbol{z}\in\mathbb{R}^m$$

✎PROOF:

$$\begin{aligned}
\frac{\partial}{\partial \boldsymbol{x}}\boldsymbol{z}^T\boldsymbol{y} &= \frac{\partial}{\partial \boldsymbol{x}}\sum_{k=1}^{m} z_k y_k \\
&= \sum_{k=1}^{m}\frac{\partial}{\partial \boldsymbol{x}} z_k y_k \\
&= \sum_{k=1}^{m}\frac{\partial z_k}{\partial \boldsymbol{x}} y_k + \sum_{k=1}^{m}\frac{\partial y_k}{\partial \boldsymbol{x}} z_k \qquad \text{by Lemma E.3 page 195} \\
&= \begin{bmatrix} \frac{\partial z_1}{\partial x_1}y_1 & + & \frac{\partial z_2}{\partial x_1}y_2 & + & \cdots & + & \frac{\partial z_n}{\partial x_1}y_n \\ \frac{\partial z_1}{\partial x_1}y_1 & + & \frac{\partial z_2}{\partial x_1}y_2 & + & \cdots & + & \frac{\partial z_n}{\partial x_1}y_n \\ \vdots & & & & \ddots & & \vdots \\ \frac{\partial z_1}{\partial x_1}y_1 & + & \frac{\partial z_2}{\partial x_1}y_2 & + & \cdots & + & \frac{\partial z_n}{\partial x_1}y_n \end{bmatrix} + \begin{bmatrix} \frac{\partial y_1}{\partial x_1}z_1 & + & \frac{\partial y_2}{\partial x_1}z_2 & + & \cdots & + & \frac{\partial y_n}{\partial x_1}z_n \\ \frac{\partial y_1}{\partial x_1}z_1 & + & \frac{\partial y_2}{\partial x_1}z_2 & + & \cdots & + & \frac{\partial y_n}{\partial x_1}z_n \\ \vdots & & & & \ddots & & \vdots \\ \frac{\partial y_1}{\partial x_1}z_1 & + & \frac{\partial y_2}{\partial x_1}z_2 & + & \cdots & + & \frac{\partial y_n}{\partial x_1}z_n \end{bmatrix} \\
&= \begin{bmatrix} \frac{\partial z_1}{\partial x_1} & \frac{\partial z_2}{\partial x_1} & \cdots & \frac{\partial z_n}{\partial x_1} \\ \frac{\partial z_1}{\partial x_1} & \frac{\partial z_2}{\partial x_1} & \cdots & \frac{\partial z_n}{\partial x_1} \\ \vdots & & \ddots & \vdots \\ \frac{\partial z_1}{\partial x_1} & \frac{\partial z_2}{\partial x_1} & \cdots & \frac{\partial z_n}{\partial x_1} \end{bmatrix}\begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_n \end{bmatrix} + \begin{bmatrix} \frac{\partial y_1}{\partial x_1} & \frac{\partial y_2}{\partial x_1} & \cdots & \frac{\partial y_n}{\partial x_1} \\ \frac{\partial y_1}{\partial x_1} & \frac{\partial y_2}{\partial x_1} & \cdots & \frac{\partial y_n}{\partial x_1} \\ \vdots & & \ddots & \vdots \\ \frac{\partial y_1}{\partial x_1} & \frac{\partial y_2}{\partial x_1} & \cdots & \frac{\partial y_n}{\partial x_1} \end{bmatrix}\begin{bmatrix} z_1 \\ z_2 \\ \vdots \\ z_n \end{bmatrix} \\
&= \frac{\partial \boldsymbol{z}}{\partial \boldsymbol{x}}\boldsymbol{y} + \frac{\partial \boldsymbol{y}}{\partial \boldsymbol{x}}\boldsymbol{z}
\end{aligned}$$

✏

Theorem E.5.

THM $$\frac{\partial}{\partial \boldsymbol{x}}\left(\boldsymbol{x}^T A\boldsymbol{x}\right) = \mathbf{A}\boldsymbol{x} + \mathbf{A}^T\boldsymbol{x} + \left[\sum_{i=1}^{n}\left(\frac{\partial}{\partial \boldsymbol{x}}\begin{bmatrix} a_{1i} & a_{2i} & \cdots & a_{ni}\end{bmatrix}\right)x_i\right]\boldsymbol{x} \qquad \forall \boldsymbol{x}\in\mathbb{R}^n, \mathbf{A}\in\mathbb{R}^n\times\mathbb{R}^n$$

✎PROOF:

$$\begin{aligned}
\frac{\partial}{\partial \boldsymbol{x}}\left(\boldsymbol{x}^T\mathbf{A}\boldsymbol{x}\right) &= \left[\frac{\partial}{\partial \boldsymbol{x}}\boldsymbol{x}\right]\mathbf{A}\boldsymbol{x} + \left[\frac{\partial}{\partial \boldsymbol{x}}\mathbf{A}\boldsymbol{x}\right]\boldsymbol{x} && \text{by Theorem E.4 page 198} \\
&= \mathbf{I}\mathbf{A}\boldsymbol{x} + \left[\mathbf{A}^T + \sum_{i=1}^{n}\left(\frac{\partial}{\partial \boldsymbol{x}}\begin{bmatrix} a_{1i} & a_{2i} & \cdots & a_{ni}\end{bmatrix}\right)x_i\right]\boldsymbol{x} && \text{by Theorem E.1 and Theorem E.2} \\
&= \mathbf{A}\boldsymbol{x} + \mathbf{A}^T\boldsymbol{x} + \left[\sum_{i=1}^{n}\left(\frac{\partial}{\partial \boldsymbol{x}}\begin{bmatrix} a_{1i} & a_{2i} & \cdots & a_{ni}\end{bmatrix}\right)x_i\right]\boldsymbol{x} && \text{by definition of identity operator } \mathbf{I}
\end{aligned}$$

✏

Theorem E.6 (Quadratic form). [7]

THM $$\mathbf{A} \text{ is independent of } \boldsymbol{x} \quad\Longrightarrow\quad \frac{\partial}{\partial \boldsymbol{x}}\left(\boldsymbol{x}^T A\boldsymbol{x}\right) = A\boldsymbol{x} + A^T\boldsymbol{x} \qquad \forall \boldsymbol{x}\in\mathbb{R}^n, \mathbf{A}\in\mathbb{R}^n\times\mathbb{R}^n$$

[6] Scharf (1991), page 274, Trees (2002), page 1398
[7] Graham (1981), page 54

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

✎PROOF:

$$\begin{aligned}\frac{\partial}{\partial \boldsymbol{x}}(\boldsymbol{x}^T\mathbf{A}\boldsymbol{x}) &= \left[\frac{\partial}{\partial \boldsymbol{x}}\boldsymbol{x}\right]\mathbf{A}\boldsymbol{x} + \left[\frac{\partial}{\partial \boldsymbol{x}}\mathbf{A}\boldsymbol{x}\right]\boldsymbol{x} && \text{by Theorem E.4 page 198}\\ &= \mathbf{I}\mathbf{A}\boldsymbol{x} + \mathbf{A}^T\boldsymbol{x} && \text{by Theorem E.1 page 195 and Theorem E.3 page 197}\end{aligned}$$

✏

Corollary E.1. [8]

COR $$\frac{\partial}{\partial \boldsymbol{x}}(\boldsymbol{x}^T\boldsymbol{x}) = 2\boldsymbol{x} \qquad \forall \boldsymbol{x} \in \mathbb{R}^n$$

✎PROOF:

$$\begin{aligned}\frac{\partial}{\partial \boldsymbol{x}}(\boldsymbol{x}^T\boldsymbol{x}) &= \frac{\partial}{\partial \boldsymbol{x}}(\boldsymbol{x}^T\mathbf{I}\boldsymbol{x}) && \text{by property of identity operator } I\\ &= \mathbf{I}\boldsymbol{x} + \mathbf{I}^T\boldsymbol{x} && \text{by previous result 3.}\\ &= \boldsymbol{x} + \boldsymbol{x} && \text{by property of identity operator } I\\ &= 2\boldsymbol{x}\end{aligned}$$

✏

Theorem E.7 (Chain rule). [9] *Let $\boldsymbol{z}$ be a function of $\boldsymbol{y}$ and $\boldsymbol{y}$ a function of $\boldsymbol{x}$ and*

$$\boldsymbol{x} \triangleq \begin{bmatrix} x_1\\ x_2\\ \vdots\\ x_n\end{bmatrix} \qquad \boldsymbol{y} \triangleq \begin{bmatrix} y_1\\ y_2\\ \vdots\\ y_m\end{bmatrix} \qquad \boldsymbol{z} \triangleq \begin{bmatrix} z_1\\ z_2\\ \vdots\\ z_k\end{bmatrix}$$

THM $$\frac{\partial}{\partial \boldsymbol{x}}\boldsymbol{z} = \frac{\partial \boldsymbol{y}}{\partial \boldsymbol{x}}\frac{\partial \boldsymbol{z}}{\partial \boldsymbol{y}}$$

✎PROOF:

$$\begin{aligned}\frac{\partial \boldsymbol{z}}{\partial \boldsymbol{x}} &= \begin{bmatrix} \frac{\partial z_1}{\partial x_1} & \frac{\partial z_2}{\partial x_1} & \cdots & \frac{\partial z_k}{\partial x_1}\\ \frac{\partial z_1}{\partial x_2} & \frac{\partial z_2}{\partial x_2} & \cdots & \frac{\partial z_k}{\partial x_2}\\ \vdots & \vdots & \ddots & \vdots\\ \frac{\partial z_1}{\partial x_n} & \frac{\partial z_2}{\partial x_n} & \cdots & \frac{\partial z_k}{\partial x_n}\end{bmatrix}\\ &= \begin{bmatrix} \sum_{j=0}^{m}\frac{\partial z_1}{\partial y_j}\frac{\partial y_j}{\partial x_1} & \sum_{j=0}^{m}\frac{\partial z_2}{\partial y_j}\frac{\partial y_j}{\partial x_1} & \cdots & \sum_{j=0}^{m}\frac{\partial z_k}{\partial y_j}\frac{\partial y_j}{\partial x_1}\\ \sum_{j=0}^{m}\frac{\partial z_1}{\partial y_j}\frac{\partial y_j}{\partial x_2} & \sum_{j=0}^{m}\frac{\partial z_2}{\partial y_j}\frac{\partial y_j}{\partial x_2} & \cdots & \sum_{j=0}^{m}\frac{\partial z_k}{\partial y_j}\frac{\partial y_j}{\partial x_2}\\ \vdots & \vdots & \ddots & \vdots\\ \sum_{j=0}^{m}\frac{\partial z_1}{\partial y_j}\frac{\partial y_j}{\partial x_n} & \sum_{j=0}^{m}\frac{\partial z_2}{\partial y_j}\frac{\partial y_j}{\partial x_n} & \cdots & \sum_{j=0}^{m}\frac{\partial z_k}{\partial y_j}\frac{\partial y_j}{\partial x_n}\end{bmatrix}\\ &= \begin{bmatrix} \frac{\partial y_1}{\partial x_1} & \frac{\partial y_2}{\partial x_1} & \cdots & \frac{\partial y_m}{\partial x_1}\\ \frac{\partial y_1}{\partial x_2} & \frac{\partial y_2}{\partial x_2} & \cdots & \frac{\partial y_m}{\partial x_2}\\ \vdots & \vdots & \ddots & \vdots\\ \frac{\partial y_1}{\partial x_n} & \frac{\partial y_2}{\partial x_n} & \cdots & \frac{\partial y_m}{\partial x_n}\end{bmatrix}\begin{bmatrix} \frac{\partial z_1}{\partial y_1} & \frac{\partial z_2}{\partial y_1} & \cdots & \frac{\partial z_k}{\partial y_1}\\ \frac{\partial z_1}{\partial y_2} & \frac{\partial z_2}{\partial y_2} & \cdots & \frac{\partial z_k}{\partial y_2}\\ \vdots & \vdots & \ddots & \vdots\\ \frac{\partial z_1}{\partial y_m} & \frac{\partial z_2}{\partial y_m} & \cdots & \frac{\partial z_k}{\partial y_m}\end{bmatrix}\\ &= \frac{\partial \boldsymbol{y}}{\partial \boldsymbol{x}}\frac{\partial \boldsymbol{z}}{\partial \boldsymbol{y}}\end{aligned}$$

[8] Graham (1981), page 54
[9] Graham (1981), pages 54–55

COPYRIGHT © 2019 DANIEL J. GREENHOE

E.2 First derivative of a matrix with respect to a scalar

Definition E.3. *Let* $x \in \mathbb{R}$, $\{y_{jk} \in \mathbb{C} | j = 1, 2, \ldots, m;\ k = 1, 2, \ldots, n\}$ *and*

$$Y = \underbrace{\begin{bmatrix} y_{11} & y_{12} & \cdots & y_{1n} \\ y_{21} & y_{22} & \cdots & y_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ y_{m1} & y_{m2} & \cdots & y_{mn} \end{bmatrix}}_{m \times n \text{ matrix}}$$

The ***derivative of*** Y ***with respect to*** x *is*

DEF

$$\frac{\mathrm{d}Y}{\mathrm{d}x} \triangleq \underbrace{\begin{bmatrix} \frac{\mathrm{d}y_{11}}{\mathrm{d}x} & \frac{\mathrm{d}y_{12}}{\mathrm{d}x} & \cdots & \frac{\mathrm{d}y_{1n}}{\mathrm{d}x} \\ \frac{\mathrm{d}y_{21}}{\mathrm{d}x} & \frac{\mathrm{d}y_{22}}{\mathrm{d}x} & \cdots & \frac{\mathrm{d}y_{2n}}{\mathrm{d}x} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\mathrm{d}y_{m1}}{\mathrm{d}x} & \frac{\mathrm{d}y_{m2}}{\mathrm{d}x} & \cdots & \frac{\mathrm{d}y_{mn}}{\mathrm{d}x} \end{bmatrix}}_{m \times n \text{ matrix}}$$

Theorem E.8. [10] *Let* $x \in \mathbb{R}$, $\{y_{jp} \in \mathbb{C} | j = 1, 2, \ldots, m;\ p = 1, 2, \ldots, n\}$, $\{w_{jp} \in \mathbb{C} | j = 1, 2, \ldots, n;\ p = 1, 2, \ldots, k\}$, *and*

$$Y = \underbrace{\begin{bmatrix} y_{11} & y_{12} & \cdots & y_{1n} \\ y_{21} & y_{22} & \cdots & y_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ y_{m1} & y_{m2} & \cdots & y_{mn} \end{bmatrix}}_{m \times n \text{ matrix}} \qquad W = \underbrace{\begin{bmatrix} w_{11} & w_{12} & \cdots & w_{1k} \\ w_{21} & w_{22} & \cdots & w_{2k} \\ \vdots & \vdots & \ddots & \vdots \\ w_{p1} & w_{p2} & \cdots & w_{pk} \end{bmatrix}}_{p \times k \text{ matrix}}$$

THM

$$\begin{aligned} \frac{\mathrm{d}}{\mathrm{d}x}\Big(Y + W\Big) &= \frac{\mathrm{d}}{\mathrm{d}x}Y + \frac{\mathrm{d}}{\mathrm{d}x}W && \text{(for } p = m,\ k = n\text{)} \\ \frac{\mathrm{d}}{\mathrm{d}x}\Big(YW\Big) &= \Big(\frac{\mathrm{d}}{\mathrm{d}x}Y\Big)W + Y\Big(\frac{\mathrm{d}}{\mathrm{d}x}W\Big) && \text{(for } p = n\text{)} \\ \frac{\mathrm{d}}{\mathrm{d}x}\Big(Y^T\Big) &= \Big(\frac{\mathrm{d}}{\mathrm{d}x}Y\Big)^T \\ \frac{\mathrm{d}}{\mathrm{d}x}\Big(Y^{-1}\Big) &= -Y^{-1}\Big(\frac{\mathrm{d}}{\mathrm{d}x}Y\Big)Y^{-1} && \text{(for } m = n \text{ and } Y \text{ invertible)} \end{aligned}$$

✎PROOF:

$$\begin{aligned} \frac{\mathrm{d}}{\mathrm{d}x}\Big(Y + W\Big) &= \frac{\mathrm{d}}{\mathrm{d}x}\left(\begin{bmatrix} y_{11} & y_{12} & \cdots & y_{1n} \\ y_{21} & y_{22} & \cdots & y_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ y_{m1} & y_{m2} & \cdots & y_{mn} \end{bmatrix} + \begin{bmatrix} w_{11} & w_{12} & \cdots & w_{1n} \\ w_{21} & w_{22} & \cdots & w_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ w_{m1} & w_{m2} & \cdots & w_{mn} \end{bmatrix}\right) \\ &= \frac{\mathrm{d}}{\mathrm{d}x}\begin{bmatrix} y_{11} + w_{11} & y_{12} + w_{12} & \cdots & y_{1n} + w_{1n} \\ y_{21} + w_{21} & y_{22} + w_{22} & \cdots & y_{2n} + w_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ y_{m1} + w_{m1} & y_{m2} + w_{m2} & \cdots & y_{mn} + w_{mn} \end{bmatrix} \end{aligned}$$

[10] Gradshteyn and Ryzhik (1980), pages 1106–1107

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

$$
\begin{aligned}
&= \begin{bmatrix} (y_{11}+w_{11})' & (y_{12}+w_{12})' & \cdots & (y_{1n}+w_{1n})' \\ (y_{21}+w_{21})' & (y_{22}+w_{22})' & \cdots & (y_{2n}+w_{2n})' \\ \vdots & \vdots & \ddots & \vdots \\ (y_{m1}+w_{m1})' & (y_{m2}+w_{m2})' & \cdots & (y_{mn}+w_{mn})' \end{bmatrix} \\
&= \begin{bmatrix} y'_{11}+w'_{11} & y'_{12}+w'_{12} & \cdots & y'_{1n}+w'_{1n} \\ y'_{21}+w'_{21} & y'_{22}+w'_{22} & \cdots & y'_{2n}+w'_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ y'_{m1}+w'_{m1} & y'_{m2}+w'_{m2} & \cdots & y'_{mn}+w'_{mn} \end{bmatrix} \\
&= \begin{bmatrix} y'_{11} & y'_{12} & \cdots & y'_{1n} \\ y'_{21} & y'_{22} & \cdots & y'_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ y'_{m1} & y'_{m2} & \cdots & y'_{mn} \end{bmatrix} + \begin{bmatrix} w'_{11} & w'_{12} & \cdots & w'_{1n} \\ w'_{21} & w'_{22} & \cdots & w'_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ w'_{m1} & w'_{m2} & \cdots & w'_{mn} \end{bmatrix} \\
&= \frac{\mathrm{d}}{\mathrm{d}x} \begin{bmatrix} y_{11} & y_{12} & \cdots & y_{1n} \\ y_{21} & y_{22} & \cdots & y_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ y_{m1} & y_{m2} & \cdots & y_{mn} \end{bmatrix} + \frac{\mathrm{d}}{\mathrm{d}x} \begin{bmatrix} w_{11} & w_{12} & \cdots & w_{1n} \\ w_{21} & w_{22} & \cdots & w_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ w_{m1} & w_{m2} & \cdots & w_{mn} \end{bmatrix} \\
&= \frac{\mathrm{d}}{\mathrm{d}x} Y + \frac{\mathrm{d}}{\mathrm{d}x} W
\end{aligned}
$$

$$
\begin{aligned}
\frac{\mathrm{d}}{\mathrm{d}x}\Big(YW\Big) &= \frac{\mathrm{d}}{\mathrm{d}x}\left(\begin{bmatrix} y_{11} & y_{12} & \cdots & y_{1n} \\ y_{21} & y_{22} & \cdots & y_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ y_{m1} & y_{m2} & \cdots & y_{mn} \end{bmatrix} \begin{bmatrix} w_{11} & w_{12} & \cdots & w_{1k} \\ w_{21} & w_{22} & \cdots & w_{2k} \\ \vdots & \vdots & \ddots & \vdots \\ w_{n1} & w_{n2} & \cdots & w_{nk} \end{bmatrix} \right) \\
&= \frac{\mathrm{d}}{\mathrm{d}x} \begin{bmatrix} \sum_{j=1}^{n} y_{1j}w_{j1} & \sum_{j=1}^{n} y_{1j}w_{j2} & \cdots & \sum_{j=1}^{n} y_{1j}w_{jk} \\ \sum_{j=1}^{n} y_{2j}w_{j1} & \sum_{j=1}^{n} y_{2j}w_{j2} & \cdots & \sum_{j=1}^{n} y_{2j}w_{jk} \\ \vdots & \vdots & \ddots & \vdots \\ \sum_{j=1}^{n} y_{mj}w_{j1} & \sum_{j=1}^{n} y_{mj}w_{j2} & \cdots & \sum_{j=1}^{n} y_{mj}w_{jk} \end{bmatrix} \\
&= \frac{\mathrm{d}}{\mathrm{d}x} \sum_{j=1}^{n} \begin{bmatrix} y_{1j}w_{j1} & y_{1j}w_{j2} & \cdots & y_{1j}w_{jk} \\ y_{2j}w_{j1} & y_{2j}w_{j2} & \cdots & y_{2j}w_{jk} \\ \vdots & \vdots & \ddots & \vdots \\ y_{mj}w_{j1} & y_{mj}w_{j2} & \cdots & y_{mj}w_{jk} \end{bmatrix} \\
&= \sum_{j=1}^{n} \frac{\mathrm{d}}{\mathrm{d}x} \begin{bmatrix} y_{1j}w_{j1} & y_{1j}w_{j2} & \cdots & y_{1j}w_{jk} \\ y_{2j}w_{j1} & y_{2j}w_{j2} & \cdots & y_{2j}w_{jk} \\ \vdots & \vdots & \ddots & \vdots \\ y_{mj}w_{j1} & y_{mj}w_{j2} & \cdots & y_{mj}w_{jk} \end{bmatrix} \\
&= \sum_{j=1}^{n} \begin{bmatrix} \frac{\mathrm{d}}{\mathrm{d}x}(y_{1j}w_{j1}) & \frac{\mathrm{d}}{\mathrm{d}x}(y_{1j}w_{j2}) & \cdots & \frac{\mathrm{d}}{\mathrm{d}x}(y_{1j}w_{jk}) \\ \frac{\mathrm{d}}{\mathrm{d}x}(y_{2j}w_{j1}) & \frac{\mathrm{d}}{\mathrm{d}x}(y_{2j}w_{j2}) & \cdots & \frac{\mathrm{d}}{\mathrm{d}x}(y_{2j}w_{jk}) \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\mathrm{d}}{\mathrm{d}x}(y_{mj}w_{j1}) & \frac{\mathrm{d}}{\mathrm{d}x}(y_{mj}w_{j2}) & \cdots & \frac{\mathrm{d}}{\mathrm{d}x}(y_{mj}w_{jk}) \end{bmatrix} \\
&= \sum_{j=1}^{n} \begin{bmatrix} \vdots & \vdots & \ddots & \vdots \\ y'_{1j}w_{j1}+y_{1j}w'_{j1} & y'_{1j}w_{j2}+y_{1j}w'_{j2} & \cdots & y'_{1j}w_{jk}+y_{1j}w'_{jk} \\ y'_{2j}w_{j1}+y_{2j}w'_{j1} & y'_{2j}w_{j2}+y_{2j}w'_{j2} & \cdots & y'_{2j}w_{jk}+y_{2j}w'_{jk} \\ y'_{mj}w_{j1}+y_{mj}w'_{j1} & y'_{mj}w_{j2}+y_{mj}w'_{j2} & \cdots & y'_{mj}w_{jk}+y_{mj}w'_{jk} \end{bmatrix}
\end{aligned}
$$

Copyright © 2019 Daniel J. Greenhoe

$$= \sum_{j=1}^{n} \left(\begin{bmatrix} y'_{1j}w_{j1} & y'_{1j}w_{j2} & \cdots & y'_{1j}w_{jk} \\ y'_{2j}w_{j1} & y'_{2j}w_{j2} & \cdots & y'_{2j}w_{jk} \\ \vdots & \vdots & \ddots & \vdots \\ y'_{mj}w_{j1} & y'_{mj}w_{j2} & \cdots & y'_{mj}w_{jk} \end{bmatrix} + \begin{bmatrix} y_{1j}w'_{j1} & y_{1j}w'_{j2} & \cdots & y_{1j}w'_{jk} \\ y_{2j}w'_{j1} & y_{2j}w'_{j2} & \cdots & y_{2j}w'_{jk} \\ \vdots & \vdots & \ddots & \vdots \\ y_{mj}w'_{j1} & y_{mj}w'_{j2} & \cdots & y_{mj}w'_{jk} \end{bmatrix} \right)$$

$$= \left(\frac{\mathrm{d}}{\mathrm{d}x}Y\right)W + Y\left(\frac{\mathrm{d}}{\mathrm{d}x}W\right)$$

$$\frac{\mathrm{d}}{\mathrm{d}x}\left(Y^T\right) = \frac{\mathrm{d}}{\mathrm{d}x}\left(\begin{bmatrix} y_{11} & y_{12} & \cdots & y_{1n} \\ y_{21} & y_{22} & \cdots & y_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ y_{m1} & y_{m2} & \cdots & y_{mn} \end{bmatrix}^T \right)$$

$$= \frac{\mathrm{d}}{\mathrm{d}x} \begin{bmatrix} y_{11} & y_{21} & \cdots & y_{n1} \\ y_{12} & y_{22} & \cdots & y_{n2} \\ \vdots & \vdots & \ddots & \vdots \\ y_{1n} & y_{2n} & \cdots & y_{nm} \end{bmatrix}$$

$$= \begin{bmatrix} y'_{11} & y'_{21} & \cdots & y'_{n1} \\ y'_{12} & y'_{22} & \cdots & y'_{n2} \\ \vdots & \vdots & \ddots & \vdots \\ y'_{1n} & y'_{2n} & \cdots & y'_{nm} \end{bmatrix}$$

$$= \begin{bmatrix} y'_{11} & y'_{12} & \cdots & y'_{1n} \\ y'_{21} & y'_{22} & \cdots & y'_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ y'_{m1} & y'_{m2} & \cdots & y'_{mn} \end{bmatrix}^T$$

$$= \left(\frac{\mathrm{d}}{\mathrm{d}x} \begin{bmatrix} y_{11} & y_{12} & \cdots & y_{1n} \\ y_{21} & y_{22} & \cdots & y_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ y_{m1} & y_{m2} & \cdots & y_{mn} \end{bmatrix} \right)^T$$

$$\frac{\mathrm{d}}{\mathrm{d}x}\left(Y^{-1}\right) = \frac{\mathrm{d}}{\mathrm{d}x}\frac{\mathrm{adj}Y}{|Y|}$$

$$\vdots$$

no proof at this time

$$\vdots$$

$$= -Y^{-1}\left(\frac{\mathrm{d}}{\mathrm{d}x}Y\right)Y^{-1}$$

E.3 Second derivative of a scalar with respect to a vector

Definition E.4. [11] *Let*

$$\boldsymbol{x} \triangleq \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}$$

[11] Lieb and Loss (2001), page 240, Horn and Johnson (1990), page 167

BY-NC-ND Copyright © 2019 Daniel J. Greenhoe

*The **Hessian matrix** of a scalar y with respect to the vector $\boldsymbol{x}$ is*

DEF

$$\frac{\partial^2 y}{\partial \boldsymbol{x}^2} = \frac{\partial}{\partial \boldsymbol{x}}\left(\frac{\partial y}{\partial \boldsymbol{x}}\right) = \frac{\partial}{\partial \boldsymbol{x}}\begin{bmatrix} \frac{\partial y}{\partial x_1} \\ \frac{\partial y}{\partial x_2} \\ \vdots \\ \frac{\partial y}{\partial x_n} \end{bmatrix} = \underbrace{\begin{bmatrix} \frac{\partial}{\partial x_1}\frac{\partial y}{\partial x_1} & \frac{\partial}{\partial x_1}\frac{\partial y}{\partial x_2} & \cdots & \frac{\partial}{\partial x_1}\frac{\partial y}{\partial x_n} \\ \frac{\partial}{\partial x_2}\frac{\partial y}{\partial x_1} & \frac{\partial}{\partial x_2}\frac{\partial y}{\partial x_2} & \cdots & \frac{\partial}{\partial x_2}\frac{\partial y}{\partial x_n} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial}{\partial x_n}\frac{\partial y}{\partial x_1} & \frac{\partial}{\partial x_n}\frac{\partial y}{\partial x_2} & \cdots & \frac{\partial}{\partial x_n}\frac{\partial y}{\partial x_n} \end{bmatrix}}_{n \times n \text{ matrix}}$$

E.4 Multiple derivatives of a vector with respect to a scalar

Definition E.5. *Let*

$$\boldsymbol{y} \triangleq \begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_m \end{bmatrix}$$

The derivative of a vector $\boldsymbol{y}$ with respect to the scalar x is

DEF

$$\begin{bmatrix} \boldsymbol{y} \\ \frac{\mathrm{d}}{\mathrm{d}x}\boldsymbol{y} \\ \frac{\mathrm{d}^2}{\mathrm{d}x^2}\boldsymbol{y} \\ \vdots \\ \frac{\mathrm{d}^n}{\mathrm{d}x^n}\boldsymbol{y} \end{bmatrix} = \underbrace{\begin{bmatrix} y_1 & y_2 & \cdots & y_m \\ \frac{\mathrm{d}}{\mathrm{d}x}y_1 & \frac{\mathrm{d}}{\mathrm{d}x}y_2 & \cdots & \frac{\mathrm{d}}{\mathrm{d}x}y_m \\ \frac{\mathrm{d}^2}{\mathrm{d}x^2}y_1 & \frac{\mathrm{d}^2}{\mathrm{d}x^2}y_2 & \cdots & \frac{\mathrm{d}^2}{\mathrm{d}x^2}y_m \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\mathrm{d}^n}{\mathrm{d}x^n}y_1 & \frac{\mathrm{d}^n}{\mathrm{d}x^n}y_2 & \cdots & \frac{\mathrm{d}^n}{\mathrm{d}x^n}y_m \end{bmatrix}}_{(n+1) \times m \text{ matrix}}$$

Copyright © 2019 Daniel J. Greenhoe

Copyright © 2019 Daniel J. Greenhoe

APPENDIX F

ALGEBRAIC STRUCTURES

❝ In this book, learned reader, you have the rules of algebra... It unties the knot not only where one term is equal to another or two to one but also where two are equal to two or three to one.... this most abstruse and unsurpassed treasury of the entire arithmetic being brought to light and, as in a theater, exposed to the sight of all...❞

Gerolamo Cardano (1501–1576), Italian mathematician, physician, and astrologer [1]

A set together with one or more operations forms several standard mathematical structures:

$$\textit{group} \supseteq \textit{ring} \supseteq \textit{commutative ring} \supseteq \textit{integral domain} \supseteq \textit{field}$$

Definition F.1. [2] *Let X be a set and $\diamond : X \times X \to X$ be an operation on X.*

DEF

*The pair $(X, \diamond)$ is a **group** if*

1.	$\exists e \in X$ such that	$e \diamond x = x \diamond e = x$	$\forall x \in X$	(IDENTITY *element*)	*and*
2.	$\exists (-x) \in X$ such that	$(-x) \diamond x = x \diamond (-x) = e$	$\forall x \in X$	(INVERSE *element*)	*and*
3.		$x \diamond (y \diamond z) = (x \diamond y) \diamond z$	$\forall x,y,z \in X$	(ASSOCIATIVE)	

Definition F.2. [3] *Let $+ : X \times X \to X$ and $* : X \times X \to X$ be operations on a set X. Furthermore, let the operation $*$ also be represented by juxtapostion as in $a * b \equiv ab$.*

DEF

*The triple $(X, +, *)$ is a **ring** if*

1.	$(X, +)$ *is a group.*			*(additive group)*	*and*
2.	$x(yz) = (xy)z$		$\forall x,y,z \in X$	(ASSOCIATIVE *with respect to* $*$)	*and*
3.	$x(y+z) = (xy) + (xz)$		$\forall x,y,z \in X$	($*$ *is* LEFT DISTRIBUTIVE *over* $+$)	*and*
4.	$(x+y)z = (xz) + (yz)$		$\forall x,y,z \in X$	($*$ *is* RIGHT DISTRIBUTIVE *over* $+$).	

Definition F.3. [4]

[1] quote: Cardano (1545), page 1
image: http://en.wikipedia.org/wiki/Image:Cardano.jpg
[2] Durbin (2000), page 29
[3] Durbin (2000), pages 114–115
[4] Durbin (2000), page 118

DEF

A triple $(X,+,*)$ *is a* ***commutative ring*** *if*

1. $(X,+,*)$ *is a* RING *and*
2. $xy = yx \quad \forall x,y \in X$ (COMMUTATIVE).

Definition F.4. [5] *Let* R *be a* COMMUTATIVE RING *(Definition F.3 page 205).*

DEF

A function $|\cdot|$ *in* $\mathbb{R}^{\mathbb{R}}$ *is an* ***absolute value*** *(or* ***modulus****) if*

1. $|x| \geq 0 \quad x \in \mathbb{R}$ (NON-NEGATIVE) *and*
2. $|x| = 0 \iff x = 0 \quad x \in \mathbb{R}$ (NONDEGENERATE) *and*
3. $|xy| = |x| \cdot |y| \quad x,y \in \mathbb{R}$ (HOMOGENEOUS / SUBMULTIPLICATIVE) *and*
4. $|x+y| \leq |x| + |y| \quad x,y \in \mathbb{R}$ (SUBADDITIVE / TRIANGLE INEQUALITY)

Definition F.5. [6]

DEF

The structure $F \triangleq (X,+,\cdot,0,1)$ *is a* ***field*** *if*

1. $(X,+,*)$ *is a ring* *(ring)* *and*
2. $xy = yx \quad \forall x,y \in X$ *(commutative with respect to* $*$*)* *and*
3. $(X \setminus \{0\},*)$ *is a group* *(group with respect to* $*$*).*

Definition F.6. [7] *Let* $\mathsf{V} = (F,+,\cdot)$ *be a vector space and* $\otimes : \mathsf{V} \times \mathsf{V} \to \mathsf{V}$ *be a vector-vector multiplication operator.*
An ***algebra*** *is any pair* $(\mathsf{V},\otimes)$ *that satisfies* ($\otimes$ *is represented by juxtaposition*)

DEF

1. $(\boldsymbol{ux})\boldsymbol{y} = \boldsymbol{u}(\boldsymbol{xy}) \quad \forall \boldsymbol{u},\boldsymbol{x},\boldsymbol{y} \in \mathsf{V}$ (ASSOCIATIVE) *and*
2. $\boldsymbol{u}(\boldsymbol{x}+\boldsymbol{y}) = (\boldsymbol{ux}) + (\boldsymbol{uy}) \quad \forall \boldsymbol{u},\boldsymbol{x},\boldsymbol{y} \in \mathsf{V}$ (LEFT DISTRIBUTIVE) *and*
3. $(\boldsymbol{u}+\boldsymbol{x})\boldsymbol{y} = (\boldsymbol{uy}) + (\boldsymbol{xy}) \quad \forall \boldsymbol{u},\boldsymbol{x},\boldsymbol{y} \in \mathsf{V}$ (RIGHT DISTRIBUTIVE) *and*
4. $\alpha(\boldsymbol{xy}) = (\alpha\boldsymbol{x})\boldsymbol{y} = \boldsymbol{x}(\alpha\boldsymbol{y}) \quad \forall \boldsymbol{x},\boldsymbol{y} \in \mathsf{V}$ *and* $\alpha \in F$ (SCALAR COMMUTATIVE) .

[5] Cohn (2002) page 312
[6] Durbin (2000), page 123, Weber (1893)
[7] Abramovich and Aliprantis (2002), page 3, Michel and Herget (1993), page 56

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

APPENDIX G

LINEAR SPACES

❝*The geometric calculus, in general, consists in a system of operations on geometric entities, and their consequences, analogous to those that algebra has on the numbers. It permits the expression in formulas of the results of geometric constructions, the representation with equations of propositions of geometry, and the substitution of a transformation of equations for a verbal argument.*❞

Giuseppe Peano (1858–1932), Italian mathematician, credited with being one of the first to introduce the concept of the *linear space* (*vector space*).[1]

G.1 Definition and basic results

A *metric space* is a *set* together with nothing else save a *metric* that gives the space a *topology* (Definition **??** page **??**). A *linear space* (next definition) in general has no topology but does have some additional *algebraic* structure (APPENDIX F page 205) that is useful in generalizing a number of mathematical concepts. If one wishes to have both algebraic structure and a topology, then this can be accomplished by appending a *topology* to a *linear space* giving a *topological linear space* (Definition **??** page **??**), a *metric* giving a *metric linear space*, an *inner product* giving an *inner product space* (Definition J.1 page 255), or a *norm* giving a *normed linear space* (Definition K.1 page 271).

Definition G.1. [2] *Let* $(\mathbb{F}, +, \cdot, 0, 1)$ *be a* FIELD *(Definition F.5 page 206). Let* X *be a set, let* $+$ *be an* OPERATOR *(Definition N.1 page 303) in* X^{X^2}*, and let* $\otimes$ *be an operator in* $X^{\mathbb{F}\times X}$.

[1] quote: Peano (1888b), page ix
image http://en.wikipedia.org/wiki/File:Giuseppe_Peano.jpg, public domain

[2] Kubrusly (2001) pages 40–41 ⟨Definition 2.1 and following remarks⟩, Haaser and Sullivan (1991), page 41, Halmos (1948), pages 1–2, Peano (1888a) ⟨Chapter IX⟩, Peano (1888b), pages 119–120, Banach (1922) pages 134–135



Figure G.1: Lattice of mathematical spaces

DEF

The structure $\boldsymbol{\Omega} \triangleq \big(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times})\big)$ *is a* ***linear space*** *over* $(\mathbb{F}, +, \cdot, 0, 1)$ *if*

1.	$\exists \mathbb{0} \in X$	such that	$\boldsymbol{x} + \mathbb{0} = \boldsymbol{x}$	$\forall \boldsymbol{x} \in X$	(+ IDENTITY)	*
2.	$\exists \boldsymbol{y} \in X$	such that	$\boldsymbol{x} + \boldsymbol{y} = \mathbb{0}$	$\forall \boldsymbol{x} \in X$	(+ INVERSE)	
3.			$(\boldsymbol{x} + \boldsymbol{y}) + \boldsymbol{z} = \boldsymbol{x} + (\boldsymbol{y} + \boldsymbol{z})$	$\forall \boldsymbol{x}, \boldsymbol{y}, \boldsymbol{z} \in X$	(+ *is* ASSOCIATIVE)	
4.			$\boldsymbol{x} + \boldsymbol{y} = \boldsymbol{y} + \boldsymbol{x}$	$\forall \boldsymbol{x}, \boldsymbol{y} \in X$	(+ *is* COMMUTATIVE)	
5.			$1 \cdot \boldsymbol{x} = \boldsymbol{x}$	$\forall \boldsymbol{x} \in X$	($\cdot$ IDENTITY)	
6.			$\alpha \cdot (\beta \cdot \boldsymbol{x}) = (\alpha \cdot \beta) \cdot \boldsymbol{x}$	$\forall \alpha, \beta \in S$ *and* $\boldsymbol{x} \in X$	($\cdot$ ASSOCIATES *with* $\cdot$)	
7.			$\alpha \cdot (\boldsymbol{x} + \boldsymbol{y}) = (\alpha \cdot \boldsymbol{x}) + (\alpha \cdot \boldsymbol{y})$	$\forall \alpha \in S$ *and* $\boldsymbol{x}, \boldsymbol{y} \in X$	($\cdot$ DISTRIBUTES *over* +)	
8.			$(\alpha + \beta) \cdot \boldsymbol{x} = (\alpha \cdot \boldsymbol{x}) + (\beta \cdot \boldsymbol{x})$	$\forall \alpha, \beta \in S$ *and* $\boldsymbol{x} \in X$	($\cdot$ PSEUDO-DISTRIBUTES *over* +)	

The set X *is called the* ***underlying set****. The elements of* X *are called* ***vectors****. The elements of* $\mathbb{F}$ *are called* ***scalars****. A* LINEAR SPACE *is also called a* ***vector space****. If* $\mathbb{F} \triangleq \mathbb{R}$*, then* $\boldsymbol{\Omega}$ *is a* ***real linear space****. If* $\mathbb{F} \triangleq \mathbb{C}$*, then* $\boldsymbol{\Omega}$ *is a* ***complex linear space****.*

Definition G.2. *Let* $\boldsymbol{L}_1 \triangleq \big(X_1, +, \cdot, (\mathbb{F}_1, \dot{+}, \dot{\times})\big)$ *and* $\boldsymbol{L}_2 \triangleq \big(X_2, +, \cdot, (\mathbb{F}_2, \dot{+}, \dot{\times})\big)$.

DEF

$\boldsymbol{\Omega}_2$ *is a* ***linear subspace*** *of* $\boldsymbol{\Omega}_1$ *if*

1. $\boldsymbol{L}_1$ *is a* LINEAR SPACE $(Definition G.1 page 207)$ *and*
2. $\boldsymbol{L}_2$ *is a* LINEAR SPACE $(Definition G.1 page 207)$ *and*
3. $\mathbb{F}_2 \subseteq \mathbb{F}_1$ *and*
4. $X_2 \subseteq X_1$ *and*

Remark G.1. [3] By the first four conditions (*]) listed in Definition G.1, $(X, +)$ is a **commutative group** (or **abelian group**).

[3] Akhiezer and Glazman (1993), page 1, Haaser and Sullivan (1991), page 41

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

Often when discussing a linear space, the operator $\cdot$ is simply expressed with juxtaposition (e.g. $\alpha\boldsymbol{x}$ is equivalent to $\alpha \cdot \boldsymbol{x}$). In doing this, there is no risk of ambiguity between scalar-vector multiplication and scalar-scalar multiplication because the operands uniquely identify the precise operator.[4]

Example G.1 (tuples in $\mathbb{F}^N$). [5] Let $(\!|x_n|\!)_1^N$ be an N*-tuple* (Definition Q.1 page 353) over a *field* (Definition F.5 page 206) $(\mathbb{F}, +, \cdot, 0, 1)$.

EX

$$\begin{aligned} \text{Let} \quad X &\triangleq \{(\!|x_n|\!)_1^N \,|\, x_n \in \mathbb{F}\} && \text{and} \\ (\!|x_n|\!)_1^N + (\!|y_n|\!)_1^N &\triangleq (\!|x_n \dot{+} y_n|\!)_1^N && \forall (\!|x_n|\!) \in X \quad \text{and} \\ \alpha \cdot (\!|x_n|\!)_1^N &\triangleq (\!|\alpha \dot{\times} x_n|\!)_1^N && \forall (\!|x_n|\!) \in X, \alpha \in \mathbb{F}. \end{aligned}$$

Then the structure $\big(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times})\big)$ is a *linear space*.

Example G.2 (real numbers). [6] Let $(\mathbb{R}, +, \cdot, 0, 1)$ be the field of real numbers.

EX

The structure $(\mathbb{R}, +, \cdot, (\mathbb{R}, +, \cdot))$ is a *linear space*.
That is, the field of real numbers forms a linear space over itself.

Example G.3 (functions). [7] Let $(\mathbb{F}, +, \cdot, 0, 1)$ be a field. Let Y^X be the set of all functions with domain X and range Y.

EX

$$\begin{aligned} \text{Let} \quad [\mathsf{f} + \mathsf{g}](x) &\triangleq \mathsf{f}(x) + \mathsf{g}(x) && \forall \mathsf{f}, \mathsf{g} \in Y^X \quad (\textit{pointwise addition}) \quad \text{and} \\ [\alpha \cdot \mathsf{f}](x) &\triangleq \alpha \cdot [\mathsf{f}(x)] && \forall \mathsf{f} \in Y^X, \alpha \in \mathbb{F}. \end{aligned}$$

Then the structure $\big(Y^X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times})\big)$ is a *linear space*.

Example G.4 (functions onto $\mathbb{F}$). [8] Let $(\mathbb{F}, +, \cdot, 0, 1)$ be a field. Let $\mathbb{F}^{\boldsymbol{X}}$ be the set of all functions with domain X and range $\mathbb{F}$.

EX

$$\begin{aligned} \text{Let} \quad [\mathsf{f} + \mathsf{g}](x) &\triangleq \mathsf{f}(x) + \mathsf{g}(x) && \forall \mathsf{f}, \mathsf{g} \in \mathbb{F}^{\boldsymbol{X}} \quad (\textit{pointwise addition}) \quad \text{and} \\ [\alpha \cdot \mathsf{f}](x) &\triangleq \alpha \cdot [\mathsf{f}(x)] && \forall \mathsf{f} \in \mathbb{F}^{\boldsymbol{X}}, \alpha \in \mathbb{F}. \end{aligned}$$

Then the structure $\big(\mathbb{F}^{\boldsymbol{X}}, +, \cdot, (\mathbb{F}, +, \cdot)\big)$ is a *linear space*.

Theorem G.1 (Additive identity properties). [9] *Let* $\big(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times})\big)$ *be a linear space,* 0 *the* ADDITIVE IDENTITY ELEMENT *(Definition F.1 page 205) with respect to* $\dot{+}$*, and* $\mathbb{0}$ *the* ADDITIVE IDENTITY ELEMENT *with respect to* $+$.

THM

1. $0\boldsymbol{x} = \mathbb{0} \qquad \forall \boldsymbol{x} \in X$
2. $\alpha\mathbb{0} = \mathbb{0} \qquad \forall \alpha \in \mathbb{F}$
3. $\alpha\boldsymbol{x} = \mathbb{0} \implies \alpha = 0 \textit{ or } \boldsymbol{x} = \mathbb{0}$
4. $\boldsymbol{x} + \boldsymbol{x} = \boldsymbol{x} \implies \boldsymbol{x} = \mathbb{0}$
5. $\alpha \neq 0 \textit{ and } \boldsymbol{x} \neq \mathbb{0} \implies \alpha\boldsymbol{x} \neq \mathbb{0}$

✎PROOF:

[4] *Operator overload* is a technique in which two fundamentally different operators or functions share the same symbol or label. It is inherent in the programming language *C++* and is therein called *operator overload*. In C++, you can define two (or more) operators or functions that share the same symbol or name, but yet are completely different. Two such operators (or functions) are distinguished from each other by the type of their operands. So for example, in C++, you can define an $m \times n$ matrix *type* and use operator overload to define a + operator that operates on this new matrix type. So if variables x and y are of floating point type and A and B are of the matrix type, you can then add either type using the same syntax style:

x+y	(add two floating point numbers)
A+B	(add two matrices)

Even though both of these operations "look" the same, they are of course fundamentally different.

[5] Kubrusly (2001) page 41 ⟨Example 2D⟩
[6] Kubrusly (2001) page 41 ⟨Example 2D⟩, Hamel (1905)
[7] Kubrusly (2001) page 42 ⟨Example 2F⟩
[8] Kubrusly (2001) page 41 ⟨Example 2E⟩
[9] Berberian (1961) page 6 ⟨Theorem 1⟩, Michel and Herget (1993) page 77

COPYRIGHT © 2019 DANIEL J. GREENHOE | CC BY-NC-ND

1. Proof that $0\boldsymbol{x} = \mathbb{0}$:

$$\begin{aligned}
0\boldsymbol{x} &= 0\boldsymbol{x} + 0\mathbb{0} && \text{by definition of } + \text{ additive identity element}\\
&= 0\boldsymbol{x} + 0\boldsymbol{x} + (-0\boldsymbol{x}) && \text{by definition of } + \text{ additive inverse}\\
&= (0+0)\boldsymbol{x} + (-0\cdot\boldsymbol{x}) && \text{by definition of } + \text{ additive identity element}\\
&= 0\boldsymbol{x} + (-0\boldsymbol{x}) && \text{by Definition G.1 property 4}\\
&= \mathbb{0} && \text{by definition of } + \text{ additive identity element}
\end{aligned}$$

2. Proof that $\alpha\mathbb{0} = \mathbb{0}$:

$$\begin{aligned}
\alpha\mathbb{0} &= \alpha(0\boldsymbol{x}) && \text{by item 1}\\
&= (\alpha 0)\boldsymbol{x} && \text{by Definition G.1 property 6}\\
&= 0\boldsymbol{x}\\
&= \mathbb{0} && \text{by item 1}
\end{aligned}$$

3. Proof that $\alpha \neq 0$ and $\boldsymbol{x} \neq \mathbb{0} \implies \alpha\boldsymbol{x} \neq \mathbb{0}$: Suppose $\alpha\boldsymbol{x} = \mathbb{0}$. Then

$$\begin{aligned}
\boldsymbol{x} &= \left(\frac{1}{\alpha}\alpha\right)\boldsymbol{x}\\
&= \frac{1}{\alpha}(\alpha\boldsymbol{x})\\
&= \frac{1}{\alpha}\mathbb{0}\\
&= \mathbb{0} && \text{by item 2}\\
&\implies \boldsymbol{x} = \mathbb{0}
\end{aligned}$$

This is a *contradiction* and so $\alpha\boldsymbol{x} \neq \mathbb{0}$.

4. Proof that $\alpha\boldsymbol{x} = \mathbb{0} \implies \alpha = 0$ or $\boldsymbol{x} = \mathbb{0}$: contrapositive argument of item 3

5. Proof that $\boldsymbol{x} + \boldsymbol{x} = \boldsymbol{x} \implies \boldsymbol{x} = \mathbb{0}$:

$$\begin{aligned}
\boldsymbol{x} &= \boldsymbol{x} + \mathbb{0} && \text{by } \textit{additive identity} \text{ property (Definition G.1 page 207)}\\
&= \boldsymbol{x} + [\boldsymbol{x} + (-\boldsymbol{x})] && \text{by } \textit{additive inverse} \text{ property (Definition G.1 page 207)}\\
&= [\boldsymbol{x} + \boldsymbol{x}] + (-\boldsymbol{x}) && \text{by } \textit{associative} \text{ property (Definition G.1 page 207)}\\
&= \boldsymbol{x} + (-\boldsymbol{x}) && \text{by left hypothesis}\\
&= \mathbb{0} && \text{by } \textit{additive inverse} \text{ property (Definition G.1 page 207)}
\end{aligned}$$

✏

Definition G.3. [10] *Let $\boldsymbol{\Omega} \triangleq \big(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times})\big)$ be a linear space with vectors $\boldsymbol{x}, \boldsymbol{y} \in X$. Let $-\boldsymbol{y}$ be the additive inverse of $\boldsymbol{y}$ such that $\boldsymbol{y} + (-\boldsymbol{y}) = \mathbb{0}$.*

DEF *The **difference** of $\boldsymbol{x}$ and $\boldsymbol{y}$ is $\boldsymbol{x} + (-\boldsymbol{y})$ and is denoted $\boldsymbol{x} - \boldsymbol{y}$.*

Theorem G.2 (Additive inverse properties). [11] *Let $\boldsymbol{\Omega} \triangleq \big(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times})\big)$ be a linear space, $\mathbb{0}$ the* ADDITIVE IDENTITY ELEMENT *with respect to $+$, and $-\boldsymbol{x}$ the* ADDITIVE INVERSE *(Definition F.1 page 205) of $\boldsymbol{x}$ with respect to $+$.*

THM

1.	$\boldsymbol{x} + \boldsymbol{y}$	$= \mathbb{0}$	$\implies \boldsymbol{x} = -\boldsymbol{y}$	$\forall \boldsymbol{x},\boldsymbol{y} \in X$	*(additive inverse is* UNIQUE*)*
2.	$(-\alpha)\boldsymbol{x}$	$= -(\alpha\boldsymbol{x}) = \alpha(-\boldsymbol{x})$		$\forall \boldsymbol{x} \in X, \alpha \in \mathbb{F}$	
3.	$\alpha(\boldsymbol{x} - \boldsymbol{y})$	$= \alpha\boldsymbol{x} - \alpha\boldsymbol{y}$		$\forall \boldsymbol{x},\boldsymbol{y} \in X, \alpha \in \mathbb{F}$	*(*DISTRIBUTIVE*)*
4.	$(\alpha - \beta)\boldsymbol{x}$	$= \alpha\boldsymbol{x} - \beta\boldsymbol{x}$		$\forall \boldsymbol{x} \in X, \alpha,\beta \in \mathbb{F}$	*(*DISTRIBUTIVE*)*

[10] Berberian (1961) page 7 ⟨Definition 1⟩

[11] Berberian (1961) page 7 ⟨Corollary 1⟩, Michel and Herget (1993) page 77, Peano (1888a) ⟨Chapter IX⟩, Peano (1888b) pages 119–120, Banach (1922) pages 134–135

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

✎PROOF:

1. Proof that $\boldsymbol{x}+\boldsymbol{y}=0 \implies \boldsymbol{x}=-\boldsymbol{y}$:

$$\begin{aligned}\boldsymbol{x} &= \boldsymbol{x}-\mathbb{0}\\ &= \boldsymbol{x}-(\boldsymbol{x}+\boldsymbol{y}) && \text{by left hypothesis}\\ &= (\boldsymbol{x}-\boldsymbol{x})-\boldsymbol{y}\\ &= \mathbb{0}-\boldsymbol{y}\\ &= -\boldsymbol{y}\end{aligned}$$

2. Proof that $(-\alpha)\boldsymbol{x}=-(\alpha\boldsymbol{x})$:

$$\begin{aligned}\mathbb{0} &= 0\boldsymbol{x} && \text{by Theorem G.1 page 209}\\ &= (\alpha-\alpha)\boldsymbol{x} && \text{by field property of } \mathbb{F}\\ &= [\alpha+(-\alpha)]\boldsymbol{x} && \text{by field property of } \mathbb{F}\\ &= \alpha\boldsymbol{x}+(-\alpha)\boldsymbol{x} && \text{by Definition G.1 page 207}\\ \implies -(\alpha\boldsymbol{x}) &= (-\alpha)\boldsymbol{x} && \text{by item (1) page 211}\end{aligned}$$

3. Proof that $\alpha(-\boldsymbol{x})=-(\alpha\boldsymbol{x})$:

$$\begin{aligned}\mathbb{0} &= \alpha\mathbb{0} && \text{by Theorem G.1 page 209}\\ &= \alpha[\boldsymbol{x}+(-\boldsymbol{x})] && \text{by definition of additive identity element } -\boldsymbol{x}\\ &= \alpha\boldsymbol{x}+\alpha(-\boldsymbol{x}) && \text{by Definition G.1 page 207}\\ &= \alpha\boldsymbol{x}+\alpha(-\boldsymbol{x})\\ \implies -(\alpha\boldsymbol{x}) &= \alpha(-\boldsymbol{x}) && \text{by item (1) page 211}\end{aligned}$$

4. Proof that $\alpha(\boldsymbol{x}-\boldsymbol{y})=\alpha\boldsymbol{x}-\alpha\boldsymbol{y}$:

$$\begin{aligned}\alpha(\boldsymbol{x}-\boldsymbol{y}) &= \alpha\big[\boldsymbol{x}+(-\boldsymbol{y})\big] && \text{by Definition G.3 page 210}\\ &= \alpha\boldsymbol{x}+\alpha(-\boldsymbol{y}) && \text{by Definition G.1 page 207}\\ &= \alpha\boldsymbol{x}+(-\alpha\boldsymbol{y}) && \text{by item (3) page 211}\\ &= \alpha\boldsymbol{x}-\alpha\boldsymbol{y} && \text{by Definition G.3 page 210}\end{aligned}$$

5. Proof that $(\alpha-\beta)\boldsymbol{x}=\alpha\boldsymbol{x}-\beta\boldsymbol{x}$:

$$\begin{aligned}(\alpha-\beta)\boldsymbol{x} &= [\alpha+(-\beta)]\boldsymbol{x} && \text{by field properties of } \mathbb{F}\\ &= \alpha\boldsymbol{x}+(-\beta)\boldsymbol{x} && \text{by Definition G.1}\\ &= \alpha\boldsymbol{x}+[-(\beta\boldsymbol{x})] && \text{by item (2) page 211}\\ &= \alpha\boldsymbol{x}-(\beta\boldsymbol{x}) && \text{by Definition G.3 page 210}\end{aligned}$$

✏

Theorem G.3. [12] *Let $\big(X,+,\cdot,(\mathbb{F},\dot{+},\dot{\times})\big)$ be a linear space, $\mathbb{0}$ the additive identity element with respect to $+$, and $-\boldsymbol{x}$ additive inverse of $\boldsymbol{x}$ with respect to $+$.*

THM						
	1.	$\alpha\boldsymbol{x}=\alpha\boldsymbol{y}$ *and* $\alpha\neq 0$	$\implies$	$\boldsymbol{x}$	$=$	$\boldsymbol{y}$	$\forall \boldsymbol{x},\boldsymbol{y}\in X$
	2.	$\alpha\boldsymbol{x}=\beta\boldsymbol{x}$ *and* $\boldsymbol{x}\neq\mathbb{0}$	$\implies$	α	$=$	β	$\forall \boldsymbol{x},\boldsymbol{y}\in X,\ \alpha,\beta\in\mathbb{F}$
	3.	$\boldsymbol{z}+\boldsymbol{x}=\boldsymbol{z}+\boldsymbol{y}$	$\implies$	$\boldsymbol{x}$	$=$	$\boldsymbol{y}$	$\forall \boldsymbol{x},\boldsymbol{y},\boldsymbol{z}\in X$

[12] Michel and Herget (1993) page 77, Peano (1888a) ⟨Chapter IX⟩, Peano (1888b) pages 119–120, Banach (1922) pages 134–135

COPYRIGHT © 2019 DANIEL J. GREENHOE | CC BY-NC-ND

✎PROOF:

1. Proof that $\alpha\boldsymbol{x} = \alpha\boldsymbol{y}$ and $\alpha \neq 0 \implies \boldsymbol{x} = \boldsymbol{y}$:

$$\begin{aligned}
\mathbb{0} &= \frac{1}{\alpha}(\mathbb{0}) && \text{by left hypothesis } (\alpha \neq 0)\\
&= \frac{1}{\alpha}\Big(\alpha\boldsymbol{x} - \alpha\boldsymbol{y}\Big) && \text{by left hypothesis } (\alpha\boldsymbol{x} = \alpha\boldsymbol{y})\\
&= \frac{1}{\alpha}\alpha\Big(\boldsymbol{x} - \boldsymbol{y}\Big) && \text{by Definition G.1 page 207}\\
&= \boldsymbol{x} - \boldsymbol{y}
\end{aligned}$$

2. Proof that $\alpha\boldsymbol{x} = \beta\boldsymbol{x}$ and $\boldsymbol{x} \neq \mathbb{0} \implies \alpha = \beta$:

$$\begin{aligned}
\mathbb{0} &= \alpha\boldsymbol{x} + (-\alpha\boldsymbol{x}) && \text{by definition of additive inverse}\\
&= \beta\boldsymbol{x} + (-\alpha\boldsymbol{x}) && \text{by left hypothesis}\\
&= \beta\boldsymbol{x} + (-\alpha)\boldsymbol{x} && \text{by Theorem G.2 page 210}\\
&= [\beta + (-\alpha)]\boldsymbol{x} && \text{by Definition G.1 page 207}\\
\implies \beta - \alpha &= 0 && \text{by Theorem G.1 page 209}\\
\implies \alpha &= \beta && \text{by field properties of } \mathbb{F}
\end{aligned}$$

3. Proof that $\boldsymbol{z} + \boldsymbol{x} = \boldsymbol{z} + \boldsymbol{y} \implies \boldsymbol{x} = \boldsymbol{y}$:

$$\begin{aligned}
\mathbb{0} &= (\boldsymbol{z} + \boldsymbol{x}) - (\boldsymbol{z} + \boldsymbol{y})\\
&= (\boldsymbol{x} + \boldsymbol{z}) - (\boldsymbol{z} + \boldsymbol{y}) && \text{by Definition G.1 property 1}\\
&= (\boldsymbol{x} + \boldsymbol{z}) + [(-1)\boldsymbol{z} + (-1)\boldsymbol{y}] && \text{by Definition G.1 property 3}\\
&= (\boldsymbol{x} + \boldsymbol{z}) + (-\boldsymbol{z} - \boldsymbol{y}) && \text{by previous result 2.}\\
&= \boldsymbol{x} + (\boldsymbol{z} - \boldsymbol{z}) - \boldsymbol{y}\\
&= \boldsymbol{x} - \boldsymbol{y}
\end{aligned}$$

✏

G.2 Order on Linear Spaces

Definition G.4. [13] *Let $\boldsymbol{\Omega} \triangleq \big(X, +, \cdot, (\mathbb{R}, \dot{+}, \dot{\times})\big)$ be a real linear space.*

DEF

*The pair $(\boldsymbol{\Omega}, \leq)$ is an **ordered linear space** if*

1. $\boldsymbol{x} \leq \boldsymbol{y} \implies \boldsymbol{x} + \boldsymbol{z} \leq \boldsymbol{y} + \boldsymbol{z} \quad \forall z{\in}X$ *and*
2. $\boldsymbol{x} \leq \boldsymbol{y} \implies \alpha\boldsymbol{x} \leq \alpha\boldsymbol{y} \quad \forall \alpha{\in}\mathbb{F}$

*A vector $\boldsymbol{x}$ is **positive** if $\mathbb{0} \leq \boldsymbol{x}$.*

*The **positive cone** $\boldsymbol{X}^+$ of $(\boldsymbol{X}, \leq)$ is the set $\boldsymbol{X}^+ \triangleq \{\boldsymbol{x} \in \boldsymbol{X} | \mathbb{0} \leq \boldsymbol{x}\}$.*

Definition G.5. [14] *Let $(\boldsymbol{X}, \leq)$ be an ordered linear space.*

DEF

*The tuple $\boldsymbol{L} \triangleq (\boldsymbol{X}, \vee, \wedge; \leq)$ is a **Riesz space** if $\boldsymbol{L}$ is a **lattice**.*

A RIESZ SPACE *is also called a **vector lattice**.*

Theorem G.4. [15] *Let $(\boldsymbol{X}, \vee, \wedge; \leq)$ be a Riesz space (Definition G.5 page 212).*

THM

$\boldsymbol{x} \vee \boldsymbol{y}$	$=$	$-\big[(-\boldsymbol{x}) \wedge (-\boldsymbol{y})\big]$	$\boldsymbol{x} \wedge \boldsymbol{y}$	$=$	$-\big[(-\boldsymbol{x}) \vee (-\boldsymbol{y})\big]$	$\forall \boldsymbol{x},\boldsymbol{y}{\in}\boldsymbol{X}$
$\boldsymbol{x} + (\boldsymbol{y} \vee \boldsymbol{z})$	$=$	$(\boldsymbol{x} + \boldsymbol{y}) \vee (\boldsymbol{x} + \boldsymbol{z})$	$\boldsymbol{x} + (\boldsymbol{y} \wedge \boldsymbol{z})$	$=$	$(\boldsymbol{x} + \boldsymbol{y}) \wedge (\boldsymbol{x} + \boldsymbol{z})$	$\forall \boldsymbol{x},\boldsymbol{y},\boldsymbol{z}{\in}\boldsymbol{X}$
$\alpha(\boldsymbol{x} \vee \boldsymbol{y})$	$=$	$(\alpha\boldsymbol{x}) \vee (\alpha\boldsymbol{y})$	$\alpha(\boldsymbol{x} \wedge \boldsymbol{y})$	$=$	$(\alpha\boldsymbol{x}) \wedge (\alpha\boldsymbol{y})$	$\forall \boldsymbol{x},\boldsymbol{y}{\in}\boldsymbol{X}, \alpha{\geq}0$
$\boldsymbol{x} + \boldsymbol{y}$	$=$	$(\boldsymbol{x} \wedge \boldsymbol{y}) + (\boldsymbol{x} \vee \boldsymbol{y})$				$\forall \boldsymbol{x},\boldsymbol{y}{\in}\boldsymbol{X}, \alpha{\in}\mathbb{F}$

[13] Aliprantis and Burkinshaw (2006) pages 1–2
[14] Aliprantis and Burkinshaw (2006) page 2
[15] Aliprantis and Burkinshaw (2006) page 3 ⟨Theorem 1.2⟩

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

✎PROOF:

1. Proof that $x \vee y = -\big[(-x) \wedge (-y)\big]$:

$\begin{aligned} (-x) \wedge (-y) &\leq -x \\ x &\leq -\big[(-x) \wedge (-y)\big] \\ x \vee y &\leq -\big[(-x) \wedge (-y)\big] \end{aligned}$	$\begin{aligned} (-x) \wedge (-y) &\leq -y \\ y &\leq -\big[(-x) \wedge (-y)\big] \end{aligned}$
$\begin{aligned} x &\leq x \vee y \\ -(x \vee y) &\leq -x \\ -(x \vee y) &\leq (-x) \wedge (-y) \\ -\big[(-x) \wedge (-y)\big] &\leq x \vee y \end{aligned}$	$\begin{aligned} y &\leq x \vee y \\ -(x \vee y) &\leq -y \end{aligned}$

2. Proof that $x \wedge y = -\big[(-x) \vee (-y)\big]$:

$x \vee y = -\big[(-x) \wedge (-y)\big]$	by item (1)
$(-x) \vee (-y) = -\big[(-(-x)) \wedge (-(-y))\big]$	replace x with $-x$ and y with y
$(-x) \vee (-y) = -\big[x \wedge y\big]$	$-(-x) = x$
$-\big[x \wedge y\big] = (-x) \vee (-y)$	by symmetry of $=$ relation
$x \wedge y = -\big[(-x) \vee (-y)\big]$	multipy both sides by -1

3. Proof that $x + (y \vee z) = (x + y) \vee (x + z)$:

$\begin{aligned} x + y &\leq x + (y \vee z) \\ (x + y) \vee (x + z) &\leq x + (y \vee z) \end{aligned}$	$x + z \leq x + (y \vee z)$
$\begin{aligned} y &= -x + (x + y) \\ &\leq -x + \big[(x + y) \vee (x + z)\big] \\ y \vee z &\leq -x + \big[(x + y) \vee (x + z)\big] \\ x + (y \vee z) &\leq (x + y) \vee (x + z) \end{aligned}$	$\begin{aligned} z &= -x + (x + z) \\ &\leq -x + \big[(x + y) \vee (x + z)\big] \end{aligned}$

4. Proof that $x + (y \wedge z) = (x + y) \wedge (x + z)$:

$\begin{aligned} x + y &\geq x + (y \wedge z) \\ (x + y) \wedge (x + z) &\geq x + (y \wedge z) \end{aligned}$	$x + z \geq x + (y \wedge z)$
$\begin{aligned} y &= -x + (x + y) \\ &\geq -x + \big[(x + y) \wedge (x + z)\big] \\ y \wedge z &\geq -x + \big[(x + y) \wedge (x + z)\big] \\ x + (y \wedge z) &\geq (x + y) \wedge (x + z) \end{aligned}$	$\begin{aligned} z &= -x + (x + z) \\ &\geq -x + \big[(x + y) \wedge (x + z)\big] \end{aligned}$

5. Proof that $\alpha(x \vee y) = (\alpha x) \vee (\alpha y)$ for $\alpha \geq 0$:

$\begin{aligned} x &\leq x \vee y \\ \alpha x &\leq \alpha(x \vee y) \\ (\alpha x) \vee (\alpha y) &\leq \alpha(x \vee y) \end{aligned}$	$\begin{aligned} y &\leq x \vee y \\ \alpha y &\leq \alpha(x \vee y) \end{aligned}$	by Definition G.4 page 212
$\begin{aligned} \alpha x &\leq (\alpha x) \vee (\alpha y) \\ x &\leq \alpha^{-1}(\alpha x) \vee (\alpha y) \\ x \vee y &\leq \alpha^{-1}(\alpha x) \vee (\alpha y) \\ \alpha(x \vee y) &\leq (\alpha x) \vee (\alpha y) \end{aligned}$	$\begin{aligned} \alpha y &\leq (\alpha x) \vee (\alpha y) \\ y &\leq \alpha^{-1}(\alpha x) \vee (\alpha y) \end{aligned}$	

6. Proof that $\alpha(x \wedge y) = (\alpha x) \wedge (\alpha y)$ for $\alpha \geq 0$:

$\begin{aligned} x &\geq x \wedge y \\ \alpha x &\geq \alpha(x \wedge y) \\ (\alpha x) \wedge (\alpha y) &\geq \alpha(x \wedge y) \end{aligned}$	$\begin{aligned} y &\geq x \wedge y \\ \alpha y &\geq \alpha(x \wedge y) \end{aligned}$	by Definition G.4 page 212

COPYRIGHT © 2019 DANIEL J. GREENHOE
BY-NC-ND

αx	$\geq$	$(\alpha x) \wedge (\alpha y)$	αy	$\geq$	$(\alpha x) \wedge (\alpha y)$	
x	$\geq$	$\alpha^{-1}(\alpha x) \wedge (\alpha y)$	y	$\geq$	$\alpha^{-1}(\alpha x) \wedge (\alpha y)$	
$x \wedge y$	$\geq$	$\alpha^{-1}(\alpha x) \wedge (\alpha y)$				
$\alpha(x \wedge y)$	$\geq$	$(\alpha x) \wedge (\alpha y)$				

7. Proof that $x + y = (x \wedge y) + (x \vee y)$:

x	$\leq$	$x \vee y$	y	$\leq$	$x \vee y$
$x + y$	$\leq$	$(x \vee y) + y$	$x + vy$	$\leq$	$x + (x \vee y)$
$x + y - (x \vee y)$	$\leq$	y	$x + vy - (x \vee y)$	$\leq$	x
$x + y - (x \vee y)$	$\leq$	$x \wedge y$			
$x + y$	$\leq$	$(x \vee y) + (x \wedge y)$			
$x \wedge y$	$\leq$	x	$x \wedge y$	$\leq$	y
0	$\leq$	$x - (x \wedge y)$	0	$\leq$	$y - (x \wedge y)$
y	$\leq$	$y + x - (x \wedge y)$	x	$\leq$	$x + y - (x \wedge y)$
y	$\leq$	$x + y - (x \wedge y)$	x	$\leq$	$x + y - (x \wedge y)$
$x \vee y$	$\leq$	$x + y - (x \wedge y)$			
$(x \wedge y) + (x \vee y)$	$\leq$	$x + y$			

✏

Definition G.6. [16] *Let* $(\boldsymbol{X}, \vee, \wedge; \leq)$ *be a* RIESZ SPACE *(Definition G.5 page 212)*.

DEF

x^+	*is defined as* $x^+ \triangleq x \vee \mathbb{0}$	*and is called the* ***positive part*** *of* x.
x^-	*is defined as* $x^- \triangleq (-x) \vee \mathbb{0}$	*and is called the* ***negative part*** *of* x.
$\lvert x \rvert$	*is defined as* $\lvert x \rvert \triangleq x \vee (-x)$	*and is called the* ***absolute value*** *of* x.

Theorem G.5. [17] *Let* $(\boldsymbol{X}, \vee, \wedge; \leq)$ *be a* RIESZ SPACE *(Definition G.5 page 212)*.

THM

$$\left.\begin{array}{rcl} y - z &=& x \quad \textit{and} \\ y \wedge z &=& \mathbb{0} \end{array}\right\} \iff \left\{\begin{array}{rcl} y &=& x^+ \quad \textit{and} \\ z &=& x^- \end{array}\right.$$

✎PROOF:

1. Proof that left hypothesis $\implies$ right hypothesis:

$$\begin{aligned}
x^+ &= x \vee \mathbb{0} && \text{by definition of } x^+ \text{ Definition G.6 page 214} \\
&= (y - z) \vee \mathbb{0} && \text{by left hypothesis} \\
&= (y - z) \vee (z - z) \\
&= (y \vee z) - z && \text{by Theorem G.4 page 212} \\
&= \left[y + z - (y \wedge z)\right] - z && \text{by Theorem G.4 page 212} \\
&= y - (y \wedge z) \\
&= y - \mathbb{0} && \text{by left hypothesis} \\
&= y \\
x^- &= (-x) \vee \mathbb{0} && \text{by definition of } x^- \text{ Definition G.6 page 214} \\
&= (z - y) \vee \mathbb{0} && \text{by left hypothesis} \\
&= (z - y) \vee (y - y) \\
&= (z \vee y) - y && \text{by Theorem G.4 page 212}
\end{aligned}$$

[16] Aliprantis and Burkinshaw (2006) page 4, Istrăţescu (1987) page 129
[17] Aliprantis and Burkinshaw (2006) page 4 ⟨Theorem 1.3⟩

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

$$
\begin{aligned}
&= \left[z + y - (z \wedge y)\right] - z && \text{by Theorem G.4 page 212}\\
&= z - (z \wedge y) \\
&= z - \mathbb{0} && \text{by left hypothesis}\\
&= z
\end{aligned}
$$

2. Proof that left hypothesis $\Longleftarrow$ right hypothesis:

$$
\begin{aligned}
y - z &= x^+ - x^- && \text{by right hypothesis}\\
&= [x \vee \mathbb{0}] - [(-x) \vee \mathbb{0}] && \text{by Definition G.6 page 214}\\
&= (x \vee \mathbb{0}) + (x \wedge \mathbb{0}) && \text{by Theorem G.4 page 212}\\
&= x && \text{by Theorem G.4 page 212}\\
y \wedge z &= x^+ \wedge x^- && \text{by right hypothesis}\\
&= \left[x^- + \left(x^+ - x^-\right)\right] \wedge [x^- + \mathbb{0}] \\
&x^- + \left[\left(x^+ - x^-\right) \wedge \mathbb{0}\right] && \text{by Theorem G.4 page 212}\\
&x^- + \left[(y - z) \wedge \mathbb{0}\right] && \text{by right hypothesis}\\
&x^- + (x \wedge \mathbb{0}) && \text{by previous result}\\
&x^- - [-x \vee \mathbb{0}] && \text{by Theorem G.4 page 212}\\
&x^- - x^- && \text{by definition of } x^- \text{ (Definition G.6 page 214)}\\
&\mathbb{0}
\end{aligned}
$$

✏

Theorem G.6. [18] *Let* $(\boldsymbol{X}, \vee, \wedge; \leq)$ *be a* RIESZ SPACE *(Definition G.5 page 212). Let* x^+ *the* POSITIVE PART *of* $x \in \boldsymbol{X}$, x^- *the* NEGATIVE PART *of* $x \in \boldsymbol{X}$, *and* $|x|$ *the* ABSOLUTE VALUE *(Definition G.6 page 214) of* $x \in \boldsymbol{X}$.

THM

$$
\begin{aligned}
|x| &= x^+ + x^- && \forall x \in \boldsymbol{X}\\
x &= (x - y)^+ + (x \wedge y) && \forall x \in \boldsymbol{X}
\end{aligned}
$$

✎PROOF:

$$
\begin{aligned}
|x| &= x \vee (-x) && \text{by definition of } |x| \text{ (Definition G.6 page 214)}\\
&= (2x - x) \vee (\mathbb{0} - x) \\
&= (-x + 2x) \vee (-x + \mathbb{0}) && \text{by } \textit{commutative} \text{ property (Definition G.1 page 207)}\\
&= (-x) + (2x \vee \mathbb{0}) && \text{by Theorem G.4 page 212}\\
&= (2x \vee 2\mathbb{0}) - x && \text{by the } \textit{commutative} \text{ property (Definition G.1 page 207)}\\
&= 2(x \vee \mathbb{0}) - x && \text{by Theorem G.4 page 212}\\
&= 2x^+ - x && \text{by definition of } x^+ \text{ (Definition G.6 page 214)}\\
&= 2x^+ - \left(x^+ - x^-\right) && \text{by 1}\\
&= x^+ + x^- \\
x &= x + \mathbb{0} && x + y - y\\
&= (x \vee y) + (x \wedge y) - y && \text{by Theorem G.4 page 212}\\
&= \left[(x - y) \vee (y - y)\right] + (x \wedge y) && \text{by Theorem G.4 page 212}\\
&= \left[(x - y) \vee \mathbb{0}\right] + (x \wedge y) \\
&= (x - y)^+ + (x \wedge y) && \text{by definition of } x^+ \text{ (Definition G.6 page 214)}
\end{aligned}
$$

✏

[18] Aliprantis and Burkinshaw (2006) page 4

COPYRIGHT © 2019 DANIEL J. GREENHOE
CC BY-NC-ND

Theorem G.7. [19] *Let* $(\boldsymbol{X}, \vee, \wedge; \leq)$ *be a* RIESZ SPACE *(Definition G.5 page 212). Let* x^+ *the* POSITIVE PART *of* $x \in \boldsymbol{X}$, x^- *the* NEGATIVE PART *of* $x \in \boldsymbol{X}$, *and* $|x|$ *the* ABSOLUTE VALUE *(Definition G.6 page 214) of* $x \in \boldsymbol{X}$.

THM

1. $x \vee y = \frac{1}{2}(x + y + |x - y|) \quad \forall x,y \in \boldsymbol{X}$
2. $x \wedge y = \frac{1}{2}(x + y - |x - y|) \quad \forall x,y \in \boldsymbol{X}$
3. $|x - y| = (x \vee y) - (x \wedge y) \quad \forall x,y \in \boldsymbol{X}$
4. $|x| \vee |y| = \frac{1}{2}(|x + y| + |x - y|) \quad \forall x,y \in \boldsymbol{X}$
5. $|x| \vee |y| = \frac{1}{2}||x + y| - |x - y|| \quad \forall x,y \in \boldsymbol{X}$

✎PROOF:

$$
\begin{aligned}
(x + y + |x - y|) &= (x + y) + \big[(x - y) \vee (y - x)\big] && \text{by Definition G.6 page 214}\\
&= \big[(x + y) + (x - y)\big] \vee \big[(x + y) + (y - x)\big] && \text{by Theorem G.4 page 212}\\
&= (2x) \vee (2y)\\
&= 2(x \vee y) && \text{by Theorem G.4 page 212}\\
(x + y - |x - y|) &= (x + y) - \big[(x - y) \vee (y - x)\big] && \text{by Definition G.6 page 214}\\
&= (x + y) - \big[(-(y - x)) \vee (-(x - y))\big]\\
&= (x + y) + \big[(y - x) \wedge (x - y)\big] && \text{by Theorem G.4 page 212}\\
&= \big[(x + y) + (y - x)\big] \wedge \big[(x + y) + (x - y)\big] && \text{by Theorem G.4 page 212}\\
&= (2y) \wedge (2x)\\
&= 2(y \wedge x) && \text{by Theorem G.4 page 212}\\
&= 2(x \wedge y)\\
|x - y| &= \frac{1}{2}(x + y + |x - y|) - \frac{1}{2}(x + y + |x - y|)\\
&= (x \vee y) - (x \wedge y) && \text{by 1 and 2}\\
|x + y| + |x - y| &= \frac{1}{2}(\mathbb{0} + |2x + 2y|) + |x - y|\\
&= \frac{1}{2}\big[(x + y) + (-x - y) + |(x + y) - (-x - y)|\big] + |x - y|\\
&= \big[(x + y) \vee (-x - y)\big] + |x - y| && \text{by 1}\\
&= \big[(x + y) + |x - y|\big] \vee \big[(-x - y) + |x - y|\big] && \text{by Theorem G.4 page 212}\\
&= 2(x \vee y) \vee 2\big[(-y) + (-x) + |(-y) - (-x)|\big] && \text{by 1}\\
&= 2(x \vee y) \vee 2\big[(-y) \vee (-x)\big] && \text{by 1}\\
&= 2([x \vee (-x)] \vee (y \vee (-y)))\\
&= 2(|x| \vee |y|) && \text{by Definition G.6 page 214}\\
||x + y| - |x - y|| &= 2(|x + y| \vee |x - y|) - (|x + y| + |x - y|) && \text{by 1}\\
&= (|x + y + x - y| + |x + y - x + y|) - 2(|x| \vee |y|) && \text{by 3}\\
&= 2(|x| + |y|) - 2(|x| \vee |y|)\\
&= 2(|x| \vee |y|) && \text{by Theorem G.4 page 212}
\end{aligned}
$$

✏

Definition G.7. [20] *Let* $(\boldsymbol{X}, \vee, \wedge; \leq)$ *be a* RIESZ SPACE *(Definition G.5 page 212). Let* x^+ *the* POSITIVE PART *of* $x \in \boldsymbol{X}$, x^- *the* NEGATIVE PART *of* $x \in \boldsymbol{X}$, *and* $|x|$ *the* ABSOLUTE VALUE *(Definition G.6 page 214) of* $x \in \boldsymbol{X}$.

DEF

*x and y are **disjoint**, denoted by $x \perp y$, if*

$$|x| \wedge |y| = \mathbb{0}.$$

*Two subsets $\boldsymbol{U}$ and $\boldsymbol{V}$ of $\boldsymbol{X}$ are **disjoint**, denoted by $\boldsymbol{U} \perp \boldsymbol{V}$ if*

$$x \perp y \qquad \forall x \in \boldsymbol{U} \text{ and } y \in \boldsymbol{V}$$

[19] Aliprantis and Burkinshaw (2006) page 5 ⟨Theorem 1.4⟩
[20] Aliprantis and Burkinshaw (2006) page 5

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

Definition G.8. [21] *Let* $(\boldsymbol{X}, \vee, \wedge; \leq)$ *be a* RIESZ SPACE *(Definition G.5 page 212). Let* $\boldsymbol{x}^+$ *the* POSITIVE PART *of* $\boldsymbol{x} \in \boldsymbol{X}$, $\boldsymbol{x}^-$ *the* NEGATIVE PART *of* $\boldsymbol{x} \in \boldsymbol{X}$, *and* $|\boldsymbol{x}|$ *the* ABSOLUTE VALUE *(Definition G.6 page 214) of* $\boldsymbol{x} \in \boldsymbol{X}$. *Let* $\boldsymbol{Y}$ *be a subset of* $\boldsymbol{X}$.

DEF

$\boldsymbol{Y}^d$ *is the* ***disjoint complement*** *of* $\boldsymbol{Y}$ *if* $\quad \boldsymbol{Y}^d \triangleq \{\boldsymbol{x} \in \boldsymbol{X} | \boldsymbol{x} \perp \boldsymbol{y} \quad \forall \boldsymbol{y} \in \boldsymbol{Y}\}$.

The quantity $\boldsymbol{Y}^{dd}$ *is defined as* $\left(\boldsymbol{Y}^d\right)^d$.

Definition G.9. [22] *Let* $(\boldsymbol{X}, \vee, \wedge; \leq)$ *be a* RIESZ SPACE *(Definition G.5 page 212). Let* $\boldsymbol{x}^+$ *the* POSITIVE PART *of* $\boldsymbol{x} \in \boldsymbol{X}$, $\boldsymbol{x}^-$ *the* NEGATIVE PART *of* $\boldsymbol{x} \in \boldsymbol{X}$, *and* $|\boldsymbol{x}|$ *the* ABSOLUTE VALUE *(Definition G.6 page 214) of* $\boldsymbol{x} \in \boldsymbol{X}$.

DEF

$$
\begin{aligned}
|A| &\triangleq \{|\boldsymbol{a}| | \boldsymbol{a} \in A\} \\
A^+ &\triangleq \left\{\boldsymbol{a}^+ | \boldsymbol{a} \in A\right\} \\
A^- &\triangleq \{\boldsymbol{a}^- | \boldsymbol{a} \in A\} \\
A \vee B &\triangleq \left\{\boldsymbol{a} \vee \boldsymbol{b} | \boldsymbol{a} \in A \text{ and } \boldsymbol{b} \in B\right\} \\
A \wedge B &\triangleq \left\{\boldsymbol{a} \wedge \boldsymbol{b} | \boldsymbol{a} \in A \text{ and } \boldsymbol{b} \in B\right\} \\
\boldsymbol{x} \vee A &\triangleq \{\boldsymbol{x} \vee \boldsymbol{a} | \boldsymbol{a} \in A\} \\
\boldsymbol{x} \wedge A &\triangleq \{\boldsymbol{x} \wedge \boldsymbol{a} | \boldsymbol{a} \in A\}
\end{aligned}
$$

[21] Aliprantis and Burkinshaw (2006) page 5
[22] Aliprantis and Burkinshaw (2006) page 7

COPYRIGHT © 2019 DANIEL J. GREENHOE

Copyright © 2019 Daniel J. Greenhoe

APPENDIX H

LINEAR COMBINATIONS

H.1 Linear combinations in linear spaces

A *linear space* (Definition G.1 page 207) in general is not equipped with a *topology*. Without a topology, it is not possible to determine whether an *infinite sum* of vectors converges. Therefore in this section (dealing with linear spaces), all definitions related to sums of vectors will be valid for *finite* sums(Definition M.1 page 289) only (finite "N").

Definition H.1. [1] *Let* $\left\{ \boldsymbol{x}_n \in X \middle|_{n=1,2,\dots,N} \right\}$ *be a set of vectors in a* LINEAR SPACE $\left(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times})\right)$.

DEF

A vector $\boldsymbol{x} \in X$ *is a* ***linear combination*** *of the vectors in* $\left\{ \boldsymbol{x}_n \right\}$ *if*

there exists $\left\{ \alpha_n \in \mathbb{F} \middle|_{n=1,2,\dots,N} \right\}$ *such that* $$\boldsymbol{x} = \sum_{n=1}^{N} \alpha_n \boldsymbol{x}_n .$$

Definition H.2. [2] *Let* $\left(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times})\right)$ *be a linear space and* Y *be a subset of* X.

DEF

The ***linear span*** *of* Y *is defined as* $$\operatorname{span} Y \triangleq \left\{ \sum_{\gamma \in \Gamma} \alpha_\gamma \boldsymbol{y}_\gamma \middle| \alpha_\gamma \in \mathbb{F},\ \boldsymbol{y}_\gamma \in Y \right\}.$$

The set Y ***spans*** *a set* A *if* $$A \subseteq \operatorname{span} Y.$$

Proposition H.1. [3] *Let* $\left\{ \boldsymbol{x}_n \in X \middle|_{n=1,2,\dots,N} \right\}$ *be a set of vectors in a* LINEAR SPACE $\boldsymbol{L} \triangleq \left(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times})\right)$.

PRP

1. $\operatorname{span}\left\{ \boldsymbol{x}_n \right\}$ *is a* LINEAR SPACE *(Definition G.1 page 207) and*
2. $\operatorname{span}\left\{ \boldsymbol{x}_n \right\}$ *is a* LINEAR SUBSPACE *of* $\boldsymbol{L}$ *(Definition G.2 page 208).*

Definition H.3. [4] *Let* $\boldsymbol{L} \triangleq \left(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times})\right)$ *be a* LINEAR SPACE.

DEF

The set $Y \triangleq \left\{ \boldsymbol{x}_n \in X \middle|_{n=1,2,\dots,N} \right\}$ *is* ***linearly independent*** *in* $\boldsymbol{L}$ *if*

$$\left\{ \sum_{n=1}^{N} \alpha_n \boldsymbol{x}_n = 0 \right\} \implies \left\{ \alpha_1 = \alpha_2 = \cdots = \alpha_N = 0 \right\}.$$

The set Y *is* ***linearly dependent*** *in* $\boldsymbol{L}$ *if* Y *is not linearly independent in* $\boldsymbol{L}$.

[1] Berberian (1961) page 11 ⟨Definition I.4.1⟩, Kubrusly (2001) page 46

[2] Michel and Herget (1993) page 86 ⟨3.3.7 Definition⟩, Kurdila and Zabarankin (2005) page 44, Searcóid (2002) page 71 ⟨Definition 3.2.5—more general definition⟩

[3] Kubrusly (2001) page 46

[4] Bachman and Narici (1966) pages 3–4, Christensen (2003) page 2, Heil (2011) page 156 ⟨Definition 5.7⟩



Figure H.1: Lattice of *linear combinations*

BY-NC-ND Copyright © 2019 Daniel J. Greenhoe

Definition H.4. [5] *Let* $\{\boldsymbol{x}_n \in X \mid_{n=1,2,\ldots,N}\}$ *be a set of vectors in a* LINEAR SPACE $\boldsymbol{L} \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}))$.

DEF

The set $\{\boldsymbol{x}_n\}$ *is a* ***Hamel basis*** *for* $\boldsymbol{L}$ *if*

1. $\{\boldsymbol{x}_n\}$ SPANS $\boldsymbol{L}$ *(Definition H.2 page 219)* *and*
2. $\{\boldsymbol{x}_n\}$ *is* LINEARLY INDEPENDENT *in* $\boldsymbol{L}$ *(Definition H.1 page 219)* .

A HAMEL BASIS *is also called a* ***linear basis***.

Definition H.5. [6] *Let* $\boldsymbol{L} \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}))$ *be a* LINEAR SPACE. *Let* $\boldsymbol{x}$ *be a* VECTOR *in* $\boldsymbol{L}$ *and* $Y \triangleq \{\boldsymbol{x}_n \in X \mid_{n=1,2,\ldots,N}\}$ *be a set of vectors in* $\boldsymbol{L}$.

DEF

The expression $\sum_{n=1}^{N} \alpha_n \boldsymbol{x}_n$ *is the* ***expansion*** *of* $\boldsymbol{x}$ *on* Y *in* $\boldsymbol{L}$ *if* $\quad \boldsymbol{x} = \sum_{n=1}^{N} \alpha_n \boldsymbol{x}_n$.

In this case, the sequence $(\!(\alpha_n)\!)_{n=1}^{N}$ *is the* ***coordinates*** *of* $\boldsymbol{x}$ *with respect to* Y *in* $\boldsymbol{L}$.

If $\alpha_N \neq 0$, *then* N *is the* ***dimension*** $\dim \boldsymbol{L}$ *of* $\boldsymbol{L}$.

Theorem H.1. [7] *Let* $\{\boldsymbol{x}_n \mid_{n=1,2,\ldots,N}\}$ *be a* HAMEL BASIS *(Definition H.4 page 221) for a* LINEAR SPACE $(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}))$.

THM

$$\left\{\boldsymbol{x} = \sum_{n=1}^{N} \alpha_n \boldsymbol{x}_n = \sum_{n=1}^{N} \beta_n \boldsymbol{x}_n\right\} \quad \Longrightarrow \quad \underbrace{\alpha_n = \beta_n \quad \forall n=1,2,\ldots,N}_{\text{coordinates of } \boldsymbol{x} \text{ are UNIQUE}} \qquad \forall \boldsymbol{x} \in X$$

✎PROOF:

$$\begin{aligned}
\mathbb{0} &= \boldsymbol{x} - \boldsymbol{x} \\
&= \sum_{n=1}^{N} \alpha_n \boldsymbol{x}_n - \sum_{n=1}^{N} \beta_n \boldsymbol{x}_n \\
&= \sum_{n=1}^{N} (\alpha_n - \beta_n) \boldsymbol{x}_n \\
&\Longrightarrow \{\boldsymbol{x}_n\} \text{ is } \textit{linearly dependent} \text{ if } (\alpha_n - \beta_n) \neq 0 \qquad \forall n = 1, 2, \ldots, N \\
&\Longrightarrow (\alpha_n - \beta_n) = 0 \qquad \forall n = 1, 2, \ldots, N \qquad \text{(because } \{\boldsymbol{x}_n\} \text{ is a } \textit{basis} \text{ and therefore must be } \textit{linearly independent}) \\
&\Longrightarrow \alpha_n = \beta_n \text{ for } n = 1, 2, \ldots, N
\end{aligned}$$

✎

Theorem H.2. [8] *Let* $\boldsymbol{L} \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}))$ *be a* LINEAR SPACE.

THM

$$\left\{\begin{array}{ll} 1. & \{\boldsymbol{x}_n \in X|_{n=1,2,\ldots,N}\} \textit{ is a } \text{HAMEL BASIS} \textit{ for } \boldsymbol{L} \qquad \textit{and} \\ 2. & \{\boldsymbol{y}_n \in X|_{n=1,2,\ldots,M}\} \textit{ is a set of } \text{LINEARLY INDEPENDENT} \textit{ vectors in } \boldsymbol{L} \end{array}\right\}$$

$$\Longrightarrow \quad \left\{\begin{array}{ll} 1. & M \leq N \qquad \textit{and} \\ 2. & M = N \Longrightarrow \{\boldsymbol{y}_n|_{n=1,2,\ldots,M}\} \textit{ is a } \text{BASIS} \textit{ for } \boldsymbol{L} \qquad \textit{and} \\ 3. & M \neq N \Longrightarrow \{\boldsymbol{y}_n|_{n=1,2,\ldots,M}\} \textit{ is } \text{NOT} \textit{ a basis for } \boldsymbol{L} \end{array}\right\}$$

✎PROOF:

[5] Hamel (1905), Bachman and Narici (1966) page 4, Kubrusly (2001) pages 48–49 ⟨Section 2.4⟩, Young (2001) page 1, Carothers (2005) page 25, Heil (2011) page 125 ⟨Definition 4.1⟩

[6] Hamel (1905), Bachman and Narici (1966) page 4, Kubrusly (2001) pages 48–49 ⟨Section 2.4⟩, Young (2001) page 1, Carothers (2005) page 25, Heil (2011) page 125 ⟨Definition 4.1⟩

[7] Michel and Herget (1993) pages 89–90 ⟨Theorem 3.3.25⟩

[8] Michel and Herget (1993) pages 90–91 ⟨Theorem 3.3.26⟩

COPYRIGHT © 2019 DANIEL J. GREENHOE

1. Proof that $\{\boldsymbol{y}_1, \boldsymbol{x}_1, \ldots, \boldsymbol{x}_{N-1}\}$ is a *basis* for $\boldsymbol{L}$:
   (a) Proof that $\{\boldsymbol{y}_1, \boldsymbol{x}_1, \ldots, \boldsymbol{x}_{N-1}\}$ *spans* $\boldsymbol{L}$:
      i. Because $\{\boldsymbol{x}_n|_{n=1,2,\ldots,N}\}$ is a *basis* for $\boldsymbol{L}$, there exists $\beta \in \mathbb{F}$ and $\{\alpha_n \in \mathbb{F}|_{n=1,2,\ldots,N}\}$ such that
      $$\beta \boldsymbol{y}_1 + \sum_{n=1}^{N} \alpha_n \boldsymbol{x}_n = 0.$$
      ii. Select an n such that $\alpha_n \neq 0$ and renumber (if necessary) the above indices such that
      $$\boldsymbol{x}_n = -\frac{\beta}{\alpha_n}\boldsymbol{y}_1 + \sum_{n=1}^{N-1} \frac{\alpha_n}{\alpha_n}\boldsymbol{x}_n.$$
      iii. Then, for any $\boldsymbol{y} \in X$, we can write
      $$\begin{aligned}\boldsymbol{y} &= \sum_{n=1}^{N} \gamma_{n\in\mathbb{Z}}\boldsymbol{x}_n \\ &= \left(\sum_{n=1}^{N-1} \gamma_{n\in\mathbb{Z}}\boldsymbol{x}_n\right) + \gamma_{n\in\mathbb{Z}}\left(-\frac{\beta}{\alpha_n}\boldsymbol{y}_1 - \sum_{n=1}^{N-1}\frac{\alpha_n}{\alpha_n}\boldsymbol{x}_n\right) \\ &= -\frac{\beta\gamma_n}{\alpha_n}\boldsymbol{y}_1 + \sum_{n=1}^{N-1}\left(\gamma_n - \frac{\alpha_n\gamma_n}{\alpha_n}\right)\boldsymbol{x}_n \\ &= \delta\boldsymbol{y}_1 + \sum_{n=1}^{N-1}\delta_{n\in\mathbb{Z}}\boldsymbol{x}_n\end{aligned}$$
      iv. This implies that $\{\boldsymbol{y}_1, \boldsymbol{x}_1, \ldots, \boldsymbol{x}_{N-1}\}$ *spans* $\boldsymbol{L}$:
   (b) Proof that $\{\boldsymbol{y}_1, \boldsymbol{x}_1, \ldots, \boldsymbol{x}_{N-1}\}$ is *linearly independent*:
      i. If $\{\boldsymbol{y}_1, \boldsymbol{x}_1, \ldots, \boldsymbol{x}_{N-1}\}$ is *linearly dependent*, then there exists $\{\epsilon, \epsilon_1, \ldots, \epsilon_{N-1}\}$ such that
      $$\epsilon\boldsymbol{y}_1 + \left(\sum_{n=1}^{N-1}\epsilon_{n\in\mathbb{Z}}\boldsymbol{x}_n\right) + 0\boldsymbol{x}_n = 0.$$
      ii. item (1(b)i) implies that the coordinate of $\boldsymbol{y}_1$ associated with $\boldsymbol{x}_n$ *is* 0.
      $$\boldsymbol{y}_1 = -\left(\sum_{n=1}^{N-1}\frac{\epsilon_n}{\epsilon}\boldsymbol{x}_n\right) + 0\boldsymbol{x}_n = 0.$$
      iii. item (1(a)i) implies that the coordinate of $\boldsymbol{y}_1$ associated with $\boldsymbol{x}_n$ is *not* 0.
      $$\boldsymbol{y}_1 = -\sum_{n=1}^{N}\frac{\alpha_n}{\beta}\boldsymbol{x}_n.$$
      iv. This implies that item (1(b)i) (that the set is linearly dependent) is *false* because item (1(b)ii) and item (1(b)iii) *contradict* each other.
      v. This implies $\{\boldsymbol{y}_1, \boldsymbol{x}_1, \ldots, \boldsymbol{x}_{N-1}\}$ is *linearly independent*.
2. Proof that $\{\boldsymbol{y}_1, \boldsymbol{y}_2, \boldsymbol{x}_1, \ldots, \boldsymbol{x}_{N-2}\}$ is a *basis*: Repeat item (1).
3. Suppose $m = n$. Proof that $\{\boldsymbol{y}_1, \boldsymbol{y}_2, \ldots, \boldsymbol{y}_M\}$ is a *basis*: Repeat item (1) $M-1$ times.
4. Proof that $M \not> N$:
   (a) Suppose that $M = N+1$.
   (b) Then because $\{\boldsymbol{y}_n|_{n=1,2,\ldots,N}\}$ is a *basis*, there exists $\{\zeta_n|_{n=1,2,\ldots,N+1}\}$ such that
   $$\sum_{n=1}^{N+1}\zeta_{n\in\mathbb{Z}}\boldsymbol{y}_{n\in\mathbb{Z}} = 0.$$
   (c) This implies that $\{\boldsymbol{y}_n|_{n=1,2,\ldots,N+1}\}$ is *linearly dependent*.

BY-NC-ND Copyright © 2019 Daniel J. Greenhoe

(d) This implies that $\{\boldsymbol{y}_n|_{n=1,2,\ldots,N+1}\}$ is *not* a basis.

(e) This implies that $M \not> N$.

5. Proof that $M \neq N \implies \{\boldsymbol{y}_n|_{n=1,2,\ldots,M}\}$ is *not* a basis for $\boldsymbol{L}$:

   (a) Proof that $M > N \implies \{\boldsymbol{y}_n|_{n=1,2,\ldots,M}\}$ is *not* a basis for $\boldsymbol{L}$: same as in item (4).

   (b) Proof that $M < N \implies \{\boldsymbol{y}_n|_{n=1,2,\ldots,M}\}$ is *not* a basis for $\boldsymbol{L}$:

      i. Suppose $m = N - 1$.

      ii. Then $\{\boldsymbol{y}_n|_{n=1,2,\ldots,N-1}\}$ is a *basis* and there exists λ such that
      $$\sum_{n=1}^{N} \lambda_{n\in\mathbb{Z}} \boldsymbol{y}_{n\in\mathbb{Z}} = 0.$$

      iii. This implies that $\{\boldsymbol{y}_n|_{n=1,2,\ldots,N}\}$ is *linearly dependent* and is *not* a basis.

      iv. But this contradicts item (3), therefore $M \neq N - 1$.

      v. Because $M = N$ yields a basis but $M = N - 1$ does not, $M < N - 1$ also does not yield a basis.

✏

Corollary H.1. [9] *Let* $\boldsymbol{L} \triangleq \big(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times})\big)$ *be a linear space.*

COR
$$\underbrace{\left\{\begin{array}{ll} 1. & \{\boldsymbol{x}_n \in X|_{n=1,2,\ldots,N}\} \text{ is a HAMEL BASIS for } \boldsymbol{L} \quad \text{and} \\ 2. & \{\boldsymbol{y}_n \in X|_{n=1,2,\ldots,M}\} \text{ is a HAMEL BASIS for } \boldsymbol{L} \end{array}\right\} \implies \{N = M\}}_{\text{(all Hamel bases for } \boldsymbol{L} \text{ have the same number of vectors)}}$$

✎PROOF: This follows from Theorem H.2 (page 221). ✏

H.2 Bases in topological linear spaces

A linear space supports the concept of the *span* of a set of vectors (Definition H.2 page 219). In a topological linear space $\boldsymbol{\Omega} \triangleq \big(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \boldsymbol{T}\big)$, a set A is said to be *total* in $\boldsymbol{\Omega}$ if the span of A is *dense* in $\boldsymbol{\Omega}$. In this case, A is said to be a *total set* or a *complete set*. However, this use of "complete" in a "*complete set*" is not equivalent to the use of "complete" in a "*complete metric space*" .[10] In this text, except for these comments and Definition H.6, "complete" refers to the metric space definition only.

If a set is both *total* and *linearly independent* (Definition H.3 page 219) in $\boldsymbol{\Omega}$, then that set is a *Hamel basis* (Definition H.4 page 221) for $\boldsymbol{\Omega}$.

Definition H.6. [11] *Let* A^- *be the* CLOSURE *of a* A *in a* TOPOLOGICAL LINEAR SPACE $\boldsymbol{\Omega} \triangleq \big(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \boldsymbol{T}$ *Let* $\operatorname{span} A$ *be the* SPAN *(Definition H.2 page 219) of a set* A.

DEF
A set of vectors A *is* ***total*** *(or* ***complete*** *or* ***fundamental****) in* $\boldsymbol{\Omega}$ *if*
$$(\operatorname{span} A)^- = \boldsymbol{\Omega} \qquad \text{(SPAN of } A \text{ is DENSE in } \boldsymbol{\Omega}\text{)}.$$

[9] Kubrusly (2001) page 52 ⟨Theorem 2.7⟩, Michel and Herget (1993) page 91 ⟨Theorem 3.3.31⟩

[10] Haaser and Sullivan (1991) pages 296–297 ⟨6·Orthogonal Bases⟩, Rynne and Youngson (2008) page 78 ⟨Remark 3.50⟩, Heil (2011) page 21 ⟨Remark 1.26⟩

[11] Young (2001) page 19 ⟨Definition 1.5.1⟩, Sohrab (2003) page 362 ⟨Definition 9.2.3⟩, Gupta (1998) page 134 ⟨Definition 2.4⟩, Bachman and Narici (1966) pages 149–153 ⟨Definition 9.3, Theorems 9.9 and 9.10⟩

COPYRIGHT © 2019 DANIEL J. GREENHOE

H.3 Schauder bases in Banach spaces

Definition H.7. [12] *Let* $\boldsymbol{B} \triangleq \big(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \|\cdot\|\big)$ *be a* BANACH SPACE. *Let* $\stackrel{\star}{=}$ *represent* STRONG CONVERGENCE *in* $\boldsymbol{B}$.

DEF

The countable set $\big\{\boldsymbol{x}_n \in X \big|_{n\in\mathbb{N}}\big\}$ *is a* ***Schauder basis*** *for* $\boldsymbol{B}$ *if for each* $\boldsymbol{x} \in X$

1. $\exists\, (\!(\alpha_n \in \mathbb{F})\!)_{n\in\mathbb{N}}$ such that $\boldsymbol{x} \stackrel{\star}{=} \sum_{n=1}^{\infty} \alpha_n \boldsymbol{x}_n$ (STRONG CONVERGENCE *in* $\boldsymbol{B}$) *and*
2. $\left\{\sum_{n=1}^{\infty} \alpha_n \boldsymbol{x}_n \stackrel{\star}{=} \sum_{n=1}^{\infty} \beta_n \boldsymbol{x}_n\right\} \implies \big\{(\!(\alpha_n)\!) = (\!(\beta_n)\!)\big\}$ (COEFFICIENT FUNCTIONALS *are* UNIQUE)

In this case, $\sum_{n=1}^{\infty} \alpha_n \boldsymbol{x}_n$ *is the* ***expansion*** *of* $\boldsymbol{x}$ *on* $\big\{\boldsymbol{x}_n \big|_{n\in\mathbb{N}}\big\}$ *and*
the elements of $(\!(\alpha_n)\!)$ *are the* ***coefficient functionals*** *associated with the basis* $\big\{\boldsymbol{x}_n\big\}$.
Coefficient functionals are also called ***coordinate functionals***.

In a Banach space, the existence of a Schauder basis implies that the space is *separable* (Theorem H.3 page 224). The question of whether the converse is also true was posed by Banach himself in 1932,[13] and became know as "*The basis problem*". This remained an open question for many years. The question was finally answered some 41 years later in 1973 by Per Enflo (University of California at Berkley), with the answer being "no". Enflo constructed a counterexample in which a separable Banach space does *not* have a Schauder basis.[14] Life is simpler in Hilbert spaces where the converse *is* true: a Hilbert space has a Schauder basis *if and only if* it is separable (Theorem H.11 page 236).

Theorem H.3. [15] *Let* $\boldsymbol{B} \triangleq \big(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \|\cdot\|\big)$ *be a* BANACH SPACE. *Let* $\mathbb{Q}$ *be the field of rational numbers.*

THM

$$\left\{\begin{array}{ll} 1. & \boldsymbol{B} \text{ has a SCHAUDER BASIS} \quad \textit{and} \\ 2. & \mathbb{Q} \text{ is DENSE in } \mathbb{F}. \end{array}\right\} \implies \big\{\, \boldsymbol{B} \text{ is SEPARABLE} \,\big\}$$

✎PROOF:

1. lemma:

$$\left|\left\{\boldsymbol{x} \,|\, \exists\, (\!(\alpha_n \mathbb{Q})\!)_{n\in\mathbb{N}} \text{ such that } \lim_{N\to\infty} \left\|\boldsymbol{x} - \sum_{n=1}^{N} \alpha_n \boldsymbol{x}_n\right\| = 0\right\}\right| = |\,\mathbb{Q}\times\mathbb{N}\,|$$
$$= |\,\mathbb{Z}\times\mathbb{Z}\,|$$
$$= |\,\mathbb{Z}\,|$$
$$= \textit{countably infinite}$$

[12] Carothers (2005) pages 24–25, Christensen (2003) pages 46–49 ⟨Definition 3.1.1 and page 49⟩, Young (2001) page 19 ⟨Section 6⟩, Singer (1970), page 17, Schauder (1927), Schauder (1928)
[13] Banach (1932a), page 111
[14] Enflo (1973), Lindenstrauss and Tzafriri (1977) pages 84–95 ⟨Section 2.d⟩
[15] Bachman et al. (2000) page 112 ⟨3.4.8⟩, Giles (2000) page 17, Heil (2011) page 21 ⟨Theorem 1.27⟩

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

2. remainder of proof:

$\boldsymbol{B}$ has a *Schauder basis* $(\!(\boldsymbol{x}_n)\!)_{n\in\mathbb{N}}$

$$\Longrightarrow \text{ for every } \boldsymbol{x}\in\boldsymbol{B}\text{, there exists } (\!(\alpha_n\in\mathbb{F})\!)_{n\in\mathbb{N}} \text{ such that } \boldsymbol{x} \triangleq \sum_{n=1}^{\infty}\alpha_n\boldsymbol{x}_n \qquad \text{by Definition H.7 page 224}$$

$$\Longrightarrow \text{ for every } \boldsymbol{x}\in\boldsymbol{B}\text{, there exists } (\!(\alpha_n\in\mathbb{F})\!)_{n\in\mathbb{N}} \text{ such that } \lim_{N\to\infty}\left\|\boldsymbol{x}-\sum_{n=1}^{N}\alpha_n\boldsymbol{x}_n\right\| = 0$$

$$\Longrightarrow \text{ for every } \boldsymbol{x}\in\boldsymbol{B}\text{, there exists } (\!(\alpha_n\in\mathbb{Q})\!)_{n\in\mathbb{N}} \text{ such that } \lim_{N\to\infty}\left\|\boldsymbol{x}-\sum_{n=1}^{N}\alpha_n\boldsymbol{x}_n\right\| = 0 \quad \text{because } \mathbb{Q}^{-}=\mathbb{F}$$

$$\Longrightarrow \boldsymbol{B} = \left\{\boldsymbol{x}\,|\,\exists\,(\!(\alpha_n\mathbb{Q})\!)_{n\in\mathbb{N}} \text{ such that } \lim_{N\to\infty}\left\|\boldsymbol{x}-\sum_{n=1}^{N}\alpha_n\boldsymbol{x}_n\right\| = 0\right\}$$

$$\Longrightarrow \boldsymbol{B} = \left\{\boldsymbol{x}\,|\,\exists\,(\!(\alpha_n\mathbb{Q})\!)_{n\in\mathbb{N}} \text{ such that } \boldsymbol{x} = \lim_{N\to\infty}\sum_{n=1}^{N}\alpha_n\boldsymbol{x}_n\right\}^{-}$$

$$\Longrightarrow \boldsymbol{B} \text{ is } \textit{separable} \qquad \text{by (1) lemma page 224}$$

✏

Definition H.8. [16] *Let* $\{\boldsymbol{x}_n|_{n\in\mathbb{N}}\}$ *and* $\{\boldsymbol{y}_n|_{n\in\mathbb{N}}\}$ *be* SCHAUDER BASES *of a* BANACH SPACE $\big(X,+,\cdot,(\mathbb{F},\dot{+},\dot{\times}),\|\cdot\|\big)$.

DEF $\{\boldsymbol{x}_n\}$ *is* ***equivalent*** *to* $\{\boldsymbol{y}_n\}$
if there exists a BOUNDED INVERTIBLE *operator* $\mathbf{R}$ *in* $\boldsymbol{X}^{\boldsymbol{X}}$ *such that* $\quad \mathbf{R}\boldsymbol{x}_n = \boldsymbol{y}_n \quad \forall n\in\mathbb{Z}$

Theorem H.4. [17] *Let* $\{\boldsymbol{x}_n|_{n\in\mathbb{N}}\}$ *and* $\{\boldsymbol{y}_n|_{n\in\mathbb{N}}\}$ *be* SCHAUDER BASES *of a* BANACH SPACE $\big(X,+,\cdot,(\mathbb{F},\dot{+},\dot{\times}),\|\cdot\|\big)$.

THM $\{\{\boldsymbol{x}_n\}$ *is* EQUIVALENT *to* $\{\boldsymbol{y}_n\}\}$

$$\Longleftrightarrow \quad \left\{\sum_{n=1}^{\infty}\alpha_n\boldsymbol{x}_n \text{ is CONVERGENT} \Longleftrightarrow \sum_{n=1}^{\infty}\alpha_n\boldsymbol{y}_n \text{ is CONVERGENT}\right\}$$

Lemma H.1. [18] *Let* $\boldsymbol{\Omega} \triangleq \big(X,+,\cdot,(\mathbb{F},\dot{+},\dot{\times}),\boldsymbol{T}\big)$ *be a topological linear space. Let* $\operatorname{span}A$ *be the* SPAN *of a set* A *(Definition H.2 page 219). Let* $\tilde{\mathsf{f}}(\omega)$ *and* $\tilde{\mathsf{g}}(\omega)$ *be the* FOURIER TRANSFORMS *(Definition O.2 page 333) of the functions* $\mathsf{f}(x)$ *and* $\mathsf{g}(x)$*, respectively, in* $\boldsymbol{L}^2_{\mathbb{R}}$ *(Definition ?? page ??). Let* $\check{\mathsf{a}}(\omega)$ *be the* DTFT *(Definition P.1 page 343) of a sequence* $(\!(a_n)\!)_{n\in\mathbb{Z}}$ *in* $\boldsymbol{\ell}^2_{\mathbb{R}}$ *(Definition Q.2 page 353).*

LEM
$$\left\{\begin{array}{ll}(1). & \{\mathbf{T}^n\mathsf{f}|_{n\in\mathbb{Z}}\} \text{ is a SCHAUDER BASIS for } \boldsymbol{\Omega} \quad \text{and}\\ (2). & \{\mathbf{T}^n\mathsf{g}|_{n\in\mathbb{Z}}\} \text{ is a SCHAUDER BASIS for } \boldsymbol{\Omega}\end{array}\right\} \Longrightarrow \left\{\begin{array}{l}\exists\,(\!(a_n)\!)_{n\in\mathbb{Z}} \quad \text{such that}\\ \tilde{\mathsf{f}}(\omega) = \check{\mathsf{a}}(\omega)\tilde{\mathsf{g}}(\omega)\end{array}\right\}$$

✎PROOF: Let $\boldsymbol{V}_0'$ be the space spanned by $\{\mathbf{T}^n\phi|_{n\in\mathbb{Z}}\}$.

$$\begin{aligned}\tilde{\mathsf{f}}(\omega) &\triangleq \tilde{\mathbf{F}}\mathsf{f} && \text{by definition of } \tilde{\mathbf{F}} \qquad \text{(Definition O.2 page 333)}\\ &= \tilde{\mathbf{F}}\sum_{n\in\mathbb{Z}}a_n\mathbf{T}\mathsf{g} && \text{by (2)}\\ &= \sum_{n\in\mathbb{Z}}a_n\tilde{\mathbf{F}}\mathbf{T}\mathsf{g}\end{aligned}$$

[16] Young (2001) page 25 ⟨Definition 1.8.1, Theorem 1.8.7⟩
[17] Young (2001) page 25 ⟨Definition 1.8.1, Theorem 1.8.7⟩
[18] Daubechies (1992), page 140

COPYRIGHT © 2019 DANIEL J. GREENHOE

$$= \underbrace{\sum_{n\in\mathbb{Z}} a_n e^{-i\omega n}}_{\check{\mathsf{a}}(\omega)} \underbrace{\tilde{\mathbf{F}}\mathsf{g}}_{\tilde{\mathsf{g}}(\omega)} \qquad \text{by Corollary ?? page ??}$$

$$= \check{\mathsf{a}}(\omega)\tilde{\mathsf{g}}(\omega) \qquad \text{by definition of } \check{\mathbf{F}} \text{ and } \tilde{\mathbf{F}} \quad \text{by (Definition P.1 page 343, Definition O.2 page 333)}$$

$$\begin{aligned}
V_0 &\triangleq \left\{ \mathsf{f}(x) | \mathsf{f}(x) = \sum_{n\in\mathbb{Z}} b_n \mathbf{T}^n \mathsf{g}(x) \right\} \\
&= \left\{ \mathsf{f}(x) | \tilde{\mathbf{F}}\mathsf{f}(x) = \tilde{\mathbf{F}} \sum_{n\in\mathbb{Z}} b_n \mathbf{T}^n \mathsf{g}(x) \right\} \\
&= \left\{ \mathsf{f}(x) | \tilde{\mathsf{f}}(\omega) = \tilde{\mathsf{b}}(\omega)\tilde{\mathsf{g}}(\omega) \right\} \\
&= \left\{ \mathsf{f}(x) | \tilde{\mathsf{f}}(\omega) = \tilde{\mathsf{b}}(\omega)\check{\mathsf{a}}(\omega)\tilde{\mathsf{f}}(\omega) \right\} \\
&= \left\{ \mathsf{f}(x) | \tilde{\mathsf{f}}(\omega) = \tilde{\mathsf{c}}(\omega)\tilde{\mathsf{f}}(\omega) \right\} \qquad \text{where } \tilde{\mathsf{c}}(\omega) \triangleq \tilde{\mathsf{b}}(\omega)\check{\mathsf{a}}(\omega) \\
&= \left\{ \mathsf{f}(x) | \mathsf{f}(x) = \sum_{n\in\mathbb{Z}} \mathsf{c}_n \mathsf{f}(x-n) \right\} \\
&\triangleq V_0'
\end{aligned}$$

✏

H.4 Linear combinations in inner product spaces

In an *inner product space, orthogonality* is a special case of *linear independence*; or alternatively, linear independence is a generalization of orthogonality (next theorem).

Theorem H.5. [19] *Let* $\left\{ \boldsymbol{x}_n \in X \middle|_{n=1,2,\ldots,N} \right\}$ *be a set of vectors in an* INNER PRODUCT SPACE *(Definition J.1 page 255)* $\left(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle | \triangledown \rangle\right)$.

THM $\left\{ \begin{array}{c} \{\boldsymbol{x}_n\} \textit{ is } \text{ORTHOGONAL} \\ \textit{(Definition J.4 page 267)} \end{array} \right\} \implies \left\{ \begin{array}{c} \{\boldsymbol{x}_n\} \textit{ is } \text{LINEARLY INDEPENDENT} \\ \textit{(Definition H.1 page 219)} \end{array} \right\}$

✎PROOF:

1. Proof using *Pythagorean theorem* (Theorem J.10 page 268):
   Let $(\!(\alpha_n)\!)_{n\in\mathbb{N}}$ be a sequence with at least one nonzero element.

$$\begin{aligned}
\left\| \sum_{n=1}^{N} \alpha_n \boldsymbol{x}_n \right\|^2 &= \sum_{n=1}^{N} \left\| \alpha_n \boldsymbol{x}_n \right\|^2 && \text{by left hypoth. and } \textit{Pythagorean Theorem} \quad \text{(Theorem J.10 page 268)} \\
&= \sum_{n=1}^{N} \left| \alpha_n \right|^2 \left\| \boldsymbol{x}_n \right\|^2 && \text{by definition of } \|\cdot\| \quad \text{(Definition K.1 page 271)} \\
&> 0
\end{aligned}$$

$$\implies \sum_{n=1}^{N} \alpha_n \boldsymbol{x}_n \neq 0$$

$$\implies (\!(\boldsymbol{x}_n)\!)_{n\in\mathbb{N}} \text{ is } \textit{linearly independent} \qquad \text{by definition of } \textit{linear independence} \quad \text{(Definition H.3 page 219)}$$

[19] Aliprantis and Burkinshaw (1998) page 283 ⟨Corollary 32.8⟩, Kubrusly (2001) page 352 ⟨Proposition 5.34⟩

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

2. Alternative proof:

$$\begin{aligned}
\sum_{n=1}^{N} \alpha_n \boldsymbol{x}_n = \mathbb{0} &\Longrightarrow \left\langle \sum_{n=1}^{N} \alpha_n \boldsymbol{x}_n \,\middle|\, \boldsymbol{x}_m \right\rangle = \langle \mathbb{0} \mid \boldsymbol{x}_m \rangle \\
&\Longrightarrow \sum_{n=1}^{N} \alpha_n \langle \boldsymbol{x}_n \mid \boldsymbol{x}_m \rangle = 0 \\
&\Longrightarrow \sum_{n=1}^{N} \alpha_n \bar{\delta}(k-m) = 0 \\
&\Longrightarrow \alpha_m = 0 \qquad \text{for } m = 1, 2, \ldots, N
\end{aligned}$$

✏

Theorem H.6 (Bessel's Equality). [20] *Let* $\left\{ \boldsymbol{x}_n \in X \,\middle|\, {\scriptstyle n=1,2,\ldots,N} \right\}$ *be a set of vectors in an* INNER PRODUCT SPACE *(Definition J.1 page 255)* $\left(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle \mid \triangledown \rangle\right)$ *and with* $\|\boldsymbol{x}\| \triangleq \sqrt{\langle \boldsymbol{x} \mid \boldsymbol{x} \rangle}$ *(Definition J.2 page 260)*.

THM

$$\left\{ \begin{array}{c} \{\boldsymbol{x}_n\} \text{ is ORTHONORMAL} \\ \text{(Definition J.4 page 267)} \end{array} \right\} \Longrightarrow \left\{ \underbrace{\left\| \boldsymbol{x} - \sum_{n=1}^{N} \langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle \boldsymbol{x}_n \right\|^2}_{\text{approximation error}} = \|\boldsymbol{x}\|^2 - \sum_{n=1}^{N} |\langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle|^2 \quad \forall \boldsymbol{x} \in X \right\}$$

✎PROOF:

$$\left\| \boldsymbol{x} - \sum_{n=1}^{N} \langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle \boldsymbol{x}_n \right\|^2$$

$$= \|\boldsymbol{x}\|^2 + \left\| \sum_{n=1}^{N} \langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle \boldsymbol{x}_n \right\|^2 - 2\Re \left\langle \boldsymbol{x} \,\middle|\, \sum_{n=1}^{N} \langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle \boldsymbol{x}_n \right\rangle \qquad \text{by } \textit{polar identity} \qquad \text{(Lemma J.1 page 259)}$$

$$= \|\boldsymbol{x}\|^2 + \left\| \sum_{n=1}^{N} \langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle \boldsymbol{x}_n \right\|^2 - 2\Re \left[\left(\sum_{n=1}^{N} \langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle \right)^* \langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle \right] \qquad \text{by property of } \langle \triangle \mid \triangledown \rangle \qquad \text{(Definition J.1 page 255)}$$

$$= \|\boldsymbol{x}\|^2 + \sum_{n=1}^{N} \left\| \langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle \boldsymbol{x}_n \right\|^2 - 2\Re \left[\left(\sum_{n=1}^{N} \langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle \right)^* \langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle \right] \qquad \text{by } \textit{Pythagorean Theorem} \qquad \text{(Theorem J.10 page 268)}$$

$$= \|\boldsymbol{x}\|^2 + \sum_{n=1}^{N} \left\| \langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle \boldsymbol{x}_n \right\|^2 - 2\Re \left(\sum_{n=1}^{N} \langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle^* \langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle \right)$$

$$= \|\boldsymbol{x}\|^2 + \sum_{n=1}^{N} \left| \langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle \right|^2 \underbrace{\|\boldsymbol{x}_n\|^2}_{1} - 2\Re \left(\sum_{n=1}^{N} \langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle^* \langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle \right) \qquad \text{by property of } \|\cdot\| \qquad \text{(Definition K.1 page 271)}$$

$$= \|\boldsymbol{x}\|^2 + \sum_{n=1}^{N} \left| \langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle \right|^2 \cdot 1 - 2\Re \left(\sum_{n=1}^{N} \langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle^* \langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle \right) \qquad \text{by def. of } \textit{orthonormality} \qquad \text{(Definition J.4 page 267)}$$

$$= \|\boldsymbol{x}\|^2 + \sum_{n=1}^{N} \left| \langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle \right|^2 - 2\Re \sum_{n=1}^{N} \left| \langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle \right|^2$$

$$= \|\boldsymbol{x}\|^2 + \sum_{n=1}^{N} \left| \langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle \right|^2 - 2 \sum_{n=1}^{N} \left| \langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle \right|^2 \qquad \text{because } |\cdot| \text{ is real}$$

[20] Bachman et al. (2000) page 103, Pedersen (2000) pages 38–39

COPYRIGHT © 2019 DANIEL J. GREENHOE

$$= \|\boldsymbol{x}\|^2 - \sum_{n=1}^{N} \left|\langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle\right|^2$$

✏

Theorem H.7 (Bessel's inequality). [21] *Let* $\left\{ \boldsymbol{x}_n \in X \middle|_{n=1,2,\ldots,N} \right\}$ *be a set of vectors in an* INNER PRODUCT SPACE *(Definition J.1 page 255)* $\left(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle \mid \triangledown \rangle\right)$ *and with* $\|\boldsymbol{x}\| \triangleq \sqrt{\langle \boldsymbol{x} \mid \boldsymbol{x} \rangle}$ *(Definition J.2 page 260)*.

THM

$$\left\{ \begin{array}{c} \{\boldsymbol{x}_n\} \text{ is ORTHONORMAL} \\ \text{(Definition J.4 page 267)} \end{array} \right\} \implies \left\{ \sum_{n=1}^{N} \left|\langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle\right|^2 \leq \|\boldsymbol{x}\|^2 \quad \forall \boldsymbol{x} \in X \right\}$$

✎PROOF:

$$0 \leq \left\| \boldsymbol{x} - \sum_{n=1}^{N} \langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle \, \boldsymbol{x}_n \right\|^2 \qquad \text{by definition of } \|\cdot\| \qquad \text{(Definition K.1 page 271)}$$

$$= \|\boldsymbol{x}\|^2 - \sum_{n=1}^{N} \left|\langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle\right|^2 \qquad \text{by } \textit{Bessel's Equality} \qquad \text{(Theorem H.6 page 227)}$$

✏

The *Best Approximation Theorem* (next) shows that

- the best sequence for representing a vector is the sequence of projections of the vector onto the sequence of basis functions
- the error of the projection is orthogonal to the projection.

Theorem H.8 (Best Approximation Theorem). [22] *Let* $\left\{ \boldsymbol{x}_n \in X \middle|_{n=1,2,\ldots,N} \right\}$ *be a set of vectors in an* INNER PRODUCT SPACE *(Definition J.1 page 255)* $\left(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle \mid \triangledown \rangle\right)$ *and with* $\|\boldsymbol{x}\| \triangleq \sqrt{\langle \boldsymbol{x} \mid \boldsymbol{x} \rangle}$ *(Definition J.2 page 260)*.

THM

$$\left\{ \begin{array}{c} \{\boldsymbol{x}_n\} \text{ is} \\ \text{ORTHONORMAL} \\ \text{(Definition J.4 page 267)} \end{array} \right\} \implies \left\{ \begin{array}{lll} 1. & \arg\min\limits_{(\alpha_n)_{n=1}^N} \left\| \boldsymbol{x} - \sum\limits_{n=1}^{N} \alpha_n \boldsymbol{x}_n \right\| = \underbrace{(\!(\langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle)\!)_{n=1}^{N}}_{\text{best } \alpha_n = \langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle} & \forall \boldsymbol{x} \in X \quad \textit{and} \\ 2. & \underbrace{\left(\sum\limits_{n=1}^{N} \langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle \, \boldsymbol{x}_n \right)}_{\textit{approximation}} \perp \underbrace{\left(\boldsymbol{x} - \sum\limits_{n=1}^{N} \langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle \, \boldsymbol{x}_n \right)}_{\textit{approximation error}} & \forall \boldsymbol{x} \in X \end{array} \right\}$$

✎PROOF:

[21] Giles (2000) pages 54–55 ⟨3.13 Bessel's inequality⟩, Bollobás (1999) page 147, Aliprantis and Burkinshaw (1998) page 284

[22] Walter and Shen (2001), pages 3–4, Pedersen (2000), page 39, Edwards (1995), pages 94–100, Weyl (1940)

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

1. Proof that $(\!(\langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle)\!)$ is the best sequence:

$$
\begin{aligned}
&\left\| \boldsymbol{x} - \sum_{n=1}^{N} \alpha_n \boldsymbol{x}_n \right\|^2 \\
&= \|\boldsymbol{x}\|^2 - 2\Re\left\langle \boldsymbol{x} \mid \sum_{n=1}^{N} \alpha_n \boldsymbol{x}_n \right\rangle + \left\| \sum_{n=1}^{N} \alpha_n \boldsymbol{x}_n \right\|^2 \\
&= \|\boldsymbol{x}\|^2 - 2\Re\left(\sum_{n=1}^{N} \alpha_n^* \langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle \right) + \sum_{n=1}^{N} \|\alpha_n \boldsymbol{x}_n\|^2 \qquad \text{by } \textit{Pythagorean Theorem} \qquad \text{(Theorem J.10 page 268)} \\
&= \|\boldsymbol{x}\|^2 - 2\Re\left(\sum_{n=1}^{N} \alpha_n^* \langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle \right) + \sum_{n=1}^{N} |\alpha_n|^2 + \underbrace{\left[\sum_{n=1}^{N} \left|\langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle\right|^2 - \sum_{n=1}^{N} \left|\langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle\right|^2 \right]}_{0} \\
&= \left[\|\boldsymbol{x}\|^2 - \sum_{n=1}^{N} |\langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle|^2 \right] + \sum_{n=1}^{N} \left[\left|\langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle\right|^2 - 2\mathbf{R}_{\mathrm{e}} \left[\alpha_n^* \langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle \right] + \left|\alpha_n\right|^2 \right] \\
&= \left[\|\boldsymbol{x}\|^2 - \sum_{n=1}^{N} |\langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle|^2 \right] + \sum_{n=1}^{N} \left[|\langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle|^2 - \alpha_n^* \langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle - \alpha_n \langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle^* + |\alpha_n|^2 \right] \\
&= \left\| \boldsymbol{x} - \sum_{n=1}^{N} \langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle \boldsymbol{x}_n \right\|^2 + \sum_{n=1}^{N} |\langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle - \alpha_n|^2 \qquad \text{by } \textit{Bessel's Equality} \qquad \text{(Theorem H.6 page 227)} \\
&\geq \left\| \boldsymbol{x} - \sum_{n=1}^{N} \langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle \boldsymbol{x}_n \right\|^2
\end{aligned}
$$

2. Proof that the approximation and approximation error are orthogonal:

$$
\begin{aligned}
\left\langle \sum_{n=1}^{N} \langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle \boldsymbol{x}_n \,\middle|\, \boldsymbol{x} - \sum_{n=1}^{N} \langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle \boldsymbol{x}_n \right\rangle &= \left\langle \sum_{n=1}^{N} \langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle \boldsymbol{x}_n \,\middle|\, \boldsymbol{x} \right\rangle - \left\langle \sum_{n=1}^{N} \langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle \boldsymbol{x}_n \,\middle|\, \sum_{n=1}^{N} \langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle \boldsymbol{x}_n \right\rangle \\
&= \sum_{n=1}^{N} \langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle^* \langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle - \sum_{n=1}^{N} \sum_{m=1}^{N} \langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle \langle \boldsymbol{x} \mid \boldsymbol{x}_m \rangle^* \langle \boldsymbol{x}_n \mid \boldsymbol{x}_m \rangle \\
&= \sum_{n=1}^{N} |\langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle|^2 - \sum_{n=1}^{N} \sum_{m=1}^{N} \langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle \langle \boldsymbol{x} \mid \boldsymbol{x}_m \rangle^* \bar{\delta}_{nm} \\
&= \sum_{n=1}^{N} |\langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle|^2 - \sum_{n=1}^{N} |\langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle|^2 \\
&= 0
\end{aligned}
$$

✍

H.5 Orthonormal bases in Hilbert spaces

Definition H.9. *Let* $\left\{ \boldsymbol{x}_n \in X \,\middle|\, {\scriptstyle n=1,2,\ldots,N} \right\}$ *be a set of vectors in an* INNER PRODUCT SPACE *(Definition J.1 page 255)* $\boldsymbol{\Omega} \triangleq \left(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle \mid \triangledown \rangle\right)$.

COPYRIGHT © 2019 DANIEL J. GREENHOE | https://github.com/dgreenhoe/pdfs/blob/master/abcstat.pdf | CC BY-NC-ND

DEF

The set $\{\boldsymbol{x}_n\}$ is an ***orthogonal basis*** *for $\boldsymbol{\Omega}$ if $\{\boldsymbol{x}_n\}$ is* ORTHOGONAL *and is a* SCHAUDER BASIS *for $\boldsymbol{\Omega}$.*
The set $\{\boldsymbol{x}_n\}$ is an ***orthonormal basis*** *for $\boldsymbol{\Omega}$ if $\{\boldsymbol{x}_n\}$ is* ORTHONORMAL *and is a* SCHAUDER BASIS *for $\boldsymbol{\Omega}$.*

Definition H.10. [23] *Let $\boldsymbol{H} \triangleq \big(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle\triangle\,|\,\triangledown\rangle\big)$ be a Hilbert space.*

DEF

Suppose there exists a set $\{\boldsymbol{x}_n \in X \,|\, {}_{n\in\mathbb{N}}\}$ such that $\boldsymbol{x} \stackrel{*}{=} \sum_{n=1}^{\infty} \langle \boldsymbol{x} \,|\, \boldsymbol{x}_n\rangle\, \boldsymbol{x}_n$.

Then the quantities $\langle \boldsymbol{x} \,|\, \boldsymbol{x}_n\rangle$ are called the ***Fourier coefficients*** *of $\boldsymbol{x}$ and the sum $\sum_{n=1}^{\infty} \langle \boldsymbol{x} \,|\, \boldsymbol{x}_n\rangle\, \boldsymbol{x}_n$ is called the* ***Fourier expansion*** *of $\boldsymbol{x}$ or the* ***Fourier series*** *for $\boldsymbol{x}$.*

Lemma H.2 (Perfect reconstruction)**.** *Let $\{\boldsymbol{x}_n \in X \,|\, {}_{n\in\mathbb{N}}\}$ be a set of vectors in a* HILBERT SPACE *$\boldsymbol{H} \triangleq \big(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle\triangle\,|\,\triangledown\rangle\big)$.*

LEM

$$\left\{\begin{array}{ll} (1). & (\boldsymbol{x}_n) \text{ is a BASIS for } \boldsymbol{H} \quad \text{and} \\ (2). & (\boldsymbol{x}_n) \text{ is ORTHONORMAL} \end{array}\right\} \implies \boldsymbol{x} \stackrel{*}{=} \overbrace{\sum_{n=1}^{\infty} \underbrace{\langle \boldsymbol{x} \,|\, \boldsymbol{x}_n\rangle}_{\text{Fourier coefficient}}\, \boldsymbol{x}_n}^{\text{Fourier expansion}} \quad \forall \boldsymbol{x}\in X$$

✎PROOF:

$$\begin{aligned}
\langle \boldsymbol{x} \,|\, \boldsymbol{x}_n\rangle &= \left\langle \sum_{m\in\mathbb{Z}} \alpha_m \boldsymbol{x}_m \,|\, \boldsymbol{x}_n \right\rangle && \text{by left hypothesis (1)} \\
&= \sum_{m\in\mathbb{Z}} \alpha_m \langle \boldsymbol{x}_m \,|\, \boldsymbol{x}_n\rangle && \text{by } \textit{homogeneous} \text{ property of } \langle\triangle\,|\,\triangledown\rangle && \text{(Definition J.1 page 255)} \\
&= \sum_{m\in\mathbb{Z}} \alpha_m \bar{\delta}_{n-m} && \text{by left hypothesis (2)} && \text{(Definition J.4 page 267)} \\
&= \alpha_n
\end{aligned}$$

✏

Proposition H.2. [24] *Let $\{\boldsymbol{x}_n \in X \,|\, {}_{n\in\mathbb{N}}\}$ be a set of vectors in a* HILBERT SPACE *$\boldsymbol{H} \triangleq \big(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle\triangle\,|\,\triangledown\rangle\big)$.*

PRP

$$\underbrace{\|\boldsymbol{x}\|^2 \stackrel{*}{=} \sum_{n=1}^{\infty} \big|\langle \boldsymbol{x} \,|\, \boldsymbol{x}_n\rangle\big|^2}_{\text{PARSEVAL FRAME}} \iff \underbrace{\boldsymbol{x} \stackrel{*}{=} \sum_{n=1}^{\infty} \langle \boldsymbol{x} \,|\, \boldsymbol{x}_n\rangle\, \boldsymbol{x}_n}_{\text{FOURIER EXPANSION (Definition H.10 page 230)}} \quad \forall \boldsymbol{x}\in X$$

✎PROOF:

[23] Fabian et al. (2010) page 27 ⟨Theorem 1.55⟩, Young (2001) page 6, Young (1980) page 6
[24] Han et al. (2007) pages 93–94 ⟨Proposition 3.11⟩

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

1. Proof that *Parseval frame* $\Longleftarrow$ *Fourier expansion*

$$\begin{aligned}
\|\boldsymbol{x}\|^2 &\triangleq \langle \boldsymbol{x} \mid \boldsymbol{x}\rangle && \text{by definition of } \|\cdot\| && \text{(Definition K.1 page 271)}\\
&= \left\langle \sum_{n=1}^{\infty} \langle \boldsymbol{x} \mid \boldsymbol{x}_n\rangle\, \boldsymbol{x} \mid \boldsymbol{x}_n \right\rangle && \text{by right hypothesis} && \\
&\overset{\star}{=} \sum_{n=1}^{\infty} \langle \boldsymbol{x} \mid \boldsymbol{x}_n\rangle \langle \boldsymbol{x} \mid \boldsymbol{x}_n\rangle && \text{by property of } \langle \triangle \mid \triangledown\rangle && \text{(Definition J.1 page 255)}\\
&\overset{\star}{=} \sum_{n=1}^{\infty} \langle \boldsymbol{x} \mid \boldsymbol{x}_n\rangle \langle \boldsymbol{x} \mid \boldsymbol{x}_n\rangle^* && \text{by property of } \langle \triangle \mid \triangledown\rangle && \text{(Definition J.1 page 255)}\\
&\overset{\star}{=} \sum_{n=1}^{\infty} \left|\langle \boldsymbol{x} \mid \boldsymbol{x}_n\rangle\right|^2 && \text{by property of } \mathbb{C} && \text{(Definition I.7 page 253)}
\end{aligned}$$

2. Proof that *Parseval frame* $\Longrightarrow$ *Fourier expansion*

   (a) Let $(\!(\boldsymbol{e}_n)\!)_{n\in\mathbb{N}}$ be the *standard othornormal basis* such that the nth element of $\boldsymbol{e}_n$ is 1 and all other elements are 0.

   (b) Let $\mathbf{M}$ be an operator in $\boldsymbol{H}$ such that $\mathbf{M}\boldsymbol{x} \triangleq \sum_{n=1}^{\infty} \langle \boldsymbol{x} \mid \boldsymbol{x}_n\rangle\, \boldsymbol{e}_n$.

   (c) lemma: $\mathbf{M}$ is *isometric*. Proof:

$$\begin{aligned}
\|\mathbf{M}\boldsymbol{x}\|^2 &= \left\| \sum_{n=1}^{\infty} \langle \boldsymbol{x} \mid \boldsymbol{x}_n\rangle\, \boldsymbol{e}_n \right\|^2 && \text{by definition of } \mathbf{M} && \text{(item (2b) page 231)}\\
&= \sum_{n=1}^{\infty} \left\|\langle \boldsymbol{x} \mid \boldsymbol{x}_n\rangle\, \boldsymbol{e}_n\right\|^2 && \text{by } \textit{Pythagorean Theorem} && \text{(Theorem J.10 page 268)}\\
&= \sum_{n=1}^{\infty} \left|\langle \boldsymbol{x} \mid \boldsymbol{x}_n\rangle\right|^2 \left\|\boldsymbol{e}_n\right\|^2 && \text{by } \textit{homogeneous} \text{ property of } \|\cdot\| && \text{(Definition K.1 page 271)}\\
&= \sum_{n=1}^{\infty} \left|\langle \boldsymbol{x} \mid \boldsymbol{x}_n\rangle\right|^2 && \text{by definition of } \textit{orthonormal} && \text{(Definition J.4 page 267)}\\
&= \|\boldsymbol{x}\|^2 && \text{by Parseval frame hypothesis} && \\
&\Longrightarrow \quad \mathbf{M} \text{ is } \textit{isometric} && \text{by definition of } \textit{isometric} && \text{(Definition N.10 page 323)}
\end{aligned}$$

   (d) Let $(\!(\boldsymbol{u}_n)\!)_{n\in\mathbb{N}}$ be an *orthornormal basis* for $\boldsymbol{H}$.

   (e) Proof for *Fourier expansion*:

$$\begin{aligned}
\boldsymbol{x} &= \sum_{n=1}^{\infty} \langle \boldsymbol{x} \mid \boldsymbol{u}_n\rangle\, \boldsymbol{u}_n && \text{by } \textit{Fourier expansion} \text{ (Proposition H.3 page 234)}\\
&= \sum_{n=1}^{\infty} \langle \mathbf{M}\boldsymbol{x} \mid \mathbf{M}\boldsymbol{u}_n\rangle\, \boldsymbol{u}_n && \text{by (2c) lemma page 231 and Theorem N.23 page 324}\\
&= \sum_{n=1}^{\infty} \left\langle \sum_{m=1}^{\infty} \langle \boldsymbol{x} \mid \boldsymbol{x}_m\rangle\, \boldsymbol{e}_m \mid \sum_{k=1}^{\infty} \langle \boldsymbol{u}_n \mid \boldsymbol{x}_k\rangle\, \boldsymbol{e}_k \right\rangle \boldsymbol{u}_n && \text{by item (2b) page 231}\\
&= \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \langle \boldsymbol{x} \mid \boldsymbol{x}_m\rangle \sum_{k=1}^{\infty} \langle \boldsymbol{u}_n \mid \boldsymbol{x}_k\rangle^* \langle \boldsymbol{e}_m \mid \boldsymbol{e}_k\rangle\, \boldsymbol{u}_n && \text{by Definition J.1 page 255}\\
&= \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \langle \boldsymbol{x} \mid \boldsymbol{x}_m\rangle \langle \boldsymbol{u}_n \mid \boldsymbol{x}_m\rangle^*\, \boldsymbol{u}_n && \text{by item (2a) page 231 and Definition J.4 page 267}
\end{aligned}$$

Copyright © 2019 Daniel J. Greenhoe

$$= \sum_{n=1}^{\infty}\sum_{m=1}^{\infty} \langle \boldsymbol{x} \mid \boldsymbol{x}_m\rangle \langle \boldsymbol{x}_m \mid \boldsymbol{u}_n\rangle \boldsymbol{u}_n \qquad \text{by Definition J.1 page 255}$$

$$= \sum_{m=1}^{\infty} \langle \boldsymbol{x} \mid \boldsymbol{x}_m\rangle \sum_{n=1}^{\infty} \langle \boldsymbol{x}_m \mid \boldsymbol{u}_n\rangle \boldsymbol{u}_n$$

$$= \sum_{m=1}^{\infty} \langle \boldsymbol{x} \mid \boldsymbol{x}_m\rangle \boldsymbol{x}_m \qquad \text{by item (2d) page 231}$$

When is a set of orthonormal vectors in a Hilbert space $\boldsymbol{H}$ *total*? Theorem H.9 (next) offers some help.

Theorem H.9 (The Fourier Series Theorem). [25] *Let* $\left\{\boldsymbol{x}_n \in X \,\middle|\, n\in\mathbb{N}\right\}$ *be a set of vectors in a* HILBERT SPACE $\boldsymbol{H} \triangleq \left(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle\triangle \mid \triangledown\rangle\right)$ *and let* $\|\boldsymbol{x}\| \triangleq \sqrt{\langle \boldsymbol{x} \mid \boldsymbol{x}\rangle}$ *(Definition J.2 page 260)*.

THM

(A) $\left\{\boldsymbol{x}_n\right\}$ *is* ORTHONORMAL *in* $\boldsymbol{H}$ $\Longrightarrow$

$$\left\{\begin{array}{llrcll}
 & (1). & (\operatorname{span}\{\boldsymbol{x}_n\})^{-} & = & \boldsymbol{H} & & (\{\boldsymbol{x}_n\} \text{ is TOTAL in } \boldsymbol{H}) \\
\Longleftrightarrow & (2). & \langle \boldsymbol{x} \mid \boldsymbol{y}\rangle & \stackrel{\star}{=} & \displaystyle\sum_{n=1}^{\infty} \langle \boldsymbol{x} \mid \boldsymbol{x}_n\rangle \langle \boldsymbol{y} \mid \boldsymbol{x}_n\rangle^* & \forall \boldsymbol{x},\boldsymbol{y}\in \boldsymbol{X} & (\text{GENERALIZED PARSEVAL'S IDENTITY}) \\
\Longleftrightarrow & (3). & \|\boldsymbol{x}\|^2 & \stackrel{\star}{=} & \displaystyle\sum_{n=1}^{\infty} \left|\langle \boldsymbol{x} \mid \boldsymbol{x}_n\rangle\right|^2 & \forall \boldsymbol{x}\in X & (\text{PARSEVAL'S IDENTITY}) \\
\Longleftrightarrow & (4). & \boldsymbol{x} & \stackrel{\star}{=} & \displaystyle\sum_{n=1}^{\infty} \langle \boldsymbol{x} \mid \boldsymbol{x}_n\rangle \boldsymbol{x}_n & \forall \boldsymbol{x}\in X & (\text{FOURIER SERIES EXPANSION})
\end{array}\right\}$$

✎PROOF:

1. Proof that (1) $\Longrightarrow$ (2):

$$\langle \boldsymbol{x} \mid \boldsymbol{y}\rangle \stackrel{\star}{=} \left\langle \sum_{n=1}^{\infty} \langle \boldsymbol{x} \mid \boldsymbol{x}_n\rangle \boldsymbol{x}_n \,\middle|\, \sum_{m=1}^{\infty} \langle \boldsymbol{y} \mid \boldsymbol{x}_m\rangle \boldsymbol{x}_m \right\rangle \qquad \text{by (A) and (1)}$$

$$\stackrel{\star}{=} \sum_{n=1}^{\infty} \langle \boldsymbol{x} \mid \boldsymbol{x}_n\rangle \left\langle \boldsymbol{x}_n \,\middle|\, \sum_{m=1}^{\infty} \langle \boldsymbol{y} \mid \boldsymbol{x}_m\rangle \boldsymbol{x}_m \right\rangle \qquad \text{by property of } \langle\cdot\mid\cdot\rangle \quad \text{(Definition J.1 page 255)}$$

$$\stackrel{\star}{=} \sum_{n=1}^{\infty} \langle \boldsymbol{x} \mid \boldsymbol{x}_n\rangle \sum_{m=1}^{\infty} \langle \boldsymbol{y} \mid \boldsymbol{x}_m\rangle^* \langle \boldsymbol{x}_n \mid \boldsymbol{x}_m\rangle \qquad \text{by property of } \langle\cdot\mid\cdot\rangle \quad \text{(Definition J.1 page 255)}$$

$$\stackrel{\star}{=} \sum_{n=1}^{\infty} \langle \boldsymbol{x} \mid \boldsymbol{x}_n\rangle \sum_{m=1}^{\infty} \langle \boldsymbol{y} \mid \boldsymbol{x}_m\rangle^* \bar{\delta}_{mn} \qquad \text{by (A)}$$

$$\stackrel{\star}{=} \sum_{n=1}^{\infty} \langle \boldsymbol{x} \mid \boldsymbol{x}_n\rangle \langle \boldsymbol{y} \mid \boldsymbol{x}_n\rangle^* \qquad \text{by definition of } \bar{\delta}_n \quad \text{(Definition J.3 page 267)}$$

[25] Bachman and Narici (1966) pages 149–155 ⟨Theorem 9.12⟩, Kubrusly (2001) pages 360–363 ⟨Theorem 5.48⟩, Aliprantis and Burkinshaw (1998) pages 298–299 ⟨Theorem 34.2⟩, Christensen (2003) page 57 ⟨Theorem 3.4.2⟩, Berberian (1961) pages 52–53 ⟨Theorem II§8.3⟩, Heil (2011) pages 34–35 ⟨Theorem 1.50⟩, Bracewell (1978) page 112 ⟨Rayleigh's theorem⟩

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

2. Proof that (2) $\Longrightarrow$ (3):

$$\begin{aligned}
\|\boldsymbol{x}\|^2 &\triangleq \langle \boldsymbol{x} \mid \boldsymbol{x} \rangle && \text{by definition of } \textit{induced norm} && \text{(Theorem J.4 page 260)}\\
&= \sum_{n=1}^{\infty} \langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle \langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle^* && \text{by (2)}\\
&= \sum_{n=1}^{\infty} \left|\langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle\right|^2
\end{aligned}$$

3. Proof that (3) $\Longleftrightarrow$ (4) *not* using (A): by Proposition H.2 page 230

4. Proof that (3) $\Longrightarrow$ (1) (proof by contradiction):

   (a) Suppose $\{\boldsymbol{x}_n\}$ is *not total*.

   (b) Then there must exist a vector $\boldsymbol{y}$ in $\boldsymbol{H}$ such that the set $B \triangleq \{\boldsymbol{x}_n\} \cup \boldsymbol{y}$ is *orthonormal*.

   (c) Then $1 = \|y\|^2 \neq \sum_{n=1}^{\infty} \left|\langle \boldsymbol{y} \mid \boldsymbol{x}_n \rangle\right|^2 = 0$.

   (d) But this contradicts (3), and so $\{\boldsymbol{x}_n\}$ must be *total* and (3) $\Longrightarrow$ (1).

5. Extraneous proof that (3) $\Longrightarrow$ (4) (this proof is not really necessary here):

$$\begin{aligned}
\left\| \boldsymbol{x} - \sum_{n=1}^{\infty} \langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle \boldsymbol{x}_n \right\|^2 &= \|\boldsymbol{x}\|^2 - \sum_{n=1}^{\infty} |\langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle|^2 && \text{by } \textit{Bessel's Equality} && \text{(Theorem H.6 page 227)}\\
&= 0 && \text{by (3)}\\
\Longrightarrow \quad \boldsymbol{x} &\stackrel{*}{=} \sum_{n=1}^{\infty} \langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle \boldsymbol{x}_n && \text{by definition of } \stackrel{*}{=}
\end{aligned}$$

6. Extraneous proof that (A) $\Longrightarrow$ (4) (this proof is not really necessary here)

   (a) The sequence $\sum_{n=1}^{N} \left|\langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle\right|^2$ is *monotonically increasing* in n.

   (b) By Bessel's inequality (page 228), the sequence is upper bounded by $\|\boldsymbol{x}\|^2$:

$$\sum_{n=1}^{N} \left|\langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle\right|^2 \leq \|\boldsymbol{x}\|^2$$

   (c) Because this sequence is both monotonically increasing and bounded in n, it must equal its bound in the limit as n approaches infinity:

$$\lim_{N\to\infty} \sum_{n=1}^{N} \left|\langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle\right|^2 = \|\boldsymbol{x}\|^2 \tag{H.1}$$

   (d) If we combine this result with *Bessel's Equality* (Theorem H.6 page 227) we have

$$\begin{aligned}
\lim_{N\to\infty} \left\| \boldsymbol{x} - \sum_{n=1}^{N} \langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle \boldsymbol{x}_n \right\|^2 &= \|\boldsymbol{x}\|^2 - \lim_{N\to\infty} \sum_{n=1}^{N} |\langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle|^2 && \text{by } \textit{Bessel's Equality} \text{ (Theorem H.6 page 227)}\\
&= \|\boldsymbol{x}\|^2 - \|\boldsymbol{x}\|^2 && \text{by equation (H.1) page 233}\\
&= 0
\end{aligned}$$

Copyright © 2019 Daniel J. Greenhoe

Proposition H.3 (Fourier expansion). *Let* $\left\{ \boldsymbol{x}_n \in X \,\middle|\, {}_{n\in\mathbb{N}} \right\}$ *be a set of vectors in a* HILBERT SPACE $\left(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle \mid \triangledown \rangle\right)$.

PRP

$$\underbrace{\{\boldsymbol{x}_n\} \text{ is an ORTHONORMAL BASIS for } \boldsymbol{H}}_{(A)} \quad \Longrightarrow \quad \left\{ \underbrace{\boldsymbol{x} \stackrel{*}{=} \sum_{n=1}^{\infty} \alpha_n \boldsymbol{x}_n}_{(1)} \quad \Longleftrightarrow \quad \underbrace{\alpha_n = \langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle}_{(2)} \right\}$$

✎PROOF:

1. Proof that (1) $\Longrightarrow$ (2): by Lemma H.2 page 230
2. Proof that (1) $\Longleftarrow$ (2):

$$\begin{aligned}
\left\| \boldsymbol{x} - \sum_{n=1}^{\infty} \alpha_n \boldsymbol{x}_{n\in\mathbb{Z}} \right\|^2 &= \left\| \boldsymbol{x} - \sum_{n=1}^{\infty} \langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle \boldsymbol{x}_{n\in\mathbb{Z}} \right\|^2 && \text{by right hypothesis} \\
&= \|\boldsymbol{x}\|^2 - \sum_{n=1}^{\infty} |\langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle|^2 && \text{by Bessel's equality} && \text{(Theorem H.6 page 227)} \\
&= 0 && \text{by } \textit{Parseval's Identity} && \text{(Theorem H.9 page 232)} \\
&\stackrel{\text{def}}{\Longleftrightarrow} \quad \boldsymbol{x} \stackrel{*}{=} \sum_{n=1}^{\infty} \langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle \boldsymbol{x}_n && \text{by definition of } \textit{strong convergence}
\end{aligned}$$

✏

Proposition H.4 (Riesz-Fischer Theorem). [26] *Let* $\left\{ \boldsymbol{x}_n \in X \,\middle|\, {}_{n\in\mathbb{N}} \right\}$ *be a set of vectors in a* HILBERT SPACE $\left(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle \mid \triangledown \rangle\right)$.

PRP

$$\underbrace{\{\boldsymbol{x}_n\} \text{ is an ORTHONORMAL BASIS for } \boldsymbol{H}}_{(A)} \quad \Longrightarrow \quad \left\{ \underbrace{\sum_{n=1}^{\infty} |\alpha_n|^2 < \infty}_{(1)} \quad \Longleftrightarrow \quad \underbrace{\begin{array}{l} \exists \boldsymbol{x}\in\boldsymbol{H} \text{ such that} \\ \alpha_n = \langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle \end{array}}_{(2)} \right\}$$

✎PROOF:

1. Proof that (1) $\Longrightarrow$ (2):
   (a) If (1) is true, then let $\boldsymbol{x} \triangleq \sum_{n\in\mathbb{N}} \alpha_n \boldsymbol{x}_n$.
   (b) Then

$$\begin{aligned}
\langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle &= \left\langle \sum_{m\in\mathbb{N}} \alpha_m \boldsymbol{x}_m \,\middle|\, \boldsymbol{x}_n \right\rangle && \text{by definition of } \boldsymbol{x} \\
&= \sum_{m\in\mathbb{N}} \alpha_m \langle \boldsymbol{x}_m \mid \boldsymbol{x}_n \rangle && \text{by } \textit{homogeneous} \text{ property of } \langle \triangle \mid \triangledown \rangle && \text{(Definition J.1 page 255)} \\
&= \sum_{m\in\mathbb{N}} \alpha_m \bar{\delta}_{mn} && \text{by (A)} \\
&= \sum_{m\in\mathbb{N}} \alpha_n && \text{by definition of } \bar{\delta} && \text{(Definition J.3 page 267)}
\end{aligned}$$

[26] Young (2001) page 6

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

2. Proof that (1) $\Longleftarrow$ (2):

$$\begin{aligned}
\sum_{n\in\mathbb{N}} |\alpha_n|^2 &= \sum_{n\in\mathbb{N}} |\langle \boldsymbol{x} \mid \boldsymbol{x}_n\rangle|^2 && \text{by (2)} \\
&\le \|\boldsymbol{x}\|^2 && \text{by } \textit{Bessel's Inequality} \quad \text{(Theorem H.7 page 228)} \\
&\le \infty
\end{aligned}$$

✏

Theorem H.10. [27]

THM

All SEPARABLE HILBERT SPACES *are* ISOMORPHIC. *That is,*

$$\left\{\begin{array}{l} \boldsymbol{X} \text{ is a separable} \\ \text{Hilbert space} \quad \textit{and} \\ \boldsymbol{Y} \text{ is a separable} \\ \text{Hilbert space} \end{array}\right\} \Longrightarrow \left\{\begin{array}{llll} \textit{there is a BIJECTIVE operator } \mathbf{M} \in \boldsymbol{Y}^{\boldsymbol{X}} \textit{ such that} & & & \\ (1).\ \boldsymbol{y} & = & \mathbf{M}\boldsymbol{x} & \forall \boldsymbol{x}\in\boldsymbol{X}, \boldsymbol{y}\in\boldsymbol{Y} \quad \textit{and} \\ (2).\ \|\mathbf{M}\boldsymbol{x}\| & = & \|\boldsymbol{x}\| & \forall \boldsymbol{x}\in\boldsymbol{X} \quad \textit{and} \\ (3).\ \langle \mathbf{M}\boldsymbol{x} \mid \mathbf{M}\boldsymbol{y}\rangle & = & \langle \boldsymbol{x} \mid \boldsymbol{y}\rangle & \forall \boldsymbol{x}\in\boldsymbol{X}, \boldsymbol{y}\in\boldsymbol{Y} \end{array}\right\}$$

✎PROOF:

1. Let $\boldsymbol{X} \triangleq \big(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle \mid \triangledown\rangle\big)$ be a *separable Hilbert space* with *orthonormal basis* $\{\boldsymbol{x}_n \mid_{n\in\mathbb{N}}\}$.
   Let $\boldsymbol{Y} \triangleq \big(Y, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle \mid \triangledown\rangle\big)$ be a *separable Hilbert space* with *orthonormal basis* $\{\boldsymbol{y}_n \mid_{n\in\mathbb{N}}\}$.

2. Proof that there exists *bijective* operator $\mathbf{M}$ and its inverse $\mathbf{M}^{-1}$ between $\{\boldsymbol{x}_n\}$ and $\{\boldsymbol{y}_n\}$:

   (a) Let $\mathbf{M}$ be defined such that $\boldsymbol{y}_n \triangleq \mathbf{M}\boldsymbol{x}_n$.

   (b) Thus $\mathbf{M}$ is a *bijection* between $\{\boldsymbol{x}_n\}$ and $\{\boldsymbol{y}_n\}$.

   (c) Because $\mathbf{M}$ is a *bijection* between $\{\boldsymbol{x}_n\}$ and $\{\boldsymbol{y}_n\}$, $\mathbf{M}$ has an inverse operator $\mathbf{M}^{-1}$ between $\{\boldsymbol{x}_n\}$ and $\{\boldsymbol{y}_n\}$ such that $\boldsymbol{x}_n = \mathbf{M}^{-1}\boldsymbol{y}_n$.

3. Proof that $\mathbf{M}$ and $\mathbf{M}^{-1}$ are *bijective* operators between $\boldsymbol{X}$ and $\boldsymbol{Y}$:

   (a) Proof that $\mathbf{M}$ maps $\boldsymbol{X}$ *into* $\boldsymbol{Y}$:

$$\begin{aligned}
\boldsymbol{x} \in \boldsymbol{X} \iff \boldsymbol{x} &\stackrel{*}{=} \sum_{n\in\mathbb{N}} \langle \boldsymbol{x} \mid \boldsymbol{x}_n\rangle \boldsymbol{x}_n && \text{by } \textit{Fourier expansion} \quad \text{(Theorem H.9 page 232)} \\
\Longrightarrow \exists \boldsymbol{y} \in \boldsymbol{Y} \quad &\text{such that} \quad \langle \boldsymbol{y} \mid \boldsymbol{y}_n\rangle = \langle \boldsymbol{x} \mid \boldsymbol{x}_n\rangle && \text{by } \textit{Riesz-Fischer Thm.} \quad \text{(Proposition H.4 page 234)} \\
\Longrightarrow & \\
\boldsymbol{y} &= \sum_{n\in\mathbb{N}} \langle \boldsymbol{y} \mid \boldsymbol{y}_n\rangle \boldsymbol{y}_n && \text{by } \textit{Fourier expansion} \quad \text{(Theorem H.9 page 232)} \\
&= \sum_{n\in\mathbb{N}} \langle \boldsymbol{x} \mid \boldsymbol{x}_n\rangle \boldsymbol{y}_n && \text{by } \textit{Riesz-Fischer Thm.} \quad \text{(Proposition H.4 page 234)} \\
&= \sum_{n\in\mathbb{N}} \langle \boldsymbol{x} \mid \boldsymbol{x}_n\rangle \mathbf{M}\boldsymbol{x}_n && \text{by definition of } \mathbf{M} \quad \text{(item (2a) page 235)} \\
&= \mathbf{M} \sum_{n\in\mathbb{N}} \langle \boldsymbol{x} \mid \boldsymbol{x}_n\rangle \boldsymbol{x}_n && \text{by prop. of linear ops.} \quad \text{(Theorem N.1 page 304)} \\
&= \mathbf{M}\boldsymbol{x} && \text{by definition of } \boldsymbol{x}
\end{aligned}$$

[27] Young (2001) page 6

COPYRIGHT © 2019 DANIEL J. GREENHOE | CC BY-NC-ND

(b) Proof that $\mathbf{M}^{-1}$ maps $\boldsymbol{Y}$ *into* $\boldsymbol{X}$:

$$\begin{aligned}
\boldsymbol{y} \in \boldsymbol{Y} &\iff \boldsymbol{y} \stackrel{*}{=} \sum_{n\in\mathbb{N}} \langle \boldsymbol{y} \mid \boldsymbol{y}_n \rangle \boldsymbol{y}_n && \text{by } \textit{Fourier expansion} && \text{(Theorem H.9 page 232)}\\
&\implies \exists \boldsymbol{x} \in \boldsymbol{X} \quad \text{such that} \quad \langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle = \langle \boldsymbol{y} \mid \boldsymbol{y}_n \rangle && \text{by } \textit{Riesz-Fischer Thm.} && \text{(Proposition H.4 page 234)}\\
&\implies\\
\boldsymbol{x} &= \sum_{n\in\mathbb{N}} \langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle \boldsymbol{x}_n && \text{by } \textit{Fourier expansion} && \text{(Theorem H.9 page 232)}\\
&= \sum_{n\in\mathbb{N}} \langle \boldsymbol{y} \mid \boldsymbol{y}_n \rangle \boldsymbol{x}_n && \text{by } \textit{Riesz-Fischer Thm.} && \text{(Proposition H.4 page 234)}\\
&= \sum_{n\in\mathbb{N}} \langle \boldsymbol{y} \mid \boldsymbol{y}_n \rangle \mathbf{M}^{-1} \boldsymbol{y}_n && \text{by definition of } \mathbf{M}^{-1} && \text{(item (2c) page 235)}\\
&= \mathbf{M}^{-1} \sum_{n\in\mathbb{N}} \langle \boldsymbol{y} \mid \boldsymbol{y}_n \rangle \boldsymbol{y}_n && \text{by prop. of } \textit{linear ops.} && \text{(Theorem N.1 page 304)}\\
&= \mathbf{M}^{-1} \boldsymbol{y} && \text{by definition of } \boldsymbol{y}
\end{aligned}$$

4. Proof for (2):

$$\begin{aligned}
\|\mathbf{M}\boldsymbol{x}\|^2 &= \left\| \mathbf{M} \sum_{n\in\mathbb{N}} \langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle \boldsymbol{x}_n \right\|^2 && \text{by } \textit{Fourier expansion} && \text{(Theorem H.9 page 232)}\\
&= \left\| \sum_{n\in\mathbb{N}} \langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle \mathbf{M}\boldsymbol{x}_n \right\|^2 && \text{by property of } \textit{linear operators} && \text{(Theorem N.1 page 304)}\\
&= \left\| \sum_{n\in\mathbb{N}} \langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle \boldsymbol{y}_n \right\|^2 && \text{by definition of } \mathbf{M} && \text{(item (2a) page 235)}\\
&= \sum_{n\in\mathbb{N}} \left| \langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle \right|^2 && \text{by } \textit{Parseval's Identity} && \text{(Proposition H.4 page 234)}\\
&= \left\| \sum_{n\in\mathbb{N}} \langle \boldsymbol{x} \mid \boldsymbol{x}_n \rangle \boldsymbol{x}_n \right\|^2 && \text{by } \textit{Parseval's Identity} && \text{(Proposition H.4 page 234)}\\
&= \|\boldsymbol{x}\|^2 && \text{by } \textit{Fourier expansion} && \text{(Theorem H.9 page 232)}
\end{aligned}$$

5. Proof for (3): by (2) and Theorem N.23 page 324

✎

Theorem H.11. [28] *Let* $\boldsymbol{H}$ *be a* HILBERT SPACE.

THM

$\boldsymbol{H}$ *has a* SCHAUDER BASIS $\iff$ $\boldsymbol{H}$ *is* SEPARABLE

Theorem H.12. [29] *Let* $\boldsymbol{H}$ *be a* HILBERT SPACE.

THM

$\boldsymbol{H}$ *has an* ORTHONORMAL BASIS $\iff$ $\boldsymbol{H}$ *is* SEPARABLE

[28] Bachman et al. (2000) page 112 ⟨3.4.8⟩, Berberian (1961) page 53 ⟨Theorem II§8.3⟩
[29] Kubrusly (2001) page 357 ⟨Proposition 5.43⟩

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

H.6 Riesz bases in Hilbert spaces

Definition H.11. [30] *Let* $\{\boldsymbol{x}_n \in X \mid n\in\mathbb{N}\}$ *be a set of vectors in a* SEPARABLE HILBERT SPACE $\boldsymbol{H} \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+},$

DEF $\{\boldsymbol{x}_n\}$ *is a* ***Riesz basis*** *for* $\boldsymbol{H}$ *if* $\{\boldsymbol{x}_n\}$ *is* EQUIVALENT *(Definition H.8 page 225) to some* ORTHONORMAL BASIS *(Definition H.9 page 229) in* $\boldsymbol{H}$.

Definition H.12. [31] *Let* $(\!(\boldsymbol{x}_n \in X)\!)_{n\in\mathbb{N}}$ *be a sequence of vectors in a* SEPARABLE HILBERT SPACE $\boldsymbol{H} \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle\triangle\,|\,\triangledown\rangle)$.

DEF *The sequence* $(\!(\boldsymbol{x}_n)\!)$ *is a* ***Riesz sequence*** *for* $\boldsymbol{H}$ *if*

$$\exists A, B \in \mathbb{R}^+ \quad \text{such that} \quad A\sum_{n=1}^{\infty}|\alpha_n|^2 \le \left\|\sum_{n=1}^{\infty}\alpha_n\boldsymbol{x}_n\right\|^2 \le B\sum_{n=1}^{\infty}|\alpha_n|^2 \qquad \forall (\!(\alpha_n)\!)\in\ell^2_{\mathbb{F}}.$$

Definition H.13. *Let* $\boldsymbol{X} \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle\triangle\,|\,\triangledown\rangle)$ *be an* INNER PRODUCT SPACE *(Definition J.1 page 255)*.

DEF *The sequences* $(\!(\boldsymbol{x}_n \in X)\!)_{n\in\mathbb{Z}}$ *and* $(\!(\boldsymbol{y}_n \in X)\!)_{n\in\mathbb{Z}}$ *are* ***biorthogonal*** *with respect to each other in* $\boldsymbol{X}$ *if* $\langle \boldsymbol{x}_n \,|\, \boldsymbol{y}_m\rangle = \bar{\delta}_{nm}$

Lemma H.3. [32] *Let* $\{\boldsymbol{x}_n \mid n\in\mathbb{N}\}$ *be a sequence in a* HILBERT SPACE $\boldsymbol{X} \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle\triangle\,|\,\triangledown\rangle)$. *Let* $\{\boldsymbol{y}_n \mid n\in\mathbb{N}\}$ *be a sequence in a* HILBERT SPACE $\boldsymbol{Y} \triangleq (Y, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle\triangle\,|\,\triangledown\rangle)$. *Let*

LEM

$$\left\{\begin{array}{lll}
(i). & \{\boldsymbol{x}_n\} \text{ is TOTAL in } \boldsymbol{X} & \text{and} \\
(ii). & \text{There exists } A>0 \text{ such that } A\sum_{n\in C}|a_n|^2 \le \left\|\sum_{n\in C}a_n\boldsymbol{x}_n\right\|^2 \text{ for finite } C & \text{and} \\
(iii). & \text{There exists } B>0 \text{ such that } \left\|\sum_{n=1}^{\infty}b_n\boldsymbol{y}_n\right\|^2 \le B\sum_{n=1}^{\infty}|b_n|^2 \quad \forall (\!(b_n)\!)_{n\in\mathbb{N}}\in\ell^2_{\mathbb{F}} &
\end{array}\right\} \Longrightarrow$$

$$\left\{\begin{array}{lll}
(1). & \mathbf{R}^\circ \text{ is a linear bounded operator that maps from } \operatorname{span}\{\boldsymbol{x}_n\} \text{ to } \operatorname{span}\{\boldsymbol{y}_n\} & \\
 & \text{where } \mathbf{R}^\circ\sum_{n\in C}c_n\boldsymbol{x}_n \triangleq \sum_{n\in C}c_n\boldsymbol{y}_n, \text{ for some sequence } (\!(c_n)\!) \text{ and finite set } C & \text{and} \\
(2). & \mathbf{R} \text{ has a unique extension to a bounded operator } \mathbf{R} \text{ that maps from } \boldsymbol{X} \text{ to } \boldsymbol{Y} & \text{and} \\
(3). & |||\mathbf{R}^\circ||| \le \frac{B}{A} & \text{and} \\
(4). & |||\mathbf{R}||| \le \frac{B}{A} &
\end{array}\right\}$$

Theorem H.13. [33] *Let* $\{\boldsymbol{x}_n \in X \mid n\in\mathbb{N}\}$ *be a set of vectors in a* SEPARABLE HILBERT SPACE $\boldsymbol{H} \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle\triangle\,|\,\triangledown\rangle)$.

THM

$$\left\{\begin{array}{l}\{\boldsymbol{x}_n\} \text{ is a RIESZ BASIS} \\ \text{for } \boldsymbol{H}\end{array}\right\} \Longleftrightarrow \left\{\begin{array}{lll}
(1). & \{\boldsymbol{x}_n\} \text{ is TOTAL in } \boldsymbol{H} & \text{and} \\
(2). & \exists A,B\in\mathbb{R}^+ \quad \text{such that} \quad \forall (\!(\alpha_n)\!)\in\ell^2_{\mathbb{F}}, & \\
 & A\sum_{n=1}^{\infty}|\alpha_n|^2 \le \left\|\sum_{n=1}^{\infty}\alpha_n\boldsymbol{x}_n\right\|^2 \le B\sum_{n=1}^{\infty}|\alpha_n|^2 &
\end{array}\right\}$$

✎PROOF:

[30] Young (2001) page 27 ⟨Definition 1.8.2⟩, Christensen (2003) page 63 ⟨Definition 3.6.1⟩, Heil (2011) page 196 ⟨Definition 7.9⟩

[31] Christensen (2003) pages 66–68 ⟨page 68 and (3.24) on page 66⟩, Wojtaszczyk (1997) page 20 ⟨Definition 2.6⟩

[32] Christensen (2003) pages 65–66 ⟨Lemma 3.6.5⟩

[33] Young (2001) page 27 ⟨Theorem 1.8.9⟩, Christensen (2003) page 66 ⟨Theorem 3.6.6⟩, Heil (2011) pages 197–198 ⟨Theorem 7.13⟩, Christensen (2008) pages 61–62 ⟨Theorem 3.3.7⟩

COPYRIGHT © 2019 DANIEL J. GREENHOE | https://github.com/dgreenhoe/pdfs/blob/master/abcstat.pdf | CC BY-NC-ND

1. Proof for ($\Longrightarrow$) case:

   (a) Proof that *Riesz basis* hypothesis $\Longrightarrow$ (1): all bases for $\boldsymbol{H}$ are *total* in $\boldsymbol{H}$.

   (b) Proof that *Riesz basis* hypothesis $\Longrightarrow$ (2):

      i. Let $(\boldsymbol{u}_n)_{n\in\mathbb{N}}$ be an *orthonormal basis* for $\boldsymbol{H}$.

      ii. Let $\mathbf{R}$ be a *bounded bijective* operator such that $\boldsymbol{x}_n = \mathbf{R}\boldsymbol{u}_n$.

      iii. Proof for upper bound B:

$$\begin{aligned}
\left\|\sum_{n=1}^{\infty}\alpha_n\boldsymbol{x}_n\right\|^2 &= \left\|\sum_{n=1}^{\infty}\alpha_n\mathbf{R}\boldsymbol{u}_n\right\|^2 && \text{by definition of } \mathbf{R} && \text{(item (1(b)ii))}\\
&= \left\|\mathbf{R}\sum_{n=1}^{\infty}\alpha_n\boldsymbol{u}_n\right\|^2 && \text{by Theorem N.1 page 304} && \\
&\le |||\mathbf{R}|||^2\left\|\sum_{n=1}^{\infty}\alpha_n\boldsymbol{u}_n\right\|^2 && \text{by Theorem N.6 page 310} && \\
&= |||\mathbf{R}|||^2\sum_{n=1}^{\infty}\left\|\alpha_n\boldsymbol{u}_n\right\|^2 && \text{by } \textit{Pythagorean Theorem} && \text{(Theorem J.10 page 268)}\\
&= |||\mathbf{R}|||^2\sum_{n=1}^{\infty}|\alpha|^2\left\|\boldsymbol{u}_n\right\|^2 && \text{by } \textit{homogeneous} \text{ property of norms} && \text{(Definition K.1 page 271)}\\
&= \underbrace{|||\mathbf{R}|||^2}_{B}\sum_{n=1}^{\infty}|\alpha|^2 && \text{by definition of } \textit{orthonormality} && \text{(Definition J.4 page 267)}
\end{aligned}$$

      iv. Proof for lower bound A:

$$\begin{aligned}
\left\|\sum_{n=1}^{\infty}\alpha_n\boldsymbol{x}_n\right\|^2 &= \frac{|||\mathbf{R}^{-1}|||^2}{|||\mathbf{R}^{-1}|||^2}\left\|\sum_{n=1}^{\infty}\alpha_n\boldsymbol{x}_n\right\|^2 && \text{because } |||\mathbf{R}^{-1}||| > 0 && \text{(Proposition N.1 page 308)}\\
&\ge \frac{1}{|||\mathbf{R}^{-1}|||^2}\left\|\mathbf{R}^{-1}\sum_{n=1}^{\infty}\alpha_n\boldsymbol{x}_n\right\|^2 && \text{by Theorem N.6 page 310} && \\
&= \frac{1}{|||\mathbf{R}^{-1}|||^2}\left\|\mathbf{R}^{-1}\sum_{n=1}^{\infty}\alpha_n\mathbf{R}\boldsymbol{u}_n\right\|^2 && \text{by definition of } \mathbf{R} && \text{(item (1(b)ii) page 238)}\\
&= \frac{1}{|||\mathbf{R}^{-1}|||^2}\left\|\mathbf{R}^{-1}\mathbf{R}\sum_{n=1}^{\infty}\alpha_n\boldsymbol{u}_n\right\|^2 && \text{by property of } \textit{linear operators} && \text{(Theorem N.1 page 304)}\\
&= \frac{1}{|||\mathbf{R}^{-1}|||^2}\left\|\sum_{n=1}^{\infty}\alpha_n\boldsymbol{u}_n\right\|^2 && \text{by definition of inverse op.} && \text{(Definition N.2 page 303)}\\
&= \frac{1}{|||\mathbf{R}^{-1}|||^2}\sum_{n=1}^{\infty}\left\|\alpha_n\boldsymbol{u}_n\right\|^2 && \text{by } \textit{Pythagorean Theorem} && \text{(Theorem J.10 page 268)}\\
&= \frac{1}{|||\mathbf{R}^{-1}|||^2}\sum_{n=1}^{\infty}\left|\alpha_n\right|^2\left\|\boldsymbol{u}_n\right\|^2 && \text{by } \|\cdot\| \textit{ homogeneous} \text{ prop.} && \text{(Definition K.1 page 271)}\\
&= \underbrace{\frac{1}{|||\mathbf{R}^{-1}|||^2}}_{A}\sum_{n=1}^{\infty}\left|\alpha_n\right|^2 && \text{by def. of } \textit{orthonormality} && \text{(Definition J.4 page 267)}
\end{aligned}$$

2. Proof for ($\Longrightarrow$) case:

BY-NC-ND | Copyright © 2019 Daniel J. Greenhoe

(a) Let $\{\boldsymbol{u}_n \mid n\in\mathbb{N}\}$ be an *orthonormal basis* for $\boldsymbol{H}$.

(b) Using (2) and Lemma H.3 (page 237), construct an bounded extension operator $\mathbf{R}$ such that $\mathbf{R}\boldsymbol{u}_n = \boldsymbol{x}_n$ for all $n \in \mathbb{N}$.

(c) Using (2) and Lemma H.3 (page 237), construct an bounded extension operator $\mathbf{S}$ such that $\mathbf{S}\boldsymbol{x}_n = \boldsymbol{u}_n$ for all $n \in \mathbb{N}$.

(d) Then, $\mathbf{RV}\boldsymbol{x} = \mathbf{VR}\boldsymbol{x} \implies \mathbf{V} = \mathbf{R}^{-1}$, and so $\mathbf{R}$ is a bounded invertible operator

(e) and $\{\boldsymbol{x}_n\}$ is a *Riesz sequence*.

✏

Theorem H.14. [34] *Let* $\boldsymbol{H} \triangleq \left(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle\triangle \mid \triangledown\rangle\right)$ *be a* SEPARABLE HILBERT SPACE.

THM

$$\left\{\begin{array}{l}(\!(\boldsymbol{x}_n \in \boldsymbol{H})\!)_{n\in\mathbb{Z}} \text{ is a} \\ \text{RIESZ BASIS for } \boldsymbol{H}\end{array}\right\} \implies \left\{\begin{array}{l}\text{There exists } (\!(\boldsymbol{y}_n \in \boldsymbol{H})\!)_{n\in\mathbb{Z}} \text{ such that} \\ (1).\quad (\!(\boldsymbol{x}_n)\!) \text{ and } (\!(\boldsymbol{y}_n)\!) \text{ are BIORTHOGONAL} \quad \text{and} \\ (2).\quad (\!(\boldsymbol{y}_n)\!) \text{ is also a RIESZ BASIS for } \boldsymbol{H} \quad \text{and} \\ (3).\quad \exists B>A>0 \quad \text{such that} \\ \qquad A\sum_{n=1}^{\infty}|a_n|^2 \le \left\|\sum_{n=1}^{\infty} a_n\boldsymbol{x}_n\right\|^2 = \left\|\sum_{n=1}^{\infty} a_n\boldsymbol{y}_n\right\|^2 \le B\sum_{n=1}^{\infty}|a_n|^2 \\ \qquad \forall (\!(a_n)\!)_{n\in\mathbb{N}} \in \ell^2_{\mathbb{F}}\end{array}\right\}$$

✎PROOF:

1. Proof for (1):

(a) Let $\boldsymbol{e}_n$ be the *unit vector* in $\boldsymbol{H}$ such that the nth element of $\boldsymbol{e}_n$ is 1 and all other elements are 0.

(b) Let $\mathbf{M}$ be an operator on $\boldsymbol{H}$ such that $\mathbf{M}\boldsymbol{e}_n = \boldsymbol{x}_n$.

(c) Note that $\mathbf{M}$ is *isometric*, and as such $\|\mathbf{M}\boldsymbol{x}\| = \|\boldsymbol{x}\| \quad \forall \boldsymbol{x}\in\boldsymbol{H}$.

(d) Let $\boldsymbol{y}_n \triangleq \left(\mathbf{M}^{-1}\right)^*$.

(e) Then,

$$\begin{aligned}\langle \boldsymbol{y}_n \mid \boldsymbol{x}_m\rangle &= \left\langle \left(\mathbf{M}^{-1}\right)^* \boldsymbol{e}_n \mid \mathbf{M}\boldsymbol{e}_m\right\rangle \\ &= \left\langle \boldsymbol{e}_n \mid \mathbf{M}^{-1}\mathbf{M}\boldsymbol{e}_m\right\rangle \\ &= \langle \boldsymbol{e}_n \mid \boldsymbol{e}_m\rangle \\ &= \bar{\delta}_{nm} \\ &\implies \{\boldsymbol{x}_n\} \text{ and } \{\boldsymbol{y}_n\} \text{ are } \textit{biorthogonal} \qquad \text{by Definition J.4 page 267}\end{aligned}$$

[34] Wojtaszczyk (1997) page 20 ⟨Lemma 2.7(a)⟩

COPYRIGHT © 2019 DANIEL J. GREENHOE | CC BY-NC-ND

2. Proof for (3):

$$\begin{aligned}
\left\|\sum_{n\in\mathbb{Z}}\alpha_n\boldsymbol{y}_n\right\| &= \left\|\sum_{n\in\mathbb{Z}}\alpha_n\left(\mathbf{M}^{-1}\right)^*\boldsymbol{e}_n\right\| && \text{by definition of } \boldsymbol{y}_n && \text{(Proposition 1d page 239)}\\
&= \left\|\left(\mathbf{M}^{-1}\right)^*\sum_{n\in\mathbb{Z}}\alpha_n\boldsymbol{e}_n\right\| && \text{by property of } \textit{linear ops.} \\
&= \left\|\sum_{n\in\mathbb{Z}}\alpha_n\boldsymbol{e}_n\right\| && \text{because } \left(\mathbf{M}^{-1}\right)^* \text{ is } \textit{isometric} && \text{(Definition N.10 page 323)}\\
&= \left\|\mathbf{M}\sum_{n\in\mathbb{Z}}\alpha_n\boldsymbol{e}_n\right\| && \text{because } \mathbf{M} \text{ is } \textit{isometric} && \text{(Definition N.10 page 323)}\\
&= \left\|\sum_{n\in\mathbb{Z}}\alpha_n\mathbf{M}\boldsymbol{e}_n\right\| && \text{by property of } \textit{linear operators} \\
&= \left\|\sum_{n\in\mathbb{Z}}\alpha_n\boldsymbol{x}_n\right\| && \text{by definition of } \mathbf{M} \\
&\Longrightarrow \{\boldsymbol{y}_n\} \text{ is a } \textit{Riesz basis} && \text{by left hypothesis}
\end{aligned}$$

3. Proof for (2): by (3) and definition of *Riesz basis* (Definition H.11 page 237)

✏

Proposition H.5. [35] *Let* $\{\boldsymbol{x}_n|_{n\in\mathbb{N}}\}$ *be a set of vectors in a* HILBERT SPACE $\boldsymbol{H} \triangleq \left(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle\triangle\,|\,\triangledown\rangle\right)$.

PRP

$$\left\{\begin{array}{l}\{\boldsymbol{x}_n\} \textit{ is a } \text{RIESZ BASIS} \textit{ for } \boldsymbol{H} \textit{ with} \\ A\sum_{n=1}^{\infty}|a_n|^2 \le \left\|\sum_{n=1}^{\infty}a_n\boldsymbol{x}_n\right\|^2 \le B\sum_{n=1}^{\infty}|a_n|^2 \\ \forall\{a_n\}\in\ell^2_{\mathbb{F}}\end{array}\right\} \Longrightarrow \left\{\begin{array}{l}\{\boldsymbol{x}_n\} \textit{ is a } \text{FRAME} \textit{ for } \boldsymbol{H} \textit{ with} \\ \underbrace{\frac{1}{B}\|\boldsymbol{x}\|^2 \le \sum_{n=1}^{\infty}|\langle\boldsymbol{x}\,|\,\boldsymbol{x}_n\rangle|^2 \le \frac{1}{A}\|\boldsymbol{x}\|^2}_{\text{STABILITY CONDITION}} \\ \forall\boldsymbol{x}\in\boldsymbol{H}\end{array}\right\}$$

✎PROOF:

1. Let $\{\boldsymbol{y}_n|_{n\in\mathbb{N}}\}$ be a *Riesz basis* that is *biorthogonal* to $\{\boldsymbol{x}_n|_{n\in\mathbb{N}}\}$ (Theorem H.14 page 239).

2. Let $\boldsymbol{x} \triangleq \sum_{n=1}^{\infty}a_n\boldsymbol{y}_n$.

3. lemma:

$$\begin{aligned}
\sum_{n=1}^{\infty}|\langle\boldsymbol{x}\,|\,\boldsymbol{x}_n\rangle|^2 &= \sum_{n=1}^{\infty}\left|\left\langle\sum_{m=1}^{\infty}a_n\boldsymbol{y}_m\,|\,\boldsymbol{x}_n\right\rangle\right|^2 && \text{by definition of } \boldsymbol{x} && \text{(item (2) page 240)}\\
&= \sum_{n=1}^{\infty}\left|\sum_{m=1}^{\infty}a_n\langle\boldsymbol{y}_m\,|\,\boldsymbol{x}_n\rangle\right|^2 && \text{by } \textit{homogeneous} \text{ property of } \langle\triangle\,|\,\triangledown\rangle && \text{(Definition J.1 page 255)}\\
&= \sum_{n=1}^{\infty}\left|\sum_{m=1}^{\infty}a_n\bar{\delta}_{mn}\right|^2 && \text{by definition of } \textit{biorthogonal} && \text{(Definition H.13 page 237)}\\
&= \sum_{n=1}^{\infty}|a_n|^2 && \text{by definition of } \bar{\delta} && \text{(Definition J.3 page 267)}
\end{aligned}$$

[35] Igari (1996) page 220 ⟨Lemma 9.8⟩, Wojtaszczyk (1997) pages 20–21 ⟨Lemma 2.7(a)⟩

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

4. Then

$$\begin{aligned}
& A\sum_{n=1}^{\infty}|a_n|^2 \le \left\|\sum_{n=1}^{\infty}a_n\boldsymbol{x}_n\right\|^2 \le B\sum_{n=1}^{\infty}|a_n|^2 && \text{by definition of } \{\boldsymbol{y}_n\} \text{ (item (1) page 240)}\\
\Longrightarrow\ & A\sum_{n=1}^{\infty}|a_n|^2 \le \left\|\sum_{n=1}^{\infty}a_n\boldsymbol{y}_n\right\|^2 \le B\sum_{n=1}^{\infty}|a_n|^2 && \text{by definition of } \{\boldsymbol{y}_n\} \text{ (item (1) page 240)}\\
\Longrightarrow\ & A\sum_{n=1}^{\infty}|a_n|^2 \le \|\boldsymbol{x}\|^2 \le B\sum_{n=1}^{\infty}|a_n|^2 && \text{by definition of } \boldsymbol{x} \text{ (item (2) page 240)}\\
\Longrightarrow\ & A\sum_{n=1}^{\infty}|\langle\boldsymbol{x}\,|\,\boldsymbol{x}_n\rangle|^2 \le \|\boldsymbol{x}\|^2 \le B\sum_{n=1}^{\infty}|\langle\boldsymbol{x}\,|\,\boldsymbol{x}_n\rangle|^2 && \text{by (3) lemma}\\
\Longrightarrow\ & \frac{1}{B}\|\boldsymbol{x}\|^2 \le \sum_{n=1}^{\infty}|\langle\boldsymbol{x}\,|\,\boldsymbol{x}_n\rangle|^2 \le \frac{1}{A}\|\boldsymbol{x}\|^2
\end{aligned}$$

✏

Theorem H.15 (Battle-Lemarié orthogonalization). [36] *Let* $\tilde{\mathsf{f}}(\omega)$ *be the* FOURIER TRANSFORM *(Definition O.2 page 333) of a function* $\mathsf{f}\in L^2_{\mathbb{R}}$.

THM

$$\left\{\begin{array}{ll} 1. & \{\mathbf{T}^n\mathsf{g}\,|\,_{n\in\mathbb{Z}}\} \text{ is a RIESZ BASIS for } L^2_{\mathbb{R}} \text{ and}\\[2ex] 2. & \tilde{\mathsf{f}}(\omega)\triangleq\dfrac{\tilde{\mathsf{g}}(\omega)}{\sqrt{2\pi\sum\limits_{n\in\mathbb{Z}}|\tilde{\mathsf{g}}(\omega+2\pi n)|^2}}\end{array}\right\}\Longrightarrow\left\{\begin{array}{l}\{\mathbf{T}^n\mathsf{f}\,|\,_{n\in\mathbb{Z}}\}\\ \text{is an ORTHONORMAL BASIS for } L^2_{\mathbb{R}}\end{array}\right\}$$

✎PROOF:

1. Proof that $\{\mathbf{T}^n\mathsf{f}\,|\,_{n\in\mathbb{Z}}\}$ is orthonormal:

$$\begin{aligned}
\tilde{\mathsf{S}}_{\phi\phi}(\omega) &= 2\pi\sum_{n\in\mathbb{Z}}\left|\tilde{\mathsf{f}}(\omega+2\pi n)\right|^2 && \text{by Theorem ?? page ??}\\
&= 2\pi\sum_{n\in\mathbb{Z}}\left|\frac{\tilde{\mathsf{g}}(\omega+2\pi n)}{\sqrt{2\pi\sum\limits_{m\in\mathbb{Z}}|\tilde{\mathsf{g}}(\omega+2\pi(n-m))|^2}}\right|^2 && \text{by left hypothesis}\\
&= \sum_{n\in\mathbb{Z}}\left|\frac{\tilde{\mathsf{g}}(\omega+2\pi n)}{\sqrt{\sum\limits_{m\in\mathbb{Z}}|\tilde{\mathsf{g}}(\omega+2\pi m)|^2}}\right|^2\\
&= \sum_{n\in\mathbb{Z}}\left|\frac{1}{\sqrt{\sum\limits_{m\in\mathbb{Z}}|\tilde{\mathsf{g}}(\omega+2\pi m)|^2}}\right|^2|\tilde{\mathsf{g}}(\omega+2\pi n)|^2\\
&= \frac{1}{\sum\limits_{m\in\mathbb{Z}}|\tilde{\mathsf{g}}(\omega+2\pi m)|^2}\sum_{n\in\mathbb{Z}}|\tilde{\mathsf{g}}(\omega+2\pi n)|^2\\
&= 1\\
\Longrightarrow\ & \{\mathsf{f}_n\,|\,_{n\in\mathbb{Z}}\} \text{ is orthonormal} && \text{by Theorem ?? page ??}
\end{aligned}$$

[36] Wojtaszczyk (1997) page 25 ⟨Remark 2.4⟩, Vidakovic (1999), page 71, Mallat (1989), page 72, Mallat (1999), page 225, Daubechies (1992) page 140 ⟨(5.3.3)⟩

COPYRIGHT © 2019 DANIEL J. GREENHOE

2. Proof that $\{\mathbf{T}^n \mathsf{f} \mid n\in\mathbb{Z}\}$ is a basis for V_0: by Lemma H.1 page 225.

✏

H.7 Frames in Hilbert spaces

Definition H.14. [37] *Let* $\{\boldsymbol{x}_n \in X \mid n\in\mathbb{N}\}$ *be a set of vectors in a* HILBERT SPACE $\boldsymbol{H} \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle\triangle\mid\triangledown\rangle)$.

DEF

The set $\{\boldsymbol{x}_n\}$ *is a* ***frame*** *for* $\boldsymbol{H}$ *if* (STABILITY CONDITION)

$$\exists A, B \in \mathbb{R}^+ \quad \text{such that} \quad A\,\|\boldsymbol{x}\|^2 \le \sum_{n=1}^{\infty} \left|\langle \boldsymbol{x} \mid \boldsymbol{x}_n\rangle\right|^2 \le B\,\|\boldsymbol{x}\|^2 \qquad \forall \boldsymbol{x}\in X.$$

The quantities A *and* B *are* ***frame bounds***.
The quantity A' *is the* ***optimal lower frame bound*** *if*
$A' = \sup\{A \in \mathbb{R}^+ \mid A \text{ is a lower frame bound}\}$.
The quantity B' *is the* ***optimal upper frame bound*** *if*
$B' = \inf\{B \in \mathbb{R}^+ \mid B \text{ is an upper frame bound}\}$.
A frame is a ***tight frame*** *if* $A = B$.
A frame is a ***normalized tight frame*** *(or a* ***Parseval frame****) if* $A = B = 1$.
A frame $\{\boldsymbol{x}_n \mid n\in\mathbb{N}\}$ *is an* ***exact frame*** *if for some* $m \in \mathbb{Z}$, $\{\boldsymbol{x}_n \mid n\in\mathbb{N}\}\setminus\{\boldsymbol{x}_m\}$ *is* NOT *a frame.*

A frame is a *Parseval frame* (Definition H.14) if it satisfies *Parseval's Identity* (Theorem H.9 page 232). All orthonormal bases are Parseval frames (Theorem H.9 page 232); but not all Parseval frames are orthonormal bases.

Definition H.15. *Let* $\{\boldsymbol{x}_n\}$ *be a* ***frame*** *(Definition H.14 page 242) for the* HILBERT SPACE $\boldsymbol{H} \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle\triangle\mid\triangledown\rangle)$. *Let* $\mathbf{S}$ *be an* OPERATOR *on* $\boldsymbol{H}$.

DEF

$\mathbf{S}$ *is a* ***frame operator*** *for* $\{\boldsymbol{x}_n\}$ *if* $\quad \mathbf{S}\mathsf{f}(x) = \sum_{n\in\mathbb{Z}} \langle \mathsf{f} \mid \boldsymbol{x}_n\rangle\, \boldsymbol{x}_n \qquad \forall \mathsf{f}\in\boldsymbol{H}.$

Theorem H.16. [38] *Let* $\mathbf{S}$ *be a* FRAME OPERATOR *(Definition H.15 page 242) of a* FRAME $\{\boldsymbol{x}_n\}$ *(Definition H.14 page 242) for the* HILBERT SPACE $\boldsymbol{H} \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle\triangle\mid\triangledown\rangle)$.

THM

(1). $\mathbf{S}$ *is* INVERTIBLE. *and*

(2). $\mathsf{f}(x) = \sum_{n\in\mathbb{Z}} \left\langle \mathsf{f} \mid \mathbf{S}^{-1}\boldsymbol{x}_n\right\rangle \boldsymbol{x}_n = \sum_{n\in\mathbb{Z}} \langle \mathsf{f} \mid \boldsymbol{x}_n\rangle\, \mathbf{S}^{-1}\boldsymbol{x}_n \quad \forall \mathsf{f} \in \boldsymbol{H}$

Theorem H.17. [39] *Let* $\{\boldsymbol{x}_n \in X \mid n=1,2,\ldots,N\}$ *be a set of vectors in a* HILBERT SPACE $\boldsymbol{H} \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle\triangle\mid\triangledown\rangle)$.

THM

$\{\boldsymbol{x}_n\}$ *is a* FRAME *for* $\operatorname{span}\{\boldsymbol{x}_n\}$.

✎PROOF:

[37] Young (2001) pages 154–155, Christensen (2003) page 88 ⟨Definitions 5.1.1, 5.1.2⟩, Heil (2011) pages 204–205 ⟨Definition 8.2⟩, Jørgensen et al. (2008) page 267 ⟨Definition 12.22⟩, Duffin and Schaeffer (1952) page 343, Daubechies et al. (1986), page 1272

[38] Christensen (2008) pages 100–102 ⟨Theorem 5.1.7⟩

[39] Christensen (2003) page 3

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

1. Upper bound: Proof that there exists B such that $\sum_{n=1}^{N} \left|\langle \boldsymbol{x} \mid \boldsymbol{x}_n\rangle\right|^2 \le B\,\|\boldsymbol{x}\|^2 \quad \forall \boldsymbol{x} \in \boldsymbol{H}$:

$$\sum_{n=1}^{N} \left|\langle \boldsymbol{x} \mid \boldsymbol{x}_n\rangle\right|^2 \le \sum_{n=1}^{N} \langle \boldsymbol{x}_n \mid \boldsymbol{x}_n\rangle \langle \boldsymbol{x} \mid \boldsymbol{x}\rangle \qquad \text{by } \textit{Cauchy-Schwarz inequality} \text{ (Theorem J.2 page 256)}$$

$$= \underbrace{\left\{\sum_{n=1}^{N} \|\boldsymbol{x}_n\|^2\right\}}_{B} \|\boldsymbol{x}\|^2$$

2. Lower bound: Proof that there exists A such that $A\,\|\boldsymbol{x}\|^2 \le \sum_{n=1}^{N} \left|\langle \boldsymbol{x} \mid \boldsymbol{x}_n\rangle\right|^2 \quad \forall \boldsymbol{x} \in \boldsymbol{H}$:

$$\sum_{n=1}^{N} \left|\langle \boldsymbol{x} \mid \boldsymbol{x}_n\rangle\right|^2 = \sum_{n=1}^{N} \left|\left\langle \boldsymbol{x}_n \mid \frac{\boldsymbol{x}}{\|\boldsymbol{x}\|}\right\rangle\right|^2 \|\boldsymbol{x}\|^2$$

$$\ge \underbrace{\left(\inf_{\boldsymbol{y}} \left\{\sum_{n=1}^{N} \left|\langle \boldsymbol{x}_n \mid \boldsymbol{y}\rangle\right|^2 \mid \|\boldsymbol{y}\| = 1\right\}\right)}_{A} \|\boldsymbol{x}\|^2$$

✏

Example H.1. Let $\left(\mathbb{R}^2, +, \cdot, (\mathbb{R}, \dot{+}, \dot{\times}), \langle \triangle \mid \triangledown\rangle\right)$ be an inner product space with $\left\langle \begin{bmatrix} x_1 \\ y_1 \end{bmatrix} \middle| \begin{bmatrix} x_2 \\ y_2 \end{bmatrix} \right\rangle \triangleq x_1x_2 + y_1y_2$. Let $\mathbf{S}$ be the *frame operator* (Definition H.15 page 242) with *inverse* $\mathbf{S}^{-1}$.

EX

Let $N \in \{3, 4, 5, \dots\}$, $\theta \in \mathbb{R}$, and $r \in \mathbb{R}^+$ $(r > 0)$.

Let $\boldsymbol{x}_n \triangleq r\begin{bmatrix} \cos(\theta + 2n\pi/N) \\ \sin(\theta + 2n\pi/N) \end{bmatrix} \quad \forall_{n\in\{0,1,\dots,N-1\}}$.

Then, $(\!(\boldsymbol{x}_0, \boldsymbol{x}_1, \dots, \boldsymbol{x}_{N-1})\!)$ is a **tight frame** for $\mathbb{R}^2$ with *frame bound* $A = \frac{Nr^2}{2}$.



Moreover, $\mathbf{S} = \frac{Nr^2}{2}\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$, $\mathbf{S}^{-1} = \frac{2}{Nr^2}\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$, and $\boldsymbol{v} = \frac{2}{Nr^2}\sum_{n=0}^{N-1} \langle \boldsymbol{v} \mid \boldsymbol{x}_n\rangle\, \boldsymbol{x}_n \quad \forall_{\boldsymbol{v}\in\mathbb{R}^2}$.

✎PROOF:

1. Proof that $(\!(\boldsymbol{x}_0, \boldsymbol{x}_1, \dots, \boldsymbol{x}_{N-1})\!)$ is a *tight frame* with *frame bound* $A = \frac{Nr^2}{2}$: Let $\boldsymbol{v} \triangleq (x, y) \in \mathbb{R}^2$.

$$\sum_{n=0}^{N-1} \left|\langle \boldsymbol{v} \mid \boldsymbol{x}_n\rangle\right|^2 \triangleq \sum_{n=0}^{N-1} \left|\boldsymbol{v}^{\mathbf{H}} r \begin{bmatrix} \cos\left(\theta + \frac{2n\pi}{N}\right) \\ \sin\left(\theta + \frac{2n\pi}{N}\right) \end{bmatrix}\right|^2 \qquad \text{by definitions of } \boldsymbol{v} \text{ of } \langle \boldsymbol{y} \mid \boldsymbol{x}\rangle$$

$$\triangleq \sum_{n=0}^{N-1} r^2 \left|x\cos\left(\theta + \frac{2n\pi}{N}\right) + y\sin\left(\theta + \frac{2n\pi}{N}\right)\right|^2 \qquad \text{by definition of } \boldsymbol{y}^{\mathbf{H}}\boldsymbol{x} \text{ operation}$$

$$= r^2x^2 \sum_{n=0}^{N-1} \cos^2\left(\theta + \frac{2n\pi}{N}\right) + r^2y^2 \sum_{n=0}^{N-1} \sin^2\left(\theta + \frac{2n\pi}{N}\right) + r^2xy \sum_{n=0}^{N-1} \cos\left(\theta + \frac{2n\pi}{N}\right)\sin\left(\theta + \frac{2n\pi}{N}\right)$$

$$= r^2x^2\frac{N}{2} + r^2y^2\frac{N}{2} + r^2xy0 \qquad \text{by Corollary ?? page ??}$$

$$= (x^2 + y^2)\frac{Nr^2}{2} = \left(\frac{Nr^2}{2}\right)\boldsymbol{v}^{\mathbf{H}}\boldsymbol{v} \triangleq \underbrace{\left(\frac{Nr^2}{2}\right)}_{A} \|\boldsymbol{v}\|^2 \qquad \text{by definition of } \|\boldsymbol{v}\|$$

2. Proof that $\mathbf{S} = \frac{Nr^2}{2}\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$:

COPYRIGHT © 2019 DANIEL J. GREENHOE
CC BY-NC-ND

(a) Let $\boldsymbol{e}_1 \triangleq \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ and $\boldsymbol{e}_2 \triangleq \begin{bmatrix} 0 \\ 1 \end{bmatrix}$.

(b) lemma: $\mathbf{S}\boldsymbol{e}_1 = \frac{Nr^2}{2}\boldsymbol{e}_1$. Proof:

$$\begin{aligned}
\mathbf{S}\boldsymbol{e}_1 &= \sum_{n=0}^{N-1} \langle \boldsymbol{e}_1 \,|\, \boldsymbol{x}_n \rangle \, \boldsymbol{x}_n \\
&= \sum_{n=0}^{N-1} r\cos\left(\theta + \frac{2n\pi}{N}\right) r \begin{bmatrix} \cos\left(\theta + \frac{2n\pi}{N}\right) \\ \sin\left(\theta + \frac{2n\pi}{N}\right) \end{bmatrix} = r^2 \sum_{n=0}^{N-1} \begin{bmatrix} \cos^2\left(\theta + \frac{2n\pi}{N}\right) \\ \cos\left(\theta + \frac{2n\pi}{N}\right)\sin\left(\theta + \frac{2n\pi}{N}\right) \end{bmatrix} \\
&= r^2 \begin{bmatrix} N/2 \\ 0 \end{bmatrix} = \frac{Nr^2}{2}\boldsymbol{e}_1 \qquad \text{by } \textit{Summation around unit circle} \text{ (Corollary ?? page ??)}
\end{aligned}$$

(c) lemma: $\mathbf{S}\boldsymbol{e}_2 = \frac{Nr^2}{2}\boldsymbol{e}_2$. Proof:

$$\begin{aligned}
\mathbf{S}\boldsymbol{e}_2 &= \sum_{n=0}^{N-1} \langle \boldsymbol{e}_2 \,|\, \boldsymbol{x}_n \rangle \, \boldsymbol{x}_n \\
&= \sum_{n=0}^{N-1} r\sin\left(\theta + \frac{2n\pi}{N}\right) r \begin{bmatrix} \cos\left(\theta + \frac{2n\pi}{N}\right) \\ \sin\left(\theta + \frac{2n\pi}{N}\right) \end{bmatrix} = r^2 \sum_{n=0}^{N-1} \begin{bmatrix} \sin\left(\theta + \frac{2n\pi}{N}\right)\cos\left(\theta + \frac{2n\pi}{N}\right) \\ \sin^2\left(\theta + \frac{2n\pi}{N}\right) \end{bmatrix} \\
&= r^2 \begin{bmatrix} 0 \\ N/2 \end{bmatrix} = \frac{Nr^2}{2}\boldsymbol{e}_2 \qquad \text{by } \textit{Summation around unit circle} \text{ (Corollary ?? page ??)}
\end{aligned}$$

(d) Complete the proof of item (2) using *Eigendecomposition* $\mathbf{S} = \mathbf{Q}\Lambda\mathbf{Q}^{-1}$:

$$\mathbf{S}\boldsymbol{e}_1 = \frac{Nr^2}{2}\boldsymbol{e}_1 \qquad \text{by (2c) lemma}$$

$$\Longrightarrow \boldsymbol{e}_1 \text{ is an } \textit{eigenvector} \text{ of } \mathbf{S} \text{ with } \textit{eigenvalue } \frac{Nr^2}{2}$$

$$\mathbf{S}\boldsymbol{e}_2 = \frac{Nr^2}{2}\boldsymbol{e}_2 \qquad \text{by (2c) lemma}$$

$$\Longrightarrow \boldsymbol{e}_2 \text{ is an } \textit{eigenvector} \text{ of } \mathbf{S} \text{ with } \textit{eigenvalue } \frac{Nr^2}{2}$$

$$\mathbf{S} = \overbrace{\underbrace{\begin{bmatrix} | & | \\ \boldsymbol{e}_1 & \boldsymbol{e}_2 \\ | & | \end{bmatrix}}_{\mathbf{Q}} \underbrace{\begin{bmatrix} \frac{Nr^2}{2} & 0 \\ 0 & \frac{Nr^2}{2} \end{bmatrix}}_{\Lambda} \underbrace{\begin{bmatrix} | & | \\ \boldsymbol{e}_1 & \boldsymbol{e}_2 \\ | & | \end{bmatrix}^{-1}}_{\mathbf{Q}^{-1}}}^{\textit{Eigendecomposition} \text{ of } \mathbf{S}} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} \frac{Nr^2}{2} & 0 \\ 0 & \frac{Nr^2}{2} \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}^{-1} = \frac{Nr^2}{2}\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

3. Proof that $\mathbf{S}^{-1} = \frac{2}{Nr^2}\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$:

$$\mathbf{S}\mathbf{S}^{-1} = \frac{Nr^2}{2}\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \frac{2}{Nr^2}\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \triangleq \mathbf{I}_2 \qquad \text{by item (2)}$$

$$\mathbf{S}^{-1}\mathbf{S} = \frac{2}{Nr^2}\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \frac{Nr^2}{2}\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \triangleq \mathbf{I}_2 \qquad \text{by item (2)}$$

4. Proof that $\boldsymbol{v} = \frac{2}{Nr^2}\sum_{n=0}^{N-1}(\boldsymbol{v}^{\mathbf{H}}\boldsymbol{x}_n)\boldsymbol{x}_n$:

$$\boldsymbol{v} = \sum_{n=0}^{N-1} \langle \boldsymbol{v} \,|\, \mathbf{S}^{-1}\boldsymbol{x}_n \rangle \, \boldsymbol{x}_n = \sum_{n=0}^{N-1} \left\langle \boldsymbol{v} \,\middle|\, \frac{2}{Nr^2}\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}\boldsymbol{x}_n \right\rangle \boldsymbol{x}_n \qquad \text{by item (3)}$$

$$= \frac{2}{Nr^2}\sum_{n=0}^{N-1} \langle \boldsymbol{v} \,|\, \boldsymbol{x}_n \rangle \, \boldsymbol{x}_n = \frac{2}{Nr^2}\sum_{n=0}^{N-1} (\boldsymbol{v}^{\mathbf{H}}\boldsymbol{x}_n)\boldsymbol{x}_n \qquad \text{by definition of } \langle \boldsymbol{y} \,|\, \boldsymbol{x} \rangle$$

BY-NC-ND Copyright © 2019 Daniel J. Greenhoe

Example H.2 (Peace Frame/Mercedes Frame). [40] Let $\left(\mathbb{R}^2, +, \cdot, (\mathbb{R}, \dot{+}, \dot{\times}), \langle \triangle \mid \triangledown \rangle\right)$ be an inner product space with $\left\langle (x_1, y_1) \mid (x_2, y_2) \right\rangle \triangleq x_1 y_1 + x_2 y_2$. Let $\mathbf{S}$ be the *frame operator* (Definition H.15 page 242) with *inverse* $\mathbf{S}^{-1}$.

EX

Let $\boldsymbol{x}_1 \triangleq \frac{1}{2}\begin{bmatrix} 0 \\ 2 \end{bmatrix}$, $\boldsymbol{x}_2 \triangleq \frac{1}{2}\begin{bmatrix} -\sqrt{3} \\ -1 \end{bmatrix}$, and $\boldsymbol{x}_3 \triangleq \frac{1}{2}\begin{bmatrix} \sqrt{3} \\ -1 \end{bmatrix}$.

Then, $(\!(\boldsymbol{x}_1, \boldsymbol{x}_2, \boldsymbol{x}_3)\!)$ is a **tight frame** for $\mathbb{R}^2$ with *frame bound* $A = \frac{3}{2}$.

Moreover, $\mathbf{S} = \frac{3}{2}\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$, $\quad \mathbf{S}^{-1} = \frac{2}{3}\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$,

and $\quad \boldsymbol{v} = \frac{2}{3}\sum_{n=1}^{3} \langle \boldsymbol{v} \mid \boldsymbol{x}_n \rangle \boldsymbol{x}_n \triangleq \frac{2}{3}\sum_{n=1}^{3} \left(\boldsymbol{v}^{\mathbf{H}} \boldsymbol{x}_n\right) \boldsymbol{x}_n \qquad \forall \boldsymbol{v} \in \mathbb{R}^2.$



✎PROOF:

1. This frame is simply a special case of the frame presented in Example H.1 (page 243) with $r = 1$, $N = 3$, and $\theta = \pi/2$.

2. Let's give it a try! Let $\boldsymbol{v} \triangleq \begin{bmatrix} 1 \\ 1 \end{bmatrix}$.

$$\begin{aligned}
\sum_{n=1}^{3} \left\langle \boldsymbol{v} \mid \mathbf{S}^{-1}\boldsymbol{x}_n \right\rangle \boldsymbol{x}_n &= \frac{2}{3}\sum_{n=1}^{3} \left(\boldsymbol{v}^{\mathbf{H}}\boldsymbol{x}_n\right)\boldsymbol{x}_n && \text{by Example H.1 page 243} \\
&= \left(\boldsymbol{v}^{\mathbf{H}}\boldsymbol{x}_1\right)\boldsymbol{x}_1 + \left(\boldsymbol{v}^{\mathbf{H}}\boldsymbol{x}_2\right)\boldsymbol{x}_2 + \left(\boldsymbol{v}^{\mathbf{H}}\boldsymbol{x}_3\right)\boldsymbol{x}_3 \\
&= \frac{2}{3}\left(\left(\boldsymbol{v}^{\mathbf{H}}\frac{1}{2}\begin{bmatrix} 0 \\ 2 \end{bmatrix}\right)\boldsymbol{x}_1 + \left(\boldsymbol{v}^{\mathbf{H}}\frac{1}{2}\begin{bmatrix} -\sqrt{3} \\ -1 \end{bmatrix}\right)\boldsymbol{x}_2 + \left(\boldsymbol{v}^{\mathbf{H}}\frac{1}{2}\begin{bmatrix} \sqrt{3} \\ -1 \end{bmatrix}\right)\boldsymbol{x}_3\right) \\
&= \frac{2}{3}\cdot\frac{1}{2}\left(\left(\boldsymbol{v}^{\mathbf{H}}\begin{bmatrix} 0 \\ 2 \end{bmatrix}\right)\boldsymbol{x}_1 + \left(\boldsymbol{v}^{\mathbf{H}}\begin{bmatrix} -\sqrt{3} \\ -1 \end{bmatrix}\right)\boldsymbol{x}_2 + \left(\boldsymbol{v}^{\mathbf{H}}\begin{bmatrix} \sqrt{3} \\ -1 \end{bmatrix}\right)\boldsymbol{x}_3\right) \\
&= \frac{1}{3}\left((2)\frac{1}{2}\begin{bmatrix} 0 \\ 2 \end{bmatrix} + \left(-\sqrt{3}-1\right)\frac{1}{2}\begin{bmatrix} -\sqrt{3} \\ -1 \end{bmatrix} + \left(\sqrt{3}-1\right)\frac{1}{2}\begin{bmatrix} \sqrt{3} \\ -1 \end{bmatrix}\right) \\
&= \frac{1}{6}\begin{bmatrix} 2(0) & + & (-\sqrt{3}-1)(-\sqrt{3}) & + & (\sqrt{3}-1)(\sqrt{3}) \\ 2(2) & + & (-\sqrt{3}-1)(-1) & + & (\sqrt{3}-1)(-1) \end{bmatrix} \\
&= \frac{1}{6}\begin{bmatrix} 0 & + & (3+\sqrt{3}) & + & (3-\sqrt{3}) \\ 4 & + & (1+\sqrt{3}) & + & (1-\sqrt{3}) \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \end{bmatrix} \triangleq \boldsymbol{v}
\end{aligned}$$

In Example H.1 (page 243) and Example H.2 (page 245), the frame operator $\mathbf{S}$ and its inverse $\mathbf{S}^{-1}$ were computed. In general however, it is not always necessary or even possible to compute these, as illustrated in Example H.3 (next).

Example H.3. [41] Let $\left(\mathbb{R}^2, +, \cdot, (\mathbb{R}, \dot{+}, \dot{\times}), \langle \triangle \mid \triangledown \rangle\right)$ be an inner product space with $\left\langle (x_1, y_1) \mid (x_2, y_2) \right\rangle \triangleq x_1 y_1 + x_2 y_2$. Let $\mathbf{S}$ be the *frame operator* (Definition H.15 page 242) with *inverse* $\mathbf{S}^{-1}$.

[40] Heil (2011) pages 204–205 ($r = 1$ case), Byrne (2005) page 80 ($r = 1$ case), Han et al. (2007) page 91 (Example 3.9, $r = \sqrt{2/3}$ case)

[41] Christensen (2003) pages 7–8 (?)

COPYRIGHT © 2019 DANIEL J. GREENHOE

EX

Let $\boldsymbol{p}$ and $\boldsymbol{q}$ be *orthonormal* vectors in $\boldsymbol{X} \triangleq \operatorname{span}\{\boldsymbol{p}, \boldsymbol{q}\}$.
Let $\boldsymbol{x}_1 \triangleq \boldsymbol{p}$, $\boldsymbol{x}_2 \triangleq \boldsymbol{p}+\boldsymbol{q}$, and $\boldsymbol{x}_3 \triangleq \boldsymbol{p}-\boldsymbol{q}$.
Then, $\{\boldsymbol{x}_1, \boldsymbol{x}_2, , \boldsymbol{x}_3\}$ is a **frame** for $\boldsymbol{X}$ with *frame bounds* $A = 0$ and $B = 5$.



Moreover,
$\mathbf{S}^{-1}\boldsymbol{x}_1 = \tfrac{1}{3}\boldsymbol{p}$ and
$\mathbf{S}^{-1}\boldsymbol{x}_2 = \tfrac{1}{3}\boldsymbol{p} + \tfrac{1}{2}\boldsymbol{q}$ and
$\mathbf{S}^{-1}\boldsymbol{x}_3 = \tfrac{1}{3}\boldsymbol{p} - \tfrac{1}{2}\boldsymbol{q}$.

✎PROOF:

1. Proof that $(\!(\boldsymbol{x}_1, \boldsymbol{x}_2, \boldsymbol{x}_3)\!)$ is a *frame* with *frame bounds* $A = 0$ and $B = 5$:

$$\begin{aligned}
\sum_{n=1}^{3} \left|\langle \boldsymbol{v} \,|\, \boldsymbol{x}_n\rangle\right|^2 &\triangleq |\langle \boldsymbol{v} \,|\, \boldsymbol{p}\rangle|^2 + |\langle \boldsymbol{v} \,|\, \boldsymbol{p}+\boldsymbol{q}\rangle|^2 + |\langle \boldsymbol{v} \,|\, \boldsymbol{p}-\boldsymbol{q}\rangle|^2 && \text{by definitions of } \boldsymbol{x}_1, \boldsymbol{x}_2, \text{ and } \boldsymbol{x}_3 \\
&= |\langle \boldsymbol{v} \,|\, \boldsymbol{p}\rangle|^2 + |\langle \boldsymbol{v} \,|\, \boldsymbol{p}\rangle + \langle \boldsymbol{v} \,|\, \boldsymbol{q}\rangle|^2 + |\langle \boldsymbol{v} \,|\, \boldsymbol{p}\rangle - \langle \boldsymbol{v} \,|\, \boldsymbol{q}\rangle|^2 && \text{by } \textit{additivity} \text{ of } \langle \triangle \,|\, \triangledown \rangle \text{ (Definition J.1 page 255)} \\
&= |\langle \boldsymbol{v} \,|\, \boldsymbol{p}\rangle|^2 + \left(|\langle \boldsymbol{v} \,|\, \boldsymbol{p}\rangle|^2 + |\langle \boldsymbol{v} \,|\, \boldsymbol{q}\rangle|^2 + \langle \boldsymbol{v} \,|\, \boldsymbol{p}\rangle \langle \boldsymbol{v} \,|\, \boldsymbol{q}\rangle^* + \langle \boldsymbol{v} \,|\, \boldsymbol{q}\rangle \langle \boldsymbol{v} \,|\, \boldsymbol{p}\rangle^*\right) \\
&\quad + \left(|\langle \boldsymbol{v} \,|\, \boldsymbol{p}\rangle|^2 + |\langle \boldsymbol{v} \,|\, \boldsymbol{q}\rangle|^2 - \langle \boldsymbol{v} \,|\, \boldsymbol{p}\rangle \langle \boldsymbol{v} \,|\, \boldsymbol{q}\rangle^* - \langle \boldsymbol{v} \,|\, \boldsymbol{q}\rangle \langle \boldsymbol{v} \,|\, \boldsymbol{p}\rangle^*\right) \\
&= 3|\langle \boldsymbol{v} \,|\, \boldsymbol{p}\rangle|^2 + 2|\langle \boldsymbol{v} \,|\, \boldsymbol{q}\rangle|^2 \\
&\boxed{\le} 3\,\|\boldsymbol{v}\|\,\|\boldsymbol{p}\| + 2\,\|\boldsymbol{v}\|\,\|\boldsymbol{q}\| && \text{by } \textit{CS Inequality} \text{ (Theorem J.2 page 256)} \\
&= \|\boldsymbol{v}\|\,(3\,\|\boldsymbol{p}\| + 2\,\|\boldsymbol{q}\|) \\
&= \boxed{5}\,\|\boldsymbol{v}\| && \text{by } \textit{orthonormality} \text{ of } \boldsymbol{p} \text{ and } \boldsymbol{q}
\end{aligned}$$

2. lemma: $\mathbf{S}\boldsymbol{p} = 3\boldsymbol{p}$, $\mathbf{S}\boldsymbol{q} = 2\boldsymbol{q}$, $\mathbf{S}^{-1}\boldsymbol{p} = \frac{1}{3}\boldsymbol{p}$, and $\mathbf{S}^{-1}\boldsymbol{q} = \frac{1}{2}\boldsymbol{q}$. Proof:

$$\begin{aligned}
\mathbf{S}\boldsymbol{p} &\triangleq \sum_{n=1}^{3} \langle \boldsymbol{p} \,|\, \boldsymbol{x}_n\rangle\, \boldsymbol{x}_n \\
&= \langle \boldsymbol{p} \,|\, \boldsymbol{p}\rangle\, \boldsymbol{p} + \langle \boldsymbol{p} \,|\, \boldsymbol{p}+\boldsymbol{q}\rangle\,(\boldsymbol{p}+\boldsymbol{q}) + \langle \boldsymbol{p} \,|\, \boldsymbol{p}-\boldsymbol{q}\rangle\,(\boldsymbol{p}-\boldsymbol{q}) \\
&= (1)\boldsymbol{p} + (1+0)(\boldsymbol{p}+\boldsymbol{q}) + (1-0)(\boldsymbol{p}-\boldsymbol{q}) \\
&= 3\boldsymbol{p} \\
\implies \mathbf{S}^{-1}\boldsymbol{p} &= \frac{1}{3}\boldsymbol{p} \\
\mathbf{S}\boldsymbol{q} &\triangleq \sum_{n=1}^{3} \langle \boldsymbol{q} \,|\, \boldsymbol{x}_n\rangle\, \boldsymbol{x}_n \\
&= \langle \boldsymbol{q} \,|\, \boldsymbol{p}\rangle\, \boldsymbol{p} + \langle \boldsymbol{q} \,|\, \boldsymbol{p}+\boldsymbol{q}\rangle\,(\boldsymbol{p}+\boldsymbol{q}) + \langle \boldsymbol{q} \,|\, \boldsymbol{p}-\boldsymbol{q}\rangle\,(\boldsymbol{p}-\boldsymbol{q}) \\
&= (0)\boldsymbol{q} + (0+1)(\boldsymbol{p}+\boldsymbol{q}) + (0-1)(\boldsymbol{p}-\boldsymbol{q}) \\
&= 2\boldsymbol{q} \\
\implies \mathbf{S}^{-1}\boldsymbol{q} &= \frac{1}{2}\boldsymbol{q}
\end{aligned}$$

3. Remark: Without knowing $\boldsymbol{p}$ and $\boldsymbol{q}$, from (2) lemma it follows that it is not possible to compute $\mathbf{S}$ or $\mathbf{S}^{-1}$ explicitly.

4. Proof that $\mathbf{S}^{-1}\boldsymbol{x}_1 = \tfrac{1}{3}\boldsymbol{p}$, $\mathbf{S}^{-1}\boldsymbol{x}_2 = \tfrac{1}{3}\boldsymbol{p} + \tfrac{1}{2}\boldsymbol{q}$ and $\mathbf{S}^{-1}\boldsymbol{x}_3 = \tfrac{1}{3}\boldsymbol{p} - \tfrac{1}{2}\boldsymbol{q}$:

$$\begin{aligned}
\mathbf{S}^{-1}\boldsymbol{x}_1 &\triangleq \mathbf{S}^{-1}\boldsymbol{p} && \text{by definition of } \boldsymbol{x}_1 \\
&= \frac{1}{3}\boldsymbol{p} && \text{by (2) lemma} \\
\mathbf{S}^{-1}\boldsymbol{x}_2 &\triangleq \mathbf{S}^{-1}(\boldsymbol{p}+\boldsymbol{q}) && \text{by definition of } \boldsymbol{x}_2 \\
&= \frac{1}{3}\boldsymbol{p} + \frac{1}{2}\boldsymbol{q} && \text{by (2) lemma}
\end{aligned}$$

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

$$\begin{aligned}
\mathbf{S}^{-1}\boldsymbol{x}_3 &\triangleq \mathbf{S}^{-1}(\boldsymbol{p}-\boldsymbol{q}) && \text{by definition of } \boldsymbol{x}_2\\
&= \frac{1}{3}\boldsymbol{p} - \frac{1}{2}\boldsymbol{q} && \text{by (2) lemma}
\end{aligned}$$

5. Check that $\boldsymbol{v} = \sum_n \langle \boldsymbol{v} \mid \boldsymbol{x}_n \rangle \boldsymbol{x}_n = \langle \boldsymbol{v} \mid \boldsymbol{p} \rangle \boldsymbol{p} + \langle \boldsymbol{v} \mid \boldsymbol{q} \rangle \boldsymbol{q}$:

$$\begin{aligned}
\boldsymbol{v} &= \sum_{n=1}^{3} \left\langle \boldsymbol{v} \mid \mathbf{S}^{-1}\boldsymbol{x}_n \right\rangle \boldsymbol{x}_n\\
&= \left\langle \boldsymbol{v} \mid \mathbf{S}^{-1}\boldsymbol{p} \right\rangle \boldsymbol{p} + \left\langle \boldsymbol{v} \mid \mathbf{S}^{-1}(\boldsymbol{p}+\boldsymbol{q}) \right\rangle (\boldsymbol{p}+\boldsymbol{q}) + \left\langle \boldsymbol{v} \mid \mathbf{S}^{-1}(\boldsymbol{p}-\boldsymbol{q}) \right\rangle (\boldsymbol{p}-\boldsymbol{q})\\
&= \left\langle \boldsymbol{v} \mid \frac{1}{3}\boldsymbol{p} \right\rangle \boldsymbol{p} + \left\langle \boldsymbol{v} \mid \frac{1}{3}\boldsymbol{p} + \frac{1}{2}\boldsymbol{q} \right\rangle (\boldsymbol{p}+\boldsymbol{q}) + \left\langle \boldsymbol{v} \mid \frac{1}{3}\boldsymbol{p} - \frac{1}{2}\boldsymbol{q}) \right\rangle (\boldsymbol{p}-\boldsymbol{q})\\
&= \left(\frac{1}{3}+\frac{1}{3}+\frac{1}{3}\right) \langle \boldsymbol{v} \mid \boldsymbol{p} \rangle \boldsymbol{p} + \left(\frac{1}{3}-\frac{1}{3}\right) \langle \boldsymbol{v} \mid \boldsymbol{p} \rangle \boldsymbol{q} + \left(\frac{1}{2}-\frac{1}{2}\right) \langle \boldsymbol{v} \mid \boldsymbol{q} \rangle \boldsymbol{p} + \left(\frac{1}{2}+\frac{1}{2}\right) \langle \boldsymbol{v} \mid \boldsymbol{q} \rangle \boldsymbol{q}\\
&= \langle \boldsymbol{v} \mid \boldsymbol{p} \rangle \boldsymbol{p} + \langle \boldsymbol{v} \mid \boldsymbol{q} \rangle \boldsymbol{q}
\end{aligned}$$

Copyright © 2019 Daniel J. Greenhoe
BY-NC-ND

Copyright © 2019 Daniel J. Greenhoe

APPENDIX I

NORMED ALGEBRAS

I.1 Algebras

All *linear spaces* (Definition G.1 page 207) are equipped with an operation by which vectors in the spaces can be added together. Linear spaces also have an operation that allows a scalar and a vector to be "multiplied" together. But linear spaces in general have no operation that allows two vectors to be multiplied together. A linear space together with such an operator is an **algebra**.[1]

There are many many possible algebras—many more than one can shake a stick at, as indicated by Michiel Hazewinkel in his book, *Handbook of Algebras*: "Algebra, as we know it today (2005), consists of many different ideas, concepts and results. A reasonable estimate of the number of these different items would be somewhere between 50,000 and 200,000. Many of these have been named and many more could (and perhaps should) have a "name" or other convenient designation."[2]

Definition I.1. [3] *Let* $\boldsymbol{A}$ *be an* ALGEBRA.

DEF *An algebra* $\boldsymbol{A}$ *is* ***unital*** *if* $\exists u \in \boldsymbol{A}$ such that $ux = xu = x \quad \forall x \in \boldsymbol{A}$

Definition I.2. [4] *Let* $\boldsymbol{A}$ *be an* UNITAL ALGEBRA *(Definition I.1 page 249) with unit* e.

DEF

The ***spectrum*** *of* $x \in \boldsymbol{A}$ *is* $\sigma(x) \triangleq \{\lambda \in \mathbb{C} | \lambda e - x \text{ is not invertible}\}$.

The ***resolvent*** *of* $x \in \boldsymbol{A}$ *is* $\rho_x(\lambda) \triangleq (\lambda e - x)^{-1} \quad \forall \lambda \notin \sigma(x)$.

The ***spectral radius*** *of* $x \in \boldsymbol{A}$ *is* $r(x) \triangleq \sup\{|\lambda| | \lambda \in \sigma(x)\}$.

[1] Fuchs (1995) page 2
[2] Hazewinkel (2000) page v
[3] Folland (1995) page 1
[4] Folland (1995) pages 3–4

I.2 Star-Algebras

Definition I.3. [5] *Let $\boldsymbol{A}$ be an* ALGEBRA.

DEF

*The pair $(\boldsymbol{A}, *)$ is a $*$-**algebra**, or **star-algebra**, if*

1. $(x+y)^* = x^* + y^* \quad \forall x,y \in \boldsymbol{A}$ (DISTRIBUTIVE) *and*
2. $(\alpha x)^* = \bar{\alpha} x^* \quad \forall x \in \boldsymbol{A}, \alpha \in \mathbb{C}$ (CONJUGATE LINEAR) *and*
3. $(xy)^* = y^* x^* \quad \forall x,y \in \boldsymbol{A}$ (ANTIAUTOMORPHIC) *and*
4. $x^{**} = x \quad \forall x \in \boldsymbol{A}$ (INVOLUTORY)

*The operator $*$ is called an **involution** on the algebra $\boldsymbol{A}$.*

Proposition I.1. [6] *Let $(\boldsymbol{A}, *)$ be an* UNITAL $*$-ALGEBRA.

PRP

$$x \text{ is invertible} \quad \Longrightarrow \quad \begin{cases} 1. & x^* \text{ is INVERTIBLE} \quad \forall x \in \boldsymbol{A} \quad \text{and} \\ 2. & \left(x^*\right)^{-1} = \left(x^{-1}\right)^* \quad \forall x \in \boldsymbol{A} \end{cases}$$

✎PROOF: Let e be the unit element of $(\boldsymbol{A}, *)$.

1. Proof that $e^* = e$:

$$\begin{aligned}
x\,e^* &= \left(x\,e^*\right)^{**} && \text{by } \textit{involutory} \text{ property of } * && \text{(Definition I.3 page 250)} \\
&= \left(x^*\,e^{**}\right)^* && \text{by } \textit{antiautomorphic} \text{ property of } * && \text{(Definition I.3 page 250)} \\
&= \left(x^*\,e\right)^* && \text{by } \textit{involutory} \text{ property of } * && \text{(Definition I.3 page 250)} \\
&= \left(x^*\right)^* && \text{by definition of } e \\
&= x && \text{by } \textit{involutory} \text{ property of } * && \text{(Definition I.3 page 250)} \\
e^*\,x &= \left(e^*\,x\right)^{**} && \text{by } \textit{involutory} \text{ property of } * && \text{(Definition I.3 page 250)} \\
&= \left(e^{**}\,x^*\right)^* && \text{by } \textit{antiautomorphic} \text{ property of } * && \text{(Definition I.3 page 250)} \\
&= \left(e\,x^*\right)^* && \text{by } \textit{involutory} \text{ property of } * && \text{(Definition I.3 page 250)} \\
&= \left(x^*\right)^* && \text{by definition of } e \\
&= x && \text{by } \textit{involutory} \text{ property of } * && \text{(Definition I.3 page 250)}
\end{aligned}$$

2. Proof that $\left(x^*\right)^{-1} = \left(x^{-1}\right)^*$:

$$\begin{aligned}
\left(x^{-1}\right)^* \left(x^*\right) &= \left[x\left(x^{-1}\right)\right]^* && \text{by } \textit{antiautomorphic} \text{ and } \textit{involution} \text{ properties of } * && \text{(Definition I.3 page 250)} \\
&= e^* \\
&= e && \text{by item (1) page 250} \\
\left(x^*\right)\left(x^{-1}\right)^* &= \left[x^{-1}\,x\right]^* && \text{by } \textit{antiautomorphic} \text{ and } \textit{involution} \text{ properties of } * && \text{(Definition I.3 page 250)} \\
&= e^* \\
&= e && \text{by item (1) page 250}
\end{aligned}$$

✏

Definition I.4. [7] *Let $(\boldsymbol{A}, \|\cdot\|)$ be a* $*$-ALGEBRA *(Definition I.3 page 250).*

DEF

- *An element $x \in \boldsymbol{A}$ is **hermitian** or **self-adjoint** if $x^* = x$.*
- *An element $x \in \boldsymbol{A}$ is **normal** if $xx^* = x^*x$.*
- *An element $x \in \boldsymbol{A}$ is a **projection** if $xx = x$ (INVOLUTORY) and $x^* = x$ (HERMITIAN).*

[5] Rickart (1960), page 178, Gelfand and Naimark (1964), page 241
[6] Folland (1995) page 5
[7] Rickart (1960), page 178, Gelfand and Naimark (1964), page 242

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

Theorem I.1. [8] *Let* $(\boldsymbol{A}, \|\cdot\|)$ *be a* $*$-ALGEBRA *(Definition I.3 page 250)*.

THM

$$\underbrace{x = x^* \text{ and } y = y^*}_{x \text{ and } y \text{ are HERMITIAN}} \quad \Longrightarrow \quad \begin{cases} x + y = (x+y)^* & (x+y \text{ is self adjoint}) \\ x^* = (x^*)^* & (x^* \text{ is self adjoint}) \\ \underbrace{xy = (xy)^*}_{(xy) \text{ is HERMITIAN}} \iff \underbrace{xy = yx}_{\text{commutative}} \end{cases}$$

✎PROOF:

$$\begin{aligned} (x+y)^* &= x^* + y^* && \text{by } \textit{distributive} \text{ property of } * && \text{(Definition I.3 page 250)} \\ &= x + y && \text{by left hypothesis} \end{aligned}$$

$$\left(x^*\right)^* = x \qquad \text{by } \textit{involutory} \text{ property of } * \qquad \text{(Definition I.3 page 250)}$$

Proof that $xy = (xy)^* \implies xy = yx$

$$\begin{aligned} xy &= (xy)^* && \text{by left hypothesis} \\ &= y^* x^* && \text{by } \textit{antiautomorphic} \text{ property of } * && \text{(Definition I.3 page 250)} \\ &= yx && \text{by left hypothesis} \end{aligned}$$

Proof that $xy = (xy)^* \impliedby xy = yx$

$$\begin{aligned} (xy)^* &= (yx)^* && \text{by left hypothesis} \\ &= x^* y^* && \text{by } \textit{antiautomorphic} \text{ property of } * && \text{(Definition I.3 page 250)} \\ &= xy && \text{by left hypothesis} \end{aligned}$$

✏

Definition I.5 (Hermitian components). [9] *Let* $(\boldsymbol{A}, \|\cdot\|)$ *be a* $*$-ALGEBRA *(Definition I.3 page 250)*.

DEF

The ***real part*** *of* x *is defined as* $\quad \mathbf{R}_{\mathrm{e}} x \triangleq \frac{1}{2}\left(x + x^*\right)$

The ***imaginary part*** *of* x *is defined as* $\quad \mathbf{I}_{\mathrm{m}} x \triangleq \frac{1}{2i}\left(x - x^*\right)$

Theorem I.2. [10] *Let* $(\boldsymbol{A}, *)$ *be a* $*$-ALGEBRA *(Definition I.3 page 250)*.

THM

$$\begin{aligned} \mathbf{R}_{\mathrm{e}} x &= \left(\mathbf{R}_{\mathrm{e}} x\right)^* && \forall x \in \boldsymbol{A} && (\mathbf{R}_{\mathrm{e}} x \text{ is HERMITIAN}) \\ \mathbf{I}_{\mathrm{m}} x &= \left(\mathbf{I}_{\mathrm{m}} x\right)^* && \forall x \in \boldsymbol{A} && (\mathbf{I}_{\mathrm{m}} x \text{ is HERMITIAN}) \end{aligned}$$

✎PROOF:

$$\begin{aligned} \left(\mathbf{R}_{\mathrm{e}} x\right)^* &= \left(\frac{1}{2}\left(x + x^*\right)\right)^* && \text{by definition of } \Re && \text{(Definition I.5 page 251)} \\ &= \frac{1}{2}\left(x^* + x^{**}\right) && \text{by } \textit{distributive} \text{ property of } * && \text{(Definition I.3 page 250)} \\ &= \frac{1}{2}\left(x^* + x\right) && \text{by } \textit{involutory} \text{ property of } * && \text{(Definition I.3 page 250)} \\ &= \mathbf{R}_{\mathrm{e}} x && \text{by definition of } \Re && \text{(Definition I.5 page 251)} \\ \left(\mathbf{I}_{\mathrm{m}} x\right)^* &= \left(\frac{1}{2i}\left(x - x^*\right)\right)^* && \text{by definition of } \Im && \text{(Definition I.5 page 251)} \end{aligned}$$

[8] Michel and Herget (1993) page 429
[9] Michel and Herget (1993) page 430, Rickart (1960), page 179, Gelfand and Naimark (1964), page 242
[10] Michel and Herget (1993) page 430, Halmos (1998) page 42

COPYRIGHT © 2019 DANIEL J. GREENHOE

$$= \frac{1}{2i}\left(x^* - x^{**}\right) \quad \text{by } \textit{distributive} \text{ property of } * \quad \text{(Definition I.3 page 250)}$$
$$= \frac{1}{2i}\left(x^* - x\right) \quad \text{by } \textit{involutory} \text{ property of } * \quad \text{(Definition I.3 page 250)}$$
$$= \mathbf{I}_{\mathsf{m}} x \quad \text{by definition of } \mathfrak{I} \quad \text{(Definition I.5 page 251)}$$

✏

Theorem I.3 (Hermitian representation). [11] *Let* $(\boldsymbol{A}, *)$ *be a* $*$-ALGEBRA *(Definition I.3 page 250)*.

THM $a = x + iy \iff x = \mathbf{R}_{\mathsf{e}} a$ *and* $y = \mathbf{I}_{\mathsf{m}} a$

✎PROOF:

Proof that $a = x + iy \implies x = \mathbf{R}_{\mathsf{e}} a$ and $y = \mathbf{I}_{\mathsf{m}} a$:

$$a = x + iy \quad \text{by left hypothesis}$$
$$\implies a^* = (x+iy)^* \quad \text{by definition of } \textit{adjoint} \quad \text{(Definition I.4 page 250)}$$
$$= x^* - iy^* \quad \text{by } \textit{distributive} \text{ property of } * \quad \text{(Definition I.3 page 250)}$$
$$= x - iy \quad \text{by Theorem I.2 page 251}$$
$$\implies x = a - iy \quad \text{by solving for } x \text{ in } a = x + iy \text{ equation}$$
$$x = a^* + iy \quad \text{by solving for } x \text{ in } a^* = x - iy \text{ equation}$$
$$\implies x + x = a + a^* \quad \text{by adding previous 2 equations}$$
$$\implies 2x = a + a^* \quad \text{by solving for } x \text{ in previous equation}$$
$$\implies x = \frac{1}{2}\left(a + a^*\right)$$
$$= \mathbf{R}_{\mathsf{e}} a \quad \text{by definition of } \mathfrak{R} \quad \text{(Definition I.5 page 251)}$$

$$iy = a - x \quad \text{by solving for } iy \text{ in } a = x + iy \text{ equation}$$
$$iy = -a^* + x \quad \text{by solving for } iy \text{ in } a = x + iy \text{ equation}$$
$$\implies iy + iy = a - a^* \quad \text{by adding previous 2 equations}$$
$$\implies y = \frac{1}{2i}\left(a - a^*\right) \quad \text{by solving for } iy \text{ in previous equations}$$
$$= \mathbf{I}_{\mathsf{m}} a \quad \text{by definition of } \mathfrak{I} \quad \text{(Definition I.5 page 251)}$$

Proof that $a = x + iy \impliedby x = \mathbf{R}_{\mathsf{e}} a$ and $y = \mathbf{I}_{\mathsf{m}} a$:

$$x + iy = \mathbf{R}_{\mathsf{e}} a + i\,\mathbf{I}_{\mathsf{m}} a \quad \text{by right hypothesis}$$
$$= \underbrace{\frac{1}{2}\left(a + a^*\right)}_{\mathbf{R}_{\mathsf{e}} a} + i\underbrace{\frac{1}{2i}\left(a - a^*\right)}_{\mathbf{I}_{\mathsf{m}} a} \quad \text{by definition of } \mathfrak{R} \text{ and } \mathfrak{I} \quad \text{(Definition I.5 page 251)}$$
$$= \left(\frac{1}{2}a + \frac{1}{2}a\right) + \cancelto{0}{\left(\frac{1}{2}a^* - \frac{1}{2}a^*\right)}$$
$$= a$$

✏

[11] Michel and Herget (1993) page 430, Rickart (1960), page 179, Gelfand and Neumark (1943b), page 7

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

I.3 Normed Algebras

Definition I.6. [12] *Let $\boldsymbol{A}$ be an algebra.*

DEF

The pair $(\boldsymbol{A}, \|\cdot\|)$ is a ***normed algebra*** *if*

$$\|xy\| \le \|x\|\,\|y\| \qquad \forall x, y \in \boldsymbol{A} \qquad \textit{(multiplicative condition)}$$

A normed algebra $(\boldsymbol{A}, \|\cdot\|)$ is a ***Banach algebra*** *if $(\boldsymbol{A}, \|\cdot\|)$ is also a Banach space.*

Proposition I.2.

PRP

$$(\boldsymbol{A}, \|\cdot\|) \text{ is a normed algebra} \quad \Longrightarrow \quad \text{multiplication is } \textbf{continuous} \text{ in } (\boldsymbol{A}, \|\cdot\|)$$

✎PROOF:

1. Define $\mathsf{f}(x) \triangleq zx$. That is, the function f represents multiplication of x times some arbitrary value z.
2. Let $\delta \triangleq \|x - y\|$ and $\epsilon \triangleq \|\mathsf{f}(x) - \mathsf{f}(y)\|$.
3. To prove that multiplication (f) is *continuous* with respect to the metric generated by $\|\cdot\|$, we have to show that we can always make ϵ arbitrarily small for some $\delta > 0$.
4. And here is the proof that multiplication is indeed continuous in $(\boldsymbol{A}, \|\cdot\|)$:

$$\begin{aligned}
\|\mathsf{f}(x) - \mathsf{f}(y)\| &\triangleq \|zx - zy\| && \text{by definition of } \mathsf{f} && \text{(item (1) page 253)}\\
&= \|z(x-y)\| \\
&\le \|z\|\,\|x-y\| && \text{by definition of } \textit{normed algebra} && \text{(Definition I.6 page 253)}\\
&\triangleq \|z\|\,\delta && \text{by definition of } \delta && \text{(item (2) page 253)}\\
&\le \epsilon && \text{for some value of } \delta > 0
\end{aligned}$$

✎

Theorem I.4 (Gelfand-Mazur Theorem)**.** [13] *Let $\mathbb{C}$ be the field of complex numbers.*

THM

$$\left.\begin{array}{l}(\boldsymbol{A}, \|\cdot\|) \text{ is a Banach algebra}\\ \text{every nonzero } x \in \boldsymbol{A} \text{ is invertible}\end{array}\right\} \quad \Longrightarrow \quad \boldsymbol{A} \equiv \mathbb{C} \quad (\boldsymbol{A} \text{ is isomorphic to } \mathbb{C})$$

I.4 C* Algebras

Definition I.7. [14]

DEF

The triple $(\boldsymbol{A}, \|\cdot\|, *)$ is a C^* ***algebra*** *if*

1. $(\boldsymbol{A}, \|\cdot\|)$ *is a Banach algebra* *and*
2. $(\boldsymbol{A}, *)$ *is a $*$-algebra* *and*
3. $\left\|x^*x\right\| = \|x\|^2 \quad \forall x \in \boldsymbol{A}$.

A C^* ***algebra*** *$(\boldsymbol{A}, \|\cdot\|, *)$ is also called a* ***C star algebra****.*

[12] Rickart (1960), page 2, Berberian (1961) page 103 ⟨Theorem IV.9.2⟩
[13] Folland (1995) page 4, Mazur (1938) ⟨(statement)⟩, Gelfand (1941) ⟨(proof)⟩
[14] Folland (1995) page 1, Gelfand and Naimark (1964), page 241, Gelfand and Neumark (1943a), Gelfand and Neumark (1943b)

COPYRIGHT © 2019 DANIEL J. GREENHOE

Theorem I.5. [15] *Let $\boldsymbol{A}$ be an algebra.*

THM $(\boldsymbol{A}, \|\cdot\|, *)$ *is a* C^* ***algebra*** $\Longrightarrow$ $\|x^*\| = \|x\|$

✎PROOF:

$$\begin{aligned}
\|x\| &= \frac{1}{\|x\|}\ \|x\|^2 \\
&= \frac{1}{\|x\|}\ \|x^*x\| && \text{by definition of } C^*\text{-}\textit{algebra} && \text{(Definition I.7 page 253)} \\
&\le \frac{1}{\|x\|}\ \|x^*\|\,\|x\| && \text{by definition of } \textit{normed algebra} && \text{(Definition I.6 page 253)} \\
&= \|x^*\| \\
\|x^*\| &\le \|x^{**}\| && \text{by previous result} \\
&= \|x\| && \text{by } \textit{involution} \text{ property of } * && \text{(Definition I.3 page 250)}
\end{aligned}$$

✏

[15] Folland (1995) page 1, Gelfand and Neumark (1943b), page 4, Gelfand and Neumark (1943a)

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

APPENDIX J

INNER PRODUCT SPACES

J.1 Definition and basic results

Definition J.1. [1] *Let* $\boldsymbol{\Omega} \triangleq \big(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times})\big)$ *be a* LINEAR SPACE *(Definition G.1 page 207)*.

DEF

A FUNCTIONAL $\langle \triangle \mid \triangledown \rangle \in \mathbb{F}^{X \times X}$ *is an* ***inner product*** *on* $\boldsymbol{\Omega}$ *if*

1.	$\langle \alpha \boldsymbol{x} \mid \boldsymbol{y} \rangle = \alpha \langle \boldsymbol{x} \mid \boldsymbol{y} \rangle$	$\forall \boldsymbol{x}, \boldsymbol{y} \in X, \forall \alpha \in \mathbb{C}$	(HOMOGENEOUS)	*and*
2.	$\langle \boldsymbol{x} + \boldsymbol{y} \mid \boldsymbol{u} \rangle = \langle \boldsymbol{x} \mid \boldsymbol{u} \rangle + \langle \boldsymbol{y} \mid \boldsymbol{u} \rangle$	$\forall \boldsymbol{x}, \boldsymbol{y}, \boldsymbol{u} \in X$	(ADDITIVE)	*and*
3.	$\langle \boldsymbol{x} \mid \boldsymbol{y} \rangle = \langle \boldsymbol{y} \mid \boldsymbol{x} \rangle^*$	$\forall \boldsymbol{x}, \boldsymbol{y} \in X$	(CONJUGATE SYMMETRIC).	*and*
4.	$\langle \boldsymbol{x} \mid \boldsymbol{x} \rangle \geq 0$	$\forall \boldsymbol{x} \in X$	(NON-NEGATIVE)	*and*
5.	$\langle \boldsymbol{x} \mid \boldsymbol{x} \rangle = 0 \iff \boldsymbol{x} = \mathbb{0}$	$\forall \boldsymbol{x} \in X$	(NON-ISOTROPIC)	

An inner product is also called a ***scalar product****.*

The tuple $\big(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle \mid \triangledown \rangle\big)$ *is called an* ***inner product space****.*

Theorem J.1. [2] *Let* $\boldsymbol{\Omega} \triangleq \big(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle \mid \triangledown \rangle\big)$ *be a* LINEAR SPACE *(Definition G.1 page 207)*.

THM

1.	$\langle \boldsymbol{x} \mid \boldsymbol{y} + \boldsymbol{z} \rangle = \langle \boldsymbol{x} \mid \boldsymbol{y} \rangle + \langle \boldsymbol{x} \mid \boldsymbol{z} \rangle$	$\forall \boldsymbol{x}, \boldsymbol{y}, \boldsymbol{z} \in X$	
2.	$\langle \boldsymbol{x} \mid \alpha \boldsymbol{y} \rangle = \alpha^* \langle \boldsymbol{x} \mid \boldsymbol{y} \rangle$	$\forall \boldsymbol{x}, \boldsymbol{y} \in X, \alpha \in \mathbb{F}$	
3.	$\langle \boldsymbol{x} \mid \mathbb{0} \rangle = \langle \mathbb{0} \mid \boldsymbol{x} \rangle = 0$	$\forall \boldsymbol{x} \in X$	
4.	$\langle \boldsymbol{x} - \boldsymbol{y} \mid \boldsymbol{z} \rangle = \langle \boldsymbol{x} \mid \boldsymbol{z} \rangle - \langle \boldsymbol{y} \mid \boldsymbol{z} \rangle$	$\forall \boldsymbol{x}, \boldsymbol{y}, \boldsymbol{z} \in X$	
5.	$\langle \boldsymbol{x} \mid \boldsymbol{y} - \boldsymbol{z} \rangle = \langle \boldsymbol{x} \mid \boldsymbol{y} \rangle - \langle \boldsymbol{x} \mid \boldsymbol{z} \rangle$	$\forall \boldsymbol{x}, \boldsymbol{y}, \boldsymbol{z} \in X$	
6.	$\langle \boldsymbol{x} \mid \boldsymbol{z} \rangle = \langle \boldsymbol{y} \mid \boldsymbol{z} \rangle$	$\forall \boldsymbol{z} \in X \neq \{\mathbb{0}\}$	$\iff \boldsymbol{x} = \boldsymbol{y}$
7.	$\langle \boldsymbol{x} \mid \boldsymbol{y} \rangle = 0$	$\forall \boldsymbol{x} \in X$	$\iff \boldsymbol{y} = \mathbb{0}$

✎PROOF:

$\langle \boldsymbol{x} \mid \boldsymbol{y} + \boldsymbol{z} \rangle = \langle \boldsymbol{y} + \boldsymbol{z} \mid \boldsymbol{x} \rangle^*$	by *conjugate symmetric* property of $\langle \triangle \mid \triangledown \rangle$	(Definition J.1 page 255)
$= (\langle \boldsymbol{y} \mid \boldsymbol{x} \rangle + \langle \boldsymbol{z} \mid \boldsymbol{x} \rangle)^*$	by *additive* property of $\langle \triangle \mid \triangledown \rangle$	(Definition J.1 page 255)
$= \langle \boldsymbol{y} \mid \boldsymbol{x} \rangle^* + \langle \boldsymbol{z} \mid \boldsymbol{x} \rangle^*$	by *distributive* property of $*$	(Definition I.3 page 250)
$= \langle \boldsymbol{x} \mid \boldsymbol{y} \rangle + \langle \boldsymbol{x} \mid \boldsymbol{z} \rangle$	by *conjugate symmetric* property of $\langle \triangle \mid \triangledown \rangle$	(Definition J.1 page 255)
$\langle \boldsymbol{x} \mid \alpha \boldsymbol{y} \rangle = \langle \alpha \boldsymbol{y} \mid \boldsymbol{x} \rangle^*$	by *conjugate symmetric* property of $\langle \triangle \mid \triangledown \rangle$	(Definition J.1 page 255)

[1] Istrăţescu (1987) page 111 ⟨Definition 4.1.1⟩, Bollobás (1999) pages 130–131, Haaser and Sullivan (1991) page 277, Aliprantis and Burkinshaw (1998), page 276, Peano (1888b) page 72

[2] Berberian (1961) page 27, Haaser and Sullivan (1991) page 277

$$
\begin{aligned}
&= (\alpha\,\langle \boldsymbol{y} \mid \boldsymbol{x}\rangle)^* && \text{by } \textit{homogeneous} \text{ property of } \langle \triangle \mid \triangledown\rangle && \text{(Definition J.1 page 255)}\\
&= \alpha^*\,\langle \boldsymbol{y} \mid \boldsymbol{x}\rangle^* && \text{by } \textit{antiautomorphic} \text{ property of } * && \text{(Definition I.3 page 250)}\\
&= \alpha^*\,\langle \boldsymbol{x} \mid \boldsymbol{y}\rangle && \text{by } \textit{conjugate symmetric} \text{ property of } \langle \triangle \mid \triangledown\rangle && \text{(Definition J.1 page 255)}\\
\langle \boldsymbol{x} \mid \mathbb{0}\rangle &= \langle \mathbb{0} \mid \boldsymbol{x}\rangle^* && \text{by } \textit{conjugate symmetric} \text{ property of } \langle \triangle \mid \triangledown\rangle && \text{(Definition J.1 page 255)}\\
&= \langle 0\cdot \boldsymbol{y} \mid \boldsymbol{x}\rangle^* && && \\
&= (0\cdot\langle \boldsymbol{y} \mid \boldsymbol{x}\rangle)^* && \text{by } \textit{homogeneous} \text{ property of } \langle \triangle \mid \triangledown\rangle && \text{(Definition J.1 page 255)}\\
&= 0 && && \\
\langle \mathbb{0} \mid \boldsymbol{x}\rangle &= \langle 0\cdot \boldsymbol{y} \mid \boldsymbol{x}\rangle && && \\
&= (0\cdot\langle \boldsymbol{y} \mid \boldsymbol{x}\rangle) && \text{by } \textit{homogeneous} \text{ property of } \langle \triangle \mid \triangledown\rangle && \text{(Definition J.1 page 255)}\\
&= 0 && && \\
\langle \boldsymbol{x}-\boldsymbol{y} \mid \boldsymbol{z}\rangle &= \langle \boldsymbol{x}+(-\boldsymbol{y}) \mid \boldsymbol{z}\rangle && \text{by definition of } + && \\
&= \langle \boldsymbol{x} \mid \boldsymbol{z}\rangle + \langle -\boldsymbol{y} \mid \boldsymbol{z}\rangle && \text{by } \textit{additive} \text{ property of } \langle \triangle \mid \triangledown\rangle && \text{(Definition J.1 page 255)}\\
&= \langle \boldsymbol{x} \mid \boldsymbol{z}\rangle - \langle \boldsymbol{y} \mid \boldsymbol{z}\rangle && \text{by } \textit{homogeneous} \text{ property of } \langle \triangle \mid \triangledown\rangle && \text{(Definition J.1 page 255)}\\
\langle \boldsymbol{x} \mid \boldsymbol{y}-\boldsymbol{z}\rangle &= \langle \boldsymbol{y}-\boldsymbol{z} \mid \boldsymbol{x}\rangle^* && \text{by } \textit{conjugate symmetric} \text{ property of } \langle \triangle \mid \triangledown\rangle && \text{(Definition J.1 page 255)}\\
&= (\langle \boldsymbol{y} \mid \boldsymbol{x}\rangle - \langle \boldsymbol{z} \mid \boldsymbol{x}\rangle)^* && \text{by 4.} && \\
&= \langle \boldsymbol{y} \mid \boldsymbol{x}\rangle^* - \langle \boldsymbol{z} \mid \boldsymbol{x}\rangle^* && \text{by } \textit{distributive} \text{ property of } * && \text{(Definition I.3 page 250)}\\
&= \langle \boldsymbol{x} \mid \boldsymbol{y}\rangle - \langle \boldsymbol{x} \mid \boldsymbol{z}\rangle && \text{by } \textit{conjugate symmetric} \text{ property of } \langle \triangle \mid \triangledown\rangle && \text{(Definition J.1 page 255)}
\end{aligned}
$$

$$
\begin{aligned}
& \langle \boldsymbol{x} \mid \boldsymbol{z}\rangle = \langle \boldsymbol{y} \mid \boldsymbol{z}\rangle && \forall \boldsymbol{z} && \\
\iff\quad & \langle \boldsymbol{x} \mid \boldsymbol{z}\rangle - \langle \boldsymbol{y} \mid \boldsymbol{z}\rangle = 0 && \forall \boldsymbol{z} && \text{by property of complex numbers}\\
\iff\quad & \langle \boldsymbol{x}-\boldsymbol{y} \mid \boldsymbol{z}\rangle = 0 && \forall \boldsymbol{z} && \text{by 4.}\\
\iff\quad & \boldsymbol{x}-\boldsymbol{y} = \mathbb{0} && \forall \boldsymbol{z} && \text{by } \textit{non-isotropic} \text{ property of } \langle \triangle \mid \triangledown\rangle \text{ (Definition J.1 page 255)}
\end{aligned}
$$

Proof that $\langle \boldsymbol{x} \mid \boldsymbol{y}\rangle = 0 \implies \boldsymbol{y} = \mathbb{0}$:

1. Suppose $\boldsymbol{y} \neq \mathbb{0}$;
2. Then $\langle \boldsymbol{y} \mid \boldsymbol{y}\rangle \neq \mathbb{0}$ by the *non-isotropic* property of $\langle \triangle \mid \triangledown\rangle$ (Definition J.1 page 255)
3. But because $\boldsymbol{y} \in X$, the left hypothesis implies that $\langle \boldsymbol{y} \mid \boldsymbol{y}\rangle = 0$.
4. This is a *contradiction*.
5. Therefore $\boldsymbol{y} \neq \mathbb{0}$ must be incorrect and $\boldsymbol{y} = \mathbb{0}$ must be correct.

Proof that $\langle \boldsymbol{x} \mid \boldsymbol{y}\rangle = 0 \impliedby \boldsymbol{y} = \mathbb{0}$:

$$
\begin{aligned}
\langle \boldsymbol{x} \mid \boldsymbol{y}\rangle &= \langle \boldsymbol{x} \mid \mathbb{0}\rangle && \text{by right hypothesis}\\
&= 0 && \text{by Theorem J.1 page 255}
\end{aligned}
$$

✏

One of the most useful and widely used inequalities in analysis is the *Cauchy-Schwarz Inequality* (sometimes also called the *Cauchy-Bunyakovsky-Schwarz Inequality*). In fact, we will use this inequality shortly to prove that every inner product space *has* a norm and therefore every inner product space *is* a normed linear space.

Theorem J.2 (Cauchy-Schwarz Inequality). [3] *Let* $\big(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle \mid \triangledown\rangle\big)$ *be an* INNER PRODUCT SPACE *and* $|\cdot| \in \mathbb{R}^{\mathbb{C}}$ *an* ABSOLUTE VALUE *function* (Definition F.4 page 206). *Let* $\|\cdot\|$ *be a function in* $\mathbb{R}^{\mathbb{F}}$ *such*

[3] Haaser and Sullivan (1991) page 278, Aliprantis and Burkinshaw (1998), page 278, Cauchy (1821) page 455, Bunyakovsky (1859) page 6, Schwarz (1885)

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

that $\|x\| \triangleq \sqrt{\langle x \mid x\rangle}$.[4]

THM

$$
\begin{array}{rcll}
|\langle x \mid y\rangle|^2 & \leq & \langle x \mid x\rangle \ \langle y \mid y\rangle & & \forall x,y \in X \\
|\langle x \mid y\rangle|^2 & = & \langle x \mid x\rangle \ \langle y \mid y\rangle & \iff \ \exists \alpha \in \mathbb{F} \ \text{ such that } \ y = \alpha x & \forall x,y \in X \\
|\langle x \mid y\rangle| & \leq & \|x\| \ \|y\| & & \forall x,y \in X \\
|\langle x \mid y\rangle| & = & \|x\| \ \|y\| & \iff \ \exists \alpha \in \mathbb{F} \ \text{ such that } \ y = \alpha x & \forall x,y \in X
\end{array}
$$

✎PROOF:

1. Proof that $|\langle x \mid y\rangle| \leq \|x\| \ \|y\|$: [5]

   (a) $y = \mathbb{0}$ case:

$$
\begin{aligned}
|\langle x \mid y\rangle|^2 &= |\langle x \mid \mathbb{0}\rangle|^2 && \text{by } y = \mathbb{0} \text{ hypothesis} \\
&= |\langle \mathbb{0} \mid x\rangle|^2 && \text{by Definition J.1 page 255} \\
&= |\langle 0\mathbb{0} \mid x\rangle|^2 && \text{by Definition G.1 page 207} \\
&= |0 \ \langle \mathbb{0} \mid x\rangle|^2 && \text{by Definition J.1 page 255} \\
&= 0 \\
&= \langle x \mid x\rangle \ \langle \mathbb{0} \mid \mathbb{0}\rangle \\
&= \langle x \mid x\rangle \ \langle y \mid y\rangle && \text{by } y = \mathbb{0} \text{ hypothesis}
\end{aligned}
$$

   (b) $y \neq \mathbb{0}$ case: Let $\lambda \triangleq \frac{\langle x \mid y\rangle}{\langle y \mid y\rangle}$.

$$
\begin{aligned}
0 &\leq \langle x - \lambda y \mid x - \lambda y\rangle && \text{by Definition J.1} \\
&= \langle x \mid x - \lambda y\rangle + \langle -\lambda y \mid x - \lambda y\rangle && \text{by Definition J.1} \\
&= \langle x - \lambda y \mid x\rangle^* - \lambda \ \langle y \mid x - \lambda y\rangle && \text{by Definition J.1} \\
&= \langle x \mid x\rangle^* + \langle -\lambda y \mid x\rangle^* - \lambda \ \langle x - \lambda y \mid y\rangle^* && \text{by Definition J.1} \\
&= \langle x \mid x\rangle^* - \lambda^* \ \langle y \mid x\rangle^* - \lambda \ \langle x \mid y\rangle^* - \lambda \ \langle -\lambda y \mid y\rangle^* && \text{by Definition J.1} \\
&= \langle x \mid x\rangle - \lambda^* \ \langle x \mid y\rangle - \lambda \ \langle x \mid y\rangle^* + \lambda\lambda^* \ \langle y \mid y\rangle^* && \text{by Definition J.1} \\
&= \langle x \mid x\rangle + \cancelto{0}{\left[\frac{\langle x \mid y\rangle}{\langle y \mid y\rangle}\lambda^* \ \langle y \mid y\rangle - \lambda^* \ \langle x \mid y\rangle\right]} - \frac{\langle x \mid y\rangle}{\langle y \mid y\rangle} \ \langle x \mid y\rangle^* && \text{by definition of } \lambda \\
&= \langle x \mid x\rangle - \frac{1}{\langle y \mid y\rangle} \ |\langle x \mid y\rangle|^2 \\
\implies |\langle x \mid y\rangle|^2 &\leq \langle x \mid x\rangle \ \langle y \mid y\rangle
\end{aligned}
$$

2. Proof that $|\langle x \mid y\rangle|^2 = \langle x \mid x\rangle \ \langle y \mid y\rangle \iff y = ax$:
   Let $\frac{1}{a} \triangleq \lambda \triangleq \frac{\langle x \mid y\rangle}{\langle y \mid y\rangle}$. Then…

$$
\begin{aligned}
& y = ax \\
\iff \quad & x = \lambda y \\
\iff \quad & x - \lambda y = \mathbb{0} \\
\iff \quad & 0 = \langle x - \lambda y \mid x - \lambda y\rangle && \text{by Definition J.1 page 255} \\
& \quad = \langle x \mid x - \lambda y\rangle + \langle -\lambda y \mid x - \lambda y\rangle && \text{by Definition J.1 page 255} \\
& \quad = \langle x - \lambda y \mid x\rangle^* - \lambda \ \langle y \mid x - \lambda y\rangle && \text{by Definition J.1 page 255} \\
& \quad \vdots && \text{(same steps as in 1(b)}
\end{aligned}
$$

[4] The function $\|\cdot\|$ is a *norm* (Theorem J.4 page 260) and is called the *norm induced by the inner product* $\langle \triangle \mid \triangledown\rangle$ (Definition J.2 page 260).

[5] Haaser and Sullivan (1991), page 278

COPYRIGHT © 2019 DANIEL J. GREENHOE | https://github.com/dgreenhoe/pdfs/blob/master/abcstat.pdf | CC BY-NC-ND

$$= \langle \boldsymbol{x} \mid \boldsymbol{x} \rangle - \frac{1}{\langle \boldsymbol{y} \mid \boldsymbol{y} \rangle} \; |\langle \boldsymbol{x} \mid \boldsymbol{y} \rangle|^2$$

$$\iff \qquad |\langle \boldsymbol{x} \mid \boldsymbol{y} \rangle|^2 = \langle \boldsymbol{x} \mid \boldsymbol{x} \rangle \; \langle \boldsymbol{y} \mid \boldsymbol{y} \rangle$$

3. Alternate proof for $|\langle \boldsymbol{x} \mid \boldsymbol{y} \rangle| \le \|\boldsymbol{x}\| \, \|\boldsymbol{y}\|$: (Note: This is essentially the same proof as used by Schwarz). [6]

   (a) Proof that $\left\{a\lambda^2 + b\lambda + c \ge 0 \qquad \forall \lambda \in \mathbb{R}\right\} \quad \Longrightarrow \quad \left\{b^2 \le 4ac\right\}$ (quadratic discriminate inequality):



   Let $k \in (0, \infty)$, and $r_1, r_2 \in \mathbb{C}$ be the roots of $a\lambda^2 + b\lambda + c = 0$. Then

$$\begin{aligned}
0 &\le a\lambda^2 + b\lambda + c && \text{by left hypothesis} \\
&= k(\lambda - r_1)(\lambda - r_2) && \text{by definition of } r_1 \text{ and } r_2 \\
&= k(\lambda^2 - r_1\lambda - r_2\lambda + r_1 r_2) \\
\Longrightarrow \qquad \lambda^2 - r_1\lambda - r_2\lambda + r_1 r_2 &\ge 0 \\
\Longrightarrow \qquad r_1 &= r_2^* && \text{because } r_1 r_2 \ge 0 \text{ for } \lambda = 0
\end{aligned}$$

   The *quadratic equation* places another constraint on r_1 and r_2:

$$\begin{aligned}
\frac{b^2 + \sqrt{b^2 - 4ac}}{2a} &= r_1 && \text{by quadratic equation} \\
&= r_2^* && \text{by previous result} \\
&= \left(\frac{b^2 - \sqrt{b^2 - 4ac}}{2a}\right)^* && \text{by quadratic equation}
\end{aligned}$$

   The only way for this to be true is if $b^2 \le 4ac$ (the **discriminate** is non-positive).

   (b) Proof that $\langle \boldsymbol{y} \mid \boldsymbol{y} \rangle \lambda^2 + 2|\langle \boldsymbol{x} \mid \boldsymbol{y} \rangle|\lambda + \langle \boldsymbol{x} \mid \boldsymbol{x} \rangle \ge 0 \quad \forall \lambda \in \mathbb{R}$:

$$\begin{aligned}
0 &\le \langle \boldsymbol{x} + \alpha \boldsymbol{y} \mid \boldsymbol{x} + \alpha \boldsymbol{y} \rangle && \text{by Definition J.1 page 255} \\
&= \langle \boldsymbol{x} \mid \boldsymbol{x} + \alpha \boldsymbol{y} \rangle + \langle \alpha \boldsymbol{y} \mid \boldsymbol{x} + \alpha \boldsymbol{y} \rangle && \text{by Definition J.1 page 255} \\
&= \langle \boldsymbol{x} \mid \boldsymbol{x} + \alpha \boldsymbol{y} \rangle + \alpha \langle \boldsymbol{y} \mid \boldsymbol{x} + \alpha \boldsymbol{y} \rangle && \text{by Definition J.1 page 255} \\
&= \langle \boldsymbol{x} + \alpha \boldsymbol{y} \mid \boldsymbol{x} \rangle^* + \alpha \langle \boldsymbol{x} + \alpha \boldsymbol{y} \mid \boldsymbol{y} \rangle^* && \text{by Definition J.1 page 255} \\
&= \langle \boldsymbol{x} \mid \boldsymbol{x} \rangle^* + \langle \alpha \boldsymbol{y} \mid \boldsymbol{x} \rangle^* + \alpha \langle \boldsymbol{x} \mid \boldsymbol{y} \rangle^* + \alpha \langle \alpha \boldsymbol{y} \mid \boldsymbol{y} \rangle^* && \text{by Definition J.1 page 255} \\
&= \langle \boldsymbol{x} \mid \boldsymbol{x} \rangle^* + \alpha^* \langle \boldsymbol{y} \mid \boldsymbol{x} \rangle^* + \alpha \langle \boldsymbol{x} \mid \boldsymbol{y} \rangle^* + \alpha\alpha^* \langle \boldsymbol{y} \mid \boldsymbol{y} \rangle^* && \text{by Definition J.1 page 255} \\
&= \langle \boldsymbol{x} \mid \boldsymbol{x} \rangle + \alpha^* \langle \boldsymbol{x} \mid \boldsymbol{y} \rangle + \left(\alpha^* \langle \boldsymbol{x} \mid \boldsymbol{y} \rangle\right)^* + |\alpha|^2 \langle \boldsymbol{y} \mid \boldsymbol{y} \rangle && \text{by Definition J.1 page 255} \\
&= \langle \boldsymbol{x} \mid \boldsymbol{x} \rangle + 2\Re\left(\alpha^* \langle \boldsymbol{x} \mid \boldsymbol{y} \rangle\right) + |\alpha|^2 \langle \boldsymbol{y} \mid \boldsymbol{y} \rangle \\
&\le \langle \boldsymbol{x} \mid \boldsymbol{x} \rangle + 2\left|\alpha^* \langle \boldsymbol{x} \mid \boldsymbol{y} \rangle\right| + |\alpha|^2 \langle \boldsymbol{y} \mid \boldsymbol{y} \rangle \\
&= \langle \boldsymbol{x} \mid \boldsymbol{x} \rangle + 2|\langle \boldsymbol{x} \mid \boldsymbol{y} \rangle||\alpha| + \langle \boldsymbol{y} \mid \boldsymbol{y} \rangle |\alpha|^2 \\
&= \langle \boldsymbol{y} \mid \boldsymbol{y} \rangle |\alpha|^2 + 2\,|\langle \boldsymbol{x} \mid \boldsymbol{y} \rangle|\,|\alpha| + \langle \boldsymbol{x} \mid \boldsymbol{x} \rangle \\
&= \underbrace{\langle \boldsymbol{y} \mid \boldsymbol{y} \rangle}_{a} \lambda^2 + \underbrace{2|\langle \boldsymbol{x} \mid \boldsymbol{y} \rangle|}_{b} \lambda + \underbrace{\langle \boldsymbol{x} \mid \boldsymbol{x} \rangle}_{c} && \text{because } \lambda \triangleq |\alpha| \in \mathbb{R}
\end{aligned}$$

[6] Aliprantis and Burkinshaw (1998), page 278, Steele (2004), page 11

BY-NC-ND Copyright © 2019 Daniel J. Greenhoe

(c) The above equation is in the quadratic form used in the lemma of part (a).

$$\left(\underbrace{2|\langle x \mid y\rangle|}_{b}\right)^2 \le 4\,\underbrace{\langle y \mid y\rangle}_{a}\;\underbrace{\langle x \mid x\rangle}_{c} \qquad \text{by the results of parts (a) and (b)}$$

$$\Longrightarrow\ |\langle x \mid y\rangle|^2 \le \langle x \mid x\rangle\ \langle y \mid y\rangle$$

4. Proof that $|\langle x \mid y\rangle| \le \|x\|\ \|y\|$:
   This follows directly from the definition $\|x\| \triangleq \sqrt{\langle x \mid x\rangle}$.

5. Proof that $|\langle x \mid y\rangle| = \|x\|\ \|y\| \iff \exists \alpha \in \mathbb{C}$ such that $y = \alpha x$:
   This follows directly from the definition $\|x\| \triangleq \sqrt{\langle x \mid x\rangle}$.

Corollary J.1. [7] *Let* $\big(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle \mid \nabla\rangle\big)$ *be an* INNER PRODUCT SPACE.

COR $\langle x \mid y\rangle$ *is* CONTINUOUS *in both* x *and* y.

✎PROOF: Let $\|x\| \triangleq \sqrt{\langle x \mid x\rangle}$.

$$\begin{aligned}
|\langle x+\epsilon \mid y\rangle - \langle x \mid y\rangle|^2 &= |\langle x+\epsilon - x \mid y\rangle|^2 && \text{by } \textit{additivity} \text{ of } \langle \triangle \mid \nabla\rangle && \text{(Definition J.1 page 255)}\\
&= |\langle \epsilon \mid y\rangle|^2 \\
&\le \|\epsilon\|^2\ \|y\| && \text{by } \textit{Cauchy-Schwarz Inequality} && \text{(Theorem J.2 page 256)}
\end{aligned}$$

J.2 Relationship between norms and inner products

J.2.1 Norms induced by inner products

Lemma J.1 (Polar Identity). [8] *Let* $\big(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle \mid \nabla\rangle\big)$ *be an* INNER PRODUCT SPACE *(Definition J.1 page 255). Let* $\Re z$ *represent the real part of* $z \in \mathbb{C}$. *Let* $\|\cdot\|$ *be a function in* $\mathbb{R}^{\mathbb{F}}$ *such that* $\|x\| \triangleq \sqrt{\langle x \mid x\rangle}$.[9]

LEM $\|x+y\|^2 = \|x\|^2 + 2\mathbf{R}_{\mathrm{e}}\left[\langle x \mid y\rangle\right] + \|y\|^2 \qquad \forall x,y \in X$

✎PROOF:

$$\begin{aligned}
\|x+y\|^2 &= \langle x+y \mid x+y\rangle && \text{by definition of } \textit{induced norm} && \text{(Theorem J.4 page 260)}\\
&= \langle x \mid x+y\rangle + \langle y \mid x+y\rangle && \text{by Definition J.1 page 255}\\
&= \langle x+y \mid x\rangle^* + \langle x+y \mid y\rangle^* && \text{by Definition J.1 page 255}\\
&= \langle x \mid x\rangle^* + \langle y \mid x\rangle^* + \langle x \mid y\rangle^* + \langle y \mid y\rangle^* && \text{by Definition J.1 page 255}\\
&= \langle x \mid x\rangle + \langle x \mid y\rangle + \langle x \mid y\rangle^* + \langle y \mid y\rangle && \text{by definition of } \textit{inner product} && \text{(Definition J.1 page 255)}\\
&= \|x\|^2 + 2\Re\,\langle x \mid y\rangle + \|y\|^2 && \text{by definition of } \textit{induced norm} && \text{(Theorem J.4 page 260)}
\end{aligned}$$

[7] Bollobás (1999) page 132, Aliprantis and Burkinshaw (1998) page 279 ⟨Lemma 32.4⟩
[8] Conway (1990) page 4, Heil (2011) page 27 ⟨Lemma 1.36(a)⟩
[9] The function $\|\cdot\|$ is a *norm* (Theorem J.4 page 260) and is called the *norm induced by the inner product* $\langle \triangle \mid \nabla\rangle$ (Definition J.2 page 260).

COPYRIGHT © 2019 DANIEL J. GREENHOE

Theorem J.3 (Minkowski's Inequality). [10] *Let* $\big(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle\triangle\,|\,\triangledown\rangle\big)$ *be an* INNER PRODUCT SPACE. *Let* $\|\cdot\|$ *be a function in* $\mathbb{R}^{\mathbb{F}}$ *such that* $\|\boldsymbol{x}\| \triangleq \sqrt{\langle \boldsymbol{x}\,|\,\boldsymbol{x}\rangle}$.[11]

THM $\|\boldsymbol{x}+\boldsymbol{y}\| \le \|\boldsymbol{x}\| + \|\boldsymbol{y}\| \quad \forall_{\boldsymbol{x},\boldsymbol{y}\in X}$

PROOF:

$$\begin{aligned}
\|\boldsymbol{x}+\boldsymbol{y}\|^2 &= \|\boldsymbol{x}\|^2 + 2\Re\langle \boldsymbol{x}\,|\,\boldsymbol{y}\rangle + \|\boldsymbol{y}\|^2 && \text{by } \textit{Polar Identity} && \text{(Lemma J.1 page 259)}\\
&\le \|\boldsymbol{x}\|^2 + 2|\langle \boldsymbol{x}\,|\,\boldsymbol{y}\rangle| + \|\boldsymbol{y}\|^2 \\
&\le \|\boldsymbol{x}\|^2 + 2\sqrt{\langle \boldsymbol{x}\,|\,\boldsymbol{x}\rangle}\sqrt{\langle \boldsymbol{y}\,|\,\boldsymbol{y}\rangle} + \|\boldsymbol{y}\|^2 && \text{by } \textit{Cauchy-Schwarz Inequality} && \text{(Theorem J.2 page 256)}\\
&= (\|\boldsymbol{x}\| + \|\boldsymbol{y}\|)^2
\end{aligned}$$

Theorem J.4 (induced norm). [12] *Let* $\boldsymbol{\Omega} \triangleq \big(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle\triangle\,|\,\triangledown\rangle\big)$ *be an* INNER PRODUCT SPACE *(Definition J.1 page 255)*.

THM $\|\boldsymbol{x}\| \triangleq \sqrt{\langle \boldsymbol{x}\,|\,\boldsymbol{x}\rangle} \quad \Longrightarrow \quad \|\cdot\|$ *is a* NORM

PROOF: For a function to be a norm, it must satisfy the four properties listed in Definition K.1 (page 271).

1. Proof that $\|\cdot\|$ is a norm:
   (a) Proof that $\|\boldsymbol{x}\| > 0$ for $\boldsymbol{x} \ne 0$ (non-negative):
   By Definition J.1 page 255, all inner products have this property.
   (b) Proof that $\|\boldsymbol{x}\| = 0 \iff \boldsymbol{x} = 0$ (non-isometric):
   By Definition J.1, all inner products have this property.
   (c) Prove $\|a\boldsymbol{x}\| = |a|\,\|\boldsymbol{x}\|$ (homogeneous):
   $$\|a\boldsymbol{x}\| \triangleq \sqrt{\langle a\boldsymbol{x}\,|\,a\boldsymbol{x}\rangle} = \sqrt{aa^*\langle \boldsymbol{x}\,|\,\boldsymbol{x}\rangle} = \sqrt{|a|^2\langle \boldsymbol{x}\,|\,\boldsymbol{x}\rangle} = |a|\,\|\boldsymbol{x}\|$$
   (d) Proof that $\|\boldsymbol{x}+\boldsymbol{y}\| \le \|\boldsymbol{x}\| + \|\boldsymbol{y}\|$ (subadditive): This is true by Minkowski's inequality page **??**
2. Proof that every inner product space is a normed linear space:
   Since every inner product induces a norm, so every inner product space has a norm (the norm induced by the inner product) and is therefore a normed linear space.

Theorem J.4 (previous theorem) demonstrates that in any inner product space $\big(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle\triangle\,|\,\triangledown\rangle\big)$, the function $\|\boldsymbol{x}\| \triangleq \sqrt{\langle \boldsymbol{x}\,|\,\boldsymbol{x}\rangle}$ is a norm. That is, $\|\boldsymbol{x}\|$ is the *norm induced by the inner product*. This norm is formally defined next.

Definition J.2. *Let* $\big(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle\triangle\,|\,\triangledown\rangle\big)$ *be an* INNER PRODUCT SPACE *(Definition J.1 page 255)*.

DEF *The* ***norm induced by the inner product*** $\langle\triangle\,|\,\triangledown\rangle$ *is defined as*
$$\|\boldsymbol{x}\| \triangleq \sqrt{\langle \boldsymbol{x}\,|\,\boldsymbol{x}\rangle}$$

[10] Aliprantis and Burkinshaw (1998) pages 278–279 ⟨Theorem 32.3⟩, Maligranda (1995), Minkowski (1910) page 115

[11] The function $\|\cdot\|$ is a *norm* (Theorem J.4 page 260) and is called the *normed induced by the inner product* $\langle\triangle\,|\,\triangledown\rangle$ (Definition J.2 page 260).

[12] Aliprantis and Burkinshaw (1998), pages 278–279, Haaser and Sullivan (1991) page 278

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

J.2.2 Inner products induced by norms

Theorem J.4 (page 260) demonstrates that if a *linear space* (Definition G.1 page 207) has an *inner product* (Definition J.1 page 255), then that inner product always induces a *norm* (Definition K.1 page 271), and the relationship between the two is simply $\|\boldsymbol{x}\| = \sqrt{\langle \boldsymbol{x} \mid \boldsymbol{x} \rangle}$ (Definition J.2 page 260). But what about the converse? What if a linear space has a norm—can that norm also induce an inner product? The answer in general is "no": Not all norms can induce an inner product. But a less harsh answer is "sometimes": Some norms **can** induce inner products. This leads to some important and interesting questions:

1. How many different inner products can be induced from a single norm? The answer turns out to be **at most** one, but maybe none (Theorem J.5 page 261).
2. When a norm *can* induce an inner product, what is that (unique) inner product? The inner product expressed in terms of the norm is given by the *Polarization Identity* (Theorem J.6 page 262).
3. Which norms can induce an inner product and which ones cannot? The answer is that norms that satisfy the *parallelogram law* (Theorem J.7 page 263) **can** induce an inner product; and the ones that don't, cannot (Theorem J.7 page 263).

Theorem J.5. [13] *Let* $\big(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \|\cdot\|\big)$ *be a* NORMED LINEAR SPACE *(Definition K.1 page 271)*.

THM

$$\underbrace{\left.\begin{array}{l} \exists\ \langle \triangle \mid \triangledown \rangle \text{ and } (\cdot \mid \cdot) \quad \text{such that} \\ \|\boldsymbol{x}\|^2 = \langle \boldsymbol{x} \mid \boldsymbol{x} \rangle = (\boldsymbol{x} \mid \boldsymbol{x}) \quad \forall x \in X \end{array}\right\}}_{\textit{If a norm is induced by two inner products...}} \implies \underbrace{\langle \boldsymbol{x} \mid \boldsymbol{y} \rangle = (\boldsymbol{x} \mid \boldsymbol{y}) \quad \forall x, y \in X}_{\textit{...then those two inner products are equivalent.}}$$

✎PROOF:

$$\begin{aligned}
2\langle \boldsymbol{x} \mid \boldsymbol{y} \rangle &= \big[\langle \boldsymbol{x} \mid \boldsymbol{y} \rangle + \langle \boldsymbol{y} \mid \boldsymbol{x} \rangle\big] + \big[\langle \boldsymbol{x} \mid \boldsymbol{y} \rangle - \langle \boldsymbol{y} \mid \boldsymbol{x} \rangle\big] \\
&= \big[\langle \boldsymbol{x} \mid \boldsymbol{y} \rangle + \langle \boldsymbol{y} \mid \boldsymbol{x} \rangle\big] + i\big[-i\langle \boldsymbol{x} \mid \boldsymbol{y} \rangle + i\langle \boldsymbol{y} \mid \boldsymbol{x} \rangle\big] \\
&= \big[\langle \boldsymbol{x} \mid \boldsymbol{y} \rangle + \langle \boldsymbol{y} \mid \boldsymbol{x} \rangle\big] + i\big[-\langle i\boldsymbol{x} \mid \boldsymbol{y} \rangle - \langle \boldsymbol{y} \mid i\boldsymbol{x} \rangle\big] \\
&= \left(\underbrace{\big[\langle \boldsymbol{x} \mid \boldsymbol{y} \rangle + \langle \boldsymbol{y} \mid \boldsymbol{x} \rangle + \langle \boldsymbol{x} \mid \boldsymbol{x} \rangle + \langle \boldsymbol{y} \mid \boldsymbol{y} \rangle\big]}_{\langle \boldsymbol{x}+\boldsymbol{y} \mid \boldsymbol{x}+\boldsymbol{y} \rangle} - \underbrace{\big[\langle \boldsymbol{x} \mid \boldsymbol{x} \rangle + \langle \boldsymbol{y} \mid \boldsymbol{y} \rangle\big]}_{\text{residue}}\right) \\
&\quad - i\left(\underbrace{\big[\langle i\boldsymbol{x} \mid \boldsymbol{y} \rangle + \langle \boldsymbol{y} \mid i\boldsymbol{x} \rangle + \langle i\boldsymbol{x} \mid i\boldsymbol{x} \rangle + \langle \boldsymbol{y} \mid \boldsymbol{y} \rangle\big]}_{\langle i\boldsymbol{x}+\boldsymbol{y} \mid i\boldsymbol{x}+\boldsymbol{y} \rangle} - \underbrace{\big[\langle i\boldsymbol{x} \mid i\boldsymbol{x} \rangle + \langle \boldsymbol{y} \mid \boldsymbol{y} \rangle\big]}_{\text{residue}}\right) \\
&= \big(\big[\langle \boldsymbol{x}+\boldsymbol{y} \mid \boldsymbol{x}+\boldsymbol{y} \rangle\big] - \big[\langle \boldsymbol{x} \mid \boldsymbol{x} \rangle + \langle \boldsymbol{y} \mid \boldsymbol{y} \rangle\big]\big) - i\big(\big[\langle i\boldsymbol{x}+\boldsymbol{y} \mid i\boldsymbol{x}+\boldsymbol{y} \rangle\big] - \big[\langle i\boldsymbol{x} \mid i\boldsymbol{x} \rangle + \langle \boldsymbol{y} \mid \boldsymbol{y} \rangle\big]\big) \\
&= \big(\big[(\boldsymbol{x}+\boldsymbol{y} \mid \boldsymbol{x}+\boldsymbol{y})\big] - \big[(\boldsymbol{x} \mid \boldsymbol{x}) + (\boldsymbol{y} \mid \boldsymbol{y})\big]\big) - i\big(\big[(i\boldsymbol{x}+\boldsymbol{y} \mid i\boldsymbol{x}+\boldsymbol{y})\big] - \big[(i\boldsymbol{x} \mid i\boldsymbol{x}) + (\boldsymbol{y} \mid \boldsymbol{y})\big]\big) \\
&= \left(\underbrace{\big[(\boldsymbol{x} \mid \boldsymbol{y}) + (\boldsymbol{y} \mid \boldsymbol{x}) + (\boldsymbol{x} \mid \boldsymbol{x}) + (\boldsymbol{y} \mid \boldsymbol{y})\big]}_{(\boldsymbol{x}+\boldsymbol{y} \mid \boldsymbol{x}+\boldsymbol{y})} - \underbrace{\big[(\boldsymbol{x} \mid \boldsymbol{x}) + (\boldsymbol{y} \mid \boldsymbol{y})\big]}_{\text{residue}}\right) \\
&\quad - i\left(\underbrace{\big[(i\boldsymbol{x} \mid \boldsymbol{y}) + (\boldsymbol{y} \mid i\boldsymbol{x}) + (i\boldsymbol{x} \mid i\boldsymbol{x}) + (\boldsymbol{y} \mid \boldsymbol{y})\big]}_{(i\boldsymbol{x}+\boldsymbol{y} \mid i\boldsymbol{x}+\boldsymbol{y})} - \underbrace{\big[(i\boldsymbol{x} \mid i\boldsymbol{x}) + (\boldsymbol{y} \mid \boldsymbol{y})\big]}_{\text{residue}}\right)
\end{aligned}$$

[13] Aliprantis and Burkinshaw (1998), page 280, Bollobás (1999), page 132, Jordan and von Neumann (1935) page 721

COPYRIGHT © 2019 DANIEL J. GREENHOE

$$
\begin{aligned}
&= \left[(\boldsymbol{x} \mid \boldsymbol{y}) + (\boldsymbol{y} \mid \boldsymbol{x})\right] + i\left[-(i\boldsymbol{x} \mid \boldsymbol{y}) - (\boldsymbol{y} \mid i\boldsymbol{x})\right] \\
&= \left[(\boldsymbol{x} \mid \boldsymbol{y}) + (\boldsymbol{y} \mid \boldsymbol{x})\right] + i\left[-i(\boldsymbol{x} \mid \boldsymbol{y}) + i(\boldsymbol{y} \mid \boldsymbol{x})\right] \\
&= \left[(\boldsymbol{x} \mid \boldsymbol{y}) + (\boldsymbol{y} \mid \boldsymbol{x})\right] + \left[(\boldsymbol{x} \mid \boldsymbol{y}) - (\boldsymbol{y} \mid \boldsymbol{x})\right] \\
&= 2(\boldsymbol{x} \mid \boldsymbol{y})
\end{aligned}
$$

✏

Theorem J.6 (Polarization Identities). [14] *Let* $\boldsymbol{\Omega} \triangleq \left(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times})\right)$ *be a linear space,* $\langle \triangle \mid \triangledown \rangle \in \mathbb{F}^{X \times X}$ *a function, and* $\|\boldsymbol{x}\| \triangleq \sqrt{\langle \boldsymbol{x} \mid \boldsymbol{x} \rangle}$.

THM

$\left(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle \mid \triangledown \rangle\right)$ *is an inner product space* $\Longrightarrow$

$$
\underbrace{4\langle \boldsymbol{x} \mid \boldsymbol{y} \rangle = \begin{cases} \|\boldsymbol{x}+\boldsymbol{y}\|^2 - \|\boldsymbol{x}-\boldsymbol{y}\|^2 + i\,\|\boldsymbol{x}+i\boldsymbol{y}\|^2 - i\,\|\boldsymbol{x}-i\boldsymbol{y}\|^2 & \text{for } \mathbb{F}=\mathbb{C} \quad \forall_{\boldsymbol{x},\boldsymbol{y}\in X} \\ \|\boldsymbol{x}+\boldsymbol{y}\|^2 - \|\boldsymbol{x}-\boldsymbol{y}\|^2 & \text{for } \mathbb{F}=\mathbb{R} \quad \forall_{\boldsymbol{x},\boldsymbol{y}\in X} \end{cases}}_{\text{inner product induced by norm}}
$$

✎PROOF:

1. These follow directly from properties of *bilinear functionals* (Theorem **??** page **??**).
2. Alternative proof for $\mathbb{F} = \mathbb{C}$ case:

$$
\begin{aligned}
&\|\boldsymbol{x}+\boldsymbol{y}\|^2 - \|\boldsymbol{x}-\boldsymbol{y}\|^2 + i\,\|\boldsymbol{x}+i\boldsymbol{y}\|^2 - i\,\|\boldsymbol{x}-i\boldsymbol{y}\|^2 \\
&= \underbrace{\|\boldsymbol{x}\|^2 + \|\boldsymbol{y}\|^2 + 2\Re\langle \boldsymbol{x} \mid \boldsymbol{y} \rangle}_{\langle \boldsymbol{x}+\boldsymbol{y} \mid \boldsymbol{x}+\boldsymbol{y} \rangle} - \underbrace{\left(\|\boldsymbol{x}\|^2 + \|-\boldsymbol{y}\|^2 + 2\Re\langle \boldsymbol{x} \mid -\boldsymbol{y} \rangle\right)}_{\langle \boldsymbol{x}-\boldsymbol{y} \mid \boldsymbol{x}-\boldsymbol{y} \rangle} \\
&+ i\underbrace{\left(\|\boldsymbol{x}\|^2 + \|i\boldsymbol{y}\|^2 + 2\Re\langle \boldsymbol{x} \mid i\boldsymbol{y} \rangle\right)}_{i\langle \boldsymbol{x}+i\boldsymbol{y} \mid \boldsymbol{x}+i\boldsymbol{y} \rangle} - i\underbrace{\left(\|\boldsymbol{x}\|^2 + \|-i\boldsymbol{y}\|^2 + 2\Re\langle \boldsymbol{x} \mid -i\boldsymbol{y} \rangle\right)}_{i\langle \boldsymbol{x}-i\boldsymbol{y} \mid \boldsymbol{x}-i\boldsymbol{y} \rangle} && \text{by Lemma J.1 page 259} \\
&= \underbrace{\|\boldsymbol{x}\|^2 + \|\boldsymbol{y}\|^2 + 2\Re\langle \boldsymbol{x} \mid \boldsymbol{y} \rangle}_{\langle \boldsymbol{x}+\boldsymbol{y} \mid \boldsymbol{x}+\boldsymbol{y} \rangle} - \underbrace{\left(\|\boldsymbol{x}\|^2 + \|\boldsymbol{y}\|^2 - 2\Re\langle \boldsymbol{x} \mid \boldsymbol{y} \rangle\right)}_{\langle \boldsymbol{x}-\boldsymbol{y} \mid \boldsymbol{x}-\boldsymbol{y} \rangle} \\
&+ i\underbrace{\left(\|\boldsymbol{x}\|^2 + \|\boldsymbol{y}\|^2 + 2\Re\langle \boldsymbol{x} \mid i\boldsymbol{y} \rangle\right)}_{i\langle \boldsymbol{x}+i\boldsymbol{y} \mid \boldsymbol{x}+i\boldsymbol{y} \rangle} - i\underbrace{\left(\|\boldsymbol{x}\|^2 + \|\boldsymbol{y}\|^2 - 2\Re\langle \boldsymbol{x} \mid i\boldsymbol{y} \rangle\right)}_{i\langle \boldsymbol{x}-i\boldsymbol{y} \mid \boldsymbol{x}-i\boldsymbol{y} \rangle} && \text{by Definition J.1 page 255} \\
&= 4\Re\langle \boldsymbol{x} \mid \boldsymbol{y} \rangle + 4i\Re\langle \boldsymbol{x} \mid i\boldsymbol{y} \rangle \\
&= \underbrace{2\left(\langle \boldsymbol{x} \mid \boldsymbol{y} \rangle + \langle \boldsymbol{x} \mid \boldsymbol{y} \rangle^*\right)}_{4\Re\langle \boldsymbol{x} \mid \boldsymbol{y} \rangle} + \underbrace{2i\left(\langle \boldsymbol{x} \mid i\boldsymbol{y} \rangle + \langle \boldsymbol{x} \mid i\boldsymbol{y} \rangle^*\right)}_{4i\Re\langle \boldsymbol{x} \mid i\boldsymbol{y} \rangle} \\
&= 2\left(\langle \boldsymbol{x} \mid \boldsymbol{y} \rangle + \langle \boldsymbol{x} \mid \boldsymbol{y} \rangle^*\right) + 2i\left(i^*\langle \boldsymbol{x} \mid \boldsymbol{y} \rangle + (i^{**})\langle \boldsymbol{x} \mid \boldsymbol{y} \rangle^*\right) \\
&= 2\left(\langle \boldsymbol{x} \mid \boldsymbol{y} \rangle + \langle \boldsymbol{x} \mid \boldsymbol{y} \rangle^*\right) + 2i\left(-i\langle \boldsymbol{x} \mid \boldsymbol{y} \rangle + i\langle \boldsymbol{x} \mid \boldsymbol{y} \rangle^*\right) && \text{by Definition J.1 page 255} \\
&= 2\langle \boldsymbol{x} \mid \boldsymbol{y} \rangle + 2\langle \boldsymbol{x} \mid \boldsymbol{y} \rangle^* + 2\langle \boldsymbol{x} \mid \boldsymbol{y} \rangle - 2\langle \boldsymbol{x} \mid \boldsymbol{y} \rangle^* \\
&= 4\langle \boldsymbol{x} \mid \boldsymbol{y} \rangle
\end{aligned}
$$

✏



In plane geometry ($\mathbb{R}^2$), the *parallelogram law* states that the sum of the squares of the lengths of the sides of a parallelogram is equal to the sum of the squares of the lengths of its diagonals. This is illustrated in the figure to the left.

[14] Berberian (1961) pages 29–30 ⟨Theorem II.3.3⟩, Istrăţescu (1987) page 110 ⟨Proposition 4.1.5⟩, Bollobás (1999), page 132, Jordan and von Neumann (1935) page 721

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

Acutally, the parallelogram law can be generalized to *any inner product space* (not just in the plane). And if the parallelogram law happens to hold true in a normed linear space, then that normed linear space is actually an *inner product space*. The parallelogram law and its relation to inner product spaces is stated in the next theorem.

Theorem J.7 (Parallelogram law). [15] *Let* $\boldsymbol{\Omega} \triangleq \big(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle \mid \triangledown \rangle\big)$ *and* $\|\boldsymbol{x}\| \triangleq \sqrt{\langle \boldsymbol{x} \mid \boldsymbol{x} \rangle}$.

THM

$$\boldsymbol{\Omega} \text{ is an inner product space} \iff \underbrace{2\,\|\boldsymbol{x}\|^2 + 2\,\|\boldsymbol{y}\|^2 = \|\boldsymbol{x}+\boldsymbol{y}\|^2 + \|\boldsymbol{x}-\boldsymbol{y}\|^2}_{\text{PARALLELOGRAM LAW / VON NEUMANN-JORDAN CONDITION}} \qquad \forall \boldsymbol{x},\boldsymbol{y} \in \boldsymbol{\Omega}$$

✎PROOF:

1. Proof that $[\exists \langle \boldsymbol{x} \mid \boldsymbol{y} \rangle$ such that $\|\boldsymbol{x}\|^2 = \langle \boldsymbol{x} \mid \boldsymbol{x} \rangle\,] \implies$ [parallelogram law is true]:

$$\begin{aligned}
\|\boldsymbol{x}+\boldsymbol{y}\|^2 + \|\boldsymbol{x}-\boldsymbol{y}\|^2 &= \big[\|\boldsymbol{x}\|^2 + \|\boldsymbol{y}\|^2 + 2\mathbf{R}_{\mathsf{e}}\big[2\langle \boldsymbol{x} \mid \boldsymbol{y}\rangle\big]\big] \qquad + \big[\|\boldsymbol{x}\|^2 + \|-\boldsymbol{y}\|^2 + 2\mathbf{R}_{\mathsf{e}}\big[2\langle \boldsymbol{x} \mid -\boldsymbol{y}\rangle\big]\big] \\
&\qquad \text{by Lemma J.1 page 259} \\
&= \big[\|\boldsymbol{x}\|^2 + \|\boldsymbol{y}\|^2 + 2\mathbf{R}_{\mathsf{e}}\big[2\langle \boldsymbol{x} \mid \boldsymbol{y}\rangle\big]\big] \qquad + \big[\|\boldsymbol{x}\|^2 + \|\boldsymbol{y}\|^2 - 2\mathbf{R}_{\mathsf{e}}\big[2\langle \boldsymbol{x} \mid \boldsymbol{y}\rangle\big]\big] \\
&= 2\,\|\boldsymbol{x}\|^2 + 2\,\|\boldsymbol{y}\|^2
\end{aligned}$$

2. Proof that $[\exists \langle \boldsymbol{x} \mid \boldsymbol{y} \rangle$ such that $\|\boldsymbol{x}\|^2 = \langle \boldsymbol{x} \mid \boldsymbol{x} \rangle\,] \impliedby$ [parallelogram law is true]:
Note that **if** an inner product exists in the norm linear space $(\boldsymbol{\Omega}, \|\cdot\|)$, **then** that norm linear space is actually an inner product space $\big(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle \mid \triangledown \rangle\big)$. And if it is an inner product space, then by Theorem J.6 page 262 that inner product must be given by the **Polarization Identity**
$\langle \boldsymbol{x} \mid \boldsymbol{y} \rangle = \|a\boldsymbol{x}+\boldsymbol{y}\|^2 - \|a\boldsymbol{x}-\boldsymbol{y}\|^2 + i\,\|a\boldsymbol{x}+i\boldsymbol{y}\|^2 - i\,\|a\boldsymbol{x}-i\boldsymbol{y}\|^2$.
Therefore, here we must use the parallelogram law to show that the bilinear function $\mathsf{f}(\boldsymbol{x},\boldsymbol{y}) \triangleq \langle \boldsymbol{x} \mid \boldsymbol{y} \rangle$ given on the left hand side of the "=" relation is indeed an inner product— that is, that it satisfies the requirements of Definition J.1 page 255.

   (a) Proof that $\langle \boldsymbol{x} \mid \boldsymbol{x} \rangle \geq 0$ (non-negative):

$$\begin{aligned}
4\langle \boldsymbol{x} \mid \boldsymbol{x} \rangle &\triangleq \|\boldsymbol{x}+\boldsymbol{x}\|^2 - \cancelto{0}{\|\boldsymbol{x}-\boldsymbol{x}\|^2} + i\,\|\boldsymbol{x}+i\boldsymbol{x}\|^2 - i\,\|\boldsymbol{x}-i\boldsymbol{x}\|^2 && \text{by } \textit{Polarization Identity} \\
&= \|2\boldsymbol{x}\|^2 - \mathbb{0} + i\,\big(\,\|\boldsymbol{x}+i\boldsymbol{x}\|^2 - \|\boldsymbol{x}-i\boldsymbol{x}\|^2\,\big) && \text{by Definition K.1 page 271} \\
&= |2|^2\,\|\boldsymbol{x}\|^2 + i\,\big(\,\|\boldsymbol{x}+i\boldsymbol{x}\|^2 - |i|\,\|\boldsymbol{x}-i\boldsymbol{x}\|^2\,\big) \\
&= 4\,\|\boldsymbol{x}\|^2 + i\big(\,\|\boldsymbol{x}+i\boldsymbol{x}\|^2 - \cancelto{0}{\|i\boldsymbol{x}+\boldsymbol{x}\|^2}\,\big) && \text{by Definition K.1 page 271} \\
&= 4\,\|\boldsymbol{x}\|^2 \\
&\geq 0 && \text{by Definition K.1 page 271}
\end{aligned}$$

   (b) Proof that $\langle \boldsymbol{x} \mid \boldsymbol{x} \rangle = 0 \iff \boldsymbol{x} = \mathbb{0}$ (non-isotropic):

$$\begin{aligned}
4\langle \boldsymbol{x} \mid \boldsymbol{x} \rangle &= 4\,\|\boldsymbol{x}\|^2 && \text{by result of part (a)} \\
&= 0 \qquad \iff \qquad \boldsymbol{x} = \mathbb{0} && \text{by Definition K.1 page 271}
\end{aligned}$$

[15] Amir (1986), page 8, Istrăţescu (1987) page 110, Day (1973), page 151, Halmos (1998), page 14, Aliprantis and Burkinshaw (1998) pages 280–281 ⟨Theorem 32.6⟩, Riesz (1934) page 36?, Jordan and von Neumann (1935) pages 721–722

COPYRIGHT © 2019 DANIEL J. GREENHOE | CC BY-NC-ND

(c) Proof that $\langle \boldsymbol{x}+\boldsymbol{u} \mid \boldsymbol{y}\rangle = \langle \boldsymbol{x} \mid \boldsymbol{y}\rangle + \langle \boldsymbol{u} \mid \boldsymbol{y}\rangle$ (additive):[16]

$$
\begin{aligned}
4\langle \boldsymbol{x}+\boldsymbol{y} \mid \boldsymbol{z}\rangle &= 8\left\langle \frac{\boldsymbol{x}+\boldsymbol{y}}{2} \mid \boldsymbol{z}\right\rangle && \text{by Definition J.1 page 255}\\
&= 2\left\|\frac{\boldsymbol{x}+\boldsymbol{y}}{2}+\boldsymbol{z}\right\|^2 - 2\left\|\frac{\boldsymbol{x}+\boldsymbol{y}}{2}-\boldsymbol{z}\right\|^2\\
&\quad + 2i\left\|\frac{\boldsymbol{x}+\boldsymbol{y}}{2}+\boldsymbol{z}\right\|^2 - 2i\left\|\frac{\boldsymbol{x}+\boldsymbol{y}}{2}-i\boldsymbol{z}\right\|^2 && \text{by } \textit{Polarization Identity}\\
&= \left(2\left\|\frac{\boldsymbol{x}+\boldsymbol{y}}{2}+\boldsymbol{z}\right\|^2 + 2\left\|\frac{\boldsymbol{x}-\boldsymbol{y}}{2}\right\|^2\right)\\
&\quad - \left(2\left\|\frac{\boldsymbol{x}+\boldsymbol{y}}{2}-\boldsymbol{z}\right\|^2 + 2\left\|\frac{\boldsymbol{x}-\boldsymbol{y}}{2}\right\|^2\right)\\
&\quad + \left(2i\left\|\frac{\boldsymbol{x}+\boldsymbol{y}}{2}+\boldsymbol{z}\right\|^2 + 2i\left\|\frac{\boldsymbol{x}-\boldsymbol{y}}{2}\right\|^2\right)\\
&\quad - \left(2i\left\|\frac{\boldsymbol{x}+\boldsymbol{y}}{2}-i\boldsymbol{z}\right\|^2 + 2i\left\|\frac{\boldsymbol{x}-\boldsymbol{y}}{2}\right\|^2\right)\\
&= \left(\|\boldsymbol{x}+\boldsymbol{z}\|^2 + \|\boldsymbol{y}+\boldsymbol{z}\|^2\right) - \left(\|\boldsymbol{x}-\boldsymbol{z}\|^2 + \|\boldsymbol{y}-\boldsymbol{z}\|^2\right)\\
&\quad + \left(i\,\|\boldsymbol{x}+\boldsymbol{z}\|^2 + i\,\|\boldsymbol{y}+\boldsymbol{z}\|^2\right) - \left(i\,\|\boldsymbol{x}-i\boldsymbol{z}\|^2 + i\,\|\boldsymbol{y}-i\boldsymbol{z}\|^2\right) && \text{by parallelogram law}\\
&= \left(\|\boldsymbol{x}+\boldsymbol{z}\|^2 - \|\boldsymbol{x}-\boldsymbol{z}\|^2 + i\,\|\boldsymbol{x}+\boldsymbol{z}\|^2 - i\,\|\boldsymbol{x}-i\boldsymbol{z}\|^2\right)\\
&\quad + \left(\|\boldsymbol{y}+\boldsymbol{z}\|^2 - \|\boldsymbol{y}-\boldsymbol{z}\|^2 + i\,\|\boldsymbol{y}+\boldsymbol{z}\|^2 - i\,\|\boldsymbol{y}-i\boldsymbol{z}\|^2\right)\\
&= 4\langle \boldsymbol{x} \mid \boldsymbol{z}\rangle + 4\langle \boldsymbol{y} \mid \boldsymbol{z}\rangle && \text{by Polarization Identity}
\end{aligned}
$$

(d) Proof that $\langle \boldsymbol{x} \mid \boldsymbol{y}\rangle = \langle \boldsymbol{y} \mid \boldsymbol{y}\rangle^*$ (*conjugate symmetric*):

$$
\begin{aligned}
4\langle \boldsymbol{x} \mid \boldsymbol{y}\rangle &\triangleq \|\boldsymbol{x}+\boldsymbol{y}\|^2 - \|\boldsymbol{x}-\boldsymbol{y}\|^2 + i\,\|\boldsymbol{x}+i\boldsymbol{y}\|^2 - i\,\|\boldsymbol{x}-i\boldsymbol{y}\|^2 && \text{by Polarization Identity}\\
&= \|\boldsymbol{y}+\boldsymbol{x}\|^2 - \|\boldsymbol{y}-\boldsymbol{x}\|^2 + i\,\|i(\boldsymbol{y}-i\boldsymbol{x})\|^2 - i\,\|-i(\boldsymbol{y}+i\boldsymbol{x})\|^2 && \text{by Definition G.1 page 207}\\
&= \|\boldsymbol{y}+\boldsymbol{x}\|^2 - \|\boldsymbol{y}-\boldsymbol{x}\|^2 + i\,\|\boldsymbol{y}-i\boldsymbol{x}\|^2 - i\,\|\boldsymbol{y}+i\boldsymbol{x}\|^2 && \text{by Definition K.1 page 271}\\
&= \left(\|\boldsymbol{y}+\boldsymbol{x}\|^2 - \|\boldsymbol{y}-\boldsymbol{x}\|^2 - i\,\|\boldsymbol{y}-i\boldsymbol{x}\|^2 + i\,\|\boldsymbol{y}+i\boldsymbol{x}\|^2\right)^*\\
&= \left(\|\boldsymbol{y}+\boldsymbol{x}\|^2 - \|\boldsymbol{y}-\boldsymbol{x}\|^2 + i\,\|\boldsymbol{y}+i\boldsymbol{x}\|^2 - i\,\|\boldsymbol{y}-i\boldsymbol{x}\|^2\right)^*\\
&\triangleq 4\langle \boldsymbol{y} \mid \boldsymbol{x}\rangle^* && \text{by } \textit{Polarization Identity}
\end{aligned}
$$

(e) Proof that $\langle \alpha\boldsymbol{x} \mid \boldsymbol{y}\rangle = \alpha\langle \boldsymbol{x} \mid \boldsymbol{y}\rangle$ (*homogeneous*): [17]

i. Proof that $\langle \alpha\boldsymbol{x} \mid \boldsymbol{y}\rangle$ is linear in α:

$$
\begin{aligned}
0 &\le \big|\|\alpha\boldsymbol{x}+\boldsymbol{y}\| - \|\beta\boldsymbol{x}+\boldsymbol{y}\|\big| && \text{by Definition F.4 page 206}\\
&\le \big|\|(\alpha\boldsymbol{x}+\boldsymbol{y}) - (\beta\boldsymbol{x}+\boldsymbol{y})\|\big| && \text{by Theorem K.2 page 272}\\
&\le \big|\|(\alpha-\beta)\boldsymbol{x}\|\big|
\end{aligned}
$$

This implies that as $a \to \beta$, $\|\alpha\boldsymbol{x}+\boldsymbol{y}\| \to \|\beta\boldsymbol{x}+\boldsymbol{y}\|$, which by definition implies that $\|\alpha\boldsymbol{x}+\boldsymbol{y}\|$ linear in α. And by the parallelogram law, $\langle \alpha\boldsymbol{x} \mid \boldsymbol{y}\rangle$ is also linear in α.

ii. Proof that $\langle n\boldsymbol{x} \mid \boldsymbol{y}\rangle = n\langle \boldsymbol{x} \mid \boldsymbol{y}\rangle$ for $n \in \mathbb{Z}$ (integer case):

A. Proof for $n = \pm 1$:

$$
\begin{aligned}
\langle n\boldsymbol{x} \mid \boldsymbol{y}\rangle &= \langle \pm 1\boldsymbol{x} \mid \boldsymbol{y}\rangle && \text{by } n = \pm 1 \text{ hypothesis}\\
&= \pm 1\langle \boldsymbol{x} \mid \boldsymbol{y}\rangle && \text{by definition of } \textit{inner product} \quad \text{(Definition J.1 page 255)}\\
&= n\langle \boldsymbol{x} \mid \boldsymbol{y}\rangle && \text{by } n = \pm 1 \text{ hypothesis}
\end{aligned}
$$

[16] Aliprantis and Burkinshaw (1998), page 281
[17] Aliprantis and Burkinshaw (1998), page 138

BY-NC-ND Copyright © 2019 Daniel J. Greenhoe

B. Proof for $n = 0$:

$$\begin{aligned}
\langle n\boldsymbol{x} \mid \boldsymbol{y}\rangle &= \langle 0\boldsymbol{x} \mid \boldsymbol{y}\rangle && \text{by } n = 0 \text{ hypothesis}\\
&= \langle \boldsymbol{x} - \boldsymbol{x} \mid \boldsymbol{y}\rangle\\
&= \langle \boldsymbol{x} \mid \boldsymbol{y}\rangle + \langle -1\boldsymbol{x} \mid \boldsymbol{y}\rangle\\
&= \langle \boldsymbol{x} \mid \boldsymbol{y}\rangle - 1\,\langle \boldsymbol{x} \mid \boldsymbol{y}\rangle\\
&= \langle \boldsymbol{x} \mid \boldsymbol{y}\rangle - \langle \boldsymbol{x} \mid \boldsymbol{y}\rangle\\
&= 0\,\langle \boldsymbol{x} \mid \boldsymbol{y}\rangle\\
&= n\,\langle \boldsymbol{x} \mid \boldsymbol{y}\rangle && \text{by } n = 0 \text{ hypothesis}
\end{aligned}$$

C. Proof for $n = \pm 2$:

$$\begin{aligned}
\langle n\boldsymbol{x} \mid \boldsymbol{y}\rangle &= \langle \pm 2\boldsymbol{x} \mid \boldsymbol{y}\rangle && \text{by } n = \pm 1 \text{ hypothesis}\\
&= \langle \pm(\boldsymbol{x} + \boldsymbol{x}) \mid \boldsymbol{y}\rangle\\
&= \pm\,\langle \boldsymbol{x} + \boldsymbol{x} \mid \boldsymbol{y}\rangle && \text{by definition of } \textit{inner product} \quad \text{(Definition J.1 page 255)}\\
&= \pm\,(\langle \boldsymbol{x} \mid \boldsymbol{y}\rangle + \langle \boldsymbol{x} \mid \boldsymbol{y}\rangle) && \text{by additive property}\\
&= \pm 2\,\langle \boldsymbol{x} \mid \boldsymbol{y}\rangle\\
&= n\,\langle \boldsymbol{x} \mid \boldsymbol{y}\rangle && \text{by } n = \pm 2 \text{ hypothesis}
\end{aligned}$$

D. Proof that [n case] $\implies$ [$n \pm 1$ case]:

$$\begin{aligned}
\langle (n \pm 1)\boldsymbol{x} \mid \boldsymbol{y}\rangle &= \langle n\boldsymbol{x} \pm 1\boldsymbol{x} \mid \boldsymbol{y}\rangle\\
&= \langle n\boldsymbol{x} \mid \boldsymbol{y}\rangle + \langle \pm 1\boldsymbol{x} \mid \boldsymbol{y}\rangle && \text{by additive property}\\
&= n\,\langle \boldsymbol{x} \mid \boldsymbol{y}\rangle \pm 1\,\langle \boldsymbol{x} \mid \boldsymbol{y}\rangle && \text{by left hypothesis}\\
&= (n \pm 1)\,\langle \boldsymbol{x} \mid \boldsymbol{y}\rangle
\end{aligned}$$

iii. Proof that $\langle q\boldsymbol{x} \mid \boldsymbol{y}\rangle = q\,\langle \boldsymbol{x} \mid \boldsymbol{y}\rangle$ for $q \in \mathbb{Q}$ (rational number case):

$$\begin{aligned}
\frac{n}{m}\,\langle \boldsymbol{x} \mid \boldsymbol{y}\rangle &= \frac{n}{m}\left\langle \frac{m}{m}\boldsymbol{x} \mid \boldsymbol{y}\right\rangle && \text{where } n, m \in \mathbb{Z} \text{ and } m \neq 0\\
&= \frac{nm}{m}\left\langle \frac{1}{m}\boldsymbol{x} \mid \boldsymbol{y}\right\rangle && \text{by previous result}\\
&= \frac{m}{m}\left\langle \frac{n}{m}\boldsymbol{x} \mid \boldsymbol{y}\right\rangle && \text{by previous result}\\
&= \left\langle \frac{n}{m}\boldsymbol{x} \mid \boldsymbol{y}\right\rangle
\end{aligned}$$

iv. Proof that $\langle r\boldsymbol{x} \mid \boldsymbol{y}\rangle = r\,\langle \boldsymbol{x} \mid \boldsymbol{y}\rangle$ for all $r \in \mathbb{R}$ (real number case):
Because $\mathbb{Q}$ is dense in $\mathbb{R}$ and because $\|\alpha\boldsymbol{x} + \boldsymbol{y}\|$ is continuous in α, so $\langle \alpha\boldsymbol{x} \mid \boldsymbol{y}\rangle = \alpha\,\langle \boldsymbol{x} \mid \boldsymbol{y}\rangle$ for all $\alpha \in \mathbb{R}$.

v. Proof that $\langle c\boldsymbol{x} \mid \boldsymbol{y}\rangle = c\,\langle \boldsymbol{x} \mid \boldsymbol{y}\rangle$ for all $c \in \mathbb{C}$ (complex number case):
No proof at this time.

✏

Remark J.1. [18] The inner product has already been defined in Definition J.1 (page 255) as a bilinear function that is *non-negative*, *non-isotropic*, *homogeneous*, *additive*, and *conjugate symmetric*. However, given a normed linear space, we could alternatively define the inner product using the *parallelogram law* (Theorem J.7 page 263) together with the *Polarization Identity* (Theorem J.6 page 262). Under this new definition, an inner product *exists* if the parallelogram law is satisfied, and is *specified*, in terms of the norm, by the Polarization Identity.

[18] Loomis (1953), pages 23–24, Kubrusly (2001) page 317

Copyright © 2019 Daniel J. Greenhoe

Of the uncountably infinite number of $\ell^p_{\mathbb{F}}$ norms, only the norm for $p = 2$ induces an inner product (Proposition J.1, next).

Proposition J.1. [19] *Let $\left\|(\!(x_n)\!)_{n\in\mathbb{Z}}\right\|_p$ be the $\ell^p_{\mathbb{F}}$ norm of the sequence $(\!(x_n)\!)$ in the space $\ell^p_{\mathbb{F}}$.*

PRP $\left\|(\!(x_n)\!)\right\|_p$ *induces an inner product* $\iff$ $p = 2$

✎PROOF:

1. Proof that $\|\cdot\|_p$ induces an inner product $\Longleftarrow$ $p = 2$ (using the *Parallelogram law* page 263):

$$
\begin{aligned}
&\|\boldsymbol{x}+\boldsymbol{y}\|_p^2 + \|\boldsymbol{x}-\boldsymbol{y}\|_p^2 \\
&= \|\boldsymbol{x}+\boldsymbol{y}\|_2^2 + \|\boldsymbol{x}-\boldsymbol{y}\|_2^2 && \text{by right hypothesis} \\
&= \left(\sum_{n\in\mathbb{Z}} |x_n+y_n|^2\right)^{\frac{2}{2}} + \left(\sum_{n\in\mathbb{Z}} |x_n-y_n|^2\right)^{\frac{2}{2}} && \text{by definition of } \|\cdot\|_p \\
&= \sum_{n\in\mathbb{Z}} (x_n+y_n)(x_n+y_n)^* + \sum_{n\in\mathbb{Z}} (x_n-y_n)(x_n-y_n)^* \\
&= \sum_{n\in\mathbb{Z}} \left(|x_n|^2 + |y_n|^2 + 2\Re(x_n y_n)\right) + \sum_{n\in\mathbb{Z}} \left(|x_n|^2 + |y_n|^2 - 2\Re(x_n y_n)\right) \\
&= 2\sum_{n\in\mathbb{Z}} |x_n|^2 + 2\sum_{n\in\mathbb{Z}} |y_n|^2 \\
&= 2\,\|\boldsymbol{x}\|_2^2 + 2\,\|\boldsymbol{y}\|_2^2 && \text{by definition of } \|\cdot\|_p \\
&= 2\,\|\boldsymbol{x}\|_p^2 + 2\,\|\boldsymbol{y}\|_p^2 && \text{by right hypothesis} \\
&\Longrightarrow \|\cdot\|_2 \text{ induces an inner product} && \text{by Theorem J.7 page 263}
\end{aligned}
$$

2. Proof that $\|\cdot\|_p$ induces an inner product $\Longrightarrow$ $p = 2$:

   (a) Let $\boldsymbol{x} \triangleq (1,0)$ and $\boldsymbol{y} \triangleq (0,1)$. Then [20]

$$
\begin{aligned}
\|\boldsymbol{x}+\boldsymbol{y}\|_p^2 + \|\boldsymbol{x}-\boldsymbol{y}\|_p^2 &= \left(\sum_{n\in\mathbb{Z}} |x_n+y_n|^p\right)^{\frac{2}{p}} + \left(\sum_{n\in\mathbb{Z}} |x_n-y_n|^p\right)^{\frac{2}{p}} && \text{by definition of } \|\cdot\|_p \\
&= \left(|1+0|^p + |0+1|^p\right)^{\frac{2}{p}} + \left(|1-0|^p + |0-1|^p\right)^{\frac{2}{p}} && \text{by definitions of } \boldsymbol{x} \text{ and } \boldsymbol{y} \\
&= 2^{\frac{2}{p}} + 2^{\frac{2}{p}} \\
&= 2\cdot 2^{\frac{2}{p}} \\
2\,\|\boldsymbol{x}\|_p^2 + 2\,\|\boldsymbol{y}\|_p^2 &= 2\left(\sum_{n\in\mathbb{Z}} |x_n|^p\right)^{\frac{2}{p}} + 2\left(\sum_{n\in\mathbb{Z}} |y_n|^p\right)^{\frac{2}{p}} && \text{by definition of } \|\cdot\|_p \\
&= 2\left(|1|^p + |0|^p\right)^{\frac{2}{p}} + 2\left(|1|^p + |0|^p\right)^{\frac{2}{p}} && \text{by definitions of } \boldsymbol{x} \text{ and } \boldsymbol{y} \\
&= 2+2 \\
&= 4 \\
2\cdot 2^{\frac{2}{p}} = 4 &= \iff 2^{\frac{2}{p}} = 2 \\
&= \Longrightarrow p = 2
\end{aligned}
$$

[19] Kubrusly (2001) pages 318–319 ⟨Example 5B⟩

[20] http://groups.google.com/group/sci.math/msg/531b1173f08871e9

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

(b) Proof that $2^{2/p}$ is monotonic decreasing in p (and so $p = 2$ is the only solution):

$$\begin{aligned}
\frac{\mathrm{d}}{\mathrm{d}p} 2^{\frac{2}{p}} &= \frac{\mathrm{d}}{\mathrm{d}p} e^{\ln 2^{\frac{2}{p}}} \\
&= \left(e^{\ln 2^{\frac{2}{p}}}\right) \frac{\mathrm{d}}{\mathrm{d}p} \ln 2^{\frac{2}{p}} \\
&= \left(2^{\frac{2}{p}}\right) \frac{\mathrm{d}}{\mathrm{d}p} (2 \ln 2) \frac{1}{p} \\
&= \left(2^{\frac{2}{p}}\right) 2 \ln 2 \left(-\frac{1}{p^2}\right) \\
&< 0 \qquad \forall p \in (0, \infty)
\end{aligned}$$



J.3 Orthogonality

Definition J.3.

DEF *The **Kronecker delta function** $\bar{\delta}_n$ is defined as* $\bar{\delta}_n \triangleq \begin{cases} 1 & \text{for } n = 0 \\ 0 & \text{for } n \neq 0 \end{cases}$ *and* $\forall n \in \mathbb{Z}$

Definition J.4. *Let $\left(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle \mid \triangledown \rangle\right)$ be an* INNER PRODUCT SPACE *(Definition J.1 page 255).*

DEF
*Two vectors $\boldsymbol{x}$ and $\boldsymbol{y}$ in X are **orthogonal** if*

$$\langle \boldsymbol{x} \mid \boldsymbol{y} \rangle = \begin{cases} 0 & \text{for } \boldsymbol{x} \neq \boldsymbol{y} \\ c \in \mathbb{F} \backslash 0 & \text{for } \boldsymbol{x} = \boldsymbol{y} \end{cases}$$

The notation $\boldsymbol{x} \perp \boldsymbol{y}$ implies $\boldsymbol{x}$ and $\boldsymbol{y}$ are ORTHOGONAL.
*A set $Y \in 2^X$ is **orthogonal** if $\boldsymbol{x} \perp \boldsymbol{y}$ $\forall_{\boldsymbol{x},\boldsymbol{y} \in Y}$.*
*A set Y is **orthonomal** if it is* ORTHOGONAL *and $\langle \boldsymbol{y} \mid \boldsymbol{y} \rangle = 1$ $\forall_{\boldsymbol{y} \in Y}$.*
*A sequence $(\boldsymbol{x}_n \in X)_{n \in \mathbb{Z}}$ is **orthogonal** if $\langle \boldsymbol{x}_n \mid \boldsymbol{x}_m \rangle = c\bar{\delta}_{nm}$ for some $c \in \mathbb{R} \backslash 0$.*
*A sequence $(\boldsymbol{x}_n \in X)_{n \in \mathbb{Z}}$ is **orthonormal** if $\langle \boldsymbol{x}_n \mid \boldsymbol{x}_m \rangle = \bar{\delta}_{nm}$.*

The definition of the orthogonality relation $\perp$ has several immediate consequences (next theorem):

Theorem J.8. [21] *Let $\left(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle \mid \triangledown \rangle\right)$ be an* INNER PRODUCT SPACE.

THM

1.	$\boldsymbol{x} \perp \boldsymbol{x} \iff \boldsymbol{x} = \mathbb{0}$	$\forall_{\boldsymbol{x} \in X}$	
2.	$\boldsymbol{x} \perp \boldsymbol{y} \implies \alpha \boldsymbol{x} \perp \boldsymbol{y}$	$\forall_{\alpha \in \mathbb{R}, \boldsymbol{x}, \boldsymbol{y} \in X}$	(HOMOGENEOUS)
3.	$\boldsymbol{x} \perp \boldsymbol{y} \iff \boldsymbol{y} \perp \boldsymbol{x}$	$\forall_{\boldsymbol{x}, \boldsymbol{y} \in X}$	(SYMMETRY)
4.	$\boldsymbol{x} \perp \boldsymbol{y}$ *and* $\boldsymbol{y} \perp \boldsymbol{z} \implies \boldsymbol{x} \perp (\boldsymbol{y} + \boldsymbol{z})$	$\forall_{\boldsymbol{x}, \boldsymbol{y}, \boldsymbol{z} \in X}$	(ADDITIVE)
5.	$\exists \beta \in \mathbb{R}$ such that $\boldsymbol{x} \perp (\beta \boldsymbol{x} + \boldsymbol{y})$	$\forall_{\boldsymbol{x} \in X \backslash \mathbb{0}, \boldsymbol{y} \in X}$	

Theorem J.9. *Let $\left(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle \mid \triangledown \rangle\right)$ be an* INNER PRODUCT SPACE.

THM

$$\left.\begin{array}{lll} 1. & \langle \boldsymbol{x} \mid \boldsymbol{y} \rangle = 0 & \textit{and} \\ 2. & \boldsymbol{x} + \boldsymbol{y} = \mathbb{0} & \end{array}\right\} \iff \left\{\begin{array}{lll} 1. & \boldsymbol{x} = \mathbb{0} & \textit{and} \\ 2. & \boldsymbol{y} = \mathbb{0} & \end{array}\right. \quad \forall_{\boldsymbol{x}, \boldsymbol{y} \in X}$$

PROOF:

[21] James (1945), page 292, Drljević (1989) page 232

COPYRIGHT © 2019 DANIEL J. GREENHOE
CC BY-NC-ND

1. Proof that $\boldsymbol{x} = \boldsymbol{y} = \mathbb{0}$:

$$\begin{aligned}
0 &= \langle \mathbb{0} \mid \mathbb{0} \rangle && \text{by } \textit{non-isotropic} \text{ property of } \langle \triangle \mid \triangledown \rangle \text{ (Definition J.1 page 255)} \\
&= \langle \boldsymbol{x}+\boldsymbol{y} \mid \boldsymbol{x}+\boldsymbol{y} \rangle && \text{by left hypothesis 2} \\
&= \langle \boldsymbol{x} \mid \boldsymbol{x}+\boldsymbol{y} \rangle + \langle \boldsymbol{y} \mid \boldsymbol{x}+\boldsymbol{y} \rangle && \text{by } \textit{additive} \text{ property of } \langle \triangle \mid \triangledown \rangle \text{ (Definition J.1 page 255)} \\
&= \langle \boldsymbol{x} \mid \boldsymbol{x} \rangle + \langle \boldsymbol{x} \mid \boldsymbol{y} \rangle + \langle \boldsymbol{x} \mid \boldsymbol{y} \rangle^* + \langle \boldsymbol{y} \mid \boldsymbol{y} \rangle && \text{by } \textit{conjugate symmetric} \text{ and } \textit{additive} \text{ properties of } \langle \triangle \mid \triangledown \rangle \\
&= \underbrace{\langle \boldsymbol{x} \mid \boldsymbol{x} \rangle}_{\geq 0} + 0 + 0 + \underbrace{\langle \boldsymbol{y} \mid \boldsymbol{y} \rangle}_{\geq 0} && \text{by left hypothesis 1} \\
\implies \boldsymbol{x} &= \mathbb{0} \text{ and } \boldsymbol{y} = \mathbb{0} && \text{by } \textit{non-negative} \text{ and } \textit{non-isotropic} \text{ properties of } \langle \triangle \mid \triangledown \rangle
\end{aligned}$$

2. Proof that $\langle \boldsymbol{x} \mid \boldsymbol{y} \rangle = 0$:

$$\begin{aligned}
\langle \boldsymbol{x} \mid \boldsymbol{y} \rangle &= \langle \mathbb{0} \mid \mathbb{0} \rangle && \text{by right hypotheses} \\
&= 0 && \text{by } \textit{non-isotropic} \text{ property of } \langle \triangle \mid \triangledown \rangle \text{ (Definition J.1 page 255)}
\end{aligned}$$

3. Proof that $\boldsymbol{x} + \boldsymbol{y} = \mathbb{0}$:

$$\begin{aligned}
\boldsymbol{x} + \boldsymbol{y} &= \mathbb{0} + \mathbb{0} && \text{by right hypotheses} \\
&= \mathbb{0}
\end{aligned}$$

✏

The *triangle inequality for vectors* in a *normed linear space* (Theorem K.1 page 271) demonstrates that $\left\| \sum_{n=1}^{N} \boldsymbol{x}_n \right\| \leq \sum_{n=1}^{N} \|\boldsymbol{x}_n\|$. The *Pythagorean Theorem* (next) demonstrates that this *inequality* becomes *equality* when the set $\{\boldsymbol{x}_n\}$ is *orthogonal*.

Theorem J.10 (Pythagorean Theorem). [22] *Let* $\{\boldsymbol{x}_n \in X \mid_{n=1,2,\ldots,N}\}$ *be a set of vectors in an* INNER PRODUCT SPACE *(Definition J.1 page 255)* $\big(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle \mid \triangledown \rangle\big)$ *and let* $\|\boldsymbol{x}\| \triangleq \sqrt{\langle \boldsymbol{x} \mid \boldsymbol{x} \rangle}$ *(Definition J.2 page 260)*.

THM

$$\{\boldsymbol{x}_n\} \textit{ is } \text{ORTHOGONAL} \iff \left\| \sum_{n=1}^{N} \boldsymbol{x}_n \right\|^2 = \sum_{n=1}^{N} \|\boldsymbol{x}_n\|^2 \quad \forall N \in \mathbb{N}$$

✎PROOF: 1. Proof for ($\implies$) case:

$$\begin{aligned}
\left\| \sum_{n=1}^{N} \boldsymbol{x}_n \right\|^2 &= \left\langle \sum_{n=1}^{N} \boldsymbol{x}_n \,\middle|\, \sum_{m=1}^{N} \boldsymbol{x}_m \right\rangle && \text{by def. of } \|\cdot\| && \text{(Definition K.1 page 271)} \\
&= \sum_{n=1}^{N} \sum_{m=1}^{N} \langle \boldsymbol{x}_n \mid \boldsymbol{x}_m \rangle && \text{by def. of } \langle \triangle \mid \triangledown \rangle && \text{(Definition J.1 page 255)} \\
&= \sum_{n=1}^{N} \sum_{m=1}^{N} \langle \boldsymbol{x}_n \mid \boldsymbol{x}_m \rangle \bar{\delta}_{n-m} && \text{by left hypothesis} \\
&= \sum_{n=1}^{N} \langle \boldsymbol{x}_n \mid \boldsymbol{x}_n \rangle && \text{by def. of } \bar{\delta} && \text{(Definition J.3 page 267)} \\
&= \sum_{n=1}^{N} \|\boldsymbol{x}_n\|^2 && \text{by def. of } \|\cdot\| && \text{(Definition K.1 page 271)}
\end{aligned}$$

[22] Aliprantis and Burkinshaw (1998) pages 282–283 ⟨Theorem 32.7⟩, Kubrusly (2001) page 324 ⟨Proposition 5.8⟩, Bollobás (1999) pages 132–133 ⟨Theorem 3⟩

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

2. Proof for ($\Longleftarrow$) case:

$$
\begin{aligned}
&4\langle \boldsymbol{x} \mid \boldsymbol{y}\rangle \\
&= \|\boldsymbol{x}+\boldsymbol{y}\|^2 - \|\boldsymbol{x}-\boldsymbol{y}\|^2 + i\,\|\boldsymbol{x}+i\boldsymbol{y}\|^2 - i\,\|\boldsymbol{x}-i\boldsymbol{y}\|^2 \quad \text{by } \textit{polarization identity} \text{ (Theorem J.6 page 262)} \\
&= \left(\|\boldsymbol{x}\|^2+\|\boldsymbol{y}\|^2\right) - \left(\|\boldsymbol{x}\|^2+\|-\boldsymbol{y}\|^2\right) + i\left(\|\boldsymbol{x}\|^2+\|i\boldsymbol{y}\|^2\right) - i\left(\|\boldsymbol{x}\|^2+\|-i\boldsymbol{y}\|^2\right) \quad \text{by right hypothesis} \\
&= \left(\|\boldsymbol{x}\|^2+\|\boldsymbol{y}\|^2\right) - \left(\|\boldsymbol{x}\|^2+|-1|^2\,\|\boldsymbol{y}\|^2\right) + i\left(\|\boldsymbol{x}\|^2+|i|^2\,\|\boldsymbol{y}\|^2\right) - i\left(\|\boldsymbol{x}\|^2+|-i|^2\,\|\boldsymbol{y}\|^2\right) \quad \text{by definition of } \|\cdot\| \\
&= \left(\|\boldsymbol{x}\|^2+\|\boldsymbol{y}\|^2\right) - \left(\|\boldsymbol{x}\|^2+\|\boldsymbol{y}\|^2\right) + i\left(\|\boldsymbol{x}\|^2+\|\boldsymbol{y}\|^2\right) - i\left(\|\boldsymbol{x}\|^2+\|\boldsymbol{y}\|^2\right) \quad \text{by def. of } |\cdot| \text{ (Definition F.4 page 206)} \\
&= 0
\end{aligned}
$$

✍

Copyright © 2019 Daniel J. Greenhoe

Copyright © 2019 Daniel J. Greenhoe

APPENDIX K

NORMED LINEAR SPACES

K.1 Definition and basic results

Definition K.1. [1] *Let* $\big(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times})\big)$ *be a* LINEAR SPACE *(Definition G.1 page 207) and* $|\cdot| \in \mathbb{R}^{\mathbb{F}}$ *the* ABSOLUTE VALUE *function (Definition F.4 page 206).*

DEF

A functional $\|\cdot\|$ *in* $\mathbb{R}^{X}$ *is a* ***norm*** *if*

1.	$\|\boldsymbol{x}\| \geq 0$	$\forall \boldsymbol{x} \in X$	(STRICTLY POSITIVE)	*and*
2.	$\|\boldsymbol{x}\| = 0 \iff \boldsymbol{x} = \mathbb{0}$	$\forall \boldsymbol{x} \in X$	(NONDEGENERATE)	*and*
3.	$\|\alpha \boldsymbol{x}\| = \lvert\alpha\rvert \, \|\boldsymbol{x}\|$	$\forall \boldsymbol{x} \in X, \alpha \in \mathbb{C}$	(HOMOGENEOUS)	*and*
4.	$\|\boldsymbol{x} + \boldsymbol{y}\| \leq \|\boldsymbol{x}\| + \|\boldsymbol{y}\|$	$\forall \boldsymbol{x}, \boldsymbol{y} \in X$	(SUBADDITIVE/TRIANGLE INQUALITY).	

A ***normed linear space*** *is the tuple* $\big(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \|\cdot\|\big)$.

The definition of the *norm* (Definition K.1 page 271) requires that any two vectors in a norm space be *sub-additive* (they satisfy the *triangle inequality* property) such that $\|\boldsymbol{x}_1 + \boldsymbol{x}_2\| \leq \|\boldsymbol{x}_1\| + \|\boldsymbol{x}_2\|$. Acutally, in **any** normed linear space, this property holds true for **any** finite number of vectors—not just two—such that $\|\boldsymbol{x}_1 + \boldsymbol{x}_2 + \cdots + \boldsymbol{x}_N\| \leq \|\boldsymbol{x}_1\| + \|\boldsymbol{x}_2\| + \cdots + \|\boldsymbol{x}_N\|$ (next theorem).

Theorem K.1 (triangle inequality)**.** [2] *Let* $(\!|\boldsymbol{x}_n \in X|\!)_1^N$ *be an* N-TUPLE *(Definition Q.1 page 353) of vectors in a* NORMED LINEAR SPACE $\big(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \|\cdot\|\big)$.

THM

$$\left\| \sum_{n=1}^{N} \boldsymbol{x}_n \right\| \leq \sum_{n=1}^{N} \|\boldsymbol{x}_n\| \qquad \forall N \in \mathbb{N}, \boldsymbol{x}_n \in V$$

✎PROOF: Proof is by induction:

[1] Aliprantis and Burkinshaw (1998), pages 217–218, Banach (1932a), page 53, Banach (1932b), page 33, Banach (1922) page 135

[2] Michel and Herget (1993), page 344, Euclid (circa 300BC) ⟨Book I Proposition 20⟩

1. Proof for the $N = 1$ case:

$$\begin{aligned}\left\|\sum_{n=1}^{1} \boldsymbol{x}_n\right\| &= \|\boldsymbol{x}_1\| \\ &= \sum_{n=1}^{1} \|\boldsymbol{x}_1\|\end{aligned}$$

2. Proof for the $N = 2$ case:

$$\begin{aligned}\left\|\sum_{n=1}^{2} \boldsymbol{x}_n\right\| &= \left\|\sum_{n=1}^{2} \boldsymbol{x}_n\right\| \\ &= \|\boldsymbol{x}_1 + \boldsymbol{x}_2\| \\ &\le \|\boldsymbol{x}_1\| + \|\boldsymbol{x}_2\| && \text{by Definition K.1 page 271 (triangle inequality)} \\ &= \sum_{n=1}^{2} \|\boldsymbol{x}_n\|\end{aligned}$$

3. Proof that [N case] $\implies$ [$N+1$ case]:

$$\begin{aligned}\left\|\sum_{n=1}^{N+1} \boldsymbol{x}_n\right\| &= \left\|\sum_{n=1}^{N} \boldsymbol{x}_n + \boldsymbol{x}_{n+1}\right\| \\ &\le \left\|\sum_{n=1}^{N} \boldsymbol{x}_n\right\| + \|\boldsymbol{x}_{n+1}\| && \text{by Definition K.1 page 271 (triangle inequality)} \\ &\le \sum_{n=1}^{N} \|\boldsymbol{x}_n\| + \|\boldsymbol{x}_{n+1}\| && \text{by left hypothesis} \\ &= \sum_{n=1}^{N+1} \|\boldsymbol{x}_n\|\end{aligned}$$

✏

Theorem K.2 (Reverse Triangle Inequality). [3] *Let* $\big(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \|\cdot\|\big)$ *be a* NORMED LINEAR SPACE *(Definition K.1 page 271)*.

THM

$$\underbrace{\big|\|\boldsymbol{x}\| - \|\boldsymbol{y}\|\big| \le \|\boldsymbol{x} - \boldsymbol{y}\|}_{\text{REVERSE TRIANGLE INEQUALITY}} \quad \le \quad \|\boldsymbol{x}\| + \|\boldsymbol{y}\| \qquad \forall \boldsymbol{x},\boldsymbol{y} \in X$$

✎PROOF:

$$\begin{aligned}\big|\|\boldsymbol{x}\| - \|\boldsymbol{y}\|\big| &= \big|\|(\boldsymbol{x} - \boldsymbol{y}) + \boldsymbol{y}\| - \|\boldsymbol{y}\|\big| \\ &\le \big|\|\boldsymbol{x} - \boldsymbol{y}\| + \|\boldsymbol{y}\| - \|\boldsymbol{y}\|\big| && \text{by Definition K.1 page 271} \\ &= \big|\|\boldsymbol{x} - \boldsymbol{y}\|\big| \\ &= \|\boldsymbol{x} - \boldsymbol{y}\| && \text{by Definition K.1 page 271}\end{aligned}$$

$$\begin{aligned}\|\boldsymbol{x} - \boldsymbol{y}\| &\le \|\boldsymbol{x} - \mathbb{0}\| + \|\mathbb{0} - \boldsymbol{y}\| && \text{by previous result with } \boldsymbol{u} = 0 \\ &= \|\boldsymbol{x}\| + |-1|\,\|\boldsymbol{y}\| && \text{by Definition K.1 page 271} \\ &= \|\boldsymbol{x}\| + \|\boldsymbol{y}\|\end{aligned}$$

✏

[3] Aliprantis and Burkinshaw (1998), page 218, Giles (2000) page 2, Banach (1922) page 136

BY-NC-ND
COPYRIGHT © 2019 DANIEL J. GREENHOE



The shortest distance between two vectors is always the difference of the vectors. This is proven in next and illustrated to the left in the Euclidean space $\mathbb{R}^2$ (the plane)

Proposition K.1. [4] *Let* $\big(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \|\cdot\|\big)$ *be a* NORMED LINEAR SPACE *(Definition K.1 page 271)*.

PRP

$$\|\boldsymbol{x}-\boldsymbol{y}\| \quad \leq \quad \|\boldsymbol{x}-\boldsymbol{u}\| + \|\boldsymbol{u}-\boldsymbol{y}\| \qquad \forall_{\boldsymbol{x},\boldsymbol{y},\boldsymbol{u}\in X}$$

✎PROOF:

$$\begin{aligned}\|\boldsymbol{x}-\boldsymbol{y}\| &= \|(\boldsymbol{x}-\boldsymbol{u})+(\boldsymbol{u}-\boldsymbol{y})\| \\ &\leq \|\boldsymbol{x}-\boldsymbol{u}\| + \|\boldsymbol{u}-\boldsymbol{y}\| \qquad \text{by Definition K.1 page 271}\end{aligned}$$

✏

Example K.1 (The usual norm). [5] Let $\mathbb{R}^{\mathbb{R}}$ be the set of all functions with domain and range the set of *real numbers* $\mathbb{R}$.

EX The **absolute value** (Definition F.4 page 206) $|\cdot| \in \mathbb{R}^{\mathbb{R}}$ is a *norm*.

Example K.2 (l_p norms). Let $(\!(x_n)\!)_{n\in\mathbb{Z}}$ be a *sequence* (Definition Q.1 page 353) of real numbers. An uncountably infinite number of norms is provided by the $\ell^{\mathsf{p}}_{\mathbb{F}}$ norms $\|(\!(x_n)\!)\|_p$:

EX $$\|(\!(x_n)\!)\|_p \triangleq \left(\sum_{n\in\mathbb{Z}} |x_n|^p\right)^{\frac{1}{p}} \qquad \text{is a norm for } p\in[1:\infty]$$

K.2 Relationship between metrics and norms

K.2.1 Metrics generated by norms

The concept of *length* is very closely related to the concept of *distance*. Thus it is not surprising that a *norm* (a "length" function) can be used to define a *metric* (a "distance" function) on any *metric linear space* (Definition ?? page ??). Another way to say this is that the norm of a normed linear space *induces* a metric on this space. And so every normed linear space also has a metric. And because every normed linear space has a metric, **every normed linear space is also a metric space**. Actually this can be generalized one step further in that every metric space is also a *topological space*. And so **every normed linear space is also a topological space.** In symbols,

$$\text{normed linear space} \quad\Longrightarrow\quad \text{metric space} \quad\Longrightarrow\quad \text{topological space.}$$

[4] Aliprantis and Burkinshaw (1998), page 218
[5] Giles (1987), page 3

COPYRIGHT © 2019 DANIEL J. GREENHOE

Theorem K.3. [6] *Let* $\mathsf{d} \in \mathbb{R}^{X \times X}$ *be a function on a* REAL *normed linear space* $\big(X, +, \cdot, (\mathbb{R}, \dot{+}, \dot{\times}), \|\cdot\|\big)$. *Let* $\mathsf{B}(\boldsymbol{x}, r) \triangleq \{\boldsymbol{y} \in \boldsymbol{X} \,|\, \|\boldsymbol{y} - \boldsymbol{x}\| < r\}$ *be the* OPEN BALL *of radius* r *centered at a point* $\boldsymbol{x}$.

THM $\mathsf{d}(\boldsymbol{x}, \boldsymbol{y}) \triangleq \|\boldsymbol{x} - \boldsymbol{y}\|$ *is a metric on* $\boldsymbol{X}$

✎PROOF: The proof follows directly from the definition of a metric (not included in this text) the definition of *norm* (Definition K.1 page 271). ✏

The previous theorem defined a metric $\mathsf{d}(\boldsymbol{x}, \boldsymbol{y})$ induced by the norm $\|\boldsymbol{x}\|$. The next definition defines this metric formally.

Definition K.2. [7] *Let* $\big(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \|\cdot\|\big)$ *be a* NORMED LINEAR SPACE *(Definition K.1 page 271)*.

DEF *The* ***metric induced by the norm*** $\|\cdot\|$ *is the function* $\mathsf{d} \in \mathbb{R}^{X}$ *such that*

$$\mathsf{d}(\boldsymbol{x}, \boldsymbol{y}) \triangleq \|\boldsymbol{x} - \boldsymbol{y}\| \qquad \forall \boldsymbol{x}, \boldsymbol{y} \in X$$

Due to its algebraic structure, every norm is *continuous* (Corollary K.1 page 274). This makes norm spaces very useful in analysis. For a function f be to *continuous*, for every $\epsilon > 0$ there must exist a $\delta > 0$ such that $|\mathsf{f}(x + \delta) - \mathsf{f}(x)| < \epsilon$. The *Reverse Triangle Inequality* (Theorem K.2 page 272) shows this to be true when $\mathsf{f}(\cdot) \triangleq \|\cdot\|$.

Corollary K.1. [8] *Let* $\boldsymbol{\Omega} \triangleq \big(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \|\cdot\|\big)$ *be a* NORMED LINEAR SPACE *(Definition K.1 page 271)*.

COR *The norm* $\|\cdot\|$ *is* CONTINUOUS *in* $\boldsymbol{\Omega}$.

✎PROOF: This follows from these concepts:

1. The fact that $\mathsf{d}(\boldsymbol{x}, \boldsymbol{y}) \triangleq \|\boldsymbol{x} - \boldsymbol{y}\|$ is a *metric* (Theorem K.3 page 274).
2. *Continuity* in a metric space.
3. The *Reverse Triangle Inequality* (Theorem K.2 page 272).

✏

Theorem K.4 (next) demonstrates that **all open or closed** balls in **any normed linear space** are *convex*. However, the converse is not true—that is, a metric not generated by a norm may still produce a ball that is convex.

Theorem K.4. [9] *Let* $\big(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \mathsf{d}\big)$ *be a* METRIC LINEAR SPACE *(Definition* **??** *page* **??***). Let* B *be the* OPEN BALL $\mathsf{B}(\boldsymbol{p}, r) \triangleq \{\boldsymbol{x} \in \boldsymbol{X} \,|\, \mathsf{d}(\boldsymbol{p}, \boldsymbol{x}) < r\}$ *(open ball with respect to metric* d *centered at point* $\boldsymbol{p}$ *and with radius* r*).*

THM

$$\left.\begin{array}{l} \exists \|\cdot\| \in \mathbb{R}^{X} \quad \text{such that} \\ \underbrace{\mathsf{d}(\boldsymbol{x}, \boldsymbol{y}) = \|\boldsymbol{y} - \boldsymbol{x}\|}_{\mathsf{d}\ \textit{is generated by a norm}} \end{array}\right\} \Longrightarrow \left\{\begin{array}{ll} 1. & \mathsf{B}(\boldsymbol{x}, r) = \boldsymbol{x} + B(\mathbb{0}, r) \\ 2. & \mathsf{B}(\mathbb{0}, r) = r\,\mathsf{B}(\mathbb{0}, 1) \\ 3. & \mathsf{B}(\boldsymbol{x}, r) \ \textit{is}\ \text{CONVEX} \\ 4. & \boldsymbol{x} \in \mathsf{B}(\mathbb{0}, r) \iff -\boldsymbol{x} \in \mathsf{B}(\mathbb{0}, r) \quad \text{(SYMMETRIC)} \end{array}\right.$$

[6] Michel and Herget (1993), page 344, Banach (1932a) page 53
[7] Giles (2000) page 1 ⟨1.1 Definition⟩
[8] Giles (2000) page 2
[9] Giles (2000) page 2 ⟨1.2 Remarks⟩, Giles (1987) pages 22–26 ⟨2.4 Theorem, 2.11 Theorem⟩

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

✎PROOF:

1. Proof that $\mathsf{d}(\boldsymbol{x}+\boldsymbol{z}, \boldsymbol{y}+\boldsymbol{v}\boldsymbol{z}) = \mathsf{d}(\boldsymbol{x}, \boldsymbol{y})$ (invariant):

$$\begin{aligned}
\mathsf{d}(\boldsymbol{x}+\boldsymbol{z}, \boldsymbol{y}+\boldsymbol{v}\boldsymbol{z}) &= \|(\boldsymbol{y}+\boldsymbol{v}\boldsymbol{z}) - (\boldsymbol{x}+\boldsymbol{z})\| && \text{by left hypothesis}\\
&= \|\boldsymbol{y}-\boldsymbol{x}\| \\
&= \mathsf{d}(\boldsymbol{x}, \boldsymbol{y}) && \text{by left hypothesis}
\end{aligned}$$

2. Proof that $\mathsf{B}(\boldsymbol{x}, r) = \boldsymbol{x} + B(0, r)$:

$$\begin{aligned}
\mathsf{B}(\boldsymbol{x}, r) &= \{\boldsymbol{y} \in \boldsymbol{X} | \mathsf{d}(\boldsymbol{x}, \boldsymbol{y}) < r\} && \text{by definition of open ball } B\\
&= \{\boldsymbol{y} \in \boldsymbol{X} | \mathsf{d}(\boldsymbol{y}-\boldsymbol{y}, \boldsymbol{y}-\boldsymbol{x}) < r\} && \text{by right result 1.}\\
&= \{\boldsymbol{y} \in \boldsymbol{X} | \mathsf{d}(\mathbb{0}, \boldsymbol{y}-\boldsymbol{x}) < r\} \\
&= \{\boldsymbol{u}+\boldsymbol{x} \in \boldsymbol{X} | \mathsf{d}(\mathbb{0}, \boldsymbol{u}) < r\} && \text{let } \boldsymbol{u} \triangleq \boldsymbol{y}-\boldsymbol{x}\\
&= \boldsymbol{x} + \{\boldsymbol{u} \in \boldsymbol{X} | \mathsf{d}(\mathbb{0}, \boldsymbol{u}) < r\} \\
&= \boldsymbol{x} + B(0, r) && \text{by definition of open ball } B
\end{aligned}$$

3. Proof that $\mathsf{B}(\mathbb{0}, r) = r\,\mathsf{B}(\mathbb{0}, 1)$:

$$\begin{aligned}
\mathsf{B}(\mathbb{0}, r) &= \{\boldsymbol{y} \in \boldsymbol{X} | \mathsf{d}(\mathbb{0}, \boldsymbol{y}) < r\} && \text{by definition of open ball } B\\
&= \left\{\boldsymbol{y} \in \boldsymbol{X} | \frac{1}{r}\mathsf{d}(\mathbb{0}, \boldsymbol{y}) < 1\right\} \\
&= \left\{\boldsymbol{y} \in \boldsymbol{X} | \frac{1}{r}\,\|\boldsymbol{y}-\mathbb{0}\| < 1\right\} && \text{by left hypothesis}\\
&= \left\{\boldsymbol{y} \in \boldsymbol{X} | \left\|\frac{1}{r}\boldsymbol{y} - \frac{1}{r}\mathbb{0}\right\| < 1\right\} && \text{by homogeneous property of } \|\cdot\| \text{ page 271}\\
&= \left\{\boldsymbol{y} \in \boldsymbol{X} | \mathsf{d}\left(\frac{1}{r}\mathbb{0}, \frac{1}{r}\boldsymbol{y}\right) < 1\right\} && \text{by left hypothesis}\\
&= \{r\boldsymbol{u} \in \boldsymbol{X} | \mathsf{d}(\mathbb{0}, \boldsymbol{u}) < 1\} && \text{let } \boldsymbol{u} \triangleq \frac{1}{r}\boldsymbol{y}\\
&= r\,\{\boldsymbol{u} \in \boldsymbol{X} | \mathsf{d}(\mathbb{0}, \boldsymbol{u}) < 1\} \\
&= r\,B(\mathbb{0}, 1) && \text{by definition of open ball } B
\end{aligned}$$

4. Proof that $\mathsf{B}(\boldsymbol{p}, r)$ is convex:
We must prove that for any pair of points $\boldsymbol{x}$ and $\boldsymbol{y}$ in the open ball $\mathsf{B}(\boldsymbol{p}, r)$, any point $\lambda\boldsymbol{x} + (1-\lambda)\boldsymbol{y}$ is also in the open ball. That is, the distance from any point $\lambda\boldsymbol{x} + (1-\lambda)\boldsymbol{y}$ to the ball's center $\boldsymbol{p}$ must be less than r.

$$\begin{aligned}
\mathsf{d}(\boldsymbol{p}, \lambda\boldsymbol{x} + (1-\lambda)\boldsymbol{y}) &= \|\boldsymbol{p} - \lambda\boldsymbol{x} - (1-\lambda)\boldsymbol{y}\| && \text{by left hypothesis}\\
&= \left\|\underbrace{\lambda\boldsymbol{p} + (1-\lambda)\boldsymbol{p}}_{\boldsymbol{p}} - \lambda\boldsymbol{x} - (1-\lambda)\boldsymbol{y}\right\| \\
&= \|\lambda\boldsymbol{p} - \lambda\boldsymbol{x} + (1-\lambda)\boldsymbol{p} - (1-\lambda)\boldsymbol{y}\| \\
&\le \|\lambda\boldsymbol{p} - \lambda\boldsymbol{x}\| + \|(1-\lambda)\boldsymbol{p} - (1-\lambda)\boldsymbol{y}\| && \text{by subadditivity property of } \|\cdot\| \text{ page 271}\\
&= |\lambda|\,\|\boldsymbol{p}-\boldsymbol{x}\| + |1-\lambda|\,\|\boldsymbol{p}-\boldsymbol{y}\| && \text{by homogeneous property of } \|\cdot\| \text{ page 271}\\
&= \lambda\,\|\boldsymbol{p}-\boldsymbol{x}\| + (1-\lambda)\,\|\boldsymbol{p}-\boldsymbol{y}\| && \text{because } 0 \le \lambda \le 1\\
&\le \lambda r + (1-\lambda)r && \text{because } \boldsymbol{x}, \boldsymbol{y} \text{ are in the ball } \mathsf{B}(\boldsymbol{p}, r)\\
&= r
\end{aligned}$$

COPYRIGHT © 2019 DANIEL J. GREENHOE
CC BY-NC-ND

5. Proof that $\boldsymbol{x} \in \mathsf{B}(\mathbb{0}, r) \iff -\boldsymbol{x} \in \mathsf{B}(\mathbb{0}, r)$ (symmetric):

$$
\begin{aligned}
\boldsymbol{x} \in \mathsf{B}(\mathbb{0}, r) &\iff \boldsymbol{x} \in \{\boldsymbol{y} \in \boldsymbol{X} \,|\, \mathsf{d}(\mathbb{0}, \boldsymbol{y}) < r\} && \text{by definition of open ball } B \\
&\iff \boldsymbol{x} \in \{\boldsymbol{y} \in \boldsymbol{X} \,|\, \|\boldsymbol{y} - \mathbb{0}\| < r\} && \text{by left hypothesis} \\
&\iff \boldsymbol{x} \in \{\boldsymbol{y} \in \boldsymbol{X} \,|\, \|\boldsymbol{y}\| < r\} \\
&\iff \boldsymbol{x} \in \{\boldsymbol{y} \in \boldsymbol{X} \,|\, \|(-1)(-\boldsymbol{y})\| < r\} \\
&\iff \boldsymbol{x} \in \{\boldsymbol{y} \in \boldsymbol{X} \,|\, |-1|\, \|-\boldsymbol{y}\| < r\} && \text{by homogeneous property of } \|\cdot\| \text{ page 271} \\
&\iff \boldsymbol{x} \in \{\boldsymbol{y} \in \boldsymbol{X} \,|\, \|-\boldsymbol{y} - \mathbb{0}\| < r\} \\
&\iff \boldsymbol{x} \in \{\boldsymbol{y} \in \boldsymbol{X} \,|\, \mathsf{d}(\mathbb{0}, -\boldsymbol{y}) < r\} && \text{by left hypothesis} \\
&\iff \boldsymbol{x} \in \{-\boldsymbol{u} \in \boldsymbol{X} \,|\, \mathsf{d}(\mathbb{0}, \boldsymbol{u}) < r\} && \text{let } \boldsymbol{u} \triangleq -\boldsymbol{y} \\
&\iff \boldsymbol{x} \in (-\{\boldsymbol{u} \in \boldsymbol{X} \,|\, \mathsf{d}(\mathbb{0}, \boldsymbol{u}) < r\}) \\
&\iff \boldsymbol{x} \in (-\mathsf{B}(\mathbb{0}, r)) \\
&\iff -\boldsymbol{x} \in \mathsf{B}(\mathbb{0}, r)
\end{aligned}
$$

✏

Theorem K.4 (page 274) demonstrates that if a metric d in a metric space $\big(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \mathsf{d}\big)$ is generated by a norm, then the ball $B(x, r)$ in that metric linear space is *convex*. However, the converse is not true. That is, it is possible for the balls in a metric space $(\boldsymbol{Y}, \mathsf{p})$ to be *convex*, but yet the metric p not be generated by a norm.



Figure K.1: Open balls in $(\mathbb{R}^0, \mathsf{d}_n)$, $(\mathbb{R}, \mathsf{d}_n)$, $(\mathbb{R}^2, \mathsf{d}_n)$, and $(\mathbb{R}^3, \mathsf{d}_n)$.

K.2.2 Norms generated by metrics

Every normed linear space is also a metric linear space (Theorem K.3 page 274). That is, a metric linear space generates a *normed linear space*. However, the converse is not true—not every metric linear space is a *normed linear space*. A characterization of metric linear spaces that *are* normed linear spaces is given by Theorem K.5 page 277.

Lemma K.1. [10] *Let* $\big(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \mathsf{d}\big)$ *be a* METRIC LINEAR SPACE. *Let* $\|\boldsymbol{x}\| \triangleq \mathsf{d}(\boldsymbol{x}, \mathbb{0})\ \forall \boldsymbol{x} \in X$.

[10] Oikhberg and Rosenthal (2007), page 599

BY-NC-ND Copyright © 2019 Daniel J. Greenhoe

LEM

$$\underbrace{\mathsf{d}(\boldsymbol{x}+\boldsymbol{z},\boldsymbol{y}+\boldsymbol{z})=\mathsf{d}(\boldsymbol{x},\boldsymbol{y}) \quad \forall \boldsymbol{x},\boldsymbol{y},\boldsymbol{z}\in X}_{\text{TRANSLATION INVARIANT}} \implies \begin{cases} 1. & \|\boldsymbol{x}\| = \|-\boldsymbol{x}\| & \forall \boldsymbol{x}\in X & \textit{and} \\ 2. & \|\boldsymbol{x}\| = 0 \iff \boldsymbol{x}=0 & \forall \boldsymbol{x}\in X & \textit{and} \\ 3. & \|\boldsymbol{x}+\boldsymbol{y}\| \le \|\boldsymbol{x}\|+\|\boldsymbol{y}\| & \forall \boldsymbol{x},\boldsymbol{y}\in X & \end{cases}$$

✎PROOF:

1. Proof that $\|\boldsymbol{x}\| = \|-\boldsymbol{x}\|$:

$$\begin{aligned} \|\boldsymbol{x}\| &= \mathsf{d}(\boldsymbol{x},\mathbb{0}) && \text{by definition of } \|\cdot\| \\ &= \mathsf{d}(\boldsymbol{x}-\boldsymbol{x},\mathbb{0}-\boldsymbol{x}) && \text{by translation invariance hypothesis} \\ &= \mathsf{d}(\mathbb{0},-\boldsymbol{x}) \\ &= \|-\boldsymbol{x}\| && \text{by definition of } \|\cdot\| \end{aligned}$$

2a. Proof that $\|\boldsymbol{x}\| = 0 \implies \boldsymbol{x} = 0$:

$$\begin{aligned} 0 &= \|\boldsymbol{x}\| && \text{by left hypothesis} \\ &= \mathsf{d}(\boldsymbol{x},\mathbb{0}) && \text{by definition of } \|\cdot\| \\ &= \mathsf{d}(\boldsymbol{x},\mathbb{0}) && \text{by definition of } \|\cdot\| \\ &\implies \boldsymbol{x} = \mathbb{0} && \text{by property of metrics} \end{aligned}$$

2b. Proof that $\|\boldsymbol{x}\| = 0 \impliedby \boldsymbol{x} = 0$:

$$\begin{aligned} \|\boldsymbol{x}\| &= \mathsf{d}(\boldsymbol{x},\mathbb{0}) && \text{by definition of } \|\cdot\| \\ &= \mathsf{d}(\mathbb{0},\mathbb{0}) && \text{by right hypothesis} \\ &= 0 && \text{by property of metrics} \end{aligned}$$

3. Proof that $\|\boldsymbol{x}+\boldsymbol{y}\| \le \|\boldsymbol{x}\|+\|\boldsymbol{y}\|$:

$$\begin{aligned} \|\boldsymbol{x}+\boldsymbol{y}\| &= \mathsf{d}(\boldsymbol{x}+\boldsymbol{y},\mathbb{0}) && \text{by definition of } \|\cdot\| \\ &= \mathsf{d}(\boldsymbol{x}+\boldsymbol{y}-\boldsymbol{y},\mathbb{0}-\boldsymbol{y}) && \text{by translation invariance hypothesis} \\ &= \mathsf{d}(\boldsymbol{x},-\boldsymbol{y}) \\ &\le \mathsf{d}(\boldsymbol{x},\mathbb{0})+\mathsf{d}(\mathbb{0},\boldsymbol{y}) && \text{by property of metrics} \\ &= \mathsf{d}(\boldsymbol{x},\mathbb{0})+\mathsf{d}(\boldsymbol{y},\mathbb{0}) && \text{by property of metrics} \\ &= \|\boldsymbol{x}\|+\|\boldsymbol{y}\| && \text{by definition of } \|\cdot\| \end{aligned}$$

✎

Theorem K.5. [11] *Let* $\big(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times})\big)$ *be a* LINEAR SPACE. *Let* $\mathsf{d}(\boldsymbol{x},\boldsymbol{y}) \triangleq \|\boldsymbol{x}-\boldsymbol{y}\| \ \forall \boldsymbol{x},\boldsymbol{y}\in X$.

THM

$$\left.\begin{array}{lll} 1. & \mathsf{d}(\boldsymbol{x}+\boldsymbol{z},\boldsymbol{y}+\boldsymbol{z})=\mathsf{d}(\boldsymbol{x},\boldsymbol{y}) \quad \forall \boldsymbol{x},\boldsymbol{y},\boldsymbol{z}\in X & \text{(TRANSLATION INVARIANT)} \quad \textit{and} \\ 2. & \mathsf{d}(\alpha\boldsymbol{x},\alpha\boldsymbol{y})=|\alpha|\mathsf{d}(\boldsymbol{x},\boldsymbol{y}) \quad \forall \boldsymbol{x},\boldsymbol{y}\in X, \alpha\in\mathbb{F} & \text{(HOMOGENEOUS)} \end{array}\right\} \iff \|\cdot\| \textit{ is a } \text{NORM}$$

✎PROOF:

1. Proof of $\implies$ assertion:
   (a) Proof that $\|\cdot\|$ is *strictly positive*: This follows directly from the definition of d.
   (b) Proof that $\|\cdot\|$ is *nondegenerate*: This follows directly from Lemma K.1 (page 276).
   (c) Proof that $\|\cdot\|$ is *homogeneous*: This follows from the second left hypothesis.

[11] Bollobás (1999), page 21

COPYRIGHT © 2019 DANIEL J. GREENHOE

(d) Proof that $\|\cdot\|$ satisfies the *triangle-inequality*: This follows directly from Lemma K.1 (page 276).

2. Proof of $\Longleftarrow$ assertion:

$$\begin{aligned}
\mathsf{d}(\boldsymbol{x}+\boldsymbol{z},\boldsymbol{y}+\boldsymbol{z}) &= \|(\boldsymbol{x}+\boldsymbol{z})-(\boldsymbol{y}+\boldsymbol{z})\| && \text{by definition of } \mathsf{d}\\
&= \|\boldsymbol{x}-\boldsymbol{y}\| \\
&= \mathsf{d}(\boldsymbol{x},\boldsymbol{y}) && \text{by definition of } \mathsf{d}\\
\mathsf{d}(\alpha\boldsymbol{x},\alpha\boldsymbol{y}) &= \|(\alpha\boldsymbol{x})-(\alpha\boldsymbol{y})\| && \text{by definition of } \mathsf{d}\\
&= \|\alpha(\boldsymbol{x}-\boldsymbol{y})\| \\
&= |\alpha|\,\|\boldsymbol{x}-\boldsymbol{y}\| && \text{by definition of } \|\cdot\| \text{ page } 271\\
&= |\alpha|\mathsf{d}(\boldsymbol{x},\boldsymbol{y}) && \text{by definition of } \mathsf{d}
\end{aligned}$$

✍

K.3 Orthogonality on normed linear spaces

Traditionally, *orthogonality* (Definition J.4 page 267) is a property defined in *inner product spaces* (Definition J.1 page 255) However, the concept of orthogonality can be extended to *normed linear spaces* (Definition K.1 page 271). Here are some examples:

① *Isosceles orthogonality*: Definition K.3 page 278
② *Pythagorean orthogonality*: Definition K.4 page 280
③ *Birkhoff orthogonality*: Definition K.5 page 280



Isosceles orthogonality (Definition K.3 page 278) can be illustrated using a *parallelogram*, as illustrated in the figure to the upper left. In this case, orthogonality implies that the parallelogram is a rectangle, which in turn implies that the lengths of the two diagonals are equal ($\|\boldsymbol{x}+\boldsymbol{y}\| = \|\boldsymbol{x}-\boldsymbol{y}\|$). Isosceles orthogonality can also be illustrated with a triangle where the sides are of lengths $\|\boldsymbol{x}+\boldsymbol{y}\|$ and $\|\boldsymbol{x}-\boldsymbol{y}\|$ and base of length $\|\boldsymbol{x}+\boldsymbol{x}\|$. In this case if $\boldsymbol{x}$ and $\boldsymbol{y}$ are orthogonal, then the triangle is *isosceles*. This is illustrated in figure to the lower left. Isosceles orthogonality is formally defined next.

Definition K.3. [12] *Let* $\big(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \|\cdot\|\big)$ *be a* NORMED LINEAR SPACE *(Definition K.1 page 271)*.

DEF

Two vectors $\boldsymbol{x}$ *and* $\boldsymbol{y}$ *are* ***orthogonal in the sense of James*** *if*

$$\|\boldsymbol{x}+\boldsymbol{y}\| = \|\boldsymbol{x}-\boldsymbol{y}\|\,.$$

This property is also called ***isosceles orthogonality*** *or* ***James orthogonality***.

Theorem K.6. *Let* $\big(X, +, \cdot, (\mathbb{R}, \dot{+}, \dot{\times}), \langle\triangle\,|\,\triangledown\rangle\big)$ *be an* INNER-PRODUCT SPACE *(Definition J.1 page 255) with induced norm* $\|\boldsymbol{x}\| \triangleq \sqrt{\langle\boldsymbol{x}\,|\,\boldsymbol{x}\rangle}$, ISOSCELES ORTHOGONALITY *(Definition K.3 page 278) relation* $\textcircled{\perp}$, *and inner-product relation* ORTHOGONALITY *(Definition J.4 page 267) relation* $\perp$.

THM

$$\underbrace{\boldsymbol{x}\,\textcircled{\perp}\,\boldsymbol{y}}_{\text{orthogonal in the sense of James}} \quad\Longleftrightarrow\quad \underbrace{\boldsymbol{x}\perp\boldsymbol{y}}_{\text{orthogonal in the sense of inner-product space}}$$

[12] James (1945) page 292 (DEFINITION 2.1), Amir (1986) page 24, Dunford and Schwartz (1957), page 93

BY-NC-ND
COPYRIGHT © 2019 DANIEL J. GREENHOE

✎PROOF:

1. Proof that $x \circledast y \implies x \perp y$:

$$
\begin{aligned}
&4\langle x \mid y\rangle \\
&= \underbrace{\|x+y\|^2 - \|x-y\|^2}_{0 \text{ by } x \circledast y \text{ hypothesis}} + i\,\|x+iy\|^2 - i\,\|x-iy\|^2 && \text{by } \textit{polarization identity} \text{ (Theorem J.6 page 262)} \\
&= 0 + i\,\|x+iy\|^2 - i\,\|x-iy\|^2 && \text{by } x \circledast y \text{ hypothesis} \\
&= i\left[\|x\|^2 + \|iy\|^2 + 2\Re\langle x \mid iy\rangle\right] \\
&\qquad - i\left[\|x\|^2 + \|-iy\|^2 + 2\Re\langle x \mid -iy\rangle\right] && \text{by } \textit{Polar Identity} \text{ (Lemma J.1 page 259)} \\
&= i\left[\|x\|^2 + \|y\|^2 + 2\Re\langle x \mid iy\rangle\right] \\
&\qquad - i\left[\|x\|^2 + \|y\|^2 + 2\Re\langle x \mid -iy\rangle\right] && \text{by Definition K.1 page 271 and Definition J.1 page 255} \\
&= 4i\Re\langle x \mid iy\rangle \\
&= 4i\Re\left[i^*\langle x \mid y\rangle\right] \\
&= 0 && \text{because inner-product space is } \textit{real} \ (\mathbb{F} = \mathbb{R})
\end{aligned}
$$

2. Proof that $x \circledast y \impliedby x \perp y$:

$$
\begin{aligned}
\|x+y\|^2 &= \|x\|^2 + \|y\|^2 + 2\Re\langle x \mid y\rangle && \text{by } \textit{Polar Identity} \text{ (Lemma J.1 page 259)} \\
&= \|x\|^2 + \|y\|^2 + 0 && \text{by } x \perp y \text{ hypothesis} \\
&= \|x\|^2 + \|y\|^2 - 2\Re\cancelto{0 \text{ when } x \perp y}{\langle x \mid y\rangle} && \text{by } x \perp y \text{ hypothesis} \\
&= \|x\|^2 + \|-y\|^2 + 2\Re\langle x \mid -y\rangle \\
&= \|x-y\|^2 && \text{by } \textit{Polar Identity} \text{ (Lemma J.1 page 259)}
\end{aligned}
$$

✏

Theorem K.7. [13] *Let* $\left(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \|\cdot\|\right)$ *be a normed linear space and with* ISOSCELES ORTHOGONALITY *(Definition K.3 page 278) relation* $\circledast$.

THM $\quad x \circledast y \iff y \circledast x \iff \alpha x \circledast \alpha y \quad \forall \alpha \in \mathbb{F}$

✎PROOF:

$$
\begin{aligned}
x \circledast y &\implies \|x+y\| &&= \|x-y\| && \text{by Definition K.3 page 278} \\
&\implies \|x+y\| &&= |-1|\ \|x-y\| \\
&\implies \|x+y\| &&= \|-(x-y)\| && \text{by Definition K.1 page 271} \\
&\implies \|y+x\| &&= \|y-x\| && \text{by Definition G.1 page 207} \\
&\implies y \circledast x && && \text{by Definition K.3 page 278}
\end{aligned}
$$

$$
\begin{aligned}
y \circledast x &\implies \|y+x\| &&= \|y-x\| && \text{by Definition K.3 page 278} \\
&\implies |\alpha|\ \|y+x\| &&= |\alpha|\ \|y-x\| \\
&\implies \|\alpha(y+x)\| &&= \|\alpha(y-x)\| && \text{by Definition K.1 page 271} \\
&\implies \|\alpha y + \alpha x\| &&= \|\alpha y - \alpha x\| \\
&\implies \|\alpha x + \alpha y\| &&= \|-(\alpha x - \alpha y)\| && \text{by Definition G.1 page 207}
\end{aligned}
$$

[13] Amir (1986) page 24

COPYRIGHT © 2019 DANIEL J. GREENHOE

$$\begin{aligned}
&\Longrightarrow \|\alpha\boldsymbol{x}+\alpha\boldsymbol{y}\| &&= |-1|\ \|\alpha\boldsymbol{x}-\alpha\boldsymbol{y}\| && \text{by Definition K.1 page 271}\\
&\Longrightarrow \|\alpha\boldsymbol{x}+\alpha\boldsymbol{y}\| &&= \|\alpha\boldsymbol{x}-\alpha\boldsymbol{y}\| && \text{by Definition F.4 page 206}\\
&\Longrightarrow \alpha\boldsymbol{x} \textcircled{\perp} \alpha\boldsymbol{y} && && \text{by Definition K.3 page 278}
\end{aligned}$$

$$\begin{aligned}
\alpha\boldsymbol{x} \textcircled{\perp} \alpha\boldsymbol{y} &\Longrightarrow \|\alpha\boldsymbol{x}+\alpha\boldsymbol{y}\| &&= \|\alpha\boldsymbol{x}-\alpha\boldsymbol{y}\| && \text{by Definition K.3 page 278}\\
&\Longrightarrow \|\alpha(\boldsymbol{x}+\boldsymbol{y})\| &&= \|\alpha(\boldsymbol{x}-\boldsymbol{y})\| && \text{by Definition G.1 page 207}\\
&\Longrightarrow |\alpha|\ \|\boldsymbol{x}+\boldsymbol{y}\| &&= |\alpha|\ \|\boldsymbol{x}-\boldsymbol{y}\| && \text{by Definition K.1 page 271}\\
&\Longrightarrow \|\boldsymbol{x}+\boldsymbol{y}\| &&= \|\boldsymbol{x}-\boldsymbol{y}\| && \text{by Definition K.1 page 271}\\
&\Longrightarrow \boldsymbol{x} \textcircled{\perp} \boldsymbol{y} && && \text{by Definition K.3 page 278}
\end{aligned}$$

✏



If a triangle in a plane has two perpendicular sides of lengths a and b and a hypotenuse of length c, then by the *Pythagorean Theorem* (Theorem J.10 page 268), $a^2+b^2=c^2$. This concept of orthogonality can be generalized to normed linear spaces. Two vectors $\boldsymbol{x}$ and $\boldsymbol{y}$ (with lengths $\|\boldsymbol{x}\|$ and $\|\boldsymbol{y}\|$) are orthogonal when $\|\boldsymbol{x}\|^2+\|\boldsymbol{y}\|^2=\|\boldsymbol{x}-\boldsymbol{y}\|^2$ ($\boldsymbol{x}-\boldsymbol{y}$ is a kind of "*hypotenuse*"). This kind of orthogonality is defined next and illustrated in the figure to the left.

Definition K.4. [14] *Let* $\big(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \|\cdot\|\big)$ *be a* NORMED LINEAR SPACE *(Definition K.1 page 271).*

DEF

Two vectors $\boldsymbol{x}$ *and* $\boldsymbol{y}$ *are* ***orthogonal in the Pythagorean sense*** *if*

$$\|\boldsymbol{x}-\boldsymbol{y}\|^2 = \|\boldsymbol{x}\|^2+\|\boldsymbol{y}\|^2.$$

This relationship is also called ***Pythagorean orthogonality****.*

Theorem K.8. [15] *Let* $\big(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle\triangle\,|\,\triangledown\rangle\big)$ *be an* INNER-PRODUCT SPACE *(Definition J.1 page 255) with induced norm* $\|\boldsymbol{x}\| \triangleq \sqrt{\langle\boldsymbol{x}\,|\,\boldsymbol{x}\rangle}$, PYTHAGOREAN ORTHOGONALITY *(Definition K.4 page 280) relation* $\textcircled{\perp}$, *and inner-product relation* ORTHOGONALITY *(Definition J.4 page 267) relation* $\perp$.

THM

$$\underbrace{\boldsymbol{x} \textcircled{\perp} \boldsymbol{y}}_{\text{orthogonal in the Pythagorean sense}} \quad\Longleftrightarrow\quad \underbrace{\boldsymbol{x} \perp \boldsymbol{y}}_{\text{orthogonal in the sense of inner-product space}}$$



Besides *isosceles orthogonality* (Definition K.3 page 278), orthogonality in normed linear spaces can be defined using *Birkhoff orthogonality*, as defined in Definition K.5 (next) and illustrated to the left.

Definition K.5. [16] *Let* $\big(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \|\cdot\|\big)$ *be a* NORMED LINEAR SPACE *(Definition K.1 page 271).*

DEF

Two vectors $\boldsymbol{x}$ *and* $\boldsymbol{y}$ *are* ***orthogonal in the sense of Birkhoff*** *if*

$$\|\boldsymbol{x}\| \le \|\boldsymbol{x}+\alpha\boldsymbol{y}\| \quad \forall\alpha\in\mathbb{F}.$$

This relationship is also called ***Birkhoff orthogonality****.*

Theorem K.9. *Let* $\big(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle\triangle\,|\,\triangledown\rangle\big)$ *be an* INNER-PRODUCT SPACE *(Definition J.1 page 255) with induced norm* $\|\boldsymbol{x}\| \triangleq \sqrt{\langle\boldsymbol{x}\,|\,\boldsymbol{x}\rangle}$, BIRKHOFF ORTHOGONALITY *relation* $\textcircled{\perp}$ *(Definition K.5 page 280), and inner-product relation* ORTHOGONALITY *relation* $\perp$ *(Definition J.4 page 267).*

THM

$$\underbrace{\boldsymbol{x} \textcircled{\perp} \boldsymbol{y}}_{\text{orthogonal in the sense of Birkhoff}} \quad\Longleftrightarrow\quad \underbrace{\boldsymbol{x} \perp \boldsymbol{y}}_{\text{orthogonal in the sense of inner-product space}}$$

[14] James (1945) page 292 ⟨DEFINITION 2.2⟩, Amir (1986) page 57, Drljević (1989) page 232
[15] Amir (1986) page 57
[16] Amir (1986) page 33, Dunford and Schwartz (1957), page 93, James (1947) page 265

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

APPENDIX L

INTERVALS AND CONVEXITY

L.1 Intervals

In the real number system, for $a \leq b$, the *interval* $[a : b]$ is the set a and b and all the numbers inbetween, as in $[a : b] \triangleq \{x \in \mathbb{R} | a \leq x \leq b\}$. This concept can be easily generalized:

- In an **ordered set**, if two elements x and y are *comparable* and $x \leq y$, then we say that x and y and all the elements inbetween, as determined by the ordering relation $\leq$, are the interval $[a : b]$.
- In a **lattice**, the concept of the *interval* can be generalized even further. In an arbitrary ordered set, the interval $[x : y]$ of item (L.1) is restricted to the case in which x and y are *comparable*. This restriction can be lifted (Definition L.2 page 281) with the additional structure of upper and lower bounds provided by lattices.
- A **metric space** in general has no *order relation* $\leq$. But intervals can still be defined (Definition L.4 page 282) in a metric space in terms of the *triangle inequality*.
- A **linear space** (Definition G.1 page 207) over a real or complex field in general has no *order relation* that compares *vectors* in the space, but the standard order relation $\leq$ for real numbers $\mathbb{R}$ can still be used (Definition L.5 page 282) to define an interval in a linear space.

Definition L.1 (intervals on ordered sets). [1] *Let* $(X, \leq)$ *be an* ORDERED SET.

DEF

The set $[x : y]$	$\triangleq$	$\{z \in X \mid x \leq z \leq y\}$	*is called a* ***closed interval***	*and*
The set $(x : y]$	$\triangleq$	$\{z \in X \mid x < z \leq y\}$	*is called a* ***half-open interval***	*and*
The set $[x : y)$	$\triangleq$	$\{z \in X \mid x \leq z < y\}$	*is called a* ***half-open interval***	*and*
The set $(x : y)$	$\triangleq$	$\{z \in X \mid x < z < y\}$	*is called an* ***open interval.***	

Definition L.2 (intervals on lattices). [2] *Let* $(X, \vee, \wedge ; \leq)$ *be a* LATTICE.

DEF

The set $[x : y]$	$\triangleq$	$\{z \in X \mid x \wedge y \leq z \leq x \vee y\}$	*is called a* ***closed interval.***
The set $(x : y]$	$\triangleq$	$\{z \in X \mid x \wedge y < z \leq x \vee y\}$	*is called a* ***half-open interval.***
The set $[x : y)$	$\triangleq$	$\{z \in X \mid x \wedge y \leq z < x \vee y\}$	*is called a* ***half-open interval.***
The set $(x : y)$	$\triangleq$	$\{z \in X \mid x \wedge y < z < x \vee y\}$	*is called an* ***open interval.***

When x and y *are* comparable and $x \leq y$, then Definition L.2 (previous) simplifies to item (L.1)

[1] Apostol (1975) page 4, Ore (1935) page 409
[2] Duthie (1942) page 2, Ore (1935) page 425 ⟨*quotient structures*⟩

(page 281).

Definition L.3. [3] *Let* $\boldsymbol{L} \triangleq (X, \vee, \wedge; \leq)$ *be a* LATTICE *with dual* $\boldsymbol{L}^*$. *Let* $[x : y]$ *be a* CLOSED INTERVAL *(Definition L.2 page 281) on set* X. *The sublattices* $\boldsymbol{L}[x : y]$ *and* $\boldsymbol{L}^*[x : y]$ *are defined as follows:*

DEF

$$\boldsymbol{L}[x : y] \triangleq \{z \in \boldsymbol{L} | z \in [x : y]\} \qquad \forall x,y \in X$$
$$\boldsymbol{L}^*[x : y] \triangleq \left\{z \in \boldsymbol{L}^* | z \in [x : y]\right\} \qquad \forall x,y \in X$$

Definition L.4. [4]

DEF

In a METRIC SPACE (X, d),
the set $[a : b]$ *is the* ***closed interval*** *from* x *to* y *and is defined as*

$$[x : y] \triangleq \{z \in X | \mathsf{d}(x, z) + \mathsf{d}(z, y) = \mathsf{d}(x, y)\}.$$

An element $z \in X$ *is* ***geodesically between*** x *and* y *if* $z \in [x : y]$.

Definition L.5. [5]

DEF

In a LINEAR SPACE $\left(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times})\right)$ *(Definition G.1 page 207),*

$$\begin{aligned}
[\boldsymbol{x} : \boldsymbol{y}] &\triangleq \{\lambda\boldsymbol{x} + (1-\lambda)\boldsymbol{y} = \boldsymbol{z} | 0 \leq \lambda \leq 1\} && \textit{is called a } \textbf{\textit{closed interval}} && \textit{and}\\
(\boldsymbol{x} : \boldsymbol{y}] &\triangleq \{\lambda\boldsymbol{x} + (1-\lambda)\boldsymbol{y} = \boldsymbol{z} | 0 < \lambda \leq 1\} && \textit{is called a } \textbf{\textit{half-open interval}} && \textit{and}\\
[\boldsymbol{x} : \boldsymbol{y}) &\triangleq \{\lambda\boldsymbol{x} + (1-\lambda)\boldsymbol{y} = \boldsymbol{z} | 0 \leq \lambda < 1\} && \textit{is called a } \textbf{\textit{half-open interval}} && \textit{and}\\
(\boldsymbol{x} : \boldsymbol{y}) &\triangleq \{\lambda\boldsymbol{x} + (1-\lambda)\boldsymbol{y} = \boldsymbol{z} | 0 < \lambda < 1\} && \textit{is called an } \textbf{\textit{open interval}}.
\end{aligned}$$

L.2 Convex sets

Using the concept of the *interval* (previous section), we can define the *convex set* (next definition).

Definition L.6. [6] *Let* X *be a* SET *in an* ORDERED SET $(X, \leq)$, *a* LATTICE $(X, \vee, \wedge; \leq)$, *a* METRIC SPACE (X, d), *or a* LINEAR SPACE $\left(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times})\right)$.

DEF

A subset $D \subseteq X$ *is a* ***convex set*** *in* X *if*

$$x, y \in D \qquad \Longrightarrow \qquad [x : y] \subseteq D.$$

A set that is ***not*** *convex is* ***concave****.*

Example L.1. Consider the *Euclidean space* $\mathbb{R}^2$ (a special case of a *linear space*).



$\Leftarrow \left\{\begin{array}{l}\text{The figure to the left is a}\\ \textit{convex} \text{ set in } \mathbb{R}^2.\end{array}\right.$

$\left.\begin{array}{l}\text{The figure to the right is a}\\ \textit{concave} \text{ set in } \mathbb{R}^2.\end{array}\right\} \Rightarrow$



Example L.2. In a *metric space*, examples of *convex sets* are convex *balls*. Examples include those balls generated by the following metrics:

- Taxi-cab metric
- Euclidean metric
- Sup metric
- Tangential metric

[3] Maeda and Maeda (1970), page 1
[4] van de Vel (1993) page 8
[5] Barvinok (2002) page 2
[6] Barvinok (2002) page 5

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

Examples of metrics generating balls which are *not* convex include the following:

- Parabolic metric
- Exponential metric

L.3 Convex functions



Figure L.1: Convex and concave functions

Definition L.7. [7] *Let* $\big(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times})\big)$ *be a* LINEAR SPACE *(Definition G.1 page 207) and* D *a* CONVEX SET *(Definition L.6 page 282) in* X.

DEF

A function $\mathsf{f} \in F^D$ *is* ***convex*** *if*

$$\mathsf{f}\big(\lambda \boldsymbol{x} + [1-\lambda]\boldsymbol{y}\big) \quad \leq \quad \lambda \mathsf{f}(\boldsymbol{x}) + (1-\lambda)\,\mathsf{f}(\boldsymbol{y}) \qquad \forall \boldsymbol{x}, \boldsymbol{y} \in D \text{ and } \forall \lambda \in (0,1)$$

A function $\mathsf{g} \in F^D$ *is* ***strictly convex*** *if*

$$\mathsf{g}\big(\lambda \boldsymbol{x} + [1-\lambda]\boldsymbol{y}\big) \quad = \quad \lambda \mathsf{g}(\boldsymbol{x}) + (1-\lambda)\,\mathsf{g}(\boldsymbol{y}) \qquad \forall \boldsymbol{x}, \boldsymbol{y} \in D,\ \boldsymbol{x} \neq \boldsymbol{y}, \text{ and } \forall \lambda \in (0,1)$$

A function $\mathsf{f} \in F^D$ *is* ***concave*** *if* $-\mathsf{f}$ *is* CONVEX.

A function $\mathsf{f} \in F^D$ *is* ***affine*** *if* f *is* CONVEX *and* CONCAVE.

Example L.3. The function $\mathsf{f}(x) = 2^x$ is a *convex function* (Definition L.7 page 283), as illustrated to the right.



Definition L.8. [8] *Let* $\big(X, +, \cdot, (\mathbb{R}, \dot{+}, \dot{\times})\big)$ *be a* LINEAR SPACE *(Definition G.1 page 207)*.

DEF

The ***epigraph*** $\mathsf{epi}(\mathsf{f})$ *and* ***hypograph*** $\mathsf{hyp}(\mathsf{f})$ *of a functional* $\mathsf{f} \in \mathbb{R}^X$ *are defined as*

$$\mathsf{epi}(\mathsf{f}) \triangleq \{(x,y) \in X \times \mathbb{R} \,|\, y \geq \mathsf{f}(x)\}$$
$$\mathsf{hyp}(\mathsf{f}) \triangleq \{(x,y) \in X \times \mathbb{R} \,|\, y \leq \mathsf{f}(x)\}$$

Example L.4.

[7] Simon (2011) page 2, Barvinok (2002) page 2, Bollobás (1999), page 3, Jensen (1906), page 176, Clarkson (1936) ⟨strictly convex⟩

[8] Beer (1993) page 13 ⟨§1.3⟩, Aubin and Frankowska (2009) page 222, Aubin (2011) page 223

Copyright © 2019 Daniel J. Greenhoe

EX



Proposition L.1. [9] *Let* $\big(X, +, \cdot, (\mathbb{R}, \dot{+}, \dot{\times})\big)$ *be a* LINEAR SPACE *(Definition G.1 page 207). Let* f *be a* FUNCTIONAL *in* $\mathbb{R}^X$.

PRP

$$\left\{\begin{array}{l}\mathsf{f} \text{ is a}\\ \text{CONVEX FUNCTION}\end{array}\right\} \iff \left\{\begin{array}{l}\text{epi}(\mathsf{f}) \text{ is a}\\ \text{CONVEX SET}\end{array}\right\}$$

Often a function can be proven to be *convex* or *concave*. *Convex* and *concave* functions are defined in Definition L.9 (page 284) (next) and illustrated in Figure L.1 (page 283).

Definition L.9. *Let*

$$\mathsf{y}(x) = \frac{\mathsf{f}(b) - \mathsf{f}(a)}{b-a}(x-a) + \mathsf{f}(a).$$

A function $\mathsf{f} : \mathbb{R} \to \mathbb{R}$ *is*

DEF

(1).	***convex***	*in* $(a : b)$ *if*	$\mathsf{f}(x)$	$\leq$	$\mathsf{y}(x)$	*for* $x \in (a : b)$
(2).	***concave***	*in* $(a : b)$ *if*	$\mathsf{f}(x)$	$\geq$	$\mathsf{y}(x)$	*for* $x \in (a : b)$
(3).	***strictly convex***	*in* $(a : b)$ *if*	$\mathsf{f}(x)$	$<$	$\mathsf{y}(x)$	*for* $x \in (a : b)$
(4).	***strictly concave***	*in* $(a : b)$ *if*	$\mathsf{f}(x)$	$>$	$\mathsf{y}(x)$	*for* $x \in (a : b)$

Theorem L.1 (Jensen's Inequality). [10] *Let* $\big(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times})\big)$ *be a* LINEAR SPACE *(Definition G.1 page 207), D a subset of X, and* f *a functional in* $\mathbb{F}^D$. *Let* $\sum$ *be the* SUMMATION OPERATOR *(Definition M.1 page 289).*

THM

$$\left\{\begin{array}{lll} 1. & D \text{ is CONVEX} & \text{and}\\ 2. & \mathsf{f} \text{ is CONVEX} & \text{and}\\ 3. & \displaystyle\sum_{n=1}^{N} \lambda_n = 1 & \text{(WEIGHTS)}\end{array}\right\} \implies \mathsf{f}\left(\sum_{n=1}^{N} \lambda_n \boldsymbol{x}_n\right) \leq \sum_{n=1}^{N} \lambda_n \mathsf{f}(\boldsymbol{x}_n) \qquad \forall \boldsymbol{x}_n \in D,\ N \in \mathbb{N}$$

✎PROOF: Proof is by induction:

1. Proof that statement is true for $N = 1$:

$$\begin{aligned}\mathsf{f}\left(\sum_{n=1}^{N=1} \lambda_n \boldsymbol{x}_n\right) &= \mathsf{f}(\lambda_1 \boldsymbol{x}_1)\\ &\leq \mathsf{f}(\lambda_1 \boldsymbol{x}_1)\\ &= \sum_{n=1}^{N=1} \lambda_n \mathsf{f}(\boldsymbol{x}_n)\end{aligned}$$

[9] Udriste (1994) page 63, Kurdila and Zabarankin (2005) page 178 ⟨Proposition 6.1.1⟩, Rockafellar (1970) page 23 ⟨Section 4 Convex Functions⟩, Çinlar and Vanderbei (2013) page 86 ⟨5.4 Theorem⟩

[10] Mitrinović et al. (2010) page 6, Bollobás (1999) page 3, Lay (1982) page 7, Jensen (1906), pages 179–180

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

2. Proof that statement is true for $N = 2$:

$$\begin{aligned}
\mathsf{f}\left(\sum_{n=1}^{N=2} \lambda_n \boldsymbol{x}_n\right) &= \mathsf{f}(\lambda_1 \boldsymbol{x}_1 + \lambda_2 \boldsymbol{x}_2) \\
&\le \lambda_1 \mathsf{f}(\boldsymbol{x}_1) + \lambda_2 \mathsf{f}(\boldsymbol{x}_2) && \text{by convexity hypothesis} \\
&= \sum_{n=1}^{N=2} \lambda_n \mathsf{f}(\boldsymbol{x}_n)
\end{aligned}$$

3. Proof that if the statement is true for N, then it is also true for $N+1$:

$$\begin{aligned}
\mathsf{f}\left(\sum_{n=1}^{N+1} \lambda_n \boldsymbol{x}_n\right) &= \mathsf{f}\left(\sum_{n=1}^{N} \lambda_n \boldsymbol{x}_n + \lambda_{N+1}\boldsymbol{x}_{N+1}\right) \\
&= \mathsf{f}\left([1-\lambda_{N+1}]\sum_{n=1}^{N} \frac{\lambda_n}{1-\lambda_{N+1}} \boldsymbol{x}_n + \lambda_{N+1}\boldsymbol{x}_{N+1}\right) \\
&\le [1-\lambda_{N+1}]\mathsf{f}\left(\sum_{n=1}^{N} \frac{\lambda_n}{1-\lambda_{N+1}} \boldsymbol{x}_n\right) + \lambda_{N+1}\mathsf{f}(\boldsymbol{x}_{N+1}) && \text{by convexity hypothesis} \\
&\le [1-\lambda_{N+1}]\sum_{n=1}^{N} \frac{\lambda_n}{1-\lambda_{N+1}} \mathsf{f}(\boldsymbol{x}_n) + \lambda_{N+1}\mathsf{f}(\boldsymbol{x}_{N+1}) && \text{by “true for } N\text{” hypothesis} \\
&= \sum_{n=1}^{N} \lambda_n \mathsf{f}(\boldsymbol{x}_n) + \lambda_{N+1}\mathsf{f}(\boldsymbol{x}_{N+1}) \\
&= \sum_{n=1}^{N+1} \lambda_n \mathsf{f}(\boldsymbol{x}_n)
\end{aligned}$$

4. Since the statement is true for $N = 1$, $N = 2$, and true for $N \implies$ true for $N+1$, then it is true for $N = 1, 2, 3, 4, \ldots$

✏

The next theorem gives another form of convex functions that is a little less intuitive but provides powerful analytic results.

Theorem L.2. *Let* $\mathsf{f} : \mathbb{R} \to \mathbb{R}$. *For every* $x_1, x_2 \in (a,b)$ *and* $\lambda \in [0,1]$

THM

$$\mathsf{f} \text{ is convex in } (a,b) \iff \mathsf{f}(\lambda x_1 + (1-\lambda)x_2) \le \lambda \mathsf{f}(x_1) + (1-\lambda)\mathsf{f}(x_2)$$

✎PROOF:

1. prove f is convex $\implies \mathsf{f}(\lambda x_1 + (1-\lambda)x_2) \le \lambda \mathsf{f}(x_1) + (1-\lambda)\mathsf{f}(x_2)$:

$$\begin{aligned}
\mathsf{f}(\lambda x_1 + (1-\lambda)x_2) &\le \frac{\mathsf{f}(b)-\mathsf{f}(a)}{b-a}[\lambda x_1 + (1-\lambda)x_2 - a] + \mathsf{f}(a) \\
&= \frac{\mathsf{f}(x_2)-\mathsf{f}(x_1)}{x_2-x_1}[\lambda x_1 + (1-\lambda)x_2 - x_1] + \mathsf{f}(x_1) \\
&= \frac{\mathsf{f}(x_2)-\mathsf{f}(x_1)}{x_2-x_1}[(x_2-x_1)(1-\lambda)] + \mathsf{f}(x_1) \\
&= (1-\lambda)\mathsf{f}(x_2) - (1-\lambda)\mathsf{f}(x_1) + \mathsf{f}(x_1) \\
&= \lambda \mathsf{f}(x_1) + (1-\lambda)\mathsf{f}(x_2)
\end{aligned}$$

COPYRIGHT © 2019 DANIEL J. GREENHOE
CC BY-NC-ND

2. prove f is convex $\Longleftarrow$ $f(\lambda x_1 + (1-\lambda)x_2) \le \lambda \mathsf{f}(x_1) + (1-\lambda)\mathsf{f}(x_2)$:

   Let $x = \lambda(b-a)+a$ Notice that as λ varies from 0 to 1, x varies from b to a. So free variable λ works as a change of variable for free variable x.

$$\lambda = \frac{x-a}{b-a}$$

$$\begin{aligned}
\mathsf{f}(x) &= \mathsf{f}(\lambda(b-a)+a) \\
&\le \lambda \mathsf{f}(b) + (1-\lambda)\mathsf{f}(a) \\
&= \lambda[\mathsf{f}(b) - \mathsf{f}(a)] + \mathsf{f}(a) \\
&= \frac{\mathsf{f}(b)-\mathsf{f}(a)}{b-a}(x-a) + \mathsf{f}(a)
\end{aligned}$$

✐

Taking the second derivative of a function provides a convenient test for whether that function is convex.

Theorem L.3. [11]

THM $\mathsf{f}''(x) > 0 \implies \mathsf{f}$ *is convex*

✎ PROOF:

$$\begin{aligned}
\mathsf{f}(x) &= \mathsf{f}(x_0) + \mathsf{f}'(x_0)(x-x_0) + \mathsf{f}''(c)(x-x_0)^2 \\
&\ge \mathsf{f}(x_0) + \mathsf{f}'(x_0)(x-x_0) \\
&= \mathsf{f}(x_0) + \mathsf{f}'(x_0)(x - \lambda x_1 - (1-\lambda)x_2)
\end{aligned}$$

$$\begin{aligned}
\mathsf{f}(x_1) &\ge \mathsf{f}(x_0) + \mathsf{f}'(x_0)(x_1 - \lambda x_1 - (1-\lambda)x_2) \\
&= \mathsf{f}(x_0) + \mathsf{f}'(x_0)(1-\lambda)(x_1 - x_2) \\
&= \mathsf{f}(x_0) - \mathsf{f}'(x_0)(1-\lambda)(x_2 - x_1)
\end{aligned}$$

$$\begin{aligned}
\mathsf{f}(x_2) &\ge \mathsf{f}(x_0) + \mathsf{f}'(x_0)(x_2 - \lambda x_1 - (1-\lambda)x_2) \\
&= \mathsf{f}(x_0) + \mathsf{f}'(x_0)\lambda(x_2 - x_1)
\end{aligned}$$

$$\begin{aligned}
\lambda \mathsf{f}(x_1) + (1-\lambda)\mathsf{f}(x_2) &\ge \lambda\left[\mathsf{f}(x_0) - \mathsf{f}'(x_0)(1-\lambda)(x_2-x_1)\right] + (1-\lambda)\left[\mathsf{f}(x_0) + \mathsf{f}'(x_0)\lambda(x_2-x_1)\right] \\
&= \lambda\left[\mathsf{f}(x_0) - \mathsf{f}'(x_0)(1-\lambda)(x_2-x_1)\right] + \left[\mathsf{f}(x_0) + \mathsf{f}'(x_0)\lambda(x_2-x_1)\right] - \lambda\left[\mathsf{f}(x_0) + \mathsf{f}'(x_0)\lambda(x_2-x_1)\right] \\
&= \mathsf{f}(x_0) \\
&= \mathsf{f}(\lambda x_1 + (1-\lambda)x_2)
\end{aligned}$$

By Theorem L.2 (page 285), $\mathsf{f}(x)$ is convex. ✐

L.4 Literature

Literature survey:

[11] Cover and Thomas (1991), pages 24–25

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

1. Abstract convexity:
   - Edelman and Jamison (1985)
   - van de Vel (1993)
   - Hörmander (1994)
2. Order convexity (lattice theory):
   - Edelman (1986)
3. Metric convexity:
   - Menger (1928)
   - Blumenthal (1970) page 41 ⟨?⟩
   - Khamsi and Kirk (2001) pages 35–38

Copyright © 2019 Daniel J. Greenhoe
BY-NC-ND

Copyright © 2019 Daniel J. Greenhoe

APPENDIX M

FINITE SUMS

❝ *I think that it was Harald Bohr who remarked to me that "all analysts spend half their time hunting through the literature for inequalities which they want to use and cannot prove."* ❞

G.H. Hardy (1877–1947) in his "Presidential Address" to the London Mathematical Society on November 8, 1928, about a remark that he suggested was from Harald Bohr (1887–1951), Danish mathematician pictured to the left. [1]

M.1 Summation

Definition M.1. [2] *Let $+$ be an addition operator on a tuple $(\!| x_n |\!)_m^N$.*

DEF

*The **summation** of $(\!| x_n |\!)$ from index m to index N with respect to $+$ is*

$$\sum_{n=m}^{N} x_n \triangleq \begin{cases} 0 & \text{for } N < m \\ \left(\sum_{n=m}^{N-1} x_n\right) + x_N & \text{for } N \geq m \end{cases}$$

Theorem M.1 (Generalized associative property). [3] *Let $+$ be an addition operator on a tuple $(\!| x_n |\!)_m^N$.*

THM

$+$ *is* ASSOCIATIVE $\Longrightarrow$

$$\underbrace{\sum_{n=m}^{L} x_n + \left(\sum_{n=L+1}^{M} x_n + \sum_{n=M+1}^{N} x_n\right) = \left(\sum_{n=m}^{L} x_n + \sum_{n=L+1}^{M} x_n\right) + \sum_{n=M+1}^{N} x_n}_{\sum_{n=m}^{N} \text{ is ASSOCIATIVE}} \qquad \text{for } m < L < M \leq N$$

✎PROOF:

[1] quote: Hardy (1929), page 64
image: http://www-history.mcs.st-andrews.ac.uk/PictDisplay/Bohr_Harald.html

[2] reference: Berberian (1961) page 8 ⟨Definition I.3.1⟩
"$\sum$" notation: Fourier (1820) page 280

[3] Berberian (1961) pages 9–10 ⟨Theorem I.3.1⟩

1. Proof for $N < m$ case: $\sum_{n=m}^{N} x_n = 0$.

2. Proof for $N = m$ case: $\sum_{n=m}^{m} x_n = \left(\sum_{n=m}^{m-1} x_n\right) + x_m = 0 + x_m = x_m$.

3. Proof for $N = m+1$ case: $\sum_{n=m}^{m+1} x_n = \left(\sum_{n=m}^{m} x_n\right) + x_{m+1} = x_m + x_{m+1}$

4. Proof for $N = m+2$ case:

$$\begin{aligned}
\sum_{n=m}^{m+2} x_n &= \left(\sum_{n=m}^{m+1} x_n\right) + x_{m+2} && \text{by Definition M.1 page 289}\\
&= \left(x_m + x_{m+1}\right) + x_{m+2} && \text{by item (3)}\\
&= x_m + \left(x_{m+1} + x_{m+2}\right) && \text{by left hypothesis}
\end{aligned}$$

5. Proof that N case $\implies$ $N+1$ case:

$$\sum_{n=m}^{N+1} x_n = \underbrace{\left(\sum_{n=m}^{N} x_n\right)}_{associative} + x_{N+1} \qquad \text{by Definition M.1 page 289}$$

$$\begin{aligned}
&= \left(\sum_{n=m}^{L} x_n + \left(\sum_{n=L+1}^{M} x_n + \sum_{n=M+1}^{N} x_n\right)\right) + x_{N+1} &&= \left(\left(\sum_{n=m}^{L} x_n + \sum_{n=L+1}^{M} x_n\right) + \sum_{n=M+1}^{N} x_n\right) + x_{N+1}\\
&= \left(\sum_{n=m}^{L} x_n + \sum_{n=L+1}^{M} x_n\right) + \left(\sum_{n=M+1}^{N} x_n + x_{N+1}\right) &&= \left(\sum_{n=m}^{L} x_n + \sum_{n=L+1}^{M} x_n\right) + \left(\sum_{n=M+1}^{N+1} x_n\right)
\end{aligned}$$

✍

M.2 Means

M.2.1 Weighted ϕ-means

Definition M.2. [4]

DEF

The $(\!|\lambda_n|\!)_1^N$ *weighted* ϕ-***mean*** *of a tuple* $(\!|x_n|\!)_1^N$ *is defined as*

$$\mathsf{M}_\phi\left((\!|x_n|\!)\right) \triangleq \phi^{-1}\left(\sum_{n=1}^{N} \lambda_n \phi(x_n)\right)$$

where ϕ *is a* CONTINUOUS *and* STRICTLY MONOTONIC *function in* $\mathbb{R}^{\mathbb{R}^+}$

and $(\!|\lambda_n|\!)_{n=1}^N$ *is a sequence of weights for which* $\sum_{n=1}^{N} \lambda_n = 1$.

[4] Bollobás (1999) page 5

BY-NC-ND Copyright © 2019 Daniel J. Greenhoe

Lemma M.1. [5] *Let* $\mathsf{M}_\phi(\langle x_n \rangle)$ *be the* $\langle \lambda_n \rangle_1^N$ *weighted* ϕ*-mean of a tuple* $\langle x_n \rangle_1^N$*. Let the property* CONVEX *be defined as in Definition L.7 (page 283).*

LEM					
	$\phi\psi^{-1}$ *is* CONVEX	*and*	ϕ *is* INCREASING	$\Longrightarrow$	$\mathsf{M}_\phi(\langle x_n \rangle) \geq \mathsf{M}_\psi(\langle x_n \rangle)$
	$\phi\psi^{-1}$ *is* CONVEX	*and*	ϕ *is* DECREASING	$\Longrightarrow$	$\mathsf{M}_\phi(\langle x_n \rangle) \leq \mathsf{M}_\psi(\langle x_n \rangle)$
	$\phi\psi^{-1}$ *is* CONCAVE	*and*	ϕ *is* INCREASING	$\Longrightarrow$	$\mathsf{M}_\phi(\langle x_n \rangle) \leq \mathsf{M}_\psi(\langle x_n \rangle)$
	$\phi\psi^{-1}$ *is* CONCAVE	*and*	ϕ *is* DECREASING	$\Longrightarrow$	$\mathsf{M}_\phi(\langle x_n \rangle) \geq \mathsf{M}_\psi(\langle x_n \rangle)$

✎PROOF:

1. Case where $\phi\psi^{-1}$ is *convex* and ϕ is *increasing*:

$$\begin{aligned}
\mathsf{M}_\phi(\langle x_n \rangle) &\triangleq \phi^{-1}\left(\sum_{n=1}^N \lambda_n \phi(x_n)\right) && \text{by definition of } \mathsf{M}_\phi && \text{(Definition M.2 page 290)}\\
&= \phi^{-1}\left(\sum_{n=1}^N \lambda_n \phi\psi^{-1}\psi(x_n)\right) && \text{by definition of } \psi^{-1} && \\
&\geq \phi^{-1}\left(\phi\psi^{-1}\sum_{n=1}^N \lambda_n \psi(x_n)\right) && \text{by } \textit{Jensen's Inequality} && \text{(Theorem L.1 page 284)}\\
&= \left(\psi^{-1}\sum_{n=1}^N \lambda_n \psi(x_n)\right) && \text{by definition of } \psi^{-1} && \\
&\triangleq \mathsf{M}_\psi(\langle x_n \rangle) && \text{by definition of } \mathsf{M}_\psi && \text{(Definition M.2 page 290)}
\end{aligned}$$

2. Case where $\phi\psi^{-1}$ is *convex* and ϕ is *decreasing*:

$$\begin{aligned}
\mathsf{M}_\phi(\langle x_n \rangle) &\triangleq \phi^{-1}\left(\sum_{n=1}^N \lambda_n \phi(x_n)\right) && \text{by definition of } \mathsf{M}_\phi && \text{(Definition M.2 page 290)}\\
&= \phi^{-1}\left(\sum_{n=1}^N \lambda_n \phi\psi^{-1}\psi(x_n)\right) && \text{by definition of } \psi^{-1} && \\
&\leq \phi^{-1}\left(\phi\psi^{-1}\sum_{n=1}^N \lambda_n \psi(x_n)\right) && \text{by } \textit{Jensen's Inequality} \text{ and because } \phi^{-1} \text{ is } \textit{decreasing} && \text{(Theorem L.1 page 284) (by hypothesis)}\\
&= \left(\psi^{-1}\sum_{n=1}^N \lambda_n \psi(x_n)\right) && \text{by definition of } \psi^{-1} && \\
&\triangleq \mathsf{M}_\psi(\langle x_n \rangle) && \text{by definition of } \mathsf{M}_\psi && \text{(Definition M.2 page 290)}
\end{aligned}$$

✏

One of the most well known inequalities in mathematics is *Minkowski's Inequality* (1910, Theorem M.5 page 297). In 1946, H.P. Mulholland submitted a result[6] that generalizes Minkowski's Inequality to an equal weighted ϕ-mean. And Milovanović and Milovanovć (1979) generalized this even further to a *weighted* ϕ-mean (Theorem M.2, next).

Theorem M.2. [7]

[5] Pečarić et al. (1992) page 107, Bollobás (1999) page 5, Hardy et al. (1952) page 75
[6] Mulholland (1950)
[7] Milovanović and Milovanović (1979), Bullen (2003) page 306 ⟨Theorem 9⟩

COPYRIGHT © 2019 DANIEL J. GREENHOE | CC BY-NC-ND

THM

$$\left\{\begin{array}{lll|lll} (1). & \phi \text{ is CONVEX} & \text{and} & (3). & \phi(0)=0 & \text{and} \\ (2). & \phi \text{ is STRICTLY MONOTONIC} & \text{and} & (4). & \log\circ\phi\circ\exp \text{ is CONVEX} & \end{array}\right\} \implies$$

$$\left\{\phi^{-1}\left(\sum_{n=1}^{N}\lambda_n\phi(x_n+y_n)\right) \le \phi^{-1}\left(\sum_{n=1}^{N}\lambda_n\phi(x_n)\right)+\phi^{-1}\left(\sum_{n=1}^{N}\lambda_n\phi(y_n)\right)\right\}$$

M.2.2 Power means

Definition M.3. [8] *Let* $\mathsf{M}_{\phi(x;r)}(\llparenthesis x_n \rrparenthesis)$ *be the* $\llparenthesis \lambda_n \rrparenthesis_1^N$ *weighted* ϕ*-mean of a* NON-NEGATIVE *tuple* $\llparenthesis x_n \rrparenthesis_1^N$ *(Definition M.2 page 290).*

DEF

A mean $\mathsf{M}_{\phi(x;r)}(\llparenthesis x_n \rrparenthesis)$ *is a* ***power mean*** *with parameter* r *if* $\phi(x) \triangleq x^r$*. That is,*

$$\mathsf{M}_{\phi(x;r)}(\llparenthesis x_n \rrparenthesis) = \left(\sum_{n=1}^{N}\lambda_n(x_n)^r\right)^{\frac{1}{r}}$$

Theorem M.3. [9] *Let* $\mathsf{M}_{\phi(x;r)}(\llparenthesis x_n \rrparenthesis)$ *be* POWER MEAN *with parameter* r *of an* N*-tuple* $\llparenthesis x_n \rrparenthesis_1^N$*. Let* $\mathbb{R}^*$ *be the set of extended real numbers* $\left(\mathbb{R}^* \triangleq \mathbb{R}\cup\{-\infty,\infty\}\right)$.[10]

THM

$$\mathsf{M}_{\phi(x;r)}(\llparenthesis x_n \rrparenthesis) \triangleq \left(\sum_{n=1}^{N}\lambda_n(x_n)^r\right)^{\frac{1}{r}} \text{ is CONTINUOUS and STRICTLY INCREASING in } \mathbb{R}^*.$$

$$\mathsf{M}_{\phi(x;r)}(\llparenthesis x_n \rrparenthesis) = \begin{cases} \min\limits_{n=1,2,\ldots,N}\llparenthesis x_n \rrparenthesis & \text{for } r=-\infty \\ \prod\limits_{n=1}^{N} x_n^{\lambda_n} & \text{for } r=0 \\ \max\limits_{n=1,2,\ldots,N}\llparenthesis x_n \rrparenthesis & \text{for } r=+\infty \end{cases}$$

✎PROOF:

1. Proof that $M_{\phi(x;r)}$ is *strictly increasing* in r:
   (a) Let r and s be such that $-\infty < r < s < \infty$.
   (b) Let $\phi_r \triangleq x^r$ and $\phi_s \triangleq x^s$. Then $\phi_r\phi_s^{-1} = x^{\frac{r}{s}}$.
   (c) The composite function $\phi_r\phi_s^{-1}$ is *convex* or *concave* depending on the values of r and s:

	$r<0$ (ϕ_r decreasing)	$r>0$ (ϕ_r increasing)
$s<0$	*convex*	(not possible)
$s>0$	*convex*	*concave*

   (d) Therefore by Lemma M.1 (page 291),
   $$-\infty < r < s < \infty \implies \mathsf{M}_{\phi(x;r)}(\llparenthesis x_n \rrparenthesis) < \mathsf{M}_{\phi(x;s)}(\llparenthesis x_n \rrparenthesis).$$
2. Proof that $\mathsf{M}_{\phi(x;r)}$ is continuous in r for $r\in\mathbb{R}\backslash 0$: The sum of continuous functions is continuous. For the cases of $r\in\{-\infty, 0, \infty\}$, see the items that follow.

[8] Bullen (2003) page 175, Bollobás (1999) page 6
[9] Bullen (2003) pages 175–177 ⟨see also page 203⟩, Bollobás (1999) pages 6–8, Besso (1879), Bienaymé (1840) page 68
[10] Rana (2002) pages 385–388 ⟨Appendix A⟩

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

3. Lemma: $\mathsf{M}_{\phi(x;-r)}\big(\llparenthesis x_n \rrparenthesis\big) = \big\{\mathsf{M}_{\phi(x;r)}\big(\llparenthesis x_n^{-1} \rrparenthesis\big)\big\}^{-1}$. Proof:

$$\begin{aligned}
\big\{\mathsf{M}_{\phi(x;r)}\big(\llparenthesis x_n^{-1} \rrparenthesis\big)\big\}^{-1} &= \left\{\left(\sum_{n=1}^{N} \lambda_n \big(x_n^{-1}\big)^r\right)^{\frac{1}{r}}\right\}^{-1} && \text{by definition of } \mathsf{M}_\phi \\
&= \left(\sum_{n=1}^{N} \lambda_n \big(x_n\big)^{-r}\right)^{\frac{1}{-r}} \\
&= \mathsf{M}_{\phi(x;-r)}\big(\llparenthesis x_n \rrparenthesis\big) && \text{by definition of } \mathsf{M}_\phi
\end{aligned}$$

4. Proof that $\lim_{r\to\infty} \mathsf{M}_\phi\big(\llparenthesis x_n \rrparenthesis\big) = \max_{n\in\mathbb{Z}} \llparenthesis x_n \rrparenthesis$:

   (a) Let $x_m \triangleq \max_{n\in\mathbb{Z}} \llparenthesis x_n \rrparenthesis$

   (b) Note that $\lim_{r\to\infty} \mathsf{M}_\phi \le \max_{n\in\mathbb{Z}} \llparenthesis x_n \rrparenthesis$ because

$$\begin{aligned}
\lim_{r\to\infty} \mathsf{M}_\phi\big(\llparenthesis x_n \rrparenthesis\big) &= \lim_{r\to\infty} \left(\sum_{n=1}^{N} \lambda_n x_n^r\right)^{\frac{1}{r}} && \text{by definition of } \mathsf{M}_\phi \\
&\le \lim_{r\to\infty} \left(\sum_{n=1}^{N} \lambda_n x_m^r\right)^{\frac{1}{r}} && \text{by definition of } x_m \text{ in item (4a) and because } \phi(x) \triangleq x^r \text{ and } \phi^{-1} \text{ are both increasing or both decreasing} \\
&= \lim_{r\to\infty} \left(x_m^r \underbrace{\sum_{n=1}^{N} \lambda_n}_{1}\right)^{\frac{1}{r}} && \text{because } x_m \text{ is a constant} \\
&= \lim_{r\to\infty} \big(x_m^r \cdot 1\big)^{\frac{1}{r}} \\
&= x_m \\
&= \max_{n\in\mathbb{Z}} \llparenthesis x_n \rrparenthesis && \text{by definition of } x_m \text{ in item (4a)}
\end{aligned}$$

   (c) But also note that $\lim_{r\to\infty} \mathsf{M}_\phi \ge \max_{n\in\mathbb{Z}} \llparenthesis x_n \rrparenthesis$ because

$$\begin{aligned}
\lim_{r\to\infty} \mathsf{M}_\phi\big(\llparenthesis x_n \rrparenthesis\big) &= \lim_{r\to\infty} \left(\sum_{n=1}^{N} \lambda_n x_n^r\right)^{\frac{1}{r}} && \text{by definition of } \mathsf{M}_\phi \\
&\ge \lim_{r\to\infty} \big(w_m x_m^r\big)^{\frac{1}{r}} && \text{by definition of } x_m \text{ in item (4a) and because } \phi(x) \triangleq x^r \text{ and } \phi^{-1} \text{ are both increasing or both decreasing} \\
&= \lim_{r\to\infty} w_m^{\frac{1}{r}} x_m^{\frac{r}{r}} \\
&= x_m \\
&= \max_{n\in\mathbb{Z}} \llparenthesis x_n \rrparenthesis && \text{by definition of } x_m \text{ in item (4a)}
\end{aligned}$$

   (d) Combining items (b) and (c) we have $\lim_{r\to\infty} \mathsf{M}_\phi = \max_{n\in\mathbb{Z}} \llparenthesis x_n \rrparenthesis$.

Copyright © 2019 Daniel J. Greenhoe
CC BY-NC-ND

5. Proof that $\lim_{r\to-\infty} \mathsf{M}_\phi\left(\llparenthesis x_n \rrparenthesis\right) = \min_{n\in\mathbb{Z}} \llparenthesis x_n \rrparenthesis$:

$$\begin{aligned}
\lim_{r\to-\infty} \mathsf{M}_{\phi(x;r)}\left(\llparenthesis x_n \rrparenthesis\right) &= \lim_{r\to\infty} \mathsf{M}_{\phi(x;-r)}\left(\llparenthesis x_n \rrparenthesis\right) && \text{by change of variable } r\\
&= \lim_{r\to\infty} \left\{\mathsf{M}_{\phi(x;r)}\left(\llparenthesis x_n^{-1} \rrparenthesis\right)\right\}^{-1} && \text{by Lemma in item (3) page 293}\\
&= \lim_{r\to\infty} \frac{1}{\mathsf{M}_{\phi(x;r)}\left(\llparenthesis x_n^{-1} \rrparenthesis\right)}\\
&= \frac{\lim_{r\to\infty} 1}{\lim_{r\to\infty} \mathsf{M}_{\phi(x;r)}\left(\llparenthesis x_n^{-1} \rrparenthesis\right)} && \text{by property of lim [11]}\\
&= \frac{1}{\max_{n\in\mathbb{Z}} \llparenthesis x_n^{-1} \rrparenthesis} && \text{by item (4)}\\
&= \frac{1}{\left(\min_{n\in\mathbb{Z}} \llparenthesis x_n \rrparenthesis\right)^{-1}}\\
&= \min_{n\in\mathbb{Z}} \llparenthesis x_n \rrparenthesis
\end{aligned}$$

6. Proof that $\lim_{r\to 0} \mathsf{M}_\phi\left(\llparenthesis x_n \rrparenthesis\right) = \prod_{n=1}^{N} x_n^{\lambda_n}$:

$$\begin{aligned}
\lim_{r\to 0} \mathsf{M}_\phi\left(\llparenthesis x_n \rrparenthesis\right) &= \lim_{r\to 0} \exp\left\{\ln\left\{\mathsf{M}_\phi\left(\llparenthesis x_n \rrparenthesis\right)\right\}\right\}\\
&= \lim_{r\to 0} \exp\left\{\ln\left\{\left(\sum_{n=1}^{N} \lambda_n \left(x_n^r\right)\right)^{\frac{1}{r}}\right\}\right\} && \text{by definition of } \mathsf{M}_\phi\\
&= \exp\left\{\frac{\frac{\partial}{\partial r}\ln\left(\sum_{n=1}^{N} \lambda_n \left(x_n^r\right)\right)}{\frac{\partial}{\partial r} r}\right\}_{r=0} && \text{by l'Hôpital's rule[12]}\\
&= \exp\left\{\frac{\sum_{n=1}^{N} \lambda_n \frac{\partial}{\partial r}\left(x_n^r\right)}{\sum_{n=1}^{N} \lambda_n \left(x_n^r\right)}\right\}_{r=0} && = \exp\left\{\frac{\sum_{n=1}^{N} \lambda_n \frac{\partial}{\partial r}\exp\left(\ln\left(x_n^r\right)\right)}{\sum_{n=1}^{N} \lambda_n}\right\}_{r=0}\\
&= \exp\left\{\frac{\sum_{n=1}^{N} \lambda_n \frac{\partial}{\partial r}\exp\left(r\ln\left(x_n\right)\right)}{1}\right\}_{r=0} && = \exp\left\{\sum_{n=1}^{N} \lambda_n \frac{\partial}{\partial r}\exp\left(r\ln\left(x_n\right)\right)\right\}_{r=0}\\
&= \exp\left\{\sum_{n=1}^{N} \lambda_n \exp\left\{r\ln x_n\right\}\ln\left(x_n\right)\right\}_{r=0} && = \exp\left\{\sum_{n=1}^{N} \lambda_n \ln\left(x_n\right)\right\}\\
&= \exp\left\{\sum_{n=1}^{N} \ln\left(x_n^{\lambda_n}\right)\right\} && = \exp\left\{\ln\prod_{n=1}^{N} x_n^{\lambda_n}\right\} = \prod_{n=1}^{N} x_n^{\lambda_n}
\end{aligned}$$

[11] Rudin (1976) page 85 ⟨4.4 Theorem⟩
[12] Rudin (1976) page 109 ⟨5.13 Theorem⟩

BY-NC-ND Copyright © 2019 Daniel J. Greenhoe

Definition M.4. *Let $(\!|x_n|\!)_1^N$ be a tuple. Let $(\!|\lambda_n|\!)_1^N$ be a tuple of weighting values.*

DEF

*The **harmonic mean** of $(\!|x_n|\!)$ is defined as* $\mu_h \triangleq \left(\sum_{n=1}^N \lambda_n \frac{1}{x_n}\right)^{-1}$ *where* $\sum_{n=1}^N \lambda_n = 1$

*The **geometric mean** of $(\!|x_n|\!)$ is defined as* $\mu_g \triangleq \sum_{n=1}^N x_n^{\lambda_n}$ *where* $\sum_{n=1}^N \lambda_n = 1$

*The **arithmetic mean** of $(\!|x_n|\!)$ is defined as* $\mu_a \triangleq \sum_{n=1}^N \lambda_n x_n$ *where* $\sum_{n=1}^N \lambda_n = 1$

*The **average** of $(\!|x_n|\!)$ is defined as* $\mu_a \triangleq \frac{1}{N}\sum_{n=1}^N x_n$

M.3 Inequalities on power means

Corollary M.1. [13] *Let $(\!|x_n|\!)_1^N$ be a tuple. Let $(\!|\lambda_n|\!)_1^N$ be a tuple of weighting values.*

COR

$$\min(\!|x_n|\!) \le \underbrace{\left(\sum_{n=1}^N \lambda_n \frac{1}{x_n}\right)^{-1}}_{\text{harmonic mean}} \le \underbrace{\prod_{n=1}^N x_n^{\lambda_n}}_{\text{geometric mean}} \le \underbrace{\sum_{n=1}^N \lambda_n x_n}_{\text{arithmetic mean}} \le \max(\!|x_n|\!) \quad \text{where} \quad \sum_{n=1}^N \lambda_n = 1$$

PROOF:

1. These five means are all special cases of the *power mean* $\mathsf{M}_{\phi(x:r)}$ (Definition M.3 page 292):
   $r = \infty$: $\max(\!|x_n|\!)$
   $r = 1$: arithmetic mean
   $r = 0$: geometric mean
   $r = -1$: harmonic mean
   $r = -\infty$: $\min(\!|x_n|\!)$

2. The inequalities follow directly from Theorem M.3 (page 292).

3. Generalized AM-GM inequality: If one is only concerned with the arithmetic mean and geometric mean, their relationship can be established directly using *Jensen's Inequality*:

$$\sum_{n=1}^N \lambda_n x_n = b^{\log_b\left(\sum_{n=1}^N \lambda_n x_n\right)} \ge b^{\left(\sum_{n=1}^N \lambda_n \log_b x_n\right)} \quad \text{by } \textit{Jensen's Inequality} \text{ (Theorem L.1 page 284)}$$

$$= \prod_{n=1}^N b^{(\lambda_n \log_b x_n)} = \prod_{n=1}^N b^{(\log_b x_n)\lambda_n} = \prod_{n=1}^N x_n^{\lambda_n}$$

[13] Bullen (2003) page 71, Bollobás (1999) page 5, Cauchy (1821), pages 457–459 ⟨Note II, theorem 17⟩, Jensen (1906) page 183

COPYRIGHT © 2019 DANIEL J. GREENHOE

Lemma M.2 (Young's Inequality). [14]

LEM

$$xy < \frac{x^p}{p} + \frac{y^q}{q} \quad \text{with } \frac{1}{p} + \frac{1}{q} = 1 \quad \forall 1 < p < \infty,\ x, y \geq 0, \quad \text{but } y \neq x^{p-1}$$

$$xy = \frac{x^p}{p} + \frac{y^q}{q} \quad \text{with } \frac{1}{p} + \frac{1}{q} = 1 \quad \forall 1 < p < \infty,\ x, y \geq 0, \quad \text{and } y = x^{p-1}$$

✎PROOF:

1. Proof that $\frac{1}{p-1} = q - 1$:

$$\begin{aligned}
\frac{1}{p} + \frac{1}{q} = 1 &\iff \frac{q}{q} + \frac{q}{p} = q \\
&\iff q\left(1 - \frac{1}{p}\right) = 1 \\
&\iff q = \frac{1}{1 - \frac{1}{p}} \\
&\iff q = \frac{p}{p-1} \\
&\iff q - 1 = \frac{p}{p-1} - \frac{p-1}{p-1} \\
&\iff q - 1 = \frac{p - (p-1)}{p-1} \\
&\iff q - 1 = \frac{1}{p-1}
\end{aligned}$$

2. Proof that $v = u^{p-1} \iff u = v^{q-1}$:

$$\begin{aligned}
u &= v^{\frac{1}{p-1}} && \text{by left hypothesis} \\
&= v^{q-1} && \text{by item (1)}
\end{aligned}$$

3. Proof that $v = u^{p-1}$ is propemonotonically increasing in u and $u = v^{q-1}$ is propemonotonically increasing in v:

$$\begin{aligned}
\frac{\mathrm{d}v}{\mathrm{d}u} &= \frac{\mathrm{d}}{\mathrm{d}u} u^{p-1} &&= (p-1)u^{p-2} &&> 0 \\
\frac{\mathrm{d}u}{\mathrm{d}v} &= \frac{\mathrm{d}}{\mathrm{d}v} v^{q-1} &&= (q-1)v^{q-2} &&> 0
\end{aligned}$$

4. Proof that $xy \leq \frac{x^p}{p} + \frac{y^q}{q}$:

$$\begin{aligned}
xy &\leq \int_0^x u^{p-1}\,\mathrm{d}u + \int_0^y v^{q-1}\,\mathrm{d}v \\
&= \left.\frac{u^p}{p}\right|_0^x + \left.\frac{v^q}{q}\right|_0^y \\
&= \frac{x^p}{p} + \frac{y^q}{q}
\end{aligned}$$



[14] Carothers (2000), page 43, Tolsted (1964), page 5, Maligranda (1995), page 257, Hardy et al. (1952) ⟨Theorem 24⟩, Young (1912) page 226

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

Theorem M.4 (Hölder's Inequality). [15] *Let* $(\!| x_n \in \mathbb{C} |\!)_1^N$ *and* $(\!| y_n \in \mathbb{C} |\!)_1^N$ *be complex* N*-tuples.*

THM

$$\underbrace{\sum_{n=1}^{N} |x_n y_n|}_{\|\boldsymbol{x}\cdot\boldsymbol{y}\|_1} \le \underbrace{\left(\sum_{n=1}^{N} |x_n|^p\right)^{\frac{1}{p}}}_{\|\boldsymbol{x}\|_p} \underbrace{\left(\sum_{n=1}^{N} |y_n|^q\right)^{\frac{1}{q}}}_{\|\boldsymbol{y}\|_q} \qquad \text{with} \qquad \frac{1}{p}+\frac{1}{q}=1 \qquad \forall\, 1<p<\infty$$

PROOF: Let $\|x_n\|_p \triangleq \left(\sum_{n=1}^{N} |x_n|^p\right)^{\frac{1}{p}}$.

$$\begin{aligned}
\sum_{n=1}^{N} |x_n y_n| &= \|(\!| x_n |\!)\|_p \, \|(\!| y_n |\!)\|_q \sum_{n=1}^{N} \frac{|x_n y_n|}{\|(\!| x_n |\!)\|_p \, \|(\!| y_n |\!)\|_q} \\
&= \|(\!| x_n |\!)\|_p \, \|(\!| y_n |\!)\|_q \sum_{n=1}^{N} \frac{|x_n|}{\|(\!| x_n |\!)\|_p} \frac{|y_n|}{\|(\!| y_n |\!)\|_q} \\
&\le \|(\!| x_n |\!)\|_p \, \|(\!| y_n |\!)\|_q \sum_{n=1}^{N} \left(\frac{1}{p}\frac{|x_n|^p}{\|(\!| x_n |\!)\|_p^p} + \frac{1}{q}\frac{|y_n|^q}{\|(\!| y_n |\!)\|_q^q}\right) && \text{by } \textit{Young's Inequality} \quad \text{(Lemma M.2 page 296)} \\
&= \|(\!| x_n |\!)\|_p \, \|(\!| y_n |\!)\|_q \left(\frac{1}{p}\cdot\frac{\sum |x_n|^p}{\|(\!| x_n |\!)\|_p^p} + \frac{1}{q}\cdot\frac{\sum |y_n|^q}{\|(\!| y_n |\!)\|_q^q}\right) \\
&= \|(\!| x_n |\!)\|_p \, \|(\!| y_n |\!)\|_q \left(\frac{1}{p}\frac{\|(\!| x_n |\!)\|_p^p}{\|(\!| x_n |\!)\|_p^p} + \frac{1}{q}\frac{\|(\!| y_n |\!)\|_q^q}{\|(\!| y_n |\!)\|_q^q}\right) && \text{by definition of } \|\cdot\| \\
&= \|(\!| x_n |\!)\|_p \, \|(\!| y_n |\!)\|_q \underbrace{\left(\frac{1}{p}+\frac{1}{q}\right)}_{1} \\
&= \|(\!| x_n |\!)\|_p \, \|(\!| y_n |\!)\|_q && \text{by } \frac{1}{p}+\frac{1}{q}=1 \text{ constraint}
\end{aligned}$$

Theorem M.5 (Minkowski's Inequality for sequences). [16] *Let* $(\!| x_n \in \mathbb{C} |\!)_1^N$ *and* $(\!| y_n \in \mathbb{C} |\!)_1^N$ *be complex* N*-tuples.*

THM

$$\left(\sum_{n=1}^{N} |x_n+y_n|^p\right)^{\frac{1}{p}} \le \left(\sum_{n=1}^{N} |x_n|^p\right)^{\frac{1}{p}} + \left(\sum_{n=1}^{N} |y_n|^p\right)^{\frac{1}{p}} \qquad \forall\, 1<p<\infty$$

PROOF:

1. Define $q \triangleq \dfrac{p}{p-1}$

[15] Bullen (2003) page 178 ⟨2.1⟩, Carothers (2000), page 44, Tolsted (1964), page 6, Maligranda (1995), page 257, Hardy et al. (1952) ⟨Theorem 11⟩, Hölder (1889)

[16] Bullen (2003) page 179, Carothers (2000), page 44, Tolsted (1964), page 7, Maligranda (1995), page 258, Hardy et al. (1952) ⟨Theorem 24⟩, Minkowski (1910) page 115

Copyright © 2019 Daniel J. Greenhoe

2. Define $\|\boldsymbol{x}\|_p \triangleq \left(\sum_{n=1}^{N} |x_n|^p\right)^{\frac{1}{p}}$.

3. Proof that $\|x_n + y_n\|_p \le \|x_n\|_p + \|y_n\|_p$:

$$\boxed{\|x_n + y_n\|_p^p}$$

$$= \sum_{n=1}^{N} |x_n + y_n|^p \qquad \text{by definition of } \|\cdot\|_p \qquad \text{(definition 2 page 298)}$$

$$= \sum_{n=1}^{N} |x_n + y_n||x_n + y_n|^{p-1} \qquad \text{by } \textit{homogeneous} \text{ property of } |\cdot|$$

$$\boxed{\le} \sum_{n=1}^{N} |x_n||x_n + y_n|^{p-1} + \sum_{n=1}^{N} |y_n||x_n + y_n|^{p-1} \qquad \text{by } \textit{subadditive} \text{ property of } |\cdot|$$

$$= \sum_{n=1}^{N} |x_n(x_n + y_n)^{p-1}| + \sum_{n=1}^{N} |y_n(x_n + y_n)^{p-1}| \qquad \text{by } \textit{homogeneous} \text{ property of } |\cdot|$$

$$\boxed{\le} \|x_n\|_p \|(x_n + y_n)^{p-1}\|_q + \|y_n\|_p \|(x_n + y_n)^{p-1}\|_q \qquad \text{by } \textit{Hölder's Inequality} \qquad \text{(Theorem M.4 page 297)}$$

$$= \left(\|x_n\|_p + \|y_n\|_p\right) \|(x_n + y_n)^{p-1}\|_q$$

$$= \left(\|x_n\|_p + \|y_n\|_p\right)\left(\sum_{n=1}^{N} |(x_n + y_n)^{p-1}|^q\right)^{\frac{1}{q}} \qquad \text{by definition of } \|\cdot\|_p \qquad \text{(definition 2 page 298)}$$

$$= \left(\|x_n\|_p + \|y_n\|_p\right)\left(\sum_{n=1}^{N} |(x_n + y_n)^{p-1}|^{\frac{p}{p-1}}\right)^{\frac{p-1}{p}} \qquad \text{by definition 1}$$

$$= \left(\|x_n\|_p + \|y_n\|_p\right)\left(\sum_{n=1}^{N} |(x_n + y_n)|^p\right)^{\frac{p-1}{p}}$$

$$= \left(\|x_n\|_p + \|y_n\|_p\right) \|x_n + y_n\|^{p-1} \qquad \text{by definition of } \|\cdot\|_p \qquad \text{(definition 2 page 298)}$$

$$\Longrightarrow \quad \boxed{\|x_n + y_n\|_p \le \|x_n\|_p + \|y_n\|_p}$$

✍

❝*Cauchy is the only one occupied with pure mathematics: Poisson, Fourier, Ampere, etc., busy themselves exclusively with magnetism and other physical subjects. Mr. Laplace writes nothing now, I believe.*❞

Niels Henrik Abel in an 1826 letter [17]

Theorem M.6 (Cauchy-Schwarz Inequality for sequences). [18] *Let* $(x_n \in \mathbb{C})_1^N$ *and* $(y_n \in \mathbb{C})_1^N$ *be complex* N*-tuples.*

[17] quote: Bell (1986) page 318 ⟨Chapter 17. "Genius and Poverty" "Abel (1802–1829)"⟩, Boyer and Merzbach (2011) page 462 ⟨without "Mr. Laplace…" portion⟩. image: http://en.wikipedia.org/wiki/File:Augustin-Louis_Cauchy_1901.jpg, public domain

[18] Aliprantis and Burkinshaw (1998), page 278, Scharz (1885), Bouniakowsky (1859), Hardy et al. (1952) page 25 ⟨Theorem 11⟩, Cauchy (1821), page 455 ⟨???⟩

BY-NC-ND Copyright © 2019 Daniel J. Greenhoe

THM

$$\left|\sum_{n=1}^{N} x_n y_n^*\right|^2 \le \left(\sum_{n=1}^{N} |x_n|^2\right)\left(\sum_{n=1}^{N} |y_n|^2\right) \qquad \forall \boldsymbol{x},\boldsymbol{y}\in\boldsymbol{X}$$

$$\left|\sum_{n=1}^{N} x_n y_n^*\right|^2 = \left(\sum_{n=1}^{N} |x_n|^2\right)\left(\sum_{n=1}^{N} |y_n|^2\right) \iff \exists a \in \mathbb{C} \quad \text{such that} \quad \boldsymbol{y} = a\boldsymbol{x} \qquad \forall \boldsymbol{x},\boldsymbol{y}\in\boldsymbol{X}$$

✎PROOF:

1. The *Cauchy-Schwarz Inequality for sequences* is a special case of the *Hölder inequality* (Theorem M.4 page 297) for $p = q = 2$.

2. Alternatively, the *Cauchy-Schwarz inequality for sequences* is a special case of the *Cauchy-Schwarz inequality for inner-product spaces*:

   (a) $\langle x_n \mid y_n\rangle \triangleq \sum_{n=1}^{N} x_n y_n$ is an inner-product and $\left(\llparenthesis x_n \rrparenthesis, \langle \triangle \mid \triangledown\rangle\right)$ is an inner-product space.

   (b) By the more general *Cauchy-Schwarz Inequality for inner-product spaces*,

$$\begin{aligned}
\left(\sum_{n=1}^{N} a_n\lambda_n\right)^2 &\triangleq \langle a_n \mid \lambda_n\rangle^2 && \text{by definition of } \langle x_n \mid y_n\rangle \\
&\boxed{\le} \|x_n\|^2 \|y_n\|^2 && \text{by } \textit{Cauchy-Schwarz Inequality for inner-product spaces} \text{ (Theorem J.2 page 256)} \\
&\triangleq \left(\sum_{n=1}^{N} x_n^2\right)\left(\sum_{n=1}^{N} y_n^2\right) && \text{by definition of } \|\cdot\|
\end{aligned}$$

3. Not only does the *Hölder inequality* imply the *Cauchy-Schwarz inequality*, but somewhat surprisingly, the converse is also true: The Cauchy-Schwarz inequality implies the Hölder inequality.[19]

✏

Proposition M.1. [20]

PRP

$$(x+y)^p \le 2^p(x^p + y^p) \qquad \forall x, y \ge 0,\ 1 < p < \infty$$

✎PROOF:

$$\begin{aligned}
(x+y)^p &\le (2\max\{x,y\})^p \\
&= 2^p(\max\{x,y\})^p \\
&= 2^p(\max\{x^p, y^p\}) \\
&\le 2^p(x^p + y^p)
\end{aligned}$$

✏

[19] Bullen (2003) pages 183–185 ⟨Theorem 5⟩
[20] Carothers (2000), page 43

COPYRIGHT © 2019 DANIEL J. GREENHOE

M.4 Power Sums

Theorem M.7 (Geometric Series). [21]

THM
$$\sum_{k=0}^{n-1} r^k = \frac{1-r^n}{1-r} \qquad \forall r \in \mathbb{C}\backslash 0$$

✎PROOF:

$$\boxed{\sum_{k=0}^{n-1} r^k} = \left(\frac{1}{1-r}\right)\left[(1-r)\sum_{k=0}^{n-1} r^k\right] = \left(\frac{1}{1-r}\right)\left[\sum_{k=0}^{n-1} r^k - r\sum_{k=0}^{n-1} r^k\right] = \left(\frac{1}{1-r}\right)\left[\sum_{k=0}^{n-1} r^k - \left(\sum_{k=0}^{n-1} r^k - 1 + r^n\right)\right]$$
$$= \left(\frac{1}{1-r}\right)\left[1-r^n\right] = \boxed{\frac{1-r^n}{1-r}}$$

✏

Lemma M.3. *Let* $\mathsf{f}(x)$ *be a function.*

LEM
$$\mathsf{S}(x) \triangleq \sum_{n\in\mathbb{Z}} \mathsf{f}(x+n\tau) = \mathsf{S}(x+\tau) \qquad (\mathsf{S}(x) \text{ is PERIODIC with period } \tau)$$

✎PROOF:

$$\mathsf{S}(x+\tau) \triangleq \sum_{n\in\mathbb{Z}} \mathsf{f}(x+\tau+n\tau) = \sum_{n\in\mathbb{Z}} \mathsf{f}(x+(n+1)\tau) = \sum_{m\in\mathbb{Z}} \mathsf{f}(x+m\tau) \quad (\text{where } m \triangleq n+1)$$
$$\triangleq \mathsf{S}(x)$$

✏

Proposition M.2 (Power Sums). [22]

PRP
$$\sum_{k=1}^{n} k = \frac{n(n+1)}{2} \qquad \forall n\in\mathbb{N} \qquad \sum_{k=1}^{n} k^3 = \frac{n^2(n+1)^2}{4} \qquad \forall n\in\mathbb{N}$$
$$\sum_{k=1}^{n} k^2 = \frac{n(n+1)(2n+1)}{6} \qquad \forall n\in\mathbb{N} \qquad \sum_{k=1}^{n} k^4 = \frac{n(n+1)(2n+1)(3n^2+3n-1)}{30} \qquad \forall n\in\mathbb{N}$$

✎PROOF:

1. Proof that $\sum_{k=1}^{n} k = \frac{n(n+1)}{2}$: (proof by induction)

$$\sum_{k=1}^{n=1} k = 1 = \frac{1(1+1)}{2} = \left.\frac{n(n+1)}{2}\right|_{n=1}$$
$$\sum_{k=1}^{n+1} k = \left(\sum_{k=1}^{n} k\right) + \left(n+1\right) = \underbrace{\left(\frac{n(n+1)}{2}\right) + \left(n+1\right)}_{\text{by left hypothesis}} = \left(n+1\right)\left(\frac{n}{2}+1\right)$$
$$= \left(n+1\right)\left(\frac{n+2}{2}\right) = \frac{(n+1)(n+2)}{2}$$

[21] Hall and Knight (1894), page 39 ⟨article 55⟩
[22] Amann and Escher (2008) pages 51–57, Menini and Oystaeyen (2004) page 91 ⟨Exercises 5.36–5.39⟩

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

2. Proof that $\sum_{k=1}^{n} k^2 = \frac{n(n+1)(2n+1)}{6}$: (proof by induction)

$$\sum_{k=1}^{n=1} k^2 = 1 = \frac{1(1+1)(2+1)}{6} = \left.\frac{n(n+1)(2n+1)}{6}\right|_{n=1}$$

$$\sum_{k=1}^{n+1} k^2 = \left(\sum_{k=1}^{n} k^2\right) + \left(n+1\right)^2 = \underbrace{\left(\frac{n(n+1)(2n+1)}{6}\right) + \left(n+1\right)^2}_{\text{by left hypothesis}} = \left(n+1\right)\left(\frac{n(2n+1)+6(n+1)}{6}\right)$$

$$= \left(n+1\right)\left(\frac{2n^2+7n+6}{6}\right) = \left(n+1\right)\left(\frac{(n+2)(2n+3)}{6}\right) = \frac{(n+1)[(n+1)+1][2(n+1)+1]}{6}$$

✍

Copyright © 2019 Daniel J. Greenhoe

Copyright © 2019 Daniel J. Greenhoe

APPENDIX N

OPERATORS ON LINEAR SPACES

❝And I am not afraid to say that there is a way to advance algebra as far beyond what Vieta and Descartes have left us as Vieta and Descartes carried it beyond the ancients....we need still another analysis which is distinctly geometrical or linear, and which will express situation directly as algebra expresses magnitude directly.❞

Gottfried Leibniz (1646–1716), German mathematician, in a September 8, 1679 letter to Christian Huygens. [1]

N.1 Operators on linear spaces

N.1.1 Operator Algebra

An operator is simply a function that maps from a linear space to another linear space (or to the same linear space).

Definition N.1. [2]

DEF *A function* $\mathbf{A}$ *in* Y^X *is an* ***operator*** *in* Y^X *if* X *and* Y *are both* LINEAR SPACES *(Definition G.1 page 207)*.

Two operators $\mathbf{A}$ and $\mathbf{B}$ in Y^X are **equal** if $\mathbf{A}x = \mathbf{B}x$ for all $x \in X$. The inverse relation of an operator $\mathbf{A}$ in Y^X always exists as a *relation* in 2^{XY}, but may not always be a *function* (may not always be an operator) in Y^X.

The operator $\mathbf{I} \in X^X$ is the *identity* operator if $\mathbf{I}x = \mathbf{I}$ for all $x \in X$.

Definition N.2. [3] *Let* X^X *be the set of all operators with from a* LINEAR SPACE X *to* X. *Let* $\mathbf{I}$ *be an operator in* X^X. *Let* $①(X)$ *be the* IDENTITY ELEMENT *in* X^X.

[1] quote: Leibniz (1679) pages 248–249
image: http://en.wikipedia.org/wiki/File:Gottfried_Wilhelm_von_Leibniz.jpg, public domain

[2] Heil (2011) page 42

[3] Michel and Herget (1993) page 411



Figure N.1: Some operator types

DEF $\mathbf{I}$ *is the* ***identity operator*** *in* $\boldsymbol{X}^{\boldsymbol{X}}$ *if* $\quad \mathbf{I} = \mathbb{1}(\boldsymbol{X})$.

N.1.2 Linear operators

Definition N.3. [4] *Let* $\boldsymbol{X} \triangleq \big(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times})\big)$ *and* $\boldsymbol{Y} \triangleq \big(Y, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times})\big)$ *be linear spaces.*

DEF *An operator* $\mathbf{L} \in \boldsymbol{Y}^{\boldsymbol{X}}$ *is* ***linear*** *if*

1. $\mathbf{L}(\boldsymbol{x}+\boldsymbol{y}) = \mathbf{L}\boldsymbol{x} + \mathbf{L}\boldsymbol{y} \quad \forall \boldsymbol{x},\boldsymbol{y} \in \boldsymbol{X}$ (ADDITIVE) *and*
2. $\mathbf{L}(\alpha\boldsymbol{x}) = \alpha\mathbf{L}\boldsymbol{x} \quad \forall \boldsymbol{x} \in \boldsymbol{X}, \ \forall \alpha \in \mathbb{F}$ (HOMOGENEOUS).

The set of all linear operators from $\boldsymbol{X}$ *to* $\boldsymbol{Y}$ *is denoted* $\mathcal{L}(\boldsymbol{X}, \boldsymbol{Y})$ *such that*

$$\mathcal{L}(\boldsymbol{X}, \boldsymbol{Y}) \triangleq \big\{\mathbf{L} \in \boldsymbol{Y}^{\boldsymbol{X}} \,|\, \mathbf{L} \text{ is linear}\big\} \quad .$$

Theorem N.1. [5] *Let* $\mathbf{L}$ *be an operator from a linear space* $\boldsymbol{X}$ *to a linear space* $\boldsymbol{Y}$*, both over a field* $\mathbb{F}$.

THM

$$\mathbf{L} \text{ is LINEAR} \quad \Longrightarrow \quad \begin{cases} 1. \ \mathbf{L}\mathbb{0} = \mathbb{0} & \text{and} \\ 2. \ \mathbf{L}(-\boldsymbol{x}) = -(\mathbf{L}\boldsymbol{x}) \quad \forall \boldsymbol{x} \in \boldsymbol{X} & \text{and} \\ 3. \ \mathbf{L}(\boldsymbol{x}-\boldsymbol{y}) = \mathbf{L}\boldsymbol{x} - \mathbf{L}\boldsymbol{y} \quad \forall \boldsymbol{x},\boldsymbol{y} \in \boldsymbol{X} & \text{and} \\ 4. \ \mathbf{L}\left(\sum_{n=1}^{N} \alpha_n \boldsymbol{x}_n\right) = \sum_{n=1}^{N} \alpha_n \big(\mathbf{L}\boldsymbol{x}_n\big) \quad \boldsymbol{x}_n \in \boldsymbol{X}, \alpha_n \in \mathbb{F} & \end{cases}$$

✎PROOF:

[4] Kubrusly (2001) page 55, Aliprantis and Burkinshaw (1998) page 224, Hilbert et al. (1927) page 6, Stone (1932) page 33

[5] Berberian (1961) page 79 ⟨Theorem IV.1.1⟩

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

1. Proof that $\mathbf{L}\mathbb{0} = \mathbb{0}$:

$$\begin{aligned}
\mathbf{L}\mathbb{0} &= \mathbf{L}(0 \cdot \mathbb{0}) && \text{by additive identity property (Theorem G.1 page 209)}\\
&= 0 \cdot (\mathbf{L}\mathbb{0}) && \text{by \textit{homogeneous} property of } \mathbf{L} \text{ (Definition N.3 page 304)}\\
&= \mathbb{0} && \text{by additive identity property (Theorem G.1 page 209)}
\end{aligned}$$

2. Proof that $\mathbf{L}(-\boldsymbol{x}) = -(\mathbf{L}\boldsymbol{x})$:

$$\begin{aligned}
\mathbf{L}(-\boldsymbol{x}) &= \mathbf{L}(-1 \cdot \boldsymbol{x}) && \text{by additive inverse property (Theorem G.2 page 210)}\\
&= -1 \cdot (\mathbf{L}\boldsymbol{x}) && \text{by \textit{homogeneous} property of } \mathbf{L} \text{ (Definition N.3 page 304)}\\
&= -(\mathbf{L}\boldsymbol{x}) && \text{by additive inverse property (Theorem G.2 page 210)}
\end{aligned}$$

3. Proof that $\mathbf{L}(\boldsymbol{x} - \boldsymbol{y}) = \mathbf{L}\boldsymbol{x} - \mathbf{L}\boldsymbol{y}$:

$$\begin{aligned}
\mathbf{L}(\boldsymbol{x} - \boldsymbol{y}) &= \mathbf{L}(\boldsymbol{x} + (-\boldsymbol{y})) && \text{by additive inverse property (Theorem G.2 page 210)}\\
&= \mathbf{L}(\boldsymbol{x}) + \mathbf{L}(-\boldsymbol{y}) && \text{by \textit{linearity} property of } \mathbf{L} \text{ (Definition N.3 page 304)}\\
&= \mathbf{L}\boldsymbol{x} - \mathbf{L}\boldsymbol{y} && \text{by 2.}
\end{aligned}$$

4. Proof that $\mathbf{L}\left(\sum_{n=1}^{N} \alpha_n \boldsymbol{x}_n\right) = \sum_{n=1}^{N} \alpha_n (\mathbf{L}\boldsymbol{x}_n)$:

   (a) Proof for $N = 1$:

$$\begin{aligned}
\mathbf{L}\left(\sum_{n=1}^{N} \alpha_n \boldsymbol{x}_n\right) &= \mathbf{L}(\alpha_1 \boldsymbol{x}_1) && \text{by } N = 1 \text{ hypothesis}\\
&= \alpha_1 (\mathbf{L}\boldsymbol{x}_1) && \text{by \textit{homogeneous} property of Definition N.3 page 304}
\end{aligned}$$

   (b) Proof that N case $\implies$ $N+1$ case:

$$\begin{aligned}
\mathbf{L}\left(\sum_{n=1}^{N+1} \alpha_n \boldsymbol{x}_n\right) &= \mathbf{L}\left(\alpha_{N+1}\boldsymbol{x}_{N+1} + \sum_{n=1}^{N} \alpha_n \boldsymbol{x}_n\right)\\
&= \mathbf{L}(\alpha_{N+1}\boldsymbol{x}_{N+1}) + \mathbf{L}\left(\sum_{n=1}^{N} \alpha_n \boldsymbol{x}_n\right) && \text{by \textit{linearity} property of Definition N.3 page 304}\\
&= \alpha_{N+1}\mathbf{L}(\boldsymbol{x}_{N+1}) + \sum_{n=1}^{N} \mathbf{L}(\alpha_n \boldsymbol{x}_n) && \text{by left } N+1 \text{ hypothesis}\\
&= \sum_{n=1}^{N+1} \mathbf{L}(\alpha_n \boldsymbol{x}_n)
\end{aligned}$$

✎

Theorem N.2. [6] *Let $\mathcal{L}(\boldsymbol{X}, \boldsymbol{Y})$ be the set of all linear operators from a linear space $\boldsymbol{X}$ to a linear space $\boldsymbol{Y}$. Let $\mathcal{N}(\mathbf{L})$ be the* NULL SPACE *of an operator $\mathbf{L}$ in $\boldsymbol{Y}^{\boldsymbol{X}}$ and $\mathcal{I}(\mathbf{L})$ the* IMAGE SET *of $\mathbf{L}$ in $\boldsymbol{Y}^{\boldsymbol{X}}$.*

THM			
	$\mathcal{L}(\boldsymbol{X}, \boldsymbol{Y})$	*is a linear space*	*(space of linear transforms)*
	$\mathcal{N}(\mathbf{L})$	*is a linear subspace of* $\boldsymbol{X}$	$\forall \mathbf{L} \in \boldsymbol{Y}^{\boldsymbol{X}}$
	$\mathcal{I}(\mathbf{L})$	*is a linear subspace of* $\boldsymbol{Y}$	$\forall \mathbf{L} \in \boldsymbol{Y}^{\boldsymbol{X}}$

✎PROOF:

[6] Michel and Herget (1993) pages 98–104, Berberian (1961) pages 80–85 ⟨Theorem IV.1.4 and Theorem IV.3.1⟩

COPYRIGHT © 2019 DANIEL J. GREENHOE

1. Proof that $\mathcal{N}(\mathbf{L})$ is a linear subspace of $\boldsymbol{X}$:
   (a) $\mathbb{0} \in \mathcal{N}(\mathbf{L}) \implies \mathcal{N}(\mathbf{L}) \neq \varnothing$
   (b) $\mathcal{N}(\mathbf{L}) \triangleq \{\boldsymbol{x} \in \boldsymbol{X} | \mathbf{L}\boldsymbol{x} = \mathbb{0}\} \subseteq \boldsymbol{X}$
   (c) $\boldsymbol{x} + \boldsymbol{y} \in \mathcal{N}(\mathbf{L}) \implies \mathbb{0} = \mathbf{L}(\boldsymbol{x} + \boldsymbol{y}) = \mathbf{L}(\boldsymbol{y} + \boldsymbol{x}) \implies \boldsymbol{y} + \boldsymbol{x} \in \mathcal{N}(\mathbf{L})$
   (d) $\alpha \in \mathbb{F},\ \boldsymbol{x} \in \boldsymbol{X} \implies \mathbb{0} = \mathbf{L}\boldsymbol{x} \implies \mathbb{0} = \alpha\mathbf{L}\boldsymbol{x} \implies \mathbb{0} = \mathbf{L}(\alpha\boldsymbol{x}) \implies \alpha\boldsymbol{x} \in \mathcal{N}(\mathbf{L})$
2. Proof that $\mathcal{I}(\mathbf{L})$ is a linear subspace of $\boldsymbol{Y}$:
   (a) $\mathbb{0} \in \mathcal{I}(\mathbf{L}) \implies \mathcal{I}(\mathbf{L}) \neq \varnothing$
   (b) $\mathcal{I}(\mathbf{L}) \triangleq \{\boldsymbol{y} \in \boldsymbol{Y} | \exists \boldsymbol{x} \in \boldsymbol{X} \quad \text{such that} \quad \boldsymbol{y} = \mathbf{L}\boldsymbol{x}\} \subseteq \boldsymbol{Y}$
   (c) $\boldsymbol{x} + \boldsymbol{y} \in \mathcal{I}(\mathbf{L}) \implies \exists \boldsymbol{v} \in \boldsymbol{X} \quad \text{such that} \quad \mathbf{L}\boldsymbol{v} = \boldsymbol{x} + \boldsymbol{y} = \boldsymbol{y} + \boldsymbol{x} \implies \boldsymbol{y} + \boldsymbol{x} \in \mathcal{I}(\mathbf{L})$
   (d) $\alpha \in \mathbb{F},\ \boldsymbol{x} \in \mathcal{I}(\mathbf{L}) \implies \exists \boldsymbol{x} \in \boldsymbol{X} \quad \text{such that} \quad \boldsymbol{y} = \mathbf{L}\boldsymbol{x} \implies \alpha\boldsymbol{y} = \alpha\mathbf{L}\boldsymbol{x} = \mathbf{L}(\alpha\boldsymbol{x}) \implies \alpha\boldsymbol{x} \in \mathcal{I}(\mathbf{L})$

✏

Example N.1. [7] Let $\mathcal{C}([a : b], \mathbb{R})$ be the set of all *continuous* functions from the closed real interval $[a : b]$ to $\mathbb{R}$.

EX $\mathcal{C}([a : b], \mathbb{R})$ is a linear space.

Theorem N.3. [8] *Let $\mathcal{L}(\boldsymbol{X}, \boldsymbol{Y})$ be the set of linear operators from a linear space $\boldsymbol{X}$ to a linear space $\boldsymbol{Y}$. Let $\mathcal{N}(\mathbf{L})$ be the* NULL SPACE *of a linear operator $\mathbf{L} \in \mathcal{L}(\boldsymbol{X}, \boldsymbol{Y})$.*

THM

$$\begin{aligned} \mathbf{L}\boldsymbol{x} = \mathbf{L}\boldsymbol{y} \quad &\iff \quad \boldsymbol{x} - \boldsymbol{y} \in \mathcal{N}(\mathbf{L}) \\ \mathbf{L} \textit{ is } \text{INJECTIVE} \quad &\iff \quad \mathcal{N}(\mathbf{L}) = \{\mathbb{0}\} \end{aligned}$$

✎PROOF:

1. Proof that $\mathbf{L}\boldsymbol{x} = \mathbf{L}\boldsymbol{y} \implies \boldsymbol{x} - \boldsymbol{y} \in \mathcal{N}(\mathbf{L})$:

$$\begin{aligned} \mathbf{L}(\boldsymbol{x} - \boldsymbol{y}) &= \mathbf{L}\boldsymbol{x} - \mathbf{L}\boldsymbol{y} && \text{by Theorem N.1 page 304} \\ &= \mathbb{0} && \text{by left hypothesis} \\ &\implies \quad \boldsymbol{x} - \boldsymbol{y} \in \mathcal{N}(\mathbf{L}) && \text{by definition of } \textit{null space} \end{aligned}$$

2. Proof that $\mathbf{L}\boldsymbol{x} = \mathbf{L}\boldsymbol{y} \impliedby \boldsymbol{x} - \boldsymbol{y} \in \mathcal{N}(\mathbf{L})$:

$$\begin{aligned} \mathbf{L}\boldsymbol{y} &= \mathbf{L}\boldsymbol{y} + \mathbb{0} && \text{by definition of linear space (Definition G.1 page 207)} \\ &= \mathbf{L}\boldsymbol{y} + \mathbf{L}(\boldsymbol{x} - \boldsymbol{y}) && \text{by right hypothesis} \\ &= \mathbf{L}\boldsymbol{y} + (\mathbf{L}\boldsymbol{x} - \mathbf{L}\boldsymbol{y}) && \text{by Theorem N.1 page 304} \\ &= (\mathbf{L}\boldsymbol{y} - \mathbf{L}\boldsymbol{y}) + \mathbf{L}\boldsymbol{x} && \text{by } \textit{associative} \text{ and } \textit{commutative} \text{ properties (Definition G.1 page 207)} \\ &= \mathbf{L}\boldsymbol{x} \end{aligned}$$

3. Proof that $\mathbf{L}$ is *injective* $\iff \mathcal{N}(\mathbf{L}) = \{\mathbb{0}\}$:

$$\begin{aligned} \mathbf{L} \text{ is } \textit{injective} &\iff \{(\mathbf{L}\boldsymbol{x} = \mathbf{L}\boldsymbol{y} \quad \iff \quad \boldsymbol{x} = \boldsymbol{y}) \quad \forall \boldsymbol{x}, \boldsymbol{y} \in X\} \\ &\iff \left\{\left[\mathbf{L}\boldsymbol{x} - \mathbf{L}\boldsymbol{y} = \mathbb{0} \quad \iff \quad (\boldsymbol{x} - \boldsymbol{y}) = \mathbb{0}\right] \quad \forall \boldsymbol{x}, \boldsymbol{y} \in X\right\} \\ &\iff \left\{\left[\mathbf{L}(\boldsymbol{x} - \boldsymbol{y}) = \mathbb{0} \quad \iff \quad (\boldsymbol{x} - \boldsymbol{y}) = \mathbb{0}\right] \quad \forall \boldsymbol{x}, \boldsymbol{y} \in X\right\} \\ &\iff \mathcal{N}(\mathbf{L}) = \{\mathbb{0}\} \end{aligned}$$

[7] Eidelman et al. (2004) page 3
[8] Berberian (1961) page 88 ⟨Theorem IV.1.4⟩

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

Theorem N.4. [9] *Let $\boldsymbol{W}$, $\boldsymbol{X}$, $\boldsymbol{Y}$, and $\boldsymbol{Z}$ be linear spaces over a field $\mathbb{F}$.*

THM

1.	$\mathbf{L}(\mathbf{MN})$	$=$	$(\mathbf{LM})\mathbf{N}$	$\forall \mathbf{L}\in\mathcal{L}(\boldsymbol{Z},\boldsymbol{W}), \mathbf{M}\in\mathcal{L}(\boldsymbol{Y},\boldsymbol{Z}), \mathbf{N}\in\mathcal{L}(\boldsymbol{X},\boldsymbol{Y})$	(ASSOCIATIVE)
2.	$\mathbf{L}\big(\mathbf{M}\mathbin{\dot{+}}\mathbf{N}\big)$	$=$	$(\mathbf{LM})\mathbin{\dot{+}}(\mathbf{LN})$	$\forall \mathbf{L}\in\mathcal{L}(\boldsymbol{Y},\boldsymbol{Z}), \mathbf{M}\in\mathcal{L}(\boldsymbol{X},\boldsymbol{Y}), \mathbf{N}\in\mathcal{L}(\boldsymbol{X},\boldsymbol{Y})$	(LEFT DISTRIBUTIVE)
3.	$\big(\mathbf{L}\mathbin{\dot{+}}\mathbf{M}\big)\mathbf{N}$	$=$	$(\mathbf{LN})\mathbin{\dot{+}}(\mathbf{MN})$	$\forall \mathbf{L}\in\mathcal{L}(\boldsymbol{Y},\boldsymbol{Z}), \mathbf{M}\in\mathcal{L}(\boldsymbol{Y},\boldsymbol{Z}), \mathbf{N}\in\mathcal{L}(\boldsymbol{X},\boldsymbol{Y})$	(RIGHT DISTRIBUTIVE)
4.	$\alpha(\mathbf{LM})$	$=$	$(\alpha\mathbf{L})\mathbf{M} = \mathbf{L}(\alpha\mathbf{M})$	$\forall \mathbf{L}\in\mathcal{L}(\boldsymbol{Y},\boldsymbol{Z}), \mathbf{M}\in\mathcal{L}(\boldsymbol{X},\boldsymbol{Y}), \alpha\in\mathbb{F}$	(HOMOGENEOUS)

PROOF:

1. Proof that $\mathbf{L}(\mathbf{MN}) = (\mathbf{LM})\mathbf{N}$: Follows directly from property of *associative* operators.

2. Proof that $\mathbf{L}\big(\mathbf{M}\mathbin{\dot{+}}\mathbf{N}\big) = (\mathbf{LM})\mathbin{\dot{+}}(\mathbf{LN})$:

$$\begin{aligned}
\big[\mathbf{L}\big(\mathbf{M}\mathbin{\dot{+}}\mathbf{N}\big)\big]\boldsymbol{x} &= \mathbf{L}\big[\big(\mathbf{M}\mathbin{\dot{+}}\mathbf{N}\big)\boldsymbol{x}\big] \\
&= \mathbf{L}\big[(\mathbf{M}\boldsymbol{x})\mathbin{\dot{+}}(\mathbf{N}\boldsymbol{x})\big] \\
&= [\mathbf{L}(\mathbf{M}\boldsymbol{x})]\mathbin{\dot{+}}[\mathbf{L}(\mathbf{N}\boldsymbol{x})] && \text{by } \textit{additive} \text{ property Definition N.3 page 304} \\
&= [(\mathbf{LM})\boldsymbol{x}]\mathbin{\dot{+}}[(\mathbf{LN})\boldsymbol{x}]
\end{aligned}$$

3. Proof that $\big(\mathbf{L}\mathbin{\dot{+}}\mathbf{M}\big)\mathbf{N} = (\mathbf{LN})\mathbin{\dot{+}}(\mathbf{MN})$: Follows directly from property of *associative* operators.

4. Proof that $\alpha(\mathbf{LM}) = (\alpha\mathbf{L})\mathbf{M}$: Follows directly from *associative* property of linear operators.

5. Proof that $\alpha(\mathbf{LM}) = \mathbf{L}(\alpha\mathbf{M})$:

$$\begin{aligned}
[\alpha(\mathbf{LM})]\boldsymbol{x} &= \alpha[(\mathbf{LM})\boldsymbol{x}] \\
&= \mathbf{L}[\alpha(\mathbf{M}\boldsymbol{x})] && \text{by } \textit{homogeneous} \text{ property Definition N.3 page 304} \\
&= \mathbf{L}[(\alpha\mathbf{M})\boldsymbol{x}] \\
&= [\mathbf{L}(\alpha\mathbf{M})]\boldsymbol{x}
\end{aligned}$$

Theorem N.5 (Fundamental theorem of linear equations)**.** *Michel and Herget (1993) page 99 Let $\boldsymbol{Y}^{\boldsymbol{X}}$ be the set of all operators from a linear space $\boldsymbol{X}$ to a linear space $\boldsymbol{Y}$. Let $\mathcal{N}(\mathbf{L})$ be the* NULL SPACE *of an operator $\mathbf{L}$ in $\boldsymbol{Y}^{\boldsymbol{X}}$ and $\mathcal{I}(\mathbf{L})$ the* IMAGE SET *of $\mathbf{L}$ in $\boldsymbol{Y}^{\boldsymbol{X}}$ (Definition ?? page ??).*

THM

$$\dim \mathcal{I}(\mathbf{L}) + \dim \mathcal{N}(\mathbf{L}) = \dim \boldsymbol{X} \qquad \forall \mathbf{L}\in\boldsymbol{Y}^{\boldsymbol{X}}$$

PROOF: Let $\{\psi_k | k = 1, 2, \ldots, p\}$ be a basis for $\boldsymbol{X}$ constructed such that $\{\psi_{p-n+1}, \psi_{p-n+2}, \ldots, \psi_p\}$ is a basis for

[9] Berberian (1961) page 88 ⟨Theorem IV.5.1⟩

COPYRIGHT © 2019 DANIEL J. GREENHOE

$\mathcal{N}(\mathbf{L})$.

Let $p \triangleq \dim \boldsymbol{X}$.
Let $n \triangleq \dim \mathcal{N}(\mathbf{L})$.

$$\begin{aligned}
\dim \mathcal{I}(\mathbf{L}) &= \dim \{\boldsymbol{y} \in \boldsymbol{Y} | \exists \boldsymbol{x} \in \boldsymbol{X} \quad \text{such that} \quad \boldsymbol{y} = \mathbf{L}\boldsymbol{x}\} \\
&= \dim \left\{ \boldsymbol{y} \in \boldsymbol{Y} | \exists(\alpha_1, \alpha_2, \dots, \alpha_p) \quad \text{such that} \quad \boldsymbol{y} = \mathbf{L} \sum_{k=1}^{p} \alpha_k \psi_k \right\} \\
&= \dim \left\{ \boldsymbol{y} \in \boldsymbol{Y} | \exists(\alpha_1, \alpha_2, \dots, \alpha_p) \quad \text{such that} \quad \boldsymbol{y} = \sum_{k=1}^{p} \alpha_k \mathbf{L}\psi_k \right\} \\
&= \dim \left\{ \boldsymbol{y} \in \boldsymbol{Y} | \exists(\alpha_1, \alpha_2, \dots, \alpha_p) \quad \text{such that} \quad \boldsymbol{y} = \sum_{k=1}^{p-n} \alpha_k \mathbf{L}\psi_k + \sum_{k=1}^{n} \alpha_k \mathbf{L}\psi_k \right\} \\
&= \dim \left\{ \boldsymbol{y} \in \boldsymbol{Y} | \exists(\alpha_1, \alpha_2, \dots, \alpha_p) \quad \text{such that} \quad \boldsymbol{y} = \sum_{k=1}^{p-n} \alpha_k \mathbf{L}\psi_k + \mathbb{0} \right\} \\
&= p - n \\
&= \dim \boldsymbol{X} - \dim \mathcal{N}(\mathbf{L})
\end{aligned}$$

Note: This "proof" may be missing some necessary detail. ✏

N.2 Operators on Normed linear spaces

N.2.1 Operator norm

Definition N.4. [10] *Let $\mathcal{L}(\boldsymbol{X}, \boldsymbol{Y})$ be the space of linear operators over normed linear spaces $\boldsymbol{X}$ and $\boldsymbol{Y}$.* [11]

DEF

*The **operator norm** $|||\cdot|||$ is defined as*

$$|||\mathbf{A}||| \triangleq \sup_{\boldsymbol{x} \in \boldsymbol{X}} \{ \|\mathbf{A}\boldsymbol{x}\| \mid \|\boldsymbol{x}\| \le 1 \} \qquad \forall \mathbf{A} \in \mathcal{L}(\boldsymbol{X}, \boldsymbol{Y})$$

*The pair $(\mathcal{L}(\boldsymbol{X}, \boldsymbol{Y}), |||\cdot|||)$ is the **normed space of linear operators** on $(\boldsymbol{X}, \boldsymbol{Y})$.*

Proposition N.1 (next) shows that the functional defined in Definition N.4 (previous) is a *norm* (Definition K.1 page 271).

Proposition N.1. [12] *Let $(\mathcal{L}(\boldsymbol{X}, \boldsymbol{Y}), |||\cdot|||)$ be the normed space of linear operators over the normed linear spaces $\boldsymbol{X} \triangleq \big(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \|\cdot\|\big)$ and $\boldsymbol{Y} \triangleq \big(Y, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \|\cdot\|\big)$.*

PRP

*The functional $|||\cdot|||$ is a **norm** on $\mathcal{L}(\boldsymbol{X}, \boldsymbol{Y})$. In particular,*

1.	$|||\mathbf{A}|||$	≥ 0	$\forall \mathbf{A} \in \mathcal{L}(\boldsymbol{X}, \boldsymbol{Y})$	(NON-NEGATIVE)	*and*
2.	$|||\mathbf{A}|||$	$= 0 \iff \mathbf{A} \mathrel{\hat{=}} \mathbb{0}$	$\forall \mathbf{A} \in \mathcal{L}(\boldsymbol{X}, \boldsymbol{Y})$	(NONDEGENERATE)	*and*
3.	$|||\alpha\mathbf{A}|||$	$= \lvert\alpha\rvert \, |||\mathbf{A}|||$	$\forall \mathbf{A} \in \mathcal{L}(\boldsymbol{X}, \boldsymbol{Y}), \alpha \in \mathbb{F}$	(HOMOGENEOUS)	*and*
4.	$\big\|\!\big\|\mathbf{A} \mathbin{\dot{+}} \mathbf{B}\big\|\!\big\|$	$\le |||\mathbf{A}||| + |||\mathbf{B}|||$	$\forall \mathbf{A} \in \mathcal{L}(\boldsymbol{X}, \boldsymbol{Y})$	(SUBADDITIVE).	

*Moreover, $(\mathcal{L}(\boldsymbol{X}, \boldsymbol{Y}), |||\cdot|||)$ is a **normed linear space**.*

[10] Rudin (1991) page 92, Aliprantis and Burkinshaw (1998) page 225
[11] The operator norm notation $|||\cdot|||$ is introduced (as a Matrix norm) in Horn and Johnson (1990), page 290
[12] Rudin (1991) page 93

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

✎PROOF:

1. Proof that $|||\mathbf{A}||| > 0$ for $\mathbf{A} \neq \mathbb{0}$:

$$\begin{aligned} |||\mathbf{A}||| &\triangleq \sup_{x\in X} \{\|\mathbf{A}x\| \mid \|x\| \leq 1\} && \text{by definition of } |||\cdot||| \text{ (Definition N.4 page 308)} \\ &> 0 \end{aligned}$$

2. Proof that $|||\mathbf{A}||| = 0$ for $\mathbf{A} \mathrel{\hat{=}} \mathbb{0}$:

$$\begin{aligned} |||\mathbf{A}||| &\triangleq \sup_{x\in X} \{\|\mathbf{A}x\| \mid \|x\| \leq 1\} && \text{by definition of } |||\cdot||| \text{ (Definition N.4 page 308)} \\ &= \sup_{x\in X} \{\|\mathbb{0}x\| \mid \|x\| \leq 1\} \\ &= 0 \end{aligned}$$

3. Proof that $|||\alpha\mathbf{A}||| = |\alpha|\,|||\mathbf{A}|||$:

$$\begin{aligned} |||\alpha\mathbf{A}||| &\triangleq \sup_{x\in X} \{\|\alpha\mathbf{A}x\| \mid \|x\| \leq 1\} && \text{by definition of } |||\cdot||| \text{ (Definition N.4 page 308)} \\ &= \sup_{x\in X} \{|\alpha|\,\|\mathbf{A}x\| \mid \|x\| \leq 1\} && \text{by definition of } |||\cdot||| \text{ (Definition N.4 page 308)} \\ &= |\alpha| \sup_{x\in X} \{\|\mathbf{A}x\| \mid \|x\| \leq 1\} && \text{by definition of sup} \\ &= |\alpha|\;\; |||\mathbf{A}||| && \text{by definition of } |||\cdot||| \text{ (Definition N.4 page 308)} \end{aligned}$$

4. Proof that $\left|\left|\left|\mathbf{A} \mathbin{\dot{+}} \mathbf{B}\right|\right|\right| \leq |||\mathbf{A}||| + |||\mathbf{B}|||$:

$$\begin{aligned} \left|\left|\left|\mathbf{A} \mathbin{\dot{+}} \mathbf{B}\right|\right|\right| &\triangleq \sup_{x\in X} \left\{\left\|(\mathbf{A} \mathbin{\dot{+}} \mathbf{B})x\right\| \mid \|x\| \leq 1\right\} && \text{by definition of } |||\cdot||| \text{ (Definition N.4 page 308)} \\ &= \sup_{x\in X} \{\|\mathbf{A}x + \mathbf{B}x\| \mid \|x\| \leq 1\} \\ &\leq \sup_{x\in X} \{\|\mathbf{A}x\| + \|\mathbf{B}x\| \mid \|x\| \leq 1\} && \text{by definition of } |||\cdot||| \text{ (Definition N.4 page 308)} \\ &\leq \sup_{x\in X} \{\|\mathbf{A}x\| \mid \|x\| \leq 1\} + \sup_{x\in X} \{\|\mathbf{B}x\| \mid \|x\| \leq 1\} \\ &\triangleq |||\mathbf{A}||| + |||\mathbf{B}||| && \text{by definition of } |||\cdot||| \text{ (Definition N.4 page 308)} \end{aligned}$$

✏

Lemma N.1. *Let* $(\mathcal{L}(\boldsymbol{X}, \boldsymbol{Y}), |||\cdot|||)$ *be the normed space of linear operators over normed linear spaces* $\boldsymbol{X} \triangleq \big(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \|\cdot\|\big)$ *and* $\boldsymbol{Y} \triangleq \big(Y, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \|\cdot\|\big)$.

LEM

$$|||\mathbf{L}||| = \sup_{x} \{\|\mathbf{L}x\| \mid \|x\| = 1\} \qquad \forall x \in \mathcal{L}(\boldsymbol{X}, \boldsymbol{Y})$$

✎PROOF: [13]

1. Proof that $\sup_x \{\|\mathbf{L}x\| \mid \|x\| \leq 1\} \geq \sup_x \{\|\mathbf{L}x\| \mid \|x\| = 1\}$:

$$\sup_{x} \{\|\mathbf{L}x\| \mid \|x\| \leq 1\} \geq \sup_{x} \{\|\mathbf{L}x\| \mid \|x\| = 1\} \qquad \text{because } A \subseteq B \implies \sup A \leq \sup B$$

[13] email

Many many thanks to former NCTU Ph.D. student Chien Yao (Chinese: 姚建; PinYin: Yáo Jiàn) for his brilliant help with this proof. (If you are viewing this text as a pdf file, zoom in on the figure to the left to see text from Chien Yao's 2007 April 16 email.)

COPYRIGHT © 2019 DANIEL J. GREENHOE | CC BY-NC-ND

2. Let the subset $Y \subsetneq X$ be defined as

$$Y \triangleq \left\{ y \in X \,\middle|\, \begin{array}{ll} 1. & \|\mathbf{L}y\| = \sup_{x \in X} \{\|\mathbf{L}x\| \mid \|x\| \le 1\} \text{ and} \\ 2. & 0 < \|y\| \le 1 \end{array} \right\}$$

3. Proof that $\sup_x \{\|\mathbf{L}x\| \mid \|x\| \le 1\} \le \sup_x \{\|\mathbf{L}x\| \mid \|x\| = 1\}$:

$$\begin{aligned}
\sup_{x \in X} \{\|\mathbf{L}x\| \mid \|x\| \le 1\} &= \|\mathbf{L}y\| && \text{by definition of set } Y \\
&= \frac{\|y\|}{\|y\|} \|\mathbf{L}y\| \\
&= \|y\| \left\| \frac{1}{\|y\|} \mathbf{L}y \right\| && \text{by homogeneous property (page 271)} \\
&= \|y\| \left\| \mathbf{L} \frac{y}{\|y\|} \right\| && \text{by homogeneous property (page 304)} \\
&\le \|y\| \sup_{y \in Y} \left\{ \left\| \mathbf{L} \frac{y}{\|y\|} \right\| \right\} && \text{by definition of supremum} \\
&= \|y\| \sup_{y \in Y} \left\{ \left\| \mathbf{L} \frac{y}{\|y\|} \right\| \,\middle|\, \left\| \frac{y}{\|y\|} \right\| = 1 \right\} && \text{because } \left\| \frac{y}{\|y\|} \right\| = 1 \text{ for all } y \in Y \\
&\le \sup_{y \in Y} \left\{ \left\| \mathbf{L} \frac{y}{\|y\|} \right\| \,\middle|\, \left\| \frac{y}{\|y\|} \right\| = 1 \right\} && \text{because } 0 < \|y\| \le 1 \\
&\le \sup_{x \in X} \{\|\mathbf{L}x\| \mid \|x\| = 1\} && \text{because } \frac{y}{\|y\|} \in X \quad \forall y \in Y
\end{aligned}$$

4. By (1) and (3),

$$\sup_{x \in X} \{\|\mathbf{L}x\| \mid \|x\| \le 1\} = \sup_{x \in X} \{\|\mathbf{L}x\| \mid \|x\| = 1\}$$

✏

Proposition N.2. [14] *Let* $\mathbf{I}$ *be the identity operator in the normed space of linear operators* $(\mathcal{L}(\boldsymbol{X}, \boldsymbol{X}), |||\cdot|||)$.

PRP $|||\mathbf{I}||| = 1$

✎PROOF:

$$\begin{aligned}
|||\mathbf{I}||| &\triangleq \sup \{\|\mathbf{I}x\| \mid \|x\| \le 1\} && \text{by definition of } |||\cdot||| \text{ (Definition N.4 page 308)} \\
&= \sup \{\|x\| \mid \|x\| \le 1\} && \text{by definition of } \mathbf{I} \text{ (Definition N.2 page 303)} \\
&= 1
\end{aligned}$$

✏

Theorem N.6. [15] *Let* $(\mathcal{L}(\boldsymbol{X}, \boldsymbol{Y}), |||\cdot|||)$ *be the normed space of linear operators over normed linear spaces* $\boldsymbol{X}$ *and* $\boldsymbol{Y}$.

THM
$$\begin{aligned}
\|\mathbf{L}x\| &\le |||\mathbf{L}|||\ \|x\| && \forall \mathbf{L} \in \mathcal{L}(\boldsymbol{X}, \boldsymbol{Y}),\ x \in \boldsymbol{X} \\
|||\mathbf{KL}||| &\le |||\mathbf{K}|||\ |||\mathbf{L}||| && \forall \mathbf{K}, \mathbf{L} \in \mathcal{L}(\boldsymbol{X}, \boldsymbol{Y})
\end{aligned}$$

[14] Michel and Herget (1993) page 410
[15] Rudin (1991) page 103, Aliprantis and Burkinshaw (1998) page 225

COPYRIGHT © 2019 DANIEL J. GREENHOE

✎PROOF:

1. Proof that $\|\mathbf{L}\boldsymbol{x}\| \le |||\mathbf{L}|||\ \|\boldsymbol{x}\|$:

$$\begin{aligned}
\|\mathbf{L}\boldsymbol{x}\| &= \frac{\|\boldsymbol{x}\|}{\|\boldsymbol{x}\|}\|\mathbf{L}\boldsymbol{x}\| \\
&= \|\boldsymbol{x}\|\left\|\frac{1}{\|\boldsymbol{x}\|}\mathbf{L}\boldsymbol{x}\right\| && \text{by property of norms} \\
&= \|\boldsymbol{x}\|\left\|\mathbf{L}\frac{\boldsymbol{x}}{\|\boldsymbol{x}\|}\right\| && \text{by property of linear operators} \\
&\triangleq \|\boldsymbol{x}\|\ \|\mathbf{L}\boldsymbol{y}\| && \text{where } y \triangleq \frac{\boldsymbol{x}}{\|\boldsymbol{x}\|} \\
&\le \|\boldsymbol{x}\| \sup_{y} \|\mathbf{L}\boldsymbol{y}\| && \text{by definition of supremum} \\
&= \|\boldsymbol{x}\| \sup_{y} \{\|\mathbf{L}\boldsymbol{y}\| \mid \|\boldsymbol{y}\| = 1\} && \text{because } \|\boldsymbol{y}\| = \left\|\frac{\boldsymbol{x}}{\|\boldsymbol{x}\|}\right\| = \frac{\|\boldsymbol{x}\|}{\|\boldsymbol{x}\|} = 1 \\
&\triangleq \|\boldsymbol{x}\|\ \|\mathbf{L}\| && \text{by definition of operator norm}
\end{aligned}$$

2. Proof that $|||\mathbf{KL}||| \le |||\mathbf{K}|||\ |||\mathbf{L}|||$:

$$\begin{aligned}
|||\mathbf{KL}||| &\triangleq \sup_{x\in\boldsymbol{X}} \{\|(\mathbf{KL})\boldsymbol{x}\| \mid \|\boldsymbol{x}\| \le 1\} && \text{by Definition N.4 page 308 } (|||\cdot|||) \\
&= \sup_{x\in\boldsymbol{X}} \{\|\mathbf{K}(\mathbf{L}\boldsymbol{x})\| \mid \|\boldsymbol{x}\| \le 1\} \\
&\le \sup_{x\in\boldsymbol{X}} \{|||\mathbf{K}|||\ \|\mathbf{L}\boldsymbol{x}\| \mid \|\boldsymbol{x}\| \le 1\} && \text{by 1.} \\
&\le \sup_{x\in\boldsymbol{X}} \{|||\mathbf{K}|||\ |||\mathbf{L}|||\ \|\boldsymbol{x}\| \mid \|\boldsymbol{x}\| \le 1\} && \text{by 1.} \\
&= \sup_{x\in\boldsymbol{X}} \{|||\mathbf{K}|||\ |||\mathbf{L}|||\ 1 \mid \|\boldsymbol{x}\| \le 1\} && \text{by definition of sup} \\
&= |||\mathbf{K}|||\ |||\mathbf{L}||| && \text{by definition of sup}
\end{aligned}$$

✏

N.2.2 Bounded linear operators

Definition N.5. [16] *Let $(\mathcal{L}(\boldsymbol{X},\boldsymbol{Y}), |||\cdot|||)$ be a normed space of linear operators.*

DEF

*An operator $\mathbf{B}$ is **bounded** if $|||\mathbf{B}||| < \infty$.*
*The quantity $\mathcal{B}(\boldsymbol{X},\boldsymbol{Y})$ is the set of all **bounded linear operators** on $(\boldsymbol{X},\boldsymbol{Y})$ such that*
$$\mathcal{B}(\boldsymbol{X},\boldsymbol{Y}) \triangleq \{\mathbf{L} \in \mathcal{L}(\boldsymbol{X},\boldsymbol{Y}) \mid |||\mathbf{L}||| < \infty\}.$$

Theorem N.7. [17] *Let $(\mathcal{L}(\boldsymbol{X},\boldsymbol{Y}), |||\cdot|||)$ be the set of linear operators over normed linear spaces $\boldsymbol{X} \triangleq \big(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \|\cdot\|\big)$ and $\boldsymbol{Y} \triangleq \big(Y, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \|\cdot\|\big)$.*

THM

The following conditions are all EQUIVALENT*:*

1. $\mathbf{L}$ *is continuous at* A SINGLE POINT $\boldsymbol{x}_0 \in X$ $\quad \forall \mathbf{L}\in\mathcal{L}(\boldsymbol{X},\boldsymbol{Y}) \quad \Longleftrightarrow$
2. $\mathbf{L}$ *is* CONTINUOUS *(at every point $\boldsymbol{x} \in X$)* $\quad \forall \mathbf{L}\in\mathcal{L}(\boldsymbol{X},\boldsymbol{Y}) \quad \Longleftrightarrow$
3. $|||\mathbf{L}||| < \infty$ *($\mathbf{L}$ is* BOUNDED*)* $\quad \forall \mathbf{L}\in\mathcal{L}(\boldsymbol{X},\boldsymbol{Y}) \quad \Longleftrightarrow$
4. $\exists M \in \mathbb{R}$ such that $\|\mathbf{L}\boldsymbol{x}\| \le M\ \|\boldsymbol{x}\|$ $\quad \forall \mathbf{L}\in\mathcal{L}(\boldsymbol{X},\boldsymbol{Y}), \boldsymbol{x}\in X$

[16] Rudin (1991) pages 92–93
[17] Aliprantis and Burkinshaw (1998) page 227

COPYRIGHT © 2019 DANIEL J. GREENHOE

✎PROOF:

1. Proof that 1 $\Longrightarrow$ 2:

$$\begin{aligned}
\epsilon &> \left\|\mathbf{L}\boldsymbol{x} - \mathbf{L}\boldsymbol{x}_0\right\| && \text{by hypothesis 1}\\
&= \left\|\mathbf{L}(\boldsymbol{x} - \boldsymbol{x}_0)\right\| && \text{by linearity (Definition N.3 page 304)}\\
&= \left\|\mathbf{L}(\boldsymbol{x} + \boldsymbol{y} - \boldsymbol{x}_0 - \boldsymbol{y})\right\| && \\
&= \left\|\mathbf{L}(\boldsymbol{x} + \boldsymbol{y}) - \mathbf{L}(\boldsymbol{x}_0 + \boldsymbol{y})\right\| && \text{by linearity (Definition N.3 page 304)}\\
&\Longrightarrow \mathbf{L} \text{ is continuous at point } \boldsymbol{x} + \boldsymbol{y} && \\
&\Longrightarrow \mathbf{L} \text{ is continuous at every point in } X && \text{(hypothesis 2)}
\end{aligned}$$

2. Proof that 2 $\Longrightarrow$ 1: obvious:

3. Proof that 4 $\Longrightarrow$ 2:[18]

$$\begin{aligned}
|||\mathbf{L}\boldsymbol{x}||| \le M\,\|\boldsymbol{x}\| &\Longrightarrow |||\mathbf{L}(\boldsymbol{x} - \boldsymbol{y})||| \le M\,\|\boldsymbol{x} - \boldsymbol{y}\| && \text{by hypothesis 4}\\
&\Longrightarrow |||\mathbf{L}\boldsymbol{x} - \mathbf{L}\boldsymbol{y}||| \le M\,\|\boldsymbol{x} - \boldsymbol{y}\| && \text{by linearity of } \mathbf{L} \text{ (Definition N.3 page 304)}\\
&\Longrightarrow |||\mathbf{L}\boldsymbol{x} - \mathbf{L}\boldsymbol{y}||| \le \epsilon \text{ whenever } M\,\|\boldsymbol{x} - \boldsymbol{y}\| < \epsilon && \\
&\Longrightarrow |||\mathbf{L}\boldsymbol{x} - \mathbf{L}\boldsymbol{y}||| \le \epsilon \text{ whenever } \|\boldsymbol{x} - \boldsymbol{y}\| < \frac{\epsilon}{M} && \text{(hypothesis 2)}
\end{aligned}$$

4. Proof that 3 $\Longrightarrow$ 4:

$$\begin{aligned}
\|\mathbf{L}\boldsymbol{x}\| &\le \underbrace{|||\mathbf{L}|||}_{M}\ \|\boldsymbol{x}\| && \text{by Theorem N.6 page 310}\\
&= M\ \|\boldsymbol{x}\| && \text{where } M \triangleq |||\mathbf{L}||| < \infty \text{ (by hypothesis 1)}
\end{aligned}$$

5. Proof that 1 $\Longrightarrow$ 3:[19]

$$\begin{aligned}
|||\mathbf{L}||| = \infty &\Longrightarrow \{\|\mathbf{L}\boldsymbol{x}\| \mid \|\boldsymbol{x}\| \le 1\} = \infty\\
&\Longrightarrow \exists\,(\boldsymbol{x}_n) \quad \text{such that} \quad \left\|\boldsymbol{x}_n\right\| = 1 \text{ and } |||\mathbf{L}||| = \left\{\left\|\mathbf{L}\boldsymbol{x}_n\right\| \mid \left\|\boldsymbol{x}_n\right\| \le 1\right\} = \infty\\
&\Longrightarrow \left\|\boldsymbol{x}_n\right\| = 1 \text{ and } \infty = |||\mathbf{L}||| = \left\|\mathbf{L}\boldsymbol{x}_n\right\|\\
&\Longrightarrow \left\|\boldsymbol{x}_n\right\| = 1 \text{ and } \left\|\mathbf{L}\boldsymbol{x}_n\right\| \ge n\\
&\Longrightarrow \frac{1}{n}\left\|\boldsymbol{x}_n\right\| = \frac{1}{n} \text{ and } \frac{1}{n}\left\|\mathbf{L}\boldsymbol{x}_n\right\| \ge 1\\
&\Longrightarrow \left\|\frac{\boldsymbol{x}_n}{n}\right\| = \frac{1}{n} \text{ and } \left\|\mathbf{L}\frac{\boldsymbol{x}_n}{n}\right\| \ge 1\\
&\Longrightarrow \lim_{n\to\infty}\left\|\frac{\boldsymbol{x}_n}{n}\right\| = 0 \text{ and } \lim_{n\to\infty}\left\|\mathbf{L}\frac{\boldsymbol{x}_n}{n}\right\| \ge 1\\
&\Longrightarrow \mathbf{L} \text{ is not continuous at } 0
\end{aligned}$$

But by hypothesis, $\mathbf{L}$ *is* continuous. So the statement $|||\mathbf{L}||| = \infty$ must be *false* and thus $|||\mathbf{L}||| < \infty$ ($\mathbf{L}$ is *bounded*).

✏

[18] Bollobás (1999), page 29
[19] Aliprantis and Burkinshaw (1998), page 227

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

N.2.3 Adjoints on normed linear spaces

Definition N.6. *Let $\mathcal{B}(\boldsymbol{X}, \boldsymbol{Y})$ be the space of* BOUNDED LINEAR OPERATORS *on normed linear spaces $\boldsymbol{X}$ and $\boldsymbol{Y}$. Let $\boldsymbol{X}^*$ be the* TOPOLOGICAL DUAL SPACE *of $\boldsymbol{X}$.*

DEF

$\mathbf{B}^*$ *is the* ***adjoint*** *of an operator* $\mathbf{B} \in \mathcal{B}(\boldsymbol{X}, \boldsymbol{Y})$ *if*

$$\mathsf{f}(\mathbf{B}\boldsymbol{x}) = \left[\mathbf{B}^*\mathsf{f}\right](\boldsymbol{x}) \qquad \forall \mathsf{f} \in \boldsymbol{X}^*,\ \boldsymbol{x} \in \boldsymbol{X}$$

Theorem N.8. [20] *Let $\mathcal{B}(\boldsymbol{X}, \boldsymbol{Y})$ be the space of* BOUNDED LINEAR OPERATORS *on* NORMED LINEAR SPACES *$\boldsymbol{X}$ and $\boldsymbol{Y}$.*

THM

$$\begin{aligned}
(\mathbf{A} \mathbin{\mathring{+}} \mathbf{B})^* &= \mathbf{A}^* \mathbin{\mathring{+}} \mathbf{B}^* && \forall \mathbf{A},\mathbf{B} \in \mathcal{B}(\boldsymbol{X},\boldsymbol{Y}) \\
(\lambda\mathbf{A})^* &= \lambda\mathbf{A}^* && \forall \mathbf{A},\mathbf{B} \in \mathcal{B}(\boldsymbol{X},\boldsymbol{Y}) \\
(\mathbf{AB})^* &= \mathbf{B}^*\mathbf{A}^* && \forall \mathbf{A},\mathbf{B} \in \mathcal{B}(\boldsymbol{X},\boldsymbol{Y})
\end{aligned}$$

✎PROOF:

$$\begin{aligned}
\left[\mathbf{A} \mathbin{\mathring{+}} \mathbf{B}\right]^*\mathsf{f}(\boldsymbol{x}) &= \mathsf{f}\left(\left[\mathbf{A} \mathbin{\mathring{+}} \mathbf{B}\right]\boldsymbol{x}\right) && \text{by definition of adjoint} && \text{(Definition N.6 page 313)} \\
&= \mathsf{f}(\mathbf{A}\boldsymbol{x} + \mathbf{B}\boldsymbol{x}) && \text{by definition of linear operators} && \text{(Definition N.3 page 304)} \\
&= \mathsf{f}(\mathbf{A}\boldsymbol{x}) + \mathsf{f}(\mathbf{B}\boldsymbol{x}) && \text{by definition of } \textit{linear functional} \\
&= \mathbf{A}^*\mathsf{f}(\boldsymbol{x}) + \mathbf{B}^*\mathsf{f}(\boldsymbol{x}) && \text{by definition of } \textit{adjoint} && \text{(Definition N.6 page 313)} \\
&= \left[\mathbf{A}^* + \mathbf{B}^*\right]\mathsf{f}(\boldsymbol{x}) && \text{by definition of } \textit{linear functional}
\end{aligned}$$

$$\begin{aligned}
[\lambda\mathbf{A}]^*\mathsf{f}(\boldsymbol{x}) &= \mathsf{f}([\lambda\mathbf{A}]\boldsymbol{x}) && \text{by definition of } \textit{adjoint} && \text{(Definition N.6 page 313)} \\
&= \lambda\mathsf{f}(\mathbf{A}\boldsymbol{x}) && \text{by definition of } \textit{linear functional} \\
&= \left[\lambda\mathbf{A}^*\right]\mathsf{f}(\boldsymbol{x}) && \text{by definition of } \textit{adjoint} && \text{(Definition N.6 page 313)}
\end{aligned}$$

$$\begin{aligned}
[\mathbf{AB}]^*\mathsf{f}(\boldsymbol{x}) &= \mathsf{f}([\mathbf{AB}]\boldsymbol{x}) && \text{by definition of } \textit{adjoint} && \text{(Definition N.6 page 313)} \\
&= \mathsf{f}(\mathbf{A}[\mathbf{B}\boldsymbol{x}]) && \text{by definition of } \textit{linear operators} && \text{(Definition N.3 page 304)} \\
&= \left[\mathbf{A}^*\mathsf{f}\right](\mathbf{B}\boldsymbol{x}) && \text{by definition of } \textit{adjoint} && \text{(Definition N.6 page 313)} \\
&= \mathbf{B}^*\left[\mathbf{A}^*\mathsf{f}\right](\boldsymbol{x}) && \text{by definition of } \textit{adjoint} && \text{(Definition N.6 page 313)} \\
&= \left[\mathbf{B}^*\mathbf{A}^*\right]\mathsf{f}(\boldsymbol{x}) && \text{by definition of } \textit{adjoint} && \text{(Definition N.6 page 313)}
\end{aligned}$$

✏

Theorem N.9. [21] *Let $\mathcal{B}(\boldsymbol{X}, \boldsymbol{Y})$ be the space of* BOUNDED LINEAR OPERATORS *on normed linear spaces $\boldsymbol{X}$ and $\boldsymbol{Y}$. Let $\mathbf{B}^*$ be the adjoint of an operator $\mathbf{B}$.*

THM

$$|||\mathbf{B}||| = \left|\left|\left|\mathbf{B}^*\right|\right|\right| \qquad \forall \mathbf{B} \in \mathcal{B}(\boldsymbol{X}, \boldsymbol{Y})$$

✎PROOF:

$$\begin{aligned}
|||\mathbf{B}||| &\triangleq \sup\{\|\mathbf{B}\boldsymbol{x}\| \mid \|\boldsymbol{x}\| \le 1\} && \text{by Definition N.4 page 308} \\
&\stackrel{?}{=} \sup\left\{|\mathsf{g}(\mathbf{B}\boldsymbol{x};\boldsymbol{y}^*)| \mid \|\boldsymbol{x}\| \le 1,\ \left\|\boldsymbol{y}^*\right\| \le 1\right\} \\
&= \sup\left\{|\mathsf{f}(\boldsymbol{x};\mathbf{B}^*\boldsymbol{y}^*)| \mid \|\boldsymbol{x}\| \le 1,\ \left\|\boldsymbol{y}^*\right\| \le 1\right\} \\
&\triangleq \sup\left\{\left\|\mathbf{B}^*\boldsymbol{y}^*\right\| \mid \|\boldsymbol{x}\| \le 1,\ \left\|\boldsymbol{y}^*\right\| \le 1\right\} \\
&= \sup\left\{\left\|\mathbf{B}^*\boldsymbol{y}^*\right\| \mid \left\|\boldsymbol{y}^*\right\| \le 1\right\} \\
&\triangleq \left|\left|\left|\mathbf{B}^*\right|\right|\right| && \text{by Definition N.4 page 308}
\end{aligned}$$

✏

[20] Bollobás (1999), page 156
[21] Rudin (1991) page 98

COPYRIGHT © 2019 DANIEL J. GREENHOE

N.2.4 More properties



❝Beginning with the third year of studies, most of my mathematical work was really started in conversations with Mazur and Banach. And according to Banach some of my own contributions were characterized by a certain "strangeness" in the formulation of problems and in the outline of possible proofs. As he told me once some years later, he was surprised how often these "strange" approaches really worked.❞

Stanislaus M. Ulam (1909–1984), Polish mathematician [22]

Theorem N.10 (Mazur-Ulam theorem)**.** [23] *Let $\phi \in \mathcal{L}(\boldsymbol{X}, \boldsymbol{Y})$ be a function on normed linear spaces $(\boldsymbol{X}, \|\cdot\|_{\boldsymbol{X}})$ and $(\boldsymbol{Y}, \|\cdot\|_{\boldsymbol{Y}})$. Let $\mathbf{I} \in \mathcal{L}(\boldsymbol{X}, \boldsymbol{X})$ be the identity operator on $(\boldsymbol{X}, \|\cdot\|_{\boldsymbol{X}})$.*

THM

$$\left.\begin{array}{ll} 1. & \underbrace{\phi^{-1}\phi = \phi\phi^{-1} = \mathbf{I}}_{\textit{bijective}} \quad \textit{and} \\ 2. & \underbrace{\|\phi\boldsymbol{x} - \phi\boldsymbol{y}\|_{\boldsymbol{Y}} = \|\boldsymbol{x} - \boldsymbol{y}\|_{\boldsymbol{X}} \quad \forall \boldsymbol{x},\boldsymbol{y} \in \boldsymbol{X}}_{\textit{isometric}} \end{array}\right\} \Longrightarrow \underbrace{\phi([1-\lambda]\boldsymbol{x} + \lambda\boldsymbol{y}) = [1-\lambda]\phi\boldsymbol{x} + \lambda\phi\boldsymbol{y} \forall \lambda \in \mathbb{R}}_{\textit{affine}}$$

✎PROOF: Proof not yet complete.

1. Let ψ be the *reflection* of $\boldsymbol{z}$ in $\boldsymbol{X}$ such that $\psi\boldsymbol{x} = 2\boldsymbol{z} - \boldsymbol{x}$

   (a) $\|\psi\boldsymbol{x} - \boldsymbol{z}\| = \|\boldsymbol{x} - \boldsymbol{z}\|$

2. Let $\lambda \triangleq \sup_{\mathrm{g}} \{\|\mathrm{g}\boldsymbol{z} - \boldsymbol{z}\|\}$

3. Proof that $\mathrm{g} \in W \implies \mathrm{g}^{-1} \in W$:

   Let $\hat{\boldsymbol{x}} \triangleq \mathrm{g}^{-1}\boldsymbol{x}$ and $\hat{\boldsymbol{y}} \triangleq \mathrm{g}^{-1}\boldsymbol{y}$.

$$\begin{aligned} \left\|\mathrm{g}^{-1}\boldsymbol{x} - \mathrm{g}^{-1}\boldsymbol{y}\right\| &= \|\hat{\boldsymbol{x}} - \hat{\boldsymbol{y}}\| && \text{by definition of } \hat{\boldsymbol{x}} \text{ and } \hat{\boldsymbol{y}} \\ &= \|\mathrm{g}\hat{\boldsymbol{x}} - \mathrm{g}\hat{\boldsymbol{y}}\| && \text{by left hypothesis} \\ &= \left\|\mathrm{g}\mathrm{g}^{-1}\boldsymbol{x} - \mathrm{g}\mathrm{g}^{-1}\boldsymbol{y}\right\| && \text{by definition of } \hat{\boldsymbol{x}} \text{ and } \hat{\boldsymbol{y}} \\ &= \|\boldsymbol{x} - \boldsymbol{y}\| && \text{by definition of } \mathrm{g}^{-1} \end{aligned}$$

[22] quote: Ulam (1991), page 33
image: http://www-history.mcs.st-andrews.ac.uk/Biographies/Ulam.html

[23] Oikhberg and Rosenthal (2007), page 598, Väisälä (2003), page 634, Giles (2000), page 11, Dunford and Schwartz (1957), page 91, Mazur and Ulam (1932)

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

4. Proof that $\mathrm{g}z = z$:

$$\begin{aligned}
2\lambda &= 2\sup\{\|\mathrm{g}z - z\|\} && \text{by definition of } \lambda \text{ item (2)}\\
&\le 2\|\mathrm{g}z - z\| && \text{by definition of sup}\\
&= \|2z - 2\mathrm{g}z\|\\
&= \|\psi\mathrm{g}z - \mathrm{g}z\| && \text{by definition of } \psi \text{ item (1)}\\
&= \left\|\mathrm{g}^{-1}\psi\mathrm{g}z - \mathrm{g}^{-1}\mathrm{g}z\right\| && \text{by item (3)}\\
&= \left\|\mathrm{g}^{-1}\psi\mathrm{g}z - z\right\| && \text{by definition of } \mathrm{g}^{-1}\\
&= \left\|\psi\mathrm{g}^{-1}\psi\mathrm{g}z - z\right\|\\
&= \left\|\mathrm{g}^{*}z - z\right\|\\
&\le \lambda && \text{by definition of } \lambda \text{ item (2)}\\
&\implies 2\lambda \le \lambda\\
&\implies \lambda = 0\\
&\implies \mathrm{g}z = z
\end{aligned}$$

5. Proof that $\phi\left(\frac{1}{2}x + \frac{1}{2}y\right) = \frac{1}{2}\phi x + \frac{1}{2}\phi y$:

$$\begin{aligned}
\phi\left(\frac{1}{2}x + \frac{1}{2}y\right) &=\\
&= \frac{1}{2}\phi x + \frac{1}{2}\phi y
\end{aligned}$$

6. Proof that $\phi([1-\lambda]x + \lambda y) = [1-\lambda]\phi x + \lambda\phi y$:

$$\begin{aligned}
\phi([1-\lambda]x + \lambda y) &=\\
&= [1-\lambda]\phi x + \lambda\phi y
\end{aligned}$$

✍

Theorem N.11 (Neumann Expansion Theorem)**.** [24] *Let* $\mathbf{A} \in \boldsymbol{X}^{\boldsymbol{X}}$ *be an operator on a linear space* $\boldsymbol{X}$*. Let* $\mathbf{A}^0 \triangleq \mathbf{I}$*.*

THM

$$\left.\begin{array}{ll} 1. & \mathbf{A} \in \mathcal{B}(\boldsymbol{X}, \boldsymbol{X}) \quad \textit{(A is bounded)}\\ 2. & |||\mathbf{A}||| < 1 \end{array}\right\} \implies \left\{\begin{array}{lll} 1. & (\mathbf{I}-\mathbf{A})^{-1} & \textit{exists}\\ 2. & |||(\mathbf{I}-\mathbf{A})^{-1}||| \le & \dfrac{1}{1-|||\mathbf{A}|||}\\ 3. & (\mathbf{I}-\mathbf{A})^{-1} = & \displaystyle\sum_{n=0}^{\infty}\mathbf{A}^n\\ & \textit{with uniform convergence} & \end{array}\right.$$

N.3 Operators on Inner product spaces

N.3.1 General Results

Theorem N.12. [25] *Let* $\mathbf{A}, \mathbf{B} \in \mathcal{B}(\boldsymbol{X}, \boldsymbol{X})$ *be* BOUNDED LINEAR OPERATORS *on an inner product space* $\boldsymbol{X} \triangleq \left(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle\triangle\,|\,\triangledown\rangle\right)$.

THM

$$\begin{array}{lllllllll} \langle\mathbf{B}x\,|\,x\rangle & = & 0 & \forall x\in X & \iff & \mathbf{B}x & = & \mathbb{0} & \forall x\in X\\ \langle\mathbf{A}x\,|\,x\rangle & = & \langle\mathbf{B}x\,|\,x\rangle & \forall x\in X & \iff & \mathbf{A} & = & \mathbf{B} & \end{array}$$

[24] Michel and Herget (1993) page 415
[25] Rudin (1991) page 310 ⟨Theorem 12.7, Corollary⟩

Copyright © 2019 Daniel J. Greenhoe

✎PROOF:

1. Proof that $\langle \mathbf{B}\boldsymbol{x} \mid \boldsymbol{x}\rangle = 0 \implies \mathbf{B}\boldsymbol{x} = \mathbb{0}$:

$$\begin{aligned}
0 &= \langle \mathbf{B}(\boldsymbol{x}+\mathbf{B}\boldsymbol{x}) \mid (\boldsymbol{x}+\mathbf{B}\boldsymbol{x})\rangle + i\,\langle \mathbf{B}(\boldsymbol{x}+i\mathbf{B}\boldsymbol{x}) \mid (\boldsymbol{x}+i\mathbf{B}\boldsymbol{x})\rangle && \text{by left hypothesis}\\
&= \left\{\left\langle \mathbf{B}\boldsymbol{x}+\mathbf{B}^2\boldsymbol{x}) \mid \boldsymbol{x}+\mathbf{B}\boldsymbol{x}\right\rangle\right\} + i\left\{\left\langle \mathbf{B}\boldsymbol{x}+i\mathbf{B}^2\boldsymbol{x}) \mid \boldsymbol{x}+i\mathbf{B}\boldsymbol{x}\right\rangle\right\} && \text{by Definition N.3 page 304}\\
&= \left\{\langle \mathbf{B}\boldsymbol{x} \mid \boldsymbol{x}\rangle + \langle \mathbf{B}\boldsymbol{x} \mid \mathbf{B}\boldsymbol{x}\rangle + \left\langle \mathbf{B}^2\boldsymbol{x} \mid \boldsymbol{x}\right\rangle + \left\langle \mathbf{B}^2\boldsymbol{x} \mid \mathbf{B}\boldsymbol{x}\right\rangle\right\} && \text{by Definition J.1 page 255}\\
&\quad + i\left\{\langle \mathbf{B}\boldsymbol{x} \mid \boldsymbol{x}\rangle - i\,\langle \mathbf{B}\boldsymbol{x} \mid \mathbf{B}\boldsymbol{x}\rangle + i\left\langle \mathbf{B}^2\boldsymbol{x} \mid \boldsymbol{x}\right\rangle - i^2\left\langle \mathbf{B}^2\boldsymbol{x} \mid \mathbf{B}\boldsymbol{x}\right\rangle\right\}\\
&= \left\{0 + \langle \mathbf{B}\boldsymbol{x} \mid \mathbf{B}\boldsymbol{x}\rangle + \left\langle \mathbf{B}^2\boldsymbol{x} \mid \boldsymbol{x}\right\rangle + 0\right\} + i\left\{0 - i\,\langle \mathbf{B}\boldsymbol{x} \mid \mathbf{B}\boldsymbol{x}\rangle + i\left\langle \mathbf{B}^2\boldsymbol{x} \mid \boldsymbol{x}\right\rangle - i^2 0\right\} && \text{by left hypothesis}\\
&= \left\{\langle \mathbf{B}\boldsymbol{x} \mid \mathbf{B}\boldsymbol{x}\rangle + \left\langle \mathbf{B}^2\boldsymbol{x} \mid \boldsymbol{x}\right\rangle\right\} + \left\{\langle \mathbf{B}\boldsymbol{x} \mid \mathbf{B}\boldsymbol{x}\rangle - \left\langle \mathbf{B}^2\boldsymbol{x} \mid \boldsymbol{x}\right\rangle\right\}\\
&= 2\,\langle \mathbf{B}\boldsymbol{x} \mid \mathbf{B}\boldsymbol{x}\rangle\\
&= 2\,\|\mathbf{B}\boldsymbol{x}\|^2\\
&\implies \mathbf{B}\boldsymbol{x} = \mathbb{0} && \text{by Definition K.1 page 271}
\end{aligned}$$

2. Proof that $\langle \mathbf{B}\boldsymbol{x} \mid \boldsymbol{x}\rangle = 0 \impliedby \mathbf{B}\boldsymbol{x} = \mathbb{0}$: by property of inner products (Theorem J.1 page 255).

3. Proof that $\langle \mathbf{A}\boldsymbol{x} \mid \boldsymbol{x}\rangle = \langle \mathbf{B}\boldsymbol{x} \mid \boldsymbol{x}\rangle \implies \mathbf{A} \mathrel{\hat{=}} \mathbf{B}$:

$$\begin{aligned}
0 &= \langle \mathbf{A}\boldsymbol{x} \mid \boldsymbol{x}\rangle - \langle \mathbf{B}\boldsymbol{x} \mid \boldsymbol{x}\rangle && \text{by left hypothesis}\\
&= \langle \mathbf{A}\boldsymbol{x} - \mathbf{B}\boldsymbol{x} \mid \boldsymbol{x}\rangle && \text{by \textit{additivity} property of } \langle \triangle \mid \triangledown\rangle \text{ (Definition J.1 page 255)}\\
&= \langle (\mathbf{A}-\mathbf{B})\boldsymbol{x} \mid \boldsymbol{x}\rangle && \text{by definition of operator addition}\\
\implies (\mathbf{A}-\mathbf{B})\boldsymbol{x} &= \mathbb{0} && \text{by item 1}\\
\implies \mathbf{A} &= \mathbf{B} && \text{by definition of operator subtraction}
\end{aligned}$$

4. Proof that $\langle \mathbf{A}\boldsymbol{x} \mid \boldsymbol{x}\rangle = \langle \mathbf{B}\boldsymbol{x} \mid \boldsymbol{x}\rangle \impliedby \mathbf{A} \mathrel{\hat{=}} \mathbf{B}$:

$$\langle \mathbf{A}\boldsymbol{x} \mid \boldsymbol{x}\rangle = \langle \mathbf{B}\boldsymbol{x} \mid \boldsymbol{x}\rangle \qquad \text{by } \mathbf{A} \mathrel{\hat{=}} \mathbf{B} \text{ hypothesis}$$

✏

N.3.2 Operator adjoint

A fundamental concept of operators on inner product spaces is the *operator adjoint* (Proposition N.3 page 316). The adjoint of an operator is a kind of generalization of the conjugate of a complex number in that

- Both are *star-algebras* (Theorem N.13 page 317).
- Both support decomposition into "real" and "imaginary" parts (Theorem I.3 page 252).

Structurally, the operator adjoint provides a convenient symmetric relationship between the *range space* and *null space* of an operator (Theorem N.14 page 318).

Proposition N.3. [26] *Let $\mathcal{B}(\boldsymbol{H},\boldsymbol{H})$ be the space of* BOUNDED LINEAR OPERATORS *(Definition N.5 page 311) on a* HILBERT SPACE $\boldsymbol{H}$.

PRP *An operator $\mathbf{B}^*$ is the **adjoint** of $\mathbf{B} \in \mathcal{B}(\boldsymbol{H},\boldsymbol{H})$ if*
$$\langle \mathbf{B}\boldsymbol{x} \mid \boldsymbol{y}\rangle = \langle \boldsymbol{x} \mid \mathbf{B}^*\boldsymbol{y}\rangle \qquad \forall \boldsymbol{x},\boldsymbol{y} \in \boldsymbol{H}.$$

[26] Michel and Herget (1993) page 220, Rudin (1991) page 311, Giles (2000), page 182, von Neumann (1929) page 49, Stone (1932) page 41

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

✎PROOF:

1. For fixed y, $f(x) \triangleq \langle x \mid y\rangle$ is a *functional* in $\mathbb{F}^X$.
2. $\mathbf{B}^*$ is the *adjoint* of $\mathbf{B}$ because

$$\begin{aligned}\langle \mathbf{B}x \mid y\rangle &\triangleq f(\mathbf{B}x)\\ &\triangleq \mathbf{B}^* f(x) && \text{by definition of } \textit{operator adjoint} && \text{(Definition N.6 page 313)}\\ &= \langle x \mid \mathbf{B}^* y\rangle\end{aligned}$$

✎

Example N.2.

EX

In matrix algebra ("linear algebra")

- The inner product operation $\langle x \mid y\rangle$ is represented by $y^H x$.
- The linear operator is represented as a matrix A.
- The operation of A on a vector x is represented as Ax.
- The adjoint of matrix A is the Hermitian matrix A^H.

✎PROOF:

$$\langle Ax \mid y\rangle \triangleq y^H Ax = [(Ax)^H y]^H = [x^H A^H y]^H = (A^H y)^H x \triangleq \left\langle x \mid A^H y\right\rangle$$

✎

Structures that satisfy the four conditions of the next theorem are known as **-algebras* ("*star-algebras*" (Definition I.3 page 250). Other structures which are *-algebras include the *field of complex numbers* $\mathbb{C}$ and any *ring of complex square* $n \times n$ *matrices*.[27]

Theorem N.13 (operator star-algebra). [28] *Let* H *be a* HILBERT SPACE *with operators* $\mathbf{A}, \mathbf{B} \in \mathcal{B}(H, H)$ *and with adjoints* $\mathbf{A}^*, \mathbf{B}^* \in \mathcal{B}(H, H)$. *Let* $\bar{\alpha}$ *be the complex conjugate of some* $\alpha \in \mathbb{C}$.

THM

The pair $(H, *)$ *is a* $*$-ALGEBRA (STAR-ALGEBRA). *In particular,*

1.	$(\mathbf{A} \mathbin{\mathring{+}} \mathbf{B})^*$	$=$	$\mathbf{A}^* + \mathbf{B}^*$	$\forall \mathbf{A},\mathbf{B} \in H$	(DISTRIBUTIVE)	*and*
2.	$(\alpha\mathbf{A})^*$	$=$	$\bar{\alpha}\mathbf{A}^*$	$\forall \mathbf{A},\mathbf{B} \in H$	(CONJUGATE LINEAR)	*and*
3.	$(\mathbf{AB})^*$	$=$	$\mathbf{B}^*\mathbf{A}^*$	$\forall \mathbf{A},\mathbf{B} \in H$	(ANTIAUTOMORPHIC)	*and*
4.	$\mathbf{A}^{**}$	$=$	$\mathbf{A}$	$\forall \mathbf{A},\mathbf{B} \in H$	(INVOLUTARY)	

✎PROOF:

$$\begin{aligned}\left\langle x \mid (\mathbf{A} \mathbin{\mathring{+}} \mathbf{B})^* y\right\rangle &= \left\langle (\mathbf{A} \mathbin{\mathring{+}} \mathbf{B})x \mid y\right\rangle && \text{by definition of adjoint} && \text{(Proposition N.3 page 316)}\\ &= \langle \mathbf{A}x \mid y\rangle + \langle \mathbf{B}x \mid y\rangle && \text{by definition of inner product} && \text{(Definition J.1 page 255)}\\ &= \langle x \mid \mathbf{A}^* y\rangle + \langle x \mid \mathbf{B}^* y\rangle && \text{by definition of operator addition}\\ &= \langle x \mid \mathbf{A}^* y + \mathbf{B}^* y\rangle && \text{by definition of inner product} && \text{(Definition J.1 page 255)}\\ &= \langle x \mid (\mathbf{A}^* + \mathbf{B}^*) y\rangle && \text{by definition of operator addition}\end{aligned}$$

$$\begin{aligned}\langle x \mid (\alpha\mathbf{A})^* y\rangle &= \langle (\alpha\mathbf{A})x \mid y\rangle && \text{by definition of adjoint} && \text{(Proposition N.3 page 316)}\\ &= \langle \alpha(\mathbf{A}x) \mid y\rangle && \text{by definition of scalar multiplication}\\ &= \alpha \langle \mathbf{A}x \mid y\rangle && \text{by definition of inner product} && \text{(Definition J.1 page 255)}\end{aligned}$$

[27] Sakai (1998) page 1

[28] Halmos (1998), pages 39–40, Rudin (1991) page 311

COPYRIGHT © 2019 DANIEL J. GREENHOE | CC BY-NC-ND

$$
\begin{aligned}
&= \alpha\,\langle \boldsymbol{x} \mid \mathbf{A}^*\boldsymbol{y}\rangle && \text{by definition of adjoint} && \text{(Proposition N.3 page 316)}\\
&= \langle \boldsymbol{x} \mid \alpha^*\mathbf{A}^*\boldsymbol{y}\rangle && \text{by definition of inner product} && \text{(Definition J.1 page 255)}\\
\\
\langle \boldsymbol{x} \mid (\mathbf{AB})^*\boldsymbol{y}\rangle &= \langle (\mathbf{AB})\boldsymbol{x} \mid \boldsymbol{y}\rangle && \text{by definition of adjoint} && \text{(Proposition N.3 page 316)}\\
&= \langle \mathbf{A}(\mathbf{B}\boldsymbol{x}) \mid \boldsymbol{y}\rangle && \text{by definition of operator multiplication} && \\
&= \langle (\mathbf{B}\boldsymbol{x}) \mid \mathbf{A}^*\boldsymbol{y}\rangle && \text{by definition of adjoint} && \text{(Proposition N.3 page 316)}\\
&= \langle \boldsymbol{x} \mid \mathbf{B}^*\mathbf{A}^*\boldsymbol{y}\rangle && \text{by definition of adjoint} && \text{(Proposition N.3 page 316)}\\
\\
\langle \boldsymbol{x} \mid \mathbf{A}^{**}\boldsymbol{y}\rangle &= \langle \mathbf{A}^*\boldsymbol{x} \mid \boldsymbol{y}\rangle && \text{by definition of adjoint} && \text{(Proposition N.3 page 316)}\\
&= \langle \boldsymbol{y} \mid \mathbf{A}^*\boldsymbol{x}\rangle^* && \text{by definition of inner product} && \text{(Definition J.1 page 255)}\\
&= \langle \mathbf{A}\boldsymbol{y} \mid \boldsymbol{x}\rangle^* && \text{by definition of adjoint} && \text{(Proposition N.3 page 316)}\\
&= \langle \boldsymbol{x} \mid \mathbf{A}\boldsymbol{y}\rangle && \text{by definition of inner product} && \text{(Definition J.1 page 255)}
\end{aligned}
$$

✎

Theorem N.14. [29] *Let $\boldsymbol{Y}^{\boldsymbol{X}}$ be the set of all operators from a linear space $\boldsymbol{X}$ to a linear space $\boldsymbol{Y}$. Let $\mathcal{N}(\mathbf{L})$ be the* NULL SPACE *of an operator $\mathbf{L}$ in $\boldsymbol{Y}^{\boldsymbol{X}}$ and $\mathcal{I}(\mathbf{L})$ the* IMAGE SET *of $\mathbf{L}$ in $\boldsymbol{Y}^{\boldsymbol{X}}$.*

THM

$$\mathcal{N}(\mathbf{A}) = \mathcal{I}(\mathbf{A}^*)^\perp$$
$$\mathcal{N}(\mathbf{A}^*) = \mathcal{I}(\mathbf{A})^\perp$$

✎ PROOF:

$$
\begin{aligned}
\mathcal{I}(\mathbf{A}^*)^\perp &= \left\{y \in \boldsymbol{H} \mid \langle \boldsymbol{y} \mid \boldsymbol{u}\rangle = 0 \quad \forall u \in \mathcal{I}(\mathbf{A}^*)\right\}\\
&= \left\{y \in \boldsymbol{H} \mid \langle \boldsymbol{y} \mid \mathbf{A}^*\boldsymbol{x}\rangle = 0 \quad \forall \boldsymbol{x} \in \boldsymbol{H}\right\}\\
&= \{y \in \boldsymbol{H} \mid \langle \mathbf{A}\boldsymbol{y} \mid \boldsymbol{x}\rangle = 0 \quad \forall \boldsymbol{x} \in \boldsymbol{H}\} && \text{by definition of } \mathbf{A}^* && \text{(Proposition N.3 page 316)}\\
&= \{y \in \boldsymbol{H} \mid \mathbf{A}\boldsymbol{y} = 0\}\\
&= \mathcal{N}(\mathbf{A}) && \text{by definition of } \mathcal{N}(\mathbf{A})
\end{aligned}
$$

$$
\begin{aligned}
\mathcal{I}(\mathbf{A})^\perp &= \{y \in \boldsymbol{H} \mid \langle \boldsymbol{y} \mid \boldsymbol{u}\rangle = 0 \quad \forall u \in \mathcal{I}(\mathbf{A})\}\\
&= \{y \in \boldsymbol{H} \mid \langle \boldsymbol{y} \mid \mathbf{A}\boldsymbol{x}\rangle = 0 \quad \forall \boldsymbol{x} \in \boldsymbol{H}\} && \text{by definition of } \mathcal{I}\\
&= \left\{y \in \boldsymbol{H} \mid \langle \mathbf{A}^*\boldsymbol{y} \mid \boldsymbol{x}\rangle = 0 \quad \forall \boldsymbol{x} \in \boldsymbol{H}\right\} && \text{by definition of } \mathbf{A}^* && \text{(Proposition N.3 page 316)}\\
&= \left\{y \in \boldsymbol{H} \mid \mathbf{A}^*\boldsymbol{y} = 0\right\}\\
&= \mathcal{N}(\mathbf{A}^*) && \text{by definition of } \mathcal{N}(\mathbf{A})
\end{aligned}
$$

✎

N.4 Special Classes of Operators

N.4.1 Projection operators

Definition N.7. [30] *Let $\mathcal{B}(\boldsymbol{X}, \boldsymbol{Y})$ be the space of* BOUNDED LINEAR OPERATORS *on normed linear spaces $\boldsymbol{X}$ and $\boldsymbol{Y}$. Let $\mathbf{P}$ be a bounded linear operator in $\mathcal{B}(\boldsymbol{X}, \boldsymbol{Y})$.*

[29] Rudin (1991) page 312

[30] Rudin (1991) page 133 ⟨5.15 Projections⟩, Kubrusly (2001) page 70, Bachman and Narici (1966) page 6, Halmos (1958) page 73 ⟨§41. Projections⟩

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

DEF $\mathbf{P}$ *is a* ***projection*** *operator if* $\mathbf{P}^2 = \mathbf{P}$.

Theorem N.15. [31] *Let* $\mathcal{B}(\boldsymbol{X}, \boldsymbol{Y})$ *be the space of* BOUNDED LINEAR OPERATORS *on normed linear spaces* $\boldsymbol{X}$ *and* $\boldsymbol{Y}$. *Let* $\mathbf{P}$ *be a bounded linear operator in* $\mathcal{B}(\boldsymbol{X}, \boldsymbol{Y})$ *with* NULL SPACE $\mathcal{N}(\mathbf{P})$ *and* IMAGE SET $\mathcal{I}(\mathbf{P})$.

THM

$$\left.\begin{array}{llll} 1. & \mathbf{P}^2 = \mathbf{P} & (\mathbf{P} \text{ is a projection operator}) & \text{and} \\ 2. & \boldsymbol{\Omega} = \boldsymbol{X} \dotplus \boldsymbol{Y} & (\boldsymbol{Y} \text{ compliments } \boldsymbol{X} \text{ in } \boldsymbol{\Omega}) & \text{and} \\ 3. & \mathbf{P}\boldsymbol{\Omega} = \boldsymbol{X} & (\mathbf{P} \text{ projects onto } \boldsymbol{X}) & \end{array}\right\} \Longrightarrow \left\{\begin{array}{llll} 1. & \mathcal{I}(\mathbf{P}) = \boldsymbol{X} & & \text{and} \\ 2. & \mathcal{N}(\mathbf{P}) = \boldsymbol{Y} & & \text{and} \\ 3. & \boldsymbol{\Omega} = \mathcal{I}(\mathbf{P}) \dotplus \mathcal{N}(\mathbf{P}) & & \end{array}\right.$$

✎PROOF:

$$\begin{aligned} \mathcal{I}(\mathbf{P}) &= \mathbf{P}\boldsymbol{\Omega} \\ &= \mathbf{P}(\boldsymbol{\Omega}_1 + \boldsymbol{\Omega}_2) \\ &= \mathbf{P}\boldsymbol{\Omega}_1 + \mathbf{P}\boldsymbol{\Omega}_2 \\ &= \boldsymbol{\Omega}_1 + \{\mathbb{0}\} \\ &= \boldsymbol{\Omega}_1 \end{aligned}$$

$$\begin{aligned} \mathcal{N}(\mathbf{P}) &= \{\boldsymbol{x} \in \boldsymbol{\Omega} | \mathbf{P}\boldsymbol{x} = \mathbb{0}\} \\ &= \left\{\boldsymbol{x} \in (\boldsymbol{\Omega}_1 + \boldsymbol{\Omega}_2) | \mathbf{P}\boldsymbol{x} = \mathbb{0}\right\} \\ &= \left\{\boldsymbol{x} \in \boldsymbol{\Omega}_1 | \mathbf{P}\boldsymbol{x} = \mathbb{0}\right\} + \left\{\boldsymbol{x} \in \boldsymbol{\Omega}_2 | \mathbf{P}\boldsymbol{x} = \mathbb{0}\right\} \\ &= \{\mathbb{0}\} + \boldsymbol{\Omega}_2 \\ &= \boldsymbol{\Omega}_2 \end{aligned}$$

⇨

Theorem N.16. [32] *Let* $\mathcal{B}(\boldsymbol{X}, \boldsymbol{Y})$ *be the space of* BOUNDED LINEAR OPERATORS *on normed linear spaces* $\boldsymbol{X}$ *and* $\boldsymbol{Y}$. *Let* $\mathbf{P}$ *be a bounded linear operator in* $\mathcal{B}(\boldsymbol{X}, \boldsymbol{Y})$.

THM

$$\underbrace{\mathbf{P}^2 = \mathbf{P}}_{\mathbf{P} \text{ is a projection operator}} \quad \Longleftrightarrow \quad \underbrace{(\mathbf{I} - \mathbf{P})^2 = (\mathbf{I} - \mathbf{P})}_{(\mathbf{I} - \mathbf{P}) \text{ is a projection operator}}$$

✎PROOF:

Proof that $\mathbf{P}^2 = \mathbf{P} \implies (\mathbf{I} - \mathbf{P})^2 = (\mathbf{I} - \mathbf{P})$:

$$\begin{aligned} (\mathbf{I} - \mathbf{P})^2 &= (\mathbf{I} - \mathbf{P})(\mathbf{I} - \mathbf{P}) \\ &= \mathbf{I}(\mathbf{I} - \mathbf{P}) + (-\mathbf{P})(\mathbf{I} - \mathbf{P}) \\ &= \mathbf{I} - \mathbf{P} - \mathbf{P}\mathbf{I} + \mathbf{P}^2 \\ &= \mathbf{I} - \mathbf{P} - \mathbf{P} + \mathbf{P} && \text{by left hypothesis} \\ &= \mathbf{I} - \mathbf{P} \end{aligned}$$

Proof that $\mathbf{P}^2 = \mathbf{P} \impliedby (\mathbf{I} - \mathbf{P})^2 = (\mathbf{I} - \mathbf{P})$:

$$\begin{aligned} \mathbf{P}^2 &= \underbrace{\mathbf{I} - \mathbf{P} - \mathbf{P} + \mathbf{P}^2}_{(\mathbf{I} - \mathbf{P})^2} - (\mathbf{I} - \mathbf{P} - \mathbf{P}) \\ &= (\mathbf{I} - \mathbf{P})^2 - (\mathbf{I} - \mathbf{P} - \mathbf{P}) \\ &= (\mathbf{I} - \mathbf{P}) - (\mathbf{I} - \mathbf{P} - \mathbf{P}) && \text{by right hypothesis} \\ &= \mathbf{P} \end{aligned}$$

⇨

[31] Michel and Herget (1993) pages 120–121
[32] Michel and Herget (1993) page 121

COPYRIGHT © 2019 DANIEL J. GREENHOE
CC BY-NC-ND

Theorem N.17. [33] *Let $\boldsymbol{H}$ be a* HILBERT SPACE *and* $\mathbf{P}$ *an operator in* $\boldsymbol{H}^{\boldsymbol{H}}$ *with adjoint* $\mathbf{P}^*$, NULL SPACE $\mathcal{N}(\mathbf{P})$, *and* IMAGE SET $\mathcal{I}(\mathbf{P})$.[34]

THM

If $\mathbf{P}$ *is a* PROJECTION OPERATOR, *then the following are equivalent:*

1.	$\mathbf{P}^* = \mathbf{P}$		(**P** *is* SELF-ADJOINT)	$\Longleftrightarrow$
2.	$\mathbf{P}^*\mathbf{P} = \mathbf{P}\mathbf{P}^*$		(**P** *is* NORMAL)	$\Longleftrightarrow$
3.	$\mathcal{I}(\mathbf{P}) = \mathcal{N}(\mathbf{P})^\perp$			$\Longleftrightarrow$
4.	$\langle \mathbf{P}x \mid x\rangle = \|\mathbf{P}x\|^2$	$\forall x \in X$		

✎PROOF: This proof is incomplete at this time.

Proof that (1) $\Longrightarrow$ (2):

$$\begin{aligned} \mathbf{P}^*\mathbf{P} &= \mathbf{P}^{**}\mathbf{P}^* && \text{by (1)} \\ &= \mathbf{P}\mathbf{P}^* && \text{by Theorem N.13 page 317} \end{aligned}$$

Proof that (1) $\Longrightarrow$ (3):

$$\begin{aligned} \mathcal{I}(\mathbf{P}) &= \mathcal{N}(\mathbf{P}^*)^\perp && \text{by Theorem N.14 page 318} \\ &= \mathcal{N}(\mathbf{P})^\perp && \text{by (1)} \end{aligned}$$

Proof that (3) $\Longrightarrow$ (4):

Proof that (4) $\Longrightarrow$ (1):

✏

N.4.2 Self Adjoint Operators

Definition N.8. [35] *Let* $\mathbf{B} \in \mathcal{B}(\boldsymbol{H}, \boldsymbol{H})$ *be a* BOUNDED *operator with adjoint* $\mathbf{B}^*$ *on a* HILBERT SPACE $\boldsymbol{H}$.

DEF

The operator $\mathbf{B}$ *is said to be* ***self-adjoint*** *or* ***hermitian*** *if* $\mathbf{B} \mathrel{\hat{=}} \mathbf{B}^*$.

Example N.3 (Autocorrelation operator). Let $\mathsf{x}(t)$ be a random process with autocorrelation

$$\mathsf{R}_{\mathsf{xx}}(t,u) \triangleq \underbrace{\mathsf{E}[\mathsf{x}(t)\mathsf{x}^*(u)]}_{\text{expectation}}.$$

Let an autocorrelation operator $\mathbf{R}$ be defined as $[\mathbf{R}\mathsf{f}](t) \triangleq \int_{\mathbb{R}} \underbrace{\mathsf{R}_{\mathsf{xx}}(t,u)}_{\text{kernel}}\mathsf{f}(u)\,\mathrm{d}u$.

EX

$\mathbf{R} = \mathbf{R}^*$ (The auto-correlation operator $\mathbf{R}$ is *self-adjoint*)

[33] Rudin (1991) page 314

[34] *null space*: Definition ?? page ??
image set: Definition ?? page ??

[35] Historical works regarding self-adjoint operators: von Neumann (1929), page 49, "linearer Operator R selbstadjungiert oder Hermitesch", Stone (1932), page 50 ⟨"self-adjoint transformations"⟩

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

Theorem N.18. [36] *Let* $\mathbf{S} : \boldsymbol{H} \to \boldsymbol{H}$ *be an operator over a* HILBERT SPACE $\boldsymbol{H}$ *with eigenvalues* $\{\lambda_n\}$ *and eigenfunctions* $\{\psi_n\}$ *such that* $\mathbf{S}\psi_n = \lambda_n\psi_n$ *and let* $\|\boldsymbol{x}\| \triangleq \sqrt{\langle \boldsymbol{x} \mid \boldsymbol{x}\rangle}$.

THM

$$\left\{\underbrace{\mathbf{S} = \mathbf{S}^*}_{\mathbf{S}\text{ is self adjoint}}\right\} \implies \left\{\begin{array}{lll} 1. & \langle \mathbf{S}\boldsymbol{x} \mid \boldsymbol{x}\rangle \in \mathbb{R} & \textit{(the hermitian quadratic form of } \mathbf{S} \textit{ is REAL-VALUED)} \\ 2. & \lambda_n \in \mathbb{R} & \textit{(eigenvalues of } \mathbf{S} \textit{ are REAL-VALUED)} \\ 3. & \lambda_n \neq \lambda_m \implies \langle \psi_n \mid \psi_m\rangle = 0 & \textit{(eigenvectors are ORTHOGONAL)} \end{array}\right\}$$

✎PROOF:

1. Proof that $\mathbf{S} = \mathbf{S}^* \implies \langle \mathbf{S}\boldsymbol{x} \mid \boldsymbol{x}\rangle \in \mathbb{R}$:

$$\begin{aligned} \langle \boldsymbol{x} \mid \mathbf{S}\boldsymbol{x}\rangle &= \langle \mathbf{S}\boldsymbol{x} \mid \boldsymbol{x}\rangle && \text{by left hypothesis} \\ &= \langle \boldsymbol{x} \mid \mathbf{S}\boldsymbol{x}\rangle^* && \text{by definition of } \langle \triangle \mid \triangledown\rangle \text{ Definition J.1 page 255} \end{aligned}$$

2. Proof that $\mathbf{S} = \mathbf{S}^* \implies \lambda_n \in \mathbb{R}$:

$$\begin{aligned} \lambda_n \|\psi_n\|^2 &= \lambda_n \langle \psi_n \mid \psi_n\rangle && \text{by definition} \\ &= \langle \lambda_n\psi_n \mid \psi_n\rangle && \text{by definition of } \langle \triangle \mid \triangledown\rangle \text{ Definition J.1 page 255} \\ &= \langle \mathbf{S}\psi_n \mid \psi_n\rangle && \text{by definition of eigenpairs} \\ &= \langle \psi_n \mid \mathbf{S}\psi_n\rangle && \text{by left hypothesis} \\ &= \langle \psi_n \mid \lambda_n\psi_n\rangle && \text{by definition of eigenpairs} \\ &= \lambda_n^* \langle \psi_n \mid \psi_n\rangle && \text{by definition of } \langle \triangle \mid \triangledown\rangle \text{ Definition J.1 page 255} \\ &= \lambda_n^* \|\psi_n\|^2 && \text{by definition} \end{aligned}$$

3. Proof that $\mathbf{S} = \mathbf{S}^* \implies [\lambda_n \neq \lambda_m \implies \langle \psi_n \mid \psi_m\rangle = 0]$:

$$\begin{aligned} \lambda_n \langle \psi_n \mid \psi_m\rangle &= \langle \lambda_n\psi_n \mid \psi_m\rangle && \text{by definition of } \langle \triangle \mid \triangledown\rangle \text{ Definition J.1 page 255} \\ &= \langle \mathbf{S}\psi_n \mid \psi_m\rangle && \text{by definition of eigenpairs} \\ &= \langle \psi_n \mid \mathbf{S}\psi_m\rangle && \text{by left hypothesis} \\ &= \langle \psi_n \mid \lambda_m\psi_m\rangle && \text{by definition of eigenpairs} \\ &= \lambda_m^* \langle \psi_n \mid \psi_m\rangle && \text{by definition of } \langle \triangle \mid \triangledown\rangle \text{ Definition J.1 page 255} \\ &= \lambda_m \langle \psi_n \mid \psi_m\rangle && \text{because } \lambda_m \text{ is real} \end{aligned}$$

This implies for $\lambda_n \neq \lambda_m \neq 0$, $\langle \psi_n \mid \psi_m\rangle = 0$.

✏

N.4.3 Normal Operators

Definition N.9. [37] *Let* $\mathcal{B}(\boldsymbol{X}, \boldsymbol{Y})$ *be the space of* BOUNDED LINEAR OPERATORS *on normed linear spaces* $\boldsymbol{X}$ *and* $\boldsymbol{Y}$. *Let* $\mathbf{N}^*$ *be the adjoint of an operator* $\mathbf{N} \in \mathcal{B}(\boldsymbol{X}, \boldsymbol{Y})$.

DEF

$$\mathbf{N} \textit{ is } \textbf{\textit{normal}} \textit{ if } \quad \mathbf{N}^*\mathbf{N} = \mathbf{N}\mathbf{N}^*.$$

[36] Lax (2002), pages 315–316, Keener (1988), pages 114–119, Bachman and Narici (1966) page 24 ⟨Theorem 2.1⟩,
Bertero and Boccacci (1998) page 225 ⟨§"9.2 SVD of a matrix …If all eigenvectors are normalized…"⟩

[37] Rudin (1991) page 312, Michel and Herget (1993) page 431, Dieudonné (1969), page 167, Frobenius (1878), Frobenius (1968), page 391

COPYRIGHT © 2019 DANIEL J. GREENHOE

Theorem N.19. [38] *Let $\mathcal{B}(\boldsymbol{H},\boldsymbol{H})$ be the space of* BOUNDED LINEAR OPERATORS *on a* HILBERT SPACE $\boldsymbol{H}$. *Let $\mathcal{N}(\mathbf{N})$ be the* NULL SPACE *of an operator $\mathbf{N}$ in $\mathcal{B}(\boldsymbol{H},\boldsymbol{H})$ and $\mathcal{I}(\mathbf{N})$ the* IMAGE SET *of $\mathbf{N}$ in $\mathcal{B}(\boldsymbol{H},\boldsymbol{H})$.*

THM
$$\underbrace{\mathbf{N}^*\mathbf{N} = \mathbf{N}\mathbf{N}^*}_{\mathbf{N}\text{ is normal}} \quad \Longleftrightarrow \quad \left\|\mathbf{N}^*\boldsymbol{x}\right\| = \|\mathbf{N}\boldsymbol{x}\| \qquad \forall \boldsymbol{x} \in \boldsymbol{H}$$

✎PROOF:

1. Proof that $\mathbf{N}^*\mathbf{N} = \mathbf{N}\mathbf{N}^* \implies \left\|\mathbf{N}^*\boldsymbol{x}\right\| = \|\mathbf{N}\boldsymbol{x}\|$:

$$\begin{aligned}
\|\mathbf{N}\boldsymbol{x}\|^2 &= \langle \mathbf{N}\boldsymbol{x} \mid \mathbf{N}\boldsymbol{x}\rangle && \text{by definition}\\
&= \langle \boldsymbol{x} \mid \mathbf{N}^*\mathbf{N}\boldsymbol{x}\rangle && \text{by Proposition N.3 page 316 (definition of } \mathbf{N}^*)\\
&= \langle \boldsymbol{x} \mid \mathbf{N}\mathbf{N}^*\boldsymbol{x}\rangle && \text{by left hypothesis } (\mathbf{N} \text{ is normal})\\
&= \langle \mathbf{N}\boldsymbol{x} \mid \mathbf{N}^*\boldsymbol{x}\rangle && \text{by Proposition N.3 page 316 (definition of } \mathbf{N}^*)\\
&= \left\|\mathbf{N}^*\boldsymbol{x}\right\|^2 && \text{by definition}
\end{aligned}$$

2. Proof that $\mathbf{N}^*\mathbf{N} = \mathbf{N}\mathbf{N}^* \impliedby \left\|\mathbf{N}^*\boldsymbol{x}\right\| = \|\mathbf{N}\boldsymbol{x}\|$:

$$\begin{aligned}
\langle \mathbf{N}^*\mathbf{N}\boldsymbol{x} \mid \boldsymbol{x}\rangle &= \langle \mathbf{N}\boldsymbol{x} \mid \mathbf{N}^{**}\boldsymbol{x}\rangle && \text{by Proposition N.3 page 316 (definition of } \mathbf{N}^*)\\
&= \langle \mathbf{N}\boldsymbol{x} \mid \mathbf{N}\boldsymbol{x}\rangle && \text{by Theorem N.13 page 317 (property of adjoint)}\\
&= \|\mathbf{N}\boldsymbol{x}\|^2 && \text{by definition}\\
&= \left\|\mathbf{N}^*\boldsymbol{x}\right\|^2 && \text{by right hypothesis } (\left\|\mathbf{N}^*\boldsymbol{x}\right\| = \|\mathbf{N}\boldsymbol{x}\|)\\
&= \langle \mathbf{N}^*\boldsymbol{x} \mid \mathbf{N}^*\boldsymbol{x}\rangle && \text{by definition}\\
&= \langle \mathbf{N}\mathbf{N}^*\boldsymbol{x} \mid \boldsymbol{x}\rangle && \text{by Proposition N.3 page 316 (definition of } \mathbf{N}^*)
\end{aligned}$$

✏

Theorem N.20. [39] *Let $\mathcal{B}(\boldsymbol{H},\boldsymbol{H})$ be the space of* BOUNDED LINEAR OPERATORS *on a* HILBERT SPACE $\boldsymbol{H}$. *Let $\mathcal{N}(\mathbf{N})$ be the* NULL SPACE *of an operator $\mathbf{N}$ in $\mathcal{B}(\boldsymbol{H},\boldsymbol{H})$ and $\mathcal{I}(\mathbf{N})$ the* IMAGE SET *of $\mathbf{N}$ in $\mathcal{B}(\boldsymbol{H},\boldsymbol{H})$.*

THM
$$\underbrace{\mathbf{N}^*\mathbf{N} = \mathbf{N}\mathbf{N}^*}_{\mathbf{N}\text{ is normal}} \quad \Longrightarrow \quad \underbrace{\mathcal{N}(\mathbf{N}^*) = \mathcal{N}(\mathbf{N})}_{\mathbf{N}\text{ and }\mathbf{N}^*\text{ have the same null space}}$$

✎PROOF:

$$\begin{aligned}
\mathcal{N}(\mathbf{N}^*) &= \left\{\boldsymbol{x} \middle| \mathbf{N}^*\boldsymbol{x} = 0 \quad \forall \boldsymbol{x} \in \boldsymbol{X}\right\} && \text{(definition of } \mathcal{N})\\
&= \left\{\boldsymbol{x} \middle| \left\|\mathbf{N}^*\boldsymbol{x}\right\| = 0 \quad \forall \boldsymbol{x} \in \boldsymbol{X}\right\} && \text{by definition of } \|\cdot\| \text{ (Definition K.1 page 271)}\\
&= \left\{\boldsymbol{x} \middle| \|\mathbf{N}\boldsymbol{x}\| = 0 \quad \forall \boldsymbol{x} \in \boldsymbol{X}\right\}\\
&= \left\{\boldsymbol{x} \middle| \mathbf{N}\boldsymbol{x} = 0 \quad \forall \boldsymbol{x} \in \boldsymbol{X}\right\} && \text{by definition of } \|\cdot\| \text{ (Definition K.1 page 271)}\\
&= \mathcal{N}(\mathbf{N}) && \text{(definition of } \mathcal{N})
\end{aligned}$$

✏

Theorem N.21. [40] *Let $\mathcal{B}(\boldsymbol{H},\boldsymbol{H})$ be the space of* BOUNDED LINEAR OPERATORS *on a* HILBERT SPACE $\boldsymbol{H}$. *Let $\mathcal{N}(\mathbf{N})$ be the* NULL SPACE *of an operator $\mathbf{N}$ in $\mathcal{B}(\boldsymbol{H},\boldsymbol{H})$ and $\mathcal{I}(\mathbf{N})$ the* IMAGE SET *of $\mathbf{N}$ in $\mathcal{B}(\boldsymbol{H},\boldsymbol{H})$.* [41]

THM
$$\left\{\underbrace{\mathbf{N}^*\mathbf{N} = \mathbf{N}\mathbf{N}^*}_{\mathbf{N}\text{ is normal}}\right\} \quad \Longrightarrow \quad \left\{\underbrace{\lambda_n \neq \lambda_m \implies \langle \psi_n \mid \psi_m\rangle = 0}_{\text{eigenfunctions associated with distinct eigenvalues are orthogonal}}\right\}$$

[38] Rudin (1991) pages 312–313
[39] Rudin (1991) pages 312–313
[40] Rudin (1991) pages 312–313
[41] *image set*: Definition ?? page ??

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

✎PROOF: The proof in (1) is flawed. This implies that (2) is also flawed. [Rudin] claims both to be true.(Rudin, 1991)313

1. Proof that $\mathbf{N}^*\mathbf{N} = \mathbf{N}\mathbf{N}^* \implies \mathbf{N}^*\psi = \lambda^*\psi$:

$$
\begin{aligned}
&\mathbf{N}\psi = \lambda\psi \\
&\Longleftarrow \\
&\quad 0 = \mathcal{N}(\mathbf{N} - \lambda\mathbf{I}) \\
&\qquad = \mathcal{N}([\mathbf{N} - \lambda\mathbf{I}]^*) && \text{by } \mathcal{N}(\mathbf{N}) = \mathcal{N}(\mathbf{N}^*) \\
&\qquad = \mathcal{N}(\mathbf{N}^* - [\lambda\mathbf{I}]^*) && \text{by Theorem N.13 page 317} \\
&\qquad = \mathcal{N}(\mathbf{N}^* - \lambda^*\mathbf{I}^*) && \text{by Theorem N.13 page 317} \\
&\qquad = \mathcal{N}(\mathbf{N}^* - \lambda^*\mathbf{I}) \\
&\Longrightarrow \\
&(\mathbf{N}^* - \lambda^*\mathbf{I})\psi = 0 \\
&\Longleftrightarrow \mathbf{N}^*\psi = \lambda^*\psi
\end{aligned}
$$

2. Proof that $\mathbf{N}^*\mathbf{N} = \mathbf{N}\mathbf{N}^* \implies [\lambda_n \neq \lambda_m \implies \langle\psi_n \,|\, \psi_m\rangle = 0]$:

$$
\begin{aligned}
\lambda_n \langle\psi_n \,|\, \psi_m\rangle &= \langle\lambda_n\psi_n \,|\, \psi_m\rangle && \text{by definition of } \langle\triangle \,|\, \triangledown\rangle \text{ Definition J.1 page 255} \\
&= \langle\mathbf{N}\psi_n \,|\, \psi_m\rangle && \text{by definition of eigenpairs} \\
&= \langle\psi_n \,|\, \mathbf{N}^*\psi_m\rangle && \text{by Proposition N.3 page 316 (definition of adjoint)} \\
&= \langle\psi_n \,|\, \lambda_m^*\psi_m\rangle && \text{by (4.)} \\
&= \lambda_m \langle\psi_n \,|\, \psi_m\rangle && \text{by definition of } \langle\triangle \,|\, \triangledown\rangle \text{ Definition J.1 page 255}
\end{aligned}
$$

This implies for $\lambda_n \neq \lambda_m \neq 0$, $\langle\psi_n \,|\, \psi_m\rangle = 0$.

⇨

N.4.4 Isometric operators

An operator on a pair of normed linear spaces is *isometric* (next definition) if it is an *isometry*.

Definition N.10. *Let* $\big(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \|\cdot\|\big)$ *and* $\big(Y, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \|\cdot\|\big)$ *be* NORMED LINEAR SPACES *(Definition K.1 page 271)*.

DEF *An operator* $\mathbf{M} \in \mathcal{L}(\boldsymbol{X}, \boldsymbol{Y})$ *is* ***isometric*** *if*
$$\|\mathbf{M}\boldsymbol{x}\| = \|\boldsymbol{x}\| \qquad \forall_{\boldsymbol{x}\in X}.$$

Theorem N.22. [42] *Let* $\big(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \|\cdot\|\big)$ *and* $\big(Y, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \|\cdot\|\big)$ *be* NORMED LINEAR SPACES. *Let* $\mathbf{M}$ *be a linear operator in* $\mathcal{L}(\boldsymbol{X}, \boldsymbol{Y})$.

THM
$$\underbrace{\|\mathbf{M}\boldsymbol{x}\| = \|\boldsymbol{x}\| \quad \forall_{\boldsymbol{x}\in X}}_{\textit{isometric in length}} \qquad \Longleftrightarrow \qquad \underbrace{\|\mathbf{M}\boldsymbol{x} - \mathbf{M}\boldsymbol{y}\| = \|\boldsymbol{x} - \boldsymbol{y}\| \quad \forall_{\boldsymbol{x},\boldsymbol{y}\in X}}_{\textit{isometric in distance}}$$

✎PROOF:

[42] Kubrusly (2001) page 239 ⟨Proposition 4.37⟩, Berberian (1961) page 27 ⟨Theorem IV.7.5⟩

COPYRIGHT © 2019 DANIEL J. GREENHOE

1. Proof that $\|\mathbf{M}\boldsymbol{x}\| = \|\boldsymbol{x}\| \implies \|\mathbf{M}\boldsymbol{x} - \mathbf{M}\boldsymbol{y}\| = \|\boldsymbol{x} - \boldsymbol{y}\|$:

$$\begin{aligned}
\|\mathbf{M}\boldsymbol{x} - \mathbf{M}\boldsymbol{y}\| &= \|\mathbf{M}(\boldsymbol{x} - \boldsymbol{y})\| && \text{by definition of linear operators (Definition N.3 page 304)}\\
&= \|\mathbf{M}\boldsymbol{u}\| && \text{let } \boldsymbol{u} \triangleq \boldsymbol{x} - \boldsymbol{y}\\
&= \|\boldsymbol{x} - \boldsymbol{y}\| && \text{by left hypothesis}
\end{aligned}$$

2. Proof that $\|\mathbf{M}\boldsymbol{x}\| = \|\boldsymbol{x}\| \impliedby \|\mathbf{M}\boldsymbol{x} - \mathbf{M}\boldsymbol{y}\| = \|\boldsymbol{x} - \boldsymbol{y}\|$:

$$\begin{aligned}
\|\mathbf{M}\boldsymbol{x}\| &= \|\mathbf{M}(\boldsymbol{x} - \mathbb{0})\|\\
&= \|\mathbf{M}\boldsymbol{x} - \mathbf{M}\mathbb{0})\| && \text{by definition of linear operators (Definition N.3 page 304)}\\
&= \|\boldsymbol{x} - \mathbb{0}\| && \text{by right hypothesis}\\
&= \|\boldsymbol{x}\|
\end{aligned}$$

✏

Isometric operators have already been defined (Definition N.10 page 323) in the more general normed linear spaces, while Theorem N.22 (page 323) demonstrated that in a normed linear space $\boldsymbol{X}$, $\|\mathbf{M}\boldsymbol{x}\| = \|\boldsymbol{x}\| \iff \|\mathbf{M}\boldsymbol{x} - \mathbf{M}\boldsymbol{y}\| = \|\boldsymbol{x} - \boldsymbol{y}\|$ for all $\boldsymbol{x}, \boldsymbol{y} \in \boldsymbol{X}$. Here in the more specialized inner product spaces, Theorem N.23 (next) demonstrates two additional equivalent properties.

Theorem N.23. [43] *Let $\mathcal{B}(\boldsymbol{X}, \boldsymbol{X})$ be the space of* BOUNDED LINEAR OPERATORS *on a normed linear space $\boldsymbol{X} \triangleq \big(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \|\cdot\|\big)$. Let $\mathbf{N}$ be a bounded linear operator in $\mathcal{L}(\boldsymbol{X}, \boldsymbol{X})$, and $\mathbf{I}$ the identity operator in $\mathcal{L}(\boldsymbol{X}, \boldsymbol{X})$. Let $\|\boldsymbol{x}\| \triangleq \sqrt{\langle \boldsymbol{x} \mid \boldsymbol{x}\rangle}$.*

THM

*The following conditions are all **equivalent**:*

1.	$\mathbf{M}^*\mathbf{M}$	$=$	$\mathbf{I}$			$\iff$
2.	$\langle \mathbf{M}\boldsymbol{x} \mid \mathbf{M}\boldsymbol{y}\rangle$	$=$	$\langle \boldsymbol{x} \mid \boldsymbol{y}\rangle$	$\forall \boldsymbol{x},\boldsymbol{y} \in X$	*($\mathbf{M}$ is surjective)*	$\iff$
3.	$\|\mathbf{M}\boldsymbol{x} - \mathbf{M}\boldsymbol{y}\|$	$=$	$\|x - y\|$	$\forall \boldsymbol{x},\boldsymbol{y} \in X$	*(isometric in distance)*	$\iff$
4.	$\|\mathbf{M}\boldsymbol{x}\|$	$=$	$\|\boldsymbol{x}\|$	$\forall \boldsymbol{x} \in X$	*(isometric in length)*	

✎PROOF:

1. Proof that (1) $\implies$ (2):

$$\begin{aligned}
\langle \mathbf{M}\boldsymbol{x} \mid \mathbf{M}\boldsymbol{y}\rangle &= \langle \boldsymbol{x} \mid \mathbf{M}^*\mathbf{M}\boldsymbol{y}\rangle && \text{by Proposition N.3 page 316 (definition of adjoint)}\\
&= \langle \boldsymbol{x} \mid \mathbf{I}\boldsymbol{y}\rangle && \text{by (1)}\\
&= \langle \boldsymbol{x} \mid \boldsymbol{y}\rangle && \text{by Definition N.2 page 303 (definition of } \mathbf{I})
\end{aligned}$$

2. Proof that (2) $\implies$ (4):

$$\begin{aligned}
\|\mathbf{M}\boldsymbol{x}\| &= \sqrt{\langle \mathbf{M}\boldsymbol{x} \mid \mathbf{M}\boldsymbol{x}\rangle} && \text{by definition of } \|\cdot\|\\
&= \sqrt{\langle \boldsymbol{x} \mid \boldsymbol{x}\rangle} && \text{by right hypothesis}\\
&= \|\boldsymbol{x}\| && \text{by definition of } \|\cdot\|
\end{aligned}$$

3. Proof that (2) $\impliedby$ (4):

$$\begin{aligned}
4\langle \mathbf{M}\boldsymbol{x} \mid \mathbf{M}\boldsymbol{y}\rangle &= \|\mathbf{M}\boldsymbol{x} + \mathbf{M}\boldsymbol{y}\|^2 - \|\mathbf{M}\boldsymbol{x} - \mathbf{M}\boldsymbol{y}\|^2 + i\,\|\mathbf{M}\boldsymbol{x} + i\mathbf{M}\boldsymbol{y}\|^2 - i\,\|\mathbf{M}\boldsymbol{x} - i\mathbf{M}\boldsymbol{y}\|^2 && \text{by polarization id.}\\
&= \|\mathbf{M}(\boldsymbol{x} + \boldsymbol{y})\|^2 - \|\mathbf{M}(\boldsymbol{x} - \boldsymbol{y})\|^2 + i\,\|\mathbf{M}(\boldsymbol{x} + i\boldsymbol{y})\|^2 - i\,\|\mathbf{M}(\boldsymbol{x} - i\boldsymbol{y})\|^2 && \text{by Definition N.3}\\
&= \|\boldsymbol{x} + \boldsymbol{y}\|^2 - \|\boldsymbol{x} - \boldsymbol{y}\|^2 + i\,\|\boldsymbol{x} + i\boldsymbol{y}\|^2 - i\,\|\boldsymbol{x} - i\boldsymbol{y}\|^2 && \text{by left hypothesis}
\end{aligned}$$

[43] Michel and Herget (1993) page 432 ⟨Theorem 7.5.8⟩, Kubrusly (2001) page 391 ⟨Proposition 5.72⟩

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

4. Proof that (3) $\iff$ (4): by Theorem N.22 page 323

5. Proof that (4) $\implies$ (1):

$$\begin{aligned}
\langle \mathbf{M}^*\mathbf{M}\boldsymbol{x} \mid \boldsymbol{x}\rangle &= \langle \mathbf{M}\boldsymbol{x} \mid \mathbf{M}^{**}\boldsymbol{x}\rangle && \text{by Proposition N.3 page 316 (definition of adjoint)}\\
&= \langle \mathbf{M}\boldsymbol{x} \mid \mathbf{M}\boldsymbol{x}\rangle && \text{by Theorem N.13 page 317 (property of adjoint)}\\
&= \|\mathbf{M}\boldsymbol{x}\|^2 && \text{by definition}\\
&= \|\boldsymbol{x}\|^2 && \text{by left hypothesis with } y = 0\\
&= \langle \boldsymbol{x} \mid \boldsymbol{x}\rangle && \text{by definition}\\
&= \langle \mathbf{I}\boldsymbol{x} \mid \boldsymbol{x}\rangle && \text{by Definition N.2 page 303 (definition of } \mathbf{I})\\
\implies \quad \mathbf{M}^*\mathbf{M} &= \mathbf{I} && \forall \boldsymbol{x} \in X
\end{aligned}$$

✏

Theorem N.24. [44] *Let $\mathcal{B}(\boldsymbol{X}, \boldsymbol{Y})$ be the space of* BOUNDED LINEAR OPERATORS *on normed linear spaces $\boldsymbol{X}$ and $\boldsymbol{Y}$. Let $\mathbf{M}$ be a bounded linear operator in $\mathcal{B}(\boldsymbol{X}, \boldsymbol{Y})$, and $\mathbf{I}$ the identity operator in $\mathcal{L}(\boldsymbol{X}, \boldsymbol{X})$. Let Λ be the set of eigenvalues of $\mathbf{M}$. Let $\|\boldsymbol{x}\| \triangleq \sqrt{\langle \boldsymbol{x} \mid \boldsymbol{x}\rangle}$.*

THM

$$\underbrace{\mathbf{M}^*\mathbf{M} = \mathbf{I}}_{\mathbf{M}\text{ is isometric}} \implies \begin{cases} |||\mathbf{M}||| = 1 & \text{(UNIT LENGTH) and}\\ |\lambda| = 1 & \forall \lambda \in \Lambda \end{cases}$$

✎PROOF:

1. Proof that $\mathbf{M}^*\mathbf{M} = \mathbf{I} \implies |||\mathbf{M}||| = 1$:

$$\begin{aligned}
|||\mathbf{M}||| &= \sup_{\boldsymbol{x}\in X}\{\|\mathbf{M}\boldsymbol{x}\| \mid \|\boldsymbol{x}\| = 1\} && \text{by Definition N.4 page 308}\\
&= \sup_{\boldsymbol{x}\in X}\{\|\boldsymbol{x}\| \mid \|\boldsymbol{x}\| = 1\} && \text{by Theorem N.23 page 324}\\
&= \sup_{\boldsymbol{x}\in X}\{1\}\\
&= 1
\end{aligned}$$

2. Proof that $|\lambda| = 1$: Let $(\boldsymbol{x}, \lambda)$ be an eigenvector-eigenvalue pair.

$$\begin{aligned}
1 &= \frac{1}{\|\boldsymbol{x}\|}\ \|\boldsymbol{x}\|\\
&= \frac{1}{\|\boldsymbol{x}\|}\ \|\mathbf{M}\boldsymbol{x}\| && \text{by Theorem N.23 page 324}\\
&= \frac{1}{\|\boldsymbol{x}\|}\ \|\lambda\boldsymbol{x}\| && \text{by definition of } \lambda\\
&= \frac{1}{\|\boldsymbol{x}\|}\ |\lambda|\ \|\boldsymbol{x}\| && \text{by homogeneous property of } \|\cdot\|\\
&= |\lambda|
\end{aligned}$$

✏

Example N.4 (One sided shift operator). [45] Let $\boldsymbol{X}$ be the set of all sequences with range $\mathbb{W}$ $(0, 1, 2, \ldots)$ and shift operators defined as

$$\begin{aligned}
&1.\quad \mathbf{S}_r\,(\!(x_0, x_1, x_2, \ldots)\!) \triangleq (\!(0, x_0, x_1, x_2, \ldots)\!) && \text{(right shift operator)}\\
&2.\quad \mathbf{S}_l\,(\!(x_0, x_1, x_2, \ldots)\!) \triangleq (\!(x_1, x_2, x_3, \ldots)\!) && \text{(left shift operator)}
\end{aligned}$$

[44] Michel and Herget (1993) page 432
[45] Michel and Herget (1993) page 441

COPYRIGHT © 2019 DANIEL J. GREENHOE | CC BY-NC-ND

EX

1. $\mathbf{S}_r$ is an isometric operator.
2. $\mathbf{S}_r^* = \mathbf{S}_l$

✎PROOF:

1. Proof that $\mathbf{S}_r^* = \mathbf{S}_l$:

$$\begin{aligned}
\langle \mathbf{S}_r (\!(x_0, x_1, x_2, \ldots)\!) \mid (\!(y_0, \boldsymbol{y}_1, \boldsymbol{y}_2, \ldots)\!) \rangle &= \langle (\!(0, x_0, x_1, x_2, \ldots)\!) \mid (\!(y_0, \boldsymbol{y}_1, \boldsymbol{y}_2, \ldots)\!) \rangle \\
&= \sum_{n=1}^{\infty} \boldsymbol{x}_{n-1}\ \boldsymbol{y}_n^* \\
&= \sum_{n=0}^{\infty} \boldsymbol{x}_n\ \boldsymbol{y}_{n+1}^* \\
&= \sum_{n=0}^{\infty} \boldsymbol{x}_n\ \boldsymbol{y}_{n+1}^* \\
&= \langle (\!(x_0, x_1, x_2, \ldots)\!) \mid (\!(y_1, \boldsymbol{y}_2, \boldsymbol{y}_3, \ldots)\!) \rangle \\
&= \left\langle (\!(x_0, x_1, x_2, \ldots)\!) \mid \underbrace{\mathbf{S}_l}_{\mathbf{S}_r^*} (\!(y_0, \boldsymbol{y}_1, \boldsymbol{y}_2, \ldots)\!) \right\rangle
\end{aligned}$$

2. Proof that $\mathbf{S}_r$ is isometric ($\mathbf{S}_r^*\mathbf{S}_r = \mathbf{I}$):

$$\begin{aligned}
\mathbf{S}_r^*\mathbf{S}_r &= \mathbf{S}_l\mathbf{S}_r && \text{by 1.} \\
&= \mathbf{I}
\end{aligned}$$

✏

N.4.5 Unitary operators

Definition N.11. [46] *Let $\mathcal{B}(\boldsymbol{X}, \boldsymbol{Y})$ be the space of* BOUNDED LINEAR OPERATORS *on normed linear spaces $\boldsymbol{X}$ and $\boldsymbol{Y}$. Let $\mathbf{U}$ be a bounded linear operator in $\mathcal{B}(\boldsymbol{X}, \boldsymbol{Y})$, and $\boldsymbol{I}$ the identity operator in $\mathcal{B}(\boldsymbol{X}, \boldsymbol{X})$.*

DEF

*The operator $\mathbf{U}$ is **unitary** if* $\mathbf{U}^*\mathbf{U} = \mathbf{U}\mathbf{U}^* = \mathbf{I}$.

Proposition N.4. *Let $\mathcal{B}(\boldsymbol{X}, \boldsymbol{Y})$ be the space of* BOUNDED LINEAR OPERATORS *on normed linear spaces $\boldsymbol{X}$ and $\boldsymbol{Y}$. Let $\mathbf{U}$ and $\mathbf{V}$ be* BOUNDED LINEAR OPERATORS *in $\mathcal{B}(\boldsymbol{X}, \boldsymbol{Y})$.*

PRP

$$\left.\begin{array}{l} \mathbf{U} \text{ is UNITARY} \quad \textit{and} \\ \mathbf{V} \text{ is UNITARY} \end{array}\right\} \implies (\mathbf{UV}) \text{ is UNITARY.}$$

[46] Rudin (1991) page 312, Michel and Herget (1993) page 431, Autonne (1901) page 209, Autonne (1902), Schur (1909), Steen (1973)

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

✎PROOF:

$$\begin{aligned}
(\mathbf{UV})(\mathbf{UV})^* &= (\mathbf{UV})\left(\mathbf{V}^*\mathbf{U}^*\right) && \text{by Theorem N.8 page 313}\\
&= \mathbf{U}\left(\mathbf{VV}^*\right)\mathbf{U}^* && \text{by associative property}\\
&= \mathbf{UIU}^* && \text{by definition of unitary operators—Definition N.11 page 326}\\
&= \mathbf{I} && \text{by definition of unitary operators—Definition N.11 page 326}
\end{aligned}$$

$$\begin{aligned}
(\mathbf{UV})^*(\mathbf{UV}) &= \left(\mathbf{V}^*\mathbf{U}^*\right)(\mathbf{UV}) && \text{by Theorem N.8 page 313}\\
&= \mathbf{V}^*\left(\mathbf{U}^*\mathbf{U}\right)\mathbf{V} && \text{by associative property}\\
&= \mathbf{V}^*\mathbf{IV} && \text{by definition of unitary operators—Definition N.11 page 326}\\
&= \mathbf{I} && \text{by definition of unitary operators—Definition N.11 page 326}
\end{aligned}$$

✏

Theorem N.25. [47] *Let $\mathcal{B}(\boldsymbol{H},\boldsymbol{H})$ be the space of* BOUNDED LINEAR OPERATORS *on a* HILBERT SPACE $\boldsymbol{H}$. *Let* $\mathbf{U}$ *be a bounded linear operator in $\mathcal{B}(\boldsymbol{H},\boldsymbol{H})$, and $\mathcal{I}(\mathbf{U})$ the* IMAGE SET *of* $\mathbf{U}$.

THM

The following conditions are ***equivalent****:*

1. $\mathbf{UU}^* = \mathbf{U}^*\mathbf{U} = \mathbf{I}$ *(unitary)* $\iff$
2. $\langle \mathbf{U}\boldsymbol{x} \mid \mathbf{U}\boldsymbol{y}\rangle = \langle \mathbf{U}^*\boldsymbol{x} \mid \mathbf{U}^*\boldsymbol{y}\rangle = \langle \boldsymbol{x} \mid \boldsymbol{y}\rangle$ *and* $\mathcal{I}(\mathbf{U}) = X$ *(surjective)* $\iff$
3. $\|\mathbf{U}\boldsymbol{x} - \mathbf{U}\boldsymbol{y}\| = \left\|\mathbf{U}^*\boldsymbol{x} - \mathbf{U}^*\boldsymbol{y}\right\| = \|\boldsymbol{x} - \boldsymbol{y}\|$ *and* $\mathcal{I}(\mathbf{U}) = X$ *(isometric in distance)* $\iff$
4. $\|\mathbf{U}\boldsymbol{x}\| = \|\boldsymbol{x}\|$ *and* $\mathcal{I}(\mathbf{U}) = X$ *(isometric in length)*

✎PROOF:

1. Proof that (1) $\implies$ (2):
   (a) $\langle \mathbf{U}\boldsymbol{x} \mid \mathbf{U}\boldsymbol{y}\rangle = \langle \mathbf{U}^*\boldsymbol{x} \mid \mathbf{U}^*\boldsymbol{y}\rangle = \langle \boldsymbol{x} \mid \boldsymbol{y}\rangle$ by Theorem N.23 (page 324).
   (b) Proof that $\mathcal{I}(\mathbf{U}) = X$:

$$\begin{aligned}
X &\supseteq \mathcal{I}(\mathbf{U}) && \text{because } \mathbf{U} \in X^X\\
&\supseteq \mathcal{I}(\mathbf{UU}^*)\\
&= \mathcal{I}(\mathbf{I}) && \text{by left hypothesis } (\mathbf{U}^*\mathbf{U} = \mathbf{UU}^* = \mathbf{I})\\
&= X && \text{by Definition N.2 page 303 (definition of } \mathbf{I})
\end{aligned}$$

2. Proof that (2) $\iff$ (3) $\iff$ (4): by Theorem N.23 page 324.
3. Proof that (3) $\implies$ (1):
   (a) Proof that $\|\mathbf{U}\boldsymbol{x} - \mathbf{U}\boldsymbol{y}\| = \|\boldsymbol{x} - \boldsymbol{y}\| \implies \mathbf{U}^*\mathbf{U} = \mathbf{I}$: by Theorem N.23 page 324
   (b) Proof that $\left\|\mathbf{U}^*\boldsymbol{x} - \mathbf{U}^*\boldsymbol{y}\right\| = \|\boldsymbol{x} - \boldsymbol{y}\| \implies \mathbf{UU}^* = \mathbf{I}$:

$$\begin{aligned}
\left\|\mathbf{U}^*\boldsymbol{x} - \mathbf{U}^*\boldsymbol{y}\right\| = \|\boldsymbol{x} - \boldsymbol{y}\| \implies \quad \mathbf{U}^{**}\mathbf{U}^* &= \mathbf{I} && \text{by Theorem N.23 page 324}\\
\mathbf{UU}^* &= \mathbf{I} && \text{by Theorem N.13 page 317}
\end{aligned}$$

✏

Theorem N.26. *Let $\mathcal{B}(\boldsymbol{H},\boldsymbol{H})$ be the space of* BOUNDED LINEAR OPERATORS *on a* HILBERT SPACE $\boldsymbol{H}$. *Let* $\mathbf{U}$ *be a bounded linear operator in $\mathcal{B}(\boldsymbol{H},\boldsymbol{H})$, $\mathcal{N}(\mathbf{U})$ the* NULL SPACE *of* $\mathbf{U}$, *and $\mathcal{I}(\mathbf{U})$ the* IMAGE SET *of* $\mathbf{U}$.

THM

$$\underbrace{\mathbf{UU}^* = \mathbf{U}^*\mathbf{U} = \mathbf{I}}_{\mathbf{U}\text{ is unitary}} \implies \begin{cases}
\mathbf{U}^{-1} = \mathbf{U}^* & \\
\mathcal{I}(\mathbf{U}) = \mathcal{I}(\mathbf{U}^*) = X & \\
\mathcal{N}(\mathbf{U}) = \mathcal{N}(\mathbf{U}^*) = \{\mathbb{0}\} & \\
|||\mathbf{U}||| = \left|||\mathbf{U}^*\right||| = 1 & \text{(UNIT LENGTH)}
\end{cases}$$

[47] Rudin (1991) pages 313–314 ⟨Theorem 12.13⟩, Knapp (2005a) page 45 ⟨Proposition 2.6⟩

COPYRIGHT © 2019 DANIEL J. GREENHOE | CC BY-NC-ND

✎PROOF:

1. Note that $\mathbf{U}$, $\mathbf{U}^*$, and $\mathbf{U}^{-1}$ are all both **isometric** and **normal**:

$$\begin{array}{lllllll}
\mathbf{U}^*\mathbf{U} & & & = & \mathbf{I} & \Longrightarrow & \mathbf{U} \text{ is isometric} \\
\mathbf{U}\mathbf{U}^* & = & \mathbf{U}^*\mathbf{U} & = & \mathbf{I} & \Longrightarrow & \mathbf{U}^* \text{ is isometric} \\
\mathbf{U}^{-1} & = & \mathbf{U}^* & & & \Longrightarrow & \mathbf{U}^{-1} \text{ is isometric} \\
 & & & & & & \\
\mathbf{U}^*\mathbf{U} & = & \mathbf{U}\mathbf{U}^* & = & & \Longrightarrow & \mathbf{U} \text{ is normal} \\
\mathbf{U}\mathbf{U}^* & = & \mathbf{U}^*\mathbf{U} & = & & \Longrightarrow & \mathbf{U}^* \text{ is normal} \\
\mathbf{U}^{-1} & = & \mathbf{U}^* & & & \Longrightarrow & \mathbf{U}^{-1} \text{ is normal}
\end{array}$$

2. Proof that $\mathbf{U}^*\mathbf{U} = \mathbf{U}\mathbf{U}^* = \mathbf{I} \implies \mathcal{I}(\mathbf{U}) = \mathcal{I}(\mathbf{U}^*) = \boldsymbol{H}$: by Theorem N.25 page 327.

3. Proof that $\mathbf{U}^*\mathbf{U} = \mathbf{U}\mathbf{U}^* = \mathbf{I} \implies \mathcal{N}(\mathbf{U}) = \mathcal{N}(\mathbf{U}^*) = \mathcal{N}(\mathbf{U}^{-1})$:

$$\begin{array}{ll}
\mathcal{N}(\mathbf{U}^*) = \mathcal{N}(\mathbf{U}) & \text{because } \mathbf{U} \text{ and } \mathbf{U}^* \text{ are both normal and by Theorem N.21 page 322} \\
\quad = \mathcal{I}(\mathbf{U})^\perp & \text{by Theorem N.14 page 318} \\
\quad = X^\perp & \text{by above result} \\
\quad = \{\mathbb{0}\} & \text{by Proposition ?? page ??}
\end{array}$$

4. Proof that $\mathbf{U}^*\mathbf{U} = \mathbf{U}\mathbf{U}^* = \mathbf{I} \implies \|\mathbf{U}^{-1}\| = \|\mathbf{U}^*\| = \|\mathbf{U}\| = 1$:
Because $\mathbf{U}$, $\mathbf{U}^*$, and $\mathbf{U}^{-1}$ are all isometric and by Theorem N.24 page 325.

✏

Example N.5. Examples of *Fredholm integral operators* include

1.	Fourier Transform	$[\tilde{\mathbf{F}}\mathsf{x}](f)$	$= \int_t \mathsf{x}(t)e^{-i2\pi ft}\,\mathrm{d}t$	$\kappa(t,f)$	$= e^{-i2\pi ft}$
2.	Inverse Fourier Transform	$[\tilde{\mathbf{F}}^{-1}\tilde{\mathsf{x}}](t)$	$= \int_f \tilde{\mathsf{x}}(f)e^{i2\pi ft}\,\mathrm{d}f$	$\kappa(f,t)$	$= e^{i2\pi ft}$
3.	Laplace operator	$[\mathbf{L}\mathsf{x}](s)$	$= \int_t \mathsf{x}(t)e^{-st}\,\mathrm{d}t$	$\kappa(t,s)$	$= e^{-st}$

Example N.6 (Translation operator). Let $\boldsymbol{X} = \boldsymbol{L}^2_{\mathbb{R}}$ and $\mathbf{T} \in \boldsymbol{X}^{\boldsymbol{X}}$ be defined as

$$\mathbf{T}\mathsf{f}(x) \triangleq \mathsf{f}(x-1) \qquad \forall \mathsf{f} \in \boldsymbol{L}^2_{\mathbb{R}} \qquad \text{(translation operator)}$$

EX

1.	$\mathbf{T}^{-1}\mathsf{f}(x)$	$=$	$\mathsf{f}(x+1)$	$\forall \mathsf{f} \in \boldsymbol{L}^2_{\mathbb{R}}$	(inverse translation operator)
2.	$\mathbf{T}^*$	$=$	$\mathbf{T}^{-1}$		($\mathbf{T}$ is invertible)
3.	$\mathbf{T}^*\mathbf{T}$	$=$	$\mathbf{T}\mathbf{T}^* = \mathbf{I}$		($\mathbf{T}$ is unitary)

✎PROOF:

1. Proof that $\mathbf{T}^{-1}\mathsf{f}(x) = \mathsf{f}(x+1)$:

$$\begin{aligned}
\mathbf{T}^{-1}\mathbf{T} &= \mathbf{I} \\
\mathbf{T}\mathbf{T}^{-1} &= \mathbf{I}
\end{aligned}$$

2. Proof that $\mathbf{T}$ is unitary:

$$\begin{aligned}
\langle \mathbf{T}\mathsf{f}(x) \,|\, \mathsf{g}(x)\rangle &= \langle \mathsf{f}(x-1) \,|\, \mathsf{g}(x)\rangle && \text{by definition of } \mathbf{T} \\
&= \int_x \mathsf{f}(x-1)\mathsf{g}^*(x)\,\mathrm{d}x \\
&= \int_x \mathsf{f}(x)\mathsf{g}^*(x+1)\,\mathrm{d}x \\
&= \langle \mathsf{f}(x) \,|\, \mathsf{g}(x+1)\rangle \\
&= \left\langle \mathsf{f}(x) \,\middle|\, \underbrace{\mathbf{T}^{-1}}_{\mathbf{T}^*}\mathsf{g}(x) \right\rangle && \text{by 1.}
\end{aligned}$$

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

Example N.7 (Dilation operator). Let $\boldsymbol{X} = \boldsymbol{L}^2_{\mathbb{R}}$ and $\mathbf{T} \in \boldsymbol{X}^{\boldsymbol{X}}$ be defined as

$$\mathbf{D}\mathsf{f}(x) \triangleq \sqrt{2}\mathsf{f}(2x) \qquad \forall \mathsf{f} \in \boldsymbol{L}^2_{\mathbb{R}} \qquad \text{(dilation operator)}$$

EX

1. $\mathbf{D}^{-1}\mathsf{f}(x) = \frac{1}{\sqrt{2}}\mathsf{f}\left(\frac{1}{2}x\right) \qquad \forall \mathsf{f} \in \boldsymbol{L}^2_{\mathbb{R}}$ (inverse dilation operator)
2. $\mathbf{D}^* = \mathbf{D}^{-1}$ ($\mathbf{D}$ is invertible)
3. $\mathbf{D}^*\mathbf{D} = \mathbf{D}\mathbf{D}^* = \mathbf{I}$ ($\mathbf{D}$ is unitary)

✎PROOF:

1. Proof that $\mathbf{D}^{-1}\mathsf{f}(x) = \frac{1}{\sqrt{2}}\mathsf{f}\left(\frac{1}{2}x\right)$:

$$\mathbf{D}^{-1}\mathbf{D} = \mathbf{I}$$
$$\mathbf{D}\mathbf{D}^{-1} = \mathbf{I}$$

2. Proof that $\mathbf{D}$ is unitary:

$$\begin{aligned}
\langle \mathbf{D}\mathsf{f}(x) \,|\, \mathsf{g}(x)\rangle &= \left\langle \sqrt{2}\mathsf{f}(2x) \,|\, \mathsf{g}(x)\right\rangle && \text{by definition of } \mathbf{D}\\
&= \int_x \sqrt{2}\mathsf{f}(2x)\mathsf{g}^*(x)\,\mathrm{d}x\\
&= \int_{u\in\mathbb{R}} \sqrt{2}\mathsf{f}(u)\mathsf{g}^*\left(\frac{1}{2}u\right)\frac{1}{2}\,\mathrm{d}u && \text{let } u \triangleq 2x \quad \Longrightarrow \quad \mathrm{d}x = \frac{1}{2}\,\mathrm{d}u\\
&= \int_{u\in\mathbb{R}} \mathsf{f}(u)\left[\frac{1}{\sqrt{2}}\mathsf{g}\left(\frac{1}{2}u\right)\right]^*\mathrm{d}u\\
&= \left\langle \mathsf{f}(x) \,\middle|\, \frac{1}{\sqrt{2}}\mathsf{g}\left(\frac{1}{2}x\right)\right\rangle\\
&= \left\langle \mathsf{f}(x) \,\middle|\, \underbrace{\mathbf{D}^{-1}}_{\mathbf{D}^*}\mathsf{g}(x)\right\rangle && \text{by 1.}
\end{aligned}$$

Example N.8 (Delay operator). Let $\boldsymbol{X}$ be the set of all sequences and $\mathbf{D} \in \boldsymbol{X}^{\boldsymbol{X}}$ be a delay operator.

EX The delay operator $\mathbf{D}\,(\!(x_n)\!)_{n\in\mathbb{Z}} \triangleq (\!(x_{n-1})\!)_{n\in\mathbb{Z}}$ is unitary.

✎PROOF: The inverse $\mathbf{D}^{-1}$ of the delay operator $\mathbf{D}$ is

$$\mathbf{D}^{-1}\,(\!(x_n)\!)_{n\in\mathbb{Z}} \triangleq (\!(x_{n+1})\!)_{n\in\mathbb{Z}}\,.$$

$$\begin{aligned}
\langle \mathbf{D}\,(\!(x_n)\!) \,|\, (\!(y_n)\!)\rangle &= \langle (\!(x_{n-1})\!) \,|\, (\!(y_n)\!)\rangle && \text{by definition of } \mathbf{D}\\
&= \sum_n \boldsymbol{x}_{n-1}\,\boldsymbol{y}_n^*\\
&= \sum_n \boldsymbol{x}_n\,\boldsymbol{y}_{n+1}^*\\
&= \langle (\!(x_n)\!) \,|\, (\!(y_{n+1})\!)\rangle\\
&= \left\langle (\!(x_n)\!) \,\middle|\, \underbrace{\mathbf{D}^{-1}}_{\mathbf{D}^*}(\!(y_n)\!)\right\rangle
\end{aligned}$$

Therefore, $\mathbf{D}^* = \mathbf{D}^{-1}$. This implies that $\mathbf{D}\mathbf{D}^* = \mathbf{D}^*\mathbf{D} = \mathbf{I}$ which implies that $\mathbf{D}$ is unitary.

COPYRIGHT © 2019 DANIEL J. GREENHOE
BY-NC-ND

Example N.9 (Fourier transform). Let $\tilde{\mathbf{F}}$ be the *Fourier Transform* and $\tilde{\mathbf{F}}^{-1}$ the *inverse Fourier Transform* operator

$$[\tilde{\mathbf{F}}\mathsf{x}](f) \triangleq \int_t \mathsf{x}(t)\underbrace{e^{-i2\pi ft}}_{\kappa(t,f)}\,\mathrm{d}t \qquad \left[\tilde{\mathbf{F}}^{-1}\tilde{\mathsf{x}}\right](t) \triangleq \int_f \tilde{\mathsf{x}}(f)\underbrace{e^{i2\pi ft}}_{\kappa^*(t,f)}\,\mathrm{d}f.$$

EX $\tilde{\mathbf{F}}^* = \tilde{\mathbf{F}}^{-1}$ (the Fourier Transform operator $\tilde{\mathbf{F}}$ is unitary)

✎PROOF:

$$\begin{aligned}
\left\langle \tilde{\mathbf{F}}\mathsf{x} \,|\, \tilde{\mathsf{y}} \right\rangle &= \left\langle \int_t \mathsf{x}(t)e^{-i2\pi ft}\,\mathrm{d}t \,|\, \tilde{\mathsf{y}}(f) \right\rangle \\
&= \int_t \mathsf{x}(t)\left\langle e^{-i2\pi ft} \,|\, \tilde{\mathsf{y}}(f)\right\rangle\,\mathrm{d}t \\
&= \int_t \mathsf{x}(t)\int_f e^{-i2\pi ft}\tilde{\mathsf{y}}^*(f)\,\mathrm{d}f\,\mathrm{d}t \\
&= \int_t \mathsf{x}(t)\left[\int_f e^{i2\pi ft}\tilde{\mathsf{y}}(f)\,\mathrm{d}f\right]^*\mathrm{d}t \\
&= \left\langle \mathsf{x}(t) \,|\, \int_f \tilde{\mathsf{y}}(f)e^{i2\pi ft}\,\mathrm{d}f \right\rangle \\
&= \left\langle \mathsf{x} \,|\, \underbrace{\tilde{\mathbf{F}}^{-1}}_{\tilde{\mathbf{F}}^*}\tilde{\mathsf{y}} \right\rangle
\end{aligned}$$

This implies that $\tilde{\mathbf{F}}$ is unitary ($\tilde{\mathbf{F}}^* = \tilde{\mathbf{F}}^{-1}$). ✏

Example N.10 (Rotation matrix). [48] Let the rotation matrix $\mathbf{R}_\theta : \mathbb{R}^2 \to \mathbb{R}^2$ be defined as

$$\mathbf{R}_\theta \triangleq \begin{bmatrix} \cos\theta & -\sin\theta \\ \sin\theta & \cos\theta \end{bmatrix}$$

EX
1. ${\mathbf{R}^{-1}}_\theta = \mathbf{R}_{-\theta}$
2. ${\mathbf{R}^*}_\theta = {\mathbf{R}^{-1}}_\theta$ ($\mathbf{R}$ is unitary)

✎PROOF:

$$\begin{aligned}
\mathbf{R}^* &= \mathbf{R}^H \\
&= \begin{bmatrix} \cos\theta & -\sin\theta \\ \sin\theta & \cos\theta \end{bmatrix}^H && \text{by definition of } \mathbf{R} \\
&= \begin{bmatrix} \cos\theta & \sin\theta \\ -\sin\theta & \cos\theta \end{bmatrix} && \text{by definition of Hermetian transpose operator } H \\
&= \begin{bmatrix} \cos(-\theta) & -\sin(-\theta) \\ \sin(-\theta) & \cos(-\theta) \end{bmatrix} && \text{by Theorem ?? page ??} \\
&= \mathbf{R}_{-\theta} && \text{by definition of } \mathbf{R} \\
&= \mathbf{R}^{-1} && \text{by 1.}
\end{aligned}$$

✏

[48] Noble and Daniel (1988), page 311

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

N.5 Operator order

Definition N.12. [49] *Let* $\mathbf{P} \in \boldsymbol{Y}^{\boldsymbol{X}}$ *be an operator.*

DEF

$\mathbf{P}$ *is* ***positive*** *if* $\langle \mathbf{P}\boldsymbol{x} \mid \boldsymbol{x}\rangle \geq 0 \;\forall \boldsymbol{x} \in \boldsymbol{X}$.
This condition is denoted $\mathbf{P} \geq 0$.

Theorem N.27. [50]

THM

$$\underbrace{\mathbf{P} \geq 0 \text{ and } \mathbf{Q} \geq 0}_{\mathbf{P} \text{ and } \mathbf{Q} \text{ are both positive}} \quad \Longrightarrow \quad \begin{cases} (\mathbf{P}+\mathbf{Q}) & \geq & 0 & & ((\mathbf{P}+\mathbf{Q}) \text{ is positive}) \\ \mathbf{A}^*\mathbf{P}\mathbf{A} & \geq & 0 & \forall \mathbf{A} \in \mathcal{B}(\boldsymbol{X},\boldsymbol{X}) & (\mathbf{A}^*\mathbf{P}\mathbf{A} \text{ is positive}) \\ \mathbf{A}^*\mathbf{A} & \geq & 0 & \forall \mathbf{A} \in \mathcal{B}(\boldsymbol{X},\boldsymbol{X}) & (\mathbf{A}^*\mathbf{A} \text{ is positive}) \end{cases}$$

✎PROOF:

$\langle (\mathbf{P}+\mathbf{Q})\boldsymbol{x} \mid \boldsymbol{x}\rangle = \langle \mathbf{P}\boldsymbol{x} \mid \boldsymbol{x}\rangle + \langle \mathbf{Q}\boldsymbol{x} \mid \boldsymbol{x}\rangle$	by additive property of $\langle \triangle \mid \triangledown\rangle$ (Definition J.1 page 255)
$\geq \langle \mathbf{P}\boldsymbol{x} \mid \boldsymbol{x}\rangle$	by left hypothesis
≥ 0	by left hypothesis
$\langle \mathbf{A}^*\mathbf{P}\mathbf{A}\boldsymbol{x} \mid \boldsymbol{x}\rangle = \langle \mathbf{P}\mathbf{A}\boldsymbol{x} \mid \mathbf{A}\boldsymbol{x}\rangle$	by definition of adjoint (Proposition N.3 page 316)
$= \langle \mathbf{P}\boldsymbol{y} \mid \boldsymbol{y}\rangle$	where $\boldsymbol{y} \triangleq \mathbf{A}\boldsymbol{x}$
≥ 0	by left hypothesis
$\langle \mathbf{I}\boldsymbol{x} \mid \boldsymbol{x}\rangle = \langle \boldsymbol{x} \mid \boldsymbol{x}\rangle$	by definition of $\mathbf{I}$ (Definition N.2 page 303)
≥ 0	by non-negative property of $\langle \triangle \mid \triangledown\rangle$ (Definition J.1 page 255)
$\Longrightarrow \mathbf{I}$ is positive	
$\langle \mathbf{A}^*\mathbf{A}\boldsymbol{x} \mid \boldsymbol{x}\rangle = \langle \mathbf{A}^*\mathbf{I}\mathbf{A}\boldsymbol{x} \mid \boldsymbol{x}\rangle$	by definition of $\mathbf{I}$ (Definition N.2 page 303)
≥ 0	by two previous results

✏

Definition N.13. [51] *Let* $\mathbf{A}, \mathbf{B} \in \mathcal{B}(\boldsymbol{X}, \boldsymbol{Y})$ *be* BOUNDED *operators.*

DEF

$\mathbf{A} \geq \mathbf{B}$ *("*$\mathbf{A}$ *is greater than or equal to* $\mathbf{B}$*") if*
$\mathbf{A} - \mathbf{B} \geq 0$ *("*$(\mathbf{A} - \mathbf{B})$ *is positive")*

[49] Michel and Herget (1993) page 429 ⟨Definition 7.4.12⟩
[50] Michel and Herget (1993) page 429
[51] Michel and Herget (1993) page 429

COPYRIGHT © 2019 DANIEL J. GREENHOE

Copyright © 2019 Daniel J. Greenhoe

APPENDIX O

FOURIER TRANSFORM

❝The analytical equations ... extend to all general phenomena. There cannot be a language more universal and more simple, more free from errors and from obscurities, ... mathematical analysis is as extensive as nature itself; ... it preserves every principle which it has once acquired; it grows and strengthens itself incessantly in the midst of the many variations and errors of the human mind. Its chief attribute is clearness; it has no marks to express confused notions. It brings together phenomena the most diverse, and discovers the hidden analogies which unite them.❞

Joseph Fourier (1768–1830) [1]

O.1 Definitions

This chapter deals with the *Fourier Transform* in the space of *Lebesgue square-integrable functions* $\boldsymbol{L}^2_{(\mathbb{R},\mathscr{B},\mu)}$, where $\mathbb{R}$ is the set of real numbers, $\mathscr{B}$ is the set of *Borel sets* on $\mathbb{R}$, μ is the standard *Borel measure* on $\mathbb{R}$, and

$$\boldsymbol{L}^2_{(\mathbb{R},\mathscr{B},\mu)} \triangleq \left\{ \mathsf{f} \in \mathbb{R}^{\mathbb{R}} \,\middle|\, \int_{\mathbb{R}} |\mathsf{f}|^2 \, \mathsf{d}\mu < \infty \right\}.$$

Furthermore, $\langle \triangle \mid \triangledown \rangle$ is the *inner product* induced by the operator $\int_{\mathbb{R}} \mathsf{d}\mu$ such that

$$\langle \mathsf{f} \mid \mathsf{g} \rangle \triangleq \int_{\mathbb{R}} \mathsf{f}(x)\mathsf{g}^*(x) \, \mathsf{d}x,$$

and $\left(\boldsymbol{L}^2_{(\mathbb{R},\mathscr{B},\mu)}, \langle \triangle \mid \triangledown \rangle \right)$ is a *Hilbert space*.

Definition O.1. *Let κ be a* FUNCTION *in $\mathbb{C}^{\mathbb{R}^2}$.*

DEF

The function κ is the ***Fourier kernel*** *if* $\quad \kappa(x,\omega) \triangleq e^{i\omega x} \quad \forall_{x,\omega \in \mathbb{R}}$

Definition O.2. [2] *Let $\boldsymbol{L}^2_{(\mathbb{R},\mathscr{B},\mu)}$ be the space of all* LEBESGUE SQUARE-INTEGRABLE FUNCTIONS.

[1] quote: Fourier (1878), pages 7–8 ⟨Preliminary Discourse⟩
image: http://en.wikipedia.org/wiki/File:Fourier2.jpg, public domain

[2] Bachman et al. (2000) page 363, Chorin and Hald (2009) page 13, Loomis and Bolker (1965), page 144, Knapp (2005b) pages 374–375, Fourier (1822), Fourier (1878) page 336?

DEF *The **Fourier Transform** operator $\tilde{\mathbf{F}}$ is defined as*

$$\left[\tilde{\mathbf{F}}\mathsf{f}\right](\omega) \triangleq \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} \mathsf{f}(x)\, e^{-i\omega x}\ \mathrm{d}x \qquad \forall \mathsf{f} \in L^2_{(\mathbb{R},\mathscr{B},\mu)}$$

*This definition of the Fourier Transform is also called the **unitary Fourier Transform**.*

Remark O.1 (**Fourier transform scaling factor**). [3] If the Fourier transform operator $\tilde{\mathbf{F}}$ and inverse Fourier transform operator $\tilde{\mathbf{F}}^{-1}$ are defined as

$$\tilde{\mathbf{F}}\mathsf{f}(x) \triangleq \mathsf{F}(\omega) \triangleq A \int_{\mathbb{R}} \mathsf{f}(x) e^{-i\omega x}\ \mathrm{d}x \qquad \text{and} \qquad \tilde{\mathbf{F}}^{-1}\tilde{\mathsf{f}}(\omega) \triangleq B \int_{\mathbb{R}} \mathsf{F}(\omega) e^{i\omega x}\ \mathrm{d}\omega$$

then A and B can be any constants as long as $AB = \frac{1}{2\pi}$. The Fourier transform is often defined with the scaling factor A set equal to 1 such that $\left[\tilde{\mathbf{F}}\mathsf{f}(x)\right](\omega) \triangleq \int_{\mathbb{R}} \mathsf{f}(x)\, e^{-i\omega x}\ \mathrm{d}x$. In this case, the inverse Fourier transform operator $\tilde{\mathbf{F}}^{-1}$ is either defined as

- $\left[\tilde{\mathbf{F}}^{-1}\mathsf{f}(x)\right](f) \triangleq \int_{\mathbb{R}} \mathsf{f}(x)\, e^{i2\pi f x}\ \mathrm{d}x$ (using oscillatory frequency free variable f) or
- $\left[\tilde{\mathbf{F}}^{-1}\mathsf{f}(x)\right](\omega) \triangleq \frac{1}{2\pi} \int_{\mathbb{R}} \mathsf{f}(x)\, e^{i\omega x}\ \mathrm{d}x$ (using angular frequency free variable ω).

In short, the 2π has to show up somewhere, either in the argument of the exponential ($e^{-i2\pi f t}$) or in front of the integral ($\frac{1}{2\pi}\int \cdots$). One could argue that it is unnecessary to burden the exponential argument with the 2π factor ($e^{-i2\pi f t}$), and thus could further argue in favor of using the angular frequency variable ω thus giving the inverse operator definition $\left[\tilde{\mathbf{F}}^{-1}\mathsf{f}(x)\right](\omega) \triangleq \frac{1}{2\pi}\int_{\mathbb{R}} \mathsf{f}(x)\, e^{-i\omega x}\ \mathrm{d}x$. But this causes a new problem. In this case, the Fourier operator $\tilde{\mathbf{F}}$ is not *unitary* (see Theorem O.2 page 334)—in particular, $\tilde{\mathbf{F}}\tilde{\mathbf{F}}^* \neq \mathbf{I}$, where $\tilde{\mathbf{F}}^*$ is the *adjoint* of $\tilde{\mathbf{F}}$; but rather, $\tilde{\mathbf{F}}\left(\frac{1}{2\pi}\tilde{\mathbf{F}}^*\right) = \left(\frac{1}{2\pi}\tilde{\mathbf{F}}^*\right)\tilde{\mathbf{F}} = \mathbf{I}$. But if we define the operators $\tilde{\mathbf{F}}$ and $\tilde{\mathbf{F}}^{-1}$ to both have the scaling factor $\frac{1}{\sqrt{2\pi}}$, then $\tilde{\mathbf{F}}$ and $\tilde{\mathbf{F}}^{-1}$ are inverses *and* $\tilde{\mathbf{F}}$ is *unitary*—that is, $\tilde{\mathbf{F}}\tilde{\mathbf{F}}^* = \tilde{\mathbf{F}}^*\tilde{\mathbf{F}} = \mathbf{I}$.

O.2 Operator properties

Theorem O.1 (Inverse Fourier transform). [4] *Let $\tilde{\mathbf{F}}$ be the Fourier Transform operator (Definition O.2 page 333). The inverse $\tilde{\mathbf{F}}^{-1}$ of $\tilde{\mathbf{F}}$ is*

THM

$$\left[\tilde{\mathbf{F}}^{-1}\tilde{\mathsf{f}}\right](x) = \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} \tilde{\mathsf{f}}(\omega) e^{i\omega x}\ \mathrm{d}\omega \qquad \forall \tilde{\mathsf{f}} \in L^2_{(\mathbb{R},\mathscr{B},\mu)}$$

Theorem O.2. *Let $\tilde{\mathbf{F}}$ be the Fourier Transform operator with inverse $\tilde{\mathbf{F}}^{-1}$ and adjoint $\tilde{\mathbf{F}}^*$.*

THM

$$\tilde{\mathbf{F}}^* = \tilde{\mathbf{F}}^{-1}$$

✎PROOF:

$$\begin{aligned}
\left\langle \tilde{\mathbf{F}}\mathsf{f} \,\middle|\, \mathsf{g} \right\rangle &= \left\langle \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} \mathsf{f}(x)\, e^{-i\omega x}\ \mathrm{d}x \,\middle|\, \mathsf{g}(\omega) \right\rangle && \text{by definition of } \tilde{\mathbf{F}} \text{ page 333} \\
&= \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} \mathsf{f}(x) \left\langle e^{-i\omega x} \,\middle|\, \mathsf{g}(\omega) \right\rangle\ \mathrm{d}x && \text{by } \textit{additive property} \text{ of } \langle \triangle \mid \triangledown \rangle \text{ page 255} \\
&= \int_{\mathbb{R}} \mathsf{f}(x) \frac{1}{\sqrt{2\pi}} \left\langle \mathsf{g}(\omega) \,\middle|\, e^{-i\omega x} \right\rangle^*\ \mathrm{d}x && \text{by } \textit{conjugate symmetric property} \text{ of } \langle \triangle \mid \triangledown \rangle \text{ page 255}
\end{aligned}$$

[3] Chorin and Hald (2009) page 13, Jeffrey and Dai (2008) pages xxxi–xxxii, Knapp (2005b) pages 374–375
[4] Chorin and Hald (2009) page 13

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

$$= \left\langle \mathsf{f}(x) \,\middle|\, \frac{1}{\sqrt{2\pi}} \left\langle \mathsf{g}(\omega) \,\middle|\, e^{-i\omega x} \right\rangle \right\rangle \qquad \text{by definition of } \langle \triangle \mid \triangledown \rangle$$

$$= \left\langle \mathsf{f} \,\middle|\, \underbrace{\tilde{\mathbf{F}}^{-1}}_{\tilde{\mathbf{F}}^*} \mathsf{g} \right\rangle \qquad \text{by Theorem O.1 page 334}$$

✏

The Fourier Transform operator has several nice properties:

- $\tilde{\mathbf{F}}$ is *unitary*[5] (Corollary O.1—next corollary).
- Because $\tilde{\mathbf{F}}$ is unitary, it automatically has several other nice properties (Theorem O.3 page 335).

Corollary O.1. *Let* $\mathbf{I}$ *be the identity operator and let* $\tilde{\mathbf{F}}$ *be the Fourier Transform operator with adjoint* $\tilde{\mathbf{F}}^*$ *and inverse* $\tilde{\mathbf{F}}^{-1}$.

COR

$$\underbrace{\tilde{\mathbf{F}}\tilde{\mathbf{F}}^* = \tilde{\mathbf{F}}^*\tilde{\mathbf{F}} = \mathbf{I}}_{\tilde{\mathbf{F}}^* = \tilde{\mathbf{F}}^{-1}} \qquad (\tilde{\mathbf{F}} \text{ is unitary})$$

✎PROOF: This follows directly from the fact that $\tilde{\mathbf{F}}^* = \tilde{\mathbf{F}}^{-1}$ (Theorem O.2 page 334). ✏

Theorem O.3. *Let* $\tilde{\mathbf{F}}$ *be the Fourier transform operator with adjoint* $\tilde{\mathbf{F}}^*$ *and inverse* $\tilde{\mathbf{F}}$. *Let* $|||\cdot|||$ *be the operator norm with respect to the vector norm* $\|\cdot\|$ *with respect to the Hilbert space* $\left(\mathbb{C}^{\mathbb{R}}, \langle \triangle \mid \triangledown \rangle\right)$. *Let* $\mathcal{R}(\mathbf{A})$ *be the range of an operator* $\mathbf{A}$.

THM

$$\begin{aligned}
\mathcal{R}(\mathbf{F}\tau) &= \mathcal{R}(\tilde{\mathbf{F}}^{-1}) &&= L^2_{\mathbb{R}} \\
|||\tilde{\mathbf{F}}||| &= |||\tilde{\mathbf{F}}^{-1}||| &&= 1 && \text{(UNITARY)} \\
\left\langle \tilde{\mathbf{F}}\mathsf{f} \,\middle|\, \tilde{\mathbf{F}}\mathsf{g} \right\rangle &= \left\langle \tilde{\mathbf{F}}^{-1}\mathsf{f} \,\middle|\, \tilde{\mathbf{F}}^{-1}\mathsf{g} \right\rangle &&= \langle \mathsf{f} \mid \mathsf{g} \rangle && \text{(PARSEVAL'S EQUATION)} \\
\left\|\tilde{\mathbf{F}}\mathsf{f}\right\| &= \left\|\tilde{\mathbf{F}}^{-1}\mathsf{f}\right\| &&= \|\mathsf{f}\| && \text{(PLANCHEREL'S FORMULA)} \\
\left\|\tilde{\mathbf{F}}\mathsf{f} - \tilde{\mathbf{F}}\mathsf{g}\right\| &= \left\|\tilde{\mathbf{F}}^{-1}\mathsf{f} - \tilde{\mathbf{F}}^{-1}\mathsf{g}\right\| &&= \|\mathsf{f} - \mathsf{g}\| && \text{(ISOMETRIC)}
\end{aligned}$$

✎PROOF: These results follow directly from the fact that $\tilde{\mathbf{F}}$ is unitary (Corollary O.1 page 335) and from the properties of unitary operators (Theorem N.26 page 327). ✏

Theorem O.4 (Shift relations)**.** *Let* $\tilde{\mathbf{F}}$ *be the Fourier transform operator.*

THM

$$\begin{aligned}
\tilde{\mathbf{F}}[\mathsf{f}(x-u)](\omega) &= e^{-i\omega u}\left[\tilde{\mathbf{F}}\mathsf{f}(x)\right](\omega) \\
\left[\tilde{\mathbf{F}}\left(e^{ivx}\mathsf{g}(x)\right)\right](\omega) &= \left[\tilde{\mathbf{F}}\mathsf{g}(x)\right](\omega - v)
\end{aligned}$$

✎PROOF:

$$\begin{aligned}
\tilde{\mathbf{F}}[\mathsf{f}(x-u)](\omega) &= \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} \mathsf{f}(x-u)e^{-i\omega x}\,\mathrm{d}x && \text{by definition of } \tilde{\mathbf{F}} && \text{(Definition O.2 page 333)} \\
&= \frac{1}{\sqrt{2\pi}} \int_{v\in\mathbb{R}} \mathsf{f}(v)e^{-i\omega(u+v)}\,\mathrm{d}v && \text{where } v \triangleq x-u \implies t = u+v \\
&= e^{-i\omega u}\,\frac{1}{\sqrt{2\pi}} \int_{v\in\mathbb{R}} \mathsf{f}(v)e^{-i\omega v}\,\mathrm{d}v \\
&= e^{-i\omega u}\,\frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} \mathsf{f}(x)e^{-i\omega x}\,\mathrm{d}x && \text{by change of variable } t = v \\
&= e^{-i\omega u}\left[\tilde{\mathbf{F}}\mathsf{f}(x)\right](\omega) && \text{by definition of } \tilde{\mathbf{F}} && \text{(Definition O.2 page 333)} \\
\left[\tilde{\mathbf{F}}\left(e^{ivx}\mathsf{g}(x)\right)\right](\omega) &= \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} e^{ivx}\mathsf{g}(x)e^{-i\omega x}\,\mathrm{d}x && \text{by definition of } \tilde{\mathbf{F}} && \text{(Definition O.2 page 333)}
\end{aligned}$$

[5] *unitary operators*: Definition N.11 page 326

COPYRIGHT © 2019 DANIEL J. GREENHOE
CC BY-NC-ND

$$= \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} g(x) e^{-i(\omega - v)x} \, dx$$

$$= \left[\tilde{\mathbf{F}} g(x)\right](\omega - v) \qquad \text{by definition of } \tilde{\mathbf{F}} \qquad \text{(Definition O.2 page 333))}$$

✏

Theorem O.5 (Complex conjugate)**.** *Let $\tilde{\mathbf{F}}$ be the Fourier Transform operator and $*$ represent the complex conjugate operation on the set of complex numbers.*

THM

$$\tilde{\mathbf{F}} f^*(-x) = -\left[\tilde{\mathbf{F}} f(x)\right]^* \quad \forall f \in L^2_{(\mathbb{R},\mathscr{B},\mu)}$$

$$f \text{ is real} \implies \tilde{f}(-\omega) = \left[\tilde{f}(\omega)\right]^* \quad \forall \omega \in \mathbb{R} \qquad \text{REALITY CONDITION}$$

✎PROOF:

$$\left[\tilde{\mathbf{F}} f^*(-x)\right](\omega) \triangleq \frac{1}{\sqrt{2\pi}} \int f^*(-x) e^{-i\omega x} \, dx \qquad \text{by definition of } \tilde{\mathbf{F}} \qquad \text{(Definition O.2 page 333)}$$

$$= \frac{1}{\sqrt{2\pi}} \int f^*(u) e^{i\omega u} (-1) \, du \qquad \text{where } u \triangleq -x \implies dx = -\, du$$

$$= -\left[\frac{1}{\sqrt{2\pi}} \int f(u) e^{-i\omega u} \, du\right]^*$$

$$\triangleq -\left[\tilde{\mathbf{F}} f(x)\right]^* \qquad \text{by definition of } \tilde{\mathbf{F}} \qquad \text{(Definition O.2 page 333)}$$

$$\tilde{f}(-\omega) \triangleq \frac{1}{\sqrt{2\pi}} \int f(x) e^{-i(-\omega)x} \, dx \qquad \text{by definition of } \tilde{\mathbf{F}} \qquad \text{(Definition O.2 page 333)}$$

$$= \left[\frac{1}{\sqrt{2\pi}} \int f^*(x) e^{-i\omega x} \, dx\right]^*$$

$$= \left[\frac{1}{\sqrt{2\pi}} \int f(x) e^{-i\omega x} \, dx\right]^* \qquad \text{by } f \text{ is real hypothesis}$$

$$\triangleq \tilde{f}^*(\omega) \qquad \text{by definition of } \tilde{\mathbf{F}} \qquad \text{(Definition O.2 page 333)}$$

✏

O.3 Convolution

Definition O.3. [6]

DEF

*The **convolution operation** is defined as*

$$\left[f \star g\right](x) \triangleq f(x) \star g(x) \triangleq \int_{u \in \mathbb{R}} f(u) g(x-u) \, du \qquad \forall f,g \in L^2_{(\mathbb{R},\mathscr{B},\mu)}$$

Theorem Q.2 (next) demonstrates that multiplication in the "time domain" is equivalent to convolution in the "frequency domain" and vice-versa.

Theorem O.6 (convolution theorem)**.** [7] *Let $\tilde{\mathbf{F}}$ be the Fourier Transform operator and $\star$ the convolution operator.*

[6] Bachman (1964), page 6, Bracewell (1978) page 108 ⟨Convolution theorem⟩
[7] Bracewell (1978) page 110

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

THM

$$\underbrace{\tilde{\mathbf{F}}\big[\mathsf{f}(x)\star\mathsf{g}(x)\big](\omega)}_{\textit{convolution in "time domain"}} = \underbrace{\sqrt{2\pi}\big[\tilde{\mathbf{F}}\mathsf{f}\big](\omega)\,\big[\tilde{\mathbf{F}}\mathsf{g}\big](\omega)}_{\textit{multiplication in "frequency domain"}} \qquad \forall \mathsf{f},\mathsf{g}\in L^2_{(\mathbb{R},\mathscr{B},\mu)}$$

$$\underbrace{\tilde{\mathbf{F}}\big[\mathsf{f}(x)\mathsf{g}(x)\big](\omega)}_{\textit{multiplication in "time domain"}} = \underbrace{\frac{1}{\sqrt{2\pi}}\big[\tilde{\mathbf{F}}\mathsf{f}\big](\omega)\star\big[\tilde{\mathbf{F}}\mathsf{g}\big](\omega)}_{\textit{convolution in "frequency domain"}} \qquad \forall \mathsf{f},\mathsf{g}\in L^2_{(\mathbb{R},\mathscr{B},\mu)}.$$

✎PROOF:

$$\begin{aligned}
\tilde{\mathbf{F}}\big[\mathsf{f}(x)\star\mathsf{g}(x)\big](\omega) &= \tilde{\mathbf{F}}\left[\int_{u\in\mathbb{R}}\mathsf{f}(u)\mathsf{g}(x-u)\,\mathrm{d}u\right](\omega) && \text{by definition of } \star \text{ (Definition O.3 page 336)}\\
&= \int_{u\in\mathbb{R}}\mathsf{f}(u)\big[\tilde{\mathbf{F}}\mathsf{g}(x-u)\big](\omega)\,\mathrm{d}u\\
&= \int_{u\in\mathbb{R}}\mathsf{f}(u)e^{-i\omega u}\,\big[\tilde{\mathbf{F}}\mathsf{g}(x)\big](\omega)\,\mathrm{d}u && \text{by Theorem O.4 page 335}\\
&= \sqrt{2\pi}\underbrace{\left(\frac{1}{\sqrt{2\pi}}\int_{u\in\mathbb{R}}\mathsf{f}(u)e^{-i\omega u}\,\mathrm{d}u\right)}_{[\tilde{\mathbf{F}}\mathsf{f}](\omega)}\big[\tilde{\mathbf{F}}\mathsf{g}\big](\omega)\\
&= \sqrt{2\pi}\big[\tilde{\mathbf{F}}\mathsf{f}\big](\omega)\,\big[\tilde{\mathbf{F}}\mathsf{g}\big](\omega) && \text{by definition of } \tilde{\mathbf{F}} \text{ (Definition O.2 page 333)}\\
\tilde{\mathbf{F}}[\mathsf{f}(x)\mathsf{g}(x)](\omega) &= \tilde{\mathbf{F}}\big[\big(\tilde{\mathbf{F}}^{-1}\tilde{\mathbf{F}}\mathsf{f}(x)\big)\,\mathsf{g}(x)\big](\omega) && \text{by definition of operator inverse (page 303)}\\
&= \tilde{\mathbf{F}}\left[\left(\frac{1}{\sqrt{2\pi}}\int_{v\in\mathbb{R}}\big[\tilde{\mathbf{F}}\mathsf{f}(x)\big](v)e^{ivx}\,\mathrm{d}v\right)\mathsf{g}(x)\right](\omega) && \text{by Theorem O.1 page 334}\\
&= \frac{1}{\sqrt{2\pi}}\int_{v\in\mathbb{R}}\big[\tilde{\mathbf{F}}\mathsf{f}(x)\big](v)\big[\tilde{\mathbf{F}}\big(e^{ivx}\,\mathsf{g}(x)\big)\big](\omega,v)\,\mathrm{d}v\\
&= \frac{1}{\sqrt{2\pi}}\int_{v\in\mathbb{R}}\big[\tilde{\mathbf{F}}\mathsf{f}(x)\big](v)\big[\tilde{\mathbf{F}}\mathsf{g}(x)\big](\omega-v)\,\mathrm{d}v && \text{by Theorem O.4 page 335}\\
&= \frac{1}{\sqrt{2\pi}}\big[\tilde{\mathbf{F}}\mathsf{f}\big](\omega)\star\big[\tilde{\mathbf{F}}\mathsf{g}\big](\omega) && \text{by definition of } \star \text{ (Definition O.3 page 336)}
\end{aligned}$$

✏

O.4 Real valued functions



Figure O.1: Fourier transform components of real-valued signal

Theorem O.7. *Let* $\mathsf{f}(x)$ *be a function in* $L^2_{\mathbb{R}}$ *and* $\tilde{\mathsf{f}}(\omega)$ *the* FOURIER TRANSFORM *of* $\mathsf{f}(x)$.

THM

$$\left\{\begin{array}{l}\mathsf{f}(x)\ \textit{is}\ \text{REAL-VALUED}\\ (\mathsf{f}\in\mathbb{R}^{\mathbb{R}})\end{array}\right\} \Longrightarrow \left\{\begin{array}{rcll}\tilde{\mathsf{f}}(\omega) &=& \tilde{\mathsf{f}}^*(-\omega) & (\text{HERMITIAN SYMMETRIC})\\ \mathbf{R}_e\left[\tilde{\mathsf{f}}(\omega)\right] &=& \mathbf{R}_e\left[\tilde{\mathsf{f}}(-\omega)\right] & (\text{SYMMETRIC})\\ \mathbf{I}_m\left[\tilde{\mathsf{f}}(\omega)\right] &=& -\mathbf{I}_m\left[\tilde{\mathsf{f}}(-\omega)\right] & (\text{ANTI-SYMMETRIC})\\ |\tilde{\mathsf{f}}(\omega)| &=& |\tilde{\mathsf{f}}(-\omega)| & (\text{SYMMETRIC})\\ \angle\tilde{\mathsf{f}}(\omega) &=& \angle\tilde{\mathsf{f}}(-\omega) & (\text{ANTI-SYMMETRIC}).\end{array}\right\}$$

COPYRIGHT © 2019 DANIEL J. GREENHOE
CC BY-NC-ND

✎PROOF:

$$\begin{array}{lllllll}
\tilde{f}(\omega) & \triangleq & [\tilde{\mathbf{F}}f(x)](\omega) & \triangleq & \left\langle f(x) \,|\, e^{i\omega x}\right\rangle & = & \left\langle f(x) \,|\, e^{i(-\omega)x}\right\rangle^* \triangleq \tilde{f}^*(-\omega) \\
\mathbf{R}_e\left[\tilde{f}(\omega)\right] & = & \mathbf{R}_e\left[\tilde{f}^*(-\omega)\right] & = & \mathbf{R}_e\left[\tilde{f}(-\omega)\right] & & \\
\mathbf{I}_m\left[\tilde{f}(\omega)\right] & = & \mathbf{I}_m\left[\tilde{f}^*(-\omega)\right] & = & -\mathbf{I}_m\left[\tilde{f}(-\omega)\right] & & \\
|\tilde{f}(\omega)| & = & |\tilde{f}^*(-\omega)| & = & |\tilde{f}(-\omega)| & & \\
\angle\tilde{f}(\omega) & = & \angle\tilde{f}^*(-\omega) & = & -\angle\tilde{f}(-\omega) & &
\end{array}$$

✏

O.5 Moment properties

Definition O.4. [8]

DEF *The quantity* M_n *is the* n***th moment*** *of a function* $f(x) \in L^2_{\mathbb{R}}$ *if*

$$\mathsf{M}_n \triangleq \int_{\mathbb{R}} x^n f(x)\,dx \qquad \text{for } n \in \mathbb{W}.$$

Lemma O.1. [9] *Let* M_n *be the* nTH MOMENT *(Definition O.4 page 338) and* $\tilde{f}(\omega) \triangleq \left[\tilde{\mathbf{F}}f\right](\omega)$ *the* FOURIER TRANSFORM *(Definition O.2 page 333) of a function* $f(x)$ *in* $L^2_{\mathbb{R}}$ *(Definition ?? page ??).*

LEM

$$\begin{array}{rcll}
\mathsf{M}_n & = & \sqrt{2\pi}(i)^n\left[\frac{d}{d\omega}\right]^n \tilde{f}(\omega)\Big|_{\omega=0} & \forall n\in\mathbb{W}, f\in L^2_{\mathbb{R}} \\
\left[\frac{d}{d\omega}\right]^n \tilde{f}(\omega)\Big|_{\omega=0} & = & \frac{1}{\sqrt{2\pi}}(-i)^n\mathsf{M}_n & \forall n\in\mathbb{W}, f\in L^2_{\mathbb{R}}
\end{array}$$

✎PROOF:

$$\begin{aligned}
\sqrt{2\pi}(i)^n\left[\left[\frac{d}{d\omega}\right]^n \tilde{f}(\omega)\right]_{\omega=0} &= \sqrt{2\pi}(i)^n\left[\left[\frac{d}{d\omega}\right]^n \frac{1}{\sqrt{2\pi}}\int_{\mathbb{R}} f(x)e^{-i\omega x}\,dx\right]_{\omega=0} && \text{by definition of } \tilde{\mathbf{F}} \text{ (Definition O.2 page 333)} \\
&= (i)^n\int_{\mathbb{R}} f(x)\left[\left[\frac{d}{d\omega}\right]^n e^{-i\omega x}\right]dx\Big|_{\omega=0} \\
&= (i)^n\int_{\mathbb{R}} f(x)\left[(-i)^n x^n e^{-i\omega x}\right]dx\Big|_{\omega=0} \\
&= (-i^2)^n\int_{\mathbb{R}} f(x)x^n\,dx \\
&= \int_{\mathbb{R}} f(x)x^n\,dx \\
&\triangleq \mathsf{M}_n && \text{by definition of } \mathsf{M}_n \text{ (Definition O.4 page 338)}
\end{aligned}$$

✏

Lemma O.2. [10] *Let* M_n *be the* nTH MOMENT *(Definition O.4 page 338) and* $\tilde{f}(\omega) \triangleq \left[\tilde{\mathbf{F}}f\right](\omega)$ *the* FOURIER TRANSFORM *(Definition O.2 page 333) of a function* $f(x)$ *in* $L^2_{\mathbb{R}}$ *(Definition ?? page ??).*

LEM

$$\mathsf{M}_n = 0 \quad\Longleftrightarrow\quad \left[\frac{d}{d\omega}\right]^n \tilde{f}(\omega)\Big|_{\omega=0} = 0 \qquad \forall n\in\mathbb{W}$$

✎PROOF:

[8] Jawerth and Sweldens (1994), pages 16–17, Sweldens and Piessens (1993), page 2, Vidakovic (1999), page 83

[9] Goswami and Chan (1999), pages 38–39

[10] Vidakovic (1999), pages 82–83, Mallat (1999), pages 241–242

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

1. Proof for ($\Longrightarrow$) case:

$$\begin{aligned}
0 &= \langle \mathsf{f}(x) \mid x^n \rangle && \text{by left hypothesis}\\
&= \sqrt{2\pi}(-i)^{-n}\left[\frac{\mathsf{d}}{\mathsf{d}\omega}\right]^n \tilde{\mathsf{f}}(\omega)\Big|_{\omega=0} && \text{by Lemma O.1 page 338}\\
\Longrightarrow\ & \left[\frac{\mathsf{d}}{\mathsf{d}\omega}\right]^n \tilde{\mathsf{f}}(\omega)\Big|_{\omega=0} = 0
\end{aligned}$$

2. Proof for ($\Longleftarrow$) case:

$$\begin{aligned}
0 &= \left[\frac{\mathsf{d}}{\mathsf{d}\omega}\right]^n \tilde{\mathsf{f}}(\omega)\Big|_{\omega=0} && \text{by right hypothesis}\\
&= \left[\frac{\mathsf{d}}{\mathsf{d}\omega}\right]^n \frac{1}{\sqrt{2\pi}}\int_{\mathbb{R}} \mathsf{f}(x)e^{-i\omega x}\,\mathsf{d}x\Big|_{\omega=0} && \text{by definition of } \tilde{\mathsf{f}}(\omega)\\
&= \frac{1}{\sqrt{2\pi}}\int_{\mathbb{R}} \mathsf{f}(x)\left[\frac{\mathsf{d}}{\mathsf{d}\omega}\right]^n e^{-i\omega x}\,\mathsf{d}x\Big|_{\omega=0}\\
&= \frac{1}{\sqrt{2\pi}}\int_{\mathbb{R}} \mathsf{f}(x)\left[(-i)^n x^n e^{-i\omega x}\right]\,\mathsf{d}x\Big|_{\omega=0}\\
&= (-i)^n\frac{1}{\sqrt{2\pi}}\int_{\mathbb{R}} \mathsf{f}(x)x^n\,\mathsf{d}x\\
&= (-i)^n\frac{1}{\sqrt{2\pi}}\langle \mathsf{f}(x) \mid x^n\rangle && \text{by definition of } \langle\cdot\mid\cdot\rangle \text{ in } \boldsymbol{L}^2_{\mathbb{R}} \text{ (Definition ?? page ??)}
\end{aligned}$$

✏

Lemma O.3 (Strang-Fix condition). [11] *Let* $\mathsf{f}(x)$ *be a function in* $\boldsymbol{L}^2_{\mathbb{R}}$ *and* M_n *the* nTH MOMENT *(Definition O.4 page 338) of* $\mathsf{f}(x)$. *Let* $\mathbf{T}$ *be the* TRANSLATION OPERATOR *(Definition ?? page ??).*

LEM

$$\underbrace{\sum_{k\in\mathbb{Z}} \mathbf{T}^k x^n \mathsf{f}(x) = \mathsf{M}_n}_{\text{STRANG-FIX CONDITION in “time”}} \quad\Longleftrightarrow\quad \underbrace{\left[\frac{\mathsf{d}}{\mathsf{d}\omega}\right]^n \tilde{\mathsf{f}}(\omega)\Big|_{\omega=2\pi k} = \frac{1}{\sqrt{2\pi}}(-i)^n\bar{\delta}_k\mathsf{M}_n}_{\text{STRANG-FIX CONDITION in “frequency”}}$$

✎PROOF:

1. Proof for ($\Longrightarrow$) case:

$$\begin{aligned}
\left[\left[\frac{\mathsf{d}}{\mathsf{d}\omega}\right]^n \tilde{\mathsf{f}}(\omega)\right]_{\omega=2\pi k} &= \sum_{k\in\mathbb{Z}}\left[\left[\frac{\mathsf{d}}{\mathsf{d}\omega}\right]^n \tilde{\mathsf{f}}(\omega)\right]_{\omega=2\pi k} e^{i2\pi kx}\bar{\delta}_k\\
&= \frac{1}{\sqrt{2\pi}}\sum_{k\in\mathbb{Z}}\left[\left[\frac{\mathsf{d}}{\mathsf{d}\omega}\right]^n \int_{\mathbb{R}} \mathsf{f}(x)e^{-i\omega x}\,\mathsf{d}x\right]_{\omega=2\pi k} e^{i2\pi kx}\bar{\delta}_k && \text{by Definition O.2 page 333}\\
&= \frac{1}{\sqrt{2\pi}}\sum_{k\in\mathbb{Z}}\left[\int_{\mathbb{R}} \mathsf{f}(x)(-ix)^n e^{-i\omega x}\,\mathsf{d}x\right]_{\omega=2\pi k} e^{i2\pi kx}\bar{\delta}_k\\
&= (-i)^n\frac{1}{\sqrt{2\pi}}\sum_{k\in\mathbb{Z}}\left[\int_{\mathbb{R}} x^n\mathsf{f}(x)e^{-i\omega x}\,\mathsf{d}x\right]_{\omega=2\pi k} e^{i2\pi kx}\bar{\delta}_k\\
&= (-i)^n\frac{1}{\sqrt{2\pi}}\sum_{k\in\mathbb{Z}}(x-k)^n\mathsf{f}(x-k)\bar{\delta}_k && \text{by } PSF \text{ (Theorem ?? page ??)}\\
&= \frac{1}{\sqrt{2\pi}}(-i)^n\bar{\delta}_k\mathsf{M}_n && \text{by left hypothesis}
\end{aligned}$$

[11] Jawerth and Sweldens (1994), pages 16–17, Sweldens and Piessens (1993), page 2, Vidakovic (1999), page 83, Mallat (1999), pages 241–243, Fix and Strang (1969)

COPYRIGHT © 2019 DANIEL J. GREENHOE

2. Proof for ($\Longleftarrow$) case:

$$\begin{aligned}
\frac{1}{\sqrt{2\pi}}(-i)^n \mathsf{M}_n &= \frac{1}{\sqrt{2\pi}} \sum_{k\in\mathbb{Z}} \left[(-i)^n \bar{\delta}_k \mathsf{M}_n\right] e^{-i2\pi kx} && \text{by definition of } \bar{\delta} && \text{(Definition J.3 page 267)}\\
&= \sum_{k\in\mathbb{Z}} \left[\left[\frac{\mathsf{d}}{\mathsf{d}\omega}\right]^n \tilde{\mathsf{f}}(\omega)\right]\Bigg|_{\omega=2\pi k} e^{-i2\pi kx} && \text{by right hypothesis}\\
&= \sum_{k\in\mathbb{Z}} \left[\left[\frac{\mathsf{d}}{\mathsf{d}\omega}\right]^n \int_{\mathbb{R}} \mathsf{f}(x)e^{-i\omega x}\,\mathsf{d}x\right]\Bigg|_{\omega=2\pi k} e^{-i2\pi kx}\\
&= \sum_{k\in\mathbb{Z}} \left[\int_{\mathbb{R}} \mathsf{f}(x)(-ix)^n e^{-i\omega x}\,\mathsf{d}x\right]\Bigg|_{\omega=2\pi k} e^{-i2\pi kx}\\
&= (-i)^n \sum_{k\in\mathbb{Z}} \left[\int_{\mathbb{R}} x^n \mathsf{f}(x)e^{-i\omega x}\,\mathsf{d}x\right]\Bigg|_{\omega=2\pi k} e^{-i2\pi kx}\\
&= (-i)^n \sum_{k\in\mathbb{Z}} (x-k)^n \mathsf{f}(x-k) && \text{by } PSF && \text{(Theorem ?? page ??)}
\end{aligned}$$

✏

O.6 Examples

Example O.1 (rectangular pulse). Let $\tilde{\mathsf{f}}(\omega)$ be the *Fourier transform* of a function $\mathsf{f}(x) \in \boldsymbol{L}^2_{\mathbb{R}}$.

EX

$$\mathsf{f}(x) = \begin{cases} c & \text{for } x \in [a : b)\\ 0 & \text{otherwise}\end{cases} \qquad \tilde{\mathsf{f}}(\omega) = \frac{c(b-a)}{\sqrt{2\pi}} e^{-i\left(\frac{a+b}{2}\omega\right)} \left[\frac{\sin\left(\frac{b-a}{2}\omega\right)}{\left(\frac{b-a}{2}\omega\right)}\right]$$





✎ PROOF:

$$\begin{aligned}
\tilde{\mathsf{f}}(\omega) &= \tilde{\mathbf{F}}[\mathsf{f}(x)](\omega) && \text{by definition of } \tilde{\mathsf{f}}(\omega)\\
&= e^{-i\left(\frac{a+b}{2}\right)\omega} \tilde{\mathbf{F}}\left[\mathsf{f}\left(x - \frac{a+b}{2}\right)\right](\omega) && \text{by } \textit{shift relation} && \text{(Theorem O.4 page 335)}\\
&= e^{-i\left(\frac{a+b}{2}\right)\omega} \tilde{\mathbf{F}}\left[c\mathbb{1}_{[a:b)}\left(x - \frac{a+b}{2}\right)\right](\omega) && \text{by definition of } \mathsf{f}(x)\\
&= e^{-i\left(\frac{a+b}{2}\right)\omega} \tilde{\mathbf{F}}\left[c\mathbb{1}_{\left[-\frac{b-a}{2}:\frac{b-a}{2}\right)}(x)\right](\omega) && \text{by definition of } \mathbb{1} && \text{(Definition ?? page ??)}\\
&= \frac{1}{\sqrt{2\pi}} e^{-i\left(\frac{a+b}{2}\right)\omega} \int_{\mathbb{R}} c\mathbb{1}_{\left[-\frac{b-a}{2}:\frac{b-a}{2}\right)}(x) e^{-i\omega x}\,\mathsf{d}x && \text{by definition of } \tilde{\mathbf{F}} && \text{(Definition O.2 page 333)}\\
&= \frac{1}{\sqrt{2\pi}} e^{-i\left(\frac{a+b}{2}\right)\omega} \int_{-\frac{b-a}{2}}^{\frac{b-a}{2}} c e^{-i\omega x}\,\mathsf{d}x && \text{by definition of } \mathbb{1} && \text{(Definition ?? page ??)}\\
&= \frac{c}{\sqrt{2\pi}} e^{-i\left(\frac{a+b}{2}\right)\omega} \frac{1}{-i\omega} e^{-i\omega x}\Bigg|_{-\frac{b-a}{2}}^{\frac{b-a}{2}}\\
&= \frac{2c}{\sqrt{2\pi}\omega} e^{-i\left(\frac{a+b}{2}\right)\omega} \left[\frac{e^{i\left(\frac{b-a}{2}\omega\right)} - e^{-i\left(\frac{b-a}{2}\omega\right)}}{2i}\right]
\end{aligned}$$

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

$$= \frac{c(b-a)}{\sqrt{2\pi}} e^{-i\left(\frac{a+b}{2}\omega\right)} \left[\frac{\sin\left(\frac{b-a}{2}\omega\right)}{\left(\frac{b-a}{2}\omega\right)} \right] \qquad \text{by } \textit{Euler formulas} \qquad \text{(Corollary ?? page ??)}$$

✏

Example O.2 (triangle). Let $\tilde{f}(\omega)$ be the *Fourier transform* of a function $f(x) \in L^2_{\mathbb{R}}$.

EX

$$f(x) = \begin{cases} c\left[1 - \frac{|2x-b-a|}{b-a}\right] & \text{for } x \in [a : b) \\ 0 & \text{otherwise} \end{cases} \qquad \tilde{f}(\omega) = \frac{c(b-a)^2}{4\sqrt{2\pi}} e^{-i\left(\frac{a+b}{2}\omega\right)} \left[\frac{\sin\left(\frac{b-a}{4}\omega\right)}{\left(\frac{b-a}{4}\omega\right)} \right]^2$$



✎PROOF:

$$\begin{aligned}
\tilde{f}(\omega) &= \tilde{\mathbf{F}}[f(x)](\omega) && \text{by definition of } \tilde{f}(\omega) \\
&= e^{-i\left(\frac{a+b}{2}\right)\omega} \tilde{\mathbf{F}}\left[f\left(x - \frac{a+b}{2}\right)\right](\omega) && \text{by } \textit{shift relation} \qquad \text{(Theorem O.4 page 335)} \\
&= \tilde{\mathbf{F}}\left[c\left(1 - \frac{|2x-b-a|}{b-a}\right)\mathbb{1}_{[a:b)}(x)\right](\omega) && \text{by definition of } f(x) \\
&= c\tilde{\mathbf{F}}\left[\mathbb{1}_{\left[\frac{a}{2}:\frac{b}{2}\right)}(x) \star \mathbb{1}_{\left[\frac{a}{2}:\frac{b}{2}\right)}(x)\right](\omega) \\
&= c\sqrt{2\pi}\tilde{\mathbf{F}}\left[\mathbb{1}_{\left[\frac{a}{2}:\frac{b}{2}\right)}\right]\tilde{\mathbf{F}}\left[\mathbb{1}_{\left[\frac{a}{2}:\frac{b}{2}\right)}\right] && \text{by } \textit{convolution theorem} \qquad \text{(Theorem Q.2 page 356)} \\
&= c\sqrt{2\pi}\left(\tilde{\mathbf{F}}\left[\mathbb{1}_{\left[\frac{a}{2}:\frac{b}{2}\right)}\right]\right)^2 \\
&= c\sqrt{2\pi}\left(\frac{\left(\frac{b}{2}-\frac{a}{2}\right)}{\sqrt{2\pi}} e^{-i\left(\frac{a+b}{4}\omega\right)} \left[\frac{\sin\left(\frac{b-a}{4}\omega\right)}{\left(\frac{b-a}{4}\omega\right)}\right]\right)^2 && \text{by } \textit{Rectangular pulse} \text{ ex.} \qquad \text{Example O.1 page 340} \\
&= \frac{c(b-a)^2}{4\sqrt{2\pi}} e^{-i\left(\frac{a+b}{2}\omega\right)} \left[\frac{\sin\left(\frac{b-a}{4}\omega\right)}{\left(\frac{b-a}{4}\omega\right)}\right]^2
\end{aligned}$$

✏

Example O.3. Let a function f be defined in terms of the cosine function (Definition ?? page ??) as follows:

EX

$$f(x) \triangleq \begin{cases} \cos^2\left(\frac{\pi}{2}x\right) & \text{for } |x| \le 1 \\ 0 & \text{otherwise} \end{cases} \qquad \tilde{f}(\omega) = \frac{1}{2\sqrt{2\pi}}\left[\underbrace{\frac{2\sin\omega}{\omega}}_{2\,\mathrm{sinc}(\omega)} + \underbrace{\frac{\sin(\omega+\pi)}{(\omega+\pi)}}_{\mathrm{sinc}(\omega+\pi)} + \underbrace{\frac{\sin(\omega-\pi)}{(\omega-\pi)}}_{\mathrm{sinc}(\omega-\pi)}\right]$$



COPYRIGHT © 2019 DANIEL J. GREENHOE

✎PROOF: Let $\mathbb{1}_A(x)$ be the *set indicator function* (Definition ?? page ??) on a set A.

$$\begin{aligned}
\tilde{\mathsf{f}}(\omega) &\triangleq \frac{1}{\sqrt{2\pi}}\int_{\mathbb{R}} \mathsf{f}(x)e^{-i\omega x}\,\mathrm{d}x && \text{by definition of } \tilde{\mathsf{f}}(\omega) \text{ (Definition O.2)}\\
&= \frac{1}{\sqrt{2\pi}}\int_{\mathbb{R}} \cos^2\left(\frac{\pi}{2}x\right)\mathbb{1}_{[-1:1]}(x)e^{-i\omega x}\,\mathrm{d}x && \text{by definition of } \mathsf{f}(x)\\
&= \frac{1}{\sqrt{2\pi}}\int_{-1}^{1} \cos^2\left(\frac{\pi}{2}x\right)e^{-i\omega x}\,\mathrm{d}x && \text{by definition of } \mathbb{1} \text{ (Definition ??)}\\
&= \frac{1}{\sqrt{2\pi}}\int_{-1}^{1} \left[\frac{e^{i\frac{\pi}{2}x}+e^{-i\frac{\pi}{2}x}}{2}\right]^2 e^{-i\omega x}\,\mathrm{d}x && \text{by Corollary ?? page ??}\\
&= \frac{1}{4\sqrt{2\pi}}\int_{-1}^{1} \left[2+e^{i\pi x}+e^{-i\pi x}\right]e^{-i\omega x}\,\mathrm{d}x\\
&= \frac{1}{4\sqrt{2\pi}}\int_{-1}^{1} 2e^{-i\omega x}+e^{-i(\omega+\pi)x}+e^{-i(\omega-\pi)x}\,\mathrm{d}x\\
&= \frac{1}{4\sqrt{2\pi}}\left[2\frac{e^{-i\omega x}}{-i\omega}+\frac{e^{-i(\omega+\pi)x}}{-i(\omega+\pi)}+\frac{e^{-i(\omega-\pi)x}}{-i(\omega-\pi)}\right]_{-1}^{1}\\
&= \frac{1}{2\sqrt{2\pi}}\left[2\frac{e^{-i\omega}-e^{+i\omega}}{-2i\omega}+\frac{e^{-i(\omega+\pi)}-e^{+i(\omega+\pi)}}{-2i(\omega+\pi)}+\frac{e^{-i(\omega-\pi)}-e^{+i(\omega-\pi)}}{-2i(\omega-\pi)}\right]_{-1}^{1}\\
&= \frac{1}{2\sqrt{2\pi}}\left[\underbrace{\frac{2\sin\omega}{\omega}}_{2\,\mathrm{sinc}(\omega)}+\underbrace{\frac{\sin(\omega+\pi)}{(\omega+\pi)}}_{\mathrm{sinc}(\omega+\pi)}+\underbrace{\frac{\sin(\omega-\pi)}{(\omega-\pi)}}_{\mathrm{sinc}(\omega-\pi)}\right]
\end{aligned}$$

✏

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

APPENDIX P

DISCRETE TIME FOURIER TRANSFORM

P.1 Definition

Definition P.1.

DEF *The **discrete-time Fourier transform** $\breve{\mathbf{F}}$ of $(\!(x_n)\!)_{n\in\mathbb{Z}}$ is defined as*

$$\left[\breve{\mathbf{F}}\,(\!(x_n)\!)\right](\omega) \triangleq \sum_{n\in\mathbb{Z}} x_n e^{-i\omega n} \qquad \forall (\!(x_n)\!)_{n\in\mathbb{Z}} \in \ell^2_{\mathbb{R}}$$

If we compare the definition of the *Discrete Time Fourier Transform* (Definition P.1 page 343) to the definition of the Z-transform (Definition Q.4 page 354), we see that the DTFT is just a special case of the more general Z-Transform, with $z = e^{i\omega}$. If we imagine $z \in \mathbb{C}$ as a complex plane, then $e^{i\omega}$ is a unit circle in this plane. The "frequency" ω in the DTFT is the unit circle in the much larger z-plane, as illustrated to the right.



P.2 Properties

Proposition P.1 (DTFT periodicity)**.** *Let $\breve{x}(\omega) \triangleq \breve{\mathbf{F}}\left[(\!(x_n)\!)\right](\omega)$ be the* DISCRETE-TIME FOURIER TRANSFORM *(Definition P.1 page 343) of a sequence $(\!(x_n)\!)_{n\in\mathbb{Z}}$ in $\ell^2_{\mathbb{R}}$.*

PRP

$$\underbrace{\breve{x}(\omega) = \breve{x}(\omega + 2\pi n)}_{\text{PERIODIC } \textit{with period } 2\pi} \qquad \forall n \in \mathbb{Z}$$

✎PROOF:

$$\begin{aligned}
\breve{x}(\omega + 2\pi n) &= \sum_{m\in\mathbb{Z}} x_m e^{-i(\omega+2\pi n)m} &&= \sum_{m\in\mathbb{Z}} x_m e^{-i\omega m} \cancelto{1}{e^{-i2\pi n m}} \\
&= \sum_{m\in\mathbb{Z}} x_m e^{-i\omega m} &&= \breve{x}(\omega)
\end{aligned}$$

⇨

Theorem P.1. *Let* $\breve{\mathsf{x}}(\omega) \triangleq \breve{\mathbf{F}}\big[(\!|\mathsf{x}[n]|\!)\big](\omega)$ *be the* DISCRETE-TIME FOURIER TRANSFORM *(Definition P.1 page 343) of a sequence* $(\!|x_n|\!)_{n\in\mathbb{Z}}$ *in* $\ell^2_{\mathbb{R}}$.

THM
$$\left\{\ \tilde{\mathsf{x}}(\omega) \triangleq \breve{\mathbf{F}}\left(\!|\mathsf{x}[n]|\!\right)\ \right\} \implies \left\{\begin{array}{llll} (1). & \breve{\mathbf{F}}\left(\!|\mathsf{x}[-n]|\!\right) & = \tilde{\mathsf{x}}(-\omega) & \textit{and} \\ (2). & \breve{\mathbf{F}}\left(\!|\mathsf{x}^*[n]|\!\right) & = \tilde{\mathsf{x}}^*(-\omega) & \textit{and} \\ (3). & \breve{\mathbf{F}}\left(\!|\mathsf{x}^*[-n]|\!\right) & = \tilde{\mathsf{x}}^*(\omega) & \end{array}\right\}$$

✎PROOF:

$$\begin{aligned}
\breve{\mathbf{F}}\left(\!|\mathsf{x}[-n]|\!\right) &\triangleq \sum_{n\in\mathbb{Z}} \mathsf{x}[-n]e^{-i\omega n} && \text{by definition of } DTFT && \text{(Definition P.1 page 343)}\\
&= \sum_{m\in\mathbb{Z}} \mathsf{x}[m]e^{i\omega m} && \text{where } m \triangleq -n \implies n = -m \\
&= \sum_{m\in\mathbb{Z}} \mathsf{x}[m]e^{-i(-\omega)m} \\
&\triangleq \tilde{\mathsf{x}}(-\omega) && \text{by left hypothesis}
\end{aligned}$$

$$\begin{aligned}
\breve{\mathbf{F}}\left(\!|\mathsf{x}^*[n]|\!\right) &\triangleq \sum_{n\in\mathbb{Z}} \mathsf{x}^*[n]e^{-i\omega n} && \text{by definition of } DTFT && \text{(Definition P.1 page 343)}\\
&= \left(\sum_{n\in\mathbb{Z}} \mathsf{x}[n]e^{i\omega n}\right)^* && \text{by } \textit{distributive} \text{ property of } *\textbf{-algebras} && \text{(Definition I.3 page 250)}\\
&= \left(\sum_{n\in\mathbb{Z}} \mathsf{x}[n]e^{-i(-\omega)n}\right)^* \\
&\triangleq \tilde{\mathsf{x}}^*(-\omega) && \text{by left hypothesis}
\end{aligned}$$

$$\begin{aligned}
\breve{\mathbf{F}}\left(\!|\mathsf{x}^*[-n]|\!\right) &\triangleq \sum_{n\in\mathbb{Z}} \mathsf{x}^*[-n]e^{-i\omega n} && \text{by definition of } DTFT && \text{(Definition P.1 page 343)}\\
&= \left(\sum_{n\in\mathbb{Z}} \mathsf{x}[-n]e^{i\omega n}\right)^* && \text{by } \textit{distributive} \text{ property of } *\textbf{-algebras} && \text{(Definition I.3 page 250)}\\
&= \left(\sum_{m\in\mathbb{Z}} \mathsf{x}[m]e^{-i\omega m}\right)^* && \text{where } m \triangleq -n \implies n = -m \\
&\triangleq \tilde{\mathsf{x}}^*(\omega) && \text{by left hypothesis}
\end{aligned}$$

✏

Theorem P.2. *Let* $\breve{\mathsf{x}}(\omega) \triangleq \breve{\mathbf{F}}\big[(\!|\mathsf{x}[n]|\!)\big](\omega)$ *be the* DISCRETE-TIME FOURIER TRANSFORM *(Definition P.1 page 343) of a sequence* $(\!|\mathsf{x}[n]|\!)_{n\in\mathbb{Z}}$ *in* $\ell^2_{\mathbb{R}}$.

THM
$$\left\{\begin{array}{lll} (1). & \tilde{\mathsf{x}}(\omega) \triangleq \breve{\mathbf{F}}\left(\!|\mathsf{x}[n]|\!\right) & \textit{and} \\ (2). & (\!|\mathsf{x}[n]|\!) \textit{ is } \text{REAL-VALUED} & \end{array}\right\} \implies \left\{\begin{array}{llllll} (1). & \breve{\mathbf{F}}\left(\!|\mathsf{x}[-n]|\!\right) & = & \tilde{\mathsf{x}}(-\omega) & & \textit{and} \\ (2). & \breve{\mathbf{F}}\left(\!|\mathsf{x}^*[n]|\!\right) & = & \tilde{\mathsf{x}}^*(-\omega) & = \tilde{\mathsf{x}}(\omega) & \textit{and} \\ (3). & \breve{\mathbf{F}}\left(\!|\mathsf{x}^*[-n]|\!\right) & = & \tilde{\mathsf{x}}^*(\omega) & = \tilde{\mathsf{x}}(-\omega) & \end{array}\right\}$$

✎PROOF:

$$\begin{aligned}
\breve{\mathbf{F}}\left(\!|\mathsf{x}[-n]|\!\right) &\triangleq \sum_{n\in\mathbb{Z}} \mathsf{x}[-n]e^{-i\omega n} && \text{by definition of } DTFT && \text{(Definition P.1 page 343)}\\
&= \sum_{m\in\mathbb{Z}} \mathsf{x}[m]e^{i\omega m} && \text{where } m \triangleq -n \implies n = -m \\
&= \sum_{m\in\mathbb{Z}} \mathsf{x}[m]e^{-i(-\omega)m}
\end{aligned}$$

COPYRIGHT © 2019 DANIEL J. GREENHOE

$$\triangleq \tilde{\mathsf{x}}(-\omega) \qquad \text{by left hypothesis}$$

$$\begin{aligned}
\boxed{\tilde{\mathsf{x}}^*(-\omega)} &= \boxed{\breve{\mathbf{F}}\left(\!\left(\mathsf{x}^*[n]\right)\!\right)} && \text{by Theorem P.1 page 344}\\
&= \breve{\mathbf{F}}\left(\!\left(\mathsf{x}[n]\right)\!\right) && \text{by \textit{real-valued} hypothesis}\\
&= \boxed{\tilde{\mathsf{x}}(\omega)} && \text{by definition of } \tilde{\mathsf{x}}(\omega) \qquad \text{(Definition P.1 page 343)}
\end{aligned}$$

$$\begin{aligned}
\boxed{\tilde{\mathsf{x}}^*(\omega)} &= \boxed{\breve{\mathbf{F}}\left(\!\left(\mathsf{x}^*[-n]\right)\!\right)} && \text{by Theorem P.1 page 344}\\
&= \breve{\mathbf{F}}\left(\!\left(\mathsf{x}[-n]\right)\!\right) && \text{by \textit{real-valued} hypothesis}\\
&= \boxed{\tilde{\mathsf{x}}(-\omega)} && \text{by result (1)}
\end{aligned}$$



Proposition P.2. *Let* $\breve{\mathsf{x}}(z)$ *be the* Z-TRANSFORM *(Definition Q.4 page 354) and* $\breve{\mathsf{x}}(\omega)$ *the* DISCRETE-TIME FOURIER TRANSFORM *(Definition P.1 page 343) of* $\left(\!\left(x_n\right)\!\right)$.

PRP

$$\underbrace{\left\{\sum_{n\in\mathbb{Z}} x_n = c\right\}}_{\text{(1) time domain}} \iff \underbrace{\left\{\breve{\mathsf{x}}(z)\Big|_{z=1} = c\right\}}_{\text{(2) z domain}} \iff \underbrace{\left\{\breve{\mathsf{x}}(\omega)\Big|_{\omega=0} = c\right\}}_{\text{(3) frequency domain}} \qquad \forall \left(\!\left(x_n\right)\!\right)_{n\in\mathbb{Z}} \in \ell^2_{\mathbb{R}}, c\in\mathbb{R}$$

✎PROOF:

1. Proof that (1) $\implies$ (2):

$$\begin{aligned}
\breve{\mathsf{x}}(z)\Big|_{z=1} &= \sum_{n\in\mathbb{Z}} x_n z^{-n}\Big|_{z=1} && \text{by definition of } \breve{\mathsf{x}}(z) \text{ (Definition Q.4 page 354)}\\
&= \sum_{n\in\mathbb{Z}} x_n && \text{because } z^n = 1 \text{ for all } n\in\mathbb{Z}\\
&= c && \text{by hypothesis (1)}
\end{aligned}$$

2. Proof that (2) $\implies$ (3):

$$\begin{aligned}
\breve{\mathsf{x}}(\omega)\Big|_{\omega=0} &= \sum_{n\in\mathbb{Z}} x_n e^{-i\omega n}\Big|_{\omega=0} && \text{by definition of } \breve{\mathsf{x}}(\omega) \qquad \text{(Definition P.1 page 343)}\\
&= \sum_{n\in\mathbb{Z}} x_n z^{-n}\Big|_{z=1}\\
&= \breve{\mathsf{x}}(z)\Big|_{z=1} && \text{by definition of } \breve{\mathsf{x}}(z) \qquad \text{(Definition Q.4 page 354)}\\
&= c && \text{by hypothesis (2)}
\end{aligned}$$

COPYRIGHT © 2019 DANIEL J. GREENHOE
CC BY-NC-ND

3. Proof that (3) $\Longrightarrow$ (1):

$$\begin{aligned}
\sum_{n\in\mathbb{Z}} x_n &= \sum_{n\in\mathbb{Z}} x_n e^{-i\omega n}\Big|_{\omega=0} \\
&= \check{\mathsf{x}}(\omega) && \text{by definition of } \check{\mathsf{x}}(\omega) && \text{(Definition P.1 page 343)} \\
&= c && \text{by hypothesis (3)}
\end{aligned}$$

✏

Proposition P.3. *If the coefficients are **real**, then the magnitude response (MR) is **symmetric**.*

✎PROOF:

$$\begin{aligned}
\left|\tilde{\mathsf{h}}(-\omega)\right| &\triangleq \left|\check{\mathsf{h}}(z)\right|_{z=e^{-i\omega}} && \triangleq \left|\sum_{m\in\mathbb{Z}} \mathsf{x}[m]z^{-m}\right|_{z=e^{-i\omega}} \\
&= \left|\sum_{m\in\mathbb{Z}} \mathsf{x}[m]e^{i\omega m}\right| && = \left|\left(\sum_{m\in\mathbb{Z}} \mathsf{x}^*[m]e^{-i\omega m}\right)^*\right| \\
&= \underbrace{\left|\left(\sum_{m\in\mathbb{Z}} \mathsf{x}[m]e^{-i\omega m}\right)^*\right|}_{\text{if } x[m] \text{ is } real} && = \left|\sum_{m\in\mathbb{Z}} \mathsf{x}[m]e^{-i\omega m}\right| \\
&\triangleq \left|\check{\mathsf{h}}(z)\right|_{z=e^{-i\omega}} && \triangleq \left|\tilde{\mathsf{h}}(\omega)\right|
\end{aligned}$$

✏

Proposition P.4. [1]

PRP

$$\underbrace{\sum_{n\in\mathbb{Z}}(-1)^n x_n = c}_{\text{(1) in "time"}} \iff \underbrace{\check{\mathsf{x}}(z)|_{z=-1} = c}_{\text{(2) in "z domain"}} \iff \underbrace{\check{\mathsf{x}}(\omega)|_{\omega=\pi} = c}_{\text{(3) in "frequency"}}$$

$$\iff \underbrace{\left(\sum_{n\in\mathbb{Z}} h_{2n}, \sum_{n\in\mathbb{Z}} h_{2n+1}\right) = \left(\frac{1}{2}\left(\sum_{n\in\mathbb{Z}} h_n + c\right), \frac{1}{2}\left(\sum_{n\in\mathbb{Z}} h_n - c\right)\right)}_{\text{(4) sum of even, sum of odd}}$$

$\forall c\in\mathbb{R}, (x_n)_{n\in\mathbb{Z}}, (y_n)_{n\in\mathbb{Z}} \in \ell^2_{\mathbb{R}}$

✎PROOF:

1. Proof that (1) $\Longrightarrow$ (2):

$$\begin{aligned}
\check{\mathsf{x}}(z)|_{z=-1} &= \sum_{n\in\mathbb{Z}} x_n z^{-n}\Big|_{z=-1} \\
&= \sum_{n\in\mathbb{Z}} (-1)^n x_n \\
&= c && \text{by (1)}
\end{aligned}$$

[1] Chui (1992) page 123

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

2. Proof that (2) $\Longrightarrow$ (3):

$$\begin{aligned}
\sum_{n\in\mathbb{Z}} x_n e^{-i\omega n}\Big|_{\omega=\pi} &= \sum_{n\in\mathbb{Z}} (-1)^n x_n \\
&= \sum_{n\in\mathbb{Z}} (-1)^{-n} x_n && = \sum_{n\in\mathbb{Z}} z^{-n} x_n\Big|_{z=-1} \\
&= c && \text{by (2)}
\end{aligned}$$

3. Proof that (3) $\Longrightarrow$ (1):

$$\begin{aligned}
\sum_{n\in\mathbb{Z}} (-1)^n x_n &= \sum_{n\in\mathbb{Z}} (-1)^{-n} x_n \\
&= \sum_{n\in\mathbb{Z}} e^{-i\omega n} x_n\Big|_{\omega=\pi} \\
&= c && \text{by (3)}
\end{aligned}$$

4. Proof that (2) $\Longrightarrow$ (4):

   (a) Define $A \triangleq \sum_{n\in\mathbb{Z}} h_{2n}$ $\qquad B \triangleq \sum_{n\in\mathbb{Z}} h_{2n+1}$.

   (b) Proof that $A - B = c$:

$$\begin{aligned}
c &= \sum_{n\in\mathbb{Z}} (-1)^n x_n && \text{by (2)} \\
&= \underbrace{\sum_{n\in\mathbb{Z}_e} (-1)^n x_n}_{\text{even terms}} + \underbrace{\sum_{n\in\mathbb{Z}_o} (-1)^n x_n}_{\text{odd terms}} \\
&= \sum_{n\in\mathbb{Z}} (-1)^{2n} x_{2n} + \sum_{n\in\mathbb{Z}} (-1)^{2n+1} x_{2n+1} \\
&= \underbrace{\sum_{n\in\mathbb{Z}} x_{2n}}_{A} - \underbrace{\sum_{n\in\mathbb{Z}} x_{2n+1}}_{B} \\
&\triangleq A - B && \text{by definitions of } A \text{ and } B
\end{aligned}$$

   (c) Proof that $A + B = \sum_{n\in\mathbb{Z}} x_n$:

$$\begin{aligned}
\sum_{n\in\mathbb{Z}} x_n &= \sum_{n \text{ even}} x_n + \sum_{n \text{ odd}} x_n \\
&= \underbrace{\sum_{n\in\mathbb{Z}} x_{2n}}_{A} + \underbrace{\sum_{n\in\mathbb{Z}} x_{2n+1}}_{B} \\
&= A + B && \text{by definitions of } A \text{ and } B
\end{aligned}$$

   (d) This gives two simultaneous equations:

$$\begin{aligned}
A - B &= c \\
A + B &= \sum_{n\in\mathbb{Z}} x_n
\end{aligned}$$

Copyright © 2019 Daniel J. Greenhoe

(e) Solutions to these equations give

$$\sum_{n\in\mathbb{Z}} x_{2n} \triangleq A \qquad = \frac{1}{2}\left(\sum_{n\in\mathbb{Z}} x_n + c\right)$$

$$\sum_{n\in\mathbb{Z}} x_{2n+1} \triangleq B \qquad = \frac{1}{2}\left(\sum_{n\in\mathbb{Z}} x_n - c\right)$$

5. Proof that (2) $\Longleftarrow$ (4):

$$\begin{aligned}
\sum_{n\in\mathbb{Z}}(-1)^n x_n &= \underbrace{\sum_{n\in\mathbb{Z}_e}(-1)^n x_n}_{\text{even terms}} + \underbrace{\sum_{n\in\mathbb{Z}_o}(-1)^n x_n}_{\text{odd terms}} \\
&= \sum_{n\in\mathbb{Z}}(-1)^{2n} x_{2n} + \sum_{n\in\mathbb{Z}}(-1)^{2n+1} x_{2n+1} \\
&= \sum_{n\in\mathbb{Z}} x_{2n} - \sum_{n\in\mathbb{Z}} x_{2n+1} \\
&= \frac{1}{2}\left(\sum_{n\in\mathbb{Z}} x_n + c\right) - \frac{1}{2}\left(\sum_{n\in\mathbb{Z}} x_n - c\right) && \text{by (3)} \\
&= c
\end{aligned}$$

✏

Lemma P.1. *Let* $\tilde{\mathsf{f}}(\omega)$ *be the* DTFT *(Definition P.1 page 343) of a sequence* $(\!(x_n)\!)_{n\in\mathbb{Z}}$.

LEM

$$\underbrace{(\!(x_n \in \mathbb{R})\!)_{n\in\mathbb{Z}}}_{\text{REAL-VALUED } \textit{sequence}} \quad \Longrightarrow \quad \underbrace{|\check{\mathsf{x}}(\omega)|^2 = |\check{\mathsf{x}}(-\omega)|^2}_{\text{EVEN}} \qquad \forall (\!(x_n)\!)_{n\in\mathbb{Z}} \in \ell^2_{\mathbb{R}}$$

✎PROOF:

$$\begin{aligned}
|\check{\mathsf{x}}(\omega)|^2 &= |\check{\mathsf{x}}(z)|^2\big|_{z=e^{i\omega}} \\
&= \check{\mathsf{x}}(z)\check{\mathsf{x}}^*(z)\big|_{z=e^{i\omega}} \\
&= \left[\sum_{n\in\mathbb{Z}} x_n z^{-n}\right]\left[\sum_{m\in\mathbb{Z}} x_m z^{-n}\right]^*\Bigg|_{z=e^{i\omega}} \\
&= \left[\sum_{n\in\mathbb{Z}} x_n z^{-n}\right]\left[\sum_{m\in\mathbb{Z}} x_m^*(z^*)^{-m}\right]\Bigg|_{z=e^{i\omega}} \\
&= \sum_{n\in\mathbb{Z}}\sum_{m\in\mathbb{Z}} x_n x_m^* z^{-n}(z^*)^{-m}\Bigg|_{z=e^{i\omega}} \\
&= \sum_{n\in\mathbb{Z}}\left[|x_n|^2 + \sum_{m>n} x_n x_m^* z^{-n}(z^*)^{-m} + \sum_{m<n} x_n x_m^* z^{-n}(z^*)^{-m}\right]\Bigg|_{z=e^{i\omega}} \\
&= \sum_{n\in\mathbb{Z}}\left[|x_n|^2 + \sum_{m>n} x_n x_m e^{i\omega(m-n)} + \sum_{m<n} x_n x_m e^{i\omega(m-n)}\right] \\
&= \sum_{n\in\mathbb{Z}}\left[|x_n|^2 + \sum_{m>n} x_n x_m e^{i\omega(m-n)} + \sum_{m>n} x_n x_m e^{-i\omega(m-n)}\right] \\
&= \sum_{n\in\mathbb{Z}}\left[|x_n|^2 + \sum_{m>n} x_n x_m \left(e^{i\omega(m-n)} + e^{-i\omega(m-n)}\right)\right]
\end{aligned}$$

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

$$= \sum_{n\in\mathbb{Z}} \left[|x_n|^2 + \sum_{m>n} x_n x_m 2\cos[\omega(m-n)] \right]$$
$$= \sum_{n\in\mathbb{Z}} |x_n|^2 + 2\sum_{n\in\mathbb{Z}}\sum_{m>n} x_n x_m \cos[\omega(m-n)]$$

Since cos is real and even, then $|\check{\mathrm{x}}(\omega)|^2$ must also be real and even. ⇨

Theorem P.3 (inverse DTFT). [2] *Let $\check{\mathrm{x}}(\omega)$ be the* DISCRETE-TIME FOURIER TRANSFORM *(Definition P.1 page 343) of a sequence $(\!(x_n)\!)_{n\in\mathbb{Z}} \in \ell^2_{\mathbb{R}}$. Let $\check{\mathrm{x}}^{-1}$ be the inverse of $\check{\mathrm{x}}$.*

THM
$$\underbrace{\left\{ \check{\mathrm{x}}(\omega) \triangleq \sum_{n\in\mathbb{Z}} x_n e^{-i\omega n} \right\}}_{\check{\mathrm{x}}(\omega) \triangleq \check{\mathbf{F}}(\!(x_n)\!)} \implies \underbrace{\left\{ x_n = \frac{1}{2\pi}\int_{\alpha-\pi}^{\alpha+\pi} \check{\mathrm{x}}(\omega) e^{i\omega n}\, d\omega \quad \forall \alpha\in\mathbb{R} \right\}}_{(\!(x_n)\!) = \check{\mathbf{F}}^{-1}\check{\mathbf{F}}(\!(x_n)\!)} \quad \forall (\!(x_n)\!)_{n\in\mathbb{Z}} \in \ell^2_{\mathbb{R}}$$

✎PROOF:

$$\frac{1}{2\pi}\int_{\alpha-\pi}^{\alpha+\pi} \check{\mathrm{x}}(\omega) e^{i\omega n}\, d\omega = \frac{1}{2\pi}\int_{\alpha-\pi}^{\alpha+\pi} \underbrace{\left[\sum_{m\in\mathbb{Z}} x_m e^{-i\omega m} \right]}_{\check{\mathrm{x}}(\omega)} e^{i\omega n}\, d\omega \qquad \text{by definition of } \check{\mathrm{x}}(\omega)$$
$$= \frac{1}{2\pi}\int_{\alpha-\pi}^{\alpha+\pi} \sum_{m\in\mathbb{Z}} x_m e^{-i\omega(m-n)}\, d\omega$$
$$= \frac{1}{2\pi}\sum_{m\in\mathbb{Z}} x_m \int_{\alpha-\pi}^{\alpha+\pi} e^{-i\omega(m-n)}\, d\omega$$
$$= \frac{1}{2\pi}\sum_{m\in\mathbb{Z}} x_m \left[2\pi \bar{\delta}_{m-n} \right]$$
$$= x_n$$

⇨

Theorem P.4 (orthonormal quadrature conditions). [3] *Let $\check{\mathrm{x}}(\omega)$ be the* DISCRETE-TIME FOURIER TRANSFORM *(Definition P.1 page 343) of a sequence $(\!(x_n)\!)_{n\in\mathbb{Z}} \in \ell^2_{\mathbb{R}}$. Let $\bar{\delta}_n$ be the* KRONECKER DELTA FUNCTION *at n (Definition J.3 page 267).*

THM
$$\sum_{m\in\mathbb{Z}} x_m y^*_{m-2n} = 0 \iff \check{\mathrm{x}}(\omega)\check{\mathrm{y}}^*(\omega) + \check{\mathrm{x}}(\omega+\pi)\check{\mathrm{y}}^*(\omega+\pi) = 0 \qquad \forall n\in\mathbb{Z}, \forall (\!(x_n)\!),(\!(y_n)\!) \in \ell^2_{\mathbb{R}}$$
$$\sum_{m\in\mathbb{Z}} x_m x^*_{m-2n} = \bar{\delta}_n \iff |\check{\mathrm{x}}(\omega)|^2 + |\check{\mathrm{x}}(\omega+\pi)|^2 = 2 \qquad \forall n\in\mathbb{Z}, \forall (\!(x_n)\!),(\!(y_n)\!) \in \ell^2_{\mathbb{R}}$$

✎PROOF: Let $z \triangleq e^{i\omega}$.

[2] J.S.Chitode (2009a) page 3-95 ⟨(3.6.2)⟩
[3] Daubechies (1992) pages 132–137 ⟨(5.1.20),(5.1.39)⟩

COPYRIGHT © 2019 DANIEL J. GREENHOE | CC BY-NC-ND

1. Proof that $2\sum_{n\in\mathbb{Z}}\left[\sum_{k\in\mathbb{Z}} x_k y^*_{k-2n}\right] e^{-i2\omega n} = \check{x}(\omega)\check{y}^*(\omega) + \check{x}(\omega+\pi)\check{y}^*(\omega+\pi)$:

$$\begin{aligned}
&2\sum_{n\in\mathbb{Z}}\left[\sum_{k\in\mathbb{Z}} x_k y^*_{k-2n}\right] e^{-i2\omega n}\\
&= 2\sum_{k\in\mathbb{Z}} x_k \sum_{n\in\mathbb{Z}} y^*_{k-2n} z^{-2n}\\
&= 2\sum_{k\in\mathbb{Z}} x_k \sum_{n \text{ even}} y^*_{k-n} z^{-n}\\
&= \sum_{k\in\mathbb{Z}} x_k \sum_{n\in\mathbb{Z}} y^*_{k-n} z^{-n}\left(1+e^{i\pi n}\right)\\
&= \sum_{k\in\mathbb{Z}} x_k \sum_{n\in\mathbb{Z}} y^*_{k-n} z^{-n} + \sum_{k\in\mathbb{Z}} x_k \sum_{n\in\mathbb{Z}} y^*_{k-n} z^{-n} e^{i\pi n}\\
&= \sum_{k\in\mathbb{Z}} x_k \sum_{m\in\mathbb{Z}} y^*_{m} z^{-(k-m)} + \sum_{k\in\mathbb{Z}} x_k \sum_{m\in\mathbb{Z}} y^*_{m} e^{-i(\omega+\pi)(k-m)} \qquad \text{where } m \triangleq k-n\\
&= \sum_{k\in\mathbb{Z}} x_k z^{-k} \sum_{m\in\mathbb{Z}} y^*_{m} z^{m} + \sum_{k\in\mathbb{Z}} x_k e^{-i(\omega+\pi)k} \sum_{m\in\mathbb{Z}} y^*_{m} e^{+i(\omega+\pi)m}\\
&= \sum_{k\in\mathbb{Z}} x_k e^{-i\omega k}\left[\sum_{m\in\mathbb{Z}} y_m e^{-i\omega m}\right]^* + \sum_{k\in\mathbb{Z}} x_k e^{-i(\omega+\pi)k}\left[\sum_{m\in\mathbb{Z}} y_m e^{-i(\omega+\pi)m}\right]^*\\
&\triangleq \check{x}(\omega)\check{y}^*(\omega) + \check{x}(\omega+\pi)\check{y}^*(\omega+\pi)
\end{aligned}$$

2. Proof that $\sum_{m\in\mathbb{Z}} x_m y^*_{m-2n} = 0 \implies \check{x}(\omega)\check{y}^*(\omega) + \check{x}(\omega+\pi)\check{y}^*(\omega+\pi) = 0$:

$$\begin{aligned}
0 &= 2\sum_{n\in\mathbb{Z}}\left[\sum_{k\in\mathbb{Z}} x_k y^*_{k-2n}\right] e^{-i2\omega n} && \text{by left hypothesis}\\
&= \check{x}(\omega)\check{y}^*(\omega) + \check{x}(\omega+\pi)\check{y}^*(\omega+\pi) && \text{by item (1)}
\end{aligned}$$

3. Proof that $\sum_{m\in\mathbb{Z}} x_m y^*_{m-2n} = 0 \impliedby \check{x}(\omega)\check{y}^*(\omega) + \check{x}(\omega+\pi)\check{y}^*(\omega+\pi) = 0$:

$$\begin{aligned}
2\sum_{n\in\mathbb{Z}}\left[\sum_{k\in\mathbb{Z}} x_k y^*_{k-2n}\right] e^{-i2\omega n} &= \check{x}(\omega)\check{y}^*(\omega) + \check{x}(\omega+\pi)\check{y}^*(\omega+\pi) && \text{by item (1)}\\
&= 0 && \text{by right hypothesis}
\end{aligned}$$

Thus by the above equation, $\sum_{n\in\mathbb{Z}}\left[\sum_{k\in\mathbb{Z}} x_k y^*_{k-2n}\right] e^{-i2\omega n} = 0$. The only way for this to be true is if $\sum_{k\in\mathbb{Z}} x_k y^*_{k-2n} = 0$.

4. Proof that $\sum_{m\in\mathbb{Z}} x_m x^*_{m-2n} = \bar{\delta}_n \implies |\check{x}(\omega)|^2 + |\check{x}(\omega'+\pi)|^2 = 2$:
Let $g_n \triangleq x_n$.

$$\begin{aligned}
2 &= 2\sum_{n\in\mathbb{Z}} \bar{\delta}_{n\in\mathbb{Z}} e^{-i2\omega n}\\
&= 2\sum_{n\in\mathbb{Z}}\left[\sum_{k\in\mathbb{Z}} x_k y^*_{k-2n}\right] e^{-i2\omega n} && \text{by left hypothesis}\\
&= \check{x}(\omega)\check{y}^*(\omega) + \check{x}(\omega+\pi)\check{y}^*(\omega+\pi) && \text{by item (1)}
\end{aligned}$$

5. Proof that $\sum_{m\in\mathbb{Z}} x_m x^*_{m-2n} = \bar{\delta}_n \impliedby |\check{x}(\omega)|^2 + |\check{x}(\omega'+\pi)|^2 = 2$:
Let $g_n \triangleq x_n$.

$$\begin{aligned}
2\sum_{n\in\mathbb{Z}}\left[\sum_{k\in\mathbb{Z}} x_k y^*_{k-2n}\right] e^{-i2\omega n} &= \check{x}(\omega)\check{y}^*(\omega) + \check{x}(\omega+\pi)\check{y}^*(\omega+\pi) && \text{by item (1)}\\
&= 2 && \text{by right hypothesis}
\end{aligned}$$

BY-NC-ND Copyright © 2019 Daniel J. Greenhoe

Thus by the above equation, $\sum_{n\in\mathbb{Z}}\left[\sum_{k\in\mathbb{Z}} x_k y^*_{k-2n}\right] e^{-i2\omega n} = 1$. The only way for this to be true is if $\sum_{k\in\mathbb{Z}} x_k y^*_{k-2n} = \bar{\delta}_n$.

✏

P.3 Derivatives

Theorem P.5. [4] *Let* $\check{x}(\omega)$ *be the* DTFT *(Definition P.1 page 343) of a sequence* $(\!(x_n)\!)_{n\in\mathbb{Z}}$.

THM

(A)	$\left[\frac{d}{d\omega}\right]^n \check{x}(\omega)\Big\vert_{\omega=0}$	$=$	0	$\Longleftrightarrow$	$\sum_{k\in\mathbb{Z}} k^n x_k$	$=$	0	*(B)*	$\forall n\in\mathbb{W}$
(C)	$\left[\frac{d}{d\omega}\right]^n \check{x}(\omega)\Big\vert_{\omega=\pi}$	$=$	0	$\Longleftrightarrow$	$\sum_{k\in\mathbb{Z}} (-1)^k k^n x_k$	$=$	0	*(D)*	$\forall n\in\mathbb{W}$

✎PROOF:

1. Proof that $(A) \Longrightarrow (B)$:

$$\begin{aligned}
0 &= \left[\frac{d}{d\omega}\right]^n \check{x}(\omega)\Big\vert_{\omega=0} && \text{by hypothesis (A)}\\
&= \left[\frac{d}{d\omega}\right]^n \sum_{k\in\mathbb{Z}} x_k e^{-i\omega k}\Bigg\vert_{\omega=0} && \text{by definition of } \check{x}(\omega) \text{ (Definition P.1 page 343)}\\
&= \sum_{k\in\mathbb{Z}} x_k \left[\frac{d}{d\omega}\right]^n e^{-i\omega k}\Bigg\vert_{\omega=0}\\
&= \sum_{k\in\mathbb{Z}} x_k \left[(-i)^n k^n e^{-i\omega k}\right]\Bigg\vert_{\omega=0}\\
&= (-i)^n \sum_{k\in\mathbb{Z}} k^n x_k
\end{aligned}$$

2. Proof that $(A) \Longleftarrow (B)$:

$$\begin{aligned}
\left[\frac{d}{d\omega}\right]^n \check{x}(\omega)\Big\vert_{\omega=0} &= \left[\frac{d}{d\omega}\right]^n \sum_{k\in\mathbb{Z}} x_k e^{-i\omega k}\Bigg\vert_{\omega=0} && \text{by definition of } \tilde{g}\\
&= \sum_{k\in\mathbb{Z}} x_k \left[\left[\frac{d}{d\omega}\right]^n e^{-i\omega k}\right]\Bigg\vert_{\omega=0}\\
&= \sum_{k\in\mathbb{Z}} x_k \left[(-i)^n k^n e^{-i\omega k}\right]\Bigg\vert_{\omega=0}\\
&= (-i)^n \sum_{k\in\mathbb{Z}} k^n x_k\\
&= 0 && \text{by hypothesis (B)}
\end{aligned}$$

[4] Vidakovic (1999), pages 82–83, Mallat (1999), pages 241–242

COPYRIGHT © 2019 DANIEL J. GREENHOE

3. Proof that $(C) \implies (D)$:

$$
\begin{aligned}
0 &= \left[\frac{\mathrm{d}}{\mathrm{d}\omega}\right]^n \check{x}(\omega)\Big|_{\omega=\pi} && \text{by hypothesis (C)}\\
&= \left[\frac{\mathrm{d}}{\mathrm{d}\omega}\right]^n \sum_{k\in\mathbb{Z}} x_k e^{-i\omega k}\Bigg|_{\omega=\pi} && \text{by definition of } \check{x} \text{ (Definition P.1 page 343)}\\
&= \sum_{k\in\mathbb{Z}} x_k \left[\frac{\mathrm{d}}{\mathrm{d}\omega}\right]^n e^{-i\omega k}\Bigg|_{\omega=\pi}\\
&= \sum_{k\in\mathbb{Z}} x_k \left[(-i)^n k^n e^{-i\omega k}\right]\Bigg|_{\omega=\pi}\\
&= \sum_{k\in\mathbb{Z}} x_k \left[(-i)^n k^n (-1)^k\right]\\
&= (-i)^n \sum_{k\in\mathbb{Z}} (-1)^k k^n x_k
\end{aligned}
$$

4. Proof that $(C) \impliedby (D)$:

$$
\begin{aligned}
\left[\frac{\mathrm{d}}{\mathrm{d}\omega}\right]^n \check{x}(\omega)\Big|_{\omega=\pi} &= \left[\frac{\mathrm{d}}{\mathrm{d}\omega}\right]^n \sum_{k\in\mathbb{Z}} x_k e^{-i\omega k}\Bigg|_{\omega=\pi} && \text{by definition of } \check{x} \text{ (Definition P.1 page 343)}\\
&= \sum_{k\in\mathbb{Z}} x_k \left[\frac{\mathrm{d}}{\mathrm{d}\omega}\right]^n e^{-i\omega k}\Bigg|_{\omega=\pi}\\
&= \sum_{k\in\mathbb{Z}} x_k \left[(-i)^n k^n e^{-i\omega k}\right]\Bigg|_{\omega=\pi}\\
&= \sum_{k\in\mathbb{Z}} x_k \left[(-i)^n k^n (-1)^k\right]\\
&= (-i)^n \sum_{k\in\mathbb{Z}} (-1)^k k^n x_k\\
&= 0 && \text{by hypothesis (D)}
\end{aligned}
$$

✏

BY-NC-ND Copyright © 2019 Daniel J. Greenhoe

APPENDIX Q

OPERATIONS ON SEQUENCES

Q.1 Convolution operator

Definition Q.1. [1] *Let X^Y be the set of all functions from a set Y to a set X. Let $\mathbb{Z}$ be the set of integers.*

DEF

*A function f in X^Y is a **sequence** over X if $Y = \mathbb{Z}$.*
A sequence may be denoted in the form $(\!(x_n)\!)_{n\in\mathbb{Z}}$ or simply as $(\!(x_n)\!)$.

Definition Q.2. [2] *Let $(\mathbb{F}, +, \cdot, 0, 1)$ be a* FIELD *(Definition F.5 page 206).*

DEF

*The **space of all absolutely square summable sequences** $\ell^2_{\mathbb{F}}$ over $\mathbb{F}$ is defined as*

$$\ell^2_{\mathbb{F}} \triangleq \left\{ (\!(x_n)\!)_{n\in\mathbb{Z}} \mid \sum_{n\in\mathbb{Z}} |x_n|^2 < \infty \right\}$$

The space $\ell^2_{\mathbb{R}}$ is an example of a *separable Hilbert space*. In fact, $\ell^2_{\mathbb{R}}$ is the *only* separable Hilbert space in the sense that all separable Hilbert spaces are isomorphically equivalent. For example, $\ell^2_{\mathbb{R}}$ is isomorphic to $L^2_{\mathbb{R}}$, the *space of all absolutely square Lebesgue integrable functions*.

Definition Q.3.

DEF

*The **convolution** operation $\star$ is defined as*

$$(\!(x_n)\!) \star (\!(y_n)\!) \triangleq \left(\left(\sum_{m\in\mathbb{Z}} x_m y_{n-m} \right) \right)_{n\in\mathbb{Z}} \qquad \forall (\!(x_n)\!)_{n\in\mathbb{Z}}, (\!(y_n)\!)_{n\in\mathbb{Z}} \in \ell^2_{\mathbb{R}}$$

Proposition Q.1. *Let $\star$ be the* CONVOLUTION OPERATOR *(Definition Q.3 page 353).*

PRP

$$(\!(x_n)\!) \star (\!(y_n)\!) = (\!(y_n)\!) \star (\!(x_n)\!) \qquad \forall (\!(x_n)\!)_{n\in\mathbb{Z}}, (\!(y_n)\!)_{n\in\mathbb{Z}} \in \ell^2_{\mathbb{R}} \qquad (\star \text{ is COMMUTATIVE})$$

[1] Bromwich (1908), page 1, Thomson et al. (2008) page 23 ⟨Definition 2.1⟩, Joshi (1997) page 31
[2] Kubrusly (2011) page 347 ⟨Example 5.K⟩

✎PROOF:

$$
\begin{aligned}
[x \star y](n) &\triangleq \sum_{m\in\mathbb{Z}} x_m y_{n-m} && \text{by Definition Q.3 page 353}\\
&= \sum_{k\in\mathbb{Z}} x_{n-k} y(k) && \text{where } k = n-m \iff m = n-k\\
&= \sum_{k\in\mathbb{Z}} x_{n-k} y(k) && \text{by change commutivity of addition}\\
&= \sum_{m\in\mathbb{Z}} x_{n-m} y_m && \text{by change of variables}\\
&= \sum_{m\in\mathbb{Z}} y_m x_{n-m} && \text{by commutative property of the field over } \mathbb{C}\\
&\triangleq (y \star x)_n && \text{by Definition Q.3 page 353}
\end{aligned}
$$

✏

Proposition Q.2. *Let $\star$ be the* CONVOLUTION OPERATOR *(Definition Q.3 page 353). Let $\ell^2_{\mathbb{R}}$ be the set of* ABSOLUTELY SUMMABLE *sequences (Definition Q.2 page 353).*

PRP

$$
\left\{ \begin{array}{lll} (A). & \mathsf{x}(n) \in \ell^2_{\mathbb{R}} & \text{and} \\ (B). & \mathsf{y}(n) \in \ell^2_{\mathbb{R}} & \end{array} \right\} \Longrightarrow \left\{ \sum_{k\in\mathbb{Z}} \mathsf{x}[k]\mathsf{y}[n+k] = \mathsf{x}[-n] \star \mathsf{y}(n) \right\}
$$

✎PROOF:

$$
\begin{aligned}
\sum_{k\in\mathbb{Z}} \mathsf{x}[k]\mathsf{y}[n+k] &= \sum_{-p\in\mathbb{Z}} \mathsf{x}[-p]\mathsf{y}[n-p] && \text{where } p \triangleq -k && \Longrightarrow k = -p\\
&= \sum_{p\in\mathbb{Z}} \mathsf{x}[-p]\mathsf{y}[n-p] && \text{by } \textit{absolutely summable} \text{ hypothesis} && \text{(Definition Q.2 page 353)}\\
&= \sum_{p\in\mathbb{Z}} \mathsf{x}'[p]\mathsf{y}[n-p] && \text{where } \mathsf{x}'[n] \triangleq \mathsf{x}[-n] && \Longrightarrow \mathsf{x}[-n] = \mathsf{x}'[n]\\
&\triangleq \mathsf{x}'[n] \star \mathsf{y}[n] && \text{by definition of } \textit{convolution} \star && \text{(Definition Q.3 page 353)}\\
&\triangleq \mathsf{x}[-n] \star \mathsf{y}[n] && \text{by definition of } \mathsf{x}'[n]
\end{aligned}
$$

✏

Q.2 Z-transform

Definition Q.4. [3]

DEF

*The **z-transform** $\mathbf{Z}$ of $(\!(x_n)\!)_{n\in\mathbb{Z}}$ is defined as*

$$
\left[\mathbf{Z}\left(\!(x_n)\!\right)\right](z) \triangleq \underbrace{\sum_{n\in\mathbb{Z}} x_n z^{-n}}_{\textit{Laurent series}} \qquad \forall (\!(x_n)\!) \in \ell^2_{\mathbb{R}}
$$

Theorem Q.1. *Let $X(z) \triangleq \mathbf{Z}\mathsf{x}[n]$ be the* Z-TRANSFORM *of $\mathsf{x}[n]$.*

THM

$$
\left\{ \check{\mathsf{x}}(z) \triangleq \mathbf{Z}\left(\!(\mathsf{x}[n])\!\right) \right\} \Longrightarrow \left\{ \begin{array}{llll}
(1). & \mathbf{Z}\left(\!(\alpha\mathsf{x}[n])\!\right) = \alpha\check{\mathsf{x}}(z) & \forall (\!(x_n)\!) \in \ell^2_{\mathbb{R}} & \text{and} \\
(2). & \mathbf{Z}\left(\!(\mathsf{x}[n-k])\!\right) = z^{-k}\check{\mathsf{x}}(z) & \forall (\!(x_n)\!) \in \ell^2_{\mathbb{R}} & \text{and} \\
(3). & \mathbf{Z}\left(\!(\mathsf{x}[-n])\!\right) = \check{\mathsf{x}}\left(\frac{1}{z}\right) & \forall (\!(x_n)\!) \in \ell^2_{\mathbb{R}} & \text{and} \\
(4). & \mathbf{Z}\left(\!(\mathsf{x}^*[n])\!\right) = \check{\mathsf{x}}^*\left(z^*\right) & \forall (\!(x_n)\!) \in \ell^2_{\mathbb{R}} & \text{and} \\
(5). & \mathbf{Z}\left(\!(\mathsf{x}^*[-n])\!\right) = \check{\mathsf{x}}^*\left(\frac{1}{z^*}\right) & \forall (\!(x_n)\!) \in \ell^2_{\mathbb{R}} &
\end{array} \right\}
$$

[3] *Laurent series*: Abramovich and Aliprantis (2002) page 49

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

✎PROOF:

$$
\begin{aligned}
\alpha\mathbb{Z}\check{\mathsf{x}}(z) &\triangleq \alpha\mathbf{Z}\left(\!\left(\mathsf{x}[n]\right)\!\right) && \text{by definition of } \check{\mathsf{x}}(z) \\
&\triangleq \alpha\sum_{n\in\mathbb{Z}}\mathsf{x}[n]z^{-n} && \text{by definition of } \mathbf{Z} \text{ operator} \\
&\triangleq \sum_{n\in\mathbb{Z}}(\alpha\mathsf{x}[n])z^{-n} && \text{by } \textit{distributive} \text{ property} \\
&\triangleq \mathbf{Z}\left(\!\left(\alpha\mathsf{x}[n]\right)\!\right) && \text{by definition of } \mathbf{Z} \text{ operator} \\
z^{-k}\check{\mathsf{x}}(z) &= z^{-k}\mathbf{Z}\left(\!\left(\mathsf{x}[n]\right)\!\right) && \text{by definition of } \check{\mathsf{x}}(z) && \text{(left hypothesis)} \\
&\triangleq z^{-k}\sum_{n=-\infty}^{n=+\infty}\mathsf{x}[n]z^{-n} && \text{by definition of } \mathbf{Z} && \text{(Definition Q.4 page 354)} \\
&= \sum_{n=-\infty}^{n=+\infty}\mathsf{x}[n]z^{-n-k} \\
&= \sum_{m-k=-\infty}^{m-k=+\infty}\mathsf{x}[m-k]z^{-m} && \text{where } m\triangleq n+k && \implies n=m-k \\
&= \sum_{m=-\infty}^{m=+\infty}\mathsf{x}[m-k]z^{-m} \\
&= \sum_{n=-\infty}^{n=+\infty}\mathsf{x}[n-k]z^{-n} && \text{where } n\triangleq m \\
&\triangleq \mathbf{Z}\left(\!\left(\mathsf{x}[n-k]\right)\!\right) && \text{by definition of } \mathbf{Z} && \text{(Definition Q.4 page 354)} \\
\mathbf{Z}\left(\!\left(\mathsf{x}^*[n]\right)\!\right) &\triangleq \sum_{n\in\mathbb{Z}}\mathsf{x}^*[n]z^{-n} && \text{by definition of } \mathbf{Z} && \text{(Definition Q.4 page 354)} \\
&\triangleq \left(\sum_{n\in\mathbb{Z}}\mathsf{x}[n](z^*)^{-n}\right)^* && \text{by definition of } \mathbf{Z} && \text{(Definition Q.4 page 354)} \\
&\triangleq \check{\mathsf{x}}^*(z^*) && \text{by definition of } \mathbf{Z} && \text{(Definition Q.4 page 354)} \\
\mathbf{Z}\left(\!\left(\mathsf{x}[-n]\right)\!\right) &\triangleq \sum_{n\in\mathbb{Z}}\mathsf{x}[-n]z^{-n} && \text{by definition of } \mathbf{Z} && \text{(Definition Q.4 page 354)} \\
&= \sum_{-m\in\mathbb{Z}}\mathsf{x}[m]z^{m} && \text{where } m\triangleq -n && \implies n=-m \\
&= \sum_{m\in\mathbb{Z}}\mathsf{x}[m]z^{m} && \text{by } \textit{absolutely summable} \text{ property} && \text{(Definition Q.2 page 353)} \\
&= \sum_{m\in\mathbb{Z}}\mathsf{x}[m]\left(\frac{1}{z}\right)^{-m} && \text{by } \textit{absolutely summable} \text{ property} && \text{(Definition Q.2 page 353)} \\
&\triangleq \check{\mathsf{x}}\left(\frac{1}{z}\right) && \text{by definition of } \mathbf{Z} && \text{(Definition Q.4 page 354)} \\
\mathbf{Z}\left(\!\left(\mathsf{x}^*[-n]\right)\!\right) &\triangleq \sum_{n\in\mathbb{Z}}\mathsf{x}^*[-n]z^{-n} && \text{by definition of } \mathbf{Z} && \text{(Definition Q.4 page 354)} \\
&= \sum_{-m\in\mathbb{Z}}\mathsf{x}^*[m]z^{m} && \text{where } m\triangleq -n && \implies n=-m \\
&= \sum_{m\in\mathbb{Z}}\mathsf{x}^*[m]z^{m} && \text{by } \textit{absolutely summable} \text{ property} && \text{(Definition Q.2 page 353)} \\
&= \sum_{m\in\mathbb{Z}}\mathsf{x}^*[m]\left(\frac{1}{z}\right)^{-m} && \text{by } \textit{absolutely summable} \text{ property} && \text{(Definition Q.2 page 353)} \\
&= \left(\sum_{m\in\mathbb{Z}}\mathsf{x}[m]\left(\frac{1}{z^*}\right)^{-m}\right)^* && \text{by } \textit{absolutely summable} \text{ property} && \text{(Definition Q.2 page 353)}
\end{aligned}
$$

COPYRIGHT © 2019 DANIEL J. GREENHOE



Figure Q.1: Direct form 1 order 2 IIR filters

$$\triangleq \check{x}^*\left(\frac{1}{z^*}\right) \qquad \text{by definition of } \mathbf{Z} \qquad \text{(Definition Q.4 page 354)}$$

✏

Theorem Q.2 (convolution theorem)**.** *Let $\star$ be the convolution operator* (Definition Q.3 page 353)*.*

THM

$$\mathbf{Z}\underbrace{\left(\langle\!\langle x_n\rangle\!\rangle \star \langle\!\langle y_n\rangle\!\rangle\right)}_{\textit{sequence convolution}} = \underbrace{\left(\mathbf{Z}\langle\!\langle x_n\rangle\!\rangle\right)\left(\mathbf{Z}\langle\!\langle y_n\rangle\!\rangle\right)}_{\textit{series multiplication}} \qquad \forall \langle\!\langle x_n\rangle\!\rangle_{n\in\mathbb{Z}}, \langle\!\langle y_n\rangle\!\rangle_{n\in\mathbb{Z}} \in \ell^2_{\mathbb{R}}$$

✎PROOF:

$$\begin{aligned}
[\mathbf{Z}(x\star y)](z) &\triangleq \mathbf{Z}\left(\sum_{m\in\mathbb{Z}} x_m y_{n-m}\right) && \text{by Definition Q.3 page 353}\\
&\triangleq \sum_{n\in\mathbb{Z}}\sum_{m\in\mathbb{Z}} x_m y_{n-m} z^{-n} && \text{by Definition Q.4 page 354}\\
&= \sum_{n\in\mathbb{Z}}\sum_{m\in\mathbb{Z}} x_m y_{n-m} z^{-n}\\
&= \sum_{m\in\mathbb{Z}}\sum_{n\in\mathbb{Z}} x_m y_{n-m} z^{-n}\\
&= \sum_{m\in\mathbb{Z}}\sum_{k\in\mathbb{Z}} x_m y_k z^{-(m+k)} && \text{where } k = n-m \iff n = m+k\\
&= \left[\sum_{m\in\mathbb{Z}} x_m z^{-m}\right]\left[\sum_{k\in\mathbb{Z}} y_k z^{-k}\right]\\
&\triangleq \left(\mathbf{Z}\langle\!\langle x_n\rangle\!\rangle\right)\left(\mathbf{Z}\langle\!\langle y_n\rangle\!\rangle\right) && \text{by Definition Q.4 page 354}
\end{aligned}$$

✏

Q.3 From z-domain back to time-domain

$$\check{y}(z) = b_0X(z) + b_1z^{-1}X(z) + b_2z^{-2}X(z) - a_1z^{-1}Y(z) + a_2z^{-2}Y(z)$$

$$y[n] = b_0x[n] + b_1x[n-1] + b_2x[n-2] - a_1y[n-1] - a_2y[n-2]$$

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

Example Q.1. See Figure Q.1 (page 356)

$$\frac{3z^2+5z+7}{2z^2+10z+12} = \frac{3z^2+5z+7}{2\left(z^2+5z+6\right)} = \frac{\left(\tfrac{3}{2}z^2+\tfrac{5}{2}z+\tfrac{7}{2}\right)}{z^2+5z+6} = \frac{\left(\tfrac{3}{2}+\tfrac{5}{2}z^{-1}+\tfrac{7}{2}z^{-2}\right)}{1+5z^{-1}+6z^{-2}}$$

Q.4 Zero locations

The system property of *minimum phase* is defined in Definition Q.5 (next) and illustrated in Figure Q.2 (page 357).



not minimum phase *minimum phase*

Figure Q.2: Minimum Phase filter

Definition Q.5. [4] *Let* $\check{x}(z) \triangleq \mathbf{Z}\left(\!\left(x_n\right)\!\right)$ *be the* Z TRANSFORM *(Definition Q.4 page 354) of a sequence* $\left(\!\left(x_n\right)\!\right)_{n\in\mathbb{Z}}$ *in* $\ell^2_{\mathbb{R}}$. *Let* $\left(\!\left(z_n\right)\!\right)_{n\in\mathbb{Z}}$ *be the* ZEROS *of* $\check{x}(z)$.

DEF

The sequence $\left(\!\left(x_n\right)\!\right)$ *is* ***minimum phase*** *if*

$$\underbrace{\left|z_n\right| < 1 \qquad \forall n \in \mathbb{Z}}_{\check{x}(z) \text{ has all its ZEROS inside the unit circle}}$$

The impulse response of a minimum phase filter has most of its energy concentrated near the beginning of its support, as demonstrated next.

Theorem Q.3 (Robinson's Energy Delay Theorem). [5] *Let* $\mathsf{p}(z) \triangleq \sum_{n=0}^{N} a_n z^{-n}$ *and* $\mathsf{q}(z) \triangleq \sum_{n=0}^{N} b_n z^{-n}$ *be polynomials.*

THM

$$\left\{\begin{array}{ll} \mathsf{p} & \text{is MINIMUM PHASE} \\ \mathsf{q} & \text{is NOT minimum phase} \end{array}\right\} \text{ and } \Longrightarrow \underbrace{\sum_{n=0}^{m-1}\left|a_n\right|^2}_{\substack{\text{"energy" of the first } m \text{ coefficients of } \mathsf{p}(z)}} \geq \underbrace{\sum_{n=0}^{m-1}\left|b_n\right|^2}_{\substack{\text{"energy" of the first } m \text{ coefficients of } \mathsf{q}(z)}} \qquad \forall 0 \leq m \leq N$$

But for more *symmetry*, put some zeros inside and some outside the unit circle.

Example Q.2. An example of a minimum phase polynomial is the Daubechies-4 scaling function. The minimum phase polynomial causes most of the energy to be concentrated near the origin, making it very *asymmetric*. In contrast, the Symlet-4 has a design very similar to that of Daubechies-4, but the selected zeros are not all within the unit circle in the complex z plane. This results in a

[4] Farina and Rinaldi (2000) page 91, Dumitrescu (2007) page 36
[5] Dumitrescu (2007) page 36, Robinson (1962), Robinson (1966) ⟨???⟩, Claerbout (1976), pages 52–53

COPYRIGHT © 2019 DANIEL J. GREENHOE
BY-NC-ND



Figure Q.3: Daubechies-4 and Symlet-4 scaling functions pole-zero plots

scaling function that is more symmetric and less contrated near the origin. Both scaling functions are illustrated in Figure Q.3 (page 358).

Q.5 Pole locations

Definition Q.6.

DEF *A filter (or system or operator)* **H** *is* ***causal*** *if its current output does not depend on future inputs.*

Definition Q.7.

DEF *A filter (or system or operator)* **H** *is* ***time-invariant*** *if the mapping it performs does not change with time.*

Definition Q.8.

DEF *An operation* **H** *is* ***linear*** *if any output* y_n *can be described as a linear combination of inputs* x_n *as in*

$$y_n = \sum_{m\in\mathbb{Z}} h(m)x(n-m) .$$

For a filter to be *stable*, place all the poles *inside* the unit circle.

Theorem Q.4. *A causal LTI filter is* ***stable*** *if all of its poles are* ***inside*** *the unit circle.*

Example Q.3. Stable/unstable filters are illustrated in Figure Q.4 (page 359).

True or False? This filter has no poles:

BY-NC-ND Copyright © 2019 Daniel J. Greenhoe



Figure Q.4: Pole-zero plot stable/unstable causal LTI filters (Example Q.3 page 358)

$H(z) = b_0 + b_1 z^{-1} + b_2 z^{-2}$



$$H(z) = b_0 + b_1 z^{-1} + b_2 z^{-2} = \frac{z^2}{z^2} \times \frac{b_0 + b_1 z^{-1} + b_2 z^{-2}}{1} = \frac{b_0 z^2 + b_1 z^1 + b_2}{z^2}$$



Q.6 Mirroring for real coefficients

If you want real coefficients, choose poles and zeros in conjugate pairs (next).

Proposition Q.3.

PRP $\left\{ \begin{array}{l} \text{ZEROS } \textit{and} \text{ POLES} \\ \textit{occur in} \text{ CONJUGATE PAIRS} \end{array} \right\} \Longrightarrow \left\{ \begin{array}{l} \text{COEFFICIENTS} \\ \textit{are} \text{ REAL.} \end{array} \right\}$

Copyright © 2019 Daniel J. Greenhoe



Figure Q.5: Conjugate pair structure yielding real coefficients

✎PROOF:

$$\begin{aligned}(z-p_1)(z-p_1^*) &= [z-(a+ib)][z-(a-ib)]\\ &= z^2+[-a+ib-ib-a]z-[ib]^2\\ &= z^2-2az+b^2\end{aligned}$$

✏

Example Q.4. See Figure Q.5 (page 360).

$$\begin{aligned}H(z) &= G\frac{[z-z_1][z-z_2]}{[z-p_1][z-p_2]} = G\frac{[z-(1+i)][z-(1-i)]}{[z-(-\tfrac{2}{3}+i\tfrac{1}{2})][z-(-\tfrac{2}{3}-i\tfrac{1}{2})]}\\ &= G\frac{z^2-z[(1-i)+(1+i)]+(1-i)(1+i)}{z^2-z[(-\tfrac{2}{3}+i\tfrac{1}{2})+(-\tfrac{2}{3}+i\tfrac{1}{2})]+(-\tfrac{2}{3}+i\tfrac{1}{2})(-\tfrac{2}{3}+i\tfrac{1}{2})}\\ &= G\frac{z^2-2z+2}{z^2-\tfrac{4}{3}z+(\tfrac{4}{3}+\tfrac{1}{4})} = G\frac{z^2-2z+2}{z^2-\tfrac{4}{3}z+\tfrac{19}{12}}\end{aligned}$$

Q.7 Rational polynomial operators

A digital filter is simply an operator on $\ell^2_{\mathbb{R}}$. If the digital filter is a causal LTI system, then it can be expressed as a rational polynomial in z as shown next.

Lemma Q.1. *A causal LTI operator* **H** *can be expressed as a rational expression* $\check{\mathsf{h}}(z)$.

$$\begin{aligned}\check{\mathsf{h}}(z) &= \frac{b_0+b_1z^{-1}+b_2z^{-2}+\cdots+b_Nz^{-N}}{1+a_1z^{-1}+a_2z^{-2}+\cdots+a_Nz^{-N}}\\ &= \frac{\sum_{n=0}^{N} b_nz^{-n}}{1+\sum_{n=1}^{N} a_nz^{-n}}\end{aligned}$$

A filter operation $\check{\mathsf{h}}(z)$ can be expressed as a product of its roots (poles and zeros).

$$\begin{aligned}\check{\mathsf{h}}(z) &= \frac{b_0+b_1z^{-1}+b_2z^{-2}+\cdots+b_Nz^{-N}}{1+a_1z^{-1}+a_2z^{-2}+\cdots+a_Nz^{-N}}\\ &= \alpha\frac{(z-z_1)(z-z_2)\cdots(z-z_N)}{(z-p_1)(z-p_2)\cdots(z-p_N)}\end{aligned}$$

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

where α is a constant, z_i are the zeros, and p_i are the poles. The poles and zeros of such a rational expression are often plotted in the z-plane with a unit circle about the origin (representing $z = e^{i\omega}$). Poles are marked with $\times$ and zeros with $\bigcirc$. An example is shown in Figure Q.6 page 361. Notice that in this figure the zeros and poles are either real or occur in complex conjugate pairs.



Figure Q.6: Pole-zero plot for rational expression with real coefficients

Q.8 Filter Banks

Conjugate quadrature filters (next definition) are used in *filter banks*. If $\check{x}(z)$ is a *low-pass filter*, then the conjugate quadrature filter of $\check{y}(z)$ is a *high-pass filter*.

Definition Q.9. [6] *Let* $(\!(x_n)\!)_{n\in\mathbb{Z}}$ *and* $(\!(y_n)\!)_{n\in\mathbb{Z}}$ *be* SEQUENCES *(Definition Q.1 page 353) in* $\ell^2_{\mathbb{R}}$ *(Definition Q.2 page 353).*

DEF

The sequence $(\!(y_n)\!)$ *is a* ***conjugate quadrature filter*** *with shift* N *with respect to* $(\!(x_n)\!)$ *if*

$$y_n = \pm(-1)^n x^*_{N-n}$$

A CONJUGATE QUADRATURE FILTER *is also called a* ***CQF*** *or a* ***Smith-Barnwell filter****.*

Any triple $\left((\!(x_n)\!), (\!(y_n)\!), N\right)$ *in this form is said to satisfy the* ***conjugate quadrature filter condition*** *or the* ***CQF condition****.*

Theorem Q.5 (CQF theorem). [7] *Let* $\check{y}(\omega)$ *and* $\check{x}(\omega)$ *be the* DTFTs *(Definition P.1 page 343) of the sequences* $(\!(y_n)\!)_{n\in\mathbb{Z}}$ *and* $(\!(x_n)\!)_{n\in\mathbb{Z}}$*, respectively, in* $\ell^2_{\mathbb{R}}$ *(Definition Q.2 page 353).*

THM

$$\underbrace{y_n = \pm(-1)^n x^*_{N-n}}_{(1)\text{ CQF in “time”}} \iff \check{y}(z) = \pm(-1)^N z^{-N}\check{x}^*\left(\frac{-1}{z^*}\right) \qquad (2)\ \text{CQF in “z-domain”}$$

$$\iff \check{y}(\omega) = \pm(-1)^N e^{-i\omega N}\check{x}^*(\omega+\pi) \qquad (3)\ \text{CQF in “frequency”}$$

$$\iff x_n = \pm(-1)^N(-1)^n y^*_{N-n} \qquad (4)\ \text{“reversed” CQF in “time”}$$

$$\iff \check{x}(z) = \pm z^{-N}\check{y}^*\left(\frac{-1}{z^*}\right) \qquad (5)\ \text{“reversed” CQF in “z-domain”}$$

$$\iff \check{x}(\omega) = \pm e^{-i\omega N}\check{y}^*(\omega+\pi) \qquad (6)\ \text{“reversed” CQF in “frequency”}$$

$\forall N\in\mathbb{Z}$

✎PROOF:

[6] Strang and Nguyen (1996) page 109, Haddad and Akansu (1992) pages 256–259 ⟨section 4.5⟩, Vaidyanathan (1993) page 342 ⟨(7.2.7), (7.2.8)⟩, Smith and Barnwell (1984a), Smith and Barnwell (1984b), Mintzer (1985)

[7] Strang and Nguyen (1996) page 109, Mallat (1999) pages 236–238 ⟨(7.58),(7.73)⟩, Haddad and Akansu (1992) pages 256–259 ⟨section 4.5⟩, Vaidyanathan (1993) page 342 ⟨(7.2.7), (7.2.8)⟩

COPYRIGHT © 2019 DANIEL J. GREENHOE

1. Proof that (1) $\implies$ (2):

$$\begin{aligned}
\check{y}(z) &= \sum_{n\in\mathbb{Z}} y_n z^{-n} && \text{by definition of } z\text{-transform} \quad \text{(Definition Q.4 page 354)}\\
&= \sum_{n\in\mathbb{Z}} \underbrace{(\pm)(-1)^n x^*_{N-n}}_{CQF} z^{-n} && \text{by (1)}\\
&= \pm \sum_{m\in\mathbb{Z}} (-1)^{N-m} x^*_m z^{-(N-m)} && \text{where } m \triangleq N-n \implies \quad n = N-m\\
&= \pm(-1)^N z^{-N} \sum_{m\in\mathbb{Z}} (-1)^{-m} x^*_m \left(z^{-1}\right)^{-m}\\
&= \pm(-1)^N z^{-N} \sum_{m\in\mathbb{Z}} x^*_m \left(-\frac{1}{z}\right)^{-m}\\
&= \pm(-1)^N z^{-N} \left[\sum_{m\in\mathbb{Z}} x_m \left(-\frac{1}{z^*}\right)^{-m}\right]^*\\
&= \pm(-1)^N z^{-N} \check{x}^*\left(\frac{-1}{z^*}\right) && \text{by definition of } z\text{-transform} \quad \text{(Definition Q.4 page 354)}
\end{aligned}$$

2. Proof that (1) $\impliedby$ (2):

$$\begin{aligned}
\check{y}(z) &= \pm(-1)^N z^{-N} \check{x}^*\left(\frac{-1}{z^*}\right) && \text{by (2)}\\
&= \pm(-1)^N z^{-N} \left[\sum_{m\in\mathbb{Z}} x_m \left(\frac{-1}{z^*}\right)^{-m}\right]^* && \text{by definition of } z\text{-transform} \quad \text{(Definition Q.4 page 354)}\\
&= \pm(-1)^N z^{-N} \left[\sum_{m\in\mathbb{Z}} x^*_m \left(-z^{-1}\right)^{-m}\right] && \text{by definition of } z\text{-transform} \quad \text{(Definition Q.4 page 354)}\\
&= \sum_{m\in\mathbb{Z}} (\pm)(-1)^{N-m} x^*_m z^{-(N-m)}\\
&= \sum_{m\in\mathbb{Z}} (\pm)(-1)^n x^*_{N-n} z^{-n} && \text{where } n = N-m \implies \quad m \triangleq N-n\\
&\implies x_n = \pm(-1)^n x^*_{N-n}
\end{aligned}$$

3. Proof that (1) $\implies$ (3):

$$\begin{aligned}
\check{y}(\omega) &\triangleq \check{x}(z)\Big|_{z=e^{i\omega}} && \text{by definition of } DTFT \quad \text{(Definition P.1 page 343)}\\
&= \left[\pm(-1)^N z^{-N} \check{x}\left(\frac{-1}{z^*}\right)\right]_{z=e^{i\omega}} && \text{by (2)}\\
&= \pm(-1)^N e^{-i\omega N} \check{x}\left(e^{i\pi} e^{i\omega}\right)\\
&= \pm(-1)^N e^{-i\omega N} \check{x}\left(e^{i(\omega+\pi)}\right)\\
&= \pm(-1)^N e^{-i\omega N} \check{x}(\omega+\pi) && \text{by definition of } DTFT \quad \text{(Definition P.1 page 343)}
\end{aligned}$$

4. Proof that (1) $\implies$ (6):

$$\begin{aligned}
\check{x}(\omega) &= \sum_{n\in\mathbb{Z}} y_n e^{-i\omega n} && \text{by definition of } DTFT \quad \text{(Definition P.1 page 343)}\\
&= \sum_{n\in\mathbb{Z}} \underbrace{\pm(-1)^n x^*_{N-n}}_{CQF} e^{-i\omega n} && \text{by (1)}\\
&= \sum_{m\in\mathbb{Z}} \pm(-1)^{N-m} x^*_m e^{-i\omega(N-m)} && \text{where } m \triangleq N-n \implies \quad n = N-m\\
&= \pm(-1)^N e^{-i\omega N} \sum_{m\in\mathbb{Z}} (-1)^m x^*_m e^{i\omega m}
\end{aligned}$$

BY-NC-ND Copyright © 2019 Daniel J. Greenhoe

$$= \pm(-1)^N e^{-i\omega N} \sum_{m\in\mathbb{Z}} e^{i\pi m} x_m^* e^{i\omega m}$$
$$= \pm(-1)^N e^{-i\omega N} \sum_{m\in\mathbb{Z}} x_m^* e^{i(\omega+\pi)m}$$
$$= \pm(-1)^N e^{-i\omega N} \left[\sum_{m\in\mathbb{Z}} x_m e^{-i(\omega+\pi)m}\right]^*$$
$$= \pm(-1)^N e^{-i\omega N} \check{x}^*(\omega+\pi) \qquad \text{by definition of } DTFT \qquad \text{(Definition P.1 page 343)}$$

5. Proof that (1) $\Longleftarrow$ (3):

$$y_n = \frac{1}{2\pi}\int_{-\pi}^{+\pi} \check{y}(\omega)e^{i\omega n}\,d\omega \qquad \text{by } \textit{inverse DTFT} \qquad \text{(Theorem P.3 page 349)}$$
$$= \frac{1}{2\pi}\int_{-\pi}^{+\pi} \underbrace{\pm(-1)^N e^{-iN\omega}\check{x}^*(\omega+\pi)}_{\text{right hypothesis}} e^{i\omega n}\,d\omega \qquad \text{by right hypothesis}$$
$$= \pm(-1)^N \frac{1}{2\pi}\int_{-\pi}^{+\pi} \check{x}^*(\omega+\pi)e^{i\omega(n-N)}\,d\omega \qquad \text{by right hypothesis}$$
$$= \pm(-1)^N \frac{1}{2\pi}\int_{0}^{2\pi} \check{x}^*(v)e^{i(v-\pi)(n-N)}\,dv \qquad \text{where } v \triangleq \omega+\pi \implies \omega = v-\pi$$
$$= \pm(-1)^N e^{-i\pi(n-N)} \frac{1}{2\pi}\int_{0}^{2\pi} \check{x}^*(v)e^{iv(n-N)}\,dv$$
$$= \pm(-1)^N \underbrace{(-1)^N}_{e^{i\pi N}}\underbrace{(-1)^n}_{e^{-i\pi n}}\left[\frac{1}{2\pi}\int_{0}^{2\pi} \check{x}(v)e^{iv(N-n)}\,dv\right]^*$$
$$= \pm(-1)^n x_{N-n}^* \qquad \text{by } \textit{inverse DTFT} \qquad \text{(Theorem P.3 page 349)}$$

6. Proof that (1) $\Longleftrightarrow$ (4):

$$\begin{aligned}
y_n = \pm(-1)^n x_{N-n}^* &\iff (\pm)(-1)^n y_n = (\pm)(\pm)(-1)^n(-1)^n x_{N-n}^* \\
&\iff \pm(-1)^n y_n = x_{N-n}^* \\
&\iff \left(\pm(-1)^n y_n\right)^* = \left(x_{N-n}^*\right)^* \\
&\iff \pm(-1)^n y_n^* = x_{N-n} \\
&\iff x_{N-n} = \pm(-1)^n y_n^* \\
&\iff x_m = \pm(-1)^{N-m} y_{N-m}^* \qquad \text{where } m \triangleq N-n \implies n = N-m \\
&\iff x_m = \pm(-1)^{N-m} y_{N-m}^* \\
&\iff x_m = \pm(-1)^N(-1)^m y_{N-m}^* \\
&\iff x_n = \pm(-1)^N(-1)^n y_{N-n}^* \qquad \text{by change of free variables}
\end{aligned}$$

7. Proofs for (5) and (6): not included. See proofs for (2) and (3).

✍

Theorem Q.6. [8] *Let* $\check{y}(\omega)$ *and* $\check{x}(\omega)$ *be the* DTFTs *(Definition P.1 page 343) of the sequences* $(\!(y_n)\!)_{n\in\mathbb{Z}}$ *and* $(\!(x_n)\!)_{n\in\mathbb{Z}}$*, respectively, in* $\ell^2_{\mathbb{R}}$ *(Definition Q.2 page 353).*

THM

Let $y_n = \pm(-1)^n x_{N-n}^*$ (CQF CONDITION, *Definition Q.9 page 361*)*. Then*

$$\left\{\begin{aligned}
&(A) \quad \left[\frac{d}{d\omega}\right]^n \check{y}(\omega)\Big|_{\omega=0} = 0 &&\iff \left[\frac{d}{d\omega}\right]^n \check{x}(\omega)\Big|_{\omega=\pi} = 0 \quad (B)\\
&&&\iff \sum_{k\in\mathbb{Z}}(-1)^k k^n x_k = 0 \quad (C)\\
&&&\iff \sum_{k\in\mathbb{Z}} k^n y_k = 0 \quad (D)
\end{aligned}\right\} \quad \forall n\in\mathbb{W}$$

[8] Vidakovic (1999), pages 82–83, Mallat (1999), pages 241–242

Copyright © 2019 Daniel J. Greenhoe

✎ PROOF:

1. Proof that (A) $\Longrightarrow$ (B):

$$
\begin{aligned}
0 &= \left[\frac{\mathrm{d}}{\mathrm{d}\omega}\right]^n \breve{\mathrm{y}}(\omega)\Big|_{\omega=0} && \text{by (A)} \\
&= \left[\frac{\mathrm{d}}{\mathrm{d}\omega}\right]^n (\pm)(-1)^N e^{-i\omega N}\breve{\mathrm{x}}^*(\omega+\pi)\Big|_{\omega=0} && \text{by } \textit{CQF theorem} \quad \text{(Theorem Q.5 page 361)} \\
&= (\pm)(-1)^N \sum_{\ell=0}^{n}\binom{n}{\ell}\left[\frac{\mathrm{d}}{\mathrm{d}\omega}\right]^{\ell}\left[e^{-i\omega N}\right]\cdot\left[\frac{\mathrm{d}}{\mathrm{d}\omega}\right]^{n-\ell}\left[\breve{\mathrm{x}}^*(\omega+\pi)\right]\Bigg|_{\omega=0} && \text{by } \textit{Leibnitz GPR} \quad \text{(Lemma ?? page ??)} \\
&= (\pm)(-1)^N \sum_{\ell=0}^{n}\binom{n}{\ell}-iN^{\ell}e^{-i\omega N}\left[\frac{\mathrm{d}}{\mathrm{d}\omega}\right]^{n-\ell}\left[\breve{\mathrm{x}}^*(\omega+\pi)\right]\Bigg|_{\omega=0} \\
&= (\pm)(-1)^N \cancelto{1}{e^{-i0N}} \sum_{\ell=0}^{n}\binom{n}{\ell}-iN^{\ell}\left[\frac{\mathrm{d}}{\mathrm{d}\omega}\right]^{n-\ell}\left[\breve{\mathrm{x}}^*(\omega+\pi)\right]\Bigg|_{\omega=0}
\end{aligned}
$$

$$
\begin{aligned}
\Longrightarrow \quad \breve{\mathrm{x}}^{(0)}(\pi) &= 0 \\
\Longrightarrow \quad \breve{\mathrm{x}}^{(1)}(\pi) &= 0 \\
\Longrightarrow \quad \breve{\mathrm{x}}^{(2)}(\pi) &= 0 \\
\Longrightarrow \quad \breve{\mathrm{x}}^{(3)}(\pi) &= 0 \\
\Longrightarrow \quad \breve{\mathrm{x}}^{(4)}(\pi) &= 0 \\
\vdots \quad & \quad \vdots \\
\Longrightarrow \quad \breve{\mathrm{x}}^{(n)}(\pi) &= 0 \quad \text{for } n = 0, 1, 2, \ldots
\end{aligned}
$$

2. Proof that (A) $\Longleftarrow$ (B):

$$
\begin{aligned}
0 &= \left[\frac{\mathrm{d}}{\mathrm{d}\omega}\right]^n \breve{\mathrm{x}}(\omega)\Big|_{\omega=\pi} && \text{by (B)} \\
&= \left[\frac{\mathrm{d}}{\mathrm{d}\omega}\right]^n (\pm)e^{-i\omega N}\breve{\mathrm{y}}^*(\omega+\pi)\Big|_{\omega=\pi} && \text{by } \textit{CQF theorem} \quad \text{(Theorem Q.5 page 361)} \\
&= (\pm) \sum_{\ell=0}^{n}\binom{n}{\ell}\left[\frac{\mathrm{d}}{\mathrm{d}\omega}\right]^{\ell}\left[e^{-i\omega N}\right]\cdot\left[\frac{\mathrm{d}}{\mathrm{d}\omega}\right]^{n-\ell}\left[\breve{\mathrm{y}}^*(\omega+\pi)\right]\Bigg|_{\omega=\pi} && \text{by } \textit{Leibnitz GPR} \quad \text{(Lemma ?? page ??)} \\
&= (\pm) \sum_{\ell=0}^{n}\binom{n}{\ell}(-iN)^{\ell}e^{-i\omega N}\left[\frac{\mathrm{d}}{\mathrm{d}\omega}\right]^{n-\ell}\left[\breve{\mathrm{y}}^*(\omega+\pi)\right]\Bigg|_{\omega=\pi} \\
&= (\pm)\cancelto{1}{e^{-i\pi N}} \sum_{\ell=0}^{n}\binom{n}{\ell}-iN^{\ell}\left[\frac{\mathrm{d}}{\mathrm{d}\omega}\right]^{n-\ell}\left[\breve{\mathrm{y}}^*(\omega+\pi)\right]\Bigg|_{\omega=\pi} \\
&= (\pm)(-1)^N \sum_{\ell=0}^{n}\binom{n}{\ell}-iN^{\ell}\left[\frac{\mathrm{d}}{\mathrm{d}\omega}\right]^{n-\ell}\left[\breve{\mathrm{y}}^*(\omega+\pi)\right]\Bigg|_{\omega=\pi}
\end{aligned}
$$

$$
\begin{aligned}
\Longrightarrow \quad \breve{\mathrm{y}}^{(0)}(0) &= 0 \\
\Longrightarrow \quad \breve{\mathrm{y}}^{(1)}(0) &= 0 \\
\Longrightarrow \quad \breve{\mathrm{y}}^{(2)}(0) &= 0 \\
\Longrightarrow \quad \breve{\mathrm{y}}^{(3)}(0) &= 0 \\
\Longrightarrow \quad \breve{\mathrm{y}}^{(4)}(0) &= 0 \\
\vdots \quad & \quad \vdots \\
\Longrightarrow \quad \breve{\mathrm{y}}^{(n)}(0) &= 0 \\
\Longrightarrow \quad \breve{\mathrm{y}}^{(n)}(0) &= 0 \quad \text{for } n = 0, 1, 2, \ldots
\end{aligned}
$$

3. Proof that (B) $\Longleftrightarrow$ (C): by Theorem P.5 page 351

4. Proof that (A) $\Longleftrightarrow$ (D): by Theorem P.5 page 351

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE



$\times$



$= \quad 1$

$$\frac{(z-z_1)(z-z_2)(z-z_3)(z-z_4)}{(z-p_1)(z-p_2)(z-p_3)(z-p_4)} \times \frac{(z-p_1)(z-p_2)(z-p_3)(z-p_4)}{(z-z_1)(z-z_2)(z-z_3)(z-z_4)} = 1$$

5. Proof that (CQF) $\not\Leftarrow$ (A): Here is a counterexample: $\check{y}(\omega) = 0$.

✏

Q.9 Inverting non-minimum phase filters

Minimum phase filters are easy to invert: each *zero* becomes a *pole* and each *pole* becomes a *zero*.



unstable

$\times$



all-pass

$=$



stable!

$$\begin{aligned}
|A(z)|_{z=e^{i\omega}} &= \frac{1}{r}\left|\frac{z-re^{i\phi}}{z-\frac{1}{r}e^{i\phi}}\right|_{z=e^{i\omega}} & &= \left|\frac{z-re^{i\phi}}{rz-e^{i\phi}}\right|_{z=e^{i\omega}} \\
&= \left|e^{i\phi}\left(\frac{e^{-i\phi}z-r}{rz-e^{i\phi}}\right)\right|_{z=e^{i\omega}} & &= \left|z\left(\frac{e^{-i\phi}-rz^{-1}}{rz-e^{i\phi}}\right)\right|_{z=e^{i\omega}} \\
&= \left|-z\left(\frac{rz^{-1}-e^{-i\phi}}{rz-e^{i\phi}}\right)\right|_{z=e^{i\omega}} & &= \left|\underbrace{e^{i\pi}}_{-1}e^{i\omega}\left(\frac{re^{-i\omega}-e^{-i\phi}}{re^{i\omega}-e^{i\phi}}\right)\right| \\
&= \left|\frac{1}{e^{-iv}}\left(\frac{re^{-i\omega}-e^{-i\phi}}{\left(re^{i\omega}-e^{i\phi}\right)^*}\right)\right| & &= \left|\frac{re^{-i\omega}-e^{-i\phi}}{re^{-i\omega}-e^{-i\phi}}\right| \\
&= 1
\end{aligned}$$

Copyright © 2019 Daniel J. Greenhoe
CC BY-NC-ND

Copyright © 2019 Daniel J. Greenhoe

APPENDIX R

B-SPLINES

R.1 Definitions

Definition R.1. *Let X be a set.*

DEF *The **step function** $\sigma \in \mathbb{R}^{\mathbb{R}}$ is defined as*

$$\sigma(x) \triangleq \mathbb{1}_{[0:\infty)}(x) \qquad \forall x \in \mathbb{R}.$$



Lemma R.1. *Let $\sigma(x)$ be the* STEP FUNCTION *(Definition R.1 page 367).*

LEM

$$\{g(x) > 0\} \quad \Longrightarrow \quad \left\{\sigma\left[g(x)f(x)\right] = \sigma[f(x)]\right\} \qquad \forall f,g \in \mathbb{R}^{\mathbb{R}}$$

✎PROOF:

$$\begin{aligned}
\sigma\left[g(x)f(x)\right] &\triangleq \mathbb{1}_{[0:\infty)}\left[g(x)f(x)\right] && \text{by definition of } \sigma(x) && \text{(Definition R.1 page 367)}\\
&\triangleq \begin{cases} 1 & \text{for } g(x)f(x) \geq 0 \\ 0 & \text{otherwise} \end{cases} && \text{by definition of } \mathbb{1}_A(x) && \text{(Definition ?? page ??)}\\
&= \begin{cases} 1 & \text{for } f(x) \geq 0 \\ 0 & \text{otherwise} \end{cases} && \text{by } g(x) > 0 \text{ hypothesis} \\
&\triangleq \mathbb{1}_{[0:\infty)}[f(x)] && \text{by definition of } \mathbb{1}_A(x) && \text{(Definition ?? page ??)}\\
&\triangleq \sigma[f(x)] && \text{by definition of } \sigma(x) && \text{(Definition R.1 page 367)}
\end{aligned}$$

✏

Definition R.2. [1] *Let $\mathbb{1}$ be the* SET INDICATOR *function (Definition ?? page ??). Let $f(x) \star g(x)$ represent the* CONVOLUTION *operation (Definition O.3 page 336).*

DEF *The n**th order cardinal B-spline** $\mathsf{N}_n(x)$ for $n \in \mathbb{W}$ is defined as*

$$\mathsf{N}_n(x) \triangleq \left\{ \begin{array}{ll} \mathbb{1}_{[0:1)}(x) & \text{for } n = 0 \\ \mathsf{N}_{n-1}(x) \star \mathsf{N}_0(x) & \text{for } n \in \mathbb{W}\backslash 0 \end{array} \right\} \qquad \forall x \in \mathbb{R}$$

Lemma R.2. [2]

[1] Chui (1992) page 85 ⟨(4.2.1)⟩, Christensen (2008) page 140, Chui (1988) page 1

[2] Christensen (2008) page 140, Chui (1992) page 85 ⟨(4.2.1)⟩, Chui (1988) page 1, Prasad and Iyengar (1997) page 145

LEM

$$\mathsf{N}_n(x) = \int_{\tau=0}^{\tau=1} \mathsf{N}_{n-1}(x-\tau)\,\mathrm{d}\tau \qquad \forall n \in \{1,2,3,\ldots\}$$

✎ PROOF:

$$\begin{aligned}
\mathsf{N}_n(x) &\triangleq \mathsf{N}_{n-1}(x) \star \mathsf{N}_0(x) && \text{by definition of } \mathsf{N}_n(x) && \text{(Definition R.2 page 367)}\\
&\triangleq \int_{\mathbb{R}} \mathsf{N}_{n-1}(x-\tau)\mathsf{N}_0(\tau)\,\mathrm{d}\tau && \text{by definition of convolution operation } \star && \text{(Definition O.3 page 336)}\\
&\triangleq \int_{\mathbb{R}} \mathsf{N}_{n-1}(x-\tau)\mathbb{1}_{[0:1)}(\tau)\,\mathrm{d}\tau && \text{by definition of } \mathsf{N}_0(x) && \text{(Definition R.2 page 367)}\\
&= \int_{[0:1)} \mathsf{N}_{n-1}(x-\tau)\,\mathrm{d}\tau && \text{by definition of } \mathbb{1}_A(x) && \text{(Definition ?? page ??)}\\
&= \int_{[0:1]} \mathsf{N}_{n-1}(x-\tau)\,\mathrm{d}\tau\\
&\triangleq \int_0^1 \mathsf{N}_{n-1}(x-\tau)\,\mathrm{d}\tau
\end{aligned}$$

✎

Lemma R.3. *Let* $\mathsf{f}(x)$ *be a* FUNCTION *in* $\mathbb{R}^{\mathbb{R}}$. *Let* $\mathsf{F}(x)$ *be the* ANTI-DERIVATIVE *of* $\mathsf{f}(x)$. *Let* $\sigma(x)$ *be the* STEP FUNCTION *(Definition R.1 page 367)*.

LEM

$$\begin{aligned}
&\int_{y=a}^{y=b} \mathsf{f}(x-y)\sigma(x-y)\,\mathrm{d}y\\
&= \left\{\begin{array}{ll} -\displaystyle\int_{y=x-a}^{y=x-b} \mathsf{f}(y)\,\mathrm{d}y & \textit{for } x \geq b\\ -\displaystyle\int_{y=x-a}^{y=0} \mathsf{f}(y)\,\mathrm{d}y & \textit{for } a \leq x \leq b\\ 0 & \textit{for } x \leq a \end{array}\right\} = \left\{\begin{array}{ll} \mathsf{F}(x-a)-\mathsf{F}(x-b) & \textit{for } x \geq b\\ \mathsf{F}(x-a)-\mathsf{F}(0) & \textit{for } a \leq x \leq b\\ 0 & \textit{for } x \leq a \end{array}\right\}\\
&= [\mathsf{F}(x-a)-\mathsf{F}(0)]\sigma(x-a) + [\mathsf{F}(0)-\mathsf{F}(x-b)]\sigma(x-b)
\end{aligned}$$

✎ PROOF:

$$\begin{aligned}
\int_{y=a}^{y=b} \mathsf{f}(x-y)\sigma(x-y)\,\mathrm{d}y &= \left\{\begin{array}{ll} \int_{y=a}^{y=b} \mathsf{f}(x-y)\,\mathrm{d}y & \text{for } x \geq b\\ \int_{y=a}^{y=x} \mathsf{f}(x-y)\,\mathrm{d}y & \text{for } a \leq x \leq b\\ 0 & \text{for } x \leq a \end{array}\right\} && \text{by definition of } \sigma \text{ (Definition R.1 page 367)}\\
&= \left\{\begin{array}{ll} -\int_{u=x-a}^{u=x-b} \mathsf{f}(u)\,\mathrm{d}u & \text{for } x \geq b\\ -\int_{u=x-a}^{u=0} \mathsf{f}(u)\,\mathrm{d}u & \text{for } a \leq x \leq b\\ 0 & \text{for } x \leq a \end{array}\right\} && \text{where } u \triangleq x-y \implies y = x-u\\
&= \left\{\begin{array}{ll} -\int_{y=x-a}^{y=x-b} \mathsf{f}(y)\,\mathrm{d}y & \text{for } x \geq b\\ -\int_{y=x-a}^{y=0} \mathsf{f}(y)\,\mathrm{d}y & \text{for } a \leq x \leq b\\ 0 & \text{for } x \leq a \end{array}\right\} && \text{by change of dummy variable } (u \to y)\\
&= \left\{\begin{array}{ll} \mathsf{F}(x-a)-\mathsf{F}(x-b) & \text{for } x \geq b\\ \mathsf{F}(x-a)-\mathsf{F}(0) & \text{for } a \leq x \leq b\\ 0 & \text{for } x \leq a \end{array}\right\} && \text{by } \textit{Fundamental Theorem of Calculus}\\
&= [\mathsf{F}(x-a)-\mathsf{F}(x-b)]\sigma(x-b) + [\mathsf{F}(x-a)-\mathsf{F}(0)][\sigma(x-a)-\sigma(x-b)]\\
&= [\mathsf{F}(x-a)-\mathsf{F}(0)]\sigma(x-a) + [\mathsf{F}(x-a)-\mathsf{F}(x-b)-\mathsf{F}(x-a)+\mathsf{F}(0)]\sigma(x-b)\\
&= [\mathsf{F}(x-a)-\mathsf{F}(0)]\sigma(x-a) + [\mathsf{F}(0)-\mathsf{F}(x-b)]\sigma(x-b)
\end{aligned}$$

✎

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE



Figure R.1: some low order B-splines (Example R.1 page 369)

Lemma R.4. *Let* $\sigma(x)$ *be the* STEP FUNCTION *(Definition R.1 page 367).*

LEM
$$\int_{\tau=0}^{\tau=1} (x-\tau-k)^n \sigma(x-\tau-k)\,d\tau = \frac{1}{n+1}\left[(x-k)^{n+1}\sigma(x-k) - (x-k-1)^{n+1}\sigma(x-k-1)\right]$$

✎PROOF:

$$\begin{aligned}
&\int_{\tau=0}^{\tau=1} (x-\tau-k)^n \sigma(x-\tau-k)\,d\tau \\
&= \int_{y=k}^{y=k+1} (x-y)^n \sigma(x-y)\,dy && \text{where } y \triangleq \tau+k \implies \tau = y-k \\
&= [F(x-k)-F(0)]\sigma(x-k) + [F(0)-F(x-k-1)]\sigma(x-k-1) && \text{by Lemma R.3 (page 368), where } f(x) \triangleq x^n \\
&= \frac{\left[(x-k)^{n+1}-0\right]\sigma(x-k) + \left[0-(x-k-1)^{n+1}\right]\sigma(x-k-1)}{n+1} && \text{because } F(x) \triangleq \int f(x)\,dx = \frac{x^{n+1}}{n+1}+c \\
&= \frac{1}{n+1}\left[(x-k)^{n+1}\sigma(x-k) - (x-k-1)^{n+1}\sigma(x-k-1)\right]
\end{aligned}$$

✏

Example R.1. [3] Let $\sigma(x)$ be the *step function* (Definition R.1 page 367). Let $\binom{n}{k}$ be the *binomial coefficient* (Definition ?? page ??). The 0*th order B-spline* (Definition R.2 page 367) $N_0(x)$ can be expressed as follows:

EX
$$N_0(x) = \left\{\begin{array}{ll} 1 & \text{for } x \in [0:1) \\ 0 & \text{otherwise} \end{array}\right\} = \left\{\sum_{k=0}^{1} (-1)^k \binom{1}{k} (x-k)^0 \sigma(x-k) \quad \forall x \in \mathbb{R}\right\}$$

The B-spline $N_0(x)$ is illustrated in Figure R.1 (page 369).

[3] Schumaker (2007) page 136 ⟨Table 1⟩

COPYRIGHT © 2019 DANIEL J. GREENHOE
CC BY-NC-ND

✎PROOF:

$$\begin{aligned}
\mathsf{N}_0(x) &= \mathbb{1}_{[0:1)}(x) && \text{by definition of } \mathsf{N}_0(x) && \text{(Definition R.2 page 367)}\\
&= \sigma(x) - \sigma(x-1) && \text{by definition of } \sigma(x) && \text{(Definition R.1 page 367)}\\
&= \left[\binom{1}{0}\sigma(x) - \binom{1}{1}\sigma(x-1)\right] && \text{by definition of } \textit{binomial coefficient } \binom{n}{k} && \text{(Definition ?? page ??)}\\
&= \sum_{k=0}^{1}(-1)^k\binom{1}{k}(x-k)^0\sigma(x-k) && \text{by definition of } \sum \text{ operator}
\end{aligned}$$

✏

Example R.2. [4] Let $\sigma(x)$ be the *step function*. Let $\binom{n}{k}$ be the *binomial coefficient*.
The 1*st order B-spline* $\mathsf{N}_1(x)$ can be expressed as follows:

EX
$$\mathsf{N}_1(x) = \left\{\begin{array}{rl} x & \text{for } x \in [0:1]\\ -x+2 & \text{for } x \in [1:2]\\ 0 & \text{otherwise}\end{array}\right\} = \left\{\sum_{k=0}^{2}(-1)^k\binom{2}{k}(x-k)\sigma(x-k) \quad \forall x\in\mathbb{R}\right\}$$

The B-spline $\mathsf{N}_1(x)$ is illustrated in Figure R.1 (page 369).

✎PROOF:

$$\begin{aligned}
\mathsf{N}_1(x) &= \int_{\tau=0}^{\tau=1} \mathsf{N}_0(x-\tau)\,\mathrm{d}\tau && \text{by Lemma R.2 page 367}\\
&= \int_{\tau=0}^{\tau=1} \sum_{k=0}^{1}(-1)^k\binom{1}{k}(x-\tau-k)^0\sigma(x-\tau-k)\,\mathrm{d}\tau && \text{by Example R.1 page 369}\\
&= \sum_{k=0}^{1}(-1)^k\binom{1}{k}\int_{\tau=0}^{\tau=1}(x-\tau-k)^0\sigma(x-\tau-k)\,\mathrm{d}\tau && \text{by } \textit{linearity} \text{ of } \int \mathrm{d}\tau \text{ operator}\\
&= \sum_{k=0}^{1}(-1)^k\binom{1}{k}\frac{1}{0+1}\left[(x-k)^{0+1}\sigma(x-k) - (x-k-1)^{0+1}\sigma(x-k-1)\right] && \text{by Lemma R.4 page 369}\\
&= \begin{pmatrix} 1\{(x-0)\sigma(x-0) - (x-1)\sigma(x-1)\}\\ -1\{(x-1)\sigma(x-1) - (x-2)\sigma(x-2)\}\end{pmatrix}\\
&= x\sigma(x) - 2(x-1)\sigma(x-1) + (x-2)\sigma(x-2)\\
&= \sum_{k=0}^{2}(-1)^k\binom{2}{k}(x-k)\sigma(x-k) && \text{by def. of } \binom{n}{k} \text{ (Definition ?? page ??)}\\
&= \left\{\begin{array}{ll} x & \text{for } x \in [0:1)\\ -x+2 & \text{for } x \in [1:2]\\ 0 & \text{otherwise}\end{array}\right\} && \text{by def. of } \sigma(x) \text{ (Definition R.1 page 367)}
\end{aligned}$$

✏

Example R.3. [5] Let $\sigma(x)$ be the *step function*. Let $\binom{n}{k}$ be the *binomial coefficient*.
The 2*nd order B-spline* $\mathsf{N}_2(x)$ can be expressed as follows:

EX
$$\mathsf{N}_2(x) = \frac{1}{2}\left\{\begin{array}{rl} x^2 & \text{for } x \in [0:1)\\ -2x^2+6x-3 & \text{for } x \in [1:2]\\ x^2-6x+9 & \text{for } x \in [2:3]\\ 0 & \text{otherwise}\end{array}\right\} = \left\{\frac{1}{2}\sum_{k=0}^{3}(-1)^k\binom{3}{k}(x-k)^2\sigma(x-k) \quad \forall x\in\mathbb{R}\right\}$$

The B-spline $\mathsf{N}_2(x)$ is illustrated in Figure R.1 (page 369).

[4] Christensen (2008) page 148 ⟨Exercise 6.2⟩, Christensen (2010) page 212 ⟨Exercise 10.2⟩, Heil (2011) pages 142–143 ⟨Definition 4.22 (The Schauder System)⟩, Schumaker (2007) page 136 ⟨Table 1⟩, Stoer and Bulirsch (2002) page 124

[5] Christensen (2008) page 148 ⟨Exercise 6.2⟩, Christensen (2010) page 212 ⟨Exercise 10.2⟩, Schumaker (2007) page 136 ⟨Table 1⟩, Stoer and Bulirsch (2002) page 124

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

✎PROOF:

$$
\begin{aligned}
\mathsf{N}_2(x) &= \int_{\tau=0}^{\tau=1} \mathsf{N}_1(x-\tau)\,\mathrm{d}\tau && \text{by Lemma R.2 page 367}\\
&= \int_{\tau=0}^{\tau=1} \sum_{k=0}^{2}(-1)^k\binom{2}{k}(x-\tau-k)\sigma(x-\tau-k)\,\mathrm{d}\tau && \text{by Example R.2 page 370}\\
&= \sum_{k=0}^{2}(-1)^k\binom{2}{k}\int_{\tau=0}^{\tau=1}(x-\tau-k)\sigma(x-\tau-k)\,\mathrm{d}\tau && \text{by \textit{linearity} of } \int \mathrm{d}\tau \text{ operator}\\
&= \sum_{k=0}^{2}(-1)^k\binom{2}{k}\frac{1}{1+1}\left[(x-k)^{1+1}\sigma(x-k)-(x-k-1)^{1+1}\sigma(x-k-1)\right] && \text{by Lemma R.4 page 369}\\
&= \frac{1}{2}\begin{pmatrix} 1 & \{(x-0)^2\sigma(x-0)-(x-1)^2\sigma(x-1)\}\\ -2 & \{(x-1)^2\sigma(x-1)-(x-2)^2\sigma(x-2)\}\\ +1 & \{(x-2)^2\sigma(x-2)-(x-3)^2\sigma(x-3)\}\end{pmatrix}\\
&= \frac{1}{2}\left[x^2\sigma(x)-3(x-1)^2\sigma(x-1)+3(x-2)^2\sigma(x-2)-(x-3)^2\sigma(x-3)\right]\\
&= \frac{1}{2}\sum_{k=0}^{3}(-1)^k\binom{3}{k}(x-k)^2\sigma(x-k) && \text{by def. of } \binom{n}{k} \text{ (Definition ?? page ??)}\\
&= \frac{1}{2}\left\{\begin{array}{rl} x^2 & \text{for } x\in[0:1)\\ -2x^2+6x-3 & \text{for } x\in[1:2]\\ x^2-6x+9 & \text{for } x\in[2:3]\\ 0 & \text{otherwise}\end{array}\right\} && \text{by def. of } \sigma(x) \text{ (Definition R.1 page 367)}
\end{aligned}
$$

The final steps of this proof can be calculated "by hand" or by using the free and open source software package *Maxima* along with the script file listed in Section **??** (page **??**). ✏

R.2 Algebraic properties

Theorem R.1 (next) presents a closed form expression for an *nth order B-spline* $\mathsf{N}_n(x)$ based on the definition of $\mathsf{N}_n(x)$ given in Definition R.2 (page 367). Alternatively, Theorem R.1 could serve as the definition and Definition R.2 as a property.

Theorem R.1. [6] *Let* $\mathsf{N}_n(x)$ *be the* nTH ORDER B-SPLINE *(Definition R.2 page 367)*.
Let $\sigma(x)$ *be the* STEP FUNCTION *(Definition R.1 page 367)*.

THM

$$\mathsf{N}_n(x) = \frac{1}{n!}\sum_{k=0}^{n+1}(-1)^k\binom{n+1}{k}(x-k)^n\sigma(x-k) \qquad \forall n\in\{0,1,2,\ldots\}=\mathbb{W}$$

✎PROOF: Proof follows by induction:

1. base case (choose one):
   - Proof for $n=0$ case: by Example R.1 (page 369).
   - Proof for $n=1$ case: by Example R.2 (page 370).
   - Proof for $n=2$ case: by Example R.3 (page 370).

[6] Christensen (2008) page 142 ⟨Theorem 6.1.3⟩, Chui (1992) page 84 ⟨(4.1.12)⟩

COPYRIGHT © 2019 DANIEL J. GREENHOE

2. inductive step—proof that n case $\implies$ $n+1$ case:

$$
\begin{aligned}
&\mathsf{N}_{n+1}(x)\\
&= \int_0^1 \mathsf{N}_n(x-\tau)\,\mathrm{d}\tau && \text{by Lemma R.2 page 367}\\
&= \int_0^1 \frac{1}{n!}\sum_{k=0}^{k=n+1}(-1)^k\binom{n+1}{k}(x-\tau-k)^n\sigma(x-\tau-k)\,\mathrm{d}\tau && \text{by induction hypothesis}\\
&= \frac{1}{n!}\sum_{k=0}^{k=n+1}(-1)^k\binom{n+1}{k}\int_0^1(x-\tau-k)^n\sigma(x-\tau-k)\,\mathrm{d}\tau && \text{by \textit{linearity} of } \int \mathrm{d}\tau \text{ operator}\\
&= \frac{1}{n!}\sum_{k=0}^{k=n+1}(-1)^k\binom{n+1}{k}\frac{1}{n+1}\left[(x-k)^{n+1}\sigma(x-k)-(x-k-1)^{n+1}\sigma(x-k-1)\right] && \text{by Lemma R.4 page 369}\\
&= \frac{1}{(n+1)!}\sum_{k=0}^{k=n+1}(-1)^k\binom{n+1}{k}\left[(x-k)^{n+1}\sigma(x-k)-(x-k-1)^{n+1}\sigma(x-k-1)\right]\\
&= \frac{1}{(n+1)!}\left[\sum_{k=0}^{k=n+1}(-1)^k\binom{n+1}{k}(x-k)^{n+1}\sigma(x-k)-\sum_{k=0}^{k=n+1}(-1)^k\binom{n+1}{k}(x-k-1)^{n+1}\sigma(x-k-1)\right]\\
&= \frac{1}{(n+1)!}\left[\sum_{k=0}^{k=n+1}(-1)^k\binom{n+1}{k}(x-k)^{n+1}\sigma(x-k)-\sum_{m=1}^{m=n+2}(-1)^{m-1}\binom{n+1}{m-1}(x-m)^{n+1}\sigma(x-m)\right]\\
&\quad \text{where } m \triangleq k+1 \implies k=m-1\\
&= \frac{1}{(n+1)!}\left(\begin{array}{l}\displaystyle\sum_{k=0}^{k=n+1}(-1)^k\binom{n+1}{k}(x-k)^{n+1}\sigma(x-k)\\ \displaystyle-\sum_{m=1}^{m=n+2}(-1)^{m-1}\left[\binom{n+2}{m}-\binom{n+1}{m}\right](x-m)^{n+1}\sigma(x-m)\end{array}\right) && \begin{array}{l}\text{by \textit{Pascal's identity} /}\\ \textit{Stifel formula}\\ \text{(Theorem ?? page ??)}\end{array}\\
&= \frac{1}{(n+1)!}\left(\begin{array}{l}\displaystyle\sum_{m=1}^{m=n+2}(-1)^m\binom{n+2}{m}(x-m)^{n+1}\sigma(x-m)\\ \displaystyle-\sum_{m=1}^{m=n+2}(-1)^m\binom{n+1}{m}(x-m)^{n+1}\sigma(x-m)\\ \displaystyle+\sum_{k=0}^{k=n+1}(-1)^k\binom{n+1}{k}(x-k)^{n+1}\sigma(x-k)\end{array}\right) && \text{note } (-1)^{m-1}=-(-1)^m\\
&= \frac{1}{(n+1)!}\left(\begin{array}{l|ll}\displaystyle\sum_{m=0}^{m=n+2}(-1)^m\binom{n+2}{m}(x-m)^{n+1}\sigma(x-m) & \text{(A)} & \text{desired } n+1 \text{ case}\\ \displaystyle-\quad(-1)^0\binom{n+2}{0}(x-0)^{n+1}\sigma(x-0) & \text{(B)} & \text{cancelled by (F)}\\ \displaystyle-\sum_{m=1}^{m=n+1}(-1)^m\binom{n+1}{m}(x-m)^{n+1}\sigma(x-m) & \text{(C)} & \text{cancelled by (E)}\\ \displaystyle-\quad(-1)^{n+2}\binom{n+1}{n+2}(x-n-2)^{n+1}\sigma(x-n-2) & \text{(D)} & \binom{n+1}{n+2}=0 \text{ by Proposition ?? page ??}\\ \displaystyle+\sum_{k=1}^{k=n+1}(-1)^k\binom{n+1}{k}(x-k)^{n+1}\sigma(x-k) & \text{(E)} & \text{cancelled by (C)}\\ \displaystyle+\quad(-1)^0\binom{n+1}{0}(x-0)^{n+1}\sigma(x-0) & \text{(F)} & \binom{n+2}{0}=\binom{n+1}{0}=1, \text{ so (F) is cancelled by (B)}\end{array}\right)\\
&= \frac{1}{(n+1)!}\sum_{m=0}^{m=n+2}(-1)^m\binom{n+2}{m}(x-m)^{n+1}\sigma(x-m) && (n+1 \text{ case})
\end{aligned}
$$

BY-NC-ND Copyright © 2019 Daniel J. Greenhoe

Example R.4. [7] Let $\mathsf{N}_3(x)$ be the 3*rd order B-spline* (Definition R.2 page 367).[8]

EX

$$\mathsf{N}_3(x) = \frac{1}{6}\left\{\begin{array}{rl} x^3 & \text{for } 0 \le x \le 1 \\ -3x^3+12x^2-12x+4 & \text{for } 1 \le x \le 2 \\ 3x^3-24x^2+60x-44 & \text{for } 2 \le x \le 3 \\ -x^3+12x^2-48x+64 & \text{for } 3 \le x \le 4 \\ 0 & \text{otherwise} \end{array}\right\}$$



✎PROOF: This expression can be calculated "by hand" using Theorem R.1 (page 371) or by using the free and open source software package *Maxima* along with the script file listed in Section **??** (page **??**).

Example R.5. Let $\mathsf{N}_4(x)$ be the 4*th order B-spline* (Definition R.2 page 367).

EX

$$\mathsf{N}_4(x) = \frac{1}{24}\left\{\begin{array}{rl} x^4 & \text{for } 0 \le x \le= 1 \\ -4x^4+20x^3-30x^2+20x-5 & \text{for } 1 \le x \le= 2 \\ 6x^4-60x^3+210x^2-300x+155 & \text{for } 2 \le x \le= 3 \\ -4x^4+60x^3-330x^2+780x-655 & \text{for } 3 \le x \le= 4 \\ x^4-20x^3+150x^2-500x+625 & \text{for } 4 \le x \le= 5 \\ 0 & \text{otherwise} \end{array}\right\}$$

✎PROOF: This expression can be calculated "by hand" using Theorem R.1 (page 371) or by using the free and open source software package *Maxima* along with the script file listed in Section **??** (page **??**).

Example R.6. Let $\mathsf{N}_5(x)$ be the 5*th order B-spline* (Definition R.2 page 367).

EX

$$\mathsf{N}_5(x) = \frac{1}{120}\left\{\begin{array}{rl} x^5 & \text{for } 0 \le x \le= 1 \\ -5x^5+30x^4-60x^3+60x^2-30x+6 & \text{for } 1 \le x \le= 2 \\ 10x^5-120x^4+540x^3-1140x^2+1170x-474 & \text{for } 2 \le x \le= 3 \\ -10x^5+180x^4-1260x^3+4260x^2-6930x+4386 & \text{for } 3 \le x \le= 4 \\ 5x^5-120x^4+1140x^3-5340x^2+12270x-10974 & \text{for } 4 \le x \le= 5 \\ -x^5+30x^4-360x^3+2160x^2-6480x+7776 & \text{for } 5 \le x \le= 6 \\ 0 & \text{otherwise} \end{array}\right\}$$

The 5*th order B-spline* $\mathsf{N}_5(x)$ is illustrated in Figure R.1 (page 369).

✎PROOF: This expression can be calculated "by hand" using Theorem R.1 (page 371) or by using the free and open source software package *Maxima* along with the script file listed in Section **??** (page **??**).

Example R.7. Let $\mathsf{N}_6(x)$ be the 6*th order B-spline* (Definition R.2 page 367).

EX

$$\mathsf{N}_6(x) = \frac{1}{720}\left\{\begin{array}{rl} x^6 & \text{for } 0 \le x \le 1 \\ -6x^6+42x^5-105x^4+140x^3-105x^2+42x-7 & \text{for } 1 \le x \le 2 \\ 15x^6-210x^5+1155x^4-3220x^3+4935x^2-3990x+1337 & \text{for } 2 \le x \le 3 \\ -20x^6+420x^5-3570x^4+15680x^3-37590x^2+47040x-24178 & \text{for } 3 \le x \le 4 \\ 15x^6-420x^5+4830x^4-29120x^3+96810x^2-168000x+119182 & \text{for } 4 \le x \le 5 \\ -6x^6+210x^5-3045x^4+23380x^3-100065x^2+225750x-208943 & \text{for } 5 \le x \le 6 \\ x^6-42x^5+735x^4-6860x^3+36015x^2-100842x+117649 & \text{for } 6 \le x \le 7 \\ 0 & \text{otherwise} \end{array}\right\}$$

The 6*th order B-spline* $\mathsf{N}_6(x)$ is illustrated in Figure R.1 (page 369).

✎PROOF: This expression can be calculated "by hand" using Theorem R.1 (page 371) or by using the free and open source software package *Maxima* along with the script file listed in Section **??** (page **??**).

[7] Schumaker (2007) page 136 ⟨Table 1⟩, Shizgal (2015) page 92 ⟨(2.199)⟩, Szabó and Horváth (2004) page 146 ⟨(4)⟩, Wei and Billings (2006) page 578 ⟨Table 1⟩, Maleknejad et al. (2013) ⟨(9)⟩

[8] For help with plotting *B-splines*, see APPENDIX S (page 399).

COPYRIGHT © 2019 DANIEL J. GREENHOE

Example R.8. Let $\mathsf{N}_7(x)$ be the 7*th order B-spline* (Definition R.2 page 367).

EX

$$7!\mathsf{N}_7(x) = 5040\mathsf{N}_7(x) =$$
$$\left\{\begin{array}{rrrrrrrrrl}
x^7 & & & & & & & & & \text{for } 0 \le x \le 1 \\
-7x^7 & +56x^6 & -168x^5 & +280x^4 & -280x^3 & +168x^2 & -56x & +8 & & \text{for } 1 \le x \le 2 \\
21x^7 & -336x^6 & +2184x^5 & -7560x^4 & +15400x^3 & -18648x^2 & +12488x & -3576 & & \text{for } 2 \le x \le 3 \\
-35x^7 & +840x^6 & -8400x^5 & +45360x^4 & -143360x^3 & +267120x^2 & -273280x & +118896 & & \text{for } 3 \le x \le 4 \\
35x^7 & -1120x^6 & +15120x^5 & -111440x^4 & +483840x^3 & -1238160x^2 & +1733760x & -1027984 & & \text{for } 4 \le x \le 5 \\
-21x^7 & +840x^6 & -14280x^5 & +133560x^4 & -741160x^3 & +2436840x^2 & -4391240x & +3347016 & & \text{for } 5 \le x \le 6 \\
7x^7 & -336x^6 & +6888x^5 & -78120x^4 & +528920x^3 & -2135448x^2 & +4753336x & -4491192 & & \text{for } 6 \le x \le 7 \\
-x^7 & +56x^6 & -1344x^5 & +17920x^4 & -143360x^3 & +688128x^2 & -1835008x & +2097152 & & \text{for } 7 \le x \le 8 \\
 & & & & & & & & 0 & \text{otherwise}
\end{array}\right\}$$

The 7*th order B-spline* $\mathsf{N}_7(x)$ is illustrated in Figure R.1 (page 369).

✎PROOF: This expression can be calculated "by hand" using Theorem R.1 (page 371) or by using the free and open source software package *Maxima* along with the script file listed in Section **??** (page **??**). ✏

Example R.9. [9] The $(n+1)^2$ coefficients of the order $n, n-1, \ldots, 0$ monomials of each *B-spline* $\mathsf{N}_n(x)$ multiplied by $n!$ induce an *integer sequence*
$\mathbb{x} \triangleq (\!(1,1,0,-1,2,1,0,0,-2,6,-3,1,-6,9,1,0,0,0,-3,12,-12,4,3,-24,60,-44,-1,12,-48,64,\ldots)\!)$
as more fully listed in Table R.1 (page 398). In this sequence $\mathbb{x} \triangleq (\!(x_0, x_1, x_2, \ldots)\!)$, the coefficients for the *order n B-spline* $\mathsf{N}_n(x)$ begin at the sequence index value

$$p \triangleq \sum_{k=0}^{n} k^2 = \frac{1}{6}n(n+1)(2n+1) \qquad \text{and end at index value } p + (n+1)^2 - 1.$$

For example, the coefficients for $\mathsf{N}_3(x)$ begin at index value $p \triangleq 0+1+4+9 = 14$ and end at index value $p + 4^2 - 1 = 29$. Using these coefficients gives the following expression for $\mathsf{N}_3(x)$:

$$\mathsf{N}_3(x) = \left[\begin{array}{rrrr|l}
1 & 0 & 0 & 0 & \text{for } 0 \le x < 1 \\
-3 & 12 & -12 & 4 & \text{for } 1 \le x < 2 \\
3 & -24 & 60 & -44 & \text{for } 2 \le x < 3 \\
-1 & 12 & -48 & 64 & \text{for } 3 \le x < 4
\end{array}\right]\left[\begin{array}{c} x^3 \\ x^2 \\ x \\ 1 \end{array}\right] = \left\{\begin{array}{rrrrl}
x^3 & & & & \text{for } 0 \le x < 1 \\
-3x^3 & +12x^2 & -12x & +4 & \text{for } 1 \le x < 2 \\
3x^3 & -24x^2 & +60x & -44 & \text{for } 2 \le x < 3 \\
-x^3 & +12x^2 & -48x & +64 & \text{for } 3 \le x < 4 \\
 & & & 0 & \text{otherwise}
\end{array}\right.$$

…which agrees with the result presented in Example R.4 (page 373).

✎PROOF:

1. The coefficients for the sequence $\mathbb{x}$ may be computed with assistance from *Maxima* together with the script file listed in Section **??** (page **??**).

2. Proof that $\sum_{k=0}^{n} k^2 = \frac{1}{6}n(n+1)(2n+1)$: The summation is a *power sum*. The relation may be proved using *induction*.[10]

   (a) Base case: n=0 case …

$$\begin{aligned}
\sum_{k=0}^{k=0} k^2 &= 0 \\
&= \frac{0(0+1)(2\cdot 0+1)}{6} \\
&= \left.\frac{n(n+1)(2n+1)}{6}\right|_{n=0}
\end{aligned}$$

[9] Greenhoe (2017b)
[10] Greenhoe (2017a), pages 186–187 ⟨Proposition 11.2 (Power Sums)⟩

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

(b) Base case: n=1 case …

$$\begin{aligned}\sum_{k=0}^{k=1} k^2 &= 0+1\\ &= \frac{1(1+1)(2\cdot 1+1)}{6}\\ &= \left.\frac{n(n+1)(2n+1)}{6}\right|_{n=1}\end{aligned}$$

(c) inductive step—proof that n case $\implies$ $n+1$ case:

$$\begin{aligned}\sum_{k=0}^{n+1} k^2 &= \left(\sum_{k=0}^{n} k^2\right) + \left(n+1\right)^2\\ &= \left(\frac{n(n+1)(2n+1)}{6}\right) + \left(n+1\right)^2 && \text{by } n \text{ case hypothesis}\\ &= \left(n+1\right)\left(\frac{n(2n+1)+6(n+1)}{6}\right)\\ &= \left(n+1\right)\left(\frac{2n^2+7n+6}{6}\right)\\ &= \left(n+1\right)\left(\frac{(n+2)(2n+3)}{6}\right)\\ &= \frac{(n+1)[(n+1)+1][2(n+1)+1]}{6}\end{aligned}$$

⇨

Theorem R.2. [11]

THM $\frac{\mathsf{d}}{\mathsf{d}x}\mathsf{N}_n(x) = \mathsf{N}_{n-1}(x) - \mathsf{N}_{n-1}(x-1) \qquad \forall n \in \mathbb{N}, \forall x \in \mathbb{R}$

✎PROOF:

1. Proof using Lemma R.2 (page 367) and the *Fundamental Theorem of Calculus*:

$$\begin{aligned}\frac{\mathsf{d}}{\mathsf{d}x}\mathsf{N}_n(x) &= \frac{\mathsf{d}}{\mathsf{d}x}\int_0^1 \mathsf{N}_{n-1}(x-\tau)\,\mathsf{d}\tau && \text{by Lemma R.2 page 367}\\ &= \frac{\mathsf{d}}{\mathsf{d}x}\int_{x-u=0}^{x-u=1} \mathsf{N}_{n-1}(u)(-1)\,\mathsf{d}u && \text{where } u \triangleq x-\tau \implies \tau = x-u\\ &= \frac{\mathsf{d}}{\mathsf{d}x}\int_{u=x-1}^{u=x} \mathsf{N}_{n-1}(u)\,\mathsf{d}u\\ &= \left\{\frac{\mathsf{d}}{\mathsf{d}x}\int \mathsf{N}_{n-1}(u)\,\mathsf{d}u\Big|_{u=x}\right\} - \left\{\frac{\mathsf{d}}{\mathsf{d}x}\int \mathsf{N}_{n-1}(u)\,\mathsf{d}u\Big|_{u=x-1}\right\} && \text{by } \textit{Fundamental Theorem of Calculus}^{12}\\ &= \left\{\mathsf{N}_{n-1}(x)\frac{\mathsf{d}}{\mathsf{d}x}(x)\right\} - \left\{\mathsf{N}_{n-1}(x-1)\frac{\mathsf{d}}{\mathsf{d}x}(x-1)\right\} && \text{by } \textit{Chain Rule}^{13}\\ &= \mathsf{N}_{n-1}(x) - \mathsf{N}_{n-1}(x-1)\end{aligned}$$

[11] Höllig (2003) page 25 ⟨3.2⟩, Schumaker (2007) page 121 ⟨Theorem 4.16⟩
[12] Hijab (2011) page 163 ⟨Theorem 4.4.3⟩
[13] Hijab (2011) pages 73–74 ⟨Theorem 3.1.2⟩

COPYRIGHT © 2019 DANIEL J. GREENHOE

2. Proof using Lemma R.2 (page 367) and *induction*:

   (a) Base case …proof for $n = 1$ case:

$$\begin{aligned}
\mathsf{N}_0(x) - \mathsf{N}_0(x-1) &= \underbrace{\sigma(x) - \sigma(x-1)}_{\mathsf{N}_0(x)} - \underbrace{[\sigma(x-1) - \sigma(x-2)]}_{\mathsf{N}_0(x-1)} && \text{by Example R.1 page 369}\\
&= \sigma(x) - 2\sigma(x-1) + \sigma(x-2)\\
&= \sum_{k=0}^{2}(-1)^k\binom{2}{k}\sigma(x-k)\\
&= \frac{\mathsf{d}}{\mathsf{d}x}\sum_{k=0}^{2}(-1)^k\binom{2}{k}(x-k)\sigma(x-k)\\
&= \frac{\mathsf{d}}{\mathsf{d}x}\mathsf{N}_1(x) && \text{by Example R.2 page 370}
\end{aligned}$$

   (b) Base case …proof for $n = 2$ case:

$$\begin{aligned}
&\mathsf{N}_1(x) - \mathsf{N}_1(x-1)\\
&= \underbrace{x\sigma(x) - 2(x-1)\sigma(x-1) + (x-2)\sigma(x-2)}_{\mathsf{N}_1(x)}\\
&\quad - \underbrace{[(x-1)\sigma(x-1) - 2(x-2)\sigma(x-2) + (x-3)\sigma(x-3)]}_{\mathsf{N}_1(x-1)} && \text{by Example R.2 page 370}\\
&= x\sigma(x) + [-2x+2-x+1]\sigma(x-1) + [x-2+2x-4]\sigma(x-2) + [-x+3]\sigma(x-3)\\
&= x\sigma(x) + [-3x+3]\sigma(x-1) + [3x-6]\sigma(x-2) + [-x+3]\sigma(x-3)\\
&= \frac{\mathsf{d}}{\mathsf{d}x}\left\{\begin{array}{l}\frac{1}{2}x^2\sigma(x) + \left[-\frac{3}{2}x^2 + 3x - \frac{1}{2}\right]\sigma(x-1) + \left[\frac{3}{2}x^2 - 6x + 3\right]\sigma(x-2)\\ \qquad + \left[-\frac{1}{2}x^2 + 3x - \frac{5}{2}\right]\sigma(x-3)\end{array}\right\}\\
&= \frac{\mathsf{d}}{\mathsf{d}x}\mathsf{N}_2(x) && \text{by Example R.3 page 370}
\end{aligned}$$

   (c) Proof that n case $\implies$ $n+1$ case:

$$\begin{aligned}
\frac{\mathsf{d}}{\mathsf{d}x}\mathsf{N}_{n+1}(x) &= \frac{\mathsf{d}}{\mathsf{d}x}\int_0^1 \mathsf{N}_n(x-\tau)\,\mathsf{d}\tau && \text{by Lemma R.2 page 367}\\
&= \int_0^1 \frac{\partial}{\partial x}\mathsf{N}_n(x-\tau)\,\mathsf{d}\tau && \text{by \textit{Leibniz Integration Rule} (Theorem ?? page ??)}\\
&= \int_0^1 \left[\mathsf{N}_{n-1}(x-\tau) - \mathsf{N}_{n-1}(x-1-\tau)\right]\mathsf{d}\tau && \text{by left hypothesis}\\
&= \int_0^1 \mathsf{N}_{n-1}(x-\tau)\,\mathsf{d}\tau - \int_0^1 \mathsf{N}_{n-1}(x-1-\tau)\,\mathsf{d}\tau\\
&= \mathsf{N}_n(x) - \mathsf{N}_n(x-1) && \text{by Lemma R.2 page 367}
\end{aligned}$$

✏

Theorem R.3 (*B-spline recursion*). [14] *Let* $\mathsf{N}_n(x)$ *be the* nTH ORDER B-SPLINE *(Definition R.2 page 367)*.

THM

$$\mathsf{N}_n(x) = \frac{x}{n}\mathsf{N}_{n-1}(x) + \frac{n+1-x}{n}\mathsf{N}_{n-1}(x-1) \quad \forall n \in \{1,2,3,\ldots\},\ \forall x \in \mathbb{R}$$

[14] Christensen (2008) page 140 ⟨Theorem 6.1.1⟩, Höllig (2003) page 27 ⟨3.4⟩, Schumaker (2007) page 120 ⟨Theorem 4.15⟩, de Boor (2001) page 90 ⟨B-Spline Property (i)⟩, Chui (1988) page 2 ⟨Theorem 1.1⟩, Wojtaszczyk (1997) page 53 ⟨Theorem 3.7⟩, Cox (1972), de Boor (1972)

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

✎PROOF:

1. Base case …proof for $n = 1$ case:

$$\frac{x}{1}\mathsf{N}_0(x) + \frac{1+1-x}{1}\mathsf{N}_0(x-1) = \frac{x}{1}\underbrace{[\sigma(x)-\sigma(x-1)]}_{\mathsf{N}_0(x)} + \frac{1+1-x}{1}\underbrace{[\sigma(x-1)-\sigma(x-2)]}_{\mathsf{N}_0(x-1)}$$
$$= x\sigma(x) + [-x-x+2]\sigma(x-1) + [x-2]\sigma(x-2)$$
$$= \mathsf{N}_1(x) \qquad \text{by Example R.2 page 370}$$

2. Induction step …proof that n case $\implies$ $n+1$ case:

$$\frac{x}{n+1}\mathsf{N}_n(x) + \frac{n+2-x}{n+1}\mathsf{N}_n(x-1) + c_1$$
$$= \int \frac{\mathsf{d}}{\mathsf{d}x}\left\{\frac{x}{n+1}\mathsf{N}_n(x) + \frac{n+2-x}{n+1}\mathsf{N}_n(x-1)\right\}\mathsf{d}x$$
$$= \int \underbrace{\frac{1}{n+1}\mathsf{N}_n(x) + \frac{x}{n+1}\frac{\mathsf{d}}{\mathsf{d}x}\mathsf{N}_n(x)}_{\frac{\mathsf{d}}{\mathsf{d}x}\frac{x}{n+1}\mathsf{N}_n(x)} + \underbrace{\frac{-1}{n+1}\mathsf{N}_n(x-1) + \frac{n+2-x}{n}\frac{\mathsf{d}}{\mathsf{d}x}\mathsf{N}_n(x-1)}_{\frac{\mathsf{d}}{\mathsf{d}x}\frac{n+2-x}{n+1}\mathsf{N}_n(x-1)}\,\mathsf{d}x$$

by *product rule*

$$= \int \frac{1}{n+1}\underbrace{\left[\frac{x}{n}\mathsf{N}_{n-1}(x) + \frac{n+1-x}{n}\mathsf{N}_{n-1}(x-1)\right]}_{\text{by } n \text{ hypothesis}} + \frac{x}{n+1}\underbrace{\left[\mathsf{N}_{n-1}(x) - \mathsf{N}_{n-1}(x-1)\right]}_{\text{by Theorem R.2 page 375}}$$
$$- \underbrace{\left[\frac{x-1}{n^2+n}\mathsf{N}_{n-1}(x-1) + \frac{n-x+2}{n(n+1)}\mathsf{N}_{n-1}(x-2)\right]}_{\text{by induction hypothesis}}$$
$$+ \frac{n+2-x}{n+1}\underbrace{\left[\mathsf{N}_{n-1}(x-1) - \mathsf{N}_{n-1}(x-2)\right]}_{\text{by Theorem R.2 page 375}}\mathsf{d}x$$
$$= \int \left[\frac{x}{n(n+1)} + \frac{x}{n+1}\right]\mathsf{N}_{n-1}(x) + \left[\frac{n-x+1}{n(n+1)} - \frac{x-1}{n(n+1)} + \frac{n+2-2x}{n+1}\right]\mathsf{N}_{n-1}(x-1)$$
$$+ \left[\frac{-n-2+x}{n(n+1)} + \frac{-n-2+x}{n+1}\right]\mathsf{N}_{n-1}(x-2)\,\mathsf{d}x$$
$$= \int \left[\frac{x+nx}{n(n+1)}\right]\mathsf{N}_{n-1}(x) + \left[\frac{n+2-2x+n(n+2-2x)}{n(n+1)}\right]\mathsf{N}_{n-1}(x-1)$$
$$+ \left[\frac{-n-2+x+n(-n-2+x)}{n(n+1)}\right]\mathsf{N}_{n-1}(x-2)\,\mathsf{d}x$$
$$= \int \left[\frac{x}{n}\right]\mathsf{N}_{n-1}(x) + \left[\frac{n+2-2x}{n}\right]\mathsf{N}_{n-1}(x-1) + \left[\frac{-n-2+x}{n}\right]\mathsf{N}_{n-1}(x-2)\,\mathsf{d}x$$
$$= \int \underbrace{\left[\frac{x}{n}\right]\mathsf{N}_{n-1}(x) + \left[\frac{n+1-x}{n}\right]\mathsf{N}_{n-1}(x-1)}_{\mathsf{N}_n(x)}$$
$$- \underbrace{\left[\frac{x-1}{n}\right]\mathsf{N}_{n-1}(x-1) - \left[\frac{n+2-x}{n}\right]\mathsf{N}_{n-1}(x-2)}_{\mathsf{N}_{n-1}(x-1)}\,\mathsf{d}x$$
$$= \int \mathsf{N}_n(x) - \mathsf{N}_n(x-1)\,\mathsf{d}x \qquad \text{by } n \text{ hypothesis}$$
$$= \int \frac{\mathsf{d}}{\mathsf{d}x}\mathsf{N}_{n+1}(x)\,\mathsf{d}x \qquad \text{by Theorem R.2 page 375}$$
$$= \mathsf{N}_{n+1}(x) + c_2$$

COPYRIGHT © 2019 DANIEL J. GREENHOE

Proof that $c_1 = c_2$: By item (2) (page 378), $\mathsf{N}_n(x) = 0$ for $x < 0$. Therefore, $c_1 = c_2$.

✏

Theorem R.4 (*B-spline general form*). [15] *Let* $\mathsf{N}_n(x)$ *be the* nTH ORDER B-SPLINE *(Definition R.2 page 367). Let* $\mathsf{supp}\,\mathsf{f}$ *be the* SUPPORT *of a function* $\mathsf{f} \in \mathbb{R}^{\mathbb{R}}$.

THM

1.	$\mathsf{N}_n(x)$	$\geq$	0	$\forall n \in \mathbb{W}, \quad \forall x \in \mathbb{R}$	(NON-NEGATIVE)
2.	$\mathsf{supp}\mathsf{N}_n(x)$	$=$	$[0 : n+1]$	$\forall n \in \mathbb{W}$	(CLOSED SUPPORT)
3.	$\int_{\mathbb{R}} \mathsf{N}_n(x)\,\mathrm{d}x$	$=$	1	$\forall n \in \mathbb{W}$	(UNIT AREA)
4.	$\mathsf{N}_n\left(\frac{n+1}{2} - x\right)$	$=$	$\mathsf{N}_n\left(\frac{n+1}{2} + x\right)$	$\forall n \in \mathbb{N} \quad \forall x \in \mathbb{R}$	(SYMMETRIC *about* $x = \frac{n+1}{2}$)

✎PROOF:

1. Proof that $\mathsf{N}_n(x) \geq 0$ (proof by induction):

   (a) base case…proof that $\mathsf{N}_0(x) \geq 0$:

   $$\begin{aligned} \mathsf{N}_0(x) &\triangleq \mathbb{1}_{[0:1)}(x) && \text{by definition of } \mathsf{N}_0(x) && \text{(Definition R.2 page 367)} \\ &\geq 0 && \text{by definition of } \mathbb{1}_A(x) && \text{(Definition ?? page ??)} \end{aligned}$$

   (b) inductive step—proof that $\{\mathsf{N}_n(x) \geq 0\} \implies \{\mathsf{N}_{n+1}(x) \geq 0\}$:

   $$\begin{aligned} \mathsf{N}_{n+1}(x) &= \int_{\tau=0}^{\tau=1} \mathsf{N}_n(x-\tau)\,\mathrm{d}\tau && \text{by Lemma R.2 page 367} \\ &\geq 0 && \text{by induction hypothesis } (\mathsf{N}_n(x) \geq 0) \end{aligned}$$

2. Proof that $\mathsf{supp}\mathsf{N}_n(x) = [0 : n+1]$ (proof by induction):

   (a) Base case …proof that $\mathsf{supp}\mathsf{N}_0 = [0 : 1]$:

   $$\begin{aligned} \mathsf{supp}\mathsf{N}_0 &\triangleq \mathsf{supp}\mathbb{1}_{[0:1)} && \text{by definition of } \mathsf{N}_0(x) && \text{(Definition R.2 page 367)} \\ &= \{[0 : 1)\}^- && \text{by definition of } \textit{support} \text{ operator} \\ &= [0 : 1] && \text{by definition of } \textit{closure} \text{ operator} \end{aligned}$$

   (b) Induction step …proof that $\{\mathsf{supp}\mathsf{N}_n = [0 : n+1]\} \implies \{\mathsf{supp}\mathsf{N}_{n+1} = [0 : n+2]\}$:

   $$\begin{aligned} \mathsf{supp}\mathsf{N}_{n+1}(x) &= \mathsf{supp} \int_{\tau=0}^{\tau=1} \mathsf{N}_n(x-\tau)\,\mathrm{d}\tau && \text{by Lemma R.2 page 367} \\ &= \mathsf{supp} \int_{[0:1]} \mathsf{N}_n(x-\tau)\,\mathrm{d}\tau && \text{by def. of } \textit{Lebesgue integration} \\ &= \left\{x \in \mathbb{R} \middle| (x-\tau) \in [0 : n+1] \text{ for some } \tau \in [0 : 1]\right\}^- && \text{by induction hypothesis} \\ &= [0 : n+1] \cup [0+1 : n+1+1]^- \\ &= [0 : n+2]^- \\ &= [0 : n+2] && \text{by property of } \textit{closure} \text{ operator} \end{aligned}$$

3. Proof that $\int_{\mathbb{R}} \mathsf{N}_n(x)\,\mathrm{d}x = 1$ (proof by induction):

[15] Christensen (2008) page 140 ⟨Theorem 6.1.1⟩, Höllig (2003) page 27 ⟨3.4⟩, Schumaker (2007) page 120 ⟨Theorem 4.15⟩, de Boor (2001) page 90 ⟨B-Spline Property (i)⟩, Chui (1988) page 2 ⟨Theorem 1.1⟩, Wojtaszczyk (1997) page 53 ⟨Theorem 3.7⟩, Cox (1972), de Boor (1972), Prasad and Iyengar (1997) pages 145–146 ⟨Theorem 6.2⟩

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

(a) Base case …proof that $\int_{\mathbb{R}} \mathsf{N}_0(x) = 1$:

$$\begin{aligned}
\int_{\mathbb{R}} \mathsf{N}_0(x)\,\mathrm{d}x &= \int_{\mathbb{R}} \mathbb{1}_{[0:1)}\,\mathrm{d}x && \text{by definition of } \mathsf{N}_0(x) && \text{(Definition R.2 page 367)}\\
&= \int_{[0:1)} 1\,\mathrm{d}x && \text{by definition of } \mathbb{1}_A(x) && \text{(Definition ?? page ??)}\\
&= \int_{[0:1]} 1\,\mathrm{d}x && \text{by property of } \textit{Lebesgue integration}\\
&= 1
\end{aligned}$$

(b) Induction step …proof that $\left\{\int_{\mathbb{R}} \mathsf{N}_n(x) = 1\right\} \implies \left\{\int_{\mathbb{R}} \mathsf{N}_{n+1} = 1\right\}$:

$$\begin{aligned}
\int_{\mathbb{R}} \mathsf{N}_{n+1}(x)\,\mathrm{d}x &= \int_{\mathbb{R}}\int_0^1 \mathsf{N}_n(x-\tau)\,\mathrm{d}\tau\,\mathrm{d}x && \text{by Lemma R.2 page 367}\\
&= \int_0^1\int_{\mathbb{R}} \mathsf{N}_n(x-\tau)\,\mathrm{d}x\,\mathrm{d}\tau\\
&= \int_0^1\int_{\mathbb{R}} \mathsf{N}_n(u)\,\mathrm{d}u\,\mathrm{d}\tau && \text{where } u \triangleq x-\tau \implies \tau = x-u\\
&= \int_0^1 1\,\mathrm{d}\tau && \text{by induction hypothesis}\\
&= 1
\end{aligned}$$

4. Proof that $\mathsf{N}_n(x)$ is *symmetric* for $n \in \{1,2,3,\ldots\}$:

(a) Note that $\mathsf{N}_0(x)$ $(n=0)$ is *not symmetric* (in particular it fails at $x = 1/2$) because
$\mathsf{N}_0\left(\frac{0+1}{2} - \frac{1}{2}\right) = \mathsf{N}_0(0) = 1 \neq 0 = \mathsf{N}_1(1) = \mathsf{N}_0\left(\frac{0+1}{2} + \frac{1}{2}\right)$

(b) Base case …proof for $n = 1$ case:

$$\begin{aligned}
\mathsf{N}_1\left(\frac{1+1}{2} - x\right) &= \mathsf{N}_1(1-x)\\
&= \left\{\begin{array}{rl} (1-x) & \text{for } 1-x \in [0:1]\\ -(1-x)+2 & \text{for } 1-x \in [1:2]\\ 0 & \text{otherwise}\end{array}\right\} && \text{by Example R.2 page 370}\\
&= \left\{\begin{array}{rl} -x+1 & \text{for } -x \in [-1:0]\\ x+1 & \text{for } -x \in [0:1]\\ 0 & \text{otherwise}\end{array}\right\}\\
&= \left\{\begin{array}{rl} x+1 & \text{for } x \in [-1:0]\\ -x+1 & \text{for } x \in [0:1]\\ 0 & \text{otherwise}\end{array}\right\}\\
&= \left\{\begin{array}{rl} (1+x) & \text{for } 1+x \in [0:1]\\ -(1+x)+2 & \text{for } 1+x \in [1:2]\\ 0 & \text{otherwise}\end{array}\right\}\\
&= \mathsf{N}_1(1+x) && \text{by Example R.2 page 370}\\
&= \mathsf{N}_1\left(\frac{1+1}{2} + x\right)
\end{aligned}$$

Copyright © 2019 Daniel J. Greenhoe

(c) Induction step …proof that $n-1$ case $\Longrightarrow$ n case:

$$\begin{aligned}
&\mathsf{N}_n\left(\frac{n+1}{2}+x\right)\\
&=\frac{\frac{n+1}{2}+x}{n}\mathsf{N}_{n-1}\left(\frac{n+1}{2}+x\right)+\frac{n+1-\left(\frac{n+1}{2}+x\right)}{n}\mathsf{N}_{n-1}\left(\frac{n+1}{2}+x-1\right) && \text{by Theorem R.3 page 376}\\
&=\frac{n+1-\left(\frac{n+1}{2}-x\right)}{n}\mathsf{N}_{n-1}\left(\frac{n}{2}+\left[x+\frac{1}{2}\right]\right)+\frac{\frac{n+1}{2}-x}{n}\mathsf{N}_{n-1}\left(\frac{n}{2}+\left[x-\frac{1}{2}\right]\right)\\
&=\frac{n+1-\left(\frac{n+1}{2}-x\right)}{n}\mathsf{N}_{n-1}\left(\frac{n}{2}-\left[x+\frac{1}{2}\right]\right)+\frac{\frac{n+1}{2}-x}{n}\mathsf{N}_{n-1}\left(\frac{n}{2}-\left[x-\frac{1}{2}\right]\right) && \text{by induction hypothesis}\\
&=\frac{n+1-\left(\frac{n+1}{2}-x\right)}{n}\mathsf{N}_{n-1}\left(\left[\frac{n+1}{2}-x\right]-1\right)+\frac{\frac{n+1}{2}-x}{n}\mathsf{N}_{n-1}\left(\frac{n+1}{2}-x\right)\\
&=\mathsf{N}_n\left(\frac{n+1}{2}-x\right) && \text{by Theorem R.3 page 376}
\end{aligned}$$

✏

R.3 Projection properties

In the case where $(\!(\mathsf{N}_n(x-k))\!)_{k\in\mathbb{Z}}$ is to be used as a basis in some subspace of $\boldsymbol{L}^2_{\mathbb{R}}$, one may want to *project* a function $\mathsf{f}(x)$ onto a basis function $\mathsf{N}_n(x-k)$. This is especially true when $(\!(\mathsf{N}_n(x-k))\!)$ is *orthogonal*; but in the case of *B-splines* this is only true when $n=0$ (Theorem R.8 page 390). Nevertheless, projection of a function onto $\mathsf{N}_n(x-k)$, or the projection of $\mathsf{N}_n(x)$ onto another basis fucntion (such as the complex exponential in the case of *Fourier analysis* as in Lemma R.5 page 382), is still useful. Projection in an *inner product space* is typically performed using the *inner product* $\langle \mathsf{f}(x) \mid \mathsf{N}_n(x-k)\rangle$; and in the space $\boldsymbol{L}^2_{\mathbb{R}}$, this inner product is typically defined as an *integral* such that

$$\langle \mathsf{f}(x) \mid \mathsf{N}_n(x-k)\rangle \triangleq \int_{\mathbb{R}} \mathsf{f}(x)\mathsf{N}_n(x-k)\,\mathrm{d}x.$$

As it turns out, there is a way to compute this inner product that only involves the function $\mathsf{f}(x)$ and the order parameter n (next theorem).

Theorem R.5. [16] *Let* f *be a* CONTINUOUS *function in* $\boldsymbol{L}^2_{\mathbb{R}}$ *and* $\mathsf{f}^{(n)}$ *the* nTH DERIVATIVE *of* $\mathsf{f}(x)$.

THM

(1). $$\int_{\mathbb{R}} \mathsf{f}(x)\mathsf{N}_n(x)\,\mathrm{d}x = \int_{[0:1)^{n+1}} \mathsf{f}(x_1+x_2+\cdots+x_{n+1})\,\mathrm{d}x_1\,\mathrm{d}x_2\cdots\mathrm{d}x_{n+1}$$

(2). $$\int_{\mathbb{R}} \mathsf{f}^{(n)}(x)\mathsf{N}_n(x)\,\mathrm{d}x = \sum_{k=0}^{n+1}(-1)^{n-k+1}\binom{n+1}{k}\mathsf{f}(k)$$

✎PROOF:

1. Proof for (1) (proof by induction):

   (a) Base case …proof for $n=0$ case:

$$\int_{\mathbb{R}} \mathsf{f}(x)\mathsf{N}_0(x)\,\mathrm{d}x = \int_{[0:1)} \mathsf{f}(x)\,\mathrm{d}x \qquad \text{by definition of } \mathsf{N}_0(x) \qquad \text{(Definition R.2 page 367)}$$

[16] Chui (1992) page 85 ⟨(4.2.2), (4.2.3)⟩, Christensen (2008) page 140 ⟨Theorem 6.1.1⟩

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

(b) Inductive step—proof that n case $\implies$ $n+1$ case:

$$\begin{aligned}
&\int_{\mathbb{R}} \mathsf{f}(x)\mathsf{N}_{n+1}(x)\,\mathrm{d}x \\
&= \int_{\mathbb{R}} \left[\int_0^1 \mathsf{N}_n(x-\tau)\,\mathrm{d}\tau\right]\mathsf{f}(x)\,\mathrm{d}x && \text{by Lemma R.2 page 367} \\
&= \int_{[0:1)}\int_{\mathbb{R}} \mathsf{N}_n(x-\tau)\mathsf{f}(x)\,\mathrm{d}x\,\mathrm{d}\tau \\
&= \int_{[0:1)}\int_{\mathbb{R}} \mathsf{N}_n(u)\mathsf{f}(u+\tau)\,\mathrm{d}u\,\mathrm{d}\tau && \text{where } u \triangleq x-\tau \implies x = u+\tau \\
&= \int_{[0:1)}\int_{[0:1)^{n+1}} \mathsf{f}(u_1+u_2+\cdots+u_{n+1}+\tau)\,\mathrm{d}u_1\ \mathrm{d}u_2\ \cdots\ \mathrm{d}u_{n+1}\ \mathrm{d}\tau && \text{by induction hypothesis} \\
&= \int_{[0:1)^{n+2}} \mathsf{f}(u_1+u_2+\cdots+u_{n+1}+u_{n+2})\,\mathrm{d}u_1\ \mathrm{d}u_2\ \cdots\ \mathrm{d}u_{n+2}\ \mathrm{d}\tau \\
&= \int_{[0:1)^{n+2}} \mathsf{f}(x_1+x_2+\cdots+x_{n+1}+x_{n+2})\,\mathrm{d}x_1\ \mathrm{d}x_2\ \cdots\ \mathrm{d}x_{n+2} && \text{by change of variables } u_k \to x_k
\end{aligned}$$

2. Proof for (2):

$$\begin{aligned}
\int_{\mathbb{R}} \mathsf{f}^{(n)}(x)\mathsf{N}_n(x)\,\mathrm{d}x &= \int_{[0:1)^{n+1}} \mathsf{f}^{(n)}\left(\sum_{k=1}^{n+1} x_k\right)\mathrm{d}x_1\ \mathrm{d}x_2\cdots\ \mathrm{d}x_{n+1} && \text{by (1)} \\
&= \sum_{k=0}^{n+1}(-1)^{n-k+1}\binom{n+1}{k}\mathsf{f}(k) && \text{by Theorem ?? page ??}
\end{aligned}$$



Figure R.2: Zero locations for B-cardinal spline $N_n(x)$ scaling coefficients

R.4 Fourier analysis

Simply put, no matter what new and fancy basis sequences are discovered, the *Fourier transform* never goes out of style. This is largely because the *kernel* of the Fourier transform—the *complex exponential* function—has two properties that makes it extremely special:

- The complex exponential is an eigenvalue of any *linear time invariant* (*LTI*) operator (Theorem ?? page ??).
- The complex exponential generates a *continuous point spectrum* for the *differential operator*.

Thus, we might expect the projection of the *B-spline* function $\mathsf{N}_n(x)$ onto the complex exponential (essentially the *Fourier transform* of $\mathsf{N}_n(x)$,…next lemma) to be useful. Such a hunch would be confirmed because it is useful for proving that

Copyright © 2019 Daniel J. Greenhoe

- the sequence $(\!(\mathsf{N}_n(x-k))\!)_{k\in\mathbb{Z}}$ is a *Riesz basis* (Lemma R.6 page 385, Theorem R.8 page 390) and
- the sequence $(\!(\mathsf{N}_n(x-k))\!)_{k\in\mathbb{Z}}$ is a *multiresolution analysis* (Theorem R.10 page 393).

Lemma R.5. [17] *Let* $\tilde{\mathbf{F}}$ *be the* FOURIER TRANSFORM *operator* (Definition O.2 page 333).

LEM

$$\tilde{\mathbf{F}}\mathsf{N}_n(\omega) = \frac{1}{\sqrt{2\pi}}\left(\frac{1-e^{-i\omega}}{i\omega}\right)^{n+1} = \frac{1}{\sqrt{2\pi}}e^{-i\frac{(n+1)\omega}{2}}\left(\frac{\sin(\omega/2)}{\omega/2}\right)^{n+1} \triangleq \frac{1}{\sqrt{2\pi}}e^{-i\frac{(n+1)\omega}{2}}\left(\operatorname{sinc}\frac{\omega}{2}\right)^{n+1}$$

✎PROOF:

1. Proof using Theorem R.5 page 380:

$$\begin{aligned}
\tilde{\mathbf{F}}\mathsf{N}_n(\omega) &= \frac{1}{\sqrt{2\pi}}\int_{\mathbb{R}} \mathsf{N}_n(x)e^{-i\omega x}\,\mathrm{d}x && \text{by definition of } \tilde{\mathbf{F}} \quad \text{(Definition O.2 page 333)}\\
&= \frac{1}{\sqrt{2\pi}}\int_{[0:1)^{n+1}} e^{-i\omega(x_1+x_2+\cdots+x_{n+1})}\,\mathrm{d}x_1\,\mathrm{d}x_2\cdots,\,\mathrm{d}x_{n+1} && \text{by Theorem R.5}\\
&= \frac{1}{\sqrt{2\pi}}\prod_{k=1}^{n+1}\left(\int_{[0:1)} e^{-i\omega x_k}\,\mathrm{d}x_k\right) && \text{because } e^{x+y}=e^xe^y\\
&= \frac{1}{\sqrt{2\pi}}\left(\int_0^1 e^{-i\omega x}\,\mathrm{d}x\right)^{n+1} = \frac{1}{\sqrt{2\pi}}\left(\left.\frac{e^{-i\omega x}}{-i\omega}\right|_0^1\right)^{n+1}\\
&= \frac{1}{\sqrt{2\pi}}\left(\frac{1-e^{-i\omega}}{i\omega}\right)^{n+1} = \frac{1}{\sqrt{2\pi}}\left[e^{-i\frac{\omega}{2}}\left(\frac{e^{i\frac{\omega}{2}}-e^{-i\frac{\omega}{2}}}{i\omega}\right)\right]^{n+1}\\
&= \frac{1}{\sqrt{2\pi}}\left[e^{-i\frac{\omega}{2}}\left(\frac{2i\sin\left(\frac{\omega}{2}\right)}{\frac{2i\omega}{2}}\right)\right]^{n+1} && \text{by } \textit{Euler formulas} \quad \text{(Corollary ?? page ??)}\\
&= \frac{1}{\sqrt{2\pi}}e^{-i\frac{(n+1)\omega}{2}}\left(\frac{\sin(\omega/2)}{\omega/2}\right)^{n+1}
\end{aligned}$$

2. Proof using *rectangular pulse* example (Example O.1 page 340) and *Convolution Theorem* (Theorem Q.2 page 356):

$$\begin{aligned}
\tilde{\mathbf{F}}\mathsf{N}_n(\omega) &= \left[\sqrt{2\pi}\right]^n\left[\tilde{\mathbf{F}}\mathsf{N}_0\right]^{n+1} && \text{by } \textit{Convolution Theorem} \quad \text{(Theorem Q.2 page 356)}\\
&= \frac{1}{\sqrt{2\pi}}\left[\sqrt{2\pi}\frac{c(b-a)}{\sqrt{2\pi}}e^{-i\left(\frac{a+b}{2}\omega\right)}\left(\frac{\sin\left(\frac{b-a}{2}\omega\right)}{\left(\frac{b-a}{2}\omega\right)}\right)\right]^{n+1} && \text{by } \textit{rectangular pulse} \text{ example with } a=0,\ b=c=1 \quad \text{(Example O.1 page 340)}\\
&= \frac{1}{\sqrt{2\pi}}\left[\sqrt{2\pi}\frac{1}{\sqrt{2\pi}}e^{-i\left(\frac{\omega}{2}\right)}\left(\frac{\sin\left(\frac{\omega}{2}\right)}{(\omega/2)}\right)\right]^{n+1}\\
&= \frac{1}{\sqrt{2\pi}}e^{-i\left(\frac{(n+1)\omega}{2}\right)}\left(\frac{\sin\left(\frac{\omega}{2}\right)}{(\omega/2)}\right)^{n+1}
\end{aligned}$$

✏

Example R.10. The *Fourier transform magnitude* $\left|\left[\tilde{\mathbf{F}}\mathsf{N}_0\right](\omega)\right|$ of the *0 order B-spline* $\mathsf{N}_0(x)$ is illustrated to the right.



[17] Christensen (2008) page 142 ⟨Corollary 6.1.2⟩

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

Example R.11. The *Fourier transform magnitude* $\left|\left[\tilde{\mathbf{F}}\mathsf{N}_1\right](\omega)\right|$ of the *1st order B-spline* $\mathsf{N}_1(x)$ is illustrated to the right.



Example R.12. The *Fourier transform magnitude* $\left|\left[\tilde{\mathbf{F}}\mathsf{N}_2\right](\omega)\right|$ of the *2nd order B-spline* $\mathsf{N}_2(x)$ is illustrated to the right.



R.5 Basis properties

R.5.1 Uniqueness properties

Coefficients of a *basis sequence* are not always *unique*. Take for example a very trivial sequence (α_1, α_2) in which the coefficients are summed. If $\mathsf{f}(x) \triangleq \alpha_1 + \alpha_2$ and $\mathsf{g}(x) \triangleq \beta_1 + \beta_2$,

$$\begin{aligned} &\text{then} \quad \{(\alpha_1,\alpha_2) = (\beta_1,\beta_2)\} \quad &&\Longrightarrow \quad \mathsf{f}(x) = \mathsf{g}(x) \\ &\text{but} \quad \mathsf{f}(x) = \mathsf{g}(x) &&\not\Longrightarrow \quad \{(\alpha_1,\alpha_2) = (\beta_1,\beta_2)\}, \end{aligned}$$

because for example if $(\alpha_1, \alpha_2) = (1, 2)$ and $(\beta_1, \beta_2) = (-6, 9)$, then $\mathsf{f}(x) = \mathsf{g}(x)$, but $(\alpha_1, \alpha_2) \neq (\beta_1, \beta_2)$. This example demonstrates that the "if and only if" condition $\Longleftrightarrow$ does not hold and coefficients are not unique in all *basis sequences*. But arguably a minimal requirement for any practical basis sequence is that the coefficients are *unique* (the "if and only if" condition $\Longleftrightarrow$ holds). And indeed, in a *B-spline* basis sequence $(\!(\mathsf{N}_n(x-k))\!)_{k\in\mathbb{Z}}$, the coefficients $(\!(\alpha_k)\!)_{k\in\mathbb{Z}}$ are *unique*, as demonstrated by Theorem R.6 (next).

Theorem R.6. [18] *Let* $\mathsf{N}_n(x)$ *be the* nTH-ORDER B-SPLINE *(Definition R.2 page 367). Let*

$$\mathsf{f}(x) \triangleq \sum_{k\in\mathbb{Z}} \alpha_k \mathsf{N}_n(x-k) \quad \textit{and} \quad \mathsf{g}(x) \triangleq \sum_{k\in\mathbb{Z}} \beta_k \mathsf{N}_n(x-k).$$

THM

$$\underbrace{\left\{\ \mathsf{f}(x) = \mathsf{g}(x) \quad \forall x \in \mathbb{R}\ \right\} \quad \Longleftrightarrow \quad \left\{(\!(\alpha_k)\!)_{k\in\mathbb{Z}} = (\!(\beta_k)\!)_{k\in\mathbb{Z}}\right\}}_{\textit{coefficients are UNIQUE}}$$

✎PROOF:

1. Proof that $\Longleftarrow$ condition holds:

$$\begin{aligned} \mathsf{f}(x) &\triangleq \sum_{k\in\mathbb{Z}} \alpha_k \mathsf{N}_n(x-k) && \text{by definition of } \mathsf{f}(x) \\ &= \sum_{k\in\mathbb{Z}} \beta_k \mathsf{N}_n(x-k) && \text{by right hypothesis} \\ &\triangleq \mathsf{g}(x) && \text{by definition of } \mathsf{g}(x) \end{aligned}$$

2. Proof that $\Longrightarrow$ condition holds (proof by contradiction):

   (a) Suppose it does *not* hold.

[18] Wojtaszczyk (1997) page 55 ⟨Theorem 3.11⟩

COPYRIGHT © 2019 DANIEL J. GREENHOE | https://github.com/dgreenhoe/pdfs/blob/master/abcstat.pdf | CC BY-NC-ND

(b) Then there exists sequences $(\!(\alpha_k)\!)_{k\in\mathbb{Z}}$ and $(\!(\beta_k)\!)_{k\in\mathbb{Z}}$ such that
$$(\!(\alpha_k)\!) - (\!(\beta_k)\!) \triangleq (\!(alpha_k - \beta_k)\!) \neq (\!(0,0,0,\cdots)\!)$$
but also such that $\mathsf{f}(x) - \mathsf{g}(x) = 0\ \forall x \in \mathbb{R}$.

(c) If this were possible, then

$$\begin{aligned}
0 &= \mathsf{f}(x) - \mathsf{g}(x) \\
&= \sum_{m\in\mathbb{Z}} \alpha_m \mathsf{N}_n(x-m) - \sum_{m\in\mathbb{Z}} \beta_m \mathsf{N}_n(x-m) \\
&= \sum_{m\in\mathbb{Z}} \big(\alpha_m - \beta_m\big)\mathsf{N}_n(x-m) \\
&= \sum_{m=0}^{m=n} \big(\alpha_m - \beta_m\big)\frac{1}{n!}\left[\sum_{k=0}^{n+1}(-1)^k\binom{n+1}{k}(x-k)^n\sigma(x-k)\right] && \text{by Theorem R.1 page 371}
\end{aligned}$$

(d) But this is *impossible* because $\mathsf{N}(x)$ is *non-negative* (Theorem R.4 page 378).

(e) Therefore, there is a contradiction, and the $\Longrightarrow$ condition *does* hold.

✏

R.5.2 Partition of unity properties

In the case in which a sequence of *B-splines* $(\!(\mathsf{N}_n(x-k))\!)_{k\in\mathbb{Z}}$ is to be used as a *basis* for some subspace of $\boldsymbol{L}^2_{\mathbb{R}}$, arguably one of the most important properties for the sequence to have is the *partition of unity* property such that $\sum_{k\in\mathbb{Z}} \mathsf{N}_n(x-k) = 1$. This allows for convenient representation of the most basic functions, such as constants.[19] As it turns out, B-splines *do* have this property (next theorem).

Theorem R.7 (*B-spline partition of unity*). [20] *Let* $\mathsf{N}_n(x)$ *be the* nTH ORDER B-SPLINE *(Definition R.2 page 367)*.

THM
$$\sum_{k\in\mathbb{Z}} \mathsf{N}_n(x-k) = 1 \quad \forall n \in \mathbb{W} \qquad \text{(PARTITION OF UNITY)}$$

✎PROOF:

1. lemma: $\sum_{k\in\mathbb{Z}} \mathsf{N}_0(x-k) = 1$. Proof:

$$\begin{aligned}
\sum_{k\in\mathbb{Z}} \mathsf{N}_0(x-k) &= \sum_{k\in\mathbb{Z}} \mathbb{1}_{[0:1)}(x-k) && \text{by definition of } \mathsf{N}_0(x) && \text{(Definition R.2 page 367)} \\
&= 1 && \text{by definition of } \mathbb{1}_A(x) && \text{(Definition ?? page ??)}
\end{aligned}$$

2. Proof for this theorem follows from the $n = 0$ case ((1) lemma page 384), the definition of $\mathsf{N}_n(x)$ (Definition R.2 page 367), and Corollary ?? (page ??).

3. Alternatively, this theorem can be proved by *induction*:

   (a) Base case ($n = 0$ case): by (1) lemma.

[19] Jawerth and Sweldens (1994) page 8

[20] Christensen (2008) page 140 ⟨Theorem 6.1.1⟩, Höllig (2003) page 27 ⟨3.4⟩, Schumaker (2007) page 120 ⟨Theorem 4.15⟩, de Boor (2001) page 90 ⟨B-Spline Property (i)⟩, Chui (1988) page 2 ⟨Theorem 1.1⟩, Wojtaszczyk (1997) page 53 ⟨Theorem 3.7⟩, Cox (1972), de Boor (1972)

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE



Figure R.3: *auto-power spectrum* $\tilde{S}_n(\omega)$ plots of *B-splines* $\mathsf{N}_n(x)$ (Lemma R.6 page 385) For C and LaTeX source code to generate such a plot, see Section **??** (page **??**).

(b) Inductive step—proof that $\sum_{k\in\mathbb{Z}} \mathsf{N}_n(x-k) = 1 \implies \sum_{k\in\mathbb{Z}} \mathsf{N}_{n+1}(x-k) = 1$:

$$\begin{aligned}
\sum_{k\in\mathbb{Z}} \mathsf{N}_{n+1}(x-k) &= \sum_{k\in\mathbb{Z}} \int_{\tau=0}^{\tau=1} \mathsf{N}_n(x-k-\tau)\,\mathrm{d}\tau && \text{by Lemma R.2 page 367} \\
&= \sum_{k\in\mathbb{Z}} \int_{x-u=0}^{x-u=1} \mathsf{N}_n(u-k)(-1)\,\mathrm{d}u && \text{where } u \triangleq x-\tau \implies \tau = x-u \\
&= \sum_{k\in\mathbb{Z}} \int_{u=x-1}^{u=x} \mathsf{N}_n(u-k)\,\mathrm{d}u \\
&= \int_{u=x-1}^{u=x} \left(\sum_{k\in\mathbb{Z}} \mathsf{N}_n(u-k)\right) \mathrm{d}u \\
&= \int_{u=x-1}^{u=x} 1\,\mathrm{d}u && \text{by induction hypothesis} \\
&= 1
\end{aligned}$$

✏

Example R.13. The *partition of unity* property for the 0 *order B-spline* $\mathsf{N}_0(x)$ (Example R.1 page 369) is illustrated to the right.



Example R.14. The *partition of unity* property for the 1*st order B-spline* $\mathsf{N}_1(x)$ (Example R.2 page 370) is illustrated to the right.



Example R.15. The *partition of unity* property for the 2*nd order B-spline* $\mathsf{N}_2(x)$ (Example R.3 page 370) is illustrated to the right.



R.5.3 Riesz basis properties

Lemma R.6. *Let* $\mathsf{N}_n(x)$ *be the* nTH ORDER B-SPLINE *(Definition R.2 page 367).*
Let $\tilde{S}_n(\omega) \triangleq 2\pi \sum_{k\in\mathbb{Z}} \left|\tilde{\mathsf{F}}\mathsf{N}_n(\omega - 2\pi k)\right|^2$ *be the* AUTO-POWER SPECTRUM *(Definition **??** page **??**) of* $\mathsf{N}_n(x)$.

COPYRIGHT © 2019 DANIEL J. GREENHOE

LEM

(1).	$0 < \tilde{S}_n(\omega) \le 1 \quad \forall \omega \in \mathbb{R} \quad , \quad \forall n \in \mathbb{W}$	(3).	$\tilde{S}_n(0) = 1 \quad \forall n \in \mathbb{W}$
(2).	$\tilde{S}_n(\omega) = 1 \quad \forall \omega \in \mathbb{R} \quad , \quad \text{for } n = 0$	(4).	$\tilde{S}_n(\pi) \le \frac{1}{3} \quad \forall n \in \mathbb{W}\setminus\{0\}$

(*Note: see illustration in Figure R.3 page 385.*)

✎PROOF:

1. lemma: $\tilde{S}_n(\omega) = \sum_{k\in\mathbb{Z}} \left[\frac{\sin\left(\frac{\omega}{2}\right)}{k\pi - \frac{\omega}{2}}\right]^{2(n+1)}$. Proof:

$$\begin{aligned}
\tilde{S}_n(\omega) &\triangleq 2\pi \sum_{k\in\mathbb{Z}} \left|\tilde{\mathbf{F}}\mathsf{N}_n(\omega - 2\pi k)\right|^2 && \text{by Definition ?? page ??}\\
&= 2\pi \sum_{k\in\mathbb{Z}} \left|\frac{1}{\sqrt{2\pi}} e^{-i\frac{(n+1)(\omega-2\pi k)}{2}} \left(\frac{\sin\left(\frac{\omega-2\pi k}{2}\right)}{\frac{\omega-2\pi k}{2}}\right)^{n+1}\right|^2 && \text{by Lemma R.5 page 382}\\
&= \sum_{k\in\mathbb{Z}} \left[\frac{\sin\left(\frac{\omega-2\pi k}{2}\right)}{\frac{\omega-2\pi k}{2}}\right]^{2(n+1)}\\
&= \sum_{k\in\mathbb{Z}} \left[\frac{\sin\left(\frac{\omega}{2} - k\pi\right)}{k\pi - \frac{\omega}{2}}\right]^{2(n+1)}\\
&= \sum_{k\in\mathbb{Z}} \left[\frac{(-1)^k \sin\left(\frac{\omega}{2}\right)}{k\pi - \frac{\omega}{2}}\right]^{2(n+1)}\\
&= \sum_{k\in\mathbb{Z}} \left[\frac{\sin\left(\frac{\omega}{2}\right)}{k\pi - \frac{\omega}{2}}\right]^{2(n+1)}
\end{aligned}$$

2. lemma (one sided series form):

$$\begin{aligned}
\tilde{S}_n(\omega) &= \sum_{k\in\mathbb{Z}} \left[\frac{\sin\left(\frac{\omega}{2}\right)}{k\pi - \frac{\omega}{2}}\right]^{2(n+1)} && \text{by (1) lemma}\\
&= \left[\frac{\sin(\omega/2)}{(\omega/2)}\right]^{2(n+1)} + \left[\frac{\sin(\omega/2)}{(\pi/2)}\right]^{2(n+1)} \left(\sum_{k=1}^{\infty} \left[\frac{1}{2k - \frac{\omega}{\pi}}\right]^{2(n+1)} + \sum_{k=1}^{\infty} \left[\frac{1}{2k + \frac{\omega}{\pi}}\right]^{2(n+1)}\right)
\end{aligned}$$

3. lemma: $\tilde{S}_n(\omega)$ is *continuous* for all $\omega \in \mathbb{R}$.
   Proof: $\sin(\omega/2)$ and $\omega/2$ are *continuous*, so $\tilde{S}_n(\omega)$ is *continuous* as well.

4. lemma: $\tilde{S}_n(\omega)$ is *periodic* with period 2π (and so we only need to examine $\tilde{S}_n(\omega)$ for $\omega \in [0 : 2\pi)$). Proof of *periodicity*: This follows directly from Proposition **??** (page **??**).

5. lemma: $\tilde{S}_n(-\omega) = \tilde{S}_n(\omega)$ (*symmetric* about 0) and $\tilde{S}_n(\pi - \omega) = \tilde{S}_n(\pi + \omega)$ (*symmetric* about π). Proof: This follows directly from Proposition **??** (page **??**).

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

6. Proof that $\tilde{S}_n(0) = 1$:

$$\begin{aligned}
\tilde{S}_n(0) &= \lim_{\omega\to 0} \tilde{S}_n(\omega) && \text{by (3) lemma}\\
&= \lim_{\omega\to 0}\left[\left[\frac{\sin(\omega/2)}{(\omega/2)}\right]^{2(n+1)} + \left[\frac{\sin(\omega/2)}{(\pi/2)}\right]^{2(n+1)}\left(\sum_{k=1}^{\infty}\left[\frac{1}{2k-\frac{\omega}{\pi}}\right]^{2(n+1)} + \sum_{k=1}^{\infty}\left[\frac{1}{2k+\frac{\omega}{\pi}}\right]^{2(n+1)}\right)\right] && \text{by (2) lemma}\\
&= \lim_{\omega\to 0}\left[\frac{\cos\left(\frac{\omega}{2}\right)}{-\frac{1}{2}}\right]^{2(n+1)} + 0 && \text{by } \textit{l'Hôpital's rule}\\
&= (-1)^{2(n+1)} = 1
\end{aligned}$$

7. Proof that $\tilde{S}_n(\pi)$ converges to some value > 0:

(a) Proof that $\tilde{S}_n(\pi) > 0$:

$$\begin{aligned}
\tilde{S}_n(\pi) &= \left[\frac{\sin(\pi/2)}{(\pi/2)}\right]^{2(n+1)} + \left[\frac{\sin(\pi/2)}{(\pi/2)}\right]^{2(n+1)}\left(\sum_{k=1}^{\infty}\left[\frac{1}{2k-\frac{\pi}{\pi}}\right]^{2(n+1)} + \sum_{k=1}^{\infty}\left[\frac{1}{2k+\frac{\pi}{\pi}}\right]^{2(n+1)}\right) && \text{by (2) lemma}\\
&= \left(\frac{2}{\pi}\right)^{2(n+1)}\left[1 + \left(\frac{1}{1}\right)^{2(n+1)} + \left(\frac{1}{3}\right)^{2(n+1)} + \left(\frac{1}{3}\right)^{2(n+1)} + \left(\frac{1}{5}\right)^{2(n+1)} + \left(\frac{1}{5}\right)^{2(n+1)} + \cdots\right]\\
&= 2\left(\frac{2}{\pi}\right)^{2(n+1)} \underbrace{\sum_{k=1}^{\infty}\left[\frac{1}{2k-1}\right]^{2(n+1)}}_{\textit{Dirichlet Lambda function } \lambda(2n+2)}\\
&> 0 \qquad \text{because } x^2 > 0 \text{ for all } x \in \mathbb{R}\setminus\{0\}
\end{aligned}$$

(b) Proof that $\tilde{S}_n(\pi)$ *converges*:

$$\begin{aligned}
\tilde{S}_n(\pi) &= 2\left(\frac{2}{\pi}\right)^{2(n+1)}\sum_{k=1}^{\infty}\left(\frac{1}{2k-1}\right)^{2(n+1)} && \text{by item (7a)}\\
&\le 2\left(\frac{2}{\pi}\right)^{2(n+1)}\sum_{k=1}^{\infty}\left(\frac{1}{k}\right)^{2(n+1)}\\
&\le 2\left(\frac{2}{\pi}\right)^{2(n+1)}\sum_{k=1}^{\infty}\left(\frac{1}{k}\right)^{2}\\
&\implies \textit{convergence} && \text{by } \textit{comparison test}
\end{aligned}$$

(c) Tighter bounds for $\tilde{S}_n(\pi)$ for certain values of $n \in \{0,1,2,3,4\}$:

$$\begin{aligned}
\tilde{S}_n(\pi) &= 2\left(\frac{2}{\pi}\right)^{2(n+1)}\sum_{k=1}^{\infty}\left(\frac{1}{2k-1}\right)^{2(n+1)} && \text{by item (7a)}\\
&= 2\left(\frac{2}{\pi}\right)^{2(n+1)} U_{2(n+1)} && \text{by Jolley (1961), pages 56–57 ⟨(307)⟩}\\
&= 2\left(\frac{2}{\pi}\right)^{2(n+1)}\left[\frac{\pi^{2(n+1)}\alpha_{n+1}}{(4)[(2n+2)!]}\right] && \text{by Jolley (1961), pages 56–57 ⟨(307)⟩}\\
&= \frac{2^{2n+1}\alpha_{n+1}}{(2n+2)!}
\end{aligned}$$

Copyright © 2019 Daniel J. Greenhoe

$$= \left\{ \begin{array}{lll} \frac{2^1(1)}{2!} & \text{for } n=0 & (\alpha_1 = 1) \\ \frac{2^5(1)}{4!} & \text{for } n=1 & (\alpha_2 = 1) \\ \frac{2^5(3)}{6!} & \text{for } n=2 & (\alpha_3 = 3) \\ \frac{2^7(17)}{8!} & \text{for } n=3 & (\alpha_4 = 17) \\ \frac{2^9(155)}{10!} & \text{for } n=4 & (\alpha_5 = 155) \end{array} \right\} \qquad \text{by Jolley (1961), page 234 } \langle(1130)\rangle$$

$$= \left\{ \begin{array}{ll} 1 & \text{for } n=0 \\ \frac{1}{3} & \text{for } n=1 \\ \frac{2}{15} & \text{for } n=2 \\ \frac{17}{315} & \text{for } n=3 \\ \frac{62}{2835} & \text{for } n=4 \end{array} \right\} = \left\{ \begin{array}{ll} 1 & \text{for } n=0 \\ 0.3333333333333\cdots & \text{for } n=1 \\ 0.1333333333333\cdots & \text{for } n=2 \\ 0.0539682539682\cdots & \text{for } n=3 \\ 0.0218694885361\cdots & \text{for } n=4 \end{array} \right\}$$

(d) Being important for the $n = 0$ case, note that[21]

$$\underbrace{\sum_{k=1}^{\infty} \left(\frac{1}{2k-1}\right)^2}_{\textit{Dirichlet Lambda function } \lambda(2)} = 1 + \frac{1}{3^2} + \frac{1}{5^2} + \frac{1}{7^2} + \cdots = \frac{\pi^2}{8}$$

(e) Proof that $\tilde{S}_n(\pi) \le \frac{1}{3}$: because $\tilde{S}_n(\pi) = \frac{1}{3}$ for $n = 1$ (item (7c) page 387) and because $\tilde{S}_n(\pi)$ is decreasing for increasing n.

8. lemma: $\tilde{S}_n(\omega)$ *converges* to some value $> 0\ \forall \omega \in \mathbb{R}$. Proof:

(a) For $\omega = 0$, $\tilde{S}_n(\omega) = 1$ by item (6).

(b) Proof that $\tilde{S}_n(\omega) > 0$ for $\omega \in (0 : 2\pi)$:

$$\tilde{S}_n(\omega) = \left[\frac{\sin(\omega/2)}{(\omega/2)}\right]^{2(n+1)} + \left[\frac{\sin(\omega/2)}{(\pi/2)}\right]^{2(n+1)} \left(\sum_{k=1}^{\infty} \left[\frac{1}{2k - \frac{\omega}{\pi}}\right]^{2(n+1)} + \sum_{k=1}^{\infty} \left[\frac{1}{2k + \frac{\omega}{\pi}}\right]^{2(n+1)} \right) \qquad \text{by (2) lemma}$$

$$> 0$$

(c) Proof that $\tilde{S}_n(\omega)$ *converges*:

i. lemma: $\displaystyle\sum_{k=1}^{\infty} \left[\frac{1}{2k \pm \frac{\omega}{\pi}}\right]^{2(n+1)}$ *converges*. Proof:

$$\lim_{b\to\infty} \int_1^b \left[\frac{1}{2y \pm \frac{\omega}{\pi}}\right]^{2(n+1)} \mathrm{d}y = \lim_{b\to\infty} \int_1^b \left[2y \pm \frac{\omega}{\pi}\right]^{-2n-2} \mathrm{d}y$$

$$= \lim_{b\to\infty} \left. \frac{\left[2y \pm \frac{\omega}{\pi}\right]^{-2n-1}}{2(-2n-1)} \right|_1^b$$

$$= \lim_{b\to\infty} \left(\frac{-1}{2(2n+1)}\right) \left[\frac{1}{\left[2b \pm \frac{\omega}{\pi}\right]^{2n+1}} - \frac{1}{\left[2 \pm \frac{\omega}{\pi}\right]^{2n+1}} \right]$$

$$= 0 + \frac{1}{2(2n+1)\left[2 \pm \frac{\omega}{\pi}\right]^{2n+1}}$$

$$< \infty \qquad \forall \omega \in [0 : 2\pi)$$

$$\Longrightarrow \sum_{k=1}^{\infty} \left[\frac{1}{2k \pm \frac{\omega}{\pi}}\right]^{2(n+1)} \quad \textit{converges} \qquad \text{by } \textit{integral test}$$

[21] Nahin (2011) page 153, Bailey et al. (2013) page 334 ⟨Catalan's Constant⟩, Bailey et al. (2011) ⟨(15)⟩, Wells (1987) page 36 ⟨Dictionary entry for π: pages 31–37⟩, Heinbockel (2010) page 94 ⟨(2.27) *Dirichlet Lambda function*⟩

BY-NC-ND Copyright © 2019 Daniel J. Greenhoe

ii. completion of proof using (8(c)i) lemma …

$$\begin{aligned}
&\tilde{S}_n(\omega)\\
&= \left[\frac{\sin(\omega/2)}{(\omega/2)}\right]^{2(n+1)} + \left[\frac{\sin(\omega/2)}{(\pi/2)}\right]^{2(n+1)}\left(\sum_{k=1}^{\infty}\left[\frac{1}{2k-\frac{\omega}{\pi}}\right]^{2(n+1)} + \sum_{k=1}^{\infty}\left[\frac{1}{2k+\frac{\omega}{\pi}}\right]^{2(n+1)}\right) && \text{by (2) lemma}\\
&\Longrightarrow \tilde{S}_n(\omega) \textit{ converges } \forall\omega \in (0:2\pi) && \text{by (8(c)i) lemma}
\end{aligned}$$

9. lemma (an expression for $\tilde{S}_n'(\omega)$):

$$\begin{aligned}
\tilde{S}_n'(\omega) &\triangleq \frac{\mathrm{d}}{\mathrm{d}\omega}\tilde{S}_n(\omega)\\
&= \frac{\mathrm{d}}{\mathrm{d}\omega}\sum_{k\in\mathbb{Z}}\left[\frac{\sin\left(\frac{\omega}{2}\right)}{k\pi-\frac{\omega}{2}}\right]^{2(n+1)} && \text{by (1) lemma page 386}\\
&= \sum_{k\in\mathbb{Z}}\frac{\mathrm{d}}{\mathrm{d}\omega}\left[\frac{\sin\left(\frac{\omega}{2}\right)}{k\pi-\frac{\omega}{2}}\right]^{2(n+1)} && \text{by } \textit{linearity} \text{ of } \frac{\mathrm{d}}{\mathrm{d}\omega} \text{ operator}\\
&= \sum_{k\in\mathbb{Z}}2(n+1)\left[\frac{\sin\left(\frac{\omega}{2}\right)}{k\pi-\frac{\omega}{2}}\right]^{2n+1}\frac{\mathrm{d}}{\mathrm{d}\omega}\left[\frac{\sin\left(\frac{\omega}{2}\right)}{k\pi-\frac{\omega}{2}}\right] && \text{by } \textit{power rule}\\
&= 2(n+1)\sum_{k\in\mathbb{Z}}\left[\frac{\sin\left(\frac{\omega}{2}\right)}{k\pi-\frac{\omega}{2}}\right]^{2n+1}\left[\frac{\frac{1}{2}\cos\left(\frac{\omega}{2}\right)\left(k\pi-\frac{\omega}{2}\right)-\sin\left(\frac{\omega}{2}\right)\left(-\frac{1}{2}\right)}{\left(k\pi-\frac{\omega}{2}\right)^2}\right] && \text{by } \textit{quotient rule}\\
&= (n+1)\sum_{k\in\mathbb{Z}}\left[\frac{\sin\left(\frac{\omega}{2}\right)}{k\pi-\frac{\omega}{2}}\right]^{2n+1}\left[\frac{\cos\left(\frac{\omega}{2}\right)\left(k\pi-\frac{\omega}{2}\right)+\sin\left(\frac{\omega}{2}\right)}{\left(k\pi-\frac{\omega}{2}\right)^2}\right]
\end{aligned}$$

10. lemma: $\tilde{S}_n'(0) = \tilde{S}_n'(\pi) = 0$. Proof: This follows from Proposition **??** (page **??**). Here is alternate proof:

$$\begin{aligned}
\tilde{S}_n'(0) &= \lim_{\omega\to 0}\tilde{S}_n'(\omega)\\
&= \lim_{\omega\to 0}(n+1)\sum_{k\in\mathbb{Z}}\left[\frac{\sin\left(\frac{\omega}{2}\right)}{k\pi-\frac{\omega}{2}}\right]^{2n+1}\left[\frac{\cos\left(\frac{\omega}{2}\right)\left(k\pi-\frac{\omega}{2}\right)+\sin\left(\frac{\omega}{2}\right)}{\left(k\pi-\frac{\omega}{2}\right)^2}\right] && \text{by (9) lemma}\\
&= \lim_{\omega\to 0}(n+1)\left[\frac{\sin\left(\frac{\omega}{2}\right)}{-\frac{\omega}{2}}\right]^{2n+1}\left[\frac{\cos\left(\frac{\omega}{2}\right)\left(-\frac{\omega}{2}\right)+\sin\left(\frac{\omega}{2}\right)}{\left(-\frac{\omega}{2}\right)^2}\right]\\
&= (n+1)\lim_{\omega\to 0}\left[\frac{\sin\left(\frac{\omega}{2}\right)}{-\frac{\omega}{2}}\right]^{2n+1}\lim_{\omega\to 0}\left[\frac{\cos\left(\frac{\omega}{2}\right)\left(-\frac{\omega}{2}\right)+\sin\left(\frac{\omega}{2}\right)}{\left(-\frac{\omega}{2}\right)^2}\right]\\
&= (n+1)[-1]^{2n+1}\lim_{\omega\to 0}\left[\frac{-\frac{1}{2}\sin\left(\frac{\omega}{2}\right)\left(-\frac{\omega}{2}\right)+\cos\left(\frac{\omega}{2}\right)\left(-\frac{1}{2}\right)+\cos\left(\frac{\omega}{2}\right)\left(\frac{1}{2}\right)}{-\frac{2}{2}\left(-\frac{\omega}{2}\right)}\right] && \text{by } \textit{l'Hôpital's rule}\\
&= (1)(0)\\
&= 0
\end{aligned}$$

Copyright © 2019 Daniel J. Greenhoe

$$\begin{aligned}
\tilde{S}_n'(\pi) &= (n+1)\sum_{k\in\mathbb{Z}}\left[\frac{\sin\left(\frac{\pi}{2}\right)}{k\pi-\frac{\pi}{2}}\right]^{2n+1}\left[\frac{\cos\left(\frac{\pi}{2}\right)\left(k\pi-\frac{\pi}{2}\right)+\sin\left(\frac{\pi}{2}\right)}{\left(k\pi-\frac{\pi}{2}\right)^2}\right] && \text{by (9) lemma}\\
&= (n+1)\sum_{k\in\mathbb{Z}}\left[\frac{1}{k\pi-\frac{\pi}{2}}\right]^{2n+1}\left[\frac{0\left(k\pi-\frac{\pi}{2}\right)+1}{\left(k\pi-\frac{\pi}{2}\right)^2}\right]\\
&= (n+1)\left(\frac{2}{\pi}\right)^{2n+3}\sum_{k\in\mathbb{Z}}\left[\frac{1}{2k-1}\right]^{2n+3}\\
&= (n+1)\left(\frac{2}{\pi}\right)^{2n+3}\left[\left(\frac{1}{1}\right)^{2n+3}+\left(\frac{1}{-1}\right)^{2n+3}+\left(\frac{1}{3}\right)^{2n+3}+\left(\frac{1}{-3}\right)^{2n+3}+\ldots\right]\\
&= (n+1)\left(\frac{2}{\pi}\right)^{2n+3}\sum_{k=1}^{\infty}(-1)^{k+1}\alpha_k \qquad \text{where } \alpha_k \triangleq \left\{\begin{array}{ll}\left(\frac{1}{k}\right)^{2n+3} & \text{for } k \text{ odd}\\ \left(\frac{1}{k-1}\right)^{2n+3} & \text{for } k \text{ even}\end{array}\right\}\\
&= 0 \qquad \text{because } \lim_{k\to\infty}\alpha_k = 0 \text{ and by } \textit{Alternating Series Test}
\end{aligned}$$

11. lemma: $\tilde{S}_n(\omega)$ is *decreasing* with respect to $\omega \in [0 : \pi]$. Proof:

$$\begin{aligned}
\tilde{S}_n'(\omega) &= (n+1)\sum_{k\in\mathbb{Z}}\left[\frac{\sin\left(\frac{\omega}{2}\right)}{k\pi-\frac{\omega}{2}}\right]^{2n+1}\left[\frac{\cos\left(\frac{\omega}{2}\right)\left(k\pi-\frac{\omega}{2}\right)+\sin\left(\frac{\omega}{2}\right)}{\left(k\pi-\frac{\omega}{2}\right)^2}\right] && \text{by (9) lemma page 389}\\
&= \underbrace{(n+1)\left(\sin\frac{\omega}{2}\right)^{2n+1}}_{\geq 0 \text{ for } \omega\in[0:2\pi)}\underbrace{\sum_{k\in\mathbb{Z}}\left[\frac{1}{k\pi-\frac{\omega}{2}}\right]^{2n+2}}_{>0}\left[\underbrace{\left(\cos\frac{\omega}{2}\right)}_{\text{sign change at } \omega=\pi}+\frac{\overbrace{\sin\frac{\omega}{2}}^{>0 \text{ for } \omega\in(0:2\pi)}}{\underbrace{k\pi-\frac{\omega}{2}}_{\textit{decreasing} \text{ w.r.t. } \omega\in\mathbb{R}}}\right]
\end{aligned}$$

12. lemma: $\tilde{S}_n(\omega)$ is *increasing* with respect to $\omega \in [\pi : 2\pi)$. Proof: This is true because $\tilde{S}_n(\omega)$ is *decreasing* in $[0 : \pi]$ ((11) lemma) and because $\tilde{S}_n(\omega)$ is *symmetric* about $\omega = \pi$ ((5) lemma).

13. Proof that $0 < \tilde{S}_n(\omega) \leq 1$:

    (a) $\tilde{S}_n(\omega) > 0$ by (8) lemma and

    (b) $\tilde{S}_n(0) = 1$ by item (6) and

    (c) $\tilde{S}_n(\omega)$ is *decreasing* from $\omega = 0$ to $\omega = \pi$ by (11) lemma and

    (d) $\tilde{S}_n(\omega)$ is *increasing* from $\omega = \pi$ to $\omega = 2\pi$ by (12) lemma and

    (e) $\tilde{S}_n(2\pi) = 1$ because $\tilde{S}_n(2\pi) = \tilde{S}_n(0)$ by (4) lemma.

✏

Theorem R.8. [22]

THM					
	1. $(\!(\mathsf{N}_n(x-k))\!)_{k\in\mathbb{Z}}$ *is a*	RIESZ BASIS	*for* $\operatorname{span}(\!(\mathsf{N}_n(x-k))\!)_{k\in\mathbb{Z}}$		$\forall n \in \mathbb{W}$
	2. $(\!(\mathsf{N}_n(x-k))\!)_{k\in\mathbb{Z}}$ *is an*	ORTHONORMAL BASIS	*for* $\operatorname{span}(\!(\mathsf{N}_n(x-k))\!)_{k\in\mathbb{Z}}$	$\Longleftrightarrow$	$n = 0$

✎PROOF:

[22] Wojtaszczyk (1997) page 56 ⟨Proposition 3.12⟩, Prasad and Iyengar (1997) page 148 ⟨Theorem 6.3⟩, Forster and Massopust (2009) page 66 ⟨Theorem 2.25⟩

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

1. Proof that $(\!(\mathsf{N}_n(x-k))\!)_{k\in\mathbb{Z}}$ is a *Riesz basis* for $\mathsf{span}(\!(\mathsf{N}_n(x-k))\!)_{k\in\mathbb{Z}}$:

$$\begin{aligned}
&0 < \tilde{S}_n(\omega) \le 1 && \text{by Lemma R.6 page 385 (1)}\\
&\implies (\!(\mathsf{N}_n(x-k))\!)_{k\in\mathbb{Z}} \text{ is a } \textit{Riesz basis} \text{ for } \mathsf{span}(\!(\mathsf{N}_n(x-k))\!)_{k\in\mathbb{Z}} && \text{by Theorem ?? page ??}
\end{aligned}$$

2. Proof that $\{n=0\} \iff (\!(\mathsf{N}_n(x-k))\!)_{k\in\mathbb{Z}}$ is an *orthonormal basis* for $\mathsf{span}(\!(\mathsf{N}_n(x-k))\!)_{k\in\mathbb{Z}}$:

$$\begin{aligned}
n=0 &\iff \tilde{S}_n(\omega) = 1 && \text{by Lemma R.6 page 385 (2), (4)}\\
&\iff (\!(\mathsf{N}_n(x-k))\!)_{k\in\mathbb{Z}} \text{ is an } \textit{orthonormal basis} \text{ for } \mathsf{span}(\!(\mathsf{N}_n(x-k))\!) && \text{by Theorem ?? page ??}
\end{aligned}$$

✏

R.6 Mutiresolution properties

R.6.1 Introduction

In 1989, Stéphane G. Mallat introduced the *Multiresolution Analysis* (MRA) structure (Definition ?? page ??) An MRA is very powerful because it can be used to approximate functions at incrementally increasing "scales" of resolution, and furthermore induces a *wavelet*. In fact, the MRA has become the dominate wavelet construction method. Moreover, P.G. Lemarié has proved that all wavelets with *compact support* are generated by an MRA. [23]

R.6.2 B-spline dyadic decomposition

One key feature of an MRA is *dyadic decomposition* such that $\mathsf{N}_n(x) = \sum_k \alpha_n \mathsf{N}_n(2x-n)$ for some sequence $(\!(\alpha_n)\!)$. As it turns out, *B-splines* also have this property (next theorem).

Theorem R.9 (*B-spline dyadic decomposition*). [24] *Let* $\mathsf{N}_n(x)$ *be the* nTH ORDER B-SPLINE.

THM
$$\mathsf{N}_n(x) = \frac{1}{2^n}\sum_{k=0}^{n+1}\binom{n+1}{k}\mathsf{N}_n(2x-k) \qquad \forall n\in\mathbb{W},\ \forall x\in\mathbb{R}$$

✎PROOF:

1. Base case ...proof for $n=0$ case:

$$\begin{aligned}
\mathsf{N}_0(x) &= \mathbb{1}_{[0:1)}(x) && \text{by definition of } \mathbb{1}_A(x) \quad \text{(Definition ?? page ??)}\\
&= \mathbb{1}_{[0:½)}(x) + \mathbb{1}_{[½:1)}(x)\\
&= \mathbb{1}_{[2\times 0:2\times ½)}(2x) + \mathbb{1}_{[2\times ½-1:2\times 1-1)}(2x-1)\\
&= \mathbb{1}_{[0:1)}(2x) + \mathbb{1}_{[0:1)}(2x-1)\\
&= \frac{1}{2^0}\sum_{k=0}^{0+1}\binom{0+1}{k}\mathsf{N}_n(2x-k)
\end{aligned}$$

[23] Mallat (1999) page 240, Definition ?? (page ??)
[24] Prasad and Iyengar (1997) pages 151–152 ⟨proof using Fourier transform⟩

COPYRIGHT © 2019 DANIEL J. GREENHOE

2. Induction step…proof that n case $\implies n+1$ case:

$$
\begin{aligned}
\mathsf{N}_{n+1}(x) &= \int_0^1 \mathsf{N}_n(x-\tau)\,\mathrm{d}\tau && \text{by Lemma R.2 page 367}\\
&= \int_0^1 \frac{1}{2^n}\sum_{k=0}^{n+1}\binom{n+1}{k}\mathsf{N}_n(2x-2\tau-k)\,\mathrm{d}\tau && \text{by induction hypothesis}\\
&= \frac{1}{2^n}\sum_{k=0}^{n+1}\binom{n+1}{k}\int_{\tau=0}^{\tau=1}\mathsf{N}_n(2x-2\tau-k)\,\mathrm{d}\tau && \text{by \textit{linearity} of } \textstyle\sum \text{ operator}\\
&= \frac{1}{2^n}\sum_{k=0}^{n+1}\binom{n+1}{k}\int_{u=0}^{u=2}\mathsf{N}_n(2x-u-k)\frac{1}{2}\,\mathrm{d}u && \text{where } u \triangleq 2\tau \implies \tau = \tfrac{1}{2}u\\
&= \frac{1}{2^{n+1}}\sum_{k=0}^{n+1}\binom{n+1}{k}\left[\int_{u=0}^{u=1}\mathsf{N}_n(2x-k-u)\,\mathrm{d}u + \int_{u=1}^{u=2}\mathsf{N}_n(2x-k-u)\,\mathrm{d}u\right]\\
&= \frac{1}{2^{n+1}}\sum_{k=0}^{n+1}\binom{n+1}{k}\left[\int_{u=0}^{u=1}\mathsf{N}_n(2x-k-u)\,\mathrm{d}u + \int_{v=0}^{v=1}\mathsf{N}_n(2x-k-v-1)\,\mathrm{d}v\right] && \begin{array}{l}\text{where } v \triangleq u-1\\ \implies u = v+1\end{array}\\
&= \frac{1}{2^{n+1}}\sum_{k=0}^{n+1}\binom{n+1}{k}\left[\mathsf{N}_n(2x-k)+\mathsf{N}_n(2x-k-1)\right] && \text{by Lemma R.2 page 367}\\
&= \frac{1}{2^{n+1}}\left[\sum_{k=0}^{n+1}\binom{n+1}{k}\mathsf{N}_n(2x-k) + \sum_{m=1}^{n+2}\binom{n+1}{m-1}\mathsf{N}_n(2x-m)\right] && \text{where } m \triangleq k+1 \implies k = m-1\\
&= \frac{1}{2^{n+1}}\left[\underbrace{\sum_{k=1}^{n+1}\left[\binom{n+1}{k}+\binom{n+1}{k-1}\right]\mathsf{N}_n(2x-k)}_{\text{common indices of above two summations}} + \underbrace{\binom{n+1}{0}\mathsf{N}_n(2x-0)}_{k=0\text{ term}} + \underbrace{\binom{n+2}{n+2}\mathsf{N}_n(2x-n-2)}_{m=n+2\text{ term}}\right]\\
&= \frac{1}{2^{n+1}}\left[\underbrace{\sum_{k=1}^{n+1}\binom{n+2}{k}\mathsf{N}_n(2x-k)}_{\text{by \textit{Stifel formual} (Theorem ?? page ??)}} + \underbrace{\binom{n+2}{0}\mathsf{N}_n(2x-0)}_{\text{because } \binom{n+1}{0}=1=\binom{n+2}{0}} + \binom{n+2}{n+2}\mathsf{N}_n(2x-n-2)\right]\\
&= \frac{1}{2^{n+1}}\sum_{k=0}^{n+2}\binom{n+2}{k}\mathsf{N}_n(2x-k)
\end{aligned}
$$

✍

Example R.16. [25]The 0 *order B-spline dyadic decomposition*

$$\mathsf{N}_0(x) = \frac{1}{1}\sum_{k=0}^{k=1}\binom{1}{k}\mathsf{N}_0(2x-k)$$

is illustrated to the right.



[25] Strang (1989) page 615 ⟨Box function⟩, Strang and Nguyen (1996) page 441 ⟨Box function⟩

BY-NC-ND Copyright © 2019 Daniel J. Greenhoe

Example R.17. [26]The 1*st order B-spline dyadic decomposition*

$$\mathsf{N}_1(x) = \frac{1}{2}\sum_{k=0}^{k=2}\binom{2}{k}\mathsf{N}_1(2x-k)$$

is illustrated to the right.



Example R.18. The 2*nd order B-spline dyadic decomposition*

$$\mathsf{N}_2(x) = \frac{1}{4}\sum_{k=0}^{k=3}\binom{3}{k}\mathsf{N}_2(2x-k)$$

is illustrated to the right.



R.6.3 B-spline MRA scaling functions

Theorem R.10. *Let f* $N_n(x)$ *be the* nTH ORDER B-SPLINE *(Definition R.2 page 367)*.
Let $V_j \triangleq \operatorname{span}\left(\left(\mathsf{N}_n\left(2^j x - k\right)\right)\right)_{k\in\mathbb{Z}}$.

THM $\left(\left(V_j\right)\right)_{j\in\mathbb{Z}}$ *is a* MULTIRESOLUTION ANALYSIS *on* $L^2_{\mathbb{R}}$ *with* SCALING FUNCTION $\phi(x) \triangleq \mathsf{N}_n(x)$

✎PROOF:

1. lemma: $(\!(\mathsf{N}_n(x-k))\!)_{k\in\mathbb{Z}}$ is a *Riesz sequence*. Proof: by Theorem R.8 (page 390).
2. lemma: $\exists\,(\!(h_k)\!)$ such that $\mathsf{N}_n(x) = \sum_{k\in\mathbb{Z}} h_k \mathsf{N}_n(2x-k)$. Proof: by Theorem R.9 (page 391). In fact, note that $h_k = \frac{1}{2^n\sqrt{2}}\binom{n+1}{k}$
3. lemma: $\tilde{\mathbf{F}}\mathsf{N}_n(\omega)$ is *continuous* at 0. Proof:

$$\begin{aligned}\tilde{\mathbf{F}}\mathsf{N}_n(\omega) &= \frac{1}{\sqrt{2\pi}}e^{-i\frac{(n+1)\omega}{2}}\left(\operatorname{sinc}\frac{\omega}{2}\right)^{n+1} && \text{by Lemma R.5 page 382}\\ &\Longrightarrow \textit{continuous} \text{ at } 0 && \text{by known property of sinc function}\end{aligned}$$

4. lemma: $\tilde{\phi}(0) \neq 0$. Proof:

$$\begin{aligned}\tilde{\mathbf{F}}\mathsf{N}_n(0) &= \frac{1}{\sqrt{2\pi}}e^{-i\frac{(n+1)\omega}{2}}\left(\operatorname{sinc}\frac{\omega}{2}\right)^{n+1}\Bigg|_{\omega=0} && \text{by Lemma R.5 page 382}\\ &= 1\cdot\frac{1}{1/2} = 2 && \text{by } \textit{l'Hôpital's rule}\\ &\neq 0\end{aligned}$$

5. The completion of this proof follows directly from (1) lemma, (2) lemma, (3) lemma, (4) lemma, and Theorem **??** (page **??**).

✏

[26] Strang (1989) page 615 ⟨Hat function⟩, Strang and Nguyen (1996) page 442 ⟨Hat function⟩, Heil (2011) page 380 ⟨Fig. 12.10⟩

COPYRIGHT © 2019 DANIEL J. GREENHOE



Figure R.4: *dilation equation* demonstrations for selected B-splines (Example R.19 page 394)

R.6.4 B-spline MRA coefficient sequences

Because each *B-spline* $\mathsf{N}_n(x)$ is the *scaling function* for an *MRA* (Theorem R.10 page 393), each *B-spline* also satisfies the *dilation equation* (Theorem **??** page **??**) such that

$$\mathsf{N}_n(x) = \sqrt{2}\sum_{k\in\mathbb{Z}} h_k \mathsf{N}(2x-k) \quad \text{where} \quad h_k = \left\{\begin{array}{ll} \frac{\sqrt{2}}{2^{n+1}}\binom{n+1}{k} & \text{for } n = 0,1,\ldots,n \\ 0 & \text{otherwise} \end{array}\right\}$$

The resulting *sequence* $(\!(h_k)\!)_{k\in\mathbb{Z}}$ is the *order n B-spline MRA coefficient sequence* induced by the *order n B-spline MRA scaling sequence* $\phi(x) \triangleq \mathsf{N}_n(x)$.[27]

Example R.19. See Figure R.4 (page 394) for some *dilation equation* demonstrations of selected B-splines.

Theorem R.11 (B-spline scaling coefficients)**.** *Let* $\left(\mathbf{L}^2_{\mathbb{R}}, (\!(\mathbf{V}_j)\!), \phi, (\!(h_n)\!)\right)$ *be an* MRA SYSTEM *(Definition* **??** *page* **??***). Let* $\mathsf{N}_n(x)$ *be a* nTH ORDER B-SPLINE *(Definition R.2 page 367).*

THM

$$\underbrace{\phi(x) \triangleq \mathsf{N}_n(x)}_{\text{(1) B-spline scaling function}} \implies (\!(h_k)\!) = \left\{\begin{array}{ll} \frac{\sqrt{2}}{2^{n+1}}\binom{n}{k} & \text{for } k = 0,1,\ldots,n \\ 0 & \text{otherwise} \end{array}\right. \qquad \text{(2) scaling sequence in "time"}$$

$$\iff \check{\mathsf{h}}(z)\Big|_{z\triangleq e^{i\omega}} = \frac{\sqrt{2}}{2^n}\left(1+z^{-1}\right)^{n+1}\Big|_{z\triangleq e^{i\omega}} \qquad \text{(3) scaling sequence in "z domain"}$$

$$\iff \tilde{\mathsf{h}}(\omega) = 2\sqrt{2}e^{-i\frac{n+1}{2}\omega}\left[\cos\left(\frac{\omega}{2}\right)\right]^{n+1} \qquad \text{(4) scaling sequence in "frequency"}$$

✎PROOF:

1. Proof that (1) $\implies$ (3): By Theorem R.10 page 393 we know that $\mathsf{N}_n(x)$ is a *scaling function* (Definition **??** page **??**). So then we know that we can use Lemma **??** page **??**.

$$\begin{aligned}
\tilde{\mathsf{h}}(\omega) &= \sqrt{2}\frac{\tilde{\phi}(2\omega)}{\tilde{\phi}(\omega)} && \text{by Lemma ?? page ??} \\
&= \sqrt{2}\frac{\tilde{\mathsf{N}}_n(2\omega)}{\tilde{\mathsf{N}}_n(\omega)} && \text{by (1)} \\
&= \sqrt{2}\frac{\frac{1}{\sqrt{2\pi}}\left(\frac{1-e^{-i2\omega}}{2i\omega}\right)^{n+1}}{\frac{1}{\sqrt{2\pi}}\left(\frac{1-e^{-i\omega}}{i\omega}\right)^{n+1}} && \text{by Lemma R.5 page 382}
\end{aligned}$$

[27] For *Octave/MatLab* code useful for plotting a function given a sequence of coefficients $(\!(h_k)\!)$, see Section **??** (page **??**).

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

$$= \frac{\sqrt{2}}{2^{n+1}}\left(\frac{1-z^{-2}}{1-z^{-1}}\right)^{n+1}\Bigg|_{z=e^{i\omega}}$$
$$= \frac{\sqrt{2}}{2^{n+1}}\left[\left(\frac{1-z^{-2}}{1-z^{-1}}\right)\left(\frac{1+z^{-1}}{1+z^{-1}}\right)\right]^{n+1}\Bigg|_{z=e^{i\omega}}$$
$$= \frac{\sqrt{2}}{2^{n+1}}\left(\frac{\left(1-z^{-2}\right)\left(1+z^{-1}\right)}{1-z^{-2}}\right)^{n+1}\Bigg|_{z=e^{i\omega}}$$
$$= \frac{\sqrt{2}}{2^{n}}\left(1+z^{-1}\right)^{n+1}\Bigg|_{z\triangleq e^{i\omega}}$$

2. Proof that (3) $\Longleftrightarrow$ (2):

$$\check{\mathsf{h}}(z)\Big|_{z\triangleq e^{i\omega}} = \frac{\sqrt{2}}{2^{n}}\left(1+z^{-1}\right)^{n+1}\Bigg|_{z\triangleq e^{i\omega}} \qquad \text{by (3)}$$
$$= \frac{\sqrt{2}}{2^{n}}\left(\sum_{k=0}^{n+1}\binom{n}{k}z^{-k}\right)\Bigg|_{z\triangleq e^{i\omega}} \qquad \text{by } \textit{binomial theorem}$$
$$\Longleftrightarrow \qquad h_k = \frac{\sqrt{2}}{2^{n+1}}\binom{n}{k} \qquad \text{by definition of } Z \textit{ transform} \text{ (Definition Q.4 page 354)}$$

3. Proof that (3) $\Longrightarrow$ (4):

$$\tilde{\mathsf{h}}(\omega) = \check{\mathsf{h}}(z)\Big|_{z\triangleq e^{i\omega}} \qquad \text{by definition of } DTFT \text{ (Definition P.1 page 343)}$$
$$= \frac{\sqrt{2}}{2^{n}}\left(1+z^{-1}\right)^{n+1}\Bigg|_{z\triangleq e^{i\omega}} \qquad \text{by (3)}$$
$$= \frac{\sqrt{2}}{2^{n}}\left(1+e^{-i\omega}\right)^{n+1} \qquad \text{by definition of } z$$
$$= \frac{\sqrt{2}}{2^{n}}\left[e^{-i\frac{1}{2}\omega}\left(e^{i\frac{\omega}{2}}+e^{-i\frac{\omega}{2}}\right)\right]^{n+1}$$
$$= \frac{\sqrt{2}}{2^{n}}e^{-i\frac{n+1}{2}\omega}\left[2\cos\left(\frac{\omega}{2}\right)\right]^{n+1}$$
$$= 2\sqrt{2}e^{-i\frac{n+1}{2}\omega}\left[\cos\left(\frac{\omega}{2}\right)\right]^{n+1}$$

4. Proof that (3) $\Longleftarrow$ (4):

$$\check{\mathsf{h}}(z)\Big|_{z\triangleq e^{i\omega}} = \check{\mathsf{h}}\left(e^{i\omega}\right)$$
$$= \tilde{\mathsf{h}}(\omega)$$
$$= 2\sqrt{2}e^{-i\frac{n+1}{2}\omega}\left[\cos\left(\frac{\omega}{2}\right)\right]^{n+1} \qquad \text{by (4)}$$
$$= \frac{\sqrt{2}}{2^{n}}e^{-i\frac{n+1}{2}\omega}\left[2\cos\left(\frac{\omega}{2}\right)\right]^{n+1}$$
$$= \frac{\sqrt{2}}{2^{n}}\left[e^{-i\frac{1}{2}\omega}\left(e^{i\frac{\omega}{2}}+e^{-i\frac{\omega}{2}}\right)\right]^{n+1}$$
$$= \frac{\sqrt{2}}{2^{n}}\left(1+e^{-i\omega}\right)^{n+1}$$
$$= \frac{\sqrt{2}}{2^{n}}\left(1+z^{-1}\right)^{n+1}\Bigg|_{z\triangleq e^{i\omega}}$$

Copyright © 2019 Daniel J. Greenhoe

Example R.20 (2 coefficient case). [28] Let $\left(\boldsymbol{L}^2_{\mathbb{R}}, (\!(\boldsymbol{V}_j)\!), \phi, (\!(h_n)\!)\right)$ be an *MRA system* (Definition ?? page ??).

EX

$$\underbrace{\left\{\begin{array}{ll} 1. & \operatorname{supp}\phi(x) = [0:1] \quad \text{and} \\ 2. & (\!(\phi(x-k))\!) \text{ forms a} \\ & \textit{partition of unity} \end{array}\right\}}_{\text{(A)}} \iff \underbrace{h_n = \left\{\begin{array}{ll} \frac{\sqrt{2}}{2} & \text{for } n=0 \\ \frac{\sqrt{2}}{2} & \text{for } n=1 \\ 0 & \text{otherwise} \end{array}\right\}}_{\text{(B)}} \iff \underbrace{\left\{\phi(x) = \mathsf{N}_0(x)\right\}}_{\text{(C)}}$$

✎PROOF:

1. Proof that (A) $\Longrightarrow$ (B):

   (a) lemma: Only h_0 and h_1 are *non-zero*; All other coefficients h_k are 0. Proof: This follows from $\operatorname{supp}\phi(x) = [0:1]$ (Definition ?? page ??) and by Theorem ?? page ??.

   (b) lemma (equations for $(\!(h_k)\!)$): Because $(\!(h_k)\!)$ is a *scaling coefficient sequence* (Definition ?? page ??), it must satisfy the *admissibility equation* (Theorem ?? page ??). And because $(\!(\phi(x-k))\!)$ forms a *partition of unity*, it must satisfy the equations given by Theorem ?? (page ??). (1a) lemma and these two constraints yield two simultaneous equations and two unknowns:

   $$\begin{array}{rcll} h_0 + h_1 & = & \sqrt{2} & (\textit{admissibility condition}) \\ h_0 - h_1 & = & 0 & (\textit{partition of unity / zero at } -1 \textit{ / vanishing 0th moment}) \end{array}$$

   (c) lemma: The equations provided by (1b) lemma can be expressed in matrix algebra form as follows…

   $$\underbrace{\begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}}_{\mathbf{A}} \begin{bmatrix} h_0 \\ h_1 \end{bmatrix} = \begin{bmatrix} \sqrt{2} \\ 0 \end{bmatrix}$$

   (d) lemma: The *inverse* $\mathbf{A}^{-1}$ of $\mathbf{A}$ can be expressed as demonstrated below…

   $$\left[\begin{array}{cc|cc} 1 & 1 & 1 & 0 \\ 1 & -1 & 0 & 1 \end{array}\right] \to \left[\begin{array}{cc|cc} 2 & 0 & 1 & 1 \\ 1 & -1 & 0 & 1 \end{array}\right] \to \left[\begin{array}{cc|cc} 2 & 0 & 1 & 1 \\ 0 & -1 & -\frac{1}{2} & \frac{1}{2} \end{array}\right] \to \left[\begin{array}{cc|cc} 1 & 0 & \frac{1}{2} & \frac{1}{2} \\ 0 & 1 & \frac{1}{2} & -\frac{1}{2} \end{array}\right]$$
   $$\Longrightarrow \mathbf{A}^{-1} = \frac{1}{2}\begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$$

   (e) Proof for the values of $(\!(h_k)\!)$ (B):

   $$\begin{aligned} \begin{bmatrix} h_0 \\ h_1 \end{bmatrix} = \mathbf{A}^{-1}\mathbf{A}\begin{bmatrix} h_0 \\ h_1 \end{bmatrix} &= \mathbf{A}^{-1}\begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}\begin{bmatrix} h_0 \\ h_1 \end{bmatrix} && \text{by (1c) lemma} \\ &= \mathbf{A}^{-1}\begin{bmatrix} \sqrt{2} \\ 0 \end{bmatrix} && \text{by (1c) lemma} \\ &= \frac{1}{2}\begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}\begin{bmatrix} \sqrt{2} \\ 0 \end{bmatrix} && \text{by (1d) lemma} \\ &= \frac{\sqrt{2}}{2}\begin{bmatrix} 1 \\ 1 \end{bmatrix} \end{aligned}$$

[28] Haar (1910), Wojtaszczyk (1997) pages 14–15 ⟨"Sources and comments"⟩

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

2. Proof that (B) $\Longrightarrow$ (C):

$$
\begin{aligned}
\text{(B)} \Longrightarrow \phi(x) &= \sum_{k\in\mathbb{Z}} h_k\sqrt{2}\phi(2x-k) && \textit{dilation equation} && \text{(Theorem ?? page ??)}\\
&= \sum_{k=0}^{k=1}\left(\frac{\sqrt{2}}{2}\right)\sqrt{2}\phi(2x-k) && \text{by item (1e) page 396}\\
&= \sum_{k=0}^{k=1}\phi(2x-k)\\
&= \sum_{k=0}^{k=1}\binom{1}{k}\phi(2x-k) && \text{by definition of } \binom{n}{k} && \text{(Definition ?? page ??)}\\
&\Longrightarrow \text{(D)} && \text{by } \textit{B-spline dyadic decomposition} && \text{(Theorem R.9 page 391)}
\end{aligned}
$$

3. Proof that (B) $\Longleftarrow$ (C):

$$
\begin{aligned}
\text{(C)} \Longrightarrow \mathsf{N}_0(x) &= \sum_{k=0}^{k=1}\binom{1}{k}\mathsf{N}_0(2x-k) && \text{by } \textit{B-spline dyadic decomposition} && \text{(Theorem R.9 page 391)}\\
&= \sum_{k=0}^{k=1}\left(\frac{\sqrt{2}}{2}\right)\sqrt{2}\mathsf{N}_0(2x-k) && \text{by definition of } \binom{n}{k} && \text{(Definition ?? page ??)}\\
&= \sum_{k\in\mathbb{Z}} h_k\sqrt{2}\mathsf{N}_0(2x-k) && \text{by definition of } \binom{n}{k} && \text{(Definition ?? page ??)}\\
&\Longrightarrow \text{(B)}
\end{aligned}
$$

4. Proof that (A) $\Longleftarrow$ (C):
   1. Proof that (C) $\Longrightarrow$ $\operatorname{supp}\phi(x) = [0 : 1]$: by Theorem R.4 (page 378)
   2. Proof that (C) $\Longrightarrow$ $(\!(\phi(x-k))\!)$ forms a *partition of unity*: by Theorem R.7 (page 384)

Copyright © 2019 Daniel J. Greenhoe
BY-NC-ND

EX

n=0,	(÷0!)	1;							
n=1,	(÷1!)	1,	0;						
		−1,	2;						
n=2,	(÷2!)	1,	0,	0;					
		−2,	6,	−3;					
		1,	−6,	9;					
n=3,	(÷3!)	1,	0,	0,	0;				
		−3,	12,	−12,	4;				
		3,	−24,	60,	−44;				
		−1,	12,	−48,	64;				
n=4,	(÷4!)	1,	0,	0,	0,	0;			
		−4,	20,	−30,	20,	−5;			
		6,	−60,	210,	−300,	155;			
		−4,	60,	−330,	780,	−655;			
		1,	−20,	150,	−500,	625;			
n=5,	(÷5!)	1,	0,	0,	0,	0,	0;		
		−5,	30,	−60,	60,	−30,	6;		
		10,	−120,	540,	−1140,	1170,	−474;		
		−10,	180,	−1260,	4260,	−6930,	4386;		
		5,	−120,	1140,	−5340,	12270,	−10974;		
		−1,	30,	−360,	2160,	−6480,	7776;		
n=6,	(÷6!)	1,	0,	0,	0,	0,	0,	0;	
		−6,	42,	−105,	140,	−105,	42,	−7;	
		15,	−210,	1155,	−3220,	4935,	−3990,	1337;	
		−20,	420,	−3570,	15680,	−37590,	47040,	−24178;	
		15,	−420,	4830,	−29120,	96810,	−168000,	119182;	
		−6,	210,	−3045,	23380,	−100065,	225750,	−208943;	
		1,	−42,	735,	−6860,	36015,	−100842,	117649;	
n=7,	(÷7!)	1,	0,	0,	0,	0,	0,	0,	0;
		−7,	56,	−168,	280,	−280,	168,	−56,	8;
		21,	−336,	2184,	−7560,	15400,	−18648,	12488,	−3576;
		−35,	840,	−8400,	45360,	−143360,	267120,	−273280,	118896;
		35,	−1120,	15120,	−111440,	483840,	−1238160,	1733760,	−1027984;
		−21,	840,	−14280,	133560,	−741160,	2436840,	−4391240,	3347016;
		7,	−336,	6888,	−78120,	528920,	−2135448,	4753336,	−4491192;
		−1,	56,	−1344,	17920,	−143360,	688128,	−1835008,	2097152;
n=8,	(÷8!)	1,	0,	0,	0,	0,	0,	0,	0,	0;
		−8,	72,	−252,	504,	−630,	504,	−252,	72,	−9
		28,	−504,	3780,	−15624,	39690,	−64008,	64260,	−36792,	9207
		−56,	1512,	−17388,	111384,	−436590,	1079064,	−1650348,	1432872,	−541917
		70,	−2520,	39060,	−340200,	1821330,	−6146280,	12800340,	−15082200,	7715619
		−56,	2520,	−49140,	541800,	−3691170,	15903720,	−42324660,	63667800,	−41503131
		28,	−1512,	35532,	−474264,	3929310,	−20674584,	67410252,	−124449192,	99584613
		−8,	504,	−13860,	217224,	−2121210,	13208328,	−51179940,	112731192,	−107948223
		1,	−72,	2268,	−40824,	459270,	−3306744,	14880348,	−38263752,	43046721

Table R.1: Coefficients of the *B-splines* $\mathsf{N}_n(x)$ multiplied by $n!$ (Example R.9 page 374)

Copyright © 2019 Daniel J. Greenhoe

APPENDIX S

SOURCE CODE

The free and open source software package Maxima has been used to compute some of the algebraic expressions for *B-spline*s used in APPENDIX R (page 367):

```
/*=====================================================================*
 * Daniel J. Greenhoe                                                  *
 * Maxima script file                                                  *
 * To execute this script, start Maxima in a command window            *
 * in the subdirectory containing this file (e.g. c:\math\maxima\)     *
 * and then after the (%i...) prompt enter                             *
 *   batchload("bspline.max")$                                         *
 * Data produced will be written to the file "bsplineout.txt".         *
 * reference: http://maxima.sourceforge.net/documentation.html         *
 *=====================================================================*/
/*-----------------------------------------
 * initialize script
 *-----------------------------------------*/
reset();
kill(all);
load(orthopoly);
display2d:false; /* 2-dimensional display */
writefile("bsplineout.txt");
/*-----------------------------------------
 * n = B-spline order parameter
 * may be set to any value in {1,2,3,...}
 *-----------------------------------------*/
n:2;
print("=====================================================================");
print("Daniel J. Greenhoe");
print("Output file for nth order B-spline Nn(x) calculation, n=",n," .");
print("Output produced using Maxima running the script file bspline.max");
print("=====================================================================");
Nnx:(1/n!)*sum((-1)^k*binomial(n+1,k)*(x-k)^n*unit_step(x-k),k,0,n+1);
print("---------------------------------------------------------------");
print("              n+1      k (n+1)       n                    ");
print(" n! Nn(x) = SUM (-1)  (    ) (x-k)  step(x-k)  ,n=",n);
print("              k=0       ( k )                            ");
print("              ",n+1,"      k (",n+1,")      ",n);
print(n,"! Nn(x) = SUM (-1)  (    ) (x-k)  step(x-k)");
print("              k=0       ( k )");
print("           = ",expand(n!*Nnx));
print("---------------------------------------------------------------");
assume(x<=0);    print(n!,"N(x)= ",expand(n!*Nnx)," for x<=0");  forget(x<=0);
for i:0 thru n step 1 do(
  assume(x>i,x<(i+1)),
  print(n!,"N(x)= ",expand(n!*Nnx)," for ",i,"<x<",i+1),
  tex(expand(n!*Nnx),"djg.tex"),/*write output in TeX format to file "djg.tex"*/
  forget(x>i,x<(i+1))
  );
assume(x>(n+1)); print(n!,"N(x)= ",expand(n!*Nnx)," for x>",n+1); forget(x>(n+1));
```

```
print("-----------------------------------------------------------");
print(" values at some specific points x:                          ");
print("-----------------------------------------------------------");
y:Nnx,x=(n+1)/2;print("N(",(n+1)/2,")= ",y," (center value)");
y:Nnx,x=(n+2)/2;print("N(",(n+2)/2,")= ",y);
y:Nnx,x=(n+3)/2;print("N(",(n+3)/2,")= ",y);
/*-----------------------------------------
 * close output file
 *-----------------------------------------*/
closefile();
```

Once the polynomial expressions for a *B-spline* have been calculated, they can be plotted within a LaTeX environment using the pst-plot package along with a LaTeX source file such as the following:[1]




```
%==========================================================================
% Daniel J. Greenhoe
% LaTeX file
% N_3(x) B-spline
% nominal unit = 10mm
%==========================================================================
\begin{pspicture}(-1,-0.5)(5,1)
  %-----------------------------------------
  % parameters
  %-----------------------------------------
  \psset{plotpoints=64,labelsep=1pt}%
  %-----------------------------------------
  % axes
  %-----------------------------------------
  \psaxes[linewidth=0.75pt,linecolor=axis,yAxis=false,ticks=x,labels=x]{<->}(0,0)(-1,0)(5,1)% x axis
  \psaxes[linewidth=0.75pt,linecolor=axis,xAxis=false,ticks=x,labels=x]{->}(0,0)(-1,0)(5,1)%  y axis
  %-----------------------------------------
  % annotation
  %-----------------------------------------
  \psline[linestyle=dashed,linewidth=0.75pt,linecolor=red](2,0)(2,0.667)%
  \psline[linestyle=dashed,linewidth=0.75pt,linecolor=red](0,0.667)(2,0.667)%
  \psline[linestyle=dashed,linewidth=0.75pt,linecolor=red](1,0)(1,0.1667)%
  \psline[linestyle=dashed,linewidth=0.75pt,linecolor=red](3,0)(3,0.1667)%
  \psline[linestyle=dashed,linewidth=0.75pt,linecolor=red](0,0.1667)(3,0.1667)%
  \uput[180](0,0.667){$\sfrac{2}{3}$}%
  \uput[180](0,0.1667){$\sfrac{1}{6}$}%
  %-----------------------------------------
  % function plot
  %-----------------------------------------
  \psplot{0}{1}{+1 x 3 exp mul                                          6 div}% for 0<=x<=1
  \psplot{1}{2}{-3 x 3 exp mul +12 x 2 exp mul add -12 x mul add  +4 add 6 div}% for 1<=x<=2
  \psplot{2}{3}{+3 x 3 exp mul -24 x 2 exp mul add +60 x mul add -44 add 6 div}% for 2<=x<=3
  \psplot{3}{4}{-1 x 3 exp mul +12 x 2 exp mul add -48 x mul add +64 add 6 div}% for 3<=x<=4
  \psline(0,0)(-0.5,0)\psline[linestyle=dotted](-0.5,0)(-0.75,0)%              % for x<=0
  \psline(4,0)(4.5,0)\psline[linestyle=dotted](4.5,0)(4.75,0)%                 % for x>=4
\end{pspicture}%
```

Alternatively, one can plot $\mathsf{N}_3(x)$ more or less directly from the equation given in Theorem R.1 (page 371) without first calculating the polynomial expressions:




```
%==========================================================================
% Daniel J. Greenhoe
% LaTeX file
% N_3(x) B-spline
% nominal unit = 10mm
%==========================================================================
\begin{pspicture}(-1,-0.5)(5,1)
  %-----------------------------------------
  % parameters
  %-----------------------------------------
  \psset{plotpoints=64,labelsep=1pt}%
```

[1] For help with PostScript® math operators, see Adobe (1999), pages 508–509 ⟨Arithmetic and Math Operators⟩.

BY-NC-ND Copyright © 2019 Daniel J. Greenhoe

```
  %-------------------------------------
  % axes
  %-------------------------------------
  \psaxes[linewidth=0.75pt,linecolor=axis,yAxis=false,ticks=x,labels=x]{<->}(0,0)(-1,0)(5,1)% x axis
  \psaxes[linewidth=0.75pt,linecolor=axis,xAxis=false,ticks=x,labels=x]{->}(0,0)(-1,0)(5,1)%  y axis
  %-------------------------------------
  % annotation
  %-------------------------------------
  \psline[linestyle=dashed,linewidth=0.75pt,linecolor=red](2,0)(2,0.667)%
  \psline[linestyle=dashed,linewidth=0.75pt,linecolor=red](0,0.667)(2,0.667)%
  \psline[linestyle=dashed,linewidth=0.75pt,linecolor=red](1,0)(1,0.1667)%
  \psline[linestyle=dashed,linewidth=0.75pt,linecolor=red](3,0)(3,0.1667)%
  \psline[linestyle=dashed,linewidth=0.75pt,linecolor=red](0,0.1667)(3,0.1667)%
  \uput[180](0,0.667){$\sfrac{2}{3}$}%
  \uput[180](0,0.1667){$\sfrac{1}{6}$}%
  %-----------------------------------------------------------------------------
  % for n=3
  %        1  n+1        (n+1)                    1   4          ( 4 )
  % N_n(x)--- SUM (-1)^k (    ) (x-k)^n s(x-k) = --- SUM  (-1)^k (    ) (x-k)^3 s(x-k)
  %        n! k=1        ( k )                    3! k=1         ( k )
  %
  % where s(x) = 0 for x<0 and 1 for x>=0  (step function)
  %-----------------------------------------------------------------------------
  \psplot{0}{1}  {1 x 0 sub 3 exp mul  6 div}%
  \psplot{1}{2}  {1 x 0 sub 3 exp mul  4 x 1 sub 3 exp mul sub  6 div}%
  \psplot{2}{3}  {1 x 0 sub 3 exp mul  4 x 1 sub 3 exp mul sub  6 x 2 sub 3 exp mul add  6 div}%
  \psplot{3}{4}  {1 x 0 sub 3 exp mul  4 x 1 sub 3 exp mul sub  6 x 2 sub 3 exp mul add  4 x 3 sub
            3 exp mul sub  6 div}%
  \psplot{4}{4.5}{1 x 0 sub 3 exp mul  4 x 1 sub 3 exp mul sub  6 x 2 sub 3 exp mul add  4 x 3 sub
            3 exp mul sub  1 x 4 sub 3 exp mul add 6 div}%
  %    N_3(x) = [(4choose0)(x-0)^3    - (4choose1)(x-1)^3      + (4choose2)(x-2)^3      -
            (4choose3)(x-3)^3      + (4choose4)(x-4)^3      ]/3!
  %           = [        1 (x-0)^3    -        4 (x-1)^3      +        6 (x-2)^3      -        4
            (x-3)^3      +        1 (x-4)^3      ]/6
  \psline(0,0)(-0.5,0)%
  \psline[linestyle=dotted](-0.5,0)(-0.75,0)%
  \psline[linestyle=dotted](4.5,0)(4.75,0)%
\end{pspicture}%
```


Copyright © 2019 Daniel J. Greenhoe

Copyright © 2019 Daniel J. Greenhoe

Back Matter

❝ It appears to me that if one wants to make progress in mathematics, one should study the masters and not the pupils.❞

Niels Henrik Abel (1802–1829), Norwegian mathematician [2]



❝ When evening comes, I return home and go to my study. On the threshold I strip naked, taking off my muddy, sweaty workaday clothes, and put on the robes of court and palace, and in this graver dress I enter the courts of the ancients and am welcomed by them, and there I taste the food that alone is mine, and for which I was born. And there I make bold to speak to them and ask the motives of their actions, and they, in their humanity reply to me. And for the space of four hours I forget the world, remember no vexation, fear poverty no more, tremble no more at death; I pass indeed into their world.❞

Niccolò Machiavelli (1469–1527), Italian political philosopher, in a 1513 letter to friend Francesco Vettori. [3]



ancient library of Alexandria



The Book Worm by *Carl Spitzweg*, circa 1850 [4]

❝ To sit alone in the lamplight with a book spread out before you, and hold intimate converse with men of unseen generations—such is a pleasure beyond compare.❞

Yoshida Kenko (Urabe Kaneyoshi) (1283? – 1350?), Japanese author and Buddhist monk [5]

[2] quote: Simmons (2007), page 187.
image: http://en.wikipedia.org/wiki/Image:Niels_Henrik_Abel.jpg, public domain

[3] quote: Machiavelli (1961), page 139?.
image: http://commons.wikimedia.org/wiki/File:Santi_di_Tito_-_Niccolo_Machiavelli%27s_portrait_headcrop.jpg, public domain

[4] http://en.wikipedia.org/wiki/File:Ancientlibraryalex.jpg, public domain http://en.wikipedia.org/wiki/File:Carl_Spitzweg_021.jpg,

Copyright © 2019 Daniel J. Greenhoe

[5] quote: Kenko (circa 1330)
image: http://en.wikipedia.org/wiki/Yoshida_Kenko

BY-NC-ND
Copyright © 2019 Daniel J. Greenhoe

BIBLIOGRAPHY

M. S. Abdaheer. *Digital Signal Processing*. University Science Press, New Delhi, 2009.

M. Abom. A note on random errors in frequency response estimators. *Journal of Sound and Vibration*, 107(2):355–358, 1986. doi: 10.1016/0022-460X(86)90244-0. URL http://adsabs.harvard.edu/abs/1986JSV...107..355A.

Yuri A. Abramovich and Charalambos D. Aliprantis. *An Invitation to Operator Theory*. American Mathematical Society, Providence, Rhode Island, 2002. ISBN 0-8218-2146-6. URL http://books.google.com/books?vid=ISBN0821821466.

Adobe. *PostScript Language Reference*. Addison-Wesley Publishing Company, Reading, Massachusetts, 1999. ISBN 0-201-37922-8. URL http://wwwimages.adobe.com/content/dam/Adobe/en/devnet/postscript/pdfs/PLRM.pdf.

N. I. Akhiezer and I. M. Glazman. *Theory of Linear Operators in Hilbert Spaces*, volume 1. Dover, New York, 1993. URL http://books.google.com/books?vid=ISBN0486677486. Translated from the original Russian text *Teoriia lineinykh operatorov v Gil'bertovom prostranstve*.

Charalambos D. Aliprantis and Owen Burkinshaw. *Principles of Real Analysis*. Acedemic Press, London, 3 edition, 1998. ISBN 9780120502578. URL http://www.amazon.com/dp/0120502577.

Charalambos D. Aliprantis and Owen Burkinshaw. *Positive Operators*. Springer, Dordrecht, 2006. ISBN 9781402050077. URL http://books.google.com/books?vid=ISBN1402050070. reprint of Academic Press 1985 edition.

Randall J. Allemang, R. Zimmerman, and David L. Brown. Technlqucs of reducing noise in frquency response measurements. *ASME Winter Annual Meeting*, 1979. session on Application of System Identlfication Techniques.

Randall J. Allemang, David L. Brown, and Robert W. Rost. *Experimental Modal Analysis And Dynamic Component Synthesis*, 2(AFWAL-TR-87-3069), December 1987. URL https://apps.dtic.mil/dtic/tr/fulltext/u2/a195145.pdf.

Theodore T. Allen. *Introduction to Engineering Statistics and Lean Six Sigma: Statistical Quality Control and Design of Experiments and Systems*. Springer, 3 edition, 2018. ISBN 9781447174202.

Herbert Amann and Joachim Escher. *Analysis II*. Birkhäauser Verlag AG, Basel–Boston–Berlin, 2008. ISBN 978-3-7643-7472-3. URL http://books.google.com/books?vid=ISBN3764374721.

Dan Amir. *Characterizations of Inner Product Spaces*, volume 20 of *Operator Theory: Advances and Applications*. Birkhäuser Verlag, Basel, 1986. ISBN 3-7643-1774-4. URL http://www.worldcat.org/isbn/3764317744.

Theodore Wilbur Anderson. *An introduction to multivariate statistical analysis*. Wiley series in probability and mathematical statistics. Wiley, 1 edition, 1958. URL https://books.google.com/books?id=7YpqAAAAMAAJ.

Theodore Wilbur Anderson. *An Introduction to Multivariate Statistical Analysis*, volume 114 of *Wiley Series in Probability and Statistics—Applied Probability and Statistics Section Series*. Wiley, 2 edition, 1984. ISBN 9780471889878. URL http://books.google.com/books?vid=ISBN9780471889878.

John E. Angus. The probability integral transform and related results. *SIAM Review*, 36(4):652–654, February 25 1994. doi: 10.1137/1036146. URL http://dx.doi.org/10.1137/1036146.

Tom M. Apostol. *Mathematical Analysis*. Addison-Wesley series in mathematics. Addison-Wesley, Reading, 2 edition, 1975. ISBN 986-154-103-9. URL http://books.google.com/books?vid=ISBN0201002884.

Jean-Pierre Aubin. *Applied Functional Analysis*, volume 47 of *Pure and Applied Mathematics: A Wiley Series of Texts, Monographs and Tracts*. John Wiley & Sons, 2 edition, September 30 2011. ISBN 9781118030974. URL http://books.google.com/books?vid=ISBN1118030974.

Jean-Pierre Aubin and Hélène Frankowska. *Set-Valued Analysis*. Modern Birkhäuser Classics. Springer, March 2 2009. ISBN 9780817648480. URL http://books.google.com/books?vid=ISBN0817648488.

Léon Autonne. Sur l'hermitien (on the hermitian). In *Comptes Rendus Des SéAnces De L'AcadéMie Des Sciences*, volume 133, pages 209–268. De L'Académie des sciences (Academy of Sciences), Paris, 1901. URL http://visualiseur.bnf.fr/Visualiseur?O=NUMM-3089. Comptes Rendus Des SéAnces De L'AcadéMie Des Sciences (Reports Of the Meetings Of the Academy of Science).

Léon Autonne. Sur l'hermitien (on the hermitian). *Rendiconti del Circolo Matematico di Palermo*, 16:104–128, 1902. Rendiconti del Circolo Matematico di Palermo (Statements of the Mathematical Circle of Palermo).

George Bachman. *Elements of Abstract Harmonic Analysis*. Academic paperbacks. Academic Press, New York, 1964. URL http://books.google.com/books?id=ZP8-AAAAIAAJ.

George Bachman and Lawrence Narici. *Functional Analysis*. Academic Press textbooks in mathematics; Pure and Applied Mathematics Series. Academic Press, 1 edition, 1966. ISBN 9780486402512. URL http://books.google.com/books?vid=ISBN0486402517. "unabridged republication" available from Dover (isbn 0486402517).

George Bachman, Lawrence Narici, and Edward Beckenstein. *Fourier and Wavelet Analysis*. Universitext Series. Springer, 2000. ISBN 9780387988993. URL http://books.google.com/books?vid=ISBN0387988998.

David H. Bailey, Jonathan M. Borwein, and Andrew Mattingly. The computation of previously inaccessible digits of π^2 and catalan's constant (2013). page 23, March 24 2011. URL https://pdfs.semanticscholar.org/0b15/22a1b2f76729e19be2be1bca9f57031a4db9.pdf.

Copyright © 2019 Daniel J. Greenhoe

David H. Bailey, Jonathan M. Borwein, and Andrew Mattingly. 20. the computation of previously inaccessible digits of π^2 and catalan's constant (2013). In *Pi: The Next Generation: A Sourcebook on the Recent History of Pi and Its Computation*, pages 324–339. Springer, 2013. ISBN 9783319323770. URL http://books.google.com/books?vid=ISBN3319323776.

Om Prakash Bajpai. *Foundations of statistics*. Asia Publishing House, 1967. URL https://books.google.com/books?id=QrPuAAAAMAAJ.

Per Bak. *How Nature Works: the science of self-organized criticality*. Springer Science & Business Media, 2013. ISBN 9781475754261. URL http://books.google.com/books?vid=ISBN1475754264.

N Balakrishnan and Chin-Diew Lai. *Continuous Bivariate Distributions*. Springer Science & Business Media, 2 edition, May 31 2009. ISBN 9780387096148. URL http://books.google.com/books?vid=ISBN0387096140.

Stefan Banach. Sur les opérations dans les ensembles abstraits et leur applications aux équations intégrales (on abstract operations and their applications to the integral equations). *Fundamenta Mathematicae*, 3:133–181, 1922. URL http://matwbn.icm.edu.pl/ksiazki/fm/fm3/fm3120.pdf.

Stefan Banach. *Théorie des opérations linéaires*. Monografje Matematyczne, Warsaw, Poland, 1932a. URL http://matwbn.icm.edu.pl/kstresc.php?tom=1&wyd=10. (Theory of linear operations).

Stefan Banach. *Theory of Linear Operations*, volume 38 of *North-Holland mathematical library*. North-Holland, Amsterdam, 1932b. ISBN 0444701842. URL http://www.amazon.com/dp/0444701842/. English translation of 1932 French edition, published in 1987.

Alexander Barvinok. *A Course in Convexity*, volume 54 of *Graduate studies in mathematics*. American Mathematical Society, 2002. ISBN 9780821872314. URL http://books.google.com/books?vid=ISBN0821872311.

Gerald Beer. *Topologies on Closed and Closed Convex Sets*, volume 268 of *Mathematics and Its Applications*. October 31 1993. ISBN 9780792325314. URL http://books.google.com/books?vid=ISBN0792325311.

Eric Temple Bell. *Men of Mathematics*. Simon & Schuster, New York, 1986. ISBN 9780671628185. URL http://books.google.com/books?vid=ISBN0671628186.

J. S. Bendat. Statistical errors in measurement of coherence functions and input/output quantities. *Journal of Sound and Vibration*, 59(3):405–421, August 08 1978. doi: 10.1016/S0022-460X(78)80007-8. URL https://doi.org/10.1016/S0022-460X(78)80007-8.

Julius S. Bendat and Allan G. Piersol. *Measurement and Analysis of Random Data*. John Wiley & Sons, 1966.

Julius S. Bendat and Allan G. Piersol. *Engineering Applications of Correlation and Spectral Analysis*. John Wiley & Sons, 1980. ISBN 9780471058878. URL http://www.amazon.com/dp/0471058874.

Julius S. Bendat and Allan G. Piersol. *Engineering Applications of Correlation and Spectral Analysis*. Wiley-Interscience, 2 edition, 1993. ISBN 9780471570554. URL http://www.amazon.com/dp/0471570559.

Julius S. Bendat and Allan G. Piersol. *Random Data: Analysis and Measurement Procedures*, volume 729 of *Wiley Series in Probability and Statistics*. John Wiley & Sons, 4 edition, 2010. ISBN 9781118210826. URL http://books.google.com/books?vid=ISBN1118210824.

Copyright © 2019 Daniel J. Greenhoe

Sterling Khazag Berberian. *Introduction to Hilbert Space*. Oxford University Press, New York, 1961. URL http://books.google.com/books?vid=ISBN0821819127.

M. Bertero and P. Boccacci. *Introduction to Inverse Problems in Imaging*. CRC Press, 1998. ISBN 9781439822067. URL http://books.google.com/books?vid=ISBN9781439822067.

D. Besso. Teoremi elementari sui massimi i minimi. *Annuari Ist. Tech. Roma*, pages 7–24, 1879. see Bullen(2003) pages 453, 203.

M. Jales Bienaymé. Société philomatique de paris—extraits des procès-verbaux. *Scéance*, pages 67–68, June 13 1840. URL http://www.archive.org/details/extraitsdesproc46183941soci. see Bullen(2003) pages 453, 203.

Patrick Billingsley. *Probability And Measure*. Wiley series in probability and mathematical statistics. Wiley, 3 edition, 1995. ISBN 978-0471007104. URL http://books.google.com/books?vid=ISBN0471007102.

Leonard Mascot Blumenthal. *Theory and Applications of Distance Geometry*. Chelsea Publishing Company, Bronx, New York, USA, 2 edition, 1970. ISBN 0-8284-0242-6. URL http://books.google.com/books?vid=ISBN0828402426.

Béla Bollobás. *Linear Analysis; an introductory course*. Cambridge mathematical textbooks. Cambridge University Press, Cambridge, 2 edition, March 1 1999. ISBN 978-0521655774. URL http://books.google.com/books?vid=ISBN0521655773.

William M. Bolstad. *Introduction to Bayesian Statistics*. Wiley, 2 edition, 2007. ISBN 9780470141151. URL http://books.google.com/books?vid=ISBN9780470141151.

V. Bouniakowsky. Sur quelques inégalités concernant les intégrales ordinaires et les intégrales aux différences finies. *Mémoires de lÁcad. de St.-Pétersbourg*, 1(9), 1859. URL http://www-stat.wharton.upenn.edu/~steele/Publications/Books/CSMC/CSMC_index.html.

Carl Benjamin Boyer and Uta C. Merzbach. *A History of Mathematics*. Wiley, New York, 3 edition, January 11 2011. ISBN 9780470525487. URL http://books.google.com/books?vid=ISBN0470525487.

Ronald Newbold Bracewell. *The Fourier transform and its applications*. McGraw-Hill electrical and electronic engineering series. McGraw-Hill, 2, illustrated, international student edition edition, 1978. ISBN 9780070070134. URL http://books.google.com/books?vid=ISBN007007013X.

Thomas John I'Anson Bromwich. *An Introduction to the Theory of Infinite Series*. Macmillan and Company, 1 edition, 1908. ISBN 9780821839768. URL http://www.archive.org/details/anintroductiont00bromgoog.

P. S. Bullen. *Handbook of Means and Their Inequalities*, volume 560 of *Mathematics and Its Applications*. Kluwer Academic Publishers, Dordrecht, Boston, 2 edition, 2003. ISBN 9781402015229. URL http://books.google.com/books?vid=ISBN1402015224.

Viktor Yakovlevich Bunyakovsky. Sur quelques inégalités concernant les intégrales ordinaires et les intégrales aux différences finies. *Mémoires de L'Académie Impériale des Sciences de St.-Pétersbourg*, 1(9):1–18, 1859. URL http://www-stat.wharton.upenn.edu/~steele/Publications/Books/CSMC/bunyakovsky.pdf. (Mémoires of the Imperial Academy of Sciences of Saint Petersburg); (On Some Inequalities concerning the Ordinary Integrals and Integrals with the Finite Differences).

BY-NC-ND
Copyright © 2019 Daniel J. Greenhoe

Charles L. Byrne. *Signal Processing: A Mathematical Approach*. AK Peters Series. A K Peters, 2005. ISBN 1568812426.

James A. Cadzow. *Foundations of Digital Signal Processing and Data Analysis*. Macmillan Publishing Company, New York, 1987. ISBN 0023180102. URL http://frontweb.vuse.vanderbilt.edu/vuse_web/directory/facultybio.asp?FacultyID=9.

Peter E. Caines. *Linear Stochastic Systems*, volume 100 of *Wiley Series in Probability and Statistics*. J. Wiley, 1988. ISBN 9780471081012. URL http://books.google.com/books?vid=ISBN0471081019.

Peter E. Caines. *Linear Stochastic Systems*, volume 77 of *Classics in Applied Mathematics*. SIAM, 2018. ISBN 9781611974713. URL http://books.google.com/books?vid=ISBN1611974712.

Florian Cajori. A history of mathematical notations; notations mainly in higher mathematics. In *A History of Mathematical Notations; Two Volumes Bound as One*, volume 2. Dover, Mineola, New York, USA, 1993. ISBN 0-486-67766-4. URL http://books.google.com/books?vid=ISBN0486677664. reprint of 1929 edition by *The Open Court Publishing Company*.

James J. Callahan. *Advanced Calculus: A Geometric View*. Springer Science & Business Media, September 9 2010. ISBN 9781441973320. URL http://books.google.com/books?vid=ISBN144197332X.

Gerolamo Cardano. *Ars Magna or the Rules of Algebra*. Dover Publications, Mineola, New York, 1545. ISBN 0486458733. URL http://www.amazon.com/dp/0486458733. English translation of the Latin *Ars Magna* edition, published in 2007.

Thomas G. Carne and C. R. Dohrmann. Improving experimental frequency response function matrices for admittance modeling. In *Proceedings of the XXIV International Modal Analysis Conference (IMAC)*, St. Louis, Missouri, USA, 2006. URL https://www.researchgate.net/publication/265790751.

N. L. Carothers. *A Short Course on Banach Space Theory*. Number 64 in London Mathematical Society Student Texts. Cambridge University Press, Cambridge, 2005. ISBN 0521842832. URL http://www.comm.musashi-tech.ac.jp/~arimoto/Drarimoto/preprints/calothers/Assemble.ps.

N.L. Carothers. *Real Analysis*. Cambridge University Press, Cambridge, 2000. ISBN 978-0521497565. URL http://books.google.com/books?vid=ISBN0521497566.

Augustin-Louis Cauchy. *Part 1: Analyse Algebrique*. Cours D'Analyse de L'école Royale Polytechnique. Debure frères, Paris, 1821. ISBN 2-87647-053-5. URL http://www.archive.org/details/coursdanalysede00caucgoog. Systems design course of the Polytechnic Royal School; 1st Part: Algebraic analysis).

Arthur Cayley. A memoir on the theory of matrices. *Philosophical Transactions of the Royal Society of London*, 148:17–37, 1858. ISSN 1364-503X. URL http://www.jstor.org/view/02610523/ap000059/00a00020/0.

Si-Wei Chen, Xue-Song Wang, and Motoyuki Sato. Polinsar complex coherence estimation based on covariance matrix similarity test. *IEEE Transactions on Geoscience and Remote Sensing*, 50(11):4699–4710, November 2012. doi: 10.1109/TGRS.2012.2192937. URL https://www.researchgate.net/profile/Si_Wei_Chen/publication/281271737.

Sung C. Choi. *Introductory applied statistics in science*. Prentice-Hall, 1978. ISBN 9780135016190. URL http://books.google.com/books?vid=ISBN9780135016190.

Copyright © 2019 Daniel J. Greenhoe

Alexandre J. Chorin and Ole H. Hald. *Stochastic Tools in Mathematics and Science*, volume 1 of *Surveys and Tutorials in the Applied Mathematical Sciences*. Springer, New York, 2 edition, 2009. ISBN 978-1-4419-1001-1. URL http://books.google.com/books?vid=ISBN9781441910011.

Ole Christensen. *An Introduction to Frames and Riesz Bases*. Applied and Numerical Harmonic Analysis. Birkhäuser, Boston/Basel/Berlin, 2003. ISBN 0-8176-4295-1. URL http://books.google.com/books?vid=ISBN0817642951.

Ole Christensen. *Frames and bases: An Introductory Course*. Applied and Numerical Harmonic Analysis. Birkhäuser, Boston/Basel/Berlin, 2008. ISBN 9780817646776. URL http://books.google.com/books?vid=ISBN0817646779.

Ole Christensen. *Functions, Spaces, and Expansions: Mathematical Tools in Physics and Engineering*. Applied and Numerical Harmonic Analysis. Birkhäuser, Boston/Basel/Berlin, 2010. ISBN 978-0-8176-4979-1. doi: 10.1007/978-0-8176-4980-7. URL http://books.google.com/books?vid=ISBN0817649794.

Charles K. Chui. *Multivariate Splines*, volume 54 of *CBMS-NSF regional conference series in applied mathematics*. SIAM, 1988. ISBN 1611970172. URL http://books.google.com/books?vid=ISBN1611970172.

Charles K. Chui. *An Introduction to Wavelets*. Academic Press, San Diego, California, USA, January 3 1992. ISBN 9780121745844. URL http://books.google.com/books?vid=ISBN0121745848.

Donato Michele Cifarelli and Eugenio Regazzini. De finetti's contribution to probability and statistics. *Statistical Science*, 11(4):253–282, 1996. URL http://www.jstor.org/stable/2246020.

Erhan Çinlar and Robert J Vanderbei. *Real and Convex Analysis*. Undergraduate Texts in Mathematics. Springer, January 4 2013. ISBN 1461452570. URL http://books.google.com/books?vid=ISBN1461452570.

Jon F. Claerbout. *Fundamentals of Geophysical Data Processing with Applications to Petroleum Prospecting*. International series in the earth and planetary sciences, Tab Mastering Electronics Series. McGraw-Hill, New York, 1976. ISBN 9780070111172. URL http://sep.stanford.edu/sep/prof/.

James A. Clarkson. Uniformly convex spaces. *Transactions of the American Mathematical Society*, 40(3):396–414, December 1936. URL http://www.jstor.org/stable/1989630.

Peter M. Clarkson. *Optimal and Adaptive Signal Processing*. Electronic Engineering Systems Series. CRC Press, 1993. ISBN 0849386098.

R. E. Cobb and L. D. Mitchell. Estimation of uncorrelated content in experimentally measured frequency response functions using three measurement channels. *Mechanical Systems and Signal Processing*, 4(6):449–461, November 1990. URL https://doi.org/10.1016/0888-3270(90)90045-M.

Richard E. Cobb. *Confidence Bands, Measurement Noise, and Multiple Input - Multiple Output Measurements Using the Three-Channel Frequency Response Function Estimator*. Ph.d. dissertation, May 1988. URL http://hdl.handle.net/10919/53675. "Dissertation submitted to the Faculty of the Virginia Polytechnic Institute and State University in partial fululllment of the requirements for the degree of Doctor of Philosophy in Mechanical Engineering".

Paul M. Cohn. *Basic Algebra; Groups, Rings and Fields*. Springer, December 6 2002. ISBN 1852335874. URL http://books.google.com/books?vid=isbn1852335874.

Copyright © 2019 Daniel J. Greenhoe

John B. Conway. *A Course in Functional Analysis*, volume 96 of *Graduate Texts in Mathematics*. Springer, 2 edition, 1990. ISBN 0-387-97245-5. URL http://books.google.com/books?vid=ISBN0387972455.

R. Courant and D. Hilbert. *Methods of Mathematical Physics*, volume 1. Interscience Publishers, New York, 1930. URL http://www.worldcat.org/isbn/0471504475.

T.M. Cover and Joy A. Thomas. *Elements of Information Theory*. John Wiley & Sons, Inc., New York, 1991. ISBN 0-471-06259-6. URL http://www.amazon.com/dp/0471062596.

M. G. Cox. The numerical evaluation of b-splines. *IMA Journal of Applied Mathematics*, 10(2):134–149, 1972. doi: 10.1093/imamat/10.2.134. URL http://imamat.oxfordjournals.org/content/10/2/134.abstract.

Guy L. Curry and Richard M. Feldman. *Manufacturing Systems Modeling and Analysis*. Technology & Engineering. Springer Science & Business Media, 2 edition, 2010. ISBN 9783642166181. URL http://books.google.com/books?vid=ISBN3642166180.

Ingrid Daubechies. *Ten Lectures on Wavelets*. Society for Industrial and Applied Mathematics, Philadelphia, 1992. ISBN 0-89871-274-2. URL http://www.amazon.com/dp/0898712742.

Ingrid Daubechies, A. Grossman, and Y. Meyer. Painless nonorthogonal expansions. *Journal of Mathematical Physics*, 27(5):1271–1283, May 1986. ISSN 0022-2488. URL link.aip.org/link/?jmp/27/1271.

Mahlon Marsh Day. *Normed Linear Spaces*, volume 21 of *Ergebnisse der Mathematik und ihrer Grenzgebiete*. Springer-Verlag, Berlin, Heidelberg, New York, 3 edition, 1973. ISBN 0387061487. URL http://books.google.com/books?id=UZ3vSAAACAAJ.

Bruno de Finetti. Sui passaggi al limite nel calcolo delle probabilità. *Rendiconti del Reale Istituto Lombardo di Scienze e Lettere*, 63:155–166, 1930a.

Bruno de Finetti. A proposito dell'estensione del teorema delle probabilita totali alle classi numerabili. *Rendiconti del Reale Istituto Lombardo di Scienze e Lettere*, 63:901–905, 1930b.

Bruno de Finetti. Ancora sull'estensione alle classi numerabili del teorema delle probabilità totali. *Rendiconti del Reale Istituto Lombardo di Scienze e Lettere*, 63:1063–1069, 1930c.

Johan de Witt. *Elementa curvarum linearum (Elements of Curves)*. 1659. Elementa curvarum linearum was an appendix to de Witt's translation of of Descartes' *La Géométrie*.

Carl R. de Boor. On calculating with b-splines. *Journal of Approximation Theory*, pages 50–62, 1972.

Carl R. de Boor. *A Practical Guide to Splines*, volume 27 of *Applied Mathematical Sciences*. Springer, revised edition, 2001. ISBN 0387953663. URL http://books.google.com/books?vid=ISBN0387953663.

F.M. Dekking, C. Kraaikamp, H.P. Lopuhaa, and L.E. Meester. *A Modern Introduction to Probability and Statistics: Understanding Why and How*. Springer Texts in Statistics. Springer Science & Business Media, 2006. ISBN 9781846281686. URL http://books.google.com/books?vid=ISBN1846281687.

René Descartes. *La géométrie*. 1637. URL http://historical.library.cornell.edu/math/math_D.html.

Copyright © 2019 Daniel J. Greenhoe

René Descartes. *Regulae ad directionem ingenii*. 1684a. URL http://www.fh-augsburg.de/~harsch/Chronologia/Lspost17/Descartes/des_re00.html.

René Descartes. *Rules for Direction of the Mind*. 1684b. URL http://en.wikisource.org/wiki/Rules_for_the_Direction_of_the_Mind.

René Descartes. *The Geometry of Rene Descartes*. Courier Dover Publications, June 1 1954. ISBN 0486600688. URL http://books.google.com/books?vid=isbn0486600688. orginally published by Open Court Publishing, Chicago, 1925; translation of La géométrie.

Luc Devroye. *Non-Uniform Random Variate Generation*. Springer-Verlag, New York, 1986. ISBN 0387963057. URL http://www.eirene.de/Devroye.pdf.

Jean Alexandre Dieudonné. *Foundations of Modern Analysis*. Academic Press, New York, 1969. ISBN 1406727911. URL http://books.google.com/books?vid=ISBN1406727911.

Hamid Drljević. On the representation of functionals and the stability of mappings in hilbert and banach spaces. In Themistocles M Rassias, editor, *Topics in mathematical analysis: a volume dedicated to the memory of A.L. Cauchy*, volume II of *Series in Pure Mathematics*, pages 231–245. World Scientific Publishing Company, 1989. ISBN 9971506661. URL http://books.google.com/books?vid=ISBN9971506661.

R. J. Duffin and A. C. Schaeffer. A class of nonharmonic fourier series. *Transactions of the American Mathematical Society*, 72(2):341–366, March 1952. ISSN 1088-6850. URL http://www.jstor.org/stable/1990760.

Bogdan Dumitrescu. *Positive Trigonometric Polynomials and Signal Processing Applications*. Signals and Communication Technology. Springer, 2007. ISBN 978-1-4020-5124-1. URL gen.lib.rus.ec/get?md5=5346e169091b2d928d8333cd053300f9.

Nelson Dunford and Jacob T. Schwartz. *Linear operators. Part 1, General Theory*, volume 7 of *Pure and applied mathematics*. Interscience Publishers, New York, 1957. ISBN 0471226394. URL http://www.amazon.com/dp/0471608483. with the assistance of William G. Bade and Robert G. Bartle.

John R. Durbin. *Modern Algebra; An Introduction*. John Wiley & Sons, Inc., 4 edition, 2000. ISBN 0-471-32147-8. URL http://www.worldcat.org/isbn/0471321478.

W.D. Duthie. Segments of ordered sets. *Transactions of the American Mathematical Society*, 51(1): 1–14, January 1942. doi: 10.2307/1989978. URL http://www.jstor.org/stable/1989978.

Jacques Dutka. On gauss' priority in the discovery of the method of least squares. *Archive for History of Exact Sciences*, 49(4):355–370, December 1995. URL https://www.jstor.org/stable/41134010.

Paul H. Edelman. Abstract convexity and meet-distributive lattices. In Ivan Rival, editor, *Combinatorics and ordered sets: Proceedings of the AMS-IMS-SIAM joint summer research conference, held August 11–17, 1985*, volume 57 of *Contemporary Mathematics*, pages 127–150, Providence RI, 1986. American Mathematical Society. ISBN 0821850512. URL http://books.google.com/books?vid=ISBN0821850512. conference held in Arcata California.

Paul H. Edelman and Robert E. Jamison. The theory of convex geometries. *Geometriae Dedicata*, 19(3):247–270, December 1985. ISSN 0046-5755. doi: 10.1007/BF00149365. URL http://www.springerlink.com/content/n4344856887387gw/.

Copyright © 2019 Daniel J. Greenhoe

Robert E. Edwards. *Functional Analysis: Theory and Applications*. Dover books on mathematics. Dover, New York, 1995. ISBN 0-486-68143-2. URL http://books.google.com/books?vid=ISBN0486681432.

Yuli Eidelman, Vitali D. Milman, and Antonis Tsolomitis. *Functional Analysis: An Introduction*, volume 66 of *Graduate Studies in Mathematics*. American Mathematical Society, 2004. ISBN 0821836463. URL http://books.google.com/books?vid=ISBN0821836463.

Per Enflo. A counterexample to the approximation problem in banach spaces. *Acta Mathematica*, 130:309–317, 1973. URL http://link.springer.com/content/pdf/10.1007/BF02392270.

Euclid. *Elements*. circa 300BC. URL http://farside.ph.utexas.edu/euclid.html.

David Ewen. *The Book of Modern Composers*. Alfred A. Knopf, New York, 1950. URL http://books.google.com/books?id=yHw4AAAAIAAJ.

David Ewen. *The New Book of Modern Composers*. Alfred A. Knopf, New York, 3 edition, 1961. URL http://books.google.com/books?id=bZIaAAAAMAAJ.

D. J. Ewins. *Modal Testing: Theory and Practice*. Research Studies Press, "reprinted with amendments april 1986" edition, April 1986. ISBN 0863800173. URL http://www.amazon.com/dp/086380036X.

Marián Fabian, Petr Habala, Petr Hájek, Vicente Montesinos, Vaclav Zizler, and Václav Zizler. *Banach Space Theory: The Basis for Linear and Nonlinear Analysis*. CMS Books in Mathematics. Springer, 2010. ISBN 1441975144. URL http://books.google.com/books?vid=isbn1441975144.

Lorenzo Farina and Sergio Rinaldi. *Positive Linear Systems: Theory and Applications*. Pure and applied mathematics. John Wiley & Sons, 1 edition, July 3 2000. ISBN 9780471384564. URL http://books.google.com/books?vid=ISBN0471384569.

Richard M. Feldman and Ciriaco Valdez-Flores. *Applied Probability and Stochastic Processes*. Technology & Engineering. Springer Science & Business Media, 2 edition, 2010. ISBN 9783642051586. URL http://books.google.com/books?vid=ISBN3642051588.

Carlos A. Felippa. *Matrix Calculus*. University of Colorado at Boulder, August 18 1999. URL http://caswww.colorado.edu/courses.d/IFEM.d/.

William Feller. *An Introduction to Probability Theory and its Applications Volume I*. Wiley series in probability and mathematical statistics. John Wiley & Sons, 3, revised edition, 1970. ISBN 9780471257080. URL http://www.amazon.com/dp/0471257087.

R. A. Fisher. On the mathematical foundations of theoretical statistics. *Philosophical Transacations of the Royal Society*, January 1922. URL https://doi.org/10.1098/rsta.1922.0009.

G.L. Fix and G. Strang. Fourier analysis of the finite element method in ritz-galerkin theory. *Studies in Applied Mathematics*, 48:265–273, 1969.

Gerald B. Folland. *A Course in Abstract Harmonic Analysis*. Studies in Advanced Mathematics. CRC Press, Boca Raton, 1995. ISBN 0-8493-8490-7. URL http://books.google.com/books?vid=ISBN0849384907.

Brigitte Forster and Peter Massopust, editors. *Four Short Courses on Harmonic Analysis: Wavelets, Frames, Time-Frequency Methods, and Applications to Signal and Image Analysis*. Applied and Numerical Harmonic Analysis. Springer, November 19 2009. ISBN 9780817648909. URL http://books.google.com/books?vid=ISBN0817648909.

Copyright © 2019 Daniel J. Greenhoe

Jean-Baptiste-Joseph Fourier. Refroidissement séculaire du globe terrestre". In M. Gaston Darboux, editor, *Œuvres De Fourier*, volume 2, pages 271–288. Ministère de L'instruction Publique, Paris, France, April 1820. URL http://gallica.bnf.fr/ark:/12148/bpt6k33707/f276.image. original paper at pages 58–70.

Jean-Baptiste-Joseph Fourier. *Théorie Analytique de la Chaleur (The Analytical Theory of Heat)*. Chez Firmin Didot, pere et fils, Paris, 1822. URL http://books.google.com/books?vid=04X2vlqZx7hydlQUWEq&id=TDQJAAAAIAAJ.

Jean-Baptiste-Joseph Fourier. *The Analytical Theory of Heat (Théorie Analytique de la Chaleur)*. Cambridge University Press, Cambridge, February 20 1878. URL http://www.archive.org/details/analyticaltheory00fourrich. 1878 English translation of the original 1822 French edition. A 2003 Dover edition is also available: isbn 0486495310.

Maurice Fréchet. Sur l'extension du théorème des probabilités totales au cas d'une suite infinie d'evénéments. *Ren- diconti del Reale Istituto Lombardo di Scienze e Lettere*, 63:899–900, 1930a.

Maurice Fréchet. Sur l'extension du théorème des probabilités totales au cas d'une suite infinie d'evénéments. *Ren- diconti del Reale Istituto Lombardo di Scienze e Lettere*, 63:1059–1062, 1930b.

Erik Ivar Fredholm. Sur une nouvelle méthode pour la résolution du problème de dirichlet (on a new method for the resolution of the problem of dirichlet). */:Oefversigt af Kongl. Sv. Vetenskaps-Academiens Förhandlingar*, 57:39–66, 1900.

Erik Ivar Fredholm. Sur une classe d'equations fonctionnelles (on a class of functional equations). *Acta Mathematica*, 27(1):365–390, December 1903. ISSN 0001-5962. doi: 10.1007/BF02421317. URL http://www.springerlink.com/content/c41371137837p252/.

Ferdinand Georg Frobenius. Uber lineare substitutionen und bilineare formen. *Journal für die reine und angewandte Mathematik (Crelle's Journal)*, 84:1–63, 1878. ISSN 0075-4102. URL http://www.digizeitschriften.de/home/services/pdfterms/?ID=509796.

Ferdinand Georg Frobenius. Uber lineare substitutionen und bilineare formen. In Jean Pierre Serre, editor, *Gesammelte Abhandlungen (Collected Papers)*, volume I, pages 343–405. Springer, Berlin, 1968. URL http://www.worldcat.org/oclc/253015. reprint of Frobenius' 1878 paper.

Jürgen Fuchs. *Affine Lie Algebras and Quantum Groups: An Introduction, With Applications in Conformal Field Theory*. Cambridge Monographs on Mathematical Physics. Cambridge University Press, 1995. ISBN 052148412X. URL http://books.google.com/books?vid=ISBN052148412X.

Israel M. Gelfand. Normierte ringe. *Mat. Sbornik*, 9(51):3–24, 1941.

Israel M. Gelfand and Mark A. Naimark. *Normed Rings with an Involution and their Representations*, pages 240–274. Chelsea Publishing Company, Bronx, 1964. ISBN 0821820222. URL http://books.google.com/books?vid=ISBN0821820222.

Israel M. Gelfand and Mark A. Neumark. On the imbedding of normed rings into the ring of operators in hilbert space. *Mat. Sbornik*, 12(54:2):197–217, 1943a.

Israel M. Gelfand and Mark A. Neumark. *On the imbedding of normed rings into the ring of operators in Hilbert Space*, pages 3–19. 1943b. ISBN 0821851756. URL http://books.google.com/books?vid=ISBN0821851756.

John Robilliard Giles. *Introduction to the Analysis of Metric Spaces*. Number 3 in Australian Mathematical Society lecture series. Cambridge University Press, Cambridge, 1987. ISBN 978-0521359283. URL http://books.google.com/books?vid=ISBN0521359287.

Copyright © 2019 Daniel J. Greenhoe

John Robilliard Giles. *Introduction to the Analysis of Normed Linear Spaces*. Number 13 in Australian Mathematical Society lecture series. Cambridge University Press, Cambridge, 2000. ISBN 0-521-65375-4. URL http://books.google.com/books?vid=ISBN0521653754.

Israel Gohberg, Seymour Goldberg, and Marinus A. Kaashoek. *Basic Classes of Linear Operators*. Birkhäuser, Basel, 1 edition, 2003. ISBN 3764369302. URL http://books.google.com/books?vid=ISBN3764369302.

Steve Goldman. *Vibration Spectrum Analysis: A Practical Approach*. Industrial Press Inc., 1999. ISBN 9780831130886. URL http://books.google.com/books?vid=ISBN0831130881.

Jaideva C. Goswami and Andrew K. Chan. *Fundamentals of Wavelets; Theory, Algorithms, and Applications*. John Wiley & Sons, Inc., 1999. ISBN 0-471-19748-3. URL http://vadkudr.boom.ru/Collection/fundwave_contents.html.

H. G. D. Goyder. Foolproof methods for frequency response measurements. Technical Report ADP003677, United Kingdom Atomic Energy Authority London (England), 1984. URL https://apps.dtic.mil/dtic/tr/fulltext/u2/a143301.pdf#page=118.

I.S. Gradshteyn and I.M. Ryzhik. *Table of Integrals, Series, and Products*. 1980. ISBN 0-12-294760-6. URL http://www.worldcat.org/isbn/0122947606.

Alexander Graham. *Kronecker Products and Matrix Calculus: With Applications*. Ellis Horwood Series; Mathematics and its Applications. Ellis Horwood Limited, Chichester, 1981. ISBN 0-85312-391-8. URL http://books.google.com/books?vid=ISBN0853123918.

Alexander Graham. *Kronecker Products and Matrix Calculus: With Applications*. Ellis Horwood Series; Mathematics and its Applications. Ellis Horwood Limited, 2018. ISBN 9780486824178. URL http://books.google.com/books?vid=ISBN9780486824178.

R.J Greechie. Orthomodular lattices admitting no states. *Journal of Combinatorial Theory, Series A*, 10(2):119–132, March 1971. URL http://dx.doi.org/10.1016/0097-3165(71)90015-X.

Daniel J. Greenhoe. *Structure and Analysis of Mathematical Spaces*, volume 2 of *Mathematical Structure and Design series*. version 0.30 edition, January 2017a. URL https://www.researchgate.net/publication/312201065.

Daniel J. Greenhoe. The sequence a(n,m) of the m polynomial coefficients of the n-th order b-spline scaled by n!, read by rows, with n in 0,1,2,... and m in 1,2,3,...,(n+1)2. *Online Encyclopedia of Integer Sequences*, July 2017b. URL https://oeis.org/A289358.

Daniel J. Greenhoe. The effects of the assorted cross-correlation definitions. *ResearchGate.com*, June 2019. doi: 10.13140/RG.2.2.15881.70240. URL https://www.researchgate.net/publication/333882235.

Amritava Gupta. *Real & Abstract Analysis*. Academic Publishers Calcutta, 1998. ISBN 9788186358443. URL http://books.google.com/books?vid=ISBN8186358447. first published in 1998, reprinted in 2000 and 2006.

Alfréd Haar. Zur theorie der orthogonalen funktionensysteme. *Mathematische Annalen*, 69:331–371, September 1910. ISSN 1432-1807. doi: 10.1007/BF01456326.

Norman B. Haaser and Joseph A. Sullivan. *Real Analysis*. Dover Publications, New York, 1991. ISBN 0-486-66509-7. URL http://books.google.com/books?vid=ISBN0486665097.

Copyright © 2019 Daniel J. Greenhoe

Paul R. Haddad and Ali N. Akansu. *Multiresolution Signal Decomposition: Transforms, Subbands, and Wavelets*. Acedemic Press, October 1 1992. ISBN 0323138365. URL http://books.google.com/books?vid=ISBN0323138365.

Henry Sinclair Hall and Samuel Ratcliffe Knight. *Higher algebra, a sequel to elementary algebra for schools*. Macmillan, London, 1894. URL http://www.archive.org/details/higheralgebraas00kniggoog.

Paul R. Halmos. *Finite Dimensional Vector Spaces*. Princeton University Press, Princeton, 1 edition, 1948. ISBN 0691090955. URL http://books.google.com/books?vid=isbn0691090955.

Paul R. Halmos. *Finite Dimensional Vector Spaces*. Springer-Verlag, New York, 2 edition, 1958. ISBN 0-387-90093-4. URL http://books.google.com/books?vid=isbn0387900934.

Paul R. Halmos. *Intoduction to Hilbert Space and the Theory of Spectral Multiplicity*. Chelsea Publishing Company, New York, 2 edition, 1998. ISBN 0821813781. URL http://books.google.com/books?vid=ISBN0821813781.

Georg Hamel. Eine basis aller zahlen und die unstetigen lösungen der funktionalgleichung $f(x+y) = f(x) + f(y)$. *Mathematische Annalen*, 60(3):459–462, 1905. URL http://gdz.sub.uni-goettingen.de/dms/load/img/?PPN=GDZPPN002260395&IDDOC=28580.

Deguang Han, Keri Kornelson, David Larson, and Eric Weber. *Frames for Undergraduates*, volume 40 of *Student Mathematical Library*. American Mathematical Society, 2007. ISBN 0821842129. URL http://books.google.com/books?vid=ISBN0821842129. Deguang Han = ???

G.H. Hardy. Prolegomena to a chapter on inequalities. *Journal of the London Mathematical Society*, 1–4:61–78, November 8 1929. URL http://jlms.oxfordjournals.org/content/vols1-4/issue13/index.dtl#PRESIDENTIAL_ADDRESS. "Presidential Address" to the London Mathematical Society.

Godfrey H. Hardy. *A Mathematician's Apology*. Cambridge University Press, Cambridge, 1940. URL http://www.math.ualberta.ca/~mss/misc/A%20Mathematician's%20Apology.pdf.

Godfrey Harold Hardy, John Edensor Littlewood, and George Pólya. *Inequalities*. Cambridge Mathematical Library. Cambridge University Press, Cambridge, 2 edition, 1952. URL http://books.google.com/books?vid=ISBN0521358809.

Simon Haykin. *Communication Systems*. John Wiley & Sons, 3 edition, 1994. ISBN 0471571768. URL http://www.amazon.com/dp/0471571768.

Simon Haykin. *Adaptive Filter Theory*. Prentice Hall, Upper Saddle River, 4 edition, September 24 2001. ISBN 978-0130901262. URL http://books.google.com/books?vid=isbn0130901261.

Michiel Hazewinkel, editor. *Handbook of Algebras*, volume 2. North-Holland, Amsterdam, 1 edition, 2000. ISBN 044450396X. URL http://books.google.com/books?vid=ISBN044450396X.

Jean Van Heijenoort. *From Frege to Gödel : A Source Book in Mathematical Logic, 1879-1931*. Harvard University Press, Cambridge, Massachusetts, 1967. URL http://www.hup.harvard.edu/catalog/VANFGX.html.

Christopher Heil. *A Basis Theory Primer*. Applied and Numerical Harmonic Analysis. Birkhäuser, Boston, expanded edition edition, 2011. ISBN 9780817646868. URL http://books.google.com/books?vid=ISBN9780817646868.

Copyright © 2019 Daniel J. Greenhoe

J. H. Heinbockel. *Introduction to Finite and Infinite Series and Related Topics*. Trafford Publishing, 2010. ISBN 9781426949548. URL http://books.google.com/books?vid=ISBN9781426949545.

Carl W. Helstrom. *Probability and Stochastic Processes for Engineers*. Maxwell Macmillan international editions in engineering. Macmillan, 2 edition, 1991. ISBN 9780023535710. URL http://books.google.com/books?vid=ISBN0023535717.

Omar Hijab. *Introduction to Calculus and Classical Analysis*. Undergraduate Texts in Mathematics. Springer, 3 edition, 2011. ISBN 9781441994875. URL http://books.google.com/books?vid=ISBN1441994874.

David Hilbert. Grundzüge einer allgemeinen theorie der linearen integralgleichungen (fundamentals of a general theory of the linear integral equations). *Mathematisch-Physikalische Klasse (Mathematical-Physical Class)*, pages 49–91, March 1904. URL http://dz-srv1.sub.uni-goettingen.de/sub/digbib/loader?did=D57552. Report 1 of 6.

David Hilbert. Grundzüge einer allgemeinen theorie der linearen integralgleichungen (fundamentals of a general theory of the linear integral equations). *Mathematisch-Physikalische Klasse (Mathematical-Physical Class)*, pages 157–228, March 1906. URL http://dz-srv1.sub.uni-goettingen.de/sub/digbib/loader?did=D58133. Report 4 of 6.

David Hilbert. *Grundzüge einer allgemeinen Theorie der linearen Integralgleichungen (Fundamentals of a general theory of the linear integral equations)*, volume 26 of *Fortschritte der mathematischen Wissenschaften in Monographien (Progress of the mathematical sciences in Monographs)*. B.G. Teubner, Leipzig und Berlin, 1912. URL http://www.worldcat.org/oclc/13468199.

David Hilbert, Lothar Nordheim, and John von Neumann. über die grundlagen der quantenmechanik (on the bases of quantum mechanics). *Mathematische Annalen*, 98:1–30, 1927. ISSN 0025-5831 (print) 1432-1807 (online). URL http://dz-srv1.sub.uni-goettingen.de/cache/toc/D27776.html.

Otto Hölder. üeber einen mittelwerthssatz. *Göttingen Nachrichten*, pages 38–47, 1889. URL http://www.digizeitschriften.de/dms/img/?PPN=GDZPPN00252421X.

Klaus Höllig. *Finite Element Methods With B-Splines*, volume 26 of *Frontiers in Applied Mathematics*. SIAM, 2003. ISBN 0-89871-533-4. URL http://books.google.com/books?vid=ISBN0898715334.

Lars Hörmander. *Notions of Convexity*. Birkhäuser, Boston, 1994. ISBN 0817637990. URL http://books.google.com/books?vid=ISBN0817637990.

Roger A. Horn and Charles R. Johnson. *Matrix Analysis*. Cambridge University Press, Cambridge, 1990. ISBN 0-521-30586-1. URL http://books.google.com/books?vid=isbn0521305861. Library: QA188H66 1985.

Alfred Edward Housman. *More Poems*. Alfred A. Knopf, 1936. URL http://books.google.com/books?id=rTMiAAAAMAAJ.

Emmanuel C. Ifeachor and Barrie W. Jervis. *Digital Signal Processing: A Practical Approach*. Electronic systems engineering series. Prentice Hall, 1993. ISBN 020154413X. URL http://www.amazon.com/dp/020154413X.

Emmanuel C. Ifeachor and Barrie W. Jervis. *Digital Signal Processing: A Practical Approach*. Electronic systems engineering series. Prentice Hall, 2002. ISBN 9780201596199. URL http://www.amazon.com/dp/0201596199.

Copyright © 2019 Daniel J. Greenhoe

Satoru Igari. *Real Analysis—With an Introduction to Wavelet Theory*, volume 177 of *Translations of mathematical monographs*. American Mathematical Society, 1996. ISBN 9780821821046. URL http://books.google.com/books?vid=ISBN0821821040.

Vasile I. Istrăţescu. *Inner Product Structures: Theory and Applications*. Mathematics and Its Applications. D. Reidel Publishing Company, 1987. ISBN 9789027721822. URL http://books.google.com/books?vid=ISBN9027721823.

Robert C. James. Orthogonality in normed linear spaces. *Duke Mathematical Journal*, 12(2):291–302, 1945. ISSN 0012-7094. doi: 10.1215/S0012-7094-45-01223-3. URL http://projecteuclid.org/euclid.dmj/1077473105.

Robert C. James. Orthogonality and linear functionals in normed linear spaces. *Transactions of the American Mathematical Society*, 61(2):265–292, March 1947. ISSN 1088-6850. doi: 10.2307/1990220. URL http://www.jstor.org/stable/1990220.

Bjorn Jawerth and Wim Sweldens. An overview of wavelet based multiresolutional analysis. *SIAM Review*, 36:377–412, September 1994. URL http://cm.bell-labs.com/who/wim/papers/papers.html#overview.

Alan Jeffrey and Hui Hui Dai. *Handbook of Mathematical Formulas and Integrals*. Handbook of Mathematical Formulas and Integrals Series. Academic Press, 4 edition, January 18 2008. ISBN 9780080556840. URL http://books.google.com/books?vid=ISBN0080556841.

Gwilym M. Jenkins and Donald G. Watts. *Spectral analysis and its applications*. Holden-Day series in time series analysis. Holden-Day, 1968. URL https://archive.org/details/spectralanalysis00jenk_0.

J. L. W. V. Jensen. Sur les fonctions convexes et les ine'galite's entre les valeurs moyennes (on the convex functions and the inequalities between the average values). *Acta Mathematica*, 30(1):175–193, December 1906. ISSN 0001-5962. doi: 10.1007/BF02418571. URL http://www.springerlink.com/content/r55q1411g840j446/.

Leonard Benjamin William Jolley. *Summation of Series*. Dover, New York, second revised edition, 1961. URL http://plouffe.fr/simon/math/SummationofSeries.pdf.

P. Jordan and J. von Neumann. On inner products in linear, metric spaces. *Annals of Mathematics*, 36(3):719–723, July 1935. doi: 10.2307/1968653. URL http://www.jstor.org/stable/1968653.

Palle E. T. Jørgensen, Kathy Merrill, Judith Packer, and Lawrence W. Baggett. *Representations, Wavelets and Frames: A Celebration of the Mathematical Work of Lawrence Baggett*. Applied and Numerical Harmonic Analysis. Birkhäuser, 2008. ISBN 9780817646820. URL http://books.google.com/books?vid=ISBN0817646825.

K. D. Joshi. *Applied Discrete Structures*. New Age International, New Delhi, 1997. ISBN 8122408265. URL http://books.google.com/books?vid=ISBN8122408265.

J.S.Chitode. *Signals And Systems*. Technical Publications, 2009a. ISBN 9788184316780. URL http://books.google.com/books?vid=ISBN818431678X.

J.S.Chitode. *Digital Signal Processing*. Technical Publications, 2009b. ISBN 9788184316469. URL http://books.google.com/books?vid=ISBN9788184316469.

Gudrun Kalmbach. *Measures and Hilbert Lattices*. World Scientific, Singapore, 1986. ISBN 9789971500092. URL http://books.google.com/books?vid=ISBN9971500094.

Copyright © 2019 Daniel J. Greenhoe

Nidal S. Kamel and K. S. Sim. Image signal-to-noise ratio and noise variance estimation using autoregressive model. *The Journal of Scanning Microscopes*, 26:277–281, September 15 2004. URL https://onlinelibrary.wiley.com/doi/pdf/10.1002/sca.4950260605.

Steven M. Kay. *Modern Spectral Estimation: Theory and Application*. Prentice-Hall signal processing series. Prentice Hall, 1988. ISBN 9788131733561. URL http://books.google.com/books?vid=ISBN8131733564.

James P. Keener. *Principles of Applied Mathematics; Transformation and Approximation*. Addison-Wesley Publishing Company, Reading, Massachusets, 1988. ISBN 0-201-15674-1. URL http://www.worldcat.org/isbn/0201156741.

Yoshida Kenko. *The Tsuredzure Gusa of Yoshida No Kaneyoshi. Being the meditations of a recluse in the 14th Century (Essays in Idleness)*. circa 1330. URL http://www.humanistictexts.org/kenko.htm. 1911 translation of circa 1330 text.

John Francis Kenney. *Mathematics of Statistics, Volume 2*. D. Van Nostrand Company, New York, second edition, second printing edition, 1947.

Mohamed A. Khamsi and W.A. Kirk. *An Introduction to Metric Spaces and Fixed Point Theory*. John Wiley, New York, 2001. ISBN 978-0471418252. URL http://books.google.com/books?vid=isbn0471418250.

K.S. Kim and Nidal S. Kamel. Image signal-to-noise ratio and noise variance estimation using autoregressive model. *The Journal of Scanning Microscopes*, 26(6):135–139, September 24 2004. URL https://doi.org/10.1002/sca.4950260605.

Anthony W Knapp. *Advanced Real Analysis*. Cornerstones. Birkhäuser, Boston, Massachusetts, USA, 1 edition, July 29 2005a. ISBN 0817643826. URL http://books.google.com/books?vid=ISBN0817643826.

Anthony W Knapp. *Basic Real Analysis*. Cornerstones. Birkhäuser, Boston, Massachusetts, USA, 1 edition, July 29 2005b. ISBN 0817632506. URL http://books.google.com/books?vid=ISBN0817632506.

Andrei N. Kolmogorov. *Grundbegriffe der Wahrscheinlichkeitsrechnung*. Springer, Berlin, 1933a.

Andrei Nikolaevich Kolmogorov. *Foundations of the theory of probability*. Chelsea Publishing Company, New Yourk, 2 edition, 1933b. ISBN B0006AUOGO. URL http://statweb.stanford.edu/~cgates/PERSI/Courses/Phil166-266/Kolmogorov-Foundations.pdf. 1956 2nd edition English translation of A. N. Kolmogorov's 1933 “Grundbegriffe der Wahrscheinlichkeitsrechnung”.

Lambert Herman Koopmans. volume 22 of *Probability and mathematical statistics*. Academic Press, 2 edition, 1995. ISBN 9780124192515.

Carlos S. Kubrusly. *The Elements of Operator Theory*. Springer, 1 edition, 2001. ISBN 9780817641740. URL http://books.google.com/books?vid=ISBN0817641742.

Carlos S. Kubrusly. *The Elements of Operator Theory*. Springer, 2 edition, 2011. ISBN 9780817649975. URL http://books.google.com/books?vid=ISBN0817649972.

Andrew Kurdila and Michael Zabarankin. *Convex Functional Analysis*. Systems & Control: Foundations & Applications. Birkhäuser, Boston, 2005. ISBN 9783764321987. URL http://books.google.com/books?vid=ISBN3764321989.

Copyright © 2019 Daniel J. Greenhoe

Peter D. Lax. *Functional Analysis*. John Wiley & Sons Inc., USA, 2002. ISBN 0-471-55604-1. URL http://www.worldcat.org/isbn/0471556041. QA320.L345 2002.

Steven R. Lay. *Convex Sets and their Applications*. John Wiley & Sons, New York, 1982. ISBN 0-471-09584-2. URL http://books.google.com/books?vid=isbn0471095842.

Q. Leclere, Bert Roozen, and Celine Sandier. On the use of the hs estimator for the experimental assessment of transmissibility matrices. *Mechanical Systems and Signal Processing*, 43(1-2):237–245, 2014. URL https://hal.archives-ouvertes.fr/hal-00911916/document. <hal-00911916>.

Gottfried Wilhelm Leibniz. Letter to christian huygens, 1679. In Leroy E. Loemker, editor, *Philosophical Papers and Letters*, volume 2 of *The New Synthese Historical Library*, chapter 27, pages 248–249. Kluwer Academic Press, Dordrecht, 2 edition, September 8 1679. ISBN 902770693X. URL http://books.google.com/books?vid=ISBN902770693X.

J. Leuridan, J. De Vis, and Van der Auweraer H. F. Lembregts D. "a comparison of some frequency response function measurement techniques. pages 908–918, 1986. URL https://www.researchgate.net/publication/283994172.

Zach Liang and George C. Lee. *Random Vibration: Mechanical, Structural, and Earthquake Engineering Applications*. Advances in Earthquake Engineering. Taylor & Francis, illustrated edition, 2015. ISBN 9781498702348. URL http://www.amazon.com/dp/1498702341.

Elliott H. Lieb and Michael Loss. *Analysis*, volume 14 of *Graduate studies in mathematics*. American Mathematical Soceity, Providence, Rhode Island, USA, 2 edition, 2001. ISBN 0821827839. URL http://books.google.com/books?vid=ISBN0821827839.

Joram Lindenstrauss and Lior Tzafriri. *Classical Banach Spaces I: Sequence Spaces*, volume 92 of *Ergebnisse der Mathematik und ihrer Grenzgebiete*. Springer-Verlag, 1977. ISBN 978-3642665592. URL http://www.amazon.com/dp/3642665594.

Anders Lindquist and Giorgio Picci. *Linear Stochastic Systems: A Geometric Approach to Modeling, Estimation and Identification*, volume 1 of *Series in Contemporary Mathematics*. Springer, 2015. ISBN 9783662457504. URL http://books.google.com/books?vid=ISBN3662457504.

Lynn H. Loomis. *An Introduction to Abstract Harmonic Analysis*. The University Series in Higher Mathematics. D. Van Nostrand Company, Toronto, 1953. URL http://books.google.com/books?id=aNg-AAAAIAAJ.

Lynn H. Loomis and Ethan D. Bolker. *Harmonic analysis*. Mathematical Association of America, 1965. URL http://books.google.com/books?id=MEfvAAAAMAAJ.

Niccolò Machiavelli. *The Literary Works of Machiavelli: Mandragola, Clizia, A Dialogue on Language, and Belfagor, with Selections from the Private Correspondence*. Oxford University Press, 1961. ISBN 0313212481. URL http://www.worldcat.org/isbn/0313212481.

Fumitomo Maeda and Shûichirô Maeda. *Theory of Symmetric lattices*, volume 173 of *Die Grundlehren der mathematischen Wissenschaften in Einzeldarstellungen*. Springer-Verlag, Berlin/New York, 1970. URL http://books.google.com/books?id=4oeBAAAAIAAJ.

Khosrow Maleknejad, Monireh Nosrati Sahlan, and Azadeh Ostad. Numerical solution of fractional integro-differential equation by using cubic b-spline wavelets. In *Proceedings of the World Congress on Engineering*, volume I, London, July 3–5 2013. URL https://pdfs.semanticscholar.org/bbe0/55f40768bbefe9e6476bdce701effb9243f7.pdf.

Copyright © 2019 Daniel J. Greenhoe

Lech Maligranda. A simple proof of the hölder and the minkowski inequality. *The American Mathematical Monthly*, 102(3):256–259, March 1995. URL http://www.jstor.org/stable/2975013.

Stéphane G. Mallat. Multiresolution approximations and wavelet orthonormal bases of $l^2(\mathbb{R})$. *Transactions of the American Mathematical Society*, 315(1):69–87, September 1989. URL http://blanche.polytechnique.fr/~mallat/papiers/math_multiresolution.pdf.

Stéphane G. Mallat. *A Wavelet Tour of Signal Processing*. Elsevier, 2 edition, September 15 1999. ISBN 0-12-466606-X. URL http://books.google.com/books?vid=ISBN012466606X.

R. Viswanathan Mandyam D. Srinath, P.K. Rajasekaran. *Introduction to Statistical Signal Processing with Applications*. Prentice Hall Inc, Upper Saddle River, 1996. ISBN 013125295X. URL http://engr.smu.edu/ee/mds/.

Dimitris G. Manolakis, Vinay K. Ingle, and Stephen M. Kogon. *Statistical and Adaptive Signal Processing*. Mc-Graw Hill, Boston, 2000. ISBN 0-07-040051-2.

Dimitris G. Manolakis, Vinay K. Ingle, and Stephen M. Kogon. *Statistical and Adaptive Signal Processing: Spectral Estimation, Signal Modeling, Adaptive Filtering, and Array Processing*. Artech House signal processing library. Artech House, illustrated, reprint edition, 2005. ISBN 9781580536103. URL http://www.amazon.com/dp/1580536107.

MatLab. cpsd: Cross power spectral density. *MatLab*, accessed 2018 November 26 2018a. URL https://www.mathworks.com/help/signal/ref/cpsd.html.

MatLab. xcorr: Cross-correlation. *MatLab*, accessed 2018 November 26 2018b. URL https://www.mathworks.com/help/signal/ref/xcorr.html.

Stefan Mazur. Sur les anneaux linéaires. *Comptes rendus de l'Académie des sciences*, 207:1025–1027, 1938.

Stefan Mazur and Stanislaus M. Ulam. Sur les transformations isométriques d'espaces vectoriels normées. *Comptes rendus de l'Académie des sciences*, 194:946–948, 1932.

Karl Menger. Untersuchungen über allgemeine metrik. *Mathematische Annalen*, 100:75–163, 1928. ISSN 0025-5831. URL http://link.springer.com/article/10.1007/BF01455705. (Investigations on general metric).

Claudia Menini and Freddy Van Oystaeyen. *Abstract Algebra; A Comprehensive Treatment*. Marcel Dekker Inc, New York, April 2004. ISBN 0-8247-0985-3. URL http://books.google.com/books?vid=isbn0824709853.

James Mercer. Functions of positive and negative type and their connection with the theory of integral equations. *Philosophical Transactions of the Royal Society of London*, 209:415–446, 1909. ISSN 02643952. URL http://www.jstor.org/stable/91043.

Anthony N. Michel and Charles J. Herget. *Applied Algebra and Functional Analysis*. Dover Publications, Inc., 1993. ISBN 0-486-67598-X. URL http://books.google.com/books?vid=ISBN048667598X. original version published by Prentice-Hall in 1981.

Gregory K. Miller. *Probability: Modeling and Applications to Random Processes*. Wiley, August 25 2006. URL http://www.amazon.com/dp/0471458929.

Gradimir V. Milovanović and Igorž. Milovanović. On a generalization of certain results of a. ostrowski and a. lupaş. *Publikacije Elektrotehničkog Fakulteta (Publications Electrical Engineering)*, (643):62–69, 1979. URL http://www.mi.sanu.ac.rs/~gvm/radovi/643.pdf.

Copyright © 2019 Daniel J. Greenhoe

Hermann Minkowski. *Geometrie der Zahlen*. Druck und Verlag von B.G. Teubner, Leipzig, 1910. URL http://www.archive.org/details/geometriederzahl00minkrich. Geometry of Numbers.

Fred Mintzer. Filters for distortion-free two-band multi-rate filter banks. *IEEE Transactions on Acoustics, Speech and Signal Processing*, 32, 1985.

Larry D. Mitchell. An improved method for the fast fourier transform (fft) calculation of the frequency response function. In M. Kunt and F. De Coulon, editors, *Proceedings of EUSIPCO-80, First European Signal Processing Conference*, Signal processing, theories and applications, pages 159–160, Lausanne, Switzerland, September 16–19 1980. Elsevier. URL http://www.worldcat.org/oclc/63416846. citation from Cobb(1988).

Larry D. Mitchell. Improved methods for the fast fourier transform (fft) calculation of the frequency response function. *Journal of Mechanical Design*, 104(2):277–279, April 01 1982. doi: 10.1115/1.3256336. URL http://mechanicaldesign.asmedigitalcollection.asme.org/article.aspx?articleid=1451670.

Dragoslav S. Mitrinović, J. E. Pečarić, and Arlington M. Fink. *Classical and New Inequalities in Analysis*, volume 61 of *Mathematics and its Applications (East European Series)*. Kluwer Academic Publishers, Dordrecht, Boston, London, 2010. ISBN 978-90-481-4225-5. URL http://www.amazon.com/dp/0792320646.

Todd K. Moon and Wynn C. Stirling. *Mathematical Methods and Algorithms for Signal Processing*. Prentice Hall, Upper Saddle River, 2000. ISBN 0-201-36186-8. URL http://books.google.com/books?vid=isbn0201361868.

H. P. Mulholland. On generalizations of minkowski's inequality in the form of a triangle inequality. *Proceedings of the London Mathematical Society*, s2-51:294–307, 1950. URL http://plms.oxfordjournals.org/content/s2-51/1/294.extract. received 1946 October 10, read 1947 June 19.

Paul J. Nahin. *Dr. Euler's Fabulous Formula: Cures Many Mathematical Ills*. Princeton Science Library. Princeton University Press, reprint, revised edition, April 25 2011. ISBN 9781400838479. URL http://books.google.com/books?vid=ISBN1400838479.

Louis Narens. *Theories of Probability: An examination of logical and qualitative foundations*, volume 2 of *Advanced series on mathematical psychology*. World Scientific, January 2007. ISBN 9789812708014. URL http://www.amazon.com/dp/9812708014.

Louis Narens. Alternative probability theories for cognitive psychology. *Topics in Cognitive Science*, 6(1):114–120, January 2014. ISSN 1756-8757. doi: 10.1111/tops.12071. URL http://onlinelibrary.wiley.com/doi/10.1111/tops.12071/full.

Oliver Nelles. *Nonlinear System Identification*. Springer, New York, 2001. ISBN 9783540673699.

Ben Noble and James W. Daniel. *Applied Linear Algebra*. Prentice-Hall, Englewood Cliffs, NJ, USA, 3 edition, 1988. ISBN 0-13-041260-0. URL http://www.worldcat.org/isbn/0130412600. Library QA184.N6 1988 512.5 87-11511.

Timur Oikhberg and Haskell Rosenthal. A metric characterization of normed linear spaces. *Rocky Mountain Journal Of Mathematics*, 37(2):597–608, 2007. URL http://www.ma.utexas.edu/users/rosenthl/pdf-papers/95-oikh.pdf.

Oystein Ore. On the foundation of abstract algebra. i. *The Annals of Mathematics*, 36(2):406–437, April 1935. URL http://www.jstor.org/stable/1968580.

BY-NC-ND
Copyright © 2019 Daniel J. Greenhoe

Brad Osgood. The fourier transform and its applications; ee261: Standford university course. university course notes, 2002. URL http://www.stanford.edu/class/ee261/.

Lincoln P. Paine. *Warships of the World to 1900*. Ships of the World Series. Houghton Mifflin Harcourt, 2000. ISBN 9780395984147. URL http://books.google.com/books?vid=ISBN9780395984149.

Endre Pap. *Null-Additive Set Functions*, volume 337 of *Mathematics and Its Applications*. Kluwer Academic Publishers, 1995. ISBN 0792336585. URL http://www.amazon.com/dp/0792336585.

Anthanasios Papoulis. *Probability, Random Variables, and Stochastic Processes*. McGraw-Hill Series in Electrical Engineering. McGraw-Hill Book Company, New York, 2 edition, 1984. ISBN 0070484686. URL http://books.google.com/books?vid=ISBN0070484686.

Anthanasios Papoulis. *Probability, Random Variables, and Stochastic Processes*. McGraw-Hill, New York, 3 edition, 1991. ISBN 0070484775. URL http://books.google.com/books?vid=ISBN0070484775.

Anthanasios Papoulis. *Probability, Random Variables, and Stochastic Processes*. McGraw-Hill series in electrical and computer engineering. Tata McGraw-Hill, New York, 4 edition, 2002. ISBN 9780071226615. URL http://books.google.com/books?vid=ISBN0071226613.

Athanasios Papoulis. *Probability & Statistics*. Prentice-Hall international editions. Prentice Hall, illustrated edition, 1990. ISBN 9780137116980. URL http://books.google.com/books?vid=ISBN0137116985.

Giuseppe Peano. *Calcolo geometrico secondo l'Ausdehnungslehre di H. Grassmann preceduto dalle operazioni della logica deduttiva*. Fratelli Bocca Editori, Torino, 1888a. Geometric Calculus: According to the *Ausdehnungslehre* of H. Grassmann.

Giuseppe Peano. *Geometric Calculus: According to the Ausdehnungslehre of H. Grassmann*. Springer (2000), 1888b. ISBN 0817641262. URL http://books.google.com/books?vid=isbn0817641262. originally published in 1888 in Italian.

Michael Pedersen. *Functional Analysis in Applied Mathematics and Engineering*. Chapman & Hall/CRC, New York, 2000. ISBN 9780849371691. URL http://books.google.com/books?vid=ISBN0849371694. Library QA320.P394 1999.

J. E. Pečarić, Frank Proschan, and Yung Liang Tong. *Convex Functions, Partial Orderings, and Statistical Applications*, volume 187 of *Mathematics in Science and Engineering*. Academic Press, San Diego, California, 1992. ISBN 978-0125492508. URL http://books.google.com/books?vid=ISBN0125492502.

Rik Pintelon and Johan Schoukens. *System Identification: A Frequency Domain Approach*. John Wiley & Sons, 2 edition, 2012. ISBN 9781118287392. URL http://books.google.com/books?vid=ISBN1118287398.

R. L. Plackett. Studies in the history of probability and statistics. xxix: The discovery of the method of least squares. *Biometrika*, 59(2):239–251, August 1972. doi: 10.2307/2334569. URL https://www.jstor.org/stable/2334569. https://doi.org/10.1093/biomet/59.2.239.

Henri Poincaré. *Calcul Des Probabilités*. De L'ecole Polytechnique, Paris, 1912. URL http://www.archive.org/details/calculdeprobabil00poinrich.

Karl Raimund Popper. *Conjectures and refutations: the growth of scientific knowledge*. Basic Books, 1962. URL https://books.google.com/books?id=OBgNAQAAIAAJ.

Copyright © 2019 Daniel J. Greenhoe

Karl Raimund Popper. *Conjectures and refutations: the growth of scientific knowledge*. Routledge, 2 edition, 1963. ISBN 9781135971304.

Lakshman Prasad and Sundararaja S. Iyengar. *Wavelet Analysis with Applications to Image Processing*. CRC Press LLC, Boca Raton, 1997. ISBN 978-0849331695. URL http://books.google.com/books?vid=ISBN0849331692. Library TA1637.P7 1997.

John G. Proakis. *Digital Communications*. McGraw Hill, 4 edition, 2001. ISBN 0-07-232111-3. URL http://www.mhhe.com/.

John G. Proakis, editor. *Algorithms for Statistical Signal Processing*. Prentice Hall, 2002. ISBN 9780130622198. URL http://books.google.com/books?vid=ISBN9780130622198.

John G. Proakis and Dimitris G. Manolakis. *Digital Signal Processing: Principles, Algorithms, and Applications*. Prentice Hall international editions. Prentice Hall, 3 edition, 1996. ISBN 9780133737622. URL http://books.google.com/books?vid=ISBN0133737624.

Inder K. Rana. *An Introduction to Measure and Integration*, volume 45 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, R.I., 2 edition, 2002. ISBN 978-0821829745. URL http://books.google.com/books?vid=ISBN0821829742.

K. Deergha Rao and M. N. S. Swamy. *Digital Signal Processing: Theory and Practice*. Springer, 2018. ISBN 9789811080814. URL http://books.google.com/books?vid=ISBN9789811080814.

Charles Earl Rickart. *General Theory of Banach Algebras*. University series in higher mathematics. D. Van Nostrand Company, Yale University, 1960. URL http://books.google.com/books?id=PVrvAAAAMAAJ.

Frigyes Riesz. Zur theorie des hilbertschen raumes. *Acta Scientiarum Mathematicarum*, 7:34–38, 1934. URL http://www.math.u-szeged.hu/acta/. The Theory of Hilbert Space.

Enders A. Robinson. *Random Wavelets and Cybernetic Systems*, volume 9 of *Griffins Statistical Monographs & Courses*. Lubrecht & Cramer Limited, London, June 1962. ISBN 0852640757. URL http://books.google.com/books?vid=ISBN0852640757.

Enders A. Robinson. Multichannel z-transforms and minimum delay. *Geophyics*, 31(3):482–500, June 1966. doi: 10.1190/1.1439788. URL http://dx.doi.org/10.1190/1.1439788.

R. Tyrrell Rockafellar. *Convex Analysis*. Princeton landmarks in mathematics. Princeton University Press, 1970. ISBN 9780691015866. URL http://books.google.com/books?vid=ISBN0691015864.

Murray Rosenblatt. Remarks on some non-parametric estimates of a density function. *Annals of Mathematical Statistics*, 27(3):832–837, September 1956. URL https://link.springer.com/content/pdf/10.1007/978-1-4419-8339-8_13.pdf. https://projecteuclid.org/download/pdf_1/euclid.aoms/1177728190.

Sheldon Ross. *A First Course in Probability*. Prentice Hall, Upper Saddle River, New Jersey, USA 07458, 5 edition, 1998. ISBN 0-13-746314-6. URL http://www.amazon.com/dp/0137463146.

George G. Roussas. *An Introduction to Probability and Statistical Inference*. Academic Press, 2 edition, October 21 2014. ISBN 9780128004371. URL http://books.google.com/books?vid=ISBN0128004371.

Walter Rudin. *Principles of Mathematical Analysis*. McGraw-Hill, New York, 3 edition, 1976. ISBN 007054235X. URL http://books.google.com/books?vid=ISBN007054235X. Library QA300.R8 1976.

Copyright © 2019 Daniel J. Greenhoe

Walter Rudin. *Functional Analysis*. McGraw-Hill, New York, 2 edition, 1991. ISBN 0-07-118845-2. URL http://www.worldcat.org/isbn/0070542252. Library QA320.R83 1991.

Bryan P. Rynne and Martin A. Youngson. *Linear Functional Analysis*. Springer undergraduate mathematics series. Springer, 2 edition, January 1 2008. ISBN 9781848000056. URL http://books.google.com/books?vid=ISBN1848000057.

Jr. S. Lawrence Marple. *Digital Spectral Analysis with Applications*. Prentice Hall, 1 edition, 1987. ISBN 9780132141499. URL http://books.google.com/books?vid=ISBN9780132141499.

Jr. S. Lawrence Marple. *Digital Spectral Analysis*. Dover Books on Electrical Engineering. Courier Dover Publications, 2019. ISBN 9780486780528. URL http://books.google.com/books?vid=ISBN9780486780528.

Shôichirô Sakai. *C*-Algebras and W*-Algebras*. Springer-Verlag, Berlin, 1 edition, 1998. ISBN 9783540636335. URL http://books.google.com/books?vid=ISBN3540636331. reprint of 1971 edition.

S. Salivahanan and A. Vallavaraj. *Digital Signal Processing*. Tata McGraw-Hill Education, 2001. ISBN 9780074639962. URL http://books.google.com/books?vid=ISBN9780074639962.

Louis L. Scharf. *Statistical Signal Processing*. Addison-Wesley Publishing Company, Reading, MA, 1991. ISBN 0-201-19038-9.

H. A. Scharz. über ein die flächen kleinsten flächeninhalts betreffendes problem der variationsrechnung. *Acta Soc. Scient. Fen.*, 15:315–362, 1885. URL http://www-stat.wharton.upenn.edu/~steele/Publications/Books/CSMC/CSMC_index.html.

Juljusz Schauder. Zur theorie stetiger abbildungen in funktionalräumen. *Mathematische Zeitschrift*, 26:47–65, 1927. URL http://eudml.org/doc/167911;jsessionid=156A34EBAB6C0E2DDAAC8C1232D23E8F.

Juljusz Schauder. Eine eigenschaft des haarschen orthogonalsystems. *Mathematische Zeitschrift*, 28:317–320, 1928.

Larry L. Schumaker. *Spline Functions: Basic Theory*. Cambridge Mathematical Library. Cambridge University Press, 3 edition, August 2007. ISBN 978-0-521-70512-7. URL http://books.google.com/books?vid=ISBN0521705126.

Isaac Schur. Uber die charakterischen wurzeln einer linearen substitution mit enier anwendung auf die theorie der integralgleichungen (over the characteristic roots of one linear substitution with an application to the theory of the integral). *Mathematische Annalen*, 66:488–510, 1909. URL http://dz-srv1.sub.uni-goettingen.de/cache/toc/D38231.html.

Karl Hermann Amandus Schwarz. *Ueber Ein Die Flachen Kleinsten Flächeninhalts; Betreffendes Problem Der Variationsrechnung (Over flattening smallest flat contents;*. October 31 1885. URL http://www-stat.wharton.upenn.edu/~steele/Publications/Books/CSMC/Schwarz.pdf. Festschrift Zum Jubelgeburtstage des Herrn Karl Weierstrass (anniversary publication to comemerate the birthdays of Mr. Karl Weierstrass).

Mícheál Ó Searcóid. *Elements of Abstract Analysis*. Springer Undergraduate Mathematics Series. Springer, 2002. ISBN 9781852334246. URL http://books.google.com/books?vid=ISBN185233424X.

Jun Shao. *Mathematical Statistics*. Springer Texts in Statistics. Springer Science & Business Media, 2003. ISBN 9780387953823. URL http://books.google.com/books?vid=ISBN0387953825.

Copyright © 2019 Daniel J. Greenhoe

Kihong Shin and Joseph Hammond. *Fundamentals of Signal Processing for Sound and Vibration Engineers*. John Wiley & Sons, 2008. ISBN 9780470725641.

Bernard Shizgal. *Spectral Methods in Chemistry and Physics: Applications to Kinetic Theory and Quantum Mechanics*. Scientific Computation. Springer, January 7 2015. ISBN 9789401794541. URL http://books.google.com/books?vid=ISBN9401794545.

Bernard. W. Silverman. *Density Estimation for Statistics and Data Analysis*, volume 26 of *Monographs on Statistics & Applied Probability*. Chapman & Hall/CRC, illustrated, reprint edition, 1986. ISBN 9780412246203. URL http://books.google.com/books?vid=ISBN9780412246203.

George Finlay Simmons. *Calculus Gems: Brief Lives and Memorable Mathematicians*. Mathematical Association of America, Washington DC, 2007. ISBN 0883855615. URL http://books.google.com/books?vid=ISBN0883855615.

Barry Simon. *Convexity: An Analytic Viewpoint*, volume 187 of *Cambridge Tracts in Mathematics*. Cambridge University Press, May 19 2011. ISBN 9781107007314. URL http://books.google.com/books?vid=ISBN1107007313.

Iván Singer. *Bases in Banach Spaces I*, volume 154 of *Grundlehren der mathematischen Wissenschaften in Einzeldarstellungen mit besonderer Berücksichtigung der Anwendungsgebiete*. Springer-Verlag New York, New York, 1970. ISBN 9780387048338.

Herbert Ellsworth Slaught and Nels Johann Lennes. *Elementary Algebra*. Allyn and Bacon, 1915. URL https://archive.org/details/elementaryalgebr00slaurich/.

M.J.T. Smith and T.P. Barnwell. A procedure for designing exact reconstruction filter banks for tree-structured subband coders. *IEEE International Conference on Acoustics, Speech and Signal Processing*, 9:421–424, 1984a. T.P. Barnwell is T.P. Barnwell III.

M.J.T. Smith and T.P. Barnwell. The design of digital filters for exact reconstruction in subband coding. *IEEE Transactions on Acoustics, Speech and Signal Processing*, 34(3):434–441, June 1984b. ISSN 0096-3518. doi: 10.1109/TASSP.1986.1164832. T.P. Barnwell is T.P. Barnwell III.

J. Laurie Snell. A conversation with joe doob. *Statistical Science*, 12(4):301–311, 1997. URL http://www.jstor.org/stable/2246220. similar article on web at http://www.dartmouth.edu/~chance/Doob/conversation.html.

J. Laurie Snell. Obituary: Joseph leonard doob. *Journal of Applied Probability*, 42(1):247–256, 2005. URL http://projecteuclid.org/euclid.jap/1110381384.

Houshang H. Sohrab. *Basic Real Analysis*. Birkhäuser, Boston, 1 edition, 2003. ISBN 978-0817642112. URL http://books.google.com/books?vid=ISBN0817642110.

H. W. Sorenson. Least-squares estimation: from gauss to kalman. *IEEE Spectrum*, pages 63–68, July 1970. URL https://www2.pv.infn.it/~fontana/download/lect/Sorenson.pdf.

J. Michael Steele. *The Cauchy-Schwarz Master Class: An Introduction to the Art of Mathematical Inequalities*. Cambridge University Press, Cambridge, April 26 2004. ISBN 052154677X. URL http://www-stat.wharton.upenn.edu/~steele/Publications/Books/CSMC/CSMC_index.htm.

Lynn Arthur Steen. Highlights in the history of spectral theory. *The American Mathematical Monthly*, 80(4):359–381, April 1973. ISSN 00029890. URL http://www.jstor.org/stable/2319079.

BY-NC-ND
Copyright © 2019 Daniel J. Greenhoe

Stephen M. Stigler. Gauss and the invention of least squares. *The Annals of Statistics*, 9(3):465–474, May 1981. URL https://projecteuclid.org/euclid.aos/1176345451. https://www.jstor.org/stable/2240811.

Josef Stoer and R. Bulirsch. *Introduction to Numerical Analysis*, volume 12 of *Texts in Applied Mathematics*. Springer Science & Business Media, 3 edition, August 21 2002. ISBN 9780387954523. URL http://books.google.com/books?vid=ISBN038795452X.

Marshall Harvey Stone. *Linear transformations in Hilbert space and their applications to analysis*, volume 15 of *American Mathematical Society. Colloquium publications*. American Mathematical Society, New York, 1932. URL http://books.google.com/books?vid=ISBN0821810154. 1990 reprint of the original 1932 edition.

Gilbert Strang. Wavelets and dilation equations: A brief introduction. *SIAM Review*, 31(4): 614–627, December 1989. ISSN 0036-1445. URL https://pdfs.semanticscholar.org/d118/a25a91488b4b37457cdc506472819de90523.pdf.

Gilbert Strang and Truong Nguyen. *Wavelets and Filter Banks*. Wellesley-Cambridge Press, Wellesley, MA, 1996. ISBN 9780961408879. URL http://books.google.com/books?vid=ISBN0961408871.

Alan Stuart and J. Keith Ord. *Kendall's Advanced Theory of Statistics Volume 2 Classical Inference and Relationship*. Hodder & Stoughton, 5 edition, 1991. ISBN 9780340560235. URL http://books.google.com/books?vid=ISBN9780340560235.

Yuri M. Suhov, Michael Kelbert, and Mark Kelbert. *Probability and Statistics by Example: Volume 1, Basic Probability and Statistics*, volume 1 of *Probability and Statistics by Example*. Cambridge University Press, illustrated, reprint edition, October 13 2005. ISBN 9780521847667.

Wim Sweldens and Robert Piessens. Wavelet sampling techniques. In *1993 Proceedings of the Statistical Computing Section*, pages 20–29. American Statistical Association, August 1993. URL http://citeseer.ist.psu.edu/18531.html.

Tamás Szabó and Gábor Horváth. An efficient hardware implementation of feed-forward neural networks. *Applied Intelligence*, 1:143–158, 2004. URL https://www.researchgate.net/profile/Gabor_Horvath/publication/263107833.

Brian S. Thomson, Andrew M. Bruckner, and Judith B. Bruckner. *Elementary Real Analysis*. www.classicalrealanalysis.com, 2 edition, 2008. ISBN 9781434843678. URL http://classicalrealanalysis.info/com/Elementary-Real-Analysis.php.

J. T. L. Thong, K. S. Sim, and J. C. H. Phang. Single-image signal-to-noise ratio estimation. *The Journal of Scanning Microscopes*, 23:328–336, December 07 2001. URL https://doi.org/10.1002/sca.4950230506.

Elmer Tolsted. An elementary derivation of cauchy, hölder, and minkowski inequalities from young's inequality. *Mathematics Magazine*, 37:2–12, 1964. URL http://mathdl.maa.org/mathDL/22/?pa=content&sa=viewDocument&nodeId=3036.

Harry L. Van Trees. *Detection, Estimation, and Modulation Theory, Part I*. Wiley-Interscience, reprint edition edition, September 27 2001. ISBN 0471095176. URL http://ece.gmu.edu/faculty_info/van.html.

Harry L. Van Trees. *Otimum Array Processing; Part IV of Detection, Estimation, and Modulation Theory*. Wiley-Interscience, New York, 2002. ISBN 0-471-09390-4. URL http://ece.gmu.edu/faculty_info/van.html.

Copyright © 2019 Daniel J. Greenhoe

Kishor S. Trivedi. *Probability and Statistics with Reliability, Queuing, and Computer Science Applications*. John Wiley & Sons, 2 edition, 2016. ISBN 9781119314202.

Howard G. Tucker. *An Introduction to Probability and Mathematical Statistics*. Academic press textbooks in mathematics. Academic Press, 1 edition, 1965.

Constantin Udriste. *Convex Functions and Optimization Methods on Riemannian Manifolds*, volume 297 of *Mathematics and Its Applications*. Springer, July 31 1994. ISBN 0792330021. URL http://books.google.com/books?vid=ISBN0792330021.

Stanislaw Marcin Ulam. *Adventures of a Mathematician*. University of California Press, Berkeley, 1991. ISBN 0520071549. URL http://books.google.com/books?vid=ISBN0520071549.

P.P. Vaidyanathan. *Multirate Systems and Filter Banks*. Prentice Hall Signal Processing Series. Prentice Hall, 1993. ISBN 0136057187. URL http://books.google.com/books?vid=ISBN0136057187.

Jussi Väisälä. A proof of the mazur-ulam theorem. *The American Mathematical Monthly*, 110(7): 633–635, August–September 2003. URL http://www.helsinki.fi/~jvaisala/mazurulam.pdf.

M.L.J. van de Vel. *Theory of Convex Structures*, volume 50 of *North-Holland Mathematical Library*. North-Holland, Amsterdam, 1993. ISBN 978-0444815057. URL http://books.google.com/books?vid=ISBN0444815058.

Saeed V. Vaseghi. *Advanced Digital Signal Processing and Noise Reduction*. John Wiley & Sons, 2 edition, 2000. ISBN 0471626929. URL http://books.google.com/books?vid=ISBN0471626929.

Brani Vidakovic. *Statistical Modeling by Wavelets*. John Wiley & Sons, Inc, New York, 1999. ISBN 0-471-29365-2. URL http://www.amazon.com/dp/0471293652.

Wolfgang von der Linden, Volker Dose, and Udo Von Toussaint. *Bayesian Probability Theory: Applications in the Physical Sciences*. Cambridge University Press, 2014. ISBN 9781107035904. URL http://books.google.com/books?vid=ISBN1107035902.

John von Neumann. Allgemeine eigenwerttheorie hermitescher funktionaloperatoren. *Mathematische Annalen*, 102(1):49–131, 1929. ISSN 0025-5831 (print) 1432-1807 (online). URL http://resolver.sub.uni-goettingen.de/purl?GDZPPN002273535. General eigenvalue theory of Hermitian functional operators.

Gilbert G. Walter and XiaoPing Shen. *Wavelets and Other Orthogonal Systems*. Chapman and Hall/CRC, New York, 2 edition, 2001. ISBN 1-58488-227-1. URL http://books.google.com/books?vid=ISBN1584882271. Library QA403.3.W34 2000.

Heinrich Martin Weber. Die allgemeinen grundlagen der galois'schen gleichungstheorie. *Mathematische Annalen*, 43(4):521–549, December 1893. URL http://resolver.sub.uni-goettingen.de/purl?GDZPPN002254670. The general foundation of Galois' equation theory.

H. L. Wei and S. A. Billings. An efficient nonlinear cardinal b-spline model for high tide forecasts at the venice lagoon. *Nonlinear Processes in Geophysics*, 13:577–584, 2006. URL https://www.nonlin-processes-geophys.net/13/577/2006/.

David Wells. *The Penguin Dictionary of Curious and Interesting Numbers*. revised edition, April 7 1987. ISBN 9780140080292. URL http://www.amazon.com/dp/0140080295.

Hermann Weyl. The method of orthogonal projection in potential theory. *Duke Mathematical Journal*, 7(1):411–444, 1940. URL http://projecteuclid.org/euclid.dmj/1077492266.

BY-NC-ND | Copyright © 2019 Daniel J. Greenhoe

P. R. White, M. H. Tan, and J. K. Hammond. Analysis of the maximum likelihood, total least squares and principal component approaches for frequency response function estimation. *Journal of Sound and Vibration*, 290(3–5):676–689, March 2006. URL https://doi.org/10.1016/j.jsv.2005.04.029.

Alfred L. Wicks and Harvard Vold. The h_s frequency response function estimator. In *Proceedings of the 4th International Modal Analysis Conference*, pages 897–899, Los Angeles Airport Marriott Hotel, California, USA, February 3–6 1986. URL https://www.tib.eu/en/search/id/TIBKAT%3A246489537/Proceedings-of-the-4th-International-Modal-Analysis/.

Samuel Stanley Wilks. *Mathematical Statistics*. Wiley series in probability and mathematical statistics. Wiley, 1 edition, 1963a.

Samuel Stanley Wilks. *Mathematical Statistics*. Wiley series in probability and mathematical statistics. Wiley, "second printing with corrections" edition, 1963b. URL https://books.google.com/books?id=-sJLAAAAMAAJ.

Peter Willett, Peter F. Swaszek, and Rick S. Blum. The good, bad, and ugly: Distributed detection of a known signal in dependent gaussian noise. *IEEE Transactions on Signal Processing*, 48(12): 3266–3279, December 2000.

P. Wojtaszczyk. *A Mathematical Introduction to Wavelets*, volume 37 of *London Mathematical Society student texts*. Cambridge University Press, February 13 1997. ISBN 9780521578943. URL http://books.google.com/books?vid=ISBN0521578949.

W. J. Yan and W. X. Ren. Operational modal parameter identification from power spectrum density transmissibility. *Computer-Aided Civil and Infrastructure Engineering*, 27(3):202–217, 2012. URL https://doi.org/10.1111/j.1467-8667.2011.00735.x.

Frank Yates and Kenneth Mather. Ronald aylmer fisher, 1890–1962. *Biographical Memoirs of Fellows of the Royal Society of London*, 9:90–120, November 1963. URL https://doi.org/10.1098/rsbm.1963.0006. print ISSN:0080-4606, online ISSN:1748-8494, https://pdfs.semanticscholar.org/6ccd/4303f68f3543fc794c376ce00395947a79b6.pdf.

Robert M. Young. *An introduction to nonharmonic Fourier series*, volume 93 of *Pure and applied mathematics*. Academic Press, 1 edition, 1980. ISBN 0127728503. URL http://books.google.com/books?vid=ISBN0127728503.

Robert M. Young. *An introduction to nonharmonic Fourier series*, volume 93 of *Pure and applied mathematics*. Academic Press, revised first edition, May 16 2001. ISBN 0127729550. URL http://books.google.com/books?vid=ISBN0127729550.

William Henry Young. On classes of summable functions and their fourier series. *Proceedings of the Royal Society of London*, 87(594):225–229, August 1912. URL http://www.archive.org/details/philtrans02496252.

Yu Zhang, Tong Zhu, and Jing Zhou. Understanding power spectrum density transmissibility. *Shock and Vibration*, 2016(8479721):9, 2016. URL http://dx.doi.org/10.1155/2016/8479721.

Mu-Qiao Zheng, She-Ping Hu, and Li Li. Improved methods for the estimate of frequency response functions. pages 917–921, 2002. URL https://pdfs.semanticscholar.org/edfc/0572a727633e729204bc9a8cbce4eb3211e5.pdf.

Copyright © 2019 Daniel J. Greenhoe
CC BY-NC-ND

Yun-Lai Zhou and Magd Abdel Wahab. Republished paper. rapid early damage detection using transmissibility with distance measure analysis under unknown excitation in long-term health monitoring. *Journal of Vibroengineering*, 20(1):823–831, 2018. doi: 10.21595/jve.2016.19718. URL https://doi.org/10.21595/jve.2016.19718.

Copyright © 2019 Daniel J. Greenhoe

REFERENCE INDEX

Abramovich and Aliprantis (2002), 206, 354
Aliprantis and Burkinshaw (2006), 212, 214–217
Abdaheer (2009), 97, 105
Abom (1986), 112
Aliprantis and Burkinshaw (1998), 226, 228, 232, 255, 256, 258–261, 263, 264, 268, 271–273, 298, 304, 308, 310–312
Adobe (1999), 400
Akhiezer and Glazman (1993), 208
Allemang et al. (1979), 111
Allemang et al. (1987), 112
Allen (2018), 6
Amann and Escher (2008), 300
Amir (1986), 263, 278–280
Anderson (1958), 186, 193
Anderson (1984), 186, 193
Angus (1994), 25
Apostol (1975), 281
Aubin and Frankowska (2009), 283
Aubin (2011), 283
Autonne (1901), 326
Autonne (1902), 326
Bachman (1964), 336
Bachman and Narici (1966), 219, 221, 223, 232, 318, 320
Bachman et al. (2000), 224, 227, 236, 333
Bailey et al. (2011), 388
Bailey et al. (2013), 388
Bajpai (1967), 98
Bak (2013), 123
Balakrishnan and Lai (2009), 25
Banach (1922), 207, 210, 211, 271, 272
Banach (1932b), 271
Banach (1932a), 224, 271, 274
Barvinok (2002), 282, 283
Beer (1993), 283
Bell (1986), 298
Bendat and Piersol (1966), 65
Bendat (1978), 111
Bendat and Piersol (1980), 65, 127
Bendat and Piersol (1993), 111, 127
Bendat and Piersol (2010), 65, 111, 112, 127
?, 44
Berberian (1961), 209, 210, 219, 232, 236, 253, 255, 262, 289, 304–307, 323
Bertero and Boccacci (1998), 320
Besso (1879), 292
Bienaymé (1840), 292
Billingsley (1995), 164
Blumenthal (1970), 287
Bollobás (1999), 189–191, 228, 255, 259, 261, 262, 268, 277, 283, 284, 290–292, 295, 312, 313
Bolstad (2007), 66
Bouniakowsky (1859), 298
Boyer and Merzbach (2011), 298
Bracewell (1978), 232, 336
Bromwich (1908), 353
Bullen (2003), 291, 292, 295–297, 299
Bunyakovsky (1859), 256
Byrne (2005), 245
Cadzow (1987), 11, 12, 44
Caines (1988), 8
Caines (2018), 8
Callahan (2010), 26
Cardano (1545), 205
Carne and Dohrmann (2006), 114
Carothers (2000), 296, 297, 299
Carothers (2005), 221, 224
Cauchy (1821), 256, 295, 298
Cayley (1858), 189
Chen et al. (2012), 132
J.S.Chitode (2009b), 105
J.S.Chitode (2009a), 349
Choi (1978), 65, 66
Chorin and Hald (2009), 333, 334
Christensen (2003), 219, 224, 232, 237, 242, 245
Christensen (2008), 237, 242, 367, 370, 371, 376, 378, 380, 382, 384
Christensen (2010), 370
Chui (1988), 367, 376, 378, 384
Chui (1992), 346, 367, 371, 380
Cifarelli and Regazzini (1996), 164
Çinlar and Vanderbei (2013), 284
Claerbout (1976), 357
Clarkson (1993), 65
Cobb (1988), 109, 112, 116
Cobb and Mitchell (1990), 116
Cohn (2002), 206
Conway (1990), 259
Courant and Hilbert (1930), 192
Cover and Thomas (1991), 6, 286
Cox (1972), 376, 378, 384
Curry and Feldman (2010), 173
Daubechies (1992), 225, 241,

349
Day (1973), 263
de Witt (1659), 191
de Boor (1972), 376, 378, 384
de Boor (2001), 376, 378, 384
de Finetti (1930a), 164
de Finetti (1930b), 164
de Finetti (1930c), 164
Dekking et al. (2006), 6
Descartes (1954), 189
Descartes (1637), 189
Devroye (1986), 25, 29
Daubechies et al. (1986), 242
Dieudonné (1969), 321
Drljević (1989), 267, 280
Duffin and Schaeffer (1952), 242
Dumitrescu (2007), 357
Dunford and Schwartz (1957), 278, 280, 314
Durbin (2000), 205, 206
Duthie (1942), 281
Dutka (1995), 69
Edelman and Jamison (1985), 287
Edelman (1986), 287
Edwards (1995), 228
Eidelman et al. (2004), 306
Enflo (1973), 224
Ewen (1950), viii
Ewen (1961), viii
Ewins (1986), 110, 132
Fabian et al. (2010), 230
Farina and Rinaldi (2000), 357
Feldman and Valdez-Flores (2010), 173
Felippa (1999), 193
Feller (1970), 167
Fisher (1922), 79
Fix and Strang (1969), 339
Folland (1995), 249, 250, 253, 254
Forster and Massopust (2009), 390
Fourier (1820), 289
Fourier (1878), 333
Fourier (1822), 333
Fréchet (1930a), 164
Fréchet (1930b), 164
Fredholm (1900), 18, 191
Fredholm (1903), 18, 191
Frobenius (1968), 321
Frobenius (1878), 321
Fuchs (1995), 249
Gelfand (1941), 253
Gelfand and Neumark (1943b), 252–254
Gelfand and Neumark (1943a), 253, 254
Gelfand and Naimark (1964), 250, 251, 253
Giles (1987), 273, 274
Giles (2000), 224, 228, 272, 274, 314, 316
Gohberg et al. (2003), 192
Goldman (1999), 111
Goswami and Chan (1999), 338
Goyder (1984), 112, 116
Gradshteyn and Ryzhik (1980), 200
Graham (2018), 193, 197
Graham (1981), 193, 197–199
Greechie (1971), 164
Greenhoe (2017a), 374
Greenhoe (2019), 13
Gupta (1998), 223
Haar (1910), 396
Haaser and Sullivan (1991), 207, 208, 223, 255–257, 260
Haddad and Akansu (1992), 361
Hall and Knight (1894), 300
Halmos (1948), 207
Halmos (1958), 318
Halmos (1998), 251, 263, 317
Hamel (1905), 209, 221
Han et al. (2007), 230, 245
Hardy (1929), 289
Hardy et al. (1952), 291, 296–298
Haykin (1994), 26
Hazewinkel (2000), 249
Heijenoort (1967), viii
Heil (2011), 219, 221, 223, 224, 232, 237, 242, 245, 259, 303, 370, 393
Heinbockel (2010), 388
Helstrom (1991), 44
Hijab (2011), 375
Hilbert (1904), 189
Hilbert (1906), 190, 191
Hilbert (1912), 189, 191
Hilbert et al. (1927), 304
Hölder (1889), 296
Höllig (2003), 375, 376, 378, 384
Hörmander (1994), 287
Horn and Johnson (1990), 74, 202, 308
Housman (1936), viii
Ifeachor and Jervis (1993), 71
Ifeachor and Jervis (2002), 71, 105
Igari (1996), 240
Istrăţescu (1987), 214, 255, 262, 263
James (1945), 267, 278, 280
James (1947), 280
Jawerth and Sweldens (1994), 338, 339, 384
Jeffrey and Dai (2008), 334
Jenkins and Watts (1968), 103
Jensen (1906), 6, 283, 284, 295
Jolley (1961), 387, 388
Jordan and von Neumann (1935), 261–263
Jørgensen et al. (2008), 242
Joshi (1997), 353
Kalmbach (1986), 164
Kamel and Sim (2004), 135
Kay (1988), 44, 65, 71, 96
Keener (1988), 18, 20, 191, 320
Kenko (circa 1330), 404
Kenney (1947), 98
Khamsi and Kirk (2001), 287
Kim and Kamel (2004), 135
Knapp (2005a), 327
Knapp (2005b), 333, 334
Kolmogorov (1933b), 164
Kolmogorov (1933a), 164
Koopmans (1995), 44
Kubrusly (2001), 207, 209, 219, 221, 223, 226, 232, 236, 265, 266, 268, 304, 318, 323, 324
Kubrusly (2011), 353
Kurdila and Zabarankin (2005), 219, 284
Lax (2002), 320
Lay (1982), 284
Leclere et al. (2014), 115, 121
Leibniz (1679), 303
Leuridan et al. (1986), 109, 111
Liang and Lee (2015), 132
Lieb and Loss (2001), 202
von der Linden et al. (2014), 174
Lindenstrauss and Tzafriri (1977), 224
Lindquist and Picci (2015), 8
Loomis (1953), 265
Loomis and Bolker (1965), 333
Machiavelli (1961), 403
Maeda and Maeda (1970), 282
Maleknejad et al. (2013), 372
Maligranda (1995), 260, 296, 297
Mallat (1989), 241
Mallat (1999), 241, 338, 339, 351, 361, 363, 391
Manolakis et al. (2005), 44
S. Lawrence Marple (1987), 13
S. Lawrence Marple (2019), 13
MatLab (2018a), 11

BY-NC-ND
COPYRIGHT © 2019 DANIEL J. GREENHOE

MatLab (2018b), 11
Mazur and Ulam (1932), 314
Mazur (1938), 253
Menger (1928), 287
Menini and Oystaeyen (2004), 300
Mercer (1909), 192
Michel and Herget (1993), 18, 189, 191, 192, 206, 209–211, 219, 221, 223, 251, 252, 271, 274, 303, 305, 307, 310, 315, 316, 319, 321, 324–326, 330, 331
Manolakis et al. (2000), 76
Miller (2006), 173
Milovanović and Milovanović (1979), 291
Minkowski (1910), 260, 297
Mintzer (1985), 361
Mitchell (1980), 112
Mitchell (1982), 112, 128
Mitrinović et al. (2010), 284
Moon and Stirling (2000), 8, 186
Mulholland (1950), 291
Nahin (2011), 388
Narens (2014), 164
Nelles (2001), 63, 75–77
Noble and Daniel (1988), 330
Greenhoe (2017b), 374
Oikhberg and Rosenthal (2007), 276, 314
Ore (1935), 281
Osgood (2002), 179
Paine (2000), vi
Pap (1995), 164
Papoulis (1984), 11, 12, 18, 26, 28, 29, 44
Papoulis (1990), 6, 26, 31, 165, 166, 175, 182
Papoulis (2002), 44
Papoulis (1991), 17, 26, 28–30, 35–37, 39, 41, 45, 50, 150, 164, 166, 167, 174, 181
Peano (1888b), 207, 210, 211, 255
Peano (1888a), 210, 211
Pečarić et al. (1992), 291
Pedersen (2000), 189, 227, 228
Pintelon and Schoukens (2012), 112, 116
Plackett (1972), 69
Poincaré (1912), 180
Popper (1962), 109
Popper (1963), 109
Prasad and Iyengar (1997), 367, 378, 390, 391
Proakis and Manolakis (1996), 105
Proakis (2002), 105
Proakis (2001), 26, 28, 29, 182–184, 186
Rana (2002), 292
Rao and Swamy (2018), 105
Rickart (1960), 250–253
Riesz (1934), 263
Robinson (1962), 357
Robinson (1966), 357
Rockafellar (1970), 284
Rosenblatt (1956), 65
Ross (1998), 6, 7, 181–183
Roussas (2014), 25
Rudin (1991), 308, 310, 311, 313, 315–318, 320–322, 326, 327
Rudin (1976), 294
Rynne and Youngson (2008), 223
Sakai (1998), 317
Salivahanan and Vallavaraj (2001), 105
Scharf (1991), 193, 195, 198
Schauder (1927), 224
Schauder (1928), 224
Schumaker (2007), 369, 370, 372, 375, 376, 378, 384
Schur (1909), 326
Scharz (1885), 298
Schwarz (1885), 256
Searcóid (2002), 219
Shao (2003), 6, 66
Shin and Hammond (2008), 109, 111, 112, 115, 116, 121, 128, 130
Shizgal (2015), 372
Silverman (1986), 65
Simmons (2007), 403
Simon (2011), 283
Singer (1970), 224
Slaught and Lennes (1915), 122
Smith and Barnwell (1984a), 361
Smith and Barnwell (1984b), 361
Snell (1997), 173
Snell (2005), 173
Sohrab (2003), 223
Sorenson (1970), 69
Mandyam D. Srinath (1996), 63, 85, 88, 89
Steele (2004), 258
Steen (1973), 189, 326
Stigler (1981), 69
Stoer and Bulirsch (2002), 370
Stone (1932), 304, 316, 320
Strang and Nguyen (1996), 361, 392, 393
Strang (1989), 392, 393
Stuart and Ord (1991), 65, 98
Suhov et al. (2005), 6
Sweldens and Piessens (1993), 338, 339
Szabó and Horváth (2004), 372
Thomson et al. (2008), 353
Thong et al. (2001), 135
Tolsted (1964), 296, 297
Trivedi (2016), 173
Tucker (1965), 98
Udriste (1994), 284
Ulam (1991), 314
Vaidyanathan (1993), 361
Väisälä (2003), 314
Trees (2001), 141
Trees (2002), 193, 195, 198
Vaseghi (2000), 103
van de Vel (1993), 282, 287
Vidakovic (1999), 241, 338, 339, 351, 363
von Neumann (1929), 191, 316, 320
Walter and Shen (2001), 228
Weber (1893), 206
Wei and Billings (2006), 372
Wells (1987), 388
Weyl (1940), 228
White et al. (2006), 112, 115, 130
Wicks and Vold (1986), 115, 130
Wilks (1963a), 98, 100
Wilks (1963b), 4, 7, 11, 98
Willett et al. (2000), 158
Wojtaszczyk (1997), 237, 239–241, 376, 378, 383, 384, 390, 396
Yan and Ren (2012), 111
Yates and Mather (1963), 11
Young (1912), 296
Young (1980), 230
Young (2001), 221, 223–225, 230, 234, 235, 237, 242
Zhang et al. (2016), 111
Zheng et al. (2002), 112, 113, 115, 135
Zhou and Wahab (2018), 111

Copyright © 2019 Daniel J. Greenhoe
CC BY-NC-ND

COPYRIGHT © 2019 DANIEL J. GREENHOE

SUBJECT INDEX

0 order B-spline, 385
0 order B-spline dyadic decomposition, 392
0th order B-spline, 369
1st order B-spline, 370, 385
1st order B-spline dyadic decomposition, 393
2nd order B-spline, 370, 385
2nd order B-spline dyadic decomposition, 393
3rd order B-spline, 372
4th order B-spline, 373
5th order B-spline, 373
6th order B-spline, 373
7th order B-spline, 373, 374
C^* algebra, **253**, **254**
C^*-algebra, 254
$\Gamma(b)$, 181
$*$-algebra, **12**, **15**, **35–37**, **250**, 250–252, 317, **344**
$\langle \triangle \mid \triangledown \rangle$, 255
αth-order modified Bessel function of the first kind, **185**
nth derivative, 380
nth moment, **338**, 338, 339
nth order B-spline, 371, 376, 378, 384, 385, 391, 393, 394
nth order cardinal B-spline, **367**
nth-order B-spline, 383
N-tuple, 209, 271
ϕ-mean, **290**
(ESD), 44
*-algebra, 70
*-algebras, 317
LaTeX, vi
TeX-Gyre Project, vi
XeLaTeX, vi
attention markers, 146, 150, 265, 314
    problem, 189, 308, 313, 320, 322
σ-additive, 164, 170
σ-additivity, 164
σ-algebra, 165, 170, 172
0 order B-spline, 382
1st order B-spline, 383
2 coefficient case, **396**
2nd order B-spline, 383
4th central moment, 99

inverse, 140

Abel, 298
Abel, Niels Henrik, 403
abelian group, 208
absolute value, x, **206**, **214**, 215–217, 256, 271, 273
absolutely summable, 15, 354, 355
ADC, 111
additive, 19, 164–167, 169, 255, 256, 265, 267, 268, 304, 307
additive identity, 12, 66, 209–211
additive identity element, 209, 210
additive inverse, 210, 211
additive property, 334
additivity, 19, 164, 165, 246, 259, 316
adjoint, 252, **313**, 313, **316**, 317, 334
admissibility condition, 396
admissibility equation, 396
Adobe Systems Incorporated, vi
affine, 190, **283**, 314
Affine equations, **197**
affine equations, 71
algebra, 205, **206**, 207, **249**, 249, 250
algebra of sets, xi, 171
algebraic, 207
algebras
    C^*-algebra, 253
    $*$-algebra, 250
algebras of sets, 171
Alternating Series Test, 390
AM-GM inequality, 295
analog-to-digital converter, 111
AND, x
anti-derivative, 368
anti-symmetric, 13, 14, 48, 337
anti-symmetric , 48
antiautomorphic, 15, 250, 251, 256, 317
antilinear, 251
antitone, 166, 169
Aristotle, 11, 17
arithmetic, 98
arithmetic mean, 95, 97, 112, **295**
arithmetic mean estimate, 65
arithmetic mean geometric mean inequality, 295
Arithmetic Mean transfer function estimate, **112**
associates, 208
associative, 5, 205, 206, 208, 210, 289, 290, 306, 307, 326
asymmetric, 357
asymptotically unbiased, 103
auto-correlation, **12**, 13, 18, 19, 48
auto-correlation function, **18**
auto-correlation operator, **18**, 20
auto-power spectrum, 385
auto-spectral density, **14**, 16

autocorrelation, 320
Avant-Garde, vi
average, **96**, 96–99, **295**
average power, **44**
AWGN, 79
Axiom of Continuity, 164

B-spline, 34, 372, 374, 380, 381, 383, 384, 391, 394, 398–400
B-spline dyadic decomposition, 391, 397
B-spline general form, 378
B-spline partition of unity, 384
B-spline recursion, 376
B-spline scaling coefficients, **394**
B-splines, 385
Bak, Per, 123
ball, 282
Banach algebra, **253**
Banach space, 224, 225
Bartlett Method, 105
basis, 20, 22, 221–223, 230, 384
  frame, 242
  orthogonal, 229
  orthonormal, 229
  Riesz, 237
  tight frame, 242
basis sequence, 383
Battle-Lemarié orthogonalization, **241**
Bayesian, 63
Bayesian estimate, **64**
Bessel's correction, 98
Bessel's Equality, **227**, 228, 229, 233
Bessel's Inequality, 235
Bessel's inequality, **228**
Best Approximation Theorem, **228**, 228
biased, 98, 99, 112
biased sample variance, **98**, 98
bijection, 235
bijective, xi, 235, 314
bilinear functionals, 262
binomial coefficient, 369, 370
Binomial Theorem, 5, 66, 100
binomial theorem, 395
biorthogonal, **237**, 239, 240
Birkhoff orthogonality, 278, **280**, 280
Blackman-Tukey Method, 105
Bohr, Harald, 289
Boole's inequality, 167
Boolean, 165, 167
Boolean algebra, 170
Boolean lattice, 164, 167
Borel algebra on $\mathbb{R}$, 170
Borel measure, 333
Borel sets, 333
bounded, xi, 166, 225, **311**, 320, 331
bounded bijective, 238
bounded lattice, 167
bounded linear operators, **311**, 313, 315, 316, 318, 319, 321, 322, 324–327
bounded operator, 311

C star algebra, **253**
calculus, 207
Cardano, Gerolamo, 205
Carl Spitzweg, 403
Cartesian product, x
Cauchy Schwartz inequality, 118, 120
Cauchy-Bunyakovsky-Schwarz Inequality, 256
Cauchy-Schwartz inequality, 9, 97
Cauchy-Schwarz Inequality, 93, 97, **256**, 256, 259, 260
Cauchy-Schwarz inequality, 243, 299
Cauchy-Schwarz Inequality for inner-product spaces, 299
Cauchy-Schwarz inequality for inner-product spaces, 299
Cauchy-Schwarz Inequality for sequences, **298**, 299
Cauchy-Schwarz inequality for sequences, 299
causal, **358**, 358
cdf, 25, **174**, 174
Chain Rule, 375
chain rule, 27, 199
channel capacity, 142
channel coding, 142
channel statistics, 142
characteristic function, x
Chebyshev's inequality, 7
chi-square distribution, **182**
Chi-squared distribution, 182
chi-squared distribution, 182
Chi-squared distribution with n degrees of freedom, **183**, 183
chi-squared distribution with n degrees of freedom, 183
closed interval, **281**, **282**, 282
closed support, 378
closure, 223, 378
coefficient functionals, **224**, 224
coefficients, 359
coherence, 131, 132
colored, 19
communication system, 126, 139
communications additive noise cross-correlation, 55
communications additive noise model, **110**, 123
Communications additive noise system, 55
communications LTI additive noise model, **127**, **135**
commutative, 12, 206, 208, 215, 306, 353, 354
commutative group, 208
commutative ring, 205, **206**, 206
compact, 191
compact support, 391
comparable, 281
comparison test, 387
complement, x
complete, **223**
complete metric space, 223
complete set, 223
complex, 120
complex coherence, **132**
complex exponential, 381
complex linear space, **208**
complex transmissibility, **114**, 114, 132
complex-valued, 114
concave, **282**, 282, **283**, 283, **284**, 284, 291, 292
conditional cumulative distribution function, **174**, 174
conditional probability, 65, 167
conditional probability density function, **174**, 174
conditional probability of x given y, **166**
conjuage symmetric, 18
conjugate linear, 250, 251, 317
conjugate pairs, 359
conjugate quadrature filter, **361**, 361
conjugate quadrature filter condition, **361**
Conjugate quadrature filters, 361
conjugate symmetric, 14, 15, 18, 48, 192, 255, 256, 264, 265, 268
conjugate symmetric property, 334
conjugate symmetry, 19
conjugate-symmetric, 15

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

consistent, 96, 97, 99, 103
constant, 12, 43, 65
constraint, 140
Continuity, 274
continuity, 175, 176
continuous, xi, 25, 168, 175, 192, 253, 259, 274, 290, 292, 306, 380, 386, 393
Continuous data whitening, 91
continuous point spectrum, 381
continuous spectrum, **190**
contradiction, 210
convergence, 387
convergence in probability, 20
convergent, 44, 225
converges, 387–389
converges absolutely, 192
Convex, 284
convex, 5, 6, 274, 276, 282, **283**, 283, **284**, 284, 291, 292
functional, 283
strictly, 283
convex function, 283, 284
convex set, **282**, 282–284
convolution, 31, 32, 35, 36, 38, 45, 57, 182, 336, **353**, 353, 354, 367
convolution operation, **336**
convolution operator, 353, 354
Convolution Theorem, 38, 382
convolution theorem, **336**, 341, **356**
coordinate functionals, **224**
coordinates, **221**
correlated, 52, 110, 111
cost function, 63, 70, 123
countably infinite, 191, 224
counting measure, xi
CQF, **361**, 361, 362
CQF condition, **361**, 363
CQF theorem, **361**, 364
Cramér-Rao bound, 90
Cramér-Rao lower bound, 88
criterion, 140
critical point, **26**, 26, 28, 29
cross spectral density, **14**
cross-correlation, **12**, 13, **18**, 18
CS Inequality, 246
CSD, **14**
cummulative distribution function, **174**, 174
cumulative distribution function, 25, 168

decision region, 157, 158
decreasing, 291, 390
definitions
C^* algebra, **253**, **254**
$*$-algebra, **12**, **15**, **35–37**, **250**, **344**
nth order cardinal B-spline, **367**
absolute value, **214**
algebra, **206**, **249**
Banach algebra, **253**
biorthogonal, **237**
Birkhoff orthogonality, **280**
bounded, **311**
bounded linear operators, **311**
C star algebra, **253**
closed interval, **281**, **282**
coefficient functionals, **224**
commutative ring, **206**
complete, **223**
complex linear space, **208**
conditional probability of x given y, **166**
continuous spectrum, **190**
convex set, **282**
coordinate functionals, **224**
coordinates, **221**
CQF, **361**
difference, **210**
disjoint, **216**
disjoint complement, **217**
eigenspace, **189**
epigraph, **283**
equal, **303**
equivalent, **225**
even/odd die toss, **169**
event, **165**
exact frame, **242**
expansion, **221**, **224**
field, **206**
Fourier coefficients, **230**
Fourier expansion, **230**
Fourier series, **230**
frame, **242**
frame bounds, **242**
fundamental, **223**
group, **205**
half-open interval, **281**, **282**
Hamel basis, **221**
hermitian, **250**
Hessian matrix, **203**
hypograph, **283**
inner product space, **255**
intervals on lattices, **281**
intervals on ordered sets, **281**
isometric, **323**
isosceles orthogonality, **278**
James orthogonality, **278**
linear basis, **221**
linear combination, **219**
linear space, **208**
linear subspace, **208**
minimum phase, **357**
multiplicative condition, **253**
negative part, **214**
norm, **271**
norm induced by the inner product, **260**
normal, **250**
normalized tight frame, **242**
normed algebra, **253**
normed linear space, **308**
normed space of linear operators, **308**
open interval, **281**, **282**
operator norm, **308**
optimal lower frame bound, **242**
optimal upper frame bound, **242**
ordered linear space, **212**
orthogonal basis, **230**
orthogonal in the Pythagorean sense, **280**
orthogonal in the sense of Birkhoff, **280**
orthogonal in the sense of James, **278**
orthonormal basis, **230**
outcome, **165**
Parseval frame, **242**
point spectrum, **190**
positive, **212**, **330**
positive cone, **212**
positive part, **214**
power mean, **292**
probability space, **165**, **170**
projection, **250**
Pythagorean orthogonality, **280**
real linear space, **208**
residual spectrum, **190**
resolvent, **249**
resolvent set, **190**
Riesz basis, **237**
Riesz space, **212**
ring, **205**

COPYRIGHT © 2019 DANIEL J. GREENHOE
BY-NC-ND

scalars, **208**
Schauder basis, **224**
self-adjoint, **250**
sequence, **353**
set projection operators, **157**
Smith-Barnwell filter, **361**
space of all absolutely square summable sequences, **353**
spans, **219**
spectral radius, **249**
spectrum, **190**, **249**
star-algebra, **250**
summation, **289**
tight frame, **242**
total, **223**
underlying set, **208**
unital, **249**
unitary, **326**
vector lattice, **212**
vector space, **208**
vectors, **208**
delay, 329
dense, 223, 224
Descartes, René, ix
detection, 63
deterministic, 69
difference, x, **210**
differentiable, 26
differentiable function, 26, 28, 29
differential operator, 381
dilation, 328
dilation equation, 394, 397
dimension, **221**
direction, 75
Dirichlet Lambda function, 387, 388
Discrete data whitening, 91
Discrete Time Fourier Series, xi
Discrete Time Fourier Transform, xii, 14, 343
discrete-time Fourier transform, **343**, 343–345, 349
discriminate, 258
disjoint, **216**
disjoint complement, **217**
distance, 273
distributes, 208
distributive, 12, 35–37, 70, 164–167, 210, 250–252, 255, 256, 317, 344, 355
distributive pseudocomplemented lattice, 164
domain, x
Double coin toss, **168**, 168
DTFT, 14–16, 109, 225, 344, 348, 351, 361–363, 395
DTFT periodicity, **343**
dyadic decomposition, 391

efficient, 85, 88, 89, 91
eigen-system, 21
Eigendecomposition, 244
eigenspace, **189**
eigenvalue, **189**, 244
eigenvector, **189**, 244
empty set, xi
energy, 357
energy spectral density, **44**
ensemble average, 44
epigraph, **283**
equal, **303**
equalities
Bessel's, 227
equality
triangle, 272
equality by definition, x
equality relation, x
equations
parallelogram law, 263
equivalent, **225**, 225, 237
ergodic in the mean, **44**, 44, 45
error vector, 70
estimate, **64**, 109, 141
Euclidean metric, 276
Euclidean space, 282
Euler formulas, 340, 382
even, 348
even/odd die, 170
even/odd die probability space, 169
even/odd die toss, **169**
event, **165**, 165, 166
events, 165
exact frame, **242**
examples
2 coefficient case, **396**
Double coin toss, **168**
even/odd die, 170
even/odd die toss, **169**
Mercedes Frame, **245**
Peace Frame, **245**
Rectangular pulse, 341
rectangular pulse, **340**, 382
single coin toss, **168**, 170
The usual norm, **273**
triangle, **341**
tuples in $\mathbb{F}^N$, **209**
excluded middle, 166, 167
exclusive OR, xi
existential quantifier, xi
exists, 44
expansion, **221**, **224**
expectation, 5, 110
expectation operator, **3**, 3, 4, 6

false, x
FFT, 13
field, 205, **206**, 207, 209, 353
field of complex numbers, 317
field of probability, 164
filter banks, 361
finite modular lattice, 164
FontLab Studio, vi
for each, xi
Fourier analysis, 380
fourier analysis, 333
Fourier coefficients, **230**
Fourier expansion, **230**, 230, 231, **234**, 235, 236
Fourier kernel, **333**
Fourier Series, xi
Fourier series, **230**
Fourier series expansion, 232
Fourier Transform, xi, xii, 45, 241, 329, 333, **334**, 337, 382
adjoint, **334**
Fourier transform, 225, 329, 338, 340, 341, 381
inverse, **334**
Fourier transform magnitude, 382, 383
Fourier transform scaling factor, 334
Fourier, Joseph, 333
frame, 240, **242**, 242, 246
frame bound, 243, 245
frame bounds, **242**, 246
frame operator, **242**, 242, 243, 245
frames, 220
Fredholm integral equation of the first kind, 18, **191**
Fredholm integral operators, 328
Fredholm operator, **191**, 191, 192
Fredholm operators, 191
Free Software Foundation, vi
Frequency Response Function, 109
Frequency Response Identification, 109
Frequency Shift Keying
coherent, 145
FRF, 109
FSK
coherent, 145
function, 26, 173, 174, 303, 333, 368
even, 348
step, 367
function overload, 13
functional, 255, 284, 317
functions, xi
0 order B-spline, 385

BY-NC-ND
COPYRIGHT © 2019 DANIEL J. GREENHOE

0th order B-spline, 369
1st order B-spline, 370, 385
2nd order B-spline, 370, 385
3rd order B-spline, 372
4th order B-spline, 373
5th order B-spline, 373
6th order B-spline, 373
7th order B-spline, 373, 374
αth-order modified Bessel function of the first kind, **185**
nth derivative, 380
nth moment, 338
nth order B-spline, 371, 376, 378, 384, 385, 391, 393, 394
nth-order B-spline, 383
ϕ-mean, **290**
(ESD), 44
0 order B-spline, 382
1st order B-spline, 383
2nd order B-spline, 383
4th central moment, 99
absolute value, **206**, 215, 256, 271
adjoint, 252
anti-derivative, 368
arithmetic, 98
arithmetic mean, 95, 97, 112, **295**
arithmetic mean estimate, 65
Arithmetic Mean transfer function estimate, **112**
auto-correlation, **12**, 13, 18, 19, 48
auto-correlation function, **18**
auto-power spectrum, 385
auto-spectral density, 16
average, **96**, 96–99, **295**
average power, **44**
B-spline, 34, 372, 374, 380, 381, 383, 384, 391, 394, 398–400
B-splines, 385
Bayesian estimate, **64**
biased sample variance, **98**, 98
binomial coefficient, 369, 370
Borel measure, 333
cdf, 25, **174**, 174
chi-square distribution, **182**
Chi-squared distribution with n degrees of freedom, **183**
coherence, 131, 132
complex coherence, **132**
complex exponential, 381
complex transmissibility, **114**, 114, 132
conditional cumulative distribution function, **174**, 174
conditional probability, 65
conditional probability density function, **174**, 174
conjugate quadrature filter, **361**
continuous point spectrum, 381
cost function, 63, 70, 123
cross-correlation, **12**, 13, **18**
cummulative distribution function, **174**, 174
cumulative distribution function, 25, 168
dilation equation, 394, 397
Dirichlet Lambda function, 387, 388
Discrete Time Fourier Transform, 343
discrete-time Fourier transform, 343–345
DTFT, 348, 351, 362, 363
eigenvector, 244
energy spectral density, **44**
error vector, 70
estimate, **64**
Fourier kernel, **333**
Fourier transform, 338, 340, 341
Fredholm integral equation of the first kind, **191**
function, 174
Gamma distribution, **182**
gamma function, **181**
Gaussian distribution, **181**, **186**
generalized probability, **164**, 165
Geometric mean, 113
geometric mean, 113, **295**
geometric mean estimator, 113
Geometric mean transfer function estimate, **113**
Hanning window, 97
Harmonic mean, 114, 115
harmonic mean, **295**
Harmonic mean transfer function estimate, **114**
impulse response, 14, 15, 35, 37, 38, 56, 109
induced norm, 233, 259
inner product, 8, 9, 97, **255**, 259, 264, 265, 333, 380
input, 14, 15, 35, 37
integer sequence, 374
Inverse Method transfer function estimate, **112**
Jensen's inequality, 5
joint cumulative distribution function, **174**, 174
Joint Gaussian distribution, 185
Joint Gaussian pdf, **186**
joint probability density function, **174**, 174
kernel, 381
Kronecker delta, 21
Kronecker delta function, **267**
kronecker delta function, 349
Least Squares Technique, 111
Least Squares transfer function estimate, **112**
linear functional, 313
magnitude, 113
MAP estimate, **64**, 79
maximum a-posteriori probability estimate, **64**
maximum likelihood estimate, **64**
mean, **3**, **12**, 12, **18**, 35
mean integrated square error, **65**
mean square error, **65**, 65, 97
Mean square estimate, **64**
measure, 3, 164, 165, 168, 174, 175
measure-theoretic, 165
mini-max estimate, **64**
ML estimate, **64**, 79, 81, 84, 86, 89
MM estimate, **64**
modulus, **206**
MS estimate, **64**
non-central chi-squared distribution, 184
noncentral chi-square distribution, **184**
noncentral chi-square with n-degrees of freedom,

COPYRIGHT © 2019 DANIEL J. GREENHOE
CC BY-NC-ND

185
norm, 8, 9, 70, 97, 308
normal distribution, **181**, **186**
normalized rms error, **65**, 65
order n B-spline, 374
order n B-spline MRA coefficient sequence, 394
order n B-spline MRA scaling sequence, 394
order-k moment estimate, **101**
ordinary coherence, **132**
ordinary transmissibility, 114
output, 14, 15, 35, 37
pdf, **174**, 174
phase, 113, 114, 120
power mean, 292, 295
power spectral density, 16, 44
power sum, 374
probability, **165**
probability density function, 3, 25, 34, 168, **174**, 174, 182
probability function, **164**
quantization noise, 111
quantum probability, **164**, 165
random process, **17**, 19, 20, 44
random sequence, **11**, 11–14, 35, 39, 110
random variable, 3–8, **17**, 25, 26, 28, 29, 31, 33, 168, 173, **174**, 174, 175, 182
Rayleigh distribution, **185**
Rice distribution, **185**
Riesz sequence, **237**
sample variance, **98**
scalar product, **255**
scaling coefficient sequence, 396
scaling function, 393, 394
Scaling transfer function estimate, **115**, 115
sequence, 11, 34, 394
set indicator function, 341
signal-to-noise ratio, 111
sine sweep, 109
SNR, 111
spectral power, 16, **44**, 111
step function, **367**, 367–371
Total Least Squares transfer function estimate, **115**
Total least squares transfer function estimate, 115
traditional probability, **164**, 165
transfer function estimate $\hat{H}_\kappa(\omega;\kappa)$, **115**
transfer function estimate $\hat{H}_c(\omega)$, **116**
translation operator, 339
Transmissibility, 114
transmissibility, 111, 113
transmissibility $\tilde{T}_{xy}(\omega)$, **111**
uniform distribution, **179**, 179
variance, **3**, **12**, 12, 95
wavelet, 391
windowed autocorrelation estimate, **103**
Z transform, 395
z transform, 357
Z-transform, 345
z-transform, 362
fundamental, **223**
Fundamental Theorem of Calculus, 368, 375
Fundamental theorem of calculus, 175, 177
Fundamental theorem of linear equations, 307

Gamma distribution, **182**, 182, 183
gamma function, **181**, 181
Gaussian, 33
Gaussian distribution, 180, **181**, **186**
Gelfand-Mazur Theorem, **253**
General ML estimation, **84**
Generalized AM-GM inequality, 295
generalized arithmetic mean geometric mean inequality, 295
Generalized associative property, **289**
Generalized Parseval's Identity, 232
generalized probability, **164**, 165
Generalized triangle inequality, 9
geodesically between, **282**
Geometric mean, 113
geometric mean, 113, **295**
geometric mean estimator, 113
Geometric mean transfer function estimate, **113**
Geometric Series, **300**
geometry, 207
GNU Octave
for, 399
step, 399
Golden Hind, vi
gradient , 75
gradient of $\boldsymbol{y}$ with respect to $\boldsymbol{x}$, **193**
gradient of $\boldsymbol{y}^T$ with respect to $\boldsymbol{x}$, **193**
greatest lower bound, xi
group, **205**, 205
Gutenberg Press, vi

Hölder inequality, 299
Hölder's Inequality, **296**, 298
half-open interval, **281**, **282**
Hamel bases, 220
Hamel basis, **221**, 221, 223
Handbook of Algebras, 249
Hanning window, 97
Hardy, G.H., 289
harmonic analysis, 333
Harmonic mean, 114, 115
harmonic mean, **295**
Harmonic mean transfer function estimate, **114**
hermitian, **250**, 250, 251, **320**
hermitian components, 252
Hermitian representation, **252**
Hermitian symmetric, 71, 337
Hermitian Transpose, 71
Hessian, 76
Hessian matrix, **202**, **203**
Heuristica, vi
high-pass filter, 361
Hilbert space, 230, 232, 234, 236, 237, 240, 242, 316, 317, 320–322, 327, 333
homogeneous, 5, 206, 230, 231, 234, 238, 240, 255, 256, 260, 264, 265, 267, 271, 277, 298, 304, 305, 307, 308
Housman, Alfred Edward, vii
hypograph, **283**
hypotenuse, 280

identities
Fourier expansion, 230, 231
Parseval frame, 230, 231
identity, 205, 208, 303
identity element, 303
identity operator, 140, 196, 303, **304**

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

if, xi
if and only if, xi
image, x
image set, 305, 307, 318–322, 327
imaginary part, xi, **251**
implied by, xi
implies, xi
implies and is implied by, xi
impossible, 384
impulse response, 14, 15, 35, 37, 38, 56, 109
inclusive OR, xi
increasing, 291, 390
independence, 81, 84, 169
independent, 7, 31, 34, 85, 142, **166**, 168, 182
indicator function, x
induced norm, 233, 259, **260**
induces, 273
induction, 374, 375, 384
inequalities
AM-GM, 295
Cauchy-Bunyakovsky-Schwarz, 256, 298
Cauchy-Bunyakovsky-Schwarz Inequality, 256
Cauchy-Schwartz inequality, 97
Cauchy-Schwarz, 256, 298
Cauchy-Schwarz Inequality, 97, **256**, 256, 259, 260
Cauchy-Schwarz inequality, 299
Cauchy-Schwarz Inequality for inner-product spaces, 299
Cauchy-Schwarz inequality for inner-product spaces, 299
Cauchy-Schwarz Inequality for sequences, **298**, 299
Cauchy-Schwarz inequality for sequences, 299
Hölder, 296
Hölder inequality, 299
Hölder's Inequality, **296**, 298
Jensen's, 284
Jensen's Inequality, 291, 295
Minkowski (sequences), 297
Minkowski's Inequality, **260**, 291
Minkowski's Inequality for sequences, **297**
quadratic discriminate inequality, **258**
triangle inequality for vectors, 268
Young, 296
Young's Inequality, **296**
inequality
Bessel's, 228
triangle, 271, 308
infinite sum, 219
injective, xi, 306
inner product, 8, 9, 97, 207, **255**, 259, 261, 264, 265, 333, 380
Polarization Identity, 262
uniqueness, 261
inner product space, 8, 9, 207, 226–229, 237, **255**, 256, 259, 260, 263, 267, 268, 278, 380
inner-product, xi
inner-product space, 278, 280
input, 14, 15, 35, 37
inside, 358
instantaneous error, 76
integer sequence, 374
integral, 380
integral domain, 205
integral test, 388
intersection, x, 165
interval, 281, 282
intervals on lattices, **281**
intervals on ordered sets, **281**
into, 235, 236
inverse, 205, 208, 243, 245, 396
inverse DTFT, **349**, 363
Inverse Fourier transform, **334**
inverse Fourier Transform, 329
Inverse Method transfer function estimate, **112**
Inverse probability integral transform, **25**
invertible, 26, 168, 225, 242, 250
involutary, 317
involution, **250**, 250, 254
involutory, 15, 250–252
irreflexive ordering relation, xi
isometric, 231, 239, 240, 314, **323**, 323, 335
isometric operator, 321, 324–326
isometry, 323
isomorphic, 235
isosceles, 278
Isosceles orthogonality, 278
isosceles orthogonality, **278**, 278–280

Jacobian matrix, **193**, 193
James orthogonality, **278**
Jensen's Inequality, 6, **284**, 291, 295
Jensen's inequality, 5, **6**, 6
join, xi
joint cumulative distribution function, **174**, 174
Joint Gaussian distribution, 185
Joint Gaussian pdf, **186**
joint probability density function, **174**, 174

Kaneyoshi, Urabe, 403
Karhunen-Loève Expansion, **20**
Kenko, Yoshida, 403
kernel, 18, 381
Kolmogorov's inequality, **7**
Kronecker delta, 21
Kronecker delta function, **267**
kronecker delta function, 196, 349
kronecker product, 194
kronecker products, 194

l'Hôpital's rule, 118, 121, 122, 294, 387, 389, 393
lattice, 212, 281, 282
lattice with negation, 164
Laurent series, 354
Law of the Unconscious Statistician, **6**
Least squares, **72**
least squares, 72, 73
Least squares estimation, 123
least squares estimation, 76
Least squares estimations, 123
Least Squares Technique, 111
Least Squares transfer function estimate, **112**
least upper bound, xi
least-squares, 123, 126
Lebesgue integral, 170
Lebesgue integration, 378, 379
Lebesgue square-integrable functions, 333
left distributive, 5, 130, 205, 206, 307
left inverse, 140
Leibnitz GPR, 364
Leibniz Integration Rule, 376
Leibniz, Gottfried, ix, 303

COPYRIGHT © 2019 DANIEL J. GREENHOE
BY-NC-ND

length, 273
linear, 4, 5, 35, 36, 54, 55, 57, 110, 120, **304**, 304, 358
linear basis, **221**
linear bounded, xi
linear combination, **219**
linear combinations, 220
linear functional, 313
linear independence, 226
linear operator, 4, 5, 52, 54, 55, 238, 240
linear operators, 236, 304, 313
linear ops., 236, 240
linear space, 207, **208**, 208, 209, 219, 221, 249, 255, 261, 271, 277, 281–284, 303
linear spaces, 303
linear span, **219**
linear subspace, **208**, 219
linear time invariant, 37, 41, 111, 123, 127, 128, 381
linear time invariant systems, 45
linear time-invariant, 57
linearity, 5, 19, 21, 22, 27, 30, 31, 45, 51, 55–57, 66, 71, 95, 99, 100, 124, 195, 305, 370, 371, 389, 392
Linearity of E, **4**
linearly dependent, **219**, 221–223
linearly independent, **219**, 221–223, 226
Lippmann, Bernard A., 180
Liquid Crystal, vi
low-pass filter, 361
lower bounded, 167
LTI, 14, 15, 35–38, 45, 56–59, 111, 112, 123, 124, 127–129, 135, 381
    operations on non-stationary random processes, 41
    operations on WSS random processes, 43

Machiavelli, Niccolò, 403
magnitude, 113, 114
MAP, 64, 79
MAP estimate, **64**, 79, 81
maps to, x
Markov's inequality, **6**, 7
matched filter, 92
MatLab, 394
matrix
    rotation, 330
matrix algebra, 70
matrix calculus, 193
matrix:quadratic form, 198, 199
Maxima, 371, 373, 374
maximal likelihood (ML), 142
maximum a-posteriori, 64
maximum a-posteriori probability estimate, **64**
maximum a-posteriori probability estimation, 79
maximum likelihood, 64
maximum likelihood estimate, **64**
maximum likelihood estimation, 79, 84
    amplitude, 85
    general, 84
    phase, 88
Mazur-Ulam theorem, 314
mean, **3**, 3, **12**, 12, **18**, 35, 98
mean integrated square error, **65**
mean square error, **65**, 65, 97
Mean square estimate, **64**
measurable space, 165
measure, 3, 164, 165, 168, 174, 175
measure space, 165, 166
measure-theoretic, 165
measurement additive noise cross-correlation, 55
measurement additive noise model, **110**, 123
Measurement additive noise system, 55
measurement LTI additive noise model, **127**, **135**
measurement system, 126
meet, xi
memoryless, 142
Mercedes Frame, **245**
Mercer's Theorem, 21, **192**, 192
metric, xi, 207, 273, 274
    Euclidean, 276
    generated by norm, 274
    induced by norm, 274
    parabolic, 276
    sup, 276
    taxi-cab, 276
metric induced by the norm, 274
metric linear space, 207, 273, 274, 276
metric space, 207, 274, 281, 282
mini-max estimate, **64**
Minimum mean square estimation, **70**
Minimum phase, 365
minimum phase, 38, 49, **357**, 357, 358
    energy, 357
minimum variance unbiased estimator, **66**
Minkowski addition, 157
Minkowski's Inequality, **260**, 291
Minkowski's Inequality for sequences, **297**
ML, 64, 79
ML amplitude estimation, **85**
ML estimate, **64**, 79, 81, 84, 86, 89
ML estimation of a function of a parameter, **89**
ML phase estimation, **88**
MM estimate, **64**
modular, 164
modulus, **206**
monotone, 164, 166, 168
monotonically decreasing, 117
MRA, 394
MRA system, 394, 396
MS estimate, **64**
multiplicative condition, **253**
Multiresolution Analysis, 391
multiresolution analysis, 382, 393
mutually exclusive, 167, 170
MVUE, **66**

negative part, **214**, 215–217
Neumann Expansion Theorem, 315
Newton's method, 75
no input noise, 129
no output noise, 129
noise
    colored, 91
non-central chi-squared distribution, 184
non-contradiction, 166
non-isometric, 260
non-isotropic, 255, 256, 263, 265, 268
non-linear, 54, 55, 110, 126
non-negative, 19–21, 206, 255, 260, 263, 265, 268, 292, 308, 378, 384
non-positive, 258
non-stationary, 44
non-structured, 141
non-zero, 396
Noncentral chi-square distribution, 184
noncentral chi-square distribution, **184**
noncentral chi-square with n-degrees of freedom, **185**
nondegenerate, 164, 166, 169, 170, 206, 271, 277, 308
nonnegative, 164
nonparametric, 63

COPYRIGHT © 2019 DANIEL J. GREENHOE
BY-NC-ND

norm, 8, 9, 70, 97, 207, 257, 259–261, **271**, 271, 273, 274, 277, 308
induced by inner product, 260
Polarization Identity, 262
usual, 273
norm induced by the inner product, **260**, 260
normal, 191, **250**, 320, **321**, 321, 322
normal distribution, 180, **181**, **186**
normal operator, 321, 326
normalized, 20, 21, 164, 166, 169
normalized rms error, **65**, 65
normalized tight frame, **242**
normed algebra, **253**, 253, 254
normed linear space, 207, 261, 268, 271–274, 276, 278, 280, **308**
normed linear spaces, 313, 323
normed space of linear operators, **308**
NOT, x
not ergodic in the mean, 44
not included
metric, 274
not invertible, 26
not necessarily linear, 37, 51
not symmetric, 379
not total, 233
null space, x, 189, 305–307, 316, 318–322, 327

O_6 lattice, 164
Octave, 394
one sided shift operator, 325
only if, xi
open ball, 274
open interval, **281**, **282**
operations
0 order B-spline dyadic decomposition, 392
1st order B-spline dyadic decomposition, 393
2nd order B-spline dyadic decomposition, 393
adjoint, **313**, 313, **316**
auto-correlation, 19
auto-correlation operator, **18**, 20
auto-spectral density, **14**
Bartlett Method, 105
Blackman-Tukey Method, 105
closure, 378
communications additive noise model, **110**
communications LTI additive noise model, **127**, **135**
Continuous data whitening, 91
convolution, 31, 32, 35, 36, 38, 57, 182, **353**, 354, 367
convolution operation, **336**
cross spectral density, **14**
CSD, **14**
detection, 63
differential operator, 381
Discrete data whitening, 91
Discrete Time Fourier Series, xi
Discrete Time Fourier Transform, xii, 14
discrete-time Fourier transform, **343**
DTFT, 14–16, 109, 225, 344, 361, 363, 395
Eigendecomposition, 244
ensemble average, 44
estimate, 109
expectation, 5, 110
expectation operator, **3**, 3, 4, 6
FFT, 13
Fourier analysis, 380
Fourier Series, xi
Fourier Transform, xi, xii, 45, 241, 329, **334**, 337
Fourier transform, 225, 381
Fourier transform magnitude, 382, 383
frame operator, **242**, 242, 243, 245
Fredholm integral equation of the first kind, 18
Fredholm operator, **191**
Frequency Response Function, 109
Frequency Response Identification, 109
FRF, 109
gradient of $\boldsymbol{y}$ with respect to $\boldsymbol{x}$, **193**
gradient of $\boldsymbol{y}^T$ with respect to $\boldsymbol{x}$, **193**
Hermitian Transpose, 71
identity operator, **304**
imaginary part, **251**
integral, 380
inverse, 243, 245, 396
inverse Fourier Transform, 329
involution, **250**
Jacobian matrix, **193**
kernel, 18
Least squares estimation, 123
Least squares estimations, 123
Lebesgue integral, 170
Lebesgue integration, 378, 379
linear operator, 4, 5, 52, 54, 55
linear operators, 313
linear span, **219**
measure, 165
measurement additive noise model, **110**
measurement LTI additive noise model, **127**, **135**
operator, 242, **303**
operator adjoint, 317
Periodogram, 105
power spectral density, **14**
project, 380
projection, **318**
PSD, **14**
rationalizing the denominator, 122
real part, **251**
sampling, 4
Sum, 32
summation operator, 284
support, 378
system identication, 109
time average, 44
unitary Fourier Transform, **334**
variance, 44
vector addition, 195
Welch Method 0% overlap, 105
Welch Method 50% overlap, 105
z-domain cross spectral density, **14**
z-domain power spectral density, **14**
Z-Transform, xii
z-transform, **354**, 354
Z-transform operator, 14
operator, 207, 242, **303**
adjoint, 251
autocorrelation, 320
bounded, 311
definition, 303

COPYRIGHT © 2019 DANIEL J. GREENHOE

delay, 329
dilation, 328
identity, 303
isometric, 321, 324–326
linear, 304
norm, 308
normal, 321, 322, 326
null space, 318
positive, 330, 331
projection, 318
range, 318
self-adjoint, 320
shift, 325
translation, 328
unbounded, 311
unitary, 321, 326, 327, 335
operator adjoint, 316, 317
operator norm, xi, **308**
operator star-algebra, 317
optimal lower frame bound, **242**
optimal receiver, 79
optimal upper frame bound, **242**
order, x, xi
order n B-spline, 374
order n B-spline MRA coefficient sequence, 394
order n B-spline MRA scaling sequence, 394
order relation, 281
order-k moment estimate, **101**
ordered linear space, **212**
ordered pair, x
ordered set, 281, 282
ordinary coherence, **132**
ordinary transmissibility, 114
orthocomplemented lattice, 166
orthogonal, 19–21, 226, 229, 230, **267**, 267, 268, 320, 380
orthogonal basis, **230**
orthogonal in the Pythagorean sense, **280**
orthogonal in the sense of Birkhoff, **280**
orthogonal in the sense of James, **278**
orthogonality, 226, 278, 280
Birkhoff, 280
inner product space, 267
isosceles, 278
James, 278
Pythagorean, 280
orthomodular, 164
orthomodular lattice, 164
orthonomal, **267**
orthonormal, 227–233, 246, **267**
orthonormal bases, 220
orthonormal basis, 144, **230**, 234–239, 241, 390, 391
Orthonormal decomposition, 92
orthonormal expansion, 230, 234
orthonormal quadrature conditions, **349**
orthonormality, 227, 238, 246
orthornormal basis, 231
outcome, **165**
outcomes, 165
output, 14, 15, 35, 37
over estimate, 112
over-estimated, 128

PAM, 152
parabolic metric, 276
parallelogram, 262, 278
Parallelogram Law, 9
Parallelogram law, **263**, 266
parallelogram law, 261–263, 265
parametric, 63
Parseval frame, 230, 231, **242**, 242
Parseval frames, 220
Parseval's equation, 335
Parseval's Identity, 232, 234, 236, 242
Parseval's Theorem, 16
partition, 167, 168
partition of unity, 34, 166, 384, 385, 396, 397
Pascal's identity, 372
Pascal, Blaise, 163
pdf, **174**, 174
Peace Frame, **245**
Per Enflo, 224
Perfect reconstruction, **230**
periodic, 300, 343, 386
periodicity, 386
Periodogram, 105
phase, 113, 114, 120
Phase Shift Keying, 149
Plancheral's formula, 45, 85
Plancherel's formula, 335
Poincaré, Henri, 180
point spectrum, **190**
pointwise addition, 209
Polar Identity, **259**, 260, 279
polar identity, 227
Polarization Identities, 262
Polarization Identity, 261, 263–265
polarization identity, 269, 279
pole, 365
poles, 49, 359
polynomial approximation, 73
Popper, Karl, 109
positive, 20, **212**, **330**
positive cone, **212**
positive definite, 20
positive part, **214**, 215–217
power mean, **292**, 292, 295
power rule, 389
power set, xi
power spectral density, **14**, 16, 44
power sum, 374
Power Sums, **300**
probability, **165**
probability density function, 3, 25, 34, 168, **174**, 174, 182
probability function, **164**
Probability integral transform, **25**
probability measure, 165
probability space, 3–5, 8, 9, 11, 17, 33, **165**, 165–169, **170**, 170, 174, 175, 182
probabilty space, 170
product rule, 195, 198, 377
project, 380
projection, **250**, **318**
projection operator, 318, 320
projection statistics
Additive Gaussian noise channel, 82
Additive noise channel, 81
Additive white Gaussian noise channel, 83
Additive white noise channel, 83
projections, 84
proper subset, x
proper superset, x
properties
σ-additive, 164, 170
σ-additivity, 164
absolute value, x
absolutely summable, 15, 354, 355
additive, 19, 164–167, 169, 255, 256, 265, 267, 268, 304, 307
additive identity, 12, 66, 210
additive inverse, 210
additivity, 19, 164, 165, 246, 259, 316
admissibility condition, 396
admissibility equation, 396
affine, 190, **283**, 314
algebra of sets, xi
AND, x

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

anti-symmetric, 13, 14, 48, 337
anti-symmetric , 48
antiautomorphic, 15, 250, 251, 256, 317
antitone, 166, 169
associates, 208
associative, 5, 205, 206, 208, 210, 289, 290, 306, 307, 326
asymptotically unbiased, 103
AWGN, 79
Bayesian, 63
biased, 98, 99, 112
bijection, 235
bijective, 235, 314
biorthogonal, 239, 240
Birkhoff orthogonality, 280
Boole's inequality, 167
Boolean, 165, 167
Boolean lattice, 167
bounded, 166, 225, 320, 331
bounded bijective, 238
Cartesian product, x
causal, **358**
characteristic function, x
closed support, 378
closure, 223
colored, 19
commutative, 12, 206, 208, 215, 306, 353, 354
compact, 191
compact support, 391
comparable, 281
complement, x
complex, 120
complex-valued, 114
concave, **282**, 282, **283**, 283, **284**, 291, 292
conjugate linear, 250, 317
conjugate quadrature filter condition, **361**
conjugate symmetric, 14, 15, 18, 48, 192, 255, 256, 264, 265, 268
conjugate symmetry, 19
conjugate-symmetric, 15
consistent, 96, 97, 99, 103
constant, 12, 43, 65
Continuity, 274
continuity, 175, 176
continuous, 25, 168, 175, 192, 259, 274, 290, 292, 306, 380, 386, 393
convergence, 387
convergence in probability, 20
convergent, 44, 225
converges, 387–389
converges absolutely, 192
convex, 5, 6, 274, 276, 282, **283**, 283, **284**, 284, 291, 292
correlated, 52, 110, 111
countably infinite, 191, 224
counting measure, xi
CQF condition, **361**, 363
critical point, **26**, 26, 28, 29
decreasing, 291, 390
dense, 223, 224
difference, x
differentiable, 26
differentiable function, 28, 29
distributes, 208
distributive, 12, 35–37, 70, 164–167, 210, 250–252, 255, 256, 317, 344, 355
domain, x
dyadic decomposition, 391
efficient, 85, 88, 89, 91
empty set, xi
equality by definition, x
equality relation, x
equivalent, 225, 237
ergodic in the mean, **44**, 44, 45
even, 348
excluded middle, 166, 167
exclusive OR, xi
existential quantifier, xi
exists, 44
false, x
for each, xi
Fredholm operator, 191, 192
Fredholm operators, 191
Gamma distribution, 182
Gaussian, 33
geodesically between, **282**
greatest lower bound, xi
hermitian, 250, 251, **320**
Hermitian symmetric, 71, 337
homogeneous, 5, 206, 230, 231, 234, 238, 240, 255, 256, 260, 264, 265, 267, 271, 277, 298, 304, 305, 307, 308
hypotenuse, 280
identity, 205, 208
identity operator, 196
if, xi
if and only if, xi
image, x
imaginary part, xi
implied by, xi
implies, xi
implies and is implied by, xi
impossible, 384
inclusive OR, xi
increasing, 291, 390
independence, 81, 84, 169
independent, 7, 31, 34, 85, **166**, 168, 182
indicator function, x
injective, 306
inner-product, xi
inside, 358
intersection, x
into, 235, 236
inverse, 205
invertible, 26, 168, 225, 242, 250
involutary, 317
involution, 250, 254
involutory, 15, 250–252
irreflexive ordering relation, xi
isometric, 231, 239, 240, 314, 323, 335
isomorphic, 235
Isosceles orthogonality, 278
isosceles orthogonality, 278–280
join, xi
kronecker delta function, 196
least upper bound, xi
least-squares, 123, 126
left distributive, 5, 130, 205, 206, 307
linear, 4, 5, 35, 36, 54, 55, 57, 110, 120, **304**, 304
linear independence, 226
linear space, 209
linear time invariant, 37, 111, 123, 127, 128, 381
linear time-invariant, 57
linearity, 5, 19, 21, 22, 27, 30, 31, 45, 51, 55–57, 66, 71, 95, 99, 100, 124, 195, 305, 370, 371, 389, 392
linearly dependent, **219**, 221–223
linearly independent,

COPYRIGHT © 2019 DANIEL J. GREENHOE
BY-NC-ND

219, 221–223, 226
lower bounded, 167
LTI, 14, 15, 35–38, 56–59, 111, 112, 123, 124, 127–129, 135, 381
maps to, x
meet, xi
metric, xi
Minimum phase, 365
minimum phase, 38, 357, 358
minimum variance unbiased estimator, **66**
modular, 164
monotone, 164, 166, 168
monotonically decreasing, 117
mutually exclusive, 167, 170
MVUE, **66**
no input noise, 129
no output noise, 129
non-contradiction, 166
non-isometric, 260
non-isotropic, 255, 256, 263, 265, 268
non-linear, 54, 55, 110, 126
non-negative, 19–21, 206, 255, 260, 263, 265, 268, 292, 308, 378, 384
non-positive, 258
non-stationary, 44
non-zero, 396
nondegenerate, 164, 166, 169, 170, 206, 271, 308
nonnegative, 164
nonparametric, 63
normal, 191, 320, **321**
normalized, 20, 21, 164, 166, 169
NOT, x
not ergodic in the mean, 44
not invertible, 26
not necessarily linear, 37, 51
not symmetric, 379
not total, 233
null space, x
only if, xi
operator norm, xi
order, x, xi
ordered pair, x
orthogonal, 19–21, 226, 230, **267**, 267, 268, 320, 380
orthogonality, 226, 278, 280
orthomodular, 164
orthonomal, **267**
orthonormal, 227, 228, 230–233, 246, **267**
orthonormality, 227, 238, 246
over estimate, 112
over-estimated, 128
parallelogram, 262, 278
parallelogram law, 265
parametric, 63
Parseval's Theorem, 16
partition of unity, 34, 166, 384, 385, 396, 397
periodic, 300, 343, 386
periodicity, 386
pointwise addition, 209
Polarization Identity, 265
positive, 20
positive definite, 20
power set, xi
proper subset, x
proper superset, x
pseudo-distributes, 208
Pythagorean orthogonality, 280
Pythagorean Theorem, 280
randomness, 173
range, x
real, 20, 274, 279, 346, 359
real part, xi
real-valued, 15, 16, 20, 70, 114, 320, 337, 344, 345, 348
reality condition, 336
reflexive ordering relation, xi
regular, 191
relation, x
relational and, x
reverse triangle inequality, 272
right distributive, 5, 205, 206, 307
ring, 71
ring of sets, xi
scalar commutative, 206
self adjoint, 19, 320
self-adjoint, 19, **320**, 320
separable, 224, 225, 236
set of algebras of sets, xi
set of rings of sets, xi
set of topologies, xi
space of linear transforms, 305
span, xi
spans, 221, 222
stability condition, 240, 242
stable, 358
stationary in correlation, 43
stationary in the 1st moment, 12
stationary in the 2nd moment, 12
stationary in the mean, 43, 44
Strang-Fix condition, 339
strictly concave, **284**
strictly convex, **283, 284**
strictly increasing, 292
strictly monotone, 168
strictly monotonic, 290, 292
strictly monotonic increasing, 119
strictly positive, 271
strong convergence, 224, 234
subadditive, 164, 206, 260, 271, 298, 308
submultiplicative, 206
subset, x
super set, x
symmetric, 13, 14, 48, 274, 337, 346, 378, 379, 386, 390
symmetric , 48
symmetric difference, x
symmetry, 267, 357
there exists, xi
tight frame, 243
time-invariant, **358**
topology of sets, xi
total, 223, 232, 233, 237, 238
translation invariant, 277
triangle inequality, 206, 281
triangle inquality, 271
true, x
unbiased, 66, 85, 86, 95–99, 129
uncorrelated, 19, 51–59, 80, 95, 96, 99, 110–112, 124, 128, 130, 133
under estimate, 112
under estimates, 112
under-estimated, 128
uniformly, 192
uniformly distributed, 25, 34
union, x
unique, 26, 210, 221, 224, 383
unit area, 378
unit length, 325, 327
unitary, 326, 335
universal quantifier, xi

BY-NC-ND
COPYRIGHT © 2019 DANIEL J. GREENHOE

upper bounded, 164
vanishing 0th moment, 396
vector norm, xi
von Neumann-Jordan condition, 263
white, **19**, 19
wide sense stationary, **12**, 12–14, 37, **43**, 43, 44, 52, 95, 96, 111
wide-sense stationary, 128
WSS, **12**, 13–15, 36–38, **43**, 43, 51, 53–59, 95, 96, 99, 111, 112, 124, 128
zero at −1, 396
zero measurement error, 111
zero measurement noise, 111
zero-mean, 7, 51–59, 124
PSD, **14**
pseudo-distributes, 208
PSF, 339, 340
PSK, 149
pstricks, vi
Pulse Amplitude Modulation, 152
Pythagorean orthogonality, 278, **280**, 280
Pythagorean Theorem, 226, 227, 229, 231, 238, **268**, 268, 280
Pythagorean theorem, 226

QAM, 147
quadratic discriminate inequality, **258**
Quadratic Equation, 131
quadratic equation, 258
Quadratic form, **198**
quadratic form, 71, 198, 199
Quadrature Amplitude Modulation, 147
quantization noise, 111
quantum probability, **164**, 165
quotes
Abel, 298
Abel, Niels Henrik, 403
Aristotle, 11, 17
Bak, Per, 123
Bohr, Harald, 289
Cardano, Gerolamo, 205
Descartes, René, ix
Fourier, Joseph, 333
Hardy, G.H., 289
Housman, Alfred Edward, vii
Kaneyoshi, Urabe, 403
Kenko, Yoshida, 403
Leibniz, Gottfried, ix, 303
Lippmann, Bernard A., 180
Machiavelli, Niccolò, 403
Pascal, Blaise, 163
Poincaré, Henri, 180
Popper, Karl, 109
Russull, Bertrand, vii
Stravinsky, Igor, vii
Ulam, Stanislaus M., 314
Quotient Rule, 130, 131
quotient rule, 389
quotient structures, 281

random process, **17**, 17, 19, 20, 44
random sequence, **11**, 11–14, 35, 39, 110
random variable, 3–8, **17**, 17, 25, 26, 28, 29, 31, 33, 168, 173, **174**, 174, 175, 182
randomness, 173
range, x
range space, 316
rational expression, 38, 49
rationalizing factor, 122
rationalizing the denominator, 122
Rayleigh distribution, **185**, 185
real, 20, 274, 279, 346, 359
real interval, 175
real linear space, **208**
real numbers, 273
real part, xi, **251**
real-valued, 15, 16, 20, 70, 114, 320, 337, 344, 345, 348
reality condition, 336
Rectangular pulse, 341
rectangular pulse, **340**, 382
recursively, 75
reflection, 314
reflexive ordering relation, xi
regular, 191
relation, x, 26, 303
relational and, x
relations, xi
order relation, 281
residual spectrum, **190**
resolvent, **249**
resolvent set, **190**, 191
Reverse Triangle Inequality, **272**, 274
reverse triangle inequality, 272
Rice distribution, **185**, 185
Riesz bases, 220
Riesz basis, **237**, 237–241, 382, 390, 391
Riesz sequence, **237**, 239, 393
Riesz space, **212**, 212, 214–217
Riesz-Fischer Theorem, **234**
Riesz-Fischer Thm., 235, 236
right distributive, 5, 205, 206, 307
right inverse, 140
ring, 71, **205**, 205, 206
absolute value, 206
commutative, 205
modulus, 206
ring of complex square $n \times n$ matrices, 317
ring of sets, xi
Robinson's Energy Delay Theorem, **357**
rotation matrix, 330
Russull, Bertrand, vii

sample variance, **98**
sampling, 4
scalar commutative, 206
scalar product, **255**
scalars, **208**
scaling coefficient sequence, 396
scaling function, **115**, 393, 394
scaling parameter, **115**
Scaling transfer function estimate, **115**, 115
Schauder bases, 220, 225
Schauder basis, **224**, 224, 225, 230, 236
self adjoint, 19, 320
self-adjoint, 19, **250**, **320**, 320
semilinear, 251
separable, 224, 225, 236
separable Hilbert space, 235, 237, 239, 353
separable Hilbert spaces, 235
sequence, 11, 34, 273, **353**, 394
sequences, 361
set, 165, 207, 282
set indicator, 367
set indicator function, 341
set of algebras of sets, xi
set of rings of sets, xi
set of topologies, xi
set projection operators, **157**
sets
open ball, 274
real numbers, 273
shift operator, 325
shift relation, 340, 341
sigma-algebra, 165
Signal matching, 92
signal to noise ratio, 92
signal-to-noise ratio, 111
sinc, 340, 341
sine sweep, 109
single coin toss, **168**, 170

COPYRIGHT © 2019 DANIEL J. GREENHOE

Smith-Barnwell filter, **361**
SNR, 111
source code, 399
space, 165
inner product, 255
linear, 207
metric, 274
normed vector, 271
vector, 207
space of all absolutely square Lebesgue integrable functions, 353
space of all absolutely square summable sequences, **353**
space of linear transforms, 305
span, xi, 223, 225
spans, **219**, 221, 222
spectral power, 16, **44**, 111
spectral radius, **249**
Spectral Theorem, **191**
spectrum, **190**, 191, **249**
stability, 358
stability condition, 240, 242
stable, 358
standard othornormal basis, 231
star-algebra, **250**, 250, 317
star-algebras, 316, 317
state , 75
stationary in correlation, 43
stationary in the 1st moment, 12
stationary in the 2nd moment, 12
stationary in the mean, 43, 44
statistical average, 76
statistics, 142
steepest descent, 75, 76
step function, **367**, 367–371
step size, 75
Stifel formual, 392
Stifel formula, 372
stochastic, 69
Strang-Fix condition, **339**, 339
Stravinsky, Igor, vii
strictly concave, **284**
strictly convex, **283**, 283, **284**
strictly increasing, 292
strictly monotone, 168
strictly monotonic, 290, 292
strictly monotonic increasing, 119
strictly positive, 271, 277
strong convergence, 224, 234
structured, 141
structures
C^* algebra, **253**
C^*-algebra, 254
$*$-algebra, **12**, **15**, **35–37**, **250**, 250–252, 317, **344**
N-tuple, 209, 271
*-algebra, 70
*-algebras, 317
σ-algebra, 165, 170, 172
absolute value, 216, 217
additive identity element, 209, 210
additive inverse, 210
adjoint, 317
algebra, **206**, 207, **249**, 249, 250
algebraic, 207
ball, 282
Banach space, 224, 225
basis, 20, 22, 221–223, 230, 384
basis sequence, 383
bijection, 235
bijective, 235
bilinear functionals, 262
Boolean algebra, 170
Boolean lattice, 164, 167
Borel algebra on $\mathbb{R}$, 170
Borel sets, 333
bounded lattice, 167
bounded linear operators, 313, 315, 316, 318, 319, 321, 322, 324–327
C star algebra, **253**
calculus, 207
closed interval, **281**, **282**, 282
coefficient functionals, 224
coefficients, 359
communication system, 126
communications additive noise model, 123
Communications additive noise system, 55
commutative ring, 205, **206**, 206
complete metric space, 223
complete set, 223
complex linear space, **208**
conditional probability, 167
conditional probability of x given y, **166**
conjugate pairs, 359
conjugate quadrature filter, 361
Conjugate quadrature filters, 361
continuous spectrum, **190**
convex function, 283, 284
convex set, **282**, 282–284
convolution operator, 353, 354
coordinates, **221**
CQF, 361, 362
critical point, 26
differentiable function, 26
discrete-time Fourier transform, 349
distributive pseudo-complemented lattice, 164
Double coin toss, 168
eigen-system, 21
eigenspace, **189**
epigraph, **283**
Euclidean space, 282
even/odd die probability space, 169
even/odd die toss, **169**
event, **165**, 165, 166
expansion, **221**
field, 205, **206**, 207, 209, 353
field of complex numbers, 317
field of probability, 164
filter banks, 361
finite modular lattice, 164
Fourier Transform, 333
frame, 240, **242**, 242, 246
frames, 220
function, 26, 173, 333, 368
functional, 255, 284, 317
geometry, 207
group, **205**, 205
half-open interval, **281**, **282**
Hamel bases, 220
Hamel basis, **221**, 221, 223
high-pass filter, 361
Hilbert space, 230, 232, 234, 236, 237, 240, 242, 316, 317, 320–322, 327, 333
hypograph, **283**
identity, 208
identity element, 303
image set, 305, 320, 322, 327
infinite sum, 219
inner product, 207, 261
inner product space, 8, 9, 207, 226–229, 237, **255**, 256, 259, 260, 263, 267, 268, 278, 380
inner-product space, 278, 280

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

integral domain, 205
interval, 281, 282
intervals on lattices, **281**
intervals on ordered sets, **281**
inverse, 208
isometry, 323
lattice, 281, 282
lattice with negation, 164
Laurent series, 354
Lebesgue square-integrable functions, 333
linear basis, **221**
linear combination, **219**
linear combinations, 220
linear operator, 238, 240
linear operators, 236
linear ops., 236, 240
linear space, 207, **208**, 208, 219, 221, 249, 255, 261, 271, 277, 281–284, 303
linear spaces, 303
linear subspace, **208**, 219
low-pass filter, 361
matrix algebra, 70
measurable space, 165
measure, 165
measure space, 165, 166
measurement additive noise model, 123
Measurement additive noise system, 55
measurement system, 126
metric, 207, 274
metric linear space, 207, 273, 274, 276
metric space, 207, 274, 281, 282
MRA, 394
MRA system, 394, 396
Multiresolution Analysis, 391
multiresolution analysis, 382, 393
negative part, 215–217
norm, 207, 257, 259–261, 271, 273, 274, 277
normalized tight frame, **242**
normed algebra, 253, 254
normed linear space, 207, 261, 268, 271–274, 276, 278, 280
normed linear spaces, 313, 323
normed space of linear operators, **308**
null space, 189, 306, 320, 327
O_6 lattice, 164
open ball, 274
open interval, **281, 282**
operator, 207
ordered set, 281, 282
orthocomplemented lattice, 166
orthogonal basis, **230**
orthomodular lattice, 164
orthonormal bases, 220
orthonormal basis, **230**, 234–239, 241, 390, 391
orthornormal basis, 231
outcome, **165**
Parseval frame, **242**
Parseval frames, 220
Parseval's equation, 335
partition, 167, 168
Plancherel's formula, 335
point spectrum, **190**
pole, 365
poles, 359
positive part, 215–217
probability space, 3–5, 8, 9, 11, 17, 33, **165**, 165–169, **170**, 170, 174, 175, 182
probabilty space, 170
projection operator, 320
quotient structures, 281
random variable, 174
real interval, 175
real linear space, **208**
relation, 26
residual spectrum, **190**
resolvent, **249**
resolvent set, **190**, 191
Riesz bases, 220
Riesz basis, 237–241, 382, 390, 391
Riesz sequence, 239, 393
Riesz space, 212, 214–217
ring, **205**, 205, 206
ring of complex square $n \times n$ matrices, 317
scalars, **208**
Schauder bases, 220, 225
Schauder basis, **224**, 224, 225, 230, 236
separable Hilbert space, 235, 237, 239, 353
separable Hilbert spaces, 235
sequence, 273
sequences, 361
set, 165, 207, 282
sigma-algebra, 165
space, 165
space of all absolutely square Lebesgue integrable functions, 353
space of all absolutely square summable sequences, **353**
span, 223, 225
spectral radius, **249**
spectrum, **190**, 191, **249**
standard othornormal basis, 231
star-algebra, **250**, 317
star-algebras, 316
support, 378
system, 14, 37, 51–59, 109, 111–115, 119, 123, 124, 126–130, 132
tight frame, **242**, 243, 245
tight frames, 220
topological dual space, 313
topological linear space, 207, 223
topology, 165, 207, 219
total set, 223
underlying set, **208**
unit vector, 239
unital $*$-algebra, 250
unital algebra, 249
vector, 221
vector space, 8, 207, **208**
vector space of random variables, 4
vectors, **208**, 281
weights, 284
zero, 365
zeros, 357, 359
subadditive, 164, 206, 260, 271, 298, 308
submultiplicative, 206
subset, x
sufficient statistic, 79
Sufficient statistic theorem, **79**
Sum, 32
sum of products, 27, 31, **167**, 175
summation, **289**
Summation around unit circle, 244
summation operator, 284
sup metric, 276
super set, x
support, 378
surjective, xi
symmetric, 13, 14, 48, 274, 337, 346, 378, 379, 386, 390

COPYRIGHT © 2019 DANIEL J. GREENHOE
BY-NC-ND

conjugate, 18
symmetric , 48
symmetric difference, x
symmetry, 267, 357
system, 14, 37, 51–59, 109, 111–115, 119, 123, 124, 126–130, 132
system identication, 109

taxi-cab metric, 276
The basis problem, 224
The Book Worm, 403
The Fourier Series Theorem, **232**
The usual norm, **273**
Theorem of Reversibility, **141**
theorems
Affine equations, **197**
affine equations, 71
Alternating Series Test, 390
B-spline dyadic decomposition, 391, 397
B-spline general form, 378
B-spline partition of unity, 384
B-spline recursion, 376
B-spline scaling coefficients, **394**
Battle-Lemarié orthogonalization, **241**
Bessel's Equality, **227**, 228, 229, 233
Bessel's Inequality, 235
Bessel's inequality, **228**
Best Approximation Theorem, **228**, 228
Binomial Theorem, 5, 66, 100
binomial theorem, 395
Cauchy Schwartz inequality, 118, 120
Cauchy-Schwartz inequality, 9
Cauchy-Schwarz Inequality, 93
Cauchy-Schwarz inequality, 243
chain rule, 27
Chebyshev's inequality, 7
communications additive noise cross-correlation, 55
comparison test, 387
Convolution Theorem, 38, 382
convolution theorem, **336**, 341, **356**
CQF theorem, **361**, 364
CS Inequality, 246
dilation equation, 394
DTFT periodicity, **343**
Euler formulas, 340, 382
Fourier expansion, 231, **234**, 235, 236
Fourier series expansion, 232
Fundamental Theorem of Calculus, 368, 375
Fundamental theorem of calculus, 175, 177
Fundamental theorem of linear equations, 307
Gelfand-Mazur Theorem, **253**
General ML estimation, **84**
Generalized AM-GM inequality, 295
Generalized associative property, **289**
Generalized Parseval's Identity, 232
Generalized triangle inequality, 9
Geometric Series, **300**
Hermitian representation, **252**
induced norm, **260**
induction, 374, 375, 384
integral test, 388
inverse DTFT, **349**, 363
Inverse Fourier transform, **334**
Inverse probability integral transform, **25**
Jensen's Inequality, 6, **284**
Jensen's inequality, **6**, 6
Karhunen-Loève Expansion, **20**
Kolmogorov's inequality, **7**
l'Hôpital's rule, 118, 121, 122, 294, 387, 389, 393
Law of the Unconscious Statistician, **6**
Least squares, **72**
Leibnitz GPR, 364
Leibniz Integration Rule, 376
Linearity of E, **4**
Markov's inequality, **6**, 7
Mazur-Ulam theorem, 314
measurement additive noise cross-correlation, 55
Mercer's Theorem, 21, **192**, 192
Minimum mean square estimation, **70**
ML amplitude estimation, **85**
ML estimation of a function of a parameter, **89**
ML phase estimation, **88**
Neumann Expansion Theorem, 315
operator star-algebra, 317
orthonormal quadrature conditions, **349**
Parallelogram Law, 9
Parallelogram law, **263**
parallelogram law, 263
Parseval's Identity, 232, 234, 236, 242
Pascal's identity, 372
Perfect reconstruction, **230**
Plancheral's formula, 45, 85
Polar Identity, **259**, 260, 279
polar identity, 227
Polarization Identities, 262
Polarization Identity, 263–265
polarization identity, 269, 279
power rule, 389
Power Sums, **300**
Probability integral transform, **25**
PSF, 339, 340
Pythagorean Theorem, 226, 227, 229, 231, 238, **268**, 268
Pythagorean theorem, 226
Quadratic Equation, 131
Quadratic form, **198**
quadratic form, 71
Quotient Rule, 130, 131
quotient rule, 389
Reverse Triangle Inequality, **272**, 274
Riesz-Fischer Theorem, **234**
Riesz-Fischer Thm., 235, 236
Robinson's Energy Delay Theorem, **357**
shift relation, 340, 341
Spectral Theorem, **191**
Stifel formual, 392
Stifel formula, 372
Strang-Fix condition, **339**
Sufficient statistic theo-

BY-NC-ND COPYRIGHT © 2019 DANIEL J. GREENHOE

rem, **79**
sum of products, 27, 31, **167**, 175
Summation around unit circle, 244
The Fourier Series Theorem, **232**
Theorem of Reversibility, **141**
triangle inequality, **271**
Wiener-Hopf equations, **71**, 71
Wold's Theorem, 179
Young's Inequality, 297
there exists, xi
tight frame, **242**, 243, 245
tight frames, 220
time average, 44
time-invariant, **358**, 358
topological dual space, 313
topological linear space, 207, 223
topological space, 165, 273
topology, 165, 207, 219
topology of sets, xi
total, **223**, 223, 232, 233, 237, 238
Total Least Squares transfer function estimate, **115**
Total least squares transfer function estimate, 115
total set, 223
traditional probability, **164**, 165
transfer function estimate $\hat{H}_\kappa(\omega;\kappa)$, **115**
transfer function estimate $\hat{H}_c(\omega)$, **116**
transform
inverse Fourier, **334**
translation, 328
translation invariant, 277
translation operator, 339
Transmissibility, 114
transmissibility, 111, 113
transmissibility $\tilde{T}_{xy}(\omega)$, **111**
triangle, **341**
triangle inequality, 206, **271**, 271, 281, 308
triangle inequality for vectors, 268
triangle inquality, 271
triangle-inequality, 278
true, x
tuples in $\mathbb{F}^N$, **209**

Ulam, Stanislaus M., 314
unbiased, 66, 85, 86, 95–99, 129
uncorrelated, 19, 51–59, 80, 95, 96, 99, 110–112, 124, 128, 130, 133
under estimate, 112
under estimates, 112
under-estimated, 128
underlying set, **208**
uniform distribution, **179**, 179
uniformly, 192
uniformly distributed, 25, 34
union, x, 165
unique, 26, 210, 221, 224, 383
unit area, 378
unit length, 325, 327
unit vector, 239
unital, **249**
unital $*$-algebra, 250
unital algebra, 249
unitary, **326**, 326, 329, 334, 335
unitary Fourier Transform, **334**
unitary operator, 321, 326
universal quantifier, xi
upper bounded, 164
usual norm, 273
Utopia, vi

values
nth moment, **338**
Bessel's correction, 98
Cramér-Rao bound, 90
Cramér-Rao lower bound, 88
dimension, **221**
eigenvalue, **189**, 244
eigenvector, **189**
frame bound, 243, 245
frame bounds, 246
MAP estimate, 81
mean, 98
ML estimate, 81
rationalizing factor, 122
scaling function, **115**
scaling parameter, **115**
sufficient statistic, 79
Vandermonde matrix, 73
vanishing 0th moment, 396
vanishing moments, 338, 351
variance, **3**, 3, **12**, 12, 44, 95
vector, 221
vector addition, 195
vector lattice, **212**
vector norm, xi
vector space, 8, 207, **208**
vector space of random variables, 4
vectors, **208**, 281
von Neumann-Jordan condition, 263

wavelet, 391
weighted, 291
weights, 284
Welch Method 0% overlap, 105
Welch Method 50% overlap, 105
white, **19**, 19
whitening filter, 38, 48
wide sense stationary, **12**, 12–14, 37, **43**, 43, 44, 52, 95, 96, 111
wide-sense stationary, 128
Wiener-Hopf equations, **71**, 71
windowed auto-correlation estimate, **103**
Wold's Theorem, 179
Wronskian, 203
WSS, **12**, 12–15, 36–38, **43**, 43, 51, 53–59, 95, 96, 99, 111, 112, 124, 128

Young's Inequality, **296**, 297

Z transform, 395
z transform, 357
z-domain cross spectral density, **14**
z-domain power spectral density, **14**
Z-Transform, xii
Z-transform, 345
z-transform, **354**, 354, 362
Z-transform operator, 14
zero, 365
zero at -1, 346, 396
zero measurement error, 111
zero measurement noise, 111
zero-mean, 7, 51–59, 124
zeros, 49, 357, 359

COPYRIGHT © 2019 DANIEL J. GREENHOE
BY-NC-ND

License

This document is provided under the terms of the Creative Commons license CC BY-NC-ND 4.0. For an exact statement of the license, see

https://creativecommons.org/licenses/by-nc-nd/4.0/legalcode

The icon appearing throughout this document is based on one that was once at

https://creativecommons.org/

where it was stated, "Except where otherwise noted, content on this site is licensed under a Creative Commons Attribution 4.0 International license."

…last page …please stop reading …

Copyright © 2019 Daniel J. Greenhoe
BY-NC-ND