







# A Book Concerning Transforms

VERSION 0.10



Daniel J. Greenhoe







title: *A Book Concerning Transforms*  
 document type: book  
 series: *Signal Processing ABCs*  
 volume: 5  
 author: Daniel J. Greenhoe  
 version: VERSION 0.10  
 time stamp: 2019 December 10 (Tuesday) 11:10am UTC  
 copyright: Copyright © 2019 Daniel J. Greenhoe  
 license: [Creative Commons](#) license [CC BY-NC-ND 4.0](#)  
 typesetting engine: Xe<sub>La</sub>TeX  
 document url: <https://github.com/dgreenhoe/pdfs/blob/master/abcstat.pdf>  
<https://www.researchgate.net/project/Signal-Processing-ABCs>

This text was typeset using  $\text{Xe}\text{\LaTeX}$ , which is part of the  $\text{T}\text{\E}$ Xfamily of typesetting engines, which is arguably the greatest development since the Gutenberg Press. Graphics were rendered using the *pstricks* and related packages, and  $\text{\LaTeX}$  graphics support.

The main roman, *italic*, and **bold** font typefaces used are all from the *Heuristica* family of typefaces (based on the *Utopia* typeface, released by *Adobe Systems Incorporated*). The math font is XITS from the XITS font project. The font used in quotation boxes is adapted from *Zapf Chancery Medium Italic*, originally from URW++ Design and Development Incorporated. The font used for the text in the title is Adventor (similar to *Avant-Garde*) from the *T<sub>E</sub>X-Gyre Project*. The font used for the ISBN in the footer of individual pages is LIQUID CRYSTAL (*Liquid Crystal*) from *FontLab Studio*. The Latin handwriting font is *Lavi* from the *Free Software Foundation*.

The ship appearing throughout this text is loosely based on the *Golden Hind*, a sixteenth century English galleon famous for circumnavigating the globe.<sup>1</sup>



---

<sup>1</sup>  Paine (2000) page 63 ⟨Golden Hind⟩

*“Here, on the level sand,  
Between the sea and land,  
What shall I build or write  
Against the fall of night?”*



*“Tell me of runes to grave  
That hold the bursting wave,  
Or bastions to design  
For longer date than mine.”*

[Alfred Edward Housman](#), English poet (1859–1936) <sup>2</sup>



*“The uninitiated imagine that one must await inspiration in order to create. That is a mistake. I am far from saying that there is no such thing as inspiration; quite the opposite. It is found as a driving force in every kind of human activity, and is in no wise peculiar to artists. But that force is brought into action by an effort, and that effort is work. Just as appetite comes by eating so work brings inspiration, if inspiration is not discernible at the beginning.”*



[Igor Fyodorovich Stravinsky](#) (1882–1971), Russian-born composer <sup>3</sup>






*“As I think about acts of integrity and grace, I realise that there is nothing in my knowledge to compare with Frege's dedication to truth. His entire life's work was on the verge of completion, much of his work had been ignored to the benefit of men infinitely less capable, his second volume was about to be published, and upon finding that his fundamental assumption was in error, he responded with intellectual pleasure clearly submerging any feelings of personal disappointment. It was almost superhuman and a telling indication of that of which men are capable if their dedication is to creative work and knowledge instead of cruder efforts to dominate and be known.”*

[Bertrand Russell](#) (1872–1970), [British mathematician](#), in a 1962 November 23 letter to Dr. van Heijenoort. <sup>4</sup>



<sup>2</sup> quote:  [Housman \(1936\)](#) page 64 <“Smooth Between Sea and Land”>,  [Hardy \(1940\)](#) <section 7>  
image: <http://en.wikipedia.org/wiki/Image:Housman.jpg>

<sup>3</sup> quote:  [Ewen \(1961\)](#) page 408,  [Ewen \(1950\)](#)  
image: [http://en.wikipedia.org/wiki/Image:Igor\\_Stravinsky.jpg](http://en.wikipedia.org/wiki/Image:Igor_Stravinsky.jpg)

<sup>4</sup> quote:  [Heijenoort \(1967\)](#) page 127  
image: <http://www-history.mcs.st-andrews.ac.uk/PictDisplay/Russell.html>



“*regula XVI. Quae vero praesentem mentis attentionem non requirunt, etiamsi ad conclusionem necessaria sint, illa melius est per brevissimas notas designare quam per integras figuras: ita enim memoria non poterit falli, nec tamen interim cogitatio distrahetur ad haec retinenda, dum aliis deducendis incumbit.*”



“*Rule XVI. As for things which do not require the immediate attention of the mind, however necessary they may be for the conclusion, it is better to represent them by very concise symbols rather than by complete figures. It will thus be impossible for our memory to go wrong, and our mind will not be distracted by having to retain these while it is taken up with deducing other matters.*”

René Descartes (1596–1650), French philosopher and mathematician <sup>5</sup>



“*In signs one observes an advantage in discovery which is greatest when they express the exact nature of a thing briefly and, as it were, picture it; then indeed the labor of thought is wonderfully diminished.*”

Gottfried Leibniz (1646–1716), German mathematician, <sup>6</sup>

## Symbol list

symbol	description	
numbers:		
$\mathbb{Z}$	integers	$\dots, -3, -2, -1, 0, 1, 2, 3, \dots$
$\mathbb{W}$	whole numbers	$0, 1, 2, 3, \dots$

...continued on next page...

<sup>5</sup>quote: Descartes (1684a) ⟨*regula XVI*⟩, translation: Descartes (1684b) ⟨*rule XVI*⟩, image: Frans Hals (circa 1650), <http://en.wikipedia.org/wiki/Descartes>, public domain

<sup>6</sup>quote: Cajori (1993) ⟨paragraph 540⟩, image: [http://en.wikipedia.org/wiki/File:Gottfried\\_Wilhelm\\_von\\_Leibniz.jpg](http://en.wikipedia.org/wiki/File:Gottfried_Wilhelm_von_Leibniz.jpg), public domain

symbol	description	
$\mathbb{N}$	natural numbers	$1, 2, 3, \dots$
$\mathbb{Z}^{\leq 0}$	non-positive integers	$\dots, -3, -2, -1, 0$
$\mathbb{Z}^-$	negative integers	$\dots, -3, -2, -1$
$\mathbb{Z}_o$	odd integers	$\dots, -3, -1, 1, 3, \dots$
$\mathbb{Z}_e$	even integers	$\dots, -4, -2, 0, 2, 4, \dots$
$\mathbb{Q}$	rational numbers	$\frac{m}{n}$ with $m \in \mathbb{Z}$ and $n \in \mathbb{Z} \setminus 0$
$\mathbb{R}$	real numbers	completion of $\mathbb{Q}$
$\mathbb{R}^+$	non-negative real numbers	$[0, \infty)$
$\mathbb{R}^{\leq 0}$	non-positive real numbers	$(-\infty, 0]$
$\mathbb{R}^+$	positive real numbers	$(0, \infty)$
$\mathbb{R}^-$	negative real numbers	$(-\infty, 0)$
$\mathbb{R}^*$	extended real numbers	$\mathbb{R}^* \triangleq \mathbb{R} \cup \{-\infty, \infty\}$
$\mathbb{C}$	complex numbers	
$\mathbb{F}$	arbitrary field	(often either $\mathbb{R}$ or $\mathbb{C}$ )
$\infty$	positive infinity	
$-\infty$	negative infinity	
$\pi$	pi	$3.14159265 \dots$
relations:		
$\mathbb{R}$	relation	
$\oslash$	relational and	
$X \times Y$	Cartesian product of $X$ and $Y$	
$(\triangle, \nabla)$	ordered pair	
$ z $	absolute value of a complex number $z$	
$=$	equality relation	
$\triangleq$	equality by definition	
$\rightarrow$	maps to	
$\in$	is an element of	
$\notin$	is not an element of	
$\mathcal{D}(\mathbb{R})$	domain of a relation $\mathbb{R}$	
$\mathcal{I}(\mathbb{R})$	image of a relation $\mathbb{R}$	
$\mathcal{R}(\mathbb{R})$	range of a relation $\mathbb{R}$	
$\mathcal{N}(\mathbb{R})$	null space of a relation $\mathbb{R}$	
set relations:		
$\subseteq$	subset	
$\subsetneq$	proper subset	
$\supseteq$	super set	
$\supsetneq$	proper superset	
$\not\subseteq$	is not a subset of	
$\not\subsetneq$	is not a proper subset of	
operations on sets:		
$A \cup B$	set union	
$A \cap B$	set intersection	
$A \triangle B$	set symmetric difference	
$A \setminus B$	set difference	
$A^c$	set complement	
$ \cdot $	set order	
$\mathbb{1}_A(x)$	set indicator function or characteristic function	
logic:		
1	“true” condition	

...continued on next page...

symbol	description	
0	“false” condition	
$\neg$	logical NOT operation	
$\wedge$	logical AND operation	
$\vee$	logical inclusive OR operation	
$\oplus$	logical exclusive OR operation	
$\Rightarrow$	“implies”;	“only if”
$\Leftarrow$	“implied by”;	“if”
$\Leftrightarrow$	“if and only if”;	“implies and is implied by”
$\forall$	universal quantifier:	“for each”
$\exists$	existential quantifier:	“there exists”
order on sets:		
$\vee$	join or least upper bound	
$\wedge$	meet or greatest lower bound	
$\leq$	reflexive ordering relation	“less than or equal to”
$\geq$	reflexive ordering relation	“greater than or equal to”
$<$	irreflexive ordering relation	“less than”
$>$	irreflexive ordering relation	“greater than”
measures on sets:		
$ X $	order or counting measure of a set $X$	
distance spaces:		
$d$	metric or distance function	
linear spaces:		
$\ \cdot\ $	vector norm	
$\ \cdot\ $	operator norm	
$\langle \triangle   \nabla \rangle$	inner-product	
$\text{span}(V)$	span of a linear space $V$	
algebras:		
$\Re$	real part of an element in a $*$ -algebra	
$\Im$	imaginary part of an element in a $*$ -algebra	
set structures:		
$\mathcal{T}$	a topology of sets	
$\mathcal{R}$	a ring of sets	
$\mathcal{A}$	an algebra of sets	
$\emptyset$	empty set	
$2^X$	power set on a set $X$	
sets of set structures:		
$\mathcal{T}(X)$	set of topologies on a set $X$	
$\mathcal{R}(X)$	set of rings of sets on a set $X$	
$\mathcal{A}(X)$	set of algebras of sets on a set $X$	
classes of relations/functions/operators:		
$2^{XY}$	set of <i>relations</i> from $X$ to $Y$	
$Y^X$	set of <i>functions</i> from $X$ to $Y$	
$S_j(X, Y)$	set of <i>surjective</i> functions from $X$ to $Y$	
$I_j(X, Y)$	set of <i>injective</i> functions from $X$ to $Y$	
$B_j(X, Y)$	set of <i>bijective</i> functions from $X$ to $Y$	
$B(\mathbf{X}, \mathbf{Y})$	set of <i>bounded</i> functions/operators from $\mathbf{X}$ to $\mathbf{Y}$	
$\mathcal{L}(\mathbf{X}, \mathbf{Y})$	set of <i>linear bounded</i> functions/operators from $\mathbf{X}$ to $\mathbf{Y}$	
$C(\mathbf{X}, \mathbf{Y})$	set of <i>continuous</i> functions/operators from $\mathbf{X}$ to $\mathbf{Y}$	
specific transforms/operators:		

...continued on next page...

symbol	description
$\tilde{\mathbf{F}}$	<i>Fourier Transform operator</i> (Definition 4.2 page 16)
$\hat{\mathbf{F}}$	<i>Fourier Series operator</i> (Definition 7.1 page 51)
$\check{\mathbf{F}}$	<i>Discrete Time Fourier Series operator</i> (Definition 6.1 page 41)
$\mathbf{Z}$	<i>Z-Transform operator</i> (Definition 5.4 page 28)
$\tilde{f}(\omega)$	<i>Fourier Transform of a function <math>f(x) \in L^2_{\mathbb{R}}</math></i>
$\check{x}(\omega)$	<i>Discrete Time Fourier Transform of a sequence <math>(x_n \in \mathbb{C})_{n \in \mathbb{Z}}</math></i>
$\check{x}(z)$	<i>Z-Transform of a sequence <math>(x_n \in \mathbb{C})_{n \in \mathbb{Z}}</math></i>

$P$ , 111  
 $T_n(x)$ , 96  
 $\mathbb{C}$ , 187  
 $\mathbb{Q}$ , 192  
 $\mathbb{R}$ , 187  
 $1$ , 188  
 $D_n$ , 102  
 $J_n$ , 111  
 $K_n$ , 109  
 $V_n$ , 111  
 $\tilde{\delta}_n$ , 160  
 $\tilde{S}_{ff}(z)$ , 179  
 $\tilde{S}_{fg}(z)$ , 179  
 $(A, \|\cdot\|, *)$ , 117  
 $(\kappa_n)_{n \in \mathbb{Z}}$ , 109  
 $\|\cdot\|$ , 124

$L^2_{(\mathbb{R}, \mathcal{B}, \mu)}$ , 69  
 $L^2_{\mathbb{R}}$ , 69  
 $PW^2_{\sigma}$ , 198  
 $\frac{d}{dx}$ , 10  
 $\exp(ix)$ , 80  
 $\tan$ , 85  
 $\star$ , 27  
 $\mathcal{L}(\mathbb{C}, \mathbb{C})$ , 198  
 $\cos$ , 85  
 $\cos(x)$ , 75  
 $\sin$ , 85  
 $\sin(x)$ , 75  
 $\tilde{S}_{ff}(\omega)$ , 179  
 $\tilde{S}_{fg}(\omega)$ , 179  
 $\tilde{F}$ , 16

$w_N$ , 103  
 $X$ , 187  
 $Y$ , 187  
 $\mathbb{C}^{\mathbb{C}}$ , 187  
 $\mathbb{R}^{\mathbb{R}}$ , 187  
 $D^*$ , 190  
 $D_{\alpha}$ , 188  
 $I_m$ , 115  
 $I$ , 120  
 $L$ , 11  
 $R_e$ , 115  
 $T^*$ , 190  
 $T$ , 188  
 $T_{\tau}$ , 188  
 $Z$ , 28  
 $\text{span}$ , 149

$Y^X$ , 187  
 $T_n$ , 96  
 $*$ , 114  
 $\hat{F}^{-1}$ , 52  
 $\hat{F}^*$ , 53  
 $\hat{F}$ , 51  
 $\|\cdot\|$ , 125  
 $\star$ , 12, 19  
 $B(X, Y)$ , 128  
 $Y^X$ , 121  
 $\rho$ , 113, 210  
 $\sigma_c$ , 210  
 $\sigma_p$ , 210  
 $\sigma_r$ , 210  
 $\sigma$ , 113, 210  
 $r$ , 113



Title page	v
Typesetting	vi
Quotes	vii
Symbol list	ix
Symbol index	xiii
Contents	xv
<b>1 Analyses and Transforms</b>	<b>1</b>
1.1 Abstract spaces	1
1.2 Lattice of subspaces	2
1.3 Analyses	3
1.4 Transform	3
1.5 Properties of subspace order structures	4
1.6 Operator inducing analyses	5
1.7 Wavelet analyses	6
<b>2 Taylor Expansions (Transforms)</b>	<b>9</b>
2.1 Introduction	9
2.2 Taylor Expansion	10
<b>3 Laplace Transform</b>	<b>11</b>
3.1 Definition	11
3.2 Shift relations	11
3.3 Convolution relations	12
3.4 Calculus relations	13
<b>4 Fourier Transform</b>	<b>15</b>
4.1 Introduction	15
4.2 Definitions	16
4.3 Operator properties	17
4.4 Shift relations	18
4.5 Convolution relations	19
4.6 Calculus relations	19
4.7 Real valued functions	20
4.8 Moment properties	20
4.9 Examples	23
<b>5 Z Transform</b>	<b>27</b>
5.1 Convolution operator	27
5.2 Z-transform	28
5.3 From z-domain back to time-domain	30
5.4 Zero locations	31
5.5 Pole locations	32
5.6 Mirroring for real coefficients	33
5.7 Rational polynomial operators	34
5.8 Filter Banks	34

5.9 Inverting non-minimum phase filters	39
<b>6 Discrete Time Fourier Transform</b>	<b>41</b>
6.1 Definition	41
6.2 Properties	41
6.3 Derivatives	49
<b>7 Fourier Series</b>	<b>51</b>
7.1 Definition	51
7.2 Inverse Fourier Series operator	52
7.3 Fourier series for compactly supported functions	54
<b>8 Fast Wavelet Transform (FWT)</b>	<b>55</b>
<b>9 KL Expansion (Transform)</b>	<b>59</b>
9.1 Definitions	59
9.2 Properties	59
9.3 Quasi-basis	63
<b>Appendices</b>	<b>65</b>
<b>A Algebraic structures</b>	<b>67</b>
<b>B Calculus</b>	<b>69</b>
<b>C Trigonometric Functions</b>	<b>73</b>
C.1 Definition Candidates	73
C.2 Definitions	75
C.3 Basic properties	75
C.4 The complex exponential	80
C.5 Trigonometric Identities	82
C.6 Planar Geometry	88
C.7 The power of the exponential	88
<b>D Trigonometric Polynomials</b>	<b>91</b>
D.1 Trigonometric expansion	91
D.2 Trigonometric reduction	96
D.3 Spectral Factorization	100
D.4 Dirichlet Kernel	101
D.5 Trigonometric summations	105
D.6 Summability Kernels	109
<b>E Normed Algebras</b>	<b>113</b>
E.1 Algebras	113
E.2 Star-Algebras	114
E.3 Normed Algebras	117
E.4 C* Algebras	117
<b>F Operators on Linear Spaces</b>	<b>119</b>
F.1 Operators on linear spaces	119
F.1.1 Operator Algebra	119
F.1.2 Linear operators	121
F.2 Operators on Normed linear spaces	124
F.2.1 Operator norm	124
F.2.2 Bounded linear operators	128
F.2.3 Adjoints on normed linear spaces	129
F.2.4 More properties	130
F.3 Operators on Inner product spaces	132
F.3.1 General Results	132
F.3.2 Operator adjoint	133
F.4 Special Classes of Operators	135
F.4.1 Projection operators	135
F.4.2 Self Adjoint Operators	137



F.4.3	Normal Operators	138
F.4.4	Isometric operators	140
F.4.5	Unitary operators	143
F.5	Operator order	148
<b>G</b>	<b>Linear Combinations</b>	<b>149</b>
G.1	Linear combinations in linear spaces	149
G.2	Bases in topological linear spaces	153
G.3	Schauder bases in Banach spaces	154
G.4	Linear combinations in inner product spaces	156
G.5	Orthonormal bases in Hilbert spaces	160
G.6	Riesz bases in Hilbert spaces	167
G.7	Frames in Hilbert spaces	172
<b>H</b>	<b>Power Spectrum Functions</b>	<b>179</b>
H.1	Correlation	179
H.2	Power Spectrum	179
<b>I</b>	<b>Transversal Operators</b>	<b>187</b>
I.1	Families of Functions	187
I.2	Definitions and algebraic properties	188
I.3	Linear space properties	189
I.4	Inner product space properties	190
I.5	Normed linear space properties	191
I.6	Fourier transform properties	194
I.7	Examples	198
<b>J</b>	<b>Continuous Random Processes</b>	<b>201</b>
J.1	Definitions	201
J.2	Properties	202
<b>K</b>	<b>Random Sequences</b>	<b>203</b>
K.1	Definitions	203
K.2	Properties	204
K.3	Wide Sense Stationary processes	204
K.4	Spectral density	207
K.5	Spectral Power	208
<b>L</b>	<b>Spectral Theory</b>	<b>209</b>
L.1	Operator Spectrum	209
L.2	Fredholm kernels	211
<b>Back Matter</b>		<b>213</b>
References		214
Reference Index		235
Subject Index		239
License		249
End of document		251



# CHAPTER 1

## ANALYSES AND TRANSFORMS



*“The analytical equations, unknown to the ancient geometers, which Descartes was the first to introduce into the study of curves and surfaces, ...they extend to all general phenomena. There cannot be a language more universal and more simple, more free from errors and from obscurities, ...mathematical analysis is as extensive as nature itself; it defines all perceptible relations, measures times, spaces, forces, temperatures ; this difficult science is formed slowly, but it preserves every principle which it has once acquired; it grows and strengthens itself incessantly in the midst of the many variations and errors of the human mind. Its chief attribute is clearness; it has no marks to express confused notions. It brings together phenomena the most diverse, and discovers the hidden analogies which unite them. ”*

Joseph Fourier (1768–1830) <sup>1</sup>

### 1.1 Abstract spaces

The **abstract space** was introduced by Maurice Fréchet in his 1906 Ph.D. thesis.<sup>2</sup> An *abstract space* in mathematics does not really have a rigorous definition; but in general it is a set together with some other unifying structure. Examples of spaces include *topological spaces*, *metric spaces*, and *linear spaces* (*vector spaces*).




<sup>1</sup> quote:  Fourier (1878) pages 7–8 (Preliminary Discourse)

image: <http://en.wikipedia.org/wiki/File:Fourier2.jpg>, public domain

<sup>2</sup>  Fréchet (1906),  Fréchet (1928). “A collection of these abstract elements will be called an abstract set. If to this set there is added some rule of association of these elements, or some relation between them, the set will be called an abstract space.”—Maurice Fréchet

## 1.2 Lattice of subspaces

An abstract space can be decomposed into one or more *subspaces*. Roughly speaking, a subspace of an abstract space is simply a subset the abstract space that has the same properties of that abstract space. The subspaces can be ordered under the ordering relation  $\subseteq$  (subset or equal to relation) to form a *lattice*.





Figure 1.1: lattice of topologies on  $X \triangleq \{x, y, z\}$  (Example 1.1 page 2)

**Example 1.1.** <sup>3</sup> The power set  $2^X$  is a *topology* on the set  $X$ . But there are also 28 other topologies on  $\{x, y, z\}$ , and these are all *subspaces* of  $2^{\{x,y,z\}}$ . Let a given topology in  $\mathcal{T}(\{x, y, z\})$  be represented by a Hasse diagram as illustrated to the right, where a circle present means the indicated set is in the topology, and a circle absent means the indicated set is not in the topology. <sup>4</sup> The lattice of the 29 topologies  $(\mathcal{T}(\{x, y, z\}), \cup, \cap; \subseteq)$  is illustrated in Figure 1.1 (page 2). The lattice of these 29 topologies is *non-distributive* (it contains the *N5 lattice*). The five topologies illustrated by red shaded nodes are also *algebras of sets*.



**Example 1.2.** The power set  $2^X$  is an *algebra of sets* on the set  $X$ . But there are also 14 other algebras of sets on  $\{w, x, y, z\}$ , and these are all *subspaces* of  $2^{\{w,x,y,z\}}$ . The *lattice of algebras of sets* on  $\{w, x, y, z\}$  is illustrated in Figure 1.2 (page 3).

A *linear subspace* is a subspace of a *linear space* (*vector space*). Linear subspaces have some special properties: Every linear subspace contains the additive identity zero vector, and every linear subspace is *convex*.

<sup>4</sup> [Isham \(1999\)](#) page 44, [Isham \(1989\)](#) page 1516, [Steiner \(1966\)](#) page 386





Figure 1.2: lattice of *algebras of sets* on  $\{w, x, y, z\}$  (Example 1.2 page 2)

*Example 1.3.* The 3-dimensional Euclidean space  $\mathbb{R}^3$  contains the 2-dimensional  $xy$ -plane and  $xz$ -plane subspaces, which in turn both contain the 1-dimensional  $x$ -axis subspace. These subspaces are illustrated in the figure to the right and in Figure 1.3 (page 4).



## 1.3 Analyses

An **analysis** of a space  $X$  is any lattice of subspaces of  $X$ . The partial or complete reconstruction of  $X$  from this set is a **synthesis**.<sup>5</sup>


*Example 1.4.* The lattices of subspaces illustrated in Figure 1.4 (page 4) are all *analyses* of  $\mathbb{R}^3$ .

## 1.4 Transform

**Definition 1.1.** A **transform** on a space  $X$  is a sequence of projection operators that induces an ANALYSIS on  $X$ .

Section 1.3 defined an **analysis** of a space  $X$  as is any lattice of subspaces of  $X$ . In like manner, an **analysis** of a function  $f(x)$  with respect to a transform  $T$  is simply the transform  $T$  of  $f$  ( $Tf$ ). Such

<sup>5</sup>The word *analysis* comes from the Greek word ἀνάλυσις, meaning “dissolution” (Perschbacher (1990) page 23 (entry 359)), which in turn means “the resolution or separation into component parts” (Black et al. (2009), <http://dictionary.reference.com/browse/dissolution>)

Figure 1.3: lattice of subspaces of  $\mathbb{R}^3$  (Example 1.3 page 3)linearly ordered analysis of  $\mathbb{R}^3$     M-3 analysis of  $\mathbb{R}^3$     wavelet-like analysis of  $\mathbb{R}^3$ Figure 1.4: some analyses of  $\mathbb{R}^3$  (Example 1.4 page 3)

an analysis or transform is often represented as the sequence of coefficients  $(\lambda_n)$  multiplying the basis vectors  $(\psi_n(x))$  such that

$$f(x) = \mathbf{T}f(x) = \sum_{n \in \mathbb{Z}} \lambda_n \psi_n(x)$$

*Example 1.5.* A *Fourier analysis* is a sequence of subspaces with sinusoidal bases. Examples of subspaces in a Fourier analysis include  $V_1 = \text{span}\{e^{ix}\}$ ,  $V_{2,3} = \text{span}\{e^{i2.3x}\}$ ,  $V_{\sqrt{2}} = \text{span}\{e^{i\sqrt{2}x}\}$ , etc. A **transform** is a set of *projection operators* that maps a family of functions (e.g.  $L^2_{\mathbb{R}}$ ) into an analysis. The *Fourier transform* for Fourier Analysis is (Definition 4.2 page 16)

$$[\tilde{\mathbf{F}}f](\omega) \triangleq \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} f(x) e^{-i\omega x} dx$$

## 1.5 Properties of subspace order structures

The ordered set of all linear subspaces of a *Hilbert space* is an *orthomodular lattice*. Orthomodular lattices (and hence Hilbert subspaces) have some special properties (next theorem). One is that they satisfy *de Morgan's law*.

**Theorem 1.1.** <sup>6</sup> Let  $L \triangleq (X, \vee, \wedge, 0, 1; \leq)$  be an algebraic structure.

<b>T H M</b>	$L \text{ is an ORTHOMODULAR LATTICE} \} \Rightarrow$	$\left\{ \begin{array}{l} 1. (x \vee y)^\perp \\ 2. (x \wedge y)^\perp \\ 3. (z^\perp \wedge y^\perp)^\perp \vee x \\ 4. x \wedge (x \vee y) \\ 5. x \vee (y \wedge y^\perp) \end{array} \right.$	$\left\{ \begin{array}{l} = x^\perp \wedge y^\perp \\ = x^\perp \vee y^\perp \\ = (x \vee y) \vee z \\ = x \\ = x \end{array} \right.$	$\forall x, y \in X$	$(\text{DE MORGAN})$	$\text{and}$

<sup>6</sup> Beran (1985) pages 30–33, Birkhoff and Neumann (1936) page 830 (L74), Beran (1976) pages 251–252






Figure 1.5: some common transforms


Most transforms have a very simple  $M$ - $n$  order structure, as illustrated to the right and in Figure 1.5 page 5. The  $M$ - $n$  lattices for  $n \geq 3$  are *modular* but not *distributive*. Analyses typically have one subspace that is a *scaling* subspace; and this subspace is often simply a family of constants (as is the case with *Fourier Analysis*). There is one notable exception to this—MRA induced *wavelet analysis*.



## 1.6 Operator inducing analyses



An *analysis* is often defined in terms of a small number (e.g. 2) operators. Two such operators are the *translation operator* (Definition 1.3 page 188).



**Example 1.6.** In *Fourier analysis*, continuous dilations (Definition 1.3 page 188) of the *complex exponential* form a *basis* (Definition G.7 page 154) for the *space of square integrable functions*  $L^2_{\mathbb{R}}$  (Definition B.1 page 69) such that  $L^2_{\mathbb{R}} = \text{span} \{D_{\omega} e^{ix} \mid \omega \in \mathbb{R}\}$ .

**Example 1.7.** In *Fourier series analysis* (Theorem 7.1 page 52), discrete dilations of the complex exponential form a basis for  $L^2_{\mathbb{R}}(0 : 2\pi)$  such that  $L^2_{\mathbb{R}}(0 : 2\pi) = \text{span} \{D_j e^{ix} \mid j \in \mathbb{Z}\}$ .





Figure 1.6: some wavelet transforms

## 1.7 Wavelet analyses

The term “wavelet” comes from the French word “*ondelette*”, meaning “small wave”. And in essence, wavelets are “small waves” (as opposed to the “long waves” of Fourier analysis) that form a basis for the Hilbert space  $L^2_{\mathbb{R}}$ .<sup>8</sup>

A **special characteristic** of wavelet analysis is that there is not just one scaling subspace, (as is with the case of Fourier and several other analyses), but an entire sequence of scaling subspaces (Figure 1.6 page 6). These scaling subspaces are *linearly ordered* with respect to the ordering relation  $\subseteq$ . In wavelet theory, this structure is called a *multiresolution analysis*, or *MRA*. The MRA was introduced by Stéphane G. Mallat in 1989. The concept of a scaling space was perhaps first introduced by Taizo Iijima in 1959 in Japan, and later as the *Gaussian Pyramid* by Burt and Adelson in the 1980s in the West.<sup>9</sup>



The MRA has become the dominate wavelet construction method. Moreover, P.G. Lemarié has proved that all wavelets with *compact support* are generated by an MRA.<sup>11</sup>


<sup>8</sup> Strang and Nguyen (1996) page ix Atkinson and Han (2009) page 191

<sup>10</sup> Mallat (1989) page 70 Iijima (1959) Burt and Adelson (1983) Adelson and Burt (1981) Lindeberg (1993) Alvarez et al. (1993) Guichard et al. (2012) Weickert (1999) (historical survey)


<sup>11</sup> Lemarié (1990), Mallat (1999) page 240



A **second special characteristic** of wavelet analysis is that it's order structure with respect to the  $\subseteq$  relation is not a simple  $M_n$  lattice (as is with the case of Fourier and several other analyses). Rather, it is a lattice of the form illustrated to the right and in Figure 1.6 (page 6). This lattice is *non-complemented*, *non-distributive*, *non-modular*, and *non-Boolean* (Proposition ?? page ??).<sup>12</sup>



In the world of mathematical structures, the order structure of wavelet analyses is quite rare, but not completely unique. One example of a system with similar structure is the set of *Primorial numbers* together with the  $|$  (“divides”) ordering relation<sup>13</sup> as illustrated to the right.



The basis sequence of most transform are fixed with no design freedom For example, the Fourier Transform uses the complex exponential, Taylor Expansion uses monomials of the form  $(x - a)^n$ . However, there are an infinite number of wavelet basis sequences—lots and lots of design freedom. For information regarding designing wavelet basis sequences, see Greenhoe (2013).

However, one arguable disadvantage is that wavelets do not support a **convolution theorem**—a theorem enjoyed by the Fourier transforms, Laplace Transform, and Z Transform. These other transforms induce a convolution theorem because they are defined in terms of an exponential (e.g.  $e^{-i\omega t}$ ,  $e^{-i\omega n}$ ,  $e^{-st}$ ,  $z^{-n}$ ), and exponentials sport the property  $a^{x+y} = a^x a^y$ .

<sup>12</sup> Greenhoe (2013) page 72 (Section 2.4.3 Order structure)

<sup>14</sup> Sloane (2014) (<http://oeis.org/A002110>), Greenhoe (2013) page 30



## CHAPTER 2

## TAYLOR EXPANSIONS (TRANSFORMS)

### 2.1 Introduction

For modeling real-world processes above the quantum level, measurements are *continuous* in time—that is, the first derivative of a function over time representing the measurement *exists*.

But even for “simple” physical systems, it is not just the first derivative that matters. For example, the classical “vibrating string” vertical displacement  $u(x, t)$  wave equation can be described as

$$\frac{\partial^2 u}{\partial x^2} - \frac{1}{c^2} \frac{\partial^2 u}{\partial t^2} = 0$$

Not only do physical systems demonstrate heavy dependence on the derivatives of their measurement functions, but also commonly exhibit *oscillation*, as demonstrated by sunspot activity over the last 300 years or earthquake activity (Figure 2.1 page 10).

In fact, derivatives and oscillations are fundamentally linked as demonstrated by the fact that all solutions of homogeneous second order differential equations are linear combinations of sine and cosine functions (Theorem C.3 page 78):

$$\left\{ \frac{d^2}{dx^2} f + f = 0 \right\} \iff \left\{ f(x) = f(0) \cos(x) + f'(0) \sin(x) \right\} \quad \forall f \in \mathcal{C}, \forall x \in \mathbb{R}$$

Derivatives are calculated *locally* about a point. Oscillations are observed *globally* over a range, and analyzed (decomposed) by projecting the function onto a sequence of basis functions—sinusoids in the case of Fourier Transform family. Projection is accomplished using inner products, and often these are calculated using *integration*. Note that derivatives and integrals are also fundamentally linked as demonstrated by the *Fundamental Theorem of Calculus*...which shows that integration can be calculated using anti-differentiation:

$$\int_a^b f(x) dx = F(b) - F(a) \quad \text{where } F(x) \text{ is the antiderivative of } f(x).$$

Brook Taylor showed that for *analytic* functions,<sup>1</sup> knowledge of the derivatives of a function at a location  $x = a$  allows you to determine (predict) arbitrarily closely all the points  $f(x)$  in the vicinity

<sup>1</sup>*analytic* functions: Functions for which all their derivatives exist.




Figure 2.1: Sunspot and earthquake measurements

of  $x = a$ :

$$f(x) = f(a) + \frac{1}{1!}f'(a)[x - a] + \frac{1}{2!}f''(a)[x - a]^2 + \frac{1}{3!}f'''(a)[x - a]^3 + \dots$$

On the other hand, the *Fourier Transform* is a kind of counter-part of the Taylor expansion:

	Taylor coefficients	Fourier coefficients
	Depend on derivatives $\frac{d^n}{dx^n}f(x)$	Depend on integrals $\int_{x \in \mathbb{R}} f(x)e^{-i\omega x} dx$
	Behavior in the vicinity of a point.	Behavior over the entire function.
	Demonstrate trends locally.	Demonstrate trends globally, such as oscillations.
	Admits <i>analytic</i> functions only.	Admits <i>non-analytic</i> functions as well.
	Function must be <i>continuous</i> .	Function can be <i>discontinuous</i> .

## 2.2 Taylor Expansion

**Theorem 2.1** (Taylor Series). <sup>4</sup> Let  $\mathcal{C}$  be the space of all ANALYTIC functions and  $\frac{d}{dx}$  in  $\mathcal{C}$  the DIFFERENTIATION OPERATOR.

THEM

A **Taylor Series** about the point  $x = a$  of a function  $f(x) \in \mathcal{C}$  is

$$f(x) = \sum_{n=0}^{\infty} \underbrace{\frac{\left[\frac{d^n}{dx^n} f\right](a)}{n!}}_{\text{coefficient}} \underbrace{(x - a)^n}_{\text{basis function}} \quad \forall a \in \mathbb{R}, f \in \mathcal{C}$$

A **Maclaurin Series** is a TAYLOR SERIES about the point  $a = 0$ .

<sup>2</sup> Robinson (1982) page 886

<sup>3</sup> Robinson (1982) page 886

<sup>4</sup> Flanigan (1983) page 221 (Theorem 15), Strichartz (1995) page 281, Sohrab (2003) page 317 (Theorem 8.4.9), Taylor (1715), Maclaurin (1742)

## CHAPTER 3

## LAPLACE TRANSFORM

### 3.1 Definition

**Definition 3.1.** Let  $\mathcal{L}^2_{(\mathbb{R}, \mathcal{B}, \mu)}$  be the space of all LEBESGUE SQUARE-INTEGRABLE FUNCTIONS.

DEF

The **Laplace Transform** operator  $\mathbf{L}$  is here defined as

$$[\mathbf{L}f](\omega) \triangleq \frac{1}{\sqrt{2\pi}} \int_{x \in \mathbb{R}} f(x) e^{-sx} dx \quad \forall f \in \mathcal{L}^2_{(\mathbb{R}, \mathcal{B}, \mu)}$$

In Definition 3.1, the scaling factor  $\frac{1}{\sqrt{2\pi}}$  is not normally found in most definitions of the Laplace Transform. However it is included here to make the operator  $\mathbf{L}$  more directly compatible with the Unitary Fourier Transform operator  $\tilde{\mathbf{F}}$  (Definition 4.2 page 16).

### 3.2 Shift relations

**Theorem 3.1** (Shift relations). Let  $\mathbf{L}$  be the LAPLACE TRANSFORM operator (Definition 3.1 page 11).

THM

$$\begin{aligned} \mathbf{L}[f(x-y)](s) &= e^{-sy} [\mathbf{L}f(x)](s) \\ [\mathbf{L}(e^{rx}g(x))](s) &= [\mathbf{L}g(x)](s-r) \end{aligned}$$

PROOF:

$\mathbf{L}[f(x-y)](s) = \frac{1}{\sqrt{2\pi}} \int_{x \in \mathbb{R}} f(x-y) e^{-sx} dx$	by definition of $\mathbf{L}$	(Definition 3.1 page 11)
$= \frac{1}{\sqrt{2\pi}} \int_{u \in \mathbb{R}} f(u) e^{-s(y+u)} du$	where $u \triangleq x-y$	$\implies x = y+u$
$= e^{-sy} \frac{1}{\sqrt{2\pi}} \int_{u \in \mathbb{R}} f(u) e^{-su} du$	by property of exponents	$a^{x+y} = a^x a^y$
$= e^{-sy} \frac{1}{\sqrt{2\pi}} \int_{x \in \mathbb{R}} f(x) e^{-sx} du$	by change of variable	$u \rightarrow x$
$= e^{-sy} [\mathbf{L}f(x)](s)$	by definition of $\mathbf{L}$	(Definition 3.1 page 11)

$$\begin{aligned}
[\mathbf{L}(e^{rx}g(x))](s) &= \frac{1}{\sqrt{2\pi}} \int_{x \in \mathbb{R}} e^{rx} g(x) e^{-sx} dx && \text{by definition of } \mathbf{L} && (\text{Definition 3.1 page 11}) \\
&= \frac{1}{\sqrt{2\pi}} \int_{x \in \mathbb{R}} g(x) e^{-(s-r)x} dx && \text{by property of exponents} && a^{x+y} = a^x a^y \\
&= [\mathbf{L}g(x)](s-r) && \text{by definition of } \mathbf{L} && (\text{Definition 3.1 page 11})
\end{aligned}$$

⇒

### 3.3 Convolution relations

#### Definition 3.2.<sup>1</sup>

DEF

The **convolution operation** is defined as

$$[f \star g](x) \triangleq f(x) \star g(x) \triangleq \int_{u \in \mathbb{R}} f(u)g(x-u) du \quad \forall f, g \in L^2_{(\mathbb{R}, \mathcal{B}, \mu)}$$

Theorem 3.2 (next) demonstrates that multiplication in the “time domain” is equivalent to convolution in the “s domain” and vice-versa.

**Theorem 3.2** (convolution theorem). *Let  $\mathbf{L}$  be the LAPLACE TRANSFORM operator (Definition 3.1 page 11) and  $\star$  the convolution operator (Definition 4.3 page 19).*

THM

$$\begin{aligned}
\underbrace{\mathbf{L}[f(x) \star g(x)](\omega)}_{\text{convolution in “time domain”}} &= \underbrace{\sqrt{2\pi}[\mathbf{L}f](s) [\mathbf{L}g](s)}_{\text{multiplication in “s domain”}} && \forall f, g \in L^2_{(\mathbb{R}, \mathcal{B}, \mu)} \\
\underbrace{\mathbf{L}[f(x)g(x)](\omega)}_{\text{multiplication in “time domain”}} &= \underbrace{\frac{1}{\sqrt{2\pi}}[\mathbf{L}f](s) \star [\mathbf{L}g](s)}_{\text{convolution in “s domain”}} && \forall f, g \in L^2_{(\mathbb{R}, \mathcal{B}, \mu)}.
\end{aligned}$$

✎ PROOF:

$$\begin{aligned}
\mathbf{L}[f(x) \star g(x)](s) &= \mathbf{L}\left[\int_{u \in \mathbb{R}} f(u)g(x-u) du\right](s) && \text{by definition of } \star && (\text{Definition 4.3 page 19}) \\
&= \int_{u \in \mathbb{R}} f(u) [\mathbf{L}g(x-u)](s) du \\
&= \int_{u \in \mathbb{R}} f(u) e^{-su} [\mathbf{L}g(x)](s) du && \text{by Fourier shift theorem} && (\text{Theorem 4.4 page 18}) \\
&= \sqrt{2\pi} \underbrace{\left(\frac{1}{\sqrt{2\pi}} \int_{u \in \mathbb{R}} f(u) e^{-su} du\right)}_{[\mathbf{L}f](s)} [\mathbf{L}g](s) \\
&= \sqrt{2\pi} [\mathbf{L}f](s) [\mathbf{L}g](s) && \text{by definition of } \mathbf{L} && (\text{Definition 4.2 page 16}) \\
\mathbf{L}[f(x)g(x)](s) &= \mathbf{L}[(\mathbf{L}^{-1}\mathbf{L}f(x)) g(x)](s) && \text{by def. of operator inverse} && (\text{Definition F.3 page 120}) \\
&= \mathbf{L}\left[\left(\frac{1}{\sqrt{2\pi}} \int_{v \in \mathbb{R}} [\mathbf{L}f(x)](v) e^{s xv} dv\right) g(x)\right](s) && \text{by Theorem 4.1 page 17} \\
&= \frac{1}{\sqrt{2\pi}} \int_{v \in \mathbb{R}} [\mathbf{L}f(x)](v) [\mathbf{L}(e^{s xv} g(x))](s, v) dv \\
&= \frac{1}{\sqrt{2\pi}} \int_{v \in \mathbb{R}} [\mathbf{L}f(x)](v) [\mathbf{L}g(x)](s-v) dv && \text{by Theorem 4.4 page 18} \\
&= \frac{1}{\sqrt{2\pi}} [\mathbf{L}f](s) \star [\mathbf{L}g](s) && \text{by definition of } \star && (\text{Definition 4.3 page 19})
\end{aligned}$$

<sup>1</sup> [Bachman \(1964\) page 6](#), [Bracewell \(1978\) page 108](#) (Convolution theorem)



## 3.4 Calculus relations

**Theorem 3.3.** Let  $\mathbf{L}$  be the LAPLACE TRANSFORM operator (Definition 3.1 page 11).

$$\boxed{\text{T H M}} \quad \left\{ \lim_{t \rightarrow -\infty} x(t) = 0 \right\} \implies \left\{ \mathbf{L} \left[ \frac{d}{dt} x(t) \right] = s[\mathbf{L}x](s) \right\}$$

PROOF:

$$\begin{aligned} \mathbf{L} \left[ \frac{d}{dt} x(t) \right] &\triangleq \int_{t \in \mathbb{R}} \underbrace{\left[ \frac{d}{dt} x(t) \right]}_{dv} \underbrace{e^{-st}}_u dt && \text{by definition of } \mathbf{L} \\ &= \underbrace{e^{-st}}_u \underbrace{x(t)}_v \Big|_{t=-\infty}^{t=+\infty} - \int_{t \in \mathbb{R}} \underbrace{x(t)}_v \underbrace{(-s)e^{-st}}_{du} dt && \text{by Integration by Parts} \\ &= \cancel{e^{-s\infty}} \overset{0}{x(\infty)} - \cancel{e^{s\infty}} \overset{0}{x(-\infty)} - (-s) \underbrace{\int_{t \in \mathbb{R}} x(t) e^{-st} dt}_{\text{Laplace Transform of } x(t)} && \text{by left hypothesis} \\ &= s[\mathbf{L}x](s) \end{aligned}$$



**Theorem 3.4.** Let  $\mathbf{L}$  be the LAPLACE TRANSFORM operator (Definition 3.1 page 11).

$$\boxed{\text{T H M}} \quad \mathbf{L} \int_{u=-\infty}^{u=t} x(u) du = \frac{1}{s} [\mathbf{L}x](s)$$

PROOF:

1. Define the Heaviside function  $h(t)$  as  $h(t) \triangleq \begin{cases} 1 & \text{for } t \geq 0 \\ 0 & \text{otherwise} \end{cases}$
2. Remainder of proof...

$$\begin{aligned} \mathbf{L} \int_{u=-\infty}^{u=t} x(u) du &\triangleq \int_{t=-\infty}^{t=+\infty} \left[ \int_{u=-\infty}^{u=t} x(u) du \right] e^{-st} dt && \text{by definition of } \mathbf{L} \\ &= \int_{t=-\infty}^{t=+\infty} \left[ \int_{u=-\infty}^{u=+\infty} x(u) h(t-u) du \right] e^{-st} dt && \left( \begin{array}{l} \text{by definition of Heaviside function} \\ \text{definition 1} \end{array} \right) \\ &= \int_{v=-\infty}^{v=+\infty} \int_{u=-\infty}^{u=+\infty} x(u) h(v) e^{-s(u+v)} du dv && \left( \begin{array}{l} \text{where } v \triangleq t-u \\ \implies t = u+v \end{array} \right) \\ &= \left[ \int_{v=-\infty}^{v=+\infty} h(v) e^{-sv} dv \right] \underbrace{\left[ \int_{u=-\infty}^{u=+\infty} x(u) e^{-su} du \right]}_{\text{Laplace Transform of } x(t)} \\ &= \left[ \int_{v=0}^{v=+\infty} e^{-sv} dv \right] [\mathbf{L}x](s) && \left( \begin{array}{l} \text{by definition of Heaviside function} \\ \text{definition 1} \end{array} \right) \end{aligned}$$

$$\begin{aligned} &= \frac{1}{-s} e^{-sv} \Big|_{v=0}^{v=\infty} [\mathbf{L}x](s) \\ &= \boxed{\frac{1}{s} [\mathbf{L}x](s)} \end{aligned}$$

by *Fundamental Theorem of Calculus*





## CHAPTER 4

## FOURIER TRANSFORM



*“Up to this point we have supposed that the function whose development is required in a series of sines of multiple arcs can be developed in a series arranged according to powers of the variable  $x$ . ... We can extend the same results to any functions, even to those which are discontinuous and entirely arbitrary. ... even entirely arbitrary functions may be developed in series of sines of multiple arcs.”*

Joseph Fourier (1768–1830) <sup>1</sup>

### 4.1 Introduction

Historically, before the Fourier Transform was the Taylor Expansion (transform). The Taylor Expansion demonstrates that for **analytic** functions knowledge of the derivatives of a function at a location  $x = a$  allows you to determine (predict) arbitrarily closely all the points  $f(x)$  in the vicinity of  $x = a$  (CHAPTER 2 page 9). But analytic functions are by definition functions for which all their derivatives exist. Thus, if a function is *discontinuous*, it is simply not a candidate for a Taylor Expansion. And some 300 years ago, mathematician giants of the day were fairly content with this.

But then in came an engineer named Joseph Fourier whose day job was working as a governor of lower Egypt under Napoleon. He claimed that, rather than expansion based on derivatives, one could expand based on integrals over sinusoids, and that this would work not just for analytic functions, but for **discontinuous** ones as well!<sup>2</sup>

Needless to say, this did not go over too well initially in the mathematical community. But over time (on the order of 200 or so years), the Fourier Transform has in many ways won the day.





<sup>1</sup> quote:  Fourier (1878) page 184,186 (§219,220)

image: <http://en.wikipedia.org/wiki/File:Fourier2.jpg>, public domain

<sup>2</sup>  Robinson (1982) page 886

<sup>3</sup> Caricature of Legendre (left) and Fourier (right), 1820, by Julien-Léopold Boilly (1796–1874). “Album de 73

## 4.2 Definitions

This chapter deals with the *Fourier Transform* in the space of *Lebesgue square-integrable functions*  $L^2_{(\mathbb{R}, \mathcal{B}, \mu)}$ , where  $\mathbb{R}$  is the set of real numbers,  $\mathcal{B}$  is the set of *Borel sets* on  $\mathbb{R}$ ,  $\mu$  is the standard *Borel measure* on  $\mathbb{R}$ , and

$$L^2_{(\mathbb{R}, \mathcal{B}, \mu)} \triangleq \left\{ f \in \mathbb{R}^{\mathbb{R}} \mid \int_{\mathbb{R}} |f|^2 d\mu < \infty \right\}.$$

Furthermore,  $\langle \triangle | \nabla \rangle$  is the *inner product* induced by the operator  $\int_{\mathbb{R}} d\mu$  such that

$$\langle f | g \rangle \triangleq \int_{\mathbb{R}} f(x) g^*(x) dx,$$

and  $(L^2_{(\mathbb{R}, \mathcal{B}, \mu)}, \langle \triangle | \nabla \rangle)$  is a *Hilbert space*.

**Definition 4.1.** Let  $\kappa$  be a FUNCTION in  $\mathbb{C}^{\mathbb{R}^2}$ .

DEF

The function  $\kappa$  is the **Fourier kernel** if  $\kappa(x, \omega) \triangleq e^{i\omega x} \quad \forall x, \omega \in \mathbb{R}$

**Definition 4.2.** <sup>4</sup> Let  $L^2_{(\mathbb{R}, \mathcal{B}, \mu)}$  be the space of all LEBESGUE SQUARE-INTEGRABLE FUNCTIONS.

DEF

The **Fourier Transform** operator  $\tilde{\mathbf{F}}$  is defined as

$$[\tilde{\mathbf{F}}f](\omega) \triangleq \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} f(x) e^{-i\omega x} dx \quad \forall f \in L^2_{(\mathbb{R}, \mathcal{B}, \mu)}$$

This definition of the Fourier Transform is also called the **unitary Fourier Transform**.

**Remark 4.1 (Fourier transform scaling factor).** <sup>5</sup> If the Fourier transform operator  $\tilde{\mathbf{F}}$  and inverse Fourier transform operator  $\tilde{\mathbf{F}}^{-1}$  are defined as

$$\tilde{\mathbf{F}}f(x) \triangleq F(\omega) \triangleq A \int_{\mathbb{R}} f(x) e^{-i\omega x} dx \quad \text{and} \quad \tilde{\mathbf{F}}^{-1}\tilde{f}(\omega) \triangleq B \int_{\mathbb{R}} F(\omega) e^{i\omega x} d\omega$$

then  $A$  and  $B$  can be any constants as long as  $AB = \frac{1}{2\pi}$ . The Fourier transform is often defined with the scaling factor  $A$  set equal to 1 such that  $[\tilde{\mathbf{F}}f(x)](\omega) \triangleq \int_{\mathbb{R}} f(x) e^{-i\omega x} dx$ . In this case, the inverse Fourier transform operator  $\tilde{\mathbf{F}}^{-1}$  is either defined as

$$\begin{aligned} \text{🐡} \quad [\tilde{\mathbf{F}}^{-1}f(x)](f) &\triangleq \int_{\mathbb{R}} f(x) e^{i2\pi f x} dx \quad (\text{using oscillatory frequency free variable } f) \text{ or} \\ \text{🐡} \quad [\tilde{\mathbf{F}}^{-1}f(x)](\omega) &\triangleq \frac{1}{2\pi} \int_{\mathbb{R}} f(x) e^{i\omega x} dx \quad (\text{using angular frequency free variable } \omega). \end{aligned}$$

In short, the  $2\pi$  has to show up somewhere, either in the argument of the exponential ( $e^{-i2\pi f t}$ ) or in front of the integral ( $\frac{1}{2\pi} \int \dots$ ). One could argue that it is unnecessary to burden the exponential argument with the  $2\pi$  factor ( $e^{-i2\pi f t}$ ), and thus could further argue in favor of using the angular frequency variable  $\omega$  thus giving the inverse operator definition  $[\tilde{\mathbf{F}}^{-1}f(x)](\omega) \triangleq \frac{1}{2\pi} \int_{\mathbb{R}} f(x) e^{-i\omega x} dx$ . But this causes a new problem. In this case, the Fourier operator  $\tilde{\mathbf{F}}$  is not *unitary* (see Theorem 4.2 page 17)—in particular,  $\tilde{\mathbf{F}}\tilde{\mathbf{F}}^* \neq \mathbf{I}$ , where  $\tilde{\mathbf{F}}^*$  is the *adjoint* of  $\tilde{\mathbf{F}}$ ; but rather,  $\tilde{\mathbf{F}}\left(\frac{1}{2\pi}\tilde{\mathbf{F}}^*\right) = \left(\frac{1}{2\pi}\tilde{\mathbf{F}}^*\right)\tilde{\mathbf{F}} = \mathbf{I}$ . But if we define the operators  $\tilde{\mathbf{F}}$  and  $\tilde{\mathbf{F}}^{-1}$  to both have the scaling factor  $\frac{1}{\sqrt{2\pi}}$ , then  $\tilde{\mathbf{F}}$  and  $\tilde{\mathbf{F}}^{-1}$  are inverses and  $\tilde{\mathbf{F}}$  is *unitary*—that is,  $\tilde{\mathbf{F}}\tilde{\mathbf{F}}^* = \tilde{\mathbf{F}}^*\tilde{\mathbf{F}} = \mathbf{I}$ .

Portraits-Charge Aquarelle's des Membres de l'Institute (watercolor portrait #29). Bibliotheque de l'Institut de France." Public domain. [https://en.wikipedia.org/wiki/File:Legendre\\_and\\_Fourier\\_\(1820\).jpg](https://en.wikipedia.org/wiki/File:Legendre_and_Fourier_(1820).jpg)

<sup>4</sup> 🐡 Bachman et al. (2000) page 363, 🐡 Chorin and Hald (2009) page 13, 🐡 Loomis and Bolker (1965) page 144, 🐡 Knapp (2005b) pages 374–375, 🐡 Fourier (1822), 🐡 Fourier (1878) page 336?

<sup>5</sup> 🐡 Chorin and Hald (2009) page 13, 🐡 Jeffrey and Dai (2008) pages xxxi–xxxii, 🐡 Knapp (2005b) pages 374–375

## 4.3 Operator properties

**Theorem 4.1** (Inverse Fourier transform).<sup>6</sup> Let  $\tilde{\mathbf{F}}$  be the Fourier Transform operator (Definition 4.2 page 16). The inverse  $\tilde{\mathbf{F}}^{-1}$  of  $\tilde{\mathbf{F}}$  is

$$\boxed{\text{THM}} \quad [\tilde{\mathbf{F}}^{-1}\tilde{f}](x) = \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} \tilde{f}(\omega) e^{i\omega x} d\omega \quad \forall \tilde{f} \in L^2_{(\mathbb{R}, \mathcal{B}, \mu)}$$

**Theorem 4.2.** Let  $\tilde{\mathbf{F}}$  be the Fourier Transform operator with inverse  $\tilde{\mathbf{F}}^{-1}$  and adjoint  $\tilde{\mathbf{F}}^*$ .

$$\boxed{\text{THM}} \quad \tilde{\mathbf{F}}^* = \tilde{\mathbf{F}}^{-1}$$

✎ PROOF:

$$\begin{aligned} \langle \tilde{\mathbf{F}}f | g \rangle &= \left\langle \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} f(x) e^{-i\omega x} dx | g(\omega) \right\rangle && \text{by definition of } \tilde{\mathbf{F}} \text{ page 16} \\ &= \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} f(x) \langle e^{-i\omega x} | g(\omega) \rangle dx && \text{by additive property of } \langle \Delta | \nabla \rangle \\ &= \int_{\mathbb{R}} f(x) \frac{1}{\sqrt{2\pi}} \langle g(\omega) | e^{-i\omega x} \rangle^* dx && \text{by conjugate symmetric property of } \langle \Delta | \nabla \rangle \\ &= \left\langle f(x) | \frac{1}{\sqrt{2\pi}} \langle g(\omega) | e^{-i\omega x} \rangle \right\rangle && \text{by definition of } \langle \Delta | \nabla \rangle \\ &= \left\langle f | \underbrace{\tilde{\mathbf{F}}^{-1}}_{\tilde{\mathbf{F}}^*} g \right\rangle && \text{by Theorem 4.1 page 17} \end{aligned}$$

⇒

The Fourier Transform operator has several nice properties:

🔥  $\tilde{\mathbf{F}}$  is unitary<sup>7</sup> (Corollary 4.1—next corollary).

🔥 Because  $\tilde{\mathbf{F}}$  is unitary, it automatically has several other nice properties (Theorem 4.3 page 17).

**Corollary 4.1.** Let  $\mathbf{I}$  be the identity operator and let  $\tilde{\mathbf{F}}$  be the Fourier Transform operator with adjoint  $\tilde{\mathbf{F}}^*$  and inverse  $\tilde{\mathbf{F}}^{-1}$ .

$$\boxed{\text{COR}} \quad \underbrace{\tilde{\mathbf{F}}\tilde{\mathbf{F}}^* = \tilde{\mathbf{F}}^*\tilde{\mathbf{F}} = \mathbf{I}}_{\tilde{\mathbf{F}}^* = \tilde{\mathbf{F}}^{-1}} \quad (\tilde{\mathbf{F}} \text{ is unitary})$$

✎ PROOF: This follows directly from the fact that  $\tilde{\mathbf{F}}^* = \tilde{\mathbf{F}}^{-1}$  (Theorem 4.2 page 17).

⇒

**Theorem 4.3.** Let  $\tilde{\mathbf{F}}$  be the Fourier transform operator with adjoint  $\tilde{\mathbf{F}}^*$  and inverse  $\tilde{\mathbf{F}}$ . Let  $\|\cdot\|$  be the operator norm with respect to the vector norm  $\|\cdot\|$  with respect to the Hilbert space  $(\mathbb{C}^{\mathbb{R}}, \langle \Delta | \nabla \rangle)$ . Let  $\mathcal{R}(\mathbf{A})$  be the range of an operator  $\mathbf{A}$ .

$$\boxed{\text{THM}} \quad \begin{aligned} \mathcal{R}(\tilde{\mathbf{F}}) &= \mathcal{R}(\tilde{\mathbf{F}}^{-1}) &&= L^2_{\mathbb{R}} \\ \|\tilde{\mathbf{F}}\| &= \|\tilde{\mathbf{F}}^{-1}\| &&= 1 && \text{(UNITARY)} \\ \langle \tilde{\mathbf{F}}f | \tilde{\mathbf{F}}g \rangle &= \langle \tilde{\mathbf{F}}^{-1}f | \tilde{\mathbf{F}}^{-1}g \rangle &&= \langle f | g \rangle && \text{(PARSEVAL'S EQUATION)} \\ \|\tilde{\mathbf{F}}f\| &= \|\tilde{\mathbf{F}}^{-1}f\| &&= \|f\| && \text{(PLANCHEREL'S FORMULA)} \\ \|\tilde{\mathbf{F}}f - \tilde{\mathbf{F}}g\| &= \|\tilde{\mathbf{F}}^{-1}f - \tilde{\mathbf{F}}^{-1}g\| &&= \|f - g\| && \text{(ISOMETRIC)} \end{aligned}$$

✎ PROOF: These results follow directly from the fact that  $\tilde{\mathbf{F}}$  is unitary (Corollary 4.1 page 17) and from the properties of unitary operators (Theorem F.26 page 144).

⇒

<sup>6</sup> Chorin and Hald (2009) page 13

<sup>7</sup> unitary operators: Definition F.14 page 143

## 4.4 Shift relations

**Theorem 4.4** (Shift relations). *Let  $\tilde{\mathbf{F}}$  be the Fourier Transform operator (Definition 4.2 page 16).*

<b>T H M</b>	$\tilde{\mathbf{F}}[f(x - y)](\omega) = e^{-i\omega y} [\tilde{\mathbf{F}}f(x)](\omega)$
	$[\tilde{\mathbf{F}}(e^{irx}g(x))](\omega) = [\tilde{\mathbf{F}}g(x)](\omega - r)$

 **PROOF:** Let  $\mathbf{L}$  be the Laplace Transform operator (Definition 3.1 page 11).

$\tilde{\mathbf{F}}[f(x - y)](\omega) = \mathbf{L}[f(x - y)](s) _{s=i\omega}$	by definition of $\mathbf{L}$	(Definition 3.1 page 11)
$= e^{-sy} [\mathbf{L}f(x)](s) _{s=i\omega}$	by Laplace shift relation	(Theorem 3.1 page 11)
$= e^{-i\omega y} [\tilde{\mathbf{F}}f(x)](\omega)$	by definition of $\tilde{\mathbf{F}}$	(Definition 4.2 page 16)
$[\tilde{\mathbf{F}}(e^{irx}g(x))](\omega) = [\mathbf{L}(e^{irx}g(x))](s) _{s=i\omega}$	by definition of $\mathbf{L}$	(Definition 3.1 page 11)
$= [[\mathbf{L}g(x)](s - r)] _{s=i\omega}$	by Laplace shift relation	(Theorem 3.1 page 11)
$= [\tilde{\mathbf{F}}g(x)](\omega - r)$	by definition of $\tilde{\mathbf{F}}$	(Definition 4.2 page 16)



**Theorem 4.5** (Complex conjugate). *Let  $\tilde{\mathbf{F}}$  be the Fourier Transform operator and  $*$  represent the complex conjugate operation on the set of complex numbers.*

<b>T H M</b>	$\tilde{\mathbf{F}}f^*(-x) = -[\tilde{\mathbf{F}}f(x)]^* \quad \forall f \in L^2_{(\mathbb{R}, \mathcal{B}, \mu)}$	
	$f \text{ is real} \implies \tilde{f}(-\omega) = [\tilde{f}(\omega)]^* \quad \forall \omega \in \mathbb{R}$	REALITY CONDITION

 **PROOF:**

$[\tilde{\mathbf{F}}f^*(-x)](\omega) \triangleq \frac{1}{\sqrt{2\pi}} \int f^*(-x) e^{-i\omega x} dx$	by definition of $\tilde{\mathbf{F}}$	(Definition 4.2 page 16)
$= \frac{1}{\sqrt{2\pi}} \int f^*(u) e^{i\omega u} (-1) du$	where $u \triangleq -x \implies dx = -du$	
$= - \left[ \frac{1}{\sqrt{2\pi}} \int f(u) e^{-i\omega u} du \right]^*$		
$\triangleq -[\tilde{\mathbf{F}}f(x)]^*$	by definition of $\tilde{\mathbf{F}}$	(Definition 4.2 page 16)
$\tilde{f}(-\omega) \triangleq \frac{1}{\sqrt{2\pi}} \int f(x) e^{-i(-\omega)x} dx$	by definition of $\tilde{\mathbf{F}}$	(Definition 4.2 page 16)
$= \left[ \frac{1}{\sqrt{2\pi}} \int f^*(x) e^{-i\omega x} dx \right]^*$		
$= \left[ \frac{1}{\sqrt{2\pi}} \int f(x) e^{-i\omega x} dx \right]^*$	by $f$ is real hypothesis	
$\triangleq \tilde{f}^*(\omega)$	by definition of $\tilde{\mathbf{F}}$	(Definition 4.2 page 16)



## 4.5 Convolution relations

### Definition 4.3. <sup>8</sup>

DEF

The **convolution operation** is defined as


$$[f \star g](x) \triangleq f(x) \star g(x) \triangleq \int_{u \in \mathbb{R}} f(u)g(x-u) du \quad \forall f, g \in L^2_{(\mathbb{R}, \mathcal{B}, \mu)}$$

Theorem 5.2 (next) demonstrates that multiplication in the “time domain” is equivalent to convolution in the “frequency domain” and vice-versa.

**Theorem 4.6** (convolution theorem). <sup>9</sup> Let  $\tilde{\mathbf{F}}$  be the Fourier Transform operator (Definition 4.2 page 16) and  $\star$  the convolution operator (Definition 4.3 page 19).

THM

$$\begin{aligned} \underbrace{\tilde{\mathbf{F}}[f(x) \star g(x)](\omega)}_{\text{convolution in “time domain”}} &= \underbrace{\sqrt{2\pi} [\tilde{\mathbf{F}}f](\omega) [\tilde{\mathbf{F}}g](\omega)}_{\text{multiplication in “frequency domain”}} && \forall f, g \in L^2_{(\mathbb{R}, \mathcal{B}, \mu)} \\ \underbrace{\tilde{\mathbf{F}}[f(x)g(x)](\omega)}_{\text{multiplication in “time domain”}} &= \underbrace{\frac{1}{\sqrt{2\pi}} [\tilde{\mathbf{F}}f](\omega) \star [\tilde{\mathbf{F}}g](\omega)}_{\text{convolution in “frequency domain”}} && \forall f, g \in L^2_{(\mathbb{R}, \mathcal{B}, \mu)}. \end{aligned}$$

 **PROOF:** Let  $\mathbf{L}$  be the Laplace Transform operator (Definition 3.1 page 11).

$$\begin{aligned} \tilde{\mathbf{F}}[f(x) \star g(x)](\omega) &= \mathbf{L}[f(x) \star g(x)](s)|_{s=i\omega} && \text{by definition of } \mathbf{L} && (\text{Definition 3.1 page 11}) \\ &= \sqrt{2\pi} [\mathbf{L}f](s) [\mathbf{L}g](s)|_{s=i\omega} && \text{by Laplace convolution result} && (\text{Theorem 3.2 page 12}) \\ &= \sqrt{2\pi} [\tilde{\mathbf{F}}f](\omega) [\tilde{\mathbf{F}}g](\omega) \\ \tilde{\mathbf{F}}[f(x)g(x)](\omega) &= \mathbf{L}[f(x)g(x)](s)|_{s=i\omega} \\ &= \frac{1}{\sqrt{2\pi}} [\mathbf{L}f](s) \star [\mathbf{L}g](s)|_{s=i\omega} \\ &= \frac{1}{\sqrt{2\pi}} [\tilde{\mathbf{F}}f](\omega) \star [\tilde{\mathbf{F}}g](\omega) \end{aligned}$$



## 4.6 Calculus relations

**Theorem 4.7.** Let  $\tilde{\mathbf{F}}$  be the FOURIER TRANSFORM operator (Definition 4.2 page 16).

THM


$$\left\{ \lim_{t \rightarrow -\infty} x(t) = 0 \right\} \implies \left\{ \tilde{\mathbf{F}} \left[ \frac{d}{dt} x(t) \right] = i\omega [\tilde{\mathbf{F}}x](\omega) \right\}$$

 **PROOF:** Let  $\mathbf{L}$  be the Laplace Transform operator (Definition 3.1 page 11).

$$\begin{aligned} \tilde{\mathbf{F}} \left[ \frac{d}{dt} x(t) \right] &\triangleq \mathbf{L} \left[ \frac{d}{dt} x(t) \right](s)|_{s=i\omega} && \text{by definitions of } \mathbf{L} \text{ and } \tilde{\mathbf{F}} && (\text{Definition 3.1 page 11}) \\ &= s[\mathbf{L}x(t)](s)|_{s=i\omega} && \text{by Theorem 3.3 page 13} \\ &= i\omega [\tilde{\mathbf{F}}x](\omega) \end{aligned}$$



<sup>8</sup>  Bachman (1964) page 6,  Bracewell (1978) page 108 (Convolution theorem)

<sup>9</sup>  Bracewell (1978) page 110

**Theorem 4.8.** Let  $\tilde{\mathbf{F}}$  be the FOURIER TRANSFORM operator (Definition 4.2 page 16).

$$\tilde{\mathbf{F}} \int_{u=-\infty}^{u=t} x(u) du = \frac{1}{i\omega} [\tilde{\mathbf{F}}x](\omega)$$

Let  $\mathbf{L}$  be the Laplace Transform operator (Definition 3.1 page 11).  $\Rightarrow$  PROOF:

$$\begin{aligned} \tilde{\mathbf{F}} \int_{u=-\infty}^{u=t} x(u) du &\triangleq \mathbf{L} \int_{u=-\infty}^{u=t} x(u) du \Big|_{s=i\omega} \\ &= \frac{1}{s} [\mathbf{L}x(t)](s) \Big|_{s=i\omega} && \text{by Theorem 3.4 page 13} \\ &= \frac{1}{i\omega} [\tilde{\mathbf{F}}x(t)](\omega) \end{aligned}$$

$\Rightarrow$

## 4.7 Real valued functions




Figure 4.1: Fourier transform components of real-valued signal

**Theorem 4.9.** Let  $f(x)$  be a function in  $L^2_{\mathbb{R}}$  and  $\tilde{f}(\omega)$  the FOURIER TRANSFORM of  $f(x)$ .

$$\left\{ \begin{array}{l} f(x) \text{ is REAL-VALUED} \\ (f \in \mathbb{R}^{\mathbb{R}}) \end{array} \right\} \Rightarrow \left\{ \begin{array}{ll} \tilde{f}(\omega) = \tilde{f}^*(-\omega) & (\text{HERMITIAN SYMMETRIC}) \\ \mathbf{R}_e[\tilde{f}(\omega)] = \mathbf{R}_e[\tilde{f}(-\omega)] & (\text{SYMMETRIC}) \\ \mathbf{I}_m[\tilde{f}(\omega)] = -\mathbf{I}_m[\tilde{f}(-\omega)] & (\text{ANTI-SYMMETRIC}) \\ |\tilde{f}(\omega)| = |\tilde{f}(-\omega)| & (\text{SYMMETRIC}) \\ \angle \tilde{f}(\omega) = \angle \tilde{f}(-\omega) & (\text{ANTI-SYMMETRIC}). \end{array} \right\}$$

$\Rightarrow$  PROOF:

$$\begin{aligned} \tilde{f}(\omega) &\triangleq [\tilde{\mathbf{F}}f(x)](\omega) \triangleq \langle f(x) | e^{i\omega x} \rangle = \langle f(x) | e^{i(-\omega)x} \rangle^* \triangleq \tilde{f}^*(-\omega) \\ \mathbf{R}_e[\tilde{f}(\omega)] &= \mathbf{R}_e[\tilde{f}^*(-\omega)] = \mathbf{R}_e[\tilde{f}(-\omega)] \\ \mathbf{I}_m[\tilde{f}(\omega)] &= \mathbf{I}_m[\tilde{f}^*(-\omega)] = -\mathbf{I}_m[\tilde{f}(-\omega)] \\ |\tilde{f}(\omega)| &= |\tilde{f}^*(-\omega)| = |\tilde{f}(-\omega)| \\ \angle \tilde{f}(\omega) &= \angle \tilde{f}^*(-\omega) = -\angle \tilde{f}(-\omega) \end{aligned}$$

$\Rightarrow$

## 4.8 Moment properties

**Definition 4.4.** <sup>10</sup>

The quantity  $M_n$  is the  $n$ th moment of a function  $f(x) \in L^2_{\mathbb{R}}$  if

$$M_n \triangleq \int_{\mathbb{R}} x^n f(x) dx \quad \text{for } n \in \mathbb{W}.$$

<sup>10</sup> [Jawerth and Sweldens \(1994\)](#) pages 16–17, [Sweldens and Piessens \(1993\)](#) page 2, [Vidakovic \(1999\)](#) page 83

**Lemma 4.1.** <sup>11</sup> Let  $M_n$  be the  $n$ TH MOMENT (Definition 4.4 page 20) and  $\tilde{f}(\omega) \triangleq [\tilde{\mathbf{F}}f](\omega)$  the FOURIER TRANSFORM (Definition 4.2 page 16) of a function  $f(x)$  in  $\mathcal{L}_{\mathbb{R}}^2$  (Definition B.1 page 69).

<b>L E M</b>	$M_n = \left. \sqrt{2\pi}(i)^n \left[ \frac{d}{d\omega} \right]^n \tilde{f}(\omega) \right _{\omega=0} \quad \forall n \in \mathbb{W}, f \in \mathcal{L}_{\mathbb{R}}^2$
	$\left[ \frac{d}{d\omega} \right]^n \tilde{f}(\omega) \Big _{\omega=0} = \frac{1}{\sqrt{2\pi}} (-i)^n M_n \quad \forall n \in \mathbb{W}, f \in \mathcal{L}_{\mathbb{R}}^2$

PROOF:

$$\begin{aligned}
 \sqrt{2\pi}(i)^n \left[ \frac{d}{d\omega} \right]^n \tilde{f}(\omega) \Big|_{\omega=0} &= \sqrt{2\pi}(i)^n \left[ \frac{d}{d\omega} \right]^n \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} f(x) e^{-i\omega x} dx \Big|_{\omega=0} && \text{by definition of } \tilde{\mathbf{F}} \quad (\text{Definition 4.2 page 16}) \\
 &= (i)^n \int_{\mathbb{R}} f(x) \left[ \frac{d}{d\omega} \right]^n e^{-i\omega x} dx \Big|_{\omega=0} \\
 &= (i)^n \int_{\mathbb{R}} f(x) [(-i)^n x^n e^{-i\omega x}] dx \Big|_{\omega=0} \\
 &= (-i^2)^n \int_{\mathbb{R}} f(x) x^n dx \\
 &= \int_{\mathbb{R}} f(x) x^n dx \\
 &\triangleq M_n && \text{by definition of } M_n \quad (\text{Definition 4.4 page 20})
 \end{aligned}$$

⇒

**Lemma 4.2.** <sup>12</sup> Let  $M_n$  be the  $n$ TH MOMENT (Definition 4.4 page 20) and  $\tilde{f}(\omega) \triangleq [\tilde{\mathbf{F}}f](\omega)$  the FOURIER TRANSFORM (Definition 4.2 page 16) of a function  $f(x)$  in  $\mathcal{L}_{\mathbb{R}}^2$  (Definition B.1 page 69).

<b>L E M</b>	$M_n = 0 \quad \iff \quad \left[ \frac{d}{d\omega} \right]^n \tilde{f}(\omega) \Big _{\omega=0} = 0 \quad \forall n \in \mathbb{W}$
----------------------	---

PROOF:



1. Proof for (  $\implies$  ) case:

$$\begin{aligned}
 0 &= \langle f(x) | x^n \rangle && \text{by left hypothesis} \\
 &= \sqrt{2\pi}(-i)^{-n} \left[ \frac{d}{d\omega} \right]^n \tilde{f}(\omega) \Big|_{\omega=0} && \text{by Lemma 4.1 page 21} \\
 &\implies \left[ \frac{d}{d\omega} \right]^n \tilde{f}(\omega) \Big|_{\omega=0} = 0
 \end{aligned}$$

2. Proof for (  $\impliedby$  ) case:

$$\begin{aligned}
 0 &= \left[ \frac{d}{d\omega} \right]^n \tilde{f}(\omega) \Big|_{\omega=0} && \text{by right hypothesis} \\
 &= \left[ \frac{d}{d\omega} \right]^n \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} f(x) e^{-i\omega x} dx \Big|_{\omega=0} && \text{by definition of } \tilde{f}(\omega) \\
 &= \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} f(x) \left[ \frac{d}{d\omega} \right]^n e^{-i\omega x} dx \Big|_{\omega=0} \\
 &= \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} f(x) [(-i)^n x^n e^{-i\omega x}] dx \Big|_{\omega=0} \\
 &= (-i)^n \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} f(x) x^n dx \\
 &= (-i)^n \frac{1}{\sqrt{2\pi}} \langle f(x) | x^n \rangle && \text{by definition of } \langle \cdot | \cdot \rangle \text{ in } \mathcal{L}_{\mathbb{R}}^2 \quad (\text{Definition B.1 page 69})
 \end{aligned}$$

<sup>11</sup>  Goswami and Chan (1999) pages 38–39

<sup>12</sup>  Vidakovic (1999) pages 82–83,  Mallat (1999) pages 241–242



**Lemma 4.3** (Strang-Fix condition).<sup>13</sup> Let  $f(x)$  be a function in  $L^2_{\mathbb{R}}$  and  $M_n$  the  $n$ TH MOMENT (Definition 4.4 page 20) of  $f(x)$ . Let  $T$  be the TRANSLATION OPERATOR (Definition 1.3 page 188).

<b>L E M</b>	$\underbrace{\sum_{k \in \mathbb{Z}} T^k x^n f(x) = M_n}_{\text{STRANG-FIX CONDITION in "time"}} \iff \underbrace{\left[ \frac{d}{d\omega} \right]^n \tilde{f}(\omega) \Big _{\omega=2\pi k} = \frac{1}{\sqrt{2\pi}} (-i)^n \bar{\delta}_k M_n}_{\text{STRANG-FIX CONDITION in "frequency"}}$
----------------------	---

PROOF:

1. Proof for ( $\implies$ ) case:

$$\begin{aligned}
 \left[ \left[ \frac{d}{d\omega} \right]^n \tilde{f}(\omega) \right]_{\omega=2\pi k} &= \sum_{k \in \mathbb{Z}} \left[ \left[ \frac{d}{d\omega} \right]^n \tilde{f}(\omega) \right]_{\omega=2\pi k} e^{i2\pi k x} \bar{\delta}_k \\
 &= \frac{1}{\sqrt{2\pi}} \sum_{k \in \mathbb{Z}} \left[ \left[ \frac{d}{d\omega} \right]^n \int_{\mathbb{R}} f(x) e^{-i\omega x} dx \right]_{\omega=2\pi k} e^{i2\pi k x} \bar{\delta}_k && \text{by definition of } \tilde{f}(\omega) \quad (\text{Definition 4.2 page 16}) \\
 &= \frac{1}{\sqrt{2\pi}} \sum_{k \in \mathbb{Z}} \left[ \int_{\mathbb{R}} f(x) (-ix)^n e^{-i\omega x} dx \right]_{\omega=2\pi k} e^{i2\pi k x} \bar{\delta}_k \\
 &= (-i)^n \frac{1}{\sqrt{2\pi}} \sum_{k \in \mathbb{Z}} \left[ \int_{\mathbb{R}} x^n f(x) e^{-i\omega x} dx \right]_{\omega=2\pi k} e^{i2\pi k x} \bar{\delta}_k \\
 &= (-i)^n \frac{1}{\sqrt{2\pi}} \sum_{k \in \mathbb{Z}} (x-k)^n f(x-k) \bar{\delta}_k && \text{by PSF} \quad (\text{Theorem 1.2 page 196}) \\
 &= \frac{1}{\sqrt{2\pi}} (-i)^n \bar{\delta}_k M_n && \text{by left hypothesis}
 \end{aligned}$$

2. Proof for ( $\impliedby$ ) case:

$$\begin{aligned}
 \frac{1}{\sqrt{2\pi}} (-i)^n M_n &= \frac{1}{\sqrt{2\pi}} \sum_{k \in \mathbb{Z}} [(-i)^n \bar{\delta}_k M_n] e^{-i2\pi k x} && \text{by definition of } \bar{\delta} \quad (\text{Definition G.12 page 160}) \\
 &= \sum_{k \in \mathbb{Z}} \left[ \left[ \frac{d}{d\omega} \right]^n \tilde{f}(\omega) \right]_{\omega=2\pi k} e^{-i2\pi k x} && \text{by right hypothesis} \\
 &= \sum_{k \in \mathbb{Z}} \left[ \left[ \frac{d}{d\omega} \right]^n \int_{\mathbb{R}} f(x) e^{-i\omega x} dx \right]_{\omega=2\pi k} e^{-i2\pi k x} \\
 &= \sum_{k \in \mathbb{Z}} \left[ \int_{\mathbb{R}} f(x) (-ix)^n e^{-i\omega x} dx \right]_{\omega=2\pi k} e^{-i2\pi k x} \\
 &= (-i)^n \sum_{k \in \mathbb{Z}} \left[ \int_{\mathbb{R}} x^n f(x) e^{-i\omega x} dx \right]_{\omega=2\pi k} e^{-i2\pi k x} \\
 &= (-i)^n \sum_{k \in \mathbb{Z}} (x-k)^n f(x-k) && \text{by PSF} \quad (\text{Theorem 1.2 page 196})
 \end{aligned}$$




<sup>13</sup> Jawerth and Sweldens (1994) pages 16–17, Sweldens and Piessens (1993) page 2, Vidakovic (1999) page 83, Mallat (1999) pages 241–243, Fix and Strang (1969)



## 4.9 Examples

**Example 4.1** (rectangular pulse). Let  $\tilde{f}(\omega)$  be the *Fourier transform* of a function  $f(x) \in L^2_{\mathbb{R}}$ .




PROOF:

$$\begin{aligned}
 \tilde{f}(\omega) &= \tilde{\mathbf{F}}[f(x)](\omega) && \text{by definition of } \tilde{f}(\omega) \\
 &= e^{-i\left(\frac{a+b}{2}\omega\right)} \tilde{\mathbf{F}}\left[f\left(x - \frac{a+b}{2}\right)\right](\omega) && \text{by shift relation (Theorem 4.4 page 18)} \\
 &= e^{-i\left(\frac{a+b}{2}\omega\right)} \tilde{\mathbf{F}}\left[c \mathbb{1}_{[a:b)}\left(x - \frac{a+b}{2}\right)\right](\omega) && \text{by definition of } f(x) \\
 &= e^{-i\left(\frac{a+b}{2}\omega\right)} \tilde{\mathbf{F}}\left[c \mathbb{1}_{\left[-\frac{b-a}{2} : \frac{b-a}{2}\right)}(x)\right](\omega) && \text{by definition of } \mathbb{1} \text{ (Definition 1.2 page 188)} \\
 &= \frac{1}{\sqrt{2\pi}} e^{-i\left(\frac{a+b}{2}\omega\right)} \int_{\mathbb{R}} c \mathbb{1}_{\left[-\frac{b-a}{2} : \frac{b-a}{2}\right)}(x) e^{-i\omega x} dx && \text{by definition of } \tilde{\mathbf{F}} \text{ (Definition 4.2 page 16)} \\
 &= \frac{1}{\sqrt{2\pi}} e^{-i\left(\frac{a+b}{2}\omega\right)} \int_{-\frac{b-a}{2}}^{\frac{b-a}{2}} c e^{-i\omega x} dx && \text{by definition of } \mathbb{1} \text{ (Definition 1.2 page 188)} \\
 &= \frac{c}{\sqrt{2\pi}} e^{-i\left(\frac{a+b}{2}\omega\right)} \frac{1}{-i\omega} e^{-i\omega x} \Big|_{-\frac{b-a}{2}}^{\frac{b-a}{2}} \\
 &= \frac{2c}{\sqrt{2\pi}\omega} e^{-i\left(\frac{a+b}{2}\omega\right)} \left[ \frac{e^{i\left(\frac{b-a}{2}\omega\right)} - e^{-i\left(\frac{b-a}{2}\omega\right)}}{2i} \right] \\
 &= \frac{c(b-a)}{\sqrt{2\pi}} e^{-i\left(\frac{a+b}{2}\omega\right)} \left[ \frac{\sin\left(\frac{b-a}{2}\omega\right)}{\left(\frac{b-a}{2}\omega\right)} \right] && \text{by Euler formulas (Corollary C.2 page 81)}
 \end{aligned}$$

⇒

**Example 4.2** (triangle). Let  $\tilde{f}(\omega)$  be the *Fourier transform* of a function  $f(x) \in L^2_{\mathbb{R}}$ .




PROOF:

$$\tilde{f}(\omega) = \tilde{\mathbf{F}}[f(x)](\omega) \quad \text{by definition of } \tilde{f}(\omega)$$

$$\begin{aligned}
&= e^{-i\left(\frac{a+b}{2}\right)\omega} \tilde{\mathbf{F}}\left[f\left(x - \frac{a+b}{2}\right)\right](\omega) && \text{by shift relation} && (\text{Theorem 4.4 page 18}) \\
&= \tilde{\mathbf{F}}\left[c\left(1 - \frac{|2x - b - a|}{b - a}\right) \mathbb{1}_{[a:b]}(x)\right](\omega) && \text{by definition of } f(x) \\
&= c \tilde{\mathbf{F}}\left[\mathbb{1}_{\left[\frac{a}{2}:\frac{b}{2}\right]}(x) \star \mathbb{1}_{\left[\frac{a}{2}:\frac{b}{2}\right]}(x)\right](\omega) \\
&= c \sqrt{2\pi} \tilde{\mathbf{F}}\left[\mathbb{1}_{\left[\frac{a}{2}:\frac{b}{2}\right]}\right] \tilde{\mathbf{F}}\left[\mathbb{1}_{\left[\frac{a}{2}:\frac{b}{2}\right]}\right] && \text{by convolution theorem} && (\text{Theorem 5.2 page 30}) \\
&= c \sqrt{2\pi} \left(\tilde{\mathbf{F}}\left[\mathbb{1}_{\left[\frac{a}{2}:\frac{b}{2}\right]}\right]\right)^2 \\
&= c \sqrt{2\pi} \left(\frac{\left(\frac{b-a}{2}\right)}{\sqrt{2\pi}} e^{-i\left(\frac{a+b}{4}\right)\omega} \left[\frac{\sin\left(\frac{b-a}{4}\omega\right)}{\left(\frac{b-a}{4}\omega\right)}\right]\right)^2 && \text{by Rectangular pulse ex.} && \text{Example 4.1 page 23} \\
&= \frac{c(b-a)^2}{4\sqrt{2\pi}} e^{-i\left(\frac{a+b}{2}\right)\omega} \left[\frac{\sin\left(\frac{b-a}{4}\omega\right)}{\left(\frac{b-a}{4}\omega\right)}\right]^2
\end{aligned}$$

⇒

**Example 4.3.** Let a function  $f$  be defined in terms of the cosine function (Definition C.1 page 75) as follows:



**PROOF:** Let  $\mathbb{1}_A(x)$  be the *set indicator function* (Definition 1.2 page 188) on a set  $A$ .

$$\begin{aligned}
\tilde{f}(\omega) &\triangleq \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} f(x) e^{-i\omega x} dx && \text{by definition of } \tilde{f}(\omega) \text{ (Definition 4.2)} \\
&= \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} \cos^2\left(\frac{\pi}{2}x\right) \mathbb{1}_{[-1:1]}(x) e^{-i\omega x} dx && \text{by definition of } f(x) \\
&= \frac{1}{\sqrt{2\pi}} \int_{-1}^1 \cos^2\left(\frac{\pi}{2}x\right) e^{-i\omega x} dx && \text{by definition of } \mathbb{1} \text{ (Definition 1.2)} \\
&= \frac{1}{\sqrt{2\pi}} \int_{-1}^1 \left[ \frac{e^{i\frac{\pi}{2}x} + e^{-i\frac{\pi}{2}x}}{2} \right]^2 e^{-i\omega x} dx && \text{by Corollary C.2 page 81} \\
&= \frac{1}{4\sqrt{2\pi}} \int_{-1}^1 [2 + e^{i\pi x} + e^{-i\pi x}] e^{-i\omega x} dx \\
&= \frac{1}{4\sqrt{2\pi}} \int_{-1}^1 2e^{-i\omega x} + e^{-i(\omega+\pi)x} + e^{-i(\omega-\pi)x} dx \\
&= \frac{1}{4\sqrt{2\pi}} \left[ 2 \frac{e^{-i\omega x}}{-i\omega} + \frac{e^{-i(\omega+\pi)x}}{-i(\omega+\pi)} + \frac{e^{-i(\omega-\pi)x}}{-i(\omega-\pi)} \right]_{-1}^1 \\
&= \frac{1}{2\sqrt{2\pi}} \left[ 2 \frac{e^{-i\omega} - e^{+i\omega}}{-2i\omega} + \frac{e^{-i(\omega+\pi)} - e^{+i(\omega+\pi)}}{-2i(\omega+\pi)} + \frac{e^{-i(\omega-\pi)} - e^{+i(\omega-\pi)}}{-2i(\omega-\pi)} \right]_{-1}^1
\end{aligned}$$

$$= \frac{1}{2\sqrt{2\pi}} \left[ \underbrace{\frac{2\sin\omega}{\omega}}_{2\operatorname{sinc}(\omega)} + \underbrace{\frac{\sin(\omega + \pi)}{(\omega + \pi)}}_{\operatorname{sinc}(\omega + \pi)} + \underbrace{\frac{\sin(\omega - \pi)}{(\omega - \pi)}}_{\operatorname{sinc}(\omega - \pi)} \right]$$





# CHAPTER 5

## Z TRANSFORM

### 5.1 Convolution operator

**Definition 5.1.**<sup>1</sup> Let  $X^Y$  be the set of all functions from a set  $Y$  to a set  $X$ . Let  $\mathbb{Z}$  be the set of integers.

DEF

A function  $f$  in  $X^Y$  is a **sequence** over  $X$  if  $Y = \mathbb{Z}$ .

A sequence may be denoted in the form  $(x_n)_{n \in \mathbb{Z}}$  or simply as  $(x_n)$ .

**Definition 5.2.**<sup>2</sup> Let  $(\mathbb{F}, +, \cdot, 0, 1)$  be a FIELD (Definition A.5 page 68).

DEF

The space of all absolutely square summable sequences  $\ell_{\mathbb{F}}^2$  over  $\mathbb{F}$  is defined as

$$\ell_{\mathbb{F}}^2 \triangleq \left\{ (x_n)_{n \in \mathbb{Z}} \mid \sum_{n \in \mathbb{Z}} |x_n|^2 < \infty \right\}$$

The space  $\ell_{\mathbb{R}}^2$  is an example of a *separable Hilbert space*. In fact,  $\ell_{\mathbb{R}}^2$  is the *only* separable Hilbert space in the sense that all separable Hilbert spaces are isomorphically equivalent. For example,  $\ell_{\mathbb{R}}^2$  is isomorphic to  $L_{\mathbb{R}}^2$ , the space of all absolutely square Lebesgue integrable functions.

**Definition 5.3.**

DEF

The **convolution** operation  $\star$  is defined as

$$(x_n) \star (y_n) \triangleq \left( \sum_{m \in \mathbb{Z}} x_m y_{n-m} \right)_{n \in \mathbb{Z}} \quad \forall (x_n)_{n \in \mathbb{Z}}, (y_n)_{n \in \mathbb{Z}} \in \ell_{\mathbb{R}}^2$$

**Proposition 5.1.** Let  $\star$  be the CONVOLUTION OPERATOR (Definition 5.3 page 27).

PRP

$$(x_n) \star (y_n) = (y_n) \star (x_n) \quad \forall (x_n)_{n \in \mathbb{Z}}, (y_n)_{n \in \mathbb{Z}} \in \ell_{\mathbb{R}}^2 \quad (\star \text{ is COMMUTATIVE})$$

<sup>1</sup> Bromwich (1908) page 1, Thomson et al. (2008) page 23 (Definition 2.1), Joshi (1997) page 31

<sup>2</sup> Kubrusly (2011) page 347 (Example 5.K)

✎ PROOF:

$$\begin{aligned}
 [x \star y](n) &\triangleq \sum_{m \in \mathbb{Z}} x_m y_{n-m} && \text{by Definition 5.3 page 27} \\
 &= \sum_{k \in \mathbb{Z}} x_{n-k} y(k) && \text{where } k \triangleq n - m \implies m = n - k \\
 &= \sum_{k \in \mathbb{Z}} x_{n-k} y(k) && \text{by commutativity of addition} \\
 &= \sum_{m \in \mathbb{Z}} x_{n-m} y_m && \text{by change of variables} \\
 &= \sum_{m \in \mathbb{Z}} y_m x_{n-m} && \text{by commutative property of the field over } \mathbb{C} \\
 &\triangleq (y \star x)_n && \text{by Definition 5.3 page 27}
 \end{aligned}$$

⇒

**Proposition 5.2.** Let  $\star$  be the CONVOLUTION OPERATOR (Definition 5.3 page 27). Let  $\ell_{\mathbb{R}}^2$  be the set of ABSOLUTELY SUMMABLE sequences (Definition 5.2 page 27).

$$\text{PRP} \left\{ \begin{array}{l} \text{(A). } x(n) \in \ell_{\mathbb{R}}^2 \text{ and} \\ \text{(B). } y(n) \in \ell_{\mathbb{R}}^2 \end{array} \right\} \implies \left\{ \sum_{k \in \mathbb{Z}} x[k]y[n+k] = x[-n] \star y(n) \right\}$$

✎ PROOF:

$$\begin{aligned}
 \sum_{k \in \mathbb{Z}} x[k]y[n+k] &= \sum_{-p \in \mathbb{Z}} x[-p]y[n-p] && \text{where } p \triangleq -k \implies k = -p \\
 &= \sum_{p \in \mathbb{Z}} x[-p]y[n-p] && \text{by absolutely summable hypothesis (Definition 5.2 page 27)} \\
 &= \sum_{p \in \mathbb{Z}} x'[p]y[n-p] && \text{where } x'[n] \triangleq x[-n] \implies x[-n] = x'[n] \\
 &\triangleq x'[n] \star y[n] && \text{by definition of convolution } \star \text{ (Definition 5.3 page 27)} \\
 &\triangleq x[-n] \star y[n] && \text{by definition of } x'[n]
 \end{aligned}$$

⇒

## 5.2 Z-transform

**Definition 5.4.** <sup>3</sup>

**DEF** The **z-transform**  $\mathbf{Z}$  of  $(x_n)_{n \in \mathbb{Z}}$  is defined as

$$[\mathbf{Z}(x_n)](z) \triangleq \underbrace{\sum_{n \in \mathbb{Z}} x_n z^{-n}}_{\text{Laurent series}} \quad \forall (x_n) \in \ell_{\mathbb{R}}^2$$

**Theorem 5.1.** Let  $X(z) \triangleq \mathbf{Z}x[n]$  be the Z-TRANSFORM of  $x[n]$ .

$$\text{THM} \left\{ \check{x}(z) \triangleq \mathbf{Z}(x[n]) \right\} \implies \left\{ \begin{array}{l} \text{(1). } \mathbf{Z}(\alpha x[n]) = \alpha \check{x}(z) \quad \forall (x_n) \in \ell_{\mathbb{R}}^2 \text{ and} \\ \text{(2). } \mathbf{Z}(x[n-k]) = z^{-k} \check{x}(z) \quad \forall (x_n) \in \ell_{\mathbb{R}}^2 \text{ and} \\ \text{(3). } \mathbf{Z}(x[-n]) = \check{x}\left(\frac{1}{z}\right) \quad \forall (x_n) \in \ell_{\mathbb{R}}^2 \text{ and} \\ \text{(4). } \mathbf{Z}(x^*[n]) = \check{x}^*\left(z^*\right) \quad \forall (x_n) \in \ell_{\mathbb{R}}^2 \text{ and} \\ \text{(5). } \mathbf{Z}(x^*[-n]) = \check{x}^*\left(\frac{1}{z^*}\right) \quad \forall (x_n) \in \ell_{\mathbb{R}}^2 \end{array} \right\}$$

<sup>3</sup>Laurent series: Abramovich and Aliprantis (2002) page 49

✎ PROOF:

$$\begin{aligned}
 \alpha \mathbb{Z} \check{x}(z) &\triangleq \alpha \mathbf{Z}(\check{x}[n]) && \text{by definition of } \check{x}(z) \\
 &\triangleq \alpha \sum_{n \in \mathbb{Z}} x[n] z^{-n} && \text{by definition of } \mathbf{Z} \text{ operator} \\
 &\triangleq \sum_{n \in \mathbb{Z}} (\alpha x[n]) z^{-n} && \text{by distributive property} \\
 &\triangleq \mathbf{Z}(\alpha x[n]) && \text{by definition of } \mathbf{Z} \text{ operator} \\
 z^{-k} \check{x}(z) &= z^{-k} \mathbf{Z}(\check{x}[n]) && \text{by definition of } \check{x}(z) \quad (\text{left hypothesis}) \\
 &\triangleq z^{-k} \sum_{n=-\infty}^{n=+\infty} x[n] z^{-n} && \text{by definition of } \mathbf{Z} \quad (\text{Definition 5.4 page 28}) \\
 &= \sum_{n=-\infty}^{n=+\infty} x[n] z^{-n-k} \\
 &= \sum_{m-k=-\infty}^{m-k=+\infty} x[m-k] z^{-m} && \text{where } m \triangleq n+k \quad \implies n = m-k \\
 &= \sum_{m=-\infty}^{m=+\infty} x[m-k] z^{-m} \\
 &= \sum_{n=-\infty}^{n=+\infty} x[n-k] z^{-n} && \text{where } n \triangleq m \\
 &\triangleq \mathbf{Z}(\check{x}[n-k]) && \text{by definition of } \mathbf{Z} \quad (\text{Definition 5.4 page 28}) \\
 \mathbf{Z}(\check{x}^*[n]) &\triangleq \sum_{n \in \mathbb{Z}} x^*[n] z^{-n} && \text{by definition of } \mathbf{Z} \quad (\text{Definition 5.4 page 28}) \\
 &\triangleq \left( \sum_{n \in \mathbb{Z}} x[n] (z^*)^{-n} \right)^* && \text{by definition of } \mathbf{Z} \quad (\text{Definition 5.4 page 28}) \\
 &\triangleq \check{x}^*(z^*) && \text{by definition of } \mathbf{Z} \quad (\text{Definition 5.4 page 28}) \\
 \mathbf{Z}(\check{x}[-n]) &\triangleq \sum_{n \in \mathbb{Z}} x[-n] z^{-n} && \text{by definition of } \mathbf{Z} \quad (\text{Definition 5.4 page 28}) \\
 &= \sum_{-m \in \mathbb{Z}} x[m] z^m && \text{where } m \triangleq -n \quad \implies n = -m \\
 &= \sum_{m \in \mathbb{Z}} x[m] z^m && \text{by absolutely summable property} \quad (\text{Definition 5.2 page 27}) \\
 &= \sum_{m \in \mathbb{Z}} x[m] \left( \frac{1}{z} \right)^{-m} && \text{by absolutely summable property} \quad (\text{Definition 5.2 page 27}) \\
 &\triangleq \check{x} \left( \frac{1}{z} \right) && \text{by definition of } \mathbf{Z} \quad (\text{Definition 5.4 page 28}) \\
 \mathbf{Z}(\check{x}^*[-n]) &\triangleq \sum_{n \in \mathbb{Z}} x^*[-n] z^{-n} && \text{by definition of } \mathbf{Z} \quad (\text{Definition 5.4 page 28}) \\
 &= \sum_{-m \in \mathbb{Z}} x^*[m] z^m && \text{where } m \triangleq -n \quad \implies n = -m \\
 &= \sum_{m \in \mathbb{Z}} x^*[m] z^m && \text{by absolutely summable property} \quad (\text{Definition 5.2 page 27}) \\
 &= \sum_{m \in \mathbb{Z}} x^*[m] \left( \frac{1}{z} \right)^{-m} && \text{by absolutely summable property} \quad (\text{Definition 5.2 page 27}) \\
 &= \left( \sum_{m \in \mathbb{Z}} x[m] \left( \frac{1}{z^*} \right)^{-m} \right)^* && \text{by absolutely summable property} \quad (\text{Definition 5.2 page 27})
 \end{aligned}$$




Figure 5.1: Direct form 1 order 2 IIR filters

$$\triangleq \check{x}^* \left( \frac{1}{z^*} \right)$$

by definition of  $\mathbf{Z}$ 

(Definition 5.4 page 28)

⇒

**Theorem 5.2** (convolution theorem). *Let  $\star$  be the convolution operator (Definition 5.3 page 27).*

<b>T H M</b>	$\underbrace{\mathbf{Z}((x_n) \star (y_n))}_{\text{sequence convolution}} = \underbrace{(\mathbf{Z}(x_n)) (\mathbf{Z}(y_n))}_{\text{series multiplication}} \quad \forall (x_n)_{n \in \mathbb{Z}}, (y_n)_{n \in \mathbb{Z}} \in \ell^2_{\mathbb{R}}$
----------------------	---

PROOF:

$$[\mathbf{Z}(x \star y)](z) \triangleq \mathbf{Z} \left( \sum_{m \in \mathbb{Z}} x_m y_{n-m} \right)$$

by definition of  $\star$ 

(Definition 5.3 page 27)

$$\triangleq \sum_{n \in \mathbb{Z}} \sum_{m \in \mathbb{Z}} x_m y_{n-m} z^{-n}$$

by definition of  $\mathbf{Z}$ 

(Definition 5.4 page 28)

$$= \sum_{n \in \mathbb{Z}} \sum_{m \in \mathbb{Z}} x_m y_{n-m} z^{-n}$$

$$= \sum_{m \in \mathbb{Z}} \sum_{n \in \mathbb{Z}} x_m y_{n-m} z^{-n}$$

$$= \sum_{m \in \mathbb{Z}} \sum_{k \in \mathbb{Z}} x_m y_k z^{-(m+k)}$$

where  $k \triangleq n - m$ 

$$\iff n = m + k$$

$$= \left[ \sum_{m \in \mathbb{Z}} x_m z^{-m} \right] \left[ \sum_{k \in \mathbb{Z}} y_k z^{-k} \right]$$

$$\triangleq [\mathbf{Z}(x_n)] [\mathbf{Z}(y_n)]$$

by definition of  $\mathbf{Z}$ 

(Definition 5.4 page 28)

⇒

## 5.3 From z-domain back to time-domain

$$\check{y}(z) = b_0 X(z) + b_1 z^{-1} X(z) + b_2 z^{-2} X(z) - a_1 z^{-1} Y(z) + a_2 z^{-2} Y(z)$$

$$y[n] = b_0 x[n] + b_1 x[n-1] + b_2 x[n-2] - a_1 y[n-1] - a_2 y[n-2]$$

*Example 5.1.* See Figure 5.1 (page 30)

$$\frac{3z^2 + 5z + 7}{2z^2 + 10z + 12} = \frac{3z^2 + 5z + 7}{2(z^2 + 5z + 6)} = \frac{(3/2)z^2 + (5/2)z + 7/2}{z^2 + 5z + 6} = \frac{(3/2 + 5/2 z^{-1} + 7/2 z^{-2})}{1 + 5z^{-1} + 6z^{-2}}$$



## 5.4 Zero locations

The system property of *minimum phase* is defined in Definition 5.5 (next) and illustrated in Figure 5.2 (page 31).




Figure 5.2: Minimum Phase filter

**Definition 5.5.** <sup>4</sup> Let  $\check{x}(z) \triangleq \mathbf{Z}(x_n)$  be the Z TRANSFORM (Definition 5.4 page 28) of a sequence  $(x_n)_{n \in \mathbb{Z}}$  in  $\ell^2_{\mathbb{R}}$ . Let  $(z_n)_{n \in \mathbb{Z}}$  be the ZEROS of  $\check{x}(z)$ .

The sequence  $(x_n)$  is **minimum phase** if

$$|z_n| < 1 \quad \forall n \in \mathbb{Z}$$

$\check{x}(z)$  has all its ZEROS inside the unit circle

The impulse response of a minimum phase filter has most of its energy concentrated near the beginning of its support, as demonstrated next.

**Theorem 5.3** (Robinson's Energy Delay Theorem). <sup>5</sup> Let  $p(z) \triangleq \sum_{n=0}^N a_n z^{-n}$  and  $q(z) \triangleq \sum_{n=0}^N b_n z^{-n}$  be polynomials.

$$\left\{ \begin{array}{l} p \text{ is MINIMUM PHASE} \\ q \text{ is NOT minimum phase} \end{array} \right. \text{ and } \left\{ \begin{array}{l} p \text{ is NOT minimum phase} \\ q \text{ is MINIMUM PHASE} \end{array} \right. \Rightarrow \underbrace{\sum_{n=0}^{m-1} |a_n|^2}_{\substack{\text{"energy" of} \\ \text{the first } m \text{ co-} \\ \text{efficients of} \\ p(z)}} \geq \underbrace{\sum_{n=0}^{m-1} |b_n|^2}_{\substack{\text{"energy" of} \\ \text{the first } m \text{ co-} \\ \text{efficients of} \\ q(z)}} \quad \forall 0 \leq m \leq N$$

But for more *symmetry*, put some zeros inside and some outside the unit circle (Figure 5.3 page 32).

**Example 5.2.** An example of a minimum phase polynomial is the Daubechies-4 scaling function. The minimum phase polynomial causes most of the energy to be concentrated near the origin, making it very *asymmetric*. In contrast, the Symlet-4 has a design very similar to that of Daubechies-4, but the selected zeros are not all within the unit circle in the complex  $z$  plane. This results in a scaling function that is more symmetric and less contrated near the origin. Both scaling functions are illustrated in Figure 5.3 (page 32).

<sup>4</sup> Farina and Rinaldi (2000) page 91, Dumitrescu (2007) page 36

<sup>5</sup> Dumitrescu (2007) page 36, Robinson (1962), Robinson (1966)  $\langle ??? \rangle$ , Claerbout (1976) pages 52–53




Figure 5.3: Daubechies-4 and Symlet-4 scaling functions pole-zero plots

## 5.5 Pole locations

### Definition 5.6.

**DEF** A filter (or system or operator)  $\mathbf{H}$  is **causal** if its current output does not depend on future inputs.

### Definition 5.7.

**DEF** A filter (or system or operator)  $\mathbf{H}$  is **time-invariant** if the mapping it performs does not change with time.

### Definition 5.8.

**DEF** An operation  $\mathbf{H}$  is **linear** if any output  $y_n$  can be described as a linear combination of inputs  $x_n$  as in

$$y_n = \sum_{m \in \mathbb{Z}} h(m)x(n - m).$$

For a filter to be *stable*, place all the poles *inside* the unit circle.

**Theorem 5.4.** A causal LTI filter is **stable** if all of its poles are **inside** the unit circle.

*Example 5.3.* Stable/unstable filters are illustrated in Figure 5.4 (page 33).

True or False? This filter has no poles:

$$H(z) = b_0 + b_1 z^{-1} + b_2 z^{-2}$$

$$H(z) = b_0 + b_1 z^{-1} + b_2 z^{-2} = \frac{z^2}{z^2} \times \frac{b_0 + b_1 z^{-1} + b_2 z^{-2}}{1} = \frac{b_0 z^2 + b_1 z^1 + b_2}{z^2}$$




Figure 5.4: Pole-zero plot stable/unstable causal LTI filters (Example 5.3 page 32)




Figure 5.5: FIR filters

## 5.6 Mirroring for real coefficients

If you want real coefficients, choose poles and zeros in conjugate pairs (next).

### Proposition 5.3.

$$\left\{ \begin{array}{l} \text{ZEROS and POLES} \\ \text{occur in CONJUGATE PAIRS} \end{array} \right\} \Rightarrow \left\{ \begin{array}{l} \text{COEFFICIENTS} \\ \text{are REAL.} \end{array} \right\}$$

PROOF:

$$\begin{aligned} (z - p_1)(z - p_1^*) &= [z - (a + ib)][z - (a - ib)] \\ &= z^2 + [-a + ib - ib - a]z - [ib]^2 \end{aligned}$$




Figure 5.6: Conjugate pair structure yielding real coefficients

$$= z^2 - 2az + b^2$$



*Example 5.4.* See Figure 5.6 (page 33).

$$\begin{aligned} H(z) &= G \frac{[z - z_1][z - z_2]}{[z - p_1][z - p_2]} = G \frac{[z - (1 + i)][z - (1 - i)]}{[z - (-\frac{2}{3} + i\frac{1}{2})][z - (-\frac{2}{3} - i\frac{1}{2})]} \\ &= G \frac{z^2 - z[(1 - i) + (1 + i)] + (1 - i)(1 + i)}{z^2 - z[(-\frac{2}{3} + i\frac{1}{2}) + (-\frac{2}{3} - i\frac{1}{2})] + (-\frac{2}{3} + i\frac{1}{2})(-\frac{2}{3} - i\frac{1}{2})} \\ &= G \frac{z^2 - 2z + 2}{z^2 - \frac{4}{3}z + (\frac{4}{9} + \frac{1}{4})} = G \frac{z^2 - 2z + 2}{z^2 - \frac{4}{3}z + \frac{19}{12}} \end{aligned}$$

## 5.7 Rational polynomial operators

A digital filter is simply an operator on  $\ell_{\mathbb{R}}^2$ . If the digital filter is a causal LTI system, then it can be expressed as a rational polynomial in  $z$  as shown next.

**Lemma 5.1.** *A causal LTI operator  $\mathbf{H}$  can be expressed as a rational expression  $\check{h}(z)$ .*

$$\begin{aligned} \check{h}(z) &= \frac{b_0 + b_1 z^{-1} + b_2 z^{-2} + \cdots + b_N z^{-N}}{1 + a_1 z^{-1} + a_2 z^{-2} + \cdots + a_N z^{-N}} \\ &= \frac{\sum_{n=0}^N b_n z^{-n}}{1 + \sum_{n=1}^N a_n z^{-n}} \end{aligned}$$

A filter operation  $\check{h}(z)$  can be expressed as a product of its roots (poles and zeros).

$$\begin{aligned} \check{h}(z) &= \frac{b_0 + b_1 z^{-1} + b_2 z^{-2} + \cdots + b_N z^{-N}}{1 + a_1 z^{-1} + a_2 z^{-2} + \cdots + a_N z^{-N}} \\ &= \alpha \frac{(z - z_1)(z - z_2) \cdots (z - z_N)}{(z - p_1)(z - p_2) \cdots (z - p_N)} \end{aligned}$$

where  $\alpha$  is a constant,  $z_i$  are the zeros, and  $p_i$  are the poles. The poles and zeros of such a rational expression are often plotted in the  $z$ -plane with a unit circle about the origin (representing  $z = e^{i\omega}$ ). Poles are marked with  $\times$  and zeros with  $\circ$ . An example is shown in Figure 5.7 page 35. Notice that in this figure the zeros and poles are either real or occur in complex conjugate pairs.

## 5.8 Filter Banks

*Conjugate quadrature filters* (next definition) are used in *filter banks*. If  $\check{x}(z)$  is a *low-pass filter*, then the conjugate quadrature filter of  $\check{y}(z)$  is a *high-pass filter*.




Figure 5.7: Pole-zero plot for rational expression with real coefficients

**Definition 5.9.** <sup>6</sup> Let  $(x_n)_{n \in \mathbb{Z}}$  and  $(y_n)_{n \in \mathbb{Z}}$  be SEQUENCES (Definition 5.1 page 27) in  $\ell^2_{\mathbb{R}}$  (Definition 5.2 page 27).

The sequence  $(y_n)$  is a **conjugate quadrature filter** with shift  $N$  with respect to  $(x_n)$  if

$$y_n = \pm(-1)^n x_{N-n}^*$$

A CONJUGATE QUADRATURE FILTER is also called a **CQF** or a **Smith-Barnwell filter**.

Any triple  $((x_n), (y_n), N)$  in this form is said to satisfy the

**conjugate quadrature filter condition** or the **CQF condition**.

**Theorem 5.5** (CQF theorem). <sup>7</sup> Let  $\check{y}(\omega)$  and  $\check{x}(\omega)$  be the DTFTs (Definition 6.1 page 41) of the sequences  $(y_n)_{n \in \mathbb{Z}}$  and  $(x_n)_{n \in \mathbb{Z}}$ , respectively, in  $\ell^2_{\mathbb{R}}$  (Definition 5.2 page 27).

T H M	$\underbrace{y_n = \pm(-1)^n x_{N-n}^*}_{(1) \text{ CQF in "time"}} \iff \check{y}(z) = \pm(-1)^N z^{-N} \check{x}^*\left(\frac{-1}{z^*}\right) \quad (2) \text{ CQF in "z-domain"}$
	$\iff \check{y}(\omega) = \pm(-1)^N e^{-i\omega N} \check{x}^*(\omega + \pi) \quad (3) \text{ CQF in "frequency"}$
	$\iff x_n = \pm(-1)^N (-1)^n y_{N-n}^* \quad (4) \text{ "reversed" CQF in "time"}$
	$\iff \check{x}(z) = \pm z^{-N} \check{y}^*\left(\frac{-1}{z^*}\right) \quad (5) \text{ "reversed" CQF in "z-domain"}$
	$\iff \check{x}(\omega) = \pm e^{-i\omega N} \check{y}^*(\omega + \pi) \quad (6) \text{ "reversed" CQF in "frequency"}$
	$\forall n \in \mathbb{Z}$

PROOF:

1. Proof that (1)  $\implies$  (2):

$$\begin{aligned}
 \check{y}(z) &= \sum_{n \in \mathbb{Z}} y_n z^{-n} && \text{by definition of } z\text{-transform} \quad (\text{Definition 5.4 page 28}) \\
 &= \sum_{n \in \mathbb{Z}} \underbrace{(\pm)(-1)^n x_{N-n}^*}_{\text{CQF}} z^{-n} && \text{by (1)} \\
 &= \pm \sum_{m \in \mathbb{Z}} (-1)^{N-m} x_m^* z^{-(N-m)} && \text{where } m \triangleq N - n \implies n = N - m \\
 &= \pm(-1)^N z^{-N} \sum_{m \in \mathbb{Z}} (-1)^{-m} x_m^* (z^{-1})^{-m} \\
 &= \pm(-1)^N z^{-N} \sum_{m \in \mathbb{Z}} x_m^* \left(-\frac{1}{z}\right)^{-m} \\
 &= \pm(-1)^N z^{-N} \left[ \sum_{m \in \mathbb{Z}} x_m \left(-\frac{1}{z^*}\right)^{-m} \right]^*
 \end{aligned}$$

<sup>6</sup> Strang and Nguyen (1996) page 109, Haddad and Akansu (1992) pages 256–259 (section 4.5), Vaidyanathan (1993) page 342 ((7.2.7), (7.2.8)), Smith and Barnwell (1984a), Smith and Barnwell (1984b), Mintzer (1985)

<sup>7</sup> Strang and Nguyen (1996) page 109, Mallat (1999) pages 236–238 ((7.58), (7.73)), Haddad and Akansu (1992) pages 256–259 (section 4.5), Vaidyanathan (1993) page 342 ((7.2.7), (7.2.8))

$$= \pm(-1)^N z^{-N} \check{x}^* \left( \frac{-1}{z^*} \right)$$

by definition of  $z$ -transform

(Definition 5.4 page 28)

2. Proof that (1)  $\Leftarrow$  (2):

$$\check{y}(z) = \pm(-1)^N z^{-N} \check{x}^* \left( \frac{-1}{z^*} \right)$$

by (2)

$$= \pm(-1)^N z^{-N} \left[ \sum_{m \in \mathbb{Z}} x_m \left( \frac{-1}{z^*} \right)^{-m} \right]^*$$

by definition of  $z$ -transform

(Definition 5.4 page 28)

$$= \pm(-1)^N z^{-N} \left[ \sum_{m \in \mathbb{Z}} x_m^* (-z^{-1})^{-m} \right]$$

by definition of  $z$ -transform

(Definition 5.4 page 28)

$$= \sum_{m \in \mathbb{Z}} (\pm)(-1)^{N-m} x_m^* z^{-(N-m)}$$

$$= \sum_{m \in \mathbb{Z}} (\pm)(-1)^n x_{N-n}^* z^{-n}$$

where  $n = N - m \implies$  $m \triangleq N - n$ 

$$\implies x_n = \pm(-1)^n x_{N-n}^*$$

3. Proof that (1)  $\implies$  (3):

$$\check{y}(\omega) \triangleq \check{x}(z) \Big|_{z=e^{i\omega}}$$

by definition of  $DTFT$  (Definition 6.1 page 41)

$$= \left[ \pm(-1)^N z^{-N} \check{x} \left( \frac{-1}{z^*} \right) \right]_{z=e^{i\omega}}$$

by (2)

$$= \pm(-1)^N e^{-i\omega N} \check{x}(e^{i\pi} e^{i\omega})$$

$$= \pm(-1)^N e^{-i\omega N} \check{x}(e^{i(\omega+\pi)})$$

$$= \pm(-1)^N e^{-i\omega N} \check{x}(\omega + \pi)$$

by definition of  $DTFT$  (Definition 6.1 page 41)4. Proof that (1)  $\implies$  (6):

$$\check{x}(\omega) = \sum_{n \in \mathbb{Z}} y_n e^{-i\omega n}$$

by definition of  $DTFT$ 

(Definition 6.1 page 41)

$$= \sum_{n \in \mathbb{Z}} \underbrace{\pm(-1)^n x_{N-n}^*}_{CQF} e^{-i\omega n}$$

by (1)

$$= \sum_{m \in \mathbb{Z}} \pm(-1)^{N-m} x_m^* e^{-i\omega(N-m)}$$

where  $m \triangleq N - n \implies$  $n = N - m$ 

$$= \pm(-1)^N e^{-i\omega N} \sum_{m \in \mathbb{Z}} (-1)^m x_m^* e^{i\omega m}$$

$$= \pm(-1)^N e^{-i\omega N} \sum_{m \in \mathbb{Z}} e^{i\pi m} x_m^* e^{i\omega m}$$

$$= \pm(-1)^N e^{-i\omega N} \sum_{m \in \mathbb{Z}} x_m^* e^{i(\omega+\pi)m}$$

$$= \pm(-1)^N e^{-i\omega N} \left[ \sum_{m \in \mathbb{Z}} x_m e^{-i(\omega+\pi)m} \right]^*$$

$$= \pm(-1)^N e^{-i\omega N} \check{x}^*(\omega + \pi)$$

by definition of  $DTFT$ 

(Definition 6.1 page 41)

5. Proof that (1)  $\Leftarrow$  (3):

$$\begin{aligned}
 y_n &= \frac{1}{2\pi} \int_{-\pi}^{+\pi} \check{y}(\omega) e^{i\omega n} d\omega && \text{by inverse DTFT} && (\text{Theorem 6.3 page 47}) \\
 &= \frac{1}{2\pi} \int_{-\pi}^{+\pi} \underbrace{\pm (-1)^N e^{-iN\omega} \check{x}^*(\omega + \pi)}_{\text{right hypothesis}} e^{i\omega n} d\omega && \text{by right hypothesis} \\
 &= \pm (-1)^N \frac{1}{2\pi} \int_{-\pi}^{+\pi} \check{x}^*(\omega + \pi) e^{i\omega(n-N)} d\omega && \text{by right hypothesis} \\
 &= \pm (-1)^N \frac{1}{2\pi} \int_0^{2\pi} \check{x}^*(v) e^{i(v-\pi)(n-N)} dv && \text{where } v \triangleq \omega + \pi \implies \omega = v - \pi \\
 &= \pm (-1)^N e^{-i\pi(n-N)} \frac{1}{2\pi} \int_0^{2\pi} \check{x}^*(v) e^{iv(n-N)} dv \\
 &= \pm (-1)^N \underbrace{(-1)^N}_{e^{i\pi N}} \underbrace{(-1)^n}_{e^{-i\pi n}} \left[ \frac{1}{2\pi} \int_0^{2\pi} \check{x}(v) e^{iv(N-n)} dv \right]^* \\
 &= \pm (-1)^n x_{N-n}^* && \text{by inverse DTFT} && (\text{Theorem 6.3 page 47})
 \end{aligned}$$

6. Proof that (1)  $\Leftrightarrow$  (4):

$$\begin{aligned}
 y_n = \pm (-1)^n x_{N-n}^* &\Leftrightarrow (\pm)(-1)^n y_n = (\pm)(\pm)(-1)^n (-1)^n x_{N-n}^* \\
 &\Leftrightarrow \pm (-1)^n y_n = x_{N-n}^* \\
 &\Leftrightarrow (\pm(-1)^n y_n)^* = (x_{N-n}^*)^* \\
 &\Leftrightarrow \pm (-1)^n y_n^* = x_{N-n} \\
 &\Leftrightarrow x_{N-n} = \pm (-1)^n y_n^* \\
 &\Leftrightarrow x_m = \pm (-1)^{N-m} y_{N-m}^* && \text{where } m \triangleq N - n \implies n = N - m \\
 &\Leftrightarrow x_m = \pm (-1)^{N-m} y_{N-m}^* \\
 &\Leftrightarrow x_m = \pm (-1)^N (-1)^m y_{N-m}^* \\
 &\Leftrightarrow x_n = \pm (-1)^N (-1)^n y_{N-n}^* && \text{by change of free variables}
 \end{aligned}$$

7. Proofs for (5) and (6): not included. See proofs for (2) and (3).

$\Rightarrow$

**Theorem 5.6.**<sup>8</sup> Let  $\check{y}(\omega)$  and  $\check{x}(\omega)$  be the DTFTs (Definition 6.1 page 41) of the sequences  $(y_n)_{n \in \mathbb{Z}}$  and  $(x_n)_{n \in \mathbb{Z}}$ , respectively, in  $\ell_{\mathbb{R}}^2$  (Definition 5.2 page 27).

T H M	Let $y_n = \pm (-1)^n x_{N-n}^*$ (CQF CONDITION, Definition 5.9 page 35). Then						
	{	(A)	$\left[ \frac{d}{d\omega} \right]^n \check{y}(\omega) \Big _{\omega=0} = 0$	$\Leftrightarrow$	$\left[ \frac{d}{d\omega} \right]^n \check{x}(\omega) \Big _{\omega=\pi} = 0$	(B)	$\forall n \in \mathbb{W}$
				$\Leftrightarrow$	$\sum_{k \in \mathbb{Z}} (-1)^k k^n x_k = 0$	(C)	
				$\Leftrightarrow$	$\sum_{k \in \mathbb{Z}} k^n y_k = 0$	(D)	

PROOF:

<sup>8</sup> Vidakovic (1999) pages 82–83, Mallat (1999) pages 241–242

1. Proof that (A)  $\implies$  (B):

$$\begin{aligned}
 0 &= \left[ \frac{d}{d\omega} \right]^n \check{y}(\omega) \Big|_{\omega=0} && \text{by (A)} \\
 &= \left[ \frac{d}{d\omega} \right]^n (\pm)(-1)^N e^{-i\omega N} \check{x}^*(\omega + \pi) \Big|_{\omega=0} && \text{by CQF theorem (Theorem 5.5 page 35)} \\
 &= (\pm)(-1)^N \sum_{\ell=0}^n \binom{n}{\ell} \left[ \frac{d}{d\omega} \right]^\ell [e^{-i\omega N}] \cdot \left[ \frac{d}{d\omega} \right]^{n-\ell} [\check{x}^*(\omega + \pi)] \Big|_{\omega=0} && \text{by Leibnitz GPR (Lemma B.2 page 71)} \\
 &= (\pm)(-1)^N \sum_{\ell=0}^n \binom{n}{\ell} (-iN)^\ell e^{-i\omega N} \left[ \frac{d}{d\omega} \right]^{n-\ell} [\check{x}^*(\omega + \pi)] \Big|_{\omega=0} \\
 &= (\pm)(-1)^N e^{-i0N} \sum_{\ell=0}^n \binom{n}{\ell} (-iN)^\ell \left[ \frac{d}{d\omega} \right]^{n-\ell} [\check{x}^*(\omega + \pi)] \Big|_{\omega=0} \\
 &\implies \check{x}^{(0)}(\pi) = 0 \\
 &\implies \check{x}^{(1)}(\pi) = 0 \\
 &\implies \check{x}^{(2)}(\pi) = 0 \\
 &\implies \check{x}^{(3)}(\pi) = 0 \\
 &\implies \check{x}^{(4)}(\pi) = 0 \\
 &\quad \vdots \\
 &\implies \check{x}^{(n)}(\pi) = 0 \quad \text{for } n = 0, 1, 2, \dots
 \end{aligned}$$

2. Proof that (A)  $\Leftarrow$  (B):

$$\begin{aligned}
 0 &= \left[ \frac{d}{d\omega} \right]^n \check{x}(\omega) \Big|_{\omega=\pi} && \text{by (B)} \\
 &= \left[ \frac{d}{d\omega} \right]^n (\pm) e^{-i\omega N} \check{y}^*(\omega + \pi) \Big|_{\omega=\pi} && \text{by CQF theorem (Theorem 5.5 page 35)} \\
 &= (\pm) \sum_{\ell=0}^n \binom{n}{\ell} \left[ \frac{d}{d\omega} \right]^\ell [e^{-i\omega N}] \cdot \left[ \frac{d}{d\omega} \right]^{n-\ell} [\check{y}^*(\omega + \pi)] \Big|_{\omega=\pi} && \text{by Leibnitz GPR (Lemma B.2 page 71)} \\
 &= (\pm) \sum_{\ell=0}^n \binom{n}{\ell} (-iN)^\ell e^{-i\omega N} \left[ \frac{d}{d\omega} \right]^{n-\ell} [\check{y}^*(\omega + \pi)] \Big|_{\omega=\pi} \\
 &= (\pm) e^{-i\pi N} \sum_{\ell=0}^n \binom{n}{\ell} (-iN)^\ell \left[ \frac{d}{d\omega} \right]^{n-\ell} [\check{y}^*(\omega + \pi)] \Big|_{\omega=\pi} \\
 &= (\pm)(-1)^N \sum_{\ell=0}^n \binom{n}{\ell} (-iN)^\ell \left[ \frac{d}{d\omega} \right]^{n-\ell} [\check{y}^*(\omega + \pi)] \Big|_{\omega=\pi} \\
 &\implies \check{y}^{(0)}(0) = 0 \\
 &\implies \check{y}^{(1)}(0) = 0 \\
 &\implies \check{y}^{(2)}(0) = 0 \\
 &\implies \check{y}^{(3)}(0) = 0 \\
 &\implies \check{y}^{(4)}(0) = 0 \\
 &\quad \vdots \\
 &\implies \check{y}^{(n)}(0) = 0 \\
 &\implies \check{y}^{(n)}(0) = 0 \quad \text{for } n = 0, 1, 2, \dots
 \end{aligned}$$


3. Proof that (B)  $\iff$  (C): by Theorem 6.5 page 49

4. Proof that (A)  $\iff$  (D): by Theorem 6.5 page 49

5. Proof that (CQF)  $\nLeftarrow$  (A): Here is a counterexample:  $\check{y}(\omega) = 0$ .








$$\frac{(z - z_1)(z - z_2)(z - z_3)(z - z_4)}{(z - p_1)(z - p_2)(z - p_3)(z - p_4)} \times \frac{(z - p_1)(z - p_2)(z - p_3)(z - p_4)}{(z - z_1)(z - z_2)(z - z_3)(z - z_4)} = 1$$

## 5.9 Inverting non-minimum phase filters

*Minimum phase* filters are easy to invert: each *zero* becomes a *pole* and each *pole* becomes a *zero*.



$$\begin{aligned}
 |A(z)|_{z=e^{i\omega}} &= \frac{1}{r} \left| \frac{z - re^{i\phi}}{z - \frac{1}{r}e^{i\phi}} \right|_{z=e^{i\omega}} = \left| \frac{z - re^{i\phi}}{rz - e^{i\phi}} \right|_{z=e^{i\omega}} \\
 &= \left| e^{i\phi} \left( \frac{e^{-i\phi}z - r}{rz - e^{i\phi}} \right) \right|_{z=e^{i\omega}} = \left| z \left( \frac{e^{-i\phi} - rz^{-1}}{rz - e^{i\phi}} \right) \right|_{z=e^{i\omega}} \\
 &= \left| -z \left( \frac{rz^{-1} - e^{-i\phi}}{rz - e^{i\phi}} \right) \right|_{z=e^{i\omega}} = \left| \underbrace{e^{i\pi}}_{-1} e^{i\omega} \left( \frac{re^{-i\omega} - e^{-i\phi}}{re^{i\omega} - e^{i\phi}} \right) \right| \\
 &= \left| \frac{1}{e^{-i\omega}} \left( \frac{re^{-i\omega} - e^{-i\phi}}{(re^{i\omega} - e^{i\phi})^*} \right) \right| = \left| \frac{re^{-i\omega} - e^{-i\phi}}{re^{-i\omega} - e^{-i\phi}} \right| = 1
 \end{aligned}$$



# CHAPTER 6

## DISCRETE TIME FOURIER TRANSFORM

### 6.1 Definition


#### Definition 6.1.

DEF

The **discrete-time Fourier transform**  $\check{\mathbf{F}}$  of  $(x_n)_{n \in \mathbb{Z}}$  is defined as

$$[\check{\mathbf{F}}((x_n))](\omega) \triangleq \sum_{n \in \mathbb{Z}} x_n e^{-i\omega n} \quad \forall (x_n)_{n \in \mathbb{Z}} \in \ell^2_{\mathbb{R}}$$

If we compare the definition of the *Discrete Time Fourier Transform* (Definition 6.1 page 41) to the definition of the Z-transform (Definition 5.4 page 28), we see that the DTFT is just a special case of the more general Z-Transform, with  $z = e^{i\omega}$ . If we imagine  $z \in \mathbb{C}$  as a complex plane, then  $e^{i\omega}$  is a unit circle in this plane. The “frequency”  $\omega$  in the DTFT is the unit circle in the much larger  $z$ -plane, as illustrated to the right.



### 6.2 Properties

**Proposition 6.1** (DTFT periodicity). Let  $\check{x}(\omega) \triangleq \check{\mathbf{F}}[(x_n)](\omega)$  be the DISCRETE-TIME FOURIER TRANSFORM (Definition 6.1 page 41) of a sequence  $(x_n)_{n \in \mathbb{Z}}$  in  $\ell^2_{\mathbb{R}}$ .

PRP

$$\underbrace{\check{x}(\omega) = \check{x}(\omega + 2\pi n)}_{\text{PERIODIC with period } 2\pi} \quad \forall n \in \mathbb{Z}$$

PROOF:

$$\begin{aligned} \check{x}(\omega + 2\pi n) &= \sum_{m \in \mathbb{Z}} x_m e^{-i(\omega + 2\pi n)m} \\ &= \sum_{m \in \mathbb{Z}} x_m e^{-i\omega m} e^{-i2\pi nm} \xrightarrow{1} \\ &= \sum_{m \in \mathbb{Z}} x_m e^{-i\omega m} \\ &= \check{x}(\omega) \end{aligned}$$

**Theorem 6.1.** Let  $\tilde{x}(\omega) \triangleq \check{\mathbf{F}}[(x[n])](\omega)$  be the DISCRETE-TIME FOURIER TRANSFORM (Definition 6.1 page 41) of a sequence  $(x_n)_{n \in \mathbb{Z}}$  in  $\ell^2_{\mathbb{R}}$ .

$$\text{THM} \quad \left\{ \begin{array}{l} \tilde{x}(\omega) \triangleq \check{\mathbf{F}}[(x[n])] \end{array} \right\} \implies \left\{ \begin{array}{l} (1). \quad \check{\mathbf{F}}[(x[-n])] = \tilde{x}(-\omega) \quad \text{and} \\ (2). \quad \check{\mathbf{F}}[(x^*[n])] = \tilde{x}^*(-\omega) \quad \text{and} \\ (3). \quad \check{\mathbf{F}}[(x^*[-n])] = \tilde{x}^*(\omega) \end{array} \right\}$$

PROOF:

$$\begin{aligned} \check{\mathbf{F}}[(x[-n])] &\triangleq \sum_{n \in \mathbb{Z}} x[-n] e^{-i\omega n} && \text{by definition of DTFT} && (\text{Definition 6.1 page 41}) \\ &= \sum_{m \in \mathbb{Z}} x[m] e^{i\omega m} && \text{where } m \triangleq -n \implies n = -m \\ &= \sum_{m \in \mathbb{Z}} x[m] e^{-i(-\omega)m} \\ &\triangleq \tilde{x}(-\omega) && \text{by left hypothesis} \end{aligned}$$

$$\begin{aligned} \check{\mathbf{F}}[(x^*[n])] &\triangleq \sum_{n \in \mathbb{Z}} x^*[n] e^{-i\omega n} && \text{by definition of DTFT} && (\text{Definition 6.1 page 41}) \\ &= \left( \sum_{n \in \mathbb{Z}} x[n] e^{i\omega n} \right)^* && \text{by distributive property of *-algebras} && (\text{Definition E.3 page 114}) \\ &= \left( \sum_{n \in \mathbb{Z}} x[n] e^{-i(-\omega)n} \right)^* \\ &\triangleq \tilde{x}^*(-\omega) && \text{by left hypothesis} \end{aligned}$$

$$\begin{aligned} \check{\mathbf{F}}[(x^*[-n])] &\triangleq \sum_{n \in \mathbb{Z}} x^*[-n] e^{-i\omega n} && \text{by definition of DTFT} && (\text{Definition 6.1 page 41}) \\ &= \left( \sum_{n \in \mathbb{Z}} x[-n] e^{i\omega n} \right)^* && \text{by distributive property of *-algebras} && (\text{Definition E.3 page 114}) \\ &= \left( \sum_{m \in \mathbb{Z}} x[m] e^{-i\omega m} \right)^* && \text{where } m \triangleq -n \implies n = -m \\ &\triangleq \tilde{x}^*(\omega) && \text{by left hypothesis} \end{aligned}$$

⇒

**Theorem 6.2.** Let  $\tilde{x}(\omega) \triangleq \check{\mathbf{F}}[(x[n])](\omega)$  be the DISCRETE-TIME FOURIER TRANSFORM (Definition 6.1 page 41) of a sequence  $(x[n])_{n \in \mathbb{Z}}$  in  $\ell^2_{\mathbb{R}}$ .

$$\text{THM} \quad \left\{ \begin{array}{l} (1). \quad \tilde{x}(\omega) \triangleq \check{\mathbf{F}}[(x[n])] \\ (2). \quad (x[n]) \text{ is REAL-VALUED} \end{array} \right\} \text{ and } \implies \left\{ \begin{array}{l} (1). \quad \check{\mathbf{F}}[(x[-n])] = \tilde{x}(-\omega) \quad \text{and} \\ (2). \quad \check{\mathbf{F}}[(x^*[n])] = \tilde{x}^*(-\omega) = \tilde{x}(\omega) \quad \text{and} \\ (3). \quad \check{\mathbf{F}}[(x^*[-n])] = \tilde{x}^*(\omega) = \tilde{x}(-\omega) \end{array} \right\}$$

PROOF:

$$\begin{aligned} \check{\mathbf{F}}[(x[-n])] &\triangleq \sum_{n \in \mathbb{Z}} x[-n] e^{-i\omega n} && \text{by definition of DTFT} && (\text{Definition 6.1 page 41}) \\ &= \sum_{m \in \mathbb{Z}} x[m] e^{i\omega m} && \text{where } m \triangleq -n \implies n = -m \\ &= \sum_{m \in \mathbb{Z}} x[m] e^{-i(-\omega)m} \end{aligned}$$

$$\triangleq \check{x}(-\omega)$$

by left hypothesis

$$\begin{aligned}\check{x}^*(-\omega) &= \check{\mathbf{F}}(\mathbf{x}^*[n]) \\ &= \check{\mathbf{F}}(\mathbf{x}[n]) \\ &= \check{x}(\omega)\end{aligned}$$

by Theorem 6.1 page 42

by *real-valued* hypothesis

by definition of  $\check{x}(\omega)$


(Definition 6.1 page 41)

$$\begin{aligned}\check{x}^*(\omega) &= \check{\mathbf{F}}(\mathbf{x}^*[-n]) \\ &= \check{\mathbf{F}}(\mathbf{x}[-n]) \\ &= \check{x}(-\omega)\end{aligned}$$

by Theorem 6.1 page 42

by *real-valued* hypothesis

by result (1)



**Proposition 6.2.** Let  $\check{x}(z)$  be the Z-TRANSFORM (Definition 5.4 page 28) and  $\check{x}(\omega)$  the DISCRETE-TIME FOURIER TRANSFORM (Definition 6.1 page 41) of  $(x_n)$ .

P R P	$\underbrace{\left\{ \sum_{n \in \mathbb{Z}} x_n = c \right\}}_{(1) \text{ time domain}} \iff \underbrace{\left\{ \check{x}(z) \Big _{z=1} = c \right\}}_{(2) \text{ } z \text{ domain}} \iff \underbrace{\left\{ \check{x}(\omega) \Big _{\omega=0} = c \right\}}_{(3) \text{ frequency domain}} \quad \forall (x_n)_{n \in \mathbb{Z}} \in \ell^2_{\mathbb{R}}, c \in \mathbb{R}$
-------------	---

PROOF:

1. Proof that (1)  $\implies$  (2):

$$\begin{aligned}\check{x}(z) \Big|_{z=1} &= \sum_{n \in \mathbb{Z}} x_n z^{-n} \Big|_{z=1} \\ &= \sum_{n \in \mathbb{Z}} x_n \\ &= c\end{aligned}$$

by definition of  $\check{x}(z)$  (Definition 5.4 page 28)

because  $z^n = 1$  for all  $n \in \mathbb{Z}$

by hypothesis (1)

2. Proof that (2)  $\implies$  (3):

$$\begin{aligned}\check{x}(\omega) \Big|_{\omega=0} &= \sum_{n \in \mathbb{Z}} x_n e^{-i\omega n} \Big|_{\omega=0} \\ &= \sum_{n \in \mathbb{Z}} x_n z^{-n} \Big|_{z=1} \\ &= \check{x}(z) \Big|_{z=1} \\ &= c\end{aligned}$$

by definition of  $\check{x}(\omega)$

(Definition 6.1 page 41)

by definition of  $\check{x}(z)$

(Definition 5.4 page 28)

by hypothesis (2)

3. Proof that (3)  $\implies$  (1):

$$\begin{aligned} \sum_{n \in \mathbb{Z}} x_n &= \sum_{n \in \mathbb{Z}} x_n e^{-i\omega n} \Big|_{\omega=0} \\ &= \check{x}(\omega) && \text{by definition of } \check{x}(\omega) && \text{(Definition 6.1 page 41)} \\ &= c && \text{by hypothesis (3)} \end{aligned}$$

$\Rightarrow$

**Proposition 6.3.** *If the coefficients are **real**, then the magnitude response (MR) is **symmetric**.*

$\Rightarrow$  PROOF:

$$\begin{aligned} |\tilde{h}(-\omega)| &\triangleq |\check{h}(z)|_{z=e^{-i\omega}} && \triangleq \left| \sum_{m \in \mathbb{Z}} x[m] z^{-m} \right|_{z=e^{-i\omega}} \\ &= \left| \sum_{m \in \mathbb{Z}} x[m] e^{i\omega m} \right| && = \left| \left( \sum_{m \in \mathbb{Z}} x^*[m] e^{-i\omega m} \right)^* \right| \\ &= \left| \underbrace{\left( \sum_{m \in \mathbb{Z}} x[m] e^{-i\omega m} \right)^*}_{\text{if } x[m] \text{ is real}} \right| && = \left| \sum_{m \in \mathbb{Z}} x[m] e^{-i\omega m} \right| \\ &\triangleq |\check{h}(z)|_{z=e^{-i\omega}} && \triangleq |\tilde{h}(\omega)| \end{aligned}$$

$\Rightarrow$

**Proposition 6.4.** <sup>1</sup>

P  
R  
P

$$\underbrace{\sum_{n \in \mathbb{Z}} (-1)^n x_n = c}_{(1) \text{ in "time"}} \iff \underbrace{\check{x}(z)|_{z=-1} = c}_{(2) \text{ in "z domain"}} \iff \underbrace{\check{x}(\omega)|_{\omega=\pi} = c}_{(3) \text{ in "frequency"}}$$

$$\iff \underbrace{\left( \sum_{n \in \mathbb{Z}} h_{2n}, \sum_{n \in \mathbb{Z}} h_{2n+1} \right) = \left( \frac{1}{2} \left( \sum_{n \in \mathbb{Z}} h_n + c \right), \frac{1}{2} \left( \sum_{n \in \mathbb{Z}} h_n - c \right) \right)}_{(4) \text{ sum of even, sum of odd}}$$

$\forall c \in \mathbb{R}, (x_n)_{n \in \mathbb{Z}}, (y_n)_{n \in \mathbb{Z}} \in \ell^2_{\mathbb{R}}$

$\Rightarrow$  PROOF:

1. Proof that (1)  $\implies$  (2):

$$\begin{aligned} \check{x}(z)|_{z=-1} &= \sum_{n \in \mathbb{Z}} x_n z^{-n} \Big|_{z=-1} \\ &= \sum_{n \in \mathbb{Z}} (-1)^n x_n \\ &= c && \text{by (1)} \end{aligned}$$

<sup>1</sup> Chui (1992) page 123

2. Proof that (2)  $\implies$  (3):

$$\begin{aligned}
 \left. \sum_{n \in \mathbb{Z}} x_n e^{-i\omega n} \right|_{\omega=\pi} &= \sum_{n \in \mathbb{Z}} (-1)^n x_n \\
 &= \sum_{n \in \mathbb{Z}} (-1)^{-n} x_n &= \sum_{n \in \mathbb{Z}} z^{-n} x_n \Big|_{z=-1} \\
 &= c && \text{by (2)}
 \end{aligned}$$

3. Proof that (3)  $\implies$  (1):

$$\begin{aligned}
 \sum_{n \in \mathbb{Z}} (-1)^n x_n &= \sum_{n \in \mathbb{Z}} (-1)^{-n} x_n \\
 &= \sum_{n \in \mathbb{Z}} e^{-i\omega n} x_n \Big|_{\omega=\pi} \\
 &= c && \text{by (3)}
 \end{aligned}$$

4. Proof that (2)  $\implies$  (4):

(a) Define  $A \triangleq \sum_{n \in \mathbb{Z}} h_{2n}$        $B \triangleq \sum_{n \in \mathbb{Z}} h_{2n+1}$ .

(b) Proof that  $A - B = c$ :

$$\begin{aligned}
 c &= \sum_{n \in \mathbb{Z}} (-1)^n x_n && \text{by (2)} \\
 &= \underbrace{\sum_{n \in \mathbb{Z}_e} (-1)^n x_n}_{\text{even terms}} + \underbrace{\sum_{n \in \mathbb{Z}_o} (-1)^n x_n}_{\text{odd terms}} \\
 &= \sum_{n \in \mathbb{Z}} (-1)^{2n} x_{2n} + \sum_{n \in \mathbb{Z}} (-1)^{2n+1} x_{2n+1} \\
 &= \underbrace{\sum_{n \in \mathbb{Z}} x_{2n}}_A - \underbrace{\sum_{n \in \mathbb{Z}} x_{2n+1}}_B \\
 &\triangleq A - B && \text{by definitions of } A \text{ and } B
 \end{aligned}$$

(c) Proof that  $A + B = \sum_{n \in \mathbb{Z}} x_n$ :

$$\begin{aligned}
 \sum_{n \in \mathbb{Z}} x_n &= \sum_{n \text{ even}} x_n + \sum_{n \text{ odd}} x_n \\
 &= \underbrace{\sum_{n \in \mathbb{Z}} x_{2n}}_A + \underbrace{\sum_{n \in \mathbb{Z}} x_{2n+1}}_B \\
 &= A + B && \text{by definitions of } A \text{ and } B
 \end{aligned}$$

(d) This gives two simultaneous equations:

$$\begin{aligned}
 A - B &= c \\
 A + B &= \sum_{n \in \mathbb{Z}} x_n
 \end{aligned}$$

(e) Solutions to these equations give

$$\begin{aligned}\sum_{n \in \mathbb{Z}} x_{2n} &\triangleq A &= \frac{1}{2} \left( \sum_{n \in \mathbb{Z}} x_n + c \right) \\ \sum_{n \in \mathbb{Z}} x_{2n+1} &\triangleq B &= \frac{1}{2} \left( \sum_{n \in \mathbb{Z}} x_n - c \right)\end{aligned}$$

5. Proof that (2)  $\iff$  (4):

$$\begin{aligned}\sum_{n \in \mathbb{Z}} (-1)^n x_n &= \underbrace{\sum_{n \in \mathbb{Z}_e} (-1)^n x_n}_{\text{even terms}} + \underbrace{\sum_{n \in \mathbb{Z}_o} (-1)^n x_n}_{\text{odd terms}} \\ &= \sum_{n \in \mathbb{Z}} (-1)^{2n} x_{2n} + \sum_{n \in \mathbb{Z}} (-1)^{2n+1} x_{2n+1} \\ &= \sum_{n \in \mathbb{Z}} x_{2n} - \sum_{n \in \mathbb{Z}} x_{2n+1} \\ &= \frac{1}{2} \left( \sum_{n \in \mathbb{Z}} x_n + c \right) - \frac{1}{2} \left( \sum_{n \in \mathbb{Z}} x_n - c \right) && \text{by (3)} \\ &= c\end{aligned}$$

$\Rightarrow$

**Lemma 6.1.** Let  $\tilde{f}(\omega)$  be the DTFT (Definition 6.1 page 41) of a sequence  $(x_n)_{n \in \mathbb{Z}}$ .

<b>L E M</b>	$\underbrace{((x_n \in \mathbb{R}))_{n \in \mathbb{Z}}}_{\text{REAL-VALUED sequence}} \implies \underbrace{ \tilde{x}(\omega) ^2 =  \tilde{x}(-\omega) ^2}_{\text{EVEN}} \quad \forall (x_n)_{n \in \mathbb{Z}} \in \ell^2_{\mathbb{R}}$
----------------------	--

$\P$  PROOF:

$$\begin{aligned}|\tilde{x}(\omega)|^2 &= |\tilde{x}(z)|^2 \Big|_{z=e^{i\omega}} \\ &= \tilde{x}(z) \tilde{x}^*(z) \Big|_{z=e^{i\omega}} \\ &= \left[ \sum_{n \in \mathbb{Z}} x_n z^{-n} \right] \left[ \sum_{m \in \mathbb{Z}} x_m z^{-n} \right]^* \Big|_{z=e^{i\omega}} \\ &= \left[ \sum_{n \in \mathbb{Z}} x_n z^{-n} \right] \left[ \sum_{m \in \mathbb{Z}} x_m^* (z^*)^{-m} \right] \Big|_{z=e^{i\omega}} \\ &= \sum_{n \in \mathbb{Z}} \sum_{m \in \mathbb{Z}} x_n x_m^* z^{-n} (z^*)^{-m} \Big|_{z=e^{i\omega}} \\ &= \sum_{n \in \mathbb{Z}} \left[ |x_n|^2 + \sum_{m>n} x_n x_m^* z^{-n} (z^*)^{-m} + \sum_{m<n} x_n x_m^* z^{-n} (z^*)^{-m} \right] \Big|_{z=e^{i\omega}} \\ &= \sum_{n \in \mathbb{Z}} \left[ |x_n|^2 + \sum_{m>n} x_n x_m e^{i\omega(m-n)} + \sum_{m<n} x_n x_m e^{i\omega(m-n)} \right] \\ &= \sum_{n \in \mathbb{Z}} \left[ |x_n|^2 + \sum_{m>n} x_n x_m e^{i\omega(m-n)} + \sum_{m>n} x_n x_m e^{-i\omega(m-n)} \right] \\ &= \sum_{n \in \mathbb{Z}} \left[ |x_n|^2 + \sum_{m>n} x_n x_m (e^{i\omega(m-n)} + e^{-i\omega(m-n)}) \right]\end{aligned}$$



$$\begin{aligned}
&= \sum_{n \in \mathbb{Z}} \left[ |x_n|^2 + \sum_{m > n} x_n x_m 2 \cos[\omega(m-n)] \right] \\
&= \sum_{n \in \mathbb{Z}} |x_n|^2 + 2 \sum_{n \in \mathbb{Z}} \sum_{m > n} x_n x_m \cos[\omega(m-n)]
\end{aligned}$$

Since  $\cos$  is real and even, then  $|\check{x}(\omega)|^2$  must also be real and even.  $\Rightarrow$

**Theorem 6.3** (inverse DTFT). <sup>2</sup> Let  $\check{x}(\omega)$  be the DISCRETE-TIME FOURIER TRANSFORM (Definition 6.1 page 41) of a sequence  $(x_n)_{n \in \mathbb{Z}} \in \ell^2_{\mathbb{R}}$ . Let  $\check{x}^{-1}$  be the inverse of  $\check{x}$ .

<b>T H M</b>	$ \left\{ \check{x}(\omega) \triangleq \sum_{n \in \mathbb{Z}} x_n e^{-i\omega n} \right\} \Rightarrow \left\{ x_n = \frac{1}{2\pi} \int_{\alpha-\pi}^{\alpha+\pi} \check{x}(\omega) e^{i\omega n} d\omega \quad \forall \alpha \in \mathbb{R} \right\} \quad \forall (x_n)_{n \in \mathbb{Z}} \in \ell^2_{\mathbb{R}} $ <div style="display: flex; justify-content: space-around; margin-top: 10px;"> <div style="text-align: center;"> <math>\check{x}(\omega) \triangleq \check{\mathbf{F}}((x_n))</math> </div> <div style="text-align: center;"> <math>(x_n) = \check{\mathbf{F}}^{-1} \check{\mathbf{F}}((x_n))</math> </div> </div>
----------------------	--

$\Rightarrow$  PROOF:

$$\begin{aligned}
\frac{1}{2\pi} \int_{\alpha-\pi}^{\alpha+\pi} \check{x}(\omega) e^{i\omega n} d\omega &= \frac{1}{2\pi} \int_{\alpha-\pi}^{\alpha+\pi} \underbrace{\left[ \sum_{m \in \mathbb{Z}} x_m e^{-i\omega m} \right]}_{\check{x}(\omega)} e^{i\omega n} d\omega && \text{by definition of } \check{x}(\omega) \\
&= \frac{1}{2\pi} \int_{\alpha-\pi}^{\alpha+\pi} \sum_{m \in \mathbb{Z}} x_m e^{-i\omega(m-n)} d\omega \\
&= \frac{1}{2\pi} \sum_{m \in \mathbb{Z}} x_m \int_{\alpha-\pi}^{\alpha+\pi} e^{-i\omega(m-n)} d\omega \\
&= \frac{1}{2\pi} \sum_{m \in \mathbb{Z}} x_m [2\pi \delta_{m-n}] \\
&= x_n
\end{aligned}$$

$\Rightarrow$

**Theorem 6.4** (orthonormal quadrature conditions). <sup>3</sup> Let  $\check{x}(\omega)$  be the DISCRETE-TIME FOURIER TRANSFORM (Definition 6.1 page 41) of a sequence  $(x_n)_{n \in \mathbb{Z}} \in \ell^2_{\mathbb{R}}$ . Let  $\bar{\delta}_n$  be the KRONECKER DELTA FUNCTION at  $n$  (Definition G.12 page 160).

<b>T H M</b>	$ \begin{aligned} \sum_{m \in \mathbb{Z}} x_m y_{m-2n}^* &= 0 && \iff && \check{x}(\omega) \check{y}^*(\omega) + \check{x}(\omega + \pi) \check{y}^*(\omega + \pi) &= 0 && \forall n \in \mathbb{Z}, \forall (x_n), (y_n) \in \ell^2_{\mathbb{R}} \\ \sum_{m \in \mathbb{Z}} x_m x_{m-2n}^* &= \bar{\delta}_n && \iff &&  \check{x}(\omega) ^2 +  \check{x}(\omega + \pi) ^2 &= 2 && \forall n \in \mathbb{Z}, \forall (x_n), (y_n) \in \ell^2_{\mathbb{R}} \end{aligned} $
----------------------	---

$\Rightarrow$  PROOF: Let  $z \triangleq e^{i\omega}$ .

<sup>2</sup> J.S.Chitode (2009) page 3-95 <(3.6.2)>

<sup>3</sup> Daubechies (1992) pages 132-137 <(5.1.20),(5.1.39)>

1. Proof that  $2 \sum_{n \in \mathbb{Z}} \left[ \sum_{k \in \mathbb{Z}} x_k y_{k-2n}^* \right] e^{-i2\omega n} = \check{x}(\omega) \check{y}^*(\omega) + \check{x}(\omega + \pi) \check{y}^*(\omega + \pi)$ :

$$\begin{aligned}
 & 2 \sum_{n \in \mathbb{Z}} \left[ \sum_{k \in \mathbb{Z}} x_k y_{k-2n}^* \right] e^{-i2\omega n} \\
 &= 2 \sum_{k \in \mathbb{Z}} x_k \sum_{n \in \mathbb{Z}} y_{k-2n}^* z^{-2n} \\
 &= 2 \sum_{k \in \mathbb{Z}} x_k \sum_{n \text{ even}} y_{k-n}^* z^{-n} \\
 &= \sum_{k \in \mathbb{Z}} x_k \sum_{n \in \mathbb{Z}} y_{k-n}^* z^{-n} (1 + e^{i\pi n}) \\
 &= \sum_{k \in \mathbb{Z}} x_k \sum_{n \in \mathbb{Z}} y_{k-n}^* z^{-n} + \sum_{k \in \mathbb{Z}} x_k \sum_{n \in \mathbb{Z}} y_{k-n}^* z^{-n} e^{i\pi n} \\
 &= \sum_{k \in \mathbb{Z}} x_k \sum_{m \in \mathbb{Z}} y_m^* z^{-(k-m)} + \sum_{k \in \mathbb{Z}} x_k \sum_{m \in \mathbb{Z}} y_m^* e^{-i(\omega+\pi)(k-m)} \quad \text{where } m \triangleq k - n \\
 &= \sum_{k \in \mathbb{Z}} x_k z^{-k} \sum_{m \in \mathbb{Z}} y_m^* z^m + \sum_{k \in \mathbb{Z}} x_k e^{-i(\omega+\pi)k} \sum_{m \in \mathbb{Z}} y_m^* e^{+i(\omega+\pi)m} \\
 &= \sum_{k \in \mathbb{Z}} x_k e^{-i\omega k} \left[ \sum_{m \in \mathbb{Z}} y_m^* e^{-i\omega m} \right]^* + \sum_{k \in \mathbb{Z}} x_k e^{-i(\omega+\pi)k} \left[ \sum_{m \in \mathbb{Z}} y_m^* e^{-i(\omega+\pi)m} \right]^* \\
 &\triangleq \check{x}(\omega) \check{y}^*(\omega) + \check{x}(\omega + \pi) \check{y}^*(\omega + \pi)
 \end{aligned}$$

2. Proof that  $\sum_{m \in \mathbb{Z}} x_m y_{m-2n}^* = 0 \implies \check{x}(\omega) \check{y}^*(\omega) + \check{x}(\omega + \pi) \check{y}^*(\omega + \pi) = 0$ :

$$\begin{aligned}
 0 &= 2 \sum_{n \in \mathbb{Z}} \left[ \sum_{k \in \mathbb{Z}} x_k y_{k-2n}^* \right] e^{-i2\omega n} && \text{by left hypothesis} \\
 &= \check{x}(\omega) \check{y}^*(\omega) + \check{x}(\omega + \pi) \check{y}^*(\omega + \pi) && \text{by item (1)}
 \end{aligned}$$

3. Proof that  $\sum_{m \in \mathbb{Z}} x_m y_{m-2n}^* = 0 \iff \check{x}(\omega) \check{y}^*(\omega) + \check{x}(\omega + \pi) \check{y}^*(\omega + \pi) = 0$ :

$$\begin{aligned}
 2 \sum_{n \in \mathbb{Z}} \left[ \sum_{k \in \mathbb{Z}} x_k y_{k-2n}^* \right] e^{-i2\omega n} &= \check{x}(\omega) \check{y}^*(\omega) + \check{x}(\omega + \pi) \check{y}^*(\omega + \pi) && \text{by item (1)} \\
 &= 0 && \text{by right hypothesis}
 \end{aligned}$$

Thus by the above equation,  $\sum_{n \in \mathbb{Z}} \left[ \sum_{k \in \mathbb{Z}} x_k y_{k-2n}^* \right] e^{-i2\omega n} = 0$ . The only way for this to be true is if  $\sum_{k \in \mathbb{Z}} x_k y_{k-2n}^* = 0$ .

4. Proof that  $\sum_{m \in \mathbb{Z}} x_m x_{m-2n}^* = \bar{\delta}_n \implies |\check{x}(\omega)|^2 + |\check{x}(\omega' + \pi)|^2 = 2$ :  
Let  $g_n \triangleq x_n$ .

$$\begin{aligned}
 2 &= 2 \sum_{n \in \mathbb{Z}} \bar{\delta}_n e^{-i2\omega n} \\
 &= 2 \sum_{n \in \mathbb{Z}} \left[ \sum_{k \in \mathbb{Z}} x_k y_{k-2n}^* \right] e^{-i2\omega n} && \text{by left hypothesis} \\
 &= \check{x}(\omega) \check{y}^*(\omega) + \check{x}(\omega + \pi) \check{y}^*(\omega + \pi) && \text{by item (1)}
 \end{aligned}$$

5. Proof that  $\sum_{m \in \mathbb{Z}} x_m x_{m-2n}^* = \bar{\delta}_n \iff |\check{x}(\omega)|^2 + |\check{x}(\omega' + \pi)|^2 = 2$ :  
Let  $g_n \triangleq x_n$ .

$$\begin{aligned}
 2 \sum_{n \in \mathbb{Z}} \left[ \sum_{k \in \mathbb{Z}} x_k y_{k-2n}^* \right] e^{-i2\omega n} &= \check{x}(\omega) \check{y}^*(\omega) + \check{x}(\omega + \pi) \check{y}^*(\omega + \pi) && \text{by item (1)} \\
 &= 2 && \text{by right hypothesis}
 \end{aligned}$$

Thus by the above equation,  $\sum_{n \in \mathbb{Z}} \left[ \sum_{k \in \mathbb{Z}} x_k y_{k-2n}^* \right] e^{-i2\omega n} = 1$ . The only way for this to be true is if  $\sum_{k \in \mathbb{Z}} x_k y_{k-2n}^* = \delta_n$ .

⇒

## 6.3 Derivatives

**Theorem 6.5.**<sup>4</sup> Let  $\check{x}(\omega)$  be the DTFT (Definition 6.1 page 41) of a sequence  $(x_n)_{n \in \mathbb{Z}}$ .

<b>T H M</b>	(A)	$\left[ \frac{d}{d\omega} \right]^n \check{x}(\omega) \Big _{\omega=0} = 0$	$\iff$	$\sum_{k \in \mathbb{Z}} k^n x_k = 0$	(B)	$\forall n \in \mathbb{W}$
	(C)	$\left[ \frac{d}{d\omega} \right]^n \check{x}(\omega) \Big _{\omega=\pi} = 0$	$\iff$	$\sum_{k \in \mathbb{Z}} (-1)^k k^n x_k = 0$	(D)	$\forall n \in \mathbb{W}$



✎ PROOF:

1. Proof that (A)  $\implies$  (B):

$$\begin{aligned}
 0 &= \left[ \frac{d}{d\omega} \right]^n \check{x}(\omega) \Big|_{\omega=0} && \text{by hypothesis (A)} \\
 &= \left[ \frac{d}{d\omega} \right]^n \sum_{k \in \mathbb{Z}} x_k e^{-i\omega k} \Big|_{\omega=0} && \text{by definition of } \check{x}(\omega) \text{ (Definition 6.1 page 41)} \\
 &= \sum_{k \in \mathbb{Z}} x_k \left[ \frac{d}{d\omega} \right]^n e^{-i\omega k} \Big|_{\omega=0} \\
 &= \sum_{k \in \mathbb{Z}} x_k [(-i)^n k^n e^{-i\omega k}] \Big|_{\omega=0} \\
 &= (-i)^n \sum_{k \in \mathbb{Z}} k^n x_k
 \end{aligned}$$

2. Proof that (A)  $\longleftarrow$  (B):

$$\begin{aligned}
 \left[ \frac{d}{d\omega} \right]^n \check{x}(\omega) \Big|_{\omega=0} &= \left[ \frac{d}{d\omega} \right]^n \sum_{k \in \mathbb{Z}} x_k e^{-i\omega k} \Big|_{\omega=0} && \text{by definition of } \check{x} \\
 &= \sum_{k \in \mathbb{Z}} x_k \left[ \left[ \frac{d}{d\omega} \right]^n e^{-i\omega k} \right] \Big|_{\omega=0} \\
 &= \sum_{k \in \mathbb{Z}} x_k [(-i)^n k^n e^{-i\omega k}] \Big|_{\omega=0} \\
 &= (-i)^n \sum_{k \in \mathbb{Z}} k^n x_k \\
 &= 0 && \text{by hypothesis (B)}
 \end{aligned}$$

<sup>4</sup>  Vidakovic (1999) pages 82–83,  Mallat (1999) pages 241–242

3. Proof that (C)  $\implies$  (D):

$$\begin{aligned}
 0 &= \left[ \frac{d}{d\omega} \right]^n \check{x}(\omega) \Big|_{\omega=\pi} && \text{by hypothesis (C)} \\
 &= \left[ \frac{d}{d\omega} \right]^n \sum_{k \in \mathbb{Z}} x_k e^{-i\omega k} \Big|_{\omega=\pi} && \text{by definition of } \check{x} \text{ (Definition 6.1 page 41)} \\
 &= \sum_{k \in \mathbb{Z}} x_k \left[ \frac{d}{d\omega} \right]^n e^{-i\omega k} \Big|_{\omega=\pi} \\
 &= \sum_{k \in \mathbb{Z}} x_k [(-i)^n k^n e^{-i\omega k}] \Big|_{\omega=\pi} \\
 &= \sum_{k \in \mathbb{Z}} x_k [(-i)^n k^n (-1)^k] \\
 &= (-i)^n \sum_{k \in \mathbb{Z}} (-1)^k k^n x_k
 \end{aligned}$$

4. Proof that (C)  $\Longleftarrow$  (D):

$$\begin{aligned}
 \left[ \frac{d}{d\omega} \right]^n \check{x}(\omega) \Big|_{\omega=\pi} &= \left[ \frac{d}{d\omega} \right]^n \sum_{k \in \mathbb{Z}} x_k e^{-i\omega k} \Big|_{\omega=\pi} && \text{by definition of } \check{x} \text{ (Definition 6.1 page 41)} \\
 &= \sum_{k \in \mathbb{Z}} x_k \left[ \frac{d}{d\omega} \right]^n e^{-i\omega k} \Big|_{\omega=\pi} \\
 &= \sum_{k \in \mathbb{Z}} x_k [(-i)^n k^n e^{-i\omega k}] \Big|_{\omega=\pi} \\
 &= \sum_{k \in \mathbb{Z}} x_k [(-i)^n k^n (-1)^k] \\
 &= (-i)^n \sum_{k \in \mathbb{Z}} (-1)^k k^n x_k \\
 &= 0 && \text{by hypothesis (D)}
 \end{aligned}$$



# CHAPTER 7

## FOURIER SERIES

“...et la nouveauté de l'objet, jointe à son importance, a déterminé la classe à couronner cet ouvrage, en observant cependant que la manière dont l'auteur parvient à ses équations n'est pas exempte de difficultés, et que son analyse, pour les intégrer, laisse encore quelque chose à désirer, soit relativement à la généralité, soit même du côté de la rigueur.”

A competition awards committee consisting of the mathematical giants [Lagrange](#), [Laplace](#), [Legendre](#), and others, commenting on [Fourier's 1807 landmark paper](#) *Dissertation on the propagation of heat in solid bodies* that introduced the *Fourier Series*.<sup>1</sup>



“...and the innovation of the subject, together with its importance, convinced the committee to crown this work. By observing however that the way in which the author arrives at his equations is not free from difficulties, and the analysis of which, to integrate them, still leaves something to be desired, either relative to generality, or even on the side of rigour.”

## 7.1 Definition

The *Fourier Series* expansion of a periodic function is simply a complex trigonometric polynomial. In the special case that the periodic function is even, then the Fourier Series expansion is a cosine polynomial.

### Definition 7.1.<sup>2</sup>

The **Fourier Series operator**  $\hat{\mathbf{F}} : L^2_{\mathbb{R}} \rightarrow \mathcal{E}^2_{\mathbb{R}}$  is defined as

$$[\hat{\mathbf{F}}f](n) \triangleq \frac{1}{\sqrt{\tau}} \int_0^{\tau} f(x) e^{-i \frac{2\pi}{\tau} nx} dx \quad \forall f \in \{f \in L^2_{\mathbb{R}} \mid f \text{ is periodic with period } \tau\}$$

<sup>1</sup> quote: [Lagrange et al. \(1812b\)](#) page 374, [Lagrange et al. \(1812a\)](#) page 112, [Kahane \(2008\)](#) page 199  
translation: assisted by [Google Translate](#), [Castanedo \(2005\)](#) (chapter 2 footnote 5)

paper: [Fourier \(1807\)](#)

<sup>2</sup> [Katznelson \(2004\)](#) page 3

## 7.2 Inverse Fourier Series operator

**Theorem 7.1.** Let  $\hat{\mathbf{F}}$  be the Fourier Series operator.

T H M

The **inverse Fourier Series operator**  $\hat{\mathbf{F}}^{-1}$  is given by

$$[\hat{\mathbf{F}}^{-1}((\tilde{x}_n)_{n \in \mathbb{Z}})](x) \triangleq \frac{1}{\sqrt{\tau}} \sum_{n \in \mathbb{Z}} \tilde{x}_n e^{i \frac{2\pi}{\tau} nx} \quad \forall (\tilde{x}_n) \in \ell^2_{\mathbb{R}}$$

✎ **PROOF:** The proof of the pointwise convergence of the Fourier Series is notoriously difficult. It was conjectured in 1913 by Nikolai Luzin that the Fourier Series for all square summable periodic functions are pointwise convergent: [Luzin \(1913\)](#)

Fifty-three years later (1966) at a conference in Moscow, Lennart Axel Edvard Carleson presented one of the most spectacular results ever in mathematics; he demonstrated that the Luzin conjecture is indeed correct. Carleson formally published his result that same year: [Carleson \(1966\)](#)

Carleson's proof is expounded upon in Reyna's (2002) 175 page book: [de Reyna \(2002\)](#)

Interestingly enough, Carleson started out trying to disprove Luzin's conjecture. Carleson said this in an interview published in 2001:<sup>3</sup> “Well, the problem of course presents itself already when you are a student and I was thinking of the problem on and off, but the situation was more interesting than that. The great authority in those days was Zygmund and he was completely convinced that what one should produce was not a proof but a counter-example. When I was a young student in the United States, I met Zygmund and I had an idea how to produce some very complicated functions for a counter-example and Zygmund encouraged me very much to do so. I was thinking about it for about 15 years on and off, on how to make these counter-examples work and the interesting thing that happened was that I suddenly realized why there should be a counter-example and how you should produce it. I thought I really understood what was the background and then to my amazement I could prove that this “correct” counter-example couldn't exist and therefore I suddenly realized that what you should try to do was the opposite, you should try to prove what was not fashionable, namely to prove convergence. The most important aspect in solving a mathematical problem is the conviction of what is the true result! Then it took like 2 or 3 years using the technique that had been developed during the past 20 years or so. It is actually a problem related to analytic functions basically even though it doesn't look that way.”

For now, if you just want some intuitive justification for the Fourier Series, and you can somehow imagine that the Dirichlet kernel generates a *comb function* of *Dirac delta* functions, then perhaps what follows may help (or not). It is certainly not mathematically rigorous and is by no means a real proof (but at least it is less than 175 pages).

$$\begin{aligned} [\hat{\mathbf{F}}^{-1} \hat{\mathbf{F}} \mathbf{x}](x) &= \hat{\mathbf{F}}^{-1} \left[ \underbrace{\frac{1}{\sqrt{\tau}} \int_0^\tau \mathbf{x}(x) e^{-i \frac{2\pi}{\tau} nx} dx}_{\hat{\mathbf{F}} \mathbf{x}} \right] && \text{by definition of } \hat{\mathbf{F}} && \text{(Definition 7.1 page 51)} \\ &= \frac{1}{\sqrt{\tau}} \sum_{n \in \mathbb{Z}} \left[ \frac{1}{\sqrt{\tau}} \int_0^\tau \mathbf{x}(u) e^{-i \frac{2\pi}{\tau} nu} du \right] e^{i \frac{2\pi}{\tau} nx} && \text{by definition of } \hat{\mathbf{F}}^{-1} \\ &= \frac{1}{\sqrt{\tau}} \sum_{n \in \mathbb{Z}} \frac{1}{\sqrt{\tau}} \int_0^\tau \mathbf{x}(u) e^{-i \frac{2\pi}{\tau} nu} e^{i \frac{2\pi}{\tau} nx} du \\ &= \frac{1}{\sqrt{\tau}} \sum_{n \in \mathbb{Z}} \frac{1}{\sqrt{\tau}} \int_0^\tau \mathbf{x}(u) e^{i \frac{2\pi}{\tau} n(x-u)} du \end{aligned}$$

<sup>3</sup> [Carleson and Engquist \(2001\)](#)

$$\begin{aligned}
&= \int_0^\tau x(u) \underbrace{\frac{1}{\tau} \sum_{n \in \mathbb{Z}} e^{i \frac{2\pi}{\tau} n(x-u)}}_{\lim_{N \rightarrow \infty} D_N(x)} du \\
&= \int_0^\tau x(u) \left[ \sum_{n \in \mathbb{Z}} \delta(x - u - n\tau) \right] du \\
&= \sum_{n \in \mathbb{Z}} \int_{u=0}^{u=\tau} x(u) \delta(x - u - n\tau) du \\
&= \sum_{n \in \mathbb{Z}} \int_{v=n\tau}^{v=n\tau+\tau} x(v - n\tau) \delta(x - v) dv && \text{where } v \triangleq u + n\tau \\
&= \sum_{n \in \mathbb{Z}} \int_{v=n\tau}^{v=(n+1)\tau} x(v - n\tau) \delta(x - v) dv && \text{where } v \triangleq u + n\tau \\
&= \sum_{n \in \mathbb{Z}} \int_{v=n\tau}^{v=(n+1)\tau} x(v) \delta(x - v) dv && \text{because } x \text{ is periodic with period } \tau \\
&= \int_{\mathbb{R}} x(v) \delta(x - v) dv \\
&= x(x) \\
&= \mathbf{I}\tilde{x}(n) && \text{by definition of } \mathbf{I} \quad (\text{Definition F.3 page 120})
\end{aligned}$$

$$\begin{aligned}
[\hat{\mathbf{F}}\hat{\mathbf{F}}^{-1}\tilde{x}](n) &= \hat{\mathbf{F}} \left[ \frac{1}{\sqrt{\tau}} \sum_{k \in \mathbb{Z}} \tilde{x}(k) e^{i \frac{2\pi}{\tau} kx} \right] && \text{by definition of } \hat{\mathbf{F}}^{-1} \\
&= \frac{1}{\sqrt{\tau}} \int_0^\tau \left[ \frac{1}{\sqrt{\tau}} \sum_{k \in \mathbb{Z}} \tilde{x}(k) e^{i \frac{2\pi}{\tau} kx} \right] e^{-i \frac{2\pi}{\tau} nx} dx && \text{by definition of } \hat{\mathbf{F}} \quad (\text{Definition 7.1 page 51}) \\
&= \frac{1}{\tau} \int_0^\tau \left[ \sum_{k \in \mathbb{Z}} \tilde{x}(k) e^{i \frac{2\pi}{\tau} (k-n)x} \right] dx \\
&= \sum_{k \in \mathbb{Z}} \tilde{x}(k) \left[ \frac{1}{\tau} \int_0^\tau e^{i \frac{2\pi}{\tau} (k-n)x} dx \right] \\
&= \sum_{k \in \mathbb{Z}} \tilde{x}(k) \frac{1}{\tau} \left[ \frac{1}{i \frac{2\pi}{\tau} (k-n)} e^{i \frac{2\pi}{\tau} (k-n)x} \right]_0^\tau \\
&= \sum_{k \in \mathbb{Z}} \tilde{x}(k) \frac{1}{i 2\pi (k-n)} [e^{i 2\pi (k-n)} - 1] \\
&= \sum_{k \in \mathbb{Z}} \tilde{x}(k) \delta(k-n) \lim_{x \rightarrow 0} \left[ \frac{e^{i 2\pi x} - 1}{i 2\pi x} \right] \\
&= \tilde{x}(n) \frac{\frac{d}{dx} (e^{i 2\pi x} - 1)}{\frac{d}{dx} (i 2\pi x)} \Big|_{x=0} && \text{by l'Hôpital's rule} \\
&= \tilde{x}(n) \frac{i 2\pi e^{i 2\pi x}}{i 2\pi} \Big|_{x=0} \\
&= \tilde{x}(n) \\
&= \mathbf{I}\tilde{x}(n) && \text{by definition of } \mathbf{I} \quad (\text{Definition F.3 page 120})
\end{aligned}$$



### Theorem 7.2.

The *Fourier Series adjoint operator*  $\hat{\mathbf{F}}^*$  is given by  
 $\hat{\mathbf{F}}^* = \hat{\mathbf{F}}^{-1}$


 PROOF:

$$\begin{aligned}
 \langle \hat{\mathbf{F}}x(x) | \tilde{y}(n) \rangle_{\mathbb{Z}} &= \left\langle \frac{1}{\sqrt{\tau}} \int_0^\tau x(x) e^{-i\frac{2\pi}{\tau}nx} dx | \tilde{y}(n) \right\rangle_{\mathbb{Z}} && \text{by definition of } \hat{\mathbf{F}} && (\text{Definition 7.1 page 51}) \\
 &= \frac{1}{\sqrt{\tau}} \int_0^\tau x(x) \left\langle e^{-i\frac{2\pi}{\tau}nx} | \tilde{y}(n) \right\rangle_{\mathbb{Z}} dx && \text{by additivity property of } \langle \triangle | \nabla \rangle \\
 &= \int_0^\tau x(x) \frac{1}{\sqrt{\tau}} \left\langle \tilde{y}(n) | e^{-i\frac{2\pi}{\tau}nx} \right\rangle_{\mathbb{Z}}^* dx && \text{by property of } \langle \triangle | \nabla \rangle \\
 &= \int_0^\tau x(x) [\hat{\mathbf{F}}^{-1}\tilde{y}(n)]^* dx && \text{by definition of } \hat{\mathbf{F}}^{-1} && (\text{Theorem 7.1 page 52}) \\
 &= \left\langle x(x) | \underbrace{\hat{\mathbf{F}}^{-1}\tilde{y}(n)}_{\hat{\mathbf{F}}^*} \right\rangle_{\mathbb{R}}
 \end{aligned}$$




The Fourier Series operator has several nice properties:

  $\hat{\mathbf{F}}$  is *unitary*<sup>4</sup> (Corollary 7.1 page 54).

 Because  $\hat{\mathbf{F}}$  is unitary, it automatically has several other nice properties such as being *isometric*, and satisfying *Parseval's equation*, satisfying *Plancherel's formula*, and more (Corollary 7.2 page 54).

**Corollary 7.1.** Let  $\mathbf{I}$  be the identity operator and let  $\hat{\mathbf{F}}$  be the Fourier Series operator with adjoint  $\hat{\mathbf{F}}^*$ .


**COR**  $\{ \hat{\mathbf{F}}\hat{\mathbf{F}}^* = \hat{\mathbf{F}}^*\hat{\mathbf{F}} = \mathbf{I} \} \quad ( \hat{\mathbf{F}} \text{ is } \textbf{unitary} \dots \text{and thus also NORMAL and ISOMETRIC} )$

 PROOF: This follows directly from the fact that  $\hat{\mathbf{F}}^* = \hat{\mathbf{F}}^{-1}$  (Theorem 7.2 page 53).



**Corollary 7.2.** Let  $\hat{\mathbf{F}}$  be the Fourier series operator with adjoint  $\hat{\mathbf{F}}^*$  and inverse  $\hat{\mathbf{F}}^{-1}$ .

**COR** 
$$\begin{aligned}
 \mathcal{R}(\hat{\mathbf{F}}) &= \mathcal{R}(\hat{\mathbf{F}}^{-1}) && = \mathcal{L}_{\mathbb{R}}^2 \\
 \|\hat{\mathbf{F}}\| &= \|\hat{\mathbf{F}}^{-1}\| && = 1 && (\text{UNITARY}) \\
 \langle \hat{\mathbf{F}}x | \hat{\mathbf{F}}y \rangle &= \langle \hat{\mathbf{F}}^{-1}x | \hat{\mathbf{F}}^{-1}y \rangle && = \langle x | y \rangle && (\text{PARSEVAL'S EQUATION}) \\
 \|\hat{\mathbf{F}}x\| &= \|\hat{\mathbf{F}}^{-1}x\| && = \|x\| && (\text{PLANCHEREL'S FORMULA}) \\
 \|\hat{\mathbf{F}}x - \hat{\mathbf{F}}y\| &= \|\hat{\mathbf{F}}^{-1}x - \hat{\mathbf{F}}^{-1}y\| && = \|x - y\| && (\text{ISOMETRIC})
 \end{aligned}$$

 PROOF: These results follow directly from the fact that  $\hat{\mathbf{F}}$  is unitary (Corollary 7.1 page 54) and from the properties of unitary operators (Theorem F.26 page 144).



## 7.3 Fourier series for compactly supported functions

**Theorem 7.3.**

**THM** The set  $\left\{ \frac{1}{\sqrt{\tau}} e^{i\frac{2\pi}{\tau}nx} \middle| n \in \mathbb{Z} \right\}$  is an ORTHONORMAL BASIS for all functions  $f(x)$  with support in  $[0 : \tau]$ .

<sup>4</sup>unitary operators: Definition F.14 page 143



# CHAPTER 8

## FAST WAVELET TRANSFORM (FWT)

The Fast Wavelet Transform can be computed using simple discrete filter operations (as a conjugate mirror filter).

**Definition 8.1** (Wavelet Transform). *Let the wavelet transform  $\mathbf{W} : \{f : \mathbb{R} \rightarrow \mathbb{C}\} \rightarrow \{w : \mathbb{Z}^2 \rightarrow \mathbb{C}\}$  be defined as <sup>1</sup>*



**DEF**  $[\mathbf{W}f](j, n) \triangleq \langle f(x) | \psi_{k,n}(x) \rangle$

**Definition 8.2.** *The following relations are defined as described below:*

**DEF**






scaling coefficients	$v_j : \mathbb{Z} \rightarrow \mathbb{C}$	such that	$v_j(n) \triangleq \langle f(x)   \phi_{j,n}(x) \rangle$
wavelet coefficients	$w_j : \mathbb{Z} \rightarrow \mathbb{C}$	such that	$w_j(n) \triangleq \langle f(x)   \psi_{j,n}(x) \rangle$
scaling filter coefficients	$\bar{h} : \mathbb{Z} \rightarrow \mathbb{C}$	such that	$\bar{h}(n) \triangleq h(-n)$
wavelet filter coefficients	$\bar{g} : \mathbb{Z} \rightarrow \mathbb{C}$	such that	$\bar{g}(n) \triangleq g(-n)$

The scaling and wavelet filter coefficients at scale  $j$  are equal to the filtered and downsampled (Theorem ?? page ??) scaling filter coefficients at scale  $j + 1$ :<sup>2</sup>

-  The convolution (Definition 5.3 page 27) of  $v_{j+1}(n)$  with  $\bar{h}(n)$  and then downsampling by 2 produces  $v_j(n)$ .
-  The convolution of  $v_{j+1}(n)$  with  $\bar{g}(n)$  and then downsampling by 2 produces  $w_j(n)$ .

This is formally stated and proved in the next theorem.

<sup>1</sup>Notice that this definition is similar to the definition of transforms of other analysis systems:

	Laplace Transform	$\mathcal{L}f(s)$	$\triangleq \langle f(x)   e^{sx} \rangle$	$\triangleq \int_x f(x) e^{-sx} dx$
	Continuous Fourier Transform	$\mathcal{F}f(\omega)$	$\triangleq \langle f(x)   e^{i\omega x} \rangle$	$\triangleq \int_x f(x) e^{-i\omega x} dx$
	Fourier Series Transform	$\mathcal{F}_s f(k)$	$\triangleq \langle f(x)   e^{i\frac{2\pi}{T} kx} \rangle$	$\triangleq \int_x f(x) e^{-i\frac{2\pi}{T} kx} dx$
	Z-Transform	$\mathcal{Z}f(z)$	$\triangleq \langle f(x)   z^n \rangle$	$\triangleq \sum_n f(x) z^{-n}$
	Discrete Fourier Transform	$\mathcal{F}_d f(k)$	$\triangleq \langle f(n)   e^{i\frac{2\pi}{N} kn} \rangle$	$\triangleq \sum_n f(x) e^{-i\frac{2\pi}{N} kn}$

<sup>2</sup> Mallat (1999) page 257,  Burrus et al. (1998) page 35

**Theorem 8.1.**

<b>T H M</b>	$v_j(n) = [\bar{h} \star v_{j+1}](2n)$
	$w_j(n) = [\bar{g} \star v_{j+1}](2n)$

 PROOF:

$$\begin{aligned}
 v_j(n) &= \langle f(x) | \phi_{j,n}(x) \rangle \\
 &= \langle f(x) | \sqrt{2^j} \phi(2^j x - n) \rangle \\
 &= \left\langle f(x) | \sqrt{2^j} \sqrt{2} \sum_m h(m) \phi(2(2^j x - n) - m) \right\rangle \\
 &= \left\langle f(x) | \sum_m h(j) \sqrt{2^{j+1}} \phi(2^{j+1} x - 2n - m) \right\rangle \\
 &= \sum_m h(m) \langle f(x) | \sqrt{2^{j+1}} \phi(2^{j+1} x - 2n - m) \rangle \\
 &= \sum_m h(m) \langle f(x) | \phi_{j+1,2n+m}(x) \rangle \\
 &= \sum_m h(m) v_{j+1}(2n + m) \\
 &= \sum_p h(p - 2n) v_{j+1}(p) \\
 &= \sum_p \bar{h}(2n - p) v_{j+1}(p) \\
 &= [\bar{h} \star v_{j+1}](2n)
 \end{aligned}$$


$$\text{let } p = 2n + m \iff m = p - 2n$$

$$\begin{aligned}
 w_j(n) &= \langle f(x) | \psi_{j,n}(x) \rangle \\
 &= \langle f(x) | \sqrt{2^j} \psi(2^j x - n) \rangle \\
 &= \left\langle f(x) | \sqrt{2^j} \sqrt{2} \sum_m g(j) \phi(2(2^j x - n) - m) \right\rangle \\
 &= \left\langle f(x) | \sum_m g(m) \sqrt{2^{j+1}} \phi(2^{j+1} x - 2n - m) \right\rangle \\
 &= \sum_m g(m) \langle f(x) | \sqrt{2^{j+1}} \phi(2^{j+1} x - 2n - m) \rangle \\
 &= \sum_m g(m) \langle f(x) | \phi_{j+1,2n+m}(x) \rangle \\
 &= \sum_m g(m) v_{j+1}(2n + m) \\
 &= \sum_p g(p - 2n) v_{j+1}(p) \\
 &= \sum_p \bar{g}(2n - p) v_{j+1}(p) \\
 &= [\bar{g} \star v_{j+1}](2n)
 \end{aligned}$$

$$\text{let } p = 2n + m \iff m = p - 2n$$



These filtering and downsampling operations are equivalent to the operations performed by a filter bank. Therefore, a filter bank can be used to implement a *Fast Wavelet Transform (FWT)*, as illustrated in Figure 8.1 (page 57).

Figure 8.1:  $k$ -Stage Fast Wavelet Transform



# CHAPTER 9

## KL EXPANSION (TRANSFORM)

### 9.1 Definitions

**Definition 9.1.** Let  $x(t)$  be random processes with continuous AUTO-CORRELATION  $R_{xx}(t, u)$  (Definition J.2 page 201) or discrete AUTO-CORRELATION  $R_{xx}(n, m)$  (Definition K.2 page 203).

The **auto-correlation operator**  $\mathbf{R}$  of  $x(t)$  is defined as

$$\mathbf{R}f \triangleq \underbrace{\int_{u \in \mathbb{R}} R_{xx}(t, u) f(u) du}_{\text{(continuous case)}} \quad \text{or} \quad \mathbf{R}f \triangleq \underbrace{\sum_{n \in \mathbb{Z}} R_{xx}(n, m) f(m)}_{\text{(discrete case)}}$$

**Definition 9.2.** Let  $x(t)$  be a RANDOM PROCESS with AUTO-CORRELATION  $R_{xx}(\tau)$  (Definition J.2 page 201).

A RANDOM PROCESS  $x(t)$  is **white** if  $R_{xx}(\tau) = \delta(\tau)$

If a random process  $x(t)$  is **white** (Definition 9.2 page 59) and the set  $\Psi = \{\psi_1(t), \psi_2(t), \dots, \psi_N(t)\}$  is **any** set of orthonormal basis functions, then the innerproducts  $\langle n(t) | \psi_n(t) \rangle$  and  $\langle n(t) | \psi_m(t) \rangle$  are **uncorrelated** for  $m \neq n$ . However, if  $x(t)$  is **colored** (not white), then the innerproducts are not in general uncorrelated. But if the elements of  $\Psi$  are chosen to be the eigenfunctions of  $\mathbf{R}$  such that  $\mathbf{R}\psi_n = \lambda_n \psi_n$ , then by Theorem J.1 (page 202), the set  $\{\psi_n(t)\}$  are **orthogonal** and the innerproducts are **uncorrelated** even though  $x(t)$  is not white. This criterion is called the Karhunen-Loève criterion for  $x(t)$ .

### 9.2 Properties

**Theorem 9.1.** Let  $\mathbf{R}$  be an AUTO-CORRELATION operator.

$$\left\{ \begin{array}{l} \text{(A). } \langle x | y \rangle \triangleq \int_{t \in \mathbb{R}} x(t) y^*(t) dt \quad \text{OR} \\ \text{(B). } \langle x | y \rangle \triangleq \sum_{n \in \mathbb{Z}} x(n) y^*(n) \end{array} \right\} \Rightarrow \left\{ \begin{array}{l} \text{(1). } \langle \mathbf{R}x | x \rangle \geq 0 \quad \text{(NON-NEGATIVE)} \\ \text{(2). } \langle \mathbf{R}x | y \rangle = \langle x | \mathbf{R}y \rangle \quad \text{(SELF-ADJOINT)} \end{array} \right\} \text{ and }$$

✎ PROOF:

1. Proof that  $\mathbf{R}$  is *non-negative* under hypothesis (A):

$$\begin{aligned}
 \langle \mathbf{R}y | y \rangle &= \left\langle \int_{u \in \mathbb{R}} R_{xx}(t, u) y(u) du | y(t) \right\rangle && \text{by definition of } \mathbf{R} && (\text{Definition 9.1 page 59}) \\
 &= \left\langle \int_{u \in \mathbb{R}} E[x(t)x^*(u)] y(u) du | y(t) \right\rangle && \text{by definition of } R_{xx}(t, u) && (\text{Definition J.2 page 201}) \\
 &= E \left[ \left\langle \int_{u \in \mathbb{R}} x(t)x^*(u) y(u) du | y(t) \right\rangle \right] && \text{by linearity of } \langle \Delta | \nabla \rangle \text{ and } \int && \\
 &= E \left[ \int_{u \in \mathbb{R}} x^*(u) y(u) du \langle x(t) | y(t) \rangle \right] && \text{by additivity property of } \langle \Delta | \nabla \rangle && \\
 &= E [\langle y(u) | x(u) \rangle \langle x(t) | y(t) \rangle] && \text{by local definition of } \langle \Delta | \nabla \rangle && \\
 &= E [\langle x(u) | y(u) \rangle^* \langle x(t) | y(t) \rangle] && \text{by conjugate symmetry prop.} && \\
 &= E |\langle x(t) | y(t) \rangle|^2 && \text{by definition of } |\cdot| && (\text{Definition A.4 page 68}) \\
 &\geq 0 && \text{by strictly positive property of norms} &&
 \end{aligned}$$

2. Proof that  $\mathbf{R}$  is *self-adjoint* under hypothesis (A):

$$\begin{aligned}
 \langle [\mathbf{R}x](t) | y \rangle &= \left\langle \int_{u \in \mathbb{R}} R_{xx}(t, u) x(u) du | y(t) \right\rangle && \text{by definition of } \mathbf{R} && (\text{Definition 9.1 page 59}) \\
 &= \int_{u \in \mathbb{R}} x(u) \langle R_{xx}(t, u) | y(t) \rangle du && \text{by additive property of } \langle \Delta | \nabla \rangle && \\
 &= \int_{u \in \mathbb{R}} x(u) \langle y(t) | R_{xx}(t, u) \rangle^* du && \text{by conjugate symmetry prop.} && \\
 &= \langle x(u) | \langle y(t) | R_{xx}(t, u) \rangle \rangle && \text{by local definition of } \langle \Delta | \nabla \rangle && \\
 &= \left\langle x(u) | \int_{t \in \mathbb{R}} y(t) R_{xx}^*(t, u) dt \right\rangle && && \\
 &= \left\langle x(u) | \int_{t \in \mathbb{R}} y(t) R_{xx}(u, t) dt \right\rangle && \text{by property of } R_{xx} && (\text{Theorem J.1 page 202}) \\
 &= \left\langle x(u) | \mathbf{R}y \right\rangle_{\mathbf{R}^*} && \text{by definition of } \mathbf{R} && (\text{Definition 9.1 page 59}) \\
 \implies \mathbf{R} &= \mathbf{R}^* && \text{by definition of adjoint } \mathbf{R}^* && (\text{Definition F.8 page 129}) \\
 \implies \mathbf{R} &\text{ is self-adjoint} && \text{by definition of self-adjoint} && (\text{Definition F.11 page 137})
 \end{aligned}$$

3. Proofs under hypothesis (B): substitute  $\sum_{n \in \mathbb{Z}}$  operator for  $\int_{t \in \mathbb{R}} dt$  operator in above proofs.

⇒

**Theorem 9.2.**<sup>1</sup> Let  $(\lambda_n)_{n \in \mathbb{Z}}$  be the eigenvalues and  $(\psi_n)_{n \in \mathbb{Z}}$  be the eigenfunctions of operator  $\mathbf{R}$  such that  $\mathbf{R}\psi_n = \lambda_n \psi_n$ .

- |             |  |   |
|-------------|--|---|
| T<br>H<br>M | (1). $\lambda_n \in \mathbb{R}$  | (REAL-VALUED)   |
|             | (2). $\lambda_n \neq \lambda_m \implies \langle \psi_n   \psi_m \rangle = 0$           | (ORTHOGONAL)  |
|             | (3). $\ \psi_n(t)\ ^2 > 0 \implies \lambda_n \geq 0$                                   | (NON-NEGATIVE)  |
|             | (4). $\ \psi_n(t)\ ^2 > 0, \langle \mathbf{R}f   f \rangle > 0 \implies \lambda_n > 0$ | ( $\mathbf{R}$ POSITIVE DEFINITE $\implies \lambda_n$ POSITIVE) |

✎ PROOF:

<sup>1</sup> Keener (1988) pages 114–119

1. Proof that eigenvalues are *real-valued*: Because  $\mathbf{R}$  is *self-adjoint*, its eigenvalues are real (Theorem F.18 page 137).
2. Proof that eigenfunctions associated with distinct eigenvalues are orthogonal: Because  $\mathbf{R}$  is *self-adjoint*, this property follows (Theorem F.18 page 137).
3. Proof that eigenvalues are *non-negative*:

$$\begin{aligned}
 0 &\leq \langle \mathbf{R}\psi_n | \psi_n \rangle && \text{by definition of } \textit{non-negative definite} \\
 &= \langle \lambda_n \psi_n | \psi_n \rangle && \text{by hypothesis} \\
 &= \lambda_n \langle \psi_n | \psi_n \rangle && \text{by definition of inner-products} \\
 &= \lambda_n \|\psi_n\|^2 && \text{by definition of norm induced by inner-product}
 \end{aligned}$$

4. Proof that eigenvalues are *positive* if  $\mathbf{R}$  is *positive definite*:

$$\begin{aligned}
 0 &< \langle \mathbf{R}\psi_n | \psi_n \rangle && \text{by definition of } \textit{positive definite} \\
 &= \langle \lambda_n \psi_n | \psi_n \rangle && \text{by hypothesis} \\
 &= \lambda_n \langle \psi_n | \psi_n \rangle && \text{by } \textit{homogeneous} \text{ property of } \langle \Delta | \nabla \rangle \\
 &= \lambda_n \|\psi_n\|^2 && \text{by } \textit{induced norm} \text{ theorem}
 \end{aligned}$$



**Theorem 9.3** (Karhunen-Loève Expansion). <sup>2</sup> Let  $\mathbf{R}$  be the AUTO-CORRELATION OPERATOR (Definition 9.1 page 59) of a RANDOM PROCESS  $x(t)$ . Let  $(\lambda_n)_{n \in \mathbb{Z}}$  be the eigenvalues of  $\mathbf{R}$  and  $(\psi_n)_{n \in \mathbb{Z}}$  are the eigenfunctions of  $\mathbf{R}$  such that  $\mathbf{R}\psi_n = \lambda_n \psi_n$ .

T H M

$$\underbrace{\|\psi_n(t)\| = 1}_{\{\psi_n(t)\} \text{ are NORMALIZED}} \implies \underbrace{E \left( \left| x(t) - \sum_{n \in \mathbb{Z}} \langle x(t) | \psi_n(t) \rangle \psi_n(t) \right|^2 \right)}_{\text{CONVERGENCE IN PROBABILITY}} = 0 \quad (\{\psi_n(t)\} \text{ is a BASIS for } x(t))$$

PROOF:

1. Define  $\dot{x}_n \triangleq \langle x(t) | \psi_n(t) \rangle$
2. Define  $\mathbf{R}x(t) \triangleq \int_{u \in \mathbb{R}} R_{xx}(t, u) x(u) du$
3. lemma:  $E[x(t)x(t)] = \sum_{n \in \mathbb{Z}} \lambda_n |\psi_n(t)|^2$ . Proof:

$$E[x(t)x(t)] = \sum_{n \in \mathbb{Z}} \lambda_n |\psi_n(t)|^2 \quad \begin{array}{l} \text{by } \textit{non-negative property} \text{ (Theorem 9.1 page 59)} \\ \text{and } \textit{Mercer's Theorem} \text{ (Theorem L.4 page 212)} \end{array}$$

<sup>2</sup> Keener (1988) pages 114–119

4. lemma:

$$\begin{aligned}
 & \mathbb{E} \left[ x(t) \left( \sum_{n \in \mathbb{Z}} \dot{x}_n \psi_n(t) \right)^* \right] \\
 & \triangleq \mathbb{E} \left[ x(t) \left( \sum_{n \in \mathbb{Z}} \int_{u \in \mathbb{R}} x(u) \psi_n^*(u) du \psi_n(t) \right)^* \right] && \text{by definition of } \dot{x} && (\text{definition 1 page 61}) \\
 & = \sum_{n \in \mathbb{Z}} \left( \int_{u \in \mathbb{R}} \mathbb{E} [x(t) x^*(u)] \psi_n(u) du \right) \psi_n^*(t) && \text{by linearity} \\
 & \triangleq \sum_{n \in \mathbb{Z}} \left( \int_{u \in \mathbb{R}} R_{xx}(t, u) \psi_n(u) du \right) \psi_n^*(t) && \text{by definition of } R_{xx}(t, u) && (\text{Definition J.2 page 201}) \\
 & \triangleq \sum_{n \in \mathbb{Z}} (\mathbf{R} \psi_n(t) \psi_n^*(t)) && \text{by definition of } \mathbf{R} && (\text{definition 2 page 61}) \\
 & = \sum_{n \in \mathbb{Z}} \lambda_n \psi_n(t) \psi_n^*(t) && \text{by property of eigen-system} \\
 & = \sum_{n \in \mathbb{Z}} \lambda_n |\psi_n(t)|^2
 \end{aligned}$$

5. lemma:

$$\begin{aligned}
 & \mathbb{E} \left[ \sum_{n \in \mathbb{Z}} \dot{x}_n \psi_n(t) \left( \sum_{m \in \mathbb{Z}} \dot{x}_m \psi_m(t) \right)^* \right] \\
 & \triangleq \mathbb{E} \left[ \sum_{n \in \mathbb{Z}} \int_{u \in \mathbb{R}} x(u) \psi_n^*(u) du \psi_n(t) \left( \sum_{m \in \mathbb{Z}} \int_{v \in \mathbb{R}} x(v) \psi_m^*(v) dv \psi_m(t) \right)^* \right] && \text{by definition of } \dot{x} && (\text{definition 1 page 61}) \\
 & = \sum_{n \in \mathbb{Z}} \sum_{m \in \mathbb{Z}} \int_{u \in \mathbb{R}} \left( \int_{v \in \mathbb{R}} \mathbb{E} [x(u) x^*(v)] \psi_m(v) dv \right) \psi_n^*(u) du \psi_n(t) \psi_m^*(t) && \text{by linearity} \\
 & = \sum_{n \in \mathbb{Z}} \sum_{m \in \mathbb{Z}} \int_{u \in \mathbb{R}} \left( \int_{v \in \mathbb{R}} R_{xx}(u, v) \psi_m(v) dv \right) \psi_n^*(u) du \psi_n(t) \psi_m^*(t) && \text{by definition of } R_{xx}(t, u) && (\text{Definition J.2 page 201}) \\
 & = \sum_{n \in \mathbb{Z}} \sum_{m \in \mathbb{Z}} \int_{u \in \mathbb{R}} (\mathbf{R} \psi_m(u)) \psi_n^*(u) du \psi_n(t) \psi_m^*(t) && \text{by definition of } \mathbf{R} && (\text{definition 2 page 61}) \\
 & = \sum_{n \in \mathbb{Z}} \sum_{m \in \mathbb{Z}} \int_{u \in \mathbb{R}} (\lambda_m \psi_m(u)) \psi_n^*(u) du \psi_n(t) \psi_m^*(t) && \text{by property of eigen-system} \\
 & = \sum_{n \in \mathbb{Z}} \sum_{m \in \mathbb{Z}} \lambda_m \left( \int_{u \in \mathbb{R}} \psi_m(u) \psi_n^*(u) du \right) \psi_n(t) \psi_m^*(t) && \text{by linearity} \\
 & = \sum_{n \in \mathbb{Z}} \sum_{m \in \mathbb{Z}} \lambda_m \|\psi(t)\|^2 \bar{\delta}_{mn} \psi_n(t) \psi_m^*(t) && \text{by orthogonal property} && (\text{Theorem 9.2 page 60}) \\
 & = \sum_{n \in \mathbb{Z}} \sum_{m \in \mathbb{Z}} \lambda_m \bar{\delta}_{mn} \psi_n(t) \psi_m^*(t) && \text{by normalized hypothesis} \\
 & = \sum_{n \in \mathbb{Z}} \lambda_n \psi_n(t) \psi_n^*(t) && \text{by definition of Kronecker delta } \bar{\delta} && (\text{Definition G.12 page 160}) \\
 & = \sum_{n \in \mathbb{Z}} \lambda_n |\psi_n(t)|^2
 \end{aligned}$$

6. Proof that  $\{\psi_n(t)\}$  is a *basis* for  $x(t)$ :

$$\mathbb{E} \left( \left| x(t) - \sum_{n \in \mathbb{Z}} \dot{x}_n \psi_n(t) \right|^2 \right)$$



$$\begin{aligned}
&= \mathbb{E} \left( \left[ x(t) - \sum_{n \in \mathbb{Z}} \dot{x}_n \psi_n(t) \right] \left[ x(t) - \sum_{m \in \mathbb{Z}} \dot{x}_m \psi_m(t) \right]^* \right) \\
&= \mathbb{E} \left( x(t)x^*(t) - x(t) \left[ \sum_{n \in \mathbb{Z}} \dot{x}_n \psi_n(t) \right]^* - x^*(t) \sum_{n \in \mathbb{Z}} \dot{x}_n \psi_n(t) + \left[ \sum_{n \in \mathbb{Z}} \dot{x}_n \psi_n(t) \right] \left[ \sum_{m \in \mathbb{Z}} \dot{x}_m \psi_m(t) \right]^* \right) \\
&= \mathbb{E}(x(t)x^*(t)) - \mathbb{E} \left[ x(t) \left[ \sum_{n \in \mathbb{Z}} \dot{x}_n \psi_n(t) \right]^* \right] - \mathbb{E} \left[ x^*(t) \sum_{n \in \mathbb{Z}} \dot{x}_n \psi_n(t) \right] + \mathbb{E} \left[ \sum_{n \in \mathbb{Z}} \dot{x}_n \psi_n(t) \left[ \sum_{m \in \mathbb{Z}} \dot{x}_m \psi_m(t) \right]^* \right] \\
&\quad \text{by linearity of } \mathbb{E} \\
&= \underbrace{\sum_{n \in \mathbb{Z}} \lambda_n |\psi_n(t)|^2}_{\text{by (3) lemma}} - \underbrace{\sum_{n \in \mathbb{Z}} \lambda_n |\psi_n(t)|^2}_{\text{by (4) lemma}} - \underbrace{\left[ \sum_{n \in \mathbb{Z}} \lambda_n |\psi_n(t)|^2 \right]^*}_{\text{by (4) lemma}} + \underbrace{\sum_{n \in \mathbb{Z}} \lambda_n |\psi_n(t)|^2}_{\text{by (5) lemma}} \\
&= 0
\end{aligned}$$



## 9.3 Quasi-basis

The *auto-correlation operator*  $\mathbf{R}$  (Definition 9.1 page 59) in the discrete case can be approximated using a *correlation matrix*. In the *zero-mean* case, this becomes

$$\mathbf{R} \triangleq \begin{bmatrix} \mathbb{E}[y_1 y_1] & \mathbb{E}[y_1 y_2] & \cdots & \mathbb{E}[y_1 y_n] \\ \mathbb{E}[y_2 y_1] & \mathbb{E}[y_2 y_2] & & \mathbb{E}[y_2 y_n] \\ \vdots & \vdots & \ddots & \vdots \\ \mathbb{E}[y_n y_1] & \mathbb{E}[y_n y_2] & \cdots & \mathbb{E}[y_n y_n] \end{bmatrix}$$

The eigen-vectors (and hence a quasi-basis) for  $\mathbf{R}$  can be found using a *Cholesky Decomposition*.

### Proposition 9.1. <sup>3</sup>

P  
R  
P

The AUTO-CORRELATION MATRIX  $\mathbf{R}$  is *Toeplitz*.

*Remark 9.1.* For more information about the properties of **Toeplitz matrices**, see

1. Grenander and Szegö (1958),
2. Widom (1965),
3. Gray (1971),
4. Smylie et al. (1973) page 408 (§“B. PROPERTIES OF THE TOEPLITZ MATRIX”),
5. GRENANDER AND SZEGÖ (1984),
6. HAYKIN AND KESLER (1979),
7. HAYKIN AND KESLER (1983),
8. S. LAWRENCE MARPLE (1987) PAGES 80–92 (§“3.8 THE TOEPLITZ MATRIX”),
9. BÖTTCHER AND SILBERMANN (1999) (ISBN:9780387985701),
10. GRAY (2006),
11. S. LAWRENCE MARPLE (2019) PAGES 80–93 (§“3.8 THE TOEPLITZ MATRIX”).

<sup>3</sup>See Clarkson (1993) page 131 (§“Appendix 3A — Positive Semi-Definite Form of the Autocorrelation Matrix”)







# APPENDIX A

## ALGEBRAIC STRUCTURES



“In this book, learned reader, you have the rules of algebra... It unties the knot not only where one term is equal to another or two to one but also where two are equal to two or three to one.... this most abstruse and unsurpassed treasury of the entire arithmetic being brought to light and, as in a theater, exposed to the sight of all...”

Gerolamo Cardano (1501–1576), Italian mathematician, physician, and astrologer <sup>1</sup>

A set together with one or more operations forms several standard mathematical structures:

*group*  $\supseteq$  *ring*  $\supseteq$  *commutative ring*  $\supseteq$  *integral domain*  $\supseteq$  *field*

**Definition A.1.** <sup>2</sup> Let  $X$  be a set and  $\diamond : X \times X \rightarrow X$  be an operation on  $X$ .

The pair  $(X, \diamond)$  is a **group** if

- |            |    |                                |   |                         |                    |     |
|------------|----|--------------------------------|---|-------------------------|--------------------|-----|
| <b>DEF</b> | 1. | $\exists e \in X$ such that    | $e \diamond x = x \diamond e = x$                       | $\forall x \in X$       | (IDENTITY element) | and |
|            | 2. | $\exists (-x) \in X$ such that | $(-x) \diamond x = x \diamond (-x) = e$                 | $\forall x \in X$       | (INVERSE element)  | and |
|            | 3. |                                | $x \diamond (y \diamond z) = (x \diamond y) \diamond z$ | $\forall x, y, z \in X$ | (ASSOCIATIVE)      |     |

**Definition A.2.** <sup>3</sup> Let  $+$  :  $X \times X \rightarrow X$  and  $*$  :  $X \times X \rightarrow X$  be operations on a set  $X$ . Furthermore, let the operation  $*$  also be represented by juxtaposition as in  $a * b \equiv ab$ .

The triple  $(X, +, *)$  is a **ring** if

- |            |    |                          |                         |   |     |
|------------|----|--------------------------|-------------------------|---|-----|
| <b>DEF</b> | 1. | $(X, +)$ is a group.     |                         | (additive group)                        | and |
|            | 2. | $x(yz) = (xy)z$          | $\forall x, y, z \in X$ | (ASSOCIATIVE with respect to $*$ )      | and |
|            | 3. | $x(y + z) = (xy) + (xz)$ | $\forall x, y, z \in X$ | ( $*$ is LEFT DISTRIBUTIVE over $+$ )   | and |
|            | 4. | $(x + y)z = (xz) + (yz)$ | $\forall x, y, z \in X$ | ( $*$ is RIGHT DISTRIBUTIVE over $+$ ). |     |

**Definition A.3.** <sup>4</sup>

<sup>1</sup> quote: Cardano (1545) page 1

image: <http://en.wikipedia.org/wiki/Image:Cardano.jpg>

<sup>2</sup> Durbin (2000) page 29

<sup>3</sup> Durbin (2000) pages 114–115

<sup>4</sup> Durbin (2000) page 118

DEF

A triple  $(X, +, *)$  is a **commutative ring** if

1.  $(X, +, *)$  is a RING and
2.  $xy = yx \quad \forall x, y \in X$  (COMMUTATIVE).

**Definition A.4.** <sup>5</sup> Let  $R$  be a COMMUTATIVE RING (Definition A.3 page 67).

DEF

A function  $|\cdot|$  in  $\mathbb{R}^{\mathbb{R}}$  is an **absolute value** (or **modulus**) if

1.  $|x| \geq 0 \quad x \in \mathbb{R}$  (NON-NEGATIVE) and
2.  $|x| = 0 \iff x = 0 \quad x \in \mathbb{R}$  (NONDEGENERATE) and
3.  $|xy| = |x| \cdot |y| \quad x, y \in \mathbb{R}$  (HOMOGENEOUS / SUBMULTIPLICATIVE) and
4.  $|x + y| \leq |x| + |y| \quad x, y \in \mathbb{R}$  (SUBADDITIVE / TRIANGLE INEQUALITY)

**Definition A.5.** <sup>6</sup>

DEF

The structure  $F \triangleq (X, +, \cdot, 0, 1)$  is a **field** if


1.  $(X, +, *)$  is a ring (ring) and
2.  $xy = yx \quad \forall x, y \in X$  (commutative with respect to  $*$ ) and
3.  $(X \setminus \{0\}, *)$  is a group (group with respect to  $*$ ).

**Definition A.6.** <sup>7</sup> Let  $V = (F, +, \cdot)$  be a vector space and  $\otimes : V \times V \rightarrow V$  be a vector-vector multiplication operator.



An **algebra** is any pair  $(V, \otimes)$  that satisfies ( $\otimes$  is represented by juxtaposition)

DEF

1.  $(ux)y = u(xy) \quad \forall u, x, y \in V$  (ASSOCIATIVE) and
2.  $u(x + y) = (ux) + (uy) \quad \forall u, x, y \in V$  (LEFT DISTRIBUTIVE) and
3.  $(u + x)y = (uy) + (xy) \quad \forall u, x, y \in V$  (RIGHT DISTRIBUTIVE) and
4.  $\alpha(xy) = (\alpha x)y = x(\alpha y) \quad \forall x, y \in V \text{ and } \alpha \in F$  (SCALAR COMMUTATIVE) .

<sup>5</sup>  Cohn (2002) page 312

<sup>6</sup>  Durbin (2000) page 123,  Weber (1893)

<sup>7</sup>  Abramovich and Aliprantis (2002) page 3,  Michel and Herget (1993) page 56

## APPENDIX B

## CALCULUS

**Definition B.1.** Let  $\mathbb{R}$  be the set of real numbers,  $\mathcal{B}$  the set of BOREL SETS on  $\mathbb{R}$ , and  $\mu$  the standard BOREL MEASURE on  $\mathcal{B}$ . Let  $\mathbb{R}^{\mathbb{R}}$  be as in Definition 1.1 page 187.

The **space of Lebesgue square-integrable functions**  $L^2_{(\mathbb{R}, \mathcal{B}, \mu)}$  (or  $L^2_{\mathbb{R}}$ ) is defined as

$$L^2_{\mathbb{R}} \triangleq L^2_{(\mathbb{R}, \mathcal{B}, \mu)} \triangleq \left\{ f \in \mathbb{R}^{\mathbb{R}} \mid \left( \int_{\mathbb{R}} |f|^2 \right)^{\frac{1}{2}} d\mu < \infty \right\}.$$

The **standard inner product**  $\langle \triangle \mid \nabla \rangle$  on  $L^2_{\mathbb{R}}$  is defined as

$$\langle f(x) \mid g(x) \rangle \triangleq \int_{\mathbb{R}} f(x) g^*(x) dx.$$

The **standard norm**  $\|\cdot\|$  on  $L^2_{\mathbb{R}}$  is defined as  $\|f(x)\| \triangleq \langle f(x) \mid f(x) \rangle^{\frac{1}{2}}$

**Definition B.2.** Let  $f(x)$  be a FUNCTION in  $\mathbb{R}^{\mathbb{R}}$ .

$$\frac{d}{dx} f(x) \triangleq f'(x) \triangleq \lim_{\varepsilon \rightarrow 0} \frac{f(x + \varepsilon) - f(x)}{\varepsilon}$$

**Proposition B.1.**

$$\left\{ \begin{array}{l} (1). \quad f(x) \text{ is CONTINUOUS} \quad \text{and} \\ (2). \quad \underbrace{f(a+x) = f(a-x)}_{\text{SYMMETRIC about a point } a} \end{array} \right\} \Rightarrow \left\{ \begin{array}{l} (1). \quad f'(a+x) = -f'(a-x) \quad (\text{ANTI-SYMMETRIC about } a) \\ (2). \quad f'(a) = 0 \end{array} \right\}$$

 PROOF:

$$\begin{aligned} f'(a+x) &= \lim_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon} [f(a+x+\varepsilon) - f(a+x-\varepsilon)] \\ &= \lim_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon} [f(a-x-\varepsilon) - f(a-x+\varepsilon)] && \text{by hypothesis (2)} \\ &= -\lim_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon} [f(a-x+\varepsilon) - f(a-x-\varepsilon)] \\ &= -f'(a-x) \end{aligned}$$

$$\begin{aligned} f'(a) &= \frac{1}{2} f'(a+0) + \frac{1}{2} f'(a-0) \\ &= \frac{1}{2} [f'(a+0) - f'(a+0)] && \text{by previous result} \end{aligned}$$

$$= 0$$



### Lemma B.1.

**L  
E  
M**

$$f(x) \text{ is INVERTIBLE} \implies \left\{ \frac{d}{dy} f^{-1}(y) = \frac{1}{\frac{d}{dx} f[f^{-1}(y)]} \right\}$$

PROOF:

$$\begin{aligned} \frac{d}{dy} f^{-1}(y) &\triangleq \lim_{\varepsilon \rightarrow 0} \frac{f^{-1}(y + \varepsilon) - f^{-1}(y)}{\varepsilon} && \text{by definition of } \frac{d}{dy} && (\text{Definition B.2 page 69}) \\ &= \lim_{\delta \rightarrow 0} \frac{1}{\left[ \frac{f(x + \delta) - f(x)}{\delta} \right]} \bigg|_{x \triangleq f^{-1}(y)} && \text{because in the limit, } \frac{\Delta y}{\Delta x} = \left( \frac{\Delta x}{\Delta y} \right)^{-1} \\ &\triangleq \frac{1}{\frac{d}{dx} f(x)} \bigg|_{x \triangleq f^{-1}(y)} && \text{by definition of } \frac{d}{dx} && (\text{Definition B.2 page 69}) \\ &= \frac{1}{\frac{d}{dx} f[f^{-1}(y)]} && \text{because } x \triangleq f^{-1}(y) \end{aligned}$$



**Theorem B.1.** <sup>1</sup> Let  $f$  be a continuous function in  $L^2_{\mathbb{R}}$  and  $f^{(n)}$  the  $n$ th derivative of  $f$ .

**T  
H  
M**

$$\int_{[0:1]^n} f^{(n)} \left( \sum_{k=1}^n x_k \right) dx_1 dx_2 \cdots dx_n = \sum_{k=0}^n (-1)^{n-k} \binom{n}{k} f(k) \quad \forall n \in \mathbb{N}$$

PROOF: Proof by induction:

1. Base case ...proof for  $n = 1$  case:

$$\begin{aligned} \int_{[0:1]} f^{(1)}(x) dx &= f(1) - f(0) && \text{by Fundamental theorem of calculus} \\ &= (-1)^{1+1} \binom{1}{1} f(1) + (-1)^{1+0} \binom{1}{0} f(0) \\ &= \sum_{k=0}^1 (-1)^{1-k} \binom{1}{k} f(k) \end{aligned}$$

<sup>1</sup> Chui (1992) page 86 (item (ii)), Prasad and Iyengar (1997) pages 145–146 (Theorem 6.2 (b))



2. Induction step ...proof that  $n$  case  $\implies n + 1$  case:

$$\begin{aligned}
 & \int_{[0:1]^{n+1}} f^{(n+1)} \left( \sum_{k=1}^{n+1} x_k \right) dx_1 dx_2 \cdots dx_{n+1} \\
 &= \int_{[0:1]^n} \left[ \int_0^1 f^{(n+1)} \left( x_{n+1} + \sum_{k=1}^n x_k \right) dx_{n+1} \right] dx_1 dx_2 \cdots dx_n \\
 &= \int_{[0:1]^n} \left[ f^{(n)} \left( x_{n+1} + \sum_{k=1}^n x_k \right) \right]_{x_{n+1}=0}^{x_{n+1}=1} dx_1 dx_2 \cdots dx_n \quad \text{by Fundamental theorem of calculus} \\
 &= \int_{[0:1]^n} \left[ f^{(n)} \left( 1 + \sum_{k=1}^n x_k \right) - f^{(n)} \left( 0 + \sum_{k=1}^n x_k \right) \right] dx_1 dx_2 \cdots dx_n \\
 &= \sum_{k=0}^n (-1)^{n-k} \binom{n}{k} f(k+1) - \sum_{k=0}^n (-1)^{n-k} \binom{n}{k} f(k) \quad \text{by induction hypothesis} \\
 &= \sum_{m=1}^{m=n+1} (-1)^{n-m+1} \binom{n}{m-1} f(m) + \sum_{k=0}^n (-1)(-1)^{n-k} \binom{n}{k} f(k) \quad \text{where } m \triangleq k+1 \implies k = m-1 \\
 &= \left[ f(n+1) + \sum_{k=1}^n (-1)^{n-k+1} \binom{n}{k-1} f(k) \right] + \left[ (-1)^{n+1} f(0) + \sum_{k=1}^n (-1)^{n-k+1} \binom{n}{k} f(k) \right] \\
 &= f(n+1) + (-1)^{n+1} f(0) + \sum_{k=1}^n (-1)^{n-k+1} \underbrace{\left[ \binom{n}{k-1} + \binom{n}{k} \right]}_{\text{use Stifel formula}} f(k) \\
 &= (-1)^0 \binom{n+1}{n+1} f(n+1) + (-1)^{n+1} \binom{n+1}{0} f(0) + \sum_{k=1}^n (-1)^{n-k+1} \binom{n+1}{k} f(k) \quad \text{by Stifel formula} \\
 &= \sum_{k=0}^{n+1} (-1)^{n-k+1} \binom{n+1}{k} f(k)
 \end{aligned}$$

$\Rightarrow$

Some proofs invoke differentiation multiple times. This is simplified thanks to the *Leibniz rule*, also called the *generalized product rule* (GPR, next lemma). The Leibniz rule is remarkably similar in form to the *binomial theorem*.

**Lemma B.2** (Leibniz rule / generalized product rule). <sup>2</sup> Let  $f(x), g(x) \in \mathcal{L}_{\mathbb{R}}^2$  with derivatives  $f^{(n)}(x) \triangleq \frac{d^n}{dx^n} f(x)$  and  $g^{(n)}(x) \triangleq \frac{d^n}{dx^n} g(x)$  for  $n = 0, 1, 2, \dots$ , and  $\binom{n}{k} \triangleq \frac{n!}{(n-k)!k!}$  (binomial coefficient). Then

$$\frac{d^n}{dx^n} [f(x)g(x)] = \sum_{k=0}^n \binom{n}{k} f^{(k)}(x) g^{(n-k)}(x)$$

Example B.1.

$$\frac{d^3}{dx^3} [f(x)g(x)] = f'''(x)g(x) + 3f''(x)g'(x) + 3f'(x)g''(x) + f(x)g'''(x)$$

**Theorem B.2** (Leibniz integration rule). <sup>3</sup>

$$\frac{d}{dx} \int_{a(x)}^{b(x)} g(t) dt = g[b(x)]b'(x) - g[a(x)]a'(x)$$

<sup>2</sup> Ben-Israel and Gilbert (2002) page 154, Leibniz (1710)

<sup>3</sup> Flanders (1973) page 615 (1.1) Talvila (2001), Knapp (2005b) page 389 (Chapter VII), ? page 422 (Leibniz Rule. Theorem 1.), <http://planetmath.org/encyclopedia/DifferentiationUnderIntegralSign.html>




# APPENDIX C

## TRIGONOMETRIC FUNCTIONS

### C.1 Definition Candidates

There are several ways of defining the sine and cosine functions, including the following:<sup>1</sup>

1. **Planar geometry:** Trigonometric functions are traditionally introduced as they have come to us historically—that is, as related to the parameters of triangles.<sup>2</sup>



$$\begin{aligned}\cos x &\triangleq \frac{x}{r} \\ \sin x &\triangleq \frac{y}{r}\end{aligned}$$

2. **Complex exponential:** The cosine and sine functions are the real and imaginary parts of the complex exponential such that<sup>3</sup>

$$\cos x \triangleq \mathbf{R}_e e^{ix} \quad \sin x \triangleq \mathbf{I}_m(e^{ix})$$

3. **Polynomial:** Let  $\sum_{n=0}^{\infty} x_n \triangleq \lim_{N \rightarrow \infty} \sum_{n=0}^N x_n$  in some topological space. The sine and cosine functions can be defined in terms of *Taylor expansions* such that<sup>4</sup>

$$\begin{aligned}\cos(x) &\triangleq \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n)!} x^{2n} = 1 - \frac{x^2}{2} + \frac{x^4}{4!} - \frac{x^6}{6!} + \dots \\ \sin(x) &\triangleq \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n+1)!} x^{2n+1} = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + \dots\end{aligned}$$

<sup>1</sup>The term *sine* originally came from the Hindu word *jiva* and later adapted to the Arabic word *jiba*. Abrabic-Latin translator [Robert of Chester](#) apparently confused this word with the Arabic word *jaiib*, which means “bay” or “inlet”—thus resulting in the Latin translation *sinus*, which also means “bay” or “inlet”. Reference: [Boyer and Merzbach \(1991\) page 252](#)

<sup>2</sup>[Abramowitz and Stegun \(1972\) page 78](#)

<sup>3</sup>[Euler \(1748\)](#)

<sup>4</sup>[Rosenlicht \(1968\) page 157](#), [Abramowitz and Stegun \(1972\) page 74](#)

4. **Product of factors:** Let  $\prod_{n=0}^{\infty} x_n \triangleq \lim_{N \rightarrow \infty} \prod_{n=0}^N x_n$  in some topological space. The sine and cosine functions can be defined in terms of a product of factors such that<sup>5</sup>

$$\cos(x) \triangleq \prod_{n=1}^{\infty} \left[ 1 - \left( \frac{x}{(2n-1)\frac{\pi}{2}} \right)^2 \right] \quad \sin(x) \triangleq x \prod_{n=1}^{\infty} \left[ 1 - \left( \frac{x}{n\pi} \right)^2 \right]$$

5. **Partial fraction expansion:** The sine function can be defined in terms of a partial fraction expansion such that<sup>6</sup>

$$\sin(x) \triangleq \frac{1}{\frac{1}{x} + 2x \sum_{n=1}^{\infty} \frac{(-1)^n}{x^2 - (n\pi)^2}} \quad \cos(x) \triangleq \underbrace{\left( \frac{1}{x} + 2x \sum_{n=1}^{\infty} \frac{1}{x^2 - (n\pi)^2} \right)}_{\cot(x)} \sin(x)$$




6. **Differential operator:** The sine and cosine functions can be defined as solutions to differential equations expressed in terms of the differential operator  $\frac{d}{dx}$  such that

$$\begin{array}{llll} \cos(x) \triangleq f(x) & \text{where} & \underbrace{\frac{d^2}{dx^2} f + f = 0}_{\text{differential equation}} & \underbrace{f(0) = 1}_{\text{1st initial condition}} & \underbrace{\left[ \frac{d}{dx} f \right](0) = 0}_{\text{2nd initial condition}} \\ \sin(x) \triangleq g(x) & \text{where} & \underbrace{\frac{d^2}{dx^2} g + g = 0}_{\text{differential equation}} & \underbrace{g(0) = 0}_{\text{1st initial condition}} & \underbrace{\left[ \frac{d}{dx} g \right](0) = 1}_{\text{2nd initial condition}} \end{array}$$

7. **Integral operator:** The sine and cosine functions can be defined as inverses of integrals of square roots of rational functions such that<sup>7</sup>

$$\begin{array}{ll} \cos(x) \triangleq f^{-1}(x) & \text{where } f(x) \triangleq \underbrace{\int_x^1 \sqrt{\frac{1}{1-y^2}} dy}_{\arccos(x)} \\ \sin(x) \triangleq g^{-1}(x) & \text{where } g(x) \triangleq \underbrace{\int_0^x \sqrt{\frac{1}{1-y^2}} dy}_{\arcsin(x)} \end{array}$$

For purposes of analysis, it can be argued that the more natural approach for defining harmonic functions is in terms of the differentiation operator  $\frac{d}{dx}$  (Definition C.1 page 75). Support for such an approach includes the following:

-  Both sine and cosine are very easily represented analytically as polynomials with coefficients involving the operator  $\frac{d}{dx}$  (Theorem C.1 page 76).
-  All solutions of homogeneous second order differential equations are linear combinations of sine and cosine (Theorem C.3 page 78).
-  Sine and cosine themselves are related to each other in terms of the differentiation operator (Theorem C.4 page 79).

<sup>5</sup>  Abramowitz and Stegun (1972) page 75

<sup>6</sup>  Abramowitz and Stegun (1972) page 75

<sup>7</sup>  Abramowitz and Stegun (1972) page 79

- 🔥 The complex exponential function is a solution of a second order homogeneous differential equation (Definition C.4 page 80).
- 🔥 Sine and cosine are orthogonal with respect to an innerproduct generated by an integral operator—which is a kind of inverse differential operator (Section C.6 page 88).

## C.2 Definitions

**Definition C.1.**<sup>8</sup> Let  $\mathcal{C}$  be the SPACE OF ALL CONTINUOUSLY DIFFERENTIABLE REAL FUNCTIONS and  $\frac{d}{dx} \in \mathcal{C}^{\mathcal{C}}$  the differentiation operator.

The function  $f \in \mathcal{C}^{\mathcal{C}}$  is the **cosine** function  $\cos(x) \triangleq f(x)$  if

1.  $\frac{d^2}{dx^2}f + f = 0$  (second order homogeneous differential equation) and
2.  $f(0) = 1$  (first initial condition) and
3.  $\left[\frac{d}{dx}f\right](0) = 0$  (second initial condition).

**Definition C.2.**<sup>9</sup> Let  $\mathcal{C}$  and  $\frac{d}{dx} \in \mathcal{C}^{\mathcal{C}}$  be defined as in definition of  $\cos(x)$  (Definition C.1 page 75).

The function  $f \in \mathcal{C}^{\mathcal{C}}$  is the **sine** function  $\sin(x) \triangleq f(x)$  if

1.  $\frac{d^2}{dx^2}f + f = 0$  (second order homogeneous differential equation) and
2.  $f(0) = 0$  (first initial condition) and
3.  $\left[\frac{d}{dx}f\right](0) = 1$  (second initial condition).

**Definition C.3.**<sup>10</sup>

Let  $\pi$  (“pi”) be defined as the element in  $\mathbb{R}$  such that

- (1).  $\cos\left(\frac{\pi}{2}\right) = 0$  and
- (2).  $\pi > 0$  and
- (3).  $\pi$  is the **smallest** of all elements in  $\mathbb{R}$  that satisfies (1) and (2).

## C.3 Basic properties

**Lemma C.1.**<sup>11</sup> Let  $\mathcal{C}$  be the SPACE OF ALL CONTINUOUSLY DIFFERENTIABLE REAL FUNCTIONS and  $\frac{d}{dx} \in \mathcal{C}^{\mathcal{C}}$  the differentiation operator.

$$\left\{ \begin{aligned} &\left\{ \frac{d^2}{dx^2}f + f = 0 \right\} \iff \\ &\left\{ \begin{aligned} f(x) &= \underbrace{[f](0) \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n}}{(2n)!}}_{\text{even terms}} + \underbrace{\left[\frac{d}{dx}f\right](0) \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{(2n+1)!}}_{\text{odd terms}} \\ &= \left( f(0) + \left[\frac{d}{dx}f\right](0)x \right) - \left( \frac{f(0)}{2!}x^2 + \frac{\left[\frac{d}{dx}f\right](0)}{3!}x^3 \right) + \left( \frac{f(0)}{4!}x^4 + \frac{\left[\frac{d}{dx}f\right](0)}{5!}x^5 \right) \dots \end{aligned} \right\} \end{aligned} \right.$$

<sup>8</sup> Rosenlicht (1968) page 157, Flanigan (1983) pages 228–229

<sup>9</sup> Rosenlicht (1968) page 157, Flanigan (1983) pages 228–229

<sup>10</sup> Rosenlicht (1968) page 158

<sup>11</sup> Rosenlicht (1968) page 156, Liouville (1839)

PROOF: Let  $f'(x) \triangleq \frac{d}{dx}f(x)$ .

$$\begin{aligned} f'''(x) &= -\left[\frac{d}{dx}f\right](x) \\ f^{(4)}(x) &= -\left[\frac{d}{dx}f\right](x) = -\left[\frac{d^2}{dx^2}f\right](x) = f(x) \end{aligned}$$

1. Proof that  $\left[\frac{d^2}{dx^2}f\right](x) + f(x) = 0 \implies f(x) = \sum_{n=0}^{\infty} (-1)^n \left[ \frac{f(0)}{(2n)!} x^{2n} + \frac{\left[\frac{d}{dx}f\right](0)}{(2n+1)!} x^{2n+1} \right]$ :

$$\begin{aligned} f(x) &= \sum_{n=0}^{\infty} \frac{f^{(n)}(0)}{n!} x^n \quad \text{by Taylor expansion} \\ &= f(0) + \left[\frac{d}{dx}f\right](0)x - \frac{\left[\frac{d^2}{dx^2}f\right](0)}{2!}x^2 - \frac{f^{(3)}(0)}{3!}x^3 + \frac{f^{(4)}(0)}{4!}x^4 + \frac{f^{(5)}(0)}{5!}x^5 - \dots \\ &= f(0) + \left[\frac{d}{dx}f\right](0)x - \frac{f(0)}{2!}x^2 - \frac{\left[\frac{d}{dx}f\right](0)}{3!}x^3 + \frac{f(0)}{4!}x^4 + \frac{\left[\frac{d}{dx}f\right](0)}{5!}x^5 - \dots \\ &= f(x) = \sum_{n=0}^{\infty} (-1)^n \left[ \frac{f(0)}{(2n)!} x^{2n} + \frac{\left[\frac{d}{dx}f\right](0)}{(2n+1)!} x^{2n+1} \right] \end{aligned}$$

2. Proof that  $\left[\frac{d^2}{dx^2}f\right](x) + f(x) = 0 \iff f(x) = \sum_{n=0}^{\infty} (-1)^n \left[ \frac{f(0)}{(2n)!} x^{2n} + \frac{\left[\frac{d}{dx}f\right](0)}{(2n+1)!} x^{2n+1} \right]$ :

$$\begin{aligned} \left[\frac{d^2}{dx^2}f\right](x) &= \frac{d}{dx} \frac{d}{dx} [f(x)] \\ &= \frac{d}{dx} \frac{d}{dx} \sum_{n=0}^{\infty} (-1)^n \left[ \frac{f(0)}{(2n)!} x^{2n} + \frac{\left[\frac{d}{dx}f\right](0)}{(2n+1)!} x^{2n+1} \right] && \text{by right hypothesis} \\ &= \sum_{n=1}^{\infty} (-1)^n \left[ \frac{(2n)(2n-1)f(0)}{(2n)!} x^{2n-2} + \frac{(2n+1)(2n)\left[\frac{d}{dx}f\right](0)}{(2n+1)!} x^{2n-1} \right] \\ &= \sum_{n=1}^{\infty} (-1)^n \left[ \frac{f(0)}{(2n-2)!} x^{2n-2} + \frac{\left[\frac{d}{dx}f\right](0)}{(2n-1)!} x^{2n-1} \right] \\ &= \sum_{n=0}^{\infty} (-1)^{n+1} \left[ \frac{f(0)}{(2n)!} x^{2n} + \frac{\left[\frac{d}{dx}f\right](0)}{(2n+1)!} x^{2n+1} \right] \\ &= -f(x) && \text{by right hypothesis} \end{aligned}$$

⇒

**Theorem C.1** (Taylor series for cosine/sine). <sup>12</sup>

<b>T H M</b>	$\cos(x) = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n}}{(2n)!} = 1 - \frac{x^2}{2} + \frac{x^4}{4!} - \frac{x^6}{6!} + \dots \quad \forall x \in \mathbb{R}$
	$\sin(x) = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{(2n+1)!} = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + \dots \quad \forall x \in \mathbb{R}$

<sup>12</sup> Rosenlicht (1968) page 157

 PROOF:

$$\cos(x) = \underbrace{f(0) \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n}}{(2n)!}}_{\text{even terms}} + \underbrace{\left[ \frac{d}{dx} f \right](0) \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{(2n+1)!}}_{\text{odd terms}}$$

by Lemma C.1 page 75

$$= 1 \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n}}{(2n)!} + 0 \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{(2n+1)!}$$

$$= \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n}}{(2n)!}$$

by cos initial conditions (Definition C.1 page 75)

$$\sin(x) = \underbrace{f(0) \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n}}{(2n)!}}_{\text{even terms}} + \underbrace{\left[ \frac{d}{dx} f \right](0) \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{(2n+1)!}}_{\text{odd terms}}$$

by Lemma C.1 page 75

$$= 0 \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n}}{(2n)!} + 1 \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{(2n+1)!}$$

$$= \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{(2n+1)!}$$

by sin initial conditions (Definition C.2 page 75)



### Theorem C.2. <sup>13</sup>

<b>T H M</b>	$\cos(0) = 1$	$\cos(-x) = \cos(x) \quad \forall x \in \mathbb{R}$
	$\sin(0) = 0$	$\sin(-x) = -\sin(x) \quad \forall x \in \mathbb{R}$

 PROOF:

$$\cos(0) = 1 - \frac{x^2}{2} + \frac{x^4}{4!} - \frac{x^6}{6!} + \dots \Big|_{x=0}$$

$$= 1$$

by Taylor series for cosine

(Theorem C.1 page 76)

$$\sin(0) = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + \dots \Big|_{x=0}$$

$$= 0$$

by Taylor series for sine

(Theorem C.1 page 76)

$$\cos(-x) = 1 - \frac{(-x)^2}{2} + \frac{(-x)^4}{4!} - \frac{(-x)^6}{6!} + \dots$$

$$= 1 - \frac{x^2}{2} + \frac{x^4}{4!} - \frac{x^6}{6!} + \dots$$

$$= \cos(x)$$

by Taylor series for cosine

(Theorem C.1 page 76)

$$\sin(-x) = (-x) - \frac{(-x)^3}{3!} + \frac{(-x)^5}{5!} - \frac{(-x)^7}{7!} + \dots$$

$$= - \left[ x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + \dots \right]$$

$$= \sin(x)$$

by Taylor series for cosine

(Theorem C.1 page 76)

by Taylor series for sine

(Theorem C.1 page 76)


by Taylor series for sine


(Theorem C.1 page 76)



### Lemma C.2. <sup>14</sup>

<b>L E M</b>	$\cos(1) > 0$	$x \in (0 : 2) \implies \sin(x) > 0$
	$\cos(2) < 0$	

<sup>13</sup>  Rosenlicht (1968) page 157

<sup>14</sup>  Rosenlicht (1968) page 158

✎ PROOF:

$$\begin{aligned} \cos(1) &= 1 - \frac{x^2}{2} + \frac{x^4}{4!} - \frac{x^6}{6!} + \dots \Big|_{x=1} && \text{by Taylor series for cosine} && (\text{Theorem C.1 page 76}) \\ &= 1 - \frac{1}{2} + \frac{1}{4!} - \frac{1}{6!} + \dots \\ &> 0 \end{aligned}$$

$$\begin{aligned} \cos(2) &= 1 - \frac{x^2}{2} + \frac{x^4}{4!} - \frac{x^6}{6!} + \dots \Big|_{x=2} && \text{by Taylor series for cosine} && (\text{Theorem C.1 page 76}) \\ &= 1 - \frac{4}{2} + \frac{16}{24} - \frac{64}{720} + \dots \\ &< 0 \end{aligned}$$

$$\begin{aligned} x \in (0 : 2) &\implies \text{each term in the sequence } \left( \left( x - \frac{x^3}{3!} \right), \left( \frac{x^5}{5!} - \frac{x^7}{7!} \right), \left( \frac{x^9}{9!} - \frac{x^{11}}{11!} \right), \dots \right) \text{ is } > 0 \\ &\implies \sin(x) > 0 \end{aligned}$$

⇒

**Proposition C.1.** Let  $\pi$  be defined as in Definition C.3 (page 75).

- P R P**
- (A). The value  $\pi$  **exists** in  $\mathbb{R}$ .  
 (B).  $2 < \pi < 4$ .

✎ PROOF:

$$\begin{aligned} \cos(1) &> 0 && \text{by Lemma C.2 page 77} \\ \cos(2) &< 0 && \text{by Lemma C.2 page 77} \\ &\implies 1 < \frac{\pi}{2} < 2 \\ &\implies 2 < \pi < 4 \end{aligned}$$

⇒

**Theorem C.3.** <sup>15</sup> Let  $\mathcal{C}$  be the space of all continuously differentiable real functions and  $\frac{d}{dx} \in \mathcal{C}^{\mathcal{C}}$  the differentiation operator. Let  $f'(0) \triangleq \left[ \frac{d}{dx} f \right](0)$ .

**T H M**

$$\left\{ \frac{d^2}{dx^2} f + f = 0 \right\} \iff \left\{ f(x) = f(0) \cos(x) + f'(0) \sin(x) \right\} \quad \forall f \in \mathcal{C}, \forall x \in \mathbb{R}$$

✎ PROOF:

1. Proof that  $\left[ \frac{d^2}{dx^2} f \right](x) = -f(x) \implies f(x) = f(0)\cos(x) + \left[ \frac{d}{dx} f \right](0)\sin(x)$ :

$$\begin{aligned} f(x) &= f(0) \underbrace{\sum_{n=0}^{\infty} (-1)^n \frac{x^{2n}}{(2n)!}}_{\cos(x)} + \left[ \frac{d}{dx} f \right](0) \underbrace{\sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{(2n+1)!}}_{\sin(x)} && \text{by left hypothesis and Lemma C.1 page 75} \\ &= f(0)\cos x + \left[ \frac{d}{dx} f \right](0)\sin x && \text{by definitions of cos and sin (Definition C.1 page 75, Definition C.2 page 75)} \end{aligned}$$

<sup>15</sup> Rosenlicht (1968) page 157. The general solution for the *non-homogeneous* equation  $\frac{d^2}{dx^2} f(x) + f(x) = g(x)$  with initial conditions  $f(a) = 1$  and  $f'(a) = \rho$  is  $f(x) = \cos(x) + \rho \sin(x) + \int_a^x g(y) \sin(x-y) dy$ . This type of equation is called a *Volterra integral equation of the second type*. References: Folland (1992) page 371, Liouville (1839). Volterra equation references: Pedersen (2000) page 99, Lalescu (1908), Lalescu (1911)



2. Proof that  $\frac{d^2}{dx^2}f = -f \iff f(x) = f(0)\cos(x) + \left[\frac{d}{dx}f\right](0)\sin(x)$  :

$$f(x) = f(0)\cos x + \left[\frac{d}{dx}f\right](0)\sin x \quad \text{by right hypothesis}$$

$$= f(0) \underbrace{\sum_{n=0}^{\infty} (-1)^n \frac{x^{2n}}{(2n)!}}_{\cos(x)} + \left[\frac{d}{dx}f\right](0) \underbrace{\sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{(2n+1)!}}_{\sin(x)}$$

$$\implies \frac{d^2}{dx^2}f + f = 0 \quad \text{by Lemma C.1 page 75}$$



**Theorem C.4.** <sup>16</sup> Let  $\frac{d}{dx} \in C^C$  be the differentiation operator.

<b>T H M</b>	$\frac{d}{dx}\cos(x) = -\sin(x) \quad \forall x \in \mathbb{R} \quad \Bigg  \quad \frac{d}{dx}\sin(x) = \cos(x) \quad \forall x \in \mathbb{R} \quad \Bigg  \quad \cos^2(x) + \sin^2(x) = 1 \quad \forall x \in \mathbb{R}$
----------------------	---

PROOF:

$$\frac{d}{dx}\cos(x) = \frac{d}{dx} \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n}}{(2n)!} \quad \text{by Taylor series} \quad (\text{Theorem C.1 page 76})$$

$$= \sum_{n=1}^{\infty} (-1)^n \frac{2nx^{2n-1}}{(2n)!} = \sum_{n=1}^{\infty} (-1)^n \frac{x^{2n-1}}{(2n-1)!} = \sum_{n=0}^{\infty} (-1)^{n+1} \frac{x^{2n}}{(2n)!}$$

$$= -\sin(x) \quad \text{by Taylor series} \quad (\text{Theorem C.1 page 76})$$

$$\frac{d}{dx}\sin(x) = \frac{d}{dx} \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{(2n+1)!} \quad \text{by Taylor series} \quad (\text{Theorem C.1 page 76})$$

$$= \sum_{n=0}^{\infty} (-1)^n \frac{(2n+1)x^{2n}}{(2n+1)!} = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n}}{(2n)!}$$

$$= \cos(x) \quad \text{by Taylor series} \quad (\text{Theorem C.1 page 76})$$

$$\frac{d}{dx} [\cos^2(x) + \sin^2(x)] = -2\cos(x)\sin(x) + 2\sin(x)\cos(x)$$

$$= 0$$

$$\implies \cos^2(x) + \sin^2(x) \text{ is constant}$$

$$\implies \cos^2(x) + \sin^2(x)$$

$$= \cos^2(0) + \sin^2(0)$$

$$= 1 + 0 = 1$$

by Theorem C.2 page 77



**Proposition C.2.**

<b>P R P</b>	$\sin\left(\frac{\pi}{2}\right) = 1$
----------------------	--------------------------------------

<sup>16</sup> Rosenlicht (1968) page 157

 PROOF:

$$\begin{aligned}
 \sin(\pi/2) &= \pm \sqrt{\sin^2(\pi/2) + 0} \\
 &= \pm \sqrt{\sin^2(\pi/2) + \cos^2(\pi/2)} && \text{by definition of } \pi && \text{(Definition C.3 page 75)} \\
 &= \pm \sqrt{1} && \text{by Theorem C.4 page 79} \\
 &= \pm 1 \\
 &= 1 && \text{by Lemma C.2 page 77}
 \end{aligned}$$



## C.4 The complex exponential

### Definition C.4.

The function  $f \in \mathbb{C}^{\mathbb{C}}$  is the **exponential function**  $\exp(ix) \triangleq f(x)$  if

DEF

1.  $\frac{d^2}{dx^2}f + f = 0$  (second order homogeneous differential equation) and
2.  $f(0) = 1$  (first initial condition) and
3.  $\left[\frac{d}{dx}f\right](0) = i$  (second initial condition).

### Theorem C.5 (Euler's identity). <sup>17</sup>

THEM

$$e^{ix} = \cos(x) + i\sin(x) \quad \forall x \in \mathbb{R}$$

 PROOF:

$$\begin{aligned}
 \exp(ix) &= f(0) \cos(x) + \left[\frac{d}{dx}f\right](0) \sin(x) && \text{by Theorem C.3 page 78} \\
 &= \cos(x) + i\sin(x) && \text{by Definition C.4 page 80}
 \end{aligned}$$



### Proposition C.3.

PRP

$$e^{-i\pi/2} = -i \mid e^{i\pi/2} = i$$

 PROOF:

$$\begin{aligned}
 e^{i\pi/2} &= \cos(\pi/2) + i\sin(\pi/2) && \text{by Euler's identity (Theorem C.5 page 80)} \\
 &= 0 + i && \text{by Theorem C.2 (page 77) and Proposition C.2 (page 79)} \\
 e^{-i\pi/2} &= \cos(-\pi/2) + i\sin(-\pi/2) && \text{by Euler's identity (Theorem C.5 page 80)} \\
 &= \cos(\pi/2) - i\sin(\pi/2) && \text{by Theorem C.2 page 77} \\
 &= 0 - i && \text{by Theorem C.2 (page 77) and Proposition C.2 (page 79)}
 \end{aligned}$$



### Corollary C.1.

COR

$$e^{ix} = \sum_{n \in \mathbb{W}} \frac{(ix)^n}{n!} \quad \forall x \in \mathbb{R}$$

<sup>17</sup>  Euler (1748),  Bottazzini (1986) page 12

✎ PROOF:

$$\begin{aligned}
 e^{ix} &= \cos(x) + i\sin(x) \\
 &= \underbrace{\sum_{n \in \mathbb{W}} \frac{(-1)^n x^{2n}}{(2n)!}}_{\cos(x)} + i \underbrace{\sum_{n \in \mathbb{W}} \frac{(-1)^n x^{2n+1}}{(2n+1)!}}_{\sin(x)} \\
 &= \sum_{n \in \mathbb{W}} \frac{(i^2)^n x^{2n}}{(2n)!} + \sum_{n \in \mathbb{W}} \frac{i(i^2)^n x^{2n+1}}{(2n+1)!} \\
 &= \sum_{n \in \mathbb{W} \cap \mathbb{Z}_e} \frac{(ix)^n}{n!} + \sum_{n \in \mathbb{W} \cap \mathbb{Z}_o} \frac{(ix)^n}{n!}
 \end{aligned}$$

by *Euler's identity*

(Theorem C.5 page 80)

by *Taylor series*

(Theorem C.1 page 76)

$$\begin{aligned}
 &= \sum_{n \in \mathbb{W}} \frac{(ix)^{2n}}{(2n)!} + \sum_{n \in \mathbb{W}} \frac{(ix)^{2n+1}}{(2n+1)!} \\
 &= \boxed{\sum_{n \in \mathbb{W}} \frac{(ix)^n}{n!}}
 \end{aligned}$$

⇒

### Corollary C.2 (Euler formulas).<sup>18</sup>

COR

$$\cos(x) = \mathbf{R}_e(e^{ix}) = \frac{e^{ix} + e^{-ix}}{2} \quad \forall x \in \mathbb{R} \quad \left| \quad \sin(x) = \mathbf{I}_m(e^{ix}) = \frac{e^{ix} - e^{-ix}}{2i} \quad \forall x \in \mathbb{R} \right.$$

✎ PROOF:

$$\begin{aligned}
 \mathbf{R}_e(e^{ix}) &\triangleq \frac{e^{ix} + (e^{ix})^*}{2} = \frac{e^{ix} + e^{-ix}}{2} \\
 &= \frac{\cos(x) + i\sin(x)}{2} + \frac{\cos(-x) + i\sin(-x)}{2} \\
 &= \frac{\cos(x) + i\sin(x)}{2} + \frac{\cos(x) - i\sin(x)}{2} \\
 \mathbf{I}_m(e^{ix}) &\triangleq \frac{e^{ix} - (e^{ix})^*}{2i} = \frac{e^{ix} - e^{-ix}}{2i} \\
 &= \frac{\cos(x) + i\sin(x)}{2i} - \frac{\cos(-x) + i\sin(-x)}{2i} \\
 &= \frac{\cos(x) + i\sin(x)}{2i} - \frac{\cos(x) - i\sin(x)}{2i}
 \end{aligned}$$

by definition of  $\Re$

(Definition E.5 page 115)

by *Euler's identity*

(Theorem C.5 page 80)

$$= \frac{\cos(x)}{2} + \frac{\cos(x)}{2} = \boxed{\cos(x)}$$

by definition of  $\Im$

(Definition E.5 page 115)

by *Euler's identity*

(Theorem C.5 page 80)

$$= \frac{i\sin(x)}{2i} + \frac{i\sin(x)}{2i} = \boxed{\sin(x)}$$

⇒

### Theorem C.6.<sup>19</sup>

THM

$$e^{(\alpha+\beta)} = e^\alpha e^\beta \quad \forall \alpha, \beta \in \mathbb{C}$$

✎ PROOF:

$$\begin{aligned}
 e^\alpha e^\beta &= \left( \sum_{n \in \mathbb{W}} \frac{\alpha^n}{n!} \right) \left( \sum_{m \in \mathbb{W}} \frac{\beta^m}{m!} \right) \\
 &= \sum_{n \in \mathbb{W}} \sum_{k=0}^n \frac{\alpha^k}{k!} \frac{\beta^{n-k}}{(n-k)!} \\
 &= \sum_{n \in \mathbb{W}} \sum_{k=0}^n \frac{n!}{n!} \frac{\alpha^k}{k!} \frac{\beta^{n-k}}{(n-k)!}
 \end{aligned}$$

by Corollary C.1 page 80

<sup>18</sup> Euler (1748), Bottazzini (1986) page 12

<sup>19</sup> Rudin (1987) page 1

$$\begin{aligned}
&= \sum_{n \in \mathbb{W}} \frac{1}{n!} \sum_{k=0}^n \frac{n!}{k!(n-k)!} \alpha^k \beta^{n-k} \\
&= \sum_{n \in \mathbb{W}} \frac{1}{n!} \sum_{k=0}^n \binom{n}{k} \alpha^k \beta^{n-k} \\
&= \sum_{n \in \mathbb{W}} \frac{(\alpha + \beta)^n}{n!} && \text{by the Binomial Theorem} \\
&= e^{\alpha + \beta} && \text{by Corollary C.1 page 80}
\end{aligned}$$



## C.5 Trigonometric Identities

**Theorem C.7** (shift identities).

<b>T H M</b>	$\cos\left(x + \frac{\pi}{2}\right) = -\sin x \quad \forall x \in \mathbb{R}$	$\sin\left(x + \frac{\pi}{2}\right) = \cos x \quad \forall x \in \mathbb{R}$
	$\cos\left(x - \frac{\pi}{2}\right) = \sin x \quad \forall x \in \mathbb{R}$	$\sin\left(x - \frac{\pi}{2}\right) = -\cos x \quad \forall x \in \mathbb{R}$

PROOF:

$$\begin{aligned}
\cos\left(x + \frac{\pi}{2}\right) &= \frac{e^{i\left(x + \frac{\pi}{2}\right)} + e^{-i\left(x + \frac{\pi}{2}\right)}}{2} && \text{by Euler formulas} && (\text{Corollary C.2 page 81}) \\
&= \frac{e^{ix} e^{i\frac{\pi}{2}} + e^{-ix} e^{-i\frac{\pi}{2}}}{2} && \text{by } e^{\alpha\beta} = e^\alpha e^\beta \text{ result} && (\text{Theorem C.6 page 81}) \\
&= \frac{e^{ix}(i) + e^{-ix}(-i)}{2} && \text{by Proposition C.3 page 80} \\
&= \frac{e^{ix} - e^{-ix}}{-2i} \\
&= -\sin x && \text{by Euler formulas} && (\text{Corollary C.2 page 81}) \\
\cos\left(x - \frac{\pi}{2}\right) &= \frac{e^{i\left(x - \frac{\pi}{2}\right)} + e^{-i\left(x - \frac{\pi}{2}\right)}}{2} && \text{by Euler formulas} && (\text{Corollary C.2 page 81}) \\
&= \frac{e^{ix} e^{-i\frac{\pi}{2}} + e^{-ix} e^{i\frac{\pi}{2}}}{2} && \text{by } e^{\alpha\beta} = e^\alpha e^\beta \text{ result} && (\text{Theorem C.6 page 81}) \\
&= \frac{e^{ix}(-i) + e^{-ix}(i)}{2} && \text{by Proposition C.3 page 80} \\
&= \frac{e^{ix} - e^{-ix}}{2i} \\
&= \sin x && \text{by Euler formulas} && (\text{Corollary C.2 page 81}) \\
\sin\left(x + \frac{\pi}{2}\right) &= \cos\left(\left[x + \frac{\pi}{2}\right] - \frac{\pi}{2}\right) && \text{by previous result} \\
&= \cos(x) \\
\sin\left(x - \frac{\pi}{2}\right) &= -\cos\left(\left[x - \frac{\pi}{2}\right] + \frac{\pi}{2}\right) && \text{by previous result} \\
&= -\cos(x)
\end{aligned}$$



**Theorem C.8** (product identities).T  
H  
M

$$\begin{aligned}
 (A). \quad \cos x \cos y &= \frac{1}{2} \cos(x-y) + \frac{1}{2} \cos(x+y) \quad \forall x, y \in \mathbb{R} \\
 (B). \quad \cos x \sin y &= -\frac{1}{2} \sin(x-y) + \frac{1}{2} \sin(x+y) \quad \forall x, y \in \mathbb{R} \\
 (C). \quad \sin x \cos y &= \frac{1}{2} \sin(x-y) + \frac{1}{2} \sin(x+y) \quad \forall x, y \in \mathbb{R} \\
 (D). \quad \sin x \sin y &= \frac{1}{2} \cos(x-y) - \frac{1}{2} \cos(x+y) \quad \forall x, y \in \mathbb{R}
 \end{aligned}$$

✎ PROOF:

1. Proof for (A) using *Euler formulas* (Corollary C.2 page 81)  
(algebraic method requiring *complex number system*  $\mathbb{C}$ ):

$$\begin{aligned}
 \cos x \cos y &= \left( \frac{e^{ix} + e^{-ix}}{2} \right) \left( \frac{e^{iy} + e^{-iy}}{2} \right) && \text{by Euler formulas} && (\text{Corollary C.2 page 81}) \\
 &= \frac{e^{i(x+y)} + e^{i(x-y)} + e^{i(-x+y)} + e^{i(-x-y)}}{4} \\
 &= \frac{e^{i(x+y)} + e^{-i(x+y)}}{4} + \frac{e^{i(x-y)} + e^{-i(x-y)}}{4} \\
 &= \frac{2\cos(x+y)}{4} + \frac{2\cos(x-y)}{4} && \text{by Euler formulas} && (\text{Corollary C.2 page 81}) \\
 &= \frac{1}{2} \cos(x+y) + \frac{1}{2} \cos(x-y)
 \end{aligned}$$

2. Proof for (A) using *Volterra integral equation* (Theorem C.3 page 78)  
(differential equation method requiring only *real number system*  $\mathbb{R}$ ):

$$\begin{aligned}
 f(x) &\triangleq \frac{1}{2} \cos(x-y) + \frac{1}{2} \cos(x+y) \\
 \Rightarrow \frac{d}{dx} f(x) &= -\frac{1}{2} \sin(x-y) - \frac{1}{2} \sin(x+y) && \text{by Theorem C.4 page 79} \\
 \Rightarrow \frac{d^2}{dx^2} f(x) &= -\frac{1}{2} \cos(x-y) - \frac{1}{2} \cos(x+y) && \text{by Theorem C.4 page 79} \\
 \Rightarrow \frac{d^2}{dx^2} f(x) + f(x) &= 0 && \text{by additive inverse property} \\
 \Rightarrow \underbrace{\frac{1}{2} \cos(x-y) + \frac{1}{2} \cos(x+y)}_{f(x)} &= \underbrace{[\frac{1}{2} \cos(0-y) + \frac{1}{2} \cos(0+y)]}_{f''(0)} \cos(x) + \underbrace{[-\frac{1}{2} \sin(0-y) - \frac{1}{2} \sin(0+y)]}_{f'(0)} \sin(x) \\
 \Rightarrow \frac{1}{2} \cos(x-y) + \frac{1}{2} \cos(x+y) &= \cos y \cos x + 0 \sin(x) \\
 \Rightarrow \cos x \cos y &= \frac{1}{2} \cos(x-y) + \frac{1}{2} \cos(x+y)
 \end{aligned}$$

3. Proof for (B) using *Euler formulas* (Corollary C.2 page 81):

$$\begin{aligned}
 \sin x \sin y &= \left( \frac{e^{ix} - e^{-ix}}{2i} \right) \left( \frac{e^{iy} - e^{-iy}}{2i} \right) && \text{by Corollary C.2 page 81} \\
 &= \frac{e^{i(x+y)} - e^{i(x-y)} - e^{i(-x+y)} + e^{i(-x-y)}}{-4} \\
 &= \frac{e^{i(x+y)} + e^{-i(x+y)} - e^{i(x-y)} - e^{-i(x-y)}}{-4} \\
 &= \frac{e^{i(x+y)} + e^{-i(x+y)}}{-4} - \frac{e^{i(x-y)} + e^{-i(x-y)}}{-4} \\
 &= \frac{2\cos(x+y)}{-4} - \frac{2\cos(x-y)}{-4} \\
 &= \frac{1}{2} \cos(x+y) - \frac{1}{2} \cos(x-y)
 \end{aligned}$$

by Corollary C.2 page 81

4. Proofs for (C) and (D) using (A) and (B):

$$\begin{aligned}
 \cos x \sin y &= \cos(x) \cos\left(y - \frac{\pi}{2}\right) && \text{by shift identities} && (\text{Theorem C.7 page 82}) \\
 &= \frac{1}{2} \cos\left(x + y - \frac{\pi}{2}\right) + \frac{1}{2} \cos\left(x - y + \frac{\pi}{2}\right) && \text{by (A)} \\
 &= \frac{1}{2} \sin(x + y) - \frac{1}{2} \sin(x - y) && \text{by shift identities} && (\text{Theorem C.7 page 82}) \\
 \sin x \cos y &= \cos y \sin x \\
 &= \frac{1}{2} \sin(y + x) - \frac{1}{2} \sin(y - x) && \text{by (B)} \\
 &= \frac{1}{2} \sin(x + y) + \frac{1}{2} \sin(x - y) && \text{by Theorem C.2 page 77}
 \end{aligned}$$

⇒

### Proposition C.4.

P R P	(A). $\cos(\pi) = -1$	(C). $\cos(2\pi) = 1$	(E). $e^{i\pi} = -1$
	(B). $\sin(\pi) = 0$	(D). $\sin(2\pi) = 0$	(F). $e^{i2\pi} = 0$

✎ PROOF:

$$\begin{aligned}
 \cos(\pi) &= -1 + 1 + \cos(\pi) \\
 &= -1 + 2\left[\frac{1}{2}\cos(\pi/2 - \pi/2) + \frac{1}{2}\cos(\pi/2 + \pi/2)\right] && \text{by } \cos(0) = 1 \text{ result} && (\text{Theorem C.2 page 77}) \\
 &= -1 + 2\cos(\pi/2)\cos(\pi/2) && \text{by product identities} && (\text{Theorem C.8 page 82}) \\
 &= -1 + 2(0)(0) && \text{by definition of } \pi && (\text{Definition C.3 page 75}) \\
 &= -1 \\
 \sin(\pi) &= 0 + \sin(\pi) \\
 &= 2\left[-\frac{1}{2}\sin(\pi/2 - \pi/2) + \frac{1}{2}\sin(\pi/2 + \pi/2)\right] && \text{by } \sin(0) = 0 \text{ result} && (\text{Theorem C.2 page 77}) \\
 &= 2\cos(\pi/2)\sin(\pi/2) && \text{by product identities} && (\text{Theorem C.8 page 82}) \\
 &= 2(0)\sin(\pi/2) && \text{by definition of } \pi && (\text{Definition C.3 page 75}) \\
 &= 0 \\
 \cos(2\pi) &= 1 + \cos(2\pi) - 1 \\
 &= 2\left[\frac{1}{2}\cos(\pi - \pi) + \frac{1}{2}\cos(\pi + \pi)\right] - 1 && \text{by } \cos(0) = 1 \text{ result} && (\text{Theorem C.2 page 77}) \\
 &= 2\cos(\pi)\cos(\pi) - 1 && \text{by product identities} && (\text{Theorem C.8 page 82}) \\
 &= 2(-1)(-1) - 1 && \text{by (A)} \\
 &= 1 \\
 \sin(2\pi) &= 0 + \sin(2\pi) \\
 &= 2\left[\frac{1}{2}\sin(\pi - \pi) + \frac{1}{2}\sin(\pi + \pi)\right] && \text{by } \sin(0) = 0 \text{ result} && (\text{Theorem C.2 page 77}) \\
 &= 2\sin(\pi)\cos(\pi) && \text{by product identities} && (\text{Theorem C.8 page 82}) \\
 &= 2(0)(-1) && \text{by (A) and (B)} \\
 &= 0 \\
 e^{i\pi} &= \cos(\pi) + i\sin(\pi) && \text{by Euler's identity} && (\text{Theorem C.5 page 80}) \\
 &= -1 + 0 \\
 &= -1 && \text{by (A) and (B)} \\
 e^{i2\pi} &= \cos(2\pi) + i\sin(2\pi) && \text{by Euler's identity} && (\text{Theorem C.5 page 80}) \\
 &= 1 + 0 \\
 &= 1 && \text{by (C) and (D)}
 \end{aligned}$$

⇒

**Theorem C.9** (double angle formulas). <sup>20</sup>T  
H  
M

(A).	$\cos(x + y) = \cos x \cos y - \sin x \sin y$	$\forall x, y \in \mathbb{R}$
(B).	$\sin(x + y) = \sin x \cos y + \cos x \sin y$	$\forall x, y \in \mathbb{R}$
(C).	$\tan(x + y) = \frac{\tan x + \tan y}{1 - \tan x \tan y}$	$\forall x, y \in \mathbb{R}$

✎ PROOF:

1. Proof for (A) using *product identities* (Theorem C.8 page 82).

$$\begin{aligned}
 \cos(x + y) &= \underbrace{\frac{1}{2}\cos(x + y) + \frac{1}{2}\cos(x + y)}_{\cos(x + y)} + \underbrace{\frac{1}{2}\cos(x - y) - \frac{1}{2}\cos(x - y)}_0 \\
 &= \left[ \frac{1}{2}\cos(x - y) + \frac{1}{2}\cos(x + y) \right] - \left[ \frac{1}{2}\cos(x - y) - \frac{1}{2}\cos(x + y) \right] \\
 &= \cos x \cos y - \sin x \sin y
 \end{aligned}$$

by Theorem C.8 page 82

2. Proof for (A) using *Volterra integral equation* (Theorem C.3 page 78):

$$\begin{aligned}
 f(x) \triangleq \cos(x + y) &\implies \frac{d}{dx}f(x) = -\sin(x + y) && \text{by Theorem C.4 page 79} \\
 &\implies \frac{d^2}{dx^2}f(x) = -\cos(x + y) && \text{by Theorem C.4 page 79} \\
 &\implies \frac{d^2}{dx^2}f(x) + f(x) = 0 && \text{by additive inverse property} \\
 &\implies \cos(x + y) = \cos y \cos x - \sin y \sin x && \text{by Theorem C.3 page 78} \\
 &\implies \cos(x + y) = \cos x \cos y - \sin x \sin y && \text{by commutative property}
 \end{aligned}$$

3. Proof for (B) and (C) using (A):

$$\begin{aligned}
 \sin(x + y) &= \cos\left(x - \frac{\pi}{2} + y\right) && \text{by shift identities (Theorem C.7 page 82)} \\
 &= \cos\left(x - \frac{\pi}{2}\right)\cos(y) - \sin\left(x - \frac{\pi}{2}\right)\sin(y) && \text{by (A)} \\
 &= \sin(x)\cos(y) + \cos(x)\sin(y) && \text{by shift identities (Theorem C.7 page 82)}
 \end{aligned}$$

$$\begin{aligned}
 \tan(x + y) &= \frac{\sin(x + y)}{\cos(x + y)} \\
 &= \frac{\sin x \cos y + \cos x \sin y}{\cos x \cos y - \sin x \sin y} && \text{by (A)} \\
 &= \left( \frac{\sin x \cos y + \cos x \sin y}{\cos x \cos y - \sin x \sin y} \right) \left( \frac{\cos x \cos y}{\cos x \cos y} \right) \\
 &= \frac{\frac{\sin x \cos y}{\cos x \cos y} + \frac{\cos x \sin y}{\cos x \cos y}}{\frac{\cos x \cos y}{\cos x \cos y} - \frac{\sin x \sin y}{\cos x \cos y}} = \frac{\frac{\sin x}{\cos x} + \frac{\sin y}{\cos y}}{1 - \frac{\sin x}{\cos x} \frac{\sin y}{\cos y}} = \frac{\tan x + \tan y}{1 - \tan x \tan y}
 \end{aligned}$$

⇒

**Theorem C.10** (trigonometric periodicity).T  
H  
M

(A).	$\cos(x + M\pi) = (-1)^M \cos(x)$	$\forall x \in \mathbb{R}, M \in \mathbb{Z}$
(B).	$\sin(x + M\pi) = (-1)^M \sin(x)$	$\forall x \in \mathbb{R}, M \in \mathbb{Z}$
(C).	$e^{i(x + M\pi)} = (-1)^M e^{ix}$	$\forall x \in \mathbb{R}, M \in \mathbb{Z}$
(D).	$\cos(x + 2M\pi) = \cos(x)$	$\forall x \in \mathbb{R}, M \in \mathbb{Z}$
(E).	$\sin(x + 2M\pi) = \sin(x)$	$\forall x \in \mathbb{R}, M \in \mathbb{Z}$
(F).	$e^{i(x + 2M\pi)} = e^{ix}$	$\forall x \in \mathbb{R}, M \in \mathbb{Z}$

<sup>20</sup>Expressions for  $\cos(\alpha + \beta)$ ,  $\sin(\alpha + \beta)$ , and  $\sin^2 x$  appear in works as early as **Ptolemy** (circa 100AD). Reference: [http://en.wikipedia.org/wiki/History\\_of\\_trigonometric\\_functions](http://en.wikipedia.org/wiki/History_of_trigonometric_functions)

✎ PROOF:

1. Proof for (A):

(a)  $M = 0$  case:  $\cos(x + 0\pi) = \cos(x) = (-1)^0 \cos(x)$

(b) Proof for  $M > 0$  cases (by induction):

i. Base case  $M = 1$ :

$$\begin{aligned} \cos(x + \pi) &= \cos x \cos \pi - \sin x \sin \pi && \text{by double angle formulas} && (\text{Theorem C.9 page 85}) \\ &= \cos x (-1) - \sin x (0) && \text{by } \cos \pi = -1 \text{ result} && (\text{Proposition C.4 page 84}) \\ &= (-1)^1 \cos x \end{aligned}$$

ii. Inductive step...Proof that  $M$  case  $\implies M + 1$  case:

$$\begin{aligned} \cos(x + [M + 1]\pi) &= \cos([x + \pi] + M\pi) \\ &= (-1)^M \cos(x + \pi) && \text{by induction hypothesis (M case)} \\ &= (-1)^M (-1) \cos(x) && \text{by base case (item (1b)i) page 86)} \\ &= (-1)^{M+1} \cos(x) \\ &\implies M + 1 \text{ case} \end{aligned}$$

(c) Proof for  $M < 0$  cases: Let  $N \triangleq -M \dots \implies N > 0$ .

$$\begin{aligned} \cos(x + M\pi) &\triangleq \cos(x - N\pi) && \text{by definition of } N \\ &= \cos(x) \cos(-N\pi) - \sin(x) \sin(-N\pi) && \text{by double angle formulas} && (\text{Theorem C.9 page 85}) \\ &= \cos(x) \cos(N\pi) + \sin(x) \sin(N\pi) && \text{by Theorem C.2 page 77} \\ &= \cos(x) \cos(0 + N\pi) + \sin(x) \sin(0 + N\pi) \\ &= \cos(x) (-1)^N \cos(0) + \sin(x) (-1)^N \sin(0) && \text{by } M \geq 0 \text{ results} && (\text{item (1b) page 86}) \\ &= (-1)^N \cos(x) && \text{by } \cos(0)=1, \sin(0)=0 \text{ results} && (\text{Theorem C.2 page 77}) \\ &\triangleq (-1)^{-M} \cos(x) && \text{by definition of } N \\ &= (-1)^M \cos(x) \end{aligned}$$

(d) Proof using complex exponential:

$$\begin{aligned} \cos(x + M\pi) &= \frac{e^{i(x+M\pi)} + e^{-i(x+M\pi)}}{2} && \text{by Euler formulas} && (\text{Corollary C.2 page 81}) \\ &= e^{iM\pi} \left[ \frac{e^{ix} + e^{-ix}}{2} \right] && \text{by } e^{\alpha\beta} = e^\alpha e^\beta \text{ result} && (\text{Theorem C.6 page 81}) \\ &= (e^{i\pi})^M \cos x && \text{by Euler formulas} && (\text{Corollary C.2 page 81}) \\ &= (-1)^M \cos x && \text{by } e^{i\pi} = -1 \text{ result} && (\text{Proposition C.4 page 84}) \end{aligned}$$

2. Proof for (B):

(a)  $M = 0$  case:  $\sin(x + 0\pi) = \sin(x) = (-1)^0 \sin(x)$

(b) Proof for  $M > 0$  cases (by induction):

i. Base case  $M = 1$ :

$$\begin{aligned} \sin(x + \pi) &= \sin x \cos \pi + \cos x \sin \pi && \text{by double angle formulas} && (\text{Theorem C.9 page 85}) \\ &= \sin x (-1) - \cos x (0) && \text{by } \sin \pi = 0 \text{ results} && (\text{Proposition C.4 page 84}) \\ &= (-1)^1 \sin x \end{aligned}$$



ii. Inductive step...Proof that  $M$  case  $\implies M + 1$  case:

$$\begin{aligned}
 \sin(x + [M + 1]\pi) &= \sin([x + \pi] + M\pi) \\
 &= (-1)^M \sin(x + \pi) && \text{by induction hypothesis (M case)} \\
 &= (-1)^M (-1) \sin(x) && \text{by base case (item (2b)i) page 86)} \\
 &= (-1)^{M+1} \sin(x) \\
 &\implies M + 1 \text{ case}
 \end{aligned}$$

(c) Proof for  $M < 0$  cases: Let  $N \triangleq -M \dots \implies N > 0$ .

$$\begin{aligned}
 \sin(x + M\pi) &\triangleq \sin(x - N\pi) && \text{by definition of } N \\
 &= \sin(x)\sin(-N\pi) - \sin(x)\sin(-N\pi) && \text{by double angle formulas (Theorem C.9 page 85)} \\
 &= \sin(x)\sin(N\pi) + \sin(x)\sin(N\pi) && \text{by Theorem C.2 page 77} \\
 &= \sin(x)\sin(0 + N\pi) + \sin(x)\sin(0 + N\pi) \\
 &= \sin(x)(-1)^N \sin(0) + \sin(x)(-1)^N \sin(0) && \text{by } M \geq 0 \text{ results (item (2b) page 86)} \\
 &= (-1)^N \sin(x) && \text{by } \sin(0)=1, \sin(0)=0 \text{ results (Theorem C.2 page 77)} \\
 &\triangleq (-1)^{-M} \sin(x) && \text{by definition of } N \\
 &= (-1)^M \sin(x)
 \end{aligned}$$

(d) Proof using complex exponential:

$$\begin{aligned}
 \sin(x + M\pi) &= \frac{e^{i(x+M\pi)} - e^{-i(x+M\pi)}}{2i} && \text{by Euler formulas (Corollary C.2 page 81)} \\
 &= e^{iM\pi} \left[ \frac{e^{ix} - e^{-ix}}{2i} \right] && \text{by } e^{\alpha\beta} = e^\alpha e^\beta \text{ result (Theorem C.6 page 81)} \\
 &= (e^{i\pi})^M \sin x && \text{by Euler formulas (Corollary C.2 page 81)} \\
 &= (-1)^M \sin x && \text{by } e^{i\pi} = -1 \text{ result (Proposition C.4 page 84)}
 \end{aligned}$$

3. Proof for (C):

$$\begin{aligned}
 e^{i(x+M\pi)} &= e^{iM\pi} e^{ix} && \text{by } e^{\alpha\beta} = e^\alpha e^\beta \text{ result (Theorem C.6 page 81)} \\
 &= (e^{i\pi})^M (e^{ix}) \\
 &= (-1)^M e^{ix} && \text{by } e^{i\pi} = -1 \text{ result (Proposition C.4 page 84)}
 \end{aligned}$$

4. Proofs for (D), (E), and (F):

$$\begin{aligned}
 \cos(i(x + 2M\pi)) &= (-1)^{2M} \cos(ix) = \cos(ix) && \text{by (A)} \\
 \sin(i(x + 2M\pi)) &= (-1)^{2M} \sin(ix) = \sin(ix) && \text{by (B)} \\
 e^{i(x+2M\pi)} &= (-1)^{2M} e^{ix} = e^{ix} && \text{by (C)}
 \end{aligned}$$


**Theorem C.11** (half-angle formulas/squared identities).

<b>T H M</b>	(A). $\cos^2 x = \frac{1}{2}(1 + \cos 2x) \quad \forall x \in \mathbb{R}$	(C). $\cos^2 x + \sin^2 x = 1 \quad \forall x \in \mathbb{R}$
	(B). $\sin^2 x = \frac{1}{2}(1 - \cos 2x) \quad \forall x \in \mathbb{R}$	

PROOF:

$$\begin{aligned}
 \cos^2 x &\triangleq (\cos x)(\cos x) = \frac{1}{2}\cos(x - x) + \frac{1}{2}\cos(x + x) && \text{by product identities (Theorem C.8 page 82)} \\
 &= \frac{1}{2}[1 + \cos(2x)] && \text{by } \cos(0) = 1 \text{ result (Theorem C.2 page 77)} \\
 \sin^2 x &= (\sin x)(\sin x) = \frac{1}{2}\cos(x - x) - \frac{1}{2}\cos(x + x) && \text{by product identities (Theorem C.8 page 82)} \\
 &= \frac{1}{2}[1 - \cos(2x)] && \text{by } \cos(0) = 1 \text{ result (Theorem C.2 page 77)} \\
 \cos^2 x + \sin^2 x &= \frac{1}{2}[1 + \cos(2x)] + \frac{1}{2}[1 - \cos(2x)] = 1 && \text{by (A) and (B)} \\
 &&& \text{note: see also Theorem C.4 page 79}
 \end{aligned}$$



## C.6 Planar Geometry

The harmonic functions  $\cos(x)$  and  $\sin(x)$  are *orthogonal* to each other in the sense


$$\begin{aligned}
 \langle \cos(x) | \sin(x) \rangle &= \int_{-\pi}^{+\pi} \cos(x) \sin(x) \, dx \\
 &= \frac{1}{2} \int_{-\pi}^{+\pi} \sin(x-x) \, dx + \frac{1}{2} \int_{-\pi}^{+\pi} \sin(x+x) \, dx && \text{by Theorem C.8 page 82} \\
 &= \frac{1}{2} \int_{-\pi}^{+\pi} \sin(0) \, dx + \frac{1}{2} \int_{-\pi}^{+\pi} \sin(2x) \, dx \\
 &= -\frac{1}{2} \cdot \frac{1}{2} \cos(2x) \Big|_{-\pi}^{+\pi} \cos(2x) \\
 &= -\frac{1}{4} [\cos(2\pi) - \cos(-2\pi)] \\
 &= 0
 \end{aligned}$$

Because  $\cos(x)$  and  $\sin(x)$  are orthogonal, they can be conveniently represented by the  $x$  and  $y$  axes in a plane—because perpendicular axes in a plane are also orthogonal. Vectors in the plane can be represented by linear combinations of  $\cos x$  and  $\sin x$ . Let  $\tan x$  be defined as

$$\tan x \triangleq \frac{\sin x}{\cos x}.$$


We can also define a value  $\theta$  to represent the angle between such a vector and the  $x$ -axis such that

$$\theta = \tan^{-1} \left( \frac{\sin \theta}{\cos \theta} \right)$$



$$\begin{array}{ll}
 \cos \theta \triangleq \frac{x}{r} & \sec \theta \triangleq \frac{r}{x} \\
 \sin \theta \triangleq \frac{y}{r} & \csc \theta \triangleq \frac{r}{y} \\
 \tan \theta \triangleq \frac{y}{x} & \cot \theta \triangleq \frac{x}{y}
 \end{array}$$

## C.7 The power of the exponential




“Gentlemen, that is surely true, it is absolutely paradoxical; we cannot understand it, and we don't know what it means. But we have proved it, and therefore we know it must be the truth.”

Benjamin Peirce (1809–1880), American Harvard University mathematician after proving  $e^{i\pi} = -1$  in a lecture. <sup>21</sup>

<sup>21</sup> quote: [Kasner and Newman \(1940\) page 104](#)

image: [http://www-history.mcs.st-andrews.ac.uk/history/PictDisplay/Peirce\\_Benjamin.html](http://www-history.mcs.st-andrews.ac.uk/history/PictDisplay/Peirce_Benjamin.html)



“Young man, in mathematics you don't understand things. You just get used to them.”

John von Neumann (1903–1957), Hungarian-American mathematician, as allegedly told to Gary Zukav by Felix T. Smith, Head of Molecular Physics at Stanford Research Institute, about a “physicist friend”.<sup>22</sup>

The following corollary presents one of the most amazing relations in all of mathematics. It shows a simple and compact relationship between the transcendental numbers  $\pi$  and  $e$ , the imaginary number  $i$ , and the additive and multiplicative identity elements 0 and 1. The fact that there is any relationship at all is somewhat amazing; but for there to be such an elegant one is truly one of the wonders of the world of numbers.

### Corollary C.3.<sup>23</sup>

C  
O  
R

 $e^{i\pi} + 1 = 0$

PROOF:

$$\begin{aligned} e^{ix} \Big|_{x=\pi} &= [\cos x + i \sin x]_{x=\pi} \\ &= -1 + i \cdot 0 \\ &= -1 \end{aligned}$$

by Euler's identity (Theorem C.5 page 80)

by Proposition C.4 page 84

⇒

There are many transforms available, several of them integral transforms  $[Af](s) \triangleq \int_t f(s) \kappa(t, s) \, ds$  using different kernels  $\kappa(t, s)$ . But of all of them, two of the most often used themselves use an exponential kernel:

- ① The *Laplace Transform* with kernel  $\kappa(t, s) \triangleq e^{st}$
- ② The *Fourier Transform* with kernel  $\kappa(t, \omega) \triangleq e^{i\omega t}$ .

Of course, the Fourier kernel is just a special case of the Laplace kernel with  $s = i\omega$  ( $i\omega$  is a unit circle in  $s$  if  $s$  is depicted as a plane with real and imaginary axes). What is so special about exponential kernels? Is it just that they were discovered sooner than other kernels with other transforms? The answer in general is “no”. The exponential has two properties that makes it extremely special:

The exponential is an eigenvalue of any *linear time invariant* (LTI) operator (Theorem C.12 page 89).

The exponential generates a *continuous point spectrum* for the *differential operator*.

**Theorem C.12.**<sup>24</sup> Let  $L$  be an operator with kernel  $h(t, \omega)$  and

$$\check{h}(s) \triangleq \langle h(t, \omega) | e^{st} \rangle \quad (\text{LAPLACE TRANSFORM}).$$

<sup>22</sup> quote: Zukav (1980) page 208

image: [http://en.wikipedia.org/wiki/John\\_von\\_Neumann](http://en.wikipedia.org/wiki/John_von_Neumann)

The quote appears in a footnote in Zukav (1980) that reads like this: Dr. Felix Smith, Head of Molecular Physics, Stanford Research Institute, once related to me the true story of a physicist friend who worked at Los Alamos after World War II. Seeking help on a difficult problem, he went to the great Hungarian mathematician, John von Neumann, who was at Los Alamos as a consultant. “Simple,” said von Neumann. “This can be solved by using the method of characteristics.” After the explanation the physicist said, “I’m afraid I don’t understand the method of characteristics.” “Young man,” said von Neumann, “in mathematics you don’t understand things, you just get used to them.”

<sup>23</sup> Euler (1748), Euler (1988) (chapter 8?), [http://www.daviddarling.info/encyclopedia/E/Eulers\\_formula.html](http://www.daviddarling.info/encyclopedia/E/Eulers_formula.html)

<sup>24</sup> Mallat (1999) page 2, ...page 2 online: <http://www.cmap.polytechnique.fr/~mallat/WTintro.pdf>

T  
H  
M

$$\left\{ \begin{array}{l} 1. \text{ L is LINEAR and} \\ 2. \text{ L is TIME-INVARIANT} \end{array} \right\} \Rightarrow \left\{ \text{Le}^{st} = \underbrace{\check{h}^*(-s)}_{\text{eigenvalue}} \underbrace{e^{st}}_{\text{eigenvector}} \right\}$$

 PROOF:

$$\begin{aligned} [\text{Le}^{st}](s) &= \langle e^{su} | h((t; u), s) \rangle \\ &= \langle e^{su} | h((t - u), s) \rangle \\ &= \langle e^{s(t-u)} | h(v, s) \rangle \\ &= e^{st} \langle e^{-sv} | h(v, s) \rangle \\ &= \langle h(v, s) | e^{-sv} \rangle^* e^{st} \\ &= \langle h(v, s) | e^{(-s)v} \rangle^* e^{st} \\ &= \check{h}^*(-s) e^{st} \end{aligned}$$


by linear hypothesis

by time-invariance hypothesis

let  $v = t - u \Rightarrow u = t - v$ by additivity of  $\langle \Delta | \nabla \rangle$ by conjugate symmetry of  $\langle \Delta | \nabla \rangle$ by definition of  $\check{h}(s)$ 


# APPENDIX D

## TRIGONOMETRIC POLYNOMIALS



“I turn aside with a shudder of horror from this lamentable plague of functions which have no derivatives.”

Charles Hermite (1822 – 1901), French mathematician, in an 1893 letter to Stieltjes, in response to the “pathological” everywhere continuous but nowhere differentiable *Weierstrass functions*  $f(x) = \sum_{n=0}^{\infty} a^n \cos(b^n \pi x)$ .<sup>1</sup>

### D.1 Trigonometric expansion

**Theorem D.1** (DeMoivre's Theorem).

**T H M**  $(re^{ix})^n = r^n(\cos nx + i \sin nx) \quad \forall r, x \in \mathbb{R}$

PROOF:

$$\begin{aligned} (re^{ix})^n &= r^n e^{inx} \\ &= r^n (\cos nx + i \sin nx) \end{aligned} \quad \text{by Euler's identity (Theorem C.5 page 80)}$$



The cosine with argument  $nx$  can be expanded as a polynomial in  $\cos(x)$  (next).

**Theorem D.2** (trigonometric expansion).<sup>2</sup>

<sup>1</sup> quote: Hermite (1893)  
translation: Lakatos (1976) page 19  
image: <http://www-groups.dcs.sx-and.ac.uk/~history/PictDisplay/Hermite.html>  
<sup>2</sup> Rivlin (1974) page 3 (1.8)

T H M

$$\begin{aligned}\cos(nx) &= \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} \sum_{m=0}^k (-1)^{k+m} \binom{n}{2k} \binom{k}{m} (\cos x)^{n-2(k-m)} & \forall n \in \mathbb{W} \text{ and } x \in \mathbb{R} \\ \sin(nx) &= \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} \sum_{m=0}^k (-1)^{k+m} \binom{n}{2k} \binom{k}{m} (\sin x)^{n-2(k-m)} & \forall n \in \mathbb{W} \text{ and } x \in \mathbb{R}\end{aligned}$$

PROOF:

$$\begin{aligned}\cos(nx) &= \Re(\cos nx + i \sin nx) \\ &= \Re(e^{inx}) \\ &= \Re[(e^{ix})^n] \\ &= \Re[(\cos x + i \sin x)^n] \\ &= \Re \left[ \sum_{k \in \mathbb{Z}} \binom{n}{k} (\cos x)^{n-k} (i \sin x)^k \right] \\ &= \Re \left[ \sum_{k \in \mathbb{Z}} i^k \binom{n}{k} \cos^{n-k} x \sin^k x \right] \\ &= \Re \left[ \sum_{k \in \{0,4,\dots,n\}} \binom{n}{k} \cos^{n-k} x \sin^k x + i \sum_{k \in \{1,5,\dots,n\}} \binom{n}{k} \cos^{n-k} x \sin^k x \right. \\ &\quad \left. - \sum_{k \in \{2,6,\dots,n\}} \binom{n}{k} \cos^{n-k} x \sin^k x + -i \sum_{k \in \{3,7,\dots,n\}} \binom{n}{k} \cos^{n-k} x \sin^k x \right] \\ &= \sum_{k \in \{0,4,\dots,n\}} \binom{n}{k} \cos^{n-k} x \sin^k x - \sum_{k \in \{2,6,\dots,n\}} \binom{n}{k} \cos^{n-k} x \sin^k x \\ &= \sum_{k \in \{0,2,\dots,n\}} \binom{n}{k} (-1)^{\frac{k}{2}} \cos^{n-k} x \sin^k x \\ &= \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} \binom{n}{2k} (-1)^k \cos^{n-2k} x \sin^{2k} x \\ &= \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} \binom{n}{2k} (-1)^k \cos^{n-2k} x (1 - \cos^2 x)^k \\ &= \left[ \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} \binom{n}{2k} (-1)^k \cos^{n-2k} x \right] \left[ \sum_{m=0}^k \binom{k}{m} (-1)^m \cos^{2m} x \right] \\ &= \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} \sum_{m=0}^k (-1)^{k+m} \binom{n}{2k} \binom{k}{m} \cos^{n-2(k-m)} x\end{aligned}$$

$$\begin{aligned}\sin(nx) &= \cos\left(nx - \frac{\pi}{2}\right) \\ &= \cos\left(n \left[x - \frac{\pi}{2n}\right]\right) \\ &= \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} \sum_{m=0}^k (-1)^{k+m} \binom{n}{2k} \binom{k}{m} \cos^{n-2(k-m)} \left(x - \frac{\pi}{2n}\right)\end{aligned}$$

$$\begin{aligned}
&= \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} \sum_{m=0}^k (-1)^{k+m} \binom{n}{2k} \binom{k}{m} \cos^{n-2(k-m)} \left( nx - \frac{\pi}{2} \right) \\
&= \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} \sum_{m=0}^k (-1)^{k+m} \binom{n}{2k} \binom{k}{m} \sin^{n-2(k-m)} (nx)
\end{aligned}$$



Example D.1.

<b>E X</b>	$\cos 5x = 16\cos^5 x - 20\cos^3 x + 5\cos x$
	$\sin 5x = 16\sin^5 x - 20\sin^3 x + 5\sin x.$


PROOF:

1. Proof using *DeMoivre's Theorem* (Theorem D.1 page 91):

$$\begin{aligned}
&\cos 5x + i \sin 5x \\
&= e^{i5x} \\
&= (e^{ix})^5 \\
&= (\cos x + i \sin x)^5 \\
&= \sum_{k=0}^5 \binom{5}{k} [\cos x]^{5-k} [i \sin x]^k \\
&= \binom{5}{0} [\cos x]^{5-0} [i \sin x]^0 + \binom{5}{1} [\cos x]^{5-1} [i \sin x]^1 + \binom{5}{2} [\cos x]^{5-2} [i \sin x]^2 + \\
&\quad \binom{5}{3} [\cos x]^{5-3} [i \sin x]^3 + \binom{5}{4} [\cos x]^{5-4} [i \sin x]^4 + \binom{5}{5} [\cos x]^{5-5} [i \sin x]^5 \\
&= 1\cos^5 x + i5\cos^4 x \sin x - 10\cos^3 x \sin^2 x - i10\cos^2 x \sin^3 x + 5\cos x \sin^4 x + i1\sin^5 x \\
&= [\cos^5 x - 10\cos^3 x \sin^2 x + 5\cos x \sin^4 x] + i [5\cos^4 x \sin x - 10\cos^2 x \sin^3 x + \sin^5 x] \\
&= [\cos^5 x - 10\cos^3 x(1 - \cos^2 x) + 5\cos x(1 - \cos^2 x)(1 - \cos^2 x)] + \\
&\quad i [5(1 - \sin^2 x)(1 - \sin^2 x) \sin x - 10(1 - \sin^2 x) \sin^3 x + \sin^5 x] \\
&= [\cos^5 x - 10(\cos^3 x - \cos^5 x) + 5\cos x(1 - 2\cos^2 x + \cos^4 x)] + \\
&\quad i [5(1 - 2\sin^2 x + \sin^4 x) \sin x - 10(\sin^3 x - \sin^5 x) + \sin^5 x] \\
&= [\cos^5 x - 10(\cos^3 x - \cos^5 x) + 5(\cos x - 2\cos^3 x + \cos^5 x)] + \\
&\quad i [5(\sin x - 2\sin^3 x + \sin^5 x) - 10(\sin^3 x - \sin^5 x) + \sin^5 x] \\
&= \underbrace{[16\cos^5 x - 20\cos^3 x + 5\cos x]}_{\cos 5x} + i \underbrace{[16\sin^5 x - 20\sin^3 x + 5\sin x]}_{\sin 5x}
\end{aligned}$$

2. Proof using trigonometric expansion (Theorem D.2 page 91):

$$\begin{aligned}
\cos 5x &= \sum_{k=0}^{\lfloor \frac{5}{2} \rfloor} \sum_{m=0}^k (-1)^{k+m} \binom{5}{2k} \binom{k}{m} (\cos x)^{5-2(k-m)} \\
&= \sum_{k=0}^2 \sum_{m=0}^k (-1)^{k+m} \binom{5}{2k} \binom{k}{m} (\cos x)^{5-2(k-m)} \\
&= (-1)^0 \binom{5}{0} \binom{0}{0} \cos^5 x + (-1)^1 \binom{5}{2} \binom{1}{0} \cos^3 x + (-1)^2 \binom{5}{4} \binom{2}{1} \cos^5 x + \\
&\quad (-1)^2 \binom{5}{4} \binom{2}{0} \cos^1 x + (-1)^3 \binom{5}{6} \binom{3}{1} \cos^3 x + (-1)^4 \binom{5}{8} \binom{4}{2} \cos^5 x
\end{aligned}$$

Figure D.1: Lattice of harmonic cosines  $\{\cos(nx) | n = 0, 1, 2, \dots\}$ 

$$\begin{aligned}
 &= +(1)(1)\cos^5 x - (10)(1)\cos^3 x + (10)(1)\cos^5 x + (5)(1)\cos x - (5)(2)\cos^3 x + (5)(1)\cos^5 x \\
 &= +(1 + 10 + 5)\cos^5 x + (-10 - 10)\cos^3 x + 5\cos x \\
 &= 16\cos^5 x - 20\cos^3 x + 5\cos x
 \end{aligned}$$

⇒

Example D.2. <sup>3</sup>

E X	$n$	$\cos nx$	polynomial in $\cos x$	$n$	$\cos nx$	polynomial in $\cos x$
	0	$\cos 0x$	$= 1$	4	$\cos 4x$	$= 8\cos^4 x - 8\cos^2 x + 1$
	1	$\cos 1x$	$= \cos^1 x$	5	$\cos 5x$	$= 16\cos^5 x - 20\cos^3 x + 5\cos x$
	2	$\cos 2x$	$= 2\cos^2 x - 1$	6	$\cos 6x$	$= 32\cos^6 x - 48\cos^4 x + 18\cos^2 x - 1$
	3	$\cos 3x$	$= 4\cos^3 x - 3\cos x$	7	$\cos 7x$	$= 64\cos^7 x - 112\cos^5 x + 56\cos^3 x - 7\cos x$

PROOF:

$$\begin{aligned}
 \cos 2x &= \sum_{k=0}^{\lfloor \frac{2}{2} \rfloor} \sum_{m=0}^k (-1)^{k+m} \binom{3}{2k} \binom{k}{m} (\cos x)^{2-2(k-m)} \\
 &= (-1)^0 \binom{3}{0} \binom{0}{0} \cos^2 x + (-1)^1 \binom{3}{2} \binom{1}{0} \cos^0 x + (-1)^2 \binom{3}{2} \binom{1}{1} \cos^2 x \\
 &= +(1)(1)\cos^2 x - (1)(1) + (1)(1)\cos^2 x \\
 &= 2\cos^2 x - 1
 \end{aligned}$$

$$\begin{aligned}
 \cos 3x &= \sum_{k=0}^{\lfloor \frac{3}{2} \rfloor} \sum_{m=0}^k (-1)^{k+m} \binom{3}{2k} \binom{k}{m} (\cos x)^{3-2(k-m)} \\
 &= (-1)^0 \binom{3}{0} \binom{0}{0} \cos^3 x + (-1)^1 \binom{3}{2} \binom{1}{0} \cos^1 x + (-1)^2 \binom{3}{2} \binom{1}{1} \cos^3 x \\
 &= +\binom{3}{0} \binom{0}{0} \cos^3 x - \binom{3}{2} \binom{1}{0} \cos^1 x + \binom{3}{2} \binom{1}{1} \cos^3 x \\
 &= +(1)(1)\cos^3 x - (3)(1)\cos^1 x + (3)(1)\cos^3 x \\
 &= 4\cos^3 x - 3\cos x
 \end{aligned}$$

$$\cos 4x = \sum_{k=0}^{\lfloor \frac{4}{2} \rfloor} \sum_{m=0}^k (-1)^{k+m} \binom{4}{2k} \binom{k}{m} (\cos x)^{4-2(k-m)}$$

<sup>3</sup> [Abramowitz and Stegun \(1972\)](#) page 795, [Guillemin \(1957\)](#) page 593 (21), [Sloane \(2014\)](#) (<http://oeis.org/A039991>), [Sloane \(2014\)](#) (<http://oeis.org/A028297>)



$$\begin{aligned}
&= \sum_{k=0}^2 \sum_{m=0}^k (-1)^{k+m} \binom{4}{2k} \binom{k}{m} (\cos x)^{4-2(k-m)} \\
&= (-1)^{0+0} \binom{4}{2 \cdot 0} \binom{0}{0} (\cos x)^{4-2(0-0)} + (-1)^{1+0} \binom{4}{2 \cdot 1} \binom{1}{0} (\cos x)^{4-2(1-0)} \\
&\quad + (-1)^{1+1} \binom{4}{2 \cdot 1} \binom{1}{1} (\cos x)^{4-2(1-1)} + (-1)^{2+0} \binom{4}{2 \cdot 2} \binom{2}{0} (\cos x)^{4-2(2-0)} \\
&\quad + (-1)^{2+1} \binom{4}{2 \cdot 2} \binom{2}{1} (\cos x)^{4-2(2-1)} + (-1)^{2+2} \binom{4}{2 \cdot 2} \binom{2}{2} (\cos x)^{4-2(2-2)} \\
&= (1)(1)\cos^4 x - (6)(1)\cos^2 x + (6)(1)\cos^4 x + (1)(1)\cos^0 x - (1)(2)\cos^2 x + (1)(1)\cos^4 x \\
&= 8\cos^4 x - 8\cos^2 x + 1
\end{aligned}$$

$$\cos 5x = 16\cos^5 x - 20\cos^3 x + 5\cos x \quad \text{see Example D.1 page 93}$$

$$\begin{aligned}
\cos 6x &= \sum_{k=0}^{\lfloor \frac{6}{2} \rfloor} \sum_{m=0}^k (-1)^{k+m} \binom{6}{2k} \binom{k}{m} (\cos x)^{6-2(k-m)} \\
&= (-1)^0 \binom{6}{0} \binom{0}{0} \cos^6 x + (-1)^1 \binom{6}{2} \binom{1}{0} \cos^4 x + (-1)^2 \binom{6}{4} \binom{2}{0} \cos^2 x + \\
&\quad (-1)^3 \binom{6}{6} \binom{3}{0} \cos^0 x + (-1)^4 \binom{6}{8} \binom{4}{0} \cos^0 x + (-1)^5 \binom{6}{10} \binom{5}{0} \cos^0 x + (-1)^6 \binom{6}{12} \binom{6}{0} \cos^0 x \\
&\quad + (-1)^1 \binom{6}{2} \binom{1}{1} \cos^4 x + (-1)^2 \binom{6}{4} \binom{2}{1} \cos^2 x + (-1)^3 \binom{6}{6} \binom{3}{1} \cos^0 x + (-1)^4 \binom{6}{8} \binom{4}{1} \cos^0 x \\
&\quad + (-1)^5 \binom{6}{10} \binom{5}{1} \cos^0 x + (-1)^6 \binom{6}{12} \binom{6}{1} \cos^0 x \\
&= (1)(1)\cos^6 x - (15)(1)\cos^4 x + (15)(1)\cos^6 x + (15)(1)\cos^2 x - (15)(2)\cos^4 x + (15)(1)\cos^6 x \\
&\quad - (1)(1)\cos^0 x + (1)(3)\cos^2 x - (1)(3)\cos^4 x + (1)(1)\cos^6 x \\
&= 32\cos^6 x - 48\cos^4 x + 18\cos^2 x - 1
\end{aligned}$$

$$\begin{aligned}
\cos 7x &= \sum_{k=0}^{\lfloor \frac{7}{2} \rfloor} \sum_{m=0}^k (-1)^{k+m} \binom{7}{2k} \binom{k}{m} (\cos x)^{7-2(k-m)} \\
&= \sum_{k=0}^3 \sum_{m=0}^k (-1)^{k+m} \binom{7}{2k} \binom{k}{m} (\cos x)^{7-2(k-m)} \\
&= (-1)^0 \binom{7}{0} \binom{0}{0} \cos^7 x + (-1)^1 \binom{7}{2} \binom{1}{0} \cos^5 x + (-1)^2 \binom{7}{4} \binom{2}{0} \cos^3 x \\
&\quad + (-1)^3 \binom{7}{6} \binom{3}{0} \cos^1 x + (-1)^4 \binom{7}{8} \binom{4}{0} \cos^0 x + (-1)^5 \binom{7}{10} \binom{5}{0} \cos^0 x + (-1)^6 \binom{7}{12} \binom{6}{0} \cos^0 x \\
&\quad + (-1)^1 \binom{7}{2} \binom{1}{1} \cos^5 x + (-1)^2 \binom{7}{4} \binom{2}{1} \cos^3 x + (-1)^3 \binom{7}{6} \binom{3}{1} \cos^1 x + (-1)^4 \binom{7}{8} \binom{4}{1} \cos^0 x \\
&\quad + (-1)^5 \binom{7}{10} \binom{5}{1} \cos^0 x + (-1)^6 \binom{7}{12} \binom{6}{1} \cos^0 x \\
&= (1)(1)\cos^7 x - (21)(1)\cos^5 x + (21)(1)\cos^7 x + (35)(1)\cos^3 x \\
&\quad - (35)(2)\cos^5 x + (35)(1)\cos^7 x - (7)(1)\cos^1 x + (7)(3)\cos^3 x \\
&\quad - (7)(3)\cos^5 x + (7)(1)\cos^7 x \\
&= (1 + 21 + 35 + 7)\cos^7 x - (21 + 70 + 21)\cos^5 x + (35 + 21)\cos^3 x - (7)\cos^1 x \\
&= 64\cos^7 x - 112\cos^5 x + 56\cos^3 x - 7\cos x
\end{aligned}$$




Figure D.2: Lattice of Chebyshev polynomials  $\{T_n(x) | n = 0, 1, 2, 3\}$

Note: Trigonometric expansion of  $\cos(nx)$  for particular values of  $n$  can also be performed with the free software package *Maxima*<sup>TM</sup> using the syntax illustrated to the right:<sup>4</sup>

```
1 trigexpand(cos(2*x));
2 trigexpand(cos(3*x));
3 trigexpand(cos(4*x));
4 trigexpand(cos(5*x));
5 trigexpand(cos(6*x));
6 trigexpand(cos(7*x));
```

### Definition D.1.

**DEF** The  $n$ th Chebyshev polynomial of the first kind is defined as

$$T_n(x) \triangleq \cos nx \quad \text{where} \quad \cos x \triangleq x$$

**Theorem D.3.**<sup>5</sup> Let  $T_n(x)$  be a CHEBYSHEV POLYNOMIAL with  $n \in \mathbb{W}$ .

**THM**  $n$  is EVEN  $\implies T_n(x)$  is EVEN.  
 $n$  is ODD  $\implies T_n(x)$  is ODD.

**Example D.3.** Let  $T_n(x)$  be a Chebyshev polynomial with  $n \in \mathbb{W}$ .

$T_0(x) = 1$	$T_4(x) = 8x^4 - 8x^2 + 1$
$T_1(x) = x$	$T_5(x) = 16x^5 - 20x^3 + 5x$
$T_2(x) = 2x^2 - 1$	$T_6(x) = 32x^6 - 48x^4 + 18x^2 - 1$
$T_3(x) = 4x^3 - 3x$	

**PROOF:** Proof of these equations follows directly from Example D.2 (page 94).


## D.2 Trigonometric reduction

Theorem D.2 (page 91) showed that  $\cos nx$  can be expressed as a polynomial in  $\cos x$ . Conversely, Theorem D.4 (next) shows that a polynomial in  $\cos x$  can be expressed as a linear combination of  $(\cos nx)_{n \in \mathbb{Z}}$ .

**Theorem D.4** (trigonometric reduction).

<sup>4</sup> [maxima](#) pages 157–158 (10.5 Trigonometric Functions)

<sup>5</sup> [Rivlin \(1974\) page 5](#) (1.13), [Süli and Mayers \(2003\) page 242](#) (Lemma 8.2), [Davidson and Donsig \(2010\) page 222](#) (exercise 10.7.A(a))

Figure D.3: Lattice of exponential cosines  $\{\cos^n x | n = 0, 1, 2, 3\}$ 

T H M

$$\begin{aligned} \cos^n x &= \frac{1}{2^n} \sum_{k=0}^n \binom{n}{k} \cos[(n-2k)x] \\ &= \begin{cases} \frac{1}{2^n} \binom{n}{\frac{n}{2}} + \frac{1}{2^{n-1}} \sum_{k=0}^{\frac{n}{2}-1} \binom{n}{k} \cos[(n-2k)x] & \text{for } n \text{ even} \\ \frac{1}{2^{n-1}} \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} \binom{n}{k} \cos[(n-2k)x] & \text{for } n \text{ odd} \end{cases} \end{aligned}$$

PROOF:

$$\begin{aligned} \cos^n x &= \left( \frac{e^{ix} + e^{-ix}}{2} \right)^n \\ &= \mathbf{R}_e \left[ \left( \frac{e^{ix} + e^{-ix}}{2} \right)^n \right] \\ &= \mathbf{R}_e \left[ \frac{1}{2^n} \sum_{k=0}^n \binom{n}{k} e^{i(n-k)x} e^{-ikx} \right] \\ &= \mathbf{R}_e \left[ \frac{1}{2^n} \sum_{k=0}^n \binom{n}{k} e^{i(n-2k)x} \right] \\ &= \mathbf{R}_e \left[ \frac{1}{2^n} \sum_{k=0}^n \binom{n}{k} (\cos[(n-2k)x] + i \sin[(n-2k)x]) \right] \\ &= \mathbf{R}_e \left[ \frac{1}{2^n} \sum_{k=0}^n \binom{n}{k} \cos[(n-2k)x] + i \frac{1}{2^n} \sum_{k=0}^n \binom{n}{k} \sin[(n-2k)x] \right] \\ &= \frac{1}{2^n} \sum_{k=0}^n \binom{n}{k} \cos[(n-2k)x] \\ &= \begin{cases} \frac{1}{2^n} \binom{n}{\frac{n}{2}} + \frac{1}{2^{n-1}} \sum_{k=0}^{\frac{n}{2}-1} \binom{n}{k} \cos[(n-2k)x] & : \quad n \text{ even} \\ \frac{1}{2^{n-1}} \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} \binom{n}{k} \cos[(n-2k)x] & : \quad n \text{ odd} \end{cases} \end{aligned}$$

Example D.4. <sup>6</sup>

<sup>6</sup> Abramowitz and Stegun (1972) page 795, Sloane (2014) (<http://oeis.org/A100257>), Sloane (2014) (<http://oeis.org/A008314>)

E X	$n$	$\cos^n x$	trigonometric reduction	$n$	$\cos^n x$	trigonometric reduction
	0	$\cos^0 x$	$= 1$	4	$\cos^4 x$	$= \frac{\cos 4x + 4\cos 2x + 3}{2^3}$
	1	$\cos^1 x$	$= \cos x$	5	$\cos^5 x$	$= \frac{\cos 5x + 5\cos 3x + 10\cos x}{2^4}$
	2	$\cos^2 x$	$= \frac{\cos 2x + 1}{2}$	6	$\cos^6 x$	$= \frac{\cos 6x + 6\cos 4x + 15\cos 2x + 10}{2^5}$
	3	$\cos^3 x$	$= \frac{\cos 3x + 3\cos x}{2^2}$	7	$\cos^7 x$	$= \frac{\cos 7x + 7\cos 5x + 21\cos 3x + 35\cos x}{2^6}$

PROOF:

$$\begin{aligned}
 \cos^0 x &= \frac{1}{2^n} \sum_{k=0}^n \binom{n}{k} \cos([n - 2k]x) \Big|_{n=0} \\
 &= \frac{1}{2^0} \sum_{k=0}^0 \binom{0}{k} \cos[(0 - 2k)x] \\
 &= \binom{0}{0} \cos[(0 - 2 \cdot 0)x] \\
 &= 1
 \end{aligned}$$

$$\begin{aligned}
 \cos^1 x &= \frac{1}{2^n} \sum_{k=0}^n \binom{n}{k} \cos([n - 2k]x) \Big|_{n=1} \\
 &= \frac{1}{2^1} \sum_{k=0}^1 \binom{1}{k} \cos[(1 - 2k)x] \\
 &= \frac{1}{2} \left[ \binom{1}{0} \cos[(1 - 2 \cdot 0)x] + \binom{1}{1} \cos[(1 - 2 \cdot 1)x] \right] \\
 &= \frac{1}{2} [1\cos x + 1\cos(-x)] \\
 &= \frac{1}{2} (\cos x + \cos x) \\
 &= \cos x
 \end{aligned}$$

$$\begin{aligned}
 \cos^2 x &= \frac{1}{2^n} \sum_{k=0}^n \binom{n}{k} \cos([n - 2k]x) \Big|_{n=2} \\
 &= \frac{1}{2^2} \sum_{k=0}^2 \binom{2}{k} \cos([2 - 2k]x) \\
 &= \frac{1}{2^2} \left[ \binom{2}{0} \cos([2 - 2 \cdot 0]x) + \binom{2}{1} \cos([2 - 2 \cdot 1]x) + \binom{2}{2} \cos([2 - 2 \cdot 2]x) \right] \\
 &= \frac{1}{2^2} [1\cos(2x) + 2\cos(0x) + 1\cos(-2x)] \\
 &= \frac{1}{2^2} [\cos(2x) + 2 + \cos(2x)] \\
 &= \frac{1}{2} [\cos(2x) + 1]
 \end{aligned}$$

$$\begin{aligned}
 \cos^3 x &= \frac{1}{2^n} \sum_{k=0}^n \binom{n}{k} \cos([n - 2k]x) \Big|_{n=3} \\
 &= \frac{1}{2^3} \sum_{k=0}^3 \binom{3}{k} \cos([3 - 2k]x)
 \end{aligned}$$

$$\begin{aligned}
&= \frac{1}{2^3} [1\cos(3x) + 3\cos(1x) + 3\cos(-1x) + 1\cos(-3x)] \\
&= \frac{1}{2^3} [\cos(3x) + 3\cos(x) + 3\cos(x) + \cos(3x)] \\
&= \frac{1}{2^2} [\cos(3x) + 3\cos(x)] \\
\cos^4 x &= \frac{1}{2^n} \sum_{k=0}^n \binom{n}{k} \cos([n-2k]x) \Big|_{n=4} \\
&= \frac{1}{2^4} \sum_{k=0}^4 \binom{4}{k} \cos([4-2k]x) \\
&= \frac{1}{2^4} [1\cos(4x) + 4\cos(2x) + 6\cos(0x) + 4\cos(-2x) + 1\cos(-4x)] \\
&= \frac{1}{2^3} [\cos(4x) + 4\cos(2x) + 3] \\
\cos^5 x &= \frac{1}{2^{5-1}} \sum_{k=0}^{\lfloor \frac{5}{2} \rfloor} \binom{5}{k} \cos[(5-2k)x] \\
&= \frac{1}{16} \sum_{k=0}^2 \binom{5}{k} \cos[(5-2k)x] \\
&= \frac{1}{16} \left[ \binom{5}{0} \cos 5x + \binom{5}{1} \cos 3x + \binom{5}{2} \cos x \right] \\
&= \frac{1}{16} [\cos 5x + 5\cos 3x + 10\cos x] \\
\cos^6 x &= \frac{1}{2^6} \binom{6}{\frac{6}{2}} + \frac{1}{2^{6-1}} \sum_{k=0}^{\frac{6}{2}-1} \binom{6}{k} \cos[(6-2k)x] \\
&= \frac{1}{2^6} \binom{6}{3} + \frac{1}{2^5} \sum_{k=0}^2 \binom{6}{k} \cos[(6-2k)x] \\
&= \frac{1}{64} 20 + \frac{1}{32} \left[ \binom{6}{0} \cos 6x + \binom{6}{1} \cos 4x + \binom{6}{2} \cos 2x \right] \\
&= \frac{1}{32} [\cos 6x + 6\cos 4x + 15\cos 2x + 10] \\
\cos^7 x &= \frac{1}{2^{7-1}} \sum_{k=0}^{\lfloor \frac{7}{2} \rfloor} \binom{7}{k} \cos[(7-2k)x] \\
&= \frac{1}{64} \sum_{k=0}^2 \binom{7}{k} \cos[(7-2k)x] \\
&= \frac{1}{64} \left[ \binom{7}{0} \cos 7x + \binom{7}{1} \cos 5x + \binom{7}{2} \cos 3x + \binom{7}{3} \cos x \right] \\
&= \frac{1}{64} [\cos 7x + 7\cos 5x + 21\cos 3x + 35\cos x]
\end{aligned}$$

Note: Trigonometric reduction of  $\cos^n(x)$  for particular values of  $n$  can also be performed with the free software package *Maxima*<sup>TM</sup> using the syntax illustrated to the right:<sup>7</sup>

```

1 trigreduce((cos(x))^2);
2 trigreduce((cos(x))^3);
3 trigreduce((cos(x))^4);
4 trigreduce((cos(x))^5);
5 trigreduce((cos(x))^6);
6 trigreduce((cos(x))^7);

```

<sup>7</sup> [http://maxima.sourceforge.net/docs/manual/en/maxima\\_15.html](http://maxima.sourceforge.net/docs/manual/en/maxima_15.html)  
 // [maxima](https://github.com/dgreenhoe/pdfs/blob/master/abctran.pdf) page 158 (10.5 Trigonometric Functions)



## D.3 Spectral Factorization

**Theorem D.5** (Fejér-Riesz spectral factorization).<sup>8</sup> Let  $[0, \infty) \not\subset \mathbb{R}$  and

$$p(e^{ix}) \triangleq \sum_{n=-N}^N a_n e^{inx} \quad (\text{Laurent trigonometric polynomial order } 2N)$$

$$q(e^{ix}) \triangleq \sum_{n=1}^N b_n e^{inx} \quad (\text{standard trigonometric polynomial order } N)$$

<b>T H M</b>	$p(e^{ix}) \in [0, \infty) \quad \forall x \in [0, 2\pi] \quad \implies \quad \left\{ \begin{array}{l} \exists (b_n)_{n \in \mathbb{Z}} \text{ such that} \\ p(e^{ix}) = q(e^{ix}) q^*(e^{ix}) \end{array} \right. \quad \forall x \in \mathbb{R}$
----------------------	--

PROOF:

1. Proof that  $a_n = a_{-n}^*$  ( $(a_n)_{n \in \mathbb{Z}}$  is *Hermitian symmetric*):


Let  $a_n \triangleq r_n e^{i\phi_n}$ ,  $r_n, \phi_n \in \mathbb{R}$ . Then

$$\begin{aligned}
 p(e^{inx}) &\triangleq \sum_{n=-N}^N a_n e^{inx} \\
 &= \sum_{n=-N}^N r_n e^{i\phi_n} e^{inx} \\
 &= \sum_{n=-N}^N r_n e^{inx + \phi_n} \\
 &= \sum_{n=-N}^N r_n \cos(nx + \phi_n) + i \sum_{n=-N}^N r_n \sin(nx + \phi_n) \\
 &= \sum_{n=-N}^N r_n \cos(nx + \phi_n) + i \underbrace{\left[ r_0 \sin(0x + \phi_0) + \sum_{n=1}^N r_n \sin(nx + \phi_n) + \sum_{n=1}^N r_{-n} \sin(-nx + \phi_{-n}) \right]}_{\text{imaginary part must equal 0 because } p(x) \in \mathbb{R}} \\
 &= \sum_{n=-N}^N r_n \cos(nx + \phi_n) + i \underbrace{\left[ r_0 \sin(\phi_0) + \sum_{n=1}^N r_n \sin(nx + \phi_n) - \sum_{n=1}^N r_{-n} \sin(nx - \phi_{-n}) \right]}_{\implies r_n = r_{-n}, \phi_n = -\phi_{-n} \implies a_n = a_{-n}^*, a_0 \in \mathbb{R}}
 \end{aligned}$$

2. Because the coefficients  $(c_n)_{n \in \mathbb{Z}}$  are *Hermitian symmetric*, the zeros of  $P(z)$  occur in *conjugate reciprocal pairs*. This means that if  $\sigma \in \mathbb{C}$  is a zero of  $P(z)$  ( $P(\sigma) = 0$ ), then  $\frac{1}{\sigma^*}$  is also a zero of  $P(z)$  ( $P\left(\frac{1}{\sigma^*}\right) = 0$ ). In the complex  $z$  plane, this relationship means zeros are reflected across the unit circle such that

$$\frac{1}{\sigma^*} = \frac{1}{(re^{i\phi})^*} = \frac{1}{r} \frac{1}{e^{-i\phi}} = \frac{1}{r} e^{i\phi}$$

<sup>8</sup> Pinsky (2002) pages 330–331



3. Because the zeros of  $p(z)$  occur in conjugate reciprocal pairs,  $p(e^{ix})$  can be factored:

$$\begin{aligned}
 p(e^{ix}) &= p(z)|_{z=e^{ix}} \\
 &= z^{-N} C \prod_{n=1}^N (z - \sigma_n) \prod_{n=1}^N \left( z - \frac{1}{\sigma_n^*} \right) \Big|_{z=e^{ix}} \\
 &= C \prod_{n=1}^N (z - \sigma_n) \prod_{n=1}^N z^{-1} \left( z - \frac{1}{\sigma_n^*} \right) \Big|_{z=e^{ix}} \\
 &= C \prod_{n=1}^N (z - \sigma_n) \prod_{n=1}^N \left( 1 - \frac{1}{\sigma_n^*} z^{-1} \right) \Big|_{z=e^{ix}} \\
 &= C \prod_{n=1}^N (z - \sigma_n) \prod_{n=1}^N (z^{-1} - \sigma_n^*) \left( -\frac{1}{\sigma_n^*} \right) \Big|_{z=e^{ix}} \\
 &= \left[ C \prod_{n=1}^N \left( -\frac{1}{\sigma_n^*} \right) \right] \left[ \prod_{n=1}^N (z - \sigma_n) \right] \left[ \prod_{n=1}^N \left( \frac{1}{z^*} - \sigma_n \right) \right]^* \Big|_{z=e^{ix}} \\
 &= \left[ C_2 \prod_{n=1}^N (z - \sigma_n) \right] \left[ C_2 \prod_{n=1}^N \left( \frac{1}{z^*} - \sigma_n \right) \right]^* \Big|_{z=e^{ix}} \\
 &= q(z) q^* \left( \frac{1}{z^*} \right) \Big|_{z=e^{ix}} \\
 &= q(e^{ix}) q^*(e^{ix})
 \end{aligned}$$

⇒

## D.4 Dirichlet Kernel



“Dirichlet alone, not I, nor Cauchy, nor Gauss knows what a completely rigorous proof is. Rather we learn it first from him. When Gauss says he has proved something it is clear; when Cauchy says it, one can wager as much pro as con; when Dirichlet says it, it is certain.”

Carl Gustav Jacob Jacobi (1804–1851), Jewish-German mathematician <sup>9</sup>

<sup>9</sup> quote: Schubring (2005) page 558

image: [http://en.wikipedia.org/wiki/File:Carl\\_Jacobi.jpg](http://en.wikipedia.org/wiki/File:Carl_Jacobi.jpg), public domain

The *Dirichlet Kernel* is critical in proving what is not immediately obvious in examining the Fourier Series—that for a broad class of periodic functions, a function can be recovered from (with uniform convergence) its Fourier Series analysis.

**Definition D.2.** <sup>10</sup>

DEF

The *Dirichlet Kernel*  $D_n \in \mathbb{R}^{\mathbb{W}}$  with period  $\tau$  is defined as

$$D_n(x) \triangleq \frac{1}{\tau} \sum_{k=-n}^n e^{i \frac{2\pi}{\tau} kx}$$

**Proposition D.1.** <sup>11</sup> Let  $D_n$  be the DIRICHLET KERNEL with period  $\tau$  (Definition D.2 page 102).

PRP

$$D_n(x) = \frac{1}{\tau} \frac{\sin\left(\frac{\pi}{\tau}[2n+1]x\right)}{\sin\left(\frac{\pi}{\tau}x\right)}$$

PROOF:

$$\begin{aligned} D_n(x) &\triangleq \frac{1}{\tau} \sum_{k=-n}^n e^{i \frac{2\pi}{\tau} kx} && \text{by definition of } D_n && (\text{Definition D.2 page 102}) \\ &= \frac{1}{\tau} \sum_{k=0}^{2n} e^{i \frac{2\pi}{\tau} (k-n)x} = \frac{1}{\tau} e^{-i \frac{2\pi n}{\tau} x} \sum_{k=0}^{2n} e^{i \frac{2\pi}{\tau} kx} = \frac{1}{\tau} e^{-i \frac{2\pi n}{\tau} x} \sum_{k=0}^{2n} \left(e^{i \frac{2\pi}{\tau} x}\right)^k \\ &= \frac{1}{\tau} e^{-i \frac{2\pi n}{\tau} x} \frac{1 - \left(e^{i \frac{2\pi}{\tau} x}\right)^{2n+1}}{1 - e^{i \frac{2\pi}{\tau} x}} && \text{by geometric series} \\ &= \frac{1}{\tau} e^{-i \frac{2\pi n}{\tau} x} \frac{1 - e^{i \frac{2\pi}{\tau} (2n+1)x}}{1 - e^{i \frac{2\pi}{\tau} x}} = \frac{1}{\tau} e^{-i \frac{2\pi n}{\tau} x} \left(\frac{e^{i \frac{\pi}{\tau} (2n+1)x}}{e^{i \frac{\pi}{\tau} x}}\right) \frac{e^{-i \frac{\pi}{\tau} (2n+1)x} - e^{i \frac{\pi}{\tau} (2n+1)x}}{e^{-i \frac{\pi}{\tau} x} - e^{i \frac{\pi}{\tau} x}} \\ &= \frac{1}{\tau} e^{-i \frac{2\pi n}{\tau} x} \left(e^{i \frac{2\pi n}{\tau} x}\right) \frac{-2i \sin\left[\frac{\pi}{\tau} (2n+1)x\right]}{-2i \sin\left[\frac{\pi}{\tau} x\right]} = \frac{1}{\tau} \frac{\sin\left[\frac{\pi}{\tau} (2n+1)x\right]}{\sin\left[\frac{\pi}{\tau} x\right]} \end{aligned}$$

⇒

**Proposition D.2.** <sup>12</sup> Let  $D_n$  be the DIRICHLET KERNEL with period  $\tau$  (Definition D.2 page 102).

PRP

$$\int_0^{\tau} D_n(x) dx = 1$$

PROOF:

$$\begin{aligned} \int_0^{\tau} D_n(x) dx &\triangleq \int_0^{\tau} \frac{1}{\tau} \sum_{k=-n}^n e^{i \frac{2\pi}{\tau} kx} dx && \text{by definition of } D_n \text{ (Definition D.2 page 102)} \\ &= \frac{1}{\tau} \sum_{k=-n}^n \int_{-\frac{\tau}{2}}^{\frac{\tau}{2}} e^{i \frac{2\pi}{\tau} kx} dx \\ &= \frac{1}{\tau} \sum_{k=-n}^n \int_{-\frac{\tau}{2}}^{\frac{\tau}{2}} \cos\left(\frac{2\pi}{\tau} kx\right) + i \sin\left(\frac{2\pi}{\tau} kx\right) dx \end{aligned}$$

<sup>10</sup> Katznelson (2004) page 14, Heil (2011) pages 443–444, Folland (1992) pages 33–34

<sup>11</sup> Katznelson (2004) page 14, Heil (2011) page 444, Folland (1992) page 34

<sup>12</sup> Bruckner et al. (1997) pages 620–621






Figure D.4:  $D_n$  function for  $N = 1, 5, 10, 20$ .  $D_n \rightarrow \text{comb.}$  (See Proposition D.1 page 102).

$$\begin{aligned}
 &= \frac{1}{\tau} \sum_{k=-n}^n \int_{-\frac{\tau}{2}}^{\frac{\tau}{2}} \cos\left(\frac{2\pi}{\tau} kx\right) dx \\
 &= \frac{1}{\tau} \sum_{k=-n}^n \left. \frac{\sin\left(\frac{2\pi}{\tau} kx\right)}{\frac{2\pi}{\tau} k} \right|_{-\frac{\tau}{2}}^{\frac{\tau}{2}} \\
 &= \frac{1}{\tau} \sum_{k=-n}^n \left[ \frac{\sin\left(\frac{2\pi}{\tau} k \frac{\tau}{2}\right)}{\frac{2\pi}{\tau} k} - \frac{\sin\left(-\frac{2\pi}{\tau} k \frac{\tau}{2}\right)}{\frac{2\pi}{\tau} k} \right] \\
 &= \frac{1}{\tau} \sum_{k=-n}^n \left[ \frac{\sin(\pi k)}{\pi k} + \frac{\sin(\pi k)}{\pi k} \right] \\
 &= \frac{1}{2} \left[ 2 \frac{\sin(\pi n)}{\pi n} \right]_{k=0} \\
 &= 1
 \end{aligned}$$

⇒

**Proposition D.3.** Let  $D_n$  be the DIRICHLET KERNEL with period  $\tau$  (Definition D.2 page 102). Let  $w_N$  (the “width” of  $D_n(x)$ ) be the distance between the two points where the center pulse of  $D_n(x)$  intersects the  $x$  axis.

P R P	$D_n(0) = \frac{1}{\tau}(2n+1)$
	$w_n = \frac{2\tau}{2n+1}$

 PROOF:

$$\begin{aligned}
 D_n(0) &= D_n(x) \Big|_{t=0} \\
 &= \frac{1}{\tau} \frac{\sin \left[ \frac{\pi}{\tau} (2n+1)x \right]}{\sin \left[ \frac{\pi}{\tau} t \right]} \Big|_{t=0} && \text{by Proposition D.1 page 102} \\
 &= \frac{1}{\tau} \frac{\frac{d}{dx} \sin \left[ \frac{\pi}{\tau} (2n+1)x \right]}{\frac{d}{dx} \sin \left[ \frac{\pi}{\tau} t \right]} \Big|_{t=0} && \text{by l'Hôpital's rule} \\
 &= \frac{1}{\tau} \frac{\frac{\pi}{\tau} (2n+1) \cos \left[ \frac{\pi}{\tau} (2n+1)x \right]}{\cos \left[ \frac{\pi}{\tau} t \right]} \Big|_{t=0} \\
 &= \frac{1}{\tau} \frac{\frac{\pi}{\tau} (2n+1)}{\frac{\pi}{\tau}} \frac{1}{1} \\
 &= \frac{1}{\tau} (2n+1)
 \end{aligned}$$

The center pulse of kernel  $D_n(x)$  intersects the  $x$  axis at

$$t = \pm \frac{\tau}{(2n+1)}$$

which implies

$$w_n = \frac{\tau}{2n+1} + \frac{\tau}{2n+1} = \frac{2\tau}{(2n+1)}.$$




**Proposition D.4.** <sup>13</sup> Let  $D_n$  be the DIRICHLET KERNEL with period  $\tau$  (Definition D.2 page 102).

P R P	$D_n(x) = D_n(-x) \quad (D_n \text{ is an EVEN function})$
-------------	--

 PROOF:

$$\begin{aligned}
 D_n(x) &= \frac{1}{\tau} \frac{\sin \left[ \frac{\pi}{\tau} (2n+1)x \right]}{\sin \left[ \frac{\pi}{\tau} t \right]} && \text{by Proposition D.1 page 102} \\
 &= \frac{1}{\tau} \frac{-\sin \left[ -\frac{\pi}{\tau} (2n+1)x \right]}{-\sin \left[ -\frac{\pi}{\tau} t \right]} && \text{because } \sin x \text{ is an } \textit{odd} \text{ function} \\
 &= \frac{1}{\tau} \frac{\sin \left[ \frac{\pi}{\tau} (2n+1)(-x) \right]}{\sin \left[ \frac{\pi}{\tau} (-x) \right]} \\
 &= D_n(-x) && \text{by Proposition D.1 page 102}
 \end{aligned}$$



<sup>13</sup>  Bruckner et al. (1997) pages 620–621

## D.5 Trigonometric summations

**Theorem D.6** (Lagrange trigonometric identities). <sup>14</sup>

T H M	$\sum_{n=0}^{N-1} \cos(nx) = \frac{1}{2} + \frac{\sin\left(\left[N - \frac{1}{2}\right]x\right)}{2\sin\left(\frac{1}{2}x\right)} = \frac{\sin\left(\left[N - \frac{1}{2}\right]x\right) + \sin\left(\frac{1}{2}x\right)}{2\sin\left(\frac{1}{2}x\right)} \quad \forall x \in \mathbb{R}$
	$\sum_{n=0}^{N-1} \sin(nx) = \frac{1}{2} \cot\left(\frac{1}{2}x\right) + \frac{\cos\left(\left[N - \frac{1}{2}\right]x\right)}{2\sin\left(\frac{1}{2}x\right)} = \frac{\cos\left(\left[N - \frac{1}{2}\right]x\right) + \cos\left(\frac{1}{2}x\right)}{2\sin\left(\frac{1}{2}x\right)} \quad \forall x \in \mathbb{R}$

✎ PROOF:

$$\begin{aligned}
 \sum_{n=0}^{N-1} \cos(nx) &= \sum_{n=0}^{N-1} \Re e^{inx} = \Re \sum_{n=0}^{N-1} e^{inx} = \Re \sum_{n=0}^{N-1} (e^{ix})^n \\
 &= \Re \left[ \frac{1 - e^{iNx}}{1 - e^{ix}} \right] && \text{by geometric series} \\
 &= \Re \left[ \left( \frac{e^{i\frac{1}{2}Nx}}{e^{i\frac{1}{2}x}} \right) \left( \frac{e^{-i\frac{1}{2}Nx} - e^{i\frac{1}{2}Nx}}{e^{-i\frac{1}{2}x} - e^{i\frac{1}{2}x}} \right) \right] \\
 &= \Re \left[ \left( e^{i\frac{1}{2}(N-1)x} \right) \left( \frac{-i\frac{1}{2}\sin\left(\frac{1}{2}Nx\right)}{-i\frac{1}{2}\sin\left(\frac{1}{2}x\right)} \right) \right] \\
 &= \cos\left(\frac{1}{2}(N-1)x\right) \left( \frac{\sin\left(\frac{1}{2}Nx\right)}{\sin\left(\frac{1}{2}x\right)} \right) \\
 &= \frac{-\frac{1}{2}\sin\left(-\frac{1}{2}x\right) + \frac{1}{2}\sin\left(\left[N - \frac{1}{2}\right]x\right)}{\sin\left(\frac{1}{2}x\right)} && \text{by product identities} \quad (\text{Theorem C.8 page 82}) \\
 &= \frac{1}{2} + \frac{\sin\left(\left[N - \frac{1}{2}\right]x\right)}{2\sin\left(\frac{1}{2}x\right)}
 \end{aligned}$$

$$\begin{aligned}
 \sum_{n=0}^{N-1} \sin(nx) &= \sum_{n=0}^{N-1} \Im e^{inx} = \Im \sum_{n=0}^{N-1} e^{inx} = \Im \sum_{n=0}^{N-1} (e^{ix})^n \\
 &= \Im \left[ \frac{1 - e^{iNx}}{1 - e^{ix}} \right] && \text{by geometric series} \\
 &= \Im \left[ \left( \frac{e^{i\frac{1}{2}Nx}}{e^{i\frac{1}{2}x}} \right) \left( \frac{e^{-i\frac{1}{2}Nx} - e^{i\frac{1}{2}Nx}}{e^{-i\frac{1}{2}x} - e^{i\frac{1}{2}x}} \right) \right] \\
 &= \Im \left[ \left( e^{i\frac{1}{2}(N-1)x} \right) \left( \frac{-\frac{1}{2}i\sin\left(\frac{1}{2}Nx\right)}{-\frac{1}{2}i\sin\left(\frac{1}{2}x\right)} \right) \right]
 \end{aligned}$$

<sup>14</sup> [Muniz \(1953\)](#) page 140 (“Lagrange's Trigonometric Identities”), [Jeffrey and Dai \(2008\)](#) pages 128–130 (2.4.1.6 Sines, Cosines, and Tagents of Multiple Angles; (14), (13))

$$\begin{aligned}
&= \sin\left(\frac{(N-1)x}{2}\right) \left( \frac{\sin\left(\frac{1}{2}Nx\right)}{\sin\left(\frac{1}{2}x\right)} \right) \\
&= \frac{\frac{1}{2}\cos\left(-\frac{1}{2}x\right) - \frac{1}{2}\cos\left(\left[N - \frac{1}{2}\right]x\right)}{\sin\left(\frac{1}{2}x\right)} && \text{by product identities} \quad (\text{Theorem C.8 page 82}) \\
&= \frac{1}{2} \cot\left(\frac{1}{2}x\right) + \frac{\cos\left(\left[N - \frac{1}{2}\right]x\right)}{2\sin\left(\frac{1}{2}x\right)}
\end{aligned}$$

Note that these results (summed with indices from  $n = 0$  to  $n = N - 1$ ) are compatible with [Muniz \(1953\)](#) page 140 (summed with indices from  $n = 1$  to  $n = N$ ) as demonstrated next:

$$\begin{aligned}
\sum_{n=0}^{N-1} \cos(nx) &= \sum_{n=1}^N \cos(nx) + [\cos(0x) - \cos(Nx)] \\
&= \left[ -\frac{1}{2} + \frac{\sin\left(\left[N + \frac{1}{2}\right]x\right)}{2\sin\left(\frac{1}{2}x\right)} \right] + [\cos(0x) - \cos(Nx)] && \text{by } \text{Muniz (1953) page 140} \\
&= \left(1 - \frac{1}{2}\right) + \frac{\sin\left(\left[N + \frac{1}{2}\right]x\right) - 2\sin\left(\frac{1}{2}x\right)\cos(Nx)}{2\sin\left(\frac{1}{2}x\right)} \\
&= \frac{1}{2} + \frac{\sin\left(\left[N + \frac{1}{2}\right]x\right) - 2\left[\sin\left(\left[\frac{1}{2} - N\right]x\right) + \sin\left(\left[\frac{1}{2} + N\right]x\right)\right]}{2\sin\left(\frac{1}{2}x\right)} && \text{by Theorem C.8 page 82} \\
&= \frac{1}{2} + \frac{\sin\left(\frac{1}{2}[2N-1]x\right)}{2\sin\left(\frac{1}{2}x\right)} && \Rightarrow \text{above result}
\end{aligned}$$

$$\begin{aligned}
\sum_{n=0}^{N-1} \sin(nx) &= \sum_{n=1}^N \sin(nx) + [\sin(0x) - \sin(Nx)] \\
&= \frac{1}{2} \cot\left(\frac{1}{2}x\right) - \frac{\cos\left(\left[N + \frac{1}{2}\right]x\right)}{2\sin\left(\frac{1}{2}x\right)} + [0 - \sin(Nx)] && \text{by } \text{Muniz (1953) page 140} \\
&= \frac{1}{2} \cot\left(\frac{1}{2}x\right) - \frac{\cos\left(\left[N + \frac{1}{2}\right]x\right) - 2\sin\left(\frac{1}{2}x\right)\sin(Nx)}{2\sin\left(\frac{1}{2}x\right)} \\
&= \frac{1}{2} \cot\left(\frac{1}{2}x\right) - \frac{\cos\left(\left[N + \frac{1}{2}\right]x\right) - \left[\cos\left(\left[\frac{1}{2} - N\right]x\right) - \cos\left(\left[\frac{1}{2} + N\right]x\right)\right]}{2\sin\left(\frac{1}{2}x\right)} \\
&= \frac{1}{2} \cot\left(\frac{1}{2}x\right) + \frac{\cos\left(\left[N - \frac{1}{2}\right]x\right)}{2\sin\left(\frac{1}{2}x\right)} && \Rightarrow \text{above result}
\end{aligned}$$

⇒

### Theorem D.7. <sup>15</sup>

<sup>15</sup> [Jeffrey and Dai \(2008\)](#) pages 128–130 ⟨2.4.1.6 Sines, Cosines, and Tagents of Multiple Angles; (16) and (17)⟩



T H M	$\sum_{n=0}^{N-1} \cos(nx + y) = \cos(y) \left[ \frac{1}{2} + \frac{\sin\left(\left[N - \frac{1}{2}\right]x\right)}{2\sin\left(\frac{1}{2}x\right)} \right] - \sin(y) \left[ \frac{1}{2} \cot\left(\frac{1}{2}x\right) + \frac{\cos\left(\left[N - \frac{1}{2}\right]x\right)}{2\sin\left(\frac{1}{2}x\right)} \right] \quad \forall x \in \mathbb{R}$
	$\sum_{n=0}^{N-1} \sin(nx + y) = \cos(y) \left[ \frac{1}{2} + \frac{\sin\left(\left[N - \frac{1}{2}\right]x\right)}{2\sin\left(\frac{1}{2}x\right)} \right] + \sin(y) \left[ \frac{1}{2} \cot\left(\frac{1}{2}x\right) + \frac{\cos\left(\left[N - \frac{1}{2}\right]x\right)}{2\sin\left(\frac{1}{2}x\right)} \right] \quad \forall x \in \mathbb{R}$

PROOF:

$$\begin{aligned} \sum_{n=0}^{N-1} \cos(nx + y) &= \sum_{n=0}^{N-1} [\cos(nx)\cos(y) - \sin(nx)\sin(y)] && \text{by double angle formulas} && (\text{Theorem C.9 page 85}) \\ &= \cos(y) \sum_{n=0}^{N-1} \cos(nx) - \sin(y) \sum_{n=0}^{N-1} \sin(nx) \\ \sum_{n=0}^{N-1} \sin(nx + y) &= \sum_{n=0}^{N-1} [\cos(nx)\cos(y) + \sin(nx)\sin(y)] && \text{by double angle formulas} && (\text{Theorem C.9 page 85}) \\ &= \cos(y) \sum_{n=0}^{N-1} \cos(nx) + \sin(y) \sum_{n=0}^{N-1} \sin(nx) \end{aligned}$$

⇒

**Corollary D.1** (Summation around unit circle).

T H M	$\sum_{n=0}^{N-1} \cos\left(\theta + \frac{2nM\pi}{N}\right) = \sum_{n=0}^{N-1} \sin\left(\theta + \frac{2nM\pi}{N}\right) = \sum_{n=0}^{N-1} \cos\left(\theta + \frac{2nM\pi}{N}\right) \sin\left(\theta + \frac{2nM\pi}{N}\right) = 0 \quad \begin{matrix} \forall \theta \in \mathbb{R} \\ \forall M \in \mathbb{N} \end{matrix}$
	$\sum_{n=0}^{N-1} \cos^2\left(\theta + \frac{2nM\pi}{N}\right) = \sum_{n=0}^{N-1} \sin^2\left(\theta + \frac{2nM\pi}{N}\right) = \frac{N}{2} \quad \begin{matrix} \forall \theta \in \mathbb{R} \\ \forall M \in \mathbb{N} \end{matrix}$

PROOF:

$$\begin{aligned} &\sum_{n=0}^{N-1} \cos\left(\theta + \frac{2nM\pi}{N}\right) \\ &= \cos(\theta) \sum_{n=0}^{N-1} \cos\left(\frac{2nM\pi}{N}\right) - \sin(\theta) \sum_{n=0}^{N-1} \sin\left(\frac{2nM\pi}{N}\right) && \text{by Theorem C.9 page 85} \\ &= \cos(\theta) \left[ \frac{1}{2} + \frac{\sin\left(\left[N - \frac{1}{2}\right] \frac{2M\pi}{N}\right)}{2\sin\left(\frac{1}{2} \frac{2M\pi}{N}\right)} \right] - \sin(\theta) \left[ \frac{1}{2} \cot\left(\frac{1}{2} \frac{2M\pi}{N}\right) + \frac{\cos\left(\left[N - \frac{1}{2}\right] \frac{2M\pi}{N}\right)}{2\sin\left(\frac{1}{2} \frac{2M\pi}{N}\right)} \right] && \text{by Theorem D.6 page 105} \\ &= \cos(\theta) \left[ \frac{1}{2} - \frac{\sin\left(\frac{M\pi}{N} - 2M\pi\right)}{2\sin\left(\frac{M\pi}{N}\right)} \right] - \sin(\theta) \left[ \frac{1}{2} \cot\left(\frac{M\pi}{N}\right) - \frac{\cos\left(\frac{M\pi}{N} - 2M\pi\right)}{2\sin\left(\frac{M\pi}{N}\right)} \right] \\ &= \cos(\theta) \left[ \frac{1}{2} - \frac{1}{2} \frac{\sin\left(\frac{M\pi}{N}\right)}{\sin\left(\frac{M\pi}{N}\right)} \right] - \sin(\theta) \left[ \frac{1}{2} \cot\left(\frac{M\pi}{N}\right) - \frac{1}{2} \cot\left(\frac{M\pi}{N}\right) \right] && \text{by trigonometric periodicity} \\ & && (\text{Theorem C.10 page 85}) \\ &= \cos(\theta)[0] - \sin(\theta)[0] \\ &= 0 \end{aligned}$$

$$\begin{aligned}
\sum_{n=0}^{N-1} \sin\left(\theta + \frac{2nM\pi}{N}\right) &= \sum_{n=0}^{N-1} \cos\left(\theta - \frac{\pi}{2} + \frac{2nM\pi}{N}\right) && \text{by shift identities} && \text{(Theorem C.7 page 82)} \\
&= \sum_{n=0}^{N-1} \cos\left(\phi + \frac{2nM\pi}{N}\right) && \text{where } \phi \triangleq \theta - \frac{\pi}{2} \\
&= 0 && \text{by previous result}
\end{aligned}$$

$$\begin{aligned}
&\sum_{n=0}^{N-1} \cos\left(\theta + \frac{2nM\pi}{N}\right) \sin\left(\theta + \frac{2nM\pi}{N}\right) \\
&= -\frac{1}{2} \sum_{n=0}^{N-1} \sin\left(\left[\theta + \frac{2nM\pi}{N}\right] - \left[\theta + \frac{2nM\pi}{N}\right]\right) + \frac{1}{2} \sum_{n=0}^{N-1} \sin\left(\left[\theta + \frac{2nM\pi}{N}\right] + \left[\theta + \frac{2nM\pi}{N}\right]\right) && \text{by Theorem C.8 page 82} \\
&= -\frac{1}{2} \sum_{n=0}^{N-1} \sin(0) + \frac{1}{2} \sum_{n=0}^{N-1} \sin\left(2\theta + \frac{4nM\pi}{N}\right) \\
&= \frac{1}{2} \sin(2\theta) \sum_{n=0}^{N-1} \sin\left(\frac{4nM\pi}{N}\right) + \frac{1}{2} \cos(2\theta) \sum_{n=0}^{N-1} \sin\left(\frac{4nM\pi}{N}\right) && \text{by Theorem C.9 page 85} \\
&= \cos(2\theta) \left[ \frac{1}{2} + \frac{\sin\left(\left[N - \frac{1}{2}\right] \frac{4M\pi}{N}\right)}{2\sin\left(\frac{1}{2} \frac{2M\pi}{N}\right)} \right] - \sin(2\theta) \left[ \frac{1}{2} \cot\left(\frac{1}{2} \frac{4M\pi}{N}\right) + \frac{\cos\left(\left[N - \frac{1}{2}\right] \frac{4M\pi}{N}\right)}{2\sin\left(\frac{1}{2} \frac{4M\pi}{N}\right)} \right] && \text{by Theorem D.6 page 105} \\
&= \cos(2\theta) \left[ \frac{1}{2} - \frac{\sin\left(\frac{2M\pi}{N} - 4M\pi\right)}{2\sin\left(\frac{2M\pi}{N}\right)} \right] - \sin(2\theta) \left[ \frac{1}{2} \cot\left(\frac{2M\pi}{N}\right) - \frac{\cos\left(\frac{2M\pi}{N} - 4M\pi\right)}{2\sin\left(\frac{2M\pi}{N}\right)} \right] \\
&= \cos(\theta) \left[ \frac{1}{2} - \frac{1}{2} \frac{\sin\left(\frac{2M\pi}{N}\right)}{\sin\left(\frac{2M\pi}{N}\right)} \right] - \sin(\theta) \left[ \frac{1}{2} \cot\left(\frac{2M\pi}{N}\right) - \frac{1}{2} \cot\left(\frac{2M\pi}{N}\right) \right] && \text{by trigonometric periodicity} \\
& && \text{(Theorem C.10 page 85)} \\
&= \cos(\theta)[0] - \sin(\theta)[0] \\
&= 0
\end{aligned}$$

$$\begin{aligned}
\sum_{n=0}^{N-1} \cos^2\left(\theta + \frac{2nM\pi}{N}\right) &= \frac{1}{2} \sum_{n=0}^{N-1} \left[ 1 + \cos\left(2\theta + \frac{4nM\pi}{N}\right) \right] && \text{by Theorem C.11 page 87} \\
&= \frac{1}{2} \sum_{n=0}^{N-1} \left[ 1 + \cos(2\theta) \cos\left(\frac{4nM\pi}{N}\right) - \sin(2\theta) \sin\left(\frac{4nM\pi}{N}\right) \right] && \text{by Theorem C.9 page 85} \\
&= \frac{1}{2} \sum_{n=0}^{N-1} 1 + \frac{1}{2} \cos(2\theta) \sum_{n=0}^{N-1} \cos\left(\frac{4nM\pi}{N}\right) - \frac{1}{2} \sin(2\theta) \sum_{n=0}^{N-1} \sin\left(\frac{4nM\pi}{N}\right) \\
&= \left[ \frac{1}{2} \sum_{n=0}^{N-1} 1 \right] + \frac{1}{2} \cos(2\theta) 0 - \frac{1}{2} \sin(2\theta) 0 && \text{by previous results} \\
&= \frac{N}{2}
\end{aligned}$$

$$\begin{aligned}
\sum_{n=0}^{N-1} \sin^2\left(\theta + \frac{2nM\pi}{N}\right) &= \sum_{n=0}^{N-1} \cos^2\left(\theta - \frac{\pi}{2} + \frac{2nM\pi}{N}\right) && \text{by shift identities (Theorem C.7 page 82)} \\
&= \sum_{n=0}^{N-1} \cos^2\left(\phi + \frac{2nM\pi}{N}\right) && \text{where } \phi \triangleq \theta - \frac{\pi}{2} \\
&= \frac{N}{2} && \text{by previous result}
\end{aligned}$$



## D.6 Summability Kernels

**Definition D.3.** <sup>16</sup> Let  $(\kappa_n)_{n \in \mathbb{Z}}$  be a sequence of CONTINUOUS  $2\pi$  PERIODIC functions.

The sequence  $(\kappa_n)_{n \in \mathbb{Z}}$  is a **summability kernel** if

1.  $\frac{1}{2\pi} \int_0^{2\pi} \kappa_n(x) \, dx = 1 \quad \forall n \in \mathbb{Z}$  and
2.  $\frac{1}{2\pi} \int_0^{2\pi} |\kappa_n(x)| \, dx \in \mathbb{R} \quad \forall n \in \mathbb{Z}$  and
3.  $\lim_{n \rightarrow \infty} \int_{\delta}^{2\pi-\delta} |\kappa_n(x)| \, dx = 0 \quad \forall n \in \mathbb{Z}, 0 < \delta < \pi$

**Theorem D.8.** <sup>17</sup> Let  $(\kappa_n)_{n \in \mathbb{Z}}$  be a sequence. Let  $\mathbb{T}$  be the quotient  $\mathbb{R}/2\pi\mathbb{Z}$ .

1.  $f \in L^1(\mathbb{T})$  and
  2.  $(\kappa_n)$  is a summability kernel
- $$\implies f(x) = \lim_{n \rightarrow \infty} \frac{1}{2\pi} \int_{\mathbb{T}} \kappa_n(x) f(x - x) \, dx$$

The *Dirichlet kernel* (Definition D.2 page 102) is *not* a summability kernel. Examples of kernels that *are* summability kernels include

1. *Fejér's kernel* (Definition D.4 page 109)
2. *de la Vallée Poussin kernel* (Definition D.5 page 111)
3. *Jackson kernel* (Definition D.6 page 111)
4. *Poisson kernel* (Definition D.7 page 111.)

**Definition D.4.** <sup>18</sup>

*Fejér's kernel*  $K_n$  is defined as

$$K_n(x) \triangleq \sum_{k=-n}^{k=n} \left(1 - \frac{|k|}{n+1}\right) e^{ikx}$$

**Proposition D.5.** <sup>19</sup> Let  $K_n$  be Fejér's kernel (Definition D.4 page 109).

$$K_n(x) = \frac{1}{n+1} \left( \frac{\sin \frac{n+1}{2} x}{\sin \frac{1}{2} x} \right)^2$$

<sup>16</sup> Cerdà (2010) page 56, Katznelson (2004) page 10, de Reyna (2002) page 21, Walnut (2002) pages 40–41, Heil (2011) page 440, Istrăţescu (1987) page 309

<sup>17</sup> Katznelson (2004) page 11

<sup>18</sup> Katznelson (2004) page 12

<sup>19</sup> Katznelson (2004) page 12, Heil (2011) page 448

 PROOF:

1. Lemma: Proof that  $\sin^2 \frac{x}{2} \equiv \frac{-1}{4}(e^{-ix} - 2 + e^{ix})$ :

$$\begin{aligned} \sin^2 \frac{x}{2} &\equiv \left( \frac{e^{-i\frac{x}{2}} - e^{+i\frac{x}{2}}}{2i} \right)^2 && \text{by Euler Formulas (Corollary C.2 page 81)} \\ &\equiv \frac{-1}{4} \left( e^{-2i\frac{x}{2}} - 2e^{-i\frac{x}{2}}e^{i\frac{x}{2}} + e^{2i\frac{x}{2}} \right) \\ &\equiv \frac{-1}{4} (e^{-ix} - 2 + e^{ix}) : \end{aligned}$$

2. Lemma:

$$2|k| - |k+1| - |k-1| = \begin{cases} -2 & \text{for } k = 0 \\ 0 & \text{for } k \in \mathbb{Z} \setminus 0 \end{cases}$$

3. Proof that  $K_n(x) = \frac{1}{n+1} \left( \frac{\sin \frac{n+1}{2}x}{\sin \frac{1}{2}x} \right)^2$ :

$$\begin{aligned} &-4(n+1) \left( \sin \frac{1}{2}x \right)^2 K_n(x) \\ &= -4(n+1) \left( \frac{-1}{4} \right) (e^{-ix} - 2 + e^{ix}) K_n(x) && \text{by item (1)} \\ &= (n+1) (e^{-ix} - 2 + e^{ix}) \sum_{k=-n}^{k=n} \left( 1 - \frac{|k|}{n+1} \right) e^{ikx} && \text{by Definition D.4} \\ &= (n+1) \frac{1}{n+1} (e^{-ix} - 2 + e^{ix}) \sum_{k=-n}^{k=n} (n+1 - |k|) e^{ikx} \\ &= e^{-ix} \sum_{k=-n}^{k=n} (n+1 - |k|) e^{ikx} - 2 \sum_{k=-n}^{k=n} (n+1 - |k|) e^{ikx} e^{ix} \sum_{k=-n}^{k=n} (n+1 - |k|) e^{ikx} \\ &= \sum_{k=-n}^{k=n} (n+1 - |k|) e^{i(k-1)x} - 2 \sum_{k=-n}^{k=n} (n+1 - |k|) e^{ikx} \sum_{k=-n}^{k=n} (n+1 - |k|) e^{i(k+1)x} \\ &= \sum_{k=-n-1}^{k=n-1} (n+1 - |k+1|) e^{ikx} - 2 \sum_{k=-n}^{k=n} (n+1 - |k|) e^{ikx} \sum_{k=-n+1}^{k=n+1} (n+1 - |k-1|) e^{ikx} \\ &= \underbrace{e^{-i(n+1)x}}_{k=-n-1} + \underbrace{2e^{-inx}}_{k=-n} + \sum_{k=-n+1}^{k=n-1} (n+1 - |k+1|) e^{ikx} + \\ &\quad \underbrace{-2e^{-inx}}_{k=-n} + \underbrace{-2e^{inx}}_{k=n} - 2 \sum_{k=-n+1}^{k=n-1} (n+1 - |k|) e^{ikx} + \\ &\quad \underbrace{e^{i(n+1)x}}_{k=n+1} + \underbrace{2e^{inx}}_{k=n} + \sum_{k=-n+1}^{k=n-1} (n+1 - |k-1|) e^{ikx} \\ &= e^{-i(n+1)x} + \sum_{k=-n+1}^{k=n-1} (n+1 - |k+1|) e^{ikx} + \\ &\quad -2 \sum_{k=-n+1}^{k=n-1} (n+1 - |k|) e^{ikx} + \\ &\quad e^{i(n+1)x} + \sum_{k=-n+1}^{k=n-1} (n+1 - |k-1|) e^{ikx} \end{aligned}$$



$$\begin{aligned}
&= e^{-i(n+1)x} + e^{i(n+1)x} + \sum_{k=-n+1}^{k=n-1} [(n+1-|k+1|) - 2(n+1-|k|) + (n+1-|k-1|)] e^{ikx} \\
&= e^{-i(n+1)x} + e^{i(n+1)x} + \sum_{k=-n+1}^{k=n-1} (2|k| - |k+1| - |k-1|) e^{ikx} \\
&= e^{-i(n+1)x} + e^{i(n+1)x} - 2 \quad \text{by item (2)} \\
&= -4 \left( \sin \frac{n+1}{2} x \right)^2 \quad \text{by item (1)}
\end{aligned}$$



**Definition D.5.** <sup>20</sup> Let  $K_n$  be FEJÉR'S KERNEL (Definition D.4 page 109).

**DEF** The *de la Vallée Poussin kernel*  $V_n$  is defined as

$$V_n(x) \triangleq 2K_{2n+1}(x) - K_n(x)$$

**Definition D.6.** <sup>21</sup> Let  $K_n$  be FEJÉR'S KERNEL (Definition D.4 page 109).

**DEF** The *Jackson kernel*  $J_n$  is defined as

$$J_n(x) \triangleq \|K_n\|^{-2} K_n^2(x)$$

**Definition D.7.** <sup>22</sup>

**DEF** The *Poisson kernel*  $P$  is defined as

$$P(r, x) \triangleq \sum_{k \in \mathbb{Z}} r^{|k|} e^{ikx}$$

<sup>20</sup> Katznelson (2004) page 16

<sup>21</sup> Katznelson (2004) page 17

<sup>22</sup> Katznelson (2004) page 16



# APPENDIX E

---

## NORMED ALGEBRAS

### E.1 Algebras

All *linear spaces* are equipped with an operation by which vectors in the spaces can be added together. Linear spaces also have an operation that allows a scalar and a vector to be “multiplied” together. But linear spaces in general have no operation that allows two vectors to be multiplied together. A linear space together with such an operator is an **algebra**.<sup>1</sup>


There are many many possible algebras—many more than one can shake a stick at, as indicated by Michiel Hazewinkel in his book, *Handbook of Algebras*: “Algebra, as we know it today (2005), consists of many different ideas, concepts and results. A reasonable estimate of the number of these different items would be somewhere between 50,000 and 200,000. Many of these have been named and many more could (and perhaps should) have a “name” or other convenient designation.”<sup>2</sup>

**Definition E.1.**<sup>3</sup> Let  $A$  be an ALGEBRA.


**DEF** An algebra  $A$  is **unital** if  $\exists u \in A$  such that  $ux = xu = x \quad \forall x \in A$

**Definition E.2.**<sup>4</sup> Let  $A$  be an UNITAL ALGEBRA (Definition E.1 page 113) with unit  $e$ .

**DEF** The **spectrum** of  $x \in A$  is  $\sigma(x) \triangleq \{\lambda \in \mathbb{C} \mid \lambda e - x \text{ is not invertible}\}.$   
 The **resolvent** of  $x \in A$  is  $\rho_x(\lambda) \triangleq (\lambda e - x)^{-1} \quad \forall \lambda \notin \sigma(x).$   
 The **spectral radius** of  $x \in A$  is  $r(x) \triangleq \sup \{|\lambda| \mid \lambda \in \sigma(x)\}.$

<sup>1</sup>  Fuchs (1995) page 2

<sup>2</sup>  Hazewinkel (2000) page v

<sup>3</sup>  Folland (1995) page 1

<sup>4</sup>  Folland (1995) pages 3–4

## E.2 Star-Algebras

**Definition E.3.**<sup>5</sup> Let  $A$  be an ALGEBRA.

The pair  $(A, *)$  is a ***\*-algebra***, or ***star-algebra***, if

DEF

1.  $(x + y)^* = x^* + y^* \quad \forall x, y \in A$  (DISTRIBUTIVE) and
2.  $(\alpha x)^* = \bar{\alpha} x^* \quad \forall x \in A, \alpha \in \mathbb{C}$  (CONJUGATE LINEAR) and
3.  $(xy)^* = y^* x^* \quad \forall x, y \in A$  (ANTIAUTOMORPHIC) and
4.  $x^{**} = x \quad \forall x \in A$  (INVOLUTORY)

The operator  $*$  is called an ***involution*** on the algebra  $A$ .

**Proposition E.1.**<sup>6</sup> Let  $(A, *)$  be an UNITAL \*-ALGEBRA.

PRP

$x$  is invertible  $\implies \begin{cases} 1. & x^* \text{ is INVERTIBLE } \forall x \in A \text{ and} \\ 2. & (x^*)^{-1} = (x^{-1})^* \quad \forall x \in A \end{cases}$

PROOF: Let  $e$  be the unit element of  $(A, *)$ .

1. Proof that  $e^* = e$ :

$$\begin{aligned}
 x e^* &= (x e^*)^{**} && \text{by involutory property of } * && (\text{Definition E.3 page 114}) \\
 &= (x^* e^{**})^* && \text{by antiautomorphic property of } * && (\text{Definition E.3 page 114}) \\
 &= (x^* e)^* && \text{by involutory property of } * && (\text{Definition E.3 page 114}) \\
 &= (x^*)^* && \text{by definition of } e \\
 &= x && \text{by involutory property of } * && (\text{Definition E.3 page 114}) \\
 e^* x &= (e^* x)^{**} && \text{by involutory property of } * && (\text{Definition E.3 page 114}) \\
 &= (e^{**} x^*)^* && \text{by antiautomorphic property of } * && (\text{Definition E.3 page 114}) \\
 &= (e x^*)^* && \text{by involutory property of } * && (\text{Definition E.3 page 114}) \\
 &= (x^*)^* && \text{by definition of } e \\
 &= x && \text{by involutory property of } * && (\text{Definition E.3 page 114})
 \end{aligned}$$




2. Proof that  $(x^*)^{-1} = (x^{-1})^*$ :

$$\begin{aligned}
 (x^{-1})^* (x^*) &= [x (x^{-1})]^* && \text{by antiautomorphic and involution properties of } * && (\text{Definition E.3 page 114}) \\
 &= e^* \\
 &= e && \text{by item (1) page 114} \\
 (x^*) (x^{-1})^* &= [x^{-1} x]^* && \text{by antiautomorphic and involution properties of } * && (\text{Definition E.3 page 114}) \\
 &= e^* \\
 &= e && \text{by item (1) page 114}
 \end{aligned}$$


$\Rightarrow$

**Definition E.4.**<sup>7</sup> Let  $(A, \|\cdot\|)$  be a \*-ALGEBRA (Definition E.3 page 114).

DEF

-  An element  $x \in A$  is ***hermitian*** or ***self-adjoint*** if  $x^* = x$ .
-  An element  $x \in A$  is ***normal*** if  $xx^* = x^*x$ .
-  An element  $x \in A$  is a ***projection*** if  $xx = x$  (INVOLUTORY) and  $x^* = x$  (HERMITIAN).

<sup>5</sup>  Rickart (1960) page 178,  Gelfand and Naimark (1964), page 241

<sup>6</sup>  Folland (1995) page 5

<sup>7</sup>  Rickart (1960) page 178,  Gelfand and Naimark (1964), page 242

**Theorem E.1.** <sup>8</sup> Let  $(A, \|\cdot\|)$  be a  $*$ -ALGEBRA (Definition E.3 page 114).

T H M	$\underbrace{x = x^* \text{ and } y = y^*}_{x \text{ and } y \text{ are HERMITIAN}} \implies \begin{cases} x + y = (x + y)^* & (x + y \text{ is self adjoint}) \\ x^* = (x^*)^* & (x^* \text{ is self adjoint}) \\ \underbrace{xy = (xy)^*}_{(xy) \text{ is HERMITIAN}} \iff \underbrace{xy = yx}_{\text{commutative}} \end{cases}$
-------------	---

PROOF:

$$\begin{aligned} (x + y)^* &= x^* + y^* && \text{by distributive property of } * && (\text{Definition E.3 page 114}) \\ &= x + y && \text{by left hypothesis} \end{aligned}$$

$$(x^*)^* = x \quad \text{by involutory property of } * \quad (\text{Definition E.3 page 114})$$

Proof that  $xy = (xy)^* \implies xy = yx$

$$\begin{aligned} xy &= (xy)^* && \text{by left hypothesis} \\ &= y^* x^* && \text{by antiautomorphic property of } * && (\text{Definition E.3 page 114}) \\ &= yx && \text{by left hypothesis} \end{aligned}$$

Proof that  $xy = (xy)^* \iff xy = yx$

$$\begin{aligned} (xy)^* &= (yx)^* && \text{by left hypothesis} \\ &= x^* y^* && \text{by antiautomorphic property of } * && (\text{Definition E.3 page 114}) \\ &= xy && \text{by left hypothesis} \end{aligned}$$

⇒

**Definition E.5** (Hermitian components). <sup>9</sup> Let  $(A, \|\cdot\|)$  be a  $*$ -ALGEBRA (Definition E.3 page 114).

D E F	<p>The <b>real part</b> of <math>x</math> is defined as <math>\mathbf{R}_e x \triangleq \frac{1}{2}(x + x^*)</math></p> <p>The <b>imaginary part</b> of <math>x</math> is defined as <math>\mathbf{I}_m x \triangleq \frac{1}{2i}(x - x^*)</math></p>
-------------	---

**Theorem E.2.** <sup>10</sup> Let  $(A, *)$  be a  $*$ -ALGEBRA (Definition E.3 page 114).

T H M	$\begin{aligned} \mathbf{R}_e x &= (\mathbf{R}_e x)^* & \forall x \in A & \quad (\mathbf{R}_e x \text{ is HERMITIAN}) \\ \mathbf{I}_m x &= (\mathbf{I}_m x)^* & \forall x \in A & \quad (\mathbf{I}_m x \text{ is HERMITIAN}) \end{aligned}$
-------------	--

PROOF:

$$\begin{aligned} (\mathbf{R}_e x)^* &= \left( \frac{1}{2}(x + x^*) \right)^* && \text{by definition of } \mathfrak{R} && (\text{Definition E.5 page 115}) \\ &= \frac{1}{2}(x^* + x^{**}) && \text{by distributive property of } * && (\text{Definition E.3 page 114}) \\ &= \frac{1}{2}(x^* + x) && \text{by involutory property of } * && (\text{Definition E.3 page 114}) \\ &= \mathbf{R}_e x && \text{by definition of } \mathfrak{R} && (\text{Definition E.5 page 115}) \\ (\mathbf{I}_m x)^* &= \left( \frac{1}{2i}(x - x^*) \right)^* && \text{by definition of } \mathfrak{I} && (\text{Definition E.5 page 115}) \end{aligned}$$

<sup>8</sup> Michel and Herget (1993) page 429

<sup>9</sup> Michel and Herget (1993) page 430, Rickart (1960) page 179, Gelfand and Naimark (1964), page 242

<sup>10</sup> Michel and Herget (1993) page 430, Halmos (1998) page 42

$$\begin{aligned}
&= \frac{1}{2i}(x^* - x^{**}) && \text{by distributive property of } * && (\text{Definition E.3 page 114}) \\
&= \frac{1}{2i}(x^* - x) && \text{by involutory property of } * && (\text{Definition E.3 page 114}) \\
&= \mathbf{I}_m x && \text{by definition of } \mathfrak{I} && (\text{Definition E.5 page 115})
\end{aligned}$$

⇒

**Theorem E.3** (Hermitian representation).<sup>11</sup> Let  $(A, *)$  be a  $*$ -ALGEBRA (Definition E.3 page 114).

T H M	$a = x + iy \quad \Longleftrightarrow \quad x = \mathbf{R}_e a \quad \text{and} \quad y = \mathbf{I}_m a$
-------------	---

✎ PROOF:

🔥 Proof that  $a = x + iy \implies x = \mathbf{R}_e a$  and  $y = \mathbf{I}_m a$ :

$$\begin{aligned}
& \implies a = x + iy && \text{by left hypothesis} \\
& \implies a^* = (x + iy)^* && \text{by definition of adjoint} && (\text{Definition E.4 page 114}) \\
& \quad = x^* - iy^* && \text{by distributive property of } * && (\text{Definition E.3 page 114}) \\
& \quad = x - iy && \text{by Theorem E.2 page 115} \\
& \implies x = a - iy && \text{by solving for } x \text{ in } a = x + iy \text{ equation} \\
& \quad x = a^* + iy && \text{by solving for } x \text{ in } a^* = x - iy \text{ equation} \\
& \implies x + x = a + a^* && \text{by adding previous 2 equations} \\
& \implies 2x = a + a^* && \text{by solving for } x \text{ in previous equation} \\
& \implies x = \frac{1}{2}(a + a^*) && \\
& \quad = \mathbf{R}_e a && \text{by definition of } \mathfrak{R} && (\text{Definition E.5 page 115}) \\
& \implies iy = a - x && \text{by solving for } iy \text{ in } a = x + iy \text{ equation} \\
& \quad iy = -a^* + x && \text{by solving for } iy \text{ in } a^* = x - iy \text{ equation} \\
& \implies iy + iy = a - a^* && \text{by adding previous 2 equations} \\
& \implies y = \frac{1}{2i}(a - a^*) && \text{by solving for } iy \text{ in previous equations} \\
& \quad = \mathbf{I}_m a && \text{by definition of } \mathfrak{I} && (\text{Definition E.5 page 115})
\end{aligned}$$

🔥 Proof that  $a = x + iy \Leftarrow x = \mathbf{R}_e a$  and  $y = \mathbf{I}_m a$ :

$$\begin{aligned}
x + iy &= \mathbf{R}_e a + i \mathbf{I}_m a && \text{by right hypothesis} \\
&= \underbrace{\frac{1}{2}(a + a^*)}_{\mathbf{R}_e a} + i \underbrace{\frac{1}{2i}(a - a^*)}_{\mathbf{I}_m a} && \text{by definition of } \mathfrak{R} \text{ and } \mathfrak{I} && (\text{Definition E.5 page 115}) \\
&= \left(\frac{1}{2}a + \frac{1}{2}a\right) + \left(\frac{1}{2}a^* - \frac{1}{2}a^*\right) && \text{red arrow points to } 0 \\
&= a
\end{aligned}$$

⇒

<sup>11</sup> Michel and Herget (1993) page 430, Rickart (1960) page 179, Gelfand and Neumark (1943b) page 7

## E.3 Normed Algebras

**Definition E.6.** <sup>12</sup> Let  $A$  be an algebra.

DEF

The pair  $(A, \|\cdot\|)$  is a **normed algebra** if

$$\|xy\| \leq \|x\| \|y\| \quad \forall x, y \in A \quad (\text{multiplicative condition})$$

A normed algebra  $(A, \|\cdot\|)$  is a **Banach algebra** if  $(A, \|\cdot\|)$  is also a Banach space.

**Proposition E.2.**

PRP

$(A, \|\cdot\|)$  is a normed algebra  $\implies$  multiplication is **continuous** in  $(A, \|\cdot\|)$

 PROOF:

1. Define  $f(x) \triangleq zx$ . That is, the function  $f$  represents multiplication of  $x$  times some arbitrary value  $z$ .
2. Let  $\delta \triangleq \|x - y\|$  and  $\epsilon \triangleq \|f(x) - f(y)\|$ .
3. To prove that multiplication ( $f$ ) is *continuous* with respect to the metric generated by  $\|\cdot\|$ , we have to show that we can always make  $\epsilon$  arbitrarily small for some  $\delta > 0$ .
4. And here is the proof that multiplication is indeed continuous in  $(A, \|\cdot\|)$ :

$$\begin{aligned} \|f(x) - f(y)\| &\triangleq \|zx - zy\| && \text{by definition of } f && (\text{item (1) page 117}) \\ &= \|z(x - y)\| \\ &\leq \|z\| \|x - y\| && \text{by definition of normed algebra} && (\text{Definition E.6 page 117}) \\ &\triangleq \|z\| \delta && \text{by definition of } \delta && (\text{item (2) page 117}) \\ &\leq \epsilon && \text{for some value of } \delta > 0 \end{aligned}$$



**Theorem E.4** (Gelfand-Mazur Theorem). <sup>13</sup> Let  $\mathbb{C}$  be the field of complex numbers.

THM

$\left. \begin{array}{l} (A, \|\cdot\|) \text{ is a Banach algebra} \\ \text{every nonzero } x \in A \text{ is invertible} \end{array} \right\} \implies A \equiv \mathbb{C} \quad (A \text{ is isomorphic to } \mathbb{C})$

## E.4 $C^*$ Algebras

**Definition E.7.** <sup>14</sup>




DEF





The triple  $(A, \|\cdot\|, *)$  is a  **$C^*$  algebra** if

1.  $(A, \|\cdot\|)$  is a Banach algebra and
2.  $(A, *)$  is a  $*$ -algebra and
3.  $\|x^*x\| = \|x\|^2 \quad \forall x \in A$ .

A  **$C^*$  algebra**  $(A, \|\cdot\|, *)$  is also called a  **$C$  star algebra**.

<sup>12</sup>  Rickart (1960) page 2,  Berberian (1961) page 103 (Theorem IV.9.2)

<sup>13</sup>  Folland (1995) page 4,  Mazur (1938) (statement),  Gelfand (1941) (proof)

<sup>14</sup>  Folland (1995) page 1,  Gelfand and Naimark (1964), page 241,  Gelfand and Neumark (1943a),  Gelfand and Neumark (1943b)

**Theorem E.5.** <sup>15</sup> *Let  $A$  be an algebra.*

<b>T H M</b>	$(A, \ \cdot\ , *) \text{ is a } C^* \text{ algebra} \quad \implies \quad \ x^*\  = \ x\ $
----------------------	--

 PROOF:

$\ x\  = \frac{1}{\ x\ } \ x\ ^2$		
$= \frac{1}{\ x\ } \ x^* x\ $	by definition of $C^*$ -algebra	(Definition E.7 page 117)
$\leq \frac{1}{\ x\ } \ x^*\  \ x\ $	by definition of <i>normed algebra</i>	(Definition E.6 page 117)
$= \ x^*\ $		
$\ x^*\  \leq \ x^{**}\ $	by previous result	
$= \ x\ $	by <i>involution</i> property of $*$	(Definition E.3 page 114)



<sup>15</sup>  Folland (1995) page 1,  Gelfand and Neumark (1943b) page 4,  Gelfand and Neumark (1943a)



# APPENDIX F

## OPERATORS ON LINEAR SPACES



*“And I am not afraid to say that there is a way to advance algebra as far beyond what Vieta and Descartes have left us as Vieta and Descartes carried it beyond the ancients... we need still another analysis which is distinctly geometrical or linear, and which will express situation directly as algebra expresses magnitude directly.”*

Gottfried Leibniz (1646–1716), German mathematician, in a September 8, 1679 letter to Christian Huygens. <sup>1</sup>

## F.1 Operators on linear spaces

### F.1.1 Operator Algebra

An operator is simply a function that maps from a linear space to another linear space (or to the same linear space).

**Definition F.1.** <sup>2</sup> Let  $(\mathbb{F}, +, \cdot, 0, 1)$  be a FIELD (Definition A.5 page 68). Let  $X$  be a set, let  $+$  be an OPERATOR (Definition F.2 page 120) in  $X^{X^2}$ , and let  $\otimes$  be an operator in  $X^{\mathbb{F} \times X}$ .

<sup>1</sup> quote: [Leibniz \(1679\) pages 248–249](#)

image: [http://en.wikipedia.org/wiki/File:Gottfried\\_Wilhelm\\_von\\_Leibniz.jpg](http://en.wikipedia.org/wiki/File:Gottfried_Wilhelm_von_Leibniz.jpg), public domain

<sup>2</sup> [Kubrusly \(2001\) pages 40–41](#) (Definition 2.1 and following remarks), [Haaser and Sullivan \(1991\) page 41](#), [Halmos \(1948\) pages 1–2](#), [Peano \(1888a\)](#) (Chapter IX), [Peano \(1888b\) pages 119–120](#), [Banach \(1922\) pages 134–135](#)




Figure F.1: Some operator types

The structure  $\Omega \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}))$  is a **linear space** over  $(\mathbb{F}, +, \cdot, 0, 1)$  if

- |    |                             |   |  |                               |    |
|----|-----------------------------|---|--|-------------------------------|----|
| 1. | $\exists 0 \in X$ such that | $x + 0 = x$   | $\forall x \in X$                                  | (+ IDENTITY)                  | *] |
| 2. | $\exists y \in X$ such that | $x + y = 0$   | $\forall x \in X$                                  | (+ INVERSE)                   |    |
| 3. |                             | $(x + y) + z = x + (y + z)$                                     | $\forall x, y, z \in X$                            | (+ is ASSOCIATIVE)            |    |
| 4. |                             | $x + y = y + x$   | $\forall x, y \in X$                               | (+ is COMMUTATIVE)            |    |
| 5. |                             | $1 \cdot x = x$   | $\forall x \in X$                                  | (· IDENTITY)                  |    |
| 6. |                             | $\alpha \cdot (\beta \cdot x) = (\alpha \cdot \beta) \cdot x$   | $\forall \alpha, \beta \in S \text{ and } x \in X$ | (· ASSOCIATES with ·)         |    |
| 7. |                             | $\alpha \cdot (x + y) = (\alpha \cdot x) + (\alpha \cdot y)$    | $\forall \alpha \in S \text{ and } x, y \in X$     | (· DISTRIBUTES over +)        |    |
| 8. |                             | $(\alpha + \beta) \cdot x = (\alpha \cdot x) + (\beta \cdot x)$ | $\forall \alpha, \beta \in S \text{ and } x \in X$ | (· PSEUDO-DISTRIBUTES over +) |    |

The set  $X$  is called the **underlying set**. The elements of  $X$  are called **vectors**. The elements of  $\mathbb{F}$  are called **scalars**. A linear space is also called a **vector space**. If  $\mathbb{F} \triangleq \mathbb{R}$ , then  $\Omega$  is a **real linear space**. If  $\mathbb{F} \triangleq \mathbb{C}$ , then  $\Omega$  is a **complex linear space**.

### Definition F2. <sup>3</sup>

**DEF** A function  $A$  in  $Y^X$  is an **operator** in  $Y^X$  if  $X$  and  $Y$  are both LINEAR SPACES (Definition F.1 page 119).

Two operators  $A$  and  $B$  in  $Y^X$  are **equal** if  $Ax = Bx$  for all  $x \in X$ . The inverse relation of an operator  $A$  in  $Y^X$  always exists as a *relation* in  $2^{X^Y}$ , but may not always be a *function* (may not always be an operator) in  $Y^X$ .

The operator  $I \in X^X$  is the *identity* operator if  $Ix = x$  for all  $x \in X$ .

**Definition F3. <sup>4</sup>** Let  $X^X$  be the set of all operators with from a LINEAR SPACE  $X$  to  $X$ . Let  $I$  be an operator in  $X^X$ . Let  $\mathbb{I}(X)$  be the IDENTITY ELEMENT in  $X^X$ .

**DEF**  $I$  is the **identity operator** in  $X^X$  if  $I = \mathbb{I}(X)$ .

<sup>3</sup> Heil (2011) page 42

<sup>4</sup> Michel and Herget (1993) page 411

## F.1.2 Linear operators

**Definition F.4.** <sup>5</sup> Let  $\mathbf{X} \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}))$  and  $\mathbf{Y} \triangleq (Y, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}))$  be linear spaces.

DEF

An operator  $\mathbf{L} \in \mathbf{Y}^{\mathbf{X}}$  is **linear** if

1.  $\mathbf{L}(\mathbf{x} + \mathbf{y}) = \mathbf{L}\mathbf{x} + \mathbf{L}\mathbf{y} \quad \forall \mathbf{x}, \mathbf{y} \in \mathbf{X} \quad (\text{ADDITIVE}) \quad \text{and}$
2.  $\mathbf{L}(\alpha \mathbf{x}) = \alpha \mathbf{L}\mathbf{x} \quad \forall \mathbf{x} \in \mathbf{X}, \quad \forall \alpha \in \mathbb{F} \quad (\text{HOMOGENEOUS}).$

The set of all linear operators from  $\mathbf{X}$  to  $\mathbf{Y}$  is denoted  $\mathcal{L}(\mathbf{X}, \mathbf{Y})$  such that  $\mathcal{L}(\mathbf{X}, \mathbf{Y}) \triangleq \{\mathbf{L} \in \mathbf{Y}^{\mathbf{X}} \mid \mathbf{L} \text{ is linear}\}$ .

**Theorem F.1.** <sup>6</sup> Let  $\mathbf{L}$  be an operator from a linear space  $\mathbf{X}$  to a linear space  $\mathbf{Y}$ , both over a field  $\mathbb{F}$ .

THM

$$\{\mathbf{L} \text{ is LINEAR}\} \implies \left\{ \begin{array}{ll} 1. \mathbf{L}\mathbf{0} &= \mathbf{0} \quad \text{and} \\ 2. \mathbf{L}(-\mathbf{x}) &= -(\mathbf{L}\mathbf{x}) \quad \forall \mathbf{x} \in \mathbf{X} \quad \text{and} \\ 3. \mathbf{L}(\mathbf{x} - \mathbf{y}) &= \mathbf{L}\mathbf{x} - \mathbf{L}\mathbf{y} \quad \forall \mathbf{x}, \mathbf{y} \in \mathbf{X} \quad \text{and} \\ 4. \mathbf{L}\left(\sum_{n=1}^N \alpha_n \mathbf{x}_n\right) &= \sum_{n=1}^N \alpha_n (\mathbf{L}\mathbf{x}_n) \quad \mathbf{x}_n \in \mathbf{X}, \alpha_n \in \mathbb{F} \end{array} \right\}$$

 PROOF:

1. Proof that  $\mathbf{L}\mathbf{0} = \mathbf{0}$ :

$$\begin{aligned} \mathbf{L}\mathbf{0} &= \mathbf{L}(\mathbf{0} \cdot \mathbf{0}) && \text{by additive identity property} \\ &= \mathbf{0} \cdot (\mathbf{L}\mathbf{0}) && \text{by homogeneous property of } \mathbf{L} \quad (\text{Definition F.4 page 121}) \\ &= \mathbf{0} && \text{by additive identity property} \end{aligned}$$

2. Proof that  $\mathbf{L}(-\mathbf{x}) = -(\mathbf{L}\mathbf{x})$ :

$$\begin{aligned} \mathbf{L}(-\mathbf{x}) &= \mathbf{L}(-1 \cdot \mathbf{x}) && \text{by additive inverse property} \\ &= -1 \cdot (\mathbf{L}\mathbf{x}) && \text{by homogeneous property of } \mathbf{L} \quad (\text{Definition F.4 page 121}) \\ &= -(\mathbf{L}\mathbf{x}) && \text{by additive inverse property} \end{aligned}$$





3. Proof that  $\mathbf{L}(\mathbf{x} - \mathbf{y}) = \mathbf{L}\mathbf{x} - \mathbf{L}\mathbf{y}$ :

$$\begin{aligned} \mathbf{L}(\mathbf{x} - \mathbf{y}) &= \mathbf{L}(\mathbf{x} + (-\mathbf{y})) && \text{by additive inverse property} \\ &= \mathbf{L}(\mathbf{x}) + \mathbf{L}(-\mathbf{y}) && \text{by linearity property of } \mathbf{L} \quad (\text{Definition F.4 page 121}) \\ &= \mathbf{L}\mathbf{x} - \mathbf{L}\mathbf{y} && \text{by item (2)} \end{aligned}$$

4. Proof that  $\mathbf{L}\left(\sum_{n=1}^N \alpha_n \mathbf{x}_n\right) = \sum_{n=1}^N \alpha_n (\mathbf{L}\mathbf{x}_n)$ :

(a) Proof for  $N = 1$ :

$$\begin{aligned} \mathbf{L}\left(\sum_{n=1}^N \alpha_n \mathbf{x}_n\right) &= \mathbf{L}(\alpha_1 \mathbf{x}_1) && \text{by } N = 1 \text{ hypothesis} \\ &= \alpha_1 (\mathbf{L}\mathbf{x}_1) && \text{by homogeneous property of } \mathbf{L} \quad (\text{Definition F.4 page 121}) \end{aligned}$$

<sup>5</sup>  Kubrusly (2001) page 55,  Aliprantis and Burkinshaw (1998) page 224,  Hilbert et al. (1927) page 6,  Stone (1932) page 33

<sup>6</sup>  Berberian (1961) page 79 (Theorem IV.1.1)

(b) Proof that  $N$  case  $\implies N + 1$  case:

$$\begin{aligned}
 \mathbf{L}\left(\sum_{n=1}^{N+1} \alpha_n \mathbf{x}_n\right) &= \mathbf{L}\left(\alpha_{N+1} \mathbf{x}_{N+1} + \sum_{n=1}^N \alpha_n \mathbf{x}_n\right) \\
 &= \mathbf{L}(\alpha_{N+1} \mathbf{x}_{N+1}) + \mathbf{L}\left(\sum_{n=1}^N \alpha_n \mathbf{x}_n\right) \quad \text{by linearity property of } \mathbf{L} \quad (\text{Definition F.4 page 121}) \\
 &= \alpha_{N+1} \mathbf{L}(\mathbf{x}_{N+1}) + \sum_{n=1}^N \mathbf{L}(\alpha_n \mathbf{x}_n) \quad \text{by left } N + 1 \text{ hypothesis} \\
 &= \sum_{n=1}^{N+1} \mathbf{L}(\alpha_n \mathbf{x}_n)
 \end{aligned}$$

$\Rightarrow$

**Theorem F.2.** <sup>7</sup> Let  $\mathcal{L}(\mathbf{X}, \mathbf{Y})$  be the set of all linear operators from a linear space  $\mathbf{X}$  to a linear space  $\mathbf{Y}$ . Let  $\mathcal{N}(\mathbf{L})$  be the NULL SPACE of an operator  $\mathbf{L}$  in  $\mathbf{Y}^{\mathbf{X}}$  and  $\mathcal{I}(\mathbf{L})$  the IMAGE SET of  $\mathbf{L}$  in  $\mathbf{Y}^{\mathbf{X}}$ .

<b>T H M</b>	$\mathcal{L}(\mathbf{X}, \mathbf{Y})$	is a linear space	(space of linear transforms)
	$\mathcal{N}(\mathbf{L})$	is a linear subspace of $\mathbf{X}$	$\forall \mathbf{L} \in \mathbf{Y}^{\mathbf{X}}$
	$\mathcal{I}(\mathbf{L})$	is a linear subspace of $\mathbf{Y}$	$\forall \mathbf{L} \in \mathbf{Y}^{\mathbf{X}}$

$\Rightarrow$  PROOF:

1. Proof that  $\mathcal{N}(\mathbf{L})$  is a linear subspace of  $\mathbf{X}$ :

- (a)  $0 \in \mathcal{N}(\mathbf{L}) \implies \mathcal{N}(\mathbf{L}) \neq \emptyset$
- (b)  $\mathcal{N}(\mathbf{L}) \triangleq \{\mathbf{x} \in \mathbf{X} \mid \mathbf{L}\mathbf{x} = 0\} \subseteq \mathbf{X}$
- (c)  $\mathbf{x} + \mathbf{y} \in \mathcal{N}(\mathbf{L}) \implies 0 = \mathbf{L}(\mathbf{x} + \mathbf{y}) = \mathbf{L}(\mathbf{y} + \mathbf{x}) \implies \mathbf{y} + \mathbf{x} \in \mathcal{N}(\mathbf{L})$
- (d)  $\alpha \in \mathbb{F}, \mathbf{x} \in \mathbf{X} \implies 0 = \mathbf{L}\mathbf{x} \implies 0 = \alpha \mathbf{L}\mathbf{x} \implies 0 = \mathbf{L}(\alpha \mathbf{x}) \implies \alpha \mathbf{x} \in \mathcal{N}(\mathbf{L})$

2. Proof that  $\mathcal{I}(\mathbf{L})$  is a linear subspace of  $\mathbf{Y}$ :

- (a)  $0 \in \mathcal{I}(\mathbf{L}) \implies \mathcal{I}(\mathbf{L}) \neq \emptyset$
- (b)  $\mathcal{I}(\mathbf{L}) \triangleq \{\mathbf{y} \in \mathbf{Y} \mid \exists \mathbf{x} \in \mathbf{X} \text{ such that } \mathbf{y} = \mathbf{L}\mathbf{x}\} \subseteq \mathbf{Y}$
- (c)  $\mathbf{x} + \mathbf{y} \in \mathcal{I}(\mathbf{L}) \implies \exists \mathbf{v} \in \mathbf{X} \text{ such that } \mathbf{L}\mathbf{v} = \mathbf{x} + \mathbf{y} = \mathbf{y} + \mathbf{x} \implies \mathbf{y} + \mathbf{x} \in \mathcal{I}(\mathbf{L})$
- (d)  $\alpha \in \mathbb{F}, \mathbf{x} \in \mathcal{I}(\mathbf{L}) \implies \exists \mathbf{x} \in \mathbf{X} \text{ such that } \mathbf{y} = \mathbf{L}\mathbf{x} \implies \alpha \mathbf{y} = \alpha \mathbf{L}\mathbf{x} = \mathbf{L}(\alpha \mathbf{x}) \implies \alpha \mathbf{x} \in \mathcal{I}(\mathbf{L})$

$\Rightarrow$

**Example F.1.** <sup>8</sup> Let  $C([a : b], \mathbb{R})$  be the set of all continuous functions from the closed real interval  $[a : b]$  to  $\mathbb{R}$ .

**E  
X**  $C([a : b], \mathbb{R})$  is a linear space.

**Theorem F.3.** <sup>9</sup> Let  $\mathcal{L}(\mathbf{X}, \mathbf{Y})$  be the set of linear operators from a linear space  $\mathbf{X}$  to a linear space  $\mathbf{Y}$ . Let  $\mathcal{N}(\mathbf{L})$  be the NULL SPACE of a linear operator  $\mathbf{L} \in \mathcal{L}(\mathbf{X}, \mathbf{Y})$ .

<b>T H M</b>	$\mathbf{L}\mathbf{x} = \mathbf{L}\mathbf{y}$	$\iff$	$\mathbf{x} - \mathbf{y} \in \mathcal{N}(\mathbf{L})$
	$\mathbf{L}$ is INJECTIVE	$\iff$	$\mathcal{N}(\mathbf{L}) = \{0\}$

<sup>7</sup> Michel and Herget (1993) pages 98–104, Berberian (1961) pages 80–85 (Theorem IV.1.4 and Theorem IV.3.1)

<sup>8</sup> Eidelman et al. (2004) page 3

<sup>9</sup> Berberian (1961) page 88 (Theorem IV.1.4)

✎ PROOF:

1. Proof that  $\mathbf{L}x = \mathbf{L}y \implies x - y \in \mathcal{N}(\mathbf{L})$ :

$$\begin{aligned} \mathbf{L}(x - y) &= \mathbf{L}x - \mathbf{L}y && \text{by Theorem F.1 page 121} \\ &= \mathbf{0} && \text{by left hypothesis} \\ \implies x - y &\in \mathcal{N}(\mathbf{L}) && \text{by definition of null space} \end{aligned}$$

2. Proof that  $\mathbf{L}x = \mathbf{L}y \iff x - y \in \mathcal{N}(\mathbf{L})$ :

$$\begin{aligned} \mathbf{L}y &= \mathbf{L}y + \mathbf{0} && \text{by definition of linear space (Definition F.1 page 119)} \\ &= \mathbf{L}y + \mathbf{L}(x - y) && \text{by right hypothesis} \\ &= \mathbf{L}y + (\mathbf{L}x - \mathbf{L}y) && \text{by Theorem F.1 page 121} \\ &= (\mathbf{L}y - \mathbf{L}y) + \mathbf{L}x && \text{by associative and commutative properties (Definition F.1 page 119)} \\ &= \mathbf{L}x \end{aligned}$$

3. Proof that  $\mathbf{L}$  is *injective*  $\iff \mathcal{N}(\mathbf{L}) = \{\mathbf{0}\}$ :

$$\begin{aligned} \mathbf{L} \text{ is injective} &\iff \{(\mathbf{L}x = \mathbf{L}y \iff x = y) \mid \forall x, y \in X\} \\ &\iff \{[\mathbf{L}x - \mathbf{L}y = \mathbf{0} \iff (x - y) = \mathbf{0}] \mid \forall x, y \in X\} \\ &\iff \{[\mathbf{L}(x - y) = \mathbf{0} \iff (x - y) = \mathbf{0}] \mid \forall x, y \in X\} \\ &\iff \mathcal{N}(\mathbf{L}) = \{\mathbf{0}\} \end{aligned}$$

⇒

**Theorem F.4.** <sup>10</sup> Let  $\mathcal{W}$ ,  $\mathcal{X}$ ,  $\mathcal{Y}$ , and  $\mathcal{Z}$  be linear spaces over a field  $\mathbb{F}$ .

<b>T H M</b>	1. $\mathbf{L}(\mathbf{M}\mathbf{N}) = (\mathbf{L}\mathbf{M})\mathbf{N}$	$\forall \mathbf{L} \in \mathcal{L}(\mathcal{Z}, \mathcal{W}), \mathbf{M} \in \mathcal{L}(\mathcal{Y}, \mathcal{Z}), \mathbf{N} \in \mathcal{L}(\mathcal{X}, \mathcal{Y})$	(ASSOCIATIVE)
	2. $\mathbf{L}(\mathbf{M} \dot{+} \mathbf{N}) = (\mathbf{L}\mathbf{M}) \dot{+} (\mathbf{L}\mathbf{N})$	$\forall \mathbf{L} \in \mathcal{L}(\mathcal{Y}, \mathcal{Z}), \mathbf{M} \in \mathcal{L}(\mathcal{X}, \mathcal{Y}), \mathbf{N} \in \mathcal{L}(\mathcal{X}, \mathcal{Y})$	(LEFT DISTRIBUTIVE)
	3. $(\mathbf{L} \dot{+} \mathbf{M})\mathbf{N} = (\mathbf{L}\mathbf{N}) \dot{+} (\mathbf{M}\mathbf{N})$	$\forall \mathbf{L} \in \mathcal{L}(\mathcal{Y}, \mathcal{Z}), \mathbf{M} \in \mathcal{L}(\mathcal{Y}, \mathcal{Z}), \mathbf{N} \in \mathcal{L}(\mathcal{X}, \mathcal{Y})$	(RIGHT DISTRIBUTIVE)
	4. $\alpha(\mathbf{L}\mathbf{M}) = (\alpha\mathbf{L})\mathbf{M} = \mathbf{L}(\alpha\mathbf{M})$	$\forall \mathbf{L} \in \mathcal{L}(\mathcal{Y}, \mathcal{Z}), \mathbf{M} \in \mathcal{L}(\mathcal{X}, \mathcal{Y}), \alpha \in \mathbb{F}$	(HOMOGENEOUS)

✎ PROOF:

1. Proof that  $\mathbf{L}(\mathbf{M}\mathbf{N}) = (\mathbf{L}\mathbf{M})\mathbf{N}$ : Follows directly from property of *associative* operators.

2. Proof that  $\mathbf{L}(\mathbf{M} \dot{+} \mathbf{N}) = (\mathbf{L}\mathbf{M}) \dot{+} (\mathbf{L}\mathbf{N})$ :

$$\begin{aligned} [\mathbf{L}(\mathbf{M} \dot{+} \mathbf{N})]x &= \mathbf{L}[(\mathbf{M} \dot{+} \mathbf{N})x] \\ &= \mathbf{L}[(\mathbf{M}x) \dot{+} (\mathbf{N}x)] \\ &= [\mathbf{L}(\mathbf{M}x)] \dot{+} [\mathbf{L}(\mathbf{N}x)] && \text{by additive property Definition F.4 page 121} \\ &= [(\mathbf{L}\mathbf{M})x] \dot{+} [(\mathbf{L}\mathbf{N})x] \end{aligned}$$

3. Proof that  $(\mathbf{L} \dot{+} \mathbf{M})\mathbf{N} = (\mathbf{L}\mathbf{N}) \dot{+} (\mathbf{M}\mathbf{N})$ : Follows directly from property of *associative* operators.

4. Proof that  $\alpha(\mathbf{L}\mathbf{M}) = (\alpha\mathbf{L})\mathbf{M}$ : Follows directly from *associative* property of linear operators.

5. Proof that  $\alpha(\mathbf{L}\mathbf{M}) = \mathbf{L}(\alpha\mathbf{M})$ :

$$\begin{aligned} [\alpha(\mathbf{L}\mathbf{M})]x &= \alpha[(\mathbf{L}\mathbf{M})x] \\ &= \mathbf{L}[\alpha(\mathbf{M}x)] && \text{by homogeneous property Definition F.4 page 121} \\ &= \mathbf{L}[(\alpha\mathbf{M})x] \\ &= [\mathbf{L}(\alpha\mathbf{M})]x \end{aligned}$$

<sup>10</sup> Berberian (1961) page 88 (Theorem IV.5.1)



**Theorem F.5** (Fundamental theorem of linear equations). [Michel and Herget \(1993\) page 99](#) Let  $\mathcal{Y}^{\mathcal{X}}$  be the set of all operators from a linear space  $\mathcal{X}$  to a linear space  $\mathcal{Y}$ . Let  $\mathcal{N}(\mathbf{L})$  be the NULL SPACE of an operator  $\mathbf{L}$  in  $\mathcal{Y}^{\mathcal{X}}$  and  $\mathcal{I}(\mathbf{L})$  the IMAGE SET of  $\mathbf{L}$  in  $\mathcal{Y}^{\mathcal{X}}$  (Definition ?? page ??).

$$\text{THM} \quad \dim \mathcal{I}(\mathbf{L}) + \dim \mathcal{N}(\mathbf{L}) = \dim \mathcal{X} \quad \forall \mathbf{L} \in \mathcal{Y}^{\mathcal{X}}$$

**PROOF:** Let  $\{\psi_k | k = 1, 2, \dots, p\}$  be a basis for  $\mathcal{X}$  constructed such that  $\{\psi_{p-n+1}, \psi_{p-n+2}, \dots, \psi_p\}$  is a basis for  $\mathcal{N}(\mathbf{L})$ .

Let  $p \triangleq \dim \mathcal{X}$ .

Let  $n \triangleq \dim \mathcal{N}(\mathbf{L})$ .

$$\begin{aligned} \dim \mathcal{I}(\mathbf{L}) &= \dim \{y \in \mathcal{Y} | \exists x \in \mathcal{X} \text{ such that } y = \mathbf{L}x\} \\ &= \dim \left\{ y \in \mathcal{Y} | \exists (\alpha_1, \alpha_2, \dots, \alpha_p) \text{ such that } y = \mathbf{L} \sum_{k=1}^p \alpha_k \psi_k \right\} \\ &= \dim \left\{ y \in \mathcal{Y} | \exists (\alpha_1, \alpha_2, \dots, \alpha_p) \text{ such that } y = \sum_{k=1}^p \alpha_k \mathbf{L} \psi_k \right\} \\ &= \dim \left\{ y \in \mathcal{Y} | \exists (\alpha_1, \alpha_2, \dots, \alpha_p) \text{ such that } y = \sum_{k=1}^{p-n} \alpha_k \mathbf{L} \psi_k + \sum_{k=1}^n \alpha_k \mathbf{L} \psi_k \right\} \\ &= \dim \left\{ y \in \mathcal{Y} | \exists (\alpha_1, \alpha_2, \dots, \alpha_p) \text{ such that } y = \sum_{k=1}^{p-n} \alpha_k \mathbf{L} \psi_k + \mathbf{0} \right\} \\ &= p - n \\ &= \dim \mathcal{X} - \dim \mathcal{N}(\mathbf{L}) \end{aligned}$$

Note: This “proof” may be missing some necessary detail.

## F.2 Operators on Normed linear spaces

### F.2.1 Operator norm

**Definition F.5.** <sup>11</sup> Let  $\mathcal{V} = (\mathcal{X}, \mathbb{F}, \hat{+}, \cdot)$  be a linear space and  $\mathbb{F}$  be a field with absolute value function  $|\cdot| \in \mathbb{R}^{\mathbb{F}}$  (Definition A.4 page 68).

A **norm** is any functional  $\|\cdot\|$  in  $\mathbb{R}^{\mathcal{X}}$  that satisfies

- |    |  |  |                                    |     |
|----|--|--|------------------------------------|-----|
| 1. | $\ \mathbf{x}\  \geq 0$  | $\forall \mathbf{x} \in \mathcal{X}$                   | (STRICTLY POSITIVE)                | and |
| 2. | $\ \mathbf{x}\  = 0 \iff \mathbf{x} = \mathbf{0}$                  | $\forall \mathbf{x} \in \mathcal{X}$                   | (NONDEGENERATE)                    | and |
| 3. | $\ a\mathbf{x}\  =  a  \ \mathbf{x}\ $                             | $\forall \mathbf{x} \in \mathcal{X}, a \in \mathbb{C}$ | (HOMOGENEOUS)                      | and |
| 4. | $\ \mathbf{x} + \mathbf{y}\  \leq \ \mathbf{x}\  + \ \mathbf{y}\ $ | $\forall \mathbf{x}, \mathbf{y} \in \mathcal{X}$       | (SUBADDITIVE/triangle inequality). |     |

A **normed linear space** is the pair  $(\mathcal{V}, \|\cdot\|)$ .

<sup>11</sup> [Aliprantis and Burkinshaw \(1998\) pages 217–218](#), [Banach \(1932a\) page 53](#), [Banach \(1932b\) page 33](#), [Banach \(1922\) page 135](#)

**Definition F.6.**<sup>12</sup> Let  $\mathcal{L}(\mathbf{X}, \mathbf{Y})$  be the space of linear operators over normed linear spaces  $\mathbf{X}$  and  $\mathbf{Y}$ .

DEF

The **operator norm**  $\|\cdot\|$  is defined as

$$\|\mathbf{A}\| \triangleq \sup_{\mathbf{x} \in \mathbf{X}} \{ \|\mathbf{A}\mathbf{x}\| \mid \|\mathbf{x}\| \leq 1 \} \quad \forall \mathbf{A} \in \mathcal{L}(\mathbf{X}, \mathbf{Y})$$

The pair  $(\mathcal{L}(\mathbf{X}, \mathbf{Y}), \|\cdot\|)$  is the **normed space of linear operators** on  $(\mathbf{X}, \mathbf{Y})$ .

Proposition F.1 (next) shows that the functional defined in Definition F.6 (previous) is a *norm* (Definition F.5 page 124).

**Proposition F.1.**<sup>14</sup> Let  $(\mathcal{L}(\mathbf{X}, \mathbf{Y}), \|\cdot\|)$  be the normed space of linear operators over the normed linear spaces  $\mathbf{X} \triangleq (\mathbf{X}, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \|\cdot\|)$  and  $\mathbf{Y} \triangleq (\mathbf{Y}, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \|\cdot\|)$ .

PRP

The functional  $\|\cdot\|$  is a **norm** on  $\mathcal{L}(\mathbf{X}, \mathbf{Y})$ . In particular,

- |    |  |   |                 |     |
|----|--|---|-----------------|-----|
| 1. | $\ \mathbf{A}\  \geq 0$  | $\forall \mathbf{A} \in \mathcal{L}(\mathbf{X}, \mathbf{Y})$                        | (NON-NEGATIVE)  | and |
| 2. | $\ \mathbf{A}\  = 0 \iff \mathbf{A} \doteq \mathbf{0}$                   | $\forall \mathbf{A} \in \mathcal{L}(\mathbf{X}, \mathbf{Y})$                        | (NONDEGENERATE) | and |
| 3. | $\ \alpha \mathbf{A}\  =  \alpha  \ \mathbf{A}\ $                        | $\forall \mathbf{A} \in \mathcal{L}(\mathbf{X}, \mathbf{Y}), \alpha \in \mathbb{F}$ | (HOMOGENEOUS)   | and |
| 4. | $\ \mathbf{A} \dot{+} \mathbf{B}\  \leq \ \mathbf{A}\  + \ \mathbf{B}\ $ | $\forall \mathbf{A} \in \mathcal{L}(\mathbf{X}, \mathbf{Y})$                        | (SUBADDITIVE).  |     |

Moreover,  $(\mathcal{L}(\mathbf{X}, \mathbf{Y}), \|\cdot\|)$  is a **normed linear space**.

**PROOF:**

1. Proof that  $\|\mathbf{A}\| > 0$  for  $\mathbf{A} \neq \mathbf{0}$ :

$$\begin{aligned} \|\mathbf{A}\| &\triangleq \sup_{\mathbf{x} \in \mathbf{X}} \{ \|\mathbf{A}\mathbf{x}\| \mid \|\mathbf{x}\| \leq 1 \} \\ &> 0 \end{aligned}$$

by definition of  $\|\cdot\|$  (Definition F.6 page 125)

2. Proof that  $\|\mathbf{A}\| = 0$  for  $\mathbf{A} \doteq \mathbf{0}$ :

$$\begin{aligned} \|\mathbf{A}\| &\triangleq \sup_{\mathbf{x} \in \mathbf{X}} \{ \|\mathbf{A}\mathbf{x}\| \mid \|\mathbf{x}\| \leq 1 \} \\ &= \sup_{\mathbf{x} \in \mathbf{X}} \{ \|\mathbf{0}\mathbf{x}\| \mid \|\mathbf{x}\| \leq 1 \} \\ &= 0 \end{aligned}$$

by definition of  $\|\cdot\|$  (Definition F.6 page 125)

3. Proof that  $\|\alpha \mathbf{A}\| = |\alpha| \|\mathbf{A}\|$ :

$$\begin{aligned} \|\alpha \mathbf{A}\| &\triangleq \sup_{\mathbf{x} \in \mathbf{X}} \{ \|\alpha \mathbf{A}\mathbf{x}\| \mid \|\mathbf{x}\| \leq 1 \} \\ &= \sup_{\mathbf{x} \in \mathbf{X}} \{ |\alpha| \|\mathbf{A}\mathbf{x}\| \mid \|\mathbf{x}\| \leq 1 \} \\ &= |\alpha| \sup_{\mathbf{x} \in \mathbf{X}} \{ \|\mathbf{A}\mathbf{x}\| \mid \|\mathbf{x}\| \leq 1 \} \\ &= |\alpha| \|\mathbf{A}\| \end{aligned}$$

by definition of  $\|\cdot\|$  (Definition F.6 page 125)

by definition of  $\|\cdot\|$  (Definition F.6 page 125)

by definition of sup

by definition of  $\|\cdot\|$  (Definition F.6 page 125)

<sup>12</sup> Rudin (1991) page 92, Aliprantis and Burkinshaw (1998) page 225

<sup>13</sup> The operator norm notation  $\|\cdot\|$  is introduced (as a Matrix norm) in

Horn and Johnson (1990) page 290

<sup>14</sup> Rudin (1991) page 93

4. Proof that  $\|A \dot{+} B\| \leq \|A\| + \|B\|$ :

$$\begin{aligned}
 \|A \dot{+} B\| &\triangleq \sup_{x \in X} \{ \|(A \dot{+} B)x\| \mid \|x\| \leq 1 \} && \text{by definition of } \|\cdot\| \text{ (Definition F.6 page 125)} \\
 &= \sup_{x \in X} \{ \|Ax + Bx\| \mid \|x\| \leq 1 \} \\
 &\leq \sup_{x \in X} \{ \|Ax\| + \|Bx\| \mid \|x\| \leq 1 \} && \text{by definition of } \|\cdot\| \text{ (Definition F.6 page 125)} \\
 &\leq \sup_{x \in X} \{ \|Ax\| \mid \|x\| \leq 1 \} + \sup_{x \in X} \{ \|Bx\| \mid \|x\| \leq 1 \} \\
 &\triangleq \|A\| + \|B\| && \text{by definition of } \|\cdot\| \text{ (Definition F.6 page 125)}
 \end{aligned}$$

⇒

**Lemma F.1.** Let  $(\mathcal{L}(X, Y), \|\cdot\|)$  be the normed space of linear operators over normed linear spaces  $X \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \|\cdot\|)$  and  $Y \triangleq (Y, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \|\cdot\|)$ .

L E M	$\ L\  = \sup_x \{ \ Lx\  \mid \ x\  = 1 \} \quad \forall x \in \mathcal{L}(X, Y)$
-------------	--

PROOF: 15

1. Proof that  $\sup_x \{ \|Lx\| \mid \|x\| \leq 1 \} \geq \sup_x \{ \|Lx\| \mid \|x\| = 1 \}$ :

$$\sup_x \{ \|Lx\| \mid \|x\| \leq 1 \} \geq \sup_x \{ \|Lx\| \mid \|x\| = 1 \} \quad \text{because } A \subseteq B \implies \sup A \leq \sup B$$

2. Let the subset  $Y \subsetneq X$  be defined as

$$Y \triangleq \left\{ y \in X \mid \begin{array}{l} 1. \ \|Ly\| = \sup_{x \in X} \{ \|Lx\| \mid \|x\| \leq 1 \} \text{ and} \\ 2. \ 0 < \|y\| \leq 1 \end{array} \right\}$$

3. Proof that  $\sup_x \{ \|Lx\| \mid \|x\| \leq 1 \} \leq \sup_x \{ \|Lx\| \mid \|x\| = 1 \}$ :

$$\begin{aligned}
 \sup_{x \in X} \{ \|Lx\| \mid \|x\| \leq 1 \} &= \|Ly\| && \text{by definition of set } Y \\
 &= \frac{\|y\|}{\|y\|} \|Ly\| \\
 &= \|y\| \left\| \frac{1}{\|y\|} Ly \right\| && \text{by homogeneous property (page 124)} \\
 &= \|y\| \left\| L \frac{y}{\|y\|} \right\| && \text{by homogeneous property (page 121)} \\
 &\leq \|y\| \sup_{y \in Y} \left\{ \left\| L \frac{y}{\|y\|} \right\| \right\} && \text{by definition of supremum} \\
 &= \|y\| \sup_{y \in Y} \left\{ \left\| L \frac{y}{\|y\|} \right\| \mid \left\| \frac{y}{\|y\|} \right\| = 1 \right\} && \text{because } \left\| \frac{y}{\|y\|} \right\| = 1 \text{ for all } y \in Y \\
 &\leq \sup_{y \in Y} \left\{ \left\| L \frac{y}{\|y\|} \right\| \mid \left\| \frac{y}{\|y\|} \right\| = 1 \right\} && \text{because } 0 < \|y\| \leq 1 \\
 &\leq \sup_{x \in X} \{ \|Lx\| \mid \|x\| = 1 \} && \text{because } \frac{y}{\|y\|} \in X \quad \forall y \in Y
 \end{aligned}$$

15

email



Many many thanks to former NCTU Ph.D. student [Chien Yao](#) (Chinese: 姚建; PinYin: Yáo Jiàn) for his brilliant help with this proof. (If you are viewing this text as a pdf file, zoom in on the figure to the left to see text from Chien Yao's 2007 April 16 email.)



4. By (1) and (3),

$$\sup_{x \in X} \{ \|Lx\| \mid \|x\| \leq 1 \} = \sup_{x \in X} \{ \|Lx\| \mid \|x\| = 1 \}$$

⇒

**Proposition F.2.** <sup>16</sup> Let  $I$  be the identity operator in the normed space of linear operators  $(\mathcal{L}(X, X), \|\cdot\|)$ .

P R P	$\ I\  = 1$
-------------	-------------

✎ PROOF:

$$\begin{aligned} \|I\| &\triangleq \sup \{ \|Ix\| \mid \|x\| \leq 1 \} && \text{by definition of } \|\cdot\| \text{ (Definition F.6 page 125)} \\ &= \sup \{ \|x\| \mid \|x\| \leq 1 \} && \text{by definition of } I \text{ (Definition F.3 page 120)} \\ &= 1 \end{aligned}$$

⇒

**Theorem F.6.** <sup>17</sup> Let  $(\mathcal{L}(X, Y), \|\cdot\|)$  be the normed space of linear operators over normed linear spaces  $X$  and  $Y$ .

T H M	$\ Lx\  \leq \ L\  \ x\  \quad \forall L \in \mathcal{L}(X, Y), x \in X$ $\ KL\  \leq \ K\  \ L\  \quad \forall K, L \in \mathcal{L}(X, Y)$
-------------	--

✎ PROOF:

1. Proof that  $\|Lx\| \leq \|L\| \|x\|$ :

$$\begin{aligned} \|Lx\| &= \frac{\|x\|}{\|x\|} \|Lx\| \\ &= \|x\| \left\| \frac{1}{\|x\|} Lx \right\| && \text{by property of norms} \\ &= \|x\| \left\| L \frac{x}{\|x\|} \right\| && \text{by property of linear operators} \\ &\triangleq \|x\| \|Ly\| && \text{where } y \triangleq \frac{x}{\|x\|} \\ &\leq \|x\| \sup_y \|Ly\| && \text{by definition of supremum} \\ &= \|x\| \sup_y \{ \|Ly\| \mid \|y\| = 1 \} && \text{because } \|y\| = \left\| \frac{x}{\|x\|} \right\| = \frac{\|x\|}{\|x\|} = 1 \\ &\triangleq \|x\| \|L\| && \text{by definition of operator norm} \end{aligned}$$

<sup>16</sup> Michel and Herget (1993) page 410

<sup>17</sup> Rudin (1991) page 103, Aliprantis and Burkinshaw (1998) page 225

2. Proof that  $\|KL\| \leq \|K\| \|L\|$ :

$$\begin{aligned}
 \|KL\| &\triangleq \sup_{x \in X} \{ \|(KL)x\| \mid \|x\| \leq 1 \} && \text{by Definition F.6 page 125 } (\|\cdot\|) \\
 &= \sup_{x \in X} \{ \|K(Lx)\| \mid \|x\| \leq 1 \} \\
 &\leq \sup_{x \in X} \{ \|K\| \|Lx\| \mid \|x\| \leq 1 \} && \text{by 1.} \\
 &\leq \sup_{x \in X} \{ \|K\| \|L\| \|x\| \mid \|x\| \leq 1 \} && \text{by 1.} \\
 &= \sup_{x \in X} \{ \|K\| \|L\| 1 \mid \|x\| \leq 1 \} && \text{by definition of sup} \\
 &= \|K\| \|L\| && \text{by definition of sup}
 \end{aligned}$$

⇒

## F.2.2 Bounded linear operators

**Definition F.7.** <sup>18</sup> Let  $(\mathcal{L}(X, Y), \|\cdot\|)$  be a normed space of linear operators.

**DEF** An operator  $B$  is **bounded** if  $\|B\| < \infty$ .  
 The quantity  $B(X, Y)$  is the set of all **bounded linear operators** on  $(X, Y)$  such that  
 $B(X, Y) \triangleq \{L \in \mathcal{L}(X, Y) \mid \|L\| < \infty\}$ .

**Theorem F.7.** <sup>19</sup> Let  $(\mathcal{L}(X, Y), \|\cdot\|)$  be the set of linear operators over normed linear spaces  $X \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \|\cdot\|)$  and  $Y \triangleq (Y, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \|\cdot\|)$ .

The following conditions are all EQUIVALENT:


- |             |   |  |        |
|-------------|---|--|--------|
| T<br>H<br>M | 1. $L$ is continuous at a SINGLE POINT $x_0 \in X$            | $\forall L \in \mathcal{L}(X, Y)$          | $\iff$ |
|             | 2. $L$ is CONTINUOUS (at every point $x \in X$ )              | $\forall L \in \mathcal{L}(X, Y)$          | $\iff$ |
|             | 3. $\ L\  < \infty$ ( $L$ is BOUNDED)                         | $\forall L \in \mathcal{L}(X, Y)$          | $\iff$ |
|             | 4. $\exists M \in \mathbb{R}$ such that $\ Lx\  \leq M \ x\ $ | $\forall L \in \mathcal{L}(X, Y), x \in X$ |        |


✎ PROOF:

1. Proof that 1  $\implies$  2:

$$\begin{aligned}
 \epsilon &> \|Lx - Lx_0\| && \text{by hypothesis 1} \\
 &= \|L(x - x_0)\| && \text{by linearity (Definition F.4 page 121)} \\
 &= \|L(x + y - x_0 - y)\| \\
 &= \|L(x + y) - L(x_0 + y)\| && \text{by linearity (Definition F.4 page 121)} \\
 \implies L &\text{ is continuous at point } x + y \\
 \implies L &\text{ is continuous at every point in } X && \text{(hypothesis 2)}
 \end{aligned}$$

2. Proof that 2  $\implies$  1: obvious:

<sup>18</sup>  Rudin (1991) pages 92–93

<sup>19</sup>  Aliprantis and Burkinshaw (1998) page 227

3. Proof that 4  $\implies$  2:<sup>20</sup>

$$\begin{aligned}
 \|Lx\| &\leq M \|x\| \implies \|L(x-y)\| \leq M \|x-y\| && \text{by hypothesis 4} \\
 &\implies \|Lx - Ly\| \leq M \|x-y\| && \text{by linearity of } L \text{ (Definition F.4 page 121)} \\
 &\implies \|Lx - Ly\| \leq \epsilon \text{ whenever } M \|x-y\| < \epsilon \\
 &\implies \|Lx - Ly\| \leq \epsilon \text{ whenever } \|x-y\| < \frac{\epsilon}{M} \quad (\text{hypothesis 2})
 \end{aligned}$$

4. Proof that 3  $\implies$  4:

$$\begin{aligned}
 \|Lx\| &\leq \underbrace{\|L\|}_{M} \|x\| && \text{by Theorem F6 page 127} \\
 &= M \|x\| && \text{where } M \triangleq \|L\| < \infty \text{ (by hypothesis 1)}
 \end{aligned}$$

5. Proof that 1  $\implies$  3:<sup>21</sup>

$$\begin{aligned}
 \|L\| = \infty &\implies \{\|Lx\| \mid \|x\| \leq 1\} = \infty \\
 &\implies \exists (x_n) \text{ such that } \|x_n\| = 1 \text{ and } \|L\| = \{\|Lx_n\| \mid \|x_n\| \leq 1\} = \infty \\
 &\implies \|x_n\| = 1 \text{ and } \infty = \|L\| = \|Lx_n\| \\
 &\implies \|x_n\| = 1 \text{ and } \|Lx_n\| \geq n \\
 &\implies \frac{1}{n} \|x_n\| = \frac{1}{n} \text{ and } \frac{1}{n} \|Lx_n\| \geq 1 \\
 &\implies \left\| \frac{x_n}{n} \right\| = \frac{1}{n} \text{ and } \left\| L \frac{x_n}{n} \right\| \geq 1 \\
 &\implies \lim_{n \rightarrow \infty} \left\| \frac{x_n}{n} \right\| = 0 \text{ and } \lim_{n \rightarrow \infty} \left\| L \frac{x_n}{n} \right\| \geq 1 \\
 &\implies L \text{ is not continuous at } 0
 \end{aligned}$$

But by hypothesis,  $L$  is continuous. So the statement  $\|L\| = \infty$  must be *false* and thus  $\|L\| < \infty$  ( $L$  is *bounded*).



## F.2.3 Adjoints on normed linear spaces

**Definition F8.** Let  $B(X, Y)$  be the space of BOUNDED LINEAR OPERATORS on normed linear spaces  $X$  and  $Y$ . Let  $X^*$  be the TOPOLOGICAL DUAL SPACE of  $X$ .

**DEF**  $B^*$  is the **adjoint** of an operator  $B \in B(X, Y)$  if

$$f(Bx) = [B^*f](x) \quad \forall f \in X^*, x \in X$$

**Theorem F8.**<sup>22</sup> Let  $B(X, Y)$  be the space of BOUNDED LINEAR OPERATORS on NORMED LINEAR SPACES  $X$  and  $Y$ .

<b>T H M</b>	$(A + B)^*$	$= A^* + B^*$	$\forall A, B \in B(X, Y)$
	$(\lambda A)^*$	$= \lambda A^*$	$\forall A, B \in B(X, Y)$
	$(AB)^*$	$= B^* A^*$	$\forall A, B \in B(X, Y)$

<sup>20</sup> Bollobás (1999) page 29

<sup>21</sup> Aliprantis and Burkinshaw (1998) page 227

<sup>22</sup> Bollobás (1999) page 156

✎ PROOF:

$$\begin{aligned}
 [A \dot{+} B]^* f(x) &= f([A \dot{+} B]x) && \text{by definition of adjoint} && (\text{Definition F.8 page 129}) \\
 &= f(Ax + Bx) && \text{by definition of linear operators} && (\text{Definition F.4 page 121}) \\
 &= f(Ax) + f(Bx) && \text{by definition of linear functional} && \\
 &= A^*f(x) + B^*f(x) && \text{by definition of adjoint} && (\text{Definition F.8 page 129}) \\
 &= [A^* + B^*]f(x) && \text{by definition of linear functional} && 
 \end{aligned}$$

$$\begin{aligned}
 [\lambda A]^* f(x) &= f([\lambda A]x) && \text{by definition of adjoint} && (\text{Definition F.8 page 129}) \\
 &= \lambda f(Ax) && \text{by definition of linear functional} && \\
 &= [\lambda A^*]f(x) && \text{by definition of adjoint} && (\text{Definition F.8 page 129})
 \end{aligned}$$

$$\begin{aligned}
 [AB]^* f(x) &= f([AB]x) && \text{by definition of adjoint} && (\text{Definition F.8 page 129}) \\
 &= f(A[Bx]) && \text{by definition of linear operators} && (\text{Definition F.4 page 121}) \\
 &= [A^*f](Bx) && \text{by definition of adjoint} && (\text{Definition F.8 page 129}) \\
 &= B^*[A^*f](x) && \text{by definition of adjoint} && (\text{Definition F.8 page 129}) \\
 &= [B^*A^*]f(x) && \text{by definition of adjoint} && (\text{Definition F.8 page 129})
 \end{aligned}$$

⇒

**Theorem F.9.** <sup>23</sup> Let  $B(X, Y)$  be the space of BOUNDED LINEAR OPERATORS on normed linear spaces  $X$  and  $Y$ . Let  $B^*$  be the adjoint of an operator  $B$ .


T H M	$   B    =    B^*    \quad \forall B \in B(X, Y)$
-------------	---

✎ PROOF:

$$\begin{aligned}
 |||B||| &\triangleq \sup \{ \|Bx\| \mid \|x\| \leq 1 \} && \text{by Definition F.6 page 125} \\
 &\stackrel{?}{=} \sup \{ \|g(Bx; y^*)\| \mid \|x\| \leq 1, \|y^*\| \leq 1 \} \\
 &= \sup \{ \|f(x; B^*y^*)\| \mid \|x\| \leq 1, \|y^*\| \leq 1 \} \\
 &\triangleq \sup \{ \|B^*y^*\| \mid \|x\| \leq 1, \|y^*\| \leq 1 \} \\
 &= \sup \{ \|B^*y^*\| \mid \|y^*\| \leq 1 \} \\
 &\triangleq |||B^*||| && \text{by Definition F.6 page 125}
 \end{aligned}$$

⇒

## F.2.4 More properties



*“Beginning with the third year of studies, most of my mathematical work was really started in conversations with Mazur and Banach. And according to Banach some of my own contributions were characterized by a certain “strangeness” in the formulation of problems and in the outline of possible proofs. As he told me once some years later, he was surprised how often these “strange” approaches really worked.”*

Stanislaus M. Ulam (1909–1984), Polish mathematician <sup>24</sup>

<sup>23</sup> Rudin (1991) page 98

**Theorem F.10** (Mazur-Ulam theorem).<sup>25</sup> Let  $\phi \in \mathcal{L}(X, Y)$  be a function on normed linear spaces  $(X, \|\cdot\|_X)$  and  $(Y, \|\cdot\|_Y)$ . Let  $I \in \mathcal{L}(X, X)$  be the identity operator on  $(X, \|\cdot\|_X)$ .

T H M	$  \left. \begin{array}{l}  1. \underbrace{\phi^{-1}\phi = \phi\phi^{-1} = I}_{\text{bijective}} \quad \text{and} \\  2. \underbrace{\ \phi x - \phi y\ _Y = \ x - y\ _X}_{\text{isometric}} \quad \forall x, y \in X  \end{array} \right\} \implies \underbrace{\phi([1 - \lambda]x + \lambda y) = [1 - \lambda]\phi x + \lambda\phi y}_{\text{affine}} \quad \forall \lambda \in \mathbb{R}  $
-------------	--

PROOF: Proof not yet complete.

1. Let  $\psi$  be the *reflection* of  $z$  in  $X$  such that  $\psi x = 2z - x$

(a)  $\|\psi x - z\| = \|x - z\|$

2. Let  $\lambda \triangleq \sup_g \{\|gz - z\|\}$

3. Proof that  $g \in W \implies g^{-1} \in W$ :

Let  $\hat{x} \triangleq g^{-1}x$  and  $\hat{y} \triangleq g^{-1}y$ .

$\ g^{-1}x - g^{-1}y\ $	by definition of $\hat{x}$ and $\hat{y}$
$= \ \hat{x} - \hat{y}\ $	by left hypothesis
$= \ g\hat{x} - g\hat{y}\ $	by definition of $\hat{x}$ and $\hat{y}$
$= \ gg^{-1}x - gg^{-1}y\ $	by definition of $g^{-1}$
$= \ x - y\ $	

4. Proof that  $gz = z$ :

$2\lambda = 2 \sup \{\ gz - z\ \}$	by definition of $\lambda$ item (2)
$\leq 2 \ gz - z\ $	by definition of sup
$= \ 2z - 2gz\ $	
$= \ \psi gz - gz\ $	by definition of $\psi$ item (1)
$= \ g^{-1}\psi gz - g^{-1}gz\ $	by item (3)
$= \ g^{-1}\psi gz - z\ $	by definition of $g^{-1}$
$= \ \psi g^{-1}\psi gz - z\ $	
$= \ g^*z - z\ $	
$\leq \lambda$	by definition of $\lambda$ item (2)
$\implies 2\lambda \leq \lambda$	
$\implies \lambda = 0$	
$\implies gz = z$	

5. Proof that  $\phi\left(\frac{1}{2}x + \frac{1}{2}y\right) = \frac{1}{2}\phi x + \frac{1}{2}\phi y$ :

$$\begin{aligned}
 \phi\left(\frac{1}{2}x + \frac{1}{2}y\right) &= \\
 &= \frac{1}{2}\phi x + \frac{1}{2}\phi y
 \end{aligned}$$

<sup>24</sup> quote: [Ulam \(1991\)](#) page 33

image: <http://www-history.mcs.st-andrews.ac.uk/Biographies/Ulam.html>

<sup>25</sup> [Oikherberg and Rosenthal \(2007\)](#) page 598, [Väisälä \(2003\)](#) page 634, [Giles \(2000\)](#) page 11, [Dunford and Schwartz \(1957\)](#) page 91, [Mazur and Ulam \(1932\)](#)

6. Proof that  $\phi([1 - \lambda]\mathbf{x} + \lambda\mathbf{y}) = [1 - \lambda]\phi\mathbf{x} + \lambda\phi\mathbf{y}$ :

$$\begin{aligned}\phi([1 - \lambda]\mathbf{x} + \lambda\mathbf{y}) &= \\ &= [1 - \lambda]\phi\mathbf{x} + \lambda\phi\mathbf{y}\end{aligned}$$

⇒

**Theorem F.11** (Neumann Expansion Theorem).<sup>26</sup> Let  $\mathbf{A} \in \mathbf{X}^{\mathbf{X}}$  be an operator on a linear space  $\mathbf{X}$ . Let  $\mathbf{A}^0 \triangleq \mathbf{I}$ .

<b>T H M</b>	$\left. \begin{array}{l} 1. \quad \mathbf{A} \in \mathcal{B}(\mathbf{X}, \mathbf{X}) \quad (\mathbf{A} \text{ is bounded}) \\ 2. \quad \ \mathbf{A}\  < 1 \end{array} \right\} \Rightarrow \left\{ \begin{array}{l} 1. \quad (\mathbf{I} - \mathbf{A})^{-1} \text{ exists} \\ 2. \quad \ (\mathbf{I} - \mathbf{A})^{-1}\  \leq \frac{1}{1 - \ \mathbf{A}\ } \\ 3. \quad (\mathbf{I} - \mathbf{A})^{-1} = \sum_{n=0}^{\infty} \mathbf{A}^n \\ \text{with uniform convergence} \end{array} \right.$
----------------------	---

## F.3 Operators on Inner product spaces

### F.3.1 General Results

**Definition F.9.**<sup>27</sup> Let  $\Omega \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}))$  be a linear space.

A function  $\langle \triangle | \nabla \rangle \in \mathbb{F}^{X \times X}$  is an **inner product** on  $\Omega$  if

- |                      |    |  |   |                        |     |
|----------------------|----|--|---|------------------------|-----|
| <b>D<br/>E<br/>F</b> | 1. | $\langle \mathbf{x}   \mathbf{x} \rangle \geq 0$   | $\forall \mathbf{x} \in X$  | (non-negative)         | and |
|                      | 2. | $\langle \mathbf{x}   \mathbf{x} \rangle = 0 \iff \mathbf{x} = \mathbf{0}$   | $\forall \mathbf{x} \in X$  | (nondegenerate)        | and |
|                      | 3. | $\langle \alpha \mathbf{x}   \mathbf{y} \rangle = \alpha \langle \mathbf{x}   \mathbf{y} \rangle$  | $\forall \mathbf{x}, \mathbf{y} \in X, \forall \alpha \in \mathbb{C}$ | (homogeneous)          | and |
|                      | 4. | $\langle \mathbf{x} + \mathbf{y}   \mathbf{u} \rangle = \langle \mathbf{x}   \mathbf{u} \rangle + \langle \mathbf{y}   \mathbf{u} \rangle$ | $\forall \mathbf{x}, \mathbf{y}, \mathbf{u} \in X$                    | (additive)             | and |
|                      | 5. | $\langle \mathbf{x}   \mathbf{y} \rangle = \langle \mathbf{y}   \mathbf{x} \rangle^*$  | $\forall \mathbf{x}, \mathbf{y} \in X$                                | (conjugate symmetric). |     |

An inner product is also called a **scalar product**.

An **inner product space** is the pair  $(\Omega, \langle \triangle | \nabla \rangle)$ .

**Theorem F.12.**<sup>28</sup> Let  $\mathbf{A}, \mathbf{B} \in \mathcal{B}(\mathbf{X}, \mathbf{X})$  be BOUNDED LINEAR OPERATORS on an inner product space  $\mathbf{X} \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle | \nabla \rangle)$ .

<b>T H M</b>	$\langle \mathbf{B}\mathbf{x}   \mathbf{x} \rangle = 0 \quad \forall \mathbf{x} \in X \iff \mathbf{B}\mathbf{x} = \mathbf{0} \quad \forall \mathbf{x} \in X$
	$\langle \mathbf{A}\mathbf{x}   \mathbf{x} \rangle = \langle \mathbf{B}\mathbf{x}   \mathbf{x} \rangle \quad \forall \mathbf{x} \in X \iff \mathbf{A} = \mathbf{B}$

PROOF:

<sup>26</sup> Michel and Herget (1993) page 415

<sup>27</sup> Haaser and Sullivan (1991) page 277, Aliprantis and Burkinshaw (1998) page 276, Peano (1888b) page 72

<sup>28</sup> Rudin (1991) page 310 (Theorem 12.7, Corollary)

1. Proof that  $\langle \mathbf{B}\mathbf{x} | \mathbf{x} \rangle = 0 \implies \mathbf{B}\mathbf{x} = \mathbf{0}$ :

$$\begin{aligned}
 0 &= \langle \mathbf{B}(\mathbf{x} + \mathbf{B}\mathbf{x}) | (\mathbf{x} + \mathbf{B}\mathbf{x}) \rangle + i \langle \mathbf{B}(\mathbf{x} + i\mathbf{B}\mathbf{x}) | (\mathbf{x} + i\mathbf{B}\mathbf{x}) \rangle && \text{by left hypothesis} \\
 &= \{ \langle \mathbf{B}\mathbf{x} + \mathbf{B}^2\mathbf{x} | \mathbf{x} + \mathbf{B}\mathbf{x} \rangle \} + i \{ \langle \mathbf{B}\mathbf{x} + i\mathbf{B}^2\mathbf{x} | \mathbf{x} + i\mathbf{B}\mathbf{x} \rangle \} && \text{by Definition F.4 page 121} \\
 &= \{ \langle \mathbf{B}\mathbf{x} | \mathbf{x} \rangle + \langle \mathbf{B}\mathbf{x} | \mathbf{B}\mathbf{x} \rangle + \langle \mathbf{B}^2\mathbf{x} | \mathbf{x} \rangle + \langle \mathbf{B}^2\mathbf{x} | \mathbf{B}\mathbf{x} \rangle \} && \text{by Definition F.9 page 132} \\
 &\quad + i \{ \langle \mathbf{B}\mathbf{x} | \mathbf{x} \rangle - i \langle \mathbf{B}\mathbf{x} | \mathbf{B}\mathbf{x} \rangle + i \langle \mathbf{B}^2\mathbf{x} | \mathbf{x} \rangle - i^2 \langle \mathbf{B}^2\mathbf{x} | \mathbf{B}\mathbf{x} \rangle \} \\
 &= \{ 0 + \langle \mathbf{B}\mathbf{x} | \mathbf{B}\mathbf{x} \rangle + \langle \mathbf{B}^2\mathbf{x} | \mathbf{x} \rangle + 0 \} + i \{ 0 - i \langle \mathbf{B}\mathbf{x} | \mathbf{B}\mathbf{x} \rangle + i \langle \mathbf{B}^2\mathbf{x} | \mathbf{x} \rangle - i^2 0 \} && \text{by left hypothesis} \\
 &= \{ \langle \mathbf{B}\mathbf{x} | \mathbf{B}\mathbf{x} \rangle + \langle \mathbf{B}^2\mathbf{x} | \mathbf{x} \rangle \} + \{ \langle \mathbf{B}\mathbf{x} | \mathbf{B}\mathbf{x} \rangle - \langle \mathbf{B}^2\mathbf{x} | \mathbf{x} \rangle \} \\
 &= 2 \langle \mathbf{B}\mathbf{x} | \mathbf{B}\mathbf{x} \rangle \\
 &= 2 \|\mathbf{B}\mathbf{x}\|^2 \\
 &\implies \mathbf{B}\mathbf{x} = \mathbf{0} && \text{by Definition F.5 page 124}
 \end{aligned}$$

2. Proof that  $\langle \mathbf{B}\mathbf{x} | \mathbf{x} \rangle = 0 \iff \mathbf{B}\mathbf{x} = \mathbf{0}$ : by property of inner products.

3. Proof that  $\langle \mathbf{A}\mathbf{x} | \mathbf{x} \rangle = \langle \mathbf{B}\mathbf{x} | \mathbf{x} \rangle \implies \mathbf{A} \doteq \mathbf{B}$ :

$$\begin{aligned}
 0 &= \langle \mathbf{A}\mathbf{x} | \mathbf{x} \rangle - \langle \mathbf{B}\mathbf{x} | \mathbf{x} \rangle && \text{by left hypothesis} \\
 &= \langle \mathbf{A}\mathbf{x} - \mathbf{B}\mathbf{x} | \mathbf{x} \rangle && \text{by additivity property of } \langle \triangle | \nabla \rangle \text{ (Definition F.9 page 132)} \\
 &= \langle (\mathbf{A} - \mathbf{B})\mathbf{x} | \mathbf{x} \rangle && \text{by definition of operator addition} \\
 \implies (\mathbf{A} - \mathbf{B})\mathbf{x} &= \mathbf{0} && \text{by item 1} \\
 \implies \mathbf{A} &= \mathbf{B} && \text{by definition of operator subtraction}
 \end{aligned}$$

4. Proof that  $\langle \mathbf{A}\mathbf{x} | \mathbf{x} \rangle = \langle \mathbf{B}\mathbf{x} | \mathbf{x} \rangle \iff \mathbf{A} \doteq \mathbf{B}$ :

$$\langle \mathbf{A}\mathbf{x} | \mathbf{x} \rangle = \langle \mathbf{B}\mathbf{x} | \mathbf{x} \rangle \quad \text{by } \mathbf{A} \doteq \mathbf{B} \text{ hypothesis}$$



## F.3.2 Operator adjoint

A fundamental concept of operators on inner product spaces is the *operator adjoint* (Proposition F.3 page 133). The adjoint of an operator is a kind of generalization of the conjugate of a complex number in that

Both are *star-algebras* (Theorem F.13 page 134).

Both support decomposition into “real” and “imaginary” parts (Theorem E.3 page 116).

Structurally, the operator adjoint provides a convenient symmetric relationship between the *range space* and *null space* of an operator (Theorem F.14 page 135).

**Proposition F.3.** <sup>29</sup> Let  $B(H, H)$  be the space of BOUNDED LINEAR OPERATORS (Definition F.7 page 128) on a HILBERT SPACE  $H$ .

**P** An operator  $\mathbf{B}^*$  is the **adjoint** of  $\mathbf{B} \in B(H, H)$  if  
**P**  $\langle \mathbf{B}\mathbf{x} | \mathbf{y} \rangle = \langle \mathbf{x} | \mathbf{B}^*\mathbf{y} \rangle \quad \forall \mathbf{x}, \mathbf{y} \in H.$   
**P**

PROOF:

<sup>29</sup> Michel and Herget (1993) page 220, Rudin (1991) page 311, Giles (2000) page 182, von Neumann (1929) page 49, Stone (1932) page 41

1. For fixed  $y$ ,  $f(x) \triangleq \langle x | y \rangle$  is a *functional* in  $\mathbb{F}^X$ .
2.  $B^*$  is the *adjoint* of  $B$  because





$$\begin{aligned}
 \langle Bx | y \rangle &\triangleq f(Bx) \\
 &\triangleq B^*f(x) && \text{by definition of operator adjoint} && (\text{Definition F.8 page 129}) \\
 &= \langle x | B^*y \rangle
 \end{aligned}$$

⇒

*Example F.2.*

In matrix algebra (“linear algebra”)

E  
X

-  The inner product operation  $\langle x | y \rangle$  is represented by  $y^H x$ .
-  The linear operator is represented as a matrix  $A$ .
-  The operation of  $A$  on a vector  $x$  is represented as  $Ax$ .
-  The adjoint of matrix  $A$  is the Hermitian matrix  $A^H$ .

✎ PROOF:

$$\langle Ax | y \rangle \triangleq y^H Ax = [(Ax)^H y]^H = [x^H A^H y]^H = (A^H y)^H x \triangleq \langle x | A^H y \rangle$$

⇒

Structures that satisfy the four conditions of the next theorem are known as *\*-algebras* (“*star-algebras*” (Definition E.3 page 114). Other structures which are *\*-algebras* include the *field of complex numbers*  $\mathbb{C}$  and any *ring of complex square*  $n \times n$  *matrices*.<sup>30</sup>

**Theorem F.13** (operator star-algebra).<sup>31</sup> *Let  $H$  be a HILBERT SPACE with operators  $A, B \in \mathcal{B}(H, H)$  and with adjoints  $A^*, B^* \in \mathcal{B}(H, H)$ . Let  $\bar{\alpha}$  be the complex conjugate of some  $\alpha \in \mathbb{C}$ .*




*The pair  $(H, *)$  is a \*-ALGEBRA (STAR-ALGEBRA). In particular,*T  
H  
M

- |    |                                   |                      |                    |     |
|----|-----------------------------------|----------------------|--------------------|-----|
| 1. | $(A \dot{+} B)^* = A^* + B^*$     | $\forall A, B \in H$ | (DISTRIBUTIVE)     | and |
| 2. | $(\alpha A)^* = \bar{\alpha} A^*$ | $\forall A, B \in H$ | (CONJUGATE LINEAR) | and |
| 3. | $(AB)^* = B^* A^*$                | $\forall A, B \in H$ | (ANTIAUTOMORPHIC)  | and |
| 4. | $A^{**} = A$                      | $\forall A, B \in H$ | (INVOLUTARY)       |     |

✎ PROOF:

$$\begin{aligned}
 \langle x | (A \dot{+} B)^* y \rangle &= \langle (A \dot{+} B)x | y \rangle && \text{by definition of adjoint} && (\text{Proposition F.3 page 133}) \\
 &= \langle Ax | y \rangle + \langle Bx | y \rangle && \text{by definition of inner product} && (\text{Definition F.9 page 132}) \\
 &= \langle x | A^* y \rangle + \langle x | B^* y \rangle && \text{by definition of operator addition} \\
 &= \langle x | A^* y + B^* y \rangle && \text{by definition of inner product} && (\text{Definition F.9 page 132}) \\
 &= \langle x | (A^* + B^*) y \rangle && \text{by definition of operator addition}
 \end{aligned}$$

$$\begin{aligned}
 \langle x | (\alpha A)^* y \rangle &= \langle (\alpha A)x | y \rangle && \text{by definition of adjoint} && (\text{Proposition F.3 page 133}) \\
 &= \langle \alpha(Ax) | y \rangle && \text{by definition of scalar multiplication} \\
 &= \alpha \langle Ax | y \rangle && \text{by definition of inner product} && (\text{Definition F.9 page 132}) \\
 &= \alpha \langle x | A^* y \rangle && \text{by definition of adjoint} && (\text{Proposition F.3 page 133}) \\
 &= \langle x | \alpha^* A^* y \rangle && \text{by definition of inner product} && (\text{Definition F.9 page 132})
 \end{aligned}$$

<sup>30</sup>  Sakai (1998) page 1<sup>31</sup>  Halmos (1998) pages 39–40,  Rudin (1991) page 311



$\langle \mathbf{x}   (\mathbf{AB})^* \mathbf{y} \rangle = \langle (\mathbf{AB})\mathbf{x}   \mathbf{y} \rangle$	by definition of adjoint	(Proposition F.3 page 133)
$= \langle \mathbf{A}(\mathbf{B}\mathbf{x})   \mathbf{y} \rangle$	by definition of operator multiplication	
$= \langle (\mathbf{B}\mathbf{x})   \mathbf{A}^* \mathbf{y} \rangle$	by definition of adjoint	(Proposition F.3 page 133)
$= \langle \mathbf{x}   \mathbf{B}^* \mathbf{A}^* \mathbf{y} \rangle$	by definition of adjoint	(Proposition F.3 page 133)
$\langle \mathbf{x}   \mathbf{A}^{**} \mathbf{y} \rangle = \langle \mathbf{A}^* \mathbf{x}   \mathbf{y} \rangle$	by definition of adjoint	(Proposition F.3 page 133)
$= \langle \mathbf{y}   \mathbf{A}^* \mathbf{x} \rangle^*$	by definition of inner product	(Definition F.9 page 132)
$= \langle \mathbf{A}\mathbf{y}   \mathbf{x} \rangle^*$	by definition of adjoint	(Proposition F.3 page 133)
$= \langle \mathbf{x}   \mathbf{A}\mathbf{y} \rangle$	by definition of inner product	(Definition F.9 page 132)



**Theorem F.14.** <sup>32</sup> Let  $\mathbf{Y}^{\mathbf{X}}$  be the set of all operators from a linear space  $\mathbf{X}$  to a linear space  $\mathbf{Y}$ . Let  $\mathcal{N}(\mathbf{L})$  be the NULL SPACE of an operator  $\mathbf{L}$  in  $\mathbf{Y}^{\mathbf{X}}$  and  $\mathcal{I}(\mathbf{L})$  the IMAGE SET of  $\mathbf{L}$  in  $\mathbf{Y}^{\mathbf{X}}$ .

<b>T H M</b>	$\mathcal{N}(\mathbf{A}) = \mathcal{I}(\mathbf{A}^*)^\perp$
	$\mathcal{N}(\mathbf{A}^*) = \mathcal{I}(\mathbf{A})^\perp$

PROOF:

$$\begin{aligned}
 \mathcal{I}(\mathbf{A}^*)^\perp &= \{y \in H \mid \langle y | u \rangle = 0 \quad \forall u \in \mathcal{I}(\mathbf{A}^*)\} \\
 &= \{y \in H \mid \langle y | \mathbf{A}^* \mathbf{x} \rangle = 0 \quad \forall \mathbf{x} \in H\} \\
 &= \{y \in H \mid \langle \mathbf{A}y | \mathbf{x} \rangle = 0 \quad \forall \mathbf{x} \in H\} && \text{by definition of } \mathbf{A}^* && \text{(Proposition F.3 page 133)} \\
 &= \{y \in H \mid \mathbf{A}y = 0\} \\
 &= \mathcal{N}(\mathbf{A}) && \text{by definition of } \mathcal{N}(\mathbf{A})
 \end{aligned}$$

$$\begin{aligned}
 \mathcal{I}(\mathbf{A})^\perp &= \{y \in H \mid \langle y | u \rangle = 0 \quad \forall u \in \mathcal{I}(\mathbf{A})\} \\
 &= \{y \in H \mid \langle y | \mathbf{A}\mathbf{x} \rangle = 0 \quad \forall \mathbf{x} \in H\} && \text{by definition of } \mathcal{I} \\
 &= \{y \in H \mid \langle \mathbf{A}^* y | \mathbf{x} \rangle = 0 \quad \forall \mathbf{x} \in H\} && \text{by definition of } \mathbf{A}^* && \text{(Proposition F.3 page 133)} \\
 &= \{y \in H \mid \mathbf{A}^* y = 0\} \\
 &= \mathcal{N}(\mathbf{A}^*) && \text{by definition of } \mathcal{N}(\mathbf{A})
 \end{aligned}$$



## F.4 Special Classes of Operators

### F.4.1 Projection operators

**Definition F.10.** <sup>33</sup> Let  $B(\mathbf{X}, \mathbf{Y})$  be the space of BOUNDED LINEAR OPERATORS on normed linear spaces  $\mathbf{X}$  and  $\mathbf{Y}$ . Let  $\mathbf{P}$  be a bounded linear operator in  $B(\mathbf{X}, \mathbf{Y})$ .

<b>D E F</b>	$\mathbf{P}$ is a <b>projection operator</b> if $\mathbf{P}^2 = \mathbf{P}$ .
----------------------	---

<sup>32</sup> Rudin (1991) page 312

<sup>33</sup> Rudin (1991) page 133 (5.15 Projections), Kubrusly (2001) page 70, Bachman and Narici (1966) page 6, Halmos (1958) page 73 (§41. Projections)

**Theorem F.15.** <sup>34</sup> Let  $\mathcal{B}(\mathbf{X}, \mathbf{Y})$  be the space of BOUNDED LINEAR OPERATORS on normed linear spaces  $\mathbf{X}$  and  $\mathbf{Y}$ . Let  $\mathbf{P}$  be a bounded linear operator in  $\mathcal{B}(\mathbf{X}, \mathbf{Y})$  with NULL SPACE  $\mathcal{N}(\mathbf{P})$  and IMAGE SET  $\mathcal{I}(\mathbf{P})$ .

<b>T H M</b>	1. $\mathbf{P}^2 = \mathbf{P}$ ( $\mathbf{P}$ is a projection operator)      and	$\Rightarrow$	1. $\mathcal{I}(\mathbf{P}) = \mathbf{X}$ and
	2. $\mathbf{\Omega} = \mathbf{X} \hat{+} \mathbf{Y}$ ( $\mathbf{Y}$ compliments $\mathbf{X}$ in $\mathbf{\Omega}$ )      and		2. $\mathcal{N}(\mathbf{P}) = \mathbf{Y}$ and
	3. $\mathbf{P}\mathbf{\Omega} = \mathbf{X}$ ( $\mathbf{P}$ projects onto $\mathbf{X}$ )		3. $\mathbf{\Omega} = \mathcal{I}(\mathbf{P}) \hat{+} \mathcal{N}(\mathbf{P})$

 PROOF:

$$\begin{aligned}
 \mathcal{I}(\mathbf{P}) &= \mathbf{P}\mathbf{\Omega} \\
 &= \mathbf{P}(\mathbf{\Omega}_1 + \mathbf{\Omega}_2) \\
 &= \mathbf{P}\mathbf{\Omega}_1 + \mathbf{P}\mathbf{\Omega}_2 \\
 &= \mathbf{\Omega}_1 + \{0\} \\
 &= \mathbf{\Omega}_1
 \end{aligned}$$


$$\begin{aligned}
 \mathcal{N}(\mathbf{P}) &= \{x \in \mathbf{\Omega} | \mathbf{P}x = 0\} \\
 &= \{x \in (\mathbf{\Omega}_1 + \mathbf{\Omega}_2) | \mathbf{P}x = 0\} \\
 &= \{x \in \mathbf{\Omega}_1 | \mathbf{P}x = 0\} + \{x \in \mathbf{\Omega}_2 | \mathbf{P}x = 0\} \\
 &= \{0\} + \mathbf{\Omega}_2 \\
 &= \mathbf{\Omega}_2
 \end{aligned}$$




**Theorem F.16.** <sup>35</sup> Let  $\mathcal{B}(\mathbf{X}, \mathbf{Y})$  be the space of BOUNDED LINEAR OPERATORS on normed linear spaces  $\mathbf{X}$  and  $\mathbf{Y}$ . Let  $\mathbf{P}$  be a bounded linear operator in  $\mathcal{B}(\mathbf{X}, \mathbf{Y})$ .

<b>T H M</b>	$\mathbf{P}^2 = \mathbf{P}$	$\iff$	$(\mathbf{I} - \mathbf{P})^2 = (\mathbf{I} - \mathbf{P})$
	$\mathbf{P}$ is a projection operator		$(\mathbf{I} - \mathbf{P})$ is a projection operator

 PROOF:

 Proof that  $\mathbf{P}^2 = \mathbf{P} \implies (\mathbf{I} - \mathbf{P})^2 = (\mathbf{I} - \mathbf{P})$ :

$$\begin{aligned}
 (\mathbf{I} - \mathbf{P})^2 &= (\mathbf{I} - \mathbf{P})(\mathbf{I} - \mathbf{P}) \\
 &= \mathbf{I}(\mathbf{I} - \mathbf{P}) + (-\mathbf{P})(\mathbf{I} - \mathbf{P}) \\
 &= \mathbf{I} - \mathbf{P} - \mathbf{P}\mathbf{I} + \mathbf{P}^2 \\
 &= \mathbf{I} - \mathbf{P} - \mathbf{P} + \mathbf{P} \quad \text{by left hypothesis} \\
 &= \mathbf{I} - \mathbf{P}
 \end{aligned}$$

 Proof that  $\mathbf{P}^2 = \mathbf{P} \iff (\mathbf{I} - \mathbf{P})^2 = (\mathbf{I} - \mathbf{P})$ :

$$\begin{aligned}
 \mathbf{P}^2 &= \underbrace{\mathbf{I} - \mathbf{P} - \mathbf{P} + \mathbf{P}^2}_{(\mathbf{I} - \mathbf{P})^2} - (\mathbf{I} - \mathbf{P} - \mathbf{P}) \\
 &= (\mathbf{I} - \mathbf{P})^2 - (\mathbf{I} - \mathbf{P} - \mathbf{P}) \\
 &= (\mathbf{I} - \mathbf{P}) - (\mathbf{I} - \mathbf{P} - \mathbf{P}) \quad \text{by right hypothesis} \\
 &= \mathbf{P}
 \end{aligned}$$



<sup>34</sup>  Michel and Herget (1993) pages 120–121


<sup>35</sup>  Michel and Herget (1993) page 121

**Theorem F.17.** <sup>36</sup> Let  $\mathbf{H}$  be a HILBERT SPACE and  $\mathbf{P}$  an operator in  $\mathbf{H}^{\mathbf{H}}$  with adjoint  $\mathbf{P}^*$ , NULL SPACE  $\mathcal{N}(\mathbf{P})$ , and IMAGE SET  $\mathcal{I}(\mathbf{P})$ .

If  $\mathbf{P}$  is a PROJECTION OPERATOR, then the following are equivalent:

T H M

- |    |  |                                 |        |
|----|--|---------------------------------|--------|
| 1. | $\mathbf{P}^* = \mathbf{P}$  | ( $\mathbf{P}$ is SELF-ADJOINT) | $\iff$ |
| 2. | $\mathbf{P}^*\mathbf{P} = \mathbf{P}\mathbf{P}^*$  | ( $\mathbf{P}$ is NORMAL)       | $\iff$ |
| 3. | $\mathcal{I}(\mathbf{P}) = \mathcal{N}(\mathbf{P})^\perp$  |                                 | $\iff$ |
| 4. | $\langle \mathbf{P}\mathbf{x}   \mathbf{x} \rangle = \ \mathbf{P}\mathbf{x}\ ^2 \quad \forall \mathbf{x} \in \mathbf{X}$ |                                 |        |

 PROOF: This proof is incomplete at this time.

Proof that (1)  $\implies$  (2):

$$\begin{aligned} \mathbf{P}^*\mathbf{P} &= \mathbf{P}^{**}\mathbf{P}^* && \text{by (1)} \\ &= \mathbf{P}\mathbf{P}^* && \text{by Theorem F.13 page 134} \end{aligned}$$

Proof that (1)  $\implies$  (3):

$$\begin{aligned} \mathcal{I}(\mathbf{P}) &= \mathcal{N}(\mathbf{P}^*)^\perp && \text{by Theorem F.14 page 135} \\ &= \mathcal{N}(\mathbf{P})^\perp && \text{by (1)} \end{aligned}$$

Proof that (3)  $\implies$  (4):

Proof that (4)  $\implies$  (1):

$\Rightarrow$

## F.4.2 Self Adjoint Operators

**Definition F.11.** <sup>37</sup> Let  $\mathbf{B} \in \mathcal{B}(\mathbf{H}, \mathbf{H})$  be a BOUNDED operator with adjoint  $\mathbf{B}^*$  on a HILBERT SPACE  $\mathbf{H}$ .

D E F

The operator  $\mathbf{B}$  is said to be **self-adjoint** or **hermitian** if  $\mathbf{B} \doteq \mathbf{B}^*$ .

**Example F.3** (Autocorrelation operator). Let  $\mathbf{x}(t)$  be a random process with autocorrelation

$$\mathbf{R}_{\mathbf{xx}}(t, u) \triangleq \underbrace{\mathbb{E}[\mathbf{x}(t)\mathbf{x}^*(u)]}_{\text{expectation}}.$$

Let an autocorrelation operator  $\mathbf{R}$  be defined as  $[\mathbf{R}\mathbf{f}](t) \triangleq \int_{\mathbb{R}} \underbrace{\mathbf{R}_{\mathbf{xx}}(t, u)}_{\text{kernel}} \mathbf{f}(u) du$ .


E X


$\mathbf{R} = \mathbf{R}^*$  (The auto-correlation operator  $\mathbf{R}$  is **self-adjoint**)





**Theorem F.18.** <sup>38</sup> Let  $\mathbf{S} : \mathbf{H} \rightarrow \mathbf{H}$  be an operator over a HILBERT SPACE  $\mathbf{H}$  with eigenvalues  $\{\lambda_n\}$  and eigenfunctions  $\{\psi_n\}$  such that  $\mathbf{S}\psi_n = \lambda_n\psi_n$  and let  $\|\mathbf{x}\| \triangleq \sqrt{\langle \mathbf{x} | \mathbf{x} \rangle}$ .

T H M

$$\left\{ \begin{array}{l} \mathbf{S} = \mathbf{S}^* \\ \mathbf{S} \text{ is self-adjoint} \end{array} \right\} \implies \left\{ \begin{array}{ll} 1. \quad \langle \mathbf{S}\mathbf{x} | \mathbf{x} \rangle \in \mathbb{R} & (\text{the hermitian quadratic form of } \mathbf{S} \text{ is REAL-VALUED}) \\ 2. \quad \lambda_n \in \mathbb{R} & (\text{eigenvalues of } \mathbf{S} \text{ are REAL-VALUED}) \\ 3. \quad \lambda_n \neq \lambda_m \implies \langle \psi_n | \psi_m \rangle = 0 & (\text{eigenvectors are ORTHOGONAL}) \end{array} \right\}$$

<sup>36</sup>  Rudin (1991) page 314

<sup>37</sup> Historical works regarding self-adjoint operators:  von Neumann (1929) page 49, “linearer Operator  $\mathbf{R}$  selbstadjungiert oder Hermitesche”,  Stone (1932) page 50 (“self-adjoint transformations”)

<sup>38</sup>  Lax (2002) pages 315–316,  Keener (1988) pages 114–119,  Bachman and Narici (1966) page 24 (Theorem 2.1),  Bertero and Boccacci (1998) page 225 (“9.2 SVD of a matrix ... If all eigenvectors are normalized...”)

 PROOF:

1. Proof that  $\mathbf{S} = \mathbf{S}^* \implies \langle \mathbf{S}\mathbf{x} | \mathbf{x} \rangle \in \mathbb{R}$ :

$$\begin{aligned} \langle \mathbf{x} | \mathbf{S}\mathbf{x} \rangle &= \langle \mathbf{S}\mathbf{x} | \mathbf{x} \rangle && \text{by left hypothesis} \\ &= \langle \mathbf{x} | \mathbf{S}\mathbf{x} \rangle^* && \text{by definition of } \langle \triangle | \nabla \rangle \text{ Definition F.9 page 132} \end{aligned}$$

2. Proof that  $\mathbf{S} = \mathbf{S}^* \implies \lambda_n \in \mathbb{R}$ :

$$\begin{aligned} \lambda_n \|\psi_n\|^2 &= \lambda_n \langle \psi_n | \psi_n \rangle && \text{by definition} \\ &= \langle \lambda_n \psi_n | \psi_n \rangle && \text{by definition of } \langle \triangle | \nabla \rangle \text{ Definition F.9 page 132} \\ &= \langle \mathbf{S}\psi_n | \psi_n \rangle && \text{by definition of eigenpairs} \\ &= \langle \psi_n | \mathbf{S}\psi_n \rangle && \text{by left hypothesis} \\ &= \langle \psi_n | \lambda_n \psi_n \rangle && \text{by definition of eigenpairs} \\ &= \lambda_n^* \langle \psi_n | \psi_n \rangle && \text{by definition of } \langle \triangle | \nabla \rangle \text{ Definition F.9 page 132} \\ &= \lambda_n^* \|\psi_n\|^2 && \text{by definition} \end{aligned}$$

3. Proof that  $\mathbf{S} = \mathbf{S}^* \implies [\lambda_n \neq \lambda_m \implies \langle \psi_n | \psi_m \rangle = 0]$ :

$$\begin{aligned} \lambda_n \langle \psi_n | \psi_m \rangle &= \langle \lambda_n \psi_n | \psi_m \rangle && \text{by definition of } \langle \triangle | \nabla \rangle \text{ Definition F.9 page 132} \\ &= \langle \mathbf{S}\psi_n | \psi_m \rangle && \text{by definition of eigenpairs} \\ &= \langle \psi_n | \mathbf{S}\psi_m \rangle && \text{by left hypothesis} \\ &= \langle \psi_n | \lambda_m \psi_m \rangle && \text{by definition of eigenpairs} \\ &= \lambda_m^* \langle \psi_n | \psi_m \rangle && \text{by definition of } \langle \triangle | \nabla \rangle \text{ Definition F.9 page 132} \\ &= \lambda_m \langle \psi_n | \psi_m \rangle && \text{because } \lambda_m \text{ is real} \end{aligned}$$

This implies for  $\lambda_n \neq \lambda_m \neq 0$ ,  $\langle \psi_n | \psi_m \rangle = 0$ .








### F.4.3 Normal Operators


**Definition F.12.** <sup>39</sup> Let  $\mathcal{B}(\mathbf{X}, \mathbf{Y})$  be the space of BOUNDED LINEAR OPERATORS on normed linear spaces  $\mathbf{X}$  and  $\mathbf{Y}$ . Let  $\mathbf{N}^*$  be the adjoint of an operator  $\mathbf{N} \in \mathcal{B}(\mathbf{X}, \mathbf{Y})$ .

**DEF**  $\mathbf{N}$  is *normal* if  $\mathbf{N}^*\mathbf{N} = \mathbf{N}\mathbf{N}^*$ .

**Theorem F.19.** <sup>40</sup> Let  $\mathcal{B}(\mathbf{H}, \mathbf{H})$  be the space of BOUNDED LINEAR OPERATORS on a HILBERT SPACE  $\mathbf{H}$ . Let  $\mathcal{N}(\mathbf{N})$  be the NULL SPACE of an operator  $\mathbf{N}$  in  $\mathcal{B}(\mathbf{H}, \mathbf{H})$  and  $\mathcal{I}(\mathbf{N})$  the IMAGE SET of  $\mathbf{N}$  in  $\mathcal{B}(\mathbf{H}, \mathbf{H})$ .

**THM**  $\underbrace{\mathbf{N}^*\mathbf{N} = \mathbf{N}\mathbf{N}^*}_{\mathbf{N} \text{ is normal}} \iff \|\mathbf{N}^*\mathbf{x}\| = \|\mathbf{N}\mathbf{x}\| \quad \forall \mathbf{x} \in \mathbf{H}$

<sup>39</sup>  Rudin (1991) page 312,  Michel and Herget (1993) page 431,  Dieudonné (1969) page 167,  Frobenius (1878),  Frobenius (1968) page 391

<sup>40</sup>  Rudin (1991) pages 312–313

 PROOF:

1. Proof that  $\mathbf{N}^*\mathbf{N} = \mathbf{N}\mathbf{N}^* \implies \|\mathbf{N}^*\mathbf{x}\| = \|\mathbf{N}\mathbf{x}\|$ :

$$\begin{aligned}
 \|\mathbf{N}\mathbf{x}\|^2 &= \langle \mathbf{N}\mathbf{x} | \mathbf{N}\mathbf{x} \rangle && \text{by definition} \\
 &= \langle \mathbf{x} | \mathbf{N}^*\mathbf{N}\mathbf{x} \rangle && \text{by Proposition F.3 page 133 (definition of } \mathbf{N}^*) \\
 &= \langle \mathbf{x} | \mathbf{N}\mathbf{N}^*\mathbf{x} \rangle && \text{by left hypothesis (N is normal)} \\
 &= \langle \mathbf{N}\mathbf{x} | \mathbf{N}^*\mathbf{x} \rangle && \text{by Proposition F.3 page 133 (definition of } \mathbf{N}^*) \\
 &= \|\mathbf{N}^*\mathbf{x}\|^2 && \text{by definition}
 \end{aligned}$$

2. Proof that  $\mathbf{N}^*\mathbf{N} = \mathbf{N}\mathbf{N}^* \iff \|\mathbf{N}^*\mathbf{x}\| = \|\mathbf{N}\mathbf{x}\|$ :

$$\begin{aligned}
 \langle \mathbf{N}^*\mathbf{N}\mathbf{x} | \mathbf{x} \rangle &= \langle \mathbf{N}\mathbf{x} | \mathbf{N}^{**}\mathbf{x} \rangle && \text{by Proposition F.3 page 133 (definition of } \mathbf{N}^*) \\
 &= \langle \mathbf{N}\mathbf{x} | \mathbf{N}\mathbf{x} \rangle && \text{by Theorem F.13 page 134 (property of adjoint)} \\
 &= \|\mathbf{N}\mathbf{x}\|^2 && \text{by definition} \\
 &= \|\mathbf{N}^*\mathbf{x}\|^2 && \text{by right hypothesis } (\|\mathbf{N}^*\mathbf{x}\| = \|\mathbf{N}\mathbf{x}\|) \\
 &= \langle \mathbf{N}^*\mathbf{x} | \mathbf{N}^*\mathbf{x} \rangle && \text{by definition} \\
 &= \langle \mathbf{N}\mathbf{N}^*\mathbf{x} | \mathbf{x} \rangle && \text{by Proposition F.3 page 133 (definition of } \mathbf{N}^*)
 \end{aligned}$$

$\Rightarrow$

**Theorem F.20.** <sup>41</sup> Let  $B(\mathbf{H}, \mathbf{H})$  be the space of BOUNDED LINEAR OPERATORS on a HILBERT SPACE  $\mathbf{H}$ . Let  $\mathcal{N}(\mathbf{N})$  be the NULL SPACE of an operator  $\mathbf{N}$  in  $B(\mathbf{H}, \mathbf{H})$  and  $\mathcal{I}(\mathbf{N})$  the IMAGE SET of  $\mathbf{N}$  in  $B(\mathbf{H}, \mathbf{H})$ .

<b>T H M</b>	$  \underbrace{\mathbf{N}^*\mathbf{N} = \mathbf{N}\mathbf{N}^*}_{\mathbf{N} \text{ is normal}} \implies \underbrace{\mathcal{N}(\mathbf{N}^*) = \mathcal{N}(\mathbf{N})}_{\mathbf{N} \text{ and } \mathbf{N}^* \text{ have the same null space}}  $
----------------------	---


 PROOF:


$$\begin{aligned}
 \mathcal{N}(\mathbf{N}^*) &= \{ \mathbf{x} | \mathbf{N}^*\mathbf{x} = 0 \quad \forall \mathbf{x} \in \mathbf{X} \} && \text{(definition of } \mathcal{N}) \\
 &= \{ \mathbf{x} | \|\mathbf{N}^*\mathbf{x}\| = 0 \quad \forall \mathbf{x} \in \mathbf{X} \} && \text{by definition of } \|\cdot\| \text{ (Definition F.5 page 124)} \\
 &= \{ \mathbf{x} | \|\mathbf{N}\mathbf{x}\| = 0 \quad \forall \mathbf{x} \in \mathbf{X} \} \\
 &= \{ \mathbf{x} | \mathbf{N}\mathbf{x} = 0 \quad \forall \mathbf{x} \in \mathbf{X} \} && \text{by definition of } \|\cdot\| \text{ (Definition F.5 page 124)} \\
 &= \mathcal{N}(\mathbf{N}) && \text{(definition of } \mathcal{N})
 \end{aligned}$$


$\Rightarrow$

**Theorem F.21.** <sup>42</sup> Let  $B(\mathbf{H}, \mathbf{H})$  be the space of BOUNDED LINEAR OPERATORS on a HILBERT SPACE  $\mathbf{H}$ . Let  $\mathcal{N}(\mathbf{N})$  be the NULL SPACE of an operator  $\mathbf{N}$  in  $B(\mathbf{H}, \mathbf{H})$  and  $\mathcal{I}(\mathbf{N})$  the IMAGE SET of  $\mathbf{N}$  in  $B(\mathbf{H}, \mathbf{H})$ .

<b>T H M</b>	$  \underbrace{\left\{ \mathbf{N}^*\mathbf{N} = \mathbf{N}\mathbf{N}^* \right\}}_{\mathbf{N} \text{ is normal}} \implies \left\{ \underbrace{\lambda_n \neq \lambda_m \implies \langle \psi_n   \psi_m \rangle = 0}_{\text{eigenfunctions associated with distinct eigenvalues are orthogonal}} \right\}  $
----------------------	---

 PROOF: The proof in (1) is flawed. This implies that (2) is also flawed. [Rudin] claims both to be true. (Rudin, 1991)313

<sup>41</sup>  Rudin (1991) pages 312–313

<sup>42</sup>  Rudin (1991) pages 312–313

1. Proof that  $\mathbf{N}^*\mathbf{N} = \mathbf{N}\mathbf{N}^* \implies \mathbf{N}^*\psi = \lambda^*\psi$ :

$$\begin{aligned}
 & \mathbf{N}\psi = \lambda\psi \\
 \iff & \\
 & 0 = \mathcal{N}(\mathbf{N} - \lambda\mathbf{I}) \\
 & = \mathcal{N}([\mathbf{N} - \lambda\mathbf{I}]^*) && \text{by } \mathcal{N}(\mathbf{N}) = \mathcal{N}(\mathbf{N}^*) \\
 & = \mathcal{N}(\mathbf{N}^* - [\lambda\mathbf{I}]^*) && \text{by Theorem F13 page 134} \\
 & = \mathcal{N}(\mathbf{N}^* - \lambda^*\mathbf{I}^*) && \text{by Theorem F13 page 134} \\
 & = \mathcal{N}(\mathbf{N}^* - \lambda^*\mathbf{I}) \\
 \implies & \\
 & (\mathbf{N}^* - \lambda^*\mathbf{I})\psi = 0 \\
 \iff & \mathbf{N}^*\psi = \lambda^*\psi
 \end{aligned}$$

2. Proof that  $\mathbf{N}^*\mathbf{N} = \mathbf{N}\mathbf{N}^* \implies [\lambda_n \neq \lambda_m \implies \langle \psi_n | \psi_m \rangle = 0]$ :

$$\begin{aligned}
 \lambda_n \langle \psi_n | \psi_m \rangle &= \langle \lambda_n \psi_n | \psi_m \rangle && \text{by definition of } \langle \triangle | \nabla \rangle \text{ Definition F9 page 132} \\
 &= \langle \mathbf{N}\psi_n | \psi_m \rangle && \text{by definition of eigenpairs} \\
 &= \langle \psi_n | \mathbf{N}^*\psi_m \rangle && \text{by Proposition F3 page 133 (definition of adjoint)} \\
 &= \langle \psi_n | \lambda_m^*\psi_m \rangle && \text{by (4.)} \\
 &= \lambda_m \langle \psi_n | \psi_m \rangle && \text{by definition of } \langle \triangle | \nabla \rangle \text{ Definition F9 page 132}
 \end{aligned}$$

This implies for  $\lambda_n \neq \lambda_m \neq 0$ ,  $\langle \psi_n | \psi_m \rangle = 0$ .

⇒

## F.4.4 Isometric operators

An operator on a pair of normed linear spaces is *isometric* (next definition) if it is an *isometry*.

**Definition F.13.** Let  $(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \|\cdot\|)$  and  $(Y, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \|\cdot\|)$  be NORMED LINEAR SPACES (Definition F.5 page 124).

**DEF** An operator  $\mathbf{M} \in \mathcal{L}(X, Y)$  is *isometric* if  $\|\mathbf{M}\mathbf{x}\| = \|\mathbf{x}\| \quad \forall \mathbf{x} \in X$ .

**Theorem F.22.**<sup>43</sup> Let  $(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \|\cdot\|)$  and  $(Y, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \|\cdot\|)$  be NORMED LINEAR SPACES. Let  $\mathbf{M}$  be a linear operator in  $\mathcal{L}(X, Y)$ .

<b>T H M</b>	$\underbrace{\ \mathbf{M}\mathbf{x}\  = \ \mathbf{x}\  \quad \forall \mathbf{x} \in X}_{\text{isometric in length}} \iff \underbrace{\ \mathbf{M}\mathbf{x} - \mathbf{M}\mathbf{y}\  = \ \mathbf{x} - \mathbf{y}\  \quad \forall \mathbf{x}, \mathbf{y} \in X}_{\text{isometric in distance}}$
----------------------	--

✎ PROOF:

1. Proof that  $\|\mathbf{M}\mathbf{x}\| = \|\mathbf{x}\| \implies \|\mathbf{M}\mathbf{x} - \mathbf{M}\mathbf{y}\| = \|\mathbf{x} - \mathbf{y}\|$ :

$$\begin{aligned}
 \|\mathbf{M}\mathbf{x} - \mathbf{M}\mathbf{y}\| &= \|\mathbf{M}(\mathbf{x} - \mathbf{y})\| && \text{by definition of linear operators (Definition F4 page 121)} \\
 &= \|\mathbf{M}\mathbf{u}\| && \text{let } \mathbf{u} \triangleq \mathbf{x} - \mathbf{y} \\
 &= \|\mathbf{x} - \mathbf{y}\| && \text{by left hypothesis}
 \end{aligned}$$

<sup>43</sup> [Kubrusly \(2001\) page 239](#) (Proposition 4.37), [Berberian \(1961\) page 27](#) (Theorem IV.7.5)

2. Proof that  $\|\mathbf{M}\mathbf{x}\| = \|\mathbf{x}\| \iff \|\mathbf{M}\mathbf{x} - \mathbf{M}\mathbf{y}\| = \|\mathbf{x} - \mathbf{y}\|$ :

$$\begin{aligned}
 \|\mathbf{M}\mathbf{x}\| &= \|\mathbf{M}(\mathbf{x} - \mathbf{0})\| \\
 &= \|\mathbf{M}\mathbf{x} - \mathbf{M}\mathbf{0}\| && \text{by definition of linear operators (Definition F.4 page 121)} \\
 &= \|\mathbf{x} - \mathbf{0}\| && \text{by right hypothesis} \\
 &= \|\mathbf{x}\|
 \end{aligned}$$

⇒

Isometric operators have already been defined (Definition F.13 page 140) in the more general normed linear spaces, while Theorem F.22 (page 140) demonstrated that in a normed linear space  $\mathbf{X}$ ,  $\|\mathbf{M}\mathbf{x}\| = \|\mathbf{x}\| \iff \|\mathbf{M}\mathbf{x} - \mathbf{M}\mathbf{y}\| = \|\mathbf{x} - \mathbf{y}\|$  for all  $\mathbf{x}, \mathbf{y} \in \mathbf{X}$ . Here in the more specialized inner product spaces, Theorem F.23 (next) demonstrates two additional equivalent properties.

**Theorem F.23.**<sup>44</sup> *Let  $\mathcal{B}(\mathbf{X}, \mathbf{X})$  be the space of BOUNDED LINEAR OPERATORS on a normed linear space  $\mathbf{X} \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \|\cdot\|)$ . Let  $\mathbf{N}$  be a bounded linear operator in  $\mathcal{L}(\mathbf{X}, \mathbf{X})$ , and  $\mathbf{I}$  the identity operator in  $\mathcal{L}(\mathbf{X}, \mathbf{X})$ . Let  $\|\mathbf{x}\| \triangleq \sqrt{\langle \mathbf{x} | \mathbf{x} \rangle}$ .*

*The following conditions are all **equivalent**:*

- |             |    |   |  |        |
|-------------|----|---|--|--------|
| T<br>H<br>M | 1. | $\mathbf{M}^*\mathbf{M} = \mathbf{I}$   |  | $\iff$ |
|             | 2. | $\langle \mathbf{M}\mathbf{x}   \mathbf{M}\mathbf{y} \rangle = \langle \mathbf{x}   \mathbf{y} \rangle$ | $\forall \mathbf{x}, \mathbf{y} \in X$ ( $\mathbf{M}$ is surjective) | $\iff$ |
|             | 3. | $\ \mathbf{M}\mathbf{x} - \mathbf{M}\mathbf{y}\  = \ \mathbf{x} - \mathbf{y}\ $                         | $\forall \mathbf{x}, \mathbf{y} \in X$ (isometric in distance)       | $\iff$ |
|             | 4. | $\ \mathbf{M}\mathbf{x}\  = \ \mathbf{x}\ $   | $\forall \mathbf{x} \in X$ (isometric in length)                     |        |

✎ PROOF:

1. Proof that (1)  $\implies$  (2):

$$\begin{aligned}
 \langle \mathbf{M}\mathbf{x} | \mathbf{M}\mathbf{y} \rangle &= \langle \mathbf{x} | \mathbf{M}^*\mathbf{M}\mathbf{y} \rangle && \text{by Proposition F.3 page 133 (definition of adjoint)} \\
 &= \langle \mathbf{x} | \mathbf{I}\mathbf{y} \rangle && \text{by (1)} \\
 &= \langle \mathbf{x} | \mathbf{y} \rangle && \text{by Definition F.3 page 120 (definition of } \mathbf{I} \text{)}
 \end{aligned}$$



2. Proof that (2)  $\implies$  (4):

$$\begin{aligned}
 \|\mathbf{M}\mathbf{x}\| &= \sqrt{\langle \mathbf{M}\mathbf{x} | \mathbf{M}\mathbf{x} \rangle} && \text{by definition of } \|\cdot\| \\
 &= \sqrt{\langle \mathbf{x} | \mathbf{x} \rangle} && \text{by right hypothesis} \\
 &= \|\mathbf{x}\| && \text{by definition of } \|\cdot\|
 \end{aligned}$$

3. Proof that (2)  $\iff$  (4):

$$\begin{aligned}
 4 \langle \mathbf{M}\mathbf{x} | \mathbf{M}\mathbf{y} \rangle &= \|\mathbf{M}\mathbf{x} + \mathbf{M}\mathbf{y}\|^2 - \|\mathbf{M}\mathbf{x} - \mathbf{M}\mathbf{y}\|^2 + i \|\mathbf{M}\mathbf{x} + i\mathbf{M}\mathbf{y}\|^2 - i \|\mathbf{M}\mathbf{x} - i\mathbf{M}\mathbf{y}\|^2 && \text{by polarization id.} \\
 &= \|\mathbf{M}(\mathbf{x} + \mathbf{y})\|^2 - \|\mathbf{M}(\mathbf{x} - \mathbf{y})\|^2 + i \|\mathbf{M}(\mathbf{x} + i\mathbf{y})\|^2 - i \|\mathbf{M}(\mathbf{x} - i\mathbf{y})\|^2 && \text{by Definition F.4} \\
 &= \|\mathbf{x} + \mathbf{y}\|^2 - \|\mathbf{x} - \mathbf{y}\|^2 + i \|\mathbf{x} + i\mathbf{y}\|^2 - i \|\mathbf{x} - i\mathbf{y}\|^2 && \text{by left hypothesis}
 \end{aligned}$$

4. Proof that (3)  $\iff$  (4): by Theorem F.22 page 140

<sup>44</sup>  Michel and Herget (1993) page 432 (Theorem 7.5.8),  Kubrusly (2001) page 391 (Proposition 5.72)

5. Proof that (4)  $\implies$  (1):

$$\begin{aligned}
 \langle \mathbf{M}^* \mathbf{M} \mathbf{x} \mid \mathbf{x} \rangle &= \langle \mathbf{M} \mathbf{x} \mid \mathbf{M}^{**} \mathbf{x} \rangle && \text{by Proposition F3 page 133 (definition of adjoint)} \\
 &= \langle \mathbf{M} \mathbf{x} \mid \mathbf{M} \mathbf{x} \rangle && \text{by Theorem F13 page 134 (property of adjoint)} \\
 &= \|\mathbf{M} \mathbf{x}\|^2 && \text{by definition} \\
 &= \|\mathbf{x}\|^2 && \text{by left hypothesis with } y = 0 \\
 &= \langle \mathbf{x} \mid \mathbf{x} \rangle && \text{by definition} \\
 &= \langle \mathbf{I} \mathbf{x} \mid \mathbf{x} \rangle && \text{by Definition F3 page 120 (definition of } \mathbf{I} \text{)} \\
 \implies \mathbf{M}^* \mathbf{M} &= \mathbf{I} && \forall \mathbf{x} \in X
 \end{aligned}$$

$\Rightarrow$

**Theorem F.24.** <sup>45</sup> Let  $B(X, Y)$  be the space of BOUNDED LINEAR OPERATORS on normed linear spaces  $X$  and  $Y$ . Let  $\mathbf{M}$  be a bounded linear operator in  $B(X, Y)$ , and  $\mathbf{I}$  the identity operator in  $\mathcal{L}(X, X)$ . Let  $\Lambda$  be the set of eigenvalues of  $\mathbf{M}$ . Let  $\|\mathbf{x}\| \triangleq \sqrt{\langle \mathbf{x} \mid \mathbf{x} \rangle}$ .

<b>T H M</b>	$  \underbrace{\mathbf{M}^* \mathbf{M} = \mathbf{I}}_{\mathbf{M} \text{ is isometric}} \implies \begin{cases} \ \mathbf{M}\  = 1 & \text{(UNIT LENGTH)} \\  \lambda  = 1 & \forall \lambda \in \Lambda \end{cases} \text{ and }  $
----------------------	--

PROOF:

1. Proof that  $\mathbf{M}^* \mathbf{M} = \mathbf{I} \implies \|\mathbf{M}\| = 1$ :

$$\begin{aligned}
 \|\mathbf{M}\| &= \sup_{\mathbf{x} \in X} \{ \|\mathbf{M} \mathbf{x}\| \mid \|\mathbf{x}\| = 1 \} && \text{by Definition F6 page 125} \\
 &= \sup_{\mathbf{x} \in X} \{ \|\mathbf{x}\| \mid \|\mathbf{x}\| = 1 \} && \text{by Theorem F23 page 141} \\
 &= \sup_{\mathbf{x} \in X} \{ 1 \} \\
 &= 1
 \end{aligned}$$

2. Proof that  $|\lambda| = 1$ : Let  $(\mathbf{x}, \lambda)$  be an eigenvector-eigenvalue pair.

$$\begin{aligned}
 1 &= \frac{1}{\|\mathbf{x}\|} \|\mathbf{x}\| \\
 &= \frac{1}{\|\mathbf{x}\|} \|\mathbf{M} \mathbf{x}\| && \text{by Theorem F23 page 141} \\
 &= \frac{1}{\|\mathbf{x}\|} \|\lambda \mathbf{x}\| && \text{by definition of } \lambda \\
 &= \frac{1}{\|\mathbf{x}\|} |\lambda| \|\mathbf{x}\| && \text{by homogeneous property of } \|\cdot\| \\
 &= |\lambda|
 \end{aligned}$$

$\Rightarrow$

**Example F4** (One sided shift operator). <sup>46</sup> Let  $X$  be the set of all sequences with range  $\mathbb{W}$   $(0, 1, 2, \dots)$  and shift operators defined as

$$\begin{aligned}
 1. \quad \mathbf{S}_r(x_0, x_1, x_2, \dots) &\triangleq (0, x_0, x_1, x_2, \dots) && \text{(right shift operator)} \\
 2. \quad \mathbf{S}_l(x_0, x_1, x_2, \dots) &\triangleq (x_1, x_2, x_3, \dots) && \text{(left shift operator)}
 \end{aligned}$$

<b>E X</b>	<ol style="list-style-type: none"> <li>1. <math>\mathbf{S}_r</math> is an isometric operator.</li> <li>2. <math>\mathbf{S}_r^* = \mathbf{S}_l</math></li> </ol>
----------------	---

<sup>45</sup> Michel and Herget (1993) page 432

<sup>46</sup> Michel and Herget (1993) page 441



 PROOF:

1. Proof that  $S_r^* = S_l$ :

$$\begin{aligned}
 \langle S_r(x_0, x_1, x_2, \dots) | (y_0, y_1, y_2, \dots) \rangle &= \langle (0, x_0, x_1, x_2, \dots) | (y_0, y_1, y_2, \dots) \rangle \\
 &= \sum_{n=1}^{\infty} x_{n-1} y_n^* \\
 &= \sum_{n=0}^{\infty} x_n y_{n+1}^* \\
 &= \sum_{n=0}^{\infty} x_n y_{n+1}^* \\
 &= \langle (x_0, x_1, x_2, \dots) | (y_1, y_2, y_3, \dots) \rangle \\
 &= \left\langle (x_0, x_1, x_2, \dots) | \underbrace{S_l(y_0, y_1, y_2, \dots)}_{S_r^*} \right\rangle
 \end{aligned}$$

2. Proof that  $S_r$  is isometric ( $S_r^* S_r = I$ ):

$$\begin{aligned}
 S_r^* S_r &= S_l S_r \\
 &= I
 \end{aligned}$$

by 1.



## F.4.5 Unitary operators

**Definition F.14.** <sup>47</sup> Let  $B(X, Y)$  be the space of BOUNDED LINEAR OPERATORS on normed linear spaces  $X$  and  $Y$ . Let  $U$  be a bounded linear operator in  $B(X, Y)$ , and  $I$  the identity operator in  $B(X, X)$ .

**DEF** The operator  $U$  is **unitary** if  $U^* U = U U^* = I$ .







**Proposition F.4.** Let  $B(X, Y)$  be the space of BOUNDED LINEAR OPERATORS on normed linear spaces  $X$  and  $Y$ . Let  $U$  and  $V$  be BOUNDED LINEAR OPERATORS in  $B(X, Y)$ .

**P R P**  $\left. \begin{array}{l} U \text{ is UNITARY} \\ V \text{ is UNITARY} \end{array} \right\} \text{ and } \Rightarrow (UV) \text{ is UNITARY.}$

 PROOF:

$$\begin{aligned}
 (UV)(UV)^* &= (UV)(V^* U^*) && \text{by Theorem F.8 page 129} \\
 &= U(VV^*)U^* && \text{by associative property} \\
 &= U I U^* && \text{by definition of unitary operators (Definition F.14 page 143)} \\
 &= I && \text{by definition of unitary operators (Definition F.14 page 143)}
 \end{aligned}$$

$$\begin{aligned}
 (UV)^*(UV) &= (V^* U^*)(UV) && \text{by Theorem F.8 page 129} \\
 &= V^*(U^* U)V && \text{by associative property} \\
 &= V^* I V && \text{by definition of unitary operators (Definition F.14 page 143)} \\
 &= I && \text{by definition of unitary operators (Definition F.14 page 143)}
 \end{aligned}$$

<sup>47</sup>  Rudin (1991) page 312,  Michel and Herget (1993) page 431,  Autonne (1901) page 209,  Autonne (1902),  Schur (1909),  Steen (1973)



**Theorem F.25.** <sup>48</sup> Let  $\mathcal{B}(\mathcal{H}, \mathcal{H})$  be the space of BOUNDED LINEAR OPERATORS on a HILBERT SPACE  $\mathcal{H}$ . Let  $\mathcal{I}(\mathbf{U})$  be the IMAGE SET of  $\mathbf{U}$ .

If  $\mathbf{U}$  is a **bounded linear operator** ( $\mathbf{U} \in \mathcal{B}(\mathcal{H}, \mathcal{H})$ ), then the following conditions are **equivalent**:

**T H M**

- |    |   |                                   |                                |
|----|---|-----------------------------------|--------------------------------|
| 1. | $\mathbf{U}\mathbf{U}^* = \mathbf{U}^*\mathbf{U} = \mathbf{I}$  | (UNITARY)                         | $\iff$                         |
| 2. | $\langle \mathbf{U}\mathbf{x}   \mathbf{U}\mathbf{y} \rangle = \langle \mathbf{U}^*\mathbf{x}   \mathbf{U}^*\mathbf{y} \rangle = \langle \mathbf{x}   \mathbf{y} \rangle$ | and $\mathcal{I}(\mathbf{U}) = X$ | (SURJECTIVE) $\iff$            |
| 3. | $\ \mathbf{U}\mathbf{x} - \mathbf{U}\mathbf{y}\  = \ \mathbf{U}^*\mathbf{x} - \mathbf{U}^*\mathbf{y}\  = \ \mathbf{x} - \mathbf{y}\ $                                     | and $\mathcal{I}(\mathbf{U}) = X$ | (ISOMETRIC IN DISTANCE) $\iff$ |
| 4. | $\ \mathbf{U}\mathbf{x}\  = \ \mathbf{x}\ $   | and $\mathcal{I}(\mathbf{U}) = X$ | (ISOMETRIC IN LENGTH)          |

PROOF:

1. Proof that (1)  $\implies$  (2):

(a)  $\langle \mathbf{U}\mathbf{x} | \mathbf{U}\mathbf{y} \rangle = \langle \mathbf{U}^*\mathbf{x} | \mathbf{U}^*\mathbf{y} \rangle = \langle \mathbf{x} | \mathbf{y} \rangle$  by Theorem F.23 (page 141).

(b) Proof that  $\mathcal{I}(\mathbf{U}) = X$ :

$$\begin{aligned}
 X &\supseteq \mathcal{I}(\mathbf{U}) && \text{because } \mathbf{U} \in X^X \\
 &\supseteq \mathcal{I}(\mathbf{U}\mathbf{U}^*) \\
 &= \mathcal{I}(\mathbf{I}) && \text{by left hypothesis } (\mathbf{U}^*\mathbf{U} = \mathbf{U}\mathbf{U}^* = \mathbf{I}) \\
 &= X && \text{by Definition F.3 page 120 (definition of } \mathcal{I})
 \end{aligned}$$

2. Proof that (2)  $\iff$  (3)  $\iff$  (4): by Theorem F.23 page 141.

3. Proof that (3)  $\implies$  (1):

(a) Proof that  $\|\mathbf{U}\mathbf{x} - \mathbf{U}\mathbf{y}\| = \|\mathbf{x} - \mathbf{y}\| \implies \mathbf{U}^*\mathbf{U} = \mathbf{I}$ : by Theorem F.23 page 141

(b) Proof that  $\|\mathbf{U}^*\mathbf{x} - \mathbf{U}^*\mathbf{y}\| = \|\mathbf{x} - \mathbf{y}\| \implies \mathbf{U}\mathbf{U}^* = \mathbf{I}$ :

$$\begin{aligned}
 \|\mathbf{U}^*\mathbf{x} - \mathbf{U}^*\mathbf{y}\| = \|\mathbf{x} - \mathbf{y}\| &\implies \mathbf{U}^{**}\mathbf{U}^* = \mathbf{I} && \text{by Theorem F.23 page 141} \\
 &\implies \mathbf{U}\mathbf{U}^* = \mathbf{I} && \text{by Theorem F.13 page 134}
 \end{aligned}$$



**Theorem F.26.** Let  $\mathcal{B}(\mathcal{H}, \mathcal{H})$  be the space of BOUNDED LINEAR OPERATORS on a HILBERT SPACE  $\mathcal{H}$ . Let  $\mathbf{U}$  be a bounded linear operator in  $\mathcal{B}(\mathcal{H}, \mathcal{H})$ ,  $\mathcal{N}(\mathbf{U})$  the NULL SPACE of  $\mathbf{U}$ , and  $\mathcal{I}(\mathbf{U})$  the IMAGE SET of  $\mathbf{U}$ .

**T H M**

$$\underbrace{\mathbf{U}\mathbf{U}^* = \mathbf{U}^*\mathbf{U} = \mathbf{I}}_{\mathbf{U} \text{ is unitary}} \implies \left\{ \begin{array}{lll} \mathbf{U}^{-1} = \mathbf{U}^* & & \text{and} \\ \mathcal{I}(\mathbf{U}) = \mathcal{I}(\mathbf{U}^*) = X & & \text{and} \\ \mathcal{N}(\mathbf{U}) = \mathcal{N}(\mathbf{U}^*) = \{0\} & & \text{and} \\ \|\mathbf{U}\| = \|\mathbf{U}^*\| = 1 & & \text{(UNIT LENGTH)} \end{array} \right\}$$

PROOF:

1. Note that  $\mathbf{U}$ ,  $\mathbf{U}^*$ , and  $\mathbf{U}^{-1}$  are all both *isometric* and *normal*:

$$\begin{aligned}
 \mathbf{U}^*\mathbf{U} &= \mathbf{I} \implies \mathbf{U} \text{ is isometric} \\
 \mathbf{U}\mathbf{U}^* &= \mathbf{U}^*\mathbf{U} = \mathbf{I} \implies \mathbf{U}^* \text{ is isometric} \\
 \mathbf{U}^{-1} &= \mathbf{U}^* \implies \mathbf{U}^{-1} \text{ is isometric}
 \end{aligned}$$

$$\begin{aligned}
 \mathbf{U}^*\mathbf{U} &= \mathbf{U}\mathbf{U}^* = \mathbf{I} \implies \mathbf{U} \text{ is normal} \\
 \mathbf{U}\mathbf{U}^* &= \mathbf{U}^*\mathbf{U} = \mathbf{I} \implies \mathbf{U}^* \text{ is normal} \\
 \mathbf{U}^{-1} &= \mathbf{U}^* \implies \mathbf{U}^{-1} \text{ is normal}
 \end{aligned}$$

<sup>48</sup> Rudin (1991) pages 313–314 (Theorem 12.13), Knapp (2005a) page 45 (Proposition 2.6)

2. Proof that  $\mathbf{U}^*\mathbf{U} = \mathbf{U}\mathbf{U}^* = \mathbf{I} \implies \mathcal{I}(\mathbf{U}) = \mathcal{I}(\mathbf{U}^*) = \mathcal{H}$ : by Theorem F.25 page 144.

3. Proof that  $\mathbf{U}^*\mathbf{U} = \mathbf{U}\mathbf{U}^* = \mathbf{I} \implies \mathcal{N}(\mathbf{U}) = \mathcal{N}(\mathbf{U}^*) = \mathcal{N}(\mathbf{U}^{-1})$ :

$$\begin{aligned}\mathcal{N}(\mathbf{U}^*) &= \mathcal{N}(\mathbf{U}) && \text{because } \mathbf{U} \text{ and } \mathbf{U}^* \text{ are both normal and by Theorem F.21 page 139} \\ &= \mathcal{I}(\mathbf{U})^\perp && \text{by Theorem F.14 page 135} \\ &= X^\perp && \text{by above result} \\ &= \{0\}\end{aligned}$$

4. Proof that  $\mathbf{U}^*\mathbf{U} = \mathbf{U}\mathbf{U}^* = \mathbf{I} \implies \|\mathbf{U}^{-1}\| = \|\mathbf{U}^*\| = \|\mathbf{U}\| = 1$ :

Because  $\mathbf{U}$ ,  $\mathbf{U}^*$ , and  $\mathbf{U}^{-1}$  are all isometric and by Theorem F.24 page 142.



Example F.5 (Rotation matrix). <sup>49</sup>

$$\begin{array}{|l|} \mathbf{E} \\ \mathbf{X} \end{array} \left\{ \mathbf{R}_\theta \triangleq \begin{bmatrix} \cos\theta & -\sin\theta \\ \sin\theta & \cos\theta \end{bmatrix} \right\} \implies \left\{ \begin{array}{l} (1). \quad \mathbf{R}_\theta^{-1} = \mathbf{R}_{-\theta} \quad \text{and} \\ (2). \quad \mathbf{R}_\theta^* = \mathbf{R}_\theta^{-1} \quad (\mathbf{R} \text{ is unitary}) \end{array} \right\}$$

rotation matrix  $\mathbf{R}_\theta : \mathbb{R}^2 \rightarrow \mathbb{R}^2$

PROOF:

$$\begin{aligned}\mathbf{R}^* &= \mathbf{R}^H \\ &= \begin{bmatrix} \cos\theta & -\sin\theta \\ \sin\theta & \cos\theta \end{bmatrix}^H && \text{by definition of } \mathbf{R} \\ &= \begin{bmatrix} \cos\theta & \sin\theta \\ -\sin\theta & \cos\theta \end{bmatrix} && \text{by definition of Hermetian transpose operator } H \\ &= \begin{bmatrix} \cos(-\theta) & -\sin(-\theta) \\ \sin(-\theta) & \cos(-\theta) \end{bmatrix} && \text{by Theorem C.2 page 77} \\ &= \mathbf{R}_{-\theta} && \text{by definition of } \mathbf{R} \\ &= \mathbf{R}^{-1} && \text{by 1.}\end{aligned}$$



Example F.6. <sup>50</sup> Let  $\mathbf{A}$  and  $\mathbf{B}$  be matrix operators.

$$\begin{array}{|l|} \mathbf{E} \\ \mathbf{X} \end{array} \left\{ \mathbf{A} \triangleq \begin{bmatrix} \cos\phi & -\sin\phi & 0 \\ \sin\phi & \cos\phi & 0 \\ 0 & 0 & 1 \end{bmatrix} \quad \mathbf{B} \triangleq \begin{bmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{bmatrix} \right.$$

$\mathbf{A}$  is a rotation operator.  $\mathbf{B}$  is a reflection operator.

Both  $\mathbf{A}$  and  $\mathbf{B}$  are unitary.

Example F.7. Examples of Fredholm integral operators include

$$\begin{array}{|l|} \mathbf{E} \\ \mathbf{X} \end{array} \begin{array}{ll} 1. \text{ Fourier Transform} & [\tilde{\mathbf{F}}\mathbf{x}](f) = \int_{t \in \mathbb{R}} \mathbf{x}(t) e^{-i2\pi f t} dt \quad \kappa(t, f) = e^{-i2\pi f t} \\ 2. \text{ Inverse Fourier Transform} & [\tilde{\mathbf{F}}^{-1}\tilde{\mathbf{x}}](t) = \int_{f \in \mathbb{R}} \tilde{\mathbf{x}}(f) e^{i2\pi f t} df \quad \kappa(f, t) = e^{i2\pi f t} \\ 3. \text{ Laplace operator} & [\mathbf{L}\mathbf{x}](s) = \int_{t \in \mathbb{R}} \mathbf{x}(t) e^{-st} dt \quad \kappa(t, s) = e^{-st} \end{array}$$

Example F.8 (Translation operator). Let  $\mathbf{X} = \mathcal{L}_{\mathbb{R}}^2$  and  $\mathbf{T} \in \mathbf{X}^{\mathbf{X}}$  be defined as

$$\mathbf{T}f(x) \triangleq f(x-1) \quad \forall f \in \mathcal{L}_{\mathbb{R}}^2 \quad (\text{translation operator})$$

<sup>49</sup> Noble and Daniel (1988) page 311

<sup>50</sup> Gel'fand (1963) page 4, Gelfand et al. (2018) page 4

<b>E X</b>	1.	$\mathbf{T}^{-1}\mathbf{f}(x) = \mathbf{f}(x+1)$	$\forall \mathbf{f} \in L^2_{\mathbb{R}}$	(inverse translation operator)
	2.	$\mathbf{T}^* = \mathbf{T}^{-1}$		( $\mathbf{T}$ is invertible)
	3.	$\mathbf{T}^*\mathbf{T} = \mathbf{T}\mathbf{T}^* = \mathbf{I}$		( $\mathbf{T}$ is unitary)

PROOF:

1. Proof that  $\mathbf{T}^{-1}\mathbf{f}(x) = \mathbf{f}(x+1)$ :

$$\begin{aligned}\mathbf{T}^{-1}\mathbf{T} &= \mathbf{I} \\ \mathbf{T}\mathbf{T}^{-1} &= \mathbf{I}\end{aligned}$$

2. Proof that  $\mathbf{T}$  is unitary:

$$\begin{aligned}\langle \mathbf{T}\mathbf{f}(x) | \mathbf{g}(x) \rangle &= \langle \mathbf{f}(x-1) | \mathbf{g}(x) \rangle && \text{by definition of } \mathbf{T} \\ &= \int_x \mathbf{f}(x-1) \mathbf{g}^*(x) \, dx \\ &= \int_x \mathbf{f}(x) \mathbf{g}^*(x+1) \, dx \\ &= \langle \mathbf{f}(x) | \mathbf{g}(x+1) \rangle \\ &= \left\langle \mathbf{f}(x) | \underbrace{\mathbf{T}^{-1}\mathbf{g}(x)}_{\mathbf{T}^*} \right\rangle && \text{by 1.}\end{aligned}$$

⇒

*Example F9* (Dilation operator). Let  $\mathbf{X} = L^2_{\mathbb{R}}$  and  $\mathbf{T} \in \mathbf{X}^{\mathbf{X}}$  be defined as

$$\mathbf{D}\mathbf{f}(x) \triangleq \sqrt{2}\mathbf{f}(2x) \quad \forall \mathbf{f} \in L^2_{\mathbb{R}} \quad (\text{dilation operator})$$

<b>E X</b>	1.	$\mathbf{D}^{-1}\mathbf{f}(x) = \frac{1}{\sqrt{2}}\mathbf{f}\left(\frac{1}{2}x\right)$	$\forall \mathbf{f} \in L^2_{\mathbb{R}}$	(inverse dilation operator)
	2.	$\mathbf{D}^* = \mathbf{D}^{-1}$		( $\mathbf{D}$ is invertible)
	3.	$\mathbf{D}^*\mathbf{D} = \mathbf{D}\mathbf{D}^* = \mathbf{I}$		( $\mathbf{D}$ is unitary)

PROOF:

1. Proof that  $\mathbf{D}^{-1}\mathbf{f}(x) = \frac{1}{\sqrt{2}}\mathbf{f}\left(\frac{1}{2}x\right)$ :

$$\begin{aligned}\mathbf{D}^{-1}\mathbf{D} &= \mathbf{I} \\ \mathbf{D}\mathbf{D}^{-1} &= \mathbf{I}\end{aligned}$$

2. Proof that  $\mathbf{D}$  is unitary:

$$\begin{aligned}\langle \mathbf{D}\mathbf{f}(x) | \mathbf{g}(x) \rangle &= \left\langle \sqrt{2}\mathbf{f}(2x) | \mathbf{g}(x) \right\rangle && \text{by definition of } \mathbf{D} \\ &= \int_x \sqrt{2}\mathbf{f}(2x) \mathbf{g}^*(x) \, dx \\ &= \int_{u \in \mathbb{R}} \sqrt{2}\mathbf{f}(u) \mathbf{g}^*\left(\frac{1}{2}u\right) \frac{1}{2} \, du && \text{let } u \triangleq 2x \implies dx = \frac{1}{2} du \\ &= \int_{u \in \mathbb{R}} \mathbf{f}(u) \left[ \frac{1}{\sqrt{2}}\mathbf{g}\left(\frac{1}{2}u\right) \right]^* \, du \\ &= \left\langle \mathbf{f}(x) | \frac{1}{\sqrt{2}}\mathbf{g}\left(\frac{1}{2}x\right) \right\rangle \\ &= \left\langle \mathbf{f}(x) | \underbrace{\mathbf{D}^{-1}\mathbf{g}(x)}_{\mathbf{D}^*} \right\rangle && \text{by 1.}\end{aligned}$$



*Example F.10 (Delay operator).* Let  $\mathbf{X}$  be the set of all sequences and  $\mathbf{D} \in \mathbf{X}^{\mathbf{X}}$  be a delay operator.

**E X** The delay operator  $\mathbf{D} ((x_n)_{n \in \mathbb{Z}}) \triangleq ((x_{n-1})_{n \in \mathbb{Z}})$  is unitary.

**PROOF:** The inverse  $\mathbf{D}^{-1}$  of the delay operator  $\mathbf{D}$  is

$$\mathbf{D}^{-1} ((x_n)_{n \in \mathbb{Z}}) \triangleq ((x_{n+1})_{n \in \mathbb{Z}}).$$

$$\begin{aligned} \langle \mathbf{D}((x_n)) | (y_n) \rangle &= \langle ((x_{n-1})) | (y_n) \rangle && \text{by definition of } \mathbf{D} \\ &= \sum_n x_{n-1} y_n^* \\ &= \sum_n x_n y_{n+1}^* \\ &= \langle ((x_n)) | ((y_{n+1})) \rangle \\ &= \left\langle ((x_n)) | \underbrace{\mathbf{D}^{-1}((y_n))}_{\mathbf{D}^*} \right\rangle \end{aligned}$$

Therefore,  $\mathbf{D}^* = \mathbf{D}^{-1}$ . This implies that  $\mathbf{D}\mathbf{D}^* = \mathbf{D}^*\mathbf{D} = \mathbf{I}$  which implies that  $\mathbf{D}$  is unitary.

*Example F.11 (Fourier transform).* Let  $\tilde{\mathbf{F}}$  be the *Fourier Transform* and  $\tilde{\mathbf{F}}^{-1}$  the *inverse Fourier Transform* operator (Theorem 7.1 page 52)

$$[\tilde{\mathbf{F}}\mathbf{x}](f) \triangleq \int_t \mathbf{x}(t) \underbrace{e^{-i2\pi ft}}_{\kappa(t, f)} dt \quad [\tilde{\mathbf{F}}^{-1}\tilde{\mathbf{x}}](t) \triangleq \int_f \tilde{\mathbf{x}}(f) \underbrace{e^{i2\pi ft}}_{\kappa^*(t, f)} df.$$

**E X**  $\tilde{\mathbf{F}}^* = \tilde{\mathbf{F}}^{-1}$  (the Fourier Transform operator  $\tilde{\mathbf{F}}$  is unitary)

**PROOF:**

$$\begin{aligned} \langle \tilde{\mathbf{F}}\mathbf{x} | \tilde{\mathbf{y}} \rangle &= \left\langle \int_t \mathbf{x}(t) e^{-i2\pi ft} dt | \tilde{\mathbf{y}}(f) \right\rangle \\ &= \int_t \mathbf{x}(t) \langle e^{-i2\pi ft} | \tilde{\mathbf{y}}(f) \rangle dt \\ &= \int_t \mathbf{x}(t) \int_f e^{-i2\pi ft} \tilde{\mathbf{y}}^*(f) df dt \\ &= \int_t \mathbf{x}(t) \left[ \int_f e^{i2\pi ft} \tilde{\mathbf{y}}(f) df \right]^* dt \\ &= \left\langle \mathbf{x}(t) | \int_f \tilde{\mathbf{y}}(f) e^{i2\pi ft} df \right\rangle \\ &= \left\langle \mathbf{x} | \underbrace{\tilde{\mathbf{F}}^{-1}\tilde{\mathbf{y}}}_{\tilde{\mathbf{F}}^*} \right\rangle \end{aligned}$$

This implies that  $\tilde{\mathbf{F}}$  is unitary ( $\tilde{\mathbf{F}}^* = \tilde{\mathbf{F}}^{-1}$ ).

## F.5 Operator order

**Definition F.15.** <sup>51</sup> Let  $\mathbf{P} \in \mathcal{Y}^{\mathcal{X}}$  be an operator.

**DEF**  $\mathbf{P}$  is **positive** if  $\langle \mathbf{P}\mathbf{x} | \mathbf{x} \rangle \geq 0 \ \forall \mathbf{x} \in \mathcal{X}$ .  
This condition is denoted  $\mathbf{P} \geq 0$ .

**Theorem F.27.** <sup>52</sup>

**THM**  $\underbrace{\mathbf{P} \geq 0 \text{ and } \mathbf{Q} \geq 0}_{\mathbf{P} \text{ and } \mathbf{Q} \text{ are both positive}} \implies \begin{cases} (\mathbf{P} + \mathbf{Q}) \geq 0 & ((\mathbf{P} + \mathbf{Q}) \text{ is positive}) \\ \mathbf{A}^* \mathbf{P} \mathbf{A} \geq 0 & \forall \mathbf{A} \in \mathcal{B}(\mathcal{X}, \mathcal{X}) \quad (\mathbf{A}^* \mathbf{P} \mathbf{A} \text{ is positive}) \\ \mathbf{A}^* \mathbf{A} \geq 0 & \forall \mathbf{A} \in \mathcal{B}(\mathcal{X}, \mathcal{X}) \quad (\mathbf{A}^* \mathbf{A} \text{ is positive}) \end{cases}$


 PROOF:

$\langle (\mathbf{P} + \mathbf{Q})\mathbf{x}   \mathbf{x} \rangle = \langle \mathbf{P}\mathbf{x}   \mathbf{x} \rangle + \langle \mathbf{Q}\mathbf{x}   \mathbf{x} \rangle$	by additive property of $\langle \triangle   \nabla \rangle$ (Definition F.9 page 132)
$\geq \langle \mathbf{P}\mathbf{x}   \mathbf{x} \rangle$	by left hypothesis
$\geq 0$	by left hypothesis
$\langle \mathbf{A}^* \mathbf{P} \mathbf{A} \mathbf{x}   \mathbf{x} \rangle = \langle \mathbf{P} \mathbf{A} \mathbf{x}   \mathbf{A} \mathbf{x} \rangle$	by definition of adjoint (Proposition F.3 page 133)
$= \langle \mathbf{P} \mathbf{y}   \mathbf{y} \rangle$	where $\mathbf{y} \triangleq \mathbf{A} \mathbf{x}$
$\geq 0$	by left hypothesis
$\langle \mathbf{I} \mathbf{x}   \mathbf{x} \rangle = \langle \mathbf{x}   \mathbf{x} \rangle$	by definition of $\mathbf{I}$ (Definition F.3 page 120)
$\geq 0$	by non-negative property of $\langle \triangle   \nabla \rangle$ (Definition F.9 page 132)
$\implies \mathbf{I}$ is positive	
$\langle \mathbf{A}^* \mathbf{A} \mathbf{x}   \mathbf{x} \rangle = \langle \mathbf{A}^* \mathbf{I} \mathbf{A} \mathbf{x}   \mathbf{x} \rangle$	by definition of $\mathbf{I}$ (Definition F.3 page 120)
$\geq 0$	by two previous results



**Definition F.16.** <sup>53</sup> Let  $\mathbf{A}, \mathbf{B} \in \mathcal{B}(\mathcal{X}, \mathcal{Y})$  be BOUNDED operators.

**DEF**  $\mathbf{A} \geq \mathbf{B}$  (“ $\mathbf{A}$  is greater than or equal to  $\mathbf{B}$ ”) if  
 $\mathbf{A} - \mathbf{B} \geq 0$  (“ $(\mathbf{A} - \mathbf{B})$  is positive”)

<sup>51</sup>  Michel and Herget (1993) page 429 (Definition 7.4.12)

<sup>52</sup>  Michel and Herget (1993) page 429

<sup>53</sup>  Michel and Herget (1993) page 429

# APPENDIX G

## LINEAR COMBINATIONS

### G.1 Linear combinations in linear spaces

A *linear space* (Definition F.1 page 119) in general is not equipped with a *topology*. Without a topology, it is not possible to determine whether an *infinite sum* of vectors converges. Therefore in this section (dealing with linear spaces), all definitions related to sums of vectors will be valid for *finite* sums only (finite “ $N$ ”).

**Definition G.1.** <sup>1</sup> Let  $\{\mathbf{x}_n \in X \mid n=1,2,\dots,N\}$  be a set of vectors in a LINEAR SPACE  $(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}))$ .

DEF

A vector  $\mathbf{x} \in X$  is a **linear combination** of the vectors in  $\{\mathbf{x}_n\}$  if

$$\text{there exists } \{\alpha_n \in \mathbb{F} \mid n=1,2,\dots,N\} \text{ such that } \mathbf{x} = \sum_{n=1}^N \alpha_n \mathbf{x}_n.$$

**Definition G.2.** <sup>2</sup> Let  $(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}))$  be a linear space and  $Y$  be a subset of  $X$ .

DEF

The **linear span** of  $Y$  is defined as  $\text{span} Y \triangleq \left\{ \sum_{\gamma \in I} \alpha_\gamma \mathbf{y}_\gamma \mid \alpha_\gamma \in \mathbb{F}, \mathbf{y}_\gamma \in Y \right\}$ .

The set  $Y$  **spans** a set  $A$  if  $A \subseteq \text{span} Y$ .

**Proposition G.1.** <sup>3</sup> Let  $\{\mathbf{x}_n \in X \mid n=1,2,\dots,N\}$  be a set of vectors in a LINEAR SPACE  $L \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}))$ .

PRP

1.  $\text{span}\{\mathbf{x}_n\}$  is a LINEAR SPACE (Definition F.1 page 119) and
2.  $\text{span}\{\mathbf{x}_n\}$  is a LINEAR SUBSPACE of  $L$ .

**Definition G.3.** <sup>4</sup> Let  $L \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}))$  be a LINEAR SPACE.

DEF

The set  $Y \triangleq \{\mathbf{x}_n \in X \mid n=1,2,\dots,N\}$  is **linearly independent** in  $L$  if

$$\left\{ \sum_{n=1}^N \alpha_n \mathbf{x}_n = 0 \right\} \implies \{\alpha_1 = \alpha_2 = \dots = \alpha_N = 0\}.$$


The set  $Y$  is **linearly dependent** in  $L$  if  $Y$  is not linearly independent in  $L$ .

<sup>1</sup> Berberian (1961) page 11 (Definition I.4.1), Kubrusly (2001) page 46

<sup>2</sup> Michel and Herget (1993) page 86 (3.3.7 Definition), Kurdila and Zabrankin (2005) page 44, Searcoid (2002) page 71 (Definition 3.2.5—more general definition)

<sup>3</sup> Kubrusly (2001) page 46

<sup>4</sup> Bachman and Narici (1966) pages 3–4, Christensen (2003) page 2, Heil (2011) page 156 (Definition 5.7)

Figure G.1: Lattice of *linear combinations*



**Definition G.4.** <sup>5</sup> Let  $\{\mathbf{x}_n \in X \mid n=1,2,\dots,N\}$  be a set of vectors in a LINEAR SPACE  $L \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}))$ .

DEF

The set  $\{\mathbf{x}_n\}$  is a **Hamel basis** for  $L$  if

1.  $\{\mathbf{x}_n\}$  SPANS  $L$  (Definition G.2 page 149) and
2.  $\{\mathbf{x}_n\}$  is LINEARLY INDEPENDENT in  $L$  (Definition G.1 page 149).

A **HAMEL BASIS** is also called a **linear basis**.

**Definition G.5.** <sup>6</sup> Let  $L \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}))$  be a LINEAR SPACE. Let  $\mathbf{x}$  be a VECTOR in  $L$  and  $Y \triangleq \{\mathbf{x}_n \in X \mid n=1,2,\dots,N\}$  be a set of vectors in  $L$ .

DEF

The expression  $\sum_{n=1}^N \alpha_n \mathbf{x}_n$  is the **expansion** of  $\mathbf{x}$  on  $Y$  in  $L$  if  $\mathbf{x} = \sum_{n=1}^N \alpha_n \mathbf{x}_n$ .

In this case, the sequence  $(\alpha_n)_{n=1}^N$  is the **coordinates** of  $\mathbf{x}$  with respect to  $Y$  in  $L$ .  
If  $\alpha_N \neq 0$ , then  $N$  is the **dimension**  $\dim L$  of  $L$ .

**Theorem G.1.** <sup>7</sup> Let  $\{\mathbf{x}_n \mid n=1,2,\dots,N\}$  be a HAMEL BASIS (Definition G.4 page 151) for a LINEAR SPACE  $(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}))$ .

THM

$$\left\{ \mathbf{x} = \sum_{n=1}^N \alpha_n \mathbf{x}_n = \sum_{n=1}^N \beta_n \mathbf{x}_n \right\} \implies \underbrace{\alpha_n = \beta_n \quad \forall n=1,2,\dots,N}_{\text{coordinates of } \mathbf{x} \text{ are UNIQUE}} \quad \forall \mathbf{x} \in X$$

 PROOF:

$$\mathbf{0} = \mathbf{x} - \mathbf{x}$$

$$= \sum_{n=1}^N \alpha_n \mathbf{x}_n - \sum_{n=1}^N \beta_n \mathbf{x}_n$$

$$= \sum_{n=1}^N (\alpha_n - \beta_n) \mathbf{x}_n$$

$$\implies \{\mathbf{x}_n\} \text{ is linearly dependent if } (\alpha_n - \beta_n) \neq 0 \quad \forall n = 1, 2, \dots, N$$

$$\implies (\alpha_n - \beta_n) = 0 \quad \forall n = 1, 2, \dots, N \quad (\text{because } \{\mathbf{x}_n\} \text{ is a basis and therefore must be linearly independent})$$

$$\implies \alpha_n = \beta_n \text{ for } n = 1, 2, \dots, N$$













**Theorem G.2.** <sup>8</sup> Let  $L \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}))$  be a LINEAR SPACE.


THM


$$\left\{ \begin{array}{l} 1. \{\mathbf{x}_n \in X \mid n=1,2,\dots,N\} \text{ is a HAMEL BASIS for } L \\ 2. \{\mathbf{y}_n \in X \mid n=1,2,\dots,M\} \text{ is a set of LINEARLY INDEPENDENT vectors in } L \end{array} \right\} \text{ and } \implies \left\{ \begin{array}{l} 1. M \leq N \\ 2. M = N \implies \{\mathbf{y}_n \mid n=1,2,\dots,M\} \text{ is a BASIS for } L \\ 3. M \neq N \implies \{\mathbf{y}_n \mid n=1,2,\dots,M\} \text{ is NOT a basis for } L \end{array} \right\} \text{ and }$$

 PROOF:

<sup>5</sup> Hamel (1905),  Bachman and Narici (1966) page 4,  Kubrusly (2001) pages 48–49 (Section 2.4),  Young (2001) page 1,  Carothers (2005) page 25,  Heil (2011) page 125 (Definition 4.1)

<sup>6</sup> Hamel (1905),  Bachman and Narici (1966) page 4,  Kubrusly (2001) pages 48–49 (Section 2.4),  Young (2001) page 1,  Carothers (2005) page 25,  Heil (2011) page 125 (Definition 4.1)

<sup>7</sup>  Michel and Herget (1993) pages 89–90 (Theorem 3.3.25)

<sup>8</sup>  Michel and Herget (1993) pages 90–91 (Theorem 3.3.26)

1. Proof that  $\{y_1, x_1, \dots, x_{N-1}\}$  is a *basis* for  $L$ :

(a) Proof that  $\{y_1, x_1, \dots, x_{N-1}\}$  *spans*  $L$ :

i. Because  $\{x_n\}_{n=1,2,\dots,N}$  is a *basis* for  $L$ , there exists  $\beta \in \mathbb{F}$  and  $\{\alpha_n \in \mathbb{F}\}_{n=1,2,\dots,N}$  such that

$$\beta y_1 + \sum_{n=1}^N \alpha_n x_n = 0.$$

ii. Select an  $n$  such that  $\alpha_n \neq 0$  and renumber (if necessary) the above indices such that

$$x_n = -\frac{\beta}{\alpha_n} y_1 + \sum_{n=1}^{N-1} \frac{\alpha_n}{\alpha_n} x_n.$$

iii. Then, for any  $y \in X$ , we can write

$$\begin{aligned} y &= \sum_{n=1}^N \gamma_n x_n \\ &= \left( \sum_{n=1}^{N-1} \gamma_n x_n \right) + \gamma_n x_n \left( -\frac{\beta}{\alpha_n} y_1 - \sum_{n=1}^{N-1} \frac{\alpha_n}{\alpha_n} x_n \right) \\ &= -\frac{\beta \gamma_n}{\alpha_n} y_1 + \sum_{n=1}^{N-1} \left( \gamma_n - \frac{\alpha_n \gamma_n}{\alpha_n} \right) x_n \\ &= \delta y_1 + \sum_{n=1}^{N-1} \delta_n x_n \end{aligned}$$

iv. This implies that  $\{y_1, x_1, \dots, x_{N-1}\}$  *spans*  $L$ :

(b) Proof that  $\{y_1, x_1, \dots, x_{N-1}\}$  is *linearly independent*:

i. If  $\{y_1, x_1, \dots, x_{N-1}\}$  is *linearly dependent*, then there exists  $\{\epsilon, \epsilon_1, \dots, \epsilon_{N-1}\}$  such that

$$\epsilon y_1 + \left( \sum_{n=1}^{N-1} \epsilon_n x_n \right) + 0 x_n = 0.$$

ii. item (1(b)i) implies that the coordinate of  $y_1$  associated with  $x_n$  is 0.

$$y_1 = -\left( \sum_{n=1}^{N-1} \frac{\epsilon_n}{\epsilon} x_n \right) + 0 x_n = 0.$$

iii. item (1(a)i) implies that the coordinate of  $y_1$  associated with  $x_n$  is *not* 0.

$$y_1 = -\sum_{n=1}^N \frac{\alpha_n}{\beta} x_n.$$

iv. This implies that item (1(b)i) (that the set is linearly dependent) is *false* because item (1(b)ii) and item (1(b)iii) *contradict* each other.

v. This implies  $\{y_1, x_1, \dots, x_{N-1}\}$  is *linearly independent*.

2. Proof that  $\{y_1, y_2, x_1, \dots, x_{N-2}\}$  is a *basis*: Repeat item (1).

3. Suppose  $m = n$ . Proof that  $\{y_1, y_2, \dots, y_M\}$  is a *basis*: Repeat item (1)  $M - 1$  times.

4. Proof that  $M \not\asymp N$ :

(a) Suppose that  $M = N + 1$ .

(b) Then because  $\{y_n\}_{n=1,2,\dots,N}$  is a *basis*, there exists  $\{\zeta_n\}_{n=1,2,\dots,N+1}$  such that

$$\sum_{n=1}^{N+1} \zeta_n y_n = 0.$$

(c) This implies that  $\{y_n\}_{n=1,2,\dots,N+1}$  is *linearly dependent*.

(d) This implies that  $\{y_n|_{n=1,2,\dots,N+1}\}$  is *not* a basis.

(e) This implies that  $M \neq N$ .

5. Proof that  $M \neq N \implies \{y_n|_{n=1,2,\dots,M}\}$  is *not* a basis for  $L$ :

(a) Proof that  $M > N \implies \{y_n|_{n=1,2,\dots,M}\}$  is *not* a basis for  $L$ : same as in item (4).

(b) Proof that  $M < N \implies \{y_n|_{n=1,2,\dots,M}\}$  is *not* a basis for  $L$ :

i. Suppose  $m = N - 1$ .

ii. Then  $\{y_n|_{n=1,2,\dots,N-1}\}$  is a *basis* and there exists  $\lambda$  such that

$$\sum_{n=1}^N \lambda_{n \in \mathbb{Z}} y_{n \in \mathbb{Z}} = 0.$$

iii. This implies that  $\{y_n|_{n=1,2,\dots,N}\}$  is *linearly dependent* and is *not* a basis.

iv. But this contradicts item (3), therefore  $M \neq N - 1$ .

v. Because  $M = N$  yields a basis but  $M = N - 1$  does not,  $M < N - 1$  also does not yield a basis.

⇒

**Corollary G.1.** <sup>9</sup> Let  $L \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}))$  be a linear space.

$$\left\{ \begin{array}{l} 1. \{x_n \in X|_{n=1,2,\dots,N}\} \text{ is a HAMEL BASIS for } L \text{ and} \\ 2. \{y_n \in X|_{n=1,2,\dots,M}\} \text{ is a HAMEL BASIS for } L \end{array} \right\} \implies \{N = M\}$$

(all Hamel bases for  $L$  have the same number of vectors)

✎ PROOF: This follows from Theorem G.2 (page 151).

⇒

## G.2 Bases in topological linear spaces

A linear space supports the concept of the *span* of a set of vectors (Definition G.2 page 149). In a topological linear space  $\Omega \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), T)$ , a set  $A$  is said to be *total* in  $\Omega$  if the span of  $A$  is *dense* in  $\Omega$ . In this case,  $A$  is said to be a *total set* or a *complete set*. However, this use of “complete” in a “complete set” is not equivalent to the use of “complete” in a “complete metric space”.<sup>10</sup> In this text, except for these comments and Definition G.6, “complete” refers to the metric space definition only.

If a set is both *total* and *linearly independent* (Definition G.3 page 149) in  $\Omega$ , then that set is a *Hamel basis* (Definition G.4 page 151) for  $\Omega$ .

**Definition G.6.** <sup>11</sup> Let  $A^-$  be the CLOSURE of a  $A$  in a TOPOLOGICAL LINEAR SPACE  $\Omega \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), T)$ . Let  $\text{span } A$  be the SPAN (Definition G.2 page 149) of a set  $A$ .

**DEF** A set of vectors  $A$  is **total** (or **complete** or **fundamental**) in  $\Omega$  if  
 $(\text{span } A)^- = \Omega$  (SPAN of  $A$  is DENSE in  $\Omega$ ).

<sup>9</sup> Kubrusly (2001) page 52 (Theorem 2.7), Michel and Herget (1993) page 91 (Theorem 3.3.31)

<sup>10</sup> Haaser and Sullivan (1991) pages 296–297 (6-Orthogonal Bases), Rynne and Youngson (2008) page 78 (Remark 3.50), Heil (2011) page 21 (Remark 1.26)

<sup>11</sup> Young (2001) page 19 (Definition 1.5.1), Sohrab (2003) page 362 (Definition 9.2.3), Gupta (1998) page 134 (Definition 2.4), Bachman and Narici (1966) pages 149–153 (Definition 9.3, Theorems 9.9 and 9.10)

## G.3 Schauder bases in Banach spaces

**Definition G.7.** <sup>12</sup> Let  $\mathbf{B} \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \|\cdot\|)$  be a BANACH SPACE. Let  $\dot{=}$  represent STRONG CONVERGENCE in  $\mathbf{B}$ .

The countable set  $\{x_n \in X \mid n \in \mathbb{N}\}$  is a **Schauder basis** for  $\mathbf{B}$  if for each  $x \in X$

1.  $\exists (\alpha_n \in \mathbb{F})_{n \in \mathbb{N}}$  such that  $x \dot{=} \sum_{n=1}^{\infty} \alpha_n x_n$  (STRONG CONVERGENCE in  $\mathbf{B}$ ) and
2.  $\left\{ \sum_{n=1}^{\infty} \alpha_n x_n \dot{=} \sum_{n=1}^{\infty} \beta_n x_n \right\} \implies \{(\alpha_n) = (\beta_n)\}$  (COEFFICIENT FUNCTIONALS are UNIQUE)

In this case,  $\sum_{n=1}^{\infty} \alpha_n x_n$  is the **expansion** of  $x$  on  $\{x_n \mid n \in \mathbb{N}\}$  and

the elements of  $(\alpha_n)$  are the **coefficient functionals** associated with the basis  $\{x_n\}$ . Coefficient functionals are also called **coordinate functionals**.

In a Banach space, the existence of a Schauder basis implies that the space is *separable* (Theorem G.3 page 154). The question of whether the converse is also true was posed by Banach himself in 1932,<sup>13</sup> and became known as “*The basis problem*”. This remained an open question for many years. The question was finally answered some 41 years later in 1973 by Per Enflo (University of California at Berkeley), with the answer being “no”. Enflo constructed a counterexample in which a separable Banach space does *not* have a Schauder basis.<sup>14</sup> Life is simpler in Hilbert spaces where the converse is true: a Hilbert space has a Schauder basis *if and only if* it is separable (Theorem G.11 page 167).

**Theorem G.3.** <sup>15</sup> Let  $\mathbf{B} \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \|\cdot\|)$  be a BANACH SPACE. Let  $\mathbb{Q}$  be the field of rational numbers.

$$\left\{ \begin{array}{l} 1. \mathbf{B} \text{ has a SCHAUDER BASIS} \text{ and} \\ 2. \mathbb{Q} \text{ is DENSE in } \mathbb{F}. \end{array} \right\} \implies \{ \mathbf{B} \text{ is SEPARABLE} \}$$

PROOF:

1. lemma:

$$\begin{aligned} \left| \left\{ x \mid \exists (\alpha_n \mathbb{Q})_{n \in \mathbb{N}} \text{ such that } \lim_{N \rightarrow \infty} \left\| x - \sum_{n=1}^N \alpha_n x_n \right\| = 0 \right\} \right| &= |\mathbb{Q} \times \mathbb{N}| \\ &= |\mathbb{Z} \times \mathbb{Z}| \\ &= |\mathbb{Z}| \\ &= \text{countably infinite} \end{aligned}$$

<sup>12</sup> Carothers (2005) pages 24–25, Christensen (2003) pages 46–49 (Definition 3.1.1 and page 49), Young (2001) page 19 (Section 6), Singer (1970) page 17, Schauder (1927), Schauder (1928)

<sup>13</sup> Banach (1932a) page 111

<sup>14</sup> Enflo (1973), Lindenstrauss and Tzafriri (1977) pages 84–95 (Section 2.d)

<sup>15</sup> Bachman et al. (2000) page 112 (3.4.8), Giles (2000) page 17, Heil (2011) page 21 (Theorem 1.27)

2. remainder of proof:

$\mathcal{B}$  has a Schauder basis  $(\mathbf{x}_n)_{n \in \mathbb{N}}$

$\Rightarrow$  for every  $\mathbf{x} \in \mathcal{B}$ , there exists  $(\alpha_n \in \mathbb{F})_{n \in \mathbb{N}}$  such that  $\mathbf{x} \triangleq \sum_{n=1}^{\infty} \alpha_n \mathbf{x}_n$  by Definition G.7 page 154

$\Rightarrow$  for every  $\mathbf{x} \in \mathcal{B}$ , there exists  $(\alpha_n \in \mathbb{F})_{n \in \mathbb{N}}$  such that  $\lim_{N \rightarrow \infty} \left\| \mathbf{x} - \sum_{n=1}^N \alpha_n \mathbf{x}_n \right\| = 0$

$\Rightarrow$  for every  $\mathbf{x} \in \mathcal{B}$ , there exists  $(\alpha_n \in \mathbb{Q})_{n \in \mathbb{N}}$  such that  $\lim_{N \rightarrow \infty} \left\| \mathbf{x} - \sum_{n=1}^N \alpha_n \mathbf{x}_n \right\| = 0$  because  $\mathbb{Q}^- = \mathbb{F}$

$\Rightarrow \mathcal{B} = \left\{ \mathbf{x} \mid \exists (\alpha_n \mathbb{Q})_{n \in \mathbb{N}} \text{ such that } \lim_{N \rightarrow \infty} \left\| \mathbf{x} - \sum_{n=1}^N \alpha_n \mathbf{x}_n \right\| = 0 \right\}$

$\Rightarrow \mathcal{B} = \left\{ \mathbf{x} \mid \exists (\alpha_n \mathbb{Q})_{n \in \mathbb{N}} \text{ such that } \mathbf{x} = \lim_{N \rightarrow \infty} \sum_{n=1}^N \alpha_n \mathbf{x}_n \right\}^-$

$\Rightarrow \mathcal{B}$  is separable by (1) lemma page 154

$\Rightarrow$

**Definition G.8.** <sup>16</sup> Let  $\{\mathbf{x}_n \mid n \in \mathbb{N}\}$  and  $\{\mathbf{y}_n \mid n \in \mathbb{N}\}$  be SCHAUDER BASES of a BANACH SPACE  $(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \|\cdot\|)$ .

**DEF**  $\{\mathbf{x}_n\}$  is **equivalent** to  $\{\mathbf{y}_n\}$   
if there exists a BOUNDED INVERTIBLE operator  $\mathbf{R}$  in  $\mathcal{X}^{\mathcal{X}}$  such that  $\mathbf{R}\mathbf{x}_n = \mathbf{y}_n \quad \forall n \in \mathbb{Z}$

**Theorem G.4.** <sup>17</sup> Let  $\{\mathbf{x}_n \mid n \in \mathbb{N}\}$  and  $\{\mathbf{y}_n \mid n \in \mathbb{N}\}$  be SCHAUDER BASES of a BANACH SPACE  $(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \|\cdot\|)$ .

**THM**  $\left\{ \begin{aligned} &\{\mathbf{x}_n\} \text{ is EQUIVALENT to } \{\mathbf{y}_n\} \\ &\iff \left\{ \sum_{n=1}^{\infty} \alpha_n \mathbf{x}_n \text{ is CONVERGENT} \iff \sum_{n=1}^{\infty} \alpha_n \mathbf{y}_n \text{ is CONVERGENT} \right\} \end{aligned} \right\}$

**Lemma G.1.** <sup>18</sup> Let  $\Omega \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), T)$  be a topological linear space. Let  $\text{span} A$  be the SPAN of a set  $A$  (Definition G.2 page 149). Let  $\tilde{f}(\omega)$  and  $\tilde{g}(\omega)$  be the FOURIER TRANSFORMS (Definition 4.2 page 16) of the functions  $f(x)$  and  $g(x)$ , respectively, in  $\mathcal{L}_{\mathbb{R}}^2$  (Definition B.1 page 69). Let  $\check{a}(\omega)$  be the DTFT (Definition 6.1 page 41) of a sequence  $(a_n)_{n \in \mathbb{Z}}$  in  $\mathcal{E}_{\mathbb{R}}^2$  (Definition 5.2 page 27).

**LEM**  $\left\{ \begin{aligned} &(1). \quad \{\mathbf{T}^n \mathbf{f} \mid n \in \mathbb{Z}\} \text{ is a SCHAUDER BASIS for } \Omega \text{ and} \\ &(2). \quad \{\mathbf{T}^n \mathbf{g} \mid n \in \mathbb{Z}\} \text{ is a SCHAUDER BASIS for } \Omega \end{aligned} \right\} \Rightarrow \left\{ \begin{aligned} &\exists (a_n)_{n \in \mathbb{Z}} \text{ such that} \\ &\tilde{f}(\omega) = \check{a}(\omega) \tilde{g}(\omega) \end{aligned} \right\}$

$\nabla$ PROOF: Let  $\mathcal{V}'_0$  be the space spanned by  $\{\mathbf{T}^n \phi \mid n \in \mathbb{Z}\}$ .

$$\begin{aligned} \tilde{f}(\omega) &\triangleq \tilde{\mathbf{F}} \mathbf{f} && \text{by definition of } \tilde{\mathbf{F}} && (\text{Definition 4.2 page 16}) \\ &= \tilde{\mathbf{F}} \sum_{n \in \mathbb{Z}} a_n \mathbf{T} \mathbf{g} && \text{by (2)} \\ &= \sum_{n \in \mathbb{Z}} a_n \tilde{\mathbf{F}} \mathbf{T} \mathbf{g} \end{aligned}$$

<sup>16</sup> Young (2001) page 25 (Definition 1.8.1, Theorem 1.8.7)

<sup>17</sup> Young (2001) page 25 (Definition 1.8.1, Theorem 1.8.7)

<sup>18</sup> Daubechies (1992) page 140

$$= \underbrace{\sum_{n \in \mathbb{Z}} a_n e^{-i\omega n}}_{\check{a}(\omega)} \tilde{\mathbf{F}} \tilde{\mathbf{g}}(\omega)$$

by Corollary I.1 page 195

$$= \check{a}(\omega) \tilde{\mathbf{g}}(\omega)$$

by definition of  $\tilde{\mathbf{F}}$  and  $\tilde{\mathbf{F}}$  by (Definition 6.1 page 41, Definition 4.2 page 16)

$$\begin{aligned} V_0 &\triangleq \left\{ f(x) \mid f(x) = \sum_{n \in \mathbb{Z}} b_n \mathbf{T}^n g(x) \right\} \\ &= \left\{ f(x) \mid \tilde{\mathbf{F}} f(x) = \tilde{\mathbf{F}} \sum_{n \in \mathbb{Z}} b_n \mathbf{T}^n g(x) \right\} \\ &= \{ f(x) \mid \tilde{f}(\omega) = \tilde{b}(\omega) \tilde{g}(\omega) \} \\ &= \{ f(x) \mid \tilde{f}(\omega) = \tilde{b}(\omega) \check{a}(\omega) \tilde{f}(\omega) \} \\ &= \{ f(x) \mid \tilde{f}(\omega) = \tilde{c}(\omega) \tilde{f}(\omega) \} \quad \text{where } \tilde{c}(\omega) \triangleq \tilde{b}(\omega) \check{a}(\omega) \\ &= \left\{ f(x) \mid f(x) = \sum_{n \in \mathbb{Z}} c_n f(x - n) \right\} \\ &\triangleq V'_0 \end{aligned}$$



## G.4 Linear combinations in inner product spaces

**Definition G.9.** Let  $(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle \mid \nabla \rangle)$  be an INNER PRODUCT SPACE (Definition F.9 page 132).

DEF

Two vectors  $\mathbf{x}$  and  $\mathbf{y}$  in  $X$  are **orthogonal** if

$$\langle \mathbf{x} \mid \mathbf{y} \rangle = \begin{cases} 0 & \text{for } \mathbf{x} \neq \mathbf{y} \\ c \in \mathbb{F} \setminus 0 & \text{for } \mathbf{x} = \mathbf{y} \end{cases}$$

In an *inner product space*, *orthogonality* is a special case of *linear independence*; or alternatively, linear independence is a generalization of orthogonality (next theorem).

**Theorem G.5.** <sup>19</sup> Let  $\{\mathbf{x}_n \in X \mid n=1,2,\dots,N\}$  be a set of vectors in an INNER PRODUCT SPACE (Definition F.9 page 132)  $(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle \mid \nabla \rangle)$ .

THM

$$\left\{ \begin{array}{l} \{\mathbf{x}_n\} \text{ is ORTHOGONAL} \\ \text{(Definition G.9 page 156)} \end{array} \right\} \implies \left\{ \begin{array}{l} \{\mathbf{x}_n\} \text{ is LINEARLY INDEPENDENT} \\ \text{(Definition G.1 page 149)} \end{array} \right\}$$

PROOF:

1. Proof using *Pythagorean theorem*:

Let  $(\alpha_n)_{n \in \mathbb{N}}$  be a sequence with at least one nonzero element.

<sup>19</sup> Aliprantis and Burkinshaw (1998) page 283 (Corollary 32.8), Kubrusly (2001) page 352 (Proposition 5.34)

$$\begin{aligned}
\left\| \sum_{n=1}^N \alpha_n \mathbf{x}_n \right\|^2 &= \sum_{n=1}^N \|\alpha_n \mathbf{x}_n\|^2 && \text{by left hypoth. and Pythagorean Theorem} \\
&= \sum_{n=1}^N |\alpha_n|^2 \|\mathbf{x}_n\|^2 && \text{by definition of } \|\cdot\| \quad (\text{Definition F.5 page 124}) \\
&> 0 \\
\Rightarrow \sum_{n=1}^N \alpha_n \mathbf{x}_n &\neq 0 \\
\Rightarrow (\mathbf{x}_n)_{n \in \mathbb{N}} &\text{ is linearly independent } && \text{by definition of linear independence} \quad (\text{Definition G.3 page 149})
\end{aligned}$$

2. Alternative proof:

$$\begin{aligned}
\sum_{n=1}^N \alpha_n \mathbf{x}_n = \mathbf{0} &\Rightarrow \left\langle \sum_{n=1}^N \alpha_n \mathbf{x}_n \mid \mathbf{x}_m \right\rangle = \langle \mathbf{0} \mid \mathbf{x}_m \rangle \\
&\Rightarrow \sum_{n=1}^N \alpha_n \langle \mathbf{x}_n \mid \mathbf{x}_m \rangle = 0 \\
&\Rightarrow \sum_{n=1}^N \alpha_n \delta(k-m) = 0 \\
&\Rightarrow \alpha_m = 0 \quad \text{for } m = 1, 2, \dots, N
\end{aligned}$$

⇒

**Theorem G.6** (Bessel's Equality).<sup>20</sup> Let  $\{\mathbf{x}_n \in X \mid n=1,2,\dots,N\}$  be a set of vectors in an INNER PRODUCT SPACE (Definition F.9 page 132)  $(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle \mid \nabla \rangle)$  and with  $\|\mathbf{x}\| \triangleq \sqrt{\langle \mathbf{x} \mid \mathbf{x} \rangle}$ .

T H M	$ \left\{ \begin{array}{l} \{\mathbf{x}_n\} \text{ is ORTHONORMAL} \\ (\text{Definition G.9 page 156}) \end{array} \right\} \Rightarrow \left\{ \underbrace{\left\  \mathbf{x} - \sum_{n=1}^N \langle \mathbf{x} \mid \mathbf{x}_n \rangle \mathbf{x}_n \right\ ^2}_{\text{approximation error}} = \ \mathbf{x}\ ^2 - \sum_{n=1}^N  \langle \mathbf{x} \mid \mathbf{x}_n \rangle ^2 \quad \forall \mathbf{x} \in X} \right\} $
-------------	--

PROOF:

$$\begin{aligned}
&\left\| \mathbf{x} - \sum_{n=1}^N \langle \mathbf{x} \mid \mathbf{x}_n \rangle \mathbf{x}_n \right\|^2 \\
&= \|\mathbf{x}\|^2 + \left\| \sum_{n=1}^N \langle \mathbf{x} \mid \mathbf{x}_n \rangle \mathbf{x}_n \right\|^2 - 2\Re \left\langle \mathbf{x} \mid \sum_{n=1}^N \langle \mathbf{x} \mid \mathbf{x}_n \rangle \mathbf{x}_n \right\rangle && \text{by polar identity} \\
&= \|\mathbf{x}\|^2 + \left\| \sum_{n=1}^N \langle \mathbf{x} \mid \mathbf{x}_n \rangle \mathbf{x}_n \right\|^2 - 2\Re \left[ \left( \sum_{n=1}^N \langle \mathbf{x} \mid \mathbf{x}_n \rangle \right)^* \langle \mathbf{x} \mid \mathbf{x}_n \rangle \right] && \text{by property of } \langle \triangle \mid \nabla \rangle \quad (\text{Definition F.9 page 132}) \\
&= \|\mathbf{x}\|^2 + \sum_{n=1}^N \|\langle \mathbf{x} \mid \mathbf{x}_n \rangle \mathbf{x}_n\|^2 - 2\Re \left[ \left( \sum_{n=1}^N \langle \mathbf{x} \mid \mathbf{x}_n \rangle \right)^* \langle \mathbf{x} \mid \mathbf{x}_n \rangle \right] && \text{by Pythagorean Theorem}
\end{aligned}$$

<sup>20</sup> Bachman et al. (2000) page 103, Pedersen (2000) pages 38–39

$$\begin{aligned}
&= \|\mathbf{x}\|^2 + \sum_{n=1}^N \left\| \langle \mathbf{x} | \mathbf{x}_n \rangle \mathbf{x}_n \right\|^2 - 2\Re \left( \sum_{n=1}^N \langle \mathbf{x} | \mathbf{x}_n \rangle^* \langle \mathbf{x} | \mathbf{x}_n \rangle \right) \\
&= \|\mathbf{x}\|^2 + \sum_{n=1}^N \underbrace{|\langle \mathbf{x} | \mathbf{x}_n \rangle|^2}_{1} \underbrace{\|\mathbf{x}_n\|^2}_1 - 2\Re \left( \sum_{n=1}^N \langle \mathbf{x} | \mathbf{x}_n \rangle^* \langle \mathbf{x} | \mathbf{x}_n \rangle \right) \quad \text{by property of } \|\cdot\| \quad (\text{Definition F.5 page 124}) \\
&= \|\mathbf{x}\|^2 + \sum_{n=1}^N |\langle \mathbf{x} | \mathbf{x}_n \rangle|^2 \cdot 1 - 2\Re \left( \sum_{n=1}^N \langle \mathbf{x} | \mathbf{x}_n \rangle^* \langle \mathbf{x} | \mathbf{x}_n \rangle \right) \quad \text{by def. of orthonormality} \quad (\text{Definition G.9 page 156}) \\
&= \|\mathbf{x}\|^2 + \sum_{n=1}^N |\langle \mathbf{x} | \mathbf{x}_n \rangle|^2 - 2\Re \sum_{n=1}^N |\langle \mathbf{x} | \mathbf{x}_n \rangle|^2 \\
&= \|\mathbf{x}\|^2 + \sum_{n=1}^N |\langle \mathbf{x} | \mathbf{x}_n \rangle|^2 - 2 \sum_{n=1}^N |\langle \mathbf{x} | \mathbf{x}_n \rangle|^2 \quad \text{because } |\cdot| \text{ is real} \\
&= \|\mathbf{x}\|^2 - \sum_{n=1}^N |\langle \mathbf{x} | \mathbf{x}_n \rangle|^2
\end{aligned}$$

⇒

**Theorem G.7** (Bessel's inequality). <sup>21</sup> Let  $\{\mathbf{x}_n \in X \mid n=1,2,\dots,N\}$  be a set of vectors in an INNER PRODUCT SPACE (Definition F.9 page 132)  $(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle | \nabla \rangle)$  and with  $\|\mathbf{x}\| \triangleq \sqrt{\langle \mathbf{x} | \mathbf{x} \rangle}$ .

<b>T H M</b>	$ \left\{ \begin{array}{l} \{\mathbf{x}_n\} \text{ is ORTHONORMAL} \\ (\text{Definition G.9 page 156}) \end{array} \right\} \implies \left\{ \sum_{n=1}^N  \langle \mathbf{x}   \mathbf{x}_n \rangle ^2 \leq \ \mathbf{x}\ ^2 \quad \forall \mathbf{x} \in X \right\} $
----------------------	---

PROOF:

$$\begin{aligned}
0 &\leq \left\| \mathbf{x} - \sum_{n=1}^N \langle \mathbf{x} | \mathbf{x}_n \rangle \mathbf{x}_n \right\|^2 && \text{by definition of } \|\cdot\| && (\text{Definition F.5 page 124}) \\
&= \|\mathbf{x}\|^2 - \sum_{n=1}^N |\langle \mathbf{x} | \mathbf{x}_n \rangle|^2 && \text{by Bessel's Equality} && (\text{Theorem G.6 page 157})
\end{aligned}$$

⇒

The *Best Approximation Theorem* (next) shows that

- the best sequence for representing a vector is the sequence of projections of the vector onto the sequence of basis functions
- the error of the projection is orthogonal to the projection.

**Theorem G.8** (Best Approximation Theorem). <sup>22</sup> Let  $\{\mathbf{x}_n \in X \mid n=1,2,\dots,N\}$  be a set of vectors in an INNER PRODUCT SPACE (Definition F.9 page 132)  $(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle | \nabla \rangle)$  and with  $\|\mathbf{x}\| \triangleq \sqrt{\langle \mathbf{x} | \mathbf{x} \rangle}$ .

<sup>21</sup> Giles (2000) pages 54–55 (3.13 Bessel's inequality), Bollobás (1999) page 147, Aliprantis and Burkinshaw (1998) page 284

<sup>22</sup> Walter and Shen (2001) pages 3–4, Pedersen (2000) page 39, Edwards (1995) pages 94–100, Weyl (1940)



T  
H  
M

$$\left\{ \begin{array}{l} \{ \mathbf{x}_n \} \text{ is} \\ \text{ORTHONORMAL} \\ \text{(Definition G.9 page 156)} \end{array} \right\} \Rightarrow \left\{ \begin{array}{l} 1. \arg \min_{(\alpha_n)_{n=1}^N} \left\| \mathbf{x} - \sum_{n=1}^N \alpha_n \mathbf{x}_n \right\| = \underbrace{(\langle \mathbf{x} | \mathbf{x}_n \rangle)_{n=1}^N}_{\text{best } \alpha_n = \langle \mathbf{x} | \mathbf{x}_n \rangle} \quad \forall \mathbf{x} \in X \quad \text{and} \\ 2. \underbrace{\left( \sum_{n=1}^N \langle \mathbf{x} | \mathbf{x}_n \rangle \mathbf{x}_n \right)}_{\text{approximation}} \perp \underbrace{\left( \mathbf{x} - \sum_{n=1}^N \langle \mathbf{x} | \mathbf{x}_n \rangle \mathbf{x}_n \right)}_{\text{approximation error}} \quad \forall \mathbf{x} \in X \end{array} \right\}$$

 PROOF:

1. Proof that  $(\langle \mathbf{x} | \mathbf{x}_n \rangle)$  is the best sequence:

$$\begin{aligned} & \left\| \mathbf{x} - \sum_{n=1}^N \alpha_n \mathbf{x}_n \right\|^2 \\ &= \|\mathbf{x}\|^2 - 2\Re \left\langle \mathbf{x} \left| \sum_{n=1}^N \alpha_n \mathbf{x}_n \right. \right\rangle + \left\| \sum_{n=1}^N \alpha_n \mathbf{x}_n \right\|^2 \\ &= \|\mathbf{x}\|^2 - 2\Re \left( \sum_{n=1}^N \alpha_n^* \langle \mathbf{x} | \mathbf{x}_n \rangle \right) + \sum_{n=1}^N \|\alpha_n \mathbf{x}_n\|^2 \quad \text{by Pythagorean Theorem} \\ &= \|\mathbf{x}\|^2 - 2\Re \left( \sum_{n=1}^N \alpha_n^* \langle \mathbf{x} | \mathbf{x}_n \rangle \right) + \sum_{n=1}^N |\alpha_n|^2 + \underbrace{\left[ \sum_{n=1}^N |\langle \mathbf{x} | \mathbf{x}_n \rangle|^2 - \sum_{n=1}^N |\langle \mathbf{x} | \mathbf{x}_n \rangle|^2 \right]}_0 \\ &= \left[ \|\mathbf{x}\|^2 - \sum_{n=1}^N |\langle \mathbf{x} | \mathbf{x}_n \rangle|^2 \right] + \sum_{n=1}^N \left[ |\langle \mathbf{x} | \mathbf{x}_n \rangle|^2 - 2\Re [\alpha_n^* \langle \mathbf{x} | \mathbf{x}_n \rangle] + |\alpha_n|^2 \right] \\ &= \left[ \|\mathbf{x}\|^2 - \sum_{n=1}^N |\langle \mathbf{x} | \mathbf{x}_n \rangle|^2 \right] + \sum_{n=1}^N \left[ |\langle \mathbf{x} | \mathbf{x}_n \rangle|^2 - \alpha_n^* \langle \mathbf{x} | \mathbf{x}_n \rangle - \alpha_n \langle \mathbf{x} | \mathbf{x}_n \rangle^* + |\alpha_n|^2 \right] \\ &= \left\| \mathbf{x} - \sum_{n=1}^N \langle \mathbf{x} | \mathbf{x}_n \rangle \mathbf{x}_n \right\|^2 + \sum_{n=1}^N |\langle \mathbf{x} | \mathbf{x}_n \rangle - \alpha_n|^2 \quad \text{by Bessel's Equality} \quad (\text{Theorem G.6 page 157}) \\ &\geq \left\| \mathbf{x} - \sum_{n=1}^N \langle \mathbf{x} | \mathbf{x}_n \rangle \mathbf{x}_n \right\|^2 \end{aligned}$$

2. Proof that the approximation and approximation error are orthogonal:

$$\begin{aligned} \left\langle \sum_{n=1}^N \langle \mathbf{x} | \mathbf{x}_n \rangle \mathbf{x}_n \left| \mathbf{x} - \sum_{n=1}^N \langle \mathbf{x} | \mathbf{x}_n \rangle \mathbf{x}_n \right. \right\rangle &= \left\langle \sum_{n=1}^N \langle \mathbf{x} | \mathbf{x}_n \rangle \mathbf{x}_n \left| \mathbf{x} \right. \right\rangle - \left\langle \sum_{n=1}^N \langle \mathbf{x} | \mathbf{x}_n \rangle \mathbf{x}_n \left| \sum_{n=1}^N \langle \mathbf{x} | \mathbf{x}_n \rangle \mathbf{x}_n \right. \right\rangle \\ &= \sum_{n=1}^N \langle \mathbf{x} | \mathbf{x}_n \rangle^* \langle \mathbf{x} | \mathbf{x}_n \rangle - \sum_{n=1}^N \sum_{m=1}^N \langle \mathbf{x} | \mathbf{x}_n \rangle \langle \mathbf{x} | \mathbf{x}_m \rangle^* \langle \mathbf{x}_n | \mathbf{x}_m \rangle \\ &= \sum_{n=1}^N |\langle \mathbf{x} | \mathbf{x}_n \rangle|^2 - \sum_{n=1}^N \sum_{m=1}^N \langle \mathbf{x} | \mathbf{x}_n \rangle \langle \mathbf{x} | \mathbf{x}_m \rangle^* \bar{\delta}_{nm} \\ &= \sum_{n=1}^N |\langle \mathbf{x} | \mathbf{x}_n \rangle|^2 - \sum_{n=1}^N |\langle \mathbf{x} | \mathbf{x}_n \rangle|^2 \\ &= 0 \end{aligned}$$



## G.5 Orthonormal bases in Hilbert spaces

**Definition G.10.** Let  $\{\mathbf{x}_n \in X \mid n=1,2,\dots,N\}$  be a set of vectors in an INNER PRODUCT SPACE (Definition F.9 page 132)  $\Omega \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \Delta \mid \nabla \rangle)$ .

DEF

The set  $\{\mathbf{x}_n\}$  is an **orthogonal basis** for  $\Omega$  if  $\{\mathbf{x}_n\}$  is ORTHOGONAL and is a SCHAUDER BASIS for  $\Omega$ .

The set  $\{\mathbf{x}_n\}$  is an **orthonormal basis** for  $\Omega$  if  $\{\mathbf{x}_n\}$  is ORTHONORMAL and is a SCHAUDER BASIS for  $\Omega$ .

**Definition G.11.** <sup>23</sup> Let  $H \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \Delta \mid \nabla \rangle)$  be a Hilbert space.

DEF

Suppose there exists a set  $\{\mathbf{x}_n \in X \mid n \in \mathbb{N}\}$  such that  $\mathbf{x} \triangleq \sum_{n=1}^{\infty} \langle \mathbf{x} \mid \mathbf{x}_n \rangle \mathbf{x}_n$ .

Then the quantities  $\langle \mathbf{x} \mid \mathbf{x}_n \rangle$  are called the **Fourier coefficients** of  $\mathbf{x}$  and the sum

$\sum_{n=1}^{\infty} \langle \mathbf{x} \mid \mathbf{x}_n \rangle \mathbf{x}_n$  is called the **Fourier expansion** of  $\mathbf{x}$  or the **Fourier series** for  $\mathbf{x}$ .

**Definition G.12.**

DEF

The **Kronecker delta function**  $\bar{\delta}_n$  is defined as  $\bar{\delta}_n \triangleq \begin{cases} 1 & \text{for } n = 0 \\ 0 & \text{for } n \neq 0 \end{cases}$  and  $\forall n \in \mathbb{Z}$

**Lemma G.2** (Perfect reconstruction). Let  $\{\mathbf{x}_n \in X \mid n \in \mathbb{N}\}$  be a set of vectors in a HILBERT SPACE  $H \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \Delta \mid \nabla \rangle)$ .

LEM

$$\left\{ \begin{array}{l} (1). \ (\mathbf{x}_n) \text{ is a BASIS for } H \\ (2). \ (\mathbf{x}_n) \text{ is ORTHONORMAL} \end{array} \right\} \text{ and } \Rightarrow \mathbf{x} \triangleq \sum_{n=1}^{\infty} \underbrace{\langle \mathbf{x} \mid \mathbf{x}_n \rangle}_{\text{Fourier coefficient}} \mathbf{x}_n \quad \forall \mathbf{x} \in X$$

Fourier expansion

PROOF:

$$\begin{aligned} \langle \mathbf{x} \mid \mathbf{x}_n \rangle &= \left\langle \sum_{m \in \mathbb{Z}} \alpha_m \mathbf{x}_m \mid \mathbf{x}_n \right\rangle && \text{by left hypothesis (1)} \\ &= \sum_{m \in \mathbb{Z}} \alpha_m \langle \mathbf{x}_m \mid \mathbf{x}_n \rangle && \text{by homogeneous property of } \langle \Delta \mid \nabla \rangle \quad (\text{Definition F.9 page 132}) \\ &= \sum_{m \in \mathbb{Z}} \alpha_m \bar{\delta}_{n-m} && \text{by left hypothesis (2)} \quad (\text{Definition G.9 page 156}) \\ &= \alpha_n \end{aligned}$$



**Proposition G.2.** <sup>24</sup> Let  $\{\mathbf{x}_n \in X \mid n \in \mathbb{N}\}$  be a set of vectors in a HILBERT SPACE  $H \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \Delta \mid \nabla \rangle)$ .

<sup>23</sup> Fabian et al. (2010) page 27 (Theorem 1.55), Young (2001) page 6, Young (1980) page 6

<sup>24</sup> Han et al. (2007) pages 93–94 (Proposition 3.11)

P  
R  
P

$$\underbrace{\|x\|^2 \triangleq \sum_{n=1}^{\infty} |\langle x | x_n \rangle|^2}_{\text{PARSEVAL FRAME}} \iff \underbrace{x \triangleq \sum_{n=1}^{\infty} \langle x | x_n \rangle x_n}_{\text{FOURIER EXPANSION (Definition G.11 page 160)}} \quad \forall x \in X$$

 PROOF:

1. Proof that *Parseval frame*  $\iff$  *Fourier expansion*

$$\begin{aligned} \|x\|^2 &\triangleq \langle x | x \rangle && \text{by definition of } \|\cdot\| \\ &= \left\langle \sum_{n=1}^{\infty} \langle x | x_n \rangle x | x_n \right\rangle && \text{by right hypothesis} \\ &\triangleq \sum_{n=1}^{\infty} \langle x | x_n \rangle \langle x | x_n \rangle && \text{by property of } \langle \Delta | \nabla \rangle \\ &\triangleq \sum_{n=1}^{\infty} \langle x | x_n \rangle \langle x | x_n \rangle^* && \text{by property of } \langle \Delta | \nabla \rangle \\ &\triangleq \sum_{n=1}^{\infty} |\langle x | x_n \rangle|^2 && \text{by property of } \mathbb{C} \quad (\text{Definition E.7 page 117}) \end{aligned}$$

2. Proof that *Parseval frame*  $\implies$  *Fourier expansion*

(a) Let  $(e_n)_{n \in \mathbb{N}}$  be the *standard orthonormal basis* such that the  $n$ th element of  $e_n$  is 1 and all other elements are 0.

(b) Let  $\mathbf{M}$  be an operator in  $\mathbf{H}$  such that  $\mathbf{M}x \triangleq \sum_{n=1}^{\infty} \langle x | x_n \rangle e_n$ .

(c) lemma:  $\mathbf{M}$  is *isometric*. Proof:

$$\begin{aligned} \|\mathbf{M}x\|^2 &= \left\| \sum_{n=1}^{\infty} \langle x | x_n \rangle e_n \right\|^2 && \text{by definition of } \mathbf{M} \quad (\text{item (2b) page 161}) \\ &= \sum_{n=1}^{\infty} \|\langle x | x_n \rangle e_n\|^2 && \text{by Pythagorean Theorem} \\ &= \sum_{n=1}^{\infty} |\langle x | x_n \rangle|^2 \|e_n\|^2 && \text{by homogeneous property of } \|\cdot\| \quad (\text{Definition F.5 page 124}) \\ &= \sum_{n=1}^{\infty} |\langle x | x_n \rangle|^2 && \text{by definition of orthonormal} \quad (\text{Definition G.9 page 156}) \\ &= \|x\|^2 && \text{by Parseval frame hypothesis} \\ \implies \mathbf{M} \text{ is isometric} &&& \text{by definition of isometric} \quad (\text{Definition F.13 page 140}) \end{aligned}$$

(d) Let  $(u_n)_{n \in \mathbb{N}}$  be an *orthonormal basis* for  $\mathbf{H}$ .

(e) Proof for *Fourier expansion*:

$$\begin{aligned}
 x &= \sum_{n=1}^{\infty} \langle x | u_n \rangle u_n && \text{by } \textit{Fourier expansion} \text{ (Proposition G.3 page 164)} \\
 &= \sum_{n=1}^{\infty} \langle Mx | Mu_n \rangle u_n && \text{by (2c) lemma page 161 and Theorem F.23 page 141} \\
 &= \sum_{n=1}^{\infty} \left\langle \sum_{m=1}^{\infty} \langle x | x_m \rangle e_m \mid \sum_{k=1}^{\infty} \langle u_n | x_k \rangle e_k \right\rangle u_n && \text{by item (2b) page 161} \\
 &= \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \langle x | x_m \rangle \sum_{k=1}^{\infty} \langle u_n | x_k \rangle^* \langle e_m | e_k \rangle u_n && \text{by Definition F.9 page 132} \\
 &= \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \langle x | x_m \rangle \langle u_n | x_m \rangle^* u_n && \text{by item (2a) page 161 and Definition G.9 page 156} \\
 &= \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \langle x | x_m \rangle \langle x_m | u_n \rangle u_n && \text{by Definition F.9 page 132} \\
 &= \sum_{m=1}^{\infty} \langle x | x_m \rangle \sum_{n=1}^{\infty} \langle x_m | u_n \rangle u_n \\
 &= \sum_{m=1}^{\infty} \langle x | x_m \rangle x_m && \text{by item (2d) page 161}
 \end{aligned}$$

⇒

When is a set of orthonormal vectors in a Hilbert space  $H$  *total*? Theorem G.9 (next) offers some help.

**Theorem G.9** (The Fourier Series Theorem).<sup>25</sup> Let  $\{x_n \in X \mid n \in \mathbb{N}\}$  be a set of vectors in a HILBERT SPACE  $H \triangleq (X, +, \cdot, (\mathbb{F}, \dagger, \dot{\times}), \langle \triangle | \nabla \rangle)$  and let  $\|x\| \triangleq \sqrt{\langle x | x \rangle}$ .

T H M	(A) $\{x_n\}$ is ORTHONORMAL in $H$ $\implies$	
	$\left\{ \begin{array}{l} \iff (1). \quad (\text{span}\{x_n\})^- = H \\ \iff (2). \quad \langle x   y \rangle \stackrel{*}{=} \sum_{n=1}^{\infty} \langle x   x_n \rangle \langle y   x_n \rangle^* \quad \forall x, y \in X \\ \iff (3). \quad \ x\ ^2 \stackrel{*}{=} \sum_{n=1}^{\infty}  \langle x   x_n \rangle ^2 \quad \forall x \in X \\ \iff (4). \quad x \stackrel{*}{=} \sum_{n=1}^{\infty} \langle x   x_n \rangle x_n \quad \forall x \in X \end{array} \right.$	$\left\{ \begin{array}{l} \{x_n\} \text{ is TOTAL in } H \\ \text{(GENERALIZED PARSEVAL'S IDENTITY)} \\ \text{(PARSEVAL'S IDENTITY)} \\ \text{(FOURIER SERIES EXPANSION)} \end{array} \right.$

✎ PROOF:

<sup>25</sup> [Bachman and Narici \(1966\) pages 149–155](#) (Theorem 9.12), [Kubrusly \(2001\) pages 360–363](#) (Theorem 5.48), [Aliprantis and Burkinshaw \(1998\) pages 298–299](#) (Theorem 34.2), [Christensen \(2003\) page 57](#) (Theorem 3.4.2), [Berberian \(1961\) pages 52–53](#) (Theorem II§8.3), [Heil \(2011\) pages 34–35](#) (Theorem 1.50), [Bracewell \(1978\) page 112](#) (Rayleigh's theorem)

1. Proof that (1)  $\implies$  (2):

$$\begin{aligned}
 \langle \mathbf{x} | \mathbf{y} \rangle &\triangleq \left\langle \sum_{n=1}^{\infty} \langle \mathbf{x} | \mathbf{x}_n \rangle \mathbf{x}_n \mid \sum_{m=1}^{\infty} \langle \mathbf{y} | \mathbf{x}_m \rangle \mathbf{x}_m \right\rangle && \text{by (A) and (1)} \\
 &\triangleq \sum_{n=1}^{\infty} \langle \mathbf{x} | \mathbf{x}_n \rangle \left\langle \mathbf{x}_n \mid \sum_{m=1}^{\infty} \langle \mathbf{y} | \mathbf{x}_m \rangle \mathbf{x}_m \right\rangle && \text{by property of } \langle \cdot | \cdot \rangle \quad (\text{Definition F.9 page 132}) \\
 &\triangleq \sum_{n=1}^{\infty} \langle \mathbf{x} | \mathbf{x}_n \rangle \sum_{m=1}^{\infty} \langle \mathbf{y} | \mathbf{x}_m \rangle^* \langle \mathbf{x}_n | \mathbf{x}_m \rangle && \text{by property of } \langle \cdot | \cdot \rangle \quad (\text{Definition F.9 page 132}) \\
 &\triangleq \sum_{n=1}^{\infty} \langle \mathbf{x} | \mathbf{x}_n \rangle \sum_{m=1}^{\infty} \langle \mathbf{y} | \mathbf{x}_m \rangle^* \bar{\delta}_{mn} && \text{by (A)} \\
 &\triangleq \sum_{n=1}^{\infty} \langle \mathbf{x} | \mathbf{x}_n \rangle \langle \mathbf{y} | \mathbf{x}_n \rangle^* && \text{by definition of } \bar{\delta}_n \quad (\text{Definition G.12 page 160})
 \end{aligned}$$

2. Proof that (2)  $\implies$  (3):

$$\begin{aligned}
 \|\mathbf{x}\|^2 &\triangleq \langle \mathbf{x} | \mathbf{x} \rangle && \text{by definition of induced norm} \\
 &= \sum_{n=1}^{\infty} \langle \mathbf{x} | \mathbf{x}_n \rangle \langle \mathbf{x} | \mathbf{x}_n \rangle^* && \text{by (2)} \\
 &= \sum_{n=1}^{\infty} |\langle \mathbf{x} | \mathbf{x}_n \rangle|^2
 \end{aligned}$$

3. Proof that (3)  $\iff$  (4) *not* using (A): by Proposition G.2 page 160

4. Proof that (3)  $\implies$  (1) (proof by contradiction):

- (a) Suppose  $\{\mathbf{x}_n\}$  is *not total*.
- (b) Then there must exist a vector  $\mathbf{y}$  in  $\mathbf{H}$  such that the set  $B \triangleq \{\mathbf{x}_n\} \cup \mathbf{y}$  is *orthonormal*.
- (c) Then  $1 = \|\mathbf{y}\|^2 \neq \sum_{n=1}^{\infty} |\langle \mathbf{y} | \mathbf{x}_n \rangle|^2 = 0$ .
- (d) But this contradicts (3), and so  $\{\mathbf{x}_n\}$  must be *total* and (3)  $\implies$  (1).

5. Extraneous proof that (3)  $\implies$  (4) (this proof is not really necessary here):

$$\begin{aligned}
 \left\| \mathbf{x} - \sum_{n=1}^{\infty} \langle \mathbf{x} | \mathbf{x}_n \rangle \mathbf{x}_n \right\|^2 &= \|\mathbf{x}\|^2 - \sum_{n=1}^{\infty} |\langle \mathbf{x} | \mathbf{x}_n \rangle|^2 && \text{by Bessel's Equality} \quad (\text{Theorem G.6 page 157}) \\
 &= 0 && \text{by (3)} \\
 \implies \mathbf{x} &\triangleq \sum_{n=1}^{\infty} \langle \mathbf{x} | \mathbf{x}_n \rangle \mathbf{x}_n && \text{by definition of } \triangleq
 \end{aligned}$$

6. Extraneous proof that (A)  $\implies$  (4) (this proof is not really necessary here)

- (a) The sequence  $\sum_{n=1}^N |\langle \mathbf{x} | \mathbf{x}_n \rangle|^2$  is *monotonically increasing* in  $n$ .
- (b) By Bessel's inequality (page 158), the sequence is upper bounded by  $\|\mathbf{x}\|^2$ :

$$\sum_{n=1}^N |\langle \mathbf{x} | \mathbf{x}_n \rangle|^2 \leq \|\mathbf{x}\|^2$$

- (c) Because this sequence is both monotonically increasing and bounded in  $n$ , it must equal its bound in the limit as  $n$  approaches infinity:

$$\lim_{N \rightarrow \infty} \sum_{n=1}^N |\langle \mathbf{x} | \mathbf{x}_n \rangle|^2 = \|\mathbf{x}\|^2 \quad (\text{G.1})$$

- (d) If we combine this result with *Bessel's Equality* (Theorem G.6 page 157) we have

$$\begin{aligned} \lim_{N \rightarrow \infty} \left\| \mathbf{x} - \sum_{n=1}^N \langle \mathbf{x} | \mathbf{x}_n \rangle \mathbf{x}_n \right\|^2 &= \|\mathbf{x}\|^2 - \lim_{N \rightarrow \infty} \sum_{n=1}^N |\langle \mathbf{x} | \mathbf{x}_n \rangle|^2 \quad \text{by Bessel's Equality (Theorem G.6 page 157)} \\ &= \|\mathbf{x}\|^2 - \|\mathbf{x}\|^2 \quad \text{by equation (G.1) page 164} \\ &= 0 \end{aligned}$$

⇒

**Proposition G.3** (Fourier expansion). *Let  $\{\mathbf{x}_n \in X \mid n \in \mathbb{N}\}$  be a set of vectors in a HILBERT SPACE  $(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \Delta | \nabla \rangle)$ .*

$$\underbrace{\{\mathbf{x}_n\} \text{ is an ORTHONORMAL BASIS for } H}_{(A)} \implies \underbrace{\left\{ \mathbf{x} \doteq \sum_{n=1}^{\infty} \alpha_n \mathbf{x}_n \right\}}_{(1)} \iff \underbrace{\left\{ \alpha_n = \langle \mathbf{x} | \mathbf{x}_n \rangle \right\}}_{(2)}$$

✎ PROOF:

1. Proof that (1)  $\implies$  (2): by Lemma G.2 page 160
2. Proof that (1)  $\impliedby$  (2):

$$\begin{aligned} \left\| \mathbf{x} - \sum_{n=1}^{\infty} \alpha_n \mathbf{x}_n \right\|^2 &= \left\| \mathbf{x} - \sum_{n=1}^{\infty} \langle \mathbf{x} | \mathbf{x}_n \rangle \mathbf{x}_n \right\|^2 \quad \text{by right hypothesis} \\ &= \|\mathbf{x}\|^2 - \sum_{n=1}^{\infty} |\langle \mathbf{x} | \mathbf{x}_n \rangle|^2 \quad \text{by Bessel's equality} \quad (\text{Theorem G.6 page 157}) \\ &= 0 \quad \text{by Parseval's Identity} \quad (\text{Theorem G.9 page 162}) \\ &\stackrel{\text{def}}{\iff} \mathbf{x} \doteq \sum_{n=1}^{\infty} \langle \mathbf{x} | \mathbf{x}_n \rangle \mathbf{x}_n \quad \text{by definition of strong convergence} \end{aligned}$$

⇒

**Proposition G.4** (Riesz-Fischer Theorem).<sup>26</sup> *Let  $\{\mathbf{x}_n \in X \mid n \in \mathbb{N}\}$  be a set of vectors in a HILBERT SPACE  $(X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \Delta | \nabla \rangle)$ .*

$$\underbrace{\{\mathbf{x}_n\} \text{ is an ORTHONORMAL BASIS for } H}_{(A)} \implies \underbrace{\left\{ \sum_{n=1}^{\infty} |\alpha_n|^2 < \infty \right\}}_{(1)} \iff \underbrace{\left\{ \exists \mathbf{x} \in H \text{ such that } \alpha_n = \langle \mathbf{x} | \mathbf{x}_n \rangle \right\}}_{(2)}$$

✎ PROOF:

<sup>26</sup> Young (2001) page 6

1. Proof that (1)  $\implies$  (2):

(a) If (1) is true, then let  $\mathbf{x} \triangleq \sum_{n \in \mathbb{N}} \alpha_n \mathbf{x}_n$ .

(b) Then

$$\begin{aligned}
 \langle \mathbf{x} | \mathbf{x}_n \rangle &= \left\langle \sum_{m \in \mathbb{N}} \alpha_m \mathbf{x}_m | \mathbf{x}_n \right\rangle && \text{by definition of } \mathbf{x} \\
 &= \sum_{m \in \mathbb{N}} \alpha_m \langle \mathbf{x}_m | \mathbf{x}_n \rangle && \text{by homogeneous property of } \langle \Delta | \nabla \rangle \quad (\text{Definition F.9 page 132}) \\
 &= \sum_{m \in \mathbb{N}} \alpha_m \bar{\delta}_{mn} && \text{by (A)} \\
 &= \sum_{m \in \mathbb{N}} \alpha_n && \text{by definition of } \bar{\delta} \quad (\text{Definition G.12 page 160})
 \end{aligned}$$

2. Proof that (1)  $\Longleftarrow$  (2):

$$\begin{aligned}
 \sum_{n \in \mathbb{N}} |\alpha_n|^2 &= \sum_{n \in \mathbb{N}} |\langle \mathbf{x} | \mathbf{x}_n \rangle|^2 && \text{by (2)} \\
 &\leq \|\mathbf{x}\|^2 && \text{by Bessel's Inequality} \quad (\text{Theorem G.7 page 158}) \\
 &\leq \infty
 \end{aligned}$$

$\Rightarrow$

### Theorem G.10. <sup>27</sup>

All SEPARABLE HILBERT SPACES are ISOMORPHIC. That is,

T H M	$  \left\{ \begin{array}{l} \mathbf{X} \text{ is a separable} \\ \text{Hilbert space} \end{array} \right. \text{ and } \left\{ \begin{array}{l} \mathbf{Y} \text{ is a separable} \\ \text{Hilbert space} \end{array} \right.  $	$\implies$	$  \left\{ \begin{array}{l} \text{there is a BIJECTIVE operator } \mathbf{M} \in \mathbf{Y}^{\mathbf{X}} \text{ such that} \\ (1). \quad \mathbf{y} = \mathbf{M}\mathbf{x} \quad \forall \mathbf{x} \in \mathbf{X}, \mathbf{y} \in \mathbf{Y} \text{ and} \\ (2). \quad \ \mathbf{M}\mathbf{x}\  = \ \mathbf{x}\  \quad \forall \mathbf{x} \in \mathbf{X} \text{ and} \\ (3). \quad \langle \mathbf{M}\mathbf{x}   \mathbf{M}\mathbf{y} \rangle = \langle \mathbf{x}   \mathbf{y} \rangle \quad \forall \mathbf{x} \in \mathbf{X}, \mathbf{y} \in \mathbf{Y} \end{array} \right.  $
-------------	--	------------	---

 PROOF:

1. Let  $\mathbf{X} \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \Delta | \nabla \rangle)$  be a *separable Hilbert space* with *orthonormal basis*  $\{\mathbf{x}_n\}_{n \in \mathbb{N}}$ .  
Let  $\mathbf{Y} \triangleq (Y, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \Delta | \nabla \rangle)$  be a *separable Hilbert space* with *orthonormal basis*  $\{\mathbf{y}_n\}_{n \in \mathbb{N}}$ .


2. Proof that there exists *bijective* operator  $\mathbf{M}$  and its inverse  $\mathbf{M}^{-1}$  between  $\{\mathbf{x}_n\}$  and  $\{\mathbf{y}_n\}$ :

(a) Let  $\mathbf{M}$  be defined such that  $\mathbf{y}_n \triangleq \mathbf{M}\mathbf{x}_n$ .

(b) Thus  $\mathbf{M}$  is a *bijection* between  $\{\mathbf{x}_n\}$  and  $\{\mathbf{y}_n\}$ .

(c) Because  $\mathbf{M}$  is a *bijection* between  $\{\mathbf{x}_n\}$  and  $\{\mathbf{y}_n\}$ ,  $\mathbf{M}$  has an inverse operator  $\mathbf{M}^{-1}$  between  $\{\mathbf{x}_n\}$  and  $\{\mathbf{y}_n\}$  such that  $\mathbf{x}_n = \mathbf{M}^{-1}\mathbf{y}_n$ .

3. Proof that  $\mathbf{M}$  and  $\mathbf{M}^{-1}$  are *bijective* operators between  $\mathbf{X}$  and  $\mathbf{Y}$ :

<sup>27</sup>  Young (2001) page 6

(a) Proof that  $\mathbf{M}$  maps  $\mathbf{X}$  into  $\mathbf{Y}$ :

$$\begin{aligned}
 \mathbf{x} \in \mathbf{X} &\iff \mathbf{x} \triangleq \sum_{n \in \mathbb{N}} \langle \mathbf{x} | \mathbf{x}_n \rangle \mathbf{x}_n && \text{by Fourier expansion} && (\text{Theorem G.9 page 162}) \\
 &\implies \exists \mathbf{y} \in \mathbf{Y} \text{ such that } \langle \mathbf{y} | \mathbf{y}_n \rangle = \langle \mathbf{x} | \mathbf{x}_n \rangle && \text{by Riesz-Fischer Thm.} && (\text{Proposition G.4 page 164}) \\
 &\implies \\
 \mathbf{y} &= \sum_{n \in \mathbb{N}} \langle \mathbf{y} | \mathbf{y}_n \rangle \mathbf{y}_n && \text{by Fourier expansion} && (\text{Theorem G.9 page 162}) \\
 &= \sum_{n \in \mathbb{N}} \langle \mathbf{x} | \mathbf{x}_n \rangle \mathbf{y}_n && \text{by Riesz-Fischer Thm.} && (\text{Proposition G.4 page 164}) \\
 &= \sum_{n \in \mathbb{N}} \langle \mathbf{x} | \mathbf{x}_n \rangle \mathbf{M} \mathbf{x}_n && \text{by definition of } \mathbf{M} && (\text{item (2a) page 165}) \\
 &= \mathbf{M} \sum_{n \in \mathbb{N}} \langle \mathbf{x} | \mathbf{x}_n \rangle \mathbf{x}_n && \text{by prop. of linear ops.} && (\text{Theorem F.1 page 121}) \\
 &= \mathbf{M} \mathbf{x} && \text{by definition of } \mathbf{x}
 \end{aligned}$$

(b) Proof that  $\mathbf{M}^{-1}$  maps  $\mathbf{Y}$  into  $\mathbf{X}$ :

$$\begin{aligned}
 \mathbf{y} \in \mathbf{Y} &\iff \mathbf{y} \triangleq \sum_{n \in \mathbb{N}} \langle \mathbf{y} | \mathbf{y}_n \rangle \mathbf{y}_n && \text{by Fourier expansion} && (\text{Theorem G.9 page 162}) \\
 &\implies \exists \mathbf{x} \in \mathbf{X} \text{ such that } \langle \mathbf{x} | \mathbf{x}_n \rangle = \langle \mathbf{y} | \mathbf{y}_n \rangle && \text{by Riesz-Fischer Thm.} && (\text{Proposition G.4 page 164}) \\
 &\implies \\
 \mathbf{x} &= \sum_{n \in \mathbb{N}} \langle \mathbf{x} | \mathbf{x}_n \rangle \mathbf{x}_n && \text{by Fourier expansion} && (\text{Theorem G.9 page 162}) \\
 &= \sum_{n \in \mathbb{N}} \langle \mathbf{y} | \mathbf{y}_n \rangle \mathbf{x}_n && \text{by Riesz-Fischer Thm.} && (\text{Proposition G.4 page 164}) \\
 &= \sum_{n \in \mathbb{N}} \langle \mathbf{y} | \mathbf{y}_n \rangle \mathbf{M}^{-1} \mathbf{y}_n && \text{by definition of } \mathbf{M}^{-1} && (\text{item (2c) page 165}) \\
 &= \mathbf{M}^{-1} \sum_{n \in \mathbb{N}} \langle \mathbf{y} | \mathbf{y}_n \rangle \mathbf{y}_n && \text{by prop. of linear ops.} && (\text{Theorem F.1 page 121}) \\
 &= \mathbf{M}^{-1} \mathbf{y} && \text{by definition of } \mathbf{y}
 \end{aligned}$$

4. Proof for (2):

$$\begin{aligned}
 \|\mathbf{M} \mathbf{x}\|^2 &= \left\| \mathbf{M} \sum_{n \in \mathbb{N}} \langle \mathbf{x} | \mathbf{x}_n \rangle \mathbf{x}_n \right\|^2 && \text{by Fourier expansion} && (\text{Theorem G.9 page 162}) \\
 &= \left\| \sum_{n \in \mathbb{N}} \langle \mathbf{x} | \mathbf{x}_n \rangle \mathbf{M} \mathbf{x}_n \right\|^2 && \text{by property of linear operators} && (\text{Theorem F.1 page 121}) \\
 &= \left\| \sum_{n \in \mathbb{N}} \langle \mathbf{x} | \mathbf{x}_n \rangle \mathbf{y}_n \right\|^2 && \text{by definition of } \mathbf{M} && (\text{item (2a) page 165}) \\
 &= \sum_{n \in \mathbb{N}} |\langle \mathbf{x} | \mathbf{x}_n \rangle|^2 && \text{by Parseval's Identity} && (\text{Proposition G.4 page 164}) \\
 &= \left\| \sum_{n \in \mathbb{N}} \langle \mathbf{x} | \mathbf{x}_n \rangle \mathbf{x}_n \right\|^2 && \text{by Parseval's Identity} && (\text{Proposition G.4 page 164}) \\
 &= \|\mathbf{x}\|^2 && \text{by Fourier expansion} && (\text{Theorem G.9 page 162})
 \end{aligned}$$

5. Proof for (3): by (2) and Theorem F23 page 141



**Theorem G.11.** <sup>28</sup> Let  $H$  be a HILBERT SPACE.

**T H M**  $H$  has a SCHAUDER BASIS  $\iff H$  is SEPARABLE

**Theorem G.12.** <sup>29</sup> Let  $H$  be a HILBERT SPACE.

**T H M**  $H$  has an ORTHONORMAL BASIS  $\iff H$  is SEPARABLE

## G.6 Riesz bases in Hilbert spaces

**Definition G.13.** <sup>30</sup> Let  $\{x_n \in X \mid n \in \mathbb{N}\}$  be a set of vectors in a SEPARABLE HILBERT SPACE  $H \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \Delta \mid \nabla \rangle)$ .

**D E F**  $\{x_n\}$  is a **Riesz basis** for  $H$  if  $\{x_n\}$  is EQUIVALENT (Definition G.8 page 155) to some ORTHONORMAL BASIS (Definition G.10 page 160) in  $H$ .

**Definition G.14.** <sup>31</sup> Let  $(x_n \in X)_{n \in \mathbb{N}}$  be a sequence of vectors in a SEPARABLE HILBERT SPACE  $H \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \Delta \mid \nabla \rangle)$ .

**D E F** The sequence  $(x_n)$  is a **Riesz sequence** for  $H$  if

$$\exists A, B \in \mathbb{R}^+ \text{ such that } A \sum_{n=1}^{\infty} |\alpha_n|^2 \leq \left\| \sum_{n=1}^{\infty} \alpha_n x_n \right\|^2 \leq B \sum_{n=1}^{\infty} |\alpha_n|^2 \quad \forall (\alpha_n)_{n \in \mathbb{N}} \in \ell_{\mathbb{F}}^2.$$

**Definition G.15.** Let  $X \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \Delta \mid \nabla \rangle)$  be an INNER PRODUCT SPACE (Definition F.9 page 132).

**D E F** The sequences  $(x_n \in X)_{n \in \mathbb{Z}}$  and  $(y_n \in X)_{n \in \mathbb{Z}}$  are **biorthogonal** with respect to each other in  $X$  if  $\langle x_n \mid y_m \rangle = \delta_{nm}$

**Lemma G.3.** <sup>32</sup> Let  $\{x_n \mid n \in \mathbb{N}\}$  be a sequence in a HILBERT SPACE  $X \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \Delta \mid \nabla \rangle)$ . Let  $\{y_n \mid n \in \mathbb{N}\}$  be a sequence in a HILBERT SPACE  $Y \triangleq (Y, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \Delta \mid \nabla \rangle)$ . Let

**L E M**  $\left\{ \begin{array}{l} \text{(i). } \{x_n\} \text{ is TOTAL in } X \\ \text{(ii). There exists } A > 0 \text{ such that } A \sum_{n \in C} |a_n|^2 \leq \left\| \sum_{n \in C} a_n x_n \right\|^2 \text{ for finite } C \\ \text{(iii). There exists } B > 0 \text{ such that } \left\| \sum_{n=1}^{\infty} b_n y_n \right\|^2 \leq B \sum_{n=1}^{\infty} |b_n|^2 \quad \forall (b_n)_{n \in \mathbb{N}} \in \ell_{\mathbb{F}}^2 \end{array} \right\} \implies$

$\left\{ \begin{array}{l} \text{(1). } \mathbf{R}^\circ \text{ is a linear bounded operator that maps from } \text{span}\{x_n\} \text{ to } \text{span}\{y_n\} \\ \text{where } \mathbf{R}^\circ \sum_{n \in C} c_n x_n \triangleq \sum_{n \in C} c_n y_n, \text{ for some sequence } (c_n) \text{ and finite set } C \\ \text{(2). } \mathbf{R} \text{ has a unique extension to a bounded operator } \mathbf{R} \text{ that maps from } X \text{ to } Y \\ \text{(3). } \|\mathbf{R}^\circ\| \leq \frac{B}{A} \\ \text{(4). } \|\mathbf{R}\| \leq \frac{B}{A} \end{array} \right\}$  and

<sup>28</sup> Bachman et al. (2000) page 112 (3.4.8), Berberian (1961) page 53 (Theorem II\$8.3)

<sup>29</sup> Kubrusly (2001) page 357 (Proposition 5.43)

<sup>30</sup> Young (2001) page 27 (Definition 1.8.2), Christensen (2003) page 63 (Definition 3.6.1), Heil (2011) page 196 (Definition 7.9)

<sup>31</sup> Christensen (2003) pages 66–68 (page 68 and (3.24) on page 66), Wojtaszczyk (1997) page 20 (Definition 2.6)

<sup>32</sup> Christensen (2003) pages 65–66 (Lemma 3.6.5)

**Theorem G.13.** <sup>33</sup> Let  $\{\mathbf{x}_n \in X \mid n \in \mathbb{N}\}$  be a set of vectors in a SEPARABLE HILBERT SPACE  $H \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \Delta \mid \nabla \rangle)$ .

T  
H  
M

$$\left\{ \begin{array}{l} \{\mathbf{x}_n\} \text{ is a RIESZ BASIS} \\ \text{for } H \end{array} \right\} \iff \left\{ \begin{array}{l} (1). \quad \{\mathbf{x}_n\} \text{ is TOTAL in } H \\ (2). \quad \exists A, B \in \mathbb{R}^+ \text{ such that } \forall (\alpha_n) \in \ell_{\mathbb{F}}^2, \\ A \sum_{n=1}^{\infty} |\alpha_n|^2 \leq \left\| \sum_{n=1}^{\infty} \alpha_n \mathbf{x}_n \right\|^2 \leq B \sum_{n=1}^{\infty} |\alpha_n|^2 \end{array} \right\} \text{ and }$$

PROOF:

1. Proof for  $(\implies)$  case:

(a) Proof that *Riesz basis* hypothesis  $\implies$  (1): all bases for  $H$  are *total* in  $H$ .

(b) Proof that *Riesz basis* hypothesis  $\implies$  (2):

i. Let  $(\mathbf{u}_n)_{n \in \mathbb{N}}$  be an *orthonormal basis* for  $H$ .

ii. Let  $\mathbf{R}$  be a *bounded bijective operator* such that  $\mathbf{x}_n = \mathbf{R}\mathbf{u}_n$ .

iii. Proof for upper bound  $B$ :

$$\begin{aligned} \left\| \sum_{n=1}^{\infty} \alpha_n \mathbf{x}_n \right\|^2 &= \left\| \sum_{n=1}^{\infty} \alpha_n \mathbf{R}\mathbf{u}_n \right\|^2 && \text{by definition of } \mathbf{R} && \text{(item (1(b)iii))} \\ &= \left\| \mathbf{R} \sum_{n=1}^{\infty} \alpha_n \mathbf{u}_n \right\|^2 && \text{by Theorem F.1 page 121} \\ &\leq \|\mathbf{R}\|^2 \left\| \sum_{n=1}^{\infty} \alpha_n \mathbf{u}_n \right\|^2 && \text{by Theorem F.6 page 127} \\ &= \|\mathbf{R}\|^2 \sum_{n=1}^{\infty} \|\alpha_n \mathbf{u}_n\|^2 && \text{by Pythagorean Theorem} \\ &= \|\mathbf{R}\|^2 \sum_{n=1}^{\infty} |\alpha_n|^2 \|\mathbf{u}_n\|^2 && \text{by homogeneous property of norms (Definition F.5 page 124)} \\ &= \underbrace{\|\mathbf{R}\|^2}_B \sum_{n=1}^{\infty} |\alpha_n|^2 && \text{by definition of orthonormality (Definition G.9 page 156)} \end{aligned}$$

iv. Proof for lower bound  $A$ :

$$\begin{aligned} \left\| \sum_{n=1}^{\infty} \alpha_n \mathbf{x}_n \right\|^2 &= \frac{\|\mathbf{R}^{-1}\|^2}{\|\mathbf{R}^{-1}\|^2} \left\| \sum_{n=1}^{\infty} \alpha_n \mathbf{x}_n \right\|^2 && \text{because } \|\mathbf{R}^{-1}\| > 0 && \text{(Proposition F.1 page 125)} \\ &\geq \frac{1}{\|\mathbf{R}^{-1}\|^2} \left\| \mathbf{R}^{-1} \sum_{n=1}^{\infty} \alpha_n \mathbf{x}_n \right\|^2 && \text{by Theorem F.6 page 127} \\ &= \frac{1}{\|\mathbf{R}^{-1}\|^2} \left\| \mathbf{R}^{-1} \sum_{n=1}^{\infty} \alpha_n \mathbf{R}\mathbf{u}_n \right\|^2 && \text{by definition of } \mathbf{R} && \text{(item (1(b)ii) page 168)} \\ &= \frac{1}{\|\mathbf{R}^{-1}\|^2} \left\| \mathbf{R}^{-1} \mathbf{R} \sum_{n=1}^{\infty} \alpha_n \mathbf{u}_n \right\|^2 && \text{by property of linear operators (Theorem F.1 page 121)} \end{aligned}$$

<sup>33</sup> Young (2001) page 27 (Theorem 1.8.9), Christensen (2003) page 66 (Theorem 3.6.6), Heil (2011) pages 197–198 (Theorem 7.13), Christensen (2008) pages 61–62 (Theorem 3.3.7)

$$\begin{aligned}
&= \frac{1}{\|\mathbf{R}^{-1}\|^2} \left\| \sum_{n=1}^{\infty} \alpha_n \mathbf{u}_n \right\|^2 && \text{by definition of inverse op.} && (\text{Definition F.3 page 120}) \\
&= \frac{1}{\|\mathbf{R}^{-1}\|^2} \sum_{n=1}^{\infty} \|\alpha_n \mathbf{u}_n\|^2 && \text{by Pythagorean Theorem} \\
&= \frac{1}{\|\mathbf{R}^{-1}\|^2} \sum_{n=1}^{\infty} |\alpha_n|^2 \|\mathbf{u}_n\|^2 && \text{by } \|\cdot\| \text{ homogeneous prop.} && (\text{Definition F.5 page 124}) \\
&= \underbrace{\frac{1}{\|\mathbf{R}^{-1}\|^2}}_A \sum_{n=1}^{\infty} |\alpha_n|^2 && \text{by def. of orthonormality} && (\text{Definition G.9 page 156})
\end{aligned}$$

2. Proof for ( $\implies$ ) case:

- Let  $\{\mathbf{u}_n | n \in \mathbb{N}\}$  be an *orthonormal basis* for  $\mathbf{H}$ .
- Using (2) and Lemma G.3 (page 167), construct an bounded extension operator  $\mathbf{R}$  such that  $\mathbf{R}\mathbf{u}_n = \mathbf{x}_n$  for all  $n \in \mathbb{N}$ .
- Using (2) and Lemma G.3 (page 167), construct an bounded extension operator  $\mathbf{S}$  such that  $\mathbf{S}\mathbf{x}_n = \mathbf{u}_n$  for all  $n \in \mathbb{N}$ .
- Then,  $\mathbf{R}\mathbf{V}\mathbf{x} = \mathbf{V}\mathbf{R}\mathbf{x} \implies \mathbf{V} = \mathbf{R}^{-1}$ , and so  $\mathbf{R}$  is a bounded invertible operator
- and  $\{\mathbf{x}_n\}$  is a *Riesz sequence*.

$\Rightarrow$

**Theorem G.14.** <sup>34</sup> Let  $\mathbf{H} \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \Delta | \nabla \rangle)$  be a SEPARABLE HILBERT SPACE.

T H M	$ \left\{ \begin{array}{l} (\mathbf{x}_n \in \mathbf{H})_{n \in \mathbb{Z}} \text{ is a} \\ \text{RIESZ BASIS for } \mathbf{H} \end{array} \right\} \implies \left\{ \begin{array}{l} \text{There exists } (\mathbf{y}_n \in \mathbf{H})_{n \in \mathbb{Z}} \text{ such that} \\ \begin{array}{ll} (1). \ (\mathbf{x}_n) \text{ and } (\mathbf{y}_n) \text{ are BIORTHOGONAL} & \text{and} \\ (2). \ (\mathbf{y}_n) \text{ is also a RIESZ BASIS for } \mathbf{H} & \text{and} \\ (3). \ \exists B > A > 0 \text{ such that} & \\ \quad A \sum_{n=1}^{\infty}  a_n ^2 \leq \left\  \sum_{n=1}^{\infty} a_n \mathbf{x}_n \right\ ^2 = \left\  \sum_{n=1}^{\infty} a_n \mathbf{y}_n \right\ ^2 \leq B \sum_{n=1}^{\infty}  a_n ^2 & \\ \quad \forall (a_n)_{n \in \mathbb{N}} \in \ell_{\mathbb{F}}^2 & \end{array} \end{array} \right\} $
-------------	--

$\pencil$  PROOF:

1. Proof for (1):

- Let  $\mathbf{e}_n$  be the *unit vector* in  $\mathbf{H}$  such that the  $n$ th element of  $\mathbf{e}_n$  is 1 and all other elements are 0.
- Let  $\mathbf{M}$  be an operator on  $\mathbf{H}$  such that  $\mathbf{M}\mathbf{e}_n = \mathbf{x}_n$ .
- Note that  $\mathbf{M}$  is *isometric*, and as such  $\|\mathbf{M}\mathbf{x}\| = \|\mathbf{x}\| \quad \forall \mathbf{x} \in \mathbf{H}$ .
- Let  $\mathbf{y}_n \triangleq (\mathbf{M}^{-1})^*$ .
- Then,

$$\begin{aligned}
\langle \mathbf{y}_n | \mathbf{x}_m \rangle &= \langle (\mathbf{M}^{-1})^* \mathbf{e}_n | \mathbf{M}\mathbf{e}_m \rangle \\
&= \langle \mathbf{e}_n | \mathbf{M}^{-1} \mathbf{M}\mathbf{e}_m \rangle \\
&= \langle \mathbf{e}_n | \mathbf{e}_m \rangle \\
&= \bar{\delta}_{nm} \\
&\implies \{\mathbf{x}_n\} \text{ and } \{\mathbf{y}_n\} \text{ are biorthogonal}
\end{aligned}$$

by Definition G.9 page 156

<sup>34</sup> Wojtaszczyk (1997) page 20 (Lemma 2.7(a))

2. Proof for (3):

$$\begin{aligned}
 \left\| \sum_{n \in \mathbb{Z}} \alpha_n \mathbf{y}_n \right\| &= \left\| \sum_{n \in \mathbb{Z}} \alpha_n (\mathbf{M}^{-1})^* \mathbf{e}_n \right\| && \text{by definition of } \mathbf{y}_n && \text{(Proposition 1d page 169)} \\
 &= \left\| (\mathbf{M}^{-1})^* \sum_{n \in \mathbb{Z}} \alpha_n \mathbf{e}_n \right\| && \text{by property of linear ops.} \\
 &= \left\| \sum_{n \in \mathbb{Z}} \alpha_n \mathbf{e}_n \right\| && \text{because } (\mathbf{M}^{-1})^* \text{ is isometric} && \text{(Definition F.13 page 140)} \\
 &= \left\| \mathbf{M} \sum_{n \in \mathbb{Z}} \alpha_n \mathbf{e}_n \right\| && \text{because } \mathbf{M} \text{ is isometric} && \text{(Definition F.13 page 140)} \\
 &= \left\| \sum_{n \in \mathbb{Z}} \alpha_n \mathbf{M} \mathbf{e}_n \right\| && \text{by property of linear operators} \\
 &= \left\| \sum_{n \in \mathbb{Z}} \alpha_n \mathbf{x}_n \right\| && \text{by definition of } \mathbf{M} \\
 &\Rightarrow \{ \mathbf{y}_n \} \text{ is a Riesz basis} && \text{by left hypothesis}
 \end{aligned}$$

3. Proof for (2): by (3) and definition of *Riesz basis* (Definition G.13 page 167)

⇒

**Proposition G.5.** <sup>35</sup> Let  $\{ \mathbf{x}_n | n \in \mathbb{N} \}$  be a set of vectors in a HILBERT SPACE  $\mathbf{H} \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle | \nabla \rangle)$ .

$$\left\{ \begin{array}{l} \{ \mathbf{x}_n \} \text{ is a RIESZ BASIS for } \mathbf{H} \text{ with} \\ A \sum_{n=1}^{\infty} |a_n|^2 \leq \left\| \sum_{n=1}^{\infty} a_n \mathbf{x}_n \right\|^2 \leq B \sum_{n=1}^{\infty} |a_n|^2 \\ \forall \{ a_n \} \in \ell_{\mathbb{F}}^2 \end{array} \right\} \Rightarrow \left\{ \begin{array}{l} \{ \mathbf{x}_n \} \text{ is a FRAME for } \mathbf{H} \text{ with} \\ \frac{1}{B} \|\mathbf{x}\|^2 \leq \sum_{n=1}^{\infty} |\langle \mathbf{x} | \mathbf{x}_n \rangle|^2 \leq \frac{1}{A} \|\mathbf{x}\|^2 \\ \underbrace{\hspace{10em}}_{\text{STABILITY CONDITION}} \\ \forall \mathbf{x} \in \mathbf{H} \end{array} \right\}$$

PROOF:

1. Let  $\{ \mathbf{y}_n | n \in \mathbb{N} \}$  be a *Riesz basis* that is *biorthogonal* to  $\{ \mathbf{x}_n | n \in \mathbb{N} \}$  (Theorem G.14 page 169).

2. Let  $\mathbf{x} \triangleq \sum_{n=1}^{\infty} a_n \mathbf{y}_n$ .

3. lemma:

$$\begin{aligned}
 \sum_{n=1}^{\infty} |\langle \mathbf{x} | \mathbf{x}_n \rangle|^2 &= \sum_{n=1}^{\infty} \left| \left\langle \sum_{m=1}^{\infty} a_m \mathbf{y}_m | \mathbf{x}_n \right\rangle \right|^2 && \text{by definition of } \mathbf{x} && \text{(item (2) page 170)} \\
 &= \sum_{n=1}^{\infty} \left| \sum_{m=1}^{\infty} a_m \langle \mathbf{y}_m | \mathbf{x}_n \rangle \right|^2 && \text{by homogeneous property of } \langle \triangle | \nabla \rangle && \text{(Definition F.9 page 132)} \\
 &= \sum_{n=1}^{\infty} \left| \sum_{m=1}^{\infty} a_n \bar{\delta}_{mn} \right|^2 && \text{by definition of biorthogonal} && \text{(Definition G.15 page 167)} \\
 &= \sum_{n=1}^{\infty} |a_n|^2 && \text{by definition of } \bar{\delta} && \text{(Definition G.12 page 160)}
 \end{aligned}$$

<sup>35</sup> Igarı (1996) page 220 (Lemma 9.8), Wojtaszczyk (1997) pages 20–21 (Lemma 2.7(a))

4. Then

$$\begin{aligned}
 A \sum_{n=1}^{\infty} |a_n|^2 &\leq \left\| \sum_{n=1}^{\infty} a_n \mathbf{x}_n \right\|^2 \leq B \sum_{n=1}^{\infty} |a_n|^2 && \text{by definition of } \{\mathbf{y}_n\} \text{ (item (1) page 170)} \\
 \Rightarrow A \sum_{n=1}^{\infty} |a_n|^2 &\leq \left\| \sum_{n=1}^{\infty} a_n \mathbf{y}_n \right\|^2 \leq B \sum_{n=1}^{\infty} |a_n|^2 && \text{by definition of } \{\mathbf{y}_n\} \text{ (item (1) page 170)} \\
 \Rightarrow A \sum_{n=1}^{\infty} |a_n|^2 &\leq \|\mathbf{x}\|^2 \leq B \sum_{n=1}^{\infty} |a_n|^2 && \text{by definition of } \mathbf{x} \text{ (item (2) page 170)} \\
 \Rightarrow A \sum_{n=1}^{\infty} |\langle \mathbf{x} | \mathbf{x}_n \rangle|^2 &\leq \|\mathbf{x}\|^2 \leq B \sum_{n=1}^{\infty} |\langle \mathbf{x} | \mathbf{x}_n \rangle|^2 && \text{by (3) lemma} \\
 \Rightarrow \frac{1}{B} \|\mathbf{x}\|^2 &\leq \sum_{n=1}^{\infty} |\langle \mathbf{x} | \mathbf{x}_n \rangle|^2 \leq \frac{1}{A} \|\mathbf{x}\|^2
 \end{aligned}$$

⇒

**Theorem G.15** (Battle-Lemarié orthogonalization). <sup>36</sup> Let  $\tilde{f}(\omega)$  be the FOURIER TRANSFORM (Definition 4.2 page 16) of a function  $f \in L^2_{\mathbb{R}}$ .

T H M	$  \left\{ \begin{array}{l} 1. \{ \mathbf{T}^n \mathbf{g}   n \in \mathbb{Z} \} \text{ is a RIESZ BASIS for } L^2_{\mathbb{R}} \text{ and} \\ 2. \tilde{f}(\omega) \triangleq \frac{\tilde{g}(\omega)}{\sqrt{2\pi \sum_{n \in \mathbb{Z}}  \tilde{g}(\omega + 2\pi n) ^2}} \end{array} \right\} \Rightarrow \left\{ \begin{array}{l} \{ \mathbf{T}^n \mathbf{f}   n \in \mathbb{Z} \} \\ \text{is an ORTHONORMAL BASIS for } L^2_{\mathbb{R}} \end{array} \right\}  $
-------	---

PROOF:

1. Proof that  $\{ \mathbf{T}^n \mathbf{f} | n \in \mathbb{Z} \}$  is orthonormal:

$$\tilde{S}_{\phi\phi}(\omega) = 2\pi \sum_{n \in \mathbb{Z}} |\tilde{f}(\omega + 2\pi n)|^2 \quad \text{by Theorem H.1 page 179}$$

$$= 2\pi \sum_{n \in \mathbb{Z}} \left| \frac{\tilde{g}(\omega + 2\pi n)}{\sqrt{2\pi \sum_{m \in \mathbb{Z}} |\tilde{g}(\omega + 2\pi(n-m))|^2}} \right|^2 \quad \text{by left hypothesis}$$

$$= \sum_{n \in \mathbb{Z}} \left| \frac{\tilde{g}(\omega + 2\pi n)}{\sqrt{\sum_{m \in \mathbb{Z}} |\tilde{g}(\omega + 2\pi m)|^2}} \right|^2$$

$$= \sum_{n \in \mathbb{Z}} \left| \frac{1}{\sqrt{\sum_{m \in \mathbb{Z}} |\tilde{g}(\omega + 2\pi m)|^2}} \right|^2 |\tilde{g}(\omega + 2\pi n)|^2$$

$$= \frac{1}{\sum_{m \in \mathbb{Z}} |\tilde{g}(\omega + 2\pi m)|^2} \sum_{n \in \mathbb{Z}} |\tilde{g}(\omega + 2\pi n)|^2$$

$$= 1$$

$$\Rightarrow \{ \mathbf{f}_n | n \in \mathbb{Z} \} \text{ is orthonormal}$$

by Theorem H.3 page 185

<sup>36</sup> Wojtaszczyk (1997) page 25 (Remark 2.4), Vidakovic (1999) page 71, Mallat (1989) page 72, Mallat (1999) page 225, Daubechies (1992) page 140 ((5.3.3))

2. Proof that  $\{\mathbf{T}^n \mathbf{f} \mid n \in \mathbb{Z}\}$  is a basis for  $V_0$ : by Lemma G.1 page 155.



## G.7 Frames in Hilbert spaces

**Definition G.16.** <sup>37</sup> Let  $\{\mathbf{x}_n \in X \mid n \in \mathbb{N}\}$  be a set of vectors in a HILBERT SPACE  $H \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle \mid \nabla \rangle)$ .

The set  $\{\mathbf{x}_n\}$  is a **frame** for  $H$  if (STABILITY CONDITION)

$$\exists A, B \in \mathbb{R}^+ \quad \text{such that} \quad A \|\mathbf{x}\|^2 \leq \sum_{n=1}^{\infty} |\langle \mathbf{x} \mid \mathbf{x}_n \rangle|^2 \leq B \|\mathbf{x}\|^2 \quad \forall \mathbf{x} \in X.$$

The quantities  $A$  and  $B$  are **frame bounds**.

The quantity  $A'$  is the **optimal lower frame bound** if

$$A' = \sup \{A \in \mathbb{R}^+ \mid A \text{ is a lower frame bound}\}.$$

The quantity  $B'$  is the **optimal upper frame bound** if

$$B' = \inf \{B \in \mathbb{R}^+ \mid B \text{ is an upper frame bound}\}.$$

A frame is a **tight frame** if  $A = B$ .

A frame is a **normalized tight frame** (or a **Parseval frame**) if  $A = B = 1$ .

A frame  $\{\mathbf{x}_n \mid n \in \mathbb{N}\}$  is an **exact frame** if for some  $m \in \mathbb{Z}$ ,  $\{\mathbf{x}_n \mid n \in \mathbb{N}\} \setminus \{\mathbf{x}_m\}$  is NOT a frame.

A frame is a *Parseval frame* (Definition G.16) if it satisfies *Parseval's Identity* (Theorem G.9 page 162). All orthonormal bases are Parseval frames (Theorem G.9 page 162); but not all Parseval frames are orthonormal bases.

**Definition G.17.** Let  $\{\mathbf{x}_n\}$  be a **frame** (Definition G.16 page 172) for the HILBERT SPACE  $H \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle \mid \nabla \rangle)$ . Let  $\mathbf{S}$  be an OPERATOR on  $H$ .

$\mathbf{S}$  is a **frame operator** for  $\{\mathbf{x}_n\}$  if  $\mathbf{S}\mathbf{f}(x) = \sum_{n \in \mathbb{Z}} \langle \mathbf{f} \mid \mathbf{x}_n \rangle \mathbf{x}_n \quad \forall \mathbf{f} \in H.$

**Theorem G.16.** <sup>38</sup> Let  $\mathbf{S}$  be a FRAME OPERATOR (Definition G.17 page 172) of a FRAME  $\{\mathbf{x}_n\}$  (Definition G.16 page 172) for the HILBERT SPACE  $H \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle \mid \nabla \rangle)$ .

- (1).  $\mathbf{S}$  is INVERTIBLE. and  
 (2).  $\mathbf{f}(x) = \sum_{n \in \mathbb{Z}} \langle \mathbf{f} \mid \mathbf{S}^{-1} \mathbf{x}_n \rangle \mathbf{x}_n = \sum_{n \in \mathbb{Z}} \langle \mathbf{f} \mid \mathbf{x}_n \rangle \mathbf{S}^{-1} \mathbf{x}_n \quad \forall \mathbf{f} \in H$

**Theorem G.17.** <sup>39</sup> Let  $\{\mathbf{x}_n \in X \mid n=1,2,\dots,N\}$  be a set of vectors in a HILBERT SPACE  $H \triangleq (X, +, \cdot, (\mathbb{F}, \dot{+}, \dot{\times}), \langle \triangle \mid \nabla \rangle)$ .

$\{\mathbf{x}_n\}$  is a FRAME for  $\text{span}\{\mathbf{x}_n\}$ .

PROOF:

<sup>37</sup> Young (2001) pages 154–155, Christensen (2003) page 88 (Definitions 5.1.1, 5.1.2), Heil (2011) pages 204–205 (Definition 8.2), Jørgensen et al. (2008) page 267 (Definition 12.22), Duffin and Schaeffer (1952) page 343, Daubechies et al. (1986) page 1272

<sup>38</sup> Christensen (2008) pages 100–102 (Theorem 5.1.7)

<sup>39</sup> Christensen (2003) page 3

1. Upper bound: Proof that there exists  $B$  such that  $\sum_{n=1}^N |\langle \mathbf{x} | \mathbf{x}_n \rangle|^2 \leq B \|\mathbf{x}\|^2 \quad \forall \mathbf{x} \in H$ :

$$\begin{aligned} \sum_{n=1}^N |\langle \mathbf{x} | \mathbf{x}_n \rangle|^2 &\leq \sum_{n=1}^N \langle \mathbf{x}_n | \mathbf{x}_n \rangle \langle \mathbf{x} | \mathbf{x} \rangle && \text{by Cauchy-Schwarz inequality} \\ &= \underbrace{\left\{ \sum_{n=1}^N \|\mathbf{x}_n\|^2 \right\}}_B \|\mathbf{x}\|^2 \end{aligned}$$

2. Lower bound: Proof that there exists  $A$  such that  $A \|\mathbf{x}\|^2 \leq \sum_{n=1}^N |\langle \mathbf{x} | \mathbf{x}_n \rangle|^2 \quad \forall \mathbf{x} \in H$ :

$$\begin{aligned} \sum_{n=1}^N |\langle \mathbf{x} | \mathbf{x}_n \rangle|^2 &= \sum_{n=1}^N \left| \left\langle \mathbf{x}_n \mid \frac{\mathbf{x}}{\|\mathbf{x}\|} \right\rangle \right|^2 \|\mathbf{x}\|^2 \\ &\geq \underbrace{\left( \inf_y \left\{ \sum_{n=1}^N |\langle \mathbf{x}_n | \mathbf{y} \rangle|^2 \mid \|\mathbf{y}\| = 1 \right\} \right)}_A \|\mathbf{x}\|^2 \end{aligned}$$


*Example G.1.* Let  $(\mathbb{R}^2, +, \cdot, (\mathbb{R}, \dot{+}, \dot{\times}), \langle \triangle | \nabla \rangle)$  be an inner product space with  $\left\langle \begin{bmatrix} x_1 \\ y_1 \end{bmatrix} \mid \begin{bmatrix} x_2 \\ y_2 \end{bmatrix} \right\rangle \triangleq x_1 x_2 + y_1 y_2$ . Let  $\mathbf{S}$  be the *frame operator* (Definition G.17 page 172) with *inverse*  $\mathbf{S}^{-1}$ .

Let  $N \in \{3, 4, 5, \dots\}$ ,  $\theta \in \mathbb{R}$ , and  $r \in \mathbb{R}^+$  ( $r > 0$ ).

Let  $\mathbf{x}_n \triangleq r \begin{bmatrix} \cos(\theta + 2n\pi/N) \\ \sin(\theta + 2n\pi/N) \end{bmatrix} \quad \forall n \in \{0, 1, \dots, N-1\}$ .

Then,  $(\mathbf{x}_0, \mathbf{x}_1, \dots, \mathbf{x}_{N-1})$  is a **tight frame** for  $\mathbb{R}^2$  with *frame bound*  $A = \frac{Nr^2}{2}$ .

Moreover,  $\mathbf{S} = \frac{Nr^2}{2} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$ ,  $\mathbf{S}^{-1} = \frac{2}{Nr^2} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$ , and  $\mathbf{v} = \frac{2}{Nr^2} \sum_{n=0}^{N-1} \langle \mathbf{v} | \mathbf{x}_n \rangle \mathbf{x}_n \quad \forall \mathbf{v} \in \mathbb{R}^2$ .



**PROOF:**

1. Proof that  $(\mathbf{x}_0, \mathbf{x}_1, \dots, \mathbf{x}_{N-1})$  is a *tight frame* with *frame bound*  $A = \frac{Nr^2}{2}$ : Let  $\mathbf{v} \triangleq (x, y) \in \mathbb{R}^2$ .

$$\begin{aligned} \sum_{n=0}^{N-1} |\langle \mathbf{v} | \mathbf{x}_n \rangle|^2 &\triangleq \sum_{n=0}^{N-1} \left| \mathbf{v}^H r \begin{bmatrix} \cos\left(\theta + \frac{2n\pi}{N}\right) \\ \sin\left(\theta + \frac{2n\pi}{N}\right) \end{bmatrix} \right|^2 && \text{by definitions of } \mathbf{v} \text{ of } \langle \mathbf{y} | \mathbf{x} \rangle \\ &\triangleq \sum_{n=0}^{N-1} r^2 \left| x \cos\left(\theta + \frac{2n\pi}{N}\right) + y \sin\left(\theta + \frac{2n\pi}{N}\right) \right|^2 && \text{by definition of } \mathbf{y}^H \mathbf{x} \text{ operation} \\ &= r^2 x^2 \sum_{n=0}^{N-1} \cos^2\left(\theta + \frac{2n\pi}{N}\right) + r^2 y^2 \sum_{n=0}^{N-1} \sin^2\left(\theta + \frac{2n\pi}{N}\right) + r^2 xy \sum_{n=0}^{N-1} \cos\left(\theta + \frac{2n\pi}{N}\right) \sin\left(\theta + \frac{2n\pi}{N}\right) \\ &= r^2 x^2 \frac{N}{2} + r^2 y^2 \frac{N}{2} + r^2 xy 0 && \text{by Corollary D.1 page 107} \\ &= (x^2 + y^2) \frac{Nr^2}{2} = \underbrace{\left( \frac{Nr^2}{2} \right)}_A \mathbf{v}^H \mathbf{v} \triangleq \underbrace{\left( \frac{Nr^2}{2} \right)}_A \|\mathbf{v}\|^2 && \text{by definition of } \|\mathbf{v}\| \end{aligned}$$

2. Proof that  $\mathbf{S} = \frac{Nr^2}{2} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$ :

(a) Let  $\mathbf{e}_1 \triangleq \begin{bmatrix} 1 \\ 0 \end{bmatrix}$  and  $\mathbf{e}_2 \triangleq \begin{bmatrix} 0 \\ 1 \end{bmatrix}$ .

(b) lemma:  $\mathbf{S}\mathbf{e}_1 = \frac{Nr^2}{2}\mathbf{e}_1$ . Proof:

$$\begin{aligned} \mathbf{S}\mathbf{e}_1 &= \sum_{n=0}^{N-1} \langle \mathbf{e}_1 | \mathbf{x}_n \rangle \mathbf{x}_n \\ &= \sum_{n=0}^{N-1} r \cos\left(\theta + \frac{2n\pi}{N}\right) r \begin{bmatrix} \cos\left(\theta + \frac{2n\pi}{N}\right) \\ \sin\left(\theta + \frac{2n\pi}{N}\right) \end{bmatrix} = r^2 \sum_{n=0}^{N-1} \begin{bmatrix} \cos^2\left(\theta + \frac{2n\pi}{N}\right) \\ \cos\left(\theta + \frac{2n\pi}{N}\right) \sin\left(\theta + \frac{2n\pi}{N}\right) \end{bmatrix} \\ &= r^2 \begin{bmatrix} N/2 \\ 0 \end{bmatrix} = \frac{Nr^2}{2} \mathbf{e}_1 \quad \text{by Summation around unit circle (Corollary D.1 page 107)} \end{aligned}$$

(c) lemma:  $\mathbf{S}\mathbf{e}_2 = \frac{Nr^2}{2}\mathbf{e}_2$ . Proof:

$$\begin{aligned} \mathbf{S}\mathbf{e}_2 &= \sum_{n=0}^{N-1} \langle \mathbf{e}_2 | \mathbf{x}_n \rangle \mathbf{x}_n \\ &= \sum_{n=0}^{N-1} r \sin\left(\theta + \frac{2n\pi}{N}\right) r \begin{bmatrix} \cos\left(\theta + \frac{2n\pi}{N}\right) \\ \sin\left(\theta + \frac{2n\pi}{N}\right) \end{bmatrix} = r^2 \sum_{n=0}^{N-1} \begin{bmatrix} \sin\left(\theta + \frac{2n\pi}{N}\right) \cos\left(\theta + \frac{2n\pi}{N}\right) \\ \sin^2\left(\theta + \frac{2n\pi}{N}\right) \end{bmatrix} \\ &= r^2 \begin{bmatrix} 0 \\ N/2 \end{bmatrix} = \frac{Nr^2}{2} \mathbf{e}_2 \quad \text{by Summation around unit circle (Corollary D.1 page 107)} \end{aligned}$$

(d) Complete the proof of item (2) using *Eigendecomposition*  $\mathbf{S} = \mathbf{Q}\mathbf{\Lambda}\mathbf{Q}^{-1}$ :

$$\mathbf{S}\mathbf{e}_1 = \frac{Nr^2}{2}\mathbf{e}_1 \quad \text{by (2c) lemma}$$

$$\Rightarrow \mathbf{e}_1 \text{ is an eigenvector of } \mathbf{S} \text{ with eigenvalue } \frac{Nr^2}{2}$$

$$\mathbf{S}\mathbf{e}_2 = \frac{Nr^2}{2}\mathbf{e}_2 \quad \text{by (2c) lemma}$$

$$\Rightarrow \mathbf{e}_2 \text{ is an eigenvector of } \mathbf{S} \text{ with eigenvalue } \frac{Nr^2}{2}$$

$$\begin{aligned} &\text{Eigendecomposition of } \mathbf{S} \\ \mathbf{S} &= \underbrace{\begin{bmatrix} | & | \\ \mathbf{e}_1 & \mathbf{e}_2 \\ | & | \end{bmatrix}}_{\mathbf{Q}} \underbrace{\begin{bmatrix} \frac{Nr^2}{2} & 0 \\ 0 & \frac{Nr^2}{2} \end{bmatrix}}_{\mathbf{\Lambda}} \underbrace{\begin{bmatrix} | & | \\ \mathbf{e}_1 & \mathbf{e}_2 \\ | & | \end{bmatrix}^{-1}}_{\mathbf{Q}^{-1}} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} \frac{Nr^2}{2} & 0 \\ 0 & \frac{Nr^2}{2} \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}^{-1} = \frac{Nr^2}{2} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \end{aligned}$$

3. Proof that  $\mathbf{S}^{-1} = \frac{2}{Nr^2} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$ :

$$\mathbf{S}\mathbf{S}^{-1} = \frac{Nr^2}{2} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \frac{2}{Nr^2} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \triangleq \mathbf{I}_2 \quad \text{by item (2)}$$

$$\mathbf{S}^{-1}\mathbf{S} = \frac{2}{Nr^2} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \frac{Nr^2}{2} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \triangleq \mathbf{I}_2 \quad \text{by item (2)}$$

4. Proof that  $\mathbf{v} = \frac{2}{Nr^2} \sum_{n=0}^{N-1} (\mathbf{v}^H \mathbf{x}_n) \mathbf{x}_n$ :

$$\mathbf{v} = \sum_{n=0}^{N-1} \langle \mathbf{v} | \mathbf{S}^{-1} \mathbf{x}_n \rangle \mathbf{x}_n = \sum_{n=0}^{N-1} \left\langle \mathbf{v} \middle| \frac{2}{Nr^2} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \mathbf{x}_n \right\rangle \mathbf{x}_n \quad \text{by item (3)}$$

$$= \frac{2}{Nr^2} \sum_{n=0}^{N-1} \langle \mathbf{v} | \mathbf{x}_n \rangle \mathbf{x}_n = \frac{2}{Nr^2} \sum_{n=0}^{N-1} (\mathbf{v}^H \mathbf{x}_n) \mathbf{x}_n \quad \text{by definition of } \langle \mathbf{y} | \mathbf{x} \rangle$$





**Example G.2** (Peace Frame/Mercedes Frame).<sup>40</sup> Let  $(\mathbb{R}^2, +, \cdot, (\mathbb{R}, \dot{+}, \dot{\times}), \langle \Delta | \nabla \rangle)$  be an inner product space with  $\langle (x_1, y_1) | (x_2, y_2) \rangle \triangleq x_1 y_1 + x_2 y_2$ . Let **S** be the *frame operator* (Definition G.17 page 172) with *inverse*  $\mathbf{S}^{-1}$ .


E  
X

$$\text{Let } \mathbf{x}_1 \triangleq \frac{1}{2} \begin{bmatrix} 0 \\ 2 \end{bmatrix}, \mathbf{x}_2 \triangleq \frac{1}{2} \begin{bmatrix} -\sqrt{3} \\ -1 \end{bmatrix}, \text{ and } \mathbf{x}_3 \triangleq \frac{1}{2} \begin{bmatrix} \sqrt{3} \\ -1 \end{bmatrix}.$$

Then,  $(\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3)$  is a **tight frame** for  $\mathbb{R}^2$  with *frame bound*  $A = \frac{3}{2}$ .

$$\text{Moreover, } \mathbf{S} = \frac{3}{2} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \quad \mathbf{S}^{-1} = \frac{2}{3} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix},$$

$$\text{and } \mathbf{v} = \frac{2}{3} \sum_{n=1}^3 \langle \mathbf{v} | \mathbf{x}_n \rangle \mathbf{x}_n \triangleq \frac{2}{3} \sum_{n=1}^3 (\mathbf{v}^H \mathbf{x}_n) \mathbf{x}_n \quad \forall \mathbf{v} \in \mathbb{R}^2.$$



**PROOF:**

1. This frame is simply a special case of the frame presented in Example G.1 (page 173) with  $r = 1$ ,  $N = 3$ , and  $\theta = \pi/2$ .
2. Let's give it a try! Let  $\mathbf{v} \triangleq \begin{bmatrix} 1 \\ 1 \end{bmatrix}$ .

$$\begin{aligned} \sum_{n=1}^3 \langle \mathbf{v} | \mathbf{S}^{-1} \mathbf{x}_n \rangle \mathbf{x}_n &= \frac{2}{3} \sum_{n=1}^3 (\mathbf{v}^H \mathbf{x}_n) \mathbf{x}_n && \text{by Example G.1 page 173} \\ &= (\mathbf{v}^H \mathbf{x}_1) \mathbf{x}_1 + (\mathbf{v}^H \mathbf{x}_2) \mathbf{x}_2 + (\mathbf{v}^H \mathbf{x}_3) \mathbf{x}_3 \\ &= \frac{2}{3} \left( \left( \mathbf{v}^H \frac{1}{2} \begin{bmatrix} 0 \\ 2 \end{bmatrix} \right) \mathbf{x}_1 + \left( \mathbf{v}^H \frac{1}{2} \begin{bmatrix} -\sqrt{3} \\ -1 \end{bmatrix} \right) \mathbf{x}_2 + \left( \mathbf{v}^H \frac{1}{2} \begin{bmatrix} \sqrt{3} \\ -1 \end{bmatrix} \right) \mathbf{x}_3 \right) \\ &= \frac{2}{3} \cdot \frac{1}{2} \left( \left( \mathbf{v}^H \begin{bmatrix} 0 \\ 2 \end{bmatrix} \right) \mathbf{x}_1 + \left( \mathbf{v}^H \begin{bmatrix} -\sqrt{3} \\ -1 \end{bmatrix} \right) \mathbf{x}_2 + \left( \mathbf{v}^H \begin{bmatrix} \sqrt{3} \\ -1 \end{bmatrix} \right) \mathbf{x}_3 \right) \\ &= \frac{1}{3} \left( (2) \frac{1}{2} \begin{bmatrix} 0 \\ 2 \end{bmatrix} + (-\sqrt{3} - 1) \frac{1}{2} \begin{bmatrix} -\sqrt{3} \\ -1 \end{bmatrix} + (\sqrt{3} - 1) \frac{1}{2} \begin{bmatrix} \sqrt{3} \\ -1 \end{bmatrix} \right) \\ &= \frac{1}{6} \begin{bmatrix} 2(0) & + & (-\sqrt{3} - 1)(-\sqrt{3}) & + & (\sqrt{3} - 1)(\sqrt{3}) \\ 2(2) & + & (-\sqrt{3} - 1)(-1) & + & (\sqrt{3} - 1)(-1) \end{bmatrix} \\ &= \frac{1}{6} \begin{bmatrix} 0 & + & (3 + \sqrt{3}) & + & (3 - \sqrt{3}) \\ 4 & + & (1 + \sqrt{3}) & + & (1 - \sqrt{3}) \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \end{bmatrix} \triangleq \mathbf{v} \end{aligned}$$



In Example G.1 (page 173) and Example G.2 (page 175), the frame operator **S** and its inverse  $\mathbf{S}^{-1}$  were computed. In general however, it is not always necessary or even possible to compute these, as illustrated in Example G.3 (next).

**Example G.3.**<sup>41</sup> Let  $(\mathbb{R}^2, +, \cdot, (\mathbb{R}, \dot{+}, \dot{\times}), \langle \Delta | \nabla \rangle)$  be an inner product space with  $\langle (x_1, y_1) | (x_2, y_2) \rangle \triangleq x_1 y_1 + x_2 y_2$ . Let **S** be the *frame operator* (Definition G.17 page 172) with *inverse*  $\mathbf{S}^{-1}$ .


<sup>40</sup> Heil (2011) pages 204–205 ( $r = 1$  case), Byrne (2005) page 80 ( $r = 1$  case), Han et al. (2007) page 91 (Example 3.9,  $r = \sqrt{2/3}$  case)

<sup>41</sup> Christensen (2003) pages 7–8 (?)

E  
X

Let  $p$  and  $q$  be *orthonormal* vectors in  $X \triangleq \text{span}\{p, q\}$ .

Let  $x_1 \triangleq p$ ,  $x_2 \triangleq p + q$ , and  $x_3 \triangleq p - q$ . Then,  $\{x_1, x_2, x_3\}$  is a **frame** for  $X$  with *frame bounds*  $A = 0$  and  $B = 5$ .



Moreover,

$$\begin{aligned} S^{-1}x_1 &= \frac{1}{3}p & \text{and} \\ S^{-1}x_2 &= \frac{1}{3}p + \frac{1}{2}q & \text{and} \\ S^{-1}x_3 &= \frac{1}{3}p - \frac{1}{2}q. \end{aligned}$$

PROOF:

1. Proof that  $(x_1, x_2, x_3)$  is a *frame* with *frame bounds*  $A = 0$  and  $B = 5$ :

$$\begin{aligned} \sum_{n=1}^3 |\langle v | x_n \rangle|^2 &\triangleq |\langle v | p \rangle|^2 + |\langle v | p + q \rangle|^2 + |\langle v | p - q \rangle|^2 && \text{by definitions of } x_1, x_2, \text{ and } x_3 \\ &= |\langle v | p \rangle|^2 + |\langle v | p \rangle + \langle v | q \rangle|^2 + |\langle v | p \rangle - \langle v | q \rangle|^2 && \text{by additivity of } \langle \triangle | \nabla \rangle \text{ (Definition F.9 page 132)} \\ &= |\langle v | p \rangle|^2 + (|\langle v | p \rangle|^2 + |\langle v | q \rangle|^2 + \langle v | p \rangle \langle v | q \rangle^* + \langle v | q \rangle \langle v | p \rangle^*) \\ &\quad + (|\langle v | p \rangle|^2 + |\langle v | q \rangle|^2 - \langle v | p \rangle \langle v | q \rangle^* - \langle v | q \rangle \langle v | p \rangle^*) \\ &= 3|\langle v | p \rangle|^2 + 2|\langle v | q \rangle|^2 \\ &\leq 3\|v\| \|p\| + 2\|v\| \|q\| && \text{by CS Inequality} \\ &= \|v\| (3\|p\| + 2\|q\|) \\ &= 5\|v\| && \text{by orthonormality of } p \text{ and } q \end{aligned}$$

2. lemma:  $Sp = 3p$ ,  $Sq = 2q$ ,  $S^{-1}p = \frac{1}{3}p$ , and  $S^{-1}q = \frac{1}{2}q$ . Proof:

$$\begin{aligned} Sp &\triangleq \sum_{n=1}^3 \langle p | x_n \rangle x_n \\ &= \langle p | p \rangle p + \langle p | p + q \rangle (p + q) + \langle p | p - q \rangle (p - q) \\ &= (1)p + (1 + 0)(p + q) + (1 - 0)(p - q) \\ &= 3p \\ \Rightarrow S^{-1}p &= \frac{1}{3}p \\ Sq &\triangleq \sum_{n=1}^3 \langle q | x_n \rangle x_n \\ &= \langle q | p \rangle p + \langle q | p + q \rangle (p + q) + \langle q | p - q \rangle (p - q) \\ &= (0)q + (0 + 1)(p + q) + (0 - 1)(p - q) \\ &= 2q \\ \Rightarrow S^{-1}q &= \frac{1}{2}q \end{aligned}$$

3. Remark: Without knowing  $p$  and  $q$ , from (2) lemma it follows that it is not possible to compute  $S$  or  $S^{-1}$  explicitly.
4. Proof that  $S^{-1}x_1 = \frac{1}{3}p$ ,  $S^{-1}x_2 = \frac{1}{3}p + \frac{1}{2}q$  and  $S^{-1}x_3 = \frac{1}{3}p - \frac{1}{2}q$ :

$$\begin{aligned} S^{-1}x_1 &\triangleq S^{-1}p && \text{by definition of } x_1 \\ &= \frac{1}{3}p && \text{by (2) lemma} \\ S^{-1}x_2 &\triangleq S^{-1}(p + q) && \text{by definition of } x_2 \\ &= \frac{1}{3}p + \frac{1}{2}q && \text{by (2) lemma} \end{aligned}$$

$$\begin{aligned} \mathbf{S}^{-1} \mathbf{x}_3 &\triangleq \mathbf{S}^{-1}(\mathbf{p} - \mathbf{q}) \\ &= \frac{1}{3}\mathbf{p} - \frac{1}{2}\mathbf{q} \end{aligned}$$

by definition of  $\mathbf{x}_2$ 

by (2) lemma

5. Check that  $\mathbf{v} = \sum_n \langle \mathbf{v} | \mathbf{x}_n \rangle \mathbf{x}_n = \langle \mathbf{v} | \mathbf{p} \rangle \mathbf{p} + \langle \mathbf{v} | \mathbf{q} \rangle \mathbf{q}$ :

$$\begin{aligned} \mathbf{v} &= \sum_{n=1}^3 \langle \mathbf{v} | \mathbf{S}^{-1} \mathbf{x}_n \rangle \mathbf{x}_n \\ &= \langle \mathbf{v} | \mathbf{S}^{-1} \mathbf{p} \rangle \mathbf{p} + \langle \mathbf{v} | \mathbf{S}^{-1}(\mathbf{p} + \mathbf{q}) \rangle (\mathbf{p} + \mathbf{q}) + \langle \mathbf{v} | \mathbf{S}^{-1}(\mathbf{p} - \mathbf{q}) \rangle (\mathbf{p} - \mathbf{q}) \\ &= \left\langle \mathbf{v} \left| \frac{1}{3}\mathbf{p} \right. \right\rangle \mathbf{p} + \left\langle \mathbf{v} \left| \frac{1}{3}\mathbf{p} + \frac{1}{2}\mathbf{q} \right. \right\rangle (\mathbf{p} + \mathbf{q}) + \left\langle \mathbf{v} \left| \frac{1}{3}\mathbf{p} - \frac{1}{2}\mathbf{q} \right. \right\rangle (\mathbf{p} - \mathbf{q}) \\ &= \left( \frac{1}{3} + \frac{1}{3} + \frac{1}{3} \right) \langle \mathbf{v} | \mathbf{p} \rangle \mathbf{p} + \left( \frac{1}{3} - \frac{1}{3} \right) \langle \mathbf{v} | \mathbf{p} \rangle \mathbf{q} + \left( \frac{1}{2} - \frac{1}{2} \right) \langle \mathbf{v} | \mathbf{q} \rangle \mathbf{p} + \left( \frac{1}{2} + \frac{1}{2} \right) \langle \mathbf{v} | \mathbf{q} \rangle \mathbf{q} \\ &= \langle \mathbf{v} | \mathbf{p} \rangle \mathbf{p} + \langle \mathbf{v} | \mathbf{q} \rangle \mathbf{q} \end{aligned}$$





# APPENDIX H

## POWER SPECTRUM FUNCTIONS

### H.1 Correlation

Definition H.1 and Definition H.2 define four quantities. In this document, the quantities' notation and terminology are similar to those used in the study of *random processes*.

**Definition H.1.** <sup>1</sup> Let  $\langle \triangle | \nabla \rangle$  be the STANDARD INNER PRODUCT in  $L^2_{\mathbb{R}}$  (Definition B.1 page 69).

**DEF**  $R_{fg}(n) \triangleq \langle f(x) | T^n g(x) \rangle, \quad n \in \mathbb{Z}; \quad f, g \in L^2_{\mathbb{F}},$  is the **cross-correlation function** of  $f$  and  $g$ .  
 $R_{ff}(n) \triangleq \langle f(x) | T^n f(x) \rangle, \quad n \in \mathbb{Z}; \quad f \in L^2_{\mathbb{F}},$  is the **autocorrelation function** of  $f$ .

**Definition H.2.** <sup>2</sup> Let  $R_{fg}(n)$  and  $R_{ff}(n)$  be the sequences defined in Definition H.1 page 179. Let  $\mathbf{Z}(x_n)$  be the Z-TRANSFORM (Definition 5.4 page 28) of a sequence  $(x_n)_{n \in \mathbb{Z}}$ .

**DEF**  $\check{S}_{fg}(z) \triangleq \mathbf{Z}[R_{fg}(n)], \quad f, g \in L^2_{\mathbb{F}},$  is the **complex cross-power spectrum** of  $f$  and  $g$ .  
 $\check{S}_{ff}(z) \triangleq \mathbf{Z}[R_{ff}(n)], \quad f \in L^2_{\mathbb{F}},$  is the **complex auto-power spectrum** of  $f$ .

### H.2 Power Spectrum

**Definition H.3.** <sup>3</sup> Let  $\check{S}_{fg}(z)$  and  $\check{S}_{ff}(z)$  be the functions defined in Definition H.2 page 179.

**DEF**  $\tilde{S}_{fg}(\omega) \triangleq \check{S}_{fg}(e^{i\omega}), \quad \forall f, g \in L^2_{\mathbb{F}},$  is the **cross-power spectrum** of  $f$  and  $g$ .  
 $\tilde{S}_{ff}(\omega) \triangleq \check{S}_{ff}(e^{i\omega}), \quad \forall f \in L^2_{\mathbb{F}},$  is the **auto-power spectrum** of  $f$ .

**Theorem H.1.** <sup>4</sup> Let  $\tilde{S}_{fg}(\omega)$  and  $\tilde{S}_{ff}(\omega)$  be defined as in Definition H.3 (page 179).

Let  $\tilde{f}(\omega)$  be the FOURIER TRANSFORM (Definition 4.2 page 16) of a function  $f(x) \in L^2_{\mathbb{F}}$ .

**THM** 
$$\begin{aligned} \tilde{S}_{fg}(\omega) &= 2\pi \sum_{n \in \mathbb{Z}} \tilde{f}(\omega + 2\pi n) \tilde{g}^*(\omega + 2\pi n) \quad \forall f, g \in L^2_{\mathbb{F}} \\ \tilde{S}_{ff}(\omega) &= 2\pi \sum_{n \in \mathbb{Z}} |\tilde{f}(\omega + 2\pi n)|^2 \quad \forall f \in L^2_{\mathbb{F}} \end{aligned}$$

<sup>1</sup> Chui (1992) page 134, Papoulis (1991) pages 294–332  $\langle (10-29), (10-169) \rangle$

<sup>2</sup> Chui (1992) page 134, Papoulis (1991) page 334  $\langle (10-178) \rangle$

<sup>3</sup> Chui (1992) page 134, Papoulis (1991) page 333  $\langle (10-179) \rangle$

<sup>4</sup> Chui (1992) page 135

✎ PROOF: Let  $z \triangleq e^{i\omega}$ .

$$\begin{aligned}
 \tilde{S}_{fg}(\omega) &\triangleq \tilde{S}_{fg}(z) && \text{by definition of } \tilde{S}_{fg} && (\text{Definition H.3 page 179}) \\
 &= \sum_{n \in \mathbb{Z}} R_{fg}(n) z^{-n} && \text{by definition of } \tilde{S}_{fg} && (\text{Definition H.2 page 179}) \\
 &= \sum_{n \in \mathbb{Z}} \langle f(x) | g(x-n) \rangle z^{-n} && \text{by definition of } \tilde{S}_{fg} && (\text{Definition H.3 page 179}) \\
 &= \sum_{n \in \mathbb{Z}} \langle \tilde{F}[f(x)] | \tilde{F}[g(x-n)] \rangle z^{-n} && \text{by unitary property of } \tilde{F} && (\text{Theorem 4.3 page 17}) \\
 &= \sum_{n \in \mathbb{Z}} \langle \tilde{f}(v) | e^{-ivn} \tilde{g}(v) \rangle z^{-n} && \text{by shift relation} && (\text{Theorem 4.4 page 18}) \\
 &= \sum_{n \in \mathbb{Z}} \sqrt{2\pi} \left[ \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} \tilde{f}(v) \tilde{g}^*(v) e^{ivn} dv \right] z^{-n} && \text{by definition of } L_{\mathbb{R}}^2 && (\text{Definition B.1 page 69}) \\
 &= \sqrt{2\pi} \sum_{n \in \mathbb{Z}} \left[ \tilde{F}^{-1} \left( \sqrt{2\pi} \tilde{f}(v) \tilde{g}^*(v) \right) \right]_{u=n} e^{-i\omega n} && \text{by Theorem 4.1 page 17} && \\
 &= 2\pi \sum_{n \in \mathbb{Z}} \tilde{f}(\omega + 2\pi n) \tilde{g}^*(\omega + 2\pi n) && \text{by IPSF with } \tau = 1 && (\text{Theorem I.3 page 197})
 \end{aligned}$$

$$\begin{aligned}
 \tilde{S}_{ff}(\omega) &= \tilde{S}_{fg}(\omega) \Big|_{g=f} && \text{by definition of } \tilde{S}_{fg}(\omega) && (\text{Definition H.3 page 179}) \\
 &= 2\pi \sum_{n \in \mathbb{Z}} \tilde{f}(\omega + 2\pi n) \tilde{g}^*(\omega + 2\pi n) \Big|_{g \triangleq f} && \text{by previous result} && \\
 &= 2\pi \sum_{n \in \mathbb{Z}} \tilde{f}(\omega + 2\pi n) \tilde{f}^*(\omega + 2\pi n) && && \\
 &= 2\pi \sum_{n \in \mathbb{Z}} |\tilde{f}(\omega + 2\pi n)|^2 && \text{because } |z|^2 \triangleq z z^* \quad \forall z \in \mathbb{C} &&
 \end{aligned}$$

⇒

**Proposition H.1.** Let  $\tilde{S}_{ff}(\omega)$  be defined as in Definition H.3 (page 179).

P R P	$\tilde{S}_{ff}(\omega) \geq 0$ (NON-NEGATIVE)
-------------	--

✎ PROOF:

$$\begin{aligned}
 \tilde{S}_{ff}(\omega) &= 2\pi \sum_{n \in \mathbb{Z}} |\tilde{f}(\omega + 2\pi n)|^2 && \text{by Theorem H.1 page 179} \\
 &\geq 0 && \text{because } |z| \geq 0 \quad \forall z \in \mathbb{C}
 \end{aligned}$$

⇒

**Proposition H.2.** Let  $\tilde{S}_{fg}(\omega)$  and  $\tilde{S}_{ff}(\omega)$  be defined as in Definition H.3 (page 179).

P R P	$\tilde{S}_{fg}(\omega + 2\pi) = \tilde{S}_{fg}(\omega)$ (PERIODIC with period $2\pi$ ) $\tilde{S}_{ff}(\omega + 2\pi) = \tilde{S}_{ff}(\omega)$ (PERIODIC with period $2\pi$ )
-------------	--

✎ PROOF:

$$\begin{aligned}
 \tilde{S}_{fg}(\omega + 2\pi) &= 2\pi \sum_{n \in \mathbb{Z}} \tilde{f}(\omega + 2\pi + 2\pi n) \tilde{g}^*(\omega + 2\pi + 2\pi n) && \text{by Theorem H.1 page 179} \\
 &= 2\pi \sum_{n \in \mathbb{Z}} \tilde{f}[\omega + 2\pi(n+1)] \tilde{g}^*[\omega + 2\pi(n+1)] \\
 &= 2\pi \sum_{m \in \mathbb{Z}} \tilde{f}[\omega + 2\pi m] \tilde{g}^*[\omega + 2\pi m] && \text{where } m \triangleq n+1 \\
 &= \tilde{S}_{fg}(\omega) && \text{by Theorem H.1 page 179} \\
 \tilde{S}_{ff}(\omega + 2\pi) &= \tilde{S}_{fg}(\omega + 2\pi) \Big|_{g=f} \\
 &= \tilde{S}_{fg}(\omega) \Big|_{g=f} && \text{by previous result} \\
 &= \tilde{S}_{ff}(\omega)
 \end{aligned}$$

⇒

**Proposition H.3.** Let  $\tilde{S}_{fg}(\omega)$  and  $\tilde{S}_{ff}(\omega)$  be defined as in Definition H.3 (page 179).

P R P	$f, g \text{ are real} \implies \tilde{S}_{fg}(-\omega) = \tilde{S}_{gf}(\omega)$	
	$f \text{ is real} \implies \tilde{S}_{ff}(-\omega) = \tilde{S}_{ff}(\omega)$	(SYMMETRIC about 0)
	$f, g \text{ are real} \implies \tilde{S}_{fg}(\pi - \omega) = \tilde{S}_{gf}(\pi + \omega)$	
	$f \text{ is real} \implies \tilde{S}_{ff}(\pi - \omega) = \tilde{S}_{ff}(\pi + \omega)$	(SYMMETRIC about $\pi$ )

✎ PROOF:

$$\begin{aligned}
 \tilde{S}_{fg}(-\omega) &= 2\pi \sum_{n \in \mathbb{Z}} \tilde{f}(-\omega + 2\pi n) \tilde{g}^*(-\omega + 2\pi n) && \text{by Theorem H.1 page 179} \\
 &= 2\pi \sum_{n \in \mathbb{Z}} \tilde{f}^*(\omega - 2\pi n) \tilde{g}(\omega - 2\pi n) && \text{by hypothesis and Theorem 4.5 page 18} \\
 &= 2\pi \sum_{m \in \mathbb{Z}} \tilde{g}(\omega + 2\pi m) \tilde{f}^*(\omega + 2\pi m) && \text{where } m \triangleq -n \\
 &= \tilde{S}_{gf}(\omega) && \text{by Theorem H.1 page 179}
 \end{aligned}$$

$$\begin{aligned}
 \tilde{S}_{fg}(\pi - \omega) &= 2\pi \sum_{n \in \mathbb{Z}} \tilde{f}(\pi - \omega + 2\pi n) \tilde{g}^*(\pi - \omega + 2\pi n) && \text{by Theorem H.1 page 179} \\
 &= 2\pi \sum_{n \in \mathbb{Z}} \tilde{f}^*(-\pi + \omega - 2\pi n) \tilde{g}(-\pi + \omega - 2\pi n) && \text{by hypothesis and Theorem 4.5 page 18} \\
 &= 2\pi \sum_{n \in \mathbb{Z}} \tilde{f}^*(\pi + \omega - 2\pi - 2\pi n) \tilde{g}(\pi + \omega - 2\pi - 2\pi n) \\
 &= 2\pi \sum_{n \in \mathbb{Z}} \tilde{f}^*(\pi + \omega + 2\pi(-n-1)) \tilde{g}(\pi + \omega + 2\pi(-n-1)) \\
 &= 2\pi \sum_{m \in \mathbb{Z}} \tilde{g}(\pi + \omega + 2\pi m) \tilde{f}^*(\pi + \omega + 2\pi m) && \text{where } m \triangleq -n-1 \\
 &= \tilde{S}_{gf}(\pi + \omega) && \text{by Theorem H.1 page 179}
 \end{aligned}$$

$$\begin{aligned}
 \tilde{S}_{ff}(-\omega) &= \tilde{S}_{fg}(-\omega) \Big|_{g \triangleq f} \\
 &= \tilde{S}_{gf}(+\omega) \Big|_{g \triangleq f} && \text{by previous result} \\
 &= \tilde{S}_{ff}(+\omega) && \text{by definition of } g \text{ (} g \triangleq f \text{)}
 \end{aligned}$$

$$\tilde{S}_{ff}(\pi - \omega) = \tilde{S}_{fg}(\pi - \omega) \Big|_{g \triangleq f}$$

$$\begin{aligned}
&= \tilde{S}_{gf}(\pi + \omega) \Big|_{g \triangleq f} \\
&= \tilde{S}_{ff}(\pi + \omega)
\end{aligned}$$

by previous result

by definition of  $g$  ( $g \triangleq f$ )

⇒

**Proposition H.4.** Let  $\tilde{S}_{ff}(\omega)$  be the AUTO-POWER SPECTRUM (Definition H.3 page 179) of a function  $f(x) \in L^2_{\mathbb{R}}$  and  $\tilde{S}'_{ff}(\omega) \triangleq \frac{d}{d\omega} \tilde{S}_{ff}(\omega)$  (Definition B.2 page 69).

P R O P	$\left\{ \begin{array}{l} \text{(a). } f \text{ is REAL and} \\ \text{(b). } \tilde{S}_{ff}(\omega) \text{ is CONTINUOUS at } \omega = 0 \end{array} \right\} \Rightarrow \left\{ \begin{array}{l} \text{(1). } \tilde{S}'_{ff}(0) = 0 \text{ and} \\ \text{(2). } \underbrace{\tilde{S}'_{ff}(\omega) = -\tilde{S}'_{ff}(-\omega)}_{\text{ANTI-SYMMETRIC about 0}} \quad \forall \omega \in \mathbb{R} \end{array} \right\}$
	$\left\{ \begin{array}{l} \text{(c). } f \text{ is REAL and} \\ \text{(d). } \tilde{S}_{ff}(\omega) \text{ is CONTINUOUS at } \omega = \pi \end{array} \right\} \Rightarrow \left\{ \begin{array}{l} \text{(3). } \tilde{S}'_{ff}(\pi) = 0 \text{ and} \\ \text{(4). } \underbrace{\tilde{S}'_{ff}(\pi + \omega) = -\tilde{S}'_{ff}(\pi - \omega)}_{\text{ANTI-SYMMETRIC about } \pi} \quad \forall \omega \in \mathbb{R} \end{array} \right\}$

PROOF: This follows from Proposition H.3 (page 181) and Proposition B.1 (page 69).

⇒

Theorem H.2 (next) is a major result and provides strong motivation for bothering with *power spectrum* functions in the first place. In particular, the *auto-power spectrum* being *bounded* provides a necessary and sufficient condition for a sequence of functions  $(\phi(x - n))_{n \in \mathbb{Z}}$  to be a *Riesz basis* (Definition G.13 page 167) for the *span*  $(\phi(x - n))$  of the sequence.

**Theorem H.2.**<sup>5</sup> Let  $\tilde{S}_{ff}(\omega)$  be defined as in Definition H.3 (page 179). Let  $\|\cdot\|$  be defined as in Definition B.1 (page 69). Let  $0 < A < B$ .

T H M	$\left\{ A \sum_{n \in \mathbb{N}}  a_n ^2 \leq \left\  \sum_{n \in \mathbb{Z}} a_n \phi(x - n) \right\ ^2 \leq B \sum_{n \in \mathbb{N}}  \alpha_n ^2 \quad \forall (a_n) \in \ell^2_{\mathbb{F}} \right\} \iff \{ A \leq \tilde{S}_{\phi\phi}(\omega) \leq B \}$
	$(\phi(x - n)) \text{ is a RIESZ BASIS for } \text{span}(\phi(x - n)) \text{ (Theorem G.13 page 168)}$

PROOF:

1. lemma:

$$\begin{aligned}
\left\| \sum_{n \in \mathbb{Z}} a_n \phi(x - n) \right\|^2 &= \left\| \tilde{\mathbf{F}} \sum_{n \in \mathbb{Z}} a_n \phi(x - n) \right\|^2 \\
&= \|\check{a}(\omega) \check{\phi}(\omega)\|^2 \\
&= \int_{\mathbb{R}} |\check{a}(\omega) \check{\phi}(\omega)|^2 d\omega \\
&= \sum_{n \in \mathbb{Z}} \int_0^{2\pi} |\check{a}(\omega + 2\pi n) \check{\phi}(\omega + 2\pi n)|^2 d\omega \\
&= \int_0^{2\pi} \sum_{n \in \mathbb{Z}} |\check{a}(\omega + 2\pi n)|^2 |\check{\phi}(\omega + 2\pi n)|^2 d\omega \\
&= \int_0^{2\pi} \sum_{n \in \mathbb{Z}} |\check{a}(\omega)|^2 |\check{\phi}(\omega + 2\pi n)|^2 d\omega \\
&= \int_0^{2\pi} |\check{a}(\omega)|^2 \frac{1}{2\pi} 2\pi \sum_{n \in \mathbb{Z}} |\check{\phi}(\omega + 2\pi n)|^2 d\omega
\end{aligned}$$

because  $\tilde{\mathbf{F}}$  is *unitary* (Theorem 4.2 page 17)

by Proposition I.13 page 195

by definition of  $\|\cdot\|$ 

by Proposition 6.1 page 41

<sup>5</sup> Wojtaszczyk (1997) pages 22–23 (Proposition 2.8), Igari (1996) page 219 (Lemma 9.6), Pinsky (2002) page 306 (Theorem 6.4.8)



$$= \frac{1}{2\pi} \int_0^{2\pi} |\check{a}(\omega)|^2 \tilde{S}_{\phi\phi}(\omega) d\omega \quad \text{by definition of } \tilde{S}_{\phi\phi}(\omega) \text{ (Theorem H.1 page 179)}$$

2. lemma:

$$\begin{aligned} \frac{1}{2\pi} \int_0^{2\pi} |\check{a}(\omega)|^2 d\omega &= \frac{1}{2\pi} \int_0^{2\pi} \left| \sum_{n \in \mathbb{Z}} a_n e^{-i\omega n} \right|^2 d\omega && \text{by def. of DTFT (Definition 6.1 page 41)} \\ &= \frac{1}{2\pi} \int_0^{2\pi} \left[ \sum_{n \in \mathbb{Z}} a_n e^{-i\omega n} \right] \left[ \sum_{m \in \mathbb{Z}} a_m e^{-i\omega m} \right]^* d\omega \\ &= \frac{1}{2\pi} \int_0^{2\pi} \left[ \sum_{n \in \mathbb{Z}} a_n e^{-i\omega n} \right] \left[ \sum_{m \in \mathbb{Z}} a_m^* e^{i\omega m} \right] d\omega \\ &= \frac{1}{2\pi} \sum_{n \in \mathbb{Z}} \sum_{m \in \mathbb{Z}} a_n a_m^* \int_0^{2\pi} e^{-i\omega(n-m)} d\omega \\ &= \frac{1}{2\pi} \sum_{n \in \mathbb{Z}} \sum_{m \in \mathbb{Z}} a_n a_m^* 2\pi \bar{\delta}_{nm} \\ &= \sum_{n \in \mathbb{Z}} |a_n|^2 && \text{by definition of } \bar{\delta} \text{ (Definition G.12 page 160)} \end{aligned}$$

3. Proof for (  $\Leftarrow$  ) case:

$$\begin{aligned} \boxed{A \sum_{n \in \mathbb{Z}} |a_n|^2} &= \frac{A}{2\pi} \int_0^{2\pi} |\check{a}(\omega)|^2 d\omega && \text{by (2) lemma page 183} \\ &= \frac{1}{2\pi} \int_0^{2\pi} |\check{a}(\omega)|^2 A d\omega \\ &\leq \frac{1}{2\pi} \int_0^{2\pi} |\check{a}(\omega)|^2 \tilde{S}_{\phi\phi}(\omega) d\omega && \text{by right hypothesis} \\ &= \left\| \sum_{n \in \mathbb{Z}} a_n \phi(x-n) \right\|^2 && \text{by (1) lemma page 182} \\ &= \frac{1}{2\pi} \int_0^{2\pi} |\check{a}(\omega)|^2 \tilde{S}_{\phi\phi}(\omega) d\omega && \text{by (1) lemma page 182} \\ &\leq \frac{1}{2\pi} \int_0^{2\pi} |\check{a}(\omega)|^2 B d\omega && \text{by right hypothesis} \\ &= \frac{B}{2\pi} \int_0^{2\pi} |\check{a}(\omega)|^2 d\omega \\ &= \boxed{B \sum_{n \in \mathbb{Z}} |a_n|^2} && \text{by (2) lemma page 183} \end{aligned}$$

4. Proof for (  $\Rightarrow$  ) case:

- (a) Let  $Y \triangleq \{\omega \in [0 : 2\pi] | \tilde{S}_{\phi\phi}(\omega) > \alpha\}$   
and  $X \triangleq \{\omega \in [0 : 2\pi] | \tilde{S}_{\phi\phi}(\omega) < \alpha\}$
- (b) Let  $\mathbb{1}_{A(x)}$  be the *set indicator* (Definition I.2 page 188) of a set  $A$ .  
Let  $(b_n)_{n \in \mathbb{Z}}$  be the *inverse DTFT* (Theorem 6.3 page 47) of  $\mathbb{1}_Y(\omega)$  such that  
 $\mathbb{1}_Y(\omega) \triangleq \sum_{n \in \mathbb{N}} b_n e^{-i\omega n} \triangleq \tilde{b}(\omega)$ .
- Let  $(a_n)_{n \in \mathbb{Z}}$  be the *inverse DTFT* (Theorem 6.3 page 47) of  $\mathbb{1}_X(\omega)$  such that  
 $\mathbb{1}_X(\omega) \triangleq \sum_{n \in \mathbb{N}} a_n e^{-i\omega n} \triangleq \check{a}(\omega)$ .

(c) Proof that  $\alpha \leq B$ :

Let  $\mu(A)$  be the *measure* of a set  $A$ .

$$\begin{aligned}
 \boxed{B} \sum_{n \in \mathbb{Z}} |b_n|^2 &\geq \left\| \sum_{n \in \mathbb{Z}} b_n \phi(x - n) \right\|^2 && \text{by left hypothesis} \\
 &= \frac{1}{2\pi} \int_0^{2\pi} |\tilde{b}(\omega)|^2 \tilde{S}_{\phi\phi}(\omega) d\omega && \text{by (1) lemma page 182} \\
 &= \frac{1}{2\pi} \int_0^{2\pi} |\mathbb{1}_Y(\omega)|^2 \tilde{S}_{\phi\phi}(\omega) d\omega && \text{by definition of } \mathbb{1}_Y(\omega) \quad (\text{item (4b) page 183}) \\
 &= \frac{1}{2\pi} \int_Y |1|^2 \tilde{S}_{\phi\phi}(\omega) d\omega && \text{by definition of } \mathbb{1}_Y(\omega) \quad (\text{item (4b) page 183}) \\
 &\geq \frac{\alpha}{2\pi} \mu(Y) && \text{by definition of } Y \quad (\text{item (4a) page 183}) \\
 &= \int_0^{2\pi} |\mathbb{1}_Y(\omega)|^2 d\omega && \text{by definition of } \mathbb{1}_Y(\omega) \quad (\text{item (4b) page 183}) \\
 &= \int_0^{2\pi} \left| \sum_{n \in \mathbb{Z}} b_n e^{-i\omega n} \right|^2 d\omega && \text{by definition of } (b_n) \quad (\text{item (4b) page 183}) \\
 &= \int_0^{2\pi} |\tilde{b}(\omega)|^2 d\omega && \text{by definition of } \tilde{b}(\omega) \quad (\text{item (4b) page 183}) \\
 &= \boxed{\alpha} \sum_{n \in \mathbb{Z}} |b_n|^2 && \text{by (2) lemma page 183}
 \end{aligned}$$

(d) Proof that  $\tilde{S}_{\phi\phi}(\omega) \leq B$ :

- (i).  $\tilde{S}_{\phi\phi}(\omega) > \alpha$  whenever  $\omega \in Y$  (item (4a) page 183).
- (ii). But even then,  $\alpha \leq B$  (item (4c) page 184).
- (iii). So,  $\tilde{S}_{\phi\phi}(\omega) \leq B$ .

(e) Proof that  $A \leq \alpha$ :

Let  $\mu(A)$  be the *measure* of a set  $A$ .

$$\begin{aligned}
 \boxed{A} \sum_{n \in \mathbb{Z}} |a_n|^2 &\leq \left\| \sum_{n \in \mathbb{Z}} a_n \phi(x - n) \right\|^2 && \text{by left hypothesis} \\
 &= \frac{1}{2\pi} \int_0^{2\pi} |\tilde{a}(\omega)|^2 \tilde{S}_{\phi\phi}(\omega) d\omega && \text{by (1) lemma page 182} \\
 &= \frac{1}{2\pi} \int_0^{2\pi} |\mathbb{1}_X(\omega)|^2 \tilde{S}_{\phi\phi}(\omega) d\omega && \text{by definition of } \mathbb{1}_X(\omega) \quad (\text{Definition 1.2 page 188}) \\
 &= \frac{1}{2\pi} \int_X |1|^2 \tilde{S}_{\phi\phi}(\omega) d\omega && \text{by definition of } \mathbb{1}_X(\omega) \quad (\text{Definition 1.2 page 188}) \\
 &\leq \frac{\alpha}{2\pi} \mu(X) && \text{by definition of } X \quad (\text{item (4a) page 183}) \\
 &= \int_0^{2\pi} |\mathbb{1}_X(\omega)|^2 d\omega && \text{by definition of } \mathbb{1}_X(\omega) \quad (\text{Definition 1.2 page 188}) \\
 &= \int_0^{2\pi} \left| \sum_{n \in \mathbb{Z}} a_n e^{-i\omega n} \right|^2 d\omega && \text{by definition of } (a_n) \quad ((2) \text{ lemma page 183}) \\
 &= \int_0^{2\pi} |\tilde{a}(\omega)|^2 d\omega && \text{by definition of } \tilde{a}(\omega) \quad ((2) \text{ lemma page 183}) \\
 &= \boxed{\alpha} \sum_{n \in \mathbb{Z}} |a_n|^2 && \text{by (2) lemma page 183}
 \end{aligned}$$

(f) Proof that  $A \leq \tilde{S}_{\phi\phi}(\omega)$ :

- (i).  $\tilde{S}_{\phi\phi}(\omega) < \alpha$  whenever  $\omega \in X$  (item (4a) page 183).
- (ii). But even then,  $A \leq \alpha$  (item (4e) page 184).
- (iii). So,  $A \leq \tilde{S}_{\phi\phi}(\omega)$ .

⇒

In the case that  $f$  and  $g$  are *orthonormal*, the spectral density relations simplify considerably (next).

**Theorem H.3.** <sup>6</sup> Let  $\tilde{S}_{ff}$  and  $\tilde{S}_{fg}$  be the SPECTRAL DENSITY FUNCTIONS (Definition H.3 page 179).

**T  
H  
M**

$$\begin{aligned} \langle f(x) | f(x-n) \rangle &= \bar{\delta}_n \quad ((f(x-n)) \text{ is ORTHONORMAL}) &\iff \tilde{S}_{ff}(\omega) &= 1 \quad \forall f \in L^2_{\mathbb{F}} \\ \langle f(x) | g(x-n) \rangle &= 0 \quad (f(x) \text{ is ORTHOGONAL to } (g(x-n))) &\iff \tilde{S}_{fg}(\omega) &= 0 \quad \forall f, g \in L^2_{\mathbb{F}} \end{aligned}$$

PROOF:

1. Proof that  $\langle f(x) | f(x-n) \rangle = \bar{\delta}_n \iff \tilde{S}_{ff}(\omega) = 1$ : This follows directly from Theorem H.2 (page 182) with  $A = B = 1$  (by Parseval's Identity Theorem G.9 page 162 since  $\{T^n f\}$  is *orthonormal*)

2. Alternate proof that  $\langle f(x) | f(x-n) \rangle = \bar{\delta}_n \implies \tilde{S}_{ff}(\omega) = 1$ :

$$\begin{aligned} \tilde{S}_{ff}(\omega) &= \sum_{n \in \mathbb{Z}} R_{ff}(n) e^{-i\omega n} && \text{by definition of } \tilde{S}_{fg} && (\text{Definition H.3 page 179}) \\ &= \sum_{n \in \mathbb{Z}} \langle f(x) | f(x-n) \rangle e^{-i\omega n} && \text{by definition of } R_{ff} && (\text{Definition H.1 page 179}) \\ &= \sum_{n \in \mathbb{Z}} \bar{\delta}_n e^{-i\omega n} && \text{by left hypothesis} \\ &= 1 && \text{by definition of } \bar{\delta} && (\text{Definition G.12 page 160}) \end{aligned}$$

3. Alternate proof that  $\langle f(x) | f(x-n) \rangle = \bar{\delta}_n \iff \tilde{S}_{ff}(\omega) = 1$ :

$$\begin{aligned} &\langle f(x) | f(x-n) \rangle \\ &= \langle \tilde{F}f(x) | \tilde{F}f(x-n) \rangle && \text{by unitary property of } \tilde{F} && (\text{Theorem 4.3 page 17}) \\ &= \langle \tilde{f}(\omega) | e^{-i\omega n} \tilde{f}(\omega) \rangle && \text{by shift property of } \tilde{F} && (\text{Theorem 4.4 page 18}) \\ &= \int_{\mathbb{R}} \tilde{f}(\omega) e^{i\omega n} \tilde{f}^*(\omega) d\omega && \text{by definition of } \langle \triangle | \nabla \rangle && (\text{Definition B.1 page 69}) \\ &= \int_{\mathbb{R}} |\tilde{f}(\omega)|^2 e^{i\omega n} d\omega \\ &= \sum_{n \in \mathbb{Z}} \int_{2\pi n}^{2\pi(n+1)} |\tilde{f}(\omega)|^2 e^{i\omega n} d\omega \\ &= \sum_{n \in \mathbb{Z}} \int_0^{2\pi} |\tilde{f}(u + 2\pi n)|^2 e^{i(u+2\pi n)n} du && \text{where } u \triangleq \omega - 2\pi n \implies \omega = u + 2\pi n \\ &= \frac{1}{2\pi} \int_0^{2\pi} \left[ 2\pi \sum_{n \in \mathbb{Z}} |\tilde{f}(u + 2\pi n)|^2 \right] e^{iun} e^{i2\pi n^2} du \\ &= \frac{1}{2\pi} \int_0^{2\pi} \tilde{S}_{ff}(\omega) e^{iun} du && \text{by Theorem H.1 page 179} \\ &= \frac{1}{2\pi} \int_0^{2\pi} e^{iun} du && \text{by right hypothesis} \\ &= \bar{\delta}_n && \text{by definition of } \bar{\delta} && (\text{Definition G.12 page 160}) \end{aligned}$$

<sup>6</sup> [Hernández and Weiss \(1996\) page 50](#) (PROPOSITION 2.1.11), [Wojtaszczyk \(1997\) PAGE 23](#) (COROLLARY 2.9), [IGARI \(1996\) PAGES 214–215](#) (LEMMA 9.2), [PINSKY \(2002\) PAGE 306](#) (COROLLARY 6.4.9)

4. Proof that  $\langle f(x) | g(x - n) \rangle = 0 \implies \tilde{S}_{fg}(\omega) = 0$ :


$$\begin{aligned}
 \tilde{S}_{fg}(\omega) &= \sum_{n \in \mathbb{Z}} R_{fg}(n) e^{-i\omega n} && \text{by definition of } \tilde{S}_{fg} && (\text{Definition H.3 page 179}) \\
 &= \sum_{n \in \mathbb{Z}} \langle f(x) | g(x - n) \rangle e^{-i\omega n} && \text{by definition of } R_{fg} && (\text{Definition H.1 page 179}) \\
 &= \sum_{n \in \mathbb{Z}} 0 e^{-i\omega n} && \text{by left hypothesis} \\
 &= 0
 \end{aligned}$$

5. Proof that  $\langle f(x) | g(x - n) \rangle = 0 \iff \tilde{S}_{fg}(\omega) = 0$ :

$$\begin{aligned}
 &\langle f(x) | g(x - n) \rangle \\
 &= \langle \tilde{F}f(x) | \tilde{F}g(x - n) \rangle && \text{by unitary property of } \tilde{F} && (\text{Theorem 4.3 page 17}) \\
 &= \langle \tilde{f}(\omega) | e^{-i\omega n} \tilde{g}(\omega) \rangle && \text{by shift property of } \tilde{F} && (\text{Theorem 4.4 page 18}) \\
 &= \int_{\mathbb{R}} \tilde{f}(\omega) e^{i\omega n} \tilde{g}^*(\omega) d\omega && \text{by definition of } \langle \Delta | \nabla \rangle && (\text{Definition B.1 page 69}) \\
 &= \int_{\mathbb{R}} \tilde{f}(\omega) \tilde{g}^*(\omega) e^{i\omega n} d\omega \\
 &= \sum_{n \in \mathbb{Z}} \int_{2\pi n}^{2\pi(n+1)} \tilde{f}(\omega) \tilde{g}^*(\omega) e^{i\omega n} d\omega \\
 &= \sum_{n \in \mathbb{Z}} \int_0^{2\pi} \tilde{f}(u + 2\pi n) \tilde{g}^*(u + 2\pi n) e^{i(u+2\pi n)n} du && \text{where } u \triangleq \omega - 2\pi n \implies \omega = u + 2\pi n \\
 &= \frac{1}{2\pi} \int_0^{2\pi} \left[ 2\pi \sum_{n \in \mathbb{Z}} \tilde{f}(u + 2\pi n) \tilde{g}^*(u + 2\pi n) \right] e^{iun} e^{i2\pi n^2} du \\
 &= \frac{1}{2\pi} \int_0^{2\pi} \tilde{S}_{fg}(u) e^{iun} du && \text{by Theorem H.1 page 179} \\
 &= \frac{1}{2\pi} \int_0^{2\pi} 0 \cdot e^{iun} du && \text{by right hypothesis} \\
 &= 0
 \end{aligned}$$

⇒

“Je me plaisois surtout aux mathématiques, à cause de la certitude et de l'évidence de leurs raisons: mais je ne remarquois point encore leur vrai usage; et, pensant qu'elles ne servoient qu'aux arts mécaniques, je m'étonnois de ce que leurs fondemens étant si fermes et si solides, on n'avoit rien bâti dessus de plus relevé.”



“I was especially delighted with the mathematics, on account of the certitude and evidence of their reasonings; but I had not as yet a precise knowledge of their true use; and thinking that they but contributed to the advancement of the mechanical arts, I was astonished that foundations, so strong and solid, should have had no loftier superstructure reared on them.”

René Descartes, philosopher and mathematician (1596–1650) <sup>1</sup>

## I.1 Families of Functions

This text is largely set in the space of *Lebesgue square-integrable functions*  $\mathcal{L}^2_{\mathbb{R}}$  (Definition B.1 page 69). The space  $\mathcal{L}^2_{\mathbb{R}}$  is a subspace of the space  $\mathbb{R}^{\mathbb{R}}$ , the set of all functions with *domain*  $\mathbb{R}$  (the set of real numbers) and *range*  $\mathbb{R}$ . The space  $\mathbb{R}^{\mathbb{R}}$  is a subspace of the space  $\mathbb{C}^{\mathbb{C}}$ , the set of all functions with *domain*  $\mathbb{C}$  (the set of complex numbers) and *range*  $\mathbb{C}$ . That is,  $\mathcal{L}^2_{\mathbb{R}} \subseteq \mathbb{R}^{\mathbb{R}} \subseteq \mathbb{C}^{\mathbb{C}}$ . In general, the notation  $Y^X$  represents the set of all functions with domain  $X$  and range  $Y$  (Definition I.1 page 187). Although this notation may seem curious, note that for finite  $X$  and finite  $Y$ , the number of functions (elements) in  $Y^X$  is  $|Y^X| = |Y|^{|X|}$ .

**Definition I.1.** *Let  $X$  and  $Y$  be sets.*

**DEF** The space  $Y^X$  represents the set of all functions with DOMAIN  $X$  and RANGE  $Y$  such that  $Y^X \triangleq \{f(x) | f(x) : X \rightarrow Y\}$

<sup>1</sup> quote: Descartes (1637b)  
translation: Descartes (1637c) (part I, paragraph 10)  
image: [http://en.wikipedia.org/wiki/File:Frans\\_Hals\\_-\\_Portret\\_van\\_Ren%C3%A9\\_Descartes.jpg](http://en.wikipedia.org/wiki/File:Frans_Hals_-_Portret_van_Ren%C3%A9_Descartes.jpg), public domain

**Definition I.2.** <sup>2</sup> Let  $X$  be a set.

DEF

The **indicator function**  $\mathbb{1} \in \{0, 1\}^{2^X}$  is defined as

$$\mathbb{1}_A(x) = \begin{cases} 1 & \text{for } x \in A \\ 0 & \text{for } x \notin A \end{cases} \quad \forall x \in X, A \in 2^X$$

The indicator function  $\mathbb{1}$  is also called the **characteristic function**.

## I.2 Definitions and algebraic properties

Much of the wavelet theory developed in this text is constructed using the **translation operator**  $\mathbf{T}$  and the **dilation operator**  $\mathbf{D}$  (next).

**Definition I.3.** <sup>3</sup>

DEF


$\mathbf{T}_\tau$  is a **translation operator** on  $\mathbb{C}^\mathbb{C}$  if  $\mathbf{T}_\tau f(x) \triangleq f(x - \tau) \quad \forall f \in \mathbb{C}^\mathbb{C}$ .

$\mathbf{D}_\alpha$  is a **dilation operator** on  $\mathbb{C}^\mathbb{C}$  if  $\mathbf{D}_\alpha f(x) \triangleq f(\alpha x) \quad \forall f \in \mathbb{C}^\mathbb{C}$ .

Moreover,  $\mathbf{T} \triangleq \mathbf{T}_1$  and  $\mathbf{D} \triangleq \sqrt{2}\mathbf{D}_2$ .

**Example I.1.** Let  $\mathbf{T}$  and  $\mathbf{D}$  be defined as in Definition I.3 (page 188).

EX



**Proposition I.1.** Let  $\mathbf{T}_\tau$  be a TRANSLATION OPERATOR (Definition I.3 page 188).

PRP

$$\sum_{n \in \mathbb{Z}} \mathbf{T}_\tau^n f(x) = \sum_{n \in \mathbb{Z}} \mathbf{T}_\tau^n f(x + \tau) \quad \forall f \in \mathbb{R}^\mathbb{R} \quad \left( \sum_{n \in \mathbb{Z}} \mathbf{T}_\tau^n f(x) \text{ is PERIODIC with period } \tau \right)$$

PROOF:

$$\begin{aligned} \sum_{n \in \mathbb{Z}} \mathbf{T}_\tau^n f(x + \tau) &= \sum_{n \in \mathbb{Z}} f(x - n\tau + \tau) && \text{by definition of } \mathbf{T}_\tau && (\text{Definition I.3 page 188}) \\ &= \sum_{m \in \mathbb{Z}} f(x - m\tau) && \text{where } m \triangleq n - 1 && \implies n = m + 1 \\ &= \sum_{m \in \mathbb{Z}} \mathbf{T}_\tau^m f(x) && \text{by definition of } \mathbf{T}_\tau && (\text{Definition I.3 page 188}) \end{aligned}$$

In a linear space, every operator has an *inverse*. Although the inverse always exists as a *relation*, it may not exist as a *function* or as an *operator*. But in some cases the inverse of an operator is itself an operator. The inverses of the operators  $\mathbf{T}$  and  $\mathbf{D}$  both exist as operators, as demonstrated next.

<sup>2</sup> Aliprantis and Burkinshaw (1998) page 126, Hausdorff (1937) page 22, de la Vallée-Poussin (1915) page 440

<sup>3</sup> Walnut (2002) pages 79–80 (Definition 3.39), Christensen (2003) pages 41–42, Wojtaszczyk (1997) page 18 (Definitions 2.3, 2.4), Kammler (2008) page A-21, Bachman et al. (2000) page 473, Packer (2004) page 260, Zay (2004) page, Heil (2011) page 250 (Notation 9.4), Casazza and Lammers (1998) page 74, Goodman et al. (1993a) page 639, Heil and Walnut (1989) page 633 (Definition 1.3.1), Dai and Lu (1996) page 81, Dai and Larson (1998) page 2

**Proposition I.2** (transversal operator inverses). *Let  $\mathbf{T}$  and  $\mathbf{D}$  be as defined in Definition I.3 page 188.*

PRP

$\mathbf{T}$  has an INVERSE  $\mathbf{T}^{-1}$  in  $\mathbb{C}^{\mathbb{C}}$  expressed by the relation

$$\mathbf{T}^{-1}\mathbf{f}(x) = \mathbf{f}(x+1) \quad \forall \mathbf{f} \in \mathbb{C}^{\mathbb{C}} \quad (\text{translation operator inverse}).$$

$\mathbf{D}$  has an INVERSE  $\mathbf{D}^{-1}$  in  $\mathbb{C}^{\mathbb{C}}$  expressed by the relation

$$\mathbf{D}^{-1}\mathbf{f}(x) = \frac{\sqrt{2}}{2} \mathbf{f}\left(\frac{1}{2}x\right) \quad \forall \mathbf{f} \in \mathbb{C}^{\mathbb{C}} \quad (\text{dilation operator inverse}).$$

✎ PROOF:

1. Proof that  $\mathbf{T}^{-1}$  is the inverse of  $\mathbf{T}$ :

$$\begin{aligned} \mathbf{T}^{-1}\mathbf{T}\mathbf{f}(x) &= \mathbf{T}^{-1}\mathbf{f}(x-1) && \text{by definition of } \mathbf{T} && (\text{Definition I.3 page 188}) \\ &= \mathbf{f}([x+1]-1) \\ &= \mathbf{f}(x) \\ &= \mathbf{f}([x-1]+1) \\ &= \mathbf{T}\mathbf{f}(x+1) && \text{by definition of } \mathbf{T} && (\text{Definition I.3 page 188}) \\ &= \mathbf{T}\mathbf{T}^{-1}\mathbf{f}(x) \\ \implies \mathbf{T}^{-1}\mathbf{T} &= \mathbf{I} = \mathbf{T}\mathbf{T}^{-1} \end{aligned}$$

2. Proof that  $\mathbf{D}^{-1}$  is the inverse of  $\mathbf{D}$ :

$$\begin{aligned} \mathbf{D}^{-1}\mathbf{D}\mathbf{f}(x) &= \mathbf{D}^{-1}\sqrt{2}\mathbf{f}(2x) && \text{by definition of } \mathbf{D} && (\text{Definition I.3 page 188}) \\ &= \left(\frac{\sqrt{2}}{2}\right)\sqrt{2}\mathbf{f}\left(2\left[\frac{1}{2}x\right]\right) \\ &= \mathbf{f}(x) \\ &= \sqrt{2}\left[\frac{\sqrt{2}}{2}\mathbf{f}\left(\frac{1}{2}[2x]\right)\right] \\ &= \mathbf{D}\left[\frac{\sqrt{2}}{2}\mathbf{f}\left(\frac{1}{2}x\right)\right] && \text{by definition of } \mathbf{D} && (\text{Definition I.3 page 188}) \\ &= \mathbf{D}\mathbf{D}^{-1}\mathbf{f}(x) \\ \implies \mathbf{D}^{-1}\mathbf{D} &= \mathbf{I} = \mathbf{D}\mathbf{D}^{-1} \end{aligned}$$

⇒

**Proposition I.3.** *Let  $\mathbf{T}$  and  $\mathbf{D}$  be as defined in Definition I.3 page 188.*

*Let  $\mathbf{D}^0 = \mathbf{T}^0 \triangleq \mathbf{I}$  be the IDENTITY OPERATOR.*

PRP

$$\mathbf{D}^j\mathbf{T}^n\mathbf{f}(x) = 2^{j/2}\mathbf{f}(2^jx - n) \quad \forall j, n \in \mathbb{Z}, \mathbf{f} \in \mathbb{C}^{\mathbb{C}}$$

## I.3 Linear space properties

**Proposition I.4.** *Let  $\mathbf{T}$  and  $\mathbf{D}$  be as in Definition I.3 page 188.*

PRP

$$\mathbf{D}^j\mathbf{T}^n[\mathbf{f}\mathbf{g}] = 2^{-j/2} [\mathbf{D}^j\mathbf{T}^n\mathbf{f}] [\mathbf{D}^j\mathbf{T}^n\mathbf{g}] \quad \forall j, n \in \mathbb{Z}, \mathbf{f} \in \mathbb{C}^{\mathbb{C}}$$


✎ PROOF:

$$\begin{aligned} \mathbf{D}^j\mathbf{T}^n[\mathbf{f}(x)\mathbf{g}(x)] &= 2^{j/2}\mathbf{f}(2^jx - n)\mathbf{g}(2^jx - n) && \text{by Proposition I.3 page 189} \\ &= 2^{-j/2}[2^{j/2}\mathbf{f}(2^jx - n)][2^{j/2}\mathbf{g}(2^jx - n)] \\ &= 2^{-j/2}[\mathbf{D}^j\mathbf{T}^n\mathbf{f}(x)][\mathbf{D}^j\mathbf{T}^n\mathbf{g}(x)] && \text{by Proposition I.3 page 189} \end{aligned}$$



In general the operators  $\mathbf{T}$  and  $\mathbf{D}$  are *noncommutative* ( $\mathbf{TD} \neq \mathbf{DT}$ ), as demonstrated by Counterexample I.1 (next) and Proposition I.5 (page 190).

Counterexample I.1.



**Proposition I.5** (commutator relation). <sup>4</sup> Let  $\mathbf{T}$  and  $\mathbf{D}$  be as in Definition I.3 page 188.

P R P

$$\begin{aligned} \mathbf{D}^j \mathbf{T}^n &= \mathbf{T}^{2^{-j/2}n} \mathbf{D}^j & \forall j, n \in \mathbb{Z} \\ \mathbf{T}^n \mathbf{D}^j &= \mathbf{D}^j \mathbf{T}^{2^j n} & \forall n, j \in \mathbb{Z} \end{aligned}$$

PROOF:

$$\begin{aligned} \mathbf{D}^j \mathbf{T}^{2^j n} f(x) &= 2^{j/2} f(2^j x - 2^j n) && \text{by Proposition I.4 page 189} \\ &= 2^{j/2} f(2^j [x - n]) && \text{by distributivity of the field } (\mathbb{R}, +, \cdot, 0, 1) \text{ (Definition A.6 page 68)} \\ &= \mathbf{T}^n 2^{j/2} f(2^j x) && \text{by definition of } \mathbf{T} \text{ (Definition I.3 page 188)} \\ &= \mathbf{T}^n \mathbf{D}^j f(x) && \text{by definition of } \mathbf{D} \text{ (Definition I.3 page 188)} \end{aligned}$$

$$\begin{aligned} \mathbf{D}^j \mathbf{T}^n f(x) &= 2^{j/2} f(2^j x - n) && \text{by Proposition I.4 page 189} \\ &= 2^{j/2} f(2^j [x - 2^{-j/2} n]) && \text{by distributivity of the field } (\mathbb{R}, +, \cdot, 0, 1) \text{ (Definition A.6 page 68)} \\ &= \mathbf{T}^{2^{-j/2}n} 2^{j/2} f(2^j x) && \text{by definition of } \mathbf{T} \text{ (Definition I.3 page 188)} \\ &= \mathbf{T}^{2^{-j/2}n} \mathbf{D}^j f(x) && \text{by definition of } \mathbf{D} \text{ (Definition I.3 page 188)} \end{aligned}$$



## I.4 Inner product space properties

In an inner product space, every operator has an *adjoint* (Proposition F.3 page 133) and this adjoint is always itself an operator. In the case where the adjoint and inverse of an operator  $\mathbf{U}$  coincide, then  $\mathbf{U}$  is said to be *unitary* (Definition F.14 page 143). And in this case,  $\mathbf{U}$  has several nice properties (see Proposition I.9 and Theorem I.1 page 193). Proposition I.6 (next) gives the adjoints of  $\mathbf{D}$  and  $\mathbf{T}$ , and Proposition I.7 (page 191) demonstrates that both  $\mathbf{D}$  and  $\mathbf{T}$  are unitary. Other examples of unitary operators include the *Fourier Transform operator*  $\tilde{\mathbf{F}}$  (Corollary 4.1 page 17) and the *rotation matrix operator* (Example F.5 page 145).

**Proposition I.6.** Let  $\mathbf{T}$  be the TRANSLATION OPERATOR (Definition I.3 page 188) with ADJOINT  $\mathbf{T}^*$  and  $\mathbf{D}$  the DILATION OPERATOR with ADJOINT  $\mathbf{D}^*$  (Definition F.8 page 129).

P R P

$$\begin{aligned} \mathbf{T}^* f(x) &= f(x + 1) & \forall f \in \mathcal{L}_{\mathbb{R}}^2 & \text{ (TRANSLATION OPERATOR ADJOINT)} \\ \mathbf{D}^* f(x) &= \frac{\sqrt{2}}{2} f\left(\frac{1}{2}x\right) & \forall f \in \mathcal{L}_{\mathbb{R}}^2 & \text{ (DILATION OPERATOR ADJOINT)} \end{aligned}$$

<sup>4</sup> Christensen (2003) page 42 (equation (2.9)), Dai and Larson (1998) page 21, Goodman et al. (1993a) page 641, Goodman et al. (1993b) page 110



 PROOF:

1. Proof that  $\mathbf{T}^*f(x) = f(x + 1)$ :

$$\begin{aligned}\langle g(x) | \mathbf{T}^*f(x) \rangle &= \langle g(u) | \mathbf{T}^*f(u) \rangle \\ &= \langle \mathbf{T}g(u) | f(u) \rangle \\ &= \langle g(u - 1) | f(u) \rangle \\ &= \langle g(x) | f(x + 1) \rangle \\ \implies \mathbf{T}^*f(x) &= f(x + 1)\end{aligned}$$

by change of variable  $x \rightarrow u$

by definition of adjoint  $\mathbf{T}^*$

(Definition F.8 page 129)

by definition of  $\mathbf{T}$

(Definition I.3 page 188)

where  $x \triangleq u - 1 \implies u = x + 1$

2. Proof that  $\mathbf{D}^*f(x) = \frac{\sqrt{2}}{2} f\left(\frac{1}{2}x\right)$ :

$$\begin{aligned}\langle g(x) | \mathbf{D}^*f(x) \rangle &= \langle g(u) | \mathbf{D}^*f(u) \rangle \\ &= \langle \mathbf{D}g(u) | f(u) \rangle \\ &= \left\langle \sqrt{2}g(2u) | f(u) \right\rangle \\ &= \int_{u \in \mathbb{R}} \sqrt{2}g(2u)f^*(u) du \\ &= \int_{x \in \mathbb{R}} g(x) \left[ \sqrt{2}f\left(\frac{x}{2}\right)\frac{1}{2} \right]^* dx \\ &= \left\langle g(x) | \frac{\sqrt{2}}{2}f\left(\frac{x}{2}\right) \right\rangle \\ \implies \mathbf{D}^*f(x) &= \frac{\sqrt{2}}{2} f\left(\frac{x}{2}\right)\end{aligned}$$

by change of variable  $x \rightarrow u$

by definition of  $\mathbf{D}^*$

(Definition F.8 page 129)

by definition of  $\mathbf{D}$

(Definition I.3 page 188)

by definition of  $\langle \Delta | \nabla \rangle$

where  $x = 2u$

by definition of  $\langle \Delta | \nabla \rangle$



**Proposition I.7.** <sup>5</sup> Let  $\mathbf{T}$  and  $\mathbf{D}$  be as in Definition I.3 (page 188).

Let  $\mathbf{T}^{-1}$  and  $\mathbf{D}^{-1}$  be as in Proposition I.2 (page 189).

<b>P R P</b>	$\mathbf{T}$ is UNITARY in $L_{\mathbb{R}}^2$ ( $\mathbf{T}^{-1} = \mathbf{T}^*$ in $L_{\mathbb{R}}^2$ ).
	$\mathbf{D}$ is UNITARY in $L_{\mathbb{R}}^2$ ( $\mathbf{D}^{-1} = \mathbf{D}^*$ in $L_{\mathbb{R}}^2$ ).

 PROOF:

$$\mathbf{T}^{-1} = \mathbf{T}^*$$

$$\implies \mathbf{T} \text{ is unitary}$$

by Proposition I.2 page 189 and Proposition I.6 page 190

by the definition of *unitary* operators (Definition F.14 page 143)

$$\mathbf{D}^{-1} = \mathbf{D}^*$$

$$\implies \mathbf{D} \text{ is unitary}$$

by Proposition I.2 page 189 and Proposition I.6 page 190



by the definition of *unitary* operators (Definition F.14 page 143)



## I.5 Normed linear space properties

**Proposition I.8.** Let  $\mathbf{D}$  be the DILATION OPERATOR (Definition I.3 page 188).

<b>P R P</b>	$\left\{ \begin{array}{l} (1). \quad \mathbf{D}f(x) = \sqrt{2}f(x) \\ (2). \quad f(x) \text{ is CONTINUOUS} \end{array} \right\}$	$\iff$	$\{f(x) \text{ is a CONSTANT}\}$	$\forall f \in L_{\mathbb{R}}^2$

<sup>5</sup>  Christensen (2003) page 41 (Lemma 2.5.1),  Wojtaszczyk (1997) page 18 (Lemma 2.5)

✎ PROOF:

1. Proof that (1)  $\Leftarrow$  *constant* property:

$$\begin{aligned} Df(x) &\triangleq \sqrt{2}f(2x) && \text{by definition of } D && (\text{Definition 1.3 page 188}) \\ &= \sqrt{2}f(x) && \text{by } \textit{constant} \text{ hypothesis} \end{aligned}$$

2. Proof that (2)  $\Leftarrow$  *constant* property:

$$\begin{aligned} \|f(x) - f(x+h)\| &= \|f(x) - f(x)\| && \text{by } \textit{constant} \text{ hypothesis} \\ &= \|0\| \\ &= 0 && \text{by } \textit{nondegenerate} \text{ property of } \|\cdot\| \\ &\leq \varepsilon \\ &\Rightarrow \forall h > 0, \exists \varepsilon \text{ such that } \|f(x) - f(x+h)\| < \varepsilon \\ &\stackrel{\text{def}}{\Leftrightarrow} f(x) \text{ is } \textit{continuous} \end{aligned}$$

3. Proof that (1,2)  $\Rightarrow$  *constant* property:

(a) Suppose there exists  $x, y \in \mathbb{R}$  such that  $f(x) \neq f(y)$ .

(b) Let  $(x_n)_{n \in \mathbb{N}}$  be a sequence with limit  $x$  and  $(y_n)_{n \in \mathbb{N}}$  a sequence with limit  $y$

(c) Then

$$\begin{aligned} 0 &< \|f(x) - f(y)\| && \text{by assumption in item (3a) page 192} \\ &= \lim_{n \rightarrow \infty} \|f(x_n) - f(y_n)\| && \text{by (2) and definition of } (x_n) \text{ and } (y_n) \text{ in item (3b) page 192} \\ &= \lim_{n \rightarrow \infty} \|f(2^m x_n) - f(2^\ell y_n)\| \quad \forall m, \ell \in \mathbb{Z} \quad \text{by (1)} \\ &= 0 \end{aligned}$$

(d) But this is a *contradiction*, so  $f(x) = f(y)$  for all  $x, y \in \mathbb{R}$ , and  $f(x)$  is *constant*.

⇒

*Remark I.1.*


**REM** In Proposition 1.8 page 191, it is not possible to remove the *continuous* constraint outright, as demonstrated by the next two counterexamples.

*Counterexample I.2.* Let  $f(x)$  be a function in  $\mathbb{R}^{\mathbb{R}}$ .

CNT

$$\text{Let } f(x) \triangleq \begin{cases} 0 & \text{for } x = 0 \\ 1 & \text{otherwise.} \end{cases}$$

$$\text{Then } Df(x) \triangleq \sqrt{2}f(2x) = \sqrt{2}f(x), \text{ but } f(x) \text{ is } \textit{not constant}.$$




*Counterexample I.3.* Let  $f(x)$  be a function in  $\mathbb{R}^{\mathbb{R}}$ .

Let  $\mathbb{Q}$  be the set of *rational numbers* and  $\mathbb{R} \setminus \mathbb{Q}$  the set of *irrational numbers*.

CNT

$$\text{Let } f(x) \triangleq \begin{cases} 1 & \text{for } x \in \mathbb{Q} \\ -1 & \text{for } x \in \mathbb{R} \setminus \mathbb{Q}. \end{cases}$$

$$\text{Then } Df(x) \triangleq \sqrt{2}f(2x) = \sqrt{2}f(x), \text{ but } f(x) \text{ is } \textit{not constant}.$$



**Proposition I.9** (Operator norm). *Let  $\mathbf{T}$  and  $\mathbf{D}$  be as in Definition I.3 page 188. Let  $\mathbf{T}^{-1}$  and  $\mathbf{D}^{-1}$  be as in Proposition I.2 page 189. Let  $\mathbf{T}^*$  and  $\mathbf{D}^*$  be as in Proposition I.6 page 190. Let  $\|\cdot\|$  and  $\langle \triangle | \nabla \rangle$  be as in Definition B.1 page 69. Let  $\|\cdot\|$  be the operator norm (Definition F.6 page 125) induced by  $\|\cdot\|$ .*

$$\|\mathbf{T}\| = \|\mathbf{D}\| = \|\mathbf{T}^*\| = \|\mathbf{D}^*\| = \|\mathbf{T}^{-1}\| = \|\mathbf{D}^{-1}\| = 1$$

PROOF: These results follow directly from the fact that  $\mathbf{T}$  and  $\mathbf{D}$  are *unitary* (Proposition I.7 page 191) and from Theorem E.25 page 144 and Theorem F.26 page 144.  $\Rightarrow$

**Theorem I.1.** *Let  $\mathbf{T}$  and  $\mathbf{D}$  be as in Definition I.3 page 188.*

*Let  $\mathbf{T}^{-1}$  and  $\mathbf{D}^{-1}$  be as in Proposition I.2 page 189. Let  $\|\cdot\|$  and  $\langle \triangle | \nabla \rangle$  be as in Definition B.1 page 69.*

<b>T H M</b>	1.	$\ \mathbf{T}f\ $	$=$	$\ \mathbf{D}f\ $	$=$	$\ f\ $	$\forall f \in L^2_{\mathbb{R}}$	(ISOMETRIC IN LENGTH)
	2.	$\ \mathbf{T}f - \mathbf{T}g\ $	$=$	$\ \mathbf{D}f - \mathbf{D}g\ $	$=$	$\ f - g\ $	$\forall f, g \in L^2_{\mathbb{R}}$	(ISOMETRIC IN DISTANCE)
	3.	$\ \mathbf{T}^{-1}f - \mathbf{T}^{-1}g\ $	$=$	$\ \mathbf{D}^{-1}f - \mathbf{D}^{-1}g\ $	$=$	$\ f - g\ $	$\forall f, g \in L^2_{\mathbb{R}}$	(ISOMETRIC IN DISTANCE)
	4.	$\langle \mathbf{T}f   \mathbf{T}g \rangle$	$=$	$\langle \mathbf{D}f   \mathbf{D}g \rangle$	$=$	$\langle f   g \rangle$	$\forall f, g \in L^2_{\mathbb{R}}$	(SURJECTIVE)
	5.	$\langle \mathbf{T}^{-1}f   \mathbf{T}^{-1}g \rangle$	$=$	$\langle \mathbf{D}^{-1}f   \mathbf{D}^{-1}g \rangle$	$=$	$\langle f   g \rangle$	$\forall f, g \in L^2_{\mathbb{R}}$	(SURJECTIVE)

PROOF: These results follow directly from the fact that  $\mathbf{T}$  and  $\mathbf{D}$  are *unitary* (Proposition I.7 page 191) and from Theorem E.25 page 144 and Theorem F.26 page 144.  $\Rightarrow$

**Proposition I.10.** *Let  $\mathbf{T}$  be as in Definition I.3 page 188. Let  $\mathbf{A}^*$  be the ADJOINT (Definition F.8 page 129) of an operator  $\mathbf{A}$ . Let the property “SELF ADJOINT” be defined as in Definition F.11 (page 137).*

$$\left( \sum_{n \in \mathbb{Z}} \mathbf{T}^n \right) = \left( \sum_{n \in \mathbb{Z}} \mathbf{T}^n \right)^* \quad \left( \text{The operator } \left[ \sum_{n \in \mathbb{Z}} \mathbf{T}^n \right] \text{ is SELF-ADJOINT} \right)$$

PROOF:

$$\begin{aligned}
 \left\langle \left( \sum_{n \in \mathbb{Z}} \mathbf{T}^n \right) f(x) \mid g(x) \right\rangle &= \left\langle \sum_{n \in \mathbb{Z}} f(x-n) \mid g(x) \right\rangle && \text{by definition of } \mathbf{T} && (\text{Definition I.3 page 188}) \\
 &= \left\langle \sum_{n \in \mathbb{Z}} f(x+n) \mid g(x) \right\rangle && \text{by commutative property} && (\text{Definition A.5 page 68}) \\
 &= \sum_{n \in \mathbb{Z}} \langle f(x+n) \mid g(x) \rangle && \text{by additive property of } \langle \triangle | \nabla \rangle && \\
 &= \sum_{n \in \mathbb{Z}} \langle f(u) \mid g(u-n) \rangle && \text{where } u \triangleq x+n && \\
 &= \left\langle f(u) \mid \sum_{n \in \mathbb{Z}} g(u-n) \right\rangle && \text{by additive property of } \langle \triangle | \nabla \rangle && \\
 &= \left\langle f(x) \mid \sum_{n \in \mathbb{Z}} g(x-n) \right\rangle && \text{by change of variable: } u \rightarrow x && \\
 &= \left\langle f(x) \mid \sum_{n \in \mathbb{Z}} \mathbf{T}^n g(x) \right\rangle && \text{by definition of } \mathbf{T} && (\text{Definition I.3 page 188}) \\
 &\Leftrightarrow \left( \sum_{n \in \mathbb{Z}} \mathbf{T}^n \right) = \left( \sum_{n \in \mathbb{Z}} \mathbf{T}^n \right)^* && \text{by definition of adjoint} && (\text{Proposition F.3 page 133}) \\
 &\Leftrightarrow \left( \sum_{n \in \mathbb{Z}} \mathbf{T}^n \right) \text{ is self-adjoint} && \text{by definition of self-adjoint} && (\text{Definition F.11 page 137})
 \end{aligned}$$

$\Rightarrow$

## I.6 Fourier transform properties

**Proposition I.11.** Let  $\mathbf{T}$  and  $\mathbf{D}$  be as in Definition I.3 page 188.

Let  $\mathbf{B}$  be the TWO-SIDED LAPLACE TRANSFORM defined as  $[\mathbf{B}f](s) \triangleq \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} f(x)e^{-sx} dx$ .

PRP

- |    |  |   |
|----|--|---|
| 1. | $\mathbf{B}\mathbf{T}^n = e^{-sn}\mathbf{B}$   | $\forall n \in \mathbb{Z}$  |
| 2. | $\mathbf{B}\mathbf{D}^j = \mathbf{D}^{-j}\mathbf{B}$   | $\forall j \in \mathbb{Z}$  |
| 3. | $\mathbf{D}\mathbf{B} = \mathbf{B}\mathbf{D}^{-1}$   | $\forall n \in \mathbb{Z}$  |
| 4. | $\mathbf{B}\mathbf{D}^{-1}\mathbf{B}^{-1} = \mathbf{B}^{-1}\mathbf{D}^{-1}\mathbf{B} = \mathbf{D}$ | $\forall n \in \mathbb{Z}$ ( $\mathbf{D}^{-1}$ is SIMILAR to $\mathbf{D}$ ) |
| 5. | $\mathbf{D}\mathbf{B}\mathbf{D} = \mathbf{D}^{-1}\mathbf{B}\mathbf{D}^{-1} = \mathbf{B}$           | $\forall n \in \mathbb{Z}$  |

 PROOF:

$$\mathbf{B}\mathbf{T}^n f(x) = \mathbf{B}f(x-n) \quad \text{by definition of } \mathbf{T} \quad (\text{Definition I.3 page 188})$$

$$= \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} f(x-n)e^{-sx} dx \quad \text{by definition of } \mathbf{B}$$

$$= \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} f(u)e^{-s(u+n)} du \quad \text{where } u \triangleq x-n$$

$$= e^{-sn} \left[ \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} f(u)e^{-su} du \right] \quad \text{by definition of } \mathbf{B}$$

$$= e^{-sn} \mathbf{B}f(x)$$

$$\mathbf{B}\mathbf{D}^j f(x) = \mathbf{B}[2^{j/2} f(2^j x)] \quad \text{by definition of } \mathbf{D} \quad (\text{Definition I.3 page 188})$$

$$= \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} [2^{j/2} f(2^j x)] e^{-sx} dx \quad \text{by definition of } \mathbf{B}$$

$$= \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} [2^{j/2} f(u)] e^{-s2^{-j} 2^j u} du \quad \text{let } u \triangleq 2^j x \implies x = 2^{-j} u$$

$$= \frac{\sqrt{2}}{2} \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} f(u) e^{-s2^{-j} u} du$$

$$= \mathbf{D}^{-1} \left[ \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} f(u) e^{-su} du \right] \quad \text{by Proposition I.6 page 190 and Proposition I.7 page 191}$$

$$= \mathbf{D}^{-j} \mathbf{B}f(x) \quad \text{by definition of } \mathbf{B}$$

$$\mathbf{D}\mathbf{B} f(x) = \mathbf{D} \left[ \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} f(x) e^{-sx} dx \right] \quad \text{by definition of } \mathbf{B}$$

$$= \frac{\sqrt{2}}{\sqrt{2\pi}} \int_{\mathbb{R}} f(x) e^{-2sx} dx \quad \text{by definition of } \mathbf{D} \quad (\text{Definition I.3 page 188})$$

$$= \frac{\sqrt{2}}{\sqrt{2\pi}} \int_{\mathbb{R}} f\left(\frac{u}{2}\right) e^{-su} \frac{1}{2} du \quad \text{let } u \triangleq 2x \implies x = \frac{1}{2}u$$

$$= \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} \left[ \frac{\sqrt{2}}{2} f\left(\frac{u}{2}\right) \right] e^{-su} du$$

$$= \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} [\mathbf{D}^{-1}f](u) e^{-su} du \quad \text{by Proposition I.6 page 190 and Proposition I.7 page 191}$$

$$= \mathbf{B}\mathbf{D}^{-1}f(x) \quad \text{by definition of } \mathbf{B}$$

$$\mathbf{B}^{-1}\mathbf{D}^{-1}\mathbf{B} = \mathbf{B}^{-1}\mathbf{B}\mathbf{D} \quad \text{by previous result}$$

$$= \mathbf{D} \quad \text{by definition of operator inverse} \quad (\text{Definition F.3 page 120})$$

$$\mathbf{B}\mathbf{D}^{-1}\mathbf{B}^{-1} = \mathbf{D}\mathbf{B}\mathbf{B}^{-1} \quad \text{by previous result}$$

$$= \mathbf{D} \quad \text{by definition of operator inverse} \quad (\text{Definition F.3 page 120})$$

$$\begin{aligned}
\mathbf{D}\mathbf{B}\mathbf{D} &= \mathbf{D}\mathbf{D}^{-1}\mathbf{B} && \text{by previous result} \\
&= \mathbf{B} && \text{by definition of operator inverse (Definition F.3 page 120)} \\
\mathbf{D}^{-1}\mathbf{B}\mathbf{D}^{-1} &= \mathbf{D}^{-1}\mathbf{D}\mathbf{B} && \text{by previous result} \\
&= \mathbf{B} && \text{by definition of operator inverse (Definition F.3 page 120)}
\end{aligned}$$

⇒

**Corollary I.1.** Let  $\mathbf{T}$  and  $\mathbf{D}$  be as in Definition I.3 page 188. Let  $\tilde{\mathbf{f}}(\omega) \triangleq \tilde{\mathbf{F}}\mathbf{f}(x)$  be the FOURIER TRANSFORM (Definition 4.2 page 16) of some function  $\mathbf{f} \in \mathcal{L}_{\mathbb{R}}^2$  (Definition B.1 page 69).

C O R	1. $\tilde{\mathbf{F}}\mathbf{T}^n = e^{-i\omega n}\tilde{\mathbf{F}}$
	2. $\tilde{\mathbf{F}}\mathbf{D}^j = \mathbf{D}^{-j}\tilde{\mathbf{F}}$
	3. $\mathbf{D}\tilde{\mathbf{F}} = \tilde{\mathbf{F}}\mathbf{D}^{-1}$
	4. $\mathbf{D} = \tilde{\mathbf{F}}\mathbf{D}^{-1}\tilde{\mathbf{F}}^{-1} = \tilde{\mathbf{F}}^{-1}\mathbf{D}^{-1}\tilde{\mathbf{F}}$
	5. $\tilde{\mathbf{F}} = \mathbf{D}\tilde{\mathbf{F}}\mathbf{D} = \mathbf{D}^{-1}\tilde{\mathbf{F}}\mathbf{D}^{-1}$

PROOF: These results follow directly from Proposition I.11 page 194 with  $\tilde{\mathbf{F}} = \mathbf{B}|_{s=i\omega}$ .

⇒

**Proposition I.12.** Let  $\mathbf{T}$  and  $\mathbf{D}$  be as in Definition I.3 page 188. Let  $\tilde{\mathbf{f}}(\omega) \triangleq \tilde{\mathbf{F}}\mathbf{f}(x)$  be the FOURIER TRANSFORM (Definition 4.2 page 16) of some function  $\mathbf{f} \in \mathcal{L}_{\mathbb{R}}^2$  (Definition B.1 page 69).

P R O P	$\tilde{\mathbf{F}}\mathbf{D}^j\mathbf{T}^n\mathbf{f}(x) = \frac{1}{2^{j/2}}e^{-i\frac{\omega}{2^j}n}\tilde{\mathbf{f}}\left(\frac{\omega}{2^j}\right)$
------------------	---

PROOF:

$$\begin{aligned}
\tilde{\mathbf{F}}\mathbf{D}^j\mathbf{T}^n\mathbf{f}(x) &= \mathbf{D}^{-j}\tilde{\mathbf{F}}\mathbf{T}^n\mathbf{f}(x) && \text{by Corollary I.1 page 195 (3)} \\
&= \mathbf{D}^{-j}e^{-i\omega n}\tilde{\mathbf{F}}\mathbf{f}(x) && \text{by Corollary I.1 page 195 (3)} \\
&= \mathbf{D}^{-j}e^{-i\omega n}\tilde{\mathbf{f}}(\omega) \\
&= 2^{-j/2}e^{-i2^{-j}\omega n}\tilde{\mathbf{f}}(2^{-j}\omega) && \text{by Proposition I.2 page 189}
\end{aligned}$$

⇒

**Proposition I.13.** Let  $\mathbf{T}$  be the translation operator (Definition I.3 page 188). Let  $\tilde{\mathbf{f}}(\omega) \triangleq \tilde{\mathbf{F}}\mathbf{f}(x)$  be the FOURIER TRANSFORM (Definition 4.2 page 16) of a function  $\mathbf{f} \in \mathcal{L}_{\mathbb{R}}^2$ . Let  $\check{\mathbf{a}}(\omega)$  be the DTFT (Definition 6.1 page 41) of a sequence  $(a_n)_{n \in \mathbb{Z}} \in \ell_{\mathbb{R}}^2$  (Definition 5.2 page 27).

P R O P	$\tilde{\mathbf{F}} \sum_{n \in \mathbb{Z}} a_n \mathbf{T}^n \phi(x) = \check{\mathbf{a}}(\omega) \tilde{\phi}(\omega) \quad \forall (a_n) \in \ell_{\mathbb{R}}^2, \phi(x) \in \mathcal{L}_{\mathbb{R}}^2$
------------------	---

PROOF:

$$\begin{aligned}
\tilde{\mathbf{F}} \sum_{n \in \mathbb{Z}} a_n \mathbf{T}^n \phi(x) &= \sum_{n \in \mathbb{Z}} a_n \tilde{\mathbf{F}}\mathbf{T}^n \phi(x) \\
&= \sum_{n \in \mathbb{Z}} a_n e^{-i\omega n} \tilde{\mathbf{F}}\phi(x) && \text{by Corollary I.1 page 195} \\
&= \left[ \sum_{n \in \mathbb{Z}} a_n e^{-i\omega n} \right] \tilde{\phi}(\omega) && \text{by definition of } \tilde{\phi}(\omega) \\
&= \check{\mathbf{a}}(\omega) \tilde{\phi}(\omega) && \text{by definition of DTFT (Definition 6.1 page 41)}
\end{aligned}$$

⇒

**Definition I.4.** Let  $L^2_{(\mathbb{R}, \mathcal{B}, \mu)}$  be the SPACE OF LEBESGUE SQUARE-INTEGRABLE FUNCTIONS (Definition B.1 page 69). Let  $\ell^2_{\mathbb{R}}$  be the SPACE OF ALL ABSOLUTELY SQUARE SUMMABLE SEQUENCES OVER  $\mathbb{R}$  (Definition B.1 page 69).

**DEF**  $S$  is the **sampling operator** in  $\ell^2_{\mathbb{R}}$  if  $[\mathbf{S}f(x)](n) \triangleq f\left(\frac{2\pi}{\tau}n\right) \quad \forall f \in L^2_{(\mathbb{R}, \mathcal{B}, \mu)}, \tau \in \mathbb{R}^+$

**Theorem I.2** (Poisson Summation Formula—PSF).<sup>6</sup> Let  $\tilde{f}(\omega)$  be the FOURIER TRANSFORM (Definition 4.2 page 16) of a function  $f(x) \in L^2_{\mathbb{R}}$ . Let  $S$  be the SAMPLING OPERATOR (Definition I.4 page 196).

**THM**

$$\underbrace{\sum_{n \in \mathbb{Z}} T_{\tau}^n f(x)}_{\text{summation in "time"}} = \underbrace{\sum_{n \in \mathbb{Z}} f(x + n\tau)}_{\text{operator notation}} = \underbrace{\sqrt{\frac{2\pi}{\tau}} \hat{\mathbf{F}}^{-1} \mathbf{S} \tilde{\mathbf{F}}[f(x)]}_{\text{summation in "frequency"}} = \frac{\sqrt{2\pi}}{\tau} \sum_{n \in \mathbb{Z}} \tilde{f}\left(\frac{2\pi}{\tau}n\right) e^{i\frac{2\pi}{\tau}nx}$$

✎ PROOF:

1. lemma: If  $h(x) \triangleq \sum_{n \in \mathbb{Z}} f(x + n\tau)$  then  $h \equiv \hat{\mathbf{F}}^{-1} \hat{\mathbf{F}}h$ . Proof:

Note that  $h(x)$  is *periodic* with period  $\tau$ . Because  $h$  is periodic, it is in the domain of  $\hat{\mathbf{F}}$  and thus  $h \equiv \hat{\mathbf{F}}^{-1} \hat{\mathbf{F}}h$ .

2. Proof of PSF (this theorem—Theorem I.2):

$$\begin{aligned} \sum_{n \in \mathbb{Z}} f(x + n\tau) &= \hat{\mathbf{F}}^{-1} \hat{\mathbf{F}} \sum_{n \in \mathbb{Z}} f(x + n\tau) && \text{by (1) lemma page 196} \\ &= \hat{\mathbf{F}}^{-1} \left[ \frac{1}{\sqrt{\tau}} \int_0^{\tau} \left( \sum_{n \in \mathbb{Z}} f(x + n\tau) \right) e^{-i\frac{2\pi}{\tau}kx} dx \right] && \text{by definition of } \hat{\mathbf{F}} \quad (\text{Definition 7.1 page 51}) \\ &\quad \underbrace{\hspace{10em}}_{\hat{\mathbf{F}}[\sum_{n \in \mathbb{Z}} f(x + n\tau)]} \\ &= \hat{\mathbf{F}}^{-1} \left[ \frac{1}{\sqrt{\tau}} \sum_{n \in \mathbb{Z}} \int_0^{\tau} f(x + n\tau) e^{-i\frac{2\pi}{\tau}kx} dx \right] \\ &= \hat{\mathbf{F}}^{-1} \left[ \frac{1}{\sqrt{\tau}} \sum_{n \in \mathbb{Z}} \int_{u=n\tau}^{u=(n+1)\tau} f(u) e^{-i\frac{2\pi}{\tau}k(u-n\tau)} du \right] && \text{where } u \triangleq x + n\tau \implies x = u - n\tau \\ &= \hat{\mathbf{F}}^{-1} \left[ \frac{1}{\sqrt{\tau}} \sum_{n \in \mathbb{Z}} \underbrace{e^{i2\pi kn}}_{=1} \int_{u=n\tau}^{u=(n+1)\tau} f(u) e^{-i\frac{2\pi}{\tau}ku} du \right] \\ &= \sqrt{\frac{2\pi}{\tau}} \hat{\mathbf{F}}^{-1} \left[ \frac{1}{\sqrt{2\pi}} \int_{u \in \mathbb{R}} f(u) e^{-i\left(\frac{2\pi}{\tau}k\right)u} du \right] && \text{by evaluation of } \hat{\mathbf{F}}^{-1} \quad (\text{Theorem 7.1 page 52}) \\ &\quad \underbrace{\hspace{10em}}_{[\tilde{\mathbf{F}}f]\left(\frac{2\pi}{\tau}k\right)} \\ &= \sqrt{\frac{2\pi}{\tau}} \hat{\mathbf{F}}^{-1} \left[ [\tilde{\mathbf{F}}f(x)]\left(\frac{2\pi}{\tau}k\right) \right] && \text{by definition of } \tilde{\mathbf{F}} \quad (\text{Definition 4.2 page 16}) \\ &= \sqrt{\frac{2\pi}{\tau}} \hat{\mathbf{F}}^{-1} \mathbf{S} \tilde{\mathbf{F}}f && \text{by definition of } S \quad (\text{Definition I.4 page 196}) \\ &= \frac{\sqrt{2\pi}}{\tau} \sum_{n \in \mathbb{Z}} \tilde{f}\left(\frac{2\pi}{\tau}n\right) e^{i\frac{2\pi}{\tau}nx} && \text{by evaluation of } \hat{\mathbf{F}}^{-1} \quad (\text{Theorem 7.1 page 52}) \end{aligned}$$

⇒

<sup>6</sup> Andrews et al. (2001) page 624, Knapp (2005b) page 389, Lasser (1996) page 254, Rudin (1987) pages 194–195, Folland (1992) page 337

**Theorem I.3** (Inverse Poisson Summation Formula—IPSF).<sup>7</sup>

Let  $\tilde{f}(\omega)$  be the FOURIER TRANSFORM (Definition 4.2 page 16) of a function  $f(x) \in L^2_{\mathbb{R}}$ .

$$\underbrace{\sum_{n \in \mathbb{Z}} T_{2\pi/\tau}^n \tilde{f}(\omega) \triangleq \sum_{n \in \mathbb{Z}} \tilde{f}\left(\omega - \frac{2\pi}{\tau}n\right)}_{\text{summation in "frequency"}} = \underbrace{\frac{\tau}{\sqrt{2\pi}} \sum_{n \in \mathbb{Z}} f(n\tau) e^{-i\omega n\tau}}_{\text{summation in "time"}}$$

PROOF:

1. lemma: If  $h(\omega) \triangleq \sum_{n \in \mathbb{Z}} \tilde{f}\left(\omega + \frac{2\pi}{\tau}n\right)$ , then  $h \equiv \hat{F}^{-1} \hat{F} h$ . Proof:

Note that  $h(\omega)$  is periodic with period  $2\pi/\tau$ :

$$h\left(\omega + \frac{2\pi}{\tau}\right) \triangleq \sum_{n \in \mathbb{Z}} \tilde{f}\left(\omega + \frac{2\pi}{\tau} + \frac{2\pi}{\tau}n\right) = \sum_{n \in \mathbb{Z}} \tilde{f}\left(\omega + (n+1)\frac{2\pi}{\tau}\right) = \sum_{n \in \mathbb{Z}} \tilde{f}\left(\omega + \frac{2\pi}{\tau}n\right) \triangleq h(\omega)$$

Because  $h$  is periodic, it is in the domain of  $\hat{F}$  and is equivalent to  $\hat{F}^{-1} \hat{F} h$ .

2. Proof of IPSF (this theorem—Theorem I.3):

$$\begin{aligned} & \sum_{n \in \mathbb{Z}} \tilde{f}\left(\omega + \frac{2\pi}{\tau}n\right) \\ &= \hat{F}^{-1} \hat{F} \sum_{n \in \mathbb{Z}} \tilde{f}\left(\omega + \frac{2\pi}{\tau}n\right) && \text{by (1) lemma page 197} \\ &= \hat{F}^{-1} \left[ \underbrace{\sqrt{\frac{\tau}{2\pi}} \int_0^{\frac{2\pi}{\tau}} \sum_{n \in \mathbb{Z}} \tilde{f}\left(\omega + \frac{2\pi}{\tau}n\right) e^{-i\omega \frac{2\pi}{\tau}k} d\omega}_{\hat{F}\left[\sum_{n \in \mathbb{Z}} \tilde{f}\left(\omega + \frac{2\pi}{\tau}n\right)\right]} \right] && \text{by definition of } \hat{F} \quad (\text{Definition 7.1 page 51}) \\ &= \hat{F}^{-1} \left[ \sqrt{\frac{\tau}{2\pi}} \sum_{n \in \mathbb{Z}} \int_0^{\frac{2\pi}{\tau}} \tilde{f}\left(\omega + \frac{2\pi}{\tau}n\right) e^{-i\omega T k} d\omega \right] \\ &= \hat{F}^{-1} \left[ \sqrt{\frac{\tau}{2\pi}} \sum_{n \in \mathbb{Z}} \int_{u=\frac{2\pi}{\tau}n}^{u=\frac{2\pi}{\tau}(n+1)} \tilde{f}(u) e^{-i\left(u - \frac{2\pi}{\tau}n\right)T k} du \right] && \text{where } u \triangleq \omega + \frac{2\pi}{\tau}n \implies \omega = u - \frac{2\pi}{\tau}n \\ &= \hat{F}^{-1} \left[ \sqrt{\frac{\tau}{2\pi}} \sum_{n \in \mathbb{Z}} e^{i2\pi n k} \int_{\frac{2\pi}{\tau}n}^{\frac{2\pi}{\tau}(n+1)} \tilde{f}(u) e^{-iu\tau k} du \right] \\ &= \hat{F}^{-1} \left[ \sqrt{\frac{\tau}{2\pi}} \int_{\mathbb{R}} \tilde{f}(u) e^{-iu\tau k} du \right] \\ &= \sqrt{\tau} \hat{F}^{-1} \left[ \underbrace{\frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} \tilde{f}(u) e^{iu(-\tau k)} du}_{[\hat{F}^{-1}\tilde{f}](-k\tau)} \right] \\ &= \sqrt{\tau} \hat{F}^{-1} [[\hat{F}^{-1}\tilde{f}](-k\tau)] && \text{by value of } \hat{F}^{-1} \quad (\text{Theorem 4.1 page 17}) \\ &= \sqrt{\tau} \hat{F}^{-1} \mathbf{S} \hat{F}^{-1} \tilde{f} && \text{by definition of } \mathbf{S} \quad (\text{Definition 1.4 page 196}) \\ &= \sqrt{\tau} \hat{F}^{-1} \mathbf{S} f(x) && \text{by definition of } \tilde{F} \quad (\text{Definition 4.2 page 16}) \\ &= \sqrt{\tau} \hat{F}^{-1} f(-k\tau) && \text{by definition of } \mathbf{S} \quad (\text{Definition 1.4 page 196}) \\ &= \sqrt{\tau} \frac{1}{\sqrt{\frac{2\pi}{\tau}}} \sum_{k \in \mathbb{Z}} f(-k\tau) e^{i2\pi \frac{1}{\tau} k \omega} && \text{by definition of } \hat{F}^{-1} \quad (\text{Theorem 7.1 page 52}) \end{aligned}$$

<sup>7</sup> Gauss (1900) page 88

$$= \frac{\tau}{\sqrt{\frac{2\pi}{\tau}}} \sum_{k \in \mathbb{Z}} f(-k\tau) e^{ik\tau\omega}$$

$$= \frac{\tau}{\sqrt{2\pi}} \sum_{m \in \mathbb{Z}} f(m\tau) e^{-i\omega m\tau}$$

by definition of  $\hat{\mathbf{F}}^{-1}$  (Theorem 7.1 page 52)let  $m \triangleq -k$ 

⇒

**Remark I.2.** The left hand side of the *Poisson Summation Formula* (Theorem I.2 page 196) is very similar to the *Zak Transform Z*:<sup>8</sup>

$$(\mathbf{Z}f)(t, \omega) \triangleq \sum_{n \in \mathbb{Z}} f(x + n\tau) e^{i2\pi n\omega}$$

**Remark I.3.** A generalization of the *Poisson Summation Formula* (Theorem I.2 page 196) is the **Selberg Trace Formula**.<sup>9</sup>

## I.7 Examples

**Example I.2** (linear functions).<sup>10</sup> Let  $\mathbf{T}$  be the *translation operator* (Definition I.3 page 188). Let  $\mathcal{L}(\mathbb{C}, \mathbb{C})$  be the set of all *linear* functions in  $L^2_{\mathbb{R}}$ .

- |                |   |
|----------------|---|
| <b>E<br/>X</b> | 1. $\{x, \mathbf{T}x\}$ is a <i>basis</i> for $\mathcal{L}(\mathbb{C}, \mathbb{C})$ and     |
|                | 2. $f(x) = f(1)x - f(0)\mathbf{T}x \quad \forall f \in \mathcal{L}(\mathbb{C}, \mathbb{C})$ |

PROOF: By left hypothesis,  $f$  is *linear*; so let  $f(x) \triangleq ax + b$

$$\begin{aligned} f(1)x - f(0)\mathbf{T}x &= f(1)x - f(0)(x - 1) \\ &= (ax + b)|_{x=1} x - (ax + b)|_{x=0} (x - 1) \\ &= (a + b)x - b(x - 1) \\ &= ax + bx - bx + b \\ &= ax + b \\ &= f(x) \end{aligned}$$

by Definition I.3 page 188

by left hypothesis and definition of  $f$ by left hypothesis and definition of  $f$ 

⇒

**Example I.3** (Cardinal Series). Let  $\mathbf{T}$  be the *translation operator* (Definition I.3 page 188). The *Paley-Wiener* class of functions  $\mathbf{PW}_{\sigma}^2$  are those functions which are “*bandlimited*” with respect to their Fourier transform (Definition 4.2 page 16). The cardinal series forms an orthogonal basis for such a space. The *Fourier coefficients* (Definition G.11 page 160) for a projection of a function  $f$  onto the Cardinal series basis elements is particularly simple—these coefficients are samples of  $f(x)$  taken at regular intervals. In fact, one could represent the coefficients using inner product notation with the *Dirac delta distribution*  $\delta$  as follows:

$$\langle f(x) | \mathbf{T}^n \delta(x) \rangle \triangleq \int_{\mathbb{R}} f(x) \delta(x - n) dx \triangleq f(n)$$

- |                |   |
|----------------|---|
| <b>E<br/>X</b> | 1. $\left\{ \mathbf{T}^n \frac{\sin(\pi x)}{\pi x} \right\}_{n \in \mathbb{N}}$ is a <i>basis</i> for $\mathbf{PW}_{\sigma}^2$ and  |
|                | 2. $f(x) = \underbrace{\sum_{n=1}^{\infty} f(n) \mathbf{T}^n \frac{\sin(\pi x)}{\pi x}}_{\text{Cardinal series}} \quad \forall f \in \mathbf{PW}_{\sigma}^2, \sigma \leq \frac{1}{2}$ |

<sup>8</sup> Janssen (1988) page 24, Zayed (1996) page 482


<sup>9</sup> Lax (2002) page 349, Selberg (1956), Terras (1999)

<sup>10</sup> Higgins (1996) page 2



*Example I.4 (Fourier Series).*

- E X**
1.  $\{\mathbf{D}_n e^{ix} \mid n \in \mathbb{Z}\}$  is a *basis* for  $L(0 : 2\pi)$  and
  2.  $f(x) = \frac{1}{\sqrt{2\pi}} \sum_{n \in \mathbb{Z}} \alpha_n \mathbf{D}_n e^{ix} \quad \forall x \in (0 : 2\pi), f \in L(0 : 2\pi)$  where
  3.  $\alpha_n \triangleq \frac{1}{\sqrt{2\pi}} \int_0^{2\pi} f(x) \mathbf{D}_n e^{-ix} dx \quad \forall f \in L(0 : 2\pi)$

 **PROOF:** See Theorem 7.1 page 52. 

*Example I.5 (Fourier Transform).* <sup>11</sup>

- E X**
1.  $\{\mathbf{D}_\omega e^{ix} \mid \omega \in \mathbb{R}\}$  is a *basis* for  $L^2_{\mathbb{R}}$  and
  2.  $f(x) = \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} \tilde{f}(\omega) \mathbf{D}_x e^{i\omega} d\omega \quad \forall f \in L^2_{\mathbb{R}}$  where
  3.  $\tilde{f}(\omega) \triangleq \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} f(x) \mathbf{D}_\omega e^{-ix} dx \quad \forall f \in L^2_{\mathbb{R}}$

*Example I.6 (Gabor Transform).* <sup>12</sup>

- E X**
1.  $\left\{ \left( \mathbf{T}_\tau e^{-\pi x^2} \right) (\mathbf{D}_\omega e^{ix}) \mid \tau, \omega \in \mathbb{R} \right\}$  is a *basis* for  $L^2_{\mathbb{R}}$  and
  2.  $f(x) = \int_{\mathbb{R}} G(\tau, \omega) \mathbf{D}_x e^{i\omega} d\omega \quad \forall x \in \mathbb{R}, f \in L^2_{\mathbb{R}}$  where
  3.  $G(\tau, \omega) \triangleq \int_{\mathbb{R}} f(x) \left( \mathbf{T}_\tau e^{-\pi x^2} \right) (\mathbf{D}_\omega e^{-ix}) dx \quad \forall x \in \mathbb{R}, f \in L^2_{\mathbb{R}}$

*Example I.7 (wavelets). Let  $\psi(x)$  be a wavelet.*

- E X**
1.  $\{\mathbf{D}^k \mathbf{T}^n \psi(x) \mid k, n \in \mathbb{Z}\}$  is a *basis* for  $L^2_{\mathbb{R}}$  and
  2.  $f(x) = \sum_{k \in \mathbb{Z}} \sum_{n \in \mathbb{Z}} \alpha_{k,n} \mathbf{D}^k \mathbf{T}^n \psi(x) \quad \forall f \in L^2_{\mathbb{R}}$  where
  3.  $\alpha_n \triangleq \int_{\mathbb{R}} f(x) \mathbf{D}^k \mathbf{T}^n \psi^*(x) dx \quad \forall f \in L^2_{\mathbb{R}}$

<sup>11</sup>cross reference: Definition 4.2 page 16

<sup>12</sup> Gabor (1946),  Qian and Chen (1996) (Chapter 3),  Forster and Massopust (2009) page 32 (Definition 1.69)



# APPENDIX J

## CONTINUOUS RANDOM PROCESSES

### J.1 Definitions

**Definition J.1.** <sup>1</sup> Let  $(\Omega, \mathbb{E}, P)$  be a PROBABILITY SPACE.

**DEF** The function  $x : \Omega \rightarrow \mathbb{R}$  is a **random variable**.  
The function  $y : \mathbb{R} \times \Omega \rightarrow \mathbb{R}$  is a **random process**.

The random process  $x(t, \omega)$ , where  $t$  commonly represents time and  $\omega \in \Omega$  is an outcome of an experiment, can take on more specialized forms depending on whether  $t$  and  $\omega$  are fixed or allowed to vary. These forms are illustrated in Figure J.1 page 201<sup>2</sup> and Figure J.2 page 202.

$x(t, \omega)$	fixed $t$	variable $t$
fixed $\omega$	number	time function
variable $\omega$	random variable	random process

Figure J.1: Specialized forms of a random process  $x(t, \omega)$

**Definition J.2.** <sup>3</sup> Let  $x(t)$  and  $y(t)$  be random processes.

**DEF** The **mean**  $\mu_x(t)$  of  $x(t)$  is  $\mu_x(t) \triangleq E[x(t)]$   
The **cross-correlation**  $R_{xy}(t)$  of  $x(t)$  and  $y(t)$  is  $R_{xy}(t, u) \triangleq E[x(t)y^*(u)]$   
The **auto-correlation function**  $R_{xx}(t)$  of  $x(t)$  is  $R_{xx}(t, u) \triangleq E[x(t)x^*(u)]$


**Remark J.1.** <sup>4</sup> The equation  $\int_{u \in \mathbb{R}} R_{xx}(t, u)f(u) du$  is a *Fredholm integral equation of the first kind* and  $R_{xx}(t, u)$  is the *kernel* of the equation.

<sup>1</sup> Papoulis (1991) page 63, Papoulis (1991) page 285

<sup>2</sup> Papoulis (1991) pages 285–286

<sup>3</sup> Papoulis (1984) page 216  $\langle R_{xy}(t_1, t_2) = E\{x(t_1)y^*(t_2)\}$  (9-35)),

<sup>4</sup> Fredholm (1900), Fredholm (1903) page 365, Michel and Herget (1993) page 97, Keener (1988) page 101

Figure J.2: Example of a random process  $x(t, \omega)$ 

## J.2 Properties

**Theorem J.1.** Let  $x(t)$  and  $y(t)$  be random processes with cross-correlation  $R_{xy}(t, u)$  and let  $R_{xx}(t, u)$  be the auto-correlation of  $x(t)$ .

<b>T H M</b>	$R_{xx}(t, u) = R_{xx}^*(u, t)$ (CONJUGATE SYMMETRIC)
	$R_{xy}(t, u) = R_{yx}^*(u, t)$

PROOF:

$$\begin{aligned}
 R_{xx}(t, u) &\triangleq E[x(t)x^*(u)] &= E[x^*(u)x(t)] &= (E[x(u)x^*(t)])^* &\triangleq R_{xx}^*(u, t) \\
 R_{xy}(t, u) &\triangleq E[x(t)y^*(u)] &= E[y^*(u)x(t)] &= (E[y(u)x^*(t)])^* &\triangleq R_{yx}^*(u, t)
 \end{aligned}$$

# APPENDIX K

## RANDOM SEQUENCES



*“We are quite in danger of sending highly trained and highly intelligent young men out into the world with tables of erroneous numbers under their arms, and with a dense fog in the place where their brains ought to be. In this century, of course, they will be working on guided missiles and advising the medical profession on the control of disease, and there is no limit to the extent to which they could impede every sort of national effort.”*

Ronald A. Fisher, (1890–1962), Statistician, at a lecture in 1958 at Michigan State University <sup>1</sup>

## K.1 Definitions

### Definition K.1.

**DEF** A **random sequence** is a SEQUENCE over a PROBABILITY SPACE (Definition ?? page ??).

### Definition K.2. <sup>2</sup> Let $x(n)$ and $y(n)$ be RANDOM SEQUENCES.

<b>DEF</b>	The <b>mean</b>	$\mu_X(n)$	of $x(n)$ is	$\mu_X(n) \triangleq E[x(n)]$
	The <b>variance</b>	$\sigma_X^2(n)$	of $x(n)$ is	$\sigma_X^2(n) \triangleq E\left([x(n) - \mu_X(n)]^2\right)$
	The <b>cross-correlation</b>	$R_{xy}(n, m)$	of $x(n)$ and $y(n)$ is	$R_{xy}(n, m) \triangleq E[x(n+m)y^*(n)]$
	The <b>auto-correlation</b>	$R_{xx}(n, m)$	of $x(n)$ is	$R_{xx}(n, m) \triangleq R_{xy}(n, m) _{y=x}$

<sup>1</sup>quote: [Yates and Mather (1963) page 107. image: <http://www.genetics.org/content/154/4/1419>

<sup>2</sup> [Papoulis (1984) page 263  $\langle R_{xy}(m) = E\{x(m)y^*(0)\} \rangle$ , [Wilks (1963) page 77 §3.4 “Moments of two-dimensional random variables”, [Cadzow (1987) page 341  $\langle r_{xy}(m) = E[x(m)y^*(0)] \rangle$ , [MatLab (2018b)  $\langle R_{xy}(m) = E\{x_{n+m}y_n^*\} \rangle$ , [MatLab (2018a)  $\langle R_{xy}(m) = E\{x_{n+m}y_n^*\} \rangle$

## K.2 Properties

### Theorem K.1.

T H M	$R_{xx}(n, m) = R_{xx}^*(n + m, -m)$
	$R_{xy}(n, m) = R_{yx}^*(n + m, -m)$

 PROOF:

$R_{xy}(n, m) \triangleq E[x(n+m)y^*(n)]$	by definition of $R_{xy}(n, m)$	(Definition K.2 page 203)
$= E[y^*(n)x(n+m)]$	by <i>commutative</i> property of $(\mathbb{C}, +, \cdot, 0, 1)$	(Definition A.5 page 68)
$= (E[y(n)x^*(n+m)])^*$	by <i>distributive</i> property of $*$ -algebras	(Definition E.3 page 114)
$= (E[y(n+m-m)x^*(n+m)])^*$	by <i>additive identity</i> property of $(\mathbb{R}, +, \cdot, 0, 1)$	(Definition A.5 page 68)
$\triangleq R_{yx}^*(n+m, -m)$	by definition of $R_{yx}(n, m)$	(Definition K.2 page 203)
$R_{xx}(n, m) = R_{xy}(n, m) _{y=x}$	by $y = x$ constraint	
$= R_{xy}^*(n+m, -m) _{y=x}$	by previous result	
$= R_{xx}^*(n+m, -m)$	by $y = x$ constraint	



## K.3 Wide Sense Stationary processes



**Definition K.3.** Let  $x(n)$  be a RANDOM SEQUENCE with MEAN  $\mu_X(n)$  and VARIANCE  $\sigma_X^2(n)$  (Definition K.2 page 203).

D E F	$x(n)$ is <b>wide sense stationary (WSS)</b> if	
	1. $\mu_X(n)$ is CONSTANT with respect to $n$	(STATIONARY IN THE 1ST MOMENT) and
	2. $\sigma_X^2(n)$ is CONSTANT with respect to $n$	(STATIONARY IN THE 2ND MOMENT)

**Definition K.4.**<sup>3</sup> Let  $x(n)$  be a RANDOM SEQUENCE with statistics  $\mu_X(n)$ ,  $\sigma_X^2(n)$ ,  $R_{xx}(n, m)$ , and  $R_{xy}(n, m)$  (Definition K.2 page 203).

D E F	$\{ x \text{ and } y \text{ are WIDE SENSE STATIONARY} \} \implies$			
	(1). The <b>mean</b>	$\mu_X$	of $x(n)$ is	$\mu_X \triangleq E[x(0)]$
	(2). The <b>variance</b>	$\sigma_X^2$	of $x(n)$ is	$\sigma_X^2 \triangleq E\left([x(0) - \mu_X]^2\right)$
	(4). The <b>cross-correlation</b>	$R_{xy}(m)$	of $x(n)$ and $y(n)$ is	$R_{xy}(m) \triangleq E[x(m)y^*(0)]$
	(3). The <b>auto-correlation</b>	$R_{xx}(m)$	of $x(n)$ is	$R_{xx}(m) \triangleq R_{xy}(m) _{y=x}$

**Remark K.1.** The  $R_{xy}(n, m)$  of Definition K.2 (page 203) and the  $R_{xy}(m)$  of Definition K.4 (page 204) (etc.) are examples of *function overload*—that is, functions that use the same mnemonic but are distinguished by different domains. Perhaps a more common example of function overload is the “+” mnemonic. Traditionally it is used with domain of the natural numbers  $\mathbb{N}$  as in  $3 + 2$ . Later it was extended for domain real numbers  $\mathbb{R}$  as in  $\sqrt{3} + \sqrt{2}$ , or even complex numbers  $\mathbb{C}$  as in

<sup>3</sup>  Papoulis (1984) page 263  $\langle R_{xy}(\tau) = E\{x(t+\tau)y^*(t)\} \rangle$ ,  Cadzow (1987) page 341  $\langle r_{xy}(n) = E[x(k+n)y^*(k)] \rangle$  (10.41)

$(\sqrt{3} + i\sqrt{2}) + (e + i\pi)$ . And it was even more dramatically extended for use with domain  $\mathbb{R}^N \times \mathbb{R}^M$  in “linear algebra” as in

$$\begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} + \begin{bmatrix} 5 & 6 \\ 7 & 8 \end{bmatrix} = \begin{bmatrix} 6 & 8 \\ 10 & 12 \end{bmatrix}$$

**Remark K.2.** <sup>4</sup> The definition for  $R_{xy}(m)$  can be defined with the conjugate  $*$  on either  $x$  or  $y$ , or on neither or both; and moreover  $x$  may either lead or lag  $y$ . In total, there are  $2 \times 2 \times 2 = 8$  different ways to define  $R_{xy}(m)$ .<sup>5</sup> and  $R_{xx}(m)$  involve complex numbers. This may seem curious when typical ADCs provide real-valued sequences. Note however that complex-valued sequences often come up in signal processing due to some common system architectures:


1. The presence of an *FFT* operator in the signal processing path
2. The *complex envelope*  $x_l(t)$  of a modulated *narrowband* communications signal  $x(t)$ .
3. Communications channel processing involving phase discrimination (e.g. PSK and QAM).

In the case of a narrowband signal  $x(t)$  modulated by a sinusoid at center frequency  $f_c$ , we have three canonical forms. These can be shown to be equivalent:

$$\begin{aligned} x(t) &\triangleq \underbrace{a(t)\cos[2\pi f_c t + \phi(t)]}_{\text{amplitude-phase form}} && \text{amplitude and phase form} \\ &= \underbrace{a(t)\cos[\phi(t)]}_{p(t)} \cos[2\pi f_c t] - \underbrace{a(t)\sin[\phi(t)]}_{q(t)} \sin[2\pi f_c t] && \text{by double angle formulas (Theorem C.9 page 85)} \\ &= \underbrace{p(t)\cos[2\pi f_c t] - q(t)\sin[2\pi f_c t]}_{\text{quadrature form}} && \text{quadrature form} \\ &= \mathbf{R}_e([p(t) + iq(t)][\cos(2\pi f_c t) + i\sin(2\pi f_c t)]) && \text{by definitions of } \mathbf{R}_e \\ &= \underbrace{\mathbf{R}_e[x_l(t)e^{i2\pi f_c t}]}_{\text{complex envelope form}} && \text{by Euler's identity (Theorem C.5 page 80)} \end{aligned}$$

Note that in these equivalent forms, the *complex envelope*  $x_l(t)$  is conveniently represented as a *complex-valued* function in terms of the *quadrature component*  $p(t)$  and the *inphase component*  $q(t)$  such that  $x_l(t) = p(t) + iq(t)$ .

In practice (with real hardware), you will likely first have access to the quadrature components  $p(t)$  and  $q(t)$ . Take for example the *Analog Devices ADL5387 Quadrature Demodulator* and evaluation board, as illustrated to the right.<sup>6</sup> Note that *quadrature component*  $p(t)$  is available at connector “Q OUTPUT” and *inphase component*  $q(t)$  is available at connector “I OUTPUT”.



$R_{xx}(n, m)$ , and  $R_{xy}$  the CROSS-CORRELATION of  $x$  and  $y$ .

PRP

$$\left\{ \begin{array}{l} x \text{ and } y \text{ are} \\ \text{WIDE SENSE STATIONARY} \\ \text{(WSS) (Definition ?? page ??)} \end{array} \right\} \Rightarrow \left\{ \begin{array}{ll} R_{xx}(n, m) = R_{xx}(m) & \forall n \in \mathbb{Z} \\ R_{xy}(n, m) = R_{xy}(m) & \forall n \in \mathbb{Z} \end{array} \right\}$$

(Definition K.2 page 203) (Definition K.4 page 204)

PROOF:

$$\begin{aligned} R_{xy}(n, m) &\triangleq E[x[n+m]y^*[n]] && \text{by definition of } R_{xy}(n, m) && \text{(Definition K.2 page 203)} \\ &= E[x[n-n+m]y^*[n-n]] && \text{by wide sense stationary hypothesis} \\ &= E[x[m]y^*[0]] \\ &\triangleq R_{xy}(m) && \text{by definition of } R_{xy}(m) && \text{(Definition K.4 page 204)} \\ R_{xx}(n, m) &= R_{xy}(n, m)|_{y=x} \\ &= R_{xy}(m)|_{y=x} && \text{by previous result} \\ &= R_{xx}(m) \end{aligned}$$




Figure K.1: auto-correlation  $R_{xx}(m)$

**Corollary K.1.** Let  $x(n)$  be a RANDOM SEQUENCE with AUTO-CORRELATION  $R_{xx}(n, m)$ ,  $y(n)$  a RANDOM SEQUENCE with AUTO-CORRELATION  $R_{yy}(n, m)$ , and  $R_{xy}(n, m)$  the CROSS-CORRELATION of  $x$  and  $y$ . Let  $S$  be a SYSTEM with input  $x(n)$  and output  $y(n)$ .

COR

$$\left\{ \begin{array}{l} \text{(A). } x \text{ is WSS} \\ \text{(B). } y \text{ is WSS} \\ \text{(C). } S \text{ is LTI} \end{array} \right\} \Rightarrow \left\{ \begin{array}{ll} (1). R_{xy}(m) = R_{yx}^*(-m) & \text{and} \\ (2). R_{xx}(m) = R_{xx}^*(-m) & \text{(CONJUGATE SYMMETRIC) and} \\ (3). R_e R_{xx}(m) = R_e R_{xx}(-m) & \text{(SYMMETRIC) and} \\ (4). I_m R_{xx}(m) = -I_m R_{xx}(-m) & \text{(ANTI-SYMMETRIC) and} \\ (5). |R_{xx}(m)| = |R_{xx}(-m)| & \text{(SYMMETRIC) and} \\ (6). \angle R_{xx}(m) = -\angle R_{xx}(-m) & \text{(ANTI-SYMMETRIC)} \end{array} \right\}$$

PROOF:

$$\begin{aligned} R_{xy}(m) &= R_{xy}(n, m) && \text{by Proposition K.1 page 205} && \text{and hypotheses (A),(B)} \\ &= R_{yx}^*(n+m, -m) && \text{by Theorem K.1 page 204} && \text{and hypothesis (B)} \\ &= R_{yx}^*(-m) && \text{by Proposition K.1 page 205} && \text{and hypothesis (A)} \\ R_{xx}(m) &= R_{xx}(n, m) && \text{by Proposition K.1 page 205} && \text{and hypothesis (A)} \\ &= R_{xx}^*(n+m, -m) && \text{by Theorem K.1 page 204} && \text{and hypothesis (B)} \\ &= R_{xx}^*(-m) && \text{by Proposition K.1 page 205} && \text{and hypothesis (A)} \end{aligned}$$



## K.4 Spectral density

**Definition K.5.** Let  $x(n)$  and  $y(n)$  be WIDE SENSE STATIONARY RANDOM SEQUENCES with auto-correlation  $R_{xx}(m)$  and cross-correlation  $R_{xy}(m)$ . Let  $\mathbf{Z}$  be the Z-TRANSFORM OPERATOR (Definition 5.4 page 28).

**DEF** The **z-domain cross spectral density (CSD)**  $\check{S}_{xy}(z)$  of  $x$  and  $y$  is  

$$\check{S}_{xy}(z) \triangleq \mathbf{Z}R_{xy}(m) \triangleq \sum_{m \in \mathbb{Z}} R_{xy}(m)z^{-m}$$
  
 The **z-domain power spectral density (PSD)**  $\check{S}_{xx}(z)$  of  $x$  is  $\check{S}_{xx}(z) \triangleq \check{S}_{xy}(z)|_{y(n)=x(n)}$

**Definition K.6.** Let  $x(n)$  and  $y(n)$  be WIDE SENSE STATIONARY RANDOM SEQUENCES with auto-correlation  $R_{xx}(m)$  and cross-correlation  $R_{xy}(m)$ . Let  $\check{\mathbf{F}}$  be the DISCRETE TIME FOURIER TRANSFORM (DTFT) operator (Definition 6.1 page 41).

**DEF** The **auto-spectral density**  $\check{S}_{xx}(z)$  of  $x$  is  $\check{S}_{xx}(z) \triangleq \check{\mathbf{F}}R_{xx}(m) \triangleq \sum_{m \in \mathbb{Z}} R_{xx}(m)e^{-i\omega m}$   
 The **cross spectral density (CSD)**  $\check{S}_{xy}(z)$  of  $x$  and  $y$  is  $\check{S}_{xy}(z) \triangleq \check{\mathbf{F}}R_{xy}(m) \triangleq \sum_{m \in \mathbb{Z}} R_{xy}(m)e^{-i\omega m}$   
 The **auto-spectral density** is also called **power spectral density (PSD)**.

**Theorem K.2.** Let  $\mathbf{S}$  be a system with IMPULSE RESPONSE  $h(n)$ , INPUT  $x(n)$ , and OUTPUT  $y(n)$ .

**THM**  $\{ x \text{ and } y \text{ are WIDE SENSE STATIONARY} \} \Rightarrow \left\{ \begin{array}{l} (1). \check{S}_{xx}(z) = \check{S}_{xx}^*\left(\frac{1}{z^*}\right) \text{ and} \\ (2). \check{S}_{yx}(z) = \check{S}_{xy}^*\left(\frac{1}{z^*}\right) \end{array} \right\}$

PROOF:

$\check{S}_{yx}(z) \triangleq \mathbf{Z}R_{yx}(m)$	by definition of $\check{S}_{xy}(z)$	(Definition K.6 page 207)
$\triangleq \sum_{m \in \mathbb{Z}} R_{yx}(m)z^{-m}$	by definition of $\mathbf{Z}$	(Definition 5.4 page 28)
$\triangleq \sum_{m \in \mathbb{Z}} R_{xy}^*(-m)z^{-m}$	by Corollary K.1 page 206	
$= \left[ \sum_{m \in \mathbb{Z}} R_{xy}(-m)(z^*)^{-m} \right]^*$	by antiautomorphic property of $*$ -algebras	(Definition E.3 page 114)
$= \left[ \sum_{p \in \mathbb{Z}} R_{xy}(p)(z^*)^p \right]^*$	where $p \triangleq -m$	$\Rightarrow m = -p$
$= \left[ \sum_{p \in \mathbb{Z}} R_{xy}(p)(z^*)^p \right]^*$	by absolutely summable property	(Definition 5.2 page 27)
$= \left[ \sum_{p \in \mathbb{Z}} R_{xy}(p)\left(\frac{1}{z^*}\right)^{-p} \right]^*$		
$= \check{S}_{xy}^*\left(\frac{1}{z^*}\right)$	by definition of $\mathbf{Z}$	(Definition 5.4 page 28)
$\check{S}_{xx}(z) = \check{S}_{xy}(z) _{y=x}$		
$= \check{S}_{yx}^*(z) _{y=x}$		
$= \check{S}_{xy}^*\left(\frac{1}{z^*}\right) _{y=x}$	by (2)—previous result	

$$= \check{S}_{xx}^* \left( \frac{1}{z^*} \right)$$



**Corollary K.2.** Let  $S$  be a system with IMPULSE RESPONSE  $h(n)$ , INPUT  $x(n)$ , and OUTPUT  $y(n)$ .

<b>COR</b>	$\left\{ \begin{array}{l} \text{(A). } h \text{ is LTI and} \\ \text{(B). } x \text{ and } y \text{ are WSS} \end{array} \right\} \implies \left\{ \begin{array}{l} \text{(1). } \check{S}_{xy}^*(\omega) = \check{S}_{yx}(\omega) \quad (\text{CONJUGATE-SYMMETRIC}) \quad \text{and} \\ \text{(2). } \check{S}_{xx}^*(\omega) = \check{S}_{xx}(\omega) \quad (\text{CONJUGATE SYMMETRIC}) \quad \text{and} \\ \text{(3). } \check{S}_{xx}(\omega) \in \mathbb{R} \quad (\text{REAL-VALUED}) \end{array} \right\}$
------------	---

PROOF:

$\begin{aligned} \check{S}_{xy}^*(\omega) &= \check{S}_{xy}^*(z) \Big _{z=e^{i\omega}} \\ &= \check{S}_{yx}^* \left( \frac{1}{z^*} \right) \Big _{z=e^{i\omega}} \\ &= \check{S}_{yx} \left( \frac{1}{z^*} \right) \Big _{z=e^{i\omega}} \\ &= \check{S}_{yx} \left( \frac{1}{e^{i\omega^*}} \right) \\ &= \check{S}_{yx}(e^{i\omega}) \\ &= \check{S}_{yx}(\omega) \end{aligned}$	<p>by definition of <i>DTFT</i></p> <p>by Theorem K.2 page 207</p> <p>by <i>involutory</i> property of <b>*-algebras</b></p>	<p>(Definition 6.1 page 41)</p>
$\begin{aligned} \check{S}_{xx}^*(\omega) &= \check{S}_{xx}^*(z) \Big _{z=e^{i\omega}} \\ &= \check{S}_{xx}^* \left( \frac{1}{z^*} \right) \Big _{z=e^{i\omega}} \\ &= \check{S}_{xx} \left( \frac{1}{z^*} \right) \Big _{z=e^{i\omega}} \\ &= \check{S}_{xx} \left( \frac{1}{e^{i\omega^*}} \right) \\ &= \check{S}_{xx}(e^{i\omega}) \\ &= \check{S}_{xx}(\omega) \end{aligned}$	<p>by definition of <i>DTFT</i></p> <p>by definition of <i>DTFT</i></p> <p>by Theorem K.2 page 207</p> <p>by <i>involutory</i> property of <b>*-algebras</b></p>	<p>(Definition 6.1 page 41)</p> <p>(Definition 6.1 page 41)</p>
$\begin{aligned} \check{S}_{xx}^*(\omega) &= \check{S}_{xy}^*(\omega) \Big _{y=x} \\ &= \check{S}_{yx}(\omega) \Big _{y=x} \\ &= \check{S}_{xx}(\omega) \end{aligned}$	<p>by definition of <i>DTFT</i></p> <p>by previous result</p>	<p>(Definition 6.1 page 41)</p>



## K.5 Spectral Power

The term “*spectral power*” is a bit of an oxymoron because “spectral” deals with leaving the time-domain for the frequency-domain, howbeit the concept of power is solidly founded on the concept of time in that power = energy per time.

However, *Parseval's Theorem* (Proposition G.2 page 160) demonstrates that power in time can also be calculated in frequency. So, it makes some sense to speak of the term “spectral power”. Moreover, one way to estimate this power is to average the Fourier Transforms of the product  $|x(n)|^2 = x(n)x^*(n)$ ...that is, to use an estimate of the auto-spectral density  $\check{S}_{xx}(\omega)$ . Thus, an alternate name for *auto-spectral density* is **power spectral density** (PSD).

# APPENDIX L SPECTRAL THEORY

## L.1 Operator Spectrum

**Definition L.1.** <sup>1</sup> Let  $\mathbf{A} \in \mathcal{B}(\mathbf{X}, \mathbf{Y})$  be an operator over the linear spaces  $\mathbf{X} = (X, F, \oplus, \otimes)$  and  $\mathbf{Y} \triangleq (Y, F, \oplus, \otimes)$ . Let  $\mathcal{N}(\mathbf{A})$  be the NULL SPACE of  $\mathbf{A}$ .

**DEF** An **eigenvalue** of  $\mathbf{A}$  is any value  $\lambda$  such that there exists  $\mathbf{x}$  such that  $\mathbf{A}\mathbf{x} = \lambda\mathbf{x}$ .  
 The **eigenspace**  $H_\lambda$  of  $\mathbf{A}$  at eigenvalue  $\lambda$  is  $\mathcal{N}(\mathbf{A} - \lambda\mathbf{I})$ .  
 An **eigenvector** of  $\mathbf{A}$  associated with eigenvalue  $\lambda$  is any element of  $\mathcal{N}(\mathbf{A} - \lambda\mathbf{I})$ .

**Example L.1.** <sup>2</sup> Let  $\mathbf{D}$  be the differntial operator.









**EX** The set  $\{e^{\lambda x} | \lambda \in \mathbb{C}\}$  are the eigenvectors of  $\mathbf{D}$ .  
 $\rho(\mathbf{D}) = \emptyset$  ( $\mathbf{D}$  has no non-spectral points whatsoever)  
 $\sigma_p(\mathbf{D}) = \sigma(\mathbf{D})$  (the spectrum of  $\mathbf{D}$  is all eigenvalues)  
 $\sigma_c(\mathbf{D}) = \emptyset$  ( $\mathbf{D}$  has no continuous spectrum)  
 $\sigma_r(\mathbf{D}) = \emptyset$  ( $\mathbf{D}$  has no resolvent spectrum)

 **PROOF:**

$$\begin{aligned}
 (\mathbf{D} - \lambda\mathbf{I})e^{\lambda x} &= \mathbf{D}e^{\lambda x} - \lambda\mathbf{I}e^{\lambda x} \\
 &= \lambda e^{\lambda x} - \lambda e^{\lambda x} \\
 &= 0
 \end{aligned}
 \qquad \forall \lambda \in \mathbb{C}$$

This theorem and proof needs more work and investigation to prove/disprove its claims.  $\Rightarrow$

**Definition L.2.** <sup>3</sup> Let  $\mathbf{A} \in \mathcal{B}(\mathbf{X}, \mathbf{Y})$  be an operator over the linear spaces  $\mathbf{X} = (X, F, \oplus, \otimes)$  and  $\mathbf{Y} \triangleq (Y, F, \oplus, \otimes)$ .

<sup>1</sup>  Bollobás (1999) page 168,  Descartes (1637a),  Descartes (1954),  Cayley (1858),  Hilbert (1904) page 67,  Hilbert (1912),  
<sup>2</sup>  Pedersen (2000) page 79  
<sup>3</sup>  Michel and Herget (1993) page 439

quantity	$\mathcal{N}(\mathbf{A} - \lambda\mathbf{I}) = \{\mathbf{0}\}$ ( $\mathbf{x} = \mathbf{0}$ is the only solution)	$\overline{\mathcal{R}(\mathbf{A} - \lambda\mathbf{I})} = \mathbf{X}$ (dense)	$(\mathbf{A} - \lambda\mathbf{I})^{-1} \in \mathcal{B}(\mathbf{X}, \mathbf{Y})$ (continuous/bounded)
$\rho(\mathbf{A})$ (resolvent set)	1	1	1
$\sigma_p(\mathbf{A})$ (point spectrum)	0		
$\sigma_r(\mathbf{A})$ (residual spectrum)	1	0	
$\sigma_c(\mathbf{A})$ (continuous spectrum)	1	1	0

Table L.1: Spectrum of an operator  $\mathbf{A}$ 

The **resolvent set**  $\rho(\mathbf{A})$  of operator  $\mathbf{A}$  is defined as

$$\rho(\mathbf{A}) \triangleq \left\{ \lambda \in F \mid \begin{array}{l} 1. \mathcal{N}(\mathbf{A} - \lambda\mathbf{I}) = \{\mathbf{0}\} \\ 2. \overline{\mathcal{R}(\mathbf{A} - \lambda\mathbf{I})} = \mathbf{X} \\ 3. (\mathbf{A} - \lambda\mathbf{I})^{-1} \in \mathcal{B}(\mathbf{X}, \mathbf{Y}) \end{array} \right\} \quad \begin{array}{l} \text{(no non-zero eigenvectors)} \\ \text{(the range is dense in } \mathbf{X} \text{).} \\ \text{(inverse is continuous/bounded).} \end{array} \quad \text{and} \quad \text{and}$$

The **spectrum**  $\sigma(\mathbf{A})$  of operator  $\mathbf{A}$  is defined as

$$\sigma(\mathbf{A}) \triangleq F \setminus \rho(\mathbf{A}).$$

**Definition L.3.** <sup>4</sup> Let  $\mathbf{A} \in \mathcal{B}(\mathbf{X}, \mathbf{Y})$  be an operator over the linear spaces  $\mathbf{X} = (X, F, \oplus, \otimes)$  and  $\mathbf{Y} \triangleq (Y, F, \oplus, \otimes)$ .

The **point spectrum**  $\sigma_p(\mathbf{A})$  of operator  $\mathbf{A}$  is defined as


$$\sigma_p(\mathbf{A}) \triangleq \left\{ \lambda \in F \mid 1. \mathcal{N}(\mathbf{A} - \lambda\mathbf{I}) \supsetneq \{\mathbf{0}\} \right\} \quad \text{(has non-zero eigenvector)}$$

The **residual spectrum**  $\sigma_r(\mathbf{A})$  of operator  $\mathbf{A}$  is defined as

$$\sigma_r(\mathbf{A}) \triangleq \left\{ \lambda \in F \mid \begin{array}{l} 1. \mathcal{N}(\mathbf{A} - \lambda\mathbf{I}) = \{\mathbf{0}\} \\ 2. \overline{\mathcal{R}(\mathbf{A} - \lambda\mathbf{I})} \neq \mathbf{X} \end{array} \right\} \quad \begin{array}{l} \text{(no non-zero eigenvectors)} \\ \text{(not dense in } \mathbf{X} \text{—has gaps).} \end{array} \quad \text{and}$$

The **continuous spectrum**  $\sigma_c(\mathbf{A})$  of operator  $\mathbf{A}$  is defined as

$$\sigma_c(\mathbf{A}) \triangleq \left\{ \lambda \in F \mid \begin{array}{l} 1. \mathcal{N}(\mathbf{A} - \lambda\mathbf{I}) = \{\mathbf{0}\} \\ 2. \overline{\mathcal{R}(\mathbf{A} - \lambda\mathbf{I})} = \mathbf{X} \\ 3. (\mathbf{A} - \lambda\mathbf{I})^{-1} \notin \mathcal{B}(\mathbf{X}, \mathbf{Y}) \end{array} \right\} \quad \begin{array}{l} \text{(no non-zero eigenvectors)} \\ \text{(dense in } \mathbf{X} \text{).} \\ \text{(not continuous / not bounded)} \end{array} \quad \text{and}$$



The spectral components' definitions are illustrated in the figure to the left and summarized in Table L.1 (page 210). Let a family of operators  $\mathbf{B}(\lambda)$  be defined with respect to an operator  $\mathbf{A}$  such that  $\mathbf{B}(\lambda) \triangleq (\mathbf{A} - \lambda\mathbf{I})$ . Normally, we might expect a “normal” or “regular” or even “mundane” operator  $\mathbf{B}(\lambda)$  to have the properties

1.  $\mathbf{B}(\lambda)\mathbf{x} = \mathbf{0}$  if and only if  $\mathbf{x} = \mathbf{0}$
2.  $\mathbf{B}(\lambda)\mathbf{x}$  spans virtually all of  $\mathbf{X}$  as we vary  $\mathbf{x}$
3.  $\mathbf{B}^{-1}(\lambda)$  is continuous.

After all, these are the properties that we would have if  $\mathbf{B}(\lambda)$  were simply an affine operator in the

<sup>4</sup> Bollobás (1999) page 168, Hilbert (1906) pages 169–172

field of real numbers— such as  $[\mathbf{B}(\lambda)](x) \triangleq [\lambda](x) = \lambda x$  which is 0 if and only if  $x = 0$ , has range  $\mathcal{R}(\lambda) = \mathbb{R}$ , and its inverse  $\lambda^{-1}x$  is continuous.

If for some  $\lambda$  the operator  $\mathbf{B}(\lambda)$  does have all these “regular” properties, then that  $\lambda$  part of the *resolvent set* of  $\mathbf{A}$  and  $\lambda$  is called *regular*. However if for some  $\lambda$  the operator  $\mathbf{B}(\lambda)$  fails any of these conditions, then that  $\lambda$  part of the *spectrum* of  $\mathbf{A}$ . And which conditions it fails determines which component of the spectrum it is in.

**Theorem L.1.**<sup>5</sup> Let  $\mathbf{A} \in \mathcal{B}(\mathbf{X}, \mathbf{Y})$  be an operator.

$$\sigma(\mathbf{A}) = \sigma_p(\mathbf{A}) \cup \sigma_c(\mathbf{A}) \cup \sigma_r(\mathbf{A})$$

**Theorem L.2** (Spectral Theorem).<sup>6</sup> Let  $\mathbf{N} \in \mathcal{Y}^{\mathbf{X}}$  be an operator.

$$\left. \begin{array}{l} \text{(A). } \underbrace{\mathbf{N}^* \mathbf{N} = \mathbf{N} \mathbf{N}^*}_{\mathbf{N} \text{ is NORMAL}} \\ \text{(B). } \mathbf{N} \text{ is COMPACT} \end{array} \right\} \Rightarrow \left\{ \begin{array}{l} \text{(1). } \mathbf{N} = \sum_n \lambda_n \mathbf{P}_n \\ \text{(2). } \sum_n \mathbf{P}_n = \mathbf{I} \\ \text{(3). } \mathbf{P}_n \mathbf{P}_m = \delta_{n-m} \mathbf{P}_n \\ \text{(4). } \dim(\mathbf{H}_n) < \infty \\ \text{(5). } \left| \{ \lambda_n \mid \lambda_n \neq 0 \} \right| \text{ is COUNTABLY INFINITE} \end{array} \right.$$

where

$$\begin{aligned} (\lambda_n)_{n \in \mathbb{Z}} &\triangleq \sigma_p(\mathbf{N}) && \text{(eigenvalues of } \mathbf{N}) \\ \mathbf{H}_n &\triangleq \mathcal{N}(\mathbf{N} - \lambda_n \mathbf{I}) && (\lambda_n \text{ is the eigenspace of } \mathbf{N} \text{ at } \lambda_n \text{ in } \mathbf{Y}) \\ \mathbf{H}_n &= \mathbf{P}_n \mathbf{Y} && (\mathbf{P}_n \text{ is the projection operator that generates } \mathbf{H}_n) \end{aligned}$$

## L.2 Fredholm kernels

**Definition L.4.**<sup>7</sup>

A **Fredholm operator**  $\mathbf{K}$  is defined as

$$[\mathbf{K}f](t) \triangleq \int_a^b \underbrace{\kappa(t, s)f(s) \, ds}_{\text{kernel}} \quad \forall f \in \mathcal{L}_2([a, b])$$

Fredholm integral equation of the first kind<sup>8</sup>

**Example L.2.** Examples of Fredholm operators include

- |                              |   |                                |
|------------------------------|---|--------------------------------|
| 1. Fourier Transform         | $[\tilde{\mathbf{F}}\mathbf{x}](f) = \int_t \mathbf{x}(t)e^{-i2\pi ft} \, dt$                     | $\kappa(t, f) = e^{-i2\pi ft}$ |
| 2. Inverse Fourier Transform | $[\tilde{\mathbf{F}}^{-1}\tilde{\mathbf{x}}](t) = \int_f \tilde{\mathbf{x}}(f)e^{i2\pi ft} \, df$ | $\kappa(f, t) = e^{i2\pi ft}$  |
| 3. Laplace operator          | $[\mathbf{L}\mathbf{x}](s) = \int_t \mathbf{x}(t)e^{-st} \, dt$                                   | $\kappa(t, s) = e^{-st}$       |
| 4. autocorrelation operator  | $[\mathbf{R}\mathbf{x}](t) = \int_s R(t, s)\mathbf{x}(s) \, ds$                                   | $\kappa(t, s) = R(t, s)$       |

**Theorem L.3.** Let  $\mathbf{K}$  be a Fredholm operator with kernel  $\kappa(t, s)$  and adjoint  $\mathbf{K}^*$ .

$$[\mathbf{K}f](t) = \int_A \kappa(t, s)f(s) \, ds \quad \Longleftrightarrow \quad [\mathbf{K}^*f](t) = \int_A \kappa^*(s, t)f(s) \, ds$$

<sup>5</sup> Michel and Herget (1993) page 440

<sup>6</sup> Michel and Herget (1993) page 457, Bollobás (1999) page 200, Hilbert (1906), Hilbert (1912), von Neumann (1929), de Witt (1659)

<sup>7</sup> Michel and Herget (1993) page 425

<sup>8</sup>The equation  $\int_u \kappa(t, s)f(s) \, ds$  is a **Fredholm integral equation of the first kind** and  $\kappa(t, u)$  is the **kernel** of the equation. References: Fredholm (1900), Fredholm (1903) page 365, Michel and Herget (1993) page 97, Keener (1988) page 101

✎ PROOF:

$$\begin{aligned}
 [\mathbf{K}f](t) &= \int_A \kappa(t, s) f(s) \, ds \\
 \Leftrightarrow \langle [\mathbf{K}f](t) \mid g(t) \rangle &= \left\langle \int_s \kappa(t, s) f(s) \, ds \mid g(t) \right\rangle && \text{by left hypothesis} \\
 &= \int_s f(s) \langle \kappa(t, s) \mid g(t) \rangle \, ds && \text{by additivity property of } \langle \Delta \mid \nabla \rangle \\
 &= \int_s f(s) \langle g(t) \mid \kappa(t, s) \rangle^* \, ds && \text{by conjugate symmetry property of } \langle \Delta \mid \nabla \rangle \\
 &= \langle f(s) \mid \langle g(t) \mid \kappa(t, s) \rangle \rangle && \text{by local definition of } \langle \Delta \mid \nabla \rangle \\
 &= \left\langle f(s) \mid \underbrace{\int_t \kappa^*(t, s) g(t) \, dt}_{[\mathbf{K}^*g](s)} \right\rangle && \text{by local definition of } \langle \Delta \mid \nabla \rangle \\
 \Leftrightarrow [\mathbf{K}^*g](s) &= \int_A \kappa^*(t, s) g(t) \, dt && \text{by right hypothesis} \\
 \Leftrightarrow [\mathbf{K}^*g](\sigma) &= \int_A \kappa^*(\tau, \sigma) g(\tau) \, d\tau && \text{by change of variable: } \tau = t, \sigma = s \\
 \Leftrightarrow [\mathbf{K}^*f](t) &= \int_A \kappa^*(s, t) f(s) \, ds && \text{by change of variable: } t = \sigma, s = \tau, f = g
 \end{aligned}$$

⇒

**Corollary L.1.** <sup>9</sup> Let  $\mathbf{K}$  be an Fredholm operator with kernel  $\kappa(t, s)$  and adjoint  $\mathbf{K}^*$ .

COR

$$\underbrace{\mathbf{K} = \mathbf{K}^*}_{\mathbf{K} \text{ is self-adjoint}} \Leftrightarrow \underbrace{\kappa(t, s) = \kappa^*(s, t)}_{\text{kernel is conjugate symmetric}}$$

✎ PROOF:

$$\begin{aligned}
 \mathbf{K} = \mathbf{K}^* &\Leftrightarrow \int_A \kappa(t, s) f(s) \, ds = \int_A \kappa^*(s, t) f(s) \, ds && \text{by Theorem L.3 page 211} \\
 &\Leftrightarrow \kappa(t, s) = \kappa^*(s, t)
 \end{aligned}$$

⇒

**Theorem L.4** (Mercer's Theorem). <sup>10</sup> Let  $\mathbf{K}$  be an Fredholm operator with kernel  $\kappa(t, s)$  and eigen-system  $((\lambda_n, \phi_n(t)))_{n \in \mathbb{Z}}$ .

THM

$$\left\{ \begin{array}{l} \text{(A). } \underbrace{\int_a^b \int_a^b \kappa(t, s) f(t) f^*(s) \, dt}_{\text{positive}} \geq 0 \quad \text{and} \\ \text{(B). } \kappa(t, s) \text{ is CONTINUOUS on } [a : b] \times [a : b] \end{array} \right\} \Rightarrow \left\{ \begin{array}{l} \text{(1). } \kappa(t, s) = \sum_n \lambda_n \phi_n(t) \phi_n^*(s) \quad \text{and} \\ \text{(2). } \kappa(t, s) \text{ CONVERGES ABSOLUTELY and UNIFORMLY on } [a : b] \times [a : b] \end{array} \right\}$$

<sup>9</sup> Michel and Herget (1993) page 430

<sup>10</sup> Gohberg et al. (2003) page 198, Courant and Hilbert (1930) pages 138–140, Mercer (1909) page 439



## Back Matter




*“It appears to me that if one wants to make progress in mathematics, one should study the masters and not the pupils.”*

Niels Henrik Abel (1802–1829), Norwegian mathematician <sup>11</sup>




*“When evening comes, I return home and go to my study. On the threshold I strip naked, taking off my muddy, sweaty workaday clothes, and put on the robes of court and palace, and in this graver dress I enter the courts of the ancients and am welcomed by them, and there I taste the food that alone is mine, and for which I was born. And there I make bold to speak to them and ask the motives of their actions, and they, in their humanity reply to me. And for the space of four hours I forget the world, remember no vexation, fear poverty no more, tremble no more at death; I pass indeed into their world.”*

Niccolò Machiavelli (1469–1527), Italian political philosopher, in a 1513 letter to friend Francesco Vettori. <sup>12</sup>



ancient library of Alexandria



The Book Worm by Carl Spitzweg, circa 1850

13

<sup>11</sup> quote: [Simmons \(2007\)](#) page 187.

image: [http://en.wikipedia.org/wiki/Image:Niels\\_Henrik\\_Abel.jpg](http://en.wikipedia.org/wiki/Image:Niels_Henrik_Abel.jpg), public domain

<sup>12</sup> quote: [Machiavelli \(1961\)](#) page 139?.


image: [http://commons.wikimedia.org/wiki/File:Santi\\_di\\_Tito\\_-\\_Niccolo\\_Machiavelli%27s\\_portrait\\_headcrop.jpg](http://commons.wikimedia.org/wiki/File:Santi_di_Tito_-_Niccolo_Machiavelli%27s_portrait_headcrop.jpg), public domain

<sup>13</sup> <http://en.wikipedia.org/wiki/File:Ancientlibraryalex.jpg>, public domain [http://en.wikipedia.org/wiki/File:Carl\\_Spitzweg\\_021.jpg](http://en.wikipedia.org/wiki/File:Carl_Spitzweg_021.jpg),



*“To sit alone in the lamplight with a book spread out before you, and hold intimate converse with men of unseen generations—such is a pleasure beyond compare.”*

[Yoshida Kenko \(Urabe Kaneyoshi\)](#) (1283? – 1350?), Japanese author and Buddhist monk <sup>14</sup>

<sup>14</sup> quote:  [Kenko \(circa 1330\)](#)  
image: [http://en.wikipedia.org/wiki/Yoshida\\_Kenko](http://en.wikipedia.org/wiki/Yoshida_Kenko)



## BIBLIOGRAPHY

- A prelude to sampling, wavelets, and tomography. In John Benedetto and Ahmed I. Zayed, editors, *Sampling, Wavelets, and Tomography*, Applied and Numerical Harmonic Analysis, pages 1–32. Springer, 2004. ISBN 9780817643041. URL <http://books.google.com/books?vid=ISBN0817643044>.
- Yuri A. Abramovich and Charalambos D. Aliprantis. *An Invitation to Operator Theory*. American Mathematical Society, Providence, Rhode Island, 2002. ISBN 0-8218-2146-6. URL <http://books.google.com/books?vid=ISBN0821821466>.
- Milton Abramowitz and Irene A. Stegun, editors. *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables*. National Bureau of Standards, 1972. URL <http://www.cs.bham.ac.uk/~aps/research/projects/as/book.php>.
- Edward H. Adelson and Peter J. Burt. Image data compression with the laplacian pyramid. In *Proceedings of the Pattern Recognition and Information Processing Conference*, pages 218–223, Dallas Texas, 1981. IEEE Computer Society Press. URL [citeseerx.ist.psu.edu/viewdoc/summary?doi=10.1.1.50.791](http://citeseerx.ist.psu.edu/viewdoc/summary?doi=10.1.1.50.791).
- Charalambos D. Aliprantis and Owen Burkinshaw. *Principles of Real Analysis*. Academic Press, London, 3 edition, 1998. ISBN 9780120502578. URL <http://www.amazon.com/dp/0120502577>.
- Luis Alvarez, Frédéric Guichard, Pierre-Louis Lions, and Jean Michel Morel. Axioms and fundamental equations of image processing. *Archive for Rational Mechanics and Analysis*, 123(3):199–257, 1993. URL <http://link.springer.com/article/10.1007/BF00375127>.
- George E. Andrews, Richard Askey, and Ranjan Roy. *Special Functions*, volume 71 of *Encyclopedia of mathematics and its applications*. Cambridge University Press, Cambridge, U.K., new edition, February 15 2001. ISBN 0521789885. URL <http://books.google.com/books?vid=ISBN0521789885>.
- Kendall E. Atkinson and Weimin Han. *Theoretical Numerical Analysis: A Functional Analysis Framework*, volume 39 of *Texts in Applied Mathematics*. Springer, 3 edition, 2009. ISBN 9781441904584. URL <http://books.google.com/books?vid=ISBN1441904581>.
- Léon Autonne. Sur l'hermitien (on the hermitian). In *Comptes Rendus Des Séances De L'Académie Des Sciences*, volume 133, pages 209–268. De L'Académie des sciences (Academy of Sciences), Paris, 1901. URL <http://visualiseur.bnf.fr/Visualiseur?O=NUMM-3089>. Comptes Rendus Des Séances De L'Académie Des Sciences (Reports Of the Meetings Of the Academy of Science).

Léon Autonne. Sur l'hermitien (on the hermitian). *Rendiconti del Circolo Matematico di Palermo*, 16:104–128, 1902. Rendiconti del Circolo Matematico di Palermo (Statements of the Mathematical Circle of Palermo).

George Bachman. *Elements of Abstract Harmonic Analysis*. Academic paperbacks. Academic Press, New York, 1964. URL <http://books.google.com/books?id=ZP8-AAAAIAAJ>.

George Bachman and Lawrence Narici. *Functional Analysis*. Academic Press textbooks in mathematics; Pure and Applied Mathematics Series. Academic Press, 1 edition, 1966. ISBN 9780486402512. URL <http://books.google.com/books?vid=ISBN0486402517>. “unabridged re-publication” available from Dover (isbn 0486402517).

George Bachman, Lawrence Narici, and Edward Beckenstein. *Fourier and Wavelet Analysis*. Universitext Series. Springer, 2000. ISBN 9780387988993. URL <http://books.google.com/books?vid=ISBN0387988998>.

Stefan Banach. Sur les opérations dans les ensembles abstraits et leur applications aux équations intégrales (on abstract operations and their applications to the integral equations). *Fundamenta Mathematicae*, 3:133–181, 1922. URL <http://matwbn.icm.edu.pl/ksiazki/fm/fm3/fm3120.pdf>.

Stefan Banach. *Théorie des opérations linéaires*. Monografie Matematyczne, Warsaw, Poland, 1932a. URL <http://matwbn.icm.edu.pl/kstresc.php?tom=1&wyd=10>. (Theory of linear operations).

Stefan Banach. *Theory of Linear Operations*, volume 38 of *North-Holland mathematical library*. North-Holland, Amsterdam, 1932b. ISBN 0444701842. URL <http://www.amazon.com/dp/0444701842/>. English translation of 1932 French edition, published in 1987.

Adi Ben-Israel and Robert P. Gilbert. *Computer-supported calculus*. Texts and monographs in symbolic computation. Springer, 2002. ISBN 3-211-82924-5. URL <http://books.google.com/books?vid=ISBN3211829245>.

Ladislav Beran. Three identities for ortholattices. *Notre Dame Journal of Formal Logic*, 17(2):251–252, 1976. doi: 10.1305/ndjfl/1093887530. URL <http://projecteuclid.org/euclid.ndjfl/1093887530>.

Ladislav Beran. *Orthomodular Lattices: Algebraic Approach*. Mathematics and Its Applications (East European Series). D. Reidel Publishing Company, Dordrecht, 1985. ISBN 90-277-1715-X. URL <http://books.google.com/books?vid=ISBN902771715X>.

Sterling Khazag Berberian. *Introduction to Hilbert Space*. Oxford University Press, New York, 1961. URL <http://books.google.com/books?vid=ISBN0821819127>.

M. Bertero and P. Boccacci. *Introduction to Inverse Problems in Imaging*. CRC Press, 1998. ISBN 9781439822067. URL <http://books.google.com/books?vid=ISBN9781439822067>.

Garrett Birkhoff and John Von Neumann. The logic of quantum mechanics. *The Annals of Mathematics*, 37(4):823–843, October 1936. URL <http://www.jstor.org/stable/1968621>.

Duncan Black, Ian Brooks, and et al. Robert Groves, editors. *Collins English Dictionary—Complete and Unabridged*. HarperCollins Publishers, 10 edition, 2009. ISBN 978-0-00-732119-3. URL <http://books.google.com/books?vid=ISBN9780007321193>.

- Béla Bollobás. *Linear Analysis; an introductory course*. Cambridge mathematical textbooks. Cambridge University Press, Cambridge, 2 edition, March 1 1999. ISBN 978-0521655774. URL <http://books.google.com/books?vid=ISBN0521655773>.
- Umberto Bottazzini. *The Higher Calculus: A History of Real and Complex Analysis from Euler to Weierstrass*. Springer-Verlag, New York, 1986. ISBN 0-387-96302-2. URL <http://books.google.com/books?vid=ISBN0387963022>.
- A. Böttcher and B. Silbermann. *Introduction to Large Truncated Toeplitz Matrices*. Springer, 1999. ISBN 9780387985701. URL <http://books.google.com/books?vid=ISBN9780387985701>.
- Carl Benjamin Boyer and Uta C. Merzbach. *A History of Mathematics*. Wiley, New York, 2 edition, 1991. ISBN 0471543977. URL <http://books.google.com/books?vid=ISBN0471543977>.
- Ronald Newbold Bracewell. *The Fourier transform and its applications*. McGraw-Hill electrical and electronic engineering series. McGraw-Hill, 2, illustrated, international student edition edition, 1978. ISBN 9780070070134. URL <http://books.google.com/books?vid=ISBN007007013X>.
- Thomas John I'Anson Bromwich. *An Introduction to the Theory of Infinite Series*. Macmillan and Company, 1 edition, 1908. ISBN 9780821839768. URL <http://www.archive.org/details/anintroductiont00bromgoog>.
- Andrew M. Bruckner, Judith B. Bruckner, and Brian S. Thomson. *Real Analysis*. Prentice-Hall, Upper Saddle River, N.J., 1997. ISBN 9780134588865. URL <http://books.google.com/books?vid=ISBN013458886X>.
- C. Sidney Burrus, Ramesh A. Gopinath, and Haitao Guo. *Introduction to Wavelets and Wavelet Transforms; A Primer*. Prentice Hall, Upper Saddle River, New Jersey, USA, 1998. ISBN 0-13-489600-9. URL <http://www.dsp.rice.edu/software/wavebook.shtml>.
- Peter J. Burt and Edward H. Adelson. The laplacian pyramid as a compact image code. *IEEE Transactions On Communications*, COM-3L(4):532–540, April 1983. URL <http://citeseer.ist.psu.edu/burt83laplacian.html>.
- Charles L. Byrne. *Signal Processing: A Mathematical Approach*. AK Peters Series. A K Peters, 2005. ISBN 1568812426.
- James A. Cadzow. *Foundations of Digital Signal Processing and Data Analysis*. Macmillan Publishing Company, New York, 1987. ISBN 0023180102. URL [http://frontweb.vuse.vanderbilt.edu/vuse\\_web/directory/facultybio.asp?FacultyID=9](http://frontweb.vuse.vanderbilt.edu/vuse_web/directory/facultybio.asp?FacultyID=9).
- Florian Cajori. A history of mathematical notations; notations mainly in higher mathematics. In *A History of Mathematical Notations; Two Volumes Bound as One*, volume 2. Dover, Mineola, New York, USA, 1993. ISBN 0-486-67766-4. URL <http://books.google.com/books?vid=ISBN0486677664>. reprint of 1929 edition by *The Open Court Publishing Company*.
- Gerolamo Cardano. *Ars Magna or the Rules of Algebra*. Dover Publications, Mineola, New York, 1545. ISBN 0486458733. URL <http://www.amazon.com/dp/0486458733>. English translation of the Latin *Ars Magna* edition, published in 2007.
- Lennart Axel Edvard Carleson. Convergence and growth of partial sums of fourier series. *Acta Mathematica*, 116(1):135–157, 1966. doi: 10.1007/BF02392815. URL <http://dx.doi.org/10.1007/BF02392815>.

- Lennart Axel Edvard Carleson and Björn Engquist. After the 'golden age': what next? lennart carleson interviewed by björn engquist. In Björn Engquist and Wilfried Schmid, editors, *Mathematics Unlimited—2001 and beyond*, pages 455–461. Springer, Berlin, 2001. ISBN 978-3-540-66913-5. URL [https://link.springer.com/chapter/10.1007/978-3-642-56478-9\\_22](https://link.springer.com/chapter/10.1007/978-3-642-56478-9_22).
- N. L. Carothers. *A Short Course on Banach Space Theory*. Number 64 in London Mathematical Society Student Texts. Cambridge University Press, Cambridge, 2005. ISBN 0521842832. URL <http://www.comm.musashi-tech.ac.jp/~arimoto/Drarimoto/preprints/calothers/Assemble.ps>.
- Peter G. Casazza and Mark C. Lammers. Bracket products for weyl-heisenberg frames. In Hans G. Feichtinger and Thomas Strohmer, editors, *Gabor Analysis and Algorithms: Theory and Applications*, Applied and Numerical Harmonic Analysis, pages 71–98. Birkhäuser, 1998. ISBN 9780817639594.
- Clémente Ibarra Castanedo. *Quantitative subsurface defect evaluation by pulsed phase thermography: depth retrieval with the phase*. PhD thesis, Université Laval, October 2005. URL <http://archimede.bibl.ulaval.ca/archimede/fichiers/23016/23016.html>. Faculte' Des Sciences Et De Génie.
- Arthur Cayley. A memoir on the theory of matrices. *Philosophical Transactions of the Royal Society of London*, 148:17–37, 1858. ISSN 1364-503X. URL <http://www.jstor.org/view/02610523/ap000059/00a00020/0>.
- Joan Cerdà. *Linear functional analysis*, volume 116 of *Graduate studies in mathematics*. American Mathematical Society, July 16 2010. ISBN 0821851152. URL <http://books.google.com/books?vid=ISBN0821851152>.
- Alexandre J. Chorin and Ole H. Hald. *Stochastic Tools in Mathematics and Science*, volume 1 of *Surveys and Tutorials in the Applied Mathematical Sciences*. Springer, New York, 2 edition, 2009. ISBN 978-1-4419-1001-1. URL <http://books.google.com/books?vid=ISBN9781441910011>.
- Ole Christensen. *An Introduction to Frames and Riesz Bases*. Applied and Numerical Harmonic Analysis. Birkhäuser, Boston/Basel/Berlin, 2003. ISBN 0-8176-4295-1. URL <http://books.google.com/books?vid=ISBN0817642951>.
- Ole Christensen. *Frames and bases: An Introductory Course*. Applied and Numerical Harmonic Analysis. Birkhäuser, Boston/Basel/Berlin, 2008. ISBN 9780817646776. URL <http://books.google.com/books?vid=ISBN0817646779>.
- Charles K. Chui. *An Introduction to Wavelets*. Academic Press, San Diego, California, USA, January 3 1992. ISBN 9780121745844. URL <http://books.google.com/books?vid=ISBN0121745848>.
- Jon F. Claerbout. *Fundamentals of Geophysical Data Processing with Applications to Petroleum Prospecting*. International series in the earth and planetary sciences, Tab Mastering Electronics Series. McGraw-Hill, New York, 1976. ISBN 9780070111172. URL <http://sep.stanford.edu/sep/prof/>.
- Peter M. Clarkson. *Optimal and Adaptive Signal Processing*. Electronic Engineering Systems Series. CRC Press, 1993. ISBN 0849386098. URL <http://books.google.com/books?vid=isbn0849386098>.
- Paul M. Cohn. *Basic Algebra; Groups, Rings and Fields*. Springer, December 6 2002. ISBN 1852335874. URL <http://books.google.com/books?vid=isbn1852335874>.
- R. Courant and D. Hilbert. *Methods of Mathematical Physics*, volume 1. Interscience Publishers, New York, 1930. URL <http://www.worldcat.org/isbn/0471504475>.



- Xingde Dai and David R. Larson. *Wandering vectors for unitary systems and orthogonal wavelets*. Number 640 in Memoirs of the American Mathematical Society. American Mathematical Society, Providence R.I., July 1998. ISBN 0821808001. URL <http://books.google.com/books?vid=ISBN0821808001>.
- Xingde Dai and Shijie Lu. Wavelets in subspaces. *Michigan Math. J.*, 43(1):81–98, 1996. doi: 10.1307/mmj/1029005391. URL <http://projecteuclid.org/euclid.mmj/1029005391>.
- Ingrid Daubechies. *Ten Lectures on Wavelets*. Society for Industrial and Applied Mathematics, Philadelphia, 1992. ISBN 0-89871-274-2. URL <http://www.amazon.com/dp/0898712742>.
- Ingrid Daubechies, A. Grossman, and Y. Meyer. Painless nonorthogonal expansions. *Journal of Mathematical Physics*, 27(5):1271–1283, May 1986. ISSN 0022-2488. URL [link.aip.org/link/?jmp/27/1271](http://link.aip.org/link/?jmp/27/1271).
- Kenneth R. Davidson and Allan P. Donsig. *Real Analysis and Applications*. Springer, 2010. ISBN 9781441900050. URL <http://books.google.com/books?vid=ISBN1441900055>.
- Charles Jean de la Vallée-Poussin. Sur l'intégrale de lebesgue. *Transactions of the American Mathematical Society*, 16(4):435–501, October 1915. URL <http://www.jstor.org/stable/1988879>.
- Juan-Arias de Reyna. *Pointwise Convergence of Fourier Series*, volume 1785 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin/Heidelberg/New York, 2002. ISBN 3540432701. URL <http://books.google.com/books?vid=ISBN3540432701>.
- Johan de Witt. Elementa curvarum linearum (elements of curves). In *La Géométrie*. 1659. Elementa curvarum linearum was an appendix to de Witt's translation of of Descartes' *La Géométrie*.
- René Descartes. *La géométrie*. 1637a. URL [http://historical.library.cornell.edu/math/math\\_D.html](http://historical.library.cornell.edu/math/math_D.html).
- René Descartes. *Discours de la méthode pour bien conduire sa raison, et chercher la verite' dans les sciences*. Jan Maire, Leiden, 1637b. URL <http://www.gutenberg.org/etext/13846>.
- René Descartes. *Discourse on the Method of Rightly Conducting the Reason in the Search for Truth in the Sciences*. 1637c. URL <http://www.gutenberg.org/etext/59>.
- René Descartes. *Regulae ad directionem ingenii*. 1684a. URL [http://www.fh-augsburg.de/~harsch/Chronologia/Lspost17/Descartes/des\\_re00.html](http://www.fh-augsburg.de/~harsch/Chronologia/Lspost17/Descartes/des_re00.html).
- René Descartes. *Rules for Direction of the Mind*. 1684b. URL [http://en.wikisource.org/wiki/Rules\\_for\\_the\\_Direction\\_of\\_the\\_Mind](http://en.wikisource.org/wiki/Rules_for_the_Direction_of_the_Mind).
- René Descartes. *The Geometry of Rene Descartes*. Courier Dover Publications, June 1 1954. ISBN 0486600688. URL <http://books.google.com/books?vid=isbn0486600688>. originally published by Open Court Publishing, Chicago, 1925; translation of *La géométrie*.
- Analog Devices. Adl5387. Technical report, 2016. URL <https://www.analog.com/media/en/technical-documentation/data-sheets/ADL5387.pdf>. “30 MHz to 2 GHz Quadrature Demodulator”.
- Jean Alexandre Dieudonné. *Foundations of Modern Analysis*. Academic Press, New York, 1969. ISBN 1406727911. URL <http://books.google.com/books?vid=ISBN1406727911>.
- R. J. Duffin and A. C. Schaeffer. A class of nonharmonic fourier series. *Transactions of the American Mathematical Society*, 72(2):341–366, March 1952. ISSN 1088-6850. URL <http://www.jstor.org/stable/1990760>.

- Bogdan Dumitrescu. *Positive Trigonometric Polynomials and Signal Processing Applications*. Signals and Communication Technology. Springer, 2007. ISBN 978-1-4020-5124-1. URL [gen.lib.rus.ec/get?md5=5346e169091b2d928d8333cd053300f9](http://gen.lib.rus.ec/get?md5=5346e169091b2d928d8333cd053300f9).
- Nelson Dunford and Jacob T. Schwartz. *Linear operators. Part 1, General Theory*, volume 7 of *Pure and applied mathematics*. Interscience Publishers, New York, 1957. ISBN 0471226394. URL <http://www.amazon.com/dp/0471608483>. with the assistance of William G. Bade and Robert G. Bartle.
- John R. Durbin. *Modern Algebra; An Introduction*. John Wiley & Sons, Inc., 4 edition, 2000. ISBN 0-471-32147-8. URL <http://www.worldcat.org/isbn/0471321478>.
- Robert E. Edwards. *Functional Analysis: Theory and Applications*. Dover books on mathematics. Dover, New York, 1995. ISBN 0-486-68143-2. URL <http://books.google.com/books?vid=ISBN0486681432>.
- Yuli Eidelman, Vitali D. Milman, and Antonis Tzolomitis. *Functional Analysis: An Introduction*, volume 66 of *Graduate Studies in Mathematics*. American Mathematical Society, 2004. ISBN 0821836463. URL <http://books.google.com/books?vid=ISBN0821836463>.
- Per Enflo. A counterexample to the approximation problem in banach spaces. *Acta Mathematica*, 130:309–317, 1973. URL <http://link.springer.com/content/pdf/10.1007/BF02392270>.
- Leonhard Euler. *Introductio in analysin infinitorum*, volume 1. Marcum-Michaelem Bousquet & Socios, Lausannæ, 1748. URL <http://www.math.dartmouth.edu/~euler/pages/E101.html>. Introduction to the Analysis of the Infinite.
- Leonhard Euler. *Introduction to the Analysis of the Infinite*. Springer, 1988. ISBN 0387968245. URL <http://books.google.com/books?vid=ISBN0387968245>. translation of 1748 *Introductio in analysin infinitorum*.
- David Ewen. *The Book of Modern Composers*. Alfred A. Knopf, New York, 1950. URL <http://books.google.com/books?id=yHw4AAAAIAAJ>.
- David Ewen. *The New Book of Modern Composers*. Alfred A. Knopf, New York, 3 edition, 1961. URL <http://books.google.com/books?id=bZiAAAAIAAJ>.
- Marián Fabian, Petr Habala, Petr Hájek, Vicente Montesinos, Vaclav Zizler, and Václav Zizler. *Banach Space Theory: The Basis for Linear and Nonlinear Analysis*. CMS Books in Mathematics. Springer, 2010. ISBN 1441975144. URL <http://books.google.com/books?vid=isbn1441975144>.
- Lorenzo Farina and Sergio Rinaldi. *Positive Linear Systems: Theory and Applications*. Pure and applied mathematics. John Wiley & Sons, 1 edition, July 3 2000. ISBN 9780471384564. URL <http://books.google.com/books?vid=ISBN0471384569>.
- G.L. Fix and G. Strang. Fourier analysis of the finite element method in ritz-galerkin theory. *Studies in Applied Mathematics*, 48:265–273, 1969.
- Harley Flanders. Differentiation under the integral sign. *The American Mathematical Monthly*, 80(6):615–627, June–July 1973. URL [http://sgpwe.izt.uam.mx/files/users/uami/jdf/proyectos/Derivar\\_inetegral.pdf](http://sgpwe.izt.uam.mx/files/users/uami/jdf/proyectos/Derivar_inetegral.pdf). <http://www.jstor.org/pss/2319163>.
- Francis J. Flanigan. *Complex Variables; Harmonic and Analytic Functions*. Dover, New York, 1983. ISBN 9780486613888. URL <http://books.google.com/books?vid=ISBN0486613887>.

- Gerald B. Folland. *Fourier Analysis and its Applications*. Wadsworth & Brooks / Cole Advanced Books & Software, Pacific Grove, California, USA, 1992. ISBN 0-534-17094-3. URL <http://www.worldcat.org/isbn/0534170943>.
- Gerald B. Folland. *A Course in Abstract Harmonic Analysis*. Studies in Advanced Mathematics. CRC Press, Boca Raton, 1995. ISBN 0-8493-8490-7. URL <http://books.google.com/books?vid=ISBN0849384907>.
- Brigitte Forster and Peter Massopust, editors. *Four Short Courses on Harmonic Analysis: Wavelets, Frames, Time-Frequency Methods, and Applications to Signal and Image Analysis*. Applied and Numerical Harmonic Analysis. Springer, November 19 2009. ISBN 9780817648909. URL <http://books.google.com/books?vid=ISBN0817648909>.
- Jean-Baptiste-Joseph Fourier. Mémoire sur la propagation de la chaleur dans les corps solides (dissertation on the propagation of heat in solid bodies). In M. Gaston Darboux, editor, *Œuvres De Fourier*, pages 215–221. Ministère de L'instruction Publique, Paris, France, 2 edition, December 21 1807. URL <http://gallica.bnf.fr/ark:/12148/bpt6k33707/f220n7>.
- Jean-Baptiste-Joseph Fourier. *Théorie Analytique de la Chaleur (The Analytical Theory of Heat)*. Chez Firmin Didot, pere et fils, Paris, 1822. URL <http://books.google.com/books?vid=04X2vlqZx7hydlQUWEq&id=TDQJAAAAIAAJ>.
- Jean-Baptiste-Joseph Fourier. *The Analytical Theory of Heat (Théorie Analytique de la Chaleur)*. Cambridge University Press, Cambridge, February 20 1878. URL <http://www.archive.org/details/analyticaltheory00fourrich>. 1878 English translation of the original 1822 French edition. A 2003 Dover edition is also available: isbn 0486495310.
- Maurice René Fréchet. Sur quelques points du calcul fonctionnel (on some points of functional calculation). *Rendiconti del Circolo Matematico di Palermo*, 22:1–74, April 22 1906. URL [https://www.lpsm.paris/pageperso/mazliak/Frechet\\_1906.pdf](https://www.lpsm.paris/pageperso/mazliak/Frechet_1906.pdf). Rendiconti del Circolo Matematico di Palermo (Statements of the Mathematical Circle of Palermo).
- Maurice René Fréchet. *Les Espaces abstraits et leur théorie considérée comme introduction a l'analyse générale*. Borel series. Gauthier-Villars, Paris, 1928. URL <http://books.google.com/books?id=9czoHQAACAAJ>. Abstract spaces and their theory regarded as an introduction to general analysis.
- Erik Ivar Fredholm. Sur une nouvelle méthode pour la résolution du problème de dirichlet (on a new method for the resolution of the problem of dirichlet). */:Oefversigt af Kongl. Sv. Vetenskaps-Academiens Förhandlingar*, 57:39–66, 1900.
- Erik Ivar Fredholm. Sur une classe d'equations fonctionnelles (on a class of functional equations). *Acta Mathematica*, 27(1):365–390, December 1903. ISSN 0001-5962. doi: 10.1007/BF02421317. URL <http://www.springerlink.com/content/c41371137837p252/>.
- Ferdinand Georg Frobenius. Über lineare substitutionen und bilineare formen. *Journal für die reine und angewandte Mathematik (Crelle's Journal)*, 84:1–63, 1878. ISSN 0075-4102. URL <http://www.digizeitschriften.de/home/services/pdfterms/?ID=509796>.
- Ferdinand Georg Frobenius. Über lineare substitutionen und bilineare formen. In Jean Pierre Serre, editor, *Gesammelte Abhandlungen (Collected Papers)*, volume I, pages 343–405. Springer, Berlin, 1968. URL <http://www.worldcat.org/oclc/253015>. reprint of Frobenius' 1878 paper.
- Jürgen Fuchs. *Affine Lie Algebras and Quantum Groups: An Introduction, With Applications in Conformal Field Theory*. Cambridge Monographs on Mathematical Physics. Cambridge University Press, 1995. ISBN 052148412X. URL <http://books.google.com/books?vid=ISBN052148412X>.

- Dennis Gabor. Theory of communication. *Journal of the Institution of Electrical Engineers*, 93(26): 429–457, November 1946. URL <http://bigwww.epfl.ch/chaudhury/gabor.pdf>.
- Carl Friedrich Gauss. *Carl Friedrich Gauss Werke*, volume 8. Königlichen Gesellschaft der Wissenschaften, B.G. Teubner In Leipzig, Göttingen, 1900. URL <http://gdz.sub.uni-goettingen.de/dms/load/img/?PPN=PPN236010751>.
- Israel M. Gelfand. Normierte ringe. *Mat. Sbornik*, 9(51):3–24, 1941.
- Israel M. Gelfand and Mark A. Naimark. Normed rings with an involution and their representations. In *Commutative Normed Rings*, number 170 in AMS Chelsea Publishing Series, pages 240–274. Chelsea Publishing Company, Bronx, 1964. ISBN 9780821820223. URL <http://books.google.com/books?vid=ISBN0821820222>.
- Israel M. Gelfand and Mark A. Neumark. On the imbedding of normed rings into the ring of operators in hilbert space. *Mat. Sbornik*, 12(54:2):197–217, 1943a.
- Israel M. Gelfand and Mark A. Neumark. On the imbedding of normed rings into the ring of operators in hilbert space. In Robert S. Doran, editor, *C\*-algebras: 1943–1993: a Fifty Year Celebration: Ams Special Session Commemorating the First Fifty Years of C\*-Algebra Theory January 13–14, 1993*, pages 3–19. 1943b. ISBN 978-0821851753. URL <http://books.google.com/books?vid=ISBN0821851756>.
- Israel M. Gelfand, R. A. Minlos, and Z. Ya. Shapiro. *Representations of the rotation and Lorentz groups and their applications*. Courier Dover Publications, reprint edition, 2018. ISBN 9780486823850.
- Izrail' Moiseevich Gel'fand. *Representations of the rotation and Lorentz groups and their applications*. Pergamon Press book, 1963. 2018 Dover edition available.
- John Robilliard Giles. *Introduction to the Analysis of Normed Linear Spaces*. Number 13 in Australian Mathematical Society lecture series. Cambridge University Press, Cambridge, 2000. ISBN 0-521-65375-4. URL <http://books.google.com/books?vid=ISBN0521653754>.
- Israel Gohberg, Seymour Goldberg, and Marinus A. Kaashoek. *Basic Classes of Linear Operators*. Birkhäuser, Basel, 1 edition, 2003. ISBN 3764369302. URL <http://books.google.com/books?vid=ISBN3764369302>.
- T. N. T. Goodman, S. L. Lee, and W. S. Tang. Wavelets in wandering subspaces. *Transactions of the A.M.S.*, 338(2):639–654, August 1993a. URL <http://www.jstor.org/stable/2154421>. Transactions of the American Mathematical Society.
- T. N. T. Goodman, S. L. Lee, and W. S. Tang. Wavelets in wandering subspaces. *Advances in Computational Mathematics* 1, pages 109–126, February 1993b.
- Jaideva C. Goswami and Andrew K. Chan. *Fundamentals of Wavelets; Theory, Algorithms, and Applications*. John Wiley & Sons, Inc., 1999. ISBN 0-471-19748-3. URL [http://vadkudr.boom.ru/Collection/fundwave\\_contents.html](http://vadkudr.boom.ru/Collection/fundwave_contents.html).
- Robert M. Gray. Toeplitz and circulant matrices: A review. Technical Report AD0727139, Stanford University California Stanford Electronics Labs, Norwell, Massachusetts, June 1971. URL <https://apps.dtic.mil/docs/citations/AD0727139>.
- Robert M. Gray. Toeplitz and circulant matrices: A review. *Foundations and Trends® in Communications and Information Theory*, 2(3):155–239, January 31 2006. doi: <http://dx.doi.org/10.1561/01000000006>. URL <https://ee.stanford.edu/~gray/toeplitz.pdf>.



- Daniel J. Greenhoe. *Wavelet Structure and Design*, volume 3 of *Mathematical Structure and Design series*. Abstract Space Publishing, August 2013. ISBN 9780983801139. URL <http://books.google.com/books?vid=ISBN0983801134>. revised online version available at <https://www.researchgate.net/publication/312529555>.
- Daniel J. Greenhoe. The effects of the assorted cross-correlation definitions. *ResearchGate.com*, June 2019a. doi: 10.13140/RG.2.2.15881.70240. URL <https://www.researchgate.net/publication/333882235>.
- Daniel J. Greenhoe. *A Book Concerning Digital Communications*, volume 4 of *Signal Processing ABCs*. 2019b. URL <https://github.com/dgreenhoe/pdfs/blob/master/abcdc.pdf>.
- Ulf Grenander and Gábor Szegő. *Toeplitz Forms and Their Applications*. California monographs in mathematical sciences. University of California Press, 1958. URL <https://www.worldcat.org/title/toeplitz-forms-and-their-applications/oclc/648601989>.
- Ulf Grenander and Gábor Szegő. *Toeplitz Forms and Their Applications*. Chelsea Publishing Company, 2 edition, 1984. ISBN 9780828403214. URL <http://books.google.com/books?vid=ISBN9780828403214>.
- Frédéric Guichard, Jean-Michel Morel, and Robert Ryan. Contrast invariant image analysis and pde's. 2012. URL <http://dev.ipol.im/~morel/JMMBook2012.pdf>.
- Ernst Adolph Guillemin. *Synthesis of Passive Networks: Theory and Methods Appropriate to the Realization and Approximation Problems*. John Wiley & Sons, 1957. ISBN 9780882754819. URL <http://books.google.com.tw/books?id=JQ4nAAAAMAAJ>.
- Amritava Gupta. *Real & Abstract Analysis*. Academic Publishers Calcutta, 1998. ISBN 9788186358443. URL <http://books.google.com/books?vid=ISBN8186358447>. first published in 1998, reprinted in 2000 and 2006.
- Norman B. Haaser and Joseph A. Sullivan. *Real Analysis*. Dover Publications, New York, 1991. ISBN 0-486-66509-7. URL <http://books.google.com/books?vid=ISBN0486665097>.
- Paul R. Haddad and Ali N. Akansu. *Multiresolution Signal Decomposition: Transforms, Subbands, and Wavelets*. Acedemic Press, October 1 1992. ISBN 0323138365. URL <http://books.google.com/books?vid=ISBN0323138365>.
- Paul R. Halmos. *Finite Dimensional Vector Spaces*. Princeton University Press, Princeton, 1 edition, 1948. ISBN 0691090955. URL <http://books.google.com/books?vid=isbn0691090955>.
- Paul R. Halmos. *Finite Dimensional Vector Spaces*. Springer-Verlag, New York, 2 edition, 1958. ISBN 0-387-90093-4. URL <http://books.google.com/books?vid=isbn0387900934>.
- Paul R. Halmos. *Intoduction to Hilbert Space and the Theory of Spectral Multiplicity*. Chelsea Publishing Company, New York, 2 edition, 1998. ISBN 0821813781. URL <http://books.google.com/books?vid=ISBN0821813781>.
- Georg Hamel. Eine basis aller zahlen und die unstetigen lösungen der funktionalgleichung  $f(x + y) = f(x) + f(y)$ . *Mathematische Annalen*, 60(3):459–462, 1905. URL <http://gdz.sub.uni-goettingen.de/dms/load/img/?PPN=GDZPPN002260395&IDDOC=28580>.
- Deguang Han, Keri Kornelson, David Larson, and Eric Weber. *Frames for Undergraduates*, volume 40 of *Student Mathematical Library*. American Mathematical Society, 2007. ISBN 0821842129. URL <http://books.google.com/books?vid=ISBN0821842129>. Deguang Han = ???

- Godfrey H. Hardy. *A Mathematician's Apology*. Cambridge University Press, Cambridge, 1940. URL <http://www.math.ualberta.ca/~mss/misc/A%20Mathematician's%20Apology.pdf>.
- Felix Hausdorff. *Set Theory*. Chelsea Publishing Company, New York, 3 edition, 1937. ISBN 0828401195. URL <http://books.google.com/books?vid=ISBN0828401195>. 1957 translation of the 1937 German *Grundzüge der Mengenlehre*.
- S. Haykin and S. Kesler. Prediction-error filtering and maximum-entropy spectral estimation. In S. Haykin, editor, *Nonlinear Methods of Spectral Analysis*, volume 34 of *Topics in Applied Physics*, pages 9–72. Springer-Verlag, 1 edition, 1979.
- S. Haykin and S. Kesler. Prediction-error filtering and maximum-entropy spectral estimation. In S. Haykin, editor, *Nonlinear Methods of Spectral Analysis*, volume 34 of *Topics in Applied Physics*, pages 9–72. Springer-Verlag, “second corrected and updated edition” edition, 1983.
- Michiel Hazewinkel, editor. *Handbook of Algebras*, volume 2. North-Holland, Amsterdam, 1 edition, 2000. ISBN 044450396X. URL <http://books.google.com/books?vid=ISBN044450396X>.
- Jean Van Heijenoort. *From Frege to Gödel: A Source Book in Mathematical Logic, 1879-1931*. Harvard University Press, Cambridge, Massachusetts, 1967. URL <http://www.hup.harvard.edu/catalog/VANFGX.html>.
- Christopher Heil. *A Basis Theory Primer*. Applied and Numerical Harmonic Analysis. Birkhäuser, Boston, expanded edition edition, 2011. ISBN 9780817646868. URL <http://books.google.com/books?vid=ISBN9780817646868>.
- Christopher E. Heil and David F. Walnut. Continuous and discrete wavelet transforms. *Society for Industrial and Applied Mathematics*, 31(4), December 1989. URL <http://citeseer.ist.psu.edu/viewdoc/download?doi=10.1.1.132.1241&rep=rep1&type=pdf>.
- Charles Hermite. Lettre à Stieltjes. In B. Baillaud and H. Bourget, editors, *Correspondance d'Hermite et de Stieltjes*, volume 2, pages 317–319. Gauthier-Villars, Paris, May 20 1893. published in 1905.
- Eugenio Hernández and Guido Weiss. *A First Course on Wavelets*. CRC Press, New York, 1996. ISBN 0849382742. URL <http://books.google.com/books?vid=ISBN0849382742>.
- John Rowland Higgins. *Sampling Theory in Fourier and Signal Analysis: Foundations*. Oxford Science Publications. Oxford University Press, August 1 1996. ISBN 9780198596998. URL <http://books.google.com/books?vid=ISBN0198596998>.
- David Hilbert. Grundzüge einer allgemeinen theorie der linearen integralgleichungen (fundamentals of a general theory of the linear integral equations). *Mathematisch-Physikalische Klasse (Mathematical-Physical Class)*, pages 49–91, March 1904. URL <http://dz-srv1.sub.uni-goettingen.de/sub/digbib/loader?did=D57552>. Report 1 of 6.
- David Hilbert. Grundzüge einer allgemeinen theorie der linearen integralgleichungen (fundamentals of a general theory of the linear integral equations). *Mathematisch-Physikalische Klasse (Mathematical-Physical Class)*, pages 157–228, March 1906. URL <http://dz-srv1.sub.uni-goettingen.de/sub/digbib/loader?did=D58133>. Report 4 of 6.
- David Hilbert. *Grundzüge einer allgemeinen Theorie der linearen Integralgleichungen (Fundamentals of a general theory of the linear integral equations)*, volume 26 of *Fortschritte der mathematischen Wissenschaften in Monographien (Progress of the mathematical sciences in Monographs)*. B.G. Teubner, Leipzig und Berlin, 1912. URL <http://www.worldcat.org/oclc/13468199>.

- David Hilbert, Lothar Nordheim, and John von Neumann. über die grundlagen der quantenmechanik (on the bases of quantum mechanics). *Mathematische Annalen*, 98:1–30, 1927. ISSN 0025-5831 (print) 1432-1807 (online). URL <http://dz-srv1.sub.uni-goettingen.de/cache/toc/D27776.html>.
- Roger A. Horn and Charles R. Johnson. *Matrix Analysis*. Cambridge University Press, Cambridge, 1990. ISBN 0-521-30586-1. URL <http://books.google.com/books?vid=isbn0521305861>. Library: QA188H66 1985.
- Alfred Edward Housman. *More Poems*. Alfred A. Knopf, 1936. URL <http://books.google.com/books?id=rTMiAAAAMAAJ>.
- Satoru Igari. *Real Analysis—With an Introduction to Wavelet Theory*, volume 177 of *Translations of mathematical monographs*. American Mathematical Society, 1996. ISBN 9780821821046. URL <http://books.google.com/books?vid=ISBN0821821040>.
- Taizo Iijima. Basic theory of pattern observation. *Papers of Technical Group on Automata and Automatic Control*, December 1959. see Weickert 1999 for historical information.
- Chris J. Isham. *Modern Differential Geometry for Physicists*. World Scientific Publishing, New Jersey, 2 edition, 1999. ISBN 9810235623. URL <http://books.google.com/books?vid=ISBN9810235623>.
- C.J. Isham. Quantum topology and quantisation on the lattice of topologies. *Classical and Quantum Gravity*, 6:1509–1534, November 1989. doi: 10.1088/0264-9381/6/11/007. URL <http://www.iop.org/EJ/abstract/0264-9381/6/11/007>.
- Vasile I. Istrăţescu. *Inner Product Structures: Theory and Applications*. Mathematics and Its Applications. D. Reidel Publishing Company, 1987. ISBN 9789027721822. URL <http://books.google.com/books?vid=ISBN9027721823>.
- A. J. E. M. Janssen. The zak transform: A signal transform for sampled time-continuous signals. *Philips Journal of Research*, 43(1):23–69, 1988.
- Bjorn Jawerth and Wim Sweldens. An overview of wavelet based multiresolutional analysis. *SIAM Review*, 36:377–412, September 1994. URL <http://cm.bell-labs.com/who/wim/papers/papers.html#overview>.
- Alan Jeffrey and Hui Hui Dai. *Handbook of Mathematical Formulas and Integrals*. Handbook of Mathematical Formulas and Integrals Series. Academic Press, 4 edition, January 18 2008. ISBN 9780080556840. URL <http://books.google.com/books?vid=ISBN0080556841>.
- Palle E. T. Jørgensen, Kathy Merrill, Judith Packer, and Lawrence W. Baggett. *Representations, Wavelets and Frames: A Celebration of the Mathematical Work of Lawrence Baggett*. Applied and Numerical Harmonic Analysis. Birkhäuser, 2008. ISBN 9780817646820. URL <http://books.google.com/books?vid=ISBN0817646825>.
- K. D. Joshi. *Applied Discrete Structures*. New Age International, New Delhi, 1997. ISBN 8122408265. URL <http://books.google.com/books?vid=ISBN8122408265>.
- J.S.Chitode. *Signals And Systems*. Technical Publications, 2009. ISBN 9788184316780. URL <http://books.google.com/books?vid=ISBN818431678X>.
- Jean-Pierre Kahane. Partial differential equations, trigonometric series, and the concept of function around 1800: a brief story about lagrange and fourier. In Dorina Mitrea and V. G. Mazía, editors, *Perspectives In Partial Differential Equations, Harmonic Analysis And Applications: a*

*Volume in Honor of Vladimir G. Maz'ya's 70th birthday*, volume 79 of *Proceedings of Symposia in Pure Mathematics*, pages 187–206. American Mathematical Society, 2008. ISBN 0821844245. URL <http://books.google.com/books?vid=ISBN0821844245>.

David W. Kammler. *A First Course in Fourier Analysis*. Cambridge University Press, 2 edition, 2008. ISBN 9780521883405. URL <http://books.google.com/books?vid=ISBN0521883407>.

Edward Kasner and James Roy Newman. *Mathematics and the Imagination*. Simon and Schuster, 1940. ISBN 0486417034. URL <http://books.google.com/books?vid=ISBN0486417034>. “unabridged and unaltered republication” available from Dover.

Yitzhak Katznelson. *An Introduction to Harmonic Analysis*. Cambridge mathematical library. Cambridge University Press, 3 edition, 2004. ISBN 0521543592. URL <http://books.google.com/books?vid=ISBN0521543592>.

James P. Keener. *Principles of Applied Mathematics; Transformation and Approximation*. Addison-Wesley Publishing Company, Reading, Massachusetts, 1988. ISBN 0201156741. URL <http://www.worldcat.org/isbn/0201156741>.

Yoshida Kenko. *The Tsuredzure Gusa of Yoshida No Kaneyoshi. Being the meditations of a recluse in the 14th Century (Essays in Idleness)*. circa 1330. URL <http://www.humanistictexts.org/kenko.htm>. 1911 translation of circa 1330 text.

Anthony W Knapp. *Advanced Real Analysis*. Cornerstones. Birkhäuser, Boston, Massachusetts, USA, 1 edition, July 29 2005a. ISBN 0817643826. URL <http://books.google.com/books?vid=ISBN0817643826>.

Anthony W Knapp. *Basic Real Analysis*. Cornerstones. Birkhäuser, Boston, Massachusetts, USA, 1 edition, July 29 2005b. ISBN 0817632506. URL <http://books.google.com/books?vid=ISBN0817632506>.

Carlos S. Kubrusly. *The Elements of Operator Theory*. Springer, 1 edition, 2001. ISBN 9780817641740. URL <http://books.google.com/books?vid=ISBN0817641742>.

Carlos S. Kubrusly. *The Elements of Operator Theory*. Springer, 2 edition, 2011. ISBN 9780817649975. URL <http://books.google.com/books?vid=ISBN0817649972>.

Andrew Kurdila and Michael Zabaranin. *Convex Functional Analysis*. Systems & Control: Foundations & Applications. Birkhäuser, Boston, 2005. ISBN 9783764321987. URL <http://books.google.com/books?vid=ISBN3764321989>.

Joseph-Louis Lagrange, Pierre-Simon Laplace, Étienne Louis Malus, René Just Haüy, and Adrien-Marie Legendre. Proclamation des prix décernés dans la séance publique de 6 janvier 1812. *Esprit des Journaux, Français et étrangers par Une Société de Gens Delettres*, 2:111–112, January 6 1812a. URL <http://books.google.com/books?id=QpUUAQAAAJ>.

Joseph-Louis Lagrange, Pierre-Simon Laplace, Étienne Louis Malus, René Just Haüy, and Adrien-Marie Legendre. Proclamation des prix décernés dans la séance publique de 6 janvier 1812. *Mercure De France, Journal Littéraire et Politique*, 50:374–375, January 6 1812b. URL <http://books.google.com/books?id=8HxBAAAAcAAJ>.

Imre Lakatos. *Proofs and Refutations: The Logic of Mathematical Discovery*. Cambridge University Press, Cambridge, 1976. ISBN 0521290384. URL <http://books.google.com/books?vid=ISBN0521290384>.



- Traian Lalescu. *Sur les équations de Volterra*. PhD thesis, University of Paris, 1908. advisor was Émile Picard.
- Traian Lalescu. *Introduction à la théorie des équations intégrales (Introduction to the Theory of Integral Equations)*. Librairie Scientifique A. Hermann, Paris, 1911. URL <http://www.worldcat.org/oclc/1278521>. first book about integral equations ever published.
- Rupert Lasser. *Introduction to Fourier Series*, volume 199 of *Monographs and textbooks in pure and applied mathematics*. Marcel Dekker, New York, New York, USA, February 8 1996. ISBN 978-0824796105. URL <http://books.google.com/books?vid=ISBN0824796101>. QA404.L33 1996.
- Peter D. Lax. *Functional Analysis*. John Wiley & Sons Inc., USA, 2002. ISBN 0-471-55604-1. URL <http://www.worldcat.org/isbn/0471556041>. QA320.L345 2002.
- Gottfried W. Leibniz. Symbolismus memorabilis calculi algebraici et infinitesimalis, in comparatione potentiarum et differentiarum; et de lege homogeneorum transcendentali. *Miscellanea Berolinensia ad incrementum scientiarum, ex scriptis Societati Regiae scientiarum*, pages 160–165, 1710. URL [http://bibliothek.bbaw.de/bibliothek-digital/digitalequellen/schriften/anzeige/index\\_html?band=01-misc/1&seite:int=184](http://bibliothek.bbaw.de/bibliothek-digital/digitalequellen/schriften/anzeige/index_html?band=01-misc/1&seite:int=184).
- Gottfried Wilhelm Leibniz. Letter to christian huygens, 1679. In Leroy E. Loemker, editor, *Philosophical Papers and Letters*, volume 2 of *The New Synthese Historical Library*, chapter 27, pages 248–249. Kluwer Academic Press, Dordrecht, 2 edition, September 8 1679. ISBN 902770693X. URL <http://books.google.com/books?vid=ISBN902770693X>.
- Pierre Gilles Lemarié, editor. *Les Ondelettes en 1989*, volume 1438 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1990. ISBN 978-3-540-52932-3. URL <http://link.springer.com/book/10.1007/BFb0083510/>.
- Tony Lindeberg. *Scale-Space Theory in Computer Vision*. The Springer International Series in Engineering and Computer Science. Springer, 1993. ISBN 9780792394181. URL <http://books.google.com/books?vid=ISBN0792394186>.
- Joram Lindenstrauss and Lior Tzafriri. *Classical Banach Spaces I: Sequence Spaces*, volume 92 of *Ergebnisse der Mathematik und ihrer Grenzgebiete*. Springer-Verlag, 1977. ISBN 978-3642665592. URL <http://www.amazon.com/dp/3642665594>.
- J. Liouville. Sur l'integration d'une classe d'équations différentielles du second ordre en quantités finies explicites. *Journal De Mathematiques Pures Et Appliquees*, 4:423–456, 1839. URL <http://gallica.bnf.fr/ark:/12148/bpt6k16383z>.
- Lynn H. Loomis and Ethan D. Bolker. *Harmonic analysis*. Mathematical Association of America, 1965. URL <http://books.google.com/books?id=MEfvAAAAMAAJ>.
- Nikolai Luzin. Sur la convergence des séries trigonom etriers de fourier. *C. R. Acad. Sci.*, 156:1655–1658, 1913.
- Niccolò Machiavelli. *The Literary Works of Machiavelli: Mandragola, Clizia, A Dialogue on Language, and Belfagor, with Selections from the Private Correspondence*. Oxford University Press, 1961. ISBN 0313212481. URL <http://www.worldcat.org/isbn/0313212481>.
- Colin Maclaurin. *Treatise of Fluxions*. W. Baynes, 1742. URL <http://www.amazon.com/dp/B000863E7M>.

- Stéphane G. Mallat. Multiresolution approximations and wavelet orthonormal bases of  $l^2(\mathbb{R})$ . *Transactions of the American Mathematical Society*, 315(1):69–87, September 1989. URL [http://blanche.polytechnique.fr/~mallat/papiers/math\\_multiresolution.pdf](http://blanche.polytechnique.fr/~mallat/papiers/math_multiresolution.pdf).
- Stéphane G. Mallat. *A Wavelet Tour of Signal Processing*. Elsevier, 2 edition, September 15 1999. ISBN 9780124666061. URL <http://books.google.com/books?vid=ISBN012466606X>.
- MatLab. cpsd: Cross power spectral density. *MatLab*, accessed 2018 November 26 2018a. URL <https://www.mathworks.com/help/signal/ref/cpsd.html>.
- MatLab. xcorr: Cross-correlation. *MatLab*, accessed 2018 November 26 2018b. URL <https://www.mathworks.com/help/signal/ref/xcorr.html>.
- maxima. *Maxima Manual version 5.28.0*. 5.28.0 edition. URL <http://maxima.sourceforge.net/documentation.html>.
- Stefan Mazur. Sur les anneaux linéaires. *Comptes rendus de l'Académie des sciences*, 207:1025–1027, 1938.
- Stefan Mazur and Stanislaus M. Ulam. Sur les transformations isométriques d'espaces vectoriels normés. *Comptes rendus de l'Académie des sciences*, 194:946–948, 1932.
- James Mercer. Functions of positive and negative type and their connection with the theory of integral equations. *Philosophical Transactions of the Royal Society of London*, 209:415–446, 1909. ISSN 02643952. URL <http://www.jstor.org/stable/91043>.
- Anthony N. Michel and Charles J. Herget. *Applied Algebra and Functional Analysis*. Dover Publications, Inc., 1993. ISBN 048667598X. URL <http://books.google.com/books?vid=ISBN048667598X>. original version published by Prentice-Hall in 1981.
- Fred Mintzer. Filters for distortion-free two-band multi-rate filter banks. *IEEE Transactions on Acoustics, Speech and Signal Processing*, 32, 1985.
- Eddie Ortiz Muniz. A method for deriving various formulas in electrostatics and electromagnetism using lagrange's trigonometric identities. *American Journal of Physics*, 21(140), 1953. doi: 10.1119/1.1933371. URL <http://dx.doi.org/10.1119/1.1933371>.
- Ben Noble and James W. Daniel. *Applied Linear Algebra*. Prentice-Hall, Englewood Cliffs, NJ, USA, 3 edition, 1988. ISBN 0130412600. URL <http://www.worldcat.org/isbn/0130412600>.
- Timur Oikhberg and Haskell Rosenthal. A metric characterization of normed linear spaces. *Rocky Mountain Journal Of Mathematics*, 37(2):597–608, 2007. URL <http://www.ma.utexas.edu/users/rosenthl/pdf-papers/95-oikh.pdf>.
- Judith Packer. Applications of the work of stone and von neumann to wavelets. In Robert S. Doran and Richard V. Kadison, editors, *Operator Algebras, Quantization, and Noncommutative Geometry: A Centennial Celebration Honoring John Von Neumann and Marshall H. Stone : AMS Special Session on Operator Algebras, Quantization, and Noncommutative Geometry, a Centennial Celebration Honoring John Von Neumann and Marshall H. Stone, January 15-16, 2003, Baltimore, Maryland*, volume 365 of *Contemporary mathematics—American Mathematical Society*, pages 253–280, Baltimore, Maryland, 2004. American Mathematical Society. ISBN 9780821834022. URL <http://books.google.com/books?vid=isbn0821834029>.
- Lincoln P. Paine. *Warships of the World to 1900*. Ships of the World Series. Houghton Mifflin Harcourt, 2000. ISBN 9780395984147. URL <http://books.google.com/books?vid=ISBN9780395984149>.

- Anthanasios Papoulis. *Probability, Random Variables, and Stochastic Processes*. McGraw-Hill Series in Electrical Engineering. McGraw-Hill Book Company, New York, 2 edition, 1984. ISBN 9780070484689. URL <http://books.google.com/books?vid=ISBN0070484686>.
- Anthanasios Papoulis. *Probability, Random Variables, and Stochastic Processes*. McGraw-Hill, New York, 3 edition, 1991. ISBN 0070484775. URL <http://books.google.com/books?vid=ISBN0070484775>.
- Giuseppe Peano. *Calcolo geometrico secondo l'Ausdehnungslehre di H. Grassmann preceduto dalle operazioni della logica deduttiva*. Fratelli Bocca Editori, Torino, 1888a. Geometric Calculus: According to the *Ausdehnungslehre* of H. Grassmann.
- Giuseppe Peano. *Geometric Calculus: According to the Ausdehnungslehre of H. Grassmann*. Springer (2000), 1888b. ISBN 0817641262. URL <http://books.google.com/books?vid=isbn0817641262>. originally published in 1888 in Italian.
- Michael Pedersen. *Functional Analysis in Applied Mathematics and Engineering*. Chapman & Hall/CRC, New York, 2000. ISBN 9780849371691. URL <http://books.google.com/books?vid=ISBN0849371694>. Library QA320.P394 1999.
- Wesley J. Perschbacher, editor. *The New Analytical Greek Lexicon*. Hendrickson Publishers, Peabody, Mass., 1990. ISBN 978-0-943575-33-9. URL <http://www.amazon.com/dp/0943575338>.
- Mark A. Pinsky. *Introduction to Fourier Analysis and Wavelets*. Brooks/Cole, Pacific Grove, 2002. ISBN 0-534-37660-6. URL <http://www.amazon.com/dp/0534376606>.
- Lakshman Prasad and Sundararaja S. Iyengar. *Wavelet Analysis with Applications to Image Processing*. CRC Press LLC, Boca Raton, 1997. ISBN 978-0849331695. URL <http://books.google.com/books?vid=ISBN0849331692>. Library TA1637.P7 1997.
- Ptolemy. *Ptolemy's Almagest*. Springer-Verlag (1984), New York, circa 100AD. ISBN 0387912207. URL <http://gallica.bnf.fr/ark:/12148/bpt6k3974x>.
- Shie Qian and Dapang Chen. *Joint time-frequency analysis: methods and applications*. PTR Prentice Hall, 1996. ISBN 9780132543842. URL <http://books.google.com/books?vid=ISBN0132543842>.
- Charles Earl Rickart. *General Theory of Banach Algebras*. University series in higher mathematics. D. Van Nostrand Company, Yale University, 1960. URL <http://books.google.com/books?id=PVrvAAAAAAAJ>.
- Theodore J. Rivlin. *The Chebyshev Polynomials*. Pure and Applied Mathematics: A Wiley-Interscience Series of Texts, Monographs and Tracts. John Wiley & Sons, New York, 1974. ISBN 0-471-72470-X. URL <http://books.google.com/books?vid=ISBN047172470X>.
- Enders A. Robinson. *Random Wavelets and Cybernetic Systems*, volume 9 of *Griffins Statistical Monographs & Courses*. Lubrecht & Cramer Limited, London, June 1962. ISBN 0852640757. URL <http://books.google.com/books?vid=ISBN0852640757>.
- Enders A. Robinson. Multichannel z-transforms and minimum delay. *Geophysics*, 31(3):482–500, June 1966. doi: 10.1190/1.1439788. URL <http://dx.doi.org/10.1190/1.1439788>.
- Enders A. Robinson. A historical perspective of spectrum estimation. *Proceedings of the IEEE*, 70(9):885–907, September 1982. doi: 10.1109/PROC.1982.12423. URL <http://www.archive.ece.cmu.edu/~ece792/handouts/Robinson82.pdf>.

- Maxwell Rosenlicht. *Introduction to Analysis*. Dover Publications, New York, 1968. ISBN 0-486-65038-3. URL <http://books.google.com/books?vid=ISBN0486650383>.
- Walter Rudin. *Real and Complex Analysis*. McGraw-Hill Book Company, New York, New York, USA, 3 edition, 1987. ISBN 9780070542341. URL <http://www.amazon.com/dp/0070542341>. Library QA300.R8 1976.
- Walter Rudin. *Functional Analysis*. McGraw-Hill, New York, 2 edition, 1991. ISBN 0-07-118845-2. URL <http://www.worldcat.org/isbn/0070542252>. Library QA320.R83 1991.
- Bryan P. Rynne and Martin A. Youngson. *Linear Functional Analysis*. Springer undergraduate mathematics series. Springer, 2 edition, January 1 2008. ISBN 9781848000056. URL <http://books.google.com/books?vid=ISBN1848000057>.
- Jr. S. Lawrence Marple. *Digital Spectral Analysis with Applications*. Prentice Hall, 1 edition, 1987. ISBN 9780132141499. URL <http://books.google.com/books?vid=ISBN9780132141499>.
- Jr. S. Lawrence Marple. *Digital Spectral Analysis*. Dover Books on Electrical Engineering. Courier Dover Publications, 2019. ISBN 9780486780528. URL <http://books.google.com/books?vid=ISBN9780486780528>.
- Shôichirô Sakai. *C\*-Algebras and W\*-Algebras*. Springer-Verlag, Berlin, 1 edition, 1998. ISBN 9783540636335. URL <http://books.google.com/books?vid=ISBN3540636331>. reprint of 1971 edition.
- Juljusz Schauder. Zur theorie stetiger abbildungen in funktionalräumen. *Mathematische Zeitschrift*, 26:47–65, 1927. URL <http://eudml.org/doc/167911;jsessionid=156A34EBAB6C0E2DDAAC8C1232D23E8F>.
- Juljusz Schauder. Eine eigenschaft des haarschen orthogonalsystems. *Mathematische Zeitschrift*, 28:317–320, 1928.
- Gert Schubring. *Conflicts Between Generalization, Rigor, and Intuition: Number Concepts Underlying the Development of Analysis in 17th–19th Century France and Germany*. Sources and studies in the history of mathematics and physical sciences. Springer, New York, 1 edition, June 2005. ISBN 0387228365. URL <http://books.google.com/books?vid=ISBN0387228365>.
- Isaac Schur. Über die charakterischen wurzeln einer linearen substitution mit enier anwendung auf die theorie der integralgleichungen (over the characteristic roots of one linear substitution with an application to the theory of the integral). *Mathematische Annalen*, 66:488–510, 1909. URL <http://dz-srv1.sub.uni-goettingen.de/cache/toc/D38231.html>.
- Mícheál Ó Searcóid. *Elements of Abstract Analysis*. Springer Undergraduate Mathematics Series. Springer, 2002. ISBN 9781852334246. URL <http://books.google.com/books?vid=ISBN185233424X>.
- Atle Selberg. Harmonic analysis and discontinuous groups in weakly symmetric riemannian spaces with applications to dirichlet series. *Journal of the Indian Mathematical Society*, 20:47–87, 1956.
- George Finlay Simmons. *Calculus Gems: Brief Lives and Memorable Mathematicians*. Mathematical Association of America, Washington DC, 2007. ISBN 0883855615. URL <http://books.google.com/books?vid=ISBN0883855615>.
- Iván Singer. *Bases in Banach Spaces I*, volume 154 of *Grundlehren der mathematischen Wissenschaften in Einzeldarstellungen mit besonderer Berücksichtigung der Anwendungsgebiete*. Springer-Verlag New York, New York, 1970. ISBN 9780387048338.



- Neil J. A. Sloane. On-line encyclopedia of integer sequences. World Wide Web, 2014. URL <http://oeis.org/>.
- M.J.T. Smith and T.P. Barnwell. A procedure for designing exact reconstruction filter banks for tree-structured subband coders. *IEEE International Conference on Acoustics, Speech and Signal Processing*, 9:421–424, 1984a. T.P. Barnwell is T.P. Barnwell III.
- M.J.T. Smith and T.P. Barnwell. The design of digital filters for exact reconstruction in subband coding. *IEEE Transactions on Acoustics, Speech and Signal Processing*, 34(3):434–441, June 1984b. ISSN 0096-3518. doi: 10.1109/TASSP.1986.1164832. T.P. Barnwell is T.P. Barnwell III.
- D. E. Smylie, G. K. C. Clarke, and T. J. Ulrych. Analysis of irregularities in the earth's rotation. In Bruce Bolt, editor, *Methods in Computational Physics: Advances in Research and Applications*, volume 13 of *Geophysics*, pages 391–340. Academic Press, 1973. ISBN 9780323148368. URL <http://books.google.com/books?vid=ISBN9780323148368>.
- Houshang H. Sohrab. *Basic Real Analysis*. Birkhäuser, Boston, 1 edition, 2003. ISBN 978-0817642112. URL <http://books.google.com/books?vid=ISBN0817642110>.
- Lynn Arthur Steen. Highlights in the history of spectral theory. *The American Mathematical Monthly*, 80(4):359–381, April 1973. ISSN 00029890. URL <http://www.jstor.org/stable/2319079>.
- Anne K. Steiner. The lattice of topologies: Structure and complementation. *Transactions of the American Mathematical Society*, 122(2):379–398, April 1966. URL <http://www.jstor.org/stable/1994555>.
- Marshall Harvey Stone. *Linear transformations in Hilbert space and their applications to analysis*, volume 15 of *American Mathematical Society. Colloquium publications*. American Mathematical Society, New York, 1932. URL <http://books.google.com/books?vid=ISBN0821810154>. 1990 reprint of the original 1932 edition.
- Gilbert Strang and Truong Nguyen. *Wavelets and Filter Banks*. Wellesley-Cambridge Press, Wellesley, MA, 1996. ISBN 9780961408879. URL <http://books.google.com/books?vid=ISBN0961408871>.
- Robert S. Strichartz. *The Way of Analysis*. Jones and Bartlett Publishers, Boston-London, 1995. ISBN 978-0867204711. URL <http://books.google.com/books?vid=ISBN0867204710>.
- Endre Süli and David F. Mayers. *An Introduction to Numerical Analysis*. Cambridge University Press, August 28 2003. ISBN 9780521007948. URL <http://books.google.com/books?vid=ISBN0521007941>.
- Wim Sweldens and Robert Piessens. Wavelet sampling techniques. In *1993 Proceedings of the Statistical Computing Section*, pages 20–29. American Statistical Association, August 1993. URL <http://citeseer.ist.psu.edu/18531.html>.
- Erik Talvila. Necessary and sufficient conditions for differentiating under the integral sign. *The American Mathematical Monthly*, 108(6):544–548, June–July 2001. URL <http://arxiv.org/abs/math/0101012>.
- Brook Taylor. *Methodus Incrementorum Directa et Inversa*. London, 1715.
- Audrey Terras. *Fourier Analysis on Finite Groups and Applications*. Number 43 in London Mathematical Society Student Texts. Cambridge University Press, Cambridge, 1999. ISBN 0-521-45718-1. URL <http://books.google.com/books?vid=ISBN0521457181>.

- Brian S. Thomson, Andrew M. Bruckner, and Judith B. Bruckner. *Elementary Real Analysis*. [www.classicalrealanalysis.com](http://www.classicalrealanalysis.com), 2 edition, 2008. ISBN 9781434843678. URL <http://classicalrealanalysis.info/com/Elementary-Real-Analysis.php>.
- Stanislaw Marcin Ulam. *Adventures of a Mathematician*. University of California Press, Berkeley, 1991. ISBN 0520071549. URL <http://books.google.com/books?vid=ISBN0520071549>.
- P.P. Vaidyanathan. *Multirate Systems and Filter Banks*. Prentice Hall Signal Processing Series. Prentice Hall, 1993. ISBN 0136057187. URL <http://books.google.com/books?vid=ISBN0136057187>.
- Jussi Väisälä. A proof of the mazur-ulam theorem. *The American Mathematical Monthly*, 110(7): 633–635, August–September 2003. URL <http://www.helsinki.fi/~jvaisala/mazurulam.pdf>.
- Brani Vidakovic. *Statistical Modeling by Wavelets*. John Wiley & Sons, Inc, New York, 1999. ISBN 9780471293651. URL <http://www.amazon.com/dp/0471293652>.
- John von Neumann. Allgemeine eigenwerttheorie hermitescher funktionaloperatoren. *Mathematische Annalen*, 102(1):49–131, 1929. ISSN 0025-5831 (print) 1432-1807 (online). URL <http://resolver.sub.uni-goettingen.de/purl?GDZPPN002273535>. General eigenvalue theory of Hermitian functional operators.
- David F. Walnut. *An Introduction to Wavelet Analysis*. Applied and numerical harmonic analysis. Springer, 2002. ISBN 0817639624. URL <http://books.google.com/books?vid=ISBN0817639624>.
- Gilbert G. Walter and XiaoPing Shen. *Wavelets and Other Orthogonal Systems*. Chapman and Hall/CRC, New York, 2 edition, 2001. ISBN 9781584882275. URL <http://books.google.com/books?vid=ISBN1584882271>.
- Heinrich Martin Weber. Die allgemeinen grundlagen der galois'schen gleichungstheorie. *Mathematische Annalen*, 43(4):521–549, December 1893. URL <http://resolver.sub.uni-goettingen.de/purl?GDZPPN002254670>. The general foundation of Galois' equation theory.
- Joachim Weickert. Linear scale-space has first been proposed in japan. *Journal of Mathematical Imaging and Vision*, 10:237–252, May 1999. URL <http://dl.acm.org/citation.cfm?id=607668>.
- Hermann Weyl. The method of orthogonal projection in potential theory. *Duke Mathematical Journal*, 7(1):411–444, 1940. URL <http://projecteuclid.org/euclid.dmj/1077492266>.
- Harold Widom. Toeplitz matrices. In Jr. I. I. Hirschmann, editor, *Studies in Real and Complex Analysis*, volume 3 of *MAA Studies in Mathematics*, pages 179–209. Mathematical Association of America; distributed by Prentice-Hall, 1965. ISBN 9780883851036. URL <https://www.worldcat.org/title/studies-in-real-and-complex-analysis/oclc/506377>.
- Samuel Stanley Wilks. *Mathematical Statistics*. Wiley series in probability and mathematical statistics. Wiley, “second printing with corrections” edition, 1963. URL <https://books.google.com/books?id=-sJLAAAAMAAJ>.
- P. Wojtaszczyk. *A Mathematical Introduction to Wavelets*, volume 37 of *London Mathematical Society student texts*. Cambridge University Press, February 13 1997. ISBN 9780521578943. URL <http://books.google.com/books?vid=ISBN0521578949>.
- Frank Yates and Kenneth Mather. Ronald aylmer fisher, 1890–1962. *Biographical Memoirs of Fellows of the Royal Society of London*, 9:90–120, November 1963. URL <https://doi.org/10.1098/rsbm.1963.0006>. print ISSN:0080-4606, online ISSN:1748-8494, <https://pdfs.semanticscholar.org/6ccd/4303f68f3543fc794c376ce00395947a79b6.pdf>.

- Robert M. Young. *An introduction to nonharmonic Fourier series*, volume 93 of *Pure and applied mathematics*. Academic Press, 1 edition, 1980. ISBN 0127728503. URL <http://books.google.com/books?vid=ISBN0127728503>.
- Robert M. Young. *An introduction to nonharmonic Fourier series*, volume 93 of *Pure and applied mathematics*. Academic Press, revised first edition, May 16 2001. ISBN 0127729550. URL <http://books.google.com/books?vid=ISBN0127729550>.
- Ahmed I. Zayed. *Handbook of Function and Generalized Function Transformations*. Mathematical Sciences Reference Series. CRC Press, Boca Raton, 1996. ISBN 0849378516. URL <http://books.google.com/books?vid=ISBN0849378516>.
- Gary Zukav. *The Dancing Wu Li Masters : An Overview of the New Physics*. Bantam Books, New York, 1980. ISBN 055326382X. URL <http://books.google.com/books?vid=ISBN055326382X>.



## REFERENCE INDEX

- Devices (2016), 205  
 Abramovich and Aliprantis (2002), 28, 68  
 Aliprantis and Burkinshaw (1998), 121, 124, 125, 127–129, 132, 156, 158, 162, 188  
 Adelson and Burt (1981), 6  
 Ptolemy (circa 100AD), 85  
 Alvarez et al. (1993), 6  
 Andrews et al. (2001), 196  
 Abramowitz and Stegun (1972), 73, 74, 94, 97  
 Atkinson and Han (2009), 6  
 Autonne (1901), 143  
 Autonne (1902), 143  
 Bachman (1964), 12, 19  
 Bachman and Narici (1966), 135, 137, 149, 151, 153, 162  
 Bachman et al. (2000), 16, 154, 157, 167, 188  
 Banach (1922), 119, 124  
 Banach (1932b), 124  
 Banach (1932a), 124, 154  
 Ben-Israel and Gilbert (2002), 71  
 Beran (1976), 4  
 Beran (1985), 4  
 Berberian (1961), 117, 121–123, 140, 149, 162, 167  
 Bertero and Boccacci (1998), 137  
 Birkhoff and Neumann (1936), 4  
 Bollobás (1999), 129, 158, 209–211  
 Bottazzini (1986), 80, 81  
 Boyer and Merzbach (1991), 73  
 Bracewell (1978), 12, 19, 162  
 Bromwich (1908), 27  
 Bruckner et al. (1997), 102, 104  
 Burrus et al. (1998), 55  
 Burt and Adelson (1983), 6  
 Byrne (2005), 175  
 Cadzow (1987), 203, 204  
 Cardano (1545), 67  
 Carleson and Engquist (2001), 52  
 Carothers (2005), 151, 154  
 Casazza and Lammers (1998), 188  
 Cayley (1858), 209  
 Cerdà (2010), 109  
 J.S.Chitode (2009), 47  
 Chorin and Hald (2009), 16, 17  
 Christensen (2003), 149, 154, 162, 167, 168, 172, 175, 188, 190, 191  
 Christensen (2008), 168, 172  
 Chui (1992), 44, 70, 179  
 Claerbout (1976), 31  
 Clarkson (1993), 63  
 Cohn (2002), 68  
 Black et al. (2009), 3  
 Courant and Hilbert (1930), 212  
 Dai and Lu (1996), 188  
 Dai and Larson (1998), 188, 190  
 Daubechies (1992), 47, 155, 171  
 Davidson and Donsig (2010), 96  
 de Witt (1659), 211  
 Descartes (1954), 209  
 Descartes (1637a), 209  
 Descartes (1637b), 187  
 Daubechies et al. (1986), 172  
 Dieudonné (1969), 138  
 Duffin and Schaeffer (1952), 172  
 Dumitrescu (2007), 31  
 Dunford and Schwartz (1957), 131  
 Durbin (2000), 67, 68  
 Lagrange et al. (1812a), 51  
 Edwards (1995), 158  
 Eidelman et al. (2004), 122  
 Enflo (1973), 154  
 Euler (1748), 73, 80, 81, 89  
 Ewen (1950), viii  
 Ewen (1961), viii  
 Fabian et al. (2010), 160  
 Farina and Rinaldi (2000), 31  
 Fix and Strang (1969), 22  
 Flanders (1973), 71  
 Flanigan (1983), 10, 75  
 Folland (1995), 113, 114, 117, 118  
 Folland (1992), 78, 102, 196  
 Forster and Massopust (2009), 199  
 Fourier (1807), 51  
 Fourier (1878), 1, 15, 16  
 Fourier (1822), 16  
 Fréchet (1906), 1  
 Fréchet (1928), 1  
 Fredholm (1900), 201, 211  
 Fredholm (1903), 201, 211  
 Frobenius (1968), 138  
 Frobenius (1878), 138  
 Fuchs (1995), 113  
 Gabor (1946), 199  
 Gauss (1900), 197  
 Gelfand (1941), 117  
 Gelfand and Neumark (1943b), 116–118  
 Gelfand and Neumark (1943a), 117, 118  
 Gel'fand (1963), 145  
 Gelfand and Naimark (1964), 114, 115, 117  
 Gelfand et al. (2018), 145  
 Giles (2000), 131, 133, 154,

- 158  
 Gohberg et al. (2003), 212  
 Goodman et al. (1993b), 190  
 Goodman et al. (1993a), 188, 190  
 Goswami and Chan (1999), 21  
 Gray (1971), 63  
 Gray (2006), 63  
 Greenhoe (2013), 7  
 Greenhoe (2019a), 205  
 Grenander and Szegö (1958), 63  
 Grenander and Szegö (1984), 63  
 Guichard et al. (2012), 6  
 Guillemin (1957), 94  
 Gupta (1998), 153  
 Haaser and Sullivan (1991), 119, 132, 153  
 Haddad and Akansu (1992), 35  
 Halmos (1948), 119  
 Halmos (1958), 135  
 Halmos (1998), 115, 134  
 Hamel (1905), 151  
 Han et al. (2007), 160, 175  
 Hausdorff (1937), 188  
 Haykin and Kesler (1979), 63  
 Haykin and Kesler (1983), 63  
 Hazewinkel (2000), 113  
 Heijenoort (1967), viii  
 Heil and Walnut (1989), 188  
 Heil (2011), 102, 109, 120, 149, 151, 153, 154, 162, 167, 168, 172, 175, 188  
 Hermite (1893), 91  
 Hernández and Weiss (1996), 185  
 Higgins (1996), 198  
 Hilbert (1904), 209  
 Hilbert (1906), 210, 211  
 Hilbert (1912), 209, 211  
 Hilbert et al. (1927), 121  
 Horn and Johnson (1990), 125  
 Housman (1936), viii  
 Igari (1996), 170, 182, 185  
 Iijima (1959), 6  
 Isham (1989), 2  
 Isham (1999), 2  
 Istrătescu (1987), 109  
 Janssen (1988), 198  
 Jawerth and Sweldens (1994), 20, 22  
 Jeffrey and Dai (2008), 16, 105, 106  
 Jørgensen et al. (2008), 172  
 Joshi (1997), 27  
 Kahane (2008), 51  
 Kammler (2008), 188  
 Kasner and Newman (1940), 88  
 Katznelson (2004), 51, 102, 109, 111  
 Keener (1988), 60, 61, 137, 201, 211  
 Kenko (circa 1330), 214  
 Knapp (2005a), 144  
 Knapp (2005b), 16, 71, 196  
 Kubrusly (2001), 119, 121, 135, 140, 141, 149, 151, 153, 156, 162, 167  
 Kubrusly (2011), 27  
 Kurdila and Zabrankin (2005), 149  
 Lakatos (1976), 91  
 Lalescu (1908), 78  
 Lalescu (1911), 78  
 Lasser (1996), 196  
 Lax (2002), 137, 198  
 Leibniz (1710), 71  
 Leibniz (1679), 119  
 Lemarié (1990), 6  
 Lindeberg (1993), 6  
 Lindenstrauss and Tzafriri (1977), 154  
 Liouville (1839), 75, 78  
 Loomis and Bolker (1965), 16  
 Machiavelli (1961), 213  
 Maclaurin (1742), 10  
 Mallat (1989), 6, 171  
 Mallat (1999), 6, 21, 22, 35, 37, 49, 55, 89, 171  
 S. Lawrence Marple (1987), 63, 205  
 S. Lawrence Marple (2019), 63, 205  
 MatLab (2018a), 203  
 MatLab (2018b), 203  
 maxima, 96, 99  
 Mazur and Ulam (1932), 131  
 Mazur (1938), 117  
 Lagrange et al. (1812b), 51  
 Mercer (1909), 212  
 Michel and Herget (1993), 68, 115, 116, 120, 122, 124, 127, 132, 133, 136, 138, 141–143, 148, 149, 151, 153, 201, 209, 211, 212  
 Mintzer (1985), 35  
 Muniz (1953), 105, 106  
 Noble and Daniel (1988), 145  
 Oikhberg and Rosenthal (2007), 131  
 Packer (2004), 188  
 Paine (2000), vi  
 Papoulis (1984), 201, 203, 204  
 Papoulis (1991), 179, 201  
 Peano (1888b), 119, 132  
 Pedersen (2000), 78, 157, 158, 209  
 Perschbacher (1990), 3  
 Pinsky (2002), 100, 182, 185  
 de la Vallée-Poussin (1915), 188  
 Prasad and Iyengar (1997), 70  
 ?, 71  
 Qian and Chen (1996), 199  
 de Reyna (2002), 109  
 Rickart (1960), 114–117  
 Rivlin (1974), 91, 96  
 Robinson (1962), 31  
 Robinson (1966), 31  
 Robinson (1982), 10, 15  
 Rosenlicht (1968), 73, 75–79  
 Rudin (1991), 125, 127, 128, 130, 132–135, 137–139, 143, 144  
 Rudin (1987), 81, 196  
 Rynne and Youngson (2008), 153  
 Sakai (1998), 134  
 Schauder (1927), 154  
 Schauder (1928), 154  
 Schubring (2005), 101  
 Schur (1909), 143  
 Searcoid (2002), 149  
 Selberg (1956), 198  
 Simmons (2007), 213  
 Singer (1970), 154  
 Smith and Barnwell (1984a), 35  
 Smith and Barnwell (1984b), 35  
 Smylie et al. (1973), 63  
 Sohrab (2003), 10, 153  
 Steen (1973), 143, 209  
 Steiner (1966), 2  
 Stone (1932), 121, 133, 137  
 Strang and Nguyen (1996), 6, 35  
 Strichartz (1995), 10  
 Süli and Mayers (2003), 96  
 Sweldens and Piessens (1993), 20, 22  
 Talvila (2001), 71  
 Taylor (1715), 10  
 Terras (1999), 198  
 Thomson et al. (2008), 27  
 Ulam (1991), 131  
 Vaidyanathan (1993), 35  
 Väisälä (2003), 131  
 Vidakovic (1999), 20–22, 37, 49, 171  
 von Neumann (1929), 133, 137, 211  
 Walnut (2002), 109, 188  
 Walter and Shen (2001), 158  
 Weber (1893), 68  
 Weyl (1940), 158  
 Widom (1965), 63  
 Wilks (1963), 203

Wojtaszczyk (1997), 167, 169–171, 182, 185, 188, 191 Yates and Mather (1963), 203	Young (1980), 160 Young (2001), 151, 153–155, 160, 164, 165, 167, 168, 172	zay (2004), 188 Zayed (1996), 198 Zukav (1980), 89
---	--	--





$C^*$  algebra, [117](#), [118](#)  
 $C^*$ -algebra, [118](#)  
 $*$ -algebra, [42](#), [114](#), [114–116](#),  
[134](#), [204](#), [208](#)  
 $n$ th moment, [20](#), [21](#), [22](#)  
 $*$ -algebras, [134](#)  
 $\text{\LaTeX}$ , [vi](#)  
 $\text{\TeX}$ -Gyre Project, [vi](#)  
 $\text{\XeTeX}$ , [vi](#)  
*attention markers*, [131](#)  
     problem, [124](#), [130](#), [137](#),  
[139](#), [209](#)  
  
 Abel, Niels Henrik, [213](#)  
 absolute value, [x](#), [68](#)  
 absolutely summable, [28](#), [29](#),  
[207](#)  
 abstract space, [1](#), [1](#)  
 additive, [60](#), [121](#), [123](#), [132](#),  
[193](#)  
 additive identity, [121](#), [204](#)  
 additive inverse, [83](#), [85](#), [121](#)  
 additive property, [17](#)  
 additivity, [60](#), [133](#), [176](#)  
 adjoint, [16](#), [60](#), [116](#), [129](#), [130](#),  
[133](#), [134](#), [190](#), [193](#)  
 Adobe Systems Incorporated, [vi](#)  
 affine, [131](#), [210](#)  
 algebra, [67](#), [68](#), [113](#), [113](#), [114](#)  
 algebra of sets, [xi](#), [2](#)  
 algebras  
      $C^*$ -algebra, [117](#)  
      $*$ -algebra, [114](#)  
 algebras of sets, [2](#), [3](#)  
 Analog Devices ADL5387  
 Quadrature Demodulator,  
[205](#)  
 analyses, [3](#)  
 analysis, [3](#), [3](#), [5](#)  
 analytic, [9](#), [10](#), [15](#)  
 AND, [xi](#)

anti-symmetric, [20](#), [69](#), [182](#),  
[206](#)  
 antiautomorphic, [114](#), [115](#),  
[134](#), [207](#)  
 antiderivative, [9](#)  
 antilinear, [115](#)  
 associates, [120](#)  
 associative, [67](#), [68](#), [120](#), [123](#),  
[143](#)  
 asymmetric, [31](#)  
 auto-correlation, [59](#), [203](#),  
[204](#), [205](#), [206](#)  
 auto-correlation function,  
[201](#)  
 auto-correlation matrix, [63](#)  
 auto-correlation operator,  
[59](#), [61](#), [63](#)  
 auto-power spectrum, [179](#),  
[182](#)  
 auto-spectral density, [207](#),  
[208](#)  
 autocorrelation, [137](#)  
 autocorrelation function,  
[179](#)  
 Avant-Garde, [vi](#)  
  
 Banach algebra, [117](#)  
 Banach space, [154](#), [155](#)  
 bandlimited, [198](#)  
 basis, [5](#), [61](#), [62](#), [151–153](#), [160](#),  
[198](#), [199](#)  
     frame, [172](#)  
     orthogonal, [160](#)  
     orthonormal, [160](#)  
     Riesz, [167](#), [168](#)  
     tight frame, [172](#)  
 Battle-Lemarié orthogonal-  
 ization, [171](#)  
 Bessel's Equality, [157](#), [158](#),  
[159](#), [163](#), [164](#)  
 Bessel's Inequality, [165](#)  
 Bessel's inequality, [158](#)

Best Approximation Theo-  
 rem, [158](#), [158](#)  
 bijection, [165](#)  
 bijective, [xi](#), [131](#), [165](#)  
 Binomial Theorem, [82](#)  
 binomial theorem, [71](#)  
 biorthogonal, [167](#), [169](#), [170](#)  
 Borel measure, [16](#), [69](#)  
 Borel sets, [16](#), [69](#)  
 bounded, [xi](#), [128](#), [137](#), [148](#),  
[155](#), [182](#)  
 bounded bijective, [168](#)  
 bounded linear operator,  
[144](#)  
 bounded linear operators,  
[128](#), [129](#), [130](#), [132](#), [133](#), [135](#),  
[136](#), [138](#), [139](#), [141–144](#)  
 bounded operator, [128](#)  
  
 C star algebra, [117](#)  
 Cardano, Gerolamo, [67](#)  
 Cardinal Series, [198](#)  
 Cardinal series, [198](#)  
 Carl Spitzweg, [213](#)  
 Cartesian product, [x](#)  
 Cauchy-Schwarz inequality,  
[173](#)  
 causal, [32](#), [32](#)  
 characteristic function, [x](#),  
[188](#)  
 Chebyshev polynomial, [96](#)  
 Chebyshev polynomial of the  
 first kind, [96](#)  
 Chebyshev polynomials, [96](#)  
 Cholesky Decomposition, [63](#)  
 closure, [153](#)  
 coefficient functionals, [154](#),  
[154](#)  
 coefficients, [33](#)  
 colored, [59](#)  
 comb function, [52](#)  
 commutative, [27](#), [28](#), [68](#), [85](#),

120, 123, 193, 204  
 commutative ring, 67, 68, 68  
 commutativity, 28  
 commutator relation, 190  
 compact, 211  
 compact support, 6  
 complement, [x](#)  
 complete, 153  
 complete metric space, 153  
 complete set, 153  
 complex auto-power spectrum, 179  
 complex cross-power spectrum, 179  
 complex envelope, 205  
 complex exponential, 5  
 complex linear space, 120  
 complex number system, 83  
 complex-valued, 205  
 conjugate symmetric, 202  
 conjugate linear, 114, 115, 134  
 conjugate pairs, 33  
 conjugate quadrature filter, 35, 35  
 conjugate quadrature filter condition, 35  
 Conjugate quadrature filters, 34  
 conjugate recipricol pairs, 100  
 conjugate symmetric, 132, 202, 206, 208, 212  
 conjugate symmetric property, 17  
 conjugate symmetry, 60  
 conjugate-symmetric, 208  
 constant, 79, 191, 192, 204  
 continuous, [xi](#), 5, 9, 10, 69, 109, 117, 122, 182, 191, 192, 212  
 continuous point spectrum, 89  
 continuous spectrum, 210  
 convergence in probability, 61  
 convergent, 155  
 converges absolutely, 212  
 convex, 2  
 convolution, 19, 27, 27, 28  
 convolution operation, 12, 19  
 convolution operator, 27, 28  
 convolution theorem, 7, 12, 19, 24, 30  
 coordinate functionals, 154  
 coordinates, 151  
 correlation matrix, 63  
 cosine, 75  
 countably infinite, 154, 211  
 counting measure, [xi](#)

CQF, 35, 35, 36  
 CQF condition, 35, 37  
 CQF theorem, 35, 38  
 cross spectral density, 207  
 cross-correlation, 201, 202, 203, 204, 206  
 cross-correlation function, 179  
 cross-power spectrum, 179  
 CS Inequality, 176  
 CSD, 207  
 de la Vallée Poussin kernel, 109, 111  
 de Morgan, 4  
 de Morgan's law, 4  
 definitions  
    $C^*$  algebra, 117, 118  
    $*$ -algebra, 42, 114, 204, 208  
   abstract space, 1  
   algebra, 68, 113  
   Banach algebra, 117  
   biorthogonal, 167  
   bounded linear operators, 128  
   C star algebra, 117  
   coefficient functionals, 154  
   commutative ring, 68  
   complete, 153  
   complex auto-power spectrum, 179  
   complex cross-power spectrum, 179  
   complex linear space, 120  
   continuous spectrum, 210  
   coordinate functionals, 154  
   coordinates, 151  
   CQF, 35  
   dilation operator inverse, 189  
   eigenspace, 209  
   equivalent, 155  
   exact frame, 172  
   expansion, 151, 154  
   exponential function, 80  
   field, 68  
   Fourier coefficients, 160  
   Fourier expansion, 160  
   Fourier series, 160  
   frame, 172  
   frame bounds, 172  
   fundamental, 153  
   group, 67  
   Hamel basis, 151  
   hermitian, 114  
   inner product space, 132

linear basis, 151  
 linear combination, 149  
 linear space, 120  
 multiplicative condition, 117  
   normal, 114  
   normalized tight frame, 172  
   normed algebra, 117  
   normed linear space, 124, 125  
   normed space of linear operators, 125  
   optimal lower frame bound, 172  
   optimal upper frame bound, 172  
   orthogonal basis, 160  
   orthonormal basis, 160  
   Parseval frame, 172  
   point spectrum, 210  
   projection, 114  
   real linear space, 120  
   residual spectrum, 210  
   resolvent, 113  
   resolvent set, 210  
   Riesz basis, 167  
   ring, 67  
   scalars, 120  
   Schauder basis, 154  
   Selberg Trace Formula, 198  
   self-adjoint, 114  
   Smith-Barnwell filter, 35  
   space of all absolutely square summable sequences, 27  
   space of Lebesgue square-integrable functions, 69  
   spans, 149  
   spectral radius, 113  
   spectrum, 113, 210  
   standard inner product, 69  
   standard norm, 69  
   star-algebra, 114  
   synthesis, 3  
   tight frame, 172  
   total, 153  
   translation operator inverse, 189  
   underlying set, 120  
   unital, 113  
   vector space, 120  
   vectors, 120  
 delay, 147  
 DeMoivre's Theorem, 91, 91, 93  
 dense, 153, 154  
 Descartes, René, [ix](#), 187

- difference, [x](#)
- differential operator, [89](#)
- differentiation operator, [10](#)
- dilation, [146](#)
- dilation operator, [5](#), [188](#), [188](#), [190](#), [191](#)
- dilation operator adjoint, [190](#)
- dilation operator inverse, [189](#)
- dimension, [151](#)
- Dirac delta, [52](#)
- Dirac delta distribution, [198](#)
- Dirichlet Kernel, [102–104](#)
- Dirichlet kernel, [109](#)
- discontinuous, [10](#), [15](#)
- discrete, [5](#)
- Discrete Time Fourier Series, [xii](#)
- Discrete Time Fourier Transform, [xii](#), [41](#), [207](#)
- discrete-time Fourier transform, [41](#), [41–43](#), [47](#)
- Dissertation on the propagation of heat in solid bodies, [51](#)
- distributes, [120](#)
- distributive, [5](#), [29](#), [42](#), [114–116](#), [134](#), [204](#)
- distributivity, [190](#)
- domain, [x](#), [187](#)
- double angle formulas, [85](#), [86](#), [87](#), [107](#), [205](#)
- DTFT, [35–37](#), [42](#), [46](#), [49](#), [155](#), [183](#), [195](#), [207](#), [208](#)
- DTFT periodicity, [41](#)
- eigen-system, [62](#)
- Eigendecomposition, [174](#)
- eigenspace, [209](#)
- eigenvalue, [174](#), [209](#)
- eigenvector, [174](#), [209](#)
- empty set, [xi](#)
- equal, [120](#)
- equalities
  - Bessel's, [157](#)
- equality by definition, [x](#)
- equality relation, [x](#)
- equivalent, [144](#), [155](#), [155](#), [167](#)
- Euler Formulas, [110](#)
- Euler formulas, [23](#), [81](#), [82](#), [83](#), [86](#), [87](#)
- Euler's identity, [80](#), [80](#), [81](#), [84](#), [205](#)
- even, [46](#), [96](#), [104](#)
- exact frame, [172](#)
- examples
  - Cardinal Series, [198](#)
  - Fourier Series, [199](#)
  - Fourier Transform, [199](#)
  - Gabor Transform, [199](#)
  - linear functions, [198](#)
  - Mercedes Frame, [175](#)
  - Peace Frame, [175](#)
  - Rectangular pulse, [24](#)
  - rectangular pulse, [23](#)
  - triangle, [23](#)
  - wavelets, [199](#)
- exclusive OR, [xi](#)
- existential quantifier, [xi](#)
- exists, [9](#)
- expansion, [151](#), [154](#)
- exponential function, [80](#)
- false, [xi](#)
- Fast Wavelet Transform, [56](#)
- fast wavelet transform, [55](#)
- Fejér's kernel, [109](#), [109](#), [111](#)
- Fejér-Riesz spectral factorization, [100](#), [100](#)
- FFT, [205](#)
- field, [27](#), [67](#), [68](#), [119](#)
- field of complex numbers, [134](#)
- filter banks, [34](#)
- FontLab Studio, [vi](#)
- for each, [xi](#)
- Fourier Analysis, [5](#)
- Fourier analysis, [4](#), [5](#)
- fourier analysis, [15](#)
- Fourier coefficients, [160](#), [198](#)
- Fourier expansion, [160](#), [160–162](#), [164](#), [166](#)
- Fourier kernel, [16](#)
- Fourier Series, [xii](#), [51](#), [199](#)
- Fourier series, [160](#)
- Fourier Series adjoint, [53](#)
- Fourier series analysis, [5](#)
- Fourier series expansion, [162](#)
- Fourier Series operator, [51](#)
- Fourier shift theorem, [12](#)
- Fourier Transform, [xii](#), [10](#), [16](#), [16](#), [19](#), [20](#), [89](#), [145](#), [147](#), [171](#), [195](#), [199](#)
  - adjoint, [17](#)
- Fourier transform, [4](#), [21](#), [23](#), [147](#), [155](#), [179](#), [196](#), [197](#)
  - inverse, [17](#)
- Fourier Transform operator, [190](#)
- Fourier transform scaling factor, [16](#)
- Fourier, Joseph, [1](#), [15](#)
- frame, [170](#), [172](#), [172](#), [176](#)
- frame bound, [173](#), [175](#)
- frame bounds, [172](#), [176](#)
- frame operator, [172](#), [172](#), [173](#), [175](#)
- frames, [150](#)
- Fredholm integral equation of the first kind, [201](#), [211](#)
- Fredholm integral operators, [145](#)
- Fredholm operator, [211](#), [211](#), [212](#)
- Fredholm operators, [211](#)
- Free Software Foundation, [vi](#)
- function, [16](#), [69](#), [120](#), [188](#)
  - characteristic, [188](#)
  - even, [46](#)
  - indicator, [188](#)
- function overload, [204](#)
- functional, [134](#)
- functions, [xi](#)
  - $n$ th moment, [21](#)
  - absolute value, [68](#)
  - adjoint, [116](#)
  - antiderivative, [9](#)
  - auto-correlation, [59](#), [203](#), [204](#), [205](#), [206](#)
  - auto-correlation function, [201](#)
  - auto-power spectrum, [179](#), [182](#)
  - auto-spectral density, [208](#)
  - autocorrelation function, [179](#)
  - Borel measure, [16](#), [69](#)
  - characteristic function, [188](#)
  - Chebyshev polynomial, [96](#)
  - Chebyshev polynomial of the first kind, [96](#)
  - comb function, [52](#)
  - complex envelope, [205](#)
  - complex exponential, [5](#)
  - conjugate quadrature filter, [35](#)
  - continuous point spectrum, [89](#)
  - cosine, [75](#)
  - cross-correlation, [201](#), [203](#), [204](#), [206](#)
  - cross-correlation function, [179](#)
  - cross-power spectrum, [179](#)
  - de la Vallée Poussin kernel, [109](#), [111](#)
  - dilation operator, [191](#)
  - Dirac delta, [52](#)
  - Dirichlet Kernel, [102](#), [104](#)
  - Dirichlet kernel, [109](#)
  - Discrete Time Fourier Transform, [41](#)
  - discrete-time Fourier transform, [41–43](#)
  - DTFT, [36](#), [46](#), [49](#)
  - eigenvector, [174](#)
  - Fejér's kernel, [109](#), [109](#), [111](#)
  - Fourier coefficients, [198](#)

Fourier kernel, **16**  
 Fourier transform, **21**,  
**23**, **179**, **196**, **197**  
 Fredholm integral equation of the first kind, **211**  
 Heaviside function, **13**  
 impulse response, **207**,  
**208**  
 indicator function, **188**  
 induced norm, **163**  
 inner product, **16**, **132**  
 inphase component,  
**205**  
 input, **207**, **208**  
 Jackson kernel, **109**, **111**  
 Kronecker delta, **62**  
 Kronecker delta function, **160**  
 kronecker delta function, **47**  
 linear functional, **130**  
 mean, **201**, **203**, **204**, **204**  
 measure, **184**  
 modulus, **68**  
 norm, **124**, **125**  
 operator norm, **125**  
 output, **207**, **208**  
 Parseval's equation, **54**  
 Plancherel's formula, **54**  
 Plancherel's formula, **54**  
 Poisson kernel, **109**, **111**  
 Poisson Summation Formula, **198**  
 power spectral density,  
**208**  
 power spectrum, **182**  
 quadrature component,  
**205**  
 random process, **59**, **61**,  
**201**  
 random sequence, **203**,  
**203–207**  
 random variable, **201**  
 Riesz sequence, **167**  
 scalar product, **132**  
 sequence, **27**, **203**  
 set indicator, **183**  
 set indicator function,  
**24**  
 sine, **75**  
 spectral density function, **185**  
 spectral power, **208**  
 standard inner product,  
**179**  
 summability kernel, **109**  
 Taylor expansion, **73**  
 translation operator, **22**,  
**188**  
 variance, **203**, **204**, **204**  
 Volterra integral equation,

**83**, **85**  
 Volterra integral equation of the second type, **78**  
 wavelet, **199**  
 z transform, **31**  
 Z-transform, **43**  
 z-transform, **35**, **36**, **179**  
 Zak Transform, **198**  
 fundamental, **153**  
 Fundamental Theorem of Calculus, **9**, **14**  
 Fundamental theorem of calculus, **70**, **71**  
 Fundamental theorem of linear equations, **124**  
 FWT, **55**, **56**  
 Gabor Transform, **199**  
 Gaussian Pyramid, **6**  
 Gelfand-Mazur Theorem, **117**  
 Generalized Parseval's Identity, **162**  
 generalized product rule, **71**,  
**71**  
 geometric series, **102**, **105**  
 globally, **9**  
 GNU Octave  
   cos, **96**, **99**  
 Golden Hind, **vi**  
 GPR, **71**  
 greatest lower bound, **xi**  
 group, **67**, **67**  
 Gutenberg Press, **vi**  
 half-angle formulas, **87**  
 Hamel bases, **150**  
 Hamel basis, **151**, **151**, **153**  
 Handbook of Algebras, **113**  
 harmonic analysis, **15**  
 Harmonic shifted orthonormality requirement, **185**  
 Heaviside function, **13**  
 Hermetian transpose, **145**  
 Hermite, Charles, **91**  
 hermitian, **114**, **114**, **115**, **137**  
 hermitian components, **116**  
 Hermitian representation, **116**  
 Hermitian symmetric, **20**,  
**100**  
 Heuristica, **vi**  
 high-pass filter, **34**  
 Hilbert space, **4**, **16**, **133**, **134**,  
**137–139**, **144**, **160**, **162**, **164**,  
**167**, **170**, **172**  
 homogeneous, **61**, **68**, **121**,  
**123–125**, **132**, **160**, **161**, **165**,  
**168–170**  
 Housman, Alfred Edward, **vii**  
 identities

Fourier expansion, **161**,  
**162**  
 Parseval frame, **161**  
 identity, **67**, **120**  
 identity element, **120**  
 identity operator, **120**, **120**,  
**189**  
 if, **xi**  
 if and only if, **xi**  
 image, **x**  
 image set, **122**, **124**, **135–139**,  
**144**  
 imaginary part, **xi**, **115**  
 implied by, **xi**  
 implies, **xi**  
 implies and is implied by, **xi**  
 impulse response, **207**, **208**  
 inclusive OR, **xi**  
 indicator function, **x**, **188**  
 induced norm, **61**, **163**  
 inequality  
   Bessel's, **158**  
   triangle, **124**, **125**  
 infinite sum, **149**  
 injective, **xi**, **122**, **123**  
 inner product, **16**, **132**  
 inner product space, **132**,  
**156–158**, **160**, **167**  
 inner-product, **xi**  
 inphase component, **205**  
 input, **207**, **208**  
 inside, **32**  
 integral domain, **67**  
 integration, **9**  
 Integration by Parts, **13**  
 intersection, **x**  
 into, **166**  
 inverse, **67**, **120**, **173**, **175**,  
**188**, **189**  
 inverse DTFT, **37**, **47**, **183**  
 inverse Fourier Series, **52**  
 Inverse Fourier Transform, **145**  
 Inverse Fourier transform, **17**  
 inverse Fourier Transform, **147**  
 Inverse Poisson Summation Formula, **197**, **197**  
 invertible, **70**, **114**, **155**, **172**  
 involutory, **134**  
 involution, **114**, **114**, **118**  
 involutory, **114–116**, **208**  
 IPSE, **180**, **197**, **197**  
 irrational numbers, **192**  
 irreflexive ordering relation,  
**xi**  
 isometric, **17**, **54**, **131**, **140**,  
**140**, **144**, **161**, **169**, **170**  
 isometric in distance, **144**,  
**193**

isometric in length, [144](#), [193](#)  
 isometric operator, [141–143](#)  
 isometry, [140](#)  
 isomorphic, [165](#)

Jackson kernel, [109](#), [111](#)  
 Jacobi, Carl Gustav Jacob, [101](#)  
 jaib, [73](#)  
 jiba, [73](#)  
 jiva, [73](#)  
 join, [xi](#)

Kaneyoshi, Urabe, [214](#)  
 Karhunen-Loève Expansion, [61](#)  
 Kenko, Yoshida, [214](#)  
 kernel, [201](#)  
 Kronecker delta, [62](#)  
 Kronecker delta function, [160](#)  
 kronecker delta function, [47](#)

l'Hôpital's rule, [53](#), [104](#)  
 Lagrange trigonometric identities, [105](#)  
 Laplace convolution, [19](#)  
 Laplace operator, [145](#)  
 Laplace shift relation, [18](#)  
 Laplace Transform, [11](#), [11–13](#), [18–20](#), [89](#)  
 Laplace transform, [89](#)  
 lattice, [2](#)  
 lattice of algebras of sets, [2](#)  
 Laurent series, [28](#)  
 least upper bound, [xi](#)  
 Lebesgue square-integrable functions, [11](#), [16](#), [187](#)  
 left distributive, [67](#), [68](#), [123](#)  
 Leibnitz GPR, [38](#)  
 Leibniz integration rule, [71](#)  
 Leibniz rule, [71](#), [71](#)  
 Leibniz, Gottfried, [ix](#), [119](#)  
 linear, [32](#), [90](#), [121](#), [121](#), [198](#)  
 linear basis, [151](#)  
 linear bounded, [xi](#)  
 linear combination, [149](#)  
 linear combinations, [150](#)  
 linear functional, [130](#)  
 linear functions, [198](#)  
 linear independence, [156](#), [157](#)  
 linear operator, [168](#), [170](#)  
 linear operators, [121](#), [130](#), [166](#)  
 linear ops., [166](#), [170](#)  
 linear space, [1](#), [2](#), [113](#), [120](#), [120](#), [149](#), [151](#)  
 linear spaces, [120](#)  
 linear span, [149](#)  
 linear subspace, [2](#), [149](#)  
 linear time invariant, [89](#)

linearity, [60](#), [62](#), [63](#), [121](#), [122](#)  
 linearly dependent, [149](#), [151–153](#)  
 linearly independent, [149](#), [151–153](#), [156](#), [157](#)  
 linearly ordered, [6](#)  
 Liquid Crystal, [vi](#)  
 locally, [9](#)  
 low-pass filter, [34](#)  
 LTI, [206](#), [208](#)

Machiavelli, Niccolò, [213](#)  
 Maclaurin Series, [10](#)  
 maps to, [x](#)  
 matrix  
   rotation, [145](#)  
 Maxima, [96](#), [99](#)  
 Mazur-Ulam theorem, [131](#)  
 mean, [201](#), [203](#), [204](#), [204](#)  
 measure, [184](#)  
 meet, [xi](#)  
 Mercedes Frame, [175](#)  
 Mercer's Theorem, [61](#), [212](#), [212](#)  
 metric, [xi](#)  
 metric space, [1](#)  
 Minimum phase, [39](#)  
 minimum phase, [31](#), [31](#), [32](#)  
 modular, [5](#)  
 modulus, [68](#)  
 MRA, [6](#)  
 multiplicative condition, [117](#)  
 multiresolution analysis, [6](#)

N5 lattice, [2](#)  
 narrowband, [205](#)  
 Neumann Expansion Theorem, [132](#)  
 non-analytic, [10](#)  
 non-Boolean, [7](#)  
 non-complemented, [7](#)  
 non-distributive, [2](#), [7](#)  
 non-homogeneous, [78](#)  
 non-modular, [7](#)  
 non-negative, [59–61](#), [68](#), [125](#), [132](#), [180](#)  
 non-negative definite, [61](#)  
 noncommutative, [190](#)  
 nondegenerate, [68](#), [124](#), [125](#), [132](#), [192](#)  
 norm, [124](#), [125](#)  
 normal, [54](#), [114](#), [137](#), [138](#), [138](#), [139](#), [144](#), [145](#), [211](#)  
 normal operator, [138](#), [143](#)  
 normalized, [61](#), [62](#)  
 normalized tight frame, [172](#)  
 normed algebra, [117](#), [117](#), [118](#)  
 normed linear space, [124](#), [125](#)

normed linear spaces, [129](#), [140](#)  
 normed space of linear operators, [125](#)  
 NOT, [xi](#)  
 not constant, [192](#)  
 not total, [163](#)  
 null space, [x](#), [122–124](#), [133](#), [135–139](#), [144](#), [209](#)

odd, [96](#), [104](#)  
 ondelette, [6](#)  
 one sided shift operator, [142](#)  
 only if, [xi](#)  
 operations  
   adjoint, [60](#), [129](#), [130](#), [133](#), [190](#), [193](#)  
   analysis, [3](#)  
   auto-correlation, [59](#)  
   auto-correlation matrix, [63](#)  
   auto-correlation operator, [59](#), [61](#), [63](#)  
   auto-spectral density, [207](#)  
   Cholesky Decomposition, [63](#)  
   convolution, [27](#), [28](#)  
   convolution operation, [12](#), [19](#)  
   correlation matrix, [63](#)  
   cross spectral density, [207](#)  
   CSD, [207](#)  
   differential operator, [89](#)  
   differentiation operator, [10](#)  
   dilation operator, [5](#), [188](#), [188](#), [190](#)  
   dilation operator adjoint, [190](#)  
   Discrete Time Fourier Series, [xii](#)  
   Discrete Time Fourier Transform, [xii](#), [207](#)  
   discrete-time Fourier transform, [41](#)  
   DTFT, [35](#), [37](#), [42](#), [155](#), [183](#), [195](#), [207](#), [208](#)  
   Eigendecomposition, [174](#)  
   Fast Wavelet Transform, [56](#)  
   FFT, [205](#)  
   Fourier Series, [xii](#), [51](#)  
   Fourier Series adjoint, [53](#)  
   Fourier Series operator, [51](#)  
   Fourier Transform, [xii](#), [10](#), [16](#), [16](#), [19](#), [20](#), [145](#), [147](#), [171](#), [195](#)



- Fourier transform, [4](#), [155](#)
- frame operator, [172](#), [172](#), [173](#), [175](#)
- Fredholm integral equation of the first kind, [201](#)
- Fredholm operator, [211](#)
- FWT, [56](#)
- Hermetian transpose, [145](#)
- identity operator, [120](#), [189](#)
- imaginary part, [115](#)
- integration, [9](#)
- inverse, [173](#), [175](#), [189](#)
- inverse DTFT, [183](#)
- inverse Fourier Series, [52](#)
- Inverse Fourier Transform, [145](#)
- inverse Fourier Transform, [147](#)
- involution, [114](#)
- kernel, [201](#)
- Laplace operator, [145](#)
- Laplace Transform, [11](#), [11–13](#), [18–20](#)
- Laplace transform, [89](#)
- linear operators, [130](#)
- linear span, [149](#)
- Maclaurin Series, [10](#)
- operator, [120](#), [172](#)
- operator adjoint, [134](#)
- power spectral density, [207](#)
- projection, [135](#)
- PSD, [207](#)
- real part, [115](#)
- reflection operator, [145](#)
- rotation matrix, [145](#)
- rotation operator, [145](#)
- sampling operator, [196](#), [196](#)
- Taylor Series, [10](#), [10](#)
- Toeplitz matrices, [63](#)
- transform, [3](#)
- translation operator, [5](#), [188](#), [188](#), [190](#)
- translation operator adjoint, [190](#)
- Unitary Fourier Transform, [11](#)
- unitary Fourier Transform, [16](#)
- z-domain cross spectral density, [207](#)
- z-domain power spectral density, [207](#)
- Z-Transform, [xii](#)
- z-transform, [28](#), [28](#)
- Z-transform operator, [207](#)
- operator, [119](#), [120](#), [172](#), [188](#)
- adjoint, [115](#)
- autocorrelation, [137](#)
- bounded, [128](#)
- definition, [120](#)
- delay, [147](#)
- dilation, [146](#)
- identity, [120](#)
- isometric, [141–143](#)
- linear, [121](#)
- norm, [125](#)
- normal, [138](#), [139](#), [143](#)
- null space, [135](#)
- positive, [148](#)
- projection, [135](#)
- range, [135](#)
- self-adjoint, [137](#)
- shift, [142](#)
- translation, [145](#)
- unbounded, [128](#)
- unitary, [17](#), [54](#), [143](#), [144](#)
- operator adjoint, [133](#), [134](#)
- operator norm, [xi](#), [125](#), [193](#)
- operator star-algebra, [134](#)
- optimal lower frame bound, [172](#)
- optimal upper frame bound, [172](#)
- order, [x](#), [xi](#)
- ordered pair, [x](#)
- orthogonal, [59](#), [60](#), [62](#), [88](#), [137](#), [156](#), [156](#), [160](#), [185](#)
- orthogonal basis, [160](#)
- orthogonality, [156](#)
- inner product space, [156](#)
- orthomodular lattice, [4](#)
- orthonormal, [157–163](#), [176](#), [185](#)
- orthonormal bases, [150](#)
- orthonormal basis, [54](#), [160](#), [164](#), [165](#), [167–169](#), [171](#)
- orthonormal expansion, [160](#), [164](#)
- orthonormal quadrature conditions, [47](#)
- orthonormality, [158](#), [168](#), [169](#), [176](#)
- orthonormal basis, [161](#)
- oscillation, [9](#)
- output, [207](#), [208](#)
- Paley-Wiener, [198](#)
- Parseval frame, [161](#), [172](#), [172](#)
- Parseval frames, [150](#)
- Parseval's equation, [17](#), [54](#)
- Parseval's Identity, [162](#), [164](#), [166](#), [172](#), [185](#)
- Parseval's Theorem, [208](#)
- Peace Frame, [175](#)
- Peirce, Benjamin, [88](#)
- Per Enflo, [154](#)
- Perfect reconstruction, [160](#)
- periodic, [41](#), [109](#), [180](#), [188](#), [196](#)
- Plancherel's formula, [54](#)
- Plancherel's formula, [17](#), [54](#)
- point spectrum, [210](#)
- Poisson kernel, [109](#), [111](#)
- Poisson Summation Formula, [196](#), [198](#)
- polar identity, [157](#)
- pole, [39](#)
- poles, [33](#)
- polynomial
- trigonometric, [91](#)
- positive, [60](#), [61](#), [148](#)
- positive definite, [60](#), [61](#)
- power set, [xi](#)
- power spectral density, [207](#), [208](#)
- power spectrum, [182](#)
- Primorial numbers, [7](#)
- probability space, [201](#), [203](#)
- product identities, [83](#), [84](#), [85](#), [87](#), [105](#), [106](#)
- projection, [114](#), [135](#)
- projection operator, [135](#), [137](#)
- projection operators, [4](#)
- proper subset, [x](#)
- proper superset, [x](#)
- properties
- absolute value, [x](#)
- absolutely summable, [28](#), [29](#), [207](#)
- additive, [60](#), [121](#), [123](#), [132](#), [193](#)
- additive identity, [121](#), [204](#)
- additive inverse, [83](#), [85](#), [121](#)
- additivity, [60](#), [133](#), [176](#)
- affine, [131](#), [210](#)
- algebra of sets, [xi](#)
- analytic, [9](#), [10](#), [15](#)
- AND, [xi](#)
- anti-symmetric, [20](#), [69](#), [182](#), [206](#)
- antiautomorphic, [114](#), [115](#), [134](#), [207](#)
- associates, [120](#)
- associative, [67](#), [68](#), [120](#), [123](#), [143](#)
- bijection, [165](#)
- bijective, [131](#), [165](#)
- biorthogonal, [169](#), [170](#)
- bounded, [128](#), [137](#), [148](#), [155](#), [182](#)
- bounded bijective, [168](#)
- Cartesian product, [x](#)
- causal, [32](#)
- characteristic function, [x](#)

- closure, 153
- colored, 59
- commutative, 27, 28, 68, 85, 120, 123, 193, 204
- commutativity, 28
- compact, 211
- compact support, 6
- complement, *x*
- complex-valued, 205
- conjugate linear, 114, 134
- conjugate quadrature filter condition, 35
- conjugate symmetric, 132, 202, 206, 208, 212
- conjugate symmetry, 60
- conjugate-symmetric, 208
- constant, 79, 191, 192, 204
- continuous, 5, 9, 10, 69, 109, 122, 182, 191, 192, 212
- convergence in probability, 61
- convergent, 155
- converges absolutely, 212
- convex, 2
- countably infinite, 154, 211
- counting measure, *xi*
- CQF condition, 35, 37
- de Morgan, 4
- de Morgan's law, 4
- dense, 153, 154
- difference, *x*
- discontinuous, 10, 15
- discrete, 5
- distributes, 120
- distributive, 5, 29, 42, 114–116, 134, 204
- distributivity, 190
- domain, *x*
- empty set, *xi*
- equal, 120
- equality by definition, *x*
- equality relation, *x*
- equivalent, 144, 155, 167
- even, 46, 96, 104
- exclusive OR, *xi*
- existential quantifier, *xi*
- exists, 9
- false, *xi*
- for each, *xi*
- Fredholm operator, 211, 212
- Fredholm operators, 211
- globally, 9
- greatest lower bound, *xi*
- hermitian, 114, 115, 137
- Hermitian symmetric, 20, 100
- homogeneous, 61, 68, 121, 123–125, 132, 160, 161, 165, 168–170
- identity, 67, 120
- if, *xi*
- if and only if, *xi*
- image, *x*
- imaginary part, *xi*
- implied by, *xi*
- implies, *xi*
- implies and is implied by, *xi*
- inclusive OR, *xi*
- indicator function, *x*
- injective, 122, 123
- inner-product, *xi*
- inside, 32
- intersection, *x*
- into, 166
- inverse, 67
- invertible, 70, 114, 155, 172
- involutary, 134
- involution, 114, 118
- involutory, 114–116, 208
- irreflexive ordering relation, *xi*
- isometric, 17, 54, 131, 140, 140, 144, 161, 169, 170, 144, 193
- isometric in distance, 193
- isometric in length, 144, 193
- isomorphic, 165
- join, *xi*
- least upper bound, *xi*
- left distributive, 67, 68, 123
- linear, 90, 121, 121, 198
- linear independence, 156, 157
- linear time invariant, 89
- linearity, 60, 62, 63, 121, 122
- linearly dependent, 149, 151–153
- linearly independent, 149, 151–153, 156, 157
- linearly ordered, 6
- locally, 9
- LTI, 206, 208
- maps to, *x*
- meet, *xi*
- metric, *xi*
- Minimum phase, 39
- minimum phase, 31, 31, 32
- modular, 5
- narrowband, 205
- non-analytic, 10
- non-Boolean, 7
- non-complemented, 7
- non-distributive, 2, 7
- non-homogeneous, 78
- non-modular, 7
- non-negative, 59–61, 68, 125, 132, 180
- non-negative definite, 61
- noncommutative, 190
- nondegenerate, 68, 124, 125, 132, 192
- normal, 54, 137, 138, 144, 145, 211
- normalized, 61, 62
- NOT, *xi*
- not constant, 192
- not total, 163
- null space, *x*
- odd, 96, 104
- only if, *xi*
- operator norm, *xi*
- order, *x*, *xi*
- ordered pair, *x*
- orthogonal, 59, 60, 62, 137, 156, 156, 160, 185
- orthogonality, 156
- orthonormal, 157–163, 176, 185
- orthonormality, 158, 168, 169, 176
- oscillation, 9
- Paley-Wiener, 198
- Parseval's Theorem, 208
- periodic, 41, 109, 180, 188, 196
- positive, 60, 61, 148
- positive definite, 60, 61
- power set, *xi*
- proper subset, *x*
- proper superset, *x*
- pseudo-distributes, 120
- range, *x*
- real, 33, 44, 182
- real part, *xi*
- real-valued, 20, 42, 43, 46, 60, 61, 137, 208
- reality condition, 18
- reflexive ordering relation, *xi*
- regular, 211
- relation, *x*
- relational and, *x*
- right distributive, 67, 68, 123
- ring of sets, *xi*
- scalar commutative, 68
- self adjoint, 137, 193
- self-adjoint, 59–61, 137, 137, 193
- separable, 154, 155, 167

- set of algebras of sets, [xi](#)
- set of rings of sets, [xi](#)
- set of topologies, [xi](#)
- shift property, [185](#), [186](#)
- similar, [194](#)
- space of linear trans-  
forms, [122](#)
- span, [xi](#)
- spans, [151](#), [152](#)
- stability condition, [170](#),  
[172](#)
- stable, [32](#)
- stationary in the 1st mo-  
ment, [204](#)
- stationary in the 2nd  
moment, [204](#)
- Strang-Fix condition, [22](#)
- strictly positive, [60](#), [124](#)
- strong convergence,  
[154](#), [164](#)
- subadditive, [68](#), [124](#), [125](#)
- submultiplicative, [68](#)
- subset, [x](#)
- summability kernel, [109](#)
- super set, [x](#)
- surjective, [144](#), [193](#)
- symmetric, [20](#), [44](#), [69](#),  
[181](#), [206](#)
- symmetric difference, [x](#)
- symmetry, [31](#)
- there exists, [xi](#)
- tight frame, [173](#)
- time-invariant, [32](#), [90](#)
- Toeplitz, [63](#)
- topology of sets, [xi](#)
- total, [153](#), [162](#), [163](#), [167](#),  
[168](#)
- triangle inequality, [68](#)
- triangle inequality, [124](#)
- true, [x](#)
- uncorrelated, [59](#)
- uniformly, [212](#)
- union, [x](#)
- unique, [151](#), [154](#)
- unit length, [142](#), [144](#)
- unitary, [17](#), [54](#), [143](#), [143](#)–  
[145](#), [180](#), [182](#), [185](#), [186](#), [190](#),  
[191](#), [193](#)
- universal quantifier, [xi](#)
- vector norm, [xi](#)
- white, [59](#), [59](#)
- wide sense stationary,  
[204](#), [204](#), [206](#), [207](#)
- WSS, [204](#), [206](#), [208](#)
- zero-mean, [63](#)
- PSD, [207](#)
- pseudo-distributes, [120](#)
- PSE, [22](#), [196](#)
- pstricks, [vi](#)
- Pythagorean Theorem, [157](#),  
[159](#), [161](#), [168](#), [169](#)
- Pythagorean theorem, [156](#)
- quadrature component, [205](#)
- quotes
  - Abel, Niels Henrik, [213](#)
  - Cardano, Gerolamo, [67](#)
  - Descartes, René, [ix](#), [187](#)
  - Fourier, Joseph, [1](#), [15](#)
  - Hermite, Charles, [91](#)
  - Housman, Alfred Ed-  
ward, [vii](#)
  - Jacobi, Carl Gustav Ja-  
cob, [101](#)
  - Kaneyoshi, Urabe, [214](#)
  - Kenko, Yoshida, [214](#)
  - Leibniz, Gottfried, [ix](#),  
[119](#)
  - Machiavelli, Niccolò,  
[213](#)
  - Peirce, Benjamin, [88](#)
  - Russull, Bertrand, [vii](#)
  - Stravinsky, Igor, [vii](#)
  - Ulam, Stanislaus M., [130](#)
  - von Neumann, John, [89](#)
- random process, [59](#), [61](#), [201](#),  
[201](#)
- random processes, [179](#)
- random sequence, [203](#), [203](#)–  
[207](#)
- random variable, [201](#), [201](#)
- range, [x](#), [187](#)
- range space, [133](#)
- rational numbers, [192](#)
- real, [33](#), [44](#), [182](#)
- real linear space, [120](#)
- real number system, [83](#)
- real part, [xi](#), [115](#)
- real-valued, [20](#), [42](#), [43](#), [46](#), [60](#),  
[61](#), [137](#), [208](#)
- reality condition, [18](#)
- Rectangular pulse, [24](#)
- rectangular pulse, [23](#)
- reflection, [131](#)
- reflection operator, [145](#)
- reflexive ordering relation, [xi](#)
- regular, [211](#)
- relation, [x](#), [120](#), [188](#)
- relational and, [x](#)
- relations, [xi](#)
  - function, [188](#)
  - operator, [188](#)
  - relation, [188](#)
- residual spectrum, [210](#)
- resolvent, [113](#)
- resolvent set, [210](#), [211](#)
- Riesz bases, [150](#)
- Riesz basis, [167](#), [168](#)–[171](#),  
[182](#)
- Riesz sequence, [167](#), [169](#)
- Riesz-Fischer Theorem, [164](#)
- Riesz-Fischer Thm., [166](#)
- right distributive, [67](#), [68](#), [123](#)
- ring, [67](#), [67](#), [68](#)
  - absolute value, [68](#)
  - commutative, [67](#)
  - modulus, [68](#)
- ring of complex square  $n \times n$   
matrices, [134](#)
- ring of sets, [xi](#)
- Robinson's Energy Delay  
Theorem, [31](#)
- rotation matrix, [145](#)
- rotation matrix operator, [190](#)
- rotation operator, [145](#)
- Russull, Bertrand, [vii](#)
- sampling operator, [196](#), [196](#)
- scalar commutative, [68](#)
- scalar product, [132](#)
- scalars, [120](#)
- scaling, [5](#)
- scaling coefficients, [55](#)
- scaling filter coefficients, [55](#)
- scaling filters, [56](#)
- Schauder bases, [150](#), [155](#)
- Schauder basis, [154](#), [154](#),  
[155](#), [160](#), [167](#)
- Selberg Trace Formula, [198](#)
- self adjoint, [137](#), [193](#)
- self-adjoint, [59](#)–[61](#), [114](#), [137](#),  
[137](#), [193](#)
- semilinear, [115](#)
- separable, [154](#), [155](#), [167](#)
- separable Hilbert space, [27](#),  
[165](#), [167](#)–[169](#)
- separable Hilbert spaces, [165](#)
- sequence, [27](#), [203](#)
- sequences, [35](#)
- set indicator, [183](#)
- set indicator function, [24](#)
- set of algebras of sets, [xi](#)
- set of rings of sets, [xi](#)
- set of topologies, [xi](#)
- shift identities, [82](#), [84](#), [85](#),  
[108](#), [109](#)
- shift operator, [142](#)
- shift property, [185](#), [186](#)
- shift relation, [23](#), [24](#), [180](#)
- similar, [194](#)
- sinc, [23](#)
- sine, [73](#), [75](#)
- sinus, [73](#)
- Smith-Barnwell filter, [35](#)
- space
  - inner product, [132](#)
  - linear, [119](#)
  - normed vector, [124](#)
  - vector, [119](#)
- space of all absolutely square  
Lebesgue integrable func-  
tions, [27](#)
- space of all absolutely square  
summable sequences, [27](#)



- space of all absolutely square summable sequences over  $\mathbb{R}$ , [196](#)
- space of all continuously differentiable real functions, [75](#)
- space of Lebesgue square-integrable functions, [69](#), [196](#)
- space of linear transforms, [122](#)
- space of square integrable functions, [5](#)
- span, [xi](#), [153](#), [155](#), [182](#)
- spans, [149](#), [151](#), [152](#)
- spectral density function, [185](#)
- spectral factorization, [100](#)
- spectral power, [208](#)
- spectral radius, [113](#)
- Spectral Theorem, [211](#)
- spectrum, [113](#), [210](#), [211](#)
- squared identities, [87](#)
- stability, [32](#)
- stability condition, [170](#), [172](#)
- stable, [32](#)
- standard inner product, [69](#), [179](#)
- standard norm, [69](#)
- standard orthonormal basis, [161](#)
- star-algebra, [114](#), [114](#), [134](#)
- star-algebras, [133](#), [134](#)
- stationary in the 1st moment, [204](#)
- stationary in the 2nd moment, [204](#)
- Stifel formula, [71](#)
- Strang-Fix condition, [22](#), [22](#)
- Stravinsky, Igor, [vii](#)
- strictly positive, [60](#), [124](#)
- strong convergence, [154](#), [164](#)
- structures
- $C^*$  algebra, [117](#)
  - $C^*$ -algebra, [118](#)
  - $*$ -algebra, [42](#), [114](#), [114](#)–[116](#), [134](#), [204](#), [208](#)
  - $*$ -algebras, [134](#)
  - abstract space, [1](#), [1](#)
  - adjoint, [134](#)
  - algebra, [68](#), [113](#), [113](#), [114](#)
  - algebra of sets, [2](#)
  - algebras of sets, [3](#)
  - analyses, [3](#)
  - analysis, [3](#), [5](#)
  - Banach space, [154](#), [155](#)
  - basis, [5](#), [61](#), [62](#), [151](#)–[153](#), [160](#), [198](#), [199](#)
  - bijection, [165](#)
  - bijective, [165](#)
  - Borel sets, [16](#), [69](#)
  - bounded linear operator, [144](#)
  - bounded linear operators, [128](#), [129](#), [130](#), [132](#), [133](#), [135](#), [136](#), [138](#), [139](#), [141](#)–[144](#)
  - $C$  star algebra, [117](#)
  - Cardinal series, [198](#)
  - coefficient functionals, [154](#)
  - coefficients, [33](#)
  - commutative ring, [67](#), [68](#), [68](#)
  - complete metric space, [153](#)
  - complete set, [153](#)
  - complex linear space, [120](#)
  - complex number system, [83](#)
  - conjugate pairs, [33](#)
  - conjugate quadrature filter, [35](#)
  - Conjugate quadrature filters, [34](#)
  - conjugate reciprocal pairs, [100](#)
  - continuous spectrum, [210](#)
  - convolution operator, [27](#), [28](#)
  - coordinates, [151](#)
  - CQE, [35](#), [35](#), [36](#)
  - Dirac delta distribution, [198](#)
  - discrete-time Fourier transform, [47](#)
  - domain, [187](#)
  - eigen-system, [62](#)
  - eigenspace, [209](#)
  - expansion, [151](#)
  - field, [27](#), [67](#), [68](#), [119](#)
  - field of complex numbers, [134](#)
  - filter banks, [34](#)
  - Fourier Analysis, [5](#)
  - Fourier analysis, [4](#), [5](#)
  - Fourier series analysis, [5](#)
  - frame, [170](#), [172](#), [172](#), [176](#)
  - frames, [150](#)
  - function, [16](#), [69](#)
  - functional, [134](#)
  - Gaussian Pyramid, [6](#)
  - group, [67](#), [67](#)
  - Hamel bases, [150](#)
  - Hamel basis, [151](#), [151](#), [153](#)
  - high-pass filter, [34](#)
  - Hilbert space, [4](#), [16](#), [133](#), [134](#), [137](#)–[139](#), [144](#), [160](#), [162](#), [164](#), [167](#), [170](#), [172](#)
  - identity, [120](#)
  - identity element, [120](#)
  - image set, [122](#), [124](#), [135](#)–[139](#), [144](#)
  - infinite sum, [149](#)
  - inner product space, [132](#), [156](#)–[158](#), [160](#), [167](#)
  - integral domain, [67](#)
  - inverse, [120](#), [188](#)
  - irrational numbers, [192](#)
  - isometry, [140](#)
  - lattice, [2](#)
  - lattice of algebras of sets, [2](#)
  - Laurent series, [28](#)
  - Lebesgue square-integrable functions, [11](#), [16](#), [187](#)
  - linear basis, [151](#)
  - linear combination, [149](#)
  - linear combinations, [150](#)
  - linear operator, [168](#), [170](#)
  - linear operators, [166](#)
  - linear ops., [166](#), [170](#)
  - linear space, [1](#), [2](#), [113](#), [120](#), [120](#), [149](#), [151](#)
  - linear spaces, [120](#)
  - linear subspace, [2](#), [149](#)
  - low-pass filter, [34](#)
  - metric space, [1](#)
  - MRA, [6](#)
  - multiresolution analysis, [6](#)
  - $N_5$  lattice, [2](#)
  - normalized tight frame, [172](#)
  - normed algebra, [117](#), [118](#)
  - normed linear space, [124](#), [125](#)
  - normed linear spaces, [129](#), [140](#)
  - normed space of linear operators, [125](#)
  - null space, [122](#), [123](#), [137](#)–[139](#), [144](#), [209](#)
  - operator, [119](#)
  - orthogonal basis, [160](#)
  - orthomodular lattice, [4](#)
  - orthonormal bases, [150](#)
  - orthonormal basis, [54](#), [160](#), [164](#), [165](#), [167](#)–[169](#), [171](#)
  - orthonormal basis, [161](#)
  - Parseval frame, [172](#)
  - Parseval frames, [150](#)
  - Parseval's equation, [17](#)
  - Plancherel's formula, [17](#)
  - point spectrum, [210](#)
  - pole, [39](#)
  - poles, [33](#)
  - Primorial numbers, [7](#)
  - probability space, [201](#)


- 203
  - projection operator, 137
  - projection operators, 4
  - random processes, 179
  - range, 187
  - rational numbers, 192
  - real linear space, 120
  - real number system, 83
  - residual spectrum, 210
  - resolvent, 113
  - resolvent set, 210, 211
  - Riesz bases, 150
  - Riesz basis, 168–171, 182
  - Riesz sequence, 169
  - ring, 67, 67, 68
  - ring of complex square  $n \times n$  matrices, 134
  - scalars, 120
  - Schauder bases, 150, 155
  - Schauder basis, 154, 154, 155, 160, 167
  - separable Hilbert space, 27, 165, 167–169
  - separable Hilbert spaces, 165
  - sequences, 35
  - Smith-Barnwell filter, 35
  - space of all absolutely square Lebesgue integrable functions, 27
  - space of all absolutely square summable sequences, 27
  - space of all absolutely square summable sequences over  $\mathbb{R}$ , 196
  - space of all continuously differentiable real functions, 75
  - space of Lebesgue square-integrable functions, 69, 196
  - space of square integrable functions, 5
  - span, 153, 155, 182
  - spectral radius, 113
  - spectrum, 113, 210, 211
  - standard orthonormal basis, 161
  - star-algebra, 114, 134
  - star-algebras, 133
  - subspace, 2
  - subspaces, 2
  - system, 206
  - tight frame, 172, 173, 175
  - tight frames, 150
  - topological dual space, 129
  - topological linear space, 153
    - topological space, 1
    - topology, 2, 149
    - total set, 153
    - translation operator, 198
    - underlying set, 120
    - unit vector, 169
    - unital  $*$ -algebra, 114
    - unital algebra, 113
    - vector, 151
    - vector space, 1, 2, 120
    - vectors, 120
    - wavelet analysis, 5
    - zero, 39
    - zeros, 31, 33
  - subadditive, 68, 124, 125
  - submultiplicative, 68
  - subset,  $x$
  - subspace, 2
  - subspaces, 2
  - summability kernel, 109, 109
  - Summation around unit circle, 107, 174
  - super set,  $x$
  - surjective,  $xi$ , 144, 193
  - symmetric, 20, 44, 69, 181, 206
    - conjugate, 202
  - symmetric difference,  $x$
  - symmetry, 31
  - synthesis, 3
  - system, 206
  - Taylor expansion, 73
  - Taylor Series, 10, 10
  - Taylor series, 79, 81
  - Taylor series for cosine, 77, 78
  - Taylor series for cosine/sine, 76
  - Taylor series for sine, 77
  - The basis problem, 154
  - The Book Worm, 213
  - The Fourier Series Theorem, 162
  - theorems
    - Battle-Lemarié orthogonization, 171
    - Bessel's Equality, 157, 158, 159, 163, 164
    - Bessel's Inequality, 165
    - Bessel's inequality, 158
    - Best Approximation Theorem, 158, 158
    - Binomial Theorem, 82
    - binomial theorem, 71
    - Cauchy-Schwarz inequality, 173
    - commutator relation, 190
    - convolution theorem, 7,
- 12, 19, 24, 30
  - CQF theorem, 35, 38
  - CS Inequality, 176
  - DeMoivre's Theorem, 91, 91, 93
  - double angle formulas, 85, 86, 87, 107, 205
  - DTFT periodicity, 41
  - Euler formulas, 23, 81, 82, 83, 86, 87
  - Euler's identity, 80, 80, 81, 84, 205
  - Fejér-Riesz spectral factorization, 100, 100
  - Fourier expansion, 162, 164, 166
  - Fourier series expansion, 162
  - Fourier shift theorem, 12
  - Fundamental Theorem of Calculus, 9, 14
  - Fundamental theorem of calculus, 70, 71
  - Fundamental theorem of linear equations, 124
  - Gelfand-Mazur Theorem, 117
  - Generalized Parseval's Identity, 162
  - generalized product rule, 71
  - half-angle formulas, 87
  - Hermitian representation, 116
  - induced norm, 61
  - Integration by Parts, 13
  - inverse DTFT, 37, 47
  - Inverse Fourier transform, 17
  - Inverse Poisson Summation Formula, 197, 197
  - IPSE, 180, 197
  - Karhunen-Loève Expansion, 61
  - l'Hôpital's rule, 53, 104
  - Lagrange trigonometric identities, 105
  - Laplace convolution, 19
  - Laplace shift relation, 18
  - Leibnitz GPR, 38
  - Leibniz integration rule, 71
  - Leibniz rule, 71, 71
  - Mazur-Ulam theorem, 131
  - Mercer's Theorem, 61, 212, 212
  - Neumann Expansion Theorem, 132
  - operator star-algebra, 134

- orthonormal quadrature conditions, [47](#)
  - Parseval's Identity, [162](#), [164](#), [166](#), [172](#), [185](#)
  - Perfect reconstruction, [160](#)
  - Poisson Summation Formula, [196](#)
  - polar identity, [157](#)
  - product identities, [83](#), [84](#), [85](#), [87](#), [105](#), [106](#)
  - PSF, [22](#), [196](#)
  - Pythagorean Theorem, [157](#), [159](#), [161](#), [168](#), [169](#)
  - Pythagorean theorem, [156](#)
  - Riesz-Fischer Theorem, [164](#)
  - Riesz-Fischer Thm., [166](#)
  - Robinson's Energy Delay Theorem, [31](#)
  - shift identities, [82](#), [84](#), [85](#), [108](#), [109](#)
  - shift relation, [23](#), [24](#), [180](#)
  - Spectral Theorem, [211](#)
  - squared identities, [87](#)
  - Stifel formula, [71](#)
  - Strang-Fix condition, [22](#)
  - Summation around unit circle, [107](#), [174](#)
  - Taylor Series, [10](#)
  - Taylor series, [79](#), [81](#)
  - Taylor series for cosine, [77](#), [78](#)
  - Taylor series for cosine/-sine, [76](#)
  - Taylor series for sine, [77](#)
  - The Fourier Series Theorem, [162](#)
  - transversal operator inverses, [189](#)
  - trigonometric expansion, [91](#)
  - trigonometric periodicity, [85](#), [107](#), [108](#)
  - trigonometric reduction, [96](#)
  - there exists, [xi](#)
  - tight frame, [172](#), [173](#), [175](#)
  - tight frames, [150](#)
  - time-invariant, [32](#), [32](#), [90](#)
- Toeplitz, [63](#)
- Toeplitz matrices, [63](#)
- topological dual space, [129](#)
- topological linear space, [153](#)
- topological space, [1](#)
- topology, [2](#), [149](#)
- topology of sets, [xi](#)
- total, [153](#), [153](#), [162](#), [163](#), [167](#), [168](#)
- total set, [153](#)
- transform, [3](#), [4](#)
  - inverse Fourier, [17](#)
- translation, [145](#)
- translation operator, [5](#), [22](#), [188](#), [188](#), [190](#), [198](#)
- translation operator adjoint, [190](#)
- translation operator inverse, [189](#)
- transversal operator inverses, [189](#)
- triangle, [23](#)
- triangle inequality, [68](#), [125](#)
- triangle inequality, [124](#)
- trigonometric expansion, [91](#)
- trigonometric periodicity, [85](#), [107](#), [108](#)
- trigonometric reduction, [96](#)
- true, [x](#)
- two-sided Laplace transform, [194](#)
- Ulam, Stanislaus M., [130](#)
- uncorrelated, [59](#)
- underlying set, [120](#)
- uniformly, [212](#)
- union, [x](#)
- unique, [151](#), [154](#)
- unit length, [142](#), [144](#)
- unit vector, [169](#)
- unital, [113](#)
- unital \*-algebra, [114](#)
- unital algebra, [113](#)
- unitary, [16](#), [17](#), [54](#), [143](#), [143–145](#), [147](#), [180](#), [182](#), [185](#), [186](#), [190](#), [191](#), [193](#)
- Unitary Fourier Transform, [11](#)
- unitary Fourier Transform, [16](#)
- unitary operator, [54](#), [143](#)
- universal quantifier, [xi](#)
- Utopia, [vi](#)
- values
  - $n$ th moment, [20](#)
  - dimension, [151](#)
  - eigenvalue, [174](#), [209](#)
  - eigenvector, [209](#)
  - frame bound, [173](#), [175](#)
  - frame bounds, [176](#)
- vanishing moments, [21](#), [49](#)
- variance, [203](#), [204](#), [204](#)
- vector, [151](#)
- vector norm, [xi](#)
- vector space, [1](#), [2](#), [120](#)
- vectors, [120](#)
- Volterra integral equation, [83](#), [85](#)
- Volterra integral equation of the second type, [78](#)
- von Neumann, John, [89](#)
- wavelet, [199](#)
- wavelet analysis, [5](#)
- wavelet coefficients, [55](#)
- wavelet filter coefficients, [55](#)
- wavelet filters, [56](#)
- wavelet transform, [55](#)
- wavelets, [199](#)
- Weierstrass functions, [91](#)
- white, [59](#), [59](#)
- wide sense stationary, [204](#), [204](#), [206](#), [207](#)
- width, [103](#)
- WSS, [204](#), [204](#), [206](#), [208](#)
- z transform, [31](#)
- z-domain cross spectral density, [207](#)
- z-domain power spectral density, [207](#)
- Z-Transform, [xii](#)
- Z-transform, [43](#)
- z-transform, [28](#), [28](#), [35](#), [36](#), [179](#)
- Z-transform operator, [207](#)
- Zak Transform, [198](#)
- zero, [39](#)
- zero at  $-1$ , [44](#)
- zero-mean, [63](#)
- zeros, [31](#), [33](#)

## License

This document is provided under the terms of the [Creative Commons](#) license [CC BY-NC-ND 4.0](#). For an exact statement of the license, see

<https://creativecommons.org/licenses/by-nc-nd/4.0/legalcode>

The icon  appearing throughout this document is based on one that was once at <https://creativecommons.org/> where it was stated, “Except where otherwise noted, content on this site is licensed under a Creative Commons Attribution 4.0 International license.”



...last page ...please stop reading ...

