
Ionisation Amplitudes in Electron-Impact Helium Collisions within the S-Wave Model

Tom Ross supervised by Professor Igor Bray

[ABSTRACT]

Contents

1	Introduction	3
2	Theory	3
2.1	Convergent Close-Coupling Method	3
2.1.1	Laguerre Basis	3
2.1.2	Target-Projectile System	4
2.1.3	Close-Coupling Equations	4
2.1.4	Transition Amplitudes	4
2.1.5	Cross Sections	4
2.1.6	S-Wave Model	4
2.2	Electron-Impact Hydrogen Scattering	4
2.2.1	Elastic Scattering	4
2.2.2	Excitation	4
2.2.3	Ionisation	4
2.3	Electron-Impact Helium Scattering	4
2.3.1	Considerations for a Two-Electron Target	4
2.3.2	Elastic Scattering	4
2.3.3	Excitation	4
2.3.4	Ionisation	5
3	Survey of Experimental Literature	5
4	Survey of Theoretical Literature	5
4.1	Electron-Impact Hydrogen Ionisation Calculations	5
4.1.1	Convergent Close-Coupling Calculations	5
4.1.2	Exterior-Complex-Scaling Calculations	5
4.1.3	Ansatz of Zatsarinny and Bartschat	5
4.2	Electron-Impact Helium Ionisation Calculations	5
4.2.1	Convergent Close-Coupling Calculations	5
4.2.2	Exterior-Complex-Scaling Calculations	5
4.2.3	Ansatz of Zatsarinny and Bartschat	5
5	Conclusion	5
6	Laguerre Radial Basis	6
A	Properties of Laguerre Basis	7
A.1	Laguerre Radial Basis	7
A.1.1	Completeness	7
A.2	Spherical Harmonic	7
A.2.1	Completeness	7

List of Figures

List of Tables

1 Introduction

Applications of Electron-Impact Hydrogen Scattering

Specific Applications of Electron-Impact Hydrogen Ionisation

Development of Quantum Scattering Theory

2 Theory

We shall describe the development of the Convergent Close-Coupling (CCC) method for generalised projectile-target scattering, before describing its application to the cases of: electron-impact hydrogen (e-H) scattering, and electron-impact helium (e-He) scattering. In particular, we shall explore the treatment of target ionisation within the CCC method.

2.1 Convergent Close-Coupling Method

In brief, the CCC method utilises the method of basis expansion to numerically solve the Lippmann-Schwinger equation, for a projectile-target system, to yield the transition amplitudes, which are convergent as the size of the basis is increased. The rate of convergence depends on many factors, such as the complexity and geometry of the projectile-target system for example, as well as the choice of basis used in the expansion. Furthermore, by selecting a complete basis, ionisation transition amplitudes can be treated in a similar manner to discrete excitation transition amplitudes.

2.1.1 Laguerre Basis

To describe the target structure, the CCC method utilises a Laguerre basis, $\{|\varphi_i\rangle\}_{i=1}^N$, for which the coordinate-space representation is of the form

$$\varphi_i(r, \Omega) = \frac{1}{r} \xi_{k_i, l_i}(r) Y_{l_i}^{m_i}(\Omega) \quad (1)$$

where $Y_{l_i}^{m_i}(\Omega)$ are the spherical harmonics, and where $\xi_{k_i, l_i}(r)$ are the Laguerre radial basis functions, which are of the form

$$\xi_{k, l}(r) = \sqrt{\frac{\lambda_l (k-1)!}{(2l+1+k)!}} (\lambda_l r)^{l+1} \exp\left(-\frac{1}{2} \lambda_l r\right) L_{k-1}^{2l+2}(\lambda_l r) \quad (2)$$

where α_l is the exponential fall-off, for each l , and where $L_{k-1}^{2l+2}(\lambda_l r)$ are the associated Laguerre polynomials. Note that we must have that $k_i \in \{1, \dots, N_l\}$, $l_i \in \{0, \dots, l_{max}\}$ and $m_i \in \{-l_i, \dots, l_i\}$, for each $i \in \{1, 2, \dots, N\}$, where N_l is the total number of radial basis functions for each l .

It is shown in [subsubsection A.1.1](#), for each l , that the Laguerre radial basis functions, $\{\xi_{k, l}(r)\}_{k=1}^\infty$, forms a complete basis for the Hilbert space $L^2_{\mathbb{C}}([0, \infty))$. Similarly, it is also shown in [subsubsection A.2.1](#), that the set of spherical harmonics, $\{Y_l^{-l}(\Omega), \dots, Y_l^l(\Omega)\}_{l=0}^\infty$, forms an orthonormal, complete basis for the Hilbert space $L^2_{\mathbb{C}}(S^2)$. Hence, the Laguerre basis functions $\{\varphi_i(r, \Omega)\}_{i=1}^\infty$, forms a complete basis for the Hilbert space $L^2_{\mathbb{C}}(\mathbb{R}^3)$ space.

Further properties of the Laguerre basis are provided in [Appendix A](#).

2.1.2 Target-Projectile System**2.1.3 Close-Coupling Equations****2.1.4 Transition Amplitudes****2.1.5 Cross Sections****Total Cross Sections****Differential Cross Sections****2.1.6 S-Wave Model****2.2 Electron-Impact Hydrogen Scattering****2.2.1 Elastic Scattering****2.2.2 Excitation****2.2.3 Ionisation****Singlet Case****Triplet Case****2.3 Electron-Impact Helium Scattering****2.3.1 Considerations for a Two-Electron Target****Pauli Exclusion Principle****Frozen-Core Model****2.3.2 Elastic Scattering****2.3.3 Excitation****Auto-Ionisation**

2.3.4 Ionisation

3 Survey of Experimental Literature

4 Survey of Theoretical Literature

4.1 Electron-Impact Hydrogen Ionisation Calculations

4.1.1 Convergent Close-Coupling Calculations

4.1.2 Exterior-Complex-Scaling Calculations

4.1.3 Ansatz of Zatsarinny and Bartschat

4.2 Electron-Impact Helium Ionisation Calculations

4.2.1 Convergent Close-Coupling Calculations

4.2.2 Exterior-Complex-Scaling Calculations

4.2.3 Ansatz of Zatsarinny and Bartschat

5 Conclusion

6 Laguerre Radial Basis

References

A Properties of Laguerre Basis

A.1 Laguerre Radial Basis

A.1.1 Completeness

A.2 Spherical Harmonic

A.2.1 Completeness