

# MATH 6370 $p$ -ADIC HODGE THEORY

DANIEL GULOTTA

## 1. MOTIVATION: COMPLEX HODGE THEORY

Cohomology is a way of measuring how many “loops” a space has. Consider the space  $\mathbb{C}^\times = \mathbb{C} \setminus \{0\}$ .

**Definition 1.1.** A 1-cochain on  $\mathbb{C}^\times$  is a function on paths in  $\mathbb{C}^\times$ .

A 1-cochain  $\varphi$  is *closed* if for any continuous map  $f$  from a triangle  $ABC$  to  $\mathbb{C}^\times$ ,  $\phi(f(AC)) = \phi(f(AB)) + \phi(f(BC))$ . It is *exact* if it is of the form

$$\psi(\text{ending point}) - \psi(\text{starting point})$$

for some function  $\psi$  on  $\mathbb{C}^\times$ .

Define

$$H_{\text{sing}}^1(\mathbb{C}^\times, \mathbb{Z}) = \{\mathbb{Z}\text{-valued closed 1-cochains}\} / \{\mathbb{Z}\text{-valued exact 1-cochains}\},$$

and define  $H_{\text{sing}}^1(\mathbb{C}^\times, \mathbb{C})$  similarly.

Then  $H_{\text{sing}}^1(\mathbb{C}^\times, \mathbb{Z})$  is a free abelian group of rank one, and

$$H_{\text{sing}}^1(\mathbb{C}^\times, \mathbb{C}) \cong H_{\text{sing}}^1(\mathbb{C}^\times, \mathbb{Z}) \otimes_{\mathbb{Z}} \mathbb{C}$$

is a  $\mathbb{C}$ -vector space of dimension 1. A class in  $H_{\text{sing}}^1(\mathbb{C}^\times, \mathbb{Z})$  has many representatives, but they all take on the same value on closed paths. There is a generator of  $H_{\text{sing}}^1(\mathbb{C}^\times, \mathbb{Z})$  that takes any path to its winding number around the origin.

**Definition 1.2.** An holomorphic 1-form on  $\mathbb{C}^\times$  is an expression of the form  $f(z) dz$ , where  $f(z)$  is an analytic function on  $\mathbb{C}^\times$ . The holomorphic functions on  $\mathbb{C}^\times$  are precisely the Laurent series

$$\sum_{n=-\infty}^{\infty} a_n z^n,$$

where  $a_n \in \mathbb{C}$  and  $|a_n| \rightarrow 0$  exponentially as  $n \rightarrow \pm\infty$ .

A holomorphic 1-form *exact* if it is of the form  $f'(z) dz$ , where  $f(z)$  is a holomorphic function. (All holomorphic 1-forms are closed.)

Define

$$H_{\text{dR}}^1(\mathbb{C}^\times) = \{\text{holomorphic 1-forms}\} / \{\text{exact holomorphic 1-forms}\}$$

Then  $H_{\text{dR}}^1(\mathbb{C}^\times)$  is a  $\mathbb{C}$ -vector space of dimension 1. The class of  $z^{-1} dz$  is a generator. There is an isomorphism of vector spaces

$$H_{\text{dR}}^1(\mathbb{C}^\times) \xrightarrow{\sim} H_{\text{sing}}^1(\mathbb{C}^\times, \mathbb{C})$$

given by

$$(1.3) \quad f(z) dz \mapsto \left( \gamma \mapsto \int_{\gamma} f(z) dz \right).$$

For any complex manifold  $X$ , one can define the singular cohomology  $H_{\text{sing}}^n(X)$  (defined using maps from simplices into  $X$ ) and the de Rham cohomology  $H_{\text{dR}}^n(X)$  (defined using holomorphic differentials on  $X$ ). There is an isomorphism

$$H_{\text{dR}}^n(X) \cong H_{\text{sing}}^n(X, \mathbb{C})$$

given by integration.

This isomorphism is functorial: if we have a holomorphic or antiholomorphic map  $\sigma: X \rightarrow Y$ , then there is a commutative square

$$\begin{array}{ccc} H_{\text{dR}}^n(Y) & \xrightarrow{\sim} & H_{\text{sing}}^n(Y, \mathbb{C}) \\ \downarrow \sigma^* & & \downarrow \sigma^* \\ H_{\text{dR}}^n(X) & \xrightarrow{\sim} & H_{\text{sing}}^n(X, \mathbb{C}) \end{array}$$

If  $\sigma$  is holomorphic, then

$$\begin{aligned} \sigma^*(f(z) dz) &= f(\sigma(z)) d\sigma(z) \\ \sigma^*(\varphi)(\gamma) &= \varphi(\sigma(\gamma)). \end{aligned}$$

If  $\sigma$  is antiholomorphic, then

$$\begin{aligned} \sigma^*(f(z) dz) &= \overline{f(\sigma(z)) d\sigma(z)} \\ \sigma^*(\varphi)(\gamma) &= \overline{\varphi(\sigma(\gamma))}. \end{aligned}$$

What is the  $p$ -adic version of this story? Let  $K$  be a  $p$ -adic field. (You can assume for now that  $K$  is  $\mathbb{Q}_p$  or a finite extension, but I will make a more general definition later.) A  $p$ -adic analogue of  $\mathbb{C}^\times$  is the rigid analytic space  $\mathbb{A}_K^1 \setminus \{0\}$ .

We will define rigid analytic spaces later. For now, we will just define the space of analytic functions on  $\mathbb{A}_K^1 \setminus \{0\}$ . Motivated by the complex case, We define this space to be the set of Laurent series

$$\sum_{n=-\infty}^{\infty} a_n z^n,$$

where  $a_n \in K$  and the  $|a_n|$ 's go to zero faster than exponentially as  $n \rightarrow \pm\infty$ . A 1-form is an analytic function multiplied by  $dz$ . Then

$$H_{\text{dR}}^1(\mathbb{A}_K^1 \setminus \{0\}) = \{1\text{-forms}\} / \{\text{exact 1-forms}\}$$

is a 1-dimensional  $K$ -vector space generated by the class of  $z^{-1} dz$ .

A  $p$ -adic analogue of singular cohomology is étale cohomology. For now, we will just give a heuristic definition. Consider the map  $\exp: \mathbb{C} \rightarrow \mathbb{C}^\times$ . Any path in  $\mathbb{C}^\times$  that starts and ends at 1 is the image of a path in  $\mathbb{C}$  that starts at 0 and ends at  $2\pi i k$ , where  $k$  is the winding number of the path. So we can identify

$$H_{\text{sing}}^1(\mathbb{C}^\times, \mathbb{Z}) \cong \text{Hom}_{\mathbb{Z}}(2\pi i \mathbb{Z}, \mathbb{Z}).$$

Unlike the complex exponential function, the  $p$ -adic exponential has a finite radius of convergence. So it is useful instead to look at the collection of maps  $z \mapsto z^n$  for each integer  $n$ . A path that starts and ends at 1 and has winding number  $k$  is the image under  $z \mapsto z^n$  of a path that starts at 1 and ends at  $e^{2\pi i k/n}$ . The collection of roots of unity  $\{e^{2\pi i k/n} | n \in \mathbb{Z}_{>0}\}$  is enough to recover  $k$ .

Let  $\mu$  be the set of all roots of unity of  $\mathbb{C}^\times$ . Then we can identify

$$H_{\text{sing}}^1(\mathbb{C}^\times, \mathbb{Z}) \cong \text{Hom}_{\text{cts}}(\mu, \mathbb{Q}/\mathbb{Z}).$$

Here,  $\mu$  has the topology inherited from  $\mathbb{C}^\times$ , and  $\mathbb{Q}/\mathbb{Z}$  has the topology inherited from  $\mathbb{R}/\mathbb{Z} \cong S^1$ .

The isomorphism  $\mathrm{Hom}_{\mathbb{Z}}(2\pi i\mathbb{Z}, \mathbb{Z}) \cong \mathrm{Hom}_{\mathrm{cts}}(\mu, \mathbb{Q}/\mathbb{Z})$  can be described as follows. Any element of the former group is multiplication by  $\frac{k}{2\pi i}$  for some integer  $k$ . Its image in the latter group is the map  $e^{2\pi i m/n} \mapsto mk/n$ .

With this in mind, we define

$$H_{\mathrm{\acute{e}t}}^1(\mathbb{A}_{\overline{K}}^1 \setminus \{0\}, \mathbb{Z}_p) = \mathrm{Hom}_{\mathbb{Z}_p}(\mu_{p^\infty}, \mathbb{Q}_p/\mathbb{Z}_p),$$

where  $\mu_{p^\infty}$  is the set of  $p$ -power roots of unity in  $\overline{K}$ . It is a free  $\mathbb{Z}_p$ -module of rank 1, and it has an action of  $\mathrm{Gal}(\overline{K}/K)$ . We will also denote this group by  $\mathbb{Z}_p(-1)$ .

We would like to compare  $H_{\mathrm{dR}}^1(\mathbb{A}_K^1 \setminus \{0\})$  and  $H_{\mathrm{\acute{e}t}}^1(\mathbb{A}_{\overline{K}}^1 \setminus \{0\}, \mathbb{Z}_p)$ . More specifically, we would like to write down an  $\mathrm{Gal}(\overline{K}/K)$ -equivariant isomorphism

$$H_{\mathrm{dR}}^1(\mathbb{A}_K^1 \setminus \{0\}) \otimes_K L \cong H_{\mathrm{\acute{e}t}}^1(\mathbb{A}_{\overline{K}}^1 \setminus \{0\}, \mathbb{Z}_p) \otimes_{\mathbb{Z}_p} L$$

for some field  $L$ .

What should  $L$  be? The field  $\overline{K}$  has a unique multiplicative absolute value extending the one on  $K$ . We write  $C = \widehat{\overline{K}}$  for the completion of  $\overline{K}$  with respect to this absolute value. The most obvious guess is that  $L = C$ .

However, it turns out that this guess does not work. We will show in a future lecture that there are no nonzero  $\mathrm{Gal}(\overline{K}/K)$ -equivariant maps

$$H_{\mathrm{dR}}^1(\mathbb{A}_K^1 \setminus \{0\}) \otimes_K C \rightarrow H_{\mathrm{\acute{e}t}}^1(\mathbb{A}_{\overline{K}}^1 \setminus \{0\}, \mathbb{Z}_p) \otimes_{\mathbb{Z}_p} C.$$

Instead, we will have to define a ring  $B_{\mathrm{dR}}^+$  that is a completion of  $\overline{K}$  with respect to a more unusual topology. The ring  $B_{\mathrm{dR}}^+$  will be a discrete valuation ring with residue field  $C$ . We will take  $L = B_{\mathrm{dR}} = \mathrm{Frac} B_{\mathrm{dR}}^+$ .

The  $p$ -adic analogues of complex manifolds are called rigid analytic spaces. If you are not familiar with rigid analytic spaces, you can just think about algebraic varieties over a  $p$ -adic field—there is an analytification functor that turns any such variety into a rigid analytic space. Given a rigid analytic space  $X$  over a  $p$ -adic field  $K$ , one can define étale cohomology groups

$$H_{\mathrm{\acute{e}t}}^i(X_{\overline{K}}, \mathbb{Z}_p)$$

and de Rham cohomology groups

$$H_{\mathrm{dR}}^i(X).$$

**Theorem 1.4** (Scholze, [Sch13]). *If  $X$  is proper and smooth, then there is a Galois equivariant isomorphism*

$$H_{\mathrm{dR}}^i(X) \otimes_K B_{\mathrm{dR}} \cong H_{\mathrm{\acute{e}t}}^i(X_{\overline{K}}, \mathbb{Z}_p) \otimes_{\mathbb{Z}_p} B_{\mathrm{dR}}.$$

Here,  $B_{\mathrm{dR}}$  is the fraction field of  $B_{\mathrm{dR}}^+$ .

If  $X$  comes from an algebraic variety, then this isomorphism was previously proved by Tsuji [Tsu99] and Faltings [Fal02]. There is also a version of the comparison theorem for certain non-proper varieties, including  $\mathbb{A}_K^1 \setminus \{0\}$ , due to Li–Pan [LP19].

In  $p$ -adic Hodge theory, we study cohomology theories in  $p$ -adic geometry and the relations between them. Because  $H_{\mathrm{\acute{e}t}}^i(X_{\overline{K}}, \mathbb{Z}_p)$  has a  $\mathrm{Gal}(\overline{K}/K)$ -action, and  $\mathrm{Gal}(\overline{K}/K)$  is much more interesting than  $\mathrm{Gal}(\mathbb{C}/\mathbb{R})$ , the study of Galois representations is an important component of the theory.

## 2. INFINITE GALOIS THEORY

**Definition 2.1.** Let  $L/K$  be extension of fields. Let  $\text{Aut}(L/K)$  denote the group of automorphisms of  $L$  fixing each element of  $K$ .

Give  $\text{Aut}(L/K)$  the weakest topology such that the stabilizer of any finite subset of  $L$  is open.

*Example 2.2.* The group  $\text{Aut}(\mathbb{Q}(\sqrt{2})/\mathbb{Q})$  has two elements. The nontrivial element sends  $\sqrt{2} \mapsto -\sqrt{2}$ .

*Example 2.3.* The group  $\text{Aut}(\mathbb{Q}(\sqrt[3]{2})/\mathbb{Q})$  is trivial. If  $\omega$  is a nontrivial cube root of unity, then  $\text{Aut}(\mathbb{Q}(\sqrt[3]{2}, \omega)/\mathbb{Q})$  permutes  $\{\sqrt[3]{2}, \omega\sqrt[3]{2}, \omega^2\sqrt[3]{2}\}$ , and this permutation action induces an isomorphism  $\text{Aut}(\mathbb{Q}(\sqrt[3]{2}, \omega)/\mathbb{Q}) \cong S_3$ .

*Example 2.4.* Let  $p$  be a prime number. For each positive integer  $n$ , there is a field  $\mathbb{F}_{p^n}$  with  $p^n$  elements. It is unique up to isomorphism. We have  $\text{Aut}(\mathbb{F}_{p^n}/\mathbb{F}_p) \cong \mathbb{Z}/n\mathbb{Z}$ , where the Frobenius automorphism  $x \mapsto x^p$  corresponds to the element  $1 \in \mathbb{Z}/n\mathbb{Z}$ . Then  $\text{Aut}(\mathbb{F}_p/\mathbb{F}_p) = \hat{\mathbb{Z}} = \varprojlim_n \mathbb{Z}/n\mathbb{Z}$ .

*Example 2.5.* For each positive integer  $n$ , there is a field  $\mathbb{Q}(\mu_{p^n})$  obtained by adjoining all  $p$ -power roots of unity to  $\mathbb{Q}$ . There is an isomorphism

$$\begin{aligned} \text{Aut}(\mathbb{Q}(\mu_{p^n})/\mathbb{Q}) &\cong (\mathbb{Z}/p^n\mathbb{Z})^\times \\ (\zeta &\mapsto \zeta^m) \leftarrow m. \end{aligned}$$

Let

$$\mathbb{Q}(\mu_{p^\infty}) = \varinjlim_n \mathbb{Q}(\mu_{p^n}).$$

Then

$$\text{Aut}(\mathbb{Q}(\mu_{p^\infty})/\mathbb{Q}) \cong \varprojlim_n \mathbb{Z}/p^n\mathbb{Z} = \mathbb{Z}_p^\times.$$

Similarly,

$$\text{Aut}(\mathbb{Q}_p(\mu_{p^\infty})/\mathbb{Q}_p) \cong \mathbb{Z}_p^\times.$$

*Example 2.6.* Let  $K$  be a field. Then  $\text{Aut}(K(t)/K) = \text{PGL}_2(K)$ , with the discrete topology. (We leave the proof as an exercise to the reader.)

**Lemma 2.7.** *If  $L/K$  is finite, then  $\text{Aut}(L/K)$  has the discrete topology.*

*Proof.* Choose a  $K$ -vector space basis for  $L$ . An automorphism of  $L$  fixing this basis must be the identity.  $\square$

**Lemma 2.8.** *If  $L/K$  is algebraic, then the map*

$$\text{Aut}(L/K) \rightarrow \varprojlim_{K'} \text{Aut}(K'/K)$$

*is an isomorphism of topological groups, where  $K'$  runs over  $\text{Aut}(L/K)$ -stable finite extensions of  $K$ .*

*Proof.* Since  $L/K$  is algebraic, any  $\alpha \in L$  has finite orbit under  $\text{Aut}(L/K)$ . So the field obtained by adjoining the orbit of  $\alpha$  to  $K$  is a finite  $\text{Aut}(L/K)$ -stable extension of  $K$ . Specifying compatible automorphisms of each  $K'$  is equivalent to specifying an automorphism of  $L$ .  $\square$

**Definition 2.9.** A topological space is *profinite* if it is the inverse limit of a collection of finite sets having the discrete topology.

**Lemma 2.10** ([Sta, Tag 08ZY]). *A topological space is profinite if and only if it is totally disconnected and compact.*

If  $H$  is a group acting on a field  $K$ , we denote by  $K^H$  the subfield of  $K$  fixed by  $H$ .

**Definition 2.11.** We say that  $L/K$  is *Galois* if it is algebraic and  $L^{\text{Aut}(L/K)} = K$ .

If  $L/K$  is Galois, then we will also denote  $\text{Aut}(L/K)$  by  $\text{Gal}(L/K)$ .

*Example 2.12.* Of the extensions mentioned in Examples 2.2–2.5,  $\mathbb{Q}(\sqrt{2})/\mathbb{Q}$ ,  $\mathbb{Q}(\sqrt[3]{2}, \omega)/\mathbb{Q}$ ,  $\overline{\mathbb{F}}_p/\mathbb{F}_p$ ,  $\mathbb{Q}(\mu_{p^\infty})/\mathbb{Q}$ ,  $\mathbb{Q}_p(\mu_{p^\infty})/\mathbb{Q}_p$  are Galois. The extension  $\mathbb{Q}(\sqrt[3]{2})/\mathbb{Q}$  is not Galois since the fixed field of the automorphism group is  $\mathbb{Q}(\sqrt[3]{2})$ . The extension  $K(t)/K$  is not Galois since it is not algebraic.

There is a more concrete characterization of Galois extensions in terms of splitting fields.

**Definition 2.13.** Let  $K$  be a field. A polynomial  $f(x) \in K[x]$  *factors completely* if it can be written in the form  $f(x) = c \prod_{i=1}^n (x - x_i)$  with  $n \in \mathbb{Z}_{\geq 0}$ ,  $c, x_1, \dots, x_n \in K$  and  $c \neq 0$ .

**Definition 2.14.** Let  $L/K$  be an algebraic extension, and let  $P \subset K[x] \setminus \{0\}$ . We say that  $L$  is a *splitting field* for  $P$  if every element of  $P$  factors completely over  $L$ , and no proper subfield of  $L$  has this property.

**Lemma 2.15.** *Every subset of  $K[x] \setminus \{0\}$  admits a splitting field. It is unique up to isomorphism.*

*Proof.* First, suppose  $P$  consists of a single element  $f(x)$ . Then we can construct a splitting field inductively as follows. Letting  $K_0 = K$ , and for  $i \geq 0$ , let  $f_i(x)$  be an irreducible factor of  $f(x)$  of degree  $> 1$  over  $K_i[x]$ , and let  $K_{i+1} = K[x]/f_i(x)$ . Eventually,  $f(x)$  factors completely in some  $K_n$ , and this  $K_n$  is a splitting field for  $\{f(x)\}$ .

If  $L$  is any splitting field of  $\{f(x)\}$ , we can construct an isomorphism  $K_n \xrightarrow{\sim} L$  as follows. For each  $i$ , we construct an embedding  $K_{i+1} \hookrightarrow L$  by sending the generator of  $K_{i+1}$  to some root of the polynomial  $f_i(x)$  in  $L$  (using the map  $K_i \hookrightarrow L$  to consider  $f_i(x)$  as an element of  $L[x]$ ). Since  $f$  does not factor completely over any subfield of  $L$ ,  $K_n \rightarrow L$  must be surjective, hence an isomorphism.

To prove the lemma for arbitrary  $P$ , we use Zorn's lemma. □

**Definition 2.16.** A polynomial  $f(x) \in K[x]$  is *separable* if  $f(x)$  and  $f'(x)$  generate the unit ideal.

**Lemma 2.17.** *Suppose the polynomial  $f(x) \in K[x]$  factors completely. The factors are distinct if and only if  $f$  is separable.*

*Proof.* Suppose  $f(x)$  is divisible by  $(x - \alpha)^2$  for some  $\alpha \in K$ . Then  $f'(x)$  is divisible by  $x - \alpha$ . So  $(f(x), f'(x)) \subset (x - \alpha)$ .

Conversely, suppose  $f(x) = \prod_{i=1}^n (x - \alpha_i)$  has no repeated factors. By the Chinese remainder theorem,  $f(x)$  and  $f'(x)$  generate the unit ideal if and only if  $f'(\alpha_i) \neq 0$  for all  $i$ . In fact, we have

$$f'(\alpha_i) = \prod_{j \neq i} (\alpha_i - \alpha_j) \neq 0.$$

□

**Lemma 2.18.** *An extension  $L/K$  is Galois iff it is the splitting field of a set of separable polynomials.*

*Proof.* Suppose  $L/K$  is Galois. Let  $\alpha \in L$ . Then  $\alpha$  is a zero of some polynomial over  $K$ . Any element of the  $\text{Gal}(L/K)$ -orbit of  $\alpha$  is also a zero of this polynomial. So the orbit is finite. Let  $\{\alpha_1 = \alpha, \alpha_2, \dots, \alpha_n\}$  be the orbit. Then

$$\prod_{i=1}^n (x - \alpha_i)$$

is a polynomial that is  $\text{Gal}(L/K)$ -invariant. Since  $L/K$  is Galois, the polynomial has coefficients in  $K$ . Then  $L$  is a splitting field for the set of all polynomials that can be constructed in this way.

Conversely, suppose  $L/K$  is the splitting field of a set of separable polynomials over  $K$ . WLOG we may assume that they are irreducible over  $K$ . If  $\alpha, \alpha' \in L$  are two roots of the same irreducible polynomial, then the construction of Lemma 2.15 produces an automorphism of  $L$  fixing  $K$  and sending  $\alpha$  to  $\alpha'$ .  $\square$

**Lemma 2.19.** *Let  $L/K$  be a Galois extension. If  $K'$  is a subfield of  $L$  containing  $K$ , then  $L/K'$  is Galois, and  $\text{Gal}(L/K')$  is closed in  $\text{Gal}(L/K)$ .*

*Proof.* By Lemma 2.18,  $L$  is a splitting field for some set of polynomials over  $K$ . Then  $L$  is a splitting field for the same set of polynomials over  $K'$ , so  $L/K'$  is Galois.

By Lemma 2.10,  $\text{Gal}(L/K')$  and  $\text{Gal}(L/K)$  are compact Hausdorff spaces. So  $\text{Gal}(L/K')$  must be closed in  $\text{Gal}(L/K)$ .  $\square$

**Lemma 2.20.** *If  $L$  is a field and  $H \subset \text{Aut } L$  is a finite subgroup, then the map  $H \rightarrow \text{Gal}(L/L^H)$  is an isomorphism.*

*Proof.* The map is injective, so it suffices to prove that  $|H| \geq |\text{Gal}(L/L^H)|$ . From the construction of Lemma 2.15, we see that  $|\text{Gal}(L/L^H)| = [L : L^H]$ . We will show that  $[L : L^H] \leq |H|$ .

Let  $H = \{\sigma_1 = 1, \sigma_2, \dots, \sigma_n\}$ . Let  $\alpha_1, \dots, \alpha_{n+1} \in L$ . The system

$$(2.21) \quad \sum_{j=1}^{n+1} \sigma_i(\alpha_j) X_j = 0$$

has  $n+1$  variables and  $n$  equations, so it has a nonzero solution. Among all solutions, choose a solution  $(c_1, \dots, c_{n+1})$  with the fewest nonzero elements. After reordering the  $\alpha_j$  and multiplying by a scalar, we may assume  $c_1 \neq 0$  and  $c_1 \in F$ . For any  $i$ ,

$$(c_1 - \sigma(c_1), \dots, c_{n+1} - \sigma(c_{n+1}))$$

is a solution to (2.21) with fewer nonzero terms, so it must be zero. So the  $c_j$ 's are all in  $F$ , and  $\alpha_1, \dots, \alpha_{n+1}$  are linearly dependent over  $F$ . Therefore,  $[L : L^H] \leq |H|$ , as desired.  $\square$

**Theorem 2.22** (Fundamental theorem of infinite Galois theory). *There is a bijection between closed subgroups  $H$  of  $\text{Gal}(L/K)$  and subfields  $K'$  of  $L$  containing  $K$ , given by*

$$\begin{aligned} H &\mapsto L^H \\ K' &\mapsto \text{Gal}(L/K'). \end{aligned}$$

*Proof.* By Lemma 2.19, for any subfield  $K'$  of  $L$  containing  $K$ ,  $L^{\text{Gal}(L/K')} = L$ .

Conversely, suppose  $H$  is a closed subgroup of  $\text{Gal}(L/K)$ , and suppose  $\sigma \in \text{Gal}(L/K) \setminus H$ . Since  $H$  is closed, we can find some finite Galois extension  $K''$  of  $K$  such that the action of  $\sigma$  on  $K''$  does not agree with the action of any element of  $H$ . By Lemma 2.20,  $\sigma$  cannot fix  $(K'')^H$ . So it cannot fix  $L^H$ . Therefore,  $H \rightarrow \text{Gal}(L/L^H)$  is an isomorphism.  $\square$

**Definition 2.23.** A *separable closure* of a field  $K$  is a splitting field for the set of all separable polynomials in  $K[x]$ .

We denote a separable closure of  $K$  by  $K^{\text{sep}}$ . We will sometimes write  $G_K$  for  $\text{Gal}(K^{\text{sep}}/K)$ . If  $K$  has characteristic zero, then a separable closure is the same thing as an algebraic closure.

### 3. ELLIPTIC CURVES

In  $p$ -adic Hodge theory, we consider étale cohomology groups  $H_{\text{ét}}^i(X_{\bar{K}}, \mathbb{Z}_p)$ . In general, it is difficult to describe these groups explicitly. In some situations, we can be more explicit. One of these is the case where  $X$  is an elliptic curve.

**Definition 3.1.** Let  $K$  be a field. An elliptic curve over  $K$  is pair  $(E, O)$ , where  $E$  is a complete smooth geometrically irreducible curve of genus 1 over  $K$  and  $O \in E(K)$ .

Sometimes, we will abuse notation and call  $E$  an elliptic curve.

If  $K$  has characteristic  $\neq 2$ , then any elliptic curve is isomorphic to one of the form

$$y^2 = x^3 + ax^2 + bx + c,$$

where  $x^3 + ax^2 + bx + c$  is separable. When  $E$  is written in this form, we usually take  $O$  to be the point at  $\infty$ .

The curve  $E$  has a group structure, meaning that there are morphisms

$$+: E \times E \rightarrow E$$

$$-: E \rightarrow E$$

$$O: \text{Spec } k \rightarrow E$$

(with  $O$  being the point chosen above), satisfying the usual group axioms.

The group structure can be described as follows. Given two points  $P_1, P_2$  on  $E$ , there is exactly one other point  $Q$  where the line through  $P_1, P_2$  intersects  $E$ . (If  $P_1 = P_2$ , we use the tangent line through  $P_1$ .) Define  $P_1 + P_2$  to be the reflection of  $Q$  about the  $x$ -axis.

Then the point at infinity is the identity, and the inverse of any point is its reflection about the  $x$ -axis.

To see that the group operation is associative, we consider line bundles on  $E$ . Let  $dx + ey = f$  be the equation of the line through  $P_1, P_2$ , and let  $g$  be the  $x$ -coordinate of the third intersection of this line with the elliptic curve. The rational function  $\frac{dx+ey-f}{x-g}$  has zeros at  $P_1, P_2$  and poles at  $P_1 + P_2$  and  $O$ . It determines an isomorphism of line bundles

$$\mathcal{O}([P_1] + [P_2] - [O]) \cong \mathcal{O}([P_1 + P_2]).$$

Given a third point  $P_3$ , we have

$$\mathcal{O}([P_1] + [P_2] + [P_3] - 2[O]) \cong \mathcal{O}([(P_1 + P_2) + P_3]) \cong \mathcal{O}([P_1 + (P_2 + P_3)])$$

Since  $E$  does not have genus 0, it cannot have a rational function with a single zero and pole, so we must have  $(P_1 + P_2) + P_3 = P_1 + (P_2 + P_3)$ .

The above proof is somewhat sketchy. A more rigorous treatment uses the Picard functor. There is a contravariant functor  $\text{Pic}: \text{Scheme} \rightarrow \text{Ab}$  that sends a scheme  $X$  to the group of isomorphism classes of line bundles on  $X$ , with the group operation being tensor product. For any map of schemes  $X \rightarrow S$ , there is a contravariant functor  $\text{Pic}_{X/S}: \text{Scheme}/S \rightarrow \text{Ab}$  that sends a scheme  $T$  over  $S$  to  $\text{Pic}(X \times_S T)/\text{Pic}(T)$ . For any curve  $X$  over  $S$ , there is a natural transformation  $X \mapsto \text{Pic}_{X/S}$ . If  $X$  is projective, then any invertible sheaf on  $X$  has a well-defined degree, so we can write

$$\text{Pic}_{X/S} = \bigsqcup_{d \in \mathbb{Z}} \text{Pic}_{X/S}^d.$$

If  $X$  is an elliptic curve, then one can show that  $X \mapsto \text{Pic}_{X/S}^1$  is an isomorphism. The point  $O$  also determines an isomorphism  $\text{Pic}_{X/S}^1 \cong \text{Pic}_{X/S}^0$ . Since  $\text{Pic}_{X/S}$  has a group structure,  $E$  does as well. For more details, see [KM85, Theorem 2.1.2].

*Remark 3.2.* If  $X$  is a singular curve defined by a Weierstrass equation  $y^2 = x^3 + ax^2 + bx + c$ , then the nonsingular locus of  $X$  still has a group structure. (If a line passes through a singular point of  $X$ , the intersection multiplicity is at least 2. So if a line intersects the curve at two nonsingular points, then the third intersection point must also be nonsingular.)

For example, the additive group  $\mathbb{G}_a = \text{Spec } K[t]$  is isomorphic to the nonsingular locus of the cuspidal cubic  $y^2 = x^3$  via the map  $t \mapsto (t^{-3}, t^{-2})$ . We leave it as an exercise to the reader to check that this map is a homomorphism of groups.

*Remark 3.3.* Although the focus of today's lecture was on elliptic curves over fields, one can define an elliptic curve over an arbitrary scheme  $S$ . It is a pair  $(E, O)$ , where  $E$  is a smooth proper curve over  $S$  with geometrically connected fibers of genus 1, and  $O: S \rightarrow E$  is a section of  $E \rightarrow S$ .

If  $E$  is a smooth degree 3 curve in  $\mathbb{P}_S^2$  equipped with a section  $O: S \rightarrow E$ , then  $(E, O)$  is an elliptic curve. Even if  $E$  is not smooth, the nonsingular locus still has a group structure with identity  $O$ .

#### 4. FORMAL GROUPS

(This class was a guest lecture by Petar Bakić. I was not at the lecture, but this is my attempt at summarizing the relevant sections of Silverman's book.)

Let  $R$  be a ring. Let

$$E = \text{Proj } R[X, Y, Z]/(X^3 + aX^2Z + bXZ^2 + cZ^3 - Y^2Z).$$

As we saw in the previous class, the nonsingular locus  $E_{\text{ns}}$  of  $E$  admits a group structure. We take the identity to be the point  $O = (0 : 1 : 0)$ , corresponding to the ideal  $(X, Z)$ .

We can consider the formal completion of  $E$  at  $O$ . The locus  $Y \neq 0$  is the affine  $\text{Spec } A$ , where

$$A = R[x, z]/(x^3 + ax^2z + bxz^2 + cz^3 - z),$$

where  $x = X/Y$ ,  $z = Z/Y$ .

The point  $O$  corresponds to the ideal  $(x, z)$ . The formal completion of  $E$  along  $O$  is  $\hat{E} = \text{Spf } \hat{A}$ , where

$$\hat{A} = \varprojlim_n A/I^n \cong R[[z]].$$



(Don't worry if you are not familiar with formal schemes, as we will just be working with the power series ring  $R[[z]]$ .) The group operation  $E_{\text{ns}} \times_R E_{\text{ns}} \rightarrow E_{\text{ns}}$  induces a group operation  $\hat{E} \times_R \hat{E} \rightarrow \hat{E}$ . This gives us a continuous map of power series rings

$$R[[z]] \cong \hat{A} \rightarrow \hat{A} \hat{\otimes}_R \hat{A} \cong R[[z_1, z_2]] .$$

Note that any map continuous map  $R[[z]] \rightarrow R[[z_1, z_2]]$  is determined by the image of  $z$ . Denote the image of  $z$  by  $F(z_1, z_2)$ . Similarly, there is an inverse map  $\hat{E} \rightarrow \hat{E}$ , which corresponds to a continuous map of power series rings  $R[[z]] \rightarrow R[[z]]$ . Let  $i(z)$  be the image of  $z$ .

Since the group operation is commutative and associative, we will have

$$F(z_1, z_2) = F(z_2, z_1) ,$$

$$F(z_1, F(z_2, z_3)) = F(F(z_1, z_2), z_3) .$$

Similarly, since  $i$  is the inverse operation, and  $z = 0$  is the identity,

$$F(z, i(z)) = 0 .$$

The above analysis leads us to consider the notion of a *formal group law*.

**Definition 4.1.** A *one-parameter commutative formal group* over a ring  $R$  is a power series  $F(X, Y) \in R[[X, Y]]$  satisfying the following conditions:

- (1)  $F(X, 0) = X$  and  $F(0, Y) = Y$
- (2)  $F(X, F(Y, Z)) = F(F(X, Y), Z)$
- (3)  $F(X, Y) = F(Y, X)$
- (4) There is a power series  $i(T) \in R[[T]]$  satisfying  $i(0) = 0$  and  $F(T, i(T)) = 0$ .

*Example 4.2.* The formal additive group  $\hat{\mathbb{G}}_a$  is defined by

$$F(X, Y) = X + Y .$$

*Example 4.3.* The formal multiplicative group  $\hat{\mathbb{G}}_m$  is defined by

$$F(X, Y) = (1 + X)(1 + Y) - 1 = X + Y + XY .$$

**Definition 4.4.** A homomorphism of formal groups  $F \rightarrow G$  is a power series  $\phi \in R[[T]]$  such that

$$\phi(F(X, Y)) = G(\phi(X), \phi(Y)) .$$

*Example 4.5.* For any integer  $m$ , we can define a multiplication-by- $m$  homomorphism  $[m]: F \rightarrow G$  inductively by

$$[0](T) = 0$$

$$[m+1](T) = F([m]T, T)$$

$$[m-1](T) = F([m]T, i(T))$$

If  $F = \mathbb{G}_a$ , then

$$[m](T) = mT .$$

If  $F = \mathbb{G}_m$ , then

$$[m](T) = (1 + T)^m - 1 .$$

**Lemma 4.6.** *Let*

$$F = a_1T + a_2T^2 + \cdots \in TR[[T]] ,$$

*with  $a_1 \in R^\times$ . Then there is a unique power series  $G \in TR[[T]]$  such that  $F(G(T)) = G(F(T)) = T$ .*

*Proof.* See [Sil09, Lemma IV.2.4].  $\square$

**Lemma 4.7.** *Let  $F$  be a formal group over  $R$ , and let  $m$  be an integer that is invertible in  $R$ . Then  $[m]$  is an automorphism.*

*Proof.* It is clear that multiplication by  $m$  sends

$$T \mapsto mT + O(T^2).$$

Any such map is an automorphism of  $R[[T]]$  by Lemma 4.6.  $\square$

Now let  $K$  be a complete discretely valued nonarchimedean field, and let  $R = \mathcal{O}_K$ . Let  $\mathfrak{m}_K$  be the maximal ideal of  $\mathcal{O}_K$ , and let  $k = \mathcal{O}_K/\mathfrak{m}_K$  be the residue field.

For every  $x \in \mathfrak{m}_K$ , there is a unique continuous homomorphism  $\mathcal{O}_K[[T]] \rightarrow \mathcal{O}_K$  sending  $T \mapsto x$ , and conversely, all continuous homomorphisms  $\mathcal{O}_K[[T]] \rightarrow \mathcal{O}_K$  are of this form. Similarly, homomorphisms  $\mathcal{O}_K[[T_1, T_2]] \rightarrow \mathcal{O}_K$  are in bijection with  $\mathfrak{m}_K \times \mathfrak{m}_K$ . If  $F$  is a formal group, we will denote by  $F(\mathfrak{m}_K)$  the set  $\mathfrak{m}_K$ , with the group operation  $+_F$  given by

$$x +_F y = F(x, y).$$

*Example 4.8.* We can identify  $\widehat{\mathbb{G}}_a(\mathfrak{m}_K)$  with the additive group  $\mathfrak{m}_K$ , and there is an exact sequence

$$0 \rightarrow \mathfrak{m}_K \rightarrow \mathcal{O}_K \rightarrow k \rightarrow 0.$$

*Example 4.9.* We can identify  $\widehat{\mathbb{G}}_m(\mathfrak{m}_K)$  with the multiplicative group  $1 + \mathfrak{m}_K$ , and there is an exact sequence

$$1 \rightarrow (1 + \mathfrak{m}_K) \rightarrow \mathcal{O}_K^\times \rightarrow k^\times \rightarrow 1.$$

**Lemma 4.10.** *For each positive integer  $n$ , the operation  $+_F$  induces the usual additive group structure on  $\mathfrak{m}_K^n/\mathfrak{m}_K^{n+1}$ .*

**Lemma 4.11.** *If  $x \in F(\mathfrak{m}_K)$  has finite order, then its order is a power of the characteristic of  $k$ .*

*Proof.* This follows from Lemma 4.7.  $\square$

**Definition 4.12.** An *invariant differential* for  $F$  is an expression of the form  $P(T) dT$ , where  $P(T) \in \mathcal{O}_K[[T]]$ , such that

$$(4.13) \quad P(F(X, Y))F^{(1,0)}(X, Y) = P(X)$$

as formal power series.

**Theorem 4.14.** *Let*

$$(4.15) \quad \omega_F = F^{(1,0)}(0, T)^{-1} dT.$$

*Then the invariant differentials for  $F$  are precisely the constant multiples of  $\omega_F$ .*

*Proof.* When  $X = 0$ , the identity (4.13) becomes

$$P(Y)F^{(1,0)}(0, Y) = P(0).$$

Since  $F^{(1,0)}(0, Y)$  has constant term 1, it is invertible. So the only possible invariant differentials are multiples of  $\omega_F$ . To see that these are actually invariant differentials, differentiate the associative law

$$F(X, F(Y, Z)) = F(F(X, Y), Z).$$

with respect to  $U$ . We obtain

$$F^{(1,0)}(X, F(Y, Z)) = F^{(1,0)}(F(X, Y), Z)F^{(1,0)}(X, Y).$$

When  $X = 0$ , this becomes

$$F^{(1,0)}(0, F(Y, Z)) = F^{(1,0)}(Y, Z)F^{(1,0)}(0, Z).$$

□

**Corollary 4.16.** *If  $\phi: F \rightarrow G$  is a homomorphism of formal group laws, then*

$$\omega_G \circ \phi = \phi'(0)\omega_F.$$

**Corollary 4.17.** *Let  $F$  be a formal group over  $\mathcal{O}_K$ , and suppose the multiplication-by- $p$  map sends*

$$T \mapsto G(T).$$

*Then  $G'(T)$  is divisible by  $p$ . Equivalently,*

$$G(T) = pH(T) + I(T^p)$$

*for some formal power series  $H, I$ .*

**Theorem 4.18.** *Let  $F$  be a formal group over  $\mathcal{O}_K$ , and let  $x \in F(\mathfrak{m}_K)$ . Suppose that  $x$  has exact order  $p^n$ , meaning that  $p^n x = 0$  but  $p^{n-1}x \neq 0$ . Then  $|x| \geq |p|^{1/(p^n - p^{n-1})}$ .*

*Proof.* We use induction on  $n$ . Suppose  $n = 1$ . Let  $G(T)$  be as in Corollary 4.17. Then  $x$  satisfies  $G(x) = 0$ . The linear term of  $G(x)$  is  $px$ . All other terms with exponent not divisible by  $p$  are also multiples of  $p$ , so they have strictly smaller absolute values. Among the terms with exponent divisible by  $p$ , the largest possible absolute value is  $|x|^p$ . So we must have  $|px| \leq |x|^p$ . We can rewrite this equality as  $|x| \geq |p|^{1/(p-1)}$ .

Now assume that all points of exact order  $n$  satisfy  $|x| \geq |p|^{1/(p^n - p^{n-1})}$ , and let  $y$  be a point of exact order  $n+1$ . In order for any of the terms of  $G(y)$  to have absolute value greater than or equal to  $|x|$ , we must have  $|y| \geq |p|^{1/(p^{n+1} - p^n)}$ . This completes the induction. □

## 5. ELLIPTIC CURVES, CONTINUED

**Definition 5.1.** Let  $(E, O), (E', O')$  be elliptic curves over  $K$ . A morphism  $(E, O) \rightarrow (E', O')$  is a morphism  $E \rightarrow E'$  sending  $O$  to  $O'$ .

**Lemma 5.2.** *Any morphism  $\phi: (E, O) \rightarrow (E', O')$  of elliptic curves is a group homomorphism.*

*Proof.* If  $\phi$  is constant, then it is a group homomorphism. Otherwise,  $\phi$  is a finite locally free morphism, so there is an induced homomorphism  $\phi_*: \text{Pic}_{E/K}^0 \rightarrow \text{Pic}_{E'/K}^0$ . Since we can identify  $E, E'$  with  $\text{Pic}_{E/K}^0, \text{Pic}_{E'/K}^0$ , respectively,  $\phi$  must also be a group homomorphism. □

We will write  $\text{Hom}(E, E')$  for the set of morphisms  $(E, O) \rightarrow (E', O')$ , and  $\text{End}(E)$  for  $\text{Hom}(E, E')$ . Lemma 5.2 implies that  $\text{End}(E)$  is a (not necessarily commutative) ring.

**Lemma 5.3.** *Let  $E$  be an elliptic curve.*

- (1)  *$\text{End } E$  has no zero divisors.*

(2) For any nonzero integer  $n$ , the multiplication by  $n$  map  $E \rightarrow E$  is not zero.

*Proof.* For the first item, observe that any nonzero element of  $\text{End } E$  is surjective, and the composition of two surjections is a surjection.

For the second item, see [Sil09, Proposition III.4.2(a)].  $\square$

Since the Picard functor is contravariant, any homomorphism  $\phi: E \rightarrow E'$  also induces a homomorphism  $\hat{\phi}: \text{Pic}_{E'/K}^0 \rightarrow \text{Pic}_{E/K}^0$ , or equivalently, a homomorphism  $\hat{\phi}: E' \rightarrow E$ .

**Lemma 5.4.**

- (1) For any  $\phi \in \text{Hom}(E, E')$ ,  $\hat{\phi}\phi$  is multiplication by  $\deg \phi$ .
- (2) For any  $\phi \in \text{Hom}(E, E')$ ,  $\psi \in \text{Hom}(E', E'')$ ,  $\widehat{\psi\phi} = \hat{\phi}\hat{\psi}$ .
- (3) For any  $\phi, \psi \in \text{Hom}(E, E')$ ,  $\widehat{\psi + \phi} = \hat{\psi} + \hat{\phi}$ .
- (4) For any integer  $n$ , the image of  $n$  in  $\text{Hom}(E, E)$  is self dual.
- (5) For any  $\phi \in \text{Hom}(E, E')$ ,  $\deg \hat{\phi} = \deg \phi$ .
- (6)  $\hat{\hat{\phi}} = \phi$

*Proof.* See [Sil09, Theorem 6.1 and 6.2].  $\square$

**Corollary 5.5.** The degree map  $\deg: \text{Hom}(E, E') \rightarrow \mathbb{Z}$  is a positive definite quadratic form.

**Corollary 5.6.** For any elliptic curve  $E$ , the multiplication by  $N$  map has degree  $N^2$ .

For any positive integer  $N$ , let

$$E(K^{\text{sep}})[N] = \{P \in E(K^{\text{sep}}) | NP = 0\}.$$

**Corollary 5.7.**

If the characteristic of  $K$  does not divide  $N$ , then  $E(K^{\text{sep}})[N] \cong (\mathbb{Z}/N\mathbb{Z})^2$ .

Note that  $\text{Gal}(K^{\text{sep}}/K)$  acts on  $E(K^{\text{sep}})[N]$ .

For any prime  $p$ , define the Tate module

$$T_p(E) = \varprojlim_n E(K^{\text{sep}})[p^n].$$

If the characteristic of  $K$  is different from  $p$ , then  $T_p(E)$  is a free  $\mathbb{Z}_p$ -module of rank 2

**Theorem 5.8.** Suppose the characteristic of  $K$  is different from  $p$ . Then the natural map

$$\text{Hom}(E, E') \otimes \mathbb{Z}_p \rightarrow \text{Hom}_{\mathbb{Z}_p}(T_p(E), T_p(E'))$$

is injective.

*Proof.* See [Sil09, Theorem III.7.4].  $\square$

**Corollary 5.9.**  $\text{Hom}(E, E')$  is a free  $\mathbb{Z}$ -module of rank at most 4.

**Corollary 5.10.**  $\text{End}(E) \otimes \mathbb{Q}$  is isomorphic to one of the following:

- (1)  $\mathbb{Q}$ ;
- (2) An imaginary quadratic extension of  $\mathbb{Q}$ ;
- (3) A quaternion algebra over  $\mathbb{Q}$ , ramified at  $p = \text{char } K$  and  $\infty$ , and at no other places.

A quaternion algebra over  $\mathbb{Q}$  is a division algebra  $D$  with center  $\mathbb{Q}$  satisfying  $[D : \mathbb{Q}] = 4$ . By “ramified at  $p$  and  $\infty$ ”, we mean that  $D \otimes \mathbb{Q}_p$  and  $D \otimes \mathbb{R}$  are division algebras, while  $D \otimes \mathbb{Q}_\ell \cong M_2(\mathbb{Q}_\ell)$  is the ring of  $2 \times 2$  matrices for all  $\ell \neq p$ .

*Remark 5.11.* If  $L$  is an extension of  $K$ , then  $\text{End}(E_L)$  can be larger than  $\text{End}(E)$ . For example, if  $E$  is the elliptic curve  $y^2 = x^3 - x$  over  $\mathbb{Q}$ , then  $\text{End } E = \mathbb{Z}$ , but  $\text{End } E_{\mathbb{Q}(i)} = \mathbb{Z}[i]$ , where  $i$  acts by  $(x, y) \mapsto (-x, iy)$ .

*Remark 5.12.* The action of  $G_K$  on  $T_p(E)$  commutes with endomorphisms of  $E$ . From the classification of Theorem 5.10, we deduce:

- If  $\text{End}(E) \otimes \mathbb{Q}$  is an imaginary quadratic extension  $F$  of  $\mathbb{Q}$ , then the map  $G_K \rightarrow \text{End } T_p(E)$  factors through  $(\mathbb{Z}_p \otimes \mathcal{O}_F)^\times$ .
- If  $\text{End}(E) \otimes \mathbb{Q}$  is a quaternion algebra, then  $G_K$  acts by scalars on  $T_p(E)$ .

*Remark 5.13.* Tate modules are closely related to étale cohomology. If  $E$  is an elliptic curve over a field  $K$ , then

$$H_{\text{ét}}^i(E_{\bar{K}}, \mathbb{Z}_p) = \begin{cases} \mathbb{Z}_p & (\text{with trivial } G_K\text{-action}), & i = 0 \\ \text{Hom}_{\mathbb{Z}_p}(T_p(E), \mathbb{Z}_p), & i = 1 \\ \mathbb{Z}_p(-1), & i = 2 \\ 0, & \text{otherwise.} \end{cases}$$

## 6. ELLIPTIC CURVES, CONTINUED

Étale cohomology is supposed to satisfy many of the same properties as singular cohomology. In particular, it is expected to satisfy Poincaré duality. For elliptic curves, this means that there should be an antisymmetric cup product map

$$H_{\text{ét}}^1(E_{\bar{K}}, \mathbb{Z}_p) \times H_{\text{ét}}^1(E_{\bar{K}}, \mathbb{Z}_p) \rightarrow H_{\text{ét}}^2(E_{\bar{K}}, \mathbb{Z}_p)$$

that is a perfect pairing. Since

$$H_{\text{ét}}^1(E_{\bar{K}}, \mathbb{Z}_p) = T_p(E)^*$$

$$H_{\text{ét}}^2(E_{\bar{K}}, \mathbb{Z}_p) = \mathbb{Z}_p(-1),$$

specifying the cup product map is equivalent to specifying a perfect pairing

$$T_p(E) \times T_p(E) \rightarrow \mathbb{Z}_p(1).$$

This map can be constructed using the Weil pairing.

For each  $N$  and each extension  $L$  of  $K$ , there is a Weil pairing

$$e_N : E(L)[N] \times E(L)[N] \rightarrow \mu_N(L).$$

It can be defined as follows. Let  $P, Q \in E(L)[N]$ . Then

$$\sum_{m=0}^{N-1} ([P + mQ] - [mQ])$$

is a principal divisor. Let  $g$  be a function with this divisor. Let  $T_Q g$  denote the translation of  $g$  by  $Q$ . Then  $T_Q g$  and  $g$  have the same divisor, so  $T_Q g = \omega g$  for some constant  $\omega$ . Since  $T_Q^N$  is the identity,  $\omega$  is an  $N$ th root of unity. We define

$$e_N(P, Q) = \omega.$$

**Theorem 6.1.** *The Weil pairing has the following properties:*

(1) *It is bilinear:*

$$e_N(P + Q, R) = e_N(P, R) + e_N(Q, R), \quad e_N(P, Q + R) = e_N(P, Q) + e_N(P, R)$$

(2) *It is alternating:*

$$e_N(P, P) = 1$$

(3) *If  $N$  does not divide the characteristic of  $K$ , then it is a perfect pairing.*

(4) *It is  $\text{Aut}(L/K)$ -invariant: for  $\sigma \in \text{Aut}(L/K)$ ,*

$$e_N(P^\sigma, Q^\sigma) = e_N(P, Q)^\sigma$$

(5) *The Weil pairings for various  $N$  are compatible: if  $P \in E(L)[MN], Q \in E(L)[M]$ ,*

$$e_{MN}(P, Q) = e_M(NP, Q).$$

If the characteristic of  $K$  is not  $p$ , then we can take inverse limits to get a perfect pairing

$$T_p(E) \times T_p(E) \rightarrow \mathbb{Z}_p(1).$$

(Recall that  $\mathbb{Z}_p(1) = \varprojlim_n \mu_{p^n}(K^{\text{sep}})$ , where  $\mu_{p^n}(K^{\text{sep}})$  is the group of  $p^n$ th roots of unity of  $K^{\text{sep}}$ .)

We are particularly interested in elliptic curves over  $p$ -adic fields. We still need to define what these are.

**Definition 6.2.** A *nonarchimedean field* is a field  $K$  that is complete with respect to a norm  $|\cdot|: K \rightarrow \mathbb{R}^{\geq 0}$  such that:

$$|xy| = |x||y| \text{ and } |x + y| \leq \max(|x|, |y|) \text{ for all } x, y \in K$$

$$|0| = 0, \quad |1| = 1$$

$$0 < |x| < 1 \text{ for some } x \in K$$

We will write

$$\mathcal{O}_K = \{x \in K \mid |x| \leq 1\}$$

$$\mathfrak{m}_K = \{x \in K \mid |x| < 1\}.$$

We say that  $K$  is *discretely valued* if the image of  $K^\times$  under  $|\cdot|$  is a discrete subset of  $\mathbb{R}^{>0}$ . Equivalently,

$$\sup_{x \in \mathfrak{m}_K} |x| < 1.$$

*Example 6.3.* Let  $p$  be a prime. We can define a norm  $|\cdot|$  on  $\mathbb{Q}$  by  $|p^m \frac{r}{s}| = p^{-m}$  for all integers  $r, s$  not divisible by  $p$ . Then  $\mathbb{Q}_p$  is defined to be the completion of  $\mathbb{Q}$  with respect to this norm. It is a nonarchimedean field.

**Lemma 6.4.** *Let  $K$  be a nonarchimedean field, and let  $L/K$  be an algebraic extension. Then there is a unique nonarchimedean absolute value on  $L$  extending the absolute value on  $K$ .*

*Proof.* See [Bos14, Theorem A.3].  $\square$

*Remark 6.5.* If  $L/K$  is finite, and the norms on  $K, L$  are denoted by  $|\cdot|_K, |\cdot|_L$ , respectively, then

$$|x|_L = |N_{L/K}(x)|_K^{1/[L:K]}.$$

It takes some work to show that the right-hand side satisfies the triangle inequality.

**Corollary 6.6.** *Any finite extension of  $\mathbb{Q}_p$  is nonarchimedean, as is the completion of any infinite algebraic extension of  $\mathbb{Q}_p$ .*

**Definition 6.7.** A field  $K$  is *perfect* if every irreducible polynomial in  $K[x]$  is separable.

**Proposition 6.8.** *Every field of characteristic 0 is perfect. A field of characteristic  $p > 0$  is perfect iff every element is a  $p$ th power.*

For the remainder of the lecture, we will fix a prime  $p$ .

**Definition 6.9.** A  $p$ -adic field is a discretely valued nonarchimedean field  $K$  of characteristic zero, such that its residue field  $\mathcal{O}_K/\mathfrak{m}_K$  is perfect of characteristic  $p$ .

*Example 6.10.* The field  $\mathbb{Q}_p$  is a  $p$ -adic field, as is any finite extension of  $\mathbb{Q}_p$ . The completion of an infinite algebraic extension of  $\mathbb{Q}_p$  may or may not be a  $p$ -adic field.

Let  $K$  be a  $p$ -adic field, and let  $k = \mathcal{O}_K/\mathfrak{m}_K$  be its residue field. There is a surjection  $\text{Gal}(\overline{K}/K) \rightarrow \text{Gal}(\overline{k}/k)$ . The kernel is called the *inertia subgroup* of  $\text{Gal}(\overline{K}/K)$ .

**Definition 6.11.** We say that a representation of  $\text{Gal}(\overline{K}/K)$  is *unramified* if it factors through  $\text{Gal}(\overline{k}/k)$ . Otherwise, we say that it is ramified.

*Example 6.12.* For  $\ell \neq p$ ,  $\mathbb{Z}_\ell(1)$  is unramified, since all  $\ell$ -power roots of unity in  $\overline{K}$  have distinct images in  $\overline{k}$ . But  $\mathbb{Z}_p(1)$  is ramified since all  $p$ -power roots of unity in  $\overline{K}$  map to 1 in  $\overline{k}$ .

## 7. ELLIPTIC CURVES OVER $p$ -ADIC FIELDS

Let  $K$  be a  $p$ -adic field.

**Definition 7.1.** We say that a proper smooth variety  $X$  over  $K$  has *good reduction* if it can be extended to a proper smooth scheme of finite type over  $\mathcal{O}_K$ .

Otherwise, we say that it has *bad reduction*.

*Example 7.2.* If  $p \neq 2$ , then the elliptic curve  $E$  over  $K$  defined by  $y^2 = x(x-1)(x+1)$  has good reduction, since  $y^2 = x(x-1)(x+1)$  defines a nonsingular curve over  $\mathcal{O}_K$ . It is possible to show that  $E$  does not have good reduction at 2.

Recall that an elliptic curve over  $\mathcal{O}_K$  is a pair  $(E, O)$ , where  $E$  is a proper smooth scheme over  $\mathcal{O}_K$  such that  $E_K$  and  $E_k$  are geometrically connected curves of genus 1, and  $O$  is an  $\mathcal{O}_K$ -point of  $E$ .

**Lemma 7.3.** *Let  $X$  be a proper scheme over  $\mathcal{O}_K$ . Then the restriction map  $X(\mathcal{O}_K) \rightarrow X(K)$  is a bijection. In particular, there is a natural reduction map  $X(K) \rightarrow X(k)$ .*

*Proof.* This follows from the valuative criterion of properness.  $\square$

**Theorem 7.4.** *If  $E$  is an elliptic curve over  $\mathcal{O}_K$ , then there is an exact sequence*

$$0 \rightarrow \hat{E}(\mathfrak{m}_K) \rightarrow E(K) \rightarrow E(k) \rightarrow 0.$$

**Corollary 7.5.** *If  $E$  is an elliptic curve over  $\mathcal{O}_K$ , and  $\ell$  is a prime different from  $\text{char } K$ , then the reduction maps*

$$E(K)[\ell^n] \rightarrow E(k)[\ell^n]$$

and

$$T_\ell(E_K) \rightarrow T_\ell(E_k)$$

are isomorphisms.

*Proof.* For each  $n$ , there is an exact sequence

$$\hat{E}(\mathfrak{m}_K)[\ell^n] \rightarrow E(K)[\ell^n] \rightarrow E(k)[\ell^n] \rightarrow \hat{E}(\mathfrak{m}_K)/\ell^n.$$

But multiplication by  $\ell$  is invertible on  $\hat{E}(\mathfrak{m}_K)$ , so the outer terms are zero. So the maps  $E(K)[\ell^n] \rightarrow E(k)[\ell^n]$  are isomorphisms. Taking the direct limit over algebraic extensions of  $K$  and inverse limit over  $n$  shows that  $T_\ell(E_K) \rightarrow T_\ell(E_k)$  is an isomorphism.  $\square$

**Corollary 7.6.** *If an elliptic curve  $E$  over  $K$  has good reduction, then  $T_\ell(E)$  is unramified for every prime  $\ell \neq p$ .*

The converse is also true, although we will not give a proof.

**Theorem 7.7** (Néron–Ogg–Shafarevich). *Let  $\ell$  be a prime different from  $p$ . An elliptic curve  $E$  over  $K$  has good reduction if and only if  $T_\ell(E)$  is unramified. More generally, an abelian variety  $X$  over  $K$  has good reduction if and only if  $T_\ell(X)$  is unramified.*

By contrast,  $T_p(X)$  is never unramified. If  $X$  is an elliptic curve, then it follows from the Weil pairing that  $\wedge^2 T_\ell(X) \cong \mathbb{Z}_\ell(1)$ , and we saw that this representation is ramified if  $\ell = p$ .

However, it is possible to determine if  $X$  has good reduction from  $T_p(X)$ . The criterion uses the notion of a crystalline representation, which we will define in a later lecture.

**Theorem 7.8.** *An abelian variety  $X$  over  $K$  has good reduction if and only if  $T_p(X) \otimes_{\mathbb{Z}_p} \mathbb{Q}_p$  is crystalline.*

Before we move on, I want to do a little bit of  $p$ -adic analysis. Previously, we claimed that  $\text{Gal}(\mathbb{Q}_p(\mu_{p^n})/\mathbb{Q}_p) \cong (\mathbb{Z}/p^n\mathbb{Z})^\times$ . We will now give a proof using the theory of Newton polygons. (It is possible to give a more elementary proof using Eisenstein’s criterion, but the method of Newton polygons is more general, so it is useful to know.)

Let  $K$  be a nonarchimedean field with absolute value  $|\cdot|$ . Define a “valuation”  $v: K \rightarrow \mathbb{R} \cup \{\infty\}$  by  $v(x) = -\log|x|$ .

**Definition 7.9.** Let  $f(x) = \sum_{i=0}^n a_i x^i$  be a polynomial with  $a_0, a_n \neq 0$ . The *Newton polygon* of  $f$  is the lower envelope of the points  $(i, v(a_i)) \in \mathbb{R}^2$ .

**Proposition 7.10.** *Let*

$$f(x) = \prod_{i=1}^n (x - \alpha_i)$$

*with*

$$v(\alpha_1) \leq v(\alpha_2) \leq \cdots \leq v(\alpha_n)$$

*Then the slope of the segment of the Newton polygon beginning at  $x = n - i$  and ending at  $x = n - i + 1$  is  $-v(\alpha_i)$ .*

*Proof.* Let  $1 \leq m \leq n$ . Then the coefficient of  $x^{n-m}$  is the sum of all products of  $m$  of the  $\alpha_i$ ’s. Each term in the sum has valuation at least  $\sum_{i=1}^m v(\alpha_i)$ , so the valuation of the sum is at least this large. If  $v(\alpha_m) < v(\alpha_{m+1})$ , then only one term has this valuation, so the sum is exactly this large.  $\square$



**Corollary 7.11.** *If  $f(x)$ ,  $g(x)$  are any polynomials with nonzero constant term, then the Newton polygon of  $f(x)g(x)$  is obtained from the Newton polygons of  $f(x)$  and  $g(x)$  by rearranging segments in order of slope.*

**Proposition 7.12.** *If a polynomial  $f(x)$  is irreducible, then its Newton polygon has only one slope. Conversely, if  $f$  has degree  $n$  and the  $y$ -coordinates of the Newton polygon at  $x = 1, \dots, n-1$  are not in the image of  $v$ , then  $f$  is irreducible.*

*Proof.* The first claim follows from Hensel's lemma; see [Bos14, Lemma 4] for the statement and proof of the lemma. The second claim follows from Corollary 7.11.  $\square$

*Example 7.13.* The group  $\text{Gal}(\mathbb{Q}_p(\mu_{p^n})/\mathbb{Q}_p)$  must send a  $p^n$ th root of unity to another  $p^n$ th root of unity, so there is a natural injection  $\text{Gal}(\mathbb{Q}_p(\mu_{p^n})/\mathbb{Q}_p) \hookrightarrow (\mathbb{Z}/p^n\mathbb{Z})^\times$ . To show that this map is an isomorphism, it suffices to show that both groups have the same number of elements.

We claim that the polynomial

$$\frac{(1+T)^{p^n} - 1}{(1+T)^{p^{n-1}} - 1}$$

is irreducible. It has integer coefficients, and the coefficient of the constant term is  $p$ . If we normalize  $v$  so that  $v(p) = 1$ , then the Newton polygon is the line segment from  $(0, 0)$  to  $(p^n - p^{n-1}, 1)$ . By Proposition 7.12, the polynomial is irreducible. So

$$|\text{Gal}(\mathbb{Q}_p(\mu_{p^n})/\mathbb{Q}_p)| = [\mathbb{Q}_p(\mu_{p^n}) : \mathbb{Q}_p] = p^n - p^{n-1} = |(\mathbb{Z}/p^n\mathbb{Z})^\times|.$$

## 8. GALOIS GROUPS OF FIELDS OF CHARACTERISTIC $p$ AND $\varphi$ -MODULES

It is difficult to write down Galois groups explicitly, which in turn makes it difficult to write down Galois representations explicitly. To deal with this problem, we will introduce  $\varphi$ -modules and  $(\varphi, \Gamma)$ -modules, which can be described more explicitly. We will show that categories of these modules are equivalent to categories of Galois representations.

We will exploit the fact that the absolute Galois groups of  $p$ -adic fields are closely related to the absolute Galois groups of characteristic  $p$  fields. For example, we have the following result, which will be proved in a later lecture.

**Theorem 8.1.** *The absolute Galois groups of  $\mathbb{Q}_p(\mu_{p^\infty})$  and  $\mathbb{F}_p((t))$  are isomorphic (as topological groups).*

Now let  $E$  be a field of characteristic  $p$ . Let  $G_E = \text{Gal}(E^{\text{sep}}/E)$ . Let  $\varphi_E: E \rightarrow E$  be the Frobenius map  $x \mapsto x^p$ .

Given an  $E$ -module  $M$ , we write  $\varphi_E^*(M)$  for its Frobenius pullback  $E \otimes_{\varphi_E, E} M$ . Any  $\varphi_E$ -semilinear map  $\varphi_M: M \rightarrow M$  determines an  $E$ -linear map  $\varphi_E^*(M) \rightarrow M$  by  $e \otimes m \mapsto e\varphi_M(m)$ .

**Definition 8.2.** A  $\varphi$ -module over  $E$  is a pair  $(M, \varphi_M)$ , where  $M$  is a finite-dimensional  $E$ -vector space and  $\varphi_M$  is a  $\varphi_E$ -semilinear endomorphism. We say that  $(M, \varphi_M)$  is *étale* if the  $E$ -linear map  $\varphi_E^*(M) \rightarrow M$  induced by  $\varphi$  is an isomorphism (equivalently, the image of  $\varphi_M$  generates  $M$  as an  $E$ -module).

We will denote the category of étale  $\varphi$ -modules over  $E$  by  $\varphi\text{-Mod}_E^{\text{ét}}$ .

Let  $\text{Rep}_{\mathbb{F}_p}(G_E)$  denote the category of continuous finite-dimensional  $\mathbb{F}_p$ -vector space representations of  $G_E$ .

**Theorem 8.3.** *The functor  $D_E: \text{Rep}_{\mathbb{F}_p}(G_E) \rightarrow \varphi\text{-Mod}_E^{\text{ét}}$  defined by*

$$V \mapsto (V \otimes_{\mathbb{F}_p} E^{\text{sep}})^{G_E}.$$

*and the functor  $V_E: \varphi\text{-Mod}_E^{\text{ét}} \rightarrow \text{Rep}_{\mathbb{F}_p}(G_E)$  defined by*

$$M \mapsto (M \otimes_E E^{\text{sep}})^{\varphi=1}.$$

*determine an equivalence of categories between  $\text{Rep}_{\mathbb{F}_p}(G_E)$  and  $\varphi\text{-Mod}_E^{\text{ét}}$ .*

*Remark 8.4.* There are a few advantages to working with  $\varphi$ -modules rather than Galois representations. One is that Galois groups are difficult to describe explicitly, while a  $\varphi$ -module is described by a matrix with coefficients in  $E$ . Another reason is that  $\varphi$ -modules are better suited to working with families of Galois representations. For example, for any  $\alpha \in \mathbb{F}_q^\times$ , there is a representation  $\text{Gal}(\overline{\mathbb{F}_p}/\mathbb{F}_p) \rightarrow \mathbb{F}_q^\times$  sending the Frobenius to  $\alpha$ . We might like to combine these into a representation  $\text{Gal}(\overline{\mathbb{F}_p}/\mathbb{F}_p) \rightarrow \mathbb{F}_p[t, t^{-1}]^\times$ , but this does not work because we cannot raise  $t$  to powers in  $\widehat{\mathbb{Z}}$ . On the other hand, the  $\varphi$ -module corresponding to  $\alpha$  is a 1-dimensional  $\mathbb{F}_q$ -vector space on which  $\varphi$  acts by  $\alpha^{-1}$ . These combine nicely into a free  $\mathbb{F}_p[t, t^{-1}]$ -module of rank 1 on which  $\varphi$  acts by  $t^{-1}$ .

*Remark 8.5.* One might think of Theorem 8.3 as a characteristic  $p$  version of the Riemann-Hilbert correspondence, with the Frobenius action replacing the connection. For a more geometric analogue, see [Kat73, Proposition 4.1.1].

**Lemma 8.6** (Galois descent). *Let  $L/K$  be a Galois extension of fields (of any characteristic). Let  $V$  be a finite-dimensional  $L$ -vector space equipped with a semilinear  $\text{Gal}(L/K)$ -action (meaning that the identities*

$$\sigma(v_1 + v_2) = \sigma(v_1) + \sigma(v_2)$$

$$\sigma(\lambda v_1) = \sigma(\lambda)\sigma(v_1)$$

*hold for each  $\sigma \in \text{Gal}(L/K)$ ,  $v_1, v_2 \in V$ ,  $\lambda \in L$ ).*

*Suppose that for each  $v \in V$ , the stabilizer of  $v$  in  $\text{Gal}(L/K)$  is open. Then the natural map*

$$L \otimes_K V^{\text{Gal}(L/K)} \rightarrow V$$

*is an isomorphism.*

*Proof.* The case of  $L/K$  finite is [Sil09, Lemma II.5.8.1]. In general, we use the fact that a basis for  $V$  has open stabilizer to reduce to the finite case.  $\square$

*Proof of Theorem 8.3.* Let  $V \in \text{Rep}_{\mathbb{F}_p}(G_E)$ . We will check that  $D_E(V) \in \varphi\text{-Mod}_E^{\text{ét}}$ , and that there is a natural isomorphism  $V_E(D_E(V)) \xrightarrow{\sim} V$ . By Galois descent, the  $\varphi$ - and  $G_E$ -equivariant map

$$(8.7) \quad D_E(V) \otimes_E E^{\text{sep}} \rightarrow V \otimes_{\mathbb{F}_p} E^{\text{sep}}$$

is an isomorphism. Therefore,  $\dim_E D_E(V) = \dim_{\mathbb{F}_p} V$ ; in particular,  $D_E(V)$  is finite dimensional.

To show that  $D_E(V)$  is étale, we just need to check that the matrix of Frobenius in some (equivalently, any) basis is invertible. By base change, the matrix of Frobenius on  $D_E(V)$  is invertible iff the matrix of Frobenius on  $D_E(V) \otimes_E E^{\text{sep}} = V \otimes_{\mathbb{F}_p} E^{\text{sep}}$  is invertible iff the matrix of Frobenius on  $V$  is invertible. The action of Frobenius on  $V$  is the identity.

Taking  $\varphi$ -invariants of (8.7) gives an isomorphism  $V_E(D_E(V)) \xrightarrow{\sim} V$ .

Now let  $M \in \varphi\text{-Mod}_E^{\text{ét}}$ . We want to show that the natural map

$$(8.8) \quad E^{\text{sep}} \otimes_{\mathbb{F}_p} V_E(M) \rightarrow E^{\text{sep}} \otimes_E M$$

is an isomorphism. First we will show that it is injective. It suffices to show that if some vectors in  $V_E(M) = (E^{\text{sep}} \otimes_E M)^{\varphi=1}$  are linearly independent over  $\mathbb{F}_p$ , then they are also linearly independent over  $E^{\text{sep}}$ . Suppose that there is a minimal counterexample  $v_1, \dots, v_r \in V_E(M)$ , with  $\sum_{i=1}^r a_i v_i = 0$  for  $a_i \in E^{\text{sep}}$ . WLOG we may take  $a_1 = 1$ . Using  $\varphi(v_i) = v_i$  and  $\varphi(a_1) = a_1$ , we obtain  $0 = \sum_{i=2}^r (a_i - \varphi(a_i)) v_i$ . By minimality of the counterexample, we must have  $(a_i - \varphi(a_i)) = 0$  for all  $i$ . Hence  $a_i \in \mathbb{F}_p$  for all  $i$ , which is a contradiction. Note that we did not need to use the fact that  $M$  is étale to prove injectivity.

Now we show that (8.8) is surjective. Let  $c_{ij}$  be the matrix coefficients of Frobenius in some basis. Let  $X$  be the scheme over  $E$  defined by the equations

$$x_i^p = \sum_j c_{ij} x_j.$$

Surjectivity of (8.8) is equivalent to  $|X(E^{\text{sep}})| = p^{\dim_E M}$ . Since  $X$  is finite locally free over  $\text{Spec } E$  of degree  $p^{\dim_E M}$ , it suffices to show that  $X$  is étale over  $E$ , or equivalently that  $\Omega_{X/E} = 0$ . The module  $\Omega_{X/E}$  is generated by the  $dx_i$  subject to the relations  $\sum_j c_{ij} x_j = 0$ . Since the  $c_{ij}$  define an invertible matrix,  $\Omega_{X/E} = 0$ . This concludes the proof that (8.8) is an isomorphism.

From (8.8), we see that  $V_E(M)$  is finite dimensional over  $\mathbb{F}_p$ . Then  $V_E(M) = (M \otimes_E F)^{\varphi=1}$  for some finite separable extension  $F/E$ , so the  $G_E$ -action on  $V_E(M)$  is continuous. Therefore  $V_E(M) \in \text{Rep}_{\mathbb{F}_p} G_E$ . Taking Galois invariants of (8.8) gives an isomorphism  $D_E(V_E(M)) \xrightarrow{\sim} M$ . Hence we have shown that the functors  $D_E$  and  $V_E$  are essential inverses of each other.  $\square$

## 9. GENERALIZATIONS OF $\varphi$ -MODULES, PERFECTOID FIELDS

As before, let  $E$  be a field of characteristic  $p$ . We will now turn our attention to  $\mathbb{Z}_p$ -representations of  $G_E$ . Let  $\text{Rep}_{\mathbb{Z}_p} G_E$  denote the category of finitely generated (not necessarily free)  $\mathbb{Z}_p$ -modules with continuous  $G_E$ -action.

**Definition 9.1.** Let  $E$  be a field of characteristic  $p$ . A *Cohen ring* for  $E$  is a complete discrete valuation ring  $\mathcal{O}_{\mathcal{E}}$  such that the residue field of  $\mathcal{O}_{\mathcal{E}}$  is  $E$ , and  $p$  is a uniformizer of  $\mathcal{O}_{\mathcal{E}}$ .

**Theorem 9.2** ([Mat86, Theorems 29.1 and 29.2]). *Any field  $E$  of characteristic  $p$  admits a Cohen ring  $\mathcal{O}_{\mathcal{E}}$ . The Cohen ring is unique up to isomorphism. The Frobenius automorphism of  $E$  lifts to an automorphism of  $\mathcal{O}_{\mathcal{E}}$ .*

*Example 9.3.* If  $E = \mathbb{F}_{p^n}$ , then  $\mathcal{O}_{\mathcal{E}}$  is the ring of integers of  $\mathbb{Q}_{p^n}$ , the unramified extension of  $\mathbb{Q}_p$  of degree  $n$ . More generally, if  $E$  is perfect, then  $\mathcal{O}_{\mathcal{E}} \cong W(E)$ , the ring of  $p$ -typical Witt vectors over  $E$ . If  $E = \mathbb{F}_p((T))$ , then we can take

$$\mathcal{O}_{\mathcal{E}} = \left\{ \sum_{n=-\infty}^{\infty} a_n T^n \mid a_n \in \mathbb{Z}_p, \lim_{n \rightarrow -\infty} a_n = 0 \right\}.$$

A commonly used choice of Frobenius action is  $T \mapsto (1+T)^p - 1$ .

**Lemma 9.4.** *There is an equivalence of categories between étale  $E$ -algebras and étale  $\mathcal{O}_{\mathcal{E}}$ -algebras.*

In particular,  $E^{\text{sep}}$  is a direct limit of étale  $E$  algebras. The corresponding direct limit of étale  $\mathcal{O}_E$ -algebras is called  $\mathcal{O}_E^{\text{sh}}$ , the strict Henselization of  $\mathcal{O}_E$ .

**Definition 9.5.** The category  $\varphi\text{-Mod}_{\mathcal{O}_E}^{\text{ét}}$  of étale  $\varphi$ -modules over  $\mathcal{O}_E$  consists of pairs  $(M, \varphi_M)$  where  $M$  is a finitely generated  $\mathcal{O}_E$ -module and  $\varphi_M$  is a  $\varphi$ -semilinear endomorphism of  $M$  such that  $\varphi_{\mathcal{O}_E}^*(M) \rightarrow M$  is an isomorphism.

**Definition 9.6.**

Now let  $\check{\mathcal{O}}_E = \widehat{\mathcal{O}_E^{\text{sh}}}$  be the completion of the strict henselization of  $\mathcal{O}_E$ . There is a unique continuous Frobenius on  $\check{\mathcal{O}}_E$  that extends the Frobenius on  $\mathcal{O}_E$  and  $E^{\text{sep}}$ .

**Theorem 9.7.** The functor  $D_{\mathcal{O}_E}: \text{Rep}_{\mathbb{Z}_p} G_E \rightarrow \varphi\text{-Mod}_{\mathcal{O}_E}^{\text{ét}}$  defined by

$$V \mapsto (V \otimes_{\mathbb{Z}_p} \check{\mathcal{O}}_E)^{G_E}$$

and the functor  $V_{\mathcal{O}_E}: \varphi\text{-Mod}_{\mathcal{O}_E}^{\text{ét}} \rightarrow \text{Rep}_{\mathbb{Z}_p} G_E$  defined by

$$M \mapsto (M \otimes_{\mathcal{O}_E} \check{\mathcal{O}}_E)^{\varphi=1}$$

determine an equivalence of categories between  $\text{Rep}_{\mathbb{Z}_p} G_E$  and  $\varphi\text{-Mod}_{\mathcal{O}_E}^{\text{ét}}$ .

Finally, we consider  $\mathbb{Q}_p$ -representations of  $G_E$ . Let  $\text{Rep}_{\mathbb{Q}_p} G_E$  denote the category of finite-dimensional  $\mathbb{Q}_p$ -vector space representation of  $G_E$ . Let  $\mathcal{E} := \mathcal{O}_E[1/p]$ ,  $\check{\mathcal{E}} := \check{\mathcal{O}}_E[1/p]$ .

**Definition 9.8.** The category  $\varphi\text{-Mod}_{\mathcal{E}}^{\text{ét}}$  of étale  $\varphi$ -modules over  $\mathcal{E}$  consists of pairs  $(M, \varphi_M)$  where  $M$  is a finite-dimensional  $\mathcal{E}$ -vector space and  $\varphi_M$  is a  $\varphi$ -semilinear endomorphism of  $M$  such that  $\varphi_{\mathcal{E}}^*(M) \rightarrow M$  is an isomorphism, and  $M$  admits a  $\varphi_M$ -stable  $\mathcal{O}_E$ -lattice.

**Theorem 9.9.** The functor  $D_{\mathcal{E}}: \text{Rep}_{\mathbb{Q}_p} G_E \rightarrow \varphi\text{-Mod}_{\mathcal{E}}^{\text{ét}}$  defined by

$$V \mapsto (V \otimes_{\mathbb{Q}_p} \check{\mathcal{E}})^{G_E}$$

and the functor  $V_{\mathcal{E}}: \varphi\text{-Mod}_{\mathcal{E}}^{\text{ét}} \rightarrow \text{Rep}_{\mathbb{Q}_p} G_E$  defined by

$$M \mapsto (M \otimes_{\mathcal{E}} \check{\mathcal{E}})^{\varphi=1}$$

determine an equivalence of categories between  $\text{Rep}_{\mathbb{Q}_p} G_E$  and  $\varphi\text{-Mod}_{\mathcal{E}}^{\text{ét}}$ .

The Galois group  $\text{Gal}(\overline{\mathbb{Q}_p}/\mathbb{Q}_p)$  contains the closed normal subgroup  $\text{Gal}(\overline{\mathbb{Q}_p}/\mathbb{Q}_p(\mu_{p^\infty})) \cong \text{Gal}(\mathbb{F}_p((t))^{\text{sep}}/\mathbb{F}_p((t)))$ . Moreover, this isomorphism can be extended to a map

$$\text{Gal}(\overline{\mathbb{Q}_p}/\mathbb{Q}_p) \hookrightarrow \text{Aut}(\mathbb{F}_p((t))^{\text{sep}}).$$

This motivates us to consider the following setup.

Let  $G$  be a profinite group containing  $G_E$  as a closed normal subgroup. Let  $\Gamma = G/G_E$ . Suppose that we are given a continuous action of  $\Gamma$  on  $\mathcal{O}_E$  that commutes with Frobenius. There is an induced action of  $G$  on  $\check{\mathcal{O}}_E$  (again because compatible endomorphisms on  $E^{\text{sep}}$  and  $\mathcal{O}_E$  extend uniquely to  $\check{\mathcal{O}}_E$ ).

**Definition 9.10.** A  $(\varphi, \Gamma)$ -module over  $\mathcal{O}_E$  is a  $\varphi$ -module over  $\mathcal{O}_E$  equipped with a semilinear  $\Gamma$ -action commuting with the  $\varphi$ -action. We say that a  $(\varphi, \Gamma)$ -module is étale if it is étale as a  $\varphi$ -module.

Write  $(\varphi, \Gamma)\text{-Mod}_{\mathcal{O}_E}^{\text{ét}}$  for the category of étale  $(\varphi, \Gamma)$ -modules over  $\mathcal{O}_E$ , and write  $\text{Rep}_{\mathbb{Z}_p} G$  for the category of finitely generated  $\mathbb{Z}_p$ -modules with  $G$ -action.

**Theorem 9.11.** *The functor  $D_{\mathcal{E}}: \text{Rep}_{\mathbb{Z}_p} G \rightarrow (\varphi, \Gamma)\text{-Mod}_{\mathcal{O}_{\mathcal{E}}}^{\text{ét}}$  defined by*

$$V \mapsto (V \otimes_{\mathbb{Z}_p} \check{\mathcal{O}}_{\mathcal{E}})^{G_E}$$

*and the functor  $V_{\mathcal{E}}: (\varphi, \Gamma)\text{-Mod}_{\mathcal{O}_{\mathcal{E}}}^{\text{ét}} \rightarrow \text{Rep}_{\mathbb{Z}_p} G$  defined by*

$$M \mapsto (M \otimes_{\mathcal{O}_{\mathcal{E}}} \check{\mathcal{O}}_{\mathcal{E}})^{\varphi=1}$$

*determine an equivalence of categories between  $\text{Rep}_{\mathbb{Z}_p} G$  and  $(\varphi, \Gamma)\text{-Mod}_{\mathcal{O}_{\mathcal{E}}}^{\text{ét}}$ .*

A similar result holds for  $\mathbb{F}_p$ - and  $\mathbb{Q}_p$ -representations.

*Example 9.12.* Let  $G = G_{\mathbb{Q}_p}$ ,  $E = \mathbb{F}_p((T))$ , and embed  $G$  in  $\text{Aut } E^{\text{sep}}$  using Theorem 8.1. We have  $\Gamma = \text{Gal}(\mathbb{Q}_p(\mu_{p^\infty})/\mathbb{Q}_p) \cong \mathbb{Z}_p^\times$ . The action of  $\Gamma$  on  $\mathbb{F}_p((T))$  is given by

$$\gamma \cdot T = (1 + T)^\gamma - 1.$$

The action of  $\Gamma$  on  $\mathcal{O}_{\mathcal{E}}$  can then also be taken to be  $\gamma \cdot T = (1 + T)^\gamma - 1$ .

We have claimed that  $\mathbb{Q}_p(\mu_{p^\infty})$  and  $\mathbb{F}_p((T))$  have isomorphic Galois groups. To prove the isomorphism, we will make use of the concept of perfectoid fields and the tilting correspondence.

Recall that a *nonarchimedean field*  $K$  is a field that is complete with respect to a nontrivial nonarchimedean absolute value  $|\cdot|$ , and that we write

$$\mathcal{O}_K := \{x \in K \mid |x| \leq 1\}$$

$$\mathfrak{m}_K := \{x \in K \mid |x| < 1\}.$$

**Definition 9.13.** A nonarchimedean field  $K$  of residue characteristic  $p$  is *perfectoid* if its value group is nondiscrete and the Frobenius map

$$\Phi: \mathcal{O}_K/p \rightarrow \mathcal{O}_K/p$$

is surjective.

*Remark 9.14.* Like most references but unlike [Ked15], we do not require that  $K$  have characteristic zero.

*Example 9.15.*

- The field  $\mathbb{C}_p$  is perfectoid. More generally, any complete algebraically closed nonarchimedean field of residue characteristic  $p$  is perfectoid.
- A nonarchimedean field of characteristic  $p$  is perfectoid if and only if it is perfect.

**Lemma 9.16.** *The field  $\mathbb{Q}_p^{\text{cyc}} := \widehat{\mathbb{Q}_p(\mu_{p^\infty})}$  is perfectoid.*

*Proof.* Let  $\{\zeta_{p^n}\}_{n \geq 0}$  denote a system of  $p$ -power roots of unity. Note that

$$\mathcal{O}_{\mathbb{Q}_p^{\text{cyc}}}/p \cong \varinjlim_n \mathbb{Z}_p[\zeta_{p^n}]/p.$$

Since  $\zeta_{p^n} = (\zeta_{p^{n+1}})^p$  for each  $n$ , the Frobenius map is surjective.  $\square$

**Definition 9.17.** Let  $K$  be a perfectoid field. The *tilt* of  $K$ , denoted  $K^\flat$ , is defined by

$$K^\flat := \varprojlim_{z \mapsto z^p} K.$$

Define addition on  $K^\flat$  by  $(a_n) + (b_n) = (c_n)$ , where

$$c_n = \lim_{m \rightarrow \infty} (a_{m+n} + b_{m+n})^{p^m}$$

and define multiplication on  $K^\flat$  by componentwise multiplication.

Define a homomorphism of multiplicative monoids  $\sharp: K^\flat \rightarrow K$  by  $(a_n)^\sharp = a_0$ .

**Lemma 9.18.**

- (1) *The limit in Definition 9.17 exists.*
- (2)  *$K^\flat$  is a field of characteristic  $p$ .*
- (3) *The function  $(a_n) \mapsto |(a_n)^\sharp| = |a_0|$  is a nonarchimedean norm on  $K^\flat$ , and  $K^\flat$  is a perfectoid field.*
- (4) *We have*

$$\mathcal{O}_{K^\flat} = \varprojlim_{z \mapsto z^p} \mathcal{O}_K,$$

and there is an isomorphism of rings

$$\mathcal{O}_{K^\flat} \cong \varprojlim_{\Phi} \mathcal{O}_K/p.$$

- (5)  $|K^\times| = |K^{\flat \times}|$ .

*Proof.* Left as an exercise to the reader. Parts (1) and (4) use the following lemma.  $\square$

**Lemma 9.19.** *Let  $R$  be a ring, let  $x, y \in R$ , and let  $n$  be a positive integer. If  $x \equiv y \pmod{p^n}$ , then  $x^p \equiv y^p \pmod{p^{n+1}}$ .*

## 10. TILTING AND UNTILTING

*Example 10.1.* We claim that  $(\mathbb{Q}_p^{\text{cyc}})^\flat$  is isomorphic to the completion of the perfection of  $\mathbb{F}_p((t))$ . We have

$$\mathcal{O}_{\mathbb{Q}_p^{\text{cyc}}}/p \cong \varinjlim_n \mathbb{Z}_p[\zeta_{p^n}]/p \cong \varinjlim_n \mathbb{Z}_p[\zeta_{p^n} - 1]/p^n.$$

The minimal polynomial of  $\zeta_{p^n} - 1$  is

$$\frac{(1+T)^{p^n} - 1}{(1+T)^{p^{n-1}} - 1} \cong T^{p^n - p^{n-1}} \pmod{p}.$$

Since  $\zeta_{p^n} - 1 \cong (\zeta_{p^{n+1}} - 1)^p \pmod{p}$  for each  $n$ , we can write

$$\mathcal{O}_{\mathbb{Q}_p^{\text{cyc}}}/p \cong \varinjlim_n \mathbb{F}_p[t^{p^{-n}}]/(t^{1-1/p}).$$

Then

$$\begin{aligned} (\mathcal{O}_{\mathbb{Q}_p^{\text{cyc}}})^\flat &= \varprojlim_{\Phi} \varinjlim_n \mathbb{F}_p[t^{p^{-n}}]/(t^{1-1/p}) = \varprojlim_m \varinjlim_n \mathbb{F}_p[t^{p^{-n}}]/(t^{p^m - p^{m-1}}) \\ &= (\mathbb{Q}_p^{\text{cyc}})^\flat = (\mathcal{O}_{\mathbb{Q}_p^{\text{cyc}}})^\flat[1/t] \end{aligned}$$

**Definition 10.2.** Let  $K$  be a perfectoid field of characteristic  $p$ . An *untilt* of  $K$  is a perfectoid field  $K^\sharp$ , along with an isomorphism  $(K^\sharp)^\flat \xrightarrow{\sim} K$ .

We would like to classify the untilts of a given characteristic  $p$  perfectoid field  $K$ . In order to do that, we will need to introduce the ring  $W(\mathcal{O}_K)$ .

**Definition 10.3.** An  $\mathbb{F}_p$ -algebra  $R$  is *perfect* if the Frobenius endomorphism of  $R$  is an isomorphism.

**Definition 10.4.** An abelian group  $A$  is  *$p$ -adically complete and separated* if  $A \rightarrow \varprojlim_n A/p^n A$  is an isomorphism.

**Definition 10.5.** A *strict  $p$ -ring* is a ring  $A$  that is  $p$ -adically complete and separated, such that  $A/pA$  is a perfect  $\mathbb{F}_p$ -algebra, and  $p$  is not a zero divisor in  $A$ .

*Example 10.6.* If  $K$  is the completion of an unramified extension of  $\mathbb{Q}_p$ , then  $\mathcal{O}_K$  is a strict  $p$ -ring. The  $p$ -adic completion of  $\mathbb{Z}[x^{p^{-\infty}}]$  is also a strict  $p$ -ring.

The main goal of this section is to prove the following theorem.

**Theorem 10.7.** *The functor  $A \mapsto A/pA$  from strict  $p$ -rings to perfect  $\mathbb{F}_p$ -algebras is an equivalence of categories.*

We will write  $W$  for the functor from perfect  $\mathbb{F}_p$ -algebras to strict  $p$ -rings determined by the above equivalence. For  $R$  a perfect  $\mathbb{F}_p$ -algebra, the ring  $W(R)$  is called the ring of  $p$ -typical Witt vectors of  $R$ .

**Lemma 10.8.** *Let  $A$  be a strict  $p$ -ring.*

- (1) *There is a unique section  $[\cdot]$  of the reduction map  $A \rightarrow A/pA$  that is a homomorphism of multiplicative monoids.*
- (2) *Every element of  $A$  can be written uniquely in the form*

$$\sum_{n=0}^{\infty} p^n [a_n], \quad a_n \in A/pA.$$

*Proof.* For the first part, observe that for any  $x \in R/pR$ , we must have

$$[x] = \lim_{n \rightarrow \infty} (x_n)^{p^n}$$

where  $x_n$  is a lift of  $x^{p^{-n}}$ . The argument that the limit exists and is independent of the choice of lifts follows from Lemma 9.19.

Since  $A$  is  $p$ -adically complete and separated, the second part follows immediately from the first.  $\square$

**Lemma 10.9.** *Let  $A$  be a strict  $p$ -ring, and let  $a, b \in A$ . Suppose that  $a = \sum_{n=0}^{\infty} [a_n]p^n$ ,  $b = \sum_{n=0}^{\infty} [b_n]p^n$ ,  $a + b = \sum_{n=0}^{\infty} [s_n]p^n$ ,  $ab = \sum_{n=0}^{\infty} [t_n]p^n$ . Then  $s_n$  and  $t_n$  are polynomials in the  $a_i^{p^{i-n}}$ ,  $b_i^{p^{i-n}}$  for  $0 \leq i \leq n$ . Furthermore,  $s_n$  is homogeneous of degree 1 (where each  $a_i$  and  $b_i$  has degree 1), and  $t_n$  is homogeneous in the  $a_i$  and  $b_i$  separately, each of degree 1.*

*Proof.* Repeatedly use the identity

$$[x + y] \equiv ([x^{p^{-n}}] + [y^{p^{-n}}])^{p^n} \pmod{p^{n+1}},$$

which follows from Lemma 9.19.  $\square$

**Proposition 10.10** ([Ked15, Lemma 1.1.6]). *Let  $A$  be a strict  $p$ -ring, and let  $B$  be a  $p$ -adically complete ring. Let  $\sharp: A/pA \rightarrow B$  be a multiplicative map that induces a homomorphism of rings  $A/pA \rightarrow B/pB$ . Then the formula*

$$\Theta \left( \sum_{n=0}^{\infty} p^n [x_n] \right) = \sum_{n=0}^{\infty} p^n x_n^{\sharp}$$

*defines a  $p$ -adically continuous homomorphism  $\Theta: A \rightarrow B$  such that  $\Theta \circ [\cdot] = \sharp$ .*

We are especially interested in applying this result in the case where  $A = W(\mathcal{O}_{K^\flat})$  and  $B = \mathcal{O}_K$  for some perfectoid field  $K$ .

*Proof of Theorem 10.7.* Full faithfulness follows from Proposition 10.10.

To prove essential surjectivity, let  $R$  be a perfect ring of characteristic  $p$ , and write  $R = \mathbb{F}_p[X^{-p^\infty}]/\bar{I}$  for some set  $X$  and ideal  $\bar{I} \subset \mathbb{F}_p[X^{-p^\infty}]$ . Let  $A_0$  be the  $p$ -adic completion of  $\mathbb{Z}_p[X^{-p^\infty}]$ ; then one can check that  $A_0$  is a strict  $p$ -ring and  $A_0/pA_0 = \mathbb{F}_p[X^{-p^\infty}]$ . Let  $I \subset A_0$  be the set of elements of the form  $\sum_{n=0}^\infty p^n[x_n]$  with  $x_n \in \bar{I}$ . Then one can check that  $I$  is an ideal of  $A_0$  and  $A := A_0/I$  is a strict  $p$ -ring with  $R = A/pA$ .  $\square$

Let  $K$  be a perfectoid field.

**Definition 10.11.** An ideal  $I$  of  $W(\mathcal{O}_{K^\flat})$  is *primitive of degree 1* if it is generated by an element of the form  $p + [\pi]\alpha$  for some  $\pi \in \mathfrak{m}_{K^\flat}$ ,  $\alpha \in W(\mathcal{O}_{K^\flat})$ .

**Proposition 10.12.** *The map*

$$\Theta: W(\mathcal{O}_{K^\flat}) \rightarrow \mathcal{O}_K$$

*defined in Proposition 10.10 has the following properties:*

- (1)  $\Theta$  is surjective.
- (2)  $\ker \Theta$  is primitive of degree 1.

*Proof.* By Lemma 9.18(4), the map  $\sharp$  is surjective mod  $p$ . So by successive approximation, every element of  $\mathcal{O}_K$  can be written as  $\sum_{n=0}^\infty a_n^\sharp p^n$  for some  $a_n \in \mathcal{O}_{K^\flat}$ . Therefore,  $\Theta$  is surjective.

If  $K$  has characteristic  $p$ , then  $\ker \Theta = (p)$  is primitive of degree 1. Now suppose  $K$  has characteristic 0. Choose  $\pi^\flat \in \mathcal{O}_{K^\flat}$  so that  $\pi := (\pi^\flat)^\sharp$  satisfies  $|\pi| = |p|$ . Choose  $x \in W(\mathcal{O}_{K^\flat})$  satisfying  $\Theta(x) = -p/\pi$ . Since  $\Theta(x)$  is a unit of  $K$ , the constant term in the Teichmüller expansion of  $x$  must be a unit; then  $x$  is also a unit. Let  $\xi = p + [\pi^\flat]x$ ; then  $\xi \in \ker \Theta$ . We claim that in fact  $\xi$  generates  $\ker \Theta$ . Observe that  $\ker \Theta \subseteq ([\pi^\flat], p) = (\xi, p)$ . So any element of  $\ker \Theta$  can be written as  $a\xi + bp$  with  $\Theta(pb) = p\Theta(b) = 0$ . Since  $p$  is not a zero divisor in  $\mathcal{O}_K$ , we get  $\Theta(b) = 0$ . By successive  $p$ -adic approximation, we see that  $\ker \Theta = (\xi)$ .  $\square$

*Remark 10.13.* If you find it dissatisfying that we used a separate argument for  $p = 0$ , see [BMS18, Lemma 3.2ii, Lemma 3.10] for a version of the argument that generalizes better. Essentially, the idea is to use Lemma 10.9 to prove that  $W(\mathcal{O}_{K^\flat})$  is complete for the  $[\pi^\flat]$ -adic topology; then we can use  $[\pi^\flat]$ -adic approximation and we can assume  $|p| \leq |\pi| < 1$  instead of  $|\pi| = |p|$ .

**Proposition 10.14** ([Ked15, Theorem 1.4.13]). *The category of perfectoid fields is equivalent to the category of pairs  $(K, I)$ , where  $K$  is a perfectoid field of characteristic  $p$  and  $I \subset W(\mathcal{O}_K)$  is an ideal that is primitive of degree 1.*

## 11. TILTING AND FIELD EXTENSIONS, GROUP COHOMOLOGY

**Corollary 11.1.** *Let  $K$  be a perfectoid field. Then tilting induces an equivalence of categories between perfectoid extensions of  $K$  and perfectoid extensions of  $K^\flat$ .*

*Moreover, if  $L/K$  is an extension of perfectoid fields, then  $L/K$  is finite iff  $L^\flat/K^\flat$  is finite.*

**Lemma 11.2.** *If  $K$  is a perfectoid field and  $K^\flat$  is algebraically closed, then so is  $K$ .*



*Proof.* Let  $P(X) = X^d + a_{d-1}X^{d-1} + \cdots + a_0 \in K[X]$  be a monic irreducible polynomial. Since  $K^\flat$  is algebraically closed,  $|K^{\flat \times}|$  is a  $\mathbb{Q}$ -vector space, so  $|K^\times|$  is as well. Therefore, by scaling the variable, we may assume that  $a_0 \in \mathcal{O}_K^\times$ . Since  $P$  is irreducible, its Newton polygon must be a straight line, so  $a_i \in \mathcal{O}_K$  for all  $i$ .

Let  $Q(X) \in \mathcal{O}_K^\flat[X]$  be a monic polynomial such that  $P$  and  $Q$  have the same image in  $(\mathcal{O}_K/p\mathcal{O}_K)[X]$ . Let  $y \in \mathcal{O}_{K^\flat}$  be a root of  $Q(X)$ . Then  $p \mid P(y^\sharp)$ . If  $P(y^\sharp) \neq 0$ , choose  $c \in \mathcal{O}_K$  so that  $|c|^d = |P(y^\sharp)|$ . Then replace  $P(X)$  with  $c^{-d}P(cX + y^\sharp)$ . By repeating this process, we find a sequence of elements of  $\mathcal{O}_K$  converging to a root of  $P$ .  $\square$

**Lemma 11.3** (Krasner's lemma). *Let  $K$  be a nonarchimedean field, and let  $\alpha, \beta \in \overline{K}$ , with  $\alpha$  separable over  $K(\beta)$ . Suppose that for all  $\sigma \in \text{Gal}(K(\beta)^{\text{sep}}/K(\beta))$ , either  $\sigma(\alpha) = \alpha$  or  $|\alpha - \sigma(\alpha)| > |\alpha - \beta|$ . Then  $\alpha \in K(\beta)$ .*

*Proof.* Let  $\sigma \in \text{Gal}(K(\beta)^{\text{sep}}/K(\beta))$ . Then

$$|\alpha - \sigma(\alpha)| \leq \max(|\alpha - \beta|, |\sigma(\alpha) - \beta|) = |\alpha - \beta|.$$

By assumption, we must have  $\sigma(\alpha) = \alpha$ . Since this holds for all  $\sigma$ , we must have  $\alpha \in K(\beta)$ .  $\square$

**Proposition 11.4.** *Any finite extension of a perfectoid field is perfectoid.*

*Proof.* Let  $K$  be a perfectoid field, and let  $C^\flat$  be the completion of an algebraic closure of  $K^\flat$ . By Corollary 11.1,  $C^\flat$  has an untilt  $C$  over  $K$ . Furthermore,  $C$  is algebraically closed by Lemma 11.2. Let  $C_0$  be the union of the untilts of all finite extensions of  $K^\flat$ ; then  $C_0$  is dense in  $C$  since the union of all finite extensions of  $C^\flat$  is dense in  $C^\flat$ . It follows from Krasner's lemma that a dense subfield of an algebraically closed nonarchimedean field is separably closed. Then  $C_0$  must contain all finite extensions of  $K$ . So any finite extension  $L/K$  is contained in a Galois extension  $M/K$  that is an untilt of some  $M^\flat/K^\flat$ . By Galois theory, any subfield of  $M$  containing  $K$  must be the untilt of a subfield of  $M^\flat$  containing  $K^\flat$ .  $\square$

**Theorem 11.5.** *Let  $K$  be a perfectoid field. There is an equivalence of categories between finite extensions of  $K$  and finite extensions of  $K^\flat$ .*

*Hence there is an injection*

$$\text{Aut}_{\text{cts}}(\overline{K}) \hookrightarrow \text{Aut}_{\text{cts}}(\overline{K^\flat})$$

*inducing an isomorphism*

$$\text{Gal}(\overline{K}/K) \cong \text{Gal}(\overline{K^\flat}/K^\flat).$$

*Proof.* Combine Corollary 11.1 and Proposition 11.4.  $\square$

**Corollary 11.6.** *The fields  $\mathbb{Q}_p(\mu_{p^\infty})$ ,  $\mathbb{Q}_p^{\text{cyc}} = \widehat{\mathbb{Q}_p(\mu_{p^\infty})}$ ,  $\mathbb{F}_p((t^{p^{-\infty}}))$ , and  $\mathbb{F}_p((t))$  have isomorphic Galois groups.*

*Proof.* By the above theorem,  $\mathbb{Q}_p^{\text{cyc}} = \widehat{\mathbb{Q}_p(\mu_{p^\infty})}$  and  $\mathbb{F}_p((t^{p^{-\infty}}))$  have isomorphic Galois groups. By Krasner's lemma, taking completions does not change the Galois group, and taking perfection also does not change the Galois group.  $\square$

Let  $G$  be a group. A  $G$ -module is an abelian group  $A$  along with a homomorphism  $G \rightarrow \text{Aut } A$ .

We write  $A^G$  for the  $G$ -invariants of  $A$ . Then  $(-)^G$  is a functor from the category of  $G$ -modules to the category of abelian groups. Observe that  $A^G = \text{Hom}_G(\mathbb{Z}, A)$ ,

where  $\mathbb{Z}$  has the trivial  $G$ -action. It is left exact, meaning that given an exact sequence

$$0 \rightarrow A \rightarrow A' \rightarrow A'',$$

the sequence

$$0 \rightarrow A^G \rightarrow (A')^G \rightarrow (A'')^G$$

is also exact. However, it is not right exact.

The forgetful functor from  $G$ -modules to abelian groups has a left adjoint  $G \mapsto \mathbb{Z}[G]$ . We have

$$\mathbb{Z}[G] = \bigoplus_{g \in G} \mathbb{Z} \cdot [g],$$

with

$$[g][h] = [gh].$$

A  $G$ -module  $A$  is called *projective* if the functor  $\text{Hom}_G(A, -)$  is exact. A (possibly infinite) direct sum of copies of  $\mathbb{Z}[G]$  is projective.

Given a projective resolution

$$\cdots \rightarrow P_n \rightarrow P_{n-1} \rightarrow \cdots \rightarrow P_0 \rightarrow \mathbb{Z} \rightarrow 0,$$

we define a cochain complex  $K^\bullet$  by  $K^n = \text{Hom}_G(P_n, A)$ , with differentials induced by the maps  $P_n \rightarrow P_{n-1}$ . Define

$$H^n(G, A) = H^n(K^\bullet).$$

A standard result in homological algebra is that this definition is independent of the resolution chosen. Then  $A^G = H^0(G, A)$ . Given any short exact sequence of  $G$ -modules

$$0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0,$$

there is a long exact sequence

$$0 \rightarrow H^0(G, A) \rightarrow H^0(G, B) \rightarrow H^0(G, C) \rightarrow H^1(G, A) \rightarrow \cdots.$$

A useful choice of projective resolution of  $\mathbb{Z}$  is the following. Let  $P_n = \mathbb{Z}[G^{n+1}]$ . We consider  $P_n$  as a  $G$ -module via

$$g[g_0, \dots, g_n] = [gg_0, \dots, gg_n].$$

The differentials are given by

$$d[g_0, \dots, g_n] = \sum_{j=0}^n (-1)^j [g_0, \dots, \hat{g}_j, \dots, g_n],$$

where  $\hat{g}_j$  indicates that  $g_j$  is omitted. (Most texts use the “bar resolution”, which is equivalent but written slightly differently. I think the description given here is a bit more elegant, but maybe the bar resolution is more convenient for calculations?)

For example,  $H^1(G, A)$  is the space of  $G$ -equivariant maps  $\phi: G \times G \rightarrow A$  satisfying

$$\phi(y, z) - \phi(x, z) + \phi(x, y) = 0$$

for all  $x, y, z \in G$ , modulo the space of maps of the form

$$\phi(x, y) = \psi(y) - \psi(x)$$

for some  $G$ -equivariant  $\psi: G \rightarrow A$ .

Since  $\psi$  is homogeneous, we can write

$$\phi(x, y) = x\tilde{\phi}(x^{-1}y),$$

where

$$\tilde{\phi}(y) = \phi(1, y).$$

Similarly

$$\psi(x) = x\tilde{\psi},$$

where

$$\tilde{\psi} = \psi(1).$$

So we can also think of  $H^1(G, A)$  as the space of maps  $\tilde{\phi}: G \rightarrow A$  satisfyig

$$y\tilde{\phi}(y^{-1}z) - \tilde{\phi}(z) + \tilde{\phi}(y) = 0$$

modulo the space of maps of the form

$$\tilde{\phi}(y) = y\tilde{\psi} - \tilde{\psi}.$$

By making the substitution  $z = yw$ , we can rewrite the constraint as

$$\tilde{\phi}(y) - \tilde{\phi}(yw) + y\tilde{\phi}(w) = 0.$$

If  $A$  has the trivial  $G$ -action, then the constraint simplifies to

$$\tilde{\phi}(y) - \tilde{\phi}(yw) + y\tilde{\phi}(w) = 0,$$

and  $\tilde{\psi} - y\tilde{\psi}$  is always zero. So in that case,

$$H^1(G, A) = \text{Hom}(G, A).$$

## 12. GROUP COHOMOLOGY

In the case  $G = \mathbb{Z}$ , there is also a projective resolution

$$0 \rightarrow \mathbb{Z}[\mathbb{Z}] \xrightarrow{[1]-[0]} \mathbb{Z}[\mathbb{Z}] \rightarrow \mathbb{Z} \rightarrow 0.$$

So for any  $\mathbb{Z}$ -module  $A$ ,

$$H^1(\mathbb{Z}, A) = A / ([1]a - a | a \in A).$$

$$H^i(\mathbb{Z}, A) = 0 \text{ for all } i > 1.$$

The above resolution is related to the fact that  $S^1 = \mathbb{R}/\mathbb{Z}$ . In general, if there is a free action of  $G$  on  $\mathbb{R}^n$  and  $\mathbb{R}^n/G$  admits a triangulation, then every  $G$ -module has a projective resolution of length  $n$ , and  $H^i(G, A) = 0$  for  $i > n$ .

The groups  $H^1(G, A)$  and  $H^2(G, A)$  admit concrete descriptions. The *semidirect product*  $A \rtimes G$  is the group with underlying set  $A \times G$ , and group operation given by

$$(a_1, g_1)(a_2, g_2) = (a_1 + g_1 a_2, g_1 g_2).$$

Consider the set of sections of the projection map  $A \rtimes G \rightarrow G$ . A section is a group homomorphism  $G \rightarrow A \rtimes G$  of the form

$$g \mapsto (s(g), g)$$

for some  $s: G \rightarrow A$ . Such a map is a group homomorphism if and only if

$$s(g_1 g_2) = s(g_1) + g_1 s(g_2)$$

for all  $g_1, g_2 \in G$ . We have

$$(a, 1)(s(g), g)(-a, 1) = (s(g) + a - ga, g),$$

Therefore, two sections are conjugate iff their difference is of the form  $g \mapsto a - ga$ . Therefore,  $H^1(G, A)$  classifies sections of the projection  $A \rtimes G \rightarrow G$  upto  $A$ -conjugacy.

An *extension of  $G$  by  $A$*  is a group  $E$  equipped with a short exact sequence  $1 \rightarrow A \rightarrow E \rightarrow G \rightarrow 1$ . There is a bijection between classes in  $H^2(G, A)$  and extensions of  $G$  by  $A$ . We omit the details.

The groups  $H^n(G, A)$  are functorial in both  $G$  and  $A$ . The functoriality in  $G$  can be described as follows. If  $f: H \rightarrow G$  is a homomorphism of groups and  $A$  is a  $G$ -module, then we write  $f^*A$  for the  $H$ -module with underlying abelian group  $A$  and  $H$ -action determined by  $f$ . The machinery of derived functors tells us that there are homomorphisms

$$H^n(G, A) \rightarrow H^n(G', f^*A).$$

If the cohomology is calculated using the resolution  $\cdots \rightarrow \mathbb{Z}[G \times G] \rightarrow \mathbb{Z}[G] \rightarrow \mathbb{Z}$  mentioned above, then this map sends a homogeneous cocycle  $\phi: G^{n+1} \rightarrow A$  to  $(f \times \cdots \times f) \circ \phi$ .

If  $f: H \rightarrow G$  is an injection, then the functor  $f^*$  is called *restriction* and is denoted by  $\text{Res}$ . We sometimes also denote the corresponding  $H^n(G, A) \rightarrow H^n(H, A)$  by  $\text{Res}$ .

The restriction functor  $\text{Res}$  has a right adjoint called  $\text{Ind}$ . Given an  $H$ -module  $B$ , the  $G$ -module  $\text{Ind } B$  is given by

$$\text{Ind } B = \text{Hom}_{\mathbb{Z}[H]}(\mathbb{Z}[G], B),$$

where we view  $\mathbb{Z}[G]$  as an  $H$ -module via multiplication on the left, and as a  $G$ -module via multiplication on the right. Recall that “right adjoint” means that for any  $G$ -module  $A$  and  $H$ -module  $B$ , there is a natural isomorphism

$$\text{Hom}_{\mathbb{Z}[H]}(\text{Res } A, B) \cong \text{Hom}_{\mathbb{Z}[G]}(A, \text{Ind } B).$$

We also have

$$H^n(H, B) = H^n(G, \text{Ind } B).$$

Modules in the image of the functor  $\text{Ind}$  are called “coinduced modules”. (There is also a functor called  $\text{ind}$  that is left adjoint to  $\text{Res}$ , and modules in the image of  $\text{ind}$  are called “induced modules”. But we will not use  $\text{ind}$ .)

Given a normal subgroup  $H$  of  $G$ , the composite

$$H^n(G/H, A^H) \rightarrow H^n(G, A^H) \rightarrow H^n(G, A)$$

is called an *inflation map* and denoted  $\text{Inf}$ .

We would like to allow our groups and modules to have a topology. The right way to do this is probably to use condensed mathematics, but as far as I know, no one has worked out the details yet. So instead, we make the following ad hoc definition. Let  $G$  be a topological group, and let  $A$  be a topological  $G$ -module (i.e. a  $G$ -module such that the  $G$ -action is continuous). Define a cochain complex  $K_{\text{cts}}$  by letting  $K_{\text{cts}}^n$  be the set of continuous  $G$ -equivariant homomorphisms  $G^{n+1} \rightarrow A$ . Define  $H_{\text{cts}}^n(G, A)$  to be the  $n$ th cohomology group of  $K_{\text{cts}}$ .

Warning: a short exact sequence of modules does not always give us a long exact sequence of cohomology groups. For example, we have a short exact sequence of abelian groups

$$0 \rightarrow \mathbb{Z} \rightarrow \mathbb{R} \rightarrow S^1 \rightarrow 0.$$

If we consider these as  $S^1$ -modules with trivial  $S^1$ -action, then

$$\begin{aligned} H_{\text{cts}}^1(S^1, \mathbb{R}) &= \text{Hom}_{\text{cts}}(S^1, \mathbb{R}) = 0 \\ H_{\text{cts}}^1(S^1, S^1) &= \text{Hom}_{\text{cts}}(S^1, S^1) = \mathbb{Z} \end{aligned}$$

$$H_{\text{cts}}^2(S^1, \mathbb{Z}) = 0.$$

This is why it would probably be better to use condensed mathematics.

However, given a short exact sequence  $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$ , we do always get a long exact sequence

$$0 \rightarrow A^G \rightarrow B^G \rightarrow C^G \rightarrow H_{\text{cts}}^1(G, A) \rightarrow H_{\text{cts}}^1(G, B) \rightarrow H_{\text{cts}}^1(G, C).$$

We get the full long exact sequence if the map  $B \rightarrow C$  admits a continuous section (as a map of topological spaces; the section need not be a  $G$ -module homomorphism).

**Lemma 12.1.** *Let  $G$  be a profinite group, and let  $A$  be a topological  $G$ -module.*

- (1) *If  $A$  has the discrete topology, then every vector in  $A$  has an open stabilizer, and*

$$H_{\text{cts}}^i(G, A) = \varprojlim_{H \subset G} H^i(G/H, A^H),$$

*where the limit runs over open normal subgroups of  $G$ .*

- (2) *If  $H_{\text{cts}}^i(G, \mathbb{F}_p)$  is finite for all  $i$ , and  $A$  is a finite free  $\mathbb{Z}_p$ -module, then*

$$H_{\text{cts}}^i(G, A) = \varprojlim_n H_{\text{cts}}^i(G, A/p^n A)$$

- (3) *If  $H_{\text{cts}}^i(G, \mathbb{F}_p)$  is finite for all  $i$ , and  $A$  is a finite-dimensional  $\mathbb{Q}_p$ -vector space, then for any  $G$ -stable lattice  $\Lambda \subset A$ ,*

$$H_{\text{cts}}^i(G, A) = H_{\text{cts}}^i(G, \Lambda) \otimes_{\mathbb{Z}_p} \mathbb{Q}_p$$

*(Since  $G$  is compact, such a lattice always exists.)*

**Lemma 12.2.**

- (1) *Let  $K$  be a finite extension of  $\mathbb{Q}_p$ . Then  $H_{\text{cts}}^i(\text{Gal}(\overline{K}/K), \mathbb{F}_p)$  is finite for all  $i$ .*
- (2) *Let  $K$  be a finite extension of  $\mathbb{Q}$ , and let  $S$  be a finite set of places of  $K$ . Let  $K_S$  be the largest algebraic extension of  $K$  unramified outside  $S$ . Then  $H_{\text{cts}}^i(\text{Gal}(K_S/K), \mathbb{F}_p)$  is finite for all  $i$ .*

### 13. AX-SEN-TATE THEOREM

Let  $K$  be a  $p$ -adic field, and let  $C := \widehat{\overline{K}}$  be the completion of its algebraic closure with respect to the norm topology. In a previous lecture, I claimed that there is no element “ $2\pi i$ ” in  $C$  so that  $G_K$  acts on “ $2\pi i$ ”  $\cdot \mathbb{Q}_p$  by the cyclotomic character  $\chi$ .

**Theorem 13.1.**  $C^{G_K} = K$ .

*Let  $K_\infty := K(\mu_{p^\infty})$ ,  $K^{\text{cyc}} := \widehat{K_\infty}$ ,  $\Gamma := \text{Gal}(K_\infty/K)$ . If  $\chi: \Gamma \rightarrow K^\times$  has infinite order, then  $C(\chi)^{G_K} = 0$ .*

**Remark 13.2.** One can also show that  $H^1(G_K, C)$  is a one-dimensional  $K$ -vector space and that  $H^1(G_K, C(\chi)) = 0$ . In the interest of space, we omit the proof. See [Tat67, §3].

For an elementary (but calculation-heavy) proof that  $C^{G_K} = K$ , see [Ax70] or [FO, Proposition 3.8].

The proof breaks down into the following steps.

**Lemma 13.3.** *Let  $k$  be the residue field of  $K$ , and let  $K_0 = W(k)[1/p]$ . Let  $\pi$  be a uniformizer of  $K$ , and let  $e = \frac{\log |p|}{\log |\pi|}$ . Then  $[K : K_0] = e$ .*

*Proof.* Every element of  $\mathcal{O}_K$  can be written as  $\sum_{i=0}^{\infty} [a_i] \pi^i$  with  $a_i \in k$ . We can write  $\pi^e = [u]p + \sum_{i=e+1}^{\infty} [b_i] \pi^i$  with  $u \in k^\times, b_i \in K$ . Then by repeated substitution, every element of  $\mathcal{O}_K$  can be written in the form  $\sum_{i=0}^{\infty} \sum_{j=0}^{e-1} [c_i] p^i \pi^j$ . Hence  $1, \pi, \dots, \pi^{e-1}$  form a basis for  $\mathcal{O}_K$  as a  $W(k)$ -module.  $\square$

**Lemma 13.4.** *The field  $K^{\text{cyc}}$  is perfectoid.*

*Proof.* Let  $k$  be the residue field of  $K$ . Then  $W(k)[1/p]^{\text{cyc}}$  is perfectoid by the same argument as in Lemma 9.16. Since  $K$  is finite extension of  $W(k)[1/p]$ , the result follows from Lemma 11.4.  $\square$

**Proposition 13.5.** *If  $L$  is a perfectoid field, then  $(\widehat{L})^{G_L} = L$ . In particular,  $C^{\text{Gal}(\overline{K}/K_\infty)} = K^{\text{cyc}}$ .*

**Proposition 13.6.**  $(K^{\text{cyc}})^\Gamma = K$ .

*If  $\chi: \Gamma \rightarrow K^\times$  has infinite order, then  $K^{\text{cyc}}(\chi)^\Gamma = 0$ .*

To prove Proposition 13.5, we will need a few lemmas.

**Lemma 13.7.** *Let  $M/L$  be a finite extension of perfectoid fields. Then  $\text{tr}_{M/L}(\mathfrak{m}_M) = \mathfrak{m}_L$ .*

*Proof.* Since  $M^\flat/L^\flat$  is separable,  $\text{tr}_{M^\flat/L^\flat}(\mathfrak{m}_{M^\flat})$  is a nonzero ideal of  $\mathcal{O}_L$ . By applying the inverse of Frobenius, we see that it must be all of  $\mathfrak{m}_{L^\flat}$ . Since there are compatible surjective ring homomorphisms  $\mathcal{O}_{M^\flat} \twoheadrightarrow \mathcal{O}_M/p\mathcal{O}_M, \mathcal{O}_{L^\flat} \twoheadrightarrow \mathcal{O}_L/p\mathcal{O}_L$ , this implies that  $\text{tr}_{M/L}(\mathfrak{m}_M) = \mathfrak{m}_L$ .  $\square$

**Lemma 13.8.** *Let  $L$  be a perfectoid field, and let  $y \in \overline{L}$ . Let  $c > 1$  be a real number. Then there exists  $z \in L$  so that*

$$|y - z| \leq c \max_{\sigma \in G_L} |\sigma y - y|.$$

*Proof.* Choose a finite extension  $M$  of  $L$  containing  $y$ . We will write  $\text{tr}$  for the trace from  $M$  to  $L$ . By Lemma 13.7, we can find  $x \in M$  with  $|x| < 1, |\text{tr } x| \geq c^{-1}$ . Let  $z = \frac{\text{tr}(xy)}{\text{tr } x}$ . Then

$$y - z = \frac{\sum_{\sigma \in \text{Gal}(M/L)} (\sigma x)(y - \sigma y)}{\text{tr } x}.$$

Hence  $|y - z| \leq c \max_{\sigma \in H_K} |\sigma y - y|$ , as desired.  $\square$

*Proof of Proposition 13.5.* Let  $x \in (\widehat{L})^{G_L}$ . Then for any real  $\epsilon > 0$ , we can find  $y \in \overline{L}$  so that  $|x - y| < \epsilon$ . By the strong triangle inequality, for all  $\sigma \in G_L$ ,

$$|\sigma y - y| \leq \max(|\sigma y - z|, |y - z|) = |y - z| < \epsilon.$$

By Lemma 13.8, for any  $c > 1$ , we can find  $z \in L$  so that  $|y - z| \leq c\epsilon$ . Hence  $|x - z| < c\epsilon$ . Since this is true for any  $\epsilon$ , and  $L$  is complete,  $x \in L$ .  $\square$

In order to prove Proposition 13.6, there is no harm in replacing  $K$  with a finite Galois extension. So we may assume  $K$  contains a  $p$ th root of unity. This implies that  $\Gamma \cong \mathbb{Z}_p$ . Let  $\gamma$  be a topological generator of  $\Gamma$ .

Let  $t: K_\infty \rightarrow K$  be the “normalized trace map” satisfying  $t|_L = \frac{1}{[L:K]} \text{tr}_{L/K}$  for every finite extension  $L/K$  inside  $K_\infty$ .

**Lemma 13.9.** *For any  $x \in K_\infty$ ,*

$$|x - t(x)| \leq |p|^{-1} |x - \gamma x|.$$

*Proof.* For each  $n$ , let  $K_n$  be the fixed field of  $p^n\Gamma$ . We will prove the inequality on each  $K_n$  by induction. The base case  $n = 0$  is trivial. Since  $1 - \gamma$  divides  $p - (1 + \gamma^{p^{n-1}} + \dots + \gamma^{p^{n-1}(p-1)})$ , we have

$$|px - \mathrm{tr}_{K_n/K_{n-1}} x| \leq |(1 - \gamma)x|.$$

or equivalently,

$$|x - p^{-1} \mathrm{tr}_{K_n/K_{n-1}} x| \leq |p^{-1}| |(1 - \gamma)x|.$$

By the induction hypothesis,

$$|p^{-1} \mathrm{tr}_{K_n/K_{n-1}} x - t(x)| \leq |p^{-1}| |(1 - \gamma)(p^{-1} \mathrm{tr}_{K_n/K_{n-1}} x)| \leq |p^{-1}| |(1 - \gamma)x|,$$

where we used the fact that  $1 - \gamma$  commutes with the normalized trace in the last inequality. Then the claim follows from the triangle inequality.  $\square$

**Corollary 13.10.** *For any  $x \in K_\infty$ ,*

$$|t(x)| \leq |p^{-1}| |x|.$$

*So the function  $t$  is continuous. Moreover,  $1 - \gamma$  is invertible on  $\ker t$ , and its inverse is continuous.*

*Hence  $t$  can be extended to a continuous function  $\hat{t}: K^{\mathrm{cyc}} \rightarrow K$ , and the restriction of  $1 - \gamma$  to  $\ker \hat{t}$  has a continuous inverse.*

*Proof of Proposition 13.6.* We have

$$K^{\mathrm{cyc}} = K \oplus \ker \hat{t}$$

$$(K^{\mathrm{cyc}})^\Gamma = K^\Gamma \oplus (\ker \hat{t})^\Gamma = K \oplus 0 = K.$$

Finally, suppose that  $\chi: \Gamma \rightarrow K^\times$  is a character of infinite order. We will show that  $K^{\mathrm{cyc}}(\chi^{-1})^\Gamma = 0$ . For sufficiently large  $n$ , we must have  $|\chi(\gamma^{p^n}) - 1| < |p|$ . Since there is no harm with replacing  $K$  by a finite Galois extension, we may assume  $|\chi(\gamma) - 1| < |p|$ . On  $\ker \hat{t}$ ,

$$\gamma - \chi(\gamma) = (\gamma - 1)(1 - (\chi(\gamma) - 1)(\gamma - 1)^{-1})$$

and  $(1 - (\chi(\gamma) - 1)(\gamma - 1)^{-1})^{-1}$  has a convergent power series, so  $\gamma - \chi(\gamma)$  is invertible. On  $K$ ,  $\gamma - \chi(\gamma) = 1 - \chi(\gamma)$  is invertible since  $\chi$  has infinite order.  $\square$

#### 14. THE RING $B_{\mathrm{dR}}$

Now we define the ring  $B_{\mathrm{dR}}$  mentioned in the first lecture.

Let  $K$  be a  $p$ -adic field, and let  $C$  be the completion of its algebraic closure with respect to the norm topology. We define the ring

$$A_{\mathrm{inf}} := W(\mathcal{O}_{C^\flat}).$$

By Proposition 10.10, there is a homomorphism

$$\Theta: A_{\mathrm{inf}} \rightarrow \mathcal{O}_C.$$

We will consider the localization

$$\Theta_{\mathbb{Q}}: A_{\mathrm{inf}}[1/p] \rightarrow C.$$

**Lemma 14.1.** *For each positive integer  $n$ ,  $(\ker \Theta_{\mathbb{Q}})^n \cap A_{\text{inf}} = (\ker \Theta)^n$ , and  $\bigcap_n (\ker \Theta_{\mathbb{Q}})^n = 0$ .*

*Let*

$$B_{\text{dR}}^+ := \varprojlim_n A_{\text{inf}}[1/p]/(\ker \Theta_{\mathbb{Q}})^n.$$

*Then  $B_{\text{dR}}^+$  is a complete discrete valuation ring with residue field  $C$ .*

*Proof.* Left as an exercise to the reader.  $\square$

Define

$$B_{\text{dR}} := \text{Frac } B_{\text{dR}}^+.$$

Define a decreasing filtration on  $B_{\text{dR}}$  by letting  $\text{Fil}^i B_{\text{dR}}$  be the fractional ideal  $(\ker \Theta_{\mathbb{Q}})^i$ . Now we will define an element  $t \in B_{\text{dR}}^+$  that is the  $p$ -adic analogue of  $2\pi i$ . Let  $\epsilon \in \mathcal{O}_{C^\flat}$  be an element with  $\epsilon_0 = 1$ ,  $\epsilon_1 \neq 1$ . Then  $[\epsilon] - 1 \in \ker \Theta$ , so it makes sense to define

$$t := \log[\epsilon] = \sum_{n=1}^{\infty} (-1)^{n+1} \frac{([\epsilon] - 1)^n}{n}.$$

**Lemma 14.2.** *For any  $a \in \mathbb{Q}_p$ ,  $\log([\epsilon^a]) = a \log[\epsilon]$ . Hence  $G_K$  acts on  $t \cdot \mathbb{Q}_p$  by the cyclotomic character.*

*Proof.* It is formal that for  $a \in \mathbb{Q}$ ,  $\log([\epsilon^a]) = a \log[\epsilon]$ .

We would like to argue that the equality holds for  $a \in \mathbb{Q}_p$  by continuity. But the  $G_K$ -action on  $B_{\text{dR}}^+$  is not jointly continuous for the  $\ker \Theta_{\mathbb{Q}}$ -adic topology, so we need to find a “better” topology.

Let  $\xi$  be a generator of  $\ker \Theta$ . Using Lemma 10.9 and the fact that the  $G_K$ -action on  $\mathcal{O}_{C^\flat}$  is jointly continuous, we can verify that the  $G_K$ -action on  $A_{\text{inf}}$  is jointly continuous for the  $(p, \xi)$ -adic topology on  $A_{\text{inf}}$ . Give  $A_{\text{inf}}/\xi^n$  the quotient topology. Extend this topology to  $A_{\text{inf}}[1/p]/\xi^n$  by letting  $A_{\text{inf}}/\xi^n$  be open. Finally, give  $B_{\text{dR}}^+ = \varprojlim_n A_{\text{inf}}[1/p]/\xi^n$  the inverse limit topology. Then  $G_K$  acts jointly continuously for this topology, and  $\log$  is continuous on the open set  $1 + \mathfrak{m}_{A_{\text{inf}}} + \xi B_{\text{dR}}^+ \subset B_{\text{dR}}^+$ , where  $\mathfrak{m}_{A_{\text{inf}}}$  is the maximal ideal of  $A_{\text{inf}}$ .  $\square$

**Lemma 14.3.**  *$t$  is a uniformizer of  $B_{\text{dR}}^+$ .*

*Proof.* It is clear that  $t \in \text{Fil}^1 B_{\text{dR}}^+$ , so we just need to check that  $t \notin \text{Fil}^2 B_{\text{dR}}^+$ . For this, it is enough to check that  $[\epsilon] - 1 \notin \text{Fil}^2 B_{\text{dR}}^+$ , or equivalently,  $[\epsilon] - 1 \notin (\ker \Theta)^2$ .

For simplicity, we will assume  $p \neq 2$ . See [BC, Proposition 4.4.8] for the case  $p = 2$ .

Recall from the proof of Proposition 10.12 that  $\ker \Theta \subset (p, [\pi^{\flat}])$ , where  $\pi^{\flat} \in \mathcal{O}_{C^\flat}$  satisfies  $|\pi^{\flat}| = |p|$ . So it is enough to check that  $[\epsilon] - 1 \notin (p, [\pi^{\flat}]^2)$ , i.e.  $|\epsilon - 1| > |p|^2$ .

Recall that if  $\zeta_{p^n}$  is a primitive  $n$ th root of unity, then  $|\zeta_{p^n} - 1| = |p|^{1/p^{n-1}(p-1)}$ . Therefore,

$$|\epsilon - 1| = \lim_{n \rightarrow \infty} |\zeta_{p^n} - 1|^{p^n} = |p|^{p/(p-1)} > |p|^2.$$

$\square$

**Proposition 14.4.** *There is a natural Galois-equivariant inclusion  $\overline{K} \hookrightarrow B_{\text{dR}}$ .*



*Proof.* Let  $\bar{k}$  be the residue field of  $\bar{K}$ . There is a natural inclusion  $\bar{k} \hookrightarrow \mathcal{O}_{C^\flat}$  sending  $x \mapsto [x^{p^{-n}}]$ , which induces inclusions  $W(\bar{k}) \hookrightarrow A_{\text{inf}}, W(\bar{k})[1/p] \hookrightarrow B_{\text{dR}}^+$ . Any  $x \in \bar{K}$  satisfies an irreducible monic polynomial over  $W(\bar{k})[1/p]$ . This polynomial splits completely in  $C$ , the residue field of  $B_{\text{dR}}^+$ , so it also splits in  $B_{\text{dR}}^+$  by Hensel's lemma. So there is a unique inclusion  $\bar{K} \hookrightarrow B_{\text{dR}}^+$  that makes the following diagram commute.

$$\begin{array}{ccc} W(\bar{k})[1/p] & \hookrightarrow & B_{\text{dR}}^+ \\ \downarrow & \nearrow & \downarrow \\ \bar{K} & \hookrightarrow & C \end{array}$$

□

**Proposition 14.5.** *The natural inclusions  $K = \bar{K}^{G_K} \hookrightarrow (B_{\text{dR}}^+)^{G_K} \hookrightarrow B_{\text{dR}}^{G_K}$  are isomorphisms.*

*Proof.* For any  $n$ , by Lemma 14.2, we have an exact sequence

$$0 \rightarrow \text{Fil}^{n+1} B_{\text{dR}} \rightarrow \text{Fil}^n B_{\text{dR}} \rightarrow C(n) \rightarrow 0.$$

(Here,  $C(n) = C \otimes_{\mathbb{Z}_p} (\mathbb{Z}_p(1)^{\otimes n})$ .) It induces an exact sequence

$$0 \rightarrow (\text{Fil}^{n+1} B_{\text{dR}})^{G_K} \rightarrow (\text{Fil}^n B_{\text{dR}})^{G_K} \rightarrow C(n)^{G_K}.$$

By Theorem 13.1,  $C^{G_K} = K$  and  $C(n)^{G_K} = 0$  if  $n \neq 0$ . We can then show that for all  $n > 1$ , the inclusion

$$(\text{Fil}^n B_{\text{dR}}^+)^{G_K} \hookrightarrow (\text{Fil}^1 B_{\text{dR}}^+)^{G_K}$$

is an isomorphism. Since  $\bigcap_n \text{Fil}^n B_{\text{dR}}^+ = 0$ ,

$$(\text{Fil}^1 B_{\text{dR}}^+)^{G_K} = 0.$$

Then the map  $(B_{\text{dR}}^+)^{G_K} \rightarrow C^{G_K} = K$  is injective. We already know that  $(B_{\text{dR}}^+)^{G_K}$  contains  $\bar{K}^{G_K} = K$ , so it must equal  $K$ . Finally, we can show that for all  $n \leq 0$ ,

$$(B_{\text{dR}}^+)^{G_K} \hookrightarrow (\text{Fil}^n B_{\text{dR}})^{G_K}$$

is an isomorphism, implying that

$$(B_{\text{dR}}^+)^{G_K} \hookrightarrow (B_{\text{dR}})^{G_K}$$

is an isomorphism. □

**Definition 14.6.** Let  $V$  be a finite-dimensional  $\mathbb{Q}_p$ -representation  $V$  of  $G_K$ . Let  $D_{\text{dR}}(V)$  be the filtered  $K$ -vector space  $(V \otimes_{\mathbb{Q}_p} B_{\text{dR}})^{G_K}$ .

**Proposition 14.7.** *Let  $V$  be a finite-dimensional  $\mathbb{Q}_p$ -representation  $V$  of  $G_K$ . Let*

$$\alpha_V: D_{\text{dR}}(V) \otimes_K B_{\text{dR}} \rightarrow V \otimes_{\mathbb{Q}_p} B_{\text{dR}}$$

*be the restriction of the map*

$$V \otimes_{\mathbb{Q}_p} B_{\text{dR}} \otimes_K B_{\text{dR}} \rightarrow V \otimes_{\mathbb{Q}_p} B_{\text{dR}}$$

*induced by multiplication on  $B_{\text{dR}}$ . Then  $\alpha_V$  is an injection. In particular,*

$$\dim_K D_{\text{dR}}(V) \leq \dim_{\mathbb{Q}_p} V$$

*with equality iff  $\alpha_V$  is an isomorphism.*

*Proof.* See [BC, Theorem 5.2.1(1)]. □

**Definition 14.8.** We say that  $V$  is *de Rham* if  $\dim_K D_{\text{dR}}(V) = \dim_{\mathbb{Q}_p} V$ .

If  $V$  is de Rham, then the *Hodge–Tate weights* of  $V$  are the integers  $i$  such that  $\text{gr}^i D_{\text{dR}}(V) \neq 0$ .

*Example 14.9.* The Hodge–Tate weight of  $\mathbb{Q}_p(n)$  is  $-n$ .

**Theorem 14.10.** *If  $X$  is a proper smooth variety over  $K$ , then  $H_{\text{ét}}^i(X_{\overline{K}}, \mathbb{Q}_p)$  is de Rham, with Hodge–Tate weights between 0 and  $i$ , inclusive.*

This is proved as part of the de Rham comparison theorem.

**Lemma 14.11.** *Let  $L$  be a finite extension of  $K$ . Then a representation of  $G_K$  is de Rham if and only if its restriction to  $G_L$  is de Rham.*

*Proof.* This follows from Galois descent.  $\square$

**Lemma 14.12.** *A tensor product of two de Rham representations is de Rham.*

*Subrepresentations and quotients of a de Rham representation are de Rham.*

*Proof.* Suppose  $V_1$  and  $V_2$  are de Rham representations. Multiplication on  $B_{\text{dR}}$  induces a map

$$D_{\text{dR}}(V_1) \otimes_K D_{\text{dR}}(V_2) \rightarrow D_{\text{dR}}(V_1 \otimes_{\mathbb{Q}_p} V_2).$$

To show that  $V_1 \otimes V_2$  is de Rham, it suffices to show that the above map is injective. Then it also suffices to check injectivity of the map

$$D_{\text{dR}}(V_1) \otimes_K D_{\text{dR}}(V_2) \rightarrow (V_1 \otimes_{\mathbb{Q}_p} V_2) \otimes_{\mathbb{Q}_p} B_{\text{dR}}$$

and likewise of the map

$$(14.13) \quad D_{\text{dR}}(V_1) \otimes_K D_{\text{dR}}(V_2) \otimes_K B_{\text{dR}} \rightarrow (V_1 \otimes_{\mathbb{Q}_p} V_2) \otimes_{\mathbb{Q}_p} B_{\text{dR}},$$

After rewriting the LHS as

$$(D_{\text{dR}}(V_1) \otimes_K B_{\text{dR}}) \otimes_{B_{\text{dR}}} (D_{\text{dR}}(V_2) \otimes_K B_{\text{dR}})$$

and the RHS as

$$(V_1 \otimes_{\mathbb{Q}_p} B_{\text{dR}}) \otimes_{B_{\text{dR}}} (V_2 \otimes_{\mathbb{Q}_p} B_{\text{dR}}),$$

we see that Proposition 14.7 implies that (14.13) is an isomorphism.

We can check injectivity after tensoring with  $B_{\text{dR}}$ . Since  $V_1$  and  $V_2$  are de Rham, we can identify

$$\begin{aligned} D_{\text{dR}}(V_1) \otimes_K D_{\text{dR}}(V_2) \otimes_K B_{\text{dR}} &\cong (D_{\text{dR}}(V_1) \otimes_K B_{\text{dR}}) \otimes_{B_{\text{dR}}} (D_{\text{dR}}(V_2) \otimes_K B_{\text{dR}}) \\ &\cong (V_1 \otimes_{\mathbb{Q}_p} B_{\text{dR}}) \otimes_{B_{\text{dR}}} (V_2 \otimes_{\mathbb{Q}_p} B_{\text{dR}}) \cong (V_1 \otimes V_2) \otimes_{\mathbb{Q}_p} B_{\text{dR}} \end{aligned}$$

Therefore, we are reduced to showing that the induced map

$$(V_1 \otimes_{\mathbb{Q}_p} V_2) \otimes_{\mathbb{Q}_p} B_{\text{dR}} \rightarrow D_{\text{dR}}(V_1 \otimes_{\mathbb{Q}_p} V_2) \otimes_{\mathbb{Q}_p} B_{\text{dR}}$$

is injective.

Suppose we have an exact sequence of representations

$$0 \rightarrow V' \rightarrow V \rightarrow V'' \rightarrow 0$$

with  $V$  de Rham. Then we have a left exact sequence

$$0 \rightarrow D_{\text{dR}}(V') \rightarrow D_{\text{dR}}(V) \rightarrow D_{\text{dR}}(V''),$$

so

$$\dim_K D_{\text{dR}}(V') + \dim_K D_{\text{dR}}(V'') \geq \dim_K D_{\text{dR}} V.$$

On the other hand,

$$\dim_K D_{\text{dR}}(V') \leq \dim_{\mathbb{Q}_p} V'$$

$$\begin{aligned}\dim_K D_{\text{dR}}(V'') &\leq \dim_{\mathbb{Q}_p} V'' \\ \dim_K D_{\text{dR}}(V) &= \dim_{\mathbb{Q}_p} V.\end{aligned}$$

So all inequalities must actually be equalities.  $\square$

*Example 14.14.* Let  $\psi: \mathbb{Z}_p^\times \rightarrow \mathbb{Z}_p^\times$  be a character, and let  $\chi: G_K \rightarrow \mathbb{Z}_p^\times$  be the cyclotomic character. The character  $\psi \circ \chi: G_K \rightarrow \mathbb{Z}_p^\times$  is de Rham if and only if  $\psi$  is a product of a finite order character and a character of the form  $z \mapsto z^n$  for some  $n \in \mathbb{Z}$ . In particular, there exist characters that are not de Rham. The “only if” direction follows from Theorem 13.1 by the same argument as in Proposition 14.5. For the “if” direction, we can use Lemma 14.11 to reduce to the case where the finite order character is trivial, and then apply Lemma 14.2.

### 15. $B_{\text{dR}}$ AND DIFFERENTIALS

It is not immediately obvious from the definition of  $B_{\text{dR}}$  that it should have anything to do with integrals of differential forms. We will now give an alternate characterization of  $B_{\text{dR}}^+$  that suggests a connection to differential forms.

Let  $k$  be the residue field of  $K$ . Let

$$A_{\text{inf},K} := A_{\text{inf}} \otimes_{W(k)} \mathcal{O}_K.$$

There is a map  $\Theta_K: A_{\text{inf},K} \rightarrow \mathcal{O}_C$ , and for each positive integer  $n$ ,  $B_{\text{dR}}^+ / \text{Fil}^n B_{\text{dR}} \cong A_{\text{inf},K} / (\ker \Theta_K)^n [1/p]$ . So

$$B_{\text{dR}}^+ \cong \varprojlim_n (A_{\text{inf},K} / (\ker \Theta_K)^n [1/p]).$$

Inductively define

$$\begin{aligned}\mathcal{O}_{\overline{K}}^{(0)} &:= \mathcal{O}_{\overline{K}} \\ \Omega^{(n)} &:= \mathcal{O}_{\overline{K}} \otimes_{\mathcal{O}_{\overline{K}}^{(n-1)}} \Omega_{\mathcal{O}_{\overline{K}}^{(n-1)}/\mathcal{O}_K} \\ \mathcal{O}_{\overline{K}}^{(n)} &:= \ker \left( d^{(n)}: \mathcal{O}_{\overline{K}}^{(n-1)} \rightarrow \Omega^{(n)} \right)\end{aligned}$$

**Theorem 15.1.**

- (1) For any nonnegative integer  $n$ , the preimage of  $A_{\text{inf},K} / (\ker \Theta_K)^{n+1}$  under the inclusion  $\overline{K} \hookrightarrow B_{\text{dR}}^+ / \text{Fil}^{n+1} B_{\text{dR}}$  is  $\mathcal{O}_{\overline{K}}^{(n)}$ .
- (2) For any nonnegative integers  $m, n$ , the map  $\mathcal{O}_{\overline{K}}^{(n)} / p^m \rightarrow A_{\text{inf},K} / ((\ker \Theta_K)^{n+1}, p^m)$  is an isomorphism.
- (3)  $B_{\text{dR}}^+$  is the completion of  $\overline{K}$  for the topology defined by letting the sets  $p^m \mathcal{O}_{\overline{K}}^{(n)}$  for nonnegative integers  $m, n$  be a basis of open neighborhoods of the identity.
- (4) For any nonnegative integer  $n$ ,  $d^{(n)}$  is surjective.

**Corollary 15.2.** The inclusion  $\overline{K} \hookrightarrow B_{\text{dR}}^+$  cannot be extended to a continuous map  $C \rightarrow B_{\text{dR}}^+$ .

*Proof.* Any  $x \in \text{Fil}^1 B_{\text{dR}}^+$  can be written as a limit of a sequence elements of  $\overline{K}$ . By continuity of the projection  $B_{\text{dR}}^+ \rightarrow C$ , any such sequence converges to 0 in  $C$ . So there cannot be any map  $C \rightarrow B_{\text{dR}}^+$  extending the inclusion  $\overline{K} \hookrightarrow B_{\text{dR}}^+$  that preserves sequential limits.  $\square$

**Lemma 15.3.** The image of  $\mathcal{O}_{\overline{K}}^{(n)}$  in  $B_{\text{dR}}^+$  is contained in  $A_{\text{inf},K} + \text{Fil}^{n+1} B_{\text{dR}}$ .

*Proof.* We will use induction on  $n$ . The case  $n = 0$  follows from the surjectivity of  $\Theta$ .

Let  $x \in \mathcal{O}_{\overline{K}}^{(n-1)}$ . By the induction hypothesis, the image of  $x$  in  $B_{\text{dR}}^+$  can be expressed as  $x_0 + \epsilon$ , with  $x_0 \in A_{\text{inf}}$  and  $\epsilon \in \text{Fil}^n B_{\text{dR}}$ . Recall that  $A_{\text{inf}} \cap \text{Fil}^n B_{\text{dR}} = (\ker \Theta_K)^n$ . Consider the map

$$\begin{aligned} \mathcal{O}_{\overline{K}}^{(n-1)} &\rightarrow \text{Fil}^n B_{\text{dR}} / ((\ker \Theta_K)^n + \text{Fil}^{n+1} B_{\text{dR}}) \\ x &\mapsto \epsilon \end{aligned}$$

This map is an  $\mathcal{O}_K$ -linear derivation taking values in a  $\mathcal{O}_{\overline{K}}$ -module. By the universal property of  $\Omega^{(n)}$ , the map factors through  $d^{(n)}$ . In particular, its kernel contains  $\ker d^{(n)} = \mathcal{O}_{\overline{K}}^{(n)}$ .  $\square$

**Lemma 15.4.** *Let  $x \in \mathcal{O}_{\overline{K}}$ . Let  $P \in \mathcal{O}_K[X]$  be a polynomial satisfying  $P(x) = 0$ , and let  $r$  be a nonnegative integer such that  $P'(x) \mid p^r$  in  $\mathcal{O}_{\overline{K}}$ . For each nonnegative integer  $n$ , let  $a_n = (2^n - 1)r$ ,  $b_n = (2^{n+1} - 2n - 1)r$ . Then for any positive integer  $m$ ,  $p^{a_n} x^m \in \mathcal{O}_{\overline{K}}^{(n)}$ , and  $x^{p^{b_n}} \in \mathcal{O}_{\overline{K}}^{(n)}$ .*

*Proof.* Use induction on  $n$ . The base case  $n = 0$  is trivial. Now assume that for some fixed  $n$  and all  $m$ ,  $p^{a_n} x^m \in \mathcal{O}_{\overline{K}}^{(n)}$ , and  $x^{p^{b_n}} \in \mathcal{O}_{\overline{K}}^{(n)}$ . By repeated use of the product rule, we see that

$$(15.5) \quad d^{(n+1)}(p^{2a_n} x^m) = m p^{a_n} x^{m-1} d^{(n+1)}(p^{a_n} x)$$

for each  $m$ . In particular, this implies

$$0 = d^{(n+1)}(p^{2a_n} P(x)) = p^{a_n} P'(x) d^{(n+1)}(p^{a_n} x).$$

Hence

$$(15.6) \quad 0 = d^{(n+1)}(p^{2a_n+r} x).$$

Then multiplying (15.5) by  $p^r$  and applying (15.6) yields

$$0 = d^{(n+1)}(p^{2a_n+r} x^m) = d^{(n+1)}(p^{a_{n+1}} x^m).$$

In the case  $m = p^{b_n}$ , since  $p^r \mid m$ , we get the stronger result

$$0 = d^{(n+1)}(p^{2a_n} x^{p^{b_n}}) = p^{2a_n} d^{(n+1)}(x^{p^{b_n}}).$$

Then

$$d^{(n+1)}(x^{p^{b_{n+1}}}) = d^{(n+1)}(x^{p^{b_n+2a_n}}) = p^{2a_n} (x^{p^{b_n}})^{p^{2a_n}-1} d^{(n+1)}(x^{p^{b_n}}) = 0.$$

$\square$

## 16. $B_{\text{dR}}$ AND DIFFERENTIALS, CONTINUED

Let  $\pi$  be a uniformizer of  $K$ .

**Lemma 16.1.** *For any  $n$ , the map  $\mathcal{O}_{\overline{K}}^{(n)} \rightarrow \mathcal{O}_C/\pi = \mathcal{O}_{\overline{K}}/\pi$  is surjective.*

*Proof.* Let  $x \in \mathcal{O}_{\overline{K}}$ , and let  $P$  be the minimal polynomial for  $x$  over  $k$ . Let  $x_m$  satisfy  $x_m^{p^m} + \pi x_m = x$ . Then  $x_m^{p^m} \equiv x \pmod{\pi}$ . We claim that for sufficiently large  $m$ ,  $x_m^{p^m} \in \mathcal{O}_{\overline{K}}^{(n)}$ . Indeed, let  $P_m(X) = P(X^{p^m} + \pi X)$ ; then  $P_m(x_m) = 0$  and  $P'_m(x_m) = (p^m x_m^{p^m-1} + \pi)P'(x)$ , so  $|P'_m(x_m)| = |\pi||P'(x)|$ . Then the claim follows from Lemma 15.4.  $\square$

By Lemma 15.3, the map  $\mathcal{O}_{\overline{K}}^{(n)} \hookrightarrow B_{\text{dR}}^+ / \text{Fil}^{n+1} B_{\text{dR}}^+$  factors through  $A_{\text{inf},K}/(\ker \Theta_K)^{n+1}$ .

**Lemma 16.2.** *For any  $n, m$ ,*

$$\mathcal{O}_{\overline{K}}^{(n)} \rightarrow A_{\text{inf},K}/(\pi^m, (\ker \Theta_K)^{n+1})$$

*is surjective.*

*Proof.* Observe that  $A_{\text{inf}}/(\pi^m, (\ker \Theta_K)^{n+1})$  is generated as an  $\mathcal{O}_K$ -module by the elements  $[x]$  for  $x \in \mathcal{O}_{K^\flat}$ . By Lemma 16.1, for any  $r$ , we can find  $\tilde{x}_r \in \mathcal{O}_{\overline{K}}^{(n)}$  such that  $\tilde{x}_r$  and  $x^{(r)}$  have the same image in  $\mathcal{O}_C/p$ . So the image of  $\tilde{x}_r$  in  $A_{\text{inf},K}/(\ker \Theta_K)^{n+1}$  will be congruent to  $[x^{p^{-r}}]$  modulo  $(\pi, (\ker \Theta_K))$ . If  $r$  is sufficiently large, then the image of  $\tilde{x}_r^{p^r}$  will be congruent to  $[x]$  modulo  $(\pi^m, (\ker \Theta_K)^n)$ . (Note that the bound on  $r$  does not depend on  $x$ , only on  $m$  and  $n$ .)  $\square$

**Lemma 16.3.** *Let  $m, n$  be nonnegative integers. Consider the map*

$$\theta_{m,n}: \mathcal{O}_{\overline{K}}^{(n)}/p^m \rightarrow \mathcal{O}_C/p^m.$$

*We have*

$$(\ker \theta_{m,n})^{n+1} = 0.$$

*Proof.* We will use induction on  $n$ . The base case  $n = 0$  is trivial. Assume  $(\ker \theta_{m,n-1})^n = 0$ . It suffices to show that for  $x \in \ker \theta_{m,n}$ ,  $y \in (\ker \theta_{m,n})^n$ ,  $xy = 0$ . Choose lifts  $\tilde{x}, \tilde{y} \in \mathcal{O}_{\overline{K}}^{(n)}$ . By the induction hypothesis,  $\tilde{y} \in p^m \mathcal{O}_{\overline{K}}^{(n-1)}$ . Then  $\tilde{x} \cdot p^{-m} \tilde{y} \in \mathcal{O}_{\overline{K}}^{(n-1)}$  and

$$d^{(n)}(\tilde{x} \cdot p^{-m} \tilde{y}) = p^{-m} \tilde{y} \cdot d^{(n)} \tilde{x} + p^{-m} \tilde{x} \cdot d^{(n)} \tilde{y} = 0.$$

So  $\tilde{x} \tilde{y}$  is a multiple of  $p^m$  in  $\mathcal{O}_{\overline{K}}^{(n)}$ , implying  $xy = 0$ .  $\square$

*Proof of Theorem 15.1(2).* First, we prove item (2), that the map

$$\mathcal{O}_{\overline{K}}^{(n)}/p^m \rightarrow A_{\text{inf},K}/(p^m, (\ker \Theta_K)^{n+1})$$

is an isomorphism. Denote this map by  $f_{m,n}$ . We will construct an inverse map

$$g_{m,n}: A_{\text{inf}}/(p^m, (\ker \Theta_K)^{n+1}) \rightarrow \mathcal{O}_{\overline{K}}^{(n)}/p^m.$$

By a generalization of Proposition 10.10, to construct a ring homomorphism

$$A_{\text{inf},K}/p^m \rightarrow \mathcal{O}_{\overline{K}}^{(n)}/p^m,$$

it suffices to construct a multiplicative map

$$\mathcal{O}_{C^\flat} \rightarrow \mathcal{O}_{\overline{K}}^{(n)}/p^m$$

such that the induced map

$$\mathcal{O}_{C^\flat} \rightarrow \mathcal{O}_{\overline{K}}^{(n)}/\pi$$

is a ring homomorphism. For  $x \in \mathcal{O}_{C^\flat}$  and  $r$  a nonnegative integer, let  $\tilde{x}_r \in \mathcal{O}_{\overline{K}}^{(n)}/\pi^m$  such that  $\tilde{x}_r$  and  $x^{(r)}$  have the same image on  $\mathcal{O}_C/\pi^m$ . It follows from 16.3 that for sufficiently large  $r$ ,  $\tilde{x}_r^{p^r}$  will not depend on the choice of  $\tilde{x}_r$ . Then we choose the map  $x \mapsto \tilde{x}_r^{p^r}$ . The associated map  $A_{\text{inf},K}/\pi^m \rightarrow \mathcal{O}_{\overline{K}}^{(n)}$  is given by

$$(16.4) \quad \sum_i [x_i] \pi^i \mapsto \sum_i \tilde{x}_{i,r}^{p^r} \pi^i.$$

By Lemma 16.3, this map actually factors through  $A_{\text{inf},K}/(p^m, (\ker \Theta_K)^{n+1})$ . So we have constructed  $g_{m,n}$ . It is clear that  $f_{m,n} \circ g_{m,n} = 1$ .

Now we prove that  $g_{m,n} \circ f_{m,n} = 1$ . Since  $\mathcal{O}_{\overline{K}}^{(n)}$  has no  $p$ -torsion,  $\widehat{\mathcal{O}_{\overline{K}}^{(n)}}$  also has no  $p$ -torsion. So it suffices to show that the induced maps

$$\begin{aligned} f_n : \widehat{\mathcal{O}_{\overline{K}}^{(n)}}[1/p] &\rightarrow B_{\text{dR}}^+ / \text{Fil}^{n+1} B_{\text{dR}}^+ \\ g_n : B_{\text{dR}}^+ / \text{Fil}^{n+1} B_{\text{dR}}^+ &\rightarrow \widehat{\mathcal{O}_{\overline{K}}^{(n)}}[1/p] \end{aligned}$$

satisfy  $g_n \circ f_n = 1$ . We can construct a map  $\overline{K} \hookrightarrow \widehat{\mathcal{O}_{\overline{K}}^{(n)}}[1/p]$  by the same method as for  $\overline{K} \rightarrow B_{\text{dR}}$ . It is not hard to see that  $g_n \circ f_n$  fixes  $K$ , so  $g_n \circ f_n$  must send  $\overline{K}$  to itself. Since  $g_n \circ f_n$  induces the identity on the residue field  $C$ , it must fix  $\overline{K}$ . But  $\overline{K}$  is dense in  $\widehat{\mathcal{O}_{\overline{K}}^{(n)}}[1/p]$ , so  $g_n \circ f_n = 1$ . This concludes the proof of item (2).  $\square$

*Proof of Theorem 15.1(1,3,4).* Next, we prove item (1), that the preimage of  $A_{\text{inf},K}/(\ker \Theta_K)^{n+1}$  under the inclusion  $\overline{K} \hookrightarrow B_{\text{dR}}^+ / \text{Fil}^{n+1} B_{\text{dR}}^+$  is  $\mathcal{O}_{\overline{K}}^{(n)}$ . Denote the preimage by  $\mathcal{O}'$ .

Let  $x \in \mathcal{O}'$ . Then by Lemma 15.4, there exists  $m$  so that  $p^m x \in \mathcal{O}_{\overline{K}}^{(n)}$ . By item (2),  $\mathcal{O}_{\overline{K}}^{(n)}/p^m \rightarrow A_{\text{inf}}/(p^m, (\ker \Theta_K)^{(n+1)})$  is injective, so  $p^m x$  must be a multiple of  $p^m$  in  $\mathcal{O}_{\overline{K}}^{(n)}$  as well. Hence  $x \in \mathcal{O}_{\overline{K}}^{(n)}$ .

Item (3) follows immediately from items (1) and (2).

Finally, we prove item (4). Each element of  $\Omega^{(n)}$  is of the form  $\sum_i x_i d^{(n)} y_i$  for  $x_i \in \mathcal{O}_{\overline{K}}$  and  $y_i \in \mathcal{O}_{\overline{K}}^{(n-1)}$ . By Lemma 15.4, we can find  $m_i$  so that  $p^{m_i} d^{(n)} y_i = 0$ , and by Lemma 16.2, we can find  $z_i \in \mathcal{O}_{\overline{K}}^{(n)}$  so that  $x_i - z_i \in p^{m_i} \mathcal{O}_{\overline{K}}$ . Then  $\sum_i x_i d^{(n)} y_i = \sum_i z_i d^{(n)} y_i = \sum_i d^{(n)} (y_i z_i)$ .  $\square$

**Lemma 16.5.** *For any abelian group  $A$ , we have an isomorphism*

$$\text{Ext}_{\mathbb{Z}}^1(\mathbb{Q}_p/\mathbb{Z}_p, A) \cong \varprojlim_n A/p^n A.$$

*The right-hand side is the  $p$ -adic completion of  $A$ .*

*Proof.* There is a projective resolution

$$0 \rightarrow \bigoplus_{i=1}^{\infty} \mathbb{Z} \rightarrow \bigoplus_{i=1}^{\infty} \mathbb{Z} \rightarrow \mathbb{Q}_p/\mathbb{Z}_p \rightarrow 0.$$

If we denote the basis elements of  $\mathbb{Z}^{\oplus \mathbb{N}}$  by  $e_i$ , then the first map is given by

$$e_i \mapsto \begin{cases} pe_1, & i = 1 \\ pe_i - e_{i-1}, & i > 1 \end{cases}$$

and the second map is given by

$$e_i \mapsto p^{-i}.$$

So  $\text{Ext}_{\mathbb{Z}}^1(\mathbb{Q}_p/\mathbb{Z}_p, A)$  is the space of sequences  $a_0, a_1, a_2, \dots \in A$  modulo sequences of the form  $(pb_0, pb_1 - b_0, pb_2 - b_1, \dots)$ . For each  $n$ ,  $\sum_{i=1}^n p^{i-1} a_i$  is determined modulo  $p^n$ . Together, these sums determine an element of  $\varprojlim_n A/p^n A$ .  $\square$

**Lemma 16.6.** *For each positive integer  $n$ , there is a natural isomorphism*

$$\text{Hom}_{\mathbb{Z}_p}(\mathbb{Q}_p, \Omega^{(n)}) \cong \text{Fil}^n B_{\text{dR}} / \text{Fil}^{n+1} B_{\text{dR}}.$$

*Proof.* By Theorem 15.1(4), there is an exact sequence

$$0 \rightarrow \mathcal{O}_{\bar{K}}^{(n)} \rightarrow \mathcal{O}_{\bar{K}}^{(n-1)} \rightarrow \Omega^{(n)} \rightarrow 0.$$

So we get an exact sequence

$$0 \rightarrow \mathrm{Hom}_{\mathbb{Z}}(\mathbb{Q}_p/\mathbb{Z}_p, \Omega^{(n)}) \rightarrow A_{\mathrm{inf}, K}/(\ker \Theta_K)^{n+1} \rightarrow A_{\mathrm{inf}, K}/(\ker \Theta_K)^n \rightarrow 0,$$

where we used Lemma 16.5 and Theorem 15.1(2) to compute the  $\mathrm{Ext}^1$  groups. Observe that  $\mathrm{Hom}_{\mathbb{Z}}(\mathbb{Q}_p/\mathbb{Z}_p, \Omega^{(n)}) \cong \mathrm{Hom}_{\mathbb{Z}_p}(\mathbb{Q}_p/\mathbb{Z}_p, \Omega^{(n)})$ , and since  $\Omega^{(n)}$  is  $p$ -power torsion,  $\mathrm{Hom}_{\mathbb{Z}_p}(\mathbb{Q}_p/\mathbb{Z}_p, \Omega^{(n)}) \otimes_{\mathbb{Z}_p} \mathbb{Q}_p \cong \mathrm{Hom}_{\mathbb{Z}_p}(\mathbb{Q}_p, \Omega^{(n)})$ . Then inverting  $p$  gives an exact sequence

$$0 \rightarrow \mathrm{Hom}_{\mathbb{Z}_p}(\mathbb{Q}_p, \Omega^{(n)}) \rightarrow B_{\mathrm{dR}}^+/\mathrm{Fil}^{n+1} B_{\mathrm{dR}}^+ \rightarrow B_{\mathrm{dR}}^+/\mathrm{Fil}^n B_{\mathrm{dR}}^+ \rightarrow 0.$$

□

## 17. SITES AND COHOMOLOGY

Recall that in the first lecture, we considered an isomorphism between the singular and de Rham cohomologies of a complex manifold. This isomorphism can be understood in terms of sheaf cohomology.

Let  $X$  be a complex manifold. A presheaf of sets on  $X$  is a contravariant functor  $\mathcal{F}$  from the category of open subsets of  $X$  to the category of sets. We say that  $\mathcal{F}$  is a sheaf if for every open  $U \subset X$  and every covering  $U = \bigcup_{i \in I} U_i$ ,

$$\mathcal{F}(U) \longrightarrow \prod_{i \in I} \mathcal{F}(U_i) \rightrightarrows \prod_{i, j \in I} \mathcal{F}(U_i \cap U_j)$$

is an equalizer diagram. In other words,  $\mathcal{F}(U)$  is the subset of  $\prod_{i \in I} \mathcal{F}(U_i)$  such that for each  $i, j \in I$ , the projections to  $\mathcal{F}(U_i)$  and  $\mathcal{F}(U_j)$  have the same image in  $\mathcal{F}(U_i \cap U_j)$ . A morphism of (pre)sheaves  $\mathcal{F} \rightarrow \mathcal{G}$  is a natural transformation of functors.

The categories of sheaves and presheaves of abelian groups on  $X$  are defined similarly. They are abelian categories.

A sheaf  $\mathcal{I}$  of abelian groups on  $X$  is called *injective* if the functor  $\mathrm{Hom}(-, \mathcal{I})$  is exact. An *injective resolution* of a sheaf  $\mathcal{F}$  of abelian groups on  $X$  is a long exact sequence

$$0 \rightarrow \mathcal{F} \rightarrow \mathcal{I}_0 \rightarrow \mathcal{I}_1 \rightarrow \cdots$$

where  $\mathcal{I}_0, \mathcal{I}_1, \dots$  are injective sheaves.

It can be shown that every sheaf of abelian groups on  $X$  admits an injection into an injective sheaf. This implies that every sheaf of abelian groups on  $X$  admits an injective resolution.

The cohomology groups  $H^n(X, \mathcal{F})$  are defined by the cohomology groups of the complex

$$0 \rightarrow \mathcal{I}_0(X) \rightarrow \mathcal{I}_1(X) \rightarrow \cdots$$

This definition is independent of the injective resolution chosen.

A sheaf  $\mathcal{G}$  is called *acyclic* if  $H^n(X, \mathcal{G}) = 0$  for all  $n > 0$ . One can actually compute the cohomology groups  $H^n(X, \mathcal{F})$  using any resolution of  $\mathcal{F}$  by acyclic sheaves.

For any complex manifold  $X$  and any ring  $A$ , the complex of singular cochains on  $X$  is an acyclic resolution of the constant sheaf  $A$ . So

$$H_{\mathrm{sing}}^i(X, A) = H_{\mathrm{sheaf}}^i(X, A).$$

If  $C^\bullet$  is a complex of sheaves on  $X$ , then an injective (resp. acyclic) of  $C^\bullet$  is a quasi-isomorphism of complexes  $C^\bullet \rightarrow \mathcal{I}^\bullet$ , where the objects of  $\mathcal{I}^\bullet$  are injective (resp. acyclic). (A quasi-isomorphism of complexes is a morphism that induces an isomorphism on cohomology.) Then the hypercohomology of  $H^*(X, C^\bullet)$  is defined to be the cohomology of the complex

$$0 \rightarrow \mathcal{I}_0(X) \rightarrow \mathcal{I}_1(X) \rightarrow \cdots$$

The de Rham complex  $\Omega_X^\bullet$  is the complex

$$0 \rightarrow \mathcal{O}_X \xrightarrow{d} \Omega_{X/\mathbb{C}}^1 \xrightarrow{d} \cdots \xrightarrow{d} \Omega_{X/\mathbb{C}}^n \rightarrow 0.$$

If  $X$  is an open subset of  $\mathbb{C}^n$ , then the  $\Omega_{X/\mathbb{C}}^i$  are already acyclic. We implicitly used this fact in the first lecture when computing the de Rham cohomology of  $\mathbb{C}^\times$ .

The analytic de Rham cohomology of  $X$  is defined to be the hypercohomology of  $\Omega_X^\bullet$ . The Poincaré lemma states that  $\mathbb{C}[0] \rightarrow \Omega_X^\bullet$  is a quasi-isomorphism. Therefore,

$$H_{\text{dR}}^i(X) = H_{\text{sheaf}}^i(X, \mathbb{C}).$$

If  $X$  is a smooth algebraic variety over a field  $K$ , then we can define the algebraic de Rham cohomology similarly. If  $X$  is a proper variety over  $\mathbb{C}$ , then there is an isomorphism

$$H_{\text{dR}}^i(X) \cong H_{\text{dR}}^i(X(\mathbb{C})),$$

where  $X(\mathbb{C})$  is considered as a complex analytic space.

However, the Poincaré lemma does not hold for the algebraic de Rham complex:  $K[0] \rightarrow \Omega_{X/K}^\bullet$  is generally not a quasi-isomorphism. For example, if  $X = \mathbb{A}_K^1 \setminus \{0\}$ , then  $\frac{dz}{z}$  does not have an antiderivative locally in the Zariski topology. The cohomology  $H^i(X, K)$  is not very interesting: it is zero if  $i > 0$ .

In practice, we usually use Čech cohomology to compute de Rham cohomology. If  $X$  is an affine algebraic variety, then the sheaves  $\Omega_{X/K}^i$  are acyclic. More generally, if  $X$  is a separated algebraic variety, then we can write  $X$  as a finite union  $X = \bigcup_{i \in I} X_i$ , where the  $X_i$ 's are affine, and the intersections of the  $X_i$ 's are also affine. Define a complex  $C^\bullet$  as follows. Let

$$C^n = \bigoplus_{J \subset I} \Omega^{n-|J|} \left( \bigcap_{j \in J} X_j \right).$$

Given  $\omega \in \Omega^{n-|J|} \left( \bigcap_{j \in J} X_j \right)$ , the  $\Omega^{n+1-|J|} \left( \bigcap_{j \in J} X_j \right)$ -part of its differential is  $d\omega$ .

Choose an ordering of  $I$ . If  $i \notin J$ , then the  $\Omega^{n+1-|J \cup \{i\}|} \left( \bigcap_{j \in J \cup \{i\}} X_j \right)$ -part of the differential of  $\omega$  is  $(-1)^{n+|\{j \in J | j < i\}|}$  times the restriction of  $\omega$ . All other parts are zero. Then the de Rham cohomology of  $X$  is the cohomology of  $C^\bullet$ .

## 18. SITES, ÉTALE AND PROÉTALE COHOMOLOGY

Étale cohomology is supposed to be an analogue of singular cohomology for more general kinds of spaces. To define étale cohomology, we need to use sites.

**Definition 18.1.** A site is a category  $\mathcal{C}$  along a set  $\text{Cov}(\mathcal{C})$  of morphisms with fixed target  $\{U_i \rightarrow U\}_{i \in I}$ , satisfying the following axioms:

- (1) If  $V \rightarrow U$  is an isomorphism then  $\{V \rightarrow U\} \in \text{Cov}(\mathcal{C})$ .



- (2) If  $\{U_i \rightarrow U\}_{i \in I} \in \text{Cov}(\mathcal{C})$  and for each  $i \in I$ ,  $\{V_{ij} \rightarrow U_i\}_{j \in J_i} \in \text{Cov}(\mathcal{C})$ , then  $\{V_{ij} \rightarrow U\}_{i \in I, j \in J_i} \in \text{Cov}(\mathcal{C})$ .
- (3) If  $\{U_i \rightarrow U\}_{i \in I} \in \text{Cov}(\mathcal{C})$  and  $V \rightarrow U$  is a morphism in  $\mathcal{C}$ , then for all  $i \in I$ ,  $U_i \times_U V$  exists, and  $\{U_i \times_U V\}_{i \in I} \in \text{Cov}(\mathcal{C})$

For any topological space  $X$ , we can define a site by taking  $\mathcal{C}$  to be the category of open sets of  $X$  and coverings to be open coverings in the usual sense. Note that the fiber product is just intersection.

Now let  $X$  be a scheme. Recall that a morphism of schemes  $Y \rightarrow Z$  is called *étale* if it is smooth of relative dimension 0. Define the *small étale site*  $X_{\text{ét}}$  as follows. The underlying category is the category of étale morphisms  $U \rightarrow X$ . A collection  $U_i \rightarrow X$  is a covering if the images of the  $U_i$  cover  $X$ . (There is also a big étale site, where the category includes all schemes over  $X$ , but morphisms are still jointly surjective collections of étale maps.)

Sheaves and cohomology can be defined on any site. One can show that if  $X$  is an algebraic variety over  $\mathbb{C}$ , then for any  $N$ ,

$$H^i(X(\mathbb{C}), \mathbb{Z}/N\mathbb{Z}) \cong H^i(X_{\text{ét}}, \mathbb{Z}/N\mathbb{Z}).$$

The reason that we need to use  $\mathbb{Z}/N\mathbb{Z}$  coefficients is that finite covers of  $X(\mathbb{C})$  are in bijection with finite covers of  $X$ , but the same is not true of infinite covers. To deal with this issue, we define

$$H_{\text{ét}}^i(X, \mathbb{Z}_p) = \varprojlim_n H^i(X, \mathbb{Z}/p^n\mathbb{Z})$$

$$H_{\text{ét}}^i(X, \mathbb{Q}_p) = H^i(X, \mathbb{Z}_p) \otimes_{\mathbb{Z}_p} \mathbb{Q}_p$$

for any algebraic variety  $X$ . Using this definition, if  $X$  is an algebraic variety over  $\mathbb{C}$ , then

$$H^i(X(\mathbb{C}), \mathbb{Z}_p) \cong H^i(X_{\text{ét}}, \mathbb{Z}_p).$$

$$H^i(X(\mathbb{C}), \mathbb{Q}_p) \cong H^i(X_{\text{ét}}, \mathbb{Q}_p).$$

Normally, we consider cohomology of varieties over an algebraically closed field, as the notion of cohomology most closely matches our geometric intuition in that case. For example, if  $K$  is a field, then

$$H^i((\text{Spec } K)_{\text{ét}}, \Lambda) = H_{\text{cts}}^i(\text{Gal}(K^{\text{sep}}/K), \Lambda).$$

So the cohomology of the spectrum of a non-algebraically closed field can be quite complicated.

If  $X$  is a variety over a field  $K$  (that is not necessarily algebraically closed), then we can consider the cohomology groups  $H^i(X_{K^{\text{sep}}}, \Lambda)$ , and these have a  $\text{Gal}(K^{\text{sep}}/K)$  action.

The cohomology  $H^i(X_{\text{ét}}, \Lambda)$  (without the base change to  $K^{\text{sep}}$ ) is sometimes useful. See for example [LZ] for an application to Euler systems.

Actually, there is a way to define  $H_{\text{ét}}^i(X, \mathbb{Z}_p)$  as cohomology on an actual site, called the “proétale site”. A morphism of schemes  $f: X \rightarrow Y$  is called *weakly étale* if  $f$  and  $\Delta_f: X \rightarrow X \times_Y X$  are both flat. Coverings in the proétale site are defined to be collections of weakly étale maps  $X_i \rightarrow Y$  such that for each affine open  $V \subset Y$ , there is a finite collection of affine opens  $U_i \subset X_i$  whose images cover  $V$ . Then

$$H_{\text{ét}}^i(X, \mathbb{Z}_p) = H^i(X_{\text{proét}}, \mathbb{Z}_p).$$

The site is called the “proétale site” because, if we have a tower

$$\cdots \rightarrow X_2 \rightarrow X_1 \rightarrow X$$

of schemes with étale and affine transition maps, then the projection

$$\varprojlim_i X_i \rightarrow X$$

is weakly étale. It turns out that the property of being proétale is not local, so it’s better to use the property of being weakly étale for defining a site.

If  $X$  is an algebraic variety over a  $p$ -adic field  $K$ , we now know the definitions of the objects appearing in the de Rham comparison isomorphism

$$H_{\mathrm{dR}}^i(X) \otimes_K B_{\mathrm{dR}} \cong H_{\mathrm{ét}}^i(X_{\overline{K}}, \mathbb{Z}_p) \otimes_{\mathbb{Z}_p} B_{\mathrm{dR}}.$$

(However, proving the theorem requires getting much deeper in to  $p$ -adic geometry.)

### 19. HODGE-TATE DECOMPOSITION FOR ABELIAN VARIETIES

Let  $X$  be a proper variety over a  $p$ -adic field  $K$ . Recall the comparison isomorphism

$$(19.1) \quad H_{\mathrm{dR}}^n(X) \otimes_K B_{\mathrm{dR}} \cong H_{\mathrm{ét}}^n(X_{\overline{K}}, \mathbb{Z}_p) \otimes_{\mathbb{Z}_p} B_{\mathrm{dR}}.$$

This is an isomorphism of filtered vector spaces with  $G_K$ -action. Here,  $B_{\mathrm{dR}}$  has the usual filtration,  $H_{\mathrm{ét}}^n(X_{\overline{K}}, \mathbb{Z}_p)$  has the trivial filtration, and the filtration on  $H_{\mathrm{dR}}^n(X)$  is given by

$$\mathrm{Fil}^i H_{\mathrm{dR}}^n(X) = \mathrm{im} \left( (H^n(X, \sigma_{\geq i} \Omega_{X/K}^\bullet) \rightarrow H^n(X, \Omega_{X/K}^\bullet) \right).$$

Here,  $\sigma_{\geq i} \Omega_{X/K}^\bullet$  is the complex

$$\Omega_{X/K}^i \rightarrow \Omega_{X/K}^{i+1} \rightarrow \cdots \rightarrow \Omega_{X/K}^{\dim X}.$$

For each  $i$ , there is a natural long exact sequence

$$\cdots \rightarrow H^n(X, \sigma_{\geq i} \Omega_{X/K}^\bullet) \rightarrow H^n(X, \sigma_{\geq i-1} \Omega_{X/K}^\bullet) \rightarrow H^n(X, \Omega_{X/K}^i[-i]) \rightarrow \cdots$$

It is possible to show that the boundary maps are zero. Hence we can show by induction that the maps

$$H^n(X, \sigma_{\geq i} \Omega_{X/K}^\bullet) \rightarrow H^n(X, \Omega_{X/K}^\bullet)$$

are injective. So we can identify  $\mathrm{Fil}^i H_{\mathrm{dR}}^n(X)$  with  $H^n(X, \sigma_{\geq i} \Omega_{X/K}^\bullet)$ , and

$$\mathrm{gr}^i H_{\mathrm{dR}}^n(X) \cong H^n(X, \Omega_{X/K}^i[-i]) = H^{n-i}(X, \Omega_{X/K}^i).$$

Taking the zeroth graded piece of (19.1) gives an isomorphism

$$(19.2) \quad \bigoplus_{i=0}^n H^{n-i}(X, \Omega_{X/K}^i) \otimes_K C(-i) \cong H_{\mathrm{ét}}^n(X_{\overline{K}}, \mathbb{Z}_p) \otimes_{\mathbb{Z}_p} C.$$

An *abelian variety* over  $K$  is proper group scheme over  $K$  that is geometrically reduced and irreducible. If  $X$  is an abelian variety, then the decomposition (19.2) can actually be made fairly explicit. In that case, all cohomology groups are wedge powers of  $H^1$ , so it suffices to consider  $n = 1$ . Furthermore,  $H_{\mathrm{ét}}^1(X_{\overline{K}}, \mathbb{Z}_p)$  is dual to the Tate module  $T_p(X) := \varprojlim_n X(\overline{K})[p^n]$ . Therefore, giving a  $C$ -linear map  $H^0(X, \Omega_{X/K}^1) \otimes_K C(-1) \rightarrow H_{\mathrm{ét}}^1(X_{\overline{K}}, \mathbb{Z}_p) \otimes_{\mathbb{Z}_p} C$  is equivalent to giving a map

$$H^0(X, \Omega_{X/K}^1) \times T_p(X) \rightarrow C(1)$$

that is  $K$ -linear in the first variable and  $\mathbb{Z}_p$ -linear in the second.

We will sketch the construction of the map, but we refer the reader to [Fon82] for the technical details.

Let  $\mathfrak{X}$  be a proper flat model of  $X$  over  $\mathcal{O}_K$ . Recall that we defined

$$\Omega^{(1)} = \Omega_{\mathcal{O}_{\overline{K}}/\mathcal{O}_K}^1.$$

We can define a map

$$\begin{aligned} H^0(\mathfrak{X}, \Omega_{\mathfrak{X}/\mathcal{O}_K}^1) \times \mathfrak{X}(\mathcal{O}_{\overline{K}}) &\rightarrow \Omega^{(1)} \\ (\omega, x) &\mapsto x^*(\omega). \end{aligned}$$

It is possible to use the group law on the generic fiber show that for some  $r$ , the restriction

$$p^r H^0(\mathfrak{X}, \Omega_{\mathfrak{X}/\mathcal{O}_K}^1) \times \mathfrak{X}(\mathcal{O}_{\overline{K}}) \rightarrow \Omega^{(1)}$$

is bilinear.

By the valuative criterion of properness,  $\mathfrak{X}(\mathcal{O}_{\overline{K}}) = X(K)$ . Using  $\mathrm{Hom}(\mathbb{Q}_p/\mathbb{Z}_p, X(K)) \cong T_p(X)$  and  $\mathrm{Hom}(\mathbb{Q}_p/\mathbb{Z}_p, \Omega^{(1)}) \cong \ker \Theta_K / (\ker \Theta_K)^2$ , we obtain a map

$$p^r H^0(\mathfrak{X}, \Omega_{\mathfrak{X}/\mathcal{O}_K}^1) \times T_p(X) \rightarrow (\ker \Theta_K) / (\ker \Theta_K)^2.$$

Finally, we invert  $p$  to get a bilinear map

$$H^0(X, \Omega_{X/K}^1) \times T_p(X) \rightarrow C(1).$$

Now we have a constructed map (which can be shown to be injective)

$$(19.3) \quad H^0(X, \Omega_{X/K}^1) \otimes_K C(-1) \hookrightarrow H_{\text{ét}}^1(X_{\overline{K}}, \mathbb{Z}_p) \otimes_{\mathbb{Z}_p} C.$$

If  $X^\vee$  is the dual abelian variety, then we get a map

$$H^0(X^\vee, \Omega_{X^\vee/K}^1) \otimes_K C(-1) \hookrightarrow H_{\text{ét}}^1(X_{\overline{K}}^\vee, \mathbb{Z}_p) \otimes_{\mathbb{Z}_p} C.$$

The Weil pairing determines a perfect pairing

$$H_{\text{ét}}^1(X_{\overline{K}}, \mathbb{Z}_p) \times H_{\text{ét}}^1(X_{\overline{K}}^\vee, \mathbb{Z}_p) \rightarrow \mathbb{Z}_p(-1)$$

and there is also a perfect pairing

$$H^1(X, \mathcal{O}_X) \times H^0(X^\vee, \Omega_{X^\vee/K}^1) \rightarrow K.$$

So we get a map

$$H^1(X, \mathcal{O}_X)^* \otimes_K C(-1) \hookrightarrow H_{\text{ét}}^1(X_{\overline{K}}, \mathbb{Z}_p)^* \otimes_{\mathbb{Z}_p} C(-1).$$

Taking the dual and twisting, we get a map

$$(19.4) \quad H_{\text{ét}}^1(X_{\overline{K}}, \mathbb{Z}_p) \otimes_{\mathbb{Z}_p} C \rightarrow H^1(X, \mathcal{O}_X) \otimes_K C.$$

The composite of (19.3) and (19.4) must be zero since  $C(1)^{G_K} = 0$ . Then, by dimension counting, the sequence

$$(19.5) \quad 0 \rightarrow H^0(X, \Omega_{X/K}^1) \otimes_K C(-1) \rightarrow H_{\text{ét}}^1(X_{\overline{K}}, \mathbb{Z}_p) \otimes_{\mathbb{Z}_p} C \rightarrow H^1(X, \mathcal{O}_X) \otimes_K C \rightarrow 0$$

must be exact. Since  $H^1(G_K, C(-1)) = 0$  (see Remark 13.2), this sequence has a  $G_K$ -equivariant splitting. Since  $C(-1)^{G_K} = 0$ , this splitting is unique.

## 20. HUBER RINGS

In the remaining part of the course, we will do some geometry. We are interested in developing a theory similar to the theory of schemes, but where the rings are allowed to have topology. Clausen–Scholze have proposed a very general framework for analytic geometry, and you have seen some of this framework in the learning seminar on real local Langlands. This theory is very technical and involves the use of condensed mathematics and  $\infty$ -categories. We will instead use the older, less general framework of adic spaces, developed by Huber. In the cases where Huber’s theory works, it should be equivalent to that of Clausen–Scholze.

An example of the type of space we want to consider is the closed  $p$ -adic unit disc over  $\mathbb{Q}_p$ . A function on the disc should be an element of

$$\mathbb{Q}_p\langle T \rangle = \left\{ \sum_{n=0}^{\infty} a_n T^n \mid a_n \in \mathbb{Q}_p, \lim_{n \rightarrow \infty} a_n = 0 \right\}.$$

Let  $\mathbb{Z}_p\langle T \rangle \subset \mathbb{Q}_p\langle T \rangle$  be the subring consisting of power series with coefficients in  $\mathbb{Z}_p$ . Then it is natural to give  $\mathbb{Q}_p\langle T \rangle$  a topology by letting the sets  $p^n \mathbb{Z}_p\langle T \rangle$  be a basis of open neighborhoods of 0.

**Definition 20.1.** A *Huber ring* is a (Hausdorff) topological ring  $A$  such that there exists an open subring  $A_0 \subset A$  and a finitely generated ideal  $I \subset A_0$  so that  $A_0$  has the  $I$ -adic topology (powers of  $I$  form a basis of neighborhoods of the identity).

We say that  $A_0$  is a *ring of definition* of  $A$  and  $I$  is an *ideal of definition* of  $A_0$ .

*Remark 20.2.* Huber rings are “solid” in the sense of Clausen–Scholze.

*Example 20.3.* The following topological rings are Huber rings.

- Any ring with the discrete topology ( $A_0 = \text{any subring}$ ,  $I = 0$ ).
- $\mathbb{Z}_p$ , ( $A_0 = \mathbb{Z}_p$ ,  $I = (p)$ )
- $\mathbb{Q}_p$ , ( $A_0 = \mathbb{Z}_p$ ,  $I = (p)$ )
- More generally, any field  $K$  with a nonarchimedean norm ( $A_0 = \mathcal{O}_K := \{x \in K \mid |x| \leq 1\}$ ,  $I = (\pi)$  for any  $\pi \in K^\times$  satisfying  $|\pi| < 1$ ). Note that if  $|K^\times|$  is not discrete, then the maximal ideal of  $\mathcal{O}_K$  is not finitely generated, so we cannot take  $I$  to be the maximal ideal of  $\mathcal{O}_K$ .
- The Tate algebra  $K\langle T \rangle = \{\sum_{n=0}^{\infty} a_n T^n \mid \lim_{n \rightarrow \infty} a_n = 0\}$  ( $A_0 = \mathcal{O}_K\langle T \rangle := \text{series with } |a_n| \leq 1 \text{ for all } n$ ,  $I = (\pi)$ )
- $\mathbb{Z}_p[[T]]$  ( $A_0 = \mathbb{Z}_p[[T]]$ ,  $I = (p, T)$ )
- $W(\mathcal{O}_{K^\flat})$  for a perfectoid field  $K$  ( $A = W(\mathcal{O}_{K^\flat})$ ,  $I = (p, [\pi])$  where  $\pi \in \mathcal{O}_{K^\flat}$  satisfies  $0 < |\pi| < 1$ )

**Definition 20.4.** Let  $A$  be a ring. A *valuation* on  $A$  is a map  $|\cdot| : A \rightarrow \Gamma \cup \{0\}$ , where  $\Gamma$  is a totally ordered abelian group (written multiplicatively), satisfying the following properties:

- (1)  $|xy| = |x||y|$  for all  $x, y \in A$
- (2)  $|x + y| \leq \max(|x|, |y|)$  for all  $x, y \in A$
- (3)  $|0| = 0$ ,  $|1| = 1$

If  $A$  is a topological ring, then we say that a valuation is *continuous* if for all  $\gamma \in \Gamma$ ,  $\{a \in A : |a| < \gamma\}$  is open.

We say that two valuations  $|\cdot|$  and  $|\cdot|'$  are *equivalent* if  $|a| \leq |b| \iff |a|' \leq |b|'$  for all  $a, b \in A$ .

**Definition 20.5.** Let  $A$  be a topological ring. Define  $\text{Cont}(A)$  to be the set of equivalence classes of continuous valuations of  $A$ .

Give  $\text{Cont}(A)$  the topology with a sub-basis of open sets consisting of sets the form

$$\{x \mid |f(x)| \leq |g(x)| \neq 0\}.$$

for  $f, g \in A$ . Here  $|f(x)|, |g(x)|$  denote the valuations of  $f$  and  $g$  under a representative of the equivalence class  $x$ .

**Definition 20.6.** Let  $A$  be a topological ring. A subset  $S$  of  $A$  is *bounded* if for all open neighborhoods  $U$  of 0, there is an open neighborhood  $V$  of 0 such that  $VS \subset U$ .

**Definition 20.7.** Let  $A$  be a Huber ring. An element  $f \in A$  is *power-bounded* if  $\{f^n \mid n \in \mathbb{N}\}$  is bounded.

We will write  $A^\circ$  for the subring of power-bounded elements of  $A$ .

**Definition 20.8.** Let  $A$  be a Huber ring. A subring  $A^+ \subset A^\circ$  is a *ring of integral elements* if it is open and integrally closed in  $A$ .

A *Huber pair* is a pair  $(A, A^+)$ , where  $A$  is a Huber ring, and  $A^+ \subset A$  is a ring of integral elements.

For a Huber pair  $(A, A^+)$ , define  $\text{Spa}(A, A^+) \subset \text{Cont}(A)$  to be the subspace consisting of those valuations  $x$  for which  $|f(x)| \leq 1$  for all  $f \in A^+$ .

*Example 20.9.* The points of  $\text{Spa}(\mathbb{Z}, \mathbb{Z})$  are as follows:

- (1)  $\eta: |n(\eta)| = \begin{cases} 1, & n \neq 0 \\ 0, & n = 0 \end{cases}$
- (2) For each prime  $p$ ,  $s_p: |n(s_p)| = \begin{cases} 1, & p \nmid n \\ 0, & p \mid n \end{cases}$
- (3) For each prime  $p$ ,  $\eta_p: |n(\eta_p)| = |n|_p$  (the usual  $p$ -adic absolute value)

The point  $s_p$  is closed, while  $\overline{\{\eta_p\}} = \{\eta_p, s_p\}$  and  $\overline{\{\eta\}} = \text{Spa}(\mathbb{Z}, \mathbb{Z})$ .

*Example 20.10.* Let  $K$  be a nonarchimedean field, and let  $\mathcal{O}_K$  be its ring of integral elements. The space  $\text{Spa}(K\langle T \rangle, \mathcal{O}_K\langle T \rangle)$  is called the *closed unit disc* over  $K$ .

Let  $C$  be an algebraically closed nonarchimedean field. Let  $k$  be the residue field of  $C$ . The points on the closed unit disc over  $C$  can be classified into five types:

- For any  $\alpha \in \mathcal{O}_C$ , there is a point  $f \mapsto |f(\alpha)|$ . These are called “Type 1” points.
- For  $\alpha \in \mathcal{O}_C$ ,  $r \in (0, 1]$ , let  $B(\alpha, r)$  denote the closed ball of radius  $r$  centered at  $\alpha$  in  $\mathcal{O}_{C_p}$ . Then

$$f \mapsto \sup_{\alpha \in B(\alpha, r)} |f(\alpha)|$$

is a point of  $D$ . Equivalently, we can describe this norm by

$$\sum_{n=0}^{\infty} a_n (T - \alpha)^n \mapsto \sup_n |a_n| r^n.$$

We call this point “Type 2” if  $r \in \mathbb{Q}_{>0}$  (i.e. it is the valuation of some element of  $\mathcal{O}_C$ ), and “Type 3” otherwise.

- The ring  $C$  is may not spherically complete, i.e. there may exist descending sequences  $B_1 \supset B_2 \supset \cdots$  of closed balls with empty intersection. (Since  $\mathbb{C}_p$  is complete, any descending sequence of discs whose radii go to zero have a common intersection point. But this need not be the case if the radii do not go to zero.) For example,  $\widehat{\mathbb{Q}}_p$  is not spherically complete.

There is a valuation

$$f \mapsto \inf_i \sup_{\alpha \in B_i} |f(\alpha)|.$$

These are called Type 4 points or “dead ends”.

- As mentioned before, we can consider the value group  $\mathbb{R}_{>0} \times \mathbb{R}_{>0}$  with lexicographic ordering. For any  $B(\alpha, r)$  with  $r$  rational, we can define the norms

$$\begin{aligned} \sum_{n=0}^{\infty} a_n (T - \alpha)^n &\mapsto \sup_n (|a_n| r^n, p^{-n}) \\ \sum_{n=0}^{\infty} a_n (T - \alpha)^n &\mapsto \sup_n (|a_n| r^n, p^n) \end{aligned}$$

(If  $r = 1$ , then the latter norm belongs to  $\text{Cont } \mathbb{C}_p \langle T \rangle$  but not to the closed unit disc.) These are called Type 5 points. Note that for  $r$  irrational, the above definition makes sense, but we end up with a norm that is equivalent to the Type 3 norm described above.

All points except for Type 2 points are closed. The closure of a Type 2 point looks like  $\mathbb{P}_{\mathbb{F}_p}^1$  (or  $\mathbb{A}_{\mathbb{F}_p}^1$  if  $r = 1$ ). The Type 2 point is the generic point and the Type 5 points are the closed points.

*Example 20.11.* The ring  $\mathcal{O}_K + \mathfrak{m}_K T \mathcal{O}_K \langle T \rangle$  is also a ring of integral elements of  $K \langle T \rangle$ . Then  $\text{Spa}(K \langle T \rangle, \mathcal{O}_K + \mathfrak{m}_K T \mathcal{O}_K \langle T \rangle)$  contains all points of  $\text{Spa}(K \langle T \rangle, \mathcal{O}_K \langle T \rangle)$ , plus one additional point where  $|T|$  is infinitesimally greater than 1.

*Example 20.12.* In the previous examples, the ring of integral elements is also a ring of definition. But this is not always the case. For example, if  $A = \mathbb{Q}_p[\epsilon]/(\epsilon^2)$ , then the only possible ring of integral elements is  $A^\circ = \mathbb{Z}_p + \mathbb{Q}_p \epsilon$ , but this ring is not a ring of definition.

**Definition 20.13.** A *spectral space* is a topological space that is homeomorphic to the spectrum of some ring. Equivalently, a spectral space is a limit of finite  $T_0$  spaces.

**Theorem 20.14.** For any Huber pair  $(A, A^+)$ ,  $\text{Spa}(A, A^+)$  is spectral.

## 21. ADIC SPACES

**Definition 21.1.** A *morphism of Huber pairs*  $(A, A^+) \rightarrow (B, B^+)$  is a continuous map  $A \rightarrow B$  such that the image of  $A^+$  is contained in  $B^+$ .

**Proposition 21.2.**  $\text{Spa}$  is a contravariant functor: any morphism of Huber pairs  $(A, A^+) \rightarrow (B, B^+)$  determines a continuous map  $\text{Spa}(B, B^+) \rightarrow \text{Spa}(A, A^+)$ .

**Proposition 21.3.** For any Huber pair, we can form the completion  $(\widehat{A}, \widehat{A}^+)$ . The induced morphism  $\text{Spa}(\widehat{A}, \widehat{A}^+) \rightarrow \text{Spa}(A, A^+)$  is a homeomorphism.

*Remark 21.4.* As you might expect, the completion of a Huber ring is complete. It is important that the ideal of definition is required to be finite—the completion of a ring with respect to an infinitely generated ideal need not be complete!

**Proposition 21.5.** *Let  $(A, A^+)$  be a complete Huber pair.*

- (1) *If  $A \neq 0$ , then  $\mathrm{Spa}(A, A^+)$  is nonempty.*
- (2) *One has  $A^+ = \{f \in A \mid |f(x)| \leq 1 \forall x \in X\}$ .*
- (3) *An element  $f \in A$  is invertible iff for all  $x \in X$ ,  $|f(x)| \neq 0$ .*

**Definition 21.6.** A *rational subset* of  $\mathrm{Spa}(A, A^+)$  is a subset defined by inequalities  $|a_1|, |a_2|, \dots, |a_n| \leq |a_0| \neq 0$ , where  $a_0, a_1, \dots, a_n \in A$  generate an open ideal.

*Remark 21.7.* It is important that  $a_0, \dots, a_n$  generate an open ideal of  $A$ . Choose a ring of definition  $A_0$  and ideal of definition  $I$ . In order for multiplication by  $1/a_0$  to be continuous, we need  $\frac{1}{a_0}I^n \subset A_0[a_1/a_0, \dots, a_n/a_0]$  for some  $n$ . It suffices to show that  $I^n \subset (a_0, \dots, a_n) \cdot A_0$  for large enough  $n$ , i.e. we need the  $A_0$ -submodule of  $A$  generated by  $a_0, \dots, a_n$  be open. One can show that this is equivalent to  $a_0, \dots, a_n$  generating an open ideal of  $A$ .

*Example 21.8.* If  $a_0, \dots, a_n$  do not generate an open ideal, then the locus  $|a_1|, \dots, |a_n| \leq |a_0| \neq 0$  may not be quasicompact, which would imply that it cannot be of the form  $\mathrm{Spa}(A, A^+)$ . For example, if  $X = \mathrm{Spa}(\mathbb{Q}_p\langle T \rangle, \mathbb{Z}_p\langle T \rangle)$  is the closed unit disc, then the locus  $|T| \neq 0$  is the punctured disc, which is not quasicompact.

**Lemma 21.9.** *Rational subsets form a basis for the topology of  $\mathrm{Spa}(A, A^+)$ .*

*Proof.* An intersection of two rational subsets is rational:

$$X\left(\frac{T}{a_0}\right) \cap X\left(\frac{S}{b_0}\right) = X\left(\frac{Tb_0 \cup a_0S \cup TS}{a_0b_0}\right).$$

It remains to show that for any  $f, g \in A$ , the open subset  $U$  defined by  $|f| \leq |g| \neq 0$  can be covered by rational subsets. Let  $a_1, \dots, a_n$  be generators of an ideal of definition of some ring of definition  $A_0$  of  $A$ . For  $r \in \mathbb{N}$ , let  $T_r = \{f, a_1^r, \dots, a_n^r\}$ . Since  $a_1^r, \dots, a_n^r$  generate an open ideal of  $A_0$ , they also generate an open ideal of  $A$ . Hence the  $X\left(\frac{T_r}{g}\right)$  are rational subsets of  $X$  that are contained in  $U$ . We claim that they cover  $U$ . Indeed, let  $x \in U$ . Since  $|g(x)| \neq 0$  and  $\lim_{r \rightarrow \infty} a_i^r = 0$  for each  $i$ , we can choose  $r$  so that  $|a_i^r(x)| \leq |g(x)|$  for all  $i$ . Then  $x \in X\left(\frac{T_r}{g}\right)$ .  $\square$

**Proposition 21.10.** *Let  $X = \mathrm{Spa}(A, A^+)$ , and let  $U$  be a rational subset of  $X$ . There exists a complete Huber pair  $(B, B^+)$  over  $(A, A^+)$  such that  $\mathrm{Spa}(B, B^+) \rightarrow \mathrm{Spa}(A, A^+)$  factors through  $U$ , and is universal for such maps. Moreover, the map  $\mathrm{Spa}(B, B^+) \rightarrow U$  is a homeomorphism.*

We construct  $(B, B^+)$  as follows. Choose a ring of definition  $A_0$  and an ideal of definition  $I$ .

Let the rational subset be defined by  $0 \neq |a_0| \geq |a_1|, \dots, |a_n|$ . Define a topology on  $A[1/a_0]$  by making  $A_0[a_1/a_0, \dots, a_n/a_0]$  a ring of definition with ideal of definition  $IA_0[a_1/a_0, \dots, a_n/a_0]$ . Then  $B$  is the completion of  $A[1/a_0]$ , and  $B^+$  is the completion of the integral closure of  $A^+[a_1/a_0, \dots, a_n/a_0]$  in  $A[1/s]$ .

**Definition 21.11.** For any Huber pair  $(A, A^+)$ , we define a presheaf of topological rings  $\mathcal{O}_X$  on  $X = \mathrm{Spa}(A, A^+)$  as follows. If  $U$  is rational, we define  $\mathcal{O}_X(U)$  to be

the ring  $B$  of Proposition 21.10. For general  $U$ , we define

$$\mathcal{O}_X(U) = \varinjlim_{W \subset U \text{ rational}} \mathcal{O}_X(W).$$

We define  $\mathcal{O}_X^+(U)$  similarly.

This presheaf  $\mathcal{O}_X$  is not a sheaf in general. If  $\mathcal{O}_X$  is a sheaf, then  $\mathcal{O}_X^+$  is also a sheaf. There are a few general classes of Huber rings that are known to be sheafy.

**Definition 21.12.** A Huber ring  $A$  is *Tate* if it has a topologically nilpotent unit.

**Definition 21.13.** A Huber ring  $A$  is *strongly Noetherian* if  $A\langle X_1, \dots, X_n \rangle$  is Noetherian for each  $n$ . Here

$$A\langle X_1, \dots, X_n \rangle = \left\{ \sum_{i \in \mathbb{N}^n} a_i X^i \mid \lim_{i \rightarrow \infty} a_i = 0 \right\}$$

**Definition 21.14.** A Huber ring  $A$  is *uniform* if  $A^\circ$  is bounded (equivalently,  $A^\circ$  is a ring of definition). The pair  $(A, A^+)$  is *stably uniform* if for every rational subset  $U \subset X = \text{Spa}(A, A^+)$ ,  $\mathcal{O}_X(U)$  is uniform.

**Lemma 21.15.** *The Huber pair  $(A, A^+)$  is sheafy if any of the following conditions is satisfied:*

- (1)  *$A$  is discrete. (schemes)*
- (2)  *$A$  is finitely generated over a Noetherian ring of definition. (Noetherian formal schemes, rigid spaces over discretely valued fields)*
- (3)  *$A$  is Tate and strongly Noetherian (rigid spaces)*
- (4)  *$A$  is Tate and  $(A, A^+)$  is stably uniform. (reduced rigid spaces, perfectoid spaces)*

**Lemma 21.16.** *The completion  $(A, A^+) \rightarrow (\hat{A}, \hat{A}^+)$  induces a canonical isomorphism  $\text{Spa}(\hat{A}, \hat{A}^+) \rightarrow \text{Spa}(A, A^+)$ .*

*Let  $X = \text{Spa}(A, A^+)$ . Then  $\mathcal{O}_X(X) \cong \hat{A}$ ,  $\mathcal{O}_X^+(X) \cong \hat{A}^+$ .*

**Definition 21.17.** A *v-ringed space* is a triple  $(X, \mathcal{O}_X, (|\cdot(x)|)_{x \in X})$  where  $X$  is a topological space,  $\mathcal{O}_X$  is a sheaf of topological rings on  $X$ , and for each  $x \in X$ ,  $|\cdot(x)|$  is an equivalence class of continuous valuations on  $\mathcal{O}_{X,x}$ . A morphism of v-ringed spaces is a morphism of ringed spaces, such that the maps on sections are continuous, and valuations are preserved.

**Definition 21.18.** An *affinoid adic space* is a v-ringed space that is isomorphic to some  $\text{Spa}(A, A^+)$ . (In particular, we require that the structure presheaf is a sheaf.)

An *adic space* is a v-ringed space that has an open covering by affinoid adic spaces.

*Remark 21.19.* Rational subsets of  $\text{Cont}(A)$  are generally not of the form  $\text{Cont}(B)$ , and rational subsets of  $\text{Spa}(A, A^\circ)$  are generally not of the form  $\text{Spa}(B, B^\circ)$ . This is why we do not try to build the theory just out of spaces of the form  $\text{Cont}(A)$ , or spaces of the form  $\text{Spa}(A, A^\circ)$ .

## 22. EXAMPLES OF ADIC SPACES, ANALYTIC ADIC SPACES

*Example 22.1.* The closed disc over  $\mathbb{C}_p$  is  $\text{Spa}(\mathbb{C}_p\langle X \rangle, \mathcal{O}_{\mathbb{C}_p}\langle X \rangle)$ . Similarly, the closed disc over  $\mathbb{Q}_p$  is  $\text{Spa}(\mathbb{Q}_p\langle X \rangle, \mathbb{Z}_p\langle X \rangle)$ .



*Example 22.2.*  $\text{Cont } \mathbb{C}_p \langle X \rangle = \text{Spa}(\mathbb{C}_p \langle X \rangle, A^+)$  where

$$A^+ = \left\{ \sum_{i=0}^{\infty} a_i X^i \mid a_0 \in \mathcal{O}_{\mathbb{C}_p}, a_i \in \mathfrak{m}_{\mathbb{C}_p} \text{ for } i > 0, \lim_{i \rightarrow \infty} a_i = 0 \right\}$$

*Example 22.3.* Let  $r \in \mathbb{Q}$ . Let  $A$  be the ring of power series  $\sum_{n=0}^{\infty} a_n T^n$  with  $a_n \in \mathbb{Q}_p$  and  $|a_n||p|^{nr} \rightarrow 0$  as  $n \rightarrow \infty$ . Let  $A^+$  be the subring consisting of series with  $|a_n||p|^{nr} \in \mathbb{Z}_p$  for all  $n$ . We give  $A$  the topology such that  $A^+$  is open in  $A$ , and  $A^+$  has the  $p$ -adic topology. Then  $\text{Spa}(A, A^+)$  is the closed disc of radius  $|p|^r$ .

*Example 22.4.* The open disc of radius 1 over  $\mathbb{Q}_p$  is the union of the closed discs of radius  $< 1$ .

*Example 22.5.* The affine line  $\mathbb{A}^1$  is the union of closed discs of radius  $p^r$ .

*Example 22.6.* The projective line  $\mathbb{P}^1$  is the union of two closed discs, glued along an annulus.

*Example 22.7.* Consider  $D = \text{Spa}(\mathbb{Z}_p[[T]], \mathbb{Z}_p[[T]])$ . It contains the open disc of radius 1 over  $\mathbb{Q}_p$ . It has exactly two additional points, corresponding to the trivial norm on the quotient  $\mathbb{F}_p$ , and the  $T$ -adic norm on  $\mathbb{F}_p[[T]]$ .

*Example 22.8.* The closed disc over  $\mathbb{Z}$  (with the discrete topology on  $\mathbb{Z}$ ) is  $\text{Spa}(\mathbb{Z}[T], \mathbb{Z}[T])$ . It represents the functor  $X \mapsto \mathcal{O}_X^+(X)$ .

*Example 22.9.* The affine line over  $\mathbb{Z}$  is  $\text{Spa}(\mathbb{Z}[T], \mathbb{Z})$ . It represents the functor  $X \mapsto \mathcal{O}_X(X)$ . Unlike the affine line over  $\mathbb{Q}_p$ , it is affinoid.

*Example 22.10.* Let  $S$  be a profinite set, and let  $A$  be the space of continuous (i.e. locally constant) functions  $S \rightarrow \mathbb{Z}$ , with the discrete topology. Then  $\text{Spa}(A, A)$  represents the functor  $X \mapsto \text{Hom}(|X|, S)$ .

**Definition 22.11.** Let  $X$  be an adic space. A point  $x \in X$  is *analytic* if it has a neighborhood of the form  $\text{Spa}(A, A^+)$ , with  $A$  Tate. We say  $X$  is *analytic* if all of its points are analytic.

**Proposition 22.12.** *Let  $(A, A^+)$  be a sheafy Huber pair. Then a point  $x \in \text{Spa}(A, A^+)$  is analytic iff the kernel of  $|\cdot|_x$  is not open.*

*Proof.* Let  $I = (a_0, \dots, a_n)$  be an ideal of definition of a ring of definition of  $A$ . Suppose that the kernel of  $|\cdot|_x$  is not open. Then  $I \not\subset \ker |\cdot|_x$ . So the  $|a_i|_x$  are not all zero. WLOG assume  $|a_0|_x \geq |a_1|_x, \dots, |a_n|_x$ . Then, on the rational subset defined by  $|a_0| \geq |a_1|, \dots, |a_n|$ ,  $a_0$  is a topologically nilpotent unit. In particular, this rational subset is the adic spectrum of a Tate ring.

Conversely, suppose that the kernel of  $|\cdot|_x$  is open. Let  $\text{Spa}(B, B^+)$  be an affinoid subspace of  $\text{Spa}(A, A^+)$  containing  $x$ . We claim that  $\text{Spa}(B, B^+)$  cannot be Tate. Let  $\text{Spa}(C, C^+)$  be a rational subset of  $\text{Spa}(A, A^+)$  containing  $x$  and contained in  $\text{Spa}(B, B^+)$ . Since the image of a topologically nilpotent unit in  $\text{Spa}(B, B^+)$  would be a topologically nilpotent unit in  $\text{Spa}(C, C^+)$ , it suffices to prove that  $C$  is not Tate.

Since the kernel of  $|\cdot|_x$  is open, it contains some power of  $I$ . Since  $|\cdot|_x$  is multiplicative, the kernel must contain  $I$ . The generators of  $I$  also generate an ideal of definition of  $C$ . Therefore, when  $|\cdot|_x$  is extended to  $C$ , its kernel is still open. Hence  $C$  cannot have a topologically nilpotent unit.  $\square$

*Example 22.13.* The adic space  $\mathrm{Spa}(\mathbb{Z}_p, \mathbb{Z}_p)$  consists of two points: a closed point  $s$  given by

$$|a|_s = \begin{cases} 1, & a \in \mathbb{Z}_p^\times \\ 0, & a \in p\mathbb{Z}_p \end{cases},$$

and a generic point  $\eta$  given by the  $p$ -adic norm. The kernel of  $|\cdot|_s$  is  $p\mathbb{Z}_p$ , which is open, so  $s$  is not analytic. The kernel of  $|\cdot|_\eta$  is 0, which is not open, so  $\eta$  is analytic.

Analytic adic spaces are somewhat nicer than general adic spaces.

*Remark 22.14.* There exist complete Huber pairs  $(A, A^+)$  such that  $\mathrm{Spa}(A, A^+)$  is analytic, but  $A$  is not Tate. See [Ked, §1].

### 23. RIGID ANALYTIC SPACES, PERFECTOID RINGS

Let  $K$  be a nonarchimedean field.

**Definition 23.1.** A *Tate algebra* over  $K$  is a Huber ring of the form  $K\langle T_1, \dots, T_n \rangle$  for some  $n$ .

**Proposition 23.2.** *Let  $A$  be a Tate algebra over  $K$ .*

- (1)  *$A$  is noetherian.*
- (2) *Every ideal of  $A$  is closed.*

*Proof.* See [Bos14, Proposition 2.2.14 and Corollary 2.3.8]. □

**Definition 23.3.** An *affinoid algebra* over  $K$  is a quotient  $A/I$ , where  $A$  is a Tate algebra and  $I$  is an ideal of  $A$ .

**Definition 23.4.** A *rigid analytic space* is an adic space over  $K$  that can be covered by open subspaces of the form  $\mathrm{Spa}(A, A^\circ)$ , where  $A$  is an affinoid algebra over  $K$ .

*Remark 23.5.* There is an older definition of rigid analytic spaces, due to Tate, that does not use the framework of adic spaces. Tate's category of quasiseparated rigid analytic spaces is equivalent to our category of quasiseparated rigid analytic spaces.

**Proposition 23.6.** *The category of rigid analytic spaces over  $K$  admits finite fiber products.*

**Definition 23.7.** A morphism  $f: X \rightarrow Y$  of rigid analytic spaces over  $K$  is *separated* if the image of the diagonal morphism  $\Delta_f: X \rightarrow X \times_Y X$  is closed.

**Definition 23.8.** A morphism  $f: X \rightarrow Y$  of rigid analytic spaces over  $K$  is *universally closed* if for every morphism of rigid spaces  $Z \rightarrow Y$  over  $K$ , the projection  $X \times_Y Z \rightarrow Z$  is closed.

**Definition 23.9.** A morphism  $f: X \rightarrow Y$  of rigid analytic spaces over  $K$  is *proper* if it is separated and universally closed.

**Definition 23.10.** A morphism  $f: X \rightarrow Y$  of rigid analytic spaces over  $K$  is *unramified* (resp. *smooth*, *étale*) if for any affinoid  $K$ -algebra  $A$  and any ideal  $I$  of  $A$  with  $I^2 = 0$ , and any morphism  $\mathrm{Spa}(A, A^\circ) \rightarrow Y$ , the natural map

$$\mathrm{Hom}_Y(\mathrm{Spa}(A, A^\circ), X) \rightarrow \mathrm{Hom}_Y(\mathrm{Spa}(A/I, (A/I)^\circ), X)$$

is injective (resp. surjective, bijective).

**Proposition 23.11.** *Let  $A$  be an affinoid algebra, and let  $f: X \rightarrow \mathrm{Spa}(A, A^\circ)$  be a morphism of rigid analytic spaces. Then  $f$  is smooth if and only if, for every  $x \in X$ , there exists an open neighborhood  $U$  of  $x$  and a commutative diagram*

$$\begin{array}{ccc} & \mathbb{A}_A^n & \\ g \nearrow & & \searrow \mathrm{pr} \\ U & \xrightarrow{f|_U} & \mathrm{Spa}(A, A^\circ) \end{array}$$

such that  $g$  is étale.

*Proof.* See [Hub96, Corollary 1.6.10].  $\square$

Rigid analytic spaces are similar to complex manifolds in some regard, but they are not sufficient for a satisfactory theory of  $p$ -adic geometry. A complex manifold can be built from unit balls, which are contractible. But the  $p$ -adic unit disc is not simply connected: the map  $z \mapsto z - z^p$  is a  $p$ -fold cover of the unit disc by itself. To get a better theory, we need to consider perfectoid spaces.

**Definition 23.12.** A complete Tate ring  $R$  is *perfectoid* if  $R$  is uniform and there exists a topologically nilpotent unit  $\varpi$  such that  $\varpi^p \mid p$  in  $R^\circ$ , and the Frobenius map

$$\Phi: R^\circ/\varpi \rightarrow R^\circ/\varpi^p$$

is an isomorphism.

**Lemma 23.13.** *A complete Tate ring  $R$  is perfectoid iff  $R$  is uniform, there exists a topologically nilpotent unit  $\varpi$  such that  $\varpi^p \mid p$  in  $R^\circ$ , and the Frobenius map  $\Phi: R^\circ/p \rightarrow R^\circ/p$  is surjective.*

*Proof.* We first show that  $\Phi: R^\circ/\varpi \rightarrow R^\circ/\varpi^p$  is always injective. Suppose that  $x \in R^\circ$  satisfies  $x^p \in \varpi^p R^\circ$ . Then  $(x/\varpi)^p \in R^\circ$ , so  $x/\varpi \in R^\circ$  as well.

It is clear that if  $\Phi$  is surjective mod  $p$ , then it is also surjective mod  $\varpi^p$ . Conversely, if it is surjective mod  $\varpi^p$ , then by successive approximation we can write  $x \in R^\circ$  as  $x_0 + \varpi^p x_1 + \varpi^{2p} x_2 + \cdots$ ; then  $(x_0 + \varpi x_1 + \varpi^2 x_2 + \cdots)^p$  is congruent to  $x$  mod  $p$ .  $\square$

*Example 23.14.* The following rings are perfectoid:

- (1)  $K = \widehat{\mathbb{Q}_p(p^{1/p^\infty})}$ , since  $K^\circ/p \cong \mathbb{F}_p[T^{1/p^\infty}]/(T)$  via  $p \mapsto T^{1/p^n} \bmod T$ .
- (2)  $K = \mathbb{Q}_p^{\mathrm{cyc}}$ , the completion of  $\mathbb{Q}_p(\mu_{p^\infty})$ , since  $K^\circ/p \cong \mathbb{F}_p[T^{1/p^\infty}]/(T^{1-1/p})$  via  $\zeta_{p^n} \mapsto 1 + T^{1/p^n} \bmod T^{1-1/p}$ .
- (3)  $K = \mathbb{F}_p((T^{1/p^\infty}))$ , the completion of  $\mathbb{F}_p((T))(T^{1/p^\infty})$ .
- (4)  $\mathbb{Q}_p^{\mathrm{cyc}} \langle T^{1/p^\infty} \rangle$ , which is constructed by taking the  $p$ -adic completion of  $\mathbb{Z}_p[T^{1/p^\infty}]$  and inverting  $p$ .

**Proposition 23.15.** *Let  $R$  be a Tate ring, and suppose that  $p = 0$  in  $R$ . Then the following are equivalent:*

- (1)  $R$  is perfectoid
- (2)  $R$  is perfect (Frobenius is an isomorphism) and complete.

*Proof.* Suppose  $R$  is perfectoid. Then  $R$  is uniform and since  $R^\circ = R^\circ/p$ ,  $\Phi: R^\circ \rightarrow R^\circ$  is surjective. Then  $\Phi: R \rightarrow R$  is surjective. Since  $R$  is uniform,  $\Phi: R \rightarrow R$  is injective.

Now assume that  $R$  is a perfect complete Tate ring. Let  $R_0$  be a ring of definition. By the Banach open mapping theorem,  $\Phi(R_0)$  is open in  $R_0$ . (Note that we can make  $\Phi$  linear by an appropriate choice of an  $R$ -module structure on the target.) So we can find a pseudouniformizer  $\varpi$  so that  $\varpi R_0 \subset \Phi(R_0)$ . Then  $\Phi^{-1}(R_0) \subset \varpi^{-1/p} R_0$ ,  $\Phi^{-2}(R_0) \subset \varpi^{-1/p-1/p^2} R_0$ , etc., so  $R'_0 := \bigcup_n \Phi^{-n} R_0$  is bounded; hence it is a ring of definition. Moreover,  $R$  is bounded since  $R^\circ \subseteq \varpi^{-1} R^{\circ\circ} \subseteq \varpi^{-1} R'_0$ .  $\square$

**Lemma 23.16.** *Let  $(R, R^+)$  be a perfectoid Huber pair, and let  $X = \mathrm{Spa}(R, R^+)$ . For any rational subset  $U$ ,  $\mathcal{O}_X(U)$  is again perfectoid. Therefore  $(R, R^+)$  is stably uniform, hence sheafy.*

**Definition 23.17.** Let  $R$  be a perfectoid Tate ring. The *tilt* of  $R$  is

$$R^\flat := \varprojlim_{x \mapsto x^p} R.$$

We define a ring structure on  $R$  as follows. Multiplication is defined in the obvious way. Addition is defined by

$$(x^{(0)}, x^{(1)}, \dots) + (y^{(0)}, y^{(1)}, \dots) = (z^{(0)}, z^{(1)}, \dots)$$

where

$$z^{(i)} = \lim_{n \rightarrow \infty} (x^{(i+n)} + y^{(i+n)})^{p^n}.$$

**Lemma 23.18.** *The above limit exists, and  $R^\flat$  is a topological  $\mathbb{F}_p$ -algebra that is a complete Tate ring. The subring  $R^{\flat\circ}$  of power-bounded elements is given by*

$$R^{\flat\circ} = \varprojlim_{x \mapsto x^p} R^\circ \cong \varprojlim_{\Phi} R^\circ/p \cong \varprojlim_{\Phi} R^\circ/\varpi$$

where  $\varpi$  is a pseudouniformizer that divides  $p$  in  $R^\circ$ .

Furthermore, there exists a pseudouniformizer  $\varpi \in R^\circ$  with  $\varpi \mid p$  in  $R^\circ$  admitting a compatible sequence of  $p$ th power roots  $\varpi^{1/p^n}$ . Then  $\varpi^\flat = (\varpi, \varpi^{1/p}, \dots)$  is a pseudouniformizer of  $R^\flat$ , and  $R^\flat = R^{\flat\circ}[1/\varpi^\flat]$ .

*Proof.* The key step is to check that  $\varprojlim_{x \mapsto x^p} R^\circ \rightarrow \varprojlim_{\Phi} R^\circ/p$  is a bijection. Suppose we have a sequence  $(\bar{x}_0, \bar{x}_1, \dots) \in \varprojlim_{\Phi} R^\circ/p$ , choose any lift  $(y_0, y_1, \dots) \in (R^\circ)^\mathbb{N}$ , and let  $x_i = \lim_{n \rightarrow \infty} y_{i+n}^{p^n}$ . Note that if  $y \equiv y' \pmod{p}$ , then  $y^{p^n} \equiv y'^{p^n} \pmod{p^{n+1}}$ . So the  $x_i$  do not depend on the choice of  $y_i$ , and  $x_{i+1}^p = x_i$  for each  $i$ .

Let  $\varpi_0$  be a pseudouniformizer of  $R$  satisfying  $\varpi_0^p \mid p$  in  $R^\circ$ . Since  $R$  is perfectoid, we can find  $(\bar{x}_0, \bar{x}_1, \dots) \in \varprojlim_{\Phi} R^\circ/p$  with  $\bar{x}_0 = \varpi_0 \bmod \varpi_0^p$ . Let  $\varpi^\flat$  be the corresponding element  $R^{\flat\circ}$ ; it is a pseudouniformizer and  $\varpi := \varpi^{\flat\sharp}$  is the desired pseudouniformizer of  $R$ .  $\square$

**Lemma 23.19.** *The set of rings of integral elements  $R^+ \subset R^\circ$  is in bijection with the set of rings of integral elements  $R^{\flat+} \subset R^{\flat\circ}$  via  $R^{\flat+} = \varprojlim_{x \mapsto x^p} R^+$ . Also,  $R^{\flat+}/\varpi^\flat = R^+/\varpi$ .*

**Theorem 23.20.** *Let  $(R, R^+)$  be a perfectoid Huber pair, with tilt  $(R^\flat, R^{\flat+})$ . The map  $\sharp: R^\flat \rightarrow R$  induces a homeomorphism  $X := \mathrm{Spa}(R, R^+) \cong X^\flat := \mathrm{Spa}(R^\flat, R^{\flat+})$ . This homeomorphism preserves rational subsets. For any rational subset  $U \subset X$  with image  $U^\flat \subset X^\flat$ ,  $\mathcal{O}_X(U)$  is perfectoid with tilt  $\mathcal{O}_{X^\flat}(U^\flat)$ .*

**Theorem 23.21.** *Let  $R$  be a perfectoid ring with tilt  $R^\flat$ . Then there is an equivalence of categories between perfectoid  $R$  algebras and perfectoid  $R^\flat$ -algebras, via  $S \mapsto S^\flat$ .*

**Definition 23.22.** Let  $(R, R^+)$  be a perfectoid Huber pair in characteristic  $p$ . An element  $\xi \in W(R^+)$  is *primitive of degree one* if  $\xi = p + [\varpi]\alpha$ , where  $\varpi \in R^+$  is a pseudouniformizer.

**Lemma 23.23.** *Let  $(R, R^+)$  be a perfectoid Huber pair in characteristic  $p$ . Let  $(R^\sharp, R^{\sharp+})$  be an untilt of  $(R, R^+)$ , i.e. a perfectoid Tate ring  $R^\sharp$  along with an isomorphism  $R^{\sharp\flat} \xrightarrow{\sim} R$  identifying  $R^{\sharp\flat+}$  with  $R^+$ .*

(1) *There is a canonical surjective ring homomorphism*

$$\begin{aligned} \theta: W(R^+) &\rightarrow R^{\sharp+} \\ \sum_{n=0}^{\infty} [r_n]p^n &\mapsto \sum_{n \geq 0} r_n^\sharp p^n. \end{aligned}$$

(2) *The kernel of  $\theta$  is generated by an element  $\xi$  that is primitive of degree one.*

**Lemma 23.24.** *A primitive degree one element of  $W(R^+)$  is not a zero divisor.*

**Theorem 23.25.** *There is an equivalence of categories between:*

- (1) *Perfectoid Tate-Huber pairs  $(S, S^+)$*
- (2) *Triples  $(R, R^+, J)$ , where  $(R, R^+)$  is a perfectoid Tate-Huber pair of characteristic  $p$  and  $J \subset W(R^+)$  is primitive of degree 1. The functors are*

$$\begin{aligned} (S, S^+) &\mapsto (S^\flat, S^{\flat+}, \ker \theta) \\ (R, R^+, J) &\mapsto (W(R^+) [[\varpi]^{-1}] / J, W(R^+) / J) \end{aligned}$$

**Definition 23.26.** Let  $R$  be a complete Tate  $\mathbb{Z}_p$ -algebra. We say that  $R$  is *sousperfectoid* if there exists a perfectoid Tate ring  $\hat{R}$  and an injection  $R \hookrightarrow \hat{R}$  that splits as a map of topological  $R$ -modules.

*Example 23.27.* (1) Any perfectoid ring is sousperfectoid.

(2) The ring  $R = \mathbb{Q}_p \langle T \rangle$  is sousperfectoid, by taking  $\hat{R} = \mathbb{Q}_p^{\text{cyc}} \langle T^{1/p^\infty} \rangle$ .

(3) Similarly,  $\mathbb{Q}_p \langle T^{1/p^\infty} \rangle$  is sousperfectoid.

**Proposition 23.28.** *Let  $R$  be a complete Tate  $\mathbb{Z}_p$ -algebra with a ring of integral elements  $R^+ \subset R$ , and assume that  $R$  is sousperfectoid.*

- (1) *If  $U \subset X = \text{Spa}(R, R^+)$  is a rational subset, then  $\mathcal{O}_X(U)$  is sousperfectoid.*
- (2) *If  $S$  is a finite étale  $R$ -algebra, then  $S$  is sousperfectoid.*
- (3) *For all  $n \geq 0$ , the ring  $R \langle T_1, \dots, T_n \rangle$  is sousperfectoid.*

**Proposition 23.29.** *Let  $(R, R^+)$  be a Tate-Huber pair such that  $(R, R^+)$  is sousperfectoid. Then  $(R, R^+)$  is stably uniform, hence sheafy.*

## 24. PERFECTOID SPACES

**Definition 24.1.** An *affinoid perfectoid space* is an adic space of the form  $\text{Spa}(R, R^+)$  with  $R$  perfectoid. A *perfectoid space* is an adic space that can be covered by affinoid perfectoid spaces.

*Remark 24.2.* It is unknown whether an adic space that is both affinoid and perfectoid is necessarily affinoid perfectoid.

The tilting construction can be glued, so for any perfectoid space  $X$ , we can define its tilt  $X^\flat$ .

**Theorem 24.3.** *For any perfectoid space  $X$ , the functor  $Y \mapsto Y^\flat$  induces an equivalence between the categories of perfectoid spaces over  $X$  and  $X^\flat$ .*

Recall the following result that was proved earlier in the course.

**Theorem 24.4** (Tate, 1967). *Let  $K = \mathbb{Q}_p^{\text{cyc}}$ . If  $L/K$  is a finite extension, then the image of  $\text{tr}: \mathcal{O}_L \rightarrow \mathcal{O}_K$  contains  $\mathfrak{m}_K$ .*

Tate's result says that any finite extension of  $\mathbb{Q}_p^{\text{cyc}}$  is almost unramified.

**Definition 24.5.** Let  $(R, R^+)$  be a perfectoid Tate-Huber pair. We say that an  $R^+$ -module  $M$  is *almost zero* if  $R^{\circ\circ}$  annihilates  $M$ .

*Example 24.6.* If  $K$  is a perfectoid field, then an  $\mathcal{O}_K$ -module is almost zero iff it is annihilated by  $\mathfrak{m}_K$ . One can show that an almost zero  $\mathcal{O}_K$ -module is a direct sum of copies of the residue field  $\mathcal{O}_K/\mathfrak{m}_K$ .

An extension of almost zero modules is again almost zero.

*Remark 24.7.* Here it is important that  $R$  is perfectoid. Note that for  $R = \mathbb{Q}_p$ ,  $R^+ = \mathbb{Z}_p$ , then  $\mathbb{Z}/p\mathbb{Z}$  is annihilated by  $R^{\circ\circ} = p\mathbb{Z}_p$ , but  $\mathbb{Z}/p^2\mathbb{Z}$  is not.

In fact, the subcategory of almost zero  $R^+$ -modules is a thick Serre subcategory, so the following definition makes sense.

**Definition 24.8.** The category of *almost  $R^\circ$ -modules*, denoted  $R^{\circ a}\text{-mod}$  is the quotient of the category of  $R^\circ$ -modules by the subcategory of almost zero modules.

If  $M$  is an  $R^\circ$ -module, then we write  $M^a$  for the corresponding  $R^{\circ a}$ -modules. One can similarly define the category of almost  $R^+$ -modules, but it is equivalent to the category of almost  $R^\circ$ -modules.

*Example 24.9.* If  $K$  is a perfectoid field, then  $\mathfrak{m}_K^a \rightarrow \mathcal{O}_K^a$  is an isomorphism of  $\mathcal{O}_K^a$ -modules.

**Theorem 24.10.** *Let  $(R, R^+)$  be a perfectoid Tate-Huber pair, and let  $X = \text{Spa}(R, R^+)$ . Then  $H^i(X, \mathcal{O}_X^+)$  is almost zero for  $i > 0$ , and  $H^0(X, \mathcal{O}_X^+) = R^+$ .*

**Theorem 24.11.** *Let  $R$  be a perfectoid Tate ring.*

- (1) *For any finite étale  $R$ -algebra  $S$ ,  $S$  is perfectoid.*
- (2) *Tilting induces an equivalence between finite étale  $R$ -algebras and finite étale  $R^\flat$  algebras.*
- (3) *(almost purity) For any finite étale  $R$ -algebra  $S$ ,  $S^\circ$  is almost finite étale over  $R^\circ$ .*

We would like to define a notion of tilting for spaces that are not perfectoid. If we just tried to tilt the rings, we would lose too much information. If  $K$  is a finite extension of  $\mathbb{Q}_p$ , then  $\varprojlim_{x \rightarrow x^p} K$  is just the residue field of  $K$ . The solution is to find a perfectoid cover, tilt the cover, and then take a quotient. So for example, the tilt of  $\text{Spa } \mathbb{Q}_p$  should be the quotient of  $(\text{Spa } \mathbb{Q}_p^{\text{cyc}})^\flat / \mathbb{Z}_p^\times$ . The quotient will live in the category of diamonds, which are similar to algebraic spaces.

**Definition 24.12.**

- (1) A morphism  $f: X \rightarrow Y$  of perfectoid spaces is *finite étale* if for all  $\mathrm{Spa}(B, B^+) \subset Y$  open, the pullback  $X \times_Y \mathrm{Spa}(B, B^+)$  is  $\mathrm{Spa}(A, A^+)$ , where  $A$  is a finite étale  $B$ -algebra, and  $A^+$  is the integral closure of the image of  $B^+$  in  $A$ .
- (2) A morphism  $f: X \rightarrow Y$  is étale if for all  $x \in X$  there exists an open  $U \ni x$  and  $V \supset f(U)$  such that there is a diagram

$$\begin{array}{ccc} U & \xrightarrow{\text{open}} & W \\ & \searrow f|_U & \swarrow \text{finite étale} \\ & & V \end{array}.$$

- (3) An *étale cover* is a jointly surjective family of étale maps.

**Proposition 24.13.** *One can construct the étale site  $X_{\text{ét}}$  from the above definitions. The site  $X_{\text{ét}}^b$  is naturally equivalent to  $X_{\text{ét}}$ . If  $X$  is affinoid perfectoid, then  $H^i(X_{\text{ét}}, \mathcal{O}_X) = 0$  and  $H^i(X_{\text{ét}}, \mathcal{O}_X^+) = 0$  for  $i > 0$ , and  $H^i(X_{\text{ét}}, \mathbb{F}_p) = 0$  for  $i > 1$ .*

## 25. PRO-ÉTALE SITE AND DIAMONDS

**Proposition 25.1.** *Let  $X_i = \mathrm{Spa}(R_i, R_i^+)$  be a directed system of affinoid perfectoid spaces. Then the limit  $\varprojlim_i X_i$  exists in the category of perfectoid spaces, and is isomorphic to  $\mathrm{Spa}(R, R^+)$ , where  $R^+$  is the completion of  $\varprojlim_i R_i^+$  for the  $\varpi$ -adic topology and  $R = R^+[1/\varpi]$ . Here  $\varpi$  is the image of an arbitrarily chosen pseudouniformizer of some  $R_i$ .*

- Remark 25.2.* (1) The  $\varpi$ -adic topology is generally weaker than the direct limit topology.
- (2) The limit generally does not exist in the category of adic spaces. Given maps  $(R_i, R_i^+) \rightarrow (S, S^+)$ , the induced map  $R \rightarrow S$  is continuous for the topology defined above if  $S$  is uniform, but not for more general  $S$ . To give a concrete example, the limit  $\varprojlim_n \mathrm{Spa}(\mathbb{Q}_p \langle p^n T \rangle, \mathbb{Z}_p \langle p^n T \rangle)$  in the category of sousperfectoid spaces is a single point  $\mathrm{Spa}(\mathbb{Q}_p, \mathbb{Z}_p)$ , but this is not the limit in the category of adic spaces, as can be seen by looking at  $\mathbb{Q}_p[\epsilon]/(\epsilon^2)$ -valued points.

**Definition 25.3.** Let  $f: Y \rightarrow X$  be a map of perfectoid spaces. We say that  $f$  is *affinoid pro-étale* if  $Y$  and  $X$  are affinoid,  $f$  can be written as a cofiltered inverse limit of étale maps  $Y_i \rightarrow X$ , with  $Y_i$  affinoid perfectoid. We say that  $f$  is *pro-étale* if for all  $y \in Y$ , there exist open sets  $V \subset Y$ ,  $U \subset X$  with  $y \in V$ ,  $f(V) \subset U$  and  $f|_V: V \rightarrow U$  is affinoid pro-étale.

*Remark 25.4.* Scholze's  $p$ -adic Hodge theory paper uses a different definition of pro-étale morphisms.

*Example 25.5.* Let  $X$  be a perfectoid space, and let  $S$  be a profinite set. Write  $S = \varprojlim_i S_i$ , where  $S_i$  are finite sets. Then we can define  $X \times \underline{S} := \varprojlim_i X \times S_i$ . The projection  $X \times \underline{S} \rightarrow X$  is pro-étale. More generally, this is true if  $X \times \underline{S} \rightarrow X$  is pro-étale if  $S$  is locally profinite.

*Example 25.6.* Let  $X$  be a perfectoid space, and let  $x = \mathrm{Spa}(K, \mathcal{O}_K)$  be a point of  $X$ . Then  $x = \varprojlim_{U \ni x} U$ . Since each  $U \hookrightarrow X$  is étale,  $x \rightarrow X$  is pro-étale. In particular, this means that pro-étale morphisms are not necessarily open.

**Definition 25.7.** Consider the following categories:

- $\text{Perfd}$ , the category of perfectoid spaces.
- $\text{Perf}$ , the category of perfectoid spaces in characteristic  $p$ .
- $X_{\text{proét}}$ , the category of perfectoid spaces proétale over a perfectoid space  $X$ .

We give each of these categories the structure of a site by saying that a collection of morphisms  $\{f_i: Y_i \rightarrow Y\}$  is a covering if the  $f_i$  are pro-étale, and for all quasi-compact open  $U \subset Y$ , there exists a finite index set  $I_U$  and quasicompact open subsets  $U_i \subset Y_i$  for  $i \in I_U$ , such that  $U = \bigcup_{i \in I_U} f_i(U_i)$ .

**Proposition 25.8.** (1) *The functors  $X \mapsto \mathcal{O}_X(X)$ ,  $X \mapsto \mathcal{O}_X^+(X)$  are sheaves on the pro-étale site of  $\text{Perfd}$ . If  $X$  is affinoid, then  $H^i(X_{\text{proét}}, \mathcal{O}_X) = 0$  and  $H^i(X_{\text{proét}}, \mathcal{O}_X^+)^a = 0$  for  $i > 0$ .*  
 (2) *For any perfectoid space  $X$ , the functor  $h_X = \text{Hom}(-, X)$  is a sheaf on the pro-étale site of  $\text{Perfd}$ .*

**Definition 25.9.** A *diamond* is a pro-étale sheaf  $D$  on  $\text{Perf}$  such that one can write  $D = X/R$ , where  $X$  is a perfectoid space and  $R \subset X \times X$  is a perfectoid space with  $s, t: R \rightarrow X$  pro-étale.

**Definition 25.10.** Let  $Y$  be a perfectoid space in characteristic  $p$ . An *untilt* of  $Y$  is a pair  $(Y^\sharp, (Y^\sharp)^\flat \xrightarrow{\sim} Y)$ , where  $Y^\sharp$  is a perfectoid space.

*Example 25.11.* Let  $\text{Spd } \mathbb{Q}_p$  be the functor that sends  $X \in \text{Perf}$  to the set of isomorphism classes of characteristic zero untilts of  $X$ . We claim that  $\text{Spd } \mathbb{Q}_p$  is a diamond.

Write  $\text{Spd } \mathbb{Q}_p^{\text{cyc}}$  for  $\text{Spa}(\mathbb{Q}_p^{\text{cyc}})^\flat$ . The tilting equivalence gives us a morphism  $\text{Spd } \mathbb{Q}_p^{\text{cyc}} \rightarrow \text{Spd } \mathbb{Q}_p$ . This is a surjection of pro-étale sheaves since every characteristic zero perfectoid space has a pro-étale cover that admits a map to  $\text{Spa } \mathbb{Q}_p^{\text{cyc}}$ . So  $\text{Spd } \mathbb{Q}_p$  is the quotient of  $\text{Spd } \mathbb{Q}_p^{\text{cyc}}$  by an equivalence relation  $R \subset \text{Spd } \mathbb{Q}_p^{\text{cyc}} \times \text{Spd } \mathbb{Q}_p^{\text{cyc}}$ . We claim that in fact  $R$  can be identified with  $\text{Spd } \mathbb{Q}_p^{\text{cyc}} \times \underline{\mathbb{Z}_p^\times}$ , where  $\underline{\mathbb{Z}_p^\times}$  is a constant group sheaf. This is equivalent to checking that  $\text{Spa } \mathbb{Q}_p^{\text{cyc}} \times \underline{\mathbb{Z}_p^\times} \rightarrow \text{Spa } \mathbb{Q}_p^{\text{cyc}} \times \text{Spa } \mathbb{Q}_p^{\text{cyc}}$  is an isomorphism of presheaves on  $\text{Perfd}$ . By Galois theory,  $\text{Spa } \mathbb{Q}_p(\mu_{p^n}) \times (\mathbb{Z}/p^n\mathbb{Z})^\times \xrightarrow{\sim} \text{Spa } \mathbb{Q}_p(\mu_{p^n}) \times \text{Spa } \mathbb{Q}_p(\mu_{p^n})$ , and then taking the inverse limit proves the claim. Then  $R \rightrightarrows \text{Spd } \mathbb{Q}_p^{\text{cyc}}$  are proétale morphisms, so  $\text{Spd } \mathbb{Q}_p$  is a diamond.

## 26. MORE EXAMPLES OF DIAMONDS

**Definition 26.1.** Let  $X$  be an analytic adic space over  $\text{Spa}(\mathbb{Z}_p, \mathbb{Z}_p)$ . Define a presheaf  $X^\diamond$  on  $\text{Perf}$  as follows. For  $Y \in \text{Perf}$ , let  $X^\diamond(Y)$  be the set of isomorphism classes of triples  $(Y^\sharp, (Y^\sharp)^\flat \xrightarrow{\sim} Y, Y^\sharp \rightarrow X)$ , where  $Y^\sharp$  is a perfectoid space.

**Theorem 26.2.**  $X^\diamond$  is a diamond.

A key lemma in the proof is the following.

**Lemma 26.3.** *Let  $R$  be a Tate  $\mathbb{Z}_p$ -algebra. Let  $R_\infty := \varinjlim_i R_i$  be a filtered direct limit of algebras  $R_i$  finite étale over  $R$ , that admits no nonsplit finite étale covers. Give  $R_\infty$  the topology making  $\varinjlim_i R_i^\circ$  open and bounded. Let  $\tilde{R}$  be the completion of  $R_\infty$ . Then  $\tilde{R}$  is perfectoid.*



*Proof.* First, find a pseudouniformizer  $\varpi \in \tilde{R}$  satisfying  $\varpi^p \mid p$ . Let  $\varpi_0$  be a pseudouniformizer of some  $R_i$ . For some integer  $N$ ,  $\varpi_0 \mid p^N$  in  $\tilde{R}^\circ$ . We can find  $\varpi \in \tilde{R}^\circ$  satisfying  $\varpi^{p^N} - \varpi_0 \varpi = \varpi_0$ ; then  $\varpi^{p^N} \mid \varpi_0$  in  $R^\circ$ .

It remains to check that  $\tilde{R}^\circ/\varpi \rightarrow \tilde{R}^\circ/\varpi^p$  is surjective. Observe that for any  $f \in R_i^\circ$ , the equation  $x^p - \varpi^p x - f = 0$  has a solution in  $\tilde{R}$ .  $\square$

**Definition 26.4.** A ring  $A$  is *seminormal* if the map  $A \rightarrow \{(x, y) \in A^2 \mid y^2 = x^3\}$  given by  $t \mapsto (t^2, t^3)$  is a bijection.

A rigid analytic space is *seminormal* if it is locally of the form  $\mathrm{Spa}(A, A^+)$  with  $A$  seminormal.

**Proposition 26.5.** For any nonarchimedean field  $K$  over  $\mathbb{Q}_p$ , the functor

$$\begin{aligned} \{\text{seminormal rigid analytic spaces}/K\} &\rightarrow \{\text{diamonds}/\mathrm{Spd} K\} \\ X &\mapsto X^\diamond \end{aligned}$$

is fully faithful.

*Example 26.6.* Here are some examples of morphisms of rigid spaces that induce isomorphisms of diamonds:

- (1) Frobenius, if  $K$  has characteristic  $p$ .
- (2)  $X^{\mathrm{red}} \rightarrow X$  for any rigid space  $X$ , since perfectoid rings do not have nilpotents.
- (3)  $\mathbb{A}^1 \rightarrow Y$ , where  $Y$  is the cuspidal cubic. To prove this, one uses the fact that a qcqs map of diamonds is an isomorphism iff it induces a bijection on  $(K, K^+)$  points for every  $(K, K^+)$ .

**Definition 26.7.** Let  $D$  be a diamond, and choose a presentation  $D = X/R$ . The underlying topological space of  $D$  is the quotient  $|D| = |X|/|R|$ .

It can be shown that this definition is independent of the presentation.

**Proposition 26.8.** If  $X$  is an analytic adic space over  $\mathbb{Z}_p$ , then there is a natural homeomorphism  $|X| \cong |X^\diamond|$ .

*Example 26.9.* Let  $\mathrm{Spd} \mathbb{Z}_p$  be the presheaf that sends  $Y \in \mathrm{Perf}$  to the set of isomorphism classes of pairs  $(Y^\sharp, (Y^\sharp)^\flat \xrightarrow{\sim} Y)$ . Then  $\mathrm{Spd} \mathbb{Z}_p$  is not a diamond (because  $\mathbb{Z}_p$  is not an analytic Huber ring). But it is a sheaf. Moreover, it turns out that  $\mathrm{Spd} \mathbb{Z}_p$  is an “absolute diamond”: its product with any diamond is again a diamond. In a later lecture, we construct, for any  $S \in \mathrm{Perf}$ , an analytic adic space whose diamond is  $\mathrm{Spd} \mathbb{Z}_p \times S$ .

*Example 26.10.* Let  $X_i$  be a directed system of rigid spaces with qcqs transition maps. Then  $\varprojlim_i X_i^\diamond$  is a diamond.

*Example 26.11.* For each  $n$ , define the sheaf  $B_{\mathrm{dR}}^+/\mathrm{Fil}^n$  on  $\mathrm{Perf}$  as follows. Let  $(R, R^+)$  be a perfectoid Huber-Tate pair in characteristic  $p$ . Then  $B_{\mathrm{dR}}^+/\mathrm{Fil}^n(R, R^+)$  will be the set of isomorphism classes of pairs of an untilt  $R^\sharp$  of  $R$  and an element of  $W(R^\circ)[1/p]/(\ker \theta)^n$  where  $\theta: W(R^\circ)[1/p] \rightarrow R^\sharp$  is the obvious map.

**Proposition 26.12.**  $B_{\mathrm{dR}}^+/\mathrm{Fil}^n$  is a diamond.

*Proof.* Let  $\widehat{\mathbb{G}_m}$  be the formal multiplicative group ( $\widehat{\mathbb{G}_m}(R, R^+) = 1 + R^{\circ\circ}$ ), and let  $\widetilde{\mathbb{G}_m}$  be its universal cover ( $\widetilde{\mathbb{G}_m}(R, R^+) = \varprojlim_{x \mapsto x^p} 1 + R^{\circ\circ}$ ). Then  $\widetilde{\mathbb{G}_m}^n \times \mathrm{Spa} \mathbb{Q}_p^{\mathrm{cyc}}$

is perfectoid, and  $(\widetilde{\mathbb{G}_m}^n \times \mathrm{Spa} \mathbb{Q}_p^{\mathrm{cyc}})^b \cong \widetilde{\mathbb{G}_m}^n \times \mathrm{Spa}(\mathbb{Q}_p^{\mathrm{cyc}})^b$ . There is a map  $\widetilde{\mathbb{G}_m}^n \times \mathrm{Spa}(\mathbb{Q}_p^{\mathrm{cyc}})^b \rightarrow B_{\mathrm{dR}}^+/\mathrm{Fil}^n$  that sends  $(z_0, z_1, \dots, z_{n-1}) \mapsto \sum_{i=0}^n (\log[z_i])(\log[\epsilon])^i \in B_{\mathrm{dR}}^+/\mathrm{Fil}^n$ . The map is pro-étale. To see this, recall that  $\log: (\mathbb{G}_m)_{\mathbb{Q}_p} \rightarrow \mathbb{A}_{\mathbb{Q}_p}^1$  is an étale covering with Galois group  $\mathbb{Q}_p/\mathbb{Z}_p$ , and  $(\widetilde{\mathbb{G}_m})_{\mathbb{Q}_p} \rightarrow (\mathbb{G}_m)_{\mathbb{Q}_p}$  is an étale covering with Galois group  $\mathbb{Z}_p$ , so  $(\widetilde{\mathbb{G}_m})_{\mathbb{Q}_p}$  is a  $\mathbb{Q}_p$ -torsor over  $\mathbb{A}_{\mathbb{Q}_p}^1$ .  $\square$

*Example 26.13.* Here is an example of a diamond that is not of the form  $X^\diamond$ . Let  $T$  be a compact Hausdorff space, and let  $\underline{T}$  be the functor on  $\mathrm{Perf}$  that sends  $X$  to the set of continuous maps  $|X| \rightarrow T$ . We claim that for any perfectoid field  $K$  of characteristic  $p$ ,  $\underline{T} \times \mathrm{Spa}(K, \mathcal{O}_K)$  is a diamond. If  $T$  is profinite, then  $\underline{T} \times \mathrm{Spa}(K, \mathcal{O}_K)$  is perfectoid. In general,  $\underline{T}$  admits a surjection from a profinite set  $S$  (take the Stone-Cech compactification of  $T$  considered as a discrete set). The equivalence relation  $R = S \times_T S$  is a closed subspace of  $S \times S$ , so it is also profinite. One can check that  $\underline{R} \times \mathrm{Spa}(K, \mathcal{O}_K) \rightrightarrows \underline{S} \times \mathrm{Spa}(K, \mathcal{O}_K)$  is pro-étale, and  $\underline{T} \times \mathrm{Spa}(K, \mathcal{O}_K)$  is isomorphic to the quotient  $(\underline{S} \times \mathrm{Spa}(K, \mathcal{O}_K))/(\underline{R} \times \mathrm{Spa}(K, \mathcal{O}_K))$ . The underlying topological space of  $\underline{T} \times \mathrm{Spa}(K, \mathcal{O}_K)$  is  $T$ . The underlying topological space of any diamond of the form  $X^\diamond$  is a locally spectral space. So  $\underline{T} \times \mathrm{Spa}(K, \mathcal{O}_K)$  is of the form  $X^\diamond$  iff  $T$  is profinite.

## 27. QUASI-PRO-ÉTALE SITE

**Definition 27.1.** A perfectoid space is *strictly totally disconnected* if it is quasi-compact and quasiseparated and every étale cover splits.

**Definition 27.2.** Let  $\text{Perf}$  be the site of perfectoid spaces of characteristic  $p$  with the pro-étale topology.

We say that a map  $f: \mathcal{F} \rightarrow \mathcal{G}$  of sheaves on  $\text{Perf}$  is étale if for every perfectoid space  $Y$  with a map  $Y \rightarrow \mathcal{G}$ , the pullback  $\mathcal{F} \times_{\mathcal{G}} Y$  is represented by a perfectoid space  $X$  and  $X \rightarrow Y$  is étale.

We say that  $f$  is *quasi-pro-étale* if it is locally separated and for all strictly totally disconnected perfectoid spaces  $Y$  with a map  $Y \rightarrow \mathcal{G}$ , the pullback  $\mathcal{F} \times_{\mathcal{G}} Y$  is represented by a perfectoid space  $X$  and  $X \rightarrow Y$  is pro-étale.

**Proposition 27.3.** A pro-étale sheaf  $Y$  on  $\text{Perf}$  is a diamond if and only if there is a surjective quasi-pro-étale map  $X \rightarrow Y$  from a perfectoid space  $X$ .

**Lemma 27.4.** Let  $f: X \rightarrow Y$  be a quasi-pro-étale map of pro-étale sheaves. If  $Y$  is a diamond, then  $X$  is also a diamond. Conversely, if  $f$  is surjective and  $X$  is a diamond, then  $Y$  is a diamond.

**Definition 27.5.** We say that a diamond  $X$  is *spatial* if it is qcqs and its underlying topological space is a spectral space. We say that  $X$  *locally spatial* if  $|X|$  can be covered by open sets  $U_i$  such that the restriction of  $X$  to each  $U_i$  is spatial.

**Definition 27.6.** Let  $X$  be a locally spatial diamond. The site  $X_{\text{ét}}$  is the site whose objects are locally separated étale maps  $X' \rightarrow X$ , where  $X'$  is a diamond, with coverings given by families of jointly surjective maps.

Let  $X$  be a diamond. The site  $X_{\text{qproét}}$  is the site whose objects are locally separated quasi-pro-étale maps  $X' \rightarrow X$ , where  $X'$  is a diamond, with coverings given by families of jointly surjective maps.

**Proposition 27.7.** If  $X$  is an analytic adic space over  $\text{Spa}(\mathbb{Z}_p, \mathbb{Z}_p)$ , then there is an equivalence of sites  $X_{\text{ét}} \cong X_{\text{ét}}^{\diamond}$ .

**Definition 27.8.** Let  $X$  be an analytic adic space over  $\text{Spa}(\mathbb{Z}_p, \mathbb{Z}_p)$ . Define  $X_{\text{qproét}} = X_{\text{qproét}}^{\diamond}$ .

**Proposition 27.9.** Let  $X$  be a diamond, and let  $\nu: X_{\text{qproét}} \rightarrow X_{\text{ét}}$  be the projection map. Then the functor  $\nu^*$  from sheaves on  $X_{\text{ét}}$  to sheaves on  $X_{\text{qproét}}$  is fully faithful.

Let  $\mathcal{F}$  be sheaf on  $X_{\text{ét}}$ . Then the natural map  $\mathcal{F} \rightarrow \nu_* \nu^* \mathcal{F}$  is an isomorphism. If  $\mathcal{F}$  is a sheaf of abelian groups, then  $R^i \nu_* \nu^* \mathcal{F} = 0$  for all  $i > 0$ .

Now let  $X$  be rigid analytic space over a nonarchimedean field  $F$  of characteristic zero. We will write  $X_{\text{qproét}}$  for  $X_{\text{qproét}}^{\diamond}$ .

**Lemma 27.10.** The site  $X_{\text{qproét}}$  has a basis consisting of affinoid perfectoid objects  $\text{Spa}(R, R^+)$ , such that there exists a filtered system of finite étale maps  $\text{Spa}(R_i, R_i^+) \rightarrow X$  under  $\text{Spa}(R^{\sharp}, R^{\sharp+}) \rightarrow X$ , for which  $\varinjlim_i R_i \rightarrow R^{\sharp}$  is injective with dense image.

We will define the following sheaves on  $X_{\text{qproét}}$ .

$$\begin{aligned} \mathcal{O}_X &= \nu^* \mathcal{O}_{X_{\text{ét}}} \\ \mathcal{O}_X^+ &= \nu^* \mathcal{O}_{X_{\text{ét}}}^+ \end{aligned}$$

Let  $\widehat{\mathcal{O}}_X$  (resp.  $\widehat{\mathcal{O}}_X^+$ ) be the sheaf that sends an affinoid perfectoid  $\text{Spa}(R, R^+)$  to  $R^{\sharp}$  (resp.  $R^{\sharp+}$ ).

*Example 27.11.* Let  $\mathbb{T} = \mathrm{Spa}(\mathbb{Q}_p \langle T^{\pm 1} \rangle)$ , and let  $\mathbb{T}_\infty$  be the perfectoid pro-étale cover  $\mathrm{Spa} \mathbb{Q}_p^{\mathrm{cyc}} \langle T^{\pm 1/p^\infty} \rangle$ . Then

$$\begin{aligned} \mathcal{O}_{\mathbb{T}}(\mathbb{T}_\infty) &= \varinjlim_n \mathbb{Q}(\mu_{p^n}) \langle T^{\pm 1/p^n} \rangle \\ \widehat{\mathcal{O}}_{\mathbb{T}}(\mathbb{T}_\infty) &= \mathbb{Q}_p^{\mathrm{cyc}} \langle T^{\pm 1/p^\infty} \rangle. \end{aligned}$$

**Lemma 27.12.**

$$\widehat{\mathcal{O}}_X^+ = \varprojlim_n \mathcal{O}_X^+ / p^n$$

**Corollary 27.13.** *For any pseudouniformizer  $\varpi$  of  $F$ ,*

$$\nu_* \widehat{\mathcal{O}}_X^+ / \varpi \cong \mathcal{O}_{X_{\mathrm{\acute{e}t}}}^+ / \varpi.$$

**Corollary 27.14.**

$$H^i(X_{\mathrm{\acute{e}t}}, \mathbb{F}_p) \cong H^i(X_{\mathrm{qpro\acute{e}t}}, \mathbb{F}_p).$$

*For any pseudouniformizer  $\varpi$  of  $K$ ,*

$$H^i(X_{\mathrm{\acute{e}t}}, \mathcal{O}_{X_{\mathrm{\acute{e}t}}}^+ / \varpi) \cong H^i(X_{\mathrm{qpro\acute{e}t}}, \widehat{\mathcal{O}}_X^+ / \varpi).$$

*Proof.* The first claim follows from Proposition 27.9, and the second follows from Corollary 27.13.  $\square$

**Theorem 27.15** (Primitive comparison theorem). *Let  $C$  be a complete algebraically closed extension of  $\mathbb{Q}_p$ , and let  $X$  be a proper smooth rigid analytic space over  $C$ . Then the natural maps*

$$\begin{aligned} H^i(X_{\mathrm{\acute{e}t}}, \mathbb{F}_p) \otimes_{\mathbb{F}_p} \mathcal{O}_C^a / p &\rightarrow H^i(X_{\mathrm{\acute{e}t}}, \mathcal{O}_{X_{\mathrm{\acute{e}t}}}^{+a} / p) \\ H^i(X_{\mathrm{qpro\acute{e}t}}, \mathbb{Z}_p) \otimes_{\mathbb{Z}_p} \mathcal{O}_C^a &\rightarrow H^i(X_{\mathrm{qpro\acute{e}t}}, \widehat{\mathcal{O}}_X^{+a}) \\ H^i(X_{\mathrm{qpro\acute{e}t}}, \mathbb{Q}_p) \otimes_{\mathbb{Q}_p} C &\rightarrow H^i(X_{\mathrm{qpro\acute{e}t}}, \widehat{\mathcal{O}}_X) \end{aligned}$$

*are isomorphisms.*

**Proposition 27.16.** *Let  $X$  be a smooth rigid analytic space over  $C$ . Let  $\lambda: X_{\mathrm{qpro\acute{e}t}} \rightarrow X_{\mathrm{an}}$  be the projection. Then  $R^j \lambda_* \widehat{\mathcal{O}}_X^+ \cong \Omega_X^j(-j)$ .*

*Sketch of proof.* Locally,  $X$  admits an étale map to a torus  $\mathbb{T} = \mathrm{Spa} C \langle T_1^{\pm 1}, \dots, T_n^{\pm 1} \rangle$ . The torus has an explicit perfectoid cover  $\mathbb{T}_\infty = \mathrm{Spa} C \langle T_1^{\pm 1/p^\infty}, \dots, T_n^{\pm 1/p^\infty} \rangle$ , which is a  $\mathbb{Z}_p(1)^n$ -torsor. We know that the structure sheaf on this perfectoid cover is acyclic. The proposition then reduces to a computation of the cohomology group

$$H_{\mathrm{cts}}^j \left( \mathbb{Z}_p(1)^n, C \left\langle T_1^{\pm 1/p^\infty}, \dots, T_n^{\pm 1/p^\infty} \right\rangle \right).$$

It is possible to show that the map

$$H_{\mathrm{cts}}^j(\mathbb{Z}_p(1)^n, \mathbb{Z}_p) \otimes_{\mathbb{Z}_p} C \langle T_1^{\pm 1}, \dots, T_n^{\pm 1} \rangle \rightarrow H_{\mathrm{cts}}^j \left( \mathbb{Z}_p^n, C \left\langle T_1^{\pm 1/p^\infty}, \dots, T_n^{\pm 1/p^\infty} \right\rangle \right)$$

is an isomorphism. Moreover, the cup product determines an isomorphism

$$H_{\mathrm{cts}}^j(\mathbb{Z}_p(1)^n, \mathbb{Z}_p) \cong \wedge^j H_{\mathrm{cts}}^1(\mathbb{Z}_p^n, \mathbb{Z}_p),$$

and there is a natural isomorphism

$$H_{\mathrm{cts}}^1(\mathbb{Z}_p(1)^n, \mathbb{Z}_p) \cong \mathrm{Hom}(\mathbb{Z}_p(1)^n, \mathbb{Z}_p).$$

$\square$

**Corollary 27.17** (Hodge–Tate decomposition). *Let  $X$  be a proper smooth rigid analytic space over a  $p$ -adic field  $K$ , and let  $C = \widehat{K}$ . Then*

$$H_{\text{ét}}^n(X_C, \mathbb{Q}_p) \otimes_{\mathbb{Q}_p} C \cong \bigoplus_{i+j=n} H^i(X, \Omega_X^j) \otimes_K C(-j).$$

*Sketch of proof.* First use the primitive comparison theorem to write  $H_{\text{ét}}^n(Y_C, \mathbb{Q}_p) = H^n(Y_{C, \text{qproét}}, \widehat{\mathcal{O}}_Y)$ . There is a Leray spectral sequence

$$H^i(Y_C, R^j \lambda_* \widehat{\mathcal{O}}_Y) \Rightarrow H^{i+j}(Y_{C, \text{proét}}, \widehat{\mathcal{O}}_Y).$$

Because there are no nonzero Galois-equivariant maps between  $C(-j)$  and  $C(-j')$  for  $j \neq j'$ , we must have

$$H^n(Y_{C, \text{qproét}}, \widehat{\mathcal{O}}_Y) = \bigoplus_{i+j=n} H^i(Y_C, R^j \lambda_* \widehat{\mathcal{O}}_Y).$$

□

## REFERENCES

- [Ax70] J. Ax. Zeros of polynomials over local fields—The Galois action. *J. Algebra*, 15:417–428, 1970.
- [BC] O. Brinon and B. Conrad. CMI summer school notes on  $p$ -adic Hodge theory. <http://math.stanford.edu/~conrad/papers/notes.pdf>.
- [BMS18] B. Bhatt, M. Morrow, and P. Scholze. Integral  $p$ -adic Hodge theory. *Publ. Math. Inst. Hautes Études Sci.*, 128:219–397, 2018.
- [Bos14] S. Bosch. *Lectures on formal and rigid geometry*, volume 2105 of *Lect. Notes Math.* Cham: Springer, 2014.
- [Fal02] G. Faltings. Almost étale extensions. In *Cohomologies  $p$ -adiques et applications arithmétiques (II)*, pages 185–270. Paris: Société Mathématique de France, 2002.
- [FO] J.-M. Fontaine and Y. Ouyang. Theory of  $p$ -adic Galois representations. <http://staff.ustc.edu.cn/~yiouyang/galoisrep.pdf>.
- [Fon82] J.-M. Fontaine. Formes différentielles et modules de Tate des variétés abéliennes sur les corps locaux. *Invent. Math.*, 65(3):379–409, 1981/82.
- [Hub96] R. Huber. *Étale cohomology of rigid analytic varieties and adic spaces*, volume E30 of *Aspects Math.* Wiesbaden: Vieweg, 1996.
- [Kat73] N. M. Katz.  $p$ -adic properties of modular schemes and modular forms. In *Modular functions of one variable, III (Proc. Internat. Summer School, Univ. Antwerp, Antwerp, 1972)*, pages 69–190. Lecture Notes in Mathematics, Vol. 350. Springer, Berlin, 1973.
- [Ked] K. S. Kedlaya. Sheaves, stacks, and shtukas. <https://swc-math.github.io/aws/2017/2017KedlayaNotes.pdf>.
- [Ked15] K. S. Kedlaya. New methods for  $(\Gamma, \varphi)$ -modules. *Res. Math. Sci.*, 2:Art. 20, 31, 2015.
- [KM85] N. M. Katz and B. Mazur. *Arithmetic moduli of elliptic curves*, volume 108 of *Ann. Math. Stud.* Princeton University Press, Princeton, NJ, 1985.
- [LP19] S. Li and X. Pan. Logarithmic de Rham comparison for open rigid spaces. *Forum Math. Sigma*, 7:53, 2019. Id/No e32.
- [LZ] D. Loeffler and S. Zerbes. Euler systems (2018 Arizona Winter School notes). <https://swc-math.github.io/aws/2018/2018LoefflerZerbesNotes.pdf>.
- [Mat86] H. Matsumura. *Commutative ring theory. Transl. from the Japanese by M. Reid*, volume 8 of *Camb. Stud. Adv. Math.* Cambridge University Press, Cambridge, 1986.
- [Sch13] P. Scholze.  $p$ -adic Hodge theory for rigid-analytic varieties. *Forum Math. Pi*, 1:77, 2013. Id/No e1.
- [Sil09] J. H. Silverman. *The arithmetic of elliptic curves*, volume 106 of *Grad. Texts Math.* New York, NY: Springer, 2nd ed. edition, 2009.
- [Sta] Stacks Project Authors. Stacks Project. <http://stacks.math.columbia.edu>.
- [Tat67] J. T. Tate.  $p$ -divisible groups. In *Proc. Conf. Local Fields (Driebergen, 1966)*, pages 158–183. Springer, Berlin, 1967.

- [Tsu99] T. Tsuji.  $p$ -adic étale cohomology and crystalline cohomology in the semi-stable reduction case. *Invent. Math.*, 137(2):233–411, 1999.