

Estimating Varying Parameters for Stock Return and of CAPM

A Data Analyst and Econometrician's View

Donghwan Kim

AMS 520

Data Science Program, Applied Math and Stats

October 19, 2023

The Objective I

A close look at model specification, data structure, and estimation

- ▶ Estimating β s in Fama-French (1992, pp. 430 - 431)
- ▶ Estimating μ s in Chitsiripanich et al. (2022, Table 1)

They don't have explicit model specifications. Some lecture notes have different model specification, which can cause confusion and misleading. In conclusion, for estimating Fama-French (1992)'s β s,

- (1) **Brunnermeier (2015, Princeton Univ.)'s lecture note is a correct one**
- (2) Different notations and conventions among Panel data econometric modeling, Statistical factor analysis, and Seemingly unrelated regression.

The Objective II

Modeling and estimating **varying intercepts and/or slopes**

► **In cross-sectional time-series or multi-level setting**

- **Mixed model** allows random components
- Andrew Gelman (Columbia University)'s work is famous
- E.g. Kim and Montgomery (2012)'s work

► In the setting of either time-series only or cross-section only

- Usually not estimable but possible with imposing more structure.
- Bayesian: Hierarchical structure with hyper-parameters
- State-space model: Hidden state

Literature Review

Estimating factor models is still an open question

- ▶ Connor, Hagmann, and Linton (2021): Semiparametric estimation of the Fama-French model
- ▶ Fama and French (2015): A five-factor asset pricing model
- ▶ Fama and French (2004): The capital asset pricing model: Theory and evidence
- ▶ Bai (2003): Inferential theory of factor models of large dimensions

Distinguish two different types of factors

- ▶ Observable factors
- ▶ Unobservable factors
 - **Statistical factor analysis** is to estimate unobservable factors.

Notations: A Review

For **Stock Return** (\mathbf{r})

- **Time-series data:** r_t , $t = 1, 2, \dots, T$ (e.g. monthly).
- **Cross-sectional:** r_j , $j = 1, \dots, J$. (e.g. portfolio)
- **Panel data:** $r_{j,t}$, $j = 1, \dots, J$ and $t = 1, 2, \dots, T$.
- **Multi-level**
 - $r_{j,i}$ Stock-level, Portfolio-level, and so on.
 - $r_{k,t}$ Daily, Monthly, and so on.
- **Multi-level cross-sectional time-series:** e.g. $r_{j,i,t}$

Estimating β_j in Fama-French (1992, pp. 430 - 431)

For each β ,

- **Data:** A sample of 330 time series ($t = 1, 2, \dots, 330$).
- **Model:** $r_t = \alpha + \beta r_t^m + \varepsilon_t$, $\varepsilon_t \sim N(0, \sigma^2)$.

$$\begin{bmatrix} r_1 \\ r_2 \\ \vdots \\ r_T \end{bmatrix} = \begin{bmatrix} 1 & r_1^m \\ 1 & r_2^m \\ \vdots & \vdots \\ 1 & r_T^m \end{bmatrix} \begin{bmatrix} \alpha \\ \beta \end{bmatrix} + \begin{bmatrix} \varepsilon_1 \\ \varepsilon_2 \\ \vdots \\ \varepsilon_T \end{bmatrix}$$

where r_t is excess return at time t .

- **Estimation:** OLS

Repeat it for all the portfolios $j = 1, 2, \dots, J$. (And then use $\hat{\beta}_{j,OLS}$ as that of each of the stocks in the portfolio j for the next estimation procedure).

(1) Lecture note 1

Estimate the CAPM time-series regression:

$$R_{it}^e = \alpha_i + \beta_{iM} R_{Mt}^e + \varepsilon_{it}$$

(2) Lecture note 2

Stage 1: Use *time series* data to obtain estimates for each individual stock's β^j

$$R_t^j - R^f = \alpha + \beta^j (R_t^m - R_t^f) + \epsilon_t^j$$

Comparisons of Model Specifications and Data Structures

Data: Cross-sectional time-series data $t = 1, 2, \dots, T$, $j = 1, 2, \dots, J$

Model Specifications:

- ▶ Model I: Same intercept and same slope

$$r_{j,t} = \alpha + \beta r_t^m + \varepsilon_{j,t}$$

- ▶ Model II: Different intercept and different slope

$$r_{j,t} = \alpha_j + \beta_j r_t^m + \varepsilon_{j,t}$$

Model I: Same intercept and slope model

Model: $r_{j,t} = \alpha + \beta r_t^m + \varepsilon_{j,t}$, $t = 1, 2, \dots, T$, $j = 1, 2, \dots, J$.

$$\begin{bmatrix} r_{1,1} \\ r_{1,2} \\ \vdots \\ r_{1,T} \\ r_{2,1} \\ r_{2,2} \\ \vdots \\ r_{2,T} \\ \vdots \\ r_{J,1} \\ r_{J,2} \\ \vdots \\ r_{J,T} \end{bmatrix} = \begin{bmatrix} 1 & r_1^m \\ 1 & r_2^m \\ \vdots & \vdots \\ 1 & r_T^m \\ 1 & r_1^m \\ 1 & r_2^m \\ \vdots & \vdots \\ 1 & r_T^m \\ \vdots & \vdots \\ 1 & r_1^m \\ 1 & r_2^m \\ \vdots & \vdots \\ 1 & r_T^m \end{bmatrix} \begin{bmatrix} \alpha \\ \beta \end{bmatrix} + \begin{bmatrix} \varepsilon_{1,1} \\ \varepsilon_{1,2} \\ \vdots \\ \varepsilon_{1,T} \\ \varepsilon_{2,1} \\ \varepsilon_{2,2} \\ \vdots \\ \varepsilon_{2,T} \\ \vdots \\ \varepsilon_{J,1} \\ \varepsilon_{J,2} \\ \vdots \\ \varepsilon_{J,T} \end{bmatrix}$$

Model II: A varying intercepts and slopes model

Model: $r_{j,t} = \alpha_j + \beta_j r_t^m + \varepsilon_{j,t}$

In matrix form, for a given j

$$\mathbf{r}_j = \alpha_j \mathbf{1}_T + \beta_j \mathbf{r}^m + \varepsilon_j$$

where $\mathbf{r}_j = [r_{j,1}, r_{j,2}, \dots, r_{j,T}]'$, $\mathbf{1}_T = [1, 1, \dots, 1]'$, $\mathbf{r}^m = [r_1^m, r_2^m, \dots, r_T^m]'$.

For all the data,

$$\mathbf{r} = (\mathbf{I}_J \otimes \mathbf{1}_T) \alpha + (\mathbf{I}_J \otimes \mathbf{r}^m) \beta + \varepsilon$$

where $\alpha = [\alpha_1, \alpha_2, \dots, \alpha_J]'$ and $\beta = [\beta_1, \beta_2, \dots, \beta_J]'$. That is,

$$\begin{bmatrix} \mathbf{r}_1 \\ \mathbf{r}_2 \\ \vdots \\ \mathbf{r}_J \end{bmatrix} = \begin{bmatrix} \mathbf{1}_T & 0 & \cdots & 0 & \mathbf{r}^m & 0 & \cdots & 0 \\ 0 & \mathbf{1}_T & \cdots & 0 & 0 & \mathbf{r}^m & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots & \ddots & \vdots & \\ 0 & 0 & \cdots & \mathbf{1}_T & 0 & 0 & \cdots & \mathbf{r}^m \end{bmatrix} \begin{bmatrix} \alpha \\ \beta \end{bmatrix} + \begin{bmatrix} \varepsilon_1 \\ \varepsilon_2 \\ \vdots \\ \varepsilon_J \end{bmatrix}$$

Estimating μ s in Chitsiripanich et al. (2022, Table 1, p.14)

For each sample mean μ (e.g. Excessive Return, alpha, \dots),

► **Data:** $\{y_t\}$ from Jan. 1973 to Dec. 2020.

► **Model:** $y_t = \mu + \varepsilon_t$, $\varepsilon_t \sim N(0, \sigma^2)$.

► **Estimation:** OLS (Ordinary Least Square)

Repeat it for each of the portfolios $j = low, 2, 3, 4, high, high - low$.

To estimate portfolio-specific sample mean, my recommendation is to use the mixed model,

$$y_{j,t} = \mu + u_j + \eta_{j,t}, \quad u_j \sim N(0, \sigma_u^2), \quad \eta_{j,t} \sim N(0, \sigma_\eta^2).$$

- Utilize the full sample, not losing the degree of freedom.
- Take into account correlation among cross-sectional units.
- Flexible modeling: For example, it can quantify the differences among portfolios with a set of dummy variables
- Estimation: GLS, Bayesian (Shrinkage happens).

References

Bai, Jushan, *Inferential Theory for Factor Models of Large Dimensions*, Econometrica, 70(1), 2003,

<https://users.ssc.wisc.edu/~bhansen/718/Bai2003.pdf>

Brunnermeier, Markus k., *Lecture 12: Factor Pricing*, Lecture Note, Department of Economics, Princeton University, 2015.

<https://www.princeton.edu/~markus/teaching/Fin501/12Lecture.pdf>

Chitsiripanich, Soros, Paoella, Marc S., Polak, Pawel, Walker, Patrick S., *Momentum Without Crashes*, Swiss Finance Institute Research Paper Series N°22-87, 2022.

Connor G., Hagmann, M., and Linton, O., *Efficient Semiparametric Estimation of the Fama-French Model and Extensions*, Econometrica, 80 (2), 2012.

Fama, Eugene. F. and French, Kenneth R. *The Capital Asset Pricing Model: Theory and Evidence*, Journal of Economic Perspectives, 18(3), 2004

References

Farago, Adam, *Lecture 4 - Size, Value, and Momentum*, Lecture Note, Department of Economics, University of Gothenburg. 2019.

Kim, Donghwan and Montgomery, Mark R., *Spatially Explicit Models of City Population Growth*, Population Association of America Annual Meeting, San Francisco, May 3-5, 2012. <https://paa2012.princeton.edu/abstracts/122246>.

Susmel, Rauli, *Lecture 8: CAPM*, Lecture Note, University of Huston.
<https://www.bauer.uh.edu/rsusmel/phd/lecture%208.pdf>