

Ethyl Acrylate



What is it?

Ethyl acrylate is an organic compound that is the ethyl ester of acrylic acid. It is a type of acrylate.

Acrylates are used to make strong glues. They are also used to mold things into plasticized shapes. Major sources of exposure include acrylic and gel nails, dental fillings, and hearing aids.

Acrylates start off as a chemical monomer that is a powder, liquid, gel, or paste. This monomer combines and hardens during a process called polymerization or curing. It is the monomer that can cause you to be allergic. Monomers are less likely to be present after curing. However, there may be leftover monomers if curing is incomplete and this can lead to allergic reactions.

Copolymers should not cause you to be allergic.

ETHYL ACRYLATE can also be known as:

- Ethyl propenoate
- Acrylic acid ethyl ester
- 2-methyl-2-propanoic acid ethyl ester

Some Possible Uses of ETHYL ACRYLATE /

Where it might be found:

- Tape (non-rubber)
- Adhesive, sealer, and stoppers
- Artificial eye lens implants
- Artificial fingernails
- Bone cement
- Conducting gel for diathermy
- Soft contact lenses
- Dental fillings, crowns, veneers, dentures, prostheses
- Electron microscopy material
- Exfoliating cleanser
- Eye glasses

- Feminine pads
- Floppy disk and hard disk manufacture
- Fragrance base
- Gel electrophoresis
- Glucose monitor
- Hairspray
- Hearing aids
- Incontinence pads
- Industrial adhesive, epoxies
- In-ear headphones
- Insecticide
- Insulin pump
- Leather finish resins
- Light curing gels
- Microporous biomaterials
- Mineral sedimentation
- Nail extender and hardener
- Nail polish (regular, gel, powder dipped)
- Paint plasticizer
- Paper coating
- Plastic foam and coating
- Plexiglass
- Printing ink, printing plates
- Rubber coating
- Soil improver
- Solvents
- Splints
- TV manufacture
- Textile coatings and sizings
- UV cured glue, ink, paint
- Water purification
- Wig fixing adhesive
- Windshield repair
- Wood lacquer
- Wound spray

DISCLAIMER: Every effort is made to ensure the accuracy of the information provided herein. However, The American Contact Dermatitis Society® makes no warranties or representations of any kind as to its accuracy or completeness. Such information is provided for informational purposes only and is not meant to be a substitute for professional medical advice, diagnosis, treatment, or care.

Ethyl Acrylate



Other products that may cross react with ETHYL ACRYLATE:

Anything that ends in –acrylate such as:

- Methyl methacrylate
- Polymethyl acrylate
- Ethylene glycol dimethacrylate (EGDMA)
- Diethylene glycol dimethacrylate (DEGDMA)
- Triethylene glycol dimethacrylate (TREGDMA)
- 1,4-butanediol dimethacrylate (BUDMA)
- Trimethylol propane trimethacrylate
- 1,6-hexanediol diacrylate (HDDA)
- Triethylene glycol diacrylate (TREGDA)
- Epoxy acrylate
- 2-hydroxyethyl methacrylate (2-HEMA)

* Copolymers that end in –acrylate should be safe for you to use.

What should I do to avoid ETHYL ACRYLATE?

Once you are allergic to one acrylate, it is common that your immune system will react to other acrylates even if you have never been exposed to them before (cross-reactions). Thus, avoidance of all monomers is necessary.

Tell your dentist, manicurist, and orthopedic surgeon about this allergy.

This allergen will penetrate rubber gloves in under 60 seconds. Two pairs of neoprene or polyvinyl alcohol gloves may protect for a few minutes. For more protection, try:

- Honeywell Silver Shield / 4H gloves (part# SSG/7) plastic gloves
(www.allerderm.com, 800-365-6868 or www.northsafety.com, 800-430-4110)

- Polyvinyl alcohol gloves (Ansell AlphaTec part# 15-554)
- Neoprene gloves (Ansell TouchNTuff #73-300 or #73-500, Ansell GAMMEX surgical gloves #20277255)
- The above gloves could be purchased from (www.grainger.com or www.airgas.com or www.nationwideindustrialsupply.com).

Dentists may use zinc phosphate or zinc oxide cement to avoid acrylates. Dentures can also be boiled in the mold to reduce allergens. Or, substitute prostheses made of thermoplastic nylon, terephthalate, or polyurethane. Use hearing aids made of silicone or polyamide nylon.

Instead of acrylic fingernails, you can try preformed plastic nails, silk wrap or dipped nails with cyanoacrylate glue if you test negative to it, or regular nail polish if there are no acrylates.

In addition, the ACDS Contact Allergen Management Program® (CAMP®) can help manage contact dermatitis. It helps to find products that are safe to use by creating a list of products that do not contain one's allergens. For more information about CAMP®, please ask your doctor or visit www.contactderm.org.

REVISION DATE: JANUARY 2026