

Lidocaine



What is it?

Lidocaine is a local anesthetic (numbing medication). It can be applied to the skin or injected.

Based on their chemical structure, anesthetics belong to one of two classes – ester or amide. Cross reactions are more likely to occur within each family. It is important to understand which family of anesthetics you are sensitive to. Lidocaine is in the amide family of anesthetics.

LIDOCAINE can also be known as:

- Lignocaine
- Xylocaine

Some Possible Uses of LIDOCAINE / Where it might be found:

- Anesthetics
- After-sun lotions
- Ear drops
- Mouth and Throat Sprays
- Topical Pain Relief Medications
- Hemorrhoid Medications
- Wound and Burn Ointments
- Desensitizing Topicals

Other products that might react like LIDOCAINE:

If you are allergic to lidocaine, you may need to avoid the entire amide aesthetic group. This may include:

- Bupivacaine
- Dibucaine
- Etidocaine
- Mepivacaine
- Prilocaine
- Pyrrocaine

Cross reactions to amide anesthetics do not always perfectly stay within the amide group. If you have an anesthetic allergy, it is best to get tested for all anesthetics that you might need to use, so that you can be sure which anesthetic is safe for you to use.

What should I do to avoid LIDOCAINE?

In general, do not use lidocaine or amide anesthetics. Please see the CAMP app for a list of products that do not contain amide anesthetics.

Tell your medical provider, dentist, nurse, masseuse, aesthetician, pharmacist, vet, and others that you are allergic to lidocaine and may need to avoid amide anesthetics.

What is okay to use?

People with any type of anesthetic sensitivity can usually use:

- Pramoxine
- Diphenhydramine
- Dyclonine
- Saline injections

These ester anesthetics may be ok to use:

- Benzocaine
- Tetracaine
- Amylocaine (e.g. *Stovaine*)
- Benoxinate (e.g. *Dorsacaine*)
- Benzamine (e.g. *B-eucaine*)
- Butacaine (e.g. *Butelline*)
- Butethamine (e.g. *Monocaine*)
- Chloroprocaine (e.g. *Nesacaine*)
- Cocaine (used in surgery)
- Hexylcaine (e.g. *Cyclaine*)
- Isobucaine (e.g. *Kincaine*)
- Meprylcaine (e.g. *Oracaine*)

DISCLAIMER: Every effort is made to ensure the accuracy of the information provided herein. However, The American Contact Dermatitis Society® makes no warranties or representations of any kind as to its accuracy or completeness. Such information is provided for informational purposes only and is not meant to be a substitute for professional medical advice, diagnosis, treatment, or care.

Lidocaine



- Metabutethamine (e.g. *Unacaine*)
- Metabutozycaine (e.g. *Primacaine*)
- Neapaine (e.g. *Amylsine*)
- Orthocaine (e.g. *Orthoform*, *Neo Orthoform*)
- Phenacaine
- Piperocaine (e.g. *Metycaine*)
- Procaine (e.g. *Novocain*)
- Proparacaine (e.g. *Alcaine*, *Opthaine*)
- Propoxycaine (e.g. *Ravocaine*)
- Risocaine with 2% procaine (e.g. *Propaesin*)

In addition, the ACDS Contact Allergen Management Program® (CAMP®) can help manage contact dermatitis. It helps to find products that are safe to use by creating a list of products that do not contain one's allergens. For more information about CAMP®, please ask your doctor or visit www.contactderm.org.

REVISION DATE: JANUARY 2026

DISCLAIMER: Every effort is made to ensure the accuracy of the information provided herein. However, The American Contact Dermatitis Society® makes no warranties or representations of any kind as to its accuracy or completeness. Such information is provided for informational purposes only and is not meant to be a substitute for professional medical advice, diagnosis, treatment, or care.