

Avhandlingsserie för  
Gymnastik- och Idrottshögskolan

Nr 999

DETERMINANTS OF INTRA-INDIVIDUAL VARIATION IN  
ADAPTABILITY TO RESISTANCE TRAINING OF DIFFERENT  
VOLUMES WITH SPECIAL REFERENCE TO SKELETAL MUSCLE  
PHENOTYPES



# Determinants of intra-individual variation in adaptability to resis- tance training of different volumes with special reference to skeletal muscle phenotypes

Daniel Hammarström

©Daniel Hammarström, Stockholm 2019

ISBN Provided by the library

Printed by Printer service, Stockholm, 2019

Distributor: Gymnastik- och idrottshögskolan

You can have a dedication here if you wish.

## THESIS FOR DOCTORAL DEGREE (Ph.D.)

### **The title of your thesis**

by

**Your name**

Thesis for Philosophy of Doctoral Degree in Sport Sciences, at The Swedish School of Sport and Health Sciences (GIH), which, according to the decision of the dean, will be publicly defended on *DATE*. The thesis defense will be held at the auditorium at The Swedish School of Sport and Health Sciences (GIH), Stockholm.

### **Opponent**

Profesor . . . .

### **Principal supervisor**

Profesor. . .

### **Co-supervisor(s)**

-Professor. . .

-Professor. . .

-Professor. . .

### **Examination board**

-Associate professor. . .

-Professor . . .

-Professor . . .

# Abstract

The preface pretty much says it all.

Second paragraph of abstract starts here.





# List of scientific papers

- I. **Hammarström D**, Øfsteng S, Koll L, Hanestadhaugen M, Hollan I, Apró W, Blomstrand E, Rønnestad B, Ellefsen S Benefits of higher resistance-training volume are related to ribosome biogenesis. *The Journal of physiology*. 2020 Feb;598(3):543-565. doi: 10.1113/JP278455.
- II. Khan Y, **Hammarström D**, Rønnestad B, Ellefsen S, Ahmad R Increased biological relevance of transcriptome analyses in human skeletal muscle using a model-specific pipeline. *BMC Bioinformatics*. 2020 Nov 30;21(1):548. doi: 10.1186/s12859-020-03866-y
- III. **Hammarström D**, Øfsteng S, Jacobsen N, Flobergseter K, Rønnestad B, Ellefsen S Ribosome accumulation during early phase resistance training. *Manuscript*



# Contents

<b>List of Tables</b> . . . . .	<b>xiii</b>
<b>List of Figures</b> . . . . .	<b>xv</b>
<b>1 Introduction</b> . . . . .	<b>1</b>
<b>2 Background</b> . . . . .	<b>3</b>
2.1 Resistance-exercise prescription, influences and challenges . . . . .	3
2.2 Adaptations to resistance training . . . . .	6
2.2.1 Muscle hypertrophy and strength . . . . .	6
2.2.2 Changes in muscle fiber contractile and metabolic characteristics with resistance training . . . . .	8
2.2.3 Connective tissue . . . . .	9
2.3 Effects of exercise prescription on muscle mass and strength . . . . .	10
2.3.1 Effects of resistance exercise volume on muscle strength and mass . . . . .	11
2.4 Molecular determinants of training-induced muscle hypertrophy . . . . .	13
2.4.1 mTOR . . . . .	14
2.4.2 Ribosomal biogenesis . . . . .	15
2.4.3 Transcriptional regulation of muscle mass . . . . .	16
2.5 Effects of exercise volume on molecular determinants of muscle growth . . . . .	17
2.5.1 From Training response . . . . .	17
2.6 From RNA-seq . . . . .	17
2.7 From tr10 . . . . .	19
<b>3 Aims</b> . . . . .	<b>21</b>
<b>4 Methods</b> . . . . .	<b>23</b>
4.1 Study participants, protocols and training interventions . . . . .	23

---

4.2	Resistance training interventions . . . . .	23
4.2.1	Ethical considerations . . . . .	24
4.3	Muscle strength assessments . . . . .	25
4.4	Measures of muscle mass . . . . .	26
4.5	Muscle tissue sampling and preparations for downstream analyses .	28
4.6	RNA analysis . . . . .	31
4.6.1	Hormonal measurements . . . . .	33
4.7	Statistics and data analysis . . . . .	33
4.7.1	Normalization . . . . .	33
4.8	Meta-analysis of within-session training volume . . . . .	35
4.8.1	Literature search, inclusion criteria and coding of studies .	35
4.8.2	Calculations of effect sizes and statistical analysis . . . . .	35
<b>5</b>	<b>Results and Discussion . . . . .</b>	<b>37</b>
5.1	Effects of different training volume on changes in muscle size and function . . . . .	37
5.2	Acute effects of different training volume on determinants of muscle protein synthesis . . . . .	39
<b>6</b>	<b>General Discussion . . . . .</b>	<b>41</b>
	<b>Conclusion . . . . .</b>	<b>43</b>
	<b>References . . . . .</b>	<b>45</b>

# List of Tables

4.1	Participant characteristics . . . . .	24
5.1	Training induced changes in muscle CSA and average strength in Study I . . . . .	38



# List of Figures

2.1	Relationship between RNA content and protein synthesis in rat skeletal muscle, data from (?) . . . . .	14
5.1	Differences in training induced changes to muscle mass and strength measures between volume conditions in Study I . . . . .	39





# 1. Introduction

Skeletal muscle health is essential for physical independence. In a lifespan perspective, measures of muscle mass and/or strength are inversely associated with mortality [1, 2, 3, 4, 5, 6, 7] and disability [8]. Besides adverse associations between low muscle mass and strength and clinical conditions, muscle weakness also accounts for increased health care costs in patient populations [9, 10]. The intercept between muscle mass, muscle function and health status is interrelated with variables such as age and primary illness or injury [11]. This highlights that interventions designed to increase muscle mass and strength are likely to prevent adverse health outcomes across the lifespan. A higher level of muscle mass and functional capacity would counteract the effects of muscle loss due to illness, age or inactivity.

Although a large degree of the observed variations in lean mass and strength are attributed to genetic components [12, 13], environmental factors also contribute, leaving a window of opportunity to increase muscle mass and functional capacity. Among factors affecting muscle mass and functioning are nutrition and pharmacological agents. However, physical activity and specifically systematic resistance training of sufficient volume, intensity and frequency provides a stimulus that promote morphological and functional changes to the human neuromuscular system without adverse side-effects. Irrespective of age, resistance training generally leads to increased muscle mass and strength [14, 15] and is considered safe when performed in a well organized manner [15, 16].

Resistance training can be modulated indefinitely through combined variations of training variables such as frequency, intensity and volume [17, 18]. Well designed training prescriptions should incorporate information about the current state and goals of the trainee to maximize the potential outcome of the training program [17, 19, 18]. Training volume has received particular attention in the scientific community for many reasons. Evidence suggests that exercise volume affects selected molecular determinants of muscle hypertrophy in a dose-dependent manner [20, 21, 22]. Such effects are believed to facilitate long-term training effects as training programs with higher volume generally result in higher gains in muscle mass

and strength with little evidence of differences between age groups or participants with different training backgrounds [23, 24, 25].

A consequence of a more extensive training program is the increased time required to complete such a program. As time constraints has been reported as a limiting factor for engaging in physical activity [26] some merit can be given to arguments against guidelines suggesting higher volume in resistance training prescription [27, 19]. From an individual perspective, training prescription that balances time-requirement with efficacy presumably increases the likelihood of participation in physical activity [26]. From a more general perspective, increased knowledge about mechanisms governing responses to physical training could improve training prescription also for individuals and populations that experience attenuated benefit of resistance training [28]. The overreaching goal of the present thesis is to contribute to understanding individualized training loads. To this end, training volume was used to study the effects of variable training stimulus in within-participant models of exercise-training.

## 2. Background

### 2.1 Resistance-exercise prescription, influences and challenges

Recommendations of systematic physical exercise with the purpose of improving health or physical performance has long been part of human culture, evident from records dating back to ancient Chinese, Indian and Greek civilizations [29]. Today's exercise-training prescription still bear traces of ideas from these eras, further developed during the renaissance and formalized in systems like German Turnen and Ling gymnastics during the nineteenth century [30]. German Turnen as a system of physical activities was established in a time when Germany developed from aristocracy to a unified nation. Turnen not only served as a system of preparing men to fight for the developing nation but also as a way of establishing a national identity. Ling gymnastics shared common origins with German Turnen and also served as a system of military preparation. However, Ling also established systems for medical, pedagogical and aesthetic gymnastics. Especially Ling's medical gymnastics was important for the development of modern exercise prescription as it was scientifically oriented, based on the physiological and medical understanding of that time [30]. The medical gymnastics of the nineteenth century is referenced in twentieth century texts on therapeutic exercise prescription [31].

With the introduction of "heavy resistance exercises" for the development of muscle strength and mass after injury, DeLorme outlined a system on which modern resistance-exercise prescription is based [32]. DeLorme published his system short after the Second World War [32] during which he, as a newly graduated physician, had been working with war injury rehabilitation [33]. Inspired by practitioners of weight training [33], DeLorme specifically emphasized high-resistance, low-repetition exercises where progression was achieved with increased resistance [32] as opposed to previous recommendations of endurance-like exercise where progression was achieved through increased number of repetitions [31]. DeLorme originally used the term "heavy resistance exercises" in contrasts to low-resistance exercises [32],

but as this could be perceived as exercises performed only with heavy weights, the system was renamed *progressive resistance exercise* to better reflect the method [34]<sup>1</sup>. Indeed, central to the system was the concept of repetition maximum [32]. Repetition maximum refers to the external resistance that can be overcome with a given number of repetitions. By adjusting external resistance to each individual's progression over the course of a training program, exercises are both individualized and progress is monitored [32]. DeLorme originally prescribed sessions of up to 100 repetitions performed in sets of 10 repetitions [32] but later revised this recommendation to three sets of 10 repetitions performed with increasing intensities [34].

Scientific inquiries into prescription of resistance training from the first part of the twentieth century concerned its therapeutic use [e.g. 32, 35] but was also evaluated in the context of improving strength and physical performance in healthy populations [e.g. 36, 37, 38]. Scientific contributions soon moved from questions regarding the effectiveness of resistance training *per se* to comparing outcomes from different modes of resistance training [39, 40, 41, 42, 43, 44]. A vocabulary for progressive resistance exercise-training developed through these investigations and parallel practice, the introduction of repetition maximum by DeLorme being one example. These concepts established as modern definitions of exercise variables enabling precise prescription of training loads for a variety of populations and training goals [18].

Although this development started after the Second World War, resistance training was not part of general exercise guidelines until much later. The American College of Sports Medicine (ACSM) position statement on exercise for healthy individuals from 1978 primarily concerned physical fitness in terms of cardio-respiratory fitness [46]. Since the updated 1990 ACSM statement, resistance training is recommended to be included as part of a sensible, general training program [47]. The introduction of resistance-training as part of the ACSM recommendation also coincides with specific recommendations on resistance training being part of

---

<sup>1</sup>In the current text, exercise is defined as an acute bout of physical activity designed to affect physical characteristics such as strength, speed or endurance. Training is defined as the systematic process of combining multiple exercise-sessions performed in sequence over time. DeLorme first used the adjective *heavy* to describe the resistance prescribed to overcome during exercises but later changed this adjective to *progressive*. In modern texts, the adjective is commonly omitted from the description and *resistance exercise/training* is used to describe strength-promoting exercises and training regimes requiring the neuromuscular system to exert force against resistance. Omitting the adjective has led to many heated debates among exercise physiologists as “endurance exercises are also performed against a[n] (external) resistance.” With no ambition to resolve any conflict in the area, *resistance exercise/training* will be used synonymous with *progressive* or *heavy resistance exercise/training*.

other consensus statements [48, Ch. 2]. Consequently, informed by epidemiological data, the most recent general guidelines for physical activity do include resistance training [49].

It could be argued that the above account reflects the fact that common understandings of *why* and *how to* exercise are influenced by societal norms and historic events such as the search of national identity in the nineteenth century or war injuries in the twentieth century [30,33]. In attempting to outline contemporary influences on exercise prescription, one could argue that the development of techniques to collect large amount of biological data is one such influence. The continuously decreasing cost of a sequenced human genome [50] can serve as an example of this development. Such molecular techniques has enabled efforts to describe mechanisms by which exercise training induce favorable adaptations. The newly established Molecular Transducers of Physical Activity Consortium is an example of a large scale effort, explicitly initiated to develop personalized exercise recommendations and identify molecular targets through which effects of exercised may be mimicked [51]. Advances in bio-medical technologies are enablers of this enterprise and the quest to *individualize* exercise based on molecular diagnostics can be seen as a motivation for modern exercise science [51, 52]. Scientific research into contemporary exercise prescription can thus be understood as a part of the era of *personalized medicine*.

A challenge facing this program is to accurately describe etiologies of response heterogeneity associated with physical training. A wide variation of individual responses are commonly observed after standardized resistance-training programs where measures of muscle strength-changes varies from -32 to +250% and muscle size-changes varies from negative to (-11%) to impressively large (+59%) [53, 14]. By relating such variations to individuals genome (DNA) [54], and messenger RNA (mRNA) profiles [55, 56] we are beginning to gain knowledge about the genetic influence on training responses. In such studies, a common strategy has been to dichotomize responses into “responders” and “non-responders” to exercise training. From a public health perspective this is probably fruitful when *non-response* is defined as the absence of meaningful health-related adaptations, or even adverse effects in response to a given training regime [52, 57]. The existence of non-responders would have large implications regarding exercise prescription on the population scale [58], and for any given individual as it, if properly diagnosed, would guide clinical decision-making.

[59]; A key aspect of successful exercise diagnostics would be to take advantage of the relationship between exercise variables (i.e. modality, intensity, volume etc.)

and exercise response for a given individual. By adapting an individuals training program based on some prior knowledge about the individual, it is possible that the response could be positively affected. Observations supporting such notion exists as an individual classified as non-responsive to a specific exercise modality (e.g. endurance training) may be classified as a responder to another (e.g. resistance training) [60]. Even changing training variables within a specific modality have been shown to convert non-responders to responders as endurance training volume was increased [61].

Although apparent reversal of non-response to exercise training has been observed by manipulating training variables, decisive indications for such manipulations are still lacking.

## 2.2 Adaptations to resistance training

### 2.2.1 Muscle hypertrophy and strength

Systematic resistance training typically increases muscle mass and strength, adaptations through which many beneficial effects on health (increased amino acid storage physical independence) and athletic performance are conveyed. Muscle growth is a well characterized response to resistance training. On the whole muscle level, healthy untrained individuals can be expected to increase their muscle mass by  $\sim 5$ -20% when training is conducted over two weeks to 6 months [14, 62, 53]. Over this time span, muscle growth is approximately linear with time [63, 64, 65] and has been detected as early as 3-4 weeks after training initiation, without apparent muscle edema [63, 64].

Relative muscle growth can be expected to be more pronounced in upper-compared to lower-body muscles when loading patterns are similar [62, 66]. This possibly relates to greater every day activity levels of lower-body muscles requiring a larger stimuli for adaptation [67]. Small, but detectable differences in muscle growth is typically seen between sexes for training induced muscle growth in the upper-body [53] but not for lower-body muscles [68]. Furthermore can hypertrophic responses be expected to be reduced with increasing age [66, 69] but increased with sufficient addition of dietary protein [70]. Additionally, training variables such as intensity, volume, frequency (reviewed below) together with other training aids, e.g. manipulation of blood flow through pressure cuffs [71] can effectively modulate resistance training induced hypertrophy. Together, this underlines that both non-modifiable (e.g. sexual dimorphism and age) and modifiable factors (e.g. training

variables and protein supplementation) affects training induced muscle hypertrophy.

Whole muscles growth in response to short term resistance training occurs primarily through growth of individual muscle fibers (muscle cells). This can be assumed as training-induced splitting of existing fibers and/or myogenesis, formation of new muscle cells are likely slow processes and any increase in the number of fibers by such mechanisms would only represent a small addition to the whole muscle mass [72, 73, 74, 67]. Growth of muscle fibers transfers to greater muscle strength through an expansion of the fibers contractile elements. The muscle cell is to a large degree occupied by myofibrils (about 80% of the cell volume [75]), which in turn contain sarcomeres, arranged in series. Upon neural activation and subsequent  $\text{Ca}^{2+}$  release, the sarcomere shortens through interaction between actin and myosin resulting in force generation [76]. With resistance training, the number of parallel myofibrils increases with the growth of individual fibers [77] leading to a greater force generating capacity of the whole muscle [75].

Measures of whole muscle size corresponds well with strength, particularly when a measure of muscle size reflects the muscles cross-sectional area [78, 79]. However, in response to resistance training, increases in strength are typically greater in magnitude than muscle growth [80, 81, 65, 82]. When relationships between resistance-training induced change in muscle size and strength in previously untrained individuals are analyzed, only a small portion of the variation in muscle strength can be accounted for by changes in muscle size ( $\sim 2.5\text{-}28\%$ ) [79, 14, 83] depending on type of measurements and statistical models used [83] This underlines that muscle hypertrophy is one contributing factor to muscle strength gains, but not the only one.

Different experimental models have shown that strength can increase without concomitant muscle hypertrophy. Getting acquainted to the actual strength test through repeated training of maximal performance produces similar gains in strength without concomitant hypertrophy [84]. If resistance training is performed unilaterally and the contralateral limb act as a control, strength gains are typically also seen here [85, 86]. Additionally, systematic imagery training, without muscle activation produces greater strength gains than control and low-intensity training conditions [87]. Together these observations indirectly points to the central nervous system and motor learning as important factors for strength gains. In addition to effects that mainly can be attributed to motor learning, resistance training leads to changed behavior of motor units, estimated from surface electromyograms [88]. Such changes could be attributed to morphological and functional changes of motoroneurons [89].

### 2.2.2 Changes in muscle fiber contractile and metabolic characteristics with resistance training

In adult human skeletal muscles, muscle fiber types can be identified based on their myosin heavy chain isoform composition. Pure fibers express a single myosin heavy chain isoform whereas hybrid fibers co-express isoforms. Primarily three myosin heavy chain protein isoforms are expressed in adult human skeletal muscle, determined transcriptionally through expression of the genes *MYH7*, *MYH2* and *MYH1* corresponding to myosin heavy chain I, IIA and IIX [90]

Single fibers of different fiber types have specific contractile properties regardless of muscular origin [91], with type II fibers displaying greater force-generating capacity and shortening velocities than type I fibers when normalized to fiber cross sectional area [92, 91] These differences directly relates to the myosin heavy chain proteins displaying different characteristics when interacting with actin [93]. Importantly, *in vitro* assays performed at physiological temperatures shows that myosin heavy chain isoforms extracted from type II fibers are two-fold faster compared to type I fibers with no difference between type IIX and IIA [93].

In addition to contractile characteristics, fiber types identified on the basis of their myosin heavy chain content also differs in metabolic profiles. Type I fibers are characterized as having lower glycolytic but higher oxidative potential compared to type II fibers [94]. Differences in metabolic profiles relates to fatigue resistance where type I fibers can maintain power output as well as ATP levels during intense exercise but type II and especially type IIX fibers fail to do so [90, 95].

Fiber type characteristics effectively modulate the muscle's ability to perform specific activities. Differences in fiber type composition within individuals, between different muscles, reflects this as anti-gravity muscle of the lower body typically express more type I fibers compared to upper-body muscles [90, 91]. Differences in fiber type composition between individuals and sexes are however to some degree genetically determined [96, 97, 98]. Among non-genetic factors influencing fiber-type composition is resistance training. Short term resistance training, designed for muscle hypertrophy and strength gains specifically converts type IIX fibers to more fatigue resistant type IIA fibers with unaltered type I fiber proportions [99, 100, 101, 102]. Such conversion is apparent both when measured on the protein and mRNA level [102, 101]. In contrast, reduced activity or inactivity readily increases the proportion of type IIX expressing fibers [103] [104].

Concomitant with resistance training-induced muscle hypertrophy, the relative mitochondrial density of the muscle decrease as myofibrillar protein fractions



increases, evident from electron microscopy [75]. However, a single session of resistance training, albeit with low resistance (30% of 1RM), was shown to increase mitochondrial, as well as myofibrillar and sarcoplasmic protein synthesis rates. When exercise was performed with slower movement speeds (longer time under tension), the increase in mitochondrial protein synthesis was shown to be greater [105]. This indicates that the magnitude of metabolic stress induced by resistance training affects the subsequent mitochondrial remodeling. Such remodeling could explain improved mitochondrial function, measured as mitochondrial respiration in response to 12-weeks of resistance training with less pronounced changes seen in mitochondrial proteins [106]. An improved mitochondrial efficiency could also be linked to fiber type transitions. Mitochondria have the ability to form dynamic networks within cells by fusion (and fission) of individual mitochondria, a characteristic important for normal function [107]. Such behavior have been shown to be fiber type specific in muscle, with oxidative fibers (type I and IIA) compared to glycolytic fibers (type IIX and IIB in mice) displaying greater, elongated mitochondrial networks [108]. In response to endurance training, fiber type switch from glycolytic to oxidative, coincided with switch to less fragmented mitochondria [108]. Such coordinated remodeling can be linked to common molecular mechanisms regulating both fiber type and mitochondrial biogenesis [109, 110].

I

### 2.2.3 Connective tissue

In addition to adaptive changes to the contracting apparatus of single muscle fibers and neural mechanisms regulating their activity, resistance training modulates bone, tendon and connective tissue. From a general perspective, tissues enabling e.g. locomotion by conveying forces produced by contracting muscles stabilizing body segments adapts in an activity specific manner [111, 112]. Specifically, short term resistance training leads to changes in mechanical properties of bone without increases in bone mineral content or density suggesting qualitative changes [113]. Similarly, tendons respond to short term resistance training by increasing stiffness at high mechanical stress levels without increased cross sectional area [114]. Interestingly, these adaptations seem to reach a plateau as no additional change in this characteristic is seen in individuals who have exercised over four years. However increased stiffness at lower mechanical stress levels have been documented [112]. Changes in tendon properties in response to resistance training may thus primarily be associated with qualitative changes [112] as consequence of

exercise and training-induced increased turnover of collagen in tendons, indicating remodelling [112, 115].

Muscle fibers are embedded in connective tissue, observed as surrounding the whole muscle (epimysium), muscle fascicles (perimysium) and muscle fibers (endomysium) [116]. Connective tissue structures constituting the endomysium connects muscle fibers to adjacent fibers, capillaries and nerves which together with higher order structures make up the extracellular matrix enabling mechanical and biochemical interaction between cell types [116, 117]. Together with the myotendinous junction, intramuscular structures (primarily perimysium) transmits forces originating from contracting muscle fibers and to tendon and bone and act as an elastic energy storage during e.g. locomotion [117]. The mechanical properties of the extracellular matrix also allows for mechanical stimuli to be converted to biochemical signaling initiating e.g. responses to exercise.

There is a general coordination between connective tissue and muscle cell remodeling in response to loading, evident coordinated responses from different cell types in response to exercise [117]. The major constituent of the extracellular matrix is collagen, produced in fibroblasts. In response to acute endurance-type exercise, collagen synthesis and muscle cell specific protein synthesis (myofibrillar and sarcoplasmic fractions) rise in a coordinated fashion [118]. Also, in response to short- and long-term resistance training with subsequent muscle hypertrophy, relative collagen content of muscle tissue remains stable [74]. However, fine tuning of such coordination could exist as contraction mode has shown to differentially affect myofibrillar protein but not collagen synthesis after acute exercise [119].

Increased remodeling of components of the extracellular matrix is a typical response to resistance training evident from gene expression studies [120, 121], and studies of acute protein synthesis [122, 119]. Such remodeling may contribute to increases in specific force (force generated per muscle cross section) seen after resistance training through improved lateral force transfer [123].

## 2.3 Effects of exercise prescription on muscle mass and strength

Precise exercise-training prescription gives information on exercises, their sequential order, intensity and volume, rest periods between efforts or sessions and the frequency at which exercise sessions are to be performed [24]. By manipulating these variables, resistance training programs can be tailored to better fit goals

and starting points of any individual. The relative importance of exercise-training variables for training outcomes has been examined in numerous studies including (but not limited to) the overall organization of exercise sessions, [124, 125] training frequency [126], and intensity [127]. It could be argued that training volume is of particular importance for muscle growth as when this variable is held constant, manipulation of other variables has little or no effect hypertrophy [128, 127]. For development of strength, factors such as intensity and within session organization of exercises is of importance [129, 130], however, when other factors are held constant, increased training volume generally leads to increased strength [129,131, 23], similarly to effects of training volume on muscle growth [24,25].

### 2.3.1 Effects of resistance exercise volume on muscle strength and mass

Exercise volume can be prescribed as the within session number of sets performed per muscle group. This unit is practical as it comparable between individuals and muscle groups [132]. Berger conducted an early study concerning effects of resistance exercise volume with the goal to determine what method most efficiently produced strength gains (in healthy young males) [133]. Berger compared one, two and three sets performed with two, six or ten repetition maximum (RM) in the bench press, three times per week, over twelve weeks. As the combined effect of three sets per session was superior regardless of the number of repetitions performed Berger concluded in favor of three sets. This conclusion was later challenged on the basis of data interpretation [27, 19]. Reviwing the study by Berger and others, Carpinelli and Otto arrived to the conclusion that there was “insufficient evidence to support the prevalent belief that a greater volume of exercise (through multiple sets) will elicit superior muscular strength or hypertrophy” [27]. This stand has since been repeatedly put forward as a criticism of higher volume training programs [134,135] and sparked considerable scientific activity. The main argument against the recommendation of additional volume in strength training programs has been the lack of statistically significant results in single studies [19,134]. Indeed, individual studies do not generally agree on dose-dependent effects of training volume on muscle mass and strength gains [136, 137, 138, 139, 140, 141, 41, 142, 143, 144, 145, 146], including studies performed within participants, where different training volumes are allocated to either extremity [147, 148]. For example, differences in strength are between volume conditions are found in older individuals [136, 137, 41] but not confirmed in another study [140]]. Studies shows that more

volume does not lead to increased muscle gains in young individuals [144, 142, 138] a conclusion challenged by others [146, 139].

As previously noted, combining the above results and additional studies, meta-analyses concluded that training volume dose-dependency exists for the development of muscle mass and strength [[129]; [131]; [23]; [24,25]. As a second argument against additional volume in resistance training recommendation has been the cost/benefit relationship of adding training volume without meaningful or substantial additional gains [19, 134], a subsequent question is, whom would benefit from greater volumes and whom would not? Schoenfeld *et al.* combined data from published studies to explore if participant characteristics of the above mentioned studies interacted with training volume in explaining study outcomes. Neither sex, muscle groups nor age interacted with volume prescription indicating that no such factor would be able refine training prescription guidelines [25]. As the number of studies used to synthesis the meta-analysis was relatively low ( $n = 15$ ) and the studies were heterogeneous in terms of e.g. outcome measurements, it may have lacked in power to detect any meaningful interactions. Additionally, included studies may not have been reporting relevant characteristics for such analysis.

Collectively, the available evidence suggest that there is overlap between training outcomes in studies where different volume has been utilized. The overlap cannot, with available data, be explained by general population characteristics such as age or sex. Studying the effect of different training volumes within participants could potentially help to define determinants of training outcomes in response to different volume conditions. Two within-participant studies have investigated the effects of training volume on strength and hypertrophy outcomes. Sooneste *et al.* compared strength outcomes in response to three- and one-set elbow flexor training for 12 weeks in young males using a within-participant protocol (arms allocated to either volume condition). The results showed general benefit of three- over one-set training for muscle hypertrophy and tended to do so also for strength gains [148]. No attempts were made to relate baseline characteristics to the magnitude of differences between volume conditions, presumably due to the small sample size ( $n = 8$ ). Mitchell *et al.* compared muscle hypertrophy and strength gains in response to three- and one-set of knee-extension exercise performed three times per week for ten weeks. The study contained an additional training condition (low intensity, 30% of 1RM performed with three sets) with participants legs assigned to either of the three conditions in a random fashion. No significant differences were reported between volume conditions for muscle mass or strength gains [147]. However, the analyses were performed without taking the correlation between individuals into


account due to the mixed design [147]. No attempts were made to relate any measured characteristic to differences in responses.

## 2.4 Molecular determinants of training-induced muscle hypertrophy

Muscle mass fluctuates as a consequence of the balance between muscle protein synthesis and breakdown. When a net-positive balance is achieved, muscle protein accumulates and muscle mass increases. Following a single bout of resistance exercise, muscle protein synthesis increases over resting levels up to 48-h post-exercise [149, 150, 151, 152, 153, 154] after being blunted during exercise [149]. Muscle protein synthesis and breakdown rates are highly correlated [155, 151] indicating that these processes are mechanistically coupled and fluctuates together. While acute resistance exercise also stimulates protein breakdown, it does so to a lesser extent, leading to an increase in the net protein balance from baseline under favorable conditions (with sufficient addition of dietary protein) [155, 151, 156, 157]. When resistance exercise is performed in the fasted state, post exercise net protein balance can be expected to be negative [[155]; [151].

The available data from acute studies suggests that resistance training leads to muscle hypertrophy through the accumulative effect of repeated bouts of anabolic stimuli. In recent years, this view has been supplemented by evidence suggesting that chronic RT leads to increased resting synthetic rate of muscle protein [154, 158, 159], which has been postulated to be associated with increased translational capacity, i.e. accumulation of ribosomes [159, 160]. This notion is supported by exercise-induced increases in total RNA, a proxy marker of ribosome abundance, which is closely connected to protein synthesis [161, 162] See Figure 2.1.

and muscle hypertrophy [163, 164, 165].



**Figure 2.1:** Data from (?). Group A were fed a diet containing protein, group B were starved or fed a diet not containing protein.

Conversely, inhibition of ribosomal RNA (rRNA) transcription and inhibition of its up-stream transcription factors act to diminish muscle cell growth [164,161,166].

Biosynthesis of novel ribosomes is a complex, highly coordinated and energy demanding process that involves synthesis of both ribosomal proteins and the four mature rRNA transcripts [167, 168, 166]. Ribosomal accumulation is believed to be determined by the rates of pre-rRNA transcription by RNA polymerase I (Pol I), which in turn is regulated by coordinated assembly of a complex of transcription factors at the rDNA promoter [168]. Specifically, activation of the upstream binding factor (UBF) through phosphorylation is needed to initiate transcription [169, 170]. Such activation is at least partly controlled by the mechanosensitive mTOR pathway, with its inhibition being associated with blocked UBF phosphorylation and subsequent rRNA transcription [171, 172]. Interestingly the availability of UBF *per se* has been shown to be a determinant of rRNA transcription [173] through control of rDNA gene activity [174].

[175] [176]

[177]

### 2.4.1 mTOR

The discovery of an immunosuppressant organic compound called rapamycin in the 1960's led to the characterization of a rapamycin sensitive protein involved in cell growth. The protein was later named mechanistic target for rapamycin

(mTOR) [178]. This protein has since been shown play a key role in skeletal muscle hypertrophy in relation to mechanical loading. Bodine *et al.* made a comprehensive characterization of mTOR-mediated skeletal muscle hypertrophy using rodent models in 2001, showing that mTOR activation was essential for load-induced hypertrophy. Additionally, using transfection techniques, they showed that constitutively activated Akt signaling led to hypertrophy in an mTOR-dependent manner, confirmed with concurrent administration of rapamycin [179].

[180]

In this context, molecular signatures conveyed by the mechanistic target of rapamycin complex 1 (mTORC1) has been in particular focus. Inhibition of mTORC1 impairs protein synthesis in humans [181] and activation of its associated downstream target S6 kinase 1 (S6K1) correlates with increases in muscle protein synthesis and subsequent muscle growth [20, 182]. In line with this, surplus exercise volume leads to greater phosphorylation of S6K1 [20, 21, 22] and is accompanied by increases in myofibrillar protein synthesis [20], fitting the notion that increased training volume provides more pronounced adaptations through repeated episodes of increased protein synthesis.

Individual response patterns to resistance training, including muscle strength and mass, correlate closely with muscle cell characteristics, measured in both rested-state and acute training-phase conditions [55, 121, 164, 182].

### 2.4.2 Ribosomal biogenesis

Conversely, inhibition of ribosomal RNA (rRNA) transcription and inhibition of its up-stream transcription factors act to diminish muscle cell growth [164,161,166].

Biosynthesis of novel ribosomes is a complex, highly coordinated and energy demanding process that involves synthesis of both ribosomal proteins and the four mature rRNA transcripts [167, 168, 166]. Ribosomal accumulation is believed to be determined by the rates of pre-rRNA transcription by RNA polymerase I (Pol I), which in turn is regulated by coordinated assembly of a complex of transcription factors at the rDNA promoter [168]. Specifically, activation of the of the upstream binding factor (UBF) through phosphorylation is needed to initiate transcription [169, 170].

[170] [183]

[184]

[153]

[185]

### 2.4.3 Transcriptional regulation of muscle mass

[186]

[187]

#### Ribosomal biogenesis as a determinant of RT-induced hypertrophy

Resistance exercise is a potent stimuli for rRNA transcription as a single session leads to increases in pre-rRNA [188,189] and repeated bouts lead to accumulation of mature rRNA thus also total RNA and presumably functional ribosomes [188, 190, 165, 164,163, 191, 159]. However, the true time course of ribosomal transcription and accumulation in response to RT remains largely unstudied, with a mere few studies having investigated exercise-induced changes in rRNA over multiple time-points, all of which are either limited to a selected few time-points or a limited time frame. For example, two consecutive bouts of electrically evoked muscle contractions were associated with increased levels of total RNA, with peak values being observed 72 hrs after the second bout [190]. Using voluntary contractions, peak values were reported after nine sessions, followed by a slight decrease to after 18 sessions [191], resembling data from our lab where five sessions of RT led to marked increase in total RNA levels (per-unit muscle tissue), whereby a numerical lowering occurred to after the last training session of the 12 wk interventions (31 sessions). [165]. Interestingly, during the initial phase of RT, total RNA accumulation seems to be volume-dependent, as three sets per exercise in leg exercises led to augmented total RNA and rRNA levels compared to one set per exercise, coinciding with the differences in muscle hypertrophy seen after 12 weeks of RT. [165]. These data suggest that ribosome accumulation reaches a plateau in the early phase of RT and that increases are sensitive to training volume in constant volume protocols.

Recent observations in humans are challenging this view by indicating that translational capacity is a limiting factor for training-induced muscle hypertrophy. First, increased abundances of ribosomal RNA (rRNA) in response to resistance training, measured as total RNA per-weight-unit muscle tissue, correlate with muscle hypertrophy [163]. In accordance with this, training-induced increases in rRNA are larger in muscle hypertrophy high-responders than in low-responders [164, 192]. Secondly, elderly participants typically show blunted ribosome biogenesis, coinciding with attenuated hypertrophic responses [188, 191]. Collectively, these observations suggest that muscle growth depends at least in part on increased translational capacity, making it a prime candidate for explaining the diverse response patterns seen in resistance training with different volume in different



individuals. To date, no study has investigated the association between training volume, ribosome biogenesis and regulation, and gross training adaptations.

## 2.5 Effects of exercise volume on molecular determinants of muscle growth

[193]

[194]

### 2.5.1 From Training response

In humans, the biological adaptation to resistance training varies with exercise-training variables such as volume, intensity, rest between repetitions and sets, selection and order of exercises, repetition velocity and frequency of training sessions [17]. In addition, genetic and epigenetic disposition and environmental factors play a role in variations in adaptations [58, 195, 70]. As time constraints often hinder participation in exercise training-programs [26], numerous studies have searched for the minimal required exercise dose to promote beneficial adaptations. Within-session volume has received particular attention, and although a handful of studies have shown that low-volume training provides similar gains in strength and muscular mass as moderate-volume training [145, 143, 147], meta-analyses conclude in favor of moderate volume protocols [23–25,196]. This apparent discrepancy of specific studies to demonstrate benefits of increased training volume is likely due to a combination of small sample sizes and substantial variation in training responses between individuals and experimental groups. In theory, within-participant designs should alleviate these limitations.

## 2.6 From RNA-seq

Skeletal muscle is a highly adaptable tissue that responds to environmental stress by altering growth rates and differentiation processes. During resistance training, signaling cascades that stimulate muscle plasticity are triggered. Upon repeated exposures, this facilitates growth and a phenotypic shift in a metabolically active direction [197], with the opposite happening during inactivity [198]. Despite this generalized view, muscle responsiveness and plasticity vary, both in response to

different resistance-training protocols [165] and, perhaps more importantly, between individuals [53, > 14]. Selected individuals show a near-complete absence of muscle growth after prolonged resistance training, which markedly reduces the beneficial outcomes of such interventions for muscle function and overall health [14, 53]. Currently, little is known about the etiology of this variation. However, it is usually associated with phenotypic traits of skeletal muscle [199, 164, 200], which implies interactions with environmental factors, genetics, epigenetics, and composites of the intra physiological milieu [70, 201]. This multifaceted origin makes the training-response-spectrum difficult to study directly, with each of the underlying factors offering limited explanatory value alone [58]. Instead, a more indirect approach is necessary, whereby the combined effects of the factors are targeted by studying global patterns of mRNA, protein expression, and skeletal muscle biology.

Previous studies have investigated transcriptome responses to acute resistance exercise [202, 120, 203] and chronic resistance training [204, 205, 120, 206, 202, 121], as well as described associations between transcriptome characteristics and degrees of muscle growth [121, 207], and function [208, 209]. Whereas these studies have merited interesting findings, they lack clear coherences in terms of differential expression events, even for classical exercise-inducible genes such as  $PGC1\alpha$  [210]. This lack of clear coherence is potentially due to a combination of issues such as differences in study design and methods for synthesis and analysis of transcriptome data. First, biologically founded variability can be attributed to differences in exercise protocols (e.g., differences in exercise-volume or intensity). This makes it difficult to discern a general transcriptome exercise response, as training variables are not standardized between studies. Biological heterogeneity is also caused by differences between research participants, affecting signal-to-noise ratios and making it difficult to discern the effects of single independent factors such as training variables. Design stage decisions such as the use of within-participant designs [165, 211] are likely to reduce this variation and to provide transcriptome data with increased biological meaningfulness. Second, technical variability can be attributed to decisions made during the bioinformatical treatment of data. As described by Conceal *et al.*, there is no optimal pipeline for sequencing technology as new tools keep evolving and emerging, different tools should be explored to an optimum pipeline for the specific type of data [212]. To exploit the potential of any study design, there is a need for identifying an appropriate pipeline for transcriptome analyses to ensure a biologically valid interpretation of data. This entails identifying potential violations of common assumptions caused by the experimental model at hand, relating to, for example, data normalization [213, 214].

For transcriptome data to provide adequate biological information about a given experimental set-up, numerous bioinformatic steps need to be adopted in a customized manner [212, 215]. Of these steps, data normalization is particularly decisive [214], as it aims to transform naïve transcript counts into biologically meaningful results. This essentially means expressing them as *per-cell* abundances [216]. For most experimental models, this is equivalent to providing transcript-to-total RNA ratios, given the fulfillment of the assumption that total RNA levels remain stable between conditions on a per-unit-cell or per-unit-tissue basis [216]. In cell models that exhibit high degrees of plasticity, gene expression events result in increased amounts of total RNA and mRNA transcripts per cell [217], specifically violating the assumption that most genes are not differentially expressed [216, 213]. We are not aware of any study that has addressed the need to account for such perspectives during transcriptome analyses of skeletal muscle subjected to mechanical stress, such as resistance training. Indeed, this assumption can be expected to be violated, as total RNA content increases markedly on a per-unit-weight basis [165], with potential global changes also occurring for the mRNA pool, though this remains unknown. The extent to which total RNA, and therefore ribosomal RNA, increases, coincides with the increase in muscle mass [165,164], underlining its importance for cellular growth but also its inevitable presence as a potential confounding factor in RNA sequencing experiments.

## 2.7 From tr10

Skeletal muscle is a critical target for interventions that promotes health across the lifespan [11], with resistance training (RT) being the advocated remedy. Prolonged RT leads to changes in the balance between muscle protein breakdown and synthesis, with one bout of resistance exercise acutely increasing protein synthesis for up to 48 hrs after exercise [151], with subsequent repeated bouts leading to accumulation of muscle protein over time [218,219].



### 3. Aims

The primary aim of this thesis was to relate the adaptive response to resistance training with low- and moderate-volume to skeletal-muscle characteristics in previously untrained individuals. The key question was whether manipulation of exercise-volume will have diverse effects in different individuals related to muscular intrinsic characteristics. A further aim was to characterize exercise-volume dependence in muscle molecular characteristics and determine a time course profile of markers of ribosomal biogenesis in response to resistance training. Based on these aims, the objectives of the present thesis were;

- to relate skeletal muscle and systemic characteristics to benefit of moderate-compared to low-volume resistance training;
- To determine volume-dependence in molecular networks related to muscle growth and remodeling in response to resistance training
- To determine a time course of markers related to ribosome biogenesis in the early phase of resistance training.



## 4. Methods

### 4.1 Study participants, protocols and training interventions

Study I was designed to examine effects of low- and moderate-volume on responses to acute exercise and long-term training within participants. Forty-one healthy individuals were recruited and 34 of these completed at least 85% of the prescribed sessions and were thus included in subsequent data analyses. Reasons for not completing the trial included injury not related to the study ( $n = 1$ ), pain or discomfort during exercises ( $n = 5$ ) and non-adherence to the study protocol. There were no differences in characteristics between participants included in or excluded from data analysis in Study I.

Study II was designed to study the effects of resistance training *per se*, and effects of variable volume on selected markers related to ribosome biogenesis. Participants were therefore recruited to a training group ( $n = 11$ ) and a non-training control group ( $n = 8$ ). Eligible for participation in both studies were young (Study I age 18-40; Study II 18-35), non-smoking men and women. Exclusion criteria included a training history of more than one weekly session during the last 12 (Study I) or six (Study II) months leading up to the study. Participants were also screened for intolerance to local anesthetic, current or previous injuries affecting their ability to perform resistance training, self-reported symptoms or history of disease, intake of medication or supplements with known effects on adaptations to training. Participant characteristics for both studies are shown in Table 4.1.

### 4.2 Resistance training interventions

Each training session started with a light standardized warm-up (5 min ergometer cycling and 10 repetitions each of push-ups, sit-ups, back-extensions and squats). Before each exercise in the main program, one set of 10 repetitions were performed in the specific exercise with approximately 50% of 1RM.

**Table 4.1:** Participant characteristics

	Sex	Age (years)	Stature (cm)	Mass (kg)	Fat mass (%)	Lean mass (%)
Study I	Included	Female	22.0 (1.3)	168 (7)	64.4 (10.4)	34.1 (5.6)
		Male	23.6 (4.1)	183 (6)	75.8 (10.7)	20.4 (6.0)
	Excluded	Female	22.9 (1.6)	166 (8)	64.6 (9.7)	28.8 (8.7)
		Male	24.3 (1.5)	189 (5)	88.2 (22.4)	24.3 (15.3)
	Training	Female	23.4 (2.9)	168 (8)	64.0 (9.2)	30.8 (7.1)
		Male	25.7 (5.8)	177 (3)	77.5 (8.0)	25.3 (3.9)
Study II	Control	Female	24.1 (3.5)	166 (4)	63.8 (0.6)	30.5 (6.4)
		Male	25.5 (5.5)	182 (5)	76.5 (7.7)	18.2 (5.1)

Data are means and (SD)

Studies were fully or partially performed as within-participant studies as each participant had their legs assigned to different training conditions (not including the control group in Study II). Allocation was performed after enrollment where each participant had their legs randomized to either low- or moderate volume (Study I), or variable or constant volume (Study II).

In Study I, the low-volume protocol consisted of a single set of each exercise and the moderate-volume consisted of three sets per exercise. Three unilateral leg exercises were used (leg press, leg curl and knee extension). The moderate volume-leg commenced all sessions and the low volume-leg performed a single set of each exercise in the rest between second and third set of the moderate volume training protocol.

In Study II, only unilateral knee-extension was performed in an effort to concentrate the stimulus to the quadriceps muscles. The constant-volume leg performed six sets of 10RM throughout the study and variable leg performed six sets in session one to four, three sets in session five to eight and nine sets in session nine to twelve with same intensity (10RM).

#### 4.2.1 Ethical considerations

Both studies were approved by the local ethics committee Lillehammer University College/Inland Norway University of Applied Sciences and the Norwegian Centre for Research Data. In accordance with the *Declaration of Helsinki*[220] the studies were pre-registered in publicly accessible databases (Study I, ClinicalTrials.gov Identifier: NCT02179307; Study II, <https://osf.io/wa96y>). Participants were informed of the study design, potential risks and sources of discomfort prior to



giving their informed consent.

### 4.3 Muscle strength assessments

Isokinetic and isometric unilateral knee-extension strength was assessed in a dynamometer (Cybex 6000, Cybex International, Medway USA). Participants were seated and secured in the dynamometer with the knee joint aligned with the rotation axis of the dynamometer. Maximal isokinetic torque was assessed at three angular speeds ( $60^\circ$ ,  $120^\circ$  and  $240^\circ \times \text{sec}^{-1}$ ). Prior to testing, participants were familiarized with the test protocol by performing three submaximal efforts at each angular speed. Participants were given two attempts at  $60^\circ \times \text{sec}^{-1}$  and three attempts at  $120$  and  $240^\circ \times \text{sec}^{-1}$  performed in immediate succession. The highest value was used for statistical analyses. After isokinetic testing, maximal voluntary contraction torque (MVC) was assessed at a knee angle of  $30^\circ$  (full extension =  $90^\circ$ ). Participants were instructed to push with maximal force against the lever for 5 sec. Participants were given two attempts, with 30 sec rest in-between. The highest value was used for downstream analyses.

Maximal strength was assessed as one repetition-maximum (1RM) in unilateral leg-press and knee-extension. The test session for each exercise started with a specific warm-up consisting of ten, six and three repetitions at 50, 75 and 85% of the anticipated maximum. Thereafter, 1RM was found by increasing the resistance progressively until the weight could not be lifted through the full range of motion. For each exercise, the highest load successfully attempted was defined as 1RM. Each participant was given four to six attempts.

At baseline, 1RM, isokinetic and isometric strength assessments were performed twice, separated by at least four days. The maximum value achieved for each of the tests was used in subsequent analysis. Strength tests were separated by at least 48 hours from preceeding training sessions. A combined measure of muscle strength was calculated as the average of all tests (1RM, isometric and isokinetic), wherein each test modality was given equal weight. A subset of the participants ( $n=18$ ) performed strength assessment during the course of the study (at week two, five and nine). For the remaining participants, ordinary training sessions were prioritised when participants missed training or testing due to illness or scheduling difficulties.

Muscle strength was assessed as maximal voluntary isokinetic ( $90^\circ \text{ sec}^{-1}$ ) and isometric ( $60^\circ$  angle, fully extended leg  $0^\circ$ ) knee extension torque. After a brief warm-up (5-min cycling, RPE 12-14), participants were seated and secured in

the individually adjusted dynamometer. Participants were instructed to gradually increase their effort during three warm-up repetitions (50, 60 and, 70% of subjective maximal effort). After a 30-sec restitution period participants were instructed to perform three repetitions with maximal effort in the concentric phase. Sixty seconds after the isokinetic test the lever automatically moved to a 60° angle and participants were instructed to push against the lever enough to see feedback from the visual feedback system. After an additional 15-sec restitution period, participants were instructed to push against the lever with maximal effort. Within the same assessment session, participants remained seated in the dynamometer for measurement performed on both legs. The first measurement was alternated between legs every other session. For statistical treatment of the data, all successful attempts were used. The last strength assessment at baseline was performed at least seven days prior to the first biopsy sampling. At least one of the baseline strength tests was performed on separate day with two sessions allowed to be performed on the same day with a short rest between assessments. Post training assessments were performed 48 hours and eight days after the last session.

## 4.4 Measures of muscle mass

In Study I muscle mass was measured by magnetic resonance imaging (MRI) and dual energy X-ray absorptiometry (DXA) prior to and after the intervention. Both MRI and DXA measurements were completed during the same visit to the laboratory. Participants were instructed to refrain from strenuous physical activity during the last 48 h leading up to the measurements. The post-training measurements were completed at least 48 h after the last strength testing session. Participants were asked to refrain from food consumption during 2 h leading up to the measurements.

MRI images were obtained from the mid-thigh and analyzed by the same investigator blinded for time (pre- and post-training) and condition (low- and moderate-volume). Multiple images were used to estimate the cross-sectional area of the extensor muscles at the same distance from the knee-joint.

See figure

In Study II, m. vastus lateralis muscle thickness was measured by B-mode ultrasonography (SmartUs EXT-1M, Telemed, Vilnius, Lithuania) using a 39 mm 12 MHz, linear array probe. At least three images were captured for each leg per time-point. Between each image acquisition the probe was relocated to the same position where on the skin and subsequently marked on a soft transparent plastic

sheet superimposed on the thigh.

Muscle thickness of *m. vastus lateralis* and *m. rectus femoris* were measured using Transverse images were obtained  $\sim 60\%$  distally from the trochanter major towards the femoral lateral epicondyle. Landmarks such as moles and scars were also marked on the plastic sheet for relocation of the scanned areas during post-training measurements. During analysis, pre and post images from the same participant were analyzed consecutively using the Fiji software,<sup>66</sup> and by two independent researchers. The average muscle thickness of the three images captured per muscle was used for further analyses.

For determination of body composition participants were scanned using DXA (Lunar Prodigy densitometer, GE Healthcare, Madison, WI, USA) with the standard scanning mode (13-25 cm). Participants were lying supine within the scanning bed reference lines, with a strap secured around the ankles to ensure a standardized body position in each scan. The scans were conducted with participants in a fasted state between 07.00-10.00 AM, with empty bladder and wearing only under-wear. Prior to each scan, a phantom scan was run to prevent baseline drifting from affecting analyses. The same technician was used at each time point. Analyses was performed using GE enCORE version 17.0 software (GE Healthcare). Region of interest was customized for covering upper thigh, marked with a square from pubic symphysis to lateral part of tuberculum major, and distal to art. genu.

Muscle thickness (MT) was measured using a B-mode ultra sound unit (SmartUS EXT-1M, Teled, Vilnius, Lithuania). Participants lay supine in a relaxed position for 20 min before assessments, with their feet strapped in a standardized position. A mark was set on the line 60% of the distance between Spinia Iliac Anterior Superior and the lateral femur condyle. MT of *m.vastus lateralis* was measured applying a water-soluble transmission gel (Aquasonic 100 Ultrasound Transmission Gel; Parker Laboratories Inc., Fairfield, NJ, USA), and a 39 mm 12 MHz ultrasound probe was placed perpendicular to the site of interest without pressing the skin. When the quality of the image was satisfactory, evident as distinct upper and lower muscle fascia, three images were captured, where the probe was relocated to the same position between each image. Position of the probe was marked on the skin and subsequently marked on a transparent paper to ensure similar probe placement for both the right and left *m.vastus lateralis* at subsequent assessments. Analyses were done in ImageJ Fiji [221] with images cropped and coded to ensure blinding of the assessor.

## 4.5 Muscle tissue sampling and preparations for downstream analyses

Muscle samples were obtained under local anesthesia (Study I, Xylocaine, 10 mg ml<sup>-1</sup> with adrenalin 5 µg ml<sup>-1</sup>, AstraZeneca, Oslo, Norway; Study II, Lidocaine Mylan, 10 mg ml<sup>-1</sup>, Mylan Ireland Ltd, Ireland) with a fine needle (12-14 gauge; Universal-plus, Medax, Italy) operated with a spring-loaded instrument (Bard Magnum, Bard Norway AS, Norway). Sampling was performed as previously described [222], with modifications. Anesthesia was injected in the subcutaneous tissue with care taken not to inject anesthesia into the muscle itself. Following pilot experiments we decided not to use an insertion cannula as described in [222] as the biopsy needle itself could be used to puncture the skin and muscle fascia. This also resulted in less discomfort. Several passes through the same skin puncture was made to obtain sufficient material for downstream analyses. A smaller needle (14 vs. 12 gauge) was used to further minimize discomfort in Study II where more biopsies were sampled over a shorter time span, with exception from when material was used for immunohistochemistry. The first biopsy was sampled at one third of the distance between the patella to the *anterior superior iliac spinae* with subsequent biopsies sampled ~2 cm proximal to previous samples. In Study II, samples obtained more than one week apart were sampled with closer proximity and distally from previous samples but never at previous sampling sites.

The micro biopsy technique produces smaller samples compared to other biopsy techniques [223], and thus requires several passes to produce sufficient material for multiple downstream experiments. However, reports confirms that the micro biopsy technique is comparable to the traditionally used Bergström technique in several measures of muscle characteristics at the same time as being well tolerated [222,224]. Any reported differences in fiber type distributions between sampling techniques have been suggested relating to differences in sampling depth [224,225].

For determination of fiber type distributions, a threshold of 200-300 fibers has been suggested as a suitable sample size per specimen as more fibers does not reduce the variation between duplicate samples [226]. In Study I, one or several pieces of muscle (total weight ~15 mg) were chosen per sampling for analysis of fiber type distributions (described in detail below). The total number of fibers were counted from these specimens (Figure ref fig). Using an average of fibers from the first sampling time point the between leg coefficient of variation was determined to 14% for Type I fibers and 11.3 for type II fibers. The between leg variation in Type I fibers is similar to what has been previously reported...

### Total RNA extraction

Total-RNA was extracted from pre-weighed, frozen muscle samples with a protocol modified from [227] using Trizol reagent (Life Technologies). Muscle tissue was homogenized in 300  $\mu$ l Trizol with mechanical disruption achieved by Zirconium Oxide Beads (0.5 mm, Next Advance, Inc., New York, USA) and a bead mill (Bullet blender, Next Advance). External, non-mammalian RNA (Lambda PolyA External Standard Kit, Takara Bio Europe, Saint-Germain-en-Laye, France) was added with the initial volume of Trizol to enable per-weight normalization in subsequent analyses. After homogenization, additional Trizol was added to a total volume of 1 ml. Phase separation was achieved by centrifugation after addition of chloroform (200  $\mu$ l). The upper phase (400  $\mu$ l in Study I; 450  $\mu$ l in Study II) was transferred to a fresh tube and RNA was precipitated using isopropanol (500  $\mu$ l). After incubation (10 min, room-temperature) and centrifugation (12000 g, 10 min at 4°C) the resulting RNA pellet was washed three times in chilled 75% ethanol.

As previously mentioned, to minimize discomfort with a larger number of biopsies sampled over a short time, a smaller needle was used for most biopsies in Study II. This generally led to a less tissue used for RNA extractions (Figure XX).

### Protein extraction immunoblotting

Aliquots of muscle-tissue (approximately 25 mg wet weight) were homogenised using a plastic pestle in ice-cold lysis buffer (2 mM HEPES pH 7.4, 1 mM EDTA, 5 mM EGTA, 10 mM MgCl<sub>2</sub>, 1% Triton X-100) spiked with protease and phosphatase inhibitors (Halt, Thermo Fischer Scientific, Life Technologies AS, Oslo Norway), incubated at 4° for 1 hr and centrifuged for 10 min at 10 000 g and 4°C, after which the supernatants were collected. Total protein concentrations were determined on a 1:10 dilution (Pierce Detergent Compatible Bradford Assay Reagent, Thermo Fischer Scientific). The remaining supernatant was diluted to 1.5  $\mu$ g  $\times$   $\mu$ l<sup>-1</sup> total protein in lysis buffer and 4X Laemmli sample buffer (Bio-Rad Laboratories AB, Oslo Norway) containing 2-Mercaptoethanol. Samples were heated to 95°C for 5 min and stored at -20°C until further processing. During analyses, protein samples (20  $\mu$ g of total protein) were separated at 300 V for 30 min using 4-20% gels (Criterion TGX, Bio-Rad), followed by wet transfer to PVDF membranes (0.2  $\mu$ m Immun-Blot, Bio-Rad) at 300 mA for 3 h. Gel electrophoresis and protein transfer were performed at 4°C. Membranes were then stained using a reversible total protein stain (Pierce Reversible Protein Stain, ThermoFischer Scientific) to ensure appropriate protein transfer. Primary antibodies were purchased from Cell

Signaling Technology (Leiden, The Netherlands): mTOR (mTOR<sup>Ser2448</sup>: #5536; pan: #4517), S6 kinase 1 (p85 S6K1<sup>Thr412</sup>: #9206; p70 S6K1<sup>Thr389</sup>: #9234; pan: #2708), ribosomal protein S6 (rpS6<sup>Ser235/236</sup>: #4858; pan: #2317). Membranes were blocked for 2 h in tris-buffered saline (TBS, 20 mM Tris, 150 mM NaCl) containing 3% bovine serum albumin and 0.1% Tween-20, followed by over-night incubation with primary antibodies targeting either the phosphorylated or non-phosphorylated epitope diluted in blocking buffer, followed by 2 h incubation with secondary horseradish peroxidase-conjugated antibodies diluted in TBS containing 0.1% Tween-20 and 5% skimmed milk. Membranes were washed in TBS containing 0.1% Tween-20 for  $6 \times 5$  min after incubation with primary antibody, and for  $8 \times 5$  min after incubation with secondary antibodies. For rpS6 and mTOR antibodies, following chemiluminescent detection (SuperSignal<sup>TM</sup> West Femto Maximum Sensitivity Substrate, ThermoFischer Scientific), membranes were incubated with hydrogen peroxide (15 min, 37°C) to inactivate the horseradish peroxidase (HRP), as described by Sennepin *et al.* [228], followed by over-night incubation with primary or secondary antibodies as described above. If the phosphorylated epitope was targeted during the first incubation, antibodies for the non-phosphorylated epitope were used in the second and vice versa. HRP inactivation did not affect the phosphospecific to non-phosphorylated signal ratios. Importantly, as this technique did not involve removing the first primary antibody, antibodies from different hosts (mouse or rabbit) were used for phosphorylated and non-phosphorylated epitopes respectively. As the antibody targeting p70 S6K1<sup>Thr389</sup> had the same host as the pan-antibody, total-protein was used to normalise chemiluminescent signals. All incubation and washing steps were performed at 4°C using an automated membrane processor (BlotCycler, Precision Biosystems, Mansfield, MA, USA), except for p70 S6K1 experiments, which were performed by hand at room temperature with incubations at 4°C. For mTOR and rpS6, total-protein and chemiluminescence quantification was calculated as the mean value of two separate experiments. S6K1 was quantified once for each phospho-specific antibody. Total-protein content was quantified using ImageJ [229], and was defined as the mean grey value of the whole well with between-well values subtracted as background. Chemiluminescence signals were quantified using Image Studio Lite (LI-COR Biotechnology, Lincoln, Nebraska USA).

Protein was extracted from Trizol preparations according to manufacturers instructions and [230] with modifications. The remaining aqueous phase was removed and DNA was precipitated by the addition of 300  $\mu$ l of absolute ethanol followed by gentle centrifugation (2000 g, 5 min at room temperature). An aliquot

of the phenol-ethanol phase, corresponding to  $\sim 1.75$  mg of tissue, was transferred to a fresh tube. After addition of at least two volumes of isopropanol and incubation (10 min at room temperature), samples were centrifuged (7500 g, 10 min  $4^{\circ}\text{C}$ ) and a pellet formed. The pellet was washed three times in 95% ethanol with each wash separated by centrifugation (5000g, 5 min at room temperature). After the last wash all liquid was removed and 45  $\mu\text{l}$  of Kopec buffer [230] was added (5% SDS, 10 mM Tris, 140 mM NaCl and 20 mM EDTA, pH 8; containing protease and phosphatase inhibitors). Pellets were incubated at  $50^{\circ}\text{C}$  for three hours after which the majority of samples were dissolved. Any undissolved material was sedimented by centrifugation (10000 g, 10 min at room temperature). Protein concentrations were measured (Pierce Detergent Compatible Bradford Assay, ThermoFisher Scientific). Samples were normalized in 4X Laemmli buffer (Bio-Rad Norway AS, Oslo, Norway), boiled ( $95^{\circ}\text{C}$ , 5 min) and stored at  $-20^{\circ}\text{C}$  before later use.

## 4.6 RNA analysis

Complementary DNA (cDNA) was synthesized in technical duplicates from 500 ng of total RNA using random hexamer and anchored Oligo-dT primers (Thermo Fisher Scientific) together with Superscript IV (Thermo Fisher Scientific) according to manufacturer's instruction. qPCR reactions were performed with diluted cDNA (2  $\mu\text{l}$ , 1:25 dilution), a SYBR-green based commercial master mix (PowerUp<sup>TM</sup> SYBR<sup>TM</sup> Green Master Mix, Thermo Fisher) and, target-specific primers (500 nM) in 10  $\mu\text{l}$  reaction volumes using a real-time detection system (QuantStudio 5 Real-Time PCR System, Thermo Fisher Scientific). Fast cycling was used (1 sec denaturing, 30 sec annealing) after UNG (2 min,  $50^{\circ}\text{C}$ ) and polymerase (2 min,  $95^{\circ}\text{C}$ ) activation. Melt curves were collected from all reactions to confirm single product amplification. Primers were further evaluated by agarose gel electrophoresis which confirmed primer sizes and non-template control experiments confirming no amplification without template. Primer sequences and their respective average performances are shown in Table 2.

Approximately 25 mg of wet muscle-tissue was homogenised in a total volume of 1 ml of TRIzol reagent (Invitrogen, Life technologies AS, Oslo, Norway) using 0.5 mm RNase-free Zirconium Oxide beads and a bead homogeniser (Bullet Blender, Next Advanced, Averill Park, NY, USA) according to the manufacturer's instructions. In order to enable analysis of target gene-expression per-unit tissue weight, an exogenous RNA control ( $\lambda$  polyA External Standard Kit, Takara Bio Inc, Shiga, Japan) was added at a fixed amount ( $0.04 \text{ ng} \times \text{ml}^{-1}$  of Trizol

reagent) per extraction prior to homogenisation, as previously described [231, 232]. Following phase-separation, 400  $\mu\text{l}$  of the upper phase was transferred to a fresh tube and RNA was precipitated using isopropanol. The resulting RNA pellet was washed three times with 70% EtOH and finally eluted in TE buffer. RNA quantity and purity was evaluated using a spectrophotometer, all samples had a 260/280  $\text{nm}$  ratio  $> 1.95$ . RNA was stored at  $-80^{\circ}\text{C}$  until further processing. In the analysis of total RNA content per-unit tissue weight, one sample was excluded prior to analysis due to negative deviation from the expected value based on the relationship between sample weight and RNA content suggesting sample loss in washing steps. RNA integrity was assessed by capillary electrophoresis (Experion Automated Electrophoresis Station using RNA StdSens Assay, Bio-Rad) with average integrity scores (RQI) Five-hundred nanograms of RNA were reverse transcribed using anchored Oligo-dT, random hexamer primers (Thermo Scientific) and SuperScript IV Reverse Transcriptase (Invitrogen) according to manufacturer's instructions. All samples were reverse transcribed in duplicates and diluted 1:50 prior to real-time polymerase chain reaction (qPCR). qPCR reactions were run on a fast-cycling real-time detection system (Applied Biosystems 7500 fast Real-Time PCR Systems, Life technologies AS), with a total volume of 10  $\mu\text{l}$ , containing 2  $\mu\text{l}$  of cDNA, specific primers (0.5  $\mu\text{M}$  final concentration) and a commercial master mix (2X SYBR Select Master Mix, Applied Biosystems, Life technologies AS). qPCR reactions consisted of 40 cycles (three seconds  $95^{\circ}\text{C}$  denaturing and 30 seconds  $60^{\circ}\text{C}$  annealing). Melt-curve analyses were performed for all reactions to verify single-product amplification. Gene-specific primers were designed for all targets using Primer-BLAST [233] and Primer3Plus [234] and ordered from Thermo Scientific, except for the external RNA control, for which primers were supplied with the kit. Raw fluorescence data was exported from the platform specific software and amplification curves were modelled with a best-fit sigmoidal model using the qpcR-package [235] written for R [**R-base?**]. Threshold cycles ( $C_t$ ) were estimated from the models by the second-derivate maximum method with technical duplicates modelled independently. Amplification efficiencies were estimated for every reaction [as described by 236, implemented in 235]. For every primer pair, mean amplification efficiencies ( $E$ ) were utilised to transform data to the linear scale using  $E^{-C_t}$ . Primer sequences and primer characteristics (i.e. average primer efficiencies and  $C_t$  values) are presented in Table 2. Gene expression data was log-transformed prior to statistical analysis. As  $C_t$ -values, but not efficiencies are related to RNA integrity [237], RQI scores were used in the statistical treatment of qPCR data to control for potential degradation effects on a by target basis (see



below).

#### 4.6.1 Hormonal measurements

Hormone analyses were performed on blood samples collected at five time points: alongside muscle biopsies (Figure 1, four sampling events) and 10 minutes after completion of the fifth training session. Samples were drawn from the antecubital vein into serum-separating tubes and kept at room temperature for 30 min before centrifugation (1500 g, 10 min). Serum was immediately aliquoted and stored at -80°C until further processing. Serum concentrations of total testosterone, cortisol, growth hormone and insulin-like growth-factor 1 (IGF-1) were measured on an Immulite 1000 analyzer, using kits from the Immulite Immunoassay System menu (Siemens Medical Solutions Diagnostics, NY, USA), performed according to manufacturer's protocols. Serum Vitamin D (S-25-OH-D) levels were measured in samples collected before and after the intervention using a electrochemiluminescence immunoassay (Roche Cobas Vitamin D total assay, Roche Diagnostics GmbH., Mannheim, Germany) using automated instrumentation (Roche Cobas 6000's module e601, Roche Diagnostics GmbH., Mannheim, Germany).

## 4.7 Statistics and data analysis

TO DO:


- For methods discussion, compare product length, efficiencies and ct values in relation to RQI-values. See Fleige 2006 for reference.

#### 4.7.1 Normalization

- An external reference gene was added at a constant amount in Trizol preps
- A normalization factor was used to express relative target gene abundance per-weight tissue.
- In qPCR the linearised expression (effectively  $\hat{c}_q$ ) was used to express the fraction of external reference per total RNA.
- In RNA-seq the external reference gene was sequenced and counts were used to express external RNA as a fraction of total RNA.
- In both cases the normalization factor was calculated as  $mw * counts$ .

A simulation to see that this is equivalent to tissue used in prep when no measurement errors exists.

```
# A tibble: 300 x 7
  mg rna.mg ext tot.rna ext.frac mg.inprep nf
  <dbl> <dbl> <dbl> <dbl> <dbl> <dbl> <dbl>
1     5   250  0.04   1250 0.0000320    4.00 0.000160
2     5   275  0.04   1375 0.0000291    3.64 0.000145
3     5   300  0.04   1500 0.0000267    3.33 0.000133
4     5   325  0.04   1625 0.0000246    3.08 0.000123
5     5   350  0.04   1750 0.0000229    2.86 0.000114
6     5   375  0.04   1875 0.0000213    2.67 0.000107
7     5   400  0.04   2000 0.0000200    2.50 0.000100
8     5   425  0.04   2125 0.0000188    2.35 0.0000941
9     5   450  0.04   2250 0.0000178    2.22 0.0000889
10    5   475  0.04   2375 0.0000168    2.11 0.0000842
# ... with 290 more rows
```



## 4.8 Meta-analysis of within-session training volume

### 4.8.1 Literature search, inclusion criteria and coding of studies

A first set of studies were coded based previously published meta-analyses [24,25]. For more recent studies, PubMed, Google Scholar and SportDiscuss searches were made with search terms being “training volume,” “resistance training,” “strength training,” “set,” “muscle strength,” “muscle hypertrophy” used in different combinations. Studies examining the effect of within-session training volume on muscle strength and mass, with all other training variables kept constant between study groups were considered for inclusion in the meta-analysis. Studies were further assessed for inclusion based on criteria being; (i) participants described as healthy without medications affecting muscle metabolism, (ii) interventions lasting at least 6 weeks and (iii) RT performed without additional stimuli (e.g. blood flow restriction) at intensities above 65% of 1RM or 20RM.

All available outcome measures of muscle mass and strength gains in response to RT were extracted from each study with exception of outcomes reported both as summaries and individual measures (e.g. muscle thickness reported as individual muscles and summarized for the whole muscle group). In such cases the summary was used as outcome. Weekly training volume was calculated as product of weekly sessions, number of sets and exercises for each muscle group assessed for muscle hypertrophy or strength gains. An intervention average of weekly sessions was used when the number of sessions per week differed over the course of the intervention. An exercise was assumed to influence an outcome when it targeted prime movers also assessed for strength or muscle hypertrophy measures. Participant characteristics were coded based on sex (male, female or mixed when a study failed to discern between male and females), age (young, middle-aged, old or mixed), body-mass index (BMI, calculated from average body mass and height when BMI values were not available), training status (trained,  $> 1$  session per week during the last 6 months leading up to the intervention; untrained  $< 1$  session). Study groups were considered independent also in studies utilizing within-participant models.

### 4.8.2 Calculations of effect sizes and statistical analysis

Group-wise effect sizes were calculated for each outcome measure based on the within-group change score pre- to post-training divided by the pre-training standard deviation (SD). Pre-training SD's were calculated as a pooled SD within outcome and study. Variances of the effect size were calculated using an average effect size

across all outcomes within muscle strength or mass, and correlations specific to each measurement type (isokinetic-, isometric- or repetition maximum strength tests; muscle thickness, magnetic resonance imaging, dual energy X-ray absorptiometry) estimated from previous studies.

A correction factor () was applied to both effect sizes and their variances.

Mixed-effects meta-regression models were used to model the effect of weekly number of sets on RT-induced muscle mass and strength gains. Models were fitted in a Bayesian framework using the brms-package [238].

## 5. Results and Discussion

### 5.1 Effects of different training volume on changes in muscle size and function

In Study I, the average increases (Table 5.1) in muscle strength and mass in each volume condition corresponded to what could be expected based on previous studies [239, 14].


Average within participant differences in responses between LOW and MOD were consistent across measures of muscle hypertrophy and strength gains (Figure 5.1). These differences were in agreement to what could be expected based on published meta-analyses [131, 24, 23, 25]. Taken together, these observations confirmed the efficacy of training programs in general and a dose-response with regard to within-session exercise volume.

In Study II, training efficacy was assessed by comparing outcomes to a non-training control group. The training group displayed increases compared to the control group for both strength muscle thickness measures.

**Table 5.1:** Training induced changes in muscle CSA and average strength in Study I

	Sex	Volume condition	Mean (SD)	Reference
CSA %-change	Female	LOW	3.05 (3.61)	
		MOD	5.02 (4.04)	
	Male	LOW	3.83 (3.50)	
		MOD	5.10 (3.71)	
CSA %-change day	Female	LOW	0.04 (0.05)	0.11 [0.04-0.26]a
		MOD	0.07 (0.05)	
	Male	LOW	0.05 (0.05)	
		MOD	0.07 (0.05)	
CSA %-change session	Female	LOW	0.11 (0.13)	0.08 (0.22)b
		MOD	0.18 (0.15)	
	Male	LOW	0.14 (0.12)	
		MOD	0.19 (0.13)	
Average strength %-change	Female	LOW	21.0 (9.8)	
		MOD	27.8 (14.4)	
	Male	LOW	19.2 (12.4)	
		MOD	23.1 (12.0)	
Average strength %-session	Female	LOW	0.77 (0.36)	0.67 (0.35)b
		MOD	1.00 (0.49)	
	Male	LOW	0.72 (0.48)	
		MOD	0.87 (0.46)	

<sup>a</sup> Estimates from Wernbom et al. [239]<sup>b</sup> Estimates from Ahtiainen et al. (ref:ahtiainen-citation)



**Figure 5.1:** Differences in training induced relative changes in muscle mass and strength measures. Estimates are derived from ANCOVA models controlling for baseline values and sex.

## 5.2 Acute effects of diffrent training volume on determi- nants of muscle protein synthesis





## 6. General Discussion



# Conclusion

If we don't want Conclusion to have a chapter number next to it, we can add the `{-}` attribute.

## **More info**

And here's some other random info: the first paragraph after a chapter title or section head *shouldn't be* indented, because indents are to tell the reader that you're starting a new paragraph. Since that's obvious after a chapter or section title, proper typesetting doesn't add an indent there.



## References

- [1] Li R, Xia J, Zhang XI, Gathirua-Mwangi W, Guo J, Li Y, et al. Associations of muscle mass and strength with all-cause mortality among US older adults. *Medicine and Science in Sports and Exercise* 2018;50:458–67. <https://doi.org/10.1249/MSS.0000000000001448>.
- [2] García-Hermoso A, Caverro-Redondo [8], Ramírez-Vélez R, Ruiz JR, Ortega FB, Lee D-C, et al. Muscular strength as a predictor of all-cause mortality in an apparently healthy population: A systematic review and meta-analysis of data from approximately 2 million men and women. *Archives of Physical Medicine and Rehabilitation* 2018;99:2100–2113.e5. <https://doi.org/https://doi.org/10.1016/j.apmr.2018.01.008>.
- [3] Fukasawa H, Kaneko M, Niwa H, Matsuyama T, Yasuda H, Kumagai H, et al. Lower thigh muscle mass is associated with all-cause and cardiovascular mortality in elderly hemodialysis patients. *European Journal of Clinical Nutrition* 2017;71:64–9. <https://doi.org/10.1038/ejcn.2016.186>.
- [4] Miyake H, Kanazawa I, Tanaka KI, Sugimoto T. Low skeletal muscle mass is associated with the risk of all-cause mortality in patients with type 2 diabetes mellitus. *Ther Adv Endocrinol Metab* 2019;10:2042018819842971. <https://doi.org/10.1177/2042018819842971>.
- [5] Ruiz JR, Sui X, Lobelo F, Morrow JR, Jackson AW, Sjöström M, et al. Association between muscular strength and mortality in men: Prospective cohort study. *BMJ* 2008;337:a439. <https://doi.org/10.1136/bmj.a439>.
- [6] Szulc P, Munoz F, Marchand F, Chapurlat R, Delmas PD. Rapid loss of appendicular skeletal muscle mass is associated with higher all-cause mortality in older men: The prospective MINOS study. *Am J Clin Nutr* 2010;91:1227–36. <https://doi.org/10.3945/ajcn.2009.28256>.
- Abramowitz MK, Hall CB, Amodu A, Sharma D, Androga L, Hawkins M. Muscle mass, BMI, and mortality among adults in the united states: A population-based cohort study. *PLoS One* 2018;13:e0194697. <https://doi.org/10.1371/journal.pone.0194697>.
- Janssen I, Heymsfield SB, Ross R. Low relative skeletal muscle mass (sarcopenia) in older persons is associated with functional impairment and physical disability. *J Am Geriatr Soc* 2002;50:889–96. <https://doi.org/10.1046/j.1532-5415.2002.50216.x>.
- Sousa AS, Guerra RS, Fonseca I, Pichel F, Ferreira S, Amaral TF. Financial impact of sarcopenia on hospitalization costs. *Eur J Clin Nutr* 2016;70:1046–51. <https://doi.org/10.1038/ejcn.2016.73>.
- Pinedo-Villanueva R, Westbury LD, Syddall HE, Sanchez-Santos MT, Dennison EM, Robinson SM, et al. Health care costs associated with muscle weakness: A UK population-based estimate. *Calcif Tissue Int* 2019;104:137–44. <https://doi.org/10.1007/s00223-018-0478-1>.
- Wolfe RR. The underappreciated role of muscle in health and disease. *Am J Clin Nutr* 2006;84:475–82.
- Arden NK, Spector TD. Genetic influences on muscle strength, lean body mass, and bone mineral density: A twin study. *Journal of Bone and Mineral Research* 1997;12:2076–81. <https://doi.org/10.1359/jbmr.1997.12.12.2076>.
- Roth SM. Genetic aspects of skeletal muscle strength and mass with relevance to sarcopenia. *BoneKEy Reports* 2012;1:58–8. <https://doi.org/10.1038/bonekey.2012.58>.
- Ahtiainen JP, Walker S, Peltonen H, Holviala J, Sillanpaa E, Karavirta L, et al. Heterogeneity in resistance training-induced muscle strength and mass responses in men and women of different ages. *Age (Dordr)* 2016;38:10. <https://doi.org/10.1007/s11357-015-9870-1>.

- [15] Grgic J, Garofolini A, Orazem J, Sabol[22] Schoenfeld BJ, Pedisic Z. Effects of resistance training on muscle size and strength in very elderly adults: A systematic review and meta-analysis of randomized controlled trials. *Sports Med* 2020. <https://doi.org/10.1007/s40279-020-01331-7>.
- [16] Faigenbaum AD, Myer GD. Resistance training among young athletes: Safety, efficacy and injury prevention effects. *British Journal of Sports Medicine* 2010;44:56. <https://doi.org/10.1136/bjsm.2009.068098>. [24]
- [17] Ratamess N, Alvar BA, Evetoch TK, Housh TJ, Kibler B, Kraemer WJ, et al. American college of sports medicine position stand. Progression models in resistance training for healthy adults. *Med Sci Sports Exerc*[26] 2009;41:687–708. <https://doi.org/10.1249/MSS.0b013e3181915670>.
- [18] Bird SP, Tarpenning KM, Marino FE. Designing resistance training programmes to enhance muscular fitness: A review of the acute programme variables. *Sports Med* 2005;35:8[26] 51.
- [19] Feigenbaum MS, Pollock ML. Prescription of resistance training for health and disease. *Med Sci Sports Exerc* 1999;31:38–45. <https://doi.org/10.1097/00005768-199901000-00008>.
- [20] Burd NA, Holwerda AM, Selby KC, W[27] DW, Staples AW, Cain NE, et al. Resistance exercise volume affects myofibrillar protein synthesis and anabolic signalling molecul[28] phosphorylation in young men. *J Physiol* 2010;588:3119–30. <https://doi.org/10.1113/jphysiol.2010.192856>.
- [21] Terzis G, Spengos K, Mascher H, Georgiadis G, Manta P, Blomstrand E. The degree of [29] S6k and S6 phosphorylation in human skeletal muscle in response to resistance exercise depends on the training volume. *Eur J Appl Physiol* 2010;110:835–43. <https://doi.org/10.1007/s00421-010-1527-2>.
- Ahtiainen JP, Walker S, Silvennoinen M, Kyrolainen H, Nindl BC, Hakkinen K, et al. Exercise type and volume alter signaling pathways regulating skeletal muscle glucose uptake and protein synthesis. *Eur J Appl Physiol* 2015;115:1835–45. <https://doi.org/10.1007/s00421-015-3155-3>.
- Krieger JW. Single versus multiple sets of resistance exercise: A meta-regression. *J Strength Cond Res* 2009;23:1890–901. <https://doi.org/10.1519/JSC.0b013e3181b370be>.
- Krieger JW. Single vs. Multiple sets of resistance exercise for muscle hypertrophy: A meta-analysis. *J Strength Cond Res* 2010;24:1150–9. <https://doi.org/10.1519/JSC.0b013e3181d4d436>.
- Schoenfeld BJ, Ogborn D, Krieger JW. Dose-response relationship between weekly resistance training volume and increases in muscle mass: A systematic review and meta-analysis. *J Sports Sci* 2016;1–0. <https://doi.org/10.1080/02640414.2016.1210197>.
- Choi J, Lee M, Lee JK, Kang D, Choi JY. Correlates associated with participation in physical activity among adults: A systematic review of reviews and update. *BMC Public Health* 2017;17:356. <https://doi.org/10.1186/s12889-017-4255-2>.
- Carpinelli RN, Otto RM. Strength training. Single versus multiple sets. *Sports Med* 1998;26:73–84.
- Pickering C, Kiely J. Do non-responders to exercise exist—and if so, what should we do about them? *Sports Medicine* 2019;49:1–7. <https://doi.org/10.1007/s40279-018-01041-1>.
- Tipton CM. The history of "exercise is medicine" in ancient civilizations. *Adv Physiol Educ* 2014;38:109–17. <https://doi.org/10.1152/advan.00136.2013>.
- Pfister G. Cultural confrontations: German turnen, swedish gymnastics and english sport—european diversity in physical activities from a historical perspective. *Culture, Sport, Society* 2003;6:61–91.

- [31] Nicoll EA. Principles of exercise therapy. *British Medical Journal* 1943;1:747–50. <https://doi.org/10.1136/bmj.1.4302.747>.
- [32] Delorme TL. RESTORATION OF MUSCLE POWER BY HEAVY-RESISTANCE EXERCISES. *JBJS* 1945;27.
- [33] Todd JS, Shurley JP, Todd TC. Thomas l. DeLorme and the science of progressive resistance exercise. *J Strength Cond Res* 2012;26:2913–23. <https://doi.org/10.1519/JSC.0b013e31825adcb4>.
- [34] Delorme TL, Watkins AL. Technics of progressive resistance exercise. *Arch Phys Med Rehabil* 1948;29:263–73.
- [35] Delorme TL, West FE, Shriber WJ. INFLUENCE OF PROGRESSIVE-RESISTANCE EXERCISES ON KNEE FUNCTION FOLLOWING FEMORAL FRACTURES. *JBJS* 1950;32.
- [36] Houtz SJ, Parrish AM, Hellebrandt FA. The influence of heavy resistance exercise on strength. *Physical Therapy* 1946;26:299–304. <https://doi.org/10.1093/ptj/26.6.299>.
- [37] Chui E. The effect of systematic weight training on athletic power. *Research Quarterly American Association for Health, Physical Education and Recreation* 1950;21:188–94. <https://doi.org/10.1080/10671188.1950.10624849>.
- [38] Capen EK. The effect of systematic weight training on power, strength, and endurance. *Research Quarterly American Association for Health, Physical Education and Recreation* 1950;21:83–93. <https://doi.org/10.1080/10671188.1950.10624835>.
- [39] Hettinger T, Müller EA. Muskelleistung und muskeltraining. *Arbeitsphysiologie* 1953;15:111–26. <https://doi.org/10.1007/BF00934143>.
- [40] Capen EK. Study of four programs of heavy resistance exercises for development of muscular strength. *Research Quarterly American Association for Health, Physical Education and Recreation* 1956;27:132–42. <https://doi.org/10.1080/10671188.1956.10612864>.
- Galvao DA, Taaffe DR. Resistance exercise dosage in older adults: Single- versus multi-set effects on physical performance and body composition. *J Am Geriatr Soc* 2005;53:2090–7. <https://doi.org/10.1111/j.1532-5415.2005.00494.x>.
- Berger RA. COMPARISON OF THE EFFECT OF VARIOUS WEIGHT TRAINING LOADS ON STRENGTH. *Res Q* 1965;36:141–6.
- Berger RA, Hardage B. Effect of maximum loads for each of ten repetitions on strength improvement. *Res Q* 1967;38:715–8.
- O'Shea P. Effects of selected weight training programs on the development of strength and muscle hypertrophy. *Res Q* 1966;37:95–102.
- Fleck SJ, Kraemer WJ. Designing resistance training programs. Fourth edition. Champaign, IL: Human Kinetics; 2014.
- American college of sports medicine position statement on the recommended quantity and quality of exercise for developing and maintaining fitness in healthy adults. *Med Sci Sports* 1978;10:vii–x.
- American college of sports medicine position stand. The recommended quantity and quality of exercise for developing and maintaining cardiorespiratory and muscular fitness in healthy adults. *Med Sci Sports Exerc* 1990;22:265–74.
- Manley AF. Physical activity and health: A report of the surgeon general. Diane Publishing; 1996.
- Bull FC, Al-Ansari SS, Biddle S, Borodulin K, Buman MP, Cardon G, et al. World health organization 2020 guidelines on physical activity and sedentary behaviour. *British Journal of Sports Medicine* 2020;54:1451–62. n.d.;2021.
- Sanford JA, Nogiec CD, Lindholm ME, Adkins JN, Amar D, Dasari S, et al. Molecular transducers of physical activity consortium (MoTrPAC): Mapping the dynamic responses to exercise. *Cell* 2020;181:1464–74. <https://doi.org/10.1016/j.cell.2020.06.004>.



- [52] Sparks LM. Exercise training response heterogeneity: Physiological and molecular insights. *Diabetologia* 2017;60:2329–36. <https://doi.org/10.1007/s00125-017-4461-6>. [60]
- [53] Hubal MJ, Gordish-Dressman H, Thompson PD, Price TB, Hoffman EP, Angelopoulos TJ, et al. Variability in muscle size and strength gain after unilateral resistance training. *Med Sci Sports Exerc* 2005;37:964–72. [61]
- [54] Pescatello LS, Devaney JM, Hubal MJ, Thompson PD, Hoffman EP. Highlights from the functional single nucleotide polymorphisms associated with human muscle size and strength or FAMuSS study. *Biomed Res Int* 2013;2013:643575. <https://doi.org/10.1155/2013/643575>. [62]
- [55] Thalacker-Mercer A, Stec M, Cui X, Crossland H, Windham S, Bamman M. Cluster analysis reveals differential transcript profiles associated with resistance training-induced human skeletal muscle hypertrophy. *Physiol Genomics* 2013;45:499–507. <https://doi.org/10.1152/physiolgenomics.00167.2012>. [63]
- [56] Stokes T, Timmons JA, Crossland H, Tripp TR, Murphy K, McGlory C, et al. Molecular transducers of human skeletal muscle remodeling under different loading states. *Cell Rep* 2020;32:107980. <https://doi.org/10.1016/j.celrep.2020.107980>. [64]
- [57] Roth SM. Perspective on the future use of genomics in exercise prescription. *J Appl Physiol* (1985) 2008;104:1243–5. <https://doi.org/10.1152/japplphysiol.01000.2007>. [65]
- [58] Timmons JA. Variability in training-induced skeletal muscle adaptation. *J Appl Physiol* (1985) 2011;110:846–53. <https://doi.org/10.1152/japplphysiol.00934.2010>. [66]
- [59] Yoshie T, Saito C, Kawano F. Early high-fat feeding improves histone modifications of skeletal muscle at middle-age in mice. *Lab Anim Res* 2020;36:25. <https://doi.org/10.1186/s42826-020-00060-2>. [67]
- Hautala AJ, Kiviniemi AM, Mäkipellio TH, Kinnunen H, Nissilä S, Huikuri HV, et al. Individual differences in the responses to endurance and resistance training. *Eur J Appl Physiol* 2006;96:535–42. <https://doi.org/10.1007/s00421-005-0116-2>.
- Montero D, Lundby C. Refuting the myth of non-response to exercise training: 'Non-responders' do respond to higher dose of training. *J Physiol* 2017;595:3377–87. <https://doi.org/10.1113/jp273480>.
- Wernbom M, Augustsson J, Thomeé R. The influence of frequency, intensity, volume and mode of strength training on whole muscle cross-sectional area in humans. *Sports Med* 2007;37:225–64.
- DeFreitas JM, Beck TW, Stock MS, Dillon MA, Kasishke 2nd P. R. An examination of the time course of training-induced skeletal muscle hypertrophy. *Eur J Appl Physiol* 2011;111:2785–90. <https://doi.org/10.1007/s00421-011-1905-4>.
- Stock MS, Mota JA, DeFranco RN, Grue KA, Jacobo AU, Chung E, et al. The time course of short-term hypertrophy in the absence of eccentric muscle damage. *Eur J Appl Physiol* 2017;117:989–1004. <https://doi.org/10.1007/s00421-017-3587-z>.
- Narici MV, Roi GS, Landoni L, Minetti AE, Cerretelli P. Changes in force, cross-sectional area and neural activation during strength training and detraining of the human quadriceps. *Eur J Appl Physiol Occup Physiol* 1989;59:310–9. <https://doi.org/10.1007/bf02388334>.
- Welle S, Totterman S, Thornton C. Effect of age on muscle hypertrophy induced by resistance training. *J Gerontol A Biol Sci Med Sci* 1996;51:M270–5. <https://doi.org/10.1093/gerona/51a.6.m270>.
- Folland JP, Williams AG. The adaptations to strength training : Morphological and neurological contributions to increased strength. *Sports Med* 2007;37:145–68.

- [68] Roberts BM, Nuckols G, Krieger JW. Sex differences in resistance training: A systematic review and meta-analysis. *J Strength Cond Res* 2020;34:1448–60. <https://doi.org/10.1519/jsc.0000000000003521>. [75]
- [69] Peterson MD, Sen A, Gordon PM. Influence of resistance exercise on lean body mass in aging adults: A meta-analysis. *Med Sci Sports Exerc* 2011;43:249–58. <https://doi.org/10.1249/MSS.0b013e3181eb6265>. [76]
- [70] Morton RW, Murphy KT, McKellar SR, Schoenfeld BJ, Henselmans M, Helms E, et al. A systematic review, meta-analysis and meta-regression of the effect of protein supplementation on resistance training-induced gains in muscle mass and strength in healthy adults. *Br J Sports Med* 2018;52:376–84. <https://doi.org/10.1136/bjsports-2017-097608>. [77]
- [71] Lixandrao ME, Ugrinowitsch C, Berton R, Vechin FC, Conceicao MS, Damas F, et al. Magnitude of muscle strength and mass adaptations between high-load resistance training versus low-load resistance training associated with blood-flow restriction: A systematic review and meta-analysis. *Sports Med* 2018;48:361–78. <https://doi.org/10.1007/s40279-017-0795-y>. [79]
- [72] Murach KA, Dungan CM, Peterson CA, Carthy JJ. Muscle fiber splitting is a physiological response to extreme loading in animals. *Exerc Sport Sci Rev* 2019;47:108–15. <https://doi.org/10.1249/jes.0000000000000181>. [80]
- [73] Sjöström M, Lexell J, Eriksson A, Taylor CC. Evidence of fibre hyperplasia in human skeletal muscles from healthy young men? A left-right comparison of the fibre number in whole anterior tibialis muscles. *Eur J Appl Physiol Occup Physiol* 1991;62:301–4. <https://doi.org/10.1007/bf00634963>. [81]
- [74] MacDougall JD, Sale DG, Alway SE, Sutton JR. Muscle fiber number in biceps brachii in bodybuilders and control subjects. *J Appl Physiol Respir Environ Exerc Physiol* 1984;57:1399–403. <https://doi.org/10.1152/jappl.1984.57.5.1399>. [82]
- Lüthi JM, Howald H, Claassen H, Rösler K, Vock P, Hoppeler H. Structural changes in skeletal muscle tissue with heavy-resistance exercise. *Int J Sports Med* 1986;7:123–7. <https://doi.org/10.1055/s-2008-1025748>.
- Sweeney HL, Hammers DW. Muscle contraction. *Cold Spring Harb Perspect Biol* 2018;10. <https://doi.org/10.1101/cshperspect.a023200>.
- Straight CR, Fedewa MV, Toth MJ, Miller MS. Improvements in skeletal muscle fiber size with resistance training are age-dependent in older adults: A systematic review and meta-analysis. *Journal of Applied Physiology* 2020;129:392–403. <https://doi.org/10.1152/japplphysiol.00170.2020>.
- Bamman MM, Newcomer BR, Larson-Meyer DE, Weinsier RL, Hunter GR. Evaluation of the strength-size relationship in vivo using various muscle size indices. *Med Sci Sports Exerc* 2000;32:1307–13. <https://doi.org/10.1097/00005768-200007000-00019>.
- Erskine RM, Fletcher G, Folland JP. The contribution of muscle hypertrophy to strength changes following resistance training. *Eur J Appl Physiol* 2014. <https://doi.org/10.1007/s00421-014-2855-4>.
- Ikai M, Fukunaga T. A study on training effect on strength per unit cross-sectional area of muscle by means of ultrasonic measurement. *Int Z Angew Physiol* 1970;28:173–80. <https://doi.org/10.1007/bf00696025>.
- Young A, Stokes M, Round JM, Edwards RH. The effect of high-resistance training on the strength and cross-sectional area of the human quadriceps. *Eur J Clin Invest* 1983;13:411–7. <https://doi.org/10.1111/j.1365-2362.1983.tb00122.x>.
- Narici MV, Hoppeler H, Kayser B, Landoni L, Claassen H, Gavardi C, et al. Human quadriceps cross-sectional area, torque and neural activation during 6 months strength training. *Acta Physiol Scand* 1996;157:175–86. <https://doi.org/10.1046/j.1365-201X.1996.483230000.x>.

- [83] Vigotsky AD, Schoenfeld BJ, Than C, Brogioni JM. Methods matter: The relationship between strength and hypertrophy depends on methods of measurement and analysis. *PeerJ* 2018;6:e5071. <https://doi.org/10.7717/peerj.5071>. [92]
- [84] Mattocks KT, Buckner SL, Jessee MB, Dankel SJ, Mouser JG, Loenneke JP. Practicing the test produces strength equivalent to higher volume training. *Med Sci Sports Exerc* 2017;49:1945–54. <https://doi.org/10.1249/mss.0000000000001300>. [93]
- [85] Carroll TJ, Herbert RD, Munn J, Lee M, Gandevia SC. Contralateral effects of unilateral strength training: Evidence and possible mechanisms. *J Appl Physiol* (1985) 2006;101:1514–22. <https://doi.org/10.1152/japplphysiol.00531.2006>. [94]
- [86] Green LA, Gabriel DA. The cross education of strength and skill following unilateral strength training in the upper and lower limbs. *J Neurophysiol* 2018;120:468–79. <https://doi.org/10.1152/jn.00116.2018>. [95]
- [87] Zijdwind I, Toering ST, Bessem B, Van Der Laan O, Diercks RL. Effects of imagery motor training on torque production of ankle plantar flexor muscles. *Muscle Nerve* 2003;28:168–73. <https://doi.org/10.1002/mus.10406>. [96]
- [88] Del Vecchio A, Casolo A, Negro F, Scorcelletti M, Bazzucchi I, Enoka R, et al. The increase in muscle force after 4 weeks of strength training is mediated by adaptations in motor unit recruitment and rate coding. *J Physiol* 2019;597:1873–87. <https://doi.org/10.1113/jp277250>. [97]
- [89] Gardiner P, Dai Y, Heckman CJ. Effects of exercise training on alpha-motoneurons. *J Appl Physiol* (1985) 2006;101:1228–36. <https://doi.org/10.1152/japplphysiol.00482.2006>. [98]
- [90] Schiaffino S, Reggiani C. Fiber types in mammalian skeletal muscles. *Physiol Rev* 2011;91:1447–531. <https://doi.org/10.1152/physrev.00031.2010>. [99]
- Harridge SD, Bottinelli R, Canepari M, Pellegrino MA, Reggiani C, Esbjornsson M, et al. Whole-muscle and single-fibre contractile properties and myosin heavy chain isoforms in humans. *Pflügers Arch* 1996;432:913–20.
- Widrick JJ, Stelzer JE, Shoepe TC, Garner DP. Functional properties of human muscle fibers after short-term resistance exercise training. *Am J Physiol Regul Integr Comp Physiol* 2002;283:R408–16. <https://doi.org/10.1152/ajpregu.00120.2002>.
- Lionikas A, Li M, Larsson L. Human skeletal muscle myosin function at physiological and non-physiological temperatures. *Acta Physiol (Oxf)* 2006;186:151–8. <https://doi.org/10.1111/j.1748-1716.2005.01516.x>.
- Essen B, Jansson E, Henriksson J, Taylor AW, Saltin B. Metabolic characteristics of fibre types in human skeletal muscle. *Acta Physiol Scand* 1975;95:153–65. <https://doi.org/10.1111/j.1748-1716.1975.tb10038.x>.
- Karatzafieri C, Haan A de, Mechelen W van, Sargeant AJ. Metabolism changes in single human fibres during brief maximal exercise. *Exp Physiol* 2001;86:411–5. <https://doi.org/10.1113/eph8602223>.
- Simoneau JA, Bouchard C. Genetic determinism of fiber type proportion in human skeletal muscle. *Faseb j* 1995;9:1091–5. <https://doi.org/10.1096/fasebj.9.11.7649409>.
- Simoneau JA, Bouchard C. Human variation in skeletal muscle fiber-type proportion and enzyme activities. *Am J Physiol* 1989;257:E567–72. <https://doi.org/10.1152/ajpendo.1989.257.4.E567>.
- Staron RS, Hagerman FC, Hikida RS, Murray TF, Hostler DP, Crill MT, et al. Fiber type composition of the vastus lateralis muscle of young men and women. *J Histochem Cytochem* 2000;48:623–9.
- Adams GR, Hather BM, Baldwin KM, Dudley GA. Skeletal muscle myosin heavy chain composition and resistance training. *J Appl Physiol* (1985) 1993;74:911–5. <https://doi.org/10.1152/jappl.1993.74.2.911>.

- [100] Williamson DL, Gallagher PM, Carroll JJ, Raue U, Trappe SW. Reduction in hybrid single muscle fiber proportions with resistance training in humans. *J Appl Physiol* (1985) 2001;91:1955–61. <https://doi.org/10.1152/jappl.2001.91.5.1955>. [108]
- [101] Andersen JL, Gruschy-Knudsen T. Rapid switch-off of the human myosin heavy chain IIX gene after heavy load muscle contractions is sustained for at least four days. *Scand J Med Sci Sports* 2018;28:371–80. <https://doi.org/10.1111/sms.12914>. [110]
- [102] Ellefsen S, Vikmoen O, Zacharoff E, Rauk I, Slettalokken G, Hammarstrom D, et al. Reliable determination of training-induced alterations in muscle fiber composition in human skeletal muscle using quantitative polymerase chain reaction. *Scand J Med Sci Sports* 2014;24:e332–42. <https://doi.org/10.1111/sms.12185>. [111]
- [103] Andersen JL, Aagaard P. Myosin heavy chain IIX overshoot in human skeletal muscle. *Muscle Nerve* 2000;23:1095–104.
- [104] Borina E, Pellegrino MA, D'Antona G, Bottinelli R. Myosin and actin content of human skeletal muscle fibers following 35 days bed rest. *Scand J Med Sci Sports* 2010;20:65–73. <https://doi.org/10.1111/j.1600-0838.2009.01029.x>. [112]
- [105] Burd NA, Andrews RJ, West DW, Li JT, JP, Cochran AJ, Hector AJ, et al. Muscle time under tension during resistance exercise stimulates differential muscle protein sub-fractional synthetic responses in men. *J Physiol* 2012;590:351–62. <https://doi.org/10.1113/jphysiol.2011.221200>. [113]
- [106] Porter C, Reidy PT, Bhattarai N, Sidossis LS, Rasmussen BB. Resistance exercise training alters mitochondrial function in human skeletal muscle. *Medicine and Science in Sports and Exercise* 2015;47:1922–31. <https://doi.org/10.1249/MSS.0000000000000605>. [114]
- [107] Chan DC. Mitochondrial fusion and fission in mammals. *Annu Rev Cell Dev Biol* 2006;22:79–99. <https://doi.org/10.1146/annurev.cellbio.22.010305.104638>.
- Mishra P, Varuzhanyan G, Pham AH, Chan DC. Mitochondrial dynamics is a distinguishing feature of skeletal muscle fiber types and regulates organellar compartmentalization. *Cell Metab* 2015;22:1033–44. <https://doi.org/10.1016/j.cmet.2015.09.027>.
- Lin J, Wu H, Tarr PT, Zhang CY, Wu Z, Boss O, et al. Transcriptional co-activator PGC-1 alpha drives the formation of slow-twitch muscle fibres. *Nature* 2002;418:797–801. <https://doi.org/10.1038/nature00904>.
- Handschin C, Chin S, Li P, Liu F, Maratos-Flier E, Lebrasseur NK, et al. Skeletal muscle fiber-type switching, exercise intolerance, and myopathy in PGC-1alpha muscle-specific knock-out animals. *J Biol Chem* 2007;282:30014–21. <https://doi.org/10.1074/jbc.M704817200>.
- Greendale GA, Huang MH, Wang Y, Finkelstein JS, Danielson ME, Sternfeld B. Sport and home physical activity are independently associated with bone density. *Med Sci Sports Exerc* 2003;35:506–12. <https://doi.org/10.1249/01.Mss.0000056725.64347.C9>.
- Magnusson SP, Narici MV, Maganaris CN, Kjaer M. Human tendon behaviour and adaptation, in vivo. *J Physiol* 2008;586:71–81. <https://doi.org/10.1113/jphysiol.2007.139105>.
- Miller LE, Nickols-Richardson SM, Wooten DF, Ramp WK, Steele CR, Cotton JR, et al. Isokinetic resistance training increases tibial bending stiffness in young women. *Calcif Tissue Int* 2009;84:446–52. <https://doi.org/10.1007/s00223-009-9247-5>.
- Massey GJ, Balshaw TG, Maden-Wilkinson TM, Folland JP. Tendinous tissue properties after short- and long-term functional overload: Differences between controls, 12 weeks and 4 years of resistance training. *Acta Physiol (Oxf)* 2018;222:e13019. <https://doi.org/10.1111/apha.13019>.

- [115] Kjaer M, Magnusson P, Krogsgaard M, Boye L, Møller J, Olesen J, Heinemeier K, et al. Extracellular matrix adaptation of tendon and skeletal muscle to exercise. *J Anat* 2006;208:445–50. <https://doi.org/10.1111/j.1469-7580.2006.00549.x>. [122]
- [116] Borg TK, Caulfield JB. Morphology of connective tissue in skeletal muscle. *Tissue Cells* 1980;12:197–207. [https://doi.org/10.1016/0040-8166\(80\)90061-0](https://doi.org/10.1016/0040-8166(80)90061-0). [123]
- [117] Kjaer M. Role of extracellular matrix in adaptation of tendon and skeletal muscle to mechanical loading. *Physiol Rev* 2004;84:649–98. <https://doi.org/10.1152/physrev.00031.2003>. [124]
- [118] Miller BF, Olesen JL, Hansen M, Døssing S, Crameri RM, Welling RJ, et al. Coordinated collagen and muscle protein synthesis in human patella tendon and quadriceps muscle after exercise. *J Physiol* 2005;567:1021–33. <https://doi.org/10.1113/jphysiol.2005.093690>. [125]
- [119] Holm L, Hall G van, Rose AJ, Miller BF, Doessing S, Richter EA, et al. Contraction intensity and feeding affect collagen and myofibrillar protein synthesis rates differently in human skeletal muscle. *Am J Physiol Endocrinol Metab* 2010;298:E257–69. <https://doi.org/10.1152/ajpendo.00609.2009>. [126]
- [120] Damas F, Ugrinowitsch C, Libardi CA, Janin PR, Hector AJ, McGlory C, et al. Resistance training in young men induces muscle transcriptome-wide changes associated with muscle structure and metabolism refining the response to exercise-induced stress. *Eur J Appl Physiol* 2018;118:2607–16. <https://doi.org/10.1007/s00421-018-3984-y>. [127]
- [121] Raue U, Trappe TA, Estrem ST, Qian HR, Helvering LM, Smith RC, et al. Transcriptome signature of resistance exercise adaptations: Mixed muscle and fiber type specific profiles in young and old adults. *J Appl Physiol* (1985) 2012;112:1625–36. <https://doi.org/10.1152/japplphysiol.00435.2011>. [128]
- Moore DR, Phillips SM, Babraj JA, Smith K, Rennie MJ. Myofibrillar and collagen protein synthesis in human skeletal muscle in young men after maximal shortening and lengthening contractions. *Am J Physiol Endocrinol Metab* 2005;288:E1153–9. <https://doi.org/10.1152/ajpendo.00387.2004>.
- Erskine RM, Jones DA, Maffulli N, Williams AG, Stewart CE, Degens H. What causes in vivo muscle specific tension to increase following resistance training? *Exp Physiol* 2011;96:145–55. <https://doi.org/10.1113/expphysiol.2010.053975>.
- Evans JW. Periodized resistance training for enhancing skeletal muscle hypertrophy and strength: A mini-review. *Frontiers in Physiology* 2019;10:13–3. <https://doi.org/10.3389/fphys.2019.00013>.
- Grgic J, Mikulic P, Podnar H, Pedisic Z. Effects of linear and daily undulating periodized resistance training programs on measures of muscle hypertrophy: A systematic review and meta-analysis. *PeerJ* 2017;5:e3695–5. <https://doi.org/10.7717/peerj.3695>.
- Schoenfeld BJ, Ogborn D, Krieger JW. Effects of resistance training frequency on measures of muscle hypertrophy: A systematic review and meta-analysis. *Sports Med* 2016;46:1689–97. <https://doi.org/10.1007/s40279-016-0543-8>.
- Schoenfeld BJ, Grgic J, Ogborn D, Krieger JW. Strength and hypertrophy adaptations between low- vs. High-load resistance training: A systematic review and meta-analysis. *J Strength Cond Res* 2017;31:3508–23. <https://doi.org/10.1519/jsc.0000000000002200>.
- Schoenfeld BJ, Ratamess NA, Peterson MD, Contreras B, Sonmez GT, Alvar BA. Effects of different volume-equated resistance training loading strategies on muscular adaptations in well-trained men. *J Strength Cond Res* 2014;28:2909–18. <https://doi.org/10.1519/JSC.0000000000000480>.

- [129] Grgic J, Schoenfeld BJ, Davies TB, Lazinik B, Krieger JW, Pedisic Z. Effect of resistance training frequency on gains in muscular strength: A systematic review and meta-analysis. *Sports Med* 2018;48:1207–20. <https://doi.org/10.1007/s40279-018-0872-x>.
- [130] Nunes JP, Grgic J, Cunha PM, Ribeiro AS, Schoenfeld BJ, Salles BF de, et al. What influence does resistance exercise order have on muscular strength gains and muscle hypertrophy? A systematic review and meta-analysis. *Eur J Sport Sci* 2020;1–9. <https://doi.org/10.1080/17461391.2020.1733672>.
- [131] Ralston GW, Kilgore L, Wyatt FB, Bales JS. The effect of weekly set volume on strength gain: A meta-analysis. *Sports Med* 2017;47:2585–601. <https://doi.org/10.1007/s40279-017-0762-7>.
- [132] Baz-Valle E, Fontes-Villalba M, Santos-Concejero J. Total number of sets as a training volume quantification method for muscle hypertrophy: A systematic review. *J Strength Cond Res* 2018. <https://doi.org/10.1519/jsc.0000000000002776>.
- [133] Berger R. Effect of varied weight training programs on strength. *Research Quarterly American Association for Health, Physical Education and Recreation* 1962;33:168–81. <https://doi.org/10.1080/10671188.1962.10613188>.
- [134] Junyoung H, Corinna NR, John DS, Sukho L. Low volume progressive single set of resistance training is as effective as high volume multiple sets of resistance protocol on muscle strength and power. *International Journal of Applied Sports Sciences : IJASS* 2015;27:33–42. <https://doi.org/10.24985/ijass.2015.27.1.33>.
- [135] Carpinelli RN. Science versus opinion. *British Journal of Sports Medicine* 2004;38:2010–2. <https://doi.org/10.1136/bjism.2003.010710>.
- Ribeiro AS, Schoenfeld BJ, Pina FLC, Souza MF, Nascimento MA, Santos L dos, et al. Resistance training in older women: Comparison of single vs. Multiple sets on muscle strength and body composition. *Isokinetics and Exercise Science* 2015;23:53–60. <https://doi.org/10.3233/IES-140564>.
- Correa CS, Teixeira BC, Cobos RC, Macedo RC, Kruger RL, Carteri RB, et al. High-volume resistance training reduces postprandial lipaemia in postmenopausal women. *J Sports Sci* 2015;33:1890–901. <https://doi.org/10.1080/02640414.2015.1017732>.
- Bottaro M, Veloso J, Wagner D, Gentil P. Resistance training for strength and muscle thickness: Effect of number of sets and muscle group trained. *Science & Sports* 2011;26:259–64. <https://doi.org/10.1016/j.scispo.2010.09.009>.
- Radaelli R, Fleck SJ, Leite T, Leite RD, Pinto RS, Fernandes L, et al. Dose response of 1, 3 and 5 sets of resistance exercise on strength, local muscular endurance and hypertrophy. *J Strength Cond Res* 2014. <https://doi.org/10.1519/JSC.0000000000000758>.
- Radaelli R, Wilhelm EN, Botton CE, Rech A, Bottaro M, Brown LE, et al. Effects of single vs. Multiple-set short-term strength training in elderly women. *Age (Dordr)* 2014;36:9720. <https://doi.org/10.1007/s11357-014-9720-6>.
- McBride JM, Blaak JB, Triplett-McBride T. Effect of resistance exercise volume and complexity on EMG, strength, and regional body composition. *Eur J Appl Physiol* 2003;90:626–32. <https://doi.org/10.1007/s00421-003-0930-3>.
- Starkey DB, Pollock ML, Ishida Y, Welsch MA, Brechue WF, Graves JE, et al. Effect of resistance training volume on strength and muscle thickness. *Med Sci Sports Exerc* 1996;28:1311–20. <https://doi.org/10.1097/00005768-199610000-00016>.

- [143] Ostrowski KJ, Wilson GJ, Weatherby [130] Murphy PW, Lyttle AD. The effect of weight training volume on hormonal output and muscular size and function. *Journal of Strength and Conditioning Research* 1997;11:148–54. <https://doi.org/Doi%2010.1519/00124278-199708000-00003>. [151]
- [144] Rhea MR, Alvar BA, Ball SD, Burkett LN. Three sets of weight training superior to 1 set with equal intensity for eliciting strength. *J Strength Cond Res* 2002;16:525–9. [152]
- [145] Cannon J, Marino FE. Early-phase neuromuscular adaptations to high- and low-volume resistance training in untrained young and older women. *J Sports Sci* 2010;28:1505–14. <https://doi.org/10.1080/02640414.2010.517544>. [153]
- [146] Ronnestad BR, Egeland W, Kvamme NH, Refsnes PE, Kadi F, Raastad T. Dissimilar effects of one- and three-set strength training on strength and muscle mass gains in upper and lower body in untrained subjects. *J Strength Cond Res* 2007;21:157–63. <https://doi.org/10.1519/00124278-200702000-00028>. [154]
- [147] Mitchell CJ, Churchward-Venne TA, West DW, Burd NA, Breen L, Baker SK, et al. Resistance exercise load does not determine training-mediated hypertrophic gains in young men. *J Appl Physiol* (1985) 2012;113:71–7. <https://doi.org/10.1152/japplphysiol.00307.2012>. [155]
- [148] Sooneste H, Tanimoto M, Kakigi R, Saga N, Katamoto S. Effects of training volume on strength and hypertrophy in young men. *J Strength Cond Res* 2013;27:8–13. <https://doi.org/10.1519/JSC.0b013e3182679215>. [156]
- [149] Dreyer HC, Fujita S, Cadenas JG, Chinkes DL, Volpi E, Rasmussen BB. Resistance exercise increases AMPK activity and reduces 4E-BP1 phosphorylation and protein synthesis in human skeletal muscle. *The Journal of Physiology* 2006;576:613–24. <https://doi.org/https://doi.org/10.1113/jphysiol.2006.113175>. [157]
- MacDougall JD, Gibala MJ, Tarnopolsky MA, MacDonald JR, Interisano SA, Yarasheski KE. The time course for elevated muscle protein synthesis following heavy resistance exercise. *Can J Appl Physiol* 1995;20:480–6. <https://doi.org/10.1139/h95-038>.
- Phillips SM, Tipton KD, Aarsland A, Wolf SE, Wolfe RR. Mixed muscle protein synthesis and breakdown after resistance exercise in humans. *Am J Physiol* 1997;273:E99–107.
- Biolo G, Maggi SP, Williams BD, Tipton KD, Wolfe RR. Increased rates of muscle protein turnover and amino acid transport after resistance exercise in humans. *Am J Physiol* 1995;268:E514–20. <https://doi.org/10.1152/ajpendo.1995.268.3.E514>.
- Chesley A, MacDougall JD, Tarnopolsky MA, Atkinson SA, Smith K. Changes in human muscle protein synthesis after resistance exercise. *J Appl Physiol* (1985) 1992;73:1383–8. <https://doi.org/10.1152/jappl.1992.73.4.1383>.
- Kim PL, Staron RS, Phillips SM. Fasted-state skeletal muscle protein synthesis after resistance exercise is altered with training. *J Physiol* 2005;568:283–90. <https://doi.org/10.1113/jphysiol.2005.093708>.
- Phillips SM, Tipton KD, Ferrando AA, Wolfe RR. Resistance training reduces the acute exercise-induced increase in muscle protein turnover. *Am J Physiol* 1999;276:E118–24. <https://doi.org/10.1152/ajpendo.1999.276.1.E118>.
- Børsheim E, Tipton KD, Wolf SE, Wolfe RR. Essential amino acids and muscle protein recovery from resistance exercise. *Am J Physiol Endocrinol Metab* 2002;283:E648–57. <https://doi.org/10.1152/ajpendo.00466.2001>.
- Tipton KD, Ferrando AA, Phillips SM, Doyle Jr D., Wolfe RR. Postexercise net protein synthesis in human muscle from orally administered amino acids. *Am J Physiol* 1999;276:E628–34. <https://doi.org/10.1152/ajpendo.1999.276.4.E628>.

- [158] Wilkinson SB, Phillips SM, Atherton P, Patel R, Yarasheski KE, Tarnopolsky MA, et al. Differential effects of resistance and endurance exercise in the fed state on signalling molecule phosphorylation and protein synthesis in human muscle. *J Physiol* 2008;586:3701–17. <https://doi.org/10.1113/jphysiol.2008.153916>.
- [159] Reidy PT, Borack MS, Markofski MM, Dickinson JM, Fry CS, Deer RR, et al. Post-absorptive muscle protein turnover affects resistance training hypertrophy. *Eur J Appl Physiol* 2017;117:853–66. <https://doi.org/10.1007/s00421-017-3566-4>.
- [160] Figueiredo VC, McCarthy JJ. The role of ribosome biogenesis in skeletal muscle hypertrophy. In: Sakuma K, editor. *The plasticity of skeletal muscle*, Singapore: Springer Singapore; 2017, p. 141–53. [https://doi.org/10.1007/978-981-10-3292-9\\_6](https://doi.org/10.1007/978-981-10-3292-9_6).
- [161] West DW, Baehr LM, Marcotte GR, Chason CM, Tolento L, Gomes AV, et al. Acute resistance exercise activates rapamycin-sensitive and -insensitive mechanisms that control translational activity and capacity in skeletal muscle. *J Physiol* 2016;594:453–68. <https://doi.org/10.1113/JP271365>.
- [162] Millward DJ, Garlick PJ, James WPT, Nnanyelugo DO, Ryatt JS. Relationship between protein synthesis and RNA content of skeletal muscle. *Nature* 1973;241:204. <https://doi.org/10.1038/241204a0>.
- [163] Figueiredo VC, Caldow MK, Massie V, Markworth JF, Cameron-Smith D, Blazevich AJ. Ribosome biogenesis adaptation in resistance training-induced human skeletal muscle hypertrophy. *Am J Physiol Endocrinol Metab* 2015;309:E72–83. <https://doi.org/10.1152/ajpendo.00050.2015>.
- [164] Stec MJ, Kelly NA, Many GM, Windham ST, Tuggle SC, Bamman MM. Ribosome biogenesis may augment resistance training-induced myofiber hypertrophy and is required for myotube growth in vitro. *Am J Physiol Endocrinol Metab* 2016;310:E652–61. <https://doi.org/10.1152/ajpendo.00486.2015>.
- Hammarström D, Øfsteng S, Koll L, Hanes-tadhaugen M, Hollan I, Apró W, et al. Benefits of higher resistance-training volume are related to ribosome biogenesis. *The Journal of Physiology* 2020;598:543–65. <https://doi.org/10.1113/JP278455>.
- Walden F von, Liu C, Aurigemma N, Nader GA. mTOR signaling regulates myotube hypertrophy by modulating protein synthesis, rDNA transcription and chromatin remodeling. *Am J Physiol Cell Physiol* 2016;ajp-cell 00144 2016. <https://doi.org/10.1152/ajpcell.00144.2016>.
- Warner JR. The economics of ribosome biosynthesis in yeast. *Trends Biochem Sci* 1999;24:437–40.
- Moss T, Langlois F, Gagnon-Kugler T, Stefanovsky V. A housekeeper with power of attorney: The rRNA genes in ribosome biogenesis. *Cell Mol Life Sci* 2007;64:29–49. <https://doi.org/10.1007/s00018-006-6278-1>.
- Tuan JC, Zhai W, Comai L. Recruitment of TATA-binding protein-TAFI complex SL1 to the human ribosomal DNA promoter is mediated by the carboxy-terminal activation domain of upstream binding factor (UBF) and is regulated by UBF phosphorylation. *Mol Cell Biol* 1999;19:2872–9. <https://doi.org/10.1128/mcb.19.4.2872>.
- Lin CH, Platt MD, Ficarro SB, Hoofnagle MH, Shabanowitz J, Comai L, et al. Mass spectrometric identification of phosphorylation sites of rRNA transcription factor upstream binding factor. *Am J Physiol Cell Physiol* 2007;292:C1617–24. <https://doi.org/10.1152/ajpcell.00176.2006>.
- Nader GA, McLoughlin TJ, Esser KA. mTOR function in skeletal muscle hypertrophy: Increased ribosomal RNA via cell cycle regulators. *Am J Physiol Cell Physiol* 2005;289:C1457–65. <https://doi.org/10.1152/ajpcell.00165.2005>.



- [172] Hannan KM, Brandenburger Y, Jenkins AT, Sharkey K, Cavanaugh A, Rothblum L, et al. mTOR-dependent regulation of ribosomal gene transcription requires S6K1 and is mediated by phosphorylation of the carboxy-terminal activation domain of the nuclear transcription factor UBF. *Mol Cell Biol* 2003;23:8862–77. <https://doi.org/10.1128/mcb.23.23.8862-8877.2003>.
- [173] Hannan RD, Stefanovsky V, Taylor L, Moss T, Rothblum LI. Overexpression of the transcription factor UBF1 is sufficient to increase ribosomal DNA transcription in neonatal cardiomyocytes: Implications for cardiac hypertrophy. *Proceedings of the National Academy of Sciences of the United States of America* 1996;93:8750–5. <https://doi.org/10.1073/pnas.93.16.8750>.
- [174] Sanij E, Poortinga G, Sharkey K, Hung S, Hloway TP, Quin J, et al. UBF levels determine the number of active ribosomal RNA genes in mammals. *J Cell Biol* 2008;183:1259–74. <https://doi.org/10.1083/jcb.200805146>.
- [175] Phillips SM, Parise G, Roy BD, Tipton KD, Wolfe RR, Tamopolsky MA. Resistance-training-induced adaptations in skeletal muscle protein turnover in the fed state. *Can J Physiol Pharmacol* 2002;80:1045–53. <https://doi.org/10.1139/y02-134>.
- [176] Tang JE, Perco JG, Moore DR, Wilkinson SB, Phillips SM. Resistance training alters the response of fed state mixed muscle protein synthesis in young men. *Am J Physiol Regul Integr Comp Physiol* 2008;294:R172–8. <https://doi.org/10.1152/ajpregu.00636.2007>.
- [177] Damas F, Phillips SM, Libardi CA, Vechin FC, Lixandrao ME, Jannig PR, et al. Resistance training-induced changes in integrated myofibrillar protein synthesis are related to hypertrophy only after attenuation of muscle damage. *The Journal of Physiology* 2016;594:5209–22. <https://doi.org/10.1113/JP272472>.
- Goodman CA. The role of mTORC1 in regulating protein synthesis and skeletal muscle mass in response to various mechanical stimuli. *Rev Physiol Biochem Pharmacol* 2014. [https://doi.org/10.1007/112\\_2013\\_17](https://doi.org/10.1007/112_2013_17).
- Bodine SC, Stitt TN, Gonzalez M, Kline WO, Stover GL, Bauerlein R, et al. Akt/mTOR pathway is a crucial regulator of skeletal muscle hypertrophy and can prevent muscle atrophy in vivo. *Nat Cell Biol* 2001;3:1014–9. <https://doi.org/10.1038/ncb1101-1014>.
- Kubica N, Bolster DR, Farrell PA, Kimball SR, Jefferson LS. Resistance exercise increases muscle protein synthesis and translation of eukaryotic initiation factor 2Bepsilon mRNA in a mammalian target of rapamycin-dependent manner. *J Biol Chem* 2005;280:7570–80. <https://doi.org/10.1074/jbc.M413732200>.
- Drummond MJ, Fry CS, Glynn EL, Dreyer HC, Dhanani S, Timmerman KL, et al. Rapamycin administration in humans blocks the contraction-induced increase in skeletal muscle protein synthesis. *J Physiol* 2009;587:1535–46. <https://doi.org/10.1113/jphysiol.2008.163816>.
- Terzis G, Georgiadis G, Stratakos G, Vogiatzis I, Kavouras S, Manta P, et al. Resistance exercise-induced increase in muscle mass correlates with p70S6 kinase phosphorylation in human subjects. *Eur J Appl Physiol* 2008;102:145–52. <https://doi.org/10.1007/s00421-007-0564-y>.
- Voit R, Hoffmann M, Grummt I. Phosphorylation by G1-specific cdk-cyclin complexes activates the nucleolar transcription factor UBF. *Embo j* 1999;18:1891–9. <https://doi.org/10.1093/emboj/18.7.1891>.
- Stefanovsky VY, Pelletier G, Hannan R, Gagnon-Kugler T, Rothblum LI, Moss T. An immediate response of ribosomal transcription to growth factor stimulation in mammals is mediated by ERK phosphorylation of UBF. *Molecular Cell* 2001;8:1063–73. [https://doi.org/10.1016/S1097-2765\(01\)00384-7](https://doi.org/10.1016/S1097-2765(01)00384-7).

- [185] Welle S, Bhatt K, Thornton CA. Stimulation of myofibrillar synthesis by exercise is mediated by more efficient translation of mRNA. *J Appl Physiol* (1985) 1999;86:1220–5. <https://doi.org/10.1152/jappl.1999.86.4.1220>.
- [186] Newlands S, Levitt LK, Robinson CS, Karpf AB, Hodgson VR, Wade RP, et al. Transcription occurs in pulses in muscle fibers. *Genes & Development* 1998;12:2748–58. <https://doi.org/10.1101/gad.12.17.2748>.
- [187] Kirby TJ, Patel RM, McClintock TS, Dupont-Versteegden EE, Peterson CA, McCarthy JJ. Myonuclear transcription is responsive to mechanical load and DNA content but uncoupled from cell size during hypertrophy. *Molecular Biology of the Cell* 2016;27:788–98. <https://doi.org/10.1091/mbc.E15-08-0585>.
- [188] Stec MJ, Mayhew DL, Bamman MM. The effects of age and resistance loading on skeletal muscle ribosome biogenesis. *J Appl Physiol* (1985) 2015;jap 00489 2015. <https://doi.org/10.1152/japplphysiol.00489.2015>.
- [189] Nader GA, Walden F von, Liu C, Lindvall J, Gutmann L, Pistilli EE, et al. Resistance exercise training modulates acute gene expression during human skeletal muscle hypertrophy. *J Appl Physiol* (1985) 2014;116:693–702. <https://doi.org/10.1152/japplphysiol.01366.2013>.
- [190] Bickel CS, Slade J, Mahoney E, Haddad F, Dudley GA, Adams GR. Time course of molecular responses of human skeletal muscle to acute bouts of resistance exercise. *J Appl Physiol* (1985) 2005;98:482–8. <https://doi.org/10.1152/japplphysiol.00895.2004>.
- [191] Brook MS, Wilkinson DJ, Mitchell WK, Lund JN, Phillips BE, Szewczyk NJ, et al. Synchronous deficits in cumulative muscle protein synthesis and ribosomal biogenesis underlie age-related anabolic resistance to exercise in humans. *J Physiol* 2016;594:7399–417. <https://doi.org/10.1113/JP272857>.
- Mobley CB, Haun CT, Roberson PA, Mumford PW, Kephart WC, Romero MA, et al. Biomarkers associated with low, moderate, and high vastus lateralis muscle hypertrophy following 12 weeks of resistance training. *PLoS One* 2018;13:e0195203. <https://doi.org/10.1371/journal.pone.0195203>.
- Kumar V, Atherton PJ, Selby A, Rankin D, Williams J, Smith K, et al. Muscle protein synthetic responses to exercise: Effects of age, volume, and intensity. *J Gerontol A Biol Sci Med Sci* 2012;67:1170–7. <https://doi.org/10.1093/gerona/gls141>.
- Kumar V, Selby A, Rankin D, Patel R, Atherton P, Hildebrandt W, et al. Age-related differences in the dose-response relationship of muscle protein synthesis to resistance exercise in young and old men. *J Physiol* 2009;587:211–7. <https://doi.org/10.1113/jphysiol.2008.164483>.
- Seaborne RA, Strauss J, Cocks M, Shepherd S, O'Brien TD, Someren KA van, et al. Human skeletal muscle possesses an epigenetic memory of hypertrophy. *Sci Rep* 2018;8:1898. <https://doi.org/10.1038/s41598-018-20287-3>.
- Rhea MR, Alvar BA, Burkett LN, Ball SD. A meta-analysis to determine the dose response for strength development. *Med Sci Sports Exerc* 2003;35:456–64. <https://doi.org/10.1249/01.MSS.0000053727.63505.D4>.
- Egan B, Zierath JR. Exercise metabolism and the molecular regulation of skeletal muscle adaptation. *Cell Metabolism* 2013;17:162–84. <https://doi.org/10.1016/j.cmet.2012.12.012>.
- Dirks ML, Wall BT, Valk B van de, Holloway TM, Holloway GP, Chabowski A, et al. One week of bed rest leads to substantial muscle atrophy and induces whole-body insulin resistance in the absence of skeletal muscle lipid accumulation. *Diabetes* 2016;65:2862–75. <https://doi.org/10.2337/db15-1661>.

- [199] Bamman MM, Petrella JK, Kim JS, Mayhew DL, Cross JM. Cluster analysis tests the importance of myogenic gene expression during myofiber hypertrophy in humans. *J Appl Physiol* (1985) 2007;102:2232–9. <https://doi.org/10.1152/japplphysiol.00024.2007>.
- [200] Davidson PK, Gallagher IJ, Hartman JW, Tarnopolsky MA, Dela F, Helge JW, et al. High responders to resistance exercise training demonstrate differential regulation of skeletal muscle microRNA expression. *J Appl Physiol* (1985) 2011;110:309–17. <https://doi.org/10.1152/japplphysiol.00901.2010>.
- [201] Brook MS, Wilkinson DJ, Phillips BE, Pefschner Schindler J, Philp A, Smith K, et al. Skeletal muscle homeostasis and plasticity in youth and ageing: Impact of nutrition and exercise. *Acta Physiol (Oxf)* 2016;216:15–41. <https://doi.org/10.1111/apha.12532>.
- [202] Gordon PM, Liu D, Sartor MA, Iglay Reger HB, Pistilli EE, Gutmann L, et al. Resistance exercise training influences skeletal muscle immune activation: A microarray analysis. *J Appl Physiol* (1985) 2012;112:443–53. <https://doi.org/10.1152/japplphysiol.00860.2011>.
- [203] Hyldahl RD, Xin L, Hubal MJ, Moeckel-Cole S, Chipkin S, Clarkson PM. Activation of nuclear factor- $\kappa$ B following muscle eccentric contractions in humans is localized primarily to skeletal muscle-residing pericytes. *The FASEB Journal* 2011;25:2956–66. <https://doi.org/10.1096/fj.10-177105>.
- [204] Robinson MM, Dasari S, Konopka AR, Johnson ML, Manjunatha S, Esponda RR, et al. Enhanced protein translation underlies improved metabolic and physical adaptations to different exercise training modes in young and old humans. *Cell Metabolism* 2017;25:581–92. <https://doi.org/10.1016/j.cmet.2017.02.009>.
- [205] Melov S, Tarnopolsky MA, Beckman K, Felkey K, Hubbard A. Resistance exercise reverses aging in human skeletal muscle. *PLOS ONE* 2007;2:e465. <https://doi.org/10.1371/journal.pone.0000465>.
- Murton AJ, Billeter R, Stephens FB, Des Etages SG, Graber F, Hill RJ, et al. Transient transcriptional events in human skeletal muscle at the outset of concentric resistance exercise training. *J Appl Physiol* (1985) 2014;116:113–25. <https://doi.org/10.1152/japplphysiol.00426.2013>.
- Phillips BE, Williams JP, Gustafsson T, Bouchard C, Rankinen T, Knudsen S, et al. Molecular networks of human muscle adaptation to exercise and age. *PLoS Genet* 2013;9:e1003389. <https://doi.org/10.1371/journal.pgen.1003389>.
- Hangelbroek RWJ, Fazelzadeh P, Tieland M, Boekschoten MV, Hooiveld GJEJ, Duynhoven JPM van, et al. Expression of protocadherin gamma in skeletal muscle tissue is associated with age and muscle weakness. *Journal of Cachexia, Sarcopenia and Muscle* 2016;7:604–14. <https://doi.org/10.1002/jcsm.12099>.
- STEPTO NK, COFFEY VG, CAREY AL, PONNAMPALAM AP, CANNY BJ, POWELL D, et al. Global gene expression in skeletal muscle from well-trained strength and endurance athletes. *Medicine & Science in Sports & Exercise* 2009;41:546–65. <https://doi.org/10.1249/MSS.0b013e31818c6be9>.
- Pillon NJ, Gabriel BM, Dollet L, Smith JAB, Sardón Puig L, Botella J, et al. Transcriptomic profiling of skeletal muscle adaptations to exercise and inactivity. *Nature Communications* 2020;11:470–0. <https://doi.org/10.1038/s41467-019-13869-w>.
- Tarnopolsky M, Phillips S, Parise G, Varbanov A, DeMuth J, Stevens P, et al. Gene expression, fiber type, and strength are similar between left and right legs in older adults. *The Journals of Gerontology: Series A* 2007;62:1088–95. <https://doi.org/10.1093/gerona/62.10.1088>.
- Conesa A, Madrigal P, Tarazona S, Gomez-Cabrero D, Cervera A, McPherson A, et al. A survey of best practices for RNA-seq data analysis. *Genome Biology* 2016;17:13–3. <https://doi.org/10.1186/s13059-016-0881-8>.

- [213] Robinson MD, Oshlack A. A scaling normalization method for differential expression analysis of RNA-seq data. *Genome Biology* 2010;11:R25. <https://doi.org/10.1186/gb-2010-11-3-r25>. [221]
- [214] Dillies M-A, Rau A, Aubert J, Hennequin [222] Antier C, Jeanmougin M, Servant N, et al. A comprehensive evaluation of normalization methods for illumina high-throughput RNA sequencing data analysis. *Briefings in Bioinformatics* 2013;14:671–83. <https://doi.org/10.1093/bib/bbs046>. [223]
- [215] Arora S, Pattwell SS, Holland EC, Bolouri H. Variability in estimated gene expression among commonly used RNA-seq pipelines. *Sci Rep* 2020;10:2734. <https://doi.org/10.1038/s41598-020-59516-z>. [224]
- [216] Lovén J, Orlando DA, Sigova AA, Lin CY, Rahl PB, Burge CB, et al. Revisiting global gene expression analysis. *Cell* 2012;151:476–82. <https://doi.org/10.1016/j.cell.2012.10.012>. [225]
- [217] Lin CY, Lovén J, Rahl PB, Paranal RM, Burge CB, Bradner JE, et al. Transcriptional amplification in tumor cells with elevated c-myc. *Cell* 2012;151:56–67. <https://doi.org/10.1016/j.cell.2012.08.026>. [226]
- [218] Phillips SM. A brief review of critical processes in exercise-induced muscular hypertrophy. *Sports Med* 2014;44 Suppl 1:S71–7. <https://doi.org/10.1007/s40279-014-0152-3>. [227]
- [219] Brook MS, Wilkinson DJ, Mitchell WK, Lund JN, Szewczyk NJ, Greenhaff PL, et al. Skeletal muscle hypertrophy adaptations predominate in the early stages of resistance exercise training, matching deuterium oxide-derived measures of muscle protein synthesis and mechanistic target of rapamycin complex 1 signaling. *FASEB J* 2015;29:4485–96. <https://doi.org/10.1096/fj.15-273755>. [228]
- [220] World medical association declaration of helsinki: Ethical principles for medical research involving human subjects. *Jama* 2013;310:2191–4. <https://doi.org/10.1001/jama.2013.281053>. [229]
- Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al. Fiji: An open-source platform for biological-image analysis. *Nat Methods* 2012;9:676–82. <https://doi.org/10.1038/nmeth.2019>. [230]
- Hayot M, Michaud A, Koechlin C, Caron MA, Leblanc P, Prefaut C, et al. Skeletal muscle microbiopsy: A validation study of a minimally invasive technique. *Eur Respir J* 2005;25:431–40. <https://doi.org/10.1183/09031936.05.00053404>. [231]
- Eklom B. The muscle biopsy technique. Historical and methodological considerations. *Scand J Med Sci Sports* 2017;27:458–61. <https://doi.org/10.1111/sms.12808>. [232]
- Bonafiglia JT, Islam H, Preobrazenski N, Drouin P, Ma A, Gerhart A, et al. A comparison of pain responses, hemodynamic reactivity and fibre type composition between bergström and microbiopsy skeletal muscle biopsies. *Current Research in Physiology* 2020;3:1–0. <https://doi.org/https://doi.org/10.1016/j.crphys.2020.05.001>. [233]
- Hughes MC, Ramos SV, Turnbull PC, Nejatbakhsh A, Baechler BL, Tahmasebi H, et al. Mitochondrial bioenergetics and fiber type assessments in microbiopsy vs. Bergstrom percutaneous sampling of human skeletal muscle. *Frontiers in Physiology* 2015;6. <https://doi.org/10.3389/fphys.2015.00360>. [234]
- Blomstrand E, Eklom B. The needle biopsy technique for fibre type determination in human skeletal muscle—a methodological study. *Acta Physiol Scand* 1982;116:437–42. <https://doi.org/10.1111/j.1748-1716.1982.tb07163.x>. [235]
- Chomczynski P, Sacchi N. Single-step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction. *Analytical Biochemistry* 1987;162:156–9. [https://doi.org/https://doi.org/10.1016/0003-2697\(87\)90021-2](https://doi.org/https://doi.org/10.1016/0003-2697(87)90021-2). [236]

- [228] Sennepin AD, Charpentier S, Normand[285] Sarre C, Legrand A, Mollet LM. Multiple reprobing of western blots after inactivation of peroxidase activity by its substrate, hydrogen peroxide. *Anal Biochem* 2009;393:129–31. <https://doi.org/10.1016/j.ab.2009.06.004>.
- [229] Rueden CT, Schindelin J, Hiner MC, DeZonia BE, Walter AE, Arena ET, et al. ImageJ2: ImageJ for the next generation[237] scientific image data. *BMC Bioinformatics* 2017;18:529. <https://doi.org/10.1186/s12859-017-1934-z>.
- [230] Kopec AM, Rivera PD, Lacagnina MJ, Han[238] sagar R, Bilbo SD. Optimized solubilization of TRIzol-precipitated protein permits western blotting analysis to maximize data available from brain tissue. *Journal of Neuroscience Methods* 2017;280:64–76. <https://doi.org/10.1016/j.jneumeth.2017.02.002>.
- [231] Ellefsen S, Stenslokken KO, Sandvik GK, Kristensen TA, Nilsson GE. Improved normalization of real-time reverse transcriptase polymerase chain reaction data using an external RNA control. *Anal Biochem* 2008;376:83–93. <https://doi.org/10.1016/j.ab.2008.01.028>.
- [232] Ellefsen S, Vikmoen O, Slettalokken G, Whist JE, Nygaard H, Hollan I, et al. Irisin and FNDC5: Effects of 12-week strength training, and relations to muscle phenotype and body mass composition in untrained women. *Eur J Appl Physiol* 2014;114:1875–88. <https://doi.org/10.1007/s00421-014-2922-x>.
- [233] Ye J, Coulouris G, Zaretskaya I, Cutcutache I, Rozen S, Madden TL. Primer-BLAST: A tool to design target-specific primers for polymerase chain reaction. *BMC Bioinformatics* 2012;13:134. <https://doi.org/10.1186/1471-2105-13-134>.
- [234] Untergasser A, Cutcutache I, Koressaar T, Ye J, Faircloth BC, Remm M, et al. Primer3—new capabilities and interfaces. *Nucleic Acids Res* 2012;40:e115. <https://doi.org/10.1093/nar/gks596>.
- Ritz C, Spiess AN. qpcR: An r package for sigmoidal model selection in quantitative real-time polymerase chain reaction analysis. *Bioinformatics* 2008;24:1549–51. <https://doi.org/10.1093/bioinformatics/btn227>.
- Tichopad A, Dilger M, Schwarz G, Pfaffl MW. Standardized determination of real-time PCR efficiency from a single reaction set-up. *Nucleic Acids Res* 2003;31:e122.
- Fleige S, Pfaffl MW. RNA integrity and the effect on the real-time qRT-PCR performance. *Mol Aspects Med* 2006;27:126–39. <https://doi.org/10.1016/j.mam.2005.12.003>.
- Bürkner P-C. Brms: An r package for bayesian multilevel models using stan. 2017 2017;80:28. <https://doi.org/10.18637/jss.v080.i01>.
- Bland M. An introduction to medical statistics. Fourth edition. Oxford ; Oxford University Press; 2015.