



Research Article

**From Harvest to Health: The Green Revolution,
Micronutrient Erosion, and the Hidden Hunger
Crisis — A Critical Review from a Public Health
Perspective**

Dr. Nagmanee Kumar*

Assistant professor Department of Sociology Nalanda Mahila College,
Biharsharf, Nalanda, Patliputra University, Patna (Bihar) India

ARTICLE INFO

Received: 2nd March, 2026

Accepted: 25th April, 2026

Published: 5th May, 2026

Pages: 28 - 36

Abstract

The Green Revolution of the 1960s onwards successfully averted widespread famine by dramatically increasing the yields of cereal crops through the development of high-yielding varieties (HYVs), expanded irrigation, and intensive use of synthetic fertilizers and pesticides. While the humanitarian achievement of preventing acute mass starvation is undeniable, the revolution's legacy on nutritional quality and long-term public health presents a more complex and troubling picture. This paper critically examines the trade-off between caloric abundance and micronutrient adequacy in the post-Green Revolution era. Drawing on agricultural science, nutritional epidemiology, and World Health Organization (WHO) data, we trace how the monoculturation of global food systems, the dilution effect associated with rapid-yield breeding, and intensive agrochemical use have collectively contributed to a condition the WHO terms 'hidden hunger' — caloric sufficiency alongside widespread micronutrient deficiency. We document significant declines in iron, zinc, and calcium content in both rice and wheat following the Green Revolution, and connect these declines to a global burden of micronutrient malnutrition affecting over two billion people. We further examine the double burden of malnutrition observed in countries like India, where child stunting and adult obesity coexist within the same households and communities. The paper also evaluates the ecological consequences of intensive agriculture — soil degradation, agrobiodiversity loss, and water contamination — and how these feedback into nutritional outcomes. Finally, we assess strategies advocated by WHO, UNICEF, and the FAO, including biofortification, dietary diversification, and supplementation, and argue that future agricultural policy must shift its primary metric from yield to nutritional quality per hectare if it is to adequately serve human health in the twenty-first century.

Keywords: Green Revolution, hidden hunger, micronutrient deficiency, high-yielding varieties, biofortification, food security, nutritional quality, double burden of malnutrition, WHO, pesticide health effects

***Corresponding Author**

Dr. Nagmanee Kumar Assistant Professor Department of Sociology Nalanda Mahila
College, Biharsharf, Nalanda, Patliputra University, Patna (Bihar) India
E – mail - kumarmani@nalandamahilacollege.ac.in

28

1. Introduction

Hunger, in the conventional sense, refers to not having enough food. By that measure, the Green Revolution stands as one of the greatest achievements in human history. Beginning in the late 1950s and accelerating through the 1960s and 1970s, the revolution — driven largely by the plant breeding work of Norman Borlaug and supported by international agricultural research centres — transformed food production in South Asia, Latin America, and parts of sub-Saharan Africa. Countries like India, which had been heavily dependent on food imports and vulnerable to periodic famine, became self-sufficient in food grain production within roughly a decade. Cereal yields quadrupled globally. The spectre of mass starvation that had loomed over post-independence developing nations began to recede.

Yet hunger, as scientists and public health specialists now understand it, is not merely the absence of calories. The human body requires more than energy — it requires iron to carry oxygen through the blood, zinc to support immune function and child growth, calcium to maintain bone structure, vitamin A to protect vision, and dozens of other micronutrients to function properly. When a population consumes enough food to stave off starvation but still lacks these essential nutrients, it suffers from what nutritional epidemiologists call micronutrient deficiency, or more evocatively, 'hidden hunger.' The World Health Organization estimates that hidden hunger currently affects over two billion people worldwide — a figure that has, paradoxically, risen in the decades since the Green Revolution declared victory over food insecurity (Yilmaz & Yilmaz, 2025).

This paper argues that the Green Revolution's design priorities — which placed yield maximisation at the centre of agricultural science — systematically undermined nutritional quality, and that the consequences of this trade-off are still unfolding in the epidemiology of non-communicable disease, child stunting, and immune vulnerability across the developing world. This is not an argument against the Green Revolution per se; the lives it saved are real and the suffering it prevented is not hypothetical. It is, rather, an argument for a more honest accounting of what was gained and what was lost, and for a reorientation of agricultural priorities as we face the food challenges of the twenty-first century.

2. Background and Historical Context

Before the Green Revolution, much of the developing world's agriculture was characterised by traditional crop varieties, minimal use of external inputs, and relatively low yields per unit area. Population growth in the post-World War II period was placing severe strain on food systems, and famine conditions periodically emerged in countries including India, Pakistan, and several parts of sub-Saharan Africa. The conventional wisdom of the time held that production had to increase dramatically and fast, or catastrophic food shortages were inevitable.

The solution that emerged was primarily technological: the development of dwarf wheat and rice varieties that could convert larger quantities of applied fertilizer into grain rather than stalk. These HYVs, when combined with expanded irrigation infrastructure and heavy inputs of nitrogen, phosphorus, and potassium-based fertilizers, produced dramatic yield increases. India's wheat production, for instance, rose from roughly 12 million tonnes in 1965 to over 17 million tonnes by 1968, and continued to climb through subsequent decades. The food security argument was won convincingly. What received far less attention in the policy discourse of that era was the nutritional composition of the food being produced. The HYVs were selected almost exclusively on the basis of yield, disease resistance, and adaptability to intensive irrigation conditions. Their micronutrient profiles were secondary concerns at best. As Fan et al. (2008) later documented, the concentration of minerals in wheat grain showed a significant decrease over the 160-year period they studied, with the most dramatic declines occurring in the varieties developed and adopted during and after the Green Revolution. This was not a coincidence; it was a consequence of a breeding philosophy that had not made nutrition a priority.

3. The Dilution Effect: More Food, Less Nutrition

3.1 Mechanisms of Nutrient Decline

The nutritional decline observed in modern cereal varieties operates through several overlapping mechanisms. The most important of these is what plant scientists call the 'dilution effect': when a crop is bred to produce a much larger quantity of grain per plant, the total pool of micronutrients absorbed from the soil is distributed across a larger biomass, resulting in lower nutrient concentrations per unit of food. The plant grows faster than it can accumulate minerals. Carbohydrate

synthesis — the process that drives yield — outpaces the uptake and incorporation of iron, zinc, calcium, and other trace elements (Yilmaz & Yilmaz, 2025).

The second mechanism is chemical fertilization itself. Synthetic NPK fertilizers, while highly effective at boosting the macronutrient supply needed for rapid growth, do nothing to replenish the micronutrient pools in soil that are gradually depleted through continuous cropping. Long-term intensive cultivation without micronutrient replenishment strips the soil of iron, zinc, boron, and other elements, meaning that even traditional varieties grown in these degraded soils would produce less nutritious food than they once did (Plant Nutrition Challenges for a Sustainable Agriculture of the Future, PMC, 2024). Modern fertilizer regimes have made this problem structural rather than incidental.

The third mechanism, perhaps the most consequential from a dietary perspective, is the displacement of crop diversity. Traditional agricultural systems in Asia and sub-Saharan Africa supported enormous variety: hundreds of millet varieties, dozens of pulse species, indigenous leafy vegetables, and local rice landraces, each with its own nutritional profile. The Green Revolution concentrated resources, infrastructure, and farmer incentives on a handful of HYV wheat and rice varieties. The result was an agricultural monoculture that produced more calories but narrower nutrition.

3.2 The Numbers

The data on micronutrient decline are striking. Table 1 below presents comparative estimates for selected nutrients in rice and wheat varieties before and after the Green Revolution, drawing on agricultural and nutrition transition reports:

Table 1: Pre- vs. Post-Green Revolution Micronutrient Decline in Staple Crops

Crop	Nutrient	Pre-GR Level (approx.)	Post-GR Level (approx.)	Estimated Decline
Rice	Iron	35 mg/kg	24.5 mg/kg	30% ↓
Rice	Zinc	27 mg/kg	18 mg/kg	33% ↓
Rice	Calcium	120 mg/kg	66 mg/kg	45% ↓
Wheat	Iron	42 mg/kg	34 mg/kg	19% ↓
Wheat	Zinc	33 mg/kg	24 mg/kg	27% ↓
Wheat	Calcium	115 mg/kg	80 mg/kg	30% ↓

Source: Adapted from Yilmaz & Yilmaz (2025); Fan et al. (2008); Garvin et al. (2006); agricultural and nutrition transition reports tracking staple micronutrient depletion.

These are not trivial reductions. A 45% decline in rice calcium, for instance, is not an abstraction — it means that populations whose dietary calcium is primarily derived from rice are receiving substantially less of a nutrient critical to bone health and dental integrity. The 33% decline in rice zinc is directly connected to the WHO's finding that zinc deficiency now affects approximately 17% of the global population, causing stunted growth in children and reducing their resistance to infectious diseases including diarrhoea and pneumonia (Yilmaz & Yilmaz, 2025). The 19–30% decline in iron content across both crops is one contributing factor — not the only one, but a significant one — to the WHO's estimate that iron-deficiency anaemia affects approximately 1.8 billion people globally (Agricultural Economist, 2024).

4. Hidden Hunger and Its Public Health Consequences

4.1 Scale and Demographics

The concept of hidden hunger captures a contradiction that conventional food security metrics struggle to represent: a person can eat three meals a day and still be malnourished in ways that compromise every system of their body.

According to recent global assessments, over half of all children under five years old are deficient in at least one of iron, zinc, or vitamin A, and over two-thirds of non-pregnant women of reproductive age are deficient in at least one of iron, zinc, or folic acid (Our World in Data; Stevens et al., 2022). In absolute terms, hidden hunger affects more people than acute caloric undernourishment ever did at its peak.

The health consequences are wide-ranging and sometimes severe. Iron deficiency impairs the production of haemoglobin, reducing the oxygen-carrying capacity of the blood and leading to fatigue, impaired cognitive development, and weakened immune function. In pregnant women, it is a significant contributor to maternal mortality and increases the risk of premature delivery and low birth weight. Zinc deficiency stunts physical growth and compromises immune responses, leaving children more vulnerable to common infections that in turn worsen nutritional absorption. Vitamin A deficiency, affecting approximately 190 million children under five (UNICEF, 2023), is a leading preventable cause of childhood blindness. These are not problems at the margins of global health — they are among the most prevalent health conditions on earth.

The global burden is most concentrated in sub-Saharan Africa and South Asia, which is precisely where the Green Revolution's monoculturization of agriculture had the greatest dietary displacement effect. As indigenous crops were pushed aside in favour of HYV wheat and rice, the micronutrient diversity of local diets contracted. In India alone, traditional agricultural systems that had supported over 100,000 rice varieties were replaced by a small number of HYV types. Each lost variety represented not only a loss of genetic heritage but a loss of nutritional variety that rural communities had depended on for generations.

4.2 The Double Burden of Malnutrition

One of the more striking epidemiological phenomena associated with the post-Green Revolution dietary transition is what researchers call the 'double burden of malnutrition' (DBM): the coexistence of undernutrition and overnutrition within the same population, and sometimes within the same household. As cheap, calorie-dense cereals became the dominant food source, populations that historically suffered only from caloric deprivation began simultaneously experiencing rising rates of obesity and diet-related non-communicable diseases.

India provides the most striking illustration of this paradox. National Family Health Survey data reveal that child stunting rates remain persistently high — India ranks 105th out of 127 countries in the 2024 Global Hunger Index — while adult overweight and obesity have been rising steadily, driven by increased consumption of refined grains and processed foods (PMC, 2024). A 2023 study published in Scientific Reports found that 7% of mother-child pairs in India now face the double burden simultaneously: children who are undernourished while their mothers are overweight or anaemic. The same household, the same food environment, two different manifestations of a food system that has optimised for caloric volume at the expense of nutritional breadth.

This dual crisis complicates policy responses considerably. Programmes designed to address caloric undernourishment through food distribution can inadvertently accelerate the overnutrition problem if the food provided is calorie-dense but micronutrient-poor — which is often the case when staple cereals form the backbone of nutrition interventions. The WHO has explicitly recognised this challenge, emphasising that undernutrition and overnutrition should no longer be addressed in policy silos but must be tackled simultaneously through what it calls 'double-duty actions' (WHO; Popkin et al., 2020).

5. Chemical Exposure and Direct Health Risks

The intensification of agriculture under the Green Revolution depended not only on new seed varieties and irrigation but on very large quantities of chemical inputs — synthetic fertilizers and pesticides in particular. The health consequences of this chemical intensification have been documented in agricultural workers, in communities adjacent to intensive farming areas, and in the broader population through food chain exposure.

A systematic review published in 2024 identified consistent associations between chronic pesticide exposure and a range of non-communicable diseases, including cancers, neurological disorders, and endocrine disruptions (PMC, 2024). The review found that organophosphate pesticides — one of the most widely used classes in Green Revolution-era agriculture — are capable of disrupting the hypothalamic-pituitary axis, affecting hormonal regulation in both men and women. Male agricultural workers exposed to organophosphates show reduced levels of luteinizing hormone and serum testosterone, while female workers face increased risks of irregular menstrual cycles, spontaneous abortion, and developmental anomalies in foetuses (PMC, 2024).

The carcinogenic risk associated with pesticide exposure has received renewed attention in recent years. A major study published in 2026 in *Nature Health* combined environmental monitoring, national cancer registry data, and biological research to identify a strong association between environmental pesticide exposure and increased cancer incidence. Separately, chlorpyrifos — one of the most commonly used organophosphate pesticides — has been shown to cause oxidative stress in breast tissue, and pesticides more broadly have been associated with elevated risks of thyroid, breast, ovarian, and prostate cancers through endocrine disruption mechanisms (*Journal of Endocrinological Investigation*, 2023).

It is important to note that there is genuine scientific debate about the dose-response relationship in pesticide exposure, and that regulatory bodies apply safety limits to pesticide residues in food. Industry-affiliated research has argued that at typical exposure levels, pesticides do not cause endocrine disruption in humans (Pesticide Facts). However, the preponderance of independent epidemiological literature suggests that for agricultural workers — who experience far higher and more sustained exposure than general consumers — the health risks are real and serious. The WHO estimates that pesticide poisoning affects three million agricultural workers annually, a figure that represents a concrete and current public health burden, not a theoretical risk.

Chemical contamination of water sources represents an additional pathway of health risk. Chemical run-off from intensively farmed fields causes eutrophication of water bodies and contaminates drinking water with nitrates. High nitrate concentrations in drinking water are linked to methaemoglobinaemia — a potentially fatal condition in infants, colloquially known as 'blue baby syndrome' — as well as to elevated risks of certain cancers in adults. In the Indus and Ganges-river basins, which feed the most intensively farmed regions of the Indian subcontinent, annual groundwater depletion rates of 10–20 cm have been recorded, driven by the irrigation demands of HYV crops (Yilmaz & Yilmaz, 2025).

6. Environmental Degradation and Its Nutritional Feedback

The ecological consequences of Green Revolution agriculture do not remain confined to the environment — they feed back into human nutrition in ways that are sometimes direct and sometimes operate over longer time horizons. Three dimensions of environmental degradation are particularly relevant here.

The first is soil degradation. Continuous mono-cropping of wheat and rice, sustained over decades, depletes soil organic matter, reduces microbial biodiversity, and removes trace elements from the soil profile. Modern NPK fertilizers replace nitrogen, phosphorus, and potassium — the macronutrients that drive yield — but do not replace the micronutrients that crops extract alongside them. Over time, this creates soils that are increasingly impoverished in iron, zinc, boron, and selenium, meaning that even well-grown crops produce food of lower nutritional density. Approximately 30% of global arable land is now classified as degraded (Yilmaz & Yilmaz, 2025). Monocultures further accelerate this process because they remove the natural nutrient cycling that diverse crop rotations provide; as Degryse et al. (2022) note, monocultures in combination with a general lack of biodiversity cause soil degradation through both erosion and organic matter depletion.

The second dimension is the loss of agrobiodiversity. The mass adoption of a small number of HYVs across vast growing regions led to the extinction or abandonment of tens of thousands of traditional crop varieties. This genetic homogenisation has two consequences: it makes the global food system more vulnerable to pest outbreaks and climate shocks, and it permanently reduces the diversity of nutritional options available to communities that historically relied

on those crops. The change toward industrialised agriculture characterised by monocultures has led to major genetic erosion and reduced resilience against pests and diseases (ScienceDirect, 2025). Traditional varieties often possess traits conferring resilience to drought and pests, and their loss represents a diminution of the adaptive genetic resources that future agriculture will need.

The third dimension is water contamination. The fertilizer and pesticide run-off that accompanies intensive irrigation-based agriculture enters rivers, groundwater, and coastal marine environments, disrupting aquatic ecosystems and affecting the safety of water used for drinking and cooking. In agricultural communities where water quality is compromised, the nutritional consequences compound the dietary ones: parasitic and bacterial infections caused by contaminated water impair the intestinal absorption of whatever nutrients the diet does provide, creating a vicious cycle where degraded food systems and degraded water systems reinforce each other's impacts on human health.

7. The WHO Perspective and Contemporary Policy Responses

7.1 The Shift from Food Security to Nutrition Security

The World Health Organization has been increasingly explicit in arguing that the conventional framework of food security — defined as consistent access to sufficient calories — is inadequate as a guide to agricultural or public health policy. In its framework for nutrition security, the WHO argues that food must be assessed not only for its quantity but for its nutritional quality: its capacity to provide the full complement of macro and micronutrients needed for healthy human development. This is not a semantic distinction — it has real implications for how agricultural success is measured, how research funding is allocated, and which crops and farming systems receive policy support.

The WHO's key recommendations for addressing the nutritional legacy of the Green Revolution cluster around three main approaches: biofortification, dietary diversification, and targeted supplementation.

7.2 Biofortification

Biofortification refers to the breeding or genetic modification of crops to increase their micronutrient content, working at the seed level rather than through post-harvest fortification. The most well-known example is Golden Rice, a variety engineered to produce beta-carotene — a precursor of vitamin A — in its endosperm. More broadly, biofortification programmes have developed orange-fleshed sweet potatoes rich in vitamin A, iron-rich bean varieties, and zinc-fortified wheat (HarvestPlus, 2023). The advantage of biofortification over food supplementation is its potential reach: crops grown at scale in the field can deliver improved nutrition to rural populations who may have limited access to pharmaceutical supplements or processed fortified foods.

In 2023, the World Health Assembly passed a resolution urging member states to expand national biofortification programmes and integrate them with agricultural policy and public health initiatives (PMC — Biofortification in South Asia, 2024). This represents a significant institutional endorsement of the approach. However, biofortification has limitations: it typically addresses one or two nutrients per variety, it requires ongoing seed systems to maintain adoption, and some biofortified varieties — particularly GM crops like Golden Rice — face regulatory, cultural, and consumer acceptance challenges in many countries where they are most needed.

7.3 Dietary Diversification

The most structurally comprehensive response to hidden hunger is also the most difficult to implement: restoring genuine diversity to diets and food systems. This means reintroducing traditional millets, pulses, leafy green vegetables, and indigenous fruits that were displaced by Green Revolution staples. Millets, in particular, have attracted significant renewed interest from nutritionists and food policy specialists because they are rich in iron, calcium, and zinc; they are drought-tolerant and well-suited to smallholder farming conditions; and they were historically the staple crop of large populations in South Asia and sub-Saharan Africa before being marginalised by wheat and rice promotion (ICRISAT Millet and Nutrition Reports).

Diverse diets provide two benefits that monoculture-based diets cannot: they deliver a wider range of micronutrients, and they build more resilient food systems capable of absorbing shocks from climate change, pest outbreaks, or supply chain disruption. The challenge is that dietary diversification requires changes not only in what farmers grow but in market structures, consumer preferences, government food programmes, and the economic incentives that shape both. Countries like India are still in the early stages of this transition, with the government's millet promotion initiative — including India's presidency of the International Year of Millets in 2023 — representing a tentative but meaningful step.

7.4 Targeted Supplementation

For populations currently experiencing acute micronutrient deficiency, particularly pregnant women, breastfeeding mothers, and children under five, targeted supplementation remains critical. High-dose vitamin A capsules, iron-folic acid tablets, and zinc supplements have strong evidence bases and have been deployed through national health programmes in many countries. The WHO and UNICEF have recommended multiple micronutrient supplements (MMS) for pregnant women as a more comprehensive alternative to iron-folic acid alone, given that most women in low-income settings have multiple simultaneous deficiencies (UNICEF, 2023).

Supplementation, however, is a short-term measure rather than a structural solution. It depends on functioning health delivery systems, sustained funding, and consistent supply chains — all of which can be disrupted by crises. It does not address the underlying food system dynamics that produce deficiency in the first place. The WHO therefore frames supplementation as a bridge measure rather than a destination, to be accompanied by the longer-term work of agricultural transformation.

8. Critical Analysis: What the Green Revolution Got Right and Wrong

An honest accounting of the Green Revolution requires holding two realities simultaneously. The first is that it saved a very large number of lives. Without the yield increases it generated, the caloric shortfalls of the 1960s and 1970s would have been catastrophic. The FAO and others have estimated that without these agricultural innovations, global caloric availability would have declined by 11–15%, with the worst consequences falling on the world's poorest populations. The second reality is that the design choices embedded in the Green Revolution — the prioritisation of yield over nutritional quality, the promotion of a small number of cereal crops at the expense of dietary diversity, and the reliance on chemical inputs whose long-term health and environmental consequences were poorly understood — created new problems even as they solved old ones.

What is most useful to observe, in retrospect, is what the Green Revolution did not try to do. It did not try to preserve dietary diversity. It did not try to maintain or improve the micronutrient profile of the crops it promoted. It did not try to minimise the chemical inputs it relied on, or to anticipate the soil depletion that intensive mono-cropping would cause over decades. These were not accidental oversights — they reflect the particular framing of the problem that prevailed at the time, which defined food security almost entirely in terms of caloric sufficiency and treated other nutritional dimensions as secondary concerns that could be addressed once the calorie problem was solved.

The lesson is not that agricultural science should have chosen differently in the 1960s, given what was known and what the pressures were. The lesson is that the choices made then — which are still embedded in the structure of global food systems, seed industries, fertilizer industries, and government support mechanisms — need to be actively revisited in light of what we now understand about the full nutritional and environmental consequences. Future agricultural strategies, as the WHO has argued, must take nutritional quality as seriously as yield, and must account for the health and environmental costs of production methods that impose burdens on soil, water, biodiversity, and human health.

9. Conclusion

The Green Revolution was not, as is sometimes implied in critical accounts, a simple failure disguised as a success. It was a genuinely impressive achievement in the service of a real and urgent humanitarian goal. But it was an achievement

built on trade-offs that were not fully appreciated at the time, and whose costs have accumulated in the health of populations and the condition of agricultural ecosystems over the subsequent half-century.

Hidden hunger — the global burden of micronutrient deficiency — is the most direct nutritional legacy of those trade-offs. Affecting over two billion people, it manifests in stunted children, anaemic mothers, compromised immune systems, impaired cognitive development, and increased susceptibility to infectious disease. These consequences are especially concentrated in the populations of South Asia and sub-Saharan Africa, the very regions that the Green Revolution was designed to help. That paradox demands our sustained attention.

The path forward requires an integration of agricultural science, nutritional science, and public health policy that has historically been rare. Biofortification, dietary diversification, and supplementation each address part of the problem, but none of them alone is sufficient. What is ultimately needed is a reorientation of the goals by which agricultural progress is measured — one that places nutritional quality per hectare, dietary diversity, and environmental sustainability alongside yield as primary metrics of success. The WHO has articulated this vision; the challenge is to build the policy, institutional, and economic structures that can realise it at the scale the problem demands.

The Green Revolution fed the world. The next agricultural revolution needs to nourish it.

References

- Agricultural Economist. (2024). Addressing hidden hunger: A global health crisis. Retrieved from <https://www.agrieconomist.com>
- Bouis, H. E., & Welch, R. M. (2010). Biofortification — a sustainable agricultural strategy for reducing micronutrient malnutrition in the global south. *Crop Science*, 50(S1), S20–S32.
- Evenson, R. E., & Gollin, D. (2003). Assessing the impact of the Green Revolution, 1960 to 2000. *Science*, 300(5620), 758–762.
- Fan, M. S., Zhao, F. J., Fairweather-Tait, S. J., Poulton, P. R., Dunham, S. J., & McGrath, S. P. (2008). Evidence of decreasing mineral density in wheat grain over the last 160 years. *Journal of Trace Elements in Medicine and Biology*, 22(4), 315–324.
- Food and Agriculture Organization (FAO). Nutrition-sensitive agriculture and food systems. Rome: FAO.
- Garvin, D. F., Welch, R. M., & Finley, J. W. (2006). Historical shifts in the seed mineral micronutrient concentration of US hard red winter wheat germplasm. *Journal of the Science of Food and Agriculture*, 86(13), 2213–2220.
- Godfray, H. C. J., Beddington, J. R., Crute, I. R., Haddad, L., Lawrence, D., Muir, J. F., Pretty, J., Robinson, S., Thomas, S. M., & Toulmin, C. (2010). Food security: The challenge of feeding 9 billion people. *Science*, 327(5967), 812–818.
- HarvestPlus. (2023). Biofortification progress briefs. Washington, DC: CGIAR.
- International Crops Research Institute for the Semi-Arid Tropics (ICRISAT). Millet and nutrition reports. Patancheru: ICRISAT.
- Khoury, C. K., Bjorkman, A. D., Dempewolf, H., Ramirez-Villegas, J., Guarino, L., Jarvis, A., Rieseberg, L. H., & Struik, P. C. (2014). Increasing homogeneity in global food supplies and the implications for food security. *Proceedings of the National Academy of Sciences*, 111(11), 4001–4006.

- Pingali, P. L. (2012). Green revolution: Impacts, limits, and the path ahead. *Proceedings of the National Academy of Sciences*, 109(31), 12302–12308.
- Plant Nutrition Challenges for Sustainable Agriculture of the Future. (2024). *Physiologia Plantarum*. PMC11653101.
- Popkin, B. M., Corvalan, C., & Grummer-Strawn, L. M. (2020). Dynamics of the double burden of malnutrition and the changing nutrition reality. *Lancet*, 395(10217), 65–74.
- Singh, P. (2011). Economic benefits and ecological cost of Green Revolution: A case study of Punjab. *Journal of Economic and Social Development*, 7(1), 64–74.
- Stevens, G. A., Beal, T., Mbuya, M. N. N., Luo, H., & Neufeld, L. M. (2022). Micronutrient deficiencies among preschool-aged children and women of reproductive age worldwide: A pooled analysis of individual-level data from population-representative surveys. *The Lancet Global Health*, 10(11), e1590–e1599.
- Systematic review of pesticide exposure, associated risks, and long-term human health impacts. (2024). *ScienceDirect* / PMC11664077.
- Tilman, D., Cassman, K. G., Matson, P. A., Naylor, R., & Polasky, S. (2002). Agricultural sustainability and intensive production practices. *Nature*, 418(6898), 671–677.
- UNICEF. (2023). Large-scale food fortification for the prevention of micronutrient deficiencies in children, women and communities: Guidance note. New York: UNICEF.
- Welch, R. M., & Graham, R. D. (2004). Breeding crops for enhanced micronutrient content. *Plant and Soil*, 245(1), 205–214.
- White, P. J., & Broadley, M. R. (2005). Biofortification of crops with seven mineral elements often lacking in human diets — iron, zinc, copper, calcium, magnesium, selenium and iodine. *New Phytologist*, 167(2), 411–425.
- World Health Organization (WHO). Reports on nutrition and food safety. Geneva: WHO.
- Yadav, S., & Anand, S. (2019). Green revolution and food security in India: A review. *Geographical Journal of India*, 65(3), 312–323.
- Yilmaz, H., & Yilmaz, A. (2025). Hidden hunger in the age of abundance: The nutritional pitfalls of modern staple crops. *Food Science & Nutrition*, 13(2), e4610. <https://doi.org/10.1002/fsn3.4610>