Smart tumor-targeted microcapsule logic gates

"SMART TUMOR TARGETED MICROCAPSULE LOGIC GATES"

Generated by LeapSpace, Sat Oct 10 2026

Smart Tumor-Targeted Microcapsule Logic Gates at a Glance

Key Findings Table

Topic	Key finding	Population/model or system focus	Evidence
Core concept	Smart tumor-targeted microcapsules use tumor-associated inputs such as acidic pH, redox/GSH, hypoxia, enzymes, and sometimes ATP to control activation and payload release.	Capsule-, nanocapsule-, and related gated delivery systems for cancer therapy	[1] [2] [3] [4] [5] [6]
Best-validated inputs	pH and redox/GSH are the most directly validated triggers for release behavior in capsule-like systems.	Release assays in acidic or reductive tumor-mimicking conditions	[7] [8] [9] [10] [11] [12] [13] [14] [15] [16] [17] [18]
Dominant logic design	AND and sequential AND architectures are the main logic format, typically using one cue for priming and another for final intracellular release.	Dual- and multi-responsive cancer nanocarriers/microcapsules	[13] [19] [20] [21] [22] [23] [24] [25]
Materials matter	Logic behavior depends strongly on shell architecture, crosslinking, multilayers, and microfluidic fabrication, not just on the trigger chemistry.	Double-emulsion, multilayer, polymersome, LbL, core-shell systems	[26] [27] [28] [29] [30] [31] [32] [33] [34]
Payload trend	The field is shifting from single-drug delivery toward coordinated chemo-immunotherapy and chemo-phototherapy.	DOX-centered combinations, STING agonists, GM-CSF, metformin, photosensitizers	[35] [36] [37] [38] [39] [40] [41] [42] [43]
In vivo promise	Responsive capsule-like systems can improve tumor accumulation, imaging-guided localization, efficacy, and safety relative to free drug or nonresponsive controls.	Mouse solid tumor models	[44] [45] [46] [47] [48] [49] [50] [51]
Main gap	Translational performance remains constrained by EPR heterogeneity, dense ECM, abnormal vasculature, premature activation risk, and manufacturing complexity.	Preclinical-to-clinical translation of solid-tumor delivery systems	[52] [53] [54] [55] [56] [57] [58] [59] [60] [61]

Direct Answer

Smart tumor-targeted microcapsule logic gates are feasible and increasingly sophisticated, but the evidence base shows a clear pattern: the most practical systems are not abstract Boolean devices so much as **programmable, multistage release capsules** that integrate tumor cues into conditional transport and release behavior [3] [5] [6] [21] [22] [38] [62]. The strongest current designs use **pH plus redox/GSH**, or **pH plus hypoxia/enzyme**, in **AND or sequential AND architectures** so that one signal first exposes, charges, swells, or destabilizes the capsule, and a second signal completes drug release inside tumor tissue or cells [9] [13] [23] [24].

If the goal is tumor selectivity, the supplied evidence supports prioritizing:

1. **Dual-input or cascade gating over single-trigger designs** for lower false activation and better intracellular release control [19] [20] [21] [22].
2. **Microfluidic core-shell or multilayer capsule fabrication** for precise shell thickness, monodispersity, and staged permeability control [27] [30] [31] [32].
3. **pH- and redox-responsive chemistries** because they have the clearest assay-level validation in the provided literature [15] [16] [17] [18].
4. **Combination payloads** where logic control helps synchronize chemotherapy with immune or phototherapeutic effects [39] [40] [42].

The main caveat is that preclinical success does **not** remove the classic delivery bottleneck: tumor penetration, heterogeneous EPR, and scale-up remain major barriers even for logic-gated systems [57] [58] [59] [60] [61].

Confidence level: High - the central patterns are repeatedly supported across the supplied reviews, trigger-validation studies, engineering papers, and in vivo reports [3] [19] [27] [35] [55].

Study Scope

This report covers tumor-targeted smart microcapsule and closely related nanocapsule/gated carrier literature reflected in the supplied evidence, spanning foundational work and recent studies through roughly 2018-2025, with emphasis on tumor microenvironment sensing, logic architectures, fabrication, release behavior, and preclinical in vivo performance [34] [47] [63] [64]. The source base includes mechanistic release assays, engineering/fabrication studies, foundational reviews, and mouse tumor-model efficacy and safety studies [15] [27] [51] [65].

Assumptions & Limitations

The evidence is strongest for **pH** and **redox/GSH** triggers, but much weaker for **ATP-responsive microcapsule release** and still sparse for direct **hypoxia-triggered capsule validation** [1] [15] [16] [17]. OR and hybrid OR/AND logic are discussed conceptually, yet the supplied set does not provide a robust catalog of explicit tumor-targeted capsule examples with quantified off-target activation [66] [67] [68] [69]. Cross-study comparison is also limited because many reports do not standardize trigger thresholds, release half-lives, or matched single-input controls [38] [70] [71] [72] [73] [74] [75].

Suggested Further Research

- Run **matched head-to-head comparisons** of single-input versus dual-input capsules using the same material, cargo, and tumor model [13] [23] [70].
- Develop **explicit OR or hybrid OR/AND capsule systems** for heterogeneous tumors and quantify false-positive activation in normal tissues [76] [77] [78] [79].
- Standardize reporting of **trigger thresholds, leakage at physiological conditions, release half-life, and false activation rates** [15] [16] [72] [75].
- Expand direct studies of **ATP- and hypoxia-responsive capsule release** beyond review-level discussion [1] [64] [80].
- Prioritize models that better capture **human ECM density, vascular heterogeneity, and immune context**, alongside scalable manufacturing studies [55] [57] [60] [61].

Full Report

Introduction and Scope

Smart tumor-targeted microcapsule logic gates are engineered carriers that translate biological or external inputs into conditional changes in permeability, structure, targeting, or release state [1] [3] [19] [81]. In cancer therapy, the operational goal is straightforward: keep the payload largely inactive during circulation, then activate it preferentially within tumors through microenvironment-specific cues such as acidity, redox imbalance, hypoxia, or enzymes [3] [38] [62] [82].

In the supplied evidence, “logic gate” rarely means a purely formal digital gate implemented in isolation. More often, it refers to **multi-input selectivity**, **cascade activation**, or **stage-coupled transport and release** [20] [21] [22]. This distinction matters, because the field’s practical progress is being driven less by symbolic logic per se than by **materials architectures that embody logical conditions** [26] [27] [33].

Confidence level: High - this framing is directly supported by both the conceptual and engineering evidence provided [3] [19] [27].

Foundational Context

What a “logic-gated” tumor capsule actually does

A smart tumor capsule generally has three linked functions:

- **Sensing:** detect one or more tumor-associated inputs such as low pH, high GSH, hypoxia, or enzyme activity [1] [2] [5].
- **Transduction:** convert that input into a material change, such as protonation, bond cleavage, charge reversal, shell swelling, crosslink rupture, or layer erosion [7] [12] [13] [83].
- **Action:** increase uptake, expose a targeting ligand, open the shell, or release the therapeutic cargo [24] [84] [85].

This logic can be arranged as:

- **Single-input gating:** one cue directly triggers release [15] [17].
- **AND gating:** two conditions are both required for full activation [19] [20] [23].
- **Sequential AND gating:** one cue primes the system, a second cue finishes activation [21] [22] [24].
- **OR-style concepts:** any one of several cues may trigger activation, mainly discussed conceptually for heterogeneous tumors [76] [77].

Taken together, these findings suggest that the field values **conditional selectivity** more than abstract gate taxonomy: designs are favored when they reduce premature release while improving uptake and intracellular delivery [3] [21] [86].

Confidence level: High - the supplied evidence consistently links gate logic to material-state transitions and controlled release outcomes [9] [24] [32].

Tumor Microenvironment Inputs and Sensing Mechanisms

Acidic pH as a release trigger

Acidic pH is the most operationalized endogenous input in the supplied literature [8] [87] [88]. Mechanistically, acidity is converted into release through **protonation-induced ionization**, **acid-labile bond cleavage**, and **pH-driven shell destabilization or permeability changes** [7] [8] [9] [10].

The validation evidence is unusually direct. In αS-AuNP microcapsules, cargo release increased about 4-fold at pH 6.5 and 10-fold at pH 6.0 relative to pH 7.5, with most release at pH 5.0 [15]. In a pH-responsive liposome, ~72.75% quercetin release occurred by 48 h at pH 5.4 versus ~5.82% over the study at pH 7.4 [17]. In a ferritin-doxorubicin system, acidic conditions yielded >65% release in 4 h and >75% by 48 h, versus <25% cumulative release at pH 7.4 [18].

These results support pH as both a **release gate** and an **upstream priming gate**, because acidity can also induce charge reversal or structural changes that improve penetration before final intracellular release [24] [84].

Redox and glutathione-responsive designs

Redox/GSH-responsive systems are the second-best validated class [89] [90] [91]. Their mechanism is usually cleavage of **disulfide/diselenide linkages** or disruption of **ROS-sensitive moieties**, producing carrier disassembly or increased permeability [11] [12] [13] [14].

A direct assay example added 10 mM GSH to a release system at 5 h in PBS pH 7.4 and tracked cargo liberation over 72 h via UV/ninhydrin methods, explicitly modeling tumor-like reductive conditions [16]. Reviews and primary examples repeatedly frame redox chemistry as a way to preserve extracellular stability yet promote rapid intracellular release after uptake [63] [92] [93].

A recurrent design pattern is **pH for external tumor priming plus GSH for intracellular completion**, which is effectively sequential AND logic [9] [13].

Hypoxia, enzymes, and ATP recognition

Hypoxia is biologically attractive because it maps to poorly oxygenated tumor regions, and the supplied sources describe low oxygen as a trigger for bond cleavage or disassembly [1] [4] [64]. One dual-responsive nanogel used acidic charge reversal to improve uptake, then hypoxia-activated cleavage of an azobenzene crosslinker for doxorubicin release [24]. However, the evidence base is thinner than for pH or GSH, and the supplied full-text validation set did not retrieve direct hypoxia-triggered microcapsule release kinetics [15] [16] [17].

Enzyme-responsive systems, especially MMP-linked designs, use protease activity to cleave peptide substrates or destabilize carrier structures [83] [94] [95] [96]. This is especially relevant to extracellular matrix remodeling and tumor-localized release [97] [98]. Cathepsins are mentioned as candidate protease triggers, but the supplied set does not provide detailed capsule-specific assay kinetics for them [95] [99] [100].

ATP appears mainly as a **recognition concept** rather than a well-validated capsule trigger in this dataset [1] [80]. An influential gatekeeper example used ATP-displaceable aptamer/Au nanoparticle caps in mesoporous silica, showing biomolecular control of pore opening, but this is not direct tumor-targeted microcapsule validation [101].

Confidence level: High for pH/redox, Medium for enzyme, Low for ATP/hypoxia capsule-specific validation - the asymmetry reflects the depth of direct assay evidence in the supplied record [15] [16] [17] [18].

Logic-Gate Architectures for Tumor Selectivity

AND-gated and dual-input systems

The dominant architecture is AND-like gating, where multiple tumor-associated conditions are required before full activation [19] [20] [21] [22]. This can reduce false positives by separating tumor arrival, penetration, uptake, and intracellular release into conditional steps [23].

Examples include pH/redox systems that improved penetration and achieved ~70% drug release at 24 h with greater apoptosis than free drug [13]. Another study used acidic-tumor charge reversal followed by hypoxia-triggered doxorubicin release, yielding stronger antitumor efficacy with reduced off-target toxicity [24]. A supramolecular nanotheranostic combined hypoxia, esterase, and pH responsiveness in a two-layer cascade logic gate for selective imaging and synergistic therapy [25].

The central advantage is not only stricter release control, but **division of labor between inputs**: one cue can optimize localization or penetration, another can optimize intracellular release [24].

Sequential and cascade gating

Sequential designs are especially important for solid tumors because they align with real delivery barriers [102] [103]. Core-shell and multilayer architectures can stage payload access so that an outer layer responds first, then an inner barrier opens later [104] [105] [106].

The supplied literature describes cascade systems where pH-triggered transformation reveals a secondary targeting moiety or enables a later enzymatic response, improving deep tumor uptake and subcellular delivery [84] [85] [107] [108]. Sequential targeting across vasculature, tumor cells, or mitochondria is also reported to improve killing of both endothelial and tumor compartments [109] [110] [111].

Taken together, these findings suggest that “logic” in many advanced cancer capsules is best understood as **barrier-matched progression logic** rather than simple simultaneous coincidence detection [21] [84].

OR and hybrid logic concepts

OR logic is conceptually attractive for heterogeneous tumors because any one of several biomarkers could trigger activation [76] [77]. The rationale is broader coverage and lower risk of escape from single-marker dependence [78] [79].

However, the supplied evidence explicitly notes a gap: no robust set of tumor-targeted capsule studies was retrieved for OR or hybrid OR/AND logic with exact biomarker combinations and quantified off-target activation [66] [67] [68] [69]. Thus, there is a real tension between the conceptual appeal of broad activation and the practical dominance of specificity-focused AND systems [19] [20] [77].

Confidence level: High for AND/sequential gating, Medium for the OR-versus-specificity tension - the former is directly evidenced, while the latter is mainly a synthesis across uneven literature coverage [13] [66].

Microcapsule Materials, Fabrication, and Gate Engineering

Emulsion and microfluidic fabrication

Emulsion templating and microfluidics are recurrent enabling methods for responsive microcapsules [112] [113] [114]. The strongest evidence supports **double-emulsion microfluidics** for monodisperse core-shell capsules with tunable shell thickness and composition [27] [30] [31].

This matters because shell thickness variability, aggregation, and poorly controlled architecture directly affect leakage and trigger sensitivity [29] [115]. More complex triple-emulsion or multicompartment systems expand loading and protection options and can support different compartments or barriers for sequential release [32] [116].

Polymer systems and biodegradation

PLGA is repeatedly emphasized as a biodegradable tumor-delivery material because of biocompatibility, controllable degradation, and surface functionalization potential [117] [118] [119] [120]. PEGylation improves circulation and can reduce opsonization, while PEG-containing polymersomes can show pH-tunable membrane permeability [33] [121] [122].

Hydrogels support localized and triggerable release, including NIR-assisted and intracellular pH-responsive behavior [123] [124]. Crosslink density is a critical parameter because membrane-cross-linked polymersomes showed superior stability plus strong pH-selective permeability when optimized appropriately [33].

The broad implication is that material choice determines not just biocompatibility, but the **sharpness, reversibility, and selectivity of gate behavior** [33] [86].

Capsule architecture and mechanical stability

Two-shell, multilayer, and LbL architectures can produce staged release and protect cargos from premature loss [26] [32] [34]. Mechanical stability depends on shell gelation, network structure, and crosslinking; thicker or better-crosslinked shells can improve roundness and structural robustness [28] [115].

Foundational release papers also show that shell morphology affects burst behavior: dense skins, multilayer coatings, and polymer blending can reduce early leakage and shape later sustained release [65] [125] [126].

Confidence level: High - the link between fabrication/architecture and functional gate behavior is one of the most consistently supported themes in the dataset [27] [32] [34].

Payload Classes, Release Kinetics, and Combination Therapy

Chemotherapeutics and release profiles

Chemotherapy remains central, especially doxorubicin-based systems [104] [127] [128]. Direct quantitative examples include DOX/salvianolic acid A carriers with >80% encapsulation efficiency for both drugs and SBN-PLGA nanoparticles with 70.19% entrapment efficiency plus sustained zero-order release behavior [71] [129].

Burst suppression is a key engineering objective. High-molecular-weight PLGA and 7-layer chitosan/alginate LbL coatings reduced early burst compared with lower-MW or uncoated systems [130]. Mechanistic release literature further shows that microsphere release reflects coupled diffusion, swelling,

erosion, hydration, and polymer scission processes, reinforcing why logic behavior must be engineered at the materials level [72] [73] [74] [75].

A notable gap is that no retrieved study provided release half-life values, limiting precise quantitative comparison [38] [70] [71].

Immunotherapy co-delivery

Combination therapy is a major growth area [35] [36] [37]. GM-CSF co-delivery in chitosan hydrogel systems increased APC recruitment, E7-specific CD8+ T-cell responses, IFN-γ, and tumor regression in cervical cancer models [39] [131]. STING-agonist combinations upregulated type I interferons and improved tumor suppression or survival, including systems paired with chemotherapy or siPD-L1 [40] [132]. Metformin-containing combinations reduced hypoxia, increased CD8+ and CD4+ T cells, reduced Tregs, enhanced cisplatin sensitivity, and improved antitumor immunity [41] [42] [43].

The common principle is synchronized local release of a cytotoxic and an immune or metabolic modulator to outperform monotherapy [39] [40] [42].

Light-, heat-, and phase-change activated release

External triggers are often layered onto endogenous gating to add physician-controlled activation [133] [134] [135]. NIR irradiation can open gold nanoparticle microcapsules through local heating and release cargo into the cytosol [136]. γ-irradiation decomposed radiosensitive microcapsules, with optimal release reported at 2 Gy [137] [138]. Focused ultrasound hyperthermia was used to increase intratumoral doxorubicin after ThermoDox administration [139] [140].

This hybrid endogenous-plus-exogenous strategy appears especially relevant where tumor heterogeneity makes sole reliance on biological triggers risky [47] [141].

Confidence level: Medium-High - payload and combination trends are well supported, but kinetic standardization is weak [35] [40] [72].

In Vivo Performance, Safety, and Translation

Tumor targeting and biodistribution

The in vivo evidence is encouraging. Ultrasound-triggered MRI-visible multilayer microcapsules retained doxorubicin in circulation for at least 48 h and, after focused ultrasound, achieved 16-fold higher tumor localization than off-target organs in a breast-cancer mouse model [47]. Stepwise targeting GPDC-MSNs accumulated up to 25-fold more in tumor than normal hepatic tissue with superior antitumor efficacy [45]. pH/NIR-responsive UCNP@SnO2-BSA nanocapsules markedly inhibited tumor growth with negligible side effects [44]. Other systems showed tumor-tropic accumulation, improved penetration, metastasis suppression, or imaging-confirmed localization [46] [48].

These reports support the view that gated capsule-like systems can improve spatial localization and therapeutic index in mouse models [44] [48] [142].

Safety and off-target toxicity

Safety readouts commonly included body weight, histology, hematology, and serum chemistry [51] [143] [144]. Several systems preserved body weight and showed no major organ damage or inflammatory infiltrates [143] [144] [145]. Hematology and serum chemistry often remained within normal limits under responsive delivery conditions, while free-drug controls were more toxic [51] [143] [144].

Comparative evidence for reduced systemic exposure is particularly important: an MMP-2/ROS-responsive hydrogel showed strong tumor suppression with a 5-fold lower incidence of systemic toxicity than intravenous doxorubicin [49], and reduction-responsive ultrasound-activated platforms reduced premature leakage and off-target toxicity [50].

Translation barriers and unmet needs

Despite strong preclinical outcomes, translation remains difficult [52] [53] [54]. Dense ECM, abnormal vasculature, and limited permeability reduce penetration and bioavailability in solid tumors [55] [56] [60] [146] [147]. EPR is real but heterogeneous and often insufficient for reliable uniform delivery [61] [148] [149] [150] [151]. The delivered tumor dose is often only a small fraction of the administered amount [59] [60].

Manufacturing and regulatory issues also persist, especially for monodisperse, multi-compartment, logic-gated systems requiring tight control over shell architecture and trigger sensitivity [57] [58] [152]. Better

preclinical models are needed because current mouse systems do not fully capture human ECM density, vascular heterogeneity, and immune context [57] [61].

Confidence level: High - the translational barriers are repeatedly and consistently stated across the supplied literature [52] [54] [57] [61].

Cross-Source Synthesis

Across the evidence base, three conclusions stand out. First, **multi-input and sequential gating has become the preferred strategy** because it better matches the stepwise barriers of tumor delivery than one-shot single-trigger release [3] [21] [38] [84]. Second, **materials engineering is inseparable from logic design**: shell thickness, crosslink density, layered architecture, and fabrication route determine whether a nominal trigger actually produces selective, burst-suppressed, staged release [27] [33] [125]. Third, **therapeutic ambition is expanding toward coordinated multimodal therapy**, especially chemo-immunotherapy and externally assisted release, but translation remains limited by biology rather than gate sophistication alone [40] [42] [59] [60].

Inferred: A particularly promising next-step architecture would combine extracellular pH priming, enzyme-mediated stromal unlocking, and intracellular GSH release, because that sequence explicitly maps onto tumor entry, matrix penetration, and cytosolic drug liberation [24] [55] [56] [95].

Confidence level: Medium-High - the general synthesis is strongly grounded, while the final design implication is an inference across multiple studies rather than a single directly validated capsule platform [55] [84].

Conclusion

Smart tumor-targeted microcapsule logic gates are best viewed as **programmable, condition-responsive release systems** whose most mature forms use **pH and redox/GSH inputs**, **AND or sequential AND logic**, and **carefully engineered shell architectures** to improve tumor selectivity and reduce premature leakage [7] [13] [19] [24] [27]. Their most convincing applications are in **combination therapy** and **image- or externally assisted delivery**, where multistage control can synchronize localization and action [40] [42] [47].

The field's core challenge is now less about proving that logic-gated release is possible and more about proving that it remains effective under the transport constraints of real human tumors and the manufacturing constraints of translation [57] [59] [61].

Overall confidence: High - the broad design trends, strongest triggers, engineering determinants, and translational barriers are consistently supported in the supplied evidence [3] [19] [27] [35] [54].

References

1. Ovais, M., Mukherjee, S., Pramanik, A., (...), Chen, C. (2020) Designing Stimuli-Responsive Upconversion Nanoparticles that Exploit the Tumor Microenvironment. *Advanced Materials*. https://doi.org/10.1002/adma.202000055
2. Todke, P., Attar, E., Chaudhari, V., (...), Devarajan, P.V. (2026) Overview of Stimuli-Responsive Nanosystems: Endogenous and Exogenous Triggers for Stimuli-Responsive Delivery for Cancer Therapy. *Advances in Stimuli-Responsive Nanosystems for Cancer Therapy*. https://doi.org/10.1007/978-981-95-2744-1_1
3. Liu, Y., Zhou, J., Huo, M. (2016) Advances in the tumor microenvironment-responsive smart drug delivery nanosystem. *Journal of China Pharmaceutical University*. https://doi.org/10.11665/j.issn.1000-5048.20160201
4. Gong, F., Yang, N., Wang, X., (...), Cheng, L. (2020) Tumor microenvironment-responsive intelligent nanoplatforms for cancer theranostics. *Nano Today*. https://doi.org/10.1016/j.nantod.2020.100851
5. Park, H., Saravanakumar, G., Kim, J., (...), Kim, W.J. (2021) Tumor Microenvironment Sensitive Nanocarriers for Bioimaging and Therapeutics. *Advanced Healthcare Materials*. https://doi.org/10.1002/adhm.202000834
6. Rajankar, N., Kumar, A., Gulbake, A. (2025) Navigating tumor microenvironment with TME-responsive nanocarriers: Strategies for targeted drug delivery in solid tumor. *Journal of Drug Delivery Science and Technology*. https://doi.org/10.1016/j.jddst.2025.107050

7. Mali, K.K., Gavhane, Y.N., Chakole, R.D. (2024) Natural Polymer-Based Nanogel for pH-Responsive Delivery of Sorafenib Tosylate in Hemangiosarcoma. *AAPS PharmSciTech*. https://doi.org/10.1208/s12249-024-02797-8
8. Jiang, J., Jin, L., Zhao, Z., (...), Li, F. (2025) Design and application of proton gradient-based pH-responsive nanomaterials in the tumor microenvironment. *Nanoscale*. https://doi.org/10.1039/d5nr02990b
9. Li, Y., Yi, M., Xiong, B., (...), Lu, B. (2024) pH-responsive degradable mesoporous organosilica nanoparticle for tumor targeting and phototherapy combined with chemotherapy. *Journal of Drug Delivery Science and Technology*. https://doi.org/10.1016/j.jddst.2024.105344
10. Zhang, R., Liu, T., Li, W., (...), Tao, Y. (2022) Tumor microenvironment-responsive BSA nanocarriers for combined chemo/chemodynamic cancer therapy. *Journal of Nanobiotechnology*. https://doi.org/10.1186/s12951-022-01442-5
11. Mirhadi, E., Mashreghi, M., Faal Maleki, M., (...), Jaafari, M.R. (2020) Redox-sensitive nanoscale drug delivery systems for cancer treatment. *International Journal of Pharmaceutics*. https://doi.org/10.1016/j.ijpharm.2020.119882
12. Liu, Z.-L., Ma, S., Sun, J.-L., (...), Chen, X.-S. (2020) Reactive oxygen species responsive core-shell nanoparticles increase tumor enrichment and endocytosis. *Acta Polymerica Sinica*. https://doi.org/10.11777/j.issn1000-3304.2020.20066
13. Badparvar, F., Poursattar Marjani, A., Salehi, R., (...), Talebi, M. (2024) Dual pH/redox-responsive size-switchable polymeric nano-carrier system for tumor microenvironment DTX release. *Journal of Biomaterials Science, Polymer Edition*. https://doi.org/10.1080/09205063.2024.2371203
14. Walvekar, P., Sikhosana, N., Adeyemi, S.A., (...), Choonara, Y.E. (2023) Redox-Responsive Delivery Nanoplatforms in Cancer Theranostics. *Site-specific Cancer Nanotheranostics: A Microenvironment-responsive Approach*. https://doi.org/10.1201/9781003368731-5
15. Nam, E.-J., Kong, S.-K., Cho, I., (...), Kim, Y.-S. (2024) Augmented anti-cancer effect of hypoxia-inducible factor 2α (HIF2α)- targeting PT2385 delivered in α-synuclein-AuNP microcapsules for clear cell renal carcinoma mouse model. *Journal of Drug Delivery Science and Technology*. https://doi.org/10.1016/j.jddst.2023.105179
16. Cui, S., Yu, Q., Su, Z., (...), Yang, D. (2025) A pH-responsive and nitric oxide-producing nanocarrier for targeted therapy of melanoma. *Chemical Engineering Journal*. https://doi.org/10.1016/j.cej.2025.169985
17. Lan, Q., Wang, M., Zhu, Y., (...), Xu, Y. (2025) Bioinspired pH-sensitive liposomes for quercetin delivery to synergize with 5- FU in gastric cancer therapy. *International Journal of Pharmaceutics: X*. https://doi.org/10.1016/j.ijpx.2025.100437
18. Zhao, W., Huang, F., Uddin, S., (...), Li, H. (2025) Computer-aided drug design of SP94 peptide-functionalized human H-chain ferritin for targeted doxorubicin delivery. *International Journal of Biological Macromolecules*. https://doi.org/10.1016/j.ijbiomac.2025.148521
19. Huang, H., Guo, Z., Zhang, C., (...), Tan, W. (2021) Logic-Gated Cell-Derived Nanovesicles via DNA-Based Smart Recognition Module. *ACS Applied Materials and Interfaces*. https://doi.org/10.1021/acsami.1c07632
20. Zhang, P., Gao, D., An, K., (...), Tan, W. (2020) A programmable polymer library that enables the construction of stimuli-responsive nanocarriers containing logic gates. *Nature Chemistry*. https://doi.org/10.1038/s41557-020-0426-3
21. Yan, C., Guo, Z., Liu, Y., (...), Zhu, W.-H. (2018) A sequence-activated and logic dual-channel fluorescent probe for tracking programmable drug release. *Chemical Science*. https://doi.org/10.1039/c8sc02079e
22. Fang, Y., Yin, W., Han, N., (...), Ruan, M. (2026) A logic-gated immunotherapeutic nanoplatform amplifying antigen release via reprogrammed cell death for durable antitumor immunity. *Chemical Engineering Journal*. https://doi.org/10.1016/j.cej.2026.176973
23. Jiang, C., Zhao, Z., East, A.K., (...), Chan, J. (2025) Logic-gated approach for targeted delivery and site-selective activation of photothermal agents in precision cancer treatment. *Chemical Science*. https://doi.org/10.1039/d4sc08228a

24. Liao, Y., Wang, Z., Guo, M., (...), Cheng, Q. (2025) Dual-responsive zwitterionic nanogels with synergistic charge reversal and hypoxia-activated drug release for precise tumor targeting and enhanced chemotherapy. *Colloids and Surfaces B: Biointerfaces*. https://doi.org/10.1016/j.colsurfb.2025.114839

25. Liu, X., Meng, C., Ji, G., (...), Ling, Y. (2021) Tumor microenvironment-activatable boolean logic supramolecular nanotheranostics based on a pillar[6]arene for tumor hypoxia imaging and multimodal synergistic therapy. *Materials Chemistry Frontiers*. https://doi.org/10.1039/d1qm00411e

26. Shallan, A.I., Priest, C. (2019) Microfluidic process intensification for synthesis and formulation in the pharmaceutical industry. *Chemical Engineering and Processing - Process Intensification*. https://doi.org/10.1016/j.cep.2019.107559

27. Wei, J., Ju, X.-J., Xie, R., (...), Chu, L.-Y. (2011) Novel cationic pH-responsive poly(N,N-dimethylaminoethyl methacrylate) microcapsules prepared by a microfluidic technique. *Journal of Colloid and Interface Science*. https://doi.org/10.1016/j.jcis.2011.01.105

28. Tu, W., Hu, W., Chen, J., (...), Qiao, D. (2025) Controllable fabrication of core-shell microcapsules using sodium alginate/gellan gum as shell material by microfluidics. *Food Hydrocolloids*. https://doi.org/10.1016/j.foodhyd.2025.111514

29. Jeong, S.-G., Choi, Y., Nam, J.-O., (...), Choi, C.-H. (2021) Surface-tension-induced double emulsion drops via phase separation of polymeric fluid confined in micromolds for capsule templates. *Journal of Colloid and Interface Science*. https://doi.org/10.1016/j.jcis.2020.08.105

30. Ju, X.-J., Chu, L.-Y. (2018) Lab-on-a-chip fabrication of polymeric microparticles for drug encapsulation and controlled release. *Microfluidics for Pharmaceutical Applications: From Nano/Micro Systems Fabrication to Controlled Drug Delivery*. https://doi.org/10.1016/B978-0-12-812659-2.00009-0

31. Sanjay, S.T., Zhou, W., Dou, M., (...), Li, X. (2018) Recent advances of controlled drug delivery using microfluidic platforms. *Advanced Drug Delivery Reviews*. https://doi.org/10.1016/j.addr.2017.09.013

32. Guo, Jianhua, Hou, Lihua, Hou, Junpeng, (...), Hu, Qingming (2020) Generation of Ultra-Thin-Shell Microcapsules Using Osmolarity-Controlled Swelling Method. *Micromachines*. https://doi.org/10.3390/mi11040444

33. Mohammed, A.A.-W., Lu, N., Wang, Y., (...), Ge, Z. (2020) Membrane-cross-linked polymersomes with tumor pH-tunable selective permeability as intelligent nanoreactors and drug delivery vehicles. *European Polymer Journal*. https://doi.org/10.1016/j.eurpolymj.2020.109982

34. Tong, W., Song, X., Gao, C. (2012) Layer-by-layer assembly of microcapsules and their biomedical applications. *Chemical Society Reviews*. https://doi.org/10.1039/c2cs35088b

35. Chen, M., Wang, C., Wang, X., (...), Liu, Z. (2024) An “AND” Logic-Gated Prodrug Micelle Locally Stimulates Antitumor Immunity. *Advanced Materials*. https://doi.org/10.1002/adma.202307818

36. Song, Y., Du, Y., Hu, C., (...), Gao, H. (2024) Metformin-Mediated Immunosuppressive Microenvironment Remodeling in Combination with Chemotherapy via a Spatial-Specific Multi-Responsive Carrier-Free Self-Assembled Nanoparticle. *Advanced Functional Materials*. https://doi.org/10.1002/adfm.202316145

37. Ni, H., Qian, S., Lu, J., (...), Zhang, J. (2023) Natural Polysaccharide Delivery Platforms with Multiscale Structure Used for Cancer Chemoimmunotherapy. *Molecular Pharmaceutics*. https://doi.org/10.1021/acs.molpharmaceut.3c00633

38. Liao, L., Cen, D., Fu, Y., (...), Han, G. (2020) Biodegradable MnFe-hydroxide nanocapsules to enable multi-therapeutics delivery and hypoxia-modulated tumor treatment. *Journal of Materials Chemistry B*. https://doi.org/10.1039/d0tb00243g

39. Khorramdelazad, H., Pirsadeghi, A., Amraei, E., (...), Mohammadi-Kordkhayli, M. (2026) Hydrogel-based delivery of cytokines in cancer immunotherapy: Latest evidence and therapeutic outcomes. *Biomedicine and Pharmacotherapy*. https://doi.org/10.1016/j.biopha.2025.118953

40. Yang, Y.-Y., Zhang, X.-R., Fu, M.-Q., (...), Yao, Y. (2026) Engineering copper-based nanoplatforms in cancer therapy. *Materials and Design*. https://doi.org/10.1016/j.matdes.2026.116337

41. Liu, Y., Sun, J., Huang, Y., (...), Li, S. (2021) Metformin-conjugated micellar system with intratumoral pH responsive de-shielding for co-delivery of doxorubicin and nucleic acid. *Biochemical Pharmacology*. https://doi.org/10.1016/j.bcp.2021.114453

42. Pourmadadi, M., Shaghaghi, M., Arshad, R., (...), Pandey, S. (2024) Polymeric and non-polymeric oxaliplatin nanomedicine for cancer therapy: A comprehensive review. *European Polymer Journal*. https://doi.org/10.1016/j.eurpolymj.2024.112870

43. Lu, Qiang, Kou, Dongquan, Lou, Shenghan, (...), Han, Peng (2024) Nanoparticles in tumor microenvironment remodeling and cancer immunotherapy. *Journal of Hematology and Oncology*. https://doi.org/10.1186/s13045-024-01535-8

44. Yang, L., Shi, R., Zhao, R., (...), Feng, L. (2022) Near-Infrared Upconversion Mesoporous Tin Dioxide Theranostic Nanocapsules for Synergetic Cancer Chemophototherapy. *ACS Applied Materials and Interfaces*. https://doi.org/10.1021/acsami.1c23174

45. Li, Y., Miao, Y., Chen, M., (...), Gan, Y. (2020) Stepwise targeting and responsive lipid-coated nanoparticles for enhanced tumor cell sensitivity and hepatocellular carcinoma therapy. *Theranostics*. https://doi.org/10.7150/thno.42008

46. Cherukula, K., Uthaman, S., Park, I.-K. (2019) “Navigate-dock-activate” anti-tumor strategy: Tumor micromilieu charge-switchable, hierarchically activated nanoplatform with ultrarapid tumor-tropic accumulation for trackable photothermal/chemotherapy. *Theranostics*. https://doi.org/10.7150/thno.33280

47. Alford, A., Rich, M., Kozlovskaya, V., (...), Kharlampieva, E. (2018) Ultrasound-Triggered Delivery of Anticancer Therapeutics from MRI-Visible Multilayer Microcapsules. *Advanced Therapeutics*. https://doi.org/10.1002/adtp.201800051

48. Wang, Z., Ju, Y., Ali, Z., (...), Hou, Y. (2019) Near-infrared light and tumor microenvironment dual responsive size-switchable nanocapsules for multimodal tumor theranostics. *Nature Communications*. https://doi.org/10.1038/s41467-019-12142-4

49. https://www.sciencedirect.com/science/article/pii/S3004926126000370

50. Guo, H., Chen, L., Yu, K., Wang, J. (2026) Ultrasound-activated biomimetic nanoplatform enables long-acting chemo-sonodynamic therapy of prostate cancer and lymphatic metastasis. *Colloids and Surfaces B: Biointerfaces*. https://doi.org/10.1016/j.colsurfb.2026.115578

51. Puja, S., Jayant, G., Rinti, B., Sathish, D. (2026) Targeted dual-drug delivery for minimal residual glioblastoma using multifunctional thermosensitive liposomal hydrogels. *Journal of Drug Delivery Science and Technology*. https://doi.org/10.1016/j.jddst.2026.108153

52. Khawar, I.A., Ghosh, T., Park, J.K., Kuh, H.-J. (2021) Tumor spheroid-based microtumor models for preclinical evaluation of anticancer nanomedicines. *Journal of Pharmaceutical Investigation*. https://doi.org/10.1007/s40005-021-00534-y

53. Hu, C., Gao, H.-L. (2020) Advances in tumor microenvironment responsive and regulatory drug delivery system. *Yaoxue Xuebao*. https://doi.org/10.16438/j.0513-4870.2019-0798

54. Priester, M.I., ten Hagen, T.L.M. (2023) Image-guided drug delivery in nanosystem-based cancer therapies. *Advanced Drug Delivery Reviews*. https://doi.org/10.1016/j.addr.2022.114621

55. Lu, W., Wang, N., Liu, X., (...), Teng, Z. (2024) Matrix-degrading soft-nanoplatform with enhanced tissue penetration for amplifying photodynamic therapeutic efficacy of breast cancer. *Journal of Materials Chemistry B*. https://doi.org/10.1039/d4tb00894d

56. Lu, W., Li, Y., Zhang, X., (...), Chen, K. (2024) Dual-modal overcoming of physical barriers for improved photodynamic cancer therapy via soft organosilica nanocapsules. *Journal of Nanobiotechnology*. https://doi.org/10.1186/s12951-024-02945-z

57. Mamidi, N., Franco de Silva, F., Salehi, A. (2025) Advanced disease therapeutics using engineered living drug delivery systems. *Nanoscale*. https://doi.org/10.1039/d4nr05298f

58. Bharti, N., Kumar, V., Mukherjee, S., (...), Kumar, M. (2025) Smart microneedle platforms for skin cancer therapy: a review of stimuli-responsive and targeted drug delivery systems. *Pharmaceutical Development and Technology*. https://doi.org/10.1080/10837450.2025.2580499

59. Wilhelm, S., Tavares, A.J., Dai, Q., (...), Chan, W.C.W. (2016) Analysis of nanoparticle delivery to tumours. *Nature Reviews Materials*. https://doi.org/10.1038/natrevmats.2016.14

60. Jain, R.K., Stylianopoulos, T. (2010) Delivering nanomedicine to solid tumors. *Nature Reviews Clinical Oncology*. https://doi.org/10.1038/nrclinonc.2010.139

61. Prabhakar, U., Maeda, H., Jain, R., (...), Blakey, D.C. (2013) Challenges and key considerations of the enhanced permeability and retention effect for nanomedicine drug delivery in oncology. *Cancer Research*. https://doi.org/10.1158/0008-5472.CAN-12-4561

62. Li, J., Wei, Z., Lin, X., (...), Liu, J. (2019) Programmable Therapeutic Nanodevices with Circular Amplification of H2O2 in the Tumor Microenvironment for Synergistic Cancer Therapy. *Advanced Healthcare Materials*. https://doi.org/10.1002/adhm.201801627

63. Mura, S., Nicolas, J., Couvreur, P. (2013) Stimuli-responsive nanocarriers for drug delivery. *Nature Materials*. https://doi.org/10.1038/nmat3776

64. Brown, J.M., Wilson, W.R. (2004) Exploiting tumour hypoxia in cancer treatment. *Nature Reviews Cancer*. https://doi.org/10.1038/nrc1367

65. Berkland, C., King, M., Cox, A., (...), Pack, D.W. (2002) Precise control of PLG microsphere size provides enhanced control of drug release rate. *Journal of Controlled Release*. https://doi.org/10.1016/S0168-3659(02)00136-0

66. Lam, C.K., Hyde, R.K., Patel, S.A. (2022) Synthetic Immunotherapy: Programming Immune Cells with Novel and Sophisticated Logic Capabilities. *Transplantation and Cellular Therapy*. https://doi.org/10.1016/j.jtct.2022.06.001

67. Yin, J., Ma, X., Hu, L., Ye, H. (2025) Translating synthetic gene circuits into the clinic: Challenges, opportunities, and future directions. *Cell Systems*. https://doi.org/10.1016/j.cels.2025.101425

68. Zhou, X., Han, Y., Fang, Y., (...), Li, N. (2025) Antibody-drug conjugates: Current challenges and innovative solutions for precision cancer therapy. *Med*. https://doi.org/10.1016/j.medj.2025.100849

69. Shen, N., Zhao, W., Chu, H., (...), Chen, X. (2025) Targeted delivery and controlled release of polymeric nanomedicines for tumor therapy. *Fundamental Research*. https://doi.org/10.1016/j.fmre.2025.01.011

70. Agbaje, K.O., Adesina, S.K., Adebayo, A.S. (2025) Development and Characterization of Optimized Drug-Loaded Niosomes for Delivery of 5-FU and Irinotecan. *Pharmaceutics*. https://doi.org/10.3390/pharmaceutics17070900

71. Rahimnia, S.M., Saeedi, M., Akbari, J., (...), Yazdian–Robati, R. (2024) Development, Optimization, and in vitro Evaluation of Silybin-loaded PLGA Nanoparticles and Decoration with 5TR1 Aptamer for Targeted Delivery to Colorectal Cancer Cells. *AAPS PharmSciTech*. https://doi.org/10.1208/s12249-024-02858-y

72. Goepferich, A. (1996) Mechanisms of polymer degradation and erosion. *Biomaterials*. https://doi.org/10.1016/0142-9612(96)85755-3

73. Fredenberg, S., Wahlgren, M., Reslow, M., Axelsson, A. (2011) The mechanisms of drug release in poly(lactic-co-glycolic acid)-based drug delivery systems - A review. *International Journal of Pharmaceutics*. https://doi.org/10.1016/j.ijpharm.2011.05.049

74. Batycky, R.P., Hanes, J., Langer, R., Edwards, D.A. (1997) A theoretical model of erosion and macromolecular drug release from biodegrading microspheres. *Journal of Pharmaceutical Sciences*. https://doi.org/10.1021/js9604117

75. Berkland, C., Kipper, M.J., Narasimhan, B., (...), Pack, D.W. (2004) Microsphere size, precipitation kinetics and drug distribution control drug release from biodegradable polyanhydride microspheres. *Journal of Controlled Release*. https://doi.org/10.1016/j.jconrel.2003.09.011

76. Bi, S., He, H., Gao, F., Zhao, Y. (2024) Ultrasensitive Photoelectrochemical Biosensor for Dual-miRNAs Detection Based on Molecular Logic Gates and Methylene Blue Sensitized ZnO@CdS@Au Nanorods. *ACS Applied Materials and Interfaces*. https://doi.org/10.1021/acsami.4c08276

77. Zhang, Y., Yang, Q., Zhu, L., (...), Luo, X. (2024) Intelligent Cell Profiling and Precision Release: Multimolecular Marker-Activated Transmembrane DNA Computing Nanosystem. *Analytical Chemistry*. https://doi.org/10.1021/acs.analchem.4c01122

78. Zhou, S., Li, F., Niu, M., (...), Yi, M. (2025) Advances in the clinical application of bispecific antibodies in cancer therapy. *iScience*. https://doi.org/10.1016/j.isci.2025.114203

79. Kleinmanns, Katrin, Fosse, Vibeke, Bjørge, Line, McCormack, Emmet (2020) The Emerging Role of CD24 in Cancer Theranostics—A Novel Target for Fluorescence Image-Guided Surgery in Ovarian Cancer and Beyond. *Journal of Personalized Medicine*. https://doi.org/10.3390/jpm10040255

80. Wang, Y., Deng, T., Liu, X., (...), Fan, X. (2024) Smart Nanoplatforms Responding to the Tumor Microenvironment for Precise Drug Delivery in Cancer Therapy. *International Journal of Nanomedicine*. https://doi.org/10.2147/IJN.S459710

81. Ruskowitz, E.R., Comerford, M.P., Badeau, B.A., Deforest, C.A. (2019) Logical stimuli-triggered delivery of small molecules from hydrogel biomaterials. *Biomaterials Science*. https://doi.org/10.1039/c8bm01304g

82. Du, J., Lane, L.A., Nie, S. (2015) Stimuli-responsive nanoparticles for targeting the tumor microenvironment. *Journal of Controlled Release*. https://doi.org/10.1016/j.jconrel.2015.08.050

83. Mdlovu, N.V., Lin, K.-S., Wu, C.-M. (2025) Nano-encapsulation as a promising approach for enhancing cancer therapy. *IoT-WSN-DT Based Medical Systems and Nanotechnology for Smart Cancer Care*. https://doi.org/10.1016/B978-0-443-33984-4.00004-3

84. Wu, H., Lu, H., Xiao, W., (...), Lin, T.-Y. (2020) Sequential Targeting in Crosslinking Nanotheranostics for Tackling the Multibarriers of Brain Tumors. *Advanced Materials*. https://doi.org/10.1002/adma.201903759

85. Chen, H., Yao, Y., Zhao, X., Tan, N. (2023) Programmable site-specific delivery of an alkaline phosphatase-activatable prodrug and a mitochondria-targeted cyclopeptide for combination therapy in colon cancer. *Biomaterials Science*. https://doi.org/10.1039/d3bm00834g

86. Sun, H., Zhong, Z. (2020) 100th anniversary of macromolecular science viewpoint: Biological stimuli-sensitive polymer prodrugs and nanoparticles for tumor-specific drug delivery. *ACS Macro Letters*. https://doi.org/10.1021/acsmacrolett.0c00488

87. Liu, Y., Si, L., Jiang, Y., (...), Hu, J. (2025) Design of pH-Responsive Nanomaterials Based on the Tumor Microenvironment. *International Journal of Nanomedicine*. https://doi.org/10.2147/IJN.S504629

88. Ali, A., Javed, R., Farhangi, S., (...), Rui, Y. (2023) Metal phenolic networks (MPNs)-based pH-sensitive stimulus responsive nanosystems for drug delivery in tumor microenvironment. *Journal of Drug Delivery Science and Technology*. https://doi.org/10.1016/j.jddst.2023.104536

89. Badparvar, F., Poursattar Marjani, A.P., Salehi, R., Ramezani, F. (2023) pH/redox responsive size-switchable intelligent nanovehicle for tumor microenvironment targeted DOX release. *Scientific Reports*. https://doi.org/10.1038/s41598-023-49446-x

90. Yang, Y.-Q., Gong, F., Bai, S., Cheng, L. (2021) Progress in tumor microenvironment responsive nano-platforms for cancer theranostics. *Yaoxue Xuebao*. https://doi.org/10.16438/j.0513-4870.2020-1223

91. Li, S., Wang, Q., Li, Z., (...), Feng, X. (2025) Glutathione-responsive polypeptide nanogel encapsulates Shikonin for breast cancer therapy. *Nanomedicine: Nanotechnology, Biology, and Medicine*. https://doi.org/10.1016/j.nano.2025.102802

92. Meng, X., Shen, Y., Zhao, H., (...), Zhao, Y. (2024) Redox-manipulating nanocarriers for anticancer drug delivery: a systematic review. *Journal of Nanobiotechnology*. https://doi.org/10.1186/s12951-024-02859-w

93. Yang, R., Ouyang, Z., Guo, H., (...), Shi, X. (2022) Microfluidic synthesis of intelligent nanoclusters of ultrasmall iron oxide nanoparticles with improved tumor microenvironment regulation for dynamic MR imaging-guided tumor photothermo-chemo-chemodynamic therapy. *Nano Today*. https://doi.org/10.1016/j.nantod.2022.101615

94. Dang, W., Cui, T., Jiao, G., (...), Liu, Z. (2025) Subcellular organelle-mediated precise delivery of nanomedicines: Advances and applications in cancer therapy. *Journal of Drug Delivery Science and Technology*. https://doi.org/10.1016/j.jddst.2025.107261

95. Yoon, Moon Sup, Lee, Jae Min, Jo, Min Jeong, (...), Shin, Dae Hwan (2025) Dual-Drug Delivery Systems Using Hydrogel–Nanoparticle Composites: Recent Advances and Key Applications. *Gels*. https://doi.org/10.3390/gels11070520

96. Guo, F., Du, Y., Wang, Y., (...), Yang, G. (2024) Targeted drug delivery systems for matrix metalloproteinase-responsive anoparticles in tumor cells: A review. *International Journal of Biological Macromolecules*. https://doi.org/10.1016/j.ijbiomac.2023.128658

97. Hu, Q., Katti, P.S., Gu, Z. (2014) Enzyme-responsive nanomaterials for controlled drug delivery. *Nanoscale*. https://doi.org/10.1039/c4nr04249b

98. Li, M., Zhao, G., Su, W.-K., Shuai, Q. (2020) Enzyme-Responsive Nanoparticles for Anti-tumor Drug Delivery. *Frontiers in Chemistry*. https://doi.org/10.3389/fchem.2020.00647

99. Shirazi, Amir Nasrolahi, Vadlapatla, Rajesh, Koomer, Ajoy, (...), Parang, Keykavous (2025) Peptide-Based Inorganic Nanoparticles as Efficient Intracellular Delivery Systems. *Pharmaceutics*. https://doi.org/10.3390/pharmaceutics17091123

100. Mohd Ali, I., Rungsardthong Ruktanonchai, U.R., Soottitantawat, A., (...), Tang, S.Y. (2025) Pickering emulsions for cancer: Formulation strategies, functional enhancements, and translational considerations. *International Journal of Pharmaceutics*. https://doi.org/10.1016/j.ijpharm.2025.125873

101. Zhu, C.-L., Lu, C.-H., Song, X.-Y., (...), Wang, X.-R. (2011) Bioresponsive controlled release using mesoporous silica nanoparticles capped with aptamer-based molecular gate. *Journal of the American Chemical Society*. https://doi.org/10.1021/ja110094g

102. Yang, B., Liang, H., Xu, J., (...), Wang, C. (2025) Multi-drug sequential release systems: Construction and application for synergistic tumor treatment. *International Journal of Pharmaceutics*. https://doi.org/10.1016/j.ijpharm.2024.125156

103. Zhang, Y., Tan, X., Ren, T., (...), Sun, H. (2018) Folate-modified carboxymethyl-chitosan/polyethylenimine/bovine serum albumin based complexes for tumor site-specific drug delivery. *Carbohydrate Polymers*. https://doi.org/10.1016/j.carbpol.2018.06.055

104. Xiao, W., Zeng, X., Lin, H., (...), Zhang, X.-Z. (2015) Dual stimuli-responsive multi-drug delivery system for the individually controlled release of anti-cancer drugs. *Chemical Communications*. https://doi.org/10.1039/c4cc08831j

105. Wang, H., Liu, R., Wang, S., (...), Zhang, Y. (2021) A highly programmable platform for sequential release of protein therapeutics. *Journal of Materials Chemistry B*. https://doi.org/10.1039/d0tb02657c

106. Jain, K.M.H., Siegel, R.A. (2023) Predetermination of burst times of elastoplastic osmotic capsules. *Journal of Controlled Release*. https://doi.org/10.1016/j.jconrel.2023.03.029

107. Jiang, X., Fan, X., Zhang, R., (...), Wu, G. (2020) In situ tumor-triggered subcellular precise delivery of multi-drugs for enhanced chemo-photothermal-starvation combination antitumor therapy. *Theranostics*. https://doi.org/10.7150/thno.52000

108. Jia, D.-Y., Li, S.-S., Zheng, Z. (2022) Strategy for Cascade Self-assembly/Disassembly of Small Molecules in Cancer Diagnosis and Treatment. *Journal of Instrumental Analysis*. https://doi.org/10.19969/j.fxcsxb.21121701

109. Cao, Y., Wang, B., Wang, Y., Lou, D. (2014) Polymer-controlled core-shell nanoparticles: A novel strategy for sequential drug release. *RSC Advances*. https://doi.org/10.1039/c4ra03610g

110. Arafa, K.K., Smyth, H.D.C., El-Sherbiny, I.M. (2021) Mitotropic triphenylphosphonium doxorubicin-loaded core-shell nanoparticles for cellular and mitochondrial sequential targeting of breast cancer. *International Journal of Pharmaceutics*. https://doi.org/10.1016/j.ijpharm.2021.120936

111. Wang, Z., Ho, P.C. (2010) Self-assembled core-shell vascular-targeted nanocapsules for temporal antivasculature and anticancer activities. *Small*. https://doi.org/10.1002/smll.201001122

112. Chu, L.-Y. (2011) Smart Microcapsules with Thermo-Responsive Hydrogel Membranes. *Advanced Topics in Science and Technology in China*. https://doi.org/10.1007/978-3-642-18114-6_4

113. Chu, L.-Y. (2011) Smart Microcapsules with pH-Responsive Hydrogel Membranes. *Advanced Topics in Science and Technology in China*. https://doi.org/10.1007/978-3-642-18114-6_8

114. Wei, J., Ju, X.-J., Zou, X.-Y., (...), Chu, L.-Y. (2014) Multi-stimuli-responsive microcapsules for adjustable controlled-release. *Advanced Functional Materials*. https://doi.org/10.1002/adfm.201303844

115. Wang, Jianmei, Li, Yan, Wang, Xueying, (...), Wang, Chengyang (2016) Droplet Microfluidics for the Production of Microparticles and Nanoparticles. *Micromachines*. https://doi.org/10.3390/mi8010022

116. Park, J.-Y., Jeong, H.-S., Lim, S.-R., (...), Lee, S.-J. (2026) Triple-emulsion microfluidic Core–Shell hydrogel microcapsules for oral pentoxifylline Delivery: Ameliorating colitis and rebalancing gut microbiome. *Materials Today Bio*. https://doi.org/10.1016/j.mtbio.2026.102881

117. Chavan, Y.R., Tambe, S.M., Jain, D.D., (...), Amin, P.D. (2022) Redefining the importance of polylactide-co-glycolide acid (PLGA) in drug delivery. *Annales Pharmaceutiques Francaises*. https://doi.org/10.1016/j.pharma.2021.11.009

118. Jiang, L., Luo, J., Hong, D., (...), Ge, J. (2022) Recent Advances of Poly(lactic-co-glycolic acid)-Based Nanoparticles for Tumor-Targeted Drug Delivery. *ChemistrySelect*. https://doi.org/10.1002/slct.202103524

119. Elmowafy, E.M., Tiboni, M., Soliman, M.E. (2019) Biocompatibility, biodegradation and biomedical applications of poly(lactic acid)/poly(lactic-co-glycolic acid) micro and nanoparticles. *Journal of Pharmaceutical Investigation*. https://doi.org/10.1007/s40005-019-00439-x

120. Sharma, P., Kaur, N., Shanavas, A. (2023) Targeting strategies using PLGA nanoparticles for efficient drug delivery. *Poly(lactic-co-glycolic acid) (PLGA) Nanoparticles for Drug Delivery*. https://doi.org/10.1016/B978-0-323-91215-0.00016-9

121. Kesharwani, P., Kumar, V., Goh, K.W., (...), Sahebkar, A. (2025) PEGylated PLGA nanoparticles: unlocking advanced strategies for cancer therapy. *Molecular Cancer*. https://doi.org/10.1186/s12943-025-02410-x

122. Elsewedy, H.S., Aldhubiab, B.E.A., Mahdy, M.A., Elnahas, H.M. (2020) Development, optimization, and evaluation of PEGylated brucine-loaded PLGA nanoparticles. *Drug Delivery*. https://doi.org/10.1080/10717544.2020.1797237

123. Hou, W., Mao, W., Sun, J., (...), Tang, S. (2024) Targeting Hydrogel for Intelligent Recognition and Spatiotemporal Control in Cell-Based Therapeutics. *Advanced Science*. https://doi.org/10.1002/advs.202404172

124. Kim, Y., Kwak, J., Lim, M., (...), Lim, K.S. (2025) Advances in PCL, PLA, and PLGA-Based Technologies for Anticancer Drug Delivery. *Pharmaceutics*. https://doi.org/10.3390/pharmaceutics17101354

125. Wang, J., Wang, B.M., Schwendeman, S.P. (2002) Characterization of the initial burst release of a model peptide from poly(D,L-lactide-co-glycolide) microspheres. *Journal of Controlled Release*. https://doi.org/10.1016/S0168-3659(02)00137-2

126. Ravivarapu, H.B., Burton, K., DeLuca, P.P. (2000) Polymer and microsphere blending to alter the release of a peptide from PLGA microspheres. *European Journal of Pharmaceutics and Biopharmaceutics*. https://doi.org/10.1016/S0939-6411(00)00099-0

127. Xiao, M., Lai, W., Wang, F., (...), Pei, H. (2019) Programming Drug Delivery Kinetics for Active Burst Release with DNA Toehold Switches. *Journal of the American Chemical Society*. https://doi.org/10.1021/jacs.9b10765

128. Tang, L., Yang, Z., Zhou, Z., (...), Chen, X. (2019) A logic-gated modular nanovesicle enables programmable drug release for on-demand chemotherapy. *Theranostics*. https://doi.org/10.7150/thno.32106

129. Zhang, B., Zhang, Y., Dang, W., (...), Liu, Z. (2022) The anti-tumor and renoprotection study of E-[c(RGDfK)2]/folic acid co-modified nanostructured lipid carrier loaded with doxorubicin hydrochloride/salvianolic acid A. *Journal of Nanobiotechnology*. https://doi.org/10.1186/s12951-022-01628-x

130. Choi, D.H., Subbiah, R., Kim, I.H., (...), Park, K. (2013) Dual growth factor delivery using biocompatible core-shell microcapsules for angiogenesis. *Small*. https://doi.org/10.1002/smll.201300427

131. Lian, Heping, Ma, Shuang, Zhao, Duoyi, (...), Zhang, Zhiyu (2022) Cytokine Therapy Combined with Nanomaterials Participates in Cancer Immunotherapy. *Pharmaceutics*. https://doi.org/10.3390/pharmaceutics14122606

132. Chen, L., Song, M.-H., Jin, M., (...), Huang, W. (2026) Thymine-functionalized bPEI nanocomplexes for synergistic STING activation and PD-L1 silencing to potentiate cancer immunotherapy. *Journal of Controlled Release*. https://doi.org/10.1016/j.jconrel.2026.115372

133. Tripathi, Diksha, Mondal, Moumita, Sakthivel, Natarajan (2025) Advanced materials for triggered release of drugs. *Medical Nanobiotechnology*. https://doi.org/10.1016/B978-0-443-21507-0.00003-9

134. Zhu, X., Sun, H., Cao, X., Xia, J. (2026) Targeting strategies of lipid nanoparticles for RNA therapeutics: From hepatic tropism to extrahepatic precision delivery. *Critical Reviews in Oncology/Hematology*. https://doi.org/10.1016/j.critrevonc.2026.105448

135. Hussain, Z., Abdulelah, A.A., Tahseen, S.H., (...), Thu, H.E. (2026) Stimuli-responsive nanomedicines for lung cancer therapy: Design strategies, advances, challenges, and future directions. *Journal of Drug Delivery Science and Technology*. https://doi.org/10.1016/j.jddst.2025.107900

136. Carregal-Romero, S., Ochs, M., Rivera Gil, P., (...), Parak, W.J. (2012) NIR-light triggered delivery of macromolecules into the cytosol. *Journal of Controlled Release*. https://doi.org/10.1016/j.jconrel.2011.12.013

137. Harada, S., Ehara, S., Ishii, K., (...), Goto, S. (2014) Imaging and treatment of malignant metastatic tumors by using radiation-sensitive, immunolabeled liquid-core microcapsules. *Nuclear Instruments and Methods in Physics Research, Section B: Beam Interactions with Materials and Atoms*. https://doi.org/10.1016/j.nimb.2013.06.045

138. Harada, S., Ehara, S., Ishii, K., (...), Ito, J. (2009) Targeted Delivery of Chemotherapeutic Agents Using Improved Radiosensitive Liquid Core Microcapsules and Assessment of Their Antitumor Effect. *International Journal of Radiation Oncology Biology Physics*. https://doi.org/10.1016/j.ijrobp.2009.02.082

139. Lyon, P.C., Griffiths, L.F., Lee, J., (...), Coussios, C.C. (2017) Clinical trial protocol for TARDOX: A phase I study to investigate the feasibility of targeted release of lyso-thermosensitive liposomal doxorubicin (ThermoDox®) using focused ultrasound in patients with liver tumours. *Journal of Therapeutic Ultrasound*. https://doi.org/10.1186/s40349-017-0104-0

140. Staruch, R.M., Ganguly, M., Tannock, I.F., (...), Chopra, R. (2012) Enhanced drug delivery in rabbit VX2 tumours using thermosensitive liposomes and MRI-controlled focused ultrasound hyperthermia. *International Journal of Hyperthermia*. https://doi.org/10.3109/02656736.2012.736670

141. https://www.sciencedirect.com/science/article/pii/S1559131X25005335

142. Da, X., Cao, B., Mo, J., (...), Yang, Y. (2024) Inhibition of growth of hepatocellular carcinoma by co-delivery of anti-PD-1 antibody and sorafenib using biomimetic nano-platelets. *BMC Cancer*. https://doi.org/10.1186/s12885-024-12006-1

143. Jing, L., Zhao, S., Sun, C., (...), Wang, W. (2026) Biomimetic self-assembled copper nanoadjuvant potentiates hepatocellular carcinoma immunotherapy via cuproptosis-driven cGAS-STING activation. *Materials Today Bio*. https://doi.org/10.1016/j.mtbio.2026.103325

144. Wu, Y., Qu, H., Li, X., (...), Wu, X. (2025) Excessive autophagy-inducing and highly penetrable biomineralized bacteria for multimodal imaging-guided and mild hyperthermia-enhanced immunogenic cell death. *Journal of Colloid and Interface Science*. https://doi.org/10.1016/j.jcis.2024.09.246

145. Mohaghegh, Neda, Zargar Balajam, Narges, Kraemer, Nick, (...), Hassani Najafabadi, Alireza (2026) Interleukin-4-Loaded lipid nanoparticles reprogram adipose tissue immunity and promote adipocyte browning for the treatment of obesity. *Acta Biomaterialia*. https://doi.org/10.1016/j.actbio.2026.09.007

146. Zhou, Y., Chen, X., Cao, J., Gao, H. (2020) Overcoming the biological barriers in the tumor microenvironment for improving drug delivery and efficacy. *Journal of Materials Chemistry B*. https://doi.org/10.1039/d0tb00649a

147. Heldin, C.-H., Rubin, K., Pietras, K., Oštman, A. (2004) High interstitial fluid pressure - An obstacle in cancer therapy. *Nature Reviews Cancer*. https://doi.org/10.1038/nrc1456

148. Matsumura, Y., Maeda, H. (1986) A New Concept for Macromolecular Therapeutics in Cancer Chemotherapy: Mechanism of Tumoritropic Accumulation of Proteins and the Antitumor Agent Smancs. *Cancer Research*. https://www.scopus.com/pages/publications/0022858683

149. Maeda, H., Wu, J., Sawa, T., (...), Hori, K. (2000) Tumor vascular permeability and the EPR effect in macromolecular therapeutics: A review. *Journal of Controlled Release*. https://doi.org/10.1016/S0168-3659(99)00248-5

150. Lammers, T., Kießling, F., Hennink, W.E., Storm, G. (2012) Drug targeting to tumors: Principles, pitfalls and (pre-) clinical progress. *Journal of Controlled Release*. https://doi.org/10.1016/j.jconrel.2011.09.063

151. Maeda, H. (2015) Toward a full understanding of the EPR effect in primary and metastatic tumors as well as issues related to its heterogeneity. *Advanced Drug Delivery Reviews*. https://doi.org/10.1016/j.addr.2015.01.002

152. Yadav, P., Yadav, A.K., Devidas Bhatia, D. (2025) Nanotechnology Meets the Tumor Microenvironment: Unlocking New Horizons in Cancer Therapy. *ACS Biomaterials Science and Engineering*. https://doi.org/10.1021/acsbiomaterials.5c01120