

Week 3 – Workshop

Exploring & Understanding Inversion



Geophysics 3A: Potential Fields & Geothermics

July 8, 2026

Duration: 3 hours

Setting: Lecture room with laptops (sphere inversion app)

Preparation

Pre-reading: Course Notes Ch. 13

Lecture videos:

Notes:

Purpose

This workshop introduces the structure of inverse problems in potential-field geophysics, contrasting forward prediction with inverse estimation. Students explore the role of misfit, data weighting, and regularization, and examine how non-linearity and non-uniqueness arise in gravity modelling. Through a sequence of activities, they classify linear vs. non-linear models, reason qualitatively about non-uniqueness, and engage in hands-on inversion of a buried sphere using grid search and Gauss–Newton methods.

Learning Outcomes

- Distinguish forward and inverse problems and articulate the role of misfit, weighting, and regularization.
- Classify simple geophysical relations as linear or non-linear and propose appropriate linearizations.
- Recognize manifestations of non-uniqueness in gravity and magnetic profiles and propose discriminating data.
- Derive or confirm analytical Jacobian expressions for the buried sphere model.
- Interpret misfit surfaces, inversion paths, and the effect of noise and damping.
- Explain parameter trade-offs and resolution limitations inherent to the buried-sphere problem.

Session Flow (170 min)

Create mixed groups (4–5 students).

0–10 min: Warm-up

What can you learn from a shadow?

- What can you determine directly from the shadow?
- What can you estimate only if additional assumptions are made?
- What information cannot be determined uniquely?
- What prior knowledge are you using when deciding what the object might be?

How does this relate to inversion?

- A shadow does not uniquely determine the object that created it.
- Likewise, a gravity anomaly does not uniquely determine the density distribution that produced it.
- The challenge of inversion is deciding which of many acceptable models is most reasonable.

10–40 min: Mini-lecture — Forward vs Inverse, Misfit, Weighting, Challenges

- Forward vs. inverse models: $m \mapsto g(m)$ vs. $d \mapsto m$.
- Misfit: residual $r = d - g(m)$; weighted least squares $\Phi_d = \|W_d r\|_2^2$.
- Role of W_d (noise) and W_m (model constraints); damping λ .
- Challenges: non-linearity (Jacobian needed), non-uniqueness (multiple m give similar $g(m)$), ill-conditioning.

40–60 min: Activity 3.A: Testing Linearity

Two conditions must be met for linearity. Let's explore how to rigorously test for linearity before we work on other functions.

60–90 min: Activity 3.B: Linear or Non-linear?

Using the equation handout:

- Identify model parameters and classify each relation as *linear*, *non-linear*, or *linearizable*.
- If non-linear, propose a linearization (log-transform, re-parameterization, Taylor expansion/Jacobian).
- Deliverable: brief notes per group.

70–85 min: Activity 3.C: Buried Sphere — Background

We're going to look at the gravity anomaly from a buried sphere. The forward problem is non-linear with respect to the model parameters, requiring a non-linear inverse solution. This activity is broken into two parts: a look at the formulation, and an investigation using a simple program.

- App overview: Data & Model view; Misfit view; controls for acquisition, noise, synthetic model.
- Construct a true model & generate data (freeze a single noise realization).

85–105 min: Buried Sphere — Jacobian Derivatives

1. Students derive/confirm analytical expressions for $\partial g / \partial z$, $\partial g / \partial R$, $\partial g / \partial \Delta \rho$.
2. Compare with the app's Jacobian; discuss relative sensitivities: amplitude $\propto R^3 \Delta \rho$, width controlled by depth z .

105–150 min: Buried Sphere — Investigation

With the use of `buried_sphere_gui.py`, we will examine a number of aspects of inversion, including

Grid search: Explore pairwise misfit slices and 3-D misfit surface. Identify local vs global minima.

Inversion path: Run Gauss–Newton with different λ ; examine step length and stability.

Effect of noise: Increase σ ; observe widening misfit basins and parameter uncertainty.

Non-uniqueness: Demonstrate parameter trade-offs (e.g. $a^3 \Delta \rho$) and resolution ellipses.

Role of Noise: Look at how noise affects results.

150–160 min: Debrief

- Key takeaways: forward/inverse distinctions, misfit and weighting, non-linearity, non-uniqueness, damping.
- Open Q&A; preview of next week's material.

160–170 min: Preview: Gravity Theory

This week we treated gravity primarily as data to be inverted. Next week we will begin developing gravity theory itself.

We will derive the gravity field from Newton's law of universal gravitation and examine how mass distributions create gravity anomalies. We will see how simple geological bodies such as spheres, cylinders, slabs, and contacts produce characteristic gravity signatures.

Along the way, we will connect the mathematical concepts introduced in Week 1—gradients, curvature, and potential fields—to one of the most important geophysical methods.

By the end of next week, you will be able to predict how changes in density, geometry, and observation distance affect the gravity field and begin interpreting gravity anomalies in geological terms.

Materials

- Linear/non-linear handout or slide deck.
- Activity D handouts (gravity and magnetic curves-only plots).
- Buried sphere app (laptop per group).
- Whiteboard or shared notepad for Jacobian derivations.

Assessment

- Activity notes (classification & argumentation).
- Jacobian derivation steps.
- Short reflection or exit-ticket on non-linearity and non-uniqueness.

References (suggested for students)**Further Reading:**

- [1] R. Blakely, *Potential Theory in Gravity and Magnetic Applications*, Cambridge University Press, 1995.
- [2] W. M. Telford, L. P. Geldart, and R. E. Sheriff, *Applied Geophysics*, 2nd ed., Cambridge University Press, 1990.
- [3] W. Menke, *Geophysical Data Analysis: Discrete Inverse Theory*, 3rd ed., Academic Press, 2012.
- [4] R. C. Aster, B. Borchers, and C. H. Thurber, *Parameter Estimation and Inverse Problems*, 3rd ed., Elsevier, 2019.
- [5] A. Tarantola, *Inverse Problem Theory and Methods for Model Parameter Estimation*, SIAM, 2005.

Workshop Notes

Workshop Notes

Activity 3.A

Testing Linearity

Purpose

Linear functions can be solved in a single inversion step using linear least squares. This method is simpler, faster, and less computationally expensive, so when possible a linear function is preferred. In this activity, you'll learn to determine whether a function is linear or non-linear.

Learning Objectives

By the end of this activity you should be able to:

- describe the two conditions for linearity, and
- be able to rigorously demonstrate whether a function is linear or non-linear using these tests.

Conditions for Linearity

There are two conditions that must be met for a function to be linear: homogeneity,

$$f(\alpha m_1) = \alpha f(m_1),$$

and additivity

$$f(m_1 + m_2) = f(m_1) + f(m_2).$$

Note: something that may take some getting used to is that our tests for linearity are with respect to model parameters that we are trying to estimate via inversion, not the experimental conditions or observation position, which is how we otherwise normally think of functions.

Box .1: Example of a linear function

For instance, $y = ax + b$ is linear with respect to the model parameters $\mathbf{m} = (a, b)$. For homogeneity,

$$\begin{aligned} y(\alpha \mathbf{m}) &= (\alpha a)x + (\alpha b) \\ &= \alpha(ax + b) \\ &= \alpha y(\mathbf{m}). \end{aligned}$$

For additivity, where $\mathbf{m}_1 = (a_1, b_1)$ and $\mathbf{m}_2 = (a_2, b_2)$,

$$\begin{aligned} y(\mathbf{m}_1 + \mathbf{m}_2) &= (a_1 + a_2)x + (b_1 + b_2) \\ &= (a_1x + b_1) + (a_2x + b_2) \\ &= y(\mathbf{m}_1) + y(\mathbf{m}_2). \end{aligned}$$

Since both homogeneity and additivity are satisfied, the function is linear.

Question 1. Linear function

Show that

$$k(\text{SiO}_2) = a(\text{SiO}_2)^2 + b(\text{SiO}_2) + c$$

is linear with respect to the model parameters, $\mathbf{m} = (a, b, c)$.


Question 2. Non-linear function

Show that

$$k(T) = k_0 \left(\frac{298}{T} \right)^n$$

is non-linear with respect to the model parameters, $\mathbf{m} = (k_0, n)$.

Question 3. Why is the first equation easier to invert than the second?



Activity 3.B

Linear or Non-linear?

Goal. For each of the following forward relations drawn from potential fields, geothermics, and physical property relations, decide whether it is (i) linear, (ii) non-linear, or (iii) non-linear but can be linearized. If non-linear, describe how you would linearize or re-parameterize it. Justify your choice with respect to the model parameters that would be inverted.

Inversion problems can be cast as $\mathbf{A}\mathbf{m} = \mathbf{d}$ for linear problems and $A(m) = d$ for non-linear problems. Where d is the data and A is the operator or design (more on that below). The difference may appear subtle, though in reality the difference is that the linear equation can be written as a linear algebra problem as the model parameters are independent and discrete. In the non-linear formulation, the one or more model parameters are dependent upon another. As a result they cannot be fully separated unless they can be linearized.

Let's explore what A , m , and d are a bit more intuitively. The data, d , are the results of an experiment. Identifying the data are generally straightforward. The operator and model parameters can take a bit more getting used to. The model parameters, m , are the physical or virtual properties of the system and initially unknown. Once you determine these properties, the forward model becomes a prediction. The operator or design matrix, A , can be thought of *as the experiment itself*, and is dependent upon the geometry, conditions of the experiment, and the physics. The model tells us what exists, the operator tells us how we observe it.

When we are doing a geophysical experiment with the Earth as the medium,

- $\mathbf{m} \rightarrow$ the model is Earth, what it is made of and its physical properties. Changing $\mathbf{m}$ changes the Earth.
- $\mathbf{A} \rightarrow$ how Earth talks to the instrument, how the Earth is sampled by the experiment, i.e., the rules of the experiment. Changing $\mathbf{A}$ changes the experiment, i.e., same Earth + different experiment $\rightarrow$ different data.
- $\mathbf{d} \rightarrow$ the data is the outcome of the experiment, what we measure

In most geophysical problems, $\mathbf{A}$ is

- gravity – distances from mass elements
- magnetics – sensor positions and dip angle
- heat flow – layer thicknesses and boundary conditions
- electromagnetics – distances from conductive/resistive bodies
- seismology – ray paths or travel times

So the entries in $\mathbf{A}$ are determined by:

- positions
- distances
- times
- physical laws
- boundary conditions

Put simply, the matrix $\mathbf{A}$ encodes the physics and geometry of how the experiment samples the model, $\mathbf{m}$, resulting in the observations, $\mathbf{d}$.

Instructions (15–20 min)

1. Work in groups of 3–4.
2. Identify the data and the experimental quantities.
3. For each equation

- identify the quantity or quantities that would be treated as the *model parameter(s)* in an inversion.
- Classify the mapping data → model as linear or non-linear in those parameters.
- If non-linear, propose a linearization (e.g., log-transform, Taylor expansion/Jacobian around p_0 , or a change of variables) and note any loss of identifiability.

Equations

- Q1. **Gravity (Bouguer slab):** We'll see this equation again as it is used to model the gravity due to an infinite slab.

$$\Delta g = 2\pi G(\rho_s - \rho_c)H$$

- Q2. **Gravity (buried sphere):** This gravity solution is one of the simplest models for a gravity anomaly. We will explore it further.

$$g(r) = \frac{4}{3}\pi a^3 G(\rho_s - \rho_c) \frac{z}{(r^2 + z^2)^{3/2}}$$

- Q3. **Thermal conductivity $k(T, X)$:** We'll use this formulation in the problem set. This equation is used to estimate thermal conductivity model for a solid solution series for a mineral.

$$k(T, X) = (a + bX + cX^2) (298/T)^n$$

- Q4. **Thermal conductivity $k(T, P)$:** This form of thermal conductivity is often used to estimate thermal conductivity for a rock.

$$k(T, P) = \frac{k_0}{A + BT} (1 + cP)$$

- Q5. **Isostasy (Airy):** Airy isostasy is an end-member isostatic model, used to explain topographic variations due to differences in crustal thickness.

$$\Delta t = \frac{\rho_c}{\rho_m - \rho_c} h$$

- Q6. **Isostatic gravity correction (column):** Gravity can be corrected for a mass deficit created by variations

in crustal thickness via an isostatic correction.

$$\Delta g_{\text{iso}} \approx 2\pi G(\rho_m - \rho_c) \Delta t - 2\pi G \rho_c h$$

- Q7. **Magnetics (2-D infinite dike):** The ocean floor records chaotic reversals in Earth's magnetic field. The remnant magnetic field is “frozen” in dikes produced at the mid-ocean ridge.

$$\Delta T(x) = C \left[\arctan\left(\frac{x-x_1}{z}\right) - \arctan\left(\frac{x-x_2}{z}\right) \right]$$

- Q8. **Layered 1-D heat with sources:** Geotherms can largely be explained by 1-D heat flow through a layered Earth. We'll see this again when we start geothermics.

$$T_i(z) = T_{i-1} + \frac{q_{i-1}}{k_i}(z_i - z_{i-1}) - \frac{A_i}{2k_i}(z_i - z_{i-1})^2$$

- Q9. **Density-porosity mix:** Sedimentary rocks include pore space (porosity) that may be filled with a fluid (air, water, brine, gas, oil, etc.).

$$\rho_{\text{bulk}} = (1 - \phi) \rho_{\text{matrix}} + \phi \rho_{\text{fluid}}$$

- Q10. **Gardner's relation (density-velocity):** Many solutions in geophysics involve trying to estimate one property from another, in this case density from seismic velocity.

$$\rho = a v^b$$

- Q11. **Gravity (discrete 2-D prisms):** While we can produce analytic equations for a variety of idealized geologic bodies with simple geometries (e.g., buried spheres as above), to produce more geologically realistic bodies we can produce structures from arbitrarily-shaped polygons.

$$g_z = 2G\Delta\rho \sum_{i=1}^N \left[(x_{i+1} - x_i) \log\left(\frac{r_{i+1}}{r_i}\right) + (z_{i+1} - z_i) \arctan\left(\frac{x_i z_{i+1} - x_{i+1} z_i}{x_i x_{i+1} + z_i z_{i+1}}\right) \right]$$

where

$$r_i = \sqrt{x_i^2 + z_i^2}, \quad (x_{N+1}, z_{N+1}) = (x_1, z_1).$$

Activity 3.C

Inversion of a Buried Sphere

Purpose

The purpose of this activity is to develop an intuitive understanding of inversion using the gravity anomaly of a buried sphere. By comparing grid-search and gradient-based approaches, you will explore how inverse methods search model space, how model parameterization influences the solution, and why non-uniqueness and noise make geophysical interpretation challenging.

Learning Outcomes

By the end of this activity, you should be able to:

- Explain the difference between forward modelling, grid search, and gradient-based inversion.
- Describe the role of the Jacobian matrix in determining model updates.
- Identify sources of non-uniqueness in geophysical inverse problems.
- Explain how model parameterization can improve or degrade inversion performance.
- Describe the influence of noise on inversion results and uncertainty.

Gravity of a Buried Sphere

We will return to how we derive the gravity due to a buried sphere later. For now, we will model the vertical gravity anomaly at the surface produced by a buried sphere of depth z (to center), radius a , and density contrast $\Delta\rho$. For a station at horizontal radius $r = \sqrt{x^2 + y^2}$,

$$g(r|z_c, a, \Delta\rho) = \underbrace{\frac{4}{3}\pi a^3 \Delta\rho}_{\text{mass}} G \frac{z}{(r^2 + z_c^2)^{3/2}}, \quad (1)$$

with G the gravitational constant.

Inverse Model

The gravity equation due to a buried sphere is non-linear, requiring a non-linear iterative solver. The code we will use employs the Gauss–Newton inverse method, which can be enhanced with Levenberg–Marquardt regularization.

We seek $\mathbf{m} = [z_c, a, \Delta\rho]^T$ that minimizes the objective function,

$$\Phi(m) = \underbrace{\|W_d(\mathbf{g}_{\text{obs}} - g(\mathbf{m}))\|_2^2}_{\text{model misfit}} + \underbrace{\lambda^2 \|\mathbf{m} - \mathbf{m}_0\|_2^2}_{\text{regularization term}},$$

where $\mathbf{g}_{\text{obs}}$ are the observed gravity data, $g(\mathbf{m})$ is the predicted gravity using Equation 1, $\mathbf{W}_d$ is a diagonal matrix of inverse variance (data uncertainty), and λ is the Levenberg–Marquardt regularization parameter. Our initial guess is given by the column vector $\mathbf{m}_0$.

To determine $\mathbf{m}$, we have to solve iteratively, with each model step being

$$\Delta\mathbf{m} = (\mathbf{J}^T \mathbf{W}_d^2 \mathbf{J} + \lambda^2 \mathbf{I})^{-1} \mathbf{J}^T \mathbf{W}_d^2 (\mathbf{g}_{\text{obs}} - g(\mathbf{m}_0)).$$

Role of λ . Small λ approaches Gauss–Newton (fast, but can be unstable if $J^\top J$ is ill-conditioned). Large λ yields smaller, more stable steps (gradient-descent-like), helpful far from the solution or with strong trade-offs.

Question 1. Jacobian

Before we play with the buried sphere program, let's derive derivatives we need for the Jacobian matrix.

The Jacobian for the buried sphere requires three derivatives:

$$\mathbf{J} = \begin{bmatrix} \frac{\partial g}{\partial z_c} & \frac{\partial g}{\partial a} & \frac{\partial g}{\partial \Delta\rho} \end{bmatrix}.$$

Derive the derivatives from Equation 1.

Exploration

The questions below relate to the `buried_sphere_gui.py` program. While the questions are about the demo, the answers are largely independent of both the problem and program.

Question 2. Grid Search

What is a grid search? Is it a forward or inverse driven solution? What does it tell us about the misfit? How do we determine the model parameters?

Question 3. Inversion

How does inversion approach the problem differently from the grid search? What path does it take to the answer?

Question 4. Comparison

How does the result of the inversion compare with that of the grid search? If the two solutions do not match, why?

Question 5. Non-uniqueness

The problem we have posed is non-unique even though we use a single "true" model to generate the data. Why is the solution non-unique?

Question 6. 2D versus 3D

Let's switch from 3 parameters to 2. The 2 parameter result is a better constrained, why? How is this improvement in constraint reflected in the misfit surfaces?

Question 7. Noise

How does noise affect the solution via inversion? How does noise contribute to non-uniqueness?

For this demonstration, repeat the inversion with 0%, 5%, and 10% Gaussian noise.

Key Ideas

Grid search is a brute-force forward modelling approach to systematically sample the model space with misfit is computed at every combination of model parameters. It can identify the global minimum within the explored model domain, but it becomes computationally expensive as the number of model parameters increases and requires a sufficiently dense sampling of the model space to accurately resolve the misfit surface. For many large difficult problems found in geophysics, a grid search is unfeasible.

Inversion uses information about the local shape of the misfit surface, typically through the Jacobian matrix, to move iteratively toward a low-misfit solution. While each iteration is computationally more expensive than a single forward calculation, inversion generally requires far fewer model evaluations than a grid search.

Best of all, inversion remains practical for geophysical problems with many parameters (tens to millions). The trade-off is that inversion explores only a small portion of model space and may converge to a local minimum rather than the global minimum.




Figure 1: Comparison of grid-search and gradient-based inversion approaches.