

MAT 202 - Probability and Random Processes

Lecture - 5 and 6



Ahmedabad
University

Dr. Dhaval Patel

Computer Science and Engineering (CSE),
SEAS-Ahmedabad University,
Ahmedabad, Gujarat, India

January 22, 2020

Outline

- **Joint and Conditional Probability**

Outline

- **Joint and Conditional Probability**
 - Motivation
 - How does one compute it?
 - Example

Outline

- **Joint and Conditional Probability**
 - Motivation
 - How does one compute it?
 - Example
- **Bayes Theorem**

Outline

- **Joint and Conditional Probability**

- Motivation
- How does one compute it?
- Example

- **Bayes Theorem**

- Theorem 2.9: Conditional Probability with given non-zero likelihood of an event
- Theorem 2.10: Total Probability Theorem
- Theorem 2.11: Bayes Theorem
- Examples

- **Independence**

Outline

- **Joint and Conditional Probability**

- Motivation
- How does one compute it?
- Example

- **Bayes Theorem**

- Theorem 2.9: Conditional Probability with given non-zero likelihood of an event
- Theorem 2.10: Total Probability Theorem
- Theorem 2.11: Bayes Theorem
- Examples

- **Independence**

- Definition
- Examples

- **Random Variables**

Outline

- **Joint and Conditional Probability**

- Motivation
- How does one compute it?
- Example

- **Bayes Theorem**

- Theorem 2.9: Conditional Probability with given non-zero likelihood of an event
- Theorem 2.10: Total Probability Theorem
- Theorem 2.11: Bayes Theorem
- Examples

- **Independence**

- Definition
- Examples

- **Random Variables**

- Definition: RV and PMF
- Types of RVs
- Examples

Introduction to Probability Theory

- Concepts of Joint Probability

① Motivation:

Introduction to Probability Theory

- Concepts of Joint Probability

- 1 **Motivation:** All events are not mutually exclusive

Introduction to Probability Theory

- Concepts of Joint Probability

- ① **Motivation:** All events are not mutually exclusive
- ② **Joint Probability of Event A and B :**
 - We are interested in calculating the probability of the intersection of two events, $A \cap B$. This probability is referred to as the joint probability of the events A and B , $Pr(A \cap B)$
 - Denoted as: $Pr(A, B)$

Introduction to Probability Theory

- Concepts of Joint Probability

① **Motivation:** All events are not mutually exclusive

② **Joint Probability of Event A and B :**

- We are interested in calculating the probability of the intersection of two events, $A \cap B$. This probability is referred to as the joint probability of the events A and B , $Pr(A \cap B)$
- Denoted as: $Pr(A, B)$
- The joint probability of the events A_1, A_2, \dots, A_M is

$$Pr(A_1 \cap A_2 \cap A_3 \cdots A_M)$$

- Simpler Notation: $Pr(A_1, A_2, A_3 \cdots A_M)$

Introduction to Probability Theory

- Concepts of Joint Probability

① **Motivation:** All events are not mutually exclusive

② **Joint Probability of Event A and B :**

- We are interested in calculating the probability of the intersection of two events, $A \cap B$. This probability is referred to as the joint probability of the events A and B , $Pr(A \cap B)$
- Denoted as: $Pr(A, B)$
- The joint probability of the events A_1, A_2, \dots, A_M is

$$Pr(A_1 \cap A_2 \cap A_3 \cdots A_M)$$

- Simpler Notation: $Pr(A_1, A_2, A_3 \cdots A_M)$

③ **How does one compute it?:**

- Recall Axiom-3: What is the value of $Pr(A, B)$?

Introduction to Probability Theory

- Concepts of Joint Probability

① **Motivation:** All events are not mutually exclusive

② **Joint Probability of Event A and B :**

- We are interested in calculating the probability of the intersection of two events, $A \cap B$. This probability is referred to as the joint probability of the events A and B , $Pr(A \cap B)$
- Denoted as: $Pr(A, B)$
- The joint probability of the events A_1, A_2, \dots, A_M is

$$Pr(A_1 \cap A_2 \cap A_3 \cdots A_M)$$

- Simpler Notation: $Pr(A_1, A_2, A_3 \cdots A_M)$

③ **How does one compute it?:**

- Recall Axiom-3: What is the value of $Pr(A, B)$?
- Answer : 0

Introduction to Probability Theory

- Concepts of Joint Probability (Continue..)

① How does one compute it?:

- We are more interested in calculating the joint probability of events that are not mutually exclusive.
- Two ways: **Classical** and **Relative Frequency approach**

Introduction to Probability Theory

- Concepts of Joint Probability (Continue..)

① How does one compute it?:

- We are more interested in calculating the joint probability of events that are not mutually exclusive.
- Two ways: **Classical** and **Relative Frequency approach**

② Classical:

- Both events A and B can be expressed in terms of atomic outcomes.

Introduction to Probability Theory

- Concepts of Joint Probability (Continue..)

① How does one compute it?:

- We are more interested in calculating the joint probability of events that are not mutually exclusive.
- Two ways: **Classical** and **Relative Frequency approach**

② Classical:

- Both events A and B can be expressed in terms of atomic outcomes.
- We then write as the set of those atomic outcomes that are common to both

Introduction to Probability Theory

- Concepts of Joint Probability (Continue..)

① How does one compute it?:

- We are more interested in calculating the joint probability of events that are not mutually exclusive.
- Two ways: **Classical** and **Relative Frequency approach**

② Classical:

- Both events A and B can be expressed in terms of atomic outcomes.
- We then write as the set of those atomic outcomes that are common to both
- Calculate the probabilities of each of these outcomes.

Introduction to Probability Theory

- Concepts of Joint Probability (Continue..)

① How does one compute it?:

- We are more interested in calculating the joint probability of events that are not mutually exclusive.
- Two ways: **Classical** and **Relative Frequency approach**

② Classical:

- Both events A and B can be expressed in terms of atomic outcomes.
- We then write as the set of those atomic outcomes that are common to both
- Calculate the probabilities of each of these outcomes.

③ **Relative Frequency**: Let $n_{A,B}$ be the number of times that A and B simultaneously occur in n trials:





















































$$Pr(A, B) = \lim_{n \rightarrow \infty} \frac{n_{A,B}}{n}$$

④ **Example**:

Introduction to Probability Theory

- Example 1: Joint Probability

Example set of 52 playing cards; 13 of each suit clubs, diamonds, hearts, and spades

	Ace	2	3	4	5	6	7	8	9	10	Jack	Queen	King
Clubs													
Diamonds													
Hearts													
Spades													

$A = \{\text{Red card selected}\}$, $B = \{\text{number card selected}\}$, $C = \{\text{heart card selected}\}$

Find: $\Pr(A)$, $\Pr(B)$, $\Pr(C)$, $\Pr(A,B)$, $\Pr(A,C)$, $\Pr(B,C)$

Introduction to Probability Theory

- Concepts of Conditional Probability

- 1 **Motivation:** Often the occurrence of one event may be dependent upon the occurrence of another

Introduction to Probability Theory

- Concepts of Conditional Probability

- 1 **Motivation:** Often the occurrence of one event may be dependent upon the occurrence of another
- 2 **Conditional Probability of Event A and B :**
 - The probability of A conditioned on knowing that B occurred is

$$Pr(A|B) = \frac{Pr(A, B)}{Pr(B)}$$

Introduction to Probability Theory

- Concepts of Conditional Probability

① **Motivation:** Often the occurrence of one event may be dependent upon the occurrence of another

② **Conditional Probability of Event A and B :**

- The probability of A conditioned on knowing that B occurred is

$$Pr(A|B) = \frac{Pr(A, B)}{Pr(B)}$$

$$Pr(B|A)Pr(A) = Pr(A|B)Pr(B)$$

Introduction to Probability Theory

- Concepts of Conditional Probability

- ① **Motivation:** Often the occurrence of one event may be dependent upon the occurrence of another

- ② **Conditional Probability of Event A and B :**

- The probability of A conditioned on knowing that B occurred is

$$Pr(A|B) = \frac{Pr(A, B)}{Pr(B)}$$

$$Pr(B|A)Pr(A) = Pr(A|B)Pr(B)$$

- The conditional probability of three events:

Introduction to Probability Theory

- Concepts of Conditional Probability

- ① **Motivation:** Often the occurrence of one event may be dependent upon the occurrence of another

- ② **Conditional Probability of Event A and B :**

- The probability of A conditioned on knowing that B occurred is

$$Pr(A|B) = \frac{Pr(A, B)}{Pr(B)}$$

$$Pr(B|A)Pr(A) = Pr(A|B)Pr(B)$$

- The conditional probability of three events:

$$Pr(A, B, C) = Pr(C|A, B)Pr(A, B) = Pr(C|A, B)Pr(B|A)Pr(A)$$

- Conditional probability of M events:

Introduction to Probability Theory

- Concepts of Conditional Probability

- 1 **Motivation:** Often the occurrence of one event may be dependent upon the occurrence of another

- 2 **Conditional Probability of Event A and B :**

- The probability of A conditioned on knowing that B occurred is

$$Pr(A|B) = \frac{Pr(A, B)}{Pr(B)}$$

$$Pr(B|A)Pr(A) = Pr(A|B)Pr(B)$$

- The conditional probability of three events:

$$Pr(A, B, C) = Pr(C|A, B)Pr(A, B) = Pr(C|A, B)Pr(B|A)Pr(A)$$

- Conditional probability of M events:

$$Pr(A_1, A_2, A_3 \cdots A_M) = P(A_M|A_1, A_2, \cdots A_{M-1}) \times \\ P(A_{M-1}|A_1, A_2, A_3 \cdots A_{M-2}) \cdots Pr(A_2|A_1)$$

Introduction to Probability Theory

- Concepts of Conditional Probability

- ① **Motivation:** Often the occurrence of one event may be dependent upon the occurrence of another

- ② **Conditional Probability of Event A and B :**

- The probability of A conditioned on knowing that B occurred is

$$Pr(A|B) = \frac{Pr(A, B)}{Pr(B)}$$

$$Pr(B|A)Pr(A) = Pr(A|B)Pr(B)$$

- The conditional probability of three events:

$$Pr(A, B, C) = Pr(C|A, B)Pr(A, B) = Pr(C|A, B)Pr(B|A)Pr(A)$$

- Conditional probability of M events:





















































$$Pr(A_1, A_2, A_3 \cdots A_M) = P(A_M|A_1, A_2, \cdots A_{M-1}) \times \\ P(A_{M-1}|A_1, A_2, A_3 \cdots A_{M-2}) \cdots Pr(A_2|A_1)$$

Introduction to Probability Theory

- Example #2: Conditional Probability: Drawing cards from a deck

Suppose now that we select two cards at random from the deck. When we select the second card, we do not return the first card to the deck. In this case, we say that we are selecting cards without replacement. **Compute the Joint Probability** of event A : First card was a spade and B : Second card was a spade

Example set of 52 playing cards; 13 of each suit clubs, diamonds, hearts, and spades

	Ace	2	3	4	5	6	7	8	9	10	Jack	Queen	King
Clubs													
Diamonds													
Hearts													
Spades													





















































Introduction to Probability Theory

- Example #3: Conditional Probability: Game of Poker

You are dealt 5 cards from a standard 52-card deck. A flush is when you are dealt all five cards of the same suit.

- (1) What is the probability you are dealt a flush in spades?
- (2) What is the probability of a flush in any suit?

Example set of 52 playing cards; 13 of each suit clubs, diamonds, hearts, and spades

	Ace	2	3	4	5	6	7	8	9	10	Jack	Queen	King
Clubs													
Diamonds													
Hearts													
Spades													





















































Introduction to Probability Theory

- Example #3: Conditional Probability: Game of Poker

You are dealt 5 cards from a standard 52-card deck. A flush is when you are dealt all five cards of the same suit.

- (1) What is the probability you are dealt a flush in spades?
- (2) What is the probability of a flush in any suit?

Example set of 52 playing cards; 13 of each suit clubs, diamonds, hearts, and spades

	Ace	2	3	4	5	6	7	8	9	10	Jack	Queen	King
Clubs													
Diamonds													
Hearts													
Spades													

Hint: Start with defining event, $A_i = i^{\text{th}}$ card dealt to us is a spade

Special Assignment

- Example : Probabilistic Modeling - Engineering Application

- Presentation: 30 Minutes (PPT or Document Projector)
- Date: January 28, 2020 (Tuesday)
- It's an open call: Any group / student can take an initiative to present!
- Reference: Chapter-2 : Point 2.9

Introduction to Probability Theory

- Bayes's Theorem

- **Motivation:** It is derived from Joint Probability and helps to compute the conditional probability

Introduction to Probability Theory

- Bayes's Theorem

- **Motivation:** It is derived from Joint Probability and helps to compute the conditional probability
- **Theorem 2.9:** For any events A and B such that $\Pr(B) \neq 0$,

$$\Pr(A|B) = \frac{\Pr(B|A)\Pr(A)}{\Pr(B)}$$

Introduction to Probability Theory

- Bayes's Theorem

- **Motivation:** It is derived from Joint Probability and helps to compute the conditional probability

- **Theorem 2.9:** For any events A and B such that $\Pr(B) \neq 0$,

$$\Pr(A|B) = \frac{\Pr(B|A)\Pr(A)}{\Pr(B)}$$

- **Theorem 2.10: Total Probability Theorem**

Let B_1, B_2, \dots, B_n be a set of mutually exclusive and exhaustive events. That is, $B_i \cap B_j = \emptyset$ for all $i \neq j$ and

$$\bigcup_{i=1}^n B_i = S \Rightarrow \sum_{i=1}^n \Pr(B_i) = 1$$

Then,

$$\Pr(A) = \sum_{i=1}^n \Pr(A|B_i)\Pr(B_i)$$

Introduction to Probability Theory

- Bayes's Theorem

- **Motivation:** It is derived from Joint Probability and helps to compute the conditional probability

- **Theorem 2.9:** For any events A and B such that $\Pr(B) \neq 0$,

$$\Pr(A|B) = \frac{\Pr(B|A)\Pr(A)}{\Pr(B)}$$

- **Theorem 2.10: Total Probability Theorem**

Let B_1, B_2, \dots, B_n be a set of mutually exclusive and exhaustive events. That is, $B_i \cap B_j = \emptyset$ for all $i \neq j$ and

$$\bigcup_{i=1}^n B_i = S \Rightarrow \sum_{i=1}^n \Pr(B_i) = 1$$

Then,

$$\Pr(A) = \sum_{i=1}^n \Pr(A|B_i)\Pr(B_i)$$

Introduction to Probability Theory

- Bayes's Theorem

- **Theorem 2.11 (Bayes Theorem):** Let B_1, B_2, \dots, B_n be a set of mutually exclusive and exhaustive events. Then,

$$Pr(B_i|A) = \frac{Pr(A|B_i)Pr(B_i)}{\sum_{i=1}^n Pr(A|B_i)Pr(B_i)}$$

Where,

$Pr(B_i)$: a priori probability (probabilities that are formed from self-evident or presupposed models) and

$Pr(B_i|A)$: a posteriori probability (probabilities that are derived or calculated after observing certain events) of event B_i given A .

Introduction to Probability Theory

- Bayes's Theorem

- **Theorem 2.11 (Bayes Theorem):** Let B_1, B_2, \dots, B_n be a set of mutually exclusive and exhaustive events. Then,

$$Pr(B_i|A) = \frac{Pr(A|B_i)Pr(B_i)}{\sum_{i=1}^n Pr(A|B_i)Pr(B_i)}$$

Where,

$Pr(B_i)$: a **priori probability** (probabilities that are formed from self-evident or presupposed models) and

$Pr(B_i|A)$: a **posteriori probability** (probabilities that are derived or calculated after observing certain events) of event B_i given A .

- **Examples:**

Introduction to Probability Theory

- Example #1: Auditorium with 30 Rows

A certain auditorium has 30 rows of seats. **Row 1** has 11 seats, while **Row 2** has 12 seats, **Row 3** has 13 seats, and so on to the back of the auditorium where **Row 30** has 40 seats.

Introduction to Probability Theory

- Example #1: Auditorium with 30 Rows

A certain auditorium has 30 rows of seats. **Row 1** has 11 seats, while **Row 2** has 12 seats, **Row 3** has 13 seats, and so on to the back of the auditorium where **Row 30** has 40 seats.

A door prize (a prize which each person present at an event has a chance to win, usually by means of a draw or raffle) is to be given away by randomly selecting a row (with equal probability of selecting any of the 30 rows) and then randomly selecting a seat within that row (with each seat in the row equally likely to be selected).

Introduction to Probability Theory

- Example #1: Auditorium with 30 Rows

A certain auditorium has 30 rows of seats. **Row 1** has 11 seats, while **Row 2** has 12 seats, **Row 3** has 13 seats, and so on to the back of the auditorium where **Row 30** has 40 seats.

A door prize (a prize which each person present at an event has a chance to win, usually by means of a draw or raffle) is to be given away by randomly selecting a row (with equal probability of selecting any of the 30 rows) and then randomly selecting a seat within that row (with each seat in the row equally likely to be selected).

Compute the Probability that Seat 15 was selected given that Row 20 was selected and also find the probability that Row 20 was selected given that Seat 15 was selected.

Introduction to Probability Theory

- Example #2: Communication System - Receiver (RX)

A communication system sends binary data 0 or 1 which is then detected at the RX. The RX occasionally makes mistakes and sometimes a 0 is sent and is detected as a 1 or a 1 can be sent and detected as a 0.

Introduction to Probability Theory

- Example #2: Communication System - Receiver (RX)

A communication system sends binary data 0 or 1 which is then detected at the RX. The RX occasionally makes mistakes and sometimes a 0 is sent and is detected as a 1 or a 1 can be sent and detected as a 0. Suppose the communication system is described by the following set of conditional probabilities:

Introduction to Probability Theory

- Example #2: Communication System - Receiver (RX)

A communication system sends binary data 0 or 1 which is then detected at the RX. The RX occasionally makes mistakes and sometimes a 0 is sent and is detected as a 1 or a 1 can be sent and detected as a 0. Suppose the communication system is described by the following set of conditional probabilities:

$$\Pr(0 \text{ received} \mid 0 \text{ transmitted}) = 0.95,$$

Introduction to Probability Theory

- Example #2: Communication System - Receiver (RX)

A communication system sends binary data 0 or 1 which is then detected at the RX. The RX occasionally makes mistakes and sometimes a 0 is sent and is detected as a 1 or a 1 can be sent and detected as a 0. Suppose the communication system is described by the following set of conditional probabilities:

$$\Pr(0 \text{ received} \mid 0 \text{ transmitted}) = 0.95,$$

$$\Pr(1 \text{ received} \mid 0 \text{ transmitted}) = 0.05$$

Introduction to Probability Theory

- Example #2: Communication System - Receiver (RX)

A communication system sends binary data 0 or 1 which is then detected at the RX. The RX occasionally makes mistakes and sometimes a 0 is sent and is detected as a 1 or a 1 can be sent and detected as a 0. Suppose the communication system is described by the following set of conditional probabilities:

$$\Pr(0 \text{ received} \mid 0 \text{ transmitted}) = 0.95,$$

$$\Pr(1 \text{ received} \mid 0 \text{ transmitted}) = 0.05$$

$$\Pr(0 \text{ received} \mid 1 \text{ transmitted}) = 0.10,$$

Introduction to Probability Theory

- Example #2: Communication System - Receiver (RX)

A communication system sends binary data 0 or 1 which is then detected at the RX. The RX occasionally makes mistakes and sometimes a 0 is sent and is detected as a 1 or a 1 can be sent and detected as a 0. Suppose the communication system is described by the following set of conditional probabilities:

$$\Pr(0 \text{ received} \mid 0 \text{ transmitted}) = 0.95,$$

$$\Pr(1 \text{ received} \mid 0 \text{ transmitted}) = 0.05$$

$$\Pr(0 \text{ received} \mid 1 \text{ transmitted}) = 0.10,$$

$$\Pr(1 \text{ received} \mid 1 \text{ transmitted}) = 0.90.$$

- 1 Find $\Pr(0 \text{ received})$ and $\Pr(1 \text{ received})$.

Introduction to Probability Theory

- Example #2: Communication System - Receiver (RX)

A communication system sends binary data 0 or 1 which is then detected at the RX. The RX occasionally makes mistakes and sometimes a 0 is sent and is detected as a 1 or a 1 can be sent and detected as a 0. Suppose the communication system is described by the following set of conditional probabilities:

$$\Pr(0 \text{ received} \mid 0 \text{ transmitted}) = 0.95,$$

$$\Pr(1 \text{ received} \mid 0 \text{ transmitted}) = 0.05$$

$$\Pr(0 \text{ received} \mid 1 \text{ transmitted}) = 0.10,$$

$$\Pr(1 \text{ received} \mid 1 \text{ transmitted}) = 0.90.$$

- 1 Find $\Pr(0 \text{ received})$ and $\Pr(1 \text{ received})$.
- 2 Suppose a 0 is detected at the RX. What is the probability that the transmitted bit was actually a 1? Also, if a 1 was detected at the RX, what is the probability that the transmitted bit was actually a 0?

Introduction to Probability Theory

- Example #2: Communication System - Receiver (RX)

A communication system sends binary data 0 or 1 which is then detected at the RX. The RX occasionally makes mistakes and sometimes a 0 is sent and is detected as a 1 or a 1 can be sent and detected as a 0. Suppose the communication system is described by the following set of conditional probabilities:

$$\Pr(0 \text{ received} \mid 0 \text{ transmitted}) = 0.95,$$

$$\Pr(1 \text{ received} \mid 0 \text{ transmitted}) = 0.05$$

$$\Pr(0 \text{ received} \mid 1 \text{ transmitted}) = 0.10,$$

$$\Pr(1 \text{ received} \mid 1 \text{ transmitted}) = 0.90.$$

- 1 Find $\Pr(0 \text{ received})$ and $\Pr(1 \text{ received})$.
- 2 Suppose a 0 is detected at the RX. What is the probability that the transmitted bit was actually a 1? Also, if a 1 was detected at the RX, what is the probability that the transmitted bit was actually a 0?
- 3 What is the probability that the detected bit is not equal to the transmitted bit? This is the overall probability of error of the RX.

Introduction to Probability Theory

- Independence : Example (Re-define) # 1: Auditorium and Definitions

- If the description of the auditorium were changed so that each row had an **equal number of seats** (e.g., say all 30 rows had 20 seats each), then ??

Introduction to Probability Theory

- Independence : Example (Re-define) # 1: Auditorium and Definitions

- If the description of the auditorium were changed so that each row had an **equal number of seats** (e.g., say all 30 rows had 20 seats each), then ??
- Let event $A = \text{Row 20 was selected}$ and $B = \text{Seat 15 was selected}$
- Question: Can event B give us any new information about the likelihood of the event A ?

Introduction to Probability Theory

- Independence : Example (Re-define) # 1: Auditorium and Definitions

- If the description of the auditorium were changed so that each row had an **equal number of seats** (e.g., say all 30 rows had 20 seats each), then ??
- Let event **A= Row 20 was selected** and **B=Seat 15 was selected**
- Question: Can event B give us any new information about the likelihood of the event A ? **No**

Introduction to Probability Theory

- Independence : Example (Re-define) # 1: Auditorium and Definitions

- If the description of the auditorium were changed so that each row had an **equal number of seats** (e.g., say all 30 rows had 20 seats each), then ??
- Let event **A= Row 20 was selected** and **B=Seat 15 was selected**
- Question: Can event B give us any new information about the likelihood of the event A ? **No**
- **Mathematical:** Two events A and B are independent if $Pr(A|B) = Pr(B)$ and $Pr(B|A) = Pr(A)$. So, the joint probability $Pr(A, B) = Pr(A)Pr(B)$.

Introduction to Probability Theory


- Independence : Example (Re-define) # 1: Auditorium and Definitions

- If the description of the auditorium were changed so that each row had an **equal number of seats** (e.g., say all 30 rows had 20 seats each), then ??
- Let event **A= Row 20 was selected** and **B=Seat 15 was selected**
- Question: Can event B give us any new information about the likelihood of the event A ? **No**
- **Mathematical:** Two events A and B are independent if $Pr(A|B) = Pr(A)$ and $Pr(B|A) = Pr(B)$. So, the joint probability $Pr(A, B) = Pr(A)Pr(B)$.
- **Definition:** For three Events: The events A , B and C are mutually independent if each pair of events is independent; that is $Pr(A, B) = Pr(A)Pr(B)$, $Pr(A, C) = Pr(A)Pr(C)$, $Pr(B, C) = Pr(B)Pr(C)$, $Pr(A, B, C) = Pr(A)Pr(B)Pr(C)$
- **Examples:**

Introduction to Probability Theory

- Independence : Example #2: Communication Network


Consider a communications network with nodes A, B, C, and D and links a_1 , a_2 , a_3 and a_4 as shown in the diagram. The probability of a link being available at any time is p . In order to send a message from node A to node D we must have a path of available links from A to D. Assuming independence of link availability, **What is the probability of being able to send a message?**



Introduction to Probability Theory

- Independence : Example # 3 : Communication Network (Modified)

Suppose we modify the communications network as shown in the diagram by adding a link from node B to node D . Assuming each link is available with probability p independent of any other link, **what is the probability of being able to send a message from node A to D ?**



Introduction to Probability Theory

- Independence : Example # 4 : True or False

Suppose two events A and B are independent.

Introduction to Probability Theory

- Independence : Example # 4 : True or False

Suppose two events A and B are independent.

(a) Is it true that A is independent of \bar{B} ? If yes, give a convincing proof, otherwise, give a counterexample.

Introduction to Probability Theory

- Independence : Example # 4 : True or False

Suppose two events A and B are independent.

- (a) Is it true that A is independent of \bar{B} ? If yes, give a convincing proof, otherwise, give a counterexample.
- (b) Is it true that \bar{A} is independent of \bar{B} ? If yes, give a convincing proof, otherwise, give a counterexample.

Introduction to Probability Theory

- Independence : Example # 4 : True or False

Suppose two events A and B are independent.

- (a) Is it true that A is independent of \bar{B} ? If yes, give a convincing proof, otherwise, give a counterexample.
- (b) Is it true that \bar{A} is independent of \bar{B} ? If yes, give a convincing proof, otherwise, give a counterexample.

Thank you