
13 changes, one network each: almost all inside the seed noise
Drop path 0.2 (R1)
Full-frame 448 × 600 (R1)
Layer-wise LR decay (R1)
Lesion-centred crop (R1)
Metadata fusion (R1)
EMA (square input) (R1)
Full frame + drop path 0.1 (R2)
Full frame + drop path 0.2 (R2)
Full frame + drop path 0.3 (R2)
Full frame + EMA (R2)
Soup of 10 full-frame runs (R3)
Soup of 6 square runs (R3)
Consistent distillation (R4)
Consistent distillation, α 0.9 (R4)
test +0.005 CV +0.007
test -0.000 CV +0.008
test +0.000 CV -0.002
test -0.002 CV -0.001
test -0.001 CV -0.001
test +0.001 CV -0.004
test +0.006 CV +0.001
test +0.002 CV +0.006
test +0.006 CV +0.002
test +0.003 CV +0.006
test +0.007 CV -0.008
test +0.005 CV -0.004
test +0.006 CV +0.019
test -0.004 CV +0.026
grey band ± 0.011: typical change from the seed alone
−0.03
−0.02
−0.01
0.00
0.01
0.02
0.03
0.04
change in balanced accuracy vs the HW X baseline (paired over the same 5 folds)
adopted, beats its control on CV and test
adopted by the CV rule
adopted by the rule, worse on test
not adopted
CV (hollow)