

# Engineering Sciences (SPI) Doctoral School

Analytic stochastic processes for signal processing

## INTRODUCTION TO THE CONCEPT OF GRAVITATIONAL ALLOCATIONS

**Diala HAWAT**

**Director: Rémi Bardenet**


**Co-Director: Raphael Lachize-Rey**

1 Point process

2 Gravitational allocation


- Homogeneous Poisson point process  $\text{PPP}(\lambda)$

- Homogeneous Poisson point process  $PPP(\lambda)$




- Homogeneous Poisson point process  $\text{PPP}(\lambda)$
- Determinantal point process

- Homogeneous Poisson point process  $\text{PPP}(\lambda)$
- Determinantal point process



- Homogeneous Poisson point process  $PPP(\lambda)$
- Determinantal point process
- Permanental point process

- Homogeneous Poisson point process  $PPP(\lambda)$
- Determinantal point process
- Permanental point process





# Transportation, Matching, Allocation:

Given two measures  $\nu$  and  $\mu$  on  $\Lambda$


- A transportation between  $\nu$  and  $\mu$  is a measure  $\rho$  on  $\Lambda \times \Lambda$  whose first marginal is  $\nu$  and the second marginal is  $\mu$ .

Given two measures  $\nu$  and  $\mu$  on  $\Lambda$

- A transportation between  $\nu$  and  $\mu$  is a measure  $\rho$  on  $\Lambda \times \Lambda$  whose first marginal is  $\nu$  and the second marginal is  $\mu$ .
- When  $\nu$  and  $\mu$  are both counting measures, the transportation will be also called a matching.

Given two measures  $\nu$  and  $\mu$  on  $\Lambda$

- A transportation between  $\nu$  and  $\mu$  is a measure  $\rho$  on  $\Lambda \times \Lambda$  whose first marginal is  $\nu$  and the second marginal is  $\mu$ .
- When  $\nu$  and  $\mu$  are both counting measures, the transportation will be also called a matching.




Given two measures  $\nu$  and  $\mu$  on  $\Lambda$

- A transportation between  $\nu$  and  $\mu$  is a measure  $\rho$  on  $\Lambda \times \Lambda$  whose first marginal is  $\nu$  and the second marginal is  $\mu$ .
- When  $\nu$  and  $\mu$  are both counting measures, the transportation will be also called a matching.
- When  $\nu$  is the Lebesgue measure and  $\mu$  is a counting measure, the transportation will be called an allocation.

Given two measures  $\nu$  and  $\mu$  on  $\Lambda$

- A transportation between  $\nu$  and  $\mu$  is a measure  $\rho$  on  $\Lambda \times \Lambda$  whose first marginal is  $\nu$  and the second marginal is  $\mu$ .
- When  $\nu$  and  $\mu$  are both counting measures, the transportation will be also called a matching.
- When  $\nu$  is the Lebesgue measure and  $\mu$  is a counting measure, the transportation will be called an allocation.





# Gravitational allocation from Lebesgue to the Poisson point process:



- Let  $\mathcal{Z}$  be a standard Poisson point process on  $\mathbb{R}^d$
- Consider the vector function

$$F(x) := \sum_{z \in \mathcal{Z}, |z-x| \uparrow} \frac{z-x}{|z-x|^d},$$

to be the force of attraction acting on each point  $x$  of  $\mathbb{R}^d \setminus \mathcal{Z}$ .

- For any  $x \in \mathbb{R}^d \setminus \mathcal{Z}$ , consider the integral curve  $Y_x(t)$ , to be unique solution of

$$\frac{dY_x(t)}{dt} = F(Y_x(t)), \quad Y_x(0) = x.$$

define for some maximal time  $\tau_x \in ]0, \infty]$ .

We call these curves the gravitational flow curves.



## Proposition:

Assume  $d \geq 3$ . Almost surely, the above series defining the force function

$$F(x) := \sum_{z \in \mathbb{Z}, |z-x| \uparrow} \frac{z-x}{|z-x|^d},$$

converges simultaneously for all  $x$  for which it is defined, and defines a translation-invariant (in distribution) vector valued random function. The random function  $F$  is almost surely continuously differentiable where it is defined.

# Alternative formulation of $F$ :

$$F(x) := \sum_{z \in \mathcal{Z}, |z-x| \uparrow} \frac{z-x}{|z-x|^d},$$

$$\sum_{z \in \mathcal{Z}, |z-u| \uparrow} \frac{z-x}{|z-x|^d} - \sum_{z \in \mathcal{Z}, |z-v| \uparrow} \frac{z-x}{|z-x|^d} = \kappa_d(u-v). \quad (1)$$

This equation lead to a new formulation of the force function  $F$  as follow:

$$F(x) := \sum_{z \in \mathbb{Z}, |z-x| \uparrow} \frac{z-x}{|z-x|^d},$$

$$F(x) := \sum_{z \in \mathbb{Z}, |z| \uparrow} \frac{z-x}{|z-x|^d} + \kappa_d x.$$

The force function was taken to be:

$$F(x) = \sum_{z \in \mathcal{Z}, |z| \uparrow} \frac{z - x}{|z - x|^d} + \kappa_d x,$$

while its the divergence is:

$$\operatorname{div}(F) = -d\kappa_d \sum_{z \in \mathcal{Z}} \delta_z + d\kappa_d,$$

- A force function  $F$  s.t.

$$F(x) = -\nabla u(x), \quad \Delta u = -\operatorname{div}(F)$$



- A force function  $F$  s.t.

$$F(x) = -\nabla u(x), \quad \Delta u = -\operatorname{div}(F)$$

- A potential function  $u$  super-harmonic on  $\mathbb{R}^d \setminus \mathcal{Z}$ , i.e.


$$\Delta u(x) \leq 0.$$

- A force function  $F$  s.t.

$$F(x) = -\nabla u(x), \quad \Delta u = -\operatorname{div}(F)$$

- A potential function  $u$  super-harmonic on  $\mathbb{R}^d \setminus \mathcal{Z}$ , i.e.

$$\Delta u(x) \leq 0.$$



- A force function  $F$  s.t.

$$F(x) = -\nabla u(x), \quad \Delta u = -\operatorname{div}(F)$$

- A potential function  $u$  super-harmonic on  $\mathbb{R}^d \setminus \mathcal{Z}$ , i.e.

$$\Delta u(x) \leq 0.$$

- The gradient curves  $Y_x(t)$  are the unique solution of the differential equation

$$\frac{dY_x(t)}{dt} = F(Y_x(t)) = -\nabla u(Y_x(t)), \quad Y_x(0) = x.$$



THANK YOU.