

Maths for ML Handbook

Equations and Formulas used in Machine Learning

Version 3.0

Index

PART A: Linear Algebra

1. Linear Equation
2. General Line Equation
3. Vector
4. Matrix Multiplication
5. Dot product
6. Applications of Dot Product
7. Shifting Lines
8. Distances

PART B: Differential Calculus

9. Functions
10. Differentiation
11. Partial Differentiation

PART C: Applications

12. Linear Binary Classifier
13. Optimization
14. Gradient Descent
15. Principal Component Analysis (PCA)

PART D: Probability and Statistics

16. Basics of Probability
17. Types of Events
18. Combinatorics
19. Bayes Theorem
20. Measure of Central Tendency
21. Measure of Variance
22. Distribution Functions
23. Binomial Distribution
24. Poisson Distribution
25. Geometric Distribution
26. Normal Distribution
27. Uniform Distribution
28. Exponential Distribution

PART A: Linear Algebra

1 Linear Equation

1.1 Slope-Intercept Equation

Slope-Intercept Form

$$y = m \cdot x + c$$

Where,

- m - Slope (angle between straight line and x-axis)
- c - intercept (distance of straight line from the origin)
- x - Independent variable
- y - Dependent variable

Slope

Given two points (x_1, y_1) and (x_2, y_2) , formula to calculate Slope m :

$$m = \frac{y_2 - y_1}{x_2 - x_1}$$

Intercept

Given two points (x_1, y_1) and (x_2, y_2) , formula to calculate Intercept c :

$$c = y_1 - (m \cdot x_1)$$

or

$$c = y_2 - (m \cdot x_2)$$

1.2 Limitations of Slope-Intercept Equation

Limitations

1. Slope m becomes undefined or ∞ when $x_2 - x_1 = 0$.
2. Not scalable for higher dimensions.

Solution

General Line Equation:

$$ax + by + c = 0$$

1.3 Geometric Relationship b/w lines

Parallel Lines

When given two lines $y = m_1x + c_1$ and $y = m_2x + c_2$ are parallel:

$$m_1 = m_2$$

Perpendicular Lines (Orthogonality)

When given two lines $y = m_1x + c_1$ and $y = m_2x + c_2$ are perpendicular:

$$m_1 \times m_2 = -1$$

2 General Line Equation

2.1 Standard Form

$$ax + by + c = 0$$

Rearranging above General Line Equation in Slope-Intercept form:

$$w_1x + w_2y + w_0 = 0$$

$$w_2y = -w_1x - w_0$$

$$y = \left(-\frac{w_1}{w_2}\right)x + \left(-\frac{w_0}{w_2}\right)$$

Where Slope and Intercept can be derived as,

$$m = -\frac{w_1}{w_2} \quad c = -\frac{w_0}{w_2}$$

2.2 Advantages

1. Scaling into higher dimensions is achieved by Linear Expansion of the General Line equation.
2. Undefined slope is solved by setting the coefficient of y i.e., b in GLE: $ax + by + c = 0$ to zero.

2.3 Scaling

Line Equation (2D Hyperplane)

General Line Equation for 2 dimensional data:

$$w_1x_1 + w_2x_2 + w_0 = 0$$

Plane Equation (3D Hyperplane)

Scaling General Line Equation for 3 dimensional data:

$$w_1x_1 + w_2x_2 + w_3x_3 + w_0 = 0$$

nD Hyperplane Equation

Scaling General Line Equation for higher dimensional ($d > 3$) data:

$$w_1x_1 + w_2x_2 + \dots + w_nx_n + w_0 = 0$$

3 Vector

3.1 Vector Representation

Mathematical notation of a vector $\vec{x}$ having d dimension containing real numbers:

$$x \in \mathbb{R}^d$$

Where d represents number of elements in the vector.

Row Vector

$$\vec{x} = [x_1 \ x_2 \ \dots \ x_d]$$

Column Vector

$$\vec{x} = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_d \end{bmatrix}$$

3.2 Types of Distances

Various methods to calculate distance between two points (x_1, y_1) and (x_2, y_2) .

1 Euclidean Distance

Formula to calculate Euclidean Distance between two points (x_1, y_1) and (x_2, y_2)

$$d = \sqrt{(y_2 - y_1)^2 + (x_2 - x_1)^2}$$

2 Manhattan Distance

Formula to calculate Manhattan Distance between two points (x_1, y_1) and (x_2, y_2)

$$d = |y_2 - y_1| + |x_2 - x_1|$$

3.3 Types of Norms

Norm

Norm of a vector represents the straight-line distance from the origin to the endpoint of the vector.

L2 Norm of a Vector

Formula for L2 Norm or Euclidean Distance of a vector of d dimension:

$$\begin{aligned}\|\vec{x}\|_2 &= \sqrt{x_1^2 + x_2^2 + x_3^2 + \dots + x_d^2} \\ &= \sqrt{\sum_{i=1}^d x_i^2}\end{aligned}$$

L1 Norm of a Vector

Formula for L1 Norm or Manhattan Distance of a vector of d dimension:

$$\begin{aligned}\|\vec{x}\|_1 &= |x_1| + |x_2| + |x_3| + \dots + |x_d| \\ &= \sum_{i=1}^d |x_i|\end{aligned}$$

4 Matrix Multiplication

4.1 Matrix Multiplication rule

Given two matrices A and B where:

- $A : m \times n$
- $B : n \times k$

Rule

Number of columns in the first matrix should be equal to the number of rows in the second matrix.

$$m \times n \Leftrightarrow n \times k = m \times k$$

The resulting matrix will be of shape: $m \times k$

4.2 Matrix Multiplication in NumPy

Given two NumPy matrices `m1` and `m2`

```
m1 @ m2  
# or  
np.matmul(m1, m2)
```

5 Dot Product

5.1 Dot Product of Vectors

Formula to calculate dot product of two vectors $\vec{a}$ and $\vec{b}$:

$$\vec{a} \cdot \vec{b} = a^T \cdot b$$

$$= \sum_{i=1}^d a_i b_i \in \mathbb{R}^1$$

Note: $\mathbb{R}^1$ represents scalar value i.e., Dot product of two vectors results in real number.

5.2 Dot Product in NumPy

Given two NumPy vectors `v1` and `v2`

```
np.dot(v1, v2)
```

6 Applications of Dot Product

6.1 Angle between Vectors

Formula to calculate angle between two vectors $\vec{a}$ and $\vec{b}$

$$\cos \theta = \frac{\vec{a}^T \cdot \vec{b}}{\|\vec{a}\| \|\vec{b}\|}$$

Cauchy-Schwarz Inequality

Absolute value of dot product of two vectors is always less than or equal to the product of their magnitudes.

$$|\vec{a} \cdot \vec{b}| \leq \|\vec{a}\| \|\vec{b}\|$$

Perpendicular Vectors

When two vectors are **perpendicular** to each other, their **dot product is zero**.

If two vectors are perpendicular to each other, then $\theta = 90^\circ$ and $\cos 90^\circ = 0$, therefore,

$$\vec{a} \cdot \vec{b} = 0$$

6.2 Machine Learning

In ML context, the dot product is used to represent the below "best fit" line or hyper-plane equation:

$$w_1x_1 + w_2x_2 + \dots + w_dx_d + w_0 = 0$$

using $\vec{w}$ and $\vec{x}$ vectors of d dimensions:

$$\vec{w} = \begin{bmatrix} w_1 \\ w_2 \\ \vdots \\ w_d \end{bmatrix} \quad \vec{x} = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_d \end{bmatrix}$$

in **Linear Algebraic** form as,

$$\vec{w}^T \cdot \vec{x} + w_0 = 0$$

Where $\vec{w}^T \cdot \vec{x}$ is the dot product between weight vector $\vec{w}$ and feature vector $\vec{x}$.

6.3 Projection of a Vector

Unit Vector

Unit vector is defined as a vector whose magnitude is equal to 1.

$$\|\vec{a}\| = 1$$

Formula to calculate unit vector $\hat{w}$ of a $\vec{w}$ vector:

$$\hat{w} = \frac{\vec{w}}{\|\vec{w}\|}$$

Scalar Projection

$$\text{scalar projection of } \vec{x} \text{ on } \vec{y} = \vec{x} \cdot \hat{y}$$

Formula to calculate scalar projection of $\vec{x}$ on $\vec{y}$:

$$\text{comp}_{\vec{y}} \vec{x} = \frac{\vec{x} \cdot \vec{y}}{\|\vec{y}\|}$$

Dot product is equal to the product of one vector and the projection of the vector on the first one.

$$\vec{a} \cdot \vec{b} = \|\vec{a}\| \times (\text{scalar projection of } \vec{b} \text{ on } \vec{a})$$

or

$$\vec{a} \cdot \vec{b} = \|\vec{b}\| \times (\text{scalar projection of } \vec{a} \text{ on } \vec{b})$$

Vector Projection

The Vector Projection of $\vec{x}$ onto $\vec{y}$ equals the Scalar Projection of $\vec{x}$ onto $\vec{y}$ times a unit vector in the direction of $\vec{y}$.

Formula to calculate vector projection of $\vec{x}$ on $\vec{y}$:

$$\begin{aligned} \text{proj}_{\vec{y}} \vec{x} &= \left(\frac{\vec{x} \cdot \vec{y}}{\|\vec{y}\|} \right) \hat{y} \\ &= \left(\frac{\vec{x} \cdot \vec{y}}{\|\vec{y}\|} \right) \frac{\vec{y}}{\|\vec{y}\|} \\ &= \frac{\vec{x} \cdot \vec{y}}{\|\vec{y}\|^2} \vec{y} \end{aligned}$$

7 Shifting Lines

7.1 One Direction

Given line equation:

$$w_1x_1 + w_2x_2 + w_0 = 0$$

Shifting lines in one direction (x-axis or y-axis) by a units:

$$\text{x-axis} \mid \text{Right } \rightarrow \quad w_1(x_1 - a) + w_2x_2 + w_0 = 0$$

$$\text{x-axis} \mid \text{Left } \leftarrow \quad w_1(x_1 + a) + w_2x_2 + w_0 = 0$$

$$\text{y-axis} \mid \text{Up } \uparrow \quad w_1x_1 + w_2(x_2 - a) + w_0 = 0$$

$$\text{y-axis} \mid \text{Down } \downarrow \quad w_1x_1 + w_2(x_2 + a) + w_0 = 0$$

7.2 Two Directions

Shifting lines in two direction (x-axis and y-axis) simultaneously by a units:

$$w_1(x_1 - a) + w_2(x_2 - a) + w_0 = 0$$


Solving above equation gives:

$$w_1x_1 + w_2x_2 + w'_0 = 0$$

Where w'_0 is:

$$w'_0 = w_0 - a(w_1 + w_2)$$

7.3 Usecase



8 Distances

8.1 Distance Between Line and Origin

Formula to calculate the distance between line (Decision Boundary) and origin using line's Normal vector $\vec{w}$

$$d = \frac{|-w_0|}{\|w\|}$$

8.2 Distance Between Point and Line

Formula to calculate the distance between a point x_0 and line (Decision Boundary) using line's Normal vector $\vec{w}$

$$d = \frac{|w^T x_0 + w_0|}{\|w\|}$$

8.3 Distance Between Parallel Lines

Same Normal Vector

Formula to calculate the distance between two parallel lines with Normal vector w having same magnitude:

$$d = \frac{|w_0^{(1)} - w_0^{(2)}|}{\|w\|}$$

Different Normal Vector

Formula to calculate the distance between two parallel lines with Normal vectors of different magnitudes where $w^{(1)} = \lambda \cdot w^{(2)}$

$$d = \left| \frac{w_0^{(1)}}{\|w^{(1)}\|} - \frac{w_0^{(2)}}{\|w^{(2)}\|} \right|$$

PART B: Differential Calculus

9 Functions


9.1 Continuous Function

Definition


Criteria for a Continuous function:

$$\lim_{x \rightarrow a^-} f(x) = \lim_{x \rightarrow a^+} f(x) = f(a)$$

Example #1



Example #2



Note: Even though $f(x) = |x|$ is continuous function, it is **not differentiable at** $x = 0$


9.2 Discontinuous Function

Definition


Criteria for a Discontinuous function:

$$\lim_{x \rightarrow a^-} f(x) \neq \lim_{x \rightarrow a^+} f(x)$$

Example #1




Example #2



10 Differentiation

10.1 Slopes


Slope of a straight Line



Formula to calculate slope of a line intersecting two points (x_1, y_1) and (x_2, y_2)

$$m = \tan \theta = \frac{y_2 - y_1}{x_2 - x_1}$$


Slope of Secant Line to a curve



Formula for slope of a secant line intersecting a curve at two points (x_1, y_1) and (x_2, y_2)

$$m = \frac{f(x_2) - f(x_1)}{x_2 - x_1}$$

Slope of Tangent Line to a curve



Formula for slope of a tangent line (a.k.a Derivative) intersecting a curve at one single point (x, y)

$$f'(x) = \frac{d}{dx} f(x) = \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x}$$

10.2 Derivatives Examples

Commonly used Derivatives

$$(1) \quad \frac{d}{dx} x^n = nx^{n-1}$$

$$(5) \quad \frac{d}{dx} \sin(x) = \cos(x)$$

$$(2) \quad \frac{d}{dx} \log(x) = \frac{1}{x}$$

$$(6) \quad \frac{d}{dx} \cos(x) = -\sin(x)$$

$$(3) \quad \frac{d}{dx} e^x = e^x$$

$$(7) \quad \frac{d}{dx} \tan(x) = \sec^2(x)$$

$$(4) \quad \frac{d}{dx} c = 0$$

$$(8) \quad \frac{d}{dx} b^x = b^x \log(b)$$

10.3 Rules of Differentiation

1 Constant Rule

$$\frac{d}{dx} [c \cdot f(x)] = c \cdot \frac{d}{dx} f(x)$$

2 Sum Rule

$$\frac{d}{dx} [f(x) + g(x)] = \frac{d}{dx} f(x) + \frac{d}{dx} g(x)$$

3 Difference Rule

$$\frac{d}{dx} [f(x) - g(x)] = \frac{d}{dx} f(x) - \frac{d}{dx} g(x)$$

4 Product Rule

$$\frac{d}{dx} [f(x) \cdot g(x)] = g(x) \frac{d}{dx} f(x) + f(x) \frac{d}{dx} g(x)$$

5 Quotient Rule

$$\frac{d}{dx} \left[\frac{f(x)}{g(x)} \right] = \frac{g(x) \frac{d}{dx} f(x) - f(x) \frac{d}{dx} g(x)}{[g(x)]^2}$$

6 Chain Rule


$$\frac{d}{dx} f(g(x)) = \frac{d}{dx} f(g(x)) \cdot \frac{d}{dx} g(x)$$

How Chain rule works?

$$\begin{aligned} \frac{d}{dx} \log(x^2) &= \frac{1}{x^2} \cdot 2x \\ &= \frac{2}{x} \end{aligned}$$

- Keep the inside x^2 as it is and take derivative of outside log i.e., $\frac{1}{x^2}$.
- Multiply it by the derivative of the inside x^2 i.e., $2x$.

10.4 Critical Points




Definition

Slope of the Tangent line to a curve (i.e., derivative of a function) will be **zero at Critical Points**.

$$f'(x) = 0$$

10.5 Local Optima


Types of Local Optima/Extrema



Two types of Local Optima/Extrema where $f'(x)$ (First derivative) is equal to zero and $f''(x)$ (Second Derivative) decides the type of Extrema:


$$f'(x) = 0, \quad f''(x) \begin{cases} f''(x) > 0 & \text{Local Minima} \\ f''(x) < 0 & \text{Local Maxima} \end{cases}$$

1 Local Minima




$$f''(x) > 0$$

2 Local Maxima



$$f''(x) < 0$$

Saddle Point



Non-optimal points where both $f'(x) = 0$ (First derivative) and $f''(x) = 0$ (Second derivative).

$$f''(x) = 0$$

10.6 Principles of Maxima and Minima

First and Second Derivative Test for Extrema

STEP 1: Compute First Derivative

Given function $f(x)$ compute its first derivative:

$$f'(x)$$

STEP 2: Find Critical Point

Equate the first derivative to zero to find the critical point:

$$f'(x) = 0$$

$$x = c$$

Critical Point :- Critical value of x , say c , at which slope: $f'(x) = 0$.

STEP 3: Classify Critical Point

To classify the Critical point from STEP 2 to the correct type of local Optima, compute derivative of $f'(x)$ i.e., second derivative of $f(x)$ from STEP 1:

$$f''(x)$$

Identify the type of Local Optima based on below rule:

$$f''(x) \begin{cases} \text{if } f''(x) > 0 : & \text{Local Minima} \\ \text{if } f''(x) < 0 : & \text{Local Maxima} \\ \text{if } f''(x) = 0 : & \text{Test Inconclusive} \end{cases}$$

STEP 4: Calculate Local Optima Value

Substitute the Critical Value c from STEP 2 in original function $f(x)$ to identify the actual value of Local Optima.

$$f(c) \in \mathbb{R}$$

11 Partial Differentiation

11.1 Partial Derivative

Partial Derivative of a general function

Formula to calculate Partial derivative of any general function $f(x, y)$ with two variables $\{x, y\}$

$$\frac{\partial}{\partial x} f(x, y) = \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x, y) - f(x, y)}{\Delta x}$$

$$\frac{\partial}{\partial y} f(x, y) = \lim_{\Delta y \rightarrow 0} \frac{f(x, y + \Delta y) - f(x, y)}{\Delta y}$$

11.2 Slope of Tangent Line (Gradient)

Gradient of a function

Given Linear Binary Classifier having feature vector $\vec{x}$ of d dimension and weight vector $\vec{w}$:

$$f(w_1, w_2, \dots, w_d, w_0) = w_1 x_1 + w_2 x_2 + \dots + w_d x_d + w_0$$

$$f(\vec{w}, w_0) = w_1 x_1 + w_2 x_2 + \dots + w_d x_d + w_0$$

Equation to calculate Gradient Vector for the above scalar function $f(\vec{w}, w_0)$

$$\nabla_{\vec{w}, w_0} f(\vec{w}, w_0) = \begin{bmatrix} \frac{\partial}{\partial w_1} f(\vec{w}, w_0) \\ \frac{\partial}{\partial w_2} f(\vec{w}, w_0) \\ \vdots \\ \frac{\partial}{\partial w_d} f(\vec{w}, w_0) \\ \frac{\partial}{\partial w_0} f(\vec{w}, w_0) \end{bmatrix}$$

Visualize Gradient: <https://www.desmos.com/3d/e3fratrwho>

PART C: Applications

Applications of Linear Algebra and Differential Calculus in Machine Learning

12 Linear Binary Classifier

12.1 Half-spaces

Positive Half-space

In Positive Half-space all data-points have **distance greater than zero** because,

$$w_1x_1 + w_2x_2 + \dots + w_dx_d + w_0 > 0$$

Negative Half-space

In Negative Half-space all data-points have **distance less than zero** because,

$$w_1x_1 + w_2x_2 + \dots + w_dx_d + w_0 < 0$$

Decision Boundary

On Decision Boundary all data-points have **distance equal to zero** because,

$$w_1x_1 + w_2x_2 + \dots + w_dx_d + w_0 = 0$$

12.2 Mathematical Notation

Dataset

Equation to represent a labeled dataset D used in Linear Binary Classifier:

$$D := \left\{ (x_i, y_i) ; x \in \mathbb{R}^d ; y \in \{-1, +1\} \right\}_{i=1}^n$$

Where,

- D is Dataset
- x are features of datatype Real numbers $\mathbb{R}$
- y are labels
- n is the total number of rows in the dataset
- d is the total number of features/columns (dimension) in the dataset

Label

Actual Label y can take values:

$$y \begin{cases} +1 & \text{in +ve Half-Space} \\ -1 & \text{in -ve Half-Space} \end{cases}$$

Classifier

Mapping function for a Linear Binary Classifier: nD Hyperplane

$$\begin{aligned} \hat{y} = f(x) &= w_1 x_1 + w_2 x_2 + \dots + w_d x_d + w_0 \\ &\text{or} \\ &= w^T x + w_0 \end{aligned}$$

Predicted label $\hat{y}$ can take values:

$$\hat{y} \begin{cases} +1 & \text{if } (w^T x + w_0) \geq 0 \\ -1 & \text{if } (w^T x + w_0) < 0 \end{cases}$$

12.3 Gain Function

Equation of Gain Function for Linear Binary Classifier:

$$g(D, \vec{w}, w_0) = \frac{1}{n} \sum_{i=1}^n \left(\frac{w^T x^{(i)} + w_0}{\|w\|} \right) \cdot y^{(i)}$$

12.4 Loss Function

Equation of Loss Function for Linear Binary Classifier:

Loss Function = - Gain Function

$$l(D, \vec{w}, w_0) = -\frac{1}{n} \sum_{i=1}^n \left(\frac{w^T x^{(i)} + w_0}{\|w\|} \right) \cdot y^{(i)}$$

12.5 Perceptron Learning Algorithm

Weight Updation Logic

Logic in Perceptron Learning Algorithm to update Weights for miss-classified data-point:


$$\text{if } \text{sign}\left(\frac{w^T x^{(i)} + w_0}{\|w\|}\right) \neq \text{sign}(y^{(i)}) :$$

$$w = w + x^{(i)} \cdot y^{(i)}$$

$$w_0 = w_0 + y^{(i)}$$

13 Optimization

13.1 ML Context



Error vs Classifier is represented as a function $f(x)$ which needs to be minimized.

13.2 Optimization of Gain Function

Equation for Optimization of Gain Function for Linear Binary Classifier:

$$\begin{aligned} w^*, w_0^* &= \arg \max_{w, w_0} g(D, \vec{w}, w_0) \\ &= \arg \max_{w, w_0} \frac{1}{n} \sum_{i=1}^n \left(\frac{w^T x^{(i)} + w_0}{\|w\|} \right) \cdot y^{(i)} \end{aligned}$$

Goal: Maximize Gain function.

13.3 Optimization of Loss Function

Equation for Optimization of Loss Function for Linear Binary Classifier:

$$\begin{aligned} w^*, w_0^* &= \arg \min_{w, w_0} l(D, \vec{w}, w_0) \\ &= \arg \min_{w, w_0} -\frac{1}{n} \sum_{i=1}^n \left(\frac{w^T x^{(i)} + w_0}{\|w\|} \right) \cdot y^{(i)} \end{aligned}$$

Goal: Minimize Loss function.

14 Gradient Descent

14.1 Gradient Descent Equation

Gradient Update Rule

$$x^n = x^o - \eta \frac{\partial}{\partial x} f(x^o)$$

Where,

- x - Some random vector (not features)
- x^n - New x values after update
- x^o - Old x values before update
- η - Learning Rate
- $f(x)$ - Objective function
- $\frac{\partial}{\partial x} f(x)$ - Partial Derivative of Objective function

Gradient Update Rule (In ML Context)

$$\begin{aligned} \vec{w}^{\text{new}} &= \vec{w}^{\text{old}} - \eta \frac{\partial}{\partial \vec{w}} f(\vec{w}^{\text{old}}) \\ w_0^{\text{new}} &= w_0^{\text{old}} - \eta \frac{\partial}{\partial w_0} f(w_0^{\text{old}}) \end{aligned}$$

Where,

- $\vec{w}^{\text{new}}$ - New Weight vector w after update
- $\vec{w}^{\text{old}}$ - Old Weight vector w before update
- w_0^{new} - New Bias w_0 value after update
- w_0^{old} - Old Bias w_0 value before update

14.2 Constrained Optimization

Constraint

Equation for Optimization/Minimization of Loss Function with a constraint that $\vec{w}$ is a unit vector:

$$w^*, w_0^* = \arg \min_{\vec{w}, w_0} l(D, \vec{w}, w_0) \quad \text{such that: } \|\vec{w}\| = 1$$

$$= \arg \min_{\vec{w}, w_0} -\frac{1}{n} \sum_{i=1}^n \left(\frac{w^T x^{(i)} + w_0}{\|w\|} \right) y^{(i)} \quad \text{such that: } \|\vec{w}\| = 1$$

Lagrangian Function

$$\mathcal{L}(x, \lambda) \equiv f(x) + \lambda \cdot g(x)$$

Where λ is called as **Lagrange Multiplier**.

Lagrange Multiplier for Constraint Optimization

Applying Lagrange Multiplier on Optimization equation:

$$\begin{aligned} w^*, w_0^* &= \arg \min_{\vec{w}, w_0, \lambda} l(D, \vec{w}, w_0, \lambda) \\ &= \arg \min_{\vec{w}, w_0, \lambda} -\frac{1}{n} \sum_{i=1}^n \left(w^T x^{(i)} + w_0 \right) y^{(i)} + \lambda (\|\vec{w}\| - 1) \end{aligned}$$

Gradient Update Rule With Lagrange Multiplier

$$\begin{aligned} \vec{w}^{t+1} &= \vec{w}^t - \eta \left[-\sum_{i=1}^n x_i \cdot y_i + \lambda \frac{\vec{w}}{\|w\|} \right] \\ w_0^{t+1} &= w_0^t - \eta \left[-\sum_{i=1}^n y_i \right] \end{aligned}$$

Where,

- λ - is the regularizer.

15 Principal Component Analysis (PCA)

15.1 PCA Linear Equation

Linear equation for PCA:

$$\text{feature}_{\text{new}} = u_1 * \text{feature}_1 + u_2 * \text{feature}_2 + \dots + u_n * \text{feature}_n$$

Where $\text{feature}_{\text{new}}$ is a linear combination of features: $\{\text{feature}_1, \text{feature}_2, \dots, \text{feature}_n\}$

Principal Components

Principal Components vector $v_{n \times d}$ is calculated as:

$$\begin{bmatrix} x_{11} & x_{12} & \cdots & x_{1d} \\ \vdots & \vdots & \vdots & \vdots \\ x_{n1} & x_{n2} & \cdots & x_{nd} \end{bmatrix}_{n \times d} \cdot \begin{bmatrix} u_{11} & u_{12} & \cdots & u_{1d} \\ \vdots & \vdots & \vdots & \vdots \\ u_{d1} & u_{d2} & \cdots & u_{dd} \end{bmatrix}_{d \times d} = \begin{bmatrix} v_{11} & v_{12} & \cdots & v_{1d} \\ \vdots & \vdots & \vdots & \vdots \\ v_{n1} & v_{n2} & \cdots & v_{nd} \end{bmatrix}_{n \times d}$$
$$X_{n \times d}^T \cdot u_{d \times d} = v_{n \times d}$$

15.2 Optimization of Variance in Projection

Equation for Optimization/maximizing variance during projection of vectors:

$$\vec{u}^* = \arg \max_{\vec{u}} \frac{1}{n} \sum_{i=1}^n \left(\frac{\vec{x}_i^T \cdot \vec{u}_i}{\|\vec{u}\|} - \mu \right)^2$$

15.3 Constraint Optimization

Constraint

Equation for maximizing variance during projection of vectors with constraint enforced on it:

$$\vec{u}^* = \arg \max_{\vec{u}} \frac{1}{n} \sum_{i=1}^n \left(\frac{\vec{x}_i^T \cdot \vec{u}_i}{\|\vec{u}\|} - \mu \right)^2 \quad \text{such that: } \|\vec{u}\| = 1$$

Equation for maximizing variance during projection of vectors with constraint optimization:

$$\vec{u}^* = \arg \max_{\vec{u}} \frac{1}{n} \sum_{i=1}^n \left(\vec{x}_i^T \cdot \vec{u}_i \right)^2 + \lambda (\|\vec{u}\| - 1)$$

PART D: Probability and Statistics

W.I.P