

Maths for ML Handbook

Equations and Formulas used in Machine Learning

1 Linear Equation

1.1 Slope-Intercept Equation

Slope-Intercept Form

$$y = m \cdot x + c$$

Where,

- m - Slope (angle between straight line and x-axis)
- c - intercept (distance of straight line from the origin)
- x - Independent variable
- y - Dependent variable

Slope

Given two points (x_1, y_1) and (x_2, y_2) , formula to calculate Slope m :

$$m = \frac{y_2 - y_1}{x_2 - x_1}$$

Intercept

Given two points (x_1, y_1) and (x_2, y_2) , formula to calculate Intercept c :

$$c = y_1 - (m \cdot x_1) \quad \text{or} \quad c = y_2 - (m \cdot x_2)$$

1.2 Geometric Relationship b/w lines

Parallel Lines

When given two lines $y = m_1x + c_1$ and $y = m_2x + c_2$ are parallel:

$$m_1 = m_2$$

Perpendicular Lines (Orthogonality)

When given two lines $y = m_1x + c_1$ and $y = m_2x + c_2$ are perpendicular:

$$m_1 \times m_2 = -1$$

1.3 General Line Equation

Standard Form

General Line Equation (GLE):

$$w_1x + w_2y + w_0 = 0$$

Rearranging above General Line Equation in Slope-Intercept form:

$$y = \left(-\frac{w_1}{w_2}\right)x_1 + \left(-\frac{w_0}{w_2}\right)$$

Where Slope and Intercept can be derived as,

$$m = -\frac{w_1}{w_2} \quad c = -\frac{w_0}{w_2}$$

Line Equation (2D Hyperplane)

$$w_1x_1 + w_2x_2 + w_0 = 0$$

Plane Equation (3D Hyperplane)

$$w_1x_1 + w_2x_2 + w_3x_3 + w_0 = 0$$

nD Hyperplane Equation

$$w_1x_1 + w_2x_2 + \dots + w_nx_n + w_0 = 0$$

2 Vector

2.1 Vector Representation

Mathematical notation of a vector $\vec{x}$ having d dimension containing real numbers:

$$\mathbf{x} \in \mathbb{R}^d$$

Where d represents number of elements in the vector.

Row Vector

$$\vec{x} = [x_1 \ x_2 \ \dots \ x_d]$$

Column Vector

$$\vec{x} = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_d \end{bmatrix}$$

2.2 Types of Distances

1 Euclidean Distance

Formula to calculate Euclidean Distance between two points (x_1, y_1) and (x_2, y_2)

$$d = \sqrt{(y_2 - y_1)^2 + (x_2 - x_1)^2}$$

2 Manhattan Distance

Formula to calculate Manhattan Distance between two points (x_1, y_1) and (x_2, y_2)

$$d = |y_2 - y_1| + |x_2 - x_1|$$

2.3 Types of Norms

L2 Norm of a Vector

Formula for L2 Norm or Euclidean Distance of a vector of d dimension:

$$\begin{aligned} \|\vec{x}\|_2 &= \sqrt{x_1^2 + x_2^2 + x_3^2 + \dots + x_d^2} \\ &= \sqrt{\sum_{i=1}^d x_i^2} \end{aligned}$$

L1 Norm of a Vector

Formula for L1 Norm or Manhattan Distance of a vector of d dimension:

$$\begin{aligned} \|\vec{x}\|_1 &= |x_1| + |x_2| + |x_3| + \dots + |x_d| \\ &= \sum_{i=1}^d |x_i| \end{aligned}$$

3 Matrix Multiplication

3.1 Matrix Multiplication rule

Given two matrices A and B where:

- $A : m \times n$
- $B : n \times k$

Rule

Number of columns in the first matrix should be equal to the number of rows in the second matrix.

$$m \times n \Leftrightarrow n \times k = m \times k$$

The resulting matrix will be of shape: $m \times k$

3.2 Matrix Multiplication in NumPy

Given two NumPy matrices `m1` and `m2`

```
m1 @ m2  
# or  
np.matmul(m1, m2)
```

4 Dot Product

4.1 Dot Product of Vectors

Formula to calculate dot product of two vectors $\vec{a}$ and $\vec{b}$:

$$\vec{a} \cdot \vec{b} = a^T \cdot b$$

$$= \sum_{i=1}^d a_i b_i$$

4.2 Dot Product in NumPy

Given two NumPy vectors `v1` and `v2`

```
np.dot(v1, v2)
```

4.3 Application of Dot Product

1 Angle between Vectors

Formula to calculate angle between two vectors $\vec{a}$ and $\vec{b}$

$$\cos \theta = \frac{\vec{a}^T \cdot \vec{b}}{\|\vec{a}\| \|\vec{b}\|}$$

2 Machine Learning

In ML context, the dot product is used to represent the below "best fit" line or hyper-plane equations:

$$w_1x_1 + w_2x_2 + w_0 = 0$$

or

$$w_1x_1 + w_2x_2 + \dots + w_dx_d + w_0 = 0$$

using $\vec{w}$ and $\vec{x}$ vectors of d dimensions:

$$\vec{w} = \begin{bmatrix} w_1 \\ w_2 \\ \vdots \\ w_d \end{bmatrix} \quad \vec{x} = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_d \end{bmatrix}$$

as,

$$\vec{w}^T \cdot \vec{x} + w_0 = 0$$

Where $\vec{w}^T \cdot \vec{x}$ is the dot product between weight vector $\vec{w}$ and feature vector $\vec{x}$.

3 Projection of a Vector

$$\text{Projection of } \vec{x} \text{ on } \vec{y} = \vec{x} \cdot \hat{y}$$

Unit Vector

Formula to calculate unit vector $\hat{w}$ of a $\vec{w}$ vector:

$$\hat{w} = \frac{\vec{w}}{\|\vec{w}\|}$$

Formula to calculate projection of $\vec{x}$ on $\vec{y}$:

$$\text{Projection of } \vec{x} \text{ on } \vec{y} = \frac{\vec{x} \cdot \vec{y}}{\|\vec{y}\|}$$

5 Shifting Lines

5.1 One Direction

Given line equation:

$$w_1x_1 + w_2x_2 + w_0 = 0$$

Shifting lines in one direction (x-axis or y-axis) by a units:

x-axis	Right $\rightarrow$	$w_1(x_1 - a)$	+	w_2x_2	+	$w_0 = 0$
x-axis	Left $\leftarrow$	$w_1(x_1 + a)$	+	w_2x_2	+	$w_0 = 0$
y-axis	Up $\uparrow$	w_1x_1	+	$w_2(x_2 - a)$	+	$w_0 = 0$
y-axis	Down $\downarrow$	w_1x_1	+	$w_2(x_2 + a)$	+	$w_0 = 0$

5.2 Two Directions

Shifting lines in two direction (x-axis and y-axis) simultaneously by a units:

$$w_1(x_1 - a) + w_2(x_2 - a) + w_0 = 0$$


Solving above equation gives:

$$w_1x_1 + w_2x_2 + w'_0 = 0$$

Where w'_0 is:

$$w'_0 = w_0 - a(w_1 + w_2)$$

5.3 Usecase



6 Distances

6.1 Distance Between Line and Origin

Formula to calculate the distance between line (Decision Boundary) and origin using line's Normal vector $\vec{w}$

$$d = \frac{-w_0}{\|w\|}$$

6.2 Distance Between Point and Line

Formula to calculate the distance between a point x_0 and line (Decision Boundary) using line's Normal vector $\vec{w}$

$$d = \frac{w^T x_0 + w_0}{\|w\|}$$

6.3 Distance Between Parallel Lines

Same Normal Vector

Formula to calculate the distance between two parallel lines with Normal vector w having same magnitude:

$$d = \frac{w_0^{(1)} - w_0^{(2)}}{\|w\|}$$

Different Normal Vector

Formula to calculate the distance between two parallel lines with Normal vectors of different magnitudes where $w^{(1)} = \lambda \cdot w^{(2)}$

$$d = \frac{w_0^{(1)}}{\|w^{(1)}\|} - \frac{w_0^{(2)}}{\|w^{(2)}\|}$$

7 Binary Classification

7.1 Half-spaces

Positive Half-space

In Positive Half-space all data-points have **distance greater than zero** because,

$$w_1x_1 + w_2x_2 + w_0 > 0$$

Negative Half-space

In Negative Half-space all data-points have **distance less than zero** because,

$$w_1x_1 + w_2x_2 + w_0 < 0$$

Decision Boundary

On Decision Boundary all data-points have **distance equal to zero** because,

$$w_1x_1 + w_2x_2 + w_0 = 0$$

7.2 Mathematical Notation

Dataset

Equation to represent a labeled dataset D used in Binary Classification:

$$D := \left\{ (x_i, y_i) ; x \in \mathbb{R}^d ; y \in \{-1, +1\} \right\}_{i=1}^n$$

Where,

- D is Dataset
- x are features
- y are labels
- n is the size of the dataset

Label

Actual Label y can take values:

$$y \begin{cases} +1 & \text{in +ve Half-Space} \\ -1 & \text{in -ve Half-Space} \end{cases}$$

Classifier

Mapping function for Binary Classification:

$$\hat{y} = f(x) = w_1x_1 + w_2x_2 + \dots + w_dx_d + w_0$$

or

$$= w^T x + w_0$$

Predicted label $\hat{y}$ can take values:

$$\hat{y} \begin{cases} +1 & \text{if } (w^T x + w_0) \geq 0 \\ -1 & \text{if } (w^T x + w_0) < 0 \end{cases}$$

7.3 Gain Function

Equation of Gain Function for Binary Classification:

$$g(D, \vec{w}, w_0) = \frac{1}{n} \sum_{i=1}^n \left(\frac{w^T x^{(i)} + w_0}{\|w\|} \right) \cdot y^{(i)}$$

7.4 Loss Function

$$\text{Loss Function} = - \text{Gain Function}$$

Equation of Loss Function for Binary Classification:

$$l(D, \vec{w}, w_0) = \frac{1}{n} \sum_{i=1}^n \left(\frac{w^T x^{(i)} + w_0}{\|w\|} \right) \cdot y^{(i)}$$

7.5 Perceptron Learning Algorithm

Weight Updation Logic

Logic in Perceptron Learning Algorithm to update Weights for miss-classified data-point:

$$\text{if } \text{sign} \left(\frac{w^T x^{(i)} + w_0}{\|w\|} \right) \neq \text{sign}(y^{(i)}) :$$

$$w = w + x^{(i)} \cdot y^{(i)}$$

$$w_0 = w_0 + y^{(i)}$$

8 Optimization

8.1 Optimization of Gain Function

Equation for Optimization of Gain Function for Binary Classification:

$$\begin{aligned}w^*, w_0^* &= \arg \max_{w, w_0} g(D, \vec{w}, w_0) \\&= \arg \max_{w, w_0} \frac{1}{n} \sum_{i=1}^n \left(\frac{w^T x^{(i)} + w_0}{\|w\|} \right) \cdot y^{(i)}\end{aligned}$$

Goal

Maximize Gain function.

8.2 Optimization of Loss Function


Equation for Optimization of Loss Function for Binary Classification:

$$\begin{aligned}w^*, w_0^* &= \arg \min_{w, w_0} l(D, \vec{w}, w_0) \\&= \arg \min_{w, w_0} \frac{1}{n} \sum_{i=1}^n \left(\frac{w^T x^{(i)} + w_0}{\|w\|} \right) \cdot y^{(i)}\end{aligned}$$

Goal

Minimize Loss function.

ML Context



Error vs Classifier is represented as a function $f(x)$ which needs to be minimized.

9 Functions


9.1 Continuous Function

Definition


Criteria for a Continuous function:

$$\lim_{x \rightarrow a^-} f(x) = \lim_{x \rightarrow a^+} f(x) = f(a)$$

Example #1



Example #2



Note:

Even though $f(x) = |x|$ is continuous function, it is **not differentiable at $x = 0$**


9.2 Discontinuous Function

Definition


Criteria for a Discontinuous function:

$$\lim_{x \rightarrow a^-} f(x) \neq \lim_{x \rightarrow a^+} f(x)$$

Example #1




Example #2



10 Differentiation

10.1 Secant


Slope of Line



Formula to calculate slope of a line intersecting two points (x_1, y_1) and (x_2, y_2)

$$m = \tan \theta = \frac{y_2 - y_1}{x_2 - x_1}$$

Slope of Secant Line




Formula for slope of a secant line intersecting a curve at two points (x_1, y_1) and (x_2, y_2)

$$m = \frac{f(x_2) - f(x_1)}{x_2 - x_1}$$

10.2 Tangent

Slope of Tangent Line (Derivative)



Formula for slope of a tangent line intersecting a curve at one single point (x, y)

$$f'(x) = \frac{d}{dx} f(x) = \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x}$$

10.3 Derivatives Examples

Commonly used Derivatives

$$(1) \quad \frac{d}{dx} x^n = nx^{n-1}$$

$$(5) \quad \frac{d}{dx} \sin(x) = \cos(x)$$

$$(2) \quad \frac{d}{dx} \log(x) = \frac{1}{x}$$

$$(6) \quad \frac{d}{dx} \cos(x) = -\sin(x)$$

$$(3) \quad \frac{d}{dx} e^x = e^x$$

$$(7) \quad \frac{d}{dx} \tan(x) = \sec^2(x)$$

$$(4) \quad \frac{d}{dx} c = 0$$

$$(8) \quad \frac{d}{dx} b^x = b^x \log(b)$$

10.4 Rules of Differentiation

Constant Rule

$$\frac{d}{dx} [c \cdot f(x)] = c \cdot \frac{d}{dx} f(x)$$

Sum Rule

$$\frac{d}{dx} [f(x) + g(x)] = \frac{d}{dx} f(x) + \frac{d}{dx} g(x)$$

Difference Rule

$$\frac{d}{dx} [f(x) - g(x)] = \frac{d}{dx} f(x) - \frac{d}{dx} g(x)$$

Product Rule

$$\frac{d}{dx} [f(x) \cdot g(x)] = g(x) \frac{d}{dx} f(x) + f(x) \frac{d}{dx} g(x)$$

Quotient Rule

$$\frac{d}{dx} \left[\frac{f(x)}{g(x)} \right] = \frac{g(x) \frac{d}{dx} f(x) - f(x) \frac{d}{dx} g(x)}{[g(x)]^2}$$

Chain Rule


$$\frac{d}{dx} f(g(x)) = \frac{d}{dx} f(g(x)) \cdot \frac{d}{dx} g(x)$$

How Chain rule works?

$$\begin{aligned} \frac{d}{dx} \log(x^2) &= \frac{1}{x^2} \cdot 2x \\ &= \frac{2}{x} \end{aligned}$$

- Keep the inside x^2 as it is and take derivative of outside $\log$ i.e., $\frac{1}{x^2}$.
- Multiply it by the derivative of the inside x^2 i.e., $2x$.

10.5 Critical Points



Definition

Slope of the Tangent line to a curve (i.e., derivative of a function) will be **zero at Critical Points**.

$$f'(x) = 0$$

10.6 Local Optima

Types of Local Optima

Optimal points where $f'(x) = 0$ (First derivative). Types:

1. Local Minima
2. Local Maxima

$$f''(x) \begin{cases} f''(x) > 0 & \text{Local Minima} \\ f''(x) < 0 & \text{Local Maxima} \end{cases}$$

Saddle Point

Non-optimal points where both $f'(x) = 0$ (First derivative) and $f''(x) = 0$ (Second derivative).

$$f''(x) = 0 \quad \text{Saddle point}$$

10.7 General Framework

First and Second Derivative Test for Extrema

STEP 1: Compute First Derivative

Given function $f(x)$ compute its first derivative:

$$f'(x)$$

STEP 2: Find Critical Point

Equate the first derivative to zero to find the critical point:

$$f'(x) = 0$$

Critical Point :- Critical value of x , say c , at which $f'(x) = 0$.

STEP 3: Classify Using Second Derivative

To classify the Critical point from STEP 2 to the correct type of local Optima, compute derivative of $f'(x)$ from STEP 1:

$$f''(x)$$

Identify the type of Local Optima based on below rule:

$$f''(x) \begin{cases} \text{if } f''(x) > 0 : & \text{Local Minima} \\ \text{if } f''(x) < 0 : & \text{Local Maxima} \\ \text{if } f''(x) = 0 : & \text{Test Inconclusive} \end{cases}$$

STEP 4: Calculate Local Optima Value

Substitute the Critical Value c from STEP 2 in original function $f(x)$ to identify the actual value of Local Optima.

$$f(c) \in \mathbb{R}$$