



Department of Physics and Electronics

**Masters Thesis Presentation**

**Investigating temperature variations of solar corona during Coronal Mass Ejections**

**By**

**V DHEERAJ SHENOY**

**USN: 22MSRPH022**

**Guide**

**M N Sundar**

**Jain University**

**Co-Guide**

**Dr. Tanmoy Samanta**

**Indian Institute of Astrophysics**

# Introduction


- Solar Corona
- Filaments/Prominences
- Flares
- Coronal Mass Ejections (CMEs)
- Stellar CMEs
- Coronal Dimming
- Emission Measure & Differential Emission Measure

# Objectives

1. Understand the temperature variation of solar corona during Coronal Mass Ejections (CME)
2. Which temperature plasma shows the most coronal dimming ?
3. Differential Emission Measure (DEM) analysis of point source converted Sun.

# Solar Corona

- Outermost part of atmosphere of Sun
- Several million degree Kelvin
- Contains forbidden Iron lines (Fe XIV)
- Very low density compared to the lower layers




Solar corona seen during total solar eclipse  
(Image credit: NASA)

Temperature variation of the Solar Corona during Coronal Mass Ejections

# Solar flares


- Explosive phenomena due to magnetic reconnection
- Accompanied by filament/prominence eruptions and CMEs



Filament



Prominence (Image credit:  
NASA/SDO)



Flaring event of 2011 August 4<sup>th</sup>

# Coronal Mass Ejections

- Plasma Ejections into Heliosphere
- Often associated with flares
- Very high association of phenomena of coronal dimming
- Made up of charged particles
- High magnetic field enclosure
- Affects the space weather, satellite communications etc.



LASCO C3 image of a CME  
(Image credit: SOHO/LASCO (ESA & NASA))

Temperature variation of the Solar Corona during Coronal Mass Ejections

# Stellar CMEs

- Like solar CMEs
- Highly energetic
- Difficult to observe
- No spatial resolution
- Indirect methods to detect stellar CMEs
  - Coronal Dimming
  - Blueshifted chromospheric lines
  - Type-II and Type-IV radio bursts etc.


# Literature Survey

- EM distributions from SDO/AIA data alone can overestimate the amount of high temperature ( $\log T > 6.4$ ) plasma in the solar corona by a factor of 3-15 (Athiray and Winebarger, 2024)
- About 73 % association of CME events with eruptive filaments has been observed (Sinha et al., 2019)
- Information about the underlying CMEs like mass and velocity of the CMEs ejected by knowing the depth and slope of the dimming curve (Mason et al., 2016)
- The association of dimming with a CMEs and association of CME with dimming has been observed to be very high (Veronig et al. 2021)
- Kinetic energy of stellar CME to the stellar flaring energy not being anywhere close to being a scaled version of the solar case (Namekata et al., 2022).
- Redshifted components of stellar filament eruptions in Sun-as-a-star analysis in H $\alpha$  spectra may develop into CMEs (Otsu et al., 2022)


# Methodology

- 1) Downloading level 1 Images of Sun
- 2) DEM is obtained for the full disk image of Sun.
- 3) *Conversion of full disk to point source*: full disk data is converted to a point source (pointification).
- 4) Calculating DEM for the point source
- 5) Compare the DEM curves of full disk and point source.
- 6) If CME is associated with dimming, find the maximum dimming temperature region.
- 7) Study the DEM profile which gives information about the coronal temperature variation

# Coronal Dimming



Dimming seen on a star. Image Credit: (Veronig et al.)




211 Å channel of AIA. Yellow boxes shows the dimming

Temperature variation of the Solar Corona during Coronal Mass Ejections

# Differential Emission Measure (DEM)

- DEM - emission measure with respect to temperature. Function of temperature.



Column emission  
measure

$$EM = \int_h n_e^2 dh$$

$$DEM(T) = n_e^2(T) \frac{dh}{dT}$$

Where  $n_e$  is the electron number density

# DEM Inversion

- Obtain the information about the plasma from the line intensities observed


$$y_i = \int_0^{\infty} K_i(T) DEM(T) dT$$

$K_i$  is the temperature response of each channel


- We can only approximate the underlying EM profile.
- We use Reiterative Maximum Likelihood (RML) method (Massa et al., 2023)

# DEM of Sun during the flaring event of August 4<sup>th</sup> 2011


log T=5.85




log T=6.0




log T=6.15




log T=6.3



Temperature variation of the Solar Corona during Coronal Mass Ejections



# “Pointification” of sun full disk image




a) shows the full disk image of Sun. (b) Image after pointification

Temperature variation of the Solar Corona during Coronal Mass Ejections

# Data

- NASA's Solar Dynamics Observatory - Atmospheric Imaging Assembly (SDO/AIA)
- Three CME event data
  - Flare associated CME (August 4<sup>th</sup> 2011)
  - Long filament eruption associated with a CME (August 31<sup>st</sup> 2012)
  - Ground Level Enhancement CME (21<sup>st</sup> October 2021)



NASA's SDO Space Observatory  
(Image Credit: NASA/SDO)




Four telescopes of SDO/AIA (Image Credit: NASA/SDO)

Temperature variation of the Solar Corona during Coronal Mass Ejections

# Results


Temperature variation of the Solar Corona during Coronal Mass Ejections

# Coronal Dimming



Temperature variation of the Solar Corona during Coronal Mass Ejections


# Filament Eruption



DEM Timeseries of August 31<sup>th</sup> 2012 CME event

Temperature variation of the Solar Corona during Coronal Mass Ejections


# GLE CME event



DEM Timeseries of August 31<sup>th</sup> 2012 CME event

Temperature variation of the Solar Corona during Coronal Mass Ejections

# DEM profiles of 4<sup>th</sup> August 2011 Event




Before event

During event

After event

Temperature variation of the Solar Corona during Coronal Mass Ejections

# DEM profiles of 31<sup>st</sup> August 2012 Event




Before event

During event

After event

Temperature variation of the Solar Corona during Coronal Mass Ejections

# DEM profiles of 28<sup>th</sup> October 2021 Event



Before event

During event

After event

Temperature variation of the Solar Corona during Coronal Mass Ejections

# Conclusion

- Coronal dimming for 4<sup>th</sup> August 2011 event is most prominent in  $\log T = [5.85, 6.15]$  range. Mass loss is more in this temperature
- Good correlation between point source and full disk DEM solutions (below  $\log T = 6.45$ ).
- Characteristic features of the CMEs on the full disk is still visible in the point source
- Variations in coronal plasma temperature has been observed in both point source and full disk image before, during and after the event

# Moving forward

- Correcting the *pointification* process by not direct averaging.
- Look at events originating from various regions on the Sun
- Statistical analysis to check the correctness of the correlation

# Acknowledgements

- *M. N. Sundar, Jain University*
- *Dr. Tanmoy Samanta, Indian Institute of Astrophysics*
- *Dr. Dhanush Shenoy for his constant support*

# Publications and Conferences

- Manuscript of this work is under preparation to be submitted to the journal (**The Astrophysical Journal**).
- The two-days Scientific Extravaganza '**MillCon-2023**' is an Internation Conclave on Millets, held at Jain (Deemed-to-be University), School of Sciences. Best paper and presentation award for the paper titled "**Could AI/ML help yield loss in Millets?**".
- Paper titled "**Crop Health Managemenet in Millets using Satellite Imagery and Neural Networks**" has been submitted to **VEGETOS: An International Journal of Plant Research & Biotechnology**, The paper is currently under review for publication.
- National Space Science Symposium - 2024 Conference , Goa University: Poster presentation for the paper titled "**Crop Health Management in Millets using Satellite Imagery and Neural Networks**"


# References

- Athiray, P. S., & Winebarger, A. R. (2024). Can emission measure distributions derived from extreme-ultraviolet images accurately constrain high-temperature plasma? *The Astrophysical Journal*, 961(2), 181. <https://doi.org/10.3847/1538-4357/ad1837>
- Cheung, M. C. M., Boerner, P., Schrijver, C. J., Testa, P., Chen, F., Peter, H., & Malanushenko, A. (2015). Thermal diagnostics with the atmospheric imaging assembly on board the solar dynamics observatory: A validated method for differential emission measure inversions. *The Astrophysical Journal*, 807(2), 143. <https://doi.org/10.1088/0004-637x/807/2/143>
- Gopalswamy, N. (2016). History and development of coronal mass ejections as a key player in solar-terrestrial relationship. *Geosci. Lett.*, 3(1).
- Hannah, I. G., & Kontar, E. P. (2012). Differential emission measures from the regularized inversion of Hinode and SDO data. *Astronomy & Astrophysics*, 539, A146. <https://doi.org/10.1051/0004-6361/201117576>
- Klein, K.-L., Musset, S., Vilmer, N., Briand, C., Krucker, S., Francesco Battaglia, A., Dresing, N., Palmroos, C., & Gary, D. E. (2022). The relativistic solar particle event on 28 October 2021: Evidence of particle acceleration within and escape from the solar corona. *Astronomy & Astrophysics*, 663, A173. <https://doi.org/10.1051/0004-6361/202243903>
- Korhonen, H., Vida, K., Leitzinger, M., Odert, P., & Kovacs, O. E. (2016). Hunting for stellar coronal mass ejections. *Geosci. Lett.*, 3(1).
- Lemen, J. R., Title, A. M., Akin, D. J., Boerner, P. F., Chou, C., Drake, J. F., Duncan, D. W., Edwards, C. G., Friedlaender, F. M., Heyman, G. F., Hurlburt, N. E., Katz, N. L., Kushner, G. D., Levay, M., Lindgren, R. W., Mathur, D. P., McFeaters, E. L., Mitchell, S., Rehse, R. A., ... Walther, N. (2011). The atmospheric imaging assembly (AIA) on the solar dynamics observatory (SDO). *Solar Physics*, 275(1–2), 17–40. <https://doi.org/10.1007/s11207-011-9776-8>
- Mason, J. P., Woods, T. N., Caspi, A., Thompson, B. J., & Hock, R. A. (2014). Mechanisms and observations of coronal dimming for the 2010 August 7 event. *The Astrophysical Journal*, 789(1), 61. <https://doi.org/10.1088/0004-637x/789/1/61>
- Mason, J. P., Woods, T. N., Webb, D. F., Thompson, B. J., Colaninno, R. C., & Vourlidas, A. (2016). Relationship of EUV irradiance coronal dimming slope and depth to coronal mass ejection speed and mass. *Astrophys. J.*, 830(1), 20.
- Massa, P., Emslie, A. G., Hannah, I. G., & Kontar, E. P. (2023). Robust construction of differential emission measure profiles using a regularized maximum likelihood method. *Astronomy & Astrophysics*, 672, A120. <https://doi.org/10.1051/0004-6361/202345883>
- Morgan, H., & Pickering, J. (2019). Sites: Solar iterative temperature emission solver for differential emission measure inversion of EUV observations. *Solar Physics*, 294(10). <https://doi.org/10.1007/s11207-019-1525-4>
- Moschou, S.-P., Drake, J. J., Cohen, O., Alvarado-Gómez, J. D., Garraffo, C., & Fraschetti, F. (2019). The stellar CME-flare relation: What do historic observations reveal? *Astrophys. J.*, 877(2), 105.
- Namekata, K., Maehara, H., Honda, S., Notsu, Y., Nogami, D., & Shibata, K. (2022). Hunting for stellar coronal mass ejections. *Geosci. Lett.*, 3(1).
- Namekata, K., Toriumi, S., Airapetian, V. S., Shoda, M., Watanabe, K., & Notsu, Y. (2023). Reconstructing the XUV spectra of active sun-like stars using solar scaling relations with magnetic flux. *Astrophys. J.*, 945(2), 147.
- Otsu, T., & Asai, A. (2024). Multiwavelength sun-as-a-star analysis of the m8.7 flare on 2022 October 2 using H and EUV spectra taken by Smart/SDDI and SDO/EVE. *The Astrophysical Journal*, 964(1), 75. <https://doi.org/10.3847/1538-4357/ad24ec>
- Otsu, T., Asai, A., Ichimoto, K., Ishii, T. T., & Namekata, K. (2022). Sun-as-a-star analyses of various solar active events using H spectral images taken by Smart/SDDI. *The Astrophysical Journal*, 939(2), 98. <https://doi.org/10.3847/1538-4357/ac9730>
- Pesnell, W. D., Thompson, B. J., & Chamberlin, P. C. (2011). The solar dynamics observatory (SDO). *Solar Physics*, 275(1–2), 3–15. <https://doi.org/10.1007/s11207-011-9841-3>
- Plowman, J., Kankelborg, C., & Martens, P. (2013). Fast differential emission measure inversion of solar coronal data. *The Astrophysical Journal*, 771(1), 2. <https://doi.org/10.1088/0004-637x/771/1/2>
- Sinha, S., Srivastava, N., & Nandy, D. (2019). Solar filament eruptions as precursors to flare-CME events: Establishing the temporal connection. *The Astrophysical Journal*, 880(2), 84. <https://doi.org/10.3847/1538-4357/ab2239>
- Su, Y., Veronig, A. M., Hannah, I. G., Cheung, M. C. M., Dennis, B. R., Holman, G. D., Gan, W., & Li, Y. (2018). Determination of differential emission measure from solar extreme ultraviolet images. *Astrophys. J. Lett.*, 856(1), L17.
- Veronig, A. M., Odert, P., Leitzinger, M., Dissauer, K., Fleck, N. C., & Hudson, H. S. (2021). Indications of stellar coronal mass ejections through coronal dimmings.

# Thank you


# Solar structure

- Interior region
  - Core
  - Radiative zone
  - Convective zone
- Visible surface
  - Photosphere
  - Atmosphere
  - Chromosphere + Corona



Structure of Sun (Image credit: NASA)

Temperature variation of the Solar Corona during Coronal Mass Ejections



# Magnetic Reconnection






Image credit: Wikipedia





AIA telescope layout. Telescope 2 has aperture blade to select between it's two wavelength channels. Rest of the channels rely on the filters in filter wheels to select the desired channels.

<b>Event</b>	<b>Pearson Correlation Coeffecient</b>		
	$\log T=[5.85, 6.15]$	$\log T=[6.15, 6.45]$	$\log T=[6.45, 6.75]$
4 <sup>th</sup> August 2011	0.9449	0.9767	0.2190
31 <sup>st</sup> August 2012	0.7027	0.9885	0.2079
28 <sup>th</sup> October 2021	0.9555	0.9577	0.2578