

Solving the Optimal Experiment Design Problem using Convex Mixed-Integer methods

Deborah Hendrych (ZIB, TUB), Mathieu Besançon (Inria, UGA), Sebastian
Pokutta (ZIB, TUB)

hendrych@zib.de

Zuse Institute Berlin

STOGO 2025 Stockholm · September 3, 2025



Berlin Mathematics Research Center



**ZUSE
INSTITUTE
BERLIN**



Think cinnamon rolls



Outline

1. The Optimal Experiment Design Problem (OEDP)
2. Integer Frank-Wolfe: `Boscia.jl`
3. A- and D-Optimal Design
 - Convergence analysis
 - Computational experiments
4. E-Optimal Design
 - Smoothing
 - Initial Experiments
5. Outlook

Table of Contents

The Optimal Experiment Design Problem (OEDP)

Integer Frank-Wolfe: `Boscia.jl`

A- and D-Optimal Design

Convergence analysis

Computational experiments

E-Optimal Design

Smoothing

Initial Experiments

Outlook

Set-up

- The ultimate goal is fitting a regression model.

$$\min_{\theta \in \mathbb{R}^n} \|A\theta - \mathbf{y}\|_2^2$$

- $A = [-\mathbf{a}_i -]_{i=1}^m$ with $\mathbf{a}_i \in \mathbb{R}^n$ is the *Experiment Matrix*. Assumed to have full column rank.
- θ is the unknown set of parameters to be found.
- \mathbf{y} is the **not-yet** measured response.
- **Issue:** Running all m experiments too costly and/or too time intensive.


Some information theory later ...

Pukelsheim 2006

Information matrix: $X(\mathbf{x}) := A^T \text{diag}(\mathbf{x})A$

D-Optimal Experiment Design

$$\begin{aligned} \min_{\mathbf{x}} \quad & -\log \det (A^{\top} \operatorname{diag}(\mathbf{x}) A) \\ \text{s.t.} \quad & \sum_{i=1}^m x_i = N \\ & \mathbf{l} \leq \mathbf{x} \leq \mathbf{u} \\ & \mathbf{x} \in \mathbb{N}_0^m, \end{aligned} \quad (\text{D})$$



Ponte, Fampa, and Lee 2025; Ahipaşaoğlu 2021; Li et al. 2024

A-Optimal Experiment Design


$$\min_{\mathbf{x}} \operatorname{tr} \left((A^T \operatorname{diag}(\mathbf{x}) A)^{-1} \right)$$

$$\text{s.t. } \sum_{i=1}^m x_i = N$$

$$\mathbf{l} \leq \mathbf{x} \leq \mathbf{u}$$

$$\mathbf{x} \in \mathbb{N}_0^m,$$


(A)



Ahipaşaoğlu 2015; Sagnol and Pauwels 2019; Nikolov, Singh, and Tantipongpipat 2022

E-Optimal Experiment Design

$$\begin{aligned} \min_{\mathbf{x}} \quad & \text{eigmax}(-A^T \text{diag}(\mathbf{x})A) \\ \text{s.t.} \quad & \sum_{i=1}^m x_i = N \\ & \mathbf{l} \leq \mathbf{x} \leq \mathbf{u} \\ & \mathbf{x} \in \mathbb{N}_0^m, \end{aligned} \quad (\text{D})$$



Ponte, Fampa, and Lee 2025; Ahipaşaoğlu 2021; Li et al. 2024

Table of Contents

The Optimal Experiment Design Problem (OEDP)

Integer Frank-Wolfe: `Boscia.jl`

A- and D-Optimal Design

Convergence analysis

Computational experiments


E-Optimal Design

Smoothing

Initial Experiments

Outlook

Integer Frank-Wolfe: Boscia.jl




Frank and Wolfe 1956; Levitin and Polyak
1966; Braun et al. 2022

$$\begin{aligned} \min_{\mathbf{x}} \quad & f(\mathbf{x}) \\ \text{s.t. } \quad & \mathbf{x} \in \mathcal{X} \\ & \mathbf{x}_j \in \mathbb{Z} \quad \forall j \in J \end{aligned}$$

- f continuously differentiable, convex, L -smooth.
- \mathcal{X} is a compact convex set, usually polyhedral.

Hendrych, Troppens, et al. 2025



Schematic of the algorithm.

Table of Contents

The Optimal Experiment Design Problem (OEDP)

Integer Frank-Wolfe: `Boscia.jl`

A- and D-Optimal Design

Convergence analysis

Computational experiments

E-Optimal Design

Smoothing

Initial Experiments

Outlook

Feasible region vs Domain




Figure: A schematic representation of the feasible region \mathcal{P} , the domain of the objective \mathcal{D} and the convex hull of vertices that are both feasible and in the domain denoted as \mathcal{W} .

Generalized self-concordance

By Carderera, Besançon, and Pokutta 2021, we also have convergence if the objective is generalized self-concordant. Self-concordance was already proved for the $-\log \det(X)$ on \mathbb{S}_{++}^n .

Generalized self-concordance

By Carderera, Besançon, and Pokutta 2021, we also have convergence if the objective is generalized self-concordant. Self-concordance was already proved for the $-\log \det(X)$ on \mathbb{S}_{++}^n .

Theorem (GSC A-criterion) Hendrych, Besançon, and Pokutta 2024

The function $g(X) = \operatorname{tr}(X^{-p})$, with $p > 0$, is $\left(3, \frac{(p+2)\sqrt[p]{a^{2pn}}}{\sqrt{p(p+1)}}\right)$ -generalized self-concordant on $\operatorname{dom}(g) = \{X \in \mathbb{S}_{++}^n \mid 0 \prec X \preceq aI\}$ where $a \in \mathbb{R}_{>0}$ bounds the maximum eigenvalue of X .

Other approaches

Co-BnB Ahipaşaoğlu 2021

- Branch-and-Bound with customized coordinate-descent-like algorithm for the node problems, exploits connection to Minimum Volume Enclosing Ellipsoid Problem.
- Developed for $m \gg n$.

Outer Approximation approaches

- Using the mixed-integer convex solver with conic certificates `Pajarito.jl` Coey, Kapelevich, and Vielma 2022a.
 1. **Direct Conic:** Natural formulation via `Hypatia.jl` Coey, Kapelevich, and Vielma 2022b.
 2. **SOCP:** as introduced in Sagnol 2011; Sagnol and Harman 2015.

Experimental Results I

Termination over time for the A-Optimal Problem




Figure: Independent data




Figure: Correlated data

Experimental Results II

Termination over time for the D-Optimal Problem




Figure: Independent data




Figure: Correlated data

Table of Contents

The Optimal Experiment Design Problem (OEDP)

Integer Frank-Wolfe: `Boscia.jl`

A- and D-Optimal Design

Convergence analysis

Computational experiments

E-Optimal Design

Smoothing

Initial Experiments

Outlook

E-Criterion and smoothing

Issue

The subgradients of $\lambda_i(X)$ are the corresponding eigenvectors.
 \Rightarrow the gradient $\nabla \lambda_i$ only exists if $\text{mult}(\lambda_i) = 1$.

E-Criterion and smoothing

Issue

The subgradients of $\lambda_i(X)$ are the corresponding eigenvectors.
 \Rightarrow the gradient $\nabla \lambda_i$ only exists if $\text{mult}(\lambda_i) = 1$.

Solution - Smoothing

We use the smoothing introduced in Nesterov 2007.

$$\max \lambda_{\min}(X) \Leftrightarrow \min \lambda_{\max}(-X) := f(X)$$

E-Criterion and smoothing

Issue

The subgradients of $\lambda_i(X)$ are the corresponding eigenvectors.
 \Rightarrow the gradient $\nabla \lambda_i$ only exists if $\text{mult}(\lambda_i) = 1$.

Solution - Smoothing


We use the smoothing introduced in Nesterov 2007.

$$\max \lambda_{\min}(X) \Leftrightarrow \min \lambda_{\max}(-X) := f(X)$$

$$f_{\mu}(X) = \mu \log \left(\sum_{i=1}^n e^{-\lambda_i(X)/\mu} \right) - \mu \log n \quad (1)$$

$$f_{\mu}(X) \leq f(X) \leq f_{\mu}(X) + \mu \log n \quad \forall X \in \mathbb{S}^n \quad (2)$$

Initial Computational Experiments I



Initial Computational Experiments II

Table: Heuristic Performance Comparison

Heuristic	Optimal Time (s)	Dual Gap
Baseline	6.092e+02 */ 5.825	3.204e-01 */ 1.794
Fedorov Fedorov 2013	1.979e+03 */ 2.236	6.231e-01 */ 2.840
Follow Subgradient	1.738e+03 */ 2.486	6.302e-01 */ 2.810
Pipage Rounding Brown, Laddha, and Singh 2024	2.114e+03 */ 1.615	9.723e-01 */ 2.765
Simple Randomized Rounding Lamperski, Yang, and Prokopyev n.d.	4.962e+02 */ 6.893	1.510e-01 */ 1.752

Initial Computational Experiments III




Table of Contents

The Optimal Experiment Design Problem (OEDP)

Integer Frank-Wolfe: `Boscia.jl`

A- and D-Optimal Design

Convergence analysis

Computational experiments

E-Optimal Design

Smoothing

Initial Experiments

Outlook

Outlook

E-Optimal Design

- Adapt lower bounds from Li 2025.
- Lower bound on the distance in the objective.
- More extensive experiments.

Outlook

E-Optimal Design

- Adapt lower bounds from Li 2025.
- Lower bound on the distance in the objective.
- More extensive experiments.

On the Optimal Design side

- Sequential designs.
- Knowledge of experiment matrix $A \rightarrow$ Robustness.
- Non-Linear regression and nearly convex functions.

Outlook

E-Optimal Design

- Adapt lower bounds from Li 2025.
- Lower bound on the distance in the objective.
- More extensive experiments.

On the Optimal Design side

- Sequential designs.
- Knowledge of experiment matrix $A \rightarrow$ Robustness.
- Non-Linear regression and nearly convex functions.

On the general Boscia.jl side

- Extended smoothness conditions.
- Handling of non-convex objectives.

Thank you for your attention!

Convex mixed-integer optimization
with Frank–Wolfe methods



Solving the Optimal Experiment
Design Problem with Mixed-Integer
Convex Methods



References I

- Ahipařaoğlu, Selin Damla (2015). “A first-order algorithm for the A-optimal experimental design problem: a mathematical programming approach”. In: *Statistics and Computing* 25.6, pp. 1113–1127.
- (2021). “A branch-and-bound algorithm for the exact optimal experimental design problem”. In: *Statistics and Computing* 31.5, p. 65.
- Braun, Gábor et al. (2022). “Conditional gradient methods”. In: *arXiv preprint arXiv:2211.14103*.
- Brown, Adam, Aditi Laddha, and Mohit Singh (2024). “Maximizing the minimum eigenvalue in constant dimension”. In: *arXiv preprint arXiv:2401.14317*.
- Carderera, Alejandro, Mathieu Besançon, and Sebastian Pokutta (2021). “Simple steps are all you need: Frank–Wolfe and generalized self-concordant functions”. In: *Advances in Neural Information Processing Systems* 34, pp. 5390–5401.
- Coey, Chris, Lea Kapelevich, and Juan Pablo Vielma (2022a). “Performance enhancements for a generic conic interior point algorithm”. In: *Mathematical Programming Computation*. DOI: <https://doi.org/10.1007/s12532-022-00226-0>.
- (2022b). “Solving natural conic formulations with Hypatia.jl”. In: *INFORMS Journal on Computing* 34.5, pp. 2686–2699. DOI: <https://doi.org/10.1287/ijoc.2022.1202>.
- Coey, Chris, Miles Lubin, and Juan Pablo Vielma (2020). “Outer approximation with conic certificates for mixed-integer convex problems”. In: *Mathematical Programming Computation* 12.2, pp. 249–293.
- Fedorov, Valerii Vadimovich (2013). *Theory of optimal experiments*. Elsevier.
- Frank, Marguerite and Philip Wolfe (1956). “An algorithm for quadratic programming”. In: *Naval research logistics quarterly* 3.1-2, pp. 95–110.



References I

- Hendrych, Deborah, Mathieu Besançon, and Sebastian Pokutta (2024). “Solving the Optimal Experiment Design Problem with Mixed-Integer Convex Methods”. en. In: Schloss Dagstuhl – Leibniz-Zentrum für Informatik. DOI: 10.4230/LIPICS.SEA.2024.16. arXiv: 2312.11200. URL: <https://drops.dagstuhl.de/entities/document/10.4230/LIPICS.SEA.2024.16>.
- Hendrych, Deborah, Hannah Troppens, et al. (June 28, 2025). “Convex Integer Optimization with Frank-Wolfe Methods”. In: *Mathematical Programming Computation*. DOI: 10.1007/s12532-025-00288-w. arXiv: 2208.11010 [math.OC]. URL: <https://link.springer.com/article/10.1007/s12532-025-00288-w>.
- Lamperski, Jourdain, Haeseong Yang, and Oleg Prokopyev (n.d.). “Simple Randomized Rounding for Max-Min Eigenvalue Augmentation”. In: *Forty-second International Conference on Machine Learning*.
- Levitin, Evgeny S and Boris T Polyak (1966). “Constrained minimization methods”. In: *USSR Computational mathematics and mathematical physics* 6.5, pp. 1–50.
- Li, Yongchun (2025). “Strong Formulations and Algorithms for Regularized A-optimal Design”. In: *arXiv preprint arXiv:2505.14957*.
- Li, Yongchun et al. (2024). “D-Optimal Data Fusion: Exact and Approximation Algorithms”. In: *INFORMS Journal on Computing* 36.1, pp. 97–120.
- Nesterov, Yurii (2007). “Smoothing technique and its applications in semidefinite optimization”. In: *Mathematical Programming* 110.2, pp. 245–259.

References III

- Nikolov, Aleksandar, Mohit Singh, and Uthaipon Tantipongpipat (2022). “Proportional volume sampling and approximation algorithms for A-optimal design”. In: *Mathematics of Operations Research* 47.2, pp. 847–877.
- Ponte, Gabriel, Marcia Fampa, and Jon Lee (2025). “Branch-and-bound for D-Optimality with fast local search and variable-bound tightening”. In: *Mathematical Programming*, pp. 1–38.
- Pukelsheim, Friedrich (2006). *Optimal design of experiments*. SIAM.
- Sagnol, Guillaume (2011). “Computing optimal designs of multiresponse experiments reduces to second-order cone programming”. In: *Journal of Statistical Planning and Inference* 141.5, pp. 1684–1708.
- Sagnol, Guillaume and Radoslav Harman (2015). “Computing exact D-optimal designs by mixed integer second-order cone programming”. In: *The Annals of Statistics* 43.5, pp. 2198–2224.
- Sagnol, Guillaume and Edouard Pauwels (2019). “An unexpected connection between Bayes A-optimal designs and the group lasso”. In: *Statistical Papers* 60.2, pp. 565–584.
- Zhao, Renbo (2025). “New Analysis of an Away-Step Frank-Wolfe Method for Minimizing Log-Homogeneous Barriers”. In: *Mathematics of Operations Research*.

Convex relaxation vs integer hull



Convergence results

Objective	Convergence	Linear Convergence
$f(\mathbf{x}) = -\log \det(X(\mathbf{x}))$	✓	✓*
$g(\mathbf{x}) = \text{tr}(X(\mathbf{x})^{-p})$	✓	Not guaranteed
$k(\mathbf{x}) = \log(\text{tr}(X(\mathbf{x})^{-p}))$	Conjecture	Not guaranteed

Table: Convergence of Frank-Wolfe on different objective functions.

*We adapted the proof from Zhao 2025 to the Blended Pairwise Conditional Gradient (BPCG).

Hendrych, D. and Besançon, M. and Pokutta, S. (2025). "Solving the Optimal Experiment Design Problem with Mixed-Integer Convex Methods." <https://arxiv.org/abs/2312.11200>

What do we need for convergence?

In short: Frank-Wolfe has to converge.

$$f(X) = -\log \det(X) \quad \text{and} \quad g(X) = \text{tr}(X^{-p}) \quad \text{and} \quad k(X) = \log \text{tr}(X^{-p})$$

We need:

- Convexity of the objective functions. ✓
- L -smoothness of the objective functions. → Issue with the domain.
- Generalized self-concordance of the objective functions.
- Strong convexity/Sharpness of the objective functions.

Hendrych, D. and Besançon, M. and Pokutta, S. (2025). "Solving the Optimal Experiment Design Problem with Mixed-integer Convex Methods." <https://arxiv.org/abs/2312.11200>

Theorem (L-smoothness for the Fusion Problems)

The functions $f_F(\mathbf{x}) = \log \det(X_C(\mathbf{x}))$, $g_F(\mathbf{x}) = \text{tr}(X_C(\mathbf{x})^{-p})$ and $k_F(\mathbf{x}) = \log(\text{tr}(X_C(\mathbf{x})^{-p}))$ are L-smooth on $x \in \mathbb{R}_{\geq 0}$.

L-smoothness

Theorem (L-smoothness for the Fusion Problems)

The functions $f_F(\mathbf{x}) = \log \det(X_C(\mathbf{x}))$, $g_F(\mathbf{x}) = \text{tr}(X_C(\mathbf{x})^{-p})$ and $k_F(\mathbf{x}) = \log(\text{tr}(X_C(\mathbf{x})^{-p}))$ are L-smooth on $x \in \mathbb{R}_{\geq 0}$.

Theorem (Local L-smoothness for the Optimal Problems)

The functions $f(X(\mathbf{x}))$, $g(X(\mathbf{x}))$ and $k(X(\mathbf{x}))$ are locally L-smooth on

$$\mathcal{L}_0 = \{\mathbf{x} \in \mathcal{D} \cap \mathcal{P} \mid (*) (\mathbf{x}) \leq (*) (\mathbf{x}_0)\}$$

where $()$ is a placeholder for each function and $x_0 \in \mathcal{D}$ an initial point.*

Sharpness

Theorem (Strong convexity)

The functions $f(X) = -\log \det(X)$ and $g(X) = \operatorname{tr}(X^{-p})$, $p > 0$, are strongly convex on $D := \{X \in \mathbb{S}_{++}^n \mid \lambda_{\max}(X) \leq \alpha\}$.

The function $k(X) = \log \operatorname{tr}(X^{-p})$, $p > 0$, is strongly convex on $D := \{X \in \mathbb{S}_{++}^n \mid \lambda_{\max}(X) \leq \alpha, \kappa(X) \leq \kappa\}$.

Sharpness

Theorem (Strong convexity)



The functions $f(X) = -\log \det(X)$ and $g(X) = \text{tr}(X^{-p})$, $p > 0$, are strongly convex on $D := \{X \in \mathbb{S}_{++}^n \mid \lambda_{\max}(X) \leq \alpha\}$.

The function $k(X) = \log \text{tr}(X^{-p})$, $p > 0$, is strongly convex on $D := \{X \in \mathbb{S}_{++}^n \mid \lambda_{\max}(X) \leq \alpha, \kappa(X) \leq \kappa\}$.

Theorem

The compositions of f , g and k with the information matrix $X(\mathbf{x})$, respectively, are sharp **on \mathcal{W}** .

In general, the compositions do not have a unique minimizer.



A Custom Branch-and-Bound for OEDP under Matrix Means (Co-BnB)

Ahipaşaoğlu 2021

- Coordinate-Descent-like algorithm for the nodes.
- Developed for $m \gg n$.

$$\begin{aligned} \max_{\mathbf{w}} \quad & \log(\phi(X(\mathbf{w}))) \\ \text{s.t.} \quad & \sum_{i=1}^m w_i = 1 \\ & \mathbf{w} \in [0, 1]^m \\ & N\mathbf{w} \in \mathbb{Z}_{\geq 0}^m \end{aligned} \tag{M-OEDP}$$

- \mathbf{w} can be interpreted as a probability distribution.
- Exploit connection to the Minimum Volume Enclosing Ellipsoid Problem (MVEP) for the termination criteria.

Direct Conic Formulation

coey2022conic; Coey, Kapelevich, and Vielma 2022a; Coey, Lubin, and Vielma 2020

- `Pajarito.jl` is a mixed-integer convex solver with conic certificates.
- `Hypatia.jl` is an interior point solver for conic optimization problems.
- D-Criterion:

$$\mathcal{K}_{\log \det} := \text{cl} \left\{ (u, v, W) \in \mathbb{R} \times \mathbb{R}_{>0} \times \mathbb{S}_{++}^n \mid u \leq v \log \det(W/v) \right\}$$

- A-Criterion: Dual of

$$\mathcal{K}_{\text{sepspec}} := \text{cl} \left\{ (u, v, w) \in \mathbb{R} \times \mathbb{R}_{>0} \times \text{int}(Q) \mid u \geq v \varphi(w/v) \right\}$$

- Q is the PSD cone and φ is the negative square root.
- The convex conjugate of the negative square root is the trace inverse.

Second-Order Conic Formulation (SOCP)

Sagnol 2011; Sagnol and Harman 2015

- In Sagnol 2011 the SOCP formulation for the continuous problem was introduced.
- The SOCP formulation of the exact formulation of OEDP was shown in Sagnol and Harman 2015.
- In theory, a nice result, but in practice, the problem size becomes much larger.
- For the A-Optimal Problem, we have $2m(n + 1)$ variables and $2(n + 1) + m$ constraints.
- For the D-Optimal Problem, we have $2m(1 + n) + n^2 + 1$ variables and $n(m + 1) + 3m + 4$ constraints.

SOCP A-Criterion

$$\begin{aligned} \min_{\substack{\mu \in \mathbb{R}^m \\ \mathbf{x} \in \mathbb{Z}^m \\ \forall i \in [m] \ Y_i \in \mathbb{R}^{1 \times n}}} \quad & \sum_{i=1}^m t_i \\ \text{s.t.} \quad & \sum_{i=1}^m A_i Z_i = I \\ & \forall i \in [m], \ \|Z_i\|_F^2 \leq t_i x_i \\ & \sum_{i=1}^m x_i = N \end{aligned} \quad (\text{A-SOCP})$$

SOCp D-Criterion

$$\min_{\substack{\mathbf{w} \in \mathbb{R}_{\geq 0}^m, \mathbf{x} \in \mathbb{Z}^m \\ J \in \mathbb{R}^{n \times n} \\ \forall i \in [m] Z_i \in \mathbb{R}^{1 \times n} \\ \forall i \in [m], j \in [n] t_{ij} \in \mathbb{R}_{\geq 0}}} \prod_{i=1}^m J_{ii}^{1/m}$$

$$\text{s.t. } \sum_{i=1}^m A_i Z_i = J$$

J lower triangle matrix

(D-SOCP)

$$\forall i \in [m], j \in [n] \|Z_i \mathbf{e}_j\|_F^2 \leq t_{ij} w_i$$

$$\forall j \in [n] \sum_{i=1}^m t_{ij} \leq J_{jj}$$

$$\mathbf{l} \leq \mathbf{x} \leq \mathbf{u} \text{ and } \sum_{i=1}^m w_i = 1$$

Effect of the conditioning of the problem




Figure: A IND




Figure: D IND




Figure: A CORR




Figure: D CORR