

2016/17 MPhil ACS / CST Part III
Category Theory and Logic (L108)
Exercise Sheet 1 – Solution Notes

Question 1

- (a) In Lecture 2 we saw that a morphism in **Set** is an isomorphism iff it is a bijection; but there is no bijection $3 \cong 2$, since any function $f : 3 \rightarrow 2$ cannot be injective.
- (b) Any function $f : Q \rightarrow P$ that is monotonic satisfies $f 0 \leq f 1$ in P and hence $f 0 = f 1$. So f is not a bijection. But any isomorphism in **Pre** is in particular an isomorphism in **Set** of the underlying sets (why?) and hence a bijection.
- (c) Recall that the set Q of rational numbers is countably infinite, that is, in bijection with \mathbb{N} ; so \mathbb{N} and Q are isomorphic in **Set**. However, as a pre-ordered set the rationals are dense: writing $x < y$ to mean $x \leq y \wedge x \neq y$, we have $(\forall x, y \in Q) x < y \Rightarrow (\exists z \in Q) x < z \wedge z < y$; whereas (\mathbb{N}, \leq) is not a dense pre-ordered set. It is not hard to see that the density property of pre-ordered sets is preserved under isomorphism. So (\mathbb{N}, \leq) cannot be isomorphic to (Q, \leq) in **Pre**.

Question 2

(a)

$$\begin{aligned}
 (g \circ f) \circ (f^{-1} \circ g^{-1}) &= (g \circ (f \circ f^{-1})) \circ g^{-1} && \text{(associativity)} \\
 &= (g \circ \text{id}_Y) \circ g^{-1} && \text{(definition of } f^{-1}) \\
 &= g \circ g^{-1} && \text{(unity)} \\
 &= \text{id}_Z && \text{(definition of } g^{-1})
 \end{aligned}$$

and a similar proof shows that $(f^{-1} \circ g^{-1}) \circ (g \circ f) = \text{id}_X$. So $g \circ f$ is an isomorphism with inverse $f^{-1} \circ g^{-1}$.

- (b) If f and $g \circ f$ have inverses $f^{-1} \in \mathbf{C}(Y, X)$ and $(g \circ f)^{-1} \in \mathbf{C}(Z, X)$, then consider $h \triangleq f \circ (g \circ f)^{-1} \in \mathbf{C}(Z, Y)$. We have

$$g \circ h = g \circ (f \circ (g \circ f)^{-1}) = (g \circ f) \circ (g \circ f)^{-1} = \text{id}_Z$$

and

$$h \circ g = (f \circ (g \circ f)^{-1}) \circ g = f \circ (g \circ f)^{-1} \circ g \circ f \circ f^{-1} = f \circ f^{-1} = \text{id}_Y$$

so that g is an isomorphism with inverse h .

- (c) No. In the category **Set** take $X = \{0\} = Z, Y = \{0, 1\}, f \in \mathbf{Set}(X, Y)$ to be the function $f 0 = 0$ and $g \in \mathbf{Set}(Y, Z)$ to be the function with constant value 0. Then neither f nor g are isomorphisms (since they are not bijections), but $g \circ f = \text{id}_X$ is one.

Question 3 The identity morphism $\text{id}_n \in \mathbf{Mat}(n, n)$ is the $n \times n$ matrix whose $(i, j)^{\text{th}}$ entry is 1 if $i = j$ and is 0 otherwise.

The morphism $\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \in \mathbf{Mat}(2, 2)$ is a non-identity isomorphism since $\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} = \text{id}_2$.

Two objects m and n are isomorphic in \mathbf{Mat} only if $m = n$. For if $M \in \mathbf{Mat}(m, n)$ is an isomorphism, then $M = \begin{pmatrix} \vec{v}_1 \\ \vdots \\ \vec{v}_m \end{pmatrix}$ consists of m rows that are linearly independent vectors $\vec{v}_1, \dots, \vec{v}_m \in \mathbb{R}^n$: for if $\lambda_1 \vec{v}_1 + \dots + \lambda_m \vec{v}_m = \vec{0} \in \mathbb{R}^n$, that is, $(\lambda_1 \dots \lambda_m) M = \vec{0}$, then applying the inverse of M we get $\vec{0} = \vec{0} M^{-1} = (\lambda_1 \dots \lambda_m) M M^{-1} = (\lambda_1 \dots \lambda_m)$. So since \mathbb{R}^n is a vector space of dimension n , we must have $m \leq n$. By a symmetric argument, $n \leq m$.

Question 4

- (a) If f is an isomorphism, its inverse f^{-1} is in particular a left inverse. If g is a left inverse for f , then for all $h, k \in \mathbf{C}(Z, X)$ we have $f \circ h = f \circ k \Rightarrow h = \text{id}_X \circ h = g \circ f \circ h = g \circ f \circ k = \text{id}_X \circ k = k$, so that f is a monomorphism.
- (b) If $h, k \in \mathbf{C}(W, X)$ satisfy $(g \circ f) \circ h = (g \circ f) \circ k$, then $f \circ h = f \circ k$ since g is a monomorphism; and then $h = k$ since f is a monomorphism.
- (c) If $h, k \in \mathbf{C}(W, X)$ satisfy $f \circ h = f \circ k$, then $(g \circ f) \circ h = (g \circ f) \circ k$ and since $g \circ f$ is a monomorphism, this implies $h = k$.
- (d) The monomorphisms in \mathbf{Set} are exactly the injective functions.

Proof. If $f \in \mathbf{Set}(X, Y)$ is injective, then for any $g, h \in \mathbf{Set}(Z, X)$, if $f \circ g = f \circ h$, then for all $z \in Z$ we have $f(gz) = f(hz)$, so $gz = hz$ (since f is injective); therefore g and h are equal functions.

Conversely, if $f \in \mathbf{Set}(X, Y)$ is a monomorphism, then for any $x, x' \in X$ let $\ulcorner x \urcorner, \ulcorner x' \urcorner \in \mathbf{Set}(1, X)$ be the functions mapping the unique element of $1 = \{0\}$ to x and x' respectively. If $fx = fx'$, then $f \circ \ulcorner x \urcorner = f \circ \ulcorner x' \urcorner \in \mathbf{Set}(1, Y)$. Since f is a monomorphism, this implies $\ulcorner x \urcorner = \ulcorner x' \urcorner$ and hence $x = \ulcorner x \urcorner 0 = \ulcorner x' \urcorner 0 = x'$. So f is injective. \square

Not every monomorphism in \mathbf{Set} is split. For example, consider the unique morphism in $\mathbf{Set}(\emptyset, 1)$ (where \emptyset denotes the empty set). This is injective (vacuously), but there is no function $1 \rightarrow \emptyset$ in \mathbf{Set} .

- (e) Consider $2 = \{0, 1\}$, $3 = \{0, 1, 2\}$ and the injective function $f \in \mathbf{Set}(2, 3)$ with $f0 = 0$ and $f1 = 1$. There are two different left inverses for f , one mapping 2 to 0 and the other mapping 2 to 1.
- (f) All morphisms in a pre-ordered set are monomorphisms, because there is at most one morphism between two objects. The only split monomorphisms are the isomorphisms (since if $f : p \rightarrow q$ and $g : q \rightarrow p$ then f and g are isomorphisms, since $g \circ f$ and

$f \circ g$ are necessarily equal to the unique morphism, namely the identity, on p and q respectively).

Question 5

- (a) Suppose $f \in \mathbf{Set}(X, Y)$ is surjective. If $g, h \in \mathbf{Set}(Y, Z)$ and $g \circ f = h \circ f$, then for all $y \in Y$, there exists $x \in X$ with $y = f x$ (since f is surjective) and hence $g y = g(f x) = (g \circ f) x = (h \circ f) x = h(f x) = h y$; therefore g and h are equal functions.

Conversely, suppose $f \in \mathbf{Set}(X, Y)$ is an epimorphism. For each $y \in Y$, consider the functions $g_y, h_y \in \mathbf{Set}(Y, \{0, 1\})$ that map y to 0 and to 1 respectively, and map all other elements of Y to 0. Since $g_y \neq h_y$ and f is an epimorphism, we must have $g_y \circ f \neq h_y \circ f$ and hence $g_y(f x) \neq h_y(f x)$, for some $x \in X$. Since g_y and h_y only take different values at y , it follows that $f x = y$. Therefore f is surjective.

- (b) Since the opposite category P^{op} of a pre-ordered set P is again a pre-ordered set, we can re-use the answer to question (4f): all the morphisms of P are epimorphisms.
- (c) In the pre-ordered set Q from question 1(b), the unique morphism $0 \rightarrow 1$ is both a monomorphism (by 4(f)) and an epimorphism (by 5(b)), but not an isomorphism, because there is no morphism from 1 to 0.

Question 6

- (a) $(1, 0, \text{id}_1)$ is a terminal object, where $1 = \{0\}$.
- (b) Consider the object $(\mathbb{N}, 0, \text{succ})$ where $\text{succ} \in \mathbf{Set}(\mathbb{N}, \mathbb{N})$ is the successor function, $\text{succ } n = n + 1$. This is initial in \mathbf{C} , because for any object (X, x_0, x_s) , the function $f : \mathbb{N} \rightarrow X$ recursively defined by

$$\begin{aligned} f 0 &= x_0 \\ f(n+1) &= x_s(f n) \end{aligned}$$

gives a morphism $f \in \mathbf{C}((\mathbb{N}, 0, \text{succ}), (X, x_0, x_s))$. It is the only such morphism, because if $g \in \mathbf{C}((\mathbb{N}, 0, \text{succ}), (X, x_0, x_s))$, then $g 0 = x_0$ and for all $n \in \mathbb{N}$, $g(n+1) = (g \circ \text{succ}) n = (x_s \circ g) n = x_s(g n)$; hence by induction on n , we have $(\forall n \in \mathbb{N}) g n = f n$.

Question 7

- (a) Each element $x \in X$ of a set $X \in \mathbf{Set}$ determines a point $\ulcorner x \urcorner : 1 \rightarrow X$ in \mathbf{Set} , namely the function mapping the unique element of $1 = \{0\}$ to x . The mapping $x \mapsto \ulcorner x \urcorner$ is injective, since $\ulcorner x \urcorner 0 = x$; furthermore for every $f \in \mathbf{Set}(X, Y)$, $f \circ \ulcorner x \urcorner = \ulcorner f x \urcorner$. So if $(\forall p \in \mathbf{Set}(1, X)) f \circ p = g \circ p$, then $(\forall x \in X) f x = g x$, that is, $f = g$.
- (b) \mathbf{Set}^{op} is not well-pointed. Note that the empty set \emptyset is a terminal object in \mathbf{Set}^{op} (because it is initial in \mathbf{Set}) and that $\mathbf{Set}^{\text{op}}(\emptyset, X) = \mathbf{Set}(X, \emptyset)$ is empty when $X \neq \emptyset$. Then for example $\text{id}_{\mathbb{N}} \neq \text{succ} \in \mathbf{Set}^{\text{op}}(\mathbb{N}, \mathbb{N})$, but $(\forall p \in \mathbf{Set}^{\text{op}}(\emptyset, \mathbb{N})) \text{id}_{\mathbb{N}} \circ p = \text{succ} \circ p$ is vacuously true.