

Power Rule In Derivatives

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$


$$\begin{aligned} f(x) &= x^2 \\ f'(x) &= 2x \end{aligned}$$

$$f(x) = x^2 + 3 \quad f(x) = x^3 \quad f(x) = x^2 + 2x + 1 \Rightarrow \text{Polynomial Equation}$$

$$f(x) = x^n \quad , \quad n \neq 0$$

$$\text{if } n=0 \quad f(x) = x^0 = 1 \Rightarrow \text{constant value}$$

$$\boxed{f'(x) = 0}$$



$$f(x) = x^n \quad \text{where } n \neq 0$$

$$\boxed{f'(x) = nx^{n-1}} \Rightarrow \text{Power Rule Polynomial Expression}$$

↓

$$\frac{\partial(f(x))}{\partial x} = nx^{n-1} \Rightarrow \boxed{\frac{\partial(x^n)}{\partial x} = nx^{n-1}}$$

$n \neq 0$

$$\begin{aligned} a+n &= 2 \quad \frac{\partial(x^3)}{\partial x} = 3 \cdot x^{3-1} &= 3x^2 \\ &= 3 \times (2)^2 = 3 \times 4 = 12 // . \end{aligned}$$

$$at x=5 \quad \frac{\partial(3x^2)}{\partial x} = 3 \cdot \frac{\partial(x^2)}{\partial x} = 3 \times 2x^{2-1} \Rightarrow 6x \quad \Rightarrow 6 \times 5 = 30.$$

$$\frac{\partial(y_n)}{\partial x} = \frac{\partial(x^{-1})}{\partial x} \Rightarrow -1 \cdot x^{-1-1} = -x^{-2} \Rightarrow \boxed{-\frac{1}{x^2}}$$

Assignment

$$f(x) = x^8 \Rightarrow f'(x) ?$$

$$f(x) = x^{-1} \Rightarrow f'(x) \quad at \quad n = -1,$$

④ Derivative Rules: Constant, Sum, difference And Constant Multiple

$$\frac{\partial}{\partial x}(x^n) = nx^{n-1}, \quad n \neq 0 \quad \{ \text{Power Rule} \} \Rightarrow \text{Polynomials}$$

$$\frac{\partial}{\partial x}(x^0) = \frac{\partial}{\partial x}[1] = 0 \quad \Rightarrow \text{Derivative of a constant is 0.}$$

$$\frac{\partial}{\partial x}[c] = 0 \quad \begin{matrix} \downarrow \\ \text{Constant} \end{matrix}$$

$$\frac{\partial}{\partial x}[cf(x)] = c \frac{\partial(f(x))}{\partial x} = c f'(x)$$

$$\frac{\partial(3x^4)}{\partial x} = 3 \cdot \frac{\partial(x^4)}{\partial x} = 3 \times 4x^{4-1} = 3 \times 4x^3 = 12x^3$$

$$at \quad x=2 \quad \frac{\partial(3x^4)}{\partial x} = 12 \times 2^3 = 12 \times 8 = 96 \quad \underline{\underline{}}$$

Assignment

$$\frac{\partial [A f(x)]}{\partial x} = \frac{\partial [2x^5]}{\partial x}$$

④ Sum of 2 function $f(x), g(x)$

$$\frac{\partial [f(x) + g(x)]}{\partial x} \Rightarrow \frac{\partial (f(x))}{\partial x} + \frac{\partial (g(x))}{\partial x}$$

$$\begin{aligned} \frac{\partial [x^4 + x^{-2}]}{\partial x} &= \frac{\partial (x^4)}{\partial x} + \frac{\partial (x^{-2})}{\partial x} \\ &= 4x^3 + (-2x^{-3}) \end{aligned}$$


$$\boxed{\frac{\partial [x^4 + x^{-2}]}{\partial x} = 4x^3 - 2x^{-3}}$$

Assignment

$$\frac{\partial (x^2 + 2)}{\partial x} \quad \text{Answer } f'(x) = \underline{\underline{2x}}.$$

$$\begin{aligned} ⑤ \quad \frac{\partial (4x^3 - 6x^2 + 2x + 100)}{\partial x} &\Rightarrow 12x^2 - 12x + 2 + 0 \\ &\quad \uparrow \\ &\Rightarrow \boxed{12x^2 - 12x + 2} \end{aligned}$$

Tangent of polynomials



$$\rightarrow f(x) = x^3 - 6x^2 + x - 7$$

$$y = f(1) = 1 - 6 + 1 - 7 = -11$$

$$f'(x) = \frac{\partial (x^3 - 6x^2 + x - 7)}{\partial x}$$

$$= 3x^2 - 12x + 1 - 0$$

$$f'(x) = 3x^2 - 12x + 1$$

$$f'(1) = 3 \times (1) - 12 + 1$$

$$= 3 - 12 + 1 = \boxed{-8} \Rightarrow \text{slope}.$$

$$y = mx + c$$

$$y = -8x + c$$

$$\text{for } x = 1$$

$$-11 = -8 + c$$

$$\boxed{c = -11 + 8} \Rightarrow \boxed{c = -3}$$

$$y = mx + c$$

$$\boxed{y = -8x - 3}$$


equation of
line for tangent

$m = -8 \Rightarrow \text{slope}$


$c = -3 \Rightarrow \text{intercept}$

IQ Dataset

IQ	Score
100	98
90	97




$\left\{ \text{Optimization} \right.$



\Rightarrow Chain Rule

(f) Derivatives for Trigonometric, logarithmic and Exponential function

Trigonometric function



$$f(x) = \sin x$$

$$f'(\sin x) = \cos x$$

$$f(x) = \cos x$$

$$f'(\cos x) = -\sin x$$

Logarithmic function

$$f(x) = \ln(x) \text{ then}$$

$$\boxed{f'(x) = \frac{1}{x}}$$

Exponential Function

$$f(x) = e^x$$

$$\boxed{f'(x) = e^x}$$

Constant

$$\text{if } f(x) = c$$

then

$$\boxed{f'(x) = 0}$$

Power Rule

$$\text{if } f(x) = x^n$$

$$\boxed{f'(x) = nx^{n-1}}$$