

Bifurcations of Hamiltonian Periodic Orbits

Dongho Lee

December 12th, 2025

SNU, QSMS

Presented in Symplectic Learning Seminar, SNU



Download slides

Well-known Classical Fact from 70s

There are 8 types of *generic* bifurcation of Hamiltonian periodic orbits.

Contents

- Dynamical Systems and Bifurcations
- Poincaré Section and Stability of Orbits
- Bifurcation and Floer Homology
- Ingredients : Classical works of Lyapunov and Meyer
- Classification of Generic Bifurcations

Main references : [Mey70], [AM78]

Dynamical Systems and Bifurcations

Toy Example : Critical Points of a Function




Figure 1: Graphs of $f_t(x) = x(x^2 + t)(x - 1)$ for varying t .

- $t < 0$, two minima (●) and one maximum. (●)
- $t = 0$, ● and ● meets at the inflection point. (●)
- $t > 0$, ● vanishes and one minimum survives. (●)

Idea. Similar thing happens for periodic orbits.

Example : Van der Pol Oscillator




Figure 2: Phase portraits of Van der Pol oscillator for varying μ .

Van der Pol oscillator is given by $\dot{x} = \mu(1 - y^2)x - y$, $\dot{y} = x$.

0 is a fixed point, and its property changes as μ varies.

Also, at $\mu = 0$, a family of periodic orbits emerge.

According to [Sma67], a **dynamical system** consists of following data.

- A space W (we use smooth manifolds)
- A group action $G \rightarrow \text{End}(W)$ (we use diffeomorphisms)

If $G = \mathbb{Z} = \langle F \rangle$, we say the system is **discrete**.

Interesting objects

- **Fixed point** $x \in W$ such that $F(x) = x$.
- **Periodic point** $x \in W$ such that $F^N(x) = x$ for some N .
- **Invariant subspace** $S \subset W$ such that $F(S) \subset S$
- Non-wandering points, recurrent points, etc.

Continuous Dynamical System

If $G = \mathbb{R} = \{F_t\}$, we say the system is **continuous**.

A vector field $\frac{d}{dt}F_t|_{t=0} = X$ consists equivalent information with F_t , which is called the infinitesimal generator of F_t .

Example. The system determined by an ODE $\dot{x} = X(x)$.

Interesting objects

- **Critical points** $x \in W$ such that $F_t(x) = x$ for any t . (or $X_x = 0$.)
- **Periodic orbits** $x \in W$ such that $F_\tau(x) = x$ for some $\tau > 0$.
- **Invariant subsets** $S \subset W$ such that $F_t(S) \subset S$ for any t (or $t > 0$).

Hamiltonian Dynamical System

Let (W, ω) be a symplectic manifold, $H : W \rightarrow \mathbb{R}$.

H defines a **Hamiltonian vector field** X_H via relation

$$i_{X_H}\omega = \omega(X_H, -) = -dH.$$

We call (W, ω, H) a **Hamiltonian dynamical system**.

- X_H preserves ω , i.e. $\mathcal{L}_{X_H}\omega = 0$.
- X_H preserves H , i.e. X_H flows on $H^{-1}(c)$ for each c .

The first example is the geodesic flow on a Riemannian manifold (M, g) ,

$$H_g(q, p) = \frac{|p|_g^2}{2}$$

which is defined on T^*M .

Example : Classical Mechanics

Consider the mechanical system in $T^*\mathbb{R}^n$ with energy

$$H(q, p) = \frac{|p|^2}{2} + V(q).$$

Corresponding Hamiltonian vector field is

$$X_H = p \frac{\partial}{\partial q} - \nabla V \frac{\partial}{\partial p},$$

which can be translated into

$$\dot{q} = p, \quad \dot{p} = \ddot{q} = -\nabla V.$$

This is Newton's 2nd law, which governs the classical mechanics.

Structural Stability

Two vector fields X, Y on W are **topologically conjugate** if there exists a homeomorphism $F : W \rightarrow W$ which carries the oriented orbits of X to the ones of Y .

If X has a neighborhood $\mathcal{U} \subset \mathcal{X}(W)$ such that if $Y \in \mathcal{U}$ then X and Y are topologically conjugate, we call X is **structurally stable**.

Idea. A small perturbation of a structurally stable vector field cannot change the dynamical system essentially.

Bifurcation

Consider a family of vector fields X_t on W , parametrized by a set I .

A point $s \in I$ is called

- a **robust point** if there exists a neighborhood U of s in I such that if $t \in U$, X_s and X_t are topologically conjugate.
- a **bifurcation point** if not.

We will only consider 1-dimensional bifurcation, i.e. I is 1-manifold.

Idea. If we can find *generic* properties of vector fields, we can talk about *generic* bifurcations.

Bifurcation




Figure 3: Very schematic picture of a bifurcation.

Poincaré Section and Stability of Orbits

Theorem (Poincaré Section and Return Map)

Let γ be a periodic orbit of X in W and $x \in \gamma$.

1. There exists a codimension 1 submanifold $S \subset W$ such that $x \in S$ and is transversal to X .
2. Define a map $\Psi : S \rightarrow W$ by

$$\Psi(y) = Fl_{\tau(y)}^X(y)$$

where $\tau(y) = \min\{t > 0 : Fl_t^X(y) \in S\}$, then this is well-defined and unique up to conjugation.

We call S **Poincaré section**, and Ψ **return map** or **Poincaré map**.

Poincaré Section




Figure 4: A Poincaré section and the return map

Hamiltonian Version of Poincaré Section

Proposition

Let $X = X_H$ be a Hamiltonian vector field and γ be a periodic orbit with regular energy c_0 . There exists a Poincaré section Y such that

- 1. $Y \subset W$ is a contact submanifold.*
- 2. $S_c = Y \cap H^{-1}(c)$ for c close to c_0 is a symplectic submanifold.*
- 3. The return map on S_c is a symplectomorphism.*

$2n$ -dimensional W	$\Leftrightarrow$	$(2n - 2)$ -dimensional S
Flow of vector field X_H	$\Leftrightarrow$	Return map Ψ
1-periodic orbit of X_H on W	$\Leftrightarrow$	a fixed point of Ψ on S
k -periodic orbit of X_H	$\Leftrightarrow$	k -family of k -periodic points of Ψ .

Aside : A Standard Picture the Poincaré Section




Figure 5: [HH64] Poincaré sections of the Hénon-Heiles system

1. Take an initial condition x_0 and compute its orbit γ .
2. Mark the intersections x_i 's of γ and S .
3. Draw a curve (if possible) between x_i 's.
4. Repeat for different initial conditions.

Characteristic Exponents of Critical Points

Let x be a critical point of X_H on W .

The **characteristic exponents** of x are the eigenvalues of $dX_H(p)$.

In symplectic case, the exponents come with pairs, $\pm\alpha, \pm\bar{\alpha} \in CE(x)$.

Let $\{\alpha_j\}$ be characteristic exponents of x .

Let's denote $\lambda_j = \exp(\alpha_j)$ for the future consistency.

We say λ_j are **characteristic multiplier** of x .

- If $\lambda_j \notin S^1$ (or $\alpha_j \notin i\mathbb{R}$) for all j , call x **hyperbolic**.
- If $\lambda_j \in S^1$ (or $\alpha_j \in i\mathbb{R}$) for all j and $\lambda_j \neq \pm 1$, call x **elliptic**.

Elliptic and Hyperbolic Critical Points

Let α be a characteristic exponent of x , $\lambda = e^\alpha$ and v be eigenvector.

Consider the linearized flow in direction v at a critical points.

- If $|\lambda| > 1$, the flow is expanding and tends away from x . (**unstable**)
- If $|\lambda| < 1$, the flow is contracting and tends closer to x . (**stable**)
- If $|\lambda| = 1$, the flow rotates around the x . (**center**)

In Hamiltonian case, the stable and unstable dimensions are the same.





Figure 6: Elliptic and hyperbolic fixed points

Rigidity of Critical Points of Symplectomorphism



Consider the elliptic critical point x with $CE(x) = \{\pm i\alpha, \pm i\beta\}$.

Under perturbation, α and β change continuously, but locked in $i\mathbb{R}$ unless $\alpha = \beta$ at some point. So it cannot suddenly become hyperbolic.

Characteristic Multipliers of Periodic Orbits

Let γ be a periodic orbit, S be a Poincaré section and $\gamma \cap S = \{x\}$.

The **characteristic multipliers** of γ are the eigenvalues of $d\Psi_x$.

The multipliers come with pairs, $\lambda^{\pm 1}$, $\bar{\lambda}^{\pm 1}$.


If $\dim S = 2$, which is of our interest, it simplifies by $\lambda \in S^1 \cup \mathbb{R} \setminus \{0\}$.

- If $\lambda_\gamma \in \mathbb{R} \setminus \{\pm 1\}$, call γ **hyperbolic**.

We say γ is **positive** or **negative** depending on the sign of λ_γ .

- If $\lambda_\gamma \in S^1 \setminus \{\pm 1\}$, call γ **elliptic**.

Characteristic Multipliers



As in the critical points, the transition between elliptic and hyperbolic periodic orbits can be done only through ± 1 .

Proposition ([BL34] Birkhoff-Lewis)

Let $\Psi : U \subset \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a symplectomorphism with $\Psi(0) = 0$.

If Ψ is an elementary twist mapping, which means that 0 is a generic elliptic fixed point (See Appendix 1), then in every neighborhood of 0,

- 1. there exists an invariant circle in U without periodic points.*
- 2. for any integer k , there exists a k -periodic point in U .*

It means the elliptic periodic orbits can be detected from nearby orbits, while the hyperbolic ones cannot be.

In this sense, the hyperbolic periodic orbits are *qualitatively invisible*.

Aside : Vague Attractor of Kolmogorov




Figure 9: [EI 03] VAK

- We have invariant tori (of positive measure) around γ .
- Between tori, there are periodic orbits. (red dots)

We say such configuration a **vague attractor of Kolmogorov**, or **VAK**.

Bifurcation and Floer Homology

Conley-Zehnder Indices

Let γ be a periodic Hamiltonian orbit of period τ , S be a Poincaré section, $\gamma \cap S = \{x\}$ and Ψ be the return map.

γ is **non-degenerate** if $\det(d\Psi|_x - \text{Id}) \neq 0$, i.e. the linearized return map doesn't have 1 as an eigenvalue.

The **Conley-Zehnder index** is an integer assigned to γ , defined by

$$\gamma_{CZ}(\gamma) = \frac{1}{2}\text{Sign}Q_0 + \sum \text{Sign}Q_t + \frac{1}{2}\text{Sign}Q_\tau \in \mathbb{Z}$$

where Q_t is a quadratic form $Q_t(v, v) = \omega(v, \dot{\Psi}(t)v)$ on $\ker(\Psi(t) - \text{Id})$, defined for t such that $\det(\Psi(t) - \text{Id}) = 0$, and Sign is the signature of a quadratic form.

$H : W \rightarrow \mathbb{R}$ be a Hamiltonian, c be a regular value.

Assume : $H^{-1}(c)$ is a contact type hypersurface, consists only of non-degenerate periodic orbits.

$\Rightarrow$ We can define the **symplectic homology** $SH^{S^1,+}(W_c)$,

- generated by periodic Hamiltonian orbits on $H^{-1}(c)$,
- graded by Conley-Zehnder indices,
- Differentials are given by Floer cylinders.

Relation between Symplectic Homology and Bifurcation

If $\text{CritVal}(H) \cap [c_1, c_2] = \emptyset$, then $SH^{S^1,+}(W_{c_1}) \simeq SH^{S^1,+}(W_{c_2})$.

$\Rightarrow$ CZ-indices of *all* periodic orbits are topologically controlled.

Assume that we're away from the critical energies.

- Assume that there exists a family of periodic orbits γ_c parametrized by the energy c . (an *orbit cylinder*)
- If γ_c is non-degenerate for $c_1 \leq c \leq c_2$, then $\mu_{CZ}(\gamma_{c_1}) = \mu_{CZ}(\gamma_{c_2})$.
- Once γ_c degenerates, $\mu_{CZ}(\gamma_c)$ might change, or the family might just vanish away.
- Since $SH^{S^1,+}(W_c)$ must be preserved, other orbits must take the charge of the changed index by being born, vanishing, or changing its own index.

Indeed, we only consider the **local Floer homology** [CFHW96].

- Compute the Floer homology of a neighborhood of an orbit.
- This can be done by finding the indices of nearby orbits and use Morse-Bott spectral sequence.
- The *local invariance* of the local Floer homology tells us that the result are same before and after the bifurcation.
- From this, we can deduce the index of the new orbits, or find out if there is a missing orbit.

Example : Hill's Lunar Problem




Figure 10: [Ayd23] Bifurcation diagram of Hill's lunar problem



Ingredients : Classical Works

Lyapunov's Theorem

Let x be a critical point of X_H . x is called $\mathcal{H}$ -**elementary** if

1. 0 is not a characteristic exponent.
2. If $\alpha \in i\mathbb{R}$ is a characteristic exponent, α has multiplicity 1.
3. If $\alpha, \beta \in i\mathbb{R}_{>0}$ are characteristic exponents, they are $\mathbb{Z}$ -linearly independent.

Theorem ([Buc70])

For generic Hamiltonian H , every critical point of X_H is $\mathcal{H}$ -elementary.

Lyapunov's Theorem

Theorem (Lyapunov)

Let x be an $\mathcal{H}$ -elementary critical point of X_H . If $\pm i\beta$ is a characteristic exponent for $\beta \in \mathbb{R}$, there exists a 2-dimensional submanifold C_β such that

- 1. $x \in C_\beta$.*
- 2. C_β is X_H -invariant submanifold.*
- 3. $T_x C_\beta$ is an eigenspace corresponds to $\pm i\beta$.*
- 4. C_β is a union of periodic orbits γ_r which converges to x as $r \rightarrow 0$.*
- 5. The period of γ_r converges to $2\pi/\beta$ as $r \rightarrow 0$.*

Simply, there exist families of periodic orbits emerging from x for each elliptic exponent.

Lyapunov Orbits

Example. Lyapunov orbits of the restricted three-body problem, emerges from the Lagrange points.




Figure 11: [ZWZZ20] Lyapunov orbits of RTBP, associated to L_1 and L_2 .

These Lagrange points have only one pair of elliptic exponents, so there is only one family of orbits.

Setting for Periodic Orbits

We consider 4-dimensional symplectic manifold W and a periodic orbit γ .

Let S be a Poincaré section, intersecting with γ at x .

Locally, 1-parameter family of flows can be projected to the same S .

$\Rightarrow$ We consider 1-parameter family of symplectomorphisms φ_s on S with a fixed point x , and see their local behavior.


Precisely, we have

1. A family of symplectomorphisms $\varphi : S \times I \rightarrow S$.
2. A point $x = x_s$ which is a fixed point of φ_s for $s \in I$.
3. For simplicity, we say (x, s) is a fixed point of φ .

Orbit Cylinder

Lemma (Orbit Cylinder)

If $\lambda_x \neq 1$, there exists neighborhoods $x \in U \subset M$ and $s \in V \subset S^1$ and a smooth map $P : V \rightarrow U$ such that $P(s) = x$ and $\{(P(t), t) : t \in V\}$ consists of fixed points.



In dimension 4, this can be interpreted as a family of orbits.

Note. This theorem holds for arbitrary dimension.

Two Types of Orbit Cylinders




Figure 12: Two possible orbit cylinders

The *height parameter* of the orbit cylinder can be chosen as the energy.

There are two possibilities; the orbit cylinder might be transversal or tangent to the energy hyperplane.

Generating Function

Now we only focus on the cases $\lambda_x^k = 1$ for some k .

We will see $\lambda_x = 1$ case in detail, and other cases briefly.

Locally, we take a fixed point at $x = (0, 0)$ and denote φ_s as

$$Q = Q(q, p), \quad P = P(q, p).$$

[Poi87] observed that

$$\Omega = (P - p)d(Q + q) - (Q - q)d(P + p)$$

is closed, so there exists a function G such that $dG = \Omega$.

We call G a **generating function**.

Generating Function

Assume that $\lambda \neq -1$, then under new coordinates

$$\xi = Q + q, \quad \eta = P + p,$$

we have

$$\frac{\partial G}{\partial \xi} = P - p, \quad \frac{\partial G}{\partial \eta} = q - Q,$$

and the fixed point of (Q, P) corresponds to the critical point of $G(\xi, \eta)$.

Generating Function

Let the 2nd-order part of G be

$$G(\xi, \eta) = \frac{1}{2}(\alpha\xi^2 + 2\beta\xi\eta + \gamma\eta^2),$$

and $4\Delta = \alpha\gamma - \beta^2 \neq -1$. The linear part of (Q, P) is given by

$$\begin{pmatrix} Q \\ P \end{pmatrix} = \frac{1}{1 + 4\Delta} \begin{pmatrix} (1 - \beta)^2 - \alpha\gamma & -2\gamma \\ 2\alpha & (1 + \beta)^2 - \alpha\gamma \end{pmatrix} \begin{pmatrix} q \\ p \end{pmatrix}.$$

The trace of given matrix is $T = 2(1 - 4\Delta)(1 + 4\Delta)^{-1}$.

- Min/Max of $G \Leftrightarrow \Delta > 0 \Leftrightarrow |T| < 2 \Leftrightarrow$ Elliptic fixed point
- Saddle of $G \Leftrightarrow \Delta < 0 \Leftrightarrow |T| > 2 \Leftrightarrow$ Hyperbolic fixed point

Meyer's Theorem - Extremal Fixed Point

We call (x, s) is **extremal fixed point** if, for generating function $G = G(\xi, \eta, t)$, at $(0, 0, 0)$,

$$G = G_\xi = G_\eta = G_{\eta\eta} = G_{\eta\xi} = 0, \\ G_{\xi\xi} \neq 0, \quad G_{\eta t} \neq 0, \quad G_{\eta\eta\eta} \neq 0.$$

Note. This is the case $\lambda_x = 1$.

Theorem ([Mey70])

Let (x, s) be an extremal fixed point. Then,

- 1. There exists a family of fixed points x_t .*
- 2. x divides x_t into two arcs: one entirely consists of hyperbolic fixed points and the other consists of elliptic fixed points.*
- 3. In any coordinate system, s is a nondegenerate maximum or minimum of the family.*

Outline of Proof

We have an equation $G_\eta = G_\xi = 0$ at $(0, 0, 0)$, and the Jacobian is

$$\begin{pmatrix} 0 & 0 & G_{\eta t} \\ 0 & G_{\xi\xi} & G_{t\xi} \end{pmatrix}.$$

This has rank 2, so we have solution $(\xi, \eta, t) = (\xi(s), s, t(s))$ for the equation, i.e.

$$G_\xi(\xi(s), s, t(s)) = G_\eta(\xi(s), s, t(s)) = 0.$$

From this, we can deduce that $t'(0) = 0$ and $t''(0) \neq 0$, and

$$\begin{pmatrix} G_{\xi\xi} & G_{\eta\xi} \\ G_{\xi\eta} & G_{\eta\eta} \end{pmatrix} (\xi(s), s, t(s)) = \begin{pmatrix} G_{\xi\xi}(0) & 0 \\ 0 & sG_{\eta\eta}(0) \end{pmatrix} + (\text{h.o.t})$$

Illustration of Extremal Fixed Point




Figure 13: Meyer's theorem for extremal fixed point

Each point is a fixed point, and the vertical axis is the energy.

This is the tangent case of the orbit cylinder.

Meyer's Theorem - Transitional Fixed Point

We call (x, s) is a **transitional fixed point** if $\lambda_x = -1$ and satisfies some non-degeneracy condition. (See Appendix 2)

Theorem ([Mey70])

Let (x, s) be a transitional fixed point and x_t be an orbit cylinder in $U \times V \subset M \times S^1$. Then we can choose V small enough so that $V \setminus \{s\} = V_H \cup V_E$, where x_s is hyperbolic (elliptic) if $s \in V_H$ (V_E). Moreover, either one of the following holds.

- 1. There exists functions $x_1^E, x_2^E : V_H \rightarrow U$ such that $(x_i^H(t), t)$ is elliptic periodic points of period 2 and $x_i^H(t) \rightarrow x$ as $t \rightarrow s$,*
- 2. There exists functions $x_1^H, x_2^H : V_E \rightarrow U$ such that $(x_i^E(t), t)$ is hyperbolic periodic points of period 2 and $x_i(t) \rightarrow x$ as $t \rightarrow s$.*

We call these **an arc of nearby hyperbolic/elliptic periodic points**.

Meyer's Theorem - Bifurcation Points

Let $\lambda_x = \exp(\pm 2\pi i l/k)$, $(l, k) = 1$, $k \geq 3$ and $0 < l < k$.

We call (x, s) is a **k -bifurcation point** if it satisfies some non-degeneracy condition. (See Appendix 3.)

Theorem ([Mey70])

Let (x, s) be k -bifurcation point. Then for small orbit cylinder $U \times V$,

- 1. ($k = 3$) There exists three arcs of nearby hyperbolic periodic points of period 3, parametrized by $V \setminus \{s\}$, converging to x .*
- 2. ($k \geq 5$) There exists k arcs of nearby hyperbolic periodic points and k arcs of nearby elliptic periodic points of period k , parametrized by $V_+ \subset V \setminus \{s\}$, converging to x .*
- 3. ($k = 4$) One of ($k = 3$) or ($k \geq 5$)-phenomena occurs.*

Let $\mathcal{G}$ be the set of 1-parameter family of symplectomorphisms on M such that

1. Each periodic point is elliptic, hyperbolic, extremal or transitional.
2. If $\lambda_x = \exp(\pm 2\pi i l/k)$ for $k \geq 3$, then (x, s) is k -bifurcation point.

Theorem ([Mey70])

$\mathcal{G}$ is a residual set in the space of 1-parameter family of symplectomorphisms.

$\Rightarrow$ Now we know the *generic* bifurcation behavior.

Classification of Generic Bifurcations


How to classify bifurcation

Let $\dim W = 4$, so the Poincaré section is 2-dimensional.

- For critical points, characteristic exponents are $\{\pm\alpha, \pm\bar{\alpha}\}$ in general.
- When there is $\alpha \in i\mathbb{R}$, Lyapunov theorem can be applied.
- If $CE(x_s) = \{\pm i\alpha_t, \pm i\beta_t\}$ and $l\alpha_s = k\beta_s$, for some s , the Lyapunov assumption ($\mathcal{H}$ -elementary) is violated.
- For periodic orbits, if $\lambda_\gamma \neq 1$, there is an orbit cylinder.
- If $\lambda_\gamma^k = 1$, k -th cover of the orbit bifurcates.
- Using Poincaré section, a set of k periodic points of period k corresponds to a periodic orbit of period k .
- The nearby hyperbolic/elliptic arc on the Poincaré section corresponds to the orbit cylinder of hyperbolic/elliptic orbits.

Bifurcation Diagram

We'll see this kind of diagrams.



- Vertical axis is the energy.
- $\bullet$ is the bifurcation point.
- $\odot$ is the critical point of H .
- Each line is a family of periodic points of the return map in Poincaré section.
- Thick line is a family of fixed points, or 1-periodic orbits.
- Thin line is a family of k -periodic orbit, where the period is written aside the line. (2 here.)
- —, $\odot$ are elliptic, —, $\odot$ are hyperbolic, and $\odot$ is saddle.

List of Generic Bifurcations

No.	Name	Case	Tool
I	Burst / Reincarnation	x with $i\alpha \in CE(x)$	Lyapunov theorem
II	Creation / Annihilation	γ with $\lambda_\gamma = 1$	Extremal f.p.
III	Subtle Doubling / Halving	$\lambda_\gamma = -1$	Transitional f.p.
IV	Materialization / Murder	$\lambda_\gamma = -1$	Transitional f.p.
V	Phantom Kiss	$\lambda_\gamma^3 = 1$ $\lambda_\gamma^4 = 1$	3-bifurcation point 4-bifurcation point
VI	Emission / Absorption	$\lambda_\gamma^k = 1$ ($k \geq 4$)	k -bifurcation point
VII	Bubble / Liberation	Perturbation of critical point	TBA
VIII	Resonance	Perturbation of critical point	TBA

I. Burst / Reincarnation

Target. Critical point x with pure imaginary characteristic exponents.

Case 1. $\{\pm i\alpha, \pm r\}$ with $\alpha, r \in \mathbb{R}$. (saddle-center)



- By Lyapunov's theorem, we have a family of orbits near x
- The characteristic multipliers of the Lyapunov orbits near x is eventually real, so the orbits are hyperbolic.
- This hyperbolic family cannot be observed in the phase portrait, so we call this a **phantom burst**.

Figure 14: Phantom burst

I. Burst / Reincarnation

Case 2. $\{\pm i\alpha, \pm i\beta\}$ with $\mathbb{Q}$ -independent $\alpha, \beta \in \mathbb{R}$. (pure center)



Figure 15: Stable burst



Figure 16: Reincarnation

- By Lyapunov's theorem, we have two families near $\odot$.
- The characteristic multipliers are unimodular, so both orbits are elliptic.
- If two orbit cylinders are on the same side respect to the energy hyperplane, we observe two families of orbits burst from the critical point. (**Stable burst**)
- Otherwise, we observe a family of orbit absorbed into the critical point and burst out again. (**Reincarnation**)

II. Creation / Annihilation

Target. Periodic orbit such that $\lambda_\gamma = 1$.




Figure 17: Creation and annihilation
(Also called birth-death)

- The vector field develops an unstable periodic orbit $\bullet$. It's a closing of a recurrent orbit, called **Pugh catastrophe**.
- $\bullet$ splits into a hyperbolic orbit and an elliptic orbit.
- In the phase portrait, we observe an elliptic orbit suddenly appeared. (**Creation**)
- If we read this in reversed time, we can observe a vanishing pair of orbits. (**Annihilation**)

III. Subtle Doubling / Subtle Halving

Target. Periodic orbit such that $\lambda_\gamma = -1$.





Figure 18: Subtle doubling and halving
(Also called period doubling)

- The elliptic orbit cylinder becomes hyperbolic at $\bullet$.
- An elliptic orbit with period 2 emerges.
- In phase portrait, we observe that the period of — is doubled. (**Subtle doubling**)
- In the opposite direction, the period is halved. (**Subtle halving**)

IV. Materialization / Murder

Target. Periodic orbit such that $\lambda_\gamma = -1$.




- Similar to subtle doubling, but with an interchange of elliptic and hyperbolic.
- We observe that — suddenly appeared. (**Materialization**)
- In the opposite direction, — suddenly disappears. (**Murder**)

Figure 19: Materialization and murder

V. Phantom Kiss

Target. $\lambda_\gamma = \exp(2\pi i l/k)$, $(l, k) = 1$ and $k = 3$ or 4 .




- A hyperbolic orbit of period 3 emerges in both directions of $\bullet$.
- We only observe *a moment of instability* of — .
- The family of hyperbolic orbit only experiences a sudden absorption to the elliptic orbit. **(Phantom kiss)**

Figure 20: Phantom kiss
(Also called touch-and-go)

VI. Emission / Absorption

Target. $\lambda_\gamma = \exp(2\pi i l/k)$, $(l, k) = 1$ and $k \geq 4$.



- A hyperbolic and an elliptic orbit of period k both emerges at $\bullet$. (**Emission**)
- An elliptic orbit of period k is observed.
- In the opposite direction, k -periodic orbits are absorbed into the original periodic orbit. (**Absorption**)

Figure 21: Emission and absorption

VII. Bubble / Liberation

Target. Fixed point 0 of parametrized Hamiltonian H_μ .

Assume that 0 is elliptic if $\mu < 0$ and hyperbolic if $\mu > 0$, i.e.

- $\mu < 0 : CE(x) = \{\pm i\alpha, \pm i\beta\}$ for $\alpha, \beta \in \mathbb{R}_{>0}$, $\alpha \neq \beta$.
- $\mu = 0 : \alpha = \beta$
- $\mu > 0 : CE(x) = \{\pm z, \pm \bar{z}\}$ where $z \notin S^1$.

If $\mu < 0$, the stable burst occurs, while nothing happens if $\mu > 0$.

Question. What happens to two elliptic orbit cylinders?

[MS71] showed that there are only two possibilities in general.

VII. Bubble / Liberation




Figure 22: Bubble

The first possibility is a **bubble**: the shrink of two orbit cylinders into a fixed point.

VII. Bubble / Liberation




Figure 23: Liberation

The second possibility is a **liberation**: two orbit cylinders patch together. This phenomenon actually happens in CRTBP at L_4 .

VIII. Resonance

Target. Fixed point 0 of parametrized Hamiltonian H_μ .

For characteristic exponents α_1 and α_2 , if there exist coprime integers k, l such that

$$k\alpha_1 = l\alpha_2,$$

then $\mathcal{H}$ -elementary assumption of Lyapunov theorem is violated.

This is called **resonance**, and there are many possible different phenomenon occurs depending on (k, l) .

[SS73] gives the classification of generic resonance.

The situation is quite similar to the case of the bubble: ‘bridges’ between two orbit cylinders are collapsed into a point.

A Life of Orbit




Figure 24: A life of an orbit

A periodic orbit can be born and die at critical point or other orbit.

For varying energy, it experiences various bifurcations, make birth of another orbits, and become from hyperbolic to elliptic or vice versa.

Closing

Higer Dimension?

Higher dimensional W

- [Mey70] said, there would be no essentially new generic behavior.
- The bifurcation would happen when one multiplier becomes a root of unity, and *generically* we don't expect the other multiplier simultaneously becomes another root of unity.

Higher dimensional I (the parameter of bifurcation)

- [Mey70] said, there are too many essentially different forms of bifurcation if we increase the dimension of parameters.
- The theory would become very wild, and there's no systemetic result as I know.

Genericity and Symmetry

Generically, a given Hamiltonian system is generic. (It's trivial.)

But the genericity (even though it's well-defined) cannot be checked because generically there are infinitely many periodic orbits.

Moreover, most of interesting systems are *not* generic, because of their symmetries in nature.

- The circular restricted three-body problem has $\mathbb{Z}_2 \times \mathbb{Z}_2$ -symmetry.
- [Ayd23] The linear symmetries of Hill's lunar problem, a limit case of of CRTBP, is given by $\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2$.
- [AFFvK13], [Lee25] There are much more linear symmetries in the rotating Kepler problem, so there is S^1 -family bifurcation in the planar case, and S^3 -family bifurcation in the spatial case.

Many works were done for symmetric Hamiltonians, including [GSM87].

Pitchfork Bifurcation




Figure 25: Pitchfork bifurcations

- The orbit changes hyperbolicity and ellipticity, and **two** orbits with **period 1** are born.
- The reason is $\mathbb{Z}_2$ -symmetry; if there is a new orbit one side, then there must be another one on the other side.
- This type of bifurcation, which is not *generic*, is observed in Hill's lunar problem, CRTBP, and many other systems with symmetry.

Further Questions

Investigating bifurcation by Conley-Zehnder index:

- Floer theory provides systematic way to study bifurcation.
- [AFvK⁺] studied Hill's lunar problem with CZ index.
- We might apply the method to three-body problem and other interesting systems to investigate periodic orbits and bifurcations.

General relation between the change of index and the bifurcation type:

- If we know the bifurcation type, what can we say about the index?
- If we detect index change, what can we say about bifurcation type?
- [FKM23] showed the invariance of χ_{SFT} explicitly in each case.

The End



Figure 26: Thank you for your attention!

Appendix 1. Twist Condition

$\Psi : 0 \in U \subset \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is α -**twist map** if 0 is a fixed point of Ψ with characteristic exponents $\pm i\alpha$ for $\alpha \in \mathbb{R}$.

If we can write

$$\Psi(z) = e^{i\alpha + \beta|z|}z + R_4(z),$$

where $R_4(z) \leq K|z|^4$ for some $K > 0$, we call such Ψ is written in (α, β) -**normal form**.

Theorem (Birkhoff-Sternberg-Moser Normal Form)

Let Ψ be α -twist mapping with $\alpha \neq 0, k\pi/2$ or $2k\pi/3$. Then there exists a chart such that Ψ can be written in (α, β) -normal form and the sign of β is independent of the change of the chart.

An α -twist map is an **elementary twist map** if $\alpha \neq 0, k\pi/2$, or $2k\pi/3$ and $\beta \neq 0$, which was mentioned in the Birkhoff-Lewis theorem.

Appendix 2. Transitional Fixed Point

A fixed point (x, s) is **transitional** if locally

$$\begin{pmatrix} Q \\ P \end{pmatrix} = \begin{pmatrix} -1 & 1 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} q \\ p \end{pmatrix} + (\text{h.o.t}),$$

and for

$$P_{qq} = 2\alpha, \quad P_{qp} = \beta, \quad P_{qt} = \gamma, \quad P_{qqq} = 6\delta, \quad Q_{qq} = 2\zeta$$

with $\eta = \delta + \alpha\zeta + \alpha\beta/2$,

$$\gamma \neq 0, \quad \alpha^2 + 2\eta \neq 0.$$

Appendix 3. Bifurcation points

Let $\lambda_x = \exp(\pm 2\pi i l/k)$, $k \geq 3$, $(l, k) = 1$ and $0 < l < k$.

[Bir27] introduced a normalization procedure, which gives local expression

$$\begin{aligned}(q, p) &= (r^{1/2} \cos \theta, r^{1/2} \sin \theta), \\(Q, P) &= (R^{1/2} \cos \Theta, R^{1/2} \sin \Theta), \\ \Theta &= \theta + \frac{2\pi l}{k} + \frac{t\alpha(t)}{k} + \sum_{j=1}^{\lfloor (k-2)/2 \rfloor} \frac{\beta_j(t)}{k} r^j \\ &\quad + \frac{\gamma(t)}{k} r^{(k-2)/2} \cos k\theta + \Theta_1(\theta, r, t), \\ R &= r + \frac{2\gamma(t)}{k} r^{k/2} \sin k\theta + R_1(\theta, r, t).\end{aligned}$$

Appendix 3. Bifurcation points

Moreover, we require for $r^{1/2} = \rho$,

$$\frac{\partial^j R_1}{\partial \rho^j}(\theta, 0, 0) = 0 \quad \text{for } j = 0, 1, \dots, k+1$$

$$\frac{\partial^i \Theta_1}{\partial \rho^i}(\theta, 0, 0) = 0 \quad \text{for } i = 0, 1, \dots, k-1.$$

We denote $\alpha(0) = \alpha$, $\beta_1(0) = \beta$, $\gamma(0) = \gamma$. If

- ($k = 3$) $\alpha, \gamma \neq 0$,
- ($k = 4$) $\alpha, \gamma, \beta \pm \gamma \neq 0$,
- ($k \geq 5$) $\alpha, \beta, \gamma \neq 0$,

we call (x, s) a **k -bifurcation point**.

References i

-  Peter Albers, Joel W. Fish, Urs Frauenfelder, and Otto van Koert, *The Conley-Zehnder indices of the rotating Kepler problem*, Math. Proc. Cambridge Philos. Soc. **154** (2013), no. 2, 243–260. MR 3021812
-  Cengiz Aydin, Urs Frauenfelder, Otto van Koert, Dayoung Koh, and Agustin Moreno, *Symplectic geometry and space mission design*.
-  Ralph Abraham and Jerrold E. Marsden, *Foundations of mechanics*, second ed., Benjamin/Cummings Publishing Co., Inc., Advanced Book Program, Reading, MA, 1978, With the assistance of Tudor Rațiu and Richard Cushman. MR 515141
-  Cengiz Aydin, *The linear symmetries of Hill's lunar problem*, Arch. Math. (Basel) **120** (2023), no. 3, 321–330. MR 4549765
-  G.D. Birkhoff, *Dynamical systems*, American Mathematical Society / Providence, Estados Unidos, American Mathematical Society, 1927.

References ii

-  G. D. Birkhoff and D. C. Lewis, *On the periodic motions near a given periodic motion of a dynamical system*, Ann. Mat. Pura Appl. **12** (1934), no. 1, 117–133. MR 1553217
-  Michael A. Buchner, *On the generic nature of property $H1$ for Hamiltonian vector fields*, Global Analysis (Proc. Sympos. Pure Math., Vols. XIV, XV, XVI, Berkeley, Calif., 1968), Proc. Sympos. Pure Math., vol. XIV-XVI, Amer. Math. Soc., Providence, RI, 1970, pp. 51–54. MR 264701
-  K. Cieliebak, A. Floer, H. Hofer, and K. Wysocki, *Applications of symplectic homology ii: Stability of the action spectrum*, Mathematische Zeitschrift **223** (1996), no. 1, 27–45.
-  M.S El Naschie, *Vak, vacuum fluctuation and the mass spectrum of high energy particle physics*, Chaos, Solitons & Fractals **17** (2003), no. 4, 797–807.

References iii

-  Urs Frauenfelder, Dayung Koh, and Agustin Moreno, *Symplectic methods in the numerical search of orbits in real-life planetary systems*, SIAM Journal on Applied Dynamical Systems **22** (2023), no. 4, 3284–3319.
-  Martin Golubitsky, Ian Stewart, and Jerrold Marsden, *Generic bifurcation of hamiltonian systems with symmetry*, Physica D: Nonlinear Phenomena **24** (1987), no. 1, 391–405.
-  Michel Hénon and Carl Heiles, *The applicability of the third integral of motion: Some numerical experiments*, Astronom. J. **69** (1964), 73–79. MR 158746
-  Dongho Lee, *Conley-zehnder indices of spatial rotating kepler problem*, 2025.
-  K. R. Meyer, *Generic bifurcation of periodic points*, Transactions of the American Mathematical Society **149** (1970), no. 1, 95–107.

References iv

-  K. R. Meyer and D. S. Schmidt, *Periodic orbits near $\mathcal{L}_4$ for mass ratios near the critical mass ratio of Routh*, Celestial Mech. **4** (1971), 99–109. MR 317617
-  H. Poincaré, *Les méthodes nouvelles de la mécanique céleste. tome III*, Les Grands Classiques Gauthier-Villars. [Gauthier-Villars Great Classics], Librairie Scientifique et Technique Albert Blanchard, Paris, 1987, Invariant intégraux. Solutions périodiques du deuxième genre. Solutions doublement asymptotiques. Reprint of the 1899 original, Bibliothèque Scientifique Albert Blanchard. MR 926908
-  S. Smale, *Differentiable dynamical systems*, Bull. Amer. Math. Soc. **73** (1967), 747–817. MR 228014
-  Dieter S. Schmidt and Daniel Sweet, *A unifying theory in determining periodic families for hamiltonian systems at resonance*, Journal of Differential Equations **14** (1973), no. 3, 597–609.

References v



Ruikang Zhang, Yue Wang, Hao Zhang, and Chen Zhang, *Transfers from distant retrograde orbits to low lunar orbits*, Celestial Mechanics and Dynamical Astronomy **132** (2020), 41.