

1833. Maximum Ice Cream Bars

Difficulty: Medium

Link: <https://leetcode.com/problems/maximum-ice-cream-bars/description/?envType=daily-question&envId=2026-06-21>

Generated at: 2026-06-21 17:27:43

Problem:

It is a sweltering summer day, and a boy wants to buy some ice cream bars.

At the store, there are `n` ice cream bars. You are given an array `costs` of length `n`, where `costs[i]` is the price of the `i`th ice cream bar in coins. The boy initially has `coins` coins to spend, and he wants to buy as many ice cream bars as possible.

Note: The boy can buy the ice cream bars in any order.

Return *the **maximum** number of ice cream bars the boy can buy with `coins` coins.*

You must solve the problem by counting sort.

Example 1:

Input: `costs = [1,3,2,4,1]`, `coins = 7`

Output: 4

Explanation: The boy can buy ice cream bars at indices 0,1,2,4 for a total price of $1 + 3 + 2 + 1 = 7$.

Example 2:

Input: `costs = [10,6,8,7,7,8]`, `coins = 5`

Output: 0

Explanation: The boy cannot afford any of the ice cream bars.

Example 3:

Input: `costs = [1,6,3,1,2,5]`, `coins = 20`

Output: 6

Explanation: The boy can buy all the ice cream bars for a total price of $1 + 6 + 3 + 1 + 2 + 5 = 18$.

Constraints:

- `costs.length == n`
- $1 \leq n \leq 10^5$
- $1 \leq \text{costs}[i] \leq 10^5$
- $1 \leq \text{coins} \leq 10^8$