

Air sampling in team congregate spaces for early detection of respiratory virus threats

Viral air sampling in team settings


David Simon Timothy Locksmith Nick Minor Eli J. O'Connor
Shelby L. O'Connor David H. O'Connor

2026-08-11

i This is the interactive version of the manuscript. The figures below are explorable — hover for per-room, per-city detail. The prose is authored in Google Docs and synced automatically; the figures are regenerated from the committed data in [analysis/](#) on every build. A **submission PDF** with static figures is produced from the same source.

Central figure

FOUR SAMPLERS



Viral genetic material was detected in air samples from the team hotel in all five cities where Team Canada trained and played



Possible actions that could be triggered by detection of viral genetic material in air samples

Central figure. Air sampling for early detection of respiratory-virus threats in an elite team during competition. Continuous bioaerosol sampling ran in up to four team congregate spaces per hotel (meal, equipment, physiotherapy, and coaches' rooms); each cartridge's filter was eluted on-site and tested on a Cepheid GeneXpert SARS-CoV-2/Flu/RSV cartridge (top). Across the five host cities, viral genetic material was detected in team-hotel air in every city; the timeline shows sampling coverage and each detection over the tournament (middle). A positive air signal can trigger low-regret responses — masking, portable air purifiers, distancing and reduced occupancy, far-UVC, reinforced ventilation, and excluding visibly ill personnel (bottom).

Abstract (233 words)

Objective. Respiratory infections are the leading cause of illness at major sporting events, yet surveillance relies on athletes recognising and reporting symptoms. We evaluated whether continuous air sampling with point-of-care molecular testing could detect respiratory-virus nucleic acids in an elite team's congregate spaces during competition, and whether the resulting signals were operationally useful.

Methods. We performed a prospective, descriptive environmental-surveillance study following the Canadian men's national soccer team across five host cities during the 2026 FIFA World Cup™

(3 June to 4 July 2026). InBio Apollo bioaerosol samplers ran continuously in up to four team-designated rooms per hotel (physiotherapy, meal, equipment, and coaches’ room or hallway). Filters were changed approximately twice daily, eluted on-site, and tested with the Cepheid Xpert® Xpress SARS-CoV-2/Flu/RSVplus assay. A sample was considered positive if any cycle-threshold (Ct) value was reported, as less than 45, for a target.

Results. Of 176 air filters, there were 15 detections of virus genetic material (11 SARS-CoV-2, 3 influenza A virus, 1 influenza B virus, 0 RSV). Detections were sparse early and clustered late in the tournament. An influenza A signal appeared the morning a player was sent home febrile, and SARS-CoV-2 signals coincided with visibly ill hotel staff, with signals falling after ill staff were excluded.

Conclusion. Air sampling with point-of-care testing is feasible in the mobile environment of an elite team and can surface behavior-independent viral signals during competition that can be intercepted prior to player infection.

Summary Box

What is already known on this topic

Respiratory infections are the leading cause of illnesses at major sporting events, and conventional surveillance depends on athletes recognising and reporting symptoms, which misses pre-symptomatic, mild, and asymptomatic infections. Air sampling can detect respiratory viruses in congregate settings but has not been evaluated in elite team housing during competition.

What this study adds

Continuous air sampling with same-day point-of-care testing was feasible across a five-city World Cup campaign and detected SARS-CoV-2, influenza A, and influenza B in shared team spaces even during the low-circulation Northern-hemisphere summer, with several signals coinciding with a febrile player and with visibly ill staff.

How this study might affect research, practice or policy

Air sampling offers a candidate behavior-independent early-warning signal that could allow teams to escalate precautions selectively, minimizing outbreaks among athletes and team staff.

Introduction

Acute respiratory illness is the most common non-injury medical problem in elite sport and is consistently the single largest illness category at major sporting events. At the FIFA World Cup Qatar 2022 it accounted for 80% (12 of 15) of all time-loss illnesses among players [1], and it was the leading illness category at the 2018 and 2020 Youth Olympic Games [2,3]. Elite athletes also carry elevated risk: at the 2019 Nordic World Ski Championships they had roughly a sevenfold higher risk of symptomatic respiratory infection than exercising controls [4], and infections spread readily within teams, most often within the same sport discipline [5,6].

Shared housing, communal meals, travel, and crowded indoor spaces are repeatedly identified as risk factors [7,8]. Team hotels concentrate these hazards, gathering susceptible athletes and staff in a closed, shared-ventilation environment. There, athletes and staff can be exposed both by one another and by other hotel occupants. Even if the athletes do not stray from the team environments, other hotel staff could bring viruses from the community into the team’s setting. The main infection

route is inhalation of virus-laden aerosols, which accumulate and travel beyond close range in poorly ventilated air. Larger droplets can transmit virus during close face-to-face contact, and, less often, so can contact with contaminated surfaces [9].

Symptom-triggered surveillance is behavior-dependent, because infections are only ascertained when a person feels unwell, chooses to seek care or testing, and is correctly diagnosed. It misses or underestimates three situations that matter in team settings. First, asymptomatic and subclinical infections are common, with roughly a fifth of influenza and SARS-CoV-2 infections estimated to be asymptomatic [10–12], and RSV frequently goes unrecognised in adults [13]. Such infections are invisible to symptom-based monitoring yet can still shed virus and seed transmission.

The second class are symptomatic infections that are never reported or are reported after onset of symptoms. In one community study, only 17% (95% CI 10–26) of PCR-confirmed influenza infections received medical attention [14]. Even with highly conditioned athletes attuned to their bodies and close monitoring from trainers, during one professional rugby tournament, symptoms were present for at least a day before being reported to the team physician in more than half of illnesses [15]. In some cases, the incentive to keep competing may work against disclosure of symptoms.

Third, for many respiratory viruses shedding, and probably transmission, begins before symptoms do. This pre-symptomatic window is short, on the order of a day or two, but it is when an infected person is already seeding a shared space while still appearing well, which is why symptom-based detection lags the true course of an outbreak. In experimental influenza A infection most shedding preceded peak symptoms [16], influenza B shedding can rise up to two days before onset [17], and SARS-CoV-2 is detectable in the throat within roughly 40 hours of inoculation, before symptoms begin [18].

Importantly, elite team athletes cannot live in a pathogen-free bubble. Team members need to travel, hotel staff provide security and meal support, and team members mingle with family members who likely spend time in the surrounding communities. These individuals are not tested for infections, but provide an avenue for a pathogen to infiltrate the team population [19,20](Carmola et al. 2023; Shamier et al. 2023).

A behavior-independent environmental signal such as viral nucleic acid in air could reveal a virus in a shared space occupied by people with asymptomatic, pre-symptomatic, or unreported infections. Respiratory viruses have long been recoverable from indoor air, and high-flow bioaerosol sampling is now a practical surveillance tool [21]. In congregate living settings, air samplers in student-dormitory HVAC returns detected SARS-CoV-2 in most cases when a positive resident lived on the same floor [22], and school air sampling has captured SARS-CoV-2 during sustained classroom transmission [23, 24]. Our own work has detected viral nucleic acids in congregate settings such as schools and healthcare facilities [25] and at international airports [26].

To our knowledge, continuous indoor air sampling for early identification of viral threats has not been evaluated in the context of congregate elite athlete team communities. In partnership with Team Canada, we performed twice-daily air sampling in four team-hotel spaces during the 2026 FIFA World Cup™ to ask three questions: (1) can respiratory-virus nucleic acids be detected from the air of large, variably ventilated shared rooms that teams occupy; (2) are SARS-CoV-2, influenza A, influenza B, and RSV detectable during a summer competition, when Northern-hemisphere respiratory viruses circulate at low levels [27,28]; and (3) can air-sampling signals provide team physicians with behavior-independent information that is timely and specific enough to inform precautions?

Methods

Reporting follows the STROBE recommendations for observational studies where applicable [29,30]; because this is a descriptive environmental-surveillance and feasibility study without individual participants, items pertaining to individual-level exposures and outcomes are not applicable.

Study design and setting

We conducted a prospective, descriptive environmental-surveillance study of respiratory-virus nucleic acids in the air of team congregate spaces occupied by the Canadian men’s national soccer team during the 2026 FIFA World Cup™. Sampling began on June 3, 2026 (during the final week of pre-tournament training) and continued as the team moved between host cities, ending on July 4, 2026 when the team was eliminated. In Montreal, Toronto, and Vancouver, samplers were placed in up to four team-designated rooms (physiotherapy, meal, equipment, and coaches’ rooms) at locations approved by the team physician. At the final two venues, room access was more limited, with three rooms sampled in Los Angeles and, in Houston, the coaches’ room was replaced by a team-occupied hallway. Room type rather than a fixed physical space was the unit of sampling, because rooms differed between hotels and varied widely in size and ventilation.

Ethics and consent

This study sampled ambient air in shared congregate spaces and did not collect specimens from, or identifiers of, any individual. The University of Wisconsin-Madison Institutional Review Board has determined that air sampling in congregate settings, where the specific occupants of the sampled space are not known, does not constitute human-subjects research. Sampling locations were used with the agreement of, and coordinated with, Team Canada medical staff.

Patient and public involvement

Team Canada’s medical staff, led by the team physician, collaborated on the design and conduct of air collection, including selecting and approving room placements and informing the operational use of results. Athletes and members of the public were not involved in setting the research question, in the analysis, or in the writing of this report.

Equity, diversity, and inclusion

The sampled population comprised the players and staff of a single national team. The spaces were shared with different hotel staff and security in each city. Because air sampling captured no individual-level data, participant demographics were neither collected nor analysed. The study team was assembled to combine environmental-surveillance, virological, and sports medicine expertise.

Air sampling

Air was collected using InBio Apollo ambient air samplers (InBio), a filtration-based active bioaerosol sampler with a manufacturer-stated average flow rate of 540 L/min (32.4 m³/h) through its allergen-capture filter [31]. Samplers were positioned on available surfaces within each room and run continuously. These samplers are relatively low cost, at roughly 650 US dollars per unit, which made it practical to deploy several in parallel across the team’s shared spaces. They are also quiet (i.e., under 40 decibels) in operation, allowing them to be positioned discreetly in occupied rooms without interfering with team activities.

Filters were changed approximately twice daily, giving two collection sessions of roughly 8-12 hours each: a daytime session (~07:00–08:00 to ~14:00–16:00) and an overnight session (~14:00–16:00 to ~07:00–08:00). When a room was inaccessible (for example, when the equipment room was locked or the coaches’ room was in use), a single filter was run for ~24 hours. Each filter carried a unique barcode; the sampler location, and the times of filter installation and removal, were recorded for every session. Occasionally, an additional sampler was added to a single room and was used as a second confirmatory air sample for the space.

Sample processing and elution

All processing was performed on-site in a hotel room. For each cassette, the filter was removed from its 3D-printed holder with forceps and transferred to a tube containing 600–800 μ L of phosphate-buffered saline with 0.1% Tween-20 (PBST); 800 μ L was used typically. The tube was held at room temperature for ~20 minutes to elute captured material, with intermittent agitation. The eluate was then transferred directly into a Cepheid Xpert® cartridge for testing.

Molecular testing

Eluates were tested on the Cepheid GeneXpert® platform using the Xpert® Xpress SARS-CoV-2/Flu/RSV plus assay, a cartridge-based multiplex real-time RT-PCR that detects SARS-CoV-2, influenza A, influenza B, and respiratory syncytial virus. Each cartridge was run according to the manufacturer’s instructions.

Definition of a positive detection

We defined a sample as positive for a given virus if the GeneXpert reported any cycle-threshold (Ct) value for that target, regardless of the instrument’s qualitative call. Because the Xpert® Xpress assay applies validated Ct cut-offs to return a qualitative “positive/negative” result optimized for clinical diagnosis from patient specimens, targets with a late Ct near the assay boundary may be reported qualitatively as negative. For air surveillance, where the goal is the earliest possible detection of viral signal rather than a clinical diagnosis, we treated any reported amplification as evidence of viral nucleic acid in the sampled air. This lower threshold appears appropriate, though it necessarily trades specificity for sensitivity. Each Xpert® cartridge includes an internal Sample Processing Control that verifies adequate processing and the absence of gross inhibition. Samples with a non-amplifying control or a probe error, or otherwise determined to be invalid, were excluded from the figures but retained in the study data table.

Target-enriched Illumina VSP2 viral sequencing

The final air-sample eluates from July 4, 2026 testing in Houston were reserved for viral whole-genome sequencing to characterise detected viruses and, where possible, resolve lineage. Nucleic acid from these eluates underwent target enrichment for respiratory viruses [specify panel/probe set and library-preparation kit] followed by sequencing on an Illumina NovaSeq [specify instrument/flow-cell and read configuration].

Illumina paired-end reads were imported into Lungfish Genome Explorer ([v0.5.0-beta25](#)) using the VSP2 target-enrichment preprocessing recipe. On import, read pairs were adapter- and quality-trimmed and deduplicated with fastp (v1.3.2; sliding-window trimming, Q15, duplicate removal), depleted of human reads with deacon (v0.15.0) against the panhuman-1 index, and overlapping pairs were merged with fastp; merged and unmerged reads were then length-filtered to 50 bp with

seqkit (v2.13.0) and deduplicated/reordered with BBTools clumpify (v39.80). The resulting single-end reads were analyzed with EsVirtu (v3.14) against its curated viral reference database (v3.2.4), which applies its own quality filtering and read deduplication during competitive alignment to detect viral taxa and generate reference-guided consensus sequences. Because reads are deduplicated both at import (fastp/clumpify) and again during EsVirtu mapping, reported read counts reflect unique reads per mapped taxa rather than raw sequenced reads.

Outcomes and analysis

The primary output was the presence or absence of each of the four target viruses in each air sample. Ct values were used as a rough proxy for the amount of viral nucleic acid, with lower Ct indicating more material. No sample-size calculation was performed.

Statistical analysis

Analyses were primarily descriptive, and reporting follows the CHAMP statement [32]. Detections were tabulated by virus, room, city, and sampling interval.

Results

Sampling overview

Over 32 days (June 3 to July 4, 2026), we collected and tested 182 air-sample filters for SARS-CoV-2, influenza A (IAV), influenza B (IBV), and RSV across five host cities: Montreal (n = 20), Toronto (n = 36), Vancouver (n = 79), Los Angeles (n = 11), and Houston (n = 36). Six filters (3 in Vancouver, 1 in Los Angeles, and 2 in Houston) were excluded from analysis because of invalid Cepheid GeneXpert tests. The remaining 176 filters were used for all analyses. Sampling covered four team spaces per hotel in Montreal, Toronto, and Vancouver (physiotherapy/treatment room, meal room, equipment/kit room, coaches' meeting room), three in Los Angeles (no coaches' room), and four in Houston (the coaches' room replaced by a team-occupied hallway). Sampling was briefly interrupted when teams and staffing equipment moved between each city.

Most samples were negative for all four targets. Across the study, 15 filters yielded a detection of at least one respiratory virus: SARS-CoV-2 in 11, IAV in 3, and IBV in 1. No RSV was detected. Detections and their cycle-threshold (Ct) values are summarized in Figure 1.

Detections by city

Detections were sparse and geographically dispersed through the first three cities. In Montreal, SARS-CoV-2 was detected in the meal room on June 5 (Ct 42.8), against an otherwise negative baseline. Toronto also yielded one detection, when influenza A was detected in the physiotherapy room from the overnight session beginning June 9 (Ct 44.9). This value fell beyond the assay's qualitative cut-off and was reported by the instrument as negative, but it met our detection definition of any reported amplification. In Vancouver, influenza B was detected in the physiotherapy room on June 17 (Ct 38.0) and influenza A was detected in the meal room on the morning of June 23 (Ct 40.6).

Detections became more frequent in the two final host cities. In Los Angeles, SARS-CoV-2 was detected in the meal room beginning on the first sampling day. On June 27, two filters were positive from the meal room in the daytime sampling interval (Ct 34.0 and 39.3). An additional filter was

collected overnight from the meal room and also tested positive for SARS-CoV-2 genetic material (Ct 38.0). There were no additional positives before the air sampling supplies were moved on June 29 to Houston, where Team Canada played their Round of 16 match. Houston accounted for 8 of the 15 detections (Figure 2). On July 1, viral genetic material was found in the air of all four sampled areas: the meal room (Ct 40.4), hallway (Ct 42.1), and the physiotherapy room (Ct 43.4) tested positive for SARS-CoV-2, with influenza A in the equipment room (Ct 42.5). On the morning of July 2, only the meal room tested positive from the previous overnight collection (Ct 42.7). No viruses were detected again until the morning of July 4 when SARS-CoV-2 returned on samples at lower Ct values. There were two meal-room samples positive (Ct 34.5 and 35.5) and a positive hallway collection (Ct 37.8). One speculative interpretation, which we advance cautiously and cannot confirm, is that the July 1 signal reflected an introduction from someone outside the team, the removal of the source individual(s) transiently reduced airborne virus, and that the lower-Ct resurgence reflected onward transmission within the team.

Community wastewater surveillance in all five host cities showed that the team’s visits fell in the seasonal trough for all four viruses, so the clustering of air detections in Los Angeles and Houston cannot be explained by unusually high background circulation (online supplemental methods and online supplementary figure 1) in these cities.

Operational context and team responses

A prespecified aim was to determine whether air-sampling signals could give team physicians actionable, behavior-independent information. In Toronto, the influenza A detection in the physiotherapy room (overnight session from June 9, Ct 44.9) coincided with a player being withdrawn from training that morning with a fever. Team staff asked the player to mask and isolate for several days [confirm details]. In Los Angeles, after SARS-CoV-2 was detected in the meal room on June 27, training staff asked visibly ill hotel personnel to leave the team spaces to reduce exposure [confirm which staff/roles, timing, and who made the request], and no respiratory virus was detected in the subsequent June 28 samples, coinciding with the removal. In Houston, following the July 1 detections across multiple team spaces, training staff again asked visibly ill hotel staff to leave, after which signal was absent on July 2 and returned on July 3 at lower Ct values.

Viral sequencing

Metagenomic sequencing of the VSP2 air samples detected human respiratory and environmental viruses across the sampled hotel spaces (Figure 3). SARS-CoV-2 was recovered from two independent meal-room air samples collected on July 4, as well as from an equipment-room sample from the preceding evening (1 unique read). Although these specific cartridges were pulled from the samplers the afternoon Team Canada was eliminated and were not themselves assayed on the GeneXpert, their sequences independently confirm the presence of SARS-CoV-2 in the hotel earlier that day. Sequencing additionally recovered respiratory viruses not targeted by the Cepheid assay, including influenza A, human coronavirus OC43, and rhinovirus, alongside ubiquitous commensal viruses such as Merkel cell polyomavirus that are detected in nearly all sequences from human-occupied congregate spaces. No viral detections were obtained from the extraction blank or the no-template controls.

A summary of the study design, detections across all five host cities, and the response actions a positive signal can trigger is presented in the Central Figure.

Discussion

Principal findings

Over 32-days of participation in a tournament and five host cities, twice-daily air sampling with same-day point-of-care testing detected respiratory-virus nucleic acid in team congregate spaces even during the Northern-hemisphere summer, when these viruses circulate at seasonal lows. Detections were infrequent but not random. They clustered in the final two host cities, and on three occasions coincided with events recognised by the team’s medical staff. In each case staff acted, and the air signal subsequently cleared or fell. Positive detection of virus genetic material in the air stood in stark contrast to the negative baseline that was routinely observed in air collected from the team spaces, particularly during early phases of the tournament. Continuous air sampling paired with point-of-care testing is therefore operationally feasible in the demanding, mobile environment of a competing national team, and the presence of distinct signals can be acted on in near-real time.

Comparison with existing literature

To our knowledge, this is the first description of viral air sampling in the context of sports medicine.

Clinical and research implications

Respiratory illness is consistently the single greatest illness burden in athlete health-surveillance programmes. Over four years of monitoring UK Olympic athletes it caused the largest share of 27,442 illness time-loss days [33], and it remains the leading medical problem at major Games even when prevention campaigns are in place, as at Paris 2024 [34]. Within a squad, congregate living amplifies risk. In a football academy, the SARS-CoV-2 attack rate was three times higher among those who both lived and trained on-site than among those who only trained there [35]. A single infection introduced into shared team spaces during a tournament can therefore threaten both athlete health and competitive availability.

Intensive infection-control measures unquestionably work in this population but are not sustainable indefinitely. Across four major winter-sport events, acute respiratory illness affected 38% of team members at PyeongChang 2018 and 26% at the 2019 World Ski Championships, but fell to 0% and 5% under the multilayered COVID-19 countermeasures at the 2021 and 2022 events, a roughly 10-fold reduction [36], and a parallel cohort showed annual illness incidence in elite skiers falling from 5.3 to 0.3 episodes per person during lockdown [7]. Those disruptive measures, such as quarantine, continuous masking, single-room housing, and restricted indoor facilities, are not compatible with normal competition across a month-long tournament, and illness returned as they relaxed [36]. An early-warning signal could instead let intensive precautions be applied selectively, escalated when risk is detected and relaxed when it is not.

For team medical practice, air sampling offers such a behavior-independent early-warning signal, one that does not depend on athletes reporting symptoms or submitting to individual testing. Even if team members are routinely tested, air sampling can assess whether other employees who occupy spaces in temporary hotel spaces could introduce viruses. This may be especially attractive in elite sports, where athletes may under-report illness to avoid being withheld from competition and where individual testing is logistically and ethically fraught during a tournament. A positive air signal can prompt low-cost, low-regret actions such as reinforcing ventilation [37], adding portable air purifiers [38], incorporating far-UVC lighting in highly trafficked spaces [39,40], masking in

the implicated space, reviewing who has access to team areas, excluding visibly ill personnel, and heightening clinical vigilance.

Limitations

First, this is a descriptive, uncontrolled study of a single team. We observed associations between air signals and events on the team, but cannot establish that air sampling caused any reduction in transmission, nor that the operational responses were effective. Second, air detection demonstrates the presence of viral nucleic acid in a shared space. It does not identify who was infected, does not distinguish infectious virus from residual nucleic acid, and documents exposure rather than transmission. Third, our positivity definition (any reported Ct) increases sensitivity at the cost of specificity, and low-level signals near the limit of detection may include contamination or transient environmental nucleic acid. Fourth, the volume of air sampled per session, number of occupants in each space, and their distance from the air samplers are all uncertain, which limits any quantitative interpretation of Ct values. Finally, sampling was opportunistic and constrained by room access, competition logistics, and the team’s elimination, which ended data collection.

Future research

The air samplers used here represent an early generation of the technology and are relatively insensitive, collecting onto a filter that must be removed and processed before any testing. The next generation of samplers is likely to integrate continuous, automated pathogen detection directly into the device. Programs such as the ARPA-H Building Resilient Environments for Air and Total Health initiative are explicitly developing autonomous indoor air biosensors that continuously monitor the biological content of the air and translate it into real-time risk estimates [41]. As these tools mature, approaches to early detection of respiratory threats in athletes should continue to improve in sensitivity, speed, and ease of deployment.

Assay breadth is likely to expand in parallel. Cepheid has described a GeneXpert Respiratory Panel prototype that runs on the same point-of-care platform used here and detects 26 respiratory pathogens, including rhinovirus, enterovirus, parainfluenza viruses, seasonal coronaviruses, adenoviruses, and metapneumoviruses alongside SARS-CoV-2, influenza, and RSV [42]. Applying a panel of this breadth to air samples would let air surveillance capture the common-cold viruses that account for much athlete illness but that our SARS-CoV-2, influenza, and RSV assay could not detect, giving better representation of overall respiratory illness burden. Broader pathogen coverage would also extend the approach beyond respiratory viruses. Enteric illness is consistently among the leading non-injury problems in elite sport, close behind respiratory illness in prospective tournament surveillance [15], and travel compounds the risk, with traveler’s diarrhoea the most common travel-related illness and one that can interfere with training and performance [43]. Norovirus in particular has disrupted mass-gathering events, including a PyeongChang 2018 outbreak that prompted surveillance and exclusion of infected food handlers [44]. Air sampling can in principle capture the genetic material of these agents too, and pathogen-agnostic sequencing of air-sample nucleic acid from congregate settings has recovered enteric viruses such as rotavirus and astrovirus alongside respiratory viruses [45]. Consequently, a longer-term goal is on-site metagenomic sequencing of air-sample nucleic acid, which would allow simultaneous, hypothesis-free detection of respiratory, enteric, and skin-sloughed pathogens from a single collection rather than testing for one predefined set of targets.

Conclusion

Twice-daily air sampling with same-day point-of-care testing was feasible in the mobile, high-pressure environment of a national team during a World Cup, and detected respiratory-virus nucleic acid in shared team spaces even in a low-circulation summer season. Several detections coincided with a febrile player and with visibly ill staff, and prompted the team’s medical staff to act. These findings position congregate air surveillance as a promising, behavior-independent early-warning tool for protecting athlete health.

Required end statements

Contributorship statement

DHO and SLO conceived and designed the study. DS facilitated access to team spaces, approved sampler placements, and provided clinical and operational context. SLO, EJO, DHO, TL, and NM conducted field air sampling, sample processing, and point-of-care testing across the host cities (SLO, EJO, and DHO in Montreal, Vancouver, Los Angeles, and Houston; TL in Toronto; NM in Vancouver). DHO and SLO drafted the manuscript, and all authors critically revised it for intellectual content, approved the final version, and agree to be accountable for all aspects of the work. Claude Opus 4.8 (high reasoning) was used for background research, literature retrieval and review, manuscript text drafting and organization, data analysis and visualization, and editing. All LLM-generated content was independently verified and further edited by co-authors.

Guarantor: Co-lead authors David H. O’Connor and Shelby L. O’Connor accept full responsibility for the work and the conduct of the study, had access to the data, and controlled the decision to publish.

Competing interests

D.H.O. and S.L.O. are managing partners of Pathogenuity LLC, a consultancy that advises on topics including environmental monitoring for pathogens.

Funding

D.H.O. and S.L.O. received support for this project from Inkfish and Heart of Racing.

Ethics approval

The University of Wisconsin–Madison Institutional Review Board determined that air sampling in congregate settings, where the specific occupants of the sampled space are not known, does not constitute human-subjects research. The study did not collect specimens from, or identifiers of, any individual.

Data sharing statement

The data and code for this study are available in the repository at <https://github.com/dholab/team-canada-world-cup-2026>. A compressed archive of the Lungfish Genome Explorer project (including imported sequencing reads) used to evaluate sequencing reads found in the July 4, 2026 samples can be accessed from https://dholk.primate.wisc.edu/_webdav/dho/public/manuscripts/team-canada-air-sampling/%40files/Team_Canada_VSP2.lungfish.zip. The original reads before import processing for Lungfish Genome Explorer will be available from SRA upon publication.

Patient and public involvement

Patient and public involvement is described in the Methods (see Patient and public involvement).

Equity, diversity and inclusion

An Equity, Diversity and Inclusion statement is provided in the Methods.

Acknowledgements

We thank the players and staff of Team Canada for their cooperation and for accommodating air sampling in their team spaces during the 2026 FIFA World Cup™.

Provenance and peer review

Not commissioned; externally peer reviewed.

Figures




Figure 1: **Respiratory-virus detections in team-space air samples across the 2026 FIFA World Cup.** Air was sampled continuously in Team Canada congregate spaces across five host cities from 3 June to 4 July 2026, and every cartridge was tested for SARS-CoV-2, influenza A, influenza B, and respiratory syncytial virus (RSV). All cities are shown on a single continuous local-time axis. Rows are the four target viruses; each rectangle is one cartridge tested for one virus, drawn to that cartridge’s exact sampling window. Rooms are merged within each city, so each box summarises all rooms sampled in that session. A box is blue if the virus was detected in any room, with fill mapped to the lowest cycle-threshold (Ct) value across rooms; white if no room detected the virus but at least one returned a valid negative result; grey if every room in that session was invalid. An asterisk beneath the axis marks a session with partial room validity. A sample counted as a detection if the instrument reported any Ct value, regardless of qualitative call. The per-room data are explorable above, and the Houston episode is shown by room in Figure 2.

The same Houston episode was examined independently by metagenomic sequencing (Figure 3), which confirms the airborne SARS-CoV-2 and recovers additional respiratory viruses that the GenExpert panel does not target.

Data and code

The underlying dataset — one row per cartridge $\times$ target virus (728 rows: 182 respiratory cartridges $\times$ 4 targets) — and the script that builds every figure are in the repository:




Figure 2: Per-room detections during the Houston sampling period. Sampling in Houston ran from the evening of 30 June to 4 July 2026 across four team spaces (physiotherapy room, meal room, equipment room, and a team-occupied hallway). Rows group the four target viruses within each room; the horizontal axis is local wall-clock time. Each rectangle is one cartridge tested for one virus, drawn to its exact sampling window. White boxes are valid negatives, grey boxes are invalid runs, and blue boxes are detections, with fill mapped to the Ct value. On 1 July, SARS-CoV-2 was detected in the meal room, the physiotherapy room, and the hallway, and influenza A in the equipment room. No virus was detected on 2 July. SARS-CoV-2 returned on 3–4 July at lower Ct, most strongly in the meal room. The two invalid runs fell on 1 July.




Figure 3: Human viruses detected by metagenomic sequencing of the Houston air samples. Six air cartridges from the Houston team spaces were pulled from the samplers on the afternoon of 4 July 2026 and sequenced, and reads were classified against a curated viral reference database with EsVirtu. These cartridges were collected the same day but are not the same cartridges tested on the GeneXpert, so sequencing is an independent line of evidence for the viruses present. Columns are the six sequenced samples, labelled by room, GeneXpert SARS-CoV-2 status where a same-room result exists, and sampling runtime. Rows are the human viruses detected, with Merkel cell polyomavirus and SARS-CoV-2 on top and the rest ordered by total distinct reads. Each cell reports distinct (deduplicated) reads for that virus in that sample, with fill mapped to that count on a light-to-deep blue scale. A cell is left blank when the virus was not detected in that sample, and SARS-CoV-2 was assessed in every sample. SARS-CoV-2 was recovered from both meal-room samples and the equipment room, confirming the airborne SARS-CoV-2 seen in these spaces on 4 July. Merkel cell polyomavirus, a commensal skin virus, was found in every sample. Sequencing additionally recovered influenza A, human coronavirus OC43, and rhinovirus, none of which are on the GeneXpert panel.

- Data: [analysis/data/cartridges_long.csv](#)
- Main-figure pipeline (Figures 1–2): [analysis/make_figures.py](#)
- Supplemental-figure pipeline (online supplemental figure 1): [analysis/make_figure3.py](#)
- Sequencing figure (Figure 3): [analysis/make_figure3_sequencing.py](#) with data [analysis/data/sequencing_detections.csv](#)


Columns: `city`, `room`, `sampler`, `cartridge`, `start`, `end`, `dur_h`, `virus`, `ct`, `qual`, `detected`, `spc`, `spc_ct`, `status`. A blank `ct` means the target did not amplify; `detected = 1` marks a reported Ct regardless of the instrument’s qualitative call; `spc/spc_ct` are the internal sample-processing control’s call and Ct; `status` flags each cartridge as `valid` or `invalid`. The figures show valid runs only, so invalid runs are drawn grey or marked with an asterisk rather than plotted as results.

The wastewater comparison in [online supplemental figure 1](#) is built from per-city extracts of the public dashboards, committed alongside the figure script so the figure is reproducible without re-downloading the large source files:

- Canada (Montreal, Toronto, Vancouver): [analysis/data/ww_canada.csv](#), from the [PHAC wastewater aggregate](#)
- Los Angeles (JWPCP): [analysis/data/ww_losangeles.csv](#), from the California CDPH/NWSS open dataset
- Houston (69th Street): [analysis/data/ww_houston.csv](#), from the Rice/HHD dashboard ArcGIS feature service

Each extract has columns `city`, `target`, `week`, `value`, `source`, `metric`; `target` uses `covN2`, `fluA`, `fluB`, `rsv`; `metric` records which non-interchangeable quantity that source reports (`pmmov_index`, `fecal_normalized`, `houston_vl_index`).

Supplemental material



Online supplemental figure 1. Community wastewater surveillance context in each host city over the 2025–2026 respiratory season. Each panel (A to E) is one host city, plotting publicly reported municipal wastewater levels for the four study targets (SARS-CoV-2, influenza A, influenza B, and RSV, with the same colour used for each virus in every panel) from August 2025 through mid-July 2026. The shaded gold band marks Team Canada’s air-sampling window in that city, widened to a minimum readable width for the shortest 1 to 2 day stays. **The four data sources report non-interchangeable quantities and are therefore each shown on their own independent vertical axis. Levels must be read within a panel and never compared between panels.** Panels A to C (Montreal, Toronto, and Vancouver) use the Public Health Agency of Canada wastewater aggregate (health-infobase), a PMMoV-normalized viral index, with the weekly median taken across contributing sub-sites (Vancouver is reported as “Metro Vancouver”). Open circles mark non-detections plotted at the source’s detection-limit substitution value. Panel D (Los Angeles) uses the California open dataset (CDPH and CDC NWSS) for the Joint Water Pollution Control Plant in Carson, which serves the Torrance area where the team stayed. Because that source reports absolute concentrations, each target is self-normalized to fecal load as copies/L divided by the reported human fecal marker concentration ($\times 10$), and influenza B was not reported at this plant during the season. Panel E (Houston) uses the Rice University and Houston Health Department dashboard for the 69th Street plant serving the downtown Main Street hotel, reporting the published viral-load index for SARS-CoV-2 only at plant level, scaled so that 100 represents the July 2020 baseline. The dotted line and hatched band mark that this public feed ends 22 June 2026, before the team’s 30 June to 4 July stay. Across all five cities the visit windows fall in the seasonal trough, at or near each site’s own lowest values of the season, indicating that no host city, Canadian or US, was experiencing an unusually high level of community respiratory-virus circulation while the team was present.