

Module 3

Integration :

Review And Some New Techniques

Sub-Module 3.1

Beta functions with properties

Example:1

❖ Evaluate $\int_0^1 \sqrt{1 - \sqrt{x}} dx \cdot \int_0^{\frac{1}{2}} \sqrt{2y - 4y^2} dy$

❖ **Solution:** Consider $I_1 = \int_0^1 \sqrt{1 - \sqrt{x}} dx$

Put $\sqrt{x} = t \quad \therefore x = t^2 \Rightarrow dx = 2t dt$

x	0	1
t	0	1

$$I_1 = \int_0^1 \sqrt{1 - t} 2t dt$$

$$= 2 \int_0^1 t^1 (1 - t)^{\frac{1}{2}} dt$$

$$= 2 \int_0^1 t^{2-1} (1 - t)^{\frac{3}{2}-1} dt$$

$$= 2\beta\left(2, \frac{3}{2}\right)$$

Consider $I_2 = \int_0^{\frac{1}{2}} \sqrt{2y - 4y^2} \, dy$

$$= \int_0^{\frac{1}{2}} \sqrt{2y(1 - 2y)} \, dy$$

Put $2y = t \therefore 2 \, dy = dt \Rightarrow dy = \frac{dt}{2}$

$$\begin{aligned} I_2 &= \int_0^1 \sqrt{t} (1 - t)^{\frac{1}{2}} \frac{dt}{2} \\ &= \frac{1}{2} \int_0^1 t^{\frac{3}{2}-1} (1 - t)^{\frac{3}{2}-1} \, dt \\ &= \frac{1}{2} \beta\left(\frac{3}{2}, \frac{3}{2}\right) \end{aligned}$$

y	0	$\frac{1}{2}$
t	0	1

$$\begin{aligned}
 I &= I_1 \cdot I_2 = 2\beta \left(2, \frac{3}{2}\right) \frac{1}{2} \beta \left(\frac{3}{2}, \frac{3}{2}\right) \\
 &= \frac{\left|\overline{2}\right|_{\frac{3}{2}}}{\left|\overline{7}\right|_{\frac{1}{2}}} \frac{\left|\overline{3}\right|_{\frac{3}{2}}}{\left|\overline{3}\right|_{\frac{1}{2}}} \\
 &= \frac{1 \times \frac{1}{2} \sqrt{\pi} \frac{1}{2} \sqrt{\pi} \frac{1}{2} \sqrt{\pi}}{\frac{5}{2} \times \frac{3}{2} \times \frac{1}{2} \times \sqrt{\pi} \times 2} \\
 &= \frac{\frac{\pi}{8}}{\frac{15}{4}} = \frac{\pi}{8} \times \frac{4}{15} = \frac{\pi}{30}
 \end{aligned}$$

Example:2

❖ Evaluate $\int_0^{2a} x^2 \sqrt{2ax - x^2} dx$

❖ **Solution:** $I = \int_0^{2a} x^2 \sqrt{x(2a - x)} dx$

$$= \int_0^{2a} x^{2+\frac{1}{2}} (2a - x)^{\frac{1}{2}} dx$$

Put $x = 2at$, $dx = 2a dt$

x	0	$2a$
t	0	1

$$I = \int_0^{2a} (2at)^{\frac{5}{2}} (2a - 2at)^{\frac{1}{2}} 2a dt$$

$$= \int_0^1 (2at)^{\frac{5}{2}} (2a)^{\frac{1}{2}} (1 - t)^{\frac{1}{2}} 2a dt = \int_0^1 (2a)^{\frac{5}{2} + \frac{1}{2} + 1} t^{\frac{5}{2}} (1 - t)^{\frac{1}{2}} dt$$

$$= 16a^4 \int_0^1 t^{\frac{7}{2}-1} (1 - t)^{\frac{3}{2}-1} dt$$

$$= 16a^4 \beta \left(\frac{7}{2}, \frac{3}{2} \right)$$

$$I = 16a^4 \beta \left(\frac{7}{2}, \frac{3}{2} \right)$$

$$= 16a^4 \frac{\left| \frac{7}{2} \right| \left| \frac{3}{2} \right|}{\left| 5 \right|}$$

$$= \frac{16a^4 \times \frac{5}{2} \times \frac{3}{2} \times \frac{1}{2} \times \sqrt{\pi} \times \frac{1}{2} \times \sqrt{\pi}}{4 \times 3 \times 2}$$

$$= \frac{16a^4 \times 15\pi}{16 \times 8 \times 3}$$

$$= \frac{5\pi a^4}{8}$$

Example:3

❖ Prove that $\int_0^1 \frac{dx}{\sqrt{1-x^n}} = \frac{2^{\frac{2}{n}-1} \left(\frac{1}{n}\right)^2}{n \left|\frac{2}{n}\right|}$

❖ **Solution:** Put $x^n = t$

❖ $x = t^{\frac{1}{n}}$

x	0	1
t	0	1

❖ $dx = \frac{1}{n} t^{\frac{1}{n}-1} dt$

❖ $I = \int_0^1 \frac{1}{n} \frac{t^{\frac{1}{n}-1} dt}{\sqrt{1-t}}$

❖ $= \frac{1}{n} \int_{t=0}^1 t^{\frac{1}{n}-1} (1-t)^{\frac{1}{2}-1} dt$

❖ $= \frac{1}{n} \beta\left(\frac{1}{n}, \frac{1}{2}\right)$

$$\diamond I = \frac{1}{n} \beta \left(\frac{1}{n}, \frac{1}{2} \right)$$

$$\diamond = \frac{1}{n} \frac{\left| \frac{1}{n} \right| \frac{1}{2}}{\left| \frac{1}{n} + \frac{1}{2} \right|}$$

$$\diamond = \frac{1}{n} \frac{\left| \frac{1}{n} \sqrt{\pi} \right|}{\left| \frac{1}{2} + \frac{1}{n} \right|} \dots\dots\dots(1)$$

$$\diamond \text{ By duplication formula, } 2^{2m-1} |\overline{m}| \overline{m + 1/2} = \sqrt{\pi} |\overline{2m}|$$

$$\diamond \text{ Put } m = \frac{1}{n}$$

$$\diamond 2^{\frac{2}{n}-1} \left| \frac{1}{n} \right| \overline{\frac{1}{n} + 1/2} = \sqrt{\pi} \left| \frac{2}{n} \right|$$

$$\diamondsuit 2^{\frac{2}{n}-1} \left| \frac{1}{n} \right| \overline{\left| \frac{1}{n} + \frac{1}{2} \right|} = \sqrt{\pi} \left| \frac{2}{n} \right|$$

$$\diamondsuit \frac{\sqrt{\pi}}{\left| \frac{1}{n} + \frac{1}{2} \right|} = \frac{2^{\frac{2}{n}-1} \left| \frac{1}{n} \right|}{\left| \frac{2}{n} \right|} \dots\dots\dots(2)$$

❖ Using (2) in (1)

$$\diamondsuit I = \frac{1}{n} \left| \frac{1}{n} \right| \frac{\sqrt{\pi}}{\left| \frac{1}{2} + \frac{1}{n} \right|}$$

$$\diamondsuit = \frac{1}{n} \left| \frac{1}{n} \right| \frac{2^{\frac{2}{n}-1} \left| \frac{1}{n} \right|}{\left| \frac{2}{n} \right|} = \frac{2^{\frac{2}{n}-1} \left(\left| \frac{1}{n} \right| \right)^2}{n \left| \frac{2}{n} \right|}$$

Example 4

❖ **Prove That** $\int_0^a x^4 (a^2 - x^2)^{\frac{1}{2}} dx = \frac{a^6 \pi}{32}$

❖ **Solution:** Put $x^2 = a^2 t$

❖ $x = a t^{\frac{1}{2}}, \quad dx = \frac{a}{2} t^{\frac{1}{2}-1} dt$

❖ $I = \int_{t=0}^1 a^4 t^2 (a^2 - a^2 t)^{\frac{1}{2}} \frac{a}{2} t^{\frac{1}{2}-1} dt$

❖ $= a^4 \times a \cdot \frac{a}{2} \int_{t=0}^1 t^2 (1 - t)^{\frac{1}{2}} t^{\frac{1}{2}-1} dt$

❖ $= \frac{a^6}{2} \int_{t=0}^1 t^{\frac{5}{2}-1} (1 - t)^{\frac{3}{2}-1} dt$

❖ $= \frac{a^6}{2} \beta \left(\frac{5}{2}, \frac{3}{2} \right) = \frac{a^6}{2} \frac{\left| \frac{5}{2} \right| \left| \frac{3}{2} \right|}{\left| 4 \right|}$

❖ $= \frac{a^6}{2} \times \frac{\frac{3}{2} \times \frac{1}{2} \sqrt{\pi} \times \frac{1}{2} \times \sqrt{\pi}}{3.2.1} = \frac{a^6 \pi}{32}$

x	0	a
t	0	1

Example:5

❖ Evaluate $\int_0^1 \frac{x^7}{\sqrt{1-x^2}} dx$

❖ **Solution:** Put $x = \sin \theta \therefore dx = \cos \theta d\theta$

❖ $I = \int_0^{\frac{\pi}{2}} \frac{\sin^7 \theta}{\sqrt{1-\sin^2 \theta}} \cos \theta d\theta$

❖ $= \int_0^{\frac{\pi}{2}} \sin^7 \theta d\theta$

❖ $= \int_0^{\frac{\pi}{2}} \sin^7 \theta \cos^0 \theta d\theta = \frac{1}{2} \beta \left(\frac{7+1}{2}, \frac{0+1}{2} \right)$

❖ $= \frac{1}{2} \beta \left(4, \frac{1}{2} \right)$

❖ $= \frac{1}{2} \times \frac{|\overline{4}|^{\frac{1}{2}}}{|\overline{\frac{9}{2}}|}$

❖ $= \frac{1}{2} \times \frac{3 \times 2 \times \sqrt{\pi}}{\frac{7}{2} \times \frac{5}{2} \times \frac{3}{2} \times \frac{1}{2} \times \sqrt{\pi}} = \frac{16}{35}$

x	0	1
θ	0	$\frac{\pi}{2}$

Example 6

❖ Evaluate $\int_0^1 \frac{x^2(4-x^4)}{\sqrt{1-x^2}} dx$

❖ Solution:

❖ $I = \int_0^1 \frac{4x^2}{\sqrt{1-x^2}} dx - \int_0^1 \frac{x^6}{\sqrt{1-x^2}} dx$

❖ Put $x = \sin \theta \therefore dx = \cos \theta d\theta$

❖ $I = \int_0^{\frac{\pi}{2}} \frac{4 \sin^2 \theta}{\cos \theta} \cos \theta d\theta - \int_0^{\frac{\pi}{2}} \frac{\sin^6 \theta}{\cos \theta} \cos \theta d\theta$

❖ $= 4 \int_0^{\frac{\pi}{2}} \sin^2 \theta \cos^0 \theta d\theta - \int_0^{\frac{\pi}{2}} \sin^6 \theta \cos^0 \theta d\theta$

❖ $= 4 \times \frac{1}{2} \beta \left(\frac{2+1}{2}, \frac{0+1}{2} \right) - \frac{1}{2} \beta \left(\frac{6+1}{2}, \frac{0+1}{2} \right)$

x	0	1
θ	0	$\frac{\pi}{2}$

$$\diamondsuit I = 2\beta\left(\frac{3}{2}, \frac{1}{2}\right) - \frac{1}{2}\beta\left(\frac{7}{2}, \frac{1}{2}\right)$$

$$\diamondsuit = 2 \frac{\left|\frac{3}{2}\right|\left|\frac{1}{2}\right|}{\left|\frac{2}{2}\right|} - \frac{1}{2} \times \frac{\left|\frac{7}{2}\right|\left|\frac{1}{2}\right|}{\left|\frac{4}{2}\right|}$$

$$\diamondsuit = 2 \times \frac{1}{2} \times \sqrt{\pi} \times \sqrt{\pi} - \frac{\frac{1}{2} \times \frac{5}{2} \times \frac{3}{2} \times \frac{1}{2} \times \sqrt{\pi} \times \sqrt{\pi}}{3 \times 2}$$

$$\diamondsuit = \pi - \frac{5\pi}{32}$$

$$\diamondsuit = \frac{27\pi}{32}$$

Example:7

❖ Evaluate $\int_0^{\pi} \frac{\sin^4 \theta}{(1+\cos \theta)^2} d\theta$

❖ Solution:

$$\text{❖ } I = \int_0^{\pi} \frac{\sin^4 \theta}{\left(2 \cos^2 \frac{\theta}{2}\right)^2} d\theta$$

$$\text{❖ } = \frac{1}{4} \int_0^{\pi} \sin^4 \theta \cos^{-4} \frac{\theta}{2} d\theta$$

$$\text{❖ } = \frac{1}{4} \int_0^{\pi} \left[\sin \left(\frac{\theta}{2} \right) \cos \left(\frac{\theta}{2} \right) \right]^4 \cos^{-4} \left(\frac{\theta}{2} \right) d\theta$$

$$\text{❖ Put } \frac{\theta}{2} = t \Rightarrow \theta = 2t$$

$$\text{❖ } d\theta = 2dt$$

x	0	1
θ	0	$\frac{\pi}{2}$

$$\diamondsuit I = \frac{1}{2} \int_0^{\frac{\pi}{2}} (2 \sin t \cos t)^4 \cos^{-4} t \, dt$$

$$\diamondsuit = 8 \int_0^{\frac{\pi}{2}} \sin^4 t \cos^0 t \, dt$$

$$\diamondsuit = 8 \times \frac{1}{2} \beta \left(\frac{4+1}{2}, \frac{0+1}{2} \right)$$

$$\diamondsuit = 4 \times \frac{\left| \frac{5}{2} \right| \left| \frac{1}{2} \right|}{\left| 3 \right|}$$

$$\diamondsuit = 4 \times \frac{3}{2} \times \frac{1}{2} \times \sqrt{\pi} \times \sqrt{\pi} \times \frac{1}{2}$$

$$\diamondsuit = \frac{3\pi}{2}$$