

Module 3

Integration :

Review And Some New Techniques

Sub-Module 3.1

Beta functions with properties

Example:8

❖ Evaluate $\int_0^1 x^4 \cos^{-1} x \, dx$

❖ **Solution:** Integrating by Parts,

$$\text{❖ } I = \cos^{-1} x \frac{x^5}{5} \Big|_0^1 - \int_0^1 \left(\frac{-1}{\sqrt{1-x^2}} \right) \frac{x^5}{5} dx$$

$$\text{❖ } = 0 + \frac{1}{5} \int_0^1 \frac{x^5}{\sqrt{1-x^2}} dx$$

$$\text{❖ Put } x = \sin \theta$$

$$\text{❖ } dx = \cos \theta \, d\theta$$

$$\text{❖ } I = \frac{1}{5} \int_0^{\frac{\pi}{2}} \frac{\sin^5 \theta}{\cos \theta} \cos \theta \, d\theta$$

$$\text{❖ } = \frac{1}{5} \times \frac{1}{2} \times \beta \left(\frac{5+1}{2}, \frac{1}{2} \right) = \frac{1}{10} \times \frac{\left| \frac{3}{2} \right| \frac{1}{2}}{\left| \frac{7}{2} \right|}$$

$$\text{❖ } = \frac{1}{10} \times \frac{2 \times \sqrt{\pi}}{\frac{5}{2} \times \frac{3}{2} \times \frac{1}{2} \times \sqrt{\pi}} = \frac{8}{75}$$

x	0	1
θ	0	$\frac{\pi}{2}$

Example:9

❖ Evaluate $\int_0^{\infty} \frac{x^2}{(1+x^6)^3} dx$

❖ **Solution:** Put $x^3 = t$

❖ $3x^2 dx = dt$

❖ $I = \frac{1}{3} \int_0^{\infty} \frac{dt}{(1+t^2)^3}$

❖ Now Put $t = \tan \theta$

❖ $dt = \sec^2 \theta d\theta$

❖ $I = \frac{1}{3} \int_0^{\frac{\pi}{2}} \frac{\sec^2 \theta d\theta}{(\sec^2 \theta)^3}$

x	0	∞
t	0	∞

x	0	1
θ	0	$\frac{\pi}{2}$

$$\diamondsuit I = \frac{1}{3} \int_0^{\frac{\pi}{2}} \frac{1}{\sec^4 \theta} d\theta$$

$$\diamondsuit = \frac{1}{3} \int_0^{\frac{\pi}{2}} \sin^0 \theta \cos^4 \theta d\theta$$

$$\diamondsuit = \frac{1}{3} \times \frac{1}{2} \beta \left(\frac{1}{2}, \frac{5}{2} \right)$$

$$\diamondsuit = \frac{1}{6} \frac{\left| \frac{1}{2} \right| \left| \frac{5}{2} \right|}{\left| \frac{3}{2} \right|}$$

$$\diamondsuit = \frac{1}{6} \times \sqrt{\pi} \times \frac{3}{2} \times \frac{1}{2} \times \sqrt{\pi} \times \frac{1}{2}$$

$$\diamondsuit = \frac{\pi}{16}$$

Example:10

❖ Evaluate $\int_0^{\frac{\pi}{6}} \cos^3 3\theta \sin^2 6\theta \, d\theta$

❖ **Solution:** Put $3\theta = t \quad \therefore d\theta = \frac{dt}{3}$

❖ When $\theta = 0, t = 0$

❖ When $\theta = \frac{\pi}{6}, t = \frac{\pi}{2}$

❖ $I = \int_0^{\frac{\pi}{2}} \cos^3 t \sin^2 2t \frac{dt}{3}$

❖ $= \frac{1}{3} \int_0^{\frac{\pi}{2}} \cos^3 t (2 \sin t \cos t)^2 \, dt$

❖ $= \frac{4}{3} \int_0^{\frac{\pi}{2}} \cos^3 t \sin^2 t \cos^2 t \, dt$

❖ $= \frac{4}{3} \int_0^{\frac{\pi}{2}} \sin^2 t \cos^5 t \, dt$

❖ $= \frac{4}{3} \times \frac{1}{2} \beta\left(\frac{3}{2}, \frac{6}{2}\right)$

❖ $= \frac{2}{3} \frac{\left|\frac{3}{2}\right|^{\frac{3}{2}}}{\left|\frac{9}{2}\right|^{\frac{9}{2}}} = \frac{\frac{2}{3} \times \frac{1}{2} \sqrt{\pi} \times 2}{\frac{7}{2} \times \frac{5}{2} \times \frac{3}{2} \times \frac{1}{2} \times \sqrt{\pi}} = \frac{32}{315}$

x	0	$\frac{\pi}{6}$
θ	0	$\frac{\pi}{2}$

Example:11

❖ Evaluate $\int_0^{2\pi} \sin^2 \theta (1 + \cos \theta)^4 d\theta$

❖ **Solution:** $I = \int_0^{2\pi} \left(2 \sin \frac{\theta}{2} \cos \frac{\theta}{2}\right)^2 \left(2 \cos^2 \frac{\theta}{2}\right)^4 d\theta$
 $= 64 \int_0^{2\pi} \sin^2 \frac{\theta}{2} \cos^2 \frac{\theta}{2} \cos^8 \frac{\theta}{2} d\theta$
 $= 64 \int_0^{2\pi} \sin^2 \frac{\theta}{2} \cos^{10} \frac{\theta}{2} d\theta$

Put $\frac{\theta}{2} = t \therefore d\theta = 2dt$

$$I = 64 \times 2 \int_0^{\pi} \sin^2 t \cos^{10} t dt$$

If $f(\pi - t) = f(t)$ then $\int_0^a f(t) dt = 2 \int_0^{a/2} f(t) dt$

$$I = 128 \times 2 \int_0^{\frac{\pi}{2}} \sin^2 t \cos^{10} t dt$$

$$= 128 \times 2 \times \frac{1}{2} \beta\left(\frac{3}{2}, \frac{11}{2}\right)$$

$$= 128 \times \frac{\left|\frac{3}{2}\right| \left|\frac{11}{2}\right|}{\left|\frac{7}{2}\right|} = \frac{128 \times \frac{1}{2} \times \sqrt{\pi} \times \frac{9}{2} \times \frac{7}{2} \times \frac{5}{2} \times \frac{3}{2} \times \frac{1}{2} \times \sqrt{\pi}}{6 \times 5 \times 4 \times 3 \times 2} = \frac{21\pi}{8}$$

θ	0	2π
t	0	π

Example:12

❖ Evaluate : $\int_0^{\pi} x \sin^7 x \cos^4 x \, dx$

❖ Hint: $\int_0^a f(x) \, dx = \int_0^a f(a-x) \, dx$

Example:13

❖ Evaluate $\int_0^{\frac{\pi}{2}} \sqrt{\tan \theta} d\theta$ and prove that the result
 $\int_0^{\frac{\pi}{2}} \sqrt{\tan \theta} d\theta \int_0^{\frac{\pi}{2}} \sqrt{\cot \theta} d\theta = \frac{\pi^2}{2}$

❖ **Solution:** Let $I_1 = \int_0^{\frac{\pi}{2}} \sqrt{\tan \theta} d\theta$

$$= \int_0^{\frac{\pi}{2}} \left[\frac{\sin \theta}{\cos \theta} \right]^{\frac{1}{2}} d\theta$$

$$= \int_0^{\frac{\pi}{2}} \sin^{1/2} \theta \cos^{-1/2} \theta d\theta$$

$$= \frac{1}{2} \beta \left(\frac{\frac{1}{2}+1}{2}, \frac{-\frac{1}{2}+1}{2} \right) = \frac{1}{2} \beta \left(\frac{3}{4}, \frac{1}{4} \right)$$

$$= \frac{1}{2} \times \frac{\left| \frac{3}{4} \right| \left| \frac{1}{4} \right|}{\left| 1 \right|} = \frac{1}{2} \left| \frac{1}{4} \right| \left| 1 - \frac{1}{4} \right| = \frac{1}{2} \frac{\pi}{\sin \frac{\pi}{4}} = \frac{\pi}{2 \times \frac{1}{\sqrt{2}}} = \frac{\pi}{\sqrt{2}}$$

$$\begin{aligned}
 \text{Let } I_2 &= \int_0^{\frac{\pi}{2}} \sqrt{\cot \theta} d\theta \\
 &= \int_0^{\frac{\pi}{2}} \left(\frac{\cos \theta}{\sin \theta} \right)^{\frac{1}{2}} d\theta \\
 &= \int_0^{\frac{\pi}{2}} \cos^{\frac{1}{2}} \theta \sin^{-\frac{1}{2}} \theta d\theta \\
 &= \frac{1}{2} \beta \left(\frac{-\frac{1}{2}+1}{2}, \frac{\frac{1}{2}+1}{2} \right) \\
 &= \frac{1}{2} \beta \left(\frac{1}{4}, \frac{3}{4} \right) \\
 &= \frac{1}{2} \left| \frac{1}{4} \right| \left| \frac{3}{4} \right| \\
 &= \frac{1}{2} \frac{\pi}{\sin \frac{\pi}{4}} = \frac{\pi}{\sqrt{2}}
 \end{aligned}$$

$$\text{Hence, } \int_0^{\frac{\pi}{2}} \sqrt{\tan \theta} d\theta \int_0^{\frac{\pi}{2}} \sqrt{\cot \theta} d\theta = \frac{\pi}{\sqrt{2}} \cdot \frac{\pi}{\sqrt{2}} = \frac{\pi^2}{2}$$

Example 14

❖ Express $\int_{-\frac{\pi}{4}}^{\frac{\pi}{4}} (\sin \theta + \cos \theta)^{\frac{1}{3}} d\theta$ as a Gamma function.

Solution: Consider $I = \int_{-\frac{\pi}{4}}^{\frac{\pi}{4}} (\sin \theta + \cos \theta)^{\frac{1}{3}} d\theta$

$$= \int_{-\frac{\pi}{4}}^{\frac{\pi}{4}} \sqrt[3]{2} \left[\sin \theta \frac{1}{\sqrt{2}} + \cos \theta \frac{1}{\sqrt{2}} \right]^{\frac{1}{3}} d\theta$$

$$= \int_{-\frac{\pi}{4}}^{\frac{\pi}{4}} 2^{\frac{1}{6}} \left[\sin \theta \cos \frac{\pi}{4} + \cos \theta \sin \frac{\pi}{4} \right]^{\frac{1}{3}} d\theta$$

$$= \int_{-\frac{\pi}{4}}^{\frac{\pi}{4}} 2^{\frac{1}{6}} \left[\sin \left(\theta + \frac{\pi}{4} \right) \right]^{\frac{1}{3}} d\theta$$

Put $\frac{\pi}{4} + \theta = t \rightarrow d\theta = dt$

When $\theta = -\frac{\pi}{4}$, $t = 0$

When $\theta = \frac{\pi}{4}$, $t = \frac{\pi}{2}$

$$\begin{aligned}
 &= \int_0^{\frac{\pi}{2}} 2^{\frac{1}{6}} [\sin t]^{\frac{1}{3}} dt \\
 &= 2^{\frac{1}{6}} \int_0^{\frac{\pi}{2}} \sin^{\frac{1}{3}} t \cos^0 t \, dt \\
 &= 2^{\frac{1}{6}} \frac{1}{2} \beta \left(\frac{\frac{1}{3}+1}{2}, \frac{1}{2} \right) \\
 &= 2^{\frac{1}{6}-1} \beta \left(\frac{2}{3}, \frac{1}{2} \right) = 2^{-\frac{5}{6}} \frac{\left| \frac{2}{3} \right| \left| \frac{1}{2} \right|}{\left| \frac{7}{6} \right|} \\
 &= 2^{\frac{1}{5/6}} \frac{\left| \frac{2}{3} \right| \sqrt{\pi}}{\left| \frac{7}{6} \right|}
 \end{aligned}$$