

Week 5 - Lecture 27

Strong Law of large numbers

- If we take sufficient samples then we can estimate the true mean

Assumption

Assumption
 \hookrightarrow The samples drawn should be independent from past pulls of the arm or pulls from other arms. (i.i.d.)

We need to know how many samples to pick before getting a good estimate of the arm, how quickly can I reach this value, since we have limited pulls.

Concentration of Measures

Let X_1, X_2, \dots, X_n be i.i.d. random variables
with mean $\mu = \mathbb{E}[X_i]$ and $\sigma^2 = \text{Var}[X_i]$
 $\quad \quad \quad \nwarrow$ common mean $\quad \quad \quad \nwarrow$ common variance

Natural estimator :

$$\hat{\mu} = \frac{1}{n} \sum_{i=1}^n x_i \quad E[\hat{\mu}] = \mu \quad \left[\begin{array}{l} \text{unbiased} \\ \text{estimator} \end{array} \right]$$

$$\text{Var} [\hat{\mu}] = \mathbb{E} [(\hat{\mu} - \mu)^2]$$

$$= \mathbb{E} \left[\left(\frac{1}{n} \sum_{i=1}^n x_i - \mu \right)^2 \right]$$


✓ This is true for any n

$$= \frac{1}{n^2} \mathbb{E} \left[\left(\sum_{i=1}^n (x_i - \mu) \right)^2 \right]$$

$$= \frac{1}{n^2} \sum_{i=1}^n \mathbb{E} \left[(x_i - \mu)^2 \right]$$

$$\text{Var}[\hat{\mu}] = \frac{\sigma^2}{n} \quad \leftarrow \text{as } n \text{ increases } \text{Var}[\hat{\mu}] \text{ decreases, error reduces with increase in number of samples}$$

Tail Probabilities



$$\text{Pr}\{|\hat{\mu} - \mu| \geq \epsilon\} \rightarrow \text{two-sided tail probability}$$

We want the tail probabilities to be small and figure out how it depends on n

$$\text{Pr}\{|\hat{\mu} - \mu| < \epsilon\} \rightarrow \text{how many samples do we need to collect so that the estimate lies in the epsilon region.}$$

Thus we need to bound these.

Inequalities

1) Markov inequality

$$\Pr \{ |x| \geq \varepsilon \} \leq \frac{\mathbb{E}[|x|]}{\varepsilon}$$

2) Chebychev inequality

$$\Pr \{ |x - \mathbb{E}[x]| \geq \varepsilon \} \leq \frac{\text{Var}(x)}{\varepsilon^2}$$

Now,

$$x = \hat{\mu} = \frac{1}{n} \sum_{i=1}^n x_i$$

$$\text{Var}(\hat{\mu}) = \frac{\sigma^2}{n\varepsilon^2}$$

← with more samples
the difference between
 μ and $\mathbb{E}[\mu]$ will reduce

Now for a fixed n , if ε is small, then the

$\frac{\sigma^2}{n\varepsilon^2}$ will become larger, then Probability will be
smaller

Central Limit Theorem

$x_1, x_2, \dots, x_n \rightarrow \text{i.i.d. sequence}$

$$\frac{\sum_{i=1}^n x_i - n\mu}{\sqrt{n\sigma^2}} \xrightarrow{d} N(0,1)$$

If $n \rightarrow \infty$ then average will converge to normal distribution asymptotically

$$\Pr \{ \hat{\mu} - \mu \geq \varepsilon \}$$

$$= \Pr \left\{ \frac{1}{n} \sum x_i - \mu \geq \varepsilon \right\}$$

$$= \Pr \left\{ \frac{\sum x_i - n\mu}{\sqrt{n\sigma^2}} \geq \frac{\varepsilon n}{\sqrt{n\sigma^2}} \right\}$$

$$= \Pr \left\{ \underbrace{\frac{\sum x_i - n\mu}{\sqrt{n\sigma^2}}}_{\text{PDF for } N(0,1)} \geq \frac{\varepsilon \sqrt{n}}{\sqrt{\sigma^2}} \right\}$$

If n is sufficiently large then this will converge to $N(0,1)$ (CLT)

$$\approx \int_{\frac{\varepsilon \sqrt{n}}{\sqrt{\sigma^2}}}^{\infty} \underbrace{\frac{1}{\sqrt{2\pi}} \exp \left\{ -\frac{x^2}{2} \right\}}_{\text{PDF for } N(0,1)} dx$$

This integration does not have closed form, thus need

$$\left[\text{Let } u = \frac{\varepsilon \sqrt{n}}{\sqrt{\sigma^2}} \right]$$

to look at bounds

$$\leq \frac{1}{\sqrt{2\pi}} \int_u \frac{x}{u} \exp\left\{-\frac{x^2}{2}\right\} dx$$

$$= \sqrt{\frac{\sigma^2}{2\pi n \varepsilon^2}} \exp\left(\frac{-n \varepsilon^2}{2\sigma^2}\right)$$

Thus $\Pr\{\hat{\mu} - \mu \geq \varepsilon\}$ is decaying

exponentially with n , (with CLT),

with Chebychev it was inversely proportional to n .

But assuming that n is sufficiently large.