

# ADVERSE DRUG REACTION IDENTIFICATION REPORT

## Task 1: ADR Identification Basics

**Project Title:** Identification and Analysis of Suspected Amoxicillin-Induced Erythema Multiforme

**Prepared by:** Dhruv Gupta

**Course:** Bachelor of Pharmacy

**Internship Task:** ADR Identification Basics

---

### 1. Introduction

An Adverse Drug Reaction (ADR) is an unintended and harmful response associated with the use of a medicinal product.

Pharmacovigilance involves the detection, assessment, understanding, and prevention of adverse effects or other medicine-related problems.

### 2. Patient Case Summary

The selected published case describes a 14-year-old male patient who developed severe erythema multiforme during treatment involving multiple medications, including amoxicillin.

The clinical investigation considered both infectious and drug-related causes.

### 3. Suspected Drug

**Drug:** Amoxicillin

**Drug class:** Aminopenicillin antibiotic

**Therapeutic use:** Treatment of susceptible bacterial infections.

### 4. Suspected Adverse Drug Reaction

**Reported reaction:** Erythema multiforme (EM)

Erythema multiforme is an inflammatory skin condition characterized by distinctive skin lesions, which may sometimes involve mucous membranes.

### 5. Clinical Timeline

| Parameter   | Case information |
|-------------|------------------|
| Patient age | 14 years         |

|                          |                                         |
|--------------------------|-----------------------------------------|
| Gender                   | Male                                    |
| Primary condition        | Lymphadenitis                           |
| Suspected drug           | Amoxicillin                             |
| Adverse reaction         | Severe erythema multiforme              |
| Other possible factors   | Infection and other medications         |
| Diagnostic investigation | Skin biopsy and allergy-related testing |

## **6. ADR Assessment**

### **A. Temporal Relationship**

The skin reaction developed during the course of medication treatment.

### **B. Dechallenge**

The case report should be reviewed for the documented response following withdrawal of the suspected medication. Do not assume improvement unless reported.

### **C. Rechallenge**

A rechallenge should not be assumed to have occurred. Record it as not reported if the article does not document one.

### **D. Alternative Causes**

The investigation considered Coxsackie A6 infection and a possible reaction to another medication.

### **E. Causality Assessment**

The published investigators considered amoxicillin-induced drug eruption based on the clinical pattern and diagnostic investigations.

For this educational analysis, the association should be described as suspected and supported by the reported findings, rather than independently assigning a definitive causality category.

## **7. Severity and Outcome**

The case involved severe erythema multiforme. Record the clinical outcome and seriousness criteria according to the original publication.

## **8. Conclusion**

This case demonstrates the importance of evaluating medication history, symptom onset, clinical presentation, and alternative causes when identifying suspected adverse drug reactions.

A structured pharmacovigilance assessment can support early recognition of drug-related harm and improve medication safety.

## **9. References**

Kurihara M, Yamanishi S, Ozaki S, Pawankar R. A 14-year-old boy with severe erythema multiforme due to amoxicillin. *Asia Pacific Allergy*. 2023;13(3):135–138.

DOI: 10.5415/apallergy.0000000000000108

PubMed: <https://pubmed.ncbi.nlm.nih.gov/37744962/>