

## COMPARISON OF LIMB LENGTH DISCREPANCY BETWEEN UNILATERAL AND BILATERAL T.K.R.

ALPA PUROHIT<sup>1</sup>

1. Vice-Principal, AIMS-Physiotherapy College, Ahmedabad, Gujarat

### Abstract

**AIMS & OBJECTIVES:** To find out the difference of leg length in unilateral & bilateral TKR & find out clinical management of leg length discrepancy for improving patients function.

**Purpose of the Study:** To find out the difference of leg length between unilateral v/s bilateral T.K.R. and to find out the patients level of discomfort during walking.

**METHODOLOGY: STUDY DESIGN:** cross sectional observational study. **Sample Population and Sample Size-** 30 patients included according to criteria. In Group –A-15, unilateral Total knee replacement patients & in Group B - 15, bilateral Total knee replacement patients were included. **Sampling Technique:** Convenient sampling according to selection criteria.

**OUTCOME MEASURES:** Limb length was measured by using Tape measurement method (TMM) in centimeters.

**DATA COLLECTION & ANALYSIS:** Data were collected from sterling hospital, Ahmedabad at post 3 months of surgery. Limb length measurement was done in the supine position with pelvis squared. The lower limbs were placed parallel to the body and limb length measurement (in centimeters) done from the anterior superior iliac spine to the medial malleolus using a measure tape. The measurement was taken twice by two different observers and the mean of the two values was recorded. Outcome parameters were analyzed with t-test in both group.

**RESULTS:** Data was analyzed by using Paired t-test. The t-value for unilateral T.K.R. was 9.73 ( $p < 0.001$ ) highly significant and bilateral T.K.R was 3.58 ( $p < 0.05$ ) respectively.

**CONCLUSION:** In unilateral T.K.R. there is a significant difference in leg length between operated & non-operated leg & in bilateral. There is no much leg length discrepancy between both operated legs. So in unilateral T.K.R., Limb length discrepancy is due to other leg's arthritis condition which may cause imbalance in gait.

**KEYWORDS:** limb length discrepancy; unilateral- bilateral T.K.R.

## INTRODUCTION

Leg-length inequality, defined as a difference in lengths of the 2 legs, is very common, occurring in up to 70% of the population<sup>1</sup>.

Improved surgical techniques and rehabilitation protocols have resulted in excellent knee function and range of motion following total knee arthroplasty<sup>2</sup>.

Limb length discrepancy and its effects on patient function have been discussed in depth in the literature with respect to hip arthroplasty but there are few studies that have examined the effect on function of limb length discrepancy following total knee arthroplasty (TKA)<sup>3,4</sup>.

Nevertheless, there remain 15-20% of patients with persistent dysfunction that is difficult to treat<sup>5-7</sup>.

Although problems after total knee arthroplasty are frequently linked to prosthetic mal-alignment, radiographic loosening, and comorbidities, some cases are related to functional problems that are less evident clinically and/or radiographically.

Functional problems following total knee arthroplasty may be incapacitating as a result of persistent pain<sup>8</sup>, instability<sup>9</sup>, and limited range of motion<sup>10,11</sup>.

Patients who experienced more pain and functional impairment after total knee arthroplasty were less likely to be satisfied with the procedure.

## AIMS & OBJECTIVES

- To find out the difference of leg length in unilateral & bilateral TKR.
- Find out solution of leg length discrepancy for improving patients function & satisfaction.

## METHODOLOGY

Sample size

Group A: 15, Unilateral Total knee replacement patients

Group B: 15, Bilateral Total knee replacement patients

## STUDY DESIGN

Cross sectional observational study

## INCLUSION CRITERIA

- Patients were operated with midline incision with primary T.K.R.
- Age-group 50-70yrs
- Post op 3 month

- Post op same exercises program in both group
- In both group pre op FFD 10-15degree & 8-10 degree varus deformity
- No anymal-alignment, component malposition, loosening, or patellar maltracking
- No hamstring tightness
- No any hip joint pathology & SI joint dysfunction

### EXCLUSION CRITERIA

- Other surgical technique
- Other than high flex knee joint
- Other than midline incision
- Post op less than 3 month
- Any mal-alignment, component malposition, loosening, or patellar maltracking
- Hamstring tightness.
- Any hip joint pathology & SI dysfunction & postural deviation.
- Marked abductor weakness.
- Pre-operative limb length discrepancy

### METHODS

Data were collected from sterling hospital, Ahmedabad at post 3 months of surgery.

Limb length measurement was done in the supine position with pelvis squared.

The lower limbs were placed parallel to the body and limb length measurement (in centimeters) done from the anterior superior iliac spine to the medial malleolus using a measure tape<sup>12</sup>.

The measurement was taken twice by two different observers and the mean of the two values was recorded.

### OUTCOME MEASURES

Limb length was measured by using Tape measurement method (TMM) in centimeters<sup>13,14</sup>.

### RESULTS

TABLE 1: VALUE OF UNILATERAL AND BILATERAL T.K.R.LEG LENGTH

Unilateral	Average Length	Bilateral	Average Length
Limb-1(Operated)	49.8	Limb-1(Operated)	47.2
Limb-2(Non-Operated)	47.4	Limb-2(Operated)	47.26




FIGURE 1: COMPARISION OF UNILATERAL AND BILATERAL T.K.R.LEG LENGTH

### DISCUSSION

A persistent limp is one of the most frustrating symptoms after total knee arthroplasty.

There are many causes of limp; however, leg length discrepancy is one of the common reason for litigation after successful total knee arthroplasty<sup>3,4,12</sup>.

Leg-length inequality may be an important risk Factor for the longer leg exhibited increased ground-reaction forces and therefore should be at greater risk for osteoarthritis.

Limb length discrepancy affects functional outcome after total knee arthroplasty<sup>7,15</sup>.

Bhave et al.<sup>7</sup> showed that the operated leg gained length compared to the contralateral un-operated leg due to correction of the varus deformity.

A number of factors may be responsible for the limb length discrepancy, including correction of the varus alignment after surgery, the amount of preoperative flexion deformity, and the postoperative flexion deformity

In present study it was obvious difference in both leg length in unilateral knee replacement patients who having limp during walking & there is no as such any major difference between limb length bilateral knee replacement.

### CONCLUSION

In unilateral T.K.R. there is a significant difference in leg length between operated & non-operated leg & in bilateral there is no much leg length discrepancy between both operated legs.

So in unilateral T.K.R., Limb length discrepancy is due to other leg's arthritis condition which may cause imbalance in gait.

### CLINICAL IMPLICATION

We can give sole compensation for correcting LLD & improving function.

## LIMITATIONS

- Sample size is small.
- Measurements of leg length, are subject to measurement error, which could result in misclassification.

## FUTURE STUDY

- Outcome measure for functional activity
- Comparison of LLD between unilateral operated patient & those same operated for second time knee replacement

## REFERENCES

1. Harvey WF, Yang M, Cooke TD, Segal NA, Lane N, Lewis CE, Felson DT. Association of leg-length inequality with knee osteoarthritis: a cohort study. *Annals of internal medicine*. 2010 Mar 2;152(5):287-95.
2. Vaidya SV, Patel MR, Panghate AN, Rathod PA. Total knee arthroplasty: limb length discrepancy and functional outcome. *Indian journal of orthopaedics*. 2010 Jul 1;44(3):300.
3. Austin MS, Hozack WJ, Sharkey PF, Rothman RH. Stability and leg length equality in total hip arthroplasty. *The Journal of arthroplasty*. 2003 Apr 30;18(3):88-90.
4. Attarian DE, Vail TP. Medicolegal aspects of hip and knee arthroplasty. *Clinical orthopaedics and related research*. 2005 Apr 1;433:72-6.
5. Jane K, Charles LN, Paul AL. Stiffness after total knee arthroplasty: prevalence of the complication and outcomes of debridement. *Journal of Bone and Joint Surgery*. 2004;86(7):1479.
6. Ranawat CS, Ranawat AS, Mehta A. Total knee arthroplasty rehabilitation protocol: What makes the difference?. *The Journal of arthroplasty*. 2003 Apr 30;18(3):27-30.
7. Bhave A, Mont M, Tennis S, Nickey M, Starr R, Etienne G. Functional problems and treatment solutions after total hip and knee joint arthroplasty. *J Bone Joint Surg Am*. 2005 Dec 1;87(suppl 2):9-21.
8. Dahlen L, Zimmerman L, Barron C. Pain perception and its relation to functional status post total knee arthroplasty: a pilot study. *Orthopaedic Nursing*. 2006 Jul 1;25(4):264-70.
9. Gonzalez MH, Mekhail AO. The failed total knee arthroplasty: evaluation and etiology. *Journal of the American Academy of Orthopaedic Surgeons*. 2004 Nov 1;12(6):436-46.
10. Pariente GM, Lombardi Jr AV, Berend KR, Mallory TH, Adams JB. Manipulation with prolonged epidural analgesia for treatment of TKA complicated by arthrofibrosis. *Surgical technology international*. 2005 Dec;15:221-4.
11. Padua R, Ceccarelli E, Bondi R, Campi A, Padua L. Range of motion correlates with patient perception of TKA outcome. *Clinical orthopaedics and related research*. 2007 Jul 1;460:174-7.
12. Iagulli ND, Mallory TH, Berend KR, Lombardi Jr AV, Russell JH, Adams JB, Groseth KL. A simple and accurate method for determining leg length in primary total hip arthroplasty. *American journal of orthopedics (Belle Mead, NJ)*. 2006 Oct;35(10):455-7.
13. EWALD FC. The Knee Society total knee arthroplasty roentgenographic evaluation and scoring system. *Clinical orthopaedics and related research*. 1989 Nov 1;248:9-12.
14. Edeen J, Sharkey PF, Alexander AH. Clinical significance of leg-length inequality after total hip arthroplasty. *American journal of orthopedics (Belle Mead, NJ)*. 1995 Apr;24(4):347-51.
15. Bhave A, Marker DR, Seyler TM, Ulrich SD, Plate JF, Mont MA. Functional problems and treatment solutions after total hip arthroplasty. *The Journal of arthroplasty*. 2007 Sep 30;22(6):116-24.