

# Supplementary Material for “Engineering momentum profiles of cold-atom beams”

D Hudson Smith<sup>1</sup> and Artem G Volosniev<sup>2</sup>

<sup>1</sup>*Clemson University, Clemson, South Carolina 29634, USA*

<sup>2</sup>*Institut für Kernphysik, Technische Universität Darmstadt, 64289 Darmstadt, Germany*

(Dated: March 27, 2019)

# IMPURITY IN A BOSE GAS

To model one impurity atom that moves through a one-dimensional environment made of  $N$  cold bosonic atoms, we employ the following Hamiltonian

$$H = -\frac{\hbar^2}{2m} \sum_{i=1}^N \frac{\partial^2}{\partial x_i^2} - \frac{\hbar^2}{2M} \frac{\partial^2}{\partial y^2} + \lambda \sum_{i>j=1}^N \delta(x_i - x_j) + g \sum_{i=1}^N \delta(x_i - y), \quad (1)$$

where  $M$  is the mass of the impurity atom, and  $m$  is the mass of a bosonic particle. The position of the impurity is  $y$ , bosons are at the coordinates  $\{x_i\}$ . We assume that the realistic boson-boson and boson-impurity interactions are well-described by the zero-range potentials of strengths  $\lambda$  and  $g$  respectively. The environment is large by assumption. To describe it, the periodic boundary conditions are used: The particles move in a ring of the circumference  $L$ , such that  $0 < x_i < L$  and  $0 < y < L$ . We are interested in the thermodynamic limit:  $N, L \rightarrow \infty$  with a fixed value of the density  $\rho = \frac{N}{L}$ .

If the system is non-interacting ( $\lambda = g = 0$ ), the eigenstates are written as  $e^{2\pi i \frac{n_1 x_1 + \dots + n_N x_N + m y}{L}}$ , where  $n_1, \dots, n_N$  and  $m$  are arbitrary integers. For non-vanishing interactions we use these functions to write an eigenfunction of the Hamiltonian as  $\Psi = \sum_{\{n_j\}, m} a_{\{n_j\}, m} e^{2\pi i \frac{\sum n_j x_j + m y}{L}}$ . Because all interactions are pairwise, the total (angular) momentum of the system must be conserved, and we write it as  $\mathcal{P} = \frac{2\pi\hbar}{L} \left( \sum_j n_j + m \right)$ . A conserved quantity ( $\mathcal{P}$ ) allows us to exclude one variable from the consideration. We write the function  $\Psi$  as  $\Psi = e^{i \frac{\mathcal{P} y}{\hbar}} \sum_{\{n_j\}, m} a_{\{n_j\}, m} e^{2\pi i \frac{\sum n_j x_j}{L}} \equiv e^{i \frac{\mathcal{P} y}{\hbar}} \psi(z_1, \dots, z_N)$  with  $z_i = L\theta(y - x_i) + x_i - y$ , where  $\theta(x)$  is the Heaviside step function, i.e.,  $\theta(x > 0) = 1$  and zero otherwise. The variables  $z_i$  are defined such that  $0 \leq z_i \leq L$  and the impurity is placed at  $z = 0$  ( $z = L$ ). Now if we insert this function into the Schrödinger equation,  $H\Psi = E\Psi$ , we obtain the following equation for  $\psi$  ( $0 < z_i < L$ )

$$-\frac{\hbar^2}{2m} \sum_i \frac{\partial^2 \psi}{\partial z_i^2} - \frac{\hbar^2}{2M} \left( \sum_i \frac{\partial}{\partial z_i} \right)^2 \psi + i \frac{\hbar \mathcal{P}}{M} \sum_i \frac{\partial \psi}{\partial z_i} + \lambda \sum_{i>j} \delta(z_i - z_j) \psi = \left( E - \frac{\mathcal{P}^2}{2M} \right) \psi, \quad (2)$$

which must be supplemented with the boundary conditions:

$$\psi(z_i = 0) = \psi(z_i = L); \quad \left. \frac{\partial \psi}{\partial z_i} \right|_{z_i=L^-}^{z_i=0^+} = \frac{2g\kappa}{\hbar^2} \psi(z_i = 0), \quad (3)$$

where  $\kappa = mM/(m + M)$  is the reduced mass.

By assumption the bosons interact weakly, such that the ansatz  $\psi = \prod_i \Phi(z_i)$  can be used to approximate the system. To minimize the expectation value of the Hamiltonian the function  $\Phi(z)$  must satisfy the following non-linear Schrödinger equation

$$-\frac{\hbar^2}{2\kappa} \frac{\partial^2 \Phi}{\partial z^2} + i \frac{\hbar \mathcal{P}}{M} \frac{\partial \Phi}{\partial z} - i \frac{\hbar^2 (N-1) A}{M} \frac{\partial \Phi}{\partial z} + \lambda (N-1) |\Phi|^2 \Phi = \mu \Phi, \quad (4)$$

where  $A = -i \int \Phi(x)^* \frac{\partial}{\partial x} \Phi(x) dx$  defines the momentum of a boson, and  $\mu$  is the Lagrange multiplier. We rewrite this equation as

$$-\frac{\partial^2 \Phi}{\partial z^2} + i v \frac{\partial \Phi}{\partial z} + \tilde{\lambda} (N-1) |\Phi|^2 \Phi = \tilde{\mu} \Phi, \quad (5)$$

where  $\tilde{\mu} = \frac{2\kappa\mu}{\hbar^2}$ ,  $\tilde{\lambda} = \frac{2\kappa\lambda}{\hbar^2}$ , and  $v = \frac{2\kappa P}{M\hbar}$  with  $P = \mathcal{P} - \hbar A(N-1)$ .  $P$  defines the momentum of the impurity in the thermodynamic limit; note that because  $A$  is determined by  $P$ , there is a unique value of  $\mathcal{P}$  for a given  $P$ . The boundary conditions for Eq. (5) read

$$\Phi(z = 0) = \Phi(z = L); \quad \left. \frac{\partial \Phi}{\partial z} \right|_{z=L^-}^{z=0^+} = \tilde{g} \Phi(0), \quad (6)$$

where  $\tilde{g} = \frac{2\kappa g}{\hbar^2}$ . The non-linear equation (5) has an analytic steady solution [1], which determines the properties of the dressed impurity in our problem. Let us first consider the non-interacting impurity  $g = 0$ . In this case the solution




FIG. 1. Panel **a)**: The density,  $|\Phi|^2 N$ , of the Bose gas for  $z_0 = a$  (red curve),  $z_0 = -a$  (blue curve) ( $a > 0$ ). Note that the minimum of the density is at  $-a$ . The shaded area is a combination of the two solutions with the singularity at  $z = 0$ . Panel **b)**: The phase,  $\phi$ , of the Bose gas for the densities from **a)**.

for  $v > 0$  is [2, 3]

$$\Phi = \sqrt{\frac{\tilde{\mu}}{\tilde{\lambda}(N-1)}} \left( 1 - \beta \operatorname{sech}^2 \left[ \sqrt{\frac{\tilde{\mu}\beta}{2}}(z + z_0) \right] \right)^{\frac{1}{2}} e^{i\phi(z)}, \quad (7)$$

$$\phi(z) = -\pi\theta(z + z_d) + \arctan \left( \frac{\sqrt{\frac{2v^2}{\tilde{\mu}}}\beta}{\exp[\sqrt{2\tilde{\mu}\beta}(z + z_0)] - 2\beta + 1} \right), \quad (8)$$

where  $\beta = 1 - v^2/(2\tilde{\mu})$ ,  $z_0$  is some parameter that determines the origin and  $z_d$  is the point where  $\arctan$  reaches  $\pi/2$ . It is worthwhile noting that the solution for  $v < 0$  is  $\Phi^*$ . The solution from Eqs. (7) and (8) is plotted in Fig. 1; for simplicity it is plotted in the interval  $-L/2 < z < L/2$ , the region  $0 < z < L$  easily follows.

To describe an interacting impurity, we combine two moving solitons with  $\pm z_0$ , which creates a singularity at  $z = 0$  [1, 4]. Therefore, a dressed impurity in our model is a topological defect with a dissipationless propagation. We write the corresponding ‘wave function’ as

$$\Phi = \sqrt{\frac{\tilde{\mu}}{\tilde{\lambda}(N-1)}} \left( 1 - \beta \operatorname{sech}^2 \left[ \sqrt{\frac{\tilde{\mu}\beta}{2}}(z \pm z_0) \right] \right)^{\frac{1}{2}} e^{i\phi(z)}, \quad (9)$$

with

$$\phi(z) = \delta\phi\theta(-z) + \arctan \left( \frac{\sqrt{\frac{2v^2}{\tilde{\mu}}}\beta}{\exp[\sqrt{2\tilde{\mu}\beta}(z \pm z_0)] - 2\beta + 1} \right), \quad (10)$$

where  $z_0 > 0$  is discussed below, the parameter  $\delta\phi$  is not important for the further derivations, it reassures that the phase is a continuous function; the plus sign in  $\pm$  corresponds to  $z > 0$  and the minus sign to  $z < 0$ . This function is illustrated in Fig. 1. The density has a non-analytic derivative at  $z = 0$ . The phase is a continuous function at  $z = 0$  (its derivative is also continuous). Note that the wave function is not periodic (see Eq. (10)). This non-periodicity is not important for our discussion, because we are interested in the behavior of the bosons close to the impurity. It suggests that a grey soliton must be formed upon a change of interaction parameters to take care of the phase slip.

The parameter  $\tilde{\mu}$  is found from the normalization condition  $\int \Phi^2 = 1$ . For  $N \rightarrow \infty$ , we obtain

$$\tilde{\mu} = \gamma\rho^2 \frac{N-1}{N} \left( 1 - 2\sqrt{2\beta_0} \frac{(\tanh(d) - 1)}{\sqrt{\gamma}N} \right), \quad (11)$$

where  $\gamma = \tilde{\lambda}/\rho$ ,  $\rho = N/L$ ,  $\beta_0 = 1 - v^2/(2\gamma\rho^2)$ , and  $d = \sqrt{\frac{\gamma\beta_0}{2}}\rho z_0$ . The equation to determine  $z_0$  is found by using the boundary conditions at  $z = \{0, L\}$

$$\frac{\tilde{g}}{\rho\sqrt{2\gamma}} = \frac{\beta_0^{\frac{3}{2}} \tanh(d)}{-\beta_0 + \cosh^2(d)}. \quad (12)$$

This equation is cubic (in  $\tanh(d)$ ), hence, the solutions can be found in a closed form. There are three solutions. However, only two will lead to the acceptable values of  $z_0$ . We will refer to these steady solutions as the ‘polaron’ and the ‘polaron-soliton’ pair, because in the limit  $g \rightarrow 0$  the former corresponds to the ground state, and the latter to a gray soliton. The ‘polaron-soliton’ pair is expected to be unstable (small perturbations will lead to a decay of this steady solution [1]), therefore, we do not consider it. The solutions merge for  $z_m$

$$\tanh^2\left(\sqrt{\frac{\gamma\beta_0}{2}}\rho z_m\right) = \frac{\sqrt{1 + \frac{4v_c^2}{\gamma\rho^2}} - (1 + \frac{v_c^2}{\gamma\rho^2})}{2\beta_0}, \quad (13)$$

which is derived by taking a derivative of Eq. (12) with respect to  $z_0$  and equating the resulting expression to zero – this determines the maximum value of  $g$  for which (for a fixed  $\beta_0$ ) there is a steady solution. Equations (12) and (13) give the equation for the critical value of  $v_c$ :

$$\frac{\tilde{g}}{\rho\sqrt{\gamma}} = \frac{3 - \sqrt{1 + \frac{4v_c^2}{\gamma\rho^2}}}{-1 + \sqrt{1 + \frac{4v_c^2}{\gamma\rho^2}}} \sqrt{\sqrt{1 + \frac{4v_c^2}{\gamma\rho^2}} - 1 - \frac{v_c^2}{\gamma\rho^2}}. \quad (14)$$

For  $v > v_c$  (see Fig. 1 of the main text) there is no steady solutions.

Now we can calculate the energy of the dressed impurity in the thermodynamic limit

$$\mathcal{E} \equiv \lim_{N \rightarrow \infty, \frac{N}{L} \rightarrow \rho} [E(c, P) - E(c = 0, P = 0)], \quad (15)$$

where

$$E(c, P) = \frac{P^2}{2M} + \mu N - \frac{\hbar^2 A^2 N(N-1)}{2M} - gN(N-1) \int_0^{L/2} |\Phi|^4 dz. \quad (16)$$

Using these expressions we derive

$$\mathcal{E} = \frac{P_I^2}{2M} + \frac{\hbar^2 \rho}{2\kappa} \frac{\sqrt{2\tilde{g}\rho\beta}}{3} [4b + (-4b + \beta \text{sech}^2(d)) \tanh(d)] + \frac{\hbar P_I}{M} \lim_{N \rightarrow \infty} AN, \quad (17)$$

where  $b = 1 + \frac{v^2}{4\tilde{g}\rho} = 1 + \frac{\kappa P_I^2}{2M^2 g \rho}$ . This energy for  $v \rightarrow 0$  can be written as

$$\mathcal{E} \simeq \epsilon + \frac{P_I^2}{2m_{\text{eff}}}, \quad (18)$$

where  $\epsilon$  is the effective energy of the dressed impurity, and  $m_{\text{eff}}$  is the effective mass.

The parameters  $m_{\text{eff}}$  and  $\epsilon$  calculated using the non-linear Schrödinger equation agree well with the results in the literature [5]. This agreement supports methods for our discussion. Still, further work is required to understand properties of a dressed impurity in our formalism. First of all, it will be interesting to investigate the critical momentum, which is supersonic in the employed model for  $g \rightarrow 0$  and for not heavy impurities. Indeed, the model we solve may have a different speed of sound, because it is equivalent to a heavy impurity in a gas of bosons with a mass  $\kappa$ . It will also be interesting to calculate the residue – the overlap between wave functions that describe a state with  $g = 0$  and an interacting state. In the present model, the impurity changes the order parameter only locally, which means a non-zero residue; see [6] for  $P = 0$ . It will be interesting to calculate the overlap using an exactly solvable model, e.g., a heavy impenetrable impurity in a Bose gas (solvable by Bethe ansatz) to check this result.

# BOUNDARY-AUGMENTED COST FUNCTION

Equation (6) shows the boundary-augmented cost function which includes a term that increases the cost for link-potential solutions that extend beyond the support region  $x \in [-x_0, x_0]$ . This added terms is

$$J_{\text{boundary}} = \alpha \sum_i^N J_{\text{boundary}}^i, \quad (19)$$

where

$$J_{\text{boundary}}^i = \int_{|x| > x_0} dx |V_i(x; A_i, \mu_i, \sigma_i)|^2. \quad (20)$$

Assuming the Gaussian potential form as in Eq. (5), this evaluates to

$$J_{\text{boundary}}^i = \frac{\sqrt{\pi}}{2} A_i^2 \sigma_i \left[ \text{erfc} \left( \frac{x_0 + \mu_i}{\sigma_i} \right) + \text{erfc} \left( \frac{x_0 - \mu_i}{\sigma_i} \right) \right], \quad (21)$$

where erfc is the complementary error function.

- 
- [1] V. Hakim, Phys. Rev. E **55**, 2835 (1997).
  - [2] T. Tsuzuki, Journal of Low Temperature Physics **4**, 441 (1971).
  - [3] M. Ishikawa and H. Takayama, Journal of the Physical Society of Japan **49**, 1242 (1980).
  - [4] M. Schecter, D. Gangardt, and A. Kamenev, New J. Phys. **18**, 065002 (2016).
  - [5] S. I. Mistakidis, A. G. Volosniev, N. T. Zinner, and P. Schmelcher, arXiv:1809.01889 (2018).
  - [6] A. G. Volosniev and H.-W. Hammer, Phys. Rev. A **96**, 031601 (R) (2017).