

A. G. Volosniev Institut für Kernphysik, Technische Universität Darmstadt, 64289 Darmstadt, Germany

D. H. Smith Department of Physics, The Ohio State University, Columbus, OH 43210, USA

## **Impenetrability in Floquet scattering in one dimension**

A. G. Volosniev D. H. Smith

(Dated: Received: date / Accepted: date)




FIG. 1. Panel **a**): The reflection probability  $|B_0|^2$  as a function of  $p/\sqrt{\omega}$  for  $g_1/\sqrt{\omega} = 0.2$  – black solid curve,  $g_1/\sqrt{\omega} = 0.4$  – red dashed curve,  $g_1/\sqrt{\omega} = 0.8$  – green dot-dashed curve, and  $g_1/\sqrt{\omega} = 1.0$  – blue dotted curve. For all curves  $g_0/\sqrt{\omega} = -1$ . Panel **b**):  $|B_0|^2$  for  $g_1/\sqrt{\omega} = 1.5$  – black solid curve,  $g_1/\sqrt{\omega} = 2.5$  – red dashed curve,  $g_1/\sqrt{\omega} = 3.5$  – green dot-dashed curve, and  $g_1/\sqrt{\omega} = 4.5$  – blue dotted curve. For all curves  $g_0 = 0$ . Panel **c**):  $|B_0|^2$  for  $g_0/\sqrt{\omega} = 0.1$  – black solid curve,  $g_0/\sqrt{\omega} = 0.5$  – red dashed curve,  $g_0/\sqrt{\omega} = 1$  – green dot-dashed curve, and  $g_0/\sqrt{\omega} = 1.5$  – blue dotted curve. For all curves  $g_1/\sqrt{\omega} = 1.5$ . Panel **d**):  $|B_0|^2$  for  $g_0/\sqrt{\omega} = -0.1$  – black solid curve,  $g_0/\sqrt{\omega} = -0.8$  – red dashed curve,  $g_0/\sqrt{\omega} = -1.4$  – green dot-dashed curve, and  $g_0/\sqrt{\omega} = -2$  – blue dotted curve. For all curves  $g_1/\sqrt{\omega} = 1.5$ .

## I. INTRODUCTION

## II. SCATTERING OFF A TIME-PERIODIC ZERO-RANGE POTENTIAL IN ONE DIMENSION

## III. CONCLUSIONS

**Acknowledgments** A. G. V. gratefully acknowledges the support of the Humboldt Foundation.

## Appendix A: Appendix

### 1. Floquet Formalism

**Scattering in time-dependent picture.** Here, for convenience of the reader, we discuss the time-dependent Schrödinger equation with a general short-range time-periodic potential

$$i\frac{\partial}{\partial t}\Psi(x,t) = H(t)\Psi(x,t), \quad H(t) = H_0 + W(x,t), \quad (\text{A1})$$

where the operators in the coordinate representation are  $H_0 = -\frac{\partial^2}{\partial x^2}$ , and  $W(x,t) = \sum_n e^{-\frac{2i\pi nt}{T}} W_n(x)$ . We assume that the incoming square-integrable wave packet is  $\Psi_0(x,t)$ , i.e.,  $\Psi(x,t \rightarrow -\infty) = \Psi_0(x,t)$ . Obviously, if  $g_n = 0, \forall n$  then the time propagation is

$\Psi_0(x, t) = e^{-iH_0 t} \Psi_0$ , where  $\Psi_0 \equiv \Psi_0(x, 0)$ . Therefore, the effect of scattering is deduced by comparing  $\Psi(x, t)$  with  $\Psi_0(x, t)$ .

To proceed we note that there is a unitary operator  $U(t, s)$  that determines the time evolution

$$\Psi(t) = U(t, s) \Psi(s). \quad (\text{A2})$$

The formal properties of  $U(t, s)$  are discussed in Ref. [? ]. From now on we reserve the letters  $t$  and  $s$  for time, also when it does not cause confusion we omit the coordinate variables. For convenience, we set  $t = 0$  to be a reference time and introduce  $\Psi \equiv \Psi(x, 0)$  such that  $\Psi(t) = U(t) \Psi$ , where  $U(t) \equiv U(t, 0)$ . To compare  $\Psi(t)$  and  $\Psi_0(t)$  one can introduce the wave operator  $\Omega(s)$ , such that  $\Psi = \Omega \Psi_0$ ,  $\Omega = \Omega(0)$ . This operator is defined as the limit

$$\Omega(s) \equiv \lim_{t \rightarrow -\infty} U^{-1}(t, s) e^{-i(t-s)H_0}. \quad (\text{A3})$$

In scattering theory for time-independent potentials  $\Omega$  is often called the Møller operator [? ]. This operator exists also for time-periodic short-range potentials as discussed in Refs. [? ? ]. One important property of  $\Omega$  is called the intertwining relation:

$$U(t) \Omega = \Omega(t) e^{-iH_0 t}. \quad (\text{A4})$$

We use this relation upon decomposing  $\Psi_0$  in the eigenbasis of  $H_0$ . For convenience we refrain from using coordinate representation and use Dirac's notation for vectors instead:  $|\Psi_0\rangle = \int_{-\infty}^{\infty} dp \phi(p) |p\rangle$ . Now if we apply the intertwining relation to this decomposition we obtain

$$\int dp \phi(p) \left( U(t) \Omega |p\rangle - e^{-ip^2 t} \Omega(t) |p\rangle \right) = 0. \quad (\text{A5})$$

Since it should be valid for all possible initial wave packets described by  $\phi(p)$  we conclude that the integrand should be zero. By differentiating both sides with respect to time, we see that the only way to satisfy this condition is to assume that  $|f_p\rangle(t) \equiv \Omega(t) |p\rangle$  obeys

$$\left( H(t) - i \frac{\partial}{\partial t} \right) |f_p\rangle(t) = p^2 |f_p\rangle(t). \quad (\text{A6})$$

Let us take a closer look at Eq. (??). According to the Floquet theorem  $U(t+T, s+T) = U(t, s)$ ; see also Ref. [? ]. Therefore,  $\Omega(T) = \Omega$ , hence  $|f_p\rangle(t+T) = |f_p\rangle(t)$  and  $|f_p\rangle(t) = \sum_n e^{-\frac{2i\pi n t}{T}} |\tilde{f}_{pn}\rangle$ . By inserting this ansatz function into Eq. (??) and projecting onto a particular mode we obtain

$$(p^2 + n\omega - H_0) |\tilde{f}_{pn}\rangle = \sum_m W_{n-m} |\tilde{f}_{pm}\rangle, \quad (\text{A7})$$

where  $\omega = 2\pi/T$ . This is a (infinite-dimensional) matrix equation, and therefore, for each  $p^2$  there is an infinite number of solutions, which can be formally written as

$$|\tilde{f}_{pn}\rangle = |p_n\rangle\alpha_n + \frac{1}{p^2 + n\omega - H_0 + i\epsilon} \sum_m W_{n-m} |\tilde{f}_{pm}\rangle, \quad (\text{A8})$$

where  $\alpha_n$  and  $\epsilon \in \mathbb{R}$  are coefficients, and  $p_n^2 = p^2 + n\omega$ . We show below that the solution that satisfies the initial condition  $\Psi(x, t \rightarrow -\infty) \rightarrow \Psi_0(x, t)$  has  $\alpha_n = \delta_{n,0}$  and  $\epsilon > 0$ . To this end we write the formal solution to Eq. (??)

$$|\Psi\rangle(t) = e^{-iH_0 t} \int_{-\infty}^{\infty} dp \phi(p) \left( 1 - i \int_{-\infty}^t dt' e^{iH_0 t'} W(t') \Omega(t') e^{-iH_0 t'} \right) |p\rangle. \quad (\text{A9})$$

This equation yields  $|\Psi\rangle$

$$|\Psi\rangle = \int_{-\infty}^{\infty} dp \phi(p) \left( |p\rangle + \sum_{m,n} \frac{1}{p^2 + (n+m)\omega - H_0 + i\delta} W_n |\tilde{f}_{pm}\rangle \right). \quad (\text{A10})$$

where  $\delta$  is a small positive quantity. To obtain  $|\Psi\rangle$  we used the prescription  $W(t) \rightarrow e^{\delta t} W(t)$ , which is justified by noticing that for  $t \rightarrow -\infty$  the square-integrable wave packet cannot be affected by the finite-range potential. Now if we look at Eqs. (??) and (??) and notice that  $|\Psi\rangle = \Omega|\Psi_0\rangle = \int dp \phi(p) \sum_n |\tilde{f}_{pn}\rangle$  we deduce that  $\alpha_n = \delta_{n,0}$  and  $\epsilon > 0$ . At  $t$  we have  $|\Psi\rangle(t) = \int dp \phi(p) e^{-ip^2 t} \sum_n e^{-i\omega n t} |\tilde{f}_{pn}\rangle$ . This provides us with the wave function  $\Psi(x, t)$  for  $x \rightarrow \infty$ , which determines characteristics of transmission

$$\Psi(x, t) = \Psi_0(x, t) - \frac{i}{2} \sum_{m,n} \int dp \phi(p) \frac{e^{ip_n x - ip_n^2 t}}{p_n} \int dx' e^{-ip_n x'} W_{n-m}(x') \langle x' | \tilde{f}_{pm} \rangle, \quad (\text{A11})$$

a similar expression can be derived for  $x \rightarrow -\infty$ . Here we use the coordinate representation of the Green's function

$$G(x, x'; k^2) \equiv \langle x | \frac{1}{H_0 - k^2 - i\epsilon} | x' \rangle = \frac{i}{2k} e^{ik|x-x'|}. \quad (\text{A12})$$

**Scattering in time-independent picture.** In this subsection we consider Eq. (??) that determines the properties of the scattering in time-independent picture in more detail. In the coordinate representation it reads

$$\tilde{f}_{pn}(x) = \delta_{n,0} e^{ip_n x} - \sum_{m=-\infty}^{\infty} \int dx' G(x, x'; p_n^2) W_{n-m}(x') \tilde{f}_{pm}(x'), \quad (\text{A13})$$

where  $\delta_{m,n}$  is Kronecker's delta. This function at  $x \rightarrow \infty$  has the form  $\delta_{n,0}e^{ip_n x} + B_n e^{ip_n x}$  where

$$B_n = -\frac{i}{2p_n} \sum_{m=-\infty}^{\infty} \int dx' e^{-ip_n x'} W_{n-m}(x') \tilde{f}_{pm}(x') \quad (\text{A14})$$

whereas at  $x \rightarrow -\infty$  it has the form  $\delta_{n,0}e^{ip_n x} + \tilde{B}_n e^{-ip_n x}$  with

$$\tilde{B}_n = -\frac{i}{2p_n} \sum_{m=-\infty}^{\infty} \int dx' e^{ip_n x'} W_{n-m}(x') \tilde{f}_{pm}(x'), \quad (\text{A15})$$

Apparently, Eqs. (??), (??) and (??) contain all information about the scattering process and can be used to derive Eq. (??) of the main text. It is worthwhile to notice that for a plane wave with a given  $p^2$  the total probability to find a particle with the energies  $p_n^2, n = 0, \pm 1, \dots$  is conserved in the scattering process (see the next subsection). This can be seen as the conservation of the quasi-energy. At the same time the total energy is not conserved and can become larger or smaller, depending on the problem.

## 2. Conservation of Flux.

The Floquet modes,  $f_p$ , fully describe the scattering process. Since they always contain scattering states, there should be no probability to find a particle close to the potential at  $t \rightarrow \infty$  (assuming a square-integrable wave function at  $t \rightarrow -\infty$ ): the total outgoing flux should be equal to the total incoming flux. Note that since the states with  $p^2 - \omega(m+n) < 0$  do not give any contribution to the fluxes, the particle will leave these modes after some time. Physically it is easily understood, since a particle in these modes can undergo a transition to a scattering state and leave the range of the potential. This appendix shows the conservation of flux explicitly in a time-independent picture. Let us start with the scattering off zero-range potential described by Eq. (??), for which the flux is

$$\vec{j} = i(\psi \vec{\nabla} \psi^* - \psi^* \vec{\nabla} \psi). \quad (\text{A16})$$

The incoming flux is along the  $x$  axis and amounts to  $2p$ . The outgoing flux consists of the two parts: the first is along the  $x$  direction and equals to  $\sum_{p_n \geq 0} 2p_n |C_n|^2$ . The second piece is along the  $(-x)$  direction and amounts to  $\sum_{p_n \geq 0} 2p_n |C_n - \delta_{n,0}|^2$ . Let us show that from Eq. (??) it follows that

$$\sum_{p_n \geq 0} p_n |C_n|^2 = p \text{Re} C_0, \quad (\text{A17})$$

which means that the total flux is conserved. To this end, we multiply Eq. (??) with  $C_n^*$ ,

$$2p_n|C_n|^2 = 2p\delta_{n,0}C_n^* - ig_0|C_n|^2 - \frac{ig_1}{2}(C_{n+1} + C_{n-1})C_n^*. \quad (\text{A18})$$

Next we conjugate Eq. (??) and then multiply with  $C_n$ ,

$$2p_n^*|C_n|^2 = 2p\delta_{n,0}C_n + ig_0|C_n|^2 + \frac{ig_1}{2}(C_{n+1}^* + C_{n-1}^*)C_n. \quad (\text{A19})$$

Now we add these two equations and sum over all states

$$\sum (p_n + p_n^*)|C_n|^2 = 2p\text{Re}C_0. \quad (\text{A20})$$

Since  $p_n + p_n^*$  is non-zero only for the scattering states we obtain Eq. (??). Similar steps can be taken to show that the total flux is conserved for any short-range potential.

- 
- [1] Cheng Chin, Rudolf Grimm, Paul Julienne, and Eite Tiesinga. Feshbach resonances in ultra-cold gases. *Rev. Mod. Phys.*, 82:1225–1286, Apr 2010.
  - [2] M. Girardeau. Relationship between systems of impenetrable bosons and fermions in one dimension. *Journal of Mathematical Physics*, 1(6):516–523, 1960.
  - [3] Belen Paredes, Artur Widera, Valentin Murg, Olaf Mandel, Simon Folling, Ignacio Cirac, Gora V. Shlyapnikov, Theodor W. Hansch, and Immanuel Bloch. Tonks-girardeau gas of ultracold atoms in an optical lattice. *Nature*, 429:277–281, 2004.
  - [4] Toshiya Kinoshita, Trevor Wenger, and David S. Weiss. Observation of a one-dimensional tonks-girardeau gas. *Science*, 305(5687):1125–1128, 2004.
  - [5] M. D. Girardeau, E. M. Wright, and J. M. Triscari. Ground-state properties of a one-dimensional system of hard-core bosons in a harmonic trap. *Phys. Rev. A*, 63:033601, Feb 2001.
  - [6] F. Deuretzbacher, K. Fredenhagen, D. Becker, K. Bongs, K. Sengstock, and D. Pfannkuche. Exact solution of strongly interacting quasi-one-dimensional spinor bose gases. *Phys. Rev. Lett.*, 100:160405, Apr 2008.
  - [7] A. G. Volosniev, D. V. Fedorov, M. Jensen, A. S. Valiente, and N. T. Zinner. Strongly interacting confined quantum systems in one dimension. *Nat. Commun.*, 5:5300, 2014.
  - [8] M. Olshanii. Atomic scattering in the presence of an external confinement and a gas of impenetrable bosons. *Phys. Rev. Lett.*, 81:938–941, Aug 1998.

- [9] Philip F. Bagwell and Roger K. Lake. Resonances in transmission through an oscillating barrier. *Phys. Rev. B*, 46:15329–15336, Dec 1992.
- [10] S. T. Thompson, E. Hodby, and C. E. Wieman. Ultracold molecule production via a resonant oscillating magnetic field. *Phys. Rev. Lett.*, 95:190404, Nov 2005.
- [11] M. Büttiker and R. Landauer. Traversal time for tunneling. *Phys. Rev. Lett.*, 49:1739–1742, Dec 1982.
- [12] M Büttiker and R Landauer. Traversal time for tunneling. *Physica Scripta*, 32(4):429, 1985.
- [13] Mathias Wagner. Photon-assisted transmission through an oscillating quantum well: A transfer-matrix approach to coherent destruction of tunneling. *Phys. Rev. A*, 51:798–808, Jan 1995.
- [14] G. Burmeister and K. Maschke. Scattering by time-periodic potentials in one dimension and its influence on electronic transport. *Phys. Rev. B*, 57:13050–13060, May 1998.
- [15] Wenjun Li and L. E. Reichl. Floquet scattering through a time-periodic potential. *Phys. Rev. B*, 60:15732–15741, Dec 1999.
- [16] Michael Henseler, Thomas Dittrich, and Klaus Richter. Classical and quantum periodically driven scattering in one dimension. *Phys. Rev. E*, 64:046218, Sep 2001.
- [17] Thomas M. Hanna, Thorsten Köhler, and Keith Burnett. Association of molecules using a resonantly modulated magnetic field. *Phys. Rev. A*, 75:013606, Jan 2007.
- [18] Christian Langmack, D. Hudson Smith, and Eric Braaten. Association of atoms into universal dimers using an oscillating magnetic field. *Phys. Rev. Lett.*, 114:103002, Mar 2015.
- [19] D. Hudson Smith. Inducing resonant interactions in ultracold atoms with a modulated magnetic field. *Phys. Rev. Lett.*, 115:193002, Nov 2015.
- [20] Smith, D. Hudson. Induced two-body scattering resonances from a square-well potential with oscillating depth. *EPJ Web of Conferences*, 113:02005, 2016.
- [21] D. H. Smith. *Resonant Floquet Scattering of Ultracold Atoms, PhD thesis*. arXiv:1611.05412, 2016.
- [22] A. G. Sykes, H. Landa, and D. S. Petrov. Two- and three-body problem with floquet-driven zero-range interactions. *Phys. Rev. A*, 95:062705, Jun 2017.
- [23] P. Hänggi. *Quantum Transport and Dissipation, chapter 5*. Wiley-VCH, Weinheim, edited by T. Dittrich, 1998.

- [24] Ranjan K. Mallik. The inverse of a tridiagonal matrix. *Linear Algebra and its Applications*, 325(13):109 – 139, 2001.
- [25] Moawwad E.A. El-Mikkawy. On the inverse of a general tridiagonal matrix. *Applied Mathematics and Computation*, 150(3):669 – 679, 2004.
- [26] G. Burmeister and K. Maschke. Bound states revealed by time-periodic perturbing scattering potentials. *Phys. Rev. B*, 59:4612–4614, Feb 1999.
- [27] U. Fano. Effects of configuration interaction on intensities and phase shifts. *Phys. Rev.*, 124:1866–1878, Dec 1961.
- [28] A. G. Volosniev, H.-W. Hammer, and N. T. Zinner. Real-time dynamics of an impurity in an ideal bose gas in a trap. *Phys. Rev. A*, 92:023623, Aug 2015.
- [29] Martin Ebert, Artem Volosniev, and Hans-Werner Hammer. Two cold atoms in a time-dependent harmonic trap in one dimension. *Annalen der Physik*, 528(9-10):693–704, 2016.
- [30] X. Y. Yin, Yangqian Yan, and D. Hudson Smith. Dynamics of small trapped one-dimensional fermi gas under oscillating magnetic fields. *Phys. Rev. A*, 94:043639, Oct 2016.
- [31] Kenji Yajima. Scattering theory for schrödinger equations with potentials periodic in time. *J. Math. Soc. Japan*, 29:729–743, 1977.
- [32] Jonh R. Taylor. *Scattering Theory. The Quantum Theory of Nonrelativistic Collisions*. Dover Publications, Inc., 2006.
- [33] J. Howland. Scattering theory for hamiltonians periodic in time. *Indiana J. Math.*, 28:471–494, 1979.