

SAT Math Mastery · Trick Pack 1A

SAT Trick Pack 1 — Part A

Time 75 minutes: 55 min solving + 20 min self-check and error log

Topics exponent rewriting · linear constraints · tiered models · right-triangle similarity · AM–GM bounds · symmetry & Vieta · statistics effects · conditional probability · similar solids · substitution · factor roots · parameters · systems

Name / Date _____

1 · SKILLS & IDEAS

- S1. Rewrite radicals and mixed bases as powers of one base, then equate exponents.
- S2. Optimise under several linear constraints: find the binding one, round in the right direction, verify the rest.
- S3. Build a layered cost model (flat fee, tiers, surcharge, tax) and respect the order of operations.
- S4. Use the altitude to the hypotenuse: similar triangles, geometric means, area ratios.
- S5. Bound a product with a fixed sum of squares ($a^2 + b^2 \geq 2ab$) and find where equality holds.
- S6. Read the axis of symmetry from equal outputs; combine with Vieta and sign facts.
- S7. Decide how a replacement changes the mean and the spread without recomputing.
- S8. Condition on a subgroup (it is the new denominator) and recover a missing count.
- S9. Scale area by k^2 and volume by k^3 ; invert the factor from an area ratio.
- S10. Substitute $u = a^x$ to turn an exponential equation into a quadratic; reject impossible roots.
- S11. Plug in a zero of a factor instead of expanding.
- S12. Parameters in linear equations, absolute-value graphs and parametrised solution sets.

2 · REQUIRED FACTS

- $\sqrt[n]{a^m} = a^{m/n}$, $a^m a^n = a^{m+n}$, $\frac{a^m}{a^n} = a^{m-n}$, $(a^m)^n = a^{mn}$; if $a > 0$, $a \neq 1$ and $a^p = a^q$ then $p = q$.
- Exponentials a^x with $a > 0$ are always positive: $a^x = \text{negative}$ has no real solution.
- Altitude to the hypotenuse splits a right triangle into two triangles similar to it and to each other.
- Replacing values keeps the mean if the total is unchanged; standard deviation falls when extreme values move toward the centre.
- Conditioning on a group makes that group's total the denominator.
- Similar solids with linear factor k : lengths $\times k$, areas $\times k^2$, volumes $\times k^3$.
- If $f(p) = f(q)$ for a quadratic, the axis is $x = \frac{p+q}{2}$; zeros sum to $-\frac{b}{a}$ and multiply to $\frac{c}{a}$.
- A linear equation whose x -terms cancel has either no solution or infinitely many, decided by the leftover constants.

3 · REQUIRED FORMULAS

$$a^{m/n} = \sqrt[n]{a^m}, \quad CD^2 = AD \cdot DB, \quad AC^2 = AD \cdot AB, \quad BC^2 = DB \cdot AB, \quad \frac{[ACD]}{[CDB]} = \frac{AD}{DB}$$

$$a^2 + b^2 \geq 2ab, \quad x_{\text{axis}} = \frac{p+q}{2} = -\frac{b}{2a}, \quad r+s = -\frac{b}{a}, \quad rs = \frac{c}{a}, \quad \frac{V_2}{V_1} = \left(\sqrt{\frac{S_2}{S_1}} \right)^3$$

4 · MINI THEORY SHEET

Common base first. Convert every radical and every base to one prime base ($4 = 2^2$, $8 = 2^3$, $\sqrt[3]{2^x} = 2^{x/3}$), simplify each side to a single power, and only then equate exponents. Two equations become a linear system in the exponents.

Constraint problems. Turn each sentence into an inequality. Test the maximum (or minimum) against *every* constraint; the one that fails first is binding. Whole-number answers: round down for a maximum, up for a minimum, then re-check.

Layered models. Write the pipeline in the order described (fee $\rightarrow$ tiers $\rightarrow$ surcharge on the stated part $\rightarrow$ tax on the stated total). Test one input by hand and compare with the formulas.

Parameters. Expand and collect. Same x -coefficient on both sides leaves a statement about constants: true means infinitely many solutions, false means none.

Substitution. If a^x appears as a^{2x} and a^x (or $a^{x+c} = a^c \cdot a^x$), set $u = a^x$ with $u > 0$.

Speed shortcuts.

Situation	Shortcut
mixed radicals and bases	one common base, then equate exponents; solve the linear system
ratio-plus-budget constraints	set the ratio constraint to equality, test each other limit
fee, tiers, surcharge, tax	one test input by hand; compare with each option
altitude to the hypotenuse	area ratio = segment ratio; $CD^2 = AD \cdot DB$
product with fixed $a^2 + b^2$	maximum at $a = b$ (rescale if a coefficient sits in front)
equal outputs of a quadratic	axis at the midpoint of the two inputs
swap values, same total	mean unchanged; spread falls if extremes move inward
“of those who ...”	the subgroup total is the denominator
areas $\rightarrow$ volumes	$k = \sqrt{\text{area ratio}}$, then k^3
A^{2x} and A^x together	$u = A^x$, factor, discard $u \leq 0$
$p(x) = (x - r)q(x)$ with unknown coefficient	substitute $x = r$
$ax + b = cx + d$ with parameter	match coefficients: same line $\Rightarrow$ infinitely many
$ x - h + v$ against a line	one solution exactly when the line hits the vertex
equations that are multiples	substitute the parametrised point into one equation: do the t -terms cancel?

5 · GUIDED EXAMPLES

Example 1 — Common base, then a linear system

If $2^x \cdot 8^y = 2^9$ and $\frac{4^x}{2^y} = 16$, find x and y .

$$2^x \cdot 2^{3y} = 2^9 \Rightarrow x + 3y = 9. \quad \frac{2^{2x}}{2^y} = 2^4 \Rightarrow 2x - y = 4.$$

From the second, $y = 2x - 4$; then $x + 6x - 12 = 9 \Rightarrow x = 3$, $y = 2$.

Example 2 — Similar solids: area $\rightarrow$ factor $\rightarrow$ volume

Two similar solids have surface areas 36 and 81. The smaller has volume 40. Find the volume of the larger.

$$k = \sqrt{\frac{81}{36}} = \frac{3}{2}. \quad \text{Volume scales by } k^3 = \frac{27}{8}: \quad 40 \cdot \frac{27}{8} = 135.$$

Example 3 — Equal outputs fix the axis

$g(x) = x^2 + bx + c$, with $g(2) = g(12)$ and $g(0) = 11$. Find the minimum value of g .

$$\text{Axis: } x = \frac{2 + 12}{2} = 7 = -\frac{b}{2} \Rightarrow b = -14. \quad c = g(0) = 11.$$

$$\text{Minimum} = g(7) = 49 - 98 + 11 = -38.$$

6 · PRACTICE SET — 15 PROBLEMS · ALL SAT CHALLENGE

Each problem states its tool. Write your work in the space below. Multiple choice: one correct option. Student-produced response (SPR): enter an exact value (fraction or decimal).

1.

SAT CHALLENGE · CALCULATOR

Real numbers x and y satisfy

$$\sqrt[3]{2^x} \cdot 4^y = 256 \quad \text{and} \quad \frac{16^x}{\sqrt[3]{4^y}} = 2^{22}.$$

What is the value of xy ?

2.

SAT CHALLENGE · CALCULATOR

A science department orders microscopes at \$240 each and slide kits at \$15 each. The order must include at least 6 slide kits for every microscope, the total cost may not exceed \$2,500, and the storage cabinet holds at most 46 items (microscopes and kits combined). What is the greatest number of microscopes the department can order?

3.

SAT CHALLENGE · CALCULATOR

A utility charges a \$15 service fee each month. Usage is billed at \$2.40 per unit for the first 20 units and \$3.60 for each unit beyond 20. A 10% fuel surcharge is added to the usage charge only (not to the service fee). Then 5% tax is applied to the sum of the service fee and the surcharged usage charge. For $u > 20$ units, which function gives the total monthly bill $B(u)$, in dollars?

(A) $B(u) = 3.78u - 9.45$

(B) $B(u) = 4.158u - 11.97$

(C) $B(u) = 4.158u - 10.395$

(D) $B(u) = 4.158u - 12.72$

4.

SAT CHALLENGE · NO CALCULATOR NEEDED

In triangle ABC , the angle at C is a right angle. Point D lies on $\overline{AB}$ so that $\overline{CD} \perp \overline{AB}$. The area of triangle ACD is 48 and the area of triangle CDB is 108. What is the length of $\overline{AB}$?

5.

SAT CHALLENGE • NO CALCULATOR NEEDED

For real numbers p and q with $p^2 + 4q^2 = 72$, the product $(2x + p)(x + q)$ is equivalent to $2x^2 + bx + c$, where b and c are constants. What is the greatest possible value of c ?

6.

SAT CHALLENGE • NO CALCULATOR NEEDED

The quadratic function $f(x) = ax^2 + bx + c$, where a , b and c are constants, has two distinct real zeros. It is known that $f(2) = f(8)$, $f(0) < 0$ and $f(5) > 0$. Consider the following statements.

- I. $b > 0$
- II. The product of the two zeros is greater than 25.
- III. Both zeros are positive.

Which of the statements must be true?

- (A) I only
- (B) II and III only
- (C) I and III only
- (D) I, II, and III

7.

SAT CHALLENGE · NO CALCULATOR NEEDED

A data set has eight values: 12, 15, 18, 20, 20, 22, 25, 28. The values 12 and 28 are removed and the values 19 and 21 are added in their place. Which of the following describes the mean and the standard deviation (SD) of the new data set compared with the original data set?

(A) Mean smaller, SD smaller

(B) Mean same, SD larger

(C) Mean same, SD smaller

(D) Mean same, SD same

8.

SAT CHALLENGE · NO CALCULATOR NEEDED

A conference has 300 participants. Each participant attended either the Technical track or the Business track, and either in person or online. Some of the data are shown.

	Technical	Business	Total
In person	70		
Online	50		
Total	120		300

Of the participants in the Business track, 35% attended online. How many of the 300 participants attended in person?

(A) 187

(B) 63

(C) 113

(D) 145

9.

SAT CHALLENGE · NO CALCULATOR NEEDED

Two similar rectangular crates have surface areas of 144 square feet and 400 square feet. The smaller crate has a volume of 108 cubic feet. What is the volume, in cubic feet, of the larger crate?

(A) 1500

(B) 180

(C) 300

(D) 500

10.

SAT CHALLENGE · NO CALCULATOR NEEDED

For a real number x , $4^x - 2^{x+2} - 32 = 0$. What is the value of 2^{x+1} ?

(A) -8 (B) 8 (C) 16 (D) 32 **11.**

SAT CHALLENGE · NO CALCULATOR NEEDED

For constants p , q and c , the equation

$$(2x - 3)(x^2 + px + q) = 2x^3 - 5x^2 + cx + 9$$

is true for all real values of x . What is the value of c ?

SAT CHALLENGE • NO CALCULATOR NEEDED

SAT CHALLENGE • NO CALCULATOR NEEDED

In the equation above, k and m are constants. If the equation has infinitely many solutions, what is the value of $m - k$?

14.

SAT CHALLENGE · NO CALCULATOR NEEDED

The equation $|x + 6| - 7 = \frac{x}{3} + c$, where c is a constant, has exactly one solution x . What is the value of $2c + x$?

(A) -11 (B) -6 (C) -5 (D) -16 **15.**

SAT CHALLENGE · NO CALCULATOR NEEDED

$$3x + 2y = 12$$

$$9x + 6y = 36$$

For the system above, which of the following ordered pairs (x, y) , where t is any real number, is a solution to the system?

(A) $(2t + 4, -3t)$ (B) $(2t + 4, 3t)$ (C) $(3t + 4, -2t)$ (D) $(4 - 2t, -3t)$