

SAT Math Mastery · Week 9A**Solids, Coordinate Geometry & Composite Shapes — Answer Key & Worked Solutions****Time** 120 minutes: 90 min solving + 30 min self-check and error log**Topics** volume and surface area of solids · scaling laws · composite, glued and inscribed solids · distance, midpoint, slope · area from coordinates · composite and shaded regions · rate of fill and unit volume · dilations**Name / Date** _____**1 · SKILLS & IDEAS**

- S1.** Volume of prisms, cylinders, cones, spheres and pyramids; recover a radius from a circumference, a volume or a surface area.
- S2.** Surface area: total vs lateral, slant height of a cone or pyramid, surface area of a prism from its three face pairs.
- S3.** Scaling laws: lengths $\times k$, areas $\times k^2$, volumes $\times k^3$; recover k from an area or volume ratio.
- S4.** Composite, glued, removed and inscribed solids (sphere in a cube, cylinder in a sphere, joined prisms hide two faces).
- S5.** Coordinate geometry: distance, midpoint (including a missing endpoint), slope, equidistant points.
- S6.** Area from coordinates: base and height on a grid, decomposition, and the shoelace formula.
- S7.** Composite and shaded plane regions: subtract a part from a whole; equilateral-triangle trapezoids; algebraic side lengths leading to a quadratic.
- S8.** Rate of fill, unit-volume conversions ($\text{m}^3 \leftrightarrow \text{L}$), and dilations in the coordinate plane.

2 · REQUIRED FACTS

- Volume of any prism or cylinder is $V = Bh$ (area of the base times the height); a cone or pyramid with the same base and height holds exactly $\frac{1}{3}$ of that.
- The slant height ℓ , the height h and the base radius r of a right cone form a right triangle: $\ell^2 = r^2 + h^2$.
- A cube's space diagonal is $s\sqrt{3}$; a rectangular box's is $\sqrt{a^2 + b^2 + c^2}$. A rectangle inscribed in a circle has its diagonal equal to the diameter.
- When two solids are glued along a face, the surface area is the sum of the two surface areas minus *twice* the area of the glued face.
- Similar figures: if lengths scale by k , areas scale by k^2 and volumes by k^3 ; angle measures never change.
- $1 \text{ m}^3 = 1000 \text{ L}$ and $1 \text{ cm}^3 = 1 \text{ mL}$; square and cubic units convert with the *square* and *cube* of the length factor.
- A dilation centered at the origin with scale factor k sends $(x, y) \rightarrow (kx, ky)$, multiplies lengths by k and areas by k^2 .

3 · REQUIRED FORMULAS

$$V_{\text{prism}} = Bh, \quad V_{\text{cyl}} = \pi r^2 h, \quad V_{\text{cone}} = \frac{1}{3}\pi r^2 h, \quad V_{\text{sphere}} = \frac{4}{3}\pi r^3, \quad V_{\text{pyramid}} = \frac{1}{3}Bh$$

$$SA_{\text{sphere}} = 4\pi r^2, \quad SA_{\text{cyl}} = 2\pi r^2 + 2\pi rh, \quad SA_{\text{cone}} = \pi r^2 + \pi r\ell, \quad SA_{\text{box}} = 2(ab + bc + ac)$$

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}, \quad M = \left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2}\right), \quad \text{Area} = \frac{1}{2} |\sum (x_i y_{i+1} - x_{i+1} y_i)|$$

4 · MINI THEORY SHEET

Choose the solid's key quantity first.

Given	Fastest route
a circumference C	$r = \frac{C}{2\pi}$, then use r^2 in the area or volume
radius and slant height of a cone	$h = \sqrt{\ell^2 - r^2}$ (look for 6-8-10, 5-12-13)
a volume $= k\pi$	divide by π first; solve r^3 or r^2h with whole-number guesses
lateral area of a pyramid	each triangle $= \frac{1}{2}(\text{side})\ell$; find ℓ , then $h = \sqrt{\ell^2 - (\text{side}/2)^2}$
a rate of fill Q per minute	height rate $= \frac{Q}{\text{base area}}$ (not divided by the radius)

Inscribed solids. Draw the cross section: a sphere in a cube touches the faces (diameter = edge); a sphere around a cube passes through the vertices (diameter = space diagonal); a cylinder in a sphere has the sphere's radius as the hypotenuse of a right triangle with the half-height and the base radius.

Scaling. If the ratio of areas is $a : b$, then $k = \sqrt{b/a}$; if the ratio of volumes is $a : b$, then $k = \sqrt[3]{b/a}$. Convert, then apply to the quantity asked.

Area from coordinates. Axis-parallel side $\rightarrow$ base and height. Otherwise use the shoelace sum, or box the figure and subtract corner triangles.

Unit volume. Convert every quantity to one unit *before* dividing; a rate in liters per minute needs a volume in liters.

Speed shortcuts.

Situation	Shortcut
ratio of volumes $a : b$	lengths $\sqrt[3]{b/a}$, areas $\left(\sqrt[3]{b/a}\right)^2$
radius $\times m$, height $\times n$	volume $\times m^2n$ (a cylinder or cone)
cube with surface area SA	face $= SA/6$, edge $= \sqrt{SA/6}$
glued prisms	$SA = SA_1 + SA_2 - 2(\text{glued face})$, or rebuild the combined box
missing endpoint	$B = 2M - A$ coordinatewise
cone or sphere with a π in the volume	cancel π at once and solve for r
solid of revolution of a right triangle	cone volume $\frac{1}{3}\pi r^2h$ with r the distance to the axis
dilation by k about the origin	multiply every coordinate by k ; area $\times k^2$

5 · GUIDED EXAMPLES

Example 1 — Composite solid: cylinder topped by a hemisphere

A silo is a cylinder of radius 3 and height 10 with a hemisphere of radius 3 on top. Find its volume.

Cylinder: $\pi r^2 h = \pi(9)(10) = 90\pi$. Hemisphere: $\frac{1}{2} \cdot \frac{4}{3}\pi r^3 = \frac{2}{3}\pi(27) = 18\pi$.

Total volume $= 90\pi + 18\pi = 108\pi$.

Example 2 — Scaling: area ratio to volume

Two similar prisms have surface areas 50 and 162. The smaller has volume 125. Find the volume of the larger.

Area ratio $\frac{162}{50} = \frac{81}{25} = k^2$, so $k = \frac{9}{5}$. Volume ratio $k^3 = \frac{729}{125}$.

Larger volume $= 125 \cdot \frac{729}{125} = 729$.

Example 3 — Area from coordinates (shoelace)

Find the area of the triangle with vertices $P(-1, 2)$, $Q(5, -2)$, $R(3, 6)$.

$\sum (x_i y_{i+1} - x_{i+1} y_i)$: $P \rightarrow Q$: $(-1)(-2) - (5)(2) = -8$; $Q \rightarrow R$: $(5)(6) - (3)(-2) = 36$; $R \rightarrow P$: $(3)(2) - (-1)(6) = 12$.

Sum $= -8 + 36 + 12 = 40$, so area $= \frac{1}{2}|40| = 20$.

6 · PRACTICE SET — 30 PROBLEMS · EASY → MEDIUM → HARD → SAT CHALLENGE

Each problem states its tool. Write your work in the space below. Multiple choice: one correct option. Student-produced response (SPR): enter an exact value (fraction or decimal).

1. EASY · NO CALCULATOR NEEDED

A right circular cylinder has a height of 5 and a base with circumference 8π . What is the volume of the cylinder?

(A) 40π

(B) 80π ✓

(C) 160π

(D) 320π

Worked solution


1. $C = 2\pi r = 8\pi$, so $r = 4$.

2. $V = \pi r^2 h = \pi(4)^2(5) = 80\pi$.

Answer: (B)

2. EASY · NO CALCULATOR NEEDED

In the xy -plane, the vertices of triangle ABC are plotted as shown. What is the area of triangle ABC ?



Worked solution

1. Read the vertices: $A(-2, 1)$, $B(5, 1)$, $C(3, 6)$.

2. $\overline{AB}$ is horizontal, so the base is $5 - (-2) = 7$.

3. The height is the vertical distance from C to line AB : $6 - 1 = 5$.

4. Area = $\frac{1}{2}(7)(5) = 17.5$.

Answer: 17.5

3. EASY · NO CALCULATOR NEEDED

Point $M(3, 5)$ is the midpoint of $\overline{AB}$. If A has coordinates $(-1, 2)$, what are the coordinates of B ?

(A) $\left(1, \frac{7}{2}\right)$

(B) $(8, 7)$

(C) $(7, 8)$ ✓

(D) $(4, 3)$

Worked solution

1. Midpoint formula: $3 = \frac{-1 + x_B}{2}$ and $5 = \frac{2 + y_B}{2}$.

2. $x_B = 6 + 1 = 7$ and $y_B = 10 - 2 = 8$.

3. $B = (7, 8)$.

Answer: (C)

4.

EASY · CALCULATOR

A cone has a volume of 14 cubic centimeters. A second cone is similar to the first, and each linear dimension of the second cone is 3 times the corresponding dimension of the first. What is the volume, in cubic centimeters, of the second cone?

Worked solution

1. Lengths scale by $k = 3$, so volumes scale by $k^3 = 27$.

2. $14 \cdot 27 = 378$.

Answer: 378

5.

EASY · NO CALCULATOR NEEDED

A cube has a total surface area of 150 square inches. What is the volume of the cube, in cubic inches?

(A) 125 ✓

(B) 150

(C) 625

(D) 25

Worked solution

1. A cube has 6 identical square faces: each face has area $\frac{150}{6} = 25$.

2. Edge length $= \sqrt{25} = 5$.


3. $V = 5^3 = 125$.

Answer: (A)

6.

MEDIUM · CALCULATOR

The cone shown has a base radius of 6 and a slant height of 10. What is the volume of the cone?

(A) 120π (B) 288π (C) 360π (D) 96π ✓**Worked solution**

1. The height, the radius and the slant height form a right triangle: $h = \sqrt{10^2 - 6^2} = \sqrt{64} = 8$.

2. $V = \frac{1}{3}\pi r^2 h = \frac{1}{3}\pi(36)(8) = 96\pi$.

Answer: (D)

7.

MEDIUM · NO CALCULATOR NEEDED

A sphere has a volume of 288π cubic centimeters. The surface area of the sphere is $k\pi$ square centimeters. What is the value of k ?

Worked solution

1. $\frac{4}{3}\pi r^3 = 288\pi \Rightarrow r^3 = 216 \Rightarrow r = 6$.

2. $SA = 4\pi r^2 = 4\pi(36) = 144\pi$.

3. $k = 144$.

Answer: 144

8.

MEDIUM · NO CALCULATOR NEEDED

A cylinder has radius r and height h . Which of the following changes to the cylinder results in a volume that is exactly twice the original volume?

I. Multiply the radius by $\sqrt{2}$ and leave the height unchanged.

II. Multiply the radius by 2 and divide the height by 2.

III. Multiply the radius by 2 and divide the height by 4.

(A) I only

(B) II only

(C) I and II only ✓

(D) I, II, and III

Worked solution

1. $V = \pi r^2 h$: the volume is multiplied by (radius factor)² · (height factor).

2. Statement I: $(\sqrt{2})^2 \cdot 1 = 2$, so the volume doubles.

3. Statement II: $2^2 \cdot \frac{1}{2} = 2$, so the volume doubles.

4. Statement III: $2^2 \cdot \frac{1}{4} = 1$, so the volume is unchanged.

5. Only I and II double the volume.

Answer: (C)

9.

MEDIUM · NO CALCULATOR NEEDED

A rectangular box has a square base with side length 6 and a height of h . The total surface area of the box is 240 square units. What is the value of h ?

Worked solution

1. Surface area = $2(\text{base}) + 4(\text{side face}) = 2(6)(6) + 4(6h) = 72 + 24h$.

2. $72 + 24h = 240 \Rightarrow 24h = 168$.

3. $h = 7$.

Answer: 7

10.

MEDIUM · NO CALCULATOR NEEDED

The endpoints of a diameter of a circle in the xy -plane are $(-3, 2)$ and $(5, 8)$. What is the radius of the circle?

(A) 100

(B) 5 ✓

(C) 10

(D) 25

Worked solution

1. Diameter = $\sqrt{(5 - (-3))^2 + (8 - 2)^2} = \sqrt{64 + 36} = 10$.

2. Radius = $\frac{10}{2} = 5$.

Answer: (B)

11.

MEDIUM · NO CALCULATOR NEEDED

Triangle ABC is similar to triangle DEF . The area of triangle ABC is 48 and the area of triangle DEF is 108. The longest side of triangle ABC has length 10. What is the length of the longest side of triangle DEF ?

Worked solution

1. Area ratio = $\frac{108}{48} = \frac{9}{4} = k^2$, so $k = \frac{3}{2}$.

2. Longest side of $DEF = 10 \cdot \frac{3}{2} = 15$.

Answer: 15

12.

MEDIUM · CALCULATOR

A cylindrical tank has an inside radius of 2 meters. Water flows into the tank at a constant rate of 0.5 cubic meter per minute. Which of the following is closest to the rate, in meters per minute, at which the water level in the tank rises?

(A) 0.08

(B) 0.16

(C) 0.25

(D) 0.04 ✓

Worked solution

1. In one minute the water added is 0.5 m^3 , spread over the circular base of area $\pi r^2 = 4\pi \text{ m}^2$.


2. Rate of rise = $\frac{0.5}{4\pi} = \frac{1}{8\pi} \approx 0.040$ meters per minute.

Answer: (D)

13.

MEDIUM · NO CALCULATOR NEEDED

A circle is inscribed in a square with side length 8, as shown. What is the area of the shaded region?

(A) $64 - 16\pi$ ✓(B) $64 - 8\pi$ (C) $64 - 4\pi$ (D) 16π **Worked solution**

1. The circle touches all four sides, so its diameter is 8 and its radius is 4.

2. Circle area = $\pi(4)^2 = 16\pi$; square area = 64.

3. Shaded area = $64 - 16\pi$.

Answer: (A)

14.

HARD · CALCULATOR

A solid sphere with a radius of 3 centimeters is placed in a cylindrical container that holds water and has an inside base radius of 5 centimeters. The sphere is completely submerged and no water spills. By how many centimeters does the water level rise?

Worked solution

1. The rise in water equals the volume of the sphere: $\frac{4}{3}\pi(3)^3 = 36\pi \text{ cm}^3$.

2. The water level rises over the base area $\pi(5)^2 = 25\pi \text{ cm}^2$.

3. Rise = $\frac{36\pi}{25\pi} = \frac{36}{25} = 1.44 \text{ cm}$.

Answer: 1.44

15.

HARD · NO CALCULATOR NEEDED

All eight vertices of a cube with edge length 6 lie on the surface of a sphere. What is the volume of the sphere?

(A) $72\sqrt{2}\pi$

(B) $108\sqrt{3}\pi$ ✓

(C) 288π

(D) 36π

Worked solution

1. The sphere's diameter is the cube's space diagonal: $6\sqrt{3}$, so $r = 3\sqrt{3}$.

2. $r^3 = (3\sqrt{3})^3 = 27 \cdot 3\sqrt{3} = 81\sqrt{3}$.


3. $V = \frac{4}{3}\pi r^3 = \frac{4}{3}\pi \cdot 81\sqrt{3} = 108\sqrt{3}\pi$.

Answer: (B)

16.

HARD · NO CALCULATOR NEEDED

Two identical rectangular blocks, each measuring 3 by 4 by 5, are glued together along a 4-by-5 face to form one larger rectangular block, as shown. What is the total surface area of the larger block?



(A) 188

(B) 94

(C) 148 ✓

(D) 168

Worked solution

1. One block: $2(3 \cdot 4 + 3 \cdot 5 + 4 \cdot 5) = 2(47) = 94$.

2. Two blocks: $2 \cdot 94 = 188$. The glued faces are hidden: subtract $2 \cdot (4 \cdot 5) = 40$.

3. $188 - 40 = 148$. (Check: the new block is $6 \times 4 \times 5$, with $2(24 + 30 + 20) = 148$.)

Answer: (C)

17.

HARD · CALCULATOR

The four lateral faces of a right square pyramid are congruent triangles with a combined area of 260. The base of the pyramid is a square with side length 10. What is the volume of the pyramid?

(A) 400 ✓

(B) 1200

(C) 1300

(D) $\frac{1300}{3}$

Worked solution

1. Each lateral face has area $\frac{260}{4} = 65 = \frac{1}{2}(10)\ell$, so the slant height is $\ell = 13$.

2. The height, half the base side (5) and the slant height form a right triangle: $h = \sqrt{13^2 - 5^2} = 12$.


3. $V = \frac{1}{3}Bh = \frac{1}{3}(100)(12) = 400$.

Answer: (A)

18.

HARD · CALCULATOR

In the xy -plane, quadrilateral $ABCD$ has the vertices shown on the grid. What is the area of quadrilateral $ABCD$?



(A) 70

(B) $\frac{63}{2}$

(C) 38

(D) 35 ✓

Worked solution

1. Read the vertices in order: $A(-4, -1)$, $B(3, -2)$, $C(5, 4)$, $D(-2, 3)$.
2. $A \rightarrow B$: $(-4)(-2) - (3)(-1) = 11$. $B \rightarrow C$: $(3)(4) - (5)(-2) = 22$.
3. $C \rightarrow D$: $(5)(3) - (-2)(4) = 23$. $D \rightarrow A$: $(-2)(-1) - (-4)(3) = 14$.
4. Sum = $11 + 22 + 23 + 14 = 70$, so the area is $\frac{1}{2}(70) = 35$.

Answer: (D)

19.

HARD · NO CALCULATOR NEEDED

A rectangle with side lengths 10 and 24 is inscribed in a circle. The area of the circle is $k\pi$. What is the value of k ?

Worked solution


1. The diagonal of an inscribed rectangle is a diameter: $d = \sqrt{10^2 + 24^2} = \sqrt{676} = 26$.
2. $r = 13$, so the area is $\pi(13)^2 = 169\pi$.
3. $k = 169$.

Answer: 169

20.

HARD · CALCULATOR

The trapezoid shown is composed of three congruent equilateral triangles. The perimeter of the trapezoid is 20. What is the area of the trapezoid?



- (A) $48\sqrt{3}$ (B) $4\sqrt{3}$
(C) $12\sqrt{3}$ ✓ (D) $16\sqrt{3}$

Worked solution

1. The perimeter consists of 5 sides each equal to the triangle side s : $5s = 20 \Rightarrow s = 4$.
2. Area of one equilateral triangle $= \frac{\sqrt{3}}{4}s^2 = \frac{\sqrt{3}}{4}(16) = 4\sqrt{3}$.
3. Three triangles: $3 \cdot 4\sqrt{3} = 12\sqrt{3}$.

Answer: (C)

21.

HARD · CALCULATOR

A rectangular pool measures 6 meters by 4 meters and is 1.5 meters deep. A pump fills the empty pool at a constant rate of 150 liters per minute. How many hours does it take to fill the pool completely? (1 cubic meter = 1000 liters)

Worked solution

1. Volume = $6 \cdot 4 \cdot 1.5 = 36 \text{ m}^3 = 36,000 \text{ liters}$.
2. Time = $\frac{36,000}{150} = 240 \text{ minutes}$.
3. $240 \div 60 = 4 \text{ hours}$.

Answer: 4

22.

HARD · NO CALCULATOR NEEDED

Two solid cones are similar. The volume of the larger cone is $\frac{27}{8}$ times the volume of the smaller cone. The surface area of the smaller cone is 100 square centimeters. What is the surface area, in square centimeters, of the larger cone?

- (A) 150 (B) 225 ✓
(C) 337.5 (D) 800

Worked solution

1. Volume ratio $k^3 = \frac{27}{8}$, so the length ratio is $k = \frac{3}{2}$.
2. Surface areas scale by $k^2 = \frac{9}{4}$.
3. $100 \cdot \frac{9}{4} = 225$.

Answer: (B)

23.

HARD · CALCULATOR

In the xy -plane, triangle ABC has vertices $A(1, 2)$, $B(7, 2)$ and $C(1, 10)$. Triangle DEF is the image of triangle ABC under a dilation centered at the origin, where D is the image of A and $D = (3, 6)$. What is the area of triangle DEF ?

(A) 648

(B) 72

(C) 144

(D) 216 ✓

Worked solution

1. $A(1, 2) \rightarrow D(3, 6)$: each coordinate is multiplied by 3, so $k = 3$.
2. Area of ABC : the base $\overline{AB} = 6$ is horizontal and the height $\overline{AC} = 8$ is vertical, so area $= \frac{1}{2}(6)(8) = 24$.
3. Areas scale by $k^2 = 9$: $24 \cdot 9 = 216$.

Answer: (D)

24.

SAT CHALLENGE · CALCULATOR

A cone-shaped paper cup is held point down. The cup has an inside radius of 6 centimeters at the top and a height of 12 centimeters. The cup contains water to a depth of 8 centimeters. All of this water is poured into an empty cylindrical glass with an inside radius of 4 centimeters. What is the depth, in centimeters, of the water in the glass?

Worked solution

1. The water forms a smaller cone similar to the cup, with height 8: its radius is $6 \cdot \frac{8}{12} = 4$.
2. Water volume $= \frac{1}{3}\pi(4)^2(8) = \frac{128\pi}{3}$.
3. In the glass, $\pi(4)^2d = \frac{128\pi}{3} \Rightarrow 16d = \frac{128}{3}$.
4. $d = \frac{8}{3} \approx 2.67$.

Answer: $\frac{8}{3}$

25.

SAT CHALLENGE · DESMOS

In the xy -plane, point P lies on the line $y = 2x + 1$ and is the same distance from $A(-1, 6)$ as from $B(5, 0)$. What is the area of triangle ABP ?

Worked solution


1. P lies on the perpendicular bisector of $\overline{AB}$. Midpoint of $\overline{AB}$: $(2, 3)$; slope of $\overline{AB} = \frac{0-6}{5-(-1)} = -1$, so the bisector has slope 1: $y = x + 1$.
2. Intersect with $y = 2x + 1$: $x + 1 = 2x + 1 \Rightarrow x = 0$, so $P = (0, 1)$.
3. Shoelace with $A(-1, 6)$, $B(5, 0)$, $P(0, 1)$: $(-1)(0) - (5)(6) = -30$; $(5)(1) - (0)(0) = 5$; $(0)(6) - (-1)(1) = 1$. Sum $= -24$.
4. Area $= \frac{1}{2}|-24| = 12$.

Answer: 12

26.

SAT CHALLENGE · CALCULATOR

A right circular cylinder is inscribed in a sphere with radius 13 so that the circumference of each base of the cylinder lies on the sphere. The base of the cylinder has radius 5. The figure shows a cross section through the center O of the sphere. What is the volume of the cylinder?

(A) 600π ✓(B) 650π (C) 300π (D) 325π **Worked solution**

1. In the cross section, a radius of the sphere to a corner of the rectangle is the hypotenuse of a right triangle with legs 5 (the base radius) and half the cylinder's height.
2. Half-height = $\sqrt{13^2 - 5^2} = \sqrt{144} = 12$, so the height is 24.
3. $V = \pi r^2 h = \pi(25)(24) = 600\pi$.

Answer: (A)

27.

SAT CHALLENGE · NO CALCULATOR NEEDED

A scale model of a water tank holds 8 liters of water. The actual tank holds 27 cubic meters of water, and the model is geometrically similar to the actual tank. The model is 36 centimeters tall. How tall, in meters, is the actual tank? (1 cubic meter = 1000 liters)

Worked solution


1. Actual capacity: $27 \text{ m}^3 = 27,000$ liters.
2. Volume ratio = $\frac{27,000}{8} = 3375 = k^3$, so $k = 15$.
3. Actual height = $36 \cdot 15 = 540 \text{ cm} = 5.4 \text{ m}$.

Answer: 5.4

28.

SAT CHALLENGE · CALCULATOR

A rectangular garden has a width of w meters and a length that is 7 meters greater than its width. A path of uniform width 2 meters surrounds the garden on all sides. The area of the path alone is 156 square meters. What is the area, in square meters, of the garden?



(A) 98

(B) 294 ✓

(C) 450

(D) 588

Worked solution

1. The outer rectangle is $(w + 4)$ by $(w + 11)$ because the path adds 2 meters on each side.

2. Path area = $(w + 4)(w + 11) - w(w + 7) = (w^2 + 15w + 44) - (w^2 + 7w) = 8w + 44$.

3. $8w + 44 = 156 \Rightarrow w = 14$.


4. Garden area = $w(w + 7) = 14 \cdot 21 = 294$.

Answer: (B)

29.

SAT CHALLENGE · CALCULATOR

In the xy -plane, triangle OQR has vertices $O(0,0)$, $Q(8,0)$ and $R(8,6)$, as shown. The triangle is rotated 360° about the line containing side $\overline{OR}$ to form a solid. What is the volume of the solid?

(A) 128π (B) $\frac{1152}{5}\pi$ (C) $\frac{384}{5}\pi$ ✓(D) 96π **Worked solution**

1. $OR = \sqrt{8^2 + 6^2} = 10$. The solid is two cones sharing a base circle whose radius is the distance from Q to line OR .

2. That radius is the altitude to the hypotenuse: $\frac{8 \cdot 6}{10} = 4.8$.

3. The two cones have heights that add to $OR = 10$, so $V = \frac{1}{3}\pi(4.8)^2(10)$.

4. $V = \frac{1}{3}\pi(23.04)(10) = 76.8\pi = \frac{384}{5}\pi$.

Answer: (C)

30.

SAT CHALLENGE · CALCULATOR

A solid metal sphere with radius 9 centimeters is melted. Exactly 90% of the metal is recast into identical solid cones, each with radius 3 centimeters and height 4 centimeters. What is the greatest number of complete cones that can be made?

Worked solution

1. Sphere volume $= \frac{4}{3}\pi(9)^3 = 972\pi$.

2. Metal used: $0.9 \cdot 972\pi = 874.8\pi$.

3. One cone: $\frac{1}{3}\pi(3)^2(4) = 12\pi$.

4. $\frac{874.8\pi}{12\pi} = 72.9$, so at most 72 complete cones can be made.

Answer: 72