

SAT Math Mastery · Week 9B**Capstone: Mixed Strategy & Timed Modules — Answer Key & Worked Solutions****Time** 120 minutes: 90 min solving + 30 min self-check and error log**Topics** mixed algebra, advanced math, data analysis and geometry · pick-a-number · back-solving · Desmos · eliminate by structure · pacing and flagging · the seven shared reasoning moves**Name / Date** _____**1 · SKILLS & IDEAS**

- S1.** Classify a problem in the first ten seconds: algebra, advanced math, data, or geometry, and choose the cheapest method.
- S2.** Pick-a-number: test equivalent expressions, percent chains and “for all x ” statements with simple values.
- S3.** Back-solving: plug the answer choices into the original condition, starting with the easiest value to test.
- S4.** Desmos: intersections, sliders for parameters, tangency, tables for exponential and sequence questions.
- S5.** Eliminate by structure: signs, intercepts, end behavior, parity, units, domain, and “could NOT be” tests.
- S6.** Shared moves: structure and solution counts (M1), feature $\leftrightarrow$ form (M2), solve for a constant (M3), interpret (M4), scaling and units (M5), test a candidate (M6), extraneous and boundary care (M7).
- S7.** Run a timed two-pass module: answer, flag, return; never leave a question blank.
- S8.** Self-check habits: reread what is asked, check units and rounding, plug the answer back in.

2 · REQUIRED FACTS

- Digital SAT Math: 2 modules of 22 questions, 35 minutes each. The second module’s difficulty depends on the first; there is no penalty for a wrong answer.
- A system of two lines has no solution if the slopes are equal and the intercepts differ, and infinitely many if the two equations are the same line.
- A quadratic $ax^2 + bx + c$ with $a > 0$ has a minimum at $x = -\frac{b}{2a}$; its discriminant $b^2 - 4ac$ decides 2, 1 or 0 real solutions.
- $y = a \cdot b^t$ multiplies by b for every 1 unit of t ; $b > 1$ is growth and $0 < b < 1$ is decay.
- Mean is pulled toward outliers; median is not; spread (standard deviation) grows when values are farther from the mean.
- In a right triangle, $\sin \theta = \frac{\text{opp}}{\text{hyp}}$, $\cos \theta = \frac{\text{adj}}{\text{hyp}}$, $\tan \theta = \frac{\text{opp}}{\text{adj}}$; areas of similar figures scale by k^2 .
- Squaring both sides of an equation can create extraneous solutions: always test in the original equation.

3 · REQUIRED FORMULAS

$$y = mx + b, \quad (x - h)^2 + (y - k)^2 = r^2, \quad y = a(x - h)^2 + k, \quad x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$\text{percent change} = \frac{\text{new} - \text{old}}{\text{old}} \times 100, \quad \text{mean} = \frac{\text{sum}}{n}, \quad P(A | B) = \frac{\text{count in both}}{\text{count in } B}$$

$$\text{arc} = r\theta, \quad \text{sector area} = \frac{1}{2}r^2\theta = \frac{1}{2}rL, \quad a^2 + b^2 = c^2, \quad V_{\text{cyl}} = \pi r^2 h, \quad V_{\text{cone}} = \frac{1}{3}\pi r^2 h$$

4 · MINI THEORY SHEET

Pacing table (this 30-problem sheet = 90 minutes of solving).

Tier	Items	Target per item	Block time	Running clock at end of block
Easy	1–5	1 min	5 min	5 min
Medium	6–13	2 min	16 min	21 min
Hard	14–23	3 min	30 min	51 min
SAT Challenge	24–30	$5\frac{1}{2}$ min	38 min	89 min

On the real test one module averages $35 \div 22 \approx 1.6$ minutes per question. Rule: if you have spent twice the target time and have no plan, flag the question, mark a guess, and move on. Use the last 5 minutes of a module for flagged questions. A blank answer scores nothing; a guess can score.

Which method first?

Signal in the problem	First move
variables in the answer choices; “equivalent to”; “for all x ”	pick-a-number : test a simple value in the stem and in every choice
percent chain with no starting amount	pick 100 (or 10 and a ratio)
numeric choices and a messy equation	back-solve : plug a choice into the stem
intersection, “how many solutions”, a constant that makes a line tangent	Desmos : graph, add a slider for the constant
table or two points of an exponential or sequence	Desmos table or ratio of outputs
graph or sign conditions	eliminate by structure : sign of a , intercepts, vertex, end behavior
“could NOT be”; “must be true”	test candidates against every condition; one counterexample kills a statement
units, “nearest whole”, “at least”	convert first; round in the direction that satisfies the condition

Eliminate by structure. Before computing, ask which choices have the wrong sign, the wrong intercept, the wrong units, the wrong degree or the wrong order of magnitude. Cross them out first.

Pick-a-number rules. Avoid 0, 1 and numbers that make two choices agree; check *every* choice, because two can match one value.

Back-solving rules. Start with the choice that is easiest to compute; if it is too big or too small, you know which way to go.

Desmos rules. Type each equation exactly as written; use a slider for any constant; use the intersection points; check that the window shows every intersection.

The seven shared moves.

Move	Typical question
M1 count of solutions $\Leftrightarrow$ structure	equal slopes, discriminant = 0, same line
M2 feature $\leftrightarrow$ form	vertex form for the vertex, factored for the roots
M3 solve for a constant	use a given point or condition to find k
M4 interpret, don’t compute	meaning of a slope, intercept or residual
M5 scaling and units	squares, cubes, rates, conversions
M6 test a candidate against all conditions	inequality systems, tables, “which could be”
M7 extraneous and boundary care	radicals, whole-number rounding, strict inequalities

5 · GUIDED EXAMPLES

Example 1 — Pick-a-number (equivalent expressions)

For $x \neq 2$, which expression is equivalent to $\frac{3x^2 - x - 10}{x - 2}$? (A) $3x + 5$ (B) $3x - 5$ (C) $3x + 2$ (D) $3x - 10$

Pick $x = 3$: the stem gives $\frac{27 - 3 - 10}{1} = 14$. Choices: (A) 14, (B) 4, (C) 11, (D) -1 . Only (A) matches.

Algebra confirms it: $3x^2 - x - 10 = (3x + 5)(x - 2)$.

Example 2 — Back-solving (radical equation)

Which value of x satisfies $\sqrt{x + 9} = x - 3$? (A) 0 (B) 4 (C) 7 (D) 9

Test (C): $\sqrt{7 + 9} = 4$ and $7 - 3 = 4$ ✓. Test (A): $\sqrt{9} = 3$ but $0 - 3 = -3 \neq 3$ (extraneous). (B): $\sqrt{13} \neq 1$. (D): $\sqrt{18} \neq 6$. Answer: (C).

Example 3 — Desmos with a slider (tangency)

The line $y = kx - 7$ touches the parabola $y = x^2 - 2x - 3$ at exactly one point, and $k > 0$. Find k .

Graph the parabola and $y = kx - 7$ with a slider for k ; they touch when the discriminant of $x^2 - (2 + k)x + 4 = 0$ is 0: $(2 + k)^2 - 16 = 0 \Rightarrow k = 2$ or $k = -6$.

With $k > 0$: $k = 2$. (The touch point is $x = 2$, $y = -3$.)

6 · PRACTICE SET — 30 PROBLEMS · EASY → MEDIUM → HARD → SAT CHALLENGE

Each problem states its tool. Write your work in the space below. Multiple choice: one correct option. Student-produced response (SPR): enter an exact value (fraction or decimal).

1. EASY · NO CALCULATOR NEEDED
If $4x - 7 = 21$, what is the value of $4x + 7$?

- (A) 28 **(B) 35 ✓**
(C) 7 (D) 14

Worked solution

1. Add 14 to both sides of the stem equation: $4x - 7 + 14 = 21 + 14$.
2. $4x + 7 = 35$. (Check: $4x = 28$, so $x = 7$ and $4(7) + 7 = 35$.)

Answer: (B)

2. EASY · CALCULATOR
The function f is defined by $f(x) = 2x^2 - 3x + 1$. What is the value of $f(-2)$?

Worked solution

1. $f(-2) = 2(-2)^2 - 3(-2) + 1$.
2. $= 2(4) + 6 + 1 = 8 + 6 + 1 = 15$.

Answer: 15


3. EASY · NO CALCULATOR NEEDED
The mean of five numbers is 12. Four of the numbers are 8, 10, 14 and 15. What is the fifth number?

Worked solution

1. Sum of all five = $5 \cdot 12 = 60$.
2. Sum of the four given numbers = $8 + 10 + 14 + 15 = 47$.
3. Fifth number = $60 - 47 = 13$.

Answer: 13

4. EASY · NO CALCULATOR NEEDED
In right triangle ABC shown, $\angle C = 90^\circ$, $AB = 17$ and $BC = 8$. What is the value of $\sin A$?



- (A) $\frac{8}{15}$ (B) $\frac{17}{8}$
(C) $\frac{8}{17}$ ✓ (D) $\frac{15}{17}$

Worked solution

1. $\sin A = \frac{\text{side opposite } A}{\text{hypotenuse}}$.
2. The side opposite $\angle A$ is $BC = 8$ and the hypotenuse is $AB = 17$.
3. $\sin A = \frac{8}{17}$.

Answer: (C)

5.

EASY · NO CALCULATOR NEEDED

For $x \neq -2$, which of the following expressions is equivalent to $\frac{2x^2 + x - 6}{x + 2}$?

(A) $2x - 3$ ✓

(B) $2x + 3$

(C) $2x - 6$

(D) $x - 3$

Worked solution

1. Factor the numerator: $2x^2 + x - 6 = (2x - 3)(x + 2)$.

2. Cancel $(x + 2)$: $\frac{(2x - 3)(x + 2)}{x + 2} = 2x - 3$.

3. (Pick $x = 1$: the stem gives $\frac{-3}{3} = -1$, and only $2x - 3$ equals -1 .)

Answer: (A)

6.

MEDIUM · NO CALCULATOR NEEDED

$$3x - 2y = 10$$

$$kx + 4y = 7$$

In the system of equations above, k is a constant. If the system has no solution, what is the value of k ?

(A) -3

(B) 6

(C) $\frac{3}{2}$

(D) -6 ✓

Worked solution

1. No solution means the lines are parallel and distinct: the slopes are equal.

2. Slopes: $\frac{3}{2}$ and $-\frac{k}{4}$. Set equal: $-\frac{k}{4} = \frac{3}{2} \Rightarrow k = -6$.

3. The right-hand sides are not in the same ratio ($\frac{10}{7} \neq -\frac{1}{2}$), so the lines are distinct and there is no solution.

Answer: (D)

7.

MEDIUM · CALCULATOR

A theater sold 40 tickets for a show. Adult tickets cost \$12 each and student tickets cost \$8 each. The total sales were \$392. How many adult tickets were sold?

(A) 12

(B) 16

(C) 18 ✓

(D) 22

Worked solution

1. Let a be the number of adult tickets, so $40 - a$ are student tickets.

2. $12a + 8(40 - a) = 392 \Rightarrow 4a + 320 = 392 \Rightarrow a = 18$.


3. (Back-solve: $a = 18$ gives $216 + 176 = 392$ ✓.)

Answer: (C)

8.

MEDIUM · DESMOS

The graph of a quadratic function in the xy -plane is shown. Which of the following could be an equation of the graph?



(A) $y = (x - 1)^2 - 4$

(B) $y = -(x - 1)^2 + 4$ ✓

(C) $y = -(x + 1)^2 + 4$

(D) $y = -(x - 4)^2 + 1$

Worked solution

1. The graph opens downward, so the coefficient of the squared term is negative. This eliminates $y = (x - 1)^2 - 4$.
2. The vertex is $(1, 4)$, so the form is $y = -(x - 1)^2 + 4$.
3. Check an x -intercept: at $x = 3$, $-(2)^2 + 4 = 0$ ✓. The other choices fail at the vertex or the intercepts.

Answer: (B)

9.

MEDIUM · NO CALCULATOR NEEDED

A bacteria culture starts with 120 cells and doubles every 6 hours. The number of cells t hours after the start is modeled by $N(t) = 120 \cdot 2^{t/6}$. After how many hours will the culture have 1920 cells?

Worked solution

1. Set $120 \cdot 2^{t/6} = 1920$, so $2^{t/6} = 16$.
2. $16 = 2^4$, so $\frac{t}{6} = 4$.
3. $t = 24$.

Answer: 24

10.

MEDIUM · CALCULATOR

A survey asked 200 students whether they prefer online or in-person classes. The results are shown in the table.

	Online	In-person	Total
Grade 11	48	32	80
Grade 12	66	54	120
Total	114	86	200

If one of the students who prefer in-person classes is selected at random, what is the probability that the student is in Grade 12?

(A) $\frac{9}{20}$

(B) $\frac{27}{100}$

(C) $\frac{3}{5}$

(D) $\frac{27}{43}$ ✓

Worked solution

1. Restrict to the in-person column: $32 + 54 = 86$ students.

2. Grade 12 within that column: 54.

3. $P = \frac{54}{86} = \frac{27}{43}$.

Answer: (D)

11.

MEDIUM · NO CALCULATOR NEEDED

The value of p is 40% greater than the value of q , and the value of q is 25% less than the value of r , where $r > 0$. The value of p is what percent of the value of r ?

Worked solution

1. Pick $r = 100$. Then $q = 100 - 25 = 75$.

2. $p = 75 + 0.40(75) = 105$.

3. $\frac{p}{r} = \frac{105}{100} = 105\%$.

Answer: 105

12.

MEDIUM · NO CALCULATOR NEEDED

The equation $x^2 + y^2 - 6x + 4y = 12$ defines a circle in the xy -plane. What is the radius of the circle?

(A) 5 ✓

(B) 25

(C) $\sqrt{17}$

(D) 12

Worked solution

1. Complete the square: $(x^2 - 6x + 9) + (y^2 + 4y + 4) = 12 + 9 + 4$.

2. $(x - 3)^2 + (y + 2)^2 = 25$.

3. $r^2 = 25$, so $r = 5$.

Answer: (A)

13.

MEDIUM · NO CALCULATOR NEEDED

In right triangle ABC , $\angle B = 90^\circ$, $\sin A = \frac{3}{5}$ and $AC = 20$. What is the length of $\overline{BC}$?

(A) 25

(B) 12 ✓

(C) 16

(D) 15

Worked solution

1. $\overline{AC}$ is the hypotenuse. $\sin A = \frac{BC}{AC} = \frac{3}{5}$.

2. $BC = \frac{3}{5} \cdot 20 = 12$.

3. (The other leg is $AB = 16$, since $12^2 + 16^2 = 20^2$.)

Answer: (B)

14.

HARD · NO CALCULATOR NEEDED

$$(a - 2)x + 3 = 5x + b$$

In the equation above, a and b are constants. If the equation has infinitely many solutions for x , what is the value of $a + b$?

Worked solution

1. Infinitely many solutions means the two sides are identical expressions in x .

2. Match the x -coefficients: $a - 2 = 5 \Rightarrow a = 7$. Match the constants: $3 = b$.

3. $a + b = 7 + 3 = 10$.

Answer: 10

15.

HARD · NO CALCULATOR NEEDED

$$y < -x + 7$$

$$y > 2x - 5$$

Which of the following ordered pairs (x, y) is a solution to the system of inequalities above?

(A) (1, 6)

(B) (0, -6)

(C) (3, 3) ✓

(D) (4, 2)

Worked solution

1. Test each pair in both inequalities.

2. (3, 3): $3 < -3 + 7 = 4$ ✓ and $3 > 2(3) - 5 = 1$ ✓.

3. (4, 2): $2 > 3$ is false. (1, 6): $6 < 6$ is false (strict). (0, -6): $-6 > -5$ is false.

Answer: (C)

16.

HARD · DESMOS

$$y = x^2 - 6x + c$$

$$y = 2x - 7$$

In the system above, c is a constant. If the system has exactly one solution, what is the value of c ?

Worked solution

1. Set the expressions equal: $x^2 - 6x + c = 2x - 7 \Rightarrow x^2 - 8x + (c + 7) = 0$.
2. Exactly one solution means the discriminant is 0: $(-8)^2 - 4(1)(c + 7) = 0$.
3. $64 = 4(c + 7) \Rightarrow c + 7 = 16 \Rightarrow c = 9$.

Answer: 9

17.

HARD · NO CALCULATOR NEEDED

What is the sum of all solutions to the equation $\sqrt{3x + 4} = x$?

(A) 4 ✓

(B) 5

(C) -1

(D) 3

Worked solution

1. Square both sides: $3x + 4 = x^2 \Rightarrow x^2 - 3x - 4 = 0 \Rightarrow (x - 4)(x + 1) = 0$.
2. Test $x = 4$: $\sqrt{16} = 4$ ✓. Test $x = -1$: $\sqrt{1} = 1 \neq -1$, so $x = -1$ is extraneous.
3. The only solution is 4, so the sum is 4.

Answer: (A)

18.

HARD · CALCULATOR

The function $f(t) = a \cdot b^t$, where a and b are positive constants, satisfies $f(1) = 12$ and $f(4) = 324$. What is the value of $f(6)$?

(A) 8748

(B) 972

(C) 1458

(D) 2916 ✓

Worked solution


1. $\frac{f(4)}{f(1)} = b^3 = \frac{324}{12} = 27$, so $b = 3$.
2. $f(1) = 3a = 12 \Rightarrow a = 4$.
3. $f(6) = 4 \cdot 3^6 = 4 \cdot 729 = 2916$.

Answer: (D)

19.

HARD · DESMOS

The scatterplot shows the number of hours studied, x , and the test score, y , for 10 students. The line of best fit shown passes through $(2, 60)$ and $(8, 90)$. The residual of a data point is its actual score minus the score predicted by the line. For which number of hours studied is the absolute value of the residual greatest?



(A) 10

(B) 2

(C) 6 ✓

(D) 9

Worked solution

1. Slope of the line: $\frac{90 - 60}{8 - 2} = 5$; $y = 5x + 50$.

2. Residual = actual – predicted: $x = 1$: $60 - 55 = 5$; $x = 2$: $55 - 60 = -5$; $x = 3$: $70 - 65 = 5$; $x = 4$: $75 - 70 = 5$; $x = 5$: $70 - 75 = -5$.

3. $x = 6$: $90 - 80 = 10$; $x = 7$: $80 - 85 = -5$; $x = 8$: $85 - 90 = -5$; $x = 9$: $100 - 95 = 5$; $x = 10$: $95 - 100 = -5$.

4. The largest absolute residual is 10, at 6 hours.

Answer: (C)

20.

HARD · NO CALCULATOR NEEDED

Data set A is 2, 4, 6, 8, 10. Data set B is 2, 4, 6, 8, 40. Which of the following statements is true?

I. The mean of B is greater than the mean of A .

II. The median of B is greater than the median of A .

III. The standard deviation of B is greater than the standard deviation of A .

(A) I only

(B) I and III only ✓

(C) II and III only

(D) I, II, and III

Worked solution

1. I: mean of $A = 6$, mean of $B = \frac{60}{5} = 12$. True.

2. II: both medians are 6. False.

3. III: the value 40 is far from the mean, so B is much more spread out than A . True.

4. Statements I and III only.

Answer: (B)

21.

HARD · CALCULATOR

A machine fills 15 bottles every 40 seconds. At this rate, how many hours will it take the machine to fill 4050 bottles?

Worked solution

1. Number of 40-second cycles: $\frac{4050}{15} = 270$.

2. Time = $270 \cdot 40 = 10,800$ seconds.


3. $\frac{10,800}{3600} = 3$ hours.

Answer: 3

22.

HARD · CALCULATOR

From point A on level ground, the angle of elevation to the top of a vertical tower is 30° . From point B , which is 20 meters closer to the base of the tower and on the same line, the angle of elevation is 45° . What is the height of the tower, in meters?



(A) 20

(B) $10(\sqrt{3} - 1)$ (C) $10\sqrt{3}$ (D) $10(\sqrt{3} + 1)$ ✓**Worked solution**

1. Let the height be h and let d be the distance from B to the base. At B : $\tan 45^\circ = \frac{h}{d} = 1$, so $d = h$.

2. At A : $\tan 30^\circ = \frac{h}{h + 20} = \frac{1}{\sqrt{3}} \Rightarrow \sqrt{3}h = h + 20$.

3. $h(\sqrt{3} - 1) = 20 \Rightarrow h = \frac{20}{\sqrt{3} - 1} = \frac{20(\sqrt{3} + 1)}{2} = 10(\sqrt{3} + 1) \approx 27.3$.

Answer: (D)

23.

HARD · NO CALCULATOR NEEDED

In triangle ABC , $AB = AC = 15$ and $BC = 24$. Point D lies on $\overline{AC}$ such that $\overline{BD} \perp \overline{AC}$. What is the length of $\overline{BD}$?

Worked solution

1. Drop the altitude from A to $\overline{BC}$: it bisects $\overline{BC}$, so $h = \sqrt{15^2 - 12^2} = 9$.

2. Area of $\triangle ABC = \frac{1}{2}(24)(9) = 108$.

3. Using $\overline{AC}$ as the base: $\frac{1}{2}(15)(BD) = 108 \Rightarrow BD = \frac{216}{15} = 14.4$.

Answer: 14.4

24.

SAT CHALLENGE · NO CALCULATOR NEEDED

$$(k + 1)x + 6y = 18$$

$$4x + (k - 1)y = 12$$

In the system above, k is a constant. For which value of k does the system have no solution?

Worked solution

1. The determinant condition for parallel lines: $(k + 1)(k - 1) = 6 \cdot 4 \Rightarrow k^2 - 1 = 24 \Rightarrow k = 5$ or $k = -5$.
2. $k = 5$: the equations are $6x + 6y = 18$ and $4x + 4y = 12$; both reduce to $x + y = 3$, the same line, so infinitely many solutions.
3. $k = -5$: $-4x + 6y = 18$ and $4x - 6y = 12$. Adding gives $0 = 30$, which is impossible, so no solution.
4. The value is $k = -5$.

Answer: -5

25.

SAT CHALLENGE · NO CALCULATOR NEEDED

A jar contains only red, blue and green marbles. The ratio of red to blue marbles is $3 : 5$, and the ratio of blue to green marbles is $2 : 3$. There are 36 more green marbles than red marbles. How many marbles are in the jar?

Worked solution

1. Make the blue part match: red : blue = $3 : 5 = 6 : 10$ and blue : green = $2 : 3 = 10 : 15$.
2. So red : blue : green = $6 : 10 : 15$. Let the parts be $6u$, $10u$, $15u$.
3. Green – red = $15u - 6u = 9u = 36 \Rightarrow u = 4$.
4. Total = $(6 + 10 + 15)u = 31 \cdot 4 = 124$.

Answer: 124

26.

SAT CHALLENGE · NO CALCULATOR NEEDED

The function $f(x) = x^2 + bx + c$ has a minimum value of -25 , and $f(0) = -21$. What is the greatest possible value of b ?

Worked solution

1. $f(0) = c = -21$.
2. The minimum value is $c - \frac{b^2}{4}$ (at $x = -\frac{b}{2}$): $-21 - \frac{b^2}{4} = -25 \Rightarrow b^2 = 16$.
3. $b = 4$ or $b = -4$; the greatest possible value is 4.

Answer: 4

27.

SAT CHALLENGE · NO CALCULATOR NEEDED

For what value of x is $2^{x+3} - 2^x = 56$?

(A) 3 ✓

(B) 4

(C) 7

(D) 8

Worked solution

1. Factor: $2^{x+3} - 2^x = 2^x(2^3 - 1) = 7 \cdot 2^x$.
2. $7 \cdot 2^x = 56 \Rightarrow 2^x = 8$.
3. $x = 3$. (Check: $2^6 - 2^3 = 64 - 8 = 56$.)

Answer: (A)

28.

SAT CHALLENGE · NO CALCULATOR NEEDED

The mean of a list of 9 numbers is 20. Two numbers are removed from the list; one of them is twice the other. The mean of the remaining 7 numbers is 18. What is the larger of the two numbers that were removed?

Worked solution


1. Sum of the original list $= 9 \cdot 20 = 180$. Sum of the remaining 7 $= 7 \cdot 18 = 126$.
2. Sum of the two removed numbers $= 180 - 126 = 54$.
3. Let them be x and $2x$: $3x = 54 \Rightarrow x = 18$.
4. The larger number is $2x = 36$.

Answer: 36

29.

SAT CHALLENGE · NO CALCULATOR NEEDED

In triangle ABC , point D lies on $\overline{AB}$ and point E lies on $\overline{AC}$ so that $\overline{DE} \parallel \overline{BC}$, as shown. $AD = 6$, $DB = 4$, and the area of triangle ADE is 18. What is the area of quadrilateral $DBCE$?

**Worked solution**

1. $AB = AD + DB = 10$. Since $\overline{DE} \parallel \overline{BC}$, $\triangle ADE \sim \triangle ABC$ with ratio $\frac{AD}{AB} = \frac{6}{10} = \frac{3}{5}$.
2. Areas scale by the square: $\frac{[ADE]}{[ABC]} = \frac{9}{25}$.
3. $[ABC] = 18 \cdot \frac{25}{9} = 50$.
4. $[DBCE] = 50 - 18 = 32$.

Answer: 32

30.

SAT CHALLENGE · CALCULATOR

A sector of a circle has area 20π and arc length 4π . What is the measure, in radians, of the central angle of the sector?

(A) $\frac{\pi}{5}$

(B) $\frac{2\pi}{5}$ ✓

(C) $\frac{4\pi}{5}$

(D) $\frac{\pi}{2}$

Worked solution

1. Sector area $= \frac{1}{2}rL$ where L is the arc length: $\frac{1}{2}r(4\pi) = 20\pi \Rightarrow r = 10$.
2. Arc length $L = r\theta \Rightarrow 4\pi = 10\theta$.
3. $\theta = \frac{4\pi}{10} = \frac{2\pi}{5}$.

Answer: (B)