



Children with Chromosome 22q11.2 Deletion Syndrome Have Smaller Putamen Than Typically Developing Children

Blake Manale, Alexis Daigle, B.S., Diana A. Hobbs, M.S., David D. Stephenson, M.A.,
Ashley F.P. Sanders, M.S., Michael J. Medina, B.S., & Elliott A. Beaton, Ph.D.
University of New Orleans, New Orleans, LA



Chromosome 22q11.2 Deletion Syndrome

- Chromosome 22q11.2 deletion syndrome (22q11.2DS) includes DiGeorge and Velocardiofacial (VCFS) syndromes.
- Occurs in between 1:2000 and 1:4000 live births via a deletion of 3Mb pairs on the long (q) arm at the 11.2 band of one copy of chromosome 22.
- 25 to 30% of children with 22q11.2DS will develop schizophrenia in adulthood. 75% have mild prodromal psychotic symptoms.
- The putamen is part of the striatum that plays a role in motor processes and learning.
- Smaller putamen volumes are associated with poorer prognosis in patients with schizophrenia.
- Recent findings suggest the putamen may affect psychotic action as it regulates dopamine.

Overview and Purpose

- Putamen volume changes over time may serve as a biomarker of schizophrenia risk.
- This is the first measure of a longitudinal study.
- Using MRI anatomical brain scans, we investigated size differences in putamen volumes between children with 22q11.2DS and those who are typically developing (TD).

Participants

- Boys and girls ages 7 to 16 years in two groups.
- 22q11.2DS: $n = 15$, mean age = 12.06 years, $SD = 2.85$
- TD: $n = 15$, mean age = 11.91 years, $SD = 2.51$

Hypothesis

- Children with 22q11.2DS will have smaller putamen compared to typically-developing controls.

Brain Imaging Method

- T1-weighted high-resolution 3D anatomical magnetic resonance images were acquired using a 3.0T Siemens Trio scanner with Echospeed gradients and a Siemens 8-channel whole head coil.
- Imaging parameters: TR = 2.17 s, TE = 4.82 ms, flip angle = 7°, NEX = 1, 192 slices, 128 x 128 acquisition matrix, FOV = 256, sagittal acquisition with 1 mm³ voxels, 5 min 4 sec duration.

Putamen Volume Measures

- The left and right putamen were manually outlined on each MR image slice with the tracers blind to diagnosis.
- For all steps of this process, Mango (Research Imaging Institute, U of Texas) was used for region-of-interest parcellation and volume calculations.




Figure 1— High-resolution T1-weighted MRI transverse image showing the region of interest. The putamen is outlined in red.

- T1-weighted whole brain images are imported into Mango (Research Imaging Institute, U of Texas).
- Putamen volume was calculated via manual slice by slice tracings of histology-based neuroanatomical regions of sagittal images in native space.
- Tracing Axial Criteria:
 - Crosshair Posterior Commissure.
 - Crosshair Anterior Commissure.
- Tracing Midline criteria:
 - Visible globus pallidus.
 - External capsule laterally bounds putamen.
 - Internal capsule medially and superiorly bounds putamen.
 - Anterior commissure inferiorly bounds putamen.

Results

- For total putamen volume, a significant difference was found between 22q11.2DS ($M = 7956.2 \text{ mm}^3$, $SD = 2214.77$) and TD groups ($M = 9984.47 \text{ mm}^3$, $SD = 1400.61$; $t(28) = 3.10$, $p < 0.01$).
- Right putamen was smaller in the 22q11.2DS group ($M = 4025.2 \text{ mm}^3$, $SD = 1107.28$) versus the TD group ($M = 5088.4 \text{ mm}^3$, $SD = 753.52$; $t(28) = 3.07$, $p < 0.01$).
- Left putamen was smaller in the 22q11.2DS groups ($M = 3931 \text{ mm}^3$, $SD = 1016.57$) versus the TD group ($M = 4896.07 \text{ mm}^3$, $SD = 673.35$; $t(28) = 3.07$, $p < 0.01$).
- Within the 22q11.2DS group, there was no difference between right and left putamen volumes ($p > 0.1$).




Figure 2 — Group mean (\pm SEM) bilateral putamen volumes.

Summary and Implications

- Children with 22q11.2DS in this study have smaller putamen than TD age-matched children.
- Smaller putamen volumes are reported in patients with schizophrenia.
- Lower volumes could be an indicator of general brain atrophy, atypical development, or both; however, given the ultra-high risk of schizophrenia, further investigation of putamen morphometry over time and in relation to prodromal symptoms is needed.

References

Bassett, A. S., Marshall, C. R., Lionel, A. C., Chow, E. W., & Scherer, S. W. (2008). Copy number variations and risk for schizophrenia in 22q11. 2 deletion syndrome. *Human Molecular Genetics*, 17(24), 4045-4053.

Buchsbaum, Monte, Shihabuddin, L., Brickman, A.M., Miozzo, R., Prikrly, R., Shaw, R., & Davis, K., (2003). Caudate and putamen volumes in good and poor outcome patients with schizophrenia. *Schizophrenia Research*, 64(1), 53-62..

Cincotta, C. M., & Seger, C. A. (2007). Dissociation between striatal regions while learning to categorize via feedback and via observation. *Journal of Cognitive Neuroscience*, 19(2), 249-265.

Williams, M. (2015). An introduction to the putamen in schizophrenia. *Orien-The CNS Journal*, 1 (2), 23-25.