

Lab Exercise #1

Dianene Khrysna G. Ticot

2025-09-29

#1a.

```
vector1 <- c (-5:5)
vector1
```

```
## [1] -5 -4 -3 -2 -1 0 1 2 3 4 5
```

#the output is a numbers from -5 to 5

#1b.

```
x <- 1:7
x
```

```
## [1] 1 2 3 4 5 6 7
```

#the output is a numbers from 1 to 7

#2

```
seq1 <- seq(1, 3, by=0.2)
seq1
```

```
## [1] 1.0 1.2 1.4 1.6 1.8 2.0 2.2 2.4 2.6 2.8 3.0
```

#the output is a sequence from 1 to 3 in steps of 0.2

#3

```
ages <- c(34, 28, 22, 36, 27, 18, 52, 39, 42, 29, 35, 31, 27,
22, 37, 34, 19, 20, 57, 49, 50, 37, 46, 25, 17, 37, 43, 53, 41, 51, 35,
24, 33, 41, 53, 40, 18, 44, 38, 41, 48, 27, 39, 19, 30, 61, 54, 58, 26,
18.)
```

#3a.

```
ages [3]
```

```
## [1] 22
```

#the value is 22

#3b.

```
ages [c(2,4)]
```

```
## [1] 28 36
```

#the values are 28 and 26

#3c.

```
ages[-1]
```

```
## [1] 28 22 36 27 18 52 39 42 29 35 31 27 22 37 34 19 20 57 49 50 37 46 25 17 37
## [26] 43 53 41 51 35 24 33 41 53 40 18 44 38 41 48 27 39 19 30 61 54 58 26 18
```

```
#all values except 34
```

```
#4
```

```
x<- c("first" = 3, "second" = 0, "third" = 9)
x
```

```
## first second third
##      3      0      9
```

```
#4a
```

```
x[c("first", "third")]
```

```
## first third
##      3      9
```

```
#the output is the elements with their names
```

```
#4b(RESULT)
```

```
#>
```

```
# > #a
```

```
# > x[c("first", "third")]
```

```
#first third
```

```
#3      9
```

```
#> #4
```

```
# > x<- c("first" = 3, "second" = 0, "third" = 9)
```

```
#> x
```

```
#first second third
```

```
#3      0      9
```

```
#>
```

```
# > #a
```

```
# > x[c("first", "third")]
```

```
#first third
```

```
#3      9
```

```
#> #the output is the elements with their names
```

```
# >
```

```
#5
```

```
x <- seq(-3:2)
```

```
#5a
```

```
x[2] <-0
```

```
x
```

```
## [1] 1 0 3 4 5 6
```

```
#a sequence from -3 to 2 where the second element was replaced with zero
```

```
#5b(RESULT)
```

```
#> #5
```

```
# > x <- seq(-3:2)
```

```
#>
```

```
# > #5a
```

```
# > x[2] <-0
```

```
#> x
```

```
#[1] 1 0 3 4 5 6
```

```
#>

#6a
month <- c("Jan", "Feb", "March", "Apr", "May", "June")
price <- c(52.50, 57.56, 60.00, 65.00, 74.25, 54.00)
purchase <- c(25, 30, 40, 50, 10, 45)

fuel.data <- data.frame(month, price, purchase)
fuel.data
```

```
##   month price purchase
## 1   Jan  52.50      25
## 2   Feb  57.56      30
## 3 March  60.00      40
## 4   Apr  65.00      50
## 5   May  74.25      10
## 6   June 54.00      45
```

```
#6b
avg_expenditure <- weighted.mean(price, purchase)
avg_expenditure
```

```
## [1] 59.309
```

```
#the average is 59.309
```

```
#7a.
#Type "rivers" in your R console. Create a vector data with 7
# elements, containing the number of elements (length) in rivers,
# their sum (sum), mean (mean), median (median), variance (var)
# standard deviation (sd), minimum (min) and maximum (max).
data <- c(length(rivers), sum(rivers), mean(rivers), median(rivers), var(rivers),
          sd(rivers), min(rivers), max(rivers))
data
```

```
## [1] 141.0000 83357.0000 591.1844 425.0000 243908.4086 493.8708
## [7] 135.0000 3710.0000
```

```
#7b. What are the results?
# the result are 141, 83357, 591.1844, 425, 243908, 493, 135, and 3710 correspondingly
```

```
#7c. Write the code and its outputs.
#data <- c(length(rivers), sum(rivers), mean(rivers), median(rivers), var(rivers),
#          + sd(rivers), min(rivers), max(rivers))
#> data <- c(length(rivers), sum(rivers), mean(rivers), median(rivers), var(rivers),
#          + sd(rivers), min(rivers), max(rivers))
#> data
#[1] 141.0000 83357.0000 591.1844 425.0000 243908.4086 493.8708 135.0000 3710.0000
```

```
#8a.
#Create vectors according to the above table. Write the codes.
Power_Ranking <- c(1:25)
```

```
Celebrity_Name <- c(
  "Tom Cruise", "Rolling Stones", "Oprah Winfrey", "U2", "Tiger Woods",
  "Steven Spielberg", "Howard Stern", "50 Cent", "Cast of the Sopranos",
```

```

"Dan Brown", "Bruce Springsteen", "Donald Trump", "Muhammad Ali",
"Paul McCartney", "George Lucas", "Elton John", "David Letterman",
"Phil Mickelson", "J.K. Rowling", "Bradd Pitt", "Peter Jackson",
"Dr. Phil McGraw", "Jay Lenon", "Celine Dion", "Kobe Bryant"
)

```

```

Pay_Millions <- c(
  67, 90, 225, 110, 90, 332, 302, 41, 52, 88, 55, 44, 55,
  40, 233, 34, 40, 47, 75, 25, 39, 45, 32, 40, 31
)

```

#8b. Modify the power ranking and pay of J.K. Rowling. Change power ranking to 15 and pay to 90. Write

```

Power_Ranking[19] <- 15
Pay_Millions[19] <- 90

```

#9c. Interpret the data.

```

Modified_Data <- data.frame(
  Ranking = Power_Ranking,
  Celebrity = Celebrity_Name,
  Pay = Pay_Millions)

```

Modified_Data

##	Ranking	Celebrity	Pay
## 1	1	Tom Cruise	67
## 2	2	Rolling Stones	90
## 3	3	Oprah Winfrey	225
## 4	4	U2	110
## 5	5	Tiger Woods	90
## 6	6	Steven Spielberg	332
## 7	7	Howard Stern	302
## 8	8	50 Cent	41
## 9	9	Cast of the Sopranos	52
## 10	10	Dan Brown	88
## 11	11	Bruce Springsteen	55
## 12	12	Donald Trump	44
## 13	13	Muhammad Ali	55
## 14	14	Paul McCartney	40
## 15	15	George Lucas	233
## 16	16	Elton John	34
## 17	17	David Letterman	40
## 18	18	Phil Mickelson	47
## 19	15	J.K. Rowling	90
## 20	20	Bradd Pitt	25
## 21	21	Peter Jackson	39
## 22	22	Dr. Phil McGraw	45
## 23	23	Jay Lenon	32
## 24	24	Celine Dion	40
## 25	25	Kobe Bryant	31