

Appendix S2 - Covariates of rodent community change


Appendix to Diaz, R. M. and Ernest, S. K. M., “Maintenance of community function through compensation breaks down over time in a desert rodent community” for review at *Ecology*.

Appendix S2 Figure S1. Changes in overall community energy use (A), NDVI (B), and local climate (C) surrounding the 2010 shift in rodent community composition. As documented in Christensen et al. (2018), the 2010 transition followed a period of low abundance community-wide (A) and low plant productivity (B). Since 2010, the site has experienced two periods of drought (C) interspersed with an unusually wet period.


Total rodent energy use (A) is calculated as the total energy use of all granivores on control plots ($Etot_C$) in each census period. The anomaly (shown) is calculated as the difference between the total energy use in each census period and the long-term mean of total energy use. Vertical dashed lines mark the dates of major transitions in the rodent community. NDVI anomaly (B) is calculated as the difference between monthly NDVI and the long-term mean for that month. NDVI data were obtained from Landsat 5, 7, and 8 using the `ndvi` function in the R package `portalr` (Maesk et al. 2006; Vermote et al. 2016; Christensen et al. 2019). Drought (C) was calculated using a 12-month Standardized Precipitation Evapotranspiration index (SPEI) for all months from 1989-2020, using the Thornthwaite method to estimate potential evapotranspiration (using the R package `SPEI`, Beguería and Vicente-Serrano 2017; Slette et al. 2019; Cárdenas et al. 2021). Values greater than 0 (blue) indicate wetter than average conditions, and values less than 0 (red) indicate drier conditions. Values between -1 and 1 (horizontal lines) are considered within normal variability for a system, while values < -1 constitute drought (Slette et al. 2019).

23 Appendix S2 Figure S1.


A



B



C



24

25

26 **References**

- 27 Beguería, S., and S. M. Vicente-Serrano. 2017. SPEI: Calculation of the Standardised
28 Precipitation-Evapotranspiration Index.
- 29 Cárdenas, P. A., E. Christensen, S. K. M. Ernest, D. C. Lightfoot, R. L. Schooley, P. Stapp, and
30 J. A. Rudgers. 2021. Declines in rodent abundance and diversity track regional climate
31 variability in North American drylands. *Global Change Biology*:gcb.15672.
- 32 Christensen, E. M., D. J. Harris, and S. K. M. Ernest. 2018. Long-term community change
33 through multiple rapid transitions in a desert rodent community. *Ecology* 99:1523–1529.
- 34 Christensen, E. M., G. M. Yenni, H. Ye, J. L. Simonis, E. K. Bledsoe, R. M. Diaz, S. D. Taylor,
35 E. P. White, and S. K. M. Ernest. 2019. portalr: an R package for summarizing and using
36 the Portal Project Data. *Journal of Open Source Software* 4:1098.
- 37 Masek, J.G., Vermote, E.F., Saleous, N., Wolfe, R., Hall, F.G., Huemmrich, F., Gao, F., Kutler,
38 J., and Lim, T.K. (2006). A Landsat surface reflectance data set for North America, 1990-
39 100, *IEEE Geoscience and Remote Sensing Letters*. 3:68-72.
- 40 Slette, I. J., A. K. Post, M. Awad, T. Even, A. Punzalan, S. Williams, M. D. Smith, and A. K.
41 Knapp. 2019. How ecologists define drought, and why we should do better. *Global*
42 *Change Biology* 25:3193–3200.
- 43 Vermote, E., Justice, C., Claverie, M., & Franch, B. (2016). Preliminary analysis of the
44 performance of the Landsat 8/OLI land surface reflectance product. *Remote Sensing of*
45 *Environment*, 185, 46-56.