Biomass analysis

Figures

Compensation and total biomass

Lines are 6-month moving averages. Horizontal lines + ribbons are means and SE or CL from GLM or GLS.

Compensation

Compensation refers to compensatory gains in biomass by small granivores on exclosure plots relative to controls. Calculated as $\frac{SmgranExclosure-SmgranControl}{DipoControl}$. Total biomass refers to the overall loss in biomass caused by kangaroo rat removal.

```
## Joining, by = "oera"
```

Total biomass ratio

```
## Joining, by = "oera"
```

Rodent community composition

C. baileyi

```
## Warning in eval(family$initialize): non-integer #successes in a binomial glm!
## Warning in eval(family$initialize): non-integer #successes in a binomial glm!
## Warning in eval(family$initialize): non-integer #successes in a binomial glm!
## Joining, by = c("period", "oplottype")
## Joining, by = c("period", "oplottype", "censusdate")
```


Dipodomys

```
## Warning in eval(family$initialize): non-integer #successes in a binomial glm!
## Joining, by = c("period", "oplottype")
## Joining, by = "period"
```

Full figure


```
## Setting row to 1
## Setting column to 1
## Setting row to 2
## Setting column to 1
## Setting row to 3
## Setting column to 1
## Setting column to 1
## Setting row to 4
## Setting column to 1
## Warning: Removed 228 row(s) containing missing values (geom_path).
```

Total biomass




В

Biomass compensation




С

Kangaroo rat (Dipodomys) biomass



D

C. baileyi biomass



Model results

Compensation & total biomass

Compensation

Table S1. Coefficients from GLS for compensation

```
## Value Std.Error t-value p-value
## (Intercept) 0.30814431 0.02905391 10.605950 6.936638e-23
## oera.L 0.07114123 0.05141306 1.383719 1.673549e-01
## oera.Q -0.27991205 0.04652521 -6.016352 4.624854e-09
```

Table S2. Estimates from GLS for compensation

```
## oera emmean SE df lower.CL upper.CL
## 1 a_pre_pb 0.1435663 0.05114193 39.16210 0.04013563 0.2469969
## 2 b_pre_reorg 0.5366915 0.04527450 41.78758 0.44531015 0.6280729
## 3 c_post_reorg 0.2441751 0.05172050 41.05328 0.13972758 0.3486227
```

Table S3. Contrasts from GLS for compensation

```
## contrast estimate SE df t.ratio p.value

## 1 a_pre_pb - b_pre_reorg -0.3931253 0.06738108 43.09746 -5.834357 0.0000

## 2 a_pre_pb - c_post_reorg -0.1006089 0.07270904 40.24488 -1.383719 0.3588

## 3 b_pre_reorg - c_post_reorg 0.2925164 0.06780029 44.29593 4.314383 0.0003
```

Total biomass

Table S4. Coefficients from GLS on total biomass ratio

```
## Value Std.Error t-value p-value
## (Intercept) 0.4553971 0.02724184 16.716827 3.547602e-46
## oera.L 0.1454493 0.04779892 3.042941 2.525739e-03
## oera.Q -0.2531409 0.04273433 -5.923594 7.735453e-09
```

Table S5. Estimates from GLS on total biomass ratio

```
## oera emmean SE df lower.CL upper.CL
## 1 a_pre_pb 0.2492046 0.04765836 33.75206 0.1523249 0.3460842
## 2 b_pre_reorg 0.6620857 0.04195145 35.90882 0.5769967 0.7471747
## 3 c_post_reorg 0.4549009 0.04802155 34.91256 0.3574032 0.5523986
```

Table S6. Contrasts from GLS on total biomass ratio

```
## contrast estimate SE df t.ratio p.value

## 1 a_pre_pb - b_pre_reorg -0.4128811 0.06217389 38.34660 -6.640747 0.0000

## 2 a_pre_pb - c_post_reorg -0.2056963 0.06759788 34.60306 -3.042941 0.0121

## 3 b_pre_reorg - c_post_reorg 0.2071848 0.06243247 39.12162 3.318542 0.0055
```

Community composition

Kangaroo rat proportional biomass

Table S7. Coefficients from GLM on Dipodomys biomass.

```
## Estimate Std. Error z value Pr(>|z|)
## (Intercept) 1.6149566 0.1644937 9.817741 9.443577e-23
## oera.L -1.1672395 0.3180813 -3.669626 2.429058e-04
## oera.Q 0.6619048 0.2473324 2.676175 7.446776e-03
```

Table S8. Estimates from GLM on Dipodomys biomass.

```
## oera prob SE df asymp.LCL asymp.UCL
## 1 a_pre_pb 0.9376458 0.02264595 Inf 0.8932605 0.9820310
## 2 b_pre_reorg 0.7454543 0.03850245 Inf 0.6699909 0.8209177
## 3 c_post_reorg 0.7426552 0.04371710 Inf 0.6569713 0.8283392
```

Table S9. Contrasts from GLM on Dipodomys biomass.

```
## contrast estimate SE df z.ratio p.value

## 1 a_pre_pb - b_pre_reorg 0.192191469 0.04466854 Inf 4.30261406 0.0001

## 2 a_pre_pb - c_post_reorg 0.194990571 0.04923437 Inf 3.96045595 0.0002

## 3 b_pre_reorg - c_post_reorg 0.002799103 0.05825481 Inf 0.04804929 0.9987
```

C. baileyi proportional biomass

```
## Warning in eval(family$initialize): non-integer #successes in a binomial glm!
## Warning in eval(family$initialize): non-integer #successes in a binomial glm!
## Warning in eval(family$initialize): non-integer #successes in a binomial glm!
## [1] 237.6847
## [1] 231.2374
## [1] 466.4937
## Analysis of Deviance Table
## Model 1: pb_prop ~ oera * oplottype
## Model 2: pb_prop ~ oera + oplottype
    Resid. Df Resid. Dev Df Deviance Pr(>Chi)
## 1
           451
                   57.835
           452
                   59.768 -1
                               -1.933
                                      0.1644
## Analysis of Deviance Table
## Model 1: pb_prop ~ oera + oplottype
```

```
## Model 2: pb_prop ~ oera
## Resid. Df Resid. Dev Df Deviance Pr(>Chi)
## 1     452     59.768
## 2     453     210.395 -1 -150.63 < 2.2e-16 ***
## ---
## Signif. codes: 0 '***' 0.001 '**' 0.05 '.' 0.1 ' ' 1</pre>
```

Note that a model fit as pb_proportional_biomass ~ era + treatment + era:treatment does not outperform a model fit without the interaction term, or pb_proportional_biomass ~ era + treatment (AIC for the no interaction model = 231 compared to 237 for the interaction model; p-value for an anova Chi-squared comparison of the two models = 0.16). We therefore use the model without the interaction term.

Table S10. Coefficients from GLM on C. baileyi biomass

```
## Estimate Std. Error z value Pr(>|z|)
## (Intercept) -1.538798 0.1671239 -9.207525 3.337295e-20
## oera.L -1.403286 0.2006948 -6.992140 2.707241e-12
## oplottype.L 2.270657 0.2298594 9.878462 5.161898e-23
```

Table S11. Estimates from GLM on C. baileyi biomass

```
## oera prob SE df asymp.LCL asymp.UCL
## 1 b_pre_reorg 0.36667107 0.04061520 Inf 0.28706674 0.4462754
## 2 c_post_reorg 0.07370848 0.01748044 Inf 0.03944746 0.1079695
```

Table S12. Contrasts from GLM on C. baileyi biomass.

```
## contrast estimate SE df z.ratio p.value
## 1 b_pre_reorg - c_post_reorg 0.2929626 0.04132914 Inf 7.088523 0
```