

Dykhne Adiabatic Approximation for Pair-creation

Making some assumptions

restart : *with* (*Physics*) : *with* (*Physics* [*Vectors*]) : *with* (*LinearAlgebra*) : *Setup* (*mathematicalnotation* = *true*) :

Setup (*realobjects* = { *e*, *m*, *c*, ϵ_0 }) :

Setup (*realobjects* = { *t*, $V_x(t)$, $V_y(t)$, $\eta(t)$, $\Omega(t)$ }) ;

$$[realobjects = \{\hat{i}, \hat{j}, \hat{k}, \hat{\phi}, \hat{r}, \hat{\rho}, \hat{\theta}, c, e, m, \phi, r, \rho, t, \theta, x, y, z, \epsilon_0, \Omega(t), V_x(t), V_y(t), \eta(t)\}] \quad (1.1)$$

alias ($V_x = V_x(t)$, $V_y = V_y(t)$, $\eta = \eta(t)$, $\Omega = \Omega(t)$)

$$V_x, V_y, \eta, \Omega \quad (1.2)$$

cRDI mathematical formalism

Cliford conjugation

$$ClifordConj := (X) \rightarrow \langle \langle X[2, 2] \mid -X[1, 2] \rangle, \langle -X[2, 1] \mid X[1, 1] \rangle \rangle :$$

Define a contravariant vector *v* from a paravector *P*

$$DefContrVect := (v, P) \rightarrow Define \left(v_{\sim mu} = Simplify \left(Array \left(1..4, (l) \rightarrow \frac{1}{2} \cdot Trace(P \cdot Psigma[l]) \right) \right) \right) :$$

Extract the 3D vector from a paravector *P*

$$ExtractVector := (P) \rightarrow \frac{1}{2} \cdot Simplify(Vector(3, (l) \rightarrow Trace(Psigma[l].P)). Vector([_i, _j, _k])) :$$

Get the Faraday matrix from the eigenspinor

$$GetFaraday := (\Lambda, W) \rightarrow w \cdot \frac{2 \cdot m \cdot c}{e} \cdot (diff(\Lambda, t).ClifordConj(\Lambda) + \Lambda.diff(W, t).ClifordConj(W) \cdot ClifordConj(\Lambda)) :$$

Extract the electric field vector from the Faraday matrix

$$GetElectricField := (F) \rightarrow \frac{1}{2} \cdot Simplify(Vector(3, (l) \rightarrow Re(Trace(Psigma[l].F))). Vector([_i, _j,$$

$_k]))):$

Extract the magnetic field vector from the Faraday matrix

$$GetMagneticField := (F) \rightarrow \frac{1}{2 \cdot c} \cdot Simplify(Vector(3, (l) \rightarrow \text{Im}(Trace(Psigma[l].F))). Vector([_i, _j, _k]))):$$

Extracting the scalar and vector parts

$$ScalarPart := (L) \rightarrow \frac{(L + CliffordConj(L))}{2} :$$

$$VectorPart := (L) \rightarrow \frac{(L - CliffordConj(L))}{2} :$$

Extract the Poynting vector

$$PoyntingCliford := (F) \rightarrow \frac{c \cdot \epsilon_0}{2} \cdot VectorPart(F.Dagger(F)) :$$

$$PoyntingVector := (F) \rightarrow ExtractVector(PoyntingCliford(F)) :$$

The radiated energy per unit of time is given by Larmor's formula

$$PowerEmitted := (U) \rightarrow \frac{-e^2}{6 \cdot c \cdot \text{Pi} \cdot \epsilon_0} \cdot \frac{w^2}{2} \cdot Trace(diff(U, t).CliffordConj(diff(U, t))) :$$

Particle in electromagnetic plane wave combined with homogeneous axial electric fields

The eigenspinor employed

$$\Lambda := MatrixExponential\left(\frac{\eta(t) \cdot Matrix(Psigma[3])}{2}\right) \cdot \begin{bmatrix} 1 & \frac{V_x(t)}{c} - \frac{I \cdot V_y(t)}{c} \\ 0 & 1 \end{bmatrix}$$

$$\Lambda := \begin{bmatrix} e^{\frac{\eta(t)}{2}} & e^{\frac{\eta(t)}{2}} \left(\frac{V_x(t)}{c} - \frac{I V_y(t)}{c} \right) \\ 0 & e^{-\frac{\eta(t)}{2}} \end{bmatrix} \quad (3.1)$$

Proper velocity

$$U := simplify(\Lambda.Dagger(\Lambda))$$

$$U := \begin{bmatrix} \frac{e^{\eta(t)} (V_y(t)^2 + V_x(t)^2 + c^2)}{c^2} & \frac{-I V_y(t) + V_x(t)}{c} \\ \frac{I V_y(t) + V_x(t)}{c} & e^{-\eta(t)} \end{bmatrix} \quad (3.2)$$

DefContrVect(u, U)

Defined objects with tensor properties

$$\left\{ \gamma_{\mu}, \sigma_{\mu}, \partial_{\mu}, g_{\mu, \nu}, u^{\mu}, \delta_{\mu, \nu}, \epsilon_{\alpha, \beta, \mu, \nu} \right\} \quad (3.3)$$

Print proper velocity components

'u[~0]'= *expand*(u[~0])

$$u^0 = \frac{1}{2 e^{\eta}} + \frac{e^{\eta} V_x^2}{2 c^2} + \frac{e^{\eta} V_y^2}{2 c^2} + \frac{e^{\eta}}{2} \quad (3.4)$$

'u[~1]'= *expand*(u[~1])

$$u^1 = \frac{V_x}{c} \quad (3.5)$$

'u[~2]'= *expand*(u[~2])

$$u^2 = \frac{V_y}{c} \quad (3.6)$$

'u[~3]'= *expand*(u[~3])

$$u^3 = \frac{e^{\eta} V_y^2}{2 c^2} + \frac{e^{\eta} V_x^2}{2 c^2} + \frac{e^{\eta}}{2} - \frac{1}{2 e^{\eta}} \quad (3.7)$$

Enforce the laboratory time parametrization (w must be fixed as shown below for t to become the laboratory time)

$w := \text{expand}\left(\frac{1}{2} \cdot \text{Trace}(U)\right)$

$$w := \frac{1}{2 e^{\eta}} + \frac{e^{\eta} V_x^2}{2 c^2} + \frac{e^{\eta} V_y^2}{2 c^2} + \frac{e^{\eta}}{2} \quad (3.8)$$

expand(u[~0] - w);

$$0 \quad (3.9)$$

▼ Circular trajectories in plane waves

$$\begin{aligned} \text{CircTrajParams} &:= \left\{ V_x = \text{diff}(r \cdot \cos(\omega \cdot t), t), V_y = \text{diff}(r \cdot \sin(\omega \cdot t), t), \eta = -\ln \left(\text{sqrt} \left(1 \right. \right. \right. \\ &\quad \left. \left. \left. + \left(\frac{r \cdot \omega}{c} \right)^2 \right) \right) \right\}; \\ \text{CircTrajParams} &:= \left\{ V_x = -r \omega \sin(\omega t), V_y = r \omega \cos(\omega t), \eta = -\frac{\ln \left(1 + \frac{r^2 \omega^2}{c^2} \right)}{2} \right\} \end{aligned} \quad (4.1)$$

Find trajectory

dsolve(*subs*(*CircTrajParams*, $w \cdot \text{diff}(x(t), t) = u[~1]$)) assuming $c > 0$;

$$x(t) = \frac{r \cos(\omega t)}{\sqrt{r^2 \omega^2 + c^2}} + _CI \quad (4.2)$$

$dsolve(subs(CircTrajParams, w \cdot diff(y(t), t) = u[\sim 2]))$ assuming $c > 0$ **and** $\omega > 0$;

$$y(t) = \frac{r \sin(\omega t)}{\sqrt{r^2 \omega^2 + c^2}} + _CI \quad (4.3)$$

$dsolve(subs(CircTrajParams, w \cdot diff(z(t), t) = u[\sim 3]))$;

$$z(t) = _CI \quad (4.4)$$

Notice that the equation for critical time does not depend on t ; thus, there will be no pair creation.

$simplify(subs(CircTrajParams, u[\sim 0]))$;

$$\frac{\sqrt{r^2 \omega^2 + c^2}}{|c|} \quad (4.5)$$

Pair creation in the homogeneous electric field

$EParams := \{V_x = 0, V_y = 0\}$;

$$EParams := \{V_x = 0, V_y = 0\} \quad (5.1)$$

Calculate the fields

$F := simplify(subs(EParams, (GetFaradayday(subs(EParams, \Lambda), IdentityMatrix(2)))))$
assuming $c > 0$ **and** $m > 0$;

$$F := \begin{bmatrix} \frac{(e^{-\eta(t)} + e^{\eta(t)}) m c \dot{\eta}(t)}{2 e} & 0 \\ 0 & -\frac{(e^{-\eta(t)} + e^{\eta(t)}) m c \dot{\eta}(t)}{2 e} \end{bmatrix} \quad (5.2)$$

$GetMagneticField(F)$;

$$0 \quad (5.3)$$

$GetElectricField(F)$

$$\frac{(e^{-\eta} + e^{\eta}) m c \left(\frac{\partial}{\partial t} \eta \right) \widehat{k}}{2 e} \quad (5.4)$$

Find η to make electric homogeneous time independent

$$odetest\left(\eta = \operatorname{arcsinh}\left(\frac{E_0 e t}{m \cdot c}\right), \operatorname{coeff}(GetElectricField(F), _k) = E_0\right); \quad (5.5)$$

Update params

$$EParams := EParams \text{ union } \left\{ \eta = \operatorname{arcsinh}\left(\frac{E_0 e t}{m \cdot c}\right) \right\};$$

$$EParams := \left\{ V_x = 0, V_y = 0, \eta = \operatorname{arcsinh} \left(\frac{E_0 e t}{m c} \right) \right\} \quad (5.6)$$

Find crytical time

$t0 := \operatorname{solve}(\operatorname{subs}(EParams, w), t);$

$$t0 := \frac{I m c}{E_0 e} \quad (5.7)$$

Calculate the pair creation rate notice $2 \cdot m c^2$ in front of w

$$\exp \left(- \frac{2 \cdot \operatorname{Im} \left(\operatorname{int} \left(\operatorname{subs}(EParams, 2 \cdot m \cdot c^2 \cdot w), t = 0 .. t0 \right) \right)}{\hbar} \right);$$

$$e^{-\frac{\pi m^2 c^3 \Re \left(\frac{1}{E_0} \right)}{e \hbar}} \quad (5.8)$$

Pair creation in two-color laser field ($\omega + 3\omega$)

$$LaserParams := \left\{ V_x = \frac{c}{\operatorname{sqrt}(g)} \cdot \cos(\omega \cdot t), V_y = 0, \eta = 0 \right\}$$

$$LaserParams := \left\{ V_x = \frac{c \cos(\omega t)}{\sqrt{g}}, V_y = 0, \eta = 0 \right\} \quad (6.1)$$

Calculate the fields

$F := \operatorname{simplify}(\operatorname{subs}(LaserParams, (\operatorname{GetFaradayday}(\operatorname{subs}(LaserParams, \Lambda), \operatorname{IdentityMatrix}(2)))))$
assuming $c > 0$ and $m > 0$;

$$F := \begin{bmatrix} 0 & -\frac{(c^2 \cos(\omega t)^2 + 2 c^2 g) m \omega \sin(\omega t)}{e c g^3 |^2} \\ 0 & 0 \end{bmatrix} \quad (6.2)$$

$B_- := \operatorname{simplify}(\operatorname{GetMagneticField}(F))$ assuming $\omega > 0$ and $g > 1$;

$$\vec{B} := -\frac{\sin(\omega t) \omega \hat{j} m (\cos(2 \omega t) + 1 + 4 g)}{4 e g^3 |^2} \quad (6.3)$$

$E_- := \operatorname{simplify}(\operatorname{GetElectricField}(F))$ assuming $\omega > 0$ and $g > 1$;

$$\vec{E} := -\frac{\sin(\omega t) \omega \hat{i} c m (\cos(2 \omega t) + 1 + 4 g)}{4 e g^3 |^2} \quad (6.4)$$

$E_- B_-;$

$$0 \quad (6.5)$$

Find crytical time

$\operatorname{solve}(\operatorname{subs}(LaserParams, w), t)$ assuming $g > 0$;

$$\frac{\frac{\pi}{2} - I \operatorname{arcsinh}(\sqrt{2} \sqrt{g})}{\omega}, \frac{\frac{\pi}{2} + I \operatorname{arcsinh}(\sqrt{2} \sqrt{g})}{\omega} \quad (6.6)$$

$t0 := \text{convert}(\text{combine}(\text{solve}(\text{subs}(LaserParams, w), t)[2]), \operatorname{arcsinh})$ assuming $g > 0$;

$$t0 := \frac{\frac{\pi}{2} - I \operatorname{arcsinh}(I \sqrt{-2g})}{\omega} \quad (6.7)$$


Calculate the pair creation rate notice $2 \cdot mc^2$ in front of w

$\text{transition} := \exp\left(-\frac{2 \cdot \operatorname{Im}(\text{int}(\text{subs}(LaserParams, 2 \cdot m \cdot c^2 \cdot w), t = 0 .. t0))}{\hbar}\right)$ assuming $r > 0$

and $c > 0$ and $\omega > 0$ and $g > 1$;

$$\text{transition} := e^{-\frac{mc^2(-2\sqrt{2}\sqrt{1+2g}g + 8\operatorname{arcsinh}(\sqrt{2}\sqrt{g})g^{3/2} + 2\operatorname{arcsinh}(\sqrt{2}\sqrt{g})\sqrt{g})}{2\omega g^{3/2}\hbar}} \quad (6.8)$$

$\text{plot}(\text{subs}(m = 1, c = 1, \omega = 1, \hbar = 1, (\text{transition})), g = 0 .. 1,);$



Get the asymptotic expression for large g

$$\begin{aligned}
 & asympt\left(-\frac{m\,c^2\,(-2\sqrt{2}\sqrt{1+2g}\,g+8\operatorname{arcsinh}(\sqrt{2}\sqrt{g})\,g^{3/2}+2\operatorname{arcsinh}(\sqrt{2}\sqrt{g})\sqrt{g})}{2\,\omega\,g^{3/2}\,\hbar},\right. \\
 & \left.g,2\right); \\
 & -\frac{m\,c^2\,(-4+8\ln(2\sqrt{2})+4\ln(g))}{2\,\omega\,\hbar}-\frac{m\,c^2\,(2\ln(2\sqrt{2})+\ln(g))}{2\,\omega\,\hbar\,g}+O\left(\frac{1}{g^2}\right)
 \end{aligned}
 \tag{6.9}$$

$$\begin{aligned}
 & simplify\left(\exp\left(-\frac{m\,c^2\,(4\ln(g))}{2\,\omega\,\hbar}\right)\right); \\
 & g-\frac{2\,m\,c^2}{\omega\,\hbar}
 \end{aligned}
 \tag{6.10}$$

$$\begin{aligned}
 & simplify\left(\exp\left(-\frac{m\,c^2\,(-4+8\ln(2\sqrt{2}))}{2\,\omega\,\hbar}\right)\right); \\
 & e^{-\frac{2\,m\,c^2\,(-1+3\ln(2))}{\omega\,\hbar}}
 \end{aligned}
 \tag{6.11}$$

$$\begin{aligned}
 & evalf(-1+3\ln(2)); \\
 & 1.079441542
 \end{aligned}
 \tag{6.12}$$