

priors on exchangeable directed graphs

Diana Cai

Department of Statistics
University of Chicago

Collaborators:

Nathanael Ackerman (Harvard)
Cameron Freer (Gamalon and MIT)

Overview


- We consider **exchangeable (dense) graphs**.
- Specifically, we are interested in **directed** graphs.
However, most work has focused on undirected graphs.
- **Aldous–Hoover** has an analogous statement for directed graphs, which is more complicated than merely using an *asymmetric* function.
- Many natural nonparametric priors on exchangeable undirected graphs **extend to the directed case**.
- This perspective leads to natural priors on other exchangeable structures, such as **tournaments** and **directed acyclic graphs**.

Exchangeable graphs


exchangeability:

- order of vertices doesn't affect distribution of graph

$$\Pr\left(\begin{array}{|c|c|c|} \hline \square & \blacksquare & \blacksquare \\ \hline \blacksquare & \square & \square \\ \hline \blacksquare & \square & \square \\ \hline \end{array}\right) = \Pr\left(\begin{array}{|c|c|c|} \hline \square & \square & \blacksquare \\ \hline \square & \square & \blacksquare \\ \hline \blacksquare & \blacksquare & \square \\ \hline \end{array}\right)$$



```
graph TD; 1((1)) --- 2((2)); 1 --- 3((3));
```



```
graph TD; 3((3)) --- 2((2)); 3 --- 1((1));
```

Exchangeable graphs

Undirected

Any exchangeable random infinite graph is obtained as a mixture of $\mathbb{G}(\infty, W)$.

$\mathbb{G}(\infty, W)$ is a sampling procedure from the **graphon** W .

$$W : [0, 1]^2 \rightarrow [0, 1]$$

$$W(x, y) = W(y, x)$$

Aldous–Hoover


Directed

Exchangeable graphs

Graphon:

$$W : [0, 1]^2 \rightarrow [0, 1]$$

$$W(x, y) = W(y, x)$$



$$W(x, y) = \frac{(1 - x) + (1 - y)}{2}$$

Exchangeable graphs

Graphon:


$$W : [0, 1]^2 \rightarrow [0, 1]$$

$$W(x, y) = W(y, x)$$

Sampling procedure:

$$U_i \sim \text{Uniform}[0, 1]$$

$$G_{ij} \sim \text{Bern}(W(U_i, U_j)), i < j \quad (\text{Set } G_{ji} = G_{ij})$$



Exchangeable graphs


Graphon:

$$W : [0, 1]^2 \rightarrow [0, 1]$$
$$W(x, y) = W(y, x)$$

Sampling procedure:

$$U_i \sim \text{Uniform}[0, 1]$$

$$G_{ij} \sim \text{Bern}(W(U_i, U_j)), i < j \quad (\text{Set } G_{ji} = G_{ij})$$



Exchangeable graphs


Graphon:

$$W : [0, 1]^2 \rightarrow [0, 1]$$
$$W(x, y) = W(y, x)$$

Sampling procedure:

$$U_i \sim \text{Uniform}[0, 1]$$

$$G_{ij} \sim \text{Bern}(W(U_i, U_j)), i < j$$



Exchangeable graphs


Graphon:

$$W : [0, 1]^2 \rightarrow [0, 1]$$
$$W(x, y) = W(y, x)$$

Sampling procedure:

$$U_i \sim \text{Uniform}[0, 1]$$

$$G_{ij} \sim \text{Bern}(W(U_i, U_j)), i < j$$



Exchangeable graphs


Graphon:

$$W : [0, 1]^2 \rightarrow [0, 1]$$
$$W(x, y) = W(y, x)$$

Sampling procedure:

$$U_i \sim \text{Uniform}[0, 1]$$

$$G_{ij} \sim \text{Bern}(W(U_i, U_j)), i < j$$




Exchangeable digraphs

Many results for exchangeable **undirected** graphs
extend to **directed** graphs.

Exchangeable digraphs

Many results for exchangeable **undirected** graphs extend to **directed** graphs.

- order of vertices doesn't affect distribution of graph

$$\Pr\left(\begin{array}{|c|c|c|} \hline \blacksquare & \square & \blacksquare \\ \hline \blacksquare & \square & \square \\ \hline \blacksquare & \blacksquare & \square \\ \hline \end{array}\right) = \Pr\left(\begin{array}{|c|c|c|} \hline \square & \blacksquare & \square \\ \hline \square & \blacksquare & \blacksquare \\ \hline \blacksquare & \blacksquare & \square \\ \hline \end{array}\right)$$


The diagram shows two directed graphs. The left graph has vertices labeled 1, 2, and 3. Vertex 1 has a self-loop and edges to vertices 2 and 3. Vertex 2 has edges to vertices 1 and 3. Vertex 3 has an edge to vertex 2. The right graph has vertices labeled 1, 2, and 3. Vertex 2 has a self-loop and edges to vertices 1 and 3. Vertex 1 has edges to vertices 2 and 3. Vertex 3 has an edge to vertex 1.

Exchangeable graphs

Undirected

Any exchangeable random infinite graph is obtained as a mixture of $\mathbb{G}(\infty, W)$.

$\mathbb{G}(\infty, W)$ is a sampling procedure from the **graphon** W .

$$W : [0, 1]^2 \rightarrow [0, 1]$$
$$W(x, y) = W(y, x)$$

Aldous–Hoover;
cf. Lovász–Szegedy

Directed

Any exchangeable random infinite **digraph** is obtained as a mixture of $\mathbb{G}(\infty, \mathbf{W})$.

$\mathbb{G}(\infty, \mathbf{W})$ is a sampling procedure from the **digraphon** \mathbf{W} .

$$\mathbf{W} = ?$$

implicit in A–H;
cf. Diaconis–Janson

Exchangeable digraphs

Can we just use a single **asymmetric** measurable function?

Exchangeable digraphs

Can we just use a single **asymmetric** measurable function?

Yes, by independently choosing each edge direction.

Exchangeable digraphs

Can we just use a single **asymmetric** measurable function?

Yes, by independently choosing each edge direction.

But this does not cover *all* directed graphs:


Exchangeable digraphs

Can we just use a single **asymmetric** measurable function?

Yes, by independently choosing each edge direction.

But this does not cover *all* directed graphs:

- **tournaments** (each pair of vertices has exactly one directed edge)




Exchangeable digraphs

Can we just use a single **asymmetric** measurable function?


Yes, by independently choosing each edge direction.

But this does not cover *all* directed graphs:

- **tournaments** (each pair of vertices has exactly one directed edge)
- **undirected** graphs (each pair has either both or no directed edges)




Digraph representation




- Digraphon: $\mathbf{W} = (W_{00}, W_{01}, W_{10}, W_{11}, w)$
 $W_{ab} : [0, 1]^2 \rightarrow [0, 1]$ and $w : [0, 1] \rightarrow \{0, 1\}$ measurable functions
- For $a, b \in \{0, 1\}$ and $x, y \in [0, 1]$,
 1. $W_{ab}(x, y) = W_{ba}(y, x)$, when $a = b$
 2. $W_{ab}(x, y) = W_{ba}(y, x)$, when $a \neq b$
 3. $\sum_{a,b} W_{a,b}(x, y) = 1$


Digraph representation




Digraph representation




Digraph representation




Digraph representation



Digraph representation



Digraph representation




- Digraphon: $\mathbf{W} = (W_{00}, W_{01}, W_{10}, W_{11}, w)$

$W_{ab} : [0, 1]^2 \rightarrow [0, 1]$ and $w : [0, 1] \rightarrow \{0, 1\}$ measurable functions

- For $a, b \in \{0, 1\}$ and $x, y \in [0, 1]$,
 1. $W_{ab}(x, y) = W_{ba}(y, x)$, when $a = b$
 2. $W_{ab}(x, y) = W_{ba}(y, x)$, when $a \neq b$
 3. $\sum_{a,b} W_{a,b}(x, y) = 1$

Digraph representation



- Digraphon: $\mathbf{W} = (W_{00}, W_{01}, W_{10}, W_{11}, w)$

$W_{ab} : [0, 1]^2 \rightarrow [0, 1]$ and $w : [0, 1] \rightarrow \{0, 1\}$ measurable functions


- For $a, b \in \{0, 1\}$ and $x, y \in [0, 1]$,

1. $W_{ab}(x, y) = W_{ba}(y, x)$, when $a = b$

2. $W_{ab}(x, y) = W_{ba}(y, x)$, when $a \neq b$

3. $\sum_{a,b} W_{a,b}(x, y) = 1$

Digraph representation



- Digraphon: $\mathbf{W} = (W_{00}, W_{01}, W_{10}, W_{11}, w)$

$W_{ab} : [0, 1]^2 \rightarrow [0, 1]$ and $w : [0, 1] \rightarrow \{0, 1\}$ measurable functions


- For $a, b \in \{0, 1\}$ and $x, y \in [0, 1]$,

1. $W_{ab}(x, y) = W_{ba}(y, x)$, when $a = b$

2. $W_{ab}(x, y) = W_{ba}(y, x)$, when $a \neq b$

3. $\sum_{a,b} W_{a,b}(x, y) = 1$

Digraph representation



- Digraphon: $\mathbf{W} = (W_{00}, W_{01}, W_{10}, W_{11}, w)$

$W_{ab} : [0, 1]^2 \rightarrow [0, 1]$ and $w : [0, 1] \rightarrow \{0, 1\}$ measurable functions


- For $a, b \in \{0, 1\}$ and $x, y \in [0, 1]$,

1. $W_{ab}(x, y) = W_{ba}(y, x)$, when $a = b$

2. $W_{ab}(x, y) = W_{ba}(y, x)$, when $a \neq b$

3. $\sum_{a,b} W_{a,b}(x, y) = 1$

Digraph representation



- Digraphon: $\mathbf{W} = (W_{00}, W_{01}, W_{10}, W_{11}, w)$

$W_{ab} : [0, 1]^2 \rightarrow [0, 1]$ and $w : [0, 1] \rightarrow \{0, 1\}$ measurable functions


- For $a, b \in \{0, 1\}$ and $x, y \in [0, 1]$,

1. $W_{ab}(x, y) = W_{ba}(y, x)$, when $a = b$

2. $W_{ab}(x, y) = W_{ba}(y, x)$, when $a \neq b$

3. $\sum_{a,b} W_{a,b}(x, y) = 1$

Digraph representation



- Digraphon: $\mathbf{W} = (W_{00}, W_{01}, W_{10}, W_{11}, w)$

$W_{ab} : [0, 1]^2 \rightarrow [0, 1]$ and $w : [0, 1] \rightarrow \{0, 1\}$ measurable functions


- For $a, b \in \{0, 1\}$ and $x, y \in [0, 1]$,

1. $W_{ab}(x, y) = W_{ba}(y, x)$, when $a = b$

2. $W_{ab}(x, y) = W_{ba}(y, x)$, when $a \neq b$

3. $\sum_{a,b} W_{a,b}(x, y) = 1$

Digraph representation



- Digraphon: $\mathbf{W} = (W_{00}, W_{01}, W_{10}, W_{11}, w)$

$W_{ab} : [0, 1]^2 \rightarrow [0, 1]$ and $w : [0, 1] \rightarrow \{0, 1\}$ measurable functions


- For $a, b \in \{0, 1\}$ and $x, y \in [0, 1]$,

1. $W_{ab}(x, y) = W_{ba}(y, x)$, when $a = b$

2. $W_{ab}(x, y) = W_{ba}(y, x)$, when $a \neq b$

3. $\sum_{a,b} W_{a,b}(x, y) = 1$

Digraph representation




- Digraphon: $\mathbf{W} = (W_{00}, W_{01}, W_{10}, W_{11}, w)$

$W_{ab} : [0, 1]^2 \rightarrow [0, 1]$ and $w : [0, 1] \rightarrow \{0, 1\}$ measurable functions

- For $a, b \in \{0, 1\}$ and $x, y \in [0, 1]$,
 1. $W_{ab}(x, y) = W_{ba}(y, x)$, when $a = b$
 2. $W_{ab}(x, y) = W_{ba}(y, x)$, when $a \neq b$
 3. $\sum_{a,b} W_{a,b}(x, y) = 1$

Digraph representation




- Digraphon: $\mathbf{W} = (W_{00}, W_{01}, W_{10}, W_{11}, w)$

$W_{ab} : [0, 1]^2 \rightarrow [0, 1]$ and $w : [0, 1] \rightarrow \{0, 1\}$ measurable functions

- For $a, b \in \{0, 1\}$ and $x, y \in [0, 1]$,
 1. $W_{ab}(x, y) = W_{ba}(y, x)$, when $a = b$
 2. $W_{ab}(x, y) = W_{ba}(y, x)$, when $a \neq b$
 3. $\sum_{a,b} W_{a,b}(x, y) = 1$

Sampling procedure



- Define $\mathbf{W}_4(x, y) := (W_{00}(x, y), W_{01}(x, y), W_{10}(x, y), W_{11}(x, y))$


$$U_i \sim \text{Uniform}[0, 1]$$

$$E_{ij} \sim \text{Categorical}(\mathbf{W}_4(U_i, U_j)), \text{ for } i < j$$

$$G_{ij}, G_{ji} = \text{directed edge(s) corresponding to } E_{ij}, \text{ for } i < j$$

$$G_{ii} = w(U_i)$$

Sampling procedure



- Define $\mathbf{W}_4(x, y) := (W_{00}(x, y), W_{01}(x, y), W_{10}(x, y), W_{11}(x, y))$


$$U_i \sim \text{Uniform}[0, 1]$$

$$E_{ij} \sim \text{Categorical}(\mathbf{W}_4(U_i, U_j)), \text{ for } i < j$$

$$G_{ij}, G_{ji} = \text{directed edge(s) corresponding to } E_{ij}, \text{ for } i < j$$

$$G_{ii} = w(U_i)$$

Sampling procedure



- Define $\mathbf{W}_4(x, y) := (W_{00}(x, y), W_{01}(x, y), W_{10}(x, y), W_{11}(x, y))$


$$U_i \sim \text{Uniform}[0, 1]$$

$$E_{ij} \sim \text{Categorical}(\mathbf{W}_4(U_i, U_j)), \text{ for } i < j$$

$$G_{ij}, G_{ji} = \text{directed edge(s) corresponding to } E_{ij}, \text{ for } i < j$$

$$G_{ii} = w(U_i)$$

Sampling procedure



- Define $\mathbf{W}_4(x, y) := (W_{00}(x, y), W_{01}(x, y), W_{10}(x, y), W_{11}(x, y))$


$$U_i \sim \text{Uniform}[0, 1]$$

$$E_{ij} \sim \text{Categorical}(\mathbf{W}_4(U_i, U_j)), \text{ for } i < j$$

$$G_{ij}, G_{ji} = \text{directed edge(s) corresponding to } E_{ij}, \text{ for } i < j$$

$$G_{ii} = w(U_i)$$

Sampling procedure



- Define $\mathbf{W}_4(x, y) := (W_{00}(x, y), W_{01}(x, y), W_{10}(x, y), W_{11}(x, y))$

$$U_i \sim \text{Uniform}[0, 1]$$

$$E_{ij} \sim \text{Categorical}(\mathbf{W}_4(U_i, U_j)), \text{ for } i < j$$


$$G_{ij}, G_{ji} = \text{directed edge(s) corresponding to } E_{ij}, \text{ for } i < j$$

$$G_{ii} = w(U_i)$$


Special cases

- Undirected graphs (graphon) (e.g., $\text{ER}(\frac{1}{2})$)


$$W : [0, 1]^2 \rightarrow [0, 1]$$
$$W(x, y) = W(y, x)$$



W_{00}



W_{01}





W_{10}



W_{11}






w




Special cases

- Tournaments (e.g., generic tournament)




Special cases


- Linear ordering (the only one, by Glasner–Weiss)




W_{00}



W_{01}






W_{10}



W_{11}




w





Special cases

- Directed acyclic graphs (DAGs)




Special cases

- Partial ordering (poset)



Priors on digraphons

- Can extend literature for graphon priors (cf. Orbanz–Roy) to directed graphs, e.g., infinite relational model




[Orbanz–Roy, 2015]

- Some of these models are already intended for directed graphs, via an **asymmetric** measurable function (to describe independent edge directions).


Directed block models

- In a block model, pairs of regions can vary in how tournament-like, as well as how dense they are:
- e.g., Directed stochastic block model [Wang–Wong, 1987]


Example of SBM digraphon, 0.7 division



Infinite digraph block model




Infinite digraph block model




1. Draw partition \sim DP-Stick(alpha)

Infinite digraph block model




1. Draw partition \sim DP-Stick(α)
2. Draw weights \sim Dirichlet(\mathbf{a})

Infinite digraph block model




1. Draw partition $\sim \text{DP-Stick}(\alpha)$
2. Draw weights $\sim \text{Dirichlet}(\mathbf{a})$

Infinite digraph block model




Tournament


Dirichlet parameter $\alpha = [0, 2, 1, 0]$



sample




Infinite digraph block model




Almost totally ordered


Dirichlet parameter $\alpha = [0, 0, 1, 0]$




sample



reordered



Infinite digraph block model



Dirichlet parameter $\alpha = [0.9, 2.0, 1.0, 0.5]$


Infinite digraph block model

- Inference via collapsed Gibbs sampling of cluster assignments

$$p(z_i | z_{-i}, G) \propto p(z_i | z_{-i}) p(G | z_i, z_{-i})$$

cluster assignment of vertex i cluster assignments of all vertices except i


- Experiments using synthetic data



Dirichlet parameter $a = [1, 1, 1, 1]$

Infinite digraph block model

- Random digraphon




Dirichlet parameter $a = [1, 1, 1, 1]$

- Random sample, 100 vertices




original ordering



resorted by increasing
uniform random variables


Infinite digraph block model

- Random digraphon




Dirichlet parameter $a = [1, 1, 1, 1]$

- Collapsed Gibbs sampling



Infinite digraph block model

- Random digraphon




Dirichlet parameter $\alpha = [1, 1, 1, 1]$


- Collapsed Gibbs sampling



original ordering




resorted by increasing
uniform random variables




resorted by inferred cluster

Infinite digraph block model


- Random digraphon: ER + tournament



Infinite digraph block model




- Schematic of the sampled digraph:




Infinite digraph block model


- Random digraphon: ER + tournament




- Collapsed Gibbs sampling for **this model**



original ordering




resorted by increasing
uniform random variables




resorted by inferred cluster

Infinite digraph block model


- Random digraphon: ER + tournament




- Collapsed Gibbs sampling for **this model**



original ordering




resorted by increasing
uniform random variables




resorted by inferred cluster

Infinite digraph block model


- Random digraphon: ER + tournament




- Collapsed Gibbs sampling for the **infinite relational model**



original ordering




resorted by increasing
uniform random variables



resorted by inferred cluster

Conclusions

- Summary
- Discussion:



- other types of block models:
 - could consider other partitions, e.g., **Pitman–Yor**
- other priors, e.g., **Gaussian process** (as in Lloyd et al.)
- Aldous–Hoover already describes exchangeable **hypergraphs**
- **sparsity**

References

- D. J. Aldous. Representations for partially exchangeable arrays of random variables. *J. Multivariate Anal.*, 11(4): 581–598, 1981.
- P. Diaconis and S. Janson. Graph limits and exchangeable random graphs. *Rend. Mat. Appl.* (7), 28(1):33–61, 2008.
- E. Glasner and B. Weiss, Minimal actions of the group $S(\mathbb{Z})$ of permutations of the integers, *Geom. Funct. Anal.* 12(5):964–988, 2002.
- D. N. Hoover. Relations on probability spaces and arrays of random variables. Preprint, Institute for Advanced Study, Princeton, NJ, 1979.
- C. Kemp, J. B. Tenenbaum, T. L. Griffiths, T. Yamada, and N. Ueda. Learning systems of concepts with an infinite relational model. In *Proc. 21st Nat. Conf. Artificial Intelligence (AAAI-06)*, 2006.
- J. R. Lloyd, P. Orbanz, Z. Ghahramani, and D. M. Roy. Random function priors for exchangeable arrays with applications to graphs and relational data. In *Adv. Neural Inform. Process. Syst. (NIPS)* 25:1007–1015, 2012.
- L. Lovász. *Large networks and graph limits*, vol. 60 of American Math. Soc. Colloq. Publ. American Math. Soc., Providence, RI, 2012.
- P. Orbanz and D. M. Roy. Bayesian models of graphs, arrays and other exchangeable random structures. *IEEE Trans. Pattern Anal. Mach. Intell.*, 37(2):437–461, 2015.
- Y. J. Wang, and G. Y. Wong. Stochastic blockmodels for directed graphs. *J. American Statist. Assoc.*, 82(397), 8–19, 1987.