Supplementary Material for: Temporal decoupling of CTLA-4 and ERK sensitivity in brain tumour signaling: a theoretical basis for sequencing checkpoint and MAPK inhibition

Table S2: Initial Concentrations of Entities

Table 1: cBioPortal-derived initial concentration values for model entities.

S.No	Entities	mRNA Expression Value (TPM)	Molarity (M)
1	CTLA-4	0.998	3×10^{-9}
2	CD80	2.190	6.64×10^{-9}
3	PP2A	59.26	1.8×10^{-7}
4	RTKs	653.92	1.98×10^{-6}
5	Grb2	85.72	2.6×10^{-7}
6	SOS	29.52	8.95×10^{-8}
7	Ras-GTP	38.99	1.18×10^{-7}
8	Raf	39.59	1.2×10^{-7}
9	MEK	53.27	1.61×10^{-7}
10	ERK	36.92	1.12×10^{-7}
11	PI3K	16.69	5.06×10^{-8}
12	SMO	25.78	7.8×10^{-8}
13	PTEN	30.10	9.12×10^{-8}
14	PDK1	13.09	3.97×10^{-8}
15	AKT	34.59	1.05×10^{-7}
16	mTOR	48.58	1.47×10^{-7}
18	NF2	40.29	1.22×10^{-7}
19	Cyclin D	59.45	1.8×10^{-7}
20	CDK4	168.40	5.1×10^{-7}
21	p16	18.52	5.61×10^{-8}
22	RB	45.66	1.38×10^{-7}
23	E2F	27.28	8.27×10^{-8}
24	G1-S	224.50	6.68×10^{-7}
25	p14	18.52	5.61×10^{-8}
26	MDM2	96.32	2.92×10^{-7}
27	p53	25.52	7.73×10^{-8}
28	Cyclin E	8.23	2.5×10^{-8}
29	CDK2	15.78	4.78×10^{-8}
30	p21	116.54	3.53×10^{-7}
31	p27	57.25	2.28×10^{-7}