



Nutritional Epidemiology

Relationship between Adherence to the 2019 Canada's Food Guide Recommendations on Healthy Food Choices and Nutrient Intakes in Older Adults

Didier Brassard¹, Stéphanie Chevalier^{1,2,*}¹ School of Human Nutrition, McGill University, Montréal, QC, Canada; ² Research Institute of the McGill University Health Centre, Montréal, QC, Canada

A B S T R A C T

Background: Following Canada's food guide (CFG) recommendations should ensure adequate nutrient intakes. Older adults have increased needs for certain nutrients and nutrient density; the extent to which adherence to CFG recommendations can help reduce inadequate nutrient intakes is unknown.

Objectives: We aimed to assess the relationship between adherence to CFG recommendations on healthy food choices and intake of key nutrients in adults 65 y and older from the Canadian Community Health Survey 2015 - Nutrition.

Methods: Secondary analysis of data from 4093 older adults of the Canadian Community Health Survey 2015 - Nutrition (mean age, 73.6 y, 54% females). Dietary intakes were measured using an interviewer-administered 24-h dietary recall, including 1 repeat in a subsample (42%). The National Cancer Institute multivariate method was used to estimate usual (i.e., long-term) dietary intakes. Adherence to CFG recommendations was measured using the Healthy Eating Food Index (HEFI)-2019 score. Simple linear and logistic regression models estimated the effect of increased HEFI-2019 score on usual nutrient intakes and the prevalence of inadequate nutrient intakes (i.e., below the estimated average requirements), respectively.

Results: Compared with the prevalence of inadequate intakes at median HEFI-2019 score (46.4/80 points), a higher HEFI-2019 (+11 points) was associated with reductions in the prevalence of inadequate intakes of magnesium, vitamin B6 and protein [−19.8% (95% confidence interval (CI): −30.8, −8.9), −12.7% (95% CI: −22.5, −3.0), and −4.7% (95% CI: −9.4, −0.1), respectively]. In contrast, data for higher HEFI-2019 scores were compatible with increased prevalence of inadequate intakes of folate, vitamin D, and calcium [4.0% (95% CI: −8.4, 16.3), 2.6% (95% CI: 1.1, 4.0), and 2.3% (95% CI: −3.0, 7.5), respectively].

Conclusions: Based on dietary intakes of Canadian older adults in 2015, increasing the degree of adherence to CFG recommendations on healthy food choices may reduce nutrient intake inadequacy for most key nutrients except folate, vitamin D, and calcium.

Keywords: older adults, 24-h dietary recalls, CFG, HEFI-2019, Canada's food guide, dietary guidelines, Healthy Eating Food Index, Canadian Community Health Survey, CCHS

Introduction

Previous editions of the Canada's food guide (CFG) informed on food choices to prevent malnutrition as the primary goal. Recommendations of the CFG-2007 were expressed in terms of number of servings to consume every day for the broad categories of foods: Vegetables and fruits, Grains, Dairy and Alternatives, and Meat and Alternatives [1,2]. The CFG-2007 recommended daily servings based on comprehensive diet simulations, thus ensuring that Dietary Reference Intakes were met

[2]. The latest CFG (2019) adopted a different approach and primarily aimed at chronic disease risk reduction [3–6]. In addition, recommendations in CFG-2019 (e.g., “eat more often ...”) are more flexible than those of CFG-2007 and rely on proportions of food categories to constitute a healthy plate without recommended quantitative daily servings.

The release of the CFG-2019 elicited positive reactions [7,8] but also raised concerns on nutrient adequacy. For example, preliminary modeling of eating patterns consistent with the CFG-2019 plate snapshot revealed that adherence may be

Abbreviations: CCHS, Canadian Community Health Survey; CFG, Canada's food guide; HEFI-2019, Healthy Eating Food Index-2019.

* Corresponding author E-mail address: stephanie.chevalier@mcgill.ca (S. Chevalier).

<https://doi.org/10.1016/j.tjnut.2023.07.005>

Received 23 February 2023; Received in revised form 24 June 2023; Accepted 11 July 2023; Available online 17 July 2023
0022-3166/© 2023 American Society for Nutrition. Published by Elsevier Inc. All rights reserved.

insufficient to meet calcium and vitamin D requirements [9]. The ability of consumers and policymakers to properly implement CFG-2019 recommendations on protein foods was also found unclear [10]. In the Canadian Community Health Survey (CCHS) 2015 – Nutrition, animal-based protein foods contributed >2 thirds of total protein intake [11] as well as to overall nutrient intakes [12], which contrasts with CFG-2019 recommendations aiming at increasing intake of plant-based foods. Accordingly, following CFG-2019 recommendations while failing to include nutrient-dense foods and beverages could amplify the proportion of the population with inadequate nutrient intakes, especially for certain age groups at higher risk. In that regard, maintaining adequate nutrient intakes is a challenge for older adults [13,14]. Older adults face social (e.g., loneliness, inability to buy or prepare foods) and physiological (e.g., changes in taste, loss of appetite, malabsorption) barriers to consuming a nutrient-dense diet [14]. Notably, calcium, vitamin D, fiber, potassium, and protein were identified as nutrients of concern among older adults in the United States 2015 Dietary Guidelines report [15]. Similarly, a high proportion of long-term care residents in Canada had inadequate calcium, folate, vitamin B6, and magnesium intakes [16]. In sum, the 1-size-fits-all recommendations in CFG-2019 may not be well-suited for older adults who face unique challenges to healthy eating compared with other strata of the population.

Whether adherence to CFG-2019 recommendations on healthy food choices fulfills nutritional needs of older Canadians is currently unknown; no evidence from contemporary eating patterns exists. The general aim of this study was, therefore, to assess the relationship between adherence to CFG-2019 recommendations, using the Healthy Eating Food Index-2019 (HEFI-2019), and intakes of key nutrients from food sources only of adults aged ≥ 65 y from the CCHS 2015 – Nutrition. The HEFI-2019 score is a metric measuring the alignment of eating patterns with CFG-2019 recommendations on food choices [17,18]. More precisely, we assessed 1) the continuous relationship between the HEFI-2019 score and nutrient intakes and 2) the estimated change in the prevalence of nutrient inadequacy according to a hypothetical increase in HEFI-2019 score. We hypothesized that higher adherence to CFG-2019 recommendations would show an inverse relationship with intakes of nutrients more commonly found or eaten in animal-based foods, such as protein, calcium, vitamin D, iron, zinc, vitamin B6, and vitamin B12, and a positive relationship with intakes of nutrients more commonly found in plant-based foods such as fiber, folate, magnesium, potassium, and vitamin A.

Methods

Study design and participants

This study is based on a sample of older adults aged 65 y or older from the CCHS 2015 – Nutrition [19]. The CCHS 2015 – Nutrition is a nationally representative survey of individuals aged 1 y and older living in private dwellings in the 10 Canadian provinces. Full-time members of the Canadian Forces and individuals living in the Territories, on reserves, in remote areas, and in institutions were not included. Data collection occurred between January 1 and December 31, 2015. Respondents aged <65 y ($n = 16,394$) and those reporting 0 energy on their first 24-h dietary recall intake were excluded ($n = 4$), yielding a final

sample of 4089 respondents. The public use microdata file of CCHS 2015 – Nutrition was obtained from Statistics Canada.

Data collection and dietary assessment

Interviewer-administered 24-h dietary recalls were used to assess dietary intakes. The interviews were mostly (98%) conducted in person for the first 24-h recall and by telephone for the second 24-h recall. All 24-h recall interviews were structured according to the Automated Multiple Pass Method [19]. Portion size estimation of foods and beverages in plates, bowls, glasses, and mugs was facilitated using a food booklet designed for the survey [19]. All respondents completed 1 24-h recall, and a subsample of 1706 respondents (42%) completed a second 24-h recall. Nutrient intakes were calculated based on the Canadian Nutrient File 2015 [20], except for intakes of free sugars, which were recently published by Health Canada [21]. Total food intakes expressed in reference amounts [22] were calculated for each respondent, 24-h dietary recall, and HEFI-2019 food and beverage categories. The interviewers measured body weight of respondents with a standard scale (LifeSource Scales Model US-321).

The HEFI 2019

The HEFI-2019 is a continuous score that measures the degree of adherence between dietary intakes and CFG 2019 recommendations on healthy food choices. Complete details about the development and evaluation of the HEFI-2019 are available elsewhere [17,18]. Briefly, the HEFI-2019 has 10 components, including 5 based on intake of foods (*vegetables and fruits*, *whole-grain foods*, *grain foods ratio*, *protein foods*, *plant-based protein foods*), 1 on beverages (*beverages*), and 4 on nutrients (*fatty acids ratio*, *saturated fats*, *free sugars*, and *sodium*). The *saturated fats*, *free sugars*, and *sodium* components are so-called “moderation” components, for which higher scores are attributed to intakes below thresholds consistent with CFG-2019 recommendations. The remaining components are “adequacy” components, for which higher scores are attributed to greater intakes. Details about each component and scoring standards are presented in [Supplementary Table 1](#). The total HEFI-2019 score sums ≤ 80 points, with higher scores indicating greater adherence to recommendations. The HEFI-2019 was evaluated using dietary intake data from the CCHS 2015 – Nutrition [18]. Notably, the variability of the HEFI-2019 score was sufficient, the HEFI-2019 score was correlated with the Healthy Eating Index-2015 ($r = 0.79$), and differences in HEFI-2019 score were found between subgroups with known diet quality differences based on sex, age, and smoking status [18]. Finally, a prospective analysis of UK Biobank data suggested that a hypothetical increase in HEFI-2019 score reduces the 11-y risk of cardiovascular disease in middle-aged and older adults [6].

Nutrient intakes

Key nutrients for older adults were selected a priori according to data availability in the CCHS 2015 – Nutrition and previous literature [13,23]. The nutrients were protein, calcium, vitamin D, iron, zinc, vitamin B6, vitamin B12, dietary folate equivalent (folate hereafter), magnesium, potassium, fiber, and retinol activity equivalent (vitamin A hereafter). Nutrient intakes from supplements were not included since CFG recommendations on healthy food choices focus on foods and beverages. The

proportion of respondents with inadequate intakes was estimated using the cut-point method and the estimated average requirements (EAR) of the Dietary Reference Intake [24]. Because there is no EAR for potassium and fiber, the prevalence of intake inadequacy cannot be assessed. Instead, the proportion of respondents with intakes above the AI value was considered.

Recognizing the consensus that recommended protein intakes for older adults should be higher than current recommendations [25–27], we also estimated the proportion of respondents with inadequate protein intakes at cut-offs higher than the current EAR. The hypothetical EAR cut-offs of 0.8 and 1.0 g of protein/kg of body weight were selected to reflect hypothetical higher RDA of 1.0 and 1.2 g/kg, respectively. The latter analysis was not pre-specified and is considered exploratory.

Statistical analyses

Sampling weights provided by Statistics Canada were used in all analyses to reflect the Canadian population of older adults in 2015, and bootstrap replicate weights for variance estimation. The sampling weights accounting for missing data on body weight were also used where appropriate. Analyses were performed in SAS Studio version 3.8 (SAS Institute) and R version 4.2.2 (R Foundation for Statistical Computing).

To assess the relationship between the HEFI-2019 score and nutrient intakes, analyses involved 4 steps, detailed below: 1) measurement error correction of dietary intakes to estimate usual intakes using the NCI Markov Chain Monte Carlo (MCMC) multivariate method; 2) linear regression of continuous nutrient intakes on total HEFI-2019 scores; 3) logistic regression of the proportion of individuals below EAR on total HEFI-2019 scores; and 4) variance estimation using bootstrap replicate weights provided by Statistics Canada.

First, the 24-h dietary recalls are mainly affected by within-individual random errors, which require correction to estimate distribution of intakes [28,29]. To account for correlated random errors of all dietary constituents of the HEFI-2019 (e.g., vegetables and fruits, free sugars) and those of nutrient intakes (e.g., protein), the NCI MCMC multivariate method was used (MULTIVAR_MCMC_MACRO_V2.1 and MULTIVAR_DISTRIB_MACRO_V2.1, 2017) [30,31]. Briefly, the multivariate method uses Monte Carlo simulation to estimate distribution of usual intakes (i.e., long-term intakes) of multiple dietary constituents correlated with each other. The multivariate method also accounts for systematic differences because of “nuisance” factors (day of the week, sequence of recall) and considers foods that are episodically consumed (e.g., plant-based protein foods) [30].

Nutrients with common food sources were modeled together to have parsimonious measurement error correction models, yielding 5 different models, each combining 1–4 nutrients with the 15 dietary constituents of the HEFI-2019 (see below). Model 1 included protein only; model 2 included calcium and vitamin D; model 3 included iron, zinc, vitamin B6, and vitamin B12; model 4 included folate, magnesium, potassium, and fiber; model 5 included vitamin A only. The 15 dietary constituents of the HEFI-2019 were vegetables and fruits, whole-grain foods, animal-based protein foods, plant-based protein foods, unsweetened milk, water, and other healthy beverages (e.g., unsweetened coffee or tea), refined grain foods, “other” foods not recommended, “other” beverages not recommended (i.e., sugary drinks, artificially sweetened beverages, vegetable and

fruit juices, sweetened milk and plant-based beverages, alcohol), free sugars, monounsaturated fats, polyunsaturated fats, saturated fats, sodium, and total energy intake. A total of 5 food and beverage categories (i.e., whole-grain foods, refined grain foods, plant-based protein foods, “other” beverages not recommended, and unsweetened milk) were considered episodically consumed since $\geq 10\%$ of respondents reported 0 consumption on the first 24-h recall [31]. All the remaining foods and nutrients were considered as consumed daily.

The measurement error correction models were stratified by sex to reflect sex-specific random variations in dietary intakes and to derive sex-specific associations [31,32]. The models also included the covariates age (indicator for 71 y or older), sequence of recall (indicator for second recall), and weekend (indicator for a 24-h recall of intakes on Friday, Saturday, or Sunday). In addition, the model for protein intake also included body weight (kg) as a covariate to derive protein intake per kg and assess the proportion of individuals below the EAR. A total of 500 pseudo-individuals were generated in the Monte Carlo simulation step of the NCI multivariate method. Simulations from each stratum (i.e., males and females) were pooled together. The HEFI-2019 scoring algorithm was then applied using the modeled dietary constituents among the 500 pseudo-individuals to derive HEFI-2019 scores. Descriptive statistics were calculated based on usual intakes among the 500 pseudo-individuals, including the prevalence of intake inadequacy, mean (SD) dietary intakes, and HEFI-2019 scores as well as correlations between dietary constituents of the HEFI-2019 and nutrient intakes.

Second, based on the Monte Carlo simulation data, simple linear regression models were used to assess the relationship between the continuous HEFI-2019 score as the independent variable and continuous nutrient intakes as the dependent variable. A restricted cubic spline transformation with 5 knots (percentiles 5, 27, 50, 73, and 95) was applied a priori to the HEFI-2019 score to assess potential non-linearity [33]. To estimate change according to feasible increases in adherence, expected nutrient intake differences were calculated according to an increase in HEFI-2019 score from the median score to the 90th percentile of the usual intake distribution. In other words, we estimated nutrient intake difference according to a hypothetical change where respondents would have had high HEFI-2019 scores compared with the HEFI-2019 score respondents had on average, taken as the median of this sample.

Third, also based on the Monte Carlo simulation data, logistic regression models were used to assess the odds of having nutrient intake below the age- and sex-specific EAR and the continuous HEFI-2019 score. A restricted cubic spline transformation was also applied to the HEFI-2019 score. Predicted odds of nutrient intake inadequacy were generated for the 90th percentile and the median. Both predicted odds were then re-expressed as risk of inadequacy ($risk = \frac{e^X}{1+e^X}$; where X corresponds to predicted odds at a given HEFI-2019 score percentile). The expected change in the prevalence of inadequate nutrient intakes, i.e., risk difference, was calculated by the difference between the estimated risk of inadequacy at the 90th percentile compared with the median HEFI-2019 score.

Fourth, steps 1–3 were repeated 500 times using bootstrap replicate weights to generate SEs and 95% CI via normal

approximation. The convergence of bootstrap SEs and the normality of bootstrap estimates were confirmed graphically. Data from 1 bootstrap replicate was removed for potassium because of non-convergence.

Results

HEFI-2019 score and dietary constituents

The mean (SD) HEFI-2019 score among all adults 65 y and older was 46.0 (8.9) (/80 points). Among age and sex subgroups, the mean HEFI-2019 score was the highest in females, 65–70 y (48.2 points), and the lowest in males, 65–70 y (44.9 points; [Figure 1](#)).

[Supplementary Tables 2 and 3](#) present the usual mean intakes and percentile of the distribution of foods and beverages, as well

as nutrients, contributing to the HEFI-2019 score, respectively. Older adults consumed 86% of their total protein foods from animal-based sources and 65% of their total grain foods intake from non-whole grain foods.

[Supplementary Figure 1](#) presents Pearson correlations between food and beverage categories contributing to the HEFI-2019 score and key nutrients. Most food and beverage categories recommended in CFG 2019 were correlated with greater nutrient intakes. Notably, the highest correlations were observed between milk and calcium ($r = 0.56$), milk and vitamin D ($r = 0.53$), vegetables and fruits and fiber ($r = 0.54$), as well as whole-grain foods and fiber ($r = 0.54$). Among foods and beverages not recommended in CFG, non-whole grain foods were the only group showing positive correlations with nutrient intakes, including folate ($r = 0.16$) and with vitamin D ($r = 0.12$).




FIGURE 1. Distribution of total HEFI-2019 score, based on usual dietary intakes in 4089 adults aged 65 y and more from the CCHS 2015 - Nutrition. Diamonds are means. Left and right whiskers indicate the 5th and 95th percentile, respectively. Dietary intakes were modeled using the NCI multivariate method to estimate usual intakes (see Methods). CCHS, Canadian Community Health Survey; HEFI-2019, Healthy Eating Food Index-2019.

Prevalence of inadequate nutrient intake

The prevalence of inadequate nutrient intake ranged from 1% (iron) to 96% (vitamin D) (Figure 2). The prevalence of inadequate intakes was high for vitamin D, calcium, and magnesium (96%, 83%, 64%, respectively) and moderately high for vitamin B6, vitamin A, folate, and zinc (38%, 36%, 30%, 28%, respectively) (Figure 2). Results were similar for most nutrients when stratified by Dietary Reference Intakes age and sex group (Supplementary Table 4).

Relationship between HEFI-2019 score and nutrient intakes

Relationships between the HEFI-2019 score and nutrient intakes are illustrated in Figure 3. Respondents with higher HEFI-2019 scores had higher intakes of fiber, magnesium, vitamin B6, potassium, and protein but lower intakes of vitamin D, vitamin B12, folate, and iron. The associations between the HEFI-2019 score and intakes of vitamin A, calcium, and zinc were weak or null. Table 1 presents the expected differences in nutrient intakes associated with an increase in HEFI-2019 score to the 90th percentile of the score distribution compared with median HEFI-2019 score. For example, increasing HEFI-2019 from the median to the 90th percentile of the score distribution was associated with a 3.4 g/d higher fiber intake (95% CI: 2.0, 4.8; Table 1). In contrast, the same HEFI-2019 score increase was associated with a 0.5 µg/d lower vitamin D intake (95% CI: −1.0, 0.0; Table 1).

Relationship between the HEFI-2019 score and nutrient intake inadequacy

Inadequate nutrient intakes

Figure 4 presents the prevalence and the difference in the prevalence of inadequate nutrient intake according to the HEFI-2019 score. An increase in HEFI-2019 score to the 90th percentile of the score distribution compared with median HEFI-2019 score was associated with a reduction of 19.8, 12.7, and 4.7% points in the prevalence of inadequate nutrient intakes for magnesium, vitamin B6, and protein, respectively. Inversely, the same HEFI-2019 score increase was associated with a minor increase in the prevalence of inadequate vitamin D intakes (prevalence difference, +2.6%; 95% CI: 1.1, 4.0%). The prevalence of inadequate iron intake was 0% and thus unaffected by hypothetical HEFI-2019 score change. For vitamin A, zinc, vitamin B12, calcium, and folate, the 95% CI was wide and compatible with both increase and decrease in the prevalence of inadequate intakes (Figure 3).

Hypothetical higher EAR for protein. Compared with the median HEFI-2019 score, scores at the 90th percentile decreased the proportion of respondents with protein intakes below hypothetical EAR of 0.8 g/kg/d and 1.0 g/kg/d (Supplementary Figure 2). For 0.8 g/kg/d, the prevalence of inadequacy decreased from 29.3% to 18.5% (prevalence difference: −10.8; 95% CI: −20.0, −1.6). For 1.0 g/kg/d, the prevalence of inadequacy decreased




FIGURE 2. Prevalence of inadequate nutrient intakes in 4089 adults aged 65 y and older from the CCHS 2015 - Nutrition. Only nutrient intakes from foods were considered (i.e., excluding intakes from dietary supplements). Inadequate intakes are intakes below the age- and sex-specific Estimated Average Requirements. Potassium and fiber are not shown because only Adequate Intake values are available for these nutrients. All dietary intakes were modeled using the NCI multivariate method to estimate usual intakes (see Methods). The 95% CI was estimated using 500 bootstrap weight replicates. CCHS, Canadian Community Health Survey.




FIGURE 3. Linear regression of nutrient intake on the total HEFI-2019 score in 4,089 adults aged 65 y or more from the CCHS 2015 - Nutrition. A positive relationship indicates that greater HEFI-2019 scores are associated with greater nutrient intake, and inversely. For visualization purposes, data points are from a random sample of 200 respondents selected proportionally to sampling weights. All dietary intakes were modeled using the NCI multivariate method to estimate usual intakes (see Methods). The 95% CI was estimated using 500 bootstrap weight replicates. CCHS, Canadian Community Health Survey; HEFI-2019, Healthy Eating Food Index-2019; DFE, Dietary Folate Equivalent; RAE, Retinol Activity Equivalent.

from 64.7% to 50.8% (prevalence difference: -13.9 ; 95% CI: $-25.0, -2.8$).

AIs for fiber and potassium

Compared with the median HEFI-2019 score, a score at the 90th percentile increased the proportion of respondents with intakes above the AI for both fiber and potassium. For fiber, the prevalence increased from 6.5% to 20.5% (prevalence difference: $+14.1$; 95% CI: 5.1, 23.0). For potassium, the prevalence increased from 25.1% to 36.8% (prevalence difference: $+11.7$; 95% CI: 2.2, 21.1).

Discussion

The objective of this study was to describe the relationship between adherence to CFG-2019 recommendations on healthy food choices, measured using the HEFI-2019, and nutrient intakes from food sources in adults 65 y and older from Canada in 2015. We found that higher adherence was associated with higher intakes of fiber, magnesium, vitamin B6, potassium, and protein. Had respondents had a higher adherence, we estimated that the prevalence of nutrient inadequacy from foods and beverages for magnesium, vitamin B6, and protein would have been 20%, 13%, and 5% lower, respectively. However, a higher adherence was not associated with higher intakes of calcium, zinc, iron, folate, vitamin B12, and vitamin D. In turn, we estimated that the high prevalence of calcium inadequacy would not have changed and that of food-based vitamin D would have been

3% higher in 2015. Overall, these results indicate that CFG-2019 recommendations are insufficient to mitigate nutrient intake inadequacy for certain key nutrients based on the eating patterns of adults aged 65 y or more from Canada in 2015. These findings partially confirm our hypothesis that nutrients typically found in animal-based foods (e.g., iron, vitamin B12, calcium, vitamin D) were inversely associated with adherence to CFG.

Few studies have examined nutrient intake inadequacy according to adherence to the CFG-2019 recommendations. Barr [9] assessed the probability of intake inadequacy upon adherence to an eating pattern consistent with foods depicted in the CFG-2019 plate snapshot. The probability of inadequacy was near 100% for calcium and vitamin D for males and females aged 71 y or more. This result is consistent with findings in the present study, where a higher adherence to recommendations on healthy food choices was insufficient to mitigate the prevalence of inadequacy for calcium and vitamin D. The lack of specific recommendations regarding dairy foods and alternatives in CFG-2019, and consequently, the absence of points in the HEFI-2019 may partly explain these findings. Among food and beverage categories contributing to the HEFI-2019, unsweetened milk had the highest correlation with intakes of calcium ($r = 0.56$) and vitamin D ($r = 0.53$; [Supplementary Figure 1](#)). Indeed, the main food source of calcium and vitamin D in Canadians' diet was the "Milk & Alternatives" food group in 2015 [34–36]. Of note, the prevalence of inadequacy for vitamin D was very high in this sample ($>95\%$), supporting the notion that meeting vitamin D recommendations through foods alone is an important

TABLE 1

Usual nutrient intake difference between Healthy Eating Food Index (HEFI)-2019 scores at the 90th percentile and the median HEFI-2019 score distribution in 4089 adults aged 65 y or more from the Canadian Community Health Survey 2015 - Nutrition

Nutrients ¹	Adults, 65 y+ (n = 4089)	Dietary reference intake group			
		Males, 65–70 y (n = 567)	Females, 65–70 y (n = 720)	Males, 71 y+(n = 1246)	Females, 71 y+(n = 1556)
Protein, g/d	3 (–2, 8)	2 (–4, 9)	5 (–1, 11)	2 (–4, 8)	4 (–2, 10)
Calcium, mg/d	4 (–50, 57)	–16 (–96, 65)	21 (–47, 89)	–14 (–88, 60)	25 (–38, 89)
Vitamin D, µg /d	–0.5 (–1.0, 0.0)	–0.8 (–1.6, –0.1)	–0.1 (–0.7, 0.5)	–0.7 (–1.5, 0.0)	–0.1 (–0.7, 0.6)
Iron, mg/d	–0.2 (–1.0, 0.5)	–0.6 (–1.8, 0.6)	0.2 (–0.7, 1.1)	–0.6 (–1.8, 0.6)	0.3 (–0.5, 1.1)
Vitamin B12, µg /d	–0.3 (–0.7, 0.1)	–0.4 (–0.9, 0.0)	–0.1 (–0.7, 0.4)	–0.4 (–0.8, 0.1)	–0.1 (–0.6, 0.4)
Vitamin B6, mg/d	0.1 (0.0, 0.2)	0.1 (0.0, 0.2)	0.1 (0.0, 0.3)	0.1 (0.0, 0.2)	0.1 (0.0, 0.2)
Zinc, mg/d	0.0 (–0.7, 0.7)	–0.4 (–1.5, 0.7)	0.4 (–0.3, 1.2)	–0.3 (–1.3, 0.6)	0.5 (–0.2, 1.2)
Folate, µg /d	–13 (–39, 14)	–15 (–56, 26)	–6 (–36, 25)	–16 (–55, 22)	–2 (–31, 27)
Fiber, g/d	3.4 (2.0, 4.8)	4.5 (2.2, 6.8)	3.0 (1.2, 4.8)	4.3 (2.2, 6.4)	3.0 (1.3, 4.7)
Magnesium, mg/d	33 (14, 52)	44 (14, 73)	30 (6, 55)	39 (13, 65)	31 (9, 53)
Potassium, mg/d	173 (17, 328)	230 (6, 455)	170 (–27, 368)	214 (3, 424)	178 (–7, 363)
Vitamin A, RAE/d	16 (–37, 69)	9 (–54, 71)	26 (–54, 106)	10 (–48, 68)	26 (–42, 95)

CCHS, Canadian Community Health Survey; HEFI-2019, Healthy Eating Food Index-2019; RAE, Retinol Activity Equivalent.

¹ Values are expected nutrient intake differences (95% CI) for a HEFI-2019 score at the 90th compared with the 50th percentile of the score distribution and were estimated using linear regression models. Estimates reflect the expected nutrient intake difference when HEFI-2019 scores are increased to correspond to the 90th percentile of the score distribution, compared with the median HEFI-2019 score. All dietary intakes were modeled using the NCI multivariate method to estimate usual intakes (see Methods). The 95% CI was estimated using 500 bootstrap weight replicates. Sample size indicates the number of respondents without applying sampling weights, but other estimates were weighted to reflect the Canadian population.

challenge [36]. Similarly, data from the CCHS indicate that calcium intake inadequacy in supplement non-users increased from 58% to 68% between 2004 and 2015, respectively [35]. All in all, higher adherence to CFG-2019 recommendations on food choices was insufficient to mitigate the high prevalence of inadequate calcium and vitamin D intakes. Thus, meeting calcium and vitamin D needs would require additional strategies, such as specific recommendations in CFG in addition to supplementation or food fortification.

A review by Fernandez et al. [10] highlighted potential gaps in the application of CFG-2019 recommendations, notably whether the CFG-2019 permitted the adequate consumption of nutrient-rich protein foods or not. Our results revealed that higher adherence to CFG-2019 recommendations slightly reduced the low proportion of older adults with intakes below the current EAR (0.66 g/kg) and reduced by 11% and 14% the proportion of those below the hypothetical EAR of 0.8 g/kg and 1.0 g/kg. Notwithstanding, the prevalence of inadequacy for the hypothetical EAR would remain considerable, with 19% and 51% inadequate protein intakes, thus representing an additional risk factor for sarcopenia, respectively [26]. Furthermore, these findings only reflect the total daily protein intake, and the impact on overall protein food quality or distribution during the day is unknown. Future studies should investigate the relationship between higher adherence to CFG-2019 and overall protein food quality and should examine how to increase consumption of high-quality plant-based protein foods. In that regard, a qualitative study among older adults highlighted that health benefits and food preparation skills would facilitate consumption of plant-based protein foods [37]. Increasing consumption of unsweetened soy beverages (currently 0 in the present study) or unsweetened milk (mean intake of 125 mL/d; [Supplementary Table 2](#)) could also contribute to higher intake of high-quality protein. Finally, we stress that the CCHS 2015 – Nutrition excluded individuals living in institutions who likely have additional needs and barriers to consuming higher protein diets [14].

The inverse associations between the HEFI-2019 and intakes of iron, folate, and vitamin B12 also raise concerns. On the one hand, the prevalence of inadequacy for these nutrients did not increase at the population level, partly because the EAR can be met at relatively low intakes for these nutrients. On the contrary, at the individual level, meeting nutrient requirements while following CFG-2019 recommendations may be more difficult in individuals at risk of inadequate intakes, e.g., older adults with loss of appetite [13,14]. Further, vitamin B12 may be particularly important for older adults, and recent evidence supports that food groups contribute differently to reducing risk of deficiency [38]. Meeting folate requirements while following CFG may also be challenging for some individuals since refined grains, not recommended in CFG, showed the highest correlations with folate intake in the present study ([Supplementary Figure 1](#)). Indeed, non-whole-grain foods are subject to mandatory fortification in Canada, which is not the case for whole-grain foods [39]. In sum, CFG-2019 recommendations should be carefully implemented to ensure meeting nutrient intakes both at the population and individual levels. Health Canada published additional information in 2022 on how to apply dietary guidelines to support health professionals and policymakers [40]. The information describes the type and frequency of foods to support nutritional needs. The extent to which the additional information helps older adults shift their eating patterns to be both consistent with CFG-2019 and provide optimal nutrient intakes remains to be determined. The findings of the present study highlight that eating patterns of older adults in 2015 failed to provide optimal intakes of calcium, folate, and vitamin D despite adhering to CFG-2019 recommendations, thereby supporting the need for additional guidance, perhaps with quantitative recommendations.

The use of national survey data to estimate the impact of higher adherence on the prevalence of nutrient inadequacy in Canada is a strength of this study. However, it must be recognized that relationships between adherence to CFG-2019 recommendations and nutrient intakes reflect the underlying eating




FIGURE 4. Prevalence of inadequate intakes and difference for HEFI-2019 scores at the 90th compared with the 50th percentile of the score distribution in 4089 adults aged 65 y or more from the CCHS 2015 - Nutrition. Inadequate intakes are intakes below the age- and sex-specific Estimated Average Requirements. Potassium and fiber are not shown because only Adequate Intake values are available for these nutrients. All dietary intakes were modeled using the NCI multivariate method to estimate usual intakes (see Methods). The 95% CI was estimated using 500 bootstrap weight replicates. CCHS, Canadian Community Health Survey; HEFI-2019, Healthy Eating Food Index-2019.

patterns of adults 65 y or older from Canada in 2015. A variety of underlying food and beverage consumption patterns may be highly consistent with CFG-2019 recommendations yet, have diverging associations with intakes of certain nutrients. As well, eating patterns of older adults, though generally stable, may evolve over time, which would result in different associations between adherence to CFG-2019 and nutrient intakes. Another key strength of this work is the use of the NCI MCMC multivariate method to account for random errors in dietary intakes measured with 24-h recall [30,31]. Using the multivariate method permitted the joint assessment of the relationship between usual nutrient intakes and adherence to CFG-2019 recommendations on healthy food choices, as measured with the HEFI-2019, and the modeling of higher adherence to CFG-2019 recommendations compared with median adherence. Limitations must be addressed. First, self-reported dietary intakes with 24-h dietary recall are affected by systematic errors or bias [41,42]. The working assumption that intakes are unbiased cannot be verified. Of note, we did not adjust for “misreporting status” to account for energy under-reporting in the present study. The proportion of under-reporters among older adults is not higher than in other age and sex group samples of the CCHS 2015 – Nutrition [43], and evidence indicates that diet quality based on the 24-h dietary recall may not be prone to large bias [44]. Second, the food composition database, the Canadian Nutrient File 2015, may not adequately reflect the nutrient profile of foods actually consumed [45]. Accordingly, estimates of the prevalence of nutrient inadequacy based on self-reported

dietary intake data are not as accurate as estimates based on biomarker data. Third, our results cannot be used to extrapolate nutrient intake inadequacy at the national level. Although calcium and vitamin D supplements could contribute to reducing inadequacies [35,36], nutrient intakes from dietary supplements were not included since the objective was to assess the relationship between nutrients and adherence to CFG-2019 recommendations on healthy food choices [4,5,17].

In conclusion, higher adherence to CFG-2019 recommendations on healthy food choices is associated with greater intakes of protein, fiber, magnesium, and many nutrients considered, based on eating patterns of adults aged 65 y or more from the CCHS 2015 – Nutrition. However, higher adherence is associated with lower intakes of iron, folate, vitamin B12, and vitamin D and not associated with calcium, zinc, or vitamin A intakes, thus insufficient to mitigate dietary inadequacy in these key nutrients. Knowledge of shortfalls of CFG-2019 recommendations regarding nutrient intake inadequacy is relevant to help the government, policymakers, and registered dietitians provide more comprehensive and appropriate recommendations to older adults. Our findings support the need for additional information, as that recently published by Health Canada, to help ensure that eating patterns are both consistent with CFG-2019 and adequate nutrient intakes [40]. Future studies should investigate how eating patterns changed after the publication of CFG-2019 and the additional information and how these changes may affect the relationship between adherence to recommendations and nutrient intakes.

Funding

The authors reported no funding received for this study.

Author contributions

The authors' responsibilities were as follows – DB, SC: designed the research; DB: performed statistical analysis; DB: wrote the first draft of the manuscript; DB, SC: critically reviewed the manuscript.

Conflict of interest

DB was a casual employee of Health Canada (2019–2020), held a doctoral training award from the Fonds de recherche du Québec – Santé (2019–2021), and holds a Canadian Institutes of Health Research (CIHR) Fellowship award (MFE-181852). SC receives research funding from the CIHR, Dairy Farmers of Canada, and the Canadian Foundation for Dietetics Research. None of these agencies has funded nor was involved in this work.

Data availability

The Canadian Community Health Survey 2015 – Nutrition Public Use Microdata File is available upon request at <https://www150.statcan.gc.ca/n1/en/catalogue/82M0024X2018001>. The analytic code will be made freely available at https://github.com/didierbrassard/hefi2019_nutrient.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.tjn.2023.07.005>.

References

- Health Canada, Eating well with Canada's food guide, Health Canada, Ottawa, 2007, pp. 1–6.
- S.W. Katamay, K.A. Esslinger, M. Vigneault, J.L. Johnston, B.A. Junkins, L.G. Robbins, et al., Eating well with Canada's Food Guide (2007): development of the food intake pattern, *Nutr. Rev.* 65 (4) (2007) 155–166.
- Health Canada, Food, nutrients and health: interim evidence update 2018 for health professionals and policy makers, Health Canada, Ottawa, 2019.
- Health Canada, Canada's food guide, Health Canada, Ottawa, 2019.
- Health Canada, Canada's dietary guidelines for health professionals and policy makers, 2019.
- D. Brassard, H.D. Manikpurage, S. Thériault, B.J. Arsenault, B. Lamarche, Greater adherence to the 2019 Canada's Food Guide recommendations on healthy food choices reduces the risk of cardiovascular disease in adults: a prospective analysis of UK Biobank data, *Am. J. Clin. Nutr.* 116 (6) (2022) 1748–1758.
- K.E. Asher, S. Doucet, A. Luke, Registered dietitians' perceptions and use of the plant-based recommendations in the 2019 Canada's Food Guide, *J. Hum. Nutr. Diet.* 34 (4) (2021) 715–723.
- A.C. Barco Leme, A. Laila, S. Hou, R.M. Fisberg, D.W.L. Ma, M. Fisberg, et al., Perceptions of the 2019 Canada's Food Guide: a qualitative study with parents from Southwestern Ontario, *Appl. Physiol. Nutr. Metab.* (2021) 1–7.
- S.I. Barr, Is the 2019 Canada's Food Guide Snapshot nutritionally adequate? *Appl. Physiol. Nutr. Metab.* 44 (12) (2019) 1387–1390.
- M.A. Fernandez, R.F. Bertolo, A.M. Duncan, S.M. Phillips, R. Elango, D.W.L. Ma, et al., Translating "protein foods" from the new Canada's Food Guide to consumers: knowledge gaps and recommendations, *Appl. Physiol. Nutr. Metab.* 45 (12) (2020) 1311–1323.
- O. Auclair, S.A. Burgos, Protein consumption in Canadian habitual diets: usual intake, inadequacy, and the contribution of animal- and plant-based foods to nutrient intakes, *Appl. Physiol. Nutr. Metab.* 46 (5) (2021) 501–510.
- H. Fabek, D. Sanchez-Hernandez, M. Ahmed, C.P.F. Marinangeli, J.D. House, G.H. Anderson, An examination of contributions of animal- and plant-based dietary patterns on the nutrient quality of diets of adult Canadians, *Appl. Physiol. Nutr. Metab.* 46 (8) (2021) 877–886.
- J. Shlisky, D.E. Bloom, A.R. Beaudreault, K.L. Tucker, H.H. Keller, Y. Freund-Levi, et al., Nutritional considerations for healthy aging and reduction in age-related chronic disease, *Adv. Nutr.* 8 (1) (2017) 17–26.
- D. Rémond, D.R. Shahar, D. Gille, P. Pinto, J. Kachal, M.A. Peyron, et al., Understanding the gastrointestinal tract of the elderly to develop dietary solutions that prevent malnutrition, *Oncotarget* 6 (16) (2015) 13858–13898, 2015.
- B.E. Millen, S. Abrams, L. Adams-Campbell, C.A. Anderson, J.T. Brenna, W.W. Campbell, et al., The 2015 dietary guidelines advisory committee scientific report: development and major conclusions, *Adv. Nutr.* 7 (3) (2016) 438–444.
- H.H. Keller, C. Lengyel, N. Carrier, S.E. Slaughter, J. Morrison, A.M. Duncan, et al., Prevalence of inadequate micronutrient intakes of Canadian long-term care residents, *Br. J. Nutr.* 119 (9) (2018) 1047–1056.
- D. Brassard, L.A. Elvidge Munene, S. St-Pierre, P.M. Guenther, S.I. Kirkpatrick, J. Slater, et al., Development of the Healthy Eating Food Index (HEFI)-2019 measuring adherence to Canada's Food Guide 2019 recommendations on healthy food choices, *Appl. Physiol. Nutr. Metab.* 47 (5) (2022) 595–610.
- D. Brassard, L.A. Elvidge Munene, S. St-Pierre, A. Gonzalez, P.M. Guenther, M. Jessri, et al., Evaluation of the Healthy Eating Food Index (HEFI)-2019 measuring adherence to Canada's Food Guide 2019 recommendations on healthy food choices, *Appl. Physiol. Nutr. Metab.* 47 (5) (2022) 582–594.
- Health Canada, Reference guide to understanding and using the data. Nutrition. 2015 Canadian Community Health Survey, 2017, pp. 1–91.
- Health Canada. [Internet], Canadian Nutrient File, 2015. Available from: <https://food-nutrition.canada.ca/cnf-fce/index-eng.jsp>. Accessed on July 2023.
- H. Rana, M.C. Mallet, A. Gonzalez, M.F. Verreault, S. St-Pierre, Free sugars consumption in Canada, *Nutrients* 13 (5) (2021) 1471.
- Health Canada, Nutrition labelling: Table of reference amounts for food, 2016. Ottawa.
- N. Dorrington, R. Fallaize, D.A. Hobbs, M. Weech, J.A. Lovegrove, A review of nutritional requirements of adults aged ≥65 Years in the UK >=65 Years in the UK, *J. Nutr.* 150 (9) (2020) 2245–2256.
- J.J. Otten, J.P. Hellwig, L.D. Meyers, Dietary reference intakes: the essential guide to nutrient requirements, National Academies Press, Washington, DC, 2006.
- S.M. Phillips, S. Chevalier, H.J. Leidy, Protein "requirements" beyond the RDA: implications for optimizing health, *Appl. Physiol. Nutr. Metab.* 41 (5) (2016) 565–572.
- J. Bauer, G. Biolo, T. Cederholm, M. Cesari, A.J. Cruz-Jentoft, J.E. Morley, et al., Evidence-based recommendations for optimal dietary protein intake in older people: a position paper from the PROT-AGE Study Group, *J. Am. Med. Dir. Assoc.* 14 (8) (2013) 542–559.
- D.A. Traylor, S.H.M. Gorissen, S.M. Phillips, Perspective: protein requirements and optimal intakes in aging: are we ready to recommend more than the recommended daily allowance? *Adv. Nutr.* 9 (3) (2018) 171–182.
- K.W. Dodd, P.M. Guenther, L.S. Freedman, A.F. Subar, V. Kipnis, D. Midthune, et al., Statistical methods for estimating usual intake of nutrients and foods: a review of the theory, *J. Am. Diet. Assoc.* 106 (10) (2006) 1640–1650.
- F.E. Thompson, S.I. Kirkpatrick, A.F. Subar, J. Reedy, T.E. Schap, M.M. Wilson, et al., The National Cancer Institute's dietary assessment primer: A resource for diet research, *J. Acad. Nutr. Diet.* 115 (12) (2015) 1986, 95.
- S. Zhang, D. Midthune, P.M. Guenther, S.M. Krebs-Smith, V. Kipnis, K.W. Dodd, et al., A new multivariate measurement error model with zero-inflated dietary data, and its application to dietary assessment, *Ann. Appl. Stat.* 5 (2B) (2011) 1456–1487.
- S.I. Kirkpatrick, P.M. Guenther, A.F. Subar, S.M. Krebs-Smith, K.A. Herrick, L.S. Freedman, et al., Using short-term dietary intake data to address research questions related to usual dietary intake among

- populations and subpopulations: assumptions, statistical techniques, and considerations, *J. Acad. Nutr. Diet.* 122 (7) (2022) 1246–1262.
- [32] K.A. Herrick, L.M. Rossen, R. Parsons, K.W. Dodd, Estimating Usual Dietary intake from National Health and Nutrition Examination Survey Data Using the National Cancer Institute Method, *Vital Health Stat 2 2* (178) (2018) 1–63.
- [33] F.E. Harrell, General aspects of fitting regression models, in: *Regression modeling strategies: with applications to linear models, logistic and ordinal regression, and survival analysis*, Edition, Springer International Publishing, Cham, 2015, pp. 13–44.
- [34] H. Vatanparast, N. Islam, M. Shafiee, Consumption of Milk and alternatives decreased among Canadians from 2004 to 2015: evidence from the Canadian community health surveys, *BMC Nutr* 7 (1) (2021) 63.
- [35] H. Vatanparast, N. Islam, R.P. Patil, M. Shafiee, S.J. Whiting, Calcium intake from food and supplemental sources decreased in the Canadian population from 2004 to 2015, *J Nutr* 150 (4) (2020) 833–841.
- [36] H. Vatanparast, R.P. Patil, N. Islam, M. Shafiee, S.J. Whiting, Vitamin D intake from supplemental sources but not from food sources has increased in the Canadian population over time, *J. Nutr.* 150 (3) (2020) 526–535.
- [37] V. Drolet-Labelle, D. Laurin, A. Bédard, V. Drapeau, S. Desroches, Beliefs underlying older adults' intention to consume plant-based protein foods: A qualitative study, *Appetite* 180 (2023) 106346.
- [38] H.H. Huang, A.A. Cohen, P. Gaudreau, C. Auray-Blais, D. Allard, M. Boutin, et al., Vitamin B-12 intake from dairy but not meat is associated with decreased risk of low vitamin B-12 status and deficiency in older adults from Quebec, Canada, *J. Nutr.* 152 (11) (2022) 2483–2492.
- [39] [Internet] Government of Canada, Foods to which vitamins, mineral nutrients and amino acids may or must be added, 2022 June 2023. Available from: <https://inspection.canada.ca/food-labels/labelling/industry/nutrient-content/reference-information/eng/1389908857542/1389908896254?chap=0>. Accessed on July 2023.
- [40] [Internet] Health Canada, Canada's Dietary Guidelines, 2022 June 2023. Available from: <https://food-guide.canada.ca/en/guidelines>. Accessed on July 2023.
- [41] L.S. Freedman, J.M. Commins, J.E. Moler, W. Willett, L.F. Tinker, A.F. Subar, et al., Pooled results from 5 validation studies of dietary self-report instruments using recovery biomarkers for potassium and sodium intake, *Am. J. Epidemiol.* 181 (7) (2015) 473–487.
- [42] L.S. Freedman, J.M. Commins, J.E. Moler, L. Arab, D.J. Baer, V. Kipnis, et al., Pooled results from 5 validation studies of dietary self-report instruments using recovery biomarkers for energy and protein intake, *Am. J. Epidemiol.* 180 (2) (2014) 172–188.
- [43] D. Garriguet, Accounting for misreporting when comparing energy intake across time in Canada, *Health Rep* 29 (5) (2018) 3–12.
- [44] S.I. Kirkpatrick, K.W. Dodd, N. Potischman, T.P. Zimmerman, D. Douglass, P.M. Guenther, et al., Healthy eating Index-2015 scores among adults based on observed vs recalled dietary intake, *J. Acad. Nutr. Diet.* 121 (11) (2021) 2233–2241.e1.
- [45] A. Satija, E. Yu, W.C. Willett, F.B. Hu, Understanding nutritional epidemiology and its role in policy, *Adv. Nutr.* 6 (1) (2015) 5–18.