Gene	Phenotype	P (burden)	Literature	Open Targets	Evidence	Why it is worth following up	Key reports
Novel on both branches							
THNSL1	Stroke	1.8×10^{-7}	**Novel**	**Novel**	insufficient	*THNSL1* has not been directly implicated in stroke; it was only among >1,900 genes altered in mouse splenic B cells after protective hypoxic preconditioning, before stroke induction, with no functional follow-up.	Monson NL et al. 2014, PMID 24485041
NAA15	Hypertension	1.2×10^{-8}	**Novel**	**Novel**	insufficient	*NAA15* haploinsufficiency causes congenital heart disease, but hypertension was reported in only 1/19 evaluated individuals ascertained for truncating variants (Cheng H et al. 2018).	Cheng H et al. 2018, PMID 29656860 Ward T et al. 2021, PMID 33557580
Hypothesized on both branches							
SIN3A	Stroke	5.5×10^{-8}	**Hypoth.**	**Hypoth.**	mechanistic / animal	Cerebral ischemia recruits *mSin3A* to REST-corepressor complexes, providing mechanistic support without direct human genetic evidence for stroke.	Noh KM et al. 2012, PMID 22371606
IRS2	Acute kidney failure	9.3×10^{-7}	**Hypoth.**	**Hypoth.**	indirect human genetic + animal	*IRS2* truncating variants increase chronic kidney disease risk independently of diabetes, and ghrelin-mediated protection from ischemic acute kidney injury is lost in *Irs2*-knockout mice.	Zhao Y et al. 2025, PMID 41073786 Takeda R et al. 2006, PMID 16306169
Hypothesized in the literature, Novel in Open Targets							
OSBP	HDL cholesterol	1.1×10^{-7}	**Hypoth.**	**Novel**	mechanistic	*OSBP* knockdown increases cholesterol efflux in CHO cells and raises *ABCA1* in J774 macrophages, providing a mechanistic link to HDL biology without direct human genetic evidence.	Bowden K, Ridgway ND. 2008, PMID 18450749
PODN	Coronary artery disease	5.4×10^{-8}	**Hypoth.**	**Novel**	animal / functional	*Podn*-null mice show excessive neointimal lesion formation after arterial injury; overexpression of human *PODN* inhibits human vascular smooth-muscle-cell migration and proliferation, and *PODN* is expressed in human atheroma.	Hutter R et al. 2013, PMID 24043300
Novel in the literature, Hypothesized in Open Targets							
CPXM2	Rheumatoid arthritis	8.6×10^{-12}	**Novel**	**Hypoth.**	transcriptomic / related disease	No retrieved literature directly links *CPXM2* to rheumatoid arthritis; Open Targets provides low-confidence transcriptomic evidence of differential *CPXM2* expression in rheumatoid arthritis and additional indirect support through ulcerative colitis, a related immune-mediated inflammatory disease.	—
SETD1A	Asthma	8.1×10^{-9}	**Novel**	**Hypoth.**	lung function	No retrieved literature directly links *SETD1A* to asthma. Open Targets provides indirect support through forced expiratory volume, with a low evidence score (0.053).	—