

Useful recurrence relations for multidimensional volumes and monomial integrals*

Nico Schlömer

June 4, 2020

This article gives closed formulas and recurrence expressions for many n -dimensional volumes and monomial integrals. The recurrence expressions are often much simpler, more instructive, and better suited for numerical computation.

n -dimensional unit cube

$$C_n = \{(x_1, \dots, x_n) : -1 \leq x_i \leq 1\}$$

- Volume.

$$|C_n| = 2^n = \begin{cases} 1 & \text{if } n = 0 \\ |C_{n-1}| \times 2 & \text{otherwise} \end{cases} \quad (1)$$

- Monomial integration.

$$\begin{aligned} I_{k_1, \dots, k_n} &= \int_{C_n} x_1^{k_1} \cdots x_n^{k_n} \\ &= \prod_{i=1}^n \frac{1 + (-1)^{k_i}}{k_i + 1} = \begin{cases} 0 & \text{if any } k_i \text{ is odd} \\ |C_n| & \text{if all } k_i = 0 \\ I_{k_1, \dots, k_{i_0}-2, \dots, k_n} \times \frac{k_{i_0}-1}{k_{i_0}+1} & \text{if } k_{i_0} > 0 \end{cases} \end{aligned} \quad (2)$$

n -dimensional unit simplex

$$T_n = \left\{ (x_1, \dots, x_n) : x_i \geq 0, \sum_{i=1}^n x_i \leq 1 \right\}$$

*The LaTeX sources of this article are on <https://github.com/nschloe/useful-recurrence-relations>

- Volume.

$$|T_n| = \frac{1}{n!} = \begin{cases} 1 & \text{if } n = 0 \\ |T_{n-1}| \times \frac{1}{n} & \text{otherwise} \end{cases} \quad (3)$$

- Monomial integration.

$$\begin{aligned} I_{k_1, \dots, k_n} &= \int_{T_n} x_1^{k_1} \cdots x_n^{k_n} \\ &= \frac{\prod_{i=1}^n \Gamma(k_i)}{\Gamma(\sum_{i=1}^n k_i)} \end{aligned} \quad (4)$$

$$= \begin{cases} |T_n| & \text{if all } k_i = 0 \\ I_{k_1, \dots, k_{i_0}-1, \dots, k_n} \times \frac{k_{i_0}}{n + \sum_{i=1}^n k_i} & \text{if } k_{i_0} > 0 \end{cases} \quad (5)$$

Remark. Note that both numerator and denominator in expression (4) will assume very large values even for polynomials of moderate degree. This can lead to difficulties when evaluating the expression on a computer; the registers will overflow. A common countermeasure is to use the log-gamma function,

$$\frac{\prod_{i=1}^n \Gamma(k_i)}{\Gamma(\sum_{i=1}^n k_i)} = \exp \left(\sum_{i=1}^n \ln \Gamma(k_i) - \ln \Gamma \left(\sum_{i=1}^n k_i \right) \right),$$

but a simpler and arguably more elegant solution is to use the recurrence (5). This holds true for all such expressions in this note.

***n*-dimensional unit sphere**

$$U_n = \left\{ (x_1, \dots, x_n) : \sum_{i=1}^n x_i^2 = 1 \right\}$$

See also [2].

- Volume.

$$|U_n| = \frac{n\sqrt{\pi}^n}{\Gamma(\frac{n}{2} + 1)} = \begin{cases} 2 & \text{if } n = 1 \\ 2\pi & \text{if } n = 2 \\ |U_{n-2}| \times \frac{2\pi}{n-2} & \text{otherwise} \end{cases} \quad (6)$$

- Monomial integral [1].

$$\begin{aligned}
I_{k_1, \dots, k_n} &= \int_{U_n} x_1^{k_1} \dots x_n^{k_n} \\
&= \frac{2 \prod_{i=1}^n \Gamma\left(\frac{k_i+1}{2}\right)}{\Gamma\left(\sum_{i=1}^n \frac{k_i+1}{2}\right)}
\end{aligned} \tag{7}$$

$$= \begin{cases} 0 & \text{if any } k_i \text{ is odd} \\
|U_n| & \text{if all } k_i = 0 \\
I_{k_1, \dots, k_{i_0}-2, \dots, k_n} \times \frac{k_{i_0}-1}{n-2+\sum_{i=1}^n k_i} & \text{if } k_{i_0} > 0 \end{cases} \tag{8}$$

n -dimensional unit ball

$$S_n = \left\{ (x_1, \dots, x_n) : \sum_{i=1}^n x_i^2 \leq 1 \right\}$$

- Volume.

$$|S_n| = \frac{\sqrt{\pi}^n}{\Gamma(\frac{n}{2} + 1)} = \begin{cases} 1 & \text{if } n = 0 \\
2 & \text{if } n = 1 \\
|S_{n-2}| \times \frac{2\pi}{n} & \text{otherwise} \end{cases} \tag{9}$$

- Monomial integral [1].

$$\begin{aligned}
I_{k_1, \dots, k_n} &= \int_{S_n} x_1^{k_1} \dots x_n^{k_n} \\
&= \frac{2^{n+p}}{n+p} |S_n| = \begin{cases} 0 & \text{if any } k_i \text{ is odd} \\
|S_n| & \text{if all } k_i = 0 \\
I_{k_1, \dots, k_{i_0}-2, \dots, k_n} \times \frac{k_{i_0}-1}{n-2+p} & \text{if } k_{i_0} > 0 \end{cases}
\end{aligned} \tag{10}$$

with $p = \sum_{i=1}^n k_i$.

n -dimensional unit ball with Gegenbauer weight

$\lambda > -1$. (Compare with (9) for $\lambda = 0$.) See A.1 for a proof.

- Volume.

$$\begin{aligned}
|G_n^\lambda| &= \int_{S^n} \left(1 - \sum_{i=1}^n x_i^2 \right)^\lambda \\
&= \frac{\Gamma(1+\lambda) \sqrt{\pi}^n}{\Gamma(1+\lambda + \frac{n}{2})} = \begin{cases} 1 & \text{for } n = 0 \\
B(\lambda + 1, \frac{1}{2}) & \text{for } n = 1 \\
|G_{n-2}^\lambda| \times \frac{2\pi}{2\lambda+n} & \text{otherwise} \end{cases}
\end{aligned} \tag{11}$$




Figure 1: The volumes of the n -dimensional ball (and sphere) mysteriously peak at 5 (and 7, respectively). The recurrence relations (6) and (9) make it obvious why: The factor $\frac{2\pi}{n}$ ($\frac{2\pi}{n-2}$) becomes smaller than 1.

- Monomial integration.

$$\begin{aligned}
 I_{k_1, \dots, k_n} &= \int_{S^n} x_1^{k_1} \cdots x_n^{k_n} \left(1 - \sum_{i=1}^n x_i^2\right)^\lambda \\
 &= \frac{\Gamma(1 + \lambda) \prod_{i=1}^n \Gamma\left(\frac{k_i+1}{2}\right)}{\Gamma\left(1 + \lambda + \sum_{i=1}^n \frac{k_i+1}{2}\right)} \quad (12)
 \end{aligned}$$

$$= \begin{cases} 0 & \text{if any } k_i \text{ is odd} \\ |G_n^\lambda| & \text{if all } k_i = 0 \\ I_{k_1, \dots, k_{i_0}-2, \dots, k_n} \times \frac{k_{i_0}-1}{2\lambda+n+\sum_{i=1}^n k_i} & \text{if } k_{i_0} > 0 \end{cases} \quad (13)$$

n -dimensional unit ball with Chebyshev-1 weight

Gegenbauer with $\lambda = -\frac{1}{2}$.

- Volume.

$$\begin{aligned}
 |G_n^{-1/2}| &= \int_{S^n} \frac{1}{\sqrt{1 - \sum_{i=1}^n x_i^2}} \\
 &= \frac{\sqrt{\pi}^{n+1}}{\Gamma\left(\frac{n+1}{2}\right)} = \begin{cases} 1 & \text{if } n = 0 \\ \pi & \text{if } n = 1 \\ |G_{n-2}^{-1/2}| \times \frac{2\pi}{n-1} & \text{otherwise} \end{cases} \quad (14)
 \end{aligned}$$

- Monomial integration.

$$\begin{aligned}
I_{k_1, \dots, k_n} &= \int_{S^n} \frac{x_1^{k_1} \dots x_n^{k_n}}{\sqrt{1 - \sum_{i=1}^n x_i^2}} \\
&= \frac{\sqrt{\pi} \prod_{i=1}^n \Gamma\left(\frac{k_i+1}{2}\right)}{\Gamma\left(\frac{1}{2} + \sum_{i=1}^n \frac{k_i+1}{2}\right)} \tag{15}
\end{aligned}$$

$$= \begin{cases} 0 & \text{if any } k_i \text{ is odd} \\ |G_n^{-1/2}| & \text{if all } k_i = 0 \\ I_{k_1, \dots, k_{i_0}-2, \dots, k_n} \times \frac{k_{i_0}-1}{n-1+\sum_{i=1}^n k_i} & \text{if } k_{i_0} > 0 \end{cases} \tag{16}$$

n -dimensional unit ball with Chebyshev-2 weight

Gegenbauer with $\lambda = +\frac{1}{2}$.

- Volume.

$$\begin{aligned}
|G_n^{+1/2}| &= \int_{S^n} \sqrt{1 - \sum_{i=1}^n x_i^2} \\
&= \frac{\sqrt{\pi}^{n+1}}{2\Gamma\left(\frac{n+3}{2}\right)} = \begin{cases} 1 & \text{if } n = 0 \\ \frac{\pi}{2} & \text{if } n = 1 \\ |G_{n-2}^{+1/2}| \times \frac{2\pi}{n+1} & \text{otherwise} \end{cases} \tag{17}
\end{aligned}$$

- Monomial integration.

$$\begin{aligned}
I_{k_1, \dots, k_n} &= \int_{S^n} x_1^{k_1} \dots x_n^{k_n} \sqrt{1 - \sum_{i=1}^n x_i^2} \\
&= \frac{\sqrt{\pi} \prod_{i=1}^n \Gamma\left(\frac{k_i+1}{2}\right)}{2\Gamma\left(\frac{3}{2} + \sum_{i=1}^n \frac{k_i+1}{2}\right)} \tag{18}
\end{aligned}$$

$$= \begin{cases} 0 & \text{if any } k_i \text{ is odd} \\ |G_n^{+1/2}| & \text{if all } k_i = 0 \\ I_{k_1, \dots, k_{i_0}-2, \dots, k_n} \times \frac{k_{i_0}-1}{n+1+\sum_{i=1}^n k_i} & \text{if } k_{i_0} > 0 \end{cases} \tag{19}$$

n -dimensional generalized Laguerre volume

$\alpha > -1$. See A.2 for a proof.

- Volume.

$$\begin{aligned}
V_n &= \int_{\mathbb{R}^n} \left(\sqrt{x_1^2 + \cdots + x_n^2} \right)^\alpha \exp \left(-\sqrt{x_1^2 + \cdots + x_n^2} \right) \\
&= \frac{2\sqrt{\pi}^n \Gamma(n + \alpha)}{\Gamma(\frac{n}{2})} = \begin{cases} 2\Gamma(1 + \alpha) & \text{if } n = 1 \\ 2\pi\Gamma(2 + \alpha) & \text{if } n = 2 \\ V_{n-2} \times \frac{2\pi(n+\alpha-1)(n+\alpha-2)}{n-2} & \text{otherwise} \end{cases} \quad (20)
\end{aligned}$$

- Monomial integration.

$$\begin{aligned}
I_{k_1, \dots, k_n} &= \int_{\mathbb{R}^n} x_1^{k_1} \cdots x_n^{k_n} \left(\sqrt{x_1^2 + \cdots + x_n^2} \right)^\alpha \exp \left(-\sqrt{x_1^2 + \cdots + x_n^2} \right) \\
&= \frac{2\Gamma(\alpha + n + \sum_{i=1}^n k_i) \left(\prod_{i=1}^n \Gamma\left(\frac{k_i+1}{2}\right) \right)}{\Gamma\left(\sum_{i=1}^n \frac{k_i+1}{2}\right)} \quad (21)
\end{aligned}$$

$$= \begin{cases} 0 & \text{if any } k_i \text{ is odd} \\ V_n & \text{if all } k_i = 0 \\ I_{k_1, \dots, k_{i_0}-2, \dots, k_n} \times \frac{(\alpha+n+p-1)(\alpha+n+p-2)(k_{i_0}-1)}{n+p-2} & \text{if } k_{i_0} > 0 \end{cases} \quad (22)$$

with $p = \sum_{k=1}^n k_i$.

***n*-dimensional Hermite (physicists')**

- Volume.

$$\begin{aligned}
V_n &= \int_{\mathbb{R}^n} \exp \left(-(x_1^2 + \cdots + x_n^2) \right) \\
&= \sqrt{\pi}^n = \begin{cases} 1 & \text{if } n = 0 \\ \sqrt{\pi} & \text{if } n = 1 \\ V_{n-2} \times \pi & \text{otherwise} \end{cases} \quad (23)
\end{aligned}$$

- Monomial integration.

$$\begin{aligned}
I_{k_1, \dots, k_n} &= \int_{\mathbb{R}^n} x_1^{k_1} \cdots x_n^{k_n} \exp \left(-(x_1^2 + \cdots + x_n^2) \right) \\
&= \prod_{i=1}^n \frac{(-1)^{k_i} + 1}{2} \times \Gamma\left(\frac{k_i + 1}{2}\right) \quad (24)
\end{aligned}$$

$$= \begin{cases} 0 & \text{if any } k_i \text{ is odd} \\ V_n & \text{if all } k_i = 0 \\ I_{k_1, \dots, k_{i_0}-2, \dots, k_n} \times \frac{k_{i_0}-1}{2} & \text{if } k_{i_0} > 0 \end{cases} \quad (25)$$

***n*-dimensional Hermite (probabilists')**

- Volume.

$$V_n = \frac{1}{\sqrt{2\pi}^n} \int_{\mathbb{R}^n} \exp\left(-\frac{1}{2}(x_1^2 + \dots + x_n^2)\right) = 1 \quad (26)$$

- Monomial integration.

$$\begin{aligned} I_{k_1, \dots, k_n} &= \frac{1}{\sqrt{2\pi}^n} \int_{\mathbb{R}^n} x_1^{k_1} \dots x_n^{k_n} \exp\left(-\frac{1}{2}(x_1^2 + \dots + x_n^2)\right) \\ &= \prod_{i=1}^n \frac{(-1)^{k_i} + 1}{2} \times \frac{2^{\frac{k_i+1}{2}}}{\sqrt{2\pi}} \Gamma\left(\frac{k_i+1}{2}\right) \end{aligned} \quad (27)$$

$$= \begin{cases} 0 & \text{if any } k_i \text{ is odd} \\ V_n & \text{if all } k_i = 0 \\ I_{k_1, \dots, k_{i_0}-2, \dots, k_n} \times (k_{i_0} - 1) & \text{if } k_{i_0} > 0 \end{cases} \quad (28)$$

A. Some proofs

A.1. Gegenbauer

Proof.

$$\begin{aligned} \int_{\mathbb{R}^n} x^{k_1} \dots x^{k_n} \left(1 - \sum_{i=1}^n x_i^2\right)^\lambda dx &= \int_{S_n} \int_0^1 r^{n-1} r^{\sum k_i} (1 - r^2)^\lambda x'^{k_1} \dots x'^{k_n} dr d\sigma(x') \\ &= \int_0^1 r^{n-1} r^{\sum k_i} (1 - r^2)^\lambda dr \times \int_{S_n} x'^{k_1} \dots x'^{k_n} d\sigma(x') \end{aligned}$$

with $x'_i = x_i/r$. The one-dimensional integral in r can be evaluated explicitly such that, with the spherical integral taken from (7),

$$\begin{aligned} I_{k_1, \dots, k_n} &= \frac{\Gamma\left(\frac{n+\sum k_i}{2}\right) \Gamma(1+\lambda)}{2\Gamma\left(\frac{n+\sum k_i}{2} + \lambda + 1\right)} \times \frac{2 \prod_{i=1}^n \Gamma\left(\frac{k_i+1}{2}\right)}{\Gamma\left(\sum_{i=1}^n \frac{k_i+1}{2}\right)} \\ &= \frac{\Gamma(1+\lambda) \prod_{i=1}^n \Gamma\left(\frac{k_i+1}{2}\right)}{\Gamma\left(\sum \frac{k_i+1}{2} + \lambda + 1\right)}. \end{aligned}$$

□

A.2. Generalized Laguerre

Proof.

$$\begin{aligned} \int_{\mathbb{R}^n} x^{k_1} \dots x^{k_n} r^\alpha \exp(-r) dx &= \int_{S_n} \int_0^\infty r^{n-1} r^{\sum k_i} r^\alpha \exp(-r) x'^{k_1} \dots x_n'^{k_n} dr d\sigma(x') \\ &= \int_0^\infty r^{n-1} r^{\sum k_i} r^\alpha \exp(-r) dr \times \int_{S_n} x'^{k_1} \dots x_n'^{k_n} d\sigma(x') \end{aligned}$$

with $x'_i = x_i/r$. The one-dimensional integral in r can be evaluated explicitly such that, with the spherical integral taken from (7),

$$I_{k_1, \dots, k_n} = \Gamma\left(\alpha + n + \sum k_i\right) \times \frac{2 \prod_{i=1}^n \Gamma\left(\frac{k_i+1}{2}\right)}{\Gamma\left(\sum_{i=1}^n \frac{k_i+1}{2}\right)}.$$

□

References

- [1] Gerald B. Folland. How to integrate a polynomial over a sphere. *The American Mathematical Monthly*, 108(5):446–448, May 2001.
- [2] Michael Hartl. The tau manifesto, 2010. URL:<https://tauday.com/tau-manifesto>.