

CMS at LHC

Diego Baron*¹

* Instituto de Física, Universidad de Antioquia, Medellín, Colombia

Synopsis For experimental particle physicists it is crucial to understand how an experiment like LHC is carried out, in order to know how and where to search for new physics. This seminar is focused in explaining how the LHC and in particular the CMS experiment works, we are going to describe the process of collision, from the acceleration chain of protons to what kind of detectors are mounted in order to reconstruct the collisions and trajectories of particles. We are also going to describe the limitations of the CMS detector and how radiation damage involves constant recalibration of the systems.

After the Superconducting Super Collider was canceled, the 16 of December of 1994 the Large Hadron Collider (LHC) was approved and thus turned into the biggest and most important particle physics experiment. LHC uses the same 27 km tunnel that was caved for LEP (Large Electron-Positron Collider), the LHC depth ranges from 50 to 175 m and is capable of accelerating protons and lead ions to an energy of 7 TeV per beam.

Collisions in this particle accelerator occur in four points, at each one, is located one of the four principal experiments conducted by CERN: ATLAS, CMS, ALICE and LHCb. We are going to put our efforts into explaining the Compact Muon Solenoid (CMS) experiment, it is the second biggest experiment on LHC and currently Universidad de Antioquia is part of the collaboration with the participation of 3 persons.

CMS is a detector located at point 5 of LHC ring, the detector has a diameter of 15m and a longitude of 29m, it weighs 14000 Tons. It is called "Compact Solenoid" because some of the detection systems are inside of its 367.5 m³ solenoid capable of producing a 3.8 T magnetic field and "Muon" due to its systems designed to reconstruct muon trajectories with very high precision [1,2,3].

The main subsystems of the CMS detector are: the tracker system, the superconducting solenoid, the electromagnetic calorimeter (ECAL), the hadronic calorimeter (HCAL) and the muon chambers (a 3-D representation is shown in Figure 1). All these systems produce an extraordinary amount of data (1 PB/s) and not all events are interesting, so an "intelligent" filter must be applied, the CMS detector has a system called the Trigger system that does this job in two stages and achieves a huge reduction of the data, by making the already mentioned selection [4].

The motivation of this talk is to present the LHC and CMS functioning, this is important because when an analysis of some particle physics model is going to be done, particle physicists have to know well what they can measure and with how many precision. As a really simple example, neutrinos escape from the detector, so you cannot look for their trajectories, another important task where it must be known very well how CMS operates, is making an accurate simulation of the detector, this has to be done every time before performing a realistic search in the detector.




Figure 1. CMS detector 3-D representation showing each of its subsystems.

References

- [1] Jose David Ruiz, 2015, Search for a vector-like quark T decaying into top+Higgs in single production mode in full hadronic final state using CMS data collected at 8 TeV.
- [2] Tanja Rommelskirchen, 2006, Study of Muonic Decays of Doubly Charged Higgs Bosons with the CMS Detector.
- [3] Werner Herr and Bruno Muratori, Concept of luminosity.
- [4] <https://cms.cern/>.

¹E-mail: diego.baron@udea.edu.co