

# ggplot\_intro

Diego Perez

2025-02-24

## Setting up my enviroment

Notes: setting up R enviroment by loading tidyverse and 'palmerpenguins' packages

```
library(tidyverse)
```

```
## -- Attaching core tidyverse packages ----- tidyverse 2.0.0 --
## v dplyr      1.1.2      v readr      2.1.4
## v forcats    1.0.0      v stringr   1.5.0
## v ggplot2    3.4.2      v tibble    3.2.1
## v lubridate  1.9.2      v tidyr     1.3.0
## v purrr      1.0.1
## -- Conflicts ----- tidyverse_conflicts() --
## x dplyr::filter() masks stats::filter()
## x dplyr::lag()     masks stats::lag()
## i Use the conflicted package (<http://conflicted.r-lib.org/>) to force all conflicts to become errors
```

```
library(palmerpenguins)
```

## Visualization


Displaying series of visualizations.

### Flipper and body mass in green

Here I displayed the reationship between flipper length and body mass.

```
ggplot(data=penguins,aes(x=flipper_length_mm,y=body_mass_g))+
  geom_point(color="green")
```

```
## Warning: Removed 2 rows containing missing values (`geom_point()`).
```




## Flipper and body mass by species

Flipper length and body mass are broken down by species

```
ggplot(data=penguins,aes(x=flipper_length_mm,y=body_mass_g))+  
  geom_point(aes(shape=species))
```

```
## Warning: Removed 2 rows containing missing values (`geom_point()`).
```



### Flipper and body mass by species and sex

Here we plot flipper length and body mass and break it down by species and sex.

```
ggplot(data=penguins,aes(x=flipper_length_mm,y=body_mass_g))+  
  geom_point(aes(color=species,  
                 shape=species))+  
  facet_wrap(~sex)
```

```
## Warning: Removed 2 rows containing missing values (`geom_point()`).
```

