



# Exploring the Photoactive Properties of MXenes for Water Splitting



UNIVERSITAT DE  
BARCELONA

Diego Ontiveros, Francesc Viñes, Carme Sousa

Institut de Química Teòrica i Computacional (IQTCUB),  
Departament de Ciència de Materials i Química Física, Universitat de Barcelona.



Institut de Química Teòrica  
i Computacional  
UNIVERSITAT DE BARCELONA





## General Properties:

- An effective photocatalyst needs good band alignment, efficient charge separation, minimal VBM/CBM overlap, anisotropic carrier mobilities, strong visible light absorption, and high STH efficiency.
- Cohesive energy, phonons → energetical and dynamical stability.

MXene	$E_{\text{coh}}$ (eV/at.)	$E_{\text{g}}^{\text{PBE0}}$ (eV)	$E_{\text{opt}}$ (eV)	Overlap (%)	$\eta_{\text{STH}}$ (%)
Zr <sub>2</sub> CO <sub>2</sub>	-7.68	2.26	2.87	27.9	2.5
Sc <sub>2</sub> CS <sub>2</sub>	-5.52	3.23	3.34	35.2	0.9
Y <sub>2</sub> CS <sub>2</sub>	-5.52	3.42	2.93	38.5	2.7
Sc <sub>2</sub> CSe <sub>2</sub>	-5.16	2.75	3.15	31.9	1.4
Y <sub>2</sub> CSe <sub>2</sub>	-5.17	3.21	2.99	34.5	2.3
Sc <sub>2</sub> CCl <sub>2</sub>	-5.41	2.48	2.30	26.2	11.0
Y <sub>2</sub> CCl <sub>2</sub>	-5.42	2.44	1.89	31.7	21.3
Sc <sub>2</sub> CBr <sub>2</sub>	-5.09	2.31	2.31	27.1	10.9
Y <sub>2</sub> CBr <sub>2</sub>	-5.13	2.36	1.85	30.7	22.6
Y <sub>2</sub> Cl <sub>2</sub>	-4.77	1.94	1.79	25.2	12.2


## Band Alignment

- $pH = 0$ : ✓ band alignment for all cases, excepting Sc<sub>2</sub>CSe<sub>2</sub> (on H<sub>x</sub> surface).
- $pH = 7$ : Some halide-terminated MXenes become unable to photocatalyze HER.
- H<sub>MX</sub>: Janus → intrinsic  $\vec{E} \rightarrow e^-/h^+$  separation.




## Optical Absorption

- H<sub>M</sub>: Good optical absorption in visible range.
- H<sub>MX</sub>: Optical absorption shifted to UV.
- $E_{\text{opt}}(\text{Y}) < E_{\text{opt}}(\text{Sc})$ .
- BSE → Exciton →  $E_{\text{b}} \approx 0.3 - 0.7$  eV.
- STH efficiency: halide (11–23%) > chalcogen (1–3%).




## Electronic Structure:

- Semiconductors  $E_{\text{g}} \approx$  visible range.
- H<sub>MX</sub> MXene structures:  $E_{\text{g}} \approx$  UV.
- Indirect  $\Gamma \rightarrow \text{M}$  (H<sub>M</sub>) or  $\Gamma \rightarrow \text{K}$  (H<sub>MX</sub>) transitions.
- VB → C and M atoms, and T at lower energies.
- CB → d orbitals of M.




## Charge Density:

- Overlap (VBM, CBM)  $\approx 25 - 38$  %.
- H<sub>M</sub>: VBM → p(C) orbitals, CBM → M layers.
- H<sub>MX</sub>: VBM → H<sub>x</sub> face, CBM → M-C layers.
- asymmetry in charge distribution.



## Charge Carrier Mobility:

- Along  $x$  (zigzag) and  $y$  (armchair).
- Anisotropic electron carrier mobility,  $\mu_e^x \neq \mu_e^y$ .
- H<sub>MX</sub>:  $\mu_e^x < \mu_e^y$ , H<sub>M</sub>:  $\mu_e^x > \mu_e^y$ .
- Isotropic hole carrier mobility,  $\mu_h^x \approx \mu_h^y$ .
- Except for Sc<sub>2</sub>CBr<sub>2</sub> → Asymmetric anisotropy both in  $e$  and  $h$  (charge separation ✓)



## CONCLUSIONS ✓

The photoactive potential of a group of 10 MXenes has been deeply explored using DFT calculations, showcasing robust stability, high charge carrier mobilities, strong visible light absorption, and promising solar-to-hydrogen efficiency. These features make them leading candidates for efficient water splitting photocatalysis, with H<sub>M</sub> structures generally outperforming H<sub>MX</sub> structures.

## REFERENCES

- [1] *Adv. Mater.* 2011, 23, 4248–4253.
- [2] *J. Mater. Chem. A*, 2023, 11, 13754–13764.
- [3] *Energy Environ. Mater.*, 2024, 0, e12774.

