



Bookmarks

- [Unit 0: Overview](#)
- [Entrance Survey](#)
- [Unit 1: Probability models and axioms](#)
- ▾ [Unit 2: Conditioning and independence](#)

[Unit overview](#)

[Lec. 2: Conditioning and Bayes' rule](#)

[Exercises 2 due Feb 2, 2017](#)
20:59 ART

[Lec. 3: Independence](#)

[Exercises 3 due Feb 2, 2017](#)
20:59 ART

[Solved problems](#)**[Problem Set 2](#)**

[Problem Set 2 due Feb 2, 2017](#)
20:59 ART

- [Unit 3: Counting](#)

Unit 2: Conditioning and independence > Lec. 2: Conditioning and Bayes' rule > Exercise: Total probability theorem

Exercise: Total probability theorem

[Bookmark this page](#)

Exercise: Total probability theorem

2.0/2.0 points (graded)

We have an infinite collection of biased coins, indexed by the positive integers. Coin i has probability 2^{-i} of being selected. A flip of coin i results in Heads with probability 3^{-i} . We select a coin and flip it. What is the probability that the result is Heads? The

geometric sum formula may be useful here: $\sum_{i=1}^{\infty} \alpha^i = \frac{\alpha}{1-\alpha}$, when $|\alpha| < 1$.

The probability that the result is Heads is:



Answer: 0.2

Answer:

We think of the selection of coin i as scenario/event A_i . By the total probability theorem, for the case of infinitely many scenarios,

$$\mathbf{P}(\text{Heads}) = \sum_{i=1}^{\infty} \mathbf{P}(A_i) \mathbf{P}(\text{Heads} \mid A_i) = \sum_{i=1}^{\infty} 2^{-i} 3^{-i} = \sum_{i=1}^{\infty} (1/6)^i = \frac{1/6}{1 - (1/6)} = \frac{1/6}{5/6} = \frac{1}{5}$$

[Submit](#)

You have used 1 of 10 attempts

© All Rights Reserved



© 2012-2017 edX Inc. All rights reserved except where noted. EdX, Open edX and the edX and Open EdX logos are registered trademarks or trademarks of edX Inc.

POWERED BY
OPENedX

