



Bookmarks

- ▶ [Unit 0: Overview](#)
- ▶ [Entrance Survey](#)
- ▶ [Unit 1: Probability models and axioms](#)
- ▶ [Unit 2: Conditioning and independence](#)

▼ [Unit 3: Counting](#)

[Lec. 4: Counting](#)

[Exercises 4 due Feb 9, 2017 20:59 ART](#)

[Solved problems](#)

[Problem Set 3](#)

[Problem Set 3 due Feb 9, 2017 20:59 ART](#)

- ▶ [Unit 4: Discrete random variables](#)

Unit 3: Counting > Lec. 4: Counting > Exercise: Coin tossing

Exercise: Coin tossing

[Bookmark this page](#)

Exercise: Coin tossing

2/2 points (graded)

Use the second method in the preceding segment to find the probability that the 6th toss out of a total of 10 tosses is Heads, given that there are exactly 2 Heads out of the 10 tosses. As in the preceding segment, continue to assume that all coin tosses are independent and that each coin toss has the same fixed probability of Heads. (In this and subsequent questions, your answer should be a number. Do not enter '!' or combinations in your answer.)

✓ Answer: 0.2

Answer:

The conditional universe consists of sequences of length 10 that contain exactly 2 Heads. There are

$$\binom{10}{2} = \frac{10!}{8!2!} = \frac{10 \cdot 9}{2} = 45$$

such sequences. Out of these 45 sequences, how many have the property that the 6th toss was Heads? There are 9 sequences with this property: the 6th toss is fixed to be Heads, and the other Heads can be any of the remaining 9 tosses. Therefore, the desired conditional probability is $9/45 = 1/5$.

You have used 2 of 10 attempts

✓ Correct (2/2 points)



© 2012-2017 edX Inc. All rights reserved except where noted. EdX, Open edX and the edX and Open EdX logos are registered trademarks or trademarks of edX Inc.

